

(43) International Publication Date
9 February 2017 (09.02.2017)(51) International Patent Classification:
A61K 35/13 (2015.01) *G01N 33/574* (2006.01)
A61P 35/00 (2006.01)(21) International Application Number:
PCT/IB2016/054584(22) International Filing Date:
29 July 2016 (29.07.2016)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/199,571 31 July 2015 (31.07.2015) US

(71) Applicant: VASCULAR BIOGENICS LTD. [IL/IL]; 6 Jonathan Netanyahu Street, 60376 Or Yehuda (IL).

(72) Inventors: MENDEL, Itzhak; Tzipora Tov 5/8 Street, 76349 Rehovot (IL). PROPHETA-MEIRAN, Oshrat; Rosh Pina 2/4 Street, 49723 Petah Tikva (IL). SALEM, Yaniv; Levy Eshkol 37/7 Street, 55510 Kyriat Ono (IL). SHOHAM, Anat; 56 Derech-Magdiel Street, 45342 Hod Hasharon (IL). YACOV, Niva; Hadar Yosef 22/15 Street, 69364 Tel Aviv (IL). BREITBART, Eyal; 27 HaYitzhar Street, 73127 Hashmonaim (IL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,


AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: MOTILE SPERM DOMAIN CONTAINING PROTEIN 2 AND CANCER

FIG. 1

(57) **Abstract:** Disclosed herein are methods of treating, reducing the incidence of, or preventing one or more activities in or of a cancer cell, methods of treating, reducing the incidence of, or preventing migration or metastasis of a cancer cell, methods of treating, reducing the incidence of, or preventing a cancer by reducing tumor associated macrophages (TAMs) or their migration, and methods of treating, reducing the incidence of, or preventing a cancer (including metastatic cancer), for example, with an inhibitor of Motile Sperm Domain containing Protein 2 (MOSPD2). Also disclosed are inhibitors of MOSPD2 (e.g., anti-MOSPD2 antibodies or antigen binding fragments thereof) and pharmaceutical compositions containing MOSPD2 inhibitors. Also disclosed are methods for the prediction, diagnosis, or prognosis of cancer, cancer metastasis, tumor progression, or tumor invasiveness in a subject.

MOTILE SPERM DOMAIN CONTAINING PROTEIN 2 AND CANCER

FIELD OF THE INVENTION

[0001] This invention relates to the treatment, prevention, or reduction of incidence of cancer and metastasis, for example, methods of treating, preventing, or reducing the incidence of one or more activities in or of a cancer cell, methods of treating, preventing, or reducing the incidence of migration or metastasis of a cancer cell, methods of treating, preventing, or reducing the incidence of cancer by regulating migration of tumor associated macrophages (TAMs), and methods of treating, preventing, or reducing the incidence of cancer (including metastatic cancer), with an inhibitor of a Motile Sperm Domain containing Protein 2 (MOSPD2). The invention also relates to pharmaceutical compositions comprising one or more inhibitors of MOSPD2, and to polypeptide inhibitors of MOSPD2 such as antibodies or antigen binding fragments thereof. The invention also relates to methods for the prediction, diagnosis, or prognosis of cancer, cancer metastasis, tumor progression, or tumor invasiveness in a subject.

BACKGROUND OF THE INVENTION

[0002] Metastasis, the spread of cancer cells from their tissue of origin to other organs, is a result of a multi-step process that involves a number of molecules. Evidence suggests that chemokines and chemokine receptors play an important role in tumor metastasis. Chemokines are small molecules that induce directional cell migration through interaction with their cognate receptors. Binding of chemokines to chemokine receptors activates signaling pathways such as the MAPK/ERK and PI3K/AKT pathways, resulting in phosphorylation of ERK and AKT, respectively.

[0003] Motile Sperm Domain containing Protein 2 (MOSPD2) is a 518-amino acid long, highly conserved protein with 90% homology between human and mouse. Bioinformatic analyses indicate that MOSPD2 contains a CRAL-TRIO region, named after the cellular retinaldehyde-binding protein (CRALBP) and the TRIO protein. MOSPD2 also contains a structurally related region to the nematode major sperm protein and one transmembrane region. A biological function for MOSPD2 has not yet been described. As detailed herein, the inventors found that MOSPD2 is essential for the migration of certain cells

(e.g., monocytes and various cancer cells) towards different chemokines (e.g., Epidermal Growth Factor (EGF)).

SUMMARY OF THE INVENTION

[0004] The invention, in some embodiments, relates to methods of treating, preventing, or reducing the incidence of metastasis of a cancer cell with an inhibitor of a Motile Sperm Domain containing Protein 2 (MOSPD2). In some embodiments, the methods include administering to a subject in need thereof an effective amount of an inhibitor of MOSPD2. In some embodiments, the methods include contacting the cancer cell with an effective amount of an inhibitor of MOSPD2. In other embodiments, the MOSPD2 is expressed by the cancer cell.

[0005] In other embodiments, the invention relates to methods of inhibiting or preventing one or more activities in or of a cancer cell with an inhibitor of MOSPD2. In some embodiments, the methods include administering to a subject in need thereof an effective amount of an inhibitor of MOSPD2. In some embodiments, the methods include contacting the cancer cell with an effective amount of an inhibitor of MOSPD2. In some embodiments, the MOSPD2 is expressed by the cancer cell. In other embodiments, the one or more activities is one or more of: MOSPD2 expression, cancer cell migration, monocyte migration associated with tumor growth, a chemokine signaling pathway, a growth factor signaling pathway, epidermal growth factor (EGF) receptor phosphorylation, extracellular-signal-regulated kinase (ERK) phosphorylation, Protein kinase B (AKT) phosphorylation, and Focal Adhesion Kinase (FAK) phosphorylation.

[0006] In other embodiments, the invention relates to methods of treating, preventing, or reducing the incidence of a cancer with an inhibitor of MOSPD2. In some embodiments, the methods include administering to a subject in need thereof an effective amount of an inhibitor of MOSPD2. In some embodiments, the methods include contacting circulating monocytes or tumor associated macrophages with an effective amount of an inhibitor of MOSPD2 to reduce the number of tumor associated macrophages near or within the cancer mass and/or to regulate migration of tumor associated macrophages. In some embodiments, the MOSPD2 is expressed by circulating monocytes or tumor associated macrophages. In some embodiments, administering the inhibitor of MOSPD2 reduces the number of tumor associated macrophages near or within a cancer mass and/or regulates

migration of tumor associated macrophages. In other embodiments, administering the inhibitor of MOSPD2 reduces the number or migration of tumor associated macrophages by at least 10% or more. In some embodiments, the methods include contacting the cancer cells with an effective amount of an inhibitor of MOSPD2. In some embodiments, the MOSPD2 is expressed by the cancer cells.

[0007] In other embodiments, the invention relates to methods of treating, preventing, or reducing the incidence of a metastatic cancer with an inhibitor of MOSPD2. In some embodiments, the methods include administering to a subject in need thereof a therapeutically effective amount of an inhibitor of MOSPD2. In some embodiments, the methods include contacting the metastatic cancer cells with an effective amount of an inhibitor of MOSPD2. In some embodiments, the MOSPD2 is expressed by the metastatic cancer cells.

[0008] In other embodiments, the methods of treating, preventing, or reducing the incidence described herein include administering a therapeutically effective amount of another anticancer drug along with the inhibitor of MOSPD2.

[0009] In other embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment thereof. In some embodiments, the antibody is a polyclonal, monoclonal, murine, human, humanized, or chimeric antibody. In some embodiments, the inhibitor of MOSPD2 is a small molecule, such as an oxidized phospholipid. In one preferred embodiment, the inhibitor of MOSPD2 is VB-201. In some embodiments, the inhibitor of MOSPD2 is an inhibitor of MOSPD2 that is not an oxidized phospholipid. In some embodiments, the inhibitor of MOSPD2 is an inhibitor of MOSPD2 that is not VB-201. In some embodiments, the inhibitor of MOSPD2 binds to MOSPD2 expressed on a cell surface (*e.g.*, a cancer cell surface).

[0010] In other embodiments, the invention also relates to polypeptides that inhibit MOSPD2 expressed by a cancer cell and pharmaceutical compositions containing a polypeptide that inhibits MOSPD2 expressed by a cancer cell. In other embodiments, the polypeptide is an antibody or antigen binding fragment thereof.

[0011] In some embodiments, the invention relates to an isolated antibody or antigen binding fragment thereof that specifically binds to MOSPD2. In other embodiments, the antibody or antigen binding fragment thereof binds to MOSPD2 with an equilibrium dissociation constant (K_D) of from about 10^{-6} M to about 10^{-12} M. In other embodiments,

the MOSPD2 is human MOSPD2. In other embodiments, the antibody or antigen binding fragment thereof specifically binds to one or more of the following amino acid regions of human MOSPD2, numbered according to SEQ ID NO:1: about 508 to about 517, about 501 to about 514, about 233 to about 241, about 509 to about 517, about 212 to about 221, about 13 to about 24, about 505 to about 517, about 505 to about 514, about 89 to about 100, about 506 to about 517, about 233 to about 245, about 504 to about 514, about 128 to about 136, about 218 to about 226, about 15 to about 24, about 83 to about 96, about 42 to about 50, about 462 to 474, about 340 to about 351, about 504 to about 517, about 462 to about 470, about 327 to about 337, about 21 to about 32, about 217 to about 226, about 510 to about 517, about 178 to about 190, about 497 to about 509, about 504 to about 516, about 64 to about 77, about 504 to about 515, about 147 to about 159, about 503 to about 515, about 88 to about 97, about 208 to about 218, about 178 to about 191, about 502 to about 515, about 503 to about 516, about 497 to about 505, about 500 to about 509, about 189 to about 202, about 189 to about 197, about 505 to about 516, about 1 to about 63, about 82 to about 239, about 93 to about 234, about 327 to about 445, about 327 to about 431, and about 497 to about 517.

[0012] In some embodiments, the antibody or antigen binding fragment thereof specifically binds to one or more of the following amino acid regions of human MOSPD2, numbered according to SEQ ID NO:1: about 505 to about 515, about 500 to about 515, about 230 to about 240, about 510 to about 520, about 210 to about 220, about 15 to about 25, about 505 to about 520, about 505 to about 515, about 90 to about 100, about 505 to about 525, about 230 to about 245, about 505 to about 510, about 130 to about 140, about 220 to about 230, about 15 to about 30, about 80 to about 95, about 40 to about 50, about 460 to about 475, about 340 to about 350, about 500 to about 515, about 460 to about 470, about 325 to about 335, about 20 to about 35, about 215 to about 225, about 510 to about 520, about 175 to about 190, about 500 to about 510, about 505 to about 530, about 60 to about 75, about 500 to about 520, about 145 to about 160, about 502 to about 515, about 85 to about 100, about 205 to about 220, about 175 to about 190, about 500 to about 505, about 500 to about 525, about 495 to about 505, about 495 to about 510, about 190 to about 200, about 190 to about 198, about 502 to about 515, about 1 to about 60, about 80 to about 240, about 90 to about 235, about 330 to about 445, about 330 to about 430, and about 495 to about 515.

[0013] In other embodiments, the invention relates to a pharmaceutical composition comprising the antibody or antigen binding fragment, and a pharmaceutically acceptable carrier.

[0014] In other embodiments, the invention relates to a method of treating, preventing, or reducing the incidence of metastasis of a cancer cell, comprising administering to a subject in need thereof a therapeutically effective amount of the antibody or antigen binding fragment thereof or the pharmaceutical composition. In other embodiments, the invention relates to a method of inhibiting or preventing one or more activities in or of a cancer cell, comprising administering to a subject in need thereof a therapeutically effective amount of the antibody or antigen binding fragment thereof or the pharmaceutical composition, wherein the one or more activities is one or more of: MOSPD2 expression, cancer cell migration, monocyte migration associated with tumor growth, a chemokine signaling pathway, a growth factor signaling pathway, EGF receptor phosphorylation, ERK phosphorylation, AKT phosphorylation, and FAK phosphorylation. In other embodiments, the invention relates to a method of treating, preventing, or reducing the incidence of a cancer, comprising administering to a subject in need thereof a therapeutically effective amount of the antibody or antigen binding fragment thereof or the pharmaceutical composition. In other embodiments, the invention relates to a method of treating, preventing, or reducing the incidence of a metastatic cancer, comprising administering to a subject in need thereof a therapeutically effective amount of the antibody or antigen binding fragment thereof or the pharmaceutical composition.

[0015] In other embodiments, the invention relates to methods for the prediction, diagnosis, or prognosis of cancer, cancer metastasis, tumor progression, or tumor invasiveness in a subject.

BRIEF DESCRIPTION OF THE DRAWINGS

[0016] Some embodiments of the invention are herein described, by way of example only, with reference to the accompanying drawings. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of embodiments of the invention.

[0017] FIG. 1 presents images showing test results of cancer cells transduced with sh-control or sh-MOSPD2 lenti-virus particles in a trans-well migration assay towards 10% fetal calf serum (FCS) and EGF (200 ng/ml). MOSPD2 expression by human MDA-231 breast cancer and A2058 melanoma cell lines was silenced using sh-MOSPD2 lenti-virus particles. Western blots in FIG. 1 show decreased MOSPD2 protein expression in cancer cell lines transduced with sh-MOSPD2. FIG. 1 shows that MOSPD2 promotes migration of metastatic breast cancer and melanoma cell lines.

[0018] FIG. 2 presents a graph showing cell proliferation rates for MDA-231 breast cancer cells transduced with sh-control or sh-MOSPD2 lenti-virus particles. MDA-231 cells were seeded as described and collected and counted every 24 hours for three consecutive days. The results are expressed as mean \pm standard deviation of triplicates. These results demonstrate that silencing of MOSPD2 did not affect cell viability or proliferation of MDA-231 cells.

[0019] FIGs. 3A-3C show *in vivo* test results of metastasis of MDA-231 breast cancer cells with or without MOSPD2 being silenced. In FIG. 3A, MDA-231 breast cancer cells transduced with sh-control or sh-MOSPD2 lenti-virus particles were injected (10^6) in the tail vein of SCID mice (n=10/group). Mice were sacrificed on day 28, their lungs harvested for H&E staining, and tumor area was determined. The results shown in FIG. 3A are expressed as mean \pm standard error of measured metastasis size (* p<0.05).

[0020] In FIGs. 3B and 3C, MDA-231 breast cancer cells transduced with sh-control or sh-MOSPD2 lenti-virus particles (n=13 and n=8 respectively) were injected (5×10^6) in the mammary fat pad of SCID mice. Mice were sacrificed on day 56. The ipsilateral inguinal lymph node was excised (FIG. 3B), the lungs were harvested for H&E staining and the tumor area was determined (FIG. 3C). The results shown in FIG. 3C are expressed as mean \pm standard error of measured metastasis size: the tumor area for sh-control transduced cells is 1376.9 ± 752.6 (n=13), compared to 550.0 ± 326.2 (n=8) for sh-MOSPD2 transduced cells..

[0021] FIGs. 3A-3C show that MOSPD2 promotes metastasis of MDA-231 breast cancer cells *in vivo*.

[0022] FIGs. 4A-4E show images comparing MOSPD2 expression levels of various human cancer tissues to those of their respective normal tissue counterparts. Slides containing various normal and cancerous human tissues were stained with control or anti-

MOSPD2 antibody. Cancer tissues that stained positively for MOSPD2 are shown. FIGs. 4A-4E show that MOSPD2 is expressed in various human cancer tissues.

[0023] FIGs. 5A and 5B show the results of cancer cells transduced with control or MOSPD2 CRISPR-CAS9 lenti-virus particles, that were tested in a trans-well migration assay in which cells were seeded at the upper compartment and attracted to the lower compartment using medium supplemented with 10% FCS and EGF (200 ng/ml). The graph shown in FIG. 5A was determined by fluorescence-activated cell sorting (FACS) with results expressed as mean \pm standard deviation of triplicates. The images shown in FIG. 5B are from visual recordation. In FIGs. 5A and 5B, MDA-231 breast cancer cells were transduced with lenti-viral particles with plasmids containing control or MOSPD2 CRISPR-CAS9 system. Western blots show decreased MOSPD2 protein expression (inset). FIGs. 5A and 5B show that CRISPR-CAS9 driven MOSPD2 gene editing inhibits breast cancer cell migration.

[0024] FIG. 5C presents Western blots showing the effect of MOSPD2 silencing by CRISPR-CAS9 driven gene editing on phosphorylation events associated with cell migration. MDA-231 breast cancer cells transduced with control or MOSPD2 CRISPR-CAS9 lenti-virus particles were incubated with 10% FCS and EGF (400 ng/ml) for 10 minutes. Phosphorylation of ERK, AKT and FAK was determined by Western blotting. HSP90 was used for loading control. FIG. 5C shows that MOSPD2 silencing by CRISPR-CAS9 driven gene editing inhibits phosphorylation events associated with cell migration.

[0025] FIG. 5D shows *in vivo* test results of metastasis of MDA-231 breast cancer cells transduced with control or MOSPD2 CRISPR-CAS9 lenti-virus particles. In FIG. 5D, 10^6 CRISPR-control or CRISPR-MOSPD2 lenti-virus transduced MDA-231 cells were injected into the tail vein of 8 weeks old female SCID mice (C.B-17/IcrHsd-Prkdc^{scid}, Harlan Israel). Mice were sacrificed after 3 weeks and their lungs were excised for histopathologic examination. FIG. 5D shows that silencing MOSPD2 by the CRISPR-CAS9 system significantly inhibits the presence of metastatic breast cancer cells in the lungs by more than 95% (metastasis area), with a p-value of 0.002.

[0026] FIG. 6 presents an image of Western Blots showing the effect of VB-201 in inhibiting EGF induced phosphorylation of AKT in MDA-231 cancer cells. As shown in FIG. 6, VB-201 at 10 μ g/ml nearly completely inhibits EGF induced phosphorylation of

AKT, with significant inhibition observed at 5 μ g/ml. HSP90 was used for loading control.

[0027] FIG. 7 lists 17 anti-MOSPD2 F(ab')₂ monoclonal antibody clones that were identified following a primary screen for binding to cells over-expressing MOSPD2. Further analysis of the clones for MOSPD2 binding with enzyme-linked immunosorbent assay (ELISA) identified 12 clones having optical density (O.D.) values greater than 5 times over background (* in FIG. 7).

[0028] FIGs. 8A-8B show binding of two representative anti-MOSPD2 F(ab')₂ monoclonal antibody (mAb) clones to cells overexpressing MOSPD2.

[0029] FIG. 9 shows binding of a representative anti-MOSPD2 F(ab')₂ mAb to MOSPD2 expressed by MDA-231 breast cancer cells.

[0030] FIGs. 10A-10B show anti-MOSPD2 F(ab')₂ mAb binds to MDA-231 cells (FIG. 10A), but does not bind to MOSPD2-silenced MDA-231 cells (FIG. 10B).

[0031] FIGs. 11A-11B show anti-MOSPD2 F(ab')₂ mAb binds to MOSPD2 on A2058 melanoma and HepG2 liver cancer cell lines.

[0032] FIG. 12 shows that incubation of MDA-231 cells with anti-MOSPD2 F(ab')₂ mAb inhibited phosphorylation of EGF receptor (p-EGF-R), AKT (p-AKT) and ERK1/2 (p-ERK1/2).

[0033] FIG. 13 shows anti-MOSPD2 F(ab')₂ mAb significantly inhibited EGF-induced trans-well migration of MDA-231 cells.

[0034] FIGs. 14A-14D show the cellular expression specificity and localization of MOSPD2.

[0035] FIGs. 15A-15C show MOSPD2 is expressed on monocytes that have infiltrated into inflamed tissues.

[0036] FIGs. 16A-16E show MOSPD2 promotes monocyte migration. FIG. 16A shows mRNA and protein expression of MOSPD2 in U937 cells transduced with sh-control or sh-MOSPD2 lentivirus particles. One of at least three experiments is shown. FIG. 16B shows three-hour trans-well migration of U937 cells transduced with sh-control or sh-MOSPD2 lentivirus particles towards RANTES (100 ng/ml). The percent of sh-MOSPD2 transduced cells relative to sh-control transduced migrating cells is presented. One of three experiments is shown. FIG. 16C shows U937 cells transduced with sh-control or sh-MOSPD2 lentivirus particles were incubated for the indicated time (min)

with RANTES, and the phosphorylation of ERK1/2 (p-ERK1/2) and AKT (p-AKT) was evaluated. HSP90 was used as loading control. FIG. 16D shows three-hour trans-well migration of U937 cells transduced with sh-control (sh-cont) or sh-MOSPD2 lentivirus particles towards MCP-3 (100 ng/ml), MCP-1 (100 ng/ml), RANTES (100 ng/ml) and SDF-1 (25 ng/ml). The percent of sh-MOSPD2 relative to sh-control transduced migrating cells is presented. One of three experiments is shown. FIG. 16E shows U937 cells transduced with sh-control (sh-cont) or sh-MOSPD2 lentivirus particles were incubated for 5 min with MCP-3 (100 ng/ml), MCP-1 (100 ng/ml), RANTES (100 ng/ml) and SDF-1 (25 ng/ml), and the phosphorylation of ERK1/2 (p-ERK1/2) and AKT (p-AKT) was evaluated. Tubulin was used as loading control.

[0037] FIGs. 17A-17B show that MOSPD2 does not affect IFN-gamma-induced phosphorylation of STAT1 (p-Stat1) or PMA-mediated phosphorylation of ERK1/2 (p-ERK1/2), respectively, supporting the specificity of the aforementioned MOSPD2 activities.

[0038] FIGs. 18A-18F show histological images of human breast cancer samples from different pathological stages or from normal tissue adjacent to a tumor (normal adjacent tissue; NAT). The slides were stained with anti-MOSPD2 antibody. FIGs 18A-18F show that MOSPD2 expression was associated with the transition of breast cancer cells from locally-restricted tumor to invasive and metastatic tumor.

[0039] FIG. 19 shows scoring of MOSPD2 expression intensity (in a scale of 0-3, where 0 is no expression and 3 is very high expression) in samples from different stages of breast cancer or normal adjacent tissue (NAT) (* p<0.001).

[0040] FIGs. 20A-20D show images comparing MOSPD2 expression level in various normal and cancerous human tissues collected from the colon (FIGs. 20A-20B) or the liver (FIGs. 20C-20D). MOSPD2 was expressed in 67% of colon adenocarcinoma and in 45% of hepatocellular carcinoma samples, while no expression was detected in normal colon and liver tissues.

[0041] FIGs. 21A-21E show images comparing MOSPD2 expression level of normal tissue, NAT and cancerous tissue at different grades collected from human liver. FIGs. 21C-21E show that MOSPD2 staining intensity was increased along with the increase in the tumor grade of hepatocellular carcinoma.

[0042] FIGs. 22A-22B show MOSPD2 scoring of MOSPD2 expression intensity in samples collected from hepatocellular carcinoma. FIG. 22A shows that MOSPD2 expression was significantly increased ($p \leq 0.001$) in samples collected from malignant hepatocellular carcinoma, compared to normal and NAT samples. FIG. 22B shows that MOSPD2 staining intensity increased significantly in correlation with the progression of hepatocellular carcinoma.

[0043] FIG. 23 presents images of Western blots showing that VB-201 binds to MOSPD2 from cell lysate of human CD14 monocytes. Labelled VB-201 or VB-221 (OB201 or OB221) was added to the cell lysate and proteins were precipitated. Samples were run on a gel and blotted against TLR2 and MOSPD2.

[0044] FIG. 24 presents images of Western blots showing that MOSPD2 promoted EGF-induced signaling events in breast cancer cells.

DETAILED DESCRIPTION OF THE INVENTION

[0045] Before explaining embodiments of the invention in detail, it is to be understood that the invention is not limited in its application to the details set forth in the following description or exemplified by the Examples. The invention is capable of other embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

General Definitions

[0046] The terms "comprises", "comprising", "includes", "including", "having", and their conjugates mean "including but not limited to."

[0047] The term "consisting of" means "including and limited to."

[0048] The term "consisting essentially of" means the specified material of a composition, or the specified steps of a method, and those additional materials or steps that do not materially affect the basic characteristics of the material or method.

[0049] The word "exemplary" is used herein to mean "serving as an example, instance or illustration." Any embodiment described as "exemplary" is not necessarily to be construed as preferred or advantageous over other embodiments and/or to exclude the incorporation of features from other embodiments.

[0050] The word "optionally" is used herein to mean "is provided in some embodiments and not provided in other embodiments." Any particular embodiment of the invention can include a plurality of "optional" features unless such features conflict.

[0051] As used herein, the singular form "a", "an" and "the" include plural references unless the context clearly dictates otherwise. For example, the term "a compound" or "at least one compound" may include a plurality of compounds, including mixtures thereof.

[0052] As used herein, the term "about" modifying an amount related to the invention refers to variation in the numerical quantity that can occur, for example, through routine testing and handling; through inadvertent error in such testing and handling; through differences in the manufacture, source, or purity of ingredients employed in the invention; and the like. Whether or not modified by the term "about", the claims include equivalents of the recited quantities. In one embodiment, the term "about" means within 10% of the reported numerical value. In another embodiment, the term "about" means within 5% of the reported numerical value.

[0053] Throughout this application, various embodiments of this invention can be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range, such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 *etc.*, as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

[0054] As used herein the term "method" refers to manners, means, techniques and procedures for accomplishing a given task including, but not limited to, those manners, means, techniques and procedures either known to, or readily developed from known manners, means, techniques and procedures by practitioners of the chemical, pharmacological, biological, biochemical and medical arts.

[0055] As used herein, the term "treating" includes abrogating, substantially inhibiting, slowing or reversing the progression of a condition, substantially ameliorating clinical or

aesthetical symptoms of a condition or substantially preventing the appearance of clinical or aesthetical symptoms of a condition.

[0056] As used herein, "MOSPD2" refers to any polypeptide classified as a Motile Sperm Domain containing Protein 2. Examples of MOSPD2 include, but are not limited to, the polypeptides of SEQ ID NOS:1-4, or any variant thereof (*e.g.*, having a sequence at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to any one of SEQ ID NOS:1-4). Other examples of MOSPD2 include, but are not limited to, a polypeptide encoded by a polynucleotide of any one of SEQ ID NOS:5-8, or any variant thereof (*e.g.*, a polynucleotide having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to any one of SEQ ID NOS:5-8). Polynucleotide sequences encoding MOSPD2 can be codon optimized for expression in a particular organism by methods known in the art. Other examples of MOSPD2 can be identified by searching public databases (*e.g.*, BLAST), as well known to one skilled in the art.

[0057] In any of the embodiments described herein, the MOSPD2 can be MOSPD2 expressed by a cancer cell, *e.g.*, a human cancer cell. Also, in any of the embodiments described herein, the MOSPD2 can be a mammalian MOSPD2 or a human MOSPD2. Non-exclusive listing of types of cancer cells include cells of bladder cancer, breast cancer, colon cancer, rectal cancer, kidney cancer, liver cancer, lung cancer, esophageal cancer, gall-bladder cancer, ovarian cancer, pancreatic cancer, stomach cancer, cervical cancer, thyroid cancer, prostate cancer, skin cancer, hematopoietic cancer, cancer of mesenchymal origin, cancer of central or peripheral nervous system, endometrial cancer, head and neck cancer, glioblastoma, and malignant ascites. In some embodiments, the cancer is a small cell lung cancer or a non-small-cell lung cancer. In some embodiments, the cancer is skin cancer, *e.g.*, squamous cell carcinoma, basal cell cancer, melanoma, dermatofibrosarcoma protuberans, Merkel cell carcinoma, Kaposi's sarcoma, keratoacanthoma, spindle cell tumors, sebaceous carcinomas, microcystic adnexal carcinoma, Paget's disease of the breast, atypical fibroxanthoma, leiomyosarcoma, or angiosarcoma. In some embodiments, the cancer is a hematopoietic cancer of lymphoid lineage, *e.g.*, leukemia, acute lymphocytic leukemia, chronic lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma or Burkitt's lymphoma. In some

embodiments, the cancer is a hematopoietic cancer of myeloid lineage, *e.g.*, fibrosarcoma, rhabdomyosarcoma, soft tissue sarcoma, or bone sarcoma. In some embodiments, the cancer is a cancer of the central or peripheral nervous system, *e.g.*, astrocytoma, neuroblastoma, glioma, or schwannomas. In some embodiments, the cancer is anal cancer, bone cancer, gastrointestinal stromal cancer, gestational trophoblastic disease, Hodgkin's lymphoma, Kaposi sarcoma, keratoacanthoma, malignant mesothelioma, multicentric castleman disease, multiple myeloma and other plasma cell neoplasms, myeloproliferative neoplasms, neuroblastoma, non-Hodgkin's lymphoma, osteosarcoma, ovarian, fallopian tube, or primary peritoneal cancer, penile cancer, retinoblastoma, rhabdomyosarcoma, seminoma, soft tissue sarcoma, stomach (gastric) cancer, testicular cancer, teratocarcinoma, thyroid follicular cancer, vaginal cancer, vulvar cancer, Wilms tumor and other childhood kidney cancers, and xeroderma pigmentosum. In some embodiments, the cancer is bladder cancer, brain cancer (*e.g.*, cerebrum astrocytoma), breast cancer, colon cancer (*e.g.*, colon adenocarcinoma), esophageal cancer (*e.g.*, esophageal adenocarcinoma), lung cancer, skin cancer (*e.g.*, melanoma), tongue cancer (*e.g.*, head and neck (tongue) cell carcinoma), kidney cancer (*e.g.*, kidney clear cell carcinoma), or hepatic cancer (*e.g.*, hepatocellular carcinoma).

[0058] As used herein, "an activity of MOSPD2" or "a MOSPD2 activity" includes any known or herein described function of a Motile Sperm Domain containing Protein 2. Such activities include, for example, regulation of cell migration (*e.g.*, leukocyte, monocyte or cancer cell migration), presence of tumor associated macrophages, chemotaxis, chemokine-induced leukocyte migration, chemokine receptor signaling pathways, growth factor signaling pathways, EGF receptor phosphorylation, ERK phosphorylation, AKT phosphorylation, FAK phosphorylation, or inflammation.

[0059] As used herein, "chemotaxis" refers to the movement of a cell in response to a chemical stimulus. Chemotaxis includes, but is not limited to, the movement of a cancer cell to a chemokine (*e.g.*, an EGF).

[0060] As used herein, "an inhibitor of MOSPD2" and "a MOSPD2 inhibitor" refer to any compound which downregulates an activity of MOSPD2. The inhibitor can be, for example, a polypeptide, DNA, or RNA. Inhibition of MOSPD2 can also occur, for example, by ectopic overexpression of MOSPD2 by infection, and it is intended that an inhibitor of MOSPD2 or a MOSPD2 inhibitor encompasses this type of inhibition. The

inhibitor can also be, for example, a molecule that specifically binds to a MOSPD2 polypeptide, a molecule that specifically binds to a ligand of a MOSPD2 polypeptide, an antisera raised against a MOSPD2 polypeptide, a soluble MOSPD2 polypeptide, or a soluble MOSPD2 polypeptide comprising, consisting essentially of, or consisting of an extracellular domain of a MOSPD2 polypeptide. The inhibitor can also be, for example, an antibody that specifically binds to a MOSPD2 polypeptide or an antigen binding fragment of an antibody that specifically binds to a MOSPD2 polypeptide. The inhibitor can also be, for example, an RNAi, miRNA, siRNA, shRNA, antisense RNA, antisense DNA, decoy molecule, decoy DNA, double-stranded DNA, single-stranded DNA, complexed DNA, encapsulated DNA, viral DNA, plasmid DNA, naked RNA, encapsulated RNA, viral RNA, double-stranded RNA, molecule capable of generating RNA interference, or combinations thereof, that hybridizes to a nucleotide sequence encoding a MOSPD2 polypeptide. The inhibitor can also be, for example, a clustered regularly interspaced short palindromic repeats CRISPR-CAS9 system. CRISPR-CAS9 systems have been described in the literature and can include, for example, CAS9 and a guide RNA. Other gene editing techniques have also been described in the literature and can also be used. The inhibitor can also be a small molecule chemical compound which downregulates an activity of MOSPD2.

[0061] An "antibody" or an "antigen binding fragment" of an antibody include, but are not limited to, polyclonal, monoclonal, murine, human, humanized, or chimeric antibodies, single chain antibodies, epitope-binding fragments, *e.g.*, Fab, Fab' and F(ab')₂, Fd, Fvs, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv), a light chain variable region (VL) or a heavy chain variable region (VH) domain, fragments comprising either a VL or VH domain, and fragments produced by a Fab expression library. An antibody or antigen binding fragment of an antibody can be of any type (*e.g.*, IgG, IgE, IgM, IgD, IgA, and IgY), class (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule. Methods for making an antigen binding fragment of an antibody are known and include, for example, chemical or protease digestion of an antibody.

[0062] A "constant region" of an antibody refers to the constant region of the antibody light chain or the constant region of the antibody heavy chain, either alone or in combination.

[0063] The term "Fc region" is used to define a C-terminal region of an immunoglobulin heavy chain. The "Fc region" may be a native sequence Fc region or a variant Fc region. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy chain Fc region is usually defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The numbering of the residues in the Fc region is that of the EU index as in Kabat. Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md., 1991. The Fc region of an immunoglobulin generally comprises two constant domains, CH2 and CH3.

[0064] By "specifically binds," it is generally meant that an antibody or fragment, variant, or derivative thereof binds to an epitope by its antigen-binding domain, and that the binding entails some complementarity between the antigen binding domain and the epitope. According to this definition, an antibody or fragment, variant, or derivative thereof is said to "specifically bind" to an epitope when it binds to that epitope via its antigen-binding domain more readily than it would bind to a random, unrelated epitope.

[0065] As used herein, an "epitope" refers to a localized region of an antigen to which an antibody can specifically bind. An epitope can be, for example, contiguous amino acids of a polypeptide (linear or contiguous epitope) or an epitope can, for example, come together from two or more non-contiguous regions of a polypeptide or polypeptides (conformational, non-linear, discontinuous, or non-contiguous epitope). In certain embodiments, the epitope to which an antibody binds can be determined by methods described in the literature and herein, *e.g.*, NMR spectroscopy, X-ray diffraction crystallography studies, ELISA assays, hydrogen/deuterium exchange coupled with mass spectrometry (*e.g.*, MALDI mass spectrometry), array-based oligo-peptide scanning assays, and/or mutagenesis mapping (*e.g.*, site-directed mutagenesis mapping).

[0066] The term "percent identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" and "sequence identity" also mean the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods and publicly available resources, including but not limited to those described in: (1) Computational Molecular

Biology (Lesk, A. M., Ed.) Oxford University: NY (1988); (2) Biocomputing: Informatics and Genome Projects (Smith, D. W., Ed.) Academic: NY (1993); (3) Computer Analysis of Sequence Data, Part I (Griffin, A. M., and Griffin, H. G., Eds.) Humania: NJ (1994); (4) Sequence Analysis in Molecular Biology (von Heinje, G., Ed.) Academic (1987); and (5) Sequence Analysis Primer (Gribskov, M. and Devereux, J., Eds.) Stockton: NY (1991).

[0067] A polynucleotide can "hybridize" to another polynucleotide, when a single-stranded form of the nucleic acid fragment can anneal to the other nucleic acid fragment under the appropriate conditions of temperature and solution ionic strength. Hybridization and washing conditions are well known and exemplified, for example, in Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory: Cold Spring Harbor, N.Y. (1989), particularly Chapter 11 and Table 11.1 therein (incorporated herein by reference in its entirety). The conditions of temperature and ionic strength determine the "stringency" of the hybridization. Stringency conditions can be adjusted to screen for moderately similar fragments (such as homologous sequences from distantly related organisms), to highly similar fragments (such as genes that duplicate functional enzymes from closely related organisms). Post-hybridization washes determine stringency conditions. One exemplary set of stringent conditions uses a series of washes starting with 6xSSC, 0.5% SDS at room temperature for 15 min, then repeated with 2xSSC, 0.5% SDS at 45°C for 30 min, and then repeated twice with 0.2xSSC, 0.5% SDS at 50°C for 30 min. Another set of exemplary stringent conditions uses higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2xSSC, 0.5% SDS was increased to 60°C. This set of stringent conditions can be modified to a "highly stringent condition" by adding two final washes in 0.1xSSC, 0.1% SDS at 65°C. An additional exemplary set of stringent conditions include hybridization at 0.1xSSC, 0.1% SDS, 65°C and washes with 2xSSC, 0.1% SDS followed by 0.1xSSC, 0.1% SDS, for example.

[0068] Hybridization requires that the two nucleic acids contain complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible. The appropriate stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of complementation, variables well known in the art. The greater the degree of similarity or homology between two

nucleotide sequences, the greater the value of Tm for hybrids of nucleic acids having those sequences. The relative stability (corresponding to higher Tm) of nucleic acid hybridizations decreases in the following order: RNA:RNA, DNA:RNA, DNA:DNA. For hybrids of greater than 100 nucleotides in length, equations for calculating Tm have been derived (see Sambrook *et al.*, 9.50-9.51). For hybridizations with shorter nucleic acids, *i.e.*, oligonucleotides, the position of mismatches becomes more important, and the length of the oligonucleotide determines its specificity (see Sambrook *et al.*, *supra*, 11.7-11.8). In one embodiment, the length for a hybridizable nucleic acid is at least about 10 nucleotides. In other embodiments, a minimum length for a hybridizable nucleic acid is at least about 15 nucleotides, or at least about 20 nucleotides.

[0069] As used herein throughout, the term "alkyl" refers to a saturated aliphatic hydrocarbon including straight chain and branched chain groups. In some embodiments, the alkyl group has 1 to 20 carbon atoms. Whenever a numerical range; *e.g.*, "1-20", is stated herein, it implies that the group, in this case the alkyl group, may contain 1 carbon atom, 2 carbon atoms, 3 carbon atoms, etc., up to and including 20 carbon atoms. In some embodiments, the alkyl is a medium size alkyl having 1 to 10 carbon atoms. In some embodiments, the alkyl is a lower alkyl having 1 to 4 carbon atoms. The alkyl group may be substituted or unsubstituted. When substituted, the substituent group can be, for example, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroalicyclic, halo, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, sulfinyl, sulfonyl, cyano, nitro, azide, sulfonyl, sulfinyl, sulfonamide, phosphonyl, phosphinyl, oxo, carbonyl, thiocarbonyl, urea, thiourea, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, C-carboxy, O-carboxy, sulfonamido, and amino, as these terms are defined herein.

[0070] A "cycloalkyl" group refers to an all-carbon monocyclic or fused ring (*i.e.*, rings which share an adjacent pair of carbon atoms) group wherein one or more of the rings does not have a completely conjugated pi-electron system. Examples, without limitation, of cycloalkyl groups are cyclopropane, cyclobutane, cyclopentane, cyclopentene, cyclohexane, cyclohexadiene, cycloheptane, cycloheptatriene, and adamantane. A cycloalkyl group may be substituted or unsubstituted. When substituted, the substituent group can be, for example, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroalicyclic, halo, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, sulfinyl, sulfonyl, cyano,

nitro, azide, sulfonyl, sulfinyl, sulfonamide, phosphonyl, phosphinyl, oxo, carbonyl, thiocarbonyl, urea, thiourea, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, C-carboxy, O-carboxy, sulfonamido, and amino, as these terms are defined herein.

[0071] An "alkenyl" group refers to an alkyl group which consists of at least two carbon atoms and at least one carbon-carbon double bond.

[0072] An "alkynyl" group refers to an alkyl group which consists of at least two carbon atoms and at least one carbon-carbon triple bond.

[0073] An "aryl" group refers to an all-carbon monocyclic or fused-ring polycyclic (*i.e.*, rings which share adjacent pairs of carbon atoms) groups having a completely conjugated pi-electron system. Examples, without limitation, of aryl groups are phenyl, naphthalenyl and anthracenyl. The aryl group may be substituted or unsubstituted. When substituted, the substituent group can be, for example, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heteroalicyclic, halo, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, sulfinyl, sulfonyl, cyano, nitro, azide, sulfonyl, sulfinyl, sulfonamide, phosphonyl, phosphinyl, oxo, carbonyl, thiocarbonyl, urea, thiourea, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, C-carboxy, O-carboxy, sulfonamido, and amino, as these terms are defined herein.

[0074] A "heteroaryl" group refers to a monocyclic or fused ring (*i.e.*, rings which share an adjacent pair of atoms) group having in the ring(s) one or more atoms, such as, for example, nitrogen, oxygen and sulfur and, in addition, having a completely conjugated pi-electron system. Examples, without limitation, of heteroaryl groups include pyrrole, furane, thiophene, imidazole, oxazole, thiazole, pyrazole, pyridine, pyrimidine, quinoline, isoquinoline and purine. The heteroaryl group may be substituted or unsubstituted. When substituted, the substituent group can be, for example, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heteroalicyclic, halo, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, sulfinyl, sulfonyl, cyano, nitro, azide, sulfonyl, sulfinyl, sulfonamide, phosphonyl, phosphinyl, oxo, carbonyl, thiocarbonyl, urea, thiourea, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, C-carboxy, O-carboxy, sulfonamido, and amino, as these terms are defined herein.

[0075] A "heteroalicyclic" group refers to a monocyclic or fused ring group having in the ring(s) one or more atoms such as nitrogen, oxygen and sulfur. The rings may also have

one or more double bonds. However, the rings do not have a completely conjugated pi-electron system. The heteroalicyclic may be substituted or unsubstituted. When substituted, the substituted group can be, for example, lone pair electrons, alkyl, alkenyl, alkyanyl, cycloalkyl, aryl, heteroaryl, heteroalicyclic, halo, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, sulfinyl, sulfonyl, cyano, nitro, azide, sulfonyl, sulfinyl, sulfonamide, phosphonyl, phosphinyl, oxo, carbonyl, thiocarbonyl, urea, thiourea, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, C-carboxy, O-carboxy, sulfonamido, and amino, as these terms are defined herein. Representative examples are piperidine, piperazine, tetrahydro furane, tetrahydropyrane, morpholino and the like.

[0076] An "alkoxy" group refers to both an -O-alkyl and an -O-cycloalkyl group, as defined herein.

[0077] An "aryloxy" group refers to both an -O-aryl and an -O-heteroaryl group, as defined herein.

[0078] A "thioalkoxy" group refers to both an -S-alkyl group, and an -S-cycloalkyl group, as defined herein.

[0079] A "thioaryloxy" group refers to both an -S-aryl and an -S-heteroaryl group, as defined herein.

[0080] A "carbonyl" group refers to a -C(=O)-R group, where R is hydrogen, alkyl, alkenyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) or heteroalicyclic (bonded through a ring carbon) as defined herein.

[0081] An "aldehyde" group refers to a carbonyl group, where R is hydrogen.

[0082] A "thiocarbonyl" group refers to a -C(=S)-R group, where R is as defined herein.

[0083] A "C-carboxy" group refers to a -C(=O)-O-R groups, where R is as defined herein.

[0084] An "O-carboxy" group refers to an RC(=O)-O- group, where R is as defined herein.

[0085] An "oxo" group refers to a =O group.

[0086] A "carboxylic acid" group refers to a C-carboxyl group in which R is hydrogen.

[0087] A "halo" group refers to fluorine, chlorine, bromine or iodine.

[0088] A "trihalomethyl" group refers to a -CX₃ group wherein X is a halo group as defined herein.

[0089] A "sulfinyl" group refers to an $-S(=O)-R$ group, where R is as defined herein.

[0090] A "sulfonyl" group refers to an $-S(=O)_2-R$ group, where R is as defined herein.

[0091] An "S-sulfonamido" group refers to a $-S(=O)_2-NR_2$ group, with each of R as is defined herein.

[0092] An "N-sulfonamido" group refers to an $RS(=O)_2-NR$ group, where each of R is as defined herein.

[0093] An "O-carbamyl" group refers to an $-OC(=O)-NR_2$ group, where each of R is as defined herein.

[0094] An "N-carbamyl" group refers to an $ROC(=O)-NR-$ group, where each of R is as defined herein.

[0095] An "O-thiocarbamyl" group refers to an $-OC(=S)-NR_2$ group, where each of R is as defined herein.

[0096] An "N-thiocarbamyl" group refers to an $ROC(=S)NR-$ group, where each of R is as defined herein.

[0097] An "amino" group refers to an $-NR_2$ group where each of R is as defined herein.

[0098] A "C-amido" group refers to a $-C(=O)-NR_2$ group, where each of R is as defined herein.

[0099] An "N-amido" group refers to an $RC(=O)-NR-$ group, where each of R is as defined herein.

[0100] An "urea" group refers to an $-NRC(=O)-NR_2$ group, where each of R is as defined herein.

[0101] A "guanidino" group refers to an $-RNC(=N)-NR_2$ group, where each of R is as defined herein.

[0102] A "guanyl" group refers to an $R_2NC(=N)-$ group, where each of R is as defined herein.

[0103] The term "phosphonyl" or "phosphonate" describes a $-P(=O)(OR)_2$ group, with each of R as defined herein.

[0104] The term "phosphate" describes an $-O-P(=O)(OR)_2$ group, with each of R as defined herein.

[0105] A "phosphoric acid" is a phosphate group in which each of R is hydrogen.

[0106] The term "phosphinyl" describes a $-PR_2$ group, with each of R as defined herein.

[0107] The term "thiourea" describes a -NR-C(=S)-NR- group, with each of R as defined herein.

[0108] The term "saccharide" refers to one or more sugar unit, either an open-chain sugar unit or a cyclic sugar unit (*e.g.*, pyranose- or furanose-based units), and encompasses any monosaccharide, disaccharide and oligosaccharide, unless otherwise indicated.

[0109] The term "salt" includes both internal salt or external salt. In some embodiments, the salt is an internal salt, *i.e.*, a zwitterion structure. In some embodiments, the salt is an external salt. In some embodiments, the external salt is a pharmaceutically acceptable salt having a suitable counter ion. Suitable counterions for pharmaceutical use are known in the art.

[0110] The term "VB-201" refers to 1-hexadecyl-2-(4'-carboxy)butyl-glycero-3-phosphocholine. According to embodiments of the present invention, VB-201 may be a chiral enantiomer of 1-hexadecyl-2-(4'-carboxy)butyl-glycero-3-phosphocholine, *i.e.*, either the (*R*)- enantiomer ((*R*)-1-hexadecyl-2-(4'-carboxy)butyl-*sn*-glycero-3-phosphocholine) or the (*S*)- enantiomer ((*S*)-1-hexadecyl-2-(4'-carboxy)butyl-*sn*-glycero-3-phosphocholine), or a mixture thereof (*e.g.*, a racemate). According to exemplary embodiments, VB-201 is (*R*)-1-hexadecyl-2-(4'-carboxy)butyl-*sn*-glycero-3-phosphocholine. As understood by those skilled in the art, designating VB-201 as the (*R*)- enantiomer or the (*S*)- enantiomer does not require 100% enantiomeric purity, but instead refers to a substantially enriched single enantiomer either as an R or S isomer (*e.g.*, having an enantiomeric excess of at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, or higher). In some embodiments, VB-201 is (*R*)-1-hexadecyl-2-(4'-carboxy)butyl-*sn*-glycero-3-phosphocholine having at least 90% enantiomeric excess.

[0111] The term "VB-221" refers to 1-(2'-octyl)dodecyl-2-(4'-carboxy)butyl-glycero-3-phosphocholine. According to embodiments of the present invention, VB-221 may be a chiral enantiomer of (1-(2'-octyl)dodecyl-2-(4'-carboxy)butyl-glycero-3-phosphocholine), *i.e.*, either the (*R*)- enantiomer or the (*S*)- enantiomer, or any mixtures thereof (*e.g.*, a racemate). According to exemplary embodiments, VB-221 is (*R*)-1-(2'-octyl)dodecyl-2-(4'-carboxy)butyl-*sn*-glycero-3-phosphocholine. Similarly, as understood by those skilled in the art, designating VB-221 as the (*R*)- enantiomer or the (*S*)- enantiomer does not require 100% enantiomeric purity, but instead refers to a substantially enriched single

enantiomer either as an R or S isomer (*e.g.*, having an enantiomeric excess of at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, or higher). In some embodiments, VB-221 is (*R*)-1-(2'-octyl)dodecyl-2-(4'-carboxy)butyl-*sn*-glycero-3-phosphocholine having at least 90% enantiomeric excess.

[0112] It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention, which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable subcombination or as suitable in any other described embodiment of the invention. Certain features described in the context of various embodiments are not to be considered essential features of those embodiments, unless the embodiment is inoperative without those elements.

MOSPD2 and Inhibitors of MOSPD2

[0113] Inhibition of MOSPD2 has been found to inhibit migration of cancer cells and monocytes towards different chemokines (*e.g.*, EGF) and block activation of chemokine receptor signaling pathways. These results indicate that MOSPD2 is pivotal for cancer cell migration and metastasis and that blocking its activity has therapeutic benefit, for example, in treating, preventing, or reducing the incidence of metastasis of cancer cells.

[0114] Embodiments of the invention relate to an inhibitor of MOSPD2, *e.g.*, MOSPD2 expressed by a cancer cell, or to methods and compositions comprising an inhibitor of MOSPD2, *e.g.*, MOSPD2 expressed by a cancer cell. In some embodiments the MOSPD2 is a mammalian MOSPD2. In other embodiments, the MOSPD2 is a human MOSPD2. In some embodiments, the inhibitor is an isolated binding molecule that inhibits MOSPD2. In other embodiments, the inhibitor is a polypeptide, DNA, or RNA. In other embodiments, the inhibitor is a polypeptide that specifically binds to MOSPD2. In other embodiments, the inhibitor is an antibody or antigen binding fragment thereof that specifically binds to MOSPD2. In other embodiments, the inhibitor is an RNA silencing agent.

[0115] In additional embodiments of the invention, inhibition of MOSPD2 and downregulation of a MOSPD2 activity can be affected on the genomic and/or the transcription level using a variety of molecules which interfere with transcription and/or translation [*e.g.*, RNA silencing agents (*e.g.*, antisense, siRNA, shRNA, micro-RNA),

Ribozyme and DNAzyme], or on the protein level using *e.g.*, antagonists, enzymes that cleave the polypeptide, small molecules that interfere with the protein's activity (*e.g.*, competitive ligands) and the like.

[0116] Following is an exemplary list of agents capable of downregulating expression level and/or activity of a target such as MOSPD2.

[0117] Inhibition of MOSPD2 can occur, for example, by ectopic overexpression of MOSPD2 by infection, and it is intended that an inhibitor of MOSPD2 or a MOSPD2 inhibitor encompasses this type of inhibition.

[0118] Downregulation of MOSPD2 can also be achieved by gene editing. Gene editing can be performed, for example, with a clustered regularly interspaced short palindromic repeats CRISPR-CAS9 system. CRISPR-CAS9 systems have been described in the literature and can include, for example, CAS9 and a guide RNA. Other gene editing techniques have also been described in the literature and can also be used.

[0119] Downregulation of MOSPD2 can also be achieved by RNA silencing. As used herein, the phrase "RNA silencing" refers to a group of regulatory mechanisms [*e.g.*, RNA interference (RNAi), transcriptional gene silencing (TGS), post-transcriptional gene silencing (PTGS), quelling, co-suppression, and translational repression] mediated by RNA molecules which result in the inhibition or "silencing" of the expression of a corresponding protein-coding gene. RNA silencing has been observed in many types of organisms, including plants, animals, and fungi.

[0120] As used herein, the term "RNA silencing agent" refers to an RNA which is capable of specifically inhibiting or "silencing" the expression of a target gene. In some embodiments, the RNA silencing agent is capable of preventing complete processing (*e.g.*, the full translation and/or expression) of an mRNA molecule through a post-transcriptional silencing mechanism. RNA silencing agents include noncoding RNA molecules, for example, RNA duplexes comprising paired strands, as well as precursor RNAs from which such small non-coding RNAs can be generated. Exemplary RNA silencing agents include dsRNAs such as siRNAs, miRNAs and shRNAs. In one embodiment, the RNA silencing agent is capable of inducing RNA interference. In another embodiment, the RNA silencing agent is capable of mediating translational repression.

[0121] RNA interference refers to the process of sequence-specific post-transcriptional gene silencing in animals mediated by short interfering RNAs (siRNAs). The corresponding process in plants is commonly referred to as post-transcriptional gene silencing or RNA silencing and is also referred to as quelling in fungi. The process of post-transcriptional gene silencing is thought to be an evolutionarily-conserved cellular defense mechanism used to prevent the expression of foreign genes and is commonly shared by diverse flora and phyla. Such protection from foreign gene expression may have evolved in response to the production of double-stranded RNAs (dsRNAs) derived from viral infection or from the random integration of transposon elements into a host genome via a cellular response that specifically destroys homologous single-stranded RNA or viral genomic RNA.

[0122] Some embodiments of the invention contemplate use of dsRNA to downregulate protein expression from mRNA.

[0123] The term "siRNA" refers to small inhibitory RNA duplexes (generally between 18-30 basepairs) that induce the RNA interference (RNAi) pathway. Typically, siRNAs are chemically synthesized as 21mers with a central 19 bp duplex region and symmetric 2-base 3'-overhangs on the termini, although it has been recently described that chemically synthesized RNA duplexes of 25-30 base length can have as much as a 100-fold increase in potency compared with 21mers at the same location. The observed increased potency obtained using longer RNAs in triggering RNAi is theorized to result from providing Dicer with a substrate (27mer) instead of a product (21mer) and that this improves the rate or efficiency of entry of the siRNA duplex into RISC.

[0124] The strands of a double-stranded interfering RNA (*e.g.*, an siRNA) may be connected to form a hairpin or stem-loop structure (*e.g.*, an shRNA or sh-RNA). Thus, as mentioned, the RNA silencing agent of some embodiments of the invention may also be a short hairpin RNA (shRNA).

[0125] The terms "shRNA" or "sh-RNA", as used herein, refer to an RNA agent having a stem-loop structure, comprising a first and second region of complementary sequence, the degree of complementarity and orientation of the regions being sufficient such that base pairing occurs between the regions, the first and second regions being joined by a loop region, the loop resulting from a lack of base pairing between nucleotides (or nucleotide analogs) within the loop region. The number of nucleotides in the loop is a number

between and including 3 to 23, or 5 to 15, or 7 to 13, or 4 to 9, or 9 to 11. Some of the nucleotides in the loop can be involved in base-pair interactions with other nucleotides in the loop.

[0126] It will be appreciated that the RNA silencing agent of some embodiments of the invention need not be limited to those molecules containing only RNA, but further encompasses chemically-modified nucleotides and non-nucleotides.

[0127] In some embodiments, the RNA silencing agent provided herein can be functionally associated with a cell-penetrating peptide. As used herein, a "cell-penetrating peptide" is a peptide that comprises a short (about 12-30 residues) amino acid sequence or functional motif that confers the energy-independent (*i.e.*, non-endocytotic) translocation properties associated with transport of the membrane-permeable complex across the plasma and/or nuclear membranes of a cell.

[0128] According to another embodiment, the RNA silencing agent may be a miRNA or a mimic thereof.

[0129] The term "microRNA", "miRNA", and "miR" are synonymous and refer to a collection of non-coding single-stranded RNA molecules of about 19-28 nucleotides in length, which regulate gene expression. miRNAs are found in a wide range of organisms and have been shown to play a role in development, homeostasis, and disease etiology.

[0130] The term "microRNA mimic" refers to synthetic non-coding RNAs that are capable of entering the RNAi pathway and regulating gene expression. miRNA mimics imitate the function of endogenous microRNAs (miRNAs) and can be designed as mature, double stranded molecules or mimic precursors (*e.g.*, or pre-miRNAs). miRNA mimics can be comprised of modified or unmodified RNA, DNA, RNA-DNA hybrids, or alternative nucleic acid chemistries (*e.g.*, LNAs or 2'-O,4'-C-ethylene-bridged nucleic acids (ENA)). For mature, double stranded miRNA mimics, the length of the duplex region can vary between 13-33, 18-24 or 21-23 nucleotides. The miRNA can also comprise a total of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 nucleotides. The sequence of the miRNA can be the first 13-33 nucleotides of the pre-miRNA. The sequence of the miRNA can also be the last 13-33 nucleotides of the pre-miRNA.

[0131] Another agent capable of downregulating a target is a DNAzyme molecule capable of specifically cleaving an mRNA transcript or DNA sequence of the target.

DNAzymes are single-stranded polynucleotides which are capable of cleaving both single and double stranded target sequences. (Breaker *et al.*, *Chemistry and Biology* 1995; 2:655; Santoro *et al.*, *Proc. Natl. Acad. Sci. USA* 1997; 94:4262.) A general model (the "10-23" model) for the DNAzyme has been proposed. "10-23" DNAzymes have a catalytic domain of 15 deoxyribonucleotides, flanked by two substrate-recognition domains of seven to nine deoxyribonucleotides each. This type of DNAzyme can effectively cleave its substrate RNA at purine:pyrimidine junctions. (Santoro *et al.*,; Khachigian, *Curr. Opin. Mol. Ther.* 2002; 4:119-121.)

[0132] Downregulation of a target can also be affected by using an antisense polynucleotide capable of specifically hybridizing with an mRNA transcript encoding the target.

[0133] Another agent capable of downregulating a target is a ribozyme molecule capable of specifically cleaving an mRNA transcript encoding a target. Ribozymes are being increasingly used for the sequence-specific inhibition of gene expression by the cleavage of mRNAs encoding proteins of interest. (Welch *et al.*, *Curr. Opin. Biotechnol.* 1998; 9:486-96.)

[0134] Another agent capable of downregulating a target is any molecule which binds to and/or cleaves the target. Such molecules can be antagonists of the target, or inhibitory peptides of the target.

[0135] It will be appreciated that a non-functional analogue of at least a catalytic or binding portion of a target can be also used as an agent which downregulates the target.


[0136] Another agent which can be used along with some embodiments of the invention to downregulate a target is a molecule which prevents target activation or substrate binding.

[0137] In some embodiments, an inhibitor of a given protein target inhibits the protein by binding to the protein, by binding to a compound which binds to the protein (e.g., a substrate, a regulatory protein) and/or by binding to an oligonucleotide (e.g., mRNA) encoding the protein.

[0138] In some embodiments, the inhibitor of MOSPD2 is a small molecule (e.g., characterized by a molecular weight of less than 800 Da). In some embodiments, the small molecule MOSPD2 inhibitor is a tocopherol or a derivative thereof (e.g., alpha-tocopherol, beta-tocopherol, gamma-tocopherol, delta-tocopherol), a triterpene (e.g.,

squalene), a vitamin A or a derivative thereof (e.g., retinaldehyde), a phosphatidylglyceride (e.g., phosphatidylinositol), or a phospholipid (e.g., phosphatidylcholine, an oxidized phospholipid).

[0139] In some embodiments, the small molecule MOSPD2 inhibitor is an oxidized phospholipid having a structure according to Formula I:



Formula I

or a pharmaceutically acceptable salt, a hydrate or a solvate thereof, wherein:


n is an integer from 1 to 6, wherein when n is 1, Cn, Bn, Rn, and Y are absent, and C₁ is attached to R'n;

each of B₁, B₂, ... B_{n-1} and B_n is independently selected from the group consisting of oxygen, sulfur, nitrogen, phosphorus and silicon, whereby each of said nitrogen, phosphorus and silicon is optionally substituted by one substituent selected from the group consisting of alkyl, halo, cycloalkyl, aryl, hydroxy, thiohydroxy, alkoxy, aryloxy, thioaryloxy, thioalkoxy and oxo;

each of A₁, A₂, ... A_{n-1} and A_n is independently selected from the group consisting of CR"R", C=O and C=S,

Y is selected from the group consisting of hydrogen, acyl, alkyl, aryl, cycloalkyl, carboxy, saccharide, phosphoric acid, phosphoryl choline, phosphoryl ethanolamine, phosphoryl serine, phosphoryl cardiolipin, phosphoryl inositol, ethylphosphocholine, phosphorylmethanol, phosphorylethanol, phosphorylpropanol, phosphorylbutanol, phosphorylethanolamine-N-lactose, phosphoethanolamine-N-glutaric acid, phosphoethanolamine-N-[methoxy(propylene glycol)], phosphoinositol-4-phosphate, phosphoinositol-4,5-biphosphonate, phosphoinositol-4,5-

bisphosphate, pyrophosphate, phosphoethanolamine-diethylenetriamine-pentaacetate, dinitrophenyl-phosphoethanolamine, phosphoglycerol and a moiety having the general formula:




wherein:

each of B' and B" is independently selected from the group consisting of sulfur and oxygen; and

each of D' and D" is independently selected from the group consisting of hydrogen, alkyl, amino substituted alkyl, cycloalkyl, phosphonate and thiophosphonate; and


each of X_1, X_2, \dots, X_{n-1} is independently a saturated or unsaturated hydrocarbon having the general Formula II:



Formula II

wherein m is an integer from 1 to 26; and

Z is selected from the group consisting of:



wherein W is selected from the group consisting of oxygen and sulfur;


wherein at least one of X_1, X_2, \dots, X_{n-1} comprises a Z other than hydrogen,

and wherein:

each of R_1 , R'_1 , R_2 , ... R_{n-1} , R_n , $R'n$, each of R'' and R''' and each of R_a , $R'a$, R_b , $R'b$, ... R_{m-1} , $R'm-1$, R_m and $R'm$ is independently selected from the group consisting of a bond, hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, halo, trihalomethyl, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, phosphonate, phosphate, phosphinyl, sulfonyl, sulfinyl, sulfonamide, amide, carbonyl, thiocarbonyl, C-carboxy, O-

carboxy, C-carbamate, N-carbamate, C-thiocarboxy, S-thiocarboxy and amino, or, alternatively, at least two of R₁, R'₁, R₂, ... R_{n-1}, R_n and R'_n and/or at least two of R_a, R'_a, R_b, R'_b, ... R_{m-1}, R'_{m-1}, R_m and R'_m form at least one four-, five- or six-membered aromatic, heteroaromatic, alicyclic or heteroalicyclic ring, or a pharmaceutically acceptable salt, a hydrate or a solvate thereof. In any of the embodiments described herein, the oxidized phospholipid can exist as a stereoisomeric mixture of any ratio, for example, as a substantially enriched single enantiomer such as an R or S isomer (e.g., having an enantiomeric excess of at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, or higher), or as a mixture of two enantiomers (e.g., a racemic mixture); and/or as a substantially enriched single diastereomer (e.g., having an diastereomeric excess of at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, or higher), or as a mixture of two or more diastereomers.

[0140] In one embodiment, the oxidized phospholipid useful in any of the methods of the present disclosure has a structure according to Formula III:



Formula III

or a pharmaceutically acceptable salt, hydrate or solvate thereof.


[0141] In Formula III, n is an integer selected from 1 to 4.

[0142] In Formula III, B₁, each B₂, and B₃ are independently selected from the group consisting of oxygen, sulfur, and NR₄, wherein R₄ is selected from hydrogen, alkyl, cycloalkyl, aryl, and acyl.

[0143] In Formula III, A₁ and each A₂ are independently selected from the group consisting of CR_eR_{ee}, CR_e=CR_{ee}, C=O and C=S, wherein R_e and R_{ee} are independently selected from hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heteroaryl.

[0144] In Formula III, Y is selected from the group consisting of hydrogen, acyl, alkyl, aryl, cycloalkyl, carboxy, saccharide, phosphoric acid, phosphoryl choline, phosphoryl

ethanolamine, phosphoryl serine, phosphoryl cardiolipin, phosphoryl inositol, ethylphosphocholine, phosphorylmethanol, phosphorylethanol, phosphorylpropanol, phosphorylbutanol, phosphorylethanolamine-N-lactose, phosphoethanolamine-N-glutaric acid, phosphoethanolamine-N-[methoxy(propylene glycol)], phosphoinositol-4-phosphate, phosphoinositol-4,5-bisphosphate, pyrophosphate, phosphoethanolamine-diethylenetriamine-pentaacetate, dinitrophenyl-phosphoethanolamine, phosphoglycerol, and a moiety having the general formula:




wherein:

each of B and B_a is independently selected from the group consisting of sulfur and oxygen; and


D and D_a are independently selected from the group consisting of hydrogen, alkyl, aminoalkyl, cycloalkyl, phosphonate and thiophosphonate.

[0145] In Formula III, X₁ and each X₂ are independently a saturated or unsaturated, linear or branched hydrocarbon, wherein at least one of X₁ and X₂ is substituted with an oxidized moiety Z selected from the group consisting of:



wherein W is oxygen or sulfur; and R_d and R_{dd} are independently selected from hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heteroaryl.


[0146] In one embodiment in Formula III, X₁ and each X₂ independently have the general Formula IV:



Formula IV

[0147] In Formula IV, m is an integer selected from 1 to 26.

[0148] In Formula IV, Z is selected from the group consisting of:



wherein W is oxygen or sulfur; and R_d and R_{dd} are independently selected from hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heteroaryl,

wherein at least one of X₁ and X₂ comprises a Z other than hydrogen.

[0149] In Formula III and Formula IV, R₁, R_{1a}, each R₂, R₃, R_{3a}, R_a, R_{aa}, each R_b, each R_{bb}, R_c and R_{cc} are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, halo, trihalomethyl, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, phosphonate, phosphate, phosphinyl, sulfonyl, sulfinyl, sulfonamide, amide, carbonyl, thiocarbonyl, C-carboxy, O-carboxy, C-carbamate, N-carbamate, C-thiocarboxy, S-thiocarboxy and amino, wherein at least two of R₁, R_{1a}, R₂, R₃ and R_{3a} are optionally joined to form a four-, five- or six-membered aromatic, heteroaromatic, alicyclic or heteroalicyclic ring, and wherein at least two of R_a, R_{aa}, R_b, R_{bb}, R_c, and R_{cc} are optionally joined to form a four-, five- or six-membered aromatic, heteroaromatic, alicyclic or heteroalicyclic ring.


[0150] In one embodiment in Formula III, n is 1 or 2. In another embodiment in Formula III, n is 1.

[0151] In one embodiment in Formula III, Y is selected from the group consisting of hydrogen, acyl, alkyl, aryl, cycloalkyl, carboxy, saccharide, phosphoric acid, phosphoryl choline, phosphoryl ethanolamine, phosphoryl serine, phosphoryl cardiolipin, phosphoryl inositol, ethylphosphocholine, phosphorylmethanol, phosphorylethanol, phosphorylpropanol, phosphorylbutanol, phosphorylethanolamine-N-lactose, phosphoethanolamine-N-glutaric acid, phosphoethanolamine-N-[methoxy(propylene glycol)], phosphoinositol-4-phosphate, phosphoinositol-4,5-bisphosphate, pyrophosphate, phosphoethanolamine-diethylenetriamine-pentaacetate, dinitrophenyl-phosphoethanolamine, and phosphoglycerol.

[0152] In another embodiment in Formula III, Y is selected from the group consisting of hydrogen, phosphoryl choline, and phosphoryl ethanolamine.

[0153] In another embodiment in Formula III, Y is selected from the group consisting of phosphoryl choline, and phosphoryl ethanolamine.

[0154] In one embodiment in Formula III, Y is phosphoryl choline.




[0155] In one embodiment in Formula III, Z is . In another embodiment in Formula III, Z is a carboxylic acid group.

[0156] In a further embodiment in Formula III, n is 1 and Y is phosphoryl choline.

[0157] In a further embodiment in Formula III, each of B₁, B₂, and B₃ is oxygen.

[0158] In a further embodiment in Formula III, n is 1, Y is phosphoryl choline, and each of B₁, B₂, and B₃ is oxygen.

[0159] In one embodiment, the oxidized phospholipid useful in any of the methods of the present disclosure has a structure according to Formula IIIa:



Formula IIIa

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

[0160] In Formula IIIa, B₁, B₂, and B₃ are independently selected from oxygen and sulfur.


[0161] In Formula IIIa, A₁ and A₂ are independently selected from the group consisting of CH₂, CH=CH, C=O and C=S.

[0162] In Formula IIIa, Y is selected from the group consisting of hydrogen, acyl, alkyl, aryl, cycloalkyl, carboxy, saccharide, phosphoric acid, phosphoryl choline, phosphoryl ethanolamine, phosphoryl serine, phosphoryl cardiolipin, phosphoryl inositol, ethylphosphocholine, phosphorylmethanol, phosphorylethanol, phosphorylpropanol, phosphorylbutanol, phosphorylethanolamine-N-lactose, phosphoethanolamine-N-glutaric acid, phosphoethanolamine-N-[methoxy(propylene glycol)], phosphoinositol-4-

phosphate, phosphoinositol-4,5-bisphosphate, pyrophosphate, phosphoethanolamine-diethylenetriamine-pentaacetate, dinitrophenyl-phosphoethanolamine, and phosphoglycerol.


[0163] In Formula IIIa, R₁, R_{1a}, R₂, R₃, and R_{3a}, are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, halo, trihalomethyl, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, phosphonate, phosphate, phosphinyl, sulfonyl, sulfinyl, sulfonamide, amide, carbonyl, thiocarbonyl, C-carboxy, O-carboxy, C-carbamate, N-carbamate, C-thiocarboxy, S-thiocarboxy and amino, wherein at least two of R₁, R_{1a}, R₂, R₃ and R_{3a} are optionally joined to form a four-, five- or six-membered aromatic, heteroaromatic, alicyclic or heteroalicyclic ring, and wherein at least two of R_a, R_{aa}, R_b, R_{bb}, R_c, and R_{cc} are optionally joined to form a four-, five- or six-membered aromatic, heteroaromatic, alicyclic or heteroalicyclic ring;

[0164] In Formula IIIa, X₁ and X₂ are independently a saturated or unsaturated, linear or branched hydrocarbon, wherein at least one of X₁ and X₂ is substituted with an oxidized moiety Z having a formula selected from:



wherein W is oxygen or sulfur; and R_d and R_{dd} are independently selected from hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heteroaryl.

[0165] In one embodiment in Formula IIIa, X₁ and X₂ independently have a structure according to Formula IVa:




[0166] In Formula IVa, m is an integer selected from 1 to 26.


[0167] In Formula IVa, R_a, R_{aa}, each R_b, each R_{bb}, R_c, and R_{cc} are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl,

heteroaryl, halo, trihalomethyl, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, phosphonate, phosphate, phosphinyl, sulfonyl, sulfinyl, sulfonamide, amide, carbonyl, thiocarbonyl, C-carboxy, O-carboxy, C-carbamate, N-carbamate, C-thiocarboxy, S-thiocarboxy and amino, wherein at least two of R_a , R_{aa} , R_b , R_{bb} , R_c , and R_{cc} are optionally joined to form a four-, five- or six-membered aromatic, heteroaromatic, alicyclic or heteroalicyclic ring.

[0168] In Formula IVa, Z is selected from the group consisting of:



wherein W is oxygen or sulfur; and R_d and R_{dd} are independently selected from hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heteroaryl, wherein at least one of X_1 and X_2 comprises a Z other than hydrogen.



[0169] In one embodiment in Formula IIIa, Z is . In another embodiment in Formula IIIa, Z is a carboxylic acid group.

[0170] In one embodiment in Formula IIIa, Y is selected from the group consisting of hydrogen, acyl, alkyl, aryl, cycloalkyl, carboxy, saccharide, phosphoric acid, phosphoryl choline, phosphoryl ethanolamine, phosphoryl serine, phosphoryl cardiolipin, phosphoryl inositol, ethylphosphocholine, phosphorylmethanol, phosphorylethanol, phosphorylpropanol, phosphorylbutanol, phosphorylethanolamine-N-lactose, phosphoethanolamine-N-glutaric acid, phosphoethanolamine-N-[methoxy(propylene glycol)], phosphoinositol-4-phosphate, phosphoinositol-4,5-bisphosphate, phosphoethanolamine-diethylenetriamine-pentaacetate, dinitrophenyl-phosphoethanolamine, and phosphoglycerol.

[0171] In one embodiment in Formula IIIa, Y is selected from the group consisting of hydrogen, phosphoryl choline, and phosphoryl ethanolamine.


[0172] In another embodiment in Formula IIIa, Y is selected from the group consisting of phosphoryl choline, and phosphoryl ethanolamine.

[0173] In one embodiment in Formula IIIa, Y is phosphoryl choline.

[0174] In a further embodiment in Formula IIIa, each of B₁, B₂, and B₃ is oxygen.

[0175] In a further embodiment in Formula IIIa, Y is phosphoryl choline, and each of B₁, B₂, and B₃ is oxygen.


[0176] In one embodiment in Formula IIIa, the oxidized phospholipid has a structure according to Formula V:



Formula V

wherein B₁, B₂, B₃, A₁, A₂, X₁, X₂, and Y are defined as for Formula IIIa.

[0177] In one embodiment, each of B₁, B₂, B₃ in Formula V is oxygen and the oxidized phospholipid has a structure according to the Formula VI:




Formula VI

[0178] In Formula VI, A₁ is selected from the group consisting of CH₂, CH=CH and C=O. In one example, A₁ in Formula VI is CH₂.

[0179] In Formula VI, A₂ is absent or CH₂.

[0180] In Formula VI, X₁ is an alkyl having from 1 to 30 carbon atoms.


[0181] In Formula VI, X₂ is 

wherein

E is absent or is an alkyl chain having from 1 to 24 carbon atoms;

F is selected from the group consisting of hydrogen, hydroxy, alkyl, alkoxy, halide, acetoxy and aryl; and

Z is selected from the group consisting of:



wherein R_d is selected from H, alkyl and aryl.

[0182] In Formula VI, Y is selected from the group consisting of hydrogen, alkyl, aryl, phosphoric acid, phosphoryl choline, phosphoryl ethanolamine, phosphoryl serine, phosphatidyl choline, phosphatidyl ethanolamine, phosphatidyl serine, phosphatidyl cardiolipin, phosphatidyl inositol, phosphoryl cardiolipin, phosphoryl inositol, ethylphosphocholine, phosphorylmethanol, phosphorylethanol, phosphorylpropanol, phosphorylbutanol, phosphorylethanolamine-N-lactose, phosphoethanolamine-N-[methoxy(propylene glycol)], phosphoinositol-4-phosphate, phosphoinositol-4,5-bisphosphate, pyrophosphate, phosphoethanolamine-diethylenetriamine-pentaacetate, dinitrophenyl-phosphoethanolamine, phosphoglycerol.

[0183] In one embodiment in Formula VI, X₁ is alkyl having from 10 to 30 carbon atoms, or from 8 to 30 carbon atoms.


[0184] In one embodiment in Formula VI, E is alkyl having from 1 to 10 carbon atoms, or from 1 to 4 carbon atoms.

[0185] In one embodiment in Formula VI, Y is phosphoryl choline.

[0186] Each carbon atom in Formula I, II, III, IIIa, V, and VI is a chiral or non-chiral carbon atom, wherein each chiral carbon atom can have an S-configuration or R-configuration.


[0187] In one preferred embodiment, the oxidized phospholipid is 1-hexadecyl-2-(4'-carboxy)butyl-glycero-3-phosphocholine. In another preferred embodiment, the oxidized phospholipid is (R)-1-hexadecyl-2-(4'-carboxy)butyl-sn-glycero-3-phosphocholine.

[0188] In one preferred embodiment, the oxidized phospholipid is



, or a pharmaceutically acceptable salt thereof.

[0189] In one preferred embodiment, the oxidized phospholipid is



, or a pharmaceutically acceptable salt thereof.

[0190] The small molecule MOSPD2 inhibitor described herein can be used alone, as a single agent, in any of the methods described herein or it can be used in combination with another agent (e.g., another MOSPD2 inhibitor or an anticancer drug).

[0191] VB-201 inhibits human monocyte chemotaxis *in vitro* and signaling pathways activated downstream of chemokine receptors. In contrast, VB-221, a derivative of VB-201, does not inhibit chemokine-induced signaling and migration in human monocytes. It was also found that ovalbumin labeled VB-201 binds and precipitates MOSPD2 from cell lysate of human CD14 monocytes. Further, HEK293 cells transfected with hemagglutinin (HA)-tagged human MOSPD2 and positively stained for HA have a strong binding to ovalbumin labeled VB-201, but not to ovalbumin labeled VB-221. These experiments and others demonstrate that 1) VB-201 binds MOSPD2; 2) that VB-201 inhibits cell chemotaxis and chemotaxis-mediated downstream pathways; and 3) that addition of VB-201 yields the same signaling effects as silencing of MOSPD2.

[0192] In any of the embodiments described herein, useful small molecule MOSPD2 inhibitors include those that are more potent inhibitors of MOSPD2 (e.g., human MOSPD2 on the cell surface of a monocyte or cancer cell) when compared to VB-221, e.g., those having a lower IC₅₀ value compared to that of VB-221. More preferably, useful small molecule MOSPD2 inhibitors include those that are equal or more potent inhibitors of MOSPD2 (e.g., human MOSPD2 on the cell surface of a monocyte or cancer

cell) when compared to VB-201, *e.g.*, those having a lower IC₅₀ value compared to that of VB-201. As understood by those skilled in the art, an IC₅₀ value indicates how much of a particular drug or other substance (inhibitor) is needed to inhibit a given biological process (or component of a process, *i.e.* an enzyme, cell, cell receptor or microorganism) by half. Methods for determining IC₅₀ values are known in the art.

[0193] When a small molecule MOSPD2 inhibitor (as described herein) in a pharmaceutical composition is administered to a subject alone as a single agent or in combination with another agent, the small molecule MOSPD2 inhibitor (*e.g.*, VB-201) is present in an amount such that the administration causes at least 10% (*e.g.*, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 99%, or higher) inhibition of one or more activities of MOSPD2 (*e.g.*, human MOSPD2) (*e.g.*, MOSPD2 expression, cancer cell migration, monocyte migration associated with tumor growth (*e.g.*, presence of tumor associated macrophages), a chemokine signaling pathway, a growth factor signaling pathway, EGF receptor phosphorylation, ERK phosphorylation, AKT phosphorylation, and/or FAK phosphorylation). In some embodiments, administration of the small molecule MOSPD2 inhibitor to a human subject causes at least 10% (*e.g.*, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 99%, or higher) inhibition of one or more activities of a human MOSPD2. In one aspect, administration of the small molecule MOSPD2 inhibitor causes from about 10% to 100%, from about 10% to about 99%, from about 10% to about 95%, from about 10% to about 90%, from about 10% to about 85%, from about 10% to about 80%, from about 10% to about 70%, from about 20% to about 99%, from about 20% to about 95%, from about 20% to about 90%, from about 20% to about 85%, from about 20% to about 80%, from about 30% to about 95%, from about 30% to about 90%, from about 30% to about 85%, from about 30% to about 80%, from about 40% to about 95%, from about 40% to about 90%, from about 40% to about 85%, from about 40% to about 80%, from about 50% to about 95%, from about 50% to about 90%, from about 50% to about 85%, from about 50% to about 80%, from about 60% to about 95%, from about 60% to about 90%, from about 60% to about 85%, or from about 60% to about 80% inhibition of one or more activities of MOSPD2, *e.g.*, regulation of cancer cell migration, monocyte migration associated with tumor growth, presence of tumor associated macrophages, chemokine

signaling pathways, growth factor signaling pathways, EGF receptor phosphorylation, ERK phosphorylation, AKT phosphorylation, and/or FAK phosphorylation. Preferably, the small molecule MOSPD2 inhibitor is VB-201.

[0194] In some embodiments, an inhibitor of a given protein inhibits the protein by binding to the protein and/or to an oligonucleotide (e.g., mRNA) encoding the protein.

[0195] In other embodiments, the MOSPD2 inhibitor is (i) an isolated binding molecule that specifically binds to a MOSPD2 polypeptide, (ii) an isolated binding molecule that specifically binds to a ligand of a MOSPD2 polypeptide, (iii) an antisera raised against a MOSPD2 polypeptide, (iv) a soluble MOSPD2 polypeptide, or (v) a soluble MOSPD2 polypeptide comprising, consisting essentially of, or consisting of an extracellular domain of a MOSPD2 polypeptide.

[0196] In still other embodiments, the inhibitor is an antibody that specifically binds to a MOSPD2 polypeptide. In other embodiments, the inhibitor is an antigen binding fragment of an antibody that specifically binds to a MOSPD2 polypeptide. In other embodiments, the antibody is a polyclonal, monoclonal, murine, human, humanized, chimeric, or single chain antibody. In other embodiments, the antigen binding fragment is a Fab, Fab', F(ab')₂, Fv, scFv, sdFv fragment, VH domain, or VL domain.

[0197] In some embodiments, an antibody or antigen binding fragment thereof described herein, which specifically binds to MOSPD2 (e.g., human MOSPD2), comprises a VH, a VL, or a VH and VL. In other embodiments, the antibody or antigen binding fragment thereof comprises a constant region.

[0198] In some embodiments, the VH, VL, or VH and VL comprise one or more complementarity determining regions (CDRs). In some embodiments, the VH comprises CDR1, CDR2, CDR3, or any combination thereof. In some embodiments, the VL comprises CDR1, CDR2, CDR3, or any combination thereof.

[0199] In some embodiments, the VH, VL, or VH and VL comprise one or more framework regions (FRs). In some embodiments, the VH comprises FR1, FR2, FR3, FR4, or any combination thereof. In some embodiments, the VL comprises FR1, FR2, FR3, FR4, or any combination thereof.

[0200] In a particular embodiment, an antibody or antigen binding fragment thereof described herein, which specifically binds to MOSPD2 (e.g., human MOSPD2),

comprises a VH comprising CDR1, CDR2, and CDR3, and a VL comprising CDR1, CDR2, and CDR3.

[0201] In other embodiments, the antibodies or antigen binding fragments thereof comprise a constant region. In some embodiments, the constant region of the light chain comprises the amino acid sequence of a human kappa light chain constant region or a human lambda light chain constant region. In some embodiments, the constant region of the heavy chain comprises the amino acid sequence of a human gamma heavy chain constant region. Non-limiting examples of human constant region sequences have been described, *e.g.*, *see* U.S. Patent No. 5,693,780 and Kabat, EA *et al.*, (1991). In some embodiments, the constant region amino acid sequence has been modified (*e.g.*, one, two or more amino acid substitutions) such that it has at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to a native human sequence. In another aspect, provided herein are antibodies or antigen binding fragments thereof that recognize or bind to an epitope of MOSPD2 (*e.g.*, an epitope of human MOSPD2). In another aspect, provided herein are antibodies or antigen binding fragments thereof that recognize or bind to the same epitope or an overlapping epitope of MOSPD2 (*e.g.*, human MOSPD2) as an antibody described herein (*e.g.*, an antibody described in Example 1 or 8). In another aspect, the antibodies or antigen binding fragments thereof recognize more than one epitope of MOSPD2 (*e.g.*, two, three, four, five or six epitopes).

[0202] In certain embodiments, an epitope of MOSPD2 can be determined by one or more methods described in the literature, *e.g.*, NMR spectroscopy, X-ray diffraction crystallography studies, ELISA assays, hydrogen/deuterium exchange coupled with mass spectrometry (*e.g.*, MALDI mass spectrometry), array-based oligo-peptide scanning assays, and/or mutagenesis mapping (*e.g.*, site-directed mutagenesis mapping). For X-ray crystallography, crystallization can be accomplished using methods described in the literature (*e.g.*, Giegé R *et al.*, (1994) *Acta Crystallogr D Biol Crystallogr* 50(Pt 4): 339-350; McPherson A (1990) *Eur J Biochem* 189: 1-23; Chayen NE (1997) *Structure* 5: 1269-1274; McPherson A (1976) *J Biol Chem* 251: 6300-6303). Antibody:antigen crystals can be studied using well known X-ray diffraction techniques and may be refined using computer software such as X-PLOR (Yale University, 1992, distributed by Molecular Simulations, Inc.; *see e.g.* *Meth Enzymol* (1985) volumes 114 & 115, eds

Wyckoff HW *et al.*; U.S. Patent Application No. 2004/0014194), and BUSTER (Bricogne G (1993) *Acta Crystallogr D Biol Crystallogr* 49(Pt 1): 37-60; Bricogne G (1997) *Meth Enzymol* 276A: 361-423, ed Carter CW; Roversi P *et al.*, (2000) *Acta Crystallogr D Biol Crystallogr* 56(Pt 10): 1316-1323). Mutagenesis mapping studies can be accomplished using methods described in the literature. *See, e.g.*, Champe M *et al.*, (1995) *supra* and Cunningham BC & Wells JA (1989) *supra* for a description of mutagenesis techniques, including alanine scanning mutagenesis techniques. In a specific embodiment, an epitope of an antibody or antigen binding fragment thereof is determined using alanine scanning mutagenesis studies. Epitope characterization of an antibody can also be determined by the methods provided in Ravn *et al.*, *Journal of Biological Chemistry* 288: 19760-19772 (2013).

[0203] In addition, antibodies or antigen binding fragments thereof that recognize or bind to the same or overlapping epitopes of MOSPD2 (*e.g.*, human MOSPD2) can be identified using routine techniques such as an immunoassay, for example, by showing the ability of one antibody to block the binding of another antibody to a target antigen, *i.e.*, a competitive binding assay. Competitive binding can be determined in an assay in which the immunoglobulin under test inhibits specific binding of a reference antibody to a common antigen, such as MOSPD2. Numerous types of competitive binding assays have been described, for example: solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition assay (*see* Stahli C *et al.*, (1983) *Methods Enzymol* 9: 242-253); solid phase direct biotin-avidin EIA (*see* Kirkland TN *et al.*, (1986) *J Immunol* 137: 3614-9); solid phase direct labeled assay, solid phase direct labeled sandwich assay (*see* Harlow E & Lane D, (1988) *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press); solid phase direct label RIA using I-125 label (*see* Morel GA *et al.*, (1988) *Mol Immunol* 25(1): 7-15); solid phase direct biotin-avidin EIA (Cheung RC *et al.*, (1990) *Virology* 176: 546-52); and direct labeled RIA (Moldenhauer G *et al.*, (1990) *Scand J Immunol* 32: 77-82). Typically, such an assay involves the use of purified antigen (*e.g.*, MOSPD2 such as human MOSPD2) bound to a solid surface or cells bearing either of these, an unlabeled test immunoglobulin and a labeled reference immunoglobulin. Competitive inhibition can be measured by determining the amount of label bound to the solid surface or cells in the presence of the test immunoglobulin. Usually the test immunoglobulin is present in excess. Usually,

when a competing antibody is present in excess, it will inhibit specific binding of a reference antibody to a common antigen by at least 50-55%, 55-60%, 60-65%, 65-70% 70-75% or more. A competition binding assay can be configured in a large number of different formats using either labeled antigen or labeled antibody. In a common version of this assay, the antigen is immobilized on a 96-well plate. The ability of unlabeled antibodies to block the binding of labeled antibodies to the antigen is then measured using radioactive or enzyme labels. For further details *see*, for example, Wagener C *et al.*, (1983) J Immunol 130: 2308-2315; Wagener C *et al.*, (1984) J Immunol Methods 68: 269-274; Kuroki M *et al.*, (1990) Cancer Res 50: 4872-4879; Kuroki M *et al.*, (1992) Immunol Invest 21: 523-538; Kuroki M *et al.*, (1992) Hybridoma 11: 391-407 and Antibodies: A Laboratory Manual, Ed Harlow E & Lane D editors *supra*, pp. 386-389.

[0204] In one embodiment, a competition assay is performed using surface plasmon resonance (BIAcore®), *e.g.*, by an "in tandem approach" such as that described by Abdiche YN *et al.*, (2009) Analytical Biochem 386: 172-180, whereby MOSPD2 antigen is immobilized on the chip surface, for example, a CM5 sensor chip and the anti-MOSPD2 antibodies are then run over the chip. To determine if an antibody or antigen binding fragment thereof competes with an anti-MOSPD2 antibody or antigen binding fragment thereof described herein, the anti-MOSPD2 antibody or antigen binding fragment thereof is first run over the chip surface to achieve saturation and then the potential, competing antibody is added. Binding of the competing antibody can then be determined and quantified relative to a non-competing control.

[0205] In certain aspects, competition binding assays can be used to determine whether an antibody or antigen binding fragment thereof is competitively blocked, *e.g.*, in a dose dependent manner, by another antibody for example, an antibody binds essentially the same epitope, or overlapping epitopes, as a reference antibody, when the two antibodies recognize identical or sterically overlapping epitopes in competition binding assays such as competition ELISA assays, which can be configured in all number of different formats, using either labeled antigen or labeled antibody. In a particular embodiment, an antibody or antigen binding fragment thereof can be tested in competition binding assays with an antibody described herein (*e.g.*, those in Example 1 or 8), or a chimeric or Fab antibody thereof, or an antibody comprising VH CDRs and VL CDRs of an antibody described herein (*e.g.*, those in Example 1 or 8).

[0206] Accordingly, in a certain aspect, provided herein are antibodies or antigen binding fragments thereof that compete (e.g., in a dose dependent manner) for binding to MOSPD2 (e.g., human MOSPD2) with an antibody described herein (e.g., Example 1 or 8), as determined using assays known to one of skill in the art or described herein (e.g., ELISA competitive assays, surface plasmon resonance or Scatchard analysis).

[0207] In some embodiments, anti-MOSPD2 antibodies or antigen binding fragments thereof of the invention specifically bind to one or more of the following amino acid regions (epitopes) of MOSPD2, numbered according to SEQ ID NO:1 (amino acid residues 1-518): 508-517, 501-514, 233-241, 509-517, 212-221, 13-24, 505-517, 505-514, 89-100, 506-517, 233-245, 504-514, 128-136, 218-226, 15-24, 83-96, 42-50, 462-474, 340-351, 504-517, 462-470, 327-337, 21-32, 217-226, 510-517, 178-190, 497-509, 504-516, 64-77, 504-515, 147-159, 503-315, 88-97, 208-218, 178-191, 502-515, 503-516, 497-505, 500-509, 189-202, 189-197, 505-516, 1-63, 82-239, 93-234, 327-445, 327-431, and 497-517.

[0208] In some embodiments, anti-MOSPD2 antibodies or antigen binding fragments thereof of the invention specifically bind to one or more of the following amino acid regions (epitopes) of MOSPD2, numbered according to SEQ ID NO:1 (amino acid residues 1-518): about 508 to about 517, about 501 to about 514, about 233 to about 241, about 509 to about 517, about 212 to about 221, about 13 to about 24, about 505 to about 517, about 505 to about 514, about 89 to about 100, about 506 to about 517, about 233 to about 245, about 504 to about 514, about 128 to about 136, about 218 to about 226, about 15 to about 24, about 83 to about 96, about 42 to about 50, about 462 to about 474, about 340 to about 351, about 504 to about 517, about 462 to about 470, about 327 to about 337, about 21 to about 32, about 217 to about 226, about 510 to about 517, about 178 to about 190, about 497 to about 509, about 504 to about 516, about 64 to about 77, about 504 to about 515, about 147 to about 159, about 503 to about 515, about 88 to about 97, about 208 to about 218, about 178 to about 191, about 502 to about 515, about 503 to about 516, about 497 to about 505, about 500 to about 509, about 189 to about 202, about 189 to about 197, about 505 to about 516, about 1 to about 63, about 82 to about 239, about 93 to about 234, about 327 to about 445, about 327 to about 431, and about 497 to about 517.

[0209] In some embodiments, anti-MOSPD2 antibodies or antigen binding fragments thereof of the invention specifically bind to one or more of the following amino acid regions (epitopes) of MOSPD2, numbered according to SEQ ID NO:1 (amino acid residues 1-518): about 505 to about 515, about 500 to about 515, about 230 to about 240, about 510 to about 520, about 210 to about 220, about 15 to about 25, about 505 to about 520, about 505 to about 515, about 90 to about 100, about 505 to about 525, about 230 to about 245, about 505 to about 510, about 130 to about 140, about 220 to about 230, about 15 to about 30, about 80 to about 95, about 40 to about 50, about 460 to about 475, about 340 to about 350, about 500 to about 515, about 460 to about 470, about 325 to about 335, about 20 to about 35, about 215 to about 225, about 510 to about 520, about 175 to about 190, about 500 to about 510, about 505 to about 530, about 60 to about 75, about 500 to about 520, about 145 to about 160, about 502 to about 515, about 85 to about 100, about 205 to about 220, about 175 to about 190, about 500 to about 505, about 500 to about 525, about 495 to about 505, about 495 to about 510, about 190 to about 200, about 190 to about 198, about 502 to about 515, about 1 to about 60, about 80 to about 240, about 90 to about 235, about 330 to about 445, about 330 to about 430, and about 495 to about 515.

[0210] In some embodiments, antibodies or antigen binding fragments of the invention bind to MOSPD2 with an antibody-antigen equilibrium dissociation constant (K_D) of from about 10^{-6} M to about 10^{-12} M, or any range of values thereof (e.g., from about 10^{-7} M to about 10^{-12} , from 10^{-8} M to about 10^{-12} M, from about 10^{-9} M to about 10^{-12} M, from about 10^{-10} M to about 10^{-12} M, from about 10^{-11} M to about 10^{-12} M, from about 10^{-6} M to about 10^{-11} M, from about 10^{-7} M to about 10^{-11} M, from about 10^{-8} M to about 10^{-11} M, from about 10^{-9} M to about 10^{-11} M, from about 10^{-10} M to about 10^{-11} M, from about 10^{-6} M to about 10^{-10} M, from about 10^{-7} M to about 10^{-10} M, from about 10^{-8} M to about 10^{-10} M, from about 10^{-9} M to about 10^{-10} M, from about 10^{-6} M to about 10^{-9} M, from about 10^{-7} M to about 10^{-9} M, from about 10^{-8} M to about 10^{-9} M, from about 10^{-6} M to about 10^{-8} M, or from about 10^{-7} M to about 10^{-8}). In other embodiments, the antibody or antigen binding fragment thereof has a K_D of about 10^{-6} M, about 10^{-7} M, about 10^{-8} M, about 10^{-9} M, about 10^{-10} M, about 10^{-11} M, or about 10^{-12} M. In some embodiments, the antibody or antigen binding fragment binds to one or more epitopes on MOSPD2. In

some embodiments, the K_D is determined by Scatchard analysis, surface plasmon resonance, or other method described herein, in some embodiments, at 37°C.

[0211] In some embodiments, antibodies or antigen binding fragments of the invention bind to MOSPD2 with a K_{on} of from about 10^3 1/Ms to about 10^6 1/Ms, or any range of values thereof (e.g., from about 10^3 1/Ms to about 10^5 1/Ms, from about 10^4 1/Ms to about 10^5 1/Ms, from about 10^4 1/Ms to about 10^6 1/Ms, from about 10^5 1/Ms to about 10^6 1/Ms, or from about 10^3 1/Ms to about 10^4 1/Ms). In other embodiments, the antibody or antigen binding fragment has a K_{on} of about 10^3 1/Ms, about 10^4 1/Ms, about 10^5 1/Ms, or about 10^6 1/Ms.

[0212] In some embodiments, antibodies or antigen binding fragments of the invention bind to MOSPD2 with a K_{off} of from about 10^{-3} 1/s to about 10^{-6} 1/s, or any range of values thereof (e.g., from about 10^{-3} 1/s to about 10^{-5} 1/s, from about 10^{-4} 1/s to about 10^{-5} 1/s, from about 10^{-4} 1/s to about 10^{-6} 1/s, from about 10^{-5} 1/s to about 10^{-6} 1/s, or from about 10^{-3} 1/s to about 10^{-4} 1/s). In other embodiments, the antibody or antigen binding fragment has a K_{off} of about 10^{-3} 1/s, about 10^{-4} 1/s, about 10^{-5} 1/s, or about 10^{-6} 1/s.

[0213] In still other embodiments, the inhibitor of MOSPD2 is an RNAi, miRNA, siRNA, shRNA, an antisense RNA, an antisense DNA, a decoy molecule, a decoy DNA, a double-stranded DNA, a single-stranded DNA, a complexed DNA, an encapsulated DNA, a viral DNA, a plasmid DNA, a naked RNA, an encapsulated RNA, a viral RNA, a double-stranded RNA, a molecule capable of generating RNA interference, or combinations thereof. In some embodiments, the inhibitor hybridizes to a nucleotide sequence encoding a MOSPD2 polypeptide. In some embodiments, the hybridization is under a stringent condition or under a highly stringent condition.

[0214] In some embodiments, the inhibitor is a clustered regularly interspaced short palindromic repeats CRISPR-CAS9 system.

[0215] In further embodiments, a MOSPD2 polypeptide has a sequence at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identical to any one of SEQ ID NOs:1-4. In other embodiments, the MOSPD2 polypeptide has a sequence at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to any one of SEQ ID NOs:1-4. In other embodiments, the MOSPD2 polypeptide has a sequence of about

75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, or about 99% identical to any one of SEQ ID NOs:1-4. In other embodiments, the MOSPD2 polypeptide has a sequence with 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity to any one of SEQ ID NOs:1-4. In other embodiments, the MOSPD2 polypeptide has a sequence with from about 75% to 100% identity to any one of SEQ ID NOs:1-4, or any range of values thereof, for example, from about 80% to 100% identity, from about 85% to 100% identity, from about 90% to 100% identity, from about 95% to 100% identity, from about 96% to 100% identity, from about 97% to 100% identity, from about 98% to 100% identity, from about 99% to about 100% identity, from about 75% to about 99% identity, from about 80% to about 99% identity, from about 85% to about 99% identity, from about 90% to about 99% identity, from about 95% to about 99% identity, from about 96% to about 99% identity, from about 97% to about 99% identity, from about 98% to about 99% identity, from about 99% to about 100% identity, from about 75% to about 95% identity, from about 80% to about 95% identity, from about 85% to about 95% identity, or from about 90% to about 95% identity to any one of SEQ ID NOs:1-4.

[0216] In further embodiments of the invention, the MOSPD2 polypeptide is encoded by a polynucleotide sequence having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identity to any one of SEQ ID NOs:5-8. In other embodiments, the MOSPD2 polypeptide is encoded by a polynucleotide sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity to any one of SEQ ID NOs:5-8. In other embodiments, the MOSPD2 polypeptide is encoded by a polynucleotide sequence having about 75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, or about 99% identity to any one of SEQ ID NOs:5-8. In other embodiments, the MOSPD2 polypeptide is encoded by a polynucleotide sequence 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to any one of SEQ ID NOs:1-4. In other embodiments, the MOSPD2 polypeptide is encoded by a polynucleotide sequence having from about 75% to 100% identity to any one of SEQ ID NOs:5-8, or any range of values thereof, for example, from about 80% to 100% identity, from about 85% to 100% identity, from about 90% to 100% identity, from about 95% to 100% identity, from about

96% to 100% identity, from about 97% to 100% identity, from about 98% to 100% identity, from about 99% to about 100% identity, from about 75% to about 99% identity, from about 80% to about 99% identity, from about 85% to about 99% identity, from about 90% to about 99% identity, from about 95% to about 99% identity, from about 96% to about 99% identity, from about 97% to about 99% identity, from about 98% to about 99% identity, from about 99% to about 100% identity, from about 75% to about 95% identity, from about 80% to about 95% identity, from about 85% to about 95% identity, or from about 90% to about 95% identity to any one of SEQ ID NOs:5-8.

[0217] In any of the embodiments described herein, an inhibitor of MOSPD2 can be an inhibitor of MOSPD2 expressed by a cancer cell, *e.g.*, a human cancer cell. Non-exclusive listing of types of cancer cells include cells of bladder cancer, breast cancer, colon cancer, rectal cancer, kidney cancer, liver cancer, lung cancer, esophageal cancer, gall-bladder cancer, ovarian cancer, pancreatic cancer, stomach cancer, cervical cancer, thyroid cancer, prostate cancer, skin cancer, hematopoietic cancer, cancer of mesenchymal origin, cancer of central or peripheral nervous system, endometrial cancer, head and neck cancer, glioblastoma, and malignant ascites. In some embodiments, the cancer is a small cell lung cancer or a non-small-cell lung cancer. In some embodiments, the cancer is skin cancer, *e.g.*, squamous cell carcinoma, basal cell cancer, melanoma, dermatofibrosarcoma protuberans, Merkel cell carcinoma, Kaposi's sarcoma, keratoacanthoma, spindle cell tumors, sebaceous carcinomas, microcystic adnexal carcinoma, Paget's disease of the breast, atypical fibroxanthoma, leiomyosarcoma, or angiosarcoma. In some embodiments, the cancer is a hematopoietic cancer of lymphoid lineage, *e.g.*, leukemia, acute lymphocytic leukemia, chronic lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma or Burkitt's lymphoma. In some embodiments, the cancer is a hematopoietic cancer of myeloid lineage, *e.g.*, fibrosarcoma, rhabdomyosarcoma, soft tissue sarcoma, or bone sarcoma. In some embodiments, the cancer is a cancer of the central or peripheral nervous system, *e.g.*, astrocytoma, neuroblastoma, glioma, or schwannomas. In some embodiments, the cancer is anal cancer, bone cancer, gastrointestinal stromal cancer, gestational trophoblastic disease, Hodgkin's lymphoma, Kaposi sarcoma, keratoacanthoma, malignant mesothelioma, multicentric castleman disease, multiple myeloma and other plasma cell neoplasms,

myeloproliferative neoplasms, neuroblastoma, non-Hodgkin's lymphoma, osteosarcoma, ovarian, fallopian tube, or primary peritoneal cancer, penile cancer, retinoblastoma, rhabdomyosarcoma, seminoma, soft tissue sarcoma, stomach (gastric) cancer, testicular cancer, teratocarcinoma, thyroid follicular cancer, vaginal cancer, vulvar cancer, Wilms tumor and other childhood kidney cancers, and xeroderma pigmentosum. In some embodiments, the cancer is bladder cancer, brain cancer (*e.g.*, cerebrum astrocytoma), breast cancer, colon cancer (*e.g.*, colon adenocarcinoma), esophageal cancer (*e.g.*, esophageal adenocarcinoma), lung cancer, skin cancer (*e.g.*, melanoma), tongue cancer (*e.g.*, head and neck (tongue) cell carcinoma), kidney cancer (*e.g.*, kidney clear cell carcinoma), or hepatic cancer (*e.g.*, hepatocellular carcinoma).

Pharmaceutical Compositions

[0218] Other embodiments of the invention relate to a pharmaceutical composition comprising an inhibitor of MOSPD2, *e.g.*, MOSPD2 expressed by a cancer cell. In some embodiments, the pharmaceutical composition comprises an inhibitor of MOSPD2, *e.g.*, MOSPD2 expressed by a cancer cell and a pharmaceutically acceptable carrier. In other embodiments, the pharmaceutical composition comprises a therapeutically effective amount of the inhibitor of MOSPD2, *e.g.*, MOSPD2 expressed by a cancer cell. Exemplary inhibitors of MOSPD2 are described herein. Suitable types of cancer are also described herein.

μg/ml, from about 2 μg/ml to about 7 μg/ml, from about 3 μg/ml to about 7 μg/ml, from about 4 μg/ml to about 7 μg/ml, from about 5 μg/ml to about 7 μg/ml, from about 6 μg/ml to about 7 μg/ml, from about 1 μg/ml to about 6 μg/ml, from about 2 μg/ml to about 6 μg/ml, from about 3 μg/ml to about 6 μg/ml, from about 4 μg/ml to about 6 μg/ml, from about 5 μg/ml to about 6 μg/ml, from about 1 μg/ml to about 5 μg/ml, from about 2 μg/ml to about 5 μg/ml, from about 3 μg/ml to about 5 μg/ml, from about 4 μg/ml to about 5 μg/ml, from about 1 μg/ml to about 4 μg/ml, from about 2 μg/ml to about 4 μg/ml, from about 3 μg/ml to about 4 μg/ml, from about 1 μg/ml to about 3 μg/ml, from about 2 μg/ml to about 3 μg/ml, or from about 1 μg/ml to about 2 μg/ml). In other embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is about 1 μg/ml, about 2 μg/ml, about 3 μg/ml, about 4 μg/ml, about 5 μg/ml, about 6 μg/ml, about 7 μg/ml, about 8 μg/ml, about 9 μg/ml, or about 10 μg/ml.

[0220] In some embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is from about 10 mg/kg to about 40 mg/kg, or any range of values thereof (e.g., from about 15 mg/kg to about 40 mg/kg, from about 20 mg/kg to about 40 mg/kg, from about 25 mg/kg to about 40 mg/kg, from about 30 mg/kg to about 40 mg/kg, from about 35 mg/kg to about 40 mg/kg, from about 10 mg/kg to about 35 mg/kg, from about 15 mg/kg to about 35 mg/kg, from about 20 mg/kg to about 35 mg/kg, from about 25 mg/kg to about 35 mg/kg, from about 30 mg/kg to about 35 mg/kg, from about 10 mg/kg to about 30 mg/kg, from about 15 mg/kg to about 30 mg/kg, from about 20 mg/kg to about 30 mg/kg, from about 25 mg/kg to about 30 mg/kg, from about 10 mg/kg to about 25 mg/kg, from about 15 mg/kg to about 25 mg/kg, from about 20 mg/kg to about 25 mg/kg, from about 10 mg/kg to about 20 mg/kg, from about 15 mg/kg to about 20 mg/kg, or from about 10 mg/kg to about 15 mg/kg). In other embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is about 10 mg/kg, about 15 mg/kg, about 20 mg/kg, about 25 mg/kg, about 30 mg/kg, about 35 mg/kg, or about 40 mg/kg.

[0221] In some embodiments, the inhibitor of MOSPD2 (e.g., an antibody or antigen binding fragment thereof) is present in an amount such that administration of the MOSPD2 inhibitor causes at least 10% (e.g., at least 20%, at least 30%, at least 40%, at

least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 99%, or higher) inhibition of one or more activities of MOSPD2 (*e.g.*, human MOSPD2, *e.g.*, MOSPD2 expressed by a human cancer cell) (*e.g.*, MOSPD2 expression, cancer cell migration, monocyte migration associated with tumor growth (*e.g.*, presence of tumor associated macrophages), a chemokine signaling pathway, a growth factor signaling pathway, EGF receptor phosphorylation, ERK phosphorylation, AKT phosphorylation, and/or FAK phosphorylation). In some embodiments, administration of the MOSPD2 inhibitor to a human subject causes at least 10% (*e.g.*, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 99%, or higher) inhibition of one or more activities of a human MOSPD2, *e.g.*, MOSPD2 expressed by a human cancer cell.

[0222] In another aspect, administration of the MOSPD2 inhibitor (*e.g.*, an anti-MOSPD2 antibody or antigen binding fragment thereof) causes from about 10% to 100%, from about 10% to about 99%, from about 10% to about 95%, from about 10% to about 90%, from about 10% to about 85%, from about 10% to about 80%, from about 10% to about 70%, from about 20% to about 99%, from about 20% to about 95%, from about 20% to about 90%, from about 20% to about 85%, from about 20% to about 80%, from about 30% to about 95%, from about 30% to about 90%, from about 30% to about 85%, from about 30% to about 80%, from about 40% to about 95%, from about 40% to about 90%, from about 40% to about 85%, from about 40% to about 80%, from about 50% to about 95%, from about 50% to about 90%, from about 50% to about 85%, from about 50% to about 80%, from about 60% to about 95%, from about 60% to about 90%, from about 60% to about 85%, or from about 60% to about 80% inhibition of one or more activities of MOSPD2, *e.g.*, regulation of cancer cell migration, monocyte migration associated with tumor growth, presence of tumor associated macrophages, chemokine signaling pathways, growth factor signaling pathways, EGF receptor phosphorylation, ERK phosphorylation, AKT phosphorylation, and/or FAK phosphorylation.

[0223] As used herein, a "pharmaceutical composition" refers to a preparation of one or more agents as described herein (*e.g.*, a MOSPD2 inhibitor, or a MOSPD2 inhibitor with one or more other agents described herein), or physiologically acceptable salts or prodrugs thereof, with other chemical components, including, but not limited to, pharmaceutically acceptable carriers, excipients, lubricants, buffering agents, antibacterial

agents, bulking agents (*e.g.*, mannitol), antioxidants (*e.g.*, ascorbic acid or sodium bisulfite), and the like. The purpose of the pharmaceutical composition is to facilitate administration of the agent(s) to a subject.

[0224] As used herein, "administration" or "administering" to a subject includes, but is not limited to, the act of a physician or other medical professional prescribing a pharmaceutical composition of the invention for a subject. Administration can be local administration, *e.g.*, intra-tumor administration.

[0225] Herein, the phrase "pharmaceutically acceptable carrier" refers to a carrier or a diluent that does not cause significant irritation to the subject and does not abrogate the biological activity and properties of the agent(s) described herein.

[0226] As used herein, the term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered.

[0227] Herein the term "excipient" refers to an inert substance added to a pharmaceutical composition to further facilitate administration of an active ingredient.

[0228] In some embodiments, a pharmaceutical composition comprising a MOSPD2 inhibitor further comprises one or more additional active agents. In some embodiments, the one or more additional active agent is an anticancer drug. In some embodiments, the one or more additional active agent is an anti-proliferative agent.

[0229] In any of the embodiments described herein, useful anticancer drugs include those known in the art, for example, those anticancer drugs approved for use by a regulatory agency such as the U.S. Food and Drug Administration (US FDA) or the like. Some of the useful anticancer drugs are listed by the U.S. National Cancer Institute at <http://www.cancer.gov/about-cancer/treatment/drugs>. Exemplary useful anticancer drugs include those approved (*e.g.*, by the US FDA) for anal cancer, bladder cancer, bone cancer, brain cancer, breast cancer, cervical cancer, colon and rectal cancer, endometrial cancer, esophageal cancer, gastrointestinal stromal cancer, gestational trophoblastic disease, head and neck cancer, Hodgkin's lymphoma, Kaposi sarcoma, kidney (renal cell) cancer, leukemia, liver cancer, lung cancer, malignant mesothelioma, melanoma, multicentric Castleman disease, multiple myeloma and other plasma cell neoplasms, myeloproliferative neoplasms, neuroblastoma, non-Hodgkin's lymphoma, ovarian, fallopian tube, or primary peritoneal cancer, pancreatic cancer, penile cancer, prostate cancer, retinoblastoma, rhabdomyosarcoma, skin cancer, soft tissue sarcoma, stomach

(gastric) cancer, testicular cancer, thyroid cancer, vaginal cancer, vulvar cancer, and Wilms tumor and other childhood kidney cancers.

[0230] In any of the embodiments described herein, the anticancer drug can be selected from the group consisting of Abiraterone Acetate, Abitrexate (Methotrexate), Abraxane (Paclitaxel Albumin-stabilized Nanoparticle Formulation), ABVD, ABVE, ABVE-PC, AC, AC-T, Adcetris (Brentuximab Vedotin), ADE, Ado-Trastuzumab Emtansine, Adriamycin (Doxorubicin Hydrochloride), Adrucil (Fluorouracil), Afatinib Dimaleate, Afinitor (Everolimus), Akynzeo (Netupitant and Palonosetron Hydrochloride), Aldara (Imiquimod), Aldesleukin, Alemtuzumab, Alimta (Pemetrexed Disodium), Aloxi (Palonosetron Hydrochloride), Ambochlorin (Chlorambucil), Amboclorin (Chlorambucil), Aminolevulinic Acid, Anastrozole, Aprepitant, Aredia (Pamidronate Disodium), Arimidex (Anastrozole), Aromasin (Exemestane), Arranon (Nelarabine), Arsenic Trioxide, Arzerra (Ofatumumab), Asparaginase Erwinia chrysanthemi, Avastin (Bevacizumab), Axitinib, Azacitidine, BEACOPP, Becenum (Carmustine), Beleodaq (Belinostat), Belinostat, Bendamustine Hydrochloride, BEP, Bevacizumab, Bexarotene, Bexxar (Tositumomab and I 131 Iodine Tositumomab), Bicalutamide, BiCNU (Carmustine), Bleomycin, Blinatumomab, Blincyto (Blinatumomab), Bortezomib, Bosulif (Bosutinib), Bosutinib, Brentuximab Vedotin, Busulfan, Busulfex (Busulfan), Cabazitaxel, Cabozantinib-S-Malate, CAF, Campath (Alemtuzumab), Camptosar (Irinotecan Hydrochloride), Capecitabine, CAPOX, Carboplatin, Carboplatin-Taxol, Carfilzomib, Carmubris (Carmustine), Carmustine, Carmustine Implant, Casodex (Bicalutamide), CeeNU (Lomustine), Ceritinib, Cerubidine (Daunorubicin Hydrochloride), Cervarix (Recombinant HPV Bivalent Vaccine), Cetuximab, Chlorambucil, Chlorambucil-Prednisone, CHOP, Cisplatin, Clafen (Cyclophosphamide), Clofarabine, CMF, Cometriq (Cabozantinib-S-Malate), COPP, COPP-ABV, Cosmegen (Dactinomycin), Crizotinib, CVP, Cyclophosphamide, Cyfos (Ifosfamide), Cyramza (Ramucirumab), Cytarabine, Cytarabine, Liposomal, Cytosar-U (Cytarabine), Cytoxan (Cyclophosphamide), Dabrafenib, Dacarbazine, Dacogen (Decitabine), Dactinomycin, Dasatinib, Daunorubicin Hydrochloride, Decitabine, Degarelix, Denileukin Diftitox, Denosumab, DepoCyt (Liposomal Cytarabine), DepoFoam (Liposomal Cytarabine), Dexrazoxane Hydrochloride, Dinutuximab, Docetaxel, Doxil (Doxorubicin Hydrochloride Liposome), Doxorubicin Hydrochloride, Doxorubicin Hydrochloride

Liposome, Dox-SL (Doxorubicin Hydrochloride Liposome), DTIC-Dome (Dacarbazine), Efudex (Fluorouracil), Elitek (Rasburicase), Ellence (Epirubicin Hydrochloride), Eloxatin (Oxaliplatin), Eltrombopag Olamine, Emend (Aprepitant), Enzalutamide, Epirubicin Hydrochloride, EPOCH, Erbitux (Cetuximab), Eribulin Mesylate, Erivedge (Vismodegib), Erlotinib Hydrochloride, Erwinaze (Asparaginase Erwinia chrysanthemi), Etopophos (Etoposide Phosphate), Etoposide, Etoposide Phosphate, Evacet (Doxorubicin Hydrochloride Liposome), Everolimus, Evista (Raloxifene Hydrochloride), Exemestane, Fareston (Toremifene), Farydak (Panobinostat), Faslodex (Fulvestrant), FEC, Femara (Letrozole), Filgrastim, Fludara (Fludarabine Phosphate), Fludarabine Phosphate, Fluoroplex (Fluorouracil), Fluorouracil, Folex (Methotrexate), Folex PFS (Methotrexate), Folfiri, Folfiri-Bevacizumab, Folfiri-Cetuximab, Folfirinox, Folfox, Folotyn (Pralatrexate), FU-LV, Fulvestrant, Gardasil (Recombinant HPV Quadrivalent Vaccine), Gardasil 9 (Recombinant HPV Nonavalent Vaccine), Gazyva (Obinutuzumab), Gefitinib, Gemcitabine Hydrochloride, Gemcitabine-Cisplatin, Gemcitabine-Oxaliplatin, Gemtuzumab Ozogamicin, Gemzar (Gemcitabine Hydrochloride), Gilotrif (Afatinib Dimaleate), Gleevec (Imatinib Mesylate), Gliadel (Carmustine Implant), Gliadel wafer (Carmustine Implant), Glucarpidase, Goserelin Acetate, Halaven (Eribulin Mesylate), Herceptin (Trastuzumab), HPV Bivalent Vaccine, Recombinant, HPV Nonavalent Vaccine, Recombinant, HPV Quadrivalent Vaccine, Recombinant, Hycamtin (Topotecan Hydrochloride), Hyper-CVAD, Ibrance (Palbociclib), Ibritumomab Tiuxetan, Ibrutinib, ICE, Iclusig (Ponatinib Hydrochloride), Idamycin (Idarubicin Hydrochloride), Idarubicin Hydrochloride, Idelalisib, Ifex (Ifosfamide), Ifosfamide, Ifosfamidum (Ifosfamide), Imatinib Mesylate, Imbruvica (Ibrutinib), Imiquimod, Inlyta (Axitinib), Intron A (Recombinant Interferon Alfa-2b), Iodine 131 Tositumomab and Tositumomab, Ipilimumab, Iressa (Gefitinib), Irinotecan Hydrochloride, Istodax (Romidepsin), Ixabepilone, Ixempra (Ixabepilone), Jakafi (Ruxolitinib Phosphate), Jevtana (Cabazitaxel), Kadcyla (Ado-Trastuzumab Emtansine), Keoxifene (Raloxifene Hydrochloride), Kepivance (Palifermin), Keytruda (Pembrolizumab), Kyprolis (Carfilzomib), Lanreotide Acetate, Lapatinib Ditosylate, Lenalidomide, Lenvatinib Mesylate, Lenvima (Lenvatinib Mesylate), Letrozole, Leucovorin Calcium, Leukeran (Chlorambucil), Leuprolide Acetate, Levulan (Aminolevulinic Acid), Linfolizin (Chlorambucil), LipoDox (Doxorubicin Hydrochloride Liposome), Liposomal

Cytarabine, Lomustine, Lupron (Leuprolide Acetate), Lupron Depot (Leuprolide Acetate), Lupron Depot-Ped (Leuprolide Acetate), Lupron Depot-3 Month (Leuprolide Acetate), Lupron Depot-4 Month (Leuprolide Acetate), Lynparza (Olaparib), Marqibo (Vincristine Sulfate Liposome), Matulane (Procarbazine Hydrochloride), Mechlorethamine Hydrochloride, Megace (Megestrol Acetate), Megestrol Acetate, Mekinist (Trametinib), Mercaptopurine, Mesna, Mesnex (Mesna), Methazolastone (Temozolomide), Methotrexate, Methotrexate LPF (Methotrexate), Mexate (Methotrexate), Mexate-AQ (Methotrexate), Mitomycin C, Mitoxantrone Hydrochloride, Mitozytrex (Mitomycin C), MOPP, Mozobil (Plerixafor), Mustargen (Mechlorethamine Hydrochloride), Mutamycin (Mitomycin C), Myleran (Busulfan), Mylosar (Azacitidine), Mylotarg (Gemtuzumab Ozogamicin), Nanoparticle Paclitaxel (Paclitaxel Albumin-stabilized Nanoparticle Formulation), Navelbine (Vinorelbine Tartrate), Nelarabine, Neosar (Cyclophosphamide), Netupitant and Palonosetron Hydrochloride, Neupogen (Filgrastim), Nexavar (Sorafenib Tosylate), Nilotinib, Nivolumab, Nolvadex (Tamoxifen Citrate), Nplate (Romiplostim), Obinutuzumab, OEPa, Ofatumumab, OFF, Olaparib, Omacetaxine Mepesuccinate, Oncaspar (Pegaspargase), Ontak (Denileukin Diftitox), Opdivo (Nivolumab), OPPA, Oxaliplatin, Paclitaxel, Paclitaxel Albumin-stabilized Nanoparticle Formulation, PAD, Palbociclib, Palifermin, Palonosetron Hydrochloride, Pamidronate Disodium, Panitumumab, Panobinostat, Paraplat (Carboplatin), Paraplatin (Carboplatin), Pazopanib Hydrochloride, Pegaspargase, Peginterferon Alfa-2b, PEG-Intron (Peginterferon Alfa-2b), Pembrolizumab, Pemetrexed Disodium, Perjeta (Pertuzumab), Pertuzumab, Platinol (Cisplatin), Platinol-AQ (Cisplatin), Plerixafor, Pomalidomide, Pomalyst (Pomalidomide), Ponatinib Hydrochloride, Pralatrexate, Prednisone, Procarbazine Hydrochloride, Proleukin (Aldesleukin), Prolia (Denosumab), Promacta (Eltrombopag Olamine), Provenge (Sipuleucel-T), Purinethol (Mercaptopurine), Purixan (Mercaptopurine), Radium 223 Dichloride, Raloxifene Hydrochloride, Ramucirumab, Rasburicase, R-CHOP, R-CVP, Recombinant Human Papillomavirus (HPV) Bivalent Vaccine, Recombinant Human Papillomavirus (HPV) Nonavalent Vaccine, Recombinant Human Papillomavirus (HPV) Quadrivalent Vaccine, Recombinant Interferon Alfa-2b, Regorafenib, R-EPOCH, Revlimid (Lenalidomide), Rheumatrex (Methotrexate), Rituxan (Rituximab), Rituximab, Romidepsin, Romiplostim, Rubidomycin (Daunorubicin Hydrochloride), Ruxolitinib Phosphate, Sclerosol

Intrapleural Aerosol (Talc), Siltuximab, Sipuleucel-T, Somatuline Depot (Lanreotide Acetate), Sorafenib Tosylate, Sprycel (Dasatinib), STANFORD V, Sterile Talc Powder (Talc), Steritalc (Talc), Stivarga (Regorafenib), Sunitinib Malate, Sutent (Sunitinib Malate), Sylatron (Peginterferon Alfa-2b), Sylvant (Siltuximab), Synovir (Thalidomide), Synribo (Omacetaxine Mepesuccinate), TAC, Tafinlar (Dabrafenib), Talc, Tamoxifen Citrate, Tarabine PFS (Cytarabine), Tarceva (Erlotinib Hydrochloride), Targretin (Bexarotene), Tasigna (Nilotinib), Taxol (Paclitaxel), Taxotere (Docetaxel), Temodar (Temozolomide), Temozolomide, Temsirolimus, Thalidomide, Thalomid (Thalidomide), Thiotepa, Toposar (Etoposide), Topotecan Hydrochloride, Toremifene, Torisel (Temsirolimus), Tositumomab and I 131 Iodine Tositumomab, Totect (Dexrazoxane Hydrochloride), TPF, Trametinib, Trastuzumab, Treanda (Bendamustine Hydrochloride), Trisenox (Arsenic Trioxide), Tykerb (Lapatinib Ditosylate), Unituxin (Dinutuximab), Vandetanib, VAMP, Vectibix (Panitumumab), VeIP, Velban (Vinblastine Sulfate), Velcade (Bortezomib), Velsar (Vinblastine Sulfate), Vemurafenib, VePesid (Etoposide), Viadur (Leuprolide Acetate), Vidaza (Azacitidine), Vinblastine Sulfate, Vincasar PFS (Vincristine Sulfate), Vincristine Sulfate, Vincristine Sulfate Liposome, Vinorelbine Tartrate, VIP, Vismodegib, Voraxaze (Glucarpidase), Vorinostat, Votrient (Pazopanib Hydrochloride), Wellcovorin (Leucovorin Calcium), Xalkori (Crizotinib), Xeloda (Capecitabine), XELIRI, XELOX, Xgeva (Denosumab), Xofigo (Radium 223 Dichloride), Xtandi (Enzalutamide), Yervoy (Ipilimumab), Zaltrap (Ziv-Aflibercept), Zelboraf (Vemurafenib), Zevalin (Ibritumomab Tiuxetan), Zinecard (Dexrazoxane Hydrochloride), Ziv-Aflibercept, Zoladex (Goserelin Acetate), Zoledronic Acid, Zolinza (Vorinostat), Zometa (Zoledronic Acid), Zydelig (Idelalisib), Zykadia (Ceritinib), and Zytiga (Abiraterone Acetate).

[0231] When two or more agents are administered as a pharmaceutical composition, each agent may optionally be administered in a separate composition and/or via a different route of administration. Possible routes of administration for each agent independently include, but are not limited to, parenteral administration, transmucosal administration, rectal administration, buccal administration and/or inhalation (e.g., as described herein).

[0232] In some embodiments, the pharmaceutical composition is suitable for systemic or local administration. In other embodiments, the pharmaceutical composition is suitable for nasal, oral, or intra-peritoneal administration. In other embodiments, the

pharmaceutical composition is suitable for intravenous administration, intramuscular administration or subcutaneous administration. In other embodiments, the pharmaceutical composition is suitable for intra-tumor administration.

Methods of Use in Cancer and Metastasis

[0233] In some embodiments, the present invention relates to the discovery that MOSPD2 expression in cancer cells is upregulated when compared to its non-cancerous counterpart. As shown in the Examples section, MOSPD2 expression was positively identified in various types of cancer cells, *e.g.*, bladder cancer, brain cancer (*e.g.*, cerebrum astrocytoma), breast cancer, colon cancer (*e.g.*, colon adenocarcinoma), esophageal cancer (*e.g.*, esophageal adenocarcinoma), lung cancer, skin cancer (*e.g.*, melanoma), tongue cancer (*e.g.*, head and neck (tongue) cell carcinoma), kidney cancer (*e.g.*, kidney clear cell carcinoma), and hepatic cancer (*e.g.*, hepatocellular carcinoma), but not in the counterpart normal, non-cancerous cells. Further, inhibition of MOSPD2 by silencing of MOSPD2 expression or administration of anti-MOSPD2 F(ab')₂ mAb in various types of cancer cells inhibits migration and metastasis of cancer cells both *in vitro* and *in vivo*.

[0234] Embodiments of the invention relate to methods for treating, preventing, or reducing the incidence of metastasis of a cancer cell with an inhibitor of MOSPD2. In some embodiments, the methods include administering to a subject in need thereof an effective amount of an inhibitor of MOSPD2. In other embodiments, the MOSPD2 is expressed by the cancer cell. Exemplary MOSPD2 inhibitors include those described herein. Suitable types of cancer are also described herein. In some embodiments, the cancer is bladder cancer, brain cancer (*e.g.*, cerebrum astrocytoma), breast cancer, colon cancer (*e.g.*, colon adenocarcinoma), esophageal cancer (*e.g.*, esophageal adenocarcinoma), lung cancer, skin cancer (*e.g.*, melanoma), tongue cancer (*e.g.*, head and neck (tongue) cell carcinoma), kidney cancer (*e.g.*, kidney clear cell carcinoma), or hepatic cancer (*e.g.*, hepatocellular carcinoma).

[0235] Embodiments of the invention also relate to methods of treating, preventing, or reducing the incidence of a cancer with an inhibitor of MOSPD2. In some embodiments, the methods include administering to a subject in need thereof a therapeutically effective amount of an inhibitor of MOSPD2. In other embodiments, the MOSPD2 is expressed by the cancer cells. In some embodiments, the methods further comprise administering a

therapeutically effective amount of an inhibitor of MOSPD2 and another anticancer drug. Exemplary MOSPD2 inhibitors and anticancer drugs include those described herein. Suitable types of cancer are also described herein. In some embodiments, the cancer is bladder cancer, brain cancer (*e.g.*, cerebrum astrocytoma), breast cancer, colon cancer (*e.g.*, colon adenocarcinoma), esophageal cancer (*e.g.*, esophageal adenocarcinoma), lung cancer, skin cancer (*e.g.*, melanoma), tongue cancer (*e.g.*, head and neck (tongue) cell carcinoma), kidney cancer (*e.g.*, kidney clear cell carcinoma), or hepatic cancer (*e.g.*, hepatocellular carcinoma).

[0236] Embodiments of the invention also relate to methods of treating, preventing, or reducing the incidence of a metastatic cancer with an inhibitor of MOSPD2. In some embodiments, the methods include administering to a subject in need thereof a therapeutically effective amount of an inhibitor of MOSPD2 and another anticancer drug. In some embodiments, the administration is local, *e.g.*, intra-tumor administration. In some embodiments, the MOSPD2 is expressed by the metastatic cancer cells. Exemplary MOSPD2 inhibitors and anticancer drugs include those described herein. Suitable types of cancer are also described herein. In some embodiments, the cancer is bladder cancer, brain cancer (*e.g.*, cerebrum astrocytoma), breast cancer, colon cancer (*e.g.*, colon adenocarcinoma), esophageal cancer (*e.g.*, esophageal adenocarcinoma), lung cancer, skin cancer (*e.g.*, melanoma), tongue cancer (*e.g.*, head and neck (tongue) cell carcinoma), kidney cancer (*e.g.*, kidney clear cell carcinoma), or hepatic cancer (*e.g.*, hepatocellular carcinoma).

[0237] In some embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody described herein, *e.g.*, having an antibody-antigen equilibrium dissociation constant (K_D) of from about 10^{-6} M to about 10^{-12} M, or any range of values thereof (*e.g.*, from about 10^{-7} M to about 10^{-12} , from 10^{-8} M to about 10^{-12} M, from about 10^{-9} M to about 10^{-12} M, from about 10^{-10} M to about 10^{-12} M, from about 10^{-11} M to about 10^{-12} M, from about 10^{-6} M to about 10^{-11} M, from about 10^{-7} M to about 10^{-11} M, from about 10^{-8} M to about 10^{-11} M, from about 10^{-9} M to about 10^{-11} M, from about 10^{-10} M to about 10^{-11} M, from about 10^{-6} M to about 10^{-10} M, from about 10^{-7} M to about 10^{-10} M, from about 10^{-8} M to about 10^{-10} M, from about 10^{-9} M to about 10^{-10} M, from about 10^{-6} M to about 10^{-9} M, from about 10^{-7} M to about 10^{-9} M, from about 10^{-8} M to about 10^{-9} M, from about 10^{-6} M to about 10^{-8} M, or from about 10^{-7} M to about 10^{-8}). In other

embodiments, the antibody or antigen binding fragment thereof has a K_D of about 10^{-6} M, about 10^{-7} M, about 10^{-8} M, about 10^{-9} M, about 10^{-10} M, about 10^{-11} M, or about 10^{-12} M. In some embodiments, the antibody or antigen binding fragment binds to one or more epitopes on MOSPD2. In some embodiments, the K_D is determined by Scatchard analysis, surface plasmon resonance, or other method described herein, in some embodiments, at 37°C.

[0238] In some embodiments, antibodies or antigen binding fragments of the invention bind to MOSPD2 with a K_{on} of from about 10^3 1/Ms to about 10^6 1/Ms, or any range of values thereof (e.g., from about 10^3 1/Ms to about 10^5 1/Ms, from about 10^4 1/Ms to about 10^5 1/Ms, from about 10^4 1/Ms to about 10^6 1/Ms, from about 10^5 1/Ms to about 10^6 1/Ms, or from about 10^3 1/Ms to about 10^4 1/Ms). In other embodiments, the antibody or antigen binding fragment has a K_{on} of about 10^3 1/Ms, about 10^4 1/Ms, about 10^5 1/Ms, or about 10^6 1/Ms.

[0239] In some embodiments, antibodies or antigen binding fragments of the invention bind to MOSPD2 with a K_{off} of from about 10^{-3} 1/s to about 10^{-6} 1/s, or any range of values thereof (e.g., from about 10^{-3} 1/s to about 10^{-5} 1/s, from about 10^{-4} 1/s to about 10^{-5} 1/s, from about 10^{-4} 1/s to about 10^{-6} 1/s, from about 10^{-5} 1/s to about 10^{-6} 1/s, or from about 10^{-3} 1/s to about 10^{-4} 1/s). In other embodiments, the antibody or antigen binding fragment has a K_{off} of about 10^{-3} 1/s, about 10^{-4} 1/s, about 10^{-5} 1/s, or about 10^{-6} 1/s.

[0240] In some embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is from about 1 μ g/ml to about 10 μ g/ml, or any range of values thereof (e.g., from about 2 μ g/ml to about 10 μ g/ml, from about 3 μ g/ml to about 10 μ g/ml, from about 4 μ g/ml to about 10 μ g/ml, from about 5 μ g/ml to about 10 μ g/ml, from about 6 μ g/ml to about 10 μ g/ml, from about 7 μ g/ml to about 10 μ g/ml, from about 8 μ g/ml to about 10 μ g/ml, from about 9 μ g/ml to about 10 μ g/ml, from about 1 μ g/ml to about 9 μ g/ml, from about 2 μ g/ml to about 9 μ g/ml, from about 3 μ g/ml to about 9 μ g/ml, from about 4 μ g/ml to about 9 μ g/ml, from about 5 μ g/ml to about 9 μ g/ml, from about 6 μ g/ml to about 9 μ g/ml, from about 7 μ g/ml to about 9 μ g/ml, from about 8 μ g/ml to about 9 μ g/ml, from about 1 μ g/ml to about 8 μ g/ml, from about 2 μ g/ml to about 8 μ g/ml, from about 3 μ g/ml to about 8 μ g/ml, from about 4 μ g/ml to about 8 μ g/ml, from about 5 μ g/ml to about 8 μ g/ml, from about 6 μ g/ml to about 8 μ g/ml, from about 7 μ g/ml to about 8 μ g/ml, from about 1 μ g/ml to about 7 μ g/ml, from about 2 μ g/ml to about 7 μ g/ml, from about 3 μ g/ml to about 7 μ g/ml, from about 4 μ g/ml to about 7 μ g/ml, from about 5 μ g/ml to about 7 μ g/ml, from about 6 μ g/ml to about 7 μ g/ml, from about 7 μ g/ml to about 7 μ g/ml, from about 1 μ g/ml to about 6 μ g/ml, from about 2 μ g/ml to about 6 μ g/ml, from about 3 μ g/ml to about 6 μ g/ml, from about 4 μ g/ml to about 6 μ g/ml, from about 5 μ g/ml to about 6 μ g/ml, from about 6 μ g/ml to about 6 μ g/ml, from about 7 μ g/ml to about 6 μ g/ml, from about 8 μ g/ml to about 6 μ g/ml, from about 9 μ g/ml to about 6 μ g/ml, from about 1 μ g/ml to about 5 μ g/ml, from about 2 μ g/ml to about 5 μ g/ml, from about 3 μ g/ml to about 5 μ g/ml, from about 4 μ g/ml to about 5 μ g/ml, from about 5 μ g/ml to about 5 μ g/ml, from about 6 μ g/ml to about 5 μ g/ml, from about 7 μ g/ml to about 5 μ g/ml, from about 8 μ g/ml to about 5 μ g/ml, from about 9 μ g/ml to about 5 μ g/ml, from about 1 μ g/ml to about 4 μ g/ml, from about 2 μ g/ml to about 4 μ g/ml, from about 3 μ g/ml to about 4 μ g/ml, from about 4 μ g/ml to about 4 μ g/ml, from about 5 μ g/ml to about 4 μ g/ml, from about 6 μ g/ml to about 4 μ g/ml, from about 7 μ g/ml to about 4 μ g/ml, from about 8 μ g/ml to about 4 μ g/ml, from about 9 μ g/ml to about 4 μ g/ml, from about 1 μ g/ml to about 3 μ g/ml, from about 2 μ g/ml to about 3 μ g/ml, from about 3 μ g/ml to about 3 μ g/ml, from about 4 μ g/ml to about 3 μ g/ml, from about 5 μ g/ml to about 3 μ g/ml, from about 6 μ g/ml to about 3 μ g/ml, from about 7 μ g/ml to about 3 μ g/ml, from about 8 μ g/ml to about 3 μ g/ml, from about 9 μ g/ml to about 3 μ g/ml, from about 1 μ g/ml to about 2 μ g/ml, from about 2 μ g/ml to about 2 μ g/ml, from about 3 μ g/ml to about 2 μ g/ml, from about 4 μ g/ml to about 2 μ g/ml, from about 5 μ g/ml to about 2 μ g/ml, from about 6 μ g/ml to about 2 μ g/ml, from about 7 μ g/ml to about 2 μ g/ml, from about 8 μ g/ml to about 2 μ g/ml, from about 9 μ g/ml to about 2 μ g/ml, from about 1 μ g/ml to about 1 μ g/ml, from about 2 μ g/ml to about 1 μ g/ml, from about 3 μ g/ml to about 1 μ g/ml, from about 4 μ g/ml to about 1 μ g/ml, from about 5 μ g/ml to about 1 μ g/ml, from about 6 μ g/ml to about 1 μ g/ml, from about 7 μ g/ml to about 1 μ g/ml, from about 8 μ g/ml to about 1 μ g/ml, from about 9 μ g/ml to about 1 μ g/ml).

μg/ml, from about 2 μg/ml to about 7 μg/ml, from about 3 μg/ml to about 7 μg/ml, from about 4 μg/ml to about 7 μg/ml, from about 5 μg/ml to about 7 μg/ml, from about 6 μg/ml to about 7 μg/ml, from about 1 μg/ml to about 6 μg/ml, from about 2 μg/ml to about 6 μg/ml, from about 3 μg/ml to about 6 μg/ml, from about 4 μg/ml to about 6 μg/ml, from about 5 μg/ml to about 6 μg/ml, from about 1 μg/ml to about 5 μg/ml, from about 2 μg/ml to about 5 μg/ml, from about 3 μg/ml to about 5 μg/ml, from about 4 μg/ml to about 5 μg/ml, from about 1 μg/ml to about 4 μg/ml, from about 2 μg/ml to about 4 μg/ml, from about 3 μg/ml to about 4 μg/ml, from about 1 μg/ml to about 3 μg/ml, from about 2 μg/ml to about 3 μg/ml, or from about 1 μg/ml to about 2 μg/ml). In other embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is about 1 μg/ml, about 2 μg/ml, about 3 μg/ml, about 4 μg/ml, about 5 μg/ml, about 6 μg/ml, about 7 μg/ml, about 8 μg/ml, about 9 μg/ml, or about 10 μg/ml.

[0241] In some embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is from about 10 mg/kg to about 40 mg/kg, or any range of values thereof (e.g., from about 15 mg/kg to about 40 mg/kg, from about 20 mg/kg to about 40 mg/kg, from about 25 mg/kg to about 40 mg/kg, from about 30 mg/kg to about 40 mg/kg, from about 35 mg/kg to about 40 mg/kg, from about 10 mg/kg to about 35 mg/kg, from about 15 mg/kg to about 35 mg/kg, from about 20 mg/kg to about 35 mg/kg, from about 25 mg/kg to about 35 mg/kg, from about 30 mg/kg to about 35 mg/kg, from about 10 mg/kg to about 30 mg/kg, from about 15 mg/kg to about 30 mg/kg, from about 20 mg/kg to about 30 mg/kg, from about 25 mg/kg to about 30 mg/kg, from about 10 mg/kg to about 25 mg/kg, from about 15 mg/kg to about 25 mg/kg, from about 20 mg/kg to about 25 mg/kg, from about 10 mg/kg to about 20 mg/kg, from about 15 mg/kg to about 20 mg/kg, or from about 10 mg/kg to about 15 mg/kg). In other embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is about 10 mg/kg, about 15 mg/kg, about 20 mg/kg, about 25 mg/kg, about 30 mg/kg, about 35 mg/kg, or about 40 mg/kg.

[0242] In some embodiments, the subject is a mammal or a human. In other embodiments the MOSPD2 is a mammalian MOSPD2 or a human MOSPD2.

Methods of Inhibiting or Preventing One or More Activities in or of a Cancer Cell

[0243] Embodiments of the invention also relate to methods of inhibiting or preventing one or more activities in or of a cancer cell comprising administering an inhibitor of MOSPD2. In some embodiments, the methods comprise administering a therapeutically effective amount of an inhibitor of MOSPD2 to a subject in need thereof. In some embodiments, the administration is local administration, *e.g.*, intra-tumor administration. Exemplary MOSPD2 inhibitors include those described herein. Suitable types of cancer are also described herein.

[0244] In some embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody described herein, *e.g.*, having an antibody-antigen equilibrium dissociation constant (K_D) of from about 10^{-6} M to about 10^{-12} M, or any range of values thereof (*e.g.*, from about 10^{-7} M to about 10^{-12} , from 10^{-8} M to about 10^{-12} M, from about 10^{-9} M to about 10^{-12} M, from about 10^{-10} M to about 10^{-12} M, from about 10^{-11} M to about 10^{-12} M, from about 10^{-6} M to about 10^{-11} M, from about 10^{-7} M to about 10^{-11} M, from about 10^{-8} M to about 10^{-11} M, from about 10^{-9} M to about 10^{-11} M, from about 10^{-10} M to about 10^{-11} M, from about 10^{-6} M to about 10^{-10} M, from about 10^{-7} M to about 10^{-10} M, from about 10^{-8} M to about 10^{-10} M, from about 10^{-9} M to about 10^{-10} M, from about 10^{-6} M to about 10^{-9} M, from about 10^{-7} M to about 10^{-9} M, from about 10^{-8} M to about 10^{-9} M, from about 10^{-6} M to about 10^{-8} M, or from about 10^{-7} M to about 10^{-8}). In other embodiments, the antibody or antigen binding fragment thereof has a K_D of about 10^{-6} M, about 10^{-7} M, about 10^{-8} M, about 10^{-9} M, about 10^{-10} M, about 10^{-11} M, or about 10^{-12} M. In some embodiments, the antibody or antigen binding fragment binds to one or more epitopes on MOSPD2. In some embodiments, the K_D is determined by Scatchard analysis, surface plasmon resonance, or other method described herein, in some embodiments, at 37°C.

[0245] In some embodiments, antibodies or antigen binding fragments of the invention bind to MOSPD2 with a K_{on} of from about 10^3 1/Ms to about 10^6 1/Ms, or any range of values thereof (*e.g.*, from about 10^3 1/Ms to about 10^5 1/Ms, from about 10^4 1/Ms to about 10^5 1/Ms, from about 10^4 1/Ms to about 10^6 1/Ms, from about 10^5 1/Ms to about 10^6 1/Ms, or from about 10^3 1/Ms to about 10^4 1/Ms). In other embodiments, the antibody or antigen binding fragment has a K_{on} of about 10^3 1/Ms, about 10^4 1/Ms, about 10^5 1/Ms, or about 10^6 1/Ms.

[0246] In some embodiments, antibodies or antigen binding fragments of the invention bind to MOSPD2 with a K_{off} of from about 10^{-3} 1/s to about 10^{-6} 1/s, or any range of values thereof (e.g., from about 10^{-3} 1/s to about 10^{-5} 1/s, from about 10^{-4} 1/s to about 10^{-5} 1/s, from about 10^{-4} 1/s to about 10^{-6} 1/s, from about 10^{-5} 1/s to about 10^{-6} 1/s, or from about 10^{-3} 1/s to about 10^{-4} 1/s). In other embodiments, the antibody or antigen binding fragment has a K_{off} of about 10^{-3} 1/s, about 10^{-4} 1/s, about 10^{-5} 1/s, or about 10^{-6} 1/s.

[0247] In some embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is from about 1 μ g/ml to about 10 μ g/ml, or any range of values thereof (e.g., from about 2 μ g/ml to about 10 μ g/ml, from about 3 μ g/ml to about 10 μ g/ml, from about 4 μ g/ml to about 10 μ g/ml, from about 5 μ g/ml to about 10 μ g/ml, from about 6 μ g/ml to about 10 μ g/ml, from about 7 μ g/ml to about 10 μ g/ml, from about 8 μ g/ml to about 10 μ g/ml, from about 9 μ g/ml to about 10 μ g/ml, from about 1 μ g/ml to about 9 μ g/ml, from about 2 μ g/ml to about 9 μ g/ml, from about 3 μ g/ml to about 9 μ g/ml, from about 4 μ g/ml to about 9 μ g/ml, from about 5 μ g/ml to about 9 μ g/ml, from about 6 μ g/ml to about 9 μ g/ml, from about 7 μ g/ml to about 9 μ g/ml, from about 8 μ g/ml to about 9 μ g/ml, from about 1 μ g/ml to about 8 μ g/ml, from about 2 μ g/ml to about 8 μ g/ml, from about 3 μ g/ml to about 8 μ g/ml, from about 4 μ g/ml to about 8 μ g/ml, from about 5 μ g/ml to about 8 μ g/ml, from about 6 μ g/ml to about 8 μ g/ml, from about 7 μ g/ml to about 8 μ g/ml, from about 1 μ g/ml to about 7 μ g/ml, from about 2 μ g/ml to about 7 μ g/ml, from about 3 μ g/ml to about 7 μ g/ml, from about 4 μ g/ml to about 7 μ g/ml, from about 5 μ g/ml to about 7 μ g/ml, from about 6 μ g/ml to about 7 μ g/ml, from about 7 μ g/ml to about 7 μ g/ml, from about 1 μ g/ml to about 6 μ g/ml, from about 2 μ g/ml to about 6 μ g/ml, from about 3 μ g/ml to about 6 μ g/ml, from about 4 μ g/ml to about 6 μ g/ml, from about 5 μ g/ml to about 6 μ g/ml, from about 1 μ g/ml to about 5 μ g/ml, from about 2 μ g/ml to about 5 μ g/ml, from about 3 μ g/ml to about 5 μ g/ml, from about 4 μ g/ml to about 5 μ g/ml, from about 1 μ g/ml to about 4 μ g/ml, from about 2 μ g/ml to about 4 μ g/ml, from about 3 μ g/ml to about 4 μ g/ml, from about 1 μ g/ml to about 3 μ g/ml, from about 2 μ g/ml to about 3 μ g/ml, or from about 1 μ g/ml to about 2 μ g/ml). In other embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is about 1 μ g/ml, about 2 μ g/ml, about 3 μ g/ml, about 4 μ g/ml, about 5 μ g/ml, about 6 μ g/ml, about 7 μ g/ml, about 8 μ g/ml, about 9 μ g/ml, or about 10 μ g/ml.

[0248] In some embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is from about 10 mg/kg to about 40 mg/kg, or any range of values thereof (e.g., from about 15 mg/kg to about 40 mg/kg, from about 20 mg/kg to about 40 mg/kg, from about 25 mg/kg to about 40 mg/kg, from about 30 mg/kg to about 40 mg/kg, from about 35 mg/kg to about 40 mg/kg, from about 10 mg/kg to about 35 mg/kg, from about 15 mg/kg to about 35 mg/kg, from about 20 mg/kg to about 35 mg/kg, from about 25 mg/kg to about 35 mg/kg, from about 30 mg/kg to about 35 mg/kg, from about 10 mg/kg to about 30 mg/kg, from about 15 mg/kg to about 30 mg/kg, from about 20 mg/kg to about 30 mg/kg, from about 25 mg/kg to about 30 mg/kg, from about 10 mg/kg to about 25 mg/kg, from about 15 mg/kg to about 25 mg/kg, from about 20 mg/kg to about 25 mg/kg, from about 10 mg/kg to about 20 mg/kg, from about 15 mg/kg to about 20 mg/kg, or from about 10 mg/kg to about 15 mg/kg). In other embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is about 10 mg/kg, about 15 mg/kg, about 20 mg/kg, about 25 mg/kg, about 30 mg/kg, about 35 mg/kg, or about 40 mg/kg.

[0249] In some embodiments, the inhibitor of MOSPD2 (e.g., an antibody or antigen binding fragment thereof) causes at least about 10% (e.g., at least about 20%, at least about 30%, at least about 40%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 99%, or higher) inhibition of one or more activities of MOSPD2 (e.g., human MOSPD2) (e.g., MOSPD2 expression, cancer cell migration, monocyte migration associated with tumor growth (e.g., presence of tumor associated macrophages), a chemokine signaling pathway, a growth factor signaling pathway, EGF receptor phosphorylation, ERK phosphorylation, AKT phosphorylation, and/or FAK phosphorylation). In other embodiments, administration of the MOSPD2 inhibitor (e.g., an antibody or antigen binding fragment thereof) causes from about 10% to 100%, from about 10% to about 99%, from about 10% to about 95%, from about 10% to about 90%, from about 10% to about 85%, from about 10% to about 80%, from about 10% to about 70%, from about 20% to about 99%, from about 20% to about 95%, from about 20% to about 90%, from about 20% to about 85%, from about 20% to about 80%, from about 30% to about 95%, from about 30% to about 90%, from about 30% to about 85%, from

about 30% to about 80%, from about 40% to about 95%, from about 40% to about 90%, from about 40% to about 85%, from about 40% to about 80%, from about 50% to about 95%, from about 50% to about 90%, from about 50% to about 85%, from about 50% to about 50% to about 80%, from about 60% to about 95%, from about 60% to about 90%, from about 60% to about 85%, or from about 60% to about 80% inhibition of one or more activities of MOSPD2 (*e.g.*, human MOSPD2) (*e.g.*, MOSPD2 expression, cancer cell migration, monocyte migration associated with tumor growth (*e.g.*, presence of tumor associated macrophages), a chemokine signaling pathway, a growth factor signaling pathway, EGF receptor phosphorylation, ERK phosphorylation, AKT phosphorylation, and/or FAK phosphorylation). In one aspect, administration of an anti-MOSPD2 antibody or antigen binding fragment thereof causes from about 10% to 100%, from about 10% to about 99%, from about 10% to about 95%, from about 10% to about 90%, from about 10% to about 85%, from about 10% to about 80%, from about 10% to about 70%, from about 20% to about 99%, from about 20% to about 95%, from about 20% to about 90%, from about 20% to about 85%, from about 20% to about 80%, from about 30% to about 95%, from about 30% to about 90%, from about 30% to about 85%, from about 30% to about 80%, from about 40% to about 95%, from about 40% to about 90%, from about 40% to about 85%, from about 40% to about 80%, from about 50% to about 95%, from about 50% to about 90%, from about 50% to about 85%, from about 50% to about 80%, from about 60% to about 95%, from about 60% to about 90%, from about 60% to about 85%, or from about 60% to about 80% inhibition of cancer cell migration (*e.g.*, EGF-induced migration).

[0250] In some embodiments, the one or more activities is one or more of: MOSPD2 expression, cancer cell migration, monocyte migration associated with tumor growth (*e.g.*, presence of tumor associated macrophages), a chemokine signaling pathway, a growth factor signaling pathway, EGF receptor phosphorylation, ERK phosphorylation, AKT phosphorylation, and/or FAK phosphorylation. In some embodiments, at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, at least ten, or all of these activities are inhibited.

[0251] In some embodiments, at least cancer cell migration and a chemokine signaling pathway are inhibited. In other embodiments, the inhibiting of a chemokine signaling pathway or a growth factor signaling pathway is the inhibiting of EGF receptor

phosphorylation, ERK phosphorylation, AKT phosphorylation, and/or FAK phosphorylation. In other embodiments, the cancer cell migration or monocyte migration associated with tumor growth (e.g., presence of tumor associated macrophages) is induced by more than one chemokine or growth factor (e.g., EGF) or chemokine receptor or growth factor receptor (e.g., EGFR).

[0252] In some embodiments, the subject is a mammal or a human. In other embodiments the MOSPD2 is a mammalian MOSPD2 or a human MOSPD2.

Methods of Reducing Tumor Associated Macrophages or Tumor Associated Macrophage Migration

[0253] Tumor associated macrophages (TAMs) are often found in close proximity or within tumor masses. TAMs are known to be important for tumor growth. TAMs mostly originate from circulating monocytes and their recruitment into tumors is driven by tumor-derived chemotactic factors. TAMs promote tumor cell proliferation and metastasis by secreting a wide range of growth and proangiogenic factors. Consequently, many tumors with a high number of TAMs have an increased tumor growth rate, local proliferation and distant metastasis. In fact, the extent of TAM infiltration has been used as an inverse prognostic predictor in breast cancer, head and neck cancer, prostate and uterine cancer (R. D. Leek, R. Landers, S. B. Fox, F. Ng, A. L. Harris, C. E. Lewis, *British journal of cancer* 1998, 77, 2246; M. R. Young, M. A. Wright, Y. Lozano, M. M. Prechel, J. Benefield, J. P. Leonetti, S. L. Collins, G. J. Petruzzelli, *International Journal of Cancer* 1997, 74, 69; I. F. Lissbrant, P. Stattin, P. Wikstrom, J. E. Damberg, L. Egevad, A. Bergh, *International journal of oncology* 2000, 17, 445; H. B. Salvesen, L. A. Akslen, *International Journal of Cancer* 1999, 84, 538). TAMs are also prominent in tumor tissues, comprising up to 80% of the cell mass in breast carcinoma.

[0254] Some embodiments of the invention relate to methods for treating, reducing the incidence of, or preventing a cancer, which include administering to a subject in need thereof an effective amount of an inhibitor of MOSPD2, e.g., MOSPD2 expressed by circulating monocytes or tumor associated macrophages, to reduce the number of tumor associated macrophages near or within the cancer mass or reduce migration of tumor associated macrophages. Some embodiments of the invention relate to methods for treating, preventing, or reducing the incidence of cancer metastasis, which include administering to a subject in need thereof an effective amount of an inhibitor of

MOSPD2, *e.g.*, MOSPD2 expressed by circulating monocytes or tumor associated macrophages, to reduce the number of tumor associated macrophages near or within the cancer mass or reduce migration of tumor associated macrophages. In some embodiments, the administration is local administration, *e.g.*, intra-tumor administration. In some embodiments, the administration is effective in reducing the number or migration of tumor associated macrophages by at least about 5%, at least about 10%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or 100%, or any number in between the aforementioned percentages, when compared to baseline.

[0255] In some embodiments, the administration is effective in reducing the number or migration of tumor associated macrophages by about 5%, about 10%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, or 100%, or any number in between the aforementioned percentages, when compared to baseline.

[0256] In other embodiments, the administration is effective in reducing the number or migration of tumor associated macrophages, when compared to baseline, by from about 5% to 100%, or from about 5% to about 95%, from about 5% to about 90%, from about 5% to about 80%, from about 5% to about 70%, from about 5% to about 60%, from about 5% to about 50%, from about 5% to about 40%, from about 5% to about 30%, from about 10% to 100%, from about 10% to about 95%, from about 10% to about 90%, from about 10% to about 80%, from about 10% to about 70%, from about 10% to about 60%, from about 10% to about 50%, from about 10% to about 40%, from about 20% to 100%, from about 20% to about 95%, from about 20% to about 90%, from about 20% to about 80%, from about 20% to about 70%, from about 20% to about 60%, from about 20% to about 50%, from about 20% to about 40%, or any other range of values described herein.

[0257] Any assay known in the art can be used to measure tumor associated macrophage density or numbers such as immunohistochemical staining of tumor sections using antibodies that specifically detect macrophages. *See e.g.*, U.S. Patent Publication Nos. 2007/0218116 and 2011/0311616. Exemplary MOSPD2 inhibitors include those

described herein. Suitable types of cancer are also described herein. In some embodiments, the cancer is a breast cancer, a head and neck cancer, a prostate cancer or a uterine cancer.

[0258] In some embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody described herein, *e.g.*, having an antibody-antigen equilibrium dissociation constant (K_D) of from about 10^{-6} M to about 10^{-12} M, or any range of values thereof (*e.g.*, from about 10^{-7} M to about 10^{-12} , from 10^{-8} M to about 10^{-12} M, from about 10^{-9} M to about 10^{-12} M, from about 10^{-10} M to about 10^{-12} M, from about 10^{-11} M to about 10^{-12} M, from about 10^{-6} M to about 10^{-11} M, from about 10^{-7} M to about 10^{-11} M, from about 10^{-8} M to about 10^{-11} M, from about 10^{-9} M to about 10^{-11} M, from about 10^{-10} M to about 10^{-11} M, from about 10^{-6} M to about 10^{-10} M, from about 10^{-7} M to about 10^{-10} M, from about 10^{-8} M to about 10^{-10} M, from about 10^{-9} M to about 10^{-10} M, from about 10^{-6} M to about 10^{-9} M, from about 10^{-7} M to about 10^{-9} M, from about 10^{-8} M to about 10^{-9} M, from about 10^{-6} M to about 10^{-8} M, or from about 10^{-7} M to about 10^{-8}). In other embodiments, the antibody or antigen binding fragment thereof has a K_D of about 10^{-6} M, about 10^{-7} M, about 10^{-8} M, about 10^{-9} M, about 10^{-10} M, about 10^{-11} M, or about 10^{-12} M. In some embodiments, the antibody or antigen binding fragment binds to one or more epitopes on MOSPD2. In some embodiments, the K_D is determined by Scatchard analysis, surface plasmon resonance, or other method described herein, in some embodiments, at 37°C.

[0259] In some embodiments, antibodies or antigen binding fragments of the invention bind to MOSPD2 with a K_{on} of from about 10^3 1/Ms to about 10^6 1/Ms, or any range of values thereof (*e.g.*, from about 10^3 1/Ms to about 10^5 1/Ms, from about 10^4 1/Ms to about 10^5 1/Ms, from about 10^4 1/Ms to about 10^6 1/Ms, from about 10^5 1/Ms to about 10^6 1/Ms, or from about 10^3 1/Ms to about 10^4 1/Ms). In other embodiments, the antibody or antigen binding fragment has a K_{on} of about 10^3 1/Ms, about 10^4 1/Ms, about 10^5 1/Ms, or about 10^6 1/Ms.

[0260] In some embodiments, antibodies or antigen binding fragments of the invention bind to MOSPD2 with a K_{off} of from about 10^{-3} 1/s to about 10^{-6} 1/s, or any range of values thereof (*e.g.*, from about 10^{-3} 1/s to about 10^{-5} 1/s, from about 10^{-4} 1/s to about 10^{-5} 1/s, from about 10^{-4} 1/s to about 10^{-6} 1/s, from about 10^{-5} 1/s to about 10^{-6} 1/s, or from

about 10^{-3} 1/s to about 10^{-4} 1/s). In other embodiments, the antibody or antigen binding fragment has a K_{off} of about 10^{-3} 1/s, about 10^{-4} 1/s, about 10^{-5} 1/s, or about 10^{-6} 1/s.

[0261] In some embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is from about 1 $\mu\text{g}/\text{ml}$ to about 10 $\mu\text{g}/\text{ml}$, or any range of values thereof (e.g., from about 2 $\mu\text{g}/\text{ml}$ to about 10 $\mu\text{g}/\text{ml}$, from about 3 $\mu\text{g}/\text{ml}$ to about 10 $\mu\text{g}/\text{ml}$, from about 4 $\mu\text{g}/\text{ml}$ to about 10 $\mu\text{g}/\text{ml}$, from about 5 $\mu\text{g}/\text{ml}$ to about 10 $\mu\text{g}/\text{ml}$, from about 6 $\mu\text{g}/\text{ml}$ to about 10 $\mu\text{g}/\text{ml}$, from about 7 $\mu\text{g}/\text{ml}$ to about 10 $\mu\text{g}/\text{ml}$, from about 8 $\mu\text{g}/\text{ml}$ to about 10 $\mu\text{g}/\text{ml}$, from about 9 $\mu\text{g}/\text{ml}$ to about 10 $\mu\text{g}/\text{ml}$, from about 1 $\mu\text{g}/\text{ml}$ to about 9 $\mu\text{g}/\text{ml}$, from about 2 $\mu\text{g}/\text{ml}$ to about 9 $\mu\text{g}/\text{ml}$, from about 3 $\mu\text{g}/\text{ml}$ to about 9 $\mu\text{g}/\text{ml}$, from about 4 $\mu\text{g}/\text{ml}$ to about 9 $\mu\text{g}/\text{ml}$, from about 5 $\mu\text{g}/\text{ml}$ to about 9 $\mu\text{g}/\text{ml}$, from about 6 $\mu\text{g}/\text{ml}$ to about 9 $\mu\text{g}/\text{ml}$, from about 7 $\mu\text{g}/\text{ml}$ to about 9 $\mu\text{g}/\text{ml}$, from about 8 $\mu\text{g}/\text{ml}$ to about 9 $\mu\text{g}/\text{ml}$, from about 1 $\mu\text{g}/\text{ml}$ to about 8 $\mu\text{g}/\text{ml}$, from about 2 $\mu\text{g}/\text{ml}$ to about 8 $\mu\text{g}/\text{ml}$, from about 3 $\mu\text{g}/\text{ml}$ to about 8 $\mu\text{g}/\text{ml}$, from about 4 $\mu\text{g}/\text{ml}$ to about 8 $\mu\text{g}/\text{ml}$, from about 5 $\mu\text{g}/\text{ml}$ to about 8 $\mu\text{g}/\text{ml}$, from about 6 $\mu\text{g}/\text{ml}$ to about 8 $\mu\text{g}/\text{ml}$, from about 7 $\mu\text{g}/\text{ml}$ to about 8 $\mu\text{g}/\text{ml}$, from about 1 $\mu\text{g}/\text{ml}$ to about 7 $\mu\text{g}/\text{ml}$, from about 2 $\mu\text{g}/\text{ml}$ to about 7 $\mu\text{g}/\text{ml}$, from about 3 $\mu\text{g}/\text{ml}$ to about 7 $\mu\text{g}/\text{ml}$, from about 4 $\mu\text{g}/\text{ml}$ to about 7 $\mu\text{g}/\text{ml}$, from about 5 $\mu\text{g}/\text{ml}$ to about 7 $\mu\text{g}/\text{ml}$, from about 6 $\mu\text{g}/\text{ml}$ to about 7 $\mu\text{g}/\text{ml}$, from about 7 $\mu\text{g}/\text{ml}$ to about 7 $\mu\text{g}/\text{ml}$, from about 1 $\mu\text{g}/\text{ml}$ to about 6 $\mu\text{g}/\text{ml}$, from about 2 $\mu\text{g}/\text{ml}$ to about 6 $\mu\text{g}/\text{ml}$, from about 3 $\mu\text{g}/\text{ml}$ to about 6 $\mu\text{g}/\text{ml}$, from about 4 $\mu\text{g}/\text{ml}$ to about 6 $\mu\text{g}/\text{ml}$, from about 5 $\mu\text{g}/\text{ml}$ to about 6 $\mu\text{g}/\text{ml}$, from about 1 $\mu\text{g}/\text{ml}$ to about 5 $\mu\text{g}/\text{ml}$, from about 2 $\mu\text{g}/\text{ml}$ to about 5 $\mu\text{g}/\text{ml}$, from about 3 $\mu\text{g}/\text{ml}$ to about 5 $\mu\text{g}/\text{ml}$, from about 4 $\mu\text{g}/\text{ml}$ to about 5 $\mu\text{g}/\text{ml}$, from about 1 $\mu\text{g}/\text{ml}$ to about 4 $\mu\text{g}/\text{ml}$, from about 2 $\mu\text{g}/\text{ml}$ to about 4 $\mu\text{g}/\text{ml}$, from about 3 $\mu\text{g}/\text{ml}$ to about 4 $\mu\text{g}/\text{ml}$, from about 1 $\mu\text{g}/\text{ml}$ to about 3 $\mu\text{g}/\text{ml}$, from about 2 $\mu\text{g}/\text{ml}$ to about 3 $\mu\text{g}/\text{ml}$, to about 3 $\mu\text{g}/\text{ml}$, or from about 1 $\mu\text{g}/\text{ml}$ to about 2 $\mu\text{g}/\text{ml}$). In other embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is about 1 $\mu\text{g}/\text{ml}$, about 2 $\mu\text{g}/\text{ml}$, about 3 $\mu\text{g}/\text{ml}$, about 4 $\mu\text{g}/\text{ml}$, about 5 $\mu\text{g}/\text{ml}$, about 6 $\mu\text{g}/\text{ml}$, about 7 $\mu\text{g}/\text{ml}$, about 8 $\mu\text{g}/\text{ml}$, about 9 $\mu\text{g}/\text{ml}$, or about 10 $\mu\text{g}/\text{ml}$.

[0262] In some embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is from about 10 mg/kg to about 40 mg/kg, or any range of values thereof (e.g., from about 15 mg/kg to about 40 mg/kg, from about 20 mg/kg to about 40 mg/kg, from about 25 mg/kg to about 40 mg/kg,

from about 30 mg/kg to about 40 mg/kg, from about 35 mg/kg to about 40 mg/kg, from about 10 mg/kg to about 35 mg/kg, from about 15 mg/kg to about 35 mg/kg, from about 20 mg/kg to about 35 mg/kg, from about 25 mg/kg to about 35 mg/kg, from about 30 mg/kg to about 35 mg/kg, from about 10 mg/kg to about 30 mg/kg, from about 15 mg/kg to about 30 mg/kg, from about 20 mg/kg to about 30 mg/kg, from about 25 mg/kg to about 30 mg/kg, from about 10 mg/kg to about 25 mg/kg, from about 15 mg/kg to about 25 mg/kg, from about 20 mg/kg to about 25 mg/kg, from about 10 mg/kg to about 20 mg/kg, from about 15 mg/kg to about 20 mg/kg, or from about 10 mg/kg to about 15 mg/kg). In other embodiments, the inhibitor of MOSPD2 is an antibody or antigen binding fragment of an antibody, and the therapeutically effective amount is about 10 mg/kg, about 15 mg/kg, about 20 mg/kg, about 25 mg/kg, about 30 mg/kg, about 35 mg/kg, or about 40 mg/kg.

[0263] In some embodiments, the subject is a mammal or a human. In other embodiments the MOSPD2 is a mammalian MOSPD2 or a human MOSPD2.

Diagnostic Methods

[0264] The inventors have discovered that MOSPD2 is expressed on the surface of different types of cancer cells and tumors, and on inflammatory cells that have infiltrated into inflamed tissues or that are associated with tumors. The inventors have also discovered that MOSPD2 expression is increased in correlation with tumor grade in various types of tumors. Therefore, in one aspect, the invention relates to a method for the prediction, diagnosis, or prognosis of cancer or cancer metastasis in a subject (e.g., breast cancer, colon cancer, liver cancer, melanoma, or other type of cancer described herein), which comprises determining the expression level of MOSPD2 in a sample of the subject. In another aspect, the invention relates to a method for the prediction, diagnosis, or prognosis of tumor progression or invasiveness in a subject, which comprises determining the expression level of MOSPD2 in a sample of the subject. In one embodiment of these methods, the expression level of MOSPD2 is the level of MOSPD2 gene expression. In another embodiment, the expression level of MOSPD2 is the level of MOSPD2 protein expression.

[0265] In one aspect, the invention relates to an *in vitro* method for the prediction, diagnosis or prognosis of cancer in a subject (e.g., breast cancer, colon cancer, liver cancer, melanoma, or other type of cancer described herein), which comprises

determining or quantifying the expression level of MOSPD2 in a sample of the subject. In another aspect, the invention relates to an *in vitro* method for the prediction, diagnosis or prognosis of cancer in a subject (e.g., breast cancer, colon cancer, liver cancer, melanoma, or other type of cancer described herein), which comprises (i) determining or quantifying the expression level of MOSPD2 in a sample of the subject, and (ii) comparing the expression level obtained in step (i) with a control or reference value, wherein an increased expression level of MOSPD2 with respect to the control or reference value is indicative of cancer or an increased risk of developing cancer. In some embodiments, if MOSPD2 expression is present in the sample of the subject, then the subject has cancer or an increased risk of cancer. In other embodiments, if MOSPD2 expression is present in the sample of the subject in an amount greater than MOSPD2 expression of the control or reference value, then the subject has cancer or an increased risk of cancer.

[0266] In one aspect, the invention relates to an *in vitro* method for the prediction, diagnosis or prognosis of cancer metastasis in a subject (e.g., breast cancer, colon cancer, liver cancer, melanoma, or other type of cancer described herein), which comprises determining or quantifying the expression level of MOSPD2 in a sample of the subject. In another aspect, the invention relates to an *in vitro* method for the prediction, diagnosis or prognosis of cancer metastasis in a subject (e.g., breast cancer, colon cancer, liver cancer, melanoma, or other type of cancer described herein), which comprises (i) determining or quantifying the expression level of MOSPD2 in a sample of the subject, and (ii) comparing the expression level obtained in step (i) with a control or reference value, wherein an increased expression level of MOSPD2 with respect to the control or reference value is indicative of cancer metastasis or an increased risk of cancer metastasis. In some embodiments, if MOSPD2 expression is present in the sample of the subject, then the subject has cancer metastasis or an increased risk of cancer metastasis. In other embodiments, if MOSPD2 expression is present in the sample of the subject in an amount greater than MOSPD2 expression of the control or reference value, then the subject has cancer metastasis or an increased risk of cancer metastasis.

[0267] In one aspect, the invention relates to an *in vitro* method for the prediction, diagnosis or prognosis of tumor progression (e.g., increased tumor grade) or invasiveness in a subject, which comprises determining or quantifying the expression level of

MOSPD2 in a sample of the subject. In another aspect, the invention relates to an *in vitro* method for the prediction, diagnosis or prognosis of tumor progression (*e.g.*, increased tumor grade) or invasiveness in a subject, which comprises (i) determining or quantifying the expression level of MOSPD2 in a sample of the subject, and (ii) comparing the expression level obtained in step (i) with a control or reference value, wherein increased expression level of MOSPD2 with respect to the control or reference value is indicative of tumor progression (*e.g.*, increased tumor grade) or invasiveness or an increased risk of tumor progression or invasiveness. In some embodiments, if MOSPD2 expression is present in the sample of the subject, then the subject has tumor progression or invasiveness or an increased risk of tumor progression or invasiveness. In other embodiments, if MOSPD2 expression is present in the sample of the subject in an amount greater than MOSPD2 expression of the control or reference value, then the subject has tumor progression or tumor invasiveness or an increased risk of tumor progression or invasiveness.

[0268] In some embodiments, the methods of the invention comprise one or more of the following additional steps: instructing a laboratory to quantify the expression level of MOSPD2 in the sample; obtaining a report of the expression level of MOSPD2 in the sample from the laboratory; and/or administering a therapeutically effective amount of an inhibitor of MOSPD2 (*e.g.*, an anti-MOSPD2 antibody or antigen binding fragment thereof) to the subject.

[0269] In some embodiments, the sample is a tissue biopsy, tumor biopsy, or blood sample from a subject.

[0270] In some embodiments, the control or reference value is the expression level of MOSPD2 in normal tissue (*e.g.*, normal adjacent tissue (NAT)). In other embodiments, the control or reference value is no detectable MOSDP2 expression or no significant MOSPD2 expression.

[0271] Methods for determining the expression level of MOSPD2 are known in the literature and described herein.

[0272] In some embodiments, the invention relates to a method for treating a cancer or cancer metastasis responsive to an inhibitor of MOSPD2 in a subject, comprising (i) determining the expression level of MOSPD2 in the subject, and when the expression level is determined to be greater than that of a control or reference value, (ii)

administering, to the subject, a therapeutically effective amount of an inhibitor of MOSPD2 (e.g., an anti-MOSPD2 antibody or antigen binding fragment thereof).

[0273] In some embodiments, the invention relates to a method for treating a MOSPD2 expressing tumor in a subject, which comprises administering a therapeutically effective amount of an inhibitor of MOSPD2 (e.g., an anti-MOSPD2 antibody or antigen binding fragment thereof). In other embodiments, the invention relates to a method of treating a subject, which comprises administering a therapeutically effective amount of an inhibitor of MOSPD2 (e.g., an anti-MOSPD2 antibody or antigen binding fragment thereof), wherein the subject has a tumor expressing MOSPD2.

[0274] In some embodiments, the invention relates to an inhibitor of MOSPD2 (e.g., an anti-MOSPD2 antibody or antigen binding fragment thereof), for use in treating cancer or cancer metastasis in a patient having a cancer cell or tumor that expresses MOSPD2.

EXAMPLES

[0275] Reference is now made to the following examples, which together with the above descriptions illustrate some embodiments of the invention in a non-limiting fashion.

Materials and Methods

MOSPD2 Silencing

[0276] The human breast cancer cell line MDA-MB-231 (hereafter MDA-231) (HTB-26) and the human malignant melanoma cell line A2058 (CRL-11147) were purchased from the American Type Culture Collection (ATCC). The cells (2×10^6 in 2ml) were placed in a 15ml tube. Lenti-virus particles expressing control short hairpin RNA (sh-RNA) (2×10^5 viral particles) or human MOSPD2 sh-RNA (2×10^6 viral particles) were applied to the cells, which were then spun for 60 min, 2000 rpm at room temperature in the presence of 8 μ g/ml polybrene (Sigma, Israel). The cells were then seeded in a 6 well plate. After 72 hour, fresh medium containing puromycin (4 μ g/ml Sigma, Israel) was added for the selection of transduced cells. For CRISPR-CAS9 mediated silencing, MDA-231 cells were transduced with CRISPR-CAS9 non-target control or CRISPR-CAS9 human MOSPD2 lenti-viral particles as described above. Single cell cloning was performed on

transduced cells to isolate cells with silenced MOSPD2 protein expression and impaired migration.

Western Blotting

[0277] sh-control or sh-MOSPD2 Lenti-virus transduced A2058 or MDA-231 cells, or control or MOSPD2 CRISPR-CAS9 lenti-viral particles transduced MDA-231 cells (10^6), were washed and resuspended in lysis buffer containing 1:100 dithiothreitol (DTT), phosphatase and protease inhibitors (Thermo Scientific). Samples were loaded onto a precast Criterion TGX gel (Bio-Rad, Hemel Hempstead, UK) and transferred onto a nitrocellulose membrane. Blots were blocked with 5% milk or bovine serum albumin (BSA) in Tris buffered saline and Tween 20 (TBST) for 1 hour, followed by incubation with primary and secondary antibodies. Membranes were developed using an ECL kit (Thermo Scientific). The following antibodies were used for immunoblotting:

[0278] Primary antibodies: Rabbit anti-MOSPD2 (1:5000) generated by Vascular Biogenics Ltd. Phospho extracellular-regulated kinase (p-ERK1/2) (Thr 183 and Tyr 185, 1:4000) was purchased from Sigma (Israel). Phospho-AKT (Ser 473, 1:1000) was purchased from Cell Signaling. Phospho-FAK (1:2000) was purchased from Abcam (Cambridge, UK). Heat shock protein (HSP) 90 (1:1000) was purchased from Santa Cruz Biotechnology (Dallas, TX).

[0279] Secondary antibodies: Horseradish peroxidase (HRP) donkey anti-rabbit (1:5000) and HRP goat anti-mouse (1:5000) antibodies were purchased from Jackson ImmunoResearch (West Grove, PA).

Q-PCR

[0280] To determine silencing efficacy, RNA was extracted from sh-control and sh-MOSPD2 Lenti-virus transduced MDA-231 cells using RNeasy mini kit (Qiagen, ValenVBA, CA). For cDNA preparation, 2 μ g of RNA was combined with qScript reaction mix and qScript reverse transcriptase (Quanta Bioscience, Gaithersburg, MD). The reaction was placed in a thermal cycler (BioRad, Hercules, CA) and a run program was set according to the manufacturer instructions. Real-time PCR reactions were performed on an Applied Biosystems 7300 real time PCR system (Grand Island, NY) using sets of primers for human MOSPD2, 28S to normalize RNA levels (BIOSEARCH

TECHNOLOGIES, Petaluma, CA) and SYBR Green PCR Master Mix (Applied Biosystems, Warrington, UK).

Immunohistochemistry staining

[0281] To assess the expression level of MOSPD2 in cancer tissues, Biomax arrays (US Biomax Rockville, MD) for breast cancer (T088B and BR2028a), for liver cancer (BC03116a), and for multiple organ tumor (MC6163) were stained with the rabbit anti-MOSPD2 antibody or control rabbit IgG (R&D Systems Cat# AB-105-C) followed by incubation with anti-Rabbit HRP (Cat #0399 DAKO, Denmark).

EXAMPLE 1

Anti-MOSPD2 Antibodies

[0282] Anti-MOSPD2 polyclonal antibodies were generated according to the following methods.

Materials and Methods

Production and purification of hemagglutinin (HA)-tagged recombinant human MOSPD2 (HA-rhMOSPD2)

[0283] Full length human MOSPD2 cDNA was inserted, using EcoRI and XbaI restriction sites, into the lentivirus plasmid vector pLVX-EF1 α -IRES-Puro (Clonetech, CA). Oligonucleotide encoding the HA-tag (YPYDVPDYA; SEQ ID NO:15) was inserted into the N-terminal region of MOSPD2 with EcoRI restriction sites. For transduction, A2058 melanoma cells (ATCC CRL-11147, VA) were spun for 60 minutes at 2000 rpm at room temperature in the presence of 8 μ g/ml polybrene (Sigma, Israel) and lentiviral particles containing HA-rhMOSPD2 expressing vector. The cells were then seeded in a 6 well plate. After 72 hours, fresh medium containing puromycin (4 μ g/ml Sigma, Israel) was added for the selection of transduced cells. To purify HA-rhMOSPD2, A2058 transduced cells were lysed with M-PER mammalian protein extraction reagent (Thermo Scientific) and passed through anti-HA agarose beads (Thermo Scientific). Glycine or sodium thiocyanate was used for the elution of HA-rhMOSPD2 from the beads, followed by thorough dialysis against PBS.

Generation and isolation of α -MOSPD2 polyclonal antibodies

[0284] Rabbits were immunized with approximately 0.5 mg of HA-rhMOSPD2 emulsified in complete freunds adjuvant followed by three boosts every three weeks with approximately 0.25 mg of HA-rhMOSPD2 emulsified in incomplete freunds adjuvant. Serum was collected one week after each boost to assess for antibody immunogenicity and titers. α -MOSPD2 antibodies were isolated from serum using protein A/G beads (SantaCruz, CA).

Results

Rabbit polyclonal α -MOSPD2 antibodies detect and precipitate endogenous human MOSPD2

[0285] Isolated α -MOSPD2 polyclonal antibodies were evaluated for their ability to detect and precipitate endogenous MOSPD2. Cell lysate was prepared from U937 cells transduced with control or sh-MOSPD2 Lenti-virus particles. Samples were analyzed by Western blot using the isolated α -MOSPD2 antibodies (diluted 1:5000). Expression of HSP90 was also determined as a loading control. Immunoprecipitation of U397 cell lysate was also performed using the isolated α -MOSPD2 antibodies or rabbit IgG (10 μ g) as a control. The resulting precipitates were analyzed by immunoblot with the isolated α -MOSPD2 antibodies, followed by incubation with goat anti-rabbit antibody-HRP (1:5000). Results show that the isolated α -MOSPD2 antibodies readily detect and immunoprecipitate endogenously expressed MOSPD2 in U937 cells.

EXAMPLE 2

MOSPD2 and Migration of Metastatic Cell Lines

[0286] In order to assess the role of MOSPD2 in cancer cell migration, MOSPD2 expression in two metastatic cell lines, A2058 melanoma and MDA-231 breast cancer, was silenced using sh-control or sh-MOSPD2 lenti-virus particles.

[0287] In particular, sh-control or sh-MOSPD2 transduced A2058 or MDA-231 cells (3×10^5) previously starved for 3 hours in 0.5% FBS/RPMI-1640 were seeded in the upper chamber of a QCM 24-well, 5 μ m pore, migration assay plate (Corning-Costar, Corning, NY), followed by incubation for 24 hours in the presence of 10% FBS/RPMI-1640 and EGF (200ng/ml, Peprotech Israel) in the lower chamber. Subsequently, the cells which

migrated to the lower compartment were stained with crystal violet before images were taken.

[0288] FIG. 1 demonstrates that sh-MOSPD2 lenti-virus particles have profoundly decreased protein expression and inhibited cancer cell migration *in vitro*.

EXAMPLE 3

MOSPD2 and Cell Proliferation

[0289] To determine whether the inhibitory effect on cell migration subsequent to MOSPD2 silencing is secondary to fundamental cell function such as proliferation, sh-control or sh-MOSPD2 lenti-virus particle transduced MDA-231 breast cancer cells were tested for proliferation over a period of 3 days.

[0290] Specifically, sh-control or sh-MOSPD2 lenti-virus transduced MDA-231 cells were seeded in 6 well plates (10^4 per well). The cells were counted by FACS every 24 hours in triplicates for 3 consecutive days.

[0291] The data shown in FIG. 2 indicate that MOSPD2 is not essential for the proliferation of these cells, suggesting a regulatory role for MODPD2 specifically in migration.

EXAMPLE 4

MOSPD2 and Cell Metastasis

[0292] To assess the role of MOSPD2 in disseminating cancer cells to organs beyond the original site of cancer, the extent of lung metastasis in sh-control or sh-MOSPD2 lenti-virus particle-transduced MDA-231 breast cancer cells were adoptively transferred into immune-deficient mice. In another model in which the site of inception occurs in the breast, immunodeficient mice were inoculated with sh-control or sh-MOSPD2 lenti-virus particle-transduced MDA-231 breast cancer cells in the mammary fat pad.

[0293] *Pathological examination:* Histology slides were stained with hematoxylin/eosin (H&E). Formalin-fixed tissue was dehydrated, embedded in paraffin, and sectioned at 4 μ m thickness. The H&E staining was calibrated on a Leica staining module. The slides were warmed to 90 °C for 7 minutes and then processed according to a fully automated protocol. After sections were dewaxed and rehydrated, slides were stained for 7 minutes

in Gill's Hematoxylin No. 3 (Surgipath), washed, dipped in acidic alcohol, and washed. After short dipping in 70% ethanol and 96% ethanol, slides were stained for 4 minutes in eosin (Sigma), and dehydrated in 96% ethanol and then twice in 100% ethanol for 1 minute each time. After a run on an automated stainer was completed, sections were cleared in xylene for 10 seconds and mounted with Entellan. Mean tumor area comprises the maximal lung tumor area measured for each mouse.

[0294] *Systemic*: 10^6 sh-control or sh-MOSPD2 lenti-virus transduced MDA-231 cells were injected into the tail vein of 8 weeks old female SCID mice (C.B-17/IcrHsd-Prkdc^{scid}, Harlan Israel). Mice were sacrificed after 4 weeks. Lungs were excised for histopathologic examination. The results in FIG. 3A show that silencing MOSPD2 expression significantly ($p=0.023$) inhibits the presence of metastatic breast cancer cells in the lungs by more than 50% (metastasis area).

[0295] *Orthotopic*: 5×10^6 sh-control or sh-MOSPD2 lenti-virus transduced MDA-231 cells were injected into the mammary fat pad of 8 weeks old female SCID mice (C.B-17/IcrHsd-Prkdc^{scid}, Harlan Israel). Mice were sacrificed after 10 weeks. Ipsilateral inguinal lymph node and the lungs were excised for examination. Macroscopic examination showed that the vast majority of lymph nodes excised from mice transferred with sh-control cells were overwhelmingly bigger than those from mice transferred with sh-MOSPD2 treated cells (FIG. 3B). Moreover, the mean metastasis area measured in the lungs of mice transferred with sh-MOSPD2 treated cells was reduced by more than 50% compared to the control group (FIG. 3C).

[0296] The ratio of MOSPD2 mRNA silencing in sh-MOSPD2 injected cells was ~80%, as determined by Q-PCR as described in the Materials and Methods.

[0297] These results demonstrate that MOSPD2 plays a major role in breast cancer metastasis.

EXAMPLE 5

MOSPD2 Expression in Various Types of Cancer

[0298] To determine whether MOSPD2 expression is associated with the transformation of cells from normal to cancerous, slides carrying normal and cancerous tissues were screened using anti-MOSPD2 antibody as described in the Materials and Methods section.

[0299] FIG. 4A shows representative staining of normal and cancerous breast tissue. While normal and cancerous breast tissues were negatively stained with control IgG antibody, anti-MOSPD2 antibody distinctively stained cancerous tissues only. Similarly, MOSPD2 is not expressed in normal bladder, brain, colon, esophagus, tongue, kidney and hepatic tissues, but is upregulated when these tissues turn cancerous (FIGs. 4B-4E). These results suggest that in various tissues, MOSPD2 expression is associated with transformation of normal tissue to cancerous tissue.

EXAMPLE 6

MOSPD2 Gene Knockdown and Cancer Cell Migration

[0300] *In vitro*: To achieve sustainable knockdown of MOSPD2, MDA-231 breast cancer cells were transduced with lenti-viral particles that contain the CRISPR-CAS9 gene editing system as described in the Materials and Methods section. Control or MOSPD2 CRISPR-CAS9 lenti-viral particles transduced MDA-231 cells were tested for migration similar to the method described in Example 2. Control or MOSPD2 CRISPR-CAS9 lenti-viral particles transduced MDA-231 cells (3×10^5) were seeded in the upper chamber, followed by incubation for 2-4 hours. Subsequently, the number of cells which migrated to the lower compartment was determined by FACS.

[0301] FIGs. 5A and 5B show that introducing the CRISPR-CAS9 system for MOSPD2 in MDA-231 cancer cells abolished protein expression and consequently profoundly inhibited migration of the cells in a trans-well assay.

[0302] To test the effects of MOSPD2 silencing by CRISPR-CAS9 on chemokine receptor-driven signaling events, phosphorylation levels of ERK, AKT and FAK were studied as described in the Materials and Methods. In accordance with the migration assay results, silencing MOSPD2 by the CRISPR-CAS9 system compared to control completely prevented phosphorylation of AKT and distinctly inhibited phosphorylation of ERK and FAK (see Western Blots in FIG. 5C) in cells exposed to EGF.

[0303] *In vivo*: 10^6 CRISPR-control or CRISPR-MOSPD2 lenti-virus transduced MDA-231 cells were injected into the tail vein of 8 weeks old female SCID mice (C.B-17/IcrHsd-Prkdcscid, Harlan Israel). Mice were then sacrificed after 3 weeks. Lungs were excised for histopathologic examination similar to the method described in Example 4. FIG. 5D shows that silencing MOSPD2 by the CRISPR-CAS9 system significantly

inhibited the presence of metastatic breast cancer cells in the lungs by more than 95% (metastasis area).

EXAMPLE 7

VB-201 and EGF Signaling Pathway

[0304] *In vitro*: To test the effect of VB-201 on epidermal growth factor (EGF)- induced phosphorylation, MDA-231 breast cancer cells (10^6) were starved for 3 hours in 0.5% FCS medium, followed by incubation with various concentrations of VB-201 (1 μ g/ml, 5 μ g/ml, and 10 μ g/ml) or a solvent control for 20 minutes. The MDA-231 cells were then activated with EGF (200 ng/ml) for 10 min. Phosphorylation of AKT was then analyzed by Western Blot. HSP90 was used for a loading control.

[0305] As shown in FIG. 6, VB-201 at 10 μ g/ml nearly completely inhibits EGF induced phosphorylation of AKT, with significant inhibition observed at 5 μ g/ml.

EXAMPLE 8

Generation of Anti-MOSPD2 (Fab')₂ Monoclonal Antibodies

[0306] Anti-MOSPD2 (Fab')₂ monoclonal antibodies (mAb) were obtained using the HuCAL PLATINUM® Platform (Bio-Rad AbD Serotec, GmnH) which contains a selection of phage displayed human Fab.

[0307] Briefly, recombinant protein of the extracellular region of MOSPD2 fused to human Fc was immobilized on a solid support. The HuCAL® library presented on phage particles was incubated with the immobilized antigen. Nonspecific antibodies were removed by extensive washing and specific antibody phages were eluted by adding a reducing agent. Antibody DNA was isolated as a pool and subcloned into an *E. coli* expression vector to generate bivalent F(ab')₂ mAb. Colonies were picked and grown in a microtiter plate. The cultures were lysed to release the antibody molecules and screened for specific antigen binding by ELISA and FACS. Unique antibodies were expressed and purified using one-step affinity chromatography, and then tested again by ELISA and FACS for specificity.

[0308] FIG. 7 lists 17 anti-MOSPD2 F(ab')₂ monoclonal antibody clones that were identified following a primary screen for binding to cells over-expressing MOSPD2.

Further analysis of the clones for MOSPD2 binding with ELISA identified 12 clones having O.D. values greater than 5 times over background (* in FIG. 7).

EXAMPLE 9

Anti-MOSPD2 F(ab')2 mAb Bind Human MOSPD2 Overexpressed on Cells

[0309] A2058 melanoma cells were transfected with HA-tagged human MOSPD2 to generate cells overexpressing MOSPD2.

[0310] Binding of the 12 antibody clones identified in Example 8 to MOSPD2 was then tested using flow cytometry with these cells. Specifically, 10^5 cells were incubated with 2.5 μ g of F(ab')₂ mAb at 4°C for 1hr in 100 μ l of FACS buffer (PBS + 2% FCS + 0.02% sodium azide). Cells were then washed, resuspended in FACS buffer and stained for 30min at 4°C with Alexa-Fluor 647-conjugated (Fab')₂ goat anti-human IgG, F(ab')₂ 1:200 (Cat# 109-606-097, Jackson Immunoresearch, PA). Cells were washed, resuspended in FACS buffer and analyzed on a FACS-Calibur device.

[0311] All clones positively stained the cells. Representative staining for 2 clones is shown in FIGs. 8A-8B. A clone that was not identified as a positive clone in Example 8 with ELISA was used as a negative control.

EXAMPLE 10

Anti-MOSPD2 F(ab')2 mAb Specifically Binds Endogenous MOSPD2 on Human Breast Cancer Cells

[0312] Anti-MOSPD2 F(ab')₂ mAb was tested for binding to surface expressed endogenous MOSPD2 on MDA-231 breast cancer cells. Cells were stained with anti-MOSPD2 F(ab')₂ mAb as described in Example 9. Staining with 2 different clones is shown in FIG. 9. The ELISA negative clone described in Example 9 was used as negative control. FIG. 9 shows that anti-MOSPD2 F(ab')₂ mAb specifically binds endogenous MOSPD2 on human breast cancer cells.

[0313] To further demonstrate antigen binding specificity, MOSPD2 gene expression was silenced in MDA-231 cells using CRISP-CAS9 lentiviral particles. MOSPD2-silenced cells and non-silenced cells were combined with anti-MOSPD2 (Fab')₂ mAb or a negative control and analyzed with FACS. FIGs. 10A-10B show anti-MOSPD2 F(ab')₂ mAb binds

to MDA-231 cells (FIG. 10A), but does not bind to MOSPD2-silenced MDA-231 cells (FIG. 10B).

EXAMPLE 11

Anti-MOSPD2 F(ab')2 mAb Binds Endogenous MOSPD2 on Melanoma and Liver Cancer Cells

[0314] Anti-MOSPD2 F(ab')₂ mAb was tested for binding to surface expressed endogenous MOSPD2 on A2058 melanoma and HepG2 liver cancer cell lines. Cells were stained with anti-MOSPD2 F(ab')₂ mAb and tested for binding to MOSPD2 as described in Examples 9 and 10. FIGs. 11A-11B show that anti-MOSPD2 F(ab')₂ mAb specifically binds endogenous MOSPD2 on melanoma and liver cancer cells.

EXAMPLE 12

Anti-MOSPD2 F(ab')2 mAb Inhibits EGF-induced Signaling in MDA-231 Cancer Cells

[0315] The effect of anti-MOSPD2 F(ab')₂ mAb on EGF-induced signaling in MDA-231 cancer cells was analyzed with Western blot. Specifically, MDA-231 cells were starved overnight with medium containing 0.5% FCS and then incubated for 1hr with anti-MOSPD2 (Fab')₂ mAb before adding EGF (100ng/ml) for 5min. Cells were washed and resuspended in lysis buffer, loaded onto a precast Criterion TGX gel (Bio-Rad, Hemel Hempstead, UK) and transferred onto nitrocellulose membrane. The membranes were blocked with 5% milk or BSA in Tris buffered saline and Tween 20 (TBST) for 1hr, and then incubated with primary and secondary antibodies. Membranes were developed using an ECL kit (Thermo Scientific). Cells that were not treated with anti-MOSPD2 F(ab')₂ mAb (unt) were analyzed as a negative control. Heat shock protein (HSP)-90 protein levels were also analyzed as a protein loading control.

[0316] The following antibodies were used:

[0317] Primary antibodies: p-ERK1/2 (cat. no. M8159; 1:10,000) from Sigma (Israel); phospho-AKT (cat no. 9271; Ser 473, 1:1000) and phospho-EGF Receptor (cat no. 2236 1:1000) from Cell Signaling; and HSP-90 (cat. no. 13119; 1:500) from Santa Cruz Biotechnology (Santa Cruz, CA).

[0318] Secondary antibodies: HRP donkey anti-rabbit (1:5000) and HRP goat anti-mouse (1:3000) from Jackson ImmunoResearch (West Grove, PA, USA).

[0319] As shown in FIG. 12, incubation of MDA-231 cells with anti-MOSPD2 (Fab')₂ mAb inhibited phosphorylation of the EGF Receptor (pEGF-R) as well as phosphorylation of AKT and ERK, which are mediators of the downstream signaling pathways associated with cell migration (p-AKT and p-ERK1/2, respectively).

EXAMPLE 13

Anti-MOSPD2 F(ab')₂ mAb Inhibits EGF-induced Migration of MDA-231 Cancer Cells

[0320] The effect of anti-MOSPD2 F(ab')₂ mAb on EGF-induced migration of MDA-231 cancer cells was analyzed with trans-well migration as explained in Example 2. MDA-231 breast cancer cells (3×10^5) were starved for 4-5hr in RPMI medium containing 0.5% FCS and then incubated for 1hr with anti-MOSPD2 F(ab')₂ mAb. EGF was dissolved and placed in the lower chamber (400 ng/ml) of a QCM 24-well migration assay plate (8 μ m pores) (Corning-Costar, Corning, NY) which contained RPMI medium with 10% FCS. Cells were seeded in the upper chamber, followed by over-night incubation, after which the number of cells that migrated to the lower compartment was determined by FACS.

[0321] As shown in FIG. 13, F(ab')₂ mAb significantly inhibited EGF-induced trans-well migration of MDA-231 breast cancer cells.

EXAMPLE 14

Defining Cellular Expression Specificity and Localization of MOSPD2

[0322] Analysis of different immune cell subpopulations indicated that MOSPD2 is expressed predominantly in CD14+ monocytes over T and B lymphocytes (FIG. 14A). To determine MOSPD2 mRNA expression level, RNA was extracted from cells using RNeasy mini kit (Qiagen, ValenVBA, CA). For cDNA preparation, 2 μ g of RNA was combined with qScript reaction mix and qScript reverse transcriptase (Quanta Bioscience, Gaithersburg, MD). The reaction was placed in a thermal cycler (BioRad, Hercules, CA) and a run was programmed according to manufacturer's instructions. Real-time PCR reactions were performed on an Applied Biosystems 7300 real time PCR system (Grand

Island, NY) using sets of primers for human MOSPD2, 28S to normalize RNA levels (BIOSEARCH TECHNOLOGIES, Petaluma, CA) and SYBR Green PCR Master Mix (Applied Biosystems, Warrington, UK).

[0323] MOSPD2 is predicted to be a plasma membrane protein with one transmembrane region and one residue-long intracellular tail. Fractionation of cellular compartments, and immunofluorescence staining of human monocytes, and flow cytometry on HEK 293 cells transfected to overexpress HA-tagged MOSPD2 (performed according to the methods described above) revealed that MOSPD2 is a cell surface protein that is expressed on the plasma membrane of human monocytes (FIGs. 14B-14D, respectively).

EXAMPLE 15

MOSPD2 is Expressed on Monocytes Infiltrated Into Inflamed Tissues

[0324] Formalin-fixed tissues were dehydrated, embedded in paraffin, and sectioned at 4 μ m. Immunostaining was fully calibrated on a Benchmark XT staining module (Ventana Medical Systems). After sections were dewaxed and rehydrated, anti-CD163 (Cell Marque, Rocklin, USA, MRQ-26) or anti-MOSPD2 diluted at 1:80 and 1:100, respectively, added rest for 40 minutes. Anti-CD163 staining was detected using UltraView universal Alkaline Phosphatase red detection kit (Ventana Medical Systems, 760-501) and anti-MOSPD2 staining was detected using UltraView universal DAB detection kit (Ventana Medical Systems, 760-500). When double staining was applied, MOSPD2 staining was performed first followed by CD163 staining. Slides were counterstained with hematoxylin (Ventana Medical Systems). After the run on the automated stainer was completed, slides were dehydrated consecutively in 70% ethanol, 95% ethanol and 100% ethanol for 10 sec each. Before cover slipping, sections were cleared in xylene for 10 sec and mounted with Entellan. MOSPD2 and CD163 stained slides were viewed using an Olympus BX51 microscope. Images were taken using a Nikon digital sight camera and NIS Elements Imaging Software.

[0325] As shown in FIGs. 15A-15C, MOSPD2 is expressed on monocytes infiltrated into a variety of inflamed tissues. FIG. 15A shows the staining of synovial membrane from a rheumatoid arthritis patient for CD163, MOSPD2, or both CD163 and MOSPD2. FIG. 15B shows the staining of atherosclerotic carotid tissue for CD163, MOSPD2, or both

CD163 and MOSPD2. FIG. 15C shows the staining of infiltrating ductal carcinoma breast tissue for MOSPD2. Dark arrows indicate positive staining for tumor cells. Light arrows indicate staining of infiltrating monocytes.

EXAMPLE 16

MOSPD2 Promotes Monocyte Migration

[0326] U937 monocytic line cells were transduced with sh-lenti control or sh-lenti MOSPD2 viral particles as described above. FIG. 16A shows the silencing efficacy of sh-lenti MOSPD2 as assessed by Q-PCR and western blot. When tested for migration, MOSPD2-silenced cells were severely impaired in their ability to migrate *in vitro* towards RANTES (CCL5) (FIG. 16B). Two major signaling pathways recognized as crucial for monocyte migration are the MEK-ERK and PI3K-AKT pathways (Di Lorenzo et al., 2009; Wain et al., 2002). FIG. 16C shows that phosphorylation of ERK and AKT in the presence of RANTES is completely suppressed in MOSPD2-silenced cells.

[0327] To ascertain whether the effect observed is restricted to only one chemokine, sh-control and sh-MOSPD2 silenced U937 cells were activated with ligands that induce migration and phosphorylation via different chemokine receptors. Silencing MOSPD2 impaired monocyte migration and ERK and AKT phosphorylation regardless of the chemokine used (FIGs. 16D and 16E, respectively).

EXAMPLE 17

MOSPD2 Does Not Affect IFN-gamma-induced Activation or PKC-mediated Activation

[0328] Targeting MOSPD2 did not compromise biological functions of monocytes other than migration. U937 monocytic line cells were transduced with sh-lenti control or sh-lenti MOSPD2 viral particles as described above and treated with IFN-gamma or with PMA. Western blot analysis of treated cells showed that silencing of MOSPD2 did not alter phosphorylation of downstream signaling markers by IFN-gamma or PMA (FIGs. 17A and 17B, respectively). These results suggest that MOSPD2 specifically promotes monocyte migration.

EXAMPLE 18***Epitope Mapping of Anti-MOSPD2 Antibodies***

[0329] To determine the epitope(s) that anti-MOSPD2 antibodies may specifically bind on human MOSPD2, binding affinities to various human MOSPD2 fragments are measured, as described herein, by capturing N-terminally biotinylated MOSPD2 fragments via a pre-immobilized streptavidin (SA) on a SA chip and measuring binding kinetics of anti-MOSPD2 antibodies titrated across the MOSPD2 surface (the BIAcore®3000™ surface plasmon resonance (SPR) system, Biacore, Inc., Piscataway NJ). BIAcore assays are conducted in HBS-EP running buffer (10 mM HEPES pH 7.4, 150 mM NaCl, 3 mM EDTA, 0.005% v/v polysorbate P20). MOSPD2 surfaces are prepared by diluting the N-biotinylated MOSPD2 to a concentration of less than 0.001 mg/mL into HBS-EP buffer and injecting it across the SA sensor chip using variable contact times. Low capacity surfaces, corresponding to capture levels <50 response units (RU) are used for high-resolution kinetic studies, whereas high capacity surfaces (about 800 RU of captured MOSPD2) are used for concentration studies, screening, and solution affinity determinations.

[0330] Kinetic data is obtained by diluting antibody G1 Fab serially in two- or three-fold increments to concentrations spanning 1 μ M-0.1 nM (aimed at 0.1-10 times estimated K_D). Samples are typically injected for 1 minute at 100 μ L/min and dissociation times of at least 10 minutes are allowed. After each binding cycle, surfaces are regenerated with 25 mM NaOH in 25% v/v ethanol, which is tolerated over hundreds of cycles. An entire titration series (typically generated in duplicate) is fit globally to a 1:1 Langmuir binding model using the BIAevaluation program. This returns a unique pair of association and dissociation kinetic rate constants (respectively, K_{on} and K_{off}) for each binding interaction, whose ratio gives the equilibrium dissociation constant ($K_D = K_{off}/K_{on}$).

[0331] Anti-MOSPD2 antibodies may bind to one or more of the following amino acid regions of human MOSPD2, numbered according to SEQ ID NO:1 (amino acid residues 1-518): 508-517, 501-514, 233-241, 509-517, 212-221, 13-24, 505-517, 505-514, 89-100, 506-517, 233-245, 504-514, 128-136, 218-226, 15-24, 83-96, 42-50, 462-474, 340-351, 504-517, 462-470, 327-337, 21-32, 217-226, 510-517, 178-190, 497-509, 504-516, 64-77, 504-515, 147-159, 503-315, 88-97, 208-218, 178-191, 502-515, 503-516, 497-505,

500-509, 189-202, 189-197, 505-516, 1-63, 82-239, 93-234, 327-445, 327-431, and 497-517.

EXAMPLE 19

Additional Anti-MOSPD2 Antibodies

[0332] Additional anti-MOSPD2 antibodies are generated that recognize one or more MOSPD2 epitopes, following the methodology described in Example 1 (polyclonal antibodies) or Example 8 (monoclonal antibodies).

[0333] Briefly, portions of MOSPD2 identified in Example 18 as MOSPD2 epitopes are fused to human Fc and immobilized on a solid support. A HuCAL® library (HuCAL PLATINUM® Platform; Bio-Rad AbD Serotec, GmnH) presented on phage particles is incubated with the immobilized antigen. Nonspecific antibodies are removed by extensive washing and specific antibody phages are eluted by adding a reducing agent. Antibody DNA is isolated as a pool and subcloned into an *E. coli* expression vector to generate bivalent F(ab')₂ mAb. Colonies are picked and grown in a microtiter plate. The cultures are lysed to release the antibody molecules and screened for specific antigen binding by ELISA and FACS. Unique antibodies are expressed and purified using one-step affinity chromatography, and then tested again by ELISA and FACS for specificity.

EXAMPLE 20

MOSPD2 Expression is Increased in Correlation with Tumor Grade in Various Types of Cancer

[0334] To determine whether MOSPD2 expression was associated with tumor progression, slides carrying normal and cancerous tissues in different tumor grades were screened using anti-MOSPD2 antibody as described in the Materials and Methods section. MOSPD2 abundance was scored according to the staining intensity on a scale from 0 to 3. In cases where intra-heterogeneity staining within a single core was observed, the score of the area with the highest coverage was assigned.

[0335] FIGs. 18A-18F show representative MOSPD2 staining in Breast cancer and control tissue. Normal adjacent tissue (NAT) served as a negative control, and the

escalating tumor stages included lobular carcinoma in situ (LCIS), intraductal carcinoma in situ (IDIS), invasive ductal carcinoma (IDC), invasive lobular carcinoma (ILC) and Metastatic invasive ductal carcinoma (MIDC). While representative NAT, LCIS and IDIS staining were negatively stained for MOSPD2, IDC, ILC and MIDC representative staining demonstrated intense positive MOSPD2 staining.

[0336] FIG. 19 demonstrates increased MOSPD2 staining intensity in invasive and metastatic breast cancer. Within NAT, only 18% percent (2/11) of samples showed a staining intensity of 1, while 21% (4/19) of in situ carcinoma samples (IDIS+LCIS) were scored 1 or 2. However, analysis of invasive and metastatic tissues demonstrated higher frequency in score of 2 and increased staining intensity up to score of 3, compared to NAT and in situ carcinoma (IDIS+LCIS). Thus, the percent of combined scores 2 and 3 for ILC, IDC and MIDC were 63% (12/19), 77% (50/65) and 81% (25/31), respectively.

[0337] MOSPD2 expression correlated with the transformation of cells from normal to cancerous in colon and in hepatic tissues as well. FIGs 20A-20D demonstrate that in 67% of colon cancer samples and in 45% of hepatocellular carcinoma samples tested, there was a positive MOSPD2 staining. No MOSPD2 staining (0%) was detected in the normal colon or liver tissues tested.

[0338] MOSPD2 expression also correlated with malignancy. FIGs 21A-21E show intense MOSPD2 staining in hepatocellular carcinoma that increased with tumor grade, while normal and NAT samples were negative for MOSPD2 staining.

[0339] FIGs. 22A-22B summarize the intensity of MOSPD2 staining in malignant liver tissues or controls from FIGs-21A-21E. MOSPD2 staining intensity was significantly increased by 3.2 or 4 fold in malignant samples in comparison to normal and NAT, respectively ($p \leq 0.001$). FIG. 22B shows the increase in MOSPD2 staining intensity in different stages of hepatocellular carcinoma.

EXAMPLE 21

VB-201 Inhibits MOSPD2

Labeling of VB-201 and VB-221

[0340] VB-201 and VB-221 were labeled with biotin as follows. VB-201, VB-221 and ovalbumin (OVA, Sigma, Israel) were dissolved in 0.1M MES buffer (Thermo Scientific,

Rockford, IL) and conjugated using EDC [1-ethyl-3-(dimethylaminopropyl) carbodiimide HCL] (Thermo Scientific) at a molar ratio of 100 (VB-201/VB-221):1 (OVA):240 (EDC) for 2-3hr at room temperature. After which, samples were transferred to 10kDa dialysis cassettes (Thermo Scientific) and dialyzed overnight against PBS. The ovalbumin bound VB-201 (OB201) and VB-221 (OB221) were then conjugated with amine-PEG2-biotin (in 0.1M MES buffer) using EDC at a molar ratio of 1 (OB201/OB221):100 (amine-PEG2-biotin):700 (EDC). The reaction was allowed to proceed for 2-3 hours at room temperature after which samples were again transferred to a 10kDa dialysis cassette and dialyzed overnight against PBS.

Precipitation

[0341] Cells were lysed using a 1% NP-40 lysis buffer containing 1:100 protease and phosphatase inhibitors, followed by 20 min incubation on ice and 15 min centrifugation at maximum speed. Samples were incubated overnight at 4°C with solvent, OB201 or OB221 in a rotator. Streptavidin agarose beads (Sigma, Israel) were added for 2 hours. Protein elution was performed with lysis buffer without DTT for 10 min at room temperature. Sample loading, transfer and immuno-blotting were performed as described above.

Results

VB-201 binds MOSPD2

[0342] It was previously shown that VB-201 inhibits migration of monocytes *in vitro* and *in vivo*. However, VB-221, a derivative of VB-201, did not inhibit chemokine-induced signaling and migration in human monocytes. Using labeled VB-201 and VB-221, proteins from human monocytes were precipitated and differential display by Mass-Spectrometry, was studied. The Mass-Spectrometry results revealed that MOSPD2 has a strong binding to VB-201 but not VB-221.

[0343] To further validate these results, labelled VB-201 and VB-221 were employed on cell lysates from human CD14 monocytes. Samples were then probed with anti MOSPD2 and TLR2. Whereas VB-201 and VB-221 precipitated TLR2 in a comparable intensity, VB-201 precipitated MOSPD2 markedly more intense than VB-221 (FIG. 23). These results also indicate that VB-201 binds MOSPD2.

EXAMPLE 22

MOSPD2 promotes EGF-induced signaling events in breast cancer cells

MOSPD2 Silencing in MDA-231 Breast Cancer Cells

[0344] EGF ligation to the EGF Receptor (EGF-R) induces a cascade of signaling that involves phosphorylation down-stream to the receptor. We investigated whether MOSPD2 is affecting signaling cascades induced by EGF. The human breast cancer cell line MDA-MB-231 (hereafter MDA-231) (HTB-26) was purchased from ATCC. The cells (2×10^6 in 2ml) were placed in a 15 ml tube. Lentiviral particles expressing CRISPR non-target control (CRISPR-Control) or CRISPR human MOSPD2 (CRISPR-MOSPD2) were applied on the cells which were then spun for 60 min, 2000 rpm in room temperature in the presence of 8 μ g/ml polybrene (Sigma, Israel). The cells were then seeded in a 6 well plate. After 72 hours, fresh medium containing puromycin (4 μ g/ml Sigma, Israel) was added for the selection of transduced cells. Single cell cloning was performed on CRISPR transduced cells to isolate cells with silenced MOSPD2 protein expression and impaired migration.

[0345] When CRISPR-Control MDA-231 cells were activated with EGF, the EGF-R and downstream signaling molecules became phosphorylated. However, in CRISPR-MOSPD2 silenced cells, a remarkable inhibition in EGF-R phosphorylation and the downstream molecules was observed (FIG. 24). These results indicate that MOSPD2 regulates EGF-induced signaling pathways in breast cancer cells.

[0346] All publications, patents and patent applications mentioned in this application are herein incorporated in their entirety by reference into the specification, to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated herein by reference. In addition, citation or identification of any reference in this application shall not be construed as an admission that such reference is available as prior art to the present invention. To the extent that section headings are used, they should not be construed as necessarily limiting.

WHAT IS CLAIMED IS:

1. A method of treating or preventing metastasis of a cancer cell, comprising administering to a subject in need thereof an effective amount of an inhibitor of Motile Sperm Domain containing Protein 2 (MOSPD2) expressed by the cancer cell.
2. A method of inhibiting or preventing one or more activities in or of a cancer cell, comprising administering to a subject in need thereof an effective amount of an inhibitor of MOSPD2 expressed by the cancer cell, wherein the one or more activities is one or more of: cancer cell migration, monocyte migration associated with tumor growth, a chemokine signaling pathway, a growth factor signaling pathway, EGF Receptor phosphorylation, ERK phosphorylation, AKT phosphorylation, and FAK phosphorylation.
3. A method of treating or preventing a cancer, comprising administering to a subject in need thereof an effective amount of an inhibitor of MOSPD2 expressed by circulating monocytes or tumor associated macrophages to reduce the number of tumor associated macrophages near or within the cancer mass or the migration of tumor associated macrophages.
4. A method of treating or preventing a cancer, comprising administering to a subject in need thereof a therapeutically effective amount of an inhibitor of MOSPD2 expressed by the cancer cells.
5. The method of claim 4, further comprising administering a therapeutically effective amount of an anticancer drug.
6. A method of treating or preventing a metastatic cancer, comprising administering to a subject in need thereof a therapeutically effective amount of an anticancer drug and an inhibitor of MOSPD2 expressed by the metastatic cancer cells.
7. A method of treating or preventing metastasis of a cancer cell, comprising contacting a cancer cell with an effective amount of an inhibitor of MOSPD2 expressed by the cancer cell.
8. A method of inhibiting or preventing one or more activities in or of a cancer cell, comprising contacting the cancer cell with an effective amount of an inhibitor of MOSPD2

expressed by the cancer cell, wherein the one or more activities is one or more of: MOSPD2 expression, cancer cell migration, a chemokine signaling pathway, a growth factor signaling pathway, EGF Receptor phosphorylation, ERK phosphorylation, AKT phosphorylation, and FAK phosphorylation.

9. A method of treating or preventing a cancer, comprising contacting circulating monocytes or tumor associated macrophages with an effective amount of an inhibitor of MOSPD2 expressed by the circulating monocytes or tumor associated macrophages to reduce the number of tumor associated macrophages near or within the cancer mass or migration of tumor associated macrophages.

10. A method of treating or preventing a cancer, comprising contacting cancer cells with a therapeutically effective amount of an inhibitor of MOSPD2 expressed by the cancer cells.

11. The method of claim 10, further comprising administering a therapeutically effective amount of another anticancer drug.

12. The method of any one of claims 1-11, wherein the inhibitor is a polypeptide, DNA, or RNA.

13. The method of any one of claims 1-11, wherein the inhibitor is (i) an isolated binding molecule that specifically binds to a MOSPD2 polypeptide, (ii) an isolated binding molecule that specifically binds to a ligand of a MOSPD2 polypeptide, (iii) an antisera raised against a MOSPD2 polypeptide, (iv) a soluble MOSPD2 polypeptide, or (v) a soluble MOSPD2 polypeptide comprising, consisting essentially of, or consisting of an extracellular domain of a MOSPD2 polypeptide.

14. The method of any one of claims 1-11, wherein the inhibitor is an antibody that specifically binds to a MOSPD2 polypeptide.

15. The method of any one of claims 1-11, wherein the inhibitor is an antigen binding fragment of an antibody that specifically binds to a MOSPD2 polypeptide.

16. The method of claim 14, wherein the antibody is a polyclonal, monoclonal, murine, human, humanized, or chimeric antibody.


17. The method of claim 15, wherein the antigen binding fragment is a Fab, Fab', F(ab')₂, Fv, scFv, sdFv fragment, VH domain, or VL domain.

18. The method of any one of claims 1-11, wherein the inhibitor is an RNAi, miRNA, siRNA, shRNA, an antisense RNA, an antisense DNA, a decoy molecule, a decoy DNA, a double-stranded DNA, a single-stranded DNA, a complexed DNA, an encapsulated DNA, a viral DNA, a plasmid DNA, a naked RNA, an encapsulated RNA, a viral RNA, a double-stranded RNA, a molecule capable of generating RNA interference, or combinations thereof, that hybridizes to a nucleotide sequence encoding a MOSPD2 polypeptide under a stringent condition, or a gene editing system.

19. The method of any one of claims 12-17, wherein the MOSPD2 polypeptide has a sequence at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to any one of SEQ ID NOs:1-4.

20. The method of any one of claims 12-17, wherein the MOSPD2 polypeptide is encoded by a sequence at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to any one of SEQ ID NOs:5-8.

21. The method of any one of claims 1-11, wherein the MOSPD2 inhibitor is an oxidized phospholipid having a structure according to Formula I:



Formula I

or a pharmaceutically acceptable salt, a hydrate or a solvate thereof,


wherein:

n is an integer from 1 to 6, wherein when n is 1, Cn, Bn, Rn, and Y are absent, and C₁ is attached to R'n;

each of B₁, B₂, ... B_{n-1} and B_n is independently selected from the group consisting of oxygen, sulfur, nitrogen, phosphorus and silicon, whereby each of said nitrogen, phosphorus and silicon is optionally substituted by one substituent selected from the group consisting of alkyl, halo, cycloalkyl, aryl, hydroxy, thiohydroxy, alkoxy, aryloxy, thioaryloxy, thioalkoxy and oxo;

each of A₁, A₂, ... A_{n-1} and A_n is independently selected from the group consisting of CR"R", C=O and C=S,

Y is selected from the group consisting of hydrogen, acyl, alkyl, aryl, cycloalkyl, carboxy, saccharide, phosphoric acid, phosphoryl choline, phosphoryl ethanolamine, phosphoryl serine, phosphoryl cardiolipin, phosphoryl inositol, ethylphosphocholine, phosphorylmethanol, phosphorylethanol, phosphorylpropanol, phosphorylbutanol, phosphorylethanolamine-N-lactose, phosphoethanolamine-N-glutaric acid, phosphoethanolamine-N-[methoxy(propylene glycol)], phosphoinositol-4-phosphate, phosphoinositol-4,5-biphosphonate, phosphoinositol-4,5-bisphosphate, pyrophosphate, phosphoethanolamine-diethylenetriamine-pentaacetate, dinitrophenyl-phosphoethanolamine, phosphoglycerol and a moiety having the general formula:




wherein:

each of B' and B'' is independently selected from the group consisting of sulfur and oxygen; and

each of D' and D'' is independently selected from the group consisting of hydrogen, alkyl, amino substituted alkyl, cycloalkyl, phosphonate and thiophosphonate; and


each of X₁, X₂, ... X_{n-1} is independently a saturated or unsaturated hydrocarbon having the general Formula II:



Formula II

wherein m is an integer from 1 to 26; and

Z is selected from the group consisting of:




wherein W is selected from the group consisting of oxygen and sulfur;

wherein at least one of X₁, X₂, ... X_{n-1} comprises a Z other than hydrogen,

and wherein:

each of R₁, R'₁, R₂, ... R_{n-1}, R_n, R'n, each of R'' and R''' and each of Ra, R'a, Rb, R'b, ... Rm-1, R'm-1, Rm and R'm is independently selected from the group consisting of a bond, hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, halo, trihalomethyl, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, phosphonate, phosphate, phosphinyl, sulfonyl, sulfinyl, sulfonamide, amide, carbonyl, thiocarbonyl, C-carboxy, O-carboxy, C-carbamate, N-carbamate, C-thiocarboxy, S-thiocarboxy and amino, or, alternatively, at least two of R₁, R'₁, R₂, ... R_{n-1}, R_n and R'n and/or at least two of Ra, R'a, Rb, R'b, ... Rm-1, R'm-1, Rm and R'm form at least one four-, five- or six-membered aromatic, heteroaromatic, alicyclic or heteroalicyclic ring, or a pharmaceutically acceptable salt, a hydrate or a solvate thereof.

22. The method of claim 21, wherein the oxidized phospholipid has a structure according to Formula III:



Formula III


or a pharmaceutically acceptable salt, hydrate or solvate thereof.

wherein n is an integer selected from 1 to 4;

B₁, each B₂, and B₃ are independently selected from the group consisting of oxygen, sulfur, and NR₄, wherein R₄ is selected from hydrogen, alkyl, cycloalkyl, aryl, and acyl;


A₁ and each A₂ are independently selected from the group consisting of CR_eR_{ee}, CR_e=CR_{ee}, C=O and C=S, wherein R_e and R_{ee} are independently selected from hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heteroaryl;

Y is selected from the group consisting of hydrogen, acyl, alkyl, aryl, cycloalkyl, carboxy, saccharide, phosphoric acid, phosphoryl choline, phosphoryl ethanolamine, phosphoryl serine, phosphoryl cardiolipin, phosphoryl inositol, ethylphosphocholine, phosphorylmethanol, phosphorylethanol, phosphorylpropanol, phosphorylbutanol, phosphorylethanolamine-N-lactose, phosphoethanolamine-N-glutaric acid, phosphoethanolamine-N-[methoxy(propylene glycol)], phosphoinositol-4-phosphate, phosphoinositol-4,5-bisphosphate, pyrophosphate, phosphoethanolamine-diethylenetriamine-pentaacetate, dinitrophenyl-phosphoethanolamine, phosphoglycerol, and a moiety having the general formula:



wherein: each of B and B_a is independently selected from the group consisting of sulfur and oxygen; and D and D_a are independently selected from the group consisting of hydrogen, alkyl, aminoalkyl, cycloalkyl, phosphonate and thiophosphonate;


X_1 and each X_2 are independently a saturated or unsaturated, linear or branched hydrocarbon, wherein at least one of X_1 and X_2 is substituted with an oxidized moiety Z selected from the group consisting of:



wherein W is oxygen or sulfur; and R_d and R_{dd} are independently selected from hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heteroaryl;

R_1 , R_{1a} , each R_2 , R_3 , and R_{3a} are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, halo, trihalomethyl, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, phosphonate, phosphate, phosphinyl, sulfonyl, sulfinyl, sulfonamide, amide, carbonyl, thiocarbonyl, C-carboxy, O-carboxy, C-carbamate, N-carbamate, C-thiocarboxy, S-thiocarboxy and amino, wherein at least two of R_1 , R_{1a} , R_2 , R_3 and R_{3a} are optionally joined to form a four-, five- or six-membered aromatic, heteroaromatic, alicyclic or heteroalicyclic ring.


23. The method of claim 22, wherein the oxidized phospholipid has a structure according to Formula III, and wherein X_1 and each X_2 in Formula III independently have the general Formula IV:



Formula IV

wherein m is an integer selected from 1 to 26;

Z is selected from the group consisting of:



wherein W is oxygen or sulfur; and R_d and R_{dd} are independently selected from hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heteroaryl, and wherein at least one of X₁ and X₂ comprises a Z other than hydrogen; and


R_a, R_{aa}, each R_b, each R_{bb}, R_c and R_{cc} are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, halo, trihalomethyl, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, phosphonate, phosphate, phosphinyl, sulfonyl, sulfinyl, sulfonamide, amide, carbonyl, thiocarbonyl, C-carboxy, O-carboxy, C-carbamate, N-carbamate, C-thiocarboxy, S-thiocarboxy and amino, wherein at least two of R_a, R_{aa}, R_b, R_{bb}, R_c, and R_{cc} are optionally joined to form a four-, five- or six-membered aromatic, heteroaromatic, alicyclic or heteroalicyclic ring.


24. The method of any one of claims 21-23, wherein n in Formula I is 3 or n in Formula III is 1.

25. The method of any one of claims 21-24, wherein Y is selected from the group consisting of hydrogen, acyl, alkyl, aryl, cycloalkyl, carboxy, saccharide, phosphoric acid, phosphoryl choline, phosphoryl ethanolamine, phosphoryl serine, phosphoryl cardiolipin, phosphoryl inositol, ethylphosphocholine, phosphorylmethanol, phosphorylethanol, phosphorylpropanol, phosphorylbutanol, phosphorylethanolamine-N-lactose, phosphoethanolamine-N-glutaric acid, phosphoethanolamine-N-[methoxy(propylene glycol)], phosphoinositol-4-phosphate, phosphoinositol-4,5-bisphosphate, pyrophosphate, phosphoethanolamine-diethylenetriamine-pentaacetate, dinitrophenyl-phosphoethanolamine, and phosphoglycerol.


26. The method of any one of claims 21-25, wherein Y is selected from the group consisting of hydrogen, phosphoryl choline, and phosphoryl ethanolamine.

27. The method of any one of claims 21-26, wherein each of B₁, B₂, and B₃ is oxygen.



28. The method of any one of claims 21-27, wherein Z is  , wherein W is oxygen.

29. The method of claim 21, wherein the oxidized phospholipid has a structure according to Formula IIIa:



Formula IIIa

or a pharmaceutically acceptable salt, hydrate or solvate thereof,

wherein B₁, B₂, and B₃ are independently selected from oxygen and sulfur,


A₁ and A₂ are independently selected from the group consisting of CH₂, CH=CH, C=O and C=S;

Y is selected from the group consisting of hydrogen, acyl, alkyl, aryl, cycloalkyl, carboxy, saccharide, phosphoric acid, phosphoryl choline, phosphoryl ethanolamine, phosphoryl serine, phosphoryl cardiolipin, phosphoryl inositol, ethylphosphocholine, phosphorylmethanol, phosphorylethanol, phosphorylpropanol, phosphorylbutanol, phosphorylethanolamine-N-lactose, phosphoethanolamine-N-glutaric acid, phosphoethanolamine-N-[methoxy(propylene glycol)], phosphoinositol-4-phosphate, phosphoinositol-4,5-bisphosphate, pyrophosphate, phosphoethanolamine-diethylenetriamine-pentaacetate, dinitrophenyl-phosphoethanolamine, and phosphoglycerol;

R₁, R_{1a}, R₂, R₃, and R_{3a}, are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, halo, trihalomethyl, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, phosphonate, phosphate, phosphinyl,

sulfonyl, sulfinyl, sulfonamide, amide, carbonyl, thiocarbonyl, C-carboxy, O-carboxy, C-carbamate, N-carbamate, C-thiocarboxy, S-thiocarboxy and amino, wherein at least two of R₁, R_{1a}, R₂, R₃ and R_{3a} are optionally joined to form a four-, five- or six-membered aromatic, heteroaromatic, alicyclic or heteroalicyclic ring,


and wherein X₁ and X₂ are independently a saturated or unsaturated, linear or branched hydrocarbon, wherein at least one of X₁ and X₂ is substituted with an oxidized moiety Z having a formula selected from:



wherein W is oxygen or sulfur; and R_d and R_{dd} are independently selected from hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heteroaryl.

30. The method of claim 29, wherein R₁, R_{1a}, R₂, R₃, and R_{3a} are each hydrogen.

31. The method of claim 29 or 30, wherein X₁ and X₂ independently have a structure according to Formula IVa:




Formula IVa


wherein m is an integer selected from 1 to 26,

R_a, R_{aa}, each R_b, each R_{bb}, R_c, and R_{cc} are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, halo, trihalomethyl, hydroxy, alkoxy, aryloxy, thiohydroxy, thioalkoxy, thioaryloxy, phosphonate, phosphate, phosphinyl, sulfonyl, sulfinyl, sulfonamide, amide, carbonyl, thiocarbonyl, C-carboxy, O-carboxy, C-carbamate, N-carbamate, C-thiocarboxy, S-thiocarboxy and amino, wherein at least two of R_a, R_{aa}, R_b, R_{bb}, R_c, and R_{cc} are optionally joined to form a four-, five- or six-membered aromatic, heteroaromatic, alicyclic or heteroalicyclic ring;

Z is selected from the group consisting of:



wherein W is oxygen or sulfur; and R_d and R_{dd} are independently selected from hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heteroaryl, wherein at least one of X₁ and X₂ comprises a Z other than hydrogen.




32. The method of any one of claims 29-31, wherein Z is , and wherein W is oxygen.

33. The method of any one of claims 29-32, wherein Y is selected from the group consisting of hydrogen, phosphoryl choline, and phosphoryl ethanolamine.


34. The method of any one of claims 29-33, wherein each of B₁, B₂, and B₃ is oxygen.


35. The method of claim 21, wherein the oxidized phospholipid has a structure according to the Formula VI:



Formula VI

wherein A₁ is selected from the group consisting of CH₂, CH=CH and C=O; A₂ is absent




or CH₂; X₁ is an alkyl having from 1 to 30 carbon atoms; X₂ is , wherein

E is absent or is an alkyl chain having from 1 to 24 carbon atoms;

F is selected from the group consisting of hydrogen, hydroxy, alkyl, alkoxy, halide, acetoxy and aryl; and

Z is selected from the group consisting of:



wherein R_d is selected from H, alkyl and aryl; and


Y is selected from the group consisting of hydrogen, alkyl, aryl, phosphoric acid, phosphoryl choline, phosphoryl ethanolamine, phosphoryl serine, phosphatidyl choline, phosphatidyl ethanolamine, phosphatidyl serine, phosphatidyl cardiolipin, phosphatidyl inositol, phosphoryl cardiolipin, phosphoryl inositol, ethylphosphocholine, phosphorylmethanol, phosphorylethanol, phosphorylpropanol, phosphorylbutanol, phosphorylethanolamine-N-lactose, phosphoethanolamine-N-[methoxy(propylene glycol)], phosphoinositol-4-phosphate, phosphoinositol-4,5-bisphosphate, pyrophosphate, phosphoethanolamine-diethylenetriamine-pentaacetate, dinitrophenyl-phosphoethanolamine, phosphoglycerol.

36. The method of claim 35, wherein X₁ is alkyl having from 10 to 30 carbon atoms.

37. The method of claim 35 or 36, wherein E is alkyl having from 1 to 10 carbon atoms.


38. The method of any one of claims 35-37, wherein Y is phosphocholine.

39. The method of any one of claims 21-38, wherein the oxidized phospholipid is



or a pharmaceutically acceptable salt thereof.

40. The method of any one of claims 21-39, wherein the oxidized phospholipid is



or a pharmaceutically acceptable salt thereof.

41. The method of any one of claims 1-11, wherein the MOSPD2 inhibitor is a tocopherol, a triterpene, a vitamin A, or a phospholipid.

42. The method of any one of claims 1-41, wherein the cancer is selected from the group consisting of bladder cancer, breast cancer, colon cancer, rectal cancer, kidney cancer, liver cancer, lung cancer, esophageal cancer, gall-bladder cancer, ovarian cancer, pancreatic cancer, stomach cancer, cervical cancer, thyroid cancer, prostate cancer, skin cancer, hematopoietic cancer, cancer of mesenchymal origin, cancer of central or peripheral nervous system, endometrial cancer, head and neck cancer, glioblastoma, and malignant ascites.

43. The method of claim 42, wherein the cancer is a small-cell lung cancer or a non-small-cell lung cancer.

44. The method of claim 42, wherein the cancer is skin cancer, wherein the skin cancer is selected from the group consisting of squamous cell carcinoma, basal cell cancer, melanoma, dermatofibrosarcoma protuberans, Merkel cell carcinoma, Kaposi's sarcoma, keratoacanthoma, spindle cell tumors, sebaceous carcinomas, microcystic adnexal carcinoma, Paget's disease of the breast, atypical fibroxanthoma, leiomyosarcoma, and angiosarcoma.

45. The method of claim 42, wherein the cancer is a hematopoietic cancer of lymphoid lineage.

46. The method of claim 45, wherein the hematopoietic cancer of lymphoid lineage is selected from the group consisting of leukemia, acute lymphocytic leukemia, chronic lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma and Burkitt's lymphoma.

47. The method of claim 42, wherein the cancer is a hematopoietic cancer of myeloid lineage.

48. The method of claim 47, wherein the hematopoietic cancer of myeloid lineage is selected from the group consisting of acute myelogenous leukemia, chronic myelogenous leukemia, myelodysplastic syndrome, and promyelocytic leukemia.

49. The method of claim 42, wherein the cancer is a cancer of mesenchymal origin selected from the group consisting of fibrosarcoma, rhabdomyosarcoma, soft tissue sarcoma, and bone sarcoma.

50. The method of claim 42, wherein the cancer is a cancer of the central or peripheral nervous system selected from the group consisting of astrocytoma, neuroblastoma, glioma, and schwannomas.

51. The method of any one of claims 1-41, wherein the cancer is selected from the group consisting of anal cancer, bone cancer, gastrointestinal stromal cancer, gestational trophoblastic disease, Hodgkin's lymphoma, Kaposi sarcoma, keratoacanthoma, malignant mesothelioma, multicentric castleman disease, multiple myeloma and other plasma cell neoplasms, myeloproliferative neoplasms, neuroblastoma, non-Hodgkin's lymphoma, osteosarcoma, ovarian, fallopian tube, or primary peritoneal cancer, penile cancer, retinoblastoma, rhabdomyosarcoma, seminoma, soft tissue sarcoma, stomach (gastric) cancer, testicular cancer, teratocarcinoma, thyroid follicular cancer, vaginal cancer, vulvar cancer, Wilms tumor and other childhood kidney cancers, and xeroderma pigmentosum.

52. The method of any one of claims 1-41, wherein the cancer is selected from the group consisting of bladder cancer, brain cancer, breast cancer, colon cancer, esophageal cancer, lung cancer, skin cancer, tongue cancer, kidney cancer, and hepatic cancer.

53. The method of any one of claims 1-52, wherein MOSPD2 is upregulated in the cancer cell compared to its non-cancerous counterpart.

54. The method of any one of claims 1-53, wherein migration of the cancer cell is inhibited.

55. The method of any one of claims 1-54, wherein ERK phosphorylation in the cancer cell is inhibited.

56. The method of any one of claims 1-55, wherein AKT phosphorylation in the cancer cell is inhibited.

57. The method of any one of claims 1-56, wherein FAK phosphorylation in the cancer cell is inhibited.

58. The method of any one of claims 1-6 and 12-57, wherein the subject is a human.

59. The method of any one of claims 4, 5, and 11-58, where in the anticancer drug is selected from the group consisting of Abiraterone Acetate, Abitrexate (Methotrexate), Abraxane (Paclitaxel Albumin-stabilized Nanoparticle Formulation), ABVD, ABVE, ABVE-PC, AC, AC-T, Adcetris (Brentuximab Vedotin), ADE, Ado-Trastuzumab Emtansine, Adriamycin (Doxorubicin Hydrochloride), Adrucil (Fluorouracil), Afatinib Dimaleate, Afinitor (Everolimus), Akynzeo (Netupitant and Palonosetron Hydrochloride), Aldara (Imiquimod), Aldesleukin, Alemtuzumab, Alimta (Pemetrexed Disodium), Aloxi (Palonosetron Hydrochloride), Ambochlorin (Chlorambucil), Amboclorin (Chlorambucil), Aminolevulinic Acid, Anastrozole, Aprepitant, Aredia (Pamidronate Disodium), Arimidex (Anastrozole), Aromasin (Exemestane), Arranon (Nelarabine), Arsenic Trioxide, Arzerra (Ofatumumab), Asparaginase Erwinia chrysanthemi, Avastin (Bevacizumab), Axitinib, Azacitidine, BEACOPP, Becenum (Carmustine), Beleodaq (Belinostat), Belinostat, Bendamustine Hydrochloride, BEP, Bevacizumab, Bexarotene, Bexxar (Tositumomab and I 131 Iodine Tositumomab), Bicalutamide, BiCNU (Carmustine), Bleomycin, Blinatumomab, Blincyto (Blinatumomab), Bortezomib, Bosulif (Bosutinib), Bosutinib, Brentuximab Vedotin, Busulfan, Busulfex (Busulfan), Cabazitaxel, Cabozantinib-S-Malate, CAF, Campath (Alemtuzumab), Camptosar (Irinotecan Hydrochloride), Capecitabine, CAPOX, Carboplatin, Carboplatin-Taxol, Carfilzomib, Carmubris (Carmustine), Carmustine, Carmustine Implant, Casodex (Bicalutamide), CeeNU (Lomustine), Ceritinib, Cerubidine (Daunorubicin Hydrochloride), Cervarix (Recombinant HPV Bivalent Vaccine), Cetuximab, Chlorambucil, Chlorambucil-Prednisone, CHOP, Cisplatin, Clafen (Cyclophosphamide), Clofarabine, CMF,Cometriq (Cabozantinib-S-Malate), COPP, COPP-ABV, Cosmegen (Dactinomycin), Crizotinib, CVP, Cyclophosphamide, Cyfos (Ifosfamide), Cyramza (Ramucirumab), Cytarabine, Cytarabine, Liposomal, Cytosar-U (Cytarabine), Cytoxan

(Cyclophosphamide), Dabrafenib, Dacarbazine, Dacogen (Decitabine), Dactinomycin, Dasatinib, Daunorubicin Hydrochloride, Decitabine, Degarelix, Denileukin Diftitox, Denosumab, DepoCyt (Liposomal Cytarabine), DepoFoam (Liposomal Cytarabine), Dexrazoxane Hydrochloride, Dinutuximab, Docetaxel, Doxil (Doxorubicin Hydrochloride Liposome), Doxorubicin Hydrochloride, Doxorubicin Hydrochloride Liposome, Dox-SL (Doxorubicin Hydrochloride Liposome), DTIC-Dome (Dacarbazine), Efudex (Fluorouracil), Elitek (Rasburicase), Ellence (Epirubicin Hydrochloride), Eloxatin (Oxaliplatin), Eltrombopag Olamine, Emend (Aprepitant), Enzalutamide, Epirubicin Hydrochloride, EPOCH, Erbitux (Cetuximab), Eribulin Mesylate, Erivedge (Vismodegib), Erlotinib Hydrochloride, Erwinaze (Asparaginase Erwinia chrysanthemi), Etopophos (Etoposide Phosphate), Etoposide, Etoposide Phosphate, Evacet (Doxorubicin Hydrochloride Liposome), Everolimus, Evista (Raloxifene Hydrochloride), Exemestane, Fareston (Toremifene), Farydak (Panobinostat), Faslodex (Fulvestrant), FEC, Femara (Letrozole), Filgrastim, Fludara (Fludarabine Phosphate), Fludarabine Phosphate, Fluoroplex (Fluorouracil), Fluorouracil, Folex (Methotrexate), Folex PFS (Methotrexate), Folfiri, Folfiri-Bevacizumab, Folfiri-Cetuximab, Folfirinox, Folflox, Folotyn (Pralatrexate), FU-LV, Fulvestrant, Gardasil (Recombinant HPV Quadrivalent Vaccine), Gardasil 9 (Recombinant HPV Nonavalent Vaccine), Gazyva (Obinutuzumab), Gefitinib, Gemcitabine Hydrochloride, Gemcitabine-Cisplatin, Gemcitabine-Oxaliplatin, Gemtuzumab Ozogamicin, Gemzar (Gemcitabine Hydrochloride), Gilotrif (Afatinib Dimaleate), Gleevec (Imatinib Mesylate), Gliadel (Carmustine Implant), Gliadel wafer (Carmustine Implant), Glucarpidase, Goserelin Acetate, Halaven (Eribulin Mesylate), Herceptin (Trastuzumab), HPV Bivalent Vaccine, Recombinant, HPV Nonavalent Vaccine, Recombinant, HPV Quadrivalent Vaccine, Recombinant, Hycamtin (Topotecan Hydrochloride), Hyper-CVAD, Ibrance (Palbociclib), Ibrutumomab Tiuxetan, Ibrutinib, ICE, Iclusig (Ponatinib Hydrochloride), Idamycin (Idarubicin Hydrochloride), Idarubicin Hydrochloride, Idelalisib, Ifex (Ifosfamide), Ifosfamide, Ifosfamidum (Ifosfamide), Imatinib Mesylate, Imbruvica (Ibrutinib), Imiquimod, Inlyta (Axitinib), Intron A (Recombinant Interferon Alfa-2b), Iodine 131 Tositumomab and Tositumomab, Ipilimumab, Iressa (Gefitinib), Irinotecan Hydrochloride, Istodax (Romidepsin), Ixabepilone, Ixempra (Ixabepilone), Jakafi (Ruxolitinib Phosphate), Jevtana (Cabazitaxel), Kadcyla (Ado-Trastuzumab Emtansine), Keoxifene (Raloxifene Hydrochloride), Kepivance (Palifermin), Keytruda (Pembrolizumab), Kyprolis (Carfilzomib), Lanreotide Acetate, Lapatinib Ditosylate, Lenalidomide, Lenvatinib Mesylate, Lenvima (Lenvatinib Mesylate), Letrozole, Leucovorin

Calcium, Leukeran (Chlorambucil), Leuprolide Acetate, Levulan (Aminolevulinic Acid), Linfozin (Chlorambucil), LipoDox (Doxorubicin Hydrochloride Liposome), Liposomal Cytarabine, Lomustine, Lupron (Leuprolide Acetate), Lupron Depot (Leuprolide Acetate), Lupron Depot-Ped (Leuprolide Acetate), Lupron Depot-3 Month (Leuprolide Acetate), Lupron Depot-4 Month (Leuprolide Acetate), Lynparza (Olaparib), Marqibo (Vincristine Sulfate Liposome), Matulane (Procarbazine Hydrochloride), Mechlorethamine Hydrochloride, Megace (Megestrol Acetate), Megestrol Acetate, Mekinist (Trametinib), Mercaptopurine, Mesna, Mesnex (Mesna), Methazolastone (Temozolomide), Methotrexate, Methotrexate LPF (Methotrexate), Mexate (Methotrexate), Mexate-AQ (Methotrexate), Mitomycin C, Mitoxantrone Hydrochloride, Mitozytrex (Mitomycin C), MOPP, Mozobil (Plerixafor), Mustargen (Mechlorethamine Hydrochloride), Mutamycin (Mitomycin C), Myleran (Busulfan), Mylosar (Azacitidine), Mylotarg (Gemtuzumab Ozogamicin), Nanoparticle Paclitaxel (Paclitaxel Albumin-stabilized Nanoparticle Formulation), Navelbine (Vinorelbine Tartrate), Nelarabine, Neosar (Cyclophosphamide), Netupitant and Palonosetron Hydrochloride, Neupogen (Filgrastim), Nexavar (Sorafenib Tosylate), Nilotinib, Nivolumab, Nolvadex (Tamoxifen Citrate), Nplate (Romiplostim), Obinutuzumab, OEPA, Ofatumumab, OFF, Olaparib, Omacetaxine Mepesuccinate, Oncaspar (Pegaspargase), Ontak (Denileukin Diftitox), Opdivo (Nivolumab), OPPA, Oxaliplatin, Paclitaxel, Paclitaxel Albumin-stabilized Nanoparticle Formulation, PAD, Palbociclib, Palifermin, Palonosetron Hydrochloride, Pamidronate Disodium, Panitumumab, Panobinostat, Paraplat (Carboplatin), Paraplatin (Carboplatin), Pazopanib Hydrochloride, Pegaspargase, Peginterferon Alfa-2b, PEG-Intron (Peginterferon Alfa-2b), Pembrolizumab, Pemetrexed Disodium, Perjeta (Pertuzumab), Pertuzumab, Platinol (Cisplatin), Platinol-AQ (Cisplatin), Plerixafor, Pomalidomide, Pomalyst (Pomalidomide), Ponatinib Hydrochloride, Pralatrexate, Prednisone, Procarbazine Hydrochloride, Proleukin (Aldesleukin), Prolia (Denosumab), Promacta (Eltrombopag Olamine), Provenge (Sipuleucel-T), Purinethol (Mercaptopurine), Purixan (Mercaptopurine), Radium 223 Dichloride, Raloxifene Hydrochloride, Ramucirumab, Rasburicase, R-CHOP, R-CVP, Recombinant Human Papillomavirus (HPV) Bivalent Vaccine, Recombinant Human Papillomavirus (HPV) Nonavalent Vaccine, Recombinant Human Papillomavirus (HPV) Quadrivalent Vaccine, Recombinant Interferon Alfa-2b, Regorafenib, R-EPOCH, Revlimid (Lenalidomide), Rheumatrex (Methotrexate), Rituxan (Rituximab), Rituximab, Romidepsin, Romiplostim, Rubidomycin (Daunorubicin Hydrochloride), Ruxolitinib Phosphate, Sclerosol Intrapleural

Aerosol (Talc), Siltuximab, Sipuleucel-T, Somatuline Depot (Lanreotide Acetate), Sorafenib Tosylate, Sprycel (Dasatinib), STANFORD V, Sterile Talc Powder (Talc), Steritalc (Talc), Stivarga (Regorafenib), Sunitinib Malate, Sutent (Sunitinib Malate), Sylatron (Peginterferon Alfa-2b), Sylvant (Siltuximab), Synovir (Thalidomide), Synribo (Omacetaxine Mepesuccinate), TAC, Tafinlar (Dabrafenib), Talc, Tamoxifen Citrate, Tarabine PFS (Cytarabine), Tarceva (Erlotinib Hydrochloride), Targretin (Bexarotene), Tasigna (Nilotinib), Taxol (Paclitaxel), Taxotere (Docetaxel), Temodar (Temozolomide), Temozolomide, Temsirolimus, Thalidomide, Thalomid (Thalidomide), Thiotepa, Toposar (Etoposide), Topotecan Hydrochloride, Toremifene, Torisel (Temsirolimus), Tositumomab and I 131 Iodine Tositumomab, Totect (Dexrazoxane Hydrochloride), TPF, Trametinib, Trastuzumab, Treanda (Bendamustine Hydrochloride), Trisenox (Arsenic Trioxide), Tykerb (Lapatinib Ditosylate), Unituxin (Dinutuximab), Vandetanib, VAMP, Vectibix (Panitumumab), VeIP, Velban (Vinblastine Sulfate), Velcade (Bortezomib), Velsar (Vinblastine Sulfate), Vemurafenib, VePesid (Etoposide), Viadur (Leuprolide Acetate), Vidaza (Azacitidine), Vinblastine Sulfate, Vincasar PFS (Vincristine Sulfate), Vincristine Sulfate, Vincristine Sulfate Liposome, Vinorelbine Tartrate, VIP, Vismodegib, Voraxaze (Glucarpidase), Vorinostat, Votrient (Pazopanib Hydrochloride), Wellcovorin (Leucovorin Calcium), Xalkori (Crizotinib), Xeloda (Capecitabine), XELIRI, XELOX, Xgeva (Denosumab), Xofigo (Radium 223 Dichloride), Xtandi (Enzalutamide), Yervoy (Ipilimumab), Zaltrap (Ziv-Aflibercept), Zelboraf (Vemurafenib), Zevalin (Ibritumomab Tiuxetan), Zinecard (Dexrazoxane Hydrochloride), Ziv-Aflibercept, Zoladex (Goserelin Acetate), Zoledronic Acid, Zolinza (Vorinostat), Zometa (Zoledronic Acid), Zydelig (Idelalisib), Zykadia (Ceritinib), and Zytiga (Abiraterone Acetate).

60. An isolated polypeptide that inhibits MOSPD2 expressed by a cancer cell.

61. The isolated polypeptide of claim 60, wherein the polypeptide (i) specifically binds to a MOSPD2 polypeptide, (ii) specifically binds to a ligand of a MOSPD2 polypeptide, (iii) is an antisera raised against a MOSPD2 polypeptide, (iv) is a soluble MOSPD2 polypeptide, or (v) is a soluble MOSPD2 polypeptide comprising, consisting essentially of, or consisting of an extracellular domain of a MOSPD2 polypeptide.

62. The isolated polypeptide of claim 61, wherein the polypeptide is an antibody that specifically binds to a MOSPD2 polypeptide.

63. The isolated polypeptide of claim 61, wherein the polypeptide is an antigen binding fragment of an antibody that specifically binds to a MOSPD2 polypeptide.

64. The isolated polypeptide of claim 62, wherein the antibody is a polyclonal, monoclonal, murine, human, humanized, or chimeric antibody.

65. The isolated polypeptide of claim 63, wherein the antigen binding fragment is a Fab, Fab', F(ab')₂, Fv, scFv, sdFv fragment, VH domain, or VL domain.

66. The isolated polypeptide of any one of claims 60-65, wherein the MOSPD2 polypeptide has a sequence at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to any one of SEQ ID NOs:1-4.

67. The isolated polypeptide of any one of claims 60-65, wherein the MOSPD2 polypeptide is encoded by a nucleotide sequence at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to any one of SEQ ID NOs:5-8.

68. A pharmaceutical composition comprising the isolated polypeptide of any one of claims 60-67, and a pharmaceutically acceptable carrier.

69. A pharmaceutical composition comprising an isolated polypeptide that inhibits MOSPD2 expressed by a cancer cell, and a pharmaceutically acceptable carrier.

70. The pharmaceutical composition of claim 68 or 69, suitable for systemic or local administration.

71. The pharmaceutical composition of claim 68 or 69, suitable for nasal, oral, intra-peritoneal, or intra-tumor administration.

72. The pharmaceutical composition of claim 68 or 69, suitable for intravenous administration, intramuscular administration, or subcutaneous administration.

73. The isolated polypeptide of any one of claims 60-67 or the pharmaceutical composition of any one of claims 62-68, wherein the cancer is selected from the group consisting of bladder cancer, breast cancer, colon cancer, rectal cancer, kidney cancer, liver cancer, lung

cancer, esophageal cancer, gall-bladder cancer, ovarian cancer, pancreatic cancer, stomach cancer, cervical cancer, thyroid cancer, prostate cancer, skin cancer, hematopoietic cancer, cancer of mesenchymal origin, cancer of central or peripheral nervous system, endometrial cancer, head and neck cancer, glioblastoma, and malignant ascites.

74. The isolated polypeptide or pharmaceutical composition of claim 73, wherein the cancer is a small-cell lung cancer or a non-small-cell lung cancer.

75. The isolated polypeptide or pharmaceutical composition of claim 73, wherein the cancer is skin cancer, wherein the skin cancer is selected from the group consisting of squamous cell carcinoma, basal cell cancer, melanoma, dermatofibrosarcoma protuberans, Merkel cell carcinoma, Kaposi's sarcoma, keratoacanthoma, spindle cell tumors, sebaceous carcinomas, microcystic adnexal carcinoma, Paget's disease of the breast, atypical fibroxanthoma, leiomyosarcoma, and angiosarcoma.

76. The isolated polypeptide or pharmaceutical composition of claim 73, wherein the cancer is a hematopoietic cancer of lymphoid lineage.

77. The isolated polypeptide or pharmaceutical composition of claim 76, wherein the hematopoietic cancer of lymphoid lineage is selected from the group consisting of leukemia, acute lymphocytic leukemia, chronic lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma, and Burkitt's lymphoma.

78. The isolated polypeptide or pharmaceutical composition of claim 73, wherein the cancer is a hematopoietic cancer of myeloid lineage.

79. The isolated polypeptide or pharmaceutical composition of claim 78, wherein the hematopoietic cancer of myeloid lineage is selected from the group consisting of acute myelogenous leukemia, chronic myelogenous leukemia, myelodysplastic syndrome, and promyelocytic leukemia.

80. The isolated polypeptide or pharmaceutical composition of claim 73, wherein the cancer is a cancer of mesenchymal origin selected from the group consisting of fibrosarcoma, rhabdomyosarcoma, soft tissue sarcoma, and bone sarcoma.

81. The isolated polypeptide or pharmaceutical composition of claim 73, wherein the cancer is a cancer of the central or peripheral nervous system selected from the group consisting of astrocytoma, neuroblastoma, glioma, and schwannomas.

82. The isolated polypeptide of any one of claims 60-67 or the pharmaceutical composition of any one of claims 68-72, wherein the cancer is selected from the group consisting of anal cancer, bone cancer, gastrointestinal stromal cancer, gestational trophoblastic disease, Hodgkin's lymphoma, Kaposi sarcoma, keratoacanthoma, malignant mesothelioma, multicentric castleman disease, multiple myeloma and other plasma cell neoplasms, myeloproliferative neoplasms, neuroblastoma, non-Hodgkin's lymphoma, osteosarcoma, ovarian, fallopian tube, or primary peritoneal cancer, penile cancer, retinoblastoma, rhabdomyosarcoma, seminoma, soft tissue sarcoma, stomach (gastric) cancer, testicular cancer, teratocarcinoma, thyroid follicular cancer, vaginal cancer, vulvar cancer, Wilms tumor and other childhood kidney cancers, and xeroderma pigmentosum.

83. The isolated polypeptide of any one of claims 60-67 or the pharmaceutical composition of any one of claims 68-72, wherein the cancer is selected from the group consisting of bladder cancer, brain cancer, breast cancer, colon cancer, esophageal cancer, lung cancer, skin cancer, tongue cancer, kidney cancer, and hepatic cancer.

84. An isolated antibody or antigen binding fragment thereof that specifically binds to MOSPD2.

85. An isolated antibody or antigen binding fragment thereof that binds to MOSPD2 with an equilibrium dissociation constant (K_D) of from about 10^{-6} M to about 10^{-12} M.

86. The isolated antibody or antigen binding fragment thereof of claim 84 or 85, wherein the MOSPD2 is human MOSPD2.

87. The isolated antibody or antigen binding fragment thereof of any one of claims 84-86, wherein the antibody or antigen binding fragment thereof specifically binds to one or more of the following amino acid regions of human MOSPD2, numbered according to SEQ ID NO:1: about 508 to about 517, about 501 to about 514, about 233 to about 241, about 509 to about 517, about 212 to about 221, about 13 to about 24, about 505 to about 517, about 505 to about 514, about 89 to about 100, about 506 to about 517, about 233 to about 245, about 504 to

about 514, about 128 to about 136, about 218 to about 226, about 15 to about 24, about 83 to about 96, about 42 to about 50, about 462 to 474, about 340 to about 351, about 504 to about 517, about 462 to about 470, about 327 to about 337, about 21 to about 32, about 217 to about 226, about 510 to about 517, about 178 to about 190, about 497 to about 509, about 504 to about 516, about 64 to about 77, about 504 to about 515, about 147 to about 159, about 503 to about 515, about 88 to about 97, about 208 to about 218, about 178 to about 191, about 502 to about 515, about 503 to about 516, about 497 to about 505, about 500 to about 509, about 189 to about 202, about 189 to about 197, about 505 to about 516, about 1 to about 63, about 82 to about 239, about 93 to about 234, about 327 to about 445, about 327 to about 431, and about 497 to about 517.

88. The isolated antibody or antigen binding fragment thereof of any one of claims 84-86, wherein the antibody or antigen binding fragment thereof specifically binds to one or more of the following amino acid regions of human MOSPD2, numbered according to SEQ ID NO:1: about 505 to about 515, about 500 to about 515, about 230 to about 240, about 510 to about 520, about 210 to about 220, about 15 to about 25, about 505 to about 520, about 505 to about 515, about 90 to about 100, about 505 to about 525, about 230 to about 245, about 505 to about 510, about 130 to about 140, about 220 to about 230, about 15 to about 30, about 80 to about 95, about 40 to about 50, about 460 to about 475, about 340 to about 350, about 500 to about 515, about 460 to about 470, about 325 to about 335, about 20 to about 35, about 215 to about 225, about 510 to about 520, about 175 to about 190, about 500 to about 510, about 505 to about 530, about 60 to about 75, about 500 to about 520, about 145 to about 160, about 502 to about 515, about 85 to about 100, about 205 to about 220, about 175 to about 190, about 500 to about 505, about 500 to about 525, about 495 to about 505, about 495 to about 510, about 190 to about 200, about 190 to about 198, about 502 to about 515, about 1 to about 60, about 80 to about 240, about 90 to about 235, about 330 to about 445, about 330 to about 430, and about 495 to about 515.

89. The isolated antibody or antigen binding fragment thereof of any one of claims 84-88, wherein the antibody is an IgG, IgM, IgE, IgA or IgD molecule, or is derived therefrom.

90. The isolated antibody or antigen binding fragment thereof of any one of claims 84-89, wherein the antibody comprises an Fc region.

89. A pharmaceutical composition comprising the antibody or antigen binding fragment thereof of any one of claims 84-88, and a pharmaceutically acceptable carrier.

90. A method of treating or preventing metastasis of a cancer cell, comprising administering to a subject in need thereof a therapeutically effective amount of the antibody or antigen binding fragment thereof of any one of claims 84-88 or the pharmaceutical composition of claim 89.

91. A method of inhibiting or preventing one or more activities in or of a cancer cell, comprising administering to a subject in need thereof a therapeutically effective amount of the antibody or antigen binding fragment thereof of any one of claims 84-88 or the pharmaceutical composition of claim 89, wherein the one or more activities is one or more of: MOSPD2 expression, cancer cell migration, monocyte migration associated with tumor growth, a chemokine signaling pathway, a growth factor signaling pathway, EGF Receptor phosphorylation, ERK phosphorylation, AKT phosphorylation, and FAK phosphorylation.

92. A method of treating or preventing a cancer, comprising administering to a subject in need thereof a therapeutically effective amount of the antibody or antigen binding fragment thereof of any one of claims 84-88 or the pharmaceutical composition of claim 89.

93. A method of treating or preventing a metastatic cancer, comprising administering to a subject in need thereof a therapeutically effective amount of the antibody or antigen binding fragment thereof of any one of claims 84-88 or the pharmaceutical composition of claim 89.

94. A method for the prediction, diagnosis, or prognosis of cancer or cancer metastasis in a subject, which comprises determining the expression level of MOSPD2 in a sample of the subject.

95. A method for the prediction, diagnosis, or prognosis of tumor progression or invasiveness in a subject, which comprises determining the expression level of MOSPD2 in a sample of the subject.

96. The method of claim 94 or 95, wherein the expression level of MOSPD2 is the level of MOSPD2 gene expression.

97. The method of claim 94 or 95, wherein the expression level of MOSPD2 is the level of MOSPD2 protein expression.

98. An *in vitro* method for the prediction, diagnosis, or prognosis of cancer in a subject, which comprises (i) determining or quantifying the expression level of MOSPD2 in a sample of the subject, and (ii) comparing the expression level obtained in step (i) with a control or reference value, wherein an increased expression level of MOSPD2 with respect to the control or reference value is indicative of cancer, an increased risk of developing cancer, or a poor cancer prognosis.

99. An *in vitro* method for the prediction, diagnosis, or prognosis of cancer metastasis in a subject, which comprises (i) determining or quantifying the expression level of MOSPD2 in a sample of the subject, and (ii) comparing the expression level obtained in step (i) with a control or reference value, wherein an increased expression level of MOSPD2 with respect to the control or reference value is indicative of cancer metastasis, an increased risk of cancer metastasis, or a poor cancer metastasis prognosis.

100. An *in vitro* method for the prediction, diagnosis, or prognosis of tumor progression or invasiveness in a subject, which comprises (i) determining or quantifying the expression level of MOSPD2 in a sample of the subject, and (ii) comparing the expression level obtained in step (i) with a control or reference value, wherein an increased expression level of MOSPD2 with respect to the control or reference value is indicative of tumor progression or invasiveness, an increased risk of tumor progression or invasiveness, or a poor tumor progression or invasiveness prognosis.

101. The method of any one of claims 98-100, wherein if MOSPD2 expression is present in the sample of the subject, then the subject has cancer, cancer metastasis, tumor progression or invasiveness, an increased risk of cancer, an increased risk of cancer metastasis, an increased risk of tumor progression or invasiveness, a poor cancer prognosis, a poor cancer metastasis prognosis, or a poor tumor progression or invasiveness prognosis.

102. The method of any one of claims 98-100, wherein if MOSPD2 expression is present in the sample of the subject in an amount greater than MOSPD2 expression of the control or reference value, then the subject has cancer, cancer metastasis, tumor progression or invasiveness, an increased risk of cancer, an increased risk of cancer metastasis, an increased risk

of tumor progression or invasiveness, a poor cancer prognosis, a poor cancer metastasis prognosis, or a poor tumor progression or invasiveness prognosis.

103. The method of any one of claims 98-102, further comprising one or more of the following steps:

instructing a laboratory to quantify the expression level of MOSPD2 in the sample;
obtaining a report of the expression level of MOSPD2 in the sample from a laboratory;
and/or

administering a therapeutically effective amount of an inhibitor of MOSPD2 to the subject.

104. The method of any one of claims 98-103, wherein the sample is a tissue biopsy, tumor biopsy, or blood sample from the subject.

105. The method of any one of claims 98-104, wherein the control or reference value is the expression level of MOSPD2 in normal tissue or normal adjacent tissue (NAT).

106. The method of any one of claims 98-104, wherein the control or reference value is no detectable MOSDP2 expression or no significant MOSPD2 expression.

107. A method for treating a cancer or cancer metastasis responsive to an inhibitor of MOSPD2, in a subject, which comprises (i) determining the expression level of MOSPD2 in the subject, and when the expression level is determined to be greater than that of a control or reference value, (ii) administering, to the subject, a therapeutically effective amount of an inhibitor of MOSPD2.

108. A method for preventing or treating a MOSPD2 expressing tumor, in a subject, which comprises administering a therapeutically effective amount of an inhibitor of MOSPD2.

109. A method for preventing or treating a tumor having MOSPD2 expressing tumor associated macrophages, in a subject, which comprises administering a therapeutically effective amount of an inhibitor of MOSPD2.

FIG. 1
sh-Control **sh-MOSPD2**





FIG. 2

FIG. 3A




FIG. 3B




FIG. 3C





FIG. 4A



FIG. 4B


FIG. 4C



9/33


FIG. 4D

Anti-MOSPD2




Normal
Tongue

Anti-MOSPD2



Normal
Kidney

Head and Neck
(Tongue)
cell carcinoma



Kidney Clear cell
carcinoma






FIG. 4E

Anti-MOSPD2



Normal
Hepatic Tissue



Hepatocellular
Carcinoma





FIG. 5A

FIG. 5B





CRISPR-Control



CRISPR-MOSPD2

FIG. 5C








FIG. 6

FIG. 7

FIG. 8






FIG. 9



FIG. 10

FIG. 11
A A2058 Melanoma
B (HepG2) Liver cancer






FIG. 13





FIG. 14







FIG. 16



26/33





FIG. 18






FIG. 19



FIG. 20

FIG. 21






FIG. 23





FIG. 24



INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB2016/054584

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 35/13; A61P 35/00; G01N 33/574 (2016.01)

CPC - A61K 35/13; C07K 2317/76; C12Q 2600/118; G01N 33/57407; G01N 33/57419 (2016.11)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC - A61K 35/13; A61P 35/00; G01N 33/574

CPC - A61K 35/13; C07K 2317/76; C12Q 2600/118; G01N 33/574; G01N 33/57407; G01N 33/57419

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
USPC - 435/6.13; 435/7.23; 435/375 (keyword delimited)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Orbit, Google Patents, Google Scholar.

Search terms used: MOSPD2, MOSPD1, MOSPD3, motile sperm domain containing, cancer, tumor, carcinoma, monocyte, macrophage, expression, blood, circulating, associated, antibody, monoclonal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2011/0257034 A1 (BARANY et al) 20 October 2011 (20.10.2011) entire document	94-102
A	WO 2012/121679 A1 (AGENCY FOR SCIENCE, TECHNOLOGY AND RESEARCH et al) 13 September 2012 (13.09.2012) entire document	1-18, 21-23, 29, 30, 35, 36, 41, 60-67, 69, 84-86, 94-102, 107-109
A	BUERGER, K. "Functional Analysis of The MOSPD Gene Family," Thesis, University of Edinburgh, 31 March 2010 (31.03.2010), originally Pgs. 1-212 (Pgs. 1-100 provided). Retrieved from the Internet:<www.era.lib.ed.ac.uk/bitstream/handle/1842/4435/Buerger2010.pdf> on 02 November 2016 (02.11.2016). entire document	1-18, 21-23, 29, 30, 35, 36, 41, 60-67, 69, 84-86, 94-102, 107-109
A	HAN et al. "Sperm and Oocyte Communication Mechanisms Controlling C. elegans Fertility," Developmental Dynamics, 23 December 2009 (23.12.2009), Vol. 239, Pgs. 1265-1281. entire document	1-18, 21-23, 29, 30, 35, 36, 41, 60-67, 69, 84-86, 94-102, 107-109
A	RU et al. "Transient receptor potential-canonical 3 modulates sperm motility and capacitation-associated protein tyrosine phosphorylation via [Ca ²⁺] mobilization," Acta Biochimica et Biophysica Sinica, 23 April 2015 (23.04.2015), Vol. 47, Pgs. 404-413. entire document	1-18, 21-23, 29, 30, 35, 36, 41, 60-67, 69, 84-86, 94-102, 107-109

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search
02 November 2016

Date of mailing of the international search report

13 DEC 2016

Name and mailing address of the ISA/US
Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, VA 22313-1450
Facsimile No. 571-273-8300

Authorized officer

Blaine R. Copenheaver

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB2016/054584

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.: because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.: 19, 20, 24-28, 31-34, 37-40, 42-59, 68, 70-83, 87, 88, 89a, 89b, 90a, 90b, 91-93, 103-106 because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.