



(51) International Patent Classification:

A61K 31/47 (2006.01) C07D 405/04 (2006.01)
 A61K 31/4709 (2006.01) C07D 405/12 (2006.01)
 C07D 215/22 (2006.01) C07D 413/04 (2006.01)
 C07D 401/04 (2006.01) C07D 413/12 (2006.01)
 C07D 401/06 (2006.01) A61P 35/00 (2006.01)
 C07D 401/12 (2006.01)

(21) International Application Number:

PCT/US2019/015297

(22) International Filing Date:

25 January 2019 (25.01.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/622,702 26 January 2018 (26.01.2018) US
 62/758,321 09 November 2018 (09.11.2018) US

(71) Applicant: **EXELIXIS, INC.** [US/US]; 1851 Harbor Bay Parkway, Alameda, CA 94502 (US).

(72) Inventors: **BANNEN, Lynne, Canne**; 66 Bridge Rd., Novato, CA 94945 (US). **BUL, Minna**; 2445 25th Ave., Oakland, CA 94601 (US). **JIANG, Faming**; 5780 Shadow Ridge Drive, Castro Valley, CA 94552 (US). **TSO, Kin**; 5000 Diamond Heights Boulevard, San Francisco, CA 94131 (US). **WANG, Yong**; 170 Harbor Way, South San Francisco, CA 94080 (US). **XU, Wei**; 327 Glasgow Circle, Danville, CA 94526 (US).

(74) Agent: **BERVEN, Heidi M.**; Honigman LLP, 650 Trade Centre Way, Suite 200, Kalamazoo, MI 49002 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,


SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: COMPOUNDS FOR THE TREATMENT OF KINASE-DEPENDENT DISORDERS



(I)

(57) Abstract: Disclosed herein are compounds of formula I. Compounds of formula I inhibit, regulate and/or modulate kinase receptor, particularly Axl and Mer signal transduction pathways related to the changes in cellular activities as mentioned above, compositions which contain these compounds, and methods of using them to treat kinase-dependent diseases and conditions. The present invention also provides methods for making compounds as mentioned above, and compositions which contain these compounds.

COMPOUNDS FOR THE TREATMENT OF KINASE-DEPENDENT DISORDERS

Field of the Invention

[0001] The invention relates to compounds that modulate cellular activities such as proliferation, differentiation, programmed cell death, migration, and chemoinvasion by modulating protein kinase enzymatic activity. Even more specifically, the invention relates to compounds which inhibit, regulate, and/or modulate Axl and Mer receptor tyrosine kinases, compositions which contain these compounds, methods of using them to treat kinase-dependent diseases and conditions, synthesis of the compounds, and processes for formulating the compounds for pharmaceutical purposes.

Cross-Reference to Related Applications

[0002] This application claims priority to U.S. Provisional Application Serial Number 62/622,702, filed January 26, 2018, and to U.S. Provisional Application Serial Number 62/758,321, filed November 9, 2018, the entire contents of which are incorporated herein

Background of the Invention

[0003] Human Axl belongs to the TAM subfamily of receptor tyrosine kinases that includes Mer. TAM kinases are characterized by an extracellular ligand binding domain consisting of two immunoglobulin-like domains and two fibronectin type III domains. Axl is overexpressed in a number of tumor cell types and was initially cloned from patients with chronic myelogenous leukemia. When overexpressed, Axl exhibits transforming potential. Axl signaling is believed to cause tumor growth through activation of proliferative and anti-apoptotic signaling pathways. Axl has been associated with cancers such as lung cancer, myeloid leukemia, uterine cancer, ovarian cancer, gliomas, melanoma, thyroid cancer, renal cell carcinoma, osteosarcoma, gastric cancer, prostate cancer, and breast cancer. The over-expression of Axl results in a poor prognosis for patients with the indicated cancers.


[0004] Activation of Mer, like Axl, conveys downstream signaling pathways that cause tumor growth and activation. Mer binds ligands such as the soluble protein Gas-6. Gas-6 binding to Mer induces autophosphorylation of Mer on its intracellular domain, resulting in downstream signal activation. Over-expression of Mer in cancer cells leads to increased metastasis, most likely by generation of soluble Mer extracellular domain protein as a decoy

receptor. Tumor cells secrete a soluble form of the extracellular Mer receptor which reduces the ability of soluble Gas-6 ligand to activate Mer on endothelial cells, leading to cancer progression.

[0005] Therefore a need exists for compounds that inhibit TAM receptor tyrosine kinases such as Axl and Mer for the treatment of selected cancers.

Summary of the Invention

[0006] In one aspect, the present invention provides a compound of formula I:



or a pharmaceutically acceptable salt thereof, wherein:

R_1 is selected from the group consisting of -H, -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₁-C₆) alkyl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C₁-C₆) alkyl;

wherein when R_1 is selected from the group consisting of -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C₁-C₆) alkyl, R_2 is -H, halo, -NR₅R₆, or optionally substituted (C₁-C₆) alkoxy;


wherein when R_1 is -H, optionally substituted (C₁-C₆) alkyl, or optionally substituted (C₁-C₆) alkoxy, R_2 is -CO-NR₅R₆; or -CO₂R₇;

or R_1 and R_2 taken together with the atoms to which they are attached to form optionally substituted cycloalkyl or optionally substituted heterocycloalkyl;

R_3 is selected from the group consisting of -H, optionally substituted (C₁-C₆) alkyl, -CN, and halo;

R_4 is -H or halo;



is optionally substituted with one, two, three, or four groups independently selected from the group consisting of halo and (C₁-C₆) alkyl, wherein “” indicate points of attachment;

R₅ and R₆ are each independently -H, optionally substituted (C₁-C₆) alkyl, or optionally substituted (C₁-C₆) alkoxy;


R₇ is -H or optionally substituted (C₁-C₆) alkyl

R₈ and R₉ are each independently -H or optionally substituted (C₁-C₆) alkyl;

or R₈ and R₉ may connect to form an optionally substituted heterocycle; and

Y is selected from the group consisting of O, S, SO, SO₂, NH, and N-(C₁-C₆ alkyl).

[0007] Another aspect provides a compound of formula A:



A

or a pharmaceutically acceptable salt thereof, wherein

(i) R₁ is selected from the group consisting of (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, -CN, -NHOH, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)NR^aR^a, C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -S(O)NR^aR^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and S(O)₂NR^aR^a; and

R₂ is selected from -H, halo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NO₂, -OR^a, -SR^a, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)R^a, -OC(O)NR^aR^a, -NHR^a, -NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -

NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and -S(O)₂NR^aR^a, wherein the (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R₁ or R₂ are each optionally substituted with 1, 2, 3, 4, or 5 independently selected R^b substituents, provided when R₁ is 5-7 membered heteroaryl or 5-7 membered heterocycloalkyl and R₂ is C₁₋₆ alkoxy, then the 5-7 membered heteroaryl or 5-7 membered heterocycloalkyl does not connect to the fused phenyl ring of the quinoline moiety through a ring nitrogen atom; or

(ii) R₁ is selected from -H, halo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NO₂, -OR^a, -SR^a, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)R^a, -OC(O)NR^aR^a, -NHR^a, -NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and -S(O)₂NR^aR^a, wherein the (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R₁ or R₂ are each optionally substituted with 1, 2, 3, 4, or 5 independently selected R^b substituents; and

R₂ is selected from the group consisting of (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -CN, -NHOH, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -S(O)NR^aR^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and S(O)₂NR^aR^a, provided when R₁ is 5-7 membered heteroaryl or 5-7 membered heterocycloalkyl and R₂ is C₁₋₆ alkoxy, then the 5-7 membered heteroaryl or 5-7 membered

heterocycloalkyl of R₁ does not connect to the fused phenyl ring of the quinoline moiety through a ring nitrogen atom,

(iii) R₁ and R₂ taken together with the atoms to which they are attached form a fused (C₃-C₇) cycloalkyl ring or a fused 4- to 10-membered heterocycloalkyl ring, wherein the fused (C₃-C₇) cycloalkyl ring and fused 4- to 10-membered heterocycloalkyl ring are each optionally substituted with 1, 2, or 3 independently selected R^b substituents, provided that the compound is not 1-[2-(4-Fluoro-phenyl)-acetyl]-cyclopropanecarboxylic acid [3-fluoro-4-(7,8,10,11,13,14-hexahydro-6,9,12,15-tetraoxa-1-aza-cyclododeca[b]naphthalen-4-yloxy)-phenyl]-amide;

R₁₀ and R₁₁ are each independently selected from the group consisting of -H, halo, (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NO₂, -OR^a, -SR^a, -NHR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)R^a, -OC(O)NR^aR^a, -NHR^a, -NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and S(O)₂NR^aR^a, wherein the (C₁-C₆) alkyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R₁ or R₂ are each optionally substituted with 1, 2, 3, 4, or 5 independently selected R^b substituents;

each R₃ is independently selected from the group consisting of -H, halo, -OH, -CN, optionally substituted (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkoxy, -NH₂, -NH(C₁-C₆)alkyl, -N(C₁-C₆ alkyl)₂, and (C₃-C₆) cycloalkyl, wherein the (C₁-C₆) alkoxy, -NH(C₁-C₆)alkyl, -N(C₁-C₆ alkyl)₂, and (C₃-C₆) cycloalkyl of R₃ are each optionally substituted with 1, 2, or 3 independently selected R^g substituents;

each R₁₄ is independently selected from the group consisting of halo, -OH, -NH₂, -CN, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, -COOH, -NH(C₁-C₆)alkyl, -N(C₁-C₆ alkyl)₂, phenyl, phenyl-(C₁-C₂) alkylene, (C₃-C₆) cycloalkyl, (C₃-C₆) cycloalkyl-(C₁-

C₄) alkylene-, 4- to 6-membered heterocycloalkyl, (4- to 6-membered heterocycloalkyl)-(C₁-C₄) alkylene-, 5- to 6-membered heteroaryl, (5- to 6-membered heteroaryl)-(C₁-C₄) alkylene-, and -OR^e, wherein the (C₁-C₆) alkyl, phenyl, phenyl-(C₁-C₂) alkylene, (C₃-C₆) cycloalkyl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, 4- to 6-membered heterocycloalkyl, (4- to 6-membered heterocycloalkyl)-(C₁-C₄) alkylene-, 5- to 6-membered heteroaryl, and (5- to 6-membered heteroaryl)-(C₁-C₄) alkylene- of R₁₄ are each optionally substituted with 1, 2, or 3 independently selected R^g substituents,

R₁₅ is H or C₁₋₆ alkyl;

each R₄ is independently selected from the group consisting of -H, halo, -OH, -COOR^e, -CONR^eR^e, -CN, -NH₂, -NH((C₁-C₆) alkyl), -N((C₁-C₆) alkyl)₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, -CONR^aR^a, -NR^aCOR^a, -NR^aCONR^aR^a, -SO₂R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, (C₃-C₆) cycloalkyl, 4- to 6-membered heterocycloalkyl, phenyl, 5- or 6-membered heteroaryl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, (4- to 6-membered heterocycloalkyl)-(C₁-C₄) alkylene-, phenyl-(C₁-C₂) alkylene, and (5- or 6-membered heteroaryl)-(C₁-C₄) alkylene-, wherein the (C₁-C₆) alkyl, (C₃-C₆) cycloalkyl, 4- to 6-membered heterocycloalkyl, phenyl, 5- or 6-membered heteroaryl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, (4- to 6-membered heterocycloalkyl)-(C₁-C₄) alkylene-, phenyl-(C₁-C₂) alkylene, and (5- or 6-membered heteroaryl)-(C₁-C₄) alkylene- of R₄ are each optionally substituted with 1, 2, or 3 independently selected R^f substituents;

each R^a is independently selected from the group consisting of -H, -CN, (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, wherein the (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R^a are each optionally substituted with 1, 2, 3, 4, or 5 independently selected R^d substituents;

each R^b is independently selected from the group consisting of halo, oxo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl, (C₃-C₁₀)

cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-10 membered heteroaryl)-(C₁-C₄) alkylene-, (4-10 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -OH, -NH₂, -NO₂, -NHOR^c, -OR^c, -SR^c, -C(O)R^c, -C(O)NR^cR^c, -C(O)OR^c, -C(O)NR^cS(O)₂R^c, -OC(O)R^c, -OC(O)NR^cR^c, -C(=NOH)R^c, -C(=NOH)NR^c, -C(=NCN)NR^cR^c, -NR^cC(=NCN)NR^cR^c, -C(=NR^c)NR^cR^c, -NR^cC(=NR^c)NR^cR^c, -NHR^c, -NR^cR^c, -NR^cC(O)R^c, -NR^cC(=NR^c)R^c, -NR^cC(O)OR^c, -NR^cC(O)NR^cR^c, -NR^cS(O)R^c, -NR^cS(O)₂R^c, -NR^cS(O)₂NR^cR^c, -S(O)R^c, -S(O)NR^cR^c, -S(O)₂R^c, -S(O)₂NR^cC(O)R^c, -Si(R^c)₃, -P(O)R^cR^c, -P(O)(OR^c)(OR^c), -B(OH)₂, -B(OR^c)₂, and -S(O)₂NR^cR^c, wherein the (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-10 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-10 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R^b are each further optionally substituted with 1, 2, or 3 independently selected R^d substituents;

each R^c is independently selected from the group consisting of -H, (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-10 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-10 membered heterocycloalkyl)-(C₁-C₄) alkylene-, wherein the (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-10 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-10 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R^c are each optionally substituted with 1, 2, 3, 4, or 5 independently selected R^f substituents;

each R^d is independently selected from the group consisting of (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, halo, (C₆-C₁₀) aryl, 5-10 membered heteroaryl, (C₃-C₁₀) cycloalkyl, 4-10 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-10 membered heteroaryl)-(C₁-C₄) alkylene-, (4-10 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NH₂, -NHOR^e, -OR^e, -SR^e, -C(O)R^e, -C(O)NR^eR^e, -C(O)OR^e, -OC(O)R^e, -OC(O)NR^eR^e, -NHR^e, -NR^eR^e, -NR^eC(O)R^e, -NR^eC(O)NR^eR^e, -NR^eC(O)OR^e, -C(=NR^e)NR^eR^e, -NR^eC(=NR^e)NR^eR^e, -NR^eC(=NOH)NR^eR^e, -NR^eC(=NCN)NR^eR^e, -S(O)R^e, -S(O)NR^eR^e, -S(O)₂R^e, -NR^eS(O)₂R^e, -NR^eS(O)₂NR^eR^e, and -S(O)₂NR^eR^e, wherein the (C₁-C₆) alkyl, (C₁-C₆)

haloalkyl, (C₆-C₁₀) aryl, 5-10 membered heteroaryl, (C₃-C₁₀) cycloalkyl, 4-10 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-10 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-10 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R^d are each optionally substituted with 1, 2, or 3 independently selected R^f substituents;

each R^e is independently selected from the group consisting of -H, (C₁-C₆) alkyl, (C₃-C₆) cycloalkyl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, (C₆-C₁₀) aryl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, 5- or 6-membered heteroaryl, (5- or 6-membered heteroaryl)-(C₁-C₄) alkylene-, 4-7-membered heterocycloalkyl, (4-7-membered heterocycloalkyl)-(C₁-C₄) alkylene-, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₂-C₄) alkenyl, and (C₂-C₄) alkynyl, wherein the (C₁-C₄) alkyl, (C₃-C₆) cycloalkyl, (C₆-C₁₀) aryl, 5 or 6-membered heteroaryl, 4-7-membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (5- or 6-membered heteroaryl)-(C₁-C₄) alkylene-, (4-7-membered heterocycloalkyl)-(C₁-C₄) alkylene-, (C₂-C₄) alkenyl, and (C₂-C₄) alkynyl of R^e are each optionally substituted with 1, 2, or 3 R^f substituents,

or any two R^a substituents together with the nitrogen atom to which they are attached form 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered heterocycloalkyl, each of which is optionally substituted with 1, 2, or 3 independently selected R^f substituents;

or any two R^c substituents together with the nitrogen atom to which they are attached form 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered heterocycloalkyl, each of which is optionally substituted with 1, 2, or 3 independently selected R^f substituents,

or any two R^e substituents together with the nitrogen atom to which they are attached form 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered heterocycloalkyl, each of which is optionally substituted with 1, 2, or 3 independently selected R^f substituents;

each R^f is independently selected from the group consisting of halo, -OH, -CN, -COOH, -NH₂, -NH-(C₁-C₆) alkyl, -N((C₁-C₆) alky)₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) alkylthio, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, phenyl, 5-6 membered heteroaryl, 4-6 membered heterocycloalkyl, and (C₃-C₆) cycloalkyl, wherein the (C₁-C₆) alkyl, phenyl, (C₃-C₆) cycloalkyl, 4-6 membered heterocycloalkyl, and 5-6 membered heteroaryl of R^f are each optionally substituted with 1, 2, or 3 substituents selected from halo, -OH, -CN, -COOH, -NH₂, (C₁-C₄) alkyl, (C₁-C₄) alkoxy, (C₁-C₄) haloalkyl, (C₁-C₄) haloalkoxy, phenyl, (C₃-C₁₀) cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl;

each R^g is independently selected from the group consisting of halo, -OH, -CN, -COOH, -COO-(C₁-C₄) alkyl, -NH₂, -NH-(C₁-C₆) alkyl, -N((C₁-C₆) alky)₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) alkylthio, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, phenyl, 5-6 membered heteroaryl, 4-6 membered heterocycloalkyl, and (C₃-C₆) cycloalkyl;

Y is selected from -O-, -S-, -SO-, -SO₂-, -NH-, and -N((C₁-C₆) alkyl)-;

the ring nitrogen atom on the quinoline moiety in Formula A is optionally oxidized;

the subscript n is an integer of 1, 2, 3, or 4;

the subscript m is an integer of 1, 2, 3, 4, or 5; and

the subscript p is an integer of 0, 1, 2, 3, or 4.

[0008] Another aspect provides methods of using compounds of formula I or a pharmaceutically acceptable salt thereof for the treatment of a disease, disorder, or syndrome mediated at least in part by modulating in vivo activity of a protein kinase.

[0009] A further aspect provides processes for making compounds of formula A and of formula I.

[00010] These and other aspects and embodiments are described below.

Detailed Description of the Invention

Abbreviations and Definitions

[00011] The following abbreviations and terms have the indicated meanings throughout:

Abbreviation	Meaning
Ac	Acetyl
anhyd	Anhydrous
Aq	Aqueous
Ar	Argon
Boc	Tert-butoxycarbonyl
Br	Broad
°C	Degrees Celsius
c-	Cyclo
calcd	Calculated
CBZ	CarboBenZoxy = benzyloxycarbonyl
d	Doublet
dd	Doublet of doublets
ddd	Doublet of doublets of doublets
dt	Doublet of triplets
DCM	Dichloromethane


Abbreviation	Meaning
DMF	<i>N,N</i> -Dimethylformamide
DMSO	Dimethyl sulfoxide
Dppf	1,1'-bis(diphenylphosphano)ferrocene
EA	Elemental Analysis
EI	Electron Impact ionization
eq or equiv	Equivalent
Fmoc	Fluorenylmethyloxycarbonyl
g	Gram(s)
h or hr	Hour(s)
HPLC	High pressure liquid chromatography
H ₂	Hydrogen
L	Liter(s)
LiHMDS	Lithium bis(trimethylsilyl)azide
M	Molar or molarity
m	Multiplet
MHz	Megahertz (frequency)
Min	Minute(s)
mL	Milliliter(s)
Mp	Melting point
m/z	Mass to charge ratio
μL	Microliter(s)
Mol	Mole(s)
MS	Mass spectral analysis
N ₂	Nitrogen
N	Normal or normality
nM	Nanomolar
NMR	Nuclear magnetic resonance spectroscopy
Pd/C	Palladium on carbon
Q	Quartet
RT	Room temperature
s	Singlet
soln	Solution
S/C	Substrate/catalyst ratio
t or tr	Triplet
THF	Tetrahydrofuran
TLC	Thin layer chromatography
v/v	Volume to volume

[00012] The symbol “-” means a single bond, and “=” means a double bond.


[00013] As used herein, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise.

[00014] When a variable is defined generically, with a number of possible substituents, each individual radical can be defined with or without the bond. For example, if R^Z can be hydrogen, this can be indicated as “-H” or “H” in the definition of R^Z .

[00015] When chemical structures are depicted or described, unless explicitly stated otherwise, all carbons are assumed to have hydrogen substitution to conform to a valence of four. For example, in the structure on the left-hand side of the schematic below, there are nine hydrogens implied. The nine hydrogens are depicted in the right-hand structure. Sometimes a particular atom in a structure is described in textual formula as having a hydrogen or hydrogens as substitution (expressly defined hydrogen), for example, $-\text{CH}_2\text{CH}_2-$. It is understood by one of ordinary skill in the art that the aforementioned descriptive techniques are common in the chemical arts to provide brevity and simplicity to description of otherwise complex structures.




[00016] If a group “R” is depicted as “floating” on a ring system, as for example in the formula:




then, unless otherwise defined, a substituent “R” may reside on any atom of the ring system, assuming replacement of a depicted, implied, or expressly defined hydrogen from one of the ring atoms, so long as a stable structure is formed.

[00017] If a group “R” is depicted as floating on a fused ring system, as for example in the formulae:




then, unless otherwise defined, a substituent “R” may reside on any atom of the fused ring system, assuming replacement of a depicted hydrogen (for example the -NH- in the formula above), implied hydrogen (for example, in the formula above, where the hydrogens are not shown but understood to be present), or expressly defined hydrogen (for example, where in the

formula above, “Z” equals =CH-) from one of the ring atoms, so long as a stable structure is formed. In the example depicted, the “R” group may reside on either the 5-membered or the 6-membered ring of the fused ring system. When a group “R” is depicted as existing on a ring system containing saturated carbons, for example in the formula:



where, in this example, “y” can be more than one, assuming each replaces a currently depicted, implied, or expressly defined hydrogen on the ring; then, unless otherwise defined, where the resulting structure is stable, two “R’s” may reside on the same carbon. A simple example is when R is a methyl group, there can exist a geminal dimethyl on a carbon of the depicted ring (an “annular” carbon). In another example, two R’s on the same carbon, including that carbon, may form a ring, thus creating a spirocyclic ring (a “spirocyclyl” group) structure with the depicted ring as for example in the formula:



[00018] “Halogen” or “halo” refers to fluorine, chlorine, bromine, or iodine.

[00019] The term “C_{n-m}” or “C_n-C_m” indicates a range which includes the endpoints, wherein n and m are integers and indicate the number of carbons. Examples include C₁₋₄, C₁-C₄, C₁₋₆, C₁-C₆, and the like.

[00020] “Alkyl” refers to a branched or straight hydrocarbon chain of one to eight carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, pentyl, hexyl, and heptyl. (C₁-C₆)alkyl is preferred. The term “C_{n-m} alkyl” or (C_n-C_m) alkyl, refers to an alkyl group having n to m carbon atoms. When optionally substituted, one or more hydrogen atoms of the alkyl group (e.g., from 1 to 4, from 1 to 2, or 1) may be replaced with a moiety as described below under “Optional Substitution.” In some aspects, the alkyl group is unsubstituted or not optionally substituted.

[00021] “Alkylene” refers to an optionally substituted bivalent saturated aliphatic radical having from 1 to 10 carbon atoms, 1 to 8 carbon atoms, 1 to 6 carbon atoms, 1 to 4 carbon atoms, or 1 to 2 carbon atoms. When optionally substituted, one or more hydrogen atoms of the alkylene group (e.g., from 1 to 4, from 1 to 2, or 1) may be replaced with a moiety as described below under “Optional Substitution.” In some aspects, the alkylene group is unsubstituted or not

optionally substituted. The term "C_{n-m} alkylene" refers to an alkylene group having n to m carbon atoms. Examples of alkylene groups include, but are not limited to, methylene, ethan-1,2-diyl, propan-1,3-diyl, propan-1,2-diyl, butan-1,4-diyl, butan-1,3-diyl, butan-1,2-diyl, 2-methyl-propan-1,3-diyl and the like.

[00022] The term "alkenyl" refers to a straight-chain or branched hydrocarbon group corresponding to an alkyl group having one or more double carbon-carbon bonds. An alkenyl group formally corresponds to an alkene with one C-H bond replaced by the point of attachment of the alkenyl group to the remainder of the compound. The term "C_{n-m} alkenyl" or (C_n-C_m) alkenyl refers to an alkenyl group having n to m carbons. In some embodiments, the alkenyl moiety contains 2 to 6, 2 to 4, or 2 to 3 carbon atoms. Example alkenyl groups include, but are not limited to, ethenyl, *n*-propenyl, isopropenyl, *n*-butenyl, *sec*-butenyl, and the like.

[00023] The term "alkynyl" refers to a straight-chain or branched hydrocarbon group corresponding to an alkyl group having one or more triple carbon-carbon bonds. An alkynyl group formally corresponds to an alkyne with one C-H bond replaced by the point of attachment of the alkyl group to the remainder of the compound. The term "C_{n-m} alkynyl" or (C_n-C_m) alkynyl refers to an alkynyl group having n to m carbons. Example alkynyl groups include, but are not limited to, ethynyl, propyn-1-yl, propyn-2-yl, and the like. In some embodiments, the alkynyl moiety contains 2 to 6, 2 to 4, or 2 to 3 carbon atoms.

[00024] "Alkoxy" refers to a moiety of the formula -OR', wherein R' is an (C₁-C₆)alkyl moiety as defined herein. The term "C_{n-m} alkoxy" or (C_n-C_m) alkoxy refers to an alkoxy group, the alkyl group of which has n to m carbons. Examples of alkoxy moieties include, but are not limited to, methoxy, ethoxy, isopropoxy, and the like.

[00025] An alkoxy group can be unsubstituted or optionally substituted. When optionally substituted, one or more hydrogen atoms of the alkoxy group (e.g., from 1 to 4, from 1 to 2, or 1) may be replaced with a moiety as described below under "Optional Substitution," with the proviso that no hydrogen atom alpha to the ether oxygen is replaced by a hydroxy, amino, or thio group. In some aspects, the alkoxy group is unsubstituted or not optionally substituted.

[00026] "Alkoxycarbonyl" refers to a group -C(O)-R' wherein R' is (C₁-C₆)alkoxy as defined herein.

[00027] The term "amino" refers to a group of formula -NH₂.

[00028] The term "carbamyl" refers to a group of formula -C(O)NH₂.

[00029] The term “carbonyl”, employed alone or in combination with other terms, refers to a -C(=O)- group, which also may be written as C(O).

[00030] The term “cyano” or “nitrile” refers to a group of formula $\text{-C}\equiv\text{N}$, which also may be written as -CN or CN.

[00031] The term “oxo” refers to an oxygen atom as a divalent substituent, forming a carbonyl group when attached to carbon, or attached to a heteroatom forming a sulfoxide or sulfone group, or an *N*-oxide group. In some embodiments, heterocyclic groups may be optionally substituted by 1 or 2 oxo (=O) substituents.

[00032] The term “sulfide” refers to a sulfur atom as a divalent substituent, forming a thiocarbonyl group (C=S) when attached to carbon.

[00033] The term “heteroatom” used herein is meant to include boron, phosphorus, sulfur, oxygen, and nitrogen.

[00034] The term “haloalkyl” as used herein refers to an alkyl group in which one or more of the hydrogen atoms has been replaced by a halogen atom. The term “C_{n-m} haloalkyl” or (C_n-C_m) haloalkyl refers to a C_{n-m} alkyl group having n to m carbon atoms and from at least one up to {2(n to m)+1} halogen atoms, which may either be the same or different. In some embodiments, the halogen atoms are fluoro atoms. In some embodiments, the haloalkyl group has 1 to 6 or 1 to 4 carbon atoms. Example haloalkyl groups include CF₃, C₂F₅, CHF₂, CCl₃, CHCl₂, C₂Cl₅, and the like. In some embodiments, the haloalkyl group is a fluoroalkyl group.

[00035] The term “haloalkoxy,” employed alone or in combination with other terms, refers to a group of formula -O-haloalkyl, wherein the haloalkyl group is as defined above. The term “C_{n-m} haloalkoxy” or (C_n-C_m) haloalkoxy refers to a haloalkoxy group, the haloalkyl group of which has n to m carbons. Example haloalkoxy groups include trifluoromethoxy and the like. In some embodiments, the haloalkoxy group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

[00036] “Aryl” means a monovalent six- to fourteen-membered, mono- or bi-carbocyclic ring (e.g., having two fused rings), wherein the monocyclic ring is aromatic and at least one of the rings in the bicyclic ring is aromatic. The term “C_{n-m} aryl” or “(C_n-C_m) aryl” refers to an aryl group having from n to m ring carbon atoms. In some embodiments, aryl groups have from 6 to about 10 carbon atoms. In some embodiments aryl groups have 6 carbon atoms. In some embodiments aryl groups have 10 carbon atoms. Unless stated otherwise, the valency of the

group may be located on any atom of any ring within the radical, valency rules permitting. Representative examples include phenyl, naphthyl, and indanyl, and the like.

[00037] An aryl group can be unsubstituted or optionally substituted. When optionally substituted, one or more hydrogen atoms of the aryl group (e.g., from 1 to 5, from 1 to 2, or 1) may be replaced with a moiety as described below under “Optional Substitution.” In some aspects, the alkoxy group is unsubstituted or not optionally substituted.

[00038] “Arylene” means a divalent six- to fourteen-membered, mono- or bi-carbocyclic ring, wherein the monocyclic ring is aromatic and at least one of the rings in the bicyclic ring is aromatic. Representative examples include phenylene, naphthylene, and indanylene, and the like.

[00039] “Cycloalkyl” refers to a non-aromatic hydrocarbon ring system (monocyclic, bicyclic, or polycyclic), including cyclized alkyl and alkenyl groups. The term “C_{n-m} cycloalkyl” or “(C_n-C_m) cycloalkyl” refers to a cycloalkyl that has n to m ring member carbon atoms. Cycloalkyl groups can include mono- or polycyclic (e.g., having 2, 3, or 4 fused rings) groups and spirocycles. Cycloalkyl groups can have 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14 ring-forming carbons (C₃₋₁₄). In some embodiments, the cycloalkyl group has 3 to 14 members, 3 to 10 members, 3 to 6 ring members, 3 to 5 ring members, or 3 to 4 ring members. In some embodiments, the cycloalkyl group is monocyclic. In some embodiments, the cycloalkyl group is monocyclic or bicyclic. In some embodiments, the cycloalkyl group is a C₃₋₆ monocyclic cycloalkyl group. Ring-forming carbon atoms of a cycloalkyl group can be optionally oxidized to form an oxo or sulfido group. Cycloalkyl groups also include cycloalkylidenes. In some embodiments, cycloalkyl is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. Examples of cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptatrienyl, norbornyl, norpinyl, norcarnyl, bicyclo[1.1.1]pentanyl, bicyclo[2.1.1]hexanyl, and the like. In some embodiments, the cycloalkyl group is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In some embodiments, cycloalkyl includes a single saturated carbocyclic ring of three to eight ring carbons, such as cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. Cycloalkyl may optionally be substituted with one or more substituents, such as one, two, or three substituents. In some embodiments, the cycloalkyl substituent is selected from the group consisting of (C₁-

C₆)alkyl, hydroxy, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, halo, amino, mono- and di(C₁-C₆)alkylamino, hetero(C₁-C₆)alkyl, acyl, aryl, and heteroaryl.

[00040] A cycloalkyl group can be unsubstituted or optionally substituted. When optionally substituted, one or more hydrogen atoms of the cycloalkyl group (e.g., from 1 to 4, from 1 to 2, or 1) may be replaced with a moiety as described below under “Optional Substitution.” In some aspects, a substituted cycloalkyl group can incorporate an exo- or endocyclic alkene (e.g., cyclohex-2-en-1-yl). In some aspects, a cycloalkyl group is unsubstituted or not optionally substituted.

[00041] “Cycloalkyloxycarbonyl” means a group -C(O)-OR’ wherein R’ is (C₃-C₆)cycloalkyl as defined herein.

[00042] “Phenyloxycarbonyl” refers to a group -C(O)-Ophenyl.

[00043] “Heteroaryl” means a monocyclic, fused bicyclic, or fused tricyclic, monovalent radical of 5 to 14 ring atoms containing one or more, preferably one, two, three, or four ring heteroatoms independently selected from -O-, -S(O)_n- (n is 0, 1, or 2), -N-, and -N(R’)-, and the remaining ring atoms being carbon, wherein the ring comprising a monocyclic radical is aromatic and wherein at least one of the fused rings comprising a bicyclic or tricyclic radical is aromatic. One or two ring carbon atoms of any nonaromatic rings comprising a bicyclic or tricyclic radical may be replaced by a -C(O)-, -C(S)-, or -C(=NH)- group. R’ is hydrogen, alkyl, hydroxy, alkoxy, acyl, or alkylsulfonyl. Unless stated otherwise, the valency may be located on any atom of any ring of the heteroaryl group, valency rules permitting. In particular, when the point of valency is located on the nitrogen, an additional nitrogen substituent is not present. More specifically, the term heteroaryl includes, but is not limited to, 1,2,4-triazolyl, 1,3,5-triazolyl, phthalimidyl, pyridinyl, pyrrolyl, imidazolyl, thienyl, furanyl, indolyl, 2,3-dihydro-1*H*-indolyl (including, for example, 2,3-dihydro-1*H*-indol-2-yl or 2,3-dihydro-1*H*-indol-5-yl, and the like), isoindolyl, indolinyl, isoindolinyl, benzimidazolyl, benzodioxol-4-yl, benzofuranyl, cinnolinyl, indoliziny, naphthyridin-3-yl, phthalazin-3-yl, phthalazin-4-yl, pteridiny, purinyl, quinazolinyl, quinoxalinyl, tetrazoyl, pyrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, isooxazolyl, oxadiazolyl, benzoxazolyl, quinolinyl, isoquinolinyl, tetrahydroisoquinolinyl (including, for example, tetrahydroisoquinolin-4-yl or tetrahydroisoquinolin-6-yl, and the like), pyrrolo[3,2-*c*]pyridinyl (including, for example, pyrrolo[3,2-*c*]pyridin-2-yl or pyrrolo[3,2-*c*]pyridin-7-yl, and the like), benzopyranyl, thiazolyl,

isothiazolyl, thiadiazolyl, benzothiazolyl, benzothienyl, and the derivatives thereof, and N-oxide or a protected derivative thereof.

[00044] A five-membered heteroaryl ring is a heteroaryl group having five ring atoms wherein one or more (e.g., 1, 2, 3, or 4) ring atoms are independently selected from N, O, and S. Exemplary five-membered ring heteroaryls include thienyl, furyl, pyrrolyl, imidazolyl, thiazolyl, oxazolyl, pyrazolyl, isothiazolyl, isoxazolyl, 1,2,3-triazolyl, tetrazolyl, 1,2,3-thiadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-triazolyl, 1,2,4-thiadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-triazolyl, 1,3,4-thiadiazolyl, and 1,3,4-oxadiazolyl.

[00045] A six-membered heteroaryl ring is a heteroaryl group having six ring atoms wherein one or more (e.g., 1, 2, 3, or 4) ring atoms are independently selected from N, O, and S. Exemplary six-membered ring heteroaryls are pyridyl, pyrazinyl, pyrimidinyl, triazinyl, and pyridazinyl.

[00046] “Heteroarylene” means a monocyclic, fused bicyclic, or fused tricyclic, divalent radical of 5 to 14 ring atoms containing one or more, preferably one, two, three, or four ring heteroatoms independently selected from -O-, -S(O)_n- (n is 0, 1, or 2), -N-, and -N(R¹⁹)-, and the remaining ring atoms being carbon, wherein the ring comprising a monocyclic radical is aromatic and wherein at least one of the fused rings comprising a bicyclic or tricyclic radical is aromatic. One or two ring carbon atoms of any nonaromatic rings comprising a bicyclic or tricyclic radical may be replaced by a -C(O)-, -C(S)-, or -C(=NH)- group. R¹⁹ is hydrogen, alkyl, or alkenyl. Unless stated otherwise, the valencies may be located on any atom of any ring of the heteroarylene group, valency rules permitting. In particular, when the point of valency is located on the nitrogen, an additional nitrogen substituent is not present. More specifically, the term heteroaryl includes, but is not limited to, thien-diyl, benzo[*d*]isoxazol-diyl, benzo[*d*]isothiazol-diyl, 1*H*-indazol-diyl (optionally substituted at the N1 position with R¹⁹), benzo[*d*]oxazol-diyl, benzo[*d*]thiazol-diyl, 1*H*-benzo[*d*]imidazol-diyl (optionally substituted at the N1 position with R¹⁹), 1*H*-benzo[*d*][1,2,3]triazol-diyl (optionally substituted at the N1 position with R¹⁹), imidazo[1,2-*a*]pyridin-diyl, cinnolin-diyl, quinolin-diyl, pyridin-diyl, 1-oxido-pyridin-diyl, [1,2,4]triazolo[4,3-*a*]pyridin-diyl, and 2,3-dihydroimidazo[1,2-*a*]pyridin-diyl, and the like.

[00047] As used herein, “heterocycloalkyl” or “heterocyclo” refer to a non-aromatic ring or ring system, which may optionally contain one or more alkenylene groups as part of the ring

structure, which has at least one heteroatom ring member independently selected from boron, nitrogen, sulfur, oxygen, and phosphorus, and which has 4-14 ring members, 4-10 ring members, 4-7 ring members, or 4-6 ring members. Included within the term “heterocycloalkyl” are monocyclic 4-, 5-, 6-, and 7-membered heterocycloalkyl groups. Heterocycloalkyl groups can include mono- or bicyclic or polycyclic (e.g., having two or three fused or bridged rings) ring systems or spirorcycles. In some embodiments, the heterocycloalkyl group is a monocyclic group having 1, 2, or 3 heteroatoms independently selected from nitrogen, sulfur, and oxygen. Ring-forming carbon atoms and heteroatoms of a heterocycloalkyl group can be optionally oxidized to form an oxo or sulfido group or other oxidized linkage (e.g., C(O), S(O), C(S), S(O)₂, N-oxide, and the like.) or a nitrogen atom can be quaternized. The heterocycloalkyl group can be attached through a ring-forming carbon atom or a ring-forming heteroatom. In some embodiments, the heterocycloalkyl group contains 0 to 3 double bonds. In some embodiments, the heterocycloalkyl group contains 0 to 2 double bonds. Also included in the definition of heterocycloalkyl are moieties that have one or more aromatic rings fused (i.e., having a bond in common with) to the heterocycloalkyl ring, e.g., benzo or thienyl derivatives of piperidine, morpholine, azepine, and the like. A heterocycloalkyl group containing a fused aromatic ring can be attached through any ring-forming atom, including a ring-forming atom of the fused aromatic ring. Examples of heterocycloalkyl groups include azetidiny, azepanyl, dihydrobenzofuranyl, dihydrofuranyl, dihydropyranyl, morpholino, 3-oxa-9-azaspiro[5.5]undecanyl, 1-oxa-8-azaspiro[4.5]decanyl, piperidiny, piperaziny, oxopiperaziny, pyranyl, pyrrolidiny, quinuclidiny, tetrahydrofuranyl, tetrahydropyranyl, 1,2,3,4-tetrahydroquinoliny, tropanyl, 4,5,6,7-tetrahydrothiazolo[5,4-c]pyridiny, and thiomorpholino.

[00048] “Heterocycloalkyl” or “heterocyclo,” can be unsubstituted or optionally substituted. When optionally substituted, one or more hydrogen atoms of the group (e.g., from 1 to 4, from 1 to 2, or 1) may be replaced with a moiety independently selected from fluoro, hydroxy, alkoxy, amino, alkylamino, acylamino, thio, and alkylthio. In some aspects, a substituted heterocycyl group can incorporate an exo- or endocyclic alkene (e.g., cyclohex-2-en-1-yl). In some aspects, the heterocycyl group is unsubstituted or not optionally substituted.

Optional Substitution

[00049] A group is optionally substituted herein unless expressly provided otherwise. The term “optionally substituted” refers to being substituted or unsubstituted. In certain embodiments, alkyl, alkenyl, alkynyl, carbocycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups are optionally substituted. “Optionally substituted” refers to a group which may be substituted or unsubstituted (e.g., “substituted” or “unsubstituted” alkyl, “substituted” or “unsubstituted” alkenyl, “substituted” or “unsubstituted” alkynyl, “substituted” or “unsubstituted” cycloalkyl, “substituted” or “unsubstituted” heterocycloalkyl, “substituted” or “unsubstituted” aryl or “substituted” or “unsubstituted” heteroaryl group). In general, the term “substituted” means that at least one hydrogen present on a group is replaced with a permissible substituent, e.g., a substituent which upon substitution results in a stable compound, e.g., a compound which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, or other reaction. Unless otherwise indicated, a “substituted” group has a substituent at one or more substitutable positions of the group, and when more than one position in any given structure is substituted, the substituent is either the same or different at each position. The term “substituted” is contemplated to include substitution with all permissible substituents of organic compounds, and includes any of the substituents described herein that results in the formation of a stable compound. The present invention contemplates any and all such combinations in order to arrive at a stable compound. For purposes of this invention, heteroatoms such as nitrogen may have hydrogen substituents and/or any suitable substituent as described herein which satisfy the valencies of the heteroatoms and results in the formation of a stable moiety. The invention is not intended to be limited in any manner by the exemplary substituents described herein.

[00050] Exemplary carbon atom substituents include, but are not limited to, halogen (halo), $-\text{CN}$, $-\text{NO}_2$, $-\text{N}_3$, $-\text{SO}_2\text{H}$, $-\text{SO}_3\text{H}$, $-\text{OH}$, $-\text{OR}^{\text{aa}}$, $-\text{ON}(\text{R}^{\text{bb}})_2$, $-\text{N}(\text{R}^{\text{bb}})_2$, $-\text{N}(\text{R}^{\text{bb}})_3^+\text{X}^-$, $-\text{N}(\text{OR}^{\text{cc}})\text{R}^{\text{bb}}$, $-\text{SH}$, $-\text{SR}^{\text{aa}}$, $-\text{SSR}^{\text{cc}}$, $-\text{C}(=\text{O})\text{R}^{\text{aa}}$, $-\text{CO}_2\text{H}$, $-\text{CHO}$, $-\text{C}(\text{OR}^{\text{cc}})_2$, $-\text{CO}_2\text{R}^{\text{aa}}$, $-\text{OC}(=\text{O})\text{R}^{\text{aa}}$, $-\text{OCO}_2\text{R}^{\text{aa}}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{OC}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{NR}^{\text{bb}}\text{C}(=\text{O})\text{R}^{\text{aa}}$, $-\text{NR}^{\text{bb}}\text{CO}_2\text{R}^{\text{aa}}$, $-\text{NR}^{\text{bb}}\text{C}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{C}(=\text{NR}^{\text{bb}})\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{bb}})\text{OR}^{\text{aa}}$, $-\text{OC}(=\text{NR}^{\text{bb}})\text{R}^{\text{aa}}$, $-\text{OC}(=\text{NR}^{\text{bb}})\text{OR}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{OC}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{NR}^{\text{bb}}\text{C}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{C}(=\text{O})\text{NR}^{\text{bb}}\text{SO}_2\text{R}^{\text{aa}}$, $-\text{NR}^{\text{bb}}\text{SO}_2\text{R}^{\text{aa}}$, $-\text{SO}_2\text{N}(\text{R}^{\text{bb}})_2$, $-\text{SO}_2\text{R}^{\text{aa}}$, $-\text{SO}_2\text{OR}^{\text{aa}}$, $-\text{OSO}_2\text{R}^{\text{aa}}$, $-\text{S}(=\text{O})\text{R}^{\text{aa}}$, $-\text{OS}(=\text{O})\text{R}^{\text{aa}}$, $-\text{Si}(\text{R}^{\text{aa}})_3$, $-\text{OSi}(\text{R}^{\text{aa}})_3$, $-\text{C}(=\text{S})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{C}(=\text{O})\text{SR}^{\text{aa}}$, $-\text{C}(=\text{S})\text{SR}^{\text{aa}}$, $-\text{SC}(=\text{S})\text{SR}^{\text{aa}}$, $-\text{SC}(=\text{O})\text{SR}^{\text{aa}}$,

$-\text{OC}(=\text{O})\text{SR}^{\text{aa}}$, $-\text{SC}(=\text{O})\text{OR}^{\text{aa}}$, $-\text{SC}(=\text{O})\text{R}^{\text{aa}}$, $-\text{P}(=\text{O})_2\text{R}^{\text{aa}}$, $-\text{OP}(=\text{O})_2\text{R}^{\text{aa}}$, $-\text{P}(=\text{O})(\text{R}^{\text{aa}})_2$,
 $-\text{OP}(=\text{O})(\text{R}^{\text{aa}})_2$, $-\text{OP}(=\text{O})(\text{OR}^{\text{cc}})_2$, $-\text{P}(=\text{O})_2\text{N}(\text{R}^{\text{bb}})_2$, $-\text{OP}(=\text{O})_2\text{N}(\text{R}^{\text{bb}})_2$, $-\text{P}(=\text{O})(\text{NR}^{\text{bb}})_2$,
 $-\text{OP}(=\text{O})(\text{NR}^{\text{bb}})_2$, $-\text{NR}^{\text{bb}}\text{P}(=\text{O})(\text{OR}^{\text{cc}})_2$, $-\text{NR}^{\text{bb}}\text{P}(=\text{O})(\text{NR}^{\text{bb}})_2$, $-\text{OP}(\text{R}^{\text{cc}})_2$, $-\text{OP}(\text{R}^{\text{cc}})_3$, $-\text{B}(\text{OR}^{\text{cc}})_2$,
 $-\text{BR}^{\text{aa}}(\text{OR}^{\text{cc}})$, C_{1-10} alkyl, C_{1-10} perhaloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, $(\text{C}_3-\text{C}_{10})$
carbocycloalkyl, 3-14 membered heterocycloalkyl, $(\text{C}_6-\text{C}_{14})$ aryl, and 5-14 membered
heteroaryl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and
heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

or two geminal hydrogens on a carbon atom are replaced with the group $=\text{O}$, $=\text{S}$,
 $=\text{NN}(\text{R}^{\text{bb}})_2$, $=\text{NNR}^{\text{bb}}\text{C}(=\text{O})\text{R}^{\text{aa}}$, $=\text{NNR}^{\text{bb}}\text{C}(=\text{O})\text{OR}^{\text{aa}}$, $=\text{NNR}^{\text{bb}}\text{S}(=\text{O})_2\text{R}^{\text{aa}}$, $=\text{NR}^{\text{bb}}$, or $=\text{NOR}^{\text{cc}}$;

each instance of R^{aa} is, independently, selected from $(\text{C}_1-\text{C}_{10})$ alkyl, $(\text{C}_1-\text{C}_{10})$
perhaloalkyl, $(\text{C}_2-\text{C}_{10})$ alkenyl, $(\text{C}_2-\text{C}_{10})$ alkynyl, $(\text{C}_3-\text{C}_{10})$ cycloalkyl, 3-14 membered
heterocycloalkyl, $(\text{C}_6-\text{C}_{14})$ aryl, and 5-14 membered heteroaryl, or two R^{aa} groups are joined to
form a 3-14 membered heterocycloalkyl or 5-14 membered heteroaryl ring, wherein each alkyl,
alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently substituted
with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

each instance of R^{bb} is, independently, selected from hydrogen, $(\text{C}_1-\text{C}_{10})$ perhaloalkyl,
 $(\text{C}_2-\text{C}_{10})$ alkenyl, $(\text{C}_2-\text{C}_{10})$ alkynyl, $(\text{C}_3-\text{C}_{10})$ cycloalkyl, C_{6-14} aryl, and 5-14 membered
heteroaryl, or two R^{bb} groups are joined to form a 3-14 membered heterocycloalkyl or 5-14
membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl,
aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

each instance of R^{cc} is, independently, selected from hydrogen, $(\text{C}_1-\text{C}_{10})$ alkyl, $(\text{C}_1-\text{C}_{10})$
perhaloalkyl, $(\text{C}_2-\text{C}_{10})$ alkenyl, $(\text{C}_2-\text{C}_{10})$ alkynyl, $(\text{C}_3-\text{C}_{10})$ cycloalkyl, 3-14 membered
heterocycloalkyl, $(\text{C}_6-\text{C}_{14})$ aryl, and 5-14 membered heteroaryl, or two R^{cc} groups are joined to
form a 3-14 membered heterocycloalkyl or 5-14 membered heteroaryl ring, wherein each alkyl,
alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently substituted
with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

each instance of R^{dd} is, independently, selected from halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{SO}_2\text{H}$,
 $-\text{SO}_3\text{H}$, $-\text{OH}$, $-\text{OR}^{\text{ee}}$, $-\text{ON}(\text{R}^{\text{ff}})_2$, $-\text{N}(\text{R}^{\text{ff}})_2$, $-\text{N}(\text{R}^{\text{ff}})_3^+\text{X}^-$, $-\text{N}(\text{OR}^{\text{ee}})\text{R}^{\text{ff}}$, $-\text{SH}$, $-\text{SR}^{\text{ee}}$, $-\text{SSR}^{\text{ee}}$,
 $-\text{C}(=\text{O})\text{R}^{\text{ee}}$, $-\text{CO}_2\text{H}$, $-\text{CO}_2\text{R}^{\text{ee}}$, $-\text{OC}(=\text{O})\text{R}^{\text{ee}}$, $-\text{OCO}_2\text{R}^{\text{ee}}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{\text{ff}})_2$, $-\text{OC}(=\text{O})\text{N}(\text{R}^{\text{ff}})_2$,
 $-\text{NR}^{\text{ff}}\text{C}(=\text{O})\text{R}^{\text{ee}}$, $-\text{NR}^{\text{ff}}\text{CO}_2\text{R}^{\text{ee}}$, $-\text{NR}^{\text{ff}}\text{C}(=\text{O})\text{N}(\text{R}^{\text{ff}})_2$, $-\text{C}(=\text{NR}^{\text{ff}})\text{OR}^{\text{ee}}$, $-\text{OC}(=\text{NR}^{\text{ff}})\text{R}^{\text{ee}}$,
 $-\text{OC}(=\text{NR}^{\text{ff}})\text{OR}^{\text{ee}}$, $-\text{C}(=\text{NR}^{\text{ff}})\text{N}(\text{R}^{\text{ff}})_2$, $-\text{OC}(=\text{NR}^{\text{ff}})\text{N}(\text{R}^{\text{ff}})_2$, $-\text{NR}^{\text{ff}}\text{C}(=\text{NR}^{\text{ff}})\text{N}(\text{R}^{\text{ff}})_2$, $-\text{NR}^{\text{ff}}\text{SO}_2\text{R}^{\text{ee}}$,

$-\text{SO}_2\text{N}(\text{R}^{\text{ff}})_2$, $-\text{SO}_2\text{R}^{\text{ee}}$, $-\text{SO}_2\text{OR}^{\text{ee}}$, $-\text{OSO}_2\text{R}^{\text{ee}}$, $-\text{S}(=\text{O})\text{R}^{\text{ee}}$, $-\text{Si}(\text{R}^{\text{ee}})_3$, $-\text{OSi}(\text{R}^{\text{ee}})_3$,
 $-\text{C}(=\text{S})\text{N}(\text{R}^{\text{ff}})_2$, $-\text{C}(=\text{O})\text{SR}^{\text{ee}}$, $-\text{C}(=\text{S})\text{SR}^{\text{ee}}$, $-\text{SC}(=\text{S})\text{SR}^{\text{ee}}$, $-\text{P}(=\text{O})_2\text{R}^{\text{ee}}$, $-\text{P}(=\text{O})(\text{R}^{\text{ee}})_2$,
 $-\text{OP}(=\text{O})(\text{R}^{\text{ee}})_2$, $-\text{OP}(=\text{O})(\text{OR}^{\text{ee}})_2$, $(\text{C}_1\text{-C}_{10})$ alkyl, $(\text{C}_1\text{-C}_{10})$ perhaloalkyl, $(\text{C}_2\text{-C}_{10})$ alkenyl, $(\text{C}_2\text{-C}_{10})$ alkynyl, $(\text{C}_3\text{-C}_{10})$ cycloalkyl, 3-10 membered heterocycloalkyl, $(\text{C}_6\text{-C}_{10})$ aryl, 5-10 membered heteroaryl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{gg} groups, or two geminal R^{dd} substituents can be joined to form $=\text{O}$ or $=\text{S}$;

each instance of R^{ee} is, independently, selected from $(\text{C}_1\text{-C}_6)$ alkyl, $(\text{C}_1\text{-C}_6)$ perhaloalkyl, $(\text{C}_2\text{-C}_6)$ alkenyl, $(\text{C}_2\text{-C}_6)$ alkynyl, $(\text{C}_3\text{-C}_{10})$ cycloalkyl, $(\text{C}_6\text{-C}_{10})$ aryl, 3-10 membered heterocycloalkyl, and 3-10 membered heteroaryl, wherein each alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{gg} groups;

each instance of R^{ff} is, independently, selected from hydrogen, $(\text{C}_1\text{-C}_6)$ alkyl, $(\text{C}_1\text{-C}_6)$ perhaloalkyl, $(\text{C}_2\text{-C}_6)$ alkenyl, $(\text{C}_2\text{-C}_6)$ alkynyl, $(\text{C}_3\text{-C}_{10})$ cycloalkyl, $(\text{C}_6\text{-C}_{10})$ aryl, and 5-10 membered heteroaryl, or two R^{ff} groups are joined to form a 3-10 membered heterocycloalkyl or 5-10 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{gg} groups; and

each instance of R^{gg} is, independently, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{SO}_2\text{H}$, $-\text{SO}_3\text{H}$, $-\text{OH}$, $-\text{OC}_{1-6}$ alkyl, $-\text{ON}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{N}(\text{C}_{1-6}$ alkyl) $_3^+\text{X}^-$, $-\text{NH}(\text{C}_{1-6}$ alkyl) $_2^+\text{X}^-$, $-\text{NH}_2(\text{C}_{1-6}$ alkyl) $^+\text{X}^-$, $-\text{NH}_3^+\text{X}^-$, $-\text{N}(\text{OC}_{1-6}$ alkyl)(C_{1-6} alkyl), $-\text{N}(\text{OH})(\text{C}_{1-6}$ alkyl), $-\text{NH}(\text{OH})$, $-\text{SH}$, $-\text{SC}_{1-6}$ alkyl, $-\text{SS}(\text{C}_{1-6}$ alkyl), $-\text{C}(=\text{O})(\text{C}_{1-6}$ alkyl), $-\text{CO}_2\text{H}$, $-\text{CO}_2(\text{C}_{1-6}$ alkyl), $-\text{OC}(=\text{O})(\text{C}_{1-6}$ alkyl), $-\text{OCO}_2(\text{C}_{1-6}$ alkyl), $-\text{C}(=\text{O})\text{NH}_2$, $-\text{C}(=\text{O})\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{OC}(=\text{O})\text{NH}(\text{C}_{1-6}$ alkyl), $-\text{NHC}(=\text{O})(\text{C}_{1-6}$ alkyl), $-\text{N}(\text{C}_{1-6}$ alkyl) $\text{C}(=\text{O})(\text{C}_{1-6}$ alkyl), $-\text{NHCO}_2(\text{C}_{1-6}$ alkyl), $-\text{NHC}(=\text{O})\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{NHC}(=\text{O})\text{NH}(\text{C}_{1-6}$ alkyl), $-\text{NHC}(=\text{O})\text{NH}_2$, $-\text{C}(=\text{NH})\text{O}(\text{C}_{1-6}$ alkyl), $-\text{OC}(=\text{NH})(\text{C}_{1-6}$ alkyl), $-\text{OC}(=\text{NH})\text{OC}_{1-6}$ alkyl, $-\text{C}(=\text{NH})\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{C}(=\text{NH})\text{NH}(\text{C}_{1-6}$ alkyl), $-\text{C}(=\text{NH})\text{NH}_2$, $-\text{OC}(=\text{NH})\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{OC}(\text{NH})\text{NH}(\text{C}_{1-6}$ alkyl), $-\text{OC}(\text{NH})\text{NH}_2$, $-\text{NHC}(\text{NH})\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{NHC}(=\text{NH})\text{NH}_2$, $-\text{NHSO}_2(\text{C}_{1-6}$ alkyl), $-\text{SO}_2\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{SO}_2\text{NH}(\text{C}_{1-6}$ alkyl), $-\text{SO}_2\text{NH}_2$, $-\text{SO}_2\text{C}_{1-6}$ alkyl, $-\text{SO}_2\text{OC}_{1-6}$ alkyl, $-\text{OSO}_2\text{C}_{1-6}$ alkyl, $-\text{SOC}_{1-6}$ alkyl, $-\text{Si}(\text{C}_{1-6}$ alkyl) $_3$, $-\text{OSi}(\text{C}_{1-6}$ alkyl) $_3$, $-\text{C}(=\text{S})\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $\text{C}(=\text{S})\text{NH}(\text{C}_{1-6}$ alkyl), $\text{C}(=\text{S})\text{NH}_2$, $-\text{C}(=\text{O})\text{S}(\text{C}_{1-6}$ alkyl), $-\text{C}(=\text{S})\text{SC}_{1-6}$ alkyl, $-\text{SC}(=\text{S})\text{SC}_{1-6}$ alkyl, $-\text{P}(=\text{O})_2(\text{C}_{1-6}$ alkyl),

$-\text{P}(=\text{O})(\text{C}_{1-6} \text{ alkyl})_2$, $-\text{OP}(=\text{O})(\text{C}_{1-6} \text{ alkyl})_2$, $-\text{OP}(=\text{O})(\text{OC}_{1-6} \text{ alkyl})_2$, $(\text{C}_1\text{-C}_6)$ alkyl, $(\text{C}_1\text{-C}_6)$ perhaloalkyl, $(\text{C}_2\text{-C}_6)$ alkenyl, $(\text{C}_2\text{-C}_6)$ alkynyl, $(\text{C}_3\text{-C}_{10})$ cycloalkyl, $(\text{C}_6\text{-C}_{10})$ aryl, 3-10 membered heterocycloalkyl, 5-10 membered heteroaryl; or two geminal R^{gg} substituents can be joined to form $=\text{O}$ or $=\text{S}$; wherein X^- is a counterion.

[00051] As noted previously, nitrogen atoms can be substituted or unsubstituted as valency permits, and include primary, secondary, tertiary, and quaternary nitrogen atoms. Exemplary nitrogen atom substituents include, but are not limited to, hydrogen, $-\text{OH}$, $-\text{OR}^{\text{aa}}$, $-\text{N}(\text{R}^{\text{cc}})_2$, $-\text{CN}$, $-\text{C}(=\text{O})\text{R}^{\text{aa}}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{CO}_2\text{R}^{\text{aa}}$, $-\text{SO}_2\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{bb}})\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{cc}})\text{OR}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{cc}})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{SO}_2\text{N}(\text{R}^{\text{cc}})_2$, $-\text{SO}_2\text{R}^{\text{cc}}$, $-\text{SO}_2\text{OR}^{\text{cc}}$, $-\text{SOR}^{\text{aa}}$, $-\text{C}(=\text{S})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{C}(=\text{O})\text{SR}^{\text{cc}}$, $-\text{C}(=\text{S})\text{SR}^{\text{cc}}$, $-\text{P}(=\text{O})_2\text{R}^{\text{aa}}$, $-\text{P}(=\text{O})(\text{R}^{\text{aa}})_2$, $-\text{P}(=\text{O})_2\text{N}(\text{R}^{\text{cc}})_2$, $-\text{P}(=\text{O})(\text{NR}^{\text{cc}})_2$, $(\text{C}_1\text{-C}_{10})$ alkyl, $(\text{C}_1\text{-C}_{10})$ perhaloalkyl, $(\text{C}_2\text{-C}_{10})$ alkenyl, $(\text{C}_2\text{-C}_{10})$ alkynyl, $(\text{C}_3\text{-C}_{10})$ cycloalkyl, 3-14 membered heterocycloalkyl, $(\text{C}_6\text{-C}_{14})$ aryl, and 5-14 membered heteroaryl, or two R^{cc} groups attached to an N atom are joined to form a 3-14 membered heterocycloalkyl or 5-14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups, and wherein R^{aa} , R^{bb} , R^{cc} and R^{dd} are as defined above.

[00052] In certain embodiments, the substituent present on the nitrogen atom is a nitrogen protecting group (also referred to herein as an “amino protecting group”). Nitrogen protecting groups include, but are not limited to, $-\text{OH}$, $-\text{OR}^{\text{aa}}$, $-\text{N}(\text{R}^{\text{cc}})_2$, $-\text{C}(=\text{O})\text{R}^{\text{aa}}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{CO}_2\text{R}^{\text{aa}}$, $-\text{SO}_2\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{cc}})\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{cc}})\text{OR}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{cc}})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{SO}_2\text{N}(\text{R}^{\text{cc}})_2$, $-\text{SO}_2\text{R}^{\text{cc}}$, $-\text{SO}_2\text{OR}^{\text{cc}}$, $-\text{SOR}^{\text{aa}}$, $-\text{C}(=\text{S})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{C}(=\text{O})\text{SR}^{\text{cc}}$, $-\text{C}(=\text{S})\text{SR}^{\text{cc}}$, $(\text{C}_1\text{-C}_{10})$ alkyl (e.g., aralkyl, heteroaralkyl), $(\text{C}_2\text{-C}_{10})$ alkenyl, $(\text{C}_2\text{-C}_{10})$ alkynyl, $(\text{C}_3\text{-C}_{10})$ cycloalkyl, 3-14 membered heterocycloalkyl, $(\text{C}_6\text{-C}_{14})$ aryl, and 5-14 membered heteroaryl groups, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aralkyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups, and wherein R^{aa} , R^{bb} , R^{cc} , and R^{dd} are as defined herein. Nitrogen protecting groups are well known in the art and include those described in detail in *Protecting Groups in Organic Synthesis*, T. W. Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, incorporated herein by reference.

[00053] For example, nitrogen protecting groups such as amide groups (e.g., $-\text{C}(=\text{O})\text{R}^{\text{aa}}$) include, but are not limited to, formamide, acetamide, chloroacetamide, trichloroacetamide, trifluoroacetamide, phenylacetamide, 3-phenylpropanamide, picolinamide, 3-

pyridylcarboxamide, *N*-benzoylphenylalanyl derivative, benzamide, *p*-phenylbenzamide, *o*-nitrophenylacetamide, *o*-nitrophenoxyacetamide, acetoacetamide, (*N*'-dithiobenzyloxyacylamino)acetamide, 3-(*p*-hydroxyphenyl)propanamide, 3-(*o*-nitrophenyl)propanamide, 2-methyl-2-(*o*-nitrophenoxy)propanamide, 2-methyl-2-(*o*-phenylazophenoxy)propanamide, 4-chlorobutanamide, 3-methyl-3-nitrobutanamide, *o*-nitrocinnamide, *N*-acetylmethionine derivative, *o*-nitrobenzamide, and *o*-(benzoyloxymethyl)benzamide.

[00054] Nitrogen protecting groups such as carbamate groups (e.g., $-C(=O)OR^{aa}$) include, but are not limited to, methyl carbamate, ethyl carbamate, 9-fluorenylmethyl carbamate (Fmoc), 9-(2-sulfo)fluorenylmethyl carbamate, 9-(2,7-dibromo)fluorenylmethyl carbamate, 2,7-di-*t*-butyl-[9-(10,10-dioxo-10,10,10,10-tetrahydrothioxanthyl)]methyl carbamate (DBD-Tmoc), 4-methoxyphenacyl carbamate (Phenoc), 2,2,2-trichloroethyl carbamate (Troc), 2-trimethylsilylethyl carbamate (Teoc), 2-phenylethyl carbamate (hZ), 1-(1-adamantyl)-1-methylethyl carbamate (Adpoc), 1,1-dimethyl-2-haloethyl carbamate, 1,1-dimethyl-2,2-dibromoethyl carbamate (DB-*t*-BOC), 1,1-dimethyl-2,2,2-trichloroethyl carbamate (TCBOC), 1-methyl-1-(4-biphenyl)ethyl carbamate (Bpoc), 1-(3,5-di-*t*-butylphenyl)-1-methylethyl carbamate (*t*-Bumeoc), 2-(2'- and 4'-pyridyl)ethyl carbamate (Pyoc), 2-(*N,N*-dicyclohexylcarboxamido)ethyl carbamate, *t*-butyl carbamate (BOC or Boc), 1-adamantyl carbamate (Adoc), vinyl carbamate (Voc), allyl carbamate (Alloc), 1-isopropylallyl carbamate (Ipaoc), cinnamyl carbamate (Coc), 4-nitrocinnamyl carbamate (Noc), 8-quinolyl carbamate, *N*-hydroxypiperidinyl carbamate, alkylthio carbamate, benzyl carbamate (Cbz), *p*-methoxybenzyl carbamate (Moz), *p*-nitrobenzyl carbamate, *p*-bromobenzyl carbamate, *p*-chlorobenzyl carbamate, 2,4-dichlorobenzyl carbamate, 4-methylsulfinylbenzyl carbamate (MsZ), 9-anthrylmethyl carbamate, diphenylmethyl carbamate, 2-methylthioethyl carbamate, 2-methylsulfonylethyl carbamate, 2-(*p*-toluenesulfonyl)ethyl carbamate, [2-(1,3-dithianyl)]methyl carbamate (Dmoc), 4-methylthiophenyl carbamate (Mtpc), 2,4-dimethylthiophenyl carbamate (Bmpc), 2-phosphonioethyl carbamate (Peoc), 2-triphenylphosphonioisopropyl carbamate (Ppoc), 1,1-dimethyl-2-cyanoethyl carbamate, *m*-chloro-*p*-acyloxybenzyl carbamate, *p*-(dihydroxyboryl)benzyl carbamate, 5-benzisoxazolylmethyl carbamate, 2-(trifluoromethyl)-6-chromonylmethyl carbamate (Troc), *m*-nitrophenyl carbamate, 3,5-dimethoxybenzyl carbamate, *o*-nitrobenzyl carbamate, 3,4-dimethoxy-6-nitrobenzyl carbamate, phenyl(*o*-

nitrophenyl)methyl carbamate, *t*-amyl carbamate, *S*-benzyl thiocarbamate, *p*-cyanobenzyl carbamate, cyclobutyl carbamate, cyclohexyl carbamate, cyclopentyl carbamate, cyclopropylmethyl carbamate, *p*-decyloxybenzyl carbamate, 2,2-dimethoxyacetylvinyl carbamate, *o*-(*N,N*-dimethylcarboxamido)benzyl carbamate, 1,1-dimethyl-3-(*N,N*-dimethylcarboxamido)propyl carbamate, 1,1-dimethylpropynyl carbamate, di(2-pyridyl)methyl carbamate, 2-furanylmethyl carbamate, 2-iodoethyl carbamate, isoborynl carbamate, isobutyl carbamate, isonicotinyl carbamate, *p*-(*p*'-methoxyphenylazo)benzyl carbamate, 1-methylcyclobutyl carbamate, 1-methylcyclohexyl carbamate, 1-methyl-1-cyclopropylmethyl carbamate, 1-methyl-1-(3,5-dimethoxyphenyl)ethyl carbamate, 1-methyl-1-(*p*-phenylazophenyl)ethyl carbamate, 1-methyl-1-phenylethyl carbamate, 1-methyl-1-(4-pyridyl)ethyl carbamate, phenyl carbamate, *p*-(phenylazo)benzyl carbamate, 2,4,6-tri-*t*-butylphenyl carbamate, 4-(trimethylammonium)benzyl carbamate, and 2,4,6-trimethylbenzyl carbamate.

[00055] Nitrogen protecting groups such as sulfonamide groups (e.g., $-S(=O)_2R^{aa}$) include, but are not limited to, *p*-toluenesulfonamide (Ts), benzenesulfonamide, 2,3,6-trimethyl-4-methoxybenzenesulfonamide (Mtr), 2,4,6-trimethoxybenzenesulfonamide (Mtb), 2,6-dimethyl-4-methoxybenzenesulfonamide (Pme), 2,3,5,6-tetramethyl-4-methoxybenzenesulfonamide (Mte), 4-methoxybenzenesulfonamide (Mbs), 2,4,6-trimethylbenzenesulfonamide (Mts), 2,6-dimethoxy-4-methylbenzenesulfonamide (iMds), 2,2,5,7,8-pentamethylchroman-6-sulfonamide (Pmc), methanesulfonamide (Ms), β -trimethylsilylethanesulfonamide (SES), 9-anthracenesulfonamide, 4-(4',8'-dimethoxynaphthylmethyl)benzenesulfonamide (DNMBS), benzylsulfonamide, trifluoromethylsulfonamide, and phenacylsulfonamide.

[00056] Other nitrogen protecting groups include, but are not limited to, phenothiazinyl-(10)-acyl derivative, *N*'-*p*-toluenesulfonylaminoacyl derivative, *N*'-phenylaminothioacyl derivative, *N*-benzoylphenylalanyl derivative, *N*-acetylmethionine derivative, 4,5-diphenyl-3-oxazolin-2-one, *N*-phthalimide, *N*-dithiasuccinimide (Dts), *N*-2,3-diphenylmaleimide, *N*-2,5-dimethylpyrrole, *N*-1,1,4,4-tetramethyldisilylazacyclopentane adduct (STABASE), 5-substituted 1,3-dimethyl-1,3,5-triazacyclohexan-2-one, 5-substituted 1,3-dibenzyl-1,3,5-triazacyclohexan-2-one, 1-substituted 3,5-dinitro-4-pyridone, *N*-methylamine, *N*-allylamine, *N*-[2-(trimethylsilyl)ethoxy]methylamine (SEM), *N*-3-acetoxypyrrolamine, *N*-(1-isopropyl-4-nitro-2-oxo-3-pyrroline-3-yl)amine, quaternary ammonium salts, *N*-benzylamine, *N*-di(4-

methoxyphenyl)methylamine, *N*-5-dibenzosuberylamine, *N*-triphenylmethylamine (Tr), *N*-[(4-methoxyphenyl)diphenylmethyl]amine (MMTr), *N*-9-phenylfluorenylamine (PhF), *N*-2,7-dichloro-9-fluorenylmethyleneamine, *N*-ferrocenylmethylamino (Fcm), *N*-2-picolylamino *N*'-oxide, *N*-1,1-dimethylthiomethyleneamine, *N*-benzylideneamine, *N*-*p*-methoxybenzylideneamine, *N*-diphenylmethyleneamine, *N*-[(2-pyridyl)mesityl]methyleneamine, *N*-(*N*',*N*'-dimethylaminomethylene)amine, *N*,*N*'-isopropylidenediamine, *N*-*p*-nitrobenzylideneamine, *N*-salicylideneamine, *N*-5-chlorosalicylideneamine, *N*-(5-chloro-2-hydroxyphenyl)phenylmethyleneamine, *N*-cyclohexylideneamine, *N*-(5,5-dimethyl-3-oxo-1-cyclohexenyl)amine, *N*-borane derivative, *N*-diphenylborinic acid derivative, *N*-[phenyl(pentaacylchromium- or tungsten)acyl]amine, *N*-copper chelate, *N*-zinc chelate, *N*-nitroamine, *N*-nitrosoamine, amine *N*-oxide, diphenylphosphinamide (Dpp), dimethylthiophosphinamide (Mpt), diphenylthiophosphinamide (Ppt), dialkyl phosphoramidates, dibenzyl phosphoramidate, diphenyl phosphoramidate, benzenesulfenamide, *o*-nitrobenzenesulfenamide (Nps), 2,4-dinitrobenzenesulfenamide, pentachlorobenzenesulfenamide, 2-nitro-4-methoxybenzenesulfenamide, triphenylmethylsulfenamide, and 3-nitropyridinesulfenamide (Npys).

[00057] In certain embodiments, the substituent present on an oxygen atom is an oxygen protecting group (also referred to herein as an “hydroxyl protecting group”). Oxygen protecting groups include, but are not limited to, $-R^{aa}$, $-N(R^{bb})_2$, $-C(=O)SR^{aa}$, $-C(=O)R^{aa}$, $-CO_2R^{aa}$, $-C(=O)N(R^{bb})_2$, $-C(=NR^{bb})R^{aa}$, $-C(=NR^{bb})OR^{aa}$, $-C(=NR^{bb})N(R^{bb})_2$, $-S(=O)R^{aa}$, $-SO_2R^{aa}$, $-Si(R^{aa})_3$, $-P(R^{cc})_2$, $-P(R^{cc})_3$, $-P(=O)_2R^{aa}$, $-P(=O)(R^{aa})_2$, $-P(=O)(OR^{cc})_2$, $-P(=O)_2N(R^{bb})_2$, and $-P(=O)(NR^{bb})_2$, wherein R^{aa} , R^{bb} , and R^{cc} are as defined herein. Oxygen protecting groups are well known in the art and include those described in detail in *Protecting Groups in Organic Synthesis*, T. W. Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, incorporated herein by reference.

[00058] Exemplary oxygen protecting groups include, but are not limited to, methyl, methoxymethyl (MOM), methylthiomethyl (MTM), *t*-butylthiomethyl, (phenyldimethylsilyl)methoxymethyl (SMOM), benzyloxymethyl (BOM), *p*-methoxybenzyloxymethyl (PMBM), (4-methoxyphenoxy)methyl (*p*-AOM), guaiacolmethyl (GUM), *t*-butoxymethyl, 4-pentenylloxymethyl (POM), siloxymethyl, 2-methoxyethoxymethyl (MEM), 2,2,2-trichloroethoxymethyl, bis(2-chloroethoxy)methyl, 2-

(trimethylsilyl)ethoxymethyl (SEMOR), tetrahydropyranyl (THP), 3-bromotetrahydropyranyl, tetrahydrothiopyranyl, 1-methoxycyclohexyl, 4-methoxytetrahydropyranyl (MTHP), 4-methoxytetrahydrothiopyranyl, 4-methoxytetrahydrothiopyranyl S,S-dioxide, 1-[(2-chloro-4-methyl)phenyl]-4-methoxypiperidin-4-yl (CTMP), 1,4-dioxan-2-yl, tetrahydrofuranyl, tetrahydrothiofuranyl, 2,3,3a,4,5,6,7,7a-octahydro-7,8,8-trimethyl-4,7-methanobenzofuran-2-yl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 1-methyl-1-methoxyethyl, 1-methyl-1-benzyloxyethyl, 1-methyl-1-benzyloxy-2-fluoroethyl, 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 2-(phenylselenyl)ethyl, *t*-butyl, allyl, *p*-chlorophenyl, *p*-methoxyphenyl, 2,4-dinitrophenyl, benzyl (Bn), *p*-methoxybenzyl, 3,4-dimethoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-halobenzyl, 2,6-dichlorobenzyl, *p*-cyanobenzyl, *p*-phenylbenzyl, 2-picolyl, 4-picolyl, 3-methyl-2-picolyl *N*-oxido, diphenylmethyl, *p,p'*-dinitrobenzhydryl, 5-dibenzosuberyl, triphenylmethyl, α -naphthylidiphenylmethyl, *p*-methoxyphenyldiphenylmethyl, di(*p*-methoxyphenyl)phenylmethyl, tri(*p*-methoxyphenyl)methyl, 4-(4'-bromophenacyloxyphenyl)diphenylmethyl, 4,4',4''-tris(4,5-dichlorophthalimidophenyl)methyl, 4,4',4''-tris(levulinoyloxyphenyl)methyl, 4,4',4''-tris(benzyloxyphenyl)methyl, 3-(imidazol-1-yl)bis(4',4''-dimethoxyphenyl)methyl, 1,1-bis(4-methoxyphenyl)-1'-pyrenylmethyl, 9-anthryl, 9-(9-phenyl)xanthenyl, 9-(9-phenyl-10-oxo)anthryl, 1,3-benzodithiolan-2-yl, benzisothiazolyl S,S-dioxido, trimethylsilyl (TMS), triethylsilyl (TES), triisopropylsilyl (TIPS), dimethylisopropylsilyl (IPDMS), diethylisopropylsilyl (DEIPS), dimethylhexylsilyl, *t*-butyldimethylsilyl (TBDMS), *t*-butyldiphenylsilyl (TBDPS), tribenzylsilyl, tri-*p*-xylylsilyl, triphenylsilyl, diphenylmethylsilyl (DPMS), *t*-butylmethoxyphenylsilyl (TBMPS), formate, benzoylformate, acetate, chloroacetate, dichloroacetate, trichloroacetate, trifluoroacetate, methoxyacetate, triphenylmethoxyacetate, phenoxyacetate, *p*-chlorophenoxyacetate, 3-phenylpropionate, 4-oxopentanoate (levulinate), 4,4-(ethylenedithio)pentanoate (levulinoyldithioacetal), pivaloate, adamantate, crotonate, 4-methoxycrotonate, benzoate, *p*-phenylbenzoate, 2,4,6-trimethylbenzoate (mesitoate), methyl carbonate, 9-fluorenylmethyl carbonate (Fmoc), ethyl carbonate, 2,2,2-trichloroethyl carbonate (Troc), 2-(trimethylsilyl)ethyl carbonate (TMSEC), 2-(phenylsulfonyl) ethyl carbonate (Psec), 2-(triphenylphosphonio) ethyl carbonate (Peoc), isobutyl carbonate, vinyl carbonate, allyl carbonate, *t*-butyl carbonate (BOC or Boc), *p*-nitrophenyl carbonate, benzyl carbonate, *p*-methoxybenzyl carbonate, 3,4-dimethoxybenzyl carbonate, *o*-nitrobenzyl carbonate, *p*-nitrobenzyl carbonate, *S*-benzyl thiocarbonate, 4-ethoxy-1-naphthyl carbonate, methyl

dithiocarbonate, 2-iodobenzoate, 4-azidobutyrate, 4-nitro-4-methylpentanoate, *o*-(dibromomethyl)benzoate, 2-formylbenzenesulfonate, 2-(methylthiomethoxy)ethyl, 4-(methylthiomethoxy)butyrate, 2-(methylthiomethoxymethyl)benzoate, 2,6-dichloro-4-methylphenoxyacetate, 2,6-dichloro-4-(1,1,3,3-tetramethylbutyl)phenoxyacetate, 2,4-bis(1,1-dimethylpropyl)phenoxyacetate, chlorodiphenylacetate, isobutyrate, monosuccinoate, (*E*)-2-methyl-2-butenolate, *o*-(methoxyacyl)benzoate, α -naphthoate, nitrate, alkyl *N,N,N',N'*-tetramethylphosphorodiamidate, alkyl *N*-phenylcarbamate, borate, dimethylphosphinothioyl, alkyl 2,4-dinitrophenylsulfenate, sulfate, methanesulfonate (mesylate), benzylsulfonate, and tosylate (Ts).

[00059] In certain embodiments, the substituent present on a sulfur atom is a sulfur protecting group (also referred to as a “thiol protecting group”). Sulfur protecting groups include, but are not limited to, $-R^{aa}$, $-N(R^{bb})_2$, $-C(=O)SR^{aa}$, $-C(=O)R^{aa}$, $-CO_2R^{aa}$, $-C(=O)N(R^{bb})_2$, $-C(=NR^{bb})R^{aa}$, $-C(=NR^{bb})OR^{aa}$, $-C(=NR^{bb})N(R^{bb})_2$, $-S(=O)R^{aa}$, $-SO_2R^{aa}$, $-Si(R^{aa})_3$, $-P(R^{cc})_2$, $-P(R^{cc})_3$, $-P(=O)_2R^{aa}$, $-P(=O)(R^{aa})_2$, $-P(=O)(OR^{cc})_2$, $-P(=O)_2N(R^{bb})_2$, and $-P(=O)(NR^{bb})_2$, wherein R^{aa} , R^{bb} , and R^{cc} are as defined herein. Sulfur protecting groups are well known in the art and include those described in detail in *Protecting Groups in Organic Synthesis*, T. W. Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, incorporated herein by reference.

[00060] As used herein, a “leaving group” (LG) is an art-understood term referring to a molecular fragment that departs with a pair of electrons in heterolytic bond cleavage, wherein the molecular fragment is an anion or neutral molecule. As used herein, a leaving group can be an atom or a group capable of being displaced by a nucleophile. See, for example, Smith, March *Advanced Organic Chemistry* 6th ed. (501-502). Exemplary leaving groups include, but are not limited to, halo (e.g., chloro, bromo, iodo), $-OR^{aa}$ (when the O atom is attached to a carbonyl group, wherein R^{aa} is as defined herein), $-O(C=O)R^{LG}$, or $-O(SO)_2R^{LG}$ (e.g., tosyl, mesyl, besyl), wherein R^{LG} is optionally substituted alkyl, optionally substituted aryl, or optionally substituted heteroaryl. In certain embodiments, the leaving group is a halogen.

[00061] The terms for which definitions are given above are specifically exemplified in the Examples.

[00062] “Yield” for each of the reactions described herein is expressed as a percentage of the theoretical yield.

[00063] “Patient” for the purposes of the present invention includes humans and any other animals, particularly mammals, and other organisms. Thus the methods are applicable to both human therapy and veterinary applications. In a preferred embodiment the patient is a mammal, and in a most preferred embodiment the patient is human. Examples of the preferred mammals include mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, and primates.

[00064] “Kinase-dependent diseases or conditions” refer to pathologic conditions that depend on the activity of one or more kinases. Kinases either directly or indirectly participate in the signal transduction pathways of a variety of cellular activities including proliferation, adhesion, migration, differentiation, and invasion. Diseases associated with kinase activities include tumor growth, the pathologic neovascularization that supports solid tumor growth, and associated with other diseases where excessive local vascularization is involved such as ocular diseases (diabetic retinopathy, age-related macular degeneration, and the like) and inflammation (psoriasis, rheumatoid arthritis, and the like).

[00065] “Therapeutically effective amount” is an amount of a compound of the invention that, when administered to a patient, ameliorates a symptom of the disease. The amount of a compound of the invention which constitutes a “therapeutically effective amount” will vary depending on the compound, the disease state and its severity, the age of the patient to be treated, and the like. The therapeutically effective amount can be determined routinely by one of ordinary skill in the art having regard to his own knowledge and to this disclosure.

[00066] “Cancer” refers to cellular-proliferative disease states, including but not limited to: Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Head and neck: squamous cell carcinomas of the head and neck, laryngeal and hypopharyngeal cancer, nasal cavity and paranasal sinus cancer, nasopharyngeal cancer, salivary gland cancer, oral and oropharyngeal cancer; Lung: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma, non-small cell lung cancer), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Colon: colorectal cancer, adenocarcinoma, gastrointestinal stromal tumors, lymphoma, carcinoids, Turcot Syndrome; Gastrointestinal: gastric cancer, gastroesophageal junction adenocarcinoma, esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma,

glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma); Breast: metastatic breast cancer, ductal carcinoma in situ, invasive ductal carcinoma, tubular carcinoma, medullary carcinoma, mucinous carcinoma, lobular carcinoma in situ, triple negative breast cancer; Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma, leukemia, renal cell carcinoma), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma, urothelial carcinoma), prostate (adenocarcinoma, sarcoma, castrate resistant prostate cancer), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma), clear cell carcinoma, papillary carcinoma; Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondilaginous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma, and giant cell tumors; Thyroid: medullary thyroid cancer, differentiated thyroid cancer, papillary thyroid cancer, follicular thyroid cancer, hurthle cell cancer, and anaplastic thyroid cancer; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial cancer), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli--Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple

myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma]; Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and Adrenal glands: neuroblastoma. Thus, the term “cancerous cell” as provided herein, includes a cell afflicted by any one of the above-identified conditions.

[00067] “Pharmaceutically acceptable salts” includes “pharmaceutically acceptable acid addition salts” and “pharmaceutically acceptable base addition salts.” “Pharmaceutically acceptable acid addition salts” refers to those salts that retain the biological effectiveness of the free bases and that are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like, as well as organic acids such as acetic acid, trifluoroacetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like.

[00068] “Pharmaceutically acceptable base addition salts” include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts, and the like. Exemplary salts are the ammonium, potassium, sodium, calcium, and magnesium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins, and the like. Exemplary organic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline, and caffeine. (See, for example, S. M. Berge, et al., “Pharmaceutical Salts,” J. Pharm. Sci., 1977;66:1-19 which is incorporated herein by reference.)

[00069] The term, “compound,” as used herein is meant to include all stereoisomers, geometric isomers, tautomers and isotopes of the structures depicted. The term is also meant to

refer to compounds of the inventions, regardless of how they are prepared, e.g., synthetically, through biological process (e.g., metabolism or enzyme conversion), or a combination thereof.

[00070] Compounds of the invention can also include all isotopes of atoms occurring in the intermediates or final compounds. Isotopes include those atoms having the same atomic number but different mass numbers. For example, isotopes of hydrogen include tritium and deuterium.


[00071] Any one of the process steps or sequences disclosed and/or claimed herein can be performed under an inert gas atmosphere, more particularly under argon or nitrogen. In addition, the methods of the present invention may be carried out as semi-continuous or continuous processes, more preferably as continuous processes.

[00072] Moreover, many of the process steps and sequences that are described herein can be telescoped.

[00073] In general, the nomenclature used in this Application is based on naming conventions adopted by the International Union of Pure and Applied Chemistry (IUPAC). Chemical structures shown herein were prepared using CHEMDRAW®. Any open valency appearing on a carbon, oxygen, or nitrogen atom in the structures herein indicates the presence of a hydrogen atom.

Embodiments of the Invention


[00074] One aspect provides a compound of formula A:



A


or a pharmaceutically acceptable salt thereof, wherein the variables and substituents in formula A are as defined in the Summary of the Invention.

[00075] In one embodiment of this aspect, the compound of formula A is a compound of formula A-1.




A-1

[00076] In another embodiment of this aspect, the compound of formula A is a compound of formula A-2.



A-2

[00077] In a further embodiment of this aspect, the compound of formula A is a compound of formula A-3:




A-3

wherein R^{a1} is -H or (C₁-C₆) alkyl.

[00078] In a further embodiment, R_1 in the compound of formula A-3 is -H.

[00079] In a further embodiment of this aspect, the compound of formula A is a compound of formula A-4:



A-4

wherein ring A is 5- to 14-membered heteroaryl; and the subscript r is 1, 2, 3, or 4.

[00080] In this embodiment, R_2 is -H.

[00081] In a further embodiment, r in formula A-4 is 1 or 2.

[00082] In a further embodiment of formula A: R_1 is -H, optionally substituted (C_1 - C_6) alkyl, halo, $-OR^a$, $-NO_2$, $-NH_2$, $-NHR^a$, $-NR^aR^a$, $-SR^a$, $-SOR^a$, or $-S(O)_2R^a$, and R_2 is selected from the group consisting of (C_2 - C_6) alkenyl, (C_2 - C_6) alkynyl, (C_6 - C_{10}) aryl, (C_3 - C_{10}) cycloalkyl, $-CN$, $-NHOR^a$, $-C(O)R^a$, $-C(O)NR^aR^a$, $-C(O)NHOR^a$, $-C(O)OR^a$, $-C(O)NR^aS(O)_2R^a$, $-OC(O)NR^aR^a$, $NR^aC(O)R^a$, $-NR^aC(=NR^a)R^a$, $-NR^aC(O)OR^a$, $-NR^aC(O)NR^aR^a$, $-C(=NR^a)R^a$, $-C(=NOH)R^a$, $-C(=NOR^a)R^a$, $-C(=NOH)NR^a$, $-C(=NCN)NR^aR^a$, $-NR^aC(=NCN)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-NR^aC(=NR^a)NR^aR^a$, $-NR^aS(O)R^a$, $-NR^aS(O)_2R^a$, $-NR^aS(O)_2NR^aR^a$, $-S(O)R^a$, $-S(O)NR^aR^a$, $-S(O)_2R^a$, $-S(O)_2NR^aC(O)R^a$, $-P(O)R^aR^a$, $-P(O)(OR^a)(OR^a)$, $-B(OH)_2$, $-B(OR^a)_2$, and $-S(O)_2NR^aR^a$.

[00083] In one embodiment of this embodiment, R_1 is -H.

[00084] In a further embodiment: R_1 is selected from the group consisting of (C_2 - C_6) alkenyl, (C_2 - C_6) alkynyl, (C_6 - C_{10}) aryl, (C_3 - C_{10}) cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, $-CN$, $-NHOR^a$, $-C(O)R^a$, $-C(O)NR^aR^a$, $-C(O)NHOR^a$, $-C(O)OR^a$, $-C(O)NR^aS(O)_2R^a$, $-OC(O)NR^aR^a$, $-NR^aC(O)R^a$, $-NR^aC(=NR^a)R^a$, $-NR^aC(O)OR^a$, $-NR^aC(O)NR^aR^a$, $-C(=NR^a)R^a$, $-C(=NOH)R^a$, $-C(=NOH)NR^a$, $-C(=NOR^a)R^a$, $-C(=NCN)NR^aR^a$, $-NR^aC(=NCN)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-NR^aC(=NR^a)NR^aR^a$, $-NR^aS(O)R^a$, $-NR^aS(O)_2R^a$, $-NR^aS(O)_2NR^aR^a$, $-S(O)R^a$, $-S(O)NR^aR^a$, $-S(O)_2R^a$, $-S(O)_2NR^aC(O)R^a$, $-P(O)R^aR^a$, $-P(O)(OR^a)(OR^a)$, $-B(OH)_2$, $-B(OR^a)_2$, and $-S(O)_2NR^aR^a$; and

R_2 is -H, optionally substituted (C_1 - C_6) alkyl, halo, $-OR^a$, $-NO_2$, $-NH_2$, $-NHR^a$, $-NR^aR^a$, $-SR^a$, $-SOR^a$, or $-S(O)_2R^a$.

[00085] In one embodiment of this embodiment, R₂ is -H.

[00086] In another embodiment:

R₁ is (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, -CN, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOR^a)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, or -S(O)₂NR^aR^a; and

R₂ is selected from the group consisting of -H, halo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NO₂, -OR^a, -SR^a, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)R^a, -OC(O)NR^aR^a, -NHR^a, -NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NOR^a)R^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂; and -S(O)₂NR^aR^a.

[00087] In one embodiment of this embodiment, R₂ is -H.

[00088] In another embodiment:


R₁ is selected from the group consisting of -H, halo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NO₂, -OR^a, -SR^a, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)R^a, -OC(O)NR^aR^a, -NHR^a, -NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOR^a)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and -S(O)₂NR^aR^a; and

In one embodiment of this embodiment, R₁ is -H.

[00089] In another further embodiment:

R₁ and R₂ taken together with the atoms to which they are attached form a fused (C₃-C₇) cycloalkyl ring or a fused 4- to 10-membered heterocycloalkyl ring; wherein the fused (C₃-C₇) cycloalkyl ring or a fused 4- to 10-membered heterocycloalkyl ring are each optionally substituted with 1, 2, or 3 independently selected R^b substituents, provided that the compound is not 1-[2-(4-Fluoro-phenyl)-acetyl]-cyclopropanecarboxylic acid [3-fluoro-4-(7,8,10,11,13,14-hexahydro-6,9,12,15-tetraoxa-1-aza-cyclododeca[b]naphthalen-4-yloxy)-phenyl]-amide.

[00090] In a further embodiment: R₁ and R₂ taken together with the atoms to which they are attached form a fused (C₃-C₇) cycloalkyl ring or a fused 4- to 10-membered heterocycloalkyl ring, wherein the fused (C₃-C₇) cycloalkyl ring or a fused 4- to 10-membered heterocycloalkyl ring are each optionally substituted with 1, 2, or 3 independently selected R^b substituents, provided that the compound is not a compound having the formula:



wherein ring E is a fused 4- to 10-membered heterocycloalkyl.

[00091] In another embodiment, R₁ in the compound of formula A, A-1, or A-3, ~~or A-4~~ is selected from the group consisting of -H, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -C(=NO-(C₁-C₆) alkyl)R^a, -CN, -C(O)OR^a, -C(O)NR^aR^a, -C(O)NHR^a, -S(O)₂NR^aR^a, phenyl, 5- to 6-membered heteroaryl, (C₃-C₆) cycloalkyl, and 4- to 6-membered heterocycloalkyl.

[00092] In another embodiment, R₁ is selected from the group consisting of -H, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -C(=NO-(C₁-C₆) alkyl)R^a, -CN, -C(O)OR^a, -C(O)NR^aR^a, -C(O)NHR^a, -S(O)₂NR^aR^a, phenyl, 5- to 6-membered heteroaryl, (C₃-C₆) cycloalkyl, and 4- to 6-membered heterocycloalkyl; and

R₂ is H, optionally substituted (C₁-C₆) alkyl, halo, -OR^a, -NO₂, -NH₂, -NHR^a, -NR^aR^a, -SR^a, -SOR^a, or -S(O)₂R^a.

[00093] In another embodiment, R₁ is selected from the group consisting of -H, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -C(=NO-(C₁-C₆) alkyl)R^a, -CN, -C(O)OR^a, -C(O)NR^aR^a,

-C(O)NHOR^a, -S(O)₂NR^aR^a, phenyl, 5- to 6-membered heteroaryl, (C₃-C₆) cycloalkyl; and 4- to 6-membered heterocycloalkyl; and

R₂ is selected from the group consisting of -H, halo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NO₂, -OR^a, -SR^a, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)R^a, -OC(O)NR^aR^a, -NHR^a, -NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NOR^a)R^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂; and -S(O)₂NR^aR^a.

[00094] In another embodiment, R₁ is selected from the group consisting of -H, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -C(=NO-(C₁-C₆) alkyl)R^a, -CN, -C(O)OR^a, -C(O)NR^aR^a, -C(O)NHOR^a, -S(O)₂NR^aR^a, phenyl, 5- to 6-membered heteroaryl, -(C₃-C₆) cycloalkyl, and 4- to 6-membered heterocycloalkyl; and

R₂ is (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -CN, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NOR^a)R^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, or -S(O)₂NR^aR^a.

[00095] In a further embodiment, R₁ is -H, R^aNHC(O)-, R^aOC(O)-, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, or -C(=NO-CH₃)R^a, and R₂ is selected from 2-methoxyethylamino, azetidin-1-yl, methylamino, 3-morpholinopropoxy, 2-methoxyethoxy, 2-hydroxyethoxy, propoxy, 2-hydroxypropoxy, methoxycarbonyl, carboxy, carbamoyl, methylcarbamoyl, (2-hydroxyethoxy)carbamoyl, (2,2-dihydroxyethoxy)carbamoyl, (oxetan-3-yloxy)carbamoyl, methoxycarbamoyl, 2-trimethylsilylethynyl, ethynyl, sulfamoyl, acetyl, and -C(=NOCH₃)CH₃.

[00096] In a further embodiment of formula A, and A-2, and R₂ is selected from the group consisting of -H, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -C(=NO-(C₁-C₆) alkyl)R^a, -CN, -C(O)OR^a, -C(O)NR^aR^a, -C(O)NHOR^a, and -S(O)₂NR^aR^a.

[00097] In a further embodiment, R_1 is -H, optionally substituted (C₁-C₆) alkyl, halo, -OR^a, -NO₂, -NH₂, -NHR^a, -NR^aR^a, -SR^a, -SOR^a, or S(O)₂R^a, and

R_2 is selected from the group consisting of -H, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -C(=NO-(C₁-C₆) alkyl)R^a, -CN, -C(O)OR^a, -C(O)NR^aR^a, -C(O)NHOR^a, -S(O)₂NR^aR^a, phenyl, and (C₃-C₆) cycloalkyl.

[00098] In a further embodiment, R_1 is selected from the group consisting of -H, optionally substituted (C₁-C₆) alkyl, halo, -OR^a, -NO₂, -NH₂, -NHR^a, -NR^aR^a, -SR^a, -SOR^a, and -S(O)₂R^a, and

R_2 is selected from the group consisting of -H, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -C(=NO-(C₁-C₆) alkyl)R^a, -CN, -C(O)OR^a, -C(O)NR^aR^a, -C(O)NHOR^a, -S(O)₂NR^aR^a, .

[00099] In a further embodiment, R_1 is selected from the group consisting of (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, -CN, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NOR^a)R^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and -S(O)₂NR^aR^a; and

R_2 is selected from the group consisting of -H, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -C(=NO-(C₁-C₆) alkyl)R^a, -CN, -C(O)OR^a, -C(O)NR^aR^a, -C(O)NHOR^a, -S(O)₂NR^aR^a, .

[000100] In a further embodiment, R_1 is selected from the group consisting of -H, halo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NO₂, -OR^a, -SR^a, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)R^a, -OC(O)NR^aR^a, -NHR^a, -NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOR^a)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and -S(O)₂NR^aR^a; and

R_2 is selected from the group consisting of -H, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, (C_2-C_6) alkynyl, $-C(=NO-(C_1-C_6) \text{ alkyl})R^a$, -CN, $-C(O)OR^a$, $-C(O)NR^aR^a$, $-C(O)NHOR^a$, $-S(O)_2NR^aR^a$.

[000101] In a further embodiment, R_1 is selected from the group consisting of 2-methoxyethylamino, azetidin-1-yl, methylamino, 3-morpholinopropoxy, 2-methoxyethoxy, 2-hydroxyethoxy, propoxy, 2-hydroxypropoxy, methoxycarbonyl, carboxy, carbamoyl, methylcarbamoyl, 2-oxazolyl, pyrazol-3-yl, pyrazol-4-yl, 4-isoxazolyl, 3,5-dimethylisoxazol-4-yl, 1-methyl-pyrazol-4-yl, 2-methyl-pyrazol-3-yl, 2-ethyl-pyrazol-3-yl, 2-(2-hydroxyethyl)-pyrazol-3-yl, 2-(2,2,2-trifluoroethyl)-pyrazol-3-yl, 2-(2-fluoroethyl)-pyrazol-3-yl, 2-(2,2-difluoroethyl)-pyrazol-3-yl, 2-trifluoromethyl-pyrazol-3-yl, 2-difluoromethyl-pyrazol-3-yl, 1-methyl-imidazol-4-yl, 1-methyl-imidazol-2-yl, 1H-imidazol-2-yl, (2-hydroxyethoxy)carbamoyl, (2,2-dihydroxyethoxy)carbamoyl, (oxetan-3-yloxy)carbamoyl, methoxycarbamoyl, 2-trimethylsilylethynyl, ethynyl, 1,3,4-oxadiazol-3-yl, 1H-1,2,3-triazol-5-yl, sulfamoyl, acetyl, and $-C(=NOCH_3)CH_3$; and

R_2 is -H, $-R^aNHC(O)-$, $-R^aOC(O)-$, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, or $-C(=NO-CH_3)R^a$.

[000102] In a further embodiment of formula A-4, the subscript r is 1 or 2.


[000103] In a further embodiment of the above aspect and embodiments, R_{10} and R_{11} are each -H.

[000104] In a further embodiment of the above aspect and embodiments, the subscript n is 1.

[000105] In a further embodiment of the above aspect and embodiments, the subscript m is 1.

[000106] In a further embodiment of the above aspect and embodiments, the subscript p is 1.

[000107] Another embodiment, a compound of formula A is a compound of formula B:



B

or a pharmaceutically acceptable salt thereof, wherein R₁ and R₂ are as defined in (i), (ii), or (iii) of formula A; and R₃, R₁₀, R₁₁, R₁₄, R₄, n, p, m and Y are as defined as follows:

each R₃ is independently selected from the group consisting of -H, halo, -OH, -CN, optionally substituted (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkoxy, -NH₂, -NH(C₁-C₆)alkyl, -N(C₁-C₆ alkyl)₂, and (C₃-C₆) cycloalkyl, wherein (C₁-C₆) alkoxy, -NH(C₁-C₆)alkyl, -N(C₁-C₆ alkyl)₂, and (C₃-C₆) cycloalkyl are each optionally substituted;

each of R₁₀ and R₁₁ is independently selected from the group consisting of -H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and (C₁-C₆) haloalkoxy;

each R₁₄ is independently selected from the group consisting of -halo, -OH, -NH₂, -CN, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, -COOH, -NH(C₁-C₆)alkyl, -N(C₁-C₆ alkyl)₂, phenyl, phenyl-(C₁-C₂) alkylene, (C₃-C₆) cycloalkyl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, 4- to 6-membered heterocycloalkyl, (4- to 6-membered heterocycloalkyl)-(C₁-C₄) alkylene-, 5- to 6-membered heteroaryl, (5- to 6-membered heteroaryl)-(C₁-C₄) alkylene-, and -OR^e, wherein the (C₁-C₆) alkyl, phenyl, phenyl-(C₁-C₂) alkylene, (C₃-C₆) cycloalkyl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, 4- to 6-membered heterocycloalkyl, (4- to 6-membered heterocycloalkyl)-(C₁-C₄) alkylene-, 5- to 6-membered heteroaryl, and (5- to 6-membered heteroaryl)-(C₁-C₄) alkylene- of R₁₄ are each optionally substituted;

each R₄ is independently selected from -H, halo, -OH, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, and (C₁-C₆) haloalkoxy, wherein the (C₁-C₆) alkyl are each (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, and (C₁-C₆) haloalkoxy are each independently optionally substituted;

each R^a is independently selected from -H, (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, wherein (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄)

alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- are each independently optionally substituted;

n, p, and m are each independently integers of 0 to 3; and

Y is selected from -O-, -S-, -SO-, -SO₂-, -NH-, and -N((C₁-C₆) alkyl)-.


[000108] In one embodiment of formula A and B:

(i) R₁ is selected from the group consisting of:

(C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl are each independently optionally substituted,



-CN, -P(O)R^aR^a, P(O)(OR^a)₂, B(OH)₂, B(OR^a)₂,

X₂R^a, wherein X₂ is -NHO-, -NH-S(O)-, -N-(C₁-C₆)alkyl-S(O)-, -NH-S(O)₂-, -N-(C₁-C₆)alkyl-S(O)₂R^a-, -NH-S(O)-NH-, -N-(C₁-C₆)alkyl-S(O)NH-, -NH-S(O)₂NH-, -N-(C₁-C₆)alkyl-S(O)₂NH-, -S(O)₂NHC(O)-, and

$$\begin{array}{c} \text{Z}^1 \\ \parallel \\ \text{R}^a - \text{Y}_2 - \text{C} - \text{Y}_1 - \text{Z}^2 \end{array}$$
 wherein “” indicates the point of attachment, wherein:

Y₁ is absent, or is -NH-, -N-(C₁-C₆) alkyl-, or -O-,

Y₂ is absent, or is -O-, -NH-, -NHO-, -N-(C₁-C₆) alkyl-, -N₂H₂-, -NH-S(O)-, or -NH-S(O)₂-, or

Y₂ is or optionally substituted , wherein “” indicates points of attachment, wherein ring A is a 3, 4, 5, 6, or 7-membered ring; and

Z¹ is O, NH, N-(C₁-C₆) alkyl, NOH, NO-(C₁-C₆) alkyl, or NCN; and

R₂ is:

-H or a group selected from the group consisting of:

(C₁-C₆) alkyl, halo, -NO₂, and X₁R^a, wherein X₁ is -O-, -S-, -SO-, -SO₂-, -SO₂NH-, -SO₂NR^a-, -NH-, and -N-(C₁-C₆) alkyl-, wherein (C₁-C₆) alkyl is optionally substituted.


[000109] In another embodiment of formula A and B:

(ii) R₁ is selected from the group consisting of:

(C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, and 4-14 membered heterocycloalkyl are each independently optionally substituted;


P(O)R^aR^a, P(O)(OR^a)(OR^a), B(OH)₂, B(OR^a)₂, CN,

X₂R^a, wherein X₂ is -NHO-, -NH-S(O)-, -N-(C₁-C₆)alkyl-S(O)-, -NH-S(O)₂-, -N-(C₁-C₆) alkyl-S(O)₂R^a-, -NH-S(O)-NH-, -N-(C₁-C₆) alkyl-S(O)NH-, -NH-S(O)₂NH-, -N-(C₁-C₆) alkyl-S(O)₂NH-, -S(O)-, -S(O)₂-, -S(O)₂NHC(O), and



Y₁ is absent, or is -NH-, -N-(C₁-C₆) alkyl-, or -O-,

Y₂ is absent, or is -O-, -NH-, -NHO-, -N-(C₁-C₆) alkyl-, -N₂H₂-, -NH-S(O)-, or -NH-S(O)₂-, or



Y₂ is optionally substituted, wherein “” indicates points of attachment, wherein ring A is a 3, 4, 5, 6, or 7-membered ring;


Z¹ is -O-, -NH-, -N-(C₁-C₆) alkyl-, -NOH-, -NO-(C₁-C₆) alkyl-, or -NCN-; and

R₂ is selected from the group consisting of

H, halo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, wherein (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- are each independently optionally substituted,


CN, NO₂, P(O)R^aR^a, P(O)(OR^a)(OR^a), B(OH)₂, B(OR^a)₂,



X_1R^a , wherein X_1 is $-O-$, $-S-$, $-NH-$, or $-N-(C_1-C_6)-$, $-NHO-$, $-NH-S(O)-$, $-N-(C_1-C_6)alkyl-S(O)-$, $-NH-S(O)_2-$, $-N-(C_1-C_6)alkyl-S(O)_2-$, $-NH-S(O)-NH-$, $-N-(C_1-C_6)alkyl-S(O)NH-$, $-NH-S(O)_2NH-$, $-N-(C_1-C_6)alkyl-S(O)_2NH-$, $-S(O)_2NHC(O)-$, $-NH-S(O)R^a-$, $-N-(C_1-C_6)alkyl-S(O)R^a-$, $-NH-S(O)_2R^a-$, $-N-(C_1-C_6)alkyl-S(O)_2R^a-$, and



Y_1 is absent or is $-NH-$, $-N-(C_1-C_6)alkyl-$, or $-O-$;

Y_2 is absent or is $-O-$, $-NH-$, $-NHO-$, $-N-(C_1-C_6)alkyl-$, $-N_2H_2-$, $-NH-S(O)-$, or $-NH-S(O)_2-$, or




Y_2 is optionally substituted , wherein ring A is a 3, 4, 5, 6, or 7-membered ring and wherein “” indicates points of attachment; and Z^1 is O, NH, $N-(C_1-C_6)alkyl$, NOH, $NO-(C_1-C_6)alkyl$, or NCN,

[000110] In another embodiment of formula A and B:

(iii) R_1 and R_2 taken together with the atoms to which they are attached form a 4- to 10-membered heterocycloalkyl ring optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of halo, $(C_1-C_6)alkyl$, $(C_1-C_6)haloalkyl$, $(C_1-C_6)haloalkoxy$, $-CN$, $-OH$, $-NH_2$, provided that the compound is not 1-[2-(4-Fluoro-phenyl)-acetyl]-cyclopropanecarboxylic acid [3-fluoro-4-(7,8,10,11,13,14-hexahydro-6,9,12,15-tetraoxa-1-aza-cyclododeca[b]naphthalen-4-yloxy)-phenyl]-amide.

[000111] In a further embodiment of the compound of formula A and B:

R_1 is $-H$, $-CN$, $(C_1-C_6)alkyl$, $(C_3-C_{10})cycloalkyl$, $(C_6-C_{10})aryl$, 4-10 membered heterocycloalkyl, 5-10 membered heteroaryl, $-S(O)_2NHR^a$, $-P(O)R^aR^a$, $-OR^a$, or




Y_1 is absent or is $-NH-$, $-N-(C_1-C_6)alkyl-$, or $-O-$;

Y_2 is absent or is $-O-$, $-NH-$, $-NHO-$, $-N-(C_1-C_6)alkyl-$, $-NH-NH-$, $-NH-S(O)-$, or $NH-S(O)_2$; and

Z^1 is $-O$, $-NH$, $-N-(C_1-C_6)alkyl$, $-N-OH$, or $-N-O(C_1-C_6)alkyl$.

[000112] In another embodiment of the of formula A and B:

R_2 is -H, halo, $-X_1R^a$, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, or



Y_1 is absent or is NH, N-(C₁-C₆) alkyl, or O;

Y_2 is absent or is O, NH, NHO, N-(C₁-C₆) alkyl, N₂H₂, NH-S(O), or NH-S(O)₂;

and

Z^1 is -O, -NH, -N-(C₁-C₆) alkyl, -NOH-, or -N-O(C₁-C₆)alkyl.


[000113] In another embodiment of formula A and B, R_3 is -H or halo.

[000114] In another embodiment of formula A and B, R_4 is -H or halo.


[000115] In another embodiment of formula A and B, wherein R_{14} is -H or halo.

[000116] In another embodiment of formula A and B, Y is -O-.

[000117] In another embodiment, the compound of B is a compound of either formula B-1 or B-2:



B-1




B-2

or a pharmaceutically acceptable salt thereof.

[000118] In one embodiment of formula B-1:


R^{a1} is optionally substituted (C₁-C₆) alkyl;


R₁ is -H, -CN, optionally substituted 3-6 membered cycloalkyl, optionally substituted phenyl, optionally substituted 4-6 membered heterocycloalkyl, optionally substituted 5-6 membered heteroaryl, -SO₂-(C₁-C₆) alkyl, -SO₂NH₂, -SO₂-NH(C₁-C₆) alkyl, P(O)((C₁-C₆) alkyl)₂, or



Y₁ is absent;

Y₂ is absent or is -O-, -NH-, -NHO-, -NH-NH-, -N-(C₁-C₆) alkyl-; or



Y₂ is optionally substituted , wherein ring A is a 3, 4, 5, 6, or 7-membered ring, wherein "∼" indicates points of attachment;


Z¹ is O, NH, N-(C₁-C₆) alkyl, NHO, or NO-(C₁-C₆) alkyl; and

R^a is -H, -(C₁-C₆) alkyl, 4-6-membered heterocycloalkyl, 3-6-membered cycloalkyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NH(C₁-C₆), -(C₂-C₆) alkylene-N(C₁-C₆)₂, or -(C₂-C₆) alkylene-N-(4-6-membered heterocycloalkyl);

[000119] In another embodiment of formula B-1:

R^{al} is (C₁-C₆) alkyl;

R₁ is -H, -CN, optionally substituted cyclopropyl, optionally substituted phenyl, optionally substituted 4-6 membered azetidiny, optionally substituted pyrrolidiny, optionally substituted piperidiny, optionally substituted oxetany, optionally substituted oxazolyl, optionally substituted pyridiny, optionally substituted imidazolyl, optionally substituted pyrrolyl, optionally substituted furnayl, optionally substituted pyrazolyl, optionally substituted oxadiazolyl, -SO₂-(C₁-C₆) alkyl, -SO₂NH₂, -SO₂-NH(C₁-C₆) alkyl, or P(O)((C₁-C₆) alkyl)₂; or



Y₁ is absent;

Y₂ is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or



Y₂ is optionally substituted azetidiny;

Z¹ is O, NH, or N-(C₁-C₆) alkyl; and

R^a is H, (C₁-C₆) alkyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NH(C₁-C₆) alkyl, -(C₂-C₆) alkylene-N((C₁-C₆) alkyl)₂, -(C₂-C₆) alkylene-heterocycloalkyl, or 4-6 membered heterocycloalkyl, wherein heterocycloalkyl is optionally substituted.

[000120] In one embodiment of formula B-2:

R^{a2} is optionally substituted (C₁-C₆) alkyl;

R₂ is  wherein “” indicates the point of attachment, wherein:

Y₁ is absent;



Y₂ is absent or is -O- or -NH-; and

Z¹ is O; and

R^a is -H or -(C₁-C₆) alkyl.

[000121] In another embodiment of formula B-2:

R^{a2} is (C₁-C₆) alkyl;

R₂ is  wherein “” indicates the point of attachment, wherein:

Y₁ is absent;

Y₂ is absent or is -O- or -NH-; and



Z¹ is O or NO-(C₁-C₆) alkyl; and

R^a is -H or -(C₁-C₆) alkyl.

[000122] In another embodiment of formula B-1, R^{a1} is methoxy.

[000123] In another embodiment of formula B-2, R^{a2} is methoxy.

[000124] In another embodiment, the compound of formula B is a compound of either formula B-3 or B-4:

**B-3****B-4**

or a pharmaceutically acceptable salt thereof.

[000125] In one embodiment of formula B-3:

R₁ is -H or (C₁-C₆) alkyl; and

Y₁ is absent;

Y₂ is absent or is -O-, -NHO-, or -NH-; and

Z¹ is O or NO-(C₁-C₆) alkyl; and

R^a is -H or -(C₁-C₆) alkyl.

[000126] In another embodiment of formula B-3:

R₁ is -H or methyl;

Y₁ is absent;

Y₂ is absent or is -O-, -NHO-, or -NH-; and

Z¹ is O or NO-Me; and

R^a is -H or Me.

[000127] In another embodiment of formula B-3:

R₁ and R^a, together with the atoms to which they are attached, form a

4-6 membered heterocycloalkyl ring optionally substituted with halo, (C₁-C₆) alkyl, or (C₁-C₆) haloalkyl.

[000128] In one embodiment of formula B-4:

Y₁ is absent;

Y₂ is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or

Y₂ is optionally substituted azetidiny;

Z¹ is O, NH, NO-(C₁-C₆) alkyl, or N-(C₁-C₆) alkyl; and

R^a is H, (C₁-C₆) alkyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NH(C₁-C₆) alkyl, -(C₂-C₆) alkylene-N((C₁-C₆) alkyl)₂, -(C₂-C₆) alkylene-optionally substituted 4-6 membered heterocycloalkyl, or optionally substituted 4-6 membered heterocycloalkyl;

R₂ is -H, -F, -Cl, -Br, -(C₁-C₆)alkoxy, -O-(C₂-C₆)alkylene-OH, -O-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl), (C₂-C₆)alkylene-O-(C₁-C₆) alkyl, -NH₂, -NH-(C₁-C₆ alkyl), -NH-(C₁-C₆)alkylene-(optionally substituted 4-6 membered heterocycloalkyl), or -NH-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl).

[000129] In another embodiment of formula B-4:

Y₁ is absent;

Y₂ is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or

Y₂ is optionally substituted azetidiny;

Z¹ is O, NH, NO-(C₁-C₆) alkyl, N-(C₁-C₆) alkyl; and


R^a is -H, methyl, ethyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NHMe, -(C₂-C₆) alkylene-N(Me)₂, -(C₁-C₆) alkylene-morpholiny, -(C₁-C₆) alkylene-piperidiny, (C₁-C₆)alkylene-(optionally substituted pyrrolidiny), optionally substituted azetidiny, or optionally substituted oxetany;

R₂ is -H, -F, -Cl, -Br, methoxy, -O-(C₂-C₆)alkylene-OH, -O-(C₂-C₆)alkylene-OMe, -NH₂, -NH-(C₁-C₆ alkyl), -NH-(C₂-C₆)alkylene-OMe, -NH-(C₂-C₆)alkylene-(optionally substituted morpholiny), or -NH-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl).

[000130] In another embodiment of formula B-4:

R₂ and R^a, together with the atoms to which they are attached, form a 4-6 membered heterocycloalkyl ring optionally substituted with halo, (C₁-C₆) alkyl, or (C₁-C₆) haloalkyl.

[000131] In another embodiment, the compound of formula B is a compound of formula B-5:

**B-5**

or a pharmaceutically acceptable salt thereof, wherein ring A in formula B-5 is an optionally substituted 5-6 membered heteroaryl or aryl.

[000132] In one embodiment of formula B-5:

Ring A is an optionally substituted (C₆-C₁₀) aryl, optionally substituted (C₃-C₁₀) cycloalkyl, optionally substituted 5-10 membered heteroaryl, or optionally substituted 4-10 membered heterocycloalkyl; and


R₂ is H, or (C₁-C₆)alkoxy.

[000133] In another embodiment of formula B-5:

Ring A is an optionally substituted phenyl, optionally substituted cyclopropyl, optionally substituted pyridyl, optionally substituted imidazolyl, optionally substituted pyrrolyl, optionally substituted furanyl, optionally substituted pyrazolyl, optionally substituted oxazolyl, optionally substituted azetidyl, or optionally substituted oxetanyl; and

R₂ is H, or methoxy.

[000134] Another embodiment of a compound of formula A and B is a compound of formula C:

**C**

or a pharmaceutically acceptable salt thereof, wherein:

Y₁ is absent;

Y₂ is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or

Y₂ is optionally substituted azetidiny;

Z is O, NH, NO-(C₁-C₆) alkyl, or N-(C₁-C₆) alkyl;

R^a is -H, (C₁-C₆) alkyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NH(C₁-C₆) alkyl, -(C₂-C₆) alkylene-N((C₁-C₆) alkyl)₂, -(C₂-C₆) alkylene-optionally substituted 4-6 membered heterocycloalkyl), or optionally substituted 4-6 membered heterocycloalkyl;

R₂ is -H, -F, -Cl, -Br, -(C₁-C₆)alkoxy, -O-(C₂-C₆)alkylene-OH, -O-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl), (C₂-C₆)alkylene-O-(C₁-C₆) alkyl, -NH₂, -NH-(C₁-C₆ alkyl), -NH-(C₁-C₆)alkylene-(optionally substituted 4-6 membered heterocycloalkyl), -NH-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl); and

n and m are each independently integers of 0 to 3

[000135] In another embodiment of formula C:

Y₁ is absent;

Y₂ is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or

Y₂ is optionally substituted azetidiny;

Z¹ is O, NH, NO-(C₁-C₆) alkyl, or N-(C₁-C₆) alkyl;

R^a is -H, methyl, ethyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NHMe, -(C₂-C₆) alkylene-N(Me)₂, -(C₁-C₆) alkylene-morpholinyl, -(C₁-C₆) alkylene-piperidinyl, (C₁-C₆)alkylene-(optionally substituted pyrrolidinyl), optionally substituted azetidiny, or optionally substituted oxetanyl;

R₂ is -H, -F, -Cl, -Br, methoxy, -O-(C₂-C₆)alkylene-OH, -O-(C₂-C₆)alkylene-OMe, -NH₂, -NH-(C₁-C₆ alkyl), -NH-(C₂-C₆)alkylene-OMe, -NH-(C₂-C₆)alkylene-(optionally substituted morpholinyl), or -NH-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl); and


and n and m or each 0 or 1.

[000136] In another embodiment of formula C:

R₂ and R^a, together with the atoms to which they are attached, form a 4-6 membered heterocycloalkyl ring optionally substituted with halo, (C₁-C₆) alkyl, and (C₁-C₆) haloalkyl; and

n and m are each independently integers of 0 to 3.

[000137] Another embodiment of formula C is a compound of formula C-1.

**C-1**

or a pharmaceutically acceptable salt thereof, wherein:

Y₂ is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or

Y₂ is optionally substituted azetidiny;


Z¹ is O, NH, NO-(C₁-C₆) alkyl, or N-(C₁-C₆) alkyl;

R^a is -H, methyl, ethyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NHMe, -(C₂-C₆) alkylene-N(Me)₂, -(C₁-C₆) alkylene-morpholinyl, -(C₁-C₆) alkylene-piperidinyl, (C₁-C₆)alkylene-(optionally substituted pyrrolidinyl), optionally substituted azetidiny, or optionally substituted oxetanyl;

R₂ is -H, -F, -Cl, -Br, methoxy, -O-(C₂-C₆)alkylene-OH, -O-(C₂-C₆)alkylene-OMe, -NH₂, -NH-(C₁-C₆ alkyl), -NH-(C₂-C₆)alkylene-OMe, -NH-(C₂-C₆)alkylene-(optionally substituted morpholinyl), or -NH-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl); and

and n and m are each indepenenently 0 or 1.

[000138] Another embodiment of formula A and B is a compound of formula D:

**D**

[000139] In one embodiment of formula D:

R₁ is -H or (C₁-C₆) alkyl; and

Y₁ is absent;

Y₂ is absent or is -O-, -NHO-, or -NH-; and

Z^1 is O or NO-(C₁-C₆) alkyl;

R^a is -H or -(C₁-C₆) alkyl; and

n and m are each independently integers of 0 to 3.

[000140] In another embodiment of formula D:

R_1 is -H or methyl;

Y_1 is absent;

Y_2 is absent or is -O-, -NHO-, or -NH-; and

Z^1 is O or NO-Me; and

R^a is -H, or -Me.


n and m are each independently integers of 0 to 1.

[000141] In another embodiment of formula D:

R_1 and R^a , together with the atoms to which they are attached, form a 4-6 membered heterocycloalkyl ring optionally substituted with halo, (C₁-C₆) alkyl, or (C₁-C₆) haloalkyl; and


n and m are each independently integers of 0 to 1.

[000142] Another embodiment of formula D is a compound of formula D-1.



D-1

[000143] Another embodiment of formula A and B is a compound of formula E:



E

or a pharmaceutically acceptable salt thereof, wherein:

Ring A is an optionally substituted (C₆-C₁₀) aryl, optionally substituted (C₃-C₁₀) cycloalkyl, optionally substituted 5-10 membered heteroaryl, or optionally substituted 4-10 membered heterocycloalkyl; and


R₂ is H or (C₁-C₆)alkoxy.

[000144] In another embodiment of formula E:

Ring A is an optionally substituted phenyl, optionally substituted cyclopropyl, optionally substituted pyridyl, optionally substituted imidazolyl, optionally substituted pyrrolyl, optionally substituted furanyl, optionally substituted pyrazolyl, optionally substituted oxazolyl, optionally substituted azetidyl, or optionally substituted oxetanyl; and

R₂ is H or methoxy.

[000145] Another embodiment of formula A and B is a compound of formula F:




F

or a pharmaceutically acceptable salt thereof, wherein


R^{al} is optionally substituted (C₁-C₆) alkyl;

R₁ is -H, -CN, optionally substituted 3-6 membered cycloalkyl, optionally substituted phenyl, optionally substituted 4-6 membered heterocycloalkyl, optionally substituted 5-6 membered heteroaryl, -SO₂-(C₁-C₆) alkyl, -SO₂NH₂, -SO₂-NH(C₁-C₆) alkyl, or P(O)((C₁-C₆) alkyl)₂, or



Y₁ is absent;

Y₂ is absent or is -O-, -NH-, -NHO-, -NH-NH-, -N-(C₁-C₆) alkyl-; or



Y₂ is optionally substituted , wherein ring A is a 3, 4, 5, 6, or 7-membered ring, wherein “wavy” indicates points of attachment;

Z^1 is O, NH, N-(C₁-C₆) alkyl, NHO, or NO-(C₁-C₆) alkyl; and

R^a is -H, -(C₁-C₆) alkyl, 4-6-membered heterocycloalkyl, 3-6-membered cycloalkyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NH(C₁-C₆), -(C₂-C₆) alkylene-N(C₁-C₆)₂, -(C₂-C₆) alkylene-N-(4-6-membered heterocycloalkyl);

[000146] In another embodiment of formula F:

R^{a1} is methyl;

R_1 is- H, -CN, optionally substituted cyclopropyl, optionally substituted phenyl, optionally substituted 4-6 membered azetidiny, optionally substituted pyrrolidiny, optionally substituted piperidiny, optionally substituted oxetany, optionally substituted oxazolyl, optionally substituted pyridiny, optionally substituted imidazolyl, optionally substituted pyrrolyl, optionally substituted furnayl, optionally substituted pyrazolyl, optionally substituted oxadiazolyl, -SO₂-(C₁-C₆) alkyl, -SO₂NH₂, -SO₂-NH(C₁-C₆) alkyl, or P(O)((C₁-C₆) alkyl)₂; or

R_1 is $\begin{array}{c} Z^1 \\ \parallel \\ R^a - Y_2 - Y_1 \end{array}$, wherein:

Y_1 is absent;


Y_2 is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or

Y_2 is optionally substituted azetidiny;

Z^1 is O, NH, or N-(C₁-C₆) alkyl; and

R^a is H, (C₁-C₆) alkyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NH(C₁-C₆) alkyl, -(C₂-C₆) alkylene-N((C₁-C₆) alkyl)₂, -(C₂-C₆) alkylene-heterocycloalkyl), and 4-6 membered heterocycloalkyl, wherein heterocycloalkyl is optionally substituted.

[000147] Another embodiment of formula A and B is a compound of formula G:

**G**

or a pharmaceutically acceptable salt thereof, wherein:

R^{a1} is optionally substituted (C₁-C₆) alkyl;

R_2 is $R^a - Y_2 - \overset{\overset{Z^1}{\parallel}}{C} - Y_1 - \text{wavy line}$ wherein “wavy line” indicates the point of attachment, wherein:

Y_1 is absent;

Y_2 is absent or is -O-, or -NH-; and

Z^1 is O; and

R^a is -H or -(C₁-C₆) alkyl.

[000148] In another embodiment of formula G:

R^{a2} is methyl;

R_2 is $R^a - Y_2 - \overset{\overset{Z^1}{\parallel}}{C} - Y_1 - \text{wavy line}$ wherein “wavy line” indicates the point of attachment, wherein:


Y_1 is absent;

Y_2 is absent or is -O-, or -NH-; and

Z^1 is O or NO-(C₁-C₆) alkyl; and

R^a is -H, or -(C₁-C₆) alkyl.

[000149] Another embodiment of formula A and B is a compound for modulating kinase activity according to Formula H:

**H**

or a pharmaceutically acceptable salt thereof, wherein:

R_1 is selected from the group consisting of -H, -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₁-C₆) alkyl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, or -(SO₂)(C₁-C₆) alkyl;



wherein when R_1 is selected from the group consisting of -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C₁-C₆) alkyl, R_2 is H, halo, NR₅R₆, or optionally substituted (C₁-C₆) alkoxy;

wherein when R_1 is -H, optionally substituted (C₁-C₆) alkyl, or optionally substituted (C₁-C₆) alkoxy, R_2 is -CO-NR₅R₆ or -CO₂R₇;

or R_1 and R_2 taken together with the atoms to which they are attached to form optionally substituted cycloalkyl or optionally substituted heterocycloalkyl;

R_3 is selected from the group consisting of -H, optionally substituted (C₁-C₆) alkyl, -CN, and halo;

R_4 is -H or halo;

 is optionally substituted with one, two, three, or four groups independently selected from the group consisting of halo and (C₁-C₆) alkyl, wherein “” indicate points of attachment;

R_5 and R_6 are each independently -H, optionally substituted (C₁-C₆) alkyl; or optionally substituted C₁-C₆ alkoxy;

R_7 is -H or optionally substituted (C₁-C₆) alkyl;


R_8 and R_9 are each independently -H or optionally substituted (C₁-C₆) alkyl; or

R_8 and R_9 may connect to form optionally substituted heterocycle; and

Y is selected from the group consisting of O, S, SO, SO₂, NH, and N-((C₁-C₆) alkyl).


[000150] In one embodiment of a compound of formula I, Y is O.


[000151] In another embodiment, R_3 is -H.












[000152] In another embodiment,  is not substituted.














[000153] In another embodiment, R_4 is halo.











- [000154] In another embodiment, R₄ is para fluoro.
- [000155] In another embodiment, R₂ is -H, halo, or optionally substituted (C₁-C₆)-alkoxy.
- [000156] In another embodiment, R₁ is -CN.
- [000157] In another embodiment, R₁ is -CO₂H.
- [000158] In another embodiment, R₁ is -CO₂-Me.
- [000159] In another embodiment, R₁ is -CO-NHR₆.
- [000160] In another embodiment, R₁ is -CO-NH₂.
- [000161] In another embodiment, R₁ is -CO-NMeR₆.
- [000162] In another embodiment, R₃ is -H or halo.
- [000163] In another embodiment, R₁ is selected from the group consisting of -CN, -



wherein “” is the point of attachment.

[000164] In another embodiment, R₁ is selected from the group consisting of , , , , , , , , , and , wherein “” is the point of attachment.

[000165] In another embodiment, R₁ is selected from the group consisting of , , , , , , , , , , , and , wherein “” is the point of attachment.

[000166] In another embodiment, R₂ is selected from the group consisting of -H, -CN, -Br, -F, -Cl, -OMe, -CH₃, , , , , -CH₂NH₂, NH₂, NHMe, , , , , and , wherein “” is the point of attachment.

[000167] In another embodiment, R₁ is -H, methyl, or methoxy.





[000168] In another embodiment, R₂ is -CO₂H.


[000169] In another embodiment, R₁ is -CO₂-Me.

[000170] In another embodiment, R₁ is -CO-NHR₆.


[000171] In another embodiment, R₁ is -CO-NH₂.

[000172] In another embodiment, R_1 is $-\text{CO}-\text{NMe}R_6$.

[000173] In another embodiment, R_1 is selected from the group consisting of , , , and .

[000174] In another embodiment, R_1 and R_2 are taken together to form .

[000175] In a further embodiment, the compound of formula I is a compound of formula I-1:











I-1







wherein R_6 is (C_1-C_6) alkyl, R_2 is (C_1-C_6) alkoxy, R_3 is $-\text{H}$ or halo, and R_4 is halo.







[000176] In another aspect, the invention provides a compound of formula A or A-I which is provided in Table 1 below.







[000177] **Table 1:** Specific compounds of the invention







No.	Structure	Name
5		methyl 4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylate
6		4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylic acid







No.	Structure	Name
7		1-N-[4-(6-carbamoyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
8		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
9		1-N-[4-[6-(ethylcarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
10		1-N-[4-[6-[2-(dimethylamino)ethylcarbamoyl]-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
11		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(2-piperidin-1-ylethylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
12		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(2-morpholin-4-ylethylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide







No.	Structure	Name
13		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(oxetan-3-yl)carbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
14		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(1-methylazetidin-3-yl)carbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
15		1-N-[4-[6-(azetidine-1-carbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
16		1-N'-(4-fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidine-1-carbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
17		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(methoxycarbonyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
20		1-N'-(4-fluorophenyl)-1-N-[4-[6-(hydroxycarbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide







No.	Structure	Name
21		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[[[(2R)-pyrrolidin-2-yl]methyl]carbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
22		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[[[(2S)-pyrrolidin-2-yl]methyl]carbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
26		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(oxetan-3-yl)oxycarbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
27		1-N'-(4-fluorophenyl)-1-N-[4-[6-(2-hydroxyethoxycarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
30		1-N-[4-[6-(2,3-dihydroxypropoxycarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide <i>Enantiomer 1</i>
31		1-N-[4-[6-(2,3-dihydroxypropoxycarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide <i>Enantiomer 2</i>







No.	Structure	Name
32		1-N'-(4-fluorophenyl)-1-N-[4-[6-(hydrazinecarbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
34		1-N-[4-(6-acetyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
35		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(E)-N-methoxy-C-methylcarbonimidoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
36		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(Z)-N-methoxy-C-methylcarbonimidoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
37		1-N-[4-(6-cyano-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
45		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide







No.	Structure	Name
50		1-N'-(4-fluorophenyl)-1-N-[4-[7-(2-hydroxyethoxy)-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
51		1-N-[4-(6-dimethylphosphoryl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
58		1-N-[4-(6-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
59		1-N'-(4-fluorophenyl)-1-N-[4-[6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
60		1-N'-(4-fluorophenyl)-1-N-[4-[6-[(1-methylazetidin-3-yl)carbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
67		1-N-[4-(6-carbamoyl-7-fluoroquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide







No.	Structure	Name
68		1-N-[4-(6-carbamoyl-7-chloroquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
69		1-N-[4-(7-bromo-6-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
70		1-N-[4-[6-carbamoyl-7-(2-methoxyethylamino)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
71		1-N-[4-[6-carbamoyl-7-(3-morpholin-4-ylpropylamino)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
72		1-N-[4-[7-(azetidin-1-yl)-6-carbamoylquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
81		4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-(methylamino)quinoline-6-carboxylic acid







No.	Structure	Name
82		1-N-[4-[6-carbamoyl-7-(methylamino)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
83		1-N'-(4-fluorophenyl)-1-N-[4-[7-(methylamino)-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
84		methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-(methylamino)quinoline-6-carboxylate
87		1-N-[4-(7-amino-6-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
88		1-N-[4-[7-amino-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
89		7-amino-4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]quinoline-6-carboxylic acid







No.	Structure	Name
90		methyl 7-amino-4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]quinoline-6-carboxylate
92		1-N'-(4-fluorophenyl)-1-N-[4-[(2-methyl-4-oxo-2,3-dihydropyrido[3,2-g][1,3]benzoxazin-6-yl)oxy]phenyl]cyclopropane-1,1-dicarboxamide
96		1-N-[4-[(2-ethyl-4-oxo-2,3-dihydropyrido[3,2-g][1,3]benzoxazin-6-yl)oxy]phenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
98		1-N-[4-[6-carbamoyl-7-(3-morpholin-4-ylpropoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
103		1-N-[4-[6-carbamoyl-7-(2-methoxyethoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
106		1-N-[4-[6-carbamoyl-7-(2-hydroxyethoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide







No.	Structure	Name
110		1-N'-(4-fluorophenyl)-1-N-[4-[7-(2-hydroxyethoxy)-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
115		1-N-[4-[6-carbamoyl-7-(2-hydroxypropoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
116		1-N'-(4-fluorophenyl)-1-N-[4-[7-(2-hydroxypropoxy)-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
125		1-N'-(4-fluorophenyl)-1-N-[4-[7-(2-hydroxypropoxy)-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
128		methyl 4-[2-chloro-4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylate
129		methyl 4-[2-fluoro-4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylate







No.	Structure	Name
130		4-[2-chloro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylic acid
131		4-[2-fluoro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylic acid
132		1-N'-[4-(6-carbamoyl-7-methoxyquinolin-4-yl)oxy-3-chlorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
133		1-N'-[4-(6-carbamoyl-7-methoxyquinolin-4-yl)oxy-3-fluorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
134		1-N'-[3-chloro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
135		1-N'-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide







No.	Structure	Name
140		methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-methylquinoline-7-carboxylate
141		4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-methylquinoline-7-carboxylic acid
142		1-N-[4-(7-carbamoyl-6-methylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
143		1-N'-(4-fluorophenyl)-1-N-[4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
150		methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-methoxyquinoline-7-carboxylate
151		4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-methoxyquinoline-7-carboxylic acid







No.	Structure	Name
152		1-N-[4-(7-carbamoyl-6-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
153		1-N'-(4-fluorophenyl)-1-N-[4-[6-methoxy-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
162		methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]quinoline-7-carboxylate
163		4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]quinoline-7-carboxylic acid
164		1-N-[4-(7-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
165		1-N'-(4-fluorophenyl)-1-N-[4-[7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide







No.	Structure	Name
166		1-N'-(4-fluorophenyl)-1-N-[4-[7-(2-hydroxyethoxycarbonyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
167		1-N'-(4-fluorophenyl)-1-N-[4-[7-(oxetan-3-yloxy)carbonyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
169		1-N-[4-[7-[(2R)-2,3-dihydroxypropoxy]carbonyl]quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
170		1-N-[4-[7-[(2S)-2,3-dihydroxypropoxy]carbonyl]quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
175		1-N-[4-[6-(3-cyano-2-fluorophenyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
176		1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-2-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide







No.	Structure	Name
177		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1-methylimidazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
180		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(5-methylfuran-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
181		tert-butyl 2-[4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinolin-6-yl]pyrrole-1-carboxylate
182		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1-methylpyrazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
183		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1,2-oxazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
184		1-N-[4-[6-(3,5-dimethyl-1,2-oxazol-4-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide







No.	Structure	Name
185		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrazol-5-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
186		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
187		1-N-[4-(6-cyclopropyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
188		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrrol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
191		1-N'-(4-fluorophenyl)-1-N-[4-[6-(1H-imidazol-2-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
192		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3-oxazol-5-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide







No.	Structure	Name
193		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(E)-methoxyiminomethyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
195		tert-butyl 3-[4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinolin-6-yl]-3-hydroxyazetidine-1-carboxylate
196		1-N'-(4-fluorophenyl)-1-N-[4-[6-(3-hydroxyoxetan-3-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
197		1-N'-(4-fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidin-3-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
198		1-N-[4-[6-(azetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
199		1-N'-(4-fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide







No.	Structure	Name
200		1-N-[4-[6-(3,3-difluoroazetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
201		1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-3-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide
202		1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-4-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide
204		1-N'-(4-fluorophenyl)-1-N-[4-[6-(1H-pyrazol-5-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
206		1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-sulfamoylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide
207		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(methylsulfamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide





No.	Structure	Name
208		1-N-[4-[6-(ethylsulfamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
209		1-N'-(4-fluorophenyl)-1-N-[4-(6-sulfamoylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide
210		1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-methylsulfonylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide
213		1-N'-(4-fluorophenyl)-1-N-[4-[7-(methoxycarbonyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
214		1-N-[4-[7-(ethylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
220		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3,4-oxadiazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide

No.	Structure	Name
221		1-N'-(4-fluorophenyl)-1-N-[4-[6-(1,3,4-oxadiazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
254		1-N'-(4-fluorophenyl)-1-N-[4-(7-sulfamoylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide
255		1-N-[4-(7-acetylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide; or
256		1-N'-(4-fluorophenyl)-1-N-[4-[7-[(E)-N-methoxy-C-methylcarbonimidoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
262		1-N-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)-1-N'-methylcyclopropane-1,1-dicarboxamide
263		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-methylcyclopropane-1,1-dicarboxamide

No.	Structure	Name
264		1-N'-(2-chloro-4-fluorophenyl)-1-N-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
265		1-N-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluoro-2-methylphenyl)cyclopropane-1,1-dicarboxamide
267		1-N'-(4-fluoro-2,6-dimethylphenyl)-1-N-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
268		1-N-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluoro-2-methoxyphenyl)cyclopropane-1,1-dicarboxamide
269		1-N-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluoro-2-(2-methoxypropan-2-yl)oxyphenyl)cyclopropane-1,1-dicarboxamide
270		1-N'-(2-cyclopropyl-4-fluorophenyl)-1-N-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide

No.	Structure	Name
273		1-N'-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-[4-(trifluoromethyl)phenyl]cyclopropane-1,1-dicarboxamide
274		1-N-(4-chlorophenyl)-1-N'-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
278		1-N'-(4-fluorophenyl)-1-N-[4-[7-[(E)-methoxyiminomethyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
279		1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(methylcarbamoylamino)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide
280		methyl N-[4-[4-[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinolin-6-yl]carbamate
281		1-N'-(4-fluorophenyl)-1-N-[4-[7-(methylcarbamoylamino)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide

No.	Structure	Name
282		methyl N-[4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]quinolin-7-yl]carbamate
283		1-N-[4-[6-(3-ethyl-1,2,4-oxadiazol-5-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
289		1-N'-[3-fluoro-4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
290		1-N'-[2,5-difluoro-4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
291		1-N'-[2-chloro-5-fluoro-4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
292		1-N-(4-fluorophenyl)-1-N'-[2,3,5-trifluoro-4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide

No.	Structure	Name
293		1-N'-[4-(7-carbamoyl-6-methylquinolin-4-yl)oxy-3-fluorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
294		1-N'-[4-(7-carbamoyl-6-methylquinolin-4-yl)oxy-2,5-difluorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
301		1-N'-[3-fluoro-4-[7-(2-hydroxyethoxycarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
302		1-N'-[2,5-difluoro-4-[7-(2-hydroxyethoxycarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide

or a pharmaceutically acceptable salt thereof.

[000178] General Administration

[000179] Administration of the compounds of the invention, or their pharmaceutically acceptable salts, in pure form or in an appropriate pharmaceutical composition, can be carried out via any of the accepted modes of administration or agents for serving similar utilities. Thus, administration can be, for example, orally, nasally, parenterally (intravenous, intramuscular, or subcutaneous), topically, transdermally, intravaginally, intravesically, intracistemally, or rectally, in the form of solid, semi-solid, lyophilized powder, or liquid dosage forms, such as, for example, tablets, suppositories, pills, soft elastic and hard gelatin capsules, powders,

solutions, suspensions, aerosols, and the like, preferably in unit dosage forms suitable for simple administration of precise dosages.

[000180] The compositions will include a conventional pharmaceutical carrier or excipient and a compound of the invention as the/an active agent, and, in addition, may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, and the like. Compositions of the invention may be used in combination with anticancer or other agents that are generally administered to a patient being treated for cancer. Adjuvants include preserving, wetting, suspending, sweetening, flavoring, perfuming, emulsifying, and dispensing agents. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate, and gelatin.

[000181] If desired, a pharmaceutical composition of the invention may also contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, antioxidants, and the like, such as, for example, citric acid, sorbitan monolaurate, triethanolamine oleate, butylated hydroxytoluene, and the like.

[000182] Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents, or vehicles include water, ethanol, polyols (propyleneglycol, polyethyleneglycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

[000183] One preferable route of administration is oral, using a convenient daily dosage regimen that can be adjusted according to the degree of severity of the disease-state to be treated.

[000184] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is admixed with at least one inert

customary excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, as for example, starches, lactose, sucrose, glucose, mannitol, and silicic acid, (b) binders, as for example, cellulose derivatives, starch, alginates, gelatin, polyvinylpyrrolidone, sucrose, and gum acacia, (c) humectants, as for example, glycerol, (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, croscarmellose sodium, complex silicates, and sodium carbonate, (e) solution retarders, as for example paraffin, (f) absorption accelerators, as for example, quaternary ammonium compounds, (g) wetting agents, as for example, cetyl alcohol, and glycerol monostearate, magnesium stearate, and the like (h) adsorbents, as for example, kaolin and bentonite, and (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents.

[000185] Solid dosage forms as described above can be prepared with coatings and shells, such as enteric coatings and others well known in the art. They may contain pacifying agents and can also be of such composition that they release the active compound or compounds in a certain part of the intestinal tract in a delayed manner. Examples of embedded compositions that can be used are polymeric substances and waxes. The active compounds can also be in microencapsulated form, if appropriate, with one or more of the above-mentioned excipients.

[000186] Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. Such dosage forms are prepared, for example, by dissolving, dispersing, and the like, a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol, and the like; solubilizing agents and emulsifiers, as for example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propyleneglycol, 1,3-butyleneglycol, and dimethylformamide; oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil, and sesame oil, glycerol, tetrahydrofurfuryl alcohol, polyethyleneglycols, and fatty acid esters of sorbitan; or mixtures of these substances, and the like, to thereby form a solution or suspension.

[000187] Suspensions, in addition to the active compounds, may contain suspending agents, as for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol, and sorbitan esters,

microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, or mixtures of these substances, and the like.

[000188] Compositions for rectal administrations are, for example, suppositories that can be prepared by mixing the compounds of the present invention with for example suitable non-irritating excipients or carriers such as cocoa butter, polyethyleneglycol, or a suppository wax, which are solid at ordinary temperatures but liquid at body temperature and therefore melt while in a suitable body cavity and release the active component therein.

[000189] Dosage forms for topical administration of a compound of this invention include ointments, powders, sprays, and inhalants. The active component is admixed under sterile conditions with a physiologically acceptable carrier and any preservatives, buffers, or propellants as may be required. Ophthalmic formulations, eye ointments, powders, and solutions are also contemplated as being within the scope of this invention.

[000190] Generally, depending on the intended mode of administration, the pharmaceutically acceptable compositions will contain about 1% to about 99% by weight of a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, and 99% to 1% by weight of a suitable pharmaceutical excipient. In one example, the composition will be between about 5% and about 75% by weight of a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, with the rest being suitable pharmaceutical excipients.

[000191] Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, 18th Ed., (Mack Publishing Company, Easton, Pa., 1990). The composition to be administered will, in any event, contain a therapeutically effective amount of a compound of the invention, or a pharmaceutically acceptable salt thereof, for treatment of a disease-state in accordance with the teachings of this invention.

[000192] The compounds of the invention, or their pharmaceutically acceptable salts, are administered in a therapeutically effective amount which will vary depending upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of the compound, the age, body weight, general health, sex, diet, mode, and time of administration, rate of excretion, drug combination, the severity of the particular disease-states, and the host undergoing therapy. The compounds of the present invention can be administered to a patient at dosage levels in the range of about 0.1 to about 1,000 mg per day.

For a normal human adult having a body weight of about 70 kilograms, a dosage in the range of about 0.01 to about 100 mg per kilogram of body weight per day is an example. The specific dosage used, however, can vary. For example, the dosage can depend on a number of factors including the requirements of the patient, the severity of the condition being treated, and the pharmacological activity of the compound being used. The determination of optimum dosages for a particular patient is well known to one of ordinary skill in the art.

[000193] Combination Therapy

[000194] A compound as disclosed herein can be administered as a single therapy or in combination (“co-administered”) with one or more additional therapies for the treatment of a disease or disorder, for instance a disease or disorder associated with hyper-proliferation such as cancer. Therapies that may be used in combination with a compound disclosed herein include: (i) surgery; (ii) radiotherapy (for example, gamma radiation, neutron beam radiotherapy, electron beam radiotherapy, proton therapy, brachytherapy, and systemic radioactive isotopes); (iii) endocrine therapy; (iv) adjuvant therapy, immunotherapy, CAR T-cell therapy; and (v) other chemotherapeutic agents.

[000195] The term “co-administered” (“co-administering”) refers to either simultaneous administration, or any manner of separate sequential administration, of a compound of Formula I’ or a salt thereof, and a further active pharmaceutical ingredient or ingredients, including cytotoxic agents and radiation treatment. If the administration is not simultaneous, the compounds are administered in a close time proximity to each other. Furthermore, it does not matter if the compounds are administered in the same dosage form, e.g. one compound may be administered topically and another compound may be administered orally.

[000196] Typically, any agent that has activity against a disease or condition being treated may be co-administered. Examples of such agents for cancer treatment can be found, for instance, at <https://www.cancer.gov/about-cancer/treatment/drugs> (last visited January 22, 2019) and in publically available sources such as Cancer Principles and Practice of Oncology by V. T. Devita and S. Hellman (editors), 11th edition (2018), Lippincott Williams & Wilkins Publishers. A person of ordinary skill in the art would be able to discern which combinations of agents would be useful based on the particular characteristics of the drugs and the disease involved.

[000197] In one embodiment, the treatment method includes the co-administration of a compound as disclosed herein or a pharmaceutically acceptable salt thereof and at least one immunotherapy. Immunotherapy (also called biological response modifier therapy, biologic therapy, biotherapy, immune therapy, or biological therapy) is treatment that uses parts of the immune system to fight disease. Immunotherapy can help the immune system recognize cancer cells, or enhance a response against cancer cells. Immunotherapies include active and passive immunotherapies. Active immunotherapies stimulate the body's own immune system while passive immunotherapies generally use immune system components created outside of the body.

[000198] Examples of active immunotherapies include, but are not limited to vaccines including cancer vaccines, tumor cell vaccines (autologous or allogeneic), dendritic cell vaccines, antigen vaccines, anti-idiotypic vaccines, DNA vaccines, viral vaccines, or Tumor-Infiltrating Lymphocyte (TIL) Vaccine with Interleukin-2 (IL-2) or Lymphokine-Activated Killer (LAK) Cell Therapy.

[000199] Examples of passive immunotherapies include but are not limited to monoclonal antibodies and targeted therapies containing toxins. Monoclonal antibodies include naked antibodies and conjugated monoclonal antibodies (also called tagged, labeled, or loaded antibodies). Naked monoclonal antibodies do not have a drug or radioactive material attached whereas conjugated monoclonal antibodies are joined to, for example, a chemotherapy drug (chemolabeled), a radioactive particle (radiolabeled), or a toxin (immunotoxin). Examples of these naked monoclonal antibody drugs include, but are not limited to Rituximab (Rituxan), an antibody against the CD20 antigen used to treat, for example, B cell non-Hodgkin lymphoma; Trastuzumab (Herceptin), an antibody against the HER2 protein used to treat, for example, advanced breast cancer; Alemtuzumab (Campath), an antibody against the CD52 antigen used to treat, for example, B cell chronic lymphocytic leukemia (B-CLL); Cetuximab (Erbix), an antibody against the EGFR protein used, for example, in combination with irinotecan to treat, for example, advanced colorectal cancer and head and neck cancers; and Bevacizumab (Avastin) which is an antiangiogenesis therapy that works against the VEGF protein and is used, for example, in combination with chemotherapy to treat, for example, metastatic colorectal cancer. Examples of the conjugated monoclonal antibodies include, but are not limited to Radiolabeled antibody Ibritumomab tiuxetan (Zevalin) which delivers radioactivity

directly to cancerous B lymphocytes and is used to treat, for example, B cell non-Hodgkin lymphoma; radiolabeled antibody Tositumomab (Bexxar) which is used to treat, for example, certain types of non-Hodgkin lymphoma; and immunotoxin Gemtuzumab ozogamicin (Mylotarg) which contains calicheamicin and is used to treat, for example, acute myelogenous leukemia (AML). BL22 is a conjugated monoclonal antibody for treating, for example, hairy cell leukemia, immunotoxins for treating, for example, leukemias, lymphomas, and brain tumors, and radiolabeled antibodies such as OncoScint for example, for colorectal and ovarian cancers and ProstaScint for example, for prostate cancers.

[000200] Further examples of therapeutic antibodies that can be used include, but are not limited to, HERCEPTINTM (Trastuzumab) (Genentech, Calif.) which is a humanized anti-HER2 monoclonal antibody for the treatment of patients with metastatic breast cancer; REOPROTM (abciximab) (Centocor) which is an anti-glycoprotein IIb/IIIa receptor on the platelets for the prevention of clot formation; ZENAPAXTM (daclizumab) (Roche Pharmaceuticals, Switzerland) which is an immunosuppressive, humanized anti-CD25 monoclonal antibody for the prevention of acute renal allograft rejection; PANOREXTM which is a murine anti-17-IA cell surface antigen IgG2a antibody (Glaxo Wellcome/Centocor); BEC2 which is a murine anti-idiotypic (GD3epitope) IgG antibody (ImClone System); IMC-C225 which is a chimeric anti-EGFR IgG antibody (ImClone System); VITAXINTM which is a humanized anti-alpha V beta 3 integrin antibody (Applied Molecular Evolution/MedImmune); Campath 1H/LDP-03 which is a humanized anti CD52 IgG1 antibody (Leukosite); Smart M195 which is a humanized anti-CD33 IgG antibody (Protein Design Lab/Kanebo); RITUXANTM which is a chimeric anti-CD20 IgG1 antibody (IDEC Pharm/Genentech, Roche/Zettyaku); LYMPHOCIDETM which is a humanized anti-CD22 IgG antibody (Immunomedics); LYMPHOCIDETM Y-90 (Immunomedics); Lymphoscan (Tc-99m-labeled; radioimaging; Immunomedics); Nuvion (against CD3; Protein Design Labs); CM3 is a humanized anti-ICAM3 antibody (ICOS Pharm); IDEC-114 is a primatized anti-CD80 antibody (IDEC Pharm/Mitsubishi); ZEVALINTM is a radiolabelled murine anti-CD20 antibody (IDEC/Schering AG); IDEC-131 is a humanized anti-CD40L antibody (IDEC/Eisai); IDEC-151 is a primatized anti-CD4 antibody (IDEC); IDEC-152 is a primatized anti-CD23 antibody (IDEC/Seikagaku); SMART anti-CD3 is a humanized anti-CD3 IgG (Protein Design Lab); 5G1.1 is a humanized anti-complement factor 5 (C5) antibody (Alexion Pharm); D2E7 is a humanized anti-TNF-alpha

antibody (CAT/BASF); CDP870 is a humanized anti-TNF-alpha. Fab fragment (Celltech); IDEC-151 is a primatized anti-CD4 IgG1 antibody (IDEC Pharm/SmithKline Beecham); MDX-CD4 is a human anti-CD4 IgG antibody (Medarex/Eisai/Genmab); CD20-sreptavidin (+biotin-⁹⁰yttrium 90; NeoRx); CDP571 is a humanized anti-TNF-alpha. IgG4 antibody (Celltech); LDP-02 is a humanized anti-alpha4 beta7 antibody (LeukoSite/Genentech); OrthoClone OKT4A is a humanized anti-CD4 IgG antibody (Ortho Biotech); ANTOVA.TM. is a humanized anti-CD40L IgG antibody (Biogen); ANTEGREN™ is a humanized anti-VLA-4 IgG antibody (Elan); and CAT-152 is a human anti-TGF-beta₂ antibody (Cambridge Ab Tech). Others are provided in later paragraphs.

[000201] Immunotherapies that can be used in combination with a compound as disclosed herein include adjuvant immunotherapies. Examples include cytokines, such as granulocyte-macrophage colony-stimulating factor (GM-CSF), granulocyte-colony stimulating factor (G-CSF), macrophage inflammatory protein (MIP)-1-alpha, interleukins (including IL-1, IL-2, IL-4, IL-6, IL-7, IL-12, IL-15, IL-18, IL-21, and IL-27), tumor necrosis factors (including TNF-alpha), and interferons (including IFN-alpha, IFN-beta, and IFN-gamma); aluminum hydroxide (alum); Bacille Calmette-Guerin (BCG); Keyhole limpet hemocyanin (KLH); Incomplete Freund's adjuvant (IFA); QS-21; DETOX; Levamisole; and Dinitrophenyl (DNP), and combinations thereof, such as, for example, combinations of, interleukins, for example, IL-2 with other cytokines, such as IFN-alpha.

[000202] In various embodiments, an immunological therapy or an immunological therapeutic agent can include, one or more of the following: an adoptive cell transfer, an angiogenesis inhibitor, Bacillus Calmette-Guerin therapy, biochemotherapy, a cancer vaccine, a chimeric antigen receptor (CAR) T-cell therapy, a cytokine therapy, gene therapy, an immune checkpoint modulator, an immunoconjugate, a radioconjugate, an oncolytic virus therapy, or a targeted drug therapy. The function or at least one of the functions of the immunological therapy or immunological therapeutic agent, collectively referred to herein as an “immunotherapeutic agent”.

[000203] The present disclosure provides a method for preventing, treating, reducing, inhibiting or controlling a neoplasia, a tumor or a cancer in a subject in need thereof, involving administering a therapeutically effective amount of a combination comprising a compound of Formula I' and an immunotherapeutic agent. In one non-limiting embodiment, the method

comprises administering a therapeutically effective amount of a combination comprising a compound of Formula I' in combination with an immunotherapeutic agent. In various embodiments, the combination provides a cooperative effect, an additive effect, or a synergistic effect in reducing the number of cancer cells when treated with the combination as compared to each treatment alone. In some embodiments, administration of a therapeutically effective amount of a combination comprising a compound of Formula I' and an immunotherapeutic agent, results in synergistic anti-tumor activity and/or antitumor activity that is more potent than the additive effect of administration of a compound of Formula I' or immunotherapeutic agent alone.

[000204] Human cancers harbor numerous genetic and epigenetic alterations, generating neoantigens potentially recognizable by the immune system (Sjoblom et al. (2006) Science 314:268-74). The adaptive immune system, comprised of T and B lymphocytes, has powerful anti-cancer potential, with a broad capacity and exquisite specificity to respond to diverse tumor antigens. Further, the immune system demonstrates considerable plasticity and a memory component. The successful harnessing of all these attributes of the adaptive immune system would make immunotherapy unique among all cancer treatment modalities.

[000205] The present disclosure provides a combination of a compound of Formula I' and an immunotherapeutic agent. These exemplified combinations can be used to treat a subject with a cancer. In various embodiments, immunotherapeutic agents that find utility in the present compositions, formulations, and methods can include one or more agents or therapies, including: an adoptive cell transfer, an angiogenesis inhibitor, Bacillus Calmette-Guerin therapy, biochemotherapy, a cancer vaccine, a chimeric antigen receptor (CAR) T-cell therapy, a cytokine therapy, gene therapy, an immune checkpoint modulator, for example an immune checkpoint inhibitor, an immunoconjugate, a radioconjugate, an oncolytic virus therapy, or a targeted drug therapy.

[000206] In certain embodiments of the present disclosure, a therapeutically effective combination comprises a compound of Formula I' and an immunotherapeutic agent. In various related embodiments, the compound of Formula I' enhances the activity of the immunotherapeutic agent.

[000207] In certain embodiments of each of the aforementioned aspects, as well as other aspects and embodiments described elsewhere herein, the immunotherapeutic agent enhances the activity of the compound of Formula I'.

[000208] In certain embodiments of each of the aforementioned aspects, as well as other aspects and embodiments described elsewhere herein, the compound of Formula I' and the immunotherapeutic agent act synergistically. In various embodiments described herein, an exemplary immunotherapeutic agent is an immune cell (e.g. T-cell, dendritic cell, a natural killer cell and the like) modulator chosen from an agonist or an activator of a costimulatory molecule, wherein the modulator is a monoclonal antibody, a bispecific antibody comprising one or more immune checkpoint antigen binding moieties, a trispecific antibody, or an immune cell-engaging multivalent antibody/fusion protein/construct known in the art). In some embodiments, the immunotherapeutic agent can be an antibody that modulates a costimulatory molecule, bind to an antigen on the surface of an immune cell, or a cancer cell. In each of these different embodiments, the antibody modulator can be a monoclonal antibody, a polyclonal antibody, a bispecific antibody, a trispecific or multispecific format antibody, a fusion protein, or a fragment thereof, for example, a Diabody, a Single-chain (sc)-diabody (scFv)₂, a Miniantibody, a Minibody, a Barnase-barstar, a scFv-Fc, a sc(Fab)₂, a Trimeric antibody construct, a Triabody antibody construct, a Trimerbody antibody construct, a Tribody antibody construct, a Collabody antibody construct, a (scFv-TNFα)₃, or a F(ab)₃/DNL antibody construct.

[000209] In certain embodiments of each of the aforementioned aspects, as well as other aspects and embodiments described elsewhere herein, the immunotherapeutic agent is an agent that modulates immune responses, for example, a checkpoint inhibitor or a checkpoint agonist. In some embodiments, the immunotherapeutic agent is an agent that enhances anti-tumor immune responses. In some embodiments, the immunotherapeutic agent is an agent that increases cell-mediated immunity. In some embodiments, the immunotherapeutic agent is an agent that increases T-cell activity. In some embodiments, the immunotherapeutic agent is an agent that increases cytolytic T-cell (CTL) activity. In some embodiments, the immunotherapeutic agent is an antibody modulator that targets PD-1, PD-L1, PD-L2, CEACAM (e.g., CEACAM-1, -3 and/or -5), CTLA-4, TIM-3, LAG-3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4, TGF beta, OX40, 41BB, LIGHT, CD40, GITR, TGF-beta, TIM-3, SIRP-

alpha, VSIG8, BTLA, SIGLEC7, SIGLEC9, ICOS, B7H3, B7H4, FAS, and/or BTNL2 among others known in the art. In some embodiments, the immunotherapeutic agent is an agent that increases natural killer (NK) cell activity. In some embodiments, the immunotherapeutic agent is an agent that inhibits suppression of an immune response. In some embodiments, the immunotherapeutic agent is an agent that inhibits suppressor cells or suppressor cell activity. In some embodiments, the immunotherapeutic agent is an agent or therapy that inhibits Treg activity. In some embodiments, the immunotherapeutic agent is an agent that inhibits the activity of inhibitory immune checkpoint receptors. In some embodiments, the combination of the present disclosure comprises a compound of Formula I' and an immunotherapeutic agent, wherein the immunotherapeutic agent includes a T cell modulator chosen from an agonist or an activator of a costimulatory molecule. In one embodiment, the agonist of the costimulatory molecule is chosen from an agonist (e.g., an agonistic antibody or antigen-binding fragment thereof, or a soluble fusion) of GITR, OX40, ICOS, SLAM (e.g., SLAMF7), HVEM, LIGHT, CD2, CD27, CD28, CDS, ICAM-1, LFA-1 (CD11a/CD18), ICOS (CD278), 4-1BB (CD137), CD30, CD40, BAFFR, CD7, NKG2C, NKp80, CD160, B7-H3, or CD83 ligand. In other embodiments, the effector cell combination includes a bispecific T cell engager (e.g., a bispecific antibody molecule that binds to CD3 and a tumor antigen (e.g., EGFR, PSCA, PSMA, EpCAM, HER2 among others).

[000210] In some embodiments, the immunotherapeutic agent is a modulator of PD-1 activity, a modulator of PD-L1 activity, a modulator of PD-L2 activity, a modulator of CTLA-4 activity, a modulator of CD28 activity, a modulator of CD80 activity, a modulator of CD86 activity, a modulator of 4-1BB activity, an modulator of OX40 activity, a modulator of KIR activity, a modulator of Tim-3 activity, a modulator of LAG3 activity, a modulator of CD27 activity, a modulator of CD40 activity, a modulator of GITR activity, a modulator of TIGIT activity, a modulator of CD20 activity, a modulator of CD96 activity, a modulator of IDO1 activity, a modulator of SIRP-alpha activity, a modulator of TIGIT activity, a modulator of VSIG8 activity, a modulator of BTLA activity, a modulator of SIGLEC7 activity, a modulator of SIGLEC9 activity, a modulator of ICOS activity, a modulator of B7H3 activity, a modulator of B7H4 activity, a modulator of FAS activity, a modulator of BTNL2 activity, a cytokine, a chemokine, an interferon, an interleukin, a lymphokine, a member of the tumor necrosis factor (TNF) family, or an immunostimulatory oligonucleotide. In some embodiments, the

immunotherapeutic agent is an immune checkpoint modulator (e.g., an immune checkpoint inhibitor e.g. an inhibitor of PD-1 activity, a modulator of PD-L1 activity, a modulator of PD-L2 activity, a modulator of CTLA-4, or a CD40 agonist (e.g., an anti-CD40 antibody molecule), (xi) an OX40 agonist (e.g., an anti-OX40 antibody molecule), or (xii) a CD27 agonist (e.g., an anti-CD27 antibody molecule). In one embodiment, the immunomodulator is an inhibitor of PD-1, PD-L1, PD-L2, CTLA-4, TIM-3, LAG-3, CEACAM (e.g., CEACAM-1, -3 and/or -5), VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4 and/or TGF beta. In one embodiment, the inhibitor of an immune checkpoint molecule inhibits PD-1, PD-L1, LAG-3, TIM-3, CEACAM (e.g., CEACAM-1, -3 and/or -5), CTLA-4, or any combination thereof.

[000211] Inhibition of an inhibitory molecule can be performed at the DNA, RNA or protein level. In embodiments, an inhibitory nucleic acid (e.g., a dsRNA, siRNA or shRNA), can be used to inhibit expression of an inhibitory molecule. In other embodiments, the inhibitor of an inhibitory signal is, a polypeptide e.g., a soluble ligand (e.g., PD-1-Ig or CTLA-4 Ig), or an antibody or antigen-binding fragment thereof, for example, a monoclonal antibody, a bispecific antibody comprising one or more immune checkpoint antigen binding moieties, a trispecific antibody, or an immune cell-engaging multivalent antibody/fusion protein/construct known in the art that binds to the inhibitory molecule; e.g., an antibody or fragment thereof (also referred to herein as "an antibody molecule") that binds to PD-1, PD-L1, PD-L2, CEACAM (e.g., CEACAM-1, -3 and/or -5), CTLA-4, TIM-3, LAG-3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4, TGF beta, or a combination thereof.

[000212] In some embodiments, where the combination comprises a compound of Formula I' and an immunotherapeutic agent, wherein the immunotherapeutic agent is a monoclonal antibody or a bispecific antibody. For example, the monoclonal or bispecific antibody may specifically bind a member of the c-Met pathway and/or an immune checkpoint modulator (e.g., the bispecific antibody binds to both a hepatocyte growth factor receptor (HGFR) and an immune checkpoint modulator described herein, such as an antibody that binds PD-1, PD-L1, PD-L2, or CTLA-4, LAG-3, OX40, 41BB, LIGHT, CD40, GITR, TGF-beta, TIM-3, SIRP-alpha, TIGIT, VSIG8, BTLA, SIGLEC7, SIGLEC9, ICOS, B7H3, B7H4, FAS, BTNL2 or CD27). In particular embodiments, the bispecific antibody specifically binds a human HGFR protein and one of PD-1, PD-L1, and CTLA-4.

[000213] In some embodiments, the immunotherapeutic agent is a cytokine, for example, a chemokine, an interferon, an interleukin, lymphokine, or a member of the tumor necrosis factor family. In some embodiments, the cytokine is IL-2, IL15, or interferon-gamma.

[000214] In some embodiments of any of the above aspects or those described elsewhere herein, the cancer is selected from the group consisting of lung cancer, pancreatic cancer, breast cancer, colon cancer, colorectal cancer, melanoma, gastrointestinal cancer, gastric cancer, renal cancer, ovarian cancer, liver cancer, endometrial cancer, kidney cancer, prostate cancer, thyroid cancer, neuroblastoma, glioma, glioblastoma, glioblastoma multiforme, cervical cancer, stomach cancer, bladder cancer, head and neck cancer, and hepatoma.

[000215] In some embodiments of any of the above aspects or those described elsewhere herein, the subject's cancer or tumor does not respond to immune checkpoint inhibition (e.g., to any immune checkpoint inhibitor described herein, such as a PD-1 antagonist or PD-L1 antagonist) or the subject's cancer or tumor has progressed following an initial response to immune checkpoint inhibition (e.g., to any immune checkpoint inhibitor described herein, such as a PD-1 antagonist or PD-L1 antagonist).

[000216] In some embodiments of any of the above aspects or those described elsewhere herein, the subject is a human.

[000217] A checkpoint inhibitor can be any molecule, agent, treatment and/or method of inhibiting an immune checkpoint, and/or promoting an inhibitor of an immune checkpoint, e.g., by promoting an intrinsic immune checkpoint inhibitor; inhibiting a transcription factor involved in the expression of an immune checkpoint; and/or by acting in concert with some additional extrinsic factor. For example, a checkpoint inhibitor could include a treatment that inhibits transcription factors involved the expression of immune checkpoint genes, or promotes the expression of transcription factors for tumor-suppressor genes, e.g., BACH2 (Luan et al., (2016). Transcription Factors and Checkpoint Inhibitor Expression with Age: Markers of Immunosenescence. Blood, 128(22), 5983). Moreover, a checkpoint inhibitor can inhibit the transcription of immune checkpoint genes; the modification and/or processing of immune checkpoint mRNA; the translation of immune checkpoint proteins; and/or molecules involved in immunity or the immune checkpoint pathway, e.g., PD-1 transcription factors such as HIF-1, STAT3, NF- κ B, and AP-1, or the activation of common oncogenic pathways such as JAK/STAT, RAS/ERK, or PI3K/AKT/mTOR (Zerdes et al., Genetic, transcriptional and post-

translational regulation of the programmed death protein ligand 1 in cancer: biology and clinical correlations, *Oncogene* volume 37, pages 4639–4661 (2018), the disclosure of which is incorporated herein by reference in its entirety).

[000218] Checkpoint inhibitors can include treatments, molecules, agents, and/or methods that regulate immune checkpoints at the transcriptional level, e.g., using the RNA-interference pathway co-suppression, and/or post-transcriptional gene silencing (PTGS) (e.g., microRNAs, miRNA; silencing-RNA, small-interfering-RNA, or short-interfering-RNA (siRNA).

Transcriptional regulation of checkpoint molecules has been shown to involve mir-16, which has been shown to target the 3'UTR of the checkpoint mRNAs CD80, CD274 (PD-L1) and CD40 (Leibowitz et al., Post-transcriptional regulation of immune checkpoint genes by mir-16 in melanoma, *Annals of Oncology* (2017) 28; v428-v448). Mir-33a has also been shown to be involved in regulating the expression of PD-1 in cases of lung adenocarcinoma (Boldini et al., Role of microRNA-33a in regulating the expression of PD-1 in lung adenocarcinoma, *Cancer Cell Int.* 2017; 17: 105, the disclosure of which is incorporated herein by reference in its entirety).

[000219] T-cell-specific aptamer–siRNA chimeras have been suggested as a highly specific method of inhibiting molecules in the immune checkpoint pathway (Hossain et al., The aptamer–siRNA conjugates: reprogramming T cells for cancer therapy, *Ther. Deliv.* 2015 Jan; 6(1): 1–4, the disclosure of which is incorporated herein by reference in its entirety).

[000220] Alternatively, members of the immune checkpoint pathway can be inhibited using treatments that affect associated pathways, e.g., metabolism. For example, oversupplying the glycolytic intermediate pyruvate in mitochondria from CAD macrophages promoted expression of PD-L1 via induction of the bone morphogenetic protein 4/phosphorylated SMAD1/5/IFN regulatory factor 1 (BMP4/p-SMAD1/5/IRF1) signaling pathway. Accordingly, implementing treatments that modulate the metabolic pathway can result in subsequent modulation of the immunoinhibitory PD-1/PD-L1 checkpoint pathway (Watanabe et al., Pyruvate controls the checkpoint inhibitor PD-L1 and suppresses T cell immunity, *J Clin Invest.* 2017 Jun 30; 127(7): 2725–2738).

[000221] Checkpoint immunity can be regulated via oncolytic viruses that selectively replicate within tumor cells and induce acute immune responses in the tumor-micro-environment, i.e., by acting as genetic vectors that carry specific agents (e.g., antibodies,

miRNA, siRNA, and the like) to cancer cells and effecting their oncolysis and secretion of cytokines and chemokines to synergize with immune checkpoint inhibition (Shi et al., Cancer Immunotherapy: A Focus on the Regulation of Immune Checkpoints, *Int J Mol Sci.* 2018 May; 19(5): 1389). Currently, there are clinical trials underway that utilize the following viruses as checkpoint inhibitors: poliovirus, measles virus, adenoviruses, poxviruses, herpes simplex virus (HSV), coxsackieviruses, reovirus, Newcastle disease virus (NDV), T-VEC (a herpes virus encoded with GM-CSF (granulocyte-macrophage colony stimulating factor)), and H101 (Shi et al., *supra*).

[000222] Checkpoint inhibitors can operate at the translational level of checkpoint immunity. The translation of mRNA into protein represents a key event in the regulation of gene expression, thus inhibition of immune checkpoint translation is a method in which the immune checkpoint pathway can be inhibited.

[000223] Inhibition of the immune checkpoint pathway can occur at any stage of the immune checkpoint translational process. For example, drugs, molecules, agents, treatments, and/or methods can inhibit the initiation process (whereby the 40S ribosomal subunit is recruited to the 5' end of the mRNA and scans the 5'UTR of the mRNA toward its 3' end. Inhibition can occur by targeting the anticodon of the initiator methionyl-transfer RNA (tRNA) (Met-tRNA_i), its base-pairing with the start codon, or the recruitment of the 60S subunit to begin elongation and sequential addition of amino acids in the translation of immune-checkpoint-specific genes. Alternatively, a checkpoint inhibitor can inhibit checkpoints at the translational level by preventing the formation of the ternary complex (TC), i.e., eukaryotic initiation factor (eIF)2 (or one or more of its α , β , and γ subunits); GTP; and Met-tRNA_i.

[000224] Checkpoint inhibition can occur via destabilization of eIF2 α by precluding its phosphorylation via protein kinase R (PKR), PERK, GCN2, or HRI, or by precluding TCs from associating with the 40S ribosome and/or other initiation factors, thus preventing the preinitiation complex (PIC) from forming; inhibiting the eIF4F complex and/or its cap-binding protein eIF4E, the scaffolding protein eIF4G, or eIF4A helicase. Methods discussing the translational control of cancer are discussed in Truitt et al., *New frontiers in translational control of the cancer genome*, *Nat Rev Cancer.* 2016 Apr 26; 16(5): 288–304, the disclosure of which is incorporated herein by reference in its entirety.

[000225] Checkpoint inhibitors can also include treatments, molecules, agents, and/or methods that regulate immune checkpoints at the cellular and/or protein level, e.g., by inhibiting an immune checkpoint receptor. Inhibition of checkpoints can occur via the use of antibodies, antibody fragments, antigen-binding fragments, small-molecules, and/or other drugs, agents, treatments, and/or methods.

[000226] Immune checkpoints refer to inhibitory pathways in the immune system that are responsible for maintaining self-tolerance and modulating the degree of immune system response to minimize peripheral tissue damage. However, tumor cells can also activate immune system checkpoints to decrease the effectiveness of immune response ('block' the immune response) against tumor tissues. In contrast to the majority of anti-cancer agents, checkpoint inhibitors do not target tumor cells directly, but rather target lymphocyte receptors or their ligands in order to enhance the endogenous antitumor activity of the immune system. (Pardoll, 2012, *Nature Reviews Cancer* 12:252-264).

[000227] Until recently, cancer immunotherapy had focused substantial effort on approaches that enhance anti-tumor immune responses by adoptive-transfer of activated effector cells, immunization against relevant antigens, or providing non-specific immune-stimulatory agents such as cytokines. In the past decade, however, intensive efforts to develop specific immune checkpoint pathway inhibitors have begun to provide new immunotherapeutic approaches for treating cancer, including the development of antibody (Ab), ipilimumab (YERVOY.RTM.), that binds to and inhibits CTLA-4 for the treatment of patients with advanced melanoma (Hodi et al. (2010) *N Engl J Med* 363:711-23) and the development of antibodies such as nivolumab and pembrolizumab (formerly lambrolizumab; USAN Council Statement (2013)

Pembrolizumab: Statement on a nonproprietary name adopted by the USAN Council (ZZ-165), Nov. 27, 2013) that bind specifically to the Programmed Death-1 (PD-1) receptor and block the inhibitory PD-1/PD-1 ligand pathway (Topalian et al. (2012a) *N Engl J Med* 366:2443-54; Topalian et al. (2012b) *Curr Opin Immunol* 24:207-12; Topalian et al. (2014) *J Clin Oncol* 32(10):1020-30; Hamid et al. (2013) *N Engl J Med* 369:134-144; Hamid and Carvajal (2013) *Expert Opin Biol Ther* 13(6):847-61; McDermott and Atkins (2013) *Cancer Med* 2(5):662-73).

[000228] PD-1 is a key immune checkpoint receptor expressed by activated T and B cells and mediates immunosuppression. Nivolumab (formerly designated 5C4, BMS-936558, MDX-1106, or ONO-4538) is a fully human IgG4 (S228P) PD-1 immune checkpoint inhibitor

antibody that selectively prevents interaction with PD-1 ligands (PD-L1 and PD-L2), thereby blocking the down-regulation of antitumor T-cell functions (U.S. Pat. No. 8,008,449; Wang et al. (2014) In vitro characterization of the anti-PD-1 antibody nivolumab, BMS-936558, and in vivo toxicology in non-human primates. Nivolumab has been approved for the treatment of patients with unresectable or metastatic melanoma and disease progression following ipilimumab and, if BRAF V600 mutation positive, a BRAF inhibitor and for the treatment of squamous non-small cell lung cancer.

[000229] Recent data suggest a secondary mechanism of anti-CTLA-4 antibodies, which could occur within the tumor itself. CTLA-4 has been found to be expressed in tumors at higher levels on regulatory T-cells (also referred to herein as "Treg cells") as compared with intra-tumoral effector T-cells (also referred to herein as "Teff cells"), resulting in the hypothesis of anti-CTLA-4 preferentially impacting the Treg cell. "Therapeutic use of anti-CTLA-4 antibodies", Christian U. Blank and Alexander Enk, International Immunology, Vol. 27, No. 1, pp. 3-10. A recent study of a PD-1 and CTLA-4 combination show that the combination blockade of the CTLA-4 and PD-1 pathways also cooperates to increase the ratio of Teff cells to both regulatory T-cells and MDSCs, thereby reducing suppression and promoting inflammation in the tumor microenvironment. "Combination of CTLA-4 and PD-1 blockade expands infiltrating T-cells and reduces regulatory T and myeloid cells within B16 melanoma tumors", Curran et al., PNAS|Mar. 2, 2010; vol. 107 (no. 9); pp. 4275-4280, the disclosure of which is incorporated herein by reference in its entirety. The combination of a checkpoint inhibitor and another therapeutic agent(s) may enhance or prolong anti-tumor response of the checkpoint inhibitor and/or effects of the therapeutic agent. In this regard, WO 2015/069770 discloses a combination treatment based on activating the adaptive immune response, in particular the combination of CTLA-4 and PD-1 inhibitors, for the treatment of cancer. The disclosure of WO 2015/069770 is incorporated by reference in its entirety in the disclosure of this application.

[000230] One mechanism by which the checkpoint blockade anti-CTLA-4 antibodies mediate anti-tumor effect is by decreasing regulatory T-cells. Due to the distinct mechanism of action of anti-CTLA-4 antibodies, they can successfully combine with the anti-PD1 checkpoint blockade antibodies which work to release the suppressive signaling conferred to effector T-cells. Dual blockade with these antibodies combine to improve anti-tumor response both

preclinically (Proc Natl Acad Sci USA 2010, 107, 4275-4280) and in the clinic (N Engl J Med 2013, 369, 122-133; N Engl J Med 2015, 372, 2006-2017).

[000231] CTLA-4 attenuates the early activation of naïve and memory T cells through interactions with its ligands B7-1 (CD80) and B7-2 (CD86) (Fig. 1A). PD-1 is a receptor expressed on the surface of activated mature T cells, activated NK cells, B cells, monocytes and multiple normal tissues and plays a crucial role in the maintenance of peripheral tolerance [20–21] (Fig. 1A). In contrast to CTLA-4, PD-1 acts via interactions with its ligands PD-L1 (also known as B7-H1 or CD274) and is involved mainly in T cell activity modulation in peripheral tissues as well as providing a major immune resistance mechanism within the tumor microenvironment.

[000232] In some embodiments, the immunotherapeutic agent is a modulator of PD-1 activity, a modulator of PD-L1 activity, a modulator of PD-L2 activity, a modulator of CTLA-4 activity, a modulator of CD28 activity, a modulator of CD80 activity, a modulator of CD86 activity, a modulator of 4-1BB activity, a modulator of OX40 activity, a modulator of KIR activity, a modulator of Tim-3 activity, a modulator of LAG3 activity, a modulator of CD27 activity, a modulator of CD40 activity, a modulator of GITR activity, a modulator of TIGIT activity, a modulator of CD20 activity, a modulator of CD96 activity, a modulator of IDO1 activity, a cytokine, a chemokine, an interferon, an interleukin, a lymphokine, a member of the tumor necrosis factor (TNF) family, or an immunostimulatory oligonucleotide. In some embodiments, the immune checkpoint modulator, i.e. is an inhibitor or antagonist, or is an activator or agonist, for example, a CD28 modulator, a 4-1BB modulator, an OX40 modulator, a CD27 modulator, a CD80 modulator, a CD86 modulator, a CD40 modulator, or a GITR modulator, a Lag-3 modulator, a 41BB modulator, a LIGHT modulator, a CD40 modulator, a GITR modulator, a TGF-beta modulator, a TIM-3 modulator, a SIRP-alpha modulator, a TIGIT modulator, a VSIG8 modulator, a BTLA modulator, a SIGLEC7 modulator, a SIGLEC9 modulator, a ICOS modulator, a B7H3 modulator, a B7H4 modulator, a FAS modulator, and/or a BTNL2 modulator. In some embodiments, the immunotherapeutic agent is an immune checkpoint modulator as described above (e.g., an immune checkpoint modulator antibody, which can be in the form of a monoclonal antibody, a bispecific antibody comprising one or more immune checkpoint antigen binding moieties, a trispecific antibody, or an immune cell-engaging multivalent antibody/fusion protein/construct known in the art).

[000233] Combination treatments with immune checkpoint inhibitor immunotherapeutic agent may include antibodies that specifically target immune system checkpoints such as CTLA4, PD1 and PD-L1 are one of the most promising new avenues of immunotherapy for cancer and other diseases. Additional checkpoint targets, such as TIM-3, LAG-3, various B-7 ligands, CHK 1 and CHK2 kinases, BTLA, A2aR, and others, are also under investigation. Currently, three checkpoint inhibitors have received rapid approval from the U.S. Food and Drug Administration for cancer treatment, including ipilimumab (Yervoy®), a CTLA-4 inhibitor, and pembrolizumab (Keytruda®) and nivolumab (Opdivo®), both PD-1 inhibitors. In addition, several checkpoint inhibitor agents are in clinical trials.

[000234] Programmed Cell Death Protein 1, (PD-1 or CD279), a 55-kD type 1 transmembrane protein, is a member of the CD28 family of T cell co-stimulatory receptors that include immunoglobulin superfamily member CD28, CTLA-4, inducible co-stimulator (ICOS), and BTLA. PD-1 is highly expressed on activated T cells and B cells. PD-1 expression can also be detected on memory T-cell subsets with variable levels of expression. Two ligands specific for PD-1 have been identified: programmed death-ligand 1 (PD-L1, also known as B7-H1 or CD274) and PD-L2 (also known as B7-DC or CD273). PD-L1 and PD-L2 have been shown to down-regulate T cell activation upon binding to PD-1 in both mouse and human systems (Okazaki et al., *Int Immunol.*, 2007; 19: 813-824). The interaction of PD-1 with its ligands, PD-L1 and PD-L2, which are expressed on antigen-presenting cells (APCs) and dendritic cells (DCs), transmits negative regulatory stimuli to down-modulate the activated T cell immune response. Blockade of PD-1 suppresses this negative signal and amplifies T cell responses.

[000235] Numerous studies indicate that the cancer microenvironment manipulates the PD-L1-/PD-1 signaling pathway and that induction of PD-L1 expression is associated with inhibition of immune responses against cancer, thus permitting cancer progression and metastasis. The PD-L1/PD-1 signaling pathway is a primary mechanism of cancer immune evasion for several reasons. First, and most importantly, this pathway is involved in negative regulation of immune responses of activated T effector cells, found in the periphery. Second, PD-L1 is up-regulated in cancer microenvironments, while PD-1 is also up-regulated on activated tumor infiltrating T cells, thus possibly potentiating a vicious cycle of inhibition. Third, this pathway is intricately involved in both innate and adaptive immune regulation

through bi-directional signaling. These factors make the PD-1/PD-L1 complex a central point through which cancer can manipulate immune responses and promote its own progression.

[000236] CTLA-4 (also known as Cytotoxic T-lymphocyte-associated protein 4, CTLA4, CTLA-4, CD152, cluster of differentiation 152; ALPS5, CD, CELIAC3, GRD4, GSE, and IDDM12). CTLA-4 is a ~24.6-kDa single-pass type I membrane protein that plays an inhibitory role in T-cell function. CTLA-4 was originally identified by differential screening of a murine cytolytic T cell cDNA library, See Brunet et al., A new member of the immunoglobulin superfamily--CTLA-4, *Nature*. 1987 Jul 16-22;328(6127):267-70. CTLA- has been shown to interact with the b7 family ligands CD80 (also known as Cluster of differentiation 80, and B7-1); and CD86 (also known as Cluster of Differentiation 86 or B7-2). See Linsley et al., CTLA-4 is a second receptor for the B cell activation antigen B7, *J Exp Med*. 1991 Sep 1;174(3):561-9. Sequence comparison between the human CTLA-4 DNA encoding region, and that of CD28, reveals significant homology between both sequences, with the greatest similarity between juxtamembrane and cytoplasmic regions; accordingly, CTLA-4 is implicated in abrogating/reducing T-cell activity, and opposes the activity of CD28. CTLA-4 deficient mice have been shown to exhibit massive lymphoproliferation. Chambers et al., Lymphoproliferation in CTLA-4-deficient mice is mediated by costimulation-dependent activation of CD4+ T cells, *Immunity*. 1997 Dec;7(6):885-95. It has been reported that CTLA-4 blockade augments T-cell responses both in vitro and in vivo, enhances an induced autoimmune disease, and exacerbates antitumor immunity. (See Luhder, *J. Exp. Med.* 1998; 187:427-432; Walunas et al., *Immunity*. 1994; 1:405-413; Kearney, *J. Immunol.* 1995; 155:1032-1036); Leach, *Science* 1996; 271:1734-1736). CTLA-4 has also been reported as having alternative and/or additional impact on the initial character of the T-cell immune response (Chambers, *Curr. Opin. Immunol.* 1997; 9:396-404; Bluestone, *J. Immunol.* 1997; 158:1989-1993; Thompson, *Immunity* 1997; 7:445-450).

[000237] The first immune-checkpoint inhibitor to be tested in a clinical trial was ipilimumab (Yervoy, Bristol-Myers Squibb), an CTLA-4 mAb. CTLA-4 belongs to the immunoglobulin superfamily of receptors, which also includes PD-1, BTLA, TIM-3, and V-domain immunoglobulin suppressor of T cell activation (VISTA). Anti-CTLA-4 mAb is a powerful checkpoint inhibitor which removes "the break" from both naive and antigen-experienced cells. Therapy enhances the antitumor function of CD8+ T cells, increases the ratio of CD8+ T cells to Foxp3+ T regulatory cells, and inhibits the suppressive function of T

regulatory cells. The major drawback to anti-CTLA-4 mAb therapy is the generation of autoimmune toxicities due to on-target effects of an over-exuberant immune system which has lost the ability to turn itself down. It has been reported that up to 25% of patients treated with ipilimumab developed serious grade 3-4 adverse events/autoimmune-type side effects including dermatitis, enterocolitis, hepatitis, endocrinopathies (including hypophysitis, thyroiditis, and adrenalitis), arthritis, uveitis, nephritis, and aseptic meningitis. In contrast to the anti-CTLA-4 experience, anti-PD-1 therapy appears to be better-tolerated and induces a relatively lower rate of autoimmune-type side effects.

[000238] In some embodiments, the immunotherapeutic agent is an agent that inhibits the activity of PD-1. In some embodiments, the immunotherapeutic agent is an agent that inhibits the activity of PD-L1 and/or PD-L2. In some embodiments, the immunotherapeutic agent is an agent that inhibits the activity of CTLA-4. In some embodiments, the immunotherapeutic agent is an agent that inhibits the activity of CD80 and/or CD86. In some embodiments, the immunotherapeutic agent is an agent that inhibits the activity of TIGIT. In some embodiments, the immunotherapeutic agent is an agent that inhibits the activity of KIR. In some embodiments, the immunotherapeutic agent is an agent that enhances or stimulates the activity of activating immune checkpoint receptors.

[000239] In some of the embodiments of the methods described herein, the immunotherapeutic agent is a PD-1 antagonist, a PD-L1 antagonist, a PD-L2 antagonist, a CTLA-4 antagonist, a CD80 antagonist, a CD86 antagonist, a KIR antagonist, a Tim-3 antagonist, a LAG3 antagonist, a TIGIT antagonist, a CD20 antagonist, a CD96 antagonist, or an IDO1 antagonist.

[000240] In some embodiments, the PD-1 antagonist is an antibody that specifically binds PD-1. In some embodiments, the antibody that binds PD-1 is pembrolizumab (KEYTRUDA®, MK-3475; Merck), pidilizumab (CT-011; Curetech Ltd.), nivolumab (OPDIVO®, BMS-936558, MDX-1106; Bristol Myer Squibb), MEDI0680 (AMP-514; AstraZeneca/MedImmune), REGN2810 (Regeneron Pharmaceuticals), BGB-A317 (BeiGene Ltd.), PDR-001 (Novartis), or STI-A1110 (Sorrento Therapeutics). In some embodiments, the antibody that binds PD-1 is described in PCT Publication WO 2014/179664, for example, an antibody identified as APE2058, APE1922, APE1923, APE1924, APE 1950, or APE1963 (Anaptysbio), or an antibody containing the CDR regions of any of these antibodies. In other

embodiments, the PD-1 antagonist is a fusion protein that includes the extracellular domain of PD-L1 or PD-L2, for example, AMP-224 (AstraZeneca/MedImmune). In other embodiments, the PD-1 antagonist is a peptide inhibitor, for example, AUNP-12 (Aurigene).

[000241] In some embodiments, the PD-L1 antagonist is an antibody that specifically binds PD-L1. In some embodiments, the antibody that binds PD-L1 is atezolizumab (RG7446, MPDL3280A; Genentech), MEDI4736 (AstraZeneca/MedImmune), BMS-936559 (MDX-1105; Bristol Myers Squibb), avelumab (MSB0010718C; Merck KGaA), KD033 (Kadmon), the antibody portion of KD033, or STI-A1014 (Sorrento Therapeutics). In some embodiments, the antibody that binds PD-L1 is described in PCT Publication WO 2014/055897, for example, Ab-14, Ab-16, Ab-30, Ab-31, Ab-42, Ab-50, Ab-52, or Ab-55, or an antibody that contains the CDR regions of any of these antibodies, the disclosure of which is incorporated herein by reference in its entirety.

[000242] In some embodiments, the CTLA-4 antagonist is an antibody that specifically binds CTLA-4. In some embodiments, the antibody that binds CTLA-4 is ipilimumab (YERVOY®; Bristol Myer Squibb) or tremelimumab (CP-675,206; Pfizer). In some embodiments, the CTLA-4 antagonist is a CTLA-4 fusion protein or soluble CTLA-4 receptor, for example, KARR-102 (Kahr Medical Ltd.).

[000243] In some embodiments, the LAG3 antagonist is an antibody that specifically binds LAG3. In some embodiments, the antibody that binds LAG3 is IMP701 (Prima BioMed), IMP731 (Prima BioMed/GlaxoSmithKline), BMS-986016 (Bristol Myer Squibb), LAG525 (Novartis), and GSK2831781 (GlaxoSmithKline). In some embodiments, the LAG3 antagonist includes a soluble LAG3 receptor, for example, IMP321 (Prima BioMed).

[000244] In some embodiments, the KIR antagonist is an antibody that specifically binds KIR. In some embodiments, the antibody that binds KIR is lirilumab (Bristol Myer Squibb/Innate Pharma).

[000245] In some embodiments, the immunotherapeutic agent used in the combinations disclosed herein (e.g., in combination with a compound of Formula I') is an activator or agonist of a costimulatory molecule. In one embodiment, the agonist of the costimulatory molecule is chosen from an agonist (e.g., an agonistic antibody or antigen-binding fragment thereof, or a soluble fusion) of OX40, CD2, CD27, CD28, CD3, ICAM-1, LFA-1 (CD11a/CD18), ICOS

(CD278), 4-1BB (CD137), GITR, CD30, CD40, BAFFR, HVEM, CD7, LIGHT, NKG2C, SLAMF7, NKp80, CD160, B7-H3, or CD83 ligand.

[000246] In some embodiments, the OX40 agonist includes OX40 ligand, or an OX40-binding portion thereof. For example, the OX40 agonist may be MEDI6383 (AstraZeneca). In some embodiments, the OX40 agonist is an antibody that specifically binds OX40. In some embodiments, the antibody that binds OX40 is MEDI6469 (AstraZeneca/MedImmune), MEDI0562 (AstraZeneca/MedImmune), or MOXR0916 (RG7888; Genentech). In some embodiments, the OX40 agonist is a vector (e.g., an expression vector or virus, such as an adenovirus) capable of expressing OX40 ligand. In some embodiments the OX40-expressing vector is Delta-24-RGDOX (DNAtrix) or DNX2401 (DNAtrix).

[000247] In some embodiments, the 4-1BB (CD137) agonist is a binding molecule, such as an anticalin. In some embodiments, the anticalin is PRS-343 (Pieris AG). In some embodiments, the 4-1BB agonist is an antibody that specifically binds 4-1BB. In some embodiments, antibody that binds 4-1BB is PF-2566 (PF-05082566; Pfizer) or urelumab (BMS-663513; Bristol Myer Squibb).

[000248] In some embodiments, the CD27 agonist is an antibody that specifically binds CD27. In some embodiments, the antibody that binds CD27 is varlilumab (CDX-1127; Celldex).

[000249] In some embodiments, the GITR agonist comprises GITR ligand or a GITR-binding portion thereof. In some embodiments, the GITR agonist is an antibody that specifically binds GITR. In some embodiments, the antibody that binds GITR is TRX518 (GITR, Inc.), MK-4166 (Merck), or INBRX-110 (Five Prime Therapeutics/Inhibrx).

[000250] TIM-3 has been identified as another important inhibitory receptor expressed by exhausted CD8⁺ T cells. In mouse models of cancer, it has been shown that the most dysfunctional tumor-infiltrating CD8⁺ T cells actually co-express PD-1 and TIM-3.

[000251] LAG-3 is another recently identified inhibitory receptor that acts to limit effector T-cell function and augment the suppressive activity of T regulatory cells. It has recently been revealed that PD-1 and LAG-3 are extensively co-expressed by tumor-infiltrating T cells in mice, and that combined blockade of PD-1 and LAG-3 provokes potent synergistic antitumor immune responses in mouse models of cancer.

[000252] PD-1 pathway blockade can be combined with vaccines or other a compound of Formula I' antibodies for improved therapeutic efficacy (Hirano, F. et al, *Cancer Res.*, 65(3): 1089-1096 (2005); Li, B. et al, *Clin. Cancer Res.*, 15: 1507-1509 (2009); and Curran, M. A. et al, *Proc. Natl. Acad. Set*, 107(9):4275-4280 (2010)).

[000253] In some embodiments, immunotherapeutic agents useful in the compositions and methods described herein may include a monoclonal antibody, a bispecific antibody comprising one or more immune checkpoint antigen binding moieties, a trispecific antibody, or an immune cell-engaging multivalent antibody/fusion protein/construct known in the art that target specifically both PD-1 and ligand PD-L1.

[000254] PD-1 (also known as Programmed Death 1, CD279, PDCD1) is a cell surface receptor with a critical role in regulating the balance between stimulatory and inhibitory signals in the immune system and maintaining peripheral tolerance (Ishida, Y et al. 1992 *EMBO J.* 11 3887; Kier, Mary E et al. 2008 *Annu Rev Immunol* 26 677-704; Okazaki, Taku et al. 2007 *International Immunology* 19 813-824). PD-1 is an inhibitory member of the immunoglobulin super-family with homology to CD28. The structure of PD-1 is a monomeric type 1 transmembrane protein, consisting of one immunoglobulin variable-like extracellular domain and a cytoplasmic domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) and an immunoreceptor tyrosine-based switch motif (ITSM). Expression of PD-1 is inducible on T cells, B cells, natural killer (NK) cells and monocytes, for example upon lymphocyte activation via T cell receptor (TCR) or B cell receptor (BCR) signalling (Kier, Mary E et al. 2008 *Annu Rev Immunol* 26 677-704; Agata, Y et al 1996 *Int Immunol* 8 765-72). PD-1 is a receptor for the ligands CD80, CD86, PD-L1 (B7-H1, CD274) and PD-L2 (B7-DC, CD273), which are cell surface expressed members of the B7 family (Freeman, Gordon et al. 2000 *J Exp Med* 192 1027; Latchman, Y et al. 2001 *Nat Immunol* 2 261). Upon ligand engagement, PD-1 recruits phosphatases such as SHP-1 and SHP-2 to its intracellular tyrosine motifs which subsequently dephosphorylate effector molecules activated by TCR or BCR signalling (Chemnitz, J et al. 2004 *J Immunol* 173 945-954; Riley, James L 2009 *Immunological Reviews* 229 114-125) In this way, PD-1 transduces inhibitory signals into T and B cells only when it is engaged simultaneously with the TCR or BCR.

[000255] PD-1 has been demonstrated to down-regulate effector T cell responses via both cell-intrinsic and cell-extrinsic functional mechanisms. Inhibitory signaling through PD-1

induces a state of unresponsiveness in T cells, resulting in the cells being unable to clonally expand or produce optimal levels of effector cytokines. PD-1 may also induce apoptosis in T cells via its ability to inhibit survival signals from co-stimulation, which leads to reduced expression of key anti-apoptotic molecules such as Bcl-XL (Kier, Mary E et al. 2008 *Annu Rev Immunol* 26 677-704). In addition to these direct effects, recent publications have implicated PD-1 as being involved in the suppression of effector cells by promoting the induction and maintenance of regulatory T cells (TREG). For example, PD-L1 expressed on dendritic cells was shown to act in synergy with TGF- β to promote the induction of CD4⁺ FoxP3⁺TREG with enhanced suppressor function (Francisco, Loise M et al. 2009 *J Exp Med* 206 3015-3029).

[000256] TIM-3 (also known as T-cell immunoglobulin and mucin-domain containing-3, TIM-3, Hepatitis A virus cellular receptor 2, HAVCR2, HAVcr-2, KIM-3, TIMD-3, TIMD3, Tim-3, and CD366) is a ~33.4-kDa single-pass type I membrane protein involved in immune responses (Sanchez-Fueyo et al., Tim-3 inhibits T helper type 1-mediated auto- and alloimmune responses and promotes immunological tolerance, *Nat. Immunol.* 4:1093-1101(2003)).

[000257] TIM-3 is selectively expressed on Th1-cells, and phagocytic cells (e.g., macrophages and dendritic cells). The use of siRNA or a blocking antibody to reduce the expression of human resulted in increased secretion of interferon γ (IFN- γ) from CD4 positive T-cells, implicating the inhibitory role of TIM-3 in human T cells. Analysis of clinical samples from autoimmune disease patients showed no expression of TIM-3 in CD4 positive cells. In particular, expression level of TIM-3 is lower and secretion of IFN- γ is higher in T cell clones derived from the cerebrospinal fluid of patients with multiple sclerosis than those in clones derived from normal healthy persons (Koguchi K et al., *J Exp Med.* 203:1413-8. (2006)).

[000258] TIM-3 is the receptor for the ligands Galectin-9, which is a member of galectin family, molecules ubiquitously expressed on a variety of cell types and which binds β -galactoside; Phosphatidyl serine (PtdSer) (DeKryff et al., T cell/transmembrane, Ig, and mucin-3 allelic variants differentially recognize phosphatidylserine and mediate phagocytosis of apoptotic cells, *J Immunol.* 2010 Feb 15;184(4):1918-30); High Mobility Group Protein 1 (also known as HMGB1, HMG1, HMG3, SBP-1, HMG-1, and high mobility group box 1) Chiba et al., Tumor-infiltrating DCs suppress nucleic acid-mediated innate immune responses through interactions between the receptor TIM-3 and the alarmin HMGB1, *Nat Immunol.* 2012 Sep;13(9):832-42); and Carcinoembryonic Antigen Related Cell Adhesion Molecule 1 (also

known as CEACAM1, BGP, BGP1, BGPI, carcinoembryonic antigen related cell adhesion molecule 1) (Huang et al., CEACAM1 regulates TIM-3-mediated tolerance and exhaustion, *Nature*. 2015 Jan 15;517(7534):386-90).

[000259] BTLA (also known as B- and T-lymphocyte attenuator, BTLA1, CD272, and B and T lymphocyte associated) is a ~27.3-kDa single-pass type I membrane protein involved in lymphocyte inhibition during immune response. BTLA is constitutively expressed in both B and T cells. BTLA interacts with HVEM (herpes virus-entry mediator), a member of the tumor-necrosis factor receptor (TNFR) family (Gonzalez et al., *Proc. Natl. Acad. Sci. USA*, 2005, 102: 1116-21). The interaction of BTLA, which belongs to the CD28 family of the immunoglobulin superfamily, and HVEM, a costimulatory tumor-necrosis factor (TNF) receptor (TNFR), is unique in that it defines a cross talk between these two families of receptors. BTLA contains a membrane proximal immunoreceptor tyrosine-based inhibitory motif (ITIM) and membrane distal immunoreceptor tyrosine-based switch motif (ITSM). Disruption of either the ITIM or ITSM abrogated the ability of BTLA to recruit either SHP1 or SHP2, suggesting that BTLA recruits SHP1 and SHP2 in a manner distinct from PD-1 and both tyrosine motifs are required to block T cell activation. The BTLA cytoplasmic tail also contains a third conserved tyrosine-containing motif within the cytoplasmic domain, similar in sequence to a Grb-2 recruitment site (YXN). Also, a phosphorylated peptide containing this BTLA N-terminal tyrosine motif can interact with GRB2 and the p85 subunit of PI3K in vitro, although the functional effects of this interaction remain unexplored in vivo (Gavrieli et al., *Biochem. Biophys Res Commun*, 2003, 312, 1236-43). BTLA is the receptor for the ligands PTPN6/SHP-1; PTPN11/SHP-2; TNFRSF14/HVEM; and B7H4.

[000260] VISTA (also known as V-domain Ig suppressor of T cell activation VSIR, B7-H5, B7H5, GI24, PP2135, SISP1, DD1alpha, VISTA, C10orf54, chromosome 10 open reading frame 54, PD-1H, and V-set immunoregulatory receptor) is a ~33.9-kDa single-pass type I membrane protein involved in T-cell inhibitory response, embryonic stem cells differentiation via BMP4 signaling inhibition, and MMP14-mediated MMP2 activation (Yoon et al., *Control of signaling-mediated clearance of apoptotic cells by the tumor suppressor p53*, *Science*. 2015 Jul 31; 349(6247): 1261669). VISTA interacts with the ligand VSIG-3 (Wang et al., VSIG-3 as a ligand of VISTA inhibits human T-cell function, *Immunology*. 2019 Jan;156(1):74-85)

[000261] LAG-3 (also known as Lymphocyte-activation gene 3, LAG3, CD223, and lymphocyte activating 3) is a ~57.4-kDa single-pass type I membrane protein involved in lymphocyte activation that also binds to HLA class-II antigens. LAG-3 is a member of the immunoglobulin supergene family, and is expressed on activated T cells (Huard et al., 1994, *Immunogenetics* 39:213), NK cells (Triebel et al., 1990, *J. Exp. Med.* 171:1393-1405), regulatory T cells (Huang et al., 2004, *Immunity* 21:503-513; Camisaschi et al., 2010, *J Immunol.* 184:6545-6551; Gagliani et al., 2013, *Nat Med* 19:739-746), and plasmacytoid dendritic cells (DCs) (Workman et al., 2009, *J Immunol* 182:1885-1891). LAG-3 is a membrane protein encoded by a gene located on chromosome 12, and is structurally and genetically related to CD4. Similar to CD4, LAG-3 can interact with MHC class II molecules on the cell surface (Baixeras et al., 1992, *J. Exp. Med.* 176:327-337; Huard et al., 1996, *Eur. J. Immunol.* 26:1180-1186). It has been suggested that the direct binding of LAG-3 to MHC class II plays a role in down-regulating antigen-dependent stimulation of CD4⁺ T lymphocytes (Huard et al., 1994, *Eur. J. Immunol.* 24:3216-3221) and LAG-3 blockade has also been shown to reinvigorate CD8⁺ lymphocytes in both tumor or self-antigen (Gross et al., 2007, *J Clin Invest.* 117:3383-3392) and viral models (Blackburn et al., 2009, *Nat. Immunol.* 10:29-37). Further, the intra-cytoplasmic region of LAG-3 can interact with LAP (LAG-3-associated protein), which is a signal transduction molecule involved in the downregulation of the CD3/TCR activation pathway (Iouzalén et al., 2001, *Eur. J. Immunol.* 31:2885-2891). Moreover, CD4⁺CD25⁺ regulatory T cells (Treg) have been shown to express LAG-3 upon activation, which contributes to the suppressor activity of Treg cells (Huang, C. et al., 2004, *Immunity* 21:503-513). LAG-3 can also negatively regulate T cell homeostasis by Treg cells in both T cell-dependent and independent mechanisms (Workman, C. J. and Vignali, D. A., 2005, *J. Immunol.* 174:688-695).

[000262] LAG-3 has been shown to interact with MHC class II molecules (Huard et al., CD4/major histocompatibility complex class II interaction analyzed with CD4- and lymphocyte activation gene-3 (LAG-3)-Ig fusion proteins, *Eur J Immunol.* 1995 Sep;25(9):2718-21).

[000263] Additionally, several kinases are known to be checkpoint inhibitors. For example, CHEK-1, CHEK-2, and ATR.

[000264] CHEK-1 (also known as CHK 1 kinase, CHK1, and checkpoint kinase 1) is a ~54.4-kDa serine/threonine-protein kinase that is involved with checkpoint-mediated cell cycle

arrest, and the activation of DNA repair in response to the DNA damage and/or unreplicated DNA.

[000265] CHEK-2 (also known as CHK2 kinase, CDS1, CHK2, HuCds1, LFS2, PP1425, RAD53, hCds1, and checkpoint kinase 2) is a ~60.9-kDa. serine/threonine-protein kinase involved in checkpoint-mediated cell cycle arrest, DNA-repair activation, and double-strand break-mediated apoptosis.

[000266] A2aR (also known as adenosine A2A receptor, ADORA2A, adenosine A2a receptor, A2aR, ADORA2, and RDC8) is a ~44.7-kDa multi-pass membrane receptor for adenosine and other ligands.

[000267] In various embodiments, the immunotherapeutic agent can comprise an antibody or an antigen binding fragment thereof. Within this definition, immune checkpoint inhibitors include bispecific antibodies and immune cell-engaging multivalent antibody/fusion protein/constructs known in the art. In some embodiments, immunotherapeutic agents which comprise bispecific antibodies may include bispecific antibodies that are bivalent and bind either the same epitope of the immune checkpoint molecule, two different epitopes of the same immune checkpoint molecule or different epitopes of two different immune checkpoints.

[000268] Persons of ordinary skill in the art can implement several bispecific antibody formats known in the field to target one or more of CTLA4, PD1, PD-L1, TIM-3, LAG-3, various B-7 ligands, B7H3, B7H4, CHK 1 and CHK2 kinases, BTLA, A2aR, OX40, 41BB, LIGHT, CD40, GITR, TGF-beta, SIRP-alpha, TIGIT, VSIG8, SIGLEC7, SIGLEC9, ICOS, FAS, BTNL2 and other for use in the combination described herein.

[000269] In various embodiments, the immunotherapeutic agent can include an immune cell-engaging multivalent antibody/fusion protein/construct.

[000270] In an embodiment of the disclosure, the checkpoint inhibitor, in combination with a compound of Formula I', is used to reduce or inhibit metastasis of a primary tumor or cancer to other sites, or the formation or establishment of metastatic tumors or cancers at other sites distal from the primary tumor or cancer thereby inhibiting or reducing tumor or cancer relapse or tumor or cancer progression.

[000271] In a further embodiment of the disclosure, there is provided a combination therapy for treating cancer, comprising a compound of Formula I' and blockade of checkpoint inhibitors

with the potential to elicit potent and durable immune responses with enhanced therapeutic benefit and more manageable toxicity.

[000272] In a further embodiment of the disclosure, there is provided a combination therapy for treating cancer, comprising a compound of Formula I' and an immune checkpoint inhibitor. In an embodiment of the disclosure is provided a method for treating cancer and/or preventing the establishment of metastases by employing a checkpoint inhibitor which act synergistically with a compound of Formula I'.

[000273] In further embodiments, methods of the disclosure include, one or more of the following: 1) reducing or inhibiting growth, proliferation, mobility or invasiveness of tumor or cancer cells that potentially or do develop metastases, 2) reducing or inhibiting formation or establishment of metastases arising from a primary tumor or cancer to one or more other sites, locations or regions distinct from the primary tumor or cancer; 3) reducing or inhibiting growth or proliferation of a metastasis at one or more other sites, locations or regions distinct from the primary tumor or cancer after a metastasis has formed or has been established, 4) reducing or inhibiting formation or establishment of additional metastasis after the metastasis has been formed or established, 5) prolonged overall survival, 6) prolonged progression free survival, or 7) disease stabilization.

[000274] In an embodiment of the disclosure, administration of the immunotherapeutic agent, in combination therapy with a compound of Formula I', provides a detectable or measurable improvement in a condition of a given subject, such as alleviating or ameliorating one or more adverse (physical) symptoms or consequences associated with the presence of a cell proliferative or cellular hyperproliferative disorder, neoplasia, tumor or cancer, or metastasis, i.e., a therapeutic benefit or a beneficial effect.

[000275] A therapeutic benefit or beneficial effect is any objective or subjective, transient, temporary, or long-term improvement in the condition or pathology, or a reduction in onset, severity, duration or frequency of adverse symptom associated with or caused by cell proliferation or a cellular hyperproliferative disorder such as a neoplasia, tumor or cancer, or metastasis. It may lead to improved survival. A satisfactory clinical endpoint of a treatment method in accordance with the disclosure is achieved, for example, when there is an incremental or a partial reduction in severity, duration or frequency of one or more associated pathologies, adverse symptoms or complications, or inhibition or reversal of one or more of the

physiological, biochemical or cellular manifestations or characteristics of cell proliferation or a cellular hyperproliferative disorder such as a neoplasia, tumor or cancer, or metastasis. A therapeutic benefit or improvement therefore may be, but is not limited to destruction of target proliferating cells (e.g., neoplasia, tumor or cancer, or metastasis) or ablation of one or more, most or all pathologies, adverse symptoms or complications associated with or caused by cell proliferation or the cellular hyperproliferative disorder such as a neoplasia, tumor or cancer, or metastasis. However, a therapeutic benefit or improvement need not be a cure or complete destruction of all target proliferating cells (e.g., neoplasia, tumor or cancer, or metastasis) or ablation of all pathologies, adverse symptoms or complications associated with or caused by cell proliferation or the cellular hyperproliferative disorder such as a neoplasia, tumor or cancer, or metastasis. For example, partial destruction of a tumor or cancer cell mass, or a stabilization of the tumor or cancer mass, size or cell numbers by inhibiting progression or worsening of the tumor or cancer, can reduce mortality and prolong lifespan even if only for a few days, weeks or months, even though a portion or the bulk of the tumor or cancer mass, size or cells remain.

[000276] Specific non-limiting examples of therapeutic benefit include a reduction in neoplasia, tumor or cancer, or metastasis volume (size or cell mass) or numbers of cells, inhibiting or preventing an increase in neoplasia, tumor or cancer volume (e.g., stabilizing), slowing or inhibiting neoplasia, tumor or cancer progression, worsening or metastasis, or inhibiting neoplasia, tumor or cancer proliferation, growth or metastasis.

[000277] In an embodiment of the disclosure, administration of the immunotherapeutic agent, in combination therapy with a compound of Formula I', provides a detectable or measurable improvement or overall response according to the irRC (as derived from time-point response assessments and based on tumor burden), including one or more of the following: (i) irCR--complete disappearance of all lesions, whether measurable or not, and no new lesions (confirmation by a repeat, consecutive assessment no less than 4 weeks from the date first documented), (ii) irPR--decrease in tumor burden $\geq 50\%$ relative to baseline (confirmed by a consecutive assessment at least 4 weeks after first documentation).

[000278] Optionally, any method described herein may not take effect immediately. For example, treatment may be followed by an increase in the neoplasia, tumor or cancer cell numbers or mass, but over time eventual stabilization or reduction in tumor cell mass, size or numbers of cells in a given subject may subsequently occur.

[000279] Additional adverse symptoms and complications associated with neoplasia, tumor, cancer and metastasis that can be inhibited, reduced, decreased, delayed or prevented include, for example, nausea, lack of appetite, lethargy, pain and discomfort. Thus, a partial or complete decrease or reduction in the severity, duration or frequency of adverse symptom or complication associated with or caused by a cellular hyperproliferative disorder, an improvement in the subjects quality of life and/or well-being, such as increased energy, appetite, psychological well-being, are all particular non-limiting examples of therapeutic benefit.

[000280] A therapeutic benefit or improvement therefore can also include a subjective improvement in the quality of life of a treated subject. In additional embodiment, a method prolongs or extends lifespan (survival) of the subject. In a further embodiment, a method improves the quality of life of the subject.

[000281] In one embodiment, administration of the immunotherapeutic agent, in combination therapy with a compound of Formula I', results in a clinically relevant improvement in one or more markers of disease status and progression selected from one or more of the following: (i): overall survival, (ii): progression-free survival, (iii): overall response rate, (iv): reduction in metastatic disease, (v): circulating levels of tumor antigens such as carbohydrate antigen 19.9 (CA19.9) and carcinoembryonic antigen (CEA) or others depending on tumor, (vii) nutritional status (weight, appetite, serum albumin), (viii): pain control or analgesic use, (ix): CRP/albumin ratio.

[000282] Treatment with a compound of Formula I' in combination with an immunotherapeutic agent gives rise to more complex immunity including not only the development of innate immunity and type-1 immunity, but also immunoregulation which more efficiently restores appropriate immune functions.

[000283] In various exemplary methods, a checkpoint inhibitor antibody (monoclonal or polyclonal, bispecific, trispecific, or an immune cell-engaging multivalent antibody/fusion protein/construct) directed to a checkpoint molecule of interest (e.g., PD-1) may be sequenced and the polynucleotide sequence may then be cloned into a vector for expression or propagation. The sequence encoding the antibody or antigen-binding fragment thereof of interest may be maintained in vector in a host cell and the host cell can then be expanded and frozen for future use. Production of recombinant monoclonal antibodies in cell culture can be

carried out through cloning of antibody genes from B cells by means known in the art. See, e.g. Tiller et al., 2008, J. Immunol. Methods 329, 112; U.S. Pat. No. 7,314,622.

[000284] In some embodiments, methods for producing the recombinant antibodies can include the steps of culturing a host cell containing isolated nucleic acid(s) encoding the antibodies of the present disclosure. Methods for culturing a host cell containing isolated nucleic acid(s) encoding the antibodies of the present disclosure can be done in a variety of ways, depending on the nature of the antibody. In some embodiments, in the case where the antibodies of the disclosure are full length traditional antibodies, for example, a heavy chain variable region and a light chain variable region under conditions such that an antibody is produced and can be isolated.

[000285] In general, nucleic acids are provided that encode the antibodies or antigen-binding fragments thereof of the present disclosure. Such polynucleotides encode for both the variable and constant regions of each of the heavy and light chains, although other combinations are also contemplated by the present disclosure. The present disclosure also contemplates oligonucleotide fragments derived from the disclosed polynucleotides and nucleic acid sequences complementary to these polynucleotides.

[000286] The polynucleotides can be in the form of RNA, DNA, cDNA, genomic DNA, nucleic acid analogs, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded, may be the coding (sense) strand or non-coding (anti-sense) strand. The coding sequence that encodes the polypeptide may be identical to the coding sequence or may be a different coding sequence, which sequence, as a result of the redundancy or degeneracy of the genetic code, encodes the same polypeptides.

[000287] In some embodiments, nucleic acid(s) encoding the antibodies of the present disclosure are incorporated into expression vectors, which can be extrachromosomal or designed to integrate into the genome of the host cell into which it is introduced. Expression vectors can contain any number of appropriate regulatory sequences (including, but not limited to, transcriptional and translational control sequences, promoters, ribosomal binding sites, enhancers, origins of replication, and the like) or other components (selection genes, and the like), all of which are operably linked as is well known in the art. In some cases two nucleic acids are used and each put into a different expression vector (e.g. heavy chain in a first expression vector, light chain in a second expression vector), or alternatively they can be put in

the same expression vector. It will be appreciated by those skilled in the art that the design of the expression vector(s), including the selection of regulatory sequences may depend on such factors as the choice of the host cell, the level of expression of protein desired, and the like.

[000288] In general, the nucleic acids and/or expression can be introduced into a suitable host cell to create a recombinant host cell using any method appropriate to the host cell selected (e.g., transformation, transfection, electroporation, infection), such that the nucleic acid molecule(s) are operably linked to one or more expression control elements (e.g., in a vector, in a construct created by processes in the cell, integrated into the host cell genome). The resulting recombinant host cell can be maintained under conditions suitable for expression (e.g. in the presence of an inducer, in a suitable non-human animal, in suitable culture media supplemented with appropriate salts, growth factors, antibiotics, nutritional supplements, and the like), whereby the encoded polypeptide(s) are produced. In some cases, the heavy chains are produced in one cell and the light chain in another.

[000289] Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC), Manassas, VA USA. including but not limited to Chinese hamster ovary (CHO) cells, HEK 293 cells, NSO cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), and a number of other cell lines. Non-mammalian cells including but not limited to bacterial, yeast, insect, and plants can also be used to express recombinant antibodies. In some embodiments, the antibodies can be produced in transgenic animals such as cows or chickens.

[000290] Exemplary and illustrative recombinant methods for antibody molecular biology, expression, purification, and screening are described, for example, in Antibody Engineering, edited by Kontermann & Dubel, Springer, Heidelberg, 2001 and 2010 Hayhurst & Georgiou, 2001, Curr. Opin. Chem. Biol. 5:683-689; Maynard & Georgiou, 2000, Annu. Rev. Biomed. Eng. 2:339-76; and Morrison, S. (1985) Science 229:1202, the disclosures of which are incorporated herein by reference in their entireties.

[000291] In various embodiments, the polynucleotide sequence encoding the selected variable heavy and light chains may be used for genetic manipulation to humanize the antibody or to improve the affinity, or other characteristics of the antibody. Antibodies may also be customized for use, for example, in dogs, cats, primate, equines and bovines.

[000292] In some embodiments, fully human antibodies may be obtained by using commercially available mice that have been engineered to express specific human immunoglobulin proteins. Transgenic animals that are designed to produce a more desirable (e.g., fully human antibodies) or more robust immune response may also be used for generation of humanized or human antibodies. Examples of such technology are Xenomouse™ from Abgenix, Inc. (Fremont, Calif.) and HuMAb-Mouse® and TC Mouse™ from Medarex, Inc. (Princeton, N.J.).

[000293] Immune checkpoint modulator antibodies of the present disclosure can be made recombinantly by first isolating the antibodies and antibody producing cells from host animals, obtaining the gene sequence, and using the gene sequence to express the antibody recombinantly in host cells (e.g., CHO cells). Another method which may be employed is to express the antibody sequence in plants (e.g., tobacco) or in yeast cells (e.g. *Pichia pastoris* or *Saccharomyces cerevisiae*). Methods for expressing antibodies recombinantly in plants or yeast have been disclosed. See, for example, Peeters, et al. Vaccine 19:2756, 2001; Lonberg, N. and D. Huszar Int. Rev. Immunol 13:65, 1995; and Horwitz, A. H. et al., Proc. Natl. Acad. Sci. 85:8678-8682; the disclosures of which are hereby incorporated by reference in their entireties. Methods for making derivatives of antibodies, e.g., domain, single chain, and the like are known in the art.

[000294] Immunoassays and flow cytometry sorting techniques such as fluorescence activated cell sorting (FACS) can also be employed to isolate antibodies that are specific for checkpoint molecules.

[000295] In some embodiments, a polynucleotide comprises a sequence encoding the heavy chain and/or the light chain variable regions of the checkpoint inhibitor antibody or antigen-binding fragment thereof of the present disclosure. The sequence encoding the antibody or antigen-binding fragment thereof of interest may be maintained in a vector in a host cell and the host cell can then be expanded and frozen for future use. Vectors (including expression vectors) and host cells are further described herein.

[000296] The disclosure includes affinity matured checkpoint modulator antibodies. For example, affinity matured antibodies can be produced by procedures known in the art (Marks et al., 1992, Bio/Technology, 10:779-783; Barbas et al., 1994, Proc Nat. Acad. Sci. USA 91:3809-3813. One way of characterizing a CDR of an antibody and/or altering (such as improving) the

binding affinity of a polypeptide, such as an antibody, termed "library scanning mutagenesis". An exemplary method for providing affinity matures antibodies and antigen-binding fragments can include replacing one or more amino acid positions in the CDR with two or more (such as 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20) amino acids using art recognized methods. a library of clones are generated, each with a complexity of two or more members (if two or more amino acids are substituted at every position). Generally, the library also includes a clone comprising the native (unsubstituted) amino acid. A small number of clones, e.g., about 20-80 clones (depending on the complexity of the library), from each library are screened for binding affinity to the target polypeptide (or other binding target), and candidates with increased, the same, decreased, or no binding are identified. Methods for determining binding affinity are well-known in the art. Binding affinity may be determined using, for example, Biacore™ surface plasmon resonance analysis, which detects differences in binding affinity of about 2-fold or greater, Kinexa® Biosensor, scintillation proximity assays, ELISA, ORIGEN® immunoassay, fluorescence quenching, fluorescence transfer, and/or yeast display. Binding affinity may also be screened using a suitable bioassay. Biacore™ is particularly useful when the starting antibody already binds with a relatively high affinity, for example a KD of about 10 nM or lower. The library of clones can then be recombinantly introduced into a selection construct using any method known in the art for selection, including phage display, yeast display, and ribosome display.

[000297] The antibodies may also be modified, e.g., in the variable domains of the heavy and/or light chains, e.g., to alter a binding property of the antibody. Changes in the variable region can alter binding affinity and/or specificity. In some embodiments, no more than one to five conservative amino acid substitutions are made within a CDR domain. In other embodiments, no more than one to three conservative amino acid substitutions are made within a CDR domain. For example, a mutation may be made in one or more of the CDR regions to increase or decrease the KD of the antibody directed to a checkpoint molecule, to increase or decrease kon or to alter the binding specificity of the antibody. Techniques in site-directed mutagenesis are well-known in the art. See, e.g., Sambrook et al. and Ausubel et al.

[000298] Pharmaceutical compositions containing a compound of Formula I' according to the present disclosure will comprise an effective amount of a compound of Formula I', an immunotherapeutic agent, and/or both, typically dispersed in a pharmaceutically acceptable

carrier. The phrases "pharmaceutically or pharmacologically acceptable" refers to molecular entities and compositions that do not produce adverse, allergic or other untoward reaction when administered to animal, such as, for example, a human, as appropriate. The preparation of an pharmaceutical composition that contains a compound of Formula I' will be known to those of skill in the art in light of the present disclosure, as exemplified by Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, Moreover, for animal (e.g., human) administration, it will be understood that preparations should meet sterility, pyrogenicity, general safety and purity standards. A specific example of a pharmacologically acceptable carrier for a combination compositions, containing a compound of Formula I' in admixture with an immunotherapeutic agent as described herein is borate buffer or sterile saline solution (0.9% NaCl).

[000299] Formulations of the an immunotherapeutic agent, for example an immune checkpoint modulator antibody used in accordance with the present disclosure can be prepared for storage by mixing an antibody having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers as amply described and illustrated in Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. [1980], in the form of lyophilized formulations or aqueous solutions and/or suspensions. Acceptable carriers, excipients, buffers or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include suitable aqueous and/or non-aqueous excipients that may be employed in the pharmaceutical compositions of the disclosure, for example, water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants, buffers such as phosphate, citrate, and other organic acids. Antioxidants may be included, for example, (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like; preservatives (such as octade-cyldimethylbenzyl ammonium

chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues). Other exemplary pharmaceutically acceptable excipients may include polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and/or non-ionic surfactants such as TWEENTM, PLURONIC^{STM} or polyethylene glycol (PEG).

[000300] In one illustrative embodiment, the pharmaceutical compositions can optionally contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents and toxicity adjusting agents, for example, sodium acetate, sodium chloride, potassium chloride, calcium chloride and sodium lactate. In some embodiments, the checkpoint inhibitor antibodies or antigen-binding fragments thereof of the present disclosure are formulated for and can be lyophilized for storage and reconstituted in a suitable excipient prior to use according to art-known lyophilization and reconstitution techniques. In one exemplary pharmaceutical composition containing one or more checkpoint inhibitor antibodies or antigen-binding fragment thereof, the composition is formulated as a sterile, preservative-free solution of one or more checkpoint inhibitor antibodies or antigen-binding fragment thereof for intravenous or subcutaneous administration. The formulation can be supplied as either a single-use, prefilled pen, as a single-use, for example containing about 1 mL prefilled glass syringe, or as a single-use institutional use vial. Preferably, the pharmaceutical composition containing the checkpoint inhibitor antibody or antigen-binding fragment thereof is clear and colorless, with a pH of about 6.9-5.0, preferably a pH of 6.5-5.0, and even more preferably a pH ranging from about 6.0 to about 5.0. In various embodiments, the formulations comprising the pharmaceutical compositions can contain from about 500 mg to about 10 mg, or from about 400 mg to about 20 mg, or from about 300 mg to about 30 mg or from about 200 mg to about 50 mg of the checkpoint inhibitor antibody or antigen-binding fragment thereof per mL of solution when reconstituted and administered to the

subject. Exemplary injection or infusion excipients can include mannitol, citric acid monohydrate, dibasic sodium phosphate dihydrate, monobasic sodium phosphate dihydrate, polysorbate 80, sodium chloride, sodium citrate and water for parenteral administration, for example, intravenously, intramuscularly, intraperitoneally, or subcutaneous administration.

[000301] In another exemplary embodiment, one or more immunotherapeutic agents, or an antigen-binding fragment thereof is formulated for intravenous or subcutaneous administration as a sterile aqueous solution containing 1-75 mg/mL, or more preferably, about 5-60 mg/mL, or yet more preferably, about 10-50 mg/mL, or even more preferably, about 10-40 mg/mL of antibody, with sodium acetate, polysorbate 80, and sodium chloride at a pH ranging from about 5 to 6. Preferably, the intravenous or subcutaneous formulation is a sterile aqueous solution containing 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 mg/mL of the immunotherapeutic agent, for example, an immune checkpoint inhibitor antibody or an antigen-binding fragment thereof, with 20 mM sodium acetate, 0.2 mg/mL polysorbate 80, and 140 mM sodium chloride at pH 5.5. Further, a solution comprising a checkpoint inhibitor antibody or an antigen-binding fragment thereof, can comprise, among many other compounds, histidine, mannitol, sucrose, trehalose, glycine, poly(ethylene)glycol, EDTA, methionine, and any combination thereof, and many other compounds known in the relevant art.

[000302] In one embodiment, a pharmaceutical composition of the present disclosure comprises the following components: 5-500 mg of an immunotherapeutic agent or antigen-binding fragment thereof of the present disclosure, 10 mM histidine, 5% sucrose, and 0.01% polysorbate 80 at pH 5.8, with or without a compound of Formula I'. This composition may be provided as a lyophilized powder. When the powder is reconstituted at full volume, the composition retains the same formulation. Alternatively, the powder may be reconstituted at half volume, in which case the composition comprises 10-500 mg of an immunotherapeutic agent or antigen-binding fragment thereof of the present disclosure, 20 mM histidine, 10% sucrose, and 0.02% polysorbate 80 at pH 5.8.

[000303] In one embodiment, part of the dose is administered by an intravenous bolus and the rest by infusion of the immunotherapeutic agent formulation. For example, from about 0.001 to about 200 mg/kg, for example, from about 0.001 mg/kg to about 100 mg/kg, or from about 0.001 mg/kg to about 50 mg/kg, or from about 0.001 mg/kg to about 10 mg/kg intravenous injection of the immunotherapeutic agent, or antigen-binding fragment thereof, may be given as

a bolus, and the rest of the antibody dose may be administered by intravenous injection. A predetermined dose of the immunotherapeutic agent, or antigen-binding fragment thereof, may be administered, for example, over a period of an hour to two hours to five hours.

[000304] In a further embodiment, part of the dose is administered by a subcutaneous injection and/or infusion in the form of a bolus and the rest by infusion of the immunotherapeutic agent formulation. In some exemplary doses, the immunotherapeutic agent formulation can be administered subcutaneously in a dose ranging from about 0.001 to about 200 mg/kg, for example, from about 0.001 mg/kg to about 100 mg/kg, or from about 0.001 mg/kg to about 50 mg/kg, or from about 0.001 mg/kg to about 10 mg/kg intravenous injection of the immunotherapeutic agent, or antigen-binding fragment thereof. In some embodiments the dose may be given as a bolus, and the rest of the immunotherapeutic agent dose may be administered by subcutaneous or intravenous injection. A predetermined dose of the immunotherapeutic agent, or antigen-binding fragment thereof, may be administered, for example, over a period of an hour to two hours to five hours.

[000305] The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to provide one or more immunotherapeutic agents with other specificities. Alternatively, or in addition, the composition may comprise an anti-inflammatory agent, a chemotherapeutic agent, a cytotoxic agent, a cytokine, a growth inhibitory agent and/or a small molecule antagonist. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[000306] The formulations to be used for in vivo administration should be sterile, or nearly so. This is readily accomplished by filtration through sterile filtration membranes.

[000307] In various embodiments, illustrative formulations of the pharmaceutical compositions described herein can be prepared using methods widely known in the field of pharmaceutical formulations. In general, such preparatory methods can include the step of bringing the active ingredient into association with a carrier or one or more other accessory ingredients, and then, if desirable, packaging the product into a desired single-or multi-dose unit.

[000308] In some embodiments, the composition comprising a compound of Formula I' can be also delivered in a vesicle, and the immunotherapeutic agent can be delivered in the same

liposome formulation, or in a separate formulation that is compatible with the liposomal formulation containing the compound of Formula I', In some illustrative examples, a liposome containing one or more liposomal surface moieties for example, polyethylene glycol, antibodies and antibody fragments thereof that target a desired tumor surface antigen, receptor, growth factor, glycoprotein, glycolipid or neoantigen, which are selectively transported into specific cells or organs, thus enhance targeted drug delivery.

[000309] In another embodiment, a compound of Formula I' can be delivered in a vesicle, in particular a liposome (see Langer, Science 249:1527-1533 (1990); Treat et al., in LIPOSOMES IN THE THERAPY OF INFECTIOUS DISEASE AND CANCER, Lopez-Berestein and Fidler (eds.), Liss, N.Y., pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*).

[000310] In yet another embodiment, a compound of Formula I', or the composition containing the combination, or a composition containing the immunotherapeutic agent, can be delivered in a controlled release system. In one embodiment, a pump can be used (see Langer, *supra*; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989)). In another embodiment, controlled release of the compound of Formula I' can comprise polymeric materials to provide sustained, intermediate, pulsatile, or alternate release (see MEDICAL APPLICATIONS OF CONTROLLED RELEASE, Langer and Wise (eds.), CRC Pres., Boca Raton, Fla. (1974); CONTROLLED DRUG BIOAVAILABILITY, DRUG PRODUCT DESIGN AND PERFORMANCE, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J. Macromol. Sci. Rev. Macromol. Chem. 23:61 (1983); see also Levy et al., Science 228:190 (1985); During et al., Ann. Neurol. 25:351(1989); Howard et al., J. Neurosurg. 71:105 (1989)). Other controlled-release systems discussed in the review by Langer (Science 249:1527-1533 (1990)) can be used.

[000311] The optimum concentration of the active ingredient(s) in the chosen medium can be determined empirically, according to procedures well known to the skilled artisan, and will depend on the ultimate pharmaceutical formulation desired and the use to be employed.

[000312] The present disclosure also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the disclosure, which at minimum will include a compound of Formula I' and one or more checkpoint inhibitor antibodies or antigen-binding fragment thereof as described

herein. In other embodiments, the kit may contain one or more further containers providing a pharmaceutically acceptable excipient, for example a diluent. In one embodiment a kit may comprise at least one container, wherein the container can include a compound of Formula I', a checkpoint inhibitor antibody or an antigen-binding fragment thereof of the present disclosure,. The kit may also include a set of instructions for preparing and administering the final pharmaceutical composition to the subject in need thereof, for the treatment of a checkpoint molecule-mediated disease or disorder.

[000313] Some embodiments of the present disclosure, the immunotherapeutic agent is a population of immune cells, which can be administered in combination with a compound of Formula I' to treat a subject with cancer. In some embodiments, the immunotherapeutic agent is a population of immune cells, such as leukocytes (nucleated white blood cells), comprising (e.g., expressing) a receptor that binds to an antigen of interest. A leukocyte of the present disclosure may be, for example, a neutrophil, eosinophil, basophil, lymphocyte or a monocyte. In some embodiments, a leukocyte is a lymphocyte. Examples of lymphocytes include T cells, B cells, Natural Killer (NK) cells or NKT cells. In some embodiments, a T-cell is a CD4⁺ Th (T helper) cell, a CD8⁺ cytotoxic T cell, a $\gamma\delta$ T cell or a regulatory (suppressor) T cell. In some embodiments, an immune cell is a dendritic cell.

[000314] Immune cells of the present disclosure, in some embodiments, are genetically engineered to express an antigen-binding receptor. A cell is considered "engineered" if it contains an engineered (exogenous) nucleic acid. Engineered nucleic acids of the present disclosure may be introduced into a cell by any known (e.g., conventional) method. For example, an engineered nucleic acid may be introduced into a cell by electroporation (see, e.g., Heiser W. C. Transcription Factor Protocols: Methods in Molecular Biology. TM. 2000; 130: 117-134), chemical (e.g., calcium phosphate or lipid), transfection (see, e.g., Lewis W. H., et al., Somatic Cell Genet. 1980 May; 6(3): 333-47; Chen C., et al., Mol Cell Biol. 1987 August; 7(8): 2745-2752), fusion with bacterial protoplasts containing recombinant plasmids (see, e.g., Schaffner W. Proc Natl Acad Sci USA. 1980 April; 77(4): 2163-7), microinjection of purified DNA directly into the nucleus of the cell (see, e.g., Capecchi M. R. Cell. 1980 November; 22(2 Pt 2): 479-88), or retrovirus transduction.

[000315] Some aspects of the present disclosure provide an "adoptive cell" approach, which involves isolating immune cells (e.g., T-cells) from a subject with cancer, genetically

engineering the immune cells (e.g., to express an antigen-binding receptor, such as a chimeric antigen receptor), expanding the cells ex vivo, and then re-introducing the immune cells into the subject. This method results in a greater number of engineered immune cells in the subject relative to what could be achieved by conventional gene delivery and vaccination methods. In some embodiments, immune cells are isolated from a subject, expanded ex vivo without genetic modification, and then re-introduced into the subject.

[000316] Immune cells of the present disclosure comprise receptors that bind to antigens, such as an antigen encoded by an exogenously delivered nucleic acid, as provided herein. In some embodiments, a leukocyte is modified (e.g., genetically modified) to express a receptor that binds to an antigen. The receptor may be, in some embodiments, a naturally-occurring antigen receptor (normally expressed on the immune cell), recombinant antigen receptor (not normally expressed on the immune cell) or a chimeric antigen receptor (CAR). Naturally-occurring and recombinant antigen receptors encompassed by the present disclosure include T cell receptors, B cell receptors, NK cell receptors, NKT cell receptors and dendritic cell receptors. A "chimeric antigen receptor" refers to an artificial immune cell receptor that is engineered to recognize and bind to an antigen expressed by tumor cells. Generally, a CAR is designed for a T cell and is a chimera of a signaling domain of the T-cell receptor (TcR) complex and an antigen-recognizing domain (e.g., a single chain fragment (scFv) of an antibody) (Enblad et al., Human Gene Therapy. 2015; 26(8):498-505), the disclosure of which is incorporated herein by reference in its entirety.

[000317] In some embodiments, an antigen binding receptor is a chimeric antigen receptor (CAR). A T cell that expressed a CAR is referred to as a "CAR T cell." A CAR T cell receptor, in some embodiments, comprises a signaling domain of the T-cell receptor (TcR) complex and an antigen-recognizing domain (e.g., a single chain fragment (scFv) of an antibody) (Enblad et al., Human Gene Therapy. 2015; 26(8):498-505) the disclosure of which is incorporated herein by reference in its entirety.

[000318] There are four generations of CARs, each of which contains different components. First generation CARs join an antibody-derived scFv to the CD3zeta (zeta. or z) intracellular signaling domain of the T-cell receptor through hinge and transmembrane domains. Second generation CARs incorporate an additional domain, e.g., CD28, 4-1BB (41BB), or ICOS, to supply a costimulatory signal. Third-generation CARs contain two costimulatory domains fused

with the TcR CD3-zeta chain. Third-generation costimulatory domains may include, e.g., a combination of CD3z, CD27, CD28, 4-1BB, ICOS, or OX40. CARs, in some embodiments, contain an ectodomain (e.g., CD3), commonly derived from a single chain variable fragment (scFv), a hinge, a transmembrane domain, and an endodomain with one (first generation), two (second generation), or three (third generation) signaling domains derived from CD3Z and/or co-stimulatory molecules (Maude et al., *Blood*. 2015; 125(26):4017-4023; Kakarla and Gottschalk, *Cancer J*. 2014; 20(2):151-155) the disclosure of which is incorporated herein by reference in its entirety.

[000319] In some embodiments, the chimeric antigen receptor (CAR) is a T-cell redirected for universal cytokine killing (TRUCK), also known as a fourth generation CAR. TRUCKs are CAR-redirectioned T-cells used as vehicles to produce and release a transgenic cytokine that accumulates in the targeted tissue, e.g., a targeted tumor tissue. The transgenic cytokine is released upon CAR engagement of the target. TRUCK cells may deposit a variety of therapeutic cytokines in the target. This may result in therapeutic concentrations at the targeted site and avoid systemic toxicity.

[000320] CARs typically differ in their functional properties. The CD3zeta signaling domain of the T-cell receptor, when engaged, will activate and induce proliferation of T-cells but can lead to anergy (a lack of reaction by the body's defense mechanisms, resulting in direct induction of peripheral lymphocyte tolerance). Lymphocytes are considered anergic when they fail to respond to a specific antigen. The addition of a costimulatory domain in second-generation CARs improved replicative capacity and persistence of modified T-cells. Similar antitumor effects are observed in vitro with CD28 or 4-1BB CARs, but preclinical in vivo studies suggest that 4-1BB CARs may produce superior proliferation and/or persistence. Clinical trials suggest that both of these second-generation CARs are capable of inducing substantial T-cell proliferation in vivo, but CARs containing the 4-1BB costimulatory domain appear to persist longer. Third generation CARs combine multiple signaling domains (costimulatory) to augment potency. Fourth generation CARs are additionally modified with a constitutive or inducible expression cassette for a transgenic cytokine, which is released by the CAR T-cell to modulate the T-cell response. See, for example, Enblad et al., *Human Gene Therapy*. 2015; 26(8):498-505; Chmielewski and Hinrich, *Expert Opinion on Biological*

Therapy. 2015; 15(8): 1145-1154 the disclosures of which are incorporated herein by reference in their entireties.

[000321] In some embodiments, an illustrative immunotherapeutic agent is a first generation chimeric antigen receptor CAR. In some embodiments, a chimeric antigen receptor is a third generation CAR. In some embodiments, a chimeric antigen receptor is a second generation CAR. In some embodiments, a chimeric antigen receptor is a third generation CAR. In some embodiments, the chimeric antigen receptor is a fourth generation CAR or a T-cell redirected for universal cytokine killing (TRUCK).

[000322] In some embodiments, a chimeric antigen receptor (CAR) comprises an extracellular domain comprising an antigen binding domain, a transmembrane domain, and a cytoplasmic domain. In some embodiments, a CAR is fully human. In some embodiments, the antigen binding domain of a CAR is specific for one or more antigens. In some embodiments, a "spacer" domain or "hinge" domain is located between an extracellular domain (comprising the antigen binding domain) and a transmembrane domain of a CAR, or between a cytoplasmic domain and a transmembrane domain of the CAR. A "spacer domain" refers to any oligopeptide or polypeptide that functions to link the transmembrane domain to the extracellular domain and/or the cytoplasmic domain in the polypeptide chain. A "hinge domain" refers to any oligopeptide or polypeptide that functions to provide flexibility to the CAR, or domains thereof, or to prevent steric hindrance of the CAR, or domains thereof. In some embodiments, a spacer domain or hinge domain may comprise up to 300 amino acids (e.g., 10 to 100 amino acids, or 5 to 20 amino acids). In some embodiments, one or more spacer domain(s) may be included in other regions of a CAR.

[000323] In some embodiments, a CAR of the disclosure comprises an antigen binding domain, such as a single chain Fv (scFv) specific for a tumor antigen. The choice of binding domain depends upon the type and number of ligands that define the surface of a target cell. For example, the antigen binding domain may be chosen to recognize a ligand that acts as a cell surface marker on target cells associated with a particular disease state, such as cancer or an autoimmune disease. Thus, examples of cell surface markers that may act as ligands for the antigen binding domain in the CAR of the present disclosure include those associated with cancer cells and/or other forms of diseased cells. In some embodiments, a CAR is engineered to target a tumor antigen of interest by way of engineering a desired antigen binding domain that

specifically binds to an antigen on a tumor cell encoded by an engineered nucleic acid, as provided herein.

[000324] An antigen binding domain (e.g., an scFv) that "specifically binds" to a target or an epitope is a term understood in the art, and methods to determine such specific binding are also known in the art. A molecule is said to exhibit "specific binding" if it reacts or associates more frequently, more rapidly, with greater duration and/or with greater affinity with a particular target antigen than it does with alternative targets. An antigen binding domain (e.g., an scFv) that specifically binds to a first target antigen may or may not specifically bind to a second target antigen. As such, "specific binding" does not necessarily require (although it can include) exclusive binding.

[000325] In some embodiments, immune cells expressing a CAR are genetically modified to recognize multiple targets or antigens, which permits the recognition of unique target or antigen expression patterns on tumor cells. Examples of CARs that can bind multiple targets include: "split signal CARs," which limit complete immune cell activation to tumors expressing multiple antigens; "tandem CARs" (TanCARs), which contain ectodomains having two scFvs; and "universal ectodomain CARs," which incorporate avidin or a fluorescein isothiocyanate (FITC)-specific scFv to recognize tumor cells that have been incubated with tagged monoclonal antibodies (Mabs).

[000326] A CAR is considered "bispecific" if it recognizes two distinct antigens (has two distinct antigen recognition domains). In some embodiments, a bispecific CAR is comprised of two distinct antigen recognition domains present in tandem on a single transgenic receptor (referred to as a TanCAR; see, e.g., Grada Z et al. *Molecular Therapy Nucleic Acids* 2013; 2:e105, incorporated herein by reference in its entirety). Thus, methods, in some embodiments, comprise delivering to a tumor a combination comprising a compound of Formula I' and an immunotherapeutic agent, wherein the immunotherapeutic agent is an engineered nucleic acid that encodes an antigen, or delivering to a tumor an engineered nucleic acid that induces expression of a self-antigen, and delivering to the tumor an immune cell expressing a bispecific CAR that binds to two antigens, one of which is encoded by the engineered nucleic acid.

[000327] In some embodiments, a CAR is an antigen-specific inhibitory CAR (iCAR), which may be used, for example, to avoid off-tumor toxicity (Fedorov, V D et al. *Sci. Transl. Med.* published online Dec. 11, 2013, incorporated herein by reference in its entirety). iCARs

contain an antigen-specific inhibitory receptor, for example, to block nonspecific immunosuppression, which may result from extra tumor target expression. iCARs may be based, for example, on inhibitory molecules CTLA-4 or PD-1. In some embodiments, these iCARs block T cell responses from T cells activated by either their endogenous T cell receptor or an activating CAR. In some embodiments, this inhibiting effect is temporary.

[000328] In some embodiments, CARs may be used in adoptive cell transfer, wherein immune cells are removed from a subject and modified so that they express receptors specific to an antigen, e.g., a tumor-specific antigen. The modified immune cells, which may then recognize and kill the cancer cells, are reintroduced into the subject (Pule, et al., *Cytotherapy*. 2003; 5(3): 211-226; Maude et al., *Blood*. 2015; 125(26): 4017-4023, each of which is incorporated herein by reference in their entireties).

[000329] According to other aspects of the disclosure, the tumor antigenic component in the vaccine of the invention is any natural or synthetic tumor-associated protein or peptide or combination of tumor-associated proteins and/or peptides or glycoproteins or glycopeptides. In still yet other aspects, the antigenic component can be patient-specific or common to many or most patients with a particular type of cancer. According to one aspect, the antigenic component consists of a cell lysate derived from tumor tissue removed from the patient being treated. In another aspect, the lysate can be engineered or synthesized from exosomes derived from tumor tissue. In yet another aspect, the antigenic component consists of a cell lysate derived from tumor tissue extracted from one or more unrelated individuals or from tumor-cell lines.

[000330] In various embodiments, an illustrative immunotherapeutic agent comprises one or more cancer vaccines, for use in combination with a compound of Formula I'. The tumor-associated antigen component of the vaccine may be manufactured by any of a variety of well-known techniques. For individual protein components, the antigenic protein is isolated from tumor tissue or a tumor-cell line by standard chromatographic means such as high-pressure liquid chromatography or affinity chromatography or, alternatively, it is synthesized by standard recombinant DNA technology in a suitable expression system, such as *E. coli*, yeast or plants. The tumor-associated antigenic protein is then purified from the expression system by standard chromatographic means. In the case of peptide antigenic components, these are generally prepared by standard automated synthesis. Proteins and peptides can be modified by addition of amino acids, lipids and other agents to improve their incorporation into the delivery system of

the vaccine (such as a multilamellar liposome). For a tumor-associated antigenic component derived from the patient's own tumor, or tumors from other individuals, or cell lines, the tumor tissue, or a single cell suspension derived from the tumor tissue, is typically homogenized in a suitable buffer. The homogenate can also be fractionated, such as by centrifugation, to isolate particular cellular components such as cell membranes or soluble material. The tumor material can be used directly or tumor-associated antigens can be extracted for incorporation in the vaccine using a buffer containing a low concentration of a suitable agent such as a detergent. An example of a suitable detergent for extracting antigenic proteins from tumor tissue, tumor cells, and tumor-cell membranes is diheptanoyl phosphatidylcholine. Exosomes derived from tumor tissue or tumor cells, whether autologous or heterologous to the patient, can be used for the antigenic component for incorporation in the vaccine or as a starting material for extraction of tumor-associated antigens.

[000331] In some embodiments of the present disclosure, a cancer vaccine, wherein the cancer vaccine includes at least one tumor-associated antigen, at least one immunostimulant, and optionally, at least one cell-based immunotherapeutic agent. In some embodiments, the immunostimulant component in the cancer vaccine of the disclosure is any Biological Response Modifier (BRM) with the ability to enhance the therapeutic cancer vaccine's effectiveness to induce humoral and cellular immune responses against cancer cells in a patient. According to one aspect, the immunostimulant is a cytokine or combination of cytokines. Examples of such cytokines include the interferons, such as IFN-gamma, the interleukins, such as IL-2, IL-15 and IL-23, the colony stimulating factors, such as M-CSF and GM-CSF, and tumor necrosis factor. According to another aspect, the immunostimulant component of the disclosed cancer vaccine includes one or more adjuvant-type immunostimulatory agents such as APC Toll-like Receptor agonists or costimulatory/cell adhesion membrane proteins, with or without immunostimulatory cytokines. Examples of Toll-like Receptor agonists include lipid A and CpG, and costimulatory/adhesion proteins such as CD80, CD86, and ICAM-1.

[000332] In some embodiments, the immunostimulant is selected from the group consisting of IFN-gamma (IFN- γ), IL-2, IL-15, IL-23, M-CSF, GM-CSF, tumor necrosis factor, lipid A, CpG, CD80, CD86, and ICAM-1, or combinations thereof. According to other aspects, the cell-based immunotherapeutic agent is selected from the group consisting of dendritic cells, tumor-infiltrating T lymphocytes, chimeric antigen receptor-modified T effector cells directed to the

patient's tumor type, B lymphocytes, natural killer cells, bone marrow cells, and any other cell of a patient's immune system, or combinations thereof. In one aspect, the cancer vaccine immunostimulant includes one or more cytokines, such as interleukin 2 (IL-2), GM-CSF, M-CSF, and interferon-gamma (IFN- γ), one or more Toll-like Receptor agonists and/or adjuvants, such as monophosphoryl lipid A, lipid A, muramyl dipeptide (MDP) lipid conjugate and double stranded RNA, or one or more costimulatory membrane proteins and/or cell adhesion proteins, such as CD80, CD86 and ICAM-1, or any combination of the above. In one aspect, the cancer vaccine includes an immunostimulant that is a cytokine selected from the group consisting of interleukin 2 (IL-2), GM-CSF, M-CSF, and interferon-gamma (IFN- γ). In another aspect, the cancer vaccine includes an immunostimulant that is a Toll-like Receptor agonist and/or adjuvant selected from the group consisting of monophosphoryl lipid A, lipid A, and muramyl dipeptide (MDP) lipid conjugate and double stranded RNA. In yet another aspect, the cancer vaccine includes an immunostimulant that is a costimulatory membrane protein and/or cell adhesion protein selected from the group consisting of CD80, CD86, and ICAM-1.

[000333] In various embodiments, an immunotherapeutic agent can include a cancer vaccine, wherein the cancer vaccine incorporates any tumor antigen that can be potentially used to construct a fusion protein according to the invention and particularly the following:

[000334] (a) cancer-testis antigens including NY-ESO-1, SSX2, SCP1 as well as RAGE, BAGE, GAGE and MAGE family polypeptides, for example, GAGE-1, GAGE-2, MAGE-1, MAGE-2, MAGE-3, MAGE-4, MAGE-5, MAGE-6, and MAGE-12, which can be used, for example, to address melanoma, lung, head and neck, NSCLC, breast, gastrointestinal, and bladder tumors; (b) mutated antigens, including p53, associated with various solid tumors, e.g., colorectal, lung, head and neck cancer; p21/Ras associated with, e.g., melanoma, pancreatic cancer and colorectal cancer; CDK4, associated with, e.g., melanoma; MUM1 associated with, e.g., melanoma; caspase-8 associated with, e.g., head and neck cancer; CIA 0205 associated with, e.g., bladder cancer; HLA-A2-R1701, beta catenin associated with, e.g., melanoma; TCR associated with, e.g., T-cell non-Hodgkin lymphoma; BCR-abl associated with, e.g., chronic myelogenous leukemia; triosephosphate isomerase; KIA 0205; CDC-27, and LDLR-FUT; (c) over-expressed antigens, including, Galectin 4 associated with, e.g., colorectal cancer; Galectin 9 associated with, e.g., Hodgkin's disease; proteinase 3 associated with, e.g., chronic myelogenous leukemia; WT 1 associated with, e.g., various leukemias; carbonic anhydrase

associated with, e.g., renal cancer; aldolase A associated with, e.g., lung cancer; PRAME associated with, e.g., melanoma; HER-2/neu associated with, e.g., breast, colon, lung and ovarian cancer; mammaglobin, alpha-fetoprotein associated with, e.g., hepatoma; KSA associated with, e.g., colorectal cancer; gastrin associated with, e.g., pancreatic and gastric cancer; telomerase catalytic protein, MUC-1 associated with, e.g., breast and ovarian cancer; G-250 associated with, e.g., renal cell carcinoma; p53 associated with, e.g., breast, colon cancer; and carcinoembryonic antigen associated with, e.g., breast cancer, lung cancer, and cancers of the gastrointestinal tract such as colorectal cancer; (d) shared antigens, including melanoma-melanocyte differentiation antigens such as MART-1/Melan A; gp100; MC1R; melanocyte-stimulating hormone receptor; tyrosinase; tyrosinase related protein-1/TRP1 and tyrosinase related protein-2/TRP2 associated with, e.g., melanoma; (e) prostate associated antigens including PAP, PSA, PSMA, PSH-P1, PSM-P1, PSM-P2, associated with e.g., prostate cancer; (f) immunoglobulin idiotypes associated with myeloma and B cell lymphomas. In certain embodiments, the one or more TAA can be selected from pi 5, Hom/Mel-40, H-Ras, E2A-PRL, H4-RET, IGH-IGK, MYL-RAR, Epstein Barr virus antigens, EBNA, human papillomavirus (HPV) antigens, including E6 and E7, hepatitis B and C virus antigens, human T-cell lymphotropic virus antigens, TSP-180, pl85erbB2, pl 80erbB-3, c-met, mn-23H1, TAG-72-4, CA 19-9, CA 72-4, CAM 17.1, NuMa, K-ras, pi 6, TAGE, PSCA, CT7, 43-9F, 5T4, 791 Tgp72, beta-HCG, BCA225, BTAA, CA 125, CA 15-3 (CA 27.29\BCAA), CA 195, CA 242, CA-50, CAM43, CD68\KP1, CO-029, FGF-5, Ga733 (EpCAM), HTgp-175, M344, MA-50, MG7-Ag, MOV18, NB/70K, NY-CO-1, RCAS1, SDCCAG16, TA-90 (Mac-2 binding protein/cyclophilin C-associated protein), TAAL6, TAG72, TLP, TPS or any combinations thereof.

[000335] In some embodiments, cancer vaccines of the present disclosure for use in combination with a compound of Formula I' can include a tumor antigen comprising the entire amino acid sequence, a portion of it, or specific immunogenic epitopes of one of the following human proteins: TCTN1 (Gene ID: ENSG00000204852), TCTN2 (Gene ID: ENSG00000168778), TCTN3 (Gene ID: ENSG00000119977), HIGD2A (Gene ID: ENSG00000146066), HIGD2B (Gene ID: ENSG00000175202), C4ORF32 (Gene ID: ENSG00000174749), FAM62A (E-SYT1, Gene ID: ENSG00000139641), COLEC11 (Gene ID: ENSG00000118004), FSTL5 (Gene ID: ENSG00000168843), FAM82A2 (Gene ID: ENSG00000137824), SCARA5 (Gene ID: ENSG00000168079), VSTM1 (Gene ID:

ENSG00000189068), RNF5 (Gene ID: ENSG00000183574), UNQ6126 (Gene ID: gi|169216088), DPY19L3 (Gene ID: ENSG00000178904), SLC39A10 (gene ID: ENSG00000196950), GPR107 (Gene ID: ENSG00000148358), COL20A1 (Gene ID: ENSG00000101203), GLT25D2 (Gene ID: ENSG00000198756), SYTL3 (Gene ID: ENSG00000164674), DENND1B (Gene ID: ENSG00000162701), C6orf98 (Gene ID: EG: 387079), FAM69B (Gene ID: ENSG00000165716), EMID1 (Gene ID: OTTHUMG00000030824), KLRG2 (GENE ID: ENSG00000188883), ERMP1 (GENE ID: ENSG00000099219), VMO1 (Gene ID: ENSG00000182853), C9orf46 (Gene ID: ENSG00000107020), F1137107 (Gene ID: ENSG00000177990), YIPF2 (Gene ID: ENSG00000130733), TRYX3 (PRSS58, ENSG00000258223.2), C14orf135 (Gene ID: ENSG00000126773), ANGPTL7 (Gene ID: ENSG00000171819), TPCN2 (Gene ID: ENSG00000162341), C18orf19 (Gene ID: ENSG00000177150), OLFML1 (Gene ID: ENSG00000183801), LYPD4 (Gene ID: ENSG00000101203), MEGF8 (Gene ID: ENSG00000105429), F1142986 (Gene ID: ENSG00000196460), SLC46A1 (Gene ID: ENSG00000076351), FAM180A (Gene ID: ENSG00000189320), CRISP-3 (GENE ID: ENSG00000096006), or combinations thereof. These tumor antigens are disclosed in WO2010/086162, WO2010/086163, WO2011/051278, WO2011/051276, WO2011/051277, WO2011/051280, WO2011/051271, WO2011/135068, WO2014/198919, the content of which is herein incorporated by reference in their entireties.

[000336] In various embodiments, an illustrative immunotherapeutic agent may include an mRNA operable to encode any one or more of the aforementioned cancer antigens useful for synthesizing a cancer vaccine. In some illustrative embodiments, the mRNA based cancer vaccine may have one or more of the following properties: a) the mRNA encoding each cancer antigen is interspersed by cleavage sensitive sites; b) the mRNA encoding each cancer antigen is linked directly to one another without a linker; c) the mRNA encoding each cancer antigen is linked to one another with a single nucleotide linker; d) each cancer antigen comprises a 20-40 amino acids and includes a centrally located SNP mutation; e) at least 40% of the cancer antigens have a highest affinity for class I MHC molecules from the subject; f) at least 40% of the cancer antigens have a highest affinity for class II MHC molecules from the subject; g) at least 40% of the cancer antigens have a predicted binding affinity of $IC_{50} > 500$ nM for HLA-A, HLA-B and/or DRB1; h) the mRNA encodes 1 to 15 cancer antigens; i) 10-60% of the cancer

antigens have a binding affinity for class I MHC and 10-60% of the cancer antigens have a binding affinity for class II MHC; and/or j) the mRNA encoding the cancer antigens is arranged such that the cancer antigens are ordered to minimize pseudo-epitopes.

[000337] In various embodiments, the combination comprising a compound of Formula I' and a cancer vaccine immunotherapeutic agent as disclosed herein can be used to illicit an immune response in a subject against a cancer antigen. The method involves administering to the subject a RNA vaccine comprising at least one RNA polynucleotide having an open reading frame encoding at least one antigenic polypeptide or an immunogenic fragment thereof, thereby inducing in the subject an immune response specific to the antigenic polypeptide or an immunogenic fragment thereof, in combination with administering a compound of Formula I' either in the same composition or a separate composition, administered at the same time, or sequentially dosed, wherein the anti-antigenic polypeptide antibody titer in the subject is increased following vaccination relative to anti-antigenic polypeptide antibody titer in a subject vaccinated with a prophylactically effective dose of a traditional vaccine against the cancer. An "anti-antigenic polypeptide antibody" is a serum antibody the binds specifically to the antigenic polypeptide.

[000338] A prophylactically effective dose is a therapeutically effective dose that prevents advancement of cancer at a clinically acceptable level. In some embodiments the therapeutically effective dose is a dose listed in a package insert for the vaccine. A traditional vaccine, as used herein, refers to a vaccine other than the mRNA vaccines of the invention. For instance, a traditional vaccine includes but is not limited to live microorganism vaccines, killed microorganism vaccines, subunit vaccines, protein antigen vaccines, DNA vaccines, and the like. In exemplary embodiments, a traditional vaccine is a vaccine that has achieved regulatory approval and/or is registered by a national drug regulatory body, for example the Food and Drug Administration (FDA) in the United States or the European Medicines Agency (EMA.)

[000339] In some embodiments the anti-antigenic polypeptide antibody titer in the subject is increased 1 log to 10 log following vaccination relative to anti-antigenic polypeptide antibody titer in a subject vaccinated with a prophylactically effective dose of a traditional vaccine against the cancer. In some embodiments the anti-antigenic polypeptide antibody titer in the subject is increased 1 log following vaccination relative to anti-antigenic polypeptide antibody titer in a subject vaccinated with a prophylactically effective dose of a traditional vaccine

against the cancer. In some embodiments the anti-antigenic polypeptide antibody titer in the subject is increased 2 log following vaccination relative to anti-antigenic polypeptide antibody titer in a subject vaccinated with a prophylactically effective dose of a traditional vaccine against the cancer.

[000340] Aspects of the invention provide nucleic acid vaccines comprising one or more RNA polynucleotides having an open reading frame encoding a first antigenic polypeptide, wherein the RNA polynucleotide is present in the formulation for in vivo administration to a host, which confers an antibody titer superior to the criterion for seroprotection for the first antigen for an acceptable percentage of human subjects. In some embodiments, the antibody titer produced by the mRNA vaccines of the invention is a neutralizing antibody titer. In some embodiments the neutralizing antibody titer is greater than a protein vaccine. In other embodiments the neutralizing antibody titer produced by the mRNA vaccines of the invention is greater than an adjuvanted protein vaccine. In yet other embodiments the neutralizing antibody titer produced by the mRNA vaccines of the invention is 1,000-10,000, 1,200-10,000, 1,400-10,000, 1,500-10,000, 1,000-5,000, 1,000-4,000, 1,800-10,000, 2000-10,000, 2,000-5,000, 2,000-3,000, 2,000-4,000, 3,000-5,000, 3,000-4,000, or 2,000-2,500. A neutralization titer is typically expressed as the highest serum dilution required to achieve a 50% reduction in the number of plaques.

[000341] In preferred aspects, RNA vaccine immunotherapeutic agents of the present disclosure (e.g., mRNA vaccines) produce prophylactically- and/or therapeutically-efficacious levels, concentrations and/or titers of antigen-specific antibodies in the blood or serum of a vaccinated subject. As defined herein, the term antibody titer refers to the amount of antigen-specific antibody produces in s subject, e.g., a human subject. In exemplary embodiments, antibody titer is expressed as the inverse of the greatest dilution (in a serial dilution) that still gives a positive result. In exemplary embodiments, antibody titer is determined or measured by enzyme-linked immunosorbent assay (ELISA). In exemplary embodiments, antibody titer is determined or measured by neutralization assay, e.g., by microneutralization assay. In certain aspects, antibody titer measurement is expressed as a ratio, such as 1:40, 1:100, and the like.

[000342] In exemplary embodiments of the invention, an efficacious vaccine produces an antibody titer of greater than 1:40, greater that 1:100, greater than 1:400, greater than 1:1000, greater than 1:2000, greater than 1:3000, greater than 1:4000, greater than 1:500, greater than

1:6000, greater than 1:7500, greater than 1:10000. In exemplary embodiments, the antibody titer is produced or reached by 10 days following vaccination, by 20 days following vaccination, by 30 days following vaccination, by 40 days following vaccination, or by 50 or more days following vaccination. In exemplary embodiments, the titer is produced or reached following a single dose of vaccine administered to the subject. In other embodiments, the titer is produced or reached following multiple doses, e.g., following a first and a second dose (e.g., a booster dose.) In exemplary aspects of the invention, antigen-specific antibodies are measured in units of g/ml or are measured in units of IU/L (International Units per liter) or mIU/ml (milli International Units per ml). In exemplary embodiments of the invention, an efficacious vaccine produces $>0.5 \mu\text{g/mL}$, $>0.1 \mu\text{g/mL}$, $>0.2 \mu\text{g/mL}$, $>0.35 \mu\text{g/mL}$, $>0.5 \mu\text{g/mL}$, $>1 \mu\text{g/mL}$, $>2 \mu\text{g/mL}$, $>5 \mu\text{g/mL}$ or $>10 \mu\text{g/mL}$. In exemplary embodiments of the invention, an efficacious vaccine produces $>10 \text{ mIU/mL}$, $>20 \text{ mIU/mL}$, $>50 \text{ mIU/mL}$, $>100 \text{ mIU/mL}$, $>200 \text{ mIU/mL}$, $>500 \text{ mIU/mL}$ or $>1000 \text{ mIU/mL}$. In exemplary embodiments, the antibody level or concentration is produced or reached by 10 days following vaccination, by 20 days following vaccination, by 30 days following vaccination, by 40 days following vaccination, or by 50 or more days following vaccination. In exemplary embodiments, the level or concentration is produced or reached following a single dose of vaccine administered to the subject. In other embodiments, the level or concentration is produced or reached following multiple doses, e.g., following a first and a second dose (e.g., a booster dose.) In exemplary embodiments, antibody level or concentration is determined or measured by enzyme-linked immunosorbent assay (ELISA). In exemplary embodiments, antibody level or concentration is determined or measured by neutralization assay, e.g., by microneutralization assay. Also provided are nucleic acid vaccines comprising one or more RNA polynucleotides having an open reading frame encoding a first antigenic polypeptide or a concatemeric polypeptide, wherein the RNA polynucleotide is present in a formulation for in vivo administration to a host for eliciting a longer lasting high antibody titer than an antibody titer elicited by an mRNA vaccine having a stabilizing element or formulated with an adjuvant and encoding the first antigenic polypeptide. In some embodiments, the RNA polynucleotide is formulated to produce a neutralizing antibodies within one week of a single administration. In some embodiments, the adjuvant is selected from a cationic peptide and an immunostimulatory nucleic acid. In some embodiments, the cationic peptide is protamine.

[000343] Immunotherapeutic agents comprising a nucleic acid vaccine comprising one or more RNA polynucleotides having an open reading frame comprising at least one chemical modification or optionally no nucleotide modification, the open reading frame encoding a first antigenic polypeptide or a concatemeric polypeptide, wherein the RNA polynucleotide is present in the formulation for in vivo administration to a host such that the level of antigen expression in the host significantly exceeds a level of antigen expression produced by an mRNA vaccine having a stabilizing element or formulated with an adjuvant and encoding the first antigenic polypeptide.

[000344] Other aspects provide nucleic acid vaccines comprising one or more RNA polynucleotides having an open reading frame comprising at least one chemical modification or optionally no nucleotide modification, the open reading frame encoding a first antigenic polypeptide or a concatemeric polypeptide, wherein the vaccine has at least 10 fold less RNA polynucleotide than is required for an unmodified mRNA vaccine to produce an equivalent antibody titer. In some embodiments, the RNA polynucleotide is present in a dosage of 25-100 micrograms.

[000345] Aspects of the invention also provide a unit of use vaccine, comprising between 10 μ g and 400 μ g of one or more RNA polynucleotides having an open reading frame comprising at least one chemical modification or optionally no nucleotide modification, the open reading frame encoding a first antigenic polypeptide or a concatemeric polypeptide, and a pharmaceutically acceptable carrier or excipient, formulated for delivery to a human subject. In some embodiments, the vaccine further comprises a cationic lipid nanoparticle.

[000346] Aspects of the invention provide methods of creating, maintaining or restoring antigenic memory to a tumor in an individual or population of individuals comprising administering to said individual or population an antigenic memory booster nucleic acid vaccine comprising (a) at least one RNA polynucleotide, said polynucleotide comprising at least one chemical modification or optionally no nucleotide modification and two or more codon-optimized open reading frames, said open reading frames encoding a set of reference antigenic polypeptides, and (b) optionally a pharmaceutically acceptable carrier or excipient. In some embodiments, the vaccine is administered to the individual via a route selected from the group consisting of intramuscular administration, intradermal administration and subcutaneous administration. In some embodiments, the administering step comprises contacting a muscle

tissue of the subject with a device suitable for injection of the composition. In some embodiments, the administering step comprises contacting a muscle tissue of the subject with a device suitable for injection of the composition in combination with electroporation.

[000347] Aspects of the invention provide methods of vaccinating a subject comprising administering to the subject a single dosage of between 25 µg /kg and 400 µg /kg of a nucleic acid vaccine comprising one or more RNA polynucleotides having an open reading frame encoding a first antigenic polypeptide or a concatemeric polypeptide in an effective amount to vaccinate the subject.

[000348] Other aspects provide nucleic acid vaccines comprising one or more RNA polynucleotides having an open reading frame comprising at least one chemical modification, the open reading frame encoding a first antigenic polypeptide or a concatemeric polypeptide, wherein the vaccine has at least 10 fold less RNA polynucleotide than is required for an unmodified mRNA vaccine to produce an equivalent antibody titer. In some embodiments, the RNA polynucleotide is present in a dosage of 25-100 micrograms.

[000349] In some embodiments, an illustrative immunotherapeutic agent can include one or more interfering RNAs that can be administered in combination with a compound of Formula I'. An "RNA interfering agent" as used herein, is defined as any agent which interferes with or inhibits expression of a target biomarker gene by RNA interference (RNAi). Such RNA interfering agents include, but are not limited to, nucleic acid molecules including RNA molecules which are homologous to the target biomarker gene of the present invention, or a fragment thereof, short interfering RNA (siRNA), and small molecules which interfere with or inhibit expression of a target biomarker nucleic acid by RNA interference (RNAi). Short interfering RNA" (siRNA), also referred to herein as "small interfering RNA" is defined as an agent which functions to inhibit expression of a target biomarker nucleic acid, e.g., by RNAi. An siRNA may be chemically synthesized, may be produced by in vitro transcription, or may be produced within a host cell. In one embodiment, siRNA is a double stranded RNA (dsRNA) molecule of about 15 to about 40 nucleotides in length, preferably about 15 to about 28 nucleotides, more preferably about 19 to about 25 nucleotides in length, and more preferably about 19, 20, 21, or 22 nucleotides in length, and may contain a 3' and/or 5' overhang on each strand having a length of about 0, 1, 2, 3, 4, or 5 nucleotides. The length of the overhang is independent between the two strands, i.e., the length of the overhang on one strand is not

dependent on the length of the overhang on the second strand. Preferably the siRNA is capable of promoting RNA interference through degradation or specific post-transcriptional gene silencing (PTGS) of the target messenger RNA (mRNA).

[000350] An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides in length. An antisense nucleic acid can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been sub-cloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

[000351] The antisense nucleic acid molecules of the present invention are typically administered to a subject or generated in situ such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a polypeptide corresponding to a selected marker of the present invention to thereby inhibit expression of the marker, e.g., by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to

form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. Examples of a route of administration of antisense nucleic acid molecules of the present invention includes direct injection at a tissue site or infusion of the antisense nucleic acid into a blood- or bone marrow-associated body fluid. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, e.g., by linking the antisense nucleic acid molecules to peptides or antibodies which bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

[000352] Antigens which can be targeted for synthesizing a corresponding antisense RNA molecule can include any antigen that is specific for one or more tumors, for example, antigens exemplified above with reference to cancer vaccines.

[000353] In some embodiments, a combination of an immunotherapeutic agent and a compound of Formula I' can include a bispecific antibody immunotherapeutic agent. The bispecific antibody can include a protein construct having a first antigen binding moiety and a second antigen binding site that binds to a cytotoxic immune cell. The first antigen binding site can bind to a tumor antigen that is specifically being treated with the combination of the present invention. For example, the first antigen binding moiety may bind to a non-limiting example of tumor antigens selected from: EGFR, HGFR, Her2, Ep-CAM, CD20, CD30, CD33, CD47, CD52, CD133, CEA, gpA33, Mucins, TAG-72, CIX, PSMA, folate-binding protein, GD2, GD3, GM2, VEGF, VEGFR, Integrin α V β 3, Integrin α 5 β 1, MUC1, ERBB2, ERBB3, MET, IGF1R, EPHA3, TRAILR1, TRAILR2, RANKL, FAP and Tenascin among others. In some embodiments, the first antigen binding moiety has specificity to a protein or a peptide that is overexpressed on a tumor cell as compared to a corresponding non-tumor cell. In some embodiments, the first antigen binding moiety has specificity to a protein that is overexpressed on a tumor cell as compared to a corresponding non-tumor cell. A "corresponding non-tumor cell" as used here, refers to a non-tumor cell that is of the same cell type as the origin of the

tumor cell. It is noted that such proteins are not necessarily different from tumor antigens. Non-limiting examples include carcinoembryonic antigen (CEA), which is overexpressed in most colon, rectum, breast, lung, pancreas and gastrointestinal tract carcinomas; heregulin receptors (HER-2, neu or c-erbB-2), which is frequently overexpressed in breast, ovarian, colon, lung, prostate and cervical cancers; epidermal growth factor receptor (EGFR), which is highly expressed in a range of solid tumors including those of the breast, head and neck, non-small cell lung and prostate; asialoglycoprotein receptor; transferrin receptor; serpin enzyme complex receptor, which is expressed on hepatocytes; fibroblast growth factor receptor (FGFR), which is overexpressed on pancreatic ductal adenocarcinoma cells; vascular endothelial growth factor receptor (VEGFR), for anti-angiogenesis gene therapy; folate receptor, which is selectively overexpressed in 90% of nonmucinous ovarian carcinomas; cell surface glycocalyx; carbohydrate receptors; and polymeric immunoglobulin receptor.

[000354] The second antigen-binding moiety is any molecule that specifically binds to an antigen or protein or polypeptide expressed on the surface of a cytotoxic immune cell (a CIK cell). Exemplary non-limiting antigens expressed on the surface of the cytotoxic immune cells suitable for use with the present disclosure may include CD2, CD3, CD4, CD5, CD8, CD11a, CD11 b, CD14, CD16a, CD27, CD28, CD45, CD45RA, CD56, CD62L, the Fc receptor, LFA, LFA-1, TCR $\alpha\beta$, CCR7, macrophage inflammatory protein 1a, perforin, PD-1, PD-L1, PD-L2, or CTLA-4, LAG-3, OX40, 41BB, LIGHT, CD40, GITR, TGF-beta, TIM-3, SIRP-alpha, TIGIT, VSIG8, BTLA, SIGLEC7, SIGLEC9, ICOS, B7H3, B7H4, FAS, BTNL2, CD27 and Fas ligand. In some embodiments, the second antigen binding moiety binds to CD3 of the cytotoxic immune cell, e.g., CIK cell. In some embodiments, the second antigen binding moiety binds to CD56 of the cytotoxic immune cell. In some embodiments, the second antigen binding moiety binds to the Fc receptor of the cytotoxic immune cell. In some embodiments, the Fc region of the bispecific antibody binds to the Fc receptor of the cytotoxic immune cell. In some embodiments, a second antigen-binding moiety is any molecule that specifically binds to an antigen expressed on the surface of a cytotoxic immune cell (e.g., a CIK cell). The second antigen binding moiety is specific for an antigen on a cytotoxic immune cell. Exemplary cytotoxic immune cells include, but are not limited to CIK cells, T-cells, CD8⁺ T cells, activated T-cells, monocytes, natural killer (NK) cells, NK T cells, lymphokine-activated killer (LAK) cells, macrophages, and dendritic cells. The second antigen binding moiety specifically

binds to an antigen expressed on the surface of a cytotoxic immune cell. Exemplary non-limiting antigens expressed on the surface of the cytotoxic immune cells suitable for modulation with the present disclosure may include CD2, CD3, CD4, CD5, CD8, CD11a, CD11b, CD14, CD16a, CD27, CD28, CD45, CD45RA, CD56, CD62L, the Fc receptor, LFA, LFA-1, TCR $\alpha\beta$, CCR7, macrophage inflammatory protein 1a, perforin, PD-1, PD-L1, PD-L2, or CTLA-4, LAG-3, OX40, 41BB, LIGHT, CD40, GITR, TGF-beta, TIM-3, SIRP-alpha, TIGIT, VSIG8, BTLA, SIGLEC7, SIGLEC9, ICOS, B7H3, B7H4, FAS, BTNL2, CD27 and Fas ligand. In other embodiments, the bispecific antibody modulator is an activator of a costimulatory molecule (e.g., an OX40 agonist). In one embodiment, the OX40 agonist is a bispecific antibody molecule to OX40 and another tumor antigen or a costimulatory antigen. The OX40 agonist can be administered alone, or in combination with other immunomodulators, e.g., in combination with an inhibitor (for example an antibody construct) of PD-1, PD-L1, CTLA-4, CEACAM (e.g., CEACAM-1, -3 and/or -5), TIM-3 or LAG-3. In some embodiments, the anti-OX40 antibody molecule is a bispecific antibody that binds to GITR and PD-1, PD-L1, CTLA-4, CEACAM (e.g., CEACAM-1, -3 and/or -5), TIM-3 or LAG-3. In one exemplary embodiment, an OX40 antibody molecule is administered in combination with an anti-PD-1 antibody molecule (e.g., an anti-PD-1 molecule as described herein). The OX40 antibody molecule and the anti-PD-1 antibody molecule may be in the form of separate antibody composition, or as a bispecific antibody molecule. In other embodiments, the OX40 agonist can be administered in combination with other costimulatory molecule, e.g., an agonist of GITR, CD2, CD27, CD28, CD3, ICAM-1, LFA-1 (CD11a/CD18), ICOS (CD278), 4-1BB (CD137), CD30, CD40, BAFFR, HVEM, CD7, LIGHT, NKG2C, SLAMF7, NKp80, CD160, B7-H3, or CD83 ligand. In some embodiments, the second antigen binding moiety binds to the Fc receptor on the cytotoxic immune cell, e.g., CIK cell.

[000355] In some embodiments, the bispecific antibody immunotherapeutic agent has specificities for a tumor antigen and a CIK cell, which brings the tumor antigen expressing tumor cell in close proximity of the CIK cell, leading to the elimination of the tumor cell through anti-tumor cytotoxicity of CIK cell. In some embodiments, the bispecific antibody has specificity for a tumor antigen but does not have specificity for a CIK cell, however, the Fc region of the bispecific antibody can bind to the Fc receptor of the CIK cell, which in turn brings the tumor cell in close proximity of the CIK cell, leading to the elimination of the tumor

cell through anti-tumor cytotoxicity of CIK cell. In some embodiments, the bispecific antibody has specificity for a CIK cell but does not have specificity for tumor cell, however, the Fc region of the bispecific antibody can bind to the Fc receptor of the tumor cell, which in turn brings the tumor cell in close proximity of the CIK cell, leading to the elimination of the tumor cell through anti-tumor cytotoxicity of CIK cell.

[000356] In some embodiments, a combination of an immunotherapeutic agent and a compound of Formula I' can include an immune cell-engaging multivalent antibody/fusion protein/construct immunotherapeutic agent. In various embodiments, an exemplary immunotherapeutic agent can include immune cell-engaging multivalent antibody/fusion protein/construct which may comprise a recombinant structure, for example, all engineered antibodies that do not imitate the original IgG structure. Here, different strategies to multimerize antibody fragments are utilized. For example, shortening the peptide linker between the V domains forces the scFv to self-associate into a dimer (diabody; 55 kDa). Bispecific diabodies are formed by the noncovalent association of two VHA-VLB and VHB-VLA fragments expressed in the same cell. This leads to the formation of heterodimers with two different binding sites. Single-chain diabodies (sc-diabodies) are bispecific molecules where the VHA-VLB and VHB-VLA fragments are linked together by an additional third linker. Tandem-diabodies (Tandabs) are tetravalent bispecific antibodies generated by two scDiabodies.

[000357] Also included are the di-diabodies known in the art. This 130-kDa molecule is formed by the fusion of a diabody to the N-terminus of the CH3 domain of an IgG, resulting in an IgG-like structure. Further diabody derivatives are the triabody and the tetra-body, which fold into trimeric and tetrameric fragments by shortening the linker to <5 or 0–2 residues. Also exemplified are (scFv)₂ constructs known as 'bispecific T cell engager' (BITE). BITEs are bispecific single-chain antibodies consisting of two scFv antibody fragments, joined via a flexible linker, that are directed against a surface antigen on target cells and CD3 on T cells. Also exemplified are bivalent (Fab)₂ and trivalent (Fab)₃ antibody formats. Also exemplified are minibodies and trimerbodies generated from scFvs. Exemplary constructs useful to target tumor antigens as can include one or more of: Diabody, Single-chain (sc)-diabody (scFv)₂, Miniantibody, Minibody, Barnase-barstar, scFv-Fc, sc(Fab)₂, Trimeric antibody constructs, Triabody antibody constructs, Trimerbody antibody constructs, Tribody antibody constructs, Collabody antibody constructs, (scFv-TNFα)₃, F(ab)₃/DNL. In each of these

exemplified constructs, at least one binding moiety may bind to an antigen or protein or polypeptide expressed on the surface of a cytotoxic immune cell, and at least one binding moiety will bind specifically to an antigen on a cytotoxic immune cell. Exemplary cytotoxic immune cells include, but are not limited to CIK cells, T-cells, CD8⁺ T cells, activated T-cells, monocytes, natural killer (NK) cells, NK T cells, lymphokine-activated killer (LAK) cells, macrophages, and dendritic cells.

[000358] In some embodiments, a combination of an immunotherapeutic agent and a compound of Formula I' can include a radioconjugate immunotherapeutic agent.

[000359] In various embodiments, a radioconjugate is a small molecule or large molecule (herein referred to as a "cell targeting agent"), for example and polypeptide, an antibody or an antibody fragment thereof, that is coupled to or otherwise affixed to a radionuclide, or a plurality of radionuclides, such that the binding of the radioconjugate to its target (a protein or molecule on or in a cancer cell), will lead to the death or morbidity of said cancer cell. In various embodiments, the radioconjugate can be a cell targeting agent labelled with a radionuclide, or the cell targeting agent may be coupled or otherwise affixed to a particle, or microparticle, or nanoparticle containing a plurality of radionuclides, wherein the radionuclides are the same or different. Methods for synthesizing radioconjugates are known in the art, and may include the class of immunoglobulin or antigen binding parts thereof, that are conjugated to a toxic radionuclide.

[000360] In some embodiments, the molecule that binds to the cancer cell can be known as a "cell targeting agent". As used herein, an exemplary cell targeting agent can allow the drug-containing nanoparticles or radionuclide to target the specific types of cells of interest. Examples of cell targeting agents include, but are not limited to, small molecules (e.g., folate, adenosine, purine) and large molecule (e.g., peptide or antibody) that bind to or target a tumor associated antigen. Examples of tumor associated antigens include, but are not limited to, adenosine receptors, alpha v beta 3, aminopeptidase P, alpha fetoprotein, cancer antigen 125, carcinoembryonic antigen, cCaveolin-1, chemokine receptors, clusterin, oncofetal antigens, CD20, epithelial tumor antigen, melanoma associated antigen, Ras, p53, Her2/Neu, ErbB2, ErbB3, ErbB4, folate receptor, prostate-specific membrane antigen, prostate specific antigen, purine receptors, radiation-induced cell surface receptor, serpin B3, serpin B4, squamous cell carcinoma antigens, thrombospondin, tumor antigen 4, tumor-associated glycoprotein 72,

tyrosinase, and tyrosine kinases. In some embodiments, the cell targeting agent is folate or a folate derivative that binds specifically to folate receptors (FRs). In some embodiments, the cell targeting agent is an antibody, a bispecific antibody, a trispecific antibody or an antigen binding construct thereof, that specifically binds to a cancer antigen selected from: EGFR, HGFR, Her2, Ep-CAM, CD20, CD30, CD33, CD47, CD52, CD133, CEA, gpA33, Mucins, TAG-72, CIX, PSMA, folate-binding protein, GD2, GD3, GM2, VEGF, VEGFR, Integrin $\alpha V\beta 3$, Integrin $\alpha 5\beta 1$, MUC1, ERBB2, ERBB3, MET, IGF1R, EPHA3, TRAILR1, TRAILR2, RANKL, FAP and Tenascin among others.

[000361] The use of folate as a targeting agent in the radioconjugate also allow both tumor cells and regulatory T (Treg) cells to be targeted for destruction. It is well accepted that high numbers of Treg cells suppress tumor immunity. Specifically, Treg cells suppress (foreign and self) reactive T cells without killing them through contact-dependent or cytokine (e.g., IL-10, TGF- β , and the like.) secretion. FR4 is selectively upregulated on Treg cells. It has been shown that antibody blockade of FR4 depleted Treg cells and provoked tumor immunity in tumor-bearing mice. Thus, folate-coated PBM nanoparticles carrying a cytotoxic agent would take FR-expressing cells for their destruction, which would both directly (i.e., BrCa cell) and indirectly (i.e., breast tumor associated and peripheral Treg cells) inhibit tumor progression.

[000362] In another further embodiment, the targeting agent is an antibody or peptide, or immune cell-engaging multivalent antibody/fusion protein/constructs capable of binding tumor associated antigens consisting of but not limited to: adenosine receptors, $\alpha v \beta 3$, aminopeptidase P, alpha fetoprotein, cancer antigen 125, carcinoembryonic antigen, caveolin-1, chemokine receptors, clusterin, oncofetal antigens, CD20, Human Growth Factor Receptor (HGFR), epithelial tumor antigen, melanoma associated antigen, MUC1, Ras, p53, Her2/Neu, ErbB2, ErbB3, ErbB4, folate receptor, prostate-specific membrane antigen, prostate specific antigen, purine receptors, radiation-induced cell surface receptor, serpin B3, serpin B4, squamous cell carcinoma antigens, thrombospondin, tumor antigen 4, tumor-associated glycoprotein 72, tyrosinase, tyrosine kinases, and the like.

[000363] In one embodiment, the treatment method includes the co-administration of a compound as disclosed herein or a pharmaceutically acceptable salt thereof and at least one cytotoxic agent. The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents a cellular function and/or causes cell death or destruction. Cytotoxic agents include,

but are not limited to, radioactive isotopes (e.g., At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³², Pb²¹² and radioactive isotopes of Lu); chemotherapeutic agents; growth inhibitory agents; enzymes and fragments thereof such as nucleolytic enzymes; and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof.

[000364] Exemplary cytotoxic agents can be selected from anti-microtubule agents, platinum coordination complexes, alkylating agents, antibiotic agents, topoisomerase II inhibitors, antimetabolites, topoisomerase I inhibitors, hormones and hormonal analogues, signal transduction pathway inhibitors, non-receptor tyrosine kinase angiogenesis inhibitors, immunotherapeutic agents, proapoptotic agents, inhibitors of LDH-A; inhibitors of fatty acid biosynthesis; cell cycle signaling inhibitors; HDAC inhibitors, proteasome inhibitors; and inhibitors of cancer metabolism.

[000365] “Chemotherapeutic agents” include chemical compounds useful in the treatment of cancer. Examples of chemotherapeutic agents include erlotinib (TARCEVA®, Genentech/OSI Pharm.), bortezomib (VELCADE®, Millennium Pharm.), disulfiram, epigallocatechin gallate, salinosporamide A, carfilzomib, 17-AAG(geldanamycin), radicicol, lactate dehydrogenase A (LDH-A), fulvestrant (FASLODEX®, AstraZeneca), sunitib (SUTENT®, Pfizer/Sugen), letrozole (FEMARA®, Novartis), imatinib mesylate (GLEEVEC®, Novartis), finasunate (VATALANIB®, Novartis), oxaliplatin (ELOXATIN®, Sanofi), 5-FU (5-fluorouracil), leucovorin, Rapamycin (Sirolimus, RAPAMUNE®, Wyeth), Lapatinib (TYKERB®, GSK572016, Glaxo Smith Kline), Lonafamib (SCH 66336), sorafenib (NEXAVAR®, Bayer Labs), gefitinib (IRESSA®, AstraZeneca), AG1478; alkylating agents such as thiotepa and CYTOXAN®; cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramidate, triethylenethiophosphoramidate and trimethylmelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including topotecan and irinotecan); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogs); cryptophycins (particularly cryptophycin 1 and cryptophycin 8); adrenocorticosteroids (including prednisone and prednisolone); cyproterone acetate; 5 alpha-reductases including finasteride and dutasteride); vorinostat, romidepsin, panobinostat, valproic acid, mocetinostat dolastatin;

aldesleukin, talc duocarmycin (including the synthetic analogs, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlomaphazine, chlorophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosoureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (e.g., calicheamicin, especially calicheamicin gamma 1I and calicheamicin omega 1I (Angew Chem. Intl. Ed. Engl. 1994 33:183-186); dynemicin, including dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabycin, caminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, ADRIAMYCIN® (doxorubicin), morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, porfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogs such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitiolesterol, mepitiolesterol, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfomithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidamnol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, Ore.); razoxane; rhizoxin; sizofuran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine;

mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxoids, e.g., TAXOL (paclitaxel; Bristol-Myers Squibb Oncology, Princeton, N.J.), ABRAXANE® (Cremophor-free), albumin-engineered nanoparticle formulations of paclitaxel (American Pharmaceutical Partners, Schaumburg, Ill.), and TAXOTERE® (docetaxel, doxetaxel; Sanofi-Aventis); chloranmbucil; GEMZAR® (gemcitabine); 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; NAVELBINE® (vinorelbine); novantrone; teniposide; edatrexate; daunomycin; aminopterin; capecitabine (XELODA®); ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylomithine (DMFO); retinoids such as retinoic acid; and pharmaceutically acceptable salts, acids and derivatives of any of the above.

[000366] Chemotherapeutic agent also includes (i) anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including NOLVADEX®; tamoxifen citrate), raloxifene, droloxifene, iodoxyfene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and FARESTON® (toremifine citrate); (ii) aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen production in the adrenal glands, such as, for example, 4(5)-imidazoles, aminoglutethimide, MEGASE® (megestrol acetate), AROMASIN® (exemestane; Pfizer), formestanie, fadrozole, RIVISOR® (vorozole), FEMARA® (letrozole; Novartis), and ARIMIDEX® (anastrozole; AstraZeneca); (iii) anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide and goserelin; buserelin, tripterelein, medroxyprogesterone acetate, diethylstilbestrol, premarin, fluoxymesterone, all transretinoic acid, fenretinide, as well as troxacitabine (a 1,3-dioxolane nucleoside cytosine analog); (iv) protein kinase inhibitors; (v) lipid kinase inhibitors; (vi) antisense oligonucleotides, particularly those which inhibit expression of genes in signaling pathways implicated in aberrant cell proliferation, such as, for example, PKC-alpha, Ralf and H-Ras; (vii) ribozymes such as VEGF expression inhibitors (e.g., ANGIOZYME®) and HER2 expression inhibitors; (viii) vaccines such as gene therapy vaccines, for example, ALLOVECTIN®, LEUVECTIN®, and VAXID®; PROLEUKIN®, rIL-2; a topoisomerase 1 inhibitor such as LURTOTECAN®; ABARELIX®; and (ix) pharmaceutically acceptable salts, acids and derivatives of any of the above.

[000367] Chemotherapeutic agents also include antibodies, as described above, including alemtuzumab (Campath), bevacizumab (AVASTIN®, Genentech); cetuximab (ERBITUX®, Imclone); panitumumab (VECTIBIX®, Amgen), rituximab (RITUXAN®, Genentech/Biogen Idec), pertuzumab (OMNITARG®, 2C4, Genentech), trastuzumab (HERCEPTIN®, Genentech), tositumomab (Bexxar, Corixa), and the antibody drug conjugate, gemtuzumab ozogamicin (MYLOTARG®, Wyeth). Additional humanized monoclonal antibodies with therapeutic potential as agents in combination with the compounds of the invention include: apolizumab, aselizumab, atlizumab, bapineuzumab, bivatuzumab mertansine, cantuzumab mertansine, cedelizumab, certolizumab pegol, cidfusituzumab, cidtuzumab, daclizumab, eculizumab, efalizumab, epratuzumab, erlizumab, felvizumab, fontolizumab, gemtuzumab ozogamicin, inotuzumab ozogamicin, ipilimumab, labetuzumab, lintuzumab, matuzumab, mepolizumab, motavizumab, motovizumab, natalizumab, nimotuzumab, nivolumab, nolovizumab, numavizumab, ocrelizumab, omalizumab, palivizumab, pascolizumab, pecfusituzumab, pectuzumab, pexelizumab, ralivizumab, ranibizumab, reslivizumab, reslizumab, resyvizumab, rovelizumab, ruplizumab, sibrotuzumab, sipilizumab, sontuzumab, tacatuzumab tetraxetan, tadocizumab, talizumab, tefibazumab, tocilizumab, toralizumab, tucotuzumab celmoleukin, tucusituzumab, umavizumab, urtoxazumab, ustekinumab, visilizumab, and the anti-interleukin-12 (ABT-8744695, Wyeth Research and Abbott Laboratories) which is a recombinant exclusively human-sequence, full-length IgG.sub.1 .lamda. antibody genetically modified to recognize interleukin-12 p40 protein.

[000368] Chemotherapeutic agents also include “tyrosine kinase inhibitors” including the EGFR inhibitors; small molecule HER2 tyrosine kinase inhibitor such as Mubritonib (TAK165, Takeda); CP-724.714, (Axon Medchem BV, an oral selective inhibitor of the ErbB2 receptor tyrosine kinase); dual-HER inhibitors such as EKB-569 (available from Wyeth) which preferentially binds EGFR but inhibits both HER2 and EGFR-overexpressing cells; lapatinib (GSK572016; available from Glaxo-SmithKline), an oral HER2 and EGFR tyrosine kinase inhibitor; PKI-166 (available from Novartis); pan-HER inhibitors such as canertinib (CI-1033; Pharmacia); Raf-1 inhibitors such as antisense agent ISIS-5132 available from ISIS Pharmaceuticals which inhibit Raf-1 signaling; non-HER targeted TK inhibitors such as imatinib mesylate (GLEEVEC®, available from Glaxo SmithKline); multi-targeted tyrosine kinase inhibitors such as sunitinib (SUTENT®, available from Pfizer); VEGF receptor tyrosine

kinase inhibitors such as vatalanib (PTK787/ZK222584, available from Novartis/Schering AG); MAPK extracellular regulated kinase 1 inhibitor CI-1040 (available from Pharmacia); quinazolines, such as PD 153035, 4-(3-chloroanilino) quinazoline; pyridopyrimidines; pyrimidopyrimidines; pyrrolopyrimidines, such as CGP 59326, CGP 60261 and CGP 62706; pyrazolopyrimidines, 4-(phenylamino)-7H-pyrrolo[2,3-d] pyrimidines; curcumin (diferuloyl methane, 4,5-bis (4-fluoroanilino)phthalimide); tyrphostines containing nitrothiophene moieties; antisense molecules (e.g. those that bind to HER-encoding nucleic acid); quinoxalines (U.S. Pat. No. 5,804,396); tryphostins (U.S. Pat. No. 5,804,396); Affinitac (ISIS 3521; Isis/Lilly); PKI166 (Novartis); Semaxinib (Pfizer); INC-1C11 (Imclone), rapamycin (sirolimus, RAPAMUNE®); or as described in any of the following patent publications: U.S. Pat. No. 5,804,396; WO 1999/09016 (American Cyanamid); WO 1998/43960 (American Cyanamid); WO 1997/38983 (Warner Lambert); WO 1999/06378 (Warner Lambert); WO 1999/06396 (Warner Lambert); WO 1996/30347 (Pfizer, Inc); WO 1996/33978 (Zeneca); WO 1996/3397 (Zeneca) and WO 1996/33980 (Zeneca). Tyrosine kinase inhibitors also include Erlotinib (Tarceva®), Gefitinib (Iressa®), Dasatinib (Sprycel®), Nilotinib (Tasigna®), Crizotinib (Xalkori®), Ruxolitinib (Jakafi®), Vemurafenib (Zelboraf®), Vandetanib (Caprelsa®), Pazopanib (Votrient®), afatinib, alisertib, amuvatinib, axitinib, bosutinib, brivanib, canertinib, cabozantinib, cediranib, crenolanib, dabrafenib, dacomitinib, danusertib, dovitinib, foretinib, ganetespib, ibrutinib, iniparib, lenvatinib, linifanib, linsitinib, masitinib, momelotinib, motesanib, neratinib, niraparib, oprozomib, olaparib, pictilisib, ponatinib, quizartinib, regorafenib, rigosertib, rucaparib, saracatinib, saridegib, tandutinib, tasocitinib, telatinib, tivantinib, tivozanib, tofacitinib, trametinib, veliparib, vismodegib, volasertib, cobimetinib (Cotellic®), and others.

[000369] Chemotherapeutic agents also include dexamethasone, interferons, colchicine, metoprine, cyclosporine, amphotericin, metronidazole, alemtuzumab, alitretinoin, allopurinol, amifostine, arsenic trioxide, asparaginase, BCG live, bevacuzimab, bexarotene, cladribine, clofarabine, darbepoetin alfa, denileukin, dexrazoxane, epoetin alfa, elotinib, filgrastim, histrelin acetate, ibritumomab, interferon alfa-2a, interferon alfa-2b, lenalidomide, levamisole, mesna, methoxsalen, nandrolone, nelarabine, nofetumomab, oprelvekin, palifermin, pamidronate, pegademase, pegaspargase, pegfilgrastim, pemetrexed disodium, plicamycin, porfimer sodium, quinacrine, rasburicase, sargramostim, temozolomide, VM-26, 6-TG,

toremifene, tretinoin, ATRA, valrubicin, zoledronate, and zoledronic acid, and pharmaceutically acceptable salts thereof.

[000370] Chemotherapeutic agents also include hydrocortisone, hydrocortisone acetate, cortisone acetate, tixocortol pivalate, triamcinolone acetonide, triamcinolone alcohol, mometasone, amcinonide, budesonide, desonide, fluocinonide, fluocinolone acetonide, betamethasone, betamethasone sodium phosphate, dexamethasone, dexamethasone sodium phosphate, fluocortolone, hydrocortisone-17-butyrate, hydrocortisone-17-valerate, aclometasone dipropionate, betamethasone valerate, betamethasone dipropionate, prednicarbate, clobetasone-17-butyrate, clobetasol-17-propionate, fluocortolone caproate, fluocortolone pivalate and fluprednidene acetate; immune selective anti-inflammatory peptides (ImSAIDs) such as phenylalanine-glutamine-glycine (FEG) and its D-isomeric form (feG) (IMULAN BioTherapeutics, LLC); anti-rheumatic drugs such as azathioprine, ciclosporin (cyclosporine A), D-penicillamine, gold salts, hydroxychloroquine, leflunomide, minocycline, sulfasalazine, tumor necrosis factor alpha (TNF alpha) blockers such as etanercept (Enbrel), infliximab (Remicade), adalimumab (1-Iumira), certolizumab pegol (Cimzia), golimumab (Simponi), Interleukin 1 (IL-1) blockers such as anakinra (Kineret), T cell costimulation blockers such as abatacept (Orencia), Interleukin 6 (IL-6) blockers such as tocilizumab (ACTEMERA®); Interleukin 13 (IL-13) blockers such as lebrikizumab; Interferon alpha (IFN) blockers such as Rontalizumab; Beta 7 integrin blockers such as rhuMab Beta7; IgE pathway blockers such as Anti-M1 prime; Secreted homotrimeric LTa3 and membrane bound heterotrimer LTa1/132 blockers such as Anti-lymphotoxin alpha (LTa); miscellaneous investigational agents such as thioplatin, PS-341, phenylbutyrate, ET-18-OCH₃, or farnesyl transferase inhibitors (L-739749, L-744832); polyphenols such as quercetin, resveratrol, piceatannol, epigallocatechin gallate, theaflavins, flavanols, procyanidins, betulinic acid and derivatives thereof; autophagy inhibitors such as chloroquine; delta-9-tetrahydrocannabinol (dronabinol, MARINOL®); beta-lapachone; lapachol; colchicines; betulinic acid; acetylcamptothecin, scopoletin, and 9-aminocamptothecin); podophyllotoxin; tegafur (UFTORAL®); bexarotene (TARGRETIN®); bisphosphonates such as clodronate (for example, BONEFOS® or OSTAC®), etidronate (DIDROCAL®), NE-58095, zoledronic acid/zoledronate (ZOMETA®), alendronate (FOSAMAX®), pamidronate (AREDIA®), tiludronate (SKELID®), or risedronate (ACTONEL®); and epidermal growth factor receptor (EGF-R); vaccines such as

THERATOPE® vaccine; perifosine, COX-2 inhibitor (e.g. celecoxib or etoricoxib), proteasome inhibitor (e.g. PS341); CCI-779; tipifarnib (R11577); orafenib, ABT510; Bcl-2 inhibitor such as oblimersen sodium (GENASENSE) pixantrone; farnesyltransferase inhibitors such as lonafamib (SCH 6636, SARASAR™); and pharmaceutically acceptable salts, acids or derivatives of any of the above; as well as combinations of two or more of the above such as CHOP, an abbreviation for a combined therapy of cyclophosphamide, doxorubicin, vincristine, and prednisolone; and FOLFOX, an abbreviation for a treatment regimen with oxaliplatin (ELOXATIN™) combined with 5-FU and leucovorin.

[000371] Chemotherapeutic agents also include Poly ADP ribose polymerase (PARP) inhibitors: olaparib (Lynparza®), rucaprib (Rubraca®) niraparib (Zejula®), talzoparib (Talzenna®).

[000372] Effective combinations of compounds of Formula I' or any formulas as described herein with other agents may be identified through preclinical and clinical testing of the combinations, and will depend on many factors, including disease type and stage of development, overall health of the patient, toxicities and side effects of the agents, and the like.

[000373] In some embodiments, compounds as disclosed herein may be used in combination therapy with any of the kinase inhibitors disclosed herein for the treatment of diseases such as cancer. Exemplary kinase inhibitors include imatinib, baricitinib, gefitinib, erlotinib, sorafenib, dasatinib, sunitinib, lapatinib, nilotinib, pirfenidone, pazopanib, crizotinib, vemurafenib, vandetanib, ruxolitinib, axitinib, bosutinib, regorafenib, tofacitinib, cabozantinib, ponatinib, trametinib, dabrafenib, afatinib, ibrutinib, ceritinib, idelalisib, nintedanib, palbociclib, lenvatinib, cobimetinib, XL-147, XL-765, XL-499, and XL-880. In some embodiments, a compound as described herein can be used in combination with a HSP90 inhibitor (e.g., XL888), liver X receptor (LXR) modulators, retinoid-related orphan receptor gamma (RORγ) *modulators*, a CK1 inhibitor, a CK1-α inhibitor, a Wnt pathway inhibitor (e.g., SST-215), or a mineralocorticoid receptor inhibitor, (e.g., esaxerenone or XL-550) for the treatment of a disease disclosed herein such as cancer.

[000374] In some embodiments, for treatment of cancer, compounds as disclosed herein may be used in combination with inhibitors of PD-1 or inhibitors of PD-L1, e.g., an anti-PD-1 monoclonal antibody or an anti-PD-L1 monoclonal antibody, for example, nivolumab (Opdivo), pembrolizumab (Keytruda, MK-3475), atezolizumab, avelumab, AMP-224, AMP-514,

PDR001, durvalumab, pidilizumab (CT-011), CK-301, BMS 936559, and MPDL3280A; CTLA-4 inhibitors, e.g., an anti-CTLA-4 antibody, for example, ipilimumab (Yervoy) and tremelimumab; and phosphatidylserine inhibitors, for example, bavituximab (PGN401); antibodies to cytokines (IL-10, TGF- β , and the like.); other anti-cancer agents such as cemiplimab.

[000375] In some embodiments, a compound as described herein can be used in combination with a vaccination protocol for the treatment of cancer. In some embodiments, a compound as described herein can be used in combination with vaccines, to stimulate the immune response to pathogens, toxins, and self antigens. Examples of pathogens for which this therapeutic approach may be particularly useful, include pathogens for which there is currently no effective vaccine, or pathogens for which conventional vaccines are less than completely effective. These include, but are not limited to, HIV, Hepatitis (A, B, & C), Influenza, Herpes, Giardia, Malaria, Leishmania, Staphylococcus aureus, Pseudomonas Aeruginosa.

[000376] In some embodiments, compounds as disclosed herein may be used in combination with inhibitors of PARP, for example, olaparib (Lynparza®), rucaprib (Rubraca®) niraparib (Zejula®), talzoparib (Talzenna®).

[000377] The amount of both the compound disclosed herein or salt thereof and the additional one or more additional therapeutic agent (in those compositions which comprise an additional therapeutic agent as described above) that may be combined with carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. In certain embodiments, compositions of this invention are formulated such that a dosage of between 0.01-100 mg/kg body weight/day of an inventive can be administered.

[000378] The additional therapeutic agent and the compound disclosed herein may act synergistically. Therefore, the amount of additional therapeutic agent in such compositions may be less than that required in a monotherapy utilizing only that therapeutic agent, or there may be fewer side effects for the patient given that a lower dose is used. In certain embodiments, in such compositions a dosage of between 0.01-10,000 μ g/kg body weight/day of the additional therapeutic agent can be administered.

[000379] Labeled Compounds and Assay Methods

[000380] Another aspect of the present invention relates to labeled compounds of the invention (radio-labeled, fluorescent-labeled, and the like.) that would be useful not only in

imaging techniques but also in assays, both in vitro and in vivo, for localizing and quantitating TAM kinases in tissue samples, including human, and for identifying TAM kinase ligands by inhibition binding of a labeled compound. Accordingly, the present invention includes TAM kinase assays that contain such labeled compounds.

[000381] The present invention further includes isotopically-labeled compounds of the invention. An “isotopically” or “radio-labeled” compound is a compound of the invention where one or more atoms are replaced or substituted by an atom having an atomic mass or mass number different from the atomic mass or mass number typically found in nature (i.e., naturally occurring). Suitable radionuclides that may be incorporated in compounds of the present invention include but are not limited to ^2H (also written as D for deuterium), ^3H (also written as T for tritium), ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{18}F , ^{35}S , ^{36}Cl , ^{82}Br , ^{75}Br , ^{76}Br , ^{77}Br , ^{123}I , ^{124}I , ^{125}I , and ^{131}I . The radionuclide that is incorporated in the instant radio-labeled compounds will depend on the specific application of that radio-labeled compound. For example, for in vitro metalloprotease labeling and competition assays, compounds that incorporate ^3H , ^{14}C , ^{82}Br , ^{125}I , ^{131}I , or ^{35}S will generally be most useful. For radio-imaging applications ^{11}C , ^{18}F , ^{125}I , ^{123}I , ^{124}I , ^{131}I , ^{75}Br , ^{76}Br , or ^{77}Br will generally be most useful.

[000382] It is understood that a “radio-labeled” or “labeled compound” is a compound that has incorporated at least one radionuclide. In some embodiments, the radionuclide is selected from the group consisting of ^3H , ^{14}C , ^{125}I , ^{35}S , and ^{82}Br .

[000383] The present invention can further include synthetic methods for incorporating radio-isotopes into compounds of the invention. Synthetic methods for incorporating radio-isotopes into organic compounds are well known in the art, and a person of ordinary skill in the art will readily recognize the methods applicable for the compounds of invention.

[000384] A labeled compound of the invention can be used in a screening assay to identify/evaluate compounds. For example, a newly synthesized or identified compound (i.e., test compound) which is labeled can be evaluated for its ability to bind a TAM by monitoring its concentration variation when contacting with the TAM kinases, through tracking of the labeling. For example, a test compound (labeled) can be evaluated for its ability to reduce binding of another compound which is known to bind to a TAM kinase (i.e., standard compound). Accordingly, the ability of a test compound to compete with the standard compound for binding to the TAM kinase directly correlates to its binding affinity. Conversely,

in some other screening assays, the standard compound is labeled, and test compounds are unlabeled. Accordingly, the concentration of the labeled standard compound is monitored in order to evaluate the competition between the standard compound and the test compound, and the relative binding affinity of the test compound is thus ascertained.

Synthesis

[000385] Compounds of this invention can be made by the synthetic procedures described below. The starting materials and reagents used in preparing these compounds are either available from commercial suppliers such as Sigma Aldrich Chemical Co. (Milwaukee, Wis.), or Bachem (Torrance, Calif.), or are prepared by methods known to those skilled in the art following procedures set forth in references such as Fieser and Fieser's Reagents for Organic Synthesis, Volumes 1-17 (John Wiley and Sons, 1991); Rodd's Chemistry of Carbon Compounds, Volumes 1-5 and Supplementals (Elsevier Science Publishers, 1989); Organic Reactions, Volumes 1-40 (John Wiley and Sons, 1991); March's Advanced Organic Chemistry, (John Wiley and Sons, 4th Edition); and Larock's Comprehensive Organic Transformations (VCH Publishers Inc., 1989). These schemes are merely illustrative of some methods by which the compounds of this invention can be synthesized, and various modifications to these schemes can be made and will be suggested to one skilled in the art having referred to this disclosure. The starting materials and the intermediates of the reaction may be isolated and purified if desired using conventional techniques, including but not limited to filtration, distillation, crystallization, chromatography, and the like. Such materials may be characterized using conventional means, including physical constants and spectral data.

[000386] Unless specified to the contrary, the reactions described herein take place at atmospheric pressure and over a temperature range from about -78 °C to about 150 °C, more preferably from about 0 °C to about 125 °C, and most preferably at about room (or ambient) temperature, e.g., about 20 °C. Unless otherwise stated (as in the case of a hydrogenation), all reactions are performed under an atmosphere of nitrogen.

[000387] The compounds disclosed and claimed herein have asymmetric carbon atoms or quaternized nitrogen atoms in their structure and may be prepared through the syntheses described herein as single stereoisomers, racemates, or mixtures of enantiomers and diastereomers. The compounds may also exist as geometric isomers. All such single

stereoisomers, racemates, and geometric isomers, and mixtures thereof are intended to be within the scope of this invention.

[000388] Some of the compounds of the invention may exist as tautomers. For example, where a ketone or aldehyde is present, the molecule may exist in the enol form; where an amide is present, the molecule may exist as the imidic acid; and where an enamine is present, the molecule may exist as an imine. All such tautomers are within the scope of the invention.

[000389] Methods for the preparation and/or separation and isolation of single stereoisomers from racemic mixtures or non-racemic mixtures of stereoisomers are well known in the art. For example, optically active (R)- and (S)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. Enantiomers (R- and S-isomers) may be resolved by methods known to one of ordinary skill in the art, for example by: formation of diastereomeric salts or complexes which may be separated, for example, by crystallization; via formation of diastereomeric derivatives which may be separated, for example, by crystallization; selective reaction of one enantiomer with an enantiomer-specific reagent, for example enzymatic oxidation or reduction, followed by separation of the modified and unmodified enantiomers; or gas-liquid or liquid chromatography in a chiral environment, for example on a chiral support, such as silica with a bound chiral ligand or in the presence of a chiral solvent. It will be appreciated that where a desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a further step may be required to liberate the desired enantiomeric form. Alternatively, specific enantiomers may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts, or solvents, or by converting one enantiomer to the other by asymmetric transformation. For a mixture of enantiomers, enriched in a particular enantiomer, the major component enantiomer may be further enriched (with concomitant loss in yield) by recrystallization.

[000390] In addition, the compounds of the present invention can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the present invention.


[000391] The methods of the present invention may be carried out as semi-continuous or continuous processes, more preferably as continuous processes.

[000392] The present invention as described above unless indicated otherwise may be carried out in the presence of a solvent or a mixture of two or more solvents. In particular the solvent is an aqueous or an organic solvent such as the ether-like solvent (e.g. tetrahydrofuran, methyltetrahydrofuran, diisopropyl ether, t-butylmethyl ether, or dibutyl ether), aliphatic hydrocarbon solvent (e.g. hexane, heptane, or pentane), saturated alicyclic hydrocarbon solvent (e.g. cyclohexane or cyclopentane), or aromatic solvent (e.g. toluene, o-, m-, or p-xylene, or t-butyl-benzene), or mixture thereof.

[000393] The starting materials and reagents, which do not have their synthetic route explicitly disclosed herein, are generally available from commercial sources or are readily prepared using methods well known to the person skilled in the art.

Processes


[000394] One aspect provides a process of making a compound of Formula I:



I

or a pharmaceutically acceptable salt thereof, comprising:


reacting a compound of formula II:



II

wherein Z is selected from the group consisting of NH₂, SH, and OH;

with a compound of formula III:



III

wherein X is a leaving group;

R₁ is selected from the group consisting of -H, -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₁-C₆) alkyl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C₁-C₆) alkyl;



wherein when R₁ is selected from the group consisting of -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C₁-C₆) alkyl, R₂ is -H, halo, -NR₅R₆, or optionally substituted (C₁-C₆) alkoxy;

wherein when R₁ is -H, optionally substituted (C₁-C₆) alkyl, or optionally substituted (C₁-C₆) alkoxy, R₂ is -CO-NR₅R₆, or -CO₂R₇;

or R₁ and R₂ taken together with the atoms to which they are attached to form optionally substituted cycloalkyl or optionally substituted heterocycloalkyl;

R₃ is selected from the group consisting of -H, optionally substituted (C₁-C₆) alkyl, -CN, and halo;

R₄ is -H or halo;

 is optionally substituted with one, two, three, or four groups independently selected from the group consisting of halo; and (C₁-C₆) alkyl, wherein “” indicate points of attachment;

R₅ and R₆ are each independently -H, optionally substituted (C₁-C₆) alkyl, or optionally substituted (C₁-C₆) alkoxy;


R₇ is -H or optionally substituted (C₁-C₆) alkyl

R₈ and R₉ are each independently -H or optionally substituted (C₁-C₆) alkyl; or

R₈ and R₉ may connect to form optionally substituted heterocycle; and

Y is selected from the group consisting of O, S, SO, SO₂, NH, and N-((C₁-C₆) alkyl).


[000395] Another aspect is a process for making a compound of Formula I:



I


or a pharmaceutically acceptable salt thereof, comprising:

reacting a compound of formula IV:



IV

with a compound of formula V:



V

R_1 is selected from the group consisting of -H, -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₁-C₆) alkyl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and (SO₂)-(C₁-C₆) alkyl;



wherein when R_1 is selected from the group consisting of -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C₁-C₆) alkyl, R_2 is -H, halo, -NR₅R₆, or optionally substituted (C₁-C₆) alkoxy;

wherein when R_1 is -H, optionally substituted (C₁-C₆) alkyl, or optionally substituted (C₁-C₆) alkoxy, R_2 is -CO-NR₅R₆, optionally substituted heteroaryl, or -CO₂R₇;

or R_1 and R_2 taken together with the atoms to which they are attached to form optionally substituted cycloalkyl or optionally substituted heterocycloalkyl;

R₃ is selected from the group consisting of -H, optionally substituted (C₁-C₆) alkyl, -CN, and halo;

R₄ is -H or halo;

 is optionally substituted with one, two, three, or four groups independently selected from the group consisting of halo, and C₁-C₆ alkyl, wherein “” indicate points of attachment;

R₅ and R₆ are each independently -H, optionally substituted (C₁-C₆) alkyl, or optionally substituted (C₁-C₆) alkoxy;


R₇ is -H or optionally substituted (C₁-C₆) alkyl;

R₈ and R₉ are each independently -H or optionally substituted (C₁-C₆) alkyl; or

R₈ and R₉ may connect to form optionally substituted heterocycle; and


Y is selected from the group consisting of O, S, SO, SO₂, NH, and N-(C₁-C₆ alkyl).

[000396] In one embodiment, compounds of Formula V are made by a process comprising reacting a compound of Formula VI:




VI

with a compound of Formula VII:



VII

to form a compound of Formula VIII:



VIII

and reducing the compound of Formula VIII to provide a compound of Formula V wherein:

R₁ is selected from the group consisting of -H, -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₁-C₆) alkyl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C₁-C₆) alkyl;

wherein when R₁ is selected from the group consisting of -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C₁-C₆) alkyl, R₂ is -H, halo, -NR₅R₆, or optionally substituted (C₁-C₆) alkoxy;

wherein when R₁ is -H, optionally substituted (C₁-C₆) alkyl, or optionally substituted (C₁-C₆) alkoxy, R₂ is -CO-NR₅R₆, or -CO₂R₇;

or R₁ and R₂ taken together with the atoms to which they are attached to form optionally substituted cycloalkyl or optionally substituted heterocycloalkyl;


R₃ is selected from the group consisting of -H, optionally substituted (C₁-C₆) alkyl, -CN, and halo; and

W is halo.

[000397] The following examples are provided for the purpose of further illustration and are not intended to limit the scope of the claimed invention.

Examples

[000398] **Example 1: Methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylate (5)**



[000399] N-(4-Fluorophenyl)-N-(4-hydroxyphenyl)cyclopropane-1,1-dicarboxamide


(3): To a solution of Compound 1 (10 g, 44.80 mmol, 1 eq) and Compound 2 (5.87 g, 53.8 mmol, 1.2 eq) in dimethyl acetamide (DMA) (60 mL) was added 3-(ethyliminomethyleneamino)-N,N-dimethyl-propan-1-amine hydrochloride (EDCI) (10.31 g, 53.8 mmol, 1.2 eq). The mixture was stirred vigorously at 20 °C until the reaction was complete. The mixture was poured into aqueous (aq) saturated NaHCO₃ (400 mL) and extracted with EtOAc (4 x 100 mL). The combined organic phases were washed with aq saturated NaCl (100 mL), dried over anhyd (anhyd) Na₂SO₄, and concentrated. Compound 3 (21 g, crude) (50% purity) was obtained. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.16 (br s, 1H), 9.72 (br s, 1H), 7.61 (dd, 2H), 7.34 (d, 2H), 7.13 (t, 2H) 6.68 (d, 2H), 1.42 (s, 4H); MS (EI) for C₁₇H₁₅FN₂O₃, found 314.9 (MH⁺).

[000400] Methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-

carbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylate (5): A mixture of Compound 3 (5.99 g, 9.5 mmol, 1.2 eq), Compound 4 (2 g, 8.0 mmol, 1.0 eq), Pd(OAc)₂ (89 mg, 397.4 μmol, 0.05 eq), *rac*-2-(Di-*tert*-butylphosphino)-1,1'-binaphthyl (TrixiePhos, 316.71 mg, 794.7 μmol, 0.1 eq), and K₃PO₄ (2.53 g, 11.9 mmol, 1.5 eq) in anisole (50 mL) was stirred at 110 °C for 2 hours (h) under an atmosphere of nitrogen. The mixture was filtered, and the filtrate was concentrated. The residue was purified by flash silica gel chromatography (1:1 petroleum ether:EtOAc to 20:1 EtOAc:MeOH). Compound 5 was obtained (2.6 g, 61.8% yield). ¹H NMR (400 MHz, CDCl₃) δ 9.38 (s, 1H), 8.80 (s, 1H), 8.63 (d, 2H), 7.64 (d, 2H), 7.54-7.41 (m, 3H),

7.18 (d, 2H), 7.09-7.01 (m, 2H), 6.43 (d, 1H), 4.05 (s, 3H), 3.97 (s, 3H), 1.78-1.72 (m, 2H), 1.69-1.63 (m, 2H); MS (EI) for $C_{29}H_{24}FN_3O_6$, found 530.0 (MH⁺).


[000401] Example 2: 4-[4-[[1-[(4-Fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylic acid (6)



[000402] 4-[4-[[1-[(4-Fluorophenyl)carbamoyl]cyclopropane-

carbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylic acid (6): To a solution of Compound **5** (1.8 g, 3.4 mmol, 1 eq) in tetrahydrofuran (THF) (15 mL) and MeOH (15 mL) was added 2 M aq NaOH (7 mL, 4.1 eq). The mixture was stirred at 6-13 °C for 4 h. The mixture was adjusted to a pH of approximately 8 with 1 M aq HCl and concentrated to remove solvent. Water (50 mL) was added, and the mixture was adjusted to a pH of approximately 6 with 1 M aq HCl. The resulting precipitate was filtered, washed with water (2 x 10 mL), and dried under vacuum. Compound **6** was obtained (1.7 g, 97.0% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.22 (s, 1H), 10.08 (s, 1H), 8.65 (d, 1H), 8.48 (s, 1H), 7.77 (d, 2H), 7.64 (dd, 2H) 7.47 (s, 1H), 7.25 (d, 2H), 7.15 (t, 2H), 6.45 (d, 1H), 3.96 (s, 3H), 1.47 (s, 4H); MS (EI) for $C_{28}H_{22}FN_3O_6$, found 516.1 (MH⁺).

[000403] Example 3: 1-N-[4-(6-Carbamoyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (7)




[000404] 1-N-[4-(6-Carbamoyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-

fluorophenyl)cyclopropane-1,1-dicarboxamide (7): A solution of Compound **6** (350 mg, 679.0 μmol, 1 eq), 1-[bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxid hexafluorophosphate (HATU) (388 mg, 1.0 mmol, 1.5 eq), and diisopropylethylamine (DIEA or DIPEA) (352 mg, 2.7 mmol, 474 μL, 4.0 eq) in dimethylformamide (DMF) (10 mL) was stirred at 6-10 °C for 1 h, after which was added NH₄Cl (73 mg, 1.4 mmol, 2.0 eq), and the mixture

was stirred at 6-10 °C for an additional 17 h. The mixture was filtered, and the resulting filtrate was concentrated and purified by prep HPLC (Column: Waters Xbridge 150mm*25mm*5μm, gradient: 32-62% of acetonitrile in 10 mM aq NH₄HCO₃, flow rate: 25mL/min). Compound 7 was obtained (90.1 mg, 25.8% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.20 (s, 1H), 10.06 (s, 1H), 8.68 (s, 1H), 8.66 (d, 1H), 7.86 (br s, 1H), 7.81-7.72 (m, 3H), 7.68-7.61 (m, 2H), 7.51 (s, 1H), 7.26 (d, 2H), 7.19-7.11 (m, 2H), 6.46 (d, 1H), 4.03 (s, 3H), 1.47 (s, 4H); MS (EI) for C₂₈H₂₃FN₄O₅, found 515.1 (MH⁺).

[000405] Example 4: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (8)



[000406] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (8): A solution of Compound 6 (300 mg, 582.0 μmol, 1 eq), HATU (332 mg, 873.2 μmol, 1.5 eq), and DIEA (301 mg, 2.3 mmol, 406 μL, 4 eq) in DMF (10 mL) was stirred at 6-10 °C for 1 h. Methanamine hydrochloride (79 mg, 1.2 mmol, 2.0 eq) was added, and the mixture was stirred at 6-10 °C for 17 h. The mixture was filtered, and the resulting filtrate was purified by prep HPLC (Column: Waters Xbridge 150mm*25mm*5μm, gradient: 33-63% of acetonitrile in 10 mM aq NH₄HCO₃, flow rate: 25mL/min). Compound 8 was obtained (105.4 mg, 34.3% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.20 (s, 1H), 10.06 (s, 1H), 8.65 (d, 1H), 8.61 (s, 1H), 8.42-8.33 (m, 1H), 7.77 (d, 2H), 7.68-7.61 (m, 2H), 7.51 (s, 1H), 7.25 (d, 2H), 7.19-7.11 (m, 2H), 6.46 (d, 1H), 4.02 (s, 3H), 2.84 (d, 3H), 1.47 (s, 4H); MS (EI) for C₂₉H₂₅FN₄O₅, found 529.1 (MH⁺).

[000407] The following compounds were prepared in a method analogous to Compound 8 in Example 4:

[000408] 1-N-[4-[6-(Ethylcarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (9): ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.20 (s, 1H), 10.06 (s, 1H), 8.65 (d, 1H), 8.56 (s, 1H), 8.40 (br t, 1H), 7.78 (br d, 2H), 7.64 (dd, 2H),

7.51 (s, 1H), 7.25 (d, 2H), 7.15 (t, 2H), 6.46 (d, 1H), 4.02 (s, 3H), 3.37-3.29 (m, 2H), 1.48 (s, 4H), 1.15 (t, 3H).; MS (EI) for $C_{30}H_{27}FN_4O_5$, found 543.2 (MH⁺).

[000409] 1-N-[4-[6-[2-(Dimethylamino)ethylcarbamoyl]-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (10): ¹H NMR (400 MHz, CDCl₃) δ 9.38 (s, 1H), 9.23 (s, 1H), 9.16 (s, 1H), 8.61 (d, 1H), 8.50 (s, 1H), 7.68 (d, 2H), 7.49-7.46 (m, 3H), 7.12 (d, 2H), 7.03 (t, 2H), 6.43 (d, 1H), 4.11 (s, 3H), 3.66 (q, 2H), 2.68 (t, 2H), 2.41 (s, 6H), 1.70 (s, 4H); MS (EI) for $C_{32}H_{32}FN_5O_5$, found 586.2 (MH⁺).

[000410] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(2-piperidin-1-ylethylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (11): ¹H NMR (400 MHz, CDCl₃) δ 9.79 (s, 1H), 9.63 (s, 1H), 9.13 (s, 1H), 8.68 (s, 1H), 8.58 (d, 1H), 7.74 (d, 2H), 7.49-7.41 (m, 3H), 7.06-6.96 (m, 4H), 6.38 (d, 1H), 4.13 (s, 3H), 3.65 (q, 2H), 2.60 (t, 2H), 2.49 (s, 4H), 1.70-1.65 (m, 8H), 1.51 (s, 2H); MS (EI) for $C_{35}H_{36}FN_5O_5$, found 626.3 (MH⁺).

[000411] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(2-morpholin-4-ylethylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (12): ¹H NMR (400 MHz, CDCl₃) δ 9.26 (s, 1H), 9.24 (s, 1H), 8.95 (s, 1H), 8.63 (d, 1H), 8.46 (s, 1H), 7.65 (d, 2H), 7.52 (s, 1H), 7.50-7.47 (m, 2H), 7.16 (d, 2H), 7.05 (t, 2H), 6.46 (d, 1H), 4.15 (s, 3H), 3.79 (t, 4H), 3.68-3.64 (m, 2H), 2.65 (t, 2H), 2.57 (s, 4H), 1.70 (d, 4H); MS (EI) for $C_{34}H_{34}FN_5O_6$, found 628.3 (MH⁺).

[000412] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(oxetan-3-ylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (13): ¹H NMR (400 MHz, CDCl₃) δ 9.34 (s, 1H), 9.22 (s, 1H), 8.92 (s, 1H), 8.64 (d, 1H), 8.47 (d, 1H), 7.65 (d, 2H), 7.55 (s, 1H), 7.50-7.46 (m, 2H), 7.14 (d, 2H), 7.05 (t, 2H), 6.47 (d, 1H), 5.36-5.27 (m, 1H), 5.06 (t, 2H), 4.68 (t, 2H), 4.17 (s, 3H), 1.71 (s, 4H); MS (EI) for $C_{31}H_{27}FN_4O_6$, found 571.2 (MH⁺).

[000413] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-[(1-methylazetidin-3-yl)carbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (14): ¹H NMR (400 MHz, CDCl₃) δ 9.60 (s, 2H), 9.11 (s, 1H), 8.58 (d, 1H), 8.34 (d, 1H), 7.67 (d, 2H), 7.53-7.37 (m, 3H), 7.06 (d, 2H), 6.98 (t, 2H), 6.40 (d, 1H), 4.82-4.66 (m, 1H), 4.12 (s, 3H), 3.73 (t, 2H), 3.10 (t, 2H), 2.38 (s, 3H), 1.74-1.60 (m, 4H); MS (EI) for $C_{32}H_{30}FN_5O_5$, found 584.9 (MH⁺).

[000414] 1-N-[4-[6-(Azetidine-1-carbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (15): ¹H NMR (400 MHz, CDCl₃) δ 9.40 (s,

1H), 9.22 (br s, 1H), 8.59 (d, 1H), 8.32 (s, 1H), 7.63 (d, 2H), 7.49 (td, 9.2 Hz, 3H), 7.13 (d, 2H), 7.04 (t, 2H), 6.43 (d, 1H), 4.28 (t, 2H), 4.08-3.98 (m, 5H), 2.37-2.32 (m, 2H), 1.77-1.64 (m, 4H); MS (EI) for C₃₁H₂₇FN₄O₅, found 554.8 (MH⁺).

[000415] 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidine-1-carbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (16): ¹H NMR (400 MHz, CDCl₃) δ 9.43 (s, 1H), 9.18 (s, 1H), 8.60 (s, 1H), 8.31 (s, 1H), 7.61 (d, 2H), 7.50-7.46 (m, 3H), 7.11 (d, 2H), 7.06-7.00 (m, 2H), 6.45 (d, 1H), 4.75-4.70 (m, 1H), 4.51-4.47 (m, 1H), 4.19-4.13 (m, 1H), 4.11-4.07 (m, 1H), 4.01 (s, 3H), 3.95-3.91 (m, 1H), 1.75-1.73 (m, 2H), 1.72-1.69 (m, 2H); MS (EI) for C₃₁H₂₇FN₄O₆, found 571.0 (MH⁺).


[000416] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(methoxycarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (17): ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.27 (br s, 1H), 10.20 (br s, 1H), 10.06 (br s, 1H), 8.65 (d, 1H), 8.44 (s, 1H), 7.76 (d, 2H), 7.63 (d, 2H), 7.49 (s, 1H), 7.25 (d, 2H), 6.46 (d, 2H), 6.93 (d, 1H), 3.98 (s, 3H), 3.74 (s, 3H), 1.47 (s, 4H); MS (EI) for C₂₉H₂₅FN₄O₆, found 545.1 (MH⁺).

[000417] *t*-Butyl (R)-2-((4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)-7-methoxyquinoline-6-carboxamido)methyl)pyrrolidine-1-carboxylate (18): MS (EI) for C₃₈H₄₀FN₅O₇, found 698.3 (MH⁺).

[000418] *t*-Butyl (S)-2-((4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)-7-methoxyquinoline-6-carboxamido)methyl)pyrrolidine-1-carboxylate (19): MS (EI) for C₃₈H₄₀FN₅O₇, found 698.3 (MH⁺).

[000419] 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(hydroxycarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide hydrochloride (20): ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.08 (s, 1H), 10.32 (s, 1H), 10.04 (s, 1H), 8.97-8.88 (m, 1H), 8.54 (s, 1H), 7.84 (d, 2H), 7.72-7.58 (m, 3H), 7.35 (d, 2H), 7.15 (t, 2H), 6.86-6.75 (m, 1H), 4.04 (s, 3H), 1.48 (d, 4H); MS (EI) for C₂₈H₂₃FN₄O₆, found 531.0 (MH⁺).

[000420] Example 5: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-[(2R)-pyrrolidin-2-yl]methylcarbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (21)




[000421] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-[(2R)-pyrrolidin-2-yl]methylcarbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (21): To a mixture of Compound **18** (67 mg, 96.0 μmol , 1 eq) in dichloromethane (DCM) (3 mL) was added trifluoroacetic acid (TFA; 1 mL) in one portion at 30 °C. The mixture was stirred at 30 °C for 0.5 h, after which the reaction mixture was concentrated under vacuum. The resulting residue was purified by prep HPLC (column: Waters Xbridge 150*25*5 μm ; mobile phase: [water (0.05% ammonia hydroxide v/v)-acetonitrile (ACN)]; B%: 50%) to give Compound **21** (24.4 mg, 42.5% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.52-9.35 (m, 2H), 9.14 (s, 1H), 8.59 (d, 1H), 8.42 (br t, 1H), 7.68 (d, 2H), 7.51-7.44 (m, 3H), 7.09 (d, 2H), 7.06-6.97 (m, 2H), 6.42 (d, 1H), 4.13 (s, 3H), 3.69 (td, 1H), 3.55-3.47 (m, 1H), 3.45-3.37 (m, 1H), 3.01 (t, 2H), 2.01-1.79 (m, 8H), 1.58-1.45 (m, 1H); MS (EI) for $\text{C}_{33}\text{H}_{32}\text{FN}_5\text{O}_5$, found 598.3 (MH^+).

[000422] The following compound was prepared from Compound **19** in a manner analogous to the method used to convert Compound **18** to Compound **21** in Example 5:

[000423] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-[(2S)-pyrrolidin-2-yl]methylcarbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (22): ^1H NMR (400 MHz, CDCl_3) δ 9.47 (s, 2H), 9.15 (s, 1H), 8.60 (d, 1H), 8.40 (t, 1H), 7.7-7.67 (m, 2H), 7.49-7.45 (m, 3H), 7.10-7.04 (m, 2H), 7.03-6.99 (m, 2H), 6.41 (d, 1H), 4.11 (s, 3H), 3.71-3.65 (m, 1H), 3.50-3.37 (m, 2H), 3.02-2.96 (m, 2H), 2.01-1.95 (m, 2H), 1.84-1.75 (m, 2H), 1.71-1.66 (m, 4H), 1.55-1.48 (m, 1H); MS (EI) for $\text{C}_{33}\text{H}_{32}\text{FN}_5\text{O}_5$, found 598.3 (MH^+).

[000424] Example 6: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(oxetan-3-yl)oxycarbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (26)



[000425] 2-(Oxetan-3-yloxy)isoindoline-1,3-dione (24): To a solution of Compound **23** (1.32 g, 8.1 mmol, 1.2 eq) and oxetan-3-ol (500 mg, 6.8 mmol, 1 eq) in THF (30 mL) were added PPh₃ (3.54 g, 13.5 mmol, 2 eq) and diisopropyl azodicarboxylate (DIAD) (2.81 g, 13.9 mmol, 2.7 mL, 2.1 eq) in portions at 0 °C. The resulting mixture was stirred at 20 °C for 15 h to give a brown solution. The mixture was concentrated with silica gel and purified by flash silica gel chromatography (0 to approximately 70% EtOAc in Petroleum ether gradient). The resulting residue was subjected to a second purification by flash silica gel chromatography using the same solvent system to obtain Compound **24** (300 mg, 10.14% yield, 50% purity). MS (EI) for C₁₁H₉NO₄, found 219.8 (MH⁺).

[000426] O-(Oxetan-3-yl)hydroxylamine (25): To a solution of Compound **24** (280 mg, 1.3 mmol, 1 eq) in DCM (5 mL) was added NH₂NH₂·H₂O (95.92 mg, 1.9 mmol, 93.13 uL, 1.5 eq) at 0 °C, and the resulting mixture stirred for 1 h to give a white suspension. The mixture was filtered and washed with DCM (5 mL), and the filtrate was concentrated. Crude Compound **25** (150 mg) was used in the next step without further purification.

[000427] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(oxetan-3-yloxy)carbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (26): To a solution of Compound **6** (100 mg, 194.0 μmol, 1 eq) and Compound **25** (20.74 mg, 232.8 μmol, 1.2 eq) in THF (3 mL) were added propylphosphonic anhydride (T₃P) (185.17 mg, 582.0 μmol, 173.06 uL, 3 eq) and DIEA (75.22 mg, 582.0 μmol, 101.37 uL, 3 eq), and the resulting mixture was stirred at 20 °C for 10 h. The mixture was concentrated, and the resulting residue was purified by prep HPLC (column: YMC-Actus Triart C18 150*30mm*5 μm; mobile phase: [water (0.05% ammonia hydroxide v/v)-ACN]; B%: 25%-65%, 10min) to give Compound **26**


(5.3 mg, 4.66% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.66 (s, 1H), 10.21 (br s, 1H), 10.06 (br s, 1H), 8.67 (d, 1H), 8.42 (s, 1H), 7.78 (d, 2H), 7.64 - 7.86 (m, 2H), 7.51 (s, 1H), 7.26 (d, 2H), 7.13 - 7.18 (m, 2H), 6.47 (d, 1H), 5.07 - 5.09 (m, 1H), 4.73 - 4.76 (m, 2H), 4.63 - 4.66 (m, 2H), 3.99 (s, 3H), 1.48 (s, 4H); MS (EI) for $\text{C}_{31}\text{H}_{27}\text{FN}_4\text{O}_7$, found 587.1 (MH⁺).

[000428] The following compounds were prepared from Compound **6** in a manner analogous to the method used to form Compound **26** from Compound **6** in the last step of Example **6**:

[000429] **1-N'-(4-Fluorophenyl)-1-N-[4-[6-(2-hydroxyethoxycarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (27)**: ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.47 (s, 1H), 10.22 (s, 1H), 10.07 (s, 1H), 8.67 (d, 1H), 8.46 (s, 1H), 7.78 (d, 2H), 7.61 - 7.67 (m, 2H), 7.51 (s, 1H), 7.26 (d, 2H), 7.12 - 7.18 (m, 2H), 6.47 (d, 1H), 4.82 (t, 1H), 4.00 (s, 3H), 3.97 (t, 2H), 3.65 (q, 2H), 1.48 (s, 4H); MS (EI) for $\text{C}_{30}\text{H}_{27}\text{FN}_4\text{O}_7$, found 575.1 (MH⁺).

[000430] **N-(4-(((6-(((2,2-Dimethyl-1,3-dioxolan-4-yl)methoxy)carbamoyl)-7-methoxyquinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (28)**. MS (EI) for $\text{C}_{34}\text{H}_{33}\text{FN}_4\text{O}_8$, found 645.0 (MH⁺).

[000431] **Example 7: 1-N-[4-[6-(2,3-Dihydroxypropoxycarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide, enantiomers 1 (30) and 2 (31)**




[000432] **1-N-[4-[6-(2,3-Dihydroxypropoxycarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide, racemate (29)**: To a solution of Compound **28** (300 mg, 465.37 μmol , 1 eq) in DCM (10 mL) was added TFA (53.06 mg, 465.37 μmol , 34.46 μL , 1 eq), and the resulting mixture was stirred at 20 °C for 10 h. Two

additional aliquots of TFA totaling 0.35 mL were added, and stirring continued at 20 °C for a total of another 5 h to give a brown solution. The reaction mixture was concentrated in vacuo, and the resulting residue was purified by prep-HPLC (column: Agela DuraShell 150mm*25mm*5μm; mobile phase: [water(0.225%FA)-ACN]; B%: 11%-51%, 10min) to give compound **29** (85 mg, 30.21%) ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.54 (br s, 1H), 10.22 (br s, 1H), 10.07 (br s, 1H), 8.67 (d, 1H), 8.49 (s, 1H), 7.77 (d, 2H), 7.64-7.67 (m, 2H), 7.52 (s, 1H), 7.26 (d, Hz, 2H), 7.14-7.16 (m, 2H), 6.47 (d, 1H), 5.00 (br s, 1H), 4.67 (br s, 1H), 3.99-4.02 (m, 4H), 3.78-3.85 (m, 2H), 3.39-3.42 (m, 2H), 1.48 (s, 4H); MS (EI) for C₃₁H₂₉FN₄O₈, found 605.4 (MH⁺).

[000433] 1-N-[4-[6-(2,3-Dihydroxypropoxycarbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide, enantiomers 1 (30) and 2 (31).


The individual enantiomers of the racemic Compound **29** (85 mg, 140.59 μmol, 1 eq) were separated using SFC (column: DAICEL CHIRALPAK AD(250mm*30mm, 10μm); mobile phase: [0.1% NH₃H₂O IPA]; B%: 45%-45%, min) to give Compound **30** (53.0 mg, 62.35%) and Compound **31** (21.3 mg, 25.06% yield). Compound **30**: ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.54 (br s, 1H), 10.22 (br s, 1H), 10.07 (br s, 1H), 8.67 (d, 1H), 8.49 (s, 1H), 7.77 (d, 2H), 7.64-7.67 (m, 2H), 7.52 (s, 1H), 7.26 (d, 2H), 7.14-7.16 (m, 2H), 6.47 (d, 1H), 5.00 (d, 1H), 4.67 (t, 1H), 3.99-4.03 (m, 4H), 3.78-3.93 (m, 2H), 3.39-3.47 (m, 2H), 1.48 (s, 4H); MS (EI) for C₃₁H₂₉FN₄O₈, found 605.3 (MH⁺). Compound **31**: ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.54 (br s, 1H), 10.23 (br s, 1H), 10.08 (br s, 1H), 8.67 (d, 1H), 8.49 (s, 1H), 7.79 (d, 2H), 7.64-7.67 (m, 2H), 7.51 (s, 1H), 7.26 (d, 2H), 7.14-7.18 (m, 2H), 6.47 (d, 1H), 5.03 (br s, 1H), 4.69 (br s, 1H), 4.00-4.04 (m, 4H), 3.76-3.86 (m, 2H), 3.52-3.56 (m, 2H), 1.48 (s, 4H); MS (EI) for C₃₁H₂₉FN₄O₈, found 605.0 (MH⁺).

[000434] Example 8: 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(hydrazinecarbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (32)



[000435] 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(hydrazinecarbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (32): To a solution of Compound **5** (100 mg, 188.85 μmol , 1 eq) in MeOH (5 mL) was added $\text{NH}_2\text{NH}_2\cdot\text{H}_2\text{O}$ (28.36 mg, 566.56 μmol , 27.54 μL , 3eq). The resulting mixture was stirred at 50 °C for 2 h and then concentrated in vacuo. The residue was triturated with MeOH (3 mL), and the resulting residue was filtered to give Compound **32** (56.6 mg, 53.77% yield). ^1H NMR (400 MHz, DMSO- d_6) δ 10.21 (s, 1H), 10.06 (s, 1H), 9.51 (br s, 1H), 8.66 (d, 1H), 8.53 (s, 1H), 7.78 (d, 2H), 7.69 - 7.60 (m, 2H), 7.51 (s, 1H), 7.26 (d, 2H), 7.16 (t, 2H), 6.48 (d, 1H), 4.62 (br d, 2H), 4.01 (s, 3H), 1.48 (s, 4H); MS (EI) for $\text{C}_{28}\text{H}_{24}\text{FN}_5\text{O}_5$, found 530.2 (MH $^+$).

[000436] Example 9: 1-N-[4-(6-Acetyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (34)




[000437] N-(4-Fluorophenyl)-N-(4-((7-methoxy-6-(methoxymethyl)carbamoyl)quinolin-4-yl)oxy)phenyl)cyclopropane-1,1-dicarboxamide (33): To a solution of Compound **6** (300 mg, 581.97 μmol , 1 eq) and N,O-dimethylhydroxylamine hydrochloride (170.30 mg, 1.75 mmol, 3 eq) in DMF (3 mL) were added HATU (442.57 mg, 1.16 mmol, 2 eq) and DIEA (225.65 mg, 1.75 mmol, 304.11 μL , 3 eq). The resulting mixture was stirred at 20 °C for 3 h to give a brown solution. The mixture was diluted with EtOAc (60 mL) and washed with water (2 x 20 mL) and aq saturated NaCl (20 mL). The organic phase was concentrated in vacuo, and the resulting residue was purified by flash silica gel chromatography (ISCO®; 5 g SepaFlash® Silica Flash Column, Eluent of 0 to approximately 20% MeOH/DCM gradient at 10 mL/min) to give Compound **33** (300 mg, 89.52% yield. MS (EI) for $\text{C}_{30}\text{H}_{27}\text{FN}_4\text{O}_6$, found 559.2 (MH $^+$).

[000438] 1-N-[4-(6-Acetyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (34): To a solution of Compound **33** (280 mg, 501.29 μmol , 1 eq) in THF (5 mL) was added MeMgBr (3 M, 1.67 mL, 10 eq) at 0 °C under an atmosphere of nitrogen. The resulting mixture was stirred at 20 °C for 15 h to give a brown suspension. The reaction mixture was quenched with aq saturated NH_4Cl (20 mL), extracted with EtOAc (3 x 15 mL), and concentrated. The resulting residue was purified by flash silica gel

chromatography (ISCO®; 4 g SepaFlash® Silica Flash Column, Eluent of 0 4% MeOH/DCM gradient @ 10 mL/min). The resulting crude product was further purified (ISCO®; 4 g SepaFlash® Silica Flash Column, Eluent of 0 to approximately 60% EtOAc/petroleum ether gradient @ 10 mL/min) to give Compound **34** (190 mg, 73.81% yield). ¹H NMR (400 MHz, CDCl₃) δ 9.40 (br s, 1H), 8.90 (br s, 1H), 8.68 (s, 1H), 7.51 (d, 1H), 7.63-7.66 (m, 2H), 7.45-7.50 (m, 3H), 7.14-7.16 (m, 2H), 7.02-7.07 (m, 2H), 6.43 (d, 1H), 4.05 (s, 3H), 2.71 (s, 3H), 1.16-1.72 (m, 4H); MS (EI) for C₂₉H₂₄FN₃O₅, found 536.1 (MH+Na)⁺.


[000439] Example 10: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-[(E)-N-methoxy-C-methylcarbonimidoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (35) and 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-[(Z)-N-methoxy-C-methylcarbonimidoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (36)



[000440] To a solution of Compound **34** (140 mg, 272.63 μmol, 1 eq) in water (2.5 mL) and EtOH (2.5 mL) was added methoxyamine hydrochloride (45.54 mg, 545.26 μmol, 41.40 uL, 2 eq), and the resulting mixture was stirred at 50 °C for 2 h to give a brown solution. The reaction mixture was diluted with EtOAc (50 mL), washed with water (2 x 20 mL), and concentrated. The resulting residue was purified by flash silica gel chromatography (ISCO®; 4g SepaFlash® Silica Flash Column, Eluent of 0 to approximately 80% EtOAc/Petroleum ether gradient @ 10 mL/min) twice. Compounds **35** and **36** were recovered as a mixture of Z and E isomers (110 mg, 69.7% yield)(MS (EI) for C₃₀H₂₇FN₄O₅, found 542.9 (MH⁺). The Z and E isomers were separated by prep-HPLC (column: Agela DuraShell 150mm*25mm*5μm;mobile phase: [water (0.05% ammonia hydroxide v/v)-ACN];B%: 47%-87%,10min) to give Compound **35** (54.7 mg, 52.10% yield) and Compound **36** (6.4 mg, 6.10% yield). Compound **35**: ¹H NMR (400 MHz,


DMSO-*d*₆) δ 10.21 (br s, 1H), 10.07 (br s, 1H), 8.63 (d, 1H), 8.09 (s, 1H), 7.77 (d, 2H), 7.65-7.68 (m, 2H), 7.49 (s, 1H), 7.77 (d, 2H), 7.16 (t, 2H), 6.44 (d, 1H), 3.99 (s, 3H), 3.91 (s, 3H), 2.16 (s, 3H), 1.48 (s, 4H). Compound **36**: ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.21 (br s, 1H), 10.07 (br s, 1H), 8.61 (d, 1H), 7.97 (s, 1H), 7.77 (d, 2H), 7.63-7.66 (m, 2H), 7.47 (s, 1H), 7.25 (d, 2H), 7.16 (t, 2H), 6.43 (d, 1H), 3.95 (s, 3H), 3.70 (s, 3H), 2.15 (s, 3H), 1.48 (s, 4H).

[000441] Example 11: 1-N-[4-(6-Cyano-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (37)



[000442] 1-N-[4-(6-Cyano-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (37): To Compound **7** (200 mg, 379.12 μ mol, 1 equiv) in MeCN (10 mL) was added POCl₃ (7.78 g, 50.74 mmol, 4.72 mL, 134 equiv). The mixture was stirred at 80 °C for 3 h. The reaction mixture was filtered, and the filtrate was concentrated and dried under vacuum. The residue was diluted with water (5 mL), the pH was adjusted to 8-9 with aq Na₂CO₃, and the residue was extracted with DCM (3 x 20 mL). The combined organic layers were concentrated and then dried under vacuum. The residue was purified by prep HPLC to give Compound **37** (80.3 mg, 42.37% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.23 (br s, 1H), 10.05 (br s, 1H), 8.79-8.72 (m, 2H), 7.79 (br d, 2H), 7.68-7.58 (m, 3H), 7.27 (d, 2H), 7.15 (t, 2H), 6.52 (d, 1H), 4.07 (s, 3H), 1.48 (s, 4H); MS (EI) for C₂₈H₂₁FN₄O₄, found 497.1 (MH⁺).

[000443] Example 12: N-(4-((6-Bromo-7-methoxyquinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (43)



[000444] 5-(((4-Bromo-3-methoxyphenyl)amino)methylene)-2,2-dimethyl-1,3-dioxane-4,6-dione (40): Compound **39** (4.10 g, 28.46 mmol, 1.15 eq) in trimethoxymethane (25 mL, 228.04 mmol, 9.22 eq) was heated to reflux at 105 °C for 1 h. Compound **38** (5 g, 24.75 mmol, 1 eq) was then added, and the reflux continued at 105 °C for another 1 h. The resulting suspension was filtered, washed with MeOH, and vacuum dried to yield Compound **40** (7.7 g, 87.4% yield), which was used in the next step without further purification. ¹H NMR (400 MHz, CDCl₃) δ 11.24-11.20 (d, 1H), 8.62-9.59 (d, 1H), 7.57-7.54 (d, 1H), 6.77-6.74 (m, 2H), 3.94 (s, 3H), 1.76 (s, 3H), 1.74 (s, 3H).

[000445] 6-Bromo-7-methoxyquinolin-4(1H)-one (41): To Ph₂O (35 mL) at 230 °C was added Compound **40** (7.7 g, 21.62 mmol, 1 eq), and the mixture was stirred for 1 h. After cooling to room temperature, the reaction mixture was poured into hexane (20 mL), and the resulting precipitate was filtered and washed with hexane. The resulting residue was dried under vacuum to give Compound **41** (6.2 g, 75.8% yield, 67.2% purity). ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.78 (s, 1H), 8.18 (s, 1H), 7.88-7.86 (d, 1H), 7.05 (s, 1H), 6.03-6.01 (d, 1H), 3.92 (s, 3H); MS (EI) for C₁₀H₈BrNO₂, found 254.2 (MH⁺).

[000446] 6-Bromo-4-chloro-7-methoxyquinoline (42): Compound **41** (6.2 g, 16.40 mmol, 1 eq) in POCl₃ (15 mL, 161.41 mmol, 9.84 eq) was stirred at 110 °C for 1 h. After cooling, the reaction mixture was cautiously poured into a mixture of aq saturated Na₂CO₃ and ice with stirring. The resulting suspension was filtered, washed with water, and dried under vacuum to give Compound **42** (7.78 g, 57.4% purity, 99.9% yield), which was used in subsequent steps without further purification. MS (EI) for C₁₀H₇BrClNO, found 272.2 (MH⁺).


[000447] N-(4-((6-Bromo-7-methoxyquinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (43): A mixture of Compound **42** (5.72 g,

12.05 mmol, 1.26 eq), Compound **3** (3.00 g, 9.54 mmol, 1 eq), and *t*-BuOK (3.21 g, 28.63 mmol, 3 eq) in DMSO (30 mL) was stirred at 150 °C for 2 h. Aq saturated NH₄Cl (200 mL) was added, and the mixture was extracted with EtOAc (3 x 250 mL). The combined organic phases were dried over anhyd Na₂SO₄ and concentrated in vacuo. The resulting residue was purified by flash silica gel chromatography (ISCO®; 40 g SepaFlash® Silica Flash Column, Eluent of 0 to approximately 100% EtOAc/Petroleum ether gradient at 40mL/min). The resulting residue was triturated with MeOH (50 mL) and filtered to give Compound **43** (3.0 g, 57.11% yield). ¹H NMR (400 MHz, CDCl₃) δ 9.33 (s, 1H), 8.66 (br s, 1H), 8.60 (d, 1H), 8.58 (s, 1H), 7.64 (d, 2H), 7.48 (dd, 2H), 7.45 (s, 1H), 7.18 (d, 2H), 7.07 (t, 2H), 6.45 (d, 1H), 4.07 (s, 3H), 1.78-1.72 (m, 2H), 1.71-1.65 (m, 2H).

[000448] The following compound was synthesized in a manner analogous to Compound **43** in the last step of Example 12, substituting Compound **42** with 6-bromo-4-chloroquinoline.


[000449] **N-(4-((6-Bromoquinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (44)**. MS (EI) for C₂₆H₁₉BrFN₃O₃, found 520.0 (MH⁺).

[000450] **Example 13: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (45)**



[000451] **1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (45)**: Compound **43** (200 mg, 363.39 μmol, 1 equiv), tributyl(oxazol-2-yl)stannane (200.0 mg, 558.5 μmol, 1.5 equiv), CuI (8 mg, 42.0 μmol, 0.12 equiv), and Pd(PPh₃)₄ (48.0 mg, 41.5 μmol, 0.11 equiv) were combined in 1,4-dioxane (5 mL) in a sealed tube, which was then heated at 100 °C under microwave irradiation for 30 min. The reaction mixture was concentrated under vacuum, and the resulting residue was purified initially by flash silica gel chromatography (0~100% EtOAc/Petroleum ether) and subsequently by prep HPLC to give Compound **45** (81.5 mg, 41.3% yield). ¹H NMR (400 MHz, CDCl₃) δ 9.36 (s, 1H), 8.98 (s, 1H), 8.75 (br s, 1H), 8.63 (d, 1H), 7.82 (s, 1H), 7.65 (d, 2H), 7.57 (s, 1H), 7.48 (dd, 2H), 7.36 (s, 1H), 7.19 (d, 2H), 7.06 (t, 2H), 6.46 (d, 1H), 4.13 (s, 3H), 1.77-1.74 (m, 2H), 1.73-1.69 (m, 2H); MS (EI) for C₃₀H₂₃FN₄O₅, found 539.1 (MH⁺).

[000452] Example 14: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(2-hydroxyethoxy)-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (50)



[000453] 6-Bromo-4-chloroquinolin-7-ol (46): A mixture of Compound **42** (5 g, 18.35 mmol, 1 eq) and BBr₃ (13.00 g, 51.89 mmol, 5 mL, 2.83 eq) in DCE (15 mL) was stirred under an atmosphere of nitrogen at 90 °C for 2.5 h. The reaction mixture was cooled to room temperature and poured into aq saturated NaHCO₃ with vigorous stirring. The resulting residue was filtered, washed with water, dissolved in methyl tert-butyl ether, and dried. The solvent was removed under reduced pressure to give Compound **46** (7g), which was used in subsequent reactions without further purification. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.03 (d, 1H), 8.49 (s, 1H), 7.93 (d, 1H), 7.69 (s, 1H), 6.08 (s, 1H); MS (EI) for C₉H₅BrClNO, found 260.2 (MH⁺).

[000454] 6-Bromo-7-(2-((tert-butyldimethylsilyl)oxy)ethoxy)-4-chloroquinoline (47): To a mixture of Compound **46** (1 g, 3.87 mmol, 1 eq) and 2-bromoethoxy-tert-butyl-dimethylsilane (1.15 g, 4.8 mmol, 1.24 eq) in DMF (10 mL) was added K₂CO₃ (1.05 g, 7.60 mmol, 1.96 eq) and NaI (720 mg, 4.80 mmol, 1.24 eq), and the mixture was heated to 80 °C with stirring for 2 h. The reaction mixture was partitioned between water (20 mL) and EtOAc (2 x 20 mL). The combined organic phases were washed with aq saturated NaCl (10 mL), dried over anhyd Na₂SO₄, and concentrated. The resulting residue was purified by flash silica gel chromatography (ISCO®; 12g SepaFlash® Silica Flash Column, Eluent of 0 to approximately 20% EtOAc/Petroleum ether gradient at 30 mL/min) to give Compound **47** (560 mg, 34.73% yield). ¹H NMR (400 MHz, CDCl₃) δ 8.71 (d, 1H), 8.45 (s, 1H), 7.47 (s, 1H), 7.37 (d, 1H), 4.29-4.26


(m, 2H), 4.13-4.11 (m, 2H), 0.93-0.92 (m, 9H), 0.16-0.14 (m, 6H); MS (EI) for $C_{17}H_{23}BrClNO_2Si$, found 417.8 (MH⁺).

[000455] N-(4-((6-Bromo-7-(2-((tert-butyldimethylsilyl)oxy)ethoxy)quinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (48): Compound **48** was made in a manner analogous to the preparation of Compound **43** from Compound **42** in Example 12. Compound **48** was recovered (400 mg, 48% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.20 (s, 1H), 10.01 (s, 1H), 8.63 (d, 1H), 8.47 (s, 1H), 7.77 (d, 2H), 7.65-7.62 (m, 2H), 7.55 (s, 1H), 7.25 (d, 2H), 7.17-7.13 (m, 2H), 6.46 (d, 1H), 4.32-4.30 (t, 2H), 4.05-4.03 (t, 2H), 1.47 (s, 4H), 0.88 (s, 9H), 0.10 (s, 6H); MS (EI) for $C_{34}H_{37}BrFN_3O_5Si$, found 696.1 (MH⁺).

[000456] N-(4-((7-(2-((tert-Butyldimethylsilyl)oxy)ethoxy)-6-(oxazol-2-yl)quinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (49): Compound **49** was made in a manner analogous to the preparation of Compound **45** from Compound **43** in Example 13. Compound **49** was recovered (400 mg, crude). ¹H NMR (400 MHz, CDCl₃) δ 10.20 (s, 1H), 10.06 (s, 1H), 8.74 (s, 1H), 8.67 (d, 1H), 8.25 (s, 1H), 7.78 (d, 2H), 7.62 - 7.59 (m, 3H), 7.43 (s, 1H), 7.28 (d, 2H), 7.19 - 7.09 (m, 2H), 6.48 (d, 1H), 4.41 - 4.25 (m, 2H), 4.04 - 4.02 (m, 2H), 1.47 (s, 4H), 0.83 (s, 9H), 0.02 (s, 6H); MS (EI) for $C_{37}H_{39}FN_4O_6Si$, found 683.3 (MH⁺).


[000457] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(2-hydroxyethoxy)-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (50): To a mixture of Compound **49** (350 mg, 512.6 μmol, 1 eq) in THF (20 mL) was added tetrabutyl ammonium fluoride (1 M, 1.40 mL, 2.73 eq), and the resulting reaction mixture was stirred for 0.5 h at 0-20 °C, after which it was poured into aq saturated NaHCO₃ with vigorous stirring. The resulting mixture was partitioned between water (20 mL) and EtOAc (2 x 20 mL). The combined organic phases were separated, washed with aq saturated NaCl (10 mL), dried over anhyd Na₂SO₄, and concentrated. The resulting residue was recrystallized from MeOH to give Compound **50** (174.3 mg, 58.6% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.20 (s, 1H), 10.06 (s, 1H), 8.77 (s, 1H), 8.67 (d, 1H), 8.30 (s, 1H), 7.78 (d, 2H), 7.66-7.63 (m, 3H), 7.45 (s, 1H), 7.28 (d, 2H), 7.17-7.13 (m, 2H), 6.47 (d, 1H), 4.97-4.94 (t, 1H), 4.32-4.30 (t, 2H), 3.86-3.82 (m, 2H), 1.47 (s, 4H); MS (EI) for $C_{31}H_{25}FN_4O_6$, found 569.2 (MH⁺).

[000458] Example 15: 1-N-[4-(6-Dimethylphosphoryl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (51)



[000459] 1-N-[4-(6-Dimethylphosphoryl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (51): To a solution of Compound **43** (100 mg, 181.7 μmol , 1 eq) in dioxane (2 mL) was added methylphosphonoylmethane (21.27 mg, 272.5 μmol , 1.5 eq), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (Xantphos) (21.03 mg, 36.3 μmol , 0.2 eq), tris(dibenzylideneacetone)dipalladium(0) ($\text{Pd}_2(\text{dba})_3$) (33.28 mg, 36.3 μmol , 0.2 eq), and Et_3N (91.93 mg, 908.5 μmol , 126.5 μL , 5 eq). The resulting mixture was stirred at 110 $^\circ\text{C}$ for 1 h under an atmosphere of nitrogen. The mixture was concentrated in vacuo and triturated with EtOH (5 mL). The resulting residue was filtered, dried, and purified by prep-HPLC (YMC-Actus Triart C18 150*30mm*5 μm ; mobile phase: [water(0.225%FA)-ACN]; B%: 20%-60%, 10min) to give Compound **51** (26.2 mg, 26.34% yield, 100% purity). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.24 (s, 1H), 10.10 (s, 1H), 8.76 (d, 1H), 8.69 (d, 1H), 7.80 (d, 2H), 7.65 (dd, 2H), 7.52 (d, 1H), 7.28 (d, 2H), 7.16 (t, 2H), 6.48 (d, 1H), 4.04 (s, 3H), 1.75 (s, 3H), 1.72 (s, 3H), 1.48 (s, 4H); MS (EI) for $\text{C}_{29}\text{H}_{27}\text{FN}_3\text{O}_5\text{P}$, found 548.1 (MH $^+$).

[000460] Example 16: Methyl 4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylate (56)



[000461] Methyl 4-(((2,2-dimethyl-4,6-dioxo-1,3-dioxan-5-ylidene)methyl)amino)benzoate (53): Compound **53** was synthesized from Compounds **52** and **39** in a manner analogous to the preparation of Compound **40** from Compounds **38** and **39** in


Example 12. Compound **53** was obtained (9 g, 89.1% yield). ^1H NMR (400 MHz, CDCl_3) δ 11.32 (br d, 1H), 8.70 (d, 1H), 8.13 (d, 2H), 7.31 (d, 2H), 3.94 (s, 3H), 1.77 (s, 6H).

[000462] Methyl 4-oxo-1,4-dihydroquinoline-6-carboxylate (54): Compound **54** was synthesized from Compound **53** in a manner analogous to the preparation of Compound **41** from Compound **40** in Example 12. Compound **54** was obtained (1.3 g, crude). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 12.03 (s, 1H), 8.74-8.64 (m, 1H), 8.14 (dd, 1H), 8.01-7.93 (m, 1H), 7.64-7.58 (m, 1H), 6.11 (d, 1H), 3.88 (s, 3H).

[000463] Methyl 4-chloroquinoline-6-carboxylate (55): Compound **55** was synthesized from Compound **54** in a manner analogous to the preparation of Compound **42** from Compound **41** in Example 12. Compound **55** was obtained (0.5 g, 35.3% yield). ^1H NMR (400 MHz, CDCl_3) δ 9.00 (d, 1H), 8.89 (d, 1H), 8.38 (dd, 1H), 8.19 (d, 1H), 7.58 (d, 1H), 4.03 (s, 3H); MS (EI) for $\text{C}_{11}\text{H}_8\text{ClNO}_2$, found 221.9 (MH $^+$).


[000464] Methyl 4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylate 56: Compound **56** was synthesized from Compounds **55** and **3** in a manner analogous to the preparation of Compound **43** from Compounds **42** and **3** in Example 12. Compound **56** was obtained (0.9 g, 99.8% yield). MS (EI) for $\text{C}_{28}\text{H}_{22}\text{FN}_3\text{O}_5$, found 500.4 (MH $^+$).

[000465] Example 17: 4-(4-(1-((4-Fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylic acid (57)



[000466] 4-(4-(1-((4-Fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylic acid (57): Compound **57** was synthesized from Compound **56** in a manner analogous to the preparation of Compound **6** from Compound **5** in Example 2. Compound **57** was obtained (0.5 g, 57.2% yield). MS (EI) for $\text{C}_{27}\text{H}_{20}\text{FN}_3\text{O}_5$, found 486.0 (MH $^+$).


[000467] Example 18: 1-N-[4-(6-Carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (58)



[000468] N-(4-((6-Carbamoylquinolin-4-yl)oxy)phenyl)-N-(4-

fluorophenyl)cyclopropane-1,1-dicarboxamide (58): Compound **58** was synthesized from Compound **57** in a manner analogous to the preparation of Compound **7** from Compound **6** in Example 3. Compound **58** was obtained (26 mg, 26.0% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.22 (s, 1H), 10.06 (s, 1H), 8.90 (d, 1H), 8.74 (d, 1H), 8.34 (br s, 1H), 8.26 (dd, 1H), 8.06 (d, 1H), 7.80 (d, 2H), 7.64 (dd, 2H), 7.56 (br s, 1H), 7.29 (d, 2H), 7.15 (t, 2H), 6.61 (d, 1H), 1.48 (s, 4H); MS (EI) for $\text{C}_{27}\text{H}_{21}\text{FN}_4\text{O}_4$, found 485.1 (MH $^+$).

[000469] Example 19: 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (59)



[000470] 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(methylcarbamoyl)quinolin-4-


yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (59): Compound **59** was synthesized from Compound **58** in a manner analogous to the preparation of Compound **8** from Compound **6** in Example 4. Compound **59** was obtained (31.1 mg, 30.3% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.22 (s, 1H), 10.05 (s, 1H), 8.86 (d, 1H), 8.81 (br d, 1H), 8.74 (d, 1H), 8.23 (dd, 1H), 8.07 (d, 1H), 7.80 (d, 2H), 7.64 (dd, 2H), 7.29 (d, 2H), 7.15 (t, 2H), 6.61 (d, 1H), 2.85 (d, 3H), 1.48 (s, 4H); MS (EI) for $\text{C}_{28}\text{H}_{23}\text{FN}_4\text{O}_4$, found 499.1 (MH $^+$).

[000471] The following compound was prepared by a method analogous to the preparation of Compound **59** in Example 19:

[000472] 1-N'-(4-Fluorophenyl)-1-N-[4-[6-[(1-methylazetidin-3-yl)carbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (60): ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.23 (s, 1H), 10.05 (s, 1H), 9.23 - 9.14 (m, 1H), 8.89 (s, 1H), 8.75 (d, 1H), 8.25 (d, 1H), 8.08 (d, 1H), 7.81 (br d, 2H), 7.68-7.62 (m, 2H), 7.30 (d, 2H), 7.16 (t, 2H), 6.61 (d, 1H), 4.60-4.44

(m, 1H), 3.69-3.60 (m, 2H), 3.31 (s, 3H), 3.20-3.09 (m, 2H), 1.48 (s, 4H); MS (EI) for $C_{31}H_{28}FN_5O_4$, found 554.1 (MH⁺).

[000473] Example 20: Methyl 7-fluoro-4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylate (65)



[000474] Methyl 4-(((2,2-dimethyl-4,6-dioxo-1,3-dioxan-5-ylidene)methyl)amino)-2-fluorobenzoate (62): Compound **62** was synthesized from Compounds **61** and **39** in a manner analogous to the preparation of Compound **40** from Compounds **38** and **39** in Example 12. Compound **62** was obtained (4.8 g, 50.2% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.28 (s, 1H), 8.65 (s, 1H), 7.91 (t, 1H), 7.72 (dd, 1H), 7.52 (dd, 1H), 3.84 (s, 3H), 1.68 (s, 6H).


[000475] Methyl 7-fluoro-4-oxo-1,4-dihydroquinoline-6-carboxylate (63): Compound **63** was synthesized from Compound **62** in a manner analogous to the preparation of Compound **41** from Compound **40** in Example 12. Compound **63** was obtained (0.8 g, 58.5% yield). MS (EI) for $C_{11}H_8FNO_3$, found 222.2 (MH⁺).

[000476] Methyl 4-chloro-7-fluoroquinoline-6-carboxylate (64): Compound **64** was synthesized from Compound **63** in a manner analogous to the preparation of Compound **42** from Compound **41** in Example 12. Compound **64** was obtained (110 mg, 31% yield). ¹H NMR (400 MHz, CDCl₃) δ 8.90 (d, 1H), 8.85 (d, 1H), 7.84 (d, 1H), 7.53 (d, 1H), 4.03 (s, 3H); MS (EI) for $C_{11}H_7ClFNO_2$, found 240.0 (MH⁺).

[000477] Methyl 4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylate (65): Compound **65** was synthesized from Compounds **64** and **3** in a manner analogous to the preparation of Compound **43** from


Compounds **42** and **3** in Example 12. Compound **65** was obtained (90 mg, 92.6% yield). MS (EI) for $C_{28}H_{21}F_2N_3O_5$, found 518.3 (MH⁺).

[000478] Example 21: 7-Fluoro-4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylic acid (66)



[000479] 7-Fluoro-4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylic acid (66): Compound **66** was synthesized from Compound **65** in a manner analogous to the preparation of Compound **6** from Compound **5** in Example 2. Compound **66** was obtained (50 mg, 57.1% yield). MS (EI) for $C_{27}H_{19}F_2N_3O_5$, found 504.4 (MH⁺).

[000480] Example 22: 1-N-[4-(6-Carbamoyl-7-fluoroquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (67)



[000481] 1-N-[4-(6-Carbamoyl-7-fluoroquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (67): Compound **67** was synthesized from Compound **66** in a manner analogous to the preparation of Compound **7** from Compound **6** in Example 3. Compound **67** was obtained (7.6 mg, 15.2% yield). ¹H NMR (400 MHz, CDCl₃) δ 9.47 (s, 1H), 9.24 (d, 1H), 8.84 (s, 1H), 8.71 (d, 1H), 7.79 (d, 1H), 7.67 (d, 2H), 7.48 (dd, 2H), 7.15 (d, 2H), 7.04 (t, 2H), 6.86 (d, 1H), 6.54 (d, 1H), 6.06 (s, 1H), 1.76 -1.68 (m, 4H); MS (EI) for $C_{27}H_{20}F_2N_4O_4$, found 503.1 (MH⁺).


[000482] The following compounds were prepared by a sequence analogous to that taken to prepare Compound **67** from Compound **61** in Examples 20-22, substituting Compound **61** with the appropriate methyl 4-amino-2-halobenzoate:

[000483] 1-N-[4-(6-Carbamoyl-7-chloroquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (68): ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.23

(s, 1H), 10.05 (s, 1H), 8.75 (d, 1H), 8.35 (s, 1H), 8.14 (s, 2H), 7.88-7.73 (m, 3H), 7.64 (dd, 2H), 7.27 (d, 2H), 7.15 (t, 2H), 6.64 (d, 1H), 1.47 (s, 4H); MS (EI) for $C_{27}H_{20}ClFN_4O_4$, found 519.4 (MH⁺).

[000484] 1-N-[4-(7-Bromo-6-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (69): ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.24 (br s, 1H), 10.07 (br s, 1H), 8.75 (d, 1H), 8.30 (d, 2H), 8.14 (br s, 1H), 7.80 (d, 2H), 7.64-7.67 (m, 2H), 7.27 (d, 2H), 7.18 (t, 3H), 6.65 (d, 1H), 1.48 (s, 4H); MS (EI) for $C_{27}H_{20}BrFN_4O_4$, found 563.3 (MH⁺).

[000485] Example 23: 1-N-[4-[6-Carbamoyl-7-(2-methoxyethylamino)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (70)



[000486] 1-N-[4-[6-Carbamoyl-7-(2-methoxyethylamino)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (70): To a mixture of Compound **69** (200 mg, 337.25 μmol, 1 eq), 2-methoxyethan-1-amine (50.66 mg, 674.51 μmol, 2 eq), and K₂CO₃ (93.22 mg, 674.51 μmol, 2 eq) in DMF (10 mL) was added bis[(Z)-1-methyl-3-oxo-but-1-en-2-yl]copper (70.62 mg, 269.80 μmol, 0.8 eq) at 10 °C under nitrogen atmosphere. The resulting mixture was stirred for 2 h at 100 °C under nitrogen atmosphere. Heating was discontinued, and once the reaction mixture reached ambient temperature, it was quenched with aq saturated NH₄OH (50 mL) and extracted with EtOAc (3 x 30 mL). The combined extracts were dried over anhyd Na₂SO₄, filtered, and concentrated. The residue was purified by prep HPLC to give Compound **70** (22.2 mg, 11.33% yield). ¹H NMR (400 MHz, CD₃OD) δ 8.62 (s, 1H), 8.43 (d, 1H), 7.72 (d, 2H), 7.56 (dd, 2H), 7.23 (d, 2H), 7.07 (t, 2H), 6.99 (s, 1H), 6.32 (d, 1H), 3.72 (t, 2H), 3.43 (s, 5H), 1.64 (s, 4H); MS (EI) for $C_{30}H_{28}FN_5O_5$, found 558.1 (MH⁺).


[000487] The following compounds were prepared in a method analogous to Compound **70** in Example 23:

[000488] 1-N-[4-[6-Carbamoyl-7-(3-morpholin-4-ylpropylamino)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (71): ¹H NMR (400 MHz, CD₃OD) δ 8.60 (s, 1H), 8.42 (d, 1H), 7.71 (d, 2H), 7.59-7.53 (m, 2H), 7.22 (d, 2H), 7.07

(t, 2H), 6.97 (s, 1H), 6.30 (d, 1H), 3.73 (t, 4H), 3.38-3.33 (m, 2H), 2.61-2.45 (m, 6H), 2.00-1.91 (m, 2H), 1.64 (s, 4H); MS (EI) for $C_{34}H_{35}FN_6O_5$, found 627.2 (MH⁺).

[000489] 1-N-[4-[7-(Azetidin-1-yl)-6-carbamoylquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (72): ¹H NMR (400 MHz, CD₃OD) δ 8.46 (d, 1H), 8.32 (s, 1H), 7.71 (d, 2H), 7.59-7.53 (m, 2H), 7.21 (d, 2H), 7.11-7.04 (m, 2H), 6.86 (s, 1H), 6.37 (d, 1H), 4.11 (t, 4H), 2.42 (m, 2H), 1.64 (s, 4H); MS (EI) for $C_{30}H_{26}FN_5O_4$, found 540.1 (MH⁺).

[000490] Example 24: Ethyl 7-bromo-4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylate (79)



[000491] Ethyl 2-bromo-4-nitrobenzoate (74): To a solution of Compound **73** (20.0 g, 79.67 mmol, 1 eq) in EtOH (78.80 g, 1.71 mol, 100.00 mL, 21.47 eq) was added H₂SO₄ (7.81 g, 79.67 mmol, 4.25 mL, 1 eq) and the reaction was stirred at 80 °C for 24 h. The reaction mixture was concentrated, and the residue was partitioned between water (200 mL) and DCM (100 mL). The phases were separated and the aq phase was extracted with DCM (3 x 100 mL). The combined organic phases were dried over anhyd Na₂SO₄ and concentrated in vacuo to afford Compound **74** (21.0 g, 91.37% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.52-8.47 (m, 1H),

8.33-8.28 (m, 1H), 7.97 (d, 1H), 4.38 (q, 2H), 1.34 (t, 3H); MS (EI) for C₉H₈BrNO₄, found 274.0 (MH⁺).

[000492] Ethyl 4-amino-2-bromobenzoate (75): To a stirred solution of Compound **74** (**2**, 5.00 g, 17.3 mmol, 1 eq) in EtOH (22 mL), THF (22 mL), and water (11 mL) were added iron (2.03 g, 36.4 mmol, 2.1 eq) and NH₄Cl (1.95 g, 36.4 mmol, 1.27 mL, 2.1 eq), and the reaction mixture was heated to 80 °C for 3 h. The reaction was repeated in the exact same manner on a scale that used 16.0 g of Compound **74**. The two reactions were combined and filtered through Celite® (diatomaceous earth), and the filter cake was washed with EtOH (3 x 30 mL) and EtOAc (3 x 30 mL). The filtrate was concentrated, and diluted with aq saturated NaHCO₃ (200 mL) ensuring the pH was approximately 8, and extracted with EtOAc (3 x 150 mL). The combined extracts were dried over anhyd Na₂SO₄ and concentrated to afford crude product (19 g). The crude product was suspended in EtOAc (5 mL) and petroleum ether (25 mL) and stirred for 20 min at room temperature. The precipitate was collected by filtration and washed with petroleum ether (2 x 10 mL) to afford Compound **75** (16.0 g, 85.54% yield, 95% purity). ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.62 (d, 1H), 6.84 (d, 1H), 6.54 (dd, 1H), 6.13 (s, 2H), 4.19 (q, 2H), 1.27 (t, 3H); MS (EI) for C₉H₁₀BrNO₂, found 243.9 (MH⁺).

[000493] Ethyl 2-bromo-4-(((2,2-dimethyl-4,6-dioxo-1,3-dioxan-5-ylidene)methyl)amino)benzoate (76): Compound **76** was synthesized from Compounds **75** and **39** in a manner analogous to the preparation of Compound **40** from Compounds **38** and **39** in Example 12. Compound **76** was obtained (23 g, 88.1% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.23 (d, 1H), 8.63 (d, 1H), 7.92 (d, 1H), 7.56 (d, 1H), 7.23 (dd, 1H), 4.40 (q, 2H), 1.76 (s, 6H), 1.41 (t, 3H).


[000494] Ethyl 7-bromo-4-oxo-1,4-dihydroquinoline-6-carboxylate (77): Compound **77** was synthesized from Compound **76** in a manner analogous to the preparation of Compound **41** from Compound **40** in Example 12. Compound **77** was obtained (20 g, 94.4% yield, 60% purity) and used in subsequent reactions without further purification. MS (EI) for C₁₂H₁₀BrNO₃, found 296.0 (MH⁺).

[000495] Ethyl 7-bromo-4-chloroquinoline-6-carboxylate (78): Compound **78** was prepared from Compound **77** as described for Compound **42** in Example 12, except that instead of using neat POCl₃, the reaction was run in MeCN with 4.21 eq of POCl₃ at 90 °C for 1.5 h

(4.2 g, 33% yield). ^1H NMR (400 MHz, CD_3OD) δ 8.84 (d, 1H), 8.62 (s, 1H), 8.45 (s, 1H), 7.55 (d, 1H), 4.50 (q, 2H), 1.47 (t, 3H); MS (EI) for $\text{C}_{12}\text{H}_9\text{BrClNO}_2$, found 313.8 (MH $^+$).

[000496] Ethyl 7-bromo-4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylate (79): Compound **79** was prepared from Compound **78** and **3** as described in Example 12 for Compound **43** except that cesium carbonate was substituted for the potassium t-butoxide and the reaction mixture was heated only to 50 °C. (6.7 g, 80.5% yield). ^1H NMR (400 MHz, CD_3OD) δ 9.55 (s, 1H), 8.79 (s, 1H), 8.70 (d, 1H), 8.55 (s, 1H), 8.40 (s, 1H), 7.67 (d, 2H), 7.47 (dd, 2H), 7.18 (d, 2H), 7.06 (t, 2H), 6.56 (d, 1H), 4.48 (q, 2H), 1.79-1.73 (m, 2H), 1.68-1.62 (m, 2H), 1.46 (t, 3H); MS (EI) for $\text{C}_{29}\text{H}_{23}\text{BrFN}_3\text{O}_5$, found 592.0 (MH $^+$).

[000497] Example 25: 4-[4-[[1-[(4-Fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-(methylamino)quinoline-6-carboxylic acid (81)




[000498] Ethyl 4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)-7-(methylamino)quinoline-6-carboxylate (80): To a mixture of Compound **79** (1.00 g, 1.60 mmol, 1 eq), methanamine hydrogen chloride (216.55 mg, 3.21 mmol, 2 eq), Cs_2CO_3 (1.57 g, 4.81 mmol, 3 eq), and Xantphos (556.73 mg, 962.17 μmol , 0.6 eq) in 1,4-dioxane (25 mL) was added $\text{Pd}_2(\text{dba})_3$ (440.54 mg, 481.08 μmol , 0.3 eq) at 10 °C under a nitrogen atmosphere. The resulting mixture was stirred for 3 h at 100 °C under a nitrogen atmosphere. The reaction mixture was allowed to cool and then diluted with water (50 mL) and EtOAc (50 mL), neutralized to a pH of approximately 8 with aq saturated NH_4Cl solution, and filtered. The filter cake was washed with EtOAc. The phases of the filtrate were

separated, and the aq phase was extracted with EtOAc (3 x 30 mL). The combined EtOAc phases were dried over anhyd Na₂SO₄ and concentrated. The residue was purified by flash silica gel chromatography (0-80% EtOAc:petroleum ether, then 0-15% EtOH in DCM) followed by further purification by prep HPLC to give Compound **80** (250 mg, 27.30% yield). ¹H NMR (400 MHz, CDCl₃) δ 10.27 (s, 1H), 9.00 (s, 2H), 8.42 (s, 2H), 7.80 (d, 2H), 7.51 (dd, 2H), 7.39 (s, 1H), 7.17 (d, 2H), 7.03 (t, 2H), 6.38 (d, 1H), 4.44 (q, 2H), 3.11 (s, 3H), 1.85 (s, 2H), 1.73 (s, 2H), 1.48-1.43 (m, 3H); MS (EI) for C₃₀H₂₇FN₄O₅, found 543.1 (MH⁺).


[000499] 4-[4-[[1-[(4-Fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-(methylamino)quinoline-6-carboxylic acid (81): A mixture of Compound **80** (250. mg, 437.4 μmol, 1 eq) and lithium hydroxide:hydrate (183.69 mg, 4.38 mmol, 10 eq) in THF (10.0 mL) and water (10.0 mL) was stirred for 20 h at 10 °C. The reaction was diluted with water (40 mL), acidified to a pH of approximately 6 with aq saturated NH₄Cl, and extracted with EtOAc (4 x 40 mL). The combined extracts were dried over anhyd Na₂SO₄ and concentrated under reduced pressure to afford crude product which was subsequently purified by prep HPLC (column: Xtimate C18 150*25mm*5μm; mobile phase: [water(0.05% HCl)-ACN]; B%: 16%-46%, 9.5 min) to give Compound **81** HCl salt (120 mg, 47.3% yield). ¹H NMR (400 MHz, CD₃OD) δ 9.13 (s, 1H), 8.57 (d, 1H), 7.81 (d, 2H), 7.56 (dd, 2H), 7.34 (d, 2H), 7.07 (t, 2H), 6.88 (s, 1H), 6.61 (d, 1H), 3.07 (s, 3H), 1.66 (s, 4H); MS (EI) for C₂₈H₂₃FN₄O₅, found 515.1 (MH⁺).

[000500] Example 26: 1-N-[4-[6-Carbamoyl-7-(methylamino)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (82)



[000501] 1-N-[4-[6-Carbamoyl-7-(methylamino)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (82): Compound **82** was synthesized from Compound **81** in a manner analogous to the preparation of Compound **7** from Compound **6** in Example 3 (29.9 mg, 24% yield). ¹H NMR (400 MHz, CD₃OD) δ 8.61 (s, 1H), 8.43 (d, 1H), 7.72 (d, 2H), 7.59 - 7.54 (m, 2H), 7.23 (d, 2H), 7.07 (t, 2H), 6.94 (s, 1H), 6.31 (d, 1H), 2.97 (s, 3H), 1.64 (s, 4H); MS (EI) for C₂₈H₂₄FN₅O₄, found 514.1 (MH⁺).


[000502] Example 27: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(methylamino)-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (83)



[000503] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(methylamino)-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (83):


Compound **83** was synthesized from Compound **81** in a manner analogous to the preparation of Compound **8** from Compound **6** in Example 4 (33.3 mg, 26% yield). ¹H NMR (400 MHz, CD₃OD) δ 8.47 (s, 1H), 8.42 (d, 1H), 7.71 (d, 2H), 7.59-7.54 (m, 2H), 7.22 (d, 2H), 7.07 (t, 2H), 6.94 (s, 1H), 6.32 (d, 1H), 2.96 (s, 3H), 2.92 (s, 3H), 1.64 (s, 4H); MS (EI) for C₂₉H₂₆FN₅O₄, found 528.1 (MH⁺).

[000504] Example 28: Methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-(methylamino)quinoline-6-carboxylate (84)



[000505] Methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-(methylamino)quinoline-6-carboxylate (84): To a solution of Compound **81** (30 mg, 55.39 μmol, 1 eq) in MeOH (5 mL) was added TMSCHN₂ (2 M, 276.97 μL, 10 eq) at 10 °C, and the reaction was stirred at 10 °C for 6 h. The reaction was repeated in an identical fashion on another 45 mg of Compound **81**. The reaction solutions from both batches were combined and concentrated in vacuo. The resulting residue was purified by prep HPLC to give Compound **84** (25.5 mg, 55.16% yield). ¹H NMR (400 MHz, CD₃OD) δ 8.99 (s, 1H), 8.45 (d, 1H), 7.72 (d, 2H), 7.57 (dd, 2H), 7.23 (d, 2H), 7.07 (t, 2H), 6.97 (s, 1H), 6.29 (d, 1H), 3.95 (s, 3H), 3.01 (s, 3H), 1.64 (s, 4H); MS (EI) for C₂₉H₂₅FN₄O₅, found 529.1 (MH⁺).


[000506] Example 29: 7-((tert-Butoxycarbonyl)amino)-4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylic acid (86)



[000507] Ethyl 7-((tert-butoxycarbonyl)amino)-4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylate (85): A mixture of Compound **79** (1.5 g, 2.53 mmol, 1 eq), NH_2Boc (355.95 mg, 3.04 mmol, 1.2 eq), Xantphos (586.03 mg, 1.01 mmol, 0.4 eq), $\text{Pd}_2(\text{dba})_3$ (463.72 mg, 506.40 μmol , 0.2 equiv), and Cs_2CO_3 (2.47 g, 7.60 mmol, 3 equiv) in dioxane (20 mL) was degassed and purged with nitrogen three times, followed by stirring at 90 °C for 16 h under an atmosphere of nitrogen. The mixture was filtered and concentrated, and the residue was purified by flash silica gel chromatography (10 – 100% EtOAc in Petroleum ether) to give Compound **85** (1.3 g, 79.14% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.28 (s, 1H), 10.23 (s, 1H), 10.06 (s, 1H), 8.94 (s, 1H), 8.72 (d, 1H), 8.69 (s, 1H), 7.79 (d, 2H), 7.64 (dd, 2H), 7.30 (d, 2H), 7.16 (t, 2H), 6.46 (d, 1H), 4.42 (q, 2H), 1.53 (s, 9H), 1.48 (s, 4H), 1.38 (t, 3H); MS (EI) for $\text{C}_{34}\text{H}_{33}\text{FN}_4\text{O}_7$, found 629.2 (MH $^+$).

[000508] 7-((tert-Butoxycarbonyl)amino)-4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-6-carboxylic acid (86): Compound **86** was synthesized from Compound **85** in a manner analogous to the preparation of Compound **6** by the hydrolysis of the methyl ester of Compound **5** in Example 2 (0.7 g, 82.4% yield). ^1H NMR (400 MHz, CD_3OD) δ 9.13 (s, 1H), 8.57 (d, 1H), 7.81 (d, 2H), 7.56 (dd, 2H), 7.34 (d, 2H), 7.07 (t, 2H), 6.88 (s, 1H), 6.61 (d, 1H), 3.07 (s, 3H), 1.66 (s, 4H); MS (EI) for $\text{C}_{32}\text{H}_{29}\text{FN}_4\text{O}_7$, found 601.1 (MH $^+$).

[000509] **Example 30: 1-N-[4-(7-Amino-6-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (87)**




[000510] **1-N-[4-(7-Amino-6-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)-cyclopropane-1,1-dicarboxamide (87):** Compound **87** was synthesized from Compound **86** in two steps. The first step followed a manner analogous to the preparation of Compound **7** from Compound **6** in Example 3. Step 2 involved TFA-mediated BOC-deprotection using a standard procedure such as that employed in Example 5 which provided the final Compound **87** (41.9 mg, 35% yield over 2 steps). ¹H NMR (400 MHz, CD₃OD) δ 8.63 (s, 1H), 8.43 (d, 1H), 7.73 (d, 2H), 7.66 - 7.52 (m, 2H), 7.32 - 7.22 (m, 2H), 7.15 (s, 1H), 7.12 - 7.05 (m, 2H), 6.31 (d, 1H), 1.66 (s, 4H); MS (EI) for C₂₇H₂₂FN₅O₄, found 500.0 (MH⁺).

[000511] **Example 31: 1-N-[4-[7-Amino-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (88)**




[000512] **1-N-[4-[7-Amino-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (88):** Compound **88** was synthesized from Compound **86** in two steps. The first step followed a manner analogous to the preparation of Compound **8** from Compound **6** in Example 4. Step 2 involved TFA-mediated BOC-deprotection using a standard procedure such as that employed in Example 5 which provided the final Compound **88** (50.9 mg, 42% yield over 2 steps). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.19 (s, 1H), 10.05 (s, 1H), 8.70 (br d, 1H), 8.52 - 8.38 (m, 2H), 7.77 (d, 2H), 7.69 - 7.59 (m, 2H), 7.30 - 7.06 (m, 5H), 6.55 (s, 2H), 6.17 (d, 1H), 2.79 (d, 3H), 1.48 (s, 4H); MS (EI) for C₂₈H₂₄FN₅O₄, found 514.1 (MH⁺).

[000513] **Example 32: 7-Amino-4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]quinoline-6-carboxylic acid (89)**




[000514] 7-Amino-4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]quinoline-6-carboxylic acid (89): Compound **89** was synthesized from Compound **86** via TFA mediated BOC-deprotection as described in Example 5. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.19 (s, 1H), 10.07 (s, 1H), 8.80 (s, 1H), 8.48 (d, 1H), 7.76 (d, 2H), 7.69 - 7.58 (m, 2H), 7.24 (d, 2H), 7.19 - 7.10 (m, 3H), 6.16 (d, 1H), 1.48 (s, 4H); MS (EI) for C₂₇H₂₁FN₄O₅, found 501.1 (MH⁺).

[000515] Example 33: Methyl 7-amino-4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]quinoline-6-carboxylate (90)




[000516] Methyl 7-amino-4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]quinoline-6-carboxylate (90): A solution of Compound **89** (100 mg, 199.81 μmol, 1 eq) in MeOH (3 mL) and H₂SO₄ (0.1 mL) was stirred at 70 °C for 40 h. The reaction was concentrated in vacuo, and the resulting residue was purified by prep HPLC to give Compound **90** (28.5 mg, 26.88% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.20 (br s, 1H), 10.07 (br s, 1H), 8.81 (s, 1H), 8.50 (d, 1H), 7.77 (br d, 2H), 7.65 (br dd, 2H), 7.25 (br d, 2H), 7.21 - 7.06 (m, 3H), 6.83 (s, 2H), 6.15 (d, 1H), 3.90 (s, 3H), 1.48 (s, 4H); MS (EI) for C₂₈H₂₃FN₄O₅, found 515.2 (MH⁺).

[000517] Example 34: N-(4-((6-Carbamoyl-7-hydroxyquinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (91)




[000518] N-(4-((6-Carbamoyl-7-hydroxyquinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (91): A mixture of Compound 7 (250 mg, 485.91 μmol , 1 eq) and BBr_3 (1.04 g, 4.15 mmol, 0.4 mL, 8.54 eq) in 1,2-dichloroethane (2 mL) was stirred under an atmosphere of nitrogen at 50 °C for 1 h. The reaction mixture was cooled to room temperature and poured into aq saturated NaHCO_3 with vigorous stirring. The resulting residue was filtered, washed with water and then methyl *t*-butyl ether, and dried. The resulting crude compound was purified by prep HPLC to give Compound **91** (69.5 mg, 27.3% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 12.91 (s, 1H), 10.20 (s, 1H), 10.04 (s, 1H), 8.97 (s, 1H), 8.84 (s, 1H), 8.60-8.59 (d, 1H), 7.79 (s, 1H), 7.77-7.64 (d, 2H), 7.63-7.62 (d, 2H), 7.30-7.26 (m, 3H), 7.17-7.13 (m, 2H), 6.63-6.33 (d, 1H), 1.47 (s, 4H); MS (EI) for $\text{C}_{27}\text{H}_{21}\text{FN}_4\text{O}_5$, found 501.1 (MH $^+$).

[000519] Example 35: 1-N'-(4-Fluorophenyl)-1-N-[4-[(2-methyl-4-oxo-2,3-dihydropyrido[3,2-g][1,3]benzoxazin-6-yl)oxy]phenyl]cyclopropane-1,1-dicarboxamide (92)



[000520] 1-N'-(4-Fluorophenyl)-1-N-[4-[(2-methyl-4-oxo-2,3-dihydropyrido[3,2-g][1,3]benzoxazin-6-yl)oxy]phenyl]cyclopropane-1,1-dicarboxamide (92): To a solution of Compound **91** (130 mg, 259.75 μmol , 1 eq) in CHCl_3 (5 mL) was added T_3P (535 mg, 1.68 mmol, 0.5 mL, 6.47 eq) and acetaldehyde (393 mg, 8.91 mmol, 0.5 mL, 34.30 eq). The resulting mixture was stirred at 50 °C for 3 h then concentrated in vacuo. The residue was purified by prep HPLC to give Compound **92** (30.7 mg, 22.45% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.22 (s, 1H), 10.07 (s, 1H), 9.05 (s, 1H), 8.78 (s, 1H), 8.70 (d, 1H), 7.72 (d, 2H), 7.65 (d, 2H), 7.50 (s, 1H), 7.28 (d, 2H), 7.17 (d, 2H), 6.50 (d, 1H), 5.62 - 5.58 (m, 1H), 1.57 (d, 3H), 1.48 (s, 4H); MS (EI) for $\text{C}_{29}\text{H}_{23}\text{FN}_4\text{O}_5$, found 527.2 (MH $^+$).

[000521] Example 36: 1-N-[4-[(2-Ethyl-4-oxo-2,3-dihydropyrido[3,2-g][1,3]benzoxazin-6-yl)oxy]phenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (96)



[000522] 4-Chloro-7-methoxyquinoline-6-carboxamide (93): To a mixture of Compound **4** (2.00 g, 7.55 mmol, 1 eq) in MeOH (20 mL) was added $\text{NH}_3 \cdot \text{H}_2\text{O}$ (10.58 g, 75.50 mmol, 11.63 mL, 10 eq) at 10 °C. The mixture was stirred for 72 h at 70 °C. The resulting precipitate was collected by filtration and dried under vacuum to give Compound **93** (1.30 g, 69.1% yield); MS (EI) for $\text{C}_{11}\text{H}_9\text{ClN}_2\text{O}_2$, found 237.0 (MH^+).


[000523] 4-Chloro-7-hydroxyquinoline-6-carboxamide (94): To a mixture of Compound **93** (1.30 g, 5.22 mmol, 1 eq) in DCM (30 mL) was added BBr_3 (6.54 g, 26.09 mmol, 2.51 mL, 5 eq) in portions at 0 °C. The mixture was stirred for 6 h at 0 °C. The reaction was quenched with aq saturated NaHCO_3 (150 mL) and filtered. The resulting residue was dried under vacuum to give Compound **94** (1.00 g, 81.77% yield). MS (EI) for $\text{C}_{10}\text{H}_7\text{ClN}_2\text{O}_2$, found 222.8 (MH^+).

[000524] 6-Chloro-2-ethyl-2,3-dihydro-4H-[1,3]oxazino[5,6-g]quinolin-4-one (95): To a solution of Compound **94** (2 g, 8.98 mmol, 1 eq) in CHCl_3 (15 mL) was added propionaldehyde (1.60 g, 27.48 mmol, 2 mL, 3.06 eq) and T_3P (10.42 g, 32.76 mmol, 9.74 mL, 3.65 eq). The resulting mixture was stirred at 50 °C for 15 h followed by further addition of propionaldehyde (1.60 g, 27.48 mmol, 2 mL, 3.06 eq) and continued stirring at 50 °C for an additional 15 h. The mixture was then concentrated to remove solvent to give crude Compound **95** as a yellow solid (2.2 g, 85.8% yield) which was used into the next step without further purification. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.16 (br s, 1H), 8.87 (d, 1H), 8.66 (s, 1H), 7.72 (d, 1H), 7.62 (s, 1H), 5.45-5.47 (m, 1H), 1.38-1.44 (m, 2H), 1.06 (t, 3H); MS (EI) for $\text{C}_{13}\text{H}_{11}\text{ClN}_2\text{O}_2$, found 262.9 (MH^+).

[000525] 1-N-[4-[(2-Ethyl-4-oxo-2,3-dihydropyrido[3,2-g][1,3]benzoxazin-6-yl)oxy]phenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (96): A solution of Compound **95** (95 mg, 361.64 μmol , 1 eq) and Compound **3** (284.17 mg, 904.10 μmol , 2.5 eq)

in chlorobenzene (10 mL) was stirred at 140 °C for 15 h. The mixture was diluted with EtOAc (60 mL) and washed with aq 2 N NaOH (3 x 20 mL). The combined water phases were extracted with EtOAc (30 mL). The combined organic phases were concentrated, and the resulting residue was purified by prep-HPLC (column: YMC-Triart Prep C18 150*40mm*7um; mobile phase: [water (0.05% ammonia hydroxide v/v)-ACN]; B%: 53%-73%, 10min) to give Compound **96** as a white solid (7.1 mg, 3.6% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.23 (s, 1H), 10.07 (s, 1H), 9.05 (s, 1H), 8.78 (s, 1H), 8.69 (s, 1H), 7.78-7.86 (m, 2H), 7.64-7.67 (m, 2H), 7.51 (s, 1H), 7.28-7.30 (m, 2H), 7.14-7.18 (m, 2H), 6.49-6.51 (m, 1H), 5.43 (s, 1H), 1.88-1.91 (m, 2H), 1.48 (s, 4H), 1.06 (t, 3H); MS (EI) for C₃₀H₂₅FN₄O₅, found 541.1 (MH⁺).

[000526] Example 37: 1-N-[4-[6-Carbamoyl-7-(3-morpholin-4-ylpropoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (98)




[000527] 4-(3-Bromopropyl)morpholine (97): 3-Morpholinopropan-1-ol (2.0 g, 13.77 mmol, 1 eq) and PPh₃ (4.0 g, 15.25 mmol, 1.11 eq) were dissolved in THF (20 mL), and the reaction mixture chilled in an ice water bath under nitrogen. CBr₄ (5.0 g, 15.08 mmol, 1.09 eq) was added in portions over 15 min. After stirring for 30 min, the mixture was warmed to 30 °C for 18 h. The reaction was quenched with water (10 mL) and EtOAc (30 mL). The layers were separated, and the organic layer was extracted with 1 N HCl (2 x 15 mL). The pH of the combined aq extracts was adjusted to 10-11 with 4 N NaOH. The aq phase was extracted with EtOAc (3 x 30 mL). The combined EtOAc extracts were dried over anhyd Na₂SO₄ and concentrated to give Compound **97** (1.8 g, 63% yield) which was used in subsequent reactions without further purification. ¹H NMR (400 MHz, CDCl₃) δ 3.72-3.69 (m, 4H), 3.49-3.46 (t, 2H), 2.50-2.43 (m, 6H), 2.06-1.99 (m, 2H).

[000528] 1-N-[4-[6-Carbamoyl-7-(3-morpholin-4-ylpropoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (98): To a mixture of Compound **91** (45 mg, 85.87 μmol, 1 eq) and Compound **97** (53.48 mg, 257.00 μmol, 2.99 eq) in DMF (2 mL) was added Cs₂CO₃ (81.18 mg, 249.14 μmol, 2.90 eq) under nitrogen. The mixture was stirred at

80 °C for 2 h. The reaction mixture was diluted with water (50 mL) and extracted with EtOAc (2 x 20 mL). The combined organic phases were separated, washed with aq saturated NaCl, dried over anhyd Na₂SO₄, concentrated, and purified by prep HPLC to give Compound **98** (16.2 mg, 29.70% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.36 (s, 1H), 9.29-9.27 (d, 1H), 9.00 (s, 1H), 8.64-8.63 (d, 1H), 7.99-7.98 (d, 1H), 7.69-7.67 (d, 2H), 7.54 (s, 1H), 7.51-7.47 (m, 2H), 7.17-7.14 (d, 2H), 7.07-7.01 (m, 2H), 6.46-6.45 (d, 1H), 5.98-5.97 (d, 1H), 4.40-4.37 (t, 2H), 3.76-3.70 (m, 4H), 2.26-2.58 (m, 2H), 2.50 (s, 4H), 2.20-2.13 (m, 2H), 1.74-1.70 (m, 4H); MS (EI) for C₃₄H₃₄FN₅O₆, found 628.5 (MH⁺).

[000529] Example 38: 1-N-[4-[6-Carbamoyl-7-(2-methoxyethoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (103)



[000530] 6-Bromo-4-chloroquinolin-7-ol (99): A mixture of Compound **42** (5 g, 18.35 mmol, 1 eq) and BBr₃ (13.0 g, 51.89 mmol, 5 mL, 2.83 eq) in 1,2-dichloroethane (15 mL) was stirred under an atmosphere of nitrogen at 90 °C for 2.5 h. The reaction mixture was cooled to room temperature and poured into aq saturated NaHCO₃ with vigorous stirring. The resulting residue was filtered, washed with water and then methyl *t*-butyl ether, and dried under vacuum to give Compound **99** (7g, crude), which was used in subsequent steps without further purification. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.04-9.03 (d, 1H), 8.49 (s, 1H), 7.93-7.92 (d, 1H), 7.69 (s, 1H), 6.08 (s, 1H); MS (EI) for C₉H₅BrClNO, found 258.2 (MH⁺).

[000531] 6-Bromo-4-chloro-7-(2-methoxyethoxy)quinoline (100): A mixture of Compound **99** (500 mg, 1.93 mmol, 1 eq), 1-bromo-2-methoxy-ethane (1.08 g, 7.73 mmol, 726.35 μ L, 4 eq), and Cs_2CO_3 (1.90 g, 5.83 mmol, 3.01 eq) in DMF (4 mL) was stirred under an atmosphere of nitrogen at 80 °C for 2 h. The reaction mixture was cooled to room temperature, and water (2 mL) was added. The resulting residue was filtered, washed with water, and dried under vacuum to give Compound **100** (590 mg, 96% yield), which was used in subsequent reactions without further purification. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.80-8.79 (d, 1H), 8.37 (s, 1H), 7.65-7.64 (d, 1H), 7.62 (s, 1H), 4.39-4.37 (t, 2H), 3.79-3.77 (t, 2H), 3.37 (s, 3H); MS (EI) for $\text{C}_{12}\text{H}_{11}\text{BrClNO}_2$, found 315.8 (MH⁺).


[000532] N-(4-((6-Bromo-7-(2-methoxyethoxy)quinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (101): Compound **101** was synthesized from Compounds **100** and **3** in a manner analogous to the preparation of Compound **43** from Compounds **42** and **3** in Example 12 (65 mg, 69.2% yield). MS (EI) for $\text{C}_{29}\text{H}_{25}\text{BrFN}_3\text{O}_5$, found 594.1 (MH⁺).

[000533] N-(4-((6-Cyano-7-(2-methoxyethoxy)quinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (102): Compound **101** (65 mg, 109.35 μ mol, 1 eq) and CuCN (15 mg, 167.48 μ mol, 36.59 μ L, 1.53 eq) were combined in N-Methyl-2-pyrrolidone (NMP) (2 mL) in a sealed tube and heated at 150 °C under microwave irradiation for 45 min. The reaction mixture was partitioned between water (10 mL), NH_4OH (5 mL), and EtOAc (20 mL), the phases were separated, and the aq phase was further extracted with EtOAc. The combined organic phases were washed with aq saturated NaCl, dried over anhyd Na_2SO_4 , and concentrated to give Compound **102** (40 mg, 68% yield) which was used in subsequent steps without further purification. MS (EI) for $\text{C}_{30}\text{H}_{25}\text{FN}_4\text{O}_5$, found 541.5 (MH⁺).

[000534] 1-N-[4-[6-Carbamoyl-7-(2-methoxyethoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (103): Compound **102** (40 mg, 74.00 μ mol, 1 eq) and NaOH (100 mg, 2.50 mmol, 33.79 eq) were dissolved in DMSO (1.5 mL) and water (0.5 mL). The resulting mixture was heated at 80 °C for 1 h. The reaction mixture was partitioned between water (10 mL) and EtOAc (20 mL), the phases were separated, and the aq phase further extracted with EtOAc. The combined organic phases were washed with aq saturated NaCl, dried over anhyd Na_2SO_4 , and concentrated. The residue was purified by flash silica gel chromatography (50~100% EtOAc/Petroleum ether) to give Compound **103** (18.6 mg,

44% yield). ^1H NMR (400 MHz, CDCl_3) δ 9.30 (s, 1H), 9.27 (s, 1H), 9.02 (s, 1H), 8.64-8.63 (d, 1H), 8.15 (s, 1H), 7.69-7.66 (d, 2H), 7.51-7.48 (m, 3H), 7.17-7.15 (d, 2H), 7.07-7.03 (m, 2H), 6.47-6.45 (d, 1H), 5.92 (s, 1H), 4.44-4.42 (t, 2H), 3.91-3.89 (t, 2H), 3.50 (s, 3H), 1.75-1.68 (m, 4H); MS (EI) for $\text{C}_{30}\text{H}_{27}\text{FN}_4\text{O}_6$, found 559.4 (MH $^+$).

[000535] Example 39: 1-N-[4-[6-Carbamoyl-7-(2-hydroxyethoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (106)




[000536] 2-((6-Carbamoyl-4-chloroquinolin-7-yl)oxy)ethyl acetate (104): To a mixture of Compound **94** (200 mg, 853.44 μmol , 1 eq) and K_2CO_3 (589.77 mg, 4.27 mmol, 5 eq) in DMF (10 mL) was added 2-bromoethyl acetate (285.05 mg, 1.71 mmol, 187.53 μL , 2.00 eq) at 10 $^\circ\text{C}$. The resulting mixture was stirred for 1 h at 70 $^\circ\text{C}$. After cooling, the reaction mixture was diluted with water (50 mL) and extracted with EtOAc (3 x 30 mL). The combined extracts were washed with aq saturated NaCl (3 x 100 mL), dried over anhyd Na_2SO_4 , and concentrated in vacuo to give Compound **104** (220 mg, 79.33% yield). MS (EI) for $\text{C}_{14}\text{H}_{13}\text{ClN}_2\text{O}_4$, found 309.0 (MH $^+$).

[000537] 2-((6-Carbamoyl-4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinolin-7-yl)oxy)ethyl acetate (105): Compound **104** (200 mg, 615.45 μmol) was converted to Compound **105** by reaction with Compound **3** using a manner analogous to the preparation of Compound **43** from Compounds **42** and **3** in Example 12. The resulting reaction mixture was treated with water at 100 $^\circ\text{C}$ for 4 h before employing the usual reaction work-up as outlined in Example 12. The resulting product was a mixture of Compound **105** and Compound **106**. The mixture was separated by flash silica gel chromatography to give Compound **105** (100 mg, 24.93% yield), which was used in the next reaction, and Compound

106 (100 mg, 25.36% yield, 85% purity), which was combined with the product of the next reaction for final purification. For Compound **105**: MS (EI) for $C_{31}H_{27}FN_4O_7$, found 587.1 (MH⁺).

[000538] 1-N-[4-[6-Carbamoyl-7-(2-hydroxyethoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (106): To a solution of Compound **105** (isolated from the previous reaction (100 mg, 153.44 μ mol, 1 eq)) in THF (5 mL) and water (1 mL) was added LiOH \cdot H₂O (64.39 mg, 1.53 mmol, 10 eq). The resulting mixture was stirred for 1 h at 70 °C. After allowing the reaction mixture to cool to room temperature, the mixture was acidified to a pH of approximately 7 with aq saturated NH₄Cl solution, diluted with water (30 mL), and extracted with EtOAc (3 x 20 mL). The combined organic phases were dried over anhyd Na₂SO₄ and concentrated in vacuo to give crude Compound **106** (90 mg, 91.56% yield, 85% purity). This material was combined with the Compound **106** isolated from the previous step and purified by flash silica gel chromatography. The resulting product was suspended in a 1:1 mixture of DCM:MeCN and stirred for 10 min. The resulting solid was filtered, washed with MeCN followed by DCM, and dried in vacuo give Compound **106** (65.8 mg, 38.7 % yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.20 (s, 1H), 10.06 (s, 1H), 8.83 (s, 1H), 8.66 (d, 1H), 7.96 (s, 1H), 7.84 (s, 1H), 7.78 (d, 2H), 7.64 (dd, 2H), 7.56 (s, 1H), 7.27 (d, 2H), 7.15 (t, 2H), 6.47 (d, 1H), 5.14 (t, 1H), 4.32 (t, 2H), 3.86 (dd, 2H), 1.47 (s, 4H); MS (EI) for $C_{29}H_{25}FN_4O_6$, found 545.1 (MH⁺).

[000539] Example 40: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(2-hydroxyethoxy)-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (110)



[000540] 4-Chloro-7-methoxy-N-methylquinoline-6-carboxamide (107): Methylamine (8 M, 50 mL, 10.07 eq) in EtOH was added to a solution of Compound **4** (10 g, 39.74 mmol, 1 eq)


in THF (150 mL) at 30 °C and stirred at that temperature for 25 h. The reaction mixture was concentrated under vacuum. The residue was slurried with warm water (100 mL), and the resulting residue was filtered and dried under vacuum to give Compound **107** (9 g, 90.4% yield). MS (EI) for $C_{12}H_{11}ClN_2O_2$, found 251.0 (MH⁺).

[000541] 4-Chloro-7-hydroxy-N-methylquinoline-6-carboxamide (108): Compound **108** was synthesized from Compound **107** in a manner analogous to the preparation of Compound **94** from Compound **93** in Example 36 (730 mg, 62% yield). MS (EI) for $C_{11}H_9ClN_2O_2$, found 237.0 (MH⁺).

[000542] 2-((4-Chloro-6-(methylcarbamoyl)quinolin-7-yl)oxy)ethyl acetate (109): Compound **109** was synthesized from Compound **108** in a manner analogous to the preparation of Compound **104** from Compound **94** in Example 39 (200 mg, crude). MS (EI) for $C_{15}H_{15}ClN_2O_4$, found 323.1 (MH⁺).

[000543] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(2-hydroxyethoxy)-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (110): t-BuOK (70 mg, 623.83 μ mol, 2.01 eq) was added to a mixture of Compound **3** (107 mg, 340.43 μ mol, 1.10 eq) and Compound **109** (100 mg, 309.84 μ mol, 1 eq) in DMSO (10 mL). The resulting mixture was heated at 100 °C with stirring for 2 h. After cooling to room temperature, the reaction mixture was diluted with water (50 mL) and extracted with EtOAc (3 x 50 mL). The combined organic phases were washed with aq saturated NaCl (2 x 100 mL), dried over anhyd Na_2SO_4 , and concentrated under vacuum. The residue was purified by flash silica gel chromatography (0-20% MeOH in EtOAc) to give the crude product, which was purified by prep-TLC (20% MeOH in DCM, $R_f=0.3$) to give the Compound **110** (54.8 mg, 100% purity). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.22 (s, 1H), 10.07 (s, 1H), 8.75 (s, 1H), 8.66 (d, 1H), 8.49 (q, 1H), 7.78 (d, 2H), 7.64 (dd, 2H), 7.57 (s, 1H), 7.27 (d, 2H), 7.15 (t, 2H), 6.46 (d, 1H), 5.19 (t, 1H), 4.32 (t, 2H), 3.86 (br d, 2H), 2.87 (d, 3H), 1.47 (s, 4H); MS (EI) for $C_{30}H_{27}FN_4O_6$, found 559.1 (MH⁺).

[000544] Example 41: 1-N-[4-[6-Carbamoyl-7-(2-hydroxypropoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (115)



[000545] 4-Chloro-7-(2-oxopropoxy)quinoline-6-carboxamide (111): To a mixture of Compound **94** (200 mg, 853.4 μ mol, 1 eq) and K_2CO_3 (589.76 mg, 4.27 mmol, 5 eq) in DMF (10 mL) was added 1-chloropropan-2-one (1.45 g, 15.7 mmol, 11.97 μ L, 18.4 eq) at 10 °C. The reaction mixture was then stirred for 1 h at 65 °C. The reaction mixture was diluted with water (50 mL) and extracted with EtOAc (3 x 30 mL). The combined extracts were washed with aq saturated NaCl (3 x 100 mL), dried over anhyd Na_2SO_4 , and concentrated. The resulting residue was purified by flash silica gel chromatography (0-10 % MeOH in DCM). The resulting yellow residue was further purified prep HPLC (column: Waters Xbridge 150mm*25mm*5 μ m; mobile phase: [water (0.05% ammonia hydroxide v/v)-ACN]; B%: 17%-47%, 7.8min) to give Compound **111** (55 mg, 30.0% yield). MS (EI) for $C_{13}H_{11}ClN_2O_3$, found 279.0 (MH⁺).

[000546] 4-Chloro-7-(2-hydroxypropoxy)quinoline-6-carboxamide (112): To a mixture of Compound **111** (50 mg, 170.44 μ mol, 1 eq) in MeOH (10 mL) was added $NaBH_4$ (12.90 mg, 340.88 μ mol, 2 eq) at 10 °C. The reaction mixture was then stirred for 1 h at 50 °C. The reaction mixture was quenched with aq saturated NH_4Cl (30 mL) and extracted with EtOAc/*i*-PrOH (v/v=3:1, 5 x 20 mL). The combined extracts were dried over anhyd Na_2SO_4 and concentrated under reduced pressure to give Compound **112** (50 mg, 99.3% yield). MS (EI) for $C_{13}H_{13}ClN_2O_3$, found 302.8 (MH+Na)⁺.

[000547] 1-((6-Chloro-7-(2-oxopropoxy)quinolin-7-yl)oxy)propan-2-yl acetate (113): To a mixture of Compound **112** (45 mg, 128.25 μ mol, 1 eq) in DCM (10 mL) were added Ac_2O (65.46 mg, 641.24 μ mol, 60.06 μ L, 5 eq) and DMAP (15.67 mg, 128.25 μ mol, 1 eq) at 10 °C. The mixture was then stirred for 16 h at 10 °C. The reaction mixture was diluted with water (30 mL) and extracted with DCM (5 x 20 mL). The combined extracts were dried over anhyd Na_2SO_4 and concentrated. The residue was purified by flash silica gel chromatography (0-3%

MeOH in EtOAc) to afford Compound **113** (45 mg, 86.98% yield, 80% purity). MS (EI) for $C_{15}H_{15}ClN_2O_4$, found 344.8 (MH+Na)⁺.


[000548] 1-((6-Carbamoyl-4-(4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinolin-7-yl)oxy)propan-2-yl acetate (114): Compound **114** was synthesized from Compounds **113** and **3** in a similar manner as described for Compound **43** in Example 12 except that cesium carbonate was substituted for potassium t-butoxide and the reaction mixture was heated to 100 °C for 2 h (40 mg, 72.9% yield). MS (EI) for $C_{32}H_{29}FN_4O_7$, found 601.1 (MH⁺).

[000549] 1-N-[4-[6-Carbamoyl-7-(2-hydroxypropoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (115): To a solution of Compound **114** (40 mg, 63.27 μmol, 1 eq) in THF (5 mL) and water (1 mL) was added LiOH·H₂O (26.55 mg, 632.71 μmol, 10 eq). The mixture was stirred for 1 h at 70 °C. After cooling to room temperature, the reaction was diluted with DCM (30 mL), dried over anhyd Na₂SO₄, and concentrated under reduced pressure. The product was suspended in MeCN (1 mL) and stirred for 5 min at 10 °C. The resulting precipitate was collected by filtration and dried under vacuum to give Compound **115** (16.7 mg, 46.78% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.20 (s, 1H), 10.06 (s, 1H), 8.81 (s, 1H), 8.66 (d, 1H), 8.02 (d, 1H), 7.84 (s, 1H), 7.78 (d, 2H), 7.64 (dd, 2H), 7.54 (s, 1H), 7.27 (d, 2H), 7.15 (t, 2H), 6.47 (d, 1H), 5.15 (d, 1H), 4.27-4.21 (m, 1H), 4.17-4.04 (m, 2H), 1.47 (s, 4H), 1.22 (d, 3H); MS (EI) for $C_{30}H_{27}FN_4O_6$, found 559.1 (MH⁺).

[000550] The following compound was prepared by a sequence analogous to that taken to prepare Compound **115** from Compound **94** in Examples 41, substituting Compound **94** with Compound **108**:

[000551] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(2-hydroxypropoxy)-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (116). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.27 (s, 1H), 10.13 (s, 1H), 8.80 (s, 1H), 8.73 (d, 1H), 8.62 (d, 1H), 7.85 (d, 2H), 7.71 (dd, 2H), 7.62 (s, 1H), 7.33 (d, 2H), 7.22 (t, 2H), 6.54 (d, 1H), 5.28 (d, 1H), 4.31 (dd, 1H), 4.23-4.10 (m, 2H), 2.94 (d, 3H), 1.54 (s, 4H), 1.29 (d, 3H); MS (EI) for $C_{31}H_{29}FN_4O_6$, found 573.1 (MH⁺).

[000552] Example 42: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(2-hydroxypropoxy)-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (125)



[000553] 4-Chloro-7-methoxyquinoline-6-carboxylic acid (117): To a solution of Compound 4 (4.0 g, 15.89 mmol, 1 *eq*) in THF (30 mL) and water (6 mL) was added LiOH.H₂O (1.33 g, 31.79 mmol, 2 *eq*), and the resulting mixture was stirred at 20 °C for 10 h. The resulting solid was filtered to give crude Compound 117 as a white solid (4.4 g) which was used in the next step without further purification. MS (EI) for C₁₁H₈ClNO₃, found 238.0 (MH⁺).

[000554] 4-Chloro-N-(2,2-dimethoxyethyl)-7-methoxyquinoline-6-carboxamide (118): To a solution of Compound 117 (4 g, 16.83 mmol, 1 *eq*) and 2,2-dimethoxyethanamine (7.08 g, 67.33 mmol, 7.34 mL, 4 *eq*) in THF (40 mL) was added T₃P (16.07 g, 50.50 mmol, 15.02 mL, 3 *eq*) and DIPEA (8.70 g, 67.33 mmol, 11.73 mL, 4 *eq*). The resulting reaction mixture was stirred at 20 °C for 15 h. Additional T₃P (5 mL) and 2,2-dimethoxyethanamine (3 mL) were added to the mixture, and stirring continued at 20 °C for 25 h. The mixture was quenched with water (30 mL) and adjusted to pH 9-10 with aq 1 M NaOH. The resulting aq mixture was extracted with EtOAc (3 x 40 mL). The combined organic extracts were washed with aq saturated NaCl (15 mL) and concentrated to give crude Compound 118 as a white solid (4.1 g,

73.5% yield) which was used in the next step without further purification. MS (EI) for $C_{15}H_{17}ClN_2O_4$, found 325.1 (MH⁺).

[000555] 2-(4-Chloro-7-methoxyquinolin-6-yl)oxazole (119): To Compound **118** (2.5 g, 7.70 mmol, 1 *eq*) at 20 °C under nitrogen was added Eaton's Reagent (7.7 wt% phosphorus pentoxide solution in methanesulfonic acid, 23.09 mmol, 46.97 mL, 3 *eq*). The resulting mixture was heated at 145 °C for 7 h. The mixture was quenched with water (100 mL) and adjusted to pH 8-9 with Et₃N. The resulting mixture was extracted with EtOAc (4 x 40 mL). The combined organic extracts were washed with aq saturated NaCl (40 mL) and concentrated. The residue was purified by flash silica gel chromatography (ISCO®; 20 g SepaFlash® Silica Flash Column, Eluent of 0~80% Ethyl acetate/Petroleum ether gradient @ 32 mL/min) to give Compound **119** as a white solid (1.38 g, 68.8% yield). ¹H NMR (400 MHz, CDCl₃) δ 8.43 (s, 1H), 8.76 (d, 1H), 7.86 (s, 1H), 7.62 (s, 1H), 7.42 (d, 1H), 7.40 (s, 1H), 4.14 (s, 3H); MS (EI) for $C_{13}H_9ClN_2O_2$, found 261.0 (MH⁺).

[000556] 4-Chloro-6-(oxazol-2-yl)quinolin-7-ol (120): Compound **120** was synthesized from Compound **119** in a manner analogous to the way Compound **94** was synthesized from Compound **93** in Example 36. MS (EI) for $C_{12}H_7ClN_2O_2$, found 246.7 (MH⁺).

[000557] 1-((4-Chloro-6-(oxazol-2-yl)quinolin-7-yl)oxy)propan-2-one (121): Compound **121** was synthesized from Compound **120** in a manner analogous to the way Compound **111** was synthesized from Compound **94** in Example 41. MS (EI) for $C_{15}H_{11}ClN_2O_3$, found 302.7 (MH⁺).

[000558] 1-((4-Chloro-6-(oxazol-2-yl)quinolin-7-yl)oxy)propan-2-ol (122): Compound **122** was synthesized from Compound **121** in a manner analogous to the way Compound **112** was synthesized from Compound **111** in Example 41. MS (EI) for $C_{15}H_{13}ClN_2O_3$, found 326.9 [M+Na]⁺.


[000559] 1-((4-Chloro-6-(oxazol-2-yl)quinolin-7-yl)oxy)propan-2-yl acetate (123): Compound **123** was synthesized from Compound **122** in a manner analogous to the way Compound **113** was synthesized from Compound **112** in Example 41. ¹H NMR (400 MHz, CDCl₃) δ 8.83 (s, 1H), 7.23 (d, 1H), 7.83 (s, 1H), 7.54 (s, 1H), 7.40 (d, 1H), 7.34 (s, 1H), 5.41 - 5.49 (m, 1H), 4.25 (d, 2H), 2.06 (s, 3H), 1.43 (d, 3H); MS (EI) for $C_{17}H_{15}ClN_2O_4$, found 347.1 (MH⁺).

[000560] 1-((4-(1-((4-Fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)-6-(oxazol-2-yl)quinolin-7-yl)oxy)propan-2-yl acetate (124):

Compound **124** was synthesized from Compounds **123** and **3** using a variation of the manner that Compound **43** was synthesized from Compounds **42** and **3** in Example 12. Cesium carbonate was substituted for the potassium t-butoxide and the reaction mixture was heated to 100 °C for 3 h. MS (EI) for C₃₄H₂₉FN₄O₇, found 625.0 (MH⁺).

[000561] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(2-hydroxypropoxy)-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (125): To a solution of Compound **124** (400 mg, 640.4 μmol, 1 *eq*) in water (5 mL) and MeOH (5 mL) was added NaOH (51.23 mg, 1.28 mmol, 2 *eq*), and the mixture was stirred at 20 °C for 5 h. The mixture was diluted with water (30 mL). The resulting precipitate was filtered and washed with water (2 x 5 mL) and MeOH (3 mL). The solid was then lyophilized to give Compound **125** as a white solid (238.5 mg, 63.9% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.22 (s, 1H), 10.07 (s, 1H), 8.79 (s, 1H), 8.68 (d, 1H), 8.32 (s, 1H), 7.79 (d, 2H), 7.65 (t, 2H), 7.60 (s, 1H), 7.47 (s, 1H), 7.29 (d, 2H), 7.18 (t, 2H), 6.49 (d, 1H), 5.02 (d, 1H), 4.15 - 4.10 (m, 3H), 1.48 (s, 4H), 1.26 (d, 3H); MS (EI) for C₃₂H₂₇FN₄O₆, found 583.3 (MH⁺).

[000562] Example 43: Methyl 4-[2-chloro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylate (128)



[000563] N-(3-Chloro-4-hydroxyphenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (127): Compound **127** was synthesized from Compounds **1** and **126** in a


manner analogous to the preparation of Compound **3** from Compounds **1** and **2** in Example 1 (1.9 g, 73.5% yield). ^1H NMR (400 MHz, CDCl_3) δ 9.03-8.85 (m, 2H), 7.68 (d, 1H), 7.50-7.40 (m, 2H), 7.18 (dd, 1H), 7.08-7.00 (m, 2H), 6.97 (d, 1H), 5.53 (s, 1H), 1.64 (s, 4H); MS (EI) for $\text{C}_{17}\text{H}_{14}\text{ClFN}_2\text{O}_3$, found 349.0 (MH $^+$).

[000564] Methyl 4-[2-chloro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylate (128**):** Compound **128** was synthesized from Compounds **127** and **4** in a manner analogous to the preparation of Compound **43** from Compounds **42** and **3** in Example 12 (0.8 g, 32.1% yield). ^1H NMR (400 MHz, CDCl_3) δ 10.08 (s, 1H), 8.83 (s, 1H), 8.63 (d, 1H), 8.20 (s, 1H), 7.92 (d, 1H), 7.55-7.48 (m, 2H), 7.48-7.41 (m, 2H), 7.21 (d, 1H), 7.10-7.02 (m, 2H), 6.30 (d, 1H), 4.04 (s, 3H), 3.97 (s, 3H), 1.85-1.78 (m, 2H), 1.66-1.58 (m, 2H); MS (EI) for $\text{C}_{29}\text{H}_{23}\text{ClFN}_3\text{O}_6$, found 564.4 (MH $^+$).

[000565] The following compound was prepared in two steps in a sequence analogous to the way Compound **128** was synthesized in Example 43, replacing Compound **126** with 4-amino-2-fluorophenol:

[000566] Methyl 4-[2-fluoro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylate (129**):** ^1H NMR (400 MHz, CDCl_3) δ 10.05 (s, 1H), 8.83 (s, 1H), 8.65 (d, 1H), 8.09 (s, 1H), 7.78 (dd, 1H), 7.50 (s, 1H), 7.48-7.42 (m, 2H), 7.27-7.19 (m, 2H), 7.10-7.04 (m, 2H), 6.40 (d, 1H), 4.05 (s, 3H), 3.97 (s, 3H), 1.84-1.81 (m, 2H), 1.63-1.60 (m, 2H); MS (EI) for $\text{C}_{29}\text{H}_{23}\text{F}_2\text{N}_3\text{O}_6$, found 548.4 (MH $^+$).

[000567] Example 44: 4-[2-Chloro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylic acid (130**)**




[000568] 4-[2-Chloro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylic acid (130**):** Compound **130** was synthesized from Compound **128** in a manner analogous to the preparation of Compound **6** from Compound **5** in Example 2 (0.55 g, 76.6% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.41 (s, 1H), 10.10 (s, 1H), 8.58 (d, 1H), 8.25 (s, 1H), 8.10 (d, 1H), 7.75-7.59 (m, 3H), 7.47-7.37 (m,

2H), 7.16 (t, 2H), 6.32 (d, 1H), 3.92 (s, 3H), 1.47 (s, 4H); MS (EI) for C₂₈H₂₁ClFN₃O₆, found 550.1 (MH⁺).

[000569] The following compound was prepared from Compound **129** by a method analogous to the preparation of Compound **130** was synthesized from Compound **128** in Example 44:

[000570] **4-[2-Fluoro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylic acid (131):** ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.20 (s, 1H), 10.42 (s, 1H), 10.01 (s, 1H), 8.72 (d, 1H), 8.59 (s, 1H), 7.93 (d, 1H), 7.64 (dd, 2H), 7.56-7.51 (m, 2H), 7.50-7.44 (m, 1H), 7.15 (t, 2H), 6.54 (d, 1H), 3.99 (s, 3H), 1.47 (d, 4H); MS (EI) for C₂₈H₂₁F₂N₃O₆, found 534.4 (MH⁺).

[000571] **Example 45: 1-N'-[4-(6-Carbamoyl-7-methoxyquinolin-4-yl)oxy-3-chlorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (132)**




[000572] **1-N'-[4-(6-Carbamoyl-7-methoxyquinolin-4-yl)oxy-3-chlorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (132):** Compound **132** was synthesized from Compound **130** in a manner analogous to the preparation of Compound **7** from Compound **6** in Example 3 (43.1 mg, 40.9% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.38 (s, 1H), 10.04 (s, 1H), 8.71 (s, 1H), 8.67 (d, 1H), 8.11 (d, 1H), 7.89 (br s, 1H), 7.78 (br s, 1H), 7.73 - 7.61 (m, 3H), 7.54 (s, 1H), 7.46 (d, 1H), 7.16 (t, 2H), 6.37 (d, 1H), 4.04 (s, 3H), 1.47 (s, 4H); MS (EI) for C₂₈H₂₂ClFN₄O₅, found 549.4 (MH⁺).

[000573] The following compound was prepared from Compound **131** by a method analogous to the preparation of Compound **132** from Compound **130** in Example 45:

[000574] **1-N'-[4-(6-Carbamoyl-7-methoxyquinolin-4-yl)oxy-3-fluorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (133):** ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.40 (s, 1H), 10.01 (s, 1H), 8.67 (m, 2H), 7.93-7.88 (m, 2H), 7.76 (s, 1H), 7.64 (dd, 2H), 7.53-7.52 (m, 2H), 7.47-7.43 (m, 1H), 7.15 (t, 2H), 6.48 (d, 1H), 4.03 (s, 3H), 1.47 (s, 4H); MS (EI) for C₂₈H₂₂F₂N₄O₅, found 533.5 (MH⁺).

[000575] Example 46: 1-N'-[3-Chloro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (134)




[000576] 1-N'-[3-Chloro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (134): Compound **134** was synthesized from Compound **130** in a manner analogous to the preparation of Compound **8** from Compound **6** in Example 4 (49.7 mg, 48.6% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.38 (s, 1H), 10.04 (s, 1H), 8.66 (d, 1H), 8.63 (s, 1H), 8.40 (br d, 1H), 8.11 (d, 1H), 7.72 - 7.61 (m, 3H), 7.54 (s, 1H), 7.46 (d, 1H), 7.20-7.12 (m, 2H), 6.37 (d, 1H), 4.03 (s, 3H), 2.85 (d, 3H), 1.48 (s, 4H); MS (EI) for $\text{C}_{29}\text{H}_{24}\text{ClFN}_4\text{O}_5$, found 563.1 (MH^+).

[000577] The following compound was prepared from Compound **131** by a method analogous to the preparation of Compound **134** from Compound **130** in Example 46:

[000578] 1-N'-[3-Fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (135): ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.40 (s, 1H), 10.01 (s, 1H), 8.67 (d, 1H), 8.61 (s, 1H), 8.39 (d, 1H), 7.91 (d, 1H), 7.64 (dd, 2H), 7.53-7.52 (m, 2H), 7.47-7.42 (m, 1H), 7.15 (t, 2H), 6.48 (d, 1H), 4.02 (s, 3H), 2.84 (d, 3H), 1.47 (s, 4H); MS (EI) for $\text{C}_{29}\text{H}_{24}\text{F}_2\text{N}_4\text{O}_5$, found 547.5 (MH^+).

[000579] Example 47: Methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-6-methylquinoline-7-carboxylate (140)




[000580] Methyl 5-(((2,2-dimethyl-4,6-dioxo-1,3-dioxan-5-ylidene)methyl)amino)-2-methylbenzoate (137): Compound **137** was synthesized from Compounds **136** and **39** in a manner analogous to the preparation of Compound **40** from Compound **38** and **39** in Example 12 (4.74 g, 81.7% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.28 (s, 1H), 8.53 (s, 1H), 8.01-7.88 (d, 1H), 7.75-7.61 (dd, 1H), 7.45-7.30 (d, 1H), 3.85 (s, 3H), 2.49 (s, 3H), 1.67 (s, 6H).

[000581] Methyl 6-methyl-4-oxo-1,4-dihydroquinoline-7-carboxylate (138): Compound **138** was synthesized from Compound **137** in a manner analogous to the preparation of Compound **41** from Compound **40** in Example 12 (625 mg, 25.9% yield). MS (EI) for C₁₂H₁₁NO₃, found 218.1 (MH⁺).

[000582] Methyl 4-chloro-6-methylquinoline-7-carboxylate (139): Compound **139** was synthesized from Compound **138** in a manner similar to that described for Compound **42** in Example 12 except Compound **138** was heated in a mixture of MeCN (10 mL) and POCl₃ (3 mL) instead of neat POCl₃ (445 mg, 65.6% yield). MS (EI) for C₁₂H₁₀ClNO₂, found 236.3 (MH⁺).

[000583] Methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-6-methylquinoline-7-carboxylate (140): Compound **140** was synthesized from Compounds **139** and **3** in a manner analogous to the preparation of Compound **43** from synthesized from Compound **42** and **3** in Example 12 (320 mg, 36.7% yield). ¹H NMR (400 MHz, CD₃OD) δ 8.66-8.61 (d, 1H), 8.52 (s, 1H), 8.28 (s, 1H), 7.78-7.71 (d, 2H), 7.60-7.54 (m, 2H), 7.28-7.22 (d, 2H), 7.11-7.03 (m, 2H), 6.71-6.67 (d, 1H), 3.98 (s, 3H), 2.76 (s, 3H), 1.64 (s, 4H); MS (EI) for C₂₉H₂₄FN₃O₅, found 514.1 (MH⁺).


[000584] Example 48: 4-[4-[[1-[(4-Fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-6-methylquinoline-7-carboxylic acid (141)



[000585] 4-[4-[[1-[(4-Fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-6-methylquinoline-7-carboxylic acid (141): Compound **141** was synthesized from Compound **140** in a manner analogous to the preparation of Compound **6** from


Compound **5** in Example 2 (100 mg, 93.5% yield). ^1H NMR (400 MHz, CD_3OD) δ 8.66-8.61 (d, 1H), 8.47 (s, 1H), 8.27 (s, 1H), 7.78-7.73 (d, 2H), 7.78-7.71 (dd, 2H), 7.28-7.24 (d, 2H), 7.11-7.03 (t, 2H), 6.73-6.69 (d, 1H), 2.76 (s, 3H), 1.65 (s, 4H); MS (EI) for $\text{C}_{28}\text{H}_{22}\text{FN}_3\text{O}_5$, found 500.5 (MH^+).

[000586] Example 49: 1-N-[4-(7-Carbamoyl-6-methylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (142)




[000587] 1-N-[4-(7-Carbamoyl-6-methylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (142): Compound **142** was synthesized from Compound **141** in a manner analogous to the preparation of Compound **7** in Example 3 (17.4 mg, 34.9% yield). ^1H NMR (400 MHz, CD_3OD) δ 8.65-8.57 (d, 1H), 8.26 (s, 1H), 8.05 (s, 1H), 7.78-7.70 (d, 2H), 7.62-7.51 (m, 2H), 7.30-7.20 (m, 2H), 7.13-7.03 (m, 2H), 6.69-6.63 (d, 1H), 2.65 (s, 3H), 1.64 (s, 4H); MS (EI) for $\text{C}_{28}\text{H}_{23}\text{FN}_4\text{O}_4$, found 499.1 (MH^+).

[000588] Example 50: 1-N'-(4-Fluorophenyl)-1-N-[4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (143)



[000589] 1-N'-(4-Fluorophenyl)-1-N-[4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (143): Compound **143** was synthesized from Compound **141** in a manner analogous to the preparation of Compound **8** in Example 4 (11.9 mg, 17.8% yield). ^1H NMR (400 MHz, CD_3OD) δ 8.64-8.55 (d, 1H), 8.25 (s, 1H), 7.97 (s, 1H), 7.78-7.68 (d, 2H), 7.61-7.50 (m, 2H), 7.28-7.18 (d, 2H), 7.11-7.02 (m, 2H), 6.70-6.61 (d, 1H), 2.98 (s, 3H), 2.60 (s, 3H), 1.64 (s, 4H); MS (EI) for $\text{C}_{29}\text{H}_{25}\text{FN}_4\text{O}_4$, found 513.1 (MH^+).

[000590] Example 51: Methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-6-methoxyquinoline-7-carboxylate (150)



[000591] Methyl 5-amino-2-methoxybenzoate (145): To a solution of Compound **144** (5.00 g, 23.68 mmol, 1 eq) in EtOH (50 mL) was added 10% Pd/C (500 mg), and the resulting mixture was degassed and purged under H₂. The reaction mixture was stirred under H₂ (15 psi) at 25 °C for 12 h. The reaction mixture was filtered, and the filtrate was concentrated to give Compound **145** (4.5 g, 99.6% yield), which was used in subsequent steps without further purification. ¹H NMR (400 MHz, CDCl₃) δ 7.16 (t, 1H), 6.86-6.82 (m, 2H), 3.88 (s, 3H), 3.83 (s, 3H), 3.09 (br s, 2H).

[000592] Methyl 5-(((2,2-dimethyl-4,6-dioxo-1,3-dioxan-5-ylidene)methyl)amino)-2-methoxybenzoate (147): To a suspension of Compound **146** (1.95 g, 10.49 mmol, 1 eq) in isopropanol (40 mL) was added Compound **145** (2 g, 10.49 mmol, 1 eq) portionwise at 20 °C. The reaction was heated at 80 °C for 30 min. The mixture was cooled to 20 °C, and the precipitate was collected, washed with tert-butyl methyl ether (50 mL), and dried to give Compound **147** (3.3 g, 93.82% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.25 (d, 1H), 8.57 (d, 1H), 7.75 (d, 1H), 7.37 (dd, 1H), 7.05 (d, 1H), 3.95 (s, 3H), 3.93 (s, 3H), 1.76 (s, 6H).

[000593] Methyl 6-methoxy-4-oxo-1,4-dihydroquinoline-7-carboxylate (148): To DOWTHERM® A (eutectic mixture of 26.5% diphenyl + 73.5% diphenyl oxide)(CAS Reg. No.


8004-13-5) (10 mL) at 220 °C was added Compound **147** (1.0 g, 2.98 mmol, 1 eq) portionwise. The reaction was stirred at 220 °C for 10 min. The reaction was cooled to 20-25 °C and then diluted with petroleum ether (50 mL). The resulting residue was collected, washed with petroleum ether (3 x 30 mL), and dried to give Compound **148** (800 mg, 69.07% yield, 60% purity). ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.22 (d, 1H), 7.94 (d, 1H), 7.92 (s, 1H), 7.61 (s, 1H), 6.06 (d, 1H), 3.87 (s, 3H), 3.84 (s, 3H).

[000594] Methyl 4-chloro-6-methoxyquinoline-7-carboxylate (149): To a mixture of Compound **148** (700 mg, 1.80 mmol, 1 eq) in SOCl₂ (10 mL) was added 5 drops of DMF. The resulting mixture was stirred at 80 °C for 1 h. The reaction mixture was concentrated in vacuo, and the resulting residue was diluted with aq saturated NaHCO₃ (20 mL) and extracted with EtOAc (3 x 30 mL). The combined organic phases were washed with aq saturated NaCl (50 mL), dried over anhyd Na₂SO₄, and concentrated. The residue was purified by flash silica gel chromatography (ISCO®; 4 g SepaFlash® Silica Flash Column, Eluent of 0~30% EtOAc/Petroleum ether gradient at 25mL/min) to give Compound **149** (220 mg, 46.14% yield, 95% purity). ¹H NMR (400 MHz, CDCl₃) δ 8.71 (d, 1H), 8.49 (s, 1H), 7.55-7.49 (m, 2H), 4.07 (s, 3H), 3.99 (s, 3H); MS (EI) for C₁₂H₁₀ClNO₃, found 252.3 (MH⁺).

[000595] Methyl 4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-methoxyquinoline-7-carboxylate (150): To a mixture of Compound **149** (190 mg, 679.47 μmol, 1 eq), Compound **3** (235 mg, 747.67 μmol, 1.1 eq), K₃PO₄ (190 mg, 895.10 μmol, 1.32 eq), and 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl (XPhos) (33 mg, 69.22 μmol, 0.1 eq) in toluene (3 mL) and 1-methyl-2-pyrrolidinone (0.5 mL) was added Pd(OAc)₂ (8 mg, 35.63 μmol, 0.05 eq) under nitrogen. The reaction was stirred at 100 °C for 3 h. The mixture was diluted with EtOAc (30 mL), and any insoluble material was filtered off. The filtrate was diluted with water (30 mL) and extracted with EtOAc (3 x 30 mL). The combined organic phases were washed with aq saturated NaCl (50 mL), dried over anhyd Na₂SO₄, and concentrated. The residue was purified by flash silica gel chromatography (ISCO®; 4 g SepaFlash® Silica Flash Column, Eluent of 0 to approximately 50% EtOAc/Petroleum ether gradient at 25mL/min) to give Compound **150** (170 mg, 46.06% yield, 97.49% purity). ¹H NMR (400 MHz, CDCl₃) δ 9.46 (s, 1H), 8.59 (d, 2H), 8.47 (s, 1H), 7.71-7.62 (m, 3H), 7.52-7.44 (m, 2H), 7.20 (d, 2H), 7.07


(t, 2H), 6.57 (d, 1H), 4.05 (s, 3H), 3.99 (s, 3H), 1.79-1.74 (m, 2H), 1.69-1.64 (m, 2H); MS (EI) for $C_{29}H_{24}FN_3O_6$, found 530.2 (MH⁺).

[000596] Example 52: 4-[4-[[1-[(4-Fluorophenyl)carbamoyl]cyclopropanecarbonyl]-amino]phenoxy]-6-methoxyquinoline-7-carboxylic acid (151)




[000597] 4-[4-[[1-[(4-Fluorophenyl)carbamoyl]cyclopropanecarbonyl]-amino]phenoxy]-6-methoxyquinoline-7-carboxylic acid (151): Compound **151** was synthesized from Compound **150** in a manner analogous to the preparation of Compound **81** from Compound **80** in Example 25 (110 mg, 75.2% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.21 (s, 1H), 10.05 (s, 1H), 8.60 (d, 1H), 8.18 (s, 1H), 7.79 (d, 2H), 7.69-7.59 (m, 3H), 7.27 (d, 2H), 7.15 (t, 2H), 6.60 (d, 1H), 3.97 (s, 3H), 1.53-1.42 (m, 4H); MS (EI) for $C_{28}H_{22}FN_3O_6$, found 516.1 (MH⁺).

[000598] Example 53: 1-N-[4-(7-Carbamoyl-6-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (152)




[000599] 1-N-[4-(7-Carbamoyl-6-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (152): Compound **152** was synthesized from Compound **151** in a manner analogous to the preparation of Compound **7** from Compound **6** in Example 3 (25.6 mg, 30.7% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.20 (s, 1H), 10.04 (s, 1H), 8.58 (d, 1H), 8.25 (s, 1H), 7.87 (br s, 1H), 7.78 (d, 2H), 7.72 (br s, 1H), 7.67-7.60 (m, 3H), 7.26 (d, 2H), 7.15 (t, 2H), 6.58 (d, 1H), 4.01 (s, 3H), 1.52-1.46 (m, 4H); MS (EI) for $C_{28}H_{23}FN_4O_5$, found 515.1 (MH⁺).

[000600] Example 54: 1-N'-(4-Fluorophenyl)-1-N-[4-[6-methoxy-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (153)




[000601] 1-N'-(4-Fluorophenyl)-1-N-[4-[6-methoxy-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (153): Compound **153** was synthesized from Compound **151** in a manner analogous to the preparation of Compound **8** from Compound **6** in Example 4 (25.2 mg, 53.1% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.23 (s, 1H), 10.07 (s, 1H), 8.58 (d, 1H), 8.39 (q, 1H), 8.20 (s, 1H), 7.79 (d, 2H), 7.69-7.60 (m, 3H), 7.26 (d, 2H), 7.15 (t, 2H), 6.58 (d, 1H), 4.00 (s, 3H), 2.84 (d, 3H), 1.52-1.44 (m, 4H); MS (EI) for C₂₉H₂₅FN₄O₅, found 529.1 (MH⁺).

[000602] Example 56: Methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]quinoline-7-carboxylate (162)




[000603] Methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]quinoline-7-carboxylate (162): Compound **162** was synthesized from Compounds **161** and **3** in a manner analogous to the preparation of Compound **43** from Compounds **42** and **3** in Example 12 (20 mg, 16% yield). ¹H NMR (400 MHz, CDCl₃) δ 9.42 (s, 1H), 8.81 (d, 1H), 8.75 (d, 1H), 8.60 (br s, 1H), 8.43 (d, 1H), 8.19 (dd, 1H), 7.69-7.63 (m, 2H), 7.51-7.44 (m, 2H), 7.23-7.18 (m, 2H), 7.07 (t, 2H), 6.63 (d, 1H), 4.03 (s, 3H), 1.79-1.74 (m, 2H), 1.69-1.65 (m, 2H); MS (EI) for C₂₈H₂₂FN₃O₅, found 500.1 (MH⁺).

[000604] Example 57: 4-[4-[[1-[(4-Fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]quinoline-7-carboxylic acid (163)




[000605] 4-[4-[[1-[(4-Fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]quinoline-7-carboxylic acid (163): Compound **163** was synthesized from Compound **162** in a manner analogous to the preparation of Compound **6** from Compound **5** in Example 2 (70 mg, 68.4% yield). ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 10.22 (s, 1H), 10.06 (s, 1H), 8.79 (d, 1H), 8.58 (s, 1H), 8.43 (d, 1H), 8.17-8.10 (m, 1H), 7.79 (d, 2H), 7.65 (dd, 2H), 7.29 (d, 2H), 7.16 (t, 2H), 6.69 (d, 1H), 1.48 (s, 4H); MS (EI) for $\text{C}_{27}\text{H}_{20}\text{FN}_3\text{O}_5$, found 486.1 (MH $^+$).

[000606] Example 58: 1-N-[4-(7-Carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (164)



[000607] 1-N-[4-(7-Carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)-cyclopropane-1,1-dicarboxamide (164): Compound **164** was synthesized from Compound **163** in a manner analogous to the preparation of Compound **7** from Compound **6** in Example 3 (17.7 mg, 56.2% yield). ^1H NMR (400 MHz, CDCl_3) δ 9.47 (s, 1H), 8.73 (d, 1H), 8.60 (br s, 1H), 8.50-8.43 (m, 2H), 8.08 (dd, 1H), 7.66 (d, 2H), 7.51-7.43 (m, 2H), 7.19 (d, 2H), 7.06 (t, 2H), 6.62 (d, 1H), 6.36 (br s, 1H), 5.73 (br s, 1H), 1.80-1.73 (m, 2H), 1.70-1.65 (m, 2H); MS (EI) for $\text{C}_{27}\text{H}_{21}\text{FN}_4\text{O}_4$, found 485.1 (MH $^+$).


[000608] Example 59: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(methycarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (165)



[000609] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (165):

Compound **165** was synthesized from Compound **163** in a manner analogous to the preparation of Compound **8** from Compound **6** in Example 4 (16.5 mg, 33.9% yield). ^1H NMR (400 MHz, CDCl_3) δ 9.45 (br s, 1H), 8.71 (d, 1H), 8.63 (br s, 1H), 8.43 (d, 1H), 8.37 (d, 1H), 8.04 (d, 1H), 7.65 (d, 2H), 7.50-7.44 (m, 2H), 7.19 (d, 2H), 7.06 (t, 2H), 6.60 (d, 1H), 6.40 (br s, 1H), 3.11 (d, 3H), 1.79 - 1.74 (m, 2H), 1.69-1.64 (m, 2H); MS (EI) for $\text{C}_{28}\text{H}_{23}\text{FN}_4\text{O}_4$, found 499.1 (MH $^+$).

[000610] Example 60: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(2-hydroxyethoxycarbamoyl)-quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (166)



[000611] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(2-hydroxyethoxycarbamoyl)-quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (166):

Compound **166** was synthesized from Compound **163** in a manner analogous to the preparation of Compound **26** from Compound **6** in Example 6 (27 mg, 46.2% yield). ^1H NMR (400 MHz, CD_3OD) δ 8.73 (d, 1H), 8.53 (d, 1H), 8.44 (d, 1H), 8.03 (dd, 1H), 7.77 (d, 2H), 7.59 (dd, 2H), 7.28 (d, 2H), 7.10 (t, 2H), 6.75 (d, 1H), 4.17 - 4.12 (m, 2H), 3.88 - 3.83 (m, 2H), 1.66 (s, 4H); MS (EI) for $\text{C}_{29}\text{H}_{25}\text{FN}_4\text{O}_6$, found 545.1 (MH $^+$).

[000612] The following compounds were prepared in a method analogous to Compound **166** in Example 60:


[000613] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(oxetan-3-ylloxycarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (167):

^1H NMR (400 MHz, CD_3OD) δ 8.72 (d, 1H), 8.50 (d, 1H), 8.43 (d, 1H), 8.02 (dd, 1H), 7.79 - 7.73 (m, 2H), 7.62 - 7.56 (m, 2H), 7.31


- 7.25 (m, 2H), 7.09 (t, 2H), 6.74 (d, 1H), 5.19 (s, 1H), 4.93 (m, 2H), 4.88 - 4.81 (m, 2H), 1.70 - 1.63 (m, 4H); MS (EI) for $C_{30}H_{25}FN_4O_6$, found 557.1 (MH⁺).

[000614] N-(4-((7-(((2,2-Dimethyl-1,3-dioxolan-4-yl)methoxy)carbamoyl)quinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (**168**): MS (EI) for $C_{29}H_{25}FN_4O_6$, found 615.3 (MH⁺).


[000615] **Example 61:** 1-N-[4-[7-[(2R)-2,3-Dihydroxypropoxy]carbamoyl]quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (**169**) and 1-N-[4-[7-[(2S)-2,3-Dihydroxypropoxy]carbamoyl]quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (**170**)



[000616] 1-N-[4-[7-[(2R)-2,3-Dihydroxypropoxy]carbamoyl]quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (**169**) and 1-N-[4-[7-[(2S)-2,3-Dihydroxypropoxy]carbamoyl]quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (**170**): Compounds **169** and **170** were synthesized from Compound **168** in a manner analogous to the preparation of Compounds **30** and **31** from Compound **28** in Example 7. Compound **169**: 1H NMR (400 MHz, CD_3OD) δ 8.73 (d, 1H), 8.52 (d, 1H), 8.44 (d, 1H), 8.02 (dd, 1H), 7.77 (d, 2H), 7.62 - 7.55 (m, 2H), 7.30 - 7.25 (m, 2H), 7.09 (t, 2H), 6.75 (d, 1H), 4.21 - 4.15 (m, 1H), 4.06 (s, 1H), 4.02 - 3.96 (m, 1H), 3.68 (m, 2H), 1.66 (d, 4H); MS (EI) for $C_{30}H_{27}FN_4O_7$, found 575.2 (MH⁺). Compound **170**: 1H NMR (400 MHz, CD_3OD) δ 8.73 (d, 1H), 8.52 (d, 1H), 8.43 (d, 1H), 8.02 (dd, 1H), 7.77 (d, 2H), 7.62 - 7.54 (m, 2H), 7.28 (d, 2H), 7.14 - 7.04 (m, 2H), 6.75 (d, 1H), 4.22 - 4.14 (m, 1H), 4.12 - 3.92 (m, 2H), 3.73 - 3.64 (m, 2H), 1.66 (s, 4H); MS (EI) for $C_{30}H_{27}FN_4O_7$, found 575.1 (MH⁺).


[000617] Example 62: 4-((6-Bromo-7-methoxyquinolin-4-yl)oxy)aniline (171)

[000618] 4-((6-Bromo-7-methoxyquinolin-4-yl)oxy)aniline (171): To a mixture of Compound **42** (1.75 g, 6.38 mmol) and 4-aminophenol **2** (1.1 g, 10 mmol) in DMA (15 mL) was added Cs_2CO_3 (3.3 g, 10 mmol) at room temperature. The mixture was stirred at 100°C for 2 h. The mixture was allowed to cool to 20°C , diluted with water, and filtered. The crude residue was purified by flash silica gel chromatography to give Compound **171** (900 mg, 40% yield). MS (EI) for $\text{C}_{16}\text{H}_{13}\text{BrN}_2\text{O}_2$, found: 345 (MH^+).

[000619] Example 63: 4-((7-Methoxy-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinolin-4-yl)oxy)aniline (172)

[000620] 4-((7-Methoxy-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinolin-4-yl)oxy)aniline (172): To a mixture of Compound **171** (200 mg, 0.58 mmol), bis(pinacolato)diboron (220 mg, 0.87 mmol), and potassium acetate (170 mg, 1.7 mmol) in 1,4-dioxane (3 mL) was added $\text{Pd}(\text{dppf})\text{Cl}_2$ (42 mg, 0.06 mmol). The resulting mixture was heated at 100°C for 2 h. After cooling, the reaction mixture was diluted with EtOAc, washed with water followed by aq saturated NaCl, concentrated, and purified by flash silica gel chromatography to give Compound **172** (43% yield). MS (EI) for $\text{C}_{22}\text{H}_{25}\text{BN}_2\text{O}_4$, found: 393 (MH^+).

[000621] Example 64: 1-N-[4-[6-(3-Cyano-2-fluorophenyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (175)



[000622] 3-(4-(4-Aminophenoxy)-7-methoxyquinolin-6-yl)-2-fluorobenzonitrile (174):

In a microwave reaction tube were mixed Compound **172** (50 mg, 0.13 mmol), Compound **173** (25 mg, 0.13 mmol), Na₂CO₃ (41 mg, 0.39 mmol), bis(di-*tert*-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II) (9 mg, 0.013 mmol), 1,4-dioxane (2 mL), and water (0.4 mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 150 °C. After cooling, the mixture was extracted with EtOAc, washed with aq saturated NaCl, and concentrated. The crude product was purified by flash column chromatography to give Compound **174**. MS (EI) for C₂₃H₁₆FN₃O₂, found: 386 (MH⁺).

[000623] 1-N-[4-[6-(3-Cyano-2-fluorophenyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (175):


To a mixture of Compound **174**, Compound **1** (22 mg, 0.1 mmol) and DIEA (25 mg, 0.2 mmol) in DCM (2 mL) was added HATU (38 mg, 0.1 mmol). The mixture was stirred at room temperature until the reaction was complete and then diluted with EtOAc, washed with aq saturated NaHCO₃, and concentrated. The crude product was purified by flash column chromatography to give Compound **175**. MS (EI) for C₃₄H₂₄F₂N₄O₄, found 591 (MH⁺).

[000624] The following compounds were prepared in a manner analogous to the method used to synthesize Compound **175** in two steps from Compound **172** in Example 64:

[000625] **1-N'-(4-Fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-2-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide (176):** 2-Bromopyridine was used in place of Compound 173. MS (EI) for $C_{32}H_{25}FN_4O_4$, found 549 (MH⁺).

[000626] **1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1-methylimidazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (177):** 4-Bromo-1-methyl-1H-imidazole was used in place of Compound 173. MS (EI) for $C_{31}H_{26}FN_5O_4$, found 552 (MH⁺).

[000627] **Example 65: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(5-methylfuran-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (180)**



[000628] **4-((7-Methoxy-6-(5-methylfuran-2-yl)quinolin-4-yl)oxy)aniline (179):** In a microwave reaction tube were mixed Compound 171 (100 mg, 0.29 mmol), Compound 178 (82 mg, 0.43 mmol), Na_2CO_3 (92 mg, 0.9 mmol), bis(di-*tert*-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II) (20 mg, 0.029 mmol), 1,4-dioxane (2.5 mL), and water (0.5 mL). The resulting mixture was irradiated in a microwave reactor for 5 min at 150 °C. After cooling, the mixture was extracted with EtOAc, washed with aq saturated NaCl, and concentrated. The crude product was purified by flash column chromatography to give Compound 179. MS (EI) for $C_{21}H_{18}N_2O_3$, found: 347 (MH⁺).

[000629] **1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(5-methylfuran-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (180):** Compound 180 was synthesized from Compound 179 in a manner analogous to the method used to synthesize Compound 175 from Compound 174 in Example 64. MS (EI) for $C_{32}H_{26}FN_3O_5$, found 552 (MH⁺).

[000630] The following compounds were prepared in a manner analogous to the method used to synthesize Compound **180** in two steps from Compound **171** in Example 65:

[000631] tert-Butyl 2-[4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]-amino]phenoxy]-7-methoxyquinolin-6-yl]pyrrole-1-carboxylate (181): *t*-Butyl 2-(trifluoro-14-borane-1-yl)-1H-pyrrole-1-carboxylate, potassium salt was used in place of Compound **178**. MS (EI) for $C_{36}H_{33}FN_4O_6$, found 637 (MH⁺).

[000632] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1-methylpyrazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (182): 1-Methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole was used in place of Compound **178**. MS (EI) for $C_{31}H_{26}FN_5O_4$, found 552 (MH⁺).

[000633] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1,2-oxazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (183): 4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole was used in place of Compound **178**. MS (EI) for $C_{30}H_{23}FN_4O_5$, found: 539 (MH⁺).


[000634] 1-N-[4-[6-(3,5-Dimethyl-1,2-oxazol-4-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (184): 3,5-Dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole was used in place of Compound **178**. MS (EI) for $C_{32}H_{27}FN_4O_5$, found 567 (MH⁺).

[000635] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrazol-5-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (185): 5-(Trifluoro-14-borane-1-yl)-1H-pyrazole, potassium salt was used in place of Compound **178**. MS (EI) for $C_{30}H_{24}FN_5O_4$, found 538 (MH⁺).

[000636] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (186): 4-(Ttrifluoro-14-borane-1-yl)-1H-pyrazole, potassium salt was used in place of Compound **178**. MS (EI) for $C_{30}H_{24}FN_5O_4$; found 538 (MH⁺).


[000637] 1-N-[4-(6-Cyclopropyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (187): Cyclopropyltrifluoro-14-borane, potassium salt was used in place of Compound **178**. MS (EI) for $C_{30}H_{26}FN_3O_4$, found 512 (MH⁺).

[000638] Example 66: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrrol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (188)



[000639] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrrol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (188): Compound **181** and excess TFA were stirred in DCM at room temperature until the reaction was complete. The mixture was concentrated and purified by prep HPLC to give Compound **188** (50% yield). MS (EI) for $C_{31}H_{25}FN_4O_4$, found: 537 (MH⁺).


[000640] Example 67: N-(4-Fluorophenyl)-N-(4-((6-formyl-7-methoxyquinolin-4-yl)oxy)phenyl)cyclopropane-1,1-dicarboxamide (190)



[000641] 4-Chloro-7-methoxyquinoline-6-carbaldehyde (189): To a -78 °C mixture of Compound **42** (320 mg, 1.17 mmol) in THF (8 mL) was added nBuLi (0.6 mL, 2.5 M in THF, 1.5 mmol). The mixture was stirred at -78 °C for 45 min, and then DMF (0.5 mL) was added. The mixture was warmed to 0 °C, and the stirring was continued for 1 h. The reaction was quenched with saturated NH₄Cl and extracted with EtOAc. The organic phase was concentrated, and the crude residue was purified by flash silica gel chromatography to give Compound **189** (130 mg, 50% yield). MS (EI) for $C_{11}H_8ClNO_2$, found 222 (MH⁺).


[000642] N-(4-Fluorophenyl)-N-(4-((6-formyl-7-methoxyquinolin-4-yl)oxy)phenyl)cyclopropane-1,1-dicarboxamide (190): Compound **190** was synthesized from Compound **189** using a modification of the method used to synthesize Compound **43** from Compound **42** in Example 12, substituting the potassium t-butoxide with cesium carbonate and using DMF as the solvent.

[000643] Example 68: 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(1H-imidazol-2-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (191)




[000644] 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(1H-imidazol-2-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (191): Compound **190** (60 mg, 0.12 mmol), glyoxal (60 mg, 40% in water) and ammonium hydroxide (130 mg, 30% in water) were mixed in iPrOH (1.5 mL). The resulting mixture was stirred at ambient (room) temperature for 12 h. The mixture was extracted with EtOAc and washed with aq saturated NaCl. The organic phase was concentrated and purified by prep HPLC to give Compound **191** (35 mg, 54% yield). MS (EI) for $C_{30}H_{24}FN_5O_4$, found 538.1 (MH⁺).

[000645] Example 69: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3-oxazol-5-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (192)




[000646] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3-oxazol-5-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (192): Compound **190** (30 mg, 0.06 mmol), TosMIC (15 mg, 0.08 mmol), and K_2CO_3 (15 mg, 0.11 mmol) were mixed in iPrOH (1 mL). The resulting mixture was stirred at 80 °C for 30 min. The mixture was concentrated to dryness. The residue was extracted with EtOAc. The organic layer was washed with water and concentrated. The crude product was purified by prep HPLC to give Compound **192** (14 mg, 43% yield). MS (EI) for $C_{30}H_{23}FN_4O_5$, found 539 (MH⁺).

[000647] Example 70: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(E)-methoxyiminomethyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (193)



[000648] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-[(E)-methoxyiminomethyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (193): To a mixture of Compound **190** (150 mg, 300.31 μmol , 1 eq) in EtOH (3 mL) and water (0.6 mL) was added O-methylhydroxylamine-HCl (41.74 mg, 500 μmol , 1.7 eq). The mixture was stirred at 50 °C for 2 h. The resulting precipitate was filtered, slurried in 50 mL of water, and filtered again. The filter cake was lyophilized to obtain Compound **193** (125.0 mg, 73.2% yield). ^1H NMR (400 MHz, DMSO- d_6) δ 10.32 (s, 1H), 10.05 (s, 1H), 8.87 (s, 1H), 8.68 (s, 1H), 8.51 (s, 1H), 7.85 (d, 2H), 7.72 - 7.58 (m, 3H), 7.37 (d, 2H), 7.16 (t, 2H), 6.72 (br d, 1H), 4.07 (s, 3H), 3.99 (s, 3H), 1.49 (s, 4H); MS (EI) for $\text{C}_{29}\text{H}_{25}\text{FN}_4\text{O}_5$, found 529.2 (MH $^+$).

[000649] Example 71: tert-Butyl 3-[4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-methoxyquinolin-6-yl]-3-hydroxyazetidine-1-carboxylate (195)



[000650] *t*-Butyl 3-(4-chloro-7-methoxyquinolin-6-yl)-3-hydroxyazetidine-1-carboxylate (194): A mixture of Compound **42** (200 mg, 0.73 mmol) and THF (4 mL) was cooled to -78°C. *n*-BuLi (0.4 mL, 1.0 mmol, 2.5 M in THF) was slowly added. The resulting mixture was stirred at -78°C for 40 min. A solution of N-Boc 3-oxoazetidine (125 mg, 0.73 mmol) in THF (0.5 mL)

was added. The reaction mixture was warmed to 0 °C, and the stirring was continued for 1 h. The reaction was quenched with water and extracted with EtOAc. The organic phase was washed with aq saturated NaCl, concentrated, and purified by flash silica gel chromatography to give Compound **194** (38% yield). MS (EI) for C₁₈H₂₁ClN₂O₄, found: 365 (MH⁺).


[000651] tert-Butyl 3-[4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-methoxyquinolin-6-yl]-3-hydroxyazetidine-1-carboxylate

(195): Compound **195** was synthesized from Compound **194** in a manner similar to the preparation of Compound **43** from Compound **42** in Example 12, except cesium carbonate was used instead of t-butoxide, DMF was the solvent, the reaction temperature was 80 °C, and the reaction time was 12 h. Compound **195** was obtained in 68% yield. MS (EI) for C₃₅H₃₅FN₄O₇, found 643 (MH⁺).

[000652] The following compound was made in a manner analogous to Compound **195** in Example 71 with oxetan-3-one replacing the Boc protected azetidin-3-one:


[000653] 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(3-hydroxyoxetan-3-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (196). MS (EI) for C₃₀H₂₆FN₃O₆, found 544 (MH⁺).

[000654] Example 72: 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidin-3-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (197)



[000655] 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidin-3-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (197): Compound **197** was synthesized from Compound **195** in a manner analogous to the preparation of Compound **188** from Compound **181** in Example 66 (55% yield). MS (EI) for C₃₀H₂₇FN₄O₅, found 543 (MH⁺).

[000656] Example 73: 1-N-[4-[6-(Azetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (198)




[000657] 1-N-[4-[6-(Azetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (198): To a mixture of Compound **43** (150 mg, 0.27 mmol), azetidine (300 mg, 5.25 mmol), and K_3PO_4 (212 mg, 1.0 mmol) in DMA (2 mL) was added bis(tri-*tert*-butylphosphine)palladium(0). The resulting mixture was stirred at 100 °C for 12 h. After cooling, water was added, and the mixture was extracted with EtOAc, washed with aq saturated NaCl and concentrated. The crude product was purified by prep HPLC to give Compound **198**. MS (EI) for $C_{30}H_{27}FN_4O_4$, found 527 (MH⁺).

[000658] The following compounds were prepared from Compound **43** in a method analogous to the synthesis of Compound **198** from Compound **43** in Example 73:

[000659] 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (199): Azetidin-3-ol was used in place of azetidine. MS (EI) for $C_{30}H_{27}FN_4O_5$, found 543 (MH⁺).

[000660] 1-N-[4-[6-(3,3-Difluoroazetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (200): The HCl salt of 3,3-difluoroazetidine was used in place of azetidine. MS (EI) for $C_{30}H_{25}F_3N_4O_4$, found 563 (MH⁺).

[000661] Example 74: 1-N'-(4-Fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-3-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide (201)



[000662] 1-N'-(4-Fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-3-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide (201): In a microwave reaction tube were mixed Compound **43** (50 mg, 0.09 mmol), 3-pyridinylboronic acid (16 mg, 0.13 mmol), Na_2CO_3 (28 mg, 0.27 mmol), Bis(di-*tert*-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II) (6 mg, 0.009 mmol), 1,4-dioxane (1.5

mL), and water (0.3 mL). The reaction mixture was irradiated in a microwave reactor for 10 min at 150 °C. After cooling, the mixture was extracted with EtOAc, washed with aq saturated NaCl, and concentrated. The crude product was purified by prep HPLC to give Compound **201**. MS (EI) for $C_{32}H_{25}FN_4O_4$, found 549 (MH⁺).


[000663] The following compounds prepared from Compound **43** in a manner analogous to the method used to prepare Compound **201** from Compound **43** in Example 74:

[000664] **1-N'-(4-Fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-4-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide (202)**: Pyridin-4-ylboronic acid was used in place of 3-pyridinylboronic acid. MS (EI) for $C_{32}H_{25}FN_4O_4$, found 549 (MH⁺).

[000665] The following compound was prepared from Compound **44** and the potassium salt of 5-(trifluoro-4-boranyl)-1H-pyrazole in a manner analogous to the preparation of Compound **201** from Compound **43** and 3-pyridinylboronic acid in Example 74:

[000666] **1-N'-(4-Fluorophenyl)-1-N-[4-[6-(1H-pyrazol-5-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (204)**: MS (EI) for $C_{29}H_{22}FN_5O_3$, found 508 (MH⁺).

[000667] **Example 75: 1-N'-(4-Fluorophenyl)-1-N-[4-(7-methoxy-6-sulfamoylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide (206)**



[000668] **S-(4-(4-(1-((4-Fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)-7-methoxyquinolin-6-yl) ethanethioate (205)**: To a mixture of Compound **43** (200 mg, 0.36 mmol), potassium ethanethioate (83 mg, 0.72 mmol), and DIEA (0.5 mL) in 1,4-dioxane (4 mL) was added $Pd_2(dba)_3$ (16 mg, 0.017 mmol) and Xantphos (20 mg, 0.035 mmol). The resulting mixture was stirred at 120 °C under microwave irradiation for

15 min. After cooling, EtOAc and water were added. Any insoluble material was filtered. The filtrate was washed with aq saturated NaCl and concentrated. The crude product was purified by flash silica gel chromatography to give Compound **205** (37% yield plus 50% of Compound **43** recovered).

[000669] 1-N'-(4-Fluorophenyl)-1-N-[4-(7-methoxy-6-sulfamoylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide (206): N-Chlorosuccinimide (80 mg, 0.6 mmol) was added to a mixture of 2N HCl (0.5 mL) and MeCN (3 mL), and the resulting mixture was cooled to 0 °C. A solution of Compound **205** (80 mg, 0.15 mmol) in MeCN (1.5 mL) was added. After stirring at 0 °C for 2 h, the reaction mixture was extracted with EtOAc. The organic phase was washed with aq saturated NaCl and concentrated. The resulting crude sulfonyl chloride was dissolved in *i*-PrOH (2 mL), to which was added excess NH₄OH at room temperature. After the reaction was complete, the mixture was concentrated, and the crude product was purified by prep HPLC to give Compound **206** (60% yield). MS (EI) for C₂₇H₂₃FN₄O₆S, found 551 (MH⁺).

[000670] The following compounds were made from Compound **205** by a manner analogous to the way Compound **206** was made from Compound **205** in the second step of Example 75:


[000671] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(methylsulfamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (207): The NH₄OH was replaced by methylamine. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.22 (s, 1H), 10.06 (s, 1H), 8.71 (d, 1H), 8.69 (s, 1H), 7.78 (d, 2H), 7.64-7.62 (m, 2H), 7.60 (s, 1H), 7.30-7.28 (m, 3H), 7.17-7.12 (m, 2H), 6.50 (d, 1H), 4.04 (s, 3H), 2.47 (s, 3H), 1.47 (s, 4H); MS (EI) for C₂₈H₂₅FN₄O₆S, found 565.1 (MH⁺)

[000672] 1-N-[4-[6-(Ethylsulfamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (208): The NH₄OH was replaced by ethylamine. MS (EI) for C₂₉H₂₇FN₄O₆S, found 579 (MH⁺).

[000673] The following compound was made from Compound **44** in a manner analogous to the way Compound **206** was made from Compound **43** in Example 75:


[000674] 1-N'-(4-Fluorophenyl)-1-N-[4-(6-sulfamoylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide (209). MS (EI) for C₂₆H₂₁FN₄O₅S, found 521 (MH⁺).

[000675] Example 76: 1-N'-(4-Fluorophenyl)-1-N-[4-(7-methoxy-6-methylsulfonylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide (210)



[000676] 1-N'-(4-Fluorophenyl)-1-N-[4-(7-methoxy-6-methylsulfonylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide (210): Compound **43** (100 mg, 0.18 mmol), sodium methanesulfinate (40 mg, 0.36 mmol), CuI (5 mg), proline (6 mg), and NaOH (3 mg) were mixed in DMSO (1 mL). The resulting mixture was stirred at 110 °C for 14 h. After cooling to room temperature, water was added. The mixture was filtered through Celite® and washed with EtOAc. The EtOAc filtrate was concentrated and purified by prep HPLC to give Compound **210** (8% yield). MS (EI) for C₂₈H₂₄FN₃O₆S, found 550 (MH⁺).

[000677] Example 77: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(methoxycarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (213)



[000678] 4-Chloro-N-methoxyquinoline-7-carboxamide (212): To a mixture of Compound **211** (60 mg, 289.00 μmol, 1 eq), HATU (180.00 mg, 473.40 μmol, 1.64 eq) and DIEA (150.00 mg, 1.16 mmol, 202.16 μL, 4.02 eq) in DMF (5 mL) was added O-methylhydroxylamine hydrochloride (54.00 mg, 646.57 μmol, 49.09 uL, 2.24 eq) in one portion. The mixture was stirred at 20-30 °C for 0.5 h and then partitioned between water (10 mL) and EtOAc (2 x 10 mL). The organic extracts were separated, washed with aq saturated NaCl (10 mL), dried with anhyd Na₂SO₄, and concentrated to give Compound **212** (140 mg, crude), which was used for the next step directly without further purification. MS (EI) for C₁₁H₉ClN₂O₂, found 236.9 (MH⁺).


[000679] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(methoxycarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (213): To a mixture of Compound **212** (140 mg, 591.58 μmol, 1 eq) and Compound **3** (280.00 mg, 890.84 μmol, 1.51 eq) in DMSO (5 mL)

was added t-BuOK (133 mg, 1.19 mmol, 2.00 eq) in one portion, and the resulting mixture was stirred at 100 °C for 1 h. The reaction mixture was partitioned between water (10 mL) and EtOAc (2 x 10 mL). The combined organic extracts were washed with aq saturated NaCl (10 mL), dried with anhyd Na₂SO₄, and concentrated. The residue was purified by prep HPLC (basic conditions; column: Waters Xbridge 150*25 5u; mobile phase: [water (0.05% ammonia hydroxide v/v)-ACN]; B%: 32%-57%, 6.5 min) to give Compound **213** (7.9 mg, 2.54% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.11 (s, 1H), 10.22 (s, 1H), 10.06 (s, 1H), 8.77 (d, 1H), 8.44 - 8.37 (m, 2H), 7.99 (d, 1H), 7.79 (d, 2H), 7.65 (dd, 2H), 7.28 (d, 2H), 7.16 (t, 2H), 6.66 (d, 1H), 3.78 (s, 3H), 1.48 (s, 4H); MS (EI) for C₂₈H₂₃FN₄O₅, found 515.1 (MH⁺).

[000680] The following compound was prepared from Compound **211** in a manner analogous to the method used to prepare Compound **213** from Compound **211** in Example 77:

[000681] **1-N-[4-[7-(Ethylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (214):** The O-methylhydroxylamine hydrochloride in the first step was replaced with ethylamine hydrochloride. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.21 (s, 1H), 10.06 (s, 1H), 8.84 (t, 1H), 8.75 (d, 1H), 8.53 (d, 1H), 8.38 (d, 1H), 8.07 (dd, 1H), 7.78 (d, 2H), 7.64 (dd, 2H), 7.28 (d, 2H), 7.15 (t, 2H), 6.63 (d, 1H), 3.31 (br s, 2H), 1.47 (s, 4H), 1.18 (t, 3H); MS (EI) for C₂₉H₂₅FN₄O₄, found 513.1 (MH⁺).

[000682] **Example 78: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3,4-oxadiazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (220)**



[000683] Methyl 7-methoxy-4-(4-nitrophenoxy)quinoline-6-carboxylate (215):

Compound **4** (300 mg, 1.2 mmol), 4-nitrophenol (200 mg, 1.42 mmol), and DIEA (230 mg, 1.78 mmol) were mixed in toluene (1 mL). The resulting mixture was stirred at 110 °C for 12 h. After cooling, water was added, and the mixture was extracted with EtOAc. The organic phase was washed with aq saturated NaCl, concentrated and purified by flash silica gel chromatography to give Compound **215** (150 mg, 36%). MS (EI) for C₁₈H₁₄N₂O₆, found: 355 (MH⁺).

[000684] 7-Methoxy-4-(4-nitrophenoxy)quinoline-6-carboxylic acid (216): Compound **216** was synthesized in a manner analogous to the preparation of Compound **81** in Example 25. MS (EI) for C₁₇H₁₂N₂O₆, found: 341 (MH⁺).

[000685] 7-Methoxy-4-(4-nitrophenoxy)quinoline-6-carbohydrazide (217): To a solution of Compound **216** (45 mg, 0.13 mmol) and DIEA (51 mg, 0.4 mmol) in DMF (2 mL) was added HATU (100 mg, 0.26 mmol). The resulting mixture was stirred at room temperature for 10 min, then excess NH₂NH₂·H₂O was added. After the reaction was complete, water was added. The precipitate was filtered and dried under vacuum to give crude Compound **217**, which was used in the next step without further purification. MS (EI) for C₁₇H₁₄N₄O₅, found: 355 (MH⁺).

[000686] 2-(7-Methoxy-4-(4-nitrophenoxy)quinolin-6-yl)-1,3,4-oxadiazole (218):

Compound **217** (66 mg, 0.18 mmol) and *p*-TsOH·H₂O (4 mg, 0.02) were mixed in triethyl orthoformate (1 mL). The resulting mixture was stirred at 120 °C for 5 h. After cooling, the mixture was concentrated, and aq saturated NaHCO₃ was added. The precipitate was filtered and washed with EtOAc to give crude Compound **218**, which was used in the next step without further purification. MS (EI) for C₁₈H₁₂N₄O₅, found: 365 (MH⁺).


[000687] 4-((7-Methoxy-6-(1,3,4-oxadiazol-2-yl)quinolin-4-yl)oxy)aniline (219): Crude Compound **218** was mixed with Fe powder (50 mg, 0.9 mmol) and NH₄Cl (20 mg, 0.36 mmol) in EtOH (2 mL). The resulting mixture was stirred at 85 °C for 3 h. After cooling, the mixture was filtered through Celite® and washed with EtOAc, and the filtrate was concentrated. The residue was purified by flash silica gel chromatography to give Compound **219** (40 mg, 66% over two steps). MS (EI) for C₁₈H₁₄N₄O₃, found: 335 (MH⁺).

[000688] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3,4-oxadiazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (220): Compound **220** was synthesized in a manner analogous to the preparation of Compound **175** in Example 64. MS (EI) for C₂₉H₂₂FN₅O₅, found: 540 (MH⁺).

[000689] The following compound was made in a manner analogous to the synthesis of Compound **220** in Example 78, replacing Compound **4** with Compound **55**:

[000690] 1-N'-(4-Fluorophenyl)-1-N-[4-[6-(1,3,4-oxadiazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (221): MS (EI) for C₂₈H₂₀FN₅O₄, found 510 (MH⁺).


[000691] Example 81: N-(4-((7-Bromoquinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (232)



[000692] N-(4-((7-Bromoquinolin-4-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (232): To a mixture of Compound **231** (1 g, 4.13 mmol) and Compound **3** (1.3 g, 4.13 mmol) in DMF (8 mL) was added Cs₂CO₃ (2.7 g, 8.26 mmol). The resulting

mixture was stirred at 85 °C for 12 h. The mixture was cooled to room temperature, diluted with water, and filtered. The residue was purified by flash column chromatography to give Compound **232** (1.5 g, 69% yield). MS (EI) for C₂₆H₁₉BrFN₃O₃, found 520, 522 (MH⁺)


[000693] Example 85: 1-N'-(4-Fluorophenyl)-1-N-[4-(7-sulfamoylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide (254)



[000694] S-(4-(4-(1-((4-Fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinolin-7-yl) ethanethioate (253): Compound **253** was synthesized from Compound **232** in a manner analogous to the preparation of Compound **205** from Compound **43** in Example 75 (170 mg, crude). MS (EI) for C₂₈H₂₂FN₃O₄S, found: 516.0 (MH⁺).

[000695] 1-N'-(4-Fluorophenyl)-1-N-[4-(7-sulfamoylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide (254): Compound **254** was synthesized from Compound **253** in a manner analogous to the preparation of Compound **206** from Compound **205** in Example 75 (170 mg, crude). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.28 (s, 1H), 10.05 (s, 1H), 8.95 (d, 1H), 8.62 (d, 1H), 8.52 (d, 1H), 8.11 (dd, Hz, 1H), 7.88 - 7.75 (m, 4H), 7.67 - 7.59 (m, 2H), 7.33 (d, 2H), 7.15 (t, 2H), 6.85 (d, 1H), 1.48 (s, 4H); MS (EI) for C₂₆H₂₁FN₄O₅S, found: 521.0 (MH⁺).

[000696] Example 86: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-[(E)-N-methoxy-C-methylcarbonimidoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (256)




[000697] 1-N-[4-(7-Acetylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (255): Compound **232** (100 mg, 192.18 μ mol, 1 eq), tributyl(1-ethoxyvinyl)stannane (138.81 mg, 384.36 μ mol, 129.73 μ L, 2 eq), and Pd(PPh₃)₂Cl₂ (13.49 mg, 19.22 μ mol, 0.1 eq) in toluene (2 mL) were degassed with nitrogen for 10 min and heated at 110 °C for 12 h under nitrogen. After cooling, KF (2 mL, 2M) was added, and the resulting mixture was stirred at room temperature for 30 min. Aq 6 M HCl (2 mL) was added to the residue, and the mixture was stirred at 25 °C for 30 min. The mixture was extracted with EtOAc. The organic layer was washed with water, aq saturated NaCl, dried (Na₂SO₄), and concentrated under reduced pressure. The residue was purified by prep-TLC (petroleum ether/ethyl acetate = 1/1), followed by re-crystallization from MeOH (2 mL) and lyophilization to give Compound **255** (12.4 mg, 10.63% yield). ¹H NMR (400 MHz, CDCl₃) δ 9.46 (s, 1H), 8.75 (d, 1H), 8.67 (s, 1H), 8.57 (br s, 1H), 8.43 (d, 1H), 8.16 (dd, 1H), 7.66 (d, 2H), 7.51 - 7.45 (m, 2H), 7.20 (d, 2H), 7.10 - 7.04 (m, 2H), 6.64 (d, 1H), 2.78 (s, 3H), 1.76 (s, 2H), 1.67 (s, 2H); MS (EI) for C₂₈H₂₂FN₃O₄, found: 484.2 (MH⁺).

[000698] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-[(E)-N-methoxy-C-methylcarbonimidoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (256): To a mixture of Compound **255** (50 mg, 103.41 μ mol, 1 eq) in EtOH (2 mL) and H₂O (0.3 mL) was added O-methylhydroxylamine (43.18 mg, 517.07 μ mol, 5 eq, HCl). The mixture was stirred at 50 °C for 1 h. The mixture was concentrated under vacuum, and the resulting residue was purified by column chromatography on silica-gel (petroleum ether/EtOAc=5/1 to 1/1), followed

by re-purification by re-crystallization from MeOH (2 mL) to give Compound **256** (27 mg, 48.90% yield). ^1H NMR (400 MHz, CDCl_3) δ 10.24 (s, 1H), 8.82 (s, 1H), 8.72 (br d, 1H), 8.48 - 8.40 (m, 2H), 8.22 (s, 1H), 7.80 (d, 2H), 7.50 - 7.42 (m, 2H), 7.23 (br d, 2H), 7.07 (t, 2H), 6.83 (br d, 1H), 4.11 (s, 3H), 2.40 (s, 3H), 1.89 - 1.81 (m, 2H), 1.68 - 1.64 (m, 2H); MS (EI) for $\text{C}_{29}\text{H}_{25}\text{FN}_4\text{O}_4$, found: 513.1 (MH^+).


[000699] Example 87: 4-(4-Amino-2-fluorophenoxy)-7-methoxy-N-methylquinoline-6-carboxamide (258)



[000700] 4-Chloro-7-methoxy-N-methylquinoline-6-carboxamide (257): Methylamine (8 M, 50 mL, 10.07 eq) in EtOH was added to a solution of Compound **4** (10 g, 39.74 mmol, 1 eq) in THF (150 mL) at $30\text{ }^\circ\text{C}$. The reaction mixture was stirred at $30\text{ }^\circ\text{C}$ for 25 h. The mixture was concentrated under vacuum. The residue was slurried with warm water (100 mL) and filtered. The filtered cake was dried under vacuum to give Compound **257** as a white solid (9 g, 90.35% yield). MS (EI) for $\text{C}_{12}\text{H}_{11}\text{ClN}_2\text{O}_2$, found: 251.0 (MH^+).

[000701] 4-(4-Amino-2-fluorophenoxy)-7-methoxy-N-methylquinoline-6-carboxamide (258): Compound **258** was synthesized from Compound **257** and 4-amino-2-fluorophenol in a manner analogous to the synthesis of Compound **171** from Compounds **42** and **2** in Example 62.

[000702] Example 88: 1-N-[3-Fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)-1-N'-methylcyclopropane-1,1-dicarboxamide (262)



[000703] Methyl 1-((4-fluorophenyl)(methyl)carbamoyl)cyclopropane-1-carboxylate (260): A solution of Compound **259** (215 mg, 1.5 mmol, 1.5 eq), HATU (568 mg, 149 μ mol, 1.5 eq) and DIEA (0.52 mL, 3 eq) in DMF (5 mL, 0.2M) was stirred at room temp for 15 min. 4-Fluoro-N-methylaniline (124 mg, 1.0 mmol, 1.0 eq) was added, and the mixture was stirred at room temp for 17 h. Water was added and the resulting mixture extracted with EtOAc. The two phases were separated and the organic layer dried over Na_2SO_4 , concentrated under reduced pressure, absorbed into silica gel, and purified by CombiFlash (60:40 Hexanes:EtOAc) to give Compound **260** (221 mg, 88.7% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.38-7.35 (m, 2H), 7.25 (m, 2H), 3.40 (s, 3H), 3.18 (s, 3H), 1.37 (d, 2H), 1.17 (bs, 2H).

[000704] 1-((4-Fluorophenyl)(methyl)carbamoyl)cyclopropane-1-carboxylic acid (261): A solution of Compound **260** (60 mg, 0.23 mmol, 1.0 eq) and NaOH (20 mg, 0.5 mmol) in MeOH and THF (1 mL each) was stirred at 50 °C overnight. The reaction mixture was cooled down and concentrated under reduced pressure. To the residue, 1N HCl was added, and the resulting mixture was extracted with DCM. The two phases were separated, and the organic layer was dried over Na_2SO_4 and concentrated under reduced pressure to give crude Compound **261** which was used for the next step without further purification (44.7 mg, 79% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.39 (br s, 2H), 7.25 (m, 2H), 3.17 (s, 3H), 1.24 (br s, 2H), 0.99 (br s, 2H).

[000705] 1-N-[3-Fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)-1-N'-methylcyclopropane-1,1-dicarboxamide (262): A solution of

Compound **261** (23.7 mg, 0.1 mmol, 1.5 eq), HATU (38 mg, 0.1 mmol, 1.5 eq) and DIEA (34 μ L, 3.0 eq) in DMF (0.4 mL) was stirred at room temperature for 15 min. Compound **258** (23.7 mg, 0.06 mmol, 1.0 eq) was added, and the mixture was stirred at room temp for 17 h. Water was added to the mixture, and the resulting solid was extracted into EtOAc. The two phases were separated, and the organic layer was dried over Na₂SO₄, concentrated under reduced pressure, absorbed onto silica gel, and purified by CombiFlash (5:95 MeOH:DCM). The resulting product was further purified by prep HPLC to give Compound **262** (7 mg, 18.8% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.94 (s, 1H), 8.79 (d, 1H), 8.59 (s, 1H), 8.38 (d, 1H), 7.50 (s, 1H), 7.37 (t, 2H), 7.22 (b m, 3H), 7.04 (t, 2H) 6.65 (d, 1H), 3.99 (s, 3H), 3.18 (s, 3H), 2.78 (d, 3H), 1.37 (d, 2H), 1.17 (b s, 2H); MS (EI) for C₃₀H₂₆F₂N₄O₅, found 561.0 (MH⁺).

[000706] The following Compound **263** was made using a variation of the method used in the synthesis of Compound **262** in Example 88:

[000707] **1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-methylcyclopropane-1,1-dicarboxamide (263)**: Compound **258** was replaced by 4-(4-aminophenoxy)-7-methoxy-N-methylquinoline-6-carboxamide in the last step of the 3-step sequence of Example 88. The 4-(4-aminophenoxy)-7-methoxy-N-methylquinoline-6-carboxamide was synthesized in the same manner as Compound **258** in Example 87, replacing the 4-amino-2-fluorophenol in the second step with 4-aminophenol. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.69 (br s, 1H), 8.67 (d, 1H), 8.62 (s, 1H), 8.38 (d, 1H), 7.52 (s, 1H), 7.47 (s, 1H), 7.29 (m, 2H), 7.18 (d, 3H), 7.11 (t, 2H) 6.47 (d, 1H), 4.03 (s, 3H), 2.85 (d, 3H), 2.08 (s, 3H), 1.37 (d, 2H), 1.24 (br s, 2H); MS (EI) for C₃₀H₂₇FN₄O₅, found 543.0 (MH⁺).

[000708] The following compounds were made from Compound **259** and Compound **258** following the same 3-step procedure used to synthesize Compound **262** from Compound **259** and Compound **258** in Example 88:

[000709] **1-N'-(2-Chloro-4-fluorophenyl)-1-N-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (264)**: The 4-fluoro-N-methylaniline in the first step was replaced with 2-chloro-4-fluoroaniline. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.36 (s, 1H), 10.25 (s, 1H), 8.61 (d, 1H), 8.54 (s, 1H), 8.31 (m, 1H), 7.84-7.79 (m, 2H), 7.47, (s, 2H), 7.40 (t, 2H) 7.18 (t, 1H), 6.42 (d, 1H), 3.96 (s, 3H), 2.77 (d, 3H), 1.56 (br s, 2H), 1.53 (br s, 2H); MS (EI) for C₂₉H₂₃ClF₂N₄O₅, found 581.0 (MH⁺).

[000710] 1-N-[3-Fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluoro-2-methylphenyl)cyclopropane-1,1-dicarboxamide (265): The 4-fluoro-N-methylaniline in the first step was replaced with 4-fluoro-2-methylaniline. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.58 (s, 1H), 9.72 (s, 1H), 8.68 (d, 1H), 8.61 (s, 1H), 8.39 (d, 1H), 7.92 (d, 1H), 7.54 (br s, 2H), 7.47 (t, 2H), 7.1 (d, 1H), 7.03 (t, 1H) 6.49 (d, 1H), 4.03 (s, 3H), 2.85 (d, 3H), 2.22 (s, 3H), 1.54 (s, 4H); MS (EI) for C₃₀H₂₆F₂N₄O₅, found 561.0 (MH⁺).

[000711] 1-N'-(4-Fluoro-2,6-dimethylphenyl)-1-N-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (267): The 4-fluoro-N-methylaniline in the first step was replaced with 4-fluoro-2,6-dimethylaniline. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.8 (s, 1H), 9.19 (s, 1H), 8.6 (d, 1H), 8.54 (s, 1H), 8.32 (d, 1H), 7.87 (d, 1H), 7.46- 7.36 (m, 3H), 6.88 (d, 2H), 6.41 (d, 1H), 3.96 (s, 3H), 2.77 (d, 3H), 2.09 (s, 6H), 1.47 (s, 4H); MS (EI) for C₃₁H₂₈F₂N₄O₅, found 575.0 (MH⁺).


[000712] 1-N-[3-Fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluoro-2-methoxyphenyl)cyclopropane-1,1-dicarboxamide (268): The 4-fluoro-N-methylaniline in the first step was replaced with 4-fluoro-2-methoxyaniline. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.2 (s, 1H), 10.17 (s, 1H), 8.68 (d, 1H), 8.62 (s, 1H), 8.39 (d, 1H), 7.98-7.96 (m, 1H), 7.86 (d, 1H), 7.54-7.47 (m, 3H), 7.01 (d, 1H), 6.77 (t, 1H) 6.51 (d, 1H), 4.04 (s, 3H), 3.85 (s, 3H), 2.86 (d, 3H), 1.62 (s, 2H), 1.58 (s, 2H); MS (EI) for C₃₀H₂₆F₂N₄O₆, found 577.0 (MH⁺).

[000713] 1-N-[3-Fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluoro-2-propan-2-yloxyphenyl)cyclopropane-1,1-dicarboxamide (269): The 4-fluoro-N-methylaniline in the first step was replaced with 4-fluoro-2-isopropoxyaniline. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.21 (s, 1H), 10.06 (s, 1H), 8.62 (d, 1H), 8.55 (s, 1H), 8.32 (d, 1H), 8.05 (t, 1H), 7.87 (d, 1H), 7.52-7.41 (m, 3H), 6.97 (d, 1H), 6.68 (t, 1H), 6.41 (d, 1H), 4.64-4.58 (m, 1H), 3.96 (s, 3H), 2.78 (d, 3H), 1.51 (s, 2H), 1.40 (s, 2H), 1.17 (s, 3H), 1.16 (s, 3H); MS (EI) for C₃₂H₃₀F₂N₄O₆, found 605 (MH⁺)

[000714] 1-N'-(2-Cyclopropyl-4-fluorophenyl)-1-N-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (270): The 4-fluoro-N-methylaniline in the first step was replaced with 2-cyclopropyl-4-fluoroaniline. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.53 (br s, 1H), 9.99 (br s, 1H), 8.68 (d, 1H), 8.62 (s, 1H), 8.39 (d, 1H), 7.91 (d, 1H), 7.61 (s, 1H), 7.54 (s, 2H), 7.49 (br s, 1H), 7.01 (t, 1H) 6.82 (d, 1H), 6.49

(d, 1H), 4.04 (s, 3H), 2.85 (d, 3H), 1.97 (br s, 1H), 1.58 (s, 4H), 0.94 (d, 2H), 0.67 (d, 2H); MS (EI) for $C_{32}H_{28}F_2N_4O_5$, found 587.4 (MH⁺).


[000715] The 2-cyclopropyl-4-fluoroaniline used in the synthesis of Compound **270** was synthesized by the following procedure:



[000716] *Tert*-Butyl (2-bromo-4-fluorophenyl)carbamate (289 mg, 1 mmol, 1.0 eq), cyclopropyl MIDA boronate (236 mg, 1.2 mmol, 1.2 eq) and K_3PO_4 (636 mg, 3.0 eq) was added to toluene:water (4:1) in a thick wall reaction tube and degassed with nitrogen for 5 min. Palladium acetate (9 mg, 4 mol percent) and 2-(dicyclohexyl)phosphino biphenyl (28 mg 8 mole percent) were added to the mixture and degassed with nitrogen for another 5 min. The tube was capped, and the mixture was heated under nitrogen at 100 °C overnight. After cooling to room temperature, the two phases were separated. The organic layer was dried over anhyd Na_2SO_4 , concentrated under reduced, absorbed into silica gel, and purified by CombiFlash using 70:30 Hexanes:DCM as gradient to give *tert*-butyl (2-cyclopropyl-4-fluorophenyl)carbamate as a colorless oil (90 mg, 36% yield).

[000717] A solution of *tert*-butyl (2-cyclopropyl-4-fluorophenyl)carbamate (86 mg, 0.3 mmol, 1.0 eq) and TFA (77mg, 2.0 eq) in DCM (1 mL) was stirred at 40 °C overnight. The reaction solution was cooled down and concentrated under reduced pressure. To the residue, DCM (4 mL) was added, and the resulting solution was washed the with aq saturated $NaHCO_3$, dried over anhyd Na_2SO_4 , and concentrated under reduced pressure to give crude 2-cyclopropyl-4-fluoroaniline as a nearly colorless oil (43 mg, 81% yield) which used in subsequent reactions without further purification. 1H NMR (400 MHz, $DMSO-d_6$) δ 7.16-7.13 (m, 1H), 7.01 (t, 1H), 6.85 (d, 1H), 1.91-1.85 (m, 1H), 1.0 (d, 2H), 0.73 (d, 2H).

[000718] **Example 89: 1-((3-Fluoro-4-((7-methoxy-6-(methylcarbamoyl)quinolin-4-yl)oxy)phenyl)-carbamoyl)-cyclopropane-1-carboxylic acid (272)**




[000719] Methyl 1-((3-fluoro-4-((7-methoxy-6-(methylcarbamoyl)quinolin-4-yl)oxy)phenyl)-carbamoyl)cyclopropane-1-carboxylate (271): A solution of Compound **259** (108 mg, 0.7 mmol, 1.5 eq), HATU (285 mg, 0.75 mmol, 1.5 eq), and DIEA (0.26 mL, 3 eq) in DMF (2.5 mL, 0.2M) was stirred at room temperature for 15 min. Compound **258** (170 mg, 0.5 mmol, 1.0 eq) was added, and the mixture was stirred at room temperature for 17 h. Water was added, and the resulting mixture extracted with EtOAc. The organic layer was separated and washed with aq saturated NaCl, dried over anhyd Na_2SO_4 , and concentrated under reduced pressure. The resulting residue was absorbed onto silica gel and purified by CombiFlash using 1:8 Hexanes:EtOAc followed by 1:6 Hexanes:DCM gradient. The resulting product was sonicated with Hexanes. The resulting solid was filtered, dissolved in DCM, and washed with aq 10% LiCl solution. Organic layer was dried over anhyd Na_2SO_4 and concentrated to give Compound **271** (90 mg, 38% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.64 (br s, 1H), 8.67 (d, 1H), 8.61 (s, 1H), 8.39 (d, 1H), 7.88 (m, 1H), 7.54 (s, 1H), 7.51-7.44 (m, 2H), 6.52 (d, 1H), 4.03 (s, 3H), 3.71 (s, 3H), 2.85 (d, 3H), 1.46-1.45 (m, 2H), 1.43-1.42 (m, 2H).

[000720] 1-((3-Fluoro-4-((7-methoxy-6-(methylcarbamoyl)quinolin-4-yl)oxy)phenyl)carbamoyl)-cyclopropane-1-carboxylic acid (272): A solution of Compound **271** (70 mg, 0.2 mmol, 1.0 eq) and NaOH (12 mg, 0.5 mmol) in MeOH and THF (1 mL each) was stirred at 50 °C overnight. The reaction solution was allowed to cool down and then concentrated under reduced pressure. To the residue, aq 1N HCl was added, and the resulting mixture was extracted with DCM. The organic phase was dried over anhyd Na_2SO_4 and

concentrated under reduced pressure to give crude Compound **272** (60 mg, 89% yield), which was used in subsequent reactions without further purification. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.85 (br s, 1H), 8.87 (d, 1H), 8.67 (s, 1H), 8.46 (d, 1H), 7.95 (m, 1H), 7.62 (s, 1H), 7.54-7.5 (m, 2H), 6.83 (d, 1H), 4.06 (s, 3H), 2.85 (d, 3H), 1.43 (s, 4H).

[000721] Example 90: 1-N'-[3-Fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-[4-(trifluoromethyl)phenyl]cyclopropane-1,1-dicarboxamide (273)




[000722] 1-N'-[3-Fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-[4-(trifluoromethyl)phenyl]cyclopropane-1,1-dicarboxamide (273): Compound **273** was synthesized from Compound **272** and 4-(trifluoromethyl)aniline using standard HATU amide bond forming techniques such as those used in Example 3, Example 4 and the first step of Example 89. ^1H NMR (400 MHz, CD_3OD) δ 8.90 (s, 1H), 8.86 (d, 1H), 7.89 (d, 1H), 7.55 (s, 1H), 7.51 (d, 2H), 7.43 (m, 2H), 7.25 (d, 2H), 6.98 (d, 1H), 4.13 (s, 3H), 2.94 (s, 3H), 1.64 - 1.54 (m, 4H). MS (EI) for $\text{C}_{30}\text{H}_{24}\text{F}_4\text{N}_4\text{O}_5$ found 597.2 (MH $^+$).

The following compound was made from Compound **272** following the same procedure used to synthesize Compound **273** from Compound **272** Example 90:

[000723] 1-N-(4-Chlorophenyl)-1-N'-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (274): The 4-(trifluoromethyl)aniline was replaced with 4-chloroaniline. ^1H NMR (400 MHz, CD_3OD) δ 8.99 (s, 1H), 8.95 (d, 1H), 7.98 (d, 1H), 7.83 (d, 2H), 7.65 (d, 3H), 7.54 (q, 2H), 7.06 (d, 1H), 4.22 (s, 3H), 3.03 (s, 3H), 1.84 - 1.64 (m, 4H). MS (EI) for $\text{C}_{29}\text{H}_{24}\text{ClFN}_4\text{O}_5$ found 563.2 (MH $^+$).

[000724] Example 91: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-[(E)-methoxyiminomethyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (278)




[000725] (4-Chloroquinolin-7-yl)methanol (275): To a solution of LiAlH₄ (116.45 mg, 3.07 mmol, 1.7 eq) in THF (5 mL) was added Compound **161** (400 mg, 1.80 mmol, 1 eq) in THF (2 mL) at 0 °C. The mixture was stirred at 20 °C for 3 h. Na₂SO₄·10H₂O (50 mg) was added, the resulting mixture was filtered, and the filtrate was dried over anhyd Na₂SO₄ and then concentrated in vacuo. The resulting residue was purified by flash silica gel chromatography (ISCO®; 4 g SepaFlash® Silica Flash Column, Eluent of 70~100% Ethyl acetate/Petroleum ether gradient @ 25 mL/min) to give Compound **275** as a light yellow solid (150 mg, 36.5% yield). ¹H NMR (400 MHz, CDCl₃) δ 8.74 (d, 1H), 8.19 (d, 1H), 8.12 (s, 1H), 7.64 (dd, 1H), 7.47 (d, 1H), 4.94 (s, 2H), 2.99 (br s, 1H).

[000726] N-(4-Fluorophenyl)-N-(4-((7-(hydroxymethyl)quinolin-4-yl)oxy)phenyl)cyclopropane-1,1-dicarboxamide (276): Compound **276** was synthesized from Compound **275** and Compound **3** in a manner analogous to the method used to synthesize Compound **43** from Compound **42** and Compound **3** in the last step of Example 12, lowering the reaction temperature to 85 °C. ¹H NMR (400 MHz, CD₃OD) δ 8.63 (d, 1H), 8.41 (d, 1H), 8.03 (s, 1H), 7.76 (d, 2H), 7.68 (dd, 1H), 7.62 - 7.55 (m, 2H), 7.30 - 7.23 (m, 2H), 7.09 (t, 2H), 6.66 (d, 1H), 4.88 (s, 2H), 1.66 (d, 4H); MS (EI) for C₂₇H₂₂FN₃O₄, found 472.1 (MH⁺).

[000727] N-(4-Fluorophenyl)-N-(4-((7-formylquinolin-4-yl)oxy)phenyl)cyclopropane-1,1-dicarboxamide (277): To a solution of Compound **276** (100 mg, 212.1 μmol, 1 eq) in DCM (5 mL) was added MnO₂ (239.72 mg, 2.76 mmol, 13 eq). The mixture was stirred at 20 °C for 1 h. The mixture was filtered and concentrated in vacuo to give Compound **277** as a yellow oil (50 mg, 40.2% yield). MS (EI) for C₂₇H₂₀FN₃O₄, found 470.0 (MH⁺).

[000728] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-[(E)-methoxyiminomethyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (278): To a mixture of Compound **277** (50 mg, 106.5 μmol , 1 eq) in EtOH (3 mL) and water (0.3 mL) was added O-methylhydroxylamine (44.47 mg, 532.5 μmol , 5 eq, HCl). The mixture was stirred at 50 °C for 2 h. The mixture was concentrated under vacuum, and EtOH (3 mL) was added to the residue. The resulting white solid was collected and then purified by prep-HPLC (YMC-Actus Triart C18 150*30mm*5 μm ; mobile phase: [water(0.225%FA)-ACN]; B%: 37%-77%, 10min) to give Compound **278** as a white solid (14.5 mg, 26.2% yield). ^1H NMR (400 MHz, DMSO- d_6) δ 10.22 (s, 1H), 10.07 (s, 1H), 8.72 (d, 1H), 8.51 (s, 1H), 8.34 (d, 1H), 8.18 (d, 1H), 7.97 (dd, 1H), 7.79 (d, 2H), 7.70 - 7.61 (m, 2H), 7.28 (d, 2H), 7.16 (t, 2H), 6.60 (d, 1H), 3.98 (s, 3H), 1.48 (s, 4H); MS (EI) for $\text{C}_{28}\text{H}_{23}\text{FN}_4\text{O}_4$, found 499.2 (MH $^+$).


[000729] Example 92: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(methylcarbamoylamino)-quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (279)



[000730] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-methoxy-6-(methylcarbamoylamino)-quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (279): To a mixture of Compound **43** (60 mg, 109.02 μmol , 1 eq) in dioxane (3 mL) was added 1-methylurea (48.46 mg, 654.10 μmol , 6 eq), $\text{Pd}_2(\text{dba})_3$ (14.97 mg, 16.35 μmol , 0.15 eq), Xantphos (18.92 mg, 32.70 μmol , 0.3 eq), and Cs_2CO_3 (99.46 mg, 305.25 μmol , 2.8 eq) under an atmosphere of nitrogen. The mixture was stirred at 100 °C for 1 h. The reaction mixture was concentrated, and water (20 mL) was added to the residue. The resulting mixture was extracted with EtOAc (3 x 25 mL). The combined organic extracts were dried over anhyd Na_2SO_4 and concentrated in vacuo, and the resulting residue was purified by prep-HPLC (column: DuraShell 150*25mm*5 μm ; mobile phase: [water (0.05% ammonia hydroxide v/v)-ACN]; B%: 30%-70%, 10min) to give Compound **279** as a white solid (12.8 mg, 20.7% yield). ^1H NMR (400 MHz, DMSO- d_6) δ 10.15 (s, 1H), 10.07 (s, 1H), 8.97 (s, 1H), 8.44 (d, 1H), 8.32 (s, 1H), 7.76 (d, 2H), 7.65 (dd, 2H),


7.40 (s, 1H), 7.24-7.12 (m, 4H), 6.95 (br d, 1H), 6.41 (d, 1H), 4.04 (s, 3H), 2.68 (s, 3H), 1.48 (br d, 4H); MS (EI) for $C_{29}H_{26}FN_5O_5$, found 544.3 (MH⁺).

[000731] Example 93: Methyl N-[4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-methoxyquinolin-6-yl]carbamate (280)



[000732] Methyl N-[4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropane-carbonyl]amino]phenoxy]-7-methoxyquinolin-6-yl]carbamate (280): Compound **280** was synthesized from Compound **43** in a manner analogous to the method used to synthesize Compound **279** from Compound **43** in Example 92, replacing the 1-methylurea with methyl carbamate. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.17 (s, 1H), 10.06 (s, 1H), 8.84 (s, 1H), 8.67 (s, 1H), 8.52 (d, 1H), 7.77 (d, 2H), 7.70-7.62 (m, 2H), 7.45 (s, 1H), 7.23 (d, 2H), 7.19-7.12 (m, 2H), 6.45 (d, 1H), 4.00 (s, 3H), 3.72 (s, 3H), 1.48 (s, 4H); MS (EI) for $C_{29}H_{25}FN_4O_6$, found 567.3 [M+Na]⁺


[000733] Example 94: 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(methylcarbamoylamino)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (281)



[000734] 1-N'-(4-Fluorophenyl)-1-N-[4-[7-(methylcarbamoylamino)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (281): A mixture of Compound **232** (100 mg, 192.18 μmol, 1 eq), methylurea (42.71 mg, 576.54 μmol, 3 eq), Xantphos (22.24 mg, 38.44 μmol, 0.2 eq), t-BuONa (36.94 mg, 384.36 μmol, 2 eq), and Pd₂(dba)₃ (17.60 mg, 19.22 μmol, 0.1 eq) in dioxane (5 mL) was degassed and purged with nitrogen three times, followed by stirring at 100 °C for 16 h. The reaction mixture was filtered and concentrated under reduced pressure. The resulting residue was purified by prep-HPLC (column: Xtimate C18 10μ 250 mm


*50mm; mobile phase: [water (0.05% ammonia hydroxide v/v)-ACN]; B%: 39%-69%, 9min) to give Compound **281** as a white solid (56.9 mg, 57.4% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.18 (s, 1H), 10.06 (s, 1H), 9.09 (s, 1H), 8.55 (d, 1H), 8.14 (d, 2H), 7.75 (d, 2H), 7.69-7.57 (m, 3H), 7.22 (d, 2H), 7.15 (t, 2H), 6.37 (d, 1H), 6.26 (d, 1H), 2.69 (d, 3H), 1.48 (s, 4H); MS (EI) for C₂₈H₂₄FN₅O₄, found 514.3 (MH⁺).

[000735] Example 95: Methyl N-[4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]-amino]phenoxy]quinolin-7-yl]carbamate (282)



[000736] Methyl N-[4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]-amino]phenoxy]quinolin-7-yl]carbamate (282): Compound **282** was synthesized from Compound **232** in a manner analogous to the method used to synthesize Compound **281** from Compound **232** in Example 94, replacing the 1-methylurea with methyl carbamate. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.82-10.72 (m, 1H), 10.40-10.33 (m, 1H), 10.12-10.04 (m, 1H), 8.89 (d, 1H), 8.59 (d, 1H), 8.50 (d, 1H), 7.93-7.81 (m, 3H), 7.69-7.60 (m, 2H), 7.38 (d, 2H), 7.16 (t, 2H), 6.78 (d, 1H), 3.79 (s, 3H), 1.50 (s, 4H); MS (EI) for C₂₈H₂₃FN₄O₅, found 515.3 (MH⁺).


[000737] Example 96: 1-N-[4-[6-(3-Ethyl-1,2,4-oxadiazol-5-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide hydrochloride (283)




[000738] 1-N-[4-[6-(3-Ethyl-1,2,4-oxadiazol-5-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide hydrochloride (283): To a solution of Compound **6** (30 mg, 0.058 mmol), (Z)-N'-hydroxypropionimidamide (7.22 mg, 0.058 mmol), and DIEA (44 μL, 0.17 mmol) in DMF (0.5 mL) was added HATU (26.45 mg, 0.070 mmol),

and the reaction was stirred at room temperature until the starting material disappeared as determined by monitoring with LC-MS. The reaction mixture was then heated to 60 °C for 2 h. Aq saturated NaHCO₃ was added to the resulting mixture, which was then extracted with DCM (3x). The combined organic extracts were washed with aq saturated NaCl, dried over Na₂SO₄, and concentrated. The resulting crude residue was subjected to HPLC purification (Gemini-NX, 10uM, 250x30 mm, C18 column; eluent: 0.1 to 100% acetonitrile in water, both eluents containing 0.1% trifluoroacetic acid, gradient elution over 15 min) and subsequently freeze-dried. The resulting powder was brought up in 20% MeOH in DCM, passed through an Agilent PL-HCO₃ ion exchange column (or brought up in DCM and washed with saturated sodium bicarbonate), and concentrated under reduced pressure. HCl (4M in Dioxane, 0.5 ml; or 1M aq, 1 mL) was added, the volatile solvents were removed under reduced pressure, and the resulting residue dried under high vacuum to give the hydrochloride salt of Compound **283** (4.1 mg, 11.7% yield). ¹H NMR (400 MHz, CD₃OD) δ 9.29 (s, 1H), 8.94 (d, 1H), 7.86 (q, 2H), 7.68 (s, 1H), 7.61 – 7.54 (m, 2H), 7.41 (d, 2H), 7.10 (t, 2H), 7.02 (d, 1H), 4.26 (s, 3H), 2.91 (q, 2H), 1.68 (s, 4H), 1.42 (t, 3H). MS (EI) for C₃₁H₂₆FN₅O₅ found 568.2 (MH⁺).

[000739] Example 97: 1-N'-[3-Fluoro-4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (289)





[000740] Methyl 4-hydroxy-6-methylquinoline-7-carboxylate (284): A mixture of Compound **139** (290 mg, 1.11 mmol, 1 *eq*) and AcONa (454.26 mg, 5.54 mmol, 5 *eq*) in AcOH (6 mL) and water (10 mL) was stirred at 90 °C for 48 h. After cooling to ambient temperature, the reaction mixture was neutralized with aq saturated NaHCO₃ and extracted with EtOAc (3 x 30 mL). The combined extracts were dried over anhyd Na₂SO₄ and concentrated under reduced pressure to give Compound **284** as a light yellow solid (200 mg, 74.8% yield) which was used directly in subsequent reactions without further purification. MS (EI) for C₁₂H₁₁NO₃, found 218.1 (MH⁺).

[000741] Methyl 4-(2-fluoro-4-nitrophenoxy)-6-methylquinoline-7-carboxylate (285): To a mixture of Compound **284** (200 mg, 828.65 μmol, 1 *eq*) and 1,2-difluoro-4-nitro-benzene (197.75 mg, 1.24 mmol, 137.32 μL, 1.5 *eq*) in ACN (15 mL) was added Cs₂CO₃ (809.97 mg, 2.49 mmol, 3 *eq*) at 25 °C, followed by stirring at 70 °C for 2 h. After being allowed to cool, the reaction mixture was diluted with water (50 mL) and extracted with EtOAc (3 x 30 mL). The combined organic extracts were washed with aq saturated NaCl (60 mL), dried over anhyd Na₂SO₄, and concentrated. The resulting residue was purified by flash chromatography on silica gel eluting with EtOAc in petroleum ether (0~60%) to give Compound **285** as a light yellow solid (120 mg, 38.6% yield). ¹H NMR (400 MHz, CDCl₃) δ 8.77 (d, 1H), 8.70 (s, 1H), 8.23-8.15 (m, 2H), 8.12 (s, 1H), 7.42-7.35 (m, 1H), 6.65 (d, 1H), 3.99 (s, 3H), 2.79 (s, 3H); MS (EI) for C₁₈H₁₃FN₂O₅, found 357.0 (MH⁺).

[000742] Methyl 4-(4-amino-2-fluorophenoxy)-6-methylquinoline-7-carboxylate (286): Compound **286** was synthesized from Compound **285** using a method analogous to that used to convert Compound **218** to Compound **219** in Example 78. MS (EI) for C₁₈H₁₅FN₂O₃, found 326.9 (MH⁺).

[000743] Methyl 4-(2-fluoro-4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)-6-methylquinoline-7-carboxylate (287): A mixture of Compound **1** (51.30 mg, 218.34 μmol, 1.5 *eq*) in SOCl₂ (8.20 g, 68.92 mmol, 5.00 mL, 473.51 *eq*) was stirred

for 1 h at 65 °C. The mixture was concentrated, and the residue was dissolved in toluene and reconcentrated. To the resulting residue was added a solution of Compound **286** (50 mg, 145.56 μmol , 1.0 *eq*) in THF (5 mL), and then was added triethylamine (44.19 mg, 436.69 μmol , 60.78 μL , 3 *eq*). The resulting mixture was stirred for 1 h at 25 °C. The reaction mixture was quenched with water (30 mL) and extracted with EtOAc (3 x 20 mL). The combined extracts were dried over anhyd Na_2SO_4 and concentrated. The resulting residue was purified by flash chromatography on silica gel eluting with (EtOAc in petroleum ether = 0~70%) to give Compound **287** as an off-white solid (70 mg, 85.95% yield). MS (EI) for $\text{C}_{29}\text{H}_{23}\text{F}_2\text{N}_3\text{O}_5$, found 532.0 (MH^+).

[000744] 4-(2-Fluoro-4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)-6-methylquinoline-7-carboxylic acid (288): A mixture of Compound **287** (70.0 mg, 125.12 μmol , 1 *eq*) and $\text{LiOH}\cdot\text{H}_2\text{O}$ (26.25 mg, 625.58 μmol , 5 *eq*) in THF (5 mL) and water (5 mL) was stirred for 4 h at 25 °C. The reaction mixture was acidified with aq 1 M HCl until a pH of ~5 was achieved. The mixture was then diluted with water (30 mL) and extracted with EtOAc (3 x 20 mL). The combined organic extracts were dried over anhyd Na_2SO_4 and concentrated to give Compound **288** as a white solid (70 mg, 86.49% yield) which was used directly in subsequent reactions without further purification. MS (EI) for $\text{C}_{28}\text{H}_{21}\text{F}_2\text{N}_3\text{O}_5$, found 518.1 (MH^+).

[000745] 1-1-N'-[3-Fluoro-4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (289): Compound **289** was synthesized from Compound **288** in a manner analogous to the preparation of Compound **8** in Example 4. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.39 (s, 1H), 9.99 (s, 1H), 8.67 (d, 1H), 8.48 (d, 1H), 8.16 (s, 1H), 7.95 (s, 1H), 7.91 (d, 1H), 7.63 (dd, 2H), 7.56-7.49 (m, 1H), 7.47-7.41 (m, 1H), 7.15 (t, 2H), 6.59 (d, 1H), 2.82 (d, 3H), 2.54 (s, 3H), 1.47 (d, 4H); MS (EI) for $\text{C}_{29}\text{H}_{24}\text{F}_2\text{N}_4\text{O}_4$, found 531.1 (MH^+).

The following compounds were made from Compound **139** using the same 6 step process used to synthesize Compound **289** in Example 97:


[000746] 1-N'-[2,5-Difluoro-4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (290): The 1,2-difluoro-4-nitrobenzene in Step 2 was replaced with 1,2,4-trifluoro-5-nitrobenzene. ^1H NMR (400 MHz, CDCl_3) δ 10.09 (br s, 1H), 8.59 (d, 1H), 8.28 (dd, 1H), 8.21 (s, 1H), 8.09 (s, 1H), 8.00 (s, 1H),

7.36-7.43 (m, 2H), 6.96-7.06 (m, 3H), 6.47 (d, 1H), 5.98 (d, 1H), 3.01 (d, 3H), 2.58 (s, 3H), 1.72-1.78 (m, 2H), 1.59-1.62 (m, 2H); MS (EI) for $C_{29}H_{23}F_3N_4O_4$, found 549.1 (MH⁺).

[000747] 1-N'-[2-Chloro-5-fluoro-4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (291): The 1,2-difluoro-4-nitrobenzene in Step 2 was replaced with 1-chloro-4,5-difluoro-2-nitrobenzene. ¹H NMR (400 MHz, CDCl₃) δ 10.08 (s, 1H), 8.66 (d, 1H), 8.61 (s, 1H), 8.44 (d, 1H), 8.15 (s, 1H), 8.07 (s, 1H), 7.52-7.45 (m, 2H), 7.36 (d, 1H), 7.06 (t, 2H), 6.54 (d, 1H), 6.15-6.07 (m, 1H), 3.08 (d, 3H), 2.65 (s, 3H), 1.82-1.72 (m, 4H); MS (EI) for $C_{29}H_{23}ClF_2N_4O_4$, found 565.1 (MH⁺).

[000748] 1-N-(4-Fluorophenyl)-1-N'-[2,3,5-trifluoro-4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide (292): The 1,2-difluoro-4-nitrobenzene in Step 2 was replaced with 1,2,3,4-tetrafluoro-5-nitrobenzene. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.33 (s, 1H), 9.77 (s, 1H), 8.71 (d, 1H), 8.54-8.48 (m, 1H), 8.18 (s, 1H), 8.08-8.02 (m, 1H), 7.98 (s, 1H), 7.63-7.56 (m, 2H), 7.23-7.14 (m, 2H), 6.89 (d, 1H), 2.83 (d, 3H), 2.55 (s, 3H), 1.72-1.65 (m, 2H), 1.62-1.57 (m, 2H); MS (EI) for $C_{29}H_{22}F_4N_4O_4$, found 567.1 (MH⁺).

[000749] Example 98: 1-N'-[4-(7-Carbamoyl-6-methylquinolin-4-yl)oxy-3-fluorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (293)




[000750] 1-N'-[4-(7-Carbamoyl-6-methylquinolin-4-yl)oxy-3-fluorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (293): Compound **293** was synthesized from Compound **288** in a manner analogous to the preparation of Compound **7** in Example 3. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.40 (s, 1H), 10.00 (s, 1H), 8.67 (d, 1H), 8.16 (s, 1H), 8.03 (s, 1H), 7.99 (s, 1H), 7.91 (d, 1H), 7.64 (dd, 2H), 7.58 (s, 1H), 7.56-7.51 (m, 1H), 7.47-7.41 (m, 1H), 7.15 (t, 2H), 6.59 (d, 1H), 2.58 (s, 3H), 1.47 (d, 4H); MS (EI) for $C_{28}H_{22}F_2N_4O_4$, found 517.1 (MH⁺).

[000751] The following compound was made using a similar process to that used to synthesize Compound **293** from Compound **288** in Example 98:

[000752] **1-N'-[4-(7-Carbamoyl-6-methylquinolin-4-yl)oxy-2,5-difluorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (294):** Compound **288** was replaced with 4-(2,5-difluoro-4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)-6-methylquinoline-7-carboxylic acid, which was made using the same multistep procedure used to synthesize Compound **288** in Example 97, replacing the 1,2-difluoro-4-nitrobenzene in Step 2 with 1,2,4-trifluoro-5-nitrobenzene. ¹H NMR (400 MHz, CDCl₃) δ 10.19 (br s, 1H), 8.67 (d, 1H), 8.36 (dd, 1H), 8.26 (br s, 1H), 8.19 (s, 2H), 7.47 (dd, 2H), 7.03-7.12 (m, 3H), 6.55 (d, 1H), 6.04 (br s, 1H), 5.78 (br s, 1H), 2.70 (s, 3H), 1.75-1.90 (m, 2H), 1.67-1.69 (m, 2H); MS (EI) for C₂₈H₂₁F₃N₄O₄, found 535.1 (MH⁺).

[000753] **Example 99: 1-N'-[3-Fluoro-4-[7-(2-hydroxyethoxycarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (301)**



[000754] Methyl 4-hydroxyquinoline-7-carboxylate (296): To a round-bottom flask was added Compound **295** (5 g, 22.32 mmol, 1 eq) and TEA (22.58 g, 223.16 mmol, 31.06 mL, 10 eq) in MeOH (60 mL) and DMSO (20 mL). DPPP (1.84 g, 4.46 mmol, 0.2 eq) and Pd(OAc)₂ (1.00 g, 4.46 mmol, 0.2 eq) were then added. The flask was purged with carbon monoxide twice and left under 30 psi of carbon monoxide while heated at 70 °C for 12 h. MeOH (100 mL) was added, the resulting suspension was filtered through a pad of Celite, and the filter cake was washed with MeOH (3 x 50 mL). The combined filtrates were concentrated, and the resulting crude product was washed with water (50 mL), triturated with MeOH (15 mL), and filtered to give Compound **296** as a yellow solid (3.5 g, 77.19% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.98 (br s, 1H), 8.23-8.15 (m, 2H), 8.01 (dd, 1H), 7.81 (br d, 1H), 6.11 (d, 1H), 3.90 (s, 3H); MS (EI) for C₁₁H₉NO₃, found 203.9 (MH⁺).

[000755] Methyl 4-(2-fluoro-4-nitrophenoxy)quinoline-7-carboxylate (297): Compound **297** was made from Compound **296** in a manner analogous to the way Compound **285** was made from Compound **284** in Example 97. ¹H NMR (400 MHz, CDCl₃) δ 8.88-8.83 (m, 2H), 8.37 (d, 1H), 8.25-8.16 (m, 3H), 7.43 (t, 1H), 6.70 (d, 1H), 4.04 (s, 3H).; MS (EI) for C₁₇H₁₁FN₂O₅, found 343.2 (MH⁺).

[000756] Methyl 4-(4-amino-2-fluorophenoxy)quinoline-7-carboxylate (298): Compound **298** was synthesized from Compound **297** using a method analogous to that used to convert Compound **218** to Compound **219** in Example 78. MS (EI) for C₁₇H₁₃FN₂O₃, found 312.9 (MH⁺).

[000757] Methyl 4-(2-fluoro-4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)phenoxy)quinoline-7-carboxylate (299): Compound **1** (428.83 mg, 1.92 mmol, 3 eq) was suspended in anhyd DCM (10 mL) at 25 °C. DMF (4.68 mg, 64.04 μmol, 4.93 μL, 0.1 eq) was added with stirring under nitrogen, followed by (COCl)₂ (290.00 mg, 2.28 mmol, 0.2 mL, 3.57 eq). The mixture was stirred at 25 °C for 30 min. Toluene (5.0 mL) was added, and the solvent was removed under reduced pressure. The resulting acyl chloride product was dissolved in anhyd THF (1.0 mL). A solution of Compound **298** (200 mg, 640.42 μmol, 1 eq) in DMA (6 mL) was added the above prepared acyl chloride in THF with stirring under nitrogen. The reaction was stirred at 25 °C for 0.5 h. The reaction mixture was poured into aq saturated NaHCO₃ (100 mL) and extracted with DCM (3 x 50 mL). The combined organic extracts were washed with aq saturated NaHCO₃ (15 mL) and then aq saturated NaCl (15 mL), dried with

anhyd Na₂SO₄, and concentrated in vacuum to give Compound **299** as a yellow solid (270 mg, 81.47% yield) which was used in subsequent reactions without further purification. MS (EI) for C₂₈H₂₁F₂N₃O₅, found 518.1 (MH⁺).

[000758] 4-(2-Fluoro-4-(1-((4-fluorophenyl)carbamoyl)cyclopropane-1-carboxamido)-phenoxy)quinoline-7-carboxylic acid (300): Compound **300** was made from Compound **299** in a manner analogous to the way Compound **288** was made from Compound **287** in Example 97. MS (EI) for C₂₇H₁₉F₂N₃O₅, found 504.1 (MH⁺).

[000759] 1-N'-[3-Fluoro-4-[7-(2-hydroxyethoxycarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (301): To a solution of Compound **300** (80 mg, 158.90 μ mol, 1 eq) in DMF (3 mL) was added HATU (72.50 mg, 190.68 μ mol, 1.2 eq) and DIEA (61.61 mg, 476.71 μ mol, 83.03 μ L, 3 eq), and the resulting mixture was stirred at 25 °C for 30 min. 2-aminooxyethanol (14.70 mg, 190.68 μ mol, 1.2 eq) was added, and the reaction mixture was stirred at 25 °C for another 2 h. The reaction mixture was poured into aq saturated NH₄Cl (50 mL) and extracted with DCM (3 x 30 mL). The combined organic extracts were concentrated under vacuum, and the resulting residue was purified by prep-HPLC (Column: Boston Prime C18 150*30mm *5 μ m, gradient: 43-63% of acetonitrile in water (0.05% NH₃H₂O), flow rate: 25 mL/min) to give Compound **301** as a white solid (46.8 mg, 52.36% yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.09 (br s, 1H), 10.41 (br s, 1H), 10.01 (br s, 1H), 8.78 (d, 1H), 8.45-8.40 (m, 2H), 8.01 (dd, 1H), 7.92 (dd, 1H), 7.68-7.60 (m, 2H), 7.56-7.51 (m, 1H), 7.50-7.44 (m, 1H), 7.20-7.11 (m, 2H), 6.68 (d, 1H), 4.80 (br s, 1H), 4.00 (t, 2H), 3.66 (t, 2H), 1.51-1.43 (m, 4H); MS (EI) for C₂₉H₂₄F₂N₄O₆, found 563.1 (MH⁺).

The following compounds were made from Compound **295** using the same 6 step process used to synthesize Compound **301** in Example 99:

[000760] 1-N'-[2,5-Difluoro-4-[7-(2-hydroxyethoxycarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (302): The 1,2-difluoro-4-nitrobenzene in Step 2 was replaced with 1,2,4-trifluoro-5-nitrobenzene. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.13 (br s, 1H), 11.17 (br s, 1H), 9.78 (br s, 1H), 8.81 (d, 1H), 8.46 (s, 1H), 8.40 (m, 1H), 8.19 (m, 1H), 8.04 (d, 1H), 7.73 (m, 1H), 7.61 (dd, 2H), 7.19 (t, 2H), 6.83 (d, 1H), 4.81 (m, 1H), 4.00 (m, 2H), 3.68 (m, 2H), 1.69 (m, 2H), 1.61 (m, 2H); MS (EI) for C₂₉H₂₃F₃N₄O₆, found 581.1 (MH⁺).

Biological Examples

[000761] Example A: Kinase Assays

[000762] Kinase activity and compound inhibition were investigated using the ^{33}P -Phosphoryl transfer radiometric kinase assay, performed using the KinaseProfilerTM service of Eurofins Pharma Discovery Services UK Limited. Dose-response experiments were performed using nine compound concentrations in a 96-well microtiter plate. For each assay, all compounds were prepared to a 50x final assay concentration (50 μM) in 100% DMSO, then diluted in a half-log series, with the final top concentration at 1 μM . This working stock of the compound was added to the assay well as the first component in the reaction, followed by the remaining components as detailed in the following assay protocols below. The positive control wells (100% kinase activity) contain all components of the reaction including 2% DMSO (control for solvent effects), except the compound of interest. Blank wells contain all components of the reaction, with the reference inhibitor, staurosporine. This reference compound was used to abolish kinase activity and generated the 0% kinase activity base-line. IC_{50} values were calculated by nonlinear regression analysis using the sigmoidal dose-response (variable slope) curve fit on XLFit version 5.3 (ID Business Solutions).

[000763] Example B: Human AXL Kinase Assay

[000764] Human Axl (residues H473-A894 with Q764R, 161nM) was incubated with 8 mM MOPS pH 7.0, 0.2 mM EDTA, 250 μM KKSRRGDYMTMQIG, 10 mM magnesium acetate, and 10 μM [γ - ^{33}P -ATP]. The reaction was initiated by the addition of the Mg/ATP mix. After incubation for 40 minutes at room temperature, the reaction was stopped by the addition of phosphoric acid to a concentration of 0.5%. A reaction aliquot of 10 μL was then spotted onto a P30 filtermat and washed four times for 4 minutes in 0.425% phosphoric acid and once in methanol prior to drying and scintillation counting. Incorporated ^{33}P was measured using the Wallac Microbeta scintillation counter (Perkin Elmer).

[000765] Example C: Human KDR Kinase Assay

[000766] Human KDR (residues K790-V1356, 55nM) was incubated with 8 mM MOPS pH 7.0, 0.2 mM EDTA, 0.33 mg/mL myelin basic protein, 10 mM magnesium acetate, and 10 μM [γ - ^{33}P -ATP]. The reaction was initiated by the addition of the Mg/ATP mix. After incubation for 40 minutes at room temperature, the reaction was stopped by the addition of phosphoric acid to a concentration of 0.5%. A reaction aliquot of 10 μL was then spotted onto a P30 filtermat

and washed four times for 4 minutes in 0.425% phosphoric acid and once in methanol prior to drying and scintillation counting. Incorporated ^{33}P was measured using the Wallac Microbeta scintillation counter (Perkin Elmer).

[000767] Example D: Human Mer Kinase Assay

[000768] Human Mer (residues R557-E882 with H628Q and R794A, 0.7nM) was incubated with 8 mM MOPS pH 7.0, 0.2 mM EDTA, 30 mM NaCl, 250 μM GGMEDIYFEFMGGKKK, 10 mM magnesium acetate and 10 μM [γ - ^{33}P -ATP]. The reaction was initiated by the addition of the Mg/ATP mix. After incubation for 40 minutes at room temperature, the reaction was stopped by the addition of phosphoric acid to a concentration of 0.5%. A reaction aliquot of 10 μL was then spotted onto a P30 filtermat and washed four times for 4 minutes in 0.425% phosphoric acid and once in methanol prior to drying and scintillation counting. Incorporated ^{33}P was measured using the Wallac Microbeta scintillation counter (Perkin Elmer).

[000769] Example E: Human Met Kinase Assay

[000770] Human Met (residues R974-S1390 with A1209G and V1290L, 3.4nM) was incubated with 8 mM MOPS pH 7.0, 0.2 mM EDTA, 250 μM KKKGQEEYVFIE, 1 mM sodium orthovanadate, 5 mM sodium-6-glycerophosphate, 10 mM magnesium acetate, and 10 μM [γ - ^{33}P -ATP]. The reaction was initiated by the addition of the Mg/ATP mix. After incubation for 40 minutes at room temperature, the reaction was stopped by the addition of phosphoric acid to a concentration of 0.5%. A reaction aliquot of 10 μL was then spotted onto a P30 filtermat and washed four times for 4 minutes in 0.425% phosphoric acid and once in methanol prior to drying and scintillation counting. Incorporated ^{33}P was measured using the Wallac Microbeta scintillation counter (Perkin Elmer).

[000771] Activity data obtained for the Example compounds using the kinase assays in Exmples A, B, D and E is provided in Table 2 (A: $\text{IC}_{50} \leq 10 \text{ nM}$; B: $10 \text{ nM} < \text{IC}_{50} \leq 100 \text{ nM}$; C: $100 \text{ nM} < \text{IC}_{50} \leq 1000 \text{ nM}$; D: $\text{IC}_{50} > 1000 \text{ nM}$).

[000772] Table 2: Activity data for selected compounds of the invention

Compound No.	Name	Axl IC_{50} (nM)	Mer IC_{50} (nM)	c-Met IC_{50} (nM)
5	methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylate	B	A	A

Compound No.	Name	Axl IC ₅₀ (nM)	Mer IC ₅₀ (nM)	c-Met IC ₅₀ (nM)
6	4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylic acid	C	B	A
7	1-N-[4-(6-carbamoyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A
8	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A
10	1-N-[4-[6-[2-(dimethylamino)ethylcarbamoyl]-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	B	A	A
11	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(2-piperidin-1-ylethylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A
12	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(2-morpholin-4-ylethylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A
13	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(oxetan-3-ylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	A
14	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(1-methylazetidin-3-yl)carbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	A
15	1-N-[4-[6-(azetidine-1-carbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	B	A	B
16	1-N'-(4-fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidine-1-carbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	A
21	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(2R)-pyrrolidin-2-yl]methylcarbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A
22	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(2S)-pyrrolidin-2-yl]methylcarbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A
37	1-N-[4-(6-cyano-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	B	A	B
45	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A

Compound No.	Name	Axl IC ₅₀ (nM)	Mer IC ₅₀ (nM)	c-Met IC ₅₀ (nM)
58	1-N-[4-(6-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	B	B	B
59	1-N'-(4-fluorophenyl)-1-N-[4-[6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	B	B
60	1-N'-(4-fluorophenyl)-1-N-[4-[6-[(1-methylazetidin-3-yl)carbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	B	B
67	1-N-[4-(6-carbamoyl-7-fluoroquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	B	B	B
68	1-N-[4-(6-carbamoyl-7-chloroquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	C	B	B
69	1-N-[4-(7-bromo-6-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	C	B	B
98	1-N-[4-[6-carbamoyl-7-(3-morpholin-4-ylpropoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A
103	1-N-[4-[6-carbamoyl-7-(2-methoxyethoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	B	A	A
128	methyl 4-[2-chloro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylate	A	A	A
130	4-[2-chloro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylic acid	B	B	B
132	1-N'-[4-(6-carbamoyl-7-methoxyquinolin-4-yl)oxy-3-chlorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A
134	1-N'-[3-chloro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A
129	methyl 4-[2-fluoro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylate	A	A	A
131	4-[2-fluoro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinoline-6-carboxylic acid	B	B	A
133	1-N'-[4-(6-carbamoyl-7-methoxyquinolin-4-yl)oxy-3-fluorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A

Compound No.	Name	Axl IC ₅₀ (nM)	Mer IC ₅₀ (nM)	c-Met IC ₅₀ (nM)
135	1-N'-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A
140	methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-methylquinoline-7-carboxylate	B	A	B
141	4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-methylquinoline-7-carboxylic acid	D	D	D
142	1-N-[4-(7-carbamoyl-6-methylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A
143	1-N'-(4-fluorophenyl)-1-N-[4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	A
150	methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-methoxyquinoline-7-carboxylate	B	A	A
151	4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-methoxyquinoline-7-carboxylic acid	C	B	B
152	1-N-[4-(7-carbamoyl-6-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	B	A	A
153	1-N'-(4-fluorophenyl)-1-N-[4-[6-methoxy-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	B
162	methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]quinoline-7-carboxylate	B	A	B
163	4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]quinoline-7-carboxylic acid	C	C	B
164	1-N-[4-(7-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A
165	1-N'-(4-fluorophenyl)-1-N-[4-[7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A
175	1-N-[4-[6-(3-cyano-2-fluorophenyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	B	A	B

Compound No.	Name	Axl IC ₅₀ (nM)	Mer IC ₅₀ (nM)	c-Met IC ₅₀ (nM)
176	1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-2-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	B
180	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(5-methylfuran-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	B
181	tert-butyl 2-[4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinolin-6-yl]pyrrole-1-carboxylate	C	B	B
182	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1-methylpyrazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	A
183	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1,2-oxazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	A
184	1-N-[4-[6-(3,5-dimethyl-1,2-oxazol-4-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	B	A	B
185	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrazol-5-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	A
186	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	A
187	1-N-[4-(6-cyclopropyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	B
188	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrrol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	A
195	tert-butyl 3-[4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-methoxyquinolin-6-yl]-3-hydroxyazetidine-1-carboxylate	B	A	B
196	1-N'-(4-fluorophenyl)-1-N-[4-[6-(3-hydroxyoxetan-3-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A
197	1-N'-(4-fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidin-3-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	A

Compound No.	Name	Axl IC ₅₀ (nM)	Mer IC ₅₀ (nM)	c-Met IC ₅₀ (nM)
198	1-N-[4-[6-(azetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	B	A	B
199	1-N'-(4-fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A
200	1-N-[4-[6-(3,3-difluoroazetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	B	A	B
201	1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-3-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A
202	1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-4-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide	B	A	A
206	1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-sulfamoylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A
210	1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-methylsulfonylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A
220	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3,4-oxadiazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide	A	A	A

[000773] Example F: AXL Autophosphorylation ELISA in A-172 Cells

[000774] A-172 glioblastoma cells (ATCC #CRL-1620) were seeded at 2.5×10^5 cells/well onto 24-well plates (Greiner #662165), in DMEM (Thermo Fisher #11995-040) containing 10% FBS (Thermo Fisher #26140-079), 1% MEM NEAA (Thermo Fisher #11140-050), 1% GlutaMax (Thermo Fisher #35050-061) and 1% Penicillin Streptomycin (Thermo Fisher #15140-122). A-172 cells were incubated at 37°C, 5% CO₂ for 24 h and then starved for 24 h in serum-free medium. Test compounds were serially diluted to produce an 8-point dose curve in fresh serum-free medium to a final concentration of 0.3% DMSO (vehicle) and added to the cells and incubated for 1 h. Cells were then stimulated with 1 µg/mL recombinant human Gas6 (R&D Systems #885-GSB-500) for 15 min, washed with cold PBS and immediately lysed with 150 µL of cold 1X lysis buffer [20 mM Tris, 137 mM sodium chloride, 2 mM EDTA, 10%

glycerol, 1% NP-40 alternative, 1 mM activated sodium orthovanadate, 1 mM PefaBloc SC (Sigma-Aldrich #11429868001), protease/phosphatase inhibitor tablet (Thermo Fisher #A32959)]. Lysates were collected and 100 μ L/well added into the human phospho-AXL DuoSet IC ELISA (R&D Systems #DYC2228-2). Assay was performed according to manufacturer's instructions and sample phospho-AXL concentrations were extrapolated using human phospho-AXL control (R&D Systems #841645) as a standard. Positive control wells (100% activity) contained Gas6-stimulated, DMSO-treated cell lysates. Negative control wells (0% activity) contained Gas6-stimulated, reference inhibitor-treated cell lysates. IC₅₀ values were calculated by nonlinear regression analysis using a 4-parameter logistic curve fit in ActivityBase XE (IDBS).

[000775] Example G: Met Autophosphorylation ELISA in PC-3 Cells

[000776] PC-3 prostate cancer cells (ATCC #CRL-1435) were seeded at 4×10^4 cells/well onto 24-well plates (Greiner #662165), in DMEM (Thermo Fisher #11995-040) containing 10% FBS (Thermo Fisher #26140-079), 1% MEM NEAA (Thermo Fisher #11140-050), 1% GlutaMax (Thermo Fisher #35050-061), and 1% Penicillin Streptomycin (Thermo Fisher #15140-122). PC-3 cells were incubated at 37°C, 5% CO₂ for 24 h and then starved for 3 h in serum-free medium. Test compounds were serially diluted to produce an 8-point dose curve in fresh serum-free medium to a final concentration of 0.3% DMSO (vehicle) and added to the cells and incubated for 1 h. Cells were then stimulated with 100 ng/mL recombinant human HGF (R&D Systems #294-HG-250) for 10 min, washed with cold PBS and immediately lysed with 130 μ L of cold 1X lysis buffer [20 mM Tris, 137 mM sodium chloride, 2 mM EDTA, 10% glycerol, 1% NP-40 alternative, 1 mM activated sodium orthovanadate, 1 mM PefaBloc SC (Sigma-Aldrich #11429868001), protease/phosphatase inhibitor tablet (Thermo Fisher #A32959)]. Lysates were clarified by centrifugation and 100 μ L/well added into the PathScan phospho-Met (panTyr) Sandwich ELISA (Cell Signaling Technology #7333). Assay was performed according to manufacturer's instructions. Positive control wells (100% activity) contained HGF-stimulated, DMSO-treated cell lysates. Negative control wells (0% activity) contained HGF-stimulated, reference inhibitor-treated cell lysates. IC₅₀ values were calculated by nonlinear regression analysis using a 4-parameter logistic curve fit in ActivityBase XE (IDBS).

[000777] Example H: KDR Autophosphorylation ELISA in HUVEC Cells

[000778] Human umbilical vein endothelial cells or HUVEC (Lonza #C2519A) were seeded at 2×10^4 cells/well onto 96-well plates (Corning #3904), in EGM-2 growth medium (Lonza #CC-3162) containing 1% Penicillin Streptomycin (Thermo Fisher #15140-122). HUVEC cells were incubated at 37°C, 5% CO₂ for 24 h and then starved for 24 h in serum-free EBM-2 basal medium (Lonza #CC-3156) containing 1% Penicillin Streptomycin. Test compounds were serially diluted to produce an 8-point dose curve in fresh serum-free medium to a final concentration of 0.3% DMSO (vehicle) and added to the cells and incubated for 1 h. Cells were then stimulated with 100 ng/mL recombinant human VEGF165 (R&D Systems #293-VE-500) for 5 min, washed with cold PBS, and immediately lysed with 130 µL of cold 1X lysis buffer [20 mM Tris, 137 mM sodium chloride, 2 mM EDTA, 10% glycerol, 1% NP-40 alternative, 1 mM activated sodium orthovanadate, 1 mM PefaBloc SC (Sigma-Aldrich #11429868001), protease/phosphatase inhibitor tablet (Thermo Fisher #A32959)]. Lysates were collected and 100 µL/well added into the human phospho-KDR DuoSet IC ELISA (R&D Systems #DYC1766-2). Assay was performed according to manufacturer's instructions and sample phospho-KDR concentrations were extrapolated using human phospho-KDR control (R&D Systems #841421) as a standard. Positive control wells (100% activity) contained VEGF165-stimulated, DMSO-treated cell lysates. Negative control wells (0% activity) contained non-stimulated cell lysates. IC₅₀ values were calculated by nonlinear regression analysis using a 4-parameter logistic curve fit in ActivityBase XE (IDBS).

[000779] Example I: Mer Autophosphorylation ELISA in Transient Transfected 293A Cells

[000780] 293A cells (Thermo Fisher #R70507) were seeded at 1.5×10^6 cells/well onto 100mm dish (Greiner #664169), in DMEM (Thermo Fisher #11995-040) containing 10% FBS (Thermo Fisher #26140-079), 1% MEM NEAA (Thermo Fisher #11140-050), 1% GlutaMax (Thermo Fisher #35050-061), and 1% Penicillin Streptomycin (Thermo Fisher #15140-122). 293A cells were incubated at 37°C, 5% CO₂ for 24 h and then transfected with 6 µg MERTK DNA (Genecopoeia #EX-Z8208-M02) using TransIT LT1 transfection reagent (Mirus-Bio #MIR2305). After 24 h incubation, the transfected 293A cells were seeded at 1×10^5 cells/well onto 96-well plates (Corning #3904) in DMEM growth medium overnight. Test compounds

were serially diluted to produce an 8-point dose curve in fresh serum-free medium to a final concentration of 0.3% DMSO (vehicle) and added to the cells and incubated for 1 h. Cells were then immediately lysed with 150 μ L of cold 1X lysis buffer [20 mM Tris, 137 mM sodium chloride, 2 mM EDTA, 10% glycerol, 1% NP-40 alternative, 1 mM activated sodium orthovanadate, 1 mM PefaBloc SC (Sigma-Aldrich #11429868001), protease/phosphatase inhibitor tablet (Thermo Fisher #A32959)]. Lysates were clarified by centrifugation and 50 μ L/well added into the human phospho-Mer DuoSet IC ELISA (R&D Systems #DYC2579-2). Assay was performed according to manufacturer's instructions and sample phospho-Mer concentrations were extrapolated using human phospho-Mer control (R&D Systems #841793) as a standard. Positive control wells (100% activity) contained DMSO-treated cell lysates. Negative control wells (0% activity) contained reference inhibitor-treated cell lysates. IC₅₀ values were calculated by nonlinear regression analysis using a 4-parameter logistic curve fit in ActivityBase XE (IDBS).

[000781] Compounds of the present disclosure, as exemplified herein, showed IC₅₀ values in the following ranges: A: IC₅₀ \leq 10 nM; B: 10 nM < IC₅₀ \leq 100 nM; C: 100 nM < IC₅₀ \leq 300 nM; D: IC₅₀ > 300 nM. "NT" means not tested

[000782] Activity data obtained for the Example compounds using cell based kinase assays in Exmples F, G, H and I is provided in Table 3.

[000783] Table 3: Cellular activity data for selected compounds of the invention

Compound No.	Axl IC ₅₀ (nM)	Mer IC ₅₀ (nM)	c-Met IC ₅₀ (nM)	KDR IC ₅₀ (nM)
5	B	NT	B	A
6	NT	NT	D	D
7	A	A	A	A
8	A	A	B	A
9	A	A	B	A
10	NT	NT	NT	NT
11	A	NT	B	A
12	A	NT	B	A
13	B	NT	B	A
14	A	NT	A	B
15	B	NT	C	D
16	B	NT	C	C
17	B	B	B	A
20	B	B	B	B
21	A	NT	B	B

Compound No.	Axl IC ₅₀ (nM)	Mer IC ₅₀ (nM)	c-Met IC ₅₀ (nM)	KDR IC ₅₀ (nM)
22	A	NT	B	B
26	B	B	B	A
27	A	B	B	A
30	C	D	C	B
31	B	D	C	B
32	A	A	B	A
34	A	A	B	A
35	C	NT	C	C
36	C	NT	C	D
37	B	NT	B	B
45	A	A	A	A
50	A	A	A	A
51	B	C	B	C
58	B	NT	C	C
59	C	NT	C	C
60	C	NT	C	D
67	C	NT	C	C
68	NT	NT	NT	NT
69	NT	NT	NT	NT
70	B	NT	B	B
71	B	B	B	B
81	D	NT	D	D
82	B	NT	B	B
83	B	NT	B	B
84	A	NT	B	A
87	C	NT	C	D
88	C	NT	C	D
89	D	NT	D	D
90	B	NT	B	B
92	B	NT	C	C
96	B	B	C	C
98	A	B	A	A
103	A	NT	B	A
106	A	NT	B	A
110	A	B	A	A
115	A	B	B	A
116	A	B	B	A
125	A	A	B	B
128	B	NT	B	A
129	A	NT	B	A
130	D	NT	D	D
131	D	NT	D	D

Compound No.	Axl IC ₅₀ (nM)	Mer IC ₅₀ (nM)	c-Met IC ₅₀ (nM)	KDR IC ₅₀ (nM)
132	A	A	A	A
133	A	A	A	A
134	A	A	A	A
135	A	A	A	A
140	B	NT	B	B
141	NT	NT	NT	NT
142	B	B	B	B
143	B	NT	B	C
150	B	NT	B	A
151	NT	NT	NT	D
152	B	NT	C	C
153	B	NT	C	B
162	D	NT	C	B
163	NT	NT	NT	NT
164	A	B	A	B
165	B	B	A	A
166	B	NT	B	C
167	C	NT	C	C
169	C	D	C	C
170	C	NT	D	C
175	B	NT	B	C
176	B	NT	B	B
177	A	NT	B	A
180	A	NT	B	B
181	NT	NT	NT	NT
182	A	A	B	A
183	B	NT	B	B
184	C	NT	B	D
185	B	A	B	A
186	A	NT	B	A
187	B	NT	B	B
188	A	NT	B	B
191	A	A	B	A
192	A	A	B	A
193	B	NT	C	C
195	C	NT	B	B
196	B	NT	B	B
197	C	NT	C	D
198	B	NT	B	B
199	A	NT	A	A
200	B	NT	B	B
201	B	NT	B	B

Compound No.	Axl IC ₅₀ (nM)	Mer IC ₅₀ (nM)	c-Met IC ₅₀ (nM)	KDR IC ₅₀ (nM)
202	B	NT	B	B
204	B	NT	B	B
206	A	B	A	A
207	B	B	B	B
208	B	B	B	B
209	C	NT	C	B
210	B	NT	B	A
213	B	B	B	C
214	B	B	B	B
220	A	A	B	A
221	B	NT	C	C
254	C	NT	C	D
255	D	NT	C	D
256	C	NT	C	C
262	D	NT	D	D
263	D	NT	D	D
264	B	B	B	B
265	A	B	B	B
267	B	NT	C	C
268	A	B	B	A
269	B	B	C	B
270	B	B	B	B
273	C	NT	C	B
274	B	A	B	A
278	C	C	C	C
279	A	A	B	A
280	A	A	B	A
281	C	NT	B	C
282	C	NT	C	D
283	B	B	B	B
289	B	B	B	B
290	A	A	A	C
291	B	B	B	D
292	B	B	B	D
293	B	B	B	B
294	A	A	A	B
301	B	B	B	B
302	B	C	A	C

[000784] Example J: Pharmacokinetic studies

[000785] Pharmacokinetic properties of select compounds were assessed in male Sprague-Dawley rats.

[000786] The non-GLP study was designed to investigate the pharmacokinetics of chosen compounds in plasma following an intravenous or oral dose administration to male Sprague Dawley rats.

[000787] Two groups of male Sprague-Dawley rats (three animals per group) received either an intravenous or oral (gavage) dose of compound at target dose levels of 3 mg/kg. Animals were observed for any clinically relevant abnormalities during dosing and at each sample collection period.

[000788] Animals in the PO group were fasted overnight prior to dose administration. Food was returned following the collection of the 4-hour blood sample. Water was not withheld.

[000789] Immediately prior to dosing, the body weight of each animal was recorded. Doses (rounded to the nearest 0.001 mL) were calculated based on the pretreatment body weight (kg) and a dose volume of 2.5 mL/kg for intravenous administration and 5 mL/kg for oral administration. Intravenous formulations were administered via a jugular vein cannula. Immediately after dosing, the cannula was flushed with saline and the line was tied off. The oral dose was administered via a ball-tipped feeding needle. Dosing syringe volumes for administration were second-person verified prior to dosing and that volume along with the results for the concentration verification analysis were used to calculate the actual dose administered. Dosing syringes were weighed immediately prior to and immediately after dosing each animal as a gravimetric check.

[000790] Serial blood samples (approximately 200 μ L per sample) were collected from each animal at 0.083 (IV dosing only), 0.25, 0.5, 1, 2, 4, 6 (PO dosing only), 8, 24, 32, 48, and 72 hours after dosing. Blood samples were collected into tubes containing K₂EDTA via the non-dosing jugular-vein cannula (JVC), which was flushed with an approximately equal volume of saline following each collection.

[000791] Blood samples were stored on wet ice until processed to plasma by centrifugation (3500 rpm at 5°C for 10 minutes) within 1 hour of collection. Plasma samples were transferred into matrix tubes and then stored in a -80°C freezer.

[000792] Plasma samples and dose formulation samples were analyzed for the compounds of interest using liquid chromatography-tandem mass spectrometry (LC-MS/MS) methods. Pharmacokinetic parameter estimates were calculated from the individual animal plasma concentration-time data using the actual dose based on the analysis of the dosing formulations,

nominal sampling times (all collections were within an acceptable range of target), and non-compartmental methods. The concentration-time data were analyzed to fit either an intravenous – bolus (IV) plasma analysis model (201) or extra-vascular (PO) dosing plasma analysis model (200) using the software WinNonlin Phoenix version 6.3 (Pharsight). The single-dose pharmacokinetic parameters assessed include, as appropriate: C_{\max} (observed peak or maximum concentration); T_{\max} (observed time of peak concentration); $T_{1/2}$ (terminal half-life); V_z (volume of distribution based on the terminal phase); V_{ss} (volume of distribution at steady state); AUC_{INF} (area under the concentration-time curve computed from time zero to infinity); AUC_{last} (area under the concentration-time curve computed from time zero to the time of the last quantifiable concentration); C_0 (back-extrapolated concentration at time zero); CL (total body clearance); V_z/F (volume of distribution for extravascular administration based on the terminal phase); CL/F (total body clearance for extravascular administration); $F\%$ (bioavailability); and MRT_{last} (mean residence time).

[000793] Areas-under-the-plasma concentration-time curves (AUC) were estimated using the linear-log trapezoidal rule. The area through the time (T_{last}) of the last observable concentration (C_{last}) is reported as AUC_{last} . AUC extrapolated to infinity, (AUC_{INF}) was estimated by adding AUC_{last} and the ratio of $C_{\text{last}}/\lambda_z$, where λ_z is the terminal rate constant. Apparent terminal half-life ($T_{1/2}$) was calculated as $\ln(2)/\lambda_z$ and determined using the slope of the log-linear terminal phase of the concentration-time curve, defined by a minimum of three plasma concentration-time points. Half-lives are reported if the correlation for the regression line, as measured by r squared, is ≥ 0.9 when rounded. After IV administration, volume of distribution (V_z) was calculated as $\text{Dose}/\lambda_z * AUC_{\text{INF-obs}}$, clearance (CL) was calculated as $\text{Dose}/AUC_{\text{INF-obs}}$ and volume of distribution at steady state (V_{ss}) was estimated as $MRT_{\text{INF}} * CL$. Mean residence time (MRT) from the time of dosing to the time of the last measurable concentration was calculated as $AUMC_{\text{last}}/AUC_{\text{last}}$. For model 200 the bioavailability (i.e. fraction of total dose that reaches the systemic circulation) cannot be calculated. Consequently, volume and clearance for this model is V_z/F or CL/F , respectively; where F is defined as bioavailability (i.e. fraction of total dose that reaches the systemic circulation; $(\text{Average } AUC_{\text{last-po}}/\text{Average } AUC_{\text{last-iv}}) * [\text{Dose}_{\text{IV}}/\text{Dose}_{\text{PO}}] * 100$).

[000794] Pharmacokinetic data produced by the procedure above for some compounds of the invention are provided in Table 4 below.

[000795] Table 4: Rat Pharmacokinetic data for selected compounds of the invention

Compound No.	Rat PK Parameters
7	IV (1 mg/kg) $t_{1/2} = 3.1 \text{ hr}$ $Cl = 89 \text{ mL/hr/kg}$ PO (3 mg/kg) $t_{1/2} = 7.3 \text{ hr}$ $C_{max} = 8.8 \text{ uM}$ $F = 72\%$
8	IV (3 mg/kg) $t_{1/2} = 5.4 \text{ hr}$ $Cl = 43 \text{ mL/hr/kg}$ PO (3 mg/kg) $t_{1/2} = 7.1 \text{ hr}$ $C_{max} = 11.4 \text{ uM}$ $F = 62\%$
9	IV (3 mg/kg) $t_{1/2} = 2.8 \text{ hr}$ $Cl = 152 \text{ mL/hr/kg}$ PO (3 mg/kg) $t_{1/2} = 2.6 \text{ hr}$ $C_{max} = 3.9 \text{ uM}$ $F = 69\%$
17	IV (3.0 mg/kg) $t_{1/2} = 3.0 \text{ hr}$ $Cl = 102 \text{ mL/hr/kg}$ PO (3.2 mg/kg) $t_{1/2} = 4.5 \text{ hr}$ $C_{max} = 3.4 \text{ uM}$ $F = 46\%$
45	IV (1 mg/kg) $t_{1/2} = 10.1 \text{ hr}$ $Cl = 41 \text{ mL/hr/kg}$ PO (3 mg/kg) $t_{1/2} = 5.7 \text{ hr}$ $C_{max} = 15.3 \text{ uM}$ $F = 133\%$

191	IV (3.0 mg/kg) t _{1/2} = 9.9 hr Cl = 63 mL/hr/kg PO (3.1 mg/kg) t _{1/2} = 6.8 hr C _{max} = 4.6 uM F = 38%
206	IV (3 mg/kg) t _{1/2} = 2.1 hr Cl = 511 mL/hr/kg PO (3 mg/kg) t _{1/2} = 2.9 hr C _{max} = 0.7 uM F = 37%


Other Embodiments

[000796] The foregoing disclosure has been described in some detail by way of illustration and example, for purposes of clarity and understanding. The invention has been described with reference to various specific and preferred embodiments and techniques. However, it should be understood that many variations and modifications can be made while remaining within the spirit and scope of the invention. It will be obvious to one of skill in the art that changes and modifications can be practiced within the scope of the appended claims. Therefore, it is to be understood that the above description is intended to be illustrative and not restrictive.

[000797] The scope of the invention should, therefore, be determined not with reference to the above description, but should instead be determined with reference to the following appended claims, along with the full scope of equivalents to which such claims are entitled.

Claims

1. A compound of formula A:



A

or a pharmaceutically acceptable salt thereof, wherein:

(i) R_1 is selected from the group consisting of (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, -CN, -NHOH, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)NR^aR^a, C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -S(O)NR^aR^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and S(O)₂NR^aR^a; and

R_2 is selected from -H, halo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NO₂, -OR^a, -SR^a, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)R^a, -OC(O)NR^aR^a, -NHR^a, -NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and -S(O)₂NR^aR^a, wherein the (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R_1 or R_2 are each optionally substituted with 1, 2, 3, 4, or 5 independently selected R^b substituents, provided when R_1 is 5-7 membered heteroaryl or 5-7 membered heterocycloalkyl and R_2 is C₁₋₆ alkoxy, then the 5-7 membered heteroaryl or 5-7 membered

heterocycloalkyl does not connect to the fused phenyl ring of the quinoline moiety through a ring nitrogen atom; or

(ii) R_1 is selected from -H, halo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NO₂, -OR^a, -SR^a, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)R^a, -OC(O)NR^aR^a, -NHR^a, -NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and -S(O)₂NR^aR^a, wherein the (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R_1 or R_2 are each optionally substituted with 1, 2, 3, 4, or 5 independently selected R^b substituents; and

R_2 is selected from the group consisting of (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -CN, -NHOH, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -S(O)NR^aR^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and S(O)₂NR^aR^a, provided when R_1 is 5-7 membered heteroaryl or 5-7 membered heterocycloalkyl and R_2 is C₁₋₆ alkoxy, then the 5-7 membered heteroaryl or 5-7 membered heterocycloalkyl of R_1 does not connect to the fused phenyl ring of the quinoline moiety through a ring nitrogen atom,

(iii) R_1 and R_2 taken together with the atoms to which they are attached form a fused (C₃-C₇) cycloalkyl ring or a fused 4- to 10-membered heterocycloalkyl ring, wherein the fused (C₃-C₇) cycloalkyl ring and fused 4- to 10-membered heterocycloalkyl ring are each optionally substituted with 1, 2, or 3 independently selected R^b substituents, provided that the compound is not 1-[2-(4-Fluoro-phenyl)-acetyl]-cyclopropanecarboxylic acid [3-fluoro-4-(7,8,10,11,13,14-hexahydro-6,9,12,15-tetraoxa-1-aza-cyclododeca[b]naphthalen-4-yloxy)-phenyl]-amide;

R₁₀ and R₁₁ are each independently selected from the group consisting of -H, halo, (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NO₂, -OR^a, -SR^a, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)R^a, -OC(O)NR^aR^a, -NHR^a, -NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and S(O)₂NR^aR^a, wherein the (C₁-C₆) alkyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R₁ or R₂ are each optionally substituted with 1, 2, 3, 4, or 5 independently selected R^b substituents;

each R₃ is independently selected from the group consisting of -H, halo, -OH, -CN, optionally substituted (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkoxy, -NH₂, -NH(C₁-C₆)alkyl, -N(C₁-C₆ alkyl)₂, and (C₃-C₆) cycloalkyl, wherein the (C₁-C₆) alkoxy, -NH(C₁-C₆)alkyl, -N(C₁-C₆ alkyl)₂, and (C₃-C₆) cycloalkyl of R₃ are each optionally substituted with 1, 2, or 3 independently selected R^g substituents;

each R₁₄ is independently selected from the group consisting of halo, -OH, -NH₂, -CN, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, -COOH, -NH(C₁-C₆)alkyl, -N(C₁-C₆ alkyl)₂, phenyl, phenyl-(C₁-C₂) alkylene, (C₃-C₆) cycloalkyl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, 4- to 6-membered heterocycloalkyl, (4- to 6-membered heterocycloalkyl)-(C₁-C₄) alkylene-, 5- to 6-membered heteroaryl, (5- to 6-membered heteroaryl)-(C₁-C₄) alkylene-, and -OR^e, wherein the (C₁-C₆) alkyl, phenyl, phenyl-(C₁-C₂) alkylene, (C₃-C₆) cycloalkyl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, 4- to 6-membered heterocycloalkyl, (4- to 6-membered heterocycloalkyl)-(C₁-C₄) alkylene-, 5- to 6-membered heteroaryl, and (5- to 6-membered heteroaryl)-(C₁-C₄) alkylene- of R₁₄ are each optionally substituted with 1, 2, or 3 independently selected R^g substituents,

R₁₅ is H or C₁₋₆ alkyl;

each R_4 is independently selected from the group consisting of -H, halo, -OH, -COOR^c, -CONR^cR^c, -CN, -NH₂, -NH((C₁-C₆) alkyl), -N((C₁-C₆) alkyl)₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, -CONR^aR^a, -NR^aCOR^a, -NR^aCONR^aR^a, -SO₂R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, (C₃-C₆) cycloalkyl, 4- to 6-membered heterocycloalkyl, phenyl, 5- or 6-membered heteroaryl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, (4- to 6-membered heterocycloalkyl)-(C₁-C₄) alkylene-, phenyl-(C₁-C₂) alkylene-, and (5- or 6-membered heteroaryl)-(C₁-C₄) alkylene-, wherein the (C₁-C₆) alkyl, (C₃-C₆) cycloalkyl, 4- to 6-membered heterocycloalkyl, phenyl, 5- or 6-membered heteroaryl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, (4- to 6-membered heterocycloalkyl)-(C₁-C₄) alkylene-, phenyl-(C₁-C₂) alkylene-, and (5- or 6-membered heteroaryl)-(C₁-C₄) alkylene- of R_4 are each optionally substituted with 1, 2, or 3 independently selected R^f substituents;

each R^a is independently selected from the group consisting of -H, -CN, (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, wherein the (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R^a are each optionally substituted with 1, 2, 3, 4, or 5 independently selected R^d substituents;

each R^b is independently selected from the group consisting of halo, oxo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-10 membered heteroaryl)-(C₁-C₄) alkylene-, (4-10 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -OH, -NH₂, -NO₂, -NHOR^c, -OR^c, -SR^c, -C(O)R^c, -C(O)NR^cR^c, -C(O)OR^c, -C(O)NR^cS(O)₂R^c, -OC(O)R^c, -OC(O)NR^cR^c, -C(=NOH)R^c, -C(=NOH)NR^c, -C(=NCN)NR^cR^c, -NR^cC(=NCN)NR^cR^c, -C(=NR^c)NR^cR^c, -NR^cC(=NR^c)NR^cR^c, -NHR^c, -NR^cR^c, -NR^cC(O)R^c, -NR^cC(=NR^c)R^c, -NR^cC(O)OR^c, -NR^cC(O)NR^cR^c, -NR^cS(O)R^c, -NR^cS(O)₂R^c, -NR^cS(O)₂NR^cR^c, -S(O)R^c, -S(O)NR^cR^c, -S(O)₂R^c, -S(O)₂NR^cC(O)R^c, -Si(R^c)₃, -P(O)R^cR^c, -P(O)(OR^c)(OR^c), -B(OH)₂, -B(OR^c)₂, and -

$S(O)_2NR^cR^e$, wherein the (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-10 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-10 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R^b are each further optionally substituted with 1, 2, or 3 independently selected R^d substituents;

each R^c is independently selected from the group consisting of -H, (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-10 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-10 membered heterocycloalkyl)-(C₁-C₄) alkylene-, wherein the (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-10 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-10 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R^c are each optionally substituted with 1, 2, 3, 4, or 5 independently selected R^f substituents;

each R^d is independently selected from the group consisting of (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, halo, (C₆-C₁₀) aryl, 5-10 membered heteroaryl, (C₃-C₁₀) cycloalkyl, 4-10 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-10 membered heteroaryl)-(C₁-C₄) alkylene-, (4-10 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NH₂, -NHO^e, -OR^e, -SR^e, -C(O)R^e, -C(O)NR^eR^e, -C(O)OR^e, -OC(O)R^e, -OC(O)NR^eR^e, -NHR^e, -NR^eR^e, -NR^eC(O)R^e, -NR^eC(O)NR^eR^e, -NR^eC(O)OR^e, -C(=NR^e)NR^eR^e, -NR^eC(=NR^e)NR^eR^e, -NR^eC(=NOH)NR^eR^e, -NR^eC(=NCN)NR^eR^e, -S(O)R^e, -S(O)NR^eR^e, -S(O)₂R^e, -NR^eS(O)₂R^e, -NR^eS(O)₂NR^eR^e, and -S(O)₂NR^eR^e, wherein the (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₆-C₁₀) aryl, 5-10 membered heteroaryl, (C₃-C₁₀) cycloalkyl, 4-10 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-10 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-10 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R^d are each optionally substituted with 1, 2, or 3 independently selected R^f substituents;

each R^e is independently selected from the group consisting of -H, (C₁-C₆) alkyl, (C₃-C₆) cycloalkyl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, (C₆-C₁₀) aryl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, 5- or 6-membered heteroaryl, (5- or 6-membered heteroaryl)-(C₁-C₄) alkylene-, 4-7-membered

heterocycloalkyl, (4-7-membered heterocycloalkyl)-(C₁-C₄) alkylene-, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₂-C₄) alkenyl, and (C₂-C₄) alkynyl, wherein the (C₁-C₄) alkyl, (C₃-C₆) cycloalkyl, (C₆-C₁₀) aryl, 5 or 6-membered heteroaryl, 4-7-membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (5- or 6-membered heteroaryl)-(C₁-C₄) alkylene-, (4-7-membered heterocycloalkyl)-(C₁-C₄) alkylene-, (C₂-C₄) alkenyl, and (C₂-C₄) alkynyl of R^e are each optionally substituted with 1, 2, or 3 R^f substituents,

or any two R^a substituents together with the nitrogen atom to which they are attached form 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered heterocycloalkyl, each of which is optionally substituted with 1, 2, or 3 independently selected R^f substituents;

or any two R^c substituents together with the nitrogen atom to which they are attached form 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered heterocycloalkyl, each of which is optionally substituted with 1, 2, or 3 independently selected R^f substituents,

or any two R^e substituents together with the nitrogen atom to which they are attached form 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered heterocycloalkyl, each of which is optionally substituted with 1, 2, or 3 independently selected R^f substituents;

each R^f is independently selected from the group consisting of halo, -OH, -CN, -COOH, -NH₂, -NH-(C₁-C₆) alkyl, -N((C₁-C₆) alky)₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) alkylthio, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, phenyl, 5-6 membered heteroaryl, 4-6 membered heterocycloalkyl, and (C₃-C₆) cycloalkyl, wherein the (C₁-C₆) alkyl, phenyl, (C₃-C₆) cycloalkyl, 4-6 membered heterocycloalkyl, and 5-6 membered heteroaryl of R^f are each optionally substituted with 1, 2, or 3 substituents selected from halo, -OH, -CN, -COOH, -NH₂, (C₁-C₄) alkyl, (C₁-C₄) alkoxy, (C₁-C₄) haloalkyl, (C₁-C₄) haloalkoxy, phenyl, (C₃-C₁₀) cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl;

each R^g is independently selected from the group consisting of halo, -OH, -CN, -COOH, -COO-(C₁-C₄) alkyl, -NH₂, -NH-(C₁-C₆) alkyl, -N((C₁-C₆) alky)₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) alkylthio, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, phenyl, 5-6 membered heteroaryl, 4-6 membered heterocycloalkyl, and (C₃-C₆) cycloalkyl;

Y is selected from -O-, -S-, -SO-, -SO₂-, -NH-, and -N((C₁-C₆) alkyl)-;


the ring nitrogen atom on the quinoline moiety in Formula A is optionally oxidized;

the subscript n is an integer of 1, 2, 3, or 4;

the subscript m is an integer of 1, 2, 3, 4, or 5; and


the subscript p is an integer of 0, 1, 2, 3, or 4.

2. The compound of claim 1, having formula A-1:




A-1.

3. The compound of claim 1, having formula A-2:



A-2.


4. The compound of claim 1, having formula A-3:



A-3

wherein R^{a1} is H or (C₁-C₆) alkyl.

5. The compound of claim 1, having formula A-4:



A-4

wherein ring A is 5- to 14-membered heteroaryl; and the subscript r is 1, 2, 3, or 4.

6. The compound of claim 1, wherein:

R_1 is selected from the group consisting of (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, -CN, -NHOH, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)NR^aR^a, C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -S(O)NR^aR^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and S(O)₂NR^aR^a; and

R_2 is selected from -H, halo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NO₂, -OR^a, -SR^a, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)R^a, -OC(O)NR^aR^a, -NHR^a, -NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and -S(O)₂NR^aR^a, wherein the (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R_1 or R_2 are each optionally substituted with 1, 2, 3, 4, or 5 independently selected R^b substituents, provided when R_1 is 5-7 membered heteroaryl or 5-7 membered heterocycloalkyl and

R₂ is C₁₋₆ alkoxy, then the 5-7 membered heteroaryl or 5-7 membered heterocycloalkyl of R₁ does not connect to the fused phenyl ring of the quinoline moiety through a ring nitrogen atom.

7. The compound of claim 1, wherein:

R₁ is selected from -H, halo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, -CN, -NO₂, -OR^a, -SR^a, -NHOR^a, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)R^a, -OC(O)NR^aR^a, -NHR^a, -NR^aR^a, -NR^aC(O)R^a, -NR^aC(=NR^a)R^a, -NR^aC(O)OR^a, -NR^aC(O)NR^aR^a, -C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -NR^aC(=NR^a)NR^aR^a, -NR^aS(O)R^a, -NR^aS(O)₂R^a, -NR^aS(O)₂NR^aR^a, -S(O)R^a, -S(O)NR^aR^a, -S(O)₂R^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and -S(O)₂NR^aR^a, wherein the (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- of R₁ or R₂ are each optionally substituted with 1, 2, 3, 4, or 5 independently selected R^b substituents; and

R₂ is selected from the group consisting of (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -CN, -NHOH, -C(O)R^a, -C(O)NR^aR^a, -C(O)NHOR^a, -C(O)OR^a, -C(O)NR^aS(O)₂R^a, -OC(O)NR^aR^a, C(=NR^a)R^a, -C(=NOH)R^a, -C(=NOH)NR^a, -C(=NCN)NR^aR^a, -NR^aC(=NCN)NR^aR^a, -C(=NR^a)NR^aR^a, -S(O)NR^aR^a, -S(O)₂NR^aC(O)R^a, -P(O)R^aR^a, -P(O)(OR^a)(OR^a), -B(OH)₂, -B(OR^a)₂, and S(O)₂NR^aR^a, provided when R₁ is 5-7 membered heteroaryl or 5-7 membered heterocycloalkyl and R₂ is C₁₋₆ alkoxy, then the 5-7 membered heteroaryl or 5-7 membered heterocycloalkyl of R₁ does not connect to the fused phenyl ring of the quinoline moiety through a ring nitrogen atom.

8. The compound of claim 1, wherein:


R₁ and R₂ taken together with the atoms to which they are attached form a fused (C₃-C₇) cycloalkyl ring or a fused 4- to 10-membered heterocycloalkyl ring, wherein the fused (C₃-C₇) cycloalkyl ring and fused 4- to 10-membered heterocycloalkyl ring are each optionally substituted

with 1, 2, or 3 independently selected R^b substituents, provided that the compound is not 1-[2-(4-Fluoro-phenyl)-acetyl]-cyclopropanecarboxylic acid [3-fluoro-4-(7,8,10,11,13,14-hexahydro-6,9,12,15-tetraoxa-1-aza-cyclododeca[b]naphthalen-4-yloxy)-phenyl]-amide .

9. The compound of claim 1, 6, 7 or 8, wherein:

R_{15} is H or CH_3 .

10. The compound of claim 1 or 9, wherein R_1 and R_2 taken together with the atoms to which they are attached form a fused (C_3 - C_7) cycloalkyl ring or a fused 4- to 10-membered heterocycloalkyl ring, wherein the fused (C_3 - C_7) cycloalkyl ring or a fused 4- to 10-membered heterocycloalkyl ring are each optionally substituted with 1, 2, or 3 independently selected R^b substituents, provided that the compound is not a compound having the formula:



wherein ring E is fused 4- to 10-membered heterocycloalkyl.

11. The compound of any of claims 2, 4, 7, and 9, wherein R_1 is -H.

12. The compound of any of claims 1, 3, 5 and 6, wherein R_2 is -H.

13. The compound of any of claims 1, 2, 4, 5, and 7-9, wherein R_1 is selected from the group consisting of -H, (C_1 - C_6) alkyl, (C_2 - C_6) alkenyl, (C_2 - C_6) alkynyl, $-C(=NO-(C_1-C_6) \text{ alkyl})R^a$, -CN, $-C(O)OR^a$, $-C(O)NR^aR^a$, $-C(O)NHOR^a$, $-S(O)_2NR^aR^a$, phenyl, 5- to 6-membered heteroaryl, (C_3 - C_6) cycloalkyl, and 4- to 6-membered heterocycloalkyl.

14. The compound of claim 13, wherein R_1 is -H, $-R^aNHC(O)-$, $R^aOC(O)-$, (C_1 - C_6) alkyl, (C_1 - C_6) alkoxy or $C(=NO-CH_3)R^a$; and

R₂ is selected from 2-methoxyethylamino, methylamino, 3-morpholinopropoxy, 2-methoxyethoxy, 2-hydroxyethoxy, propoxy, 2-hydroxypropoxy, methoxycarbonyl, carboxy, carbamoyl, methylcarbamoyl, (2-hydroxyethoxy)carbamoyl, (2,2-dihydroxyethoxy)carbamoyl, (oxetan-3-yloxy)carbamoyl, methoxycarbamoyl, 2-trimethylsilylethynyl, ethynyl, sulfamoyl, acetyl, and -C(=NOCH₃)CH₃.

15. The compound of any of claims 1, 3, 5, and 6, wherein R₂ is selected from the group consisting of -H, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -C(=NO-(C₁-C₆) alkyl)R^a, -CN, -C(O)OR^a, -C(O)NR^aR^a, -C(O)NHOR^a, and -S(O)₂NR^aR^a.

16. The compound claim 15, wherein R₁ is selected from 2-methoxyethylamino, azetidin-1-yl, methylamino, 3-morpholinopropoxy, 2-methoxyethoxy, 2-hydroxyethoxy, propoxy, 2-hydroxypropoxy, methoxycarbonyl, carboxy, carbamoyl, methylcarbamoyl, 2-oxazolyl, pyrazol-3-yl, pyrazol-4-yl, 4-isoxazolyl, 3,5-dimethylisoxazol-4-yl, 1-methyl-pyrazol-4-yl, 2-methyl-pyrazol-3-yl, 2-ethyl-pyrazol-3-yl, 2-(2-hydroxyethyl)-pyrazol-3-yl, 2-(2,2,2-trifluoroethyl)-pyrazol-3-yl, 2-(2-fluoroethyl)-pyrazol-3-yl, 2-(2,2-difluoroethyl)-pyrazol-3-yl, 2-trifluoromethyl-pyrazol-3-yl, 2-difluoromethyl-pyrazol-3-yl, 1-methyl-imidazol-4-yl, 1-methyl-imidazol-2-yl, 1H-imidazol-2-yl, (2-hydroxyethoxy)carbamoyl, (2,2-dihydroxyethoxy)carbamoyl, (oxetan-3-yloxy)carbamoyl, methoxycarbamoyl, 2-trimethylsilylethynyl, ethynyl, 1,3,4-oxadiazol-3-yl, 1H-1,2,3-triazol-5-yl, sulfamoyl, acetyl, and -C(=NOCH₃)CH₃, and R₂ is -H, -R^aNHC(O)-, -R^aOC(O)-, -(C₁-C₆) alkyl, (C₁-C₆) alkoxy, or -C(=NO-CH₃)R^a.

17. The compound of claim 5, wherein the subscript r is 1 or 2.


18. The compound of any of claims 1-17, wherein R₁₀ and R₁₁ are each H.

19. The compound of any of claims 1-18, wherein the subscript n is 1.

20. The compound of any of claims 1-19, wherein the subscript m is 1.

21. The compound of any of claims 1-20, wherein the subscript p is 1.

22. The compound of any of claims 1-21, having formula B:



B

or a pharmaceutically acceptable salt thereof, wherein:

(i) R_2 is:

-H or a group selected from the group consisting of:


(C₁-C₆) alkyl, halo, -NO₂, X₁R^a, wherein X₁ is -O-, -S-, -SO-, -SO₂-, -SO₂NH-, -SO₂NR^a-, -NH-, and -N-(C₁-C₆) alkyl-, wherein (C₁-C₆) alkyl is optionally substituted; and

R_1 is selected from the group consisting of:

(C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl are each independently optionally substituted;

-CN, -P(O)R^aR^a, P(O)(OR^a)₂, B(OH)₂, B(OR^a)₂,


X₂R^a, wherein X₂ is -NHO-, -NH-S(O)-, -N-(C₁-C₆)alkyl-S(O)-, -NH-S(O)₂-, -N-(C₁-C₆) alkyl-S(O)₂R^a-, -NH-S(O)-NH-, -N-(C₁-C₆) alkyl-S(O)NH-, -NH-S(O)₂NH-, -N-(C₁-C₆) alkyl-S(O)₂NH-, -S(O)₂NHC(O)-; and




wherein “” indicates the point of attachment, wherein:

Y₁ is absent, or is -NH-, -N-(C₁-C₆) alkyl-, or -O-;

Y₂ is absent, or is -O-, -NH-, -NHO-, -N-(C₁-C₆) alkyl-, -N₂H₂-, -NH-S(O)-, -NH-S(O)₂-; or



Y_2 is or optionally substituted , wherein “” indicates points of attachment, wherein ring A is a 3, 4, 5, 6, or 7-membered ring; and

Z^1 is O, NH, N-(C₁-C₆) alkyl, NOH, NO-(C₁-C₆) alkyl, or NCN; or


(ii)

R_1 is selected from the group consisting of:

(C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl are each independently optionally substituted;


P(O) R^aR^a , P(O)(OR^a)(OR^a), B(OH)₂, B(OR^a)₂, CN,


X_2R^a , wherein X_2 is -NHO-, -NH-S(O)-, -N-(C₁-C₆)alkyl-S(O)-, -NH-S(O)₂-, -N-(C₁-C₆) alkyl-S(O)₂ R^a -, -NH-S(O)-NH-, -N-(C₁-C₆) alkyl-S(O)NH-, -NH-S(O)₂NH-, -N-(C₁-C₆) alkyl-S(O)₂NH-, -S(O)-, -S(O)₂-, -S(O)₂NHC(O); and



Y_1 is absent, or is -NH-, -N-(C₁-C₆) alkyl-, or -O-;

Y_2 is absent, or is -O-, -NH-, -NHO-, -N-(C₁-C₆) alkyl-, -N₂H₂-, -NH-S(O)-, or -NH-S(O)₂; or



Y_2 is optionally substituted , wherein “” indicates points of attachment, wherein ring A is a 3, 4, 5, 6, or 7-membered ring; and


Z^1 is -O-, -NH-, -N-(C₁-C₆) alkyl-, -NOH-, -NO-(C₁-C₆) alkyl-, or -NCN-; or

R_2 is selected from the group consisting of:

H, halo, (C₁-C₆) alkyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, wherein (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- are each independently optionally substituted,



CN, NO₂, P(O)R^aR^a, P(O)(OR^a)(OR^a), B(OH)₂, B(OR^a)₂,

X₁R^a, wherein X₁ is -O-, -S-, -NH-, or -N-(C₁-C₆)-, -NHO-, -NH-S(O)-, -N-(C₁-C₆)alkyl-S(O)-, -NH-S(O)₂-, -N-(C₁-C₆) alkyl-S(O)₂-, -NH-S(O)-NH-, -N-(C₁-C₆) alkyl-S(O)NH-, -NH-S(O)₂NH-, -N-(C₁-C₆) alkyl-S(O)₂NH-, -S(O)₂NHC(O)-, -NH-S(O)R^a-, -N-(C₁-C₆) alkyl-S(O)R^a-, -NH-S(O)₂R^a-, or -N-(C₁-C₆) alkyl-S(O)₂R^a-;



Y₁ is absent or is -NH-, -N-(C₁-C₆) alkyl-, or -O-;

Y₂ is absent or is -O-, -NH-, -NHO-, -N-(C₁-C₆) alkyl-, -N₂H₂-, -NH-S(O)-, or -NH-S(O)₂-; or

Y₂ is optionally substituted , wherein ring A is a 3, 4, 5, 6, or 7-membered ring and wherein “” indicates points of attachment; and

Z¹ is O, NH, N-(C₁-C₆) alkyl, NOH, NO-(C₁-C₆) alkyl, or NCN; or

(iii)

R₁ and R₂ taken together with the atoms to which they are attached form a 4- to 10-membered heterocycloalkyl ring optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of halo, (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, -CN, -OH, -NH₂, provided that the compound is not 1-[2-(4-Fluoro-phenyl)-acetyl]-cyclopropanecarboxylic acid [3-fluoro-4-

(7,8,10,11,13,14-hexahydro-6,9,12,15-tetraoxa-1-aza-cyclododeca[b]naphthalen-4-yloxy)-phenyl]-amide;

and R₁₀, R₁₁, R₃, R₁₄, R₄, n, p, m and Y are as defined as follows:

each R₃ is independently selected from the group consisting of -H, halo, -OH, -CN, optionally substituted (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkoxy, -NH₂, -NH(C₁-C₆)alkyl, -N(C₁-C₆ alkyl)₂, and (C₃-C₆) cycloalkyl, wherein (C₁-C₆) alkoxy, -NH(C₁-C₆)alkyl, -N(C₁-C₆ alkyl)₂, and (C₃-C₆) cycloalkyl are each optionally substituted;

each of R₁₀ and R₁₁ is independently selected from the group consisting of -H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and (C₁-C₆) haloalkoxy;

each R₁₄ is independently selected from the group consisting of -halo, -OH, -NH₂, -CN, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, -COOH, -NH(C₁-C₆)alkyl, -N(C₁-C₆ alkyl)₂, phenyl, phenyl-(C₁-C₂) alkylene, (C₃-C₆) cycloalkyl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, 4- to 6-membered heterocycloalkyl, (4- to 6-membered heterocycloalkyl)-(C₁-C₄) alkylene-, 5- to 6-membered heteroaryl, (5- to 6-membered heteroaryl)-(C₁-C₄) alkylene-, and -OR^e, wherein the (C₁-C₆) alkyl, phenyl, phenyl-(C₁-C₂) alkylene, (C₃-C₆) cycloalkyl, (C₃-C₆) cycloalkyl-(C₁-C₄) alkylene-, 4- to 6-membered heterocycloalkyl, (4- to 6-membered heterocycloalkyl)-(C₁-C₄) alkylene-, 5- to 6-membered heteroaryl, and (5- to 6-membered heteroaryl)-(C₁-C₄) alkylene- of R₁₄ are each optionally substituted;

each R₄ is independently selected from -H, halo, -OH, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, and (C₁-C₆) haloalkoxy, wherein the (C₁-C₆) alkyl are each (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, and (C₁-C₆) haloalkoxy are each independently optionally substituted;

each R^a is independently selected from -H, (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene-, wherein (C₁-C₆) alkyl, (C₁-C₆) haloalkyl, (C₆-C₁₀) aryl, (C₃-C₁₀) cycloalkyl, 5-14 membered heteroaryl, 4-14 membered


heterocycloalkyl, (C₆-C₁₀) aryl-(C₁-C₄) alkylene-, (C₃-C₁₀) cycloalkyl-(C₁-C₄) alkylene-, (5-14 membered heteroaryl)-(C₁-C₄) alkylene-, and (4-14 membered heterocycloalkyl)-(C₁-C₄) alkylene- are each independently optionally substituted;

n, p, and m are each independently integers of 0 to 3; and

Y is selected from -O-, -S-, -SO-, -SO₂-, -NH-, and -N((C₁-C₆) alkyl)-.

23. The compound of claim 22, wherein:

R₁ is -H, -CN, (C₁-C₆) alkyl, (C₃-C₁₀) cycloalkyl, (C₆-C₁₀) aryl, 4-10 membered heterocycloalkyl, 5-10 membered heteroaryl, -S(O)₂NHR^a, -P(O)R^aR^a, -OR^a, or




Y₁ is absent or is -NH-, -N-(C₁-C₆) alkyl-, or -O-;

Y₂ is absent or is -O-, -NH-, -NHO-, -N-(C₁-C₆) alkyl-, -NH-NH-, -NH-S(O)-, or NH-S(O)₂; and

Z¹ is -O-, -NH-, -N-(C₁-C₆) alkyl-, -N-OH, or -N-O(C₁-C₆)alkyl;

24. The compound of claim 23, wherein:

R₂ is -H, halo, -X₁R^a, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, or



Y₁ is absent or is NH, N-(C₁-C₆) alkyl, or O;


Y₂ is absent or is O, NH, NHO, N-(C₁-C₆) alkyl, N₂H₂, NH-S(O), or NH-S(O)₂; and

Z¹ is -O-, -NH-, -N-(C₁-C₆) alkyl-, -NOH-, or -N-O(C₁-C₆)alkyl.


25. The compound of any of claims 22-24, wherein R₃ is -H or halo.

26. The compound of any of claims 22-25, wherein R₄ is -H or halo.

27. The compound of any of claims 22-26, wherein, wherein R_{14} is -H or halo.
28. The compound of any of claims 22-27, wherein Y is -O-.
29. The compound of any of claims 22-28, which is a compound of either formula B-1 or B-2:




B-1




B-2

or a pharmaceutically acceptable salt thereof.

30. The compound of claim 29 of formula B-1, wherein:
- R^{a1} is optionally substituted (C_1 - C_6) alkyl;
- R_1 is -H, -CN, optionally substituted 3-6 membered cycloalkyl, optionally substituted phenyl, optionally substituted 4-6 membered heterocycloalkyl, optionally substituted 5-6 membered heteroaryl, - SO_2 -(C_1 - C_6) alkyl, - SO_2NH_2 , - SO_2 -NH(C_1 - C_6) alkyl, or $P(O)((C_1$ - C_6) alkyl) $_2$; or



Y_2 is absent or is -O-, -NH-, -NHO-, -NH-NH-, or -N-(C₁-C₆) alkyl-; or



Y_2 is optionally substituted, wherein “” indicates points of attachment, wherein ring A is a 3, 4, 5, 6, or 7-membered ring, wherein “” indicates points of attachment,


Z^1 is O, NH, N-(C₁-C₆) alkyl, NHO, or NO-(C₁-C₆) alkyl; and

R^a is -H, -(C₁-C₆) alkyl, 4-6-membered heterocycloalkyl, 3-6-membered cycloalkyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NH(C₁-C₆), -(C₂-C₆) alkylene-N(C₁-C₆)₂, or -(C₂-C₆) alkylene-N-(4-6-membered heterocycloalkyl);

31. The compound of claim 30 of formula B-1, wherein:

R^{a1} is (C₁-C₆) alkyl;

R_1 is- H, -CN, optionally substituted cyclopropyl, optionally substituted phenyl, optionally substituted 4-6 membered azetidiny, optionally substituted pyrrolidinyl, optionally substituted piperidinyl, optionally substituted oxetanyl, optionally substituted oxazolyl, optionally substituted pyridinyl, optionally substituted imidazolyl, optionally substituted pyrrolyl, optionally substituted furnayl, optionally substituted pyrazolyl, optionally substituted oxadiazolyl, -SO₂-(C₁-C₆) alkyl, -SO₂NH₂, -SO₂-NH(C₁-C₆) alkyl, or P(O)((C₁-C₆) alkyl)₂; or



Y_1 is absent;

Y_2 is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or



Y_2 is optionally substituted azetidiny;

Z is O, NH, or N-(C₁-C₆) alkyl; and

R^a is H, (C₁-C₆) alkyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NH(C₁-C₆) alkyl, -(C₂-C₆) alkylene-N((C₁-C₆) alkyl)₂, -(C₂-C₆) alkylene-heterocycloalkyl, and 4-6 membered heterocycloalkyl, wherein heterocycloalkyl is optionally substituted.

32. The compound of claim 30 of formula B-2, wherein:

R^{a2} is optionally substituted (C₁-C₆) alkyl;

R_2 is  wherein “” indicates the point of attachment, wherein:

Y_1 is absent;



Y_2 is absent or is -O- or -NH-; and

Z^1 is O; and

R^a is -H or -(C₁-C₆) alkyl.

33. The compound of claim 32 of formula B-2, wherein:

R^{a2} is (C₁-C₆) alkyl;

R_2 is  wherein “” indicates the point of attachment, wherein:

Y_1 is absent;

Y_2 is absent or is -O- or -NH-; and


Z^1 is O or NO-(C₁-C₆) alkyl; and

R^a is -H or -(C₁-C₆) alkyl.


34. The compound of any of claims 29-31 of formula B-1, R^{a1} is methoxy.

35. The compound of any of claims 29 and 32-33 of formula B-2, R^{a2} is methoxy.

36. The compound of any of claims 22-28, which is a compound of either formula B-3 or B-4:



B-3



B-4

or a pharmaceutically acceptable salt thereof.

37. The compound of claim 36 which is a compound of formula B-3, wherein:

R_1 is -H or (C₁-C₆) alkyl; and

Y_1 is absent;

Y_2 is absent or is -O-, -NHO-, or -NH-; and

Z^1 is O or NO-(C₁-C₆) alkyl; and

R^a is -H or -(C₁-C₆) alkyl.

38. The compound of claim 37 which is a compound of formula B-3, wherein:

R_1 is -H or methyl;

Y_1 is absent;

Y_2 is absent or is -O-, -NHO-, or -NH-; and

Z^1 is O or NO-Me; and

R^a is -H, or Me.

39. The compound of claim 36 which is a compound of formula B-3, wherein:

R_1 and R^a , together with the atoms to which they are attached, form a 4-6 membered heterocycloalkyl ring optionally substituted with halo; (C₁-C₆) alkyl; and (C₁-C₆) haloalkyl.

40. The compound of claim 36 which is a compound of formula B-4, wherein:

Y_1 is absent;

Y_2 is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or

Y_2 is optionally substituted azetidiny;

Z^1 is O, NH, NO-(C₁-C₆) alkyl, N-(C₁-C₆) alkyl; and

R^a is H, (C₁-C₆) alkyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NH(C₁-C₆) alkyl, -(C₂-C₆) alkylene-N((C₁-C₆) alkyl)₂, -(C₂-C₆) alkylene-optionally substituted 4-6 membered heterocycloalkyl), and optionally substituted 4-6 membered heterocycloalkyl;

R_2 is -H, F, Cl, Br, (C₁-C₆)alkoxy, -O-(C₂-C₆)alkylene-OH, -O-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl), (C₂-C₆)alkylene-O-(C₁-C₆) alkyl, -NH₂, -NH-(C₁-C₆ alkyl), -NH-(C₁-C₆)alkylene-(optionally substituted 4-6 membered heterocycloalkyl), -NH-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl).

41. The compound of claim 40 which is a compound of formula B-4, wherein:

Y_1 is absent;

Y_2 is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or

Y_2 is optionally substituted azetidiny;

Z^1 is O, NH, NO-(C₁-C₆) alkyl, N-(C₁-C₆) alkyl; and


R^a is H, methyl, ethyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NHMe, -(C₂-C₆) alkylene-N(Me)₂, -(C₁-C₆) alkylene-morpholinyl, -(C₁-C₆) alkylene-piperidinyl, (C₁-C₆)alkylene-(optionally substituted pyrrolidinyl), optionally substituted azetidiny, or optionally substituted oxetanyl;

R_2 is -H, H, F, Cl, Br, methoxy, -O-(C₂-C₆)alkylene-OH, O-(C₂-C₆)alkylene-OMe, -NH₂, -NH-(C₁-C₆ alkyl), -NH-(C₂-C₆)alkylene-OMe, -NH-(C₂-C₆)alkylene-(optionally substituted morpholinyl), or -NH-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl).

42. The compound of claim 36 which is a compound of formula B-4, wherein:

R_2 and R^a , together with the atoms to which they are attached, form a 4-6 membered heterocycloalkyl ring optionally substituted with halo, (C_1-C_6) alkyl, and (C_1-C_6) haloalkyl.

43. The compound of claims 22-28 which is a compound of formula B-5:



B-5

or a pharmaceutically acceptable salt thereof, wherein ring A in formula B-5 is an optionally substituted 5-10 membered heteroaryl or C_{6-10} aryl, .

44. The compound of claim 43 which is a compound of formula B-5, wherein:

Ring A is an optionally substituted (C_6-C_{10}) aryl, optionally substituted (C_3-C_{10}) cycloalkyl, optionally substituted 5-10 membered heteroaryl, or optionally substituted 4-10 membered heterocycloalkyl; and


R_2 is H, or (C_1-C_6) alkoxy.

45. The compound of claim 44 which is a compound of formula B-5, wherein:

Ring A is an optionally substituted phenyl, optionally substituted cyclopropyl, optionally substituted pyridyl, optionally substituted imidazolyl, optionally substituted pyrrolyl, optionally substituted furanyl, optionally substituted pyrazolyl, optionally substituted oxazolyl, optionally substituted azetidiny, or optionally substituted oxetanyl; and

R_2 is H or methoxy.

46. The compound of any one of claims 22-28, which is a compound of formula C:



C

or a pharmaceutically acceptable salt thereof, wherein:

Y₁ is absent;

Y₂ is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or

Y₂ is optionally substituted azetidiny;

Z¹ is O, NH, NO-(C₁-C₆) alkyl, or N-(C₁-C₆) alkyl;

R^a is H, (C₁-C₆) alkyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NH(C₁-C₆) alkyl, -(C₂-C₆) alkylene-N((C₁-C₆) alkyl)₂, -(C₂-C₆) alkylene-optionally substituted 4-6 membered heterocycloalkyl), or optionally substituted 4-6 membered heterocycloalkyl;

R₂ is -H, F, Cl, Br, (C₁-C₆)alkoxy, -O-(C₂-C₆)alkylene-OH, -O-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl), (C₂-C₆)alkylene-O-(C₁-C₆) alkyl, -NH₂, -NH-(C₁-C₆ alkyl), -NH-(C₁-C₆)alkylene-(optionally substituted 4-6 membered heterocycloalkyl), or -NH-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl); and

n and m are each independently integers of 0 to 3

47. The compound of claim 46, wherein:

Y₁ is absent;

Y₂ is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or


Y₂ is optionally substituted azetidiny;

Z¹ is O, NH, NO-(C₁-C₆) alkyl, or N-(C₁-C₆) alkyl;

R^a is H, methyl, ethyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NHMe, -(C₂-C₆) alkylene-N(Me)₂, -(C₁-C₆) alkylene-morpholinyl, -(C₁-C₆) alkylene-piperidinyl, (C₁-C₆)alkylene-(optionally substituted pyrrolidinyl), optionally substituted azetidiny, or optionally substituted oxetany;

R_2 is -H, H, F, Cl, Br, methoxy, -O-(C₂-C₆)alkylene-OH, O-(C₂-C₆)alkylene-OMe, -NH₂, -NH-(C₁-C₆ alkyl), -NH-(C₂-C₆)alkylene-OMe, -NH-(C₂-C₆)alkylene-(optionally substituted morpholinyl), or -NH-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl); and
and n and m or each 0 or 1.

48. The compound of any one of claims 46-47, which is a compound of formula C-1.



C-1

or a pharmaceutically acceptable salt thereof, wherein:

Y_2 is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or

Y_2 is optionally substituted azetidiny,


Z^1 is O, NH, NO-(C₁-C₆) alkyl, or N-(C₁-C₆) alkyl;

R^a is H, methyl, ethyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NHMe, -(C₂-C₆) alkylene-N(Me)₂, -(C₁-C₆) alkylene-morpholinyl, -(C₁-C₆) alkylene-piperidinyl, (C₁-C₆)alkylene-(optionally substituted pyrrolidinyl), optionally substituted azetidiny, or optionally substituted oxetany;

R_2 is -H, H, F, Cl, Br, methoxy, -O-(C₂-C₆)alkylene-OH, O-(C₂-C₆)alkylene-OMe, -NH₂, -NH-(C₁-C₆ alkyl), -NH-(C₂-C₆)alkylene-OMe, -NH-(C₂-C₆)alkylene-(optionally substituted morpholinyl), or -NH-(C₂-C₆)alkylene-O-(C₁-C₆ alkyl); and

and n and m are each indepenenently 0 or 1.


49. The compound of any one of claims 22-28, which is a compound of formula D:



D


50. The compound of any one of claim 49, wherein:
- R_1 is -H, or (C_1-C_6) alkyl; and
 - Y_1 is absent;
 - Y_2 is absent or is -O-, -NHO-, or -NH-; and
 - Z^1 is O or NO- (C_1-C_6) alkyl,
 - R^a is -H or $-(C_1-C_6)$ alkyl; and
 - n and m are each independently integers of 0 to 3.
51. The compound of any one of claim 50, wherein:
- R_1 is -H or methyl;
 - Y_1 is absent;
 - Y_2 is absent or is -O-, -NHO-, or -NH-; and
 - Z^1 is O or NO-Me; and
 - R^a is -H or Me.
 - n and m are each independently integers of 0 to 1.

52. The compound of any one of claims 49-51, which a compound of formula D-1:



D-1.

53. The compound of any one of claims 22-28, which is a compound of formula E:



E

or a pharmaceutically acceptable salt thereof, wherein:

Ring A is an optionally substituted (C₆-C₁₀) aryl, optionally substituted (C₃-C₁₀) cycloalkyl, optionally substituted 5-10 membered heteroaryl, or optionally substituted 4-10 membered heterocycloalkyl; and


R₂ is H or (C₁-C₆)alkoxy.

54. The compound of claim 53, wherein:

Ring A is an optionally substituted phenyl, optionally substituted cyclopropyl, optionally substituted pyridyl, optionally substituted imidazolyl, optionally substituted pyrrolyl, optionally substituted furanyl, optionally substituted pyrazolyl, optionally substituted oxazolyl, optionally substituted azetidiny, or optionally substituted oxetanyl; and

R₂ is H or methoxy.


55. The compound of any one of claims 22-28, which is a compound of formula F:



or a pharmaceutically acceptable salt thereof, wherein


R^{a1} is optionally substituted (C₁-C₆) alkyl;

R_1 is -H, -CN, optionally substituted 3-6 membered cycloalkyl, optionally substituted phenyl, optionally substituted 4-6 membered heterocycloalkyl, optionally substituted 5-6 membered heteroaryl, -SO₂-(C₁-C₆) alkyl, -SO₂NH₂, -SO₂-NH(C₁-C₆) alkyl, or P(O)((C₁-C₆) alkyl)₂; or



Y_1 is absent;

Y_2 is absent or is -O-, -NH-, -NHO-, -NH-NH-, -N-(C₁-C₆) alkyl-; or



Y_2 is optionally substituted , wherein ring A is a 3, 4, 5, 6, or 7-membered ring, wherein “” indicates points of attachment;


Z^1 is O, NH, N-(C₁-C₆) alkyl, NHO, or NO-(C₁-C₆) alkyl; and

R^a is -H, -(C₁-C₆) alkyl, 4-6-membered heterocycloalkyl, 3-6-membered cycloalkyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NH(C₁-C₆), -(C₂-C₆) alkylene-N(C₁-C₆)₂, -(C₂-C₆) alkylene-N-(4-6-membered heterocycloalkyl);

56. The compound of any one of claim 55, wherein:

R^{a1} is methyl;

R₁ is- H, -CN, optionally substituted cyclopropyl, optionally substituted phenyl, optionally substituted 4-6 membered azetidiny, optionally substituted pyrrolidiny, optionally substituted piperidiny, optionally substituted oxetany, optionally substituted oxazolyl, optionally substituted pyridiny, optionally substituted imidazolyl, optionally substituted pyrrolyl, optionally substituted furnayl, optionally substituted pyrazolyl, optionally substituted oxadiazolyl, -SO₂-(C₁-C₆) alkyl, -SO₂NH₂, -SO₂-NH(C₁-C₆) alkyl, or P(O)((C₁-C₆) alkyl)₂; or



Y₁ is absent;


Y₂ is O, NH, NHO, NH-NH, or N-(C₁-C₆) alkyl; or

Y₂ is optionally substituted azetidiny;

Z¹ is O, NH, or N-(C₁-C₆) alkyl; and

R^a is H, (C₁-C₆) alkyl, -(C₂-C₆) alkylene-OH, -CH₂CHOH-(C₂-C₆) alkylene-OH, -(C₂-C₆) alkylene-NH₂, -(C₂-C₆) alkylene-NH(C₁-C₆) alkyl, -(C₂-C₆) alkylene-N((C₁-C₆) alkyl)₂, -(C₂-C₆) alkylene-heterocycloalkyl), and 4-6 membered heterocycloalkyl, wherein heterocycloalkyl is optionally substituted.


57. The compound of any one of claims 22-28, which is a compound of formula G:



G

or a pharmaceutically acceptable salt thereof, wherein:

R^{a1} is optionally substituted (C₁-C₆) alkyl;



Y₁ is absent;


Y_2 is absent or is -O- or -NH-; and

Z^1 is O; and

R^a is -H or -(C₁-C₆) alkyl.

58. The compound of claim 57, wherein:

R^{a2} is methyl;

R_2 is R^a - Y_2 - $\overset{\overset{Z^1}{\parallel}}{C}$ - Y_1 wherein “” indicates the point of attachment, wherein:


Y_1 is absent;

Y_2 is absent or is -O- or -NH-; and

Z^1 is O or NO-(C₁-C₆) alkyl; and

R^a is -H or -(C₁-C₆) alkyl.

59. The compound any one of claims 1-58 which is a compound of formula H:



H

or a pharmaceutically acceptable salt thereof, wherein:

R_1 is selected from the group consisting of -H, -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₃-C₆ heterocycloalkyl, SO₂NR₈R₉, and (SO₂)-C₁-C₆ alkyl;



wherein when R_1 is selected from the group consisting of -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₃-C₆ heterocycloalkyl, SO₂NR₈R₉, and (SO₂)-C₁-C₆ alkyl, R_2 is H, halo, NR₅R₆, or optionally substituted C₁-C₆ alkoxy;

wherein when R₁ is H, optionally substituted C₁-C₆ alkyl, or optionally substituted C₁-C₆ alkoxy, R₂ is -CO-NR₅R₆ or -CO₂R₇;

or R₁ and R₂ taken together with the atoms to which they are attached to form optionally substituted cycloalkyl or optionally substituted heterocycloalkyl;

R₃ is selected from the group consisting of -H, optionally substituted C₁₋₆ alkyl, -CN, and halo;

R₄ is -H or halo;

 is optionally substituted with one, two, three, or four groups independently selected from the group consisting of halo and C₁-C₆ alkyl, wherein “” indicate points of attachment;

R₅ and R₆ are each independently H, optionally substituted C₁-C₆ alkyl, or optionally substituted C₁-C₆ alkoxy;

R₇ is H or optionally substituted C₁-C₆ alkyl;


R₈ and R₉ are each independently H and optionally substituted C₁-C₆ alkyl or R₈; or

R₈ and R₉ may connect to form optionally substituted heterocycle; and

Y is selected from the group consisting of O, S, SO, SO₂, NH, and N-(C₁-C₆ alkyl).

60. The compound of claim 59, wherein Y is O.

61. The compound of claim 60, wherein R₃ is H.


62. The compound of claim 61, wherein  is not substituted.


63. The compound of claim 62, wherein R₄ is halo.

64. The compound of claim 63, wherein R₄ is para fluoro.


65. The compound of any one of claims 59-64, wherein R₂ is -H, halo, or optionally substituted (C₁-C₆)-alkoxy.

66. The compound of claim 63, wherein R_1 is $-\text{CN}$.
67. The compound of claim 63, wherein R_1 is $-\text{CO}_2\text{H}$.
68. The compound of claim 63, wherein R_1 is $-\text{CO}_2\text{-Me}$.
69. The compound of claim 63, wherein R_1 is $-\text{CO-NHR}_6$.
70. The compound of claim 69, wherein R_1 is $-\text{CO-NH}_2$.
71. The compound of claim 69, wherein R_1 is $-\text{CO-NMeR}_6$.
72. The compound of any one of claims 1, 2, 4, 5, 7, 8, 9, and 63, wherein R_1 is selected







73. The compound of claim 72, wherein R_1 is selected from the group consisting of




74. The compound of claim 72, wherein R_1 selected from the group consisting of




75. The compound of any one of claims 1, 2, 3, 5, 6, 9, and 63-74, wherein R_2 is selected from the group consisting of H, -CN, Br, F, Cl, -OMe, CH_3 , $Me-O-CH_2-CH_2-O-$, $HO-CH_2-CH_2-O-$,



76. The compound of any one of claims 59-65, wherein R_1 is -H, methyl, or methoxy.
77. The compound of claim 76, wherein R_2 is $-\text{CO}_2\text{H}$.
78. The compound of any one of claims 59-65, wherein R_1 is $-\text{CO}_2\text{-Me}$.
79. The compound of any one of claims 59-650, wherein R_1 is $-\text{CO-NHR}_6$.
80. The compound of claim 79, wherein R_1 is $-\text{CO-NH}_2$.
81. The compound of any one of claims 59-65, wherein R_1 is $-\text{CO-NMeR}_6$.
82. The compound of any one of claims 59-65, wherein R_1 is selected from the group



83. The compound of claim 64, which is a compound of formula I-A:



I-A

wherein R_6 is $(\text{C}_1\text{-C}_6)$ alkyl, R_2 is $(\text{C}_1\text{-C}_6)$ alkoxy, R_3 is -H or halo, and R_4 is halo.

84. The compound of any one of claims 1-83 which is selected from the group consisting of:

Cmpd No	Name
5	methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-

Cmpd No	Name
6	4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-
7	1-N-[4-(6-carbamoyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
8	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
9	1-N-[4-[6-(ethylcarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
10	1-N-[4-[6-[2-(dimethylamino)ethylcarbamoyl]-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
11	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(2-piperidin-1-ylethylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-
12	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(2-morpholin-4-ylethylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-
13	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(oxetan-3-ylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
14	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(1-methylazetidin-3-yl)carbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
15	1-N-[4-[6-(azetidine-1-carbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
16	1-N'-(4-fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidine-1-carbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
17	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(methoxycarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
21	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(2R)-pyrrolidin-2-yl]methylcarbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-
22	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(2S)-pyrrolidin-2-yl]methylcarbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-
37	1-N-[4-(6-cyano-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
45	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
58	1-N-[4-(6-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
59	1-N'-(4-fluorophenyl)-1-N-[4-[6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
60	1-N'-(4-fluorophenyl)-1-N-[4-[6-[(1-methylazetidin-3-yl)carbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
67	1-N-[4-(6-carbamoyl-7-fluoroquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
68	1-N-[4-(6-carbamoyl-7-chloroquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

Cmpd No	Name
69	1-N-[4-(7-bromo-6-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
70	1-N-[4-[6-carbamoyl-7-(2-methoxyethylamino)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
71	1-N-[4-[6-carbamoyl-7-(3-morpholin-4-ylpropylamino)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
81	4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-
82	1-N-[4-[6-carbamoyl-7-(methylamino)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide
83	1-N'-(4-fluorophenyl)-1-N-[4-[7-(methylamino)-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-
84	methyl 4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-
87	1-N-[4-(7-amino-6-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
88	1-N-[4-[7-amino-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
89	7-amino-4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]quinoline-
90	methyl 7-amino-4-[4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]quinoline-
92	1-N'-(4-fluorophenyl)-1-N-[4-[(2-methyl-4-oxo-2,3-dihydropyrido[3,2-g][1,3]benzoxazin-6-yl)oxy]phenyl]cyclopropane-1,1-dicarboxamide;
98	1-N-[4-[6-carbamoyl-7-(3-morpholin-4-ylpropoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
103	1-N-[4-[6-carbamoyl-7-(2-methoxyethoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
106	1-N-[4-[6-carbamoyl-7-(2-hydroxyethoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
128	methyl 4-[2-chloro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-
130	4-[2-chloro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-
132	1-N'-[4-(6-carbamoyl-7-methoxyquinolin-4-yl)oxy-3-chlorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
134	1-N'-[3-chloro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
129	methyl 4-[2-fluoro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-
131	4-[2-fluoro-4-[[1-[(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-

Cmpd No	Name
133	1-N'-[4-(6-carbamoyl-7-methoxyquinolin-4-yl)oxy-3-fluorophenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
135	1-N'-[3-fluoro-4-[7-methoxy-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
140	methyl 4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-
141	4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-
142	1-N-[4-(7-carbamoyl-6-methylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
143	1-N'-(4-fluorophenyl)-1-N-[4-[6-methyl-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
150	methyl 4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-
151	4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-6-
152	1-N-[4-(7-carbamoyl-6-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
153	1-N'-(4-fluorophenyl)-1-N-[4-[6-methoxy-7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
162	methyl 4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]quinoline-
163	4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]quinoline-
164	1-N-[4-(7-carbamoylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
165	1-N'-(4-fluorophenyl)-1-N-[4-[7-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
175	1-N-[4-[6-(3-cyano-2-fluorophenyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
176	1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-2-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide;
177	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1-methylimidazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
180	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(5-methylfuran-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
181	tert-butyl 2-[4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-
182	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1-methylpyrazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
183	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1,2-oxazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;

Cmpd No	Name
184	1-N-[4-[6-(3,5-dimethyl-1,2-oxazol-4-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
185	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrazol-5-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
186	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrazol-4-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
187	1-N-[4-(6-cyclopropyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
188	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1H-pyrrol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
191	1-N'-(4-fluorophenyl)-1-N-[4-[6-(1H-imidazol-2-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
192	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3-oxazol-5-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
195	tert-butyl 3-[4-[4-[[1-(4-fluorophenyl)carbamoyl]cyclopropanecarbonyl]amino]phenoxy]-7-
196	1-N'-(4-fluorophenyl)-1-N-[4-[6-(3-hydroxyoxetan-3-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
197	1-N'-(4-fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidin-3-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
198	1-N-[4-[6-(azetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
199	1-N'-(4-fluorophenyl)-1-N-[4-[6-(3-hydroxyazetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
200	1-N-[4-[6-(3,3-difluoroazetidin-1-yl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
201	1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-3-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide;
202	1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-pyridin-4-ylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide;
204	1-N'-(4-fluorophenyl)-1-N-[4-[6-(1H-pyrazol-5-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
206	1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-sulfamoylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide;
209	1-N'-(4-fluorophenyl)-1-N-[4-(6-sulfamoylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide;
210	1-N'-(4-fluorophenyl)-1-N-[4-(7-methoxy-6-methylsulfonylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide;
220	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(1,3,4-oxadiazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
221	1-N'-(4-fluorophenyl)-1-N-[4-[6-(1,3,4-oxadiazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;

or a pharmaceutically acceptable salt thereof.


85. The compound of any one of claims 1-83 which is selected from the group consisting of:

Cmpd No	Name
27	1-N'-(4-fluorophenyl)-1-N-[4-[6-(2-hydroxyethoxycarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
20	1-N'-(4-fluorophenyl)-1-N-[4-[6-(hydroxycarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
26	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(oxetan-3-yl)oxycarbamoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
30	1-N-[4-[6-(2,3-dihydroxypropoxycarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide; <i>Enantiomer 1</i>
31	1-N-[4-[6-(2,3-dihydroxypropoxycarbamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide; <i>Enantiomer 2</i>
32	1-N'-(4-fluorophenyl)-1-N-[4-[6-(hydrazinecarbonyl)-7-methoxyquinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
34	1-N-[4-(6-acetyl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
35	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(E)-N-methoxy-C-methylcarbonimidoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
36	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(Z)-N-methoxy-C-methylcarbonimidoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
50	1-N'-(4-fluorophenyl)-1-N-[4-[7-(2-hydroxyethoxy)-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
51	1-N-[4-(6-dimethylphosphoryl-7-methoxyquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
96	1-N-[4-[(2-ethyl-4-oxo-2,3-dihydropyrido[3,2-g][1,3]benzoxazin-6-yl)oxy]phenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
110	1-N'-(4-fluorophenyl)-1-N-[4-[7-(2-hydroxyethoxy)-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
115	1-N-[4-[6-carbamoyl-7-(2-hydroxypropoxy)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
116	1-N'-(4-fluorophenyl)-1-N-[4-[7-(2-hydroxypropoxy)-6-(methylcarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
125	1-N'-(4-fluorophenyl)-1-N-[4-[7-(2-hydroxypropoxy)-6-(1,3-oxazol-2-yl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;

Cmpd No	Name
166	1-N'-(4-fluorophenyl)-1-N-[4-[7-(2-hydroxyethoxycarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
169	1-N-[4-[7-[[[(2R)-2,3-dihydroxypropoxy]carbamoyl]quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
170	1-N-[4-[7-[[[(2S)-2,3-dihydroxypropoxy]carbamoyl]quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
167	1-N'-(4-fluorophenyl)-1-N-[4-[7-(oxetan-3-yloxycarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
193	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-[(E)-methoxyiminomethyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
207	1-N'-(4-fluorophenyl)-1-N-[4-[7-methoxy-6-(methylsulfamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
208	1-N-[4-[6-(ethylsulfamoyl)-7-methoxyquinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
213	1-N'-(4-fluorophenyl)-1-N-[4-[7-(methoxycarbamoyl)quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;
214	1-N-[4-[7-(ethylcarbamoyl)quinolin-4-yl]oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
254	1-N'-(4-fluorophenyl)-1-N-[4-(7-sulfamoylquinolin-4-yl)oxyphenyl]cyclopropane-1,1-dicarboxamide;
255	1-N-[4-(7-acetylquinolin-4-yl)oxyphenyl]-1-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide; and
256	1-N'-(4-fluorophenyl)-1-N-[4-[7-[(E)-N-methoxy-C-methylcarbonimidoyl]quinolin-4-yl]oxyphenyl]cyclopropane-1,1-dicarboxamide;

or a pharmaceutically acceptable salt thereof.


86. A process for making a compound of Formula I:



I


or a pharmaceutically acceptable salt thereof, comprising:

reacting a compound of formula II:



II

wherein Z is selected from the group consisting of NH₂, SH, and OH;
with a compound of formula III:



III

wherein X is a leaving group;

R₁ is selected from the group consisting of -H, -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₁-C₆) alkyl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C₁-C₆) alkyl;

wherein when R₁ is selected from the group consisting of -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C₁-C₆) alkyl, R₂ is -H, halo, -NR₅R₆, or optionally substituted (C₁-C₆) alkoxy;


wherein when R₁ is -H, optionally substituted (C₁-C₆) alkyl, or optionally substituted (C₁-C₆) alkoxy, R₂ is -CO-NR₅R₆ or -CO₂R₇;

or R₁ and R₂ taken together with the atoms to which they are attached to form optionally substituted cycloalkyl or optionally substituted heterocycloalkyl;

R₃ is selected from the group consisting of -H, optionally substituted (C₁-C₆) alkyl; -CN; and halo;

R₄ is -H or halo;



is optionally substituted with one two, three, or four groups independently selected from the group consisting of halo and (C₁-C₆) alkyl, wherein “” indicate points of attachment;

R_5 and R_6 are each independently -H, optionally substituted (C_1 - C_6) alkyl, or optionally substituted (C_1 - C_6) alkoxy;


R_7 is -H or optionally substituted (C_1 - C_6) alkyl;

R_8 and R_9 are each independently -H and optionally substituted (C_1 - C_6) alkyl; or

R_8 and R_9 may connect to form optionally substituted heterocycle; and


Y is selected from the group consisting of O, S, SO, SO₂, NH, and N-((C_1 - C_6) alkyl).

87. A process for making a compound of Formula I:




or a pharmaceutically acceptable salt thereof, comprising:

reacting a compound of formula IV:



with a compound of formula V:



R_1 is selected from the group consisting of -H, -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C_1 - C_6) alkyl, optionally substituted (C_3 - C_8) cycloalkyl, optionally substituted (C_3 - C_6) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C_1 - C_6) alkyl;



wherein when R_1 is selected from the group consisting of $-\text{CN}$, $-\text{CO}-\text{NR}_5\text{R}_6$, $-\text{CO}_2\text{R}_7$, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted ($\text{C}_3\text{-C}_8$) cycloalkyl, optionally substituted ($\text{C}_3\text{-C}_6$) heterocycloalkyl, $-\text{SO}_2\text{NR}_8\text{R}_9$, and $-(\text{SO}_2)-(\text{C}_1\text{-C}_6)$ alkyl, R_2 is $-\text{H}$, halo, $-\text{NR}_5\text{R}_6$, or optionally substituted ($\text{C}_1\text{-C}_6$) alkoxy;

wherein when R_1 is $-\text{H}$, optionally substituted ($\text{C}_1\text{-C}_6$) alkyl, or optionally substituted ($\text{C}_1\text{-C}_6$) alkoxy, R_2 is $-\text{CO}-\text{NR}_5\text{R}_6$ or $-\text{CO}_2\text{R}_7$;

or R_1 and R_2 taken together with the atoms to which they are attached to form optionally substituted cycloalkyl or optionally substituted heterocycloalkyl;

R_3 is selected from the group consisting of $-\text{H}$, optionally substituted ($\text{C}_1\text{-C}_6$) alkyl, $-\text{CN}$, and halo;

R_4 is $-\text{H}$ or halo;

 is optionally substituted with one, two, three, or four groups independently selected from the group consisting of halo, and ($\text{C}_1\text{-C}_6$) alkyl, wherein “” indicate points of attachment;

R_5 and R_6 are each independently $-\text{H}$, optionally substituted ($\text{C}_1\text{-C}_6$) alkyl, or optionally substituted ($\text{C}_1\text{-C}_6$) alkoxy;


R_7 is $-\text{H}$ or optionally substituted ($\text{C}_1\text{-C}_6$) alkyl;

R_8 and R_9 are each independently $-\text{H}$ and optionally substituted ($\text{C}_1\text{-C}_6$ alkyl); or

R_8 and R_9 may connect to form optionally substituted heterocycle; and


Y is selected from the group consisting of O , S , SO , SO_2 , NH , and $\text{N}-((\text{C}_1\text{-C}_6) \text{ alkyl})$.

88. The process of claim 87, further comprising reacting a compound of Formula VI:




VI

with a compound of Formula VII:



VII

to form a compound of Formula VIII:



VIII

and reducing the compound of Formula VIII to provide a compound of Formula V, wherein:

R_1 is selected from the group consisting of -H, -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₁-C₆) alkyl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C₁-C₆) alkyl;

wherein when R_1 is selected from the group consisting of -CN, -CO-NR₅R₆, -CO₂R₇, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted (C₃-C₈) cycloalkyl, optionally substituted (C₃-C₆) heterocycloalkyl, -SO₂NR₈R₉, and -(SO₂)-(C₁-C₆) alkyl, R_2 is -H, halo, -NR₅R₆, or optionally substituted (C₁-C₆) alkoxy;

wherein when R_1 is -H, optionally substituted (C₁-C₆) alkyl, or optionally substituted (C₁-C₆) alkoxy, R_2 is -CO-NR₅R₆, or -CO₂R₇;

or R_1 and R_2 taken together with the atoms to which they are attached to form optionally substituted cycloalkyl or optionally substituted heterocycloalkyl;

R_3 is selected from the group consisting of -H, optionally substituted (C₁-C₆) alkyl, -CN, and halo; and

W is halo.

89. A pharmaceutical composition comprising a compound of any one of claims 1-85, or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

90. A method of treating a disease, disorder, or syndrome mediated at least in part by modulating in vivo activity of a protein kinase, comprising administering to a subject in need thereof a compound of any one of claims 1-85 or a pharmaceutical composition of claim 89.

91. The method of claim 90, wherein the disease, disorder, or syndrome mediated at least in part by modulating in vivo activity of a protein kinase is cancer.

92. A method for inhibiting a protein kinase, the method comprising contacting the protein kinase with a compound of any one of claims 1-85.

93. The method of any one of claims 90-92, wherein the protein kinase is Axl, Mer, c-Met, KDR, or a combination thereof.

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2019/015297

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K31/47 A61K31/4709 C07D215/22 C07D401/04 C07D401/06
C07D401/12 C07D405/04 C07D405/12 C07D413/04 C07D413/12
A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K C07D A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CN 104 817 497 B (NANJING ZHONGRUIYUAN BIOTECHNOLOGY CO LTD) 8 March 2017 (2017-03-08) the whole document	1,3,7,9, 18-20, 29,35, 72,87-93
X	CN 105 797 123 A (WANG CHAO) 27 July 2016 (2016-07-27) the whole document	1,10, 18-20, 59,89,90
X	CN 106 400 155 A (PHIFER SPECIAL TEXTILE CO LTD) 15 February 2017 (2017-02-15) the whole document	1,10, 18-22,59
	-/--	

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

3 May 2019

Date of mailing of the international search report

13/05/2019

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Rufet, Jacques

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2019/015297

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2010/045095 A1 (XI NING [US]) 22 April 2010 (2010-04-22) claims 1,10, 21,23-40 -----	1-9, 11-58, 60-89
X	EP 2 769 976 A1 (BEIJING KONRUNS PHARMACEUTICAL CO LTD [CN] ET AL.) 27 August 2014 (2014-08-27) claims 1,18-28 -----	1,89-93
X	US 2018/009758 A1 (HORN GERALD [US]) 11 January 2018 (2018-01-11) compound 4; paragraph [0074] - paragraph [0075]; claim 13 -----	1,3,89, 90,92,93
X	WO 2012/006960 A1 (ZHEJIANG BETA PHARMA INC [CN]; HU SHAOJING [CN] ET AL.) 19 January 2012 (2012-01-19) Scheme 1; claims 1-84; examples 3,13 -----	1,9,59, 89-93
X	WO 2012/034055 A2 (ADVENCHEN LAB LLC [US]) 15 March 2012 (2012-03-15) claims 1-11 -----	1,3, 89-93

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2019/015297

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-93

A compound of formula A

1.1. claims: 2(completely); 1, 4-7, 9, 11-93(partially)

A compound of formula A(i)

1.2. claims: 3(completely); 1, 4-7, 9, 11-93(partially)

A compound of formula A(ii)

1.3. claims: 8, 10(completely); 1, 22, 86-93(partially)

A compound of formula A(iii)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2019/015297

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
CN 104817497	B	08-03-2017	NONE
CN 105797123	A	27-07-2016	NONE
CN 106400155	A	15-02-2017	NONE
WO 2010045095	A1	22-04-2010	AU 2009303602 A1 22-04-2010 BR PI0920261 A2 04-08-2015 CA 2740746 A1 22-04-2010 EP 2340244 A1 06-07-2011 EP 2532657 A2 12-12-2012 ES 2599458 T3 01-02-2017 HK 1172896 A1 18-08-2017 JP 5734193 B2 17-06-2015 JP 6058711 B2 11-01-2017 JP 2012505888 A 08-03-2012 JP 2015120733 A 02-07-2015 KR 20110075016 A 05-07-2011 RU 2011119478 A 27-11-2012 US 2010093727 A1 15-04-2010 US 2012315247 A1 13-12-2012 WO 2010045095 A1 22-04-2010
EP 2769976	A1	27-08-2014	AU 2011377440 A1 15-05-2014 CA 2858033 A1 28-03-2013 CN 102408411 A 11-04-2012 EP 2769976 A1 27-08-2014 HK 1171009 A1 03-07-2015 JP 5795826 B2 14-10-2015 JP 2014526522 A 06-10-2014 KR 20140069188 A 09-06-2014 NZ 623923 A 29-05-2015 RU 2014112345 A 27-10-2015 SG 11201400769P A 30-07-2014 TW 201313683 A 01-04-2013 US 2014221425 A1 07-08-2014 WO 2013040801 A1 28-03-2013
US 2018009758	A1	11-01-2018	NONE
WO 2012006960	A1	19-01-2012	AU 2011278832 A1 28-02-2013 CA 2805148 A1 19-01-2012 EP 2593462 A1 22-05-2013 ES 2598530 T3 27-01-2017 JP 5770281 B2 26-08-2015 JP 2013531016 A 01-08-2013 KR 20130058036 A 03-06-2013 SG 187064 A1 28-02-2013 US 2013123286 A1 16-05-2013 WO 2012006960 A1 19-01-2012
WO 2012034055	A2	15-03-2012	AU 2011299082 A1 02-05-2013 BR 112013005523 A2 03-05-2016 CA 2810528 A1 15-03-2012 CN 103328447 A 25-09-2013 CN 105153028 A 16-12-2015 EP 2621904 A2 07-08-2013 HK 1218758 A1 10-03-2017

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2019/015297

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		JP 5939254 B2	22-06-2016
		JP 2013537197 A	30-09-2013
		KR 20140021509 A	20-02-2014
		KR 20180021901 A	05-03-2018
		NZ 609283 A	27-02-2015
		US 2012123126 A1	17-05-2012
		WO 2012034055 A2	15-03-2012
