



(86) **Date de dépôt PCT/PCT Filing Date:** 2013/06/17
(87) **Date publication PCT/PCT Publication Date:** 2013/12/27
(85) **Entrée phase nationale/National Entry:** 2015/12/18
(86) **N° demande PCT/PCT Application No.:** US 2013/046176
(87) **N° publication PCT/PCT Publication No.:** 2013/192109
(30) **Priorités/Priorities:** 2012/06/17 (US61/660,757);
2012/12/06 (US61/734,331); 2013/03/13 (US61/780,948)

(51) **Cl.Int./Int.Cl. A61K 31/202** (2006.01),
A61P 3/06 (2006.01), **A61P 9/00** (2006.01)

(71) **Demandeur/Applicant:**
MATINAS BIOPHARMA, INC., US

(72) **Inventeurs/Inventors:**
BOBOTAS, GEORGE, US;
FAWZY, ABDEL AZIZ, US

(74) **Agent:** GOWLING LAFLEUR HENDERSON LLP

(54) **Titre : COMPOSITIONS D'ACIDE PENTANOIQUE OMEGA-3 ET LEURS PROCEDES D'UTILISATION**

(54) **Title: OMEGA-3 PENTAENOIC ACID COMPOSITIONS AND METHODS OF USE**

(57) **Abrégé/Abstract:**

Orally administrable composition comprising fatty acids, wherein at least 50% by weight of the fatty acids comprise omega-3-fatty acids, salts or derivatives thereof, wherein the omega-3 fatty acids comprise eicosapentaenoic acid (EPA; C20:5-n3), docosapentaenoic acid (DPA; C22:5-n3), and docosahexaenoic acid (DHA; C22:6-n3), wherein the ratio of DHA to EPA (DHA:EPA) is less than 1:20, and wherein the ratio of DHA to DPA (DHA:DPA) is less than 2:1 are provided. These compositions can be used for the treatment or prophylaxis of dyslipidemic, cardiovascular, CNS, inflammatory, and other diseases/conditions or risk factors therefore.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
27 December 2013 (27.12.2013)

WIPO | PCT

(10) International Publication Number

WO 2013/192109 A1

(51) International Patent Classification:

A61K 31/225 (2006.01) A61K 31/202 (2006.01)

A61K 31/40 (2006.01)

(21) International Application Number:

PCT/US2013/046176

(22) International Filing Date:

17 June 2013 (17.06.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/660,757 17 June 2012 (17.06.2012) US

61/734,331 6 December 2012 (06.12.2012) US

61/780,948 13 March 2013 (13.03.2013) US

(71) Applicant: MATINAS BIOPHARMA, INC. [US/US];
1245 North Florida Avenue, Tarpon Springs, Florida
34689 (US).(72) Inventors: BOBOTAS, George; 1245 N. Florida Ave.,
Tarpon Springs, Florida 34689 (US). FAWZY, Abdel
Aziz; 12525 Colony Preserve Drive, Boynton Beach, Flor-
ida 33436 (US).(74) Agents: KIM, Yelee et al.; Arent Fox, LLP, 1717 K Street,
N.W., Washington, District of Columbia 20036-5342 (US).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))

(54) Title: OMEGA-3 PENTAENOIC ACID COMPOSITIONS AND METHODS OF USE

(57) Abstract: Orally administrable composition comprising fatty acids, wherein at least 50% by weight of the fatty acids comprise omega-3-fatty acids, salts or derivatives thereof, wherein the omega-3 fatty acids comprise eicosapentaenoic acid (EPA; C20:5-n3), docosapentaenoic acid (DPA; C22:5-n3), and docosahexaenoic acid (DHA; C22:6-n3), wherein the ratio of DHA to EPA (DHA:EPA) is less than 1:20, and wherein the ratio of DHA to DPA (DHA:DPA) is less than 2:1 are provided. These compositions can be used for the treatment or prophylaxis of dyslipidemic, cardiovascular, CNS, inflammatory, and other diseases/conditions or risk factors therefore.

WO 2013/192109 A1

Omega-3 Pentaenoic Acid Compositions and Methods of Use

FIELD OF INVENTION

[0001] The present invention relates to a method utilizing formulations comprising omega-3 fatty acids for the treatment of patients by administering an effective amount of a pharmaceutical formulation of the present invention to a subject prone to or afflicted with such disease, and wherein the disease is an amenable disease selected from the group consisting of: hypertriglyceridemia; hypercholesterolemia; mixed dyslipidemia; coronary heart disease (CHD); vascular disease; cardiovascular disease; acute coronary syndrome; atherosclerotic disease and related conditions; heart failure; cardiac arrhythmias; coagulatory conditions associated with cardiac arrhythmias; ischemic dementia; vascular dementia; hypertension; coagulation related disorders; nephropathy; kidney or urinary tract disease; retinopathy; cognitive and other CNS disorders; autoimmune diseases; inflammatory diseases; asthma or other respiratory disease; dermatological disease; metabolic syndrome; diabetes, diabetes mellitus or other form of metabolic disease; liver disease; non-alcoholic fatty liver disease; disease of the gastrointestinal tract; disease of the male or female reproductive system or related secondary sexual organs; a cancer of any type, including lymphomas and myelomas; an infection caused by a virus, bacterium, fungus, protozoa or other organism; and the treatment and/or prevention and/or reduction of cardiac events and/or cardiovascular events and/or vascular events and/or symptoms. The present invention also relates to treatment of such conditions in with concomitant treatments regimes or combination products with other active pharmaceutical ingredients.

BACKGROUND OF THE INVENTION

[0002] In humans, cholesterol and triglycerides are part of lipoprotein complexes in the bloodstream, and can be separated via ultracentrifugation into high-density lipoprotein (HDL), intermediate-density lipoprotein (IDL), low-density lipoprotein (LDL) and very-low-density lipoprotein (VLDL) fractions. Cholesterol and triglycerides are synthesized in the liver, incorporated into VLDL, and released into the plasma. High levels of total cholesterol (total-C), LDL-cholesterol, and apolipoprotein B (a membrane complex for LDL-cholesterol and VLDL-cholesterol, as well as IDL-cholesterol in rare individuals suffering from a disorder resulting in significant IDL-

cholesterol levels) promote human atherosclerosis; these elevated levels are often referred to as hypercholesterolemia. Decreased levels of HDL-cholesterol and its transport complex, apolipoprotein A, as well as elevated levels of apolipoprotein C-III and serum triglycerides (TG) are also associated with the development of atherosclerosis. Further, cardiovascular morbidity and mortality in humans can vary directly with the level of total-C, LDL-cholesterol and TG and inversely with the level of HDL-cholesterol. In addition, researchers have found that non-HDL-cholesterol is an important indicator of hypertriglyceridemia (elevated triglycerides), vascular disease, atherosclerotic disease and related conditions. Therefore, non-HDL-cholesterol and fasting TG reduction has also been specified as a treatment objective in NCEP ATP III. Fasting TG is commonly used as a key measure for TG in lipid management, because it minimizes the confounding factor of TG recently absorbed from meals, including the high variability of the content of meals and high variability of post-meal (post-prandial) spikes in TG. In some preferred embodiments, we refer to fasting TG levels when we refer to triglycerides or TG.

[0003] The NCEP ATPIII treatment guidelines identify HMG-CoA reductase inhibitors ("statins") as the primary treatment option for hypercholesterolemia. In patients with TG<500mg/dL, LDL-cholesterol is the primary treatment parameter. Many patients, however, have increased LDL-cholesterol combined with high TG and low HDL-cholesterol, a condition also known as mixed dyslipidemia. Patients with hypercholesterolemia or mixed dyslipidemia often present with high blood levels of LDL-cholesterol (i.e. greater than 190 mg/dl) and TG (i.e. levels of 200 mg/dl or higher). The use of diet and single-drug therapy does not always decrease LDL-cholesterol and TG adequately enough to reach targeted values in patients with mixed dyslipidemia with or without a concomitant increase in triglycerides. In these patients, a combined therapy regimen of a statin and a second anti-dyslipidemic agent is often desired. This second agent has historically been a fibrate (i.e. gemfibrozil, bezafibrate, or fenofibrate) or extended release niacin. Over the few years, the use omega-3 fatty acid concentrates in combination with a statin has been growing rapidly due to concerns about the lack of outcome benefits with fibrates (i.e. the FIELD study) or extended release niacin (i.e. the AIM-HIGH study). In patients with isolated hypertriglyceridemia, the use of omega-3 fatty acid concentrates has also grown versus fibrates and extended release niacin.

[0004] Marine oils, also commonly referred to as fish oils, are a good source of the two main omega-3 fatty acids, eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), which have been found to regulate lipid metabolism. Omega-3 fatty acids have been found to have beneficial effects on the risk factors for cardiovascular diseases, especially mild hypertension, hypertriglyceridemia and on the coagulation factor VII phospholipid complex activity. Omega-3 fatty acids lower serum triglycerides (TG), increase serum HDL-cholesterol, lower systolic and diastolic blood pressure and the pulse rate, and lower the activity of the blood coagulation factor VII-phospholipid complex. Further, omega-3 fatty acids seem to be well tolerated, without giving rise to any severe side effects.

[0005] The table directly below lists the most common omega-3 fatty acids, including their 3-letter abbreviation code. In this application, the use of any of the 3-letter abbreviations shall refer to the omega-3 fatty acid, unless otherwise indicated (e.g. DPA or DPA 22:5 (n-3) or DPA 22:5-n3 or DPA 22:5n3 or DPA-n3, which all refer to the omega-3 isomer of docosapentaenoic acid).

Common Name for Omega-3 Fatty Acid (+abbreviation)	Codified Lipid Name	Chemical Name
Hexadecatrienoic acid (HTA)	16:3 (n-3)	all-cis-7,10,13-hexadecatrienoic acid
α -linolenic acid (ALA)	18:3 (n-3)	all-cis-9,12,15-octadecatrienoic acid
Stearidonic acid (SDA)	18:4 (n-3)	all-cis-6,8,12,15-octadecatetraenoic acid
Eicosatrienoic acid (ETE)	20:3 (n-3)	all-cis-11,14,17-eicosatrienoic acid
Eicosatetraenoic acid (ETA)	20:4 (n-3)	all-cis-8,11,14,17-eicosatetraenoic acid
Eicosapentaenoic acid (EPA)	20:5 (n-3)	all-cis-5,8,11,14,17-eicosapentaenoic acid
Heineicosapentaenoic acid (HPA)	21:5 (n-3)	all-cis-6,9,12,15,18-heineicosapentaenoic acid
Docosapentaenoic acid (DPA) or Clupanodonic acid	22:5 (n-3)	all-cis-7,10,13,16,19-docosapentaenoic acid
Docosahexaenoic acid (DHA)	22:6 (n-3)	all-cis-4,7,10,13,16,19-docosahexaenoic acid
Tetracosapentaenoic acid (TPA)	24:5 (n-3)	all-cis-9,12,15,18,21-tetracosapentaenoic acid
Tetracosahexaenoic acid (THA) or Nisinic acid	24:6 (n-3)	all-cis-6,9,12,15,18,21-tetracosahexaenoic acid

[0006] One form of omega-3 fatty acids is a concentrate of omega-3, long chain, polyunsaturated fatty acids from fish oil containing DHA ethyl esters, EPA ethyl esters as well as ethyl esters of other omega-3 fatty acids (described in USP35 for LOVAZA®) and is sold under the trademarks OMACOR® and LOVAZA®. Such a form of omega-3 fatty acid comprises at least 90% omega-3 fatty acids of which at least 80% EPA+DHA (in a ratio of 1.2:1) and is described, for example, in U.S. Pat. Nos. 5,502,077, 5,656,667 and 5,698,594. LOVAZA® (omega-3-acid ethyl esters) is indicated for the treatment of patients with hypertriglyceridemia with TG levels of 500mg/dL or higher.

[0007] Another form of omega-3 fatty acid concentrate is sold under the trademark EPADEL® for the treatment of dyslipidemia. This product is described as 98% EPA ethyl ester in Lancet (Vol.369; March 31, 2007; 1090-1098) reporting on a large outcome study with EPADEL®. EPADEL® is known to contain less than 1% of any fatty acid other than EPA.

[0008] Similar to EPADEL®, another form of omega-3 fatty acid concentrate also consists almost entirely of EPA ethyl ester and is known under its developmental stage name AMR101 or its trade name VASCEPA®. This product is described in US patent application 2010/0278879 as comprising at least 95% EPA (typically referred to as 97% or at least 96% in company releases and references) and less than 1% of any other fatty acid. AMR101 was previously under development for the treatment of Huntington's Disease but failed in phase III clinical development. Subsequently, AMR101 was entered in a development program for hypertriglyceridemia and mixed dyslipidemia.

[0009] Yet another concentrate of omega-3, long chain, polyunsaturated fatty acids from fish oil containing approximately 75% DHA and EPA as free fatty acids is known under its developmental stage name EPANOVA™. This product is described as comprising approximately 55% EPA and 20% DHA. EPANOVA™ was previously under development for the treatment of Crohn's Disease but failed in phase III clinical development. Subsequently, EPANOVA™ was entered in a development program for hypertriglyceridemia and mixed dyslipidemia.

[0010] Generally, the bioavailability and therapeutic effect of omega-3 fatty acid compositions is dose dependent, i.e., the higher the dose, the greater the therapeutic effect and bioavailability. However, the effect of each specific omega-3 fatty acid composition may be different, and therefore the level of therapeutic effect of one composition at a given dose cannot necessarily be inferred from the level of therapeutic effects of other omega-3 fatty acid compositions at the same or similar dose.

[0011] For instance, in the MARINE study, it was found that four 1-gram capsules of AMR101/VASCEPA® significantly reduced fasting TG in patients with very high triglycerides (TG>500mg/dL) (March 2011, ACC poster reporting top-line results of the MARINE study), similar to four 1-gram capsules of LOVAZA® but in a less potent manner (LOVAZA® prescribing information, December 2010). In this same study, AMR101 slightly and non-significantly changed LDL-C while LOVAZA® shows a

WO 2013/192109

PCT/US2013/046176

large significant increase in this same population, putting the latter at a disadvantage. Table A directly below compares these profiles.

	LOVAZA - 4 gram/day		Vascepa - 4 gram/day		Vascepa - 2 gram/day	
	% change vs. Placebo	p-value	% change vs. Placebo	p-value	% change vs. Placebo	p-value
TG	-51.6	$p < 0.05$	-33.1	$p < 0.05$	-19.7	$p < 0.05$
Total-C	-8.0	$p < 0.05$	-16.3	$p < 0.0001$	-6.8	$p = 0.2348$
LDL-C	49.3	$p < 0.05$	-2.3	NS	5.2	NS
VLDL-C	-40.8	$p < 0.05$	-28.6	$p < 0.0002$	-15.3	$p = 0.038$
Non-HDL-C	-30.2	$p < 0.05$	-17.7	$p < 0.0001$	-8.1	$p = 0.382$
Apo-B	NR		-8.5	$p = 0.0019$	-2.6	NS
HDL-C	9.1	$p < 0.05$	-3.6	NS	1.5	NS

NR = Not Reported; NS = Not Significant

[0012] In another study with AMR101/VASCEPA®, the ANCHOR study, it was found that four 1-gram capsules of AMR101 significantly reduced fasting TG in patients on statin therapy with high triglycerides (TG 200-499 mg/dL), similar to four 1-gram capsules of LOVAZA® but in a less potent manner (Study in table 3, LOVAZA® prescribing information, December 2010). In this same study, AMR101 decreased LDL-C at 4 gr/day while LOVAZA® shows a significant LDL-C increase in this same population. AMR101 is also more potent than LOVAZA® in reducing non-HDL-cholesterol in this population. Table B directly below compares these profiles.

	LOVAZA - 4 gram/day		Vascepa - 4 gram/day		Vascepa - 2 gram/day	
	% change vs. Placebo	p-value	% change vs. Placebo	p-value	% change vs. Placebo	p-value
TG	-23.2	$p < 0.0001$	-21.5	$p < 0.0001$	-10.1	$p = 0.0035$
Total-C	-3.1	$p < 0.05$	NR	$p < 0.0001$	NR	$p = 0.0019$
LDL-C	3.5	$p < 0.05$	-6.3	$p = 0.0067$	-3.6	NS
VLDL-C	-20.3	$p < 0.05$	-24.4	$p < 0.0001$	-20.5	$p = 0.0083$
Non-HDL-C	-6.8	$p < 0.0001$	-13.5	$p < 0.0001$	-6.5	$p = 0.0054$
Apo-B	-2.3	$p < 0.05$	-9.3	$p < 0.0001$	-3.8	$p = 0.0170$
HDL-C	4.6	$p < 0.05$	-4.5	$p = 0.0013$	-2.2	NS

NS = Not Significant

[0013] The resulting lipid profile of AMR101 versus LOVAZA® in highly similar patient populations indicates that there are significant benefits of using an almost pure EPA oil composition as opposed to an omega-3 mixture as in LOVAZA®. These benefits translate into better non-HDL- and LDL-Cholesterol reduction with the pure EPA form, where these benefits are less or, in the case of the LDL-C effect, the opposite.

[0014] The recently released results from Omthera's EVOLVE trial with EPANOVA™, in patients with very high triglycerides (TG ≥ 500 mg/dL), described a

TG reduction of 31% versus baseline for the 4 gram per day dose and 26% versus baseline for the 2 gram per day dose, with 10% and 8% non-HDL reduction respectively. It appears that the TG-reducing potency of EPANOVA™ is similar to the potency of AMR101. No data were reported by Omthera on the LDL-C effect in the EVOLVE trial.

[0015] The recently released results from Omthera's ESPRIT trial with EPANOVA™, in patients with high triglycerides (TG 200-499 mg/dL) while on statin therapy, described a TG reduction of 21% versus baseline for the 4 gram per day dose and 15% versus baseline for the 2 gram per day dose, with 7% and 4% non-HDL reduction respectively. It appears that the TG-reducing potency of EPANOVA™ is similar to the potency of AMR101. No data were reported by Omthera on the LDL-C effect in the ESPRIT trial.

[0016] From the comparison of LOVAZA® versus AMR101 data, there appears to be a benefit of using pure EPA concentrates for dyslipidemia treatment over omega-3 mixtures with regard to LDL-Cholesterol and non-HDL-cholesterol effects. With the NCEP ATP III guidelines placing LDL-cholesterol and non-HDL-cholesterol reduction at the top of the treatment hierarchy for patients with TG<500 mg/dL, AMR101 is clearly superior to LOVAZA® in this patient category.

[0017] In another example, in the ECLIPSE Study,, the bioavailability of EPANOVA™ is compared to LOVAZA® under high fat meal and low fat meal dosing conditions.

[0018] In the ECLIPSE study it is found that EPANOVA™ is significantly more bioavailable than LOVAZA® after single dose administration (four capsules of 1 gram for both products), both by C_{max} (maximum concentration) and AUC (area under curve) measures (see Table C below, where C_{max} and AUC are estimated from the data points in Figures 1 and 2). Relative to LOVAZA® under high fat meal conditions, EPANOVA™ is 1.17 x more bioavailable by C_{max} and 1.27 by AUC comparison. Under low fat meal conditions, LOVAZA® has only 15% AUC and 12% C_{max} of the bioavailability versus LOVAZA® under high fat meal conditions, whereas EPANOVA™ under low fat meal conditions has 78% AUC and 53% C_{max} of the bioavailability versus LOVAZA® under high fat meal conditions. EPANOVA™ under low fat meal conditions has 62% AUC and 46% C_{max} of the bioavailability versus EPANOVA™ under high fat meal conditions.

Table 2. Comparison of bioavailability of EPA+DHA in Plasma for Lovaza (4g) and Epanova (4g) under high-fat and low-fat meal dosing conditions.

	L20/ASA - High Fat		L20/ASA - Low Fat		Epanova - High Fat		Epanova - Low Fat	
Cross EPA+DHA	385	nmol/ml	85	nmol/ml	450	nmol/ml	395	nmol/ml
Est. AUC ₀₋₂₄ EPA+DHA	3080	nmol*hr/ml	965	nmol*hr/ml	2820	nmol*hr/ml	3425	nmol*hr/ml
Time EPA+DHA	5	hrs	10	hrs	5	hrs	5	hrs
Multiple of Lovaza-HF AUC	1.00	x	0.35	x	1.27	x	0.78	x
Multiple of LF vs. HF AUC	NA		0.35 x Lovaza-HF AUC		NA		0.62 x Epanova-HF AUC	
Multiple of Lovaza-HF Cross	1.00	x	0.12	x	1.27	x	0.55	x
Multiple of LF vs. HF Cross	NA		0.12 x Lovaza-HF Cross		NA		0.46 x Epanova-HF Cross	
Low fat meal - AUC vs. Low	NA		1.00	x	NA		0.28	x
Low fat meal - Cross vs. Low	NA		1.00	x	NA		0.56	x
High fat meal - AUC vs. Low	1.00	x	NA		1.27	x	NA	
High fat meal - Cross vs. Low	1.00	x	NA		1.27	x	NA	

[0019] Omega-3 fatty acids are known to be "essential fatty acids". There are two series of essential fatty acids (EFAs) in humans. They are termed "essential" because they cannot be synthesized de novo in mammals. These fatty acids can be interconverted within a series, but the omega-6 (n-6) series cannot be converted to the omega-3 series nor can the omega-3 (n-3) series be converted to the omega-6 series in humans. The main EFAs in the diet are linoleic acid of the omega-6 series and alpha-linolenic acid of the omega-3 series. However, to fulfill most of their biological effects these "parent" EFAs must be metabolised to the other longer chain fatty acids. Each fatty acid probably has a specific role in the body. The scientific literature suggests that particularly important in the n-6 series are dihomo-gammalinolenic acid (DGLA, 20:3-n6) and arachidonic acid (ARA, 20:4-n6), while particularly important in the n-3 series are eicosapentaenoic acid (EPA, 20:5-n3) and docosahexaenoic acid (DHA, 22:6-n3).

[0020] U.S. Patent No. 6,479,544 describes an invention in which it is found that ARA is highly desirable rather than undesirable and it may be helpful to administer ARA in association with EPA. This invention provides pharmaceutical formulations containing eicosapentaenoic acid or any appropriate derivative (hereinafter collectively referred to as EPA) and arachidonic acid (ARA), as set out in the granted claims for this patent. ARA may be replaced by one or more of its precursors, DGLA or GLA. In this reference, the ratio of EPA to ARA is preferably between 1:1 and 20:1.

[0021] Patent application PCT/GB 2004/000242 describes the treatment or prevention of psoriasis with a formulation comprising more than 95% EPA and less than 2% DHA. In another embodiment of this invention the EPA is replaced with DPA.

[0022] Patent application PCT/NL 2006/050291 (WO/2007/058538, GB 0301701.9) describes combinations of indigestible oligosaccharides and long chain polyunsaturated fatty acids such as ARA, EPA, DHA, and combinations thereof to improve intestinal barrier integrity, improving barrier function, stimulating gut maturation and/or reducing intestinal barrier permeability.

[0023] Given the highly beneficial efficacy and side-effect profile of omega-3 fatty acid concentrates, these compositions are increasingly popular for the treatment of patients with dyslipidemias. However, with the increased popularity of omega-3 fatty acid concentrates, there is an unmet medical need for omega-3 fatty acid containing compositions with improved bioavailability and a more optimal ratio of potency in reducing TG versus the resulting cholesterol profile. Specifically, agents with both a higher potency than AMR101/EPADEL® and lesser increase in LDL-C or further decrease in LDL-C and non-HDL-C than LOVAZA® are required.

SUMMARY OF THE INVENTION

[0024] The present invention provides an orally administrable composition comprising fatty acids, wherein at least 50% by weight of the fatty acids comprise omega-3-fatty acids, salts, esters, or derivatives thereof, wherein the omega-3 fatty acids comprise eicosapentaenoic acid (EPA; C20:5-n3), docosapentaenoic acid (DPA; C22:5-n3), and docosahexaenoic acid (DHA; C22:6-n3), wherein the ratio of DHA to EPA (DHA:EPA) is less than 1:20, and wherein the ratio of DHA to DPA (DHA:DPA) is less than 2:1. In some embodiments, the orally administrable composition comprises additional fatty acids, such as heneicosapentaenoic acid (HPA), arachidonic acid (ARA), and omega-6-docosapentaenoic acid (n-6 DPA), tetracosapentaenoic acid (TPA), and/or gamma-linoleic acid (GLA). These compositions can be used for the treatment or prophylaxis of dyslipidemic, cardiovascular, central nervous system (CNS), inflammatory, and other diseases/conditions or risk factors therefore. These compositions may also be used for the reduction of symptoms associated with these conditions.

[0025] The present invention also provides compositions having an improved profile versus very high purity EPA compositions which contain low levels of omega-6 fatty acids and low levels of omega-3 fatty acids other than EPA, ETA, HPA, TPA, and DPA.

DETAILED DESCRIPTION OF THE INVENTION

[0026] The present invention provides an orally administrable composition comprising fatty acids, wherein at least 50% by weight of the fatty acids comprise omega-3-fatty acids, salts, esters, or derivatives thereof, wherein the omega-3 fatty acids comprise eicosapentaenoic acid (EPA; C20:5-n3), docosapentaenoic acid (DPA; C22:5-n3), and docosahexaenoic acid (DHA; C22:6-n3), wherein the ratio of DHA to EPA (DHA:EPA) is less than 1:20, and wherein the ratio of DHA to DPA (DHA:DPA) is less than 2:1.

[0027] In some embodiments, the compositions of the present invention comprise at least 50% omega-3 fatty acids, alternatively at least 55%, alternatively at least 60%, alternatively at least 65%, alternatively at least 70%, alternatively at least 75%, alternatively at least 80%, alternatively at least 85%, alternatively at least 95%, most preferably at least 90% omega-3 fatty acids of the total amount of fatty acids.

[0028] In other embodiments, EPA and DPA are jointly present in the compositions of the present invention at between 55% and 100% of total fatty acids, alternatively between 60% and 100%, alternatively between 65% and 100%, alternatively between 70% and 100%, alternatively between 75% and 100%, alternatively between 80% and 100%, alternatively between 85% and 95%, alternatively between 85% and 97%, alternatively between 88% and 95%, alternatively between 88% and 97%, alternatively between 90% and 95%, alternatively between 90% and 97% of the total amount of fatty acids.

[0029] The fatty acids, such as EPA and DPA, may be present in free fatty acid form, or as a salt, ester, or derivative. The fatty acids are preferably composed as a triglyceride, an ester (such as an ethyl ester) or free fatty acid. Other forms of the fatty acids which may be useful include salts, esters of any type, amides, mono-, di- or triglycerides, phospholipids or any other form which can lead to metabolization of the fatty acids (such as EPA and/or DPA), or the incorporation of the fatty acids (such as EPA and/or DPA) into body fluids, tissues or organs.

[0030] Omega-3 fatty acids may be grouped by the number of double bonds contained in the fatty acid chain. For instance, hexadecatrienoic acid (HTA), alpha-linolenic acid (ALA) and eicosatrienoic acid (ETE) are omega-3-trienoic acids; stearidonic acid (SDA) and eicosatetraenoic acid (ETA) are omega-3-tetraenoic acids; EPA, heneicosapentaenoic acid (HPA), DPA and tetracosapentaenoic acid

(TPA) are omega-3-pentaenoic acids; and DHA and tetracosahexaenoic acid (THA) are omega-3-hexaenoic acids. In some preferred embodiments, the term omega-3-pentaenoic acids will refer to a mixture of at least two omega-3 pentaenoic acids in a ratio of at least 1:25, more preferably in a ratio of at least 1:50, more preferably in a ratio of at least 1:75, more preferably in a ratio of at least 1:100, more preferably in a ratio of at least 1:125, more preferably in a ratio of at least 1:150, more preferably in a ratio of at least 1:200. In some embodiments, the ratio refers to the ratio of the least prevalent omega-3 pentaenoic acid in the mixture to the most prevalent omega-3 pentaenoic acid in the mixture.

[0031] In some embodiments, the compositions of the present invention comprise EPA, HPA, DPA and TPA, alternatively EPA and DPA, and alternatively the compositions of the present invention comprise EPA, HPA and DPA.

[0032] In some embodiments, the omega-3-pentaenoic acids in the compositions of the present invention comprise no more than 99.5% of a single omega-3-pentaenoic acid, alternatively no more than 99%; alternatively no more than 98.5%; alternatively no more than 98%; alternatively no more than 97.5%; alternatively no more than 96%; alternatively no more than 95%; alternatively no more than 94%; alternatively no more than 93%; alternatively no more than 92%; alternatively no more than 91%; alternatively no more than 90%; alternatively no more than 88%; alternatively no more than 85%; alternatively no more than 80%; alternatively no more than 75%; alternatively no more than 70%; alternatively no more than 65%; alternatively no more than 60%; alternatively no more than 55%; alternatively no more than 50%; alternatively no more than 45%; alternatively no more than 40%; alternatively no more than 30%.

[0033] In some embodiments, the compositions of the present invention wherein at least 10%, alternatively at least 20%, alternatively at least 25%, alternatively at least 35%, alternatively at least 50%, alternatively at least 60%, alternatively at least 65%, alternatively at least 70%, alternatively at least 75%, by weight of the fatty acids comprise omega-3-pentaenoic acids, salts, esters, or derivatives thereof.

[0034] In some embodiments, the compositions of the present invention comprise at least 0.01% HPA of total fatty acids in the composition, alternatively at least 0.05% HPA, alternatively at least 0.10% HPA, alternatively at least 0.15% HPA, alternatively at least 0.2% HPA, alternatively at least 0.3% HPA, alternatively at least 0.4% HPA, alternatively at least 0.5% HPA, alternatively at least 0.75% HPA,

alternatively at least 1% HPA, alternatively at least 1.5% HPA, alternatively at least 2% HPA, alternatively at least 2.5% HPA, alternatively at least 3% HPA, alternatively at least 3.5% HPA, alternatively at least 4% HPA, alternatively at least 4.5% HPA, alternatively at least 5% HPA, alternatively at least 6% HPA, alternatively at least 7% HPA, alternatively the compositions of the present invention comprise at least 9% HPA of total fatty acids in the composition.

In some embodiments, the compositions of the present invention comprise no more than 20% HPA of total fatty acids in the composition, alternatively no more than 15% HPA, alternatively no more than 12% HPA, alternatively no more than 10% HPA, alternatively no more than 8% HPA, alternatively no more than 7% HPA, alternatively no more than 6% HPA, alternatively no more than 5% HPA, alternatively no more than 4% HPA, alternatively no more than 3% HPA, alternatively no more than 2% HPA, alternatively no more than 1.5% HPA, alternatively the compositions of the present invention comprise at least 1% HPA of total fatty acids in the composition. In some embodiments, the compositions of the present invention comprise 1% to 20% of the total fatty acids in the composition.

[0035] In some embodiments, the compositions of the present invention comprise no more than 10% omega-3 fatty acids that are not omega-3-pentaenoic acids, alternatively no more than 9%, alternatively no more than 8%, alternatively no more than 7%, alternatively no more than 6%, alternatively no more than 5%, alternatively no more than 4.5%, alternatively no more than 4%, alternatively no more than 3.5%, alternatively no more than 3%, alternatively no more than 2.5%, alternatively no more than 2%, alternatively no more than 1.5%, alternatively no more than 1.25%, alternatively no more than 1%, alternatively no more than 0.75%, alternatively no more than 0.5%, alternatively no more than 0.4%, alternatively no more than 0.3%, alternatively no more than 0.2%, alternatively the compositions of the present invention comprise no more than 0.1% omega-3 fatty acids that are not omega-3-pentaenoic acids.

[0036] In the embodiments of the present invention, the compositions comprise EPA and DPA in an EPA:DPA ratio between 99:1 and 1:99 EPA:DPA, alternatively between 90:1 and 1:90, alternatively between 60:1 and 1:60, alternatively between 60:1 and 1:20, alternatively between 60:1 and 1:4, alternatively between 40:1 and 1:20, alternatively between 30:1 and 1:20, alternatively between 30:1 and 1:10, alternatively between 30:1 and 1:5, alternatively between 40:1 and 1:4, alternatively

between 30:1 and 1:4, alternatively between 30:1 and 1:2, alternatively between 30:1 and 1:1, alternatively between 30:1 and 2:1, alternatively between 30:1 and 5:1, alternatively between 20:1 and 1:20, alternatively between 20:1 and 1:10, alternatively between 20:1 and 1:5, alternatively between 20:1 and 1:2, alternatively between 20:1 and 1:1, alternatively between 20:1 and 2:1, alternatively between 20:1 and 5:1, alternatively between 20:1 and 10:1, alternatively between 20:1 and 10:1, alternatively between 30:1 and 10:1, alternatively between 60:1 and 10:1, alternatively comprise EPA and DPA in an EPA:DPA ratio between 40:1 and 10:1. In some embodiments, the ratio of EPA:DPA is greater than 1:1, preferably greater than 2:1, and more preferably greater than 5:1. In some embodiments, the ratio of EPA:DPA is 1:1 to 25:1, preferably 5:1 to 20:1, more preferably 8:1 to 15:1, even more preferably 9:1 to 13:1, even more most preferably about 10:1 to 11:1, and most preferably about 10:1.

[0037] In some embodiments of the present invention, the compositions comprise EPA in an amount between 55% and 95% relative to the total amount of fatty acids present in the composition, alternatively between 60% and 95%, alternatively between 65% and 95%, alternatively between 70% and 95%, alternatively between 75% and 95%, alternatively between 90% and 95%, alternatively between 80% and 95%, alternatively between 90% and 95%, alternatively between 55% and 90%, alternatively between 60% and 90%, alternatively between 65% and 90%, alternatively between 70% and 90%, alternatively between 75% and 90%, alternatively between 80% and 90%, alternatively between 85% and 90%, alternatively between 55% and 92%, alternatively between 60% and 92%, alternatively between 65% and 92%, alternatively between 70% and 92%, alternatively between 75% and 92%, alternatively between 80% and 92%, alternatively between 85% and 92%, alternatively between 55% and 93%, alternatively between 60% and 93%, alternatively between 65% and 93%, alternatively between 70% and 93%, alternatively between 75% and 93%, alternatively between 80% and 93%, alternatively between 85% and 93%, alternatively more than 85%, alternatively more than 85%, alternatively between 85% and 95% EPA relative to the total amount of fatty acids present in the composition.

[0038] In other embodiments of the present invention, the compositions comprise DPA in an amount between 1% and 99% relative to the total amount of fatty acids

present in the composition, alternatively between 1% and 95%, alternatively
 alternatively between 1% and 90%, alternatively alternatively between 1% and 85%,
 alternatively between 1% and 80%, alternatively alternatively between 1% and 75%,
 alternatively between 1% and 70%, alternatively between 1% and 65%,
 alternatively between 1% and 60%, alternatively between 1% and 55%,
 alternatively between 1% and 50%, alternatively between 1% and 45%,
 alternatively between 1% and 40%, alternatively between 1% and 35%,
 alternatively between 1% and 30%, alternatively between 1% and 25%,
 alternatively between 1% and 20%, alternatively between 1% and 15%,
 alternatively between 1% and 10%, alternatively between 1% and 5%, alternatively
 between 2% and 99%, alternatively between 2% and 95%, alternatively between 2%
 and 90%, alternatively between 2% and 85%, alternatively between 2% and 80%,
 alternatively between 2% and 75%, alternatively between 2% and 70%,
 alternatively between 2% and 65%, alternatively between 2% and 60%,
 alternatively between 2% and 55%, alternatively between 2% and 50%,
 alternatively between 2% and 45%, alternatively between 2% and 40%,
 alternatively between 2% and 35%, alternatively between 2% and 30%,
 alternatively between 2% and 25%, alternatively between 2% and 20%,
 alternatively between 2% and 15%, alternatively between 2% and 10%,
 alternatively between 2% and 5%, alternatively between 3% and 99%, alternatively
 between 3% and 95%, alternatively between 3% and 90%, alternatively between
 3% and 85%, alternatively between 3% and 80%, alternatively between 3% and
 75%, alternatively between 3% and 70%, alternatively between 3% and 65%,
 alternatively between 3% and 60%, alternatively between 3% and 55%,
 alternatively between 3% and 50%, alternatively between 3% and 45%,
 alternatively between 3% and 40%, alternatively between 3% and 35%,
 alternatively between 3% and 30%, alternatively between 3% and 25%,
 alternatively between 3% and 20%, alternatively between 3% and 15%,
 alternatively between 3% and 10%, alternatively between 3% and 5%, alternatively
 between 4% and 99%, alternatively between 4% and 95%, alternatively between 4%
 and 90%, alternatively between 4% and 85%, alternatively between 4% and 80%,
 alternatively between 4% and 75%, alternatively between 4% and 70%,
 alternatively between 4% and 65%, alternatively between 4% and 60%,
 alternatively between 4% and 55%, alternatively between 4% and 50%,

alternatively between 4% and 45%, alternatively between 4% and 40%,
 alternatively between 4% and 35%, alternatively between 4% and 30%,
 alternatively between 4% and 25%, alternatively between 4% and 20%,
 alternatively between 4% and 15%, alternatively between 4% and 10%,
 alternatively between 4% and 5%, alternatively between 5% and 99%, alternatively
 between 5% and 95%, alternatively between 5% and 90%, alternatively between
 5% and 85%, alternatively between 5% and 80%, alternatively between 5% and
 75%, alternatively between 5% and 70%, alternatively between 5% and 65%,
 alternatively between 5% and 60%, alternatively between 5% and 55%,
 alternatively between 5% and 50%, alternatively between 5% and 45%,
 alternatively between 5% and 40%, alternatively between 5% and 35%,
 alternatively between 5% and 30%, alternatively between 5% and 25%,
 alternatively between 5% and 20%, alternatively between 5% and 15%,
 alternatively between 5% and 12%, alternatively between 5% and 10%,
 alternatively between 6% and 99%, alternatively between 6% and 95%, alternatively
 between 6% and 90%, alternatively between 6% and 85%, alternatively between
 6% and 80%, alternatively between 6% and 75%, alternatively between 6% and
 70%, alternatively between 6% and 65%, alternatively between 6% and 60%,
 alternatively between 6% and 55%, alternatively between 6% and 50%,
 alternatively between 6% and 45%, alternatively between 6% and 40%,
 alternatively between 6% and 35%, alternatively between 6% and 30%,
 alternatively between 6% and 25%, alternatively between 6% and 20%,
 alternatively between 6% and 15%, alternatively between 6% and 12%,
 alternatively between 6% and 11%, alternatively between 6% and 10%,
 alternatively between 7% and 99%, alternatively between 7% and 95%, alternatively
 between 7% and 90%, alternatively between 7% and 85%, alternatively between
 7% and 80%, alternatively between 7% and 75%, alternatively between 7% and
 70%, alternatively between 7% and 65%, alternatively between 7% and 60%,
 alternatively between 7% and 55%, alternatively between 7% and 50%,
 alternatively between 7% and 45%, alternatively between 7% and 40%,
 alternatively between 7% and 35%, alternatively between 7% and 30%,
 alternatively between 7% and 25%, alternatively between 7% and 20%,
 alternatively between 7% and 15%, alternatively between 7% and 12%,
 alternatively between 7% and 11%, alternatively between 7% and 10%,

alternatively between 8% and 99%, alternatively between 8% and 95%, alternatively between 8% and 90%, alternatively between 8% and 85%, alternatively between 8% and 80%, alternatively between 8% and 75%, alternatively between 8% and 70%, alternatively between 8% and 65%, alternatively between 8% and 60%, alternatively between 8% and 55%, alternatively between 8% and 50%, alternatively between 8% and 45%, alternatively between 8% and 40%, alternatively between 8% and 35%, alternatively between 8% and 30%, alternatively between 8% and 25%, alternatively between 8% and 20%, alternatively between 8% and 15%, alternatively between 8% and 12%, alternatively between 2% and 99%, alternatively between 9% and 95%, alternatively between 9% and 90%, alternatively between 9% and 85%, alternatively between 9% and 80%, alternatively between 9% and 75%, alternatively between 9% and 70%, alternatively between 9% and 65%, alternatively between 9% and 60%, alternatively between 9% and 55%, alternatively between 9% and 50%, alternatively between 9% and 45%, alternatively between 9% and 40%, alternatively between 9% and 35%, alternatively between 9% and 30%, alternatively between 9% and 25%, alternatively between 9% and 20%, alternatively between 9% and 15%, alternatively between 9% and 12%, relative to the total amount of fatty acids present in the composition.

[0039] In some embodiments, a relatively small amount of DHA as compared to EPA is present. In some embodiments, the compositions of the present invention comprise no more than 1:5 of DHA:EPA, alternatively no more than 1:6 of DHA:EPA, alternatively no more than 1:7 of DHA:EPA, alternatively no more than 1:8 of DHA:EPA, alternatively no more than 1:9 of DHA:EPA, alternatively no more than 1:10 of DHA:EPA, alternatively no more than 1:12 of DHA:EPA, alternatively no more than 1:15 of DHA:EPA, alternatively no more than 1:20 of DHA:EPA, alternatively no more than 1:25 of DHA:EPA, alternatively no more than 1:30 of DHA:EPA, alternatively no more than 1:40 of DHA:EPA, alternatively no more than 1:50 of DHA:EPA, alternatively no more than 1:75 of DHA:EPA, alternatively no more than 1:90 of DHA:EPA, alternatively no more than 1:99 of DHA:EPA. Alternatively, DHA may be present in the compositions of this invention at a relative amount of ratio less than 1% than the amount of EPA. Alternatively, DHA may be present in the compositions of this invention at a DHA:EPA ratio of less than 1:99.

[0040] In some embodiments, a relatively small amount of DHA relative to the total amount of fatty acids present in the composition is present. In some embodiments, the compositions of the present invention comprise no more than 20% DHA, alternatively no more than 15% DHA, alternatively no more than 12% DHA, alternatively no more than 10% DHA, alternatively no more than 8% DHA, alternatively no more than 7% DHA, alternatively no more than 6% DHA, alternatively no more than 5% DHA, alternatively no more than 4% DHA, alternatively no more than 3% DHA, alternatively no more than 2% DHA, alternatively no more than 1% DHA relative to the total amount of fatty acids present in the composition.

[0041] In some embodiments, the ratio of EPA:HPA is about 1500:1 to 25:1, alternatively 1000:1 to 50:1, alternatively 800:1 to 60:1, alternatively 500:1 to 50:1, alternatively 250:1 to 75:1, and alternatively 100:1 to 80:1. In some preferred embodiments, the ratio of EPA:HPA is about 85:1. In some preferred embodiments, the ratio of EPA:HPA is about 30:1. In some embodiments, the ratio of DPA:HPA is about 250:1 to 1:1, alternatively 200:1 to 2:1, alternatively 150:1 to 3:1, alternatively 100:1 to 4:1, alternatively 50:1 to 5:1, alternatively 25:1 to 6:1, and alternatively 10:1 to 7:1. In some preferred embodiments, the ratio of DPA:HPA is about 8:1. In some embodiments, the ratio of DPA:HPA is about 3:0.

[0042] In other embodiments, a relatively small amount of DHA as compared to DPA is present. In these embodiments, the compositions of the present invention comprise no more than 15:1 of DHA:DPA, alternatively no more than 12:1 of DHA:DPA, alternatively no more than 10:1 of DHA:DPA, alternatively no more than 8:1 of DHA:DPA, alternatively no more than 5:1 of DHA:DPA, alternatively no more than 3:1 of DHA:DPA, alternatively no more than 2:1 of DHA:DPA, alternatively no more than 1:1 of DHA:DPA, alternatively no more than 1:2 of DHA:DPA, alternatively no more than 1:3 of DHA:DPA, alternatively no more than 1:5 of DHA:DPA, alternatively no more than 1:8 of DHA:DPA, alternatively no more than 1:10 of DHA:DPA, alternatively no more than 1:15 of DHA:DPA, alternatively no more than 1:20 of DHA:DPA, alternatively no more than 1:25 of DHA:DPA, alternatively no more than 1:50 of DHA:DPA, alternatively no more than 1:75 of DHA:DPA, alternatively no more than 1:90 of DHA:DPA, alternatively no more than 1:95 of DHA:DPA, alternatively no more than 1:100 of DHA:DPA. In some embodiments, the ratio of DHA:DPA is preferably less than 2:1.

[0043] In other embodiments, a relatively small amount of DHA as compared to HPA is present. In these embodiments, the compositions of the present invention comprise no more than 15:1 of DHA:HPA, alternatively no more than 12:1 of DHA:HPA, alternatively no more than 10:1 of DHA:HPA, alternatively no more than 8:1 of DHA:HPA, alternatively no more than 5:1 of DHA:HPA, alternatively no more than 3:1 of DHA:HPA, alternatively no more than 2:1 of DHA:HPA, alternatively no more than 1:1 of DHA:HPA, alternatively no more than 1:2 of DHA:HPA, alternatively no more than 1:3 of DHA:HPA, alternatively no more than 1:5 of DHA:HPA, alternatively no more than 1:8 of DHA:HPA, alternatively no more than 1:10 of DHA:HPA, alternatively no more than 1:15 of DHA:HPA, alternatively no more than 1:20 of DHA:HPA, alternatively no more than 1:25 of DHA:HPA, alternatively no more than 1:50 of DHA:HPA, alternatively no more than 1:75 of DHA:HPA, alternatively no more than 1:90 of DHA:HPA, alternatively no more than 1:95 of DHA:HPA, alternatively no more than 1:100 of DHA:HPA.

[0044] In yet other embodiments, the compositions of the present invention comprise no more than 10% omega-6 fatty acids relative to the total amount of fatty acids, alternatively no more than 9%, alternatively no more than 8%, alternatively no more than 7%, alternatively no more than 6%, alternatively no more than 5%, alternatively no more than 4.5%, alternatively no more than 4%, alternatively no more than 3.5%, alternatively no more than 3%, alternatively no more than 2.5%, alternatively no more than 2%, alternatively no more than 1.7%, alternatively no more than 1.5%, alternatively no more than 1.2%, alternatively no more than 1%, alternatively no more than 0.5% omega-6 fatty acids versus the total amount of fatty acids comprised by the compositions of the present invention.

[0045] Omega-6 fatty acids include, but are not limited to: linoleic acid (LA; C18:2-n6); gamma-linoleic acid (GLA; C18:3-n6); eicosadienoic acid (C20:2-n6); dihomo-gamma-linoleic acid (DGLA; C20:3-n6); arachidonic acid (ARA; C20:4-n6); and omega-6 docosapentaenoic acid (DPA; C22:5-n6).

[0046] In further embodiments, the compositions of the present invention comprise no more than 10% omega-6 fatty acids relative to the total amount of omega-3 fatty acids plus omega-6 fatty acids, alternatively no more than 9%, alternatively no more than 8%, alternatively no more than 7%, alternatively no more than 6%, alternatively no more than 5%, alternatively no more than 4.5%, alternatively no more than 4%, alternatively no more than 3.5%, alternatively no more than 3%, alternatively no

more than 2.5%, alternatively no more than 2%, alternatively no more than 1.7%, alternatively no more than 1.5%, alternatively no more than 1.2%, alternatively no more than 1%, alternatively no more than 0.5% omega-6 fatty acids versus the total amount of omega-3 fatty acids plus omega-6 fatty acids comprised by the compositions of the present invention.

[0047] In yet other embodiments, the compositions of the present invention comprise no more than 8% arachidonic acid (ARA; C20:4-n6) relative to the total amount of omega-3 fatty acids plus omega-6 fatty acids, alternatively no more than 7%, alternatively no more than 6%, alternatively no more than 5%, alternatively no more than 4.5%, alternatively no more than 4%, alternatively no more than 3.5%, alternatively no more than 3%, alternatively no more than 2.5%, alternatively no more than 2%, alternatively no more than 1.7%, alternatively no more than 1.5%, alternatively no more than 1.2%, alternatively no more than 1%, alternatively no more than 0.5% arachidonic acid (ARA; C20:4-n6) versus the total amount of omega-3 fatty acids plus omega-6 fatty acids comprised by the compositions of the present invention.

[0048] In some embodiments, a relatively small amount of omega-3 fatty acids in aggregate other than EPA, ETA, HPA and DPA (alternatively indicated as non-EPA, non-ETA, non-HPA and non-DPA omega-3 fatty acids in aggregate) relative to the total amount of fatty acids present in the composition is present. In some embodiments, the compositions of the present invention comprise no more than 20% non-EPA, non-ETA, non-HPA and non-DPA omega-3 fatty acids, alternatively no more than 15% non-EPA, non-ETA, non-HPA and non-DPA omega-3 fatty acids, alternatively no more than 12% non-EPA, non-ETA, non-HPA and non-DPA omega-3 fatty acids, alternatively no more than 10% non-EPA, non-ETA, non-HPA and non-DPA omega-3 fatty acids, alternatively no more than 8% non-EPA, non-ETA, non-HPA and non-DPA omega-3 fatty acids, alternatively no more than 7% non-EPA, non-ETA, non-HPA and non-DPA omega-3 fatty acids, alternatively no more than 6% non-EPA, non-ETA, non-HPA and non-DPA omega-3 fatty acids, alternatively no more than 5% non-EPA, non-ETA, non-HPA and non-DPA omega-3 fatty acids, alternatively no more than 4% non-EPA, non-ETA, non-HPA and non-DPA omega-3 fatty acids, alternatively no more than 3% non-EPA, non-ETA, non-HPA and non-DPA omega-3 fatty acids, alternatively no more than 2% non-EPA, non-ETA, non-HPA and non-DPA omega-3 fatty acids, alternatively no more than 1% non-EPA,

non-ETA, non-HPA and non-DPA omega-3 fatty acids in aggregate relative to the total amount of fatty acids present in the composition.

[0049] In some embodiments, a relatively small amount of the sum of ALA, SDA and DHA relative to the total amount of fatty acids present in the composition is present, while at the same time large amounts of the sum of EPA, DPA-n3, HPA and ETA are present. In some embodiments, the compositions of the present invention comprise no more than 20% of the sum of ALA, SDA and DHA, alternatively no more than 15% of the sum of ALA, SDA and DHA, alternatively no more than 12% of the sum of ALA, SDA and DHA, alternatively no more than 10% of the sum of ALA, SDA and DHA, alternatively no more than 8% of the sum of ALA, SDA and DHA, alternatively no more than 7% of the sum of ALA, SDA and DHA, alternatively no more than 6% of the sum of ALA, SDA and DHA, alternatively no more than 5% of the sum of ALA, SDA and DHA, alternatively no more than 4% of the sum of ALA, SDA and DHA, alternatively no more than 3% of the sum of ALA, SDA and DHA, alternatively no more than 2% of the sum of ALA, SDA and DHA, alternatively no more than 1% of the sum of ALA, SDA and DHA relative to the total amount of fatty acids present in the composition, while at the same time contain more than 40% the sum of EPA, DPAn-3, HPA and ETA, alternatively more than 50% the sum of EPA, DPAn-3, HPA and ETA, , alternatively more than 60% the sum of EPA, DPAn-3, HPA and ETA, alternatively more than 70% the sum of EPA, DPAn-3, HPA and ETA, alternatively more than 75% the sum of EPA, DPAn-3, HPA and ETA, alternatively more than 80% the sum of EPA, DPAn-3, HPA and ETA, alternatively more than 85% the sum of EPA, DPAn-3, HPA and ETA, alternatively more than 90% the sum of EPA, DPAn-3, HPA and ETA, alternatively more than 95% the sum of EPA, DPAn-3, HPA and ETA, alternatively between 80% and 98% the sum of EPA, DPAn-3, HPA and ETA, alternatively between 80% and 95% the sum of EPA, DPAn-3, HPA and ETA, alternatively between 85% and 98% the sum of EPA, DPAn-3, HPA and ETA, alternatively between 85% and 96% the sum of EPA, DPAn-3, HPA and ETA, alternatively between 90% and 98% the sum of EPA, DPAn-3, HPA and ETA, alternatively between 90% and 97% the sum of EPA, DPAn-3, HPA and ETA, alternatively between 90% and 96% the sum of EPA, DPAn-3, HPA and ETA, alternatively between 90% and 95% the sum of EPA, DPAn-3, HPA and ETA, relative to the total amount of fatty acids present in the composition is present.

[0050] In further embodiments, the compositions of the present invention comprise no more than 8% arachidonic acid (ARA; C20:4-n6) relative to the total amount of fatty acids, alternatively no more than 7%, alternatively no more than 6%, alternatively no more than 5%, alternatively no more than 4.5%, alternatively no more than 4%, alternatively no more than 3.5%, alternatively no more than 3%, alternatively no more than 2.5%, alternatively no more than 2%, alternatively no more than 1.7%, alternatively no more than 1.5%, alternatively no more than 1.2%, alternatively no more than 1%, alternatively no more than 0.5% arachidonic acid (ARA; C20:4-n6) relative to the total amount of fatty acids comprised by the compositions of the present invention.

[0051] In other embodiments, the compositions of the present invention comprise no more than 2.5% arachidonic acid (ARA; C20:4-n6), no more than 0.4% omega-6-docosapentaenoic acid (DPA; C22:5-n6) and no more than 0.2% gamma-linoleic acid (GLA; C18:3-n6) relative to the total amount of fatty acids comprised by the compositions of the present invention.

[0052] Further embodiments provide fatty acid compositions comprising no more than 2.5% arachidonic acid (ARA; C20:4-n6), no more than 0.3% omega-6-docosapentaenoic acid (DPA; C22:5-n6) and no more than 0.1% gamma-linoleic acid (GLA; C18:3-n6) relative to the total amount of fatty acids comprised by the compositions of the present invention.

[0053] In yet other embodiments, the active ingredient of the formulations of the present invention consists essentially wholly of the EPA and DPA or precursors thereof (ethyl ester, triglyceride, or any other pharmaceutically acceptable salt or derivative thereof). In that case, no large amounts (preferably less than 15%, alternatively less than 12%, alternatively less than 10%, alternatively less than 9%, alternatively less than 8%, alternatively less than 7%, alternatively less than 6%, alternatively less than 5%, alternatively less than 4%, alternatively less than 3%, alternatively less than 2%, alternatively less than 1%, alternatively less than 0.5%, alternatively less than 0.25%) of any other fatty acids are present.

[0054] In further embodiments, the active ingredient of the formulations of the present invention consists essentially wholly of omega-3-pentaenoic acids or precursors thereof (ethyl ester, triglyceride, or any other pharmaceutically acceptable salt or derivative thereof). In that case, no large amounts (preferably less than 15%, alternatively less than 12%, alternatively less than 10%, alternatively less than 9%,

alternatively less than 4%, alternatively less than 4%, alternatively less than 4%, alternatively less than 4%, alternatively less than 4%, alternatively less than 3%, alternatively less than 2%, alternatively less than 1%, alternatively less than 0.5%, alternatively less than 0.25%) of any other fatty acids are present.

[0055] The fatty acid percentage is determined on a weight/weight, mol/mol, or chromatography area percent basis relative to all fatty acids present in the composition as determined by methods such as disclosed in the European Pharmacopeia monograph for omega-3 fatty acid concentrates, European Pharmacopeia monograph for omega-3-acid ethyl esters 90%, or European Pharmacopeia monograph method 2.4.29, USP monograph for fish oil dietary supplements, USP 35 omega-3-acid ethyl esters (LOVAZA®) monograph, or any essentially equivalent methods (whether by gas chromatography, HPLC, FPLC or any other chromatographic method).

[0056] In some embodiments, the fatty acid percentage is determined not as a percentage of all fatty acids present in the composition but as a specific type of fatty acid ethyl esters as percentage of all fatty acid ethyl esters present in the composition, thus excluding from the fatty acid percentage determination such fatty acids present as, for instance: free fatty acids; mono-, di-, and tri-glycerides; or fatty acids present in phospholipids (such as phosphatidylserine or phosphatidylcholine) or polysorbates (such as Tween 80, Tween 20, or polysorbate 40).

[0057] In other embodiments, the fatty acid percentage is determined not as a percentage of all fatty acids present in the composition but as a specific type of free fatty acid as percentage of all free fatty acids present in the composition, thus excluding from the fatty acid percentage determination such fatty acids present as, for instance: fatty acid ethyl esters; mono-, di-, and tri-glycerides; or fatty acids present in phospholipids (such as phosphatidylserine or phosphatidylcholine) or polysorbates (such as Tween 80, Tween 20, or polysorbate 40).

[0058] In yet other embodiments, the fatty acid percentage is determined not as a percentage of all fatty acids present in the composition but as a specific type of glycerol fatty acid ester as percentage of all glycerol fatty acid esters present in the composition, thus excluding from the fatty acid percentage determination such fatty acids present as, for instance: fatty acid ethyl esters; free fatty acids; or fatty acids present in phospholipids (such as phosphatidylserine or phosphatidylcholine) or polysorbates (such as Tween 80, Tween 20, or polysorbate 40).

[0059] In further embodiments, the fatty acid percentage is determined not as a percentage of all fatty acids present in the composition but as di- or tri-fatty acid esters with glycerol as percentage of all glycerol di- and tri-fatty acid esters present in the composition, thus excluding from the fatty acid percentage determination such fatty acids present as, for instances: glycerol-mono-fatty acid esters; fatty acid ethyl esters; free fatty acids; or fatty acids present in phospholipids (such as phosphatidylserine or phosphatidylcholine) or polysorbates (such as Tween 80, Tween 20, or polysorbate 40).

[0060] In yet other embodiments, the fatty acid percentage is determined not as a percentage of all fatty acids present in the composition but as a tri-fatty acid esters with glycerol as percentage of all glycerol tri-fatty acid esters present in the composition, thus excluding from the fatty acid percentage determination such fatty acids present as, for instance: mono- and di-fatty acid esters of glycerol; fatty acid ethyl esters; free fatty acids; or fatty acids present in phospholipids (such as phosphatidylserine or phosphatidylcholine) or polysorbates (such as Tween 80, Tween 20, or polysorbate 40).

[0061] The EPA, HPA, DPA, or omega-3-pentaenoic acids may be derived from any appropriate source including plant seed oils, microbial oils from algae or fungal or marine oils from fish or other marine animals. Certain species are a particular good source of oils containing DPA, for example seal oil. They may be used in the form of the natural oil, if that oil meets the required purity requirements of the present invention, or may be purified to give products containing the fatty acid composition of the present invention.

[0062] The compositions of the present invention may be produced through a range of the methods. Such methods may include: distillation, including short path distillation; urea precipitation; enzymatic conversion concentration; conventional chromatography; HPLC/FPLC; supercritical carbon dioxide extraction; supercritical carbon dioxide chromatography; simulated moving bed chromatography; supercritical carbon dioxide simulated moving bed chromatography; or chemical conversion methods such as iodolactonization. Such methods are generally known to those skilled in the art of purifying and isolating omega-3 fatty acids.

[0063] Typically, the omega-3 fatty acid concentration/purification process is initiated by esterifying the fatty acids comprised by the marine oil raw material (such as crude fish oil) with ethanol (to form fatty acid ethyl esters) in order to separate

omega-3 fatty acids from other fatty acids covalently bound together in the natural triglyceride molecules of the source oil. Subsequently, the material may be distilled once or several times to achieve omega-3-acid ethyl ester concentrations above 60%-70%. Alternatively, enzymatic concentration, urea precipitation or supercritical extraction may be used alone or in conjunction with distillation to reach omega-3 levels above 70%-90%. In order to prepare a highly pure concentrate of a single omega-3 fatty acid, methods such as chromatography, supercritical chromatography, simulated moving bed chromatography, supercritical simulated moving bed chromatography, or chemical conversion methods such as iodolactolization are typically most practical to reach levels above 50%, alternatively above 60%, alternatively above 70%, alternatively above 80%, alternatively above 90%, alternatively above 95%, of a single omega-3 fatty acid such as ETA, EPA, HPA, DPA, TPA, or DHA.

[0064] Those skilled in the art will be able to design processes suited to prepare a certain omega-3 fatty acid composition as desired, based on the methods described above. Such processes are flexible enough to affect the relative proportions between the long chain C18, C20, C21 and C22 fatty acids which occur naturally in available fish oil raw materials and other marine oils. It provides not only for the concentration of the individual omega-3 fatty acids, but the ratio between them will remain within a pattern of variation caused by variations in nature. However, suitable methods compensate for sometimes extreme variations which may occur naturally. Thus, for those skilled in the art, it will be possible to make a product with a constant and predetermined composition.

[0065] EPA is relatively abundant in fish oils or other marine oils and can be relatively easily obtained through the application of concentration and purification technologies from such fish or marine oils. DPA and HPA are present at much lower concentrations. In order to prepare the compositions of the present invention, DPA or HPA may be concentrated and purified from fish or other marine oils according to the methods referred to above, either alone or DPA combined with EPA and/or HPA. Alternatively, the DPA or HPA may be chemically prepared from a high purity EPA concentrate by elongation of the EPA fatty-acid chain with two or one hydrogen-saturated carbons (C2-elongation or C1-elongation) on the carboxyl side of the molecule (for instance with a method similar to or alternate methods with equivalent results such as described by Kuklev DV and Smith WL in Chem Phys Lipids, 2006;

144(2); 172-177). In another alternative approach, a high purity EPA concentrate may be partially converted to DPA (or HPA) using a method for C2-elongation (or C1-elongation) of EPA similar to those described above, thus directly yielding compositions of the present invention or intermediates therefore.

[0066] Once the oils containing one or more of the desired fatty acids have been obtained, and purified as necessary, these oils may be blended to give the desirable relative amounts of EPA, DPA, HPA, DHA, TPA, other omega-3 fatty acids and omega-6 fatty acids to obtain the compositions of the present invention described in detail above.

[0067] Fish oils may also contain by-products and contaminants such as pesticides, chlorinated or brominated hydrocarbons, heavy metals, cholesterol and vitamins. During the production of the concentrate, the concentrations of these components are significantly reduced compared to untreated fish oils. Such reduction is inherent due to the nature of purification methods and their ability to concentrate of several or specific omega-3 fatty acids, thus removing other compounds.

[0068] Triglycerides comprising more than 60% of the omega-3 fatty acids in the composition may be produced from ethyl esters and glycerol by well known, published, or alternative chemical synthetic or enzymatic procedures. The free acids may be produced from ethyl esters by well known hydrolyzation or saponification procedures. Methods for converting ethyl esters to triglycerides, free fatty acids, and other molecular forms comprising fatty acids, are generally known to those skilled in the art chemically or enzymatically converting omega-3 fatty acids from one form to another.

[0069] In some embodiments, the compositions of the present invention have improved pharmacological features as demonstrated by improved bioavailability in a mammal of EPA, HPA, DPA, DHA, EPA+DHA, EPA+DPA or EPA+HPA+DPA combined, total omega-3-pentaenoic acids, or of total omega-3 fatty acids. Key parameters for determining bioavailability are maximum concentration of a therapeutic compound or a metabolite thereof (C_{max}); the time from administration to maximum concentration (T_{max}); and the area under the concentration curve over time (AUC). Such parameters may be determined under single dose or multiple dose administration regimens. Methods to determine comparative bioavailability in mammals are generally known to those skilled in the art. When comparing T_{max}, C_{max}, and AUC for embodiments of the present invention to LOVAZA®,

EPANOVA™, and AMR101 throughout this application, such comparison will be on the basis of an equal dose of 4 capsules of 1 gram each for each of these products. The comparative parameters, however, do apply to all essentially equivalent dosing modes comparing embodiments of the present invention to LOVAZA®, EPANOVA™, and AMR101.

[0070] Meal conditions during administration to a subject of omega-3 fatty acid compositions or omega-3 fatty acid formulations are of special significance for absorption and bioavailability of omega-3 fatty acids. The meal conditions typically considered are: fasting (no food at all prior for 8-12 hours prior to administration and 2-3 hours post administration of the treatment); a low fat meal (a meal typically containing less than 25 gram of fat [350-600 Kcal] consumed just before or after the administration of the treatment; typically within a 15-30 minute range); or a high fat meal (a meal containing 40 gram to 75 gram of fat [700-1000 Kcal] consumed just before or after the administration of the treatment; typically within a 15-30 minute range).

[0071] In some embodiments of the present invention, compositions of the present invention are more rapidly absorbed as measured by the time to reach the maximum concentration (Tmax) in blood, serum or plasma of EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids. In preferred embodiments of the present invention, Tmax under high fat meal administration conditions is less than 8 hours, alternatively less than 6 hours, alternatively approximately 5 hours, alternatively 4 hours or less. In other preferred embodiments of the present invention, Tmax under low fat meal administration conditions is less than 8 hours, alternatively less than 6 hours, alternatively approximately 5 hours, alternatively 4 hours or less. In yet other preferred embodiments of the present invention, Tmax under fasting administration conditions is less than 8 hours, alternatively less than 6 hours, alternatively approximately 5 hours, alternatively 4 hours or less. In further embodiments of the present invention, Tmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, or total omega-3 fatty acids are equal or less than Tmax for LOVAZA® for EPA, DPA, DHA, EPA+DPA, EPA+DHA, or total omega-3 fatty acids under high fat meal, low fat meal, and fasting administration conditions. In other embodiments of the present invention, Tmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, or total omega-3 fatty acids are less than

Tmax for LOVAZA® for EPA, DPA, DHA, EPA+DPA, EPA+DHA, or total omega-3 fatty acids under either low fat meal, fasting, or both administration conditions.

[0072] In yet other embodiments of the present invention, Tmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, or total omega-3 fatty acids are equal or less than than Tmax for AMR101 for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under high fat meal, low fat meal, and fasting administration conditions. Finally, in other embodiments of the present invention, Tmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, or total omega-3 fatty acids are less than Tmax for AMR101 for EPA+DHA and to EPA, DPA, DHA, EPA+DPA, EPA+DHA, or total omega-3 fatty acids under either low fat meal, fasting, or both administration conditions.

[0073] In other embodiments of the present invention, compositions of the present invention are better absorbed than LOVAZA® as measured by the maximum concentration (Cmax) in blood, serum or plasma of EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids.

[0074] In preferred embodiments of the present invention, Cmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under high fat meal administration conditions are preferably at least 1.1 x (110% of) Cmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for LOVAZA®, alternatively at least 1.2 x (120% of), alternatively at least 1.3 x (130% of), alternatively at least 1.4 x (140% of), alternatively at least 1.5 x (150% of) Cmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for LOVAZA®.

[0075] In other preferred embodiments of the present invention, Cmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under low fat meal administration conditions are preferably at least 1.5 x (150% of) Cmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for LOVAZA®, alternatively at least 2 x (200% of), alternatively at least 3 x (300% of), alternatively at least 4 x (400% of), alternatively at least 5 x (500% of), alternatively at least 6 x (600% of) Cmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for LOVAZA®.

[0076] In other preferred embodiments of the present invention, C_{max} for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under fasting administration conditions are preferably at least 1.5 x (150% of) C_{max} for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for LOVAZA®, alternatively at least 2 x (200% of), alternatively at least 3 x (300% of), alternatively at least 4 x (400% of), alternatively at least 5 x (500% of), alternatively at least 6 x (600% of), alternatively at least 7 x (700% of) C_{max} for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for LOVAZA®.

[0077] In other embodiments of the present invention, compositions of the present invention are better absorbed than LOVAZA® as measured by the area under the concentration curve over time (AUC) in blood, serum or plasma of EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids.

[0078] In preferred embodiments of the present invention, AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under high fat meal administration conditions are preferably at least 1.1 x (110% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for LOVAZA®, alternatively at least 1.2 x (120% of), alternatively at least 1.3 x (130% of), alternatively at least 1.4 x (140% of), alternatively at least 1.5 x (150% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for LOVAZA®.

[0079] In other preferred embodiments of the present invention, AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under low fat meal administration conditions are preferably at least 1.5 x (150% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for LOVAZA®, alternatively at least 2 x (200% of), alternatively at least 3 x (300% of), alternatively at least 4 x (400% of), alternatively at least 5 x (500% of), alternatively at least 6 x (600% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, or total omega-3-pentaenoic acids, or total omega-3 fatty acids for LOVAZA®.

[0080] In other preferred embodiments of the present invention, AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under fasting administration conditions are preferably at least 1.5 x

(150% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for LOVAZA®, alternatively at least 2 x (200% of), alternatively at least 3 x (300% of), alternatively at least 4 x (400% of), alternatively at least 5 x (500% of), alternatively at least 6 x (600% of), alternatively at least 7 x (700% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for LOVAZA®.

[0081] In other embodiments of the present invention, compositions of the present invention are better absorbed than AMR101 as measured by the maximum concentration (C_{max}) in blood, serum or plasma of EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids.

[0082] In preferred embodiments of the present invention, C_{max} for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under high fat meal administration conditions are preferably at least 1.1 x (110% of) C_{max} for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for AMR101, alternatively at least 1.2 x (120% of), alternatively at least 1.3 x (130% of), alternatively at least 1.4 x (140% of), alternatively at least 1.5 x (150% of) C_{max} for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for AMR101.

[0083] In other preferred embodiments of the present invention, C_{max} for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under low fat meal administration conditions are preferably at least 1.5 x (150% of) C_{max} for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for AMR101, alternatively at least 2 x (200% of), alternatively at least 3 x (300% of), alternatively at least 4 x (400% of), alternatively at least 5 x (500% of), alternatively at least 6 x (600% of) C_{max} for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for AMR101.

[0084] In other preferred embodiments of the present invention, C_{max} for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under fasting administration conditions are preferably at least 1.5 x (150% of) C_{max} for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for AMR101, alternatively at least 2 x (200% of), alternatively at least 3 x (300% of), alternatively at least 4 x (400% of),

alternatively at least 5 x (500% of), alternatively at least 6 x (600% of), alternatively at least 7 x (700% of) C_{max} for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for AMR101.

[0085] In other embodiments of the present invention, compositions of the present invention are better absorbed than AMR101 as measured by the area under the concentration curve over time (AUC) in blood, serum or plasma of EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids. In preferred embodiments of the present invention, AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under high fat meal administration conditions are preferably at least 1.1 x (110% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for AMR101, alternatively at least 1.2 x (120% of), alternatively at least 1.3 x (130% of), alternatively at least 1.4 x (140% of), alternatively at least 1.5 x (150% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for AMR101.

[0086] In other preferred embodiments of the present invention, AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under low fat meal administration conditions are preferably at least 1.5 x (150% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for AMR101, alternatively at least 2 x (200% of), alternatively at least 3 x (300% of), alternatively at least 4 x (400% of), alternatively at least 5 x (500% of), alternatively at least 6 x (600% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for AMR101.

[0087] In other preferred embodiments of the present invention, AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under fasting administration conditions are preferably at least 1.5 x (150% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for AMR101, alternatively at least 2 x (200% of), alternatively at least 3 x (300% of), alternatively at least 4 x (400% of), alternatively at least 5 x (500% of), alternatively at least 6 x (600% of), alternatively at least 7 x (700% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for AMR101.

[0088] In other embodiments of the present invention, compositions of the present invention are better absorbed than EPANOVA™ as measured by the maximum concentration (Cmax) in blood, serum or plasma of EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids.

[0089] In preferred embodiments of the present invention, Cmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under high fat meal administration conditions are preferably approximately 1.0 x (100% of) Cmax (or non-significant difference) for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for EPANOVA™, alternatively at least 1.05 x (105% of), alternatively at least 1.1 x (110% of), alternatively at least 1.2 x (120% of), alternatively at least 1.3 x (130% of) Cmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for EPANOVA™.

[0090] In other preferred embodiments of the present invention, Cmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under low fat meal administration conditions are preferably approximately 1.0 x (100% of) Cmax (or non-significant difference) for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for EPANOVA™, alternatively at least 1.05 x (105% of), alternatively at least 1.1 x (110% of), alternatively at least 1.2 x (120% of), alternatively at least 1.3 x (130% of) Cmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for EPANOVA™.

[0091] In other preferred embodiments of the present invention, Cmax for EPA+DHA, total omega-3-pentaenoic acids, and total omega-3 fatty acids under fasting administration conditions are preferably approximately 1.0 x (100% of) Cmax (or non-significant difference) for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for EPANOVA™, alternatively at least 1.05 x (105% of), alternatively at least 1.1 x (110% of), alternatively at least 1.2 x (120% of), alternatively at least 1.3 x (130% of) Cmax for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for EPANOVA™.

[0092] In other embodiments of the present invention, compositions of the present invention are better absorbed than EPANOVA™ as measured by the area under the

concentration curve over time (AUC) in blood, serum or plasma of EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids. In preferred embodiments of the present invention, AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under high fat meal administration conditions are preferably approximately 1.0 x (100% of) AUC (or non-significant difference) for EPA, DPA, DHA, EPA+DPA, EPA+DHA, or total omega-3-pentaenoic acids, or total omega-3 fatty acids for EPANOVA™, alternatively at least 1.05 x (105% of), alternatively at least 1.1 x (110% of), alternatively at least 1.2 x (120% of), alternatively at least 1.3 x (130% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for EPANOVA™.

[0093] In other preferred embodiments of the present invention, AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under low fat meal administration conditions are preferably approximately 1.0 x (100% of) AUC (or non-significant difference) for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for EPANOVA™, alternatively at least 1.05 x (105% of), alternatively at least 1.1 x (110% of), alternatively at least 1.2 x (120% of), alternatively at least 1.3 x (130% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for EPANOVA™.

[0094] In other preferred embodiments of the present invention, AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids under fasting administration conditions are preferably approximately 1.0 x (100% of) AUC (or non-significant difference) for EPA, DPA, DHA, EPA+DPA, EPA+DHA, or total omega-3-pentaenoic acids, or total omega-3 fatty acids for EPANOVA™, alternatively at least 1.05 x (105% of), alternatively at least 1.1 x (110% of), alternatively at least 1.2 x (120% of), alternatively at least 1.3 x (130% of) AUC for EPA, DPA, DHA, EPA+DPA, EPA+DHA, total omega-3-pentaenoic acids, or total omega-3 fatty acids for EPANOVA™.

[0095] In some embodiments, the improved bioavailability features described above are apparent upon single dose administration, while in other embodiments the improved bioavailability features described above are apparent after multiple dose administration of formulations according to the present invention as compared to referenced comparator products above or substantial equivalent forms thereof.

[0096] The compositions of the present invention may be used for the treatment of patients by administering an effective amount of such compositions to a subject in need thereof, such as a subject prone to or afflicted with a disease or condition or in need of treatment for a disease or condition. The present invention provides methods of treating, preventing, and reducing symptoms associated with a disease or condition comprising administration of a composition of the present invention. Exemplary diseases or conditions include, but are not limited to: hypertriglyceridemia (for example, by those skilled in the art typically established by assessing fasting triglyceride (TG) levels); hypertriglyceridemia with $TG \geq 500 \text{ mg/dL}$ (VHTG); hypertriglyceridemia with TG 200-499mg/dL; hypertriglyceridemia with TG 200-499mg/dL while on statin treatment (HTG); hypercholesterolemia; mixed dyslipidemia; coronary heart disease (CHD); vascular disease; atherosclerotic disease and related conditions; heart failure; cardiac arrhythmias; blood coagulatory conditions associated with cardiac arrhythmias; hypertension; coagulation related disorders, including post-surgical deep vein thrombosis or other high risk thrombosis conditions; nephropathy; kidney or urinary tract disease; retinopathy; cognitive, psychiatric, neurological and other CNS disorders, including but not limited to schizophrenia, depression, bipolar disorder and any form of dementia (including ischemic dementia and vascular dementia); autoimmune diseases; inflammatory diseases; asthma, COPD or other respiratory disease; dermatological disease; metabolic syndrome; diabetes or other forms of metabolic disease; liver diseases including fatty liver disease; diseases affecting the senses, including those affecting vision and hearing; diseases of the gastrointestinal tract; diseases of the male or female reproductive system or related secondary sexual organs; a cancer of any type, including lymphomas, myelomas and solid tumor cancers; any infections caused by a virus, bacterium, fungus, protozoa or other organism. The present invention also provides for the treatment and/or prevention of cardiac events and/or cardiovascular events and/or vascular events and/or symptoms. The present invention also provides for the reduction of number of such events, as well as a reduction or amelioration of symptoms associated with such events.

[0097] Cardiovascular and/or cardiac events may include, but are not limited to: myocardial infarction, ischemic cardiac attack, ischemic attack, acute angina, hospitalization due to acute angina, stroke, transient ischemic cerebral attack, cardiac revascularization, cardiac revascularization with stent placement, carotid

artery revascularization, carotid artery revascularization with stent placement, peripheral artery revascularization, peripheral artery revascularization with stent placement, plaque rupture, death due to cardiovascular event, and hospitalization due to cardiovascular event. Cardiovascular and/or cardiac events may also include other events deemed to fall in such category by those skilled in the art.

[0098] The present invention provides methods of treatment for hypertriglyceridemia (either $TG \geq 500\text{mg/dL}$, $TG \geq 200\text{mg/dL}$, $TG \geq 150\text{mg/dL}$, $TG 200-499\text{mg/dL}$, or $TG 150-199\text{mg/dL}$), mixed dyslipidemia, or any other diseases or medical conditions as specified above, by dosing to a subject in need thereof omega-3 docosapentaenoic acid (DPA-n3) or its glycerol or ethyl esters. Such method of treatment provides a dose of at least 60mg DPA-N3 per day, alternatively at least 80mg DPA-N3 per day, alternatively at least 90mg DPA-N3 per day, alternatively at least 120mg DPA-N3 per day, alternatively at least 150mg DPA-N3 per day, alternatively at least 160mg DPA-N3 per day, alternatively at least 180mg DPA-N3 per day, alternatively at least 200mg DPA-N3 per day, alternatively at least 250mg DPA-N3 per day, alternatively at least 300mg DPA-N3 per day, alternatively at least 350mg DPA-N3 per day, alternatively at least 400mg DPA-N3 per day, alternatively at least 500mg DPA-N3 per day, alternatively at least 600mg DPA-N3 per day, alternatively at least 800mg DPA-N3 or its glycerol or ethyl esters per day.

[0099] In some embodiments, the compositions of the present invention, which may comprise significant amounts of omega-3 docosapentaenoic acid (DPA-n3) or its glycerol or ethyl esters, may be used for the treatment of hypertriglyceridemia (either $TG \geq 500\text{mg/dL}$, $TG \geq 200\text{mg/dL}$, $TG \geq 150\text{mg/dL}$, $TG 200-499\text{mg/dL}$, or $TG 150-199\text{mg/dL}$), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such method of treatment provides to a subject in need thereof a dose of at least 20mg DPA-N3 per day, alternatively at least 25mg DPA-N3 per day, alternatively at least 30mg DPA-N3 per day, alternatively at least 40mg DPA-N3 per day, alternatively at least 50mg DPA-N3 per day, alternatively at least 60mg DPA-N3 per day, alternatively at least 70mg DPA-N3 per day, alternatively at least 80mg DPA-N3 per day, alternatively at least 90mg DPA-N3 per day, alternatively at least 100mg DPA-N3 per day, alternatively at least 120mg DPA-N3 per day, alternatively at least 150mg DPA-N3 per day, alternatively at least 160mg DPA-N3 per day, alternatively at least 180mg DPA-N3 per day, alternatively at least 200mg DPA-N3 per day, alternatively at least 250mg DPA-N3 per day, alternatively at least 300mg

DPA-N3 per day, alternatively at least 350mg DPA-N3 per day, alternatively at least 400mg DPA-N3 per day, alternatively at least 500mg DPA-N3 per day, alternatively at least 600mg DPA-N3 per day, alternatively alternatively at least 800mg DPA-N3 or its glycerol or ethyl esters per day.

[00100] In other embodiments, the compositions of the present invention, which may comprise significant amounts of omega-3 docosapentaenoic acid (DPA-n3) or its glycerol or ethyl esters and which comprise relatively small amounts of omega-3 docosahexaenoic acid (DHA-n3), may be used for the treatment of hypertnglycemia (either TG $\geq$ 500mg/dL, TG $\geq$ 200mg/dL, TG $\geq$ 150mg/dL, TG200-499mg/dL, or TG150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such method of treatment provides to a subject in need thereof a dose of at least 20mg DPA-N3 per day, alternatively at least 25mg DPA-N3 per day, alternatively at least 30mg DPA-N3 per day, alternatively at least 40mg DPA-N3 per day, alternatively at least 50mg DPA-N3 per day, alternatively at least 60mg DPA-N3 per day, alternatively at least 80mg DPA-N3 per day, alternatively at least 90mg DPA-N3 per day, alternatively at least 120mg DPA-N3 per day, alternatively at least 150mg DPA-N3 per day, alternatively at least 160mg DPA-N3 per day, alternatively at least 180mg DPA-N3 per day, alternatively at least 200mg DPA-N3 per day, alternatively at least 250mg DPA-N3 per day, alternatively at least 300mg DPA-N3 per day, alternatively at least 350mg DPA-N3 per day, alternatively at least 400mg DPA-N3 per day, alternatively at least 500mg DPA-N3 per day, alternatively at least 600mg DPA-N3 per day, alternatively alternatively at least 800mg DPA-N3 or its glycerol or ethyl esters per day, while providing less than 1500mg of DHA, alternatively less than 1200mg of DHA, alternatively less than 1000mg of DHA, alternatively less than 800mg of DHA, alternatively less than 700mg of DHA, alternatively less than 600mg of DHA, alternatively less than 500mg of DHA, alternatively less than 400mg of DHA, alternatively less than 350mg of DHA, alternatively less than 300mg of DHA, alternatively less than 250mg of DHA, alternatively less than 200mg of DHA alternatively less than 150mg of DHA, alternatively less than 120mg of DHA, alternatively less than 100 mg of DHA, alternatively less than 80 mg of DHA, alternatively less than 60 mg of DHA, alternatively less than 40 mg of DHA, alternatively less than 30 mg of DHA, alternatively less than 25 mg of DHA, alternatively less than 20 mg of DHA or its glycerol or ethyl esters per day.

[0100] In further embodiments, the compositions of the present invention, which may comprise significant amounts of omega-3 docosapentaenoic acid (DPA-n3) or its glycerol or ethyl esters and which comprise relatively small amounts of omega-3 docosahexaenoic acid (DHA-n3), may be used for the treatment of hypertriglyceridemia (either $TG \geq 500\text{mg/dL}$, $TG \geq 200\text{mg/dL}$, $TG \geq 150\text{mg/dL}$, $TG 200-499\text{mg/dL}$, or $TG 150-199\text{mg/dL}$), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such method of treatment provides to a subject in need thereof a dose of at least 30mg DPA-N3 per day, alternatively at least 40mg DPA-N3 per day, alternatively at least 50mg DPA-N3 per day, alternatively at least 60mg DPA-N3 per day, alternatively at least 80mg DPA-N3 per day, alternatively at least 90mg DPA-N3 per day, alternatively at least 120mg DPA-N3 per day, alternatively at least 150mg DPA-N3 per day, alternatively at least 160mg DPA-N3 per day, alternatively at least 180mg DPA-N3 per day, alternatively at least 200mg DPA-N3 per day, alternatively at least 250mg DPA-N3 per day, alternatively at least 300mg DPA-N3 per day, alternatively at least 350mg DPA-N3 per day, alternatively at least 400mg DPA-N3 per day, alternatively at least 500mg DPA-N3 per day, alternatively at least 600mg DPA-N3 per day, alternatively at least 800mg DPA-N3 per day, alternatively at least 1000mg DPA-N3 per day, alternatively at least 1200mg DPA-N3 per day, alternatively at least 1500mg DPA-N3 or its glycerol or ethyl esters per day, while providing a relatively small amount of DHA-N3 such that the DHA:DPA dose ratio is no more than 15:1 of DHA:DPA, alternatively no more than 12:1 of DHA:DPA, alternatively no more than 10:1 of DHA:DPA, alternatively no more than 8:1 of DHA:DPA, alternatively no more than 5:1 of DHA:DPA, alternatively no more than 3:1 of DHA:DPA, alternatively no more than 2:1 of DHA:DPA, alternatively no more than 1:1 of DHA:DPA, alternatively no more than 1:2 of DHA:DPA, alternatively no more than 1:3 of DHA:DPA, alternatively no more than 1:5 of DHA:DPA, alternatively no more than 1:8 of DHA:DPA, alternatively no more than 1:10 of DHA:DPA, alternatively no more than 1:15 of DHA:DPA, alternatively a relative daily dose of no more than 1:20 of DHA:DPA.

[0101] The present invention provides methods of reducing triglycerides and treating hypertriglyceridemia (either $TG \geq 500\text{mg/dL}$, $TG \geq 200\text{mg/dL}$, $TG \geq 150\text{mg/dL}$, $TG 200-499\text{mg/dL}$, or $TG 150-199\text{mg/dL}$), mixed dyslipidemia, or any other diseases or medical conditions as specified above, by dosing to a subject in need thereof omega-3 docosapentaenoic acid (HPA-n3) or its glycerol or ethyl esters. Such

method of treatment provides a dose of at least 10 mg HPA-N3 per day, alternatively at least 15 mg HPA-N3 per day, alternatively at least 20 mg HPA-N3 per day, alternatively at least 25 mg HPA-N3 per day, alternatively at least 30mg HPA-N3 per day, alternatively at least 40mg HPA-N3 per day, alternatively at least 50mg HPA-N3 per day, alternatively at least 60mg HPA-N3 per day, alternatively at least 70mg HPA-N3 per day, alternatively at least 80mg HPA-N3 per day, alternatively at least 90mg HPA-N3 per day, alternatively at least 100mg HPA-N3 per day, alternatively at least 120mg HPA-N3 per day, alternatively at least 150mg HPA-N3 per day, alternatively at least 160mg HPA-N3 per day, alternatively at least 180mg HPA-N3 per day, alternatively at least 200mg HPA-N3 per day, alternatively at least 250mg HPA-N3 per day, alternatively at least 300mg HPA-N3 per day, alternatively at least 350mg HPA-N3 per day, alternatively at least 400mg HPA-N3 per day, alternatively at least 500mg HPA-N3 per day, alternatively at least 600mg HPA-N3 per day, alternatively at least 800mg HPA-N3 or its glycerol or ethyl esters per day.

[0102] In some embodiments, the compositions of the present invention, which may comprise significant amounts of omega-3 heneicosapentaenoic acid (HPA-n3) or its glycerol or ethyl esters, may be used for the treatment of hypertriglyceridemia (either TG $\geq$ 500mg/dL, TG $\geq$ 200mg/dL, TG $\geq$ 150mg/dL, TG200-499mg/dL, or TG150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such method of treatment provides to a subject in need thereof a dose of at least 10 mg HPA-N3 per day, alternatively at least 15 mg HPA-N3 per day, alternatively at least 20 mg HPA-N3 per day, alternatively at least 25 mg HPA-N3 per day, alternatively at least 30 mg HPA-N3 per day, alternatively at least 40mg HPA-N3 per day, alternatively at least 50mg HPA-N3 per day, alternatively at least 60mg HPA-N3 per day, alternatively at least 70mg HPA-N3 per day, alternatively at least 80mg HPA-N3 per day, alternatively at least 90mg HPA-N3 per day, alternatively at least 100mg HPA-N3 per day, alternatively at least 120mg HPA-N3 per day, alternatively at least 150mg HPA-N3 per day, alternatively at least 160mg HPA-N3 per day, alternatively at least 180mg HPA-N3 per day, alternatively at least 200mg HPA-N3 per day, alternatively at least 250mg HPA-N3 per day, alternatively at least 300mg HPA-N3 per day, alternatively at least 350mg HPA-N3 per day, alternatively at least 400mg HPA-N3 per day, alternatively at least 500mg HPA-N3 per day, alternatively at least 600mg HPA-N3 per day, alternatively alternatively at least 800mg HPA-N3 or its glycerol or ethyl esters per day.

[0103] In other embodiments, the compositions of the present invention, which may comprise significant amounts of omega-3 docosapentaenoic acid (HPA-n3) or its glycerol or ethyl esters and which comprise relatively small amounts of omega-3 docosahexaenoic acid (DHA-n3), may be used for the treatment of hypertriglyceridemia (either $TG \geq 500\text{mg/dL}$, $TG \geq 200\text{mg/dL}$, $TG \geq 150\text{mg/dL}$, $TG 200-499\text{mg/dL}$, or $TG 150-199\text{mg/dL}$), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such method of treatment provides to a subject in need thereof a dose of at least 20mg HPA-N3 per day, alternatively at least 25mg HPA-N3 per day, alternatively at least 30mg HPA-N3 per day, alternatively at least 40mg HPA-N3 per day, alternatively at least 50mg HPA-N3 per day, alternatively at least 60mg HPA-N3 per day, alternatively at least 70mg HPA-N3 per day, alternatively at least 80mg HPA-N3 per day, alternatively at least 90mg HPA-N3 per day, alternatively at least 100mg HPA-N3 per day, alternatively at least 120mg HPA-N3 per day, alternatively at least 150mg HPA-N3 per day, alternatively at least 160mg HPA-N3 per day, alternatively at least 180mg HPA-N3 per day, alternatively at least 200mg HPA-N3 per day, alternatively at least 250mg HPA-N3 per day, alternatively at least 300mg HPA-N3 per day, alternatively at least 350mg HPA-N3 per day, alternatively at least 400mg HPA-N3 per day, alternatively at least 500mg HPA-N3 per day, alternatively at least 600mg HPA-N3 per day, alternatively alternatively at least 800mg HPA-N3 or its glycerol or ethyl esters per day, while providing less than 1500mg of DHA, alternatively less than 1200mg of DHA, alternatively less than 1000mg of DHA, alternatively less than 800mg of DHA, alternatively less than 700mg of DHA, alternatively less than 600mg of DHA, alternatively less than 500mg of DHA, alternatively less than 400mg of DHA, alternatively less than 350mg of DHA, alternatively less than 300mg of DHA, alternatively less than 250mg of DHA, alternatively less than 200mg of DHA or its glycerol or ethyl esters per day, alternatively less than 150mg of DHA, alternatively less than 120mg of DHA, alternatively less than 100mg of DHA, alternatively less than 80mg of DHA, alternatively less than 60mg of DHA, alternatively less than 40mg of DHA, alternatively less than 30mg of DHA, alternatively less than 25mg of DHA, alternatively less than 20mg of DHA or its glycerol or ethyl esters per day.

[0104] In further embodiments, the compositions of the present invention, which may comprise significant amounts of omega-3 heneicosapentaenoic acid (HPA-n3) or its glycerol or ethyl esters and which comprise relatively small amounts of omega-

3 docosahexaenoic acid (DHA-n3), may be used for the treatment of hypertriglyceridemia (either TG $\geq$ 500mg/dL, TG $\geq$ 200mg/dL, TG $\geq$ 150mg/dL, TG200-499mg/dL, or TG150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such method of treatment provides to a subject in need thereof a dose of at least 10mg HPA-N3 per day, alternatively at least 15mg HPA-N3 per day, alternatively at least 20mg HPA-N3 per day, alternatively at least 25mg HPA-N3 per day, alternatively at least 30mg HPA-N3 per day, alternatively at least 40mg HPA-N3 per day, alternatively at least 50mg HPA-N3 per day, alternatively at least 60mg HPA-N3 per day, alternatively at least 70mg HPA-N3 per day, alternatively at least 80mg HPA-N3 per day, alternatively at least 90mg HPA-N3 per day, alternatively at least 100mg HPA-N3 per day, alternatively at least 120mg HPA-N3 per day, alternatively at least 150mg HPA-N3 per day, alternatively at least 160mg HPA-N3 per day, alternatively at least 180mg HPA-N3 per day, alternatively at least 200mg HPA-N3 per day, alternatively at least 250mg HPA-N3 per day, alternatively at least 300mg HPA-N3 per day, alternatively at least 350mg HPA-N3 per day, alternatively at least 400mg HPA-N3 per day, alternatively at least 500mg HPA-N3 per day, alternatively at least 600mg HPA-N3 per day, alternatively at least 800mg HPA-N3 per day, alternatively at least 1000mg HPA-N3 per day, alternatively at least 1200mg HPA-N3 per day, alternatively at least 1500mg HPA-N3 or its glycerol or ethyl esters per day, while providing a relatively small amount of DHA-N3 such that the DHA:HPA dose ratio is no more than 15:1 of DHA:HPA, alternatively no more than 12:1 of DHA:HPA, alternatively no more than 10:1 of DHA:HPA, alternatively no more than 8:1 of DHA:HPA, alternatively no more than 5:1 of DHA:HPA, alternatively no more than 3:1 of DHA:HPA, alternatively no more than 2:1 of DHA:HPA, alternatively no more than 1:1 of DHA:HPA, alternatively no more than 1:2 of DHA:HPA, alternatively no more than 1:3 of DHA:HPA, alternatively no more than 1:5 of DHA:HPA, alternatively no more than 1:8 of DHA:HPA, alternatively no more than 1:10 of DHA:HPA, alternatively no more than 1:15 of DHA:HPA, alternatively a relative daily dose of no more than 1:20 of DHA:HPA.

[0105] The present invention, incorporates methods of treatment for hypertriglyceridemia (either TG $\geq$ 500mg/dL, TG $\geq$ 200mg/dL, TG $\geq$ 150mg/dL, TG200-499mg/dL, or TG150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions as specified above, by dosing to a subject in need thereof omega-3-pentaenoic acids or their glycerol or ethyl esters. These methods provide

to a subject in need thereof significant amounts of omega-3-pentaenoic acids or their glycerol or ethyl esters and while providing only small amounts of omega-3 docosahexaenoic acid (DHA-n3). Such methods of treatment provides to a subject in need thereof a dose of at least 100mg omega-3-pentaenoic acids per day, alternatively at least 200mg omega-3-pentaenoic acids per day, alternatively at least 300mg omega-3-pentaenoic acids per day, alternatively at least 500mg omega-3-pentaenoic acids per day, alternatively at least 700mg omega-3-pentaenoic acids per day, alternatively at least 900mg omega-3-pentaenoic acids per day, alternatively at least 1000mg omega-3-pentaenoic acids per day, alternatively at least 1500mg omega-3-pentaenoic acids per day, alternatively at least 1900mg omega-3-pentaenoic acids per day, alternatively at least 2000mg omega-3-pentaenoic acids per day, alternatively at least 2500mg omega-3-pentaenoic acids per day, alternatively at least 2900mg omega-3-pentaenoic acids per day, alternatively at least 3000mg omega-3-pentaenoic acids per day, alternatively at least 3500mg omega-3-pentaenoic acids per day, alternatively at least 3900mg omega-3-pentaenoic acids per day, alternatively at least 4000mg omega-3-pentaenoic acids per day, alternatively at least 4100mg omega-3-pentaenoic acids per day, alternatively at least 4500mg omega-3-pentaenoic acids per day, alternatively at least 5000mg omega-3-pentaenoic acids per day, alternatively at least 5500mg omega-3-pentaenoic acids per day, alternatively at least 6100mg omega-3-pentaenoic acids per day or their glycerol or ethyl esters, while providing less than 1900mg of DHA, alternatively less than 1500mg of DHA, alternatively less than 1200mg of DHA, alternatively less than 1000mg of DHA, alternatively less than 800mg of DHA, alternatively less than 700mg of DHA, alternatively less than 600mg of DHA, alternatively less than 500mg of DHA, alternatively less than 400mg of DHA, alternatively less than 350mg of DHA, alternatively less than 300mg of DHA, alternatively less than 250mg of DHA, alternatively less than 200mg of DHA, alternatively less than 150mg of DHA, alternatively less than 120mg of DHA, alternatively less than 100mg of DHA, alternatively less than 80mg of DHA, alternatively less than 60mg of DHA, alternatively less than 40mg of DHA, alternatively less than 30mg of DHA, alternatively less than 25mg of DHA, alternatively less than 20mg of DHA or its glycerol or ethyl esters per day.

[0106] In other embodiments, the compositions of the present invention, which comprise significant amounts of omega-3-pentaenoic acids or their glycerol or ethyl

esters and which comprise relatively small amounts of omega-3 docosahexaenoic acid (DHA-n3), may be used for the treatment of hypertriglyceridemia (either $TG \geq 500\text{mg/dL}$, $TG \geq 200\text{mg/dL}$, $TG \geq 150\text{mg/dL}$, $TG 200-499\text{mg/dL}$, or $TG 150-199\text{mg/dL}$), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such method of treatment provides to a subject in need thereof a dose of at least 100mg omega-3-pentaenoic acids per day, alternatively at least 200mg omega-3-pentaenoic acids per day, alternatively at least 300mg omega-3-pentaenoic acids per day, alternatively at least 500mg omega-3-pentaenoic acids per day, alternatively at least 700mg omega-3-pentaenoic acids per day, alternatively at least 900mg omega-3-pentaenoic acids per day, alternatively at least 1000mg omega-3-pentaenoic acids per day, alternatively at least 1500mg omega-3-pentaenoic acids per day, alternatively at least 1900mg omega-3-pentaenoic acids per day, alternatively at least 2000mg omega-3-pentaenoic acids per day, alternatively at least 2500mg omega-3-pentaenoic acids per day, alternatively at least 2900mg omega-3-pentaenoic acids per day, alternatively at least 3000mg omega-3-pentaenoic acids per day, alternatively at least 3500mg omega-3-pentaenoic acids per day, alternatively at least 3900mg omega-3-pentaenoic acids per day, alternatively at least 4000mg omega-3-pentaenoic acids per day or their glycerol or ethyl esters, while providing a relatively small amount of DHA-N3 such that the DHA:DPA ratio is no more than 15:1 of DHA:DPA, alternatively no more than 12:1 of DHA:DPA, alternatively no more than 10:1 of DHA:DPA, alternatively no more than 8:1 of DHA:DPA, alternatively no more than 5:1 of DHA:DPA, alternatively no more than 3:1 of DHA:DPA, alternatively no more than 2:1 of DHA:DPA, alternatively no more than 1:1 of DHA:DPA, alternatively no more than 1:2 of DHA:DPA, alternatively no more than 1:3 of DHA:DPA, alternatively no more than 1:5 of DHA:DPA, alternatively no more than 1:8 of DHA:DPA, alternatively no more than 1:10 of DHA:DPA, alternatively no more than 1:15 of DHA:DPA, alternatively a relative amount of no more than 1:20 of DHA:DPA.

[0107] In other embodiments, the compositions of the present invention, which may comprise significant amounts of omega-3-pentaenoic acids or their glycerol or ethyl esters and which comprise relatively small amounts of omega-3 docosahexaenoic acid (DHA-n3), may be used for the reduction of triglycerides and treatment of hypertriglyceridemia (either $TG \geq 500\text{mg/dL}$, $TG \geq 200\text{mg/dL}$, $TG \geq 150\text{mg/dL}$, $TG 200-499\text{mg/dL}$, or $TG 150-199\text{mg/dL}$), mixed dyslipidemia, or any other diseases or

medical conditions specified above. Such method of treatment provides to a subject in need thereof a dose of at least 100mg omega-3-pentaenoic acids per day, alternatively at least 200mg omega-3-pentaenoic acids per day, alternatively at least 300mg omega-3-pentaenoic acids per day, alternatively at least 500mg omega-3-pentaenoic acids per day, alternatively at least 700mg omega-3-pentaenoic acids per day, alternatively at least 900mg omega-3-pentaenoic acids per day, alternatively at least 1000mg omega-3-pentaenoic acids per day, alternatively at least 1500mg omega-3-pentaenoic acids per day, alternatively at least 1900mg omega-3-pentaenoic acids per day, alternatively at least 2000mg omega-3-pentaenoic acids per day, alternatively at least 2500mg omega-3-pentaenoic acids per day, alternatively at least 2900mg omega-3-pentaenoic acids per day, alternatively at least 3000mg omega-3-pentaenoic acids per day, alternatively at least 3500mg omega-3-pentaenoic acids per day, alternatively at least 3900mg omega-3-pentaenoic acids per day, alternatively at least 4000mg omega-3-pentaenoic acids per day or their glycerol or ethyl esters, while providing a relatively small amount of DHA-N3 such that the DHA:HPA ratio is no more than 15:1 of DHA:HPA, alternatively no more than 12:1 of DHA:HPA, alternatively no more than 10:1 of DHA:HPA, alternatively no more than 8:1 of DHA:HPA, alternatively no more than 5:1 of DHA:HPA, alternatively no more than 3:1 of DHA:HPA, alternatively no more than 2:1 of DHA:HPA, alternatively no more than 1:1 of DHA:HPA, alternatively no more than 1:2 of DHA:HPA, alternatively no more than 1:3 of DHA:HPA, alternatively no more than 1:5 of DHA:HPA, alternatively no more than 1:8 of DHA:HPA, alternatively no more than 1:10 of DHA:HPA, alternatively no more than 1:15 of DHA:HPA, alternatively a relative amount of no more than 1:20 of DHA:HPA.

[0108] In further embodiments, the compositions of the present invention, which may comprise significant amounts of omega-3-pentaenoic acids or their glycerol or ethyl esters and which comprise relatively small amounts of omega-3 docosahexaenoic acid (DHA-n3), may be used for the reduction of triglycerides and treatment of hypertriglyceridemia (either $TG \geq 500\text{mg/dL}$, $TG \geq 200\text{mg/dL}$, $TG \geq 150\text{mg/dL}$, $TG 200-499\text{mg/dL}$, or $TG 150-199\text{mg/dL}$), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such method of treatment provides to a subject in need thereof a dose of at least 100mg omega-3-pentaenoic acids per day, alternatively at least 200mg omega-3-pentaenoic acids per day, alternatively at least 300mg omega-3-pentaenoic acids per day, alternatively at least

500mg omega-3-pentaenoic acids per day, alternatively at least 700mg omega-3-pentaenoic acids per day, alternatively at least 900mg omega-3-pentaenoic acids per day, alternatively at least 1000mg omega-3-pentaenoic acids per day, alternatively at least 1500mg omega-3-pentaenoic acids per day, alternatively at least 1900mg omega-3-pentaenoic acids per day, alternatively at least 2000mg omega-3-pentaenoic acids per day, alternatively at least 2500mg omega-3-pentaenoic acids per day, alternatively at least 2900mg omega-3-pentaenoic acids per day, alternatively at least 3000mg omega-3-pentaenoic acids per day, alternatively at least 3500mg omega-3-pentaenoic acids per day, alternatively at least 3900mg omega-3-pentaenoic acids per day, alternatively at least 4000mg omega-3-pentaenoic acids per day, alternatively at least 4100mg omega-3-pentaenoic acids per day, alternatively at least 4500mg omega-3-pentaenoic acids per day, alternatively at least 5000mg omega-3-pentaenoic acids per day, alternatively at least 5500mg omega-3-pentaenoic acids per day, alternatively at least 6000mg omega-3-pentaenoic acids per day or their glycerol or ethyl esters, while providing a relatively small amount of DHA:N3 as compared to the amount of omega-3-pentaenoic acids (N3-5enoicFA) such that the DHA:N3-5enoicFA ratio is no more than 15:1 of DHA:N3-5enoicFA, alternatively no more than 12:1 of DHA:N3-5enoicFA, alternatively no more than 10:1 of DHA:N3-5enoicFA, alternatively no more than 8:1 of DHA:N3-5enoicFA, alternatively no more than 5:1 of DHA:N3-5enoicFA, alternatively no more than 3:1 of DHA:N3-5enoicFA, alternatively no more than 2:1 of DHA:N3-5enoicFA, alternatively no more than 1:1 of DHA:N3-5enoicFA, alternatively no more than 1:2 of DHA:N3-5enoicFA, alternatively no more than 1:3 of DHA:N3-5enoicFA, alternatively no more than 1:5 of DHA:N3-5enoicFA, alternatively no more than 1:8 of DHA:N3-5enoicFA, alternatively no more than 1:10 of DHA:N3-5enoicFA, alternatively no more than 1:15 of DHA:N3-5enoicFA, alternatively a relative amount of no more than 1:20 of DHA:N3-5enoicFA.

[0109] In some embodiments, the improved profile of the compositions of the present invention may be demonstrated upon treatment of a subject by differentially altering the ration between blood platelets and fragments thereof (also known as platelet microparticles). Such fragments may be evaluated as a whole or examined and described as fragment sub-categories.

[0110] In other embodiments, the improved profile of the compositions of the present invention may be demonstrated upon treatment of a subject by differentially

altering the surface charge of blood platelets and fragments thereof, either in resting state (non-activated platelets) or activated stage.

[0111] In yet other embodiments, treatment of a subject or patient with compositions of the present invention affect the coagulatory cascade and differentially alter coagulation or bleeding times or platelet aggregation times and density.

[0112] In further embodiments, treatment with compositions of the present invention improves the vascular healing process in response to atherogenic disease. Such healing may be demonstrated by reduced stenosis and/or restenosis over time, reduced or lesser increase in intima-media thickness (IMT) of the arterial wall, larger lumen size and/or larger vascular diameter at vascular sites with stenosis or clot built-up, as determined by either by intravascular ultrasound (IVUS), radiographic, radiologic, non-invasive ultrasound, tomography, magnetic resonance interference (MRI), or other acceptable methods. In other embodiments, such improved healing may be demonstrated by the vascular wall composition, such as a reduced foam cell presence or fibrillated tissue in the vessel wall. In yet other embodiments, such improved vascular healing is demonstrated by improved inflammatory markers in the vascular wall.

[0113] The improved profile resulting from treatment with the compositions of the present invention may also be demonstrated by a differentiated impact on blood/serum/plasma lipid and lipoprotein levels in a mammal; these include, but are not limited to: Triglycerides (TG), total-cholesterol, non-HDL-cholesterol, LDL-cholesterol, VLDL-cholesterol, apolipoprotein B, apolipoprotein A, apolipoprotein C-III, HDL-cholesterol, and Lp-PLA2. The compositions of the present invention may also be used to provide a beneficial impact on the one or more of the following: apolipoprotein A-I (apo A-I), apolipoprotein B (apo B), apo A-I/apo B ratio, lipoprotein(a) (Lp[a]), lipoprotein-associated phospholipase A2 (Lp-PLA2), low density lipoprotein (LDL) particle number and size, oxidized LDL, C-reactive protein (CRP), high sensitivity C-reactive protein (HSCRP), intracellular adhesion molecule-1 (ICAM-1), E-selectin, P-selectin, vascular cell adhesion molecule 1 (VCAM-1) or cluster of differentiation 106 (CD106), interleukin -13 (IL-13), interleukin-2 (IL-2), interleukin-6 (IL-6), interleukin-8 (IL-8), interleukin-10 (IL-10), interleukin-12 (IL-12), interleukin-15 (IL-15), interleukin-18 (IL-18), tumor necrosis factor-alpha (TNF- α), tumor necrosis factor-beta (TNF- β), plasminogen activator inhibitor-1 (PAI-1).

homocysteine, thromboxane B2 (TXB2), thromboxane A2 (TXA2), 2,3-dinor thromboxane B2, free fatty acids (FFA), serum amyloid A1, serum amyloid A2, serum amyloid A3, serum amyloid A4, thiobarbituric acid (TBA) reacting material, adiponectin (GBP-28), hemoglobin A1c (HbA1c), macrophage colony stimulating factor (M-CSF), granulocyte macrophage colony stimulating factor (GM-CSF), fibrinogen, fibrin D-dimer, platelet derived-microparticles, mean platelet volume (MPV), platelet subpopulations, heart rate, systolic and diastolic blood pressure, nuclear factor kappa-light-chain enhancer of activated B cells (NF- κ B), adenosine diphosphate induced platelet aggregation, platelet endothelial cell adhesion molecule (PECAM-1), vitronectin receptor (α , β), and glycoprotein IIb/IIIa (gplIIB/IIIa). The compositions of the present invention may also be used in methods of treating, preventing, and reducing symptoms associated with conditions associated with the above.

[0114] Methods to determine comparative blood/serum/plasma lipid and lipoprotein levels and therapeutic effects on these levels in mammals are generally known to those skilled in the art and are typically based on fasting lipid and lipoprotein levels. Differences of active treatment versus placebo are generally assessed on a group of subjects versus another group of subjects basis, with significant changes noted if the p-value for the appropriate statistical comparison is equal to or less than 0.05. P-values larger than 0.05 but equal to or less than 0.10 may be considered borderline significant (BS). P-values larger than 0.10 are generally considered not significant (NS). In one embodiment, treatment with the compositions of the present invention is more potent than other omega-3 compositions known in the prior art (such as LOVAZA®, EPANOVA™ or AMR101) in reducing as compared to placebo or baseline: TG levels, Total-cholesterol levels, non-HDL-cholesterol levels, VLDL-cholesterol levels, LDL-cholesterol levels, apolipoprotein B levels, apolipoprotein C-III levels, Lp-PLA2 levels, or any combinations thereof. In other embodiments, such more potent effects in reducing these parameters are achieved in patients with baseline TG over 500 mg/dL, in patients on statin treatment with baseline TG in the 200-499 mg/dL range, in patients not on statin treatment with baseline LDL-cholesterol of 190 mg/dL or higher and with TG in the 300-700 mg/dL range, in patients not on statin treatment with baseline LDL-cholesterol of 190 mg/dL or higher and with TG in the 350-700 mg/dL range, in patients not on statin treatment with baseline LDL-cholesterol of 190 mg/dL or higher and with TG in the 300-750 mg/dL

range, in patients not on statin treatment with baseline LDL-cholesterol of 190 mg/dL or higher and with TG in the 350-750 mg/dL range, or in patients not on statin treatment with baseline non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 300-700 mg/dL range, or in patients not on statin treatment with baseline non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 350-700 mg/dL range, or in patients not on statin treatment with baseline non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 300-750 mg/dL range, or in patients not on statin treatment with baseline non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 350-750 mg/dL range.

[0115] In a further embodiment, treatment with the compositions of the present invention together with statin therapy is more potent than other omega-3 compositions known in the prior art (such as LOVAZA®, EPANOVA™ or AMR101) in reducing as compared to placebo or baseline: TG levels, Total-cholesterol levels, non-HDL-cholesterol levels, VLDL-cholesterol levels, LDL-cholesterol levels, apolipoprotein B levels, apolipoprotein C-III levels, Lp-PLA2 levels, or any combinations thereof. In other embodiments, such more potent effects in reducing these parameters are achieved in patients with baseline TG over 500 mg/dL, in patients on statin treatment with baseline TG in the 200-499 mg/dL range, in patients not on baseline statin treatment with baseline LDL-cholesterol of 190 mg/dL or higher and with TG in the 300-700 mg/dL range, in patients not on baseline statin treatment with baseline LDL-cholesterol of 190 mg/dL or higher and with TG in the 350-700 mg/dL range, in patients not on baseline statin treatment with baseline LDL-cholesterol of 190 mg/dL or higher and with TG in the 300-750 mg/dL range, in patients not on baseline statin treatment with baseline LDL-cholesterol of 190 mg/dL or higher and with TG in the 350-750 mg/dL range, or in patients not on baseline statin treatment with baseline non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 300-700 mg/dL range, or in patients not on baseline statin treatment with baseline non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 350-700 mg/dL range, or in patients not on baseline statin treatment with baseline non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 300-750 mg/dL range, or in patients not on baseline statin treatment with baseline Non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 350-750 mg/dL range.

[0116] The present invention also provides methods of reducing triglycerides levels in a subject, wherein the non-HDL cholesterol levels, such as LDL-cholesterol levels,

of the subject are reduced or not significantly increased from, for example, baseline levels before treatment. In some embodiments, treatment with the compositions of the present invention results in a minor (less than 10% change from baseline, alternatively less than 5%) and/or non-significant change in non-HDL cholesterol levels (such as LDL-cholesterol levels) as compared to placebo in patients with baseline TG levels above 500 mg/dL. In a further embodiment, treatment with the compositions of the present invention results in reductions of LDL-cholesterol levels as compared to placebo in patients with baseline TG levels above 500 mg/dL. In some embodiments, the methods involve coadministration of a statin.

In another embodiment, treatment with the compositions of the present invention as compared to placebo does not increase LDL-cholesterol levels in patients with baseline TG levels of 200-499 mg/dL while on statin therapy.

[0117] In yet another embodiment, treatment with the compositions of the present invention as compared to placebo results in significant reductions in LDL-cholesterol levels in patients with baseline TG levels of 200-499 mg/dL while on statin therapy.

[0118] In a further embodiment, the compositions of the present invention as compared to placebo result in significant reductions in LDL-cholesterol levels in patients not on statin treatment with LDL-cholesterol of 190 mg/dL or higher and with TG in the 300-700 mg/dL range, in patients not on statin treatment with LDL-cholesterol of 190 mg/dL or higher and with TG in the 350-700 mg/dL range, in patients not on statin treatment with LDL-cholesterol of 190 mg/dL or higher and with TG in the 300-750 mg/dL range, or in patients not on statin treatment with LDL-cholesterol of 190 mg/dL or higher and with TG in the 350-750 mg/dL range. Finally, another embodiment, the compositions of the present invention as compared to placebo result in significant reductions in LDL-cholesterol levels in patients not on statin treatment with non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 300-700 mg/dL range, in patients not on statin treatment with Non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 350-700 mg/dL range, in patients not on statin treatment with non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 300-750 mg/dL range, or in patients not on statin treatment with Non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 350-750 mg/dL range.

[0119] In another embodiment, treatment with the compositions of the present invention together with statin therapy results in significant reductions in LDL-

cholesterol levels as compared to placebo in patients not on statin treatment at baseline with baseline LDL-cholesterol of 190 mg/dL or higher and with TG in the 300-700 mg/dL range, in patients not on baseline statin treatment with baseline LDL-cholesterol of 190 mg/dL or higher and with TG in the 350-700 mg/dL range, in patients not on baseline statin treatment with baseline LDL-cholesterol of 190 mg/dL or higher and with TG in the 300-750 mg/dL range, or in patients not on baseline statin treatment with baseline LDL-cholesterol of 190 mg/dL or higher and with TG in the 350-750 mg/dL range.

[0120] Finally, another embodiment, treatment with the compositions of the present invention together with statin therapy results in significant reductions in LDL-cholesterol levels as compared to placebo in patients not on baseline statin treatment with baseline non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 300-700 mg/dL range, in patients not on baseline statin treatment with baseline non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 350-700 mg/dL range, in patients not on baseline statin treatment with baseline non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 300-750 mg/dL range, or in patients not on baseline statin treatment with baseline Non-HDL-cholesterol of 200 mg/dL or higher and with TG in the 350-750 mg/dL range.

[0121] In another embodiment, the compositions of the present invention are more potent than other omega-3 compositions known in the prior art (such as LOVAZA®, EPANOVA™ or AMR101) in increasing as compared to placebo or baseline HDL-cholesterol levels, apolipoprotein-A levels, or a combination thereof.

[0122] In yet another embodiment, the compositions of the present invention are more potent than other omega-3 compositions known in the prior art (such as LOVAZA®, EPANOVA™ or AMR101) in decreasing as compared to placebo or baseline Apolipoprotein-B (Apo-B) levels, Apolipoprotein-CIII levels, Lp-PLA2 levels or any combination thereof.

[0123] In further embodiments, the compositions of the present invention as compared to placebo or baseline are more potent than other omega-3 compositions known in the prior art (such as LOVAZA®, EPANOVA™ or AMR101) in reducing TG while causing a lesser increase in LDL-cholesterol, a lesser non-significant increase in LDL-cholesterol, no increase in LDL-cholesterol at all, or a more potent reduction in LDL-cholesterol at in patients with baseline TG levels above 500 mg/dL.

[0124] In some embodiments, the use of the compositions of the present invention may allow for a reduction in the dose of the statin required for a subject. For example, the coadministration of the composition of the present invention in a subject receiving statin therapy may allow for the reduction of the dose of the statin, compared to subject not being co-administered a composition of the present invention. In some embodiments, the dose of the statin may be reduced by at least 10%, alternatively at least 25%, alternatively at least 50%, or alternatively at least 75%.

[0125] In some embodiments, the use of the compositions of the present invention may reduce the time needed for a subject to reach the recommended blood levels. For example, the administration of compositions of the present invention may allow a subject to reach goal lipid levels, for example, those described in the NCEP ATP III Guidelines, or any levels recommended by a health care practitioner. In some embodiments, the reduction of time is greater than 5%, alternatively greater than 15%, alternatively greater than 25%, alternatively greater than 50%, and alternatively greater than 75%.

[0126] The compositions of the present invention are also useful to treat coronary heart disease (CHD), vascular disease, atherosclerotic disease or related conditions. The compositions of the present invention may also be use for the treatment and/or prevention and/or reduction of cardiac events and/or cardiovascular events and/or vascular events and/or symptoms. Determination of such cardiovascular diseases/conditions and prevention of events/symptoms in mammals and methods to determine treatment and preventative/therapeutic effects therefore are generally know to those skilled in the art.

[0127] The present invention also relates to treatment of such conditions in with concomitant treatments regimes or combination products with other active pharmaceutical ingredients. Such concomitant or fixed combination treatments may include a statin, an anticoagulant (such as aspirin or clopidogrel), an antihypertensive (such as a diuretic, beta-blocker, calcium channel blocker, ACE-inhibitor, angiotensin II receptor (ARB) antagonist), or other treatments for cardiovascular diseases.

The present invention also includes pharmaceutical compositions, for example, a unit dosage, comprising one or more HMG-CoA reductase inhibitors ("statins") and the omega-3 fatty acid composition of the present invention. The present invention

may incorporate now known or future known statins in an amount generally recognized as safe. There are currently seven statins that are widely available: atorvastatin, rosuvastatin, fluvastatin, lovastatin, pravastatin, pitavastatin, and simvastatin. An eighth statin, cerivastatin, has been removed from the U.S. market at the time of this writing. However, it is conceivable to one skilled in the art that cerivastatin may be used in conjunction with some embodiments of the present invention if cerivastatin is ultimately determined to be safe and effective in certain treatment regimens. Such statins are typically used at their common daily doses, which include, but are not limited to lovastatin 10mg, 20mg, 40mg; pravastatin 10mg, 20mg, 40mg, 80mg; simvastatin 5mg, 10mg, 20mg, 40mg, 80mg; fluvastatin 20mg, 40mg, 80mg; atorvastatin 10mg, 20mg, 40mg, 80mg; rosuvastatin 5mg, 10mg, 20mg, 40mg; and pitavastatin 1mg, 2mg, 4mg, 8mg.

[0128] Generally, the effect of statins is dose dependent, i.e., the higher the dose, the greater the therapeutic effect. However, the effect of each statin is different, and therefore the level of therapeutic effect of one statin cannot be necessarily be directly correlated to the level of therapeutic effects of other statins. For example, bioavailability varies widely among the statins. Specifically, it has been shown that simvastatin is less than 5% bioavailable, while fluvastatin is approximately 24% bioavailable. Statins are absorbed at rates ranging from about 30% with lovastatin to 98% with fluvastatin. First-pass metabolism occurs in all statins except pravastatin. Pravastatin is also the least protein-bound of the statins (about 50%), compared with the others, which are more than 90% protein-bound. Accordingly, the statins possess distinct properties from one another. The combination products of this invention involving each statin or a plurality of statins are also distinct.

[0129] The present invention also includes methods of treatment, comprising dosing of one or more statins and the omega-3 fatty acid composition of the present invention, either as concomitant therapy or in a fixed dose combination product comprising both a statin and the composition of the present invention. This method of treatment combines the administration of one or more statins at its common dose or an alternative dose with the composition of the present invention.

[0130] In some embodiments, the compositions of the present invention, which comprise significant amounts of omega-3 docosapentaenoic acid (DPA-n3) or its glycerol or ethyl esters, together with a common or alternative statin dose, may be used for the treatment of hypertriglyceridemia (either TG $\geq$ 500mg/dL, TG $\geq$ 200mg/dL,

TG $\geq$ 150mg/dL, TG 200-499mg/dL, or TG 150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such methods of treatment provide to a subject in need thereof a dose of at least 30 mg DPA-N3 per day, alternatively at least 40mg DPA-N3 per day, alternatively at least 50mg DPA-N3 per day, alternatively at least 60mg DPA-N3 per day, alternatively at least 70mg DPA-N3 per day, alternatively at least 80mg DPA-N3 per day, alternatively at least 90mg DPA-N3 per day, alternatively at least 100mg DPA-N3 per day, alternatively at least 120mg DPA-N3 per day, alternatively at least 150mg DPA-N3 per day, alternatively at least 160mg DPA-N3 per day, alternatively at least 180mg DPA-N3 per day, alternatively at least 200mg DPA-N3 per day, alternatively at least 250mg DPA-N3 per day, alternatively at least 300mg DPA-N3 per day, alternatively at least 350mg DPA-N3 per day, alternatively at least 400mg DPA-N3 per day, alternatively at least 500mg DPA-N3 per day, alternatively at least 600mg DPA-N3 per day, alternatively at least 800mg DPA-N3 or its glycerol or ethyl esters per day together with a common or alternative statin dose. In some embodiments, the compositions of the present invention, which comprise significant amounts of omega-3 docosapentaenoic acid (HPA-N3) or its glycerol or ethyl esters, together with a common or alternative statin dose, may be used for the treatment of hypertriglyceridemia (either TG $\geq$ 500mg/dL, TG $\geq$ 200mg/dL, TG $\geq$ 150mg/dL, TG200-499mg/dL, or TG150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such methods of treatment provide to a subject in need thereof a dose of at least 10mg HPA-N3 per day, alternatively at least 15mg HPA-N3 per day, alternatively at least 20mg HPA-N3 per day, alternatively at least 25mg HPA-N3 per day, alternatively at least 30mg HPA-N3 per day, alternatively at least 40mg HPA-N3 per day, alternatively at least 50mg HPA-N3 per day, alternatively at least 60mg HPA-N3 per day, alternatively at least 80mg HPA-N3 per day, alternatively at least 90mg HPA-N3 per day, alternatively at least 120mg HPA-N3 per day, alternatively at least 150mg HPA-N3 per day, alternatively at least 160mg HPA-N3 per day, alternatively at least 180mg HPA-N3 per day, alternatively at least 200mg HPA-N3 per day, alternatively at least 250mg HPA-N3 per day, alternatively at least 300mg HPA-N3 per day, alternatively at least 350mg HPA-N3 per day, alternatively at least 400mg HPA-N3 per day, alternatively at least 500mg HPA-N3 per day, alternatively at least 600mg HPA-N3 per day, alternatively at least

800mg HPA-N3 or its glycerol or ethyl esters per day together with a common or alternative statin dose.

[0131] In other embodiments, the compositions of the present invention, which comprise significant amounts of omega-3 docosapentaenoic acid (DPA-n3) or its glycerol or ethyl esters and which comprise relatively small amounts of omega-3 docosahexaenoic acid (DHA-n3), together with a common or alternative statin dose, may be used for the treatment of hypertriglyceridemia (either $TG \geq 500\text{mg/dL}$, $TG \geq 200\text{mg/dL}$, $TG \geq 150\text{mg/dL}$, TG 200-499mg/dL, or TG 150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such method of treatment provides to a subject in need thereof a common or alternative statin dose together with a dose of at least 30mg DPA-N3 per day, alternatively at least 40mg DPA-N3 per day, alternatively at least 50mg DPA-N3 per day, alternatively at least 60mg DPA-N3 per day, alternatively at least 80mg DPA-N3 per day, alternatively at least 90mg DPA-N3 per day, alternatively at least 120mg DPA-N3 per day, alternatively at least 150mg DPA-N3 per day, alternatively at least 160mg DPA-N3 per day, alternatively at least 180mg DPA-N3 per day, alternatively at least 200mg DPA-N3 per day, alternatively at least 250mg DPA-N3 per day, alternatively at least 300mg DPA-N3 per day, alternatively at least 350mg DPA-N3 per day, alternatively at least 400mg DPA-N3 per day, alternatively at least 500mg DPA-N3 per day, alternatively at least 600mg DPA-N3 per day, alternatively at least 800mg DPA-N3 or its glycerol or ethyl esters per day, while providing less than 2000mg of DHA, alternatively less than 1900mg of DHA, alternatively less than 1500mg of DHA, alternatively less than 1200mg of DHA, alternatively less than 1000mg of DHA, alternatively less than 800mg of DHA, alternatively less than 700mg of DHA, alternatively less than 600mg of DHA, alternatively less than 500mg of DHA, alternatively less than 400mg of DHA, alternatively less than 350mg of DHA, alternatively less than 300mg of DHA, alternatively less than 250mg of DHA, alternatively less than 200mg of DHA, alternatively less than 150mg of DHA, alternatively less than 120mg of DHA, alternatively less than 100mg of DHA, alternatively less than 80mg of DHA, alternatively less than 60mg of DHA, alternatively less than 50mg of DHA, alternatively less than 40mg of DHA, alternatively less than 30mg of DHA, alternatively less than 25mg of DHA, alternatively less than 20mg of DHA or its glycerol or ethyl esters per day.

[0132] In other embodiments, the compositions of the present invention, which comprise significant amounts of omega-3 docosapentaenoic acid (HPA-n3) or its glycerol or ethyl esters and which comprise relatively small amounts of omega-3 docosahexaenoic acid (DHA-n3), together with a common or alternative statin dose, may be used for the treatment of hypertriglyceridemia (either TG $\geq$ 500mg/dL, TG $\geq$ 200mg/dL, TG $\geq$ 150mg/dL, TG 200-499mg/dL, or TG 150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such method of treatment provides to a subject in need thereof a common or alternative statin dose together with a dose of at least 10mg HPA-N3 per day, alternatively at least 15mg HPA-N3 per day, alternatively at least 20mg HPA-N3 per day, alternatively at least 25mg HPA-N3 per day, alternatively at least 30mg HPA-N3 per day, alternatively at least 40mg HPA-N3 per day, alternatively at least 50mg HPA-N3 per day, alternatively at least 60mg HPA-N3 per day, alternatively at least 80mg HPA-N3 per day, alternatively at least 90mg HPA-N3 per day, alternatively at least 120mg HPA-N3 per day, alternatively at least 150mg HPA-N3 per day, alternatively at least 160mg HPA-N3 per day, alternatively at least 180mg HPA-N3 per day, alternatively at least 200mg HPA-N3 per day, alternatively at least 250mg HPA-N3 per day, alternatively at least 300mg HPA-N3 per day, alternatively at least 350mg HPA-N3 per day, alternatively at least 400mg HPA-N3 per day, alternatively at least 500mg HPA-N3 per day, alternatively at least 600mg HPA-N3 per day, alternatively at least 800mg HPA-N3 or its glycerol or ethyl esters per day, while providing less than 2000mg of DHA, alternatively less than 1900mg of DHA, alternatively less than 1500mg of DHA, alternatively less than 1200mg of DHA, alternatively less than 1000mg of DHA, alternatively less than 800mg of DHA, alternatively less than 700mg of DHA, alternatively less than 600mg of DHA, alternatively less than 500mg of DHA, alternatively less than 400mg of DHA, alternatively less than 350mg of DHA, alternatively less than 300mg of DHA, alternatively less than 250mg of DHA, alternatively less than 200mg of DHA, alternatively less than 150mg of DHA, alternatively less than 120mg of DHA, alternatively less than 100mg of DHA, alternatively less than 80mg of DHA, alternatively less than 60mg of DHA, alternatively less than 50mg of DHA, alternatively less than 40mg of DHA, alternatively less than 30mg of DHA, alternatively less than 25mg of DHA, alternatively less than 20mg of DHA or its glycerol or ethyl esters per day.

[0133] In further embodiments, the compositions of the present invention, which comprise significant amounts of omega-3 docosapentaenoic acid (DPA-n3) or its glycerol or ethyl esters and which comprise relatively small amounts of omega-3 docosahexaenoic acid (DHA-n3), together with a common or alternative statin dose, may be used for the treatment of hypertriglyceridemia (either $TG \geq 500\text{mg/dL}$, $TG \geq 200\text{mg/dL}$, $TG \geq 150\text{mg/dL}$, TG 200-499mg/dL, or TG 150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such method of treatment provides to a subject in need thereof a common or alternative statin dose together with a dose of at least 30mg DPA-N3 per day, alternatively at least 40mg DPA-N3 per day, alternatively at least 50mg DPA-N3 per day, alternatively at least 60mg DPA-N3 per day, alternatively at least 70mg DPA-N3 per day, alternatively at least 80mg DPA-N3 per day, alternatively at least 90mg DPA-N3 per day, alternatively at least 100mg DPA-N3 per day, alternatively at least 120mg DPA-N3 per day, alternatively at least 150mg DPA-N3 per day, alternatively at least 160mg DPA-N3 per day, alternatively at least 180mg DPA-N3 per day, alternatively at least 200mg DPA-N3 per day, alternatively at least 250mg DPA-N3 per day, alternatively at least 300mg DPA-N3 per day, alternatively at least 350mg DPA-N3 per day, alternatively at least 400mg DPA-N3 per day, alternatively at least 500mg DPA-N3 per day, alternatively at least 600mg DPA-N3 per day, alternatively at least 800mg DPA-N3 per day, alternatively at least 1000mg DPA-N3 per day, alternatively at least 1200mg DPA-N3 per day, alternatively at least 1500mg DPA-N3 or its glycerol or ethyl esters per day, while providing a relatively small amount of DHA-N3 such that the DHA:DPA dose ratio is no more than 15:1 of DHA:DPA, alternatively no more than 12:1 of DHA:DPA, alternatively no more than 10:1 of DHA:DPA, alternatively no more than 8:1 of DHA:DPA, alternatively no more than 5:1 of DHA:DPA, alternatively no more than 3:1 of DHA:DPA, alternatively no more than 2:1 of DHA:DPA, alternatively no more than 1:1 of DHA:DPA, alternatively no more than 1:2 of DHA:DPA, alternatively no more than 1:3 of DHA:DPA, alternatively no more than 1:5 of DHA:DPA, alternatively no more than 1:8 of DHA:DPA, alternatively no more than 1:10 of DHA:DPA, alternatively no more than 1:15 of DHA:DPA, alternatively a relative daily dose of no more than 1:20 of DHA:DPA.

[0134] In further embodiments, the compositions of the present invention, which comprise significant amounts of omega-3 docosapentaenoic acid (HPA-n3) or its glycerol or ethyl esters and which comprise relatively small amounts of omega-3

docosahexaenoic acid (DHA-n3), together with a common or alternative statin dose, may be used for the treatment of hypertriglyceridemia (either $TG \geq 500\text{mg/dL}$, $TG \geq 200\text{mg/dL}$, $TG \geq 150\text{mg/dL}$, TG 200-499mg/dL, or TG 150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such method of treatment provides to a subject in need thereof a common or alternative statin dose together with a dose of at least 10mg HPA-N3 per day, alternatively at least 15mg HPA-N3 per day, alternatively at least 20mg HPA-N3 per day, alternatively at least 25mg HPA-N3 per day, alternatively at least 30mg HPA-N3 per day, alternatively at least 40mg HPA-N3 per day, alternatively at least 50mg HPA-N3 per day, alternatively at least 60mg HPA-N3 per day, alternatively at least 70mg HPA-N3 per day, alternatively at least 80mg HPA-N3 per day, alternatively at least 90mg HPA-N3 per day, alternatively at least 100mg HPA-N3 per day, alternatively at least 120mg HPA-N3 per day, alternatively at least 150mg HPA-N3 per day, alternatively at least 160mg HPA-N3 per day, alternatively at least 180mg HPA-N3 per day, alternatively at least 200mg HPA-N3 per day, alternatively at least 250mg HPA-N3 per day, alternatively at least 300mg HPA-N3 per day, alternatively at least 350mg HPA-N3 per day, alternatively at least 400mg HPA-N3 per day, alternatively at least 500mg HPA-N3 per day, alternatively at least 600mg HPA-N3 per day, alternatively at least 800mg HPA-N3 per day, alternatively at least 1000mg HPA-N3 per day, alternatively at least 1200mg HPA-N3 per day, alternatively at least 1500mg HPA-N3 or its glycerol or ethyl esters per day, while providing a relatively small amount of DHA-N3 such that the DHA:HPA dose ratio is no more than 15:1 of DHA:HPA, alternatively no more than 12:1 of DHA:HPA, alternatively no more than 10:1 of DHA:HPA, alternatively no more than 8:1 of DHA:HPA, alternatively no more than 5:1 of DHA:HPA, alternatively no more than 3:1 of DHA:HPA, alternatively no more than 2:1 of DHA:HPA, alternatively no more than 1:1 of DHA:HPA, alternatively no more than 1:2 of DHA:HPA, alternatively no more than 1:3 of DHA:HPA, alternatively no more than 1:5 of DHA:HPA, alternatively no more than 1:8 of DHA:HPA, alternatively no more than 1:10 of DHA:HPA, alternatively no more than 1:15 of DHA:HPA, alternatively a relative daily dose of no more than 1:20 of DHA:HPA.

[0135] In yet other embodiments, the compositions of the present invention, which comprise significant amounts of omega-3-pentaenoic acids or their glycerol or ethyl esters and which comprise relatively small amounts of omega-3 docosahexaenoic

acid (DHA-n3), together with a common or alternative statin dose, may be used for the treatment of hypertriglyceridemia (either $TG \geq 500\text{mg/dL}$, $TG \geq 200\text{mg/dL}$, $TG \geq 150\text{mg/dL}$, $TG 200-499\text{mg/dL}$, or $TG 150-199\text{mg/dL}$), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such methods of treatment provides to a subject in need thereof with a common or alternative statin dose and a dose of at least 100mg omega-3-pentaenoic acids per day, alternatively at least 200mg omega-3-pentaenoic acids per day, alternatively at least 300mg omega-3-pentaenoic acids per day, alternatively at least 500mg omega-3-pentaenoic acids per day, alternatively at least 700mg omega-3-pentaenoic acids per day, alternatively at least 900mg omega-3-pentaenoic acids per day, alternatively at least 1000mg omega-3-pentaenoic acids per day, alternatively at least 1500mg omega-3-pentaenoic acids per day, alternatively at least 1900mg omega-3-pentaenoic acids per day, alternatively at least 2000mg omega-3-pentaenoic acids per day, alternatively at least 2500mg omega-3-pentaenoic acids per day, alternatively at least 2900mg omega-3-pentaenoic acids per day, alternatively at least 3000mg omega-3-pentaenoic acids per day, alternatively at least 3500mg omega-3-pentaenoic acids per day, alternatively at least 3900mg omega-3-pentaenoic acids per day, alternatively at least 4000mg omega-3-pentaenoic acids per day, alternatively at least 4100mg omega-3-pentaenoic acids per day, alternatively at least 4500mg omega-3-pentaenoic acids per day, alternatively at least 5000mg omega-3-pentaenoic acids per day, alternatively at least 5500mg omega-3-pentaenoic acids per day, alternatively at least 6000mg omega-3-pentaenoic acids per day or their glycerol or ethyl esters, while providing less than 1900mg of DHA, alternatively less than 1500mg of DHA, alternatively less than 1200mg of DHA, alternatively less than 1000mg of DHA, alternatively less than 800mg of DHA, alternatively less than 700mg of DHA, alternatively less than 600mg of DHA, alternatively less than 500mg of DHA, alternatively less than 400mg of DHA, alternatively less than 350mg of DHA, alternatively less than 300mg of DHA, alternatively less than 250mg of DHA, alternatively less than 200mg of DHA, alternatively less than 150mg of DHA, alternatively less than 120mg of DHA, alternatively less than 100mg of DHA, alternatively less than 80mg of DHA, alternatively less than 60mg of DHA, alternatively less than 50mg of DHA, alternatively less than 40mg of DHA, alternatively less than 30mg of DHA,

alternatively less than 25mg of DHA, alternatively less than 20mg of DHA or its glycerol or ethyl esters per day.

[0136] In other embodiments, the compositions of the present invention, which comprise significant amounts of omega-3-pentaenoic acids including DPA or their glycerol or ethyl esters and which comprise relatively small amounts of omega-3 docosahexaenoic acid (DHA-n3), together with a common or alternative statin dose, may be used for the treatment of hypertriglyceridemia (either TG $\geq$ 500mg/dL, TG $\geq$ 200mg/dL, TG $\geq$ 150mg/dL, TG 200-499mg/dL, or TG 150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such methods of treatment provides to a subject in need thereof with a common or alternative statin dose and a dose of at least 100mg omega-3-pentaenoic acids per day, alternatively at least 200mg omega-3-pentaenoic acids per day, alternatively at least 300mg omega-3-pentaenoic acids per day, alternatively at least 500mg omega-3-pentaenoic acids per day, alternatively at least 700mg omega-3-pentaenoic acids per day, alternatively at least 900mg omega-3-pentaenoic acids per day, alternatively at least 1000mg omega-3-pentaenoic acids per day, alternatively at least 1500mg omega-3-pentaenoic acids per day, alternatively at least 1900mg omega-3-pentaenoic acids per day, alternatively at least 2000mg omega-3-pentaenoic acids per day, alternatively at least 2500mg omega-3-pentaenoic acids per day, alternatively at least 2900mg omega-3-pentaenoic acids per day, alternatively at least 3000mg omega-3-pentaenoic acids per day, alternatively at least 3500mg omega-3-pentaenoic acids per day, alternatively at least 3900mg omega-3-pentaenoic acids per day, alternatively at least 4000mg omega-3-pentaenoic acids per day, alternatively at least 4100mg omega-3-pentaenoic acids per day, alternatively at least 4500mg omega-3-pentaenoic acids per day, alternatively at least 5000mg omega-3-pentaenoic acids per day, alternatively at least 5500mg omega-3-pentaenoic acids per day, alternatively at least 6000mg omega-3-pentaenoic acids per day or their glycerol or ethyl esters, while providing a relatively small amount of DHA-N3 such that the DHA:DPA ratio is no more than 15:1 of DHA:DPA, alternatively no more than 12:1 of DHA:DPA, alternatively no more than 10:1 of DHA:DPA, alternatively no more than 8:1 of DHA:DPA, alternatively no more than 5:1 of DHA:DPA, alternatively no more than 3:1 of DHA:DPA, alternatively no more than 2:1 of DHA:DPA, alternatively no more than 1:1 of DHA:DPA, alternatively no more than 1:2 of DHA:DPA, alternatively no more

than 1:3 of DHA:DPA, alternatively no more than 1:5 of DHA:DPA, alternatively no more than 1:8 of DHA:DPA, alternatively no more than 1:10 of DHA:DPA, alternatively no more than 1:15 of DHA:DPA, alternatively a relative amount of no more than 1:20 of DHA:DPA

[0137] In yet other embodiments, the compositions of the present invention, which comprise significant amounts of omega-3-pentaenoic acids including HPA or their glycerol or ethyl esters and which comprise relatively small amounts of omega-3 docosahexaenoic acid (DHA-n3), together with a common or alternative statin dose, may be used for the treatment of hypertriglyceridemia (either TG $\geq$ 500mg/dL, TG $\geq$ 200mg/dL, TG $\geq$ 150mg/dL, TG 200-499mg/dL, or TG 150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such methods of treatment provides to a subject in need thereof with a common or alternative statin dose and a dose of at least 100mg omega-3-pentaenoic acids per day, alternatively at least 200mg omega-3-pentaenoic acids per day, alternatively at least 300mg omega-3-pentaenoic acids per day, alternatively at least 500mg omega-3-pentaenoic acids per day, alternatively at least 700mg omega-3-pentaenoic acids per day, alternatively at least 900mg omega-3-pentaenoic acids per day, alternatively at least 1000mg omega-3-pentaenoic acids per day, alternatively at least 1500mg omega-3-pentaenoic acids per day, alternatively at least 1900mg omega-3-pentaenoic acids per day, alternatively at least 2000mg omega-3-pentaenoic acids per day, alternatively at least 2500mg omega-3-pentaenoic acids per day, alternatively at least 2900mg omega-3-pentaenoic acids per day, alternatively at least 3000mg omega-3-pentaenoic acids per day, alternatively at least 3500mg omega-3-pentaenoic acids per day, alternatively at least 3900mg omega-3-pentaenoic acids per day, alternatively at least 4000mg omega-3-pentaenoic acids per day, alternatively at least 4100mg omega-3-pentaenoic acids per day, alternatively at least 4500mg omega-3-pentaenoic acids per day, alternatively at least 5000mg omega-3-pentaenoic acids per day, alternatively at least 5500mg omega-3-pentaenoic acids per day, alternatively at least 6000mg omega-3-pentaenoic acids per day or their glycerol or ethyl esters, while providing a relatively small amount of DHA-N3 such that the DHA:HPA ratio is no more than 15:1 of DHA:HPA, alternatively no more than 12:1 of DHA:HPA, alternatively no more than 10:1 of DHA:HPA, alternatively no more than 8:1 of DHA:HPA, alternatively no more than 5:1 of DHA:HPA, alternatively no more than 3:1 of

DHA:HPA, alternatively no more than 2:1 of DHA:HPA, alternatively no more than 1:1 of DHA:HPA, alternatively no more than 1:2 of DHA:HPA, alternatively no more than 1:3 of DHA:HPA, alternatively no more than 1:5 of DHA:HPA, alternatively no more than 1:8 of DHA:HPA, alternatively no more than 1:10 of DHA:HPA, alternatively no more than 1:15 of DHA:HPA, alternatively a relative amount of no more than 1:20 of DHA:HPA. In further embodiments, the compositions of the present invention, which comprise significant amounts of omega-3-pentaenoic acids or their glycerol or ethyl esters and which comprise relatively small amounts of omega-3 docosahexaenoic acid (DHA-n3), together with a common or alternative statin dose, may be used for the treatment of hypertriglyceridemia (either TG $\geq$ 500mg/dL, TG $\geq$ 200mg/dL, TG $\geq$ 150mg/dL, TG 200-499mg/dL, or TG 150-199mg/dL), mixed dyslipidemia, or any other diseases or medical conditions specified above. Such methods of treatment provides to a subject in need thereof with a common or alternative statin dose and a dose of at least 100mg omega-3-pentaenoic acids per day, alternatively at least 200mg omega-3-pentaenoic acids per day, alternatively at least 300mg omega-3-pentaenoic acids per day, alternatively at least 500mg omega-3-pentaenoic acids per day, alternatively at least 700mg omega-3-pentaenoic acids per day, alternatively at least 900mg omega-3-pentaenoic acids per day, alternatively at least 1000mg omega-3-pentaenoic acids per day, alternatively at least 1500mg omega-3-pentaenoic acids per day, alternatively at least 1900mg omega-3-pentaenoic acids per day, alternatively at least 2000mg omega-3-pentaenoic acids per day, alternatively at least 2500mg omega-3-pentaenoic acids per day, alternatively at least 2900mg omega-3-pentaenoic acids per day, alternatively at least 3000mg omega-3-pentaenoic acids per day, alternatively at least 3500mg omega-3-pentaenoic acids per day, alternatively at least 3900mg omega-3-pentaenoic acids per day, alternatively at least 4000mg omega-3-pentaenoic acids per day, alternatively at least 4100mg omega-3-pentaenoic acids per day, alternatively at least 4500mg omega-3-pentaenoic acids per day, alternatively at least 5000mg omega-3-pentaenoic acids per day, alternatively at least 5500mg omega-3-pentaenoic acids per day, alternatively at least 6000mg omega-3-pentaenoic acids per day or their glycerol or ethyl esters, while providing a relatively small amount of DHA-N3 as compared to the amount of omega-3-pentaenoic acids (N3-5enoicFA) such that the DHA:N3-5enoic FA ratio is no more than 15:1 of DHA:N3-5enoicFA, alternatively no more than 12:1 of DHA:N3-5enoic

FA, alternatively no more than 10:1 of DHA:N3-5enoic FA, alternatively no more than 8:1 of DHA:N3-5enoic FA, alternatively no more than 5:1 of DHA:N3-5enoic FA, alternatively no more than 3:1 of DHA:N3-5enoic FA, alternatively no more than 2:1 of DHA:N3-5enoic FA, alternatively no more than 1:1 of DHA:N3-5enoic FA, alternatively no more than 1:2 of DHA:N3-5enoicFA, alternatively no more than 1:3 of DHA:N3-5enoic FA, alternatively no more than 1:5 of DHA:N3-5enoic FA, alternatively no more than 1:8 of DHA:N3-5enoic FA, alternatively no more than 1:10 of DHA:N3-5enoic FA, alternatively no more than 1:15 of DHA:N3-5enoic FA, alternatively a relative amount of no more than 1:20 of DHA:N3-5enoic FA.

[0138] In some embodiments, the composition of the present invention further comprises TPA at concentration of at least 0.05%. In some embodiments, the TPA concentration is about 0.01% to about 5%, alternatively about 0.05% to about 2%, alternatively about 0.1% to about 1%, alternatively about 0.2% to about 0.6%, alternatively about 0.4% to about 0.6%, alternatively about 0.5%.

[0139] The compositions of the present invention may also be taken as a general nutritional supplement.

[0140] On a EPA daily dose basis, the compositions of the present invention are preferably provided in a dose of between 100 mg and 10,000 mg/day, alternatively between 200 mg and 8,000 mg/day, alternatively between 300 mg and 6,000 mg/day, alternatively between 400 mg and 5,000 mg/day, alternatively between 500 mg and 4,000 mg/day.

[0141] On a EPA+DPA daily dose basis, the compositions of the present invention are preferably provided in a dose of between 100 mg and 10,100 mg/day, alternatively between 200 mg and 8,100 mg/day, alternatively between 300 mg and 6,100 mg/day, alternatively between 400 mg and 5,100 mg/day, alternatively between 500 mg and 4,100 mg/day.

[0142] On a EPA+HPA+DPA daily dose basis, the compositions of the present invention are preferably provided in a dose of between 100 mg and 10,100 mg/day, alternatively between 200 mg and 8,100 mg/day, alternatively between 300 mg and 6,100 mg/day, alternatively between 400 mg and 5,100 mg/day, alternatively between 500 mg and 4,100 mg/day.

[0143] On an omega-3-pentaenoic acid daily dose basis, the compositions of the present invention are preferably provided in a dose of between 100 mg and 10,100 mg/day, alternatively between 200 mg and 8,100 mg/day, alternatively between 300

mg and 6,100 mg/day, alternatively between 400 mg and 5,100 mg/day, alternatively between 500 mg and 4,100 mg/day.

[0144] The formulation may be a single daily dose preparation to give in one dose the above intakes, or may be in convenient divided doses, for example, a daily dose formed of two to four soft gelatin or other dosage forms, each containing 300-1500 mg of EPA, EPA+DPA, EPA+DPA+HPA, or omega-3-pentaenoic acids in any form embodied in the present invention.

[0145] Flavourants or emulsifiers may be included, for instance, to make the preparation palatable. Other conventional additives, diluents and excipients may be present. The preparation for ingestion may be in the form of a capsule, a dry powder, a tablet, a solution, an oil, an emulsion or any other appropriate form. The capsules may be hard or soft gelatin capsules, agar capsules, or any other appropriate capsule.

[0146] Use of the formulations of the invention in the manufacture of a medicament for the treatment or prevention of any disease or disorder, including those mentioned above, is included in the present invention.

[0147] The omega-3 fatty acid composition optionally includes chemical antioxidants, such as alpha tocopherol, which are administered in pure form or suspended in a vegetable oil, such as soybean oil or corn oil.

[0148] The blended fatty acid compositions may then be incorporated into any appropriate dosage form for oral, enteral, parenteral, rectal, vaginal, dermal or other route of administration. Soft or hard gelatin capsules, flavoured oil blends, emulsifiers or other liquid forms, and microencapsulate powders or other dry form vehicles are all appropriate ways of administering the products.

[0149] The formulated final drug product containing the omega-3 fatty acid composition may be administered to a mammal or patient in need thereof in a capsule, a tablet, a powder that can be dispersed in a beverage, or another solid oral dosage form, a liquid, a soft gel capsule or other convenient dosage form such as oral liquid in a capsule, as known in the art. In some embodiments, the capsule comprises a hard gelatin. The combination product may also be contained in a liquid suitable for injection or infusion.

[0150] Example pharmaceutical grade finished dosage forms: (a) Soft or hard gelatin capsules each containing 500 mg or 1000 mg of a mix 20 parts of EPA as a free fatty acid to 1 parts of DPA as a free fatty acid; (b) As in (a) but where the EPA

and DPA free fatty acids are replaced with the fatty acids in any other appropriate bioassimilable form such as the ethyl esters; (c) As in (a)-(b) but where the material is in the form of a microencapsulated powder which can be used as a powder or compressed into tablets. Such powders may be prepared by a variety of technologies known to those skilled in the art; (d) As in (a)-(b) but where the formulation is a liquid or emulsion, appropriately flavoured for palatable oral administration; (e) As in (a)-(b) but where the material is formulated into a pharmaceutically acceptable vehicle appropriate for topical application such as a cream or ointment.

[0151] The omega-3 compositions of the present invention may also be administered with a combination of one or more non-active pharmaceutical ingredients (also known generally herein as "excipients"). Non-active ingredients, for example, serve to solubilize, suspend, thicken, dilute, emulsify, stabilize, preserve, protect, color, flavor, and fashion the active ingredients into an applicable and efficacious preparation that is safe, convenient, and otherwise acceptable for use. Thus, the non-active ingredients may include colloidal silicon dioxide, crospovidone, lactose monohydrate, lecithin, microcrystalline cellulose, polyvinyl alcohol, povidone, sodium lauryl sulfate, sodium stearyl fumarate, talc, titanium dioxide and xanthum gum.

[0152] The term "pharmaceutically acceptable vehicle," as used herein, includes any of the following: a solution where the first API and optional other ingredients are wholly dissolved in a solubilizer (e.g., a pharmaceutically acceptable solvent or mixture of solvents), wherein the solution remains in clear liquid form at about room temperature; a suspension; an oil; or a semi-solid, wherein the first API and optionally other ingredients are dissolved wholly or partially in a solubilizer (an emulsion, cream, etc.).

[0153] A "pharmaceutical grade finished dosage form" as used herein may be construed as a unit dose form suitable for administration to, for example, human or animal subjects, and having content uniformity acceptable to regulatory authorities. For example, under the USP requirements for content uniformity, a pharmaceutical grade finished dosage form should have an amount of API within the range of 85% to 115% of the desired dosage and an RSD less than or equal to 6.0%. In addition, a pharmaceutical grade finished dosage form must be stable (i.e., have a "shelf life") for a pharmaceutically acceptable duration of time, preferably at least six months.

WO 2013/192109

PCT/US2013/046176

alternatively at least one year, or at least two years, when stored at room temperature (about 23 degree Celcius to 27 degree Celcius, preferably about 25 degree Celcius) and 60% relative humidity. Typically, stability is determined by physical appearance and/or chemical modification of the ingredients, in accordance with standards well-known in the pharmaceutical arts, including those documented in ICH guidelines.

[0154] The omega-3 fatty acid dosage form optionally includes chemical antioxidants, such as alpha tocopherol, oils, such as soybean oil and partially hydrogenated vegetable oil, and lubricants such as fractionated coconut oil, lecithin and a mixture of the same.

EXAMPLES

[0155] Example 1

[0156] A composition according to the present invention is prepared by mixing and homogenizing in a ratio of 98:2 the intermediates MEGAPEX E90D00EE (90% EPA ethyl ester,) and MAXOMEGA DPA95 FFA (95% DPA synthetic fatty acid produced from EPA ethyl ester concentrate) converted to ethyl ester, respectively. These intermediates were prepared and commercially offered for sale by Chemport Korea (MEGAPEX) and Equateq Ltd from Scotland, UK (MAXOMEGA). The relative amounts of fatty acids present in the starting intermediates and in the resulting novel composition are listed in Table 1 below. The resulting novel composition comprises 89.10% EPA, 1.95% DPA, 0.19% HPA, 91.24% omega-3-pentaenoic acids, less than 0.01% DHA, 91.24% omega-3-pentaenoic acids, 93.09% total omega-3 fatty acids, 3.15% ARA and 3.57% omega-6 fatty acids (all Area%).

WO 2013/192109

PCT/US2013/046176

Table 1. Fatty acid Composition (Area %) of intermediates and novel composition according to Example 1

Fatty Acid	98.0%	2.0%	Novel Composition
	Megapex E90D00EE	Maxomega DPA95FFA => EE	
c18:0	0.05	0	0.05
c18:1n7	0.06	0	0.06
c18:1n7	0.02	0	0.02
c18:2n6	0.01	0	0.01
c18:3n6	0.02	0	0.02
c18:3n3	0.03	0	0.03
c18:4n3	0.42	0	0.41
c18:4n1	0.07	0	0.07
c20:0	0	0	0.00
c20:1n11	0	0	0.00
c20:1n9	0	0	0.00
c20:1n7	0	0	0.00
c20:2n6	0.25	0	0.25
c20:3n9	0	0	0.00
c20:3n6	0.15	0	0.15
c21:0	0	0	0.00
c20:4n6	3.21	0	3.15
c20:5n3	0	0	0.00
c20:4n3	1.44	0	1.41
c20:5n3	90.92	0	89.10
c22:0	0.3	0	0.29
c22:1n11	0.07	0	0.07
c22:1n9	0.18	0	0.18
c22:1n7	0.19	0	0.19
c21:5n3	0.19	0	0.19
c22:5n6	0	0	0.00
c22:5n3	0	97.27	1.95
c22:6n3	0	0	0.00
c24:0	0	0.33	0.01
OTHER	2.42	2.4	2.42
	100	100	100

[0157] Example 2

[0158] A composition according to the present invention is prepared by mixing and homogenizing in a ratio of 96:4 the intermediates MEGAPEX E90D00EE (90% EPA ethyl ester,) and MAXOMEGA DPA95 FFA (≥95% DPA synthetic fatty acid produced from EPA ethyl ester concentrate), converted to ethyl ester, respectively. These intermediates were prepared and commercially offered for sale by Chemport Korea (MEGAPEX) and Equateq Ltd from Scotland, UK (MAXOMEGA). The relative amounts of fatty acids present in the starting intermediates and in the resulting novel composition is listed in Table 2 below. The resulting novel composition comprises

WO 2013/192109

PCT/US2013/046176

87.28% EPA, 3.89% DPA, 0.18% HPA, 91.35% omega-3-pentaenoic acids, less than 0.01% DHA, 93.17% total omega-3 fatty acids and 3.49% omega-6 fatty acids (all Area%).

Table 2. Fatty acid Composition (Area %) of intermediates and novel composition according to Example 2

Fatty Acid	96.0%	4.0%	Novel Composition
	Megapep E90D00EE	Maxomega DPA95FFA => EE	
c18:0	0.05	0	0.05
c18:1n9	0.06	0	0.06
c18:1n7	0.02	0	0.02
c18:2n6	0.01	0	0.01
c18:3n6	0.02	0	0.02
c18:3n3	0.03	0	0.03
c18:4n3	0.42	0	0.40
c18:4n1	0.07	0	0.07
c20:0	0	0	0.00
c20:1n11	0	0	0.00
c20:1n9	0	0	0.00
c20:1n7	0	0	0.00
c20:2n6	0.25	0	0.24
c20:3n9	0	0	0.00
c20:3n6	0.15	0	0.14
c21:0	0	0	0.00
c20:4n6	3.21	0	3.08
c20:3n3	0	0	0.00
c20:4n3	1.44	0	1.38
c20:5n3	90.92	0	87.28
c22:0	0.3	0	0.29
c22:1n11	0.07	0	0.07
c22:1n9	0.18	0	0.17
c22:1n7	0.19	0	0.18
c21:5n3	0.19	0	0.18
c22:5n6	0	0	0.00
c22:5n3	0	97.77	3.89
c22:6n3	0	0	0.00
c24:0	0	0.33	0.01
OTHER	2.42	2.4	2.43
	100	100	100

[0159] Example 3

[0160] A composition according to the present prevention is prepared by mixing and homogenizing in a ratio of 94:6 the intermediates MEGAPEX E90D00EE (90% EPA

WO 2013/192109

PCT/US2013/046176

ethyl ester,) and MAXOMEGA DPA95 FFA (295% DPA synthetic fatty acid produced from EPA ethyl ester concentrate) converted to ethyl ester, respectively. These intermediates were prepared and commercially offered for sale by Chemport Korea (MEGAPEX) and Equateq Ltd from Scotland, UK (MAXOMEGA). The relative amounts of fatty acids present in the starting intermediates and in the resulting novel composition are listed in table 3 below. The resulting novel composition comprises 85.46% EPA, 5.84% DPA, 0.18% HPA, 91.48% omega-3-pentaenoic acids, less than 0.01% DHA, 93.26% total omega-3 fatty acids, 3.02% ARA, and 3.42% omega-6 fatty acids (all Area%).

Table 3. Fatty acid Composition (Area %) of intermediates and novel composition according to Example 3

Fatty Acid	94.0%	6.0%	Novel Composition
	Megapex E90DCEEE	Maxomega DPA95FFA => EE	
c18:0	0.05	0	0.05
c18:1n9	0.06	0	0.06
c18:1n7	0.02	0	0.02
c18:2n6	0.01	0	0.01
c18:3n6	0.02	0	0.02
c18:3n3	0.03	0	0.03
c18:4n3	0.42	0	0.39
c18:4n1	0.07	0	0.07
c20:0	0	0	0.00
c20:1n11	0	0	0.00
c20:1n9	0	0	0.00
c20:1n7	0	0	0.00
c20:2n6	0.25	0	0.24
c20:3n3	0	0	0.00
c20:3n6	0.15	0	0.14
c21:0	0	0	0.00
c20:4n6	3.21	0	3.02
c20:3n3	0	0	0.00
c20:4n3	1.44	0	1.35
c20:5n3	90.92	0	85.46
c22:0	0.3	0	0.28
c22:1n11	0.07	0	0.07
c22:1n9	0.18	0	0.17
c22:1n7	0.19	0	0.18
c21:5n3	0.19	0	0.18
c22:5n6	0	0	0.00
c22:5n3	0	97.27	5.84
c22:6n3	0	0	0.00
c24:0	0	0.33	0.02
OTHER	2.42	2.3	2.42
	100	100	100

[0161] Example 4

[0162] A composition according to the present invention is prepared by mixing and homogenizing in a ratio of 75:25 the intermediates MEGAPEX E90D00EE (90% EPA ethyl ester,) and MAXOMEGA DPA95 FFA ($\geq 95\%$ DPA synthetic fatty acid produced from EPA ethyl ester concentrate, converted to ethyl ester, respectively. These intermediates were prepared and commercially offered for sale by Chemport Korea (MEGAPEX) and Equateq Ltd from Scotland, UK (MAXOMEGA). The relative amounts of fatty acids present in the starting intermediates and in the resulting novel composition is listed in table 4 below. The resulting novel composition comprises 68.10% EPA, 24.32% DPA, 0.19% HPA, 92.65% omega-3-pentaenoic acids, less than 0.01% DHA, 94.07% total omega-3 fatty acids, 2.41% ARA and 2.73% omega-6 fatty acids (all Area%).

WO 2013/192109

PCT/US2013/046176

Table 4. Fatty acid Composition (Area %) of intermediates and novel composition according to Example 4

Fatty Acid	75.0%	25.0%	Novel Composition
	Megapeks E90000EE	Maxomega DPA95FFA => EE	
c18:0	0.05	0	0.04
c18:1n9	0.06	0	0.05
c18:1n7	0.02	0	0.02
c18:2n6	0.01	0	0.01
c18:3n6	0.02	0	0.02
c18:3n3	0.03	0	0.02
c18:4n3	0.42	0	0.32
c18:4n1	0.07	0	0.05
c20:0	0	0	0.00
c20:1n11	0	0	0.00
c20:1n9	0	0	0.00
c20:1n7	0	0	0.00
c20:2n6	0.25	0	0.19
c20:3n9	0	0	0.00
c20:3n6	0.15	0	0.11
c21:0	0	0	0.00
c20:4n6	3.21	0	2.41
c20:3n3	0	0	0.00
c20:4n3	1.44	0	1.08
c20:5n3	90.92	0	68.19
c22:0	0.3	0	0.23
c22:1n11	0.07	0	0.05
c22:1n9	0.18	0	0.14
c22:1n7	0.19	0	0.14
c21:5n3	0.19	0	0.14
c22:5n6	0	0	0.00
c22:5n3	0	97.27	24.32
c22:6n3	0	0	0.00
c24:0	0	0.33	0.08
OTHER	2.42	2.4	2.42
	100	100	100

[0163] Example 5

[0164] A composition according to the present invention is prepared by mixing and homogenizing in a ratio of 60:40 the intermediates KD-PharmaKD-PUR 900EE and MAXOMEGA DPA95 FFA converted to ethyl ester, respectively. These intermediates were prepared and commercially offered for sale by KD-Pharma Germany (KD-Pharma) and Equateq Ltd from Scotland, UK (MAXOMEGA). The relative amounts of fatty acids present in the starting intermediates and in the resulting novel composition is listed in table 5 below. The resulting novel

WO 2013/192109

PCT/US2013/046176

composition comprises 55.74% EPA, 39.26% DPA, 2.39% HPA, 97.44% omega-3-pentaenoic acids, and 98.06% total omega-3 fatty acids (all Area%).

Table 5. Fatty acid Composition (Area %) of intermediates and novel composition according to Example 5

Fatty Acid	60.0% KD-Pur 900EE	40.0% Maxomega DPA95FFA => EE	Novel Composition
c18:0	0	0	0.00
c18:1n7	0	0	0.00
c18:1n6	0	0	0.00
c18:2n6	0	0	0.00
c18:3n6	0	0	0.00
c18:3n3	0	0	0.00
c18:4n3	0	0	0.00
c18:4n1	0	0	0.00
c20:0	0	0	0.00
c20:1n11	0	0	0.00
c20:1n9	0	0	0.00
c20:1n7	0	0	0.00
c20:2n6	0	0	0.00
c20:3n9	0	0	0.00
c20:3n6	0	0	0.00
c21:0	0	0	0.00
c20:4n6	0	0	0.00
c20:3n3	0	0	0.00
c20:4n3	1.04	0	0.62
c20:5n3	92.99	0	55.79
c22:0	0	0	0.00
c22:1n11	0	0	0.00
c22:1n9	0	0	0.00
c22:1n7	0	0	0.00
c21:5n3	3.98	0	2.39
c22:5n6	0	0	0.00
c22:5n3	0.58	97.27	39.26
c22:6n3	0	0	0.00
c24:0	0	0.33	0.13
OTHER	1.41	2.4	1.81
	100.00	100	100.00

[0165] Example 6

[0166] A composition according to the present invention is prepared by mixing and homogenizing in a ratio of 96:4 the intermediates KD-PUR 900EE KD-Pharma and MAXOMEGA DPA95 FFA converted to ethyl ester, respectively. These intermediates were prepared and commercially offered for sale by KD-Pharma Germany (KD-Pharma) and Equateq Ltd from Scotland, UK (MAXOMEGA). The

WO 2013/192109

PCT/US2013/046176

relative amounts of fatty acids present in the starting intermediates and in the resulting novel composition is listed in table 6 below. The resulting novel composition comprises 89.27% EPA, 4.45% DPA, 3.82% HPA, 97.54% omega-3-pentaenoic acids, and 98.54% total omega-3 fatty acids (all Area%).

Table 6. Fatty acid Composition (Area %) of intermediates and novel composition according to Example 6

Fatty Acid	96.0%	4.0%	Novel Composition
	ED-Pur 90XEE	Maxomega DPA95FFA -> EE	
c18:0	0	0	0.00
c18:1n7	0	0	0.00
c18:1n7	0	0	0.00
c18:2n6	0	0	0.00
c18:3n6	0	0	0.00
c18:3n3	0	0	0.00
c18:4n3	0	0	0.00
c18:4n1	0	0	0.00
c20:0	0	0	0.00
c20:1n11	0	0	0.00
c20:1n9	0	0	0.00
c20:1n7	0	0	0.00
c20:2n6	0	0	0.00
c20:3n9	0	0	0.00
c20:3n6	0	0	0.00
c21:0	0	0	0.00
c20:4n6	0	0	0.00
c20:3n3	0	0	0.00
c20:4n3	1.04	0	1.00
c20:5n3	92.99	0	89.27
c22:0	0	0	0.00
c22:1n11	0	0	0.00
c22:1n9	0	0	0.00
c22:1n7	0	0	0.00
c21:5n3	3.98	0	3.82
c22:5n6	0	0	0.00
c22:5n3	0.58	97.27	4.45
c22:6n3	0	0	0.00
c24:0	0	0.33	0.01
OTHER	1.41	2.4	1.45
	100.00	100	100.00

[0167] Example 7

[0168] The ethyl ester composition of Example 4 may be converted into a free fatty acid composition with essentially the same fatty acid composition according to "Conversion Method EE to FFA" below. This method is indiscriminate with respect to

the type, degree of saturation or length of fatty acid if performed for an adequate amount of time under the described conditions.

Conversion Method EE to FFA

1. Fatty Acid Ethyl Ester (FAEE GMP, approx. 3mmol/g) oil is brought into a closed heated/cooled reaction chamber under nitrogen atmosphere (preferably with pressure control), and heated to 50-60 degree Celcius under stirring.
2. 2M NaOH solution in water is added under firm stirring to ensure phase mixing (est. 2-3 x FAEE w/w) and stir until no ethyl ester is presence (est. 2-4 hrs). Test ethyl ester presence at lab scale/in process with TLC (hexanes/EtOAc 9:1) and with EP GC method to confirm reaction completion under GMP.
3. Under cooling (keep mixture below 70 degree Celcius), add 6M HCl in water (est. <1 hr) until slightly acid (~pH3-4). It may be necessary to control pressure to prevent excessive foaming. Then halt stirring, give time to let phases separate, and remove water phase from bottom (keep oil protected from oxygen, apply nitrogen atmosphere blanket).
4. Add demineralized water (est. 2-3 x FAEE w/w) and wash out NaCl and ethanol from oil under firm stirring (est. ~1hr). Halt stirring, give time to let phases separate, and remove water phase from bottom (keep oil protected from oxygen, apply nitrogen atmosphere blanket).
5. Repeat Step 4 several times (~2x) to remove ethanol and NaCl.
6. Remove water and remaining ethanol [determine in-process controls], confirm under GMP with USP residual solvent method (target: ethanol < 100ppm) by stirring oil while applying vacuum 10-50 mbar (with solvent trap) and heat oil (70-80 degree celcius) until water/ethanol target is met (est. 2-4 hrs).
7. Add anti-oxidants (i.e. alpha-D-tocopherol, USP, target 4 mg/g) and/or other excipients.
8. All reagents and excipients USP grade.

[0169] Example 8

[0170] The ethyl ester composition of Example 3 is converted into a free fatty acid composition with essentially the same fatty acid composition according to "Conversion Method EE to FFA" above. This method is indiscriminate with respect to the type, degree of saturation or length of fatty acid if performed for an adequate amount of time under the described conditions.

[0171] Example 9

[0172] The ethyl ester composition of Example 8 is converted into a free fatty acid composition with essentially the same fatty acid composition according to "Conversion Method EE to FFA" above. This method is indiscriminate with respect to the type, degree of saturation or length of fatty acid if performed for an adequate amount of time under the described conditions.

[0173] Example 10

[0174] The composition of Example 4 is formulated into a soft gelatin capsule. Prior to encapsulation, an anti-oxidant preparation (composed of 4000 mg alpha-D-tocopherol in one liter of corn oil; corn oil is a triglyceride low in omega-3) is added to the composition of Example 4, by mixing and homogenizing 100mL of this anti-oxidant preparation into 100 liters of the oil composition of Example 4 followed by thorough homogenization. The resulting pre-encapsulation formulated oil contains approximately 4mg/gram alpha-D-tocopherol. Subsequently, the formulated oil is encapsulated into soft gelatin capsules with printed logo according to general methods typically used by Accucaps in Canada for fish oils or by any other documented and operational encapsulation method. The fill mass of the oil is 1.08 gram/capsule, providing a dose of approximately 1000mg omega-3-pentaenoic-acids ethyl esters per capsule. Finally, the capsules are bottled in HDPE bottles with induction seal and child resistant cap.

[0175] Example 11

[0176] The composition of Example 8 is formulated into a soft gelatin capsule. Prior to encapsulation, an anti-oxidant preparation (composed of 4000 mg alpha-D-tocopherol in one liter of corn oil; corn oil is a triglyceride low in omega-3) is added to the composition of Example 4, by mixing and homogenizing 100mL of this anti-

oxidant preparation into 100 liters of the oil composition of Example 4 followed by thorough homogenization. The resulting pre-encapsulation formulated oil contains approximately 4mg/gram alpha-D-tocopherol. Subsequently, the formulated oil is encapsulated into soft gelatin capsules with printed logo according to general methods typically used by Banner in High Point, NC, for fish oils or by any other documented and operational encapsulation method. The fill mass of the oil is 1.09 gram/capsule, providing a dose of approximately 1000mg omega-3-pentaenoic-acids per capsule. Finally, the capsules are bottled in HDPE bottles with induction seal and child resistant cap.

[0177] Example 12

[0178] The composition of Example 5 is formulated into a soft gelatin capsule. Prior to encapsulation, an anti-oxidant preparation (composed of 4000 mg alpha-D-tocopherol in one liter of corn oil; corn oil is a triglyceride low in omega-3) is added to the composition of Example 4, by mixing and homogenizing 100mL of this anti-oxidant preparation into 100 liters of the oil composition of Example 4 followed by thorough homogenization. The resulting pre-encapsulation formulated oil contains approximately 4mg/gram alpha-D-tocopherol. Subsequently, the formulated oil is encapsulated into soft gelatin capsules with printed logo according to general methods typically used by Catalent in St.Petersburg, FL, for fish oils or by any other documented and operational encapsulation method. The fill mass of the oil is 1.05 gram/capsule, providing a dose of approximately 1000mg omega-3-pentaenoic-acids ethyl esters per capsule. Finally, the capsules are bottled in HDPE bottles with induction seal and child resistant cap.

[0179] Example 13

[0180] The composition of Example 9 is formulated into a soft gelatin capsule. Prior to encapsulation, an anti-oxidant preparation (composed of 4000 mg alpha-D-tocopherol in one liter of corn oil; corn oil is a triglyceride low in omega-3) is added to the composition of Example 4, by mixing and homogenizing 100mL of this anti-oxidant preparation into 100 liters of the oil composition of Example 4 followed by thorough homogenization. The resulting pre-encapsulation formulated oil contains approximately 4mg/gram alpha-D-tocopherol. Subsequently, the formulated oil is encapsulated into soft gelatin capsules with printed logo according to general

methods typically used by Banner in High Point, NC, for fish oils or by any other documented and operational encapsulation method. The fill mass of the oil is 1.08 gram/capsule, providing a dose of approximately 1000mg omega-3-pentaenoic-acids per capsule. Finally, the capsules are bottled in HDPE bottles with induction seal and child resistant cap.

[0181] Example 14

[0182] A patient is diagnosed with severe hypertriglyceridemia (TG>500mg/dL). Thereupon, the patient may be initiated on daily treatment with one of the encapsulated compositions according to Examples 10, 11, 12 or 13. Four capsules per day are administered to this patient (4g/d).

[0183] Example 15

[0184] A patient is treated as per Example 14. The treatment results in significant reduction of TG as well as non-HDL- and VLDL-cholesterol levels while the LDL-cholesterol level changes insignificantly.

[0185] Example 16

[0186] A patient is treated as per Example 14. The treatment results in significant reduction of TG as well as non-HDL-, LDL- and VLDL-cholesterol levels.

[0187] Example 17

[0188] A patient already undergoing treatment with a statin is diagnosed with high triglycerides (TG between 200 and 500mg/dL). Thereupon, the patient is initiated on daily treatment with one of the encapsulated compositions according to Examples 10, 11, 12 or 13. Four capsules per day are administered to this patient (4 g/d).

[0189] Example 18

[0190] A patient is treated as per Example 17. The treatment results in significant reduction of TG as well as non-HDL-, VLDL- and LDL-cholesterol levels.

[0191] Example 19

[0192] A patient is diagnosed with mixed dyslipidemia (TG between 200 and 700 mg/dL and LDL-cholesterol above 190 mg/dL). Thereupon, the patient is initiated on

WO 2013/192109

PCT/US2013/046176

concomitant daily treatment with a statin and one of the encapsulated compositions according to Examples 10, 11, 12 or 13. Four capsules per day are administered to this patient (4 g/d).

[0193] Example 20

[0194] A patient is treated as per Example 19. The treatment results in significant reduction of TG as well as non-HDL-, VLDL- and LDL-cholesterol levels.

[0195] Example 21

[0196] A patient is diagnosed with mixed dyslipidemia (TG between 200 and 700 mg/dL and non-HDL-cholesterol above 200 mg/dL). Thereupon, the patient is initiated on concomitant daily treatment with a statin and one of the encapsulated compositions according to Examples 10, 11, 12 or 13. Four capsules per day are administered to this patient (4 g/d).

[0197] Example 22

[0198] A patient is treated as per Example 21. The treatment results in significant reduction of TG as well as non-HDL-, VLDL- and LDL-cholesterol levels.

[0199] Example 23

[0200] A patient is diagnosed to be at significant risk for a cardiovascular event according to the NCEP guidelines and has TG levels above 150mg/dL. Thereupon, the patient is initiated on daily treatment with one of the encapsulated compositions according to Examples 10, 11, 12 or 13. Four capsules per day are administered to this patient (4 g/d).

[0201] Example 24

[0202] A patient is treated as per Example 23. The treatment results in significant reduction of TG as well as non-HDL-, VLDL- and LDL-cholesterol levels.

[0203] Example 25

[0204] A patient diagnosed as per Example 14, 17, 19, 21 or 23 is treated with 3 capsules per day (instead of 4) of one of the encapsulated compositions according to

Examples 10, 11, 12 or 13. The treatment results in significant reduction of TG as well as non-HDL- and VLDL-cholesterol levels.

[0205] Example 26

[0206] A patient diagnosed as per Example 14, 17, 19, 21 or 23 is treated with 3 capsules per day (instead of 4) of one of the encapsulated compositions according to Examples 10, 11, 12 or 13. The treatment results in significant reduction of TG as well as non-HDL-, VLDL- and LDL-cholesterol levels.

[0207] Example 27

[0208] A patient diagnosed as per Example 14, 17, 19, 21 or 23 is treated with 2 capsules per day (instead of 3 or 4) of one of the encapsulated compositions according to Examples 10, 11, 12 or 13. The treatment results in significant reduction of TG as well as non-HDL- and VLDL-cholesterol levels.

[0209] Example 28

[0210] A patient diagnosed as per Example 14, 17, 19, 21 or 23 is treated with 2 capsules per day (instead of 3 or 4) of one of the encapsulated compositions according to Examples 10, 11, 12 or 13. The treatment results in significant reduction of TG as well as non-HDL-, VLDL- and LDL-cholesterol levels.

[0211] Example 29

[0212] The following is an example of a preferred embodiment of the present invention.

COMPOSITION 1			
Composition	Minimum (mg/g)	Maximum (mg/g)	Target (mg/g)
Omega-3 pentaenoic acid	880	980	930
Eicosapentaenoic acid (EPA)	800	950	850
Heptacosapentaenoic acid (HPA)	5	60	30
Docosapentaenoic acid (DPA)	60	100	80
Docosahexaenoic acid (DHA)		25	<10

WO 2013/192109

PCT/US2013/046176

In COMPOSITION 1, the EPA:HPA ratio is between 13 and 190, the EPA:DPA ratio is between 8 and 15, the HPA:DPA ratio between 0.05 and 1, the DPA:DHA ratio more than 2.4, preferably more than 4, more preferably more than 6, most preferably more than 10, and the EPA:DHA ratio more than 32, preferably more than 38, more preferably more than 80, most preferably more than 95. The EPA, HPA, DPA and DHA may be composed as a glyceride (such as triglyceride), an ester (such as ethyl ester), or a free fatty acid.

[0213] Example 30

[0214] The following is an example of a preferred embodiment of the present invention.

COMPOSITION 2			
Composition	Minimum (mg/g)	Maximum (mg/g)	Target (mg/g)
Omega-3 pentaenoic acid	900	980	940
Eicosapentaenoic acid (EPA)	15	60	30
Heneicosapentaenoic acid (HPA)	5	60	30
Docosapentaenoic acid (DPA)	800	950	880
Docosahexaenoic acid (DHA)		25	<10

In COMPOSITION 2, the EPA:HPA ratio is between 0.25 and 12, the DPA:EPA ratio is between 13 and 83, the DPA:HPA ratio between 13 and 190, the DPA:DHA ratio more than 32, preferably more than 38, more preferably more than 80, most preferably more than 95, and the EPA:DHA ratio more than 00.6, preferably more than 1.5, more preferably more than 2.4, most preferably more than 6. The EPA, HPA, DPA and DHA may be composed as a glyceride (such as triglyceride), an ester (such as ethyl ester), or a free fatty acid.

[0215] Example 31

[0216] The following is an example of an embodiment of the present invention.

COMPOSITION 3			
Composition	Minimum	Maximum	Target (mg/g)

	(mg/g)	(mg/g)	
Docosapentaenoic acid (DPA n-3)	800	990	920

The DPA may be composed as a glyceride (such as triglyceride), an ester (such as ethyl ester), or a free fatty acid.

[0217] Example 32

[0218] A mixture of DPA and EPA was prepared by combining 1g DPA Ethyl Ester (SE-133-III) with 10g EPA Ethyl Ester, 914 mg/g (KD Pharma FM13001) in 150ml of 95% ethanol/water containing 35ml of 2M sodium hydroxide. This reaction mixture was stirred overnight at ambient temperature. Tlc analysis showed complete conversion of the ethyl esters to the corresponding acids. The reaction mixture was cooled in an ice bath, acidified with 6N hydrochloric acid and concentrated on a rotavap under reduced pressure. Water and ethyl acetate were added, the phases separated and the aqueous residue extracted with ethyl acetate. The ethyl acetate extracts were combined, dried over sodium sulfate and concentrated to dryness on a rotavap under reduced pressure. Yield: 9.83 g. The ethyl ester mixture was then converted to the free fatty acids as described in example 7.

[0219] A representative sample of this ethyl ester composition was analysed using split inject by capillary gas chromatography by a 30 meter x 0.25 mm Restek Stabilwax column using temperature programming.

[0220] Example 33

[0221] The following describes studies (Study #1 and Study #2) to determine the effect of compositions of the present invention.

[0222] STUDY #1 - Zucker rats

Male, eight to nine week old non-diabetic Zucker rats are used, with eight rats (n=8) per group. Examples of compositions of the present invention that are evaluated included COMPOSITION 1 of Example 29 and COMPOSITION 3 of Example 31. In addition to exemplary compositions of the present invention, VASCEPA® (icosapentyl ethyl; ethyl ester of eicosapentaenoic acid) and atorvastatin are used as reference compounds. Vegetable oil is used as a diluent, and to prepare the atorvastatin for dosing. A separate group of animals receiving vegetable oil alone is

used as the untreated control group. Animals receive daily doses of respective solutions by oral gavage. The study is conducted in 2 phases. In the first phase, COMPOSITION 1, COMPOSITION 3, and VASCEPA® are administered at 200 mg/kg, 400 mg/kg, and 1000 mg/kg. An additional 50 mg/kg dose group is included for the evaluation of MAT 8003 potency. Animals are dosed daily for 14 days. COMPOSITION 1 and COMPOSITION 3 are used in their free acid forms. The second phase is initiated on day 15, with the groups receiving Formulation 1 and Formulation 3 and VASCEPA® at doses of 400 mg/kg being co-administered atorvastatin. Another group receiving atorvastatin is included as a reference control. This second phase consisted of 14 days of daily, oral administration. Plasma triglycerides are measured in the fasting state on day 0, 7 and 14; and for those groups included in the second phase on days 21 and 28. In addition, total cholesterol, LDL, HDL, VLDL and NEFA (non-esterified free fatty acids) are quantified or calculated as secondary lipid parameters, along with assessment of glucose, ALT and AST concentrations. All testing is performed using standard techniques on a chemistry analyzer.

[0223] STUDY #2 - Humans

The overall pharmacokinetics of one of COMPOSITION 1, 2 or 3 are evaluated versus a reference compound after administration under fasting or fed conditions in normal, mostly healthy volunteers in a standard, 4-way cross-over trial design format. VASCEPA® is used as a reference compound. A total of 48 subjects are separated into 2 groups of 24. Each subject serves as his or her own internal control for comparison purposes under this 4-way crossover design. Inclusion criteria for tested subjects include volunteers between ages 18-65, with a BMI of 30-35 (alternatively a BMI of 27-35) and triglyceride levels less than 350 mg/dL, who consume no more than 1 fish meal per week and who are not currently prescribed pharmaco-therapy for lowering triglycerides, including but not limited to fibrates, omega-3 agents, and niacin. Volunteers on stable anti-hypertensive, anti-diabetic and thyroid therapy are allowed for consideration. Any person on stable statin therapy is considered if their triglyceride levels are less than 350 mg/dL. However, the total composition of subjects in the study with this particular profile is limited to no more than 30%.

WO 2013/192109

PCT/US2013/046176

[0224] Volunteers self-administering omega-3 non-prescription dietary supplements are asked to refrain from their use 2 weeks prior to the initiation of the study until study completion.

[0225] Pharmacokinetic Study#2 Design

[0226] The effect of oral administration of the compounds tested in this study is evaluated under fed versus fasting administration conditions, in order to determine drug pharmacokinetics, as well as to understand the effects of food on drug pharmacokinetics. COMPOSITION 1 or VASCEPA® are dosed at 4 grams/day in the morning by administration of 4 capsules containing 1 gram of each compound. Several days prior to pharmacokinetic evaluations, volunteers are housed at the testing facility in order to ensure well-controlled experimental conditions.

[0227] Compounds are given to volunteers following an overnight fast, with plasma samples obtained prior to dosing and at various time points after day 1 and day 14 dosing. Volunteers are allowed access to water, and well-defined meals at certain times. Afterwards, compound administration is stopped for a 2-4 week washout period, and the groups are switched with respect to which compound they would receive, meaning that the group initially receiving COMPOSITION 1 is switched to receive VASCEPA®, and vice versa. Fasting pharmacokinetics are determined using the procedure described above. Following completion of the second 14-day dosing cycle, compound dosing is stopped for a 2-4-week washout period prior to the initiation of the a similar cycle as above, now with COMPOSITION 1 and VASCEPA® administered together with a meal.

[0228] Plasma levels of omega-3 fatty acids of interest from the study are determined utilizing an analytical LC/MS technique under GLP laboratory conditions in order to determine C_{max}, T_{max} and AUC for the omega-3 fatty acids of interest, including EPA, DPA, HPA, DHA, and other omega-3 fatty acids.

[0229] Results

[00010] The results of the study show that COMPOSITION 1 has a better bioavailability (as measured by AUC and C_{max}) than Vascepa. This effect is seen under fasting and/or fed administration conditions.

[00011] In an alternate study design, this Study #2 is conducted with a certain dose level of COMPOSITION 2 OR COMPOSITION 3 instead if COMPOSITION 1.

[0100] Example 34

[0101] The following describes a study (STUDY #3) to determine the effect of compositions of the present invention.

[0102] A Multi-Center, Placebo-Controlled, Randomized Double-Blind, 12-Week Study to Evaluate the Efficacy and Safety of COMPOSITION 1 2 or 3 in Patients With Fasting Triglyceride Levels ≥ 500 mg/dL and ≤ 2000 mg/dL:

[0103] This Phase 3, multi-center study consists of a 6- to 8-week screening/washout period (to include a diet and lifestyle stabilization period), which includes a fasting triglyceride (TG) qualifying period of 2-3 weeks, followed by a 12-week double-blind treatment period. Patients on statin therapy (with or without ezetimibe) at screening are evaluated by the investigator as to whether this therapy could be safely discontinued at screening, or if it is to be continued. Patients on any other dyslipidemia therapy need to discontinue these in order to qualify for the study. If statin therapy (with or without ezetimibe) is to be continued, dose(s) must be stable for ≥ 4 weeks prior to the fasting TG baseline qualifying measurements for randomization. The screening visit is to occur at either 6 weeks before randomization for patients not on lipid-altering therapy at screening or for patients who do not need to discontinue their current dyslipidemia therapy, or at 8 weeks before randomization for patients who require washout of their current dyslipidemia therapy at screening.

[0104] The population for this study is men and women >18 years of age with a body mass index (BMI) ≤ 45 kg/m². Patients on lipid-lowering therapy and patients not on lipid-lowering therapy are eligible to enroll in the study. Patients had to have an average TG level ≥ 500 mg/dL and ≤ 2000 mg/dL during the screening period to be eligible for randomization.

[0105] After confirmation of qualifying fasting TG values, eligible patients will enter a 12-week randomized, double-blind treatment period. At Week 0, patients will be randomly assigned to 1 of the following treatment groups: COMPOSITION 1 (2 g daily), COMPOSITION 1 (3 g daily), COMPOSITION 1 (4 g daily), or placebo. The daily dose may be taken as either a single dose or distributed over two doses per day.

[0106] Approximately 80 patients per treatment group will be randomized in this study. Stratification will be by baseline fasting TG level (≤ 750 mg/dL or >750 mg/dL), gender, and the use of statin therapy at randomization. During the double-blind

treatment period, patients return to the site at Week 4, Week 11, and Week 12 for efficacy and safety evaluations.

[0107] The primary objective of the study is to determine the efficacy of COMPOSITION 1 at a 2 g daily dose, 3 g daily dose and 4 g daily dose, compared to placebo, in lowering fasting TG levels in patients with fasting TG levels ≥ 500 mg/dL and ≤ 2000 mg/dL.

[0108] The secondary and exploratory objectives of the study are as follows:

[0109] 1. To determine the safety and tolerability of COMPOSITION 1 2 g daily, 3 g daily and 4 g daily;

[0110] 2. To determine the effect of COMPOSITION 1 on lipid profiles, including total cholesterol (TC), non-high-density lipoprotein cholesterol (non-HDL-C), low-density lipoprotein cholesterol (LDL-C), high-density lipoprotein cholesterol (HDL-C), and very low-density lipoprotein cholesterol (VLDL-C);

[0111] 3. To determine the effect of COMPOSITION 1 on apolipoprotein A-I (apo A-I), apolipoprotein B (apo B), apo A-I/apo B ratio, lipoprotein(a) (Lp[a]), and lipoprotein-associated phospholipase A2 (Lp-PLA2);

[0112] 4. To determine the effect of COMPOSITION 1 on low-density lipoprotein (LDL) particle number and size, on oxidized LDL and on C-reactive protein (CRP);

[0113] 5. To determine the effect of COMPOSITION 1 on intracellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule 1 (VCAM-1), interleukin-1 β (IL-1 β), interleukin-2 (IL-2), interleukin-6 (IL-6), interleukin-8 (IL-8), interleukin-10 (IL-10), interleukin-12 (IL-12), interleukin-15 (IL-15), interleukin-18 (IL-18), tumor necrosis factor-alpha (TNF- α), tumor necrosis factor-beta (TNF- β) and plasminogen activator inhibitor-1 (PAI-1);

[0114] 6. To determine the effects of COMPOSITION 1 on nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B), vitronectin receptor (α v β 3), glycoprotein IIb/IIIa (gpIIb/IIIa) and other platelet and thrombogenic factors.

[0115] 7. To determine the effects of COMPOSITION 1 on E-selectin, P-selectin, homocysteine, thromboxane B2 (TXB2), thromboxane A2 (TXA2), thromboxane B23 (TXB3), thromboxane A3 (TXA3), 2,3-dinor thromboxane B2, free fatty acids (FFA or NEFA), serum amyloid A1, serum amyloid A2, serum amyloid A3, serum amyloid A4, thiobarbituric acid (TBA) reacting material, adiponectin (GBP-28), hemoglobin A1c (HbA1c), fasting insulin, fasting glucagon, fasting plasma glucose, fasting plasma

fructosamine, macrophage colony stimulating factor (M-CSF) and granulocyte macrophage colony stimulating factor (GM-CSF).

[0116] 8. To determine the effects of COMPOSITION 1 on fibrinogen, fibrin D-dimer, platelet derived-microparticles, mean platelet volume (MPV), platelet subpopulations, adenosine diphosphate induced platelet aggregation, platelet endothelial cell adhesion molecule (PECAM-1), heart rate, and systolic and diastolic blood pressure.

[0117] 9. To investigate the relationship between changes in fatty acid concentrations (including EPA, DHA and DPA) in plasma and red blood cell membranes and the reduction in fasting TG levels.

[0118] 10. To investigate the relationship between changes in fatty acid concentrations (including EPA, DHA and DPA) in plasma and red blood cell membranes and the reduction in fasting TG levels..

[0119] The primary efficacy variable for the double-blind treatment period is percent change in fasting TG from baseline to the Week 12 endpoint.

[0120] The secondary efficacy variable for the double-blind treatment period includes the following: Percent changes in fasting Non-HDL-C, LDL-C, VLDL-C, HDL-C, Lp-PLA2, and apo B from baseline to Week 12 endpoint.

[0121] Statistical methods for efficacy evaluations will be conducted on the intent-to-treat (ITT) and on the per-protocol population. Descriptive statistics for the baseline and post-baseline measurements, the percent changes, or changes from baseline are to be presented by treatment group and by visit for all efficacy variables.

[0122] The primary and secondary efficacy analyses will be performed using an analysis of covariance (ANCOVA) model with treatment, gender, and the use of statin therapy at randomization as factors and baseline fasting TG value as a covariate.

[0123] In an alternate study design, this Study #3 is conducted with one or more dose levels of COMPOSITION 2 OR 3 instead if COMPOSITION 1.

[0124] Example 35

[0125] The following describes a study (STUDY #4) to determine the effect of compositions of the present invention.

[0126] A Multi-Center, Placebo-Controlled, Randomized, Double-Blind, 6- to 12-Week Study to Evaluate the Efficacy and Safety of COMPOSITION 1, 2 or 3 in Statin-Treated Patients With High Fasting Triglyceride Levels ≥ 200 mg/dL and ≤ 499 mg/dL.

[0127] This multi-center study consists of a 4- to 6-week screening and washout period (to include a diet and lifestyle stabilization period, and to wash-out any non-statin/ezetimibe dyslipidemia medications), which also includes a 2-3 week fasting triglyceride (TG) level qualifying period, followed by a 6- to 12-week double-blind treatment period. Patients on statin therapy (with or without ezetimibe) at screening are evaluated by the investigator as to whether this therapy does maintain low-density lipoprotein (LDL) levels of ≥ 40 mg/dl and < 100 mg/dl. At screening, statin therapy (with or without ezetimibe) is to be initiated, in those patients who are not on statin therapy in order to achieve LDL levels of ≥ 40 mg/dl and < 100 mg/dl. Dose(s) of statin therapy must be stable for ≥ 4 weeks prior to the TG baseline qualifying measurements for randomization.

[0128] The population for this study is men and women > 18 years of age with a body mass index (BMI) ≤ 45 kg/m². Patients on lipid-lowering therapy and patients not on lipid-lowering therapy are eligible to enroll in the study. Patients had to have an average fasting TG level ≥ 200 mg/dL and ≤ 499 mg/dL during the qualifying period to be eligible for randomization.

[0129] After confirmation of qualifying fasting TG values, eligible patients will enter a 6- to 12-week randomized, double-blind treatment period. At Week 0, patients will be randomly assigned to one of the following treatment groups: COMPOSITION 1 at a 2 gram daily dose, COMPOSITION 1 at a 3 gram daily dose, COMPOSITION 1 at a 4 gram daily dose, or placebo. The daily dose may be taken as either a single dose or distributed over two doses per day.

[0130] Approximately 220 patients per treatment group will be randomized in this study. Stratification will be by gender. During the double-blind treatment period, patients will return to the site at Week 3 or 4, one week prior to the last week of randomized treatment period, and at the end of the randomized treatment period for efficacy and safety evaluations.

[0131] The primary objective of the study is to determine the efficacy of COMPOSITION 1 at 2 grams daily, 3 grams daily and 4 grams daily, compared to

placebo, in lowering fasting TG levels in statin-treated patients with fasting TG levels ≥ 200 mg/dL and ≤ 499 mg/dL.

[0132] The secondary and exploratory objectives of the study may include but are not limited to the following objectives:

[0133] 1. To determine the safety and tolerability of COMPOSITION 1 or 2 at 2 g daily, 3 g daily and 4 g daily;

[0134] 2. To determine the effect of COMPOSITION 1 at on lipid profiles, including total cholesterol (TC), non-high-density lipoprotein cholesterol (non-HDL-C) low-density lipoprotein cholesterol (LDL-C), high-density lipoprotein cholesterol (HDL-C), and very low-density lipoprotein cholesterol (VLDL-C);

[0135] 3. To determine the effect of COMPOSITION 1 on apolipoprotein A-I (apo A-I), apolipoprotein B (apo B), apo A-I/apo B ratio, lipoprotein(a) (Lp[a]), and lipoprotein-associated phospholipase A2 (Lp-PLA2);

[0136] 4. To determine the effect of COMPOSITION 1 on low-density lipoprotein (LDL) particle number and size, on oxidized LDL, high-sensitivity C-reactive protein (HSCRP), and on C-reactive protein (CRP);

[0137] 5. To determine the effect of COMPOSITION 1 on intracellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule 1 (VCAM_1), interleukin-1 β (IL-1 β), interleukin-2 (IL-2), interleukin-6 (IL-6), interleukin-8 (IL-8), interleukin-10 (IL-10), interleukin-12 (IL-12), interleukin-15 (IL-15), interleukin-18 (IL-18), tumor necrosis factor-alpha (TNF- α), tumor necrosis factor-beta (TNF- β) and plasminogen activator inhibitor-1 (PAI-1);

[0138] 6. To determine the effects of COMPOSITION 1 on nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B), vitronectin receptor (α v β 3), glycoprotein IIb/IIIa (gpIIb/IIIa) and other platelet and thrombogenic factors;

[0139] 7. To determine the effects of COMPOSITION 1 on E-selectin, P-selectin, homocysteine, thromboxane B2 (TXB2), thromboxane A2 (TXA2), thromboxane B23 (TXB3), thromboxane A3 (TXA3), 2,3-dinor thromboxane B2, free fatty acids (FFA or NEFA), serum amyloid A1, serum amyloid A2, serum amyloid A3, serum amyloid A4, thiobarbituric acid (TBA) reacting material, adiponectin (GBP-28), hemoglobin A1c (HbA1c), fasting insulin, fasting glucagon, fasting plasma glucose, fasting plasma fructosamine, macrophage colony stimulating factor (M-CSF) and granulocyte macrophage colony stimulating factor (GM-CSF).

[0140] 8. To determine the effects of COMPOSITION 1 on fibrinogen, fibrin D-dimer, platelet derived-microparticles, mean platelet volume (MPV), platelet subpopulations, adenosine diphosphate induced platelet aggregation, platelet endothelial cell adhesion molecule (PECAM-1), heart rate, and systolic and diastolic blood pressure.

[0141] 9. To determine the effects of COMPOSITION 1 on fatty acid concentrations (including EPA, DHA and DPA) in plasma and red blood cell membranes;

[0142] 10. To investigate the relationship between changes in fatty acid concentrations (including EPA, DHA and DPA) in plasma and red blood cell membranes and the reduction in fasting TG levels.

[0143] The primary efficacy variable for the double-blind treatment period is percent change in fasting TG from baseline to the Week 6 to 12 endpoint.

[0144] The secondary efficacy variable for the double-blind treatment period include but are not limited to the following: Percent changes in fasting Non-HDL-C, LDL-C, VLDL-C, HDL-C, Lp-PLA2, and apo B from baseline to Week 6 to 12 endpoint.

[0145] Statistical methods for efficacy evaluations will be conducted on the intent-to-treat (ITT) and on the per-protocol population. Descriptive statistics for the baseline and post-baseline measurements, the percent changes, or changes from baseline are to be presented by treatment group and by visit for all efficacy variables.

[0146] The primary and secondary efficacy analyses will be performed using an analysis of covariance (ANCOVA) model with treatment, gender, the type of statin therapy and diagnosis of diabetes at randomization as factors and baseline fasting TG value as a covariate.

[0147] In an alternate study design, this Study #4 is conducted with one or more dose levels of COMPOSITION 2 OR 3 instead of COMPOSITION 1.

[0148] Example 36

[0149] The following describes a study (STUDY #5) to determine the effect of compositions of the present invention.

[0150] The impact on fasting triglyceride levels and other pharmacodynamic endpoints of one of COMPOSITION 1, 2 or 3 are evaluated versus a reference compound after administration under fasting or fed conditions in normal, mostly healthy volunteers in a standard, 4-way cross-over trial design format. VASCEPA®

is used as a reference compound. A total of 48 subjects are separated into 2 groups of 24. Each subject serves as his or her own internal control for comparison purposes under this 4-way crossover design. Inclusion criteria for tested subjects include volunteers between ages 18-65, with a BMI of 30-35 (alternatively a BMI of 27-35) and triglyceride levels less than 350 mg/dL, who consume no more than 1 fish meal per week and who are not currently prescribed pharmacotherapy for lowering triglycerides, including but not limited to fibrates, omega-3 agents, and niacin. Volunteers on stable anti-hypertensive, anti-diabetic and thyroid therapy are allowed for consideration. Any person on stable statin therapy is considered if their triglyceride levels are less than 350 mg/dL. However, the total composition of subjects in the study with this particular profile is limited to no more than 30%.

[0151] Volunteers self-administering omega-3 non-prescription dietary supplements are asked to refrain from their use 2 weeks prior to the initiation of the study until study completion. Subjects using any other non-steroidal anti-inflammatory agents other than acetaminophen are asked to abstain and switch to acetaminophen for relief of pain, or are excluded from study consideration. Subjects are excluded if they receive any type of hormone therapy, weight loss agents, HIV therapy, beta-blockers, or are diagnosed with known cardiovascular disease, including heart failure, arrhythmia, any incidence of acute coronary syndrome, myocardial infarct, coronary artery bypass graft surgery, and/or angioplasty.

[0152] STUDY #5 Design

[0153] The effect of oral administration of the compounds tested in this study is evaluated under fed versus fasting administration conditions, in order to determine drug pharmacodynamics and effects on lipids. COMPOSITION 1 or VASCEPA® are dosed at 4 grams/day in the morning by administration of 4 capsules containing 1 gram of each compound.

[0154] Compounds are given to volunteers following an overnight fast, with plasma samples obtained prior to dosing on day 1 and day 14 dosing. Volunteers are allowed access to water, and well-defined meals at certain times. Afterwards, compound administration is stopped for a 2-4 week washout period, and the groups are switched with respect to which compound they would receive, meaning that the group initially receiving COMPOSITION 1 is switched to receive VASCEPA®, and vice versa. Following completion of the second 14-day dosing cycle, compound dosing is stopped for a 2-4-week washout period prior to the initiation of the a similar

cycle as above, now with COMPOSITION 1 and VASCEPA® administered together with a meal.

[0155] Baseline plasma levels of fasting serum triglycerides are determined on day -1 and just prior to initiation and completion of either the fasting or fed multi-dosing period. Additional lipid and other parameters (see below under "Results") that are analyzed included total cholesterol, LDL, HDL, VLDL, and NEFA as previously described.

[0156] Effects on platelet function, such as clotting time and PAF-induced aggregation are also determined. Standard physiological, plasma and urinary safety markers, including but not limited to electrolytes, ALT, AST, BUN, glucose, blood pressure, weight etc. are monitored in accordance with standard good clinical trial guidelines.

[0157] Results

[00012] The results of the study show that COMPOSITION 1 has a better fasting triglyceride lowering effect than Vascepa. This effect is seen under fasting and/or fed administration conditions. Administration of COMPOSITION 1 have a beneficial effect, versus baseline and versus Vascepa, on other lipid parameters (such as HDL cholesterol, total cholesterol, non-HDL cholesterol, VLDL cholesterol), on platelet function, and one or more of the following: apolipoprotein A-I (apo A-I), apolipoprotein B (apo B), apo A-I/apo B ratio, lipoprotein(a) (Lp[a]), lipoprotein-associated phospholipase A2 (Lp-PLA2), low density lipoprotein (LDL) particle number and size, oxidized LDL, C-reactive protein (CRP), high sensitivity C-reactive protein (HSCRP), intracellular adhesion molecule-1 (ICAM-1), E-selectin, P-selectin, vascular cell adhesion molecule 1 (VCAM-1) or cluster of differentiation 106 (CD106), interleukin-1 β (IL-1 β), interleukin-2 (IL-2), interleukin-6 (IL-6), interleukin-8 (IL-8), interleukin-10 (IL-10), interleukin-12 (IL-12), interleukin-15 (IL-15), interleukin-18 (IL-18), tumor necrosis factor-alpha (TNF- α), tumor necrosis factor-beta (TNF- β), plasminogen activator inhibitor-1 (PAI-1), homocysteine, thromboxane B2 (TXB2), thromboxane A2 (TXA2), 2,3-dinor thromboxane B2, free fatty acids (FFA), serum amyloid A1, serum amyloid A2, serum amyloid A3, serum amyloid A4, thiobarbituric acid (TBA) reacting material, adiponectin (GBP-28), hemoglobin A1c (HbA1c), macrophage colony stimulating factor (M-CSF), granulocyte macrophage colony stimulating factor (GM-CSF), fibrinogen, fibrin D-dimer, platelet derived-microparticles, mean platelet volume (MPV), platelet subpopulations, heart rate,

systolic and diastolic blood pressure, nuclear factor kappa-light-chain enhancer of activated B cells (NF- κ B), adenosine diphosphate induced platelet aggregation, platelet endothelial cell adhesion molecule (PECAM-1), vitronectin receptor ($\alpha_v\beta_3$), and glycoprotein IIb/IIIa (gpIIb/IIIa). This effect is more beneficial than that observed with VASCEPA®. Administration of COMPOSITION 1 has a beneficial impact, or a minimal impact, or no impact, on other non-HDL lipid parameters, such as LDL cholesterol versus baseline and Vascepa. In an alternate study design, this Study #5 is conducted with a certain dose level of COMPOSITION 2 OR COMPOSITION 3 instead of COMPOSITION 1.

[0100] DESCRIPTION OF THE PREFERRED EMBODIMENTS

1. A fatty acid composition comprising at least 50% omega-3-fatty acids, salts or derivatives thereof, while comprising eicosapentaenoic acid (EPA; C20:5-n3) and docosapentaenoic acid (DPA; C22:5-n3) and wherein the EPA:DHA ratio is higher than 20:1.
2. A fatty acid composition comprising at least 60% omega-3-fatty acids, salts or derivatives thereof, while comprising eicosapentaenoic acid (EPA; C20:5-n3) and docosapentaenoic acid (DPA; C22:5-n3) and wherein the EPA:DHA ratio is higher than 20:1.
3. A fatty acid composition comprising at least 70% omega-3-fatty acids, salts or derivatives thereof, while comprising eicosapentaenoic acid (EPA; C20:5-n3) and docosapentaenoic acid (DPA; C22:5-n3) and wherein the EPA:DHA ratio is higher than 20:1.
4. A fatty acid composition comprising at least 75% omega-3-fatty acids, salts or derivatives thereof, while comprising eicosapentaenoic acid (EPA; C20:5-n3) and docosapentaenoic acid (DPA; C22:5-n3) and wherein the EPA:DHA ratio is higher than 20:1.
5. A fatty acid composition comprising at least 80% omega-3-fatty acids, salts or derivatives thereof, while comprising eicosapentaenoic acid (EPA; C20:5-n3) and docosapentaenoic acid (DPA; C22:5-n3) and wherein the EPA:DHA ratio is higher than 20:1.

6. A fatty acid composition comprising at least 85% omega-3-fatty acids, salts or derivatives thereof, while comprising eicosapentaenoic acid (EPA; C20:5-n3) and docosapentaenoic acid (DPA; C22:5-n3) and wherein the EPA:DHA ratio is higher than 20:1.
7. A fatty acid composition comprising at least 90% omega-3-fatty acids, salts or derivatives thereof, while comprising eicosapentaenoic acid (EPA; C20:5-n3) and docosapentaenoic acid (DPA; C22:5-n3) and wherein the EPA:DHA ratio is higher than 20:1.
8. A fatty acid composition comprising at least 95% omega-3-fatty acids, salts or derivatives thereof, while comprising eicosapentaenoic acid (EPA; C20:5-n3) and docosapentaenoic acid (DPA; C22:5-n3) and wherein the EPA:DHA ratio is higher than 20:1.
9. A composition according to one of the preferred embodiments 1 through 8, comprising at least 2% docosapentaenoic acid (DPA; C22:5-n3).
10. A composition according to one of the preferred embodiments 1 through 8, comprising at least 4% docosapentaenoic acid (DPA; C22:5-n3).
11. A composition according to one of the preferred embodiments 1 through 8, comprising at least 5% docosapentaenoic acid (DPA; C22:5-n3).
12. A composition according to one of the preferred embodiments 1 through 8, comprising at least 6% docosapentaenoic acid (DPA; C22:5-n3).
13. A composition according to one of the preferred embodiments 1 through 8, comprising at least 7% docosapentaenoic acid (DPA; C22:5-n3).
14. A composition according to one of the preferred embodiments 1 through 8, comprising at least 8% docosapentaenoic acid (DPA; C22:5-n3).
15. A composition according to one of the preferred embodiments 1 through 8, comprising at least 10% docosapentaenoic acid (DPA; C22:5-n3).
16. A composition according to one of the preferred embodiments 1 through 8, comprising at least 12% docosapentaenoic acid (DPA; C22:5-n3).
17. A composition according to one of the preferred embodiments 1 through 8, comprising at least 15% docosapentaenoic acid (DPA; C22:5-n3).
18. A composition according to one of the preferred embodiments 1 through 17, comprising no more than 95% EPA.
19. A composition according to one of the preferred embodiments 1 through 17, comprising no more than 10% omega-6 fatty acids.

20. A composition according to one of the preferred embodiments 1 through 17, comprising no more than 7% omega-6 fatty acids.
21. A composition according to one of the preferred embodiments 1 through 17, comprising no more than 5% omega-6 fatty acids.
22. A composition according to one of the preferred embodiments 1 through 17, comprising no more than 3% omega-6 fatty acids.
23. A composition according to one of the preferred embodiments 1 through 22, comprising no more than 5% arachidonic acid (C22:4-n6).
24. A composition according to one of the preferred embodiments 1 through 22, comprising no more than 4% arachidonic acid (C22:4-n6).
25. A composition according to one of the preferred embodiments 1 through 22, comprising no more than 3% arachidonic acid (C22:4-n6).
26. A composition according to one of the preferred embodiments 1 through 22, comprising no more than 2% arachidonic acid (C22:4-n6).
27. A composition according to one of the preferred embodiments 1 through 22, comprising no more than 1% arachidonic acid (C22:4-n6).
28. A composition according to one of the preferred embodiments 1 through 27, also comprising heneicosapentaenoic acid (C21:5-n3).
29. A composition according to one of the preferred embodiments 1 through 27, comprising at least 0.01% heneicosapentaenoic acid (C21:5-n3).
30. A composition according to one of the preferred embodiments 1 through 27, comprising at least 0.1% heneicosapentaenoic acid (C21:5-n3).
31. A composition according to one of the preferred embodiments 1 through 27, comprising at least 0.3% heneicosapentaenoic acid (C21:5-n3).
32. A composition according to one of the preferred embodiments 1 through 27, comprising at least 0.5% heneicosapentaenoic acid (C21:5-n3).
33. A composition according to one of the preferred embodiments 1 through 27, comprising at least 1% heneicosapentaenoic acid (C21:5-n3).
34. A composition according to one of the preferred embodiments 1 through 27, comprising at least 2% heneicosapentaenoic acid (C21:5-n3).
35. A composition according to one of the preferred embodiments 1 through 27, comprising at least 3% heneicosapentaenoic acid (C21:5-n3).
36. A composition according to one of the preferred embodiments 1 through 27, comprising at least 4% heneicosapentaenoic acid (C21:5-n3).

37. A composition according to one of the preferred embodiments 1 through 27, comprising at least 5% heneicosapentaenoic acid (C_{21:5}-n₃).
38. A composition according to one of the preferred embodiments 1 through 37, comprising no more than 5% omega-3 fatty acids that are not omega-3-pentaenoic acids.
39. A composition according to one of the preferred embodiments 1 through 37, comprising no more than 4% omega-3 fatty acids that are not omega-3-pentaenoic acids.
40. A composition according to one of the preferred embodiments 1 through 37, comprising no more than 3% omega-3 fatty acids that are not omega-3-pentaenoic acids.
41. A composition according to one of the preferred embodiments 1 through 37, comprising no more than 2% omega-3 fatty acids that are not omega-3-pentaenoic acids.
42. A composition according to one of the preferred embodiments 1 through 37, comprising no more than 1.5% omega-3 fatty acids that are not omega-3-pentaenoic acids.
43. A composition according to one of the preferred embodiments 1 through 37, comprising no more than 1.25% omega-3 fatty acids that are not omega-3-pentaenoic acids.
44. A composition according to one of the preferred embodiments 1 through 37, comprising no more than 1% omega-3 fatty acids that are not omega-3-pentaenoic acids.
45. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 99:1 and 1:99.
46. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 60:1 and 1:60.
47. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 50:1 and 1:10.
48. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 40:1 and 1:3.
49. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 40:1 and 1:2.

50. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 40:1 and 1:1.
51. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 30:1 and 1:1.
52. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 20:1 and 1:1.
53. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 10:1 and 1:1.
54. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 5:1 and 1:1.
55. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 10:1 and 2:1.
56. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 20:1 and 2:1.
57. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 30:1 and 2:1.
58. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 40:1 and 2:1.
59. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 50:1 and 2:1.
60. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 10:1 and 3:1.
61. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 20:1 and 3:1.
62. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 30:1 and 3:1.
63. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 40:1 and 3:1.
64. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 50:1 and 3:1.
65. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 60:1 and 3:1.
66. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 10:1 and 5:1.

67. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 20:1 and 5:1.
68. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 30:1 and 5:1.
69. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 40:1 and 5:1.
70. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 50:1 and 5:1.
71. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 60:1 and 5:1.
72. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 20:1 and 10:1.
73. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 30:1 and 10:1.
74. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 40:1 and 10:1.
75. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 50:1 and 10:1.
76. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 60:1 and 10:1.
77. A composition according to one of the preferred embodiments 1 through 44, wherein the EPA:DPA ratio is between 100:1 and 10:1.
78. A composition according to one of the preferred embodiments 1 through 44, comprising between 55% and 95% EPA.
79. A composition according to one of the preferred embodiments 1 through 44, comprising between 60% and 95% EPA.
80. A composition according to one of the preferred embodiments 1 through 44, comprising between 65% and 95% EPA.
81. A composition according to one of the preferred embodiments 1 through 44, comprising between 70% and 95% EPA.
82. A composition according to one of the preferred embodiments 1 through 44, comprising between 75% and 95% EPA.
83. A composition according to one of the preferred embodiments 1 through 44, comprising between 80% and 95% EPA.

84. A composition according to one of the preferred embodiments 1 through 44, comprising between 85% and 95% EPA.
85. A composition according to one of the preferred embodiments 1 through 44, comprising between 90% and 95% EPA.
86. A composition according to one of the preferred embodiments 1 through 44, comprising between 1% and 3% DPA.
87. A composition according to one of the preferred embodiments 1 through 44, comprising between 1% and 5% DPA.
88. A composition according to one of the preferred embodiments 1 through 44, comprising between 2% and 10% DPA.
89. A composition according to one of the preferred embodiments 1 through 44, comprising between 3% and 20% DPA.
90. A composition according to one of the preferred embodiments 1 through 44, comprising between 3% and 30% DPA.
91. A composition according to one of the preferred embodiments 1 through 44, comprising between 3% and 50% DPA.
92. A composition according to one of the preferred embodiments 1 through 44, comprising between 3% and 75% DPA.
93. A composition according to one of the preferred embodiments 1 through 44, comprising between 3% and 90% DPA.
94. A fatty acid composition according to one of the preferred embodiments 1 through 93, in which the fatty acids are present as ethyl esters.
95. A fatty acid composition according to one of the preferred embodiments 1 through 93, in which the fatty acids are present as free fatty acids.
96. A fatty acid composition according to one of the preferred embodiments 1 through 93, in which the fatty acids are present as esters in di-glyceride form.
97. A fatty acid composition according to one of the preferred embodiments 1 through 93, in which the fatty acids are present as esters in triglyceride form.
98. A fatty acid composition according to one of the preferred embodiments 94 through 97, also comprising a suitable anti-oxidant in a concentration sufficient to protect the fatty acids of the composition from oxidation.
99. A pharmaceutically suitable formulation comprising one of the compositions according to preferred embodiments 94 through 98, in which the amount of

eicosapentaenoic acid plus docosapentaenoic acid is present in an amount between 100 and 10,000 mg.

100. A pharmaceutically suitable formulation or dosage form comprising one of the compositions according to preferred embodiments 94 through 98, in which the amount of eicosapentaenoic acid plus docosapentaenoic acid is present in an amount between 250 and 1,250 mg.
101. A pharmaceutically suitable formulation or dosage form comprising one of the compositions according to preferred embodiments 94 through 98, in which the amount of eicosapentaenoic acid plus docosapentaenoic acid is present in an amount between 500 and 1,100 mg.
102. A pharmaceutically suitable formulation or dosage form comprising one of the compositions according to preferred embodiments 94 through 98, in which the amount of eicosapentaenoic acid plus docosapentaenoic acid is present in an amount between 100 and 10,000 mg.
103. A method of administration or treatment to a subject of a formulation or dosage form according to one of the preferred embodiments 94 through 102 at a daily dose between 100 and 10,000 mg.
104. A method of administration or treatment to a subject of a formulation or dosage form according to one of the preferred embodiments 94 through 102 at a daily dose between 500 and 5,000 mg.
105. A method of administration or treatment to a subject of a formulation or dosage form according to one of the preferred embodiments 94 through 102 at a daily dose between 1,500 and 4,100 mg.
106. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with very high triglycerides (equal to or more than 500 mg/dL).
107. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with high triglycerides (equal to or more than 200 mg/dL but less than 500 mg/dL).
108. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient already undergoing treatment with a statin and then diagnosed with high triglycerides (equal to or more than 200 mg/dL but less than 500 mg/dL).

109. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with mixed dyslipidemia with TG 200-499 mg/dL and LDL-cholesterol equal to or more than 190 mg/dL.
110. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with mixed dyslipidemia with TG 300-700 mg/dL and LDL-cholesterol equal to or more than 190 mg/dL.
111. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with mixed dyslipidemia with TG 200-499 mg/dL and non-HDL-cholesterol equal to or more than 200 mg/dL.
112. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with mixed dyslipidemia with TG 300-700 mg/dL and non-HDL-cholesterol equal to or more than 200 mg/dL.
113. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with mixed dyslipidemia with TG 200-499 mg/dL and LDL-cholesterol equal to or more than 160 mg/dL.
114. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with mixed dyslipidemia with TG 300-700 mg/dL and LDL-cholesterol equal to or more than 160 mg/dL.
115. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with mixed dyslipidemia with TG 200-499 mg/dL and non-HDL-cholesterol equal to or more than 160 mg/dL.
116. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with mixed dyslipidemia with TG 300-700 mg/dL and non-HDL-cholesterol equal to or more than 160 mg/dL.
117. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with mixed dyslipidemia with TG 200-499 mg/dL and LDL-cholesterol equal to or more than 130 mg/dL.
118. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with mixed dyslipidemia with TG 300-700 mg/dL and LDL-cholesterol equal to or more than 130 mg/dL.
119. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with mixed dyslipidemia with TG 200-499 mg/dL and non-HDL-cholesterol equal to or more than 130 mg/dL.

120. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with mixed dyslipidemia with TG 300-700 mg/dL and non-HDL-cholesterol equal to or more than 130 mg/dL.
121. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed/assessed to be at substantially elevated risk for cardiovascular events.
122. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with diabetes.
123. A method of treatment according to preferred embodiments 103 through 105, in which the subject is a patient diagnosed with pre-diabetes or metabolic syndrome.
124. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant reduction of blood, serum or plasma triglyceride levels.
125. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant reduction of blood, serum or plasma triglyceride levels while not significantly increasing blood, serum or plasma LDL-cholesterol levels.
126. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant reduction of blood, serum or plasma total-cholesterol levels.
127. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant reduction of blood, serum or plasma non-HDL-cholesterol levels.
128. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant reduction of blood, serum or plasma LDL-cholesterol levels.
129. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant reduction of blood, serum or plasma VLDL-cholesterol levels.
130. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant reduction of blood, serum or plasma VLDL-cholesterol levels while not significantly increasing blood, serum or plasma LDL-cholesterol levels.

131. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant reduction of blood, serum or plasma apo-B levels.
132. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant reduction of blood, serum or plasma apo-C-III levels.
133. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant reduction of blood, serum or plasma LP-PLA2 levels.
134. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant reduction of blood, serum or plasma hs-CRP levels.
135. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant increase of blood, serum or plasma HDL-cholesterol levels.
136. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant increase of blood, serum or plasma apo-A levels.
137. A method of treatment according to one of the preferred embodiments 103 through 123, in which the treatment results in significant reduction of the risk of suffering certain cardiovascular events.
138. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in a Tmax of 8 hours or less
139. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in a Tmax of 6 hours or less
140. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in a Tmax of 5 hours or less
141. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in a Tmax of 4 hours or less

142. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in a Tmax of 8 hours or less
143. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in a Tmax of 6 hours or less
144. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in a Tmax of 5 hours or less
145. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in a Tmax of 4 hours or less
146. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Tmax less than the EPA+DPA+DHA Tmax for LOVAZA® under the same administration conditions
147. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Tmax less than the EPA+DPA+DHA Tmax for LOVAZA® under the same administration conditions
148. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Tmax less than the EPA+DPA Tmax for AMR101 under the same administration conditions
149. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Tmax less than the EPA+DPA Tmax for AMR101 under the same administration conditions
150. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an Total Omega-3 FA Tmax less than the Total Omega-3 FA Tmax for LOVAZA® under the same administration conditions
151. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an Total Omega-

- 3 FA Tmax less than the Total Omega-3 FA Tmax for LOVAZA® under the same administration conditions
152. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an Total Omega-3 FA Tmax less than the Total Omega-3 FA Tmax for AMR101 under the same administration conditions
153. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an Total Omega-3 FA Tmax less than the Total Omega-3 FA Tmax for AMR101 under the same administration conditions
154. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 110% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions
155. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 120% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions
156. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 130% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions
157. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 140% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions
158. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 150% of the

EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions

159. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 150% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions
160. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 200% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions
161. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 300% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions
162. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 400% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions
163. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 500% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions
164. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 600% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions
165. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an

EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 150% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions

166. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 200% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions

167. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 300% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions

168. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 400% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions

169. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 500% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions

170. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 600% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions

171. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 700% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for LOVAZA® under the same administration conditions

172. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 110% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
173. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 120% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
174. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 130% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
175. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 140% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
176. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 150% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
177. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 150% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
178. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 200% of the

EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions

179. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 300% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
180. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 400% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
181. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 500% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
182. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 600% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
183. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 150% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
184. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 200% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
185. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an

EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 300% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions

186. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 400% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
187. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 500% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
188. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 600% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
189. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 700% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for LOVAZA® under the same administration conditions
190. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 110% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
191. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 120% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions

192. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 130% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
193. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 140% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
194. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 150% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
195. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 150% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
196. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 200% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
197. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 300% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
198. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 400% of the EPA+DPA

Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions

199. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 500% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
200. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 600% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
201. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 150% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
202. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 200% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
203. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 300% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
204. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 400% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
205. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 500% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
206. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA

- Cmax or Total Omega-3 FA Cmax of at least 600% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
207. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA Cmax or Total Omega-3 FA Cmax of at least 700% of the EPA+DPA Cmax or Total Omega-3 FA Cmax for AMR101 under the same administration conditions
208. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 110% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
209. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 120% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
210. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 130% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
211. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 140% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
212. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 150% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
213. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 150% of the EPA+DPA

AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions

214. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 200% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
215. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 300% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
216. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 400% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
217. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 500% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
218. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 600% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
219. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 150% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
220. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA

AUC or Total Omega-3 FA AUC of at least 200% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions

221. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 300% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
222. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 400% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
223. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 500% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
224. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 600% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
225. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA AUC or Total Omega-3 FA AUC of at least 700% of the EPA+DPA AUC or Total Omega-3 FA AUC for AMR101 under the same administration conditions
226. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 100% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions
227. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 105% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions
228. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an

EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 110% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions

229. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 120% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions

230. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 130% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions

231. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 140% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions

232. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 100% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions

233. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 105% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions

234. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 110% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions

235. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 120% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions
236. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 130% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions
237. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 140% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions
238. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 100% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions
239. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 105% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions
240. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 110% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions
241. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 120% of the

EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions

242. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 130% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions

243. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 140% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions

244. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax of at least 150% of the EPA+DPA+DHA Cmax or Total Omega-3 FA Cmax for EPANOVA™ under the same administration conditions

245. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 100% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions

246. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 105% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions

247. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 110% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions

248. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an

EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 120% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions

249. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 130% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions
250. A method of administration or treatment under high fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 140% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions
251. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 100% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions
252. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 105% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions
253. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 110% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions
254. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 120% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions

255. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 130% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions
256. A method of administration or treatment under low fat meal conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 140% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions
257. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 100% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions
258. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 105% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions
259. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 110% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions
260. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 120% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions
261. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 130% of the

EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions

262. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 140% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions

263. A method of administration or treatment under fasting conditions according to one of the preferred embodiments 103 through 105 resulting in an EPA+DPA+DHA AUC or Total Omega-3 FA AUC of at least 150% of the EPA+DPA+DHA AUC or Total Omega-3 FA AUC for EPANOVA™ under the same administration conditions.

264. The composition of claim 1, wherein the ratio of EPA to DPA (EPA:DPA) is between 15:1 to 8:1.

265. An orally administrable composition comprising fatty acids, wherein at least 50% by weight of the fatty acids comprise omega-3-pentaenoic acids, salts, esters, or derivatives thereof, wherein the composition comprises eicosapentaenoic acid (EPA), docosapentaenoic acid (DPA), and docosahexaenoic acid (DHA), and wherein the ratio of DHA to EPA (DHA:EPA) is less than 1:20, and wherein the ratio of DHA to DPA (DHA:DPA) is less than 2:1.

WHAT IS CLAIMED:

Claim 1. An orally administrable composition comprising fatty acids, wherein at least 50% by weight of the fatty acids comprise omega-3 fatty acids, salts, esters, or derivatives thereof, wherein the omega-3 fatty acids comprise eicosapentaenoic acid (EPA), docosapentaenoic acid (DPA), and docosahexaenoic acid (DHA), and wherein the ratio of DHA to EPA (DHA:EPA) is less than 1:20, and wherein the ratio of DHA to DPA (DHA:DPA) is less than 2:1.

Claim 2. The orally administrable composition of claim 1, wherein the ratio of DHA:EPA is less than 1:50.

Claim 3. The orally administrable composition of claim 1, wherein the ratio of DHA:EPA is less than 1:99.

Claim 4. The orally administrable composition of claim 1, wherein the ratio of DHA:DPA is less than 1:1.

Claim 5. The orally administrable composition of claim 1, wherein the ratio of DHA:DPA is less than 1:5.

Claim 6. The orally administrable composition of claim 1, wherein the ratio of EPA to DPA (EPA:DPA) is between 30:1 and 1:1.

Claim 7. The orally administrable composition of claim 1, wherein the ratio of EPA to DPA (EPA:DPA) is between 20:1 to 5:1.

Claim 8. The orally administrable composition of claim 1, wherein the composition comprises EPA in an amount between 55% and 95% of the total amount of fatty acids.

Claim 9. The orally administrable composition of claim 1, wherein the composition comprises EPA in an amount between 65% and 90% of the total amount of fatty acids.

WO 2013/192109

PCT/US2013/046176

Claim 10. The orally administrable composition of claim 1, wherein the composition comprises DPA in an amount between 1% and 50% of the total amount of fatty acids.

Claim 11. The orally administrable composition of claim 1, wherein the composition comprises HPA in an amount between 1% and 20% of the total amount of fatty acids.

Claim 12. The orally administrable composition of claim 1, wherein the composition comprises DPA in an amount between 1% and 50% and comprises HPA in an amount between 1% and 20% of the total amount of fatty acids.

Claim 13. The orally administrable composition of claim 1, wherein the composition comprises DPA in an amount between 2% and 40% of the total amount of fatty acids.

Claim 14. The orally administrable composition of claim 1, wherein at least 90% by weight of the fatty acids comprise omega-3 fatty acids.

Claim 15. The orally administrable composition of claim 1, wherein the composition further comprises heneicosapentaenoic acid (HPA) in an amount of at least 0.10% of the total amount of fatty acids.

Claim 16. The orally administrable composition of claim 1, wherein the composition further comprises an HMG-CoA reductase inhibitor.

Claim 17. A method of treating a condition in a subject in need thereof, comprising administering the composition of claim 1, wherein the condition is selected from the group consisting of: hypertriglyceridemia; hypercholesterolemia; mixed dyslipidemia; coronary heart disease (CHD); vascular disease; cardiovascular disease; acute coronary syndrome; atherosclerotic disease and related conditions; heart failure; cardiac arrhythmias; coagulatory conditions associated with cardiac arrhythmias; ischemic dementia; vascular dementia; hypertension; coagulation related disorders; nephropathy; kidney or urinary tract disease; retinopathy; cognitive

WO 2013/192109

PCT/US2013/046176

and other CNS disorders; autoimmune diseases; inflammatory diseases; asthma or other respiratory disease; dermatological disease; metabolic syndrome; diabetes, diabetes mellitus or other form of metabolic disease; liver disease; non-alcoholic fatty liver disease; disease of the gastrointestinal tract; disease of the male or female reproductive system or related secondary sexual organs; a cancer of any type, including lymphomas and myelomas; and an infection caused by a virus, bacterium, fungus, protozoa or other organism.

Claim 18. The method of claim 17, comprising further administering an HMG-CoA reductase inhibitor.

Claim 19. A method for treating cardiac events, cardiovascular events, vascular events or reducing symptoms associated with cardiac events, cardiovascular events, vascular events in a subject in need thereof, comprising administering the composition of claim 1 to the subject.

Claim 20. The method of claim 19, comprising further administering an HMG-CoA reductase inhibitor.

Claim 21. A method of reducing cardiac events, cardiovascular events, or vascular events in a subject in need thereof, comprising administering the composition of claim 1 to the subject.

Claim 22. The method of claim 21, comprising further administering an HMG-CoA reductase inhibitor.

Claim 23. A method of reducing triglyceride levels in a subject in need thereof, comprising administering to the subject the composition of claim 1.

Claim 24. The method of claim 23, comprising further administering an HMG-CoA reductase inhibitor.

Claim 25. The method of any one of claims 17-24, wherein the LDL-cholesterol level of the subject is not substantially increased.

Claim 26. The method of any one of claims 17-24, wherein the LDL-cholesterol level of the subject is reduced.

Claim 27. The method of any one of claims 17-24, wherein the triglyceride level of the subject is 500 mg/dL or higher.

Claim 28. The method of any one of claims 17-24, wherein the triglyceride level of the subject is between 200 mg/dL and 499 mg/dL.

Claim 29. The composition of claim 1, wherein the composition comprises an amount of EPA and DPA of between 500 and 1,100 mg.

Claim 30. A method of reducing triglyceride levels under fasting conditions comprising administering to the subject the composition of claim 1, resulting in an AUC of EPA+DPA of at least 150% of the AUC of EPA+DPA for AMR101 under the same administration conditions.

Claim 31. A method of reducing triglyceride levels under fasting conditions comprising administering to the subject the composition of claim 1, resulting in an Cmax of EPA+DPA of at least 150% of the Cmax of EPA+DPA for AMR101 under the same administration conditions.