



(86) Date de dépôt PCT/PCT Filing Date: 1994/09/14
 (87) Date publication PCT/PCT Publication Date: 1995/03/23
 (45) Date de délivrance/Issue Date: 2011/11/15
 (85) Entrée phase nationale/National Entry: 1996/03/13
 (86) N° demande PCT/PCT Application No.: GB 1994/002006
 (87) N° publication PCT/PCT Publication No.: 1995/007985
 (30) Priorités/Priorities: 1993/09/14 (GB9318984.3);
 1994/04/29 (GB9408602.2)

(51) Cl.Int./Int.Cl. *C12N 15/19* (2006.01),
A61K 38/19 (2006.01), *A61K 39/395* (2006.01),
C07K 14/52 (2006.01), *C07K 14/715* (2006.01),
C07K 16/24 (2006.01), *C07K 16/28* (2006.01),
C12P 21/02 (2006.01), *G01N 33/15* (2006.01),
G01N 33/48 (2006.01), *G01N 33/566* (2006.01)

(72) Inventeurs/Inventors:
 WILLIAMS, TIMOTHY JOHN, GB;
 JOSE, PETER JOHN, GB;
 GRIFFITHS-JOHNSON, DAVID A., GB;
 HSUAN, JOHN JUSTIN, GB

(73) Propriétaire/Owner:
 IMPERIAL INNOVATIONS LIMITED, GB

(74) Agent: RIDOUT & MAYBEE LLP

(54) Titre : CYTOKINE CHIMIOTACTIQUE APPELEE EOTAXINE, POUR EOSINOPHILES
 (54) Title: EOTAXIN: EOSINOPHIL CHEMOTACTIC CYTOKINE

(57) **Abrégé/Abstract:**

A chemoattractant protein called "eotaxin" is capable of attracting eosinophils and of inducing eosinophil accumulation and/or activation in vitro and in vivo. Various types of agents that inhibit or otherwise hinder the production, release or activity of eotaxin may be used therapeutically in the treatment of asthma and other inflammatory diseases.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :
C12N 15/19, C07K 14/52, C12P 21/02,
C07K 14/715, 16/28, A61K 38/19, 38/17,
G01N 33/68, A61K 39/395

A1

(11) International Publication Number: **WO 95/07985**

(43) International Publication Date: 23 March 1995 (23.03.95)

(21) International Application Number: PCT/GB94/02006

(22) International Filing Date: 14 September 1994 (14.09.94)

(30) Priority Data:
9318984.3 14 September 1993 (14.09.93) GB
9408602.2 29 April 1994 (29.04.94) GB

(71) Applicants (for all designated States except US): NATIONAL HEART & LUNG INSTITUTE [GB/GB]; Dovehouse Street, London SW3 6LY (GB). LUDWIG INSTITUTE FOR CANCER RESEARCH [GB/GB]; Hedges House, 153-155 Regent Street, London W1R 7FD (GB).

(72) Inventors; and
(75) Inventors/Applicants (for US only): WILLIAMS, Timothy, John [GB/GB]; National Heart & Lung Institute, Dovehouse Street, London SW3 6LY (GB). JOSE, Peter, John [GB/GB]; National Heart & Lung Institute, Dovehouse Street, London SW3 6LY (GB). GRIFFITHS-JOHNSON, David, A. [GB/GB]; National Heart & Lung Institute, Dovehouse Street, London SW3 6LY (GB). HSUAN, John, Justin [GB/GB]; Ludwig Institute for Cancer Research, Hedges House, 153-155 Regent Street, London W1R 7FD (GB).

(74) Agents: SILVESTON, Judith et al.; Abel & Imray, Northumberland House, 303-306 High Holborn, London WC1V 7LH (GB).

(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD).

Published

With international search report.



Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

2171761

(54) Title: EOTAXIN: EOSINOPHIL CHEMOTACTIC CYTOKINE

(57) Abstract

A chemoattractant protein called "eotaxin" is capable of attracting eosinophils and of inducing eosinophil accumulation and/or activation in vitro and in vivo. Various types of agents that inhibit or otherwise hinder the production, release or activity of eotaxin may be used therapeutically in the treatment of asthma and other inflammatory diseases.



EOTAXIN : EOSINOPHIL CHEMOTACTIC CYTOKINE

The present invention relates to a chemotactic cytokine.

5 The accumulation of eosinophil leukocytes is a characteristic feature of IgE-mediated allergic reactions such as allergic asthma, rhinitis and eczema. Eosinophil accumulation also occurs in non-allergic asthma. The immediate broncho-
 10 constriction in response to a provoking stimulus in the asthmatic involves mast cell activation and the release of constrictor mediators. This is followed after several hours in some individuals by a late bronchoconstrictor response associated with a massive influx of eosinophils (1). Repeated provocation results in chronic inflammation in the airways and
 15 a marked hyper-responsiveness to constrictor mediators. The magnitude of both the late response and the chronic hyper-responsiveness correlates with the numbers of eosinophils present in the lung (2,3).

20 The present invention provides a chemoattractant protein capable of attracting eosinophils and of inducing eosinophil accumulation and/or activation in vitro and in vivo. The chemoattractant protein of the present invention is designated "eotaxin".

25

Eotaxins are proteins of the C-C branch of the platelet factor 4 superfamily of chemotactic cytokines. Within the C-C branch of the platelet factor 4 superfamily of chemotactic cytokines, or chemokines, certain members have the property of attracting
 30 eosinophils in vitro and some may induce eosinophil accumulation in vivo. For example, the chemokines RANTES and MIP-1 α attract eosinophils in vitro while MCP-1 and MIP-1 β do not. ("RANTES" denotes Regulated upon Activation in Normal T cells Expressed and Secreted, "MIP" denotes Macrophage
 35 Inflammatory Protein, and "MCP" denotes Monocyte Chemo-attractant Protein.)

Naturally-occurring cytokines within the platelet factor 4 superfamily of chemotactic cytokines may have marked inter-species variations in

the amino acid sequence of the protein, and in the carbohydrate modifications of the protein, while retaining the same characteristic functional properties. Similar variations in structure may occur in cytokines obtained from different individuals within the same species. Many chemokines within the C-C branch of the platelet factor 4 superfamily show promiscuity of receptor binding, and the ability of different chemokines to bind to the same receptor is not necessarily dependent on a high degree of homology at the amino acid level. Accordingly, both interspecies and intraspecies variations in protein length, amino acid sequence and carbohydrate modifications are generally to be expected for eotaxins.

The ability to attract eosinophils and to induce eosinophil accumulation and/or activation in vitro and in vivo is a characteristic property of eotaxins. Furthermore, eotaxins generally show substantially no attractive effect for neutrophils in vivo. The eosinophil chemoattractant effect may be an inter-species effect, for example, guinea-pig eotaxin appears to be potent in inducing chemotaxis of human eosinophils in vitro.

An eotaxin may be obtained from an appropriate body fluid, for example, from bronchoalveolar lavage fluid obtained from a human or non-human subject, particularly an allergic subject after an allergen challenge, either experimentally induced or naturally incurred. Other sources of eotaxins are, for example, inflammatory exudate fluids and in vitro cultures of macrophages, lymphocytes, neutrophils, mast cells, airway epithelial cells, connective tissue cells, vascular endothelial cells and eosinophils themselves.

For example, an eotaxin may be obtained from a sensitised guinea-pig after allergen challenge. Guinea-pig models are useful as they share many common features with the asthmatic response in man. Eotaxin obtainable from bronchoalveolar

2171761

lavage fluid of a sensitised guinea-pig by sequential HPLC purification generally has a molecular weight in the range of from 6-16 kDa. (As indicated above, intraspecies molecular weight variations of this order of magnitude are observed in 5 members of the platelet factor 4 superfamily.)

The amino acid sequence of a guinea-pig eotaxin is set out in SEQ.ID. NO. 1, SEQ.ID. NO. 2 and in Figures 7 and 8 of the accompanying drawings. Other guinea-pig eotaxins will 10 generally have at least 50% overall homology with the sequence shown in SEQ.ID. NO. 1 (Figure 7) at the amino acid level. The homology may be at least 60%, for example at least 70%, for example at least 80% with the sequence set out in SEQ.ID. NO. 1 and in Figure 7.

15

Percentage homology in the present case is calculated on the basis of amino acids that are identical in corresponding positions in the two sequences under investigation.

Conservative substitutions are not taken into account. In the 20 calculation of percentage homology of a putative eotaxin molecule under investigation with the sequence shown in SEQ.ID. NO. 1 (Figure 7) or with SEQ.ID. NO. 2 (Figure 8) if the molecule under investigation has a different length from the eotaxin set out in SEQ.ID. NO. 1 or SEQ.ID. NO. 2 (Figure 25 7 or Figure 8), then the calculation is based on the amino acids in the portion of the molecule under investigation that overlaps with the sequence shown in SEQ.ID. NO. 1 (Figure 7) or SEQ.ID. NO. 2 (Figure 8). Software packages for the alignment of amino acid sequences and the calculation of 30 homology are available commercially, for example, the "Bestfit" program available from Genetics Computer Group Sequence Analysis Software, Madison, Wisconsin, U.S.A..

Unless specified otherwise, the specific values of percentage 35 homology between eotaxin and other chemotactic cytokines given in the present specification have been calculated on the basis of the eotaxin set out in SEQ.ID. NO. 1 (Figure 7).

2171761

As indicated above, eotaxins obtainable from species other than guinea-pigs, for example humans, will exhibit inter-species differences of the type demonstrated by other members of the C-C branch of the platelet factor 4 superfamily of chemokines, for example, differences in protein length, amino acid sequence and carbohydrate modifications. There may, for example, be variations in the C- and/or N-terminal residues. For example, it is expected that the molecular weight of an eotaxin from a species other than guinea-pig will generally fall within the range of from 6 kDa to 16 kDa, but in some cases an eotaxin may have a molecular weight less than 6 kDa or more than 16 kDa.

Similarly, it is expected that in general an eotaxin from a species other than guinea-pig will have at least 40% overall homology with the sequence set out in SEQ.ID. NO. 1 and in Figure 7. The homology may be at least 50%, for example at least 60%, for example at least 70%, for example at least 80% with the sequence set out in SEQ.ID. NO. 1 and in Figure 7. There may, however, be eotaxins from species other than guinea-pigs that have less than 40% homology with SEQ.ID. NO. 1 (Figure 7).

Eotaxins may be identified by any one or more of the characteristics set out above, in particular by their ability to attract and/or active eosinophils in vitro and cause their accumulation and/or activation in vivo. A characteristic that assists the identification of a molecule as an eotaxin is the lack of attractive effect on neutrophils.

30

The present invention provides a method of determining the ability of a substance to induce eosinophil accumulation and/or activation in vivo, that is to say, a method for testing putative eotaxins, which comprises administering the substance, generally intradermally, to a test animal previously treated with labelled, for example ¹¹¹In-labelled, eosinophils and subsequently determining the number of

labelled eosinophils at a skin site.

One in vitro method that may be used to test a putative eotaxin for the ability to attract and/or activate eosinophils in vitro is the ability of the substance to increase eosinophil intracellular calcium levels. Other general methods for determining chemotactic activity in vitro may be used to test putative eotaxins in vitro.

10 Confirmation that an eosinophil attractant is an eotaxin may also be made by consideration of sequence homology of that protein with the sequence set out in SEQ.ID. NO. 1 (Figure 7) and/or with the sequence set in SEQ.ID. NO. 2 (Figure 8) and/or by consideration of the structural relationship between
15 the protein and the guinea-pig eotaxin.

As mentioned above, RANTES and MIP-1 α are both eosinophil activators. Eotaxin has functional similarities but low structural homology with RANTES and MIP-1 α (31% homology with
20 MIP-1 α at the amino acid level calculated on the basis of SEQ.ID. No. 1 (Figure 7); 32% homology when calculated on the basis of the overlapping sequences and 26% homology with RANTES at the amino acid level calculated on the basis of
25 SEQ.ID. No. 1 (Figure 7); 27% homology when calculated on the basis of the overlapping sequences). An eotaxin can be distinguished from RANTES and MIP-1 α not only by the degree of homology but also by the overall differences in sequence and structure.

30 In addition to full-length eotaxin molecules, the present invention also provides molecules that comprise less than a full length eotaxin sequence. Such molecules (called "fragments" herein) may be polypeptides or peptides. For use as an eotaxin substitute, a fragment should retain one or
35 more of the biological activities of the parent molecule.

Eosinophils contain an armoury of chemicals necessary for

killing parasites. These chemicals have been implicated in the damage to airway epithelium that occurs in asthma and may relate to the observed changes in airway function (26,27). From our studies we suggest that eotaxins should be considered
5 as important mediators of eosinophil accumulation in vivo. Macrophages, lymphocytes, neutrophils, mast cells, airway epithelial cells, connective tissue cells, vascular endothelial cells and eosinophils themselves are likely candidates as the source of the eosinophil chemoattractant
10 activity generated in the lung. Platelets may also have a role as it has been shown that they can release C-C chemokines (22). Further, an early platelet deposition may be involved in the subsequent eosinophil accumulation in vivo (28,29) and there is evidence that platelet-activating factor induces the
15 synthesis of an unidentified eosinophil chemoattractant in vivo (30). In this respect, it is of interest that platelet-derived growth factor can induce gene expression of C-C chemokines in fibroblasts (31). Furthermore, the C-C chemokines have been implicated in wound healing (18). This may be
20 important in the sub-epithelial basement membrane fibrosis that is a prominent feature of the asthmatic lung. Thus, eotaxins may be involved in both eosinophil accumulation and in chronic structural changes in the lung.

25 Eotaxins may have an important role in asthma and in other diseases having an inflammatory component where eosinophil accumulation and/or activation is a prominent feature, for example, rhinitis and eczema, especially allergic eczema. Accordingly, agents that inhibit or otherwise hinder the
30 production, release or action of eotaxins have potential as selective therapeutic agents. Such agents and their therapeutic use are part of the present invention.

Such agents include inhibitors that affect the interaction of
35 an eotaxin with eotaxin receptors, for example, by binding to an eotaxin or to an eotaxin receptor. An example of such an inhibitor is receptors themselves which, on administration,

can bind an eotaxin and prevent its interaction with naturally-occurring receptors. Such inhibitory receptors may be soluble or insoluble. Receptors which are not involved in cell activation may be bound to, or induced on, cells. Such 5 receptors may also be used to remove endogenous eotaxin.

Further examples of agents that affect the interaction of eotaxins with eotaxin receptors are receptor antagonists, and antibodies, both antibodies directed against (capable of 10 binding with) an eotaxin and antibodies directed against an eotaxin receptor, especially monoclonal antibodies. Any other agent that inhibits or otherwise hinders the binding of an eotaxin to an eotaxin receptor also has therapeutic potential, for example, any other agent that binds to an eotaxin or to an 15 eotaxin receptor. Further agents that have therapeutic potential are those that prevent or reduce activation of eotaxin receptors.

Further agents that inhibit or otherwise hinder the action of 20 eotaxins are those that change the structure of an eotaxin such that it is no longer able to bind to an eotaxin receptor, for example, an enzyme or other agent that degrades eotaxin specifically.

25 Receptor promiscuity is common among chemokines, so although it is essential that a receptor is capable of binding an eotaxin, the receptor need not necessarily be eotaxin-specific. For example, a receptor may bind MIP-1 α , RANTES and/or other eosinophil attractant chemokines as well as an 30 eotaxin.

As indicated above, possibilities for therapeutic intervention include the use of a receptor to which an eotaxin binds, especially a soluble receptor. It may be advantageous to use 35 an eotaxin-specific receptor. Further possibilities for therapeutic intervention include receptor antagonists, for example, based on 3-dimensional structures or the amino acid

217/1961

sequences of eotaxins and/or of eotaxin receptors, and agents found to inhibit eotaxin or other agonists binding to or activating eotaxin receptors. For example, a receptor antagonist or an agonist inhibitor may be a polypeptide in which the sequence of a full-length naturally-occurring eotaxin has been modified, for example, by amino acid substitution, or may be a fragment of an eotaxin (that is to say, a polypeptide or small peptide comprising part of the amino acid sequence of a naturally-occurring eotaxin), or a modified fragment of an eotaxin, for example, modified by amino acid substitution.

Furthermore, knowledge of the sequence and/or structure of eotaxins either alone or in combination with knowledge of the sequence and/or structure of other chemokines that bind to the same receptor(s) as eotaxins, provides useful information for the design of therapeutic agents.

Agents that prevent or inhibit eotaxin synthesis or release may also be used therapeutically. Such agents and their use are also part of the present invention.

All inhibitors of eotaxin activity, synthesis and release, including soluble receptors, antibodies, antagonists and inhibitors of agonist binding, and their use are part of the present invention.

The present invention accordingly provides an agent that inhibits or otherwise hinders the production, release or action of an eotaxin, especially an agent as described above, for use as a medicament. The invention also provides the use of an agent that inhibits or otherwise hinders the production, release or action of an eotaxin, especially an agent as described above, in the manufacture of a medicament for the treatment of asthma or another disease having an inflammatory component, particularly with accumulation of eosinophils, for example, rhinitis or eczema, especially allergic eczema.

The use of the structural and sequence information relating to eotaxins in the design of therapeutically and diagnostically useful agents, for example, in computer-aided design based on the three dimensional structure of eotaxins is part of the present invention.

Putative inhibitors of eotaxin activity may be screened using in vivo and in vitro assays based on inhibition of chemoattraction and/or accumulation and/or activation of eosinophils by eotaxins. Some general methods for testing the activity of a compound for an inhibitory effect on the activity of a chemoattractant cytokine in vitro are known. Such assays may be used to determine the inhibitory action of a putative inhibitor on in vitro effects induced in eosinophils by eotaxins.

Assays that are suitable for screening putative eotaxin inhibitors include, for example, inhibition in vitro of elevation of intracellular calcium levels induced in cells by eotaxin. The method of the present invention for determining the ability of a substance to induce eosinophil accumulation and/or activation in vivo, that is to say, a method for testing putative eotaxins, may also be used to determine the ability of a substance to inhibit eosinophil accumulation and/or activation induced in vivo by an eotaxin: an animal is pretreated with labelled eosinophils, an eotaxin and a putative inhibitor are administered, and the number of labelled eosinophils at a skin site are determined subsequently. The eotaxin is generally administered intradermally, and the putative inhibitor may be administered by the same route or by a different route, for example systemically.

Examples of in vitro and in vivo assays both for the determination of eotaxin activity and for the determination of eotaxin inhibitory activity are described herein. For example, Example 1 gives a detailed protocol for the in vivo

2171761

assay of the present invention, and Example 4 gives detailed protocols of various assays. The assays described herein may be used as such, or may be modified as required. Assays may be used alone or in combination to establish eotaxin and
5 eotaxin-inhibitory activity. A putative inhibitors may be any of the types of molecules described above, including receptors, for example, soluble receptors, antibodies, and antagonists and inhibitors of agonist binding. Methods for testing putative inhibitors of eotaxins are also part of the
10 present invention.

A further aspect of the present invention is a pharmaceutical preparation comprising, as active ingredient, an agent that inhibits or otherwise hinders the production, release or
15 action of an eotaxin, in admixture or in conjunction with a pharmaceutically suitable carrier. Such agents are described above and include, for example, an inhibitor of eotaxin synthesis or release, a soluble eotaxin receptor, an eotaxin receptor antagonist or an inhibitor of an eotaxin receptor
20 agonist, an antibody against eotaxin or an antibody against an eotaxin receptor.

The invention further provides a method of treating asthma and other inflammatory diseases, comprising the administration of
25 an effective amount of an agent that inhibits or otherwise hinders the production, release or action of an eotaxin. The agent may be as described above, for example, an inhibitor of eotaxin synthesis or release, a soluble eotaxin receptor, an eotaxin receptor antagonist or an inhibitor of an eotaxin
30 receptor agonist, or an antibody against eotaxin or against an eotaxin receptor.

The present invention also provides assays for eotaxins and for anti-eotaxin antibodies, especially immunoassays and in
35 particular ELISAs (enzyme-linked immunosorbent assays). The invention provides, for example, an immunoassay for an antigen, characterised in that the antigen is an eotaxin, and

2171761

also provides an immunoassay for an antibody, characterised in that the antibody is an anti-eotaxin antibody. The invention also provides assays for eotaxins that are analogous to immunoassays for eotaxins but that use a specific-binding partner other than an antibody. In such specific-binding partner assays an eotaxin receptor may be used instead of an anti-eotaxin antibody.

In an immunoassay, an anti-eotaxin antibody may, for example, be coated on a solid surface to enable capture and hence detection of eotaxin. An anti-eotaxin antibody may be used in an assay for the detection of antibodies to eotaxin, for example, in a competitive antibody assay. A labelled eotaxin or a derivative thereof, for example, a recombinant eotaxin or a synthetic peptide comprising part of the amino acid sequence of an eotaxin may be used in a competitive antigen assay for eotaxin or may be used to coat a solid surface in a capture assay for antibodies to eotaxin. The many different types of assay format are well described in the literature of the art, see for example "ELISA and Other Solid Phase Immunoassays, Theoretical and Practical Aspects" Eds. Kemeny D.M. & Challacombe S.J., John Wiley, 1988. (36). Assays using an eotaxin receptor instead of an anti-eotaxin antibody may be carried out analogously.

25

The present invention provides a process for the production of an eotaxin, which comprises obtaining bronchoalveolar lavage fluid obtained from a human or non-human animal challenged with a provoking stimulus, for example, from a human suffering from allergic or non-allergic asthma, or other lung disease, or a guinea-pig sensitised with a foreign protein, and isolating a fraction showing eosinophil chemoattractant activity. One method of isolating an eotaxin-containing fraction of bronchoalveolar lavage fluid is sequential cation exchange, size exclusion and reversed phase HPLC systems.

35

The desired fraction generally contains a polypeptide having a molecular weight in the range from 6-16kDa. Purity may be

2171761

verified by SDS-PAGE. If desired, the authenticity of the substance obtained may be determined by comparison of the amino acid sequence thereof with the amino acid sequence set out in SEQ.ID. NO. 1 or SEQ.ID. NO. 2 (Figure 7 or Figure 8).

5

Eotaxins may be obtained according to the above procedure from other sources, for example, from inflammatory exudate fluids, or from in vitro cultures of macrophages, lymphocytes, neutrophils, mast cells, airway epithelial cells, connective
10 tissue cells, vascular endothelial cells and eosinophils themselves.

Alternatively, a full-length eotaxin, or a part (fragment) of an eotaxin, for example, a polypeptide or peptide fragment,
15 may be produced by chemical synthesis, for example, according to the Merryfield technique. A further method for producing a full-length eotaxin or a part thereof is by recombinant DNA technology. All methods of producing eotaxin are part of the present invention.

20

To produce a full-length eotaxin polypeptide or a polypeptide (or peptide) fragment by recombinant DNA technology, a nucleic acid sequence encoding the polypeptide is inserted into an expression vector under the control of appropriate control
25 sequences. A recombinant polypeptide may then be expressed using a prokaryotic expression system, for example, in *E. coli*, or using a eukaryotic cell system, in which case the resulting polypeptide may be glycosylated. Such techniques are standard see, for example Sambrook, J., Fritsch, E.F. and
30 Maniatis T., *Molecular Cloning A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 1989 (37).

A nucleic acid encoding all or part of an eotaxin polypeptide may be obtained by screening a library prepared from suitable
35 cells, for example, cells from allergen-challenged guinea-pig or human lung. Screening may be carried out using a probe comprising sequences characteristic of an eotaxin, in

particular, sequences that distinguish the eotaxin from other related cytokines, for example, RANTES and MIP-1 α . It may be preferable to use a long probe, for example, a probe comprising a nucleic acid sequence encoding a full-length eotaxin polypeptide.

Alternatively, the polypeptide sequence of an eotaxin may be used to design oligonucleotide primers, for example, degenerate primers. Primers may, for example, comprise bases less specific in their interaction than the naturally-occurring bases, for example, inosine may be used. Examples of primers are the sense sequence

5' TGC TGT TTC CGI GTI ACI AAC AAA (SEQ.ID.NO. 3)

based on the amino acid sequence CCFRVTNK, and the anti-sense sequence

5' CAT CTT GTC IGG CTT IAT TTC (SEQ.ID. NO. 4)

based on the amino acid sequence EIKPDKM. Such primers may be used for amplification of reverse transcribed mRNA by the polymerase chain reaction. This provides cDNA probes for screening libraries, for example, as described above, to isolate full length eotaxin clones. Primers may include codons chosen on the basis of known species preference.

As indicated above, derivatives of naturally occurring eotaxins are also part of the present invention. Such derivatives include polypeptides that have one or more of the following modifications relative to a naturally-occurring eotaxin:

- (i) elongation or shortening at the C-terminus;
- 30 (ii) elongation or shortening at the N-terminus;
- (iii) deletion and/or insertion of internal sequences;
- (iv) amino acid substitutions, for example at the C- and/or N-terminus; and
- (v) a different pattern of glycosylation. (There is, for 35 example, a potential O-glycosylation site at residue 70.)

Such derivatives may function as agonists for structure/

2171761

activity relationship studies or as receptor antagonists.

Anti-eotaxin antibodies and anti-eotaxin receptor-antibodies, both polyclonal and monoclonal, may be produced according to 5 standard techniques, for example, Kohler & Milstein (38). A full-length eotaxin may be used as antigen, or it may be preferred to use a fragment of an eotaxin in order to produce an antibody to a specific antigenic determinant. A naturally occurring eotaxin may be used as antigen. Alternatively, a 10 recombinant or synthetic eotaxin or eotaxin polypeptide or peptide may be used.

Figures 1 to 18 of the accompanying drawings illustrate the present invention. The following is a brief description of 15 the Figures. A more detailed description of the Figures is given below and in the Examples section of this specification.

Figure 1: Protocol for the generation and assay of eosinophil chemoattractant activity in vivo.

20 Figure 2: Time course of generation of eosinophil chemoattractant activity in lungs of sensitized guinea-pigs after antigen challenge.

Figures 3, 4 and 5 all relate to the purification of eotaxin from bronchoalveolar lavage (BAL) fluids:

25 Figure 3: Final reversed phase HPLC profile showing absorbance at 214nm and the acetonitrile gradient.

Figure 4: Measurement of ^{111}In -eosinophil chemoattractant activity of the fractions obtained on HPLC chromatography.

Figure 5: Measurement of ^{111}In -neutrophil chemoattractant activity; lack of significant activity observed. 30

Figure 6: SDS-PAGE analysis of C18 reversed phase HPLC fractions 50-56.

Figures 7 and 8: Amino acid sequences of an isolated guinea-pig eotaxin.

35 Figure 9: Comparison of the eotaxin sequence of Figure 7 with human MCP-1, MCP-2, MCP-3 (21), guinea-pig MCP-1 (24), human MIP-1 α , MIP-1 β and RANTES (18) showing conserved

residues (shaded).

Figure 10: Comparison of ^{111}In -eosinophil accumulation in vivo using guinea-pig eotaxin and the recombinant human proteins RANTES, MIP-1 α and MCP-1. Figure 11 (Inset):

5 Inhibition of the binding of ^{125}I -RANTES to guinea-pig eosinophils in vitro induced by eotaxin and RANTES but not by a member of the C-X-C branch of chemokines, IL-8.

Figures 12, 13 and 14 illustrate the potential of blocking the response to guinea-pig eotaxin. This makes use of the fact
10 that human RANTES, whilst competing with eotaxin for binding sites, does not activate guinea-pig eosinophils.

Figure 12: Elevation of intracellular calcium levels in human eosinophils in vitro induced by eotaxin, RANTES and, at high concentration only, MCP-1.

15 Figure 13: Elevation of intracellular calcium levels in guinea-pig eosinophils in vitro induced by eotaxin, but not human RANTES and MCP-1.

Figure 14: Representative traces showing inhibition of guinea-pig eotaxin-induced elevation of intracellular
20 calcium levels in guinea-pig eosinophils in vitro by the previous exposure of the same cells to human RANTES. Inset:
Figure 15: Values (mean \pm SEM) for four separate eosinophil preparations.

Figure 16: Dose response curve showing inhibition of guinea-
25 pig eotaxin-induced elevation of intracellular calcium levels in guinea-pig eosinophils in vitro by the previous exposure of the same cells to human RANTES.

Figure 17: Histogram showing comparison of in vivo response of guinea-pig eosinophils to eotaxin, RANTES and RANTES plus
30 eotaxin co-injected intradermally.

Figure 18: Histogram showing number of eosinophils and neutrophils in bronchoalveolar lavage fluid (BAL) after administration of guinea-pig eotaxin as an aerosol to guinea-
pig airways.

35

As in allergic asthmatic patients, exposure of sensitised guinea-pigs to aerosolised antigen results in an immediate

2171761

phase of bronchoconstriction with associated mast cell degranulation followed, in some individuals, by a late phase of bronchoconstriction and airway hyper-responsiveness (4-7). Although clearly no one model mimics all the features of the human disease, the guinea-pig model shares common features with the asthmatic response in man and has been extensively used to investigate possible mechanisms (7). In particular, in both guinea-pig and man, the immediate response to allergen triggers the subsequent accumulation in the lung of high numbers of eosinophils. Experiments were designed to detect the appearance in the lung of chemoattractants that may be responsible for the accumulation of eosinophils. A strategy was employed that has previously been applied to the identification of neutrophil chemoattractants in inflammatory exudates (8-10).

Guinea-pigs were sensitised with intraperitoneal ovalbumin on day 1 followed by a short exposure of their lungs to aerosolised ovalbumin at day 8. On day 15-21 animals were challenged with aerosolised ovalbumin and killed at different intervals using a barbiturate overdose. Immediately after death the airways were lavaged with saline. The broncho-alveolar lavage (BAL) fluid was centrifuged to remove cells, and supernatants were either stored at -20°C for assay or subjected to purification. Eosinophil chemoattractant activity in BAL fluid samples or HPLC fractions was tested by injecting them intradermally into assay guinea-pigs previously given intravenous injections of ^{111}In -eosinophils (11,12). After a 2 or 4h interval, assay animals were killed and the punched out skin sites were counted in a gamma-counter (Figure 1).

Figure 2 shows the time-course of appearance of eosinophil chemoattractant activity in BAL fluid. Significant activity was observed 30min after antigen challenge. Activity increased up to 3h, remained high at 6h, but was not significant in 24h samples. Control samples (BAL fluid from

sham-sensitised/challenged or sensitised/sham-challenged guinea-pigs) taken at 3h had no significant activity.

Eosinophil chemoattractant activity, which we termed eotaxin, was purified from 3h BAL fluid by sequential cation exchange, size exclusion and reversed phase HPLC using the *in vivo* ^{111}In -eosinophil accumulation assay to measure the activity of fractions throughout. Eotaxin eluted as a single discrete peak of bioactivity from both the cation exchange and the size exclusion steps, indicating a strongly cationic protein of 7-14kDa. Reversed phase chromatography separated eosinophil chemoattractant activity into two peaks (fractions 51+52 and fraction 54), which were associated with discrete peaks of protein absorbance (Figures 3 and 4). Selectivity for eosinophils was shown by the lack of significant neutrophil chemoattractant activity in these fractions as measured by the accumulation of ^{111}In -neutrophils in the skin assay (Figure 5). Histological examination of skin injected with eotaxin (2pmol) demonstrated eosinophil accumulation at 4 and 24h, particularly around small blood vessels (haematoxylin and eosin stains).

SDS-PAGE analysis revealed a single protein band in each of fractions 51, 52 and 54 (Figure 6). The protein in fractions 51 and 52 was slightly larger than that in fraction 54. This was confirmed by mass analysis in which the major signals were at approximately 8.81 and 8.38 kDa respectively (see Example 3). N-terminal sequencing of fractions 51, 52 and 54 revealed identical amino acid sequences (see Figure 7). The N-terminal 37 residue sequence of eotaxin as set out in SEQ.ID. NO. 1 and Figure 7 shows 57% homology with human monocyte chemotactic protein (MCP-1 (13), otherwise known as MCAF (14) and JE (15)). Tryptic peptides of fraction 54 were also sequenced and readily aligned by comparison with human MCP-1 to give the virtually complete sequence of eotaxin with an overall homology of 53%, see Figure 7. It is likely that the variations in molecular mass reflect differential

glycosylation as the four mass signals obtained (two major and two minor, see Example 3) are all different from each other by multiples of approximately 220 mass units. The sequence contains no N-glycosylation sites, but a potential O-glycosylation site at position 70 has been identified (see Example 3). Human MCP-1 also exhibits heterogeneity on SDS-PAGE due to differences in the O-linked carbohydrate modification (16).

The platelet factor 4 superfamily of chemotactic cytokines, or chemokines, is characterised by four conserved cysteines. The relative position of the two N-terminal cysteines allows the subdivision of this superfamily into the C-X-C chemokines (eg. IL-8 (17)) which are predominantly neutrophil chemoattractants and the C-C chemokines (eg. MCP-1, RANTES, MIP-1 α and MIP-1 β (18)) which are chemotactic for leukocytes other than neutrophils. Eotaxin is a member of the C-C branch of chemokines. The greatest homology (53%) is with human MCP-1 which, in the limited in vitro studies to date, has been reported to be inactive on human eosinophils (19,20) and with the recently described human MCP-2 (54%) and MCP-3 (51% homology calculated on the basis of SEQ.ID. No. 1 (Figure 7); 54% homology when calculated on the basis of the overlapping sequences) (21). Homology with other human C-C chemokines (Figure 9) is: MIP-1 β (37% calculated on the basis of SEQ.ID. No. 1 (Figure 7); 39% when calculated on the basis of the overlapping sequences), MIP-1 α (31% calculated on the basis of SEQ.ID. No. 1 (Figure 7); 32% when calculated on the basis of the overlapping sequences) and RANTES (26% calculated on the basis of SEQ.ID. No. 1 (Figure 7); 27% when calculated on the basis of the overlapping sequences). The latter two proteins have recently been shown to be potent eosinophil activators in vitro (20,22) whereas MIP-1 β activates lymphocytes in vitro (23) but apparently not eosinophils (20). Eotaxin shows the greatest structural homology with human MCP-1, MCP-2 and MCP-3. Eotaxin has functional similarities, but relatively low homology, when compared with RANTES and MIP-1 α . Eotaxin is clearly a distinct molecule from guinea-pig MCP-1; the latter

has recently been cloned (24) and it has only a 43% homology with the eotaxin sequence, see Figure 9. Guinea-pig MCP-1 was shown to be chemotactic for monocytes but was not tested on eosinophils (24). Interestingly, eotaxin has a 41% homology with a C-C protein whose gene is expressed in mouse mast cells and upregulated 2h after the interaction between IgE and antigen (25). No functional activity has been reported for this protein but it is distinct (51% homology) from mouse MCP-1/JE (25).

10

Guinea-pig eotaxin was very potent in inducing eosinophil accumulation in vivo: 1-2pmol/skin site giving a $730 \pm 140\%$ response (mean \pm s.e.m., n=18 animals) compared with saline-injected sites. Further, marked eosinophil accumulation was seen within 30 minutes of intradermal injection (Figure 10). This is supported by experiments in vitro. The effect of eotaxin on eosinophils in vitro was shown by (i) inhibition of binding of ^{125}I -RANTES to guinea-pig cells (Figure 11), (ii) elevation of cytoplasmic calcium in human and guinea-pig cells (Figures 12 and 13) and (iii) chemotaxis of human eosinophils in a Boyden chamber system: the chemotactic responses to eotaxin and RANTES were of similar magnitude over the range 0.1-3.0nM. In contrast to eotaxin, human recombinant MCP-1 and RANTES did not induce guinea-pig eosinophil responses in vivo or in vitro (Figures 10 and 13). This may reflect a species difference although RANTES did bind to guinea-pig eosinophils without inducing activation (Figure 11, inset in Figure 10). Eotaxin has a similar potency to RANTES on human eosinophils whereas MCP-1 is either inactive (20) or active only at high doses (Figure 12). Thus, eotaxin has a potent direct effect on both human and guinea-pig eosinophils.

Responses of guinea-pig eosinophils to guinea-pig eotaxin in vitro and in vivo can be inhibited by human RANTES. Figures 14, 15 and 16 show the response of FURA-2-loaded guinea-pig eosinophils to guinea-pig eotaxin before and after the

2171761

addition of human RANTES. The eotaxin-induced increase in intracellular calcium concentration is reduced to a substantial degree when the cells are first exposed to human RANTES, which itself fails to induce a response. Figure 17 5 shows that human RANTES, when coinjected with the eotaxin, reduces the accumulation of guinea-pig eosinophils induced by eotaxin in guinea-pig skin in vivo. The results support our observations that eotaxin exhibits competitive binding with radiolabelled human RANTES on guinea-pig eosinophils and that 10 eotaxin is a potent functional stimulant whereas RANTES is not. Accordingly, RANTES appears to act as a receptor antagonist for eotaxin on guinea-pig eosinophils.

Figure 18 shows that eosinophils accumulate in guinea-pig 15 airways in vivo within 24 hours of the administration of aerosolized guinea-pig eotaxin, whereas substantially no accumulation of neutrophils is observed.

The following Examples illustrate the present invention.

20

EXAMPLE 1

Production and testing of bronchoalveolar lavage fluid

METHODS: Male Dunkin Hartley guinea-pigs (300-400g) were 25 sensitised with intraperitoneal ovalbumin (OA, 1mg) on day 1 followed by exposure to aerosolised antigen (2% OA for 5min using an ultrasonic nebuliser) on day 8 (6). On day 15-21, animals were pretreated with an antihistamine to prevent acute fatality (pyrilamine, 10mg kg⁻¹, i.p.) and challenged by 30 exposure to aerosolised antigen (1% OA for 20min). At different times after antigen challenge animals were treated with atropine (0.06mg kg⁻¹, i.p.) to prevent bronchoconstriction and killed with a barbiturate overdose. Bronchoalveolar lavage was performed with 4ml saline. Samples were 35 centrifuged to remove cells and the supernatant stored at -20°C prior to assay. BAL samples were assayed by intradermal injection (0.1ml) into guinea-pigs previously given

- 21 -

intravenous injections of 5×10^6 ^{111}In -eosinophils (elicited in donor animals by repeated intraperitoneal injections of horse serum and purified on discontinuous Percoll* gradients, >94% purity) (11,12). After 4 hours, assay animals were killed and the skin sites punched out for gamma-counting. Data are presented as the mean \pm s.e.m. and were tested by one way analysis of variance. A p value of <0.05 was considered statistically significant.

RESULTS: Figure 1 shows, schematically, procedures for the generation and measurement of eosinophil chemoattractant activity in vivo. Figure 2 illustrates the time course of generation of eosinophil chemoattractant activity in lungs of sensitised guinea-pigs after antigen challenge (filled squares, $n=4-10$). Activity was measured in an in vivo skin assay of ^{111}In -eosinophil accumulation in unsensitised guinea-pigs (3 test animals per BAL sample). Significant activity was seen in bronchoalveolar lavage samples taken 0.5, 1.5, 3 and 6h after antigen challenge (compared to responses to intradermal saline, shown as the dotted line). No significant activity was observed in 24h samples. No activity was seen in lavage samples obtained 3h after sham (saline) challenge of sensitised animals (filled circles, $n=5$) or antigen challenge of sham (saline) sensitised animals (filled diamonds, $n=5$).

25

EXAMPLE 2

Purification of eotaxin from bronchoalveolar lavage fluids

METHODS: Bronchoalveolar lavage (BAL) fluid collected from sensitised guinea-pigs (each lavaged with 4ml, followed by 2x10ml, saline) 3h after antigen challenge (1% OA, 5min exposure) was applied to a cation exchange HPLC column (Ultropac* TSK535CM, 7.5x150mm). Activity was eluted at approx 1.4M ammonium acetate, pH5.5, and lyophilised for size exclusion HPLC (Ultropac TSK columns SWP, 7.5x75mm, G4000SW, 7.5x600mm, and G2000SW, 7.5x600mm, in series, equilibrated in 0.08% TFA). Activity eluted at approx 7-14 kDa. This was

* trade-mark

applied to a wide pore (300A) Vydac* C18 reversed phase HPLC column (4x250mm) in 0.08% TFA, eluted with a linear gradient of acetonitrile (0-80% ACN in 0.08% TFA, over 80mins, at 1ml min⁻¹) and 0.5min fractions were collected. Aliquots (4%) of 5 each fraction were lyophilised in the presence of carrier protein (BSA, <0.1ng endotoxin mg⁻¹) and redissolved in 0.8ml saline for testing in the skin bioassays of ¹¹¹In-eosinophil and ¹¹¹In-neutrophil accumulation over 2 hours (n=4). Eosinophils (99%, 0.5% neutrophils) were elicited with 10 repeated intra-peritoneal injections of horse serum (11,12) and neutrophils (99.4%, 0.6% eosinophils) were elicited with a single intraperitoneal injection of casein (5% w/v, 15ml) 17h prior to purification on discontinuous Percoll gradients and labelling with ¹¹¹In.

15

RESULTS: The results are presented in Figures 3, 4 and 5. Figure 3 shows the final reversed phase HPLC profile showing absorbance at 214nm and the acetonitrile gradient. Figure 4 shows that eosinophil chemoattractant activity was seen in 2 20 peaks, corresponding to fractions 51+52 and fraction 54, which corresponded to discreet peaks of absorbance. Figure 5 shows that no significant neutrophil chemoattractant activity was detected in these fractions. In contrast, guinea-pig C5a des Arg (30% zymosan-activated plasma (11), approx 10pmol/site) 25 induced the accumulation of both ¹¹¹In-eosinophils (5211±893) and ¹¹¹In-neutrophils (9872±473). Fractions 50, 53, 55 and 56 consistently gave little or no activity in the guinea-pig skin bioassays of leukocyte accumulation. No significant protein absorbance was detected in the remainder of the gradient (up 30 to 80% acetonitrile).

Example 3

Purity, mass analysis and protein sequence of eotaxin

35 METHODS: 2% aliquots of each fraction were lyophilised, redissolved in 10µl SDS buffer, heated (95°C, 5 min) and 0.3µl run on 8-25% gradient gels in a Pharmacia Phast System. Gels

* trade-mark

were visualised with silver staining. Mass analysis was performed on fractions 51, 52 and 54 using a Finnigan MAT Lasermat with α -cyano-4-hydroxycinnamic acid and sinapinic acid matrices. Mass measurements were calibrated internally using protein standards. 5% aliquots of each bioactive fraction (51, 52 and 54) were applied directly to automated N-terminal sequence analysis using fast cycles on an Applied Biosystems 477A containing a microcartridge essentially as described (32). The amino-terminal 37, 35 and 29 residues were obtained for fractions 51, 52 and 54 respectively. No differences between corresponding positions were found. The apparent initial yields of these three analyses were all approximately 7-8pmol. Thus fractions 51, 52 and 54 contained approx 200pmol each, assuming 70-80% sequencing yields. Gaps were found at positions 8,9 and 33, consistent with the presence of cysteine residues at these positions.

Approximately 30pmol of fraction 54 was reduced and alkylated by sequential treatment with 1mM dithiothreitol for 5min at 50°C and then 10mM acrylamide for 30min at 37°C before digestion with alkylated trypsin (Promega) in 20mM Tris/HCl, pH8.8, containing 0.5% Thesit. Peptides were separated using a Reliasil C18 (300Å, 5µm) column (1x150mm) developed with a linear acetonitrile concentration gradient in 0.08% trifluoroacetic acid at 50µl/min on a Microm HPLC system. Purified peptides were subjected to N-terminal sequence analysis as above, but all four cysteine residues were positively identified as the PTH-cys-S-B-propionamide derivative (33). Position 70 gave no PTH derivative in peptides T6 and T7 and is a probable position of O-glycosylation.

RESULTS: The results are presented in Figures 6, 7, 8 and 9. Figure 6 is a photograph of the SDS-PAGE gel. For reference, human IL-8 (72 amino acids, approx 8kDa) was run in lanes A, B and C (12, 2.4 and 0.5ng/0.3µl/lane respectively). Laser desorption time of flight mass analysis gave signals at 8.81kDa (major) and 9.03kDa (minor) for each of fractions 51

2171761

and 52. Fraction 54 gave signals at 8.38kDa (major) and 8.15kDa (minor).

Figures 7 and SEQ.ID. NO. 1 show the amino acid sequence of eotaxin, which was determined by sequencing the intact molecule as well as peptides derived from digestion with trypsin (T). N-terminal analyses showed the highest homology with human MCP-1 (57%) and the tryptic peptides were readily aligned by comparison with the human MCP-1 sequence. Figure 8 shows the amino acids sequence as confirmed by nucleic acid sequencing. Figure 9 is a comparison of the eotaxin sequence with human MCP-1, MCP-2, MCP-3 (21), guinea-pig MCP-1 (24), human MIP-1 α , MIP-1 β and RANTES (18) showing conserved residues (shaded).

15 Example 4

In vitro and in vivo testing of eotaxin

METHODS:

(i) Eotaxin was a pool of both peaks of bioactivity (fractions 51+52 and fraction 54) from the final reversed phase HPLC separation described in Example 2 (see Figures 3, 4 and 5). ^{111}In -eosinophil accumulation in guinea-pig skin was measured over 4h as described in Example 1 (see Figures 1 and 2). In the same animals (n=4) additional sites were injected with eotaxin 30 minutes before killing.

(ii) For the binding studies, 4×10^5 eosinophils were incubated with 0.1nM ^{125}I -RANTES and various concentrations of cold ligand (50 μl at 0 $^\circ\text{C}$ for 2h). The Hank's buffered salt solution contained 30mM HEPES, 10mM EDTA, 0.1% sodium azide and 1% BSA at pH7.5. Results are the mean of two assays each done in triplicate.

(iii) For measurement of intracellular calcium levels human and guinea-pig eosinophils (10^7 cells/ml in $\text{Ca}^{2+}/\text{Mg}^{2+}$ -free PBS + 0.1% BSA) were loaded with fura-2-acetoxymethyl ester (2.5 μM , 30min at 37 $^\circ\text{C}$). After two washes cells were resuspended at 10^6 cells/ml in $\text{Ca}^{2+}/\text{Mg}^{2+}$ -free PBS containing 10mM HEPES, 0.25% BSA and 10mM glucose (pH7.4). Aliquots were

dispensed into quartz cuvettes and the external $[Ca^{2+}]$ adjusted to 1mM with $CaCl_2$. Changes in fluorescence were monitored at 37°C using a Perkin Elmer LS50 spectrophotometer at excitation wavelengths 340nm and 380nm and emission wavelength 510nm. $[Ca^{2+}]_i$ levels were calculated as described previously (34) using the ratio of the two fluorescence readings and a K_d for Ca^{2+} binding at 37°C of 224nM. Peripheral human eosinophils were prepared as described previously (35) by density centrifugation on Percoll followed by immunomagnetic removal of $CD16^+$ neutrophils using the MACS system. Guinea-pig eosinophils were prepared as described in Example 1 (11,12).

(iv) To test for suppression of eosinophil accumulation in vivo using human RANTES, accumulation of ^{111}I -labelled guinea-pig eosinophils in skin sites was measured as described above. Guinea-pigs were injected with 1.8pmol eotaxin (n=2), 100pmol RANTES (n=1) or both 1.8pmol eotaxin and 100pmol RANTES (n=2). Intradermal saline was used as control.

(v) For testing of receptor antagonist activity, the fura-2-acetoxymethyl ester-loaded guinea-pig eosinophils were stimulated with 3nM eotaxin with or without pretreatment with human RANTES.

(vi) To investigate the effect of aerosol exposure of guinea-pigs to eotaxin, naive guinea-pigs were exposed to an aerosol of either eotaxin or a control medium (n=8 per group).

Exposure was performed by placing two animals in a single chamber and nebulising 24pmol of eotaxin dissolved in 10ml PBS containing BSA carrier protein at 80µg/ml over a period of 35-40 minutes. Control animals received aerosolised PBS/BSA in the same manner. The animals were killed 20 hours after exposure to eotaxin or control medium. BAL fluid (5x10ml HBSS, 10mM EDTA, pH 7.35) was recovered and centrifuged (300g, 20min, 4°C). Total BAL cell count was determined by haemocytometer and differential cell counts performed on stained (DiffQuick)* cytopspin preparations (3 per animal, 400 cell counts per slide).

RESULTS: The results are presented in Figures 10, 11, 12,

* trade-mark

2171761

13, 14, 15, 16, 17 and 18.

Figure 10 shows that guinea-pig eotaxin (1.6pmol) induces significant ^{111}In -eosinophil accumulation in vivo 30min (open squares, $p < 0.01$) and 4h (filled squares, $p < 0.01$) after 5 intradermal injection. In contrast, the recombinant human proteins, RANTES, MIP-1 α and MCP-1, at doses up to 100pmol, were without effect over 4 hours. Figure 11 inset shows that eotaxin and RANTES, but not the C-X-C chemokine IL-8, inhibit the binding of ^{125}I -RANTES ($B_0 = 14.4\%$) to guinea-pig 10 eosinophils in vitro. Figure 17 shows (a) that human RANTES does not induce significant eosinophil accumulation and (b) that human RANTES inhibits to a substantial degree the eosinophil accumulation induced by eotaxin, which suggests that RANTES acts as a receptor antagonist for eotaxin in vivo.

15

Figure 12 shows that eotaxin, RANTES and, at high concentration only, MCP-1 induce elevation of intracellular calcium levels in human eosinophils in vitro. Traces are with eosinophils from one donor. In two other donors 2nM eotaxin 20 gave a mean calcium elevation of 61nM. In the three donors (97.3 \pm 1.2% eosinophils) responses to 10nM RANTES were 194 \pm 74nM $[\text{Ca}^{2+}]_i$ and responses to 100nM MCP-1 were 93 \pm 38nM $[\text{Ca}^{2+}]_i$. Figure 13 shows that guinea-pig eotaxin, but not human RANTES or MCP-1, elevates intracellular calcium levels in guinea-pig 25 eosinophils in vitro. Traces are with cells from one donor. In three donors (97.5 \pm 0.8% eosinophils) responses were: 2nM eotaxin, 90 \pm 13nM $[\text{Ca}^{2+}]_i$; 100nM RANTES, 2.0 \pm 1.7nM $[\text{Ca}^{2+}]_i$; 100nM MCP-1, 3.3 \pm 0.7nM $[\text{Ca}^{2+}]_i$.

30 Figure 14 and the inset Figure 15 show that prior treatment of guinea-pig eosinophils with RANTES (100nM) inhibits the increase in intracellular free calcium levels seen when eosinophils are treated with eotaxin (3nM) alone. RANTES appears to be acting as a receptor antagonist in guinea-pig 35 eosinophils. Figure 16 is a dose-response curve using 3nM eotaxin and increasing amounts of RANTES.

Figure 18 shows that guinea-pig eotaxin, administered as an aerosol, induces eosinophil accumulation in guinea-pig airways in vivo, whereas substantially no accumulation of neutrophils is observed.

5

REFERENCES

1. de Monchy, J. G. R., Kauffman, H. F., Venge, P., et al. Am. Rev. Respir. Dis. 131, 373-376 (1985).
2. Bousquet, J., Chanez, P., Lacoste, J. Y., et al. N. Engl. J. Med. 323, 1033-1039 (1990).
3. Bradley, B. L., Azzawi, M., Jacobson, M., et al. J. Allergy Clin. Immunol. 88, 661-674 (1991).
4. Dunn, C. J., Elliott, G. A., Oostveen, J. A. & Richards, I. M. Am. Rev. Respir. Dis. 137, 541-547 (1988).
5. Sanjar, S., Aoki, S., Kristersson, A., Smith, D. & Morley, J. Br. J. Pharmacol. 99, 679-686 (1990).
6. Griffiths-Johnson, D. A. & Karol, M. H. Toxicology 65, 283-294 (1991).
7. Campos, M. G. & Church, M. K. Clin. Exp. Allergy 22, 665-666 (1992).
8. Collins, P. D., Jose, P. J. & Williams, T. J. J. Immunol. 146, 677-684 (1991).
9. Beaubien, B. C., Collins, P. D., Jose, P. J., et al. Biochem. J. 271, 797-801 (1990).
10. Jose, P. J., Collins, P. D., Perkins, J. A., et al. Biochem. J. 278, 493-497 (1991).
11. Faccioli, L. H., Nourshargh, S., Moqbel, R., et al. Immunology 73, 222-227 (1991)
12. Weg, V. B., Williams, T. J., Lobb, R. R. & Nourshargh, S. J. Exp. Med. 177, 561-566 (1993).
13. Yoshimura, T., Yuhki, N., Moore, S. K., Appella, E., Lerman, M. I. & Leonard, E. J. FEBS Letts. 244, 487-493 (1989).
14. Furutani, Y., Nomura, H., Notake, M., et al. Biochem. Biophys. Res. Commun. 159, 249-255 (1989).
15. Rollins, B. J., Morrison, E. D. & Stiles, C. D. Proc. Natl. Acad. Sci. USA 85, 3738-3742 (1988).
16. Jiang, Y., Valente, A. J., Williamson, M. J., Zhang, L.

2171761

- & Graves, D. T. *J. Biol. Chem.* 265, 18318-18321 (1990).
17. Jiang, Y., Valente, A. J., Williamson, M. J., Zhang, L. & Graves, D. T. *J. Biol. Chem.* 265, 18318-18321 (1990).
18. Schall, T. J. *Cytokine* 3, 165-183 (1991).
- 5 19. Leonard, E. J. & Yoshimura, T. *Immunology Today* 11, 97-101 (1990).
20. Rot, A., Krieger, M., Brunner, T., Bischoff, S. C., Schall, T. J. & Dahinden, C. A. *J. Exp. Med.* 176, 1489-1495 (1992).
- 10 21. Van Damme, J., Proost, P., Lenaerts, J-P. & Opdenakker, G. J. *J. Exp. Med.* 176, 59-65 (1992).
22. Kameyoshi, Y., Dorschner, A., Mallet, A. I., Christophers, E. & Schroder, J-M. *J. Exp. Med.* 176, 587-592 (1992).
- 15 23. Tanaka, Y., Adams, D. H., Hubscher, S., Hirano, H., Siebenlist, U. & Shaw, S. *Nature* 361, 79-82 (1993).
24. Yoshimura, T. *J. Immunol.* 150, 5025-5032 (1993).
25. Kulmburg, P. A., Huber, N. E., Scheer, B. J., Wrann, M. & Baumruker, T. *J. Exp. Med.* 176, 1773-1778 (1992).
- 20 26. Frigas, E. & Gleich, G. J. *J. Allergy Clin. Immunol.* 77, 527-537 (1986).
27. Wegner, C. D., Gundel, R. H., Reilly, P., Haynes, N., Letts, L. G. & Rothlein, R. *Science* 247, 456-459 (1990).
28. Lellouch-Tubiana, A., Lefort, J., Simon, M-T., Pfister, A. & Vargaftig, B. B. *Am. Rev. Respir. Dis.* 137, 948-954 (1988).
29. Coyle, A. J., Page, C. P., Atkinson, L., Flanagan, R. & Metzger, W. J. *Am. Rev. Respir. Dis.* 142, 587-593 (1990).
- 30 30. Silva, P. M. R., Martins, M. A., Castro-Faria-Neto, H. C., Cordeiro, R. S. B. & Vargaftig, B. B. *J. Pharmacol. Exp. Ther.* 257, 1039 (1991).
31. Cochran, B. H., Reffel, A. C. & Stiles, C. D. *Cell* 33, 939-947 (1983).
32. Totty, N. F., Waterfield, M. D. & Hsuan, J. J. *Protein-Science* 1, 1215-1224 (1992).
33. Brune, D. C. *Anal. Biochem.* 207, 285-290 (1992).
34. Gryniewicz, G., Poenie, M. & Tsien, R. Y. *J. Biol. Chem.*

260, 3440-3450 (1985).

35. Anwar, A. R. F., Moqbel, R., Walsh, G. M., Kay, A. B. & Wardlaw, A. J. J. Exp. Med. 177, 839-843 (1993).

36. ELISA and Other Solid Phase Immunoassays, Theoretical and 5 Practical Aspects" Eds. Kemeny D.M. & Challacombe S.J., John Wiley, 1988.

37. Sambrook, J., Fritsch, E.F. and Maniatis T., Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory Press, 1989.

10 38. Kohler & Milstein, Nature 256, 495-497 (1975).

2171761

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: National Heart & Lung Institute
(B) STREET: Dovehouse Street
(C) CITY: London
(E) COUNTRY: England
(F) POSTAL CODE (ZIP): SW3 6LY
(G) TELEPHONE: 071 352 8121
(H) TELEFAX: 071 376 3442

(A) NAME: Ludwig Institute for Cancer Research
(B) STREET: Hedges House, 153-155 Regent Street
(C) CITY: London
(E) COUNTRY: England
(F) POSTAL CODE (ZIP): W1R 7FD
(G) TELEPHONE: 071 494 0025
(H) TELEFAX: 071 494 1404

(ii) TITLE OF INVENTION: CHEMOTACTIC CYTOKINE

(iii) NUMBER OF SEQUENCES: 4

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

(vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: GB 9318984
(B) FILING DATE: 14-SEP-1993

(vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: GB 9408602
(B) FILING DATE: 29-APR-1994

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 73 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Cavia porcellus
(D) DEVELOPMENTAL STAGE: Adult
(F) TISSUE TYPE: Bronchoalveolar lavage fluid

- 31 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

His	Pro	Gly	Ile	Pro	Ser	Ala	Cys	Cys	Phe	Arg	Val	Thr	Asn	Lys	Lys
1				5					10					15	
Ile	Ser	Phe	Gln	Arg	Leu	Lys	Ser	Tyr	Lys	Ile	Ile	Thr	Ser	Ser	Lys
			20					25					30		
Cys	Pro	Gln	Thr	Ala	Ile	Val	Phe	Glu	Ile	Lys	Pro	Asp	Lys	Met	Ile
		35					40					45			
Cys	Ala	Asp	Pro	Lys	Xaa	Xaa	Trp	Val	Gln	Asp	Ala	Lys	Lys	Tyr	Leu
	50					55					60				
Asp	Gln	Ile	Ser	Gln	Xaa	Thr	Lys	Pro							
65					70										

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 73 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Cavia cobaya
- (F) TISSUE TYPE: Bronchial lavage fluid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

His	Pro	Gly	Ile	Pro	Ser	Ala	Cys	Cys	Phe	Arg	Val	Thr	Asn	Lys	Lys
1				5					10					15	
Ile	Ser	Phe	Gln	Arg	Leu	Lys	Ser	Tyr	Lys	Ile	Ile	Thr	Ser	Ser	Lys
			20					25					30		
Cys	Pro	Gln	Thr	Ala	Ile	Val	Phe	Glu	Ile	Lys	Pro	Asp	Lys	Met	Ile
		35					40					45			
Cys	Ala	Asp	Pro	Lys	Lys	Lys	Trp	Val	Gln	Asp	Ala	Lys	Lys	Tyr	Leu
	50					55					60				
Asp	Gln	Ile	Ser	Gln	Thr	Thr	Lys	Pro							
65					70										

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

2171761

- 32 -

- (ii) MOLECULE TYPE: DNA (primer)
- (iii) HYPOTHETICAL: YES
- (iii) ANTI-SENSE: NO

- (ix) FEATURE:
 - (A) NAME/KEY: modified_base: N is inosine
 - (B) LOCATION: 12

- (ix) FEATURE:
 - (A) NAME/KEY: modified_base: N is inosine
 - (B) LOCATION: 15

- (ix) FEATURE:
 - (A) NAME/KEY: modified_base: N is inosine
 - (B) LOCATION: 18

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

TGCTGTTTCC GNGTNACNAA CAAA

24

- (2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (primer)
- (iii) HYPOTHETICAL: YES
- (iii) ANTI-SENSE: YES

- (ix) FEATURE:
 - (A) NAME/KEY: modified_base: N is inosine
 - (B) LOCATION: 10

- (ix) FEATURE:
 - (A) NAME/KEY: modified_base: N is inosine
 - (B) LOCATION: 16

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

CATCTTGTCN GGCTTNATTT C

21

CLAIMS

1. An isolated chemoattractant protein capable of attracting eosinophils and of inducing eosinophil accumulation or activation *in vitro* and *in vivo* and which shows substantially no attractive effect for neutrophils *in vivo*, consisting of an amino acid sequence which is at least 60% identical to the amino acid sequence set out in SEQ ID NO. 1 or SEQ ID NO. 2, calculated by the BESTFIT™ computer program.
2. A chemoattractant protein as claimed in claim 1, having at least 80% identity with the amino acid sequence set out in SEQ ID NO. 1 or SEQ ID NO. 2.
3. A chemoattractant protein as claimed in claim 1, consisting of or comprising an amino acid sequence as set out in SEQ ID NO. 1 or SEQ ID NO. 2.
4. A process for the production of a chemoattractant protein as claimed in any one of claims 1 to 3, which comprises isolating a fraction from broncho-alveolar lavage fluid or an inflammatory exudate obtained from a human or a non-human animal challenged with a provoking stimulus, wherein said fraction demonstrates eosinophil chemoattractant activity *in vitro* and *in vivo* and shows substantially no attractive effect for neutrophils *in vivo*.
5. A process for the production of a chemoattractant protein as claimed in any one of claims 1 to 3, which comprises culturing *in vitro* macrophages, lymphocytes, neutrophils, mast cells, airway epithelial cells, connective tissue cells, vascular endothelial cells or eosinophils obtained from a human or a non-human animal, and isolating from the cells or from the cell culture fluid a fraction that demonstrates eosinophil chemoattractant activity *in vitro* and *in vivo* and which shows substantially no attractive effect for neutrophils *in vivo* and further isolating the protein from said fraction.
6. An agent that inhibits or otherwise hinders the production, release or action of a chemoattractant protein as claimed in any one of claims 1 to 3, said agent being an antibody which specifically binds to the chemoattractant protein as claimed in claim 3.

7. An agent as claimed in claim 6, being a monoclonal antibody.
8. Use of an agent according to claim 6 for the treatment of an inflammatory disease.
9. Use according to claim 8, wherein said inflammatory disease is asthma.
10. Use of an antibody according to claim 6 for the preparation of a medicament for the treatment of an inflammatory disease.
11. Use according to claim 10, wherein said inflammatory disease is asthma.
12. Use according to claim 8, wherein said inflammatory disease is rhinitis or eczema.
13. A pharmaceutical preparation which comprises, as active ingredient, an antibody as claimed in claim 6, in admixture or conjunction with a pharmaceutically suitable carrier.
14. An assay for an antigen or for one member of a specific binding pair, characterised in that the antigen is a chemoattractant protein as claimed in any one of claims 1 to 3, comprising the use of an antibody against said chemoattractant protein to detect said antigen.
15. An assay for an antibody, characterised in that the antibody is an antibody that forms a complex with a chemoattractant protein as claimed in any one of claims 1 to 3, comprising the use of said chemoattractant protein to detect said antibody.
16. A method of testing a compound for an inhibitory effect on the activity of a chemoattractant cytokine *in vitro*, characterised in that the chemoattractant cytokine is a chemoattractant protein as claimed in any one of claims 1 to 3, comprising determining the inhibitory effect of said compound on said chemoattractant protein.

2171761

Antigen Challenge



Lavage



Bioassay




Figure 1




Figure 2




Figure 3




Figure 4





Figure 5

2171761



A 50 51 52 53 54 55 56 B C
fraction

Figure 6

10 20 30 40 50 60 70
 HPGIPSACCFRVTNKKISFQRLKSYKIITSSKCPQTAIVFEIKPDKMICADPKXXWVQDAKKYLDQISQXTKP
 N-terminal analyses T4 T5 T6
T1 T2 T3 T7

Figure 7

10 20 30 40 50 60 70
 HPGIPSACCFRVTNKKISFORLKSXKIITSSKCPQTAIVFEIKPKDMICADPKKXWVQDAKKYLDQISQTTKP

Figure 8

HPGIPSACCFRVTNKKISFORLKSXKIITSSKCPQTAIVFEIKPKDMICADPKXWVQDAKKYLDQISQTTKP
 QPDAINAEVTCYNEFTRKISVORLASVRRITSSKCPKEAVIEKTI VAKEICADPKOKHVODSMDHLDKOTQTPET
 DSVSIPITCCFNVINRKIPIORLESYTRITNIQCPKEAVIEKTKRGKEVCADPKERHVODSMDHLDQIFONLKP
 KSTTECYHFINKKIPKORLESYRRTSSHCEREAVIDE K DKEICADPTOKHVODFMEHLDKKTQTPKL
 GVNTF TCCYTF NKQIPLKRVKGERITSSRCPQEAVIDERTLKNKEVCADPTOKHVODYIAKLDORTQOKON
 SLAADTEACCFSYTSRQXPONFIADY FETSSQSKPGVIEFLTERSRQVCADPSEEHVOKYVSDIELSA
 PMGSDPETAACCFSYTARKLPRNFVVDY YETSSLSQPAVVEPOTKRSKQVCADPSESEHVQYVYDIELN
 SPYSSDITPCCFAYIARPLPRAHIKEY FYTSKCSNPAVVEVTRKNRQVCANPEKKVYREYINSEMS

Figure_9




Figure 10




Figure 12




Figure 13





Figure 15




Figure 14




Figure 16



Figure 17




Figure 18