



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|---|--|---|---|
| (51) International Patent Classification ⁶ : A61K 9/127, 31/70, C07H 21/04, C12N 15/00 | | A1 | (11) International Publication Number: WO 98/14172 |
| | | | (43) International Publication Date: 9 April 1998 (09.04.98) |
| (21) International Application Number: PCT/US97/18348 | | (81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). | |
| (22) International Filing Date: 3 October 1997 (03.10.97) | | | |
| (30) Priority Data: 08/726,211 4 October 1996 (04.10.96) US | | Published <i>With international search report.</i> | |
| (71) Applicant (for all designated States except US): BOARD OF REGENTS, THE UNIVERSITY OF TEXAS SYSTEM [US/US]; 201 West 7th Street, Austin, TX 78701 (US). | | | |
| (72) Inventors; and | | | |
| (75) Inventors/Applicants (for US only): TORMO, Mar [ES/ES]; Guillem Sorolla, 36, E-Valencia (ES). TARA, Ana, M. [PT/US]; 7500 Kirby Drive #311, Houston, TX 77030 (US). LOPEZ-BERESTEIN, Gabriel [US/US]; 122 Bellaire Court, Bellaire, TX 77602 (US). MCDONNEL, Timothy, J. [US/US]; 14103 Manderly, Houston, TX 77077 (US). | | | |
| (74) Agent: CORDER, Timothy, S.; Arnold, White & Durkee, P.O. Box 4433, Houston, TX 77210 (US). | | | |

(54) Title: INHIBITION OF BCL-2 PROTEIN EXPRESSION BY LIPOSOMAL ANTISENSE OLIGODEOXYNUCLEOTIDES

(57) Abstract

The present invention provides novel compositions and methods for use in the treatment of Bcl-2-associated diseases like cancer, specifically, in the treatment of follicular lymphoma (FL). The compositions contain antisense oligonucleotides that hybridize to Bcl-2 nucleic acids, the gene products of which are known to interact with the tumorigenic protein Bcl-2. Used alone, or in conjunction with other antisense oligonucleotides, these compositions inhibit the proliferation of FL cancer cells.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|---------------------------------------|----|---|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

DESCRIPTIONINHIBITION OF BCL-2 PROTEIN EXPRESSION BY
LIPOSOMAL ANTISENSE OLIGODEOXYNUCLEOTIDESBACKGROUND OF THE INVENTION5 **A. Field of the Invention**

The present invention relates to the field of cancer therapy, specifically, the treatment of Bcl-2 diseases. More particularly, these treatments involve the use of antisense oligodeoxynucleotides and liposomal formulations thereof.

B. Related Art

10 Bcl-2 has been linked to a wide variety of diseases such as hematologic malignancies, both leukemias and lymphomas, including follicular and nonfollicular lymphomas, chronic lymphocytic leukemia, and plasma cell dyscrasias (Campos *et al.*, *Blood*, 84:595, 1994); solid tumors like those associated with breast, prostate and colon cancer; and immune disorders. One particular Bcl-2-related disease is Follicular non-Hodgkin Lymphoma (FL). FL is the most
15 common lymphoid malignancy in Europe and the United States. Typically it is an indolent, low grade disease consisting of an accumulation of small, resting B cells. Although the response to chemotherapy is initially good, relapses are inevitable with the transformation to a more aggressive histological type and the development of drug resistance (Aisenberg, *J. Clin. Oncol.*, 13:2656, 1995; Johnson *et. al.*, *J. Clin. Oncol.*, 13:140, 1995). In over 90% of FL patients, a
20 t(14;18) translocation is found, which results in the juxtaposition of the *bcl-2* gene from chromosome 18q21 with the immunoglobulin heavy chain gene locus on chromosome 14q323 (Tsujimoto *et. al.*, *Science*, 229:1390, 1985; Graninger *et. al.*, *J. Clin. Invest.*, 80:1512, 1987). As a consequence, the *bcl-2* gene is under the influence of immunoglobulin heavy chain enhancer, and the Bcl-2 protein is overexpressed (Bakhshi *et. al.*, *Cell*, 41:899, 1985; Tsujimoto *et. al.*,
25 *Oncogene*, 2:3, 1987). Bcl-2 tumorigenic potential is related to its capacity of interfering with physiological death responses, thereby enhancing the longevity of the cell (Nuñez *et. al.*, *J. Immunol.*, 144:3602, 1990). The Bcl-2 protein blocks apoptotic stimuli such as growth factor

deprivation, radiation, heat-shock, virus, and most of the chemotherapeutic agents (Reed, *Hematol. Oncol. Clin. North Am.*, 9:451, 1995; Hockenberry *et. al*, *Nature*, 348:334, 1990). In *bcl-2*-Ig-transgenic mice, a polyclonal follicular lymphoproliferation consisting of an expansion of mature B lymphocytes is initially observed (McDonnell *et. al*, *Cell*, 57:79, 1989). 5 Subsequently, a monoclonal high grade large immunoblastic type lymphomas develop with 50% of them presenting rearrangement of C-MYC. This suggests that a second genetic alteration is necessary for the development and progression of malignant lymphoma (McDonnell and Korsmeyer, *Nature*, 349:254, 1991).

Recently, an expanding family of Bcl-2-related proteins has been identified. This 10 includes Bax, Bcl-X_L, Bcl-X_S, Bad, Bak, Mcl-1, A-1, and several open reading frames in DNA viruses (Oltvai *et. al*, *Cell*, 74:609, 1993; Boise *et. al*, *Cell*, 74:597, 1993; Yang *et. al*, *Cell*, 80:285, 1995; Chittenden *et. al*, *Nature*, 374:733, 1995; Kiefer *et. al*, *Nature*, 374: 736, 1995; Kozopas *et. al* *Proc. Nat'l Acad. Sci. USA*, 90:3516, 1993; Lin *et. al*, *J. Immunol.*, 151:1979, 1993; Pearson *et. al*, *Virology*, 160:151, 1987; Neilan *et. al*, *J. Virol.*, 67:4391, 1993). 15 Membership in the Bcl-2 family of proteins is principally defined by homology within the BH1 and BH2 domains, which help regulate dimerization between the members (Sato *et. al*, *Proc. Nat'l Acad. Sci. USA*, 91:9238, 1994). Bax, which shares 21% amino-acid identity with Bcl-2, can bind to Bcl-2 protein and neutralize its ability to block cell death. Thus, the ratio of Bcl-2 to Bax is thought to determine the cell's susceptibility to death following an apoptotic stimulus 20 (Oltvai *et. al*, 1993; Yin *et. al*, *Nature*, 369: 321, 1994).

Phosphodiester antisense oligodeoxynucleotides complementary to specific sequences 25 of the translation-initiation site of Bcl-2 mRNA are able to inhibit the production of the Bcl-2 protein and the growth of t(14;18) translocation bearing cells (Kitada *et. al*, *Antisense Res. Dev.*, 3:157, 1993). However, the therapeutic use of antisense oligonucleotides has been hampered by their low cellular uptake and their rapid degradation by nucleases and other serum or cellular components. Phosphorothioate oligonucleotides, which are resistant to nuclease degradation, 30 were found to inhibit FL cell growth at concentrations 10 times lower than phosphodiester oligonucleotides (Reed *et. al*, *Cancer Research*, 50: 6565, 1990a; Cotter *et. al*, *Oncogene*, 9:3049, 1994). However, this approach suffers from low cellular uptake of the oligonucleotides. For example, Reed *et. al* had to use concentrations of greater than 25μM of phosphorothioates to achieve 50% growth inhibitions of cell lines derived from B-cell lymphomas, such as 697 and

Su-Dhl-4 cells. Liposomal incorporation has led to enhanced uptake of oligonucleotides into leukemic cells (Akhtar *et. al*, *Nucleic Acids Res.*, 19:5551, 1991; Tari *et. al*, *Blood*, 84:601, 1994). The use of cationic lipids by Reed *et. al* to deliver phosphorothioate antisense oligonucleotides allowed them to reduce the concentration of oligonucleotides to 0.075 to 0.3 5 μ M and still induce growth inhibition in Su-Dhl-4 cells.

There is still, however, a great need for methods and compositions for the treatment of Bcl-2 associated diseases such as hematologic malignancies, both leukemias and lymphomas, including follicular and nonfollicular lymphomas, chronic lymphocytic leukemia, and plasma 10 cell dyscrasias; solid tumors like those associated with breast, prostate and colon cancer; and immune disorders.

SUMMARY OF THE INVENTION

The present invention is designed to overcome the shortcomings of the prior art by providing improved compositions and methods for the treatment of Bcl-2 associated diseases, such as hematologic malignancies, both leukemias and lymphomas, including follicular and 15 nonfollicular lymphomas, chronic lymphocytic leukemia, and plasma cell dyscrasias; solid tumors like those associated with breast, prostate and colon cancer; and immune disorders, using novel antisense oligonucleotides to target specific nucleic acids in the cells of patients.

Thus, in one embodiment, there is provided a composition comprising a polynucleotide that hybridizes to a Bcl-2-encoding polynucleotide under intracellular conditions. Intracellular 20 conditions as defined herein include monovalent cation at a concentration of approximately 160 mM (10 mM Na^+ ; 150 mM K^+) and a concentration of divalent cation of approximately 20 mM (18 mM Mg^+ ; 2 mM Ca^{++}). Intracellular conditions may also include a protein concentration, which would serve to decrease the volume of hybridization and, therefore, increase the effective concentration of nucleic acid species, of about 150 mg/ml.

25 The polynucleotides of the present invention may be oligonucleotides having a length of about 8-50 bases. The polynucleotides may be phosphodiester polynucleotides as found in nature, or they may preferably be derivatized polynucleotides or even polynucleotide analogs. Exemplary polymers would be p-ethoxy or methylphosphonate polynucleotides. Other analogs

that may be used include, but are not limited to C-5 propyne pyrimidine polynucleotides (Wagner *et al. Science*, 260:1510-1513, 1993), or polynucleotides as described in US Patent 5,138,045 or European Patent Application EP 431,523, each incorporated herein by reference. Preferred analogs are nuclease resistant, have a high melting temperature, (bind tightly to RNA), and in certain embodiments preferred polynucleotides are hydrophobic for more efficient association with a lipid formulation such as a liposome.

In a further embodiment, the polynucleotide hybridizes to bcl-2 mRNA and preferable to the translation initiation site of Bcl-2 mRNA. It is understood however, that the antisense molecules of the present invention may hybridize to any area of the bcl-2 transcript that is effective to downregulate expression of the Bcl-2 protein. There are certain advantages to targeting the first open reading frame, and as stated, the transcription initiation site is particularly preferred. In certain specific embodiments, the polynucleotide may be an oligonucleotide having the sequence ^{5'}CAGCGTGCGCCATCCTTC^{3'} (SEQ ID NO:1).

The polynucleotide preparation of the present invention is preferably associated with a lipid. A polynucleotide associated with a lipid may be described as encapsulated in the aqueous interior of a liposome, interspersed within the lipid bilayer of a liposome, attached to a liposome via a linking molecule that is associated with both the liposome and the polynucleotide, complexed with a lipid, dispersed in a solution containing a lipid, mixed with a lipid, combined with a lipid, contained as a suspension in a lipid, contained or complexed with a micelle, or otherwise associated with a lipid.

The term "lipids" as used in this specification and the claims denotes any form of both naturally occurring and synthetic lipids or liposomes. Lipids are fatty substances and are well-known by those of skill in the art. The lipids of the present invention are not limited to any particular structure in solution. For example, they may be present in a bilayer structure, as micelles, or with a "collapsed" structure. They may also simply be interspersed in a solution, possibly forming aggregates which are not uniform in either size or shape. The lipids are preferably neutral in net charge and may advantageously be comprised of the lipid dioleoylphosphatidylcholine, however other lipids such as other phosphatidylcholines, phosphatidylglycerols, and phosphatidylethanolamines may also be employed.

In yet another embodiment, there is provided a composition comprising a polynucleotide that hybridizes to a Bcl-2-encoding polynucleotide.

5 In still yet another embodiment, there is provided a composition comprising an expression construct that encodes a polynucleotide that hybridizes to a Bcl-2-encoding polynucleotide, wherein said first polynucleotide is under the control of a promoter that is active in eukaryotic cells.

This invention also comprises a method for inhibiting proliferation of a cancer cell comprising contacting said cancer cell with a composition comprising at least a polynucleotide that hybridizes to a Bcl-2-encoding nucleic acid. This method may be applied advantageously 10 to a variety of cancer cells as shown in the Examples below. The methods of the invention are shown to be effective in cancer cells that overexpress or express high levels of bcl-2 and also express the Bax protein, as well as in cells in which translocation of p53 into the nucleus upon genotoxic damage is inhibited by bcl-2. Such cells would include, but not be limited to 15 hematologic malignancies, both leukemias and lymphomas, including follicular and nonfollicular lymphomas, chronic lymphocytic leukemia, and plasma cell dyscrasias; solid tumors like those associated with breast, prostate and colon cancer; and immune disorders. The composition may comprise a neutral lipid which is associated with the polynucleotide, such as a 20 polynucleotide encapsulated in a liposome. In a specific embodiment, the contacting takes place in a patient, and in certain embodiments in a human patient. The composition may advantageously be delivered to a human patient in a volume of 0.50-10.0 ml per dose or in an amount of 5-30 mg polynucleotide per m². In a particular regimen, the composition is 25 administered 3 times per week for 8 weeks. However, various dosages may be used as determined by an attending physician based on the particular disease, age and condition of the patient and other factors that may affect a physicians decision.

25 This invention relates to antisense technology that may be employed to treat Bcl-2-associated disease. In one embodiment it encompasses a composition comprising a polynucleotide that hybridizes to a Bcl-2-encoding polynucleotide and a neutral lipid associated 30 with the polynucleotide, and preferably those that exclude cationic lipids. The polynucleotide may be an oligonucleotide having a length of between about 8 and about 50 bases. However, oligonucleotides of other lengths may also be useful. The polynucleotide may also hybridize to

the translation initiation site of Bcl-2 mRNA. An example of a useful polynucleotide is an oligonucleotide comprising the sequence CAGCGTGCGCCATCCTTC (SEQ ID NO:1).

5 Compositions of the present invention also include compositions wherein liposomes are formed from the lipid. In some cases, it may be useful to have a composition in which the polynucleotide is encapsulated in the liposome. Lipids that are particularly useful in the present invention include phosphatidylcholines, phosphatidylglycerols, and phosphatidylethanolamines, one example being the lipid dioleoylphosphatidylcholine, although any acceptable neutral lipids may be used.

10 An embodiment of this invention is a composition comprising an expression construct that encodes a polynucleotide that hybridizes to a Bcl-2-encoding polynucleotide, wherein said polynucleotide is under the control of a promoter that is active in eukaryotic cells.

15 Another embodiment encompasses a method of inhibiting a Bcl-2-associated disease comprising obtaining a polynucleotide that hybridizes to a Bcl-2-encoding polynucleotide, mixing the polynucleotide with a neutral lipid to form a polynucleotide/lipid association, and administering said association to a cell. The cell may be a cancer cell, such as a follicular lymphoma cell. This method may employ a polynucleotide comprising an oligonucleotide having a length of between about 8 and about 50 bases. The lipid may comprise a liposome. If so, the liposome may further encapsulate the polynucleotide.

20 This embodiment also includes methods wherein the contacting takes place in an animal, such as a human. For example, the composition may be delivered to said human in a volume of 0.50-10.0 ml per dose or in an amount of from about 5 to about 30 mg polynucleotide per m². It may also be administered three times per week for eight weeks.

25 Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

5 The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein:

10 **FIG. 1:** Growth inhibition of lymphoid cells by liposomal-bcl-2-antisense oligonucleotides ("L-bcl-2"). Final concentrations of 3 $\mu\text{mol/L}$ (dots), 4 $\mu\text{mol/L}$ (horizontal), 5 $\mu\text{mol/L}$ (vertical) and 6 $\mu\text{mol/L}$ (diagonal) of L-bcl-2 were added to Johnson, Jurkat, Raji and Daudi cells. After 5 days, the viability of the tumoral cells was measured by alamarBlue dye. Viability was expressed as percent of untreated cells.

15 **FIG. 2:** Non-specific toxicity in lymphoid cells at 6 $\mu\text{mol/L}$ of liposomal oligonucleotides. Empty liposomes (diagonal) and two different liposomal control oligonucleotides ("L-control oligos") (gray, black) were added to Johnson, Jurkat, Raji and Daudi cells at 6 $\mu\text{mol/L}$ final concentration. After 5 days, the viability of the tumoral cells was measured by alamar blue dye. Viability was expressed as percent of untreated cells.

20 **FIG. 3:** Western blot analysis of Bcl-2 protein in the four cell lines. Johnson, Jurkat, Daudi and Raji cells were lysed in sample buffer and normalized for total protein content. Twenty-five grams of total protein was loaded in each lane. The membranes were incubated with hamster anti-human bcl-2 monoclonal antibody. In Johnson cells, a cell line bearing the t(14;18) translocation, overexpression of Bcl-2 protein is observed. In Jurkat and Raji cells, which lack the t(14;18) translocation, expression of Bcl-2 is low. In Daudi cells, Bcl-2 expression is not observed.

FIG. 4A & FIG. 4B: Specific inhibition of Bcl-2 protein in Johnson and Jurkat cells by L-bcl-2.

25 **FIG. 4A:** 1×10^5 Johnson cells/mL in 3 mL were treated with 3 and 4 $\mu\text{mol/L}$ of L-bcl-2 or L-control oligos. After 3 days of culture, protein-containing lysates were prepared and 5 μg of total protein were subjected to SDS-PAGE and transferred to nitrocellulose membranes. Blots were cut into sections and incubated with antibodies specific for either Bcl-2 or Actin

(left). To estimate the inhibition of bcl-2 protein, data were quantified by scanning densitometry and expressed as ratio of Bcl-2:Actin (right). L-bcl-2:(filled squares); L-control:(open circles).

5 **FIG. 4B:** 1×10^5 Jurkat cells/mL in 3 mL were treated with 3 and 4 $\mu\text{mol/L}$ of L-bcl-2 or L-control oligonucleotides. After 3 days of culture, protein-containing lysates were prepared and 20 μg of total protein were subjected to SDS-PAGE and transferred to nitrocellulose membranes. Blots were cut into sections and incubated with antibodies specific for either Bcl-2 or Actin (left). To estimate the inhibition of bcl-2 protein, data were quantified by scanning densitometry and expressed as ratio of Bcl-2:Actin (right). L-bcl-2:(filled squares); L-control:(open circles).

10

FIG. 5: Western blot analysis of Bax protein in the four cell lines. Johnson, Jurkat, Daudi and Raji cells were lysed in sample buffer and normalized for total protein content. Fifty μg of total protein was loaded in each lane. The membranes were incubated with rabbit anti-human Bax polyclonal antibody.

15 **FIG. 6A & FIG. 6B:** Bcl-2/Bax ratio decreases in Johnson cells by L-bcl-2.

20 **FIG. 6A:** 1×10^5 Johnson cells/mL in 3 mL were treated with 2, 3 and 4 $\mu\text{mol/L}$ of L-bcl-2 or L-control oligos. After 3 days of culture, protein-containing lysates were subjected to SDS-PAGE and transferred to nitrocellulose membranes. Blots were cut into sections, and incubated with antibodies specific for either Bax or Actin. This experiment was made using the same lysates obtained in the experiment shown in FIG. 4A and FIG. 4B.

FIG. 6B: Data were quantified by scanning densitometry and expressed as ratio of Bcl-2:Bax. L-bcl-2:(filled squares); L-control oligo:(open circles).

25 **FIG. 7:** DNA fragmentation in Johnson cells incubated with L-Bcl-2. Johnson cells were incubated with 4 $\mu\text{mol/L}$ of L-bcl-2 and two L-control oligos. After 3 days of incubation, DNA was extracted, electrophoresed through a 2% agarose gel and stained with ethidium bromide. Lanes 1, untreated cells; lane 2, cells treated with L-control (scrambled) oligo; lane 3, cells treated with L-control (random) oligo; lane 4, cells treated with L-bcl-2.

FIG. 8A & FIG. 8B: Apoptotic Johnson cells incubated with L-bcl-2.

FIG. 8A: Fluorescent photograph of Johnson cells dyed with the DNA-binding dye, acridine orange, after 3 days of incubation with 5 μ mol/L of L-bcl-2 (right) or without liposomal oligonucleotides ("L-OS") (left).

5 **FIG. 8B:** Apoptotic index of Johnson cells treated with 4 and 5 μ M of L-bcl-2 (dots), L-control oligo (horizontal) or empty liposomes (solid). Apoptotic index= (total no. of cells with apoptotic nuclei/total no. of cell counted) x 100%.

FIG 9A & FIG. 9B: Evaluation of cell death following irradiation of LNCaP control and bcl-2 transfected cells.

10 **FIG. 9A:** Fluorescence microscopic evaluation of cell death induction following irradiation of LNCaP control and bcl-2 transfected cells (LNCaP-bcl-2). Control untreated (upper left), LNCaP-bcl-2 cells untreated (lower left), control cells 24 hours following 20 Gy γ -irradiation (upper right), and LNCaP-bcl-2 cells 24 hours following irradiation (lower right). Cells exhibited the characteristic features of apoptosis are commonly observed in control, but 15 not LNCaP-bcl-2 cells following irradiation.

FIG. 9B: Flow cytometric analysis of cell death induction following 20 Gy of γ -irradiation in LNCaP cells. Apoptotic cells comprise approximately 30% of the LNCaP vector control cell population 8 hours following irradiation and < 5% in the irradiated LNCaP-bcl-2 cells.

20 **FIG. 10A & FIG. 10B & FIG. 10C: p53 induction, subcellular localization and transcriptional control.**

25 **FIG. 10 A:** Western blot analysis of p53 protein induction and nuclear import following γ -irradiation in LNCaP control and LNCaP-bcl-2 cells. Subconfluent cultures of control LNCaP and LNCaP-bcl-2 cells were irradiated with 20Gy. Extracts of whole cells or nuclei isolated from whole cells were prepared 2 and 4 h after irradiation. Equivalent mounts of lysates were analyzed by immunoblotting with p53 antibody (Santa Cruz). Corresponding densitometric scans indicates that the mount of p53 protein induced following irradiation is approximately equivalent in whole cell extracts from LNCaP control and LNCaP-bcl-2 cells. However, nuclear

- 10 -

accumulation of p53 protein is only observed in nuclei isolated from irradiated LNCaP control cells.

FIG. 10B: Confocal microscopic analysis of p53 subcellular localization following irradiation. LNCaP control (left) and LNCaP-bcl-2 (right) cells were irradiated with 20 Gy, 5 fixed after 4 hours, and p53 protein imaged by scanning confocal laser microscopy. Nuclear localization of p53 protein is only observed in LNCaP control cells.

FIG. 10C: Bcl-2 inhibition of transcriptional activation by wt-p53. NIH3T3 cells were transfected with the effector wild type (P53 WT) or mutant p53 (P53 MUT) plasmid (10 μ g), reporter plasmid P2mdm2-Luc (4 μ g) and β -galactosidase (β gal) expression plasmid (3 μ g) with 10 or without the bcl-2 expression vector (BCL-2) (20 μ g) using the calcium-phosphate method. Co-transfection with empty effector vector (VECTOR) served as a negative control. Data represent the fold increase in luciferase activity. Bcl-2 significantly inhibited the ability of wild type p53 protein to transactivate the mdm2 promoter (* $p \leq 0.02$).

FIG. 11A & FIG. 11B: Downregulation of bcl-2 in RKO colon cancer cells by antisense oligonucleotides and localization of p53.

FIG. 11A: Selective downregulation of bcl-2 protein in RKO colon cancer cells. 20 Western blotting of whole cell extracts (40 μ g) were analyzed by immunoblotting for bcl-2 protein. A graphic representation of the relative amount of bcl-2 protein after normalization for protein loading is shown. Treatment with antisense bcl-2 oligonucleotides, but not control oligonucleotides or empty liposomes, resulted in a reduction in the amount of bcl-2 protein.

FIG. 11B: Confocal microscopy of p53 protein in irradiated RKO cells treated with control oligonucleotides (top) or antisense-bcl-2 oligonucleotides (bottom). Significant nuclear localization of p53 protein following irradiation is observed only in antisense-bcl-2 treated RKO cells.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

5 *bcl-2* is an oncogene with tumorigenic potential due to its capacity to block programmed cell death. The present invention employs liposomal antisense oligodeoxynucleotides to inhibit the production of Bcl-2 so that tumor cells can regain the capacity to enter programmed cell death. The present invention may also be used to treat hematologic malignancies, both leukemias and lymphomas, including follicular and nonfollicular lymphomas, chronic lymphocytic leukemia, and plasma cell dyscrasias; solid tumors like those associated with breast, prostate and colon cancer; and immune disorders, which are associated with Bcl-2 expression. Such diseases would include those involving cells that overexpress, or express a 10 high level, of Bcl-2 and which also express Bax.

15 The present invention relates to antisense oligonucleotides and polynucleotides directed to portions of the *bcl-2* gene and their use in the treatment of Bcl-2 related diseases. A specific type of cancer that may be treated by the methods of the present invention is FL. Over 90% of follicular lymphoma patients have a t(14;18) translocation which results in the translocation of the *bcl-2* gene from its normal location in chromosome 18 to the immunoglobulin heavy chain gene locus on chromosome 14. In consequence, the *bcl-2* gene is under the influence of the immunoglobulin heavy chain enhancer, and the Bcl-2 protein is overexpressed. Since *bcl-2* is an oncogene with tumorigenic potential due to its capacity to block programmed cell death, a potential therapy for these follicular lymphomas is to inhibit the production of the Bcl-2 protein. 20 The present invention hopes to succeed where other approaches have failed by incorporating stable, nuclease resistant antisense oligonucleotides specific for the first open reading frame, and more specifically, for the translation initiation site of the Bcl-2 mRNA, into neutral liposomes for delivery into the cell to inhibit the production of Bcl-2 protein.

25 In particular, it is contemplated that by using these antisense molecules, either alone or in conjunction with other antisense molecules, it is possible to effectively treat FL, and possibly other Bcl-2 cancers. For example, as shown herein, liposomal *bcl-2* antisense oligonucleotides (L-*bcl-2*) inhibit the growth of FL cells and other cells which overexpress Bcl-2 protein. In the practice of the invention, the oligo- or polynucleotides themselves, or expression vectors encoding therefor, may be employed. The preferred method of delivering these nucleic acids is

via liposomes, and in particular liposomes that are made of neutral lipids. The invention, in its various embodiments, is described in greater detail, below.

A. Polynucleotides and Oligonucleotides

The term "antisense" is intended to refer to polynucleotide molecules complementary to a portion of a Bcl-2 RNA, or the DNA's corresponding thereto. "Complementary" polynucleotides are those which are capable of base-pairing according to the standard Watson-Crick complementarity rules. That is, the larger purines will base pair with the smaller pyrimidines to form combinations of guanine paired with cytosine (G:C) and adenine paired with either thymine (A:T) in the case of DNA, or adenine paired with uracil (A:U) in the case of RNA. Inclusion of less common bases such as inosine, 5-methylcytosine, 6-methyladenine, hypoxanthine and others in hybridizing sequences may offer certain advantages in some cases and does not interfere with pairing.

Targeting double-stranded (ds) DNA with polynucleotides leads to triple-helix formation; targeting RNA will lead to double-helix formation. Antisense polynucleotides, when introduced into a target cell, specifically bind to their target polynucleotide and interfere with transcription, RNA processing, transport, translation and/or stability. Antisense RNA constructs, or DNA encoding such antisense RNA's, may be employed to inhibit gene transcription or translation or both within a host cell, either *in vitro* or *in vivo*, such as within a host animal, including a human subject.

The intracellular concentration of monovalent cation is approximately 160 mM (10 mM Na⁺; 150 mM K⁺). The intracellular concentration of divalent cation is approximately 20 mM (18 mM Mg²⁺; 2 mM Ca²⁺). The intracellular protein concentration, which would serve to decrease the volume of hybridization and, therefore, increase the effective concentration of nucleic acid species, is 150 mg/ml. Constructs can be tested *in vitro* under conditions that mimic these *in vivo* conditions.

Antisense constructs may be designed to bind to the promoter and other control regions, exons, introns or even exon-intron boundaries of a gene. It is contemplated that the most effective antisense constructs for the present invention will include regions complementary to the mRNA start site. One can readily test such constructs simply by testing the constructs *in*

5 *vitro* to determine whether levels of the target protein are affected. Such testing may be performed by transferring naked polynucleic acid molecules to the cell by various techniques known in the art, or by first associating the nucleic acid molecules with one or more lipids. Similarly, detrimental non-specific inhibition of protein synthesis also can be measured by determining target cell viability *in vitro*.

10 As used herein, the terms "complementary" or "antisense" mean polynucleotides that are substantially complementary over their entire length and have very few base mismatches. For example, sequences of fifteen bases in length may be termed complementary when they have a complementary nucleotide for thirteen or fourteen positions out of fifteen. Naturally, sequences 15 which are "completely complementary" or "full length complements" will be sequences which are entirely complementary throughout their entire length and have no base mismatches.

15 Other sequences with lower degrees of homology also are contemplated. For example, an antisense construct which has limited regions of high homology, but also contains a non-homologous region (e.g., a ribozyme) could be designed. These molecules, though having less than 50% homology, would bind to target sequences under appropriate conditions.

20 The polynucleotides according to the present invention may encode a *bcl-2* gene or a portion of that gene that is sufficient to effect antisense inhibition of protein expression. The polynucleotides may be derived from genomic DNA, *i.e.*, cloned directly from the genome of a particular organism. In other embodiments, however, the polynucleotides may be complementary DNA (cDNA). cDNA is DNA prepared using messenger RNA (mRNA) as template. Thus, a cDNA does not contain any interrupted coding sequences and usually 25 contains almost exclusively the coding region(s) for the corresponding protein. In other embodiments, the antisense polynucleotide may be produced synthetically.

25 It may be advantageous to combine portions of the genomic DNA with cDNA or synthetic sequences to generate specific constructs. For example, where an intron is desired in the ultimate construct, a genomic clone will need to be used. The cDNA or a synthesized polynucleotide may provide more convenient restriction sites for the remaining portion of the construct and, therefore, would be used for the rest of the sequence.

5 The DNA and protein sequences for Bcl-2 are published in literature by Tsujimoto and Croce, (*Proc. Natl. Acad. Sci. USA*, 83:5214, 1986) (SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, & SEQ ID NO:7) which is incorporated herein by reference. It is contemplated that natural variants of Bcl-2 exist that have different sequences than those disclosed herein. Thus,

10 the present invention is not limited to use of the provided polynucleotide sequence for Bcl-2 but, rather, includes use of any naturally-occurring variants. Depending on the particular sequence of such variants, they may provide additional advantages in terms of target selectivity, *i.e.*, avoid unwanted antisense inhibition of related transcripts. The present invention also encompasses chemically synthesized mutants of these sequences.

15 10 As stated above, although the antisense sequences may be full length genomic or cDNA copies, or large fragments thereof, they also may be shorter fragments, or "oligonucleotides," defined herein as polynucleotides of 50 or less bases. Although shorter oligomers (8-20) are easier to make and increase *in vivo* accessibility, numerous other factors are involved in determining the specificity of base-pairing. For example, both binding affinity and sequence

20 15 specificity of an oligonucleotide to its complementary target increase with increasing length. It is contemplated that oligonucleotides of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45 or 50 base pairs or larger may be used. While all or part of the gene sequence may be employed in the context of antisense construction, statistically, any sequence of 17 bases long should occur only once in the human genome and, therefore, suffice to specify a unique target sequence.

25 In certain embodiments, one may wish to employ antisense constructs which include other elements, for example, those which include C-5 propyne pyrimidines. Oligonucleotides which contain C-5 propyne analogues of uridine and cytidine have been shown to bind RNA with high affinity and to be potent antisense inhibitors of gene expression (Wagner *et al.*, 1993). In the practice of the invention, derivatized polynucleotides or phosphodiester analogues such as p-ethoxy or methyl phosphonate oligonucleotides may be used. Such polynucleotides may offer certain advantages such as nuclease resistance. It is a discovery of the present inventors that p-ethoxy oligonucleotides, which are hydrophobic are also advantageous in that they are more efficiently incorporated into liposomes.

As an alternative to targeted antisense delivery, targeted ribozymes may be used. The term "ribozyme" refers to an RNA-based enzyme capable of targeting and cleaving particular base sequences in both DNA and RNA. Ribozymes can either be targeted directly to cells, in the form of RNA oligonucleotides incorporating ribozyme sequences, or introduced into the cell 5 as an expression vector encoding the desired ribozymal RNA. Ribozymes may be used and applied in much the same way as described for antisense polynucleotide. Ribozyme sequences also may be modified in much the same way as described for antisense polynucleotide. For example, one could incorporate non-Watson-Crick bases, or make mixed RNA/DNA oligonucleotides, or modify the phosphodiester backbone.

10 Alternatively, the antisense oligo- and polynucleotides of the present invention may be provided as mRNA via transcription from expression constructs that carry nucleic acids encoding the oligo- or polynucleotides. Throughout this application, the term "expression construct" is meant to include any type of genetic construct containing a nucleic acid encoding an antisense product in which part or all of the nucleic acid sequence is capable of being 15 transcribed. Typical expression vectors include bacterial plasmids or phage, such as any of the pUC or BluescriptTM plasmid series or, as discussed further below, viral vectors adapted for use in eukaryotic cells.

20 In preferred embodiments, the nucleic acid encodes an antisense oligo- or polynucleotide under transcriptional control of a promoter. A "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell, or introduced synthetic machinery, required to initiate the specific transcription of a gene. The phrase "under transcriptional control" means that the promoter is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation.

25 The term promoter will be used here to refer to a group of transcriptional control modules that are clustered around the initiation site for RNA polymerase II. Much of the thinking about how promoters are organized derives from analyses of several viral promoters, including those for the HSV thymidine kinase (tk) and SV40 early transcription units. These studies, augmented by more recent work, have shown that promoters are composed of discrete 30 functional modules, each consisting of approximately 7-20 bp of DNA, and containing one or more recognition sites for transcriptional activator or repressor proteins.

At least one module in each promoter functions to position the start site for RNA synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as the promoter for the mammalian terminal deoxynucleotidyl transferase gene and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps 5 to fix the place of initiation.

Additional promoter elements regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have recently been shown to contain functional elements downstream of the start site as well. The spacing between promoter elements frequently is flexible, so that 10 promoter function is preserved when elements are inverted or moved relative to one another. In the tk promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either cooperatively or independently to activate transcription.

The particular promoter that is employed to control the expression of a nucleic acid 15 encoding the inhibitory peptide is not believed to be important, so long as it is capable of expressing the peptide in the targeted cell. Thus, where a human cell is targeted, it is preferable to position the nucleic acid coding the inhibitory peptide adjacent to and under the control of a promoter that is active in the human cell. Generally speaking, such a promoter might include either a human or viral promoter.

20 In various embodiments, the human cytomegalovirus (CMV) immediate early gene promoter, the SV40 early promoter and the Rous sarcoma virus long terminal repeat can be used to obtain high-level expression of various proteins. The use of other viral or mammalian cellular or bacterial phage promoters which are well-known in the art to achieve expression of peptides according to the present invention is contemplated as well, provided that the levels of 25 expression are sufficient for a given purpose.

By employing a promoter with well-known properties, the level and pattern of expression of an antisense oligo- or polynucleotide can be optimized. Further, selection of a promoter that is regulated in response to specific physiologic signals can permit inducible expression of an inhibitory protein. For example, a nucleic acid under control of the human 30 PAI-1 promoter results in expression inducible by tumor necrosis factor. Tables 1 and 2 list

several elements/promoters which may be employed, in the context of the present invention, to regulate the expression of antisense constructs. This list is not intended to be exhaustive of all the possible elements involved in the promotion of expression but, merely, to be exemplary thereof.

5 Enhancers were originally detected as genetic elements that increased transcription from a promoter located at a distant position on the same molecule of DNA. This ability to act over a large distance had little precedent in classic studies of prokaryotic transcriptional regulation. Subsequent work showed that regions of DNA with enhancer activity are organized much like promoters. That is, they are composed of many individual elements, each of which binds to one
10 or more transcriptional proteins.

15 The basic distinction between enhancers and promoters is operational. An enhancer region as a whole must be able to stimulate transcription at a distance; this need not be true of a promoter region or its component elements. On the other hand, a promoter must have one or more elements that direct initiation of RNA synthesis at a particular site and in a particular orientation, whereas enhancers lack these specificities. Promoters and enhancers are often overlapping and contiguous, often seeming to have a very similar modular organization.

20 Below is a list of viral promoters, cellular promoters/enhancers and inducible promoters/enhancers that could be used in combination with the nucleic acid encoding an NF-IL6 inhibitory peptide in an expression construct (Table 1 and Table 2). Additionally any promoter/enhancer combination (as per the Eukaryotic Promoter Data Base EPDB) also could be used to drive expression of a nucleic acid according to the present invention. Use of a T3, T7 or SP6 cytoplasmic expression system is another possible embodiment. Eukaryotic cells can support cytoplasmic transcription from certain bacterial promoters if the appropriate bacterial polymerase is provided, either as part of the delivery complex or as an additional genetic
25 expression construct.

TABLE 1

| PROMOTER | |
|--------------------------------|--|
| Immunoglobulin Heavy Chain | c-HA-ras |
| Immunoglobulin Light Chain | Insulin |
| T-Cell Receptor | Neural Cell Adhesion Molecule (NCAM) |
| HLA DQ α and DQ β | α 1-Antitrypsin |
| β -Interferon | H2B (TH2B) Histone |
| Interleukin-2 | Mouse or Type I Collagen |
| Interleukin-2 Receptor | Glucose-Regulated Proteins (GRP94 and GRP78) |
| MHC Class II 5 | Rat Growth Hormone |
| MHC Class II HLA-DR α | Human Serum Amyloid A (SAA) |
| β -Actin | Troponin I (TN I) |
| Muscle Creatine Kinase | Platelet-Derived Growth Factor |
| Prealbumin (Transthyretin) | Duchenne Muscular Dystrophy |
| Elastase <i>I</i> | SV40 |
| Metallothionein | Polyoma |
| Collagenase | Retroviruses |
| Albumin Gene | Papilloma Virus |
| α -Fetoprotein | Hepatitis B Virus |
| τ -Globin | Human Immunodeficiency Virus |
| β -Globin | Cytomegalovirus |
| c-fos | Gibbon Ape Leukemia Virus |

TABLE 2

| Element | Inducer |
|------------------------------------|--|
| MT II | Phorbol Ester (TPA) Heavy metals |
| MMTV (mouse mammary tumor virus) | Glucocorticoids |
| β-Interferon | poly(rI)X poly(rC) |
| Adenovirus 5 <u>E2</u> | Ela |
| c-jun | Phorbol Ester (TPA), H ₂ O ₂ |
| Collagenase | Phorbol Ester (TPA) |
| Stromelysin | Phorbol Ester (TPA), IL-1 |
| SV40 | Phorbol Ester (TPA) |
| Murine MX Gene | Interferon, Newcastle Disease Virus |
| GRP78 Gene | A23187 |
| α-2-Macroglobulin | IL-6 |
| Vimentin | Serum |
| MHC Class I Gene H-2kB | Interferon |
| HSP70 | Ela, SV40 Large T Antigen |
| Proliferin | Phorbol Ester (TPA) |
| Tumor Necrosis Factor | PHA |
| Thyroid Stimulating Hormone α Gene | Thyroid Hormone |

In certain embodiments of this invention, the delivery of a nucleic acid in a cell may be identified *in vitro* or *in vivo* by including a marker in the expression construct. The marker would result in an identifiable change to the transfected cell permitting easy identification of expression. Enzymes such as herpes simplex virus thymidine kinase (*tk*) (eukaryotic) or chloramphenicol acetyltransferase (CAT) (prokaryotic) may be employed.

One also may include a polyadenylation signal to effect proper polyadenylation of the transcript. The nature of the polyadenylation signal is not believed to be crucial to the successful practice of the invention, and any such sequence may be employed. Examples include the SV40, globin or adenovirus polyadenylation signals. Also contemplated as an

element of the expression cassette is a terminator. These elements can serve to enhance message levels and to minimize read through from the cassette into other sequences.

B. Lipid Formulations

In a preferred embodiment of the invention, the antisense oligo- or polynucleotides and expression vectors may be associated with a lipid. A polynucleotide associated with a lipid may be encapsulated in the aqueous interior of a liposome, interspersed within the lipid bilayer of a liposome, attached to a liposome via a linking molecule that is associated with both the liposome and the polynucleotide, entrapped in a liposome, complexed with a liposome, dispersed in a solution containing a lipid, mixed with a lipid, combined with a lipid, contained as a suspension in a lipid, contained or complexed with a micelle, or otherwise associated with a lipid. The lipid or lipid/oligonucleotide associated compositions of the present invention are not limited to any particular structure in solution. For example, they may be present in a bilayer structure, as micelles, or with a "collapsed" structure. They may also simply be interspersed in a solution, possibly forming aggregates which are not uniform in either size or shape. The lipids may be in the form of liposomes known in the art as multiple-lamellar vesicles, small or large unilamellar vesicles or other types of vesicles known in the art.

Lipids are fatty substances which may be naturally occurring or synthetic lipids. For example, lipids include the fatty droplets that naturally occur in the cytoplasm as well as the class of compounds which are well known to those of skill in the art which contain long-chain aliphatic hydrocarbons and their derivatives, such as fatty acids, alcohols, amines, amino alcohols, and aldehydes. An example is the lipid dioleoylphosphatidylcholine.

"Liposome" is a generic term encompassing a variety of single and multilamellar lipid vehicles formed by the generation of enclosed lipid bilayers or aggregates. Liposomes may be characterized as having vesicular structures with a phospholipid bilayer membrane and an inner aqueous medium. Multilamellar liposomes have multiple lipid layers separated by aqueous medium. They form spontaneously when phospholipids are suspended in an excess of aqueous solution. The lipid components undergo self-rearrangement before the formation of closed structures and entrap water and dissolved solutes between the lipid bilayers (Ghosh and Bachhawat, *In: Wu G. Wu C ed.. Liver diseases, targeted diagnosis and therapy using specific*

receptors and ligands, New York: Marel Dekker, pp. 87-104, 1991). However, the present invention also encompasses compositions that have different structures in solution than the normal vesicular structure. For example, the lipids may assume a micellar structure or merely exist as nonuniform aggregates of lipid molecules. Also contemplated are 5 lipofectamine-nucleic acid complexes.

Liposome-mediated polynucleotide delivery and expression of foreign DNA *in vitro* has been very successful. Wong *et al.*, (*Gene*, 10:87-94, 1980) demonstrated the feasibility of liposome-mediated delivery and expression of foreign DNA in cultured chick embryo, HeLa and hepatoma cells. Nicolau *et al.*, (*Methods Enzymol.*, 149:157-176, 1987) accomplished 10 successful liposome-mediated gene transfer in rats after intravenous injection.

In certain embodiments of the invention, the lipid may be associated with a hemagglutinating virus (HVJ). This has been shown to facilitate fusion with the cell membrane and promote cell entry of liposome-encapsulated DNA (Kaneda *et al.*, *Science*, 243:375-378, 1989). In other embodiments, the lipid may be complexed or employed in conjunction with 15 nuclear non-histone chromosomal proteins (HMG-1) (Kato *et al.*, *J. Biol. Chem.*, 266:3361-3364, 1991). In yet further embodiments, the lipid may be complexed or employed in conjunction with both HVJ and HMG-1. In that such expression vectors have been successfully employed in transfer and expression of a polynucleotide *in vitro* and *in vivo*, then they are applicable for the present invention. Where a bacterial promoter is employed in the DNA 20 construct, it also will be desirable to include within the liposome an appropriate bacterial polymerase.

Phospholipids are used for preparing the liposomes according to the present invention and can carry a net positive charge, a net negative charge or are neutral, with neutral phospholipids being the most preferred. Diacetyl phosphate is often employed to confer a 25 negative charge on a liposome, and stearylamine to confer a positive charge on a liposome.

Lipids suitable for use according to the present invention can be obtained from commercial sources. For example, dimyristyl phosphatidylcholine ("DMPC") can be obtained from Sigma Chemical Co., dicetyl phosphate ("DCP") is obtained from K & K Laboratories (Plainview, NY); cholesterol ("Chol") is obtained from Calbiochem-Behring; dimyristoyl 30 phosphatidylglycerol ("DMPG") and other lipids may be obtained from Avanti Polar Lipids,

Inc. (Birmingham, Ala.). Stock solutions of lipids in chloroform or chloroform/methanol can be stored at about -20°C. Preferably, chloroform is used as the only solvent since it is more readily evaporated than methanol.

5 Phospholipids from natural sources, such as egg or soybean phosphatidylcholine, brain phosphatidic acid, brain or plant phosphatidylinositol, heart cardiolipin and plant or bacterial phosphatidylethanolamine are preferably not used as the primary phosphatide, *i.e.*, constituting 50% or more of the total phosphatide composition, because of the instability and leakiness of the resulting liposomes.

10 Liposomes used according to the present invention can be made by different methods. The size of the liposomes varies depending on the method of synthesis. A liposome suspended in an aqueous solution is generally in the shape of a spherical vesicle, having one or more concentric layers of lipid bilayer molecules. Each layer consists of a parallel array of molecules represented by the formula XY, wherein X is a hydrophilic moiety and Y is a hydrophobic moiety. In aqueous suspension, the concentric layers are arranged such that the hydrophilic 15 moieties tend to remain in contact with an aqueous phase and the hydrophobic regions tend to self-associate. For example, when aqueous phases are present both within and without the liposome, the lipid molecules may form a bilayer, known as a lamella, of the arrangement XY-YX. Aggregates of lipids may form when the hydrophilic and hydrophobic parts of more than one lipid molecule become associated with each other. The size and shape of these 20 aggregates will depend upon many different variables, such as the nature of the solvent and the presence of other compounds in the solution.

25 Liposomes within the scope of the present invention can be prepared in accordance with known laboratory techniques. In one preferred embodiment, liposomes are prepared by mixing liposomal lipids, in a solvent in a container, *e.g.*, a glass, pear-shaped flask. The container should have a volume ten-times greater than the volume of the expected suspension of 30 liposomes. Using a rotary evaporator, the solvent is removed at approximately 40°C under negative pressure. The solvent normally is removed within about 5 min. to 2 hours, depending on the desired volume of the liposomes. The composition can be dried further in a desiccator under vacuum. The dried lipids generally are discarded after about 1 week because of a tendency to deteriorate with time.

Dried lipids can be hydrated at approximately 25-50 mM phospholipid in sterile, pyrogen-free water by shaking until all the lipid film is resuspended. The aqueous liposomes can be then separated into aliquots, each placed in a vial, lyophilized and sealed under vacuum.

5 In the alternative, liposomes can be prepared in accordance with other known laboratory procedures: the method of Bangham *et al.*, (*J. Mol. Biol.*, 13:238, 1965), the contents of which are incorporated herein by reference; the method of Gregoriadis, as described in *DRUG CARRIERS IN BIOLOGY AND MEDICINE*, G. Gregoriadis ed. (1979) pp. 287-341, the contents of which are incorporated herein by reference; the method of Deamer and Uster, (LIPOSOMES, M. Ostro ed. 1983), the contents of which are incorporated by reference; and the 10 reverse-phase evaporation method as described by Szoka and Papahadjopoulos, (*Proc. Nat'l Acad. Sci. U.S.A.* 75:4194-98 1978). The aforementioned methods differ in their respective abilities to entrap aqueous material and their respective aqueous space-to-lipid ratios.

15 The dried lipids or lyophilized liposomes prepared as described above may be dehydrated and reconstituted in a solution of inhibitory peptide and diluted to an appropriate concentration with an suitable solvent, *e.g.*, DPBS. The mixture is then vigorously shaken in a vortex mixer. Unencapsulated nucleic acid is removed by centrifugation at 29,000 \times g and the 20 liposomal pellets washed. The washed liposomes are resuspended at an appropriate total phospholipid concentration, *e.g.*, about 50-200 mM. The amount of nucleic acid encapsulated can be determined in accordance with standard methods. After determination of the amount of nucleic acid encapsulated in the liposome preparation, the liposomes may be diluted to appropriate concentrations and stored at 4°C until use.

25 P-ethoxy oligonucleotides, nuclease resistant analogues of phosphodiesters, are preferred because they are stable in serum and effectively transported into the cellular cytoplasm. In a preferred embodiment, the lipid dioleoylphosphatidylchoine is employed. However other lipids such as other phosphatidylcholines, phosphatidylglycerols, and phosphatidylethanolamines may also be useful. Nuclease-resistant oligonucleotides are mixed 30 with lipids in the presence of excess t-butanol. The mixture is vortexed before being frozen in an acetone/dry ice bath. The frozen mixture is lyophilized and hydrated with Hepes-buffered saline (1 mM Hepes, 10 mM NaCl, pH 7.5) overnight, and then the liposomes are sonicated in a bath type sonicator for 10 to 15 min. The size of the liposomal-oligonucleotides typically

ranges between 200-300 nm in diameter as determined by the submicron particle sizer autodilute model 370 (Nicomp, Santa Barbara, CA).

C. Alternative Delivery Systems

Retroviruses. The retroviruses are a group of single-stranded RNA viruses characterized by an ability to convert their RNA to double-stranded DNA in infected cells by a process of reverse-transcription (Coffin, *In: Virology*, Fields *et al.* (eds.), New York: Raven Press, pp. 1437-1500, 1990). The resulting DNA then stably integrates into cellular chromosomes as a provirus and directs synthesis of viral proteins. The integration results in the retention of the viral gene sequences in the recipient cell and its descendants. The retroviral genome contains three genes - *gag*, *pol*, and *env* - that code for capsid proteins, polymerase enzyme, and envelope components, respectively. A sequence found upstream from the *gag* gene, termed Ψ , functions as a signal for packaging of the genome into virions. Two long terminal repeat (LTR) sequences are present at the 5' and 3' ends of the viral genome. These contain strong promoter and enhancer sequences and are also required for integration in the host cell genome (Coffin, 1990).

In order to construct a retroviral vector, a nucleic acid encoding a Bcl-2 antisense construct is inserted into the viral genome in the place of certain viral sequences to produce a virus that is replication-defective. In order to produce virions, a packaging cell line containing the *gag*, *pol* and *env* genes but without the LTR and Ψ components is constructed (Mann *et al.*, *Cell*, 33:153-159, 1983). When a recombinant plasmid containing an inserted DNA, together with the retroviral LTR and Ψ sequences, is introduced into this cell line (by calcium phosphate precipitation for example), the Ψ sequence allows the RNA transcript of the recombinant plasmid to be packaged into viral particles, which are then secreted into the culture media (Nicolas and Rubenstein, *In: Vectors: A survey of molecular cloning vectors and their uses*, Rodriguez and Denhardt, (eds.), Stoneham: Butterworth, pp. 494-513, 1988; Temin, *In: Gene Transfer*, Kucherlapati (ed.), New York: Plenum Press, pp. 149-188, 1986; Mann *et al.*, 1983). The media containing the recombinant retroviruses is then collected, optionally concentrated, and used for gene transfer. Retroviral vectors are able to infect a broad variety of cell types. However, integration and stable expression require the division of host cells (Paskind *et al.*, *Virology*, 67:242-248, 1975).

5 Adenoviruses: Human adenoviruses are double-stranded DNA tumor viruses with genome sizes of approximate 36 kB. As a model system for eukaryotic gene expression, adenoviruses have been widely studied and well characterized, which makes them an attractive system for development of adenovirus as a gene transfer system. This group of viruses is easy
10 to grow and manipulate, and they exhibit a broad host range *in vitro* and *in vivo*. In lytically infected cells, adenoviruses are capable of shutting off host protein synthesis, directing cellular machineries to synthesize large quantities of viral proteins, and producing copious amounts of virus.

10 The E1 region of the genome includes E1A and E1B which encode proteins responsible for transcription regulation of the viral genome, as well as a few cellular genes. E2 expression, including E2A and E2B, allows synthesis of viral replicative functions, *e.g.* DNA-binding protein, DNA polymerase, and a terminal protein that primes replication. E3 gene products prevent cytolysis by cytotoxic T cells and tumor necrosis factor and appear to be important for viral propagation. Functions associated with the E4 proteins include DNA replication, late gene
15 expression, and host cell shutoff. The late gene products include most of the virion capsid proteins, and these are expressed only after most of the processing of a single primary transcript from the major late promoter has occurred. The major late promoter (MLP) exhibits high efficiency during the late phase of the infection (Stratford-Perricaudet and Perricaudet, pp. 51-61, *In: Human Gene Transfer*, Eds, O. Cohen-Haguenauer and M. Boiron Editions John Libbey
20 Exrotex, France, 1991).

25 As only a small portion of the viral genome appears to be required *in cis* adenovirus-derived vectors offer excellent potential for the substitution of large DNA fragments when used in connection with cell lines such as 293 cells. Ad5-transformed human embryonic kidney cell lines (Graham *et al.*, *J. Gen. Virol.*, 36:59-72, 1977) have been developed to provide the essential viral proteins *in trans*.

Particular advantages of an adenovirus system for delivering foreign proteins to a cell include (i) the ability to substitute relatively large pieces of viral DNA by foreign DNA; (ii) the structural stability of recombinant adenoviruses; (iii) the safety of adenoviral administration to humans; and (iv) lack of any known association of adenoviral infection with cancer or

malignancies; (v) the ability to obtain high titers of the recombinant virus; and (vi) the high infectivity of adenovirus.

Further advantages of adenovirus vectors over retroviruses include the higher levels of gene expression. Additionally, adenovirus replication is independent of host gene replication, 5 unlike retroviral sequences. Because adenovirus transforming genes in the E1 region can be readily deleted and still provide efficient expression vectors, oncogenic risk from adenovirus vectors is thought to be negligible (Grunhaus & Horwitz, *Seminar in Virology*, 3:237-252, 1992).

In general, adenovirus gene transfer systems are based upon recombinant, engineered adenovirus which is rendered replication-incompetent by deletion of a portion of its genome, 10 such as E1, and yet still retains its competency for infection. Sequences encoding relatively large foreign proteins can be expressed when additional deletions are made in the adenovirus genome. For example, adenoviruses deleted in both E1 and E3 regions are capable of carrying up to 10 kB of foreign DNA and can be grown to high titers in 293 cells (Stratford-Perricaudet and Perricaudet, 1991). Surprisingly persistent expression of transgenes following adenoviral 15 infection has also been reported.

Other Viral Vectors as Expression Constructs. Other viral vectors may be employed as expression constructs in the present invention. Vectors derived from viruses such as vaccinia virus (Ridgeway, Stoneham: Butterworth, pp. 467-492, 1988; Baichwal and Sugden, *In: Kucherlapati R, (ed.) Gene transfer*. New York: Plenum Press, pp. 117-148, 1986; Coupar *et al.*, 20 *Gene*, 68:1-10, 1988) adeno-associated virus (AAV) (Ridgeway, 1988; Baichwal and Sugden, 1986; Hermonat and Muzycska, *Proc. Nat'l Acad. Sci. USA*, 81:6466-6470, 1984) and herpes viruses may be employed. They offer several attractive features for various mammalian cells (Friedman *et al.*, *Genes Devel.* 3:1314, 1989; Ridgeway, 1988; Baichwal and Sugden, 1986; Coupar *et al.*, 1988; Horwich *et al.*, *J. Virol.* 64:642-650, 1990).

25 With the recent recognition of defective hepatitis B viruses, new insight was gained into the structure-function relationship of different viral sequences. *in vitro* studies showed that the virus could retain the ability for helper-dependent packaging and reverse transcription despite the deletion of up to 80% of its genome (Horwich *et al.*, 1990). This suggested that large portions of the genome could be replaced with foreign genetic material. The hepatotropism and 30 persistence (integration) were particularly attractive properties for liver-directed gene transfer.

Chang *et al.* (*Hepatology*, 14:134A, 1991) recently introduced the chloramphenicol acetyltransferase (CAT) gene into duck hepatitis B virus genome in the place of the polymerase, surface, and pre-surface coding sequences. It was cotransfected with wild-type virus into an avian hepatoma cell line. Culture media containing high titers of the recombinant virus were used to infect primary duckling hepatocytes. Stable CAT gene expression was detected for at least 24 days after transfection (Chang *et al.*, 1991).

Non-viral Methods. Several non-viral methods for the transfer of expression vectors into cultured mammalian cells also are contemplated by the present invention. These include calcium phosphate precipitation (Graham and van der Eb, *Virology*, 52:456-467, 1973; Chen and Okayama, *Mol. Cell Biol.*, 7:2745-2752, 1987; Rippe *et al.*, *Mol. Cell Biol.*, 10:689-695, 1990) DEAE-dextran (Gopal, *Mol. Cell Biol.*, 5:1188-1190, 1985), electroporation (Tur-Kaspa *et al.*, *Mol. Cell Biol.*, 6:716-718, 1986; Potter *et al.*, *Proc. Nat'l Acad. Sci. USA*, 81:7161-7165, 1984), direct microinjection (Harland and Weintraub, *J. Cell Biol.*, 101:1094-1099, 1985), DNA-loaded liposomes (Nicolau and Sene, *Biochem. Biophys. Acta*, 721:185-190, 1982; Fraley *et al.*, *Proc. Nat'l Acad. Sci. USA*, 76:3348-3352, 1979) and lipofectamine-DNA complexes, cell sonication (Fecheimer *et. al.*, *Proc. Nat'l Acad. Sci. USA*, 76:3348-52, 1979), gene bombardment using high velocity microprojectiles (Yang *et al.*, *Proc. Nat'l Acad. Sci. USA*, 87:9568-9572, 1990), polycations and receptor-mediated transfection (Wu and Wu, *J. Biol. Chem.*, 262:4429-4432, 1987; Wu and Wu, *Biochemistry*, 27:887-892, 1988). Some of these techniques may be successfully adapted for *in vivo* or *ex vivo* use.

In one embodiment of the invention, the expression construct may simply consist of naked recombinant vector. Transfer of the construct may be performed by any of the methods mentioned above which physically or chemically permeabilize the cell membrane. For example, Dubensky *et al.*, (*Proc. Nat'l Acad. Sci. USA*, 81:7529-7533, 1984) successfully injected polyomavirus DNA in the form of CaPO₄ precipitates into liver and spleen of adult and newborn mice demonstrating active viral replication and acute infection. Benvenisty and Neshif (*Proc. Nat'l Acad. Sci. USA*, 83:9551, 1986) also demonstrated that direct intraperitoneal injection of CaPO₄ precipitated plasmids results in expression of the transfected genes. It is envisioned that DNA encoding an Bcl-2 construct may also be transferred in a similar manner *in vivo*.

Another embodiment of the invention for transferring a naked DNA expression vector into cells may involve particle bombardment. This method depends on the ability to accelerate DNA coated microprojectiles to a high velocity allowing them to pierce cell membranes and enter cells without killing them (Klein *et al.*, *Nature*, 327:70-73, 1987). Several devices for accelerating small particles have been developed. One such device relies on a high voltage discharge to generate an electrical current, which in turn provides the motive force (Yang *et al.*, 1990). The microprojectiles used have consisted of biologically inert substances such as tungsten or gold beads.

Selected organs including the liver, skin, and muscle tissue of rats and mice have been bombarded *in vivo* (Yang *et al.*, 1990; Zelenin *et al.*, *FEBS Lett.*, 280:94-96, 1991). This may require surgical exposure of the tissue or cells, to eliminate any intervening tissue between the gun and the target organ. DNA encoding a Bcl-2 construct may be delivered via this method.

D. Pharmaceutical Compositions and Routes of Administration

Where clinical application of liposomes containing antisense oligo- or polynucleotides or expression vectors is undertaken, it will be necessary to prepare the liposome complex as a pharmaceutical composition appropriate for the intended application. Generally, this will entail preparing a pharmaceutical composition that is essentially free of pyrogens, as well as any other impurities that could be harmful to humans or animals. One also will generally desire to employ appropriate buffers to render the complex stable and allow for uptake by target cells.

Aqueous compositions of the present invention comprise an effective amount of the antisense expression vector encapsulated in a liposome as discussed above, further dispersed in pharmaceutically acceptable carrier or aqueous medium. Such compositions also are referred to as inocula. The phrases "pharmaceutically" or "pharmacologically acceptable" refer to compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, or a human, as appropriate.

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with

the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients also can be incorporated into the compositions.

5 Solutions of therapeutic compositions can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions also can be prepared in glycerol, liquid polyethylene glycols, mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

10 The therapeutic compositions of the present invention are advantageously administered in the form of injectable compositions either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. These preparations also may be emulsified. A typical composition for such purpose comprises a pharmaceutically acceptable carrier. For instance, the composition may contain 10 mg, 25 mg, 50 mg or up to about 100 mg of human serum albumin per milliliter of phosphate buffered saline. Other pharmaceutically acceptable carriers include aqueous solutions, non-toxic excipients, including salts, preservatives, buffers and the like.

15 Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oil and injectable organic esters such as ethyloleate. Aqueous carriers include water, alcoholic/aqueous solutions, saline solutions, parenteral vehicles such as sodium chloride, Ringer's dextrose, *etc.* Intravenous vehicles include fluid and nutrient replenishers. Preservatives include antimicrobial agents, anti-oxidants, chelating agents and inert gases. The 20 pH and exact concentration of the various components the pharmaceutical composition are adjusted according to well known parameters.

25 Additional formulations are suitable for oral administration. Oral formulations include such typical excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. The compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders. When the route is topical, the form may be a cream, ointment, salve or spray.

The therapeutic compositions of the present invention may include classic pharmaceutical preparations. Administration of therapeutic compositions according to the

present invention will be via any common route so long as the target tissue is available via that route. This includes oral, nasal, buccal, rectal, vaginal or topical. Topical administration would be particularly advantageous for the treatment of skin cancers, to prevent chemotherapy-induced alopecia or other dermal hyperproliferative disorder. Alternatively, administration may be by 5 orthotopic, intradermal subcutaneous, intramuscular, intraperitoneal or intravenous injection. Such compositions would normally be administered as pharmaceutically acceptable compositions that include physiologically acceptable carriers, buffers or other excipients. For treatment of conditions of the lungs, the preferred route is aerosol delivery to the lung. Volume of the aerosol is between about 0.01 ml and 0.5 ml. Similarly, a preferred method for treatment 10 of colon-associated disease would be via enema. Volume of the enema is between about 1 ml and 100 ml.

An effective amount of the therapeutic composition is determined based on the intended goal. The term "unit dose" or "dosage" refers to physically discrete units suitable for use in a subject, each unit containing a predetermined quantity of the therapeutic composition calculated 15 to produce the desired responses, discussed above, in association with its administration, *i.e.*, the appropriate route and treatment regimen. The quantity to be administered, both according to number of treatments and unit dose, depends on the protection desired.

Precise amounts of the therapeutic composition also depend on the judgment of the practitioner and are peculiar to each individual. Factors affecting the dose include the physical 20 and clinical state of the patient, the route of administration, the intended goal of treatment (alleviation of symptoms *versus* cure) and the potency, stability and toxicity of the particular therapeutic substance. For the instant application, it is envisioned that the amount of therapeutic peptide included in a unit dose will range from about 5-30 mg of polynucleotide.

E. Examples

25 EXAMPLE 1: Synthesis of Oligonucleotides.

Nuclease-resistant p-ethoxy oligonucleotides, non-ionic phosphodiester analogs, were purchased from Oligo Therapeutics (Willsonville, OR). An oligonucleotide specific for the translation initiation site of human Bcl-2 mRNA : ^{5'}CAGCGTGCGCCATCCTTC^{3'} (SEQ ID NO:1) was used as antisense oligonucleotide. Two different control oligonucleotides were

used: ^{5'}ACGGTCCGCCACTCCTCCCC^{3'} (SEQ ID NO:2) (scrambled version of Bcl-2 antisense oligonucleotide) and the random sequence ^{5'}CTGAAGGGCTTCTTCC^{3'} (SEQ ID NO:3).

EXAMPLE 2: Preparation of liposomal oligonucleotides (L-OS)

5 P-ethoxy-oligonucleotides dissolved in distilled water were added to phospholipids (Avanti Polar Lipids, Alabaster, AL) in the presence of excess tert-butanol ($\geq 95\%$ by volume). The mixture was frozen in a dry ice/acetone bath, lyophilized overnight and finally hydrated with HEPES buffered saline (1 mmol/L Hepes and 10 mmol/L NaCl) at a final oligonucleotide concentration of 0.1 mmol/L. Liposomal oligonucleotides (L-OS) were sonicated for 12
10 minutes in a bath-type sonicator. The average diameter of the particles was $100\text{ nm} \pm 50\text{ nm}$ as determined in a NICOMP particle sizing system (Santa Barbara, CA).

EXAMPLE 3: Oligonucleotide Inhibition of Protein Expression

Cell Lines

15 Johnson cells, a human transformed FL cell line bearing the t(14;18) translocation which overexpresses Bcl-2 protein, were used. Raji and Jurkat cells, a human Burkitt lymphoma cell line and a human acute T cell leukemia cell line, respectively, were also used. Both lines express the Bcl-2 protein but they lack the t(14;18) translocation. Daudi cells, a human Burkitt lymphoma cell line which does not express the Bcl-2 protein, was used as a negative control cell line. Johnson, Raji and Jurkat cells were grown in RPMI 1640 media (GIBCO, Grand
20 Island, NY) supplemented with 10% heat-inactivated fetal bovine serum (FBS). Daudi cells were grown in RPMI 1640 media supplemented with 20% heat-inactivated FBS.

Delivery of L-OS to cells

25 Ten thousand cells/well were seeded in a 96-well plate in 0.1 mL of the respective medium. Cells were incubated with L-OS at final concentration of 2 to 8 $\mu\text{mol/L}$ at 37°C in a 5% CO_2 incubator. Each study was done in triplicate and repeated at least 3 times.

Cell viability assay

The viability of the neoplastic cells was measured by the alamarBlue dye (Alamar, Sacramento, CA). After 5 days of incubation with L-OS, 40 μ L of cells/well were aliquoted and added to 140 μ L of fresh medium. Twenty μ L of alamarBlue dye were added to each well. 5 After incubation for 12 hours at 37°C, the plates were read directly on a microplate reader (Molecular Devices, CA) at 570 and 595 nm. The difference in absorbance between 570 and 595 nm was taken as the overall absorbance value of the cells. All experiments were analyzed by t-test in which the viabilities of the cells treated with the L-OS were compared with those of the untreated controls.

10 *Western Blots for Bcl-2 and Bax protein*

One hundred thousand cells/well were seeded in a 6-well plate in 3 mL of the respective medium, treated with 2, 3 and 4 μ mol/L of L-OS and incubated at 37°C. Untreated cells were also maintained in culture. Samples were removed on day 3 after the addition of the L-OS and lysed in 100 μ L of lysis buffer (1% Triton, 150 mmol/L NaCl and 25 mmol/L Tris pH 7.4) at 15 0°C for 30 minutes. After centrifugation at 12,000 \times g for 10 minutes, the supernatants were recovered and normalized for total protein content (5 μ g/lane of Johnson cells lysate and 20 μ g/lane of Jurkat cells lysate for Bcl-2 analysis, and 25 μ g/lane of Johnson cells for Bax analysis). The lysates were mixed with sample buffer containing 1% of sodium dodecyl sulfate (SDS) and 1% 2- β -mercaptoethanol and boiled for 5 minutes. SDS-PAGE was run on 10% 20 polyacrylamide gels, electrophoretically transferred to nitrocellulose membranes and blocked in 10% non-fat dry milk. Filters were cut in 2 portions: the bottom section was incubated with the 6C8 hamster anti-human-Bcl-2 monoclonal antibody or rabbit anti-human-Bax polyclonal antibody (Hockenberry *et. al*, *Nature*, 348:334, 1990), and the top section was incubated with mouse anti-actin monoclonal antibody (Amersham) at room temperature for 2 hours. After 25 washing and incubation with a peroxidase-labeled antihamster (Kirkegaard & Perry laboratories), antirabbit (Santa Cruz) or antimouse (Amersham) secondary antibody, blots were developed by enhanced chemiluminescence system (ECL, Amersham). To estimate the inhibition of Bcl-2 protein and the ratio of Bcl-2/Bax proteins, densitometric scans were performed on western blots on a Gilford Response Gel Scanner (CIBA Corning, Medfield, MA). Area integration of

absorbance peaks at 500 nm was used to determine the ratio of Bcl-2:Actin and Bcl-2:Bax proteins.

Analysis of apoptosis

To qualitatively determine the internucleosomal DNA cleavage associated with apoptosis, DNA fragmentation analysis by agarose gel electrophoresis was performed (Duke *et. al* In: Coligan *et. al* (eds) Current protocols in immunology, vol 1., New York: John Wiley & sons, p 3.17.1, 1991). In brief, 1×10^6 Johnson cells were cultured in 10 mL of medium in a 75 sq. cm tissue culture flask, treated with 4 $\mu\text{mol/L}$ of L-OS, and incubated at 37°C. Untreated cells were also maintained in culture. Samples were removed on day 3 after treatment, washed in PBS and pellet by centrifuging 10 minutes at 200 x g. The pellets were lysed in 0.5 mL of lysis buffer (10 mmol/L Tris pH 7.4, 1 mmol/L EDTA pH 8.0 and 0.2% Triton X-100) and fragmented DNA were separated from intact chromatin by microcentrifuging for 10 minutes at 13,000 x g. DNA of the supernatants was precipitated in 0.7 mL ice-cold isopropanol overnight at -20°C, resuspended in 30 μL of TE buffer (10 mmol/L Tris pH 7.4, 1 mmol/L EDTA pH 8.0) and incubated in 10 μL of RNase (10 $\mu\text{g/mL}$ solution) at 60°C for 1 hour. Twenty μL of sample per well were electrophoresed on a 2% agarose gel and visualized by ethidium bromide staining.

To quantitatively determine the extent of apoptosis, the fluorescent DNA-binding dye acridine orange (Duke *et. al*, 1991) was used. Briefly, 5 $\mu\text{mol/L}$ final concentration of L-OS were added to 1×10^5 cells/well plated in a 24-well plate in 1 mL of medium. After 3 days of incubation at 37°C, the cells were washed with PBS and resuspended at 1×10^6 cells/mL. Twenty-five μL of cell suspension were mixed with 1 μL of acridine orange dye (100 $\mu\text{g/mL}$, Sigma Chemicals, St. Louis, MO) and observed in a fluorescent microscope. The percentage of apoptotic cells (apoptotic index) was obtained using a hemocytometer. Apoptotic index= (total no. of cells with apoptotic nuclei/total no. of cells counted) x 100%.

Effect of L-bcl-2-antisense oligonucleotides ("L-bcl-2") on lymphoma cell growth

Five days after the addition of L-bcl-2 to the cells, the viability of tumoral cells was assessed. Cell growth was inhibited in a concentration-dependent manner in Johnson cells, which bear the t(14;18) translocation and express very high levels of Bcl-2. A concentration of

6 μ mol/L L-bcl-2 resulted in complete loss of viability of Johnson cells within 5 days (FIG. 1). Similar dose-dependent decrease in cell viabilities could be seen in three separate studies. In contrast, after treatment with 6 μ mol/L()L-bcl-2, the viabilities of Jurkat, Raji and Daudi cells decreased by only 23%, 0% and 35%, respectively (FIG. 1).

5 *Effect of liposomal control oligonucleotides (L-control oligos) on lymphoma cell growth*

Two control oligonucleotides were used to determine the specificity of the inhibition observed. When L-control oligos or empty liposomes were added to Johnson cells, cell growth inhibition was not observed. Jurkat, Raji and Daudi cells were also treated with L-control oligos and empty liposomes. Non-specific toxicity could be observed when greater than 6 μ mol/L of 10 L-OS were used, but not with empty liposomes (FIG. 2).

Selective inhibition of Bcl-2 protein by L-bcl-2-antisense oligonucleotides

In order to determine whether the cytotoxic effect of L-bcl-2 in Johnson cells was caused by a decrease in Bcl-2 protein, the Bcl-2 protein expression in these cells after treatment with L-bcl-2 as well as the effects of L-bcl-2 in the other cell lines which overexpress Bcl-2 15 protein (FIG. 3). was also determined.

When Johnson cells were treated with 2, 3 and 4 mmol/L of L-bcl-2, the ratios of Bcl-2/Actin protein were inhibited by 28, 57 and 64%, respectively. Bcl-2 protein expression was not inhibited in cells treated with the same doses of L-control oligos.

When Jurkat cells were treated with 3 and 4 mmol/L of L-bcl-2, the ratios of Bcl-2/Actin protein were inhibited by 44% and 50%, respectively. Bcl-2 protein was not significantly inhibited when the same doses of L-control oligos were used (FIG. 4).

The ratios of Bcl-2/Bax protein in Johnson and Jurkat cells before and after treatment with L-bcl-2 and L-control oligos were also determined. Johnson, Raji and Daudi cells expressed Bax protein but Jurkat cells did not express (FIG. 5). When Johnson cells were treated with 2, 3 and 4 mmol/L of L-bcl-2, the ratio of Bax/Actin was not modified, but the ratio of Bcl-2/Bax decreased by 10%, 40% and 50%, respectively. These protein ratios were unmodified after treatment with the same doses of L-control oligos (FIG. 6).

Inhibition of Bcl-2 protein leads to apoptosis in the FL cells

Whether the growth inhibitory effects seen in Johnson cells may be related to induction of apoptosis was also studied. After 3 days of incubation with L-bcl-2, the typical internucleosomal DNA degradation pattern was observed, whereas cells incubated with L-control oligos did not show the internucleosomal DNA pattern (FIG. 7). Subsequently, the quantity of apoptosis by acridine orange was assessed. After 3 days of exposure to 4 and 5 $\mu\text{mol/L}$ of L-bcl-2, apoptotic cells were evident. The apoptotic index of untreated Johnson cells was 3% while that of Johnson cells treated with 4 and 5 $\mu\text{mol/L}$ of L-bcl-2 were 43% and 61%, respectively. Significant increase in apoptotic index was not seen in cells treated with 10 liposomal control oligonucleotides or empty liposomes (FIG. 8).

L-bcl-2 selectively downregulates the expression of Bcl-2 protein and cell growth in a dose-dependent manner

The inhibition in cell growth was seen only in the FL cell line which bears the t(14;18) translocation, while cell growth inhibition was not seen in cell lines that lack the Bcl-2 expression (Daudi cells) or the t(14;18) translocation (Raji and Jurkat cells). There was no non-specific toxicity in Johnson cells exposed to two different control oligonucleotides. The growth inhibitory effects could be observed starting at a concentration of 3 $\mu\text{mol/L}$ of L-bcl-2, and the inhibitory effects were maximal at 6 $\mu\text{mol/L}$ concentration. One of the mechanisms by which L-bcl-2 exerts growth inhibition in Johnson cells might be through induction of apoptosis, since 15 treated cells showed a typical DNA internucleosomal degradation pattern, and an increased apoptotic index as measured by acridine orange. By day 3, 61% of Johnson cells treated with 5 $\mu\text{mol/L}$ of L-bcl-2 were in apoptosis as compared with 15% of cells treated with L-control oligos. Apoptosis was not observed in the other cell types.

Thus, the inhibition of Bcl-2 protein leads to cell growth inhibition in cells that are 25 dependent on the presence of Bcl-2 protein for maintaining viability. Gene transfer experiments have provided evidence that Bcl-2 plays an important role in maintaining lymphoid cell survival in vitro, although other autocrine growth factors may also be involved (Vaux *et. al.*, *Nature* (London), 355: 440, 1988; Reed *et al.*, *Proc. Nat'l Acad. Sci. USA*, 87:3660, 1990b; Blagosklonny and Neckers, *Eur. Cytokine Network*, 6:21, 1995). Using phosphorothionate antisense 30 oligonucleotides, Cotter and co-workers observed growth inhibition in DoHH2 cell line which

has the t(14;18) translocation and overexpress Bcl-2 protein, but not in FC11 cell line which overexpresses Bcl-2 protein without the t(14;18) translocation (Cotter *et. al.*, 1994). Cells that overexpress Bcl-2 and lack the t(14;18) translocation may need an apoptotic stimulus, like growth factor deprivation or treatment with chemotherapeutic drugs, to be driven into apoptosis and growth arrest (Reed, 1995). Antisense oligonucleotides may be used to reverse the chemotherapeutic resistance of those cells that also overexpress high levels of Bcl-2 without the t(14;18) translocation (Kitada *et al.*, *Antisense Res. Dev.*, 4:71, 1994). It is understood that such a combination of L-Bcl-2 of the present disclosure with an apoptotic stimulus would be encompassed by the spirit and scope of the appended claims.

Bax, a promoter of apoptotic cell death, may be a common partner involved in heterodimerization and regulation of other Bcl-2 family member functions (Sedlak *et. al.*, *Proc. Nat'l Acad. Sci. USA*, 92:7834, 1995). It has been suggested that the equilibrium in the formation of Bcl-2:Bax heterodimers and Bax:Bax homodimers appears to be central in the molecular regulation of apoptosis (Yin *et. al.*, 1994). Moreover, in a recent study, the ratio of Bcl-2:Bax correlated with cell death in IL-3-dependent FL5.12 cells. When approximately half or more of Bax was heterodimerized with Bcl-2, apoptosis was inhibited (Yang *et. al.*, 1995). The apoptotic death observed in Johnson cells, after incubation with L-bcl-2, could be due to decrease in Bcl-2:Bax ratio and formation of more Bax:Bax homodimers.

Another explanation is that other oncogenes and tumor suppressor genes such as C-MYC and p53 may be involved in the survival of the other cell lines. C-MYC, for example, is typically expressed in Burkitt lymphomas and in some transformed FL (McDonnell and Korsmeyer, 1991). Mutations of p53 gene, a suppressor gene involved in numerous human tumors, may also be involved in these cell lines; p53 gene encodes a DNA-binding protein that functions at least in part as a transcription factor to induce cell cycle arrest and apoptosis by upregulation of Bax (Vogelstein and Kinzler, *Cell*, 70:523, 1992; Miyashita *et. al.* *Oncogene*, 9:1799, 1994; Miyashita and Reed, *Cell*, 80:293, 1995). Probably, inhibition of Bcl-2 expression alone is not enough to induce apoptosis and growth inhibition in cells other than those cells with a high expression level of Bcl-2 such as those with a t(14;18) translocation. However, as described herein, a decrease in the ratio of Bcl-2/Bax by L-bcl-2 forms the basis for a molecular approach to follicular lymphoma therapy as well as other Bcl-2 diseases in cells in which Bax is expressed.

EXAMPLE 4: *Bcl-2 Regulation of p53 nuclear import*

The human prostate carcinoma cell line, LNCaP, which possesses a wild-type p53 gene (Carroll *et. al.*, *Prostate*, 23, 123-134, 1993), was selected to investigate the inhibition of p53-dependent programmed cell death by bcl-2. Stable bcl-2 expressing LNCaP prostate carcinoma 5 cell lines were generated and confirmed by Western blotting. Expression of bcl-2 conferred significant resistance to apoptosis induction following γ -irradiation compared to control clones as assessed by morphologic and flow cytometric analysis (FIG. 9A and 9B). No significant cell 10 death induction was observed in bcl-2 expressing LNCaP cells up to 48 hours following irradiation. Additionally, there was no difference in cell cycle distribution between LNCaP and 15 LNCaP-bcl-2 cells prior to irradiation and no evidence that the distribution of cells within the cell cycle was altered following irradiation. Approximately 60% of unirradiated control cells reside in G₀/G₁, 18% in S phase, and 22% in G₂/M compared to 59% of unirradiated LNCaP-bcl-2 cells in G₀/G₁, 21% in S phase, and 20% in G₂/M. These values were not significantly different from those observed at 30 minutes, 1 hour, 2 hours, 4 hours, and 8 hours following 15 irradiation.

Apoptosis induced in response to genotoxic damage is considered to be p53-dependent. In LNCaP cells, total cellular p53 protein was induced to approximately equivalent levels in bcl-2 expressing clones and control transfectants within 4 hours following 20 Gy γ -irradiation (FIG. 10A). Western blot analysis using isolated nuclei revealed that levels of p53 protein in the 20 nucleus increased within 2 hours of irradiation in control LNCaP, but not in bcl-2 expressing LNCaP, cells (FIG. 10A). This observation suggests that the nuclear import of p53 following DNA damage may be impaired in the context of high levels of bcl-2 protein.

Scanning confocal laser microscopy was used to further characterize p53 nuclear import following irradiation. The inhibition of nuclear p53 import in the bcl-2 cells was confirmed by 25 confocal microscopy using antibodies which recognize the p53 protein (FIG. 10B). Thus, by two independent techniques p53 nuclear import was demonstrated to be significantly inhibited in bcl-2 expressing cells following cell death induction by ionizing radiation.

In order to determine whether the transactivating ability of p53 was affected in the context of bcl-2 protein, NIH3T3 cells were transiently transfected with an mdm-2 promoter-30 luciferase reporter construct which possesses functional p53 binding sites. Luciferase activity

increased approximately 8-fold following co-transfection with a wild-type, but not mutant, p53 expression plasmid (FIG. 10C). Co-transfection of bcl-2 and wild-type p53 expression plasmids resulted in a 2-4 fold decrease in luciferase activity compared to wild-type p53 alone ($p \leq 0.02$).

Additional studies were undertaken using the RKO colon carcinoma cell line to assess whether the ability of bcl-2 to inhibit the nuclear import of wild-type p53 was specific for the LNCaP prostate cancer cell line. RKO cells possess a wild-type p53 gene (Nagasawa *et al.* *CancerRes.*, 55, 1842-1846, 1995) and also express bcl-2 protein. To downregulate bcl-2 expression in RKO cells, bcl-2 specific antisense oligonucleotides were delivered by liposomes. Bcl-2 protein levels were reduced 3-fold compared to RKO cells treated with empty liposomes or liposomes containing control oligonucleotides (FIG. 11A). Confocal microscopy was used to image p53 protein 4 hours following 10 Gy of γ -radiation in RKO cells treated with liposomes containing antisense bcl-2 oligonucleotides, or control oligonucleotides. RKO cells treated with control oligonucleotides showed that most of the p53 protein remained localized in the cytosol (FIG. 11B). In contrast, RKO cells in which bcl-2 had been downregulated by antisense oligonucleotides exhibited high levels of p53 protein within the nucleus and significant ($p \leq 0.005$) cell death induction compared to RKO cells treated with control oligonucleotides (FIG. 11C). These findings suggest that the ability of bcl-2 to modulate the import of wild-type p53 protein in response to DNA damage is not cell-type specific.

Irradiation and calcein-AM staining

Cells were preloaded with calcein-AM (Molecular Probes, Eugene, OR) at 1 mg/ml for 20 minutes in RPMI containing 10% FBS. Cells were given 20Gy of radiation using a ^{137}Cs γ -source and calcein-generated fluorescence was visualized using epifluorescence optics and a FITC filter (530 nm emission). The 20Gy dose of radiation was selected as the minimum dose required to induce maximum levels of apoptosis in LNCaP cells.

Flow cytometric analysis

Flow cytometric analysis of cell cycle and cell death induction in LNCaP cells. Single cell suspensions were fixed in 70% ethanol and incubated with 50 mg/ml propidium iodide (PI) and 20 mg/ml RNase for 15 min. at 37°C. Flow analysis was done with a EPICS Profile I at 488 nm excitation and collected for PI fluorescence using Elite Software 4.0 (Coulter Corp,

Miami, FL) and the Multi Cycle DNA Analysis program software (Phoenix Flow Systems, San Diego, CA). -

Transfections and Luciferase assays

LNCaP cells were transfected with the splenic focus forming virus expression plasmid 5 with, or without (control), the bcl-2 cDNA insert as previously described (Marin, *et al.* *Oncogene*, 12, 2259-66, 1996; Tu *et al.* *Cancer Lett.* 93, 147-155, 1995). The effector plasmids LTRXA and LTRKH expression plasmids have been described previously (Defile *et al.*, 1993) and represent wild-type and mutant p53, respectively. The reporter plasmid P2mdm2-Luc was made by cloning a 1Kb *Xhol*-fragment containing the p53 responsive element from the mouse 10 mdm2 gene into the *Smal* site of the pA3-luciferase plasmid. NIH3T3 cells were plated at a density of 0.5×10^6 cells per plate 24 hours before transfection. The effector wild type or mutant p53 plasmid (10 μ g), reporter plasmid P2mdm2-Luc (4 μ g) and β -galactosidase (β gal) 15 expression plasmid (3 μ g) were cotransfected with or without bcl-2 vector (20 μ g) following the calcium-phosphate method. The total amount of DNA transfected was normalized adding p- GEM plasmid up to 37 μ g of total DNA for all the transfections. At 48 hours after the 15 transfection the cells were harvested. Extracts were made and assayed for luciferase activity.

Immunofluorescence staining and confocal microscopy

LNCaPcontrol and LNCaP-bcl-2 cells were grown on laminin coated cover slides and 20 irradiated with 20 Gy. After 4h, cells were washed twice with PBS, then fixed in 4% paraformaldehyde for ten minutes and washed twice in PBS. Cells were blocked with 10% goat serum in PBS, incubated with p53 (AB-2, Calbiochem) antibody in 10% goat serum (1:75), washed twice, and incubated with fluorescein isothiocyanate (FITC)-labeled secondary antibody in 10% goat serum (1:200). Imaging was done using a Zeiss scanning confocal laser 25 microscope.

RKO cells were grown on laminin coated cover slides and incubated with liposomal 25 oligonucleotide formulations at a final concentration of 10 μ M at 37°C in a 5% CO₂ incubator for 3 days. 4h after irradiation with 10 Gy, cells were washed twice with PBS, then fixed in 4% paraformaldehyde for ten minutes and washed twice in PBS. Cells were blocked with 10 % goat serum in PBS, incubated with p53 (AB-2, Calbiochem) antibody in 10% goat serum (1:75),

washed twice and incubated with fluorescein isothiocyanate (FITC)-labeled secondary antibody in 10% goat serum (1:200). Confocal microscopy was done using a Zeiss scanning laser confocal microscope.

Antisense & DNA methods

5 P-ethoxy-oligonucleotides, a non-ionic and nuclease-resistant phosphodiester analog, were purchased from Oligo Therapeutics (Willsonville, OR). An oligonucleotide specific for the translation initiation site of human bcl-2 mRNA: 5'CAGCGTGCGCCATCCTTC3' (SEQ ID NO:1) was used as antisense oligonucleotide. The control oligonucleotide used was a scrambled version of Bcl-2 antisense oligonucleotide 5'ACGGTCCGCCACTCCTTCCC3' (SEQ ID NO:2). P-ethoxy-oligonucleotides, dissolved in DMSO, were added to phospholipids (Avanti 10 Polar Lipids, Alabaster, AL) in the presence of excess tert-butanol. The mixture was frozen in a dry ice/acetone bath, lyophilized overnight and hydrated with 0.9% saline at a final oligonucleotide concentration of 0.1 mmol/L. Empty liposomes were prepared identically as above, except that oligonucleotides were not included in the preparation. 0.25 x 10⁵ cells/mL 15 were seeded in a 24-well plate in 0.5 mL of the respective medium. Cells were incubated with antisense, control oligonucleotides and empty liposomes at final concentration of 10 µM at 37°C in a 5% CO₂ incubator for 3 days. Western blotting of whole cell extracts (40 µg) of control and bcl-2 transfected clones were analyzed by immunoblotting for bcl-2 protein using an anti-bcl-2 monoclonal antibody (Santa Cruz Biotechnology Inc, Santa Cruz, CA.).

20 *Analysis of protein expression in cell and nuclear extracts*

Western blot analysis of p53 protein induction and nuclear import following γ-irradiation. Subconfluent cultures of control LNCaP and LNCaP-bcl-2 cells were irradiated with 20Gy. Extracts of nuclei were prepared by scraping cell monolayers into hypotonic lysis buffer (100 mM Hepes, pH 7.4, 1.5 mM MgCl₂, 10 mM KCl, 0.5 mM β-mercaptoethanol and 5 ng/ml 25 leupeptin). After 10 minutes on ice, NP-40 was added to 0.625% and the crude nuclear pellet was recovered by centrifugation at 2000 x g for 5 minutes. The nuclear pellets were lysed in SDS-PAGE sample loading buffer. Extracts of nuclei and whole cells were prepared 2 and 4 h after irradiation. Equivalent amounts of lysates were analyzed by immunoblotting with p53 antibody (Santa Cruz).

Extracts of nuclei were prepared by scraping cell monolayers into hypotonic lysis buffer (100 mM Hepes, pH 7.4, 1.5 mM MgCl₂, 10 mM KCl, 0.5 mM β -mercaptoethanol and 5 ng/ml leupeptin). After 10 minutes on ice, NP-40 was added to 0.625% and the crude nuclear pellet was recovered by centrifugation at 2000 x g for 5 minutes. The nuclear pellets were lysed in 5 SDS-PAGE sample loading buffer.

EXAMPLE 5: In vivo Testing

In an initial round of *in vivo* trials, 30 nude (nu/nu) mice, aged about 5-6 weeks were each injected intraperitoneally with 3 million Johnson cells, follicular lymphoma cells bearing the t(14;18) translocation. Three groups of 10 mice each were used: untreated mice (group I), 10 liposomal p-ethoxy Bcl-2 antisense treated mice (group II) and liposomal Bcl-2 control treated mice (group III). One week after tumor implantation, groups II and III mice were administered a biweekly intravenous injection of liposomal Bcl-2 antisense oligos, or liposomal Bcl-2 control oligos. The oligo dose is 15 mg/kg of mouse body weight.

Some mice were observed to reach the moribund state (which is defined as tumor size exceeding 1.5 cm³) by day 50. On day 70 after implantation (63 days of treatment) six mice 15 (60%) in groups I and III had reached moribund state and were sacrificed, while only 2 mice (20%) in group II had reached moribund state and were sacrificed. One other mouse reached moribund state by day 77 (70 days of treatment). The study was terminated on day 78. Tissue from liver, kidney, spleen, heart, lungs, stomach, intestines and bone marrow were collected for 20 histopathology studies. Thus preliminary *in vivo* results indicate that the antisense Bcl-2 delivered in neutral liposomes are effective in inhibiting follicular lymphoma in mice.

EXAMPLE 6: Clinical Trials

This example is concerned with the development of human treatment protocols using the 25 lipid-associated oligo- and polynucleotide compositions. These lipid formulations will be of use in the clinical treatment of various *bcl-2*-overexpressing cancers and diseases in which transformed or cancerous cells play a role. Such treatment will be particularly useful tools in anti-tumor therapy, for example, in treating patients with FL. This treatment will also be useful in treating other conditions that are mediated by *bcl-2* over-expression and resistant to conventional regimens and treatments such as hematologic malignancies, both leukemias and

lymphomas, including follicular and nonfollicular lymphomas, chronic lymphocytic leukemia, and plasma cell dyscrasias; solid tumors like those associated with breast, prostate and colon cancer; and immune disorders.

5 The various elements of conducting a clinical trial, including patient treatment and monitoring, will be known to those of skill in the art in light of the present disclosure. The following information is being presented as a general guideline for use in establishing lipid-associated oligo-and polynucleotide compositions alone or in combinations with anti-cancer drugs in clinical trials.

10 Candidates for the phase 1 clinical trial will be patients on which all conventional therapies have failed. Liposomal Bcl-2 antisense oligos will be administered to them intravenously on a tentative weekly basis. To monitor disease course and evaluate the anti-tumor responses, it is contemplated that the patients should be examined for appropriate tumor markers every month. To assess the effectiveness of the drug, the following parameters will be monitored: tumor size and bone marrow infiltration of the cancer cells. Tests that will be used to 15 monitor the progress of the patients and the effectiveness of the treatments include: physical exam, X-ray, blood work and other clinical laboratory methodologies. In addition, peripheral blood and bone marrow samples will be drawn to assess the modification of the target protein expression. The doses given in the phase 1 study will be escalated as is done in standard phase 1 clinical phase trials, i.e. doses will be escalated until maximal tolerable ranges are reached.

20 Clinical responses may be defined by acceptable measure. For example, a complete response may be defined by complete disappearance of evidence of cancer cells for at least 2 months. Whereas a partial response may be defined by a 50% reduction of cancer cells for at least 2 months.

EXAMPLE 7: Human Treatment and Clinical Protocols

25 This example describes a protocol to facilitate the treatment of *bcl-2*-mediated diseases using lipid-associated oligo- or polynucleotide compositions alone or in combination with anti-cancer drugs.

Typically, patients that are candidates for treatment are those with FL although patients with hematologic malignancies, both leukemias and lymphomas; solid tumors like those associated with breast, prostate and colon cancer; and immune disorders may also be treated with the methods of this invention. The typical course of treatment will vary depending upon the 5 individual patient and disease being treated in ways known to those of skill in the art. For example, a patient with FL might be treated in eight week cycles, although longer duration may be used if no adverse effects are observed with the patient, and shorter terms of treatment may result if the patient does not tolerate the treatment as hoped. Each cycle will consist of between 20 and 10 35 individual doses spaced equally, although this too may be varied depending on the clinical situation.

A patient presenting a *bcl-2*-mediated condition, like FL, may be treated using the following protocol. Patients may, but need not, have received previous chemo-, radio- or gene therapeutic treatments. Optimally the patient will exhibit adequate bone marrow function (defined as peripheral absolute granulocyte count of $> 2,000/\text{mm}^3$ and platelet count of 100, 15 000/ mm^3 , adequate liver function (bilirubin 1.5mg/dl) and adequate renal function (creatinine 1.5mg/dl).

The over-expression of *bcl-2* is typically monitored before, during, and after the therapy. A composition of the present invention is typically administered orally or parenterally in dosage 20 unit formulations containing standard, well known non-toxic physiologically acceptable carriers, adjuvants, and vehicles as desired. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intra-arterial injection, or infusion techniques. The lipid-associated oligo- or poly-nucleotide compositions may be delivered to the patient before, after or concurrently with the other anti-cancer agents.

A typical treatment course may comprise about six doses delivered over a 7 to 21 day 25 period. Upon election by the clinician the regimen may be continued with six doses every three weeks or on a less frequent (monthly, bimonthly, quarterly etc.) basis. Of course, these are only exemplary times for treatment, and the skilled practitioner will readily recognize that many other time-courses are possible.

To kill *bcl-2*-overexpressing cancer cells using the methods and compositions described 30 in the present invention one will generally contact a target cell with the lipid-associated

formulations described previously. These compositions will be provided in an amount effective to kill or inhibit the proliferation of the cell.

Regional delivery of the lipid-associated formulations will be an efficient method for delivering a therapeutically effective dose to counteract the clinical disease. Alternatively systemic delivery may be appropriate. The therapeutic composition of the present invention may be administered to the patient directly at the site of the tumor. This is in essence a topical treatment of the surface of the cancer. The volume of the composition should usually be sufficient to ensure that the entire surface of the tumor is contacted by the lipid-associated oligo- or poly-nucleotide composition.

In one embodiment, administration simply entails injection of the therapeutic composition into the tumor. In another embodiment, a catheter is inserted into the site of the tumor and the cavity may be continuously perfused for a desired period of time.

Of course, the above-described treatment regimes may be altered in accordance with the knowledge gained from clinical trials such as those described in Example 5. Those of skill in the art will be able to take the information disclosed in this specification and optimize treatment regimes based on the clinical trials described in the specification.

- 45 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

5 (A) NAME: Board of Regents, The University of Texas System
(B) STREET: 201 W. 7th Street
(C) CITY: Austin
(D) STATE: Texas
10 (E) COUNTRY: US
(F) POSTAL CODE (ZIP): 78701
(G) TELEPHONE: (512) 418-3000
(H) TELEFAX: (713) 789-2679

15 (ii) TITLE OF INVENTION: INHIBITION OF Bcl-2 PROTEIN EXPRESSION BY LIPOSOMAL ANTISENSE OLIGODEOXYNUCLEOTIDES

20 (iii) NUMBER OF SEQUENCES: 7

(iv) COMPUTER READABLE FORM:

25 (A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(vi) PRIOR APPLICATION DATA:

30 (A) APPLICATION NUMBER: US 08/726,211
(B) FILING DATE: 04-OCT-1996

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

40 CAGCGTGCGC CATCCTTC

18

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

45 (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

50 ACGGTCCGCC ACTCCTTCCC

20

- 46 -

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

5 (A) LENGTH: 16 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

10 CTGAAGGGCT TCTTCC

16

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 5086 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 20 (D) TOPOLOGY: linear

(ix) FEATURE:

25 (A) NAME/KEY: CDS
 (B) LOCATION: 1459..2175

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

| | | |
|----|--|-----|
| 25 | GC GCCC GCCC CTCC GCGCCG CCTGCC GCGCCGCGC GCTCCGCCC GCCGCTCTCC | 60 |
| 30 | GTGGCCCCGC CGCGCTGCCG CCGCCGCCGC TGCCAGCGAA GGTGCCGGGG CTCCGGGCC | 120 |
| | TCCCTGCCGG CGGCCGTCAG CGCTCGGAGC GAACTGCCGCG ACGGGAGGTC CGGGAGGCGA | 180 |
| 35 | CCGTAGTCGC GCCGCCGCGC AGGACCAGGA GGAGGAGAAA GGGTGCAGCAG CCCGGAGGCG | 240 |
| | GGGTGCAGCG GTGGGGTGCA GCGGAAGAGG GGGTCCAGGG GGGAGAACTT CGTAGCAGTC | 300 |
| | ATCCTTTTA GGAAAAGAGG GAAAAAATAA AACCTCCCC CACCACCTCC TTCTCCCCAC | 360 |
| 40 | CCCTCGCCGC ACCACACACA GCGCGGGCTT CTAGCGCTCG GCACCGGCGG GCCAGGCGCG | 420 |
| | TCCTGCCTTC ATTTATCCAG CAGCTTTCG GAAAATGCAT TTGCTGTTCG GAGTTAAC | 480 |
| 45 | AGAAGACGAT TCCTGCCTCC GTCCCCGGCT CCTTCATCGT CCCATCTCCC CTGTCTCTCT | 540 |
| | CCTGGGGAGG CGTGAAGCGG TCCC GTGGAT AGAGATTCA GCGCTGTGTCC GCGCGTGTGT | 600 |
| | GCGCGCGTAT AAATTGCCGA GAAGGGAAA ACATCACAGG ACTTCTGCGA ATACCGGACT | 660 |
| 50 | GAAAATTGTA ATTCACTCTGC CGCCGCCGCT GCCAAAAAAA AACTCGAGCT CTTGAGATCT | 720 |
| | CCGGTTGGGA TTCCCTGCCGA TTGACATTTC TGTGAAGCAG AAGTCTGGGA ATCGATCTGG | 780 |
| 55 | AAATCCTCCT AATTTTACT CCCTCTCCCC CCGACTCCTG ATTCAATTGGG AAGTTTCAAA | 840 |
| | TCAGCTATAA CTGGAGAGTG CTGAAGATTG ATGGGATCGT TGCCTTATGC ATTTGTTTG | 900 |
| | GTTTTACAAA AAGGAAACTT GACAGAGGAT CATGCTGTAC TTAAAAAATA CAAGTAAGTC | 960 |

| | | |
|----|---|------|
| | TCGCACAGGA AATTGGTTTA ATGTAACCTT CAATGGAAAC CTTTGAGATT TTTTACTTAA | 1020 |
| 5 | AGTGCATTCG AGTAAATTAA ATTTCCAGGC AGCTTAATAC ATTGTTTTTA GCCGTGTTAC | 1080 |
| | TTGTAGTGTG TATGCCCTGC TTTCACTCAG TGTGTACAGG GAAACGCACC TGATTTTTA | 1140 |
| | CTTATTAGTT TGTTTTCTT TTAACCTTTC AGCATCACAG AGGAAGTAGA CTGATATTAA | 1200 |
| 10 | CAATACTTAC TAATAATAAC GTGCCTCATG AAATAAAGAT CCGAAAGGAA TTGGAATAAA | 1260 |
| | AATTCCTGC GTCTCATGCC AAGAGGGAAA CACCAGAAC AAGTGTCCG CGTGATTGAA | 1320 |
| 15 | GACACCCCT CGTCCAAGAA TGCAAAGCAC ATCCAATAAA ATAGCTGGAT TATAACTCCT | 1380 |
| | CTTCTTCTC TGGGGGCCGT GGGGTGGGAG CTGGGGCGAG AGGTGCCGTT GGCCCCCGTT | 1440 |
| | GCTTTCTC TGGGAAGG ATG GCG CAC GCT GGG AGA ACG GGG TAC GAC AAC | 1491 |
| 20 | Met Ala His Ala Gly Arg Thr Gly Tyr Asp Asn | |
| | 1 5 10 | |
| | CGG GAG ATA GTG ATG AAG TAC ATC CAT TAT AAG CTG TCG CAG AGG GGC | |
| | Arg Glu Ile Val Met Lys Tyr Ile His Tyr Lys Leu Ser Gln Arg Gly | |
| | 15 20 25 | 1539 |
| 25 | TAC GAG TGG GAT GCG GGA GAT GTG GGC GCC GCG CCC CCG GGG GCC GCC | |
| | Tyr Glu Trp Asp Ala Gly Asp Val Gly Ala Ala Pro Pro Gly Ala Ala | |
| | 30 35 40 | 1587 |
| 30 | CCC GCA CCG GGC ATC TTC TCC TCC CAG CCC GGG CAC ACG CCC CAT CCA | |
| | Pro Ala Pro Gly Ile Phe Ser Ser Gln Pro Gly His Thr Pro His Pro | |
| | 45 50 55 | 1635 |
| 35 | GCC GCA TCC CGC GAC CCG GTC GCC AGG ACC TCG CCG CTG CAG ACC CCG | |
| | Ala Ala Ser Arg Asp Pro Val Ala Arg Thr Ser Pro Leu Gln Thr Pro | |
| | 60 65 70 75 | 1683 |
| 40 | GCT GCC CCC GGC GCC GCG GGG CCT GCG CTC AGC CCG GTG CCA CCT | |
| | Ala Ala Pro Gly Ala Ala Gly Pro Ala Leu Ser Pro Val Pro Pro | |
| | 80 85 90 | 1731 |
| | GTG GTC CAC CTG GCC CTC CGC CAA GCC GGC GAC GAC TTC TCC CGC CGC | |
| | Val Val His Leu Ala Leu Arg Gln Ala Gly Asp Asp Phe Ser Arg Arg | |
| 45 | 95 100 105 | 1779 |
| | TAC CGC GGC GAC TTC GCC GAG ATG TCC AGC CAG CTG CAC CTG ACG CCC | |
| | Tyr Arg Gly Asp Phe Ala Glu Met Ser Ser Gln Leu His Leu Thr Pro | |
| | 110 115 120 | 1827 |
| 50 | TTC ACC GCG CGG GGA CGC TTT GCC ACG GTG GTG GAG GAG CTC TTC AGG | |
| | Phe Thr Ala Arg Gly Arg Phe Ala Thr Val Val Glu Glu Leu Phe Arg | |
| | 125 130 135 | 1875 |
| 55 | GAC GGG GTG AAC TGG GGG AGG ATT GTG GCC TTC TTT GAG TTC GGT GGG | |
| | Asp Gly Val Asn Trp Gly Arg Ile Val Ala Phe Phe Glu Phe Gly Gly | |
| | 140 145 150 155 | 1923 |

| | | |
|----|---|----------------|
| | GTC ATG TGT GTG GAG AGC GTC AAC CGG GAG ATG TCG CCC CTG GTG GAC Val Met Cys Val Glu Ser Val Asn Arg Glu Met Ser Pro Leu Val Asp 160 165 170 | 1971 |
| 5 | AAC ATC GCC CTG TGG ATG ACT GAG TAC CTG AAC CGG CAC CTG CAC ACC Asn Ile Ala Leu Trp Met Thr Glu Tyr Leu Asn Arg His Leu His Thr 175 180 185 | 2019 |
| 10 | TGG ATC CAG GAT AAC GGA GGC TGG GAT GCC TTT GTG GAA CTG TAC GGC Trp Ile Gln Asp Asn Gly Gly Trp Asp Ala Phe Val Glu Leu Tyr Gly 190 195 200 | 2067 |
| 15 | CCC AGC ATG CGG CCT CTG TTT GAT TTC TCC TGG CTG TCT CTG AAG ACT Pro Ser Met Arg Pro Leu Phe Asp Phe Ser Trp Leu Ser Leu Lys Thr 205 210 215 | 2115 |
| 20 | CTG CTC AGT TTG GCC CTG GTG GGA GCT TGC ATC ACC CTG GGT GCC TAT Leu Leu Ser Leu Ala Leu Val Gly Ala Cys Ile Thr Leu Gly Ala Tyr 220 225 230 235 | 2163 |
| | CTG AGC CAC AAG TGAAGTCAAC ATGCCTGCC CAAACAAATA TGCAAAAGGT Leu Ser His Lys | 2215 |
| 25 | TCACTAAAGC AGTAGAAATA ATATGCATTG TCAGTGATGT ACCATGAAAC AAAGCTGCAG GCTGTTAACG AAAAAATAAC ACACATATAA ACATCACACA CACAGACAGA CACACACACA CACACAATT AACAGTCTTC AGGCAAAACG TCGAATCAGC TATTACTGC CAAAGGGAAA | 2275 2335 2395 |
| 30 | TATCATTAT TTTTACATT ATTAAGAAAA AAGATTATT TATTTAAGAC AGTCCCATCA AAACTCCGTC TTTGGAAATC CGACCACTAA TTGCCAAACA CCGCTTCGTG TGGCTCCACC | 2455 2515 |
| 35 | TGGATGTTCT GTGCCTGTAA ACATAGATTC GCTTTCCATG TTGTTGGCCG GATCACCATC TGAAGAGCAG ACGGATGGAA AAAGGACCTG ATCATTGGGG AAGCTGGCTT TCTGGCTGCT | 2575 2635 |
| 40 | GGAGGCTGGG GAGAAGGTGT TCATTCACCTT GCATTTCTTT GCCCTGGGG CGTGATATTA ACAGAGGGAG GGTTCCCGTG GGGGAAAGTC CATGCCTCCC TGGCCTGAAG AAGAGACTCT | 2695 2755 |
| | TTGCATATGA CTCACATGAT GCATACCTGG TGGGAGGAAA AGAGTTGGGA ACTTCAGATG | 2815 |
| 45 | GACCTAGTAC CCACTGAGAT TTCCACGCCG AAGGACAGCG ATGGGAAAAA TGCCCTTAAA TCATAGGAAA GTATTTTTT AAGCTACCAA TTGTGCCGAG AAAAGCATTT TAGCAATTAA | 2875 2935 |
| 50 | TACAATATCA TCCAGTACCT TAAACCTGA TTGTGTATAT TCATATATTT TGGATACGCA CCCCCAACT CCCAATACTG GCTCTGTCTG AGTAAGAAC AGAACCTCT GGAACTTGAG | 2995 3055 |
| | GAAGTGAACA TTTCGGTGAC TTCCGATCAG GAAGGCTAGA GTTACCCAGA GCATCAGGCC | 3115 |
| 55 | GCCACAAAGTG CCTGCTTTA GGAGACCGAA GTCCGCAGAA CCTACCTGTG TCCCAGCTTG GAGGCCTGGT CCTGGAACTG AGCCGGGCC TCACTGGCCT CCTCCAGGGA TGATCAACAG | 3175 3235 |

| | | |
|----|---|------|
| | GGTAGTGTGG TCTCCGAATG TCTGGAAGCT GATGGATGGA GCTCAGAATT CCACTGTCAA | 3295 |
| | GAAAGAGCAG TAGAGGGGTG TGGCTGGGCC TGTCACCCTG GGGCCCTCCA GGTAGGCCCG | 3355 |
| 5 | TTTCACGTG GAGCATAGGA GCCACGACCC TTCTTAAGAC ATGTATCACT GTAGAGGGAA | 3415 |
| | GGAACAGAGG CCCTGGGCCT TCCTATCAGA AGGACATGGT GAAGGCTGGG AACGTGAGGA | 3475 |
| 10 | GAGGCAATGG CCACGGCCA TTTTGGCTGT AGCACATGGC ACGTTGGCTG TGTGGCCTTG | 3535 |
| | GCCACCTGTG AGTTTAAAGC AAGGCTTAA ATGACTTTGG AGAGGGTCAC AAATCCTAAA | 3595 |
| | AGAAGCATTG AAGTGAGGTG TCATGGATTA ATTGACCCCT GTCTATGGAA TTACATGTAA | 3655 |
| 15 | AACATTATCT TGTCACTGTA GTTTGGTTTT ATTTGAAAAC CTGACAAAAA AAAAGTTCCA | 3715 |
| | GGTGTGGAAT ATGGGGGTTA TCTGTACATC CTGGGGCATT AAAAAAAAAT CAATGGTGGG | 3775 |
| 20 | GAACATATAAA GAAGTAACAA AAGAAGTGAC ATCTTCAGCA AATAAACTAG GAAATTTTT | 3835 |
| | TTTCTTCCAG TTTAGAATCA GCCTTGAAAC ATTGATGGAA TAACTCTGTG GCATTATTGC | 3895 |
| | ATTATATACC ATTTATCTGT ATTAACCTTG GAATGTACTC TGTCAATGT TTAATGCTGT | 3955 |
| 25 | GGTTGATATT TCGAAAGCTG CTTTAAAAAA ATACATGCAT CTCAGCGTTT TTTTGTFFFF | 4015 |
| | AATTGTATTT AGTTATGGCC TATACACTAT TTGTGAGCAA AGGTGATCGT TTTCTGTTTG | 4075 |
| 30 | AGATTTTTAT CTCTTGATTC TTCAAAAGCA TTCTGAGAAG GTGAGATAAG CCCTGAGTCT | 4135 |
| | CAGCTACCTA AGAAAAACCT GGATGTCACT GGCCACTGAG GAGCTTGTT TCAACCAAGT | 4195 |
| | CATGTGCATT TCCACGTCAA CAGAATTGTT TATTGTGACA GTTATATCTG TTGTCCCTTT | 4255 |
| 35 | GACCTTGTTC CTTGAAGGTT TCCTCGTCCC TGGGCAATTG CGCATTAAAT TCATGGTATT | 4315 |
| | CAGGATTACA TGCATGTTG GTTAAACCCA TGAGATTCAT TCAGTTAAAA ATCCAGATGG | 4375 |
| 40 | CGAATGACCA GCAGATTCAA ATCTATGGTG GTTGACCTT TAGAGASTTG CTTTACGTGG | 4435 |
| | CCTGTTCAA CACAGACCCA CCCAGAGCCC TCCTGCCCTC CTTCCGCGGG GGCTTTCTCA | 4495 |
| | TGGCTGTCCT TCAGGGTCTT CCTGAAATGC AGTGGTCGTT ACGCTCCACC AAGAAAGCAG | 4555 |
| 45 | GAAACCTGTG GTATGAAGCC AGACCTCCCC GGCGGGCCTC AGGGAACAGA ATGATCAGAC | 4615 |
| | CTTTGAATGA TTCTAATTTT TAAGCAAAAT ATTATTTAT GAAAGGTTA CATTGTCAA | 4675 |
| | GTGATGAATA TGGAATATCC AATCCTGTGC TGCTATCCTG CCAAAATCAT TTTAATGGAG | 4735 |
| 50 | TCAGTTGCA GTATGCTCCA CGTGGTAAGA TCCTCCAAGC TGCTTAGAA GTAACAATGA | 4795 |
| | AGAACGTGGA CGTTTTAAT ATAAAGCCTG TTTTGTCTTT TGTTGTTGTT CAAACGGGAT | 4855 |
| 55 | TCACAGAGTA TTTGAAAAAT GTATATATAT TAAGAGGTCA CGGGGGCTAA TTGCTAGCTG | 4915 |
| | GCTGCCCTTT GCTGTGGGGT TTTGTTACCT GGTTTAATA ACAGTAAATG TGCCCAGCCT | 4975 |

- 50 -

CTTGGCCCCA GAACTGTACA GTATTGTGGC TGCACCTTGCT CTAAGAGTAG TTGATGTTGC 5035
ATTTTCCTTA TTGTTAAAAA CATGTTAGAA GCAATGAATG TATATAAAAG C 5086

5

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 239 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

15 Met Ala His Ala Gly Arg Thr Gly Tyr Asp Asn Arg Glu Ile Val Met
1 5 10 15
Lys Tyr Ile His Tyr Lys Leu Ser Gln Arg Gly Tyr Glu Trp Asp Ala
20 20 25 30
Gly Asp Val Gly Ala Ala Pro Pro Gly Ala Ala Pro Ala Pro Gly Ile
35 35 40 45
25 Phe Ser Ser Gln Pro Gly His Thr Pro His Pro Ala Ala Ser Arg Asp
50 50 55 60
Pro Val Ala Arg Thr Ser Pro Leu Gln Thr Pro Ala Ala Pro Gly Ala
65 65 70 75 80
30 Ala Ala Gly Pro Ala Leu Ser Pro Val Pro Pro Val Val His Leu Ala
85 85 90 95
Leu Arg Gln Ala Gly Asp Asp Phe Ser Arg Arg Tyr Arg Gly Asp Phe
35 100 105 110
Ala Glu Met Ser Ser Gln Leu His Leu Thr Pro Phe Thr Ala Arg Gly
115 115 120 125
40 Arg Phe Ala Thr Val Val Glu Glu Leu Phe Arg Asp Gly Val Asn Trp
130 130 135 140
Gly Arg Ile Val Ala Phe Phe Glu Phe Gly Gly Val Met Cys Val Glu
145 145 150 155 160
45 Ser Val Asn Arg Glu Met Ser Pro Leu Val Asp Asn Ile Ala Leu Trp
165 165 170 175
Met Thr Glu Tyr Leu Asn Arg His Leu His Thr Trp Ile Gln Asp Asn
50 180 185 190
Gly Gly Trp Asp Ala Phe Val Glu Leu Tyr Gly Pro Ser Met Arg Pro
195 195 200 205
55 Leu Phe Asp Phe Ser Trp Leu Ser Leu Lys Thr Leu Leu Ser Leu Ala
210 215 220

- 51 -

Leu Val Gly Ala Cys Ile Thr Leu Gly Ala Tyr Leu Ser His Lys
225 230 235

5 (2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 911 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 147.761

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

TGATTGAAGA CACCCCTCG TCCAAGAATG CAAAGCACAT CCAATAAAAT AGCTGGATTA 20
 TAACTCCTCT TCTTTCTCTG GGGGCCGTGG GGTGGGAGCT GGGGCGAGAG GTGCCGTTGG 120
 CCCCCGTTGC TTTTCCTCTG GGAAGG ATG GCG CAC GCT GGG AGA ACG GGG TAC 173
 Met Ala His Ala Gly Arg Thr Gly Tyr
 1 5
 GAC AAC CGG GAG ATA GTG ATG AAG TAC ATC CAT TAT AAG CTG TCG CAG 221
 Asp Asn Arg Glu Ile Val Met Lys Tyr Ile His Tyr Lys Leu Ser Gln
 10 15 20 25
 AGG GGC TAC GAG TGG GAT GCG GGA GAT GTG GGC GCC GCG CCC CCG GGG 269
 Arg Gly Tyr Glu Trp Asp Ala Gly Asp Val Gly Ala Ala Pro Pro Gly
 30 35 40
 GCC GCC CCC GCA CCG GGC ATC TTC TCC TCC CAG CCC GGG CAC ACG CCC 317
 Ala Ala Pro Ala Pro Gly Ile Phe Ser Ser Gln Pro Gly His Thr Pro
 45 50 55
 CAT CCA GCC GCA TCC CGC GAC CCG GTC GCC AGG ACC TCG CCG CTG CAG 365
 His Pro Ala Ala Ser Arg Asp Pro Val Ala Arg Thr Ser Pro Leu Gln
 60 65 70
 ACC CCG GCT GCC CCC GGC GCC GCG GGG CCT GCG CTC AGC CCG GTG 413
 Thr Pro Ala Ala Pro Gly Ala Ala Gly Pro Ala Leu Ser Pro Val
 75 80 85
 CCA CCT GTG GTC CAC CTG GCC CTC CGC CAA GCC GGC GAC GAC TTC TCC 461
 Pro Pro Val Val His Leu Ala Leu Arg Gln Ala Gly Asp Asp Phe Ser
 90 95 100 105
 CGC CGC TAC CGC GGC GAC TTC GCC GAG ATG TCC AGC CAG CTG CAC CTG 509
 Arg Arg Tyr Arg Gly Asp Phe Ala Glu Met Ser Ser Gln Leu His Leu
 110 115 120
 ACG CCC TTC ACC GCG CGG GGA CGC TTT GCC ACG GTG GTG GAG GAG CTC 557
 Thr Pro Phe Thr Ala Arg Gly Arg Phe Ala Thr Val Val Glu Glu Leu
 125 130 135

- 52 -

| | | | |
|---|---|-----|-----|
| 140 | 145 | 150 | 605 |
| 155 | 160 | 165 | 653 |
| 170 | 175 | 180 | 701 |
| 190 | 195 | 200 | 749 |
| 205 | | | 801 |
| GGCTCCTGGG CAATGGGAGG CTGTGGAGCC GGCGAAATAA AATCAGAGTT GTTGCTTCCC | | | 861 |
| GGCGTGTCCC TACCTCCTCC TCTGGACAAA GCGTTCACTC CCAACCTGAC | | | 911 |
| 25 (2) INFORMATION FOR SEQ ID NO: 7: | | | |
| (i) SEQUENCE CHARACTERISTICS: | | | |
| (A) LENGTH: 205 amino acids | | | |
| 30 | (B) TYPE: amino acid | | |
| | (C) TOPOLOGY: linear | | |
| (ii) MOLECULE TYPE: protein | | | |
| (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7: | | | |
| 35 | Met Ala His Ala Gly Arg Thr Gly Tyr Asp Asn Arg Glu Ile Val Met | | |
| 1 | 5 | 10 | 15 |
| 40 | Lys Tyr Ile His Tyr Lys Leu Ser Gln Arg Gly Tyr Glu Trp Asp Ala | | |
| | 20 | 25 | 30 |
| Gly Asp Val Gly Ala Ala Pro Pro Gly Ala Ala Pro Ala Pro Gly Ile | | | |
| | 35 | 40 | 45 |
| 45 | Phe Ser Ser Gln Pro Gly His Thr Pro His Pro Ala Ala Ser Arg Asp | | |
| | 50 | 55 | 60 |
| 50 | Pro Val Ala Arg Thr Ser Pro Leu Gln Thr Pro Ala Ala Pro Gly Ala | | |
| | 65 | 70 | 75 |
| 55 | Ala Ala Gly Pro Ala Leu Ser Pro Val Pro Pro Val Val His Leu Ala | | |
| | 85 | 90 | 95 |
| 55 | Leu Arg Gln Ala Gly Asp Asp Phe Ser Arg Arg Tyr Arg Gly Asp Phe | | |
| | 100 | 105 | 110 |

- 53 -

Ala Glu Met Ser Ser Gln Leu His Leu Thr Pro Phe Thr Ala Arg Gly
115 120 125

5 Arg Phe Ala Thr Val Val Glu Glu Leu Phe Arg Asp Gly Val Asn Trp
130 135 140

Gly Arg Ile Val Ala Phe Phe Glu Phe Gly Gly Val Met Cys Val Glu
145 150 155 160

10 Ser Val Asn Arg Glu Met Ser Pro Leu Val Asp Asn Ile Ala Leu Trp
165 170 175

Met Thr Glu Tyr Leu Asn Arg His Leu His Thr Trp Ile Gln Asp Asn
180 185 190

15 Gly Gly Trp Val Gly Ala Ser Gly Asp Val Ser Leu Gly
195 200 205

CLAIMS:

1. A composition comprising a first polynucleotide that hybridizes to a second, Bcl-2-encoding polynucleotide under intracellular conditions and a neutral lipid associated with said first polynucleotide, and wherein said composition contains no cationic lipid.

5 2. The composition of claim 1, wherein said first polynucleotide is an oligonucleotide having a length of between about 8 and about 50 bases.

3. The composition of claim 1, wherein the first polynucleotide is complementary to the translation initiation site of Bcl-2 mRNA.

4. The composition of claim 3, wherein the polynucleotide is an oligonucleotide comprising the sequence CAGCGTGCGCCATCCTTC (SEQ ID NO:1).

5. The composition of claim 1, comprising a liposome formed from the lipid.

6. The composition of claim 5, wherein the first polynucleotide is encapsulated in the liposome.

7. The composition of claim 1, wherein the lipid is a phosphatidylcholine, a phosphatidylglycerol, or a phosphatidylethanolamine.

8. The composition of claim 7, wherein the lipid is dioleoylphosphatidylcholine.

9. A composition comprising an expression construct that encodes a first polynucleotide that hybridizes to a second, Bcl-2-encoding polynucleotide under intracellular conditions, wherein said first polynucleotide is under the control of a promoter that is active in eukaryotic cells, and wherein said construct is associated with a neutral lipid, and further wherein said composition contains no cationic lipid.

10. A method of inhibiting a Bcl-2-associated disease comprising obtaining a first polynucleotide that hybridizes to a second, Bcl-2-encoding polynucleotide under intracellular conditions, mixing the first polynucleotide with a neutral lipid to form a polynucleotide/lipid association, and administering said association to a cell, wherein said cell expresses both Bcl-2 and Bax.

11. The method of claim 10, wherein the cell is a cancer cell.
12. The method of claim 11, wherein said cancer cell is a follicular lymphoma cell.
13. The method of claim 10, wherein said first polynucleotide is an oligonucleotide having a length of between about 8 and about 50 bases.
- 5 14. The method of claim 10, comprising a liposome formed from the lipid.
15. The method of claim 14, wherein the liposome encapsulates the first polynucleotide.
16. The method of claim 10, wherein said contacting takes place in an animal.
17. The method of claim 16, wherein said animal is a human.
18. The method of claim 17, wherein said composition is delivered to said human in a volume of 0.50-10.0 ml per dose.
- 10 19. The method of claim 17, wherein said composition is delivered to said human in an amount of from about 5 to about 30 mg polynucleotide per m².
20. The method of claim 19, wherein said composition is administered three times per week for eight weeks.




FIG. 1




FIG. 2

3/11




FIG. 3

4/11




FIG. 4A






FIG. 4B



5/11




FIG. 5

6/11




FIG. 6A




FIG. 6B

7/11



FIG. 7

8/11




FIG. 8A




FIG. 8B

9/11




FIG. 9A





FIG. 9B

10/11

WHOLE CELLS

| CONTROL | | BCL-2 | | | |
|---------|---|-------|---|---|---|
| 0 | 2 | 4 | 0 | 2 | 4 |
| ■ | ■ | ■ | ■ | ■ | ■ |




■ bcl-2 cells
 ▲ contr. cells

FIG. 10A

NUCLEI

| CONTROL | | BCL-2 | | | |
|---------|---|-------|---|---|---|
| 0 | 2 | 4 | 0 | 2 | 4 |
| ■ | ■ | ■ | ■ | ■ | ■ |



HOLD INCREASE





FIG. 10C



10B

bcl-2

contr.

time (min)

nt 5 10 20 nt 5 10 20

Ik B

FIG. 10C

11/11




FIG. 11B




FIG. 11A

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/18348

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 9/127, 31/70; C07H 21/04; C12N 15/00
 US CL : 424/450; 435/172.3, 375; 514/44; 536/24.5

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/450; 435/172.3, 375; 514/44; 536/24.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN: Medline, Biosis, Embase, CAPLus, WPIDS, JAPIO, PATOSEP, PATOSWO
 search terms: bcl-2, antisense, lipid, liposome

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| X | WO 93/20200 A1 (IMPERIAL CANCER RESEARCH TECHNOLOGY LIMITED) 14 October 1993, page 7, lines 10-29; page 15, lines 16-23; page 18, lines 26-30; page 59, lines 6-7. | 1-6, 9 |
| -- | | -- |
| Y | | 7, 8 |
| X | WO 95/08350 A1 (BROTMAN, HARRIS, F. et al.) 30 March 1995, page 3, lines 2-22; page 13, lines 2-5; page 14, lines 16-25. | 1-6 |
| -- | | -- |
| Y | | 7, 8 |
| X | ALMAZAN et al. Methylphosphonate-containing oligonucleotides efficiently and specifically inhibit Bcl-2 and erbB-2 expression in vitro. Proc. Amer. Assoc. Cancer Res. March 1996, Vol. 37, page 353 Abstract No. 2407, see entire document. | 1, 2 |

 Further documents are listed in the continuation of Box C.

See patent family annex.

| | |
|--|--|
| Special categories of cited documents: | |
| *A* document defining the general state of the art which is not considered to be of particular relevance | *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| *B* earlier document published on or after the international filing date | *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| *L* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| *O* document referring to an oral disclosure, use, exhibition or other means | *Z* document member of the same patent family |
| *P* document published prior to the international filing date but later than the priority date claimed | |

Date of the actual completion of the international search

10 DECEMBER 1997

Date of mailing of the international search report

14 JAN 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ROBERT SCHWARTZMAN

Telephone No. (703) 308-0196



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/18348

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| X | TORMO et al. Antitumor activity of liposomal-bcl-2-antisense oligonucleotides in follicular lymphoma. Proc. Amer. Assoc. Cancer Res. March 1996, Vol.37, page 173, Abstract No. 1190, see entire document. | 1, 2, 5, 6 |
| -- | | -- |
| Y | LEDLEY, F.D. Non-viral gene therapy. Curr. Opin. Biotechnol. 1994, Vol. 5, pages 626-636, see entire document. | 7, 8 |
| A | ROJANASAKUL, Y. Antisense oligonucleotide therapeutics: drug delivery and targeting. Adv. Drug Delivery Rev. 1996, Vol.18, pages 115-131, see entire document. | 1-20 |