



(51) International Patent Classification:

A01N 47/22 (2006.01) A01N 57/16 (2006.01)
A01N 47/24 (2006.01) A01N 57/28 (2006.01)
A01N 57/14 (2006.01) A01N 63/00 (2006.01)

(21) International Application Number:

PCT/EP2013/061023

(22) International Filing Date:

29 May 2013 (29.05.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12169936.7 30 May 2012 (30.05.2012) EP
12197141.0 14 December 2012 (14.12.2012) EP

(71) Applicant: **BAYER CROPSCIENCE AG** [DE/DE]; Alfred-Nobel-Strasse 50, 40789 Monheim (DE).

(72) Inventors: **HELLWEGE, Elke**; Louveciennestraße 91, 40764 Langenfeld (DE). **ANDERSCH, Wolfram**; Schلودerdicher Weg 77, 51468 Bergisch-Gladbach (DE). **STENZEL, Klaus**; Seesener Straße 17, 40595 Düsseldorf (DE). **SPRINGER, Bernd**; Haferkamp 8, 51061 Köln (DE).

(74) Agent: **GILLE HRABAL**; Brucknerstr. 20, 40593 Düsseldorf (DE).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: COMPOSITIONS COMPRISING A BIOLOGICAL CONTROL AGENT AND AN INSECTICIDE

(57) Abstract: The present invention relates to a composition comprising at least one biological control agent selected from the group consisting of *Bacillus chitinosporus* AQ746 (NRRL Accession No. B-21618), *Bacillus mycoides* AQ726 (NRRL Accession No. B-21664), *Bacillus pumilus* (NRRL Accession No. B-30087), *Bacillus pumilus* AQ717 (NRRL Accession No. B-21662), *Bacillus sp.* AQ175 (ATCC Accession No. 55608), *Bacillus sp.* AQ1 77 (ATCC Accession No. 55609), *Bacillus sp.* AQ1 78 (ATCC Accession No. 53522), *Bacillus subtilis* AQ743 (NRRL Accession No. B-21665), *Bacillus subtilis* AQ713 (NRRL Accession No. B-21661), *Bacillus subtilis* AQ153 (ATCC Accession No. 55614), *Bacillus thuringiensis* BD#32 (NRRL Accession No. B-21530), *Bacillus thuringiensis* AQ52 (NRRL Accession No. B-21619), *Muscodor albus* 620 (NRRL Accession No. 30547), *Muscodor rosaeus* A3-5 (NRRL Accession No. 30548), *Rhodococcus globerulus* AQ719 (NRRL Accession No. B-21663), *Streptomyces galbus* (NRRL Accession No. 30232), *Streptomyces sp.* (NRRL Accession No. B-30145), *Bacillus thuringiensis subspec. kurstaki* BMP 1 23, *Bacillus subtilis* AQ30002 (NRRL Accession No. B-50421), and *Bocillus subtilis* AQ 30004 (NRRL Accession No. B-50455) and/or a mutant of these strains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens and at least one insecticide selected from the group consisting of acetylcholinesterase (AChE) inhibitors, GABA-gated chloride channel antagonists, chloride channel activators and nicotinic acetylcholine receptor (nAChR) channel blockers in a synergistically effective amount. Furthermore, the present invention relates to the use of this composition as well as a method for reducing overall damage of plants and plant parts.



WO 2013/178658 A1

COMPOSITIONS COMPRISING A BIOLOGICAL CONTROL AGENT AND AN INSECTICIDE

The present invention relates to a composition comprising at least one biological control agent selected from specific microorganisms and/or a mutant of these strains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens and at least one specific insecticide in a synergistically effective amount. Furthermore, the present invention relates to the use of this composition as well as a method for reducing overall damage of plants and plant parts.

Synthetic insecticides or fungicides often are non-specific and therefore can act on organisms other than the target ones, including other naturally occurring beneficial organisms. Because of their chemical nature, they may be also toxic and non-biodegradable. Consumers worldwide are increasingly conscious of the potential environmental and health problems associated with the residuals of chemicals, particularly in food products. This has resulted in growing consumer pressure to reduce the use or at least the quantity of chemical (i. e. synthetic) pesticides. Thus, there is a need to manage food chain requirements while still allowing effective pest control.

A further problem arising with the use of synthetic insecticides or fungicides is that the repeated and exclusive application of an insecticide or fungicides often leads to selection of resistant microorganisms. Normally, such strains are also cross-resistant against other active ingredients having the same mode of action. An effective control of the pathogens with said active compounds is then not possible any longer. However, active ingredients having new mechanisms of action are difficult and expensive to develop.

The risk of resistance development in pathogen populations as well as environmental and human health concerns have fostered interest in identifying alternatives to synthetic insecticides and fungicides for managing plant diseases. The use of biological control agents (BCAs) is one alternative. However, the effectiveness of most BCAs is not at the same level as for conventional insecticides and fungicides, especially in case of severe infection pressure. Consequently, known biological control agents, their mutants and

metabolites produced by them are, in particular in low application rates, not entirely satisfactory.

Thus, there is a constant need for developing new, alternative plant protection agents which in some areas at least help to fulfill the above-mentioned requirements.

- 5 WO 2009/037242 A2 relates to a fungicidal composition of one of two specific fungicidal bacterial strains, namely *Bacillus subtilis* and *Bacillus pumilus*, and a synthetic fungicide for controlling phytopathogenic harmful fungi. However, the control of insects is not mentioned at all.

- 10 WO 2010/108973 A2 describes a method for controlling harmful fungi comprising different sequential treatment blocks of plants with at least one fungicidal biological control agent and at least one synthetic fungicide. Consequently, the control of insects is not addressed in this patent application.

- 15 In view of this, it was in particular an object of the present invention to provide compositions which exhibit activity against insects, mites, nematodes and/or phytopathogens. Moreover, it was a further particular object of the present invention, to reduce the application rates and broaden the activity spectrum of the biological control agents and the insecticides, and thereby to provide a composition which, preferably at a reduced total amount of active compounds applied, has improved activity against insects, mites, nematodes and/or phytopathogens. In particular, it was a
20 further object of the present invention to provide a composition which, when applied to a crop, results in a decreased amount of residues in the crop, thereby reducing the risk of resistance formation and nevertheless provides efficient disease control.

- Accordingly, it was found that these objects at least partly are solved by the compositions according to the invention as defined in the following. The composition
25 according to the present invention preferably fulfills the above-described needs. It has been surprisingly discovered that the application of the composition according to the present invention in a simultaneous or sequential way to plants, plant parts, harvested fruits, vegetables and/or plant's locus of growth preferably allows better control of insects, mites, nematodes and/or phytopathogens than it is possible with the strains,
30 their mutants and/or their metabolites produced by the strains on the one hand and with the individual insecticides on the other hand, alone (synergistic mixtures). By applying the biological control agent and the insecticide according to the invention

the activity against insects, mites, nematodes and/or phytopathogens is preferably increased in a superadditive manner.

As a consequence, the composition according to the present invention preferably allows a reduced total amount of active compounds to be used and thus the crops which have been treated by this composition preferably show a decreased amount of residues in the crop. Accordingly, the risk of resistance formation of harmful microorganisms is decreased.

The present invention is directed to a composition comprising at least one biological control agent selected from the group consisting of *Bacillus chittinosporus* AQ746 (NRRL Accession No. B-21618), *Bacillus mycolides* AQ726 (NRRL Accession No. B-21664), *Bacillus pumilus* (NRRL Accession No. B-30087), *Bacillus pumilus* AQ717 (NRRL Accession No. B-21662), *Bacillus sp.* AQ175 (ATCC Accession No. 55608), *Bacillus sp.* AQ177 (ATCC Accession No. 55609), *Bacillus sp.* AQ178 (ATCC Accession No. 53522), *Bacillus subtilis* AQ743 (NRRL Accession No. B-21665), *Bacillus subtilis* AQ713 (NRRL Accession No. B-21661), *Bacillus subtilis* AQ153 (ATCC Accession No. 55614), *Bacillus thuringiensis* BD#32 (NRRL Accession No. B-21530), *Bacillus thuringiensis* AQ52 (NRRL Accession No. B-21619), *Muscodora albus* 620 (NRRL Accession No. 30547), *Muscodora roseus* A3-5 (NRRL Accession No. 30548), *Rhodococcus globerulus* AQ719 (NRRL Accession No. B-21663), *Streptomyces galbus* (NRRL Accession No. 30232), *Streptomyces sp.* (NRRL Accession No. B-30145), *Bacillus thuringiensis* subspec. *kurstaki* BMP 123, *Bacillus subtilis* AQ30002 (NRRL Accession No. B-50421), and *Bacillus subtilis* AQ 30004 (NRRL Accession No. B-50455) and/or a mutant of these strains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens and at least one insecticide selected from the group consisting of acetylcholinesterase (AChE) inhibitors, GABA-gated chloride channel antagonists, chloride channel activators and nicotinic acetylcholine receptor (nAChR) channel blockers in a synergistically effective amount.

Furthermore, the present invention relates to a kit of parts comprising at least one of the specific biological control agents and at least one of the specific insecticides. The present invention is further directed to the use of said composition for reducing overall damage of plants and plant parts as well as losses in harvested fruits or vegetables caused by insects, mites, nematodes and/or phytopathogens.

Moreover, the present invention provides a method for reducing overall damage of plants and plant parts as well as losses in harvested fruits or vegetables caused by insects, mites, nematodes and/or phytopathogens.

Biological control agents

5 In general a "pesticidal" means the ability of a substance to increase mortality or inhibit the growth rate of plant pests. The term is used herein, to describe the property of a substance to exhibit activity against insects, mites, nematodes and/or phytopathogens. In the sense of the present invention the term "pests" include insects, mites, nematodes and/or phytopathogens.

10 As used herein, "biological control" is defined as control of a pathogen and/or insect and/or an acarid and/or a nematode by the use of a second organism. Known mechanisms of biological control include enteric bacteria that control root rot by out-competing fungi for space on the surface of the root. Bacterial toxins, such as antibiotics, have been used to control pathogens. The toxin can be isolated and
15 applied directly to the plant or the bacterial species may be administered so it produces the toxin in situ.

The term "metabolite" refers to any compound, substance or byproduct of a fermentation of a said microorganism that has pesticidal activity.

The term "mutant" refers to a variant of the parental strain as well as methods for
20 obtaining a mutant or variant in which the pesticidal activity is greater than that expressed by the parental strain. The "parent strain" is defined herein as the original strain before mutagenesis. To obtain such mutants the parental strain may be treated with a chemical such as N-methyl-N'-nitro-N-nitrosoguanidine, ethylmethanesulfone, or by irradiation using gamma, x-ray, or UV-irradiation, or by other means well known to
25 those skilled in the art.

A "variant" is a strain having all the identifying characteristics of the NRRL or ATCC Accession Numbers as indicated in this text and can be identified as having a genome that hybridizes under conditions of high stringency to the genome of the NRRL or ATCC Accession Numbers.

30 "Hybridization" refers to a reaction in which one or more polynucleotides react to form a complex that is stabilized via hydrogen bonding between the bases of the nucleotide

residues. The hydrogen bonding may occur by Watson-Crick base pairing, Hoogsteen binding, or in any other sequence-specific manner. The complex may comprise two strands forming a duplex structure, three or more strands forming a multi-stranded complex, a single self-hybridizing strand, or any combination of these. Hybridization reactions can be performed under conditions of different "stringency". In general, a low stringency hybridization reaction is carried out at about 40 °C in 10 X SSC or a solution of equivalent ionic strength/temperature. A moderate stringency hybridization is typically performed at about 50 °C in 6 X SSC, and a high stringency hybridization reaction is generally performed at about 60 °C in 1 X SSC.

10 A variant of the indicated NRRL or ATCC Accession Number may also be defined as a strain having a genomic sequence that is greater than 85%, more preferably greater than 90% or more preferably greater than 95% sequence identity to the genome of the indicated NRRL or ATCC Accession Number. A polynucleotide or polynucleotide region (or a polypeptide or polypeptide region) has a certain percentage (for example, 80%, 15 85%, 90%, or 95%) of "sequence identity" to another sequence means that, when aligned, that percentage of bases (or amino acids) are the same in comparing the two sequences. This alignment and the percent homology or sequence identity can be determined using software programs known in the art, for example, those described in Current Protocols in Molecular Biology (F. M. Ausubel et al., eds., 1987) Supplement 30, 20 section 7. 7. 18, Table 7. 7. 1.

NRRL is the abbreviation for the Agricultural Research Service Culture Collection, an international depositary authority for the purposes of depositing microorganism strains under the Budapest treaty on the international recognition of the deposit of microorganisms for the purposes of patent procedure, having the address National 25 Center for Agricultural Utilization Research, Agricultural Research service, U.S. Department of Agriculture, 1815 North university Street, Peroira, Illinois 61604 USA.

ATCC is the abbreviation for the American Type Culture Collection, an international depositary authority for the purposes of depositing microorganism strains under the Budapest treaty on the international recognition of the deposit of microorganisms for the purposes of patent procedure, having the address ATCC Patent Depository, 10801 30 University Blvd., Manassas, VA 10110 USA.

The biological control agents used in the present invention are known in the art as follows:

Bacillus chitinosporus AQ746 (NRRL Accession No. B-21618) (in the following sometimes referred to as B1) is known from WO 98/21966 A2. It is specifically active against nematodes and insects and produces non-exotoxin, non-proteinaceous, active metabolites in its supernatant. Those metabolites are active against nematodes and cockroaches, but inactive against flies, corn rootworm or beet armyworm.

Bacillus mycoides AQ726 (NRRL Accession No. B-21664) (in the following sometimes referred to as B2) and its water-soluble metabolites kill or stunt insects such as corn rootworm larvae and aphids (WO 99/09820 A1).

As described in WO 00/58442 A1 *Bacillus pumilus* QST2808 (NRRL Accession No. B-30087) (in the following sometimes referred to as B3) is able to inhibit a broad range of fungal plant diseases in vivo. Moreover, the combination of this strain with *Bacillus thuringiensis* enhances the insecticidal activity of the latter. Commercially available formulations of this strain are sold under the tradenames SONATA® and BALLAD® Plus from AgraQuest, Inc. USA.

Bacillus pumilus AQ717 (NRRL Accession B-21662) (in the following sometimes referred to as B4) is known from WO 99/10477 A1. It produces a metabolite that exhibits pesticidal activity against corn rootworms, nematodes and beet armyworms.

The bacterial strains *Bacillus sp.* AQ175 (ATCC Accession No. 55608) (in the following sometimes referred to as B5), *Bacillus sp.* AQ 177 (ATCC Accession No. 55609) (in the following sometimes referred to as B6) and *Bacillus sp.* AQ178 (ATCC Accession No. 53522) (in the following sometimes referred to as B7) described in WO 98/21967 A1 are effective in treating and protecting plants from aboveground fungal and bacterial infections.

The metabolite-producing strain *Bacillus subtilis* AQ743 (NRRL Accession No. B-21665) (in the following sometimes referred to as B8) kills or stunts corn rootworm larvae, beet armyworm larvae, fly adults and nematodes (cf. WO 99/09819).

Bacillus subtilis AQ713 (Accession No. B-21661), also named *Bacillus subtilis* QST713, (in the following sometimes referred to as B9) exhibits broad fungicidal and bactericidal activity and also exhibits corn rootworm activity (WO 98/50422 A1). Commercially available formulation of this strain are available under the tradenames SERENADE® Max, SERENADE® Soil, SERENADE® Aso, SERENADE® CPB and RHAPSODY® from AgraQuest, Inc. USA.

Bacillus subtilis AQ153 (ATCC Accession No. 55614) (in the following sometimes referred to as B10) as described in WO 98/21964 A1 is effective in inhibiting growth of plant pathogenic bacteria and fungi.

5 *Bacillus thuringiensis* BD#32 (NRRL Accession No. B-21530) (in the following sometimes referred to as B11) exhibits insecticidal activity (US 5,645,831 A). It produces a non-exotoxin, solvent-extractable, non-proteinaceous metabolite that is 100% effective in killing corn rootworm. The biopesticide produced by this bacterial strain is active against corn rootworm but inactive against flies.

10 According to WO 98/21965 A1 the antibiotic producing strain *Bacillus thuringiensis* AQ52 (NRRL Accession No. B-21619) (in the following sometimes referred to as B12) exhibits broad fungicidal and bactericidal activity.

15 WO 02/02082898 A1 describes endophytic fungi including *Muscodor albus* 620, also known as *Muscodor albus* QST 20799 (NRRL Accession No. 30547) (in the following sometimes referred to as B13) and *Muscodor roseus* A3-5 (NRRL Accession No. 30548) (in the following sometimes referred to as B14) that produce a mixture of volatile antibiotics with activity against fungi, bacteria, insects, mites, and nematodes.

Rhodococcus globerulus AQ719 (NRRL Accession No. B-21663) (in the following sometimes referred to as B15) produces metabolites that exhibits pesticidal activity against corn rootworms (US 6,027,723 A).

20 WO 01/79480 A2 describes a strain of *Streptomyces galbus* (NRRL Accession No. 30232) (in the following sometimes referred to as B16) which shows insecticidal activity against Lepidoptera.

25 The *Streptomyces* sp. strain described in WO 02/26041 A2 (NRRL Accession No. B-30145) (in the following sometimes referred to as B17) exhibits antifungal activity on specific plant pathogens such as *Alternaria*, *Phytophthora*, *Botrytis*, *Rhizoctonia* and *Sclerotinia*.

30 Commercially available formulation of *Bacillus thuringiensis* subspec. *kurstaki* BMP 123 (in the following sometimes referred to as B18) are available under the tradename BARITONE® from AgraQuest, Inc. USA. It exhibits insecticidal activity and is effective on lepidopterous insects, including loopers, armyworms and moths. BARITONE® is distributed subject to EPA Reg. No. 62637-5-69592.

The strains *Bacillus subtilis* AQ30002 (also known as QST30002) (NRRL Accession No. B-50421, deposited on October 5, 2010) (in the following sometimes referred to as B19) and *Bacillus subtilis* AQ30004 (also known as QST30004) (NRRL Accession No. B-50455, deposited on October 5, 2010) (in the following sometimes referred to as B20) are known from WO 2012/087980 A1, which is incorporated herein by reference. As described therein, these BCAs exhibit a broad fungicidal and bactericidal activity. B19 and B20 have a mutation in the *swrA* gene that results in impaired swarming ability and enhanced plant health promotion compared to a strain containing a wildtype *swrA* gene. The mutation causes these BCAs to form a more robust biofilm than the wildtype strain, thereby enhancing its fungicidal and bactericidal activity.

In a preferred embodiment the composition of the present invention is characterized in that the biological control agent is selected from the group consisting of *Bacillus pumilus* (NRRL Accession No. B-30087) and *Bacillus subtilis* AQ713 (NRRL Accession No. B-21661) and/or a mutant of these stains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens.

In another preferred embodiment the composition of the present invention is characterized in that the biological control agent is selected from the group consisting of *Bacillus subtilis* AQ30002 (also known as QST30002) (NRRL Accession No. B-50421), *Bacillus subtilis* AQ30004 (also known as QST30004) (NRRL Accession No. B-50455, or a *Bacillus subtilis* strain having a mutation in the *swrA* gene that results in impaired swarming ability and enhanced plant health promotion compared to a strain containing a wildtype *swrA* gene, and/or a mutant of these stains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens.

In another preferred embodiment the composition of the present invention comprises a combination of at least two biological control agents selected from the group consisting of *Bacillus chittinosporus* AQ746 (NRRL Accession No. B-21618), *Bacillus mycoides* AQ726 (NRRL Accession No. B-21664), *Bacillus pumilus* (NRRL Accession No. B-30087), *Bacillus pumilus* AQ717 (NRRL Accession No. B-21662), *Bacillus sp.* AQ175 (ATCC Accession No. 55608), *Bacillus sp.* AQ177 (ATCC Accession No. 55609), *Bacillus sp.* AQ178 (ATCC Accession No. 53522), *Bacillus subtilis* AQ743 (NRRL Accession No. B-

21665), *Bacillus subtilis* AQ713 (NRRL Accession No. B-21661), *Bacillus subtilis* AQ153 (ATCC Accession No. 55614), *Bacillus thuringiensis* BD#32 (NRRL Accession No. B-21530), *Bacillus thuringiensis* AQ52 (NRRL Accession No. B-21619), *Muscodora albus* 620 (NRRL Accession No. 30547), *Muscodora roseus* A3-5 (NRRL Accession No. 30548),
5 *Rhodococcus globerulus* AQ719 (NRRL Accession No. B-21663), *Streptomyces galbus* (NRRL Accession No. 30232), *Streptomyces* sp. (NRRL Accession No. B-30145), *Bacillus thuringiensis* subspec. *kurstaki* BMP 123, *Bacillus subtilis* AQ30002 (NRRL Accession No. B-50421), and *Bacillus subtilis* AQ 30004 (NRRL Accession No. B-50455) and/or a mutant of these strains having all the identifying characteristics of the respective strain, and/or a
10 metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens.

According to one embodiment of the present invention the biological control agent comprises not only the isolated, pure cultures of the respective microorganisms, but also their suspensions in a whole broth culture or a metabolite-containing supernatant or
15 a purified metabolite obtained from whole broth culture of the strain. "Whole broth culture" refers to a liquid culture containing both cells and media. "Supernatant" refers to the liquid broth remaining when cells grown in broth are removed by centrifugation, filtration, sedimentation, or other means well known in the art.

The above-mentioned metabolites produced by the nonpathogenic microorganisms
20 include antibiotics, enzymes, siderophores and growth promoting agents, for example zwittermicin-A, kanosamine, polyoxine, enzymes such as α -amylase, chitinases, and pectinases, phytohormones and precursors thereof, such as auxines, gibberlin-like substacnes, cytokinin-like compounds, lipopeptides such as iturins, plipastatins or surfactins, e.g. agrastatin A, bacillomycin D, bacilysin, difficidin, macrolactin, fengycin,
25 bacilysin and bacilaene. Preferred metabolites of the above listed are lipopeptides, in particular those produced by *Bacillus pumilus* (NRRL Accession No. B-30087) or *Bacillus subtilis* AQ713 (NRRL Accession No. B-21661). Especially preferred metabolites are Iturin A, Surfactin, Plipstatin and Agrastatin A . An even more preferred metabolite is agrastatin A.

30 According to the invention, the biological control agent may be employed or used in any physiologic state such as active or dormant.

Insecticides

"Insecticides" as well as the term "insecticidal" refers to the ability of a substance to increase mortality or inhibit growth rate of insects. As used herein, the term "insects" includes all organisms in the class "Insecta". The term "pre-adult" insects refers to any form of an organism prior to the adult stage, including, for example, eggs, larvae, and nymphs.

"Nematicides" and "nematicidal" refers to the ability of a substance to increase mortality or inhibit the growth rate of nematodes. In general, the term "nematode" comprises eggs, larvae, juvenile and mature forms of said organism.

10 "Acaricide" and "acaricidal" refers to the ability of a substance to increase mortality or inhibit growth rate of ectoparasites belonging to the class Arachnida, sub-class Acari.

The active ingredients specified herein by their "common name" are known and described, for example, in the Pesticide Manual ("The Pesticide Manual", 14th Ed., British Crop Protection Council 2006) or can be searched in the internet (e.g. <http://www.alanwood.net/pesticides>).

The at least one insecticide according to the present invention is selected from the group consisting of acetylcholinesterase (AChE) inhibitors, GABA-gated chloride channel antagonists, chloride channel activators, and nicotinic acetylcholine receptor (nAChR) channel blockers.

20 In a preferred embodiment the acetylcholinesterase (AChE) inhibitor is selected from the group consisting of carbamates, and organophosphates. Still preferably, the GABA-gated chloride channel antagonist is selected from the group consisting of cyclodiene organochlorines, and phenylpyrazoles (fiproles). Still preferably, the chloride channel activator is an avermectin/milbemycin.

25 In a more preferred embodiment the acetylcholinesterase (AChE) inhibitor is selected from the group consisting of Alanycarb (I1), Aldicarb (I2), Bendiocarb (I3), Benfuracarb (I4), Butocarboxim (I5), Butoxycarboxim (I6), Carbaryl (I7), Carbofuran (I8), Carbosulfan (I9), Ethiofencarb (I10), Fenobucarb (I11), Formetanate (I12), Furathiocarb (I13), Isoprocarb (I14), Methiocarb (I15), Methomyl (I16), Metolcarb (I17), Oxamyl (I18),
30 Pirimicarb (I19), Propoxur (I20), Thiodicarb (I21), Thiofanox (I22), Triazamate (I23), Trimethacarb (I24), XMC (I25), and Xylylcab (I26); Acephate (I27), Azamethiphos (I28),

Azinphos-ethyl (129), Azinphos-methyl (130), Cadusafos (131), Chlorethoxyfos (132), Chlorfenvinphos (133), Chlormephos (134), Chlorpyrifos (135), Chlorpyrifos-methyl (136), Coumaphos (137), Cyanophos (138), Demeton-S-methyl (139), Diazinon (140), Dichlorvos/DDVP (141), Dicrotophos (142), Dimethoate (143), Dimethylvinphos (144),
5 Disulfoton (145), EPN (146), Ethion (147), Ethoprophos (148), Famphur (149), Fenamiphos (150), Fenitrothion (151), Fenthion (152), Fosthiazate (153), Heptenophos (154), Imicyafos (155), Isofenphos (156), Isopropyl O-(methoxyaminothio-phosphoryl) salicylate (157), Isoxathion (158), Malathion (159), Mecarbam (160), Methamidophos (161), Methidathion (162), Mevinphos (163), Monocrotophos (164), Naled (165), Omethoate (166), Oxydemeton-methyl (167), Parathion (168), Parathion-methyl (169), Phenthoate (170), Phorate (171),
10 Phosalone (172), Phosmet (173), Phosphamidon (174), Phoxim (175), Pirimiphos-methyl (176), Profenofos (177), Propetamphos (178), Prothiofos (179), Pyraclofos (180), Pyridaphenthion (181), Quinalphos (182), Sulfotep (183), Tebupirimfos (184), Temephos (185), Terbufos (186), Tetrachlorvinphos (187), Thiometon (188), Triazophos (189), Trichlorfon (190),
15 and Vamidothion (191).

Still preferably, the GABA-gated chloride channel antagonist is selected from the group consisting of Chlordane (192), Endosulfan (193), Ethiprole (194), and Fipronil (195).

Still preferably, the chloride channel activator is selected from the group consisting of Abamectin (1152), Emamectin benzoate (1153), Lepimectin (1154), and Milbernectin
20 (1155).

Still preferably, the nicotinic acetylcholine receptor (nAChR) channel blocker is selected from the group consisting of Bensultap (1187), Cartap hydrochloride (1188), Thiocyclam (1189), and Thiosultap-sodium (1190).

According to a preferred embodiment of the present invention the at least one
25 insecticide is selected from the group consisting of Acephate (127), Chlorpyrifos (135), Carbofuran (18), Methiocarb (115), Profenofos (177), Thiodicarb (121), Ethiprole (194), Fipronil (195), Abamectin (1152), Emamectin-benzoate (1153), and Milbernectin (1155).

According to another preferred embodiment of the present invention the at least one
30 insecticide is selected from the group consisting of Methiocarb (115), Thiodicarb (121), Fipronil (195) and Abamectin (1152).

In one embodiment of the present invention, the insecticide, e.g., the insecticide for use in seed treatment, is selected from the group consisting of Carbofuran (I8), Methiocarb (I15), Thiodicarb (I21) Ethiprole (I94), Fipronil (I95), and Abamectin (I152).

In one embodiment of the present invention the composition comprises a further
5 insecticide which is different from the insecticide and the biological control agent as defined above.

Preferably, this further insecticide is selected from the group consisting of

(1) Acetylcholinesterase (AChE) inhibitors, for example

10 carbamates, e.g. Alanycarb (I1), Aldicarb (I2), Bendiocarb (I3), Benfuracarb (I4), Butocarboxim (I5), Butoxycarboxim (I6), Carbaryl (I7), Carbofuran (I8), Carbosulfan (I9), Ethiofencarb (I10), Fenobucarb (I11), Formetanate (I12), Furathiocarb (I13), Isoprocarb (I14), Methiocarb (I15), Methomyl (I16), Metolcarb (I17), Oxamyl (I18), Pirimicarb (I19), Propoxur (I20), Thiodicarb (I21), Thiofanox (I22), Triazamate (I23), Trimethacarb (I24), XMC (I25), and Xylycarb (I26); or

15 organophosphates, e.g. Acephate (I27), Azamethiphos (I28), Azinphos-ethyl (I29), Azinphos-methyl (I30), Cadusafos (I31), Chlorethoxyfos (I32), Chlorfenvinphos (I33), Chlormephos (I34), Chlorpyrifos (I35), Chlorpyrifos-methyl (I36), Coumaphos (I37), Cyanophos (I38), Demeton-S-methyl (I39), Diazinon (I40), Dichlorvos/DDVP (I41), Dicrotophos (I42), Dimethoate (I43), Dimethylvinphos (I44), Disulfoton (I45), EPN (I46),
20 Ethion (I47), Ethoprophos (I48), Famphur (I49), Fenamiphos (I50), Fenitrothion (I51), Fenthion (I52), Fosthiazate (I53), Heptenophos (I54), Imicyafos (I55), Isofenphos (I56), Isopropyl O-(methoxyaminothio-phosphoryl) salicylate (I57), Isoxathion (I58), Malathion (I59), Mecarbarn (I60), Methamidophos (I61), Methidathion (I62), Mevinphos (I63), Monocrotophos (I64), Naled (I65), Omethoate (I66), Oxydemeton-methyl (I67), Parathion
25 (I68), Parathion-methyl (I69), Phenthoate (I70), Phorate (I71), Phosalone (I72), Phosmet (I73), Phosphamidon (I74), Phoxim (I75), Pirimiphos-methyl (I76), Profenofos (I77), Propetamphos (I78), Prothiofos (I79), Pyraclofos (I80), Pyridaphenthion (I81), Quinalphos (I82), Sulfotep (I83), Tebupirimfos (I84), Temephos (I85), Terbufos (I86), Tetrachlorvinphos (I87), Thiometon (I88), Triazophos (I89), Trichlorfon (I90), and Vamidothion (I91);

30 (2) GABA-gated chloride channel antagonists, for example

cyclodiene organochlorines, e.g. Chlordane (I92) and Endosulfan (I93); or

phenylpyrazoles (fiproles), e.g. Ethiprole (I194) and Fipronil (I195);

(3) Sodium channel modulators / voltage-dependent sodium channel blockers, for example pyrethroids, e.g. Acrinathrin (I196), Allethrin (I197), d-cis-trans Allethrin (I198), d-trans Allethrin (I199), Bifenthrin (I100), Bioallethrin (I101), Bioallethrin S-cyclopentenyl isomer (I102), Bioresmethrin (I103), Cycloprothrin (I104), Cyfluthrin (I105), beta-Cyfluthrin (I106), Cyhalothrin (I107), lambda-Cyhalothrin (I108), gamma-Cyhalothrin (I109), Cypermethrin (I110), alpha-Cypermethrin (I111), beta-Cypermethrin (I112), theta-Cypermethrin (I113), zeta-Cypermethrin (I114), Cyphenothrin [(1R)-trans isomers] (I115), Deltamethrin (I116), Empenthrin [(EZ)-(1R) isomers] (I117), Esfenvalerate (I118), Etofenprox (I119), Fenpropathrin (I120), Fenvalerate (I121), Flucythrinate (I122), Flumethrin (I123), tau-Fluvalinate (I124), Halfenprox (I125), Imiprothrin (I126), Kadethrin (I127), Permethrin (I128), Phenothrin [(1R)-trans isomer] (I129), Prallethrin (I130), Pyrethrine (pyrethrum) (I131), Resmethrin (I132), Silafluofen (I133), Tefluthrin (I134), Tetramethrin (I135), Tetramethrin [(1R) isomers] (I136), Tralomethrin (I137), and Transfluthrin (I138); or DDT (I139); or Methoxychlor (I140);

(4) Nicotinic acetylcholine receptor (nAChR) agonists, for example neonicotinoids, e.g. Acetamiprid (I141), Clothianidin (I142), Dinotefuran (I143), Imidacloprid (I144), Nitenpyram (I145), Thiacloprid (I146), and Thiamethoxam (I147); or Nicotine (I148); or Sulfoxaflor (I149).

(5) Nicotinic acetylcholine receptor (nAChR) allosteric activators, for example spinosyns, e.g. Spinetoram (I150) and Spinosad (I151);

(6) Chloride channel activators, for example avermectins/milbemycins, e.g. Abamectin (I152), Emamectin benzoate (I153), Lepimectin (I154), and Milbemectin (I155);

(7) Juvenile hormone mimics, for example juvenile hormone analogues, e.g. Hydroprene (I156), Kinoprene (I157), and Methoprene (I158); or Fenoxycarb (I159); or Pyriproxyfen (I160);

(8) Miscellaneous non-specific (multi-site) inhibitors, for example alkyl halides, e.g. Methyl bromide (I161) and other alkyl halides; or Chloropicrin (I162); or Sulfuryl fluoride (I163); or Borax (I164); or Tartar emetic (I165);

(9) Selective homopteran feeding blockers, e.g. Pymetrozine (I166); or Flonicamid (I167);

- (10) Mite growth inhibitors, e.g. Clofentezine (I168), Hexythiazox (I169), and Diflovidazin (I170); or Etoxazole (I171);
- (11) Microbial disruptors of insect midgut membranes, e.g. *Bacillus thuringiensis subspecies israelensis* (I172), *Bacillus thuringiensis subspecies aizawai* (I173), *Bacillus thuringiensis subspecies kurstaki* (I174), *Bacillus thuringiensis subspecies tenebrionis* (I175), and B.t. crop proteins: Cry1Ab, Cry1Ac, Cry1Fa, Cry1A.105, Cry2Ab, Vip3A, mCry3A, Cry3Ab, Cry3Bb, Cry34 Ab1/35Ab1 (I176); or *Bacillus sphaericus* (I177);
- 5 (12) Inhibitors of mitochondrial ATP synthase, for example Diafenthuron (I178); or organotin miticides, e.g. Azocyclotin (I179), Cyhexatin (I180), and Fenbutatin oxide
10 (I181); or Propargite (I182); or Tetradifon (I183);
- (13) Uncouplers of oxidative phosphorylation via disruption of the proton gradient, for example Chlorfenapyr (I184), DNOC (I185), and Sulfluramid (I186);
- (14) Nicotinic acetylcholine receptor (nAChR) channel blockers, for example Bensultap (I187), Cartap hydrochloride (I188), Thiocyclam (I189), and Thiosultap-sodium (I190);
- 15 (15) Inhibitors of chitin biosynthesis, type 0, for example Bistrifluron (I191), Chlorfluazuron (I192), Diflubenzuron (I193), Flucycloxuron (I194), Flufenoxuron (I195), Hexaflumuron (I196), Lufenuron (I197), Novaluron (I198), Noviflumuron (I199), Teflubenzuron (I200), and Triflumuron (I201);
- (16) Inhibitors of chitin biosynthesis, type 1, for example Buprofezin (I202);
- 20 (17) Moulting disruptors, for example Cyromazine (I203);
- (18) Ecdysone receptor agonists, for example Chromafenozide (I204), Halofenozide (I205), Methoxyfenozide (I206), and Tebufenozide (I207);
- (19) Octopamine receptor agonists, for example Amitraz (I208);
- (20) Mitochondrial complex III electron transport inhibitors, for example Hydramethylnon
25 (I209); or Acequinocyl (I210); or Fluacrypyrim (I211);
- (21) Mitochondrial complex I electron transport inhibitors, for example
- METI acaricides, e.g. Fenazaquin (I212), Fenpyroximate (I213), Pyrimidifen (I214), Pyridaben (I215), Tebufenpyrad (I216), and Tolfenpyrad (I217); or Rotenone (Derris) (I218);

(22) Voltage-dependent sodium channel blockers, e.g. Indoxacarb (I219); or Metaflumizone (I220);

(23) Inhibitors of acetyl CoA carboxylase, for example tetronic and tetramic acid derivatives, e.g. Spirodiclofen (I221), Spiromesifen (I222), and Spirotetramat (I223);

5 (24) Mitochondrial complex IV electron transport inhibitors, for example phosphines, e.g. Aluminium phosphide (I224), Calcium phosphide (I225), Phosphine (I226), and Zinc phosphide (I227); or Cyanide (I228);

(25) Mitochondrial complex II electron transport inhibitors, for example beta-ketonitrile derivatives, e.g. Cyenopyrafen (I229) and Cyflumetofen (I230);

10 (28) Ryanodine receptor modulators, for example diamides, e.g. Chlorantraniliprole (I231), Cyantraniliprole (I232), and Flubendiamide (I233);

Further active ingredients with unknown or uncertain mode of action, for example Amidoflumeť (I234), Azadirachtin (I235), Benclonthiaz (I236), Benzoximate (I237), Bifenazate (I238), Bromopropylate (I239), Chinomethionat (I240), Cryolite (I241), Dicofol (I242), Diflovidazin (I243), Fluensulfone (I244), Flufenerim (I245), Flutiprole (I246),
 15 Fluopyram (I247), Fufenozide (I248), Imidaclothiz (I249), Iprodione (I250), Meperfluthrin (I251), Pyridalyl (I252), Pyrifluquinazon (I253), Tetramethylfluthrin (I254), and iodomethane (I255); furthermore products based on *Bacillus firmus* (including but not limited to strain CNCM I-1582, such as, for example, VOTIVOe, BioNem) (I256) or one of the following

20 known active compounds: 3-bromo-N-{2-bromo-4-chloro-6-[[1-cyclopropylethyl]carbonyl]phenyl}-1-(3-chloropyridin-2-yl)-1H-pyrazole-5-carboxamide (I257) (known from WO2005/077934), 4-[[[6-bromopyridin-3-yl]methyl][(2-fluoroethyl)amino]furan-2(5H)-one (I258) (known from WO2007/115644), 4-[[[6-fluoropyridin-3-yl]methyl][(2,2-difluoroethyl)amino]furan-2(5H)-one (I259) (known from
 25 WO2007/115644), 4-[[[2-chloro-1,3-thiazol-5-yl]methyl][(2-fluoroethyl)amino]furan-2(5H)-one (I260) (known from WO2007/115644), 4-[[[6-chloropyridin-3-yl]methyl][(2-fluoroethyl)amino]furan-2(5H)-one (I261) (known from WO2007/115644), Flupyradifurone (I262), 4-[[[6-chlor-5-fluoropyridin-3-yl]methyl][(methyl)amino]furan-2(5H)-one (I263) (known from
 30 WO2007/115643), 4-[[[5,6-dichloropyridin-3-yl]methyl][(2-fluoroethyl)amino]furan-2(5H)-one (I264) (known from WO2007/115646), 4-[[[6-chloro-5-fluoropyridin-3-yl]methyl][(cyclopropyl)amino]furan-2(5H)-one (I265) (known from
 WO2007/115643), 4-[[[6-chloropyridin-3-yl]methyl][(cyclopropyl)amino]furan-2(5H)-one

(I266) (known from EP-A-0 539 588), 4-[[[6-chloropyridin-3-yl)methyl](methyl)amino]furan-2(5H)-one (I267) (known from EP-A-0 539 588), {[1-(6-chloropyridin-3-yl)ethyl](methyl)oxido- λ 4-sulfanylidene}cyanamide (I268) (known from WO2007/149134) and its diastereomers {[[1R]-1-(6-chloropyridin-3-yl)ethyl](methyl)oxido- λ 4-sulfanylidene}cyanamide (A) (I269), and {[[1S]-1-(6-chloropyridin-3-yl)ethyl](methyl)oxido- λ 4-sulfanylidene}cyanamide (B) (I270) (also known from WO2007/149134) as well as diastereomers [(R)-methyl(oxido){[1R]-1-[6-(trifluoromethyl)pyridin-3-yl]ethyl}- λ 4-sulfanylidene]cyanamide (A1) (I271), and [(S)-methyl(oxido){[1S]-1-[6-(trifluoromethyl)pyridin-3-yl]ethyl}- λ 4-sulfanylidene]cyanamide (A2) (I272), referred to as group of diastereomers A (known from WO2010/074747, WO2010/074751), [(R)-methyl(oxido){[1S]-1-[6-(trifluoromethyl)pyridin-3-yl]ethyl}- λ 4-sulfanylidene]cyanamide (B1) (I273), and [(S)-methyl(oxido){[1R]-1-[6-(trifluoromethyl)pyridin-3-yl]ethyl}- λ 4-sulfanylidene]cyanamide (B2) (I274), referred to as group of diastereomers B (also known from WO2010/074747, WO2010/074751), and 11-(4-chloro-2,6-dimethylphenyl)-12-hydroxy-1,4-dioxo-9-azadispiro[4.2.4.2]tetradec-11-en-10-one (I275) (known from WO2006/089633), 3-(4'-fluoro-2,4-dimethylbiphenyl-3-yl)-4-hydroxy-8-oxa-1-azaspiro[4.5]dec-3-en-2-one (I276) (known from WO2008/067911), 1-{2-fluoro-4-methyl-5-[[2,2,2-trifluoroethyl]sulfinyl]phenyl}-3-(trifluoromethyl)-1H-1,2,4-triazol-5-amine (I277) (known from WO2006/043635), Afidopyropen [(3S,4aR,12R,12aS,12bS)-3-[[cyclopropylcarbonyl]oxy]-6,12-dihydroxy-4,12b-dimethyl-11-oxo-9-(pyridin-3-yl)-1,3,4,4a,5,6,6a,12,12a,12b-decahydro-2H,11H-benzo[f]pyrano[4,3-b]chromen-4-yl]methyl cyclopropanecarboxylate (I278) (known from WO2008/066153), 2-cyano-3-(difluoromethoxy)-N,N-dimethylbenzenesulfonamide (I279) (known from WO2006/056433), 2-cyano-3-(difluoromethoxy)-N-methylbenzenesulfonamide (I280) (known from WO2006/100288), 2-cyano-3-(difluoromethoxy)-N-ethylbenzenesulfonamide (I281) (known from WO2005/035486), 4-(difluoromethoxy)-N-ethyl-N-methyl-1,2-benzothiazol-3-amine 1,1-dioxide (I282) (known from WO2007/057407), N-[1-(2,3-dimethylphenyl)-2-(3,5-dimethylphenyl)ethyl]-4,5-dihydro-1,3-thiazol-2-amine (I283) (known from WO2008/104503), {1'-[[2E]-3-(4-chlorophenyl)prop-2-en-1-yl]-5-fluorospiro[indole-3,4'-piperidin]-1(2H)-yl}-(2-chloropyridin-4-yl)methanone (I284) (known from WO2003/106457), 3-(2,5-dimethylphenyl)-4-hydroxy-8-methoxy-1,8-diazaspiro[4.5]dec-3-en-2-one (I285) (known from WO2009/049851), 3-(2,5-dimethylphenyl)-8-methoxy-2-oxo-1,8-diazaspiro[4.5]dec-3-en-4-yl ethyl carbonate (I286) (known from WO2009/049851), 4-(but-2-yn-1-yloxy)-6-(3,5-dimethylpiperidin-1-yl)-5-fluoropyrimidine (I287) (known from WO2004/099160), (2,2,3,3,4,4,5,5-

octafluoropentyl)(3,3,3-trifluoropropyl)malononitrile (I288) (known from WO2005/063094),
(2,2,3,3,4,4,5,5-octafluoropentyl)(3,3,4,4,4-pentafluorobutyl)malononitrile (I289) (known
from WO2005/063094), 8-[2-(cyclopropylmethoxy)-4-(trifluoromethyl)phenoxy]-3-[6-
(trifluoromethyl)pyridazin-3-yl]-3-azabicyclo[3.2.1]octane (I290) (known from
5 WO2007/040280), Flometoquin (I291), PF1364 (CAS-Reg.No. 1204776-60-2) (I292)
(known from JP2010/018586), 5-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4,5-dihydro-
1,2-oxazol-3-yl]-2-(1H-1,2,4-triazol-1-yl)benzonitrile (I293) (known from WO2007/075459),
5-[5-(2-chloropyridin-4-yl)-5-(trifluoromethyl)-4,5-dihydro-1,2-oxazol-3-yl]-2-(1H-1,2,4-
triazol-1-yl)benzonitrile (I294) (known from WO2007/075459), 4-[5-(3,5-dichlorophenyl)-5-
10 (trifluoromethyl)-4,5-dihydro-1,2-oxazol-3-yl]-2-methyl-N-{2-oxo-2-[(2,2,2-
trifluoroethyl)amino]ethyl}benzamide (I295) (known from WO2005/085216), 4-[[[6-
chloropyridin-3-yl)methyl](cyclopropyl)amino]-1,3-oxazol-2(5H)-one (I296), 4-[[[6-
chloropyridin-3-yl)methyl](2,2-difluoroethyl)amino]-1,3-oxazol-2(5H)-one (I297), 4-[[[6-
chloropyridin-3-yl)methyl](ethyl)amino]-1,3-oxazol-2(5H)-one (I298), 4-[[[6-chloropyridin-
15 3-yl)methyl](methyl)amino]-1,3-oxazol-2(5H)-one (I299) (all known from
WO2010/005692), Pyflubumide N-[4-(1,1,1,3,3,3-hexafluoro-2-methoxypropan-2-yl)-3-
isobutylphenyl]-N-isobutyryl-1,3,5-trimethyl-1H-pyrazole-4-carboxamide (I300) (known
from WO2002/096882), methyl 2-[2-([3-bromo-1-(3-chloropyridin-2-yl)-1H-pyrazol-5-
yl]carbonyl)amino)-5-chloro-3-methylbenzoyl]-2-methylhydrazinecarboxylate (I301)
20 (known from WO2005/085216), methyl 2-[2-([3-bromo-1-(3-chloropyridin-2-yl)-1H-
pyrazol-5-yl]carbonyl)amino)-5-cyano-3-methylbenzoyl]-2-ethylhydrazinecarboxylate
(I302) (known from WO2005/085216), methyl 2-[2-([3-bromo-1-(3-chloropyridin-2-yl)-1H-
pyrazol-5-yl]carbonyl)amino)-5-cyano-3-methylbenzoyl]-2-methylhydrazinecarboxylate
(I303) (known from WO2005/085216), methyl 2-[3,5-dibromo-2-([3-bromo-1-(3-
25 chloropyridin-2-yl)-1H-pyrazol-5-yl]carbonyl)amino]benzoyl]-1,2-
diethylhydrazinecarboxylate (I304) (known from WO2005/085216), methyl 2-[3,5-
dibromo-2-([3-bromo-1-(3-chloropyridin-2-yl)-1H-pyrazol-5-yl]carbonyl)amino]benzoyl]-
2-ethylhydrazinecarboxylate (I305) (known from WO2005/085216), (5RS,7RS;5RS,7SR)-1-
(6-chloro-3-pyridylmethyl)-1,2,3,5,6,7-hexahydro-7-methyl-8-nitro-5-propoxyimidazo[1,2-
30 a]pyridine (I306) (known from WO2007/101369), 2-{6-[2-(5-fluoropyridin-3-yl)-1,3-thiazol-
5-yl]pyridin-2-yl}pyrimidine (I307) (known from WO2010/006713), 2-{6-[2-(pyridin-3-yl)-
1,3-thiazol-5-yl]pyridin-2-yl}pyrimidine (I308) (known from WO2010/006713), 1-(3-
chloropyridin-2-yl)-N-[4-cyano-2-methyl-6-(methylcarbamoyl)phenyl]-3-[[5-
(trifluoromethyl)-1H-tetrazol-1-yl]methyl]-1H-pyrazole-5-carboxamide (I309) (known from
35 WO2010/069502), 1-(3-chloropyridin-2-yl)-N-[4-cyano-2-methyl-6-

(methylcarbamoyl)phenyl]-3-[[5-(trifluoromethyl)-2H-tetrazol-2-yl]methyl]-1H-pyrazole-5-carboxamide (I310) (known from WO2010/069502), N-[2-(tert-butylcarbamoyl)-4-cyano-6-methylphenyl]-1-(3-chloropyridin-2-yl)-3-[[5-(trifluoromethyl)-1H-tetrazol-1-yl]methyl]-1H-pyrazole-5-carboxamide (I311) (known from WO2010/069502), N-[2-(tert-butylcarbamoyl)-4-cyano-6-methylphenyl]-1-(3-chloropyridin-2-yl)-3-[[5-(trifluoromethyl)-2H-tetrazol-2-yl]methyl]-1H-pyrazole-5-carboxamide (I312) (known from WO2010/069502), (1E)-N-[[6-chloropyridin-3-yl]methyl]-N'-cyano-N-(2,2-difluoroethyl)ethanimidamide (I313) (known from WO2008/009360), N-[2-(5-amino-1,3,4-thiadiazol-2-yl)-4-chloro-6-methylphenyl]-3-bromo-1-(3-chloropyridin-2-yl)-1H-pyrazole-5-carboxamide (I314) (known from CN102057925), and methyl 2-[3,5-dibromo-2-{{[3-bromo-1-(3-chloropyridin-2-yl)-1H-pyrazol-5-yl]carbonyl}amino}benzoyl]-2-ethyl-1-methylhydrazinecarboxylate (I315) (known from WO2011/049233).

In a preferred embodiment of the present invention the insecticide is a synthetic insecticide. As used herein, the term "synthetic" defines a compound that has not been obtained from a biological control agent. Especially a synthetic insecticide or fungicide is no metabolite of the biological control agents according to the present invention.

Compositions according to the present invention

According to the present invention the composition comprises at least one biological control agent selected from the group consisting of *Bacillus chitinosporus* AQ746 (NRRL Accession No. B-21618), *Bacillus mycolides* AQ726 (NRRL Accession No. B-21664), *Bacillus pumilus* (NRRL Accession No. B-30087), *Bacillus pumilus* AQ717 (NRRL Accession No. B-21662), *Bacillus sp.* AQ175 (ATCC Accession No. 55608), *Bacillus sp.* AQ177 (ATCC Accession No. 55609), *Bacillus sp.* AQ178 (ATCC Accession No. 53522), *Bacillus subtilis* AQ743 (NRRL Accession No. B-21665), *Bacillus subtilis* AQ713 (NRRL Accession No. B-21661), *Bacillus subtilis* AQ153 (ATCC Accession No. 55614), *Bacillus thuringiensis* BD#32 (NRRL Accession No. B-21530), *Bacillus thuringiensis* AQ52 (NRRL Accession No. B-21619), *Muscodor albus* 620 (NRRL Accession No. 30547), *Muscodor roseus* A3-5 (NRRL Accession No. 30548), *Rhodococcus globerulus* AQ719 (NRRL Accession No. B-21663), *Streptomyces galbus* (NRRL Accession No. 30232), *Streptomyces sp.* (NRRL Accession No. B-30145), *Bacillus thuringiensis subspec. kurstaki* BMP 123, *Bacillus subtilis* AQ30002 (NRRL Accession No. B-50421), and *Bacillus subtilis* AQ 30004 (NRRL Accession No. B-50455) and/or a mutant of these strains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits

activity against insects, mites, nematodes and/or phytopathogens and at least one insecticide selected from the group consisting of acetylcholinesterase (AChE) inhibitors, GABA-gated chloride channel antagonists, chloride channel activators and nicotinic acetylcholine receptor (nAChR) channel blockers, in a synergistically effective amount.

- 5 A "synergistically effective amount" according to the present invention represents a quantity of a combination of a biological control agent and an insecticide that is statistically significantly more effective against insects, mites, nematodes and/or phytopathogens than the biological control agent or the insecticide only.

In a preferred embodiment the composition according to the present invention
10 comprises the following combinations:

B1+11, B1+12, B1+13, B1+14, B1+15, B1+16, B1+17, B1+18, B1+19, B1+B10, B1+111,
B1+112, B1+113, B1+114, B1+115, B1+116, B1+117, B1+118, B1+119, B1+120, B1+121,
B1+ 122, B1+123, B1+124, B1+125, B1+126, B1+127, B1+128, B1+129, B1+130, B1+131,
B1+132, B1+133, B1+134, B1+135, B1+136, B1+137, B1+138, B1+139, B1+140, B1+141,
15 B1+142, B1+143, B1+144, B1+145, B1+146, B1+147, B1+148, B1+149, B1+150, B1+151,
B1+152, B1+153, B1+154, B1+155, B1+156, B1+157, B1+158, B1+159, B1+160, B1+161,
B1+ 162, B1+163, B1+164, B1+165, B1+166, B1+167, B1+168, B1+169, B1+170, B1+171,
B1+172, B1+173, B1+174, B1+175, B1+176, B1+177, B1+178, B1+179, B1+180, B1+181,
B1+182, B1+183, B1+184, B1+185, B1+186, B1+187, B1+188, B1+189, B1+190, B1+191,
20 B1+192, B1+193, B1+194, B1+195, B1+1152, B1+1153, B1+1154, B1+1155, B1+1187,
B1+1188, B1+1189, B1+1190;

B2+11, B2+12, B2+13, B2+14, B2+15, B2+16, B2+17, B2+18, B2+19, B2+B20, B2+111,
B2+112, B2+113, B2+114, B2+115, B2+116, B2+117, B2+118, B2+119, B2+120, B2+121,
B2+ 122, B2+123, B2+124, B2+125, B2+126, B2+127, B2+128, B2+129, B2+130, B2+131,
25 B2+132, B2+133, B2+134, B2+135, B2+136, B2+137, B2+138, B2+139, B2+140, B2+141,
B2+142, B2+143, B2+144, B2+145, B2+146, B2+147, B2+148, B2+149, B2+150, B2+151,
B2+152, B2+153, B2+154, B2+155, B2+156, B2+157, B2+158, B2+159, B2+160, B2+161,
B2+ 162, B2+163, B2+164, B2+165, B2+166, B2+167, B2+168, B2+169, B2+170, B2+171,
B2+172, B2+173, B2+174, B2+175, B2+176, B2+177, B2+178, B2+179, B2+180, B2+181,
30 B2+182, B2+183, B2+184, B2+185, B2+186, B2+187, B2+188, B2+189, B2+190, B2+191,
B2+192, B2+193, B2+194, B2+195, B2+1152, B2+1153, B2+1154, B2+1155, B2+1187,
B2+1188, B2+1189, B2+1190;

B3+11, B3+12, B3+13, B3+14, B3+15, B3+16, B3+17, B3+18, B3+19, B3+B30, B3+111,
B3+112, B3+113, B3+114, B3+115, B3+116, B3+117, B3+118, B3+119, B3+120, B3+121,
B3+ 122, B3+123, B3+124, B3+125, B3+126, B3+127, B3+128, B3+129, B3+130, B3+131,
B3+132, B3+133, B3+134, B3+135, B3+136, B3+137, B3+138, B3+139, B3+140, B3+141,
5 B3+142, B3+143, B3+144, B3+145, B3+146, B3+147, B3+148, B3+149, B3+150, B3+151,
B3+152, B3+153, B3+154, B3+155, B3+156, B3+157, B3+158, B3+159, B3+160, B3+161,
B3+ 162, B3+163, B3+164, B3+165, B3+166, B3+167, B3+168, B3+169, B3+170, B3+171,
B3+172, B3+173, B3+174, B3+175, B3+176, B3+177, B3+178, B3+179, B3+180, B3+181,
B3+182, B3+183, B3+184, B3+185, B3+186, B3+187, B3+188, B3+189, B3+190, B3+191,
10 B3+192, B3+193, B3+194, B3+195, B3+1152, B3+1153, B3+1154, B3+1155, B3+1187,
B3+1188, B3+1189, B3+1190;

B4+11, B4+12, B4+13, B4+14, B4+15, B4+16, B4+17, B4+18, B4+19, B4+B40, B4+111,
B4+112, B4+113, B4+114, B4+115, B4+116, B4+117, B4+118, B4+119, B4+120, B4+121,
B4+ 122, B4+123, B4+124, B4+125, B4+126, B4+127, B4+128, B4+129, B4+130, B4+131,
15 B4+132, B4+133, B4+134, B4+135, B4+136, B4+137, B4+138, B4+139, B4+140, B4+141,
B4+142, B4+143, B4+144, B4+145, B4+146, B4+147, B4+148, B4+149, B4+150, B4+151,
B4+152, B4+153, B4+154, B4+155, B4+156, B4+157, B4+158, B4+159, B4+160, B4+161,
B4+ 162, B4+163, B4+164, B4+165, B4+166, B4+167, B4+168, B4+169, B4+170, B4+171,
B4+172, B4+173, B4+174, B4+175, B4+176, B4+177, B4+178, B4+179, B4+180, B4+181,
20 B4+182, B4+183, B4+184, B4+185, B4+186, B4+187, B4+188, B4+189, B4+190, B4+191,
B4+192, B4+193, B4+194, B4+195, B4+1152, B4+1153, B4+1154, B4+1155, B4+1187,
B4+1188, B4+1189, B4+1190;

B5+11, B5+12, B5+13, B5+14, B5+15, B5+16, B5+17, B5+18, B5+19, B5+B50, B5+111,
B5+112, B5+113, B5+114, B5+115, B5+116, B5+117, B5+118, B5+119, B5+120, B5+121,
25 B5+ 122, B5+123, B5+124, B5+125, B5+126, B5+127, B5+128, B5+129, B5+130, B5+131,
B5+132, B5+133, B5+134, B5+135, B5+136, B5+137, B5+138, B5+139, B5+140, B5+141,
B5+142, B5+143, B5+144, B5+145, B5+146, B5+147, B5+148, B5+149, B5+150, B5+151,
B5+152, B5+153, B5+154, B5+155, B5+156, B5+157, B5+158, B5+159, B5+160, B5+161,
B5+ 162, B5+163, B5+164, B5+165, B5+166, B5+167, B5+168, B5+169, B5+170, B5+171,
30 B5+172, B5+173, B5+174, B5+175, B5+176, B5+177, B5+178, B5+179, B5+180, B5+181,
B5+182, B5+183, B5+184, B5+185, B5+186, B5+187, B5+188, B5+189, B5+190, B5+191,
B5+192, B5+193, B5+194, B5+195, B5+1152, B5+1153, B5+1154, B5+1155, B5+1187,
B5+1188, B5+1189, B5+1190;

B6+11, B6+12, B6+13, B6+14, B6+15, B6+16, B6+17, B6+18, B6+19, B6+B60, B6+111,
B6+112, B6+113, B6+114, B6+115, B6+116, B6+117, B6+118, B6+119, B6+120, B6+121,
B6+ 122, B6+123, B6+124, B6+125, B6+126, B6+127, B6+128, B6+129, B6+130, B6+131,
B6+132, B6+133, B6+134, B6+135, B6+136, B6+137, B6+138, B6+139, B6+140, B6+141,
5 B6+142, B6+143, B6+144, B6+145, B6+146, B6+147, B6+148, B6+149, B6+150, B6+151,
B6+152, B6+153, B6+154, B6+155, B6+156, B6+157, B6+158, B6+159, B6+160, B6+161,
B6+ 162, B6+163, B6+164, B6+165, B6+166, B6+167, B6+168, B6+169, B6+170, B6+171,
B6+172, B6+173, B6+174, B6+175, B6+176, B6+177, B6+178, B6+179, B6+180, B6+181,
B6+182, B6+183, B6+184, B6+185, B6+186, B6+187, B6+188, B6+189, B6+190, B6+191,
10 B6+192, B6+193, B6+194, B6+195, B6+1152, B6+1153, B6+1154, B6+1155, B6+1187,
B6+1188, B6+1189, B6+1190;

B7+11, B7+12, B7+13, B7+14, B7+15, B7+16, B7+17, B7+18, B7+19, B7+B70, B7+111,
B7+112, B7+113, B7+114, B7+115, B7+116, B7+117, B7+118, B7+119, B7+120, B7+121,
B7+ 122, B7+123, B7+124, B7+125, B7+126, B7+127, B7+128, B7+129, B7+130, B7+131,
15 B7+132, B7+133, B7+134, B7+135, B7+136, B7+137, B7+138, B7+139, B7+140, B7+141,
B7+142, B7+143, B7+144, B7+145, B7+146, B7+147, B7+148, B7+149, B7+150, B7+151,
B7+152, B7+153, B7+154, B7+155, B7+156, B7+157, B7+158, B7+159, B7+160, B7+161,
B7+ 162, B7+163, B7+164, B7+165, B7+166, B7+167, B7+168, B7+169, B7+170, B7+171,
B7+172, B7+173, B7+174, B7+175, B7+176, B7+177, B7+178, B7+179, B7+180, B7+181,
20 B7+182, B7+183, B7+184, B7+185, B7+186, B7+187, B7+188, B7+189, B7+190, B7+191,
B7+192, B7+193, B7+194, B7+195, B7+1152, B7+1153, B7+1154, B7+1155, B7+1187,
B7+1188, B7+1189, B7+1190;

B8+11, B8+12, B8+13, B8+14, B8+15, B8+16, B8+17, B8+18, B8+19, B8+B80, B8+111,
B8+112, B8+113, B8+114, B8+115, B8+116, B8+117, B8+118, B8+119, B8+120, B8+121,
25 B8+ 122, B8+123, B8+124, B8+125, B8+126, B8+127, B8+128, B8+129, B8+130, B8+131,
B8+132, B8+133, B8+134, B8+135, B8+136, B8+137, B8+138, B8+139, B8+140, B8+141,
B8+142, B8+143, B8+144, B8+145, B8+146, B8+147, B8+148, B8+149, B8+150, B8+151,
B8+152, B8+153, B8+154, B8+155, B8+156, B8+157, B8+158, B8+159, B8+160, B8+161,
B8+ 162, B8+163, B8+164, B8+165, B8+166, B8+167, B8+168, B8+169, B8+170, B8+171,
30 B8+172, B8+173, B8+174, B8+175, B8+176, B8+177, B8+178, B8+179, B8+180, B8+181,
B8+182, B8+183, B8+184, B8+185, B8+186, B8+187, B8+188, B8+189, B8+190, B8+191,
B8+192, B8+193, B8+194, B8+195, B8+1152, B8+1153, B8+1154, B8+1155, B8+1187,
B8+1188, B8+1189, B8+1190;

B9+I1, B9+I2, B9+I3, B9+I4, B9+I5, B9+I6, B9+I7, B9+I8, B9+I9, B9+B90, B9+I11,
B9+I12, B9+I13, B9+I14, B9+I15, B9+I16, B9+I17, B9+I18, B9+I19, B9+I20, B9+I21,
B9+ I22, B9+I23, B9+I24, B9+I25, B9+I26, B9+I27, B9+I28, B9+I29, B9+I30, B9+I31,
B9+I32, B9+I33, B9+I34, B9+I35, B9+I36, B9+I37, B9+I38, B9+I39, B9+I40, B9+I41,
5 B9+I42, B9+I43, B9+I44, B9+I45, B9+I46, B9+I47, B9+I48, B9+I49, B9+I50, B9+I51,
B9+I52, B9+I53, B9+I54, B9+I55, B9+I56, B9+I57, B9+I58, B9+I59, B9+I60, B9+I61,
B9+ I62, B9+I63, B9+I64, B9+I65, B9+I66, B9+I67, B9+I68, B9+I69, B9+I70, B9+I71,
B9+I72, B9+I73, B9+I74, B9+I75, B9+I76, B9+I77, B9+I78, B9+I79, B9+I80, B9+I81,
B9+I82, B9+I83, B9+I84, B9+I85, B9+I86, B9+I87, B9+I88, B9+I89, B9+I90, B9+I91,
10 B9+I92, B9+I93, B9+I94, B9+I95, B9+I152, B9+I153, B9+I154, B9+I155, B9+I187,
B9+I188, B9+I189, B9+I190;

B10+I1, B10+I2, B10+I3, B10+I4, B10+I5, B10+I6, B10+I7, B10+I8, B10+I9, B10+B100,
B10+I11, B10+I12, B10+I13, B10+I14, B10+I15, B10+I16, B10+I17, B10+I18, B10+I19,
B10+I20, B10+I21, B10+ I22, B10+I23, B10+I24, B10+I25, B10+I26, B10+I27, B10+I28,
15 B10+I29, B10+I30, B10+I31, B10+I32, B10+I33, B10+I34, B10+I35, B10+I36, B10+I37,
B10+I38, B10+I39, B10+I40, B10+I41, B10+I42, B10+I43, B10+I44, B10+I45, B10+I46,
B10+I47, B10+I48, B10+I49, B10+I50, B10+I51, B10+I52, B10+I53, B10+I54, B10+I55,
B10+I56, B10+I57, B10+I58, B10+I59, B10+I60, B10+I61, B10+ I62, B10+I63, B10+I64,
B10+I65, B10+I66, B10+I67, B10+I68, B10+I69, B10+I70, B10+I71, B10+I72, B10+I73,
20 B10+I74, B10+I75, B10+I76, B10+I77, B10+I78, B10+I79, B10+I80, B10+I81, B10+I82,
B10+I83, B10+I84, B10+I85, B10+I86, B10+I87, B10+I88, B10+I89, B10+I90, B10+I91,
B10+I92, B10+I93, B10+I94, B10+I95, B10+I152, B10+I153, B10+I154, B10+I155,
B10+I187, B10+I188, B10+I189, B10+I190;

B11+I1, B11+I2, B11+I3, B11+I4, B11+I5, B11+I6, B11+I7, B11+I8, B11+I9, B11+B110,
25 B11+I11, B11+I12, B11+I13, B11+I14, B11+I15, B11+I16, B11+I17, B11+I18, B11+I19,
B11+I20, B11+I21, B11+ I22, B11+I23, B11+I24, B11+I25, B11+I26, B11+I27, B11+I28,
B11+I29, B11+I30, B11+I31, B11+I32, B11+I33, B11+I34, B11+I35, B11+I36, B11+I37,
B11+I38, B11+I39, B11+I40, B11+I41, B11+I42, B11+I43, B11+I44, B11+I45, B11+I46,
B11+I47, B11+I48, B11+I49, B11+I50, B11+I51, B11+I52, B11+I53, B11+I54, B11+I55,
30 B11+I56, B11+I57, B11+I58, B11+I59, B11+I60, B11+I61, B11+ I62, B11+I63, B11+I64,
B11+I65, B11+I66, B11+I67, B11+I68, B11+I69, B11+I70, B11+I71, B11+I72, B11+I73,
B11+I74, B11+I75, B11+I76, B11+I77, B11+I78, B11+I79, B11+I80, B11+I81, B11+I82,
B11+I83, B11+I84, B11+I85, B11+I86, B11+I87, B11+I88, B11+I89, B11+I90, B11+I91,

B11+I92, B11+I93, B11+I94, B11+I95, B11+I152, B11+I153, B11+I154, B11+I155,
B11+I187, B11+I188, B11+I189, B11+I190;

B12+I1, B12+I2, B12+I3, B12+I4, B12+I5, B12+I6, B12+I7, B12+I8, B12+I9, B12+B120,
B12+I11, B12+I12, B12+I13, B12+I14, B12+I15, B12+I16, B12+I17, B12+I18, B12+I19,
5 B12+I20, B12+I21, B12+ I22, B12+I23, B12+I24, B12+I25, B12+I26, B12+I27, B12+I28,
B12+I29, B12+I30, B12+I31, B12+I32, B12+I33, B12+I34, B12+I35, B12+I36, B12+I37,
B12+I38, B12+I39, B12+I40, B12+I41, B12+I42, B12+I43, B12+I44, B12+I45, B12+I46,
B12+I47, B12+I48, B12+I49, B12+I50, B12+I51, B12+I52, B12+I53, B12+I54, B12+I55,
B12+I56, B12+I57, B12+I58, B12+I59, B12+I60, B12+I61, B12+ I62, B12+I63, B12+I64,
10 B12+I65, B12+I66, B12+I67, B12+I68, B12+I69, B12+I70, B12+I71, B12+I72, B12+I73,
B12+I74, B12+I75, B12+I76, B12+I77, B12+I78, B12+I79, B12+I80, B12+I81, B12+I82,
B12+I83, B12+I84, B12+I85, B12+I86, B12+I87, B12+I88, B12+I89, B12+I90, B12+I91,
B12+I92, B12+I93, B12+I94, B12+I95, B12+I152, B12+I153, B12+I154, B12+I155,
B12+I187, B12+I188, B12+I189, B12+I190;

15 B13+I1, B13+I2, B13+I3, B13+I4, B13+I5, B13+I6, B13+I7, B13+I8, B13+I9, B13+B130,
B13+I11, B13+I12, B13+I13, B13+I14, B13+I15, B13+I16, B13+I17, B13+I18, B13+I19,
B13+I20, B13+I21, B13+ I22, B13+I23, B13+I24, B13+I25, B13+I26, B13+I27, B13+I28,
B13+I29, B13+I30, B13+I31, B13+I32, B13+I33, B13+I34, B13+I35, B13+I36, B13+I37,
B13+I38, B13+I39, B13+I40, B13+I41, B13+I42, B13+I43, B13+I44, B13+I45, B13+I46,
20 B13+I47, B13+I48, B13+I49, B13+I50, B13+I51, B13+I52, B13+I53, B13+I54, B13+I55,
B13+I56, B13+I57, B13+I58, B13+I59, B13+I60, B13+I61, B13+ I62, B13+I63, B13+I64,
B13+I65, B13+I66, B13+I67, B13+I68, B13+I69, B13+I70, B13+I71, B13+I72, B13+I73,
B13+I74, B13+I75, B13+I76, B13+I77, B13+I78, B13+I79, B13+I80, B13+I81, B13+I82,
B13+I83, B13+I84, B13+I85, B13+I86, B13+I87, B13+I88, B13+I89, B13+I90, B13+I91,
25 B13+I92, B13+I93, B13+I94, B13+I95, B13+I152, B13+I153, B13+I154, B13+I155,
B13+I187, B13+I188, B13+I189, B13+I190;

B14+I1, B14+I2, B14+I3, B14+I4, B14+I5, B14+I6, B14+I7, B14+I8, B14+I9, B14+B140,
B14+I11, B14+I12, B14+I13, B14+I14, B14+I15, B14+I16, B14+I17, B14+I18, B14+I19,
B14+I20, B14+I21, B14+ I22, B14+I23, B14+I24, B14+I25, B14+I26, B14+I27, B14+I28,
30 B14+I29, B14+I30, B14+I31, B14+I32, B14+I33, B14+I34, B14+I35, B14+I36, B14+I37,
B14+I38, B14+I39, B14+I40, B14+I41, B14+I42, B14+I43, B14+I44, B14+I45, B14+I46,
B14+I47, B14+I48, B14+I49, B14+I50, B14+I51, B14+I52, B14+I53, B14+I54, B14+I55,
B14+I56, B14+I57, B14+I58, B14+I59, B14+I60, B14+I61, B14+ I62, B14+I63, B14+I64,

B14+I65, B14+I66, B14+I67, B14+I68, B14+I69, B14+I70, B14+I71, B14+I72, B14+I73,
B14+I74, B14+I75, B14+I76, B14+I77, B14+I78, B14+I79, B14+I80, B14+I81, B14+I82,
B14+I83, B14+I84, B14+I85, B14+I86, B14+I87, B14+I88, B14+I89, B14+I90, B14+I91,
B14+I92, B14+I93, B14+I94, B14+I95, B14+I152, B14+I153, B14+I154, B14+I155,
5 B14+I187, B14+I188, B14+I189, B14+I190;

B15+I1, B15+I2, B15+I3, B15+I4, B15+I5, B15+I6, B15+I7, B15+I8, B15+I9, B15+I10,
B15+I11, B15+I12, B15+I13, B15+I14, B15+I15, B15+I16, B15+I17, B15+I18, B15+I19,
B15+I20, B15+I21, B15+I22, B15+I23, B15+I24, B15+I25, B15+I26, B15+I27, B15+I28,
B15+I29, B15+I30, B15+I31, B15+I32, B15+I33, B15+I34, B15+I35, B15+I36, B15+I37,
10 B15+I38, B15+I39, B15+I40, B15+I41, B15+I42, B15+I43, B15+I44, B15+I45, B15+I46,
B15+I47, B15+I48, B15+I49, B15+I50, B15+I51, B15+I52, B15+I53, B15+I54, B15+I55,
B15+I56, B15+I57, B15+I58, B15+I59, B15+I60, B15+I61, B15+I62, B15+I63, B15+I64,
B15+I65, B15+I66, B15+I67, B15+I68, B15+I69, B15+I70, B15+I71, B15+I72, B15+I73,
B15+I74, B15+I75, B15+I76, B15+I77, B15+I78, B15+I79, B15+I80, B15+I81, B15+I82,
15 B15+I83, B15+I84, B15+I85, B15+I86, B15+I87, B15+I88, B15+I89, B15+I90, B15+I91,
B15+I92, B15+I93, B15+I94, B15+I95, B15+I152, B15+I153, B15+I154, B15+I155,
B15+I187, B15+I188, B15+I189, B15+I190;

B16+I1, B16+I2, B16+I3, B16+I4, B16+I5, B16+I6, B16+I7, B16+I8, B16+I9, B16+I10,
B16+I11, B16+I12, B16+I13, B16+I14, B16+I15, B16+I16, B16+I17, B16+I18, B16+I19,
20 B16+I20, B16+I21, B16+I22, B16+I23, B16+I24, B16+I25, B16+I26, B16+I27, B16+I28,
B16+I29, B16+I30, B16+I31, B16+I32, B16+I33, B16+I34, B16+I35, B16+I36, B16+I37,
B16+I38, B16+I39, B16+I40, B16+I41, B16+I42, B16+I43, B16+I44, B16+I45, B16+I46,
B16+I47, B16+I48, B16+I49, B16+I50, B16+I51, B16+I52, B16+I53, B16+I54, B16+I55,
B16+I56, B16+I57, B16+I58, B16+I59, B16+I60, B16+I61, B16+I62, B16+I63, B16+I64,
25 B16+I65, B16+I66, B16+I67, B16+I68, B16+I69, B16+I70, B16+I71, B16+I72, B16+I73,
B16+I74, B16+I75, B16+I76, B16+I77, B16+I78, B16+I79, B16+I80, B16+I81, B16+I82,
B16+I83, B16+I84, B16+I85, B16+I86, B16+I87, B16+I88, B16+I89, B16+I90, B16+I91,
B16+I92, B16+I93, B16+I94, B16+I95, B16+I152, B16+I153, B16+I154, B16+I155,
B16+I187, B16+I188, B16+I189, B16+I190;

30 B17+I1, B17+I2, B17+I3, B17+I4, B17+I5, B17+I6, B17+I7, B17+I8, B17+I9, B17+I10,
B17+I11, B17+I12, B17+I13, B17+I14, B17+I15, B17+I16, B17+I17, B17+I18, B17+I19,
B17+I20, B17+I21, B17+I22, B17+I23, B17+I24, B17+I25, B17+I26, B17+I27, B17+I28,
B17+I29, B17+I30, B17+I31, B17+I32, B17+I33, B17+I34, B17+I35, B17+I36, B17+I37,

B17+I38, B17+I39, B17+I40, B17+I41, B17+I42, B17+I43, B17+I44, B17+I45, B17+I46,
B17+I47, B17+I48, B17+I49, B17+I50, B17+I51, B17+I52, B17+I53, B17+I54, B17+I55,
B17+I56, B17+I57, B17+I58, B17+I59, B17+I60, B17+I61, B17+ I62, B17+I63, B17+I64,
B17+I65, B17+I66, B17+I67, B17+I68, B17+I69, B17+I70, B17+I71, B17+I72, B17+I73,
5 B17+I74, B17+I75, B17+I76, B17+I77, B17+I78, B17+I79, B17+I80, B17+I81, B17+I82,
B17+I83, B17+I84, B17+I85, B17+I86, B17+I87, B17+I88, B17+I89, B17+I90, B17+I91,
B17+I92, B17+I93, B17+I94, B17+I95, B17+I152, B17+I153, B17+I154, B17+I155,
B17+I187, B17+I188, B17+I189, B17+I190;

B18+I1, B18+I2, B18+I3, B18+I4, B18+I5, B18+I6, B18+I7, B18+I8, B18+I9, B18+B180,
10 B18+I11, B18+I12, B18+I13, B18+I14, B18+I15, B18+I16, B18+I17, B18+I18, B18+I19,
B18+I20, B18+I21, B18+ I22, B18+I23, B18+I24, B18+I25, B18+I26, B18+I27, B18+I28,
B18+I29, B18+I30, B18+I31, B18+I32, B18+I33, B18+I34, B18+I35, B18+I36, B18+I37,
B18+I38, B18+I39, B18+I40, B18+I41, B18+I42, B18+I43, B18+I44, B18+I45, B18+I46,
B18+I47, B18+I48, B18+I49, B18+I50, B18+I51, B18+I52, B18+I53, B18+I54, B18+I55,
15 B18+I56, B18+I57, B18+I58, B18+I59, B18+I60, B18+I61, B18+ I62, B18+I63, B18+I64,
B18+I65, B18+I66, B18+I67, B18+I68, B18+I69, B18+I70, B18+I71, B18+I72, B18+I73,
B18+I74, B18+I75, B18+I76, B18+I77, B18+I78, B18+I79, B18+I80, B18+I81, B18+I82,
B18+I83, B18+I84, B18+I85, B18+I86, B18+I87, B18+I88, B18+I89, B18+I90, B18+I91,
B18+I92, B18+I93, B18+I94, B18+I95, B18+I152, B18+I153, B18+I154, B18+I155,
20 B18+I187, B18+I188, B18+I189, B18+I190;

B19+I1, B19+I2, B19+I3, B19+I4, B19+I5, B19+I6, B19+I7, B19+I8, B19+I9, B19+B190,
B19+I11, B19+I12, B19+I13, B19+I14, B19+I15, B19+I16, B19+I17, B19+I18, B19+I19,
B19+I20, B19+I21, B19+ I22, B19+I23, B19+I24, B19+I25, B19+I26, B19+I27, B19+I28,
B19+I29, B19+I30, B19+I31, B19+I32, B19+I33, B19+I34, B19+I35, B19+I36, B19+I37,
25 B19+I38, B19+I39, B19+I40, B19+I41, B19+I42, B19+I43, B19+I44, B19+I45, B19+I46,
B19+I47, B19+I48, B19+I49, B19+I50, B19+I51, B19+I52, B19+I53, B19+I54, B19+I55,
B19+I56, B19+I57, B19+I58, B19+I59, B19+I60, B19+I61, B19+ I62, B19+I63, B19+I64,
B19+I65, B19+I66, B19+I67, B19+I68, B19+I69, B19+I70, B19+I71, B19+I72, B19+I73,
B19+I74, B19+I75, B19+I76, B19+I77, B19+I78, B19+I79, B19+I80, B19+I81, B19+I82,
30 B19+I83, B19+I84, B19+I85, B19+I86, B19+I87, B19+I88, B19+I89, B19+I90, B19+I91,
B19+I92, B19+I93, B19+I94, B19+I95, B19+I152, B19+I153, B19+I154, B19+I155,
B19+I187, B19+I188, B19+I189, B19+I190;

B20+11, B20+12, B20+13, B20+14, B20+15, B20+16, B20+17, B20+18, B20+19, B20+B200,
B20+111, B20+112, B20+113, B20+114, B20+115, B20+116, B20+117, B20+118, B20+119,
B20+120, B20+121, B20+122, B20+123, B20+124, B20+125, B20+126, B20+127, B20+128,
B20+129, B20+130, B20+131, B20+132, B20+133, B20+134, B20+135, B20+136, B20+137,
5 B20+138, B20+139, B20+140, B20+141, B20+142, B20+143, B20+144, B20+145, B20+146,
B20+147, B20+148, B20+149, B20+150, B20+151, B20+152, B20+153, B20+154, B20+155,
B20+156, B20+157, B20+158, B20+159, B20+160, B20+161, B20+162, B20+163, B20+164,
B20+165, B20+166, B20+167, B20+168, B20+169, B20+170, B20+171, B20+172, B20+173,
B20+174, B20+175, B20+176, B20+177, B20+178, B20+179, B20+180, B20+181, B20+182,
10 B20+183, B20+184, B20+185, B20+186, B20+187, B20+188, B20+189, B20+190, B20+191,
B20+192, B20+193, B20+194, B20+195, B20+1152, B20+1153, B20+1154, B20+1155,
B20+1187, B20+1188, B20+1189, B20+1190.

Most preferably, the composition according to the present invention is selected from the group of combinations consisting of:

15 B1+18, B1+115, B1+121, B1+127, B1+135, B1+177, B1+194, B1+195, B1+1152, B1+1153,
B1+1155,
B2+18, B2+115, B2+121, B2+127, B2+135, B2+177, B2+194, B2+195, B2+1152, B2+1153,
B2+1155,
B3+18, B3+115, B3+121, B3+127, B3+135, B3+177, B3+194, B3+195, B3+1152, B3+1153,
20 B3+1155,
B4+18, B4+115, B4+121, B4+127, B4+135, B4+177, B4+194, B4+195, B4+1152, B4+1153,
B4+1155,
B5+18, B5+115, B5+121, B5+127, B5+135, B5+177, B5+194, B5+195, B5+1152, B5+1153,
B5+1155,
25 B6+18, B6+115, B6+121, B6+127, B6+135, B6+177, B6+194, B6+195, B6+1152, B6+1153,
B6+1155,
B7+18, B7+115, B7+121, B7+127, B7+135, B7+177, B7+194, B7+195, B7+1152, B7+1153,
B7+1155,
B8+18, B8+115, B8+121, B8+127, B8+135, B8+177, B8+194, B8+195, B8+1152, B8+1153,
30 B8+1155,

B9+18, B9+115, B9+121, B9+127, B9+135, B9+177, B9+194, B9+195, B9+1152, B9+1153, B9+1155,

B10+18, B10+115, B10+121, B10+127, B10+135, B10+177, B10+194, B10+195, B10+1152, B10+1153, B10+1155,

5 B11+18, B11+115, B11+121, B11+127, B11+135, B11+177, B11+194, B11+195, B11+1152, B11+1153, B11+1155,

B12+18, B12+115, B12+121, B12+127, B12+135, B12+177, B12+194, B12+195, B12+1152, B12+1153, B12+1155,

10 B13+18, B13+115, B13+121, B13+127, B13+135, B13+177, B13+194, B13+195, B13+1152, B13+1153, B13+1155,

B14+18, B14+115, B14+121, B14+127, B14+135, B14+177, B14+194, B14+195, B14+1152, B14+1153, B14+1155,

B15+18, B15+115, B15+121, B15+127, B15+135, B15+177, B15+194, B15+195, B15+1152, B15+1153, B15+1155,

15 B16+18, B16+115, B16+121, B16+127, B16+135, B16+177, B16+194, B16+195, B16+1152, B16+1153, B16+1155,

B17+18, B17+115, B17+121, B17+127, B17+135, B17+177, B17+194, B17+195, B17+1152, B17+1153, B17+1155,

20 B18+18, B18+115, B18+121, B18+127, B18+135, B18+177, B18+194, B18+195, B18+1152, B18+1153, B18+1155,

B19+18, B19+115, B19+121, B19+127, B19+135, B19+177, B19+194, B19+195, B19+1152, B19+1153, B19+1155,

B20+18, B20+115, B20+121, B20+127, B20+135, B20+177, B20+194, B20+195, B20+1152, B20+1153, B20+1155.

25 Still preferably, the composition according to the present invention is selected from the group of combinations consisting of:

B1+I15, B1+I21, B1+I95, B1+I152, B2+I15, B2+I21, B2+I95, B2+I152, B3+I15, B3+I21, B3+I95, B3+I152, B4+I15, B4+I21, B4+I95, B4+I152, B5+I15, B5+I21, B5+I95, B5+I152, B6+I15, B6+I21, B6+I95, B6+I152, B7+I15, B7+I21, B7+I95, B7+I152, B8+I15, B8+I21, B8+I95, B8+I152, B9+I15, B9+I21, B9+I95, B9+I152, B10+I15, 5 B10+I21, B10+I95, B10+I152, B11+I15, B11+I21, B11+I95, B11+I152, B12+I15, B12+I21, B12+I95, B12+I152, B13+I15, B13+I21, B13+I95, B13+I152, B14+I15, B14+I21, B14+I95, B14+I152, B15+I15, B15+I21, B15+I95, B15+I152, B16+I15, B16+I21, B16+I95, B16+I152, B17+I15, B17+I21, B17+I95, B17+I152, B18+I15, B18+I21, B18+I95, B18+I152, B19+I15, B19+I21, B19+I95, B19+I152, B20+I15, 10 B20+I21, B20+I95, and B20+I152.

Still preferably, the composition according to the present invention is selected from the group of combinations consisting of:

B16+I15, B9+I15, B3+I15, B9+I21, B3+I21, B3+I95, B9+I95, B16+I95, B19+I95, B9+I152, B3+I152, and B19+I152. In a preferred embodiment of the present invention 15 the composition further comprises at least one fungicide, with the proviso that the biological control agent and the fungicide are not identical.

Fungicides

In general, "fungicidal" means the ability of a substance to increase mortality or inhibit the growth rate of fungi.

20 The term "fungus" or "fungi" includes a wide variety of nucleated sporebearing organisms that are devoid of chlorophyll. Examples of fungi include yeasts, molds, mildews, rusts, and mushrooms.

Preferably, the fungicide is selected so as not to have any fungicidal activity against the biological control agent according to the present invention.

25 According to one embodiment of the present invention preferred fungicides are selected from the group consisting of

(1) Inhibitors of the ergosterol biosynthesis, for example (F1) aldimorph (1704-28-5), (F2) azaconazole (60207-31-0), (F3) bitertanol (55179-31-2), (F4) bromuconazole (116255-48-2), (F5) cyproconazole (113096-99-4), (F6) diclobutrazole (75736-33-3), (F7) 30 difenoconazole (119446-68-3), (F8) diniconazole (83657-24-3), (F9) diniconazole-M

- (83657-18-5), (F10) dodemorph (1593-77-7), (F11) dodemorph acetate (31717-87-0), (F12) epoxiconazole (106325-08-0), (F13) etaconazole (60207-93-4), (F14) fenarimol (60168-88-9), (F15) fenbuconazole (114369-43-6), (F16) fenhexamid (126833-17-8), (F17) fenpropidin (67306-00-7), (F18) fenpropimorph (67306-03-0), (F19) fluquinconazole
- 5 (136426-54-5), (F20) flurprimidol (56425-91-3), (F21) flusilazole (85509-19-9), (F22) flutriafol (76674-21-0), (F23) furconazole (112839-33-5), (F24) furconazole-cis (112839-32-4), (F25) hexaconazole (79983-71-4), (F26) imazalil (60534-80-7), (F27) imazalil sulfate (58594-72-2), (F28) imibenconazole (86598-92-7), (F29) ipconazole (125225-28-7), (F30) metconazole (125116-23-6), (F31) myclobutanil (88671-89-0), (F32) naftifine
- 10 (65472-88-0), (F33) nuarimol (63284-71-9), (F34) oxpoconazole (174212-12-5), (F35) paclobutrazol (76738-62-0), (F36) pefurazoate (101903-30-4), (F37) penconazole (66246-88-6), (F38) piperalin (3478-94-2), (F39) prochloraz (67747-09-5), (F40) propiconazole (60207-90-1), (F41) prothioconazole (178928-70-6), (F42) pyributicarb (88678-67-5), (F43) pyrifenoxy (88283-41-4), (F44) quinconazole (103970-75-8), (F45)
- 15 simeconazole (149508-90-7), (F46) spiroxamine (118134-30-8), (F47) tebuconazole (107534-96-3), (F48) terbinafine (91161-71-6), (F49) tetraconazole (112281-77-3), (F50) triadimefon (43121-43-3), (F51) triadimenol (89482-17-7), (F52) tridemorph (81412-43-3), (F53) triflumizole (68694-11-1), (F54) triforine (26644-46-2), (F55) triticonazole (131983-72-7), (F56) uniconazole (83657-22-1), (F57) uniconazole-p (83657-17-4), (F58) viniconazole
- 20 (77174-66-4), (F59) voriconazole (137234-62-9), (F60) 1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)cycloheptanol (129586-32-9), (F61) methyl 1-(2,2-dimethyl-2,3-dihydro-1H-inden-1-yl)-1H-imidazole-5-carboxylate (110323-95-0), (F62) N'-{5-(difluoromethyl)-2-methyl-4-[3-(trimethylsilyl)propoxy]phenyl}-N-ethyl-N-methylimidoformamide, (F63) N-ethyl-N-methyl-N'-{2-methyl-5-(trifluoromethyl)-4-[3-
- 25 (trimethylsilyl)propoxy]phenyl}imidoformamide, (F64) O-[1-(4-methoxyphenoxy)-3,3-dimethylbutan-2-yl] 1H-imidazole-1-carbothioate (111226-71-2);
- (2) inhibitors of the respiratory chain at complex I or II, for example (F65) bixafen (581809-46-3), (F66) boscalid (188425-85-6), (F67) carboxin (5234-68-4), (F68) diflumetorim (130339-07-0), (F69) fenfuram (24691-80-3), (F70) fluopyram (658066-35-
- 30 4), (F71) flutolanil (66332-96-5), (F72) fluxapyroxad (907204-31-3), (F73) furametpyr (123572-88-3), (F74) furmecycloz (60568-05-0), (F75) isopyrazam (mixture of syn-epimeric racemate 1RS,4SR,9RS and anti-epimeric racemate 1RS,4SR,9SR) (881685-58-1), (F76) isopyrazam (anti-epimeric racemate 1RS,4SR,9SR), (F77) isopyrazam (anti-epimeric enantiomer 1R,4S,9S), (F78) isopyrazam (anti-epimeric enantiomer 1S,4R,9R),
- 35 (F79) isopyrazam (syn epimeric racemate 1RS,4SR,9RS), (F80) isopyrazam (syn-epimeric

enantiomer 1R,4S,9R), (F81) isopyrazam (syn-epimeric enantiomer 1S,4R,9S), (F82) mepronil (55814-41-0), (F83) oxycarboxin (5259-88-1), (F84) penflufen (494793-67-8), (F85) penthiopyrad (183675-82-3), (F86) sedaxane (874967-67-6), (F87) thifluzamide (130000-40-7), (F88) 1-methyl-N-[2-(1,1,2,2-tetrafluoroethoxy)phenyl]-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide, (F89) 3-(difluoromethyl)-1-methyl-N-[2-(1,1,2,2-tetrafluoroethoxy)phenyl]-1H-pyrazole-4-carboxamide, (F90) 3-(difluoromethyl)-N-[4-fluoro-2-(1,1,2,3,3,3-hexafluoropropoxy)phenyl]-1-methyl-1H-pyrazole-4-carboxamide, (F91) N-[1-(2,4-dichlorophenyl)-1-methoxypropan-2-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide (1092400-95-7), (F92) 5,8-difluoro-N-[2-(2-fluoro-4-[[4-(trifluoromethyl)pyridin-2-yl]oxy}phenyl)ethyl]quinazolin-4-amine (1210070-84-0), (F93) benzovindiflupyr, (F94) N-[(1S,4R)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (F95) N-[(1R,4S)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (F96) 3-(Difluormethyl)-1-methyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazol-4-carboxamid, (F97) 1,3,5-Trimethyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazol-4-carboxamid, (F98) 1-Methyl-3-(trifluormethyl)-N-(1,3,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazol-4-carboxamid, (F99) 1-Methyl-3-(trifluormethyl)-N-[(1S)-1,3,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (F100) 1-Methyl-3-(trifluormethyl)-N-[(1R)-1,3,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (F101) 3-(Difluormethyl)-1-methyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (F102) 3-(Difluormethyl)-1-methyl-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (F103) 1,3,5-Trimethyl-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (F104) 1,3,5-Trimethyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid;

(3) inhibitors of the respiratory chain at complex III, for example (F105) ametoctradin (865318-97-4), (F106) amisulbrom (348635-87-0), (F107) azoxystrobin (131860-33-8), (F108) cyazofamid (120116-88-3), (F109) coumethoxystrobin (850881-30-0), (F110) coumoxystrobin (850881-70-8), (F111) dimoxystrobin (141600-52-4), (F112) enestroburin (238410-11-2), (F113) famoxadone (131807-57-3), (F114) fenamidone (161326-34-7), (F115) fenoxystrobin (918162-02-4), (F116) fluoxastrobin (361377-29-9), (F117) kresoxim-methyl (143390-89-0), (F118) metominostrobin (133408-50-1), (F119) oryastrobin (189892-69-1), (F120) picoxystrobin (117428-22-5), (F121) pyraclostrobin (175013-18-0), (F122) pyrametostrobin (915410-70-7), (F123) pyraoxystrobin (862588-11-2), (F124) pyribencarb (799247-52-2), (F125) triclopyricarb (902760-40-1), (F126) trifloxystrobin

- (141517-21-7), (F127) (2E)-2-(2-{{6-(3-chloro-2-methylphenoxy)-5-fluoropyrimidin-4-yl}oxy}phenyl)-2-(methoxyimino)-N-methylethanamide, (F128) (2E)-2-(methoxyimino)-N-methyl-2-(2-{{{{(1E)-1-[3-(trifluoromethyl)phenyl]ethylidene}amino)oxy}methyl}phenyl)ethanamide, (F129) (2E)-2-
- 5 (methoxyimino)-N-methyl-2-{2-[[E)-{{1-[3-(trifluoromethyl)phenyl]ethoxy}imino}methyl]phenyl}ethanamide (158169-73-4), (F130) (2E)-2-{2-{{{{(1E)-1-[3-{{(E)-1-fluoro-2-phenylethenyl}oxy}phenyl]ethylidene}amino}oxy}methyl}phenyl)-2-(methoxyimino)-N-methylethanamide (326896-28-0), (F131) (2E)-2-{2-[[{{(2E,3E)-4-(2,6-dichlorophenyl)but-3-en-2-ylidene}amino}oxy}methyl]phenyl)-2-(methoxyimino)-N-methylethanamide,
- 10 (F132) 2-chloro-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)pyridine-3-carboxamide (119899-14-8), (F133) 5-methoxy-2-methyl-4-(2-{{{{(1E)-1-[3-(trifluoromethyl)phenyl]ethylidene}amino)oxy}methyl}phenyl)-2,4-dihydro-3H-1,2,4-triazol-3-one, (F134) methyl (2E)-2-{2-[[{cyclopropyl}[[4-
- 15 methoxyphenyl]imino}methyl]sulfanyl)methyl]phenyl}-3-methoxyprop-2-enoate (149601-03-6), (F135) N-(3-ethyl-3,5,5-trimethylcyclohexyl)-3-(formylamino)-2-hydroxybenzamide (226551-21-9), (F136) 2-{2-[[2,5-dimethylphenoxy}methyl]phenyl}-2-methoxy-N-methylacetamide (173662-97-0), (F137) (2R)-2-{2-[[2,5-dimethylphenoxy}methyl]phenyl}-2-methoxy-N-methylacetamide (394657-24-0);
- 20 (4) Inhibitors of the mitosis and cell division, for example (F138) benomyl (17804-35-2), (F139) carbendazim (10605-21-7), (F140) chlorfenazole (3574-96-7), (F141) diethofencarb (87130-20-9), (F142) ethaboxam (162650-77-3), (F143) fluopicolide (239110-15-7), (F144) fuberidazole (3878-19-1), (F145) pencycuron (66063-05-6), (F146) thiabendazole (148-79-8), (F147) thiophanate-methyl (23564-05-8), (F148) thiophanate
- 25 (23564-06-9), (F149) zoxamide (156052-68-5), (F150) 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine (214706-53-3), (F151) 3-chloro-5-(6-chloropyridin-3-yl)-6-methyl-4-(2,4,6-trifluorophenyl)pyridazine (1002756-87-7);
- (5) Compounds capable to have a multisite action, like for example (F152) bordeaux mixture (8011-63-0), (F153) captafol (2425-06-1), (F154) captan (133-06-2), (F155) chlorothalonil (1897-45-6), (F156) copper hydroxide (20427-59-2), (F157) copper naphthenate (1338-02-9), (F158) copper oxide (1317-39-1), (F159) copper oxychloride (1332-40-7), (F160) copper(2+) sulfate (7758-98-7), (F161) dichlofluanid (1085-98-9), (F162) dithianon (3347-22-6), (F163) dodine (2439-10-3), (F164) dodine free base, (F165) ferbam (14484-64-1), (F166) fluorofolpet (719-96-0), (F167) folpet (133-07-3), (F168)

guazatine (108173-90-6), (F169) guazatine acetate, (F170) iminoctadine (13516-27-3),
(F171) iminoctadine albesilate (169202-06-6), (F172) iminoctadine triacetate (57520-17-
9), (F173) mancozeb (53988-93-5), (F174) mancozeb (8018-01-7), (F175) maneb
(12427-38-2), (F176) metiram (9006-42-2), (F177) metiram zinc (9006-42-2), (F178)
5 oxine-copper (10380-28-6), (F179) propamidine (104-32-5), (F180) propineb (12071-83-
9), (F181) sulphur and sulphur preparations including calcium polysulphide (7704-34-9),
(F182) thiram (137-26-8), (F183) tolyfluanid (731-27-1), (F184) zineb (12122-67-7), (F185)
ziram (137-30-4);

(6) Compounds capable to induce a host defense, like for example (F186) acibenzolar-
10 S-methyl (135158-54-2), (F187) isotianil (224049-04-1), (F188) probenazole (27605-76-1),
(F189) tiadinil (223580-51-6);

(7) Inhibitors of the amino acid and/or protein biosynthesis, for example (F190) andoprim
(23951-85-1), (F191) blastidn-S (2079-00-7), (F192) cyprodinil (121552-61-2), (F193)
kasugamycin (6980-18-3), (F194) kasugamycin hydrochloride hydrate (19408-46-9),
15 (F195) mepanipyrim (110235-47-7), (F196) pyrimethanil (53112-28-0), (F197) 3-(5-fluoro-
3,3,4,4-tetramethyl-3,4-dihydroisoquinolin-1-yl)quinoline (861647-32-7);

(8) Inhibitors of the ATP production, for example (F198) fentin acetate (900-95-8), (F199)
fentin chloride (639-58-7), (F200) fentin hydroxide (76-87-9), (F201) silthiofam (175217-
20-6);

20 (9) Inhibitors of the cell wall synthesis, for example (F202) benthialdicarb (177406-68-7),
(F203) dimethomorph (110488-70-5), (F204) flumorph (211867-47-9), (F205) iprovaldicarb
(140923-17-7), (F206) mandipropamid (374726-62-2), (F207) polyoxins (11113-80-7),
(F208) polyoxorim (22976-86-9), (F209) validamycin A (37248-47-8), (F210) vallfenalate
(283159-94-4; 283159-90-0);

25 (10) Inhibitors of the lipid and membrane synthesis, for example (F211) biphenyl (92-52-
4), (F212) chloroneb (2675-77-6), (F213) dicloran (99-30-9), (F214) edifenphos (17109-
49-8), (F215) etridiazole (2593-15-9), (F216) iodocarb (55406-53-6), (F217) iprobenfos
(26087-47-8), (F218) isoprothiolane (50512-35-1), (F219) propamocarb (25606-41-1),
(F220) propamocarb hydrochloride (25606-41-1), (F221) prothiocarb (19622-08-3),
30 (F222) pyrazophos (13457-18-6), (F223) quintozene (82-68-8), (F224) tecnazene (117-18-
0), (F225) tolclofos-methyl (57018-04-9);

(11) Inhibitors of the melanine biosynthesis, for example (F226) carpropamid (104030-54-8), (F227) diclocymet (139920-32-4), (F228) fenoxanil (115852-48-7), (F229) phthalide (27355-22-2), (F230) pyroquilon (57369-32-1), (F231) tricyclazole (41814-78-2), (F232) 2,2,2-trifluoroethyl {3-methyl-1-[(4-methylbenzoyl)amino]butan-2-yl}carbamate (851524-22-6);

(12) Inhibitors of the nucleic acid synthesis, for example (F233) benalaxyl (71626-11-4), (F234) benalaxyl-M (kiralaxyl) (98243-83-5), (F235) bupirimate (41483-43-6), (F236) clozylacon (67932-85-8), (F237) dimethirimol (5221-53-4), (F238) ethirimol (23947-60-6), (F239) furalaxyl (57646-30-7), (F240) hymexazol (10004-44-1), (F241) metalaxyl (57837-19-1), (F242) metalaxyl-M (mefenoxam) (70630-17-0), (F243) ofurace (58810-48-3), (F244) oxadixyl (77732-09-3), (F245) oxolinic acid (14698-29-4);

(13) Inhibitors of the signal transduction, for example (F246) chlozollinate (84332-86-5), (F247) fenpiclonil (74738-17-3), (F248) fludioxonil (131341-86-1), (F249) Iprodione (36734-19-7), (F250) procymidone (32809-16-8), (F251) quinoxifen (124495-18-7), (F252) vinclozolin (50471-44-8);

(14) Compounds capable to act as an uncoupler, like for example (F253) binapacryl (485-31-4), (F254) dinocap (131-72-6), (F255) ferimzone (89269-64-7), (F256) fluazinam (79622-59-6), (F257) meptyldinocap (131-72-6);

(15) Further compounds, like for example (F258) benthiazole (21564-17-0), (F259) bethoxazin (163269-30-5), (F260) capsimycin (70694-08-5), (F261) carvone (99-49-0), (F262) chinomethionat (2439-01-2), (F263) pyriofenone (chlazafenone) (688046-61-9), (F264) cufraneb (11096-18-7), (F265) cyflufenamid (180409-60-3), (F266) cymoxanil (57966-95-7), (F267) cyprosulfamide (221667-31-8), (F268) dazomet (533-74-4), (F269) debacarb (62732-91-6), (F270) dichlorophen (97-23-4), (F271) diclomezine (62865-36-5), (F272) difenzoquat (49866-87-7), (F273) difenzoquat methylsulphate (43222-48-6), (F274) diphenylamine (122-39-4), (F275) ecomate, (F276) fenpyrazamine (473798-59-3), (F277) flumetover (154025-04-4), (F278) fluoroimide (41205-21-4), (F279) flusulfamide (106917-52-6), (F280) fluticanil (304900-25-2), (F281) fosetyl-aluminium (39148-24-8), (F282) fosetyl-calcium, (F283) fosetyl-sodium (39148-16-8), (F284) hexachlorobenzene (118-74-1), (F285) irumamycin (81604-73-1), (F286) methasulfocarb (66952-49-6), (F287) methyl isothiocyanate (556-61-6), (F288) metrafenone (220899-03-6), (F289) mildiomycin (67527-71-3), (F290) natamycin (7681-93-8), (F291) nickel dimethyldithiocarbamate (15521-65-0), (F292) nitrothal-isopropyl (10552-74-6), (F293)

octhillinone (26530-20-1), (F294) oxamocarb (917242-12-7), (F295) oxyfenthilin (34407-87-9), (F296) pentachlorophenol and salts (87-86-5), (F297) phenothrin, (F298) phosphorous acid and its salts (13598-36-2), (F299) propamocarb-fosetylalate, (F300) propanosine-sodium (88498-02-6), (F301) proquinazid (189278-12-4), (F302) pyrimorph
5 (868390-90-3), (F303) (2E)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one (1231776-28-5), (F304) (2Z)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one (1231776-29-6), (F305) pyrrolnitrine (1018-71-9), (F306) tebufloquin (376645-78-2), (F307) tecloftalam (76280-91-6), (F308) tolifenide (304911-98-6), (F309) triazoxide (72459-58-6), (F310) trichlamide (70193-21-4), (F311)
10 zarilamid (84527-51-5), (F312) (3S,6S,7R,8R)-8-benzyl-3-[(3-[(isobutyryloxy)methoxy]-4-methoxypyridin-2-yl)carbonylamino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl 2-methylpropanoate (517875-34-2), (F313) 1-(4-{4-[(5R)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone (1003319-79-6), (F314) 1-(4-{4-[(5S)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone (1003319-80-9), (F315) 1-(4-{4-[5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone (1003318-67-9), (F316) 1-(4-methoxyphenoxy)-3,3-dimethylbutan-2-yl 1H-imidazole-1-carboxylate (111227-17-9), (F317) 2,3,5,6-tetrachloro-4-
20 (methylsulfonyl)pyridine (13108-52-6), (F318) 2,3-dibutyl-6-chlorothieno[2,3-d]pyrimidin-4(3H)-one (221451-58-7), (F319) 2,6-dimethyl-1H,5H-[1,4]dithiino[2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetrone, (F320) 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-(4-{4-[(5R)-5-phenyl-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone (1003316-53-7), (F321) 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-(4-{4-[(5S)-5-phenyl-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone (1003316-54-8), (F322) 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-{4-[4-(5-phenyl-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidin-1-yl}ethanone (1003316-51-5), (F323) 2-butoxy-6-iodo-3-propyl-4H-chromen-4-one, (F324) 2-chloro-5-[2-chloro-1-(2,6-difluoro-4-methoxyphenyl)-4-methyl-1H-imidazol-5-yl]pyridine, (F325) 2-phenylphenol and salts (90-43-7), (F326) 3-
30 (4,4,5-trifluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline (861647-85-0), (F327) 3,4,5-trichloropyridine-2,6-dicarbonitrile (17824-85-0), (F328) 3-[5-(4-chlorophenyl)-2,3-dimethyl-1,2-oxazolidin-3-yl]pyridine, (F329) 3-chloro-5-(4-chlorophenyl)-4-(2,6-difluorophenyl)-6-methylpyridazine, (F330) 4-(4-chlorophenyl)-5-(2,6-difluorophenyl)-3,6-dimethylpyridazine, (F331) 5-amino-1,3,4-thiadiazole-2-thiol, (F332) 5-chloro-N'-phenyl-N'-(prop-2-yn-1-yl)thiophene-2-sulfonohydrazide (134-31-6), (F333) 5-fluoro-2-[(4-

- fluorobenzyl)oxy]pyrimidin-4-amine (1174376-11-4), (F334) 5-fluoro-2-[(4-methylbenzyl)oxy]pyrimidin-4-amine (1174376-25-0), (F335) 5-methyl-6-octyl[1,2,4]triazolo[1,5-a]pyrimidin-7-amine, (F336) ethyl (2Z)-3-amino-2-cyano-3-phenylprop-2-enoate, (F337) N'-(4-[[3-(4-chlorobenzyl)-1,2,4-thiadiazol-5-yl]oxy]-2,5-dimethylphenyl)-N-ethyl-N-methylimidoforamide, (F338) N-(4-chlorobenzyl)-3-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]propanamide, (F339) N-[(4-chlorophenyl)(cyano)methyl]-3-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]propanamide, (F340) N-[(5-bromo-3-chloropyridin-2-yl)methyl]-2,4-dichloropyridine-3-carboxamide, (F341) N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2,4-dichloropyridine-3-carboxamide, (F342) N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2-fluoro-4-iodopyridine-3-carboxamide, (F343) N-{(E)-[(cyclopropylmethoxy)imino][6-(difluoromethoxy)-2,3-difluorophenyl]methyl}-2-phenylacetamide (221201-92-9), (F344) N-{(Z)-[(cyclopropylmethoxy)imino][6-(difluoromethoxy)-2,3-difluorophenyl]methyl}-2-phenylacetamide (221201-92-9), (F345) N'-(4-[[3-tert-butyl-4-cyano-1,2-thiazol-5-yl]oxy]-2-chloro-5-methylphenyl)-N-ethyl-N-methylimidoforamide, (F346) N-methyl-2-(1-[[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl]piperidin-4-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)-1,3-thiazole-4-carboxamide (922514-49-6), (F347) N-methyl-2-(1-[[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl]piperidin-4-yl)-N-[(1R)-1,2,3,4-tetrahydronaphthalen-1-yl]-1,3-thiazole-4-carboxamide (922514-07-6), (F348) N-methyl-2-(1-[[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl]piperidin-4-yl)-N-[(1S)-1,2,3,4-tetrahydronaphthalen-1-yl]-1,3-thiazole-4-carboxamide (922514-48-5), (F349) pentyl {6-[[{[(1-methyl-1H-tetrazol-5-yl)(phenyl)methylidene]amino}oxy)methyl]pyridin-2-yl}carbamate, (F350) phenazine-1-carboxylic acid, (F351) quinolin-8-ol (134-31-6), (F352) quinolin-8-ol sulfate (2:1) (134-31-6), (F353) tert-butyl {6-[[{[(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino}oxy)methyl]pyridin-2-yl}carbamate;
- (16) Further compounds, like for example (F354) 1-methyl-3-(trifluoromethyl)-N-[2'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (F355) N-(4'-chlorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (F356) N-(2',4'-dichlorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (F357) 3-(difluoromethyl)-1-methyl-N-[4'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (F358) N-(2',5'-difluorobiphenyl-2-yl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide, (F359) 3-(difluoromethyl)-1-methyl-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (F360) 5-fluoro-1,3-dimethyl-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (F361) 2-chloro-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide, (F362) 3-(difluoromethyl)-N-[4'-(3,3-dimethylbut-

1-yn-1-yl)biphenyl-2-yl]-1-methyl-1H-pyrazole-4-carboxamide, (F363) N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide, (F364) 3-(difluoromethyl)-N-(4'-ethynylbiphenyl-2-yl)-1-methyl-1H-pyrazole-4-carboxamide, (F365) N-(4'-ethynylbiphenyl-2-yl)-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide, (F366) 2-chloro-N-(4'-ethynylbiphenyl-2-yl)pyridine-3-carboxamide, (F367) 2-chloro-N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide, (F368) 4-(difluoromethyl)-2-methyl-N-[4'-(trifluoromethyl)biphenyl-2-yl]-1,3-thiazole-5-carboxamide, (F369) 5-fluoro-N-[4'-(3-hydroxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1,3-dimethyl-1H-pyrazole-4-carboxamide, (F370) 2-chloro-N-[4'-(3-hydroxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide, (F371) 3-(difluoromethyl)-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1-methyl-1H-pyrazole-4-carboxamide, (F372) 5-fluoro-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1,3-dimethyl-1H-pyrazole-4-carboxamide, (F373) 2-chloro-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide, (F374) (5-bromo-2-methoxy-4-methylpyridin-3-yl)(2,3,4-trimethoxy-6-methylphenyl)methanone, (F375) N-[2-(4-{{[3-(4-chlorophenyl)prop-2-yn-1-yl]oxy}-3-methoxyphenyl]ethyl}-N2-(methylsulfonyl)valinamide (220706-93-4), (F376) 4-oxo-4-[(2-phenylethyl)amino]butanoic acid, (F377) but-3-yn-1-yl {6-[[{[(Z)-(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino}oxy)methyl]pyridin-2-yl} carbamate, (F378) 4-Amino-5-fluorpyrimidin-2-ol (mesomere Form: 6-Amino-5-fluorpyrimidin-2(1H)-on), (F379) propyl 3,4,5-trihydroxybenzoate and (F380) Oryzastrobin.

All named fungicides of the classes (1) to (16) (i. e. F1 to F380) can, if their functional groups enable this, optionally form salts with suitable bases or acids.

In a preferred embodiment of the present invention the at least fungicide is a synthetic fungicide.

25 In one embodiment of the present invention the composition comprises two or more fungicides. In a preferred embodiment the composition comprises two or more of the above-mentioned preferred fungicides.

According to a preferred embodiment of the present invention the fungicide is selected from the group consisting of (1) Inhibitors of the ergosterol biosynthesis, for example (F3) bitertanol, (F4) bromuconazole (116255-48-2), (F5) cyproconazole (113096-99-4), (F7) difenoconazole (119446-68-3), (F12) epoxiconazole (106325-08-0), (F16) fenhexamid (126833-17-8), (F17) fenpropidin (67306-00-7), (F18) fenpropimorph (67306-03-0), (F19) fluquinconazole (136426-54-5), (F22) flutriafol, (F26) imazalil, (F29) ipconazole (125225-

28-7), (F30) metconazole (125116-23-6), (F31) myclobutanil (88671-89-0), (F37) penconazole (66246-88-6), (F39) prochloraz (67747-09-5), (F40) propiconazole (60207-90-1), (F41) prothioconazole (178928-70-6), (F44) quinconazole (103970-75-8), (F46) spiroxamine (118134-30-8), (F47) tebuconazole (107534-96-3), (F51) triadimenol (89482-17-7), (F55) triticonazole (131983-72-7);

(2) inhibitors of the respiratory chain at complex I or II, for example (F65) bixafen (581809-46-3), (F66) boscalid (188425-85-6), (F67) carboxin (5234-68-4), (F70) fluopyram (658066-35-4), (F71) flutolanil (66332-96-5), (F72) fluxapyroxad (907204-31-3), (F73) furametpyr (123572-88-3), (F75) isopyrazam (mixture of syn-epimeric racemate 1RS,4SR,9RS and anti-epimeric racemate 1RS,4SR,9SR) (881685-58-1), (F76) isopyrazam (anti-epimeric racemate 1RS,4SR,9SR), (F77) isopyrazam (anti-epimeric enantiomer 1R,4S,9S), (F78) isopyrazam (anti-epimeric enantiomer 1S,4R,9R), (F79) isopyrazam (syn epimeric racemate 1RS,4SR,9RS), (F80) isopyrazam (syn-epimeric enantiomer 1R,4S,9R), (F81) isopyrazam (syn-epimeric enantiomer 1S,4R,9S), (F84) penflufen (494793-67-8), (F85) penthiopyrad (183675-82-3), (F86) sedaxane (874967-67-6), (F87) thifluzamide (130000-40-7), (F91) N-[1-(2,4-dichlorophenyl)-1-methoxypropan-2-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide (1092400-95-7), (F98) 1-Methyl-3-(trifluormethyl)-N-(1,3,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazol-4-carboxamid, (F99) 1-Methyl-3-(trifluormethyl)-N-[(1S)-1,3,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (F100) 1-Methyl-3-(trifluormethyl)-N-[(1R)-1,3,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (F101) 3-(Difluormethyl)-1-methyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (F102) 3-(Difluormethyl)-1-methyl-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid;

(3) inhibitors of the respiratory chain at complex III, for example (F105) ametocradin (865318-97-4), (F106) amisulbrom (348635-87-0), (F107) azoxystrobin (131860-33-8), (F108) cyazofamid (120116-88-3), (F111) dimoxystrobin (141600-52-4), (F112) enestroburin (238410-11-2), (F113) famoxadone (131807-57-3), (F114) fenamidone (161326-34-7), (F116) fluoxastrobin (361377-29-9), (F117) kresoxim-methyl (143390-89-0), (F118) metominostrobin (133408-50-1), (F119) oryastrobin (189892-69-1), (F120) picoxystrobin (117428-22-5), (F121) pyraclostrobin (175013-18-0), (F124) pyribencarb (799247-52-2), (F126) trifloxystrobin (141517-21-7);

(4) inhibitors of the mitosis and cell division, for example (F139) carbendazim (10605-21-7), (F140) chlorfenazole (3574-96-7), (F141) diethofencarb (87130-20-9), (F142)

ethaboxam (162650-77-3), (F143) fluopicolide, (F144) fuberidazole (3878-19-1), (F145) pencycuron (66063-05-6), (F147) thiophanate-methyl (23564-05-8), (F149) zoxamide (156052-68-5);

5 (5) Compounds capable to have a multisite action, like for example (F154) captan (133-06-2), (F155) chlorothalonil (1897-45-6), (F156) copper hydroxide (20427-59-2), (F159) copper oxychloride (1332-40-7), (F162) dithianon (3347-22-6), (F163) dodine (2439-10-3), (F167) folpet (133-07-3), (F168) guazatine (108173-90-6), (F172) iminoctadine triacetate (57520-17-9), (F174) mancozeb (8018-01-7), (F180) propineb (12071-83-9), (F181) sulphur and sulphur preparations including calcium polysulphide
10 (7704-34-9), (F182) thiram (137-26-8);

(6) Compounds capable to induce a host defense, like for example (F186) acibenzolar-S-methyl (135158-54-2), (F187) isotianil (224049-04-1), (F189) tiadinil (223580-51-6);

(7) Inhibitors of the amino acid and/or protein biosynthesis, for example (F192) cyprodinil (121552-61-2), (F196) pyrimethanil (53112-28-0);

15 (9) Inhibitors of the cell wall synthesis, for example (F202) benthiallalicarb (177406-68-7), (F203) dimethomorph (110488-70-5), (F205) iprovalicarb (140923-17-7), (F206) mandipropamid (374726-62-2), (F210) valifenalate (283159-94-4; 283159-90-0);

(10) Inhibitors of the lipid and membrane synthesis, for example (F216) iodocarb (55406-53-6), (F217) iprobenfos (26087-47-8), (F220) propamocarb hydrochloride (25606-41-1),
20 (F225) tolclofos-methyl;

(11) Inhibitors of the melanine biosynthesis, for example (F226) carpropamid

(12) Inhibitors of the nucleic acid synthesis, for example (F233) benalaxyl (71626-11-4), (F234) benalaxyl-M (kiralaxyl) (98243-83-5), (F239) furalaxyl (57646-30-7), (F240) hymexazol (10004-44-1), (F241) metalaxyl (57837-19-1), (F242) metalaxyl-M
25 (mefenoxam) (70630-17-0), (F244) oxadixyl (77732-09-3);

(13) Inhibitors of the signal transduction, for example (F247) fenpiclonil (74738-17-3), (F248) fludioxonil (131341-86-1), (F249) iprodione (36734-19-7), (F251) quinoxifen (124495-18-7), (F252) vinclozolin (50471-44-8);

(14) Compounds capable to act as an uncoupler, like for example (F256) fluazinam
30 (79622-59-6);

(15) Further compounds, like for example (F266) cymoxanil (57966-95-7), (F280) flutianil (304900-25-2), (F281) fosetyl-aluminium (39148-24-8), (F286) methasulfocarb (66952-49-6), (F287) methyl isothiocyanate (556-61-6), (F288) metrafenone (220899-03-6), (F298) phosphorous acid and its salts (13598-36-2), (F301) proquinazid (189278-12-4),
5 (F309) triazoxide (72459-58-6) and (F319) 2,6-dimethyl-1H,5H-[1,4]dithiino[2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetrone.

In one embodiment of the present invention, the fungicide, e.g., the fungicide for use in seed treatment is selected from the group consisting of Carbendazim (F139), Carboxin (F67), Difenoconazole (F7), Fludioxonil (F248), Fluquinconazole (F19), Fluxapyroxad (F72),
10 Ipconazole (F29), Isotianil (F187), Mefenoxam (F242), Metalaxyl (F241), Pencycuron (F145), Penflufen (F84), Prothioconazole (F41), Prochloraz (F39), Pyraclostrobin (F121), Sedaxane (F86), Silthiofam (F201), Tebuconazole (F47), Thiram (F182), Trifloxystrobin (F126), and Trifliconazole (F55).

Further additives

One aspect of the present invention is to provide a composition as described above additionally comprising at least one auxiliary selected from the group consisting of extenders, solvents, spontaneity promoters, carriers, emulsifiers, dispersants, frost
5 protectants, thickeners and adjuvants. Those compositions are referred to as formulations.

Accordingly, in one aspect of the present invention such formulations, and application forms prepared from them, are provided as crop protection agents and/or pesticidal agents, such as drench, drip and spray liquors, comprising the composition of the
10 invention. The application forms may comprise further crop protection agents and/or pesticidal agents, and/or activity-enhancing adjuvants such as penetrants, examples being vegetable oils such as, for example, rapeseed oil, sunflower oil, mineral oils such as, for example, liquid paraffins, alkyl esters of vegetable fatty acids, such as rapeseed oil or soybean oil methyl esters, or alkanol alkoxylates, and/or spreaders such as, for
15 example, alkylsiloxanes and/or salts, examples being organic or inorganic ammonium or phosphonium salts, examples being ammonium sulphate or diammonium hydrogen phosphate, and/or retention promoters such as dioctyl sulphosuccinate or hydroxypropylguar polymers and/or humectants such as glycerol and/or fertilizers such as ammonium, potassium or phosphorous fertilizers, for example.

20 Examples of typical formulations include water-soluble liquids (SL), emulsifiable concentrates (EC), emulsions in water (EW), suspension concentrates (SC, SE, FS, OD), water-dispersible granules (WG), granules (GR) and capsule concentrates (CS); these and other possible types of formulation are described, for example, by Crop Life International and in Pesticide Specifications, Manual on development and use of FAO
25 and WHO specifications for pesticides, FAO Plant Production and Protection Papers – 173, prepared by the FAO/WHO Joint Meeting on Pesticide Specifications, 2004, ISBN: 9251048576. The formulations may comprise active agrochemical compounds other than one or more active compounds of the invention.

The formulations or application forms in question preferably comprise auxiliaries, such as
30 extenders, solvents, spontaneity promoters, carriers, emulsifiers, dispersants, frost protectants, biocides, thickeners and/or other auxiliaries, such as adjuvants, for example. An adjuvant in this context is a component which enhances the biological effect of the formulation, without the component itself having a biological effect.

Examples of adjuvants are agents which promote the retention, spreading, attachment to the leaf surface, or penetration.

5 These formulations are produced in a known manner, for example by mixing the active compounds with auxiliaries such as, for example, extenders, solvents and/or solid carriers and/or further auxiliaries, such as, for example, surfactants. The formulations are prepared either in suitable plants or else before or during the application.

10 Suitable for use as auxiliaries are substances which are suitable for imparting to the formulation of the active compound or the application forms prepared from these formulations (such as, e.g., usable crop protection agents, such as spray liquors or seed dressings) particular properties such as certain physical, technical and/or biological properties.

15 Suitable extenders are, for example, water, polar and nonpolar organic chemical liquids, for example from the classes of the aromatic and non-aromatic hydrocarbons (such as paraffins, alkylbenzenes, alkylnaphthalenes, chlorobenzenes), the alcohols and polyols (which, if appropriate, may also be substituted, etherified and/or esterified), the ketones (such as acetone, cyclohexanone), esters (including fats and oils) and (poly)ethers, the unsubstituted and substituted amines, amides, lactams (such as N-alkylpyrrolidones) and lactones, the sulphones and sulphoxides (such as dimethyl sulphoxide).

20 If the extender used is water, it is also possible to employ, for example, organic solvents as auxiliary solvents. Essentially, suitable liquid solvents are: aromatics such as xylene, toluene or alkylnaphthalenes, chlorinated aromatics and chlorinated aliphatic hydrocarbons such as chlorobenzenes, chloroethylenes or methylene chloride, aliphatic hydrocarbons such as cyclohexane or paraffins, for example petroleum
25 fractions, mineral and vegetable oils, alcohols such as butanol or glycol and also their ethers and esters, ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents such as dimethylformamide and dimethyl sulphoxide, and also water. Preferred auxiliary solvents are selected from the group consisting of acetone and N,N'-dimethylformamide.

30 In principle it is possible to use all suitable solvents. Suitable solvents are, for example, aromatic hydrocarbons, such as xylene, toluene or alkylnaphthalenes, for example, chlorinated aromatic or aliphatic hydrocarbons, such as chlorobenzene,

chloroethylene or methylene chloride, for example, aliphatic hydrocarbons, such as cyclohexane, for example, paraffins, petroleum fractions, mineral and vegetable oils, alcohols, such as methanol, ethanol, isopropanol, butanol or glycol, for example, and also their ethers and esters, ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, for example, strongly polar solvents, such as dimethyl sulphoxide, and water.

All suitable carriers may in principle be used. Suitable carriers are in particular: for example, ammonium salts and ground natural minerals such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite or diatomaceous earth, and ground synthetic minerals, such as finely divided silica, alumina and natural or synthetic silicates, resins, waxes and/or solid fertilizers. Mixtures of such carriers may likewise be used. Carriers suitable for granules include the following: for example, crushed and fractionated natural minerals such as calcite, marble, pumice, sepiolite, dolomite, and also synthetic granules of inorganic and organic meals, and also granules of organic material such as sawdust, paper, coconut shells, maize cobs and tobacco stalks.

Liquefied gaseous extenders or solvents may also be used. Particularly suitable are those extenders or carriers which at standard temperature and under standard pressure are gaseous, examples being aerosol propellants, such as halogenated hydrocarbons, and also butane, propane, nitrogen and carbon dioxide.

Examples of emulsifiers and/or foam-formers, dispersants or wetting agents having ionic or nonionic properties, or mixtures of these surface-active substances, are salts of polyacrylic acid, salts of lignosulphonic acid, salts of phenolsulphonic acid or naphthalenesulphonic acid, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, with substituted phenols (preferably alkylphenols or arylphenols), salts of sulphosuccinic esters, taurine derivatives (preferably alkyltaurates), phosphoric esters of polyethoxylated alcohols or phenols, fatty acid esters of polyols, and derivatives of the compounds containing sulphates, sulphonates and phosphates, examples being alkylaryl polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates, protein hydrolysates, lignin-sulphite waste liquors and methylcellulose.

The presence of a surface-active substance is advantageous if one of the active compounds and/or one of the inert carriers is not soluble in water and if application takes place in water. Preferred emulsifiers are alkylaryl polyglycol ethers.

Further auxiliaries that may be present in the formulations and in the application forms derived from them include colorants such as inorganic pigments, examples being iron oxide, titanium oxide, Prussian Blue, and organic dyes, such as alizarin dyes, azo dyes and metal phthalocyanine dyes, and nutrients and trace nutrients, such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

Stabilizers, such as low-temperature stabilizers, preservatives, antioxidants, light stabilizers or other agents which improve chemical and/or physical stability may also be present. Additionally present may be foam-formers or defoamers.

Furthermore, the formulations and application forms derived from them may also comprise, as additional auxiliaries, stickers such as carboxymethylcellulose, natural and synthetic polymers in powder, granule or latex form, such as gum arabic, polyvinyl alcohol, polyvinyl acetate, and also natural phospholipids, such as cephalins and lecithins, and synthetic phospholipids. Further possible auxiliaries include mineral and vegetable oils.

There may possibly be further auxiliaries present in the formulations and the application forms derived from them. Examples of such additives include fragrances, protective colloids, binders, adhesives, thickeners, thixotropic substances, penetrants, retention promoters, stabilizers, sequestrants, complexing agents, humectants and spreaders. Generally speaking, the active compounds may be combined with any solid or liquid additive commonly used for formulation purposes.

Suitable retention promoters include all those substances which reduce the dynamic surface tension, such as dioctyl sulphosuccinate, or increase the viscoelasticity, such as hydroxypropylguar polymers, for example.

Suitable penetrants in the present context include all those substances which are typically used in order to enhance the penetration of active agrochemical compounds into plants. Penetrants in this context are defined in that, from the (generally aqueous) application liquor and/or from the spray coating, they are able to penetrate the cuticle of the plant and thereby increase the mobility of the active compounds in the cuticle. This property can be determined using the method described in the literature (Baur et al., 1997, Pesticide Science 51, 131-152). Examples include alcohol alkoxylates such as coconut fatty ethoxylate (10) or isotridecyl ethoxylate (12), fatty acid esters such as rapeseed or soybean oil methyl esters, fatty amine alkoxylates such as tallowamine

ethoxylate (15), or ammonium and/or phosphonium salts such as ammonium sulphate or diammonium hydrogen phosphate, for example.

The formulations preferably comprise between 0.00000001% and 98% by weight of active compound or, with particular preference, between 0.01% and 95% by weight of active compound, more preferably between 0.5% and 90% by weight of active compound, based on the weight of the formulation. The content of the active compound is defined as the sum of the at least one specified biological control agent and the at least one specified insecticide.

The active compound content of the application forms (crop protection products) prepared from the formulations may vary within wide ranges. The active compound concentration of the application forms may be situated typically between 0.00000001% and 95% by weight of active compound, preferably between 0.00001% and 1% by weight, based on the weight of the application form. Application takes place in a customary manner adapted to the application forms.

15 **Kit of parts**

Furthermore, in one aspect of the present invention a kit of parts is provided comprising at least one biological control agent selected from the group consisting of *Bacillus chittinoporos* AQ746 (NRRL Accession No. B-21618), *Bacillus mycoides* AQ726 (NRRL Accession No. B-21664), *Bacillus pumilus* (NRRL Accession No. B-30087), *Bacillus pumilus* AQ717 (NRRL Accession No. B-21662), *Bacillus sp.* AQ175 (ATCC Accession No. 55608), *Bacillus sp.* AQ177 (ATCC Accession No. 55609), *Bacillus sp.* AQ178 (ATCC Accession No. 53522), *Bacillus subtilis* AQ743 (NRRL Accession No. B-21665), *Bacillus subtilis* AQ713 (NRRL Accession No. B-21661), *Bacillus subtilis* AQ153 (ATCC Accession No. 55614), *Bacillus thuringiensis* BD#32 (NRRL Accession No. B-21530), *Bacillus thuringiensis* AQ52 (NRRL Accession No. B-21619), *Muscodor albus* 620 (NRRL Accession No. 30547), *Muscodor roseus* A3-5 (NRRL Accession No. 30548), *Rhodococcus globerulus* AQ719 (NRRL Accession No. B-21663), *Streptomyces galbus* (NRRL Accession No. 30232), *Streptomyces sp.* (NRRL Accession No. B-30145), *Bacillus thuringiensis* subspec. *kurstaki* BMP 123, *Bacillus subtilis* AQ30002 (NRRL Accession No. B-50421), and *Bacillus subtilis* AQ 30004 (NRRL Accession No. B-50455) and/or a mutant of these strains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens and at least one insecticide selected from the group consisting of

acetylcholinesterase (AChE) inhibitors, GABA-gated chloride channel antagonists, chloride channel activators and nicotinic acetylcholine receptor (nAChR) channel blockers, in a synergistically effective amount, in a spatially separated arrangement.

5 In a further embodiment of the present invention the above-mentioned kit of parts further comprises at least one fungicide, with the proviso that the biological control agent and the fungicide are not identical. The fungicide can be present either in the biological control agent component of the kit of parts or in the insecticide component of the kit of parts being spatially separated or in both of these components. Preferably, the fungicide is present in the insecticide component.

10 Moreover, the kit of parts according to the present invention can additionally comprise at least one auxiliary selected from the group consisting of extenders, solvents, spontaneity promoters, carriers, emulsifiers, dispersants, frost protectants, thickeners and adjuvants as mentioned below. This at least one auxiliary can be present either in the biological control agent component of the kit of parts or in the insecticide component of the kit of parts being spatially separated or in both of these components.

Use of the composition

In another aspect of the present invention the composition as described above is used for reducing overall damage of plants and plant parts as well as losses in harvested fruits or vegetables caused by insects, nematodes and/or phytopathogens.

20 Furthermore, in another aspect of the present invention the composition as described above increases the overall plant health.

The term "plant health" generally comprises various sorts of improvements of plants that are not connected to the control of pests. For example, advantageous properties that may be mentioned are improved crop characteristics including: emergence, crop yields, protein content, oil content, starch content, more developed root system, improved root growth, improved root size maintenance, improved root effectiveness, improved stress tolerance (e.g. against drought, heat, salt, UV, water, cold), reduced ethylene (reduced production and/or inhibition of reception), tillering increase, increase in plant height, bigger leaf blade, less dead basal leaves, stronger tillers, greener leaf color, pigment content, photosynthetic activity, less input needed (such as fertilizers or water), less seeds needed, more productive tillers, earlier flowering, early grain maturity,

less plant verse (lodging), increased shoot growth, enhanced plant vigor, increased plant stand and early and better germination.

With regard to the use according to the present invention, improved plant health preferably refers to improved plant characteristics including: crop yield, more developed root system (improved root growth), improved root size maintenance, improved root effectiveness, tillering increase, increase in plant height, bigger leaf blade, less dead basal leaves, stronger tillers, greener leaf color, photosynthetic activity, more productive tillers, enhanced plant vigor, and increased plant stand.

With regard to the present invention, improved plant health preferably especially refers to improved plant properties selected from crop yield, more developed root system, improved root growth, improved root size maintenance, improved root effectiveness, tillering increase, and increase in plant height.

The effect of a composition according to the present invention on plant health as defined herein can be determined by comparing plants which are grown under the same environmental conditions, whereby a part of said plants is treated with a composition according to the present invention and another part of said plants is not treated with a composition according to the present invention. Instead, said other part is not treated at all or treated with a placebo (i.e., an application without a composition according to the invention such as an application without all active ingredients (i.e. without a biological control agent as described herein and without an insecticide as described herein), or an application without a biological control agent as described herein, or an application without an insecticide as described herein.

The composition according to the present invention may be applied in any desired manner, such as in the form of a seed coating, soil drench, and/or directly in-furrow and/or as a foliar spray and applied either pre-emergence, post-emergence or both. In other words, the composition can be applied to the seed, the plant or to harvested fruits and vegetables or to the soil wherein the plant is growing or wherein it is desired to grow (plant's locus of growth).

Reducing the overall damage of plants and plant parts often results in healthier plants and/or in an increase in plant vigor and yield.

Preferably, the composition according to the present invention is used for treating conventional or transgenic plants or seed thereof.

In another aspect of the present invention a method for reducing overall damage of plants and plant parts as well as losses in harvested fruits or vegetables caused by insects, nematodes and/or phytopathogens is provided comprising the step of simultaneously or sequentially applying at least one biological control agent selected from the group consisting of *Bacillus chitinosporus* AQ746 (NRRL Accession No. B-21618), *Bacillus mycoides* AQ726 (NRRL Accession No. B-21664), *Bacillus pumilus* (NRRL Accession No. B-30087), *Bacillus pumilus* AQ717 (NRRL Accession No. B-21662), *Bacillus* sp. AQ175 (ATCC Accession No. 55608), *Bacillus* sp. AQ177 (ATCC Accession No. 55609), *Bacillus* sp. AQ178 (ATCC Accession No. 53522), *Bacillus subtilis* AQ743 (NRRL Accession No. B-21665), *Bacillus subtilis* AQ713 (NRRL Accession No. B-21661), *Bacillus subtilis* AQ 153 (ATCC Accession No. 55614), *Bacillus thuringiensis* BD#32 (NRRL Accession No. B-21530), *Bacillus thuringiensis* AQ52 (NRRL Accession No. B-21619), *Muscodor albus* 620 (NRRL Accession No. 30547), *Muscodor roseus* A3-5 (NRRL Accession No. 30548), *Rhodococcus globerulus* AQ719 (NRRL Accession No. B-21663), *Streptomyces galbus* (NRRL Accession No. 30232), *Streptomyces* sp. (NRRL Accession No. B-30145), *Bacillus thuringiensis* subspec. *kurstaki* BMP 123, *Bacillus subtilis* AQ30002 (NRRL Accession No. B-50421), and *Bacillus subtilis* AQ 30004 (NRRL Accession No. B-50455) and/or a mutant of these strains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens and at least one insecticide selected from the group consisting of acetylcholinesterase (AChE) inhibitors, GABA-gated chloride channel antagonists, chloride channel activators and nicotinic acetylcholine receptor (nAChR) channel blockers, and optionally at least one fungicide on the plant, plant parts, harvested fruits, vegetables and/or plant's locus of growth in a synergistically effective amount, with the proviso that the biological control agent and the fungicide are not identical.

In another preferred embodiment of the present method the at least one fungicide is a synthetic fungicide.

The method of the present invention includes the following application methods, namely both of the at least one biological control agent and the at least one insecticide mentioned before may be formulated into a single, stable composition with an agriculturally acceptable shelf life (so called "solo-formulation"), or being combined before or at the time of use (so called "combined-formulations").

If not mentioned otherwise, the expression "combination" stands for the various combinations of the at least one biological control agent and the at least one insecticide, and optionally the at least one fungicide, in a solo-formulation, in a single "ready-mix" form, in a combined spray mixture composed from solo-formulations, such as a "tank-mix", and especially in a combined use of the single active ingredients when applied in a sequential manner, i.e. one after the other within a reasonably short period, such as a few hours or days, e.g. 2 hours to 7 days. The order of applying the composition according to the present invention is not essential for working the present invention. Accordingly, the term "combination" also encompasses the presence of the at least one biological control agent and the at least one insecticide, and optionally the at least one fungicide on or in a plant to be treated or its surrounding, habitat or storage space, e.g. after simultaneously or consecutively applying the at least one biological control agent and the at least one insecticide, and optionally the at least one fungicide to a plant its surrounding, habitat or storage space.

If the at least one biological control agent and the at least one insecticide, and optionally the at least one fungicide are employed or used in a sequential manner, it is preferred to treat the plants or plant parts (which includes seeds and plants emerging from the seed), harvested fruits and vegetables according to the following method: Firstly applying the at least one insecticide and optionally the at least one fungicide on the plant or plant parts, and secondly applying the biological control agent to the same plant or plant parts. The time periods between the first and the second application within a (crop) growing cycle may vary and depend on the effect to be achieved. For example, the first application is done to prevent an infestation of the plant or plant parts with insects, nematodes and/or phytopathogens (this is particularly the case when treating seeds) or to combat the infestation with insects, nematodes and/or phytopathogens (this is particularly the case when treating plants and plant parts) and the second application is done to prevent or control the infestation with insects, nematodes and/or phytopathogens. Control in this context means that the biological control agent is not able to fully exterminate the pests or phytopathogenic fungi but is able to keep the infestation on an acceptable level.

By following the before mentioned steps, a very low level of residues of the at least one specified insecticide, and optionally at least one fungicide on the treated plant, plant parts, and the harvested fruits and vegetables can be achieved.

If not mentioned otherwise the treatment of plants or plant parts (which includes seeds and plants emerging from the seed), harvested fruits and vegetables with the composition according to the invention is carried out directly or by action on their surroundings, habitat or storage space using customary treatment methods, for example dipping, spraying, atomizing, irrigating, evaporating, dusting, fogging, broadcasting, foaming, painting, spreading-on, watering (drenching), drip irrigating. It is furthermore possible to apply the at least one biological control agent, the at least one insecticide, and optionally the at least one fungicide as solo-formulation or combined-formulations by the ultra-low volume method, or to inject the composition according to the present invention as a composition or as solo-formulations into the soil (in-furrow).

The term "plant to be treated" encompasses every part of a plant including its root system and the material - e.g., soil or nutrition medium - which is in a radius of at least 10 cm, 20 cm, 30 cm around the caulis or bole of a plant to be treated or which is at least 10 cm, 20 cm, 30 cm around the root system of said plant to be treated, respectively.

The amount of the biological control agent which is used or employed in combination with the specified insecticide, optionally in the presence of a fungicide, depends on the final formulation as well as size or type of the plant, plant parts, seeds, harvested fruits and vegetables to be treated. Usually, the biological control agent to be employed or used according to the invention is present in about 2 % to about 80 % (w/w), preferably in about 5 % to about 75 % (w/w), more preferably about 10 % to about 70 % (w/w) of its solo-formulation or combined-formulation with the at least one insecticide, and optionally the fungicide.

In a preferred embodiment the biological control agent or e.g. their spores are present in a solo-formulation or the combined-formulation in a concentration of at least 10^5 colony forming units per gram preparation (e. g. cells/g preparation, spores/g preparation), such as $10^5 - 10^{12}$ cfu/g, preferably $10^6 - 10^{11}$ cfu/g, more preferably $10^7 - 10^{10}$ cfu/g and most preferably $10^9 - 10^{10}$ cfu/g at the time point of applying biological control agents on a plant or plant parts such as seeds, fruits or vegetables. Also references to the concentration of biological control agents in form of, e.g., spores or cells - when discussing ratios between the amount of a preparation of at least one biological control agent and the amount of the specified insecticide - are made in view of the time point when the biological control agent is applied on a plant or plant parts such as seeds, fruits or vegetables.

Also the amount of the at least one insecticide which is used or employed in combination with the specified biological control agent, optionally in the presence of a fungicide, depends on the final formulation as well as size or type of the plant, plant parts, seeds, harvested fruit or vegetable to be treated. Usually, the insecticide to be employed or used according to the invention is present in about 0.1 % to about 80 % (w/w), preferably 1 % to about 60 % (w/w), more preferably about 10 % to about 50 % (w/w) of its solo-formulation or combined-formulation with the biological control agent, and optionally the fungicide.

The at least one biological control agent and at least one insecticide, and if present also the fungicide are used or employed in a synergistic weight ratio. The skilled person is able to find out the synergistic weight ratios for the present invention by routine methods. The skilled person understands that these ratios refer to the ratio within a combined-formulation as well as to the calculative ratio of the at least one biological control agent described herein and the specified insecticide when both components are applied as mono-formulations to a plant to be treated. The skilled person can calculate this ratio by simple mathematics since the volume and the amount of the biological control agent and insecticide, respectively, in a mono-formulation is known to the skilled person.

The ratio can be calculated based on the amount of the at least one insecticide, at the time point of applying said component of a combination according to the invention to a plant or plant part and the amount of a biological control agent shortly prior (e.g., 48 h, 24 h, 12 h, 6 h, 2 h, 1 h) or at the time point of applying said component of a combination according to the invention to a plant or plant part.

The application of the at least one biological control agent and the at least one insecticide according to the present invention to a plant or a plant part can take place simultaneously or at different times as long as both components are present on or in the plant after the application(s). In cases where the biological control agent and the insecticide are applied at different times and the insecticide is applied noticeable prior to the biological control agent, the skilled person can determine the concentration of the specified insecticide on/in a plant by chemical analysis known in the art, at the time point or shortly before the time point of applying the biological control agent. Vice versa, when the biological control agent is applied to a plant first, the concentration of a biological control agent can be determined using test which are also known in the art, at the time point or shortly before the time point of applying the insecticide.

In particular, in one embodiment the synergistic weight ratio of the at least one biological control agent/spore preparation and the at least one insecticide lies in the range of 1 : 500 to 1000 : 1, preferably in the range of 1 : 500 to 500 : 1, more preferably in the range of 1 : 500 to 300 : 1. It has to be noted that these ratio ranges refer to the biological control agent/spores preparation (to be combined with at least one insecticide or a preparation of at least one insecticide) of around 10^{10} cells/spores per gram preparation of said cells/spores. For example, a ratio of 100:1 means 100 weight parts of a biological control agent/spore preparation having a cell/ spore concentration of 10^{10} cells/spores per gram preparation and 1 weight part of the insecticide are combined (either as a solo formulation, a combined formulation or by separate applications to plants so that the combination is formed on the plant).

In another embodiment, the synergistic weight ratio of the at least one biological control agent/spore preparation to the insecticide is in the range of 1 : 100 to 20.000 : 1, preferably in the range of 1:50 to 10.000:1 or even in the range of 1:50 to 1000:1. Once again the mentioned ratios ranges refer to biological control agent/spore preparations of biological control agents of around 10^{10} cells or spores per gram preparation of said biological control agent. In particular, in this embodiment the biological control agent preferably is selected from the group consisting of *Muscodora albus* 620 (NRRL Accession No. 30547) and *Muscodora roseus* A3-5 (NRRL Accession No. 30548).

Still in another embodiment, the synergistic weight ratio of the at least one biological control agent/spore preparation to the insecticide is in the range of 1 : 10 to 10.000 : 1, preferably in the range of 1:1 to 7500:1 or even in the range of 10:1 to 6500:1. Once again the mentioned ratios ranges refer to biological control agent/spore preparations of biological control agents of around 10^{10} cells or spores per gram preparation of said biological control agent. In particular, in this embodiment the biological control agent preferably is *Bacillus subtilis* QST 713 (also referred to as B9). Most preferably, when B9 is used as a BCA, the synergistic weight ratio of at least B9 to the insecticide is selected from 20:1, 200:1, 250:1, and 6250:1.

Still in another embodiment, the synergistic weight ratio of the at least one biological control agent/spore preparation to the insecticide is in the range of 1 : 10 to 5000 : 1, preferably in the range of 1:5 to 2500:1 or even in the range of 1:1 to 2000:1. Once again the mentioned ratios ranges refer to biological control agent/spore preparations of biological control agents of around 10^{10} cells or spores per gram preparation of said

biological control agent. In particular, in this embodiment the biological control agent preferably is *Bacillus pumilus* QST 2808 (also referred to as B3). Most preferably, when B3 is used as a BCA, the synergistic weight ratio of at least B3 to the insecticide is selected from 1:1, 5:1, 62.5:1, and 1562.5:1.

- 5 Still in another embodiment, the synergistic weight ratio of the at least one biological control agent/spore preparation to the insecticide is in the range of 1 : 10 to 2500 : 1, preferably in the range of 1:5 to 1000:1 or even in the range of 1:1 to 750:1. Once again the mentioned ratios ranges refer to biological control agent/spore preparations of biological control agents of around 10^{10} cells or spores per gram preparation of said
- 10 biological control agent. In particular, in this embodiment the biological control agent preferably is *Streptomyces galbus* which is mentioned above as B16. Most preferably, when B16 is used as a BCA, the synergistic weight ratio of at least B16 to the insecticide is selected from 500:1, and 4:1.

- Still in another embodiment, the synergistic weight ratio of the at least one biological control agent/spore preparation to the insecticide is in the range of 1 : 10 to 1500 : 1, preferably in the range of 1:1 to 1000:1 or even in the range of 100:1 to 900:1. Once again the mentioned ratios ranges refer to biological control agent/spore preparations of biological control agents of around 10^{10} cells or spores per gram preparation of said
- 15 biological control agent. In particular, in this embodiment the biological control agent preferably is *Bacillus subtilis* AQ30002 which is mentioned above as B19. Most preferably, when B19 is used as a BCA, the synergistic weight ratio of at least B19 to the insecticide is selected from 125:1, and 800:1.

- The cell/spore concentration of preparations can be determined by applying methods known in the art. To compare weight ratios of the biological control agent/ spore
- 25 preparation to the insecticide, the skilled person can easily determine the factor between a preparation having a biological control agent/spore concentration different from 10^{10} cells/spores per gram cell/spore preparation and a preparation having a biological control agent/ spore concentration of 10^{10} cells/spores per gram preparation to calculate whether a ratio of a biological control agent/spore preparation to the
- 30 insecticide is within the scope of the above listed ratio ranges.

In one embodiment of the present invention, the concentration of the biological control agent after dispersal is at least 50 g/ha, such as 50 – 7500 g/ha, 50 – 2500 g/ha, 50 – 1500 g/ha; at least 250 g/ha (hectare), at least 500 g/ha or at least 800 g/ha.

The application rate of composition to be employed or used according to the present invention may vary. The skilled person is able to find the appropriate application rate by way of routine experiments.

Seed treatment

- 5 In another aspect of the present invention a seed treated with the composition as described above is provided.

The control of insects, nematodes and/or phytopathogens by treating the seed of plants has been known for a long time and is a subject of continual improvements. Nevertheless, the treatment of seed entails a series of problems which cannot always
10 be solved in a satisfactory manner. Thus, it is desirable to develop methods for protecting the seed and the germinating plant that remove the need for, or at least significantly reduce, the additional delivery of crop protection compositions in the course of storage, after sowing or after the emergence of the plants. It is desirable, furthermore, to optimize the amount of active ingredient employed in such a way as to
15 provide the best-possible protection to the seed and the germinating plant from attack by insects, nematodes and/or phytopathogens, but without causing damage to the plant itself by the active ingredient employed. In particular, methods for treating seed ought also to take into consideration the intrinsic insecticidal and/or nematocidal properties of pest-resistant or pest-tolerant transgenic plants, in order to achieve
20 optimum protection of the seed and of the germinating plant with a minimal use of crop protection compositions.

The present invention therefore also relates in particular to a method for protecting seed and germinating plants from attack by pests, by treating the seed with at least one biological control agent as defined above and/or a mutant of it having all identifying
25 characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens and at least one insecticide as defined above and optionally at least one fungicide of the invention. The method of the invention for protecting seed and germinating plants from attack by pests encompasses a method in which the seed is treated
30 simultaneously in one operation with the at least one biological control agent and the at least one insecticide, and optionally the at least one fungicide. It also encompasses a method in which the seed is treated at different times with the at least one biological

control agent and the at least one insecticide, and optionally the at least one fungicide.

The invention relates to the use of the composition of the invention for treating seed for the purpose of protecting the seed and the resultant plant against insects, mites,
5 nematodes and/or phytopathogens.

The invention also relates to seed which at the same time has been treated with at least one biological control agent and at least one insecticide according to the present invention, and optionally at least one fungicide. The invention further relates to seed which has been treated at different times with the at least one biological control agent
10 and the at least one insecticide, and optionally the at least one fungicide. In the case of seed which has been treated at different times with the at least one biological control agent and the at least one insecticide, and optionally the at least one fungicide, the individual active ingredients in the composition of the invention may be present in different layers on the seed.

15 Furthermore, the invention relates to seed which, following treatment with the composition of the invention, is subjected to a film-coating process in order to prevent dust abrasion of the seed.

One of the advantages of the present invention is that, owing to the particular systemic properties of the compositions of the invention, the treatment of the seed with these
20 compositions provides protection from insects, nematodes and/or phytopathogens not only to the seed itself but also to the plants originating from the seed, after they have emerged. In this way, it may not be necessary to treat the crop directly at the time of sowing or shortly thereafter.

A further advantage is to be seen in the fact that, through the treatment of the seed
25 with composition of the invention, germination and emergence of the treated seed may be promoted.

It is likewise considered to be advantageous composition of the invention may also be used, in particular, on transgenic seed.

It is also stated that the composition of the invention may be used in combination with
30 agents of the signalling technology, as a result of which, for example, colonization with

symbionts is improved, such as rhizobia, mycorrhiza and/or endophytic bacteria, for example, is enhanced, and/or nitrogen fixation is optimized.

The compositions of the invention are suitable for protecting seed of any variety of plant which is used in agriculture, in greenhouses, in forestry or in horticulture. More particularly, the seed in question is that of cereals (e.g. wheat, barley, rye, oats and millet), maize, cotton, soybeans, rice, potatoes, sunflower, coffee, tobacco, canola, oilseed rape, beets (e.g. sugar beet and fodder beet), peanuts, vegetables (e.g. tomato, cucumber, bean, brassicas, onions and lettuce), fruit plants, lawns and ornamentals. Particularly important is the treatment of the seed of cereals (such as wheat, barley, rye and oats) maize, soybeans, cotton, canola, oilseed rape and rice.

As already mentioned above, the treatment of transgenic seed with the composition of the invention is particularly important. The seed in question here is that of plants which generally contain at least one heterologous gene that controls the expression of a polypeptide having, in particular, insecticidal and/or nematicidal properties. These heterologous genes in transgenic seed may come from microorganisms such as *Bacillus*, *Rhizobium*, *Pseudomonas*, *Serratia*, *Trichoderma*, *Clavibacter*, *Glomus* or *Gliocladium*. The present invention is particularly suitable for the treatment of transgenic seed which contains at least one heterologous gene from *Bacillus* sp. With particular preference, the heterologous gene in question comes from *Bacillus thuringiensis*.

For the purposes of the present invention, the composition of the invention is applied alone or in a suitable formulation to the seed. The seed is preferably treated in a condition in which its stability is such that no damage occurs in the course of the treatment. Generally speaking, the seed may be treated at any point in time between harvesting and sowing. Typically, seed is used which has been separated from the plant and has had cobs, hulls, stems, husks, hair or pulp removed. Thus, for example, seed may be used that has been harvested, cleaned and dried to a moisture content of less than 15% by weight. Alternatively, seed can also be used that after drying has been treated with water, for example, and then dried again.

When treating seed it is necessary, generally speaking, to ensure that the amount of the composition of the invention, and/or of other additives, that is applied to the seed is selected such that the germination of the seed is not adversely affected, and/or that the plant which emerges from the seed is not damaged. This is the case in particular with active ingredients which may exhibit phytotoxic effects at certain application rates.

The compositions of the invention can be applied directly, in other words without comprising further components and without having been diluted. As a general rule, it is preferable to apply the compositions in the form of a suitable formulation to the seed. Suitable formulations and methods for seed treatment are known to the skilled person and are described in, for example, the following documents: US 4,272,417 A, 5 US 4,245,432 A, US 4,808,430 A, US 5,876,739 A, US 2003/0176428 A1, WO 2002/080675 A1, WO 2002/028186 A2.

The combinations which can be used in accordance with the invention may be converted into the customary seed-dressing formulations, such as solutions, emulsions, 10 suspensions, powders, foams, slurries or other coating compositions for seed, and also ULV formulations.

These formulations are prepared in a known manner, by mixing composition with customary adjuvants, such as, for example, customary extenders and also solvents or diluents, colorants, wetters, dispersants, emulsifiers, antifoams, preservatives, secondary 15 thickeners, stickers, gibberellins, and also water.

Colorants which may be present in the seed-dressing formulations which can be used in accordance with the invention include all colorants which are customary for such purposes. In this context it is possible to use not only pigments, which are of low solubility in water, but also water-soluble dyes. Examples include the colorants known under the 20 designations Rhodamin B, C.I. Pigment Red 112 and C.I. Solvent Red 1.

Wetters which may be present in the seed-dressing formulations which can be used in accordance with the invention include all of the substances which promote wetting and which are customary in the formulation of active agrochemical ingredients. Use may be made preferably of alkylnaphthalenesulphonates, such as diisopropyl- or 25 diisobutyl-naphthalenesulphonates.

Dispersants and/or emulsifiers which may be present in the seed-dressing formulations which can be used in accordance with the invention include all of the nonionic, anionic and cationic dispersants that are customary in the formulation of active agrochemical ingredients. Use may be made preferably of nonionic or anionic 30 dispersants or of mixtures of nonionic or anionic dispersants. Suitable nonionic dispersants are, in particular, ethylene oxide-propylene oxide block polymers, alkylphenol polyglycol ethers and also tristyrylphenol polyglycol ethers, and the

phosphated or sulphated derivatives of these. Suitable anionic dispersants are, in particular, lignosulphonates, salts of polyacrylic acid, and arylsulphonate-formaldehyde condensates.

5 Antifoams which may be present in the seed-dressing formulations which can be used in accordance with the invention include all of the foam inhibitors that are customary in the formulation of active agrochemical ingredients. Use may be made preferably of silicone antifoams and magnesium stearate.

10 Preservatives which may be present in the seed-dressing formulations which can be used in accordance with the invention include all of the substances which can be employed for such purposes in agrochemical compositions. Examples include dichlorophen and benzyl alcohol hemiformal.

15 Secondary thickeners which may be present in the seed-dressing formulations which can be used in accordance with the invention include all substances which can be used for such purposes in agrochemical compositions. Those contemplated with preference include cellulose derivatives, acrylic acid derivatives, xanthan, modified clays and highly disperse silica.

20 Stickers which may be present in the seed-dressing formulations which can be used in accordance with the invention include all customary binders which can be used in seed-dressing products. Preferred mention may be made of polyvinylpyrrolidone, polyvinyl acetate, polyvinyl alcohol and tylose.

25 Gibberellins which may be present in the seed-dressing formulations which can be used in accordance with the invention include preferably the gibberellins A1, A3 (= gibberellic acid), A4 and A7, with gibberellic acid being used with particular preference. The gibberellins are known (cf. R. Wegler, "Chemie der Pflanzenschutz- und Schädlingsbekämpfungsmittel", Volume 2, Springer Verlag, 1970, pp. 401-412).

30 The seed-dressing formulations which can be used in accordance with the invention may be used, either directly or after prior dilution with water, to treat seed of any of a wide variety of types. Accordingly, the concentrates or the preparations obtainable from them by dilution with water may be employed to dress the seed of cereals, such as wheat, barley, rye, oats and triticale, and also the seed of maize, rice, oilseed rape, peas, beans, cotton, sunflowers and beets, or else the seed of any of a very wide variety of vegetables. The seed-dressing formulations which can be used in

accordance with the invention, or their diluted preparations, may also be used to dress seed of transgenic plants. In that case, additional synergistic effects may occur in interaction with the substances formed through expression.

For the treatment of seed with the seed-dressing formulations which can be used in accordance with the invention, or with the preparations produced from them by addition of water, suitable mixing equipment includes all such equipment which can typically be employed for seed dressing. More particularly, the procedure when carrying out seed dressing is to place the seed in a mixer, to add the particular desired amount of seed-dressing formulations, either as such or following dilution with water beforehand, and to carry out mixing until the distribution of the formulation on the seed is uniform. This may be followed by a drying operation.

The application rate of the seed-dressing formulations which can be used in accordance with the invention may be varied within a relatively wide range. It is guided by the particular amount of the at least one biological control agent and the at least one insecticide in the formulations, and by the seed. The application rates in the case of the composition are situated generally at between 0.001 and 50 g per kilogram of seed, preferably between 0.01 and 15 g per kilogram of seed.

The composition according to the invention, in combination with good plant tolerance and favourable toxicity to warm-blooded animals and being tolerated well by the environment, are suitable for protecting plants and plant organs, for increasing harvest yields, for improving the quality of the harvested material and for controlling animal pests, in particular insects, arachnids, helminths, nematodes and molluscs, which are encountered in agriculture, in horticulture, in animal husbandry, in forests, in gardens and leisure facilities, in protection of stored products and of materials, and in the hygiene sector. They can be preferably employed as plant protection agents. In particular, the present invention relates to the use of the composition according to the invention as insecticide and/or fungicide.

The present composition preferably is active against normally sensitive and resistant species and against all or some stages of development. The abovementioned pests include:

pests from the phylum Arthropoda, especially from the class Arachnida, for example, *Acarus* spp., *Aceria sheldoni*, *Aculops* spp., *Aculus* spp., *Amblyomma* spp.,

Amphitetranychus viennensis, Argas spp., Boophilus spp., Brevipalpus spp., Bryobia graminum, Bryobia praetiosa, Centruroides spp., Chorioptes spp., Dermanyssus gallinae, Dermatophagoides pteronyssinus, Dermatophagoides farinae, Dermacentor spp., Eotetranychus spp., Epitrimerus pyri, Eutetranychus spp., Eriophyes spp.,
 5 Glycyphagus domesticus, Halotydeus destructor, Hemitarsonemus spp., Hyalomma spp., Ixodes spp., Latrodectus spp., Loxosceles spp., Metatetranychus spp., Neutrombicula autumnalis, Nuphessa spp., Oligonychus spp., Ornithodoros spp., Ornithonyssus spp., Panonychus spp., Phyllocoptruta oleivora, Polyphagotarsonemus latus, Psoroptes spp., Rhipicephalus spp., Rhizoglyphus spp., Sarcoptes spp., Scorpio
 10 maurus, Steneotarsonemus spp., Steneotarsonemus spinki, Tarsonemus spp., Tetranychus spp., Trombicula alfreddugesi, Vaejovis spp., Vasates lycopersici;

from the class Chilopoda, for example, Geophilus spp., Scutigera spp.;

from the order or the class Collembola, for example, Onychiurus armatus;

from the class Diplopoda, for example, Blaniulus guttulatus;

15 from the class Insecta, e.g. from the order Blattodea, for example, Blattella asahinai, Blattella germanica, Blatta orientalis, Leucophaea maderae, Panchlora spp., Parcoblatta spp., Periplaneta spp., Supella longipalpa;

from the order Coleoptera, for example, Acalymma vittatum, Acanthoscelides obtectus, Adoretus spp., Agelastica alni, Agriotes spp., Alphitobius diaperinus,
 20 Amphimallon solstitialis, Anobium punctatum, Anoplophora spp., Anthonomus spp., Anthrenus spp., Apion spp., Apogonia spp., Atomaria spp., Attagenus spp., Bruchidius obtectus, Bruchus spp., Cassida spp., Cerotoma trifurcata, Ceutorhynchus spp., Chaetocnema spp., Cleonus mendicus, Conoderus spp., Cosmopolites spp., Costelytra zealandica, Ctenicera spp., Curculio spp., Cryptolestes ferrugineus,
 25 Cryptorhynchus lapathi, Cylindrocopturus spp., Dermestes spp., Diabrotica spp., Dichocrocis spp., Dieladspa armigera, Diloboderus spp., Epilachna spp., Epitrix spp., Faustinus spp., Gibbium psylloides, Gnathocerus cornutus, Hellula undalis, Heteronychus arator, Heteronyx spp., Hylamorpha elegans, Hylotrupes bajulus, Hypera postica, Hypomeces squamosus, Hypothenemus spp., Lachnosterna consanguinea,
 30 Lasioderma serricorne, Latheticus oryzae, Lathridius spp., Lema spp., Leptinotarsa decemlineata, Leucoptera spp., Lissorhoptrus oryzophilus, Lixus spp., Luperodes spp., Lyctus spp., Megascelis spp., Melanotus spp., Meligethes aeneus, Melolontha spp.,

Migdolus spp., Monochamus spp., Naupactus xanthographus, Necrobia spp., Niptus hololeucus, Oryctes rhinoceros, Oryzaephilus surinamensis, Oryzaphagus oryzae, Otiorrhynchus spp., Oxycetonia jucunda, Phaedon cochleariae, Phyllophaga spp., Phyllophaga helleri, Phyllostreta spp., Popillia japonica, Premnotypes spp., Prostephanus truncatus, Psylliodes spp., Ptinus spp., Rhizobius ventralis, Rhizopertha dominica, Sitophilus spp., Sitophilus oryzae, Sphenophorus spp., Stegobium paniceum, Sternechus spp., Symphyletes spp., Tanymecus spp., Tenebrio molitor, Tenebrioides mauretanicus, Tribolium spp., Trogoderma spp., Tychius spp., Xylotrechus spp., Zabrus spp.;

from the order Diptera, for example, Aedes spp., Agromyza spp., Anastrepha spp., Anopheles spp., Asphondylia spp., Bactrocera spp., Bibio hortulanus, Calliphora erythrocephala, Calliphora vicina, Ceratitis capitata, Chironomus spp., Chrysomyia spp., Chrysops spp., Chrysozona pluvialis, Cochliomyia spp., Contarinia spp., Cordylobia anthropophaga, Cricotopus sylvestris, Culex spp., Culicoides spp., Culiceta spp., Cuterebra spp., Dacus oleae, Dasyneura spp., Della spp., Dermatobia hominis, Drosophila spp., Echinocnemus spp., Fannia spp., Gasterophilus spp., Glossina spp., Haematopota spp., Hydrellia spp., Hydrellia griseola, Hylemya spp., Hippobosca spp., Hypoderma spp., Liriomyza spp., Lucilia spp., Lutzomyia spp., Mansonia spp., Musca spp., Oestrus spp., Oscinella frit, Paratanytarsus spp., Paralauterborniella subcincta, Pegomyia spp., Phlebotomus spp., Phorbia spp., Phormia spp., Piophilina casei, Prodiplosis spp., Psila rosae, Rhagoletis spp., Sarcophaga spp., Simulium spp., Stomoxys spp., Tabanus spp., Tetanops spp., Tipula spp.;

from the order Heteroptera, for example, Anasa tristis, Antestiopsis spp., Boisea spp., Blissus spp., Calocoris spp., Campylomma livida, Cavelerius spp., Cimex spp., Collaria spp., Creontiades dilutus, Dasynus piperis, Dichelops furcatus, Diconocoris hewetti, Dysdercus spp., Euschistus spp., Eurygaster spp., Heliopeltis spp., Horcias nobilellus, Leptocorisa spp., Leptocorisa varicornis, Leptoglossus phyllopus, Lygus spp., Macropes excavatus, Miridae, Monalonion atratum, Nezara spp., Oebalus spp., Pentomidae, Piesma quadrata, Piezodorus spp., Psallus spp., Pseudacysta perseae, Rhodnius spp., Sahlbergella singularis, Scaptocoris castanea, Scotinophora spp., Stephanitis nashi, Tibraca spp., Triatoma spp.;

from the order Homoptera, for example, Acizzia acaciaebaileyanae, Acizzia dodonaeae, Acizzia uncatoides, Acrida turrita, Acyrthosipon spp., Acrogonia spp., Aeneolamia spp., Agonosцена spp., Aleyrodes proletella, Aleurolobus barodensis,

- Aleurothrixus floccosus, Allocaridara malayensis, Amrasca spp., Anuraphis cardui,
 Aonidiella spp., Aphanostigma piri, Aphis spp., Arboridia apicalis, Aryanilla spp.,
 Aspidiella spp., Aspidiotus spp., Atanus spp., Aulacorthum solani, Bemisia tabaci,
 Blastopsylla occidentalis, Boreioglycaspis melaleucae, Brachycaudus helichrysi,
 5 Brachycolus spp., Brevicoryne brassicae, Cacopsylla spp., Calligypona marginata,
 Carneiocephala fulgida, Ceratovacuna lanigera, Cercopidae, Ceroplastes spp.,
 Chaetosiphon fragaefolii, Chionaspis tegalensis, Chlorita onukii, Chondraxis rosea,
 Chromaphis juglandicola, Chrysomphalus ficus, Cicadulina mbila, Cocomytilus halli,
 Coccus spp., Cryptomyzus ribis, Cryptoneossa spp., Ctenarytaina spp., Dalbulus spp.,
 10 Dialeurodes citri, Diaphorina citri, Diaspis spp., Drosicha spp., Dysaphis spp.,
 Dymicoccus spp., Empoasca spp., Eriosoma spp., Erythroneura spp., Eucalyptolyma
 spp., Euphyllura spp., Euscelis bilobatus, Ferrisia spp., Geococcus coffeae, Glycaspis
 spp., Heteropsylla cubana, Heteropsylla spinulosa, Homalodisca coagulata,
 Hyalopterus arundinis, Icerya spp., Idiocerus spp., Idioscopus spp., Laodelphax
 15 striatellus, Lecanium spp., Lepidosaphes spp., Lipaphis erysimi, Macrosiphum spp.,
 Macrosteles facifrons, Mahanarva spp., Melanaphis sacchari, Metcalfiella spp.,
 Metopolophium dirhodum, Monella costalis, Monelliopsis pecanis, Myzus spp.,
 Nasonovia ribisnigri, Nephrotettix spp., Nettigoniclla spectra, Nilaparvata lugens,
 Oncometopia spp., Orthezia praelonga, Oxya chinensis, Pachyopsylla spp., Parabemisia
 20 myricae, Paratrioza spp., Parlatoria spp., Pemphigus spp., Peregrinus maidis,
 Phenacoccus spp., Phloeomyzus passerinii, Phorodon humuli, Phylloxera spp., Pinnaspis
 aspidistrae, Planococcus spp., Prosopidopsylla flava, Protopulvinaria pyriformis,
 Pseudaulacaspis pentagona, Pseudococcus spp., Psyllopsis spp., Psylla spp.,
 Pteromalus spp., Pyrilla spp., Quadraspidiotus spp., Quesada gigas, Rastrococcus spp.,
 25 Rhopalosiphum spp., Saissetia spp., Scaphoideus titanus, Schizaphis graminum,
 Selenaspidus articulatus, Sogata spp., Sogatella furcifera, Sogatodes spp.,
 Stictocephala festina, Siphoninus phillyreae, Tenalaphara malayensis,
 Tetranocephala spp., Tinocallis caryaefoliae, Tomaspis spp., Toxoptera spp.,
 Trialeurodes vaporariorum, Trioza spp., Typhlocyba spp., Unaspis spp., Viteus vitifolii,
 30 Zyginia spp.;

from the order Hymenoptera, for example, Acromyrmex spp., Athalia spp., Atta spp.,
 Diprion spp., Hoplocampa spp., Lasius spp., Monomorium pharaonis, Sirex spp.,
 Solenopsis invicta, Tapinoma spp., Urocerus spp., Vespa spp., Xeris spp.;

from the order Isopoda, for example, *Armadillidium vulgare*, *Oniscus asellus*, *Porcellio scaber*;

from the order Isoptera, for example, *Coptotermes* spp., *Cornitermes cumulans*, *Cryptotermes* spp., *Incisitermes* spp., *Microtermes obesi*, *Odontotermes* spp.,
5 *Reticulitermes* spp.;

from the order Lepidoptera, for example, *Achroia grisella*, *Acronicta major*, *Adoxophyes* spp., *Aedia leucomelas*, *Agrotis* spp., *Alabama* spp., *Amyelois transitella*, *Anarsia* spp., *Anticarsia* spp., *Argyroplote* spp., *Barathra brassicae*, *Borbo cinnara*, *Bucculatrix thurberiella*, *Bupalus piniarius*, *Busseola* spp., *Cacoecia* spp., *Caloptilia theivora*, *Capua*
10 *reticulana*, *Carpocapsa pomonella*, *Carposina niponensis*, *Chelmatobia brumata*, *Chilo* spp., *Choristoneura* spp., *Clysia ambiguella*, *Cnaphalocerus* spp., *Cnaphalocrocis medinalis*, *Cnephasia* spp., *Conopomorpha* spp., *Conotrachelus* spp., *Copitarsia* spp., *Cydia* spp., *Dalaca noctuides*, *Diaphania* spp., *Diatraea saccharalis*, *Earias* spp., *Ecdytolopha aurantium*, *Elasmopalpus lignosellus*, *Eldana saccharina*,
15 *Ephestia* spp., *Epinotia* spp., *Epiphyas postvittana*, *Etiella* spp., *Eulia* spp., *Eupoecilia ambiguella*, *Euproctis* spp., *Euxoa* spp., *Feltia* spp., *Galleria mellonella*, *Gracillaria* spp., *Grapholitha* spp., *Hedylepta* spp., *Helicoverpa* spp., *Heliothis* spp., *Hofmannophila pseudospretella*, *Homoeosoma* spp., *Homona* spp., *Hyponomeuta padella*, *Kakivoria flavofasciata*, *Laphygma* spp., *Laspeyresia molesta*, *Leucinodes orbonalis*, *Leucoptera*
20 spp., *Lithocolletis* spp., *Lithophane antennata*, *Lobesia* spp., *Loxagrotis albicosta*, *Lymantria* spp., *Lyonetia* spp., *Malacosoma neustria*, *Maruca testulalis*, *Mamstra brassicae*, *Melanitis leda*, *Mocis* spp., *Monopis obviella*, *Mythimna separata*, *Nemapogon cloacellus*, *Nymphula* spp., *Oiketicus* spp., *Oria* spp., *Orthaga* spp., *Ostrinia* spp., *Oulema oryzae*, *Panolis flammea*, *Parnara* spp., *Pectinophora* spp.,
25 *Perilleucoptera* spp., *Phthorimaea* spp., *Phyllocnistis citrella*, *Phyllonorycter* spp., *Pieris* spp., *Platynota stultana*, *Plodia interpunctella*, *Plusia* spp., *Plutella xylostella*, *Prays* spp., *Prodenia* spp., *Protoparce* spp., *Pseudaletia* spp., *Pseudaletia unipuncta*, *Pseudoplusia includens*, *Pyrausta nubilalis*, *Rachiplusia nu*, *Schoenobius* spp., *Scirpophaga* spp., *Scirpophaga innotata*, *Scotia segetum*, *Sesamia* spp., *Sesamia inferens*, *Sparganothis*
30 spp., *Spodoptera* spp., *Spodoptera praefica*, *Stathmopoda* spp., *Stomopteryx subsecivella*, *Synanthedon* spp., *Tecia solanivora*, *Thermesia gemmatalis*, *Tinea cloacella*, *Tinea pellionella*, *Tineola bisselliella*, *Tortrix* spp., *Trichophaga tapetzella*, *Trichoplusia* spp., *Tryporyza incertulas*, *Tuta absoluta*, *Virachola* spp.;

from the order Orthoptera or Saltatoria, for example, *Acheta domesticus*, *Dichroplus* spp., *Gryllotalpa* spp., *Hieroglyphus* spp., *Locusta* spp., *Melanoplus* spp., *Schistocerca gregaria*;

5 from the order Phthiraptera, for example, *Damalinea* spp., *Haematopinus* spp., *Linognathus* spp., *Pediculus* spp., *Ptirus pubis*, *Trichodectes* spp.;

from the order Psocoptera for example *Lepinatus* spp., *Liposcelis* spp.;

from the order Siphonaptera, for example, *Ceratophyllus* spp., *Ctenocephalides* spp., *Pulex irritans*, *Tunga penetrans*, *Xenopsylla cheopsis*;

10 from the order Thysanoptera, for example, *Anaphothrips obscurus*, *Baliothrips biformis*, *Drepanothrips reuteri*, *Enneothrips flavens*, *Frankliniella* spp., *Heliethrips* spp., *Hercinothrips femoralis*, *Rhipiphorocthris cruentatus*, *Scirtothrips* spp., *Taeniothrips cardamomi*, *Thrips* spp.;

from the order Zygentoma (=Thysanura), for example, *Ctenolepisma* spp., *Lepisma saccharina*, *Lepismodes inquilinus*, *Thermobia domestica*;

15 from the class Symphyla, for example, *Scutigera* spp.;

pests from the phylum Mollusca, especially from the class Bivalvia, for example, *Dreissena* spp., and from the class Gastropoda, for example, *Arion* spp., *Biomphalaria* spp., *Bulinus* spp., *Deroceras* spp., *Galba* spp., *Lymnaea* spp., *Oncomelania* spp., *Pomacea* spp., *Succinea* spp.;

20 animal pests from the phyla Plathelminthes and Nematoda, for example, *Ancylostoma duodenale*, *Ancylostoma ceylanicum*, *Ancylostoma braziliense*, *Ancylostoma* spp., *Ascaris* spp., *Brugia malayi*, *Brugia timori*, *Bunostomum* spp., *Chabertia* spp., *Clonorchis* spp., *Cooperia* spp., *Dicrocoelium* spp., *Dictyocaulus filaria*, *Diphylobothrium latum*, *Dracunculus medinensis*, *Echinococcus granulosus*,
 25 *Echinococcus multilocularis*, *Enterobius vermicularis*, *Fasciola* spp., *Haemonchus* spp., *Heterakis* spp., *Hymenolepis nana*, *Hyostrogylus* spp., *Loa Loa*, *Nematodirus* spp., *Oesophagostomum* spp., *Opisthorchis* spp., *Onchocerca volvulus*, *Ostertagia* spp., *Paragonimus* spp., *Schistosoma* spp., *Strongyloides fuelleborni*, *Strongyloides stercoralis*, *Strongyloides* spp., *Taenia saginata*, *Taenia solium*, *Trichinella spiralis*,

Trichinella nativa, Trichinella britovi, Trichinella nelsoni, Trichinella pseudopsiralis, Trichostrongylus spp., Trichuris trichuria, Wuchereria bancrofti;

phytoparasitic pests from the phylum Nematoda, for example, Aphelenchoides spp., Bursaphelenchus spp., Ditylenchus spp., Globodera spp., Heterodera spp., Longidorus
 5 spp., Meloidogyne spp., Pratylenchus spp., Radopholus spp., Trichodorus spp., Tylenchulus spp., Xiphinema spp., Helicotylenchus spp., Tylenchorhynchus spp., Scutellonema spp., Paratrichodorus spp., Meloinema spp., Paraphelenchus spp., Aglenchus spp., Belonolaimus spp., Nacobbus spp., Rotylenchulus spp., Rotylenchus
 10 spp., Neotylenchus spp., Paraphelenchus spp., Dolichodorus spp., Hoplolaimus spp., Punctodera spp., Criconemella spp., Quinisulcius spp., Hemicyclophora spp., Anguina spp., Subanguina spp., Hemicriconemoides spp., Psilenchus spp., Pseudohalenchus spp., Criconemoides spp., Cacopaurus spp., Hirschmaniella spp., *Tetylenchus* spp.,

It is furthermore possible to control organisms from the subphylum Protozoa, especially from the order Coccidia, such as Eimeria spp.

15 The present composition preferably is active against Myzus persicae, Tetranychus urticae, Phaedon cochleariae, Nilaparvata lugens, and/or Spodoptera frugiperda.

Furthermore, in case the biological control agent exhibits fungicidal activity and/or the composition additionally comprises a fungicide, the composition according to the present invention has potent microbicidal activity and can be used for control of
 20 unwanted microorganisms, such as fungi and bacteria, in crop protection and in the protection of materials.

The invention also relates to a method for controlling unwanted microorganisms, characterized in that the inventive composition is applied to the phytopathogenic fungi, phytopathogenic bacteria and/or their habitat.

25 Fungicides can be used in crop protection for control of phytopathogenic fungi. They are characterized by an outstanding efficacy against a broad spectrum of phytopathogenic fungi, including soilborne pathogens, which are in particular members of the classes *Plasmodiophoromycetes*, *Peronosporomycetes* (Syn. *Oomycetes*), *Chytridiomycetes*, *Zygomycetes*, *Ascomycetes*, *Basidiomycetes* and *Deuteromycetes*
 30 (Syn. *Fungi imperfecti*). Some fungicides are systemically active and can be used in plant protection as foliar, seed dressing or soil fungicide. Furthermore, they are suitable for combating fungi, which inter alia infest wood or roots of plant.

Bactericides can be used in crop protection for control of *Pseudomonadaceae*, *Rhizobiaceae*, *Enterobacteriaceae*, *Corynebacteriaceae* and *Streptomycetaceae*.

Non-limiting examples of pathogens of fungal diseases which can be treated in accordance with the invention include:

5 diseases caused by powdery mildew pathogens, for example *Blumeria* species, for example *Blumeria graminis*; *Podosphaera* species, for example *Podosphaera leucotricha*; *Sphaerotheca* species, for example *Sphaerotheca fuliginea*; *Uncinula* species, for example *Uncinula necator*;

diseases caused by rust disease pathogens, for example *Gymnosporangium* species, 10 for example *Gymnosporangium sabiniae*; *Hemileia* species, for example *Hemileia vastatrix*; *Phakopsora* species, for example *Phakopsora pachyrhizi* and *Phakopsora melbomiae*; *Puccinia* species, for example *Puccinia recondite*, *P. triticea*, *P. graminis* or *P. striiformis*; *Uromyces* species, for example *Uromyces appendiculatus*;

diseases caused by pathogens from the group of the Oomycetes, for example *Albugo* 15 species, for example *Albugo candida*; *Bremia* species, for example *Bremia lactucae*; *Peronospora* species, for example *Peronospora pisi* or *P. brassicae*; *Phytophthora* species, for example *Phytophthora infestans*; *Plasmopara* species, for example *Plasmopara viticola*; *Pseudoperonospora* species, for example *Pseudoperonospora humuli* or *Pseudoperonospora cubensis*; *Pythium* species, for example *Pythium* 20 *ultimum*;

leaf blotch diseases and leaf wilt diseases caused, for example, by *Alternaria* species, for example *Alternaria solani*; *Cercospora* species, for example *Cercospora beticola*; *Cladosporium* species, for example *Cladosporium cucumerinum*; *Cochliobolus* species, for example *Cochliobolus sativus* (conidia form: *Drechlera*, Syn: 25 *Helminthosporium*), *Cochliobolus miyabeanus*; *Colletotrichum* species, for example *Colletotrichum lindemuthianum*; *Cycloconium* species, for example *Cycloconium oleaginum*; *Diaporthe* species, for example *Diaporthe citri*; *Elsinoe* species, for example *Elsinoe fawcettii*; *Gloeosporium* species, for example *Gloeosporium laeticolor*; *Glomerella* species, for example *Glomerella cingulata*; *Guignardia* species, 30 for example *Guignardia bidwellii*; *Leptosphaeria* species, for example *Leptosphaeria maculans*, *Leptosphaeria nodorum*; *Magnaporthe* species, for example *Magnaporthe grisea*; *Microdochium* species, for example *Microdochium nivale*; *Mycosphaerella*

species, for example *Mycosphaerella graminicola*, *M. arachidicola* and *M. fijiensis*; *Phaeosphaeria* species, for example *Phaeosphaeria nodorum*; *Pyrenophora* species, for example *Pyrenophora teres*, *Pyrenophora tritici repentis*; *Ramularia* species, for example *Ramularia collo-cygni*, *Ramularia areola*; *Rhynchosporium* species, for example *Rhynchosporium secalis*; *Septoria* species, for example *Septoria apii*, *Septoria lycopersii*; *Typhula* species, for example *Typhula incarnata*; *Venturia* species, for example *Venturia inaequalis*;

root and stem diseases caused, for example, by *Corticium* species, for example *Corticium graminearum*; *Fusarium* species, for example *Fusarium oxysporum*; *Gaeumannomyces* species, for example *Gaeumannomyces graminis*; *Rhizoctonia* species, such as, for example *Rhizoctonia solani*; *Sarocladium* diseases caused for example by *Sarocladium oryzae*; *Sclerotium* diseases caused for example by *Sclerotium oryzae*; *Tapesia* species, for example *Tapesia acuformis*; *Thielaviopsis* species, for example *Thielaviopsis basicola*;

ear and panicle diseases (including corn cobs) caused, for example, by *Alternaria* species, for example *Alternaria* spp.; *Aspergillus* species, for example *Aspergillus flavus*; *Cladosporium* species, for example *Cladosporium cladosporioides*; *Claviceps* species, for example *Claviceps purpurea*; *Fusarium* species, for example *Fusarium culmorum*; *Gibberella* species, for example *Gibberella zeae*; *Monographella* species, for example *Monographella nivalis*; *Septoria* species, for example *Septoria nodorum*;

diseases caused by smut fungi, for example *Sphacelotheca* species, for example *Sphacelotheca relliana*; *Tilletia* species, for example *Tilletia caries*, *T. controversa*; *Urocystis* species, for example *Urocystis occulta*; *Ustilago* species, for example *Ustilago nuda*, *U. nuda tritici*;

fruit rot caused, for example, by *Aspergillus* species, for example *Aspergillus flavus*; *Botrytis* species, for example *Botrytis cinerea*; *Penicillium* species, for example *Penicillium expansum* and *P. purpurogenum*; *Sclerotinia* species, for example *Sclerotinia sclerotiorum*; *Verticillium* species, for example *Verticillium albo-atrum*;

seed and soilborne decay, mould, wilt, rot and damping-off diseases caused, for example, by *Alternaria* species, caused for example by *Alternaria brassicicola*; *Aphanomyces* species, caused for example by *Aphanomyces euteiches*; *Ascochyta* species, caused for example by *Ascochyta lentis*; *Aspergillus* species, caused for

example by *Aspergillus flavus*; *Cladosporium* species, caused for example by *Cladosporium herbarum*; *Cochliobolus* species, caused for example by *Cochliobolus sativus*; (Conidiaform: *Drechslera*, *Bipolaris* Syn: *Helminthosporium*); *Colletotrichum* species, caused for example by *Colletotrichum coccodes*; *Fusarium* species, caused
5 for example by *Fusarium culmorum*; *Gibberella* species, caused for example by *Gibberella zeae*; *Macrophomina* species, caused for example by *Macrophomina phaseolina*; *Monographella* species, caused for example by *Monographella nivalls*; *Penicillium* species, caused for example by *Penicillium expansum*; *Phoma* species, caused for example by *Phoma lingam*; *Phomopsis* species, caused for example by
10 *Phomopsis sojae*; *Phytophthora* species, caused for example by *Phytophthora cactorum*; *Pyrenophora* species, caused for example by *Pyrenophora graminea*; *Pyricularia* species, caused for example by *Pyricularia oryzae*; *Pythium* species, caused for example by *Pythium ultimum*; *Rhizoctonia* species, caused for example by *Rhizoctonia solani*; *Rhizopus* species, caused for example by *Rhizopus oryzae*;
15 *Sclerotium* species, caused for example by *Sclerotium rolfsii*; *Septoria* species, caused for example by *Septoria nodorum*; *Typhula* species, caused for example by *Typhula incarnata*; *Verticillium* species, caused for example by *Verticillium dahliae*;

cancers, galls and witches' broom caused, for example, by *Nectria* species, for example *Nectria galligena*;

20 wilt diseases caused, for example, by *Monilinia* species, for example *Monilinia laxa*;

leaf blister or leaf curl diseases caused, for example, by *Exobasidium* species, for example *Exobasidium vexans*;

Taphrina species, for example *Taphrina deformans*;

25 decline diseases of wooden plants caused, for example, by Esca disease, caused for example by *Phaemoniella clamydospora*, *Phaeoacremonium aleophilum* and *Fomitiporia mediterranea*; *Eutypa* dieback, caused for example by *Eutypa lata* ; *Ganoderma* diseases caused for example by *Ganoderma boninense*; *Rigidoporus* diseases caused for example by *Rigidoporus lignosus*;

30 diseases of flowers and seeds caused, for example, by *Botrytis* species, for example *Botrytis cinerea*;

diseases of plant tubers caused, for example, by *Rhizoctonia* species, for example *Rhizoctonia solani*; *Helminthosporium* species, for example *Helminthosporium solani*;

Club root caused, for example, by *Plasmodiophora* species, for example *Plasmodiophora brassicae*;

- 5 diseases caused by bacterial pathogens, for example *Xanthomonas* species, for example *Xanthomonas campestris* pv. *oryzae*; *Pseudomonas* species, for example *Pseudomonas syringae* pv. *lachrymans*; *Erwinia* species, for example *Erwinia amylovora*.

The following diseases of soya beans can be controlled with preference:

- 10 Fungal diseases on leaves, stems, pods and seeds caused, for example, by *Alternaria* leaf spot (*Alternaria spec. atrans tenuissima*), Anthracnose (*Colletotrichum gloeosporoides dematium* var. *truncatum*), brown spot (*Septoria glycines*), cercospora leaf spot and blight (*Cercospora kikuchii*), choanephora leaf blight (*Choanephora infundibulifera trispora* (Syn.)), dactulophora leaf spot (*Dactulophora glycines*), downy
15 mildew (*Peronospora manshurica*), drechslera blight (*Drechslera glycini*), frog-eye leaf spot (*Cercospora sojae*), leptosphaerulina leaf spot (*Leptosphaerulina trifolii*), phyllosticta leaf spot (*Phyllosticta sojaecola*), pod and stem blight (*Phomopsis sojae*), powdery mildew (*Microsphaera diffusa*), pyrenochaeta leaf spot (*Pyrenochaeta glycines*), rhizoctonia aerial, foliage, and web blight (*Rhizoctonia solani*), rust
20 (*Phakopsora pachyrhizi*, *Phakopsora meibomia*), scab (*Sphaceloma glycines*), stemphyllium leaf blight (*Stemphyllium botryosum*), target spot (*Corynespora cassicola*).

- Fungal diseases on roots and the stem base caused, for example, by black root rot (*Calonectria crotalariae*), charcoal rot (*Macrophomina phaseolina*), fusarium blight or wilt, root rot, and pod and collar rot (*Fusarium oxysporum*, *Fusarium orthoceras*,
25 *Fusarium semitectum*, *Fusarium equiseti*), mycoleptodiscus root rot (*Mycoleptodiscus terrestris*), neocosmospora (*Neocosmospora vasinfecta*), pod and stem blight (*Diaporthe phaseolorum*), stem canker (*Diaporthe phaseolorum* var. *caulivora*), phytophthora rot (*Phytophthora megasperma*), brown stem rot (*Phialophora gregata*), pythium rot (*Pythium aphanidermatum*, *Pythium irregulare*, *Pythium debaryanum*,
30 *Pythium myriotylum*, *Pythium ultimum*), rhizoctonia root rot, stem decay, and damping-off (*Rhizoctonia solani*), sclerotinia stem decay (*Sclerotinia sclerotiorum*), sclerotinia southern blight (*Sclerotinia rolfsii*), thielaviopsis root rot (*Thielaviopsis basicola*).

The inventive compositions can be used for curative or protective/preventive control of phytopathogenic fungi. The invention therefore also relates to curative and protective methods for controlling phytopathogenic fungi by the use of the inventive composition, which is applied to the seed, the plant or plant parts, the fruit or the soil in which the plants grow.

The fact that the composition is well tolerated by plants at the concentrations required for controlling plant diseases allows the treatment of above-ground parts of plants, of propagation stock and seeds, and of the soil.

According to the invention all plants and plant parts can be treated. By plants is meant all plants and plant populations such as desirable and undesirable wild plants, cultivars and plant varieties (whether or not protectable by plant variety or plant breeder's rights). Cultivars and plant varieties can be plants obtained by conventional propagation and breeding methods which can be assisted or supplemented by one or more biotechnological methods such as by use of double haploids, protoplast fusion, random and directed mutagenesis, molecular or genetic markers or by bioengineering and genetic engineering methods. By plant parts is meant all above ground and below ground parts and organs of plants such as shoot, leaf, blossom and root, whereby for example leaves, needles, stems, branches, blossoms, fruiting bodies, fruits and seed as well as roots, corms and rhizomes are listed. Crops and vegetative and generative propagating material, for example cuttings, corms, rhizomes, runners and seeds also belong to plant parts.

The inventive composition, when it is well tolerated by plants, has favourable homeotherm toxicity and is well tolerated by the environment, is suitable for protecting plants and plant organs, for enhancing harvest yields, for improving the quality of the harvested material. It can preferably be used as crop protection composition. It is active against normally sensitive and resistant species and against all or some stages of development.

Plants which can be treated in accordance with the invention include the following main crop plants: maize, soya bean, alfalfa, cotton, sunflower, *Brassica* oil seeds such as *Brassica napus* (e.g. canola, rapeseed), *Brassica rapa*, *B. juncea* (e.g. (field) mustard) and *Brassica carinata*, *Arecaceae* sp. (e.g. oilpalm, coconut), rice, wheat, sugar beet, sugar cane, oats, rye, barley, millet and sorghum, triticale, flax, nuts, grapes and vine and various fruit and vegetables from various botanic taxa, e.g. *Rosaceae* sp.

(e.g. pome fruits such as apples and pears, but also stone fruits such as apricots, cherries, almonds, plums and peaches, and berry fruits such as strawberries, raspberries, red and black currant and gooseberry), *Ribesioideae sp.*, *Juglandaceae sp.*, *Betulaceae sp.*, *Anacardiaceae sp.*, *Fagaceae sp.*, *Moraceae sp.*, *Oleaceae sp.* (e.g. olive tree), *Actinidaceae sp.*, *Lauraceae sp.* (e.g. avocado, cinnamon, camphor), *Musaceae sp.* (e.g. banana trees and plantations), *Rubiaceae sp.* (e.g. coffee), *Theaceae sp.* (e.g. tea), *Sterculiaceae sp.*, *Rutaceae sp.* (e.g. lemons, oranges, mandarins and grapefruit); *Solanaceae sp.* (e.g. tomatoes, potatoes, peppers, capsicum, aubergines, tobacco), *Liliaceae sp.*, *Compositae sp.* (e.g. lettuce, artichokes and chicory – including root chicory, endive or common chicory), *Umbelliferae sp.* (e.g. carrots, parsley, celery and celeriac), *Cucurbitaceae sp.* (e.g. cucumbers – including gherkins, pumpkins, watermelons, calabashes and melons), *Alliaceae sp.* (e.g. leeks and onions), *Cruciferae sp.* (e.g. white cabbage, red cabbage, broccoli, cauliflower, Brussels sprouts, pak choi, kohlrabi, radishes, horseradish, cress and chinese cabbage), *Leguminosae sp.* (e.g. peanuts, peas, lentils and beans – e.g. common beans and broad beans), *Chenopodiaceae sp.* (e.g. Swiss chard, fodder beet, spinach, beetroot), *Linaceae sp.* (e.g. hemp), *Cannabaceae sp.* (e.g. cannabis), *Malvaceae sp.* (e.g. okra, cocoa), *Papaveraceae* (e.g. poppy), *Asparagaceae* (e.g. asparagus); useful plants and ornamental plants in the garden and woods including turf, lawn, grass and *Stevia rebaudiana*; and in each case genetically modified types of these plants

Preferably, plants which can be treated in accordance with the invention are selected from the group consisting of fruit and vegetables from various botanic taxa, e.g. *Rosaceae sp.* (e.g. pome fruits such as apples and pears, but also stone fruits such as apricots, cherries, almonds, plums and peaches, and berry fruits such as strawberries, raspberries, red and black currant and gooseberry), *Ribesioideae sp.*, *Juglandaceae sp.*, *Betulaceae sp.*, *Anacardiaceae sp.*, *Fagaceae sp.*, *Moraceae sp.*, *Oleaceae sp.* (e.g. olive tree), *Actinidaceae sp.*, *Lauraceae sp.* (e.g. avocado, cinnamon, camphor), *Musaceae sp.* (e.g. banana trees and plantations), *Rubiaceae sp.* (e.g. coffee), *Theaceae sp.* (e.g. tea), *Sterculiaceae sp.*, *Rutaceae sp.* (e.g. lemons, oranges, mandarins and grapefruit); *Solanaceae sp.* (e.g. tomatoes, potatoes, peppers, capsicum, aubergines, tobacco), *Liliaceae sp.*, *Compositae sp.* (e.g. lettuce, artichokes and chicory – including root chicory, endive or common chicory), *Umbelliferae sp.* (e.g. carrots, parsley, celery and celeriac), *Cucurbitaceae sp.* (e.g. cucumbers – including gherkins, pumpkins, watermelons, calabashes and melons),

Alliaceae sp. (e.g. leeks and onions), *Cruciferae* sp. (e.g. white cabbage, red cabbage, broccoli, cauliflower, Brussels sprouts, pak choi, kohlrabi, radishes, horseradish, cress and chinese cabbage), *Leguminosae* sp. (e.g. peanuts, peas, lentils and beans – e.g. common beans and broad beans), *Chenopodiaceae* sp. (e.g. Swiss chard, fodder beet, spinach, beetroot), *Linaceae* sp. (e.g. hemp), *Cannabaceae* sp. (e.g. cannabis), *Malvaceae* sp. (e.g. okra, cocoa), *Papaveraceae* (e.g. poppy), *Asparagaceae* (e.g. asparagus); useful plants and ornamental plants in the garden and woods including turf, lawn, grass and *Stevia rebaudiana*; and in each case genetically modified types of these plants.

10 More preferably, plants which can be treated in accordance with the invention are selected from the group consisting of Chinese cabbage (*Brassica pekinensis*), French beans (*Phaseolus vulgaris*), Rice plants (*Oryza sativa*), and Cotton leaves (*Gossypium hirsutum*).

Depending on the plant species or plant cultivars, their location and growth conditions (soils, climate, vegetation period, diet), using or employing the composition according to the present invention the treatment according to the invention may also result in super-additive ("synergistic") effects. Thus, for example, by using or employing inventive composition in the treatment according to the invention, reduced application rates and/or a widening of the activity spectrum and/or an increase in the activity better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, bigger fruits, larger plant height, greener leaf color, earlier flowering, higher quality and/or a higher nutritional value of the harvested products, higher sugar concentration within the fruits, better storage stability and/or processability of the harvested products are possible, which exceed the effects which were actually to be expected.

At certain application rates of the inventive composition in the treatment according to the invention may also have a strengthening effect in plants. The defense system of the plant against attack by unwanted phytopathogenic fungi and/ or microorganisms and/or viruses is mobilized. Plant-strengthening (resistance-inducing) substances are to be understood as meaning, in the present context, those substances or combinations of substances which are capable of stimulating the defense system of plants in such a way that, when subsequently inoculated with unwanted phytopathogenic fungi and/or microorganisms and/or viruses, the treated plants display a substantial degree of

resistance to these phytopathogenic fungi and/or microorganisms and/or viruses, Thus, by using or employing composition according to the present invention in the treatment according to the invention, plants can be protected against attack by the abovementioned pathogens within a certain period of time after the treatment. The
5 period of time within which protection is effected generally extends from 1 to 10 days, preferably 1 to 7 days, after the treatment of the plants with the active compounds.

Plants and plant cultivars which are also preferably to be treated according to the invention are resistant against one or more biotic stresses, i.e. said plants show a better defense against animal and microbial pests, such as against nematodes, insects,
10 mites, phytopathogenic fungi, bacteria, viruses and/or viroids.

Plants and plant cultivars which may also be treated according to the invention are those plants which are resistant to one or more abiotic stresses, i. e. that already exhibit an increased plant health with respect to stress tolerance. Abiotic stress conditions may include, for example, drought, cold temperature exposure, heat exposure, osmotic
15 stress, flooding, increased soil salinity, increased mineral exposure, ozon exposure, high light exposure, limited availability of nitrogen nutrients, limited availability of phosphorus nutrients, shade avoidance. Preferably, the treatment of these plants and cultivars with the composition of the present invention additionally increases the overall plant health (cf. above).

20 Plants and plant cultivars which may also be treated according to the invention, are those plants characterized by enhanced yield characteristics i. e. that already exhibit an increased plant health with respect to this feature. Increased yield in said plants can be the result of, for example, improved plant physiology, growth and development, such as water use efficiency, water retention efficiency, improved nitrogen use,
25 enhanced carbon assimilation, improved photosynthesis, increased germination efficiency and accelerated maturation. Yield can furthermore be affected by improved plant architecture (under stress and non-stress conditions), including but not limited to, early flowering, flowering control for hybrid seed production, seedling vigor, plant size, internode number and distance, root growth, seed size, fruit size, pod size, pod or ear
30 number, seed number per pod or ear, seed mass, enhanced seed filling, reduced seed dispersal, reduced pod dehiscence and lodging resistance. Further yield traits include seed composition, such as carbohydrate content, protein content, oil content and composition, nutritional value, reduction in anti-nutritional compounds, improved processability and better storage stability. Preferably, the treatment of these plants and

cultivars with the composition of the present invention additionally increases the overall plant health (cf. above).

Plants that may be treated according to the invention are hybrid plants that already express the characteristic of heterosis or hybrid vigor which results in generally higher yield, vigor, health and resistance towards biotic and abiotic stress factors. Such plants are typically made by crossing an inbred male-sterile parent line (the female parent) with another inbred male-fertile parent line (the male parent). Hybrid seed is typically harvested from the male sterile plants and sold to growers. Male sterile plants can sometimes (e.g. in corn) be produced by detasseling, i.e. the mechanical removal of the male reproductive organs (or males flowers) but, more typically, male sterility is the result of genetic determinants in the plant genome. In that case, and especially when seed is the desired product to be harvested from the hybrid plants it is typically useful to ensure that male fertility in the hybrid plants is fully restored. This can be accomplished by ensuring that the male parents have appropriate fertility restorer genes which are capable of restoring the male fertility in hybrid plants that contain the genetic determinants responsible for male-sterility. Genetic determinants for male sterility may be located in the cytoplasm. Examples of cytoplasmic male sterility (CMS) were for instance described in Brassica species. However, genetic determinants for male sterility can also be located in the nuclear genome. Male sterile plants can also be obtained by plant biotechnology methods such as genetic engineering. A particularly useful means of obtaining male-sterile plants is described in WO 89/10396 in which, for example, a ribonuclease such as barnase is selectively expressed in the tapetum cells in the stamens. Fertility can then be restored by expression in the tapetum cells of a ribonuclease inhibitor such as barstar.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may be treated according to the invention are herbicide-tolerant plants, i.e. plants made tolerant to one or more given herbicides. Such plants can be obtained either by genetic transformation, or by selection of plants containing a mutation imparting such herbicide tolerance.

Herbicide-tolerant plants are for example glyphosate-tolerant plants, i.e. plants made tolerant to the herbicide glyphosate or salts thereof. Plants can be made tolerant to glyphosate through different means. For example, glyphosate-tolerant plants can be obtained by transforming the plant with a gene encoding the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS). Examples of such EPSPS genes are

the *AroA* gene (mutant CT7) of the bacterium *Salmonella typhimurium*, the CP4 gene of the bacterium *Agrobacterium sp*, the genes encoding a Petunia EPSPS, a Tomato EPSPS, or an Eleusine EPSPS. It can also be a mutated EPSPS. Glyphosate-tolerant plants can also be obtained by expressing a gene that encodes a glyphosate oxidoreductase enzyme. Glyphosate-tolerant plants can also be obtained by expressing a gene that encodes a glyphosate acetyl transferase enzyme. Glyphosate-tolerant plants can also be obtained by selecting plants containing naturally-occurring mutations of the above-mentioned genes.

Other herbicide resistant plants are for example plants that are made tolerant to herbicides inhibiting the enzyme glutamine synthase, such as bialaphos, phosphinothricin or glufosinate. Such plants can be obtained by expressing an enzyme detoxifying the herbicide or a mutant glutamine synthase enzyme that is resistant to inhibition. One such efficient detoxifying enzyme is an enzyme encoding a phosphinothricin acetyltransferase (such as the *bar* or *pat* protein from *Streptomyces* species). Plants expressing an exogenous phosphinothricin acetyltransferase are also described.

Further herbicide-tolerant plants are also plants that are made tolerant to the herbicides inhibiting the enzyme hydroxyphenylpyruvatedioxygenase (HPPD). Hydroxyphenylpyruvatedioxygenases are enzymes that catalyze the reaction in which para-hydroxyphenylpyruvate (HPP) is transformed into homogentisate. Plants tolerant to HPPD-inhibitors can be transformed with a gene encoding a naturally-occurring resistant HPPD enzyme, or a gene encoding a mutated HPPD enzyme. Tolerance to HPPD-inhibitors can also be obtained by transforming plants with genes encoding certain enzymes enabling the formation of homogentisate despite the inhibition of the native HPPD enzyme by the HPPD-inhibitor. Tolerance of plants to HPPD inhibitors can also be improved by transforming plants with a gene encoding an enzyme prephenate dehydrogenase in addition to a gene encoding an HPPD-tolerant enzyme.

Still further herbicide resistant plants are plants that are made tolerant to acetolactate synthase (ALS) inhibitors. Known ALS-inhibitors include, for example, sulfonylurea, imidazolinone, triazolopyrimidines, pyrimidinyoxy(thio)benzoates, and/or sulfonylaminocarbonyltriazolinone herbicides. Different mutations in the ALS enzyme (also known as acetohydroxyacid synthase, AHAS) are known to confer tolerance to different herbicides and groups of herbicides. The production of sulfonylurea-tolerant plants and imidazolinone-tolerant plants is described in WO 1996/033270. Other

imidazolinone-tolerant plants are also described. Further sulfonylurea- and imidazolinone-tolerant plants are also described in for example WO 2007/024782.

Other plants tolerant to imidazolinone and/or sulfonylurea can be obtained by induced mutagenesis, selection in cell cultures in the presence of the herbicide or mutation breeding as described for example for soybeans, for rice, for sugar beet, for lettuce, or for sunflower.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are insect-resistant transgenic plants, i.e. plants made resistant to attack by certain target insects. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such insect resistance.

An "insect-resistant transgenic plant", as used herein, includes any plant containing at least one transgene comprising a coding sequence encoding:

- 1) An insecticidal crystal protein from *Bacillus thuringiensis* or an insecticidal portion thereof, such as the insecticidal crystal proteins listed online at: http://www.lifesci.sussex.ac.uk/Home/Neil_Crickmore/Bt/, or insecticidal portions thereof, e.g., proteins of the Cry protein classes Cry1Ab, Cry1Ac, Cry1F, Cry2Ab, Cry3Aa, or Cry3Bb or insecticidal portions thereof; or
- 2) a crystal protein from *Bacillus thuringiensis* or a portion thereof which is insecticidal in the presence of a second other crystal protein from *Bacillus thuringiensis* or a portion thereof, such as the binary toxin made up of the Cry34 and Cry35 crystal proteins; or
- 3) a hybrid insecticidal protein comprising parts of different insecticidal crystal proteins from *Bacillus thuringiensis*, such as a hybrid of the proteins of 1) above or a hybrid of the proteins of 2) above, e.g., the Cry1A.105 protein produced by corn event MON98034 (WO 2007/027777); or
- 4) a protein of any one of 1) to 3) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target insect species affected, and/or because of changes introduced into the encoding DNA during cloning or transformation, such as the Cry3Bb1 protein in corn events MON863 or MON88017, or the Cry3A protein in corn event MIR604;

5) an insecticidal secreted protein from *Bacillus thuringiensis* or *Bacillus cereus*, or an insecticidal portion thereof, such as the vegetative insecticidal (VIP) proteins listed at:

http://www.lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/vip.html, e.g. proteins from the VIP3Aa protein class; or

6) secreted protein from *Bacillus thuringiensis* or *Bacillus cereus* which is insecticidal in the presence of a second secreted protein from *Bacillus thuringiensis* or *B. cereus*, such as the binary toxin made up of the VIP1A and VIP2A proteins; or

7) hybrid insecticidal protein comprising parts from different secreted proteins from *Bacillus thuringiensis* or *Bacillus cereus*, such as a hybrid of the proteins in 1) above or a hybrid of the proteins in 2) above; or

8) protein of any one of 1) to 3) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target insect species affected, and/or because of changes introduced into the encoding DNA during cloning or transformation (while still encoding an insecticidal protein), such as the VIP3Aa protein in cotton event COT102.

Of course, an insect-resistant transgenic plant, as used herein, also includes any plant comprising a combination of genes encoding the proteins of any one of the above classes 1 to 8. In one embodiment, an insect-resistant plant contains more than one transgene encoding a protein of any one of the above classes 1 to 8, to expand the range of target insect species affected when using different proteins directed at different target insect species, or to delay insect resistance development to the plants by using different proteins insecticidal to the same target insect species but having a different mode of action, such as binding to different receptor binding sites in the insect.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are tolerant to abiotic stresses. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such stress resistance. Particularly useful stress tolerance plants include:

a. plants which contain a transgene capable of reducing the expression and/or the activity of poly(ADP-ribose)polymerase (PARP) gene in the plant cells or plants

b. plants which contain a stress tolerance enhancing transgene capable of reducing the expression and/or the activity of the poly(ADP-ribose)glycohydrolase (PARG) encoding genes of the plants or plants cells.

5 c. plants which contain a stress tolerance enhancing transgene coding for a plant-functional enzyme of the nicotinamide adenine dinucleotide salvage synthesis pathway including nicotinamidase, nicotinate phosphoribosyltransferase, nicotinic acid mononucleotide adenylyl transferase, nicotinamide adenine dinucleotide synthetase or nicotine amide phosphoribosyltransferase.

10 Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention show altered quantity, quality and/or storage-stability of the harvested product and/or altered properties of specific ingredients of the harvested product such as :

15 1) transgenic plants which synthesize a modified starch, which in its physical-chemical characteristics, in particular the amylose content or the amylose/amylopectin ratio, the degree of branching, the average chain length, the side chain distribution, the viscosity behaviour, the gelling strength, the starch grain size and/or the starch grain morphology, is changed in comparison with the synthesised starch in wild type plant cells or plants, so that this is better suited for special applications.

20 2) transgenic plants which synthesize non starch carbohydrate polymers or which synthesize non starch carbohydrate polymers with altered properties in comparison to wild type plants without genetic modification. Examples are plants producing polyfructose, especially of the inulin and levan-type, plants producing alpha 1,4 glucans, plants producing alpha-1,6 branched alpha-1,4-glucans, plants producing alternan,

25 3) transgenic plants which produce hyaluronan.

Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are plants, such as cotton plants, with altered fiber characteristics. Such plants can be obtained by genetic transformation or by selection of plants contain a mutation imparting such altered fiber characteristics and include:

30 a) Plants, such as cotton plants, containing an altered form of cellulose synthase genes,

- b) Plants, such as cotton plants, containing an altered form of rsw2 or rsw3 homologous nucleic acids,
- c) Plants, such as cotton plants, with increased expression of sucrose phosphate synthase,
- 5 d) Plants, such as cotton plants, with increased expression of sucrose synthase,
- e) Plants, such as cotton plants, wherein the timing of the plasmodesmatal gating at the basis of the fiber cell is altered, e.g. through downregulation of fiberselective β 1,3-glucanase,
- 10 f) Plants, such as cotton plants, having fibers with altered reactivity, e.g. through the expression of N-acetylglucosaminetransferase gene including nodC and chitinsynthase genes.

Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are plants, such as oilseed rape or related Brassica plants, with altered oil profile characteristics.

- 15 Such plants can be obtained by genetic transformation or by selection of plants contain a mutation imparting such altered oil characteristics and include:

- a) Plants, such as oilseed rape plants, producing oil having a high oleic acid content,
- b) Plants such as oilseed rape plants, producing oil having a low linolenic acid content,
- 20 c) Plant such as oilseed rape plants, producing oil having a low level of saturated fatty acids.

Particularly useful transgenic plants which may be treated according to the invention are plants which comprise one or more genes which encode one or more toxins, such as the following which are sold under the trade names YIELD GARD[®] (for example maize, cotton, soya beans), KnockOut[®] (for example maize), BiteGard[®] (for example maize), Bt-Xtra[®] (for example maize), StarLink[®] (for example maize), Bollgard[®] (cotton), NucoIn[®] (cotton), NucoIn 33B[®] (cotton), NatureGard[®] (for example maize), Protecta[®] and NewLeaf[®] (potato). Examples of herbicide-tolerant plants which may be mentioned

25 are maize varieties, cotton varieties and soya bean varieties which are sold under the trade names Roundup Ready[®] (tolerance to glyphosate, for example maize, cotton, soya bean), Liberty Link[®] (tolerance to phosphinotricin, for example oilseed rape), IMI[®] (tolerance to imidazolinones) and STS[®] (tolerance to sulphonylureas, for example maize). Herbicide-resistant plants (plants bred in a conventional manner for herbicide

30

tolerance) which may be mentioned include the varieties sold under the name Clearfield® (for example maize).

Particularly useful transgenic plants which may be treated according to the invention are plants containing transformation events, or a combination of transformation events,
5 and that are listed for example in the databases for various national or regional regulatory agencies including Event 1143-14A (cotton, insect control, not deposited, described in WO 06/128569); Event 1143-51B (cotton, insect control, not deposited, described in WO 06/128570); Event 1445 (cotton, herbicide tolerance, not deposited, described in US-A 2002-120964 or WO 02/034946); Event 17053 (rice, herbicide
10 tolerance, deposited as PTA-9843, described in WO 10/117737); Event 17314 (rice, herbicide tolerance, deposited as PTA-9844, described in WO 10/117735); Event 281-24-236 (cotton, insect control - herbicide tolerance, deposited as PTA-6233, described in WO 05/103266 or US-A 2005-216969); Event 3006-210-23 (cotton, insect control - herbicide tolerance, deposited as PTA-6233, described in US-A 2007-143876 or WO
15 05/103266); Event 3272 (corn, quality trait, deposited as PTA-9972, described in WO 06/098952 or US-A 2006-230473); Event 40416 (corn, insect control - herbicide tolerance, deposited as ATCC PTA-11508, described in WO 11/075593); Event 43A47 (corn, insect control - herbicide tolerance, deposited as ATCC PTA-11509, described in WO 11/075595); Event 5307 (corn, insect control, deposited as ATCC PTA-9561,
20 described in WO 10/077816); Event ASR-368 (bent grass, herbicide tolerance, deposited as ATCC PTA-4816, described in US-A 2006-162007 or WO 04/053062); Event B16 (corn, herbicide tolerance, not deposited, described in US-A 2003-126634); Event BPS-CV127-9 (soybean, herbicide tolerance, deposited as NCIMB No. 41603, described in WO 10/080829); Event CE43-67B (cotton, insect control, deposited as DSM ACC2724,
25 described in US-A 2009-217423 or WO 06/128573); Event CE44-69D (cotton, insect control, not deposited, described in US-A 2010-0024077); Event CE44-69D (cotton, insect control, not deposited, described in WO 06/128571); Event CE46-02A (cotton, insect control, not deposited, described in WO 06/128572); Event COT102 (cotton, insect control, not deposited, described in US-A 2006-130175 or WO 04/039986); Event
30 COT202 (cotton, insect control, not deposited, described in US-A 2007-067868 or WO 05/054479); Event COT203 (cotton, insect control, not deposited, described in WO 05/054480); Event DAS40278 (corn, herbicide tolerance, deposited as ATCC PTA-10244, described in WO 11/022469); Event DAS-59122-7 (corn, insect control - herbicide tolerance, deposited as ATCC PTA 11384 , described in US-A 2006-070139); Event DAS-
35 59132 (corn, insect control - herbicide tolerance, not deposited, described in WO

09/100188); Event DAS68416 (soybean, herbicide tolerance, deposited as ATCC PTA-10442, described in WO 11/066384 or WO 11/066360); Event DP-098140-6 (corn, herbicide tolerance, deposited as ATCC PTA-8296, described in US-A 2009-137395 or WO 08/112019); Event DP-305423-1 (soybean, quality trait, not deposited, described in 5 US-A 2008-312082 or WO 08/054747); Event DP-32138-1 (corn, hybridization system, deposited as ATCC PTA-9158, described in US-A 2009-0210970 or WO 09/103049); Event DP-356043-5 (soybean, herbicide tolerance, deposited as ATCC PTA-8287, described in US-A 2010-0184079 or WO 08/002872); Event EE-1 (brinjal, insect control, not deposited, described in WO 07/091277); Event FI117 (corn, herbicide tolerance, 10 deposited as ATCC 209031, described in US-A 2006-059581 or WO 98/044140); Event GA21 (corn, herbicide tolerance, deposited as ATCC 209033, described in US-A 2005-086719 or WO 98/044140); Event GG25 (corn, herbicide tolerance, deposited as ATCC 209032, described in US-A 2005-188434 or WO 98/044140); Event GHB119 (cotton, insect control - herbicide tolerance, deposited as ATCC PTA-8398, described in WO 15 08/151780); Event GHB614 (cotton, herbicide tolerance, deposited as ATCC PTA-6878, described in US-A 2010-050282 or WO 07/017186); Event GJ11 (corn, herbicide tolerance, deposited as ATCC 209030, described in US-A 2005-188434 or WO 98/044140); Event GM RZ13 (sugar beet, virus resistance , deposited as NCIMB-41601, described in WO 10/076212); Event H7-1 (sugar beet, herbicide tolerance, deposited as 20 NCIMB 41158 or NCIMB 41159, described in US-A 2004-172669 or WO 04/074492); Event JOPLIN1 (wheat, disease tolerance, not deposited, described in US-A 2008-064032); Event LL27 (soybean, herbicide tolerance, deposited as NCIMB41658, described in WO 06/108674 or US-A 2008-320616); Event LL55 (soybean, herbicide tolerance, deposited as NCIMB 41660, described in WO 06/108675 or US-A 2008- 25 196127); Event LLCotton25 (cotton, herbicide tolerance, deposited as ATCC PTA-3343, described in WO 03/013224 or US-A 2003-097687); Event LLRICE06 (rice, herbicide tolerance, deposited as ATCC-23352, described in US 6,468,747 or WO 00/026345); Event LLRICE601 (rice, herbicide tolerance, deposited as ATCC PTA-2600, described in US-A 2008-2289060 or WO 00/026356); Event LY038 (corn, quality trait, deposited as 30 ATCC PTA-5623, described in US-A 2007-028322 or WO 05/061720); Event MIR162 (corn, insect control, deposited as PTA-8166, described in US-A 2009-300784 or WO 07/142840); Event MIR604 (corn, insect control, not deposited, described in US-A 2008-167456 or WO 05/103301); Event MON15985 (cotton, insect control, deposited as ATCC PTA-2516, described in US-A 2004-250317 or WO 02/100163); Event MON810 (corn, 35 insect control, not deposited, described in US-A 2002-102582); Event MON863 (corn,

insect control, deposited as ATCC PTA-2605, described in WO 04/011601 or US-A 2006-095986); Event MON87427 (corn, pollination control, deposited as ATCC PTA-7899, described in WO 11/062904); Event MON87460 (corn, stress tolerance, deposited as ATCC PTA-8910, described in WO 09/111263 or US-A 2011-0138504); Event MON87701
5 (soybean, insect control, deposited as ATCC PTA-8194, described in US-A 2009-130071 or WO 09/064652); Event MON87705 (soybean, quality trait - herbicide tolerance, deposited as ATCC PTA-9241, described in US-A 2010-0080887 or WO 10/037016); Event MON87708 (soybean, herbicide tolerance, deposited as ATCC PTA9670, described in WO 11/034704); Event MON87754 (soybean, quality trait, deposited as
10 ATCC PTA-9385, described in WO 10/024976); Event MON87769 (soybean, quality trait, deposited as ATCC PTA-8911, described in US-A 2011-0067141 or WO 09/102873); Event MON88017 (corn, insect control - herbicide tolerance, deposited as ATCC PTA-5582, described in US-A 2008-028482 or WO 05/059103); Event MON88913 (cotton, herbicide tolerance, deposited as ATCC PTA-4854, described in WO 04/072235 or US-A
15 2006-059590); Event MON89034 (corn, insect control, deposited as ATCC PTA-7455, described in WO 07/140256 or US-A 2008-260932); Event MON89788 (soybean, herbicide tolerance, deposited as ATCC PTA-6708, described in US-A 2006-282915 or WO 06/130436); Event MS11 (oilseed rape, pollination control - herbicide tolerance, deposited as ATCC PTA-850 or PTA-2485, described in WO 01/031042); Event MS8
20 (oilseed rape, pollination control - herbicide tolerance, deposited as ATCC PTA-730, described in WO 01/041558 or US-A 2003-188347); Event NK603 (corn, herbicide tolerance, deposited as ATCC PTA-2478, described in US-A 2007-292854); Event PE-7 (rice, insect control, not deposited, described in WO 08/114282); Event RF3 (oilseed rape, pollination control - herbicide tolerance, deposited as ATCC PTA-730, described in
25 WO 01/041558 or US-A 2003-188347); Event RT73 (oilseed rape, herbicide tolerance, not deposited, described in WO 02/036831 or US-A 2008-070260); Event T227-1 (sugar beet, herbicide tolerance, not deposited, described in WO 02/44407 or US-A 2009-265817); Event T25 (corn, herbicide tolerance, not deposited, described in US-A 2001-029014 or WO 01/051654); Event T304-40 (cotton, insect control - herbicide tolerance, deposited as ATCC PTA-8171, described in US-A 2010-077501 or WO 08/122406); Event
30 T342-142 (cotton, insect control, not deposited, described in WO 06/128568); Event TC1507 (corn, insect control - herbicide tolerance, not deposited, described in US-A 2005-039226 or WO 04/099447); Event VIP1034 (corn, insect control - herbicide tolerance, deposited as ATCC PTA-3925., described in WO 03/052073), Event 32316
35 (corn, insect control-herbicide tolerance, deposited as PTA-11507, described in WO

11/084632), Event 4114 (corn, insect control-herbicide tolerance, deposited as PTA-11506, described in WO 11/084621).

Particularly useful transgenic plants which may be treated according to the invention are plants containing transformation events, or combination of transformation events, that are listed for example in the databases from various national or regional regulatory agencies (see for example http://gmoinfo.jrc.it/gmp_browse.aspx and <http://www.agbios.com/dbase.php>).

Examples

For all examples the efficiencies of the compositions comprising at least one BCA and at least one specific insecticide has been determined by the "Colby-formula": The expected efficacy of a given combination of two compounds is calculated as follows (see Colby, S.R., „Calculating Synergistic and antagonistic Responses of Herbicide Combinations“, Weeds 15, pp. 20-22, 1967):

If

X is the efficacy expressed in % mortality of the untreated control for test compound A at a concentration of m ppm respectively m g/ha,

Y is the efficacy expressed in % mortality of the untreated control for test compound B at a concentration of n ppm respectively n g/ha,

E is the efficacy expressed in % mortality of the untreated control using the mixture of A and B at m and n ppm respectively m and n g /ha,

then is $E = X + Y - (X \times Y / 100)$

If the observed insecticidal efficacy of the combination is higher than the one calculated as „E“, then the combination of the two compounds is more than additive, i.e., there is a synergistic effect.

25

In the following the following compounds / abbreviation for compounds are used:

Serenade Max™ is a commercial product which contains the strain *Bacillus subtilis* QST 713 (also referred to as B9 according to the present invention).

Sonata™ is a commercial product which contains the strain *Bacillus pumilus* QST 2808 also referred to as B3 according to the present invention)

30

The strain *Bacillus subtilis* AQ30002 which is mentioned above as B19, is referred to in the following table as QST3002. A solution comprising $8.5 \cdot 10^8$ CFU/g (1.34 %) of this strain has been used.

All ratios given below refer to biological control agent/spore preparations of the respective biological control agents of around 10^{10} cells or spores per gram preparation of said biological control agent (see definition for ratios above).

Example A

Myzus persicae – spray test (MYZUPE)

Solvent: 78.0 parts by weight acetone

10 1.5 parts by weight dimethylformamide

Emulsifier: 0.5 parts by weight alkylaryl/polyglycoether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration. To produce a suitable preparation of a spore suspension the spores are diluted with emulsifier-containing water to the desired concentration.

Chinese cabbage (*Brassica pekinensis*) leaf– disks infected with all instars of the green peach aphid (*Myzus persicae*), are sprayed with a preparation of the active ingredient of the desired concentration.

20 After the specified period of time, mortality in % is determined. 100 % means all aphids have been killed; 0 % means none of the aphids have been killed. The mortality values determined thus are recalculated using the Colby-formula (see above).

According to the present application in this test e.g. the following combinations show a synergistic effect in comparison to the single compounds:

25

Table A: *Myzus persicae* – test

<u>Active Ingredient</u>	<u>Concentration in g/ha</u>	<u>Efficacy in % after 6d</u>
Sonata™ (B3)	100 50	0 0
<i>Streptomyces galbus</i> AQ 6047 (B16)	3000 2000	17.5 0
Fipronil (I95)	100 4	0 0
Sonata™ + Fipronil (1 : 1) according to the invention	100 + 100	obs.* cal.** 90 0
<i>Streptomyces galbus</i> + Fipronil (500 : 1) according to the invention	2000 + 4	obs.* cal.** 70 0
Methiocarb (I15)	500	0
<i>Streptomyces galbus</i> + Methiocarb (4 : 1) according to the invention	2000 + 500	obs.* cal.** 90 0

*obs. = observed insecticidal efficacy, ** cal. = efficacy calculated with Colby-formula

Example B

Tetranychus urticae – spray test, OP-resistant

- 5 Solvent: 78.0 parts by weight acetone
1.5 parts by weight dimethylformamide
Emulsifier: 0.5 parts by weight alkylaryl polyglycoether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration. To produce a suitable preparation of a spore suspension the spores are diluted with emulsifier-containing water to the desired concentration.

French beans (*Phaseolus vulgaris*) which are heavily infested with all stages of the two spotted spider mite (*Tetranychus urticae*), are sprayed with a preparation of the active ingredient of the desired concentration.

After the specified period of time, mortality in % is determined. 100% means all spider mites have been killed and 0% means none of the spider mites have been killed. The mortality values determined thus are recalculated using the Colby-formula (see above).

According to the present application in this test e.g. the following combinations show a synergistic effect in comparison to the single compounds:

Table B: Tetranychus urticae – test

<u>Active Ingredient</u>	<u>Concentration in g ai/ha</u>	<u>Efficacy in % after 2d</u>
Serenade Max™ (B9)	200	0
Sonata™ (B3)	50	0
Abamectin (I152)	0.032	0
Serenade Max™ + Abamectin (6250 : 1) according to the invention	200 + 0.032	obs.* cal.** 70 0
Sonata™ + Abamectin (1562.5 : 1) according to the invention	50 + 0.032	obs.* cal.** 70 0

*obs. = observed insecticidal efficacy, ** cal. = efficacy calculated with Colby-formula

Example C

Phaedon cochleariae – spray test

- 10 Solvent: 78.0 parts by weight of acetone
 1.5 parts by weight of dimethylformamide
Emulsifier: 0.5 parts by weight of alkylaryl polyglycoether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and emulsifier, and the concentrate is diluted with emulsifier- containing water to the desired concentration. To produce a suitable preparation of a spore suspension the spores are diluted with emulsifier- containing water to the desired concentration.

Chinese cabbage (*Brassica pekinensis*) leaf- disks are sprayed with a preparation of the active ingredient of the desired concentration. Once dry, the leaf disks are infested with mustard beetle larvae (*Phaedon cochleariae*).

After the specified period of time, mortality in % is determined. 100 % means all beetle larvae have been killed and 0 % means none of the beetle larvae have been killed. The mortality values determined thus are recalculated using the Colby-formula (see above).

- 5 According to the present application in this test e.g. the following combinations show a synergistic effect in comparison to the single compounds:

Table C: Phaedon cochleariae – test

<u>Active Ingredient</u>	<u>Concentration in g al/ha</u>	<u>Efficacy in % after 6d</u>
Serenade Max™ (B9)	400 200	0 0
Sonata™ (B3)	100 50	0 0
QST30002 (B19)	200 100	0 0
Methiocarb (I15)	20	0
Serenade Max™ + Methiocarb (20 : 1) according to the invention	400 + 20	obs.* cal.** 33 0
Sonata™ + Methiocarb (5 : 1) according to the invention	100 + 20	obs.* cal.** 33 0
Abamectin (I152)	0.8	0
QST30002 + Abamectin (125 : 1) according to the invention	100 + 0.8	obs.* cal.** 33 0
Flipronil (I95)	0.25	33
QST30002 + Flipronil (800 : 1) according to the invention	200 + 0.25	obs.* cal.** 67 33

*obs. = observed insecticidal efficacy, ** cal. = efficacy calculated with Colby-formula

Example D

10 Nilaparvata lugens – spray test (NILALU)

Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 2 parts by weight of alkylaryl polyglycoether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and emulsifier, and the

- 15 concentrate is diluted with emulsifier-containing water to the desired concentration. To

produce a suitable preparation of a spore suspension the spores are diluted with emulsifier- containing water to the desired concentration. Ammonium salt and/or penetration enhancer in a dosage of 1000 ppm are added to the desired concentration if necessary.

- 5 Rice plants (*Oryza sativa*) are treated by being sprayed with the desired concentration of the active compound and are infested with the brown planthopper (*Nilaparvata lugens*).

After the specified period of time, mortality in % is determined. 100 % means all planthoppers have been killed and 0 % means none of the planthoppers have been
 10 killed.

According to the present application in this test e.g. the following combinations show a synergistic effect in comparison to the single compounds:

Table D: Nilaparvata lugens – test

<u>Active ingredient</u>	<u>Concentration in ppm</u>	<u>Efficacy in % after 3d</u>
Serenade Max™ (B9)	200	10
Fipronil (195)	1	0
Serenade Max™ + Fipronil (200 : 1) according to the invention	200 + 1	obs.* cal.** 40 10

*obs. = observed insecticidal efficacy, ** cal. = efficacy calculated with Colby-formula

15 **Example E**

Spodoptera frugiperda – spray test

- Solvent: 7 parts by weight of dimethylformamide
 Emulsifier: 2 parts by weight of alkylaryl polyglycoether

To produce a suitable preparation of active compound, 1 part by weight of active
 20 compound is mixed with the stated amount of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration. To produce a suitable preparation of a spore suspension the spores are diluted with emulsifier- containing water to the desired concentration. Ammonium salt and/or

penetration enhancer in a dosage of 1000 ppm are added to the desired concentration if necessary.

Cotton leaves (*Gossypium hirsutum*) are treated by being sprayed with the preparation of the active compound of the desired concentration and are infested with caterpillars of the fall army worm (*Spodoptera frugiperda*).

After the specified period of time, mortality in % is determined. 100 % means all the caterpillars have been killed; 0 % means none of the caterpillars have been killed. The mortality values determined thus are recalculated using the Colby-formula (see above).

According to the present application in this test e.g. the following combinations show a synergistic effect in comparison to the single compounds:

Table E: *Spodoptera frugiperda* – spray test

<u>Active Ingredient</u>	<u>Concentration In ppm</u>	<u>Efficacy In % after 1d</u>
Serenade Max™ (B9)	200	0
Sonata™ (B3)	50	0
Thiodicarb (I21)	0.8	50
Serenade Max™ + Thiodicarb (250 : 1) according to the invention	200 + 0.8	obs.* cal.** 80 50
Sonata™ + Thiodicarb (62.5 : 1) according to the invention	50 + 0.8	obs.* cal.** 80 50

*obs. = observed insecticidal efficacy, ** cal. = efficacy calculated with Colby-formula

Claims

1. A composition comprising at least one biological control agent selected from the group consisting of

5 *Bacillus chitinosporus* AQ746 (NRRL Accession No. B-21618), *Bacillus mycolides* AQ726 (NRRL Accession No. B-21664), *Bacillus pumilus* (NRRL Accession No. B-30087), *Bacillus pumilus* AQ717 (NRRL Accession No. B-21662), *Bacillus* sp. AQ175 (ATCC Accession No. 55608), *Bacillus* sp. AQ177 (ATCC Accession No. 55609), *Bacillus* sp. AQ178 (ATCC Accession No. 53522), *Bacillus subtilis* AQ743 (NRRL Accession No. B-21665), *Bacillus subtilis* AQ713 (NRRL Accession No. B-21661), *Bacillus subtilis* AQ153 (ATCC Accession
10 No. 55614), *Bacillus thuringiensis* BD#32 (NRRL Accession No. B-21530), *Bacillus thuringiensis* AQ52 (NRRL Accession No. B-21619), *Muscodora albus* 620 (NRRL Accession No. 30547), *Muscodora roseus* A3-5 (NRRL Accession No. 30548), *Rhodococcus globerulus* AQ719 (NRRL Accession No. B-21663), *Streptomyces galbus* (NRRL Accession No. 30232), *Streptomyces* sp. (NRRL Accession No. B-30145), *Bacillus thuringiensis*
15 *subspec. kurstaki* BMP 123, *Bacillus subtilis* AQ30002 (NRRL Accession No. B-50421), and *Bacillus subtilis* AQ 30004 (NRRL Accession No. B-50455),

and/or a mutant of these strains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens

20 and at least one insecticide selected from the group consisting of acetylcholinesterase (AChE) inhibitors, GABA-gated chloride channel antagonists, chloride channel activators and nicotinic acetylcholine receptor (nAChR) channel blockers,

in a synergistically effective amount.

2. The composition according to claim 1, wherein

25 the acetylcholinesterase (AChE) inhibitor is selected from the group consisting of carbamates, and organophosphates, and/or

the GABA-gated chloride channel antagonist is selected from the group consisting of cyclodiene organochlorines, and phenylpyrazoles (fiproles), and/or

the chloride channel activator is an avermectin/milbemycin.

3. The composition according to claim 2, wherein the acetylcholinesterase (AChE) inhibitor is selected from the group consisting of Alanycarb, Aldicarb, Bendiocarb, Benfuracarb, Butocarboxim, Butoxycarboxim, Carbaryl, Carbofuran, Carbosulfan, Ethiofencarb, Fenobucarb, Formetanate, Furathiocarb, Isoprocarb, Methiocarb, 5 Methomyl, Metolcarb, Oxamyl, Pirimicarb, Propoxur, Thiodicarb, Thiofanox, Triazamate, Trimethacarb, XMC, Xyllycarb, Acephate, Azamethiphos, Azinphos-ethyl, Azinphos-methyl, Cadusafos, Chlorethoxyfos, Chlorfenvinphos, Chlormephos, Chlorpyrifos, Chlorpyrifos-methyl, Coumaphos, Cyanophos, Demeton-S-methyl, Diazinon, Dichlorvos/DDVP, Dicrotophos, Dimethoate, Dimethylvinphos, Disulfoton, EPN, Ethion, 10 Ethoprophos, Famphur, Fenamiphos, Fenitrothion, Fenthion, Fosthiazate, Heptenophos, Imicyafos, Isofenphos, Isopropyl O-(methoxyaminothio-phosphoryl) salicylate, Isoxathion, Malathion, Mecarbam, Methamidophos, Methidathion, Mevinphos, Monocrotophos, Naled, Omethoate, Oxydemeton-methyl, Parathion, Parathion-methyl, Phenthoate, Phorate, Phosalone, Phosmet, Phosphamidon, Phoxim, Pirimiphos-methyl, Profenofos, 15 Propetamphos, Prothiofos, Pyraclofos, Pyridaphenthion, Quinalphos, Sulfotep, Tebupirimfos, Temephos, Terbufos, Tetrachlorvinphos, Thiometon, Triazophos, Trichlorfon, and Vamidothion, and/or

the GABA-gated chloride channel antagonist is selected from the group consisting of Chlordane, Endosulfan, Ethiprole, and Fipronil, and/or

20 the chloride channel activator is selected from the group consisting of Abamectin, Enamectin benzoate, Lepimectin, and Milbemectin, and/or

the nicotinic acetylcholine receptor (nAChR) channel blocker is selected from the group consisting of Bensultap, Cartap hydrochloride, Thiocyclam, and Thiosultap-sodium.

4. The composition according to claim 3, wherein the acetylcholinesterase (AChE) 25 inhibitor is selected from the group consisting of Acephate, Chlorpyrifos, Carbofuran, Methiocarb, Profenofos, and Thiodicarb, and/or

the GABA-gated chloride channel antagonist is selected from the group consisting of Ethiprole, and Fipronil, and/or

30 the chloride channel activator is selected from the group consisting Abamectin, Enamectin-benzoate, and Milbemectin.

5. The composition according to any one of claims 1 to 4, further comprising at least one fungicide, with the proviso that the biological control agent and the fungicide are not identical.

6. The composition according to any one of claims 1 to 5, wherein the fungicide is
 5 selected from the group consisting of inhibitors of the ergosterol biosynthesis, inhibitors of the respiratory chain at complex I or II, inhibitors of the respiratory chain at complex III, inhibitors of the mitosis and cell division, compounds capable to induce a host
 10 defense, inhibitors of the amino acid and/or protein biosynthesis, inhibitors of the ATP production, inhibitors of the cell wall synthesis, inhibitors of the lipid and membrane synthesis, inhibitors of the melanine biosynthesis, inhibitors of the nucleic acid synthesis,
 15 inhibitors of the signal transduction, compounds capable to act as an uncoupler such as binapacryl, dinocap, ferimzone, fluazinam, meptyldinocap and further compounds, like for example benthiazole, bethoxazin, capsimycin, carvone, chinomethionat,
 20 pyriofenone (chlazafenone), cufraneb, cyflufenamid, cymoxanil, cyprosulfamide, dazomet, debacarb, dichlorophen, diclomezine, difenzoquat, difenzoquat methylsulphate, diphenylamine, ecomate, fenpyrazamine, flumetover, fluoroimide, flusulfamide, flutianil, fosetyl-aluminium, fosetyl-calcium, fosetyl-sodium, hexachlorobenzene, irumamycin, methasulfocarb, methyl isothiocyanate, metrafenone, mildiomyacin, natamycin, nickel dimethyldithiocarbamate, nitrothal-
 25 isopropyl, octhillinone, oxamocarb, oxyfenthiin, pentachlorophenol and salts, phenothrin, phosphorous acid and its salts, propamocarb-fosetylolate, propanosine-sodium, proquinazid, pyrimorph, (2E)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one, (2Z)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one, pyrrolnitrine, tebufloquin, tecloftalam, tolnifanide,
 30 triazoxide, trichlamide, zarilamid, (3S,6S,7R,8R)-8-benzyl-3-[[3-[(isobutyryloxy)methoxy]-4-methoxypyridin-2-yl]carbonyl]amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl 2-methylpropanoate, 1-(4-{4-[(5R)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, 1-(4-{4-[(5S)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-
 35 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, 1-(4-{4-[5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, 1-(4-methoxyphenoxy)-3,3-dimethylbutan-2-yl 1H-imidazole-1-carboxylate, 2,3,5,6-tetrachloro-4-(methylsulfonyl)pyridine, 2,3-dibutyl-6-chlorothieno[2,3-d]pyrimidin-4(3H)-one, 2,6-dimethyl-1H,5H-[1,4]dithiino[2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetrone, 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-(4-{4-

[(5R)-5-phenyl-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl]piperidin-1-yl)ethanone, 2-[5-
 methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-(4-{4-[(5S)-5-phenyl-4,5-dihydro-1,2-oxazol-3-
 yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone, 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-
 yl]-1-(4-[4-(5-phenyl-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidin-1-yl)ethanone,
 5 2-butoxy-6-iodo-3-propyl-4H-chromen-4-one, 2-chloro-5-[2-chloro-1-(2,6-difluoro-4-
 methoxyphenyl)-4-methyl-1H-imidazol-5-yl]pyridine, 2-phenylphenol and salts, 3-(4,4,5-
 trifluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinolone, 3,4,5-trichloropyridine-2,6-
 dicarbonitrile, 3-[5-(4-chlorophenyl)-2,3-dimethyl-1,2-oxazolidin-3-yl]pyridine, 3-chloro-5-
 (4-chlorophenyl)-4-(2,6-difluorophenyl)-6-methylpyridazine, 4-(4-chlorophenyl)-5-(2,6-
 10 difluorophenyl)-3,6-dimethylpyridazine, 5-amino-1,3,4-thiadiazole-2-thiol, 5-chloro-N'-
 phenyl-N'-(prop-2-yn-1-yl)thiophene-2-sulfonylhydrazide, 5-fluoro-2-[(4-
 fluorobenzyl)oxy]pyrimidin-4-amine, 5-fluoro-2-[(4-methylbenzyl)oxy]pyrimidin-4-amine, 5-
 methyl-6-octyl[1,2,4]triazolo[1,5-a]pyrimidin-7-amine, ethyl (2Z)-3-amino-2-cyano-3-
 phenylprop-2-enoate, N'-(4-{[3-(4-chlorobenzyl)-1,2,4-thiadiazol-5-yl]oxy}-2,5-
 15 dimethylphenyl)-N-ethyl-N-methylimidoforamide, N-(4-chlorobenzyl)-3-[3-methoxy-4-
 (prop-2-yn-1-yloxy)phenyl]propanamide, N-[(4-chlorophenyl)(cyano)methyl]-3-[3-
 methoxy-4-(prop-2-yn-1-yloxy)phenyl]propanamide, N-[(5-bromo-3-chloropyridin-2-
 yl)methyl]-2,4-dichloropyridine-3-carboxamide, N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-
 2,4-dichloropyridine-3-carboxamide, N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2-fluoro-4-
 20 iodopyridine-3-carboxamide, N-{(E)-[(cyclopropylmethoxy)imino][6-(difluoromethoxy)-
 2,3-difluorophenyl]methyl}-2-phenylacetamide, N-{(Z)-[(cyclopropylmethoxy)imino][6-
 (difluoromethoxy)-2,3-difluorophenyl]methyl}-2-phenylacetamide, N'-{4-[(3-tert-butyl-4-
 cyano-1,2-thiazol-5-yl)oxy]-2-chloro-5-methylphenyl}-N-ethyl-N-methylimidoforamide,
 N-methyl-2-(1-{[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-N-
 25 (1,2,3,4-tetrahydronaphthalen-1-yl)-1,3-thiazole-4-carboxamide, N-methyl-2-(1-{[5-
 methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-N-[(1R)-1,2,3,4-
 tetrahydronaphthalen-1-yl]-1,3-thiazole-4-carboxamide, N-methyl-2-(1-{[5-methyl-3-
 (trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-N-[(1S)-1,2,3,4-
 tetrahydronaphthalen-1-yl]-1,3-thiazole-4-carboxamide, pentyl {6-[[[1-methyl-1H-
 30 tetrazol-5-yl](phenyl)methylidene]amino}oxy)methyl]pyridin-2-yl}carbamate, phenazine-
 1-carboxylic acid, quinolin-8-ol, quinolin-8-ol sulfate (2:1), tert-butyl {6-[[[1-methyl-1H-
 tetrazol-5-yl](phenyl)methylene]amino}oxy)methyl]pyridin-2-yl}carbamate, 1-methyl-3-
 (trifluoromethyl)-N-[2'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, N-(4'-
 chlorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, N-(2',4'-
 35 dichlorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, 3-

(difluoromethyl)-1-methyl-N-[4'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, N-(2',5'-difluorobiphenyl-2-yl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide, 3-(difluoromethyl)-1-methyl-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, 5-fluoro-1,3-dimethyl-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, 2-chloro-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide, 3-(difluoromethyl)-N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]-1-methyl-1H-pyrazole-4-carboxamide, N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide, 3-(difluoromethyl)-N-(4'-ethynylbiphenyl-2-yl)-1-methyl-1H-pyrazole-4-carboxamide, N-(4'-ethynylbiphenyl-2-yl)-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide, 2-chloro-N-(4'-ethynylbiphenyl-2-yl)pyridine-3-carboxamide, 2-chloro-N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide, 4-(difluoromethyl)-2-methyl-N-[4'-(trifluoromethyl)biphenyl-2-yl]-1,3-thiazole-5-carboxamide, 5-fluoro-N-[4'-(3-hydroxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1,3-dimethyl-1H-pyrazole-4-carboxamide, 2-chloro-N-[4'-(3-hydroxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide, 3-(difluoromethyl)-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1-methyl-1H-pyrazole-4-carboxamide, 5-fluoro-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1,3-dimethyl-1H-pyrazole-4-carboxamide, 2-chloro-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide, (5-bromo-2-methoxy-4-methylpyridin-3-yl)(2,3,4-trimethoxy-6-methylphenyl)methanone, N-[2-(4-{{[3-(4-chlorophenyl)prop-2-yn-1-yl]oxy}-3-methoxyphenyl]ethyl}-N2-(methylsulfonyl)valinamide 4-oxo-4-[[2-phenylethyl]amino]butanoic acid, but-3-yn-1-yl {6-[[{[(Z)-(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino}oxy)methyl]pyridin-2-yl}carbamate, 4-Amino-5-fluoropyrimidin-2-ol, propyl 3,4,5-trihydroxybenzoate and oryzastrobin.

7. The composition according to any one of claims 1 to 6, wherein the biological control agent is selected from the group consisting of *Bacillus pumilus* (NRRL Accession No. B-30087), *Bacillus subtilis* AQ30002 (NRRL Accession No. B-50421), and *Bacillus subtilis* AQ713 (NRRL Accession No. B-21661) and/or a mutant of these strains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens.

8. The composition according to any one of claims 1 to 7 additionally comprising at least one auxiliary selected from the group consisting of extenders, solvents, spontaneity promoters, carriers, emulsifiers, dispersants, frost protectants, thickeners and adjuvants.

9. A seed treated with the composition according to any one of claims 1 to 8.
10. A seed according to claim 9, wherein the insecticide in the composition is selected from the group consisting of Carbofuran, Methiocarb, Thiodicarb, Ethiprole, Fipronil, and Abamectin.
- 5 11. A use of the composition according to any one of claims 1 to 8 as insecticide and/or fungicide.
12. The use according to claim 11 for reducing overall damage of plants and plant parts as well as losses in harvested fruits or vegetables caused by insects, mites, nematodes and/or phytopathogens.
- 10 13. A method for reducing overall damage of plants and plant parts as well as losses in harvested fruits or vegetables caused by insects, mites, nematodes and/or phytopathogens comprising the step of simultaneously or sequentially applying at least one biological control agent selected from the group consisting of
- 15 *Bacillus chitinosporus* AQ746 (NRRL Accession No. B-21618), *Bacillus mycooides* AQ726 (NRRL Accession No. B-21664), *Bacillus pumilus* (NRRL Accession No. B-30087), *Bacillus pumilus* AQ717 (NRRL Accession No. B-21662), *Bacillus* sp. AQ175 (ATCC Accession No. 55608), *Bacillus* sp. AQ177 (ATCC Accession No. 55609), *Bacillus* sp. AQ178 (ATCC Accession No. 53522), *Bacillus subtilis* AQ743 (NRRL Accession No. B-21665), *Bacillus subtilis* AQ713 (NRRL Accession No. B-21661), *Bacillus subtilis* AQ153 (ATCC Accession
- 20 No. 55614), *Bacillus thuringiensis* BD#32 (NRRL Accession No. B-21530), *Bacillus thuringiensis* AQ52 (NRRL Accession No. B-21619), *Muscodora albus* 620 (NRRL Accession No. 30547), *Muscodora roseus* A3-5 (NRRL Accession No. 30548), *Rhodococcus globerulus* AQ719 (NRRL Accession No. B-21663), *Streptomyces galbus* (NRRL Accession No. 30232), *Streptomyces* sp. (NRRL Accession No. B-30145), *Bacillus thuringiensis*
- 25 *subspec. kurstaki* BMP 123, *Bacillus subtilis* AQ30002 (NRRL Accession No. B-50421), and *Bacillus subtilis* AQ 30004 (NRRL Accession No. B-50455),
- and/or a mutant of these strains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens

and at least one insecticide selected from the group consisting of acetylcholinesterase (AChE) inhibitors, GABA-gated chloride channel antagonists, chloride channel activators and nicotinic acetylcholine receptor (nAChR) channel blockers,

and optionally at least one fungicide on the plant, plant parts, harvested fruits,
5 vegetables and/or plant's locus of growth in a synergistically effective amount, with the proviso that the biological control agent and optional fungicide are not identical.

14. The method according to claim 13, wherein

the acetylcholinesterase (AChE) inhibitor is selected from the group consisting of carbamates, and organophosphates, and/or

10 the GABA-gated chloride channel antagonist is selected from the group consisting of cyclodiene organochlorines, and phenylpyrazoles (fiproles), and/or

the chloride channel activator is an avermectin/milbemycin.

15. Kit-of-parts comprising at least one biological control agent selected from the group consisting of

15 *Bacillus chittinosporus* AQ746 (NRRL Accession No. B-21618), *Bacillus mycoides* AQ726 (NRRL Accession No. B-21664), *Bacillus pumilus* (NRRL Accession No. B-30087), *Bacillus pumilus* AQ717 (NRRL Accession No. B-21662), *Bacillus sp.* AQ175 (ATCC Accession No. 55608), *Bacillus sp.* AQ177 (ATCC Accession No. 55609), *Bacillus sp.* AQ178 (ATCC Accession No. 53522), *Bacillus subtilis* AQ743 (NRRL Accession No. B-21665), *Bacillus*
20 *subtilis* AQ713 (NRRL Accession No. B-21661), *Bacillus subtilis* AQ153 (ATCC Accession No. 55614), *Bacillus thuringiensis* BD#32 (NRRL Accession No. B-21530), *Bacillus thuringiensis* AQ52 (NRRL Accession No. B-21619), *Muscodora albus* 620 (NRRL Accession No. 30547), *Muscodora roseus* A3-5 (NRRL Accession No. 30548), *Rhodococcus globerulus* AQ719 (NRRL Accession No. B-21663), *Streptomyces galbus* (NRRL Accession
25 No. 30232), *Streptomyces sp.* (NRRL Accession No. B-30145), *Bacillus thuringiensis* subspec. *kurstaki* BMP 123, *Bacillus subtilis* AQ30002 (NRRL Accession No. B-50421), and *Bacillus subtilis* AQ 30004 (NRRL Accession No. B-50455),

and/or a mutant of these strains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits
30 activity against insects, mites, nematodes and/or phytopathogens,

and at least one insecticide selected from the group consisting of acetylcholinesterase (AChE) inhibitors, GABA-gated chloride channel antagonists, chloride channel activators and nicotinic acetylcholine receptor (nAChR) channel blockers,

in a synergistically effective amount in a spatially separated arrangement.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2013/061023

A. CLASSIFICATION OF SUBJECT MATTER
 INV. A01N47/22 A01N47/24 A01N57/14 A01N57/16 A01N57/28
 A01N63/00
 ADD.
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6 277 624 B1 (OUTTRUP HELLE [DK] ET AL) 21 August 2001 (2001-08-21) column 4, line 25 - line 32 column 5, line 59 - column 6, line 33 column 8, line 12 - line 18 column 8, line 47 - line 50 -----	1-4, 11-14
X	US 2008/064735 A1 (KERN MANFRED [DE]) 13 March 2008 (2008-03-13) paragraph [0006] - paragraph [0007] paragraph [0009] paragraph [0046] ----- -/--	1-5,8-14

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
---	---

Date of the actual completion of the international search 30 July 2013	Date of mailing of the international search report 29/10/2013
---	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Staber, Brigitte
--	--

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2013/061023

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ANNIE E.S. FOO, H.H. YAP: "Interactive effects of Bacillus thuringiensis H-14 with Chlorpyrifos and Permethrin", TROPICAL BIOMEDICINE, vol. 4, 1 January 1987 (1987-01-01), pages 47-50, XP009171556, page 47 table 1	1-4, 11-13
X	----- V. A. VASSILIOU: "Effectiveness of Insecticides in Controlling the First and Second Generations of the Lobesia botrana (Lepidoptera: Tortricidae) in Table Grapes", JOURNAL OF ECONOMIC ENTOMOLOGY, vol. 104, no. 2, 1 April 2011 (2011-04-01), pages 580-585, XP055073281, ISSN: 0022-0493, DOI: 10.1603/EC10343 page 581 page 583	1-4, 11-14
X	----- D.W. ATWOOD, S. Y. YOUNG III, T.J. KRING: "Mortality of Tobacco Budworm Larvae", J. ENTOMOL. SCI., vol. 33(2), 1 April 1998 (1998-04-01), pages 136-141, XP009171573, abstract	1-4
X	----- MORRIS O N: "LONG TERM STUDY OF THE EFFECTIVENESS OF AERIAL APPLICATION OF BACILLUS THURINGIENSIS - ACEPHATE COMBINATIONS AGAIST THE SPRUCE BUDWORM, CHRORISTONEURA FUMIFERANA (LEPIDOPTERA: TORTRICIDAE)", CANADIAN ENTOMOLOGIST, OTTAWA, CA, vol. 109, no. 9, 1 January 1977 (1977-01-01), pages 1239-1248, XP009171565, ISSN: 0008-347X the whole document	1-4,8, 11-14
X	----- WO 2009/060012 A2 (BASF SE [DE]; VOESTE DIRK [DE]; BRAHM LUTZ [DE]; WUTS MATHIJS [DE]; HA) 14 May 2009 (2009-05-14) page 1, line 28 - line 31 page 12, line 1 - line 13 claims 1, 20	1-15
X,P	----- WO 2012/135763 A1 (AGRAQUEST INC [US]; KLEIN CLARK D [US]; WILLIS REBECCA [US]; JOSE LUIS) 4 October 2012 (2012-10-04) paragraph [0009] - paragraph [0011] paragraph [0090] - paragraph [0091] paragraph [0182]	1-5,8,9, 11-14

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2013/061023

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

see annexe

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-15(partially)

A composition comprising at least one biological control agent selected from the group consisting of *Bacillus chitinosporus* AQ746 (Accession No. B-21618), *Bacillus mycoides* AQ726 (Accession No. B-21664), *Bacillus pumilus* (NRRL Accession No. B-30087), *Bacillus pumilus* AQ717 (NRRL Accession No. B-21662), *Bacillus* sp. AQ175 (ATCC Accession No. 55608), *Bacillus* sp. AQ177 (ATCC Accession No. 55609), *Bacillus* sp. AQ178 (ATCC Accession No. 53522), *Bacillus subtilis* AQ743 (NRRL Accession No. B-21665), *Bacillus subtilis* AQ713 (Accession No. B-21661), *Bacillus subtilis* AQ153 (ATCC Accession No. 55614), *Bacillus thuringiensis* BD#32 (NRRL Accession No. B-21530), *Bacillus thuringiensis* AQ52 (NRRL Accession No. B-21619), *Muscodor albus* 620 (Accession No. 30547), *Muscodor roseus* A3-5 (NRRL Accession No. 30548), *Rhodococcus* AQ719 (NRRL Accession No. B-21663), *Streptomyces galbus* (NRRL Accession No. 30232), *Streptomyces* sp. (NRRL Accession No. B-30145), *Bacillus thuringiensis* subspec. *kurstaki* BMP 123, *Bacillus subtilis* AQ30002 (NRRL Accession No. B-50421), and *Bacillus subtilis* AQ 30004 (NRRL Accession No. B-50455), and/or a mutant of these strains having all the identifying characteristics of the respective strain, and/or a metabolite produced by the respective strain that exhibits activity against insects, mites, nematodes and/or phytopathogens and at least one insecticide selected from the group consisting of acetylcholinesterase (AChE) inhibitors in a synergistically effective amount.

2. claims: 1-9, 11-15(all partially)

A composition comprising at least one biological control agent selected from the bacterial strains as specified under non-unity subject 1 and at least one insecticide selected from the group consisting of GABA-gated chloride channel antagonists in a synergistically effective amount.

3. claims: 1-9, 11-15(all partially)

A composition comprising at least one biological control agent selected from the bacterial strains as specified under non-unity subject 1 and at least one insecticide selected from the group consisting of chloride channel activators in a synergistically effective amount.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

4. claims: 1, 3, 5-9, 11-13, 15(all partially)

A composition comprising at least one biological control agent selected from the bacterial strains as specified under non-unity subject 1 and at least one insecticide selected from the group consisting of nicotinic acetylcholine receptor (nAChR) channel blockers in a synergistically effective amount.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/EP2013/061023

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
US 6277624	B1	21-08-2001	US 6277624 B1	21-08-2001
			US 2002004237 A1	10-01-2002

US 2008064735	A1	13-03-2008	AU 2002242678 A1	19-08-2002
			BR 0206926 A	25-02-2004
			DE 10104871 A1	08-08-2002
			EP 1367900 A2	10-12-2003
			EP 1922931 A2	21-05-2008
			EP 1967068 A2	10-09-2008
			JP 2004517945 A	17-06-2004
			KR 20030074780 A	19-09-2003
			MX PA03006934 A	18-11-2003
			US 2004078843 A1	22-04-2004
			US 2008064735 A1	13-03-2008
			WO 02062144 A2	15-08-2002

WO 2009060012	A2	14-05-2009	AR 069215 A1	06-01-2010
			WO 2009060012 A2	14-05-2009

WO 2012135763	A1	04-10-2012	NONE	
