



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2682015 A1 2008/10/02

(21) **2 682 015**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2008/03/27
(87) Date publication PCT/PCT Publication Date: 2008/10/02
(85) Entrée phase nationale/National Entry: 2009/09/25
(86) N° demande PCT/PCT Application No.: EP 2008/053592
(87) N° publication PCT/PCT Publication No.: 2008/116890
(30) Priorité/Priority: 2007/03/28 (EP07105104.9)

(51) Cl.Int./Int.Cl. *A61K 45/06* (2006.01),
A61K 31/496 (2006.01), *A61P 15/10* (2006.01),
A61P 15/12 (2006.01)
(71) Demandeur/Applicant:
BOEHRINGER INGELHEIM INTERNATIONAL GMBH,
DE
(72) Inventeur/Inventor:
BOECK, GEORG, DE
(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : NOUVELLES COMPOSITIONS PHARMACEUTIQUES
(54) Title: NEW PHARMACEUTICAL COMPOSITIONS

(57) **Abrégé/Abstract:**

The invention relates to new pharmaceutical compositions for the treatment of sexual disorders and methods for the preparation thereof. In a preferred embodiment, the instant invention is directed to pharmaceutical combinations comprising flibanserin as one active ingredient in combination with at least one additional active ingredient selected from the group consisting of compounds used for female hormone replacement therapy 2a, compounds used for menopause problems 2b, and contraceptives 2c.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 October 2008 (02.10.2008)

PCT

(10) International Publication Number
WO 2008/116890 A3

(51) International Patent Classification:

A61K 45/06 (2006.01) A61P 15/10 (2006.01)
A61K 31/496 (2006.01) A61P 15/12 (2006.01)

(21) International Application Number:

PCT/EP2008/053592

(22) International Filing Date: 27 March 2008 (27.03.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

07105104.9 28 March 2007 (28.03.2007) EP

(71) Applicant (for all designated States except DE, US):

BOEHRINGER INGELHEIM INTERNATIONAL GMBH [DE/DE]; Binger Strasse 173, 55216 Ingelheim am Rhein (DE).

(71) Applicant (for DE only): **BOEHRINGER INGELHEIM**

PHARMA GMBH & CO. KG [DE/DE]; Binger Strasse 173, 55216 Ingelheim Am Rhein (DE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **BOECK, Georg** [DE/DE]; Ahornweg 48, 88471 Laupheim (DE).

(74) Agent: **HAMMANN, Heinz**; Binger Str. 173, 55216 Ingelheim Am Rhein (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available):

ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

6 August 2009

(54) Title: PHARMACEUTICAL COMPOSITIONS COMPRISING FLIBANSERIN AND A FURTHER AGENT IN THE TREATMENT OF SEXUAL DISORDERS

(57) Abstract: The invention relates to new pharmaceutical compositions for the treatment of sexual disorders and methods for the preparation thereof. In a preferred embodiment, the instant invention is directed to pharmaceutical combinations comprising flibanserin as one active ingredient in combination with at least one additional active ingredient selected from the group consisting of compounds used for female hormone replacement therapy 2a, compounds used for menopause problems 2b, and contraceptives 2c.



WO 2008/116890 A3


New pharmaceutical compositions

The invention relates to new pharmaceutical combinations comprising flibanserin as one active ingredient in combination with at least one additional active ingredient for the treatment of sexual disorders and methods for the preparation thereof.

Background of the invention

The instant invention is directed to pharmaceutical combinations comprising a therapeutically effective amount of flibanserin **1** as one active ingredient in combination with a therapeutically effective amount of at least one additional active ingredient **2** selected from the group consisting of compounds used for female hormone replacement therapy **2a**, compounds used for menopause problems **2b**, and contraceptives **2c**, for the treatment of sexual disorders and methods for the preparation thereof.

The compound 1-[2-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)ethyl]-2,3-dihydro-1H-benzimidazol-2-one (flibanserin) is disclosed in form of its hydrochloride in European Patent Application EP-A-526434 and has the following chemical structure:



1 x HCl

Flibanserin shows affinity for the 5-HT_{1A} and 5-HT₂-receptor. It is therefore a promising therapeutic agent for the treatment of a variety of diseases, for instance depression, schizophrenia, Parkinson, anxiety, sleep disturbances, sexual and mental disorders and age associated memory impairment.

One embodiment of the invention is directed to pharmaceutical compositions comprising a therapeutically effective amount of flibanserin **1** in combination with a therapeutically effective amount of one or more, preferably one active ingredient **2** selected from the group consisting of compounds used for female hormone replacement therapy **2a**, compounds used for menopause problems **2b**, and contraceptives **2c**.

A preferred embodiment of the present invention relates to pharmaceutical compositions comprising a therapeutically effective amount of flibanserin **1** in

combination with a therapeutically effective amount of one or more, preferably one contraceptive **2c**.

The compositions according to the invention may contain flibanserin **1** and the one
5 or more additional active ingredient **2** in a single formulation or in separate formulations. If flibanserin and the one or more additional active ingredient are present in separate formulations these separate formulations may be administered simultaneously or sequentially.

10 The combinations of the present invention are especially useful for female patients who are in need of hormone replacement therapy, female patients with menopause problems and females receiving contraceptives and, in addition, suffer from sexual diseases. By combination of flibanserin with those other drugs, dosing is simplified, is more convenient and therefore leads to better compliance with therapy.

15 A preferred embodiment according to the invention is directed to pharmaceutical compositions comprising a therapeutically effective amount of flibanserin **1** and a therapeutically effective amount of one or more, preferably one compound used for female hormone replacement therapy **2a**, optionally in combination with a
20 pharmaceutically acceptable excipient. Examples of suitable compounds used for female hormone replacement therapy **2a** include chlormadinone acetate, dienogest, dydrogesterone, estradiol valerate, medroxyprogesterone, medroxyprogesterone acetate, norethisterone and norethisterone acetate, optionally in form of the pharmaceutically acceptable acid addition salts, in form of the hydrates and/or
25 solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof.

Another preferred embodiment according to the invention is directed to pharmaceutical compositions comprising a therapeutically effective amount of
30 flibanserin **1** and a therapeutically effective amount of one or more, preferably one compound used for menopause problems **2b**, optionally in combination with a pharmaceutically acceptable excipient. Examples of suitable compounds used for menopause problems **2b** include estradiol valerate, medroxyprogesterone acetate, norgestrel, prasterone, and progesterone, optionally in form of the
35 pharmaceutically acceptable salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof.

A more preferred embodiment according to the invention is directed to pharmaceutical compositions comprising a therapeutically effective amount of flibanserin **1** and a therapeutically effective amount of one or more, preferably one contraceptive **2c**, optionally in combination with a pharmaceutically acceptable
5 excipient. Examples of suitable contraceptives **2c** include chlormadinone acetate, dienogest, drospirenone, etonogestrel, gestodene, medroxyprogesterone acetate, norelgestromine, norethisterone, norethisterone enantate, and norgestimate, optionally in form of the pharmaceutically acceptable salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures
10 of the individual enantiomers or racemates thereof.

Flibanserin **1** may be used in form of the free base, optionally in form of its pharmaceutically acceptable acid addition salts and/or optionally in form of the hydrates and/or solvates thereof. Suitable acid addition salts include for example
15 those of the acids selected from, succinic acid, hydrobromic acid, acetic acid, fumaric acid, maleic acid, methanesulphonic acid, lactic acid, phosphoric acid, hydrochloric acid, sulphuric acid, tartaric acid and citric acid. Mixtures of the abovementioned acid addition salts may also be used. From the aforementioned acid addition salts the hydrochloride and the hydrobromide, particularly the
20 hydrochloride, are preferred. If flibanserin **1** is used in form of the free base, it is preferably used in form of flibanserin polymorph A as disclosed in WO 03/014079.

The active ingredients **2** which are suitable to be combined with flibanserin within the teaching of the instant invention and which are mentioned hereinbefore may also be
25 capable of forming acid addition salts with pharmaceutically acceptable acids. Representative salts include the following: Acetate, Benzenesulfonate, Benzoate, Bicarbonate, Bisulfate, Bitartrate, Borate, Bromide, Camsylate, Carbonate, Chloride, Clavulanate, Citrate, Dihydrochloride, Edetate, Edisylate, Estolate, Esylate, Fumarate, Gluceptate, Gluconate, Glutamate, Glycolylarsanilate,
30 Hexylresorcinate, Hydrabamine, Hydrobromide, Hydrochloride, Hydroxynaphthoate, Iodide, Isothionate, Lactate, Lactobionate, Laurate, Malate, Maleate, Mandelate, Mesylate, Methylbromide, Methylnitrate, Methylsulfate, Mucate, Napsylate, Nitrate, N-methylglucamine ammonium salt, Oleate, Oxalate, Pamoate (Embonate), Palmitate, Pantothenate, Phosphate/diphosphate, Polygalacturonate, Salicylate, Stearate,
35 Sulfate, Subacetate, Succinate, Tannate, Tartrate, Teoclate, Tosylate, Triethiodide and Valerate.

Furthermore, where the compounds **2** carry an acidic moiety, suitable

pharmaceutically acceptable salts thereof may include alkali metal salts, e.g., sodium or potassium salts; alkaline earth metal salts, e. g., calcium or magnesium salts; and salts formed with suitable organic ligands, e.g., quaternary ammonium salts.

5

The compounds 2 may have chiral centers and occur as racemates, racemic mixtures and as individual diastereomers, or enantiomers with all isomeric forms being included in the present invention. Therefore, where a compound is chiral, the separate enantiomers, substantially free of the other, are included within the scope
10 of the invention. Further included are all mixtures of the two enantiomers. Also included within the scope of the invention are polymorphs and hydrates of the compounds of the instant invention.

The present invention includes within its scope prodrugs of the compounds 1 and 2.
15 In general, such prodrugs will be functional derivatives of the compounds of this invention which are readily convertible in vivo into the required compound.

The term "therapeutically effective amount" shall mean that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue,
20 system, animal or human that is being sought by a researcher or clinician.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in
25 the specified amounts.

In the combination of the present invention, the components 1 and 2 may be administered separately or together in one pharmaceutical composition. In addition, the administration of one element of the combination of the present invention may be
30 prior to, concurrent to, or subsequent to the administration of the other element of the combination.

The elements of the combination of 1 and 2 may be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous or subcutaneous injection, or
35 implant), buccal, nasal, vaginal, rectal, sublingual, or topical (e.a.. ocular eyedrop) routes of administration and may be formulated, alone or together, in suitable dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles appropriate for each route of

administration.

The pharmaceutical compositions for the administration of the components 1 and 2 of this invention may conveniently be presented in dosage unit form and
5 may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredient into association with the carrier which is constituted of one or more accessory ingredients. In general, the pharmaceutical compositions are prepared by uniformly and intimately bringing the active ingredients into association with a liquid carrier or a finely divided solid carrier
10 or both, and then, if necessary, shaping the product into the desired dosage form. In the pharmaceutical compositions the active compounds are included in an amount sufficient to produce the desired pharmacologic effect.

The pharmaceutical compositions containing the active ingredients 1 and 2,
15 separately or together, that are suitable for oral administration may be in the form of discrete units such as hard or soft capsules, tablets, troches or lozenges, each containing a predetermined amount of the active ingredients; in the form of a dispersible powder or granules; in the form of a solution or a suspension in an aqueous liquid or non-aqueous liquid; in the form of syrups or elixirs; or in the form
20 of an oil-in-water emulsion or a water-in-oil emulsion.

Dosage forms intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical formulations and such compositions.

25

The excipients used may be for example, (a) inert diluents such as mannitol, sorbitol, calcium carbonate, pregelatinized starch, lactose, calcium phosphate or sodium phosphate; (b) granulating and disintegrating agents, such as povidone, copovidone, hydroxypropylmethylcellulose, corn starch, alginic acid, crospovidone,
30 sodiumstarchglycolate, croscarmellose, or polacrilin potassium ; (c) binding agents such as microcrystalline cellulose or acacia ; and (d) lubricating agents such as magnesium stearate, stearic acid, fumaric acid or talc.

In some cases, formulations for oral use may be in the form of hardgelatin or HPMC
35 capsules wherein the active ingredient 1 or 2, separately or together, is mixed with an inert solid diluent, for example pregelatinized starch, calcium carbonate, calcium phosphate or kaolin, or dispensed via a pellet formulation. They may also be in the form of soft gelatin capsules wherein the active ingredient is mixed with water or an

oil medium, for example peanut oil, liquid paraffin, medium chain triglycerides or olive oil.

The tablets, capsules or pellets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a delayed action or sustained action over a longer period. For example, a time delay material such as celluloseacetate phtalate or hydroxypropylcellulose acetate succinate or sustained release material such as ethylcellulose or ammoniomethacrylate copolymer (type B) may be employed.

10

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Besides such inert diluents, compositions can also include adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, perfuming and preserving agents.

15

Aqueous suspensions normally contain the active materials 1 and 2, separately or together, in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients may be (a) suspending agents such as hydroxy ethylcellulose, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; (b) dispersing or wetting agents which may be (b.1) a naturally-occurring phosphatide such as lecithin, (b.2) a condensation product of an alkylene oxide with a fatty acid, for example, polyoxyethylene stearate, (b.3) a condensation product of ethylene oxide with a long chain aliphatic alcohol, for example heptadecaethyleneoxycetanol, (b.4) a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol such as polyoxyethylene sorbitol monooleate, or (b.5) a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride, for example polyoxyethylene sorbitan monooleate.

20

25

30

The aqueous suspensions may also contain one or more preservatives, for example, ethyl or n-propyl p-hydroxybenzoate; one or more coloring agents; one or more flavoring agents; and one or more sweetening agents, such as sucrose or saccharin.

35

Oily suspensions may be formulated by suspending the active ingredients 1 and 2, separately or together, in a vegetable oil, for example arachis oil, olive oil, sesame

oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents and flavoring agents may be added to provide a palatable oral preparation. These compositions may be prepared by the addition of an antioxidant
5 such as ascorbic acid.

Dispersible powders and granules are suitable for the preparation of an aqueous suspension. They provide the active ingredients 1 and 2, separately or together, in admixture with a dispersing or wetting agent, a suspending agent and one or more
10 preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example, those sweetening, flavoring and coloring agents described above may also be present.

15 The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil such as olive oil or arachis oils, or a mineral oil such as liquid paraffin or a mixture thereof.

Suitable emulsifying agents may be (a) naturally-occurring gums such as gum acacia
20 and gum tragacanth, (b) naturally-occurring phosphatides such as soybean and lecithin, (c) esters or partial esters derived from fatty acids and hexitol anhydrides, for example, sorbitan monooleate, (d) condensation products of said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

25 Syrups and elixirs may be formulated with sweetening agents, for example, glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a preservative and flavoring and coloring agents.

30 The pharmaceutical compositions containing 1 and 2, separately or together, may be in the form of a sterile injectable aqueous or oleagenous suspension or solution. The suspension may be formulated according to known methods using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or
35 suspension in a non toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane-diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending

medium. For this purpose any bland fixed oil may be employed including synthetic mono-or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

- 5 Preparations according to this invention containing 1 and 2, separately or together, for parenteral administration include sterile aqueous or non-aqueous solutions, suspension, or emulsions.

Examples of non-aqueous solvents or vehicles are propylene glycol, polyethylene
10 glycol, vegetable oils, such as olive oil and corn oil, gelatin, and injectable organic esters such as ethyl oleate. Such dosage forms may also contain adjuvants such as preserving, wetting, emulsifying, and dispersing agents. They may be sterilized by, for example, filtration through a bacteria-retaining filter, by incorporating sterilizing agents into the compositions, by irradiating the compositions, or by heating the
15 compositions. They can also be manufactured in the form of sterile solid compositions which can be reconstituted in sterile water, or some other sterile injectable medium immediately before use. The combination of this invention may also be administered in the form of suppositories for rectal administration. This composition can be prepared by mixing the drugs with a suitable non-irritating
20 excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter, hard fat, and polyethylene glycols. Compositions for buccal, nasal or sublingual administration are also prepared with standard excipients well known in the art.

25

For topical administration the combinations of this invention containing 1 and 2, separately or together, may be formulated in liquid or semi-liquid preparations such as liniments, lotions, applications; oil-in-water or water-in-oil emulsions such as creams, ointments, jellies or pastes, including tooth-pastes; or solutions or
30 suspensions such as drops, and the like.

The dosage of the active ingredients in the compositions of this invention may be varied. However, it is necessary that the amount of the active ingredients 1 and 2 be such that a suitable dosage form is obtained. The selected dosage and the dosage
35 form depend upon the desired therapeutic effect, on the route of administration and on the duration of the treatment. Dosage ranges in the combination are approximately one tenth to one times the clinically effective ranges required to

induce the desired therapeutic effect, respectively when the compounds are used singly.

Within the instant invention flibanserin **1** is preferably administered in such an amount that per single dosage between 5 to 200 mg of flibanserin **1** are applied. Preferred are ranges of between 10 to 150 mg, particular preferred 20 to 100 mg of flibanserin **1**. Suitable dosage forms may contain for instance 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100 mg of flibanserin **1**. The aforementioned values are based on flibanserin **1** in form of the free base. If flibanserin **1** is applied in form of one of its acid addition salts, the corresponding values are readily calculable from the aforementioned values.

Within the instant invention the compounds used for female hormone replacement therapy **2a** are preferably administered in a range of between about 0.001 mg per kg of bodyweight per day (mg/kg/day) to about 100 mg/kg/day, preferably 0.01 to 50 mg/kg/day, and most preferably 0.1 to 30 mg/kg/day. Intravenously, the most preferred doses will range from about 0.1 to about 10 mg/kg/minute during a constant rate infusion. Advantageously, the compounds **2a** of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Suitable dosage forms may contain for instance 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, 0.95, 1, 1.05, 1.1, 1.15, 1.2, 1.25, 1.3, 1.35, 1.4, 1.45, 1.5, 1.55, 1.6, 1.65, 1.7, 1.75, 1.8, 1.85, 1.9, 1.95, 2, 2.05, 2.1, 2.15, 2.2, 2.25, 2.3, 2.35, 2.4, 2.45, 2.5, 2.55, 2.6, 2.65, 2.7, 2.75, 2.8, 2.85, 2.9, 2.95, 3, 3.05, 3.1, 3.15, 3.2, 3.25, 3.3, 3.35, 3.4, 3.45, 3.5, 3.55, 3.6, 3.65, 3.7, 3.75, 3.8, 3.85, 3.9, 3.95, 4, 4.05, 4.1, 4.15, 4.2, 4.25, 4.3, 4.35, 4.4, 4.45, 4.5, 4.55, 4.6, 4.65, 4.7, 4.75, 4.8, 4.85, 4.9, 4.95, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 mg of **2a**. Advantageously, the compounds **2a** of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily.

Within the instant invention the compounds used for menopause problems **2b** are preferably administered in a range of between about 0.001 mg per kg of bodyweight per day (mg/kg/day) to about 100 mg/kg/day, preferably 0.01 to 50 mg/kg/day, and most preferably 0.1 to 10 mg/kg/day. Intravenously, the most preferred doses will range from about 0.1 to about 10 mg/kg/minute during a constant rate infusion. Advantageously, the compounds **2b** of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses

of two, three or four times daily. Suitable dosage forms may contain for instance 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, 0.95, 1, 1.05, 1.1, 1.15, 1.2, 1.25, 1.3, 1.35, 1.4, 1.45, 1.5, 1.55, 1.6, 1.65, 1.7, 1.75, 1.8, 1.85, 1.9., 1.95, 2, 2.05, 2.1, 2.15, 2.2, 2.25, 2.3, 2.35, 2.4, 2.45, 2.5, 2.55, 2.6, 5 2.65, 2.7, 2.75, 2.8, 2.85, 2.9., 2.95, 3, 3.05, 3.1, 3.15, 3.2, 3.25, 3.3, 3.35, 3.4, 3.45, 3.5, 3.55, 3.6, 3.65, 3.7, 3.75, 3.8, 3.85, 3.9., 3.95, 4, 4.05, 4.1, 4.15, 4.2, 4.25, 4.3, 4.35, 4.4, 4.45, 4.5, 4.55, 4.6, 4.65, 4.7, 4.75, 4.8, 4.85, 4.9., 4.95, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 mg of **2b**.

Advantageously, the compounds **2b** of the present invention may be administered in 10 a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily.

Within the instant invention the contraceptives **2c** are preferably administered in a range of between about 0.001 mg per kg of bodyweight per day (mg/kg/day) to about 15 100 mg/kg/day, preferably 0.01 to 10 mg/kg/day, and most preferably 0.1 to 1 mg/kg/day. Advantageously, the compounds **2c** of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Suitable dosage forms may contain for instance 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 20 0.8, 0.85, 0.9, 0.95, 1, 1.05, 1.1, 1.15, 1.2, 1.25, 1.3, 1.35, 1.4, 1.45, 1.5, 1.55, 1.6, 1.65, 1.7, 1.75, 1.8, 1.85, 1.9., 1.95, 2, 2.05, 2.1, 2.15, 2.2, 2.25, 2.3, 2.35, 2.4, 2.45, 2.5, 2.55, 2.6, 2.65, 2.7, 2.75, 2.8, 2.85, 2.9., 2.95, 3, 3.05, 3.1, 3.15, 3.2, 3.25, 3.3, 3.35, 3.4, 3.45, 3.5, 3.55, 3.6, 3.65, 3.7, 3.75, 3.8, 3.85, 3.9., 3.95, 4, 4.05, 4.1, 4.15, 4.2, 4.25, 4.3, 4.35, 4.4, 4.45, 4.5, 4.55, 4.6, 4.65, 4.7, 4.75, 4.8, 4.85, 4.9., 4.95, 5, 25 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 mg of **2c**. Advantageously, the compounds **2c** of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily.

30 However, it should be apparent to one skilled in the art that the exact dosage and frequency of administration will depend on the particular compounds of the invention administered, the particular condition being treated, the severity of the condition being treated, the age, weight, general physical condition of the particular patient, and other medication the individual may be taking as is well known to administering 35 physicians who are skilled in this art.

In another embodiment the invention relates to a method for the treatment of sexual disorders, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a therapeutically effective amount of 2, selected from the group consisting of compounds used for female hormone replacement therapy 2a, compounds used for menopause problems 2b, and contraceptives 2c, separately or together within one pharmaceutical composition.

The generic term "Sexual disorders" includes Sexual Desire Disorders (i.e. Hypoactive Sexual Desire Disorder, Sexual Aversion Disorder), Sexual Arousal Disorders (i.e. Female Sexual Arousal Disorder, Male Erectile Disorder), Orgasmic Disorders (i.e. Female Orgasmic Disorder, Male Orgasmic Disorder, Premature Ejaculation) Sexual Pain Disorders (i.e. Dyspareunia, Vaginismus), Sexual Dysfunction due to a General Medical Condition, Substance-Induced Sexual Dysfunction, and Sexual Dysfunction not otherwise specified (Diagnostic and Statistical Manual of Mental Disorders, 4th edition, Text Revision. Washington DC, American Psychiatric Association, 2000).

In another preferred embodiment the invention relates to a method for the treatment of disorders selected from the group consisting of hypoactive sexual desire disorder (HSDD), sexual aversion disorder, loss of sexual desire, lack of sexual desire, decreased sexual desire, inhibited sexual desire, loss of libido, libido disturbance, and frigidity, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable acid addition salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof, separately or together within one pharmaceutical composition.

In another preferred embodiment the invention relates to a method for the treatment of disorders selected from the group consisting of hypoactive sexual desire disorder (HSDD), sexual aversion disorder, loss of sexual desire, lack of sexual desire, decreased sexual desire, inhibited sexual desire, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable acid addition salts, in form of the hydrates and/or solvates and optionally

in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof, separately or together within one pharmaceutical composition.

In another preferred embodiment the invention relates to a method for treatment of disorders selected from the group of hypoactive sexual desire disorder (HSDD), decreased sexual desire and inhibited sexual desire, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable acid addition salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof, separately or together within one pharmaceutical composition.

In another preferred embodiment the invention relates to a method for treatment of premenstrual disorders, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable acid addition salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof, separately or together within one pharmaceutical composition.

In another preferred embodiment the invention relates to a method for treatment of premenstrual disorders selected from the group consisting of premenstrual dysphoria, premenstrual syndrome, premenstrual dysphoric disorder, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable acid addition salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof, separately or together within one pharmaceutical composition.

In another preferred embodiment the invention relates to a method for treatment of sexual arousal disorder in females, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable acid addition salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof, separately or together within one pharmaceutical composition.

In another preferred embodiment the invention relates to a method for treatment of orgasmic disorder in females, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable acid addition salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof, separately or together within one pharmaceutical composition.

In another preferred embodiment the invention relates to a method for treatment of sexual pain disorders in females, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable acid addition salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof, separately or together within one pharmaceutical composition.

In another preferred embodiment the invention relates to a method for treatment of sexual pain disorders selected from the group consisting of dyspareunia in females, vaginismus in females, and noncoital sexual pain disorder in females, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable acid addition salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof, separately or together within one pharmaceutical composition.

In another preferred embodiment the invention relates to a method for treatment of sexual dysfunction due to a general medical condition in females, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable acid addition salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof, separately or together within one pharmaceutical composition.

In another preferred embodiment the invention relates to a method for treatment of substance-induced sexual dysfunction in females, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a

therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable acid addition salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof, separately or together within one pharmaceutical composition.

5

The beneficial effects of the compounds of formula (I) can be observed regardless of whether the above mentioned sexual disorders existed lifelong or was acquired, is of the „generalized type“ or „situational type“ and independent of etiologic origin (organic - both, physically and drug induced-, psychogen (due to psychological factors), a combination of organic - both, physically and drug induced-, and psychogen (due to combined factors), or unknown) origin. The term “lifelong” refers to such sexual disorders of the present invention, which have been present since the onset of sexual functioning. The term “acquired” refers to such sexual disorders of the present invention which developed only after a period of normal sexual functioning. The „generalized type“ refers to such sexual disorders of the present invention wherein the disorder is not limited to certain types of stimulation, situations, or partners. The „situational type“ applies to such sexual disorders of the present invention wherein the disorder is limited to certain types of stimulation, situations, or partners. The subtype due to “psychological factors” applies when psychological factors are judged to have the major role in the onset, severity, exacerbation, or maintenance of the Sexual Disorder, and general medical conditions and substance play no role in the etiology of the sexual disorder. Finally the subtype due to “combined factors” applies when 1) psychological factors are judged to have a role in the onset, severity, exacerbation, or maintenance of the sexual disorder, and 2) a general medical condition or substance use is also judged to be contributory but is not sufficient to account for a Sexual Disorder (Diagnostic and Statistical Manual of Mental Disorders, 4th edition, Text Revision. Washington DC, American Psychiatric Association, 2000).

The beneficial effects of the compounds of formula (I) can also be observed regardless of whether the females suffering from above mentioned diseases are in the pre-menopausal, peri-menopausal or post-menopausal state.

30

Another embodiment of the present invention relates to the use of the combinations of a therapeutically effective amount of flibanserin 1, and of a therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable salts, in form of the hydrates and/or solvates and optionally in the form of the individual
5 optical isomers, mixtures of the individual enantiomers or racemates thereof, for the manufacture of a medicament for the treatment of any of the aforementioned disorders.

Another embodiment of the present invention relates to the use of a therapeutically
10 effective amount of flibanserin 1, for the manufacture of a medicament for the treatment of any of the aforementioned disorders in combination with a therapeutically effective amount of 2, optionally in form of the pharmaceutically acceptable salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates
15 thereof.

In a preferred embodiment, the invention relates to any of the above mentioned methods and uses wherein the therapeutically effective amount of 2 is selected from compounds used for female hormone replacement therapy 2a.

20 In a more preferred embodiment, the invention relates to any of the above mentioned methods and uses wherein the therapeutically effective amount of compounds used for female hormone replacement therapy 2a are selected from the group consisting of chlormadinone acetate, dienogest, dydrogesterone, estradiol valerate,
25 medroxyprogesterone, medroxyprogesterone acetate, norethisterone and norethisterone acetate.

In a preferred embodiment, the invention relates to any of the above mentioned methods and uses wherein the therapeutically effective amount of 2 is selected from
30 compounds used for menopause problems 2b.

In a more preferred embodiment, the invention relates to any of the above mentioned methods and uses wherein the therapeutically effective amount of 2 is selected from compounds used for menopause problems 2b are selected from the group consisting
35 of estradiol valerate, medroxyprogesterone acetate, norgestrel, prasterone, and progesterone.

In a preferred embodiment, the invention relates to any of the above mentioned methods and uses wherein the therapeutically effective amount of **2** is selected from
5 contraceptives **2c**.

In a more preferred embodiment, the invention relates to any of the above mentioned methods and uses wherein the therapeutically effective amount of **2** is selected from
10 contraceptives **2c** are selected from the group consisting of chlormadinone acetate, dienogest, drospirenone, etonogestrel, gestodene, medroxyprogesterone acetate, norelgestromine, norethisterone, norethisterone enantate, and norgestimate.

The following examples demonstrate possible pharmaceutical compositions comprising flibanserin in combination with one of the aforementioned combination
15 partners **2**.

Example N°1 – Combination 1

Core

<u>Constituents</u>	mg/tablet
Flibanserin (free base)	50.000
Dienogest	2.000
Anhydrous dibasic calcium phosphate	100.000
Microcrystalline cellulose	203.090
HPMC (Methocel E5)	6.615
Croscarmellose sodium	8.820
Magnesium stearate	2.250

20 **Coating**

<u>Constituents</u>	mg/ tablet
HPMC (Methocel E5)	4.320
Polyethylene Glycol 6000	1.260
Titanium dioxide	1.800
Talc	1.542
Iron oxide red	0.078

Total Film coated tablet	381.775
---------------------------------	----------------

Example N°2 – Combination 2**Core**

<u>Constituents</u>	mg/tablet
Flibanserin (free base)	50.000
Estradiol valerate	2.000
Lactose monohydrate	133.750
Microcrystalline cellulose	40.000
Hydroxypropylcellulose	2.500
Corn starch	12.500
Magnesium stearate	1.250

5

Coating

<u>Constituents</u>	mg/ tablet
HPMC (e.g. Pharmacoat 606)	2.400
Polyethylene Glycol 6000	0.700
Titanium dioxide	1.000
Talc	0.857
Iron oxide yellow	0.043

Total Film coated tablet	247.000
---------------------------------	----------------

Example N°3 – Combination 310 **Core**

<u>Constituents</u>	mg/tablet
Flibanserin (free base)	50.000
Chlormadinone acetate	1.000
Lactose monohydrate	143.490
Microcrystalline cellulose	47.810
HPMC (e.g. Pharmacoat 606)	2.500
Carboxymethylcellulose sodium	5.000
Mannitol	60.000

Corn starch	36.500
Povidone	1.000
Colloidal silicon dioxide	1.000
Magnesium stearate	1.700

Coating

<u>Constituents</u>	mg/ tablet
HPMC (e.g. Methocel E5)	3.360
Polyethylene Glycol 6000	0.980
Titanium dioxide	1.400
Talc	1.200
Iron oxide red	0.060
Total Film coated bilayer tablet	357.000

- 5 The following examples show preferred pharmaceutical compositions of flibanserin, if the combinations according to the invention are administered in separate dosage units.

Example N°4 - Composition10 Core

<u>Constituents</u>	mg/tablet
Flibanserin (free base)	25.000
Lactose monohydrate	71.720
Microcrystalline cellulose	23.905
HPMC (Methocel E5)	1.250
Carboxymethylcellulose sodium	2.500
Magnesium stearate	0.625

Coating

<u>Constituents</u>	mg/ tablet
HPMC (Methocel E5)	1.440
Polyethylene Glycol 6000	0.420
Titanium dioxide	0.600

Talc	0.514
Iron oxide red	0.026
Total Film coated tablet	128.000

Example N°5 - Composition5 Core

<u>Constituents</u>	mg/tablet
Flibanserin (free base)	50.000
Lactose monohydrate	143.440
Microcrystalline cellulose	47.810
HPMC (e.g. Pharmacoat 606)	2.500
Carboxymethylcellulose sodium	5.000
Magnesium stearate	1.250

Coating

<u>Constituents</u>	mg/ tablet
HPMC (e.g. Pharmacoat 606)	2.400
Polyethylene Glycol 6000	0.700
Titanium dioxide	1.000
Talc	0.857
Iron oxide red	0.043

Total Film coated tablet	255.000
---------------------------------	----------------

10

Example N°6 - CompositionCore

<u>Constituents</u>	mg/tablet
Flibanserin (free base)	100.000
Lactose monohydrate	171.080
Microcrystalline cellulose	57.020
HPMC (e.g. Methocel E5)	3.400
Carboxymethylcellulose sodium	6.800

Magnesium stearate	1.700
--------------------	-------

Coating

<u>Constituents</u>	mg/ tablet
HPMC (e.g. Methocel E5)	3.360
Polyethylene Glycol 6000	0.980
Titanium dioxide	1.400
Talc	1.200
Iron oxide red	0.060
Total Film coated tablet	347.000

5 **Example N°7** – CompositionCore

<u>Constituents</u>	mg/tablet
Flibanserin (free base)	2.000
Dibasic Calciumphosphate, anhydrous	61.010
Microcrystalline cellulose	61.010
HPMC (Methocel E5)	1.950
Carboxymethylcellulose sodium	2.600
Colloidal silicon dioxide	0.650
Magnesium stearate	0.780

Coating

<u>Constituents</u>	mg/ tablet
HPMC (Methocel E5)	1.440
Polyethylene Glycol 6000	0.420
Titanium dioxide	0.600
Talc	0.514
Iron oxide red	0.026
Total Film coated tablet	133.000

Example N°8 – Composition

Core

<u>Constituents</u>	mg/tablet
Flibanserin (free base)	100.000
Dibasic Calciumphosphate, anhydrous	69.750
Microcrystalline cellulose	69.750
HPMC (e.g. Methocel E5)	2.750
Carboxymethylcellulose sodium	5.000
Colloidal silicon dioxide	1.250
Magnesium stearate	1.500

Coating

<u>Constituents</u>	mg/ tablet
HPMC (e.g. Methocel E5)	2.400
Polyethylene Glycol 6000	0.700
Titanium dioxide	1.043
Talc	0.857

5

Total Film coated tablet	255.000
---------------------------------	----------------

Example N°9 – Composition10 Core

<u>Constituents</u>	mg/tablet
Flibanserin (free base)	20.000
Lactose monohydrate	130.000
Microcrystalline cellulose	43.100
Hydroxypropyl Cellulose (e.g. Klucel LF)	1.900
Sodium Starch Glycolate	4.000
Magnesium stearate	1.000

Coating

<u>Constituents</u>	mg/ tablet
----------------------------	-------------------

HPMC (e.g. Methocel E5)	2.400
Polyethylene Glycol 6000	0.700
Titanium dioxide	1.043
Talc	0.857

Total Film coated tablet	205.000
---------------------------------	----------------

Claims

- 1) Pharmaceutical compositions comprising a therapeutically effective amount of flibanserin 1 as one active ingredient in combination with a therapeutically effective amount of at least one additional active ingredient 2, selected from the group consisting of compounds used for female hormone replacement therapy 2a, compounds used for menopause problems 2b, and contraceptives 2c.
- 2) Pharmaceutical compositions according to claim 1, comprising a therapeutically effective amount of flibanserin 1 and a therapeutically effective amount of one or more, preferably one contraceptive 2c, optionally in combination with a pharmaceutically acceptable excipient.
- 3) Pharmaceutical compositions according to claim 2, wherein the contraceptive 2c is selected from chlormadinone acetate, dienogest, drospirenone, etonogestrel, gestodene, medroxyprogesterone acetate, norelgestromine, norethisterone, norethisterone enantate, and norgestimate, optionally in form of the pharmaceutically acceptable salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof.
- 4) Pharmaceutical compositions according to one or more of claims 1 to 3, comprising the active ingredients 1 and 2 together in one dosage form.
- 5) Pharmaceutical compositions according to one or more of claims 1 to 3, comprising the active ingredients 1 and 2 separately, each in one dosage form.
- 6) Method for the treatment of sexual disorders, comprising the administration of a therapeutically effective amount of flibanserin 1, in combination with a therapeutically effective amount of 2, selected from the group consisting of compounds used for female hormone replacement therapy 2a, compounds used for menopause problems 2b, and contraceptives 2c, separately or together within one pharmaceutical composition.
- 7) Use of a combination of a therapeutically effective amount of flibanserin 1, and of a therapeutically effective amount of 2, selected from the group consisting of compounds used for female hormone replacement therapy 2a, compounds used for menopause problems 2b, and contraceptives 2c, optionally in form of

the pharmaceutically acceptable salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof, for the manufacture of a medicament for the treatment of sexual disorders.

5

- 8) Use of a therapeutically effective amount of flibanserin 1, for the manufacture of a medicament for the treatment of sexual disorders in combination with a therapeutically effective amount of 2, selected from the group consisting of compounds used for female hormone replacement therapy 2a, compounds used for menopause problems 2b, and contraceptives 2c, optionally in form of the pharmaceutically acceptable salts, in form of the hydrates and/or solvates and optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates thereof.

10

- 9) Method or use according to one or more of the claims 6 to 9, characterized in that the sexual disorder is Hypoactive Sexual Desire Disorder.

15

20