

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 May 2001 (31.05.2001)

(10) International Publication Number
WO 01/37630 A2

PCT

(51) International Patent Classification: Not classified

(21) International Application Number: PCT/US00/32112

(22) International Filing Date: 21 November 2000 (21.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 60/167,257 22 November 1999 (22.11.1999) US

(71) Applicant (for all designated States except US): EPIX MEDICAL, INC. [US/US]; 71 Rogers Street, Cambridge, MA 02142 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): WEISSKOFF, Robert, M. [US/US]; 18 Preston Road, Lexington, MA 02173 (US). KNIGHT, Stephen, C. [US/US]; 164 Lakeview Avenue, Cambridge, MA 02138 (US). CARTER, Wayne [US/US]; 9 Lake Place, Branford, CT 06405 (US).

(74) Agents: LIEBKE, Hope; Fish & Neave, 1251 Avenue of the Americas, New York, NY 10020 et al. (US).


(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With declaration under Article 17(2)(a); without classification and without abstract; title not checked by the International Searching Authority.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/37630 A2

(54) Title: IMAGING SEXUAL RESPONSE

(57) Abstract:

IMAGING SEXUAL RESPONSE

Technical Field of the Invention

The present invention relates to methods for contrast-enhanced imaging of sexual response. In 5 particular, the present invention relates to methods of magnetic resonance imaging (MRI), computerized X-ray tomography (CT), ultrasound, and optical imaging using contrast agents to measure blood flow during sexual response, especially female sexual response. The 10 invention relates to methods of analyzing normal sexual function and sexual dysfunction and provides a means for evaluating and screening potential therapeutic compounds for enhancing sexual function or treating sexual dysfunction.

15 Background of the Invention

Diagnostic imaging techniques, such as magnetic resonance imaging (MRI), X-ray, nuclear radiopharmaceutical imaging including PET and SPECT, ultraviolet/visible/infrared light, and ultrasound, 20 have been used in medical diagnosis for a number of years. In some cases, contrast media improves the image quality or provides specific additional useful information.

- 2 -

The contrast agent must interfere with the wavelength of electromagnetic radiation used in the imaging technique, alter the physical properties of tissue to yield an altered signal, or, as in the case 5 of radiopharmaceuticals, provide the source of radiation itself. Commonly used imaging materials include organic molecules, metal ions, salts or chelates, particles (particularly iron particles), or labeled peptides, proteins, polymers or liposomes. 10 After administration, the agent may non-specifically diffuse throughout body compartments prior to being metabolized and/or excreted; these agents are generally known as non-specific agents. Alternatively, the agent may have a specific affinity for a particular body 15 compartment, cell, organ, or tissue; these agents can be referred to as targeted agents.

In recent years, a number of contrast agents have been developed that may be used to enhance imaging of the blood pool. See, e.g., WO 96/23526, herein 20 incorporated by reference in its entirety. Such diagnostic imaging contrast agents may comprise an image-enhancing moiety, a plasma protein binding moiety and a blood half-life extending moiety and exhibit improved blood retention. Although a number of methods 25 using such contrast agents have been described, prior to this application it was not known if normal sexual function or sexual dysfunction could be measured with such types of agents.

The Diagnostic and Statistical Manual of 30 Mental Disorders, 4th edition, Washington DC, American Psychiatric Association, 1996 (DSM-IV, herein incorporated by reference) describes a number of sexual dysfunctions. These include sexual desire disorders such as hypoactive sexual desire disorder and sexual

- 3 -

aversion disorder; sexual arousal disorders such as female sexual arousal disorder and male erectile disorder; orgasmic disorders such as female orgasmic disorder (formerly, inhibited female orgasm), male 5 orgasmic disorder (formerly, inhibited male orgasm), and premature ejaculation, dyspareunia, vaginismus and sexual dysfunction not otherwise specified. The DSM-IV describes several subtypes that apply to primary sexual dysfunctions including lifelong type, acquired type, 10 generalized type, situational type, and sexual dysfunction due to psychological factors or due to combined factors. Sexual dysfunction may also be due to general medical condition or may be substance induced, such as by alcohol, antidepressants and 15 antihypertensives.

A number of compounds have been described to treat sexual dysfunction. For instance, ViagraTM (sildenafil citrate, Pfizer, Inc.) has been used to treat male erectile disorder. Physiologically, sexual 20 stimulation causes local release of nitric oxide (NO) in the corpus cavernosum. NO then activates the enzyme guanylate cyclase, which results in increased levels of cGMP, producing smooth muscle relaxation in the corpus cavernosum and allowing inflow of blood. Sildenafil 25 citrate enhances the effect of NO by specifically inhibiting PDE5, which is responsible for degradation of cGMP in the corpus cavernosum.

Female sexual arousal disorder is a poorly understood condition for which there is no reliable 30 diagnostic test. The underlying pathophysiology of this condition is thought to include an inadequate vascular arousal response to sexual stimulation. Vaginal photoplethysmography is the most widely

reported method for assessing the vascular component of the arousal response to sexual stimulation (by measuring a change in vaginal pulse amplitude). However, vaginal photoplethysmography has the 5 disadvantage of not providing an absolute measure of either vaginal blood volume or blood flow because all changes are observed without a quantifiable reference baseline. Furthermore, photoplethysmography results are highly variable and depend on exact placement of 10 the probe, among other parameters. Thus, there remains a need to develop better methods and techniques for measuring blood flow in sexual response, preferably female sexual response, and, more particularly, in female sexual arousal disorders.

15

Summary of the Invention

In one embodiment, the instant invention provides a method for contrast-enhanced diagnostic imaging of normal sexual function or sexual dysfunction. In a preferred embodiment, MRI, 20 ultrasound, computed X-ray tomography (CT), or optical imaging is used to image sexual dysfunction. In a more preferred embodiment, the sexual dysfunction is a female sexual arousal disorder.

In another embodiment, the instant invention 25 provides a method for measuring the therapeutic effect of a compound on sexual dysfunction. In a preferred embodiment, MRI or optical imaging is used to measure the effects of the therapeutic compounds. In another preferred embodiment, the sexual dysfunction treated is 30 female sexual arousal disorder. In an even more preferred embodiment, the compound tested is sildenafil citrate (ViagraTM).

- 5 -

In another embodiment, the instant invention provides a method for screening for potential therapeutic compounds for efficacy in the treatment of sexual dysfunction. In a preferred embodiment, the 5 compounds are screened for treating female sexual arousal disorder. In another preferred embodiment, MRI or optical imaging is used to screen the compounds.

Brief Description of the Drawings

Fig. 1 shows by MRI using contrast agent MS-10 325 a time course of normal female sexual arousal.

The top left-hand panel of Fig. 1 shows an MRI image before administration of MS-325.

The top middle and right-hand panel of Fig. 1 show MRI images during administration of MS-325 while 15 the female subject watches neutral video.

The bottom three panels of Fig. 1 show MRI images during administration of MS-325 while the female subject watches erotic video.

Fig. 2, left panel, shows the blood flow 20 image area ("Region of Interest" or ROI) that is quantitated (labeled with a box and number).

Fig. 2, right panel, shows a graph in which the signal intensity of the quantitated image area, as shown in Fig. 2, left panel, is plotted against time.

25 Detailed Description of the Invention

In order that the invention herein described may be more fully understood, the following detailed description is set forth.

A. Definitions

30 As used herein, the term "sexual dysfunction" encompasses those disorders including those defined by the DSM-IV, and includes, without limitation, sexual

desire disorders such as hypoactive sexual desire disorder and sexual aversion disorder; sexual arousal disorders such as female sexual arousal disorder and male erectile disorder; orgasmic disorders such as 5 female orgasmic disorder (formerly, inhibited female orgasm), male orgasmic disorder (formerly, inhibited male orgasm), and premature ejaculation, dyspareunia, vaginismus and sexual dysfunction not otherwise specified. In a preferred embodiment, the sexual 10 dysfunction is a sexual arousal disorder. In an even more preferred embodiment, the sexual dysfunction is female sexual arousal disorder. The etiology may be psychogenic, anatomical, or pathophysiological.

The term "diagnostic medical imaging", or 15 simply "medical imaging" or "diagnostic imaging" refers to a method of graphically or pictorially investigating an animal or human body for the purposes of studying the body's anatomy or physiology or an abnormality thereof. Generally, medical imaging is conducted by a 20 radiologist or someone under his or her supervision. Typical methods of medical imaging which are contemplated by the present invention include, among others, computerized X-ray tomography (CT), ultrasound, magnetic resonance imaging (MRI), and optical imaging.

25 As used herein, the term "contrast agent" comprises those agents that enhance image contrast during diagnostic imaging. In a preferred embodiment, a contrast agent is one that enhances contrast in MRI or optical imaging. In another preferred embodiment, 30 the contrast agent is one that enhances blood flow/blood pool imaging in MRI or optical imaging. In a more preferred embodiment, the contrast agent is one that exhibits improved blood retention and comprises an image-enhancing (or signal-generating) moiety (IEM), a

plasma protein binding moiety (PPBM) and a blood half-life extending moiety (BHEM). See, e.g., WO 96/23526, *supra*. In an even more preferred embodiment, the contrast agent is MS-325 (MS-325, EPIX Medical, Inc.);

5 see, e.g., WO 96/23526) or MultiHance™ (Bracco Pharmaceuticals, Inc.). Other MRI contrast agents that are suitable for use in the invention include Magnevist™ (Schering AG), ProHance™ (Bracco Pharmaceuticals, Inc.), Gadomer-17 (Schering AG),

10 B22956/1 (Bracco Pharmaceuticals, Inc.) and the contrast agents disclosed in US Patent 5,649,537, MP-2269 (Mallinckrodt, Inc.) and the contrast agents disclosed in international patent application (PCT) WO 98/20908, P760 and P775 (Guerbet SA), Clariscan™

15 (Nycomed Amersham), Combidex™ and Ferridex™ (Advanced Magnetics, Inc.), gadolinium texaphrin (Pharmacyclics, Inc.), and Eovist (Schering AG). Contrast agents for use in ultrasound imaging of sexual response include SHU-555c (Schering AG), AI-700 (Acusphere, Inc.), NC-

20 100100 (Nycomed Amersham), Sonovist™ (Schering AG), Optison™ and Albunex™ (Molecular Biosystems, Inc. , and Echogen™ (Sonus Pharmaceuticals, Inc.). Contrast agents for use in CT include iohexol, iopamidol, iopromide, iopentol, and ioxaglate. Contrast agents

25 that have affinities for endogenous blood components, especially human serum albumin, are particularly preferred. Contrast agents for use in optical imaging include indocyanine green and those disclosed in international patent application (PCT) WO 97/13490.

30 *B. Method for Measuring Physiological Sexual Response*

In one aspect, the present invention provides a method for accurately monitoring normal sexual function or sexual dysfunction in a subject. In

- 8 -

particular, the invention provides a diagnostic imaging method that involves the use of a contrast agent that measures one or more parameters of sexual function. The use of the contrast agents allow for "real-time" 5 monitoring and quantifying sexual function. In a preferred embodiment, the invention provides a diagnostic imaging method comprising the use of a contrast agent to measure and quantify female sexual arousal. The imaging methods useful in this invention 10 are MRI, including magnetic resonance spectroscopy techniques, ultrasound, computerized X-ray tomography (CT), and optical imaging.

The contrast agent is preferably a blood pool (also known as blood flow) contrast agent. 15 Furthermore, the contrast agent is preferably administered by intravenous or intra-arterial injection so that the amount of contrast agent in the blood is high relative to the amount of contrast agent in the interstitium or within cells. Contrast agents which 20 extravasate slowly, examples of which are MS-325, Gadomer-17, and P760, are preferred. Medical imaging and testing should be conducted shortly after administration of the contrast agent. As time progresses, the contrast agent may extravasate and 25 reduce the amount of contrast between blood and tissue, or it may be excreted. In the case of MS-325, imaging and testing may be conducted up to one hour after administration. When Magnevist™ is employed, the time window is 30 minutes.

30 The method for contrast-enhanced diagnostic imaging of normal sexual function or sexual dysfunction comprises the steps of:

(a) administering a contrast agent to a subject;

- 9 -

(b) subjecting the subject to diagnostic medical imaging complementary to said contrast agent's intended use;

(c) measuring an imaging signal
5 characteristic of the contrast agent;
(d) subjecting the subject to a stimulus; and
(e) monitoring an imaging signal
characteristic of the contrast agent to assess the sexual function after stimulus.

10 The steps (b) through (e) of the method may be repeated several times if sufficient contrast agent is still present in the blood of the body. The contrast agent may be re-administered, i.e. step (a) may be repeated, if the method is to be repeated, and
15 the contrast agent's effect has diminished substantially. One skilled in the art will readily recognize if enough contrast agent is circulating within the body and if more should be administered.

Several regions of the body may be imaged,
20 including the head, legs, pelvic area, and torso. The region preferably comprises the brain, spine, central nervous system (CNS) or other nervous system of the subject, and more preferably the pelvis or genitalia of the subject.

25 In a preferred embodiment, the contrast agent is a blood pool or blood flow contrast agent. In a more preferred embodiment, the contrast agent is one capable of enhancing blood flow or blood pool contrast in MRI or optical imaging. Suitable blood pool
30 contrast agents include those described in WO 96/23526.

In a preferred embodiment, the sexual function that is assessed is sexual arousal and the stimulus is one designed to cause sexual arousal, including somatosensory, audiovisual, physical, and

- 10 -

pharmacological stimuli. In a more preferred embodiment, the stimulus is one that causes sexual arousal in women without sexual dysfunction.

In one preferred embodiment of the method of measuring sexual response in women, the relative size of the clitoris is determined with and without stimulation. In this embodiment, the diagnostic imaging method is MRI and the size of the clitoris is measured by counting the number of pixels comprising the clitoris in steps (c) and (e). An increase in the number of pixels is equivalent to an increase in the size of the clitoris where all other MRI imaging and data processing parameters are the same. An increase in the size of clitoris is indicative of sexual arousal. The relative size of the clitoris in steps (c) and (e) of the method may also be determined by CT by counting the number of voxels comprising the clitoris.

In an alternative embodiment, the sexual response of women is determined by measuring the relative increase in blood volume to the clitoris. In the above method, the patient is administered MS-325 in step (a). In step (c), a region of interest (ROI) within the clitoris is selected and the signal measured prior to administration of contrast agent. A region of interest within an easily imaged major blood vessel is also selected as an internal standard against which blood flow is calculated. The femoral artery or vein is most preferred. The MRI signal in this artery is also determined before contrast agent administration. Contrast agent is administered at time equals zero and signal intensity data from the two ROIs are collected regularly for approximately five to ten minutes during which time the subject is not exposed to the sexual

- 11 -

stimulus. Stimulation is then applied, and image data acquisition from the two ROIs is continued. The data from the two ROIs are used to calculate a relative blood volume within the clitoral ROI according to the 5 following formula:

$$BV(t) = [S_c(t) - S_c(0)] / [S_c(t) - S_c(0)],$$

where t is time, $BV(t)$ is the relative blood volume within the clitoral ROI at time t , $S_c(t)$ and $S_c(0)$ are the average MRI signals within the clitoral ROI at time 10 t and immediately before contrast agent administration respectively, and $S_r(t)$ and $S_r(0)$ are the average MRI signals within the reference artery ROI at time t and immediately before contrast agent administration respectively. The $BV(t)$ data are then fit to the 15 following curve:

$$BV(t) = C1 + C2 \cdot t + C3 \cdot E(t),$$

where $C1$, $C2$, and $C3$ are fitted constants and $E(t)$ is a function that equals zero for times, t , when sexual stimulus is absent, and equals unity when sexual 20 stimulus is applied. The ratio $C3/C1$ is an indicator of increase in relative blood volume upon stimulation. A statistical test, regression for example, may be conducted to reject the null hypothesis, $C3=zero$, within desired confidence limits, which indicates a 25 complete lack of increase in blood flow, and therefore any physiological sexual response, upon stimulation. $C2$ is relative to the rate of extravasation of the contrast agent during the examination, and can be used to correct for extravasation of the agent if such 30 occurs. The contrast agent MS-325 has a stable blood concentration over typical stimulation periods and is therefore a preferred contrast agent in this embodiment.

- 12 -

In another embodiment of this invention, the method is used to quantify normal sexual function in healthy human adults. In another embodiment of this invention, the method is used to measure sexual 5 dysfunction. In a preferred embodiment of this invention, the technique is used to quantify the arousal response in healthy female volunteers. In another preferred embodiment, the technique is used to diagnose subjects with female sexual arousal disorder.

10 The invention may be practiced using subjects of either sex and of any age. The invention may be used to determine normal sexual function or may be used to study or diagnose sexual dysfunction by comparison to the mean of quantified arousal of normal subjects. 15 Subjects are chosen depending upon the parameters of a particular study or clinical dysfunction presented. In an alternative embodiment, the invention may be practiced using animals to determine normal sexual function or sexual dysfunction, see *infra*.

20 For studies of female sexual arousal, subjects may include both pre-menopausal and post-menopausal women. Post-menopausal women may include those who use estrogen-replacement therapy and those who do not. Subjects may include those who have had 25 disease or who have undergone surgery or trauma to any portion of the female reproductive system or genitalia; including those who have or who have had any type of sexually transmitted diseases (STDs). Such diseases include, without limitation, bacterial, viral or fungal infections; for example, Chlamydia, HIV, herpes, gonorrhea, syphilis, or other diseases affecting the reproductive organs. Other such diseases include, without limitation, endometriosis, diabetes, hypertension, endocrine disorders, or cancer of any of 30

- 13 -

the reproductive organs. Surgery and trauma includes, without limitation, hysterectomy, ovariectomy, and surgery for removal of fibroid tumors or prolapsed uterus.

5 *C. Therapy Assessment Methods*

In another aspect, the instant invention provides a method for measuring the therapeutic effect of a compound on sexual dysfunction. The method comprises the steps of:

- 10 (a) administering a contrast agent to a subject;
- (b) subjecting the subject to diagnostic medical imaging complementary to said contrast agent's intended use;
- 15 (c) measuring an imaging signal characteristic of the contrast agent to establish a baseline;
- (d) subjecting the subject to a stimulus that will assess the sexual function;
- 20 (e) monitoring an imaging signal characteristic of the contrast agent to assess the sexual function after stimulus;
- (f) administering an effective amount of a compound expected to provide a therapeutic effect on
- 25 the sexual function; and
- (g) determining if the sexual function has been enhanced as a result of treatment with the compound.

Comparing the imaging data in the absence and 30 in the presence of a candidate therapeutic compound gives an indication of its efficacy. In the above method, any of the steps may be repeated using the same subject. In a particularly preferred embodiment, this

- 14 -

method is practiced on a population of subjects with a reproducible and predictable induced sexual impairment.

In a preferred embodiment, the contrast agent is a blood pool or blood flow contrast agent. In a 5 more preferred embodiment, the contrast agent is one capable of enhancing blood flow or blood pool contrast in MRI or optical imaging. Suitable blood pool contrast agents include those described in WO 96/23526.

In a preferred embodiment, the sexual 10 function that is assessed is sexual arousal and the stimulus is one designed to cause sexual arousal. In a more preferred embodiment, the stimulus is one that causes sexual arousal in women.

In a preferred embodiment, the therapeutic 15 compound is one that enhances sexual arousal in a woman with female sexual arousal disorder. In a preferred embodiment, the therapeutic compound is one used to treat sexual arousal disorders in either men or women.

The therapeutic compound may be sildenafil citrate 20 (Viagra™) or other compounds used to treat male erectile dysfunction. Other drugs in development for the treatment of sexual dysfunction include Cialis™ (Eli Lilly and ICOS Corp.), Befar™ (NexMed), Vardenafil™ (Bayer), apomorphine (TAP Pharmaceuticals), 25 Alprostadil™ (MacroChem), desmethylsibutramine (Sepracor), PT 14 and PT 141 (Palatin Technologies), SC-113 (Scotia Holdings), PNU-83757 (Pharmacia), NMI-187 (NitroMed), liposomal alprostadil (BioSphere and Harvard Scientific Corp.), and Amesergide™ (Eli Lilly).

30 *D. Screening Methods for New Therapeutic Compounds*

In another aspect, the instant invention provides a method for screening potential compounds to

- 15 -

treat sexual dysfunction. The method comprises the steps of:

(a) administering a contrast agent to a non-human or human animal or animal subject;

5 (b) subjecting the subject to diagnostic medical imaging complementary to said contrast agent's intended use;

(c) measuring an imaging signal characteristic of the contrast agent to establish a 10 baseline;

(d) subjecting the subject to a stimulus that will assess the sexual function;

15 (e) monitoring an imaging signal characteristic of the contrast agent to assess the sexual function after stimulus;

(f) administering a compound that is being tested to determine if it has an effect on the sexual function; and

20 (g) determining if the sexual function has been modified as a result of treatment with the compound by comparing the results of steps (a)-(e) with those of step (f).

In a preferred embodiment, the contrast agent is a blood pool or blood flow contrast agent. In a 25 more preferred embodiment, the contrast agent is one capable of enhancing blood flow or blood pool contrast in MRI or optical imaging. Suitable blood pool contrast agents include those described in WO 96/23526.

30 In a preferred embodiment, the sexual function that is assessed is sexual arousal and the stimulus is one designed to cause sexual arousal.

For non-human animal subjects, the stimulus may be any type of signal that elicits a sexual response. These stimuli may include auditory signals,

- 16 -

e.g., recorded or live vocalizations; olfactory signals, e.g., pheromones or other distinctive signaling odors; sensory cues, such as genital manipulation including masturbation, or visual cues.

5 Stimuli that elicit a sexual response are known in the art and may be adapted for use in this method.

In a preferred embodiment of this method, compounds are tested first for safety and efficacy in non-human animal subjects. After these preliminary 10 studies, compounds that appear to be safe and efficacious in non-human animals are tested for safety and efficacy in human subjects. In another preferred embodiment, compounds that have been approved for use in humans for other indications may be tested using the 15 method of this invention to determine whether they may be used to treat sexual dysfunction or to enhance sexual function.

E. Contrast Agent Dosages and Imaging

A number of different contrast agents may be 20 used according to this invention. In general, the contrast agents will be water soluble. The contrast agents may also comprise a pharmaceutically acceptable salt. Examples of such salts are provided in WO 96/22526. Contrast agents may be administered by any 25 method known in the art, such as those disclosed in WO 96/22526. Contrast agents may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir in dosage formulations containing 30 conventional non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles. As discussed above, in a preferred embodiment, the contrast agent is administered intravenously.

- 17 -

Dosage depends on the sensitivity of the diagnostic imaging instrumentation, as well as the composition of the contrast agent. For example, for MRI imaging, a contrast agent containing a highly paramagnetic substance, e.g., gadolinium (III), generally requires a lower dosage than a contrast agent containing a paramagnetic substance with a lower magnetic moment, e.g., iron (III). Preferably, dosage will be in the range of about 0.001 to 1 mmol/kg body weight per day of the active metal-ligand complex. More preferably, dosage will be in the range of about 0.005 and about 0.2 mmol/kg body weight per day. In a more preferred embodiment, the dosage will be 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09 or 0.10 mmol/kg body weight per day administered intravenously.

Those skilled in the art will readily determine an appropriate dosage based on the recommendation of the contrast agent manufacturer or supplier and the empirical ability to undertake the methods described herein. It should be understood, however, that a specific dosage regimen for any particular subject will also depend upon a variety of factors, including the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician.

If the application of this invention is MR imaging, following administration of the appropriate dosage of the contrast agent, MR imaging is carried out. The choice of pulse sequence (inversion recovery, IR; spin echo, SE, echo planar, EPI; time-of-flight, TOF; turbo-flash; gradient echo, GE) and the values of the imaging parameters (echo time, TE; inversion time, TI; repetition time, TR; flip angle, etc.) will be

- 18 -

governed by the diagnostic information sought. In general, if one desires to obtain T_2 -weighted images, then TE should be minimized to maximize T_2 -weighting. Conversely, if one desires to measure T_1 , then TE should 5 be greater than 30 milliseconds and the TR maximized or flip angle minimized to minimize competing T_1 effects. TI and TR are generally on the order of about 5-1000 and 2-1000 milliseconds, respectively.

In order that this invention may be more 10 fully understood, the following examples are set forth. These examples are for the purpose of illustration only and are not to be construed as limiting the scope of the invention in any way.

EXAMPLE 1

15 Imaging of the genitalia was performed in four pre-menopausal and four post-menopausal healthy, sexually functional women subjects after administration of MS-325 (MS-325, EPIX Medical, Inc.). All volunteers signed an IRB-approved informed consent form for the 20 study.

Six of the subjects received an intravenous injection of 0.05 mmol/kg of MS-325 and two received an intravenous injection of 0.02 mmol/kg of MS-325. Dynamic MRI (3D SPGR, 11/1.7/35°) of the perineum was 25 performed on all subjects while they viewed videos containing first neutral material, then erotic material and finally neutral material via a fiberoptic audiovideo system. Each segment of neutral or erotic material was up to 15 minutes in length. Subjects also 30 filled out questionnaires in response to the videos

- 19 -

that had them subjectively rate their degree of sexual arousal.

Serial imaging was performed at least every 3 minutes, beginning just prior to contrast injection and 5 continuing up to 45 minutes post injection. Signal intensity measurements over time were made of the vaginal wall and mucosa, clitoris, skeletal muscle, and femoral artery. Clitoral volume was measured at each time point. Relative regional blood volume was 10 determined from signal intensity versus time curves and compared with subjective measures of arousal.

All subjects tolerated the procedure and all reported sexual arousal on subjective questionnaires.

Baseline (unstimulated) MS-325 images revealed strong 15 enhancement of the genitalia. When compared to a neutral stimulus, erotic stimulus produced significant increases in image enhancement, increases in clitoral blood volume ($43\% \pm 5\%$) and in clitoral size ($85\% \pm 5\%$). See Figs. 1 and 2. In four subjects studied up 20 to 10 minutes following termination of the erotic stimulus, these changes decreased, but did not return to baseline levels during the observation period.

In these studies, MRI was compared with vaginal photoplethysmography to confirm that MRI 25 accurately assessed the vascular component of the arousal response.

EXAMPLE 2

Imaging of genitalia is performed on female subjects who have or who are suspected of having sexual 30 arousal dysfunction. The subjects receive an intravenous injection of 0.01 to 0.1 mmol/kg of MS-325. Dynamic MRI is performed as described above while they are exposed to neutral material and then erotic

- 20 -

material. In one embodiment, the subjects view videos containing first neutral material, then erotic material and finally neutral material via a fiberoptic audiovideo system. Subjects may fill out 5 questionnaires in response to the erotic material to allow them to subjectively rate their degree of sexual arousal.

Serial imaging is performed as described above. Signal intensity measurements over time are 10 made of the vaginal wall and mucosa, clitoris, skeletal muscle, and femoral vein. Clitoral volume is measured at each time point. Relative regional blood volume is determined from signal intensity versus time curves and compared with subjective measures of arousal.

15 The clitoral volume and size of the female subjects may be compared to those of sexually functional women who have been similarly tested. For instance, mean values can be determined to compare rates of blood flow or changes in clitoral volume 20 between sexually functional subjects and those suspected of having sexual dysfunction. A lack of change in clitoral volume and size in the female subjects may be indicative of sexual arousal dysfunction.

25

EXAMPLE 3

Imaging is performed as described in Example 2. After a diagnosis of sexual arousal dysfunction has been made in a subject or group of subjects, a therapeutic compound for sexual arousal disorder is 30 administered. The therapeutic compound may be one that is known to treat sexual arousal dysfunction or may be one that is being screened for its ability to treat sexual arousal disorder. In one embodiment, the

- 21 -

therapeutic compound is Viagra™. After treatment with the therapeutic compound, imaging is performed again as described in Example 2. By comparing the change in clitoral blood volume and size before treatment to that 5 after treatment, one may determine if the therapeutic compound is effective for treating sexual arousal disorder in a particular subject or group of subjects.

All publications and patent applications cited in this specification are herein incorporated by 10 reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and example for purposes of 15 clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

- 22 -

CLAIMS

We claim:

1. A method for contrast-enhanced diagnostic imaging of normal sexual function or sexual dysfunction
5 comprising the steps of:

- (a) administering a contrast agent to a subject;
- (b) subjecting the subject to diagnostic medical imaging complementary to said contrast agent's 10 intended use;
- (c) measuring an imaging signal characteristic of the contrast agent to establish a baseline;
- (d) subjecting the subject to a stimulus; and
- 15 (e) monitoring an imaging signal characteristic of the contrast agent to assess the sexual function after stimulus.

2. The method according to claim 1, wherein the contrast agent enhances blood flow or blood pool 20 contrast in MRI, CT, ultrasound, or optical imaging.

3. The method according to claim 2, wherein the contrast agent is MS-325.

4. A method for measuring the therapeutic effect of a compound on sexual dysfunction comprising 25 the steps of:

- (a) administering a contrast agent to a subject;
- (b) subjecting the subject to diagnostic medical imaging complementary to said contrast agent's 30 intended use;

- 23 -

- (c) measuring an imaging signal characteristic of the contrast agent to establish a baseline;
- (d) subjecting the subject to a stimulus that 5 will assess the sexual function;
- (e) monitoring an imaging signal characteristic of the contrast agent to assess the sexual function after stimulus;
- (f) administering an effective amount of a 10 compound expected to provide a therapeutic effect on the sexual function; and
- (g) determining if the sexual function has been enhanced as a result of treatment.

5. The method according to claim 4, wherein 15 the contrast agent is MS-325.

6. The method according to claim 5, wherein the compound is Viagra™.

7. A method for screening potential compounds to treat sexual dysfunction comprising the 20 steps of:

- (a) administering a contrast agent to a non-human animal or animal;
- (b) subjecting the subject to diagnostic medical imaging complementary to said contrast agent's 25 intended use;
- (c) measuring an imaging signal characteristic of the contrast agent to establish a baseline;
- (d) subjecting the subject to a stimulus;

- 24 -

(e) monitoring an imaging signal characteristic of the contrast agent to assess the sexual function after stimulus;

5 (f) administering a compound that is being tested to determine if it has an effect on the sexual function;

(g) determining if the sexual function has been modified as a result of treatment with the compounda.

10 6. The method according to claim 7, wherein the contrast agent is MS-325.




FIGURE 2

