



US 20090011462A1

(19) United States

(12) Patent Application Publication
Vind et al.(10) Pub. No.: US 2009/0011462 A1
(43) Pub. Date: Jan. 8, 2009

(54) POLYPEPTIDES HAVING LIPASE ACTIVITY AND POLYNUCLEOTIDES ENCODING SAME

(75) Inventors: Jesper Vind, Værloese (DK); Kim Borch, Birkeroed (DK); Mikael Mikkelsen, Smorum (DK); Jurgen Carsten Franz, Knotzel (DK)

Correspondence Address:
NOVOZYMES NORTH AMERICA, INC.
500 FIFTH AVENUE, SUITE 1600
NEW YORK, NY 10110 (US)

(73) Assignee: Novozymes A/S, Bagsvaerd (DK)

(21) Appl. No.: 12/161,083

(22) PCT Filed: Jan. 22, 2007

(86) PCT No.: PCT/US07/60825

§ 371 (c)(1),
(2), (4) Date: Aug. 14, 2008

ID NO 1: SSSSTODYRLRSEAEIKANTPTYTLASNA
ID NO 2: SSSSTODYRLRSEAEIKANTPTYTLASNA
ID NO 3: SSSDGKRAATSSQELRLTPTYTLASNA
ID NO 4: SSSDGKKAUATTAAQIQTPTYKGTAAIA
ID NO 5: TAGHIALAASITQ GISESDLYSRL VEMATISQAA
ID NO 6: TAGHIALAASITQ GISESDLYSRL VEMATISQAA
ID NO 7: AVGWTTTDFSSNEFKYIQLQADAA
ID NO 8: TTTTDTDFSSNEFKYIQLQADAA
ID NO 9: DPTTQLEDEKEWVQYAAAT
ID NO 10: DWSITSELDQSEFSSWQYAAAS
ID NO 11: SVSITSLDELQLEQASNSAA
ID NO 12: SVSITSLDELQLEQASNSAA
ID NO 13: DVSSQDLLNNLDFQASNSAA
ID NO 14: EVSQDLPNQFRLFAQYSAAS
ID NO 15: FQDAYTASHADLIVYKATAGCA

ID NO 1: YCRTVIFG GRASCPHQGAS NLQITKTTST LIDTDTNVLVAV
ID NO 2: YCRTVIFG QGWSCEMCDVAP NLNATKTTST LIDTDTNVLVAV
ID NO 3: YCRTVIFG ATWDCLHEDATE DLKLIKTTST LIDTDTNVLVAV
ID NO 4: YCRSVUGS NKUDCVCQCCQKHWL DKLKTTTFSL LLSDTMVGWLR
ID NO 5: YDADLNSPST LIXKGBKTTYNSQTDINGWILR
ID NO 6: YDADLNSPST LIXKGBKTTYNSQTDINGWILR
ID NO 7: MC NSEAAA GKKITGSNNNGCETVCGNGATATVTFV VGSKTSIGGGVAT
ID NO 8: YG NFTPTAV GKPVHGSAGNCSDIEKDAIRVGSV VGTKTGIGAYVAT
ID NO 9: YCPNNVYAKD GKLKNSCVNCSDVDERAAGTCKLSPS DDTTTTGTAGFVAV
ID NO 10: YSEADYTAQD GDLKLNSKGNCPBVEATGATVSYDFS DTTTTTGTAGFVAV
ID NO 11: YCSNNNID SD DSENNTCTANACCEBSEVEBAS TTNLLEFDLTDNGGTAGFLA
ID NO 12: YCSNNNID SD DSENNTCTADACEBSEVEBAS TTNLLEFDLTDNGGTAGFLA
ID NO 13: YCENRNL ST GTKLKNSCVNCSDVLEAAGTCSLDENESSYNGTAGFLA
ID NO 14: YCCKNNNAPD GTNTTCTGNAECPSEVNGADTFLYSSPF DSGVGDVGTGFLA
ID NO 15: YQTDIANWAS RTVEQDFTLSSPFW DTLKGSSGYIAF

ID NO 1: GEKEKRTIYVV FRGTSIRNA IADIPVVEVN YPPV NGA KVKGCFDLSW
ID NO 2: GEKEKRTIYVV FRGTSIRNA IADIPVVEVN YPPV NGA KVKGCFDLSY
ID NO 3: GEKEKRTIYVV FRGTSIRNA IADIPVVEVN YPPV SGT KVKGCFDLSY
ID NO 4: SDKRTKTYVV FRGTSIRNA DEDIVEDNSD YPPV NGA KVKGCFDLSY
ID NO 5: DDKSKRTIYVV FRGTSIRNA QDIPVVEVN YPPV NGA KVKGCFDLSY
ID NO 6: DDKSKRTIYVV FRGTSIRNA QDIPVVEVN YPPV NGA KVKGCFDLSY
ID NO 7: DSKARKEIIVV FRGTSIRNA LCTDIPVVEVN YPPV NGA KVKGCFDLSY
ID NO 8: DPKRKEIIVV FRGTSIRNA LCTDIPVVEVN YPPV NGA KVKGCFDLSY
ID NO 9: DPKRKEIIVV FRGTSIRNA LCTDIPVVEVN YPPV NGA KVKGCFDLSY
ID NO 10: DPKRKEIIVV FRGTSIRNA LCTDIPVVEVN YPPV NGA KVKGCFDLSY
ID NO 11: DPKRKEIIVV FRGTSIRNA LCTDIPVVEVN YPPV NGA KVKGCFDLSY
ID NO 12: DPKRKEIIVV FRGTSIRNA LCTDIPVVEVN YPPV NGA KVKGCFDLSY
ID NO 13: DPKRKEIIVV FRGTSIRNA LCTDIPVVEVN YPPV NGA KVKGCFDLSY
ID NO 14: DPKRKEIIVV FRGTSIRNA LCTDIPVVEVN YPPV NGA KVKGCFDLSY
ID NO 15: DPKRKEIIVV FRGTSIRNA LCTDIPVVEVN YPPV NGA KVKGCFDLSY

ID NO 1: NIVQDQKLVAE VKAQDRLRPG YKIVVYGHSL GGATAVSLCALDLYHNGHA
ID NO 2: NIVQDQKLVAE VKAQDRLRPG YKIVVYGHSL GGATAVSLCALDLYHNGHA
ID NO 3: GEVONELVAT VLDQEQKOPYR YKIVVYGHSL GGATAVSLCALDLYGREPGL
ID NO 4: GEVONELVAT VLDQEQKOPYR YKIVVYGHSL GGATAVSLCALDLYGREPGL
ID NO 5: YEVQDQVSEL VKGQVSYQPD YALIVVYGHSL GASLAALTAQQLSATYD
ID NO 6: YEVQDQVSEL VKGQVSYQPD YALIVVYGHSL GASLAALTAQQLSATYD
ID NO 7: NEISQSQAA VASARKANPS ENVIESTYGHSL GGAVAVLAAANLRVCG
ID NO 8: ENVAANVKA VSAAKTANPT FKFVYVGHSL GGAVAVTAAAYLRDQHG
ID NO 9: KVVEDRDIKNT LOELKEPHSD YKIVVYGHSL GAAATASLAADLRTKNN
ID NO 10: KLVREDRDIKNT LOELKEPHSD YKIVVYGHSL GAAATASLAADLRTKNN
ID NO 11: ESDARDEYIISK IKSAMSTYSC YTLYFTFORSL GGALATLGATVLRNEDY
ID NO 12: ESDARDEYIISK IKSAMSTYSC YTLYFTFORSL GGALATLGATVLRNEDY
ID NO 13: SEIADTITSK VESALDSHED YSLVLAGHNS GAAALALATAALRNSGH

Related U.S. Application Data

(60) Provisional application No. 60/761,106, filed on Jan. 23, 2006.

Publication Classification

(51) Int. Cl.
C12N 9/20 (2006.01)
C12P 21/02 (2006.01)
C12N 15/55 (2006.01)
C12N 15/74 (2006.01)
C12N 15/80 (2006.01)
C12N 1/00 (2006.01)

(52) U.S. Cl. 435/69.1; 435/198; 536/23.2; 435/320.1;
435/243

ABSTRACT

The present invention relates to polypeptide having lipase activity and which further has a RP of at least 0.8 and a BR of at least 1.1 at the test conditions given in the specification.

ID NO 14: RSVADTLRQK VEDAVRSHPD YEVVFTQHNSL GGALATVAGDLRGMG
ID NO 15: LVSKQQVQEA VDSLLAKCPD ATISFFQHSL GGALACISMVDTAQRHNGI

ID NO 1: NIEIYTOG QPRIGTAPAS NYVIGT KTSYQRQLVHEDIVPHL
ID NO 2: NIEIYTOG QPRIGTAPAS NYVIGT KTSYQRQLVHEDIVPHL
ID NO 3: SSNLSTLFTQG QPRNGDPAF NVVST GIVFRTVHEDIVPHL
ID NO 4: SSNLSTLFTQG QPRNGDPAF NVVST GIVFRTVHEDIVPHL
ID NO 5: NIRLYTFC EPRIS NQAFR SYMNDIAFQASSFDTTQYFATVTHDQFPL
ID NO 6: NIRLYTFC EPRIS NQAFR SYMNDIAFQASSFDTTQYFATVTHDQFPL
ID NO 7: PUDVTTYQ SPRVNGAQLS AVFENQ AGTATVHEDIVPHL
ID NO 8: PUDVTTYQ SPRVNGAQLS AVFENQ AGTATVHEDIVPHL
ID NO 9: DAILYNAA APRVANKPLA EPTINO GNTYRFTMNEBVPVPL
ID NO 10: SAKRIVYA SPRVNGAQLA KITVQO GMMFRTHTHEDIVPHL
ID NO 11: SVELTIVYQ SPRVNGAQLA EPTINO GSCAMPFRVTHLNDIVPHV
ID NO 12: SVELTIVYQ SPRVNGAQLA EPTINO GSCAMPFPVTHLNDIVPHV
ID NO 13: SVELTIVYQ SPRVNGAQLA EPTINO NGKGNVYRVTNTDIVPHL
ID NO 14: DIDVFTSIC APRVNGAQLA EPTIVO TGDTLYA LTHNTDIVPHL
ID NO 15: KMMFPTQG QRTQNGQPA EYVENVL GHVFRVAVVYRHDIVPHM

ID NO 1: PPGAFQFPLA GREEPMWKK DSSLRVCNPNGIETDNCNSNTV
ID NO 2: PPGAFQFPLA GREEPMWKK DSSLRVCNPNGIETDNCNSNTV
ID NO 3: PPAAFQFPLA GREEWVITD NSPBTQVCTSDLETSDCNSNTV
ID NO 4: PPGAFQFPLA GREEWVITD GTSNVQICTEIETKDCNSNTV
ID NO 5: PPEVQCYANG VYETVWSV DPPSQAQTFVCTGDEVOCCCE AQGQGQ
ID NO 6: PPEVQCYANG VYETVWSV DPPSQAQTFVCTGDEVOCCCE AQGQGQ
ID NO 7: PPEVQCYANG VYETVWSV TPEFWLSCGGGDXWVDTTISDN KVCEGAAANLQ CRGGT1
ID NO 8: PPEVQCYANG VYETVWSV TPEFWLSCGGGDXWVDTTISDN KVCEGAAANLQ CRGGT1
ID NO 9: PPLTNGWYHII SPEYYVTA EINNTTVDNQVTVLVDGCVAFV CRGGT1
ID NO 10: PPLTNGWYHII SPEYYVTA EINNTTVDNQVTVLVDGCVAFV CRGGT1
ID NO 11: PPMDFQFSCQ SPEWVITS GNGASVIAASDIEVIEGELNSTA CRGGT1
ID NO 12: PPMDFQFSCQ SPEWVITS GNGASVIAASDIEVIEGELNSTA CRGGT1
ID NO 13: PPTILQYHNF SPEYYVTS ADEATVTTIDTVEVHINADATO CRGGT1
ID NO 14: PPTILQYHNF SPEYYVTS GTLVFPVTRNDIVKLEKIDATO GNNCPN
ID NO 15: PPMDFQFSCQ SPEWVITS DENIKSCXEGHNLTCELGPV

ID NO 1: PFT SVHDLNLSVHDMTGL CL
ID NO 2: PFT SVHDLNLSVHDMTGL CL
ID NO 3: PFT SVHDLNLSVHDMTGL CL
ID NO 4: PFT SVHDLNLSVHDMTGL CL
ID NO 5: VN WNRHTTYC GMTCGACTW
ID NO 6: VN WNRHTTYC GMTCGACTW
ID NO 7: GL DIANLHYY QATDA CHAGGFSWR R
ID NO 8: GL DIANLHYY QATDA CAPTAIPW K R
ID NO 9: GOLBLLAQHSHVYKIIHADACKGQPLPLR
ID NO 10: LELBLAIIWYF VQDVA GKGQPLPFK R
ID NO 11: TV SVLHLWYF FAISE CLL
ID NO 12: TV SVLHLWYF FAISE CLL
ID NO 13: GT SIDAHNRW YIYIE CS
ID NO 14: IP DIPAHRLWYF GLIOT CL
ID NO 15: FSEI NAKCHSEYD GMW

ID NO:	Micro organism	SEQ ID NO.:
1.	Absidia reflexa	3
2.	Absidia corymbifera	4
3.	Rhizomucor miehei	5
4.	Rhizopus delemar (oryzae)	6
5.	Aspergillus niger	7
6.	Aspergillus tubingensis	8
7.	Fusarium oxysporum	9
8.	Fusarium heterosporum	10
9.	Aspergillus oryzae	11
10.	Penicillium camemberti	12

ID NO 1:	SSSSTQDYIASEAEIKAFTFTYTLSANA				
ID NO 2:	SSSTQDYIASEAEIKAFTFTYTLSANA				
ID NO 3:	SIDGGIRAATSQEINELTYTTLANS				
ID NO 4:	SASDGCKVVAATTQIQEFTKYAGIATA				
ID NO 5:	TAGRALAASQ CISEDLYSRL VEMATISQAA				
ID NO 6:	TAGHALAASQ CISEDLYSRL VEMATISQAA				
ID NO 7:	AVGTTTDFSNFKFYIONGAAA				
ID NO 8:	TVTTQDLSNFRFYILQHADAA				
ID NO 9:	DIPTTQLEDKFFWVQYAAAT				
ID NO 10:	DVSTISELDQFEPWVQYAAAS				
ID NO 11:	SVSTSTLDELQLFQWSAAA				
ID NO 12:	SVSTSTLDELQLFQWSAAA				
ID NO 13:	DVSSSLNNNNLDFQYSSAAA				
ID NO 14:	BVSQDLFNQFNLFQYSSAAA				
ID NO 15:	PODAYTASHALLVKYATYAGLA				
ID NO 1:	YCRTVPG	GRWSCPNCVGAS	NLOITKTFST	LITDTNVLVAV	
ID NO 2:	YCRTVPG	GOWSCPNCVDAP	NLNITKTFPT	LITDTNVLVAV	
ID NO 3:	YCRTVPG	ATWDQIHCDATE	DLKIIKWTST	LIDDTNANVAR	
ID NO 4:	YCRSVVPG	NKWDXCVQQQKWNVP	DGKLIITPTTS	LSDTNGYVLR	
ID NO 5:	YADLCNIPST			IIKGKEIYNSQTDINGWILR	
ID NO 6:	YADLCNIPST			IIKGKEIYNSQTDINGWILR	
ID NO 7:	YC NSEAAA	GSKITCSNNNCPTVGNGATIVTSE	VGSXTGIGGYVAT		
ID NO 8:	YC NFNTAV	GKPVHCSAGNCVDIEKDAAIIVVGSV	VGTTGIGAYVAT		
ID NO 9:	YCPNNYVAKD	GEKLNCVGNCVDVEAAGSTVILSFS	DDTITDAGFVAV		
ID NO 10:	YYEADYTAOV	GDKLSCSKGNCPPEVATGATVSYDFS	DSTITDTAGYIAV		
ID NO 11:	YCSNNID SK	DSNLTCTANACPSVEAESTMLLEFDL	TNDFCGTAGFLAA		
ID NO 12:	YCSNNID SD	DSNVTCTADACPSVEAESTMLLEFDL	TNNFGGTAGFLAA		
ID NO 13:	YCDENLN ST	GTKLTCVGNCPLVEAASTOSLDRTNESSYYGPNAGYLA			
ID NO 14:	YCGKNNIDPAA	GTMNITCTGNACPEVEKADATFLYSCFE	DSCVGDVTGFLA		
ID NO 15:	YQTTDAWPAS		RTVPKDTTLISSFD	HTLGKSSGYIAF	
ID NO 1:	GEKEKTIIVVV	FRGTSSIRNA	IADIVFVSVN	YPPV	NGA KVHKGFLLDSY
ID NO 2:	GENEKTIVVV	FRGTSSIRNA	IADIVFVSVN	YPPV	NGA KVHKGFLLDSY
ID NO 3:	GDSEKTIIVIV	FRGSSSIERNW	IADLTFTVSVS	YPPV	SGT KVHKGFLLDSY
ID NO 4:	SDKQKPIVVLV	FRGTNSFRSA	ITDIVEFNSD	YKPV	KGA KVHAGFLSSY
ID NO 5:	DDSSKEIIITV	FRGTGSDTINL	QLDTINYTLTP	FDTLPQCNNGC	EVHGGYYIGW
ID NO 6:	DDGSKKEIIITV	FRGTGSDTINL	QLDTINYTLTP	FDTLPQCNSC	EVHGGYYIGW
ID NO 7:	DSARKEIIVVS	FRGSINIRNW	LTNLDPEG	QR DCSL	VSGC CVMSGFQRAN
ID NO 8:	DNARKEIIVVS	VRGSINVRNW	ITDNENFG	QK TCDSL	VAGC CVNTGFLLDAN
ID NO 9:	DNTNKAIVVA	FRGSYSIRNW	VTDNATPF	QT DPGC	CGGC KABLGFWWTAW
ID NO 10:	DNTNSAVVLA	FRGSYSVRNW	VADATEV	HT NFGL	CGGC LAGLGFWSESW
ID NO 11:	DNTNKRLLVVA	FRGSSTLLENW	IANLDPILED	NDL CTGC	KVHTGFWICAN
ID NO 12:	DNTNKRLLVVA	FRGSSTLLENW	IADLDPILOD	NDL CTGC	KVHTGFWKAN
ID NO 13:	DNTNKLLLVLS	FRGSADLANW	VANLNFGLED	ASDL CGSG	EVHSGFWKAN
ID NO 14:	DNTNKLIVLVS	FRGSRSIENW	IGNLNFDLKE	INDI CGSG	RGDGFTSSW
ID NO 15:	NEPCKEITVA	YRGTDLSLIDW	LTNLFQKTA	WPAN ISNS	LVHEGFLINAY
ID NO 1:	NEVQDKLVAE	VIAQLDNRPG	YKIVVFTGHS	GGATAVL SALDLYNNHGA	
ID NO 2:	NEVQDKLVAE	VKAQLDNRPG	YKIVVFTGHS	GGATAVL SALDLYNNHGD	
ID NO 3:	GEVQNLVELAT	VLDQFKQYPS	YKVAFTGHS	GGATALL CALDLYQREEGLS	
ID NO 4:	EQVVNDYFPV	VOEQLTAMPT	YKIVVFTGHS	GGAQALLAAGMDLYQRESPRLS	
ID NO 5:	VSVQDQVESL	VKQQVSYQPD	YALTVTGHS	GASLAALTAAGL SATYD	
ID NO 6:	ISVQDQVESL	VQQQVSQFPD	YALTVTGHS	GASLAALTAAGL SATYD	
ID NO 7:	NEISSQATAA	VASANKANP	FNVLISTGHS	GGAVAVLAAANLRVGCT	
ID NO 8:	EEVAANVKAA	VSAAKTANP	FKVIVFTGHS	GGAVAVTIAAAYLRKDGF	
ID NO 9:	KVVRDRIIKI	LDELKEHSD	YKIVVVGHS	GAIAIASLAADLRKJNY	
ID NO 10:	KLVRDIDIKE	LKEVVAQNP	YELVYVGHS	GAAVATLAATDLRGKGY	
ID NO 11:	ESAAADELTSK	IKSAMSTYSG	YTLYFTGHS	CGALATLGATVLRNDGY	
ID NO 12:	EAAAONLTSK	IKSAMSTYSG	YTLYFTGHS	CGALATLGATVLRNDGY	
ID NO 13:	SEIADTTTSK	VESALSDHSD	YSLVLTGHSY	GAALAALAATALRNSH	

Figure 1 (cont.)

ID NO 14: RSVADTLRCK VEDAVREHPD YRVVFTGHS L GGALATVAGADLRGNGY
 ID NO 15: LVSMQQVQEA VDSLLAKCPO ATISFTGHS L GGALACISMVDTAQRHRCI

ID NO 1:	NIEIYTQG QPRIGIIPASA NYVIGT	KIPYQRLVNERDTIVPHL
ID NO 2:	NIEIYTQG QPRIGIIPASA NYVIGT	KIPYQRLVNERDTIVPHL
ID NO 3:	SSNLFLYTYQG QPRVGDPAFA NYVYST	GIEVRYTNERDIVPHL
ID NO 4:	PXNLISIFTVQG QPRVGNPTEA YYVEST	GIPPORTVHKRDIVPHL
ID NO 5:	NIRLYTFC EPRSGNQAF A SYMNDAFQASSPDTTQYFRVTHANDGIPNL	
ID NO 6:	NIRLYTFC EPRS NQAF A SYMNDAFQASSPDTTQYFRVTHANDGIPNL	
ID NO 7:	PVDIYTG SDRVGNQAL A AFVSNQ	AGGEYRVTHADDPPVPRL
ID NO 8:	PFDLVYTG SDRVGNDFFA NFVTQQ	TGAERYRVTHADDPPVPRL
ID NO 9:	DALLYAYA APRVANKPLA EFTITNO	GNNYRFTHNDPPVPKL
ID NO 10:	SAKLYAYA SDRVGNQAL A KYITAQ	CNNPRFTHTNDPPVPKL
ID NO 11:	SVELYTYG CPRIGNYALA EHITSQ	GSGANFRVTHLNDIVPRV
ID NO 12:	SVELYTYG CPRVGNYALA EHITSQ	GSGANFPVTHLNDIVPRV
ID NO 13:	SVELYNYG QPRLGNEALA TYITDQ	NKGGNYRVIETNDIVPKL
ID NO 14:	DIDVFSYG APRVGNQAF A EFTVQ	TGGFLYRITHNDIVPKL
ID NO 15:	KNQMFYTG QPRTGQAF A EYVENL	CHPVFRVVYRMDIVPRM

ID NO 1:	PPGANGFLRA GEEFWIMK	DSSLRVCPNGIETDNCNSIV
ID NO 2:	PPGAGFGLRA GEEFWIMK	DSSLRVCPNGIETDNCNSIV
ID NO 3:	PPAAFGFLRA GEEYNITD	NSPETVQVCTSDETSDCSNSTV
ID NO 4:	PPQSGFGLHP QVSEWIKS	GTSNVQICTSEIETKDCSNSTV
ID NO 5:	PPVEQGYANG QVEYWSV	DPYSAQNTFVCTGDEVQCCE AQGGQG
ID NO 6:	PPADEGYAHG VVEYWSV	DPYSAQNTFVCTGDEVQCCE AQGGQG
ID NO 7:	PPFLINGYAHG TPEFWLSCGGGDVKVDTTISDVVKVCEGAANLG CNGGTL	
ID NO 8:	PPIVFGYRST SPEYWIITS QPLDHDYVTEIKVCEGIANVM CNGGTI	
ID NO 9:	PLLTMGYVHII SPEYVITA	PDNTTVTDNQVTVLDGYVNFK GNTGTS
ID NO 10:	PLLSMGYVHV SPEYWITS	PNNATVSTSDIKVLDGDEVSTD GNTGTC
ID NO 11:	PPMDJFGFSQP SPEYWITS	QNGASVTASDIEVLEGINSTA GNAGEA
ID NO 12:	PPMDNCGPSQP SPEYWITS	QIGASVTASDIEVLEGINSTA GNAGEA
ID NO 13:	PPTLLGYHHP SPEYVIISS	ADEATVTTIDVTEVTGIDATG GNDGTD
ID NO 14:	PPREFGYHS SPEYWIKS	GTLVFPVTRNDIVKIEGIDATG CNOOPN
ID NO 15:	PPMDLGPQHH QGEVWYBG	DENIKPCKGEGENLTCELOV

ID NO 1:	PFT SVIDHLSYLDMMNTGL CL
ID NO 2:	PFT SVIDHLSYLDMMNTGL CL
ID NO 3:	PFT SVLDHLSYFGINTGL CT
ID NO 4:	PFT SILDHLSYFDINECCS CL
ID NO 5:	VN NAHTTYF GMTSGACTW
ID NO 6:	VN NAHTTYF GMTSGACTW
ID NO 7:	GL DIAAHLYHF QAITDA CNAGGFSWR R
ID NO 8:	GL DILAHNTYF QSMAT CAPIAIPWK R
ID NO 9:	GGLPDLAFLASHVWYFIHADACKGPGLPLR
ID NO 10:	LPLLTIDPEAHNHWYF VQVDA CKXGPCLPFK R
ID NO 11:	TV SVALAHLWYF FALSE CLL
ID NO 12:	TV DVALAHLWYF FALSE CLL
ID NO 13:	GT SIDAHRWYF IYISE CS
ID NO 14:	LP DIFAHLWYF GLIGT CL
ID NO 15:	PSEL NAKDMSEYP GMH

ID NO:	Micro organism	SEQ ID NO.:
1.	<i>Absidia reflexa</i>	3
2.	<i>Absidia corymbifera</i>	4
3.	<i>Rhizmucor miehei</i>	5
4.	<i>Rhizopus delemar (oryzea)</i>	6
5.	<i>Aspergillus niger</i>	7
6.	<i>Aspergillus tubingensis</i>	8
7.	<i>Fusarium oxysporum</i>	9
8.	<i>Fusarium heterosporum</i>	10
9.	<i>Aspergillus oryzae</i>	11
10.	<i>Penicillium camembertii</i>	12

11.	<i>Aspergillus foetidus</i>	13
12	<i>Aspergillus niger</i>	14
13.	<i>Aspergillus oryzae</i>	15
14.	<i>Thermomyces lanuginosus</i>	2
15.	<i>Landerina penisapora</i>	16

Figure 1. Alignment of lipase sequences.

POLYPEPTIDES HAVING LIPASE ACTIVITY AND POLYNUCLEOTIDES ENCODING SAME

FIELD OF THE INVENTION

[0001] The present invention relates to polypeptides having lipase activity and polynucleotides encoding same.

BACKGROUND OF THE INVENTION

[0002] Lipases are useful, e.g., as detergent enzymes to remove lipid or fatty stains from clothes and other textiles, as additives to dough for bread and other baked products. Thus, a lipase derived from *Thermomyces lanuginosus* (synonym *Humicola lanuginosa*, EP 258 068 and EP 305 216) is sold for detergent use under the tradename Lipolase® (product of Novo Nordisk A/S). WO 0060063 describes variants of the *T. lanuginosus* lipase with a particularly good first-wash performance in a detergent solution. WO 970407a WO 9707202 and WO 0032758 also disclose variants of the *T. Lanuginosus* lipase. WO 02062973 discloses *T. lanuginosus* lipase with a C-terminal extension with reduced tendency to form odor.

SUMMARY OF THE INVENTION

[0003] The present invention relates to isolated polypeptides having lipase activity selected from the group consisting of lipases having an Average Relative Performance (RP) of at least 0.8 and a Benefit-Risk (BR) of at least 1.1 at the test conditions given in the specification.

[0004] The present invention also relates to lipase variants with reduced potential for odor generation and to a method of preparing them. If particularly relates to variants of the *Thermomyces lanuginosus* lipase having a preference for long fatty acid chains while at the same time having a good relative performance.

[0005] In a further aspect the invention relates to an isolated polynucleotide comprising a nucleotide sequence which encodes the polypeptide, a nucleic acid construct comprising the polynucleotide, a recombinant expression vector comprising the nucleic acid construct and a recombinant host cell comprising the nucleic acid construct.

[0006] The present invention also relates to a method for producing the lipases of the invention.

SEQUENCE LISTING

[0007] SEQ ID NO: 1 shows the DNA sequence encoding lipase from *Thermomyces lanuginosus*.

SEQ ID NO: 2 shows the amino acid sequence of a lipase from *Thermomyces lanuginosus*.

[0008] SEQ ID NO: 3-SEQ ID NO: 16 show sequences for alignment in FIG. 1

[0009] SEQ ID NO: 17 and SEQ ID NO: 18 show sequences used for alignment example.

DEFINITIONS

[0010] Lipase activity: The term "lipase activity" is defined herein as a carboxylic ester hydrolase activity which catalyzes the hydrolysis of triacylglycerol under the formation of diacylglycerol and a carboxylate. For purposes of the present invention, lipase activity is determined according to the procedure described in "Lipase activity" in "Materials and Methods". One unit of lipase activity is defined as the amount of enzyme capable of releasing 1.0 micro mole of butyric acid per minute at 30° C., pH 7.

[0011] The polypeptides of the present invention have at least 70%, such at least 75% or 80% or 85% or 90%, more preferably at least 95%, even more preferably 96% or 97%, most preferably 98% or 99%, and even most preferably at least 100% of the lipase activity measured as Relative Performance of the polypeptide consisting of the amino acid sequence shown as the mature polypeptide of SEQ ID NO: 2, with the substitutions T231R+N233R.

[0012] Isolated polypeptide: The term "isolated polypeptide" as used herein refers to a polypeptide which is at least 20% pure, preferably at least 40% pure, more preferably at least 60% pure, even more preferably at least 80% pure, most preferably at least 90% pure, and even most preferably at least 95% pure, as determined by SDS-PAGE.

[0013] Substantially pure polypeptide: The term "substantially pure polypeptide" denotes herein a polypeptide preparation which contains at most 10%, preferably at most 8%, more preferably at most 6%, more preferably at most 5%, more preferably at most 4%, at most 3%, even more preferably at most 2%, most preferably at most 1%, and even most preferably at most 0.5% by weight of other polypeptide material with which it is natively associated. It is, therefore, preferred that the substantially pure polypeptide is at least 92% pure, preferably at least 94% pure, more preferably at least 95% pure, more preferably at least 98% pure, more preferably at least 96% pure, more preferably at least 97% pure, more preferably at least 98% pure, even more preferably at least 99%, most preferably at least 99.5% pure, and even most preferably 100% pure by weight of the total polypeptide material present in the preparation.

[0014] The polypeptides of the present invention are preferably in a substantially pure form. In particular, it is preferred that the polypeptides are in "essentially pure form", i.e., that the polypeptide preparation is essentially free of other polypeptide material with which it is natively associated. This can be accomplished, for example, by preparing the polypeptide by means of well-known recombinant methods or by classical purification methods.

[0015] Herein, the term "substantially pure polypeptide" is synonymous with the terms "isolated polypeptide" and "polypeptide in isolated form."

[0016] Identity: The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "identity".

[0017] For purposes of the present invention, the alignment of two amino acid sequences is determined by using the Needle program from the EMBOSS package (<http://emboss.org>) version 2.8.0. The Needle program implements the global alignment algorithm described in Needleman, S. B. and Wunsch, C. D. (1970) J. Mol. Biol. 48, 443-453. The substitution matrix used is BLOSUM62, gap opening penalty is 10, and gap extension penalty is 0.5.

[0018] The degree of identity between an amino acid sequence of the present invention ("invention sequence"; e.g. amino acids 1 to 269 of SEQ ID NO: 2) and a different amino acid sequence ("foreign sequence") is calculated as the number of exact matches in an alignment of the two sequences, divided by the length of the "invention sequence" or the length of the "foreign sequence", whichever is the shortest. The result is expressed in percent identity.

[0019] An exact match occurs when the "invention sequence" and the "foreign sequence" have identical amino acid residues in the same positions of the overlap (in the alignment example below this is represented by "|"). The

length of a sequence is the number of amino acid residues in the sequence (e.g. the length of SEQ ID NO: 2 is 269).

[0020] In the alignment example below, the overlap is the amino acid sequence “HTWGER-NL” of Sequence A; or the amino acid sequence “HGWGEDANL” of Sequence B. In the example a gap is indicated by a “-”.

Alignment Example

[0021]

Sequence A:	ACMSHTWGER-NL
Sequence B:	HGWGEDANLAMNPS

[0022] Polypeptide Fragment: The term “polypeptide fragment” is defined herein as a polypeptide having one or more amino acids deleted from the amino and/or carboxyl terminus of SEQ ID NO: 2 or a homologous sequence thereof, wherein the fragment has lipase activity.

[0023] Subsequence: The term “subsequence” is defined herein as a nucleotide sequence having one or more nucleotides deleted from the 5' and/or 3' end of SEQ ID NO: 1 or a homologous sequence thereof, wherein the subsequence encodes a polypeptide fragment having lipase activity.

[0024] Allelic variant: The term “allelic variant” denotes herein any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. An allelic variant of a polypeptide is a polypeptide encoded by an allelic variant of a gene.

[0025] Substantially pure polynucleotide: The term “substantially pure polynucleotide” as used herein refers to a polynucleotide preparation free of other extraneous or unwanted nucleotides and in a form suitable for use within genetically engineered protein production systems. Thus, a substantially pure polynucleotide contains at most 10%, preferably at most 8%, more preferably at most 6%, more preferably at most 5%, more preferably at most 4%, more preferably at most 3%, even more preferably at most 2%, most preferably at most 1%, and even most preferably at most 0.5% by weight of other polynucleotide material with which it is natively associated. A substantially pure polynucleotide may, however, include naturally occurring 5' and 3' untranslated regions, such as promoters and terminators. It is preferred that the substantially pure polynucleotide is at least 90% pure, preferably at least 92% pure, more preferably at least 94% pure, more preferably at least 95% pure, more preferably at least 96% pure, more preferably at least 97% pure, even more preferably at least 98% pure, most preferably at least 99%, and even most preferably at least 99.5% pure by weight. The polynucleotides of the present invention are preferably in a substantially pure form. In particular, it is preferred that the polynucleotides disclosed herein are in “essentially pure form”, i.e., that the polynucleotide preparation is essentially free of other polynucleotide material with which it is natively associated. Herein, the term “substantially pure polynucleotide” is synonymous with the terms “isolated polynucleotide” and “polynucleotide in isolated form.” The polynucleotides may be of genomic, cDNA, RNA, semisynthetic, synthetic origin, or any combinations thereof.

[0026] cDNA: The term “cDNA” is defined herein as a DNA molecule which can be prepared by reverse transcription from a mature, spliced, mRNA molecule obtained from a eukaryotic cell. cDNA lacks intron sequences that are usually present in the corresponding genomic DNA. The initial, primary RNA transcript is a precursor to mRNA which is processed through a series of steps before appearing as mature spliced mRNA. These steps include the removal of intron sequences by a process called splicing. cDNA derived from mRNA lacks, therefore, any intron sequences.

[0027] Nucleic acid construct: The term “nucleic acid construct” as used herein refers to a nucleic acid molecule, either single- or double-stranded, which is isolated from a naturally occurring gene or which is modified to contain segments of nucleic acids in a manner that would not otherwise exist in nature. The term nucleic acid construct is synonymous with the term “expression cassette” when the nucleic acid construct contains the control sequences required for expression of a coding sequence of the present invention.

[0028] Control sequence: The term “control sequences” is defined herein to include all components, which are necessary or advantageous for the expression of a polynucleotide encoding a polypeptide of the present invention. Each control sequence may be native or foreign to the nucleotide sequence encoding the polypeptide. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the nucleotide sequence encoding a polypeptide.

[0029] Operably linked: The term “operably linked” denotes herein a configuration in which a control sequence is placed at an appropriate position relative to the coding sequence of the polynucleotide sequence such that the control sequence directs the expression of the coding sequence of a polypeptide.

[0030] Coding sequence: When used herein the term “coding sequence” means a nucleotide sequence, which directly specifies the amino acid sequence of its protein product. The boundaries of the coding sequence are generally determined by an open reading frame, which usually begins with the ATG start codon or alternative start codons such as GTG and TTG. The coding sequence may a DNA, cDNA, or recombinant nucleotide sequence.

[0031] Expression: The term “expression” includes any step involved in the production of the polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

[0032] Expression vector: The term “expression vector” is defined herein as a linear or circular DNA molecule that comprises a polynucleotide encoding a polypeptide of the invention, and which is operably linked to additional nucleotides that provide for its expression.

[0033] Host cell: The term “host cell”, as used herein, includes any cell type which is susceptible to transformation, transfection, transduction, and the like with a nucleic acid construct comprising a polynucleotide of the present invention.

[0034] Modification: The term “modification” means herein any chemical modification of the polypeptide consisting of the mature polypeptide of SEQ ID NO: 2 as well as genetic manipulation of the DNA encoding that polypeptide. The modification(s) can be substitution(s), deletion(s) and/or insertion(s) of the amino acid(s) as well as replacement(s) of amino acid side chain(s).

[0035] Artificial variant: When used herein, the term “artificial variant” means a polypeptide having lipase activity produced by an organism expressing a modified nucleotide sequence of SEQ ID NO: 1. The modified nucleotide sequence is obtained through human intervention by modification of the nucleotide sequence disclosed in SEQ ID NO: 1.

[0036] Relative performance (RP): The term relative performance reflects performance of the enzyme variant compared to a reference enzyme when measured as the brightness of the color of the textile samples washed with that specific enzyme variant.

[0037] Benefit-Risk factor (BR): The Benefit-Risk factor describes the wash performance compared to risk for odor when the substrate is removed.

[0038] Conventions for Designation of Variants:

[0039] In describing lipase variants according to the invention, the following nomenclature is used for ease of reference:

[0040] Original Amino Acid(s):Position(s):Substituted Amino Acid(s)

[0041] According to this nomenclature, for instance the substitution of glutamic acid for glycine in position 195 is shown as G195E. A deletion of glycine in the same position is shown as G195*, and insertion of an additional amino acid residue such as lysine is shown as G195GK.

[0042] Where a specific lipase contains a “deletion” in comparison with other lipases and an insertion is made in such a position this is indicated as *36D for insertion of an aspartic acid in position 36.

[0043] Multiple mutations are separated by pluses, i.e.: R170Y+G195E, representing mutations in positions 170 and 195 substituting tyrosine and glutamic acid for arginine and glycine, respectively.

[0044] X231 indicates the amino acid in a parent polypeptide corresponding to position 231, when applying the described alignment procedure. X231R indicates that the amino acid is replaced with R. For SEQ ID NO: 2 X is T, and T231R thus indicates a substitution of T in position 231 with R. Where the amino acid in a position (e.g. 231) may be substituted by another amino acid selected from a group of amino acids, e.g. the group consisting of R and P and Y, this will be indicated by X231R/P/Y.

[0045] In all cases, the accepted IUPAC single letter or triple letter amino acid abbreviation is employed.

DETAILED DESCRIPTION OF THE INVENTION

[0046] Polypeptides having Lipase Activity

[0047] The present invention relates to isolated polypeptides having lipase activity selected from the group consisting of lipases having a RP of at least 0.8 and a BR of at least 1.1 at the test conditions given in the specification.

[0048] in a preferred embodiment the lipase has a RP of at least 0.9, such as 1.0 or 1.1. In an even more preferred embodiment the lipase has a RP of at least 1.2, such as 1.3 or even 1.4.

[0049] In another preferred embodiment the lipase has a BR of at least 1.2, such as 1.3 or even 1.4. In an even more preferred embodiment the lipase has a BR of at least 1.5, such as 1.6 or even 1.7.

[0050] In a further aspect the lipase of the present invention further has a relative LU/A280 less than 1, such as less than 0.95 at the test conditions given in the specification. In a preferred embodiment the relative LU/A280 is less than 0.90, such as less than 0.85 or even less than 0.80.

[0051] In a further aspect, the present invention relates to isolated polypeptides having an amino acid sequence which is comprised by or comprises SEQ ID NO: 2, or an allelic variant thereof, and which further has BR of at least 1.1 and RP of at least 0.8. In another aspect, the present invention relates to isolated polypeptides having an amino acid sequence which is comprised by or comprises the mature part of SEQ ID NO: 2, or an allelic variant thereof, and which further has BR of at least 1.1 and RP of at least 0.8

[0052] In a still further aspect, the present invention relates to isolated polypeptides having an amino acid sequence which has a degree of identity to the mature polypeptide of SEQ ID NO: 2 (i.e., the mature polypeptide) of at least 80%, such as at least 85% or 90%, or at least 95%, preferably at least 97%, most preferably at least 98%, and even most preferably at least 99%, which have lipase activity (hereinafter “homologous polypeptides”). In a preferred aspect, the homologous polypeptides have an amino acid sequence which differs by ten amino acids, by nine amino acids, by eight amino acids, by seven amino acids, by six amino acids, preferably by five amino acids, more preferably by four amino acids, even more preferably by three amino acids, most preferably by two amino acids, and even most preferably by one amino acid from the mature polypeptide of SEQ ID NO: 2.

[0053] In a further aspect, the present invention relates to isolated polypeptides having lipase activity which are encoded by polynucleotides which hybridize under very low stringency conditions, preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with (i) nucleotides 644 to 732 of SEQ ID NO: 1, (ii) the cDNA sequence contained in nucleotides 644 to 732 of SEQ ID NO: 1, (iii) a subsequence of (i) or (ii), or (iv) a complementary strand of (i), (ii), or (iii) (J. Sambrook, E. F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning, A Laboratory Manual* 2d edition, Cold Spring Harbor, N.Y.). A subsequence of SEQ ID NO: 1 contains at least 100 contiguous nucleotides or preferably at least 200 contiguous nucleotides. Moreover, the subsequence may encode a polypeptide fragment which has lipase activity.

[0054] The nucleotide sequence of SEQ ID NO: 1 or a subsequence thereof, as well as the amino acid sequence of SEQ ID NO: 2 or a fragment thereof, may be used to design a nucleic acid probe to identify and clone DNA encoding polypeptides having lipase activity from strains of different genera or species according to methods well known in the art. In particular, such probes can be used for hybridization with the genomic or cDNA of the genus or species of interest, following standard Southern blotting procedures, in order to identify and isolate the corresponding gene therein. Such probes can be considerably shorter than the entire sequence, but should be at least 14, preferably at least 25, more preferably at least 35, and most preferably at least 70 nucleotides in

length. It is, however, preferred that the nucleic acid probe is at least 100 nucleotides in length. For example, the nucleic acid probe may be at least 200 nucleotides, preferably at least 300 nucleotides, more preferably at least 400 nucleotides, or most preferably at least 500 nucleotides in length. Even longer probes may be used, e.g., nucleic acid probes which are at least 600 nucleotides, at least preferably at least 700 nucleotides, or more preferably at least 800 nucleotides in length. Both DNA and RNA probes can be used. The probes are typically labeled for detecting the corresponding gene (for example, with ^{32}P , ^{3}H , ^{35}S , biotin, or avidin). Such probes are encompassed by the present invention.

[0055] A genomic DNA or cDNA library prepared from such other organisms may, therefore, be screened for DNA which hybridizes with the probes described above and which encodes a polypeptide having lipase activity. Genomic or other DNA from such other organisms may be separated by agarose or polyacrylamide gel electrophoresis, or other separation techniques, DNA from the libraries or the separated DNA may be transferred to and immobilized on nitrocellulose or other suitable carrier material. In order to identify a clone or DNA which is homologous with SEQ ID NO: 1 or a subsequence thereof, the carrier material is used in a Southern blot.

[0056] For purposes of the present invention, hybridization indicates that the nucleotide sequence hybridizes to a labeled nucleic acid probe corresponding to the nucleotide sequence shown in SEQ ID NO: 1, its complementary strand, or a subsequence thereof, under very low to very high stringency conditions. Molecules to which the nucleic acid probe hybridizes under these conditions can be detected using X-ray film.

[0057] In a still further aspect the present invention relates to artificial variants comprising a conservative substitution, deletion, and/or insertion of one or more amino acids of SEQ ID NO: 2 or the mature polypeptide thereof, said variants having a BR of at least 1.1 and a RP of at least 0.8. Preferably, amino acid changes are of a minor nature, that is conservative amino acid substitutions or insertions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of one to about 30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to about 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

[0058] Examples of conservative substitutions are within the group of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine, isoleucine and valine), aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino acids (glycine, alanine, serine, threonine and methionine). Amino acid substitutions which do not generally alter specific activity are known in the art and are described, for example, by H. Neurath and R. L. Hill, 1979, in, *The Proteins*, Academic Press, New York. The most commonly occurring exchanges are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly.

[0059] In addition to the 20 standard amino acids, non-standard amino acids (such as 4-hydroxyproline, 6-N-methyl lysine, 2-aminoisobutyric acid, isovaline, and alpha-methyl serine) may be substituted for amino acid residues of a wild-type polypeptide. A limited number of non-conservative

amino acids, amino acids that are not encoded by the genetic code, and unnatural amino acids may be substituted for amino acid residues. "Unnatural amino acids" have been modified after protein synthesis, and/or have a chemical structure in their side chain(s) different from that of the standard amino acids. Unnatural amino acids can be chemically synthesized, and preferably, are commercially available, and include pipolic acid, thiazolidine carboxylic acid, dehydroproline, 3- and 4-methylproline, and 3,3-dimethylproline.

[0060] Alternatively, the amino acid changes are of such a nature that the physico-chemical properties of the polypeptides are altered. For example, amino acid changes may improve the thermal stability of the polypeptide, alter the substrate specificity, change the pH optimum, and the like.

[0061] Essential amino acids in the parent polypeptide can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, 1989, *Science* 244: 1081-1085), in the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for biological activity (i.e., lipase activity) to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., 1996, *J. Biol. Chem.* 271: 4699-4708. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction, or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., 1992, *Science* 255:306-312; Smith et al., 1992, *J. Mol. Biol.* 224: 899-904; Wlodaver et al., 1992, FEES Lett. 309:59-64. The identities of essential amino acids can also be inferred from analysis of identities with polypeptides which are related to a polypeptide according to the invention.

[0062] Single or multiple amino acid substitutions can be made and tested using known methods of mutagenesis, recombination, and/or shuffling, followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer, 1988, *Science* 241: 53-57; Bowie and Sauer, 1989, *Proc. Natl. Acad. Sci. USA* 86: 2152-2156; WO 95/17413; or WO 95/22625. Other methods that can be used include error-prone PCR, phage display (e.g., Lowman et al., 1991, *Biochem.* 30: 10832-10837; U.S. Pat. No. 5,223,409; WO 92/06204), and region-directed mutagenesis (Derbyshire et al., 1986, *Gene* 46:145; Ner et al., 1988, *DNA* 7:127).

[0063] Mutagenesis/shuffling methods can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides expressed by host cells. (Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using standard methods in the art. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of interest, and can be applied to polypeptides of unknown structure.

[0064] The total number of amino acid substitutions, deletions and/or insertions of amino acids 1 to 291 of SEQ ID NO: 2 is 10, preferably 9, more preferably 8, more preferably 7, more preferably at most 6, more preferably at most 5, more preferably 4, even more preferably 3, most preferably 2, and even most preferably 1.

Identification of Regions and Substitutions

[0065] Substitutions covered by the present application may be identified as disclosed in this section.

[0066] The positions referred to in Region I through Region IV below are the positions of the amino acid residues in SEQ ID NQ: 2. To find the corresponding (or homologous) positions in a different lipase, the procedure described in "Homology and alignment" is used.

Substitutions in Region I

[0067] Region I consists of amino acid residues surrounding the N-terminal residue E1. In this region it is preferred to substitute an amino acid of the parent lipase with a more positive amino acid.

[0068] Amino acid residues corresponding to the following positions are comprised by Region I: 2 to 11 and 223-239. The following positions are of particular interest: 4, 8, 11, 223, 229, 231, 233, 234, 236.

[0069] In particular the following substitutions have been identified: X4V, X231R and X233R.

[0070] In a preferred embodiment the variant lipase has at least 80%, such as 85% or 90%, such as at least 95% or 98% or 99% identity to SEQ ID NO:2

Substitutions in Region II

[0071] Region II consists of amino acid residues in contact with substrate on one side of the acyl chain and one side of the alcohol part. In this region it is preferred to substitute an amino acid of the parent lipase with a more positive amino acid or with a less hydrophobic amino acid.

[0072] Amino acid residues corresponding to the following positions are comprised by Region II: 202 to 211 and 249 to 269. The following positions are of particular interest: 202, 210, 211, 253, 254, 255, 256.

[0073] In particular the following substitutions have been identified: X202G, X255Y/V and X256K/R.

[0074] In a preferred embodiment the variant lipase has at least 80%, such as 85% or 90%, such as at least 95% or 98% or 99% identity to SEQ ID NO: 2

Substitutions in Region III

[0075] Region II consists of amino acid residues that forms a flexible structure and thus allowing the substrate to get into the active site. In this region it is preferred to substitute an amino acid of the parent lipase with a more positive amino acid or a less hydrophobic amino acid.

[0076] Amino acid residues corresponding to the following positions are comprised by Region III: 82 to 102. The following positions are of particular interest 86, 87, 90, 91, 95, 96, 99.

[0077] In particular the following substitutions have been identified: X86V and X90A/R.

[0078] In a preferred embodiment the variant lipase has at least 80%, such as 85% or 90%, such as at least 95% or 98% or 99% identity to SEQ ID NO: 2

Substitutions in Region IV

[0079] Region IV consists of amino acid residues that binds electrostatically to a surface. In this region it is preferred to substitute an amino acid of the parent lipase with a more positive amino acid.

[0080] Amino acid residues corresponding to the following positions are comprised by Region IV: 27 and 54 to 62. The following positions are of particular interest: 27, 56, 57, 58, 60.

[0081] In particular the following substitutions have been identified: X27R, X58N/AG/T/P and X60V/S/G/N/R/K/A/L.

[0082] In a preferred embodiment the variant lipase has at least 80%, such as 85% or 90%, such as at least 95% or 98% or 99% identity to SEQ ID NO: 2

Amino Acids at Other Positions

[0083] The parent lipase may optionally comprise substitution of other amino acids, i particularly less than 10 or less than 5 such substitutions. Examples are substitutions corresponding to one or more of the positions 24, 46, 74, 81, 83, 127, 131, 137, 147, 150, 203, 206, 211, 263, 264, 265, 267 and 269 of SEQ ID NO: 2. In a particular embodiment there is a substitution in at least one of the positions corresponding to position 81, 147, 150, 227 and 249. In a preferred embodiment the at least one substitution is selected from the group consisting of X81Q/E, X147M/Y, X150G, X227G and X249R/I/L.

[0084] Further substitutions may, e.g., be made according to principles known in the art, e.g. substitutions described in WO 92/05249, WO 94/25577, WO 95/22615, WO 97/04079 and WO 97/07202.

Homology and Alignment

[0085] For purposes of the present invention, the degree of homology may be suitably determined by means of computer programs known in the art, such as GAP provided in the GCG program package (Program Manual for the Wisconsin Package, Version 8, August 1994, Genetics Computer Group, 575 Science Drive, Madison, Wis., USA 53711) (Needleman, S. B. and Wunsch, C. D., (1970), Journal of Molecular Biology, 48, 443-45), using GAP with the following settings for polypeptide sequence comparison: GAP creation penalty of 3.0 and GAP extension penalty of 0.1.

[0086] In the present invention, corresponding (or homologous) positions in the lipase sequences of *Absidia reflexa*, *Absidia corymbifera*, *Rhizomucor miehei*, *Rhizopus delemar*, *Aspergillus niger*, *Aspergillus tubingensis*, *Fusarium oxysporum*, *Fusarium heterosporum*, *Aspergillus oryzae*, *Penicillium camembertii*, *Aspergillus foetidus*, *Aspergillus niger*, *Thermomyces lanuginosus* (synonym: *Humicola lanuginosa*) and *Landerine penispora* are defined by the alignment shown in FIG. 1.

[0087] To find the homologous positions in lipase sequences not shown in the alignment, the sequence of interest is aligned to the sequences shown in FIG. 1. The new sequence is aligned to the present alignment in FIG. 1 by using the GAP alignment to the most homologous sequence found by the GAP program. GAP is provided in the GCG program package (Program Manual for the Wisconsin Package, Version 8, August 1994, Genetics Computer Group, 575 Science Drive, Madison, Wis., USA 53711) (Needleman, S. B. and Wunsch, C. D., (1970), Journal of Molecular Biology, 48, 443-45). The following settings are used for polypeptide sequence comparison: GAP creation penalty of 3.0 and GAP extension penalty of 0.1.

[0088] Any suitable parent lipase may be used. In a preferred embodiment, the parent lipase has a homology of at least 50% with the *T. lanuginosus* lipase (SEQ ID NO: 2), particularly at least 55%, at least 60%, at least 75%, at least 85%, at least 90%, more than 95%, 96%, 97% or more than

98% or 99%. In a particular embodiment the parent lipase is identical to the *T. lanuginosus* lipase (SEQ ID NO: 2).

Sources of Polypeptides Having Lipase Activity

[0089] A polypeptide of the present invention may be obtained from microorganisms of any genus. For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the polypeptide encoded by a nucleotide sequence is produced by the source or by a strain in which the nucleotide sequence from the source has been inserted. In a preferred aspect, the polypeptide obtained from a given source is secreted extracellularly.

[0090] A polypeptide of the present invention may be a bacterial polypeptide. For example, the polypeptide may be a gram positive bacterial polypeptide such as a *Bacillus* polypeptide, e.g., a *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus coagulans*, *Bacillus lautus*, *Bacillus lenth*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus stearothermophilus*, *Bacillus subtilis*, or *Bacillus thuringiensis* polypeptide; or a *Streptomyces* polypeptide, e.g., a *Streptomyces lividans* or *Streptomyces murinus* polypeptide; or a gram negative bacterial polypeptide, e.g., an *E. coli* or a *Pseudomonas* sp. polypeptide.

[0091] A polypeptide of the present invention may also be a fungal polypeptide, and more preferably a yeast polypeptide such as a *Candida*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* polypeptide; or more preferably a filamentous fungal polypeptide such as an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Cryptococcus*, *Filobasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Piromyces*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, or *Trichoderma* polypeptide.

[0092] In a preferred aspect, the polypeptide is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis*, or *Saccharomyces oviformis* polypeptide having lipase activity.

[0093] In another preferred aspect, the polypeptide is an *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, *Aspergillus turbigensis*, *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium gramininum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, *Fusarium venenatum*, *Humicola insolens*, *Thermomyces lanuginosus* (synonym: *Humicola lanuginose*), *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpurogenum*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* polypeptide.

[0094] In another preferred aspect, the polypeptide is a *Thermomyces* polypeptide.

[0095] In a more preferred aspect, the polypeptide is a *Thermomyces lanuginosus* polypeptide, e.g., the polypeptide of SEQ ID NO: 2 with mutations as disclosed in the present application.

[0096] It will be understood that for the aforementioned species, the invention encompasses both the perfect and imperfect states, and other taxonomic equivalents, e.g., anamorphs, regardless of the species name by which they are known. Those skilled in the art will readily recognize the identity of appropriate equivalents.

[0097] Strains of these species are readily accessible to the public in a number of culture collections, such as the American Type Culture Collection (ATCC), Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Centraalbureau Voor Schimmelcultures (CBS), and Agricultural Research Service Patent Culture Collection, Northern Regional Research Center (NRRL).

[0098] Furthermore, such polypeptides may be identified and obtained from other sources including microorganisms isolated from nature (e.g., soil, composts, water, etc.) using the above-mentioned probes. Techniques for isolating microorganisms from natural habitats are well known in the art. The polynucleotide may then be obtained by similarly screening a genomic or cDNA library of another microorganism. Once a polynucleotide sequence encoding a polypeptide has been detected with the probe(s), the polynucleotide can be isolated or cloned by utilizing techniques which are well known to those of ordinary skill in the art (see, e.g., Sambrook et al., 1989, *supra*).

[0099] Polypeptides of the present invention also include fused polypeptides or cleavable fusion polypeptides in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide or fragment thereof. A fused polypeptide is produced by fusing a nucleotide sequence (or a portion thereof) encoding another polypeptide to a nucleotide sequence (or a portion thereof) of the present invention. Techniques for producing fusion polypeptides are known in the art, and include ligating the coding sequences encoding the polypeptides so that they are in frame and that expression of the fused polypeptide is under control of the same promoter(s) and terminator.

Polynucleotides

[0100] The present invention also relates to isolated polynucleotides having a nucleotide sequence which encode a polypeptide of the present invention. The present invention also encompasses nucleotide sequences which encode a polypeptide being a variant of the amino acid sequence of SEQ ID NO: 2 or the mature polypeptide thereof, which differ from the encoding polynucleotide by virtue of the degeneracy of the genetic code. The present invention also relates to subsequences of SEQ ID NO: 1 which encode fragments of SEQ ID NO: 2 that have lipase activity, said fragments having a BR of at least 1.1 and a RP of at least 0.8.

[0101] The techniques used to isolate or clone a polynucleotide encoding a polypeptide are known in the art and include isolation from genomic DNA, preparation from cDNA, or a combination thereof. The cloning of the polynucleotides of the present invention from such genomic DNA can be effected, e.g., by using the well known polymerase chain reaction (PCR) or antibody screening of expression libraries to detect cloned DNA fragments with shared structural features. See, e.g., Innis et al., 1990, PCR: *A Guide to Methods and Application*, Academic Press, New York. Other nucleic

acid amplification procedures such as ligase chain reaction (LCR), ligated activated transcription (LAT) and nucleotide sequence-based amplification (NASBA) may be used. The polynucleotides may be cloned from a strain of *Thermomyces*, or another or related organism and thus, for example, may be an allelic or species variant of the polypeptide encoding region of the nucleotide sequence.

[0102] The present invention also relates to polynucleotides having nucleotide sequences which have a degree of identity to the mature polypeptide coding sequence of SEQ ID NO: 1 of at least 60%, preferably at least 65%, more preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, even more preferably at least 95%, and most preferably at least 97%, 98%, or 99% identity, which encode an active polypeptide having lipase activity and BR of at least 1.1 and RP of at least 0.8.

[0103] Modification of a nucleotide sequence encoding a polypeptide of the present invention may be necessary for the synthesis of polypeptides substantially similar to the polypeptide. The term "substantially similar" to the polypeptide refers to non-naturally occurring forms of the polypeptide. These polypeptides may differ in some engineered way from the polypeptide isolated from its native source, e.g., artificial variants that differ in specific activity, thermo stability, pH optimum, or the like. The variant sequence may be constructed on the basis of the nucleotide sequence presented as the polypeptide encoding region of SEQ ID NO: 1, e.g., a subsequence thereof, and/or by introduction of nucleotide substitutions which do not give rise to another amino acid sequence of the polypeptide encoded by the nucleotide sequence, but which correspond to the codon usage of the host organism intended for production of the enzyme, or by introduction of nucleotide substitutions which may give rise to a different amino acid sequence. For a general description of nucleotide substitution, see, e.g., Ford et al., 1991, *Protein Expression and Purification* 2: 95-107.

[0104] It will be apparent to those skilled in the art that such substitutions can be made outside the regions critical to the function of the molecule and still result in an active polypeptide. Amino acid residues essential to the activity of the polypeptide encoded by an isolated polynucleotide of the invention, and therefore preferably not subject to substitution, may be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (see, e.g., Cunningham and Wells, 1989, *Science* 244: 1081-1085). In the latter technique, mutations are introduced at every positively charged residue in the molecule, and the resultant mutant molecules are tested for lipase activity to identify amino acid residues that are critical to the activity of the molecule. Sites of substrate-enzyme interaction can also be determined by analysis of the three-dimensional structure as determined by such techniques as nuclear magnetic resonance analysis, crystallography or photoaffinity labelling (see, e.g., de Vos et al., 1992, *Science* 255: 306-312; Smith et al., 1992, *Journal of Molecular Biology* 224: 899-904; Wlodaver et al., 1992, *FEBS Letters* 309: 59-64).

[0105] The present invention also relates to isolated polynucleotides encoding a polypeptide of the present invention, which hybridize under very low stringency conditions, preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions

with (i) of SEQ ID NO: 1, (ii) the cDNA sequence contained in SEQ ID NO: 1, or (iii) a complementary strand of (i) or (ii); or allelic variants and subsequences thereof (Sambrook et al., 1989, *supra*), as defined herein.

[0106] The present invention also relates to isolated polynucleotides obtained by (a) hybridizing a population of DNA under very low, low, medium, medium-high, high, or very high stringency conditions with (i) nucleotides SEQ ID NO: 1, (ii) the cDNA sequence contained in nucleotides of SEQ ID NO: 1, or (iii) a complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having lipase activity.

Nucleic Acid Constructs

[0107] The present invention also relates to nucleic acid constructs comprising an isolated polynucleotide of the present invention operably linked to one or more control sequences which direct the expression of the coding sequence in a suitable host cell under conditions compatible with the control sequences.

[0108] An isolated polynucleotide encoding a polypeptide of the present invention may be manipulated in a variety of ways to provide for expression of the polypeptide. Manipulation of the polynucleotide's sequence prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotide sequences utilizing recombinant DNA methods are well known in the art.

[0109] The control sequence may be an appropriate promoter sequence, a nucleotide sequence which is recognized by a host cell for expression of a polynucleotide encoding a polypeptide of the present invention. The promoter sequence contains transcriptional control sequences which mediate the expression of the polypeptide. The promoter may be any nucleotide sequence which shows transcriptional activity in the host cell of choice including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

[0110] Examples of suitable promoters for directing the transcription of the nucleic acid constructs of the present invention, especially in a bacterial host cell, are the promoters obtained from the *E. coli* lac operon, *Streptomyces* coelicolor agarase gene (dagA), *Bacillus subtilis* levansucrase gene (sacB), *Bacillus licheniformis* alpha-amylase gene (amyL), *Bacillus stearothermophilus* maltogenic amylase gene (amyM), *Bacillus amyloliquefaciens* alpha-amylase gene (amyQ), *Bacillus licheniformis* penicillinase gene (penP), *Bacillus subtilis* xylA and xylB genes, and prokaryotic beta-lactamase gene (Villa-Kamaroff et al., 1978, *Proceedings of the National Academy of Sciences USA* 75: 3727-3781), as well as the tac promoter (DeBoer et al., 1983, *Proceedings of the National Academy of Sciences USA* 80: 21-25). Further promoters are described in "Useful proteins from recombinant bacteria" in *Scientific American*, 1980, 242: 74-94; and in Sambrook et al., 1989, *supra*.

[0111] Examples of suitable promoters for directing the transcription of the nucleic acid constructs of the present invention in a filamentous fungal host cell are promoters obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (glaA), *Rhizomucor miehei* lipase, *Aspergillus*

oryzae alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Aspergillus nidulans* acetamidase, *Fusarium venenatum* amyloglucosidase (WO 00/56900), *Fusarium venenatum* Daria (WO 00/56900), *Fusarium venenatum* Quinn (WO 00/56900), *Fusarium oxysporum* trypsin-like protease (WO 96/00787), *Trichoderma reesei* beta-glucosidase, *Trichoderma reesei* cellobiohydrolase I, *Trichoderma reesei* endoglucanase I, *Trichoderma reesei* endoglucanase II, *Trichoderma reesei* endoglucanase III, *Trichoderma reesei* endoglucanase IV, *Trichoderma reesei* endoglucanase V, *Trichoderma reesei* xylanase I, *Trichoderma reesei* xylanase II, *Trichoderma reesei* beta-xylosidase, as well as the NA2-tpi promoter (a hybrid of the promoters from the genes for *Aspergillus niger* neutral alpha-amylase and *Aspergillus oryzae* triose phosphate isomerase); and mutant, truncated, and hybrid promoters thereof.

[0112] In a yeast host, useful promoters are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* galactokinase (GAL1), *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH1,ADH2/GAP), *Saccharomyces cerevisiae* triose phosphate isomerase (TPI), *Saccharomyces cerevisiae* metallothioneine (CUP1), and *Saccharomyces cerevisiae* 3-phosphoglycerate kinase. Other useful promoters for yeast host cells are described by Romanos et al., 1992, *Yeast* 8: 423-488.

[0113] The control sequence may also be a suitable transcription terminator sequence, a sequence recognized by a host cell to terminate transcription. The terminator sequence is operably linked to the 3' terminus of the nucleotide sequence encoding the polypeptide. Any terminator which is functional in the host cell of choice may be used in the present invention.

[0114] Preferred terminators for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* alpha-glucosidase, and *Fusarium oxysporum* trypsin-like protease.

[0115] Preferred terminators for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase, *Saccharomyces cerevisiae* cytochrome C (CYC1), and *Saccharomyces cerevisiae* glyceraldehyde-3-phosphate dehydrogenase. Other useful terminators for yeast host cells are described by Romanos et al., 1992, supra.

[0116] The control sequence may also be a suitable leader sequence, a nontranslated region of an mRNA which is important for translation by the host cell. The leader sequence is operably linked to the 5' terminus of the nucleotide sequence encoding the polypeptide. Any leader sequence that is functional in the host cell of choice may be used in the present invention.

[0117] Preferred leaders for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase and *Aspergillus nidulans* triose phosphate isomerase.

[0118] Suitable leaders for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* 3-phosphoglycerate kinase, *Saccharomyces cerevisiae* alpha-factor, and *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP).

[0119] The control sequence may also be a polyadenylation sequence, a sequence operably linked to the 3' terminus of the nucleotide sequence and which, when transcribed, is recog-

nized by the host cell as a signal to add polyadenosine residues to transcribed mRNA. Any polyadenylation sequence which is functional in the host cell of choice may be used in the present invention.

[0120] Preferred polyadenylation sequences for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Fusarium oxysporum* trypsin-like protease, and *Aspergillus niger* alpha-glucosidase.

[0121] Useful polyadenylation sequences for yeast host cells are described by Guo and Sherman, 1995, *Molecular Cellular Biology* 15: 5983-5990.

[0122] The control sequence may also be a signal peptide coding region that codes for an amino acid sequence linked to the amino terminus of a polypeptide and directs the encoded polypeptide into the cell's secretory pathway. The 5' end of the coding sequence of the nucleotide sequence may inherently contain a signal peptide coding region naturally linked in translation reading frame with the segment of the coding region which encodes the secreted polypeptide. Alternatively, the 5' end of the coding sequence may contain a signal peptide coding region which is foreign to the coding sequence. The foreign signal peptide coding region may be required where the coding sequence does not naturally contain a signal peptide coding region. Alternatively, the foreign signal peptide coding region may simply replace the natural signal peptide coding region in order to enhance secretion of the polypeptide. However, any signal peptide coding region which directs the expressed polypeptide into the secretory pathway of a host cell of choice may be used in the present invention.

[0123] Effective signal peptide coding regions for bacterial host cells are the signal peptide coding regions obtained from the genes for *Bacillus* NCIB 11837 maltogenic amylase, *Bacillus stearothermophilus* alpha-amylase, *Bacillus licheniformis* subtilisin, *Bacillus licheniformis* beta-lactamase, *Bacillus stearothermophilus* neutral proteases (nprT, nprS, nprM), and *Bacillus subtilis* prsA. Further signal peptides are described by Simonen and Palva, 1993, *Microbiological Reviews* 57: 109-137.

[0124] Effective signal peptide coding regions for filamentous fungal host cells are the signal peptide coding regions obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* neutral amylase, *Aspergillus niger* glucoamylase, *Rhizomucor miehei* aspartic proteinase, *Humicola insolens* cellulase, and *Humicola lanuginosa* lipase.

[0125] Useful signal peptides for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* alpha-factor and *Saccharomyces cerevisiae* invertase. Other useful signal peptide coding regions are described by Romanos et al., 1992, supra.

[0126] The control sequence may also be a propeptide coding region that codes for an amino acid sequence positioned at the amino terminus of a polypeptide. The resultant polypeptide is known as a proenzyme or propolypeptide (or a zymogen in some cases). A propolypeptide is generally inactive and can be converted to a mature active polypeptide by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide. The propeptide coding region may be obtained from the genes for *Bacillus subtilis* alkaline protease (aprE), *Bacillus subtilis* neutral protease (nprT), *Saccharo-*

myces cerevisiae alpha-factor, *Rhizomucor miehei* aspartic proteinase, and *Myceliophthora thermophila* laccase (WO 95/33836).

[0127] Where both signal peptide and propeptide regions are present at the amino terminus of a polypeptide, the propeptide region is positioned next to the amino terminus of a polypeptide and the signal peptide region is positioned next to the amino terminus of the propeptide region.

[0128] It may also be desirable to add regulatory sequences which allow the regulation of the expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those which cause the expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Regulatory systems in prokaryotic systems include the lac, lac, and trp operator systems. In yeast, the ADH2 system or GAL1 system may be used. In filamentous fungi, the TAKA alpha-amylase promoter, *Aspergillus niger* glucoamylase promoter, and *Aspergillus oryzae* glucoamylase promoter may be used as regulatory sequences. Other examples of regulatory sequences are those which allow for gene amplification. In eukaryotic systems, these include the dihydrofolate reductase gene which is amplified in the presence of methotrexate, and the metallothionein genes which are amplified with heavy metals. In these cases, the nucleotide sequence encoding the polypeptide would be operably linked with the regulatory sequence.

Expression Vectors

[0129] The present invention also relates to recombinant expression vectors comprising a polynucleotide of the present invention, a promoter, and transcriptional and translation stop signals. The various nucleic acids and control sequences described above may be joined together to produce a recombinant expression vector which may include one or more convenient restriction sites to allow for insertion or substitution of the nucleotide sequence encoding the polypeptide at such sites. Alternatively, a nucleotide sequence of the present invention may be expressed by inserting the nucleotide sequence or a nucleic acid construct comprising the sequence into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression.

[0130] The recombinant expression vector may be any vector (e.g., a plasmid or virus) which can be conveniently subjected to recombinant DNA procedures and can bring about expression of the nucleotide sequence. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vectors may be linear or closed circular plasmids.

[0131] The vector may be an autonomously replicating vector, i.e., a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one which, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. Furthermore, a single vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the host cell, or a transposon may be used.

[0132] The vectors of the present invention preferably contain one or more selectable markers which permit easy selection of transformed cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like.

[0133] A conditionally essential gene may function as a non-antibiotic selectable marker. Non-limiting examples of bacterial conditionally essential non-antibiotic selectable markers are the dal genes from *Bacillus subtilis*, *Bacillus licheniformis*, or other *Bacilli*, that are only essential when the bacterium is cultivated in the absence of D-alanine. Also the genes encoding enzymes involved in the turnover of UDP-galactose can function as conditionally essential markers in a cell when the cell is grown in the presence of galactose or grown in a medium which gives rise to the presence of galactose. Non-limiting examples of such genes are those from *B. subtilis* or *B. licheniformis* encoding UTP-dependent phosphorylase (EC 2.7.7.10), UDP-glucose-dependent uridylyltransferase (EC 2.7.7.12), or UDP-galactose epimerase (EC 5.1.3.2). Also a xylose isomerase gene such as *xylA*, of *Bacilli* can be used as selectable markers in cells grown in minimal medium with xylose as sole carbon source. The genes necessary for utilizing gluconate, *gntK* and *gntP* can also be used as selectable markers in cells grown in minimal medium with gluconate as sole carbon source. Other examples of conditionally essential genes are known in the art. Antibiotic selectable markers confer antibiotic resistance to such antibiotics as ampicillin, kanamycin, chloramphenicol, erythromycin, tetracycline, neomycin, hygromycin or methotrexate.

[0134] Suitable markers for yeast host cells are ADE2, HIS3, LEU2, LYS2, MET3, TRP1, and URA3. Selectable markers for use in a filamentous fungal host cell include, but are not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hph* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase), *sC* (sulfate adenyltransferase), and *trpC* (anthranilate synthase), as well as equivalents thereof. Preferred for use in an *Aspergillus* cell are the *amdS* and *pyrG* genes of *Aspergillus nidulans* or *Aspergillus oryzae* and the *bar* gene of *Streptomyces hygroscopicus*.

[0135] The vectors of the present invention preferably contain an element(s) that permits integration of the vector into the host cell's genome or autonomous replication of the vector in the cell independent of the genome.

[0136] For integration into the host cell genome, the vector may rely on the polynucleotide's sequence encoding the polypeptide or any other element of the vector for integration into the genome by homologous or nonhomologous recombination. Alternatively, the vector may contain additional nucleotide sequences for directing integration by homologous recombination into the genome of the host cell at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements should preferably contain a sufficient number of nucleic acids, such as 100 to 10,000 base pairs, preferably 400 to 10,000 base pairs, and most preferably 800 to 10,000 base pairs, which have a high degree of identity with the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational

elements may be non-encoding or encoding nucleotide sequences. On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination.

[0137] For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. The origin of replication may be any plasmid replicator mediating autonomous replication which functions in a cell. The term "origin of replication" or "plasmid replicator" is defined herein as a nucleotide sequence that enables a plasmid or vector to replicate in vivo.

[0138] Examples of bacterial origins of replication are the origins of replication of plasmids pBR322, pUC19, pACYC177, and pACYC184 permitting replication in *E. coli*, and pUB110, pE194, pTA1060, and pAMβ1 permitting replication in *Bacillus*.

[0139] Examples of origins of replication for use in a yeast host cell are the 2 micron origin of replication, ARS1, ARS4, the combination of ARS1 and CEN3, and the combination of ARS4 and CEN6.

[0140] Examples of origins of replication useful in a filamentous fungal cell are AMA1 and ANS1 (Gems et al, 1991, *Gene* 98: 61-67; Cullen et al., 1987, *Nucleic Acids Research* 15: 9163-9175; WO 00/24883). Isolation of the AMA1 gene and construction of plasmids or vectors comprising the gene can be accomplished according to the methods disclosed in WO 00/24883.

[0141] More than one copy of a polynucleotide of the present invention may be inserted into the host cell to increase production of the gene product. An increase in the copy number of the polynucleotide can be obtained by integrating at least one additional copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the polynucleotide where cells containing amplified copies of the selectable marker gene, and thereby additional copies of the polynucleotide, can be selected for by cultivating the cells in the presence of the appropriate selectable agent.

[0142] The procedures used to ligate the elements described above to construct the recombinant expression vectors of the present invention are well known to one skilled in the art (see, e.g., Sambrook et al, 1989, supra).

Host Cells

[0143] The present invention also relates to recombinant host cells, comprising a polynucleotide of the present invention, which are advantageously used in the recombinant production of the polypeptides. A vector comprising a polynucleotide of the present invention is introduced into a host cell so that the vector is maintained as a chromosomal integrant or as a self-replicating extra-chromosomal vector as described earlier. The term "host cell" encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication. The choice of a host cell will to a large extent depend upon the gene encoding the polypeptide and its source.

[0144] The host cell may be a unicellular microorganism, e.g., a prokaryote, or a non-unicellular microorganism, e.g., a eukaryote.

[0145] Useful unicellular microorganisms are bacterial cells such as gram positive bacteria including, but not limited to, a *Bacillus* cell, e.g., *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus laetus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus stearothermophilus*, *Bacillus subtilis*, and *Bacillus thuringiensis*; or

a *Streptomyces* cell, e.g., *Streptomyces lividans* and *Streptomyces murinus*, or gram negative bacteria such as *E. coli* and *Pseudomonas* sp. In a preferred aspect, the bacterial host cell is a *Bacillus lentinus*, *Bacillus licheniformis*, *Bacillus stearothermophilus*, or *Bacillus subtilis* cell, in another preferred aspect, the *Bacillus* cell is an alkaiophilic *Bacillus*.

[0146] The introduction of a vector into a bacterial host cell may, for instance, be effected by protoplast transformation (see, e.g., Chang and Cohen, 1979, *Molecular General Genetics* 168: 111-115), using competent cells (see, e.g., Young and Spizizin, 1961, *Journal of Bacteriology* 81: 823-829, or Dubnau and Davidoff-Abelson, 1971, *Journal of Molecular Biology* 56: 209-221), electroporation (see, e.g., Shigekawa and Dower, 1988, *Biotechniques* 6: 742-751), or conjugation (see, e.g., Koehler and Thorne, 1987, *Journal of Bacteriology* 169: 5771-5278).

[0147] The host cell may also be a eukaryote, such as a mammalian, insect, plant, or fungal cell.

[0148] In a preferred aspect, the host cell is a fungal cell. "Fungi" as used herein includes the phyla Ascomycota, Basidiomycota, Chytridiomycota, and Zygomycota (as defined by Hawksworth et al., In, *Ainsworth and Bisby's Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK) as well as the Oomycota (as cited in Hawksworth et al., 1995, *supra*, page 171) and all mitosporic fungi (Hawksworth et al., 1995, *supra*).

[0149] In a more preferred aspect, the fungal host cell is a yeast cell. "Yeast" as used herein includes ascosporogenous yeast (Endomycetales), basidiosporogenous yeast, and yeast belonging to the Fungi imperfecti (Blastomycetes). Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in *Biology and Activities of Yeast* (Skinner, F. A., Passmore, S. M., and Davenport, R. R., eds, *Soc. App. Bacteriol. Symposium Series* No. 9, 1980).

[0150] In an even more preferred aspect, the yeast host cell is a *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* cell.

[0151] In a most preferred aspect, the yeast host cell is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluveri*, *Saccharomyces norbensis* or *Saccharomyces uvarum* cell. In another most preferred aspect, the yeast host cell is a *Kluyveromyces lactis* cell. In another most preferred aspect the yeast host cell is a *Yarrowia lipolytica* cell.

[0152] In another more preferred aspect, the fungal host cell is a filamentous fungal cell. "Filamentous fungi" include all filamentous forms of the subdivision Eumycota and Oomycota (as defined by Hawksworth et al., 1995, *supra*). The filamentous fungi are generally characterized by a mycelial wall composed of chitin, cellulose, glucan, chitosan, mannan, and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic, in contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative.

[0153] in an even more preferred aspect, the filamentous fungal host cell is an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Bjerkandera*, *Ceriporiopsis*, *Coprinus*, *Coriolus*, *Cryptococcus*, *Filobasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Phlebia*, *Piromyces*, *Pleurotus*, *Schizophyllum*, *Talaromyces*, *Thermovoascus*, *Thielavia*, *Tolypocladium*, *Trametes*, or *Trichoderma* cell.

[0154] In a most preferred aspect, the filamentous fungal host cell is an *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger* or *Aspergillus oryzae* cell. In another most preferred aspect, the filamentous fungal host cell is a *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarchchromum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, or *Fusarium venenatum* cell. In another most preferred aspect, the filamentous fungal host cell is a *Bjerkandera adusta*, *Ceriporiopsis aneirina*, *Ceriporiopsis aneirina*, *Ceriporiopsis caregeia*, *Ceriporiopsis gilvescens*, *Ceriporiopsis pannocinta*, *Ceriporiopsis rivulosa*, *Ceriporiopsis subrufa*, or *Ceriporiopsis subvermispora*, *Coprinus cinereus*, *Coriolus hirsutus*, *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Phlebia radiata*, *Pleurotus eryngii*, *Thielavia terrestris*, *Trametes villosa*, *Trametes versicolor*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* strain cell

[0155] Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known per se. Suitable procedures for transformation of *Aspergillus* and *Trichoderma* host cells are described in EP 238 023 and Yelton et al., 1984, *Proceedings of the National Academy of Sciences USA* 81: 1470-1474. Suitable methods for transforming *Fusarium* species are described by Maiardier et al., 1989, *Gene* 78: 147-156, and WO 96/00787. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J. N. and Simon, M. I., editors, *Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology*, Volume 194, pp 182-187, Academic Press, Inc., New York; Ito et al., 1983, *Journal of Bacteriology* 153: 163; and Hinnen et al., 1978, *Proceedings of the National Academy of Sciences USA* 75: 1920.

Methods of Production

[0156] The present invention also relates to methods for producing a polypeptide of the present invention, comprising (a) cultivating a cell, which in its wild-type form is capable of producing the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide. Preferably, the cell is of the genus *Aspergillus*, and more preferably *Aspergillus Oryzae*.

[0157] The present invention also relates to methods for producing a polypeptide of the present invention, comprising (a) cultivating a host cell under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

[0158] The present invention also relates to methods for producing a polypeptide of the present invention, comprising (a) cultivating a host cell under conditions conducive for production of the polypeptide, wherein the host cell comprises a mutant nucleotide sequence having at least one mutation in the mature polypeptide coding region of SEQ ID NO: 1, wherein the mutant nucleotide sequence encodes a polypeptide which is a lipase comprised by or comprising the polypeptide of SEQ ID NO: 2, and (b) recovering the polypeptide, in a preferred embodiment the nucleotide sequence encodes a polypeptide which is a lipase comprised

by or comprising the mature part of the polypeptide of SEQ ID NO: 2, and (b) recovering the polypeptide.

[0159] In the production methods of the present invention, the cells are cultivated in a nutrient medium suitable for production of the polypeptide using methods well known in the art. For example, the cell may be cultivated by shake flask cultivation, and small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions allowing the polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared according to published compositions (e.g., in catalogues of the American Type Culture Collection), if the polypeptide is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the polypeptide is not secreted, it can be recovered from cell lysates.

[0160] The polypeptides may be detected using methods known in the art that are specific for the polypeptides. These detection methods may include use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, an enzyme assay may be used to determine the activity of the polypeptide as described herein.

[0161] The resulting polypeptide may be recovered using methods known in the art. For example, the polypeptide may be recovered from the nutrient medium by conventional procedures including, but not limited to, centrifugation, filtration, extraction, spray-drying, evaporation, or precipitation.

[0162] The polypeptides of the present invention may be purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (e.g., preparative isoelectric focusing), differential solubility (e.g., ammonium sulfate precipitation), SDS-PAGE, or extraction (see, e.g., *Protein Purification*, J.-C. Janson and Lars Ryden, editors, VCH Publishers, New York, 1989).

Compositions

[0163] The present invention also relates to compositions comprising a polypeptide of the present invention. Preferably, the compositions are enriched in such a polypeptide. The term "enriched" indicates that the lipase activity of the composition has been increased, e.g., with an enrichment factor of 1.1.

[0164] The composition may comprise a polypeptide of the present invention as the major enzymatic component, e.g., a mono-component composition. Alternatively, the composition may comprise multiple enzymatic activities, such as an aminopeptidase, amylase, carboxypeptidase, catalase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, haloperoxidase, invertase, laccase, lipase, mannosidase, oxidase, pectinolytic enzyme, peptidoglytanase, peroxidase, phytase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transglutaminase, or xylanase. The additional enzyme(s) may be produced, for example, by a microorganism belonging to the genus *Aspergillus*, preferably *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, or *Aspergillus oryzae*; *Fusarium*, preferably *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium cul-*

morum, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sulphureum*, *Fusarium toruloseum*, *Fusarium trichothecionides*, or *Fusarium venenatum*; *Humicola*, preferably *Humicola insolens* or *Humicola lanuginosa*; or *Trichoderma*, preferably *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride*.

[0165] The polypeptide compositions may be prepared in accordance with methods known in the art and may be in the form of a liquid or a dry composition. For instance, the polypeptide composition may be in the form of a granulate or a microgranulate. The polypeptide to be included in the composition may be stabilized in accordance with methods known in the art.

[0166] Examples are given below of preferred uses of the polypeptide compositions of the invention. The dosage of the polypeptide composition of the invention and other conditions under which the composition is used may be determined on the basis of methods known in the art.

DETERGENT APPLICATIONS

[0167] The enzyme of the invention may be added to and thus become a component of a detergent composition.

[0168] The detergent composition of the invention may for example be formulated as a hand or machine laundry detergent composition including a laundry additive composition suitable for pre-treatment of stained fabrics and a rinse added fabric softener composition, or be formulated as a detergent composition for use in general household hard surface cleaning operations, or be formulated for hand or machine dish washing operations.

Enzymes

[0169] In a specific aspect, the invention provides a detergent additive comprising the enzyme of the invention. The detergent additive as well as the detergent composition may comprise one or more other enzymes such as a protease, a lipase, a cutinase, an amylase, a carbohydrase, a cellulase, a pectinase, a mannanase, an arabinase, a galactanase, a xylanase, an oxidase, e.g., a laccase, and/or a peroxidase.

[0170] In general the properties of the chosen enzyme(s) should be compatible with the selected detergent, (i.e. pH-optimum, compatibility with other enzymatic and non-enzymatic ingredients, etc.), and the enzyme(s) should be present in effective amounts.

Proteases:

[0171] Suitable proteases include those of animal, vegetable or microbial origin. Microbial origin is preferred. Chemically modified or protein engineered mutants are included. The protease may be a serine protease or a metallo protease, preferably an alkaline microbial protease or a trypsin-like protease. Examples of alkaline proteases are subtilisins, especially those derived from *Bacillus*, e.g., subtilisin Novo, subtilisin Carlsberg, subtilisin 309, subtilisin 147 and subtilisin 168 (described in WO 89/06279). Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the *Fusarium* protease described in WO 89/06270 and WO 94/25583.

[0172] Examples of useful proteases are the variants described in WO 92/19729, WO 98/20115, WO 98/20116, and WO 98/34946, especially the variants with substitutions in one or more of the following positions: 27, 36, 57, 68, 76,

87, 97, 101, 104, 106, 120, 123, 167, 170, 194, 206, 218, 222, 224, 235, 245, 252 and 274, and amongst other variants with the following mutations: (K27R, V104Y, N123S, T124A), (N76D, S103A, V104I), or (S101G, S103A, V104I, G159D, A232V, G236H, Q245R, N248D, N252K).

[0173] Preferred commercially available protease enzymes include Alcalase™, Savinase™, Primase™, Duralase™, Esperase™, Coronase™, Polarzyme™ and Kannase™ (Novozymes A/S), Maxatase™, Maxacal™, Maxapem™, Properase™, Purafect™, Purafect Prime™, Purafect OxP™, FN2, FN3 and FN4 (Genencor international Inc.).

Lipases:

[0174] Lipases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful lipases include lipases from *Humicola* (synonym *Thermomyces*), e.g. from *H. lanuginosa* (synonymous *T. lanuginosus*) as described in EP 258 068 and EP 305 216 or from *H. insolens* as described in WO 96/13580, a *Pseudomonas* lipase, e.g. from *P. alcaligenes* or *P. pseudoalcaligenes* (EP 218 272), *P. cepacia* (EP 331 376), *P. stutzeri* (GB 1,372,034), *P. fluorescens*, *Pseudomonas* sp. strain SD 705 (WO 95/06720 and WO 96/27002), *P. wisconsinensis* (WO 96/12012), a *Bacillus* lipase, e.g. from *B. subtilis* (Dartois et al., (1993), Biocfiemica et Biophysica Acta, 1131, 253-360), *B. stearothermophilus* (JP 64/744992) or *B. pumilus* (WO 91/16422).

[0175] Other examples are lipase variants such as those described in WO 92/05249., WO 94/01541, EP 407 225, EP 260 105, WO 95/35381, WO 96/00292, WO 95/30744, WO 94/25578, WO 95/14783, WO 95/22615, WO 97/04079 and WO 97/07202.

[0176] Other commercially available lipase enzymes include Lipolase™, Lipolase Ultra™ and Lipex™ (Novozymes A/S).

[0177] Preferred lipases are lipases of the present invention.

Amylases:

[0178] Suitable amylases (α and/or β) include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Amylases include, for example, α -amylases obtained from *Bacillus*, e.g. a special strain of *B. licheniformis*, described in more detail in GB 1,296,839.

[0179] Examples of useful amylases are the variants described in WO 94/02597, WO 94/18314, WO 96/23873, and WO 97/43424, especially the variants with substitutions in one or more of the following positions: 15, 23, 105, 106, 124, 128, 133, 154, 156, 181, 188, 190, 197, 202, 208, 209, 243, 264, 304, 305, 391, 408, and 444.

[0180] Commercially available amylases are Duramyl™, Termamyl™, Stainzyme™, Stainzyme Ultra™, Fungamyl™ and BAN™ (Novozymes A/S), Rapidase™ and Purastar™ (from Genencor International Inc.).

Cellulases:

[0181] Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Suitable cellulases include cellulases from the genera *Bacillus*, *Pseudomonas*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*, e.g. the fungal cellulases produced from *Humicola insolens*, *Myceliophthora thermophila* and *Fusarium oxysporum* disclosed in U.S. Pat. No. 4,435,307, U.S. Pat. No. 5,648,263, U.S. Pat. No. 5,691,178, U.S. Pat. No. 5,776,757 and WO 89/09259.

[0182] Especially suitable cellulases are the alkaline or neutral cellulases having colour care benefits. Examples of such cellulases are cellulases described in EP 0 495 257, EP 0 531 372, WO 96/11262, WO 96/29397, WO 98/08940. Other examples are cellulase variants such as those described in WO 94/07998, EP 0 531 315, U.S. Pat. No. 5,457,046, U.S. Pat. No. 5,686,593, U.S. Pat. No. 5,763,254, WO 95/24471, WO 98/12307 and PCT/DK98/00299.

[0183] Commercially available cellulases include RenozymeTM, CelluzymeTM, and CarezymeTM (Novozymes A/S), ClazinaseTM, and Puradax HATM (Genencor International Inc.), and KAC-500(B)TM (Kao Corporation).

Peroxidases/Oxidases:

[0184] Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from Coprinus, e.g. from C. cinereus, and variants thereof as those described in WO 93/24618, WO 95/10602, and WO 98/15257.

[0185] Commercially available peroxidases include GuardzymeTM (Novozymes A/S).

Detergents

[0186] The detergent enzyme(s) may be included in a detergent composition by adding separate additives containing one or more enzymes, or by adding a combined additive comprising all of these enzymes. A detergent additive of the invention, i.e. a separate additive or a combined additive, can be formulated e.g. as a granulate, a liquid, a slurry, etc. Preferred detergent additive formulations are granulates, in particular non-dusting granulates, liquids, in particular stabilized liquids, or slurries.

[0187] Non-dusting granulates may be produced, e.g., as disclosed in U.S. Pat. Nos. 4,106,991 and 4,661,452 and may optionally be coated by methods known in the art. Examples of waxy coating materials are poly(ethylene oxide) products (polyethylene glycol, PEG) with mean molar weights of 1000 to 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB1483591. Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid according to established methods. Protected enzymes may be prepared according to the method disclosed in EP 238,216.

[0188] The detergent composition of the invention may be in any convenient form, e.g., a bar, a

[0189] tablet, a powder, a granule, a paste or a liquid. A liquid detergent may be aqueous, typically containing up to 70% water and 0-30% organic solvent, or non-aqueous.

[0190] The detergent composition comprises one or more surfactants, which may be non-ionic including semi-polar and/or anionic and/or cationic and/or zwitterionic. The surfactants are typically present at a level of from 0% to 60% by weight.

[0191] When included therein the detergent will usually contain from about 0% to about 40% of an anionic surfactant such as linear alkylbenzenesulfonate, alpha-olefinsulfonate, alkyl sulfate (fatty alcohol sulfate), alcohol ethoxysulfate, secondary alkanesulfonate, alpha-sulfo fatty acid methyl ester, alky- or alkenylsuccinic acid or soap.

[0192] When included therein the detergent will usually contain from about 0% to about 40% of a non-ionic surfactant such as alcohol ethoxylate, nonylphenol ethoxylate, alkylpolyglycoside, alkyldimethylamineoxide, ethoxylated fatty acid monoethanolamide, fatty acid monoethanolamide, polyhydroxy alkyl fatty acid amide, or N-acyl N-alkyl derivatives of glucosamine ("glucamides").

[0193] The detergent may contain 0-65% of a detergent builder or complexing agent such as zeolite, diphosphate, triphosphate, phosphonate, carbonate, citrate, nitrilotriacetic acid, ethylenediaminetetraacetic acid, diethylenetriamine-pentaacetic acid, alkyl- or alkenylsuccinic acid, soluble silicates or layered silicates (e.g. SKS-6 from Hoechst).

[0194] The detergent may comprise one or more polymers. Examples are carboxymethylcellulose, poly(vinylpyrrolidone), poly(ethylene glycol), polyvinyl alcohol, poly(vinylpyridine-N-oxide), poly(vinylimidazole), polycarboxylates such as polyacrylates, maleic/acrylic acid copolymers and lauryl methacrylate/acrylic acid copolymers.

[0195] The detergent may contain a bleaching system which may comprise a H₂O₂ source such as perborate or percarbonate which may be combined with a peracid-forming bleach activator such as tetraacetyl ethylenediamine or nonanoyloxybenzenesulfonate. Alternatively, the bleaching system may comprise peroxyacids of e.g. the amide, imide, or sulfone type.

[0196] The enzyme(s) of the detergent composition of the invention may be stabilized using conventional stabilizing agents, e.g., a polyol such as propylene glycol or glycerol, a sugar or sugar alcohol, lactic acid, boric acid, or a boric acid derivative, e.g., an aromatic borate ester, or a phenyl boronic acid derivative such as 4-formylphenyl boronic acid, and the composition may be formulated as described in e.g. WO 92/19709 and WO 92/19708.

[0197] The detergent may also contain other conventional detergent ingredients such as e.g. fabric conditioners including clays, foam boosters, suds suppressors, anti-corrosion agents, soil-suspending agents, anti-soil redeposition agents, dyes, bactericides, optical brighteners, hydrotropes, tarnish inhibitors, or perfumes.

[0198] It is at present contemplated that in the detergent compositions any enzyme, in particular the enzyme of the invention, may be added in an amount corresponding to 0.001-100 mg of enzyme protein per litre of wash liquor, such as 0.01-50 mg or 0.03-30 mg, preferably 0.05-5 mg of enzyme protein per litre of wash liquor, in particular 0.1-1 mg of enzyme protein per litre of wash liquor.

[0199] The enzyme of the invention may additionally be incorporated in the detergent formulations disclosed in WO 97/07202, WO 04/041979, WO 04/074419, which is hereby incorporated as reference.

Uses

[0200] The present invention is also directed to methods for using the polypeptides having lipase activity.

[0201] The variants of the invention can be used in known applications of lipolytic enzymes by analogy with the prior art, e.g.:

[0202] A variant with lipase activity can be used in the pulp and paper industry, to remove pitch or to remove ink from used paper. WO 9213130, WO 9207138, JP 2180984 A, EP 374700.

[0203] A variant with phospholipase and/or galactolipase activity can be used in the preparation of dough, bread and cakes, e.g. to increase dough stability and dough handling properties, or to improve the elasticity of the bread or cake. WO 94/04035, WO 00/32758.

[0204] A variant with phospholipase activity can be used in a process for reducing the content of phospholipid in an edible oil. U.S. Pat. No. 5,264,367 (Metallgesellschaft, Röhm); K. Dahlke & H. Buchold, INFORM, 6 (12), 1284-91 (1995); H. Buchold, Fat Sci. Technol., 95 (8), 300-304 (1993); JP-A 2-153997 (Showa Sangyo); or EP 654,527 (Metallgesellschaft, Röhm).

[0205] A variant with lysophospholipase activity can be used to improve the filterability of an aqueous solution or slurry of carbohydrate origin, e.g. starch hydrolysate, especially a wheat starch hydrolysate. EP 219,269.

[0206] A variant with phospholipase activity can be used for the preparation of lyso-phospholipid, e.g. lyso-lecithin (EP 870840, JP-A 10-42884, JP-A 4-135456 or JP-A 2-49593) of for the production of mayonnaise (EP 628256, EP 398686 or EP 319064).

[0207] A variant with phospholipase activity may also be used in the processing of dairy and other food products, e.g. as described in EP 587,862 (Nestlé), EP 426,211 (Unilever), EP 166,284 (Nestlé), JP-A 57-189638 (Yakult) or U.S. Pat. No. 4,119,564 (Unilever).

[0208] A variant with phospholipase activity can be used in the leather industry. GB2233685, EP 505920.

[0209] A variant with lipase activity may be used for removing fatty matter containing hydrophobic esters (e.g. triglycerides) during the finishing of textiles. WO 93/13256.

[0210] The present invention is further described by the following examples which should not be construed as limiting the scope of the invention.

EXAMPLES

[0211] Chemicals used as buffers and substrates were commercial products of at least reagent grade.

Media and Solutions

[0212]

Product	Tradename
LAS:	Surfac PS
Zeolite A	Wessalith P

Materials

[0213]

Product	Supplier
EMPA221	EMPA St. Gallen, Lerchfeldstrasse 5, CH-9014 St. Gallen, Switzerland

Example 1

Production of Enzyme

[0214] A plasmid containing the gene encoding the lipase is constructed and transformed into a suitable host cell using standard methods of the art.

[0215] Fermentation is carried out as a fed-batch fermentation using a constant medium temperature of 34° C. and a start volume of 1.2 liter. The initial pH of the medium is set to 6.5. Once the pH has increased to 7.0 this value is maintained

through addition of 10% H3PO4. The level of dissolved oxygen in the medium is controlled by varying the agitation rate and using a fixed aeration rate of 1.0 liter air per liter medium per minute. The feed addition rate is maintained at a constant level during the entire fed-batch phase.

[0216] The batch medium contains maltose syrup as carbon source, urea and yeast extract as nitrogen source and a mixture of trace metals and salts. The feed added continuously during the fed-batch phase contains maltose syrup as carbon source whereas yeast extract and urea is added in order to assure a sufficient supply of nitrogen.

[0217] Purification of the lipase may be done by use of standard methods known in the art, e.g. by filtering the fermentation supernatant and subsequent hydrophobic chromatography and ion exchange chromatography, e.g. as described in EP 0 851 913 EP, Example 3.

Example 2

[0218] AMSA—Automated Mechanical Stress Assay - for calculation of RP

[0219] The enzyme variants of the present application are tested using the Automatic Mechanical Stress Assay (AMSA). With the AMSA test the wash performance of a large quantity of small volume enzyme-detergent solutions can be examined. The AMSA plate has a number of slots for test solutions and a lid firmly squeezing the textile swatch to be washed against all the slot openings. During the washing time, the plate, test solutions, textile and lid are vigorously shaken to bring the test solution in contact with the textile and apply mechanical stress. For further description see WO 02/42740 especially the paragraph "Special method embodiments" at page 23-24. The containers, which contain the detergent test solution, consist of cylindrical holes (6 mm diam, 10 mm depth) in a metal plate. The stained fabric (test material) lies on the top of the metal plates and is used as a lid and seal on the containers. Another metal plate lies on the top of the stained fabric to avoid any spillage from each container. The two metal plate together with the stained fabric are vibrated up and down at a frequency of 30 Hz with an amplitude of 2 mm.

[0220] The assay is conducted under the experimental conditions specified below:

TABLE 1

Test solution	0.5 g/l LAS 0.52 g/l Na2CO3 1.07 g/l Zeolite A 0.52 g/l Na3Citrate
Test solution volume	160 micro l
pH	As is (~9.9)
Wash time	20 minutes
Temperature	30° C.
Water hardness	15° dH
Enzyme concentration in test solution	Ratio of Ca ²⁺ /Mg ²⁺ /NaHCO ₃ : 4:1:7.5 0.125, 0.25, 0.50, 1.0 mg, ep/l
Drying	Wash performance: After washing the textile pieces is immediately flushed in tap water and air-dried at 85 C. in 5 min Odour: After washing the textile pieces is immediately flushed in tap water and dried at room temperature (20° C.) for 2 hours
Test material	Cream turmeric swatch as described below (EMPA221 used as cotton textile)

[0221] Cream-turmeric swatches were prepared by mixing 5 g of turmeric (Santa Maria, Denmark) with 100 g cream (38% fat, Aria, Denmark) at 50° C., the mixture was left at this temperature for about 20 minutes and filtered (50° C.) to remove any un-dissolved particles. The mixture is cooled to 20° C. and woven cotton swatches, EMPA221, were immersed in the cream-turmeric mixture and afterwards allowed to dry at room temperature over night and frozen until use. The preparation of cream-turmeric swatches is disclosed in the patent WO 2008/125437.

[0222] The performance of the enzyme variant is measured as the brightness of the colour of the textile samples washed with that specific enzyme variant. Brightness can also be expressed as the intensity of the light reflected from the textile sample when illuminated with white light. When the textile is stained the intensity of the reflected light is lower, than that of a clean textile. Therefore the intensity of the reflected light can be used to measure wash performance of an enzyme variant.

[0223] Color measurements are made with a professional flatbed scanner (PFU DL2400pro), which is used to capture an image of the washed textile samples. The scans are made with a resolution of 200 dpi and with an output color depth of 24 bits. In order to get accurate results, the scanner is frequently calibrated with a Kodak reflective IT8 target

[0224] To extract a value for the light intensity from the scanned images, a special designed software application is used (Novozymes Color Vector Analyzer). The program retrieves the 24 bit pixel values from the image and converts them into values for red, green and blue (RGB). The intensity value (Int) is calculated by adding the RGB values together as vectors and then taking the length of the resulting vector:

$$\text{Int} = \sqrt{r^2 + g^2 + b^2}$$

[0225] The wash performance (P) of the variants is calculated in accordance with the below formula:

$$P = \text{Int}(v) - \text{Int}(r)$$

where

Int(v) is the light intensity value of textile surface washed with enzyme, and

Int(r) is the light intensity value of textile surface washed without enzyme.

[0226] A relative performance score is given as the result of the AMSA wash in accordance with the definition:

[0227] Relative Performance scores (RP) are summing up the performances (P) of the tested

[0228] enzyme variants against the reference enzyme:

[0229] $RP = P(\text{test enzyme}) / P(\text{reference enzyme})$.

RPavg indicates the average relative performance compared to the reference enzyme at all three enzyme concentrations (0.125, 0.25, 0.5, 1.0 mg ep/l)

[0230] $RP_{avg} = \text{avg}(RP(0.125), RP(0.25), RP(0.5), RP(1.0))$

[0231] A variant is considered to exhibit improved wash performance, if it performs better than the reference.

[0232] In the context of the present invention the reference enzyme is the mature part of SEQ ID NO: 2 with the substitutions T231R+N233R.

Example 3

GC—Gas Chromatograph—For Calculation of Risk Factor

[0233] The butyric acid release from the lipase washed swatches were measured by Solid Phase Micro Extraction Gas Chromatography (SPME-GC) using the following method. Four textile pieces (5 mm in diameter), washed in the

specified solution in Table 1 containing 1 mg/L lipase, were transferred to a Gas Chromatograph (GC) vial. The samples were analysed on a Varian 3800 GC equipped with a Stabil-wax- DA w/Integra-Guard column (30 m, 0.32 mm ID and 0.25 micro-m df) and a Carboxen PDMS SPME fibre (75 micro-m). Each sample was preincubated for 10 min at 40° C. followed by 20 min sampling with the SPME fibre in the head-space over the textile pieces. The sample was subsequently injected onto the column (injector temperature=250° C.). Column flow=2 ml Helium/min. Column oven temperature gradient: 0 min=40° C., 2 min=40° C., 22 min=240° C., 32 min=240° C. The butyric acid was detected by FID detection and the amount of butyric acid was calculated based on a butyric acid standard curve.

[0234] The Risk Performance Odour, R, of a lipase variant is the ratio between the amount of released butyric acid from the lipase variant washed swatch and the amount of released butyric acid from a swatch washed with the mature part of the lipase of SEQ ID NO: 2, after both values have been corrected for the amount of released butyric acid from a non-lipase washed swatch. The risk (R) of the variants is calculated in accordance with the below formula:

[0235] $\text{Odour} = \text{measured in micro g butyric acid developed at 1 mg enzyme protein/l corrected for blank}$

[0236] $\text{Alpha}_{\text{test enzyme}} = \text{Odour}_{\text{test enzyme}} - \text{Blank}$

[0237] $\text{Alpha}_{\text{reference enzyme}} = \text{Odour}_{\text{reference enzyme}} - \text{Blank}$

[0238] $R = \text{Alpha}_{\text{test enzyme}} / \text{Alpha}_{\text{reference enzyme}}$

[0239] A variant is considered to exhibit reduced odor compared to the reference, if the R factor is lower than 1.

Example 4

Activity (LU) Relative to Absorbance at 280 nm

[0240] The activity of a lipase relative to the absorbance at 280 nm is determined by the following assay:

[0241] LIU/A280:

[0242] A substrate for lipase is prepared by emulsifying tributyrin (glycerin tributyrate) using gum Arabic as emulsifier. The hydrolysis of tributyrin at 30° C. at pH 7 or 9 is followed in a pH-stat titration experiment. One unit of lipase activity (1 LU) equals the amount of enzyme capable of releasing 1 micro mol butyric acid/min at pH 7.

[0243] The absorbance of the purified lipase at 280 nm is measured (A280) and the ratio LU/A280 is calculated. The relative LU/A280 is calculated as the LU/A280 of the variant divided by the LU/A280 of a reference enzyme. In the context of the present invention the reference enzyme is the mature part of SEQ ID NO: 2 with the mutations T231R and N233R.

Example 5

BR—Benefit Risk

[0244] The Benefit Risk factor describing the performance compared to the reduced risk for odour smell is thus defined as;

$$BR = RP_{avg} / R$$

[0245] A variant is considered to exhibit improved wash performance and reduced odor, if the BR factor is higher than 1.

[0246] Applying the above methods the following results were obtained:

TABLE 2

Variant	Mutations in mature part of polypeptide of SEQ ID NO: 2	RP	BR	LU/A280
1	I202G + T231R + N233R	0.84	1.41	not determined
2	I86V + L227G + T231R + N233R + P256K	1.08	1.52	1700
3	Q4V + S58N + V60S + T231R + N233R	0.87	1.73	1950
4	S58N + V60S + I90R + T231R, N233R	1.06	1.27	2250
5	I255Y + T231R + N233R	1.19	1.17	3600
6	I90A + T231R + N233R + I255V	1.13	1.14	2700
Reference	T231R + N233R	1.00	1.00	3650
7	G91A + E99K + T231R + N233R + Q249R + 270H + 271T + 272P + 273S + 274S + 275G + 276R + 277G + 278G + 279H + 280R	0.43	not determined	850
8	G91A + E99K + T231R, N233R + Q249R + 270H + 271T + 272P + 273S + 274S + 275G + 276R + 277G + 278G	0.13	not determined	500

[0247] The reference lipase and variants 7 and 8 in Table 2 are described in WO 2000/060063.

[0248] The invention described and claimed herein is not to be limited in scope by the specific aspects herein disclosed, since these aspects are intended as illustrations of several aspects of the invention. Any equivalent aspects are intended to be within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and

described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims, in the case of conflict, the present disclosure including definitions will control.

[0249] Various references are cited herein, the disclosures of which are incorporated by reference in their entireties.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 18

<210> SEQ ID NO 1

<211> LENGTH: 873

<212> TYPE: DNA

<213> ORGANISM: Thermomyces lanuginosus

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1)..(873)

<220> FEATURE:

<221> NAME/KEY: sig_peptide

<222> LOCATION: (1)..(51)

<220> FEATURE:

<221> NAME/KEY: propep

<222> LOCATION: (52)..(66)

<220> FEATURE:

<221> NAME/KEY: mat_peptide

<222> LOCATION: (67)..()

<400> SEQUENCE: 1

```

atg agg agc tcc ctt gtg ctg ttc ttt gtc tct gcg tgg acg gcc ttg      48
Met Arg Ser Ser Leu Val Leu Phe Phe Val Ser Ala Trp Thr Ala Leu
-20          -15          -10          -5

gcc agt cct att cgt cga gag gtc tcg cag gat ctg ttt aac cag ttc      96
Ala Ser Pro Ile Arg Arg Glu Val Ser Gln Asp Leu Phe Asn Gln Phe
-1   1          5          10

aat ctc ttt gca cag tat tct gca gcc gca tac tgc gga aaa aac aat      144
Asn Leu Phe Ala Gln Tyr Ser Ala Ala Ala Tyr Cys Gly Lys Asn Asn
15          20          25

gat gcc cca gct ggt aca aac att acg tgc acg gga aat gcc tgc ccc      192
Asp Ala Pro Ala Gly Thr Asn Ile Thr Cys Thr Gly Asn Ala Cys Pro

```

-continued

30	35	40	
gag gta gag aag gcg gat gca acg ttt ctc tac tcg ttt gaa gac tct			240
Glu Val Glu Lys Ala Asp Ala Thr Phe Leu Tyr Ser Phe Glu Asp Ser			
45	50	55	60
gga gtg ggc gat gtc acc ggc ttc ctt gct ctc gac aac acg aac aaa			288
Gly Val Gly Asp Val Thr Gly Phe Leu Ala Leu Asp Asn Thr Asn Lys			
65	70	75	
ttg atc gtc ctc tct ttc cgt ggc tct cgt tcc ata gag aac tgg atc			336
Leu Ile Val Leu Ser Phe Arg Gly Ser Arg Ser Ile Glu Asn Trp Ile			
80	85	90	
ggg aat ctt aac ttc gac ttg aaa gaa ata aat gac att tgc tcc ggc			384
Gly Asn Leu Asn Phe Asp Leu Lys Glu Ile Asn Asp Ile Cys Ser Gly			
95	100	105	
tgc agg gga cat gac ggc ttc act tcg tcc tgg agg tct gta gcc gat			432
Cys Arg Gly His Asp Gly Phe Thr Ser Ser Trp Arg Ser Val Ala Asp			
110	115	120	
acg tta agg cag aag gtg gag gat gct gtg agg gag cat ccc gac tat			480
Thr Leu Arg Gln Lys Val Glu Asp Ala Val Arg Glu His Pro Asp Tyr			
125	130	135	140
cgc gtg gtg ttt acc gga cat agc ttg ggt gca ttg gca act gtt			528
Arg Val Val Phe Thr Gly His Ser Leu Gly Gly Ala Leu Ala Thr Val			
145	150	155	
gcc gga gca gac ctg cgt gga aat ggg tat gat atc gac gtg ttt tca			576
Ala Gly Ala Asp Leu Arg Gly Asn Gly Tyr Asp Ile Asp Val Phe Ser			
160	165	170	
tat ggc gcc ccc cga gtc gga aac agg gct ttt gca gaa ttc ctg acc			624
Tyr Gly Ala Pro Arg Val Gly Asn Arg Ala Phe Ala Glu Phe Leu Thr			
175	180	185	
gta cag acc ggc gga aca ctc tac cgc att acc cac acc aat gat att			672
Val Gln Thr Gly Thr Leu Tyr Arg Ile Thr His Thr Asn Asp Ile			
190	195	200	
gtc cct aga ctc ccg ccg cgc gaa ttc ggt tac agc cat tct agc cca			720
Val Pro Arg Leu Pro Pro Arg Glu Phe Gly Tyr Ser His Ser Ser Pro			
205	210	215	220
gag tac tgg atc aaa tct gga acc ctt gtc ccc gtc acc cga aac gat			768
Glu Tyr Trp Ile Lys Ser Gly Thr Leu Val Pro Val Thr Arg Asn Asp			
225	230	235	
atc gtg aag ata gaa ggc atc gat gcc acc ggc ggc aat aac cag cct			816
Ile Val Lys Ile Glu Gly Ile Asp Ala Thr Gly Gly Asn Asn Gln Pro			
240	245	250	
aac att ccg gat atc cct gcg cac cta tgg tac ttc ggg tta att ggg			864
Asn Ile Pro Asp Ile Pro Ala His Leu Trp Tyr Phe Gly Leu Ile Gly			
255	260	265	
aca tgt ctt			873
Thr Cys Leu			
270			
<210> SEQ ID NO 2			
<211> LENGTH: 291			
<212> TYPE: PRT			
<213> ORGANISM: Thermomyces lanuginosus			
<400> SEQUENCE: 2			
Met Arg Ser Ser Leu Val Leu Phe Phe Val Ser Ala Trp Thr Ala Leu			
-20	-15	-10	-5
Ala Ser Pro Ile Arg Arg Glu Val Ser Gln Asp Leu Phe Asn Gln Phe			
-1	1	5	10
Asn Leu Phe Ala Gln Tyr Ser Ala Ala Tyr Cys Gly Lys Asn Asn			

-continued

15	20	25
Asp Ala Pro Ala Gly Thr Asn Ile Thr Cys Thr Gly Asn Ala Cys Pro		
30	35	40
Glu Val Glu Lys Ala Asp Ala Thr Phe Leu Tyr Ser Phe Glu Asp Ser		
45	50	55
Gly Val Gly Asp Val Thr Gly Phe Leu Ala Leu Asp Asn Thr Asn Lys		
65	70	75
Leu Ile Val Leu Ser Phe Arg Gly Ser Arg Ser Ile Glu Asn Trp Ile		
80	85	90
Gly Asn Leu Asn Phe Asp Leu Lys Glu Ile Asn Asp Ile Cys Ser Gly		
95	100	105
Cys Arg Gly His Asp Gly Phe Thr Ser Ser Trp Arg Ser Val Ala Asp		
110	115	120
Thr Leu Arg Gln Lys Val Glu Asp Ala Val Arg Glu His Pro Asp Tyr		
125	130	135
Arg Val Val Phe Thr Gly His Ser Leu Gly Gly Ala Leu Ala Thr Val		
145	150	155
Ala Gly Ala Asp Leu Arg Gly Asn Gly Tyr Asp Ile Asp Val Phe Ser		
160	165	170
Tyr Gly Ala Pro Arg Val Gly Asn Arg Ala Phe Ala Glu Phe Leu Thr		
175	180	185
Val Gln Thr Gly Gly Thr Leu Tyr Arg Ile Thr His Thr Asn Asp Ile		
190	195	200
Val Pro Arg Leu Pro Pro Arg Glu Phe Gly Tyr Ser His Ser Ser Pro		
205	210	215
220		
Glu Tyr Trp Ile Lys Ser Gly Thr Leu Val Pro Val Thr Arg Asn Asp		
225	230	235
Ile Val Lys Ile Glu Gly Ile Asp Ala Thr Gly Gly Asn Asn Gln Pro		
240	245	250
Asn Ile Pro Asp Ile Pro Ala His Leu Trp Tyr Phe Gly Leu Ile Gly		
255	260	265
Thr Cys Leu		
270		
<210> SEQ ID NO 3		
<211> LENGTH: 265		
<212> TYPE: PRT		
<213> ORGANISM: Absidia reflexa		
<400> SEQUENCE: 3		
Ser Ser Ser Thr Gln Asp Tyr Arg Ile Ala Ser Glu Ala Glu Ile		
1	5	10
15		
Lys Ala His Thr Phe Tyr Thr Ala Leu Ser Ala Asn Ala Tyr Cys Arg		
20	25	30
Thr Val Ile Pro Gly Gly Arg Trp Ser Cys Pro His Cys Gly Val Ala		
35	40	45
Ser Asn Leu Gln Ile Thr Lys Thr Phe Ser Thr Leu Ile Thr Asp Thr		
50	55	60
Asn Val Leu Val Ala Val Gly Glu Lys Glu Lys Thr Ile Tyr Val Val		
65	70	75
80		
Phe Arg Gly Thr Ser Ser Ile Arg Asn Ala Ile Ala Asp Ile Val Phe		
85	90	95
Val Pro Val Asn Tyr Pro Pro Val Asn Gly Ala Lys Val His Lys Gly		

-continued

100	105	110	
Phe Leu Asp Ser Tyr Asn Glu Val Gln Asp Lys Leu Val Ala Glu Val			
115	120	125	
Lys Ala Gln Leu Asp Arg His Pro Gly Tyr Lys Ile Val Val Thr Gly			
130	135	140	
His Ser Leu Gly Gly Ala Thr Ala Val Leu Ser Ala Leu Asp Leu Tyr			
145	150	155	160
His His Gly His Ala Asn Ile Glu Ile Tyr Thr Gln Gly Gln Pro Arg			
165	170	175	
Ile Gly Thr Pro Ala Phe Ala Asn Tyr Val Ile Gly Thr Lys Ile Pro			
180	185	190	
Tyr Gln Arg Leu Val His Glu Arg Asp Ile Val Pro His Leu Pro Pro			
195	200	205	
Gly Ala Phe Gly Phe Leu His Ala Gly Glu Glu Phe Trp Ile Met Lys			
210	215	220	
Asp Ser Ser Leu Arg Val Cys Pro Asn Gly Ile Glu Thr Asp Asn Cys			
225	230	235	240
Ser Asn Ser Ile Val Pro Phe Thr Ser Val Ile Asp His Leu Ser Tyr			
245	250	255	
Leu Asp Met Asn Thr Gly Leu Cys Leu			
260	265		

<210> SEQ ID NO 4
 <211> LENGTH: 264
 <212> TYPE: PRT
 <213> ORGANISM: Absidia corymbifera

<400> SEQUENCE: 4

Ser Ser Ser Thr Gln Asp Tyr Arg Ile Ala Ser Glu Ala Glu Ile Lys			
1	5	10	15
Ala His Thr Phe Tyr Thr Ala Leu Ser Ala Asn Ala Tyr Cys Arg Thr			
20	25	30	
Val Ile Pro Gly Gly Gln Trp Ser Cys Pro His Cys Asp Val Ala Pro			
35	40	45	
Asn Leu Asn Ile Thr Lys Thr Phe Thr Thr Leu Ile Thr Asp Thr Asn			
50	55	60	
Val Leu Val Ala Val Gly Glu Asn Glu Lys Thr Ile Tyr Val Val Phe			
65	70	75	80
Arg Gly Thr Ser Ser Ile Arg Asn Ala Ile Ala Asp Ile Val Phe Val			
85	90	95	
Pro Val Asn Tyr Pro Pro Val Asn Gly Ala Lys Val His Lys Gly Phe			
100	105	110	
Leu Asp Ser Tyr Asn Glu Val Gln Asp Lys Leu Val Ala Glu Val Lys			
115	120	125	
Ala Gln Leu Asp Arg His Pro Gly Tyr Lys Ile Val Val Thr Gly His			
130	135	140	
Ser Leu Gly Gly Ala Thr Ala Val Leu Ser Ala Leu Asp Leu Tyr His			
145	150	155	160
His Gly His Asp Asn Ile Glu Ile Tyr Thr Gln Gly Gln Pro Arg Ile			
165	170	175	
Gly Thr Pro Glu Phe Ala Asn Tyr Val Ile Gly Thr Lys Ile Pro Tyr			
180	185	190	

-continued

Gln	Arg	Leu	Val	Asn	Glu	Arg	Asp	Ile	Val	Pro	His	Leu	Pro	Pro	Gly
195					200							205			
Ala	Phe	Gly	Phe	Leu	His	Ala	Gly	Glu	Glu	Phe	Trp	Ile	Met	Lys	Asp
210					215						220				
Ser	Ser	Leu	Arg	Val	Cys	Pro	Asn	Gly	Ile	Glu	Thr	Asp	Asn	Cys	Ser
225					230				235			240			
Asn	Ser	Ile	Val	Pro	Phe	Thr	Ser	Val	Ile	Asp	His	Leu	Ser	Tyr	Leu
									250			255			
Asp	Met	Asn	Thr	Gly	Leu	Cys	Leu								
					260										

<210> SEQ ID NO 5

<211> LENGTH: 269

<212> TYPE: PRT

<213> ORGANISM: Rhizomucor miehei

<400> SEQUENCE: 5

Ser	Ile	Asp	Gly	Gly	Ile	Arg	Ala	Ala	Thr	Ser	Gln	Glu	Ile	Asn	Glu
1					5				10			15			
Leu	Thr	Tyr	Tyr	Thr	Leu	Ser	Ala	Asn	Ser	Tyr	Cys	Arg	Thr	Val	
					20				25			30			
Ile	Pro	Gly	Ala	Thr	Trp	Asp	Cys	Ile	His	Cys	Asp	Ala	Thr	Glu	Asp
	35				40				45						
Leu	Lys	Ile	Ile	Lys	Thr	Trp	Ser	Thr	Leu	Ile	Tyr	Asp	Thr	Asn	Ala
	50				55				60						
Met	Val	Ala	Arg	Gly	Asp	Ser	Glu	Lys	Thr	Ile	Tyr	Ile	Val	Phe	Arg
	65				70				75				80		
Gly	Ser	Ser	Ser	Ile	Arg	Asn	Trp	Ile	Ala	Asp	Leu	Thr	Phe	Val	Pro
	85				90				95						
Val	Ser	Tyr	Pro	Pro	Val	Ser	Gly	Thr	Lys	Val	His	Lys	Gly	Phe	Leu
	100				105				110						
Asp	Ser	Tyr	Gly	Glu	Val	Gln	Asn	Glu	Leu	Val	Ala	Thr	Val	Leu	Asp
	115				120				125						
Gln	Phe	Lys	Gln	Tyr	Pro	Ser	Tyr	Lys	Val	Ala	Val	Thr	Gly	His	Ser
	130				135				140						
Leu	Gly	Gly	Ala	Thr	Ala	Leu	Leu	Cys	Ala	Leu	Asp	Leu	Tyr	Gln	Arg
	145				150				155				160		
Glu	Glu	Gly	Leu	Ser	Ser	Ser	Asn	Leu	Phe	Leu	Tyr	Thr	Gln	Gly	Gln
	165				170				175						
Pro	Arg	Val	Gly	Asp	Pro	Ala	Phe	Ala	Asn	Tyr	Val	Val	Ser	Thr	Gly
	180				185				190						
Ile	Pro	Tyr	Arg	Arg	Thr	Val	Asn	Glu	Arg	Asp	Ile	Val	Pro	His	Leu
	195				200				205						
Pro	Pro	Ala	Ala	Phe	Gly	Phe	Leu	His	Ala	Gly	Glu	Glu	Tyr	Trp	Ile
	210				215				220						
Thr	Asp	Asn	Ser	Pro	Glu	Thr	Val	Gln	Val	Cys	Thr	Ser	Asp	Leu	Glu
	225				230				235				240		
Thr	Ser	Asp	Cys	Ser	Asn	Ser	Ile	Val	Pro	Phe	Thr	Ser	Val	Leu	Asp
	245				250				255						
His	Leu	Ser	Tyr	Phe	Gly	Ile	Asn	Thr	Gly	Leu	Cys	Thr			
	260				265										

<210> SEQ ID NO 6

-continued

<211> LENGTH: 271
 <212> TYPE: PRT
 <213> ORGANISM: Rhizopus oryzae

 <400> SEQUENCE: 6

```

Ser Ala Ser Asp Gly Gly Lys Val Val Ala Ala Thr Thr Ala Gln Ile
1           5           10          15

Gln Glu Phe Thr Lys Tyr Ala Gly Ile Ala Ala Thr Ala Tyr Cys Arg
20          25          30

Ser Val Val Pro Gly Asn Lys Trp Asp Cys Val Gln Cys Gln Lys Trp
35          40          45

Val Pro Asp Gly Lys Ile Ile Thr Thr Phe Thr Ser Leu Leu Ser Asp
50          55          60

Thr Asn Gly Tyr Val Leu Arg Ser Asp Lys Gln Lys Thr Ile Tyr Leu
65          70          75          80

Val Phe Arg Gly Thr Asn Ser Phe Arg Ser Ala Ile Thr Asp Ile Val
85          90          95

Phe Asn Phe Ser Asp Tyr Lys Pro Val Lys Gly Ala Lys Val His Ala
100         105         110

Gly Phe Leu Ser Ser Tyr Glu Gln Val Val Asn Asp Tyr Phe Pro Val
115         120         125

Val Gln Glu Gln Leu Thr Ala His Pro Thr Tyr Lys Val Ile Val Thr
130         135         140

Gly His Ser Leu Gly Gly Ala Gln Ala Leu Leu Ala Gly Met Asp Leu
145         150         155         160

Tyr Gln Arg Glu Pro Arg Leu Ser Pro Lys Asn Leu Ser Ile Phe Thr
165         170         175

Val Gly Gly Pro Arg Val Gly Asn Pro Thr Phe Ala Tyr Tyr Val Glu
180         185         190

Ser Thr Gly Ile Pro Phe Gln Arg Thr Val His Lys Arg Asp Ile Val
195         200         205

Pro His Val Pro Pro Gln Ser Phe Gly Phe Leu His Pro Gly Val Glu
210         215         220

Ser Trp Ile Lys Ser Gly Thr Ser Asn Val Gln Ile Cys Thr Ser Glu
225         230         235         240

Ile Glu Thr Lys Asp Cys Ser Asn Ser Ile Val Pro Phe Thr Ser Ile
245         250         255

Leu Asp His Leu Ser Tyr Phe Asp Ile Asn Glu Gly Ser Cys Leu
260         265         270
  
```

<210> SEQ ID NO 7
 <211> LENGTH: 267
 <212> TYPE: PRT
 <213> ORGANISM: Aspergillus niger

 <400> SEQUENCE: 7

```

Thr Ala Gly His Ala Leu Ala Ala Ser Thr Gln Gly Ile Ser Glu Asp
1           5           10          15

Leu Tyr Ser Arg Leu Val Glu Met Ala Thr Ile Ser Gln Ala Ala Tyr
20          25          30

Ala Asp Leu Cys Asn Ile Pro Ser Thr Ile Ile Lys Gly Glu Lys Ile
35          40          45

Tyr Asn Ser Gln Thr Asp Ile Asn Gly Trp Ile Leu Arg Asp Asp Ser
50          55          60
  
```

-continued

Ser Lys Glu Ile Ile Thr Val Phe Arg Gly Thr Gly Ser Asp Thr Asn
 65 70 75 80
 Leu Gln Leu Asp Thr Asn Tyr Thr Leu Thr Pro Phe Asp Thr Leu Pro
 85 90 95
 Gln Cys Asn Gly Cys Glu Val His Gly Gly Tyr Tyr Ile Gly Trp Val
 100 105 110
 Ser Val Gln Asp Gln Val Glu Ser Leu Val Lys Gln Gln Val Ser Gln
 115 120 125
 Tyr Pro Asp Tyr Ala Leu Thr Val Thr Gly His Ser Leu Gly Ala Ser
 130 135 140
 Leu Ala Ala Leu Thr Ala Ala Gln Leu Ser Ala Thr Tyr Asp Asn Ile
 145 150 155 160
 Arg Leu Tyr Thr Phe Gly Glu Pro Arg Ser Gly Asn Gln Ala Phe Ala
 165 170 175
 Ser Tyr Met Asn Asp Ala Phe Gln Ala Ser Ser Pro Asp Thr Thr Gln
 180 185 190
 Tyr Phe Arg Val Thr His Ala Asn Asp Gly Ile Pro Asn Leu Pro Pro
 195 200 205
 Val Glu Gln Gly Tyr Ala His Gly Gly Val Glu Tyr Trp Ser Val Asp
 210 215 220
 Pro Tyr Ser Ala Gln Asn Thr Phe Val Cys Thr Gly Asp Glu Val Gln
 225 230 235 240
 Cys Cys Glu Ala Gln Gly Gly Gln Gly Val Asn Asn Ala His Thr Thr
 245 250 255
 Tyr Phe Gly Met Thr Ser Gly Ala Cys Thr Trp
 260 265

<210> SEQ ID NO 8
 <211> LENGTH: 266
 <212> TYPE: PRT
 <213> ORGANISM: Aspergillus tubingensis

<400> SEQUENCE: 8

Thr Ala Gly His Ala Leu Ala Ala Ser Thr Gln Gly Ile Ser Glu Asp
 1 5 10 15
 Leu Tyr Ser Arg Leu Val Glu Met Ala Thr Ile Ser Gln Ala Ala Tyr
 20 25 30
 Ala Asp Leu Cys Asn Ile Pro Ser Thr Ile Ile Lys Gly Glu Lys Ile
 35 40 45
 Tyr Asn Ser Gln Thr Asp Ile Asn Gly Trp Ile Leu Arg Asp Asp Ser
 50 55 60
 Ser Lys Glu Ile Ile Thr Val Phe Arg Gly Thr Gly Ser Asp Thr Asn
 65 70 75 80
 Leu Gln Leu Asp Thr Asn Tyr Thr Leu Thr Pro Phe Asp Thr Leu Pro
 85 90 95
 Gln Cys Asn Ser Cys Glu Val His Gly Gly Tyr Tyr Ile Gly Trp Ile
 100 105 110
 Ser Val Gln Asp Gln Val Glu Ser Leu Val Gln Gln Val Ser Gln
 115 120 125
 Phe Pro Asp Tyr Ala Leu Thr Val Thr Gly His Ser Leu Gly Ala Ser
 130 135 140
 Leu Ala Ala Leu Thr Ala Ala Gln Leu Ser Ala Thr Tyr Asp Asn Ile

-continued

145	150	155	160
Arg Leu Tyr Thr Phe Gly Glu Pro Arg Ser Asn Gln Ala Phe Ala Ser			
165	170	175	
Tyr Met Asn Asp Ala Phe Gln Ala Ser Ser Pro Asp Thr Thr Gln Tyr			
180	185	190	
Phe Arg Val Thr His Ala Asn Asp Gly Ile Pro Asn Leu Pro Pro Ala			
195	200	205	
Asp Glu Gly Tyr Ala His Gly Val Val Glu Tyr Trp Ser Val Asp Pro			
210	215	220	
Tyr Ser Ala Gln Asn Thr Phe Val Cys Thr Gly Asp Glu Val Gln Cys			
225	230	235	240
Cys Glu Ala Gln Gly Gln Gly Val Asn Asn Ala His Thr Thr Tyr			
245	250	255	
Phe Gly Met Thr Ser Gly His Cys Thr Trp			
260	265		

<210> SEQ ID NO 9

<211> LENGTH: 276

<212> TYPE: PRT

<213> ORGANISM: Fusarium oxysporum

<400> SEQUENCE: 9

Ala Val Gly Val Thr Thr Asp Phe Ser Asn Phe Lys Phe Tyr Ile			
1	5	10	15
Gln His Gly Ala Ala Ala Tyr Cys Asn Ser Glu Ala Ala Gly Ser			
20	25	30	
Lys Ile Thr Cys Ser Asn Asn Gly Cys Pro Thr Val Gln Gly Asn Gly			
35	40	45	
Ala Thr Ile Val Thr Ser Phe Val Gly Ser Lys Thr Gly Ile Gly Gly			
50	55	60	
Tyr Val Ala Thr Asp Ser Ala Arg Lys Glu Ile Val Val Ser Phe Arg			
65	70	75	80
Gly Ser Ile Asn Ile Arg Asn Trp Leu Thr Asn Leu Asp Phe Gly Gln			
85	90	95	
Glu Asp Cys Ser Leu Val Ser Gly Cys Gly Val His Ser Gly Phe Gln			
100	105	110	
Arg Ala Trp Asn Glu Ile Ser Ser Gln Ala Thr Ala Val Ala Ser			
115	120	125	
Ala Arg Lys Ala Asn Pro Ser Phe Asn Val Ile Ser Thr Gly His Ser			
130	135	140	
Leu Gly Gly Ala Val Ala Val Leu Ala Ala Asn Leu Arg Val Gly			
145	150	155	160
Gly Thr Pro Val Asp Ile Tyr Thr Tyr Gly Ser Pro Arg Val Gly Asn			
165	170	175	
Ala Gln Leu Ser Ala Phe Val Ser Asn Gln Ala Gly Gly Glu Tyr Arg			
180	185	190	
Val Thr His Ala Asp Asp Pro Val Pro Arg Leu Pro Pro Leu Ile Phe			
195	200	205	
Gly Tyr Arg His Thr Thr Pro Glu Phe Trp Leu Ser Gly Gly Gly			
210	215	220	
Asp Lys Val Asp Tyr Thr Ile Ser Asp Val Lys Val Cys Glu Gly Ala			
225	230	235	240

-continued

Ala Asn Leu Gly Cys Asn Gly Gly Thr Leu Gly Leu Asp Ile Ala Ala
245 250 255

His Leu His Tyr Phe Gln Ala Thr Asp Ala Cys Asn Ala Gly Gly Phe
260 265 270

Ser Trp Arg Arg
275

<210> SEQ_ID NO 10

<211> LENGTH: 273

<212> TYPE: PRT

<213> ORGANISM: Fusarium heterosporum

<400> SEQUENCE: 10

Thr Val Thr Thr Gln Asp Leu Ser Asn Phe Arg Phe Tyr Leu Gln His
1 5 10 15

Ala Asp Ala Ala Tyr Cys Asn Phe Asn Thr Ala Val Gly Lys Pro Val
20 25 30

His Cys Ser Ala Gly Asn Cys Pro Asp Ile Glu Lys Asp Ala Ala Ile
35 40 45

Val Val Gly Ser Val Val Gly Thr Lys Thr Gly Ile Gly Ala Tyr Val
50 55 60

Ala Thr Asp Asn Ala Arg Lys Glu Ile Val Val Ser Val Arg Gly Ser
65 70 75 80

Ile Asn Val Arg Asn Trp Ile Thr Asn Phe Asn Phe Gly Gln Lys Thr
85 90 95

Cys Asp Leu Val Ala Gly Cys Gly Val His Thr Gly Phe Leu Asp Ala
100 105 110

Trp Glu Glu Val Ala Ala Asn Val Lys Ala Ala Val Ser Ala Ala Lys
115 120 125

Thr Ala Asn Pro Thr Phe Lys Phe Val Val Thr Gly His Ser Leu Gly
130 135 140

Gly Ala Val Ala Thr Ile Ala Ala Tyr Leu Arg Lys Asp Gly Phe
145 150 155 160

Pro Phe Asp Leu Tyr Thr Tyr Gly Ser Pro Arg Val Gly Asn Asp Phe
165 170 175

Phe Ala Asn Phe Val Thr Gln Gln Thr Gly Ala Glu Tyr Arg Val Thr
180 185 190

His Gly Asp Asp Pro Val Pro Arg Leu Pro Pro Ile Val Phe Gly Tyr
195 200 205

Arg His Thr Ser Pro Glu Tyr Trp Leu Asn Gly Gly Pro Leu Asp Lys
210 215 220

Asp Tyr Thr Val Thr Glu Ile Lys Val Cys Glu Gly Ile Ala Asn Val
225 230 235 240

Met Cys Asn Gly Gly Thr Ile Gly Leu Asp Ile Leu Ala His Ile Thr
245 250 255

Tyr Phe Gln Ser Met Ala Thr Cys Ala Pro Ile Ala Ile Pro Trp Lys
260 265 270

Arg

<210> SEQ_ID NO 11

<211> LENGTH: 278

<212> TYPE: PRT

<213> ORGANISM: Aspergillus oryzae

-continued

<400> SEQUENCE: 11

```

Asp Ile Pro Thr Thr Gln Leu Glu Asp Phe Lys Phe Trp Val Gln Tyr
1          5          10          15

Ala Ala Ala Thr Tyr Cys Pro Asn Asn Tyr Val Ala Lys Asp Gly Glu
20         25          30

Lys Leu Asn Cys Ser Val Gly Asn Cys Pro Asp Val Glu Ala Ala Gly
35         40          45

Ser Thr Val Lys Leu Ser Phe Ser Asp Asp Thr Ile Thr Asp Thr Ala
50         55          60

Gly Phe Val Ala Val Asp Asn Thr Asn Lys Ala Ile Val Val Ala Phe
65         70          75          80

Arg Gly Ser Tyr Ser Ile Arg Asn Trp Val Thr Asp Ala Thr Phe Pro
85         90          95

Gln Thr Asp Pro Gly Leu Cys Asp Gly Cys Lys Ala Glu Leu Gly Phe
100        105         110

Trp Thr Ala Trp Lys Val Val Arg Asp Arg Ile Ile Lys Thr Leu Asp
115        120         125

Glu Leu Lys Pro Glu His Ser Asp Tyr Lys Ile Val Val Val Gly His
130        135         140

Ser Leu Gly Ala Ala Ile Ala Ser Leu Ala Ala Ala Asp Leu Arg Thr
145        150         155         160

Lys Asn Tyr Asp Ala Ile Leu Tyr Ala Tyr Ala Ala Pro Arg Val Ala
165        170         175

Asn Lys Pro Leu Ala Glu Phe Ile Thr Asn Gln Gly Asn Asn Tyr Arg
180        185         190

Phe Thr His Asn Asp Asp Pro Val Pro Lys Leu Pro Leu Leu Thr Met
195        200         205

Gly Tyr Val His Ile Ser Pro Glu Tyr Tyr Ile Thr Ala Pro Asp Asn
210        215         220

Thr Thr Val Thr Asp Asn Gln Val Thr Val Leu Asp Gly Tyr Val Asn
225        230         235         240

Phe Lys Gly Asn Thr Gly Thr Ser Gly Gly Leu Pro Asp Leu Leu Ala
245        250         255

Phe His Ser His Val Trp Tyr Phe Ile His Ala Asp Ala Cys Lys Gly
260        265         270

Pro Gly Leu Pro Leu Arg
275

```

<210> SEQ ID NO 12

<211> LENGTH: 278

<212> TYPE: PRT

<213> ORGANISM: Penicillium camemberti

<400> SEQUENCE: 12

```

Asp Val Ser Thr Ser Glu Leu Asp Gln Phe Glu Phe Trp Val Gln Tyr
1          5          10          15

Ala Ala Ala Ser Tyr Tyr Glu Ala Asp Tyr Thr Ala Gln Val Gly Asp
20         25          30

Lys Leu Ser Cys Ser Lys Gly Asn Cys Pro Glu Val Glu Ala Thr Gly
35         40          45

Ala Thr Val Ser Tyr Asp Phe Ser Asp Ser Thr Ile Thr Asp Thr Ala
50         55          60

```

-continued

Gly Tyr Ile Ala Val Asp His Thr Asn Ser Ala Val Val Leu Ala Phe
65 70 75 80

Arg Gly Ser Tyr Ser Val Arg Asn Trp Val Ala Asp Ala Thr Phe Val
85 90 95

His Thr Asn Pro Gly Leu Cys Asp Gly Cys Leu Ala Glu Leu Gly Phe
100 105 110

Trp Ser Ser Trp Lys Leu Val Arg Asp Asp Ile Ile Lys Glu Leu Lys
115 120 125

Glu Val Val Ala Gln Asn Pro Asn Tyr Glu Leu Val Val Val Gly His
130 135 140

Ser Leu Gly Ala Ala Val Ala Thr Leu Ala Ala Thr Asp Leu Arg Gly
145 150 155 160

Lys Gly Tyr Pro Ser Ala Lys Leu Tyr Ala Tyr Ala Ser Pro Arg Val
165 170 175

Gly Asn Ala Ala Leu Ala Lys Tyr Ile Thr Ala Gln Gly Asn Asn Phe
180 185 190

Arg Phe Thr His Thr Asn Asp Pro Val Pro Lys Leu Pro Leu Leu Ser
195 200 205

Met Gly Tyr Val His Val Ser Pro Glu Tyr Trp Ile Thr Ser Pro Asn
210 215 220

Asn Ala Thr Val Ser Thr Ser Asp Ile Lys Val Ile Asp Gly Asp Val
225 230 235 240

Ser Phe Asp Gly Asn Thr Gly Thr Gly Leu Pro Leu Leu Thr Asp Phe
245 250 255

Glu Ala His Ile Trp Tyr Phe Val Gln Val Asp Ala Gly Lys Gly Pro
260 265 270

Gly Leu Pro Phe Lys Arg
275

<210> SEQ ID NO 13
<211> LENGTH: 270
<212> TYPE: PRT
<213> ORGANISM: Aspergillus foetidus

<400> SEQUENCE: 13

Ser Val Ser Thr Ser Thr Leu Asp Glu Leu Gln Leu Phe Ala Gln Trp
1 5 10 15

Ser Ala Ala Ala Tyr Cys Ser Asn Asn Ile Asp Ser Lys Asp Ser Asn
20 25 30

Leu Thr Cys Thr Ala Asn Ala Cys Pro Ser Val Glu Glu Ala Ser Thr
35 40 45

Thr Met Leu Leu Glu Phe Asp Leu Thr Asn Asp Phe Gly Gly Thr Ala
50 55 60

Gly Phe Leu Ala Ala Asp Asn Thr Asn Lys Arg Leu Val Val Ala Phe
65 70 75 80

Arg Gly Ser Ser Thr Ile Glu Asn Trp Ile Ala Asn Leu Asp Phe Ile
85 90 95

Leu Glu Asp Asn Asp Asp Leu Cys Thr Gly Cys Lys Val His Thr Gly
100 105 110

Phe Trp Lys Ala Trp Glu Ser Ala Ala Asp Glu Leu Thr Ser Lys Ile
115 120 125

Lys Ser Ala Met Ser Thr Tyr Ser Gly Tyr Thr Leu Tyr Phe Thr Gly
130 135 140

-continued

His Ser Leu Gly Gly Ala Leu Ala Thr Leu Gly Ala Thr Val Leu Arg
 145 150 155 160
 Asn Asp Gly Tyr Ser Val Glu Leu Tyr Thr Tyr Gly Cys Pro Arg Ile
 165 170 175
 Gly Asn Tyr Ala Leu Ala Glu His Ile Thr Ser Gln Gly Ser Gly Ala
 180 185 190
 Asn Phe Arg Val Thr His Leu Asn Asp Ile Val Pro Arg Val Pro Pro
 195 200 205
 Met Asp Phe Gly Phe Ser Gln Pro Ser Pro Glu Tyr Trp Ile Thr Ser
 210 215 220
 Gly Asn Gly Ala Ser Val Thr Ala Ser Asp Ile Glu Val Ile Glu Gly
 225 230 235 240
 Ile Asn Ser Thr Ala Gly Asn Ala Gly Glu Ala Thr Val Ser Val Leu
 245 250 255
 Ala His Leu Trp Tyr Phe Phe Ala Ile Ser Glu Cys Leu Leu
 260 265 270

<210> SEQ_ID NO 14
 <211> LENGTH: 270
 <212> TYPE: PRT
 <213> ORGANISM: Aspergillus niger
 <400> SEQUENCE: 14

Ser Val Ser Thr Ser Thr Leu Asp Glu Leu Gln Leu Phe Ser Gln Trp
 1 5 10 15
 Ser Ala Ala Ala Tyr Cys Ser Asn Asn Ile Asp Ser Asp Asp Ser Asn
 20 25 30
 Val Thr Cys Thr Ala Asp Ala Cys Pro Ser Val Glu Glu Ala Ser Thr
 35 40 45
 Lys Met Leu Leu Glu Phe Asp Leu Thr Asn Asn Phe Gly Gly Thr Ala
 50 55 60
 Gly Phe Leu Ala Ala Asp Asn Thr Asn Lys Arg Leu Val Val Ala Phe
 65 70 75 80
 Arg Gly Ser Ser Thr Ile Lys Asn Trp Ile Ala Asp Leu Asp Phe Ile
 85 90 95
 Leu Gln Asp Asn Asp Asp Leu Cys Thr Gly Cys Lys Val His Thr Gly
 100 105 110
 Phe Trp Lys Ala Trp Glu Ala Ala Asp Asn Leu Thr Ser Lys Ile
 115 120 125
 Lys Ser Ala Met Ser Thr Tyr Ser Gly Tyr Thr Leu Tyr Phe Thr Gly
 130 135 140
 His Ser Leu Gly Gly Ala Leu Ala Thr Leu Gly Ala Thr Val Leu Arg
 145 150 155 160
 Asn Asp Gly Tyr Ser Val Glu Leu Tyr Thr Tyr Gly Cys Pro Arg Val
 165 170 175
 Gly Asn Tyr Ala Leu Ala Glu His Ile Thr Ser Gln Gly Ser Gly Ala
 180 185 190
 Asn Phe Pro Val Thr His Leu Asn Asp Ile Val Pro Arg Val Pro Pro
 195 200 205
 Met Asp Phe Gly Phe Ser Gln Pro Ser Pro Glu Tyr Trp Ile Thr Ser
 210 215 220
 Gly Thr Gly Ala Ser Val Thr Ala Ser Asp Ile Glu Leu Ile Glu Gly

-continued

225	230	235	240
Ile Asn Ser Thr Ala Gly Asn Ala Gly Glu Ala Thr Val Asp Val Leu			
245	250	255	
Ala His Leu Trp Tyr Phe Phe Ala Ile Ser Glu Cys Leu Leu			
260	265	270	
<210> SEQ ID NO 15			
<211> LENGTH: 269			
<212> TYPE: PRT			
<213> ORGANISM: Aspergillus oryzae			
<400> SEQUENCE: 15			
Asp Val Ser Ser Leu Leu Asn Asn Leu Asp Leu Phe Ala Gln Tyr			
1	5	10	15
Ser Ala Ala Ala Tyr Cys Asp Glu Asn Leu Asn Ser Thr Gly Thr Lys			
20	25	30	
Leu Thr Cys Ser Val Gly Asn Cys Pro Leu Val Glu Ala Ala Ser Thr			
35	40	45	
Gln Ser Leu Asp Glu Phe Asn Glu Ser Ser Tyr Gly Asn Pro Ala			
50	55	60	
Gly Tyr Leu Ala Ala Asp Glu Thr Asn Lys Leu Leu Val Leu Ser Phe			
65	70	75	80
Arg Gly Ser Ala Asp Leu Ala Asn Trp Val Ala Asn Leu Asn Phe Gly			
85	90	95	
Leu Glu Asp Ala Ser Asp Leu Cys Ser Gly Cys Glu Val His Ser Gly			
100	105	110	
Phe Trp Lys Ala Trp Ser Glu Ile Ala Asp Thr Ile Thr Ser Lys Val			
115	120	125	
Glu Ser Ala Leu Ser Asp His Ser Asp Tyr Ser Leu Val Leu Thr Gly			
130	135	140	
His Ser Tyr Gly Ala Ala Leu Ala Ala Leu Ala Ala Thr Ala Leu Arg			
145	150	155	160
Asn Ser Gly His Ser Val Glu Leu Tyr Asn Tyr Gly Gln Pro Arg Leu			
165	170	175	
Gly Asn Glu Ala Leu Ala Thr Tyr Ile Thr Asp Gln Asn Lys Gly Gly			
180	185	190	
Asn Tyr Arg Val Thr His Thr Asn Asp Ile Val Pro Lys Leu Pro Pro			
195	200	205	
Thr Leu Leu Gly Tyr His His Phe Ser Pro Glu Tyr Tyr Ile Ser Ser			
210	215	220	
Ala Asp Glu Ala Thr Val Thr Thr Asp Val Thr Glu Val Thr Gly			
225	230	235	240
Ile Asp Ala Thr Gly Asn Asp Gly Thr Asp Gly Thr Ser Ile Asp			
245	250	255	
Ala His Arg Trp Tyr Phe Ile Tyr Ile Ser Glu Cys Ser			
260	265		
<210> SEQ ID NO 16			
<211> LENGTH: 251			
<212> TYPE: PRT			
<213> ORGANISM: Landerina penisapora			
<400> SEQUENCE: 16			
Pro Gln Asp Ala Tyr Thr Ala Ser His Ala Asp Leu Val Lys Tyr Ala			

-continued

1	5	10	15												
Thr	Tyr	Ala	Gly	Leu	Ala	Tyr	Gln	Thr	Thr	Asp	Ala	Trp	Pro	Ala	Ser
				20			25					30			
Arg	Thr	Val	Pro	Lys	Asp	Thr	Thr	Leu	Ile	Ser	Ser	Phe	Asp	His	Thr
					35		40				45				
Leu	Lys	Gly	Ser	Ser	Gly	Tyr	Ile	Ala	Phe	Asn	Glu	Pro	Cys	Lys	Glu
					50		55			60					
Ile	Ile	Val	Ala	Tyr	Arg	Gly	Thr	Asp	Ser	Leu	Ile	Asp	Trp	Leu	Thr
					65		70		75			80			
Asn	Leu	Asn	Phe	Asp	Lys	Thr	Ala	Trp	Pro	Ala	Asn	Ile	Ser	Asn	Ser
					85		90			95					
Leu	Val	His	Glu	Gly	Phe	Leu	Asn	Ala	Tyr	Leu	Val	Ser	Met	Gln	Gln
					100		105			110					
Val	Gln	Glu	Ala	Val	Asp	Ser	Leu	Leu	Ala	Lys	Cys	Pro	Asp	Ala	Thr
					115		120			125					
Ile	Ser	Phe	Thr	Gly	His	Ser	Leu	Gly	Gly	Ala	Leu	Ala	Cys	Ile	Ser
					130		135			140					
Met	Val	Asp	Thr	Ala	Gln	Arg	His	Arg	Gly	Ile	Lys	Met	Gln	Met	Phe
					145		150		155		160				
Thr	Tyr	Gly	Gln	Pro	Arg	Thr	Gly	Asn	Gln	Ala	Phe	Ala	Glu	Tyr	Val
					165		170			175					
Glu	Asn	Leu	Gly	His	Pro	Val	Phe	Arg	Val	Val	Tyr	Arg	His	Asp	Ile
					180		185			190					
Val	Pro	Arg	Met	Pro	Pro	Met	Asp	Leu	Gly	Phe	Gln	His	His	Gly	Gln
					195		200			205					
Glu	Val	Trp	Tyr	Glu	Gly	Asp	Glu	Asn	Ile	Lys	Phe	Cys	Lys	Gly	Glu
					210		215			220					
Gly	Glu	Asn	Leu	Thr	Cys	Glu	Leu	Gly	Val	Pro	Phe	Ser	Glu	Leu	Asn
					225		230			235			240		
Ala	Lys	Asp	His	Ser	Glu	Tyr	Pro	Gly	Met	His					
					245		250								

<210> SEQ ID NO 17
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 17

Ala Cys Met Ser His Thr Trp Gly Glu Arg Asn Leu
 1 5 10

<210> SEQ ID NO 18
 <211> LENGTH: 14
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 18

His Gly Trp Gly Glu Asp Ala Asn Leu Ala Met Asn Pro Ser
 1 5 10

1. A polypeptide having lipase activity and which further has an Average Relative Performance (RPavg) of at least 0.8 and a Benefit Risk (BR) of at least 1.1 at the test conditions given in the specification.
2. The polypeptide of claim 1, which further has a relative LU/A280 less than 1.00 at the test conditions given in the specification.
3. The polypeptide of claim 1 which is a bacterial polypeptide.
4. The polypeptide of claim 1 which is a fungal polypeptide.
5. The polypeptide of claim 4 which is a *Thermomyces* polypeptide.
6. The polypeptide of claim 5 which is a *Thermomyces lanuginosus* polypeptide.
7. The polypeptide of claim 1 which is a variant of a lipase comprised by the polypeptide of SEQ ID NO: 2.
8. The polypeptide of claim 1 which is a variant of a lipase comprised by the mature part of the polypeptide of SEQ ID NO: 2.
9. The polypeptide of claim 1 which is a variant of a lipase comprising the polypeptide of SEQ ID NO: 2.
10. The polypeptide of claim 1 which is a variant of a lipase comprising the mature part of the polypeptide of SEQ ID NO: 2.
11. The polypeptide of claim 1, which is encoded by a polynucleotide which hybridizes under at least high stringency conditions with nucleotides 644 to 732 of SEQ ID NO: 1 or a complementary strand hereto.
12. An isolated polynucleotide comprising a nucleotide sequence which encodes the polypeptide of any of the preceding claims.
13. A nucleic acid construct comprising the polynucleotide of claim 12 operably linked to one or more control sequences that direct the production of the polypeptide in an expression host.
14. A recombinant expression vector comprising the nucleic acid construct of claim 13.
15. A recombinant host cell comprising the nucleic acid construct of claim 14.
16. A method for producing the polypeptide of claim 1 comprising
 - (a) cultivating a cell, which in its wild-type form is capable of producing the polypeptide, under conditions conducive for production of the polypeptide; and
 - (b) recovering the polypeptide.
17. A method for producing the polypeptide of claim 1 comprising
 - (a) cultivating a host cell comprising a nucleic acid construct comprising a nucleotide sequence encoding the polypeptide under conditions conducive for production of the polypeptide; and
 - (b) recovering the polypeptide.

* * * * *