



US 20050251871A1

(19) **United States**(12) **Patent Application Publication** (10) **Pub. No.: US 2005/0251871 A1**
Chiaur et al. (43) **Pub. Date: Nov. 10, 2005**(54) **NOVEL UBIQUITIN LIGASES AS
THERAPEUTIC TARGETS**(52) **U.S. Cl.** **800/8**; 435/6; 435/69.1; 435/226;
435/320.1; 435/325; 536/23.2(75) Inventors: **Dah Shiam Chiaur**, New York, NY
(US); **Michele Pagano**, New York, NY
(US); **Esther Latres**, New York, NY
(US)(57) **ABSTRACT**

Correspondence Address:

JONES DAY**222 EAST 41ST ST****NEW YORK, NY 10017 (US)**(73) Assignee: **New York University**(21) Appl. No.: **10/632,150**(22) Filed: **Jul. 30, 2003****Related U.S. Application Data**(62) Division of application No. 09/385,219, filed on Aug.
27, 1999, now Pat. No. 6,720,181.(60) Provisional application No. 60/098,355, filed on Aug.
28, 1998. Provisional application No. 60/118,568,
filed on Feb. 3, 1999. Provisional application No.
60/124,449, filed on Mar. 15, 1999.**Publication Classification**(51) **Int. Cl.⁷** **C12Q 1/68**; A01K 67/00;
C07H 21/04; C12N 9/64

The present invention relates to the discovery, identification and characterization of nucleotides that encode novel substrate-targeting subunits of ubiquitin ligases. The invention encompasses nucleotides encoding novel substrate-targeting subunits of ubiquitin ligases: FBP1, FBP2, FBP3, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25, transgenic mice, knock-out mice, host cell expression systems and proteins encoded by the nucleotides of the present invention. The present invention relates to screening assays that use the novel substrate-targeting subunits to identify potential therapeutic agents such as small molecules, compounds or derivatives and analogues of the novel ubiquitin ligases which modulate activity of the novel ubiquitin ligases for the treatment of proliferative and differentiative disorders, such as cancer, major opportunistic infections, immune disorders, certain cardiovascular diseases, and inflammatory disorders. The invention further encompasses therapeutic protocols and pharmaceutical compositions designed to target ubiquitin ligases and their substrates for the treatment of proliferative disorders.

SKP2**	LP	--	DE	LL	GL	IF	SC	--	LC	PE	--	LL	KV	SG	VCK	RR	WY	R	LA	S	--	DE	S	L	W	Q																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																				
Fbp1**	LP	AR	LD	HA	EN	IL	SY	--	LD	AK	S	--	LC	AA	EL	VCK	KE	WY	R	VT	S	--	DC	M	L	W	K																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																			
Fbp2**	LP	--	LE	LS	F	Y	LL	KW	--	LD	PQ	T	--	LL	TC	CL	VS	SK	Q	W	N	K	V	I	S	--	AC	TE	V	W	Q																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																															
Fbp3a**	LL	--	QD	II	IL	QV	F	KY	--	LP	LD	--	RA	HA	SS	QV	CR	NR	W	N	Q	V	F	H	--	MP	DL	L	W	R																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																
Fbp3b**	LP	--	HH	VV	LL	Q	IF	QY	--	LP	LD	--	RA	CA	SS	V	CR	RR	W	N	E	V	F	H	--	J	SD	L	W	R																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																
Fbp4*	LP	--	ID	VQ	LY	IL	SE	--	LS	PH	D	--	LC	QL	GS	T	N	HY	W	N	E	T	V	R	--	NP	I	L	W	R																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																
Fbp5*	LR	--	HH	VA	T	IL	LA	Q	--	LS	DM	D	--	LI	NV	SK	V	ST	T	W	K	I	L	E	D	--	DK	G	A	F	Q																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																															
Fbp6*	LP	--	DN	IL	LE	F	TH	--	VP	AR	QL	--	LL	NC	R	V	CS	L	W	R	D	L	I	D	--	LL	T	L	W	K																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																
Fbp7**	LP	--	LE	KL	LR	IF	R	--	--	--	--	--	LD	V	R	S	V	LS	A	V	CR	D	L	F	T	A	S	N	--	DP	L	L	W	R																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												
Fbp8**	LP	--	PE	LS	F	T	IL	SY	--	LN	AT	D	--	LC	LA	S	--	--	--	--	--	--	--	--	--	--	DE	L	L	W	Q																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																															
Fbp9**	LP	--	GE	VL	E	Y	IL	CC	G	S	--	LT	AA	D	--	IG	R	V	S	T	CR	R	L	E	L	C	Q	S	--	SG	K	V	W	K																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												
Fbp10	LA	--	--	--	--	--	--	--	--	--	--	--	LR	V	A	C	V	CR	L	W	R	E	C	V	R	R	V	L	R	T	H	R	S	V	T	W	I																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
Fbp11	LP	--	DE	VV	L	K	IF	SY	--	LE	QD	--	LC	RA	A	C	V	CR	K	F	S	E	L	A	N	--	DP	N	L	W	K																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																															
Fbp12	LP	--	LE	L	W	R	M	IL	AY	--	CH	LD	--	LC	R	CS	L	V	CR	A	W	Y	E	L	I	L	S	--	LD	S	T	R	W	R																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												
Fbp13*	LP	--	TD	PL	LL	IL	SE	--	LD	YR	D	--	LI	NC	C	Y	V	S	R	L	S	Q	L	S	S	--	HD	P	L	W	R																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																															
Fbp14	W	A	W	GE	K	V	L	S	N	I	S	A	L	--	TD	L	GV	W	V	LC	G	S	W	R	R	H	V	G	--	AG	L	C	W	A																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												
Fbp15*	LP	--	EP	LL	LR	V	LA	A	--	LP	AA	EL	--	V	Q	AC	R	V	CL	R	W	K	E	L	V	D	--	GA	P	L	W	L																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																														
Fbp16*	LP	--	PE	L	VE	H	IL	SE	--	LP	VR	D	--	LV	AL	G	Q	T	CR	Y	F	H	E	V	C	D	--	GE	G	V	W	R																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																														
Fbp17**	LP	--	EV	LL	LL	H	M	CS	Y	--	LD	M	R	A	--	LG	R	L	A	Q	V	Y	R	W	L	H	F	T	I	N	C	--	DL	L	R	Q	I	A	W	A																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																						
Fbp18*	LP	--	LM	LN	N	IL	Y	R	--	SD	GW	D	--	IT	T	L	G	Q	V	I	P	T	L	Y	M	L	SE	--	DR	Q	L	W	K																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
Fbp19*	LP	--	DH	S	M	V	Q	IF	SE	--	LP	T	N	Q	--	LC	R	CA	R	V	CR	RR	W	Y	N	L	AW	--	DP	R	L	W	R																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
Fbp20	LP	--	LE	IL	V	Q	IF	GL	--	LV	AA	D	G	P	M	P	F	LC	R	AA	R	V	CR	RR	W	Q	E	A	S	--	QP	A	L	W	H																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
Fbp21*	LP	--	PE	V	M	L	S	IF	SY	--	LN	P	QE	--	LC	R	CS	Q	V	S	M	K	W	S	Q	L	L	K	--	T	G	S	L	W	K																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
Fbp22*	LP	--	KE	LL	L	R	IF	SE	--	LD	I	V	T	--	LC	R	CA	Q	I	S	K	A	W	N	I	L	L	A	L	--	DC	S	N	W	Q																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
Fbp23**	LP	--	YE	LI	Q	L	IL	NH	--	LT	L	P	D	--	LC	R	L	A	Q	T	CK	L	L	S	Q	H	CC	D	X	I	H	L	N	L	Q	P	Y	W	A																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
Fbp24**	LP	--	ME	V	L	M	Y	IF	R	W	--	V	V	SS	D	L	D	R	S	L	E	Q	L	S	L	V	CR	GF	Y	I	C	A	R	--	DP	E	L	W	R																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
Fbp25	LP	--	PE	I	Q	A	K	F	L	C	V	L	E	R	T	C	P	S	K	E	K	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

FIG. 1




FIG.2

10 20 30 40 50 60
MDPAEAVLQEKALKFMNSSEREDCNNGEPPRKIIPEKNSLRQTYNSCARLCLNQETVCLA

70 80 90 100 110 120
STAMKTENCVAKTKLANGTSSMIVPKQRKLSASYEKEKELCVKYFEQWSESDQVEFVEHL

130 140 150 160 170 180
ISQMCHYQHGHINSYLPMLQRDFITALPARGLDHIAENILSYLDAKSLCAAELVCKEWY

190 200 210 220 230 240
RVTSDGMLWKKLIERMVRTDSLWRGLAERRGWGQYLFKNKPPDGNAPPNSFYRALYPKII

250 260 270 280 290 300
QDIETIESNWRCGRHSLQRIHCRSETSKGVYCLQYDDQKIVSGLRDNTIKIWDKNTLECK

310 320 330 340 350 360
RILTGHTGSVLCLQYDERVIIITGSSDSTVRVWDVNTGEMLNTLIHHCEAVLHLRFNNGMM

370 380 390 400 410 420
VTCSKDRSIAVWDMASPTDITLRRVLVGHRAAVNVVDFDDKYIVSASGDRTIKVWNTSTC

430 440 450 460 470 480
EFVRTLNHGKRGIAQLQYRDRLVVGSSDNTIRLWDIECGACLRVLEGHEELVRCIRFDN

490 500 510 520 530 540
KRIVSGAYDGKIKVWDLVAALDPRAPAGTLCLRTLVEHSGRVFRLQFDEFQIVSSSHDDT

550 560
ILIWDFLNDPAAQAEPSPRSPRTYTYISR

FIG.3A

10 20 30 40 50 60 70 80 90
TGCCTGGCTGGGCTGGACCAAGGGGGCCCCGGCGGAGAGCGGACCGAGTGGCCTCGGCGATTATGGACCGCGCGAGCGGTGCTGC
100 110 120 130 140 150 160 170 180
AAGAGAAGGCACTCAAGTTTATGAATTCCTCAGAGAGAGAAGACTGTAATAATGGCGAACCCTTAGGAAGATAATACCAGAGAAGAATTCACT
190 200 210 220 230 240 250 260 270 280
TAGACAGACATACAACAGCTGTGCCAGACTCTGCTTAACCAAGAAACAGTATGTTAGCAAGCACTGCTATGAAGACTGAGAATTGTTGGCC
290 300 310 320 330 340 350 360 370
AAAACAAAACCTTGCCAAATGGCACTTCCAGTATGATTTGCCCCAAGCAACGGAACCTCTCAGCAAGCTATGAAAAGGAAAAGGAACCTGTGTCTCA
380 390 400 410 420 430 440 450 460 470
AATACITTTGAGCAGTGGTCAGAGTCAGATCAAGTGAATTTGTGGAACATCTTATATCCCAAATGTGTCATTACCAACATGGGCACATAAACTC
480 490 500 510 520 530 540 550 560
GTATCTTAAACCTATGTTGCAGAGAGATTTTCATAACCTGCTCTGCCAGCTCGGGGATTTGGAATCAATATCCCTGAGAACATCTGTCTATACCTGGAT
570 580 590 600 610 620 630 640 650
GCCAAATCACTATGTCGCTGAACTTGTTGCAAGGAATGGTACCGAGTGACCTCTGATGGCATGCTGTGGAAGAAGCTTATCGAGAGAAATGG
660 670 680 690 700 710 720 730 740 750
TCAGGACAGATTCTCTGTGGAGAGGCCCTGGCAGAACGAAGAGGATGGGGACAGTATTTATTCAAAAACAACCTCTCTGACGGGAATGCTCTCTCC
760 770 780 790 800 810 820 830 840
CAACTCTTTTATAGAGCACTTTATCCTAAAAATTATACAAGACATTGAGACAAATAGAATCTAATTGGAGATGTGGAAGACATAGTTTACAGAGA
850 860 870 880 890 900 910 920 930 940
ATTCACTGCCGAAGTGAACAAGCAAGGAGTTTACTGTTTACAGTATGATGATCAGAAAAATAGTAAGCGGCCCTTCGAGACAACACAATCAAGA

FIG.3B

950 960 970 980 990 1000 1010 1020 1030
TC TGGGATAAAAACACATTGGGAATGCAAGCGAATTCACAGGCCATACAGGTT CAGTCCCTCIGTCICCCAGTATGATGAGAGAGTGATCATAAC
1040 1050 1060 1070 1080 1090 1100 1110 1120
AGGATCATCGGATCCACGGTCAGAGTGTGGGATGTAAATACAGGTGAAATGCTAAACACGTTGATT CACCAATTGTGAAGCAGTTCTGCACCTTG
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
CGTTTCAATAATGGCATGATGGTGACCTGCTCCAAGATCGTTCCATTGCTGTATGGGATATGGCCCTCCCAACTGACATTACCCCTCCGGAGGG
1230 1240 1250 1260 1270 1280 1290 1300 1310
TGCTGGTCGGACACCGAGCGTCGTCAATGTGTGACACTTGTGACACAAGTACATTGTTCTGCACTCGGGATAGAACTATAAAGGTATGGAA
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
CACAAAGTACTTGTGAATTTGTAAGGACCTTAAATGGACACAAACGAGGCCATTGCCGTGTTTCAGTACAGGGACAGGCTGGTAGTGAGTGGCTCA
1420 1430 1440 1450 1460 1470 1480 1490 1500
TC TGACAACACTATCAGATTATGGGACATAGAAATGTGGTGCATGTTTACGAGTGTAGAGGCCCATGAGGAATTGGTCCGTTGTATTCGATTTG
1510 1520 1530 1540 1550 1560 1570 1580 1590
ATAACAAGAGGATAGTCAGTGGGGCCATGATGGGAAAAATTAAAGTGTGGGATCTTGTGGCTGCTTGGACCCCCCGTCTCTCCAGGGGACACT
1600 1610 1620 1630 1640 1650 1660 1670 1680 1690
CTGTCTACGGACCCCTTGTGGAGCATTCGGGAAGAGTTTTTCGACTACAGTTTGATGAATTCAGATTGTCAGTAGTTCACATGATGACACAATC

FIG. 3C

1700 1710 1720 1730 1740 1750 1760 1770 1780
CTCATCTGGGACTTCCTAAATGATCCAGCTGCCCAAGCTGAACCCCCCGTTCCCTTCTCGAACAATACACCTACATCTCCAGATAAATAACCA
1790 1800 1810 1820 1830 1840 1850 1860 1870 1880
TACACTGACCTCATACTTGCCCGAGGACCCATTAAAGTTGGGTATTTAACGTATCTGCCAATACCAGGATGAGCAACAACAGTAACAATCAAAAC
1890 1900 1910 1920 1930 1940 1950 1960 1970
TACTGCCCAGTTTCCCTGGACTAGCCGAGGAGCAGGGCTTTGAGACTCCTGTGGGACACAGTTGGTCTGCAGTCGGCCCCCAGGACGGTCTIACTC
1980 1990 2000 2010 2020 2030 2040 2050 2060
AGCACAACTGACTGCTTCAGTGTCTATCAGAAGATGTCCTCTATCAATTGIGAATGATTGGAACTTTAAACCTCCCCCTCCTCTCCTCCTTT
2070 2080 2090 2100 2110 2120 2130 2140 2150
CACCTCTGCACCTAGTTTTTCCCATTTGTTCCAGACAAGGTGACTTATAAATATATTTAGTGTTTTGCCAGAAAAA

FIG.3D

10 20 30 40 50 60
MERKDFETWLDNISVTFLSLTDLQKNETLDHLISLSGAVQLRHLSNNLETLLKRDFLKLL
70 80 90 100 110 120
PLELSFYLLKWLDPQTLLTCCLVSKQWNKVISACTEVWQTACKNLGWQIDDSVQDALHWK
130 140 150 160 170 180
KVYLKAILRMKQLEDHEAFETSSLIGH SARVYALYYKDGLLCTGSDDL SAKLWDVSTGQC
190 200 210 220 230 240
VYGIQHTTCAAVKFDEQKLVTGSFDNTVACWEWSSGARTQHFRGHTGAVFSVDYNDELDI
250 260 270 280 290 300
LVSGSADFTVKVWALSAGTCLNTLTGHTEWVTKVVLQKCKVKSLLHSPGDYILLSADKYE
310 320 330 340 350 360
IKIWPIGREINCKCLKTSLVSEDRSICLQPRLHFDGKYIVCSSALGLYQWDFASYDILRV
370 380 390 400 410 420
IKTPEIANLALLGFGDIFALLFDNRYLYIMDLRTESLISRWPLPEYRESKRGSSFLAGEH

PG

FIG.4A

10 20 30 40 50 60 70 80 90
ATCGAGAGAAAGGACITTTGAGACATGGCTTGATAACATTTCGTACATTTCCTCTGACGGACTTGCAGAAAAATGAAACTCTGGATCACC

100 110 120 130 140 150 160 170 180
TGATTAGTCTGAGTGGGCGAGTCCAGCTCAGGCATCTCTCCAAATAACCTAGAGACTCTCTCAAGCGGGACTTCCCTCAAACCTCCTTCCCCCTGGA

190 200 210 220 230 240 250 260 270 280
GCTCAGTTTTTATTGTTAAATGGCTCGATCCTCAGACTTTACTCAGATGCTGCCCTCTCTAAACAGTGGAAATAAGGTGATAAGTGCCTGT

290 300 310 320 330 340 350 360 370
ACAGAGGTGTGGCAGACTGCATGTAAAAATTTGGGCTGGCAGATAGATGATTCTGTTCAAGGACGCTTGGCACTGGAAGAAGGTTTATTTGAAGG

380 390 400 410 420 430 440 450 460 470
CTATTTTGAGAAATGAAGCAACTGGAGGACCATGAAGCCTTTGAACCTCGTCATTAAATGGCACACAGTCCCAGAGIGTATGCCACTTTACTACAA

480 490 500 510 520 530 540 550 560
AGATGGACTTCTCTGTACAGGGTCAGATGACTTGCTGCAAGCTGTGGATGTGAGCACAGGGCAGTGGTTTATGGCATCCAGACCCACACT

570 580 590 600 610 620 630 640 650
TGTCAGCGGTGAAGTTGATGAACAGAGCTTGTGACAGGCTCCTTTGACACACACTGTGGCTTGTGGGAAATGGAGTTCCGGAGCCAGGACCC

660 670 680 690 700 710 720 730 740 750
AGCACTTTCGGGGGCACACGGGGCGGTATTTAGCGTGGACTACAATGATGAACCTGGATATCTTGGTGAGCGGCTCTGCAGACTTCACCTGTGAA

760 770 780 790 800 810 820 830 840
AGTATGGGCTTTATCTGCTGGGACATGCCCTGAACACACTCACCGGGCACACGGAAATGGGTACCAAGGTAGTTTTGCAGAAATGCCAAAGTCAAG

850 860 870 880 890 900 910 920 930 940
TCTCTCTTGCACAGTCCCTGGAGACTACATCCTCTTAAGTGCAGACAAAATATGAGATTGAAGATTGGCCAAATGGGAGAGAAATCAACTGTAACT

FIG. 4B

950 960 970 980 990 1000 1010 1020 1030
GCTTAAAGACATTGTCCTCTGAGGATAGAAGTATCTGCCCTGCAGCCCAAGACTTCATTTTGATGGCAAAATACATTGCTCTGTAGTTCAGCACT

1040 1050 1060 1070 1080 1090 1100 1110 1120
TGGTCTCTACCAGTGGGACTTGGCCAGTTAIGATAATCTCAGGGTCATCAAGACTCCTGAGATAGCAAACTTGGCCCTTGGCTTTGGAGAT

1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
ATCTTTGCCCCTGGCTTTTGACAACCGCTACCCTGTACATCATGGACTTGGCGACAGAGAGCCCTGATTAGTCCCTGGCCCTCTGCCAGAGTACAGGG

1230 1240 1250 1260 1270 1280 1290 1300 1310
AATCAAAGAGAGGGCTCAAGCTTCCTGGCAGGCGAACAATCCTGGCTGAATGGACTGGATGGCCACAATGACACGGGCTTGGTCTTTGCCACCAGC

1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
ATGCCGTGACCACAGTATTACCTGGTGTGTGGAAGGAGCACGGCTGACACCATGAGCCACCACCGCTGACTGACTTTGGGTGCCGGGGGCTGCC

1420 1430 1440 1450 1460 1470
GGTTTGGGTGCACCTCTGCGGCACGGGACTGCGATGAACCAAAAGTTCTCACCTAATGGTATCATCA

FIG.4C

10 20 30 40 50 60
MKRGGROSDRNSSEEGTAEKSKKLRTTNEHSQTCDWGNLLQDIILQVFKYLPLLDRAHAS

70 80 90 100 110 120
QVCRNWNQVFHMPDLWRCFEFELNQPATSYLKATHPELIKQIIKRHSNHLQYVSFKVDSS

130 140 150 160 170 180
KESAEAACDILSQLVNCSLKTGLISTARPSFMDLPKSHFISALTVVFVNSKSLSSLKID

190 200 210 220 230 240
DTPVDDPSLKVLVANNSDTLKLKMS SCPHVSPAGILCVADQCHGLRELALNYHLLSDEL

250 260 270 280 290 300
LLALSSEKHVRLEHLRIDVVS ENPGQTHFHTIQSSWDAFIRHSPKVNLMYFFLYEEEF

310 320 330 340 350 360
DPFFRYEIPATHLYFGRSVSKDVLGRVGMTCPRLVELVVCANGLRPLDEELIRIAERCKN

370 380 390 400 410 420
LSAIGLGECEVSCSAFVEFVKMCGGRLSQLSIMEEVLIPDQKYSLEQIHWEVSKHLGRVW

FPDMMPTW

FIG.5A

10 20 30 40 50 60 70 80 90
CGGGTGGTGTGTGGGGAAGCCGCCCGGAGCAGGATGAACGAGGAGGAGAGATAGTACCGTAATTTCATCAGAAGAAGAACTGCAGA
100 110 120 130 140 150 160 170 180
GAAATCCAAGAACTGAGGACTACAAATGAGCATTCACAGCTTGCTGATGGGGTAATCTCCTTCAGGACATTATCTCCAAGTATTAAATAT
190 200 210 220 230 240 250 260 270 280
TTGCCCTCTTTCACCGGGCTCATGCTTCACAAGTTTGGCCGAAC TGGAACCGGATTTTACAATGCCCTGACTTGTGGAGAIGTTTGAATTG
290 300 310 320 330 340 350 360 370
AACTGAATCAGCCAGCTACATCTTATTGAAGCTACCCATCCAGAGCTGATCAAAACAGATTATTAAAGACATTCAAAACCATCTACAATATGT
380 390 400 410 420 430 440 450 460 470
CAGCTTCAAGTGGACAGCAGCAAGGAATCAGCTGAAGCAGCTTGTGATATACTATCCAACTTGTGAATTGCTCTTTAAAAACACTTGGACTT
480 490 500 510 520 530 540 550 560
ATTTCAACTGCTCGACCAAGCTTTATGGATTTACCAAGTCTCAGCTTTATCTCTGCCACTGACAGTGTGTTGGTAAACTCCAAATCCCTGCTCTT
570 580 590 600 610 620 630 640 650
CCCTTAAGATAGATGATCTCCAGTAGATGATCCATCTCAAGTACTAGTGGCCCAACAATAGTGATACACATCAAGCTGTGAAAAATGACGAG
660 670 680 690 700 710 720 730 740 750
CTGTCCCTCATGTCTCTCCAGCAGGTATCCTTTGTGTGGCTGATCAGTGTACCGGCTTAAGAGAAGTAGCCCTGAAC TACCAC TTTATTGAGTGAT
760 770 780 790 800 810 820 830 840
GAGTTGTTACTTGCCATTGCTCTCGAAAAACATGTTCCATTAGAACATTTGGCATTGATGTAGTCAGTGAGAAATCCTTGGACACACACACTTCC
850 860 870 880 890 900 910 920 930 940
ATACTATTCAAGAGTAGCTGGGATGCTTTCATCAGACATTCACCCCAAGTGAAC TTAGTGATTTTTTTTTTATGAAGAAGAAATTGA

FIG.5B

950 960 970 980 990 1000 1010 1020 1030
CCCCCTTCCTTGGCTAIGAAATACCTGCCACCCCACTGCTACTTTGGGAGATCAGTAAGCAAAGATGTGCTTGGCCGTGTGGGAATGACATGCCCCT

1040 1050 1060 1070 1080 1090 1100 1110 1120
AGACTGGTTGAAC TAGTGTGTCCAAATGGATTACCGCCCACTTGAAGAGTTAATTCGCATTCAGAACGTTGCAAAAATTGTCAGCTA

1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
TTGGACTAGCGGAATGTGAAGTCTCATGTAGTGCCTTTGTTGAGTTTGTGAAGATGTGTGGTGGCCGCCCTAICICAATTATCCATTATGGAAGA

1230 1240 1250 1260 1270 1280 1290 1300 1310
AGTACTAATTCCTGACCACCAAGTATAGTTTGGAGCAGATTCACTGGGAAGTGTCCAAGCATCTTGGTAGGGTGTGGTTTCCCGACATGATGCCC

1320 1330 1340 1350 1360 1370 1380 1390 1400
ACTTGGTAAAACTGCATGATGAATAGCACCTTAATTCAAGCAAATGTATTATAATTAAAGTTTATTTCCTGTAAAAAATAAAAAA

FIG.5C

10	20	30	40	50	60
MKRNSLSVENKIVQLSGAAKQPKVGFYSSLNQTHHTVLLDWGSLPHHVVLQIFQYLP LL					
70	80	90	100	110	120
DRACASSVCRRWNEVFHISDLWRKF EFELNQSATSSFKSTHPDL IQQIIKKHFAHLQYVS					
130	140	150	160	170	180
FKVDSSAESAEAACDILSQLVNCSIQTLGLISTAKPSFMNVSESHFVSALT VVF INSKSL					
190	200	210	220	230	240
SSIKIEDTPVDDPSLKILVANNSDTLR LPKMSSCPHVSSDGILCVADRCQGLRELALNYY					
250	260	270	280	290	300
ILTDELF LALSSETHVNLEHLRIDV VSENPQQIKFHAVKKHSDALIKHSPRVNVVMHFF					
310	320	330	340	350	360
LYEEEFETFFKEETPVTHLYFGRSVSKVVLGRVGLNCPRLIELVVCANDLQPLDNELICI					
370	380	390	400	410	420
AEHCTNLTALGLSKCEVSCSAFIRFVRLCERRLTQLSMEEVLIPDEDYSLDEI HTEVSK					
430					
YLGRVWF PDVMPLW					

FIG.6A

10	20	30	40	50	60	
ACATTTTCTAATGTTTACAGAATGAAGAGGAACAGTTTATCTGTTGAGAATAAAATTGTCCAGTTGTCA						
70	80	90	100	110	120	130
GGAGCAGCGAAACAGCCAAAAGTTGGGTTCTACTCTTCTCTCAACCAGACTCATACACACACGGTTCTT						
140	150	160	170	180	190	200
CTAGACTGGGGGAGTTTGCCTCACCATGTAGTATTACAAATTTTTCAGTATCTTCCTTTACTAGATCGG						
210	220	230	240	250	260	270
GCCTGTGCATCTTCTGTATGTAGGAGGTGGAATGAAGTTTTTCATATTTCTGACCTTTGGAGAAAGTTT						
280	290	300	310	320	330	340
GAATTTGAACTGAACCAGTCAGTACTTCATCTTTAAGTCCACTCATCCTGATCTCATTGAGCAGATC						
350	360	370	380	390	400	410
ATTA AAAAGCATTTTGCTCATCTTCAGTATGTCAGCTTTAAGGTTGACAGTAGCGCTGAGTCAGCAGAA						
420	430	440	450	460	470	480
GCTGCCTGTGATATACTCTCTCAGCTGGTAAATTGTTCCATCCAGACCTTGGGCTTGATTTCAACAGCC						
490	500	510	520	530	540	550
AAGCCAAGTTTCATGAATGTGTCGGAGTCTCATTTTGTGTCAGCACTTACAGTTGTTTTATCAACTCA						
560	570	580	590	600	610	620
AAATCATTATCATCAATCAAAATTGAAGATACACCAGTGGATGATCCTTCATTGAAGATTCTTGTGGCC						
630	640	650	660	670	680	690
AATAATAGTGACACTCTAAGACTCCCAAAGATGAGTAGCTGTCCTCATGTTTCATCTGATGGAATTCTT						
700	710	720	730	740	750	
TGTGTAGCTGACCGTTGTCAAGGCCTTAGAGAACTGGCGTTGAATTATTACATCCTAACTGATGAAC TT						
760	770	780	790	800	810	820
TTCCTTGCACTCTCAAGCGAGACTCATGTTAACCTTGAACATCTTGAATTGATGTTGTGAGTGAAAAT						
830	840	850	860	870	880	890
CCTGGACAGATTAAATTTTCATGCTGTAAAAAACACAGTTGGGATGCACTTATTAACATTCCCCTAGA						
900	910	920	930	940	950	960
GTTAATGTTGTTATGCACTTCTTCTATATGAAGAGGAATTCGAGACGTTCTTCAAAGAAGAAACCCCT						

FIG.6B

970 980 990 1000 1010 1020 1030
GTTACTCACCTTTATTTGGTCGTTCACTCAGCAAAGTGGTTTTAGGACGGTAGGTCTCAACTGTCCT

1040 1050 1060 1070 1080 1090 1100
CGACTGATTGAGTTAGTGGTGTGTGCTAATGATCTTCAGCCTCTTGATAATGAACCTATTTGTATTGCT

1110 1120 1130 1140 1150 1160 1170
GAACACTGTACAAACCTAACAGCCTTGGGCCTCAGCAAATGTGAAGTTAGCTGCAGTGCCTTCATCAGG

1180 1190 1200 1210 1220 1230 1240
TTTGTAAGACTGTGTGAGAGAAGGTTAACACAGCTCTCTGTAATGGAGGAAGTTTGATCCCTGATGAG

1250 1260 1270 1280 1290 1300 1310
GATTATAGCCTAGATGAAATTCACACTGAAGTCTCCAAATACCTGGGAAGAGTATGGTTCCTGATGTG

1230
ATGCCTCTCTGG

FIG.6C

10	20	30	40	50	60
MAGSEPRSGTNSPPPPFSDWGRLEAAILSGWKTFWQSVSKDRVARTTSREEVDEAASTLT					
70	80	90	100	110	120
RLPIDVQLYILSFLSPHDL CQLGSTNHYWNETVRNPILWRYFLLRDLPSWSSVDWKS LPY					
130	140	150	160	170	180
LQILKKPISEVSDGAFFDYMAVYLMCCPYTRASKSSRPMYGAVTSFLHSLIIPNEPRFA					
190	200	210	220	230	240
LFGPRLEQLNTSLVLSLLSSEELCPTAGLPQRQIDGIGSGVNFQLNNQHKFNILILYSTT					
250	260	270	280	290	300
RKERDRAREEHTSAVNKMF SRHNEGDDRPGSRYSVIPQIQKLCEVVDGF IYVANA EAHKR					
310	320	330	340	350	360
HEWQDEF SHIMAMTDPAFGSSGRPLLVLSCISQGDVKRMPCFYLAHELHLNLLNHPWL VQ					
370	380	390	400	410	420
DTEAETLTGFLNGIEWILEEVESKRAR*FSFQILGTETI*NLLRS*CEYLLSQPTLSCL					
430	440	450	460	470	480
FADRLSFGQL*LLCFLYYFYFLP*IN YKKRVSVLVFSPKMNL*TFFW*FLYFLSF*KY*I					

L

FIG.7A

10	20	30	40	50	60	
ATGGCGGGAAGCGAGCCGCGCAGCGGAACAAATTCGCCGCCGCCGCCCTTCAGCGACTGGGGCCGCCTG						
70	80	90	100	110	120	130
GAGGCGGCCATCCTCAGCGGCTGGAAGACCTTCTGGCAGTCAGTGAGCAAGGATAGGGTGGCGCGTACG						
140	150	160	170	180	190	200
ACCTCCCGGGAGGAGGTGGATGAGGCGGCCAGCACCTGACGCGGCTGCCGATTGATGTACAGCTATAT						
210	220	230	240	250	260	270
ATTTTGTCTTTCTTTACCTCATGATCTGTGTCAGTTGGGAAGTACAAATCATTATTGGAATGAACT						
280	290	300	310	320	330	340
GTAAGAAATCCAATTCTGTGGAGATACTTTTGTGTAGGGATCTTCCTTCTGGTCTTCTGTGACTGG						
350	360	370	380	390	400	410
AAGTCTCTTCCATATCTACAAATCTTAAAAAGCCTATATCTGAGGTCTCTGATGGTGCATTTTTTGAC						
420	430	440	450	460	470	480
TACATGGCAGTCTATCTAATGTGCTGTCCATACACAAGAAGAGCTTCAAATCCAGCCGTCCTATGTAT						
490	500	510	520	530	540	550
GGAGCTGTCACTTCTTTTTTACACTCCCTGATCATTCCCAATGAACCTCGATTTGCTCTGTTTGACCA						
560	570	580	590	600	610	620
CGTTTGGAACAATTGAATACCTCTTTGGTGTGAGCTTGCTGCTTCAGAGGAACCTTGCCCAACAGCT						
630	640	650	660	670	680	690
GGTTTGCCTCAGAGGCAGATTGATGGTATTGGATCAGGAGTCAATTTTCAGTTGAACAACCAACATAAA						
700	710	720	730	740	750	
TTCAACATTCTAATCTTATATTCAACTACCAGAAAGGAAAGAGATAGAGCAAGGGAAGAGCATACAAGT						
760	770	780	790	800	810	820
GCAGTTAACAAGATGTTTCAGTCGACACAATGAAGGTGATGATCGACCAGGAAGCCGTTACAGTGTGATT						
830	840	850	860	870	880	890
CCACAGATTCAAAAAGTGTGTGAAGTTGTAGATGGGTTTCATCTATGTTGCAAATGCTGAAGCTCATAAA						
900	910	920	930	940	950	960
AGACATGAATGGCAAGATGAATTTTCTCATATTATGGCAATGACAGATCCAGCCTTTGGGTCTTCGGGA						

FIG.7B

970 980 990 1000 1010 1020 1030
AGACCATTGTTGGTTTTATCTTGATTTCTCAAGGGGATGTAAAAAGAATGCCCTGTTTTATTGGCT

1040 1050 1060 1070 1080 1090 1100
CATGAGCTGCATCTGAATCTTCTAAATCACCCATGGCTGGTCCAGGATACAGAGGCTGAAACTCTGACT

1110 1120 1130 1140 1150 1160 1170
GGTTTTTTGAATGGCATTGAGTGGATTCTTGAAGAAGTGAATCTAAGCGTGCAAGATGATTCTCTTTT

1180 1190 1200 1210 1220 1230 1240
CAGATCTTGGGAACTGAAACCATTTGAAATTTATTACTAAGGTCGTGATGTGAATATTTGCTCAGTCAG

1250 1260 1270 1280 1290 1300 1310
CCCACCTTGTCCTGCCTTTTTGCAGATAGGCTTTCATTTGGACAGCTATAACTGCTGTGTTTTTTATAT

1320 1330 1340 1350 1360 1370 1380
TATTTTACTTTTTACCATAAATCAATTACAAGAAAAGAGTTTCAGTCCTAGTATTTAGCCCCAAAATG

1390 1400 1410 1420 1430 1440
AACCTTTAAACATTTTTTTGGTAATTTTTATATTTCTGTCTTTTTAAAAATATTAAATTTTGG

FIG.7C

10 20 30 40 50 60
MSRRPCSCALRPPRCSCSASPASVTAAGRPRPSDSCKEESSTLSVKMKCDFNCNHVHSGL

70 80 90 100 110 120
KLVKPDDIGRLVSYTPAYLEGSCKDCIKDYERLSCIGSPIVSPRIVQLETESKRLHNKEN

130 140 150 160 170 180
QHVQQTLLNSTNEIEALETSRLYEDSGYSSFSLQSGLSEHEEGSLLEENFGDSLQSCLLQI

190 200 210 220 230 240
QSPDQYPNKNLLPVLHFEKVVCSTLKKNAKRNPKVDREMLKEIIARGNFRQLNIIGRKMG

250 260 270 280 290 300
LECVDILSELFRRGLRHVLATILAQLSDMDLINVSKVSTTWKKILEDDKGAFQLYSKAIQ

310 320 330 340 350 360
RVTENNNKFSPHASTREYVMFRTPLASVQKSAAQTSKKDAQTKLSNQGQKQGSTYSRHN

370 380 390 400 410 420
EFSEVAKTLKKNESLKACIRCNSPAKYDCYLQRATCKREGCGFDYCTKCLCNYHTTKDCS

430 440
DGKLLKASCKIGPLPGTKKSKKNLRRRL

FIG.8A

10 20 30 40 50 60 70 80 90
AGGTTGCTCAGCTGCCCGGAGCGGTTCCCTCCACCTGAGGCAGACACCACCTCGGTTGGCATGAGCGCGCGCCCTGCAGCTCGGCCCTACGG

100 110 120 130 140 150 160 170 180
CCACCCCGCTGCTCCTGCAGCGCCAGCCCGGCGGAGTGACAGCGCCGCGCGCCCTCGACCTCGGATAGTTGTAAAGAAGAAAGTTCTACCC

190 200 210 220 230 240 250 260 270 280
TTTCTGTCAAAATGAAGTGTGATTTTAATTGTAACCATGTTTCATTCGGACTTAAACTGTTAAACCTGATGACATTGGAAGACTAGTTTCCTA

290 300 310 320 330 340 350 360 370
CACCCCTGCATATCTGGAAGGTTCCCTGTAAGACTGCATTAAAGACTATGAAGGCTGTCATGTTATGGGTCACCGATTGTAGCCCTAGGATT

380 390 400 410 420 430 440 450 460 470
GTACAAC TTGAAC TGAAGCAAGCGCTTGCATACAAGCAAAATCAACATGTGCAACAGACACTTAATAGTACAAATGAATAGAACACTAG

480 490 500 510 520 530 540 550 560
AGACCAGTAGACTTTATGAAGACAGTGGCTATTCCTCATTTTCTCTACAAGTGGCTCAGTGAACATGAAGAAGTAGCCTCCTGGAGGAGAA

570 580 590 600 610 620 630 640 650
TTTCGGTGACAGICTACAATCCCTGCTACAAATACAAGCCCGACACCAATATCCCAACAACAACTTGGTCCAGTCTTTCATTTTGAAAAA

660 670 680 690 700 710 720 730 740 750
GTGGTTGTTCAACATTAAAAAAGAAATGCCAAACGAAATCCCTAAAGTAGATCGGGAGATGCCGAAGCAATTAAGCCAGAGCAAAATTTIAGAC

760 770 780 790 800 810 820 830 840
TGCAGATATAATTGCCAGAAAAATGGGCTAGAAATGTTAGATATTCAGCGCAACTCTTTGGAAGGGGACTCAGACATGCTTTAGCAACTAT

850 860 870 880 890 900 910 920 930 940
TTTAGCACAAC TCACTGCACTGCACTTAATCAATGTGCTCTAAGTGAGCACAAC TTGCAAGAAGATCCTAGAAGATGATAAGGGGGCATTCGAC

FIG.8B

950 960 970 980 990 1000 1010 1020 1030
 TTGTACAGTAAAGCAATACAAGAGTTACCGAAAACAATAAATTTTCACCTCATGCTTCAACCAGAGAATATGTTATGTTCAGAACCCAC
 1040 1050 1060 1070 1080 1090 1100 1110 1120
 TGGCTTCTGTTTCAAGAAATCAGCAGCCAGACTTCTCTCAAAAAAGATGCTCAAACCAAGTTATCCAATCAAGGTGATCAGAAAGGTTCTACTTA
 1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
 TAGTCGACACAATGAATTTCTGAGGTTGCCAAGACATTGAAAAAGAACGAAAGCCTCAAAGCCTGTATTGCTGTAATTCACCTGCAAAATAT
 1230 1240 1250 1260 1270 1280 1290 1300 1310
 GATTGCTATTTACAACGGGCAACCTGCAACGAGAAGGCTGTGGATTGATTATGTACGAAGTGCTCTGTAATTATCATACTACTAAAGACT
 1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
 GTTCAGATGCCAAGCTCCTCAAAGCCAGTTGTAATAAGTCCCTGCCTGGTACAAAGAAAAGCAAAAAGAATTTACGAAGATTGTGATCTCT
 1420 1430 1440 1450 1460 1470 1480 1490 1500
 TATTAATCAATTGTTACTGATCATGAATGTTAGTTAGAAAAATGTTAGGTTTTAACTTAAAAAAATTTGATTGTGATTTTCAATTTTATGTTG
 1510 1520 1530 1540 1550 1560 1570 1580 1590
 AAATCGGTGTAGTATCCTGAGGTTTTTTTCCCCCAGAAGATAAGAGGATAGACAACCTCTTAAATATTTTTACAATTTAATGAGAAAAAGT
 1600 1610 1620 1630 1640 1650 1660 1670 1680 1690
 TTAATTTCTCAATACAATCAAACAATTTAAATATTTTAAGAAAAAGGAAAAGTAGATAGTACTGAGGGTAAAAAAAATTGATTCAA
 1700 1710 1720 1730 1740 1750 1760 1770 1780
 TTTTATGGTAAAGGAAACCATGCAATTTTACCTAGACAGTCTTAAATATGCTGGTTTTCCATCTGTTAGCATTTCAGACATTTTATGTTCTC
 1790 1800 1810 1820 1830 1840 1850 1860 1870 1880
 CTTACTCAATTGATACCAACAGAAATATCACTTCTGGAGTCTATTAAATGTGTGTACCTTTCTAAAGCTTTTTTTCATTGTGTGTATTTC
 1890 1900 1910 1920 1930 1940 1950 1960 1970
 CAAGAAAGTATCCTTTGTAAAACTTGCTTGTTCCTTATTTCTGAAATCTGTTTTAATATTTTGTATACATGTAATATTTCTGTATTTT
 1980 1990 2000 2010 2020 2030 2040 2050 2060
 TATATGTCAAAGAATATGCTCTTGTATGTACATATAAAAAATAAATTTTGCTCAATAAAATGTAAGCTTAAAAAAAAAAAAAAAAAACTCGAG
 2070
 ACTAGTGC

FIG.8C

10 20 30 40 50 60
ARSGASALRRRRVQVWVLSRPPPGGDSFRTRRPQRGPGPGGSQAMDAPHSKAALDSINE

70 80 90 100 110 120
LPDNILLELFTHVPARQLLLNCRLVCSLWRDLIDLLTLWKRKCLRKGFI TKDWDQPVADW

130 140 150 160 170 180
KIFYFLRSLHRNLLRNPCAENDMFAWQIDFNGGDRWKVDSLPGAHGTEFPDPKVKKSFT

190 200 210 220 230 240
SYELCLKWELVDLLADRYWEELLDTFRPDIVVKDWF AARADCGCTYQLKVQLASADYFVL

250 260 270 280 290 300
ASFEPPTVTIQQWNNATWTEVSYTFSYPRGVRYILFQHGGRTQYWAGWYGPRTNSSI

310 320 330
VVSPKMTRNQASSEAQPGQKHGQEEAAQSPYGAVVQIF

FIG.9A

10 20 30 40 50 60 70 80 90
GCGGTTGGGAGCTTCGGCCCTGCGTAGGAGGGGGTGCAGGTGTGGGTGCTGAGCGCGCGCGCGCTGGAGGGGAGACAGCTTCAGGACAC

100 110 120 130 140 150 160 170 180
GCAGCGCGCAGGAGGGCCCGGGGGATCCCAAGGCATGGACGCTCCCACTCCAAAGCAGCCCTGGACAGCATTAACGAGCTGCCCGCA

190 200 210 220 230 240 250 260 270 280
TAACATCCGTGCTGGAGCTGTTACGCACGTGCCCCGGCCGCGAGCTGCTGTAAGTGGCGGCTGGTCTGGAGCCCTCTGGCGGGACCTCATCGAC

290 300 310 320 330 340 350 360 370
CTCCTGACCCCTCGGAAACGCAAGTCCCTGCGAAAGGGCTTCATCACCAGGACTGGGACCAAGCCCGTGGCGGACTGGAAAATCTTCTACTTCC

380 390 400 410 420 430 440 450 460 470
TACGGAGCCTGCATAGGAACCTCCTGCGCAACCCGTGCTGTAAGCATATGTTGTCATGGCAATTTGATTCAATGGTGGGACCGCTGGAA

480 490 500 510 520 530 540 550 560
GGTGGATAGCCTCCCTGGAGCCCAAGGACAGAAATTTCCCTGACCCCAAGTCAAGAAGTCTTTGTACACATCCTACGAACGTGCGCTCAAGTGG

570 580 590 600 610 620 630 640 650
GAGCTGGTGGACCTTCTAGCCGACCGCTACTGGGAGGAGTACTAGACACATTCGCGCGGACATCGTGGTTAAGGACTGGTTTGTGTCAGAG

660 670 680 690 700 710 720 730 740 750
CCGACTGTGGCTGCACCTACCAACTCAAAGTGCAGCTGGCCTGGGCTGACTACTTGTGTTGGCCTCCTTCGAGCCCCCACCCTGTGACCATCCA

760 770 780 790 800 810 820 830 840
ACAGTGAACAATGCCACATGGACAGAGGTCTCCTACACCTTCTCAGACTACCCCGGGGTGTCGGCTACATCCTCTCCAGCATGGGGGAGG

850 860 870 880 890 900 910 920 930 940
GACACCCAGTACTGGGCAGGCTGGTATGGGCCCCGAGTCACCAACAGCAGGATTTGCTCAGCCCCCAAGATGACCAGGAACCAAGGCTCGTCCG

FIG.9B

950 960 970 980 990 1000 1010 1020 1030
AGGCTCAGCCCTGGGCAGAGCAATGGACAGGAGGAGGCTGCCCAATGCCCTACGGAGCTGTGTCCAGATTTTCIGACAGCTGTCCATCCCTGTG
1040 1050 1060 1070 1080 1090 1100 1110 1120
TCGGGTACGCCAGAGGTTCCATCCAGGCAGGAGCTGAGCATGGGTGGGCAGTGAGGTCCCTGTACCAGCGACTCCTGCCCCGGTTCAACCCCTA
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
CCAGCTTGTTGTTAACTTACTGTACATAGCTCTGACGTTTTGTTGTAATAAATGTTTTICAGGCCGGGCACCTGTGGCTCAGCCCTGTAAATCCCAG
1230 1240 1250 1260 1270 1280 1290 1300 1310
CACTTTGGGAGACCGAGGAGGTGGATCAGGAGGTACGAGACAGAGACCATCCTGGCCCAACACGGTGAAACCCCTGTCTCTACTAAAAATACAA
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
AAATTAGCCGGCGGTGGTGGCGGGCGCCTGTAGTCCCAGCTACTCGGGAGGCTGATGCAGAGAATGGCGTGAACCCCGGAAGGCAGAGCTTGC
1420 1430 1440 1450 1460 1470 1480 1490 1500
AGTGAGCCCGAGATCAGGCCACATGCACCTCCAGCCCTGGGTGGGTGACAGAGCGGAGACTCTGGCTCATAAAAATAATAATAATAAAAAATA
1510 1520 1530
AATGGTTTTCAGTAAAAAATAAAAAAATAAAAAAATAAAAAAATAAAAAAATAAAAAAATAAAAAAATAAAAAAATAAAAAAATAAAAAAATA

FIG.9C

10 20 30 40 50 60
MSNTRFTITLNYKDPLTGDEETLASYGIVSGDLICLI LHDDIPPPNIPSSDSEHSSLQN

70 80 90 100 110 120
NEQPSLATSSNQTSIQDEQPSDSFQQQAAQSGVWNDDSM LGPSQNF EAESI QDNAHMAEG

130 140 150 160 170 180
TGFYPSEPLLCSESV EQVPHSLETLYQSADCS DANDALIVL I HLLMLES GYIPQGTEAK

190 200 210 220 230 240
ALSLPEKWKLSGVYKLQYMHHLCEGSSATLTCVPLGNLIVVNATLKINNEIRSVKRLQLL

250 260 270 280 290 300
PESFICKEKLGENVANIYKDLQKLSRLFKDQLVYPLLAFTRQALNLPNVFGLVVLPLELK

310 320 330 340 350 360
LRIFRLLDVRSVLSLSAVCRDLFTASNDPLLWRF LYLRDFRDNTVRVQD TDWKEL YRKRH

370 380 390 400 410 420
IQRKESPKGRFVLLLPSSTHTIPFYPNLHPRFPSSRLPPGIIGGEYDQRPTLPYVGDP

430 440 450 460 470 480
ISSLIPGPGETPSQLPPLRPRFDPVGPLPGPNPILPGRGGPNDRFPFRPSRGRPTDGRLS

FM

FIG.10A

10 20 30 40 50 60 70 80 90
TGAATTCCCATGGACCATGCTCTAATAACCGGATTACAAATTACATTGAACCTACAAGGATCCCCCTCAGTGGAGATGAAGAGACCTTGGCTTCATA
100 110 120 130 140 150 160 170 180
TGGGATTGTTTCTGGGACATTGATATGTTTGATTCTTCAGGATGACATCCACCGCCTAATAATACCTTCATCCACAGATTACAGACCATTCCTTCA
190 200 210 220 230 240 250 260 270 280
CTCCAGAACAAATGAGCAACCCCTCTTTGGCCACCAGCTCCAATCAGACTAGCATACAGGATGAACAACCAAGTGATTTCATTCCAGGACAGGCAG
290 300 310 320 330 340 350 360 370
CCCAGTCTGGTGTGGAAATGACGACAGTAATGTTAGGGCCCTAGTCAAAAATTTGAAGCTGAGTCAATTCACAGATAATGGGCATATGGCAGAGGG
380 390 400 410 420 430 440 450 460 470
CACAGGTTTCTATCCCTCAGAACCCCTGCTCTGTAGTGAATCGGTGGAAGGGCAAGTGCCACATTCATTAGAGACCTTGATCAATCAGCTGAC
480 490 500 510 520 530 540 550 560
TGTTCTGATGCCAATGATCGGTTGATAGTGTGATACATCTTCATGTTGGAGTCAGGTTACATAACCTCAGGGCACCCGAAGCCAAAGCACTGT
570 580 590 600 610 620 630 640 650
CCCTGCCGGAGAAGTGAAGTTGACGGGGGTGTATAAGCTGCAGTACATGCATCATCTCTGGAGGGCAGCTCCGCTACTCTCACCTGTGTGCC
660 670 680 690 700 710 720 730 740 750
TTTGGGAAACCTGATTGTTGTAAATGCTACACTAAAAATCAACAATGAGATTAGAAGTGTGAAAAGATTGCAGCTGCTACCAGAACTCTTTTATT
760 770 780 790 800 810 820 830 840
TGCAAGAGAAACTAGCGGAAAATGTAGCCCAACATATACAAGAATCTTCAGAAACTCTCTCGCCCTCTTTAAAGACCAGCTGGTGTATCCTCTTC
850 860 870 880 890 900 910 920 930 940
TGGCTTTTACCCGACAAGCACTGAACCTACCAAATGTAATTTGGGTGGTGGTCTCCCTCCCAATTGGAACCTGAAACTAGCACTACGGATCTTCCGACTTCTGGA

FIG.10B

950 960 970 980 990 1000 1010 1020 1030
TGTTGGTCCGCTCTGCTTTGCTGCGGTTTGCTGACCTCTTACTGCTTCAAAATGACCCACATCCCTGGAGGTTTTTATATCTGCGTGAT
1040 1050 1060 1070 1080 1090 1100 1110 1120
TTTCGAGACAATACTGTCAGAGTTCAAGACACAGATTGGAAGAACTGTACAGGAAGAGGCCACATACAAAGAAAAGAAATCCCCGAAAGGCGGT
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
TTGTGCTGCTCCTGCCATCGTCAACCCACACCAATTCCATTCTATCCCAACCCCTTGCAACCTAGGCCATTTCCTAGCTCCCGCCTTCCTCCAGG
1230 1240 1250 1260 1270 1280 1290 1300 1310
AATTATCGGGGGTGAAATATGACCAAGACCAACACTTCCCTATGTTGGAGACCCCAATCAGTTCACATTCCTGGTCCCTGGGAGACGCCACG
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
CAGTTACCTCCACTGAGACCACGCTTTGATCCAGTTGGCCCACTTCCAGGACCTAACCCCACTCTGCCAGGGGAGGGGCCCCCAATGACAGAT
1420 1430 1440 1450 1460 1470 1480 1490 1500
TTCCCTTTAGACCCAGCAGGGGTCGGCCAACTGATGGCCGCCGTGTCATTCACTGATGATTGTTGTAATTTTCATTTCTCGAGCTCCATTGTTTT
1510 1520 1530 1540 1550 1560 1570 1580 1590
TGTTTCTAAACTACAGATGTCACCTCCTTGGGGTGGTGAATCTCGAGTGTTATTTTCTGATTGCTGGTGTGAGAGTTGCACTCCCAAGAACCTTTT
1600 1610 1620 1630 1640 1650 1660 1670 1680 1690
AAGAGATACATTTATAGCCCTAGGGGTGGTATGACCCCAAAGGTTCCTCTGTGACAAGGTGGCCCTTGGCAATAGTTGGCTGCCAATCTCCCTGC
1700 1710 1720 1730 1740 1750 1760
TCTTGGTCTCCTCTAGATTGAAGTTTGTTTCTGATGCTGCTTACCAGATTAAAAAAAAGTCTAAATT

FIG.10C

10	20	30	40	50	60
ETSKLG*SAVLAPAAGGTLSSSEGRSAVSGILIAVTSTGVDK*SLNQLLHGLGTSSRLSHF					
70	80	90	100	110	120
PFG*KSPPRGQFVAAAVEIAGRSGLQMGQGLWRVVRNQQLQQEGYSEQGYLTREQSRRMA					
130	140	150	160	170	180
ASNISNTNHRKQVQGGIDIYHLLKARKSKEQEGFINLEMLPPELSFTILSYLNATDLCLA					
190	200	210	220	230	240
SCVWQDLANDELLWQGLCKSTWGHCSIYNKNPPLGFSFRKXYMQLDEGSLTFNANPDEGV					
250	260	270	280	290	300
NYFMSKGILDDSPKEIAKFIFCTRNLNWKRLIYLDERRDVLDDLVTLNHFRNQFLPNAL					
310	320	330	340	350	360
REFFRHIHAPEERGEYLETLITKFSHRFCACNPDLRELGLSPDAVYVLCYSLILLSIDL					
370	380	390	400	410	420
TSPHVKNKMSKREFIRNTRRAAQNISSEDFVGHLNYDNIYLIGHVAA*KAQLLGLQFLLQTK					
430	440	450	460	470	480
ATQGLSRYGGYISAGHCSLSIQSSFVQPFLLPFSILVISLGN*IILQNFS*FCLSRFA					
490	500	510	520	530	540
QSRATV*HSC*RMIN*HYTLKDGVFVH*ICLKNFIFHFSLYKYHVMCTYLTKEIYSHNYF					
550	560	570	580	590	600
IVKILTKVFPFLSN*VLKFI*F*SETIVXVKVRSDFRQKPIPASFSFKL*RVLICYYITM					
610	620	630	640	650	
QNWQLFL*YKFI*FFILKTGLIKSR*VL*TI*DF*NIKIYDLHS*E*NKIXLELW					

FIG.11A

10 20 30 40 50 60 70 80 90
GGAACGTC AAAAT TGGGATAGTCGGCAGTTCTGCCCCCTGCAGCTGGAGGTACCCCTGAGTTCTGAGGGTGGTAGTCTGTTTCTGGTATCTC
100 110 120 130 140 150 160 170 180
ATCGCGTCACCTCTACCGGTGTTGGACAAGTAAAGTTTGAATCAGCTTCTCCATGGCCCTGGGCAACAGTTCCCGGCTGAGGCCATTTTCCCTTTTG
190 200 210 220 230 240 250 260 270 280
GCTAAAGTCCCGCCCAAGAGGCCAATTCTGCGCGCGCGTGGAGATCGCAGGTGGCTCAGGCTTGCAGATGGGTCAAGGGTTGTGGAGAGT
290 300 310 320 330 340 350 360 370
GGTCAGAAACAGCAGCTGCAACAAGAGGCTACAGTGAGCAAGGCTACCTCACCAGAGAGCAGAGCAGAGAAATGGCTGGCAGCAACATTTCT
380 390 400 410 420 430 440 450 460 470
AACACCAATCATCGTAACAAGTCCAAGGAGGCATTGACATATATCATCTTTTGAAGGCAAGCAATCGAAGAACAGGAAAGGATTCAATTAAT
480 490 500 510 520 530 540 550 560
TGGAAATGTTGGCTCCCTGAGCTAAGCTTTACCATCTTGCTCTACCCTGAAATGCAACAGCTTGGCTTGGCTTCAATGTTTGGCAGGACCTTGC
570 580 590 600 610 620 630 640 650
GAAATGATGAACCTTCCTGCGCAAGGTTGTCGCAATCCACTTGGGGTGACGTGTTCCATATACAATAAGAACCCACCTTTAGGATTTCTTTTAGA
660 670 680 690 700 710 720 730 740 750
AAATGTATATCCAGCTGGATGAAGGCAGCCCTCACCCTTTAATGCCAACCCAGATGAGGAGTGAACCTACCTTAAATGTCGAAGGTAATCTGGAAG
760 770 780 790 800 810 820 830 840
ATTGCGCAAGGAAATAGCAAAAGTTTATCTTCGTACAAGAACACTAAATTCGAAAAAACTGAGAAATCTATCTTGTGTAAGGAGAGATGCTCT
850 860 870 880 890 900 910 920 930 940
GGATGACCTTGTAAACATTGCATAATTTTAGAAATCAGTTCTTCCCAATGCACCTGAGAGAAATTTTTCGTATATCCATGCCCTGAAGAGCGT

FIG.11B

950 960 970 980 990 1000 1010 1020 1030
GGAGAGTATCTTGAAGCTCTTATAACAAGTCTCACATAGATTCTGTGCTTGCAACCCGATTTAATGCGAGAACTTGGCCCTTAGTCCCTGATG
1040 1050 1060 1070 1080 1090 1100 1110 1120
CIGCTAIGTACTGTGCTACTCTTTGATTCTACTTTCCATTGACCTCAGTACCCCTCACTGAGCAATGAAGAAATAAAATGTCAAAAAGCGAATTATTTCG
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
AAATACCGTCGGCTGCTCAAAATATTAGTGAAGATTTGTAGGCACTTTAAGACAATACTACCTTATTGGCCAATGTCGCTGCATAAAAA
1230 1240 1250 1260 1270 1280 1290 1300 1310
GCACAATTGCTAGGACTTCAGTTTTTACTTCAGACTAAAGCTACCCAAGGACTTAGCAGATAATGGGGTTACATCAGTGGTCATTGTAGCC
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
TGAGTATACAATCAAGCTTCAGTGTGCAACCTTTTTTCTTTGGCACTTTCTAATTTAGTAAATTTCCCTTGGGGAACATAAATAATTTTGCAGAA
1420 1430 1440 1450 1460 1470 1480 1490 1500
TTTTTCCTAATTTTGTATTACAGTTTTGCACAAAGCAGAGCCACTGTCTAACACAGCTGTAAACGAATGATAAACTGACATTATACTCTAAAA
1510 1520 1530 1540 1550 1560 1570 1580 1590
GATGGTGATTTGTGCATTAGATTGCCGTGAAAACTTTATCCATTCCATTCCTTTATACAAATACCAATGTAATGTGTACATAATTTAACTAAAG
1600 1610 1620 1630 1640 1650 1660 1670 1680 1690
AGATTATAGTCATAAATTAATTTATTGTAAAGATTTTAAGTAAAGTTTTTCCCTTTTCTCICCAACIGAGTCTCIGAAAATTTATTGTGATTCGATC

FIG.11C

1700 1710 1720 1730 1740 1750 1760 1770 1780
TGAAC TATTG TCTCY CGTAA AGTTAG AICTG ACTTC AGRCAG AACC AATACC AGCTT CCTTT CCTTT AAAC TTGA AGAG TGTG ATTGT
1790 1800 1810 1820 1830 1840 1850 1860 1870 1880
TACTA TATTAC TATG CAAAC TGGC AGTTAT TTTTAT ATATA TAAAT TTAAT TTTG ATTTT TTTT AAAC TGGT TTAAT CAAG CTCGGT
1890 1900 1910 1920 1930 1940 1950 1960 1970
AAGTCC TTTAA ACCATT TAGGAT TTTTAA AACATC AAAATTT ATGAT TTACA TTCA TAGGA TAAAA TAAAT ATYATT AGAA CTCGGT

FIG.11D

10	20	30	40	50	60
MAAAVDSAMEVVPALAEAAPEVAGLSCLVNLPGEVLEYILCCGSLTAADIGRVSSSTCR					
70	80	90	100	110	120
RLRELCQSSGKVWKEQFRVRWPSLMKHYSPTDYVNWLEEYKVRQKAGLEARKIVASFSCR					
130	140	150	160	170	180
FFSEHVPCNGFSDIENLEGPEIFFEDELVCILNMEGRKALTWKYYAKKILYYLRQQKILN					
190	200	210	220	230	240
NLKAFLQQPDDYESYLEGAVYIDQYCNPLSDISLKDIIQAQIDSIVELVCKTLRGINSRHP					
250	260	270	280	290	300
SLAFKAGESSMIMEIELQSQVLDAMNYVLYDQLKFKGNRMDYYNALNLYMHQVLIRRTGI					
310	320	330	340	350	360
PISMSLLYLTIIARQLGVPLEPVNFPSHFLLRWCQGAEGATLDIFDYIYIDAFGKGKQLTV					
370	380	390	400	410	420
KECEYLIGQHVTAALYGVVNVKKVLQRMVGNLLSLGKREGIDQSYQLLRDSL DLYLAMYP					
430	440	450	460	470	480
DQVQLLLLQARLYFHLGIWPEKVL DILQHIQTLDPGQHGA VG YLVQHTLEHIERKKEEVG					
490	500	510	520	530	540
VEVKLRSDKHRDVCYSIGLIMKHKRYGYNCVIYGDPTCMMGHEWIRNMNVHSLPHGHH					
550	560	570	580	590	600
QPFYNVLVEDGSCRYAAQENLEYNVEPQEISHPDVG RYFSEFTGTHYIPNAELEIRYPED					
610	620				
LEFVYETVQNIYSAKKENIDE					

FIG.12A

10 20 30 40 50 60 70 80 90 100 110 120 130
 CATGGCGGGCAGCAGTCGACAGGGCGATGGAGGTGGCGGGGGCTGGCGAGAGGGCGGGCGGAGGTAGCGGGCTCAGCTGGCTGGTCAACCTGCCGGCTGAGGTGGCTGGAGTACATCTCTGCTGGCGCTCG
 140 150 160 170 180 190 200 210 220 230 240 250 260 270
 CTGACGGGGCGGACATCGGGCGCTCTCCAGCACTGCCGGGGCTGCCGAGCTGCCAGAGCAGCGGGAAGGTGTCGAGAGCAGTTCGGGGTGAGGTGGCGTCCCTATGAAACACTACAGCCCCACCGACT
 280 290 300 310 320 330 340 350 360 370 380 390 400 410
 ACGTCAATTGGTTGGAAGAGTATAAGTTCCGCCAAAAGCTGGGTTAGAGCGCGGAAGATGTAGCCCTGCTCTCAAGAGGTTCCTTTCAGAGCAGCTTCCTTGTAATGGCTTCAGTGACATTGAGAACCTTGAAGG
 420 430 440 450 460 470 480 490 500 510 520 530 540 550
 ACCAGAGATTTTITGAGGATGAACGGTGTGTATCTTAATATGGAAGGAAGAAAGCTTTGACCTGGAATACTAGGCAAAAAAATCTTTACTAGCTGGCGCAACAGAGATCTTAAATAATCTTAAGSCCTTT
 560 570 580 590 600 610 620 630 640 650 660 670 680 690
 CTTACGAGCCAGATGACTGAGTGGTATCTTGAAGTGGTGATATATGACCAGTACTGCAATCCCTCTCTCGGAGATCAGCCTCAAGAGATCCAGCCCCAAATTCAGAGCATCGTGGAGCTTGTTCGAAAAACCC
 700 710 720 730 740 750 760 770 780 790 800 810 820 830
 TTGGGGCATAACAGTGGCCACCCAGCTTGGCCCTTCAGGAGGTGAATCATCCATGATATGGAATAGAACTCAGAGCCAGGTGGCTGGATGCCATGAACATATGCTTTACGACCAACTGAAGTTCAAGGGGAA
 840 850 860 870 880 890 900 910 920 930 940 950 960 970
 TCGAATGCACTTACTATAATGCCCTCAACTTATATGCAATCAGGTTTGTGATTGGCAGAACAGGAATCCCAATCAGCATGTCTCTGCTATTTGACAATTCCTGGGAGTTGGAGTCCCACTGGAGCCTGTCAACTTC
 980 990 1000 1010 1020 1030 1040 1050 1060 1070 1080 1090 1100 1110
 CCAAGTCACTTCTTATTAGGTGGTCCGAGCGGAGAGGGGGACCCCTGGACATCTTTGACTACATCTACATAATGCTTTTGGGAAGGGAAGCAGCTGACAGTGAAGAATGGCAGTACTTGAICGGGGCAGCAGG
 1120 1130 1140 1150 1160 1170 1180 1190 1200 1210 1220 1230 1240 1250
 TGACTGAGGACGTATGGGGTGGTCAATGTCAAGAGGTGTACAGAGAAATGGTGGGAACCTGTTAAGCCTGGGGAAGCGGCAAGGCCATCGACCAATCATACCAAGTCCCTGAGAGACTCGCTGGATCTCTATCTGGC
 1260 1270 1280 1290 1300 1310 1320 1330 1340 1350 1360 1370 1380 1390
 AATGTACCGGGACCGAGGTGCAGCTTCCTCTCTCCAAAGCCAGGCTTTACTTCCACCTTGGGAAATCTGGCCAGAGAGGTGCTTGACATCTCTCCAGCAGATCCAAACCCCTAGACCGGGGAGCAGCGGGCGGTGGGCTAC

FIG. 12B

10 20 30 40 50 60
RSTGFRRAGEEWSR+XLAASPGXLRRPAXTFVLSNLAEVVERVLTF LPAKALLRVACVCR
70 80 90
LWRECVRRVLRTHRSVTWISAGLAEGHLXGH

FIG.13A

10 20 30 40 50 60
CCGTAGTACTGGNTTCCGGCGGGCTGGTGAGGAATGGAGCCGGTAGNTGCTTGCGGCGAG
70 80 90 100 110 120
TCCCGGGNTCCTCCGTAGACCCGCGGANACCTTCGTGTTGAGTAACCTGGCGGAGGTGGT
130 140 150 160 170 180
GGAGCGTGTGCTCACCTTCCTGCCCCGCAAGGCGTTGCTGCGGGTGGCCTGCGTGTGCCG
190 200 210 220 230 240
CTTATGGAGGGAGTGTGTGCGCAGAGTATTGCGGACCCATCGGAGCGTAACCTGGATCTC
250 260 270
CGCAGGCCTGGCGGAGGCCGCCACCTGGNGGGGCATT

FIG.13B

10	20	30	40	50	60
RPRPVQQQQQPPQQPPPPQQQQPPPPPPQQQQQPPPPPPPPPLPQERNNVG					
70	80	90	100	110	120
ERDDDVADMAEESGPGAQNSPYQLRRKTL L PKRTACPTKNSMEGASTSTTENFGHRAK					
130	140	150	160	170	180
RARVSGKSQDL SAAPAEQYLQEKL PDEVVLKIFSYLLEQDLCRAACVCKRF SELANDPNL					
190					
WKRL YMEVF EYTRPMMH					

FIG.14A

10	20	30	40	50	60
GCGGCCGCGCCCGGTGCAGCAACAGCAGCAGCAGCCCCCGCAGCAGCCGCCGCCGCGCAGCC					
70	80	90	100	110	120
GCCCCAGCAGCAGCCGCCCGCAGCAGCAGCCTCCGCCGCCGCCGAGCAGCAGCAGCAGCA					
130	140	150	160	170	180
GCAGCCTCCGCCGCCGCCACCGCCGCTCCGCCGCTGCCTCAGGAGCGGAACAACGTCCG					
190	200	210	220	230	240
CGAGCGGGATGATGATGTGCCTGCAGATATGGTTGCAGAAGAATCAGGTCCTGGTGACA					
250	260	270	280	290	300
AAATAGTCCATACCAACTTCGTAGAAAACTCTTTTGCCGAAAAGAACAGCGTGTCAC					
310	320	330	340	350	360
AAAGAACAGTATGGAGGGCGCCTCAACTTCAACTACAGAAAACCTTTGGTCATCGTGCAAA					
370	380	390	400	410	420
ACGTGCAAGAGTGTCTGGAAAATCACAAGATCTATCAGCAGCACCTGCTGAACAGTATCT					
430	440	450	460	470	480
TCAGGAGAACTGCCAGATGAAGTGGTTCTAAAAATCTTCTTACTTGCTGGAACAGGA					
490	500	510	520	530	540
TCTTTGTAGAGCAGCTTGATGTATGTAAACGCTTCAGTGAACCTTGCTAATGATCCCAATTT					
550	560	570	580	590	
GTGGAACGATTATATATGGAAGTATTTGAATATACTCGCCCTATGATGCAT					

FIG.14B

10 20 30 40 50 60
RPRPGLRGGRAPCEVTMEAGGLPLELWRMILAYLHLPDLGRCSLVCRAWYELILSLDSTR
70 80 90 100 110 120
WRQLCLGCTECRHPNWPNQPDVEPESWREAFKQHYLASKTWTKNALDLESSICFSLFRRR
130 140 150 160 170
RERRTLSVGPGREFDSLGSALAMASLYDRIVLFPGVYEEQGEIILKVPVEIVGQGKLG

FIG.15A

10 20 30 40 50 60
GCGGCCGCGGCCCGGACTCCGCGGTGGGCGAGCGCCCTGTGAGGTGACCATGGAGGCTGG
70 80 90 100 110 120
TGGCCTCCCCTTGGAGCTGTGGCGCATGATCTTAGCCTACTTGCACCTTCCCGACCTGGG
130 140 150 160 170 180
CCGCTGCAGCCTGGTATGCAGGGCCTGGTATGAACTGATCCTCAGTCTCGACAGCACCCG
190 200 210 220 230 240
CTGGCGGCAGCTGTGTCTGGGTTGCACCGAGTGCCGCCATCCCAATTGGCCCAACCAGCC
250 260 270 280 290 300
AGATGTGGAGCCTGAGTCTTGGAGAGAAGCCTTCAAGCAGCATTACCTTGCATCCAAGAC
310 320 330 340 350 360
ATGGACCAAGAATGCCTTGGACTTGGAGTCTTCCATCTGCTTTTCTCTATTCCGCCGGAG
370 380 390 400 410 420
GAGGGAACGACGTACCCTGAGTGTGGGCCAGGCCGTGAGTTTGACAGCCTGGGCAGTGC
430 440 450 460 470 480
CTTGGCCATGGCCAGCCTGTATGACCGAATTGTGCTCTTCCCAGGTGTGTACGAAGAGCA
490 500 510 520 530
AGGTGAAATCATCTTGAAGGTGCCTGTGGAGATTGTAGGGCAGGGGAAGTTGGGTGA

FIG.15B

10 20 30 40 50 60
ETETAPLTLESLPTDPLLLILSFLDYRDLINCCYVSRRLSQLSSHDP LWRRHCKKYWLIS

70 80 90 100 110 120
EEETQKNQCWKSLEIDTYSDVGRYIDHYAAIKKASGMISRNWSPGVLGWVLSLKEGCS

130 140 150 160 170 180
RGRPRCCGSADWAASFLLDYRCSYRIHNGQKLVGSGYWEAWHCLITIVLKIC*TSIQLP

190 200 210 220 230 240
EIPAETGTEILSPFNFCIHTGLSQYIAVEAAEG*NKNEVFYQCQTVVERVFKYGIKMCSDG

250
CINGMH*VFS

FIG.16A

10	20	30	40	50	60
GAGACCGAGACGGCGCCGCTGACCCTAGAGTCGCTGCCACCGATCCCCTGCTCCTCATC					
70	80	90	100	110	120
TTATCCTTTTTGGACTATCGGGATCTAATCAACTGTTGTTATGTCAGTCGAAGATTAAGC					
130	140	150	160	170	180
CAGCTATCAAGTCATGATCCGCTGTGGAGAAGACATTGCAAAAAATACTGGCTGATATCT					
190	200	210	220	230	240
GAGGAAGAGAAAACACAGAAGAATCAGTGTTGGAAATCTCTTTCATAGATACTTACTCT					
250	260	270	280	290	300
GATGTAGGAAGATACATTGACCATTATGCTGCTATTA AAAAGGCCCTCGGAATGATCTCA					
310	320	330	340	350	360
AGAAATATTTGGAGCCCAGGTGTCCTCGGATGGGTTTTATCTCTGAAAGAGGGGTGCTCG					
370	380	390	400	410	420
AGAGGAAGACCTCGATGCTGTGGAAGCGCAGATTGGGCTGCAAGTTTCCTGGACGATTAT					
430	440	450	460	470	480
CGATGTTTCATACCGAATTCACAATGGACAGAAGTTAGTTGGTTCCTGGGGTTATTGGGAA					
490	500	510	520	530	540
GCATGGCACTGTCTAATCACTATCGTTCTGAAGATTTGTTAGACGTCGATACAGCTGCCG					
550	560	570	580	590	600
GAGATTCCAGCAGAGACAGGGACTGAAATACTGTCTCCCTTTAACTTTGCATACATACT					
610	620	630	640	650	660
GGTTTGAGTCAGTACATAGCAGTGGAAGCTGCAGAGGGTTGAAACAAAATGAAGTTTTC					
670	680	690	700	710	720
TACCAATGTCAGACAGTAGAACGTGTGTTTAAATATGGCATTAAAGATGTGTTCTGATGGT					
730	740	750			
TGTATAAATGGCATGCATTAGGTATTTTCAG					

FIG.16B

10 20 30 40 50 60
GSGFRAGGWPLTMPGKHQHFQEPEVGCCGKYFLFGFNIVFWLGALFLAIGLWAWGEKGV
70 80 90 100 110 120
LSNISALTDLGGLDPVWLVCGSWRRHVGAGLCWAAIGALRENTFLLKFFXXFLGLIFFLE
LA

FIG.17A

10 20 30 40 50 60
GGCTCCGGTTTCCGGGCCGGCGGTGCCCGCTACCATGCCCGGNAAGCACCAGCATTTTC
70 80 90 100 110 120
CAGGAACCTGAGGTCGGCTGCTGCGGAAATACTTCCTGTTTGGCTTCAACATTGTCTTC
130 140 150 160 170 180
TGGGTGCTGGGAGCCCTGTTCTGCTATCGGCCTCTGGGCCTGGGGTGAGAAGGGCGTT
190 200 210 220 230 240
CTCTCGAACATCTCAGCGCTGACAGATCTGGGAGGCCTTGACCCCGTGTGGCTTGTTTGT
250 260 270 280 290 300
GGTAGTTGGAGCGTCATGTCGGTGCTGGGCTTTGCTGGGCTGCAATTGGGGCCCTCCGG
310 320 330 340 350 360
GAGAACACCTTCCTGCTCAAGTTTTCTNCGNGTTCCTCGGTCTCATCTTCTTCCTGGAG
CTGGCAAC

FIG.17B

10 20 30 40 50 60
AAAAAAYLDELPEPLLLRVLAALPAAELVQACRLVCLRWKELVDGAPLWLLKCQQEGLVP

70 80 90 100 110 120
EGGVEEERDHWQQFYFLSKRRRNLLRNPCGEEDLEGWCDVEHGGDGWRVEELPGDSGVEF

130 140 150 160 170 180
THDESVKKYFASSFEWCRKAQVIDLQAEGYWEELDTTQPAIVVKDWYSGRSDAGCLYEL

190 200 210 220 230 240
TVKLLSEHENVLAEFSSGQVAVPQSDGGGWMEISHTFTDYGPGVRFVRFEHGGQGSVYW

250
KGWFGARVTNSSVWVEP*

FIG.18A

10 20 30 40 50 60
GCGGCGGCCGCCGCGTACCTGGACGAGCTGCCCCAGCCGCTGCTGCTGCGCTGCTGGCCGCACTG
70 80 90 100 110 120 130
CCGCGCGCCGAGCTGGTGCAGGCCTGCCGCCGCTGGTGTGCCTGCGCTGGAAGGAGCTGGTGGACGGCGCC
140 150 160 170 180 190 200
CCGCTGTGGCTGCTCAAGTGCCAGCAGGAGGGGCTGGTCCCCGAGGGCGGCGTGGAGGAGGAGCGCGAC
210 220 230 240 250 260 270
CACTGGCAGCAGTTCTACTTCCTGAGCAAGCGGCCGCAACCTTCTGCGTAACCCGTGTGGGAAGAG
280 290 300 310 320 330 340
GACTTGAAGGCTGGTGTGACGTGGAGCATGGTGGGACGGCTGGAGGTGGAGGAGCTGCCTGGAGAC
350 360 370 380 390 400 410
AGTGGGTGGAGTTCACCCACGATGAGAGCGTCAAGAAGTACTTCGCCTCCTCTTTGAGTGGTGTGCG
420 430 440 450 460 470 480
AAAGCACAGGTCATTGACCTGCAGGCTGAGGGCTACTGGGAGGAGCTGCTGGACACGACTCAGCCGGCC
490 500 510 520 530 540 550
ATCGTGGTGAAGGACTGGTACTCGGGCCGACGACGCTGGTTGCCTCTACGAGCTCACCCTTAAGCTA
560 570 580 590 600 610 620
CTGTCCGAGCACGAGAACGTGCTGGCTGAGTTCAGCAGCGGGCAGGTGGCAGTGCCCCAAGACAGTGAC
630 640 650 660 670 680 690
GGCGGGGGCTGGATGGAGATCTCCACACCTTCACCGACTACGGGCCGGGCGTCCGCTTCGTCCGCTTC
700 710 720 730 740 750
GAGCACGGGGGGCAGGGCTCCGTCTACTGGAAGGGCTGGTTCGGGGCCGGGTGACCAACAGCAGCGTG
760 770
TGGGTAGAACCCTGA

FIG.18B

10 20 30 40 50 60
MGEKAVPLLRRRRVKRSCPSGSELGVEEKRGKGNPISIQLFPELVEHIISFLPVRDLV

70 80 90 100 110 120
ALGQTCRYFHEVCDGEGVWRRICRRLSPRLQDQDTKGLYFQAFGGRRRCLSKSVAPLLAH

130 140 150 160 170 180
GYRRFLPTKDHVFILDYVGTLFFLKNALVSTLGQMQRACRYVVLRCGAKDFASDPRCD

190 200 210 220 230 240
TVYRKLYVLATREPQEVVGTTSRACDCVEVYLQSSGQRVFKMTFHHSMTFKQIVLVGQ

250 260 270 280 290 300
ETQRALLLLTEEGKIYSLVVNETQLDQPRSYTVQLALRKVSHYLPRLRVACMTSNQSSTL

310
YVTDPILCWLQPPWPGG

FIG.19A

10 20 30 40 50 60
 ATGGGCGAGAAGGCGGTCCCTTTGCTAAGGAGGAGGCGGTGAAGAGAAGCTGCCCTTCTTGTGGCTCG
 70 80 90 100 110 120 130
 GAGCTTGGGGTTGAAGAGAAGAGGGGAAAGGAAATCCGATTTCCATCCAGTTGTTCCCCCAGAGCTG
 140 150 160 170 180 190 200
 GTGGAGCATATCATCTCATTCCCTCCAGTCAGAGACCTTGTTGCCCTCGGCCAGACCTGCCGCTACTTC
 210 220 230 240 250 260 270
 CACGAAGTGTGGCATGGGGAAGGCGTGTGGAGACGCATCTGTGCGAGACTCAGTCCGCGCCTCCAAGAT
 280 290 300 310 320 330 340
 CAGGACACGAAGGGCCTGTATTTCCAGGCATTTGGAGGCCGCCCGCATGTCTCAGCAAGAGCGTGGCC
 350 360 370 380 390 400 410
 CCCTTGCTAGCCACGGCTACCGCCGCTTCTTGCCACCAAGGATCACGTCTTCATTCTTGACTACGTG
 420 430 440 450 460 470 480
 GGGACCCTCTTCTTCTCAAAAATGCCCTGGTCTCCACCCTCGGCCAGATGCAGTGAAGCGGGCCTGT
 490 500 510 520 530 540 550
 CGCTATGTTGTGTTGTGTCGTGGAGCCAAGGATTTGCCTCGGACCCAAGGTGTGACACAGTTTACCGT
 560 570 580 590 600 610 620
 AAATACCTCTACGTCTTGGCCACTCGGGAGCCCGAGGAAGTGGTGGGTACCACCAGCAGCCGGGCCTGT
 630 640 650 660 670 680 690
 GACTGTGTTGAGGTCTATCTGCAGTCTAGTGGGCAGCGGGTCTTCAAGATGACATTCCACCACTCAATG
 700 710 720 730 740 750
 ACCTTCAAGCAGATCGTGCTGGTTGGTCAGGAGACCCAGCGGGCTCTACTGCTCCTCACAGAGGAAGGA
 760 770 780 790 800 810 820
 AAGATCTACTCTTTGGTAGTGAATGAGACCCAGCTTGACCAGCCACGCTCCTACACGGTTCAGCTGGCC
 830 840 850 860 870 880 890
 CTGAGGAAGGTGTCCCACTACCTGCCTCACCTGCGGTGGCCTGCATGACTTCCAACCAGAGCAGCACC
 900 910 920 930 940 950
 CTCTACGTCACAGATCCTATTCTGTGCTCTTGCTACAACCACCTTGGCCTGGTGGATGA

FIG.19B

10	20	30	40	50	60
RGGSEGRGRGREKRARGARRKRKQGGREARAADGEGSGPGAEGARTRPREEAEGGGSV					
70	80	90	100	110	120
EEGARGIIKGDEGSVGAGKEAQGRKYGKEEWRVRARRREGARPGRVQGGQVWAYIPGT					
130	140	150	160	170	180
GAAMAAAAREEEEEAARESAACPAAGPALWRLPEVLLHMC SYLDMRALGRLAQVYRWLW					
190	200	210	220	230	240
HFTNCDLLRRQIAWASLNSGFTRLGTNLMTSVPVKVSQNWIVGCCREGILLKWRC SQMPW					
250	260	270	280	290	300
MQLEDDALYISQANFILAYQFRPDGASLNRQLGVSAGHDEDVCHFVLATSHIVSAGGDG					
310	320	330	340	350	360
KIGLGKIHSTFAAKYWAHEQEVNCDCKGGIISFGSRDRTAKWPLASGQLGQCLYTIQT					
370	380	390	400	410	420
EDQIWSVAIRPLLSSFVTGTACCGHFSPLKIWDLNSGQLMTHLDRDFPPRAGVLDVYES					
430	440	450	460	470	480
PFALLSCGYDTYVRYWDCRTSVRKCVMEWEPHNSTLYCLQTDGNHLLATGSSFYSVRL					
490	500	510	520	530	
WDRHQRACPHTFPLTSTRLGSPVYCLHLTTKHLAALSYNLHVLDIQNP*					

FIG.20A

10 20 30 40 50 60 70 80 90
CGACGGGAAGCGAAGCGGAAGAGGAAGCGGAAGCGGCAAGCGGAAGCGGCGGAAGCGGAAGCGGCGG
100 110 120 130 140 150 160 170 180
CAGACGGCGAAGGAGCGCGCGCGGGCTGAGCGGGAGCGAGGACAGCCCAAGAGAGGAGGAGCGGAGCGGAGCGGAGG
190 200 210 220 230 240 250 260 270 280
GGCGAGAGGCATCATCAAGGAGATGAGGGAGCGTAGGGCCCGGAAGAGGCCACAAGGAAGTAATGCGAAGGAAATGCGAGGTCAGG
290 300 310 320 330 340 350 360 370
GCTAGCGCGCGGAGGGCGCCAGCGCGGAAGAGTACAAGGACAAGGAGGTACGTTTGGGCCCTACATCCCGGGGACAGGGCGGCCATGGCGG
380 390 400 410 420 430 440 450 460 470
CGCGAGCCAGGAGGAGGAGGAGCGCGCTCGGGAGTCAGCCGCCCTGCCCGGCTGCGGGGCCACGGCTCTGGCGCCTCGCGGAAGTGTCT
480 490 500 510 520 530 540 550 560
GCTGCACATGTGCTCCTACCTCGACATGCGGGCCCTCGCGCGCTGCGCCAGGTGTACCGCTGGCTGTGGCACTTCACCAACTGCGACCTGCTC
570 580 590 600 610 620 630 640 650
CGCGCCAGATAGCCTGGGCTCGCTCAACICCGGCTTACCGCGCTCGGCACCAACCTGATGACCAGTGTCCCAGTGAAGGTGCTCAGAACT
660 670 680 690 700 710 720 730 740 750
GGATAGTGGGTGCTGCCGAGAGGGGATTCTGCTGAAGTGGAGATGCAGTCAGATGCCCCTGGATGCACCTAGAGGATGATGCTTTGTACATATC
760 770 780 790 800 810 820 830 840
CCAGGCTAATTTTCATCCTGGCCTACCAGTTCCGTCCAGATGGTCCAGCTTGAACCGTCAGCCCTCGGAGTCTCTGCTGGGCATGATGAGGAC
850 860 870 880 890 900 910 920 930 940
GTTTGGCACCTTGCTGCGCCACCTCGCATATTGTGTCAGTGCAGGAGGAGATGGGAAGATTGGCCCTGGTAAGATTACACAGCACCTTCGCTGCCA

FIG.20B

950 960 970 980 990 1000 1010 1020 1030
AGTACTGGGCTCATGAACAGGAGGTGAACCTGTGTGGATTGCAAGGGGGCATATATCATTTGGCTCCAGGGACAGGACGGCCAAGGTGTGGCC
1040 1050 1060 1070 1080 1090 1100 1110 1120
TTTGGCCTCAGGCCAGCTGGGGCAGTGTTTATACACCATCCAGACTGAAGACCAAACTCTGGTCTGTGCTATCAGGCCATTACTCAGCTCTTTT
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
GTGACAGGGACGGCTTGTTGTGGGCACCTTCACCCCTGAAATCTGGGACCTCAACAGTGGGACGCTGATGACACACTTGGACAGAGACTTTC
1230 1240 1250 1260 1270 1280 1290 1300 1310
CCCCAAGGGCTGGGGTGCTGGATGCATATATGAGTCCCTTTGGCACTGCTCTCCTGTGGCTATGACACCTATGTTGGCTACTGGGACTGCCG
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
CACCAGTGTCGGAAATGTGTCATGGAGTGGGAGGAGCCCCACAACAGCACCTGTACTGCCCTGCAGACAGATGGCAACCCTTGCTTGGCCACA
1420 1430 1440 1450 1460 1470 1480 1490 1500
GGTTCCCTTCTATAGCGTTGTACGGCTGTGGGACCGGACCAAGGGCCCTGCCCGCACACCTTCCCGCTGACGTCCGCCCTCGGCAGCC
1510 1520 1530 1540 1550 1560 1570 1580 1590
CTGTGTACTGCCCTGCATCTCACCACCAAGCATCTCTATGCTGGCGTGTCTTACAACCTCCACGTCCTGGATATTCAAACCCGTGA

FIG. 20C

10 20 30 40 50 60
L I L T S V L L F Q R H G Y C T L G E A F N R L D F S S A I Q D I R T F N Y V V K L L Q L I A K S Q L T S L S G V A Q K
70 80 90 100 110 120
N Y F N I L D K I V Q K V L D D H H N P R L I K D L L Q D L S S T L C I L I R G V G K S V L V G N I N I W I C R L E T I
130 140 150 160 170 180
L A W Q Q Q L Q D L Q M T K Q V N N G L T L S D L P L H M L N N I L Y R F S D G W D I I T L G Q V T P T L Y M L S E D R
190 200 210 220 230 240
Q L W K K L C Q Y H F A E K Q F C R H L I L S E K G H I E W K L M Y F A L Q K H Y P A K E Q Y G D T L H F C R H C S I L
250 260 270
F W K D S G H P C T A A D P D S C F T P V S P Q H F I D L F K F

FIG.21A

10 20 30 40 50 60
 GCATTGCTATAATTTTACTATACTCTCATCTAAATCTAAAATCAGTCTTCAAATAAAAAACAAATTGTC
 70 80 90 100 110 120 130
 CTTTGCCAAAAATTTTTTAATCGCACAATTAATTGACATTAAGTCCCAATTCTTTTTGGCTAATTGAC
 140 150 160 170 180 190 200
 TAATTTTAAGTCTGTGTGCTTTTCCAGAGGCATGGCTATTGCACCTTGGGAGAAGCCTTTAATCGGT
 210 220 230 240 250 260 270
 TAGACTTCTCAAGTGCAATTCAGATATCCGAACGTTCAATTATGTGGTCAAAGTGTTCAGCTAATTG
 280 290 300 310 320 330 340
 CAAAATCCAGTTAACTTCATTGAGTGGCGTGGCACAGAAGAATTACTTCAACATTTTGGATAAAATCG
 350 360 370 380 390 400 410
 TTCAAAGGTTCTTGATGACCACCACAATCCTCGCTTAATCAAAGATCTTCTGCAAGACCTAAGCTCTA
 420 430 440 450 460 470 480
 CCCTCTGCATTCTTATTAGAGGAGTAGGGAAGTCTGTATTAGTGGGAAACATCAATATTTGGATTGGCC
 490 500 510 520 530 540 550
 GATTAGAACTATTCTCGCCTGGCAACAACAGCTACAGGATCTTCAGATGACTAAGCAAGTGAACAATG
 560 570 580 590 600 610 620
 GCCTCACCTCAGTGACCTTCTCTGCACATGCTGAACAACATCCTATACCGTTCTCAGACGGATGGG
 630 640 650 660 670 680 690
 ACATCATCACCTTAGGCCAGGTGACCCACGTTGTATATGCTTAGTGAAGACAGACAGCTGTGGAAGA
 700 710 720 730 740 750
 AGCTTTGTCAGTACCATTTTGCTGAAAAGCAGTTTGTAGACATTTGATCCTTTCAGAAAAAGGTCATA
 760 770 780 790 800 810 820
 TTGAATGGAAGTTGATGTACTTTGCACTTCAGAAACATTACCCAGCGAAGGAGCAGTACGGAGACACAC
 830 840 850 860 870 880 890
 TGCATTTCTGTGGCACTGCAGCATTCTTTTTGGAAGGACTCAGGACACCCCTGCACGGCGGCCGACC
 900 910 920 930 940 950 960
 CTGACAGCTGCTTACGCCTGTGTCTCCGCAGCACTTCATCGACCTTTCAAGTTTTAAGGGCTGCCCC

FIG.21B

970 980 990 1000 1010 1020 1030
TGCCATCCCTATTGGAGATTGTGAATCCTGCTGTCTGTGCAGGGCTCATAGTGAGTGTCTGTGAGGTG

1040 1050 1060 1070 1080 1090 1100
GGTGGAGACTCCTCGGAAGCCCCTGCTTCCAGAAAGCCTGGGAAGAACTGCCCTTCTGCAAAGGGGGGA

1110 1120 1130 1140 1150 1160 1170
CTGCATGGTTGCATTTTCATCACTGAAAGTCAGAGGCCAAGGAAATCATTCTACTTCTTTAAAACTC

1180 1190 1200 1210
CTTCTAAGCATATTAAAATGTGAAATTTGCGTACTCTCTC

FIG.21C

10	20	30	40	50	60
YGSEKGSSSISSDVSSSTDHTPTKAQKNVATSESDLSMRTLSTPSPALICPPNLPGFQ					
70	80	90	100	110	120
NGRGSSTSSSSI TGETVAMVHSPPTRLTHPLIRLASRPQKEQASIDRLPDHSMVQIFSF					
130	140	150	160	170	180
LPTNQLCRCARVCRRWYNLAWDPRLWRTIRLTGETINVDRALKVLTRRLCQDTPNVCLML					
190	200	210	220	230	240
ETVTVSGCRRRLTDRGLYTI AQCCPELRRLEVSGCYNISNEAVFDVVS LCPNLEHLDVSGC					
250	260	270	280	290	300
SKVTCISLTREASIKLSPLHGKQISIRYLDMTDCFVLEDEGLHTIAAHCTQLTHLYLRRC					
310	320	330	340	350	360
VRLTDEGLRYLVIYCASIKELSVSDCRFVSDFGLREIAKLESRLRYLSIAHCGRVTDVGI					
370	380	390	400	410	420
RYVAKYCSKLRYLNARGCEGITDHGVEYLAKNCTKLKSLDIGKCPLVSDTGLECLALNCF					
430	440	450	460	470	480
NLKRLSLKSCESI TGQGLQIVAANCFDLQTLNVQDCEVSVEALRFVKRHCKRCVIEHTNP					

AFF

FIG.22A

10 20 30 40 50 60 70 80 90 100 110 120 130
 AGTACGGCAGTGAGGGCAAGGCAGCTCGAGCATCTCATCTGAGGTGAGTTCAAGTACAGATCACAGGCCCACTAAAGCCCAAGAAATGTGGCTACCAAGCAAGAACTCCGACCTGAGCATGGCGCACACTGAGCACGGCC
 140 150 160 170 180 190 200 210 220 230 240 250 260 270
 CAGCCAGCCCTGATATGTCCACCAATCTCCAGGATTTTCAGAAATGGAAGGGGCTGGTCCACCTCCTCGTCTCTCCATACCGGGGAGACGGTGGCCATGGTGCACACTCCCGGCCCGGACCGGCTTACACACCCGGCTC
 280 290 300 310 320 330 340 350 360 370 380 390 400 410
 ATCCGGCTGGCTCCAGACCCCAAGAGGAGGCGGAGGATAGACCGGCTCCCGGACCACTCCATGGTGGAGATCTTCTCCTTCCTGGCCCAACCAAGCAGCTGTGCGGCTGGCGCGGAGTGTGCGCGCGCTGGTACAAC
 420 430 440 450 460 470 480 490 500 510 520 530 540 550
 TGGCTGGGACCGCGGCTCTGGAGGACTATCCGGCTGACGGGCGAGACCATCAACGTGGACCGCGCCCTCAAGGTGCTGACCGCGAGACTCTGCCAGGACACCCCAAGGTGTGCTCATGCTGGAACCGTAACTGT
 560 570 580 590 600 610 620 630 640 650 660 670 680 690
 CAGTGGCTGCAGGGGCTCACAGACGAGGCTGTACACCATCGCCCGAGTGTGCCCGCACTGAGCGGACTGGAAGTCTCAGGCTGTACAATACTCCAAGAGGGGCTCTTTGATGTGGTGTCCCTCTGCCCTAAAT
 700 710 720 730 740 750 760 770 780 790 800 810 820 830
 CTGAGGACCTGGATGTCTAGGATGCTCCAAAGTGACCTGACATCAGCTTGACCCCGGAGGGCTCCATTAAACTGTCACCCCTTGCAATGGCAAGAGATTTCATCCGCTACCTGGACATGACGGACTGCTTGGTGGTGG
 840 850 860 870 880 890 900 910 920 930 940 950 960 970
 AGCAAGGGCTGCACACCATCGCGGGCCACTGCAGGAGCTACCCACCTCTACCTGCGCGGCTGGCTCGGCTGACCGAGGAGGGCTGGCTACCTGGTGATCTAC TGGCGCTCCATCAAGGAGGTGAGGCTCAG
 980 990 1000 1010 1020 1030 1040 1050 1060 1070 1080 1090 1100 1110
 CCAGTGGCGCTTGGTACGGACTTTCGGGCTGGGGAGATCGCCCAAGCTGGAGTCCCGGCTGGGTAACCTGAGCATCGCGCACTGGCGCGGGTCAACGAGTGGGCA TCGGCTAGCTGGCCCAAGTACTGACGCAAGCTG
 1120 1130 1140 1150 1160 1170 1180 1190 1200 1210 1220 1230 1240 1250
 CCTACCTCAACGGAGGGCTGGAGGGCATCACGGACCAAGGTGAGGTAACCTGGCCAGAACTGCACCAAACTCAAAATCCCTGGATATCGGCAAAATGCGCTTTGGTATCCGACAGGGGCTGGAGTGGCTGGCGCC
 1260 1270 1280 1290 1300 1310 1320 1330 1340 1350 1360 1370 1380 1390
 TGAAGTCTTCAACCTCAAGGGGCTCAGGCTCAAGTCTCGGAGAGCATCACCGGCGGAGGGCTTGCAGATGCTGGCGGCAACTGCTTTGACCTCCAGAGGCTGAATGTCCAGGACTGGAGGTCTCCGTGGAGGGCCCT

FIG. 22B

2790 2800 2810 2820 2830 2840 2850 2860 2870 2880 2890 2900 2910
 TGTAGTGTTAATTGTCGAATGGCACCCTGTGTACCTCCTCATGTCGTCGCGTGTTCACCAAGAATGCCAAGCAGACTTCCAGGTGTTAAATCTCTGCTACACAAATGCCAGATGAATGCCAAGG
 2920 2930 2940 2950 2960 2970 2980 2990 3000 3010 3020 3030 3040 3050
 GAACACACTGACATGACTTAGACTCTGGTCCACCACCAGACCCCTGGAAAGGAATACTAAATCATTTACAGGTATGGATTTAAATGGATGAACCTTCAAAATTAATCTATTGGATAGAAGTCTATATTCTAGCCCTC
 3060 3070 3080 3090 3100 3110 3120 3130 3140 3150 3160 3170 3180 3190
 ATTTGCATGAAGTCAGATAGCCAGAGAAATTCATTTGCTGGTTTTCAGCAAAATTCACCTTGTCTTTGCTAATAACACATGGCCCTTTCACAGATTATCTCTAGCCCAAGCCCAACCTTTGTTACGTTGAATCCCTC
 3200 3210 3220 3230 3240 3250 3260 3270 3280 3290 3300 3310 3320 3330
 ATTTATTTCTTCTCAAAATGCCCATTAICCAAAATGCAGAACCTCTGCATCTCCAGGAGTTAAGCTGAATTTGTCAAACTTAGACACCCCTTGACAACTGCACACTGCTACTGTAGGCTCCTGTGCACTACTGCTGCTC
 3340 3350 3360 3370 3380 3390 3400 3410 3420 3430 3440 3450 3460 3470
 TGTGGGGATGGAGAGTTAGTGTGATGAGGTGGTGTCTGCCAGGAGTTTCTTCAACATCATGGCTCCCATCCAATCAACATCATCAAAATACATGTGTAATCAAGGCTCTGTGCCATGGGGAAATGAATCAT
 3480 3490 3500 3510 3520 3530 3540 3550 3560 3570 3580 3590 3600 3610
 TTAGCTAGGCCAGCATCTAGTGAAGCCACAGAGTTTAAACCATGAAGAAGTTGAAGCAGCATTCCTCAGCTCTGTGACTGTGACCCATTGGAAGTTTCAGGATTTGGGTCTCACAAAGGATTGTCCCTAATCC
 3620 3630 3640 3650 3660 3670 3680 3690 3700 3710 3720 3730 3740 3750
 TTGGCCCTGGGGTCTTCCGAGTGAGTGTTTAATACTCTGAGAAATGAGCAGGAGATCCAGAGAAATGAATCCCTGACCGCATCACCCTAAACTGCTTCCAAACATGACAGACAAGCTGACTGTTTACACTGATTGCCCA
 3760 3770 3780 3790 3800 3810 3820 3830 3840 3850 3860 3870 3880 3890
 GCACATACCGCTTGGCAGTTTCTCTTCTCCAGTCTCCTGTTCAATCTCTTCCCTTGGGGTGGGAATCTATGATGAGGTTACTGGGGAACAGCTCAGCAGATTTTGGAGACCAAAACCAAGGTCTC
 3900 3910 3920 3930 3940 3950 3960 3970 3980 3990 4000 4010 4020 4030
 ACTAGGAAATTAATCTGTTTAAACATGCTTCCCTGGCTCTGCTAAATGAAGTCAATGTTGTTGTTGTTTAAATCTAATGTTCAAACTCAGCTGCGTGGTGTATGAATCTAGAAAGCCCTTAATTA

FIG. 22D

CTACCAAGAAATAAGCAATAIGTCTG

10 20 30 40 50 60
AAAPAPAPAPTPTPEEGPDAGWCDRIPLLEILVQIFGLLVAADGMPFLGRAARVCRRWQE

70 80 90 100 110 120
AASQPALWHTVTLSSPLVGRPAKGGVKAEEKLLASLEWLMPNRFSQLQRLTLIHWKSQVH

130 140 150 160 170 180
PVLKLVGECCPRLTFLKLSGCHGVTADALVMLAKACCQLHSLDLQHSMVESTAVVSFLEE

190 200 210 220 230 240
AGSRMRKLWLTYSQTTAILGALLGSCCPQLQVLEVSTGINRNSIPLQLPVEALQKGCPQ

250 260 270 280
LQVLRLLNLMWLPKPPGRGVAPGPGFPSLEELCLASSTCNFVS

FIG.23A

```

      10      20      30      40      50      60
TGCGGCGCGCCCGCACCCGCACCGGCACCCACGCCACGCCGAGGAAGGGCCCGACGCGGGCTGGGG

70      80      90      100     110     120     130
AGACCGCATTCCCTTGAAATCCTGGTGCAGATTTTCGGGTTGTTGGTGGCGGCGGACGGCCCCATGCC

140     150     160     170     180     190     200
CTTCCTGGGCAGGGCTGCGCGCGTGTGCCGCCGCTGGCAGGAGGCCGCTTCCCAACCCGCGCTCTGGCA

210     220     230     240     250     260     270
CACCGTGACCCGTGTCGTCCCGCTGGTCGGCCGCGCTGCCAAGGGCGGGGTCAAGGCGGAGAAGAAGCT

280     290     300     310     320     330     340
CCTTGCTTCCCTGGAGTGGCTTATGCCCAATCGGTTTTACAGCTCCAGAGGCTGACCCATCCACTG

350     360     370     380     390     400     410
GAAGTCTCAGGTACACCCCGTGTTGAAGCTGGTAGGTGAGTGCTGTCTCGGCTCACTTTCCTCAAGCT

420     430     440     450     460     470     480
CTCCGGCTGCCACGGTGTGACTGCTGACGCTCTGGTCATGCTAGCCAAAGCCTGCTGCCAGCTCCATAG

490     500     510     520     530     540     550
CCTGGACCTACAGCACTCCATGGTGGAGTCCACAGCTGTGGTGAGCTTCTTGGAGGAGGCAGGGTCCCG

560     570     580     590     600     610     620
AATGCGCAAGTTGTGGCTGACCTACAGCTCCCAGACGACAGCCATCCTGGGCGCATTGCTGGGCAGCTG

630     640     650     660     670     680     690
CTGCCCCCAGCTCCAGGTCCTGGAGGTGAGCACCGGCATCAACGTAATAGCATTCCCTTCAGCTGCC

700     710     720     730     740     750
TGTCGAGGCTCTGCAGAAAGGCTGCCCTCAGCTCCAGGTGCTGCGGCTGTTGAACCTGATGTGGCTGCC

760     770     780     790     800     810     820
CAAGCCTCCGGGACGAGGGTGGCTCCCGACCAGGCTTCCCTAGCCTAGAGGAGCTCTGCCTGGCGAG

830     840     850
CTCAACCTGCAACTTTGTGAGC

```

FIG.23B

10	20	30	40	50	60
QHCSQKDTAELLRGLSLWNHAEERQKFFKYSVDEKSDKEAEVSEHSTGITHLPPEVMLSI					
70	80	90	100	110	120
FSYLN PQELCRCSQVSMKWSQLTKTGSLWKHLYPVHWARGDWYSGPATELDTEPDDEWVK					
130	140	150	160	170	180
NRKDESRAFHEWDEDADIDESEESAESIAISIAQMEKRLLHGLIHNVL P YVGTSVKTLV					
190	200	210	220	230	240
LAYSSAVSSKMVRQILELCPNLEHLDLTQTDISDSAFDSWSWLGCCQSLRHLDLSGCEKI					
250	260	270	280	290	300
TDVALEKISRALGILTSHQSGFLKTSTSKITSTAWKNKIDITMQSTKQYACLHDLTNKGIG					
310	320	330	340	350	360
EEIDNEHPWTKPVSSENFTSPYVWMLDAEDLADIEDTVEWRHRNVESLCVMETASNFCSS					
370	380	390	400	410	420
TSGCFSKDIVGLRTSVCWQQHCASPAFAYCGHSFCCTGTALRTMSSLP ESSAMCRKAART					
430	440	450	460	470	480
RLPRGKDLIYFGSEKSDQETGRVLLFLSLSGCYQITDHGLRVLTGGGLPYLEHLNLSGC					
490	500	510	520	530	540
LTI TGAGLQDLVSACPSLNDEYFYCDNINGPHADTASGCQNLQCGFRACCRSGE*PLTS					
550	560	570	580	590	
DLCLLHLAEQAFFHALYS*HISCVNHPFLSVTCFGPIXYNFRNLNYQXIVML					

FIG.24A

10 20 30 40 50 60 70 80 90
ACAACACTGCTCTCAGAAAGGATAC TGCAGAACTCCTTAGAGGCTTAGCCCTATGGAATCATGCTGAAGAGCGACAGAARTTTTTAAATATCC

100 110 120 130 140 150 160 170 180
GTGGATGAAAAAGTCAGATAAAGAAGCAGAAGTGTCAGAAACACTCCACAGGTATAACCCCATCTTCCCTGAGGTAATGCTGCTCAATTTTCAGCT

190 200 210 220 230 240 250 260 270 280
ATCTTAATCCTCAAGAGTTATGTCGATGCAGTCAAGTAAGCATGAAATGGTCTCAGCTGACAAAAACGGGATCGCTTTTGGAAACATCTTTACCC

290 300 310 320 330 340 350 360 370
TGTTTCATTGGGCCAGAGTGACTGGTATAGTGGTCCCGCAACTGAACCTGATGATGAATGGTGAATAATAGGAAAGATGAA

380 390 400 410 420 430 440 450 460 470
AGTCGTGCTTTTCATGAGTGGGATGAAGATGCTGACATTGATGAATCTGAAGAGTCTCGGAGGAATCAATTCCTATCAGCAATGCACAAATGG

480 490 500 510 520 530 540 550 560
AAAAACGTTTACTCCATGGCTTAATTCATAACGTTCTACCATAATGTTGGTACTCTGTAAAAACCTTAGTATTAGCATACAGCTCTGCAGTTTC

570 580 590 600 610 620 630 640 650
CAGCAAAATGGTTAGCCAGATTTTAGAGCTTTGTCCTAACCTGGAGCACTCGGATCTTACCCAGACATGACATTTTCAGATTTTCGCAATTTGACAGT

660 670 680 690 700 710 720 730 740 750
TGGICTTGGCTTGGTTGCTGCCAGAGCTTCGGCATCTTGATCTGCTGCTGGTTGTGAGAAAAATCACAGATGCGCCCTAGAGAAGATTTCCAGAG

760 770 780 790 800 810 820 830 840
CTCTTGGAAATTCGACATCTCATCAAAAGTGGCTTTTGTGAAACATCTACAAGCAAAATTACTTCAACTGCGTGGAAAAATAAAGACATTACCAT

850 860 870 880 890 900 910 920 930 940
GCAGTCCACCAAGCAGTATGCCCTGTTTGCACGATTAACATAACAAGGGCATTTGGAGAAGAAATAGATAATGAACACCCCTGGACTAAGCCTGTT

FIG. 24B

950 960 970 980 990 1000 1010 1020 1030
TCTTCTGAGAAATTCACCTTCCTTATGTGTGGATGTTAGATGCTGAAGATTTGGCTGATATTTGAAGATACTGTGGAAATGGAGACATAGAAATG
1040 1050 1060 1070 1080 1090 1100 1110 1120
TTGAAAGTCTTTTGTCTAATGGAACAGCATCCAACCTTAGTTGTTCCACCCTCTGGTTGTTTAGTAAGGACATTTGTTGGACTAAGGACTAGTGT
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
CTGTTGGCAGCAGCATTTGTGCTTCCTCAGCCTTTGGGTATTGTGGTCACTCAATTTTGTGTACAGGAACAGCCTTTAAGAACTAIGTCATCACTC
1230 1240 1250 1260 1270 1280 1290 1300 1310
CCAGAACTCTTCGCAATGTGTAGAAAGCAGCAAGCACTAGATTGCCTAGCGGAAAGACTTAAATTTACTTTGGGAGTGAAGAAATCTGATCAAG
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
AGACTGGAGGTGACTTCGTCTCAGTTTATCTGGATGTTATCAGATCAGACAGACCATGGTCTCAGGGTTTTGACTCTGGGAGGAGGGCTGCC
1420 1430 1440 1450 1460 1470 1480 1490 1500
TTATTTGGAGCACCTTAAATCCTCTGCTTACTATACTGGTGCAGGCCCTGCAGGATTTGGTTTACCATGTGCTTCTCIGAAATGATGAA
1510 1520 1530 1540 1550 1560 1570 1580 1590
TACTTTTACTACTGTGACAACATTAACGGTCCCTCATGCTGATACCGCCAGTGGATGCCAGAAATTTGCAGTGTGGTTTTCCAGCCCTGCTGCCGCT
1600 1610 1620 1630 1640 1650 1660 1670 1680 1690
CTGGCGAATGACCCCTTGACTTCGTGATCTTTGTCTACTTCACTTTAGCTGAGCAGGCTTCTTTTCATGCACCTTACTCATAGCACATTTCTTGTGT
1700 1710 1720 1730 1740 1750 1760 1770
TAACCATCCCTTTTGGCGGTGACTGTGTTGGGCCCCATTNYTTACAACCTTCAGAAATCTTAATTACCAGTGRATGTAAATGTG

FIG. 24C

10	20	30	40	50	60
RVTSGCGLARGSSAMVFSNNDEGL INKKLPKELLRIFSFLDIVTLRCRAQISKAWNILA					
70	80	90	100	110	120
LDGSNWQRIDLFNFQIDVEGRVVENISKRCVGFLRKLSLRGCIGVGDSSLKTF AQNCRNI					
130	140	150	160	170	180
EHLNLNGCTKITDSTCYSLSRFC SKL KHLXLTSCVSI TNSSLKG ISEGCRNLEYLNL SWC					
190	200	210	220	230	240
DQITKDGIEALVRGCRGLKALLRGCTQLEDEALKHIQNYCHELVSLNLQSCSRITDEGV					
250	260	270	280	290	300
VQICRGCHRLQALCLSGCSNLTDASLTALGLNCPRLQILEAARCSHLTDAGFTLLARNCH					
310	320	330	340	350	360
ELEKMDLEXCILITDSTLIQLSIHCPKLQALSLSHC ELIXDDGILHLSNSTCGHERLRVL					
370	380	390	400	410	420
ELDNCLLITDVALXHLENCRGLERLELYDCQQVTRAGIKRMRAQLPHVKVHAYFAPVTPP					
430	440	450	460	470	480
TAVAGSGQRLCRCCVIL*QQLPGPKG**GILSSRRPESS*PTPPSPNLLILHWERHLQFP					
490	500	510	520	530	540
NRHLSRFKNGEDKKGFI SNI*HHIVT*NMALT*LVLLLPSSLMSSLTSTHLLL*YL*RLI					
550					
ILKTDQGTGPASKYINCVQ*					

FIG.25A

10 20 30 40 50 60 70 80 90
TTTTACTGTACAGTTGATGTAATTTTGATGCTGGGCCCTGCTCTGCTCTTGAGGATTATTAACCTTTAGAGGTATCAGAGAAGCAAAATGGG

100 110 120 130 140 150 160 170 180
TACTGGTGAGGCTGCTCATTAGGGAAGAGGGCAAAAGGAGGCACTAGCTAGGTACAGGCCATGTTTCAGGTACACAATGTGATGTCAGATGTGCT

190 200 210 220 230 240 250 260 270 280
TATAAATCCCTTCTTGCTTCGCCATTCTTAATCTTGATAGGTGCCTGTTGGGAAACTGTAAATGCCCTTCCCAATGGAGAATCAACAGATTG

290 300 310 320 330 340 350 360 370
GGTGATGGTGGAGTCGGTCAGGAAGACTCAGGTCTTCTACAGGAAGGATGCCCTCATCACCCCTTNGGCCCAGGCAGCTGCTGTCAGAGAATGA

380 390 400 410 420 430 440 450 460 470
CACAGCACCTGCACAGTCGCTGCCACTTCCTGCCACATGCTGCGGTGACGGGAGCAAGTAGGCGTGGACTTTGACATGAGGGAGCTG

480 490 500 510 520 530 540 550 560
AGCCCGCATCCGCTTGATGCCCTGCACGGGTAAACCTGCTGGCAGTCGTACAGCTCGAGGGCTCCAGGCCCTCGGCAGTTCTCTAGGTGTYCCAGG

570 580 590 600 610 620 630 640 650
GCCACATCAGTGATGAGGAGGCAGTTGTCCAACATCCAGTACCCCGCAGCCCTCTCATGCCCCACAGGTACGTGCTCAGGTGCAGGATCCCATCAT

660 670 680 690 700 710 720 730 740 750
CTGKGATGAGTTCACAGTGGGACAGGCTCAGGGCTTGCAGTTTAGGACAGTGAAATGGAGAGCTGGATGAGTGTGCTGTGCGTTATCAGGATGCA

760 770 780 790 800 810 820 830 840
WTCCTCAAGATCCATCTCTCCAAATTCGTGGCAATTCGGAGCTAAAAGTGTAACCTGCGTCAGTCAAAATGGGAGCATCGGGCAGCCCTCCCAA

FIG.25B

850 860 870 880 890 900 910 920 930 940
ATTGCCAGTCGCGGACAGTTCAAACCCAGGGCTGTAAGAGAGGCATCTGTGAGGTTGCTGCAACCCGAAAGGCAGAGAGCCTGTAGCCGGGTGAC

950 960 970 980 990 1000 1010 1020 1030
AGCCCCIGCATATCTGCACCACACCTTCATCCGTGATACGTGAGCAGGACTGCAAGTTGAGGCTCACAAGCTCATGCGAGTAATTCIGAAITG

1040 1050 1060 1070 1080 1090 1100 1110 1120
TTTCAGAGCTTCATCTTCTAACTGTGTGAGCCCCCTCAGGAGCAGGGCTTTCAGGCCCTCGACAACCTTCGCACCAGTGCCTCGATGCCATCCTTC

1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
GTGATCTGATCACACCAAGAGAGGTTCAGGTACTCCAGGTTTCGGCAGCCCCCTCAGTGTATCCCTTCAAGGAGGCTGTTTGTAAATAGACACACAGG

1230 1240 1250 1260 1270 1280 1290 1300 1310
AGGTCAGAWCCAGATGTTTCAGCTTGGAACAGAACTGCTAAGGCTATAACACCGTGCCTGTCAGTGAATTTTGTGCAATCCATTGAGGTTCAAAATG

1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
TTCAAATGTTTCGGCAGTCTGTGCAAGGTCTTCAAGGAGGAAATCCCCAACACCAATGCAGCCCTCGCAAGCTGAGCTTCCCTCAGGAATCCAACG

1420 1430 1440 1450 1460 1470 1480 1490 1500
CATCGCTTCGAGATATTTCCACCACCTCGACCCCTACATCTATTTGAAAAGTTAAAAAGATCTATTTCTTTGCCAGTTGCTTCCATCCAGGGCTA

1510 1520 1530 1540 1550 1560 1570 1580 1590
AGATGTTCCAAGCCTTGGAATCTGTGCACATCGGCACAAAAGTTACTATATCCAAGAAGGAAAAATATTTCTTAACAGAAGTTCCTTTGGGTAACCTT

1600 1610 1620 1630 1640 1650 1660 1670 1680
TTTGTTAATAAGGCCTTCATCATGTTTGAGAAAAACCATGGCCGGAAGAGCGCGGAGCGGACAGCCCAAGTCAACACGCGC

FIG.25C

10	20	30	40	50	60
MSPVFPMLTVLTMFYIICLRRRARTATRGEMMNTTHRAIESNSQTSPLNAE VVQYAKEVVD					
70	80	90	100	110	120
FSSHYGSENSMSYTMNLAGVPNVFPSSGDF TQTAVFRTYGTWWDQCPSASLPFKRTPPN					
130	140	150	160	170	180
FQSQDYVELTFEQQVYPTAVHVLE TYHPGAVIRILACSANPYSPNPPAEVRWEILWSERP					
190	200	210	220	230	240
TKVNASQARQFKPCIKQINFPTNLIRLEVNSSLLEYTELDAVVLHGVKDKPVL SLKTSL					
250	260	270	280	290	300
IDMNDIEDDAYAEKDGCGMDSL NKKFSSAVLGEGPNNGYFDKLPYELIQLILNHLTLPDL					
310	320	330	340	350	360
CRLAQTCKL LSQHCCDPLQYIHLNLQPYWAKLDDTSLEFLQSRCTLVQWLNLSWTGNRGF					
370	380	390	400	410	420
ISVAGFSRFLKVCGSELVRLELSCSHFLNETCLEVISEMCPNLQALNLSSCDKLPPQAFN					
430	440	450	460	470	480
HIAKLCSLKRLVLYRTKVEQTALLSILNFCSELQHL SLGSCVMIEDYDV IASMI GAKCKK					
490	500	510	520	530	540
LRTL DLWRCKNITENGIAELASGCPLLEELDLGWCPTLQSSTGCFTRLAHQLPNLQKFL					
550	560	570	580	590	600
TANRSVCDTDIDELACNCTRLQQLDILGTRMVSPASLRKLL ESCKDLSLLDVSFCSQIDN					
610	620				
RAVLELNASFPKVF IKKSFTQ					

FIG.26A

10 20 30 40 50 60 70 80 90
ATGTCACCGGTCCTTCCCATGTTAACAGTTCGACCATGTTTATTATATATGCCCTTCGGCCCGAGCCAGGACAGCTACAAAGAGGAGAAATGA

100 110 120 130 140 150 160 170 180
TGAACACCCATAGAGCTATAGAATCAAACAGCCAGACTTCCCCTCTCAATGCAGAGGTAGTCCAGTATGCCAAAGAAGTAGTGGATTTCAGTTTC

190 200 210 220 230 240 250 260 270 280
CCATTATGGAAGTGAGAATAGTAGTCCTATACTATGTGGAATTTGGCTGGTGTACCAAAATGATTCCCAAGTTCCTGGTACTTTACTCAGACA

290 300 310 320 330 340 350 360 370
GCTGTGTTTCGAACTTATGGGACATGGTGGATCAGTGTCTTAGTCTTCCTTGGCATTCAAGAGGACGCCACCTAATTTTCAGAGCCAGCACT

380 390 400 410 420 430 440 450 460 470
ATGTGGAACTTACTTTTGAACAACAGGGTGTATCCTACAGCTGTACATGTTCTTAGAAACCTATCATCCCGGAGCAGTCATTAGAAATTCGCTTG

480 490 500 510 520 530 540 550 560
TTCIGCAAATCCTTATTCCCCAAATCCACCAGCTGAAGTAAGATGGGAGATTCTTTGGTCAGAGAGACCTACGAAGGTGAATGCTTCCCCAAGCT

570 580 590 600 610 620 630 640 650
CGCCAGTTTAAACCTTGATTAAAGCAGATAAAATTTCCCCACAAAATCTTATACGACTGGAAGTAAGTAGTTCCTCTCGGAATATTACACTGAAT

660 670 680 690 700 710 720 730 740 750
TAGATGCAGTTGTGCTACATGGTGTGAAGGACAAGCCAGTGCCTTCTCTCAAGACTTCACCTTATTGCATGAATGATATAGAAGATGATGCCCTA

760 770 780 790 800 810 820 830 840
TGCAGAAAAGGATGGTTGTGGAATGGACAGTCTTAACAAAAAGTTTAGCAGTGTCTTCCTCGGGGAAGGCCAAATAATGGGTAATTTTGATAAA

850 860 870 880 890 900 910 920 930 940
CTACCTTATGAGCTTATTCAGCTGATTCIGAATCATCTTACACTACCAGACCTGTGTAGATTAGCACAGACTTGCAAACTACTGAGCCAGCATT

FIG. 26B

950 960 970 980 990 1000 1010 1020 1030
GCTGTGATCCTCTGCAATACATCCACCTCAATCTGCAACCATACTGGGCAAACTAGATGACACTTCTCTGGAATTTCTACAGTCTCGCTGCAC
1040 1050 1060 1070 1080 1090 1100 1110 1120
TCTTGTCAGTGGCTTAATTTATCTTGGACTGGCAATAGAGGCTTCATCTCTGTTGCAGGATTTAGCAGGTTTCTGAAGGTTTGTGGATCCGAA
1130 1140 1150 1160 1170 1180 1190 1200 1210 1220
TTAGTAGCGCTTGAATTGCTCTTGCAGCCACCTTCTTAAAGAACTTGCCTTAGAAGTTATTTCTGAGATGTGTCCTCAAAATCTACAGGCCTTAAATC
1230 1240 1250 1260 1270 1280 1290 1300 1310
TCTCCTCCTGTGATAAGCTACCACCTCAAGCTTTCACACACATTGCCAAAGTTATGCCAGCCTTAAACGACTTGTCTCTATCGAACAAGTAGA
1320 1330 1340 1350 1360 1370 1380 1390 1400 1410
GCAACACGCACTGCTCAGCATTTTGAACCTTCTGTTTCAGAGCTTCAGCACCTCAGTTAGGCAGTTGTGTCATGATTGAAGACTATGATGTGATA
1420 1430 1440 1450 1460 1470 1480 1490 1500
GCTAGCATGATAGGAGCCAAGTGTAAAAAACTCCGGACCTGGATCTGTGGAGATGTAAGAAATATTACTGAGAAATAGCAGAACTGGCTT
1510 1520 1530 1540 1550 1560 1570 1580 1590
CTGGGTGTCCTCACTACTGGAGGAGCTTGACCTTGGCTGGTGGCCCAACTCTGCAGAGCAGCACCAGCGGTGCTTCACCAGACTGGCACACCAGCTCCC
1600 1610 1620 1630 1640 1650 1660 1670 1680 1690
AAACTTGCAAAAAATCTTTCTTACACCTAATAGATCTGCTGTGACACAGACATTGATGAATTGGCATGTAAATTGTACCAGGTTACAGCAGCTG
1700 1710 1720 1730 1740 1750 1760 1770 1780
GACATATTAGGAACAAGAAATGGTAAGTCGGCATCCTTAAGAAAACTCCTGGAATCTTGTAAAGATCTTTTCTTACTTGATGTGCTCTCTGTT
1790 1800 1810 1820 1830 1840 1850 1860
CGCAGATTGATAACAGAGCTGTGCTAGAACTGAATGCAAGCTTTCCTCAAAAGTGTTCATAAAAAAGAGCTTTTACTCAGTGA

FIG. 26C

10 20 30 40 50 60
MQLVPDIEFKITYTRSPDGDGVGNSYIEDNDDDSKMADLLSYFQQQLTFQESVLKLCQPE
70 80 90 100 110 120
LESSQIHISVLPMEVLMYIFRWVSSDLRLSLEQLSLVCRGFYICARDPEIWRACLKV
130 140 150 160 170 180
WGRSCIKLVPYTSWREMFLERPRVRF DGVYISKTTYIRQGEQSLDGFYRAWHQVEYYRYI
190 200 210 220 230 240
RFFPDGHVMMLTTPPEPQSIVPRLRTRNTRTDAILLGHYRLSQDTDNQTKVFAVITKKKE
250 260 270 280 290 300
EKPLDYKYRYFRRVPVQEQSFHVGLQLCSSGHQRFNKL IWIHHSCHITYKSTGETAVS
310 320
AFEIDKMYTPLFFARVRSYTAFSERPL

FIG.27A


```

      10      20      30      40      50      60
ATGCAACTTGTACCTGATATAGAGTTCAAGATTACTTATACCCGGTCTCCAGATGGTGTATGGCGTTGGA

70      80      90      100     110     120     130
AACAGCTACATTGAAGATAATGATGATGACAGCAAAATGGCAGATCTCTTGTCTACTTCCAGCAGCAA

140     150     160     170     180     190     200
CTCACATTTTCAGGAGTCTGTGCTTAACTGTGTGACGCTGAGCTTGAGAGCAGTCAGATTCACATATCA

210     220     230     240     250     260     270
GTGCTGCCAATGGAGGTCTGATGTACATCTTCCGATGGGTGGTGTCTAGTGACTTGGACCTCAGATCA

280     290     300     310     320     330     340
TTGGAGCAGTTGTGCTGGTGTGCAGAGGATTCTACATCTGTGCCAGAGACCCTGAAATATGGCGTCTG

350     360     370     380     390     400     410
GCCTGCTTGAAAGTTTGGGGCAGAAGCTGTATTAACTTGTTCGTACACGTCCTGGAGAGAGATGTTT

420     430     440     450     460     470     480
TTAGAACGGCCTCGTGTTTCGGTTTGATGGCGTGTATATCAGTAAAACCATATATTCGTCAAGGGGAA

490     500     510     520     530     540     550
CAGTCTCTTGATGGTTTTCTATAGAGCCTGGCACCAAGTGAATATTACAGGTACATAAGATTCTTTCCT

560     570     580     590     600     610     620
GATGGCCATGTGATGATGTTGACAACCCCTGAAGAGCCTCAGTCCATTGTTCCACGTTTAAGAACTAGG

630     640     650     660     670     680     690
AATACCAGGACTGATGCAATTCTACTGGGTCATATCGCTTGTCACAAGACACAGACAATCAGACCAA

700     710     720     730     740     750
GTATTTGCTGTAATAACTAAGAAAAAGAAGAAAAACCACTTGACTATAAATACAGATATTTTCGTCGT

760     770     780     790     800     810     820
GTCCCTGTACAAGAAGCAGATCAGAGTTTTTCATGTGGGGCTACAGCTATGTTCCAGTGGTCACCAGAGG

830     840     850     860     870     880     890
TTCAACAACTCATCTGGATACATCATTCTTGTCACATTACTTACAAATCAACTGGTGAGACTGCAGTC

900     910     920     930     940     950     960
AGTGCTTTTGAGATTGACAAGATGTACACCCCTTGTTCCTCGCCAGAGTAAGGAGCTACACAGCTTTC

970     980
TCAGAAAGGCCTCTGTAG

```

FIG.27B

10 20 30 40 50 60
AALDPDLENDFFVRKTGAFHANPYVLRAFEDFRKFSEQDDSVERRDIILQCREGELVLPD
70 80 90 100 110 120
LEKDDMIVRRIPAKKEVPLSGAPDRYHPVPFPEPWTLPPEIQAKFLCVLERTCPSKEKS
130 140 150 160 170 180
NSCRILVPSYRQKKDDMLTRKIQSWKLGTTVPPISFTPGPCSEADLKRWEAIREASRLRH
190 200 210 220 230 240
KKRLMVERLFQKIYGENGSKSMSDVSAEDVQNLRLRYEEMQKIKSQLKEQDQKWQDDLA
250
KWKDRRKSYTSDLQK

FIG.28A

10 20 30 40 50 60
 GCAGCCCTGGATCCTGACTTAGAGAATGATGATTTCTTTGTCAGAAAGACTGGGGCTTTCCATGCAAAT
 70 80 90 100 110 120 130
 CCATATGTTCTCCGAGCTTTTGAAGACTTTAGAAAGTTCTCTGAGCAAGATGATTCTGTAGAGCGAGAT
 140 150 160 170 180 190 200
 ATAATTTTACAGTGTAGAGAAGGTGAACTTGTACTTCCGGATTTGAAAAAGATGATATGATTGTTCCG
 210 220 230 240 250 260 270
 CGAATCCACGACAGAGAAGAAAGAAGTGCCGCTGTCTGGGGCCCCAGATAGATACCACCCAGTCCCTTTT
 280 290 300 310 320 330 340
 CCCGAACCCTGGACTCTTCCTCCAGAAATTCAAGCAAAATTTCTCTGTGTACTTGAAAGGACATGCCCA
 350 360 370 380 390 400 410
 TCCAAAGAAAAAGTAATAGCTGTAGAATATTAGTTCCTTCATATCGGCAGAAGAAAGATGACATGCTG
 420 430 440 450 460 470 480
 ACACGTAAGATTGAGTCTGGAACTGGGAACTACCGTGCCTCCCATCAGTTTCACNCCTGGCCCCCTGC
 490 500 510 520 530 540 550
 AGTGAGGCTGACTTGAAGAGATGGGAGGCCATCCGGGAGGCCAGCAGACTCAGGCACAAGAAAAGGCTG
 560 570 580 590 600 610 620
 ATGGTGGAGAGACTCTTTCAAAGATTTATGGTGAGAATGGGAGTAAGTCCATGAGTGATGTCAGCGCA
 630 640 650 660 670 680 690
 GAAGATGTTCAAAGCTTGGCTCAGCTGCGTTACGAGGAGATGCAGAAAATAAAATCACAATTAAGAA
 700 710 720 730 740 750
 CAAGATCAGAAATGGCAGGATGACCTTGCAAAATGGAAAGATCGTCGAAAAAGTTACACTTCAGATCTG
 760
 CAGAAG

FIG.28B




FIG.29




FIG.30




FIG.31

FIG. 32A FIG. 32C FIG. 32E FIG. 32G FIG. 32I FIG. 32K



FIG. 32B FIG. 32D FIG. 32F FIG. 32H FIG. 32J FIG. 32L




FIG.33




FIG. 34A



FIG. 34B




FIG. 34C



FIG. 34D

FIG. 34E






FIG.35A




FIG.35B




FIG.35C




FIG. 36B

FIG. 36A




FIG.37A




FIG.37B




FIG.37C






FIG.38








FIG. 41B

FIG. 41A




FIG.42




FIG. 43B

FIG. 43A

1 2
FIG. 43C




FIG.44A




FIG.44B




FIG.44C

NOVEL UBIQUITIN LIGASES AS THERAPEUTIC TARGETS

[0001] This application claims priority under 35 U.S.C. §119(e) to U.S. Application No. 60/098,355, filed Aug. 28, 1998, Application No. 60/118,568, filed Feb. 3, 1999, and Application No. 60/124,449 filed Mar. 15, 1999, the contents of which are incorporated herein by reference in their entirety.

1. INTRODUCTION

[0002] The present invention relates to the discovery, identification and characterization of nucleotide sequences that encode novel substrate-targeting subunits of ubiquitin ligases. The invention encompasses nucleic acid molecules comprising nucleotide sequences encoding novel substrate-targeting subunits of ubiquitin ligases: FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP11, FBP12, FBP13, FBP14, FBP15, FBP17, FBP18, FBP20, FBP21, FBP22, FBP23, AND FBP25, transgenic mice, knock-out mice, host cell expression systems and proteins encoded by the nucleotides of the present invention. The present invention relates to screening assays to identify potential therapeutic agents such as small molecules, compounds or derivatives and analogues of the novel ubiquitin ligases which modulate activity of the novel ubiquitin ligases for the treatment of proliferative and differentiative disorders, such as cancer, major opportunistic infections, immune disorders, certain cardiovascular diseases, and inflammatory disorders. The invention further encompasses therapeutic protocols and pharmaceutical compositions designed to target ubiquitin ligases and their substrates for the treatment of proliferative disorders.

2. BACKGROUND OF THE INVENTION

[0003] 2.1 Cell Cycle Regulatory Proteins

[0004] The eukaryotic cell cycle is regulated by a family of serine/threonine protein kinases called cyclin dependent kinases (Cdks) because their activity requires the association with regulatory subunits named Cyclins (Hunter & Pines, 1994, Cell 79:573). Cdks also associate with Cdk inhibitors (Cis) which mediate cell cycle arrest in response to various antiproliferative signals. So far, based on their sequence homology, two families of Ckis have been identified in mammalian cells: the Cip/Kip family, which includes p21, p27 and p57; and the Ink family, which includes p15, p16, p18, and p20 (Sherr & Roberts, 1999, Genes & Dev. 13: 1501).

[0005] 2.2 The Ubiquitin Pathway

[0006] Ubiquitin-mediated proteolysis is an important pathway of non-lysosomal protein degradation which controls the timed destruction of many cellular regulatory proteins including, p27, p53, p300, cyclins, E2F, STAT-1, c-Myc, c-Jun, EGF receptor, Ikb α , NFkB and β -catenin (reviewed in Pagano, 1997, FASEB J. 11:1067). Ubiquitin is an evolutionary highly conserved 76-amino acid polypeptide which is abundantly present in all eukaryotic cells. The ubiquitin pathway leads to the covalent attachment of poly-ubiquitin chain to target substrates which are then degraded by the multi-catalytic proteasome complex (see Pagano, supra, for a recent review). Many of the steps regulating protein ubiquitination are known. Initially the ubiquitin

activating enzyme (E1), forms a high energy thioester with ubiquitin which is, in turn, transferred to a reactive cysteine residue of one of many ubiquitin conjugating enzymes (Ubes or E2s). The final transfer of ubiquitin to an e-amino group of a reactive lysine residue in the target protein occurs in a reaction that may or may not require an ubiquitin ligase (E3) protein. The large number of ubiquitin ligases ensures the high level of substrate specificity.

[0007] 2.3 The Ubiquitin Pathway and the Regulation of the G1 Phase by F Box Proteins

[0008] Genetic and biochemical studies in several organisms have shown that the G1 phase of the cell cycle is regulated by the ubiquitin pathway. Proteolysis of cyclins, Ckis and other G1 regulatory proteins is controlled in yeast by the ubiquitin conjugating enzyme Ubc3 (also called Cdc34) and by an E3 ubiquitin ligase formed by three subunits: Cdc53, Skp1 and one of many F box proteins (reviewed in E. Patton et al., 1998, TIG. 14:6). The F box proteins (FBPs) are so called because they contain a motif, the F box, that was first identified in Cyclin F, and that is necessary for FBP interaction with Skp1 (Bai, et al., 1996, Cell 86:263). In addition, F box proteins also contain either WD-40 domains or Leucine-Rich Repeats (LRR) protein-protein interaction domains. Cdc53 (also called Cul A) and Skp1 appear to participate in the formation of at least three distinct E3, each containing a different F box protein. Because these ligases are similar protein modules composed of Skp1, Cul A, and an F box protein, they have been named SCF. The interaction of the ligase with its substrates occurs via the F box subunit. The three SCFs identified so far in *S. cerevisiae* are: SCF^{Cdc4} (which recruits the Ckis Sic1 and Far1, the replication factor Cdc6, and the transcriptional activator Gcn4, as substrates through the F box protein Cdc4), SCF^{Grr1} (which recruits the G1 cyclins Cln1 and Cln2 as substrates through the F box protein GRR1), and SCF^{Met30} (which recruits the G1 cyclin Cln3 as a substrate throughout the F box protein MET30; see Pagano and Patton, supra, for recent reviews).

[0009] The intracellular level of the human Cki p27 is mainly regulated by degradation and it is known that the ubiquitin system controls p27 degradation (Pagano et al., 1995, Science 269:682). Similarly, degradation of other G1 human regulatory proteins (Cyclin E, Cyclin D1, p21, E2F, β -catenin) is controlled by the ubiquitin-pathway (reviewed in M. Pagano, supra). Yet, the specific enzymes involved in the degradation of G1 regulatory proteins have not been identified.

[0010] A family of 6 genes (CUL1, 2, 3, 4a, 4b, and 5) homologous to *S. cerevisiae* cul A have been identified by searching the EST database (Kipreos, et al., 1996, Cell 85:829). Human Skp1 and the F box protein Skp2 (that contains five LRRs) were identified as two proteins associated in vivo with Cyclin A and thus designated as S-phase kinase-associated protein 1 and 2 (Zhang, et al., 1995, Cell 82:915).

[0011] 2.4 Deregulation of the Ubiquitin Pathway in Cancer and other Proliferative Disorders

[0012] Cancer develops when cells multiply too quickly. Cell proliferation is determined by the net balance of positive and negative signals. When positive signals overcome or when negative signals are absent, the cells multiply too quickly and cancer develops.

[0013] Ordinarily cells precisely control the amount of any given protein and eliminate the excess or any unwanted protein. To do so, the cell specifically tags the undesired protein with a long chain of molecules called ubiquitin. These molecules are then recognized and destroyed by a complex named proteasome. However, all this mechanism goes awry in tumors leading to the excessive accumulation of positive signals (oncogenic proteins), or resulting in the abnormal degradation of negative regulators (tumor suppressor proteins). Thus, without tumor suppressor proteins or in the presence of too much oncogenic proteins, cells multiply ceaselessly, forming tumors (reviewed by Ciechanover, 1998, *EMBO J.* 17: 7151; Spataro, 1998, *Br. J. Cancer* 77: 448). For example, abnormal ubiquitin-mediated degradation of the p53 tumor suppressor (reviewed by J. Brown and M. Pagano, 1997, *Biochim. Biophys. Acta* 1332: 1), the putative oncogene β -catenin (reviewed by Peifer, 1997, *Science* 275:1752) and the Cki p27 (reviewed in Ciechanover, supra; Spataro, supra; Lloyd, 1999, *Am. J. Pathol.* 154: 313) have been correlated with tumorigenesis, opening to the hypothesis that some genes encoding ubiquitinating enzymes may be mutated in tumors.

[0014] Initial evidence indicates that human F-box proteins play a role in the ubiquitination of G1 regulatory proteins as their homologs do in yeast (see below). Unchecked degradation of cell cycle regulatory proteins has been observed in certain tumors and it is possible that deregulated ubiquitin ligase play a role in the altered degradation of cell cycle regulators. A well understood example is that of Mdm2, a ubiquitin ligase whose overexpression induces low levels of its substrate, the tumor suppressor p53.

3. SUMMARY OF THE INVENTION

[0015] The present invention relates to novel F box proteins and therapeutic protocols and pharmaceutical compositions designed to target the novel F box proteins and their interactions with substrates for the treatment of proliferative and differentiative disorders. The present invention also relates to screening assays to identify substrates of the novel F box proteins and to identify agents which modulate or target the novel ubiquitin ligases and interactions with their substrates. The invention further relates to screening assays based on the identification of novel substrates of known F box proteins, such as the two novel substrates of the known F box protein Skp2, E2F and p27. The screening assays of the present invention may be used to identify potential therapeutic agents for the treatment of proliferative or differentiative disorders and other disorders that related to levels of expression or enzymatic activity of F box proteins.

[0016] The invention is based in part, on the Applicants' discovery, identification and characterization of nucleic acids comprising nucleotide sequences that encode novel ubiquitin ligases with F box motifs. These twenty-six novel substrate-targeting subunits of ubiquitin ligase complexes, FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25, described herein, were first identified based on their interaction with components of the ubiquitin ligase complex (FBPL, FBP2, FBP3a, FBP4, FBP5, FBP6 and FBP7) or by sequence comparison of these proteins with nucleotide sequences present in DNA databases (FBP3b, FBP8, FBP9, FBP10, FBP11, FBP12,

FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25). These novel substrate-targeting subunits of ubiquitin ligase complexes each contain an F box motif through which they interact with the other components of the ubiquitin ligase complex. In addition, some of these FBPs contain WD-40 domains and LRRs (which appear to be involved in their interaction with substrates), while other FBPs contain potential protein-protein interaction modules not yet identified in FBPs, such as leucine zippers, ring fingers, helix-loop-helix motifs, proline rich motifs and SH2 domains. The invention is also based, in part, on the Applicants' discovery and identification of FBP specific substrates p27 and β -catenin and on methods to identify novel FBP substrates. Some of the genes encoding the novel F box proteins were also mapped to chromosome sites frequently altered in breast, prostate and ovarian cancer, nasopharyngeal and small cell lung carcinomas, gastric hepatocarcinomas, Burkitt's lymphoma and parathyroid adenomas. Finally, the invention is also based, in part, on the Applicants' generation of transgenic mice expressing wild type or dominant negative versions of FBP proteins and on the generation of FBP knock-out mice.

[0017] The invention encompasses the following nucleotide sequences, host cells expressing such nucleotide sequences, and the expression products of such nucleotide sequences: (a) nucleotide sequences that encode mammalian FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP11, FBP12, FBP13, FBP14, FBP15, FBP17, FBP18, FBP20, FBP21, FBP22, FBP23, and FBP25, including the human nucleotides, and their gene products; (b) nucleotides that encode portions of the novel substrate-targeting subunits of ubiquitin ligase complexes, and the polypeptide products specified by such nucleotide sequences, including but not limited to F box motifs, the substrate binding domains; WD-40 domains; and leucine rich repeats, etc.; (c) nucleotides that encode mutants of the novel ubiquitin ligases in which all or part of the domain is deleted or altered, and the polypeptide products specified by such nucleotide sequences; (d) nucleotides that encode fusion proteins containing the novel ubiquitin ligases or one of its domains fused to another polypeptide.

[0018] The invention further encompasses agonists and antagonists of the novel substrate-targeting subunits of ubiquitin ligase complexes, including small molecules, large molecules, mutants that compete with native F box binding proteins, and antibodies as well as nucleotide sequences that can be used to inhibit ubiquitin ligase gene expression (e.g., antisense and ribozyme molecules, and gene regulatory or replacement constructs) or to enhance ubiquitin ligase gene expression (e.g., expression constructs that place the ubiquitin ligase gene under the control of a strong promoter system), and transgenic animals that express a ubiquitin ligase transgene or knock-outs that do not express the novel ubiquitin ligases.

[0019] Further, the present invention also relates to methods for the use of the genes and/or gene products of novel substrate-targeting subunits of ubiquitin ligase complexes for the identification of compounds which modulate, i.e., act as agonists or antagonists, of ubiquitin ligase activity. Such compounds can be used as agents to control proliferative or differentiative disorders, e.g. cancer. In particular, the present invention encompasses methods to inhibit the inter-

action between β -catenin and FBP1 or p27 and Skp2. In fact, agents able to block these interactions can be used to modulate cell proliferation and/or growth.

[0020] Still further, the invention encompasses screening methods to identify derivatives and analogues of the novel substrate-targeting subunits of ubiquitin ligase complexes which modulate the activity of the novel ligases as potential therapeutics for proliferative or differentiative disorders. The invention provides methods of screening for proteins that interact with novel components of the ubiquitin ligase complex, including FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBPL11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25 or derivatives, fragments or domains thereof, such as the F box motif. In accordance with the invention, the screening methods may utilize known assays to identify protein-protein interactions including phage display assays or the yeast two-hybrid assay system or variations thereof.

[0021] In addition, the present invention is directed to methods that utilize FBP gene sequences and/or FBP gene product sequences for the diagnostic evaluation, genetic testing and/or prognosis of an FBP-related disorder, such as a proliferative disorder. For example, the invention relates to methods for diagnosing FBP-related disorders, e.g., proliferative disorders, wherein such methods can comprise measuring FBP gene expression in a patient sample, or detecting an FBP mutation that correlates with the presence or development of such a disorder, in the genome of a mammal suspected of exhibiting such a disorder. In particular, the invention encompasses methods for determining if a subject (e.g., a human patient) is a risk for a disorder characterized by one or more of: (i) a mutation of an FBP gene encoding a protein represented in part A of FIGS. 3-28, or a homolog thereof; (ii) the mis-expression of an FBP gene; (iii) the mis-expression of an FBP protein.

[0022] The invention is illustrated by way of working examples which demonstrate the identification and characterization of the novel substrate-targeting subunits of ubiquitin ligase complexes. The working examples of the present invention further demonstrate the identification of the specific interaction of (i) FBP1 with β -catenin and (ii) the known FBP, Skp2, with the cell-cycle regulatory proteins E2F and p27. These interactions suggest that β -catenin is a specific substrate of FBP1, while E2F and p27 are substrates of Skp2. In fact, the working examples of the present invention further demonstrate that β -catenin is a specific substrate of FBP1, while p27 is substrates of Skp2. The identification of proteins interacting with the novel FBPs will be possible using the methods described herein or with a different approach.

[0023] 3.1 Definitions

[0024] As used herein, the term "F-box motif" refers to a stretch of approximately 40 amino acid that was identified as being necessary for the interaction of F-box containing proteins with Skp1. The consensus sequence of an F-box motif is described in Bai et al., 1996, Cell 86:263-274, incorporated herein by reference in its entirety.

[0025] As used herein the term "F-box protein" (FBP) refers to peptide, polypeptide or protein which contains an F-box motif.

[0026] Although, FBPs are substrate-targeting subunits of ubiquitin ligase complexes, as used herein the term "ubiquitin ligase" refers to a peptide, polypeptide or protein that contains an F-box motif and interacts with Skp1.

[0027] As used herein, the term "functionally equivalent to an FBP gene product" refers to a gene product that exhibits at least one of the biological activities of the endogenous FBP gene product. For example, a functionally equivalent FBP gene product is one that is capable of interacting with Skp1 so as to become associated with a ubiquitin ligase complex. Such a ubiquitin ligase complex may be capable of ubiquitinating a specific cell-cycle regulatory protein, such as a cyclin or cki protein.

[0028] As used herein, the term "to target" means to inhibit, block or prevent gene expression, enzymatic activity, or interaction with other cellular factors.

[0029] As used herein, the term "therapeutic agent" refers to any molecule, compound or treatment that alleviates or assists in the treatment of a proliferative disorder or related disorder.

[0030] As used herein, the terms "WD-40 domain", "Leucine Rich Repeat", "Leucine Zipper", "Ring finger", "Helix-loop-helix motif", "Proline rich motif", and "SH2 domain" refer to domains potentially involved in mediating protein-protein interactions. The "WD-40 domain" refers to a consensus sequence of forty amino acid repeats which is rich in tryptophan and aspartic acid residues and is commonly found in the beta subunits of trimeric G proteins (see Neer et al., 1994 Nature 371:297-300 and references therein, which are incorporated herein by reference in their entirety). An "LRR" or a "Leucine Rich Repeat" is a leucine rich sequence also known to be involved in mediating protein-protein interactions (see Kobe & Deisenhofer, 1994, Trends Biochem. Sci. 19:415-421 which are incorporated herein by reference in their entirety). A "leucine zipper" domain refers to a domain comprising a stretch of amino acids with a leucine residue in every seventh position which is present in a large family of transcription factors (see Landshultz et al., 1988, Science 240:1759-64; see also Sudol et al., 1996, Trends Biochem. 21:1-3, and Koch et al., 1991, Science 252:668-74).

4. BRIEF DESCRIPTION OF THE FIGURES

[0031] FIG. 1. Alignment of the conserved F-box motif amino acid residues in the human F-box proteins FBP1 (SEQ ID NO:15), FBP2 (SEQ ID NO:16), FBP3a (SEQ ID NO:17), FBP3b (SEQ ID NO:78), FBP4 (SEQ ID NO:18), FBP5 (SEQ ID NO:19), FBP6 (SEQ ID NO:20), FBP7 (SEQ ID NO:21), Skp2 (SEQ ID NO:22), FBP8 (SEQ ID NO:61), FBP9 (SEQ ID NO:62), FBP10 (SEQ ID NO:63), FBP11 (SEQ ID NO:64), FBP12 (SEQ ID NO:65), FBP13 (SEQ ID NO:79), FBP14 (SEQ ID NO:66), FBP15 (SEQ ID NO:67), FBP16 (SEQ ID NO:68), FBP17 (SEQ ID NO:69), FBP18 (SEQ ID NO:70), FBP19 (SEQ ID NO:71), FBP20 (SEQ ID NO:72), FBP21 (SEQ ID NO:73), FBP22 (SEQ ID NO:74), FBP23 (SEQ ID NO:75), FBP24 (SEQ ID NO:76), FBP25 (SEQ ID NO:77). Alignment of the F-boxes of a previously known FBP, Skp2, with the F-boxes of FBPs identified through a two-hybrid screen (designated by the pound symbol) or BLAST searches (designated by a cross) was performed using the Clustal W method (MacVector(tm)) followed by manual re-adjustment. Identical resi-

dues in at least 15 F-boxes are shaded in dark gray, while similar residues are shaded in light gray. One asterisk indicates the presence in the cDNA of a STOP codon followed by a polyA tail, while potential full length clones are designated with two asterisks. The asterisks on the bottom of the figure indicate the amino acid residues mutated in FBP3a (see FIG. 29).

[0032] **FIG. 2.** Schematic representation of FBPs. Putative protein-protein interaction domains in human FBPs are represented (see key-box for explanation). FBPs identified by a two-hybrid screen are designated by the pound symbol, FBPs identified through BLAST searches by a cross. The double slash indicates that the corresponding cDNAs are incomplete at the 5' end; the asterisks indicate the presence in the cDNA of a STOP codon followed by a polyA tail.

[0033] **FIG. 3 A-B.** A. Amino acid sequence of human F-box protein FBP1 (SEQ ID NO:2). B. Corresponding cDNA (SEQ ID NO:1).

[0034] **FIG. 4 A-B.** A. Amino acid sequence of human F-box protein FBP2 (SEQ ID NO:4). B. Corresponding cDNA (SEQ ID NO:3).

[0035] **FIG. 5 A-B.** A. Amino acid sequence of human F-box protein FBP3a (SEQ ID NO:6). B. Corresponding cDNA (SEQ ID NO:5).

[0036] **FIG. 6 A-B.** A. Amino acid sequence of human F-box protein FBP3b (SEQ ID NO:24). B. Corresponding cDNA (SEQ ID NO:23).

[0037] **FIG. 7 A-B.** A. Amino acid sequence of human F-box protein FBP4 (SEQ ID NO:8). B. Corresponding cDNA (SEQ ID NO:7).

[0038] **FIG. 8 A-B.** A. Amino acid sequence of human F-box protein FBP5 (SEQ ID NO:10). B. Corresponding cDNA (SEQ ID NO:9).

[0039] **FIG. 9 A-B.** A. Amino acid sequence of human F-box protein FBP6 (SEQ ID NO: 12). B. Corresponding cDNA (SEQ ID NO:11).

[0040] **FIG. 10 A-B.** A. Amino acid sequence of human F-box protein FBP7 (SEQ ID NO:14). B. Corresponding cDNA (SEQ ID NO:13).

[0041] **FIG. 11 A-B.** A. Amino acid sequence of human F-box protein FBP8 (SEQ ID NO:26). B. Corresponding cDNA (SEQ ID NO:25).

[0042] **FIG. 12 A-B.** A. Amino acid sequence of human F-box protein FBP9 (SEQ ID NO:28). B. Corresponding cDNA (SEQ ID NO:27).

[0043] **FIG. 13 A-B.** A. Amino acid sequence of human F-box protein FBP10 (SEQ ID NO:30). B. Corresponding cDNA (SEQ ID NO:29).

[0044] **FIG. 14 A-B.** A. Amino acid sequence of human F-box protein FBP11 (SEQ ID NO:32). B. Corresponding cDNA (SEQ ID NO:31).

[0045] **FIG. 15 A-B.** A. Amino acid sequence of human F-box protein FBP12 (SEQ ID NO:34). B. Corresponding cDNA (SEQ ID NO:33).

[0046] **FIG. 16 A-B.** A. Amino acid sequence of human F-box protein FBP13 (SEQ ID NO:36). B. Corresponding cDNA (SEQ ID NO:35).

[0047] **FIG. 17 A-B.** A. Amino acid sequence of human F-box protein FBP14 (SEQ ID NO:38). B. Corresponding cDNA (SEQ ID NO:37).

[0048] **FIG. 18 A-B.** A. Amino acid sequence of human F-box protein FBP15 (SEQ ID NO:40). B. Corresponding cDNA (SEQ ID NO:39).

[0049] **FIG. 19 A-B.** A. Amino acid sequence of human F-box protein FBP16 (SEQ ID NO:42). B. Corresponding cDNA (SEQ ID NO:41).

[0050] **FIG. 20 A-B.** A. Amino acid sequence of human F-box protein FBP17 (SEQ ID NO:44). B. Corresponding cDNA (SEQ ID NO:43).

[0051] **FIG. 21 A-B.** A. Amino acid sequence of human F-box protein FBP18 (SEQ ID NO:46). B. Corresponding cDNA (SEQ ID NO:45).

[0052] **FIG. 22 A-B.** A. Amino acid sequence of human F-box protein FBP19 (SEQ ID NO:48). B. Corresponding cDNA (SEQ ID NO:47).

[0053] **FIG. 23 A-B.** A. Amino acid sequence of human F-box protein FBP20 (SEQ ID NO:50). B. Corresponding cDNA (SEQ ID NO:49).

[0054] **FIG. 24 A-B.** A. Amino acid sequence of human F-box protein FBP21 (SEQ ID NO:52). B. Corresponding cDNA (SEQ ID NO:51).

[0055] **FIG. 25 A-B.** A. Amino acid sequence of human F-box protein FBP22 (SEQ ID NO:54). B. Corresponding cDNA (SEQ ID NO:53).

[0056] **FIG. 26 A-B.** A. Amino acid sequence of human F-box protein FBP23 (SEQ ID NO:56). B. Corresponding cDNA (SEQ ID NO:55).

[0057] **FIG. 27 A-B.** A. Amino acid sequence of human F-box protein FBP24 (SEQ ID NO:58). B. Corresponding cDNA (SEQ ID NO:57).

[0058] **FIG. 28A-B.** A. Amino acid sequence of human F-box protein FBP25 (SEQ ID NO:60). B. Corresponding cDNA (SEQ ID NO:59).

[0059] **FIG. 29.** FBPs interact specifically with Skp1 through their F-box. The cDNAs of FBPs (wild type and mutants) were transcribed and translated in vitro (IVT) in the presence of 35S-methionine. Similar amounts of IVT proteins (indicated at the top of each lane) were subjected to a histidine-tagged pull-down assay using Nickel-agarose beads to which either His-tagged-Skp1 (lanes 1, 3, 4, 6-10, 12, 15, 17, 19 and 21), His-tagged-Elongin C (lanes 2, 5, 11, 14, 16, 18, 19 and 22), or His-tagged p27 (lane 12) were pre-bound. Bound IVT proteins were analyzed by SDS-PAGE and autoradiography. The arrows on the left side of the panels point to the indicated FBPs. The apparent molecular weights of the protein standards are indicated on the right side of the panels.

[0060] **FIG. 30.** FBP1, FBP2, FBP3a, FBP4 and FBP7 form novel SCFs with endogenous Skp1 and Cull in vivo. HeLa cells were transfected with mammalian expression plasmids encoding Flag-tagged versions of FBPI (lane 1), (ΔF)FBP1 (lane 2), FBP4 (lane 3), FBP7 (lane 5), FBP2 (lane 7), (ΔF)FBP2 (lane 8), FBP3a (lane 9), (ΔF)FBP3a (lane 10), or with an empty vector (lanes 4 and 6). Cells were lysed and extracts were subjected to immunoprecipitation

with a rabbit anti-Flag antibody (lanes 1-8). Immunoprecipitates were then immunoblotted with a mouse anti-Cull monoclonal antibody, a rabbit anti-Skp1 polyclonal antibody or a rabbit anti-Cul2 polyclonal antibody, as indicated. The last lane contains 25 μ g of extracts from non-transfected HeLa cells; lane 9 contains recombinant Cul1, Skp1, or Cul2 proteins used as markers. The slower migrating bands detected with the antibodies to Cul1 and Cul2 are likely generated by the covalent attachment of a ubiquitin-like molecule to these two cullins, as already described for the yeast cullin Cdc53 and mammalian Cul4a.

[0061] FIG. 31. FBP1, FBP2, FBP3a, FBP4 and FBP7 associate with a ubiquitin ligase activity. HeLa cells were transfected with mammalian expression plasmids encoding human Skp1, Cul1 and Flag-tagged versions of FBP1 (lane 3), (Δ F)FBP1 (lane 4), FBP2 (lanes 2 and 5), (Δ F)FBP2 (lane 6), FBP7 (lane 7), FBP3a (lanes 8 and 13), (Δ F)FBP3a (lane 9), a non relevant Flag-tagged protein (Irf3, lane 10), FBP4 (lanes 11 and 12) or with an empty vector (lane 1). Cells were lysed and extracts were subjected to immunoprecipitation with a rabbit anti-Flag antibody. Immunoprecipitates were incubated in the presence of purified recombinant E1 and Ubc4 (lanes 1-11) or Ubc2 (lanes 12 and 13) and a reaction mix containing biotinylated ubiquitin. Reaction in lane 2 contained also NEM. Ubiquitinated proteins were visualized by blotting with HRP-streptavidin. The racket on the left side of the panels marks a smear of ubiquitinated proteins produced in the reaction, the asterisk indicates ubiquitin conjugated with E1 that were resistant to oiling.

[0062] FIG. 32. Subcellular localization of FBPs. HeLa cells were transfected with mammalian expression plasmids encoding Flag-tagged versions of FBP1 (a-b), FBP2 (c-d), FBP3a (e-f), FBP4 (g-h), (Δ F)FBP2 (i-j), or (Δ F)FBP3a (k-l). After 24 hours, cells were subjected to immunofluorescence with a rabbit anti-Flag antibody (a, c, e, g, i, k) to stain FBPs and bisbenzimidazole (b, d, f, h, j, l) to stain nuclei.

[0063] FIG. 33. Abundance of FBP transcripts in human tissues. Membranes containing electrophoretically fractionated poly(A)⁺ mRNA from different human tissues were hybridized with specific probes prepared from FBP1, FBP2, FBP3a, FBP4, SKP2, and β -ACTIN cDNAs. The arrows on the left side of the figure point to the major transcripts as described in the text.

[0064] FIG. 34 A-E. FISH localization of FBP genes. Purified phage DNA containing a genomic probe was labeled with digoxigenin dUTP and detected with Cy3-conjugated antibodies. The signals corresponding to the locus of the genomic probe (red) are seen against the DAPI-Actinomycin D stained normal human chromosomes (blue-white). Panel A shows localization of FBP1 to 10q24, B shows localization of FBP2 to 9q34, C shows localization of FBP3a to 13q22, D shows localization of FBP4 to 5p12, and E shows localization of FBP5 to 6q25-26. Arrows point to FBP-specific FISH signals.

[0065] FIG. 35A-C. FBP1 associates with β -catenin. A. Extracts from baculovirus-infected insect cells expressing either β -catenin alone (lane 1) or in combination with Flag-tagged FBP1 (lane 2) were immunoprecipitated (IP) with a rabbit anti-Flag antibody (r α -Flag), followed by immunoblotting with anti-Flag (m α -Flag) and anti- β -catenin mouse antibodies, as indicated. Lanes 3 and 4 contain 25

μ g of extracts from infected insect cells immunoblotted with the same antibodies. B. Extracts from baculovirus-infected insect cells expressing cyclin D1, Flag-FBP1 in the absence (lanes 1-3) or in the presence of Skp1 (lanes 4-6) were immunoprecipitated with normal rabbit IgG (r-IgG, lanes 1 and 4), rabbit anti-Flag antibody (r α -Flag, lanes 2 and 5), or rabbit anti-cyclin D1 antibody (r α -D1, lanes 3 and 6). Immunoprecipitates were then immunoblotted with anti-Flag (m α -Flag) and cyclin D1 (m α -D1) mouse antibodies, as indicated. The last lane contains 25 μ g of a representative extract from infected insect cells immunoblotted with the same antibodies. C. 293 cells were transfected with mammalian expression plasmids encoding HA-tagged β -catenin alone or in combination with either Flag-tagged FBP1 or Flag-tagged (Δ F)FBP1. Cells were lysed and extracts were subjected to immunoprecipitation with a rabbit anti-Flag antibody (r α -Flag, lanes 4-6) and immunoblotted with rat anti-HA (α -HA) and mouse anti-Flag (m α -Flag) antibodies, as indicated. The first three lanes contain 25 μ g of extracts from transfected 293 cells immunoblotted with the same antibodies. Transfecting high levels of β -catenin expression vector, the associations of β -catenin with FBP1 and (Δ F)FBP1 could be determined independently of β -catenin levels.

[0066] FIG. 36 A-B. Stabilization of β -catenin by a dominant negative (Δ F)FBP1 mutant. A. Human 293 cells were transfected with mammalian expression plasmids encoding HA-tagged β -catenin alone or in combination with either Flag-tagged (Δ F)FBP1 or Flag-tagged (Δ F)FBP2. Cells were lysed and extracts were subjected to immunoblotting with rat anti-HA and rabbit anti-Flag (r α -Flag) antibody, as indicated. B. Pulse chase analysis of β -catenin turnover rate. HA-tagged β -catenin in combination with either an empty vector, FBP1, or (Δ F)FBP1 was co-transfected in 293 cells. 24 hours later cells were labeled with ³⁵S-methionine for 30 minutes and chased with medium for the indicated times. Extracts were then subjected to immunoprecipitation with a rat anti-HA antibody.

[0067] FIG. 37A-C. Binding of phosphorylated p27 to Skp2. A. A panel of in vitro translated [³⁵S]FBPs were used in binding reactions with beads coupled to the phosphopeptide NAGSVEQT*PKKPGRLRRRQT, corresponding to the carboxy terminus of the human p27 with a phosphothreonine at position 187 (T*). Beads were washed with RIPA buffer and bound proteins were eluted and subjected to electrophoresis and autoradiography (Upper Panel). Bottom Panel: 10% of the in vitro translated [³⁵S]FBP inputs. B. HeLa cell extracts were incubated with beads coupled to the phospho-p27 peptide (lane 2), an identical except unphosphorylated p27 peptide (lane 1) or the control phosphopeptide AEIGVGAY*GTVYKARDPHS, corresponding to an amino terminal peptide of human Cdk4 with a phosphotyrosine at position 17 (Y*) (lane 3). Beads were washed with RIPA buffer and bound proteins were immunoblotted with antibodies to the proteins indicated on the left of each panel. A portion of the HeLa extract (25 μ g) was used as a control (lane 4). The slower migrating band in Cull is likely generated by the covalent attachment of a ubiquitin-like molecule, as already described for other cullins 48. C. One μ l of in vitro translated [³⁵S] wild type p27 (WT, lanes 1-4) or p27(T187A) mutant (T187A, lanes 5-6) were incubated for 30 minutes at 30°C in 10 μ l of kinase buffer. Where indicated, ~2.5 pmole of recombinant purified cyclin E/Cdk2 or ~1 pmole Skp2 (in Skp1/Skp2 complex) were

added. Samples were then incubated with 6 μ l of Protein-A beads to which antibodies to Skp2 had been covalently linked. Beads were washed with RIPA buffer and bound proteins subjected to electrophoresis and autoradiography. Lanes 1-6: Skp2-bound proteins; Lanes 7 and 8: 7.5% of the in vitro translated [35S] protein inputs.

[0068] FIG. 38. In vivo binding of Skp2 to p27. Extracts from HeLa cells (lanes 1-2 and 5-6) or 1MR90 fibroblasts (lanes 9-10) were immunoprecipitated with different affinity purified (AP) antibodies to Skp2 or with purified control IgG fractions. Lane 1: extract immunoprecipitated with a goat IgG (G-IgG); lane 2: with an AP goat antibody to an N-terminal Skp2 peptide (G-a-Skp2); lanes 5 and 9: with a rabbit IgG (R-IgG); lanes 6 and 10: with an AP rabbit antibody to Skp2 (R-c-Skp2). Immunoprecipitates were immunoblotted with antibodies to the proteins indicated on the left of each panel. Lanes 1-4 in the bottom panel were immunoblotted with a phospho-site p27 specific antibody. Lanes 3, 7, and 11 contain 25 μ g of cell extracts; Lanes 4, 8, and 12 contain the relevant recombinant proteins used as markers. The altered migration of some markers is due to the presence of tags on the recombinant proteins.

[0069] FIG. 39 A-B. Skp2 and cyclin E/Cdk2 complex are rate-limiting for p27 ubiquitination in G1 extracts. A. In vitro ubiquitin ligation (lanes 1-12 and 17-20) and degradation (lanes 13-16) of p27 were carried out with extracts from asynchronously growing (Asyn. ext., lanes 2-3) or G1-arrested (G1 ext., lanes 4-20) HeLa cells. Lane 1 contains no extract. Recombinant purified proteins were supplemented as indicated. Reactions were performed using wild-type p27 (lanes 1-18) or p27(T187A) mutant (T187A, lanes 19-20). Lanes 1-8, 9-12, and 17-20 are from three separate experiments. The bracket on the left side of the panels marks a ladder of bands >27,000 corresponding to polyubiquitinated p27. The asterisk indicates a non-specific band present in most samples. B. Immunoblot analysis of levels of Skp2 and p27 in extracts from asynchronous (lane 1) or G1-arrested (lane 2) HeLa cells.

[0070] FIG. 40 A-C. Skp2 is required for p27-ubiquitin ligation activity. A. Immunodepletion. Extracts from asynchronous HeLa cells were untreated (lane 2) or immunodepleted with pre-immune serum (lane 3), anti-Skp2 antibody pre-incubated with 2 μ g of purified GST (lane 4), or anti-Skp2 antibody pre-incubated with 2 μ g of purified GST-Skp2 (lane 5). Lane 1 contains no extract. Samples (30 μ g of protein) were assayed for p27 ubiquitination in the presence of cyclin E/Cdk2. The bracket on the left side of the panels marks a ladder of bands >27,000 corresponding to polyubiquitinated p27. The asterisk indicates a non-specific band present in all samples. B. Reconstitution. The restoration of p27 ubiquitination activity in Skp2-immunodepleted extracts was tested by the addition of the indicated purified proteins. All samples contained 30 μ g of Skp2-depleted extract (Skp2-depl. ext.) and cyclin E/Cdk2. C. Immunopurification. Extracts from asynchronous HeLa cells were immunoprecipitated with a rabbit anti-Skp2 antibody (lanes 3 and 5) or pre-immune serum (PI, lanes 2 and 4). Total extract (lane 1) and immuno-beads (lanes 2-5) were added with p27, recombinant purified cyclin E/Cdk2 and ubiquitination reaction mix. Samples in lanes 4 and 5 were supplemented with recombinant purified E1 and Ubc3. All samples were then assayed for p27 ubiquitination.

[0071] FIG. 41 A-B. In vivo role of Skp2 in p27 degradation. A. Stabilization of p27 by a dominant negative (Δ F)Skp2 mutant in vivo. NIH-3T3 cells were transfected with mammalian expression vectors encoding human p27 alone (lane 2), p27 in combination with either (Δ F)Skp2 (lane 3), or (Δ F)FBP1 (lane 4). Lane 1: untransfected cells. Cells were lysed and extracts were subjected to immunoblotting with antibodies to p27, Skp2 or Flag [to detect Flag-tagged (Δ F)FBP1]. Exogenous human p27 protein migrates more slowly than the endogenous murine p27. B. Pulse chase analysis of p27 turnover rate. Human p27 in combination with either an empty vector, or (Δ F)Skp2 was transfected in NIH-3T3 cells. Twenty-four hours later, cells were labeled with [35S]-methionine for 20 minutes and chased with medium for the indicated times. Extracts were then subjected to immunoprecipitation with a mouse anti-p27 antibody.

[0072] FIG. 42. Stabilization of cellular p27 by antisense oligonucleotides targeting SKP2 mRNA. HeLa cells were treated for 16-18 hours with two different anti-sense oligodeoxynucleotides (AS) targeting two different regions of SKP2 mRNA. Lanes 2, 6, 12 and 16: AS targeting the N-terminal SKP2 region (NT); Lanes 4 and 8: AS targeting the C-terminal SKP2 region (CT); Lanes 1, 3, 5, 7, 11 and 15: control oligodeoxynucleotides pairs (Ctrl). Lanes 1-4, and 5-8 are from two separate experiments. Lanes 11-12 and 15-16: HeLa cells were blocked in G1/S with either Hydroxyurea or Aphidicolin treatment respectively, for 24 hours. Cells were then transfected with oligodeoxynucleotides, lysed after 12 hours (before cells had re-entered G1) and immunoblotted with antibodies to Skp2 (top panels) and p27 (bottom panels). Lanes 9 and 13: Untransfected HeLa cells; Lanes 10 and 14: Untransfected HeLa cells treated with drugs as transfected cells.

[0073] FIG. 43 A-C. Timing of Skp2 action in the process of p27 degradation. A. IMR90 fibroblasts were synchronized in G0/G1 by serum deprivation, reactivated with serum, and sampled at the indicated intervals. Protein extracts were analyzed by immunoblot with the antibodies to the indicated proteins. The Skp2 doublet was likely generated by phosphorylation since was consistently observed using a 12.5% gel only when cell lysis was performed in the presence of okadaic acid. B. HeLa cells blocked in mitosis with nocodazole were shaken off, released in fresh medium and sampled at the indicated intervals. Protein extracts were analyzed by immunoblotting with the antibodies to the indicated proteins. C. Extracts from G1 (3 hours after release from nocodazole block) (lane 1) and S-phase (12 hours after release from the nocodazole block) (lane 2) HeLa cells were either immunoprecipitated with an anti-p27 antibody (top two panels) or with an anti-Skp2 antibody (bottom three panels) and then immunoblotted with the antibodies to the indicated proteins.

[0074] FIG. 44 A-C. Western blot analysis of Skp2/E2F interaction assay. Details of the Western Blot experiments are given in the Example in Section 9.

5. DETAILED DESCRIPTION OF THE INVENTION

[0075] The present invention relates to novel F-box proteins and to novel substrates of F-box proteins. The present invention relates to screening assays designed to identify

substrates of the novel F-box proteins and to identify small molecules and compounds which modulate the interaction and/or activity of the F-box proteins and their substrates.

[0076] The present invention relates to screening assays to identify substrates of the novel F-box proteins and to identify potential therapeutic agents. The present invention further relates to screening assays based on the identification of novel substrates of both novel and known F-box proteins. The screening assays of the present invention may be used to identify potential therapeutic agents which may be used in protocols and as pharmaceutical compositions designed to target the novel ubiquitin ligases and interactions with their substrates for the treatment of proliferative disorders. In one particular embodiment the present invention relates to screening assays and potential therapeutic agents which target the interaction of FBP with novel substrates β -catenin, p27 and E2F as identified by Applicants.

[0077] The invention further encompasses the use of nucleotides encoding the novel F-box proteins, proteins and peptides, as well as antibodies to the novel ubiquitin ligases (which can, for example, act as agonists or antagonists), antagonists that inhibit ubiquitin ligase activity or expression, or agonists that activate ubiquitin ligase activity or increase its expression. In addition, nucleotides encoding the novel ubiquitin ligases and proteins are useful for the identification of compounds which regulate or mimic their activity and therefore are potentially effective in the treatment of cancer and tumorigenesis.

[0078] In particular, the invention described in the subsections below encompasses FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25 polypeptides or peptides corresponding to functional domains of the novel ubiquitin ligases (e.g., the F-box motif, the substrate binding domain, and leucine-rich repeats), mutated, truncated or deleted (e.g. with one or more functional domains or portions thereof deleted), ubiquitin ligase fusion proteins, nucleotide sequences encoding such products, and host cell expression systems that can produce such ubiquitin ligase products.

[0079] The present invention provides methods of screening for peptides and proteins that interact with novel components of the ubiquitin ligase complex, including FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25 or derivatives, fragments or analogs thereof. Preferably, the method of screening is a yeast two-hybrid assay system or a variation thereof, as further described below. Derivatives (e.g., fragments) and analogs of a protein can be assayed for binding to a binding partner by any method known in the art, for example, the modified yeast two-hybrid assay system described below, immunoprecipitation with an antibody that binds to the protein in a complex followed by analysis by size fractionation of the immunoprecipitated proteins (e.g., by denaturing or non-denaturing polyacrylamide gel electrophoresis), Western analysis, non-denaturing gel electrophoresis, etc.

[0080] The present invention relates to screening assays to identify agents which modulate the activity of the novel ubiquitin ligases. The invention encompasses both in vivo

and in vitro assays to screen small molecules, compounds, recombinant proteins, peptides, nucleic acids, antibodies etc. which modulate the activity of the novel ubiquitin ligases and thus, identify potential therapeutic agents for the treatment of proliferative or differentiative disorders. In one embodiment, the present invention provides methods of screening for proteins that interact with the novel ubiquitin ligases.

[0081] The invention also encompasses antibodies and anti-idiotypic antibodies, antagonists and agonists, as well as compounds or nucleotide constructs that inhibit expression of the ubiquitin ligase gene (transcription factor inhibitors, antisense and ribozyme molecules, or gene or regulatory sequence replacement constructs), or promote expression of the ubiquitin ligase (e.g., expression constructs in which ubiquitin ligase coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, etc.). The invention also relates to host cells and animals genetically engineered to express the human (or mutants thereof) or to inhibit or "knock-out" expression of the animal's endogenous ubiquitin ligase.

[0082] Finally, the ubiquitin ligase protein products and fusion protein products, (i.e., fusions of the proteins or a domain of the protein, e.g., F-box motif), antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate the ubiquitization pathway can be used for therapy of proliferative or differentiative diseases. Thus, the invention also encompasses pharmaceutical formulations and methods for treating cancer and tumorigenesis.

[0083] Various aspects of the invention are described in greater detail in the subsections below.

[0084] 5.1 FBP Genes

[0085] The invention provides nucleic acid molecules comprising seven novel nucleotide sequences, and fragments thereof, FBP1, FBP2, FBP3a, FBP4, FBP5, FBP6, and FBP7, nucleic acids which are novel genes identified by the interaction of their gene products with Skp1, a component of the ubiquitin ligase complex. The invention further provides fourteen novel nucleic acid molecules comprising the nucleotide sequences of FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP11, FBP12, FBP13, FBP14, FBP15, FBP17, FBP18, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25, which Nucleic acid sequences of the identified FBP genes are described herein.

[0086] As used herein, "an FBP gene" refers to:

[0087] (a) a nucleic acid molecule containing the DNA sequences of FBP1, shown in **FIG. 3** (SEQ ID NO:1), the DNA sequences of FBP2, shown in **FIG. 4** (SEQ ID NO:3), the DNA sequences of FBP3a, shown in **FIG. 5** (SEQ ID NO:5), the DNA sequences of FBP3b, shown in **FIG. 6** (SEQ ID NO:23), the DNA sequences of FBP4, shown in **FIG. 7** (SEQ ID NO:7), the DNA sequences of FBP5, shown in **FIG. 8** (SEQ ID NO:9), the DNA sequences of FBP6, shown in **FIG. 9** (SEQ ID NO:11), the DNA sequences of FBP7, shown in **FIG. 10** (SEQ ID NO: 13), the DNA sequences of FBP8, shown in **FIG. 11** (SEQ ID NO:25), the DNA sequences of FBP9, shown in **FIG. 12** (SEQ ID NO:27), the DNA sequences of FBP10, shown in

FIG. 13 (SEQ ID NO:29), the DNA sequences of FBP11, shown in **FIG. 14** (SEQ ID NO:31), the DNA sequences of FBP12, shown in **FIG. 15** (SEQ ID NO:33), the DNA sequences of FBP13, shown in **FIG. 16** (SEQ ID NO:35), the DNA sequences of FBP14, shown in **FIG. 17** (SEQ ID NO:37), the DNA sequences of FBP15, shown in **FIG. 18** (SEQ ID NO:39), the DNA sequences of FBP16, shown in **FIG. 19** (SEQ ID NO:41), the DNA sequences of FBP17, shown in **FIG. 20** (SEQ ID NO:43), the DNA sequences of FBP18, shown in **FIG. 21** (SEQ ID NO:45), the DNA sequences of FBP19, shown in **FIG. 22** (SEQ ID NO:47), the DNA sequences of FBP20, shown in **FIG. 23** (SEQ ID NO:49), the DNA sequences of FBP21, shown in **FIG. 24** (SEQ ID NO:51), the DNA sequences of FBP22, shown in **FIG. 25** (SEQ ID NO:53), the DNA sequences of FBP23, shown in **FIG. 26** (SEQ ID NO:55), the DNA sequences of FBP24, shown in **FIG. 27** (SEQ ID NO:57), the DNA sequences of FBP25, shown in **FIG. 28** (SEQ ID NO:59).

[0088] (b) any DNA sequence that encodes a polypeptide containing: the amino acid sequence of FBP1 shown in **FIG. 3A** (SEQ ID NO:2), the amino acid sequence of FBP2, shown in **FIG. 4A** (SEQ ID NO:4), the amino acid sequence of FBP3a shown in **FIG. 5A** (SEQ ID NO:6), the amino acid sequence of FBP3b shown in **FIG. 6A** (SEQ ID NO:24), the amino acid sequence of FBP4 shown in **FIG. 7A** (SEQ ID NO:8), the amino acid sequence of FBP5 shown in **FIG. 8A** (SEQ ID NO:10), or the amino acid sequence of FBP6 shown in **FIG. 9A** (SEQ ID NO:12), the amino acid sequences of FBP7, shown in **FIG. 10** (SEQ ID NO:14), the amino acid sequences of FBP8, shown in **FIG. 11** (SEQ ID NO:26), the amino acid sequences of FBP9, shown in **FIG. 12** (SEQ ID NO:28), the amino acid sequences of FBP10, shown in **FIG. 13** (SEQ ID NO:30), the amino acid sequences of FBP11, shown in **FIG. 14** (SEQ ID NO:32), the amino acid sequences of FBP12, shown in **FIG. 15** (SEQ ID NO:34), the amino acid sequences of FBP13, shown in **FIG. 16** (SEQ ID NO:36), the amino acid sequences of FBP14, shown in **FIG. 17** (SEQ ID NO:38), the amino acid sequences of FBP15, shown in **FIG. 18** (SEQ ID NO:40), the amino acid sequences of FBP16, shown in **FIG. 19** (SEQ ID NO:42), the amino acid sequences of FBP17, shown in **FIG. 20** (SEQ ID NO:44), the amino acid sequences of FBP18, shown in **FIG. 21** (SEQ ID NO:46), the amino acid sequences of FBP19, shown in **FIG. 22** (SEQ ID NO:48), the amino acid sequences of FBP20, shown in **FIG. 23** (SEQ ID NO:50), the amino acid sequences of FBP21, shown in **FIG. 24** (SEQ ID NO:52), the amino acid sequences of FBP22, shown in **FIG. 25** (SEQ ID NO:54), the amino acid sequences of FBP23, shown in **FIG. 26** (SEQ ID NO:56), the amino acid sequences of FBP24, shown in **FIG. 27** (SEQ ID NO:58), the amino acid sequences of FBP25, shown in **FIG. 28** (SEQ ID NO:60).

[0089] (c) any DNA sequence that hybridizes to the complement of the DNA sequences that encode any of the amino acid sequences of (SEQ ID NO: 2, 4, 6,

8, 10, 12 or 14) or **FIG. 15** under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaOH, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65 C, and washing in 0.1×SSC/0.1% SDS at 68 C (Ausubel F. M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. 1, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at p. 2.10.3); and/or

[0090] (d) any DNA sequence that hybridizes to the complement of the DNA sequences that encode any of the amino acid sequences in (SEQ ID NO: 2, 4, 6, 8, 10, 12 or 14) or **FIG. 15**, under less stringent conditions, such as moderately stringent conditions, e.g., washing in 0.2×SSC/0.1% SDS at 42 C (Ausubel et al., 1989, supra), and encodes a gene product functionally equivalent to an FBP gene product.

[0091] It is understood that the FBP gene sequences of the present invention do not encompass the previously described genes encoding other mammalian F-box proteins, Skp2, Elongin A, Cyclin F, mouse Md6, (see Pagano, 1997, supra; Zhang et al., 1995, supra; Bai et al., 1996, supra; SkoNvyra et al., 1997, supra). It is further understood that the nucleic acid molecules of the invention do not include nucleic acid molecules that consist solely of the nucleotide sequence in GenBank Accession Nos. AC002428, AI457595, AI105408, H66467, T47217, H38755, THC274684, AI750732, AA976979, AI571815, T57296, Z44228, Z45230, N42405, AA018063, AI751015, AI400663, T74432, AA402415, AI826000, AM590138, AF174602, Z45775, AF174599, THC288870, AI017603, AF174598, THC260994, AI475671, AA768343, AF174595, THC240016, N70417, T10511, AF174603, EST04915, AA147429, AI192344, AF174594, AI147207, AI279712, AA593015, AA644633, AA335703, N26196, AF174604, AF053356, AF174606, AA836036, AA853045, AI479142, AA772788, AA039454, AA397652, AA463756, AA007384, AA749085, AI640599, THC253263, AB020647, THC295423, AA434109, AA370939, AA215393, THC271423, AF052097, THC288182, AL049953, CAB37981, AL022395, AL031178, THC 197682, and THC205131.

[0092] FBP sequences of the present invention are derived from a eukaryotic genome, preferably a mammalian genome, and more preferably a human or murine genome. Thus, the nucleotide sequences of the present invention do not encompass those derived from yeast genomes. In a specific embodiment, the nucleotides of the present invention encompass any DNA sequence derived from a mammalian genome which hybridizes under highly stringent conditions to SEQ ID NO: 1, 3, 5, 7, 9, 11 or 13, or to DNA sequence shown in **FIG. 14**, encodes a gene product which contains an F-box motif and binds to Skp1. In a specific embodiment, the nucleotides of the present invention encompass any DNA sequence derived from a mammalian genome which hybridize under highly stringent conditions to SEQ ID NO: 1, 3, 5, 7, 9, 11 or 13 encodes a gene product which contains an F-box motif and another domain selected from the group comprising WD-40, leucine rich region, leucine zipper motif, or other protein-protein interaction domain, and binds to Skp-1 and is at least 300 or 400 nucleotides in length.

[0093] FBP sequences can include, for example, either eukaryotic genomic DNA (cDNA) or cDNA sequences.

When referring to a nucleic acid which encodes a given amino acid sequence; therefore, it is to be understood that the nucleic acid need not only be a cDNA molecule, but can also, for example, refer to a cDNA sequence from which an mRNA species is transcribed that is processed to encode the given amino acid sequence.

[0094] As used herein, an FBP gene may also refer to degenerate variants of DNA sequences (a) through (d).

[0095] The invention also includes nucleic acid molecules derived from mammalian nucleic acids, preferably DNA molecules, that hybridize to, and are therefore the complements of, the DNA sequences (a) through (d), in the preceding paragraph. Such hybridization conditions may be highly stringent or less highly stringent, as described above. In instances wherein the nucleic acid molecules are deoxyoligonucleotides ("oligos"), highly stringent conditions may refer, e.g., to washing in 6×SSC/0.05% sodium pyrophosphate at 37 C (for 14-base oligos), 48 C (for 17-base oligos), 55 C (for 20-base oligos), and 60 C (for 23-base oligos). These nucleic acid molecules may encode or act as FBP gene antisense molecules, useful, for example, in FBP gene regulation (for and/or as antisense primers in amplification reactions of FBP gene nucleic acid sequences). With respect to FBP gene regulation, such techniques can be used to regulate, for example, an FBP-regulated pathway, in order to block cell proliferation associated with cancer. Further, such sequences may be used as part of ribozyme and/or triple helix sequences, also useful for FBP gene regulation. Still further, such molecules may be used as components of diagnostic methods whereby, for example, the presence of a particular FBP allele responsible for causing an FBP-related disorder, e.g., proliferative or differentiative disorders such as tumorigenesis or cancer, may be detected.

[0096] The invention also encompasses:

[0097] (a) DNA vectors that contain any of the foregoing FBP coding sequences and/or their complements (i.e., antisense);

[0098] (b) DNA expression vectors that contain any of the foregoing FBP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences; and

[0099] (c) genetically engineered host cells that contain any of the foregoing FBP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell.

[0100] As used herein, regulatory elements include but are not limited to inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include but are not limited to the cytomegalovirus hCMV immediate early gene, the early or late promoters of SV40 adenovirus, the lac system, the trp system, the TAC system, the TRC system, the major operator and promoter regions of phage λ , the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase, the promoters of acid phosphatase, and the promoters of the yeast-mating factors.

[0101] The invention further includes fragments of any of the DNA sequences disclosed herein.

[0102] In one embodiment, the FBP gene sequences of the invention are mammalian gene sequences, with human sequences being preferred.

[0103] In yet another embodiment, the FBP gene sequences of the invention are gene sequences encoding FBP gene products containing polypeptide portions corresponding to (that is, polypeptide portions exhibiting amino acid sequence similarity to) the amino acid sequence depicted in FIGS. 2, 4-9 or 15, wherein the corresponding portion exhibits greater than about 50% amino acid identity with the depicted sequence, averaged across the FBP gene product's entire length.

[0104] In specific embodiments, F-box encoding nucleic acids comprise the cDNA sequences of SEQ ID NOs: 1, 3, 5, 23, 7, 9, 11, 13, 15, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, or 59, nucleotide sequence of FIGS. 3B, 4B, 5B, 6B, 7B, 8B, 9B, 10B, 11B, 12B, 13B, 14B, 15B, 16B, 17B, 18B, 19B, 20B, 21B, 22B, 23B, 24B, 25B, 26B, 27B, or 28B, respectively, or the coding regions thereof, or nucleic acids encoding an F-box protein (e.g., a protein having the sequence of SEQ ID NOs: 2, 4, 6, 24, 8, 10, 12, 14, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 68, or 60, or as shown in FIGS. 3A, 4A, 5A, 6A, 7A, 8A, 9A, 10A, 11A, 12A, 13A, 14A, 15A, 16A, 17A, 18A, 19A, 20A, 21A, 22A, 23A, 24A, 25A, 26A, 27A, or 28A, respectively).

[0105] The invention further provides nucleotide fragments of nucleotide sequences encoding FBP1, FBP2, FBP3a, FBP4, FBP5, FBP6, or FBP7 (SEQ ID NOs: 1, 3, 5, 7, 9, 11 and 13, respectively) of the invention. Such fragments consist of at least 8 nucleotides (i.e., a hybridizable portion) of an FBP gene sequence; in other embodiments, the nucleic acids consist of at least 25 (continuous) nucleotides, 50 nucleotides, 100 nucleotides, 150 nucleotides, or 200 nucleotides of an F-box sequence, or a full-length F-box coding sequence. In another embodiment, the nucleic acids are smaller than 35, 200 or 500 nucleotides in length. Nucleic acids can be single or double stranded. The invention also relates to nucleic acids hybridizable to or complementary to the foregoing sequences. In specific aspects, nucleic acids are provided which comprise a sequence complementary to at least 10, 25, 50, 100, or 200 nucleotides or the entire coding region of an F-box gene.

[0106] The invention further relates to the human genomic sequences of nucleic acids. In specific embodiments, F-box encoding nucleic acids comprise the genomic sequences of SEQ ID NOs: 1, 3, 5, 7, 9, 11 or 13 or the coding regions thereof, or nucleic acids encoding an FBP protein (e.g., a protein having the sequence of SEQ ID Nos: 2, 4, 6, 8, 10, 12 or 14). The invention provides purified nucleic acids consisting of at least 8 nucleotides (i.e., a hybridizable portion) of an FBP gene sequence; in other embodiments, the nucleic acids consist of at least 25 (continuous) nucleotides, 50 nucleotides, 100 nucleotides, 150 nucleotides, or 200 nucleotides of an FBP gene sequence or a full-length FBP gene coding sequence. In another embodiment, the nucleic acids are smaller than 35, 200 or 500 nucleotides in length. Nucleic acids can be single or double stranded. The invention also relates to nucleic acids hybridizable to or complementary to the foregoing sequences. In specific aspects, nucleic acids are provided which comprise a

sequence complementary to at least 10, 25, 50, 100, or 200 nucleotides or the entire coding region of an FBP gene sequence.

[0107] In addition to the human FBP nucleotide sequences disclosed herein, other FBP gene sequences can be identified and readily isolated, without undue experimentation, by molecular biological techniques well known in the art, used in conjunction with the FBP gene sequences disclosed herein. For example, additional human FBP gene sequences at the same or at different genetic loci as those disclosed in SEQ ID Nos: 1, 3, 5, 7, 9, 11 or 13 can be isolated readily. There can exist, for example, genes at other genetic or physical loci within the human genome that encode proteins that have extensive homology to one or more domains of the FBP gene products and that encode gene products functionally equivalent to an FBP gene product. Further, homologous FBP gene sequences present in other species can be identified and isolated readily.

[0108] The FBP nucleotide sequences of the invention further include nucleotide sequences that encode polypeptides having at least 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, or higher amino acid sequence identity to the polypeptides encoded by the FBP nucleotide sequences of SEQ ID No. 1, 3, 5, 7, 9, 11 or 13.

[0109] To determine the percent identity of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., $\% \text{ identity} = \frac{\text{\# of identical overlapping positions}}{\text{total \# of overlapping positions}} \times 100\%$). In one embodiment, the two sequences are the same length.

[0110] The determination of percent identity between two sequences can also be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, Proc. Natl. Acad. Sci. USA 87:2264-2268, modified as in Karlin and Altschul, 1993, Proc. Natl. Acad. Sci. USA 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, et al., 1990, J. Mol. Biol. 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score=100, wordlength=12 to obtain nucleotide sequences homologous to a nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3 to obtain amino acid sequences homologous to a protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., 1997, Nucleic Acids Res. 25:3389-3402. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules (Altschul et al.,

1997, supra). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used (see <http://www.ncbi.nlm.nih.gov>). Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, Proc. Natl. Acad. Sci. USA 87:2264-2268, modified as in Karlin and Altschul, 1993, Proc. Natl. Acad. Sci. USA 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, et al., 1990, J. Mol. Biol. 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score=100, wordlength=12 to obtain nucleotide sequences homologous to a nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3 to obtain amino acid sequences homologous to a protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., 1997, Nucleic Acids Res. 25:3389-3402. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules (Altschul et al., 1997, supra). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used (see <http://www.ncbi.nlm.nih.gov>). Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, 1988, *CABIOS* 4:11-17. Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

[0111] The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent identity, typically only exact matches are counted.

[0112] With respect to identification and isolation of FBP gene sequences present at the same genetic or physical locus as those sequences disclosed herein, such sequences can, for example, be obtained readily by utilizing standard sequencing and bacterial artificial chromosome (BAC) technologies.

[0113] With respect to the cloning of an FBP gene homologue in human or other species (e.g., mouse), the isolated FBP gene sequences disclosed herein may be labeled and used to screen a cDNA library constructed from mRNA obtained from appropriate cells or tissues (e.g., brain tissues) derived from the organism (e.g., mouse) of interest. The hybridization conditions used should be of a lower stringency when the cDNA library is derived from an organism different from the type of organism from which the labeled sequence was derived.

[0114] Alternatively, the labeled fragment may be used to screen a genomic library derived from the organism of interest, again, using appropriately stringent conditions. Low stringency conditions are well known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook, et al., 1989, *Molecular*

Cloning, A Laboratory Manual, Second Edition, Cold Spring Harbor Press, N.Y.; and Ausubel, et al., supra. Further, an FBP gene homologue may be isolated from, for example, human nucleic acid, by performing PCR using two degenerate oligonucleotide primer pools designed on the basis of amino acid sequences within any FBP gene product disclosed herein.

[0115] The PCR product may be subcloned and sequenced to ensure that the amplified sequences represent the sequences of an FBP gene nucleic acid sequence. The PCR fragment may then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment may be labeled and used to screen a bacteriophage cDNA library. Alternatively, the labeled fragment may be used to isolate genomic clones via the screening of a genomic library.

[0116] PCR technology may also be utilized to isolate full length cDNA sequences. For example, RNA may be isolated, following standard procedures, from an appropriate cellular or tissue source (i.e., one known, or suspected, to express the FBP gene, such as, for example, blood samples or brain tissue samples obtained through biopsy or post-mortem). A reverse transcription reaction may be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" with guanines using a standard terminal transferase reaction, the hybrid may be digested with RNAase H, and second strand synthesis may then be primed with a poly-C primer. Thus, cDNA sequences upstream of the amplified fragment may easily be isolated. For a review of cloning strategies that may be used, see e.g., Sambrook et al., supra.

[0117] FBP gene sequences may additionally be used to identify mutant FBP gene alleles. Such mutant alleles may be isolated from individuals either known or proposed to have a genotype that contributes to the symptoms of an FBP gene disorder, such as proliferative or differentiative disorders involved in tumorigenesis or causing cancer, for example. Mutant alleles and mutant allele products may then be utilized in the therapeutic, diagnostic and prognostic systems described below. Additionally, such FBP gene sequences can be used to detect FBP gene regulatory (e.g., promoter) defects which can be associated with an FBP disorder, such as proliferative or differentiative disorders involved in tumorigenesis or causing cancer, for example.

[0118] FBP alleles may be identified by single strand conformational polymorphism (SSCP) mutation detection techniques, Southern blot, and/or PCR amplification techniques. Primers can routinely be designed to amplify overlapping regions of the whole FBP sequence including the promoter region. In one embodiment, primers are designed to cover the exon-intron boundaries such that, first, coding regions can be scanned for mutations. Genomic DNA isolated from lymphocytes of normal and affected individuals is used as PCR template. PCR products from normal and affected individuals are compared, either by single strand conformational polymorphism (SSCP) mutation detection techniques and/or by sequencing. SSCP analysis can be performed as follows: 100 ng of genomic DNA is amplified in a 10 μ l reaction, adding 10 pmols of each primer, 0.5 U of Taq DNA polymerase (Promega), 1 μ Ci of α -[³²P]dCTP

(NEN; specific activity, 3000 Ci/mmol), in 2.5 μ M dNTPs (Pharmacia), 10 mM Tris-HCl (pH 8.8), 50 mM KCl, 1 mM MgCl₂, 0.01% gelatin, final concentration. Thirty cycles of denaturation (94° C.), annealing (56° C. to 64° C., depending on primer melting temperature), and extension (72° C.) is carried out in a thermal-cycler (MJ Research, Boston, Mass., USA), followed by a 7 min final extension at 72° C. Two microliters of the reaction mixture is diluted in 0.1% SDS, 10 mM EDTA and then mixed 1:1 with a sequencing stop solution containing 20 mM NaOH. Samples are heated at 95 C for 5 min, chilled on ice for 3 min and then 31 will be loaded onto a 6% acrylamide/TBE gel containing 5% (v/v) glycerol. Gels are run at 8 W for 12-15 h at room temperature. Autoradiography is performed by exposure to film at -70 C with intensifying screens for different periods of time. The mutations responsible for the loss or alteration of function of the mutant FBP gene product can then be ascertained.

[0119] Alternatively, a cDNA of a mutant FBP gene may be isolated, for example, using PCR. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known or suspected to be expressed in an individual putatively carrying the mutant FBP allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal gene. Using these two primers, the product is then amplified via PCR, cloned into a suitable vector, and subjected to DNA sequence analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the mutant FBP allele to that of the normal FBP allele, the mutation(s) responsible for the loss or alteration of function of the mutant FBP gene product can be ascertained.

[0120] Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of or known to carry a mutant FBP allele, or a cDNA library can be constructed using RNA from a tissue known, or suspected, to express a mutant FBP allele. An unimpaired FBP gene or any suitable fragment thereof may then be labeled and used as a probe to identify the corresponding mutant FBP allele in such libraries. Clones containing the mutant FBP gene sequences may then be purified and subjected to sequence analysis according to methods well known to those of skill in the art.

[0121] Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known, or suspected, to express a mutant FBP allele in an individual suspected of or known to carry such a mutant allele. In this manner, gene products made by the putatively mutant tissue may be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against the normal FBP gene product, as described, below, in Section 5.3. (For screening techniques, see, for example, Harlow and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.)

[0122] Nucleic acids encoding derivatives and analogs of FBP proteins, and FBP antisense nucleic acids can be isolated by the methods recited above. As used herein, a "nucleic acid encoding a fragment or portion of an F-box protein" shall be construed as referring to a nucleic acid

encoding only the recited fragment or portion of the FBP and not the other contiguous portions of the FBP protein as a continuous sequence.

[0123] Fragments of FBP gene nucleic acids comprising regions conserved between (i.e., with homology to) other FBP gene nucleic acids, of the same or different species, are also provided. Nucleic acids encoding one or more FBP domains can be isolated by the methods recited above.

[0124] In cases where an FBP mutation results in an expressed gene product with altered function (e.g., as a result of a missense or a frameshift mutation), a polyclonal set of anti-FBP gene product antibodies are likely to cross-react with the mutant FBP gene product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well known to those of skill in the art.

[0125] 5.2 Proteins and Polypeptides of FBP Genes

[0126] The amino acid sequences depicted in **FIGS. 1, 2**, and parts B of **FIGS. 3 to 28** represent FBP gene products. The FBP1 gene product, sometimes referred to herein as a "FBP1 protein", includes those gene products encoded by the FBP1 gene sequences described in Section 5.1, above. Likewise, the FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25 gene products, referred to herein as an FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25 proteins, include those gene products encoded by the FBP2, FBP3, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25 genes. In accordance with the present invention, the nucleic acid sequences encoding the FBP gene products are derived from eukaryotic genomes, including mammalian genomes. In a preferred embodiment the nucleic acid sequences encoding the FBP gene products are derived from human or murine genomes.

[0127] FBP gene products, or peptide fragments thereof, can be prepared for a variety of uses. For example, such gene products, or peptide fragments thereof, can be used for the generation of antibodies, in diagnostic and prognostic assays, or for the identification of other cellular or extracellular gene products involved in the ubiquitination pathway and thereby implicated in the regulation of cell cycle and proliferative disorders.

[0128] In addition, FBP gene products of the present invention may include proteins that represent functionally equivalent (see Section 5.1 for a definition) gene products. FBP gene products of the invention do not encompass the previously identified mammalian F-box proteins Skp2, Cyclin F, Elongin A, or mouse Md6 (see Pagano, 1997, supra; Zhang et al., 1995 supra; Bai et al., 1996 supra; Skowyra et al., 1997, supra).

[0129] Functionally equivalent FBP gene products may contain deletions, including internal deletions, additions, including additions yielding fusion proteins, or substitutions of amino acid residues within and/or adjacent to the amino acid sequence encoded by the FBP gene sequences

described, above, in Section 5.1, but that result in a "silent" change, in that the change produces a functionally equivalent FBP gene product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

[0130] Alternatively, where alteration of function is desired, deletion or non-conservative alterations can be engineered to produce altered FBP gene products. Such alterations can, for example, alter one or more of the biological functions of the FBP gene product. Further, such alterations can be selected so as to generate FBP gene products that are better suited for expression, scale up, etc. in the host cells chosen. For example, cysteine residues can be deleted or substituted with another amino acid residue in order to eliminate disulfide bridges.

[0131] The FBP gene products, peptide fragments thereof, and fusion proteins thereof, may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing the FBP gene polypeptides, peptides, fusion peptide and fusion polypeptides of the invention by expressing nucleic acid containing FBP gene sequences are described herein. Methods that are well known to those skilled in the art can be used to construct expression vectors containing FBP gene product coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. See, for example, the techniques described in Sambrook, et al., supra, and Ausubel, et al., supra. Alternatively, RNA capable of encoding FBP gene product sequences may be chemically synthesized using, for example, synthesizers. See, for example, the techniques described in "Oligonucleotide Synthesis", 1984, Gait, ed., IRL Press, Oxford.

[0132] A variety of host-expression vector systems may be utilized to express the FBP gene coding sequences of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells that may, when transformed or transfected with the appropriate nucleotide coding sequences, exhibit the FBP gene product of the invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing FBP gene product coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing the FBP gene product coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing the FBP gene product coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g. cauliflower mosaic virus, CAMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing FBP gene product coding

sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

[0133] In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the FBP gene product being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of FBP protein or for raising antibodies to FBP protein, for example, vectors that direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2, 1791), in which the FBP gene product coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye and Inouye, 1985, Nucleic Acids Res. 13, 3101-3109; Van Heeke and Schuster, 1989, J. Biol. Chem. 264, 5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

[0134] In an insect system, *Autographa californica*, nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. **The FBP gene coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of FBP gene coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed (e.g., see Smith et al., 1983, J. Virol. 46: 584; Smith, U.S. Pat. No. 4,215,051).**

[0135] In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the FBP gene coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing FBP gene product in infected hosts. (e.g., See Logan and Shenk, 1984, Proc. Natl. Acad. Sci. USA 81, 3655-3659). Specific initiation signals may also be required for efficient translation of inserted FBP gene product coding sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire FBP gene, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational

control signals may be needed. However, in cases where only a portion of the FBP gene coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner, et al., 1987, Methods in Enzymol. 153, 516-544).

[0136] In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells that possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, and WI38.

[0137] For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines that stably express the FBP gene product may be engineered. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci that in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines that express the FBP gene product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the FBP gene product.

[0138] A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska and Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48, 2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22, 817) genes can be employed in tk-, hgprt- or aprt-cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77, 3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78, 1527); gpt, which confers resistance to mycophenolic acid (Mulligan and Berg, 1981, Proc. Natl. Acad. Sci. USA 78, 2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981,

3. *Mol. Biol.* 150, 1); and hygromycin, which confers resistance to hygromycin (Santerre, et al., 1984, *Gene* 30, 147).

[0139] Alternatively, any fusion protein may be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht, et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, *Proc. Natl. Acad. Sci. USA* 88, 8972-8976). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni²⁺ nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

[0140] The FBP gene products can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, goats, sheep, and non-human primates, e.g., baboons, monkeys, and chimpanzees may be used to generate FBP transgenic animals. The term "transgenic," as used herein, refers to animals expressing FBP gene sequences from a different species (e.g., mice expressing human FBP sequences), as well as animals that have been genetically engineered to overexpress endogenous (i.e., same species) FBP sequences or animals that have been genetically engineered to no longer express endogenous FBP gene sequences (i.e., "knock-out" animals), and their progeny.

[0141] In particular, the present invention relates to FBPI knockout mice. The present invention also relates to transgenic mice which express human wild-type FBP1 and Skp2 gene sequences in addition to mice engineered to express human mutant FBPI and Skp2 gene sequences deleted of their F-box domains. Any technique known in the art may be used to introduce an FBP gene transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to pronuclear microinjection (Hoppe and Wagner, 1989, U.S. Pat. No. 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten, et al., 1985, *Proc. Natl. Acad. Sci., USA* 82, 6148-6152); gene targeting in embryonic stem cells (Thompson, et al., 1989, *Cell* 56, 313-321); electroporation of embryos (Lo, 1983, *Mol. Cell. Biol.* 3, 1803-1814); and sperm-mediated gene transfer (Lavitrano et al., 1989, *Cell* 57, 717-723) (For a review of such techniques, see Gordon, 1989, *Transgenic Animals*, *Intl. Rev. Cytol.* 115, 171-229).

[0142] Any technique known in the art may be used to produce transgenic animal clones containing an FBP transgene, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal or adult cells induced to quiescence (Campbell, et al., 1996, *Nature* 380, 64-66; Wilmut, et al., *Nature* 385, 810-813).

[0143] The present invention provides for transgenic animals that carry an FBP transgene in all their cells, as well as animals that carry the transgene in some, but not all their cells, i.e., mosaic animals. The transgene may be integrated as a single transgene or in concatamers, e.g., head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko, et al., 1992, *Proc. Natl. Acad. Sci. USA* 89, 6232-

6236). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. Examples of regulatory sequences that can be used to direct tissue-specific expression of an FBP transgene include, but are not limited to, the elastase I gene control region which is active in pancreatic acinar cells (Swift et al., 1984, *Cell* 38:639-646; Omitz et al., 1986, Cold Spring Harbor Symp. Quant. Biol. 50:399-409; MacDonald, 1987, *Hepatology* 7:42S-51 S); the insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-122); immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., 1984, *Cell* 38:647-658; Adams et al., 1985, *Nature* 318:533-538; Alexander et al., 1987, *Mol. Cell. Biol.* 7:1436-1444); albumin gene control region which is active in liver (Pinkert et al., 1987, *Genes and Devel.* 1:268-276) alpha-fetoprotein gene control region which is active in liver (Knunlauf et al., 1985, *Mol. Cell. Biol.* 5:1639-1648; Hammer et al., 1987, *Science* 235:53-58); alpha-1-antitrypsin gene control region which is active in liver (Kelsey et al., 1987, *Genes and Devel.* 1:161-171); beta-globin gene control region which is active in myeloid cells (Magram et al., 1985, *Nature* 315:338-340; Kollias et al., 1986, *Cell* 46:89-94); myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead et al., 1987, *Cell* 48:703-712); myosin light chain-2 gene control region which is active in skeletal muscle (Shani, 1985, *Nature* 314:283-286); and gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason et al., 1986, *Science* 234:1372-1378). Promoters isolated from the genome of viruses that grow in mammalian cells, (e.g., vaccinia virus 7.5K, SV40, HSV, adenoviruses MLP, MMTV, LTR and CMV promoters) may be used, as well as promoters produced by recombinant DNA or synthetic techniques.

[0144] When it is desired that the FBP gene transgene be integrated into the chromosomal site of the endogenous FBP gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous FBP gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous FBP gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous FBP gene in only that cell type, by following, for example, the teaching of Gu, et al. (Gu, et al., 1994, *Science* 265, 103-106). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

[0145] Once transgenic animals have been generated, the expression of the recombinant FBP gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques that include but are not limited to Northern blot analysis of tissue samples obtained from the animal, in situ hybridization analysis, and RT-PCR (reverse transcriptase PCR). Samples of FBP gene-expressing tissue, may also be evaluated immunocytochemically

using antibodies specific for the FBP transgene product. Transgenic mice harboring tissue-directed transgenes can be used to test the effects of FBP gene expression in the intact animal. In one embodiment, transgenic mice harboring a human FBP1 transgene in the mammary gland can be used to assess the role of FBPs in mouse mammary development and tumorigenesis. In another embodiment, transgenic mice can be generated that overexpress the human FBP1 dominant negative mutant form (F-box deleted) in the mammary gland. In a specific embodiment, for example, the MMTV LTR promoter (mouse mammary tumor virus long terminal repeat) can be used to direct integration of the transgene in the mammary gland. An MMTV/FBP1 fusion gene can be constructed by fusing sequences of the MMTV LTR promoter to nucleotide sequences upstream of the first ATG of FBP1 gene. An SV40 polyadenylation region can also be fused to sequences downstream of the FBP1 coding region. Transgenic mice are generated by methods well known in the art (Gordon, 1989, *Transgenic Animals*, Intl. Rev. Cytol. 115, 171-229). Briefly, immature B6D2F1 female mice are superovulated and mated to CD-1 males. The following morning the females are examined for the presence of vaginal plugs, and fertilized ova are recovered and microinjected with a plasmid vector. Approximately 2000 copies of the material are microinjected into each pronucleus. Screening of founder animals is performed by extraction of DNA from spleen and Southern hybridization using the MMTV/FBP1 as a probe. Screening of offspring is performed by PCR of tail DNA. Once transgenic pedigrees are established, the expression pattern of the transgene is determined by Northern blot and RT-PCR analysis in different organs in order to correlate it with subsequent pathological changes.

[0146] The resulting transgenic animals can then be examined for the role of FBP genes in tumorigenesis. In one embodiment, for example, FBP transgenes can be constructed for use as a breast cancer model. Overexpression of FBP1 genes in such mice is expected to increase β -catenin ubiquitination and degradation, resulting in a tumor suppressor phenotype. Conversely, overexpression of the FBP1 deletion mutant is expected to result in stabilization of β -catenin and induce proliferation of mammary gland epithelium. These phenotypes can be tested in both female and male transgenic mice, by assays such as those described in Sections 5.4, 5.5 and 7.

[0147] In another specific embodiment, transgenic mice are generated that express FBP1 transgenes in T-lymphocytes. In this embodiment, a CD2/FBP1 fusion gene is constructed by fusion of the CD2 promoter, which drives expression in both CD4 positive and negative T-cells, to sequences located upstream of the first ATG of an FBP gene, e.g., the wild-type and mutant FBP1 genes. The construct can also contain an SV40 polyadenylation region downstream of the FBP gene. After generation and testing of transgenic mice, as described above, the expression of the FBP transgene is examined. The transgene is expressed in thymus and spleen. Overexpression of wild-type FBP1 is expected to result in a phenotype. For example, possible expected phenotypes of FBP1 transgenic mice include increased degradation of IKBA, increased activation of NF κ B, or increased cell proliferation. Conversely, overexpression of the dominant negative mutant, FBP1, lacking the F-box domain, can be expected to have the opposite effect, for example, increased stability of IKBA, decreased activa-

tion of NF κ B, or decreased cell proliferation. Such transgenic phenotypes can be tested by assays such as those used in Section 5.4 and 5.5.

[0148] In another specific embodiment, the SKP2 gene is expressed in T-lymphocytes of transgenic mice. Conversely, the F-box deletion form acts as dominant negative, stabilizing p27 and inhibiting T-cell activation. Construction of the CD2/SKP2 fusion genes and production of transgenic mice are as described above for CD2/FBP fusion genes, using wild-type and mutant SKP2 cDNA, instead of FBP1 cDNA, controlled by the CD2 promoter. Founders and their progeny are analyzed for the presence and expression of the SKP2 transgene and the mutant SKP2 transgene. Expression of the transgene in spleen and thymus is analyzed by Northern blot and RT-PCR.

[0149] In another specific embodiment, transgenic mice are constructed by inactivation of the FBP1 locus in mice. Inactivation of the FBP1 locus in mice by homologous recombination involves four stages: 1) the construction of the targeting vector for FBP1; 2) the generation of ES+/- cells; 3) the production of knock-out mice; and 4) the characterization of the phenotype. A 129 SV mouse genomic phage library is used to identify and isolate the mouse FBP1 gene. Bacteriophages are plated at an appropriate density and an imprint of the pattern of plaques can be obtained by gently layering a nylon membrane onto the surface of agarose dishes. Bacteriophage particles and DNA are transferred to the filter by capillary action in an exact replica of the pattern of plaques. After denaturation, the DNA is bound to the filter by baking and then hybridized with ³²P labeled-FBP1 cDNA. Excess probe is washed away and the filters were then exposed for autoradiography. Hybridizing plaques, identified by aligning the film with the original agar plate, were picked for a secondary and a tertiary screening to obtain a pure plaque preparation. Using this method, positive phage which span the region of interest, for example, the region encoding the F-box, are isolated. Using PCR, Southern hybridization, restriction mapping, subcloning and DNA sequencing the partial structure of the wild-type FBP1 gene can be determined.

[0150] To inactivate the Fbp1 locus by homologous recombination, a gene targeting vector in which exon 3 in the Fbp1 locus is replaced by a selectable marker, for example, the neoR gene, in an antisense orientation can be constructed. Exon 3 encodes the F-box motif which is known to be critical for Fbp1 interaction with Skp1. The targeting construct possesses a short and a long arm of homology flanking a selectable marker gene. One of the vector arms is relatively short (2 kb) to ensure efficient amplification since homologous recombinant ES clones will be screened by PCR. The other arm is >6 kb to maximize the frequency of homologous recombination. A thymidine kinase (tk) gene, included at the end of the long homology arm of the vector provides an additional negative selection marker (using gancyclovir) against ES clones which randomly integrate the targeting vector. Since homologous recombination occurs frequently using linear DNA, the targeting vector is linearized prior to transfection of ES cells. Following electroporation and double drug selection of embryonic stem cell clones, PCR and Southern analysis is used to determine whether homologous recombination has occurred at the FBP1 locus. Screening by PCR is advantageous because a larger number of colonies can be analyzed

with this method than with Southern analysis. In addition, PCR screening allows rapid elimination of negative clones thus to avoid feeding and subsequently freezing all the clones while recombinants are identified. This PCR strategy for detection of homologous recombinants is based on the use of a primer pair chosen such that one primer anneals to a sequence specific to the targeting construct, e.g., sequences of the neomycin gene or other selectable marker, and not in the endogenous locus, and the other primer anneals to a region outside the construct, but within the endogenous locus. Southern analysis is used to confirm that a homologous recombination event has occurred (both at the short arm of homology and at the long arm of homology) and that no gene duplication events have occurred during the recombination.

[0151] Such FBP1 knockout mice can be used to test the role of Fbp1 in cellular regulation and control of proliferation. In one embodiment, phenotype of such mice lacking Fbp1 is cellular hyperplasia and increased tumor formation. In another embodiment, FBP1 null mice phenotypes include, but are not limited to, increased β -catenin activity, stabilization of β -catenin, increased cellular proliferation, accumulation of IK-Ba, decreased NF-KB activity, deficient immune response, inflammation, or increased cell death or apoptotic activity. Alternatively, a deletion of the of the FBP1 gene can result in an embryonic lethality. In this case, heterozygous mice at the FBPI allele can be tested using the above assays, and embryos of null FBP mice can be tested using the assays described above.

[0152] Transgenic mice bearing FBP transgenes can also be used to screen for compounds capable of modulating the expression of the FBP gene and/or the synthesis or activity of the FBP1 gene or gene product. Such compounds and methods for screening are described.

[0153] 5.3 Generation of Antibodies to F-Box Proteins and their Derivatives

[0154] According to the invention, F-box motif, its fragments or other derivatives, or analogs thereof, may be used as an immunogen to generate antibodies which immunospecifically bind such an immunogen. Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain, Fab fragments, and an Fab expression library. In a specific embodiment, antibodies to a human FBP protein are produced. In another embodiment, antibodies to a domain (e.g., the F-box domain or the substrate-binding domain) of an FBP are produced.

[0155] Various procedures known in the art may be used for the production of polyclonal antibodies to an FBP or derivative or analog. In a particular embodiment, rabbit polyclonal antibodies to an epitope of an FBP encoded by a sequence of FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25, or a subsequence thereof, can be obtained (Pagano, M., 1995, "From peptide to purified antibody", in *Cell Cycle: Materials and Methods*. M. Pagano, ed. Springer-Verlag. 217-281). For the production of antibody, various host animals can be immunized by injection with the native FBP, or a synthetic version, or derivative (e.g., fragment) thereof, including but not limited to rabbits, mice, rats, etc. Various adjuvants may be used to increase the immunological response, depending

on the host species, and including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, key-hole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *corynebacterium parvum*.

[0156] For preparation of monoclonal antibodies directed toward an FBP sequence or analog thereof, any technique which provides for the production of antibody molecules by continuous cell lines in culture may be used. For example, the hybridoma technique originally developed by Kohler and Milstein (1975, *Nature* 256:495-497), as well as the trioma technique, the human B-cell hybridoma technique (Kozbor et al., 1983, *Immunology Today* 4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., 1985, in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). In an additional embodiment of the invention, monoclonal antibodies can be produced in germ-free animals utilizing recent technology (PCT/US90/02545). According to the invention, human antibodies may be used and can be obtained by using human hybridomas (Cote et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80:2026-2030) or by transforming human B cells with EBV virus in vitro (Cole et al., 1985, in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, pp. 77-96). In fact, according to the invention, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, *Proc. Natl. Acad. Sci. U.S.A.* 81:6851-6855; Neuberger et al., 1984, *Nature* 312:604-608; Takeda et al., 1985, *Nature* 314:452-454) by splicing the genes from a mouse antibody molecule specific for FBP together with genes from a human antibody molecule of appropriate biological activity can be used; such antibodies are within the scope of this invention.

[0157] According to the invention, techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778) can be adapted to produce FBP-specific single chain antibodies. An additional embodiment of the invention utilizes the techniques described for the construction of Fab expression libraries (Huse et al., 1989, *Science* 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity for FBPs, derivatives, or analogs.

[0158] Antibody fragments which contain the idiotype of the molecule can be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragment which can be produced by pepsin digestion of the antibody molecule; the Fab' fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragment, the Fab fragments which can be generated by treating the antibody molecule with papain and a reducing agent, and Fv fragments.

[0159] In the production of antibodies, screening for the desired antibody can be accomplished by techniques known in the art, e.g. ELISA (enzyme-linked immunosorbent assay). For example, to select antibodies which recognize a specific domain of an FBP, one may assay generated hybridomas for a product which binds to an FBP fragment containing such domain. For selection of an antibody that specifically binds a first FBP homolog but which does not specifically bind a different FBP homolog, one can select on

the basis of positive binding to the first FBP homolog and a lack of binding to the second FBP homolog.

[0160] Antibodies specific to a domain of an FBP are also provided, such as an F-box motif.

[0161] The foregoing antibodies can be used in methods known in the art relating to the localization and activity of the FBP sequences of the invention, e.g., for imaging these proteins, measuring levels thereof in appropriate physiological samples, in diagnostic methods, etc.

[0162] In another embodiment of the invention (see *infra*), anti-FBP antibodies and fragments thereof containing the binding domain are used as therapeutics.

[0163] 5.4 Screening Assays for the Identification of Agents that Interact with F-Box Proteins and/or Interfere with their Enzymatic Activities

[0164] Novel components of the ubiquitin ligase complex, including FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25, interact with cellular proteins to regulate cellular proliferation. One aspect of the present invention provides methods for assaying and screening fragments, derivatives and analogs of the novel components to identify polypeptides or peptides or other compounds that interact with the novel ubiquitin ligases such as potential substrates of ubiquitin ligase activity. The present invention also provides screening assays to identify compounds that modulate or inhibit the interaction of the novel FBPs with other subunits or numbers of the ubiquitin ligase complex, such as Skp1, or ubiquitinating enzymes with which the novel FBPs interact.

[0165] In yet another embodiment, the assays of the present invention may be used to identify polypeptides or peptides or other compounds which inhibit or modulate the interaction between the novel ubiquitin ligases or known (e.g., Skp1) components of the ubiquitin ligase complex with novel or known substrates. By way of example, but not by limitation, the screening assays described herein may be used to identify peptides or proteins that interfere with the interaction between known ubiquitin ligase component, Skp2, and its novel substrate, p27. In another example, compounds that interfere with the interaction between FBP1 and its novel substrate, β -catenin, are identified using the screening assay. In another example, compounds that interfere with the interaction between Skp2 and another putative substrate, E2F, are identified using the screening assay. In yet another example, compounds that interfere with the interaction between FBP1 and another putative substrate, IKB α , are identified using the screening assay.

[0166] In yet another embodiment, the assays of the present invention may be used to identify polypeptides or peptides which inhibit or activate the enzymatic activators of the novel FBPs.

[0167] 5.4.1 Assays for Protein-Protein Interactions

[0168] Derivatives, analogs and fragments of proteins that interact with the novel components of the ubiquitin ligase complex of the present invention can be identified by means of a yeast two hybrid assay system (Fields and Song, 1989, *Nature* 340:245-246 and U.S. Pat. No. 5,283,173). Because the interactions are screened for in yeast, the intermolecular

protein interactions detected in this system occur under physiological conditions that mimic the conditions in mammalian cells (Chien et al., 1991, *Proc. Natl. Acad. Sci. U.S.A.* 88:9578-9581).

[0169] Identification of interacting proteins by the improved yeast two hybrid system is based upon the detection of expression of a reporter gene, the transcription of which is dependent upon the reconstitution of a transcriptional regulator by the interaction of two proteins, each fused to one half of the transcriptional regulator. The "bait" (i.e., the novel components of the ubiquitin ligase complex of the present invention or derivatives or analogs thereof) and "prey" (proteins to be tested for ability to interact with the bait) proteins are expressed as fusion proteins to a DNA binding domain, and to a transcriptional 20 regulatory domain, respectively, or vice versa. In various specific embodiments, the prey has a complexity of at least about 50, about 100, about 500, about 1,000, about 5,000, about 10,000, or about 50,000; or has a complexity in the range of about 25 to about 100,000, about 100 to about 100,000, about 50,000 to about 100,000, or about 100,000 to about 500,000. For example, the prey population can be one or more nucleic acids encoding mutants of a protein (e.g., as generated by site-directed mutagenesis or another method of making mutations in a nucleotide sequence). Preferably, the prey populations are proteins encoded by DNA, e.g., cDNA or genomic DNA or synthetically-generated DNA. For example, the populations can be expressed from chimeric genes comprising cDNA sequences from an un-characterized sample of a population of cDNA from mRNA.

[0170] In a specific embodiment, recombinant biological libraries expressing random peptides can be used as the source of prey nucleic acids.

[0171] In general, proteins of the bait and prey populations are provided as fusion (chimeric) proteins (preferably by recombinant expression of a chimeric coding sequence) comprising each protein contiguous to a pre-selected sequence. For one population, the pre-selected sequence is a DNA binding domain. The DNA binding domain can be any DNA binding domain, as long as it specifically recognizes a DNA sequence within a promoter. For example, the DNA binding domain is of a transcriptional activator or inhibitor. For the other population, the pre-selected sequence is an activator or inhibitor domain of a transcriptional activator or inhibitor, respectively. The regulatory domain alone (not as a fusion to a protein sequence) and the DNA-binding domain alone (not as a fusion to a protein sequence) preferably do not detectably-interact (so as to avoid false positives in the assay). The assay system further includes a reporter gene operably linked to a promoter that contains a binding site for the DNA binding domain of the transcriptional activator (or inhibitor). Accordingly, in the present method of the present invention, binding of a ubiquitin ligase fusion protein to a prey fusion protein leads to reconstitution of a transcriptional activator (or inhibitor) which activates (or inhibits) expression of the reporter gene. The activation (or inhibition) of transcription of the reporter gene occurs intracellularly, e.g., in prokaryotic or eukaryotic cells, preferably in cell culture.

[0172] The promoter that is operably linked to the reporter gene nucleotide sequence can be a native or non-native promoter of the nucleotide sequence, and the DNA binding

site(s) that are recognized by the DNA binding domain portion of the fusion protein can be native to the promoter (if the promoter normally contains such binding site(s)) or non-native to the promoter.

[0173] Alternatively, the transcriptional activation binding site of the desired gene(s) can be deleted and replaced with GAL4 binding sites (Bartel et al., 1993, *BioTechniques* 14:920-924, Chasman et al., 1989, *Mol. Cell. Biol.* 9:4746-4749). The reporter gene preferably contains the sequence encoding a detectable or selectable marker, the expression of which is regulated by the transcriptional activator, such that the marker is either turned on or off in the cell in response to the presence of a specific interaction. Preferably, the assay is carried out in the absence of background levels of the transcriptional activator (e.g., in a cell that is mutant or otherwise lacking in the transcriptional activator).

[0174] The activation domain and DNA binding domain used in the assay can be from a wide variety of transcriptional activator proteins, as long as these transcriptional activators have separable binding and transcriptional activation domains. For example, the GALA protein of *S. cerevisiae* (Ma et al., 1987, *Cell* 48:847-853), the GCN4 protein of *S. cerevisiae* (Hope & Struhl, 1986, *Cell* 46:885-894), the ARD1 protein of *S. cerevisiae* (Thukral et al., 1989, *Mol. Cell. Biol.* 9:2360-2369), and the human estrogen receptor (Kumar et al., 1987, *Cell* 51:941-951), have separable DNA binding and activation domains. The DNA binding domain and activation domain that are employed in the fusion proteins need not be from the same transcriptional activator. In a specific embodiment, a GALA or LEXA DNA binding domain is employed. In another specific embodiment, a GAIA or herpes simplex virus VP16 (Triezenberg et al., 1988, *Genes Dev.* 2:730-742) activation domain is employed. In a specific embodiment, amino acids 1-147 of GAL4 (Ma et al., 1987, *Cell* 48:847-853; Ptashne et al., 1990, *Nature* 346:329-331) is the DNA binding domain, and amino acids 411-455 of VP16 (Triezenberg et al., 1988, *Genes Dev.* 2:730-742; Cress et al., 1991, *Science* 251:87-90) comprise the activation domain.

[0175] In a preferred embodiment, the yeast transcription factor GAL4 is reconstituted by protein-protein interaction and the host strain is mutant for GAL4. In another embodiment, the DNA-binding domain is Ace1N and/or the activation domain is Ace1, the DNA binding and activation domains of the Ace1 protein, respectively. Ace1 is a yeast protein that activates transcription from the CUP1 operon in the presence of divalent copper. CUP1 encodes metallothionein, which chelates copper, and the expression of CUP1 protein allows growth in the presence of copper, which is otherwise toxic to the host cells. The reporter gene can also be a CUP1-lacZ fusion that expresses the enzyme beta-galactosidase (detectable by routine chromogenic assay) upon binding of a reconstituted Ace1N transcriptional activator (see Chaudhuri et al., 1995, *FEBS Letters* 357:221-226). In another specific embodiment, the DNA binding domain of the human estrogen receptor is used, with a reporter gene driven by one or three estrogen receptor response elements (Le Douarin et al., 1995, *Nucl. Acids. Res.* 23:876-878). The DNA binding domain and the transcriptional activator/inhibitor domain each preferably has a nuclear localization signal (see Ylikomi et al., 1992, *EMBO*

J. 11:3681-3694, Dingvall and Laskey, 1991, *TIBS* 16:479-481) functional in the cell in which the fusion proteins are to be expressed.

[0176] To facilitate isolation of the encoded proteins, the fusion constructs can further contain sequences encoding affinity tags such as glutathione-S-transferase or maltose-binding protein or an epitope of an available antibody, for affinity purification (e.g., binding to glutathione, maltose, or a particular antibody specific for the epitope, respectively) (Allen et al., 1995, *TIBS* 20:511-516). In another embodiment, the fusion constructs further comprise bacterial promoter sequences for recombinant production of the fusion protein in bacterial cells.

[0177] The host cell in which the interaction assay occurs can be any cell, prokaryotic or eukaryotic, in which transcription of the reporter gene can occur and be detected, including, but not limited to, mammalian (e.g., monkey, mouse, rat, human, bovine), chicken, bacterial, or insect cells, and is preferably a yeast cell. Expression constructs encoding and capable of expressing the binding domain fusion proteins, the transcriptional activation domain fusion proteins, and the reporter gene product(s) are provided within the host cell, by mating of cells containing the expression constructs, or by cell fusion, transformation, electroporation, microinjection, etc.

[0178] Various vectors and host strains for expression of the two fusion protein populations in yeast are known and can be used (see e.g., U.S. Pat. No. 5,1468,614; Bartel et al., 1993, "Using the two-hybrid system to detect protein-protein interactions" In: *Cellular Interactions in Development*, Hartley, ed., Practical Approach Series xviii, IRL Press at Oxford University Press, New York, N.Y., pp. 153-179; Fields and Stemglanz, 1994, *Trends In Genetics* 10:286-292).

[0179] If not already lacking in endogenous reporter gene activity, cells mutant in the reporter gene may be selected by known methods, or the cells can be made mutant in the target reporter gene by known gene-disruption methods prior to introducing the reporter gene (Rothstein, 1983, *Meth. Enzymol.* 101:202-211).

[0180] In a specific embodiment, plasmids encoding the different fusion protein populations can be introduced simultaneously into a single host cell (e.g., a haploid yeast cell) containing one or more reporter genes, by co-transformation, to conduct the assay for protein-protein interactions. Or, preferably, the two fusion protein populations are introduced into a single cell either by mating (e.g., for yeast cells) or cell fusions (e.g., of mammalian cells). In a mating type assay, conjugation of haploid yeast cells of opposite mating type that have been transformed with a binding domain fusion expression construct (preferably a plasmid) and an activation (or inhibitor) domain fusion expression construct (preferably a plasmid), respectively, will deliver both constructs into the same diploid cell. The mating type of a yeast strain may be manipulated by transformation with the HO gene (Herskowitz and Jensen, 1991, *Meth. Enzymol.* 194:132-146).

[0181] In a preferred embodiment, a yeast interaction mating assay is employed using two different types of host cells, strain-type a and alpha of the yeast *Saccharomyces cerevisiae*. The host cell preferably contains at least two

reporter genes, each with one or more binding sites for the DNA-binding domain (e.g., of a transcriptional activator). The activator domain and DNA binding domain are each parts of chimeric proteins formed from the two respective populations of proteins. One strain of host cells, for example the α strain, contains fusions of the library of nucleotide sequences with the DNA-binding domain of a transcriptional activator, such as GAL4. The hybrid proteins expressed in this set of host cells are capable of recognizing the DNA-binding site in the promoter or enhancer region in the reporter gene construct. The second set of yeast host cells, for example, the β strain, contains nucleotide sequences encoding fusions of a library of DNA sequences fused to the activation domain of a transcriptional activator.

[0182] In another embodiment, the fusion constructs are introduced directly into the yeast chromosome via homologous recombination. The homologous recombination for these purposes is mediated through yeast sequences that are not essential for vegetative growth of yeast, e.g., the MER2, MER1, ZIPI, REC102, or ME14 gene.

[0183] Bacteriophage vectors can also be used to express the DNA binding domain and/or activation domain fusion proteins. Libraries can generally be prepared faster and more easily from bacteriophage vectors than from plasmid vectors.

[0184] In a specific embodiment, the present invention provides a method of detecting one or more protein-protein interactions comprising (a) recombinantly expressing a novel ubiquitin ligase component of the present invention or a derivative or analog thereof in a first population of yeast cells being of a first mating type and comprising a first fusion protein containing the sequence of a novel ubiquitin ligase component of the present invention and a DNA binding domain, wherein said first population of yeast cells contains a first nucleotide sequence operably linked to a promoter driven by one or more DNA binding sites recognized by said DNA binding domain such that an interaction of said first fusion protein with a second fusion protein, said second fusion protein comprising a transcriptional activation domain, results in increased transcription of said first nucleotide sequence; (b) negatively selecting to eliminate those yeast cells in said first population in which said increased transcription of said first nucleotide sequence occurs in the absence of said second fusion protein; (c) recombinantly expressing in a second population of yeast cells of a second mating type different from said first mating type, a plurality of said second fusion proteins, each second fusion protein comprising a sequence of a fragment, derivative or analog of a protein and an activation domain of a transcriptional activator, in which the activation domain is the same in each said second fusion protein; (d) mating said first population of yeast cells with said second population of yeast cells to form a third population of diploid yeast cells, wherein said third population of diploid yeast cells contains a second nucleotide sequence operably linked to a promoter driven by a DNA binding site recognized by said DNA binding domain such that an interaction of a first fusion protein with a second fusion protein results in increased transcription of said second nucleotide sequence, in which the first and second nucleotide sequences can be the same or different; and (e) detecting said increased transcription of said first and/or second nucleotide sequence, thereby detecting an interaction between a first fusion protein and a second fusion protein.

[0185] 5.4.2 Assays to Identify F-Box Protein Interactions with known Proteins Including Potential Substrates

[0186] The cellular abundance of cell-cycle regulatory proteins, such as members of the cyclin family or the Cki inhibitory proteins, is regulated by the ubiquitin pathway. The enzymes responsible for the ubiquitination of mammalian cell cycle regulation are not known. In yeast, SCF complexes represent the ubiquitin ligases for cell cycle regulators. The F-box component of the ubiquitin ligase complexes, such as the novel F-box proteins of the invention, determines the specificity of the target of the ubiquitin ligase complex. The invention therefore provides assays to screen known molecules for specific binding to F-box protein nucleic acids, proteins, or derivatives under conditions conducive to binding, and then molecules that specifically bind to the FBP protein are identified.

[0187] In a specific embodiment, the invention provides a method for studying the interaction between the F-box protein FBP1 and the Cull/Skp1 complex, and its role in regulating the stability of β -catenin. Protein-protein interactions can be probed in vivo and in vitro using antibodies specific to these proteins, as described in detail in the experiments in Section 8.

[0188] In another specific embodiment, the invention provides for a method for detecting the interaction between the F-box protein Skp2 and E2F-1, a transcription factor involved in cell cycle progression. Insect cells can be infected with baculoviruses co-expressing Skp2 and E2F-1, and cell extracts can be prepared and analyzed for protein-protein interactions. As described in detail in Section 7, this assay has been used successfully to identify potential targets, such as E2F, for known F-box proteins, such as Skp2. This assay can be used to identify other Skp2 targets, as well as targets for novel F-box proteins.

[0189] In another specific embodiment, methods for detecting the interaction between Skp2 and p27, a cell cycle regulated cyclin-dependent kinase (Cdk) inhibitor, are provided. The interaction between Skp2 and p27 may be targeted to identify modulators of Skp2 activity, including its interaction with cell cycle regulators, such as p27. The ubiquitination of Skp2-specific substrates, such as p27 may be used as a means of measuring the ability of a test compound to modulate Skp2 activity. In another embodiment of the screening assays of the present invention, immunodepletion assays, as described in Section 9, can be used to identify modulators of the Skp2/p27 interaction. In particular, Section 9 describes a method for detection of ubiquitination activity in vitro using p27 as a substrate, which can also be used to identify modulators of the Skp2-dependent ubiquitination of p27. In another embodiment of the screening assays of the present invention, antisense oligonucleotides, as described in Section 5.7.1, can be used as inhibitors of the Skp2 activity. Such identified modulators of p27 ubiquitination/degradation and of the Skp2/p27 interaction can be useful in anti-cancer therapies.

[0190] The invention further provides methods for screening ubiquitin ligase complexes having novel F-box proteins (or fragments thereof) as one of their components for ubiquitin ligase activity using known cell-cycle regulatory molecules as potential substrates for ubiquitination. For example, cells engineered to express FBP nucleic acids can be used to recombinantly produce FBP proteins either wild-

type or dominant negative mutants in cells that also express a putative ubiquitin-ligase substrate molecule. Such candidates for substrates of the novel FBP of the present invention include, but are not limited to, such potential substrates as IKB α , β -catenin, myc, E2F-1, p27, p21, cyclin A, cyclin B, cycD1, cyclin E and p53. Then the extracts can be used to test the association of F-box proteins with their substrates, (by Western blot immunoassays) and whether the presence of the FBP increases or decreases the level of the potential substrates.

[0191] 5.5 Assays for the Identification of Compounds that Modulate the Activity of F-Box Proteins

[0192] The present invention relates to in vitro and in vivo assay systems described in the subsections below, which can be used to identify compounds or compositions that modulate the interaction of known FBPs with novel substrates and novel components of the ubiquitin ligase complex. The screening assays of the present invention may also be used to identify compounds or compositions that modulate the interaction of novel FBPs with their identified substrates and components of the ubiquitin ligase complex.

[0193] Methods to screen potential agents for their ability to disrupt or moderate FBP expression and activity can be designed based on the Applicants' discovery of novel FBPs and their interaction with other components of the ubiquitin ligase complex as well as its known and potential substrates. For example, candidate compounds can be screened for their ability to modulate the interaction of an FBP and Skp1, or the specific interactions of Skp2 with E2F-1, Skp2 with p27, or the FBP1/Cul1/Skp1 complex with β -catenin. In principle, many methods known to those of skill in the art, can be readily adapted in designed the assays of the present invention.

[0194] The screening assays of the present invention also encompass high-throughput screens and assays to identify modulators of FBP expression and activity. In accordance with this embodiment, the systems described below may be formulated into kits. To this end, cells expressing FBP and components of the ubiquitination ligase complex and the ubiquitination pathway, or cell lysates, thereof can be packaged in a variety of containers, e.g., vials, tubes, microtitre well plates, bottles, and the like. Other reagents can be included in separate containers and provided with the kit; e.g., positive control samples, negative control samples, buffers, cell culture media, etc.

[0195] The invention provides screening methodologies useful in the identification of proteins and other compounds which bind to, or otherwise directly interact with, the FBP genes and their gene products. Screening methodologies are well known in the art (see e.g., PCT International Publication No. WO 96/34099, published Oct. 31, 1996, which is incorporated by reference herein in its entirety). The proteins and compounds include endogenous cellular components which interact with the identified genes and proteins in vivo and which, therefore, may provide new targets for pharmaceutical and therapeutic interventions, as well as recombinant, synthetic, and otherwise exogenous compounds which may have binding capacity and, therefore, may be candidates for pharmaceutical agents. Thus, in one series of embodiments, cell lysates or tissue homogenates may be screened for proteins or other compounds which bind to one of the normal or mutant FBP genes and FBP proteins.

[0196] Alternatively, any of a variety of exogenous compounds, both naturally occurring and/or synthetic (e.g., libraries of small molecules or peptides), may be screened for binding capacity. All of these methods comprise the step of mixing an FBP protein or fragment with test compounds, allowing time for any binding to occur, and assaying for any bound complexes. All such methods are enabled by the present disclosure of substantially pure FBP proteins, substantially pure functional domain fragments, fusion proteins, antibodies, and methods of making and using the same.

[0197] 5.5.1 Assays for F-Box Protein Agonists and Antagonists

[0198] FBP nucleic acids, F-box proteins, and derivatives can be used in screening assays to detect molecules that specifically bind to FBP nucleic acids, proteins, or derivatives and thus have potential use as agonists or antagonists of FBPs, in particular, molecules that thus affect cell proliferation. In a preferred embodiment, such assays are performed to screen for molecules with potential utility as anti-cancer drugs or lead compounds for drug development. The invention thus provides assays to detect molecules that specifically bind to FBP nucleic acids, proteins, or derivatives. For example, recombinant cells expressing FBP nucleic acids can be used to recombinantly produce FBP proteins in these assays, to screen for molecules that bind to an FBP protein. Similar methods can be used to screen for molecules that bind to FBP derivatives or nucleic acids. Methods that can be used to carry out the foregoing are commonly known in the art. The assays of the present invention may be first optimized on a small scale (i.e., in test tubes), and then scaled up for high-throughput assays. The screening assays of the present may be performed in vitro, i.e. in test tubes, using purified components or cell lysates. The screening assays of the present invention may also be carried out in intact cells in culture and in animal models. In accordance with the present invention, test compounds which are shown to modulate the activity of the FBP as described herein in vitro, will further be assayed in vivo, including cultured cells and animal models to determine if the test compound has the similar effects in vivo and to determine the effects of the test compound on cell cycle progression, the accumulation or degradation of positive and negative regulators, cellular proliferation etc.

[0199] In accordance with the present invention, screening assays may be designed to detect molecules which act as agonists or antagonists of the activity of the novel F-box proteins. In accordance with this aspect of the invention, the test compound may be added to an assay system to measure its effect on the activity of the novel FBP, i.e., ubiquitination of its substrates, interaction with other components of the ubiquitin ligase complex, etc. These assays should be conducted both in the presence and absence of the test compound.

[0200] In accordance with the present invention, ubiquitination activity of a novel FBP in the presence or absence of a test compound can be measured in vitro using purified components of the ubiquitination pathway or may be measured using crude cellular extracts obtained from tissue culture cells or tissue samples. In another embodiment of the aspect of the present invention the screening may be performed by adding the test agent to in vitro translation systems such as a rabbit reticulocyte lysate (RRL) system

and then proceeding with the established analysis. As another alternative, purified or partially purified components which have been determined to interact with one another by the methods described above can be placed under conditions in which the interaction between them would normally occur, with and without the addition of the test agent, and the procedures previously established to analyze the interaction can be used to assess the impact of the test agent. In this approach, the purified or partially purified components may be prepared by fractionation of extracts of cells expressing the components of the ubiquitin ligase complex and pathway, or they may be obtained by expression of cloned genes or cDNAs or fragments thereof, optionally followed by purification of the expressed material.

[0201] Within the broad category of in vitro selection methods, several types of method are likely to be particularly convenient and/or useful for screening test agents. These include but are not limited to methods which measure a binding interaction between two or more components of the ubiquitin ligase complex or interaction with the target substrate, methods which measure the activity of an enzyme which is one of the interacting components, and methods which measure the activity or expression of "reporter" protein, that is, an enzyme or other detectable or selectable protein, which has been placed under the control of one of the components.

[0202] Binding interactions between two or more components can be measured in a variety of ways. One approach is to label one of the components with an easily detectable label, place it together with the other component(s) in conditions under which they would normally interact, perform a separation step which separates bound labeled component from unbound labeled component, and then measure the amount of bound component. The effect of a test agent included in the binding reaction can be determined by comparing the amount of labeled component which binds in the presence of this agent to the amount which binds in its absence.

[0203] In another embodiment, screening can be carried out by contacting the library members with an FBP protein (or nucleic acid or derivative) immobilized on a solid phase and harvesting those library members that bind to the protein (or nucleic acid or derivative). Examples of such screening methods, termed "panning" techniques are described by way of example in Parmley & Smith, 1988, *Gene* 73:305-318; Fowlkes et al., 1992, *BioTechniques* 13:422-427; PCT Publication No. WO 94/18318; and in references cited hereinabove.

[0204] In another embodiment, the two-hybrid system for selecting interacting proteins or peptides in yeast (Fields & Song, 1989, *Nature* 340:245-246; Chien et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:9578-9582) can be used to identify molecules that specifically bind to an FBP protein or derivative.

[0205] Alternatively, test methods may rely on measurements of enzyme activity, such as ubiquitination of the target substrate. Once a substrate of a novel FBP is identified or a novel putative substrate of a known FBP is identified, such as the novel substrates of Skp2, E2F and p27, these components may be used in assays to determine the effect of a test compound on the ubiquitin ligase activity of the ubiquitin ligase complex.

[0206] In one embodiment, the screening assays may be conducted with a purified system in the presence and absence of test compound. Purified substrate is incubated together with purified ubiquitin ligase complex, ubiquitin conjugating enzymes, ubiquitin activating enzymes and ubiquitin in the presence or in the absence of test compound. Ubiquitination of the substrate is analyzed by immunoassay (see Pagano et al., 1995, *Science* 269:682-685). Briefly, ubiquitination of the substrate can be performed in vitro in reactions containing 50-200 ng of proteins in 50 mM Tris pH 7.5, 5 mM MgCl₂, 2 mM ATPy-S, 0.1 mM DTT and 5 μ M of biotinylated ubiquitin. Total reactions (30 μ l) can be incubated at 25° C. for up to 3 hours in the presence or absence of test compound and then loaded on an 8% SDS gel or a 4-20% gradient gel for analysis. The gels are run and proteins are electrophoretically transferred to nitrocellulose. Ubiquitination of the substrate can be detected by immunoblotting. Ubiquitinated substrates can be visualized using Extravidin-HRP (Sigma), or by using a substrate-specific antibody, and the ECL detection system (NEN).

[0207] In another embodiment, ubiquitination of the substrate may be assayed in intact cells in culture or in animal models in the presence and absence of the test compound. For example, the test compound may be administered directly to an animal model or to crude extracts obtained from animal tissue samples to measure ubiquitination of the substrate in the presence and absence of the test compounds. For these assays, host cells to which the test compound is added may be genetically engineered to express the FBP components of the ubiquitin ligase pathway and the target substrate, the expression of which may be transient, induced or constitutive, or stable. For the purposes of the screening methods of the present invention, a wide variety of host cells may be used including, but not limited to, tissue culture cells, mammalian cells, yeast cells, and bacteria. Each cell type has its own set of advantages and drawbacks. Mammalian cells such as primary cultures of human tissue cells may be a preferred cell type in which to carry out the assays of the present invention, however these cell types are sometimes difficult to cultivate. Bacteria and yeast are relatively easy to cultivate but process proteins differently than mammalian cells. This ubiquitination assay may be conducted as follows: first, the extracts are prepared from human or animal tissue. To prepare animal tissue samples preserving ubiquitinating enzymes, 1 g of tissue can be sectioned and homogenized at 15,000 r.p.m. with a Brinkmann Polytron homogenizer (PT 3000, Westbury, N.Y.) in 1 ml of ice-cold double-distilled water. The sample is frozen and thawed 3 times. The lysate is spun down at 15,000 r.p.m. in a Beckman JA-20.1 rotor (Beckman Instruments, Palo Alto, Calif.) for 45 min at 4° C. The supernatant is retrieved and frozen at -80° C. This method of preparation of total extract preserves ubiquitinating enzymes (Loda et al. 1997, *Nature Medicine* 3:231-234, incorporated by reference herein in its entirety).

[0208] Purified recombinant substrate is added to the assay system and incubated at 37° C. for different times in 30 μ l of ubiquitination mix containing 100 μ g of protein tissue homogenates, 50 mM Tris-HCl (pH 8.0), 5 mM MgCl₂, and 1 mM DTT, 2 mM ATP, 10 mM creatine phosphokinase, 10 mM creatine phosphate and 5 μ M biotinylated ubiquitin. The substrate is then re-purified with antibodies or affinity chromatography. Ubiquitination of the substrate is measured by

immunoassays with either antibodies specific to the substrates or with Extravidin-HRP.

[0209] In addition, *Drosophila* can be used as a model system in order to detect genes that phenotypically interact with FBP. For example, overexpression of FBP in *Drosophila* eye leads to a smaller and rougher eye. Mutagenesis of the fly genome can be performed, followed by selecting flies in which the mutagenesis has resulted in suppression or enhancement of the small rough eye phenotype; the mutated genes in such flies are likely to encode proteins that interact/bind with FBP. Active compounds identified with methods described above will be tested in cultured cells and/or animal models to test the effect of blocking in vivo FBP activity (e.g. effects on cell proliferation, accumulation of substrates, etc.).

[0210] In various other embodiments, screening the can be accomplished by one of many commonly known methods. See, e.g., the following references, which disclose screening of peptide libraries: Pammlay & Smith, 1989, Adv. Exp. Med. Biol. 251:215-218; Scott & Smith, 1990, Science 249:386-390; Fowlkes et al., 1992, BioTechniques 13:422-427; Oldenburg et al., 1992, Proc. Natl. Acad. Sci. USA 89:5393-5397; Yu et al., 1994, Cell 76:933-945; Staudt et al., 1988, Science 241:577-580; Bock et al., 1992, Nature 355:564-566; Tuerk et al., 1992, Proc. Natl. Acad. Sci. USA 89:6988-6992; Ellington et al., 1992, Nature 355:850-852; U.S. Pat. No. 5,096,815, U.S. Pat. No. 5,223,409, and U.S. Pat. No. 5,198,346, all to Ladner et al.; Rebar & Pabo, 1993, Science 263:671-673; and PCT Publication No. WO 94/18318.

[0211] Compounds, peptides, and small molecules can be used in screening assays to identify candidate agonists and antagonists. In one embodiment, peptide libraries may be used to screen for agonists or antagonists of the FBP of the present invention diversity libraries, such as random or combinatorial peptide or non-peptide libraries can be screened for molecules that specifically bind to FBP. Many libraries are known in the art that can be used, e.g., chemically synthesized libraries, recombinant (e.g., phage display libraries), and in vitro translation-based libraries.

[0212] Examples of chemically synthesized libraries are described in Fodor et al., 1991, Science 251:767-773; Houghten et al., 1991, Nature 354:84-86; Lam et al., 1991, Nature 354:82-84; Medynski, 1994, Biofrechnology 12:709-710; Gallop et al., 1994, J. Medicinal Chemistry 37(9):1233-1235; Ohlmeyer et al., 1993, Proc. Natl. Acad. Sci. USA 90:10922-10926; Erb et al., 1994, Proc. Natl. Acad. Sci. USA 91:11422-11426; Houghten et al., 1992, Biotechniques 13:412; Jayawickreme et al., 1994, Proc. Natl. Acad. Sci. USA 91:1614-1618; Salmon et al., 1993, Proc. Natl. Acad. Sci. USA 90:11708-11712; PCT Publication No. WO 93/20242; and Brenner and Lerner, 1992, Proc. Natl. Acad. Sci. USA 89:5381-5383.

[0213] Examples of phage display libraries are described in Scott & Smith, 1990, Science 249:386-390; Devlin et al., 1990, Science, 249:404-406; Christian, et al., 1992, J. Mol. Biol. 227:711-718; Lenstra, 1992, J. Immunol. Meth. 152:149-157; Kay et al., 1993, Gene 128:59-65; and PCT Publication No. WO 94/18318 dated Aug. 18, 1994.

[0214] In vitro translation-based libraries include but are not limited to those described in PCT Publication No. WO

91/05058 dated Apr. 18, 1991; and Mattheakis et al., 1994, Proc. Natl. Acad. Sci. USA 91:9022-9026.

[0215] By way of examples of non-peptide libraries, a benzodiazepine library (see e.g., Bunin et al., 1994, Proc. Natl. Acad. Sci. USA 91:4708-4712) can be adapted for use. Peptoid libraries (Simon et al., 1992, Proc. Natl. Acad. Sci. USA 89:9367-9371) can also be used. Another example of a library that can be used, in which the amide functionalities in peptides have been permethylated to generate a chemically transformed combinatorial library, is described by Ostresh et al. (1994, Proc. Natl. Acad. Sci. USA 91:11138-11142).

[0216] 5.5.2 Assays for the Identification of Compounds that Modulate the Interaction of F-box Proteins with Other Proteins

[0217] Once a substrate or interacting protein is identified, as described in detail in Section 5.4, then one can assay for modulators of the F-box protein interaction with such a protein. The present invention provides for methods of detecting agonists and antagonists of such interactions.

[0218] In one embodiment, the invention encompasses methods to identify modulators, such as inhibitors or agonists, of the interaction between the F-box protein Skp2 and E2F-1, identified in Section 7 and FIG. 10. Such methods comprise both in vivo and in vitro assays for modulator activity. For example, in an in vivo assay, insect cells can be co-infected with baculoviruses co-expressing Skp2 and E2F-1 as well as potential modulators of the Skp2/E2F-1 interaction. The screening methods of the present invention encompass in vitro assays which measure the ability of a test compound to inhibit the enzymatic activity of Skp2 as described above in Section 5.5.1. Cell extracts can be prepared and analyzed for protein-protein interactions by gel electrophoresis and detected by immunoblotting, as described in detail in Section 7 and presented in FIG. 10. Alternatively, an in vitro protein-protein interaction assay can be used. Recombinant purified Skp2, E2F-1, and putative agonist or antagonist molecules can be incubated together, under conditions that allow binding to occur, such as 37 C for 30 minutes. Protein-protein complex formation can be detected by gel analysis, such as those described herein in Section 7. This assay can be used to identify modulators of interactions of known FBP, such as Skp2 with novel substrates.

[0219] In another embodiment, the invention provides for a method for identification of modulators of F-box protein/Skp1 interaction. Such agonist and antagonists can be identified in vivo or in vitro. For example, in an in vitro assay to identify modulators of F-box protein/Skp1 interactions, purified Skp1 and the novel FBP can be incubated together, under conditions that allow binding occur, such as 37C for 30 minutes. In a parallel reaction, a potential agonist or antagonist, as described above in Section 5.5.1, is added either before or during the box protein/Skp1 incubation. Protein-protein interactions can be detected by gel analysis, such as those described herein in Section 7. Modulators of FBP activities and interactions with other proteins can be used as therapeutics using the methods described herein, in Section 5.7.

[0220] These assays may be carried out utilizing any of the screening methods described herein, including the following

in vitro assay. The screening can be performed by adding the test agent to intact cells which express components of the ubiquitin pathway, and then examining the component of interest by whatever procedure has been established. Alternatively, the screening can be performed by adding the test agent to in vitro translation reactions and then proceeding with the established analysis. As another alternative, purified or partially purified components which have been determined to interact with one another by the methods described above can be placed under conditions in which the interaction between them would normally occur, with and without the addition of the test agent, and the procedures previously established to analyze the interaction can be used to assess the impact of the test agent. In this approach, the purified or partially purified components may be prepared by fractionation of extracts of cells expressing the components of the ubiquitin ligase complex and pathway, or they may be obtained by expression of cloned genes or cDNAs or fragments thereof, optionally followed by purification of the expressed material.

[0221] Within the broad category of in vitro selection methods, several types of method are likely to be particularly convenient and/or useful for screening test agents. These include but are not limited to methods which measure a binding interaction between two or more components of the ubiquitin ligase complex or interaction with the target substrate, methods which measure the activity of an enzyme which is one of the interacting components, and methods which measure the activity or expression of "reporter" protein, that is, an enzyme or other detectable or selectable protein, which has been placed under the control of one of the components.

[0222] Binding interactions between two or more components can be measured in a variety of ways. One approach is to label one of the components with an easily detectable label, place it together with the other component(s) in conditions under which they would normally interact, perform a separation step which separates bound labeled component from unbound labeled component, and then measure the amount of bound component. The effect of a test agent included in the binding reaction can be determined by comparing the amount of labeled component which binds in the presence of this agent to the amount which binds in its absence.

[0223] The separation step in this type of procedure can be accomplished in various ways. In one approach, (one of) the binding partner(s) for the labeled component can be immobilized on a solid phase prior to the binding reaction, and unbound labeled component can be removed after the binding reaction by washing the solid phase. Attachment of the binding partner to the solid phase can be accomplished in various ways known to those skilled in the art, including but not limited to chemical cross-linking, non-specific adhesion to a plastic surface, interaction with an antibody attached to the solid phase, interaction between a ligand attached to the binding partner (such as biotin) and a ligand-binding protein (such as avidin or streptavidin) attached to the solid phase, and so on.

[0224] Alternatively, the separation step can be accomplished after the labeled component had been allowed to interact with its binding partner(s) in solution. If the size differences between the labeled component and its binding

partner(s) permit such a separation, the separation can be achieved by passing the products of the binding reaction through an ultrafilter whose pores allow passage of unbound labeled component but not of its binding partner(s) or of labeled component bound to its partner(s). Separation can also be achieved using any reagent capable of capturing a binding partner of the labeled component from solution, such as an antibody against the binding partner, a ligand-binding protein which can interact with a ligand previously attached to the binding partner, and so on.

[0225] 5.6 Methods and Compositions for Diagnostic Use of F-Box Proteins, Derivatives, and Modulators

[0226] Cell cycle regulators are the products of oncogenes (cyclins, β -catenin, etc.), or tumor suppressor genes (ckis, p53, etc.) The FBPs, part of ubiquitin ligase complexes, might therefore be products of oncogenes or tumor suppressor genes, depending on which cell cycle regulatory proteins for which they regulate cellular abundance.

[0227] FBP proteins, analogues, derivatives, and subsequences thereof, FBP nucleic acids (and sequences complementary thereto), anti-FBP antibodies, have uses in diagnostics. The FBP and FBP nucleic acids can be used in assays to detect, prognose, or diagnose proliferative or differentiative disorders, including tumorigenesis, carcinomas, adenomas etc. The novel FBP nucleic acids of the present invention are located at chromosome sites associated with karyotypic abnormalities and loss of heterozygosity. The FBPI nucleic acid of the present invention is mapped and localized to chromosome position 10q24, the loss of which has been demonstrated in 10% of human prostate tumors and small cell lung carcinomas (SCLC), suggesting the presence of a tumor suppressor gene at this location. In addition, up to 7% of childhood acute T-cell leukemia is accompanied by a translocation involving 10q24 as a breakpoint, either t(10;14)(q24;q11) or t(7;10)(q35;q24). 9q34 region (where FBP2 is located) has been shown to be a site of loss of heterozygosity (LOH) in human ovarian and bladder cancers. The FBP2 nucleic acid of the present invention is mapped and localized to chromosome position 9q34 which has been shown to be a site of loss of heterozygosity (LOH) in human ovarian and bladder cancers. The FBP3 nucleic acid of the present invention is mapped and localized to chromosome position 13q22, a region known to contain a putative tumor suppressor gene with loss of heterozygosity in approx. 75% of human SCLC. The FBP4 nucleic acid of the present invention is mapped and localized to chromosome position 5p12, a region shown to be a site of karyotypic abnormalities in a variety of tumors, including human breast cancer and nasopharyngeal carcinomas. The FBP5 nucleic acid of the present invention is mapped and localized to chromosome position 6q25-26, a region shown to be a site of loss of heterozygosity in human ovarian, breast and gastric cancers hepatocarcinomas, Burkitt's lymphomas, gliomas, and parathyroid adenomas. The FBP7 nucleic acid of the present invention is mapped and localized to chromosome position 15q 5 a region which contains a tumor suppressor gene associated with progression to a metastatic stage in breast and colon cancers and a loss of heterozygosity in parathyroid adenomas.

[0228] The molecules of the present invention can be used in assays, such as immunoassays, to detect, prognose, diagnose, or monitor various conditions, diseases, and disorders

affecting FBP expression, or monitor the treatment thereof. In particular, such an immunoassay is carried out by a method comprising contacting a sample derived from a patient with an anti-FBP antibody under conditions such that immunospecific binding can occur, and detecting or measuring the amount of any immunospecific binding by the antibody. In a specific aspect, such binding of antibody, in tissue sections, can be used to detect aberrant FBP localization or aberrant (e.g., low or absent) levels of FBP. In a specific embodiment, antibody to FBP can be used to assay a patient tissue or serum sample for the presence of FBP where an aberrant level of FBP is an indication of a diseased condition. By "aberrant levels," is meant increased or decreased levels relative to that present, or a standard level representing that present, in an analogous sample from a portion of the body or from a subject not having the disorder.

[0229] The immunoassays which can be used include but are not limited to competitive and non-competitive assay systems using techniques such as western blots, immunohistochemistry radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few.

[0230] FBP genes and related nucleic acid sequences and subsequences, including complementary sequences, can also be used in hybridization assays. FBP nucleic acid sequences, or subsequences thereof comprising about at least 8 nucleotides, can be used as hybridization probes. Hybridization assays can be used to detect, prognose, diagnose, or monitor conditions, disorders, or disease states associated with aberrant changes in FBP expression and/or activity as described supra. In particular, such a hybridization assay is carried out by a method comprising contacting a sample containing nucleic acid with a nucleic acid probe capable of hybridizing to FBP DNA or RNA, under conditions such that hybridization can occur, and detecting or measuring any resulting hybridization.

[0231] In specific embodiments, diseases and disorders involving overproliferation of cells can be diagnosed, or their suspected presence can be screened for, or a predisposition to develop such disorders can be detected, by detecting decreased levels of FBP protein, FBP RNA, or FBP functional activity (e.g., ubiquitin ligase target binding activity, F-box domain binding activity, ubiquitin ligase activity etc.), or by detecting mutations in FBP RNA, DNA or FBP protein (e.g., translocations in FBP nucleic acids, truncations in the FBP gene or protein, changes in nucleotide or amino acid sequence relative to wild-type FBP) that cause decreased expression or activity of FBP. Such diseases and disorders include but are not limited to those described in Section 5.7.3. By way of example, levels of FBP protein can be detected by immunoassay, levels of FBP RNA can be detected by hybridization assays (e.g., Northern blots, in situ-hybridization), FBP activity can be assayed by measuring ubiquitin ligase activity in E3 ubiquitin ligase complexes formed in vivo or in vitro, F-box domain binding activity can be assayed by measuring binding to Skp1 protein by binding assays commonly known in the art. translocations, deletions and point mutations in FBP nucleic acids can be detected by Southern blotting, FISH, RFLP analysis, SSCP,

PCR using primers that preferably generate a fragment spanning at least most of the FBP gene, sequencing of FBP genomic DNA or cDNA obtained from the patient, etc.

[0232] In a preferred embodiment, levels of FBP mRNA or protein in a patient sample are detected or measured, in which decreased levels indicate that the subject has, or has a predisposition to developing, a malignancy or hyperproliferative disorder; in which the decreased levels are relative to the levels present in an analogous sample from a portion of the body or from a subject not having the malignancy or hyperproliferative disorder, as the case may be.

[0233] In another specific embodiment, diseases and disorders involving a deficiency in cell proliferation or in which cell proliferation is desirable for treatment, are diagnosed, or their suspected presence can be screened for, or a predisposition to develop such disorders can be detected, by detecting increased levels of FBP protein, FBP RNA, or FBP functional activity (e.g., ubiquitin ligase activity, Skp1 binding activity, etc.), or by detecting mutations in FBP RNA, DNA or protein (e.g., translocations in FBP nucleic acids, truncations in the gene or protein, changes in nucleotide or amino acid sequence relative to wild-type FBP) that cause increased expression or activity of FBP. Such diseases and disorders include but are not limited to those described in Section 5.7.3. By way of example, levels of FBP protein, levels of FBP RNA, ubiquitin ligase activity, FBP binding activity, and the presence of translocations or point mutations can be determined as described above.

[0234] In a specific embodiment, levels of FBP mRNA or protein in a patient sample are detected or measured, in which increased levels indicate that the subject has, or has a predisposition to developing, a growth deficiency or degenerative or hypoproliferative disorder; in which the increased levels are relative to the levels present in an analogous sample from a portion of the body or from a subject not having the growth deficiency, degenerative, or hypoproliferative disorder, as the case may be.

[0235] Kits for diagnostic use are also provided, that comprise in one or more containers an anti-FBP antibody, and, optionally, a labeled binding partner to the antibody. Alternatively, the anti-FBP antibody can be labeled (with a detectable marker, e.g., a chemiluminescent, enzymatic, fluorescent, or radioactive moiety). A kit is also provided that comprises in one or more containers a nucleic acid probe capable of hybridizing to FBP RNA. In a specific embodiment, a kit can comprise in one or more containers a pair of primers (e.g., each in the size range of 6-30 nucleotides) that are capable of priming amplification [e.g., by polymerase chain reaction (see e.g., Innis et al., 1990, PCR Protocols, Academic Press, Inc., San Diego, Calif.), ligase chain reaction (see EP 320,308) use of Q replicase, cyclic probe reaction, or other methods known in the art] under appropriate reaction conditions of at least a portion of a FBP nucleic acid. A kit can optionally further comprise in a container a predetermined amount of a purified FBP protein or nucleic acid, e.g., for use as a standard or control.

[0236] 5.7 Methods and Compositions for Therapeutic use of F-Box Proteins, Derivatives, and Modulators

[0237] Described below are methods and compositions for the use of F-box proteins in the treatment of proliferative disorders and oncogenic disease symptoms may be amelio-

rated by compounds that activate or enhance FBP activity, and whereby proliferative disorders and cancer may be ameliorated.

[0238] In certain instances, compounds and methods that increase or enhance the activity of an FBP can be used to treat proliferative and oncogenic disease symptoms. Such a case may involve, for example, a proliferative disorder that is brought about, at least in part, by a reduced level of FBP gene expression, or an aberrant level of an FBP gene product's activity. For example, decreased activity or under-expression of an FBP component of a ubiquitin ligase complex whose substrate is a positive cell-cycle regulator, such as a member of the Cyclin family, will result in increased cell proliferation. As such, an increase in the level of gene expression and/or the activity of such FBP gene products would bring about the amelioration of proliferative disease symptoms.

[0239] In another instance, compounds that increase or enhance the activity of an FBP can be used to treat proliferative and oncogenic disease symptoms resulting from defects in the expression or activity of other genes and gene products involved in cell cycle control, such as FBP substrate molecules. For example, an increase in the expression or activity of a positive cell-cycle positive molecule, such as a member of the Cyclin family, may result in its over-activity and thereby lead to increased cell proliferation. Compounds that increase the expression or activity of the FBP component of a ubiquitin ligase complex whose substrate is such a cell-cycle positive regulator will lead to ubiquitination of the defective molecule, and thereby result in an increase in its degradation. Disease symptoms resulting from such a defect may be ameliorated by compounds that compensate the disorder by increased FBP activity. Techniques for increasing FBP gene expression levels or gene product activity levels are discussed in Section 5.7, below.

[0240] Alternatively, compounds and methods that reduce or inactivate FBP activity may be used therapeutically to ameliorate proliferative and oncogenic disease symptoms. For example, a proliferative disorder may be caused, at least in part, by a defective FBP gene or gene product that leads to its overactivity. Where such a defective gene product is a component of a ubiquitin ligase complex whose target is a cell-cycle inhibitor molecule, such as a Cki, an overactive FBP will lead to a decrease in the level of cell-cycle molecule and therefore an increase in cell proliferation. In such an instance, compounds and methods that reduce or inactivate FBP function may be used to treat the disease symptoms.

[0241] In another instance, compounds and methods that reduce the activity of an FBP can be used to treat disorders resulting from defects in the expression or activity of other genes and gene products involved in cell cycle control, such as FBP substrate molecules. For example, a defect in the expression or activity of a cell-cycle negative regulatory molecule, such as a Cki, may lead to its under-activity and thereby result in increased cell proliferation. Reduction in the level and/or activity of an FBP component whose substrate was such molecule would decrease the ubiquitination and thereby increase the level of such a defective molecule. Therefore, compounds and methods aimed at reducing the expression and/or activity of such FBP mol-

ecules could thereby be used in the treatment of disease symptoms by compensating for the defective gene or gene product.

[0242] Techniques for the reduction of target gene expression levels or target gene product activity levels are discussed in Section 5.7 below.

[0243] 5.7.1 Therapeutic Use of Inhibitory Antisense, Ribozyme and Triple Helix Molecules and Identified Agonists and Antagonists

[0244] In another embodiment, symptoms of certain FBP disorders, such as such as proliferative or differentiative disorders causing tumorigenesis or cancer, may be ameliorated by decreasing the level of FBP gene expression and/or FBP gene product activity by using FBP gene sequences in conjunction with well-known antisense, gene "knock-out" ribozyme and/or triple helix methods to decrease the level of FBP gene expression. Among the compounds that may exhibit the ability to modulate the activity, expression or synthesis of the FBP gene, including the ability to ameliorate the symptoms of an FBP disorder, such as cancer, are antisense, ribozyme, and triple helix molecules. Such molecules may be designed to reduce or inhibit either unimpaired, or if appropriate, mutant target gene activity. Techniques for the production and use of such molecules are well known to those of skill in the art. For example, antisense targeting SKP2 mRNA stabilize the Skp2-substrate p27, as described in Section X (Figure X).

[0245] Antisense RNA and DNA molecules act to directly block the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense approaches involve the design of oligonucleotides that are complementary to a target gene mRNA. The antisense oligonucleotides will bind to the complementary target gene mRNA transcripts and prevent translation. Absolute complementarity, although preferred, is not required.

[0246] A sequence "complementary" to a portion of an RNA, as referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double-stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with an RNA it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

[0247] In one embodiment, oligonucleotides complementary to non-coding regions of the FBP gene could be used in an antisense approach to inhibit translation of endogenous FBP mRNA. Antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

[0248] In an embodiment of the present invention, oligonucleotides complementary to the nucleic acids encoding the F-box motif as indicated in **FIGS. 2 and 4-9**.

[0249] Regardless of the choice of target sequence, it is preferred that in vitro studies are first performed to quantify the ability of the antisense oligonucleotide to inhibit gene expression. It is preferred that these studies utilize controls that distinguish between antisense gene inhibition and nonspecific biological effects of oligonucleotides. It is also preferred that these studies compare levels of the target RNA or protein with that of an internal control RNA or protein. Additionally, it is envisioned that results obtained using the antisense oligonucleotide are compared with those obtained using a control oligonucleotide. It is preferred that the control oligonucleotide is of approximately the same length as the test oligonucleotide and that the nucleotide sequence of the oligonucleotide differs from the antisense sequence no more than is necessary to prevent specific hybridization to the target sequence.

[0250] The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors in vivo), or agents facilitating transport across the cell membrane (see, e.g. Letsinger, et al., 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86, 6553-6556; Lemaire, et al., 1987, *Proc. Natl. Acad. Sci.* 84, 648-652; PCT Publication No. WO88/09810, published Dec. 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication No. WO89/10134, published Apr. 25, 1988), hybridization-triggered cleavage agents (see, e.g., Krol et al., 1988, *BioTechniques* 6, 958-976) or intercalating agents (see, e.g., Zon, 1988, *Pharm. Res.* 5, 539-549). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

[0251] The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

[0252] The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

[0253] In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group consisting of a phosphorothio-

ate (S-ODNs), a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

[0254] In yet another embodiment, the antisense oligonucleotide is an -anomeric oligonucleotide. An -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual -units, the strands run parallel to each other (Gautier, et al., 1987, *Nucl. Acids Res.* 15, 6625-6641). The oligonucleotide is a 2-O-methylribonucleotide (Inoue, et al., 1987, *Nucl. Acids Res.* 15, 6131-6148), or a chimeric RNA-DNA analogue (Inoue, et al., 1987, *FEBS Lett.* 215, 327-330).

[0255] Oligonucleotides of the invention may be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein, et al. (1988, *Nucl. Acids Res.* 16, 3209), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin, et al., 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85, 7448-7451), etc.

[0256] While antisense nucleotides complementary to the target gene coding region sequence could be used, those complementary to the transcribed, untranslated region are most preferred.

[0257] In one embodiment of the present invention, gene expression downregulation is achieved because specific target mRNAs are digested by RNase H after they have hybridized with the antisense phosphorothioate oligonucleotides (S-ODNs). Since no rules exist to predict which antisense S-ODNs will be more successful, the best strategy is completely empirical and consists of trying several antisense S-ODNs. Antisense phosphorothioate oligonucleotides (S-ODNs) will be designed to target specific regions of mRNAs of interest. Control S-ODNs consisting of scrambled sequences of the antisense S-ODNs will also be designed to assure identical nucleotide content and minimize differences potentially attributable to nucleic acid content. All S-ODNs will be synthesized by Oligos Etc. (Wilsonville, Oreg.). In order to test the effectiveness of the antisense molecules when applied to cells in culture, such as assays for research purposes or ex vivo gene therapy protocols, cells will be grown to 60-80% confluence on 100 mm tissue culture plates, rinsed with PBS and overlaid with lipofection mix consisting of 8 ml Opti-MEM, 52.81 Lipofectin, and a final concentration of 200 nM S-ODNs. Lipofections will be carried out using Lipofectin Reagent and Opti-MEM (Gibco BRL). Cells will be incubated in the presence of the lipofection mix for 5 hours. Following incubation the medium will be replaced with complete DMEM. Cells will be harvested at different time points post-lipofection and protein levels will be analyzed by Western blot.

[0258] Antisense molecules should be targeted to cells that express the target gene, either directly to the subject in vivo or to cells in culture, such as in ex vivo gene therapy protocols. A number of methods have been developed for delivering antisense DNA or RNA to cells; e.g., antisense molecules can be injected directly into the tissue site, or modified antisense molecules, designed to target the desired cells (e.g., antisense linked to peptides or antibodies that

specifically bind receptors or antigens expressed on the target cell surface) can be administered systemically.

[0259] However, it is often difficult to achieve intracellular concentrations of the antisense sufficient to suppress translation of endogenous mRNAs. Therefore a preferred approach utilizes a recombinant DNA construct in which the antisense oligonucleotide is placed under the control of a strong pol III or pol II promoter. The use of such a construct to transfect target cells in the patient will result in the transcription of sufficient amounts of single stranded RNAs that will form complementary base pairs with the endogenous target gene transcripts and thereby prevent translation of the target gene mRNA. For example, a vector can be introduced e.g., such that it is taken up by a cell and directs the transcription of an antisense RNA. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in mammalian cells. Expression of the sequence encoding the antisense RNA can be by any promoter known in the art to act in mammalian, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include but are not limited to: the SV40 early promoter region (Bemoist and Chambon, 1981, *Nature* 290, 304-310), the promoter contained in the 3 long terminal repeat of Rous sarcoma virus (Yamamoto, et al., 1980, *Cell* 22, 787-797), the herpes thymidine kinase promoter (Wagner, et al., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78, 1441-1445), the regulatory sequences of the metallothionein gene (Brinster, et al., 1982, *Nature* 296, 39-42), etc. Any type of plasmid, cosmid, YAC or viral vector can be used to prepare the recombinant DNA construct which can be introduced directly into the tissue site. Alternatively, viral vectors can be used that selectively infect the desired tissue, in which case administration may be accomplished by another route (e.g., systemically).

[0260] Ribozyme molecules designed to catalytically cleave target gene mRNA transcripts can also be used to prevent translation of target gene mRNA and, therefore, expression of target gene product (see, e.g., PCT International Publication WO90/11364, published Oct. 4, 1990; Sarver, et al., 1990, *Science* 247, 1222-1225). In an embodiment of the present invention, oligonucleotides which hybridize to the FBP gene are designed to be complementary to the nucleic acids encoding the F-box motif as indicated in **FIGS. 2 and 4-9**.

[0261] Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. (For a review, see Rossi, 1994, *Current Biology* 4, 469-471). The mechanism of ribozyme action involves sequence specific hybridization of the ribozyme molecule to complementary target RNA, followed by an endonucleolytic cleavage event. The composition of ribozyme molecules must include one or more sequences complementary to the target gene mRNA, and must include the well known catalytic sequence responsible for mRNA cleavage. For this sequence, see, e.g., U.S. Pat. No. 5,093,246, which is incorporated herein by reference in its entirety.

[0262] While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy target gene

mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Myers, 1995, *Molecular Biology and Biotechnology: A Comprehensive Desk Reference*, VCH Publishers, New York, (see especially **FIG. 4**, page 833) and in Haseloff & Gerlach, 1988, *Nature*, 334, 585-591, which is incorporated herein by reference in its entirety.

[0263] Preferably the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the target gene mRNA, i.e., to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

[0264] The ribozymes of the present invention also include RNA endoribonucleases (hereinafter "Cech-type ribozymes") such as the one that occurs naturally in *Tetrahymena thermophila* (known as the IVS, or L-19 IVS RNA) and that has been extensively described by Thomas Cech and collaborators (Zaug, et al., 1984, *Science*, 224, 574-578; Zaug and Cech, 1986, *Science*, 231, 470-475; Zaug, et al., 1986, *Nature*, 324, 429-433; published International patent application No. WO 88/04300 by University Patents Inc.; Been & Cech, 1986, *Cell*, 47, 207-216). The Cech-type ribozymes have an eight base pair active site which hybridizes to a target RNA sequence whereafter cleavage of the target RNA takes place. The invention encompasses those Cech-type ribozymes which target eight base-pair active site sequences that are present in the target gene.

[0265] As in the antisense approach, the ribozymes can be composed of modified oligonucleotides (e.g., for improved stability, targeting, etc.) and should be delivered to cells that express the target gene in vivo. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous target gene messages and inhibit translation. Because ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

[0266] Endogenous target gene expression can also be reduced by inactivating or "knocking out" the target gene or its promoter using targeted homologous recombination (e.g., see Smithies, et al., 1985, *Nature* 317, 230-234; Thomas & Capecchi, 1987, *Cell* 51, 503-512; Thompson, et al., 1989, *Cell* 5, 313-321; each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional target gene (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous target gene (either the coding regions or regulatory regions of the target gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express the target gene in vivo. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the target gene. Such approaches are particularly suited modifications to ES (embryonic stem) cells can be used to generate animal offspring with an inactive target gene (e.g., see Thomas & Capecchi, 1987 and Thompson,

1989, supra). However this approach can be adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site in vivo using appropriate viral vectors.

[0267] Alternatively, endogenous target gene expression can be reduced by targeting deoxyribonucleotide sequences complementary to the regulatory region of the target gene (i.e., the target gene promoter and/or enhancers) to form triple helical structures that prevent transcription of the target gene in target cells in the body. (See generally, Helene, 1991, *Anticancer Drug Des.*, 6(6), 569-584; Helene, et al., 1992, *Ann. N.Y. Acad. Sci.*, 660, 27-36; and Maher, 1992, *Bioassays* 14(12), 807-815).

[0268] Nucleic acid molecules to be used in triple helix formation for the inhibition of transcription should be single stranded and composed of deoxynucleotides. The base composition of these oligonucleotides must be designed to promote triple helix formation via Hoogsteen base pairing rules, which generally require sizeable stretches of either purines or pyrimidines to be present on one strand of a duplex. Nucleotide sequences may be pyrimidine-based, which will result in TAT and CGC+ triplets across the three associated strands of the resulting triple helix. The pyrimidine-rich molecules provide base complementarity to a purine-rich region of a single strand of the duplex in a parallel orientation to that strand. In addition, nucleic acid molecules may be chosen that are purine-rich, for example, contain a stretch of G residues. These molecules will form a triple helix with a DNA duplex that is rich in GC pairs, in which the majority of the purine residues are located on a single strand of the targeted duplex, resulting in GGC triplets across the three strands in the triplex.

[0269] Alternatively, the potential sequences that can be targeted for triple helix formation may be increased by creating a so called "switchback" nucleic acid molecule. Switchback molecules are synthesized in an alternating 5'-3',3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex.

[0270] In instances wherein the antisense, ribozyme, and/or triple helix molecules described herein are utilized to inhibit mutant gene expression, it is possible that the technique may so efficiently reduce or inhibit the transcription (triple helix) and/or translation (antisense, ribozyme) of mRNA produced by normal target gene alleles that the possibility may arise wherein the concentration of normal target gene product present may be lower than is necessary for a normal phenotype. In such cases, to ensure that substantially normal levels of target gene activity are maintained, therefore, nucleic acid molecules that encode and express target gene polypeptides exhibiting normal target gene activity may, be introduced into cells via gene therapy methods such as those described, below, in Section 5.7.2 that do not contain sequences susceptible to whatever antisense, ribozyme, or triple helix treatments are being utilized. Alternatively, in instances whereby the target gene encodes an extracellular protein, it may be preferable to co-administer normal target gene protein in order to maintain the requisite level of target gene activity.

[0271] Anti-sense RNA and DNA, ribozyme, and triple helix molecules of the invention may be prepared by any

method known in the art for the synthesis of DNA and RNA molecules, as discussed above. These include techniques for chemically synthesizing oligodeoxyribonucleotides and oligoribonucleotides well known in the art such as for example solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences may be incorporated into a wide variety of vectors that incorporate suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters. Alternatively, antisense cDNA constructs that synthesize anti sense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines.

[0272] 5.7.2 Gene Replacement Therapy

[0273] With respect to an increase in the level of normal FBP gene expression and/or FBP gene product activity, FBP gene nucleic acid sequences, described, above, in Section 5.1 can, for example, be utilized for the treatment of proliferative disorders such as cancer. Such treatment can be administered, for example, in the form of gene replacement therapy. Specifically, one or more copies of a normal FBP gene or a portion of the FBP gene that directs the production of an FBP gene product exhibiting normal FBP gene function, may be inserted into the appropriate cells within a patient, using vectors that include, but are not limited to adenovirus, adeno-associated virus, and retrovirus vectors, in addition to other particles that introduce DNA into cells, such as liposomes.

[0274] For FBP genes that are expressed in all tissues or are preferentially expressed, such as FBP1 gene is expressed preferably in the brain, such gene replacement therapy techniques should be capable delivering FBP gene sequences to these cell types within patients. Thus, in one embodiment, techniques that are well known to those of skill in the art (see, e.g., PCT Publication No. WO89/10134, published Apr. 25, 1988) can be used to enable FBP gene sequences to cross the blood-brain barrier readily and to deliver the sequences to cells in the brain. With respect to delivery that is capable of crossing the blood-brain barrier, viral vectors such as, for example, those described above, are preferable.

[0275] In another embodiment, techniques for delivery involve direct administration of such FBP gene sequences to the site of the cells in which the FBP gene sequences are to be expressed.

[0276] Additional methods that may be utilized to increase the overall level of FBP gene expression and/or FBP gene product activity include the introduction of appropriate FBP-expressing cells, preferably autologous cells, into a patient at positions and in numbers that are sufficient to ameliorate the symptoms of an FBP disorder. Such cells may be either recombinant or non-recombinant.

[0277] Among the cells that can be administered to increase the overall level of FBP gene expression in a patient are cells that normally express the FBP gene.

[0278] Alternatively, cells, preferably autologous cells, can be engineered to express FBP gene sequences, and may then be introduced into a patient in positions appropriate for the amelioration of the symptoms of an FBP disorder or a proliferative or differentiative disorders, e.g., cancer and

tumorigenesis. Alternately, cells that express an unimpaired FBP gene and that are from a MHC matched individual can be utilized, and may include, for example, brain cells. The expression of the FBP gene sequences is controlled by the appropriate gene regulatory sequences to allow such expression in the necessary cell types. Such gene regulatory sequences are well known to the skilled artisan. Such cell-based gene therapy techniques are well known to those skilled in the art, see, e.g., Anderson, U.S. Pat. No. 5,399,349.

[0279] When the cells to be administered are non-autologous cells, they can be administered using well known techniques that prevent a host immune response against the introduced cells from developing. For example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

[0280] Additionally, compounds, such as those identified via techniques such as those described, above, in Section 5.5, that are capable of modulating FBP gene product activity can be administered using standard techniques that are well known to those of skill in the art. In instances in which the compounds to be administered are to involve an interaction with brain cells, the administration techniques should include well known ones that allow for a crossing of the blood-brain barrier.

[0281] 5.7.3 Target Proliferative Cell Disorders

[0282] With respect to specific proliferative and oncogenic disease associated with ubiquitin ligase activity, the diseases that can be treated or prevented by the methods of the present invention include but are not limited to: human sarcomas and carcinomas, e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, retinoblastoma; leukemias, e.g., acute lymphocytic leukemia and acute myelocytic leukemia (myeloblastic, promyelocytic, myelomonocytic, monocytic and erythroleukemia); chronic leukemia (chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia); and polycythemia vera, lymphoma (Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, and heavy chain disease.

[0283] Diseases and disorders involving a deficiency in cell proliferation or in which cell proliferation is desired for treatment or prevention, and that can be treated or prevented

by inhibiting FBP function, include but are not limited to degenerative disorders, growth deficiencies, hypoproliferative disorders, physical trauma, lesions, and wounds; for example, to promote wound healing, or to promote regeneration in degenerated, lesioned or injured tissues, etc. In a specific embodiment, nervous system disorders are treated. In another specific embodiment, a disorder that is not of the nervous system is treated.

[0284] 5.8 Pharmaceutical Preparations and Methods of Administration

[0285] The compounds that are determined to affect FBP gene expression or gene product activity can be administered to a patient at therapeutically effective doses to treat or ameliorate a cell proliferative disorder. A therapeutically effective dose refers to that amount of the compound sufficient to result in amelioration of symptoms of such a disorder.

[0286] 5.8.1 Effective Dose

[0287] Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds that exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

[0288] The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (i.e., the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0289] 5.8.2 Formulations and Use

[0290] Pharmaceutical compositions for use in accordance with the present invention may be formulated in conventional manner using one or more physiologically acceptable carriers or excipients.

[0291] Thus, the compounds and their physiologically acceptable salts and solvates may be formulated for administration by inhalation or insufflation (either through the mouth or the nose) or oral, buccal, parenteral or rectal administration.

[0292] For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavoring, coloring and sweetening agents as appropriate.

[0293] Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

[0294] For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

[0295] For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g., gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[0296] The compounds may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multidose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

[0297] The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

[0298] In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or

intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[0299] The compositions may, if desired, be presented in a pack or dispenser device that may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration.

6. EXAMPLE

Identification and Characterization of Novel Ubiquitin Ligase F-Box Proteins and Genes

[0300] The following studies were carried out to identify novel F-box proteins which may act to recruit novel specific substrates to the ubiquitination pathways. Studies involving several organisms have shown that some FBPs play a crucial role in the controlled degradation of important cellular regulatory proteins (e.g., cyclins, cdk-inhibitors, β -catenin, IKBa, etc.). These FBPs are subunits of ubiquitin protein SCF ligases formed by three basic subunits: a cullin subunit (called Cdc53 in *S. cerevisiae* and Cull in humans); Skp1; and one of many FBPs. SCF ligases target ubiquitin conjugating enzymes (either Ubc3 or Ubc4) to specific substrates which are recruited by different FBPs. Schematically, the Ubc is bound to the ligase through the cullin subunit while the substrate interacts with the FBP subunit. Although FBPs can bind the cullin subunit directly, the presence of fourth subunit, Skp1, which simultaneously can bind the cullin N-terminus and the F-box of the FBP, stabilizes the complex. Thus, the substrate specificity of the ubiquitin ligase complex is provided by the F-box subunit.

[0301] 6.1 Materials and Methods used for the Identification and Characterization of Novel F-Box Genes

[0302] Yeast Two-Hybrid Screening In order to clone the human genes encoding F-box proteins, proteins associated with Skp1 were identified using a modified yeast 2-hybrid system (Vidal et al., 1996, Proc. Nat. Acad. Sci., 93:10315-20; Vidal et al., 1996, Proc. Nat. Acad. Sci., 93:10321-26). This modified system takes advantage of using three reporter genes expressed from three different Gal4 binding site promoters, thereby decreasing the number of false positive interactions. This multiple reporter gene assay facilitates identification of true interactors.

[0303] Human Skp 1 was used as a bait to search for proteins that interact with Skp 1, such as novel F-box proteins and the putative human homolog of Cdc4. The plasmids pPC97—CYH2 and pPC86 plasmids, encoding the DNA binding domain (DB, aa 1-147) and the transcriptional activation domain (AD, aa 768-881) of yeast GAL4, and containing LEU2 and TRP I as selectable markers, respectively, were used (Chevray and Nathans, 1992, Proc. Nat. Acad. Sci., 89:5789-93; Vidal et al., supra).

[0304] An in-frame fusion between Skp1 and DB was obtained by homologous recombination of the PCR product described below. The following 2 oligonucleotides were designed and obtained as purified primers from Gene Link

Inc.: 5'-AGT-AGT-AAC-AAA-GGT-CAA-AGA-CAG-TTG-ACT-GTA-TCG-TCG-AGG-ATG-CCT-TCA-ATT-AAG-TT (SEQ ID NO: 80); 3'-GCG-GTT-ACT-TAC-TTA-GAG-CTC-GAC-GTC-TTA-CTT-ACT-TAG-CTC-ACT-TCT-CTT-CAC-ACC-A (SEQ ID NO: 81). The 5' primer corresponds to a sequence located in the DB of the pPC97-CYH2 plasmid (underlined) flanked by the 5' sequence of the *skp1* gene. The 3' primer corresponds to a sequence located by polylinker of the pPC97-CYH2 plasmid (underlined) flanked by the 3' sequence of the *skp1* gene. These primers were used in a PCR reaction containing the following components: 100 ng DNA template (*skp1* pET plasmid), 1 μ M of each primer, 0.2 mM dNTP, 2 mM MgCl₂, 10 mM KCl, 20 mM Tris Cl pH 8.0, 0.1% Triton X-100, 6 mM (NH₄)₂SO₄, 10 μ g/ml nuclease-free BSA, 1 unit of Pfu DNA polymerase (4' at 94° C., 1' at 50° C, 10' at 72° C. for 28 cycles). Approximately 100 ng of PCR product were transformed into yeast cells (MaV103 strain; Vidal et al., 1996, Proc. Natl. Acad. Sci. U.S.A. 93:10315-10320; Vidal et al., 1996, Proc. Natl. Acad. Sci. U.S.A. 93:10321-10326) in the presence or in the absence of 100 ng of pPC97-CYH2 plasmid previously digested with BglI and SalI. As a result of the homologous recombination, only yeast cells containing the pPC97-CYH2 plasmid homologously recombined with *skp1* cDNA, grew in the absence of leucine. Six colonies were isolated and analyzed by immunoblotting for the expression of Skp1, as described (Vidal et al., supra). All 6 colonies, but not control colonies, expressed a Mr 36,000 fusion-protein that was recognized by our affinity purified anti-Skp1 antibody.

[0305] The AD fusions were generated by cloning cDNA fragments in the frame downstream of the AD domains and constructs were confirmed by sequencing, immunoblot, and interaction with Skp 1. The pPC86-Skp2s (pPC86) include: pPC86-Skp2, and pPC86-Skp2-CT (aa 181-435 of Skp2). The first fusion represents our positive control since Skp2 is a known interactor of Skp1 (Zhang, et al, 1995, Cell, 82: 915-25); the latter fusion was used as a negative control since it lacked the F-box required for the interaction with Skp1.

[0306] MaV103 strain harboring the DB-*skp1* fusions was transformed with an activated T-cell cDNA library (Alala 2; Hu, et al., Genes & Dev. 11: 2701-14) in pPC86 using the standard lithium acetate method. Transformants were first plated onto synthetic complete (SC)-Leu-Trp plates, followed by replica plating onto (SC)-Leu-Trp-His plates containing 20 mM 3-aminotriazole (3-AT) after 2 days. Yeast colonies grown out after additional 34 days of incubation were picked as primary positives and further tested in three reporter assays: i) growth on SC-Leu-Trp-His plates supplemented with 20 mM 3-AT; ii)-galactosidase activity; and iii) URA3 activation on SC-Leu-Trp plates containing 0.2% 5-fluoroorotic acid, as a counterselection method. Of the 3 \times 10⁶ yeast transformants screened AD plasmids were rescued from the fifteen selected positive colonies after all three. MaV103 cells were re-transformed with either rescued AD plasmids and the DBskp1 fusion or rescued AD plasmid and the pPC97-CYH2vector without a cDNA insert as control. Eleven AD plasmids from colonies that repeatedly tested positive in all three reporter assays (very strong interactors) and four additional AD plasmids from clones that were positive on some but not all three reporter assays (strong interactors) were recovered and sequenced with the automated ABI 373 DNA sequencing system.

[0307] Cloning of full length FBPs Two of the clones encoding FBP4 and FBP5 appeared to be full-length, while full length clones of 4 other cDNAs encoding FBP1, FBP2, FBP3 and FBP7 were obtained with RACE using Marathon-Ready cDNA libraries (Clontech, cat. # 7406, 7445, 7402) according to manufacturer's instructions. A full-length clone encoding FBP6 was not obtained. Criteria for full length clones included at least two of the following: i) the identification of an ORF yielding a sequence related to known F-box proteins; ii) the presence of a consensus Kozak translation initiation sequence at a putative initiator methionine codon; iii) the identification of a stop codon in the same reading frame but upstream of the putative initiation codon; iv) the inability to further increase the size of the clone by RACE using three different cDNA libraries.

[0308] Analysis by Immunoblotting of Protein from Yeast Extracts Yeast cells were grown to mid-logarithmic phase, harvested, washed and resuspended in buffer (50 mM Tris pH 8.0, 20% glycerol, 1 mM EDTA, 0.1% Triton X-100, 5 mM MgCl₂, 10 mM β -mercaptoethanol, 1 mM PMSF, 1 mg/ml Leupeptin, 1 mg/ml Pepstatin) at a cell density of about 109 cells/ml. Cells were disrupted by vortexing in the presence of glass beads for 10 min at 40C. Debris was pelleted by centrifugation at 12,000 RPM for 15 min at 40C. Approximately 50 g of proteins were subjected to immunoblot analysis as described (Vidal et al., 1996a, supra; Vidal et al., 1996b, supra).

[0309] DNA database searches and analysis of protein motifs ESTs (expressed sequence tags) with homology to FBP genes were identified using BLAST, PSI-BLAST (<http://www.ncbi.nlm.nih.gov/BLAST/>) and TGI Sequence Search (http://www.tigr.org/cgi-bin/BlastSearch/blast_tgi.cgi). ESTs that overlapped more than 95% in at least 100 bps were assembled into novel contiguous ORFs using Sequencher 3.0. Protein domains were identified with ProfileScan Server (http://www.isrec.isb-sib.ch/software/PFS-CAN_form.html), BLOCKS Sercher (http://www.blocks.fhcrc.org/blocks_search.html) and IMB Jena (<http://genome.imb-ena.de/cgi-bin/GDEWWW/menu.cgi>).

[0310] Construction of F-box mutants Delta-F-box mutants [(Δ F)FBP1, residues 32-179; (Δ F)FBP2, residues 60-101; (Δ F)FBP3a, residues 40-76; (Δ F)FBP4, residues 55-98] were obtained by deletion with the appropriate restriction enzymes with conservation of the reading frame. (Δ F)Skp2 mutant was obtained by removing a DNA fragment (nucleotides 338-997) with BspEI and XbaI restriction enzymes, and replacing it with a PCR fragment containing nucleotides 457 to 997. The final construct encoded a protein lacking residues 113-152. The leucine 51-to-alanine FBP3a mutant [FBP3a(L51A)] and the tryptophan 76-to-alanine FBP3a mutant [FBP3a(W76A)] were generated by oligonucleotide-directed mutagenesis using the polymerase chain reaction of the QuikChange site-directed mutagenesis kit (Stratagene). All mutants were sequenced in their entirety.

[0311] Recombinant proteins cDNA fragments encoding the following human proteins: Flag-tagged FBP1, Flag-tagged (Δ F)FBP1, Flag-tagged FBP3a, Skp2, HA-tagged Cull, HA-tagged Cul2, (β -catenin, His-tagged cyclin D1, Skp1, His-tagged Skp1, His-tagged Elongin C were inserted into the baculovirus expression vector pBacpak-8 (Clontech) and cotransfected into Sf9 cells with linearized baculovirus DNA using the BaculoGold transfection kit

(Pharrningen). Recombinant viruses were used to infect 5B cells and assayed for expression of their encoded protein by immunoblotting as described above. His-proteins were purified with Nickel-agarose (Invitrogen) according to the manufacturer's instructions.

[0312] Antibodies. Anti-Cull antibodies was generated by injecting rabbits and mice with the following amino acid peptide: (C)DGEKDTYSYLA (SEQ ID NO: 82). This peptide corresponds to the carboxy-terminus of human Cull and is not conserved in other cullins. Anti-Cu12 antibodies was generated by injecting rabbits with the following amino acid peptide: (C)ESSFSLNMNFSSKRTKFKITSMQ (SEQ ID NO: 83). This peptide is located 87 amino acids from the carboxy-terminus of human Cu12 and is not conserved in other cullins. The anti-Skp1 antibody was generated by injecting rabbits with the peptide (C)EEAQVRKENQW (SEQ ID NO: 84), corresponding to the carboxy-terminus of human Skp1. The cysteine residues (C) were added in order to couple the peptides to keyhole limpet hemocyanin (KLH). All of the antibodies were generated, affinity-purified (AP) and characterized as described (Pagano, M., ed., 1995, "From Peptide to Purified Antibody", in *Cell Cycle: Materials and Methods*, Springer-Verlag, 217-281). Briefly, peptides whose sequence showed high antigenic index (high hydrophilicity, good surface probability, good flexibility, and good secondary structure) were chosen. Rabbits and mice were injected with peptide-KLH mixed with complete Freund's adjuvant. Subsequently they were injected with the peptide in incomplete Freund's adjuvant, every 2 weeks, until a significant immunoreactivity was detected by immunoprecipitation of 35S-methionine labeled HeLa extract. These antisera recognized bands at the predicted size in both human extracts and a extracts containing recombinant proteins.

[0313] Monoclonal antibody (Mab) to Ubc3 was generated and characterized in collaboration with Zymed Inc. Mab to cyclin B (cat # sc-245) was from Santa Cruz; Mabs to p21 (cat # C24420) and p27 (cat # K25020) from Transduction lab. (Mabs) cyclin E, (Faha, 1993, *J. of Virology* 67: 2456); AP rabbit antibodies to human p27, Skp2, Cdk2 (Pagano, 1992, *EMBO J.* 11:761), and cyclinA (Pagano, 1992, *EMBO J.* 11:761), and phospho-site p27 specific antibody, were obtained or generated by standard methods. Where indicated, an AP goat antibody to an N-terminal Skp2 peptide (Santa Cruz, cat # sc-1567) was used. Rat anti-HA antibody was from Boehringer Mannheim (cat. #1867423), rabbit anti-HA antibody was from Santa Cruz (cat. # sc-805), mouse anti-Flag antibody was from Kodak (cat. # IB313010), rabbit anti-Flag antibody was from Zymed (cat. #71-5400), anti-Skp1 and anti-(β -catenin mouse antibodies were from Transduction Laboratories (cat. # C19220 and P46020, respectively). The preparation, purification and characterization of a Mab to human cyclin D1 (clone AM29, cat. #33-2500) was performed in collaboration with Zymed Inc. Antiserum to human cyclin D1 was produced as described (Ohtsubo et al., 1995, *Mol Cell Biol*, 15, 2612-2624).

[0314] Extract preparation and cell synchronization Protein extraction was performed as previously described (Pagano, 1993, *J. Cell Biol.* 121: 101) with the only difference that 1 μ M okadaic acid was present in the lysis buffer. Human lung fibroblasts IMR-90 were synchronized in G0/G1 by serum starvation for 48 hours and the restimulated

to re-enter the cell cycle by serum readdition. HeLa cells were synchronized by mitotic shake-off as described (Pagano, 1992, *EMBO J.* 11: 761). Synchronization was monitored by flow cytometry. For in vitro ubiquitination and degradation assays, G1 HeLa cells were obtained with a 48-hour lovastatin treatment and protein extraction performed as described below.

[0315] Immunoprecipitation and Immunoblotting. Cell extracts were prepared by addition of 3-5 volumes of standard lysis buffers (Pagano et al., 1992, *Science* 255, 1144-1147), and conditions for immunoprecipitation were as described (Jenkins and Xiong, 1995; Pagano et al., 1992a *Science* 255-1144-1147). Proteins were transferred from gel to a nitrocellulose membrane (Novex) by wet blotting as described (Tam et al., 1994 *Oncogene* 9,2663). Filters were subjected to immunoblotting using a chemiluminescence (DuPont-NEN) detection system according to the manufacturer's instructions

[0316] Protein extraction for in vitro ubiquitination assay Logarithmically growing, HeLa-S3 cells were collected at a density of 6×10^5 cells/ml. Approx. 4 ml of HeLa S3 cell pellet were suspended in 6 ml of ice-cold buffer consisting of 20 mM Tris-HCl (pH 7.2), 2 mM DTT, 0.25 mM EDTA, 10 μ g/ml leupeptin, and 10 μ g/ml pepstatin. The suspension was transferred to a cell nitrogen-disruption bomb (Parr, Moline, Ill., cat #4639) that had been rinsed thoroughly and chilled on ice before use. The bomb chamber was connected to a nitrogen tank and the pressure was brought slowly to 1000 psi. The chamber was left on ice under the same pressure for 30 minutes and then the pressure was released slowly. The material was transferred to an Eppendorf tube and centrifuged in a microcentrifuge at 10,000 g for 10 minutes. The supernatant (S-10) was divided into smaller samples and frozen at -80°C.

[0317] In vitro ubiquitination The ubiquitination assay was performed as described (Lyapina, 1998, *Proc Natl Acad Sci USA*, 95: 7451). Briefly, immuno-beads containing Flag-tagged FBPs immunoprecipitated with anti-Flag antibody were added with purified recombinant human E1 and E2 enzymes (Ubc2, Ubc3 or Ubc4) to a reaction mix containing biotinylated-ubiquitin. Samples were then analyzed by blotting with HRP-streptavidin. E1 and E2 enzymes and biotinylated-ubiquitin were produced as described (Pagano, 1995, *Science* 269: 682).

[0318] Transient transfections cDNA fragments encoding the following human proteins: FBP1, (Δ F)FBP1, FBP2, (Δ F)FBP2, FBP3a, (Δ F)FBP3a, FBP3a(L51A), FBP3a(W76A), FBP4, (Δ F)FBP4, Skp2, (Δ F)Skp2, HA-tagged β -catenin, untagged β -catenin, Skp1, cyclin D1 were inserted into the mammalian expression vector pcDNA3 (Invitrogen) in frame with a Flag-tag at their C-terminus. Cells were transfected with FuGENE transfection reagent (Boehringer, cat. #1-814-443) according to the manufacturer's instruction.

[0319] Immunofluorescence Transfected cell monolayers growing on glass coverslips were rinsed in PBS and fixed with 4% paraformaldehyde in PBS for 10 minutes at 4° C. followed by permeabilization for 10 minutes with 0.25% Triton X-100 in PBS. Other fixation protocols gave comparable results. Immunofluorescence stainings were performed using 1 μ g/ml rabbit anti-Flag antibody as described (Pagano, 1994, *Genes & Dev.*, 8:1627).

[0320] Northern Blot Analysis Northern blots were performed using human multiple-tissue mRNAs from Clontech Inc. Probes were radiolabeled with [α - 32 P] dCTP (Amersham Inc.) using a random primer DNA labeling kit (Gibco BRL) (2 \times 10⁶ cpm/ml). Washes were performed with 0.2 \times SSC, 0.1% SDS, at 55-60° C. FBP1 and FBP3a probes were two HindIII restriction fragments (nucleotides 1-571 and 1-450, respectively), FBP2, FBP4, and FBP1 probes were their respective full-length cDNAs, and P-ACTIN probe was from Clontech Inc.

[0321] Fluorescence in situ hybridization (FISH) Genomic clones were isolated by high-stringency screening (65° C., 0.2 \times SSC, 0.1% SDS wash) of a λ FIX II placenta human genomic library (Stratagene) with cDNA probes obtained from the 2-hybrid screening. Phage clones were confirmed by high-stringency Southern hybridization and partial sequence analysis. Purified whole phage DNA was labeled and FISH was performed as described (M. Pagano, ed., 1994, in *Cell Cycle: Materials and Methods*, 29).

[0322] 6.2 Results

[0323] 6.2.1 Characterization of novel F-box Proteins and their activity in vivo An improved version of the yeast two-hybrid system was used to search for interactors of human Skp1. The MaV103 yeast strain harboring the Gal4 DB-Skp1 fusion protein as bait was transformed with an activated T-cell cDNA library expressing Gal4 AD fusion proteins as prey. After initial selection and re-transformation steps, 3 different reporter assays were used to obtain 13 positive clones that specifically interact with human Skp1. After sequence analysis, the 13 rescued cDNAs were found to be derived from 7 different open reading frames all encoding FBPs. These novel FBPs were named as follows: FBP1, shown in **FIG. 3** (SEQ ID NO:1); FBP2, shown in **FIG. 4** (SEQ ID NO:3), FBP3a, shown in **FIG. 5** (SEQ ID NO:5), FBP4, shown in **FIG. 7** (SEQ ID NO:7), FBP5, shown in **FIG. 8** (SEQ ID NO:9), FBP6, shown in **FIG. 9** (SEQ ID NO:11), FBP7, shown in **FIG. 10** (SEQ ID NO:13). One of the seven FBPs, FBP1 (SEQ ID NO:1) was also identified by others while our screen was in progress (Margottin et al., 1998, *Molecular Cell*, 1:565-74).

[0324] BLAST programs were used to search for predicted human proteins containing an F-box in databases available through the National Center for Biotechnology Information and The Institute for Genomic Research. The alignment of the F-box motifs from these predicted human FBPs is shown in **FIG. 1**. Nineteen previously uncharacterized human FBPs were identified by aligning available sequences (GenBank Accession Nos. AC002428, AI457595, AM105408, H66467, T47217, H38755, THC274684, AI750732, AA976979, AI571815, T57296, Z44228, Z45230, N42405, AA018063, AI751015, AI400663, T74432, AA402415, AI826000, AI590138, AF474602, Z45775, AF174599, THC288870, AI017603, AF174598, THC260994, AI475671, AA768343, AF174595, THC240016, N70417, T10511, AF174603, EST04915, AA147429, A1192344, AF174594, AI147207, A1279712, AA593015, AA644633, AA335703, N26196, AF174604, AF053356, AF174606, AA836036, AA853045, AI479142, AA772788, AA039454, AA397652, AA463756, AA007384, AA749085, AI640599, THC253263, AB020647, THC295423, AA434109, AA370939, AA215393, THC271423, AF052097, THC288182, AL049953, CAB37981, AL022395,

AL031178, THC197682, and THC205131), with the nucleotide sequences derived from the F-box proteins disclosed above.

[0325] The nineteen previously uncharacterized FBP nucleotide sequences thus identified were named as follows: FBP3b, shown in **FIG. 6** (SEQ ID NO:23); FBP8, shown in **FIG. 11** (SEQ ID NO:25); FBP9, shown in **FIG. 12** (SEQ ID NO:27); FBP10, shown in **FIG. 13** (SEQ ID NO:29); FBP11, shown in **FIG. 14** (SEQ ID NO:31); FBP12, shown in **FIG. 15** (SEQ ID NO:33); FBP13, shown in **FIG. 16** (SEQ ID NO:35); FBP14, shown in **FIG. 17** (SEQ ID NO:37); FBP15, shown in **FIG. 18** (SEQ ID NO:39); FBP16, shown in **FIG. 19** (SEQ ID NO:41); FBP17, shown in **FIG. 20** (SEQ ID NO:43); FBP18, shown in **FIG. 21** (SEQ ID NO:45); FBP19, shown in **FIG. 22** (SEQ ID NO:47); FBP20, shown in **FIG. 23** (SEQ ID NO:49); FBP21, shown in **FIG. 24** (SEQ ID NO:51); FBP22, shown in **FIG. 25** (SEQ ID NO:53); FBP23, shown in **FIG. 26** (SEQ ID NO:55); FBP24, shown in **FIG. 27** (SEQ ID NO:57); and FBP25, shown in **FIG. 28** (SEQ ID NO:59). The alignment of the F-box motifs from these predicted human FBPs is shown in **FIG. 1A**. Of these sequences, the nucleotide sequences of fourteen identified FBPs, FBP3b (SEQ ID NO:23), FBP8 (SEQ ID NO:25), FBP11 (SEQ ID NO:31), FBP12 (SEQ ID NO:33), FBP13 (SEQ ID NO:35), FBP14 (SEQ ID NO:37), FBP15 (SEQ ID NO:39), FBP17 (SEQ ID NO:43), FBP18 (SEQ ID NO:45), FBP20 (SEQ ID NO:49), FBP21 (SEQ ID NO:51), FBP22 (SEQ ID NO:53), FBP23 (SEQ ID NO:55), and FBP25 (SEQ ID NO:59) were not previously assembled and represent novel nucleic acid molecules. The five remaining sequences, FBP9 (SEQ ID NO:27), FBP10 (SEQ ID NO:29), FBP16 (SEQ ID NO:41), FBP19 (SEQ ID NO:47), and FBP24 (SEQ ID NO:57) were previously assembled and disclosed in the database, but were not previously recognized as F-box proteins.

[0326] Computer analysis of human FBPs revealed several interesting features (see the schematic representation of FBPs in **FIG. 2**). Three FBPs contain WD40 domains; seven FBPs contain LRRs, and six FBPs contain other potential protein-protein interaction modules not yet identified in FBPs, such as leucine zippers, ring fingers, helix-loop-helix domains, proline rich motifs and SH2 domains.

[0327] As examples of the human FBP family, a more detailed characterization of some FBPs was performed. To confirm the specificity of interaction between the novel FBPs and human Skp1, eight in vitro translated FBPs were tested for binding to His-tagged-Skp1 pre-bound to Nickel-agarose beads. As a control Elongin C was used, the only known human Skp1 homolog. All 7 FBPs were able to bind His-Skp1 beads but not to His-tagged-Elongin C beads (**FIG. 29**). The small amount of FBPs that bound to His-tagged-Elongin C beads very likely represents non-specific binding since it was also present when a non-relevant protein (His-tagged-p27) bound to Nickel-agarose beads was used in pull-down assays (see as an example, **FIG. 29**, lane 12).

[0328] F-box deletion mutants, (Δ F)FBP1, (Δ F)FBP2, (Δ F)FBP3a, and mutants containing single point mutations in conserved amino acid residues of the F-box, FBP3a(L51A) and FBP3a(W76A) were constructed. Mutants lacking the F-box and those with point mutations

lost their ability to bind Skp1 (**FIG. 29**), confirming that human FBPs require the integrity of their F-box to specifically bind Skp1.

[0329] In order to determine whether FBP1, FBP2, FBP3a, FBP4 and FBP7 interact with human Skp1 and Cull in vivo (as Skp2 is known to do), flag-tagged-FBP1, (ΔF) FBP1, $-FBP2$, (ΔF) FBP2, $-FBP3a$, (ΔF) FBP3a, $-FBP4$ and $-FBP7$ were expressed in HeLa cells from which cell extracts were made and subjected to immunoprecipitation with an anti-Flag antibody. As detected in immunoblots with specific antibodies to Cull, Cul2 (another human cullin), and Skp1, the anti-Flag antibody co-precipitated Cull and Skp1, but not Cul2, exclusively in extracts from cells expressing wild-type FBPs (**FIG. 30** and data not shown). These data indicate that as in yeast, the human Skp1/cullin complex forms a scaffold for many FBPs.

[0330] The binding of FBPs to the Skp1/Cul1 complex is consistent with the possibility that FBPs associate with a ubiquitin ligation activity. To test this possibility, Flag-tagged were expressed in HeLa cells, FBPs together with human Skp1 and Cull. Extracts were subjected to immunoprecipitation with an anti-Flag antibody and assayed for ubiquitin ligase activity in the presence of the human ubiquitin-activating enzyme (E1) and a human Ubc. All of the wild type FBPs tested, but not FBP mutants, associated with a ubiquitin ligase activity which produced a high molecular weight smear characteristic of ubiquitinated proteins (**FIG. 31**). The ligase activity was N-ethylmaleimide (NEM) sensitive (**FIG. 31**, lane 2) and required the presence of both Ubc4 and E1. Results similar to those with Ubc4 were obtained using human Ubc3, whereas Ubc2 was unable to sustain the ubiquitin ligase activity of these SCFs (**FIG. 31**, lanes 12, 13).

[0331] Using indirect immunofluorescence techniques, the subcellular distribution of FBP1, FBP2, FBP3a, FBP4 and FBP7 was studied in human cells. Flag-tagged-versions of these proteins were expressed in HeLa, U20S, and 293T cells and subjected to immunofluorescent staining with an anti-Flag antibody. FBP1, FBP4 and FBP7 were found to be distributed both in the cytoplasm and in the nucleus, while FBP2 was detected mainly in the cytoplasm and FBP3a mainly in the nucleus. **FIG. 32** shows, as an example, the subcellular localization of FBP1, FBP2, FBP3a, FBP4 observed in HeLa cells. The localization of (ΔF) FBP1, (ΔF) FBP2, (ΔF) FBP3a mutants was identical to those of the respective wild-type proteins (**FIG. 32**) demonstrating that the F-box and the F-box-dependent binding to Skp1 do not determine the subcellular localization of FBPs. Immunofluorescence stainings were in agreement with the results of biochemical subcellular fractionation.

[0332] 6.2.2 Northern Blot Analysis of Novel Ubiquitin Ligase Gene Transcripts

[0333] RNA blot analysis was performed on poly(A)+ mRNA from multiple normal human tissues (heart, brain, placenta, lung, liver, skeletal, muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocytes, see **FIG. 33**). FBP1 mRNA transcripts (a major band of 7-kb and two minor bands of 3.5- and ~2.5 kb) were expressed in all of the 16 human tissues tested but were more prevalent in brain and testis. Testis was the only tissue expressing the smaller FBP1 mRNA forms in amounts equal to, if not in excess of, the 7

kb form. FBP2 transcripts (~7.7-kb and ~2.4-kb) were expressed in all tissues tested, yet the ratio of the FBP2 transcripts displayed some tissue differences. An approximately 4 kb FBP3a transcript was present in all tissues tested and two minor FBP3a forms of approximately 3 kb and 2 kb became visible, upon longer exposure, especially in the testis. An approximately 4.8 kb FBP4 transcript was expressed in all normal human tissues tested, but was particularly abundant in heart and pancreas. Finally, the pattern of expression of the new FBPs was compared to that of FBP1 whose mRNA species (a major band ~4 kb and a minor band of ~8.5 kb) were found in all tissues but was particularly abundant in placenta.

[0334] 6.2.3 Chromosomal Localization Of The Human FBP Genes Unchecked degradation of cellular regulatory proteins (e.g., p53, p27, β -catenin) has been observed in certain tumors, suggesting the hypothesis that deregulated ubiquitin ligases play a role in this altered degradation (reviewed in A. Ciechanover, 1998, *Embo J*, 17: 7151). A well understood example is that of MDM2, a proto-oncogene encoding a ubiquitin ligase whose overexpression destabilize its substrate, the tumor suppressor p53 (reviewed by Brown and Pagano, 1997, *Biochim Biophys Acta*, 1332: 1, 1998). To map the chromosomal localization of the human FBP genes and to determine if these positions coincided with loci known to be altered in tumors or in inherited disease, fluorescence in situ hybridization (FISH) was used. The FBP1 gene was mapped and localized to 10q24 (**FIG. 34A**), FBP2 to 9q34 (**FIG. 34B**), FBP3a to 13q22 (**FIG. 34C**), FBP4 to 5p 12 (**FIG. 34D**) and FBP5 to 6q25-26 (**FIG. 34E**). FBP genes (particularly FBP1, FBP3a, and FBP5) are localized to chromosomal loci frequently altered in tumors (for references and details see Online Mendelian Inheritance in Man database, <http://www3.ncbi.nlm.nih.gov/omim/>). In particular, loss of 10q24 (where FBP1 is located) has been demonstrated in approx. 10% of human prostate tumors and small cell lung carcinomas (SCLC), suggesting the presence of a tumor suppressor gene at this location. In addition, up to 7% of childhood acute T-cell leukemia is accompanied by a translocation involving 10q24 as a breakpoint, either t(10; 14)(q24;q11) or t(7; 10)(q35;q24). Although rarely, the 9q34 region (where FBP2 is located) has been shown to be a site of loss of heterozygosity (LOH) in human ovarian and bladder cancers. LOH is also observed in the region. Finally, 6q25-26 (where FBP5 is located) has been shown to be a site of loss of heterozygosity in human ovarian, breast and gastric cancers hepatocarcinomas, Burkitt's lymphomas, and parathyroid adenomas.

7. EXAMPLE

FBP1 Regulates the Stability of β -Catenin

[0335] Deregulation of β -catenin proteolysis is associated with malignant transformation. *Xenopus* Slimb and *Drosophila* FBP1 negatively regulate the Wnt/ β -catenin signaling pathway (Jiang and Struhl, 1998, supra; Marikawa and Elinson, 1998). Since ubiquitin ligase complexes physically associate with their substrates, the studies in this Example were designed to determine whether FBP1 can interact with β -catenin. The results show that FBP1 forms a novel ubiquitin ligase complex that regulates the in vivo stability of β -catenin. Thus, the identification of FBP1 as a component of the novel ubiquitin ligase complex that ubiquitinates

β -catenin, provides new targets that can be used in screens for agonists, antagonists, ligands, and novel substrates using the methods of the present invention. Molecules identified by these assays are potentially useful drugs as therapeutic agents against cancer and proliferative disorders.

[0336] 7.1 Materials and Methods for Identification of FBP1 Function

[0337] Recombinant proteins, Construction of F-box mutants, Antibodies, Transient transfections, Immunoprecipitation, Immunoblotting, Cell culture and Extract preparation Details of the methods are described in Section 6.1, supra.

[0338] 7.2 Results

[0339] 7.2.1 Human FBP1 Interacts With β -Catenin

[0340] Flag-tagged FBP1 and β -catenin viruses were used to co-infect insect cells, and extracts were analyzed by immunoprecipitation followed by immunoblotting. β -catenin was co-immunoprecipitated by an anti-Flag antibody (FIG. 35A), indicating that in intact cells β -catenin and FBP1 physically interact. It has been shown that binding of the yeast FBP Cdc4 to its substrate Sic1 is stabilized by the presence of Skp1 (Skowrya et al., 1997, Cell, 91, 209-219). Simultaneous expression of human Skp1 had no effect on the strength of the interaction between FBP1 and β -catenin. To test the specificity of the FBP1/ β -catenin interaction, cells were co-infected with human cyclin D1 and FBP1 viruses. The choice of this cyclin was dictated by the fact that human cyclin D1 can form a complex with the Skp2 ubiquitin ligase complex (Skp1-Cull-Skp2; Yu et al., 1998, Proc. Natl. Acad. Sci. U.S.A., 95:11324-9). Under the same conditions used to demonstrate the formation of the FBP1/ β -catenin complex, cyclin D1 could not be co-immunoprecipitated with Flag-tagged FBP1, and anti-cyclin D1 antibodies were unable to co-immunoprecipitate FBP1 (FIG. 35B, lanes 1-3). Co-expression of Skp1 (FIG. 35B, lanes 4-6) or Cdk4 with FBP1 and cyclin D1 did not stimulate the association of cyclin D1 with FBP1.

[0341] Mammalian expression plasmids carrying HA-tagged β -catenin and Flag-tagged FBP1 (wild type or mutant) were then co-transfected in human 293 cells. β -catenin was detected in anti-Flag immunoprecipitates when co-expressed with either wild type or (Δ F)FBP1 mutant (FIG. 35C, lanes 4-6), confirming the presence of a complex formed between β -catenin and FBP1 in human cells.

[0342] 7.2.2 F-box Deleted FBP1 Mutant Stabilizes P-Catenin In Vivo The association of (Δ F)FBP1 to β -catenin suggested that (Δ F)FBP1 might act as a dominant negative mutant in vivo by being unable to bind Skp1/Cul1 complex, on the one hand, while retaining the ability to bind p catenin, on the other. HA-tagged p-catenin was co-expressed together with Flag-tagged (Δ F)FBP1 or with another F-box deleted FBP, (Δ F)FBP2. FBP2 was also obtained with our screening for Skp1-interactors; and, like FBP1, contains several WD-40 domains. The presence of (Δ F)FBP1 specifically led to the accumulation of higher quantities of β -catenin (FIG. 36A). To determine whether this accumulation was due to an increase in β -catenin stability, we measured the half-life of β -catenin using pulse chase analysis. Human 293 cells were transfected with HA-tagged β -catenin alone or in combination with the wild type or mutant FBP1. While wild type Fbp1 had little effect on the

degradation of β -catenin, the F-box deletion mutant prolonged the half life of β -catenin from 1 to 4 hours (FIG. 36B).

[0343] FBP1 is also involved in CD4 degradation induced by the HIV-1 Vpu protein (Margottin et al., supra). It has been shown that Vpu recruits FBP1 to DC4 and (Δ F) FBP1 inhibits Vpu-mediated CD4 regulation. In addition, FBP1-ubiquitin ligase complex also controls the stability of IKB α (Yaron et al., 1998, Nature, 396: 590). Thus, the interactions between FBP1 and β -catenin, Vpu protein, CD4, and IKB α are potential targets that can be used to screen for agonists, antagonists, ligands, and novel substrates using the methods of the present invention.

8. EXAMPLE

Methods for Identifying p27 as a Substrate of the FBP Skp2

[0344] Degradation of the mammalian G1 cyclin-dependent kinase (Cdk) inhibitor p27 is required for the cellular transition from quiescence to the proliferative state. The ubiquitination and degradation of p27 depend upon its phosphorylation by cyclin/Cdk complexes. Skp2, an F-box protein essential for entry into S phase, specifically recognizes p27 in a phosphorylation-dependent manner. Furthermore, both in vivo and in vitro, Skp2 is a rate-limiting component of the machinery that ubiquitinates and degrades phosphorylated p27. Thus, p27 degradation is subject to dual control by the accumulation of both Skp2 and cyclins following mitogenic stimulation.

[0345] This Example discloses novel assays that have been used to identify the interaction of Skp2 and p27 in vitro. First, an in vitro ubiquitination assay performed using p27 as a substrate is described. Second, Skp2 is depleted from cell extracts using anti-Skp2 antibody, and the effect on p27 ubiquitin ligase activity is assayed. Purified Skp2 is added back to such immunodepleted extracts to restore p27 ubiquitination and degradation. Also disclosed is the use of a dominant negative mutant, (Δ F)Skp2, which interferes with p27 ubiquitination and degradation.

[0346] The assays described herein can be used to test for compounds that inhibit cell proliferation. The assays can be carried out in the presence or absence of molecules, compounds, peptides, or other agents described in Section 5.5. Agents that either enhance or inhibit the interactions or the ubiquitination activity can be identified by an increase or decrease the formation of a final product are identified. Such agents can be used, for example, to inhibit Skp2-regulated p27 ubiquitination and degradation in vivo. Molecules identified by these assays are potentially useful drugs as therapeutic agents against cancer and proliferative disorders.

[0347] Dominant negative mutants, for example the mutant (Δ F)Skp2, and antisense oligos targeting SKP2, mRNA interfere with p27 ubiquitination and degradation, and can be used in gene therapies against cancer. The assays described herein can also be used to identify novel substrates of the novel FBP proteins, as well as modulators of novel ubiquitin ligase complex—substrate interactions and activities.

[0348] 8.1 Materials and Methods for Identification of p27 as a Skp2 Substrate

[0349] Protein extraction for in vitro ubiquitination assay. Approx. 4 ml of HeLa S3 cell pellet were suspended in 6 ml of ice-cold buffer consisting of 20 mM Tris-HCl (pH 7.2), 2 mM DTT, 0.25 mM EDTA, 10 μ g/ml leupeptin, and 10 μ g/ml pepstatin. The suspension was transferred to a cell nitrogen-disruption bomb (Parr, Moline, Ill., cat #4639) that had been rinsed thoroughly and chilled on ice before use. The bomb chamber was connected to a nitrogen tank and the pressure was brought slowly to 1000 psi. The chamber was left on ice under the same pressure for 30 minutes and then the pressure was released slowly. The material was transferred to an Eppendorf tube and centrifuged in a microcentrifuge at 10,000 g for 10 minutes. The supernatant (S-10) was divided into smaller samples and frozen at -80° C. This method of extract preparation based on the use of a cell nitrogen-disruption bomb extract preserves the activity to in vitro ubiquitinate p27 better than the method previously described (Pagano et al., 1995, Science 269:682-685).

[0350] Reagents and antibodies Ubiquitin aldehyde (Hershko & Rose, 1987, Proc. Natl. Acad. Sci. USA 84:1829-33), methyl-ubiquitin (Hershko & Heller, 1985, Biochem. Biophys. Res. Commun. 128:1079-86) and p13 beads (Brizuela et al., 1987, EMBO J. 6:3507-3514) were prepared as described. β , γ -imidoadenosine-50-triphosphate (AMP-PNP), staurosporine, hexokinase, and deoxy-glucose were from Sigma; lovastatin obtained from Merck; flavopiridol obtained from Hoechst Marion Roussel. The phospho-site p27 specific antibody was generated in collaboration with Zymed Inc. by injecting rabbits with the phospho-peptide NAGSVEQT*PKKPGLRRRQT (SEQ ID NO: 85), corresponding to the carboxy terminus of the human p27 with a phosphothreonine at position 187 (T*). The antibody was then purified from serum with two rounds of affinity chromatography using both phospho- and nonphospho-peptide chromatography. All the other antibodies are described in Section 6.1.

[0351] Immunodepletion Assays For immunodepletion assays, 3 μ l of an Skp2 antiserum was adsorbed to 15 μ l Affi-Prep Protein-A beads (BioRad), at 4° C. for 90 min. The beads were washed and then mixed (4° C., 2 hours) with 40 μ l of HeLa extract (approximately 400 μ g of protein). Beads were removed by centrifugation and supernatants were filtered through a 0.45- μ Microspin filter (Millipore). Immunoprecipitations and immunoblots were performed as described (M. Pagano, et al., 1995, supra. Rabbit polyclonal antibody against purified GST-Skp2 was generated, affinity-purified (AP) and characterized as described (M. Pagano, in Cell Cycle-Materials and Methods, M. Pagano Ed. (Springer, N.Y., 1995), chap. 24; E. Harlow and D. Lane, in Using antibodies. A Laboratory Manual (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1998), in collaboration with Zymed Inc. (cat #51-1900). Monoclonal antibodies (Mabs) to human Cull, and cyclin E, (Faha et al., 1993, J. of Virology 67:2456); AP rabbit antibodies to human p27, Skp1 (Latres et al., 1999, Oncogene 18:849), Cdk2 (Pagano, et al., 1992, Science 255:1144) and phospho-site p27 specific antibody. Mab to cyclin B was from Santa Cruz (cat # sc-245); Mabs to p21 (cat # C24420) and p27 (cat # K25020) Transduction lab; anti-Flag rabbit antibody

from Zymed (cat #71-5400). An AP goat antibody to an N-terminal Skp2 peptide (Santa Cruz, cat # sc-1567) was used.

[0352] Construction of Skp2 F-box mutant (Δ F)Skp2 mutant was obtained by removing a DNA fragment (nucleotides 338-997) with BspEI and XbaI restriction enzymes, and replacing it with a PCR fragment containing nucleotides 457 to 997. The final construct encoded a protein lacking residues 113-152.

[0353] Recombinant proteins cDNA fragments encoding the following human proteins: Flag-tagged FBP1, Flag-tagged (Δ F)FBP1, Flag-tagged FBP3a, Skp2, HA-tagged Cull, HA-tagged Cul2, β -catenin, His-tagged cyclin D1, Skp1, His-tagged Skp1, His-tagged Elongin C were inserted into the baculovirus expression vector pBacpak-8 (Clontech) and cotransfected into Sf9 cells with linearized baculovirus DNA using the BaculoGold transfection kit (Pharmingen). Baculoviruses expressing human His-tagged cyclin E and HA-tagged Cdk2 were supplied by D. Morgan (Desai, 1992, Molecular Biology of the Cell 3: 571). Recombinant viruses were used to infect 5B cells and assayed for expression of their encoded protein by immunoblotting as described above. His-proteins were purified with Nickel-agarose (Invitrogen) according to the manufacturer's instructions. The different complexes were formed by co-expression of the appropriate baculoviruses and purified by nickel-agarose chromatography, using the His tag at the 5' of Skp1 and cyclin E. Unless otherwise stated, recombinant proteins were added to incubations at the following amounts: cyclin E/Cdk2, \sim 0.5 pmol; Skp1, \sim 0.5 pmol; Skp2, \sim 0.1 pmol; FBP1, \sim 0.1 pmol; FBP3a, \sim 0.1 pmol, Cull, \sim 0.1 pmol. The molar ratio of Skp1/Skp2, Skp1/FPB1, Skp1/FPB3a, and Skp1/Cull in the purified preparations was \sim 5.

[0354] Extract preparation and cell synchronization, transient Transfections, Immunoprecipitation and Immunoblotting Methods were carried out as described in Section 6.1, supra.

[0355] 8.2 Results

[0356] 8.2.1 p27 In Vitro Ubiquitination Assay

[0357] In an exemplary in vitro ubiquitination assay, logarithmically growing, HeLa-S3 cells were collected at a density of 6×10^5 cells/ml. Cells are arrested in G1 by 48-hour treatment with 70 μ M lovastatin as described (O'Connor & Jackman, 1995 in Cell Cycle-Materials and Methods, M. Pagano, ed., Springer, N.Y., chap. 6). 1 μ l of in vitro translated [35S]p27 is incubated at 30° C. for different times (0-75 minutes) in 10 μ l of ubiquitination mix containing: 40 mM Tris pH 7.6, 5 mM $MgCl_2$, 1 mM DTT, 10% glycerol, 1 μ M ubiquitin aldehyde, 1 mg/ml methyl ubiquitin, 10 mM creatine phosphate, 0.1 mg/ml creatine phosphokinase, 0.5 mM ATP, 1 μ M okadaic acid, 20-30 μ g HeLa cell extract. Ubiquitin aldehyde can be added to the ubiquitination reaction to inhibit the isopeptidases that would remove the chains of ubiquitin from p27. Addition of methyl ubiquitin competes with the ubiquitin present in the cellular extracts and terminates p27 ubiquitin chains. Such chains appear as discrete bands instead of a high molecular smear. These shorter polyubiquitin chains have lower affinity for the proteasome and therefore are more stable. Reactions are terminated with Laemmli sample buffer containing β -mercaptoethanol and the products can be analyzed on protein gels under denaturing conditions.

[0358] Polyubiquitinated p27 forms are identified by autoradiography. p27 degradation assay is performed in a similar manner, except that (i) Methylated ubiquitin and ubiquitin aldehyde were omitted; (ii) The concentration of HeLa extract is approximately 7 $\mu\text{g}/\mu\text{l}$; (iii) Extracts are prepared by hypotonic lysis (Pagano et al., 1995, *Science* 269:682), which preserves proteasome activity better than the nitrogen bomb disruption procedure. In the absence of methyl ubiquitin, p27 degradation activity, instead of p27 ubiquitination activity, can be measured.

[0359] The samples are immunoprecipitated with an antibody to p27 followed by a subsequent immunoprecipitation with an anti-ubiquitin antibody and run on an 8% SDS gel. The high molecular species as determined by this assay are ubiquitinated. As a control, a p27 mutant lacking all 13 lysines was used. This mutant form of p27 is not ubiquitinated and runs at higher molecular weight on the 8% SDS gel.

[0360] 8.2.2 p27-Skp2 Interaction Assays and p27-Skp2 Immunodepletion Assay

[0361] The recruitment of specific substrates by yeast and human FBPs to Skp1/cullin complexes is phosphorylation-dependent. Accordingly, peptides derived from IBA and p-catenin bind to FBP1 specifically and in a phosphorylation-dependent manner (Yaron, 1998, *Nature* 396: 590; Winston et al., 1999, *Genes Dev.* 13: 270). A p27 phosphopeptide with a phosphothreonine at position 187 was assayed for its ability to bind to human FBPs, including Skp2 and the FBPI, FBP2, FBP3a, FBP4, FBPS, FBP6, and FBP7, isolated by using a 2-hybrid screen using Skp1 as bait, as described in Section 6, above. Four of these FBPs contain potential substrate interaction domains, such as WD-40 domains in FBP1 and FBP2, and leucine-rich repeats in Skp2 and FBP3a. The phospho-p27 peptide was immobilized to Sepharose beads and incubated with these seven in vitro translated FBPs (FIG. 37A). Only one FBP, Skp2, was able to bind to the phospho-T187 p27 peptide. Then, beads linked to p27 peptides (in either phosphorylated or unphosphorylated forms) or with an unrelated phosphopeptide were incubated with HeLa cell extracts. Proteins stably associated with the beads were examined by immunoblotting. Skp2 and its associated proteins, Skp1 and Cull, were readily detected as proteins bound to the phospho-p27 peptide but not to control peptides (FIG. 37B).

[0362] To further study p27 association to Skp2, in vitro translated p27 was incubated with either Skp1/Skp2 complex, cyclin E/Cdk2 complex, or the combination of both complexes under conditions in which p27 is phosphorylated on T187 by cyclin E/Cdk2 (Montagnoli, A., et al., 1999, *Genes & Dev* 13: 1181). Samples were then immunoprecipitated with an anti-Skp2 antibody. p27 was co-immunoprecipitated with Skp2 only in the presence of cyclin E/Cdk2 complex (FIG. 37C). Notably, under the same conditions, a T187-to-alanine p27 mutant, p27(T187A), was not co-immunoprecipitated by the anti-Skp2 antibody. Finally, we tested Skp2 and p27 association in vivo. Extracts from HeLa cells and IMR90 human diploid fibroblasts were subjected to immunoprecipitation with two different antibodies to Skp2 and then immunoblotted. p27 and Cull, but not cyclin D1 and cyclin B1, were specifically detected in Skp2 immunoprecipitates (FIG. 38). Importantly, using a phospho-T187 site p27 specific antibody we demonstrated that the Skp2-

bound p27 was phosphorylated on T187 (FIG. 38, lane 2, bottom panel). Furthermore, an anti-peptide p27 antibody specifically co-immunoprecipitated Skp2. These results indicate that the stable interaction of p27 with Skp2 was highly specific and dependent upon phosphorylation of p27 on T187.

[0363] A cell-free assay for p27 ubiquitination which faithfully reproduced the cell cycle stage-specific ubiquitination and degradation of p27 has been developed (Montagnoli et al., supra). Using this assay, a p27-ubiquitin ligation activity is higher in extracts from asynchronously growing cells than in those from G1-arrested cells (FIG. 39A, lanes 2 and 4). In accordance with previous findings (Montagnoli, A., et al., supra), the addition of cyclin E/Cdk2 stimulated the ubiquitination of p27 in both types of extracts (FIG. 39A, lanes 3 and 5). However, this stimulation was much lower in extracts from G1-arrested cells than in those from growing cells, suggesting that in addition to cyclin E/Cdk2, some other component of the p27-ubiquitin ligation system is rate-limiting in G1. This component could be Skp2 since, in contrast to other SCF subunits, its levels are lower in extracts from G1 cells than in those from asynchronous cells and are inversely correlated with levels of p27 (FIGS. 39B and 43). Skp2 was thus tested to determine if it is a rate-limiting component of a p27 ubiquitin ligase activity. The addition of recombinant purified Skp1/Skp2 complex alone to G1 extracts did not stimulate p27 ubiquitination significantly (FIG. 39A, lane 6). In contrast, the combined addition of Skp1/Skp2 and cyclin E/Cdk2 complexes strongly stimulated p27 ubiquitination in G1 extracts (FIG. 39A, lane 7). Similarly, the combined addition of Skp1/Skp2 and cyclin E/Cdk2 strongly stimulated p27 proteolysis as measured by a degradation assay (FIG. 39A, lanes 13-16). Since the Skp1/Skp2 complex used for these experiments was isolated from insect cells co-expressing baculovirus His-tagged-Skp1 and Skp2 (and co-purified by nickel-agarose chromatography), it was possible that an insect-derived F-box protein co-purified with His-Skp1 and was responsible for the stimulation of p27 ubiquitination in G1 extracts. This possibility was eliminated by showing that the addition of a similar amount of His-tagged-Skp1, expressed in the absence of Skp2 in insect cells and purified by the same procedure, did not stimulate p27 ubiquitination in the presence of cyclin E/Cdk2 (FIG. 39A, lane 8). Furthermore, we found that neither FBP1 nor FBP3a could replace Skp2 for the stimulation of p27-ubiquitin ligation in G1 extracts (FIG. 39A, lanes 9-12). Stimulation of p27-ubiquitination in G1 extracts by the combined addition of Skp1/Skp2 and cyclin E/Cdk2 could be observed only with wild-type p27, but not with the p27(T187A) mutant (lanes 17-20), indicating that phosphorylation of p27 on T187 is required for the Skp2-mediated ubiquitination of p27. These findings indicated that both cyclin E/Cdk2 and Skp1/Skp2 complexes are rate-limiting for p27 ubiquitination and degradation in the G1 phase.

[0364] To further investigate the requirement of Skp2 for p27 ubiquitin ligation, Skp2 was specifically removed from extracts of asynchronously growing cells by immunodepletion with an antibody to Skp2. The immunodepletion procedure efficiently removed most of Skp2 from these extracts and caused a drastic reduction of p27-ubiquitin ligation activity (FIG. 40A, lane 4) as well as of p27 degradation activity. This effect was specific as shown by the following observations: (i) Similar treatment with pre-immune serum

did not inhibit p27-ubiquitination (**FIG. 40A**, lane 3); (ii) Pre-incubation of anti-Skp2 antibody with recombinant GST-Skp2 (lane 5), but not with a control protein (lane 4), prevented the immunodepletion of p27-ubiquitination activity from extracts; (iii) p27-ubiquitinating activity could be restored in Skp2-depleted extracts by the addition of His-Skp1/Skp2 complex (**FIG. 40B**, lane 3) but not His-Skp1 (lane 2), His-Skp1/Cull complex (lane 4), or His-Skp1/FBP1.

[0365] We then immunoprecipitated Skp2 from HeLa extracts and tested whether this immunoprecipitate contained a p27 ubiquitinating activity. The anti-Skp2 beads, but not a immunoprecipitate made with a pre-immune (PI) serum, was able to induce p27 ubiquitination in the presence of cyclin E/Cdk2 (**FIG. 40C**, lanes 2 and 3). The addition of purified recombinant E1 ubiquitin-activating enzyme, and purified recombinant Ubc3 did not greatly increase the ability of the Skp2 immunoprecipitate to sustain p27 ubiquitination, (**FIG. 40C**, lane 5), likely due to the presence of both proteins in the rabbit reticulocyte lysate used for p27 in vitro translation.

[0366] 8.2.3 F-Box Deleted SKP2 Mutant Stabilizes p27 In Vivo

[0367] Skp2 also targets p27 for ubiquitin-mediated degradation in vivo. The F-box-deleted FBP1 mutant, (Δ F)FBP1, acts in vivo as a dominant negative mutant, most likely because without the F-box is unable to bind Skp1/Cull complex but retains the ability to bind its substrates. Therefore, once expressed in cells, (Δ F)Fb sequesters β -catenin and IKBa and causes their stabilization. An F-box deleted Skp2 mutant, (Δ F)Skp2, was constructed. p27 was expressed in murine cells either alone or in combination with (Δ F)Skp2 or (Δ F)FBP1 (see **FIG. 41**). The presence of (Δ F)Skp2 led to the accumulation of higher quantities of p27. To determine whether this accumulation was due to an increase in p27 stability, the half-life of p27 was measured using pulse chase analysis (for details, see Section 8, above). Indeed, (Δ F)Skp2 prolonged p27 half-life from less than 1 hour to 3 hours. Since in these experiments the efficiency of transfection was approximately 10%, (Δ F)Skp2 affected only the stability of co-expressed human exogenous p27, but not of murine endogenous p27.

[0368] 8.2.4 SKP2 Antisense Experiments

[0369] SKP2 mRNA was targeted with antisense oligonucleotides to determine whether a decrease in Skp2 levels would influence the abundance of endogenous p27. Two different antisense oligos, but not control oligodeoxynucleotides induced a decrease in Skp2 protein levels (**FIG. 42**). Concomitant with the Skp2 decrease, there was a substantial increase in the level of endogenous p27 protein. Similar results were obtained with cells blocked at the G1/S transition with hydroxyurea or aphidicolin treatment (lanes 9-16). Thus, the effect of the SKP2 antisense oligos on p27 was not a secondary consequence of a possible block in G1 due to the decrease in Skp2 levels.

[0370] Antisense experiments were performed as described in (Yu, 1998, Proc. Natl. Acad. Sci. U.S.A. 95: 11324). Briefly, four oligodeoxynucleotides that contain a phosphorothioate backbone and C-5 propyne pyrimidines were synthesized (Keck Biotechnology Resource Laboratory at Yale University): (1) 5'-CCTGGGGGATGTTCTCA-

3' (SEQ ID NO: 86) (the antisense direction of human Skp2 cDNA nucleotides 180-196); (2) 5'-GGCTTCCGGGCATTAG-3' (SEQ ID NO: 87) [the scrambled control of (1)]; (3) 5'-CATCTGGCACGATTCCA-3' (SEQ ID NO: 88) (the antisense direction of Skp2 cDNA nucleotides 1137-1153); (4) 5'-CCGCTCATCGTATGACA-3' (89) [the scrambled control for (3)]. The oligonucleotides were delivered into HeLa cells using Cytofectin GS (Glen Research) according to the manufacturers instructions. The cells were then harvested between 16 and 18 hours postransfection.

9. EXAMPLE

Assay to Identify an FBP Interaction with a Cell Cycle Regulatory Protein (e.g., SKP2 with E2F)

[0371] The following study was conducted to identify novel substrates of the known FBP, Skp2.

[0372] As shown in **FIG. 44**, E2F-1, but not other substrates of the ubiquitin pathway assayed, including p53 and Cyclin B, physically associates with Skp2. Extracts of insect cells infected with baculoviruses co-expressing Skp2 and E2F-1, (lanes 1,4 and 5), or Skp2 and hexa-histidine p53 (His-p53) (lanes 2,6,7,10 and 11), or Skp2 and His-Cyclin B (lanes 3,8,9,12, and 13) were either directly immunoblotted with an anti-serum to Skp2 (lanes 1-3) or first subjected to immunoblotted with an anti-serum to Skp2 (lanes 1-3) or first subjected to immunoprecipitation with the indicated antibodies and then immunoblotted with an anti-serum to Skp2 (lanes 4-13). Antibodies used in the immunoprecipitations are: normal purified mouse immunoglobulins (IgG) (lane 4,6,10 and 12), purified mouse monoclonal anti-E2F-1 antibody (KH-95, from Santa Cruz) (lane 5), purified mouse monoclonal anti-p53 antibody (DO-1, from Oncogene Science) (lane 7), purified rabbit IgG (lane 8), purified rabbit polyclonal anti-Cyclin B antibody (lane 9), purified mouse monoclonal anti-His antibody (clone 34660, from Qiagen) (lanes 11 and 13).

[0373] As shown in **FIG. 44B**, Skp2 physically associates with E2F-1 but not with other substrates of the ubiquitin pathway (p53 and Cyclin B). Extracts of insect cells infected with baculoviruses co-expressing Skp2 and E2F-1 (lanes 1-3), or Skp2 and His-p53 (lanes 4-6), or Skp2 and His-Cyclin B (lanes 7-9) were either directly immunoblotted with antibodies to the indicated proteins (lanes 1,4 and 7) or first subjected to immunoprecipitation with the indicated anti-sera and then immunoblotted with antibodies to the indicated proteins (lanes 2,3,5,6,8 and 9). Anti-sera used in the immunoprecipitations are: anti-Skp2 serum (lanes 2,5 and 8), and normal rabbit serum (NRS) (lane 3,6 and 9).

[0374] As shown in **FIG. 44C**, E2F-1 physically associates with Skp2 but not with another F-box protein (FBP1). Extracts of insect cells infected with baculoviruses co-expressing Skp2 and E2F-1 (lanes 1,3 and 4), or Flag-tagged-FBP1 and E2F-1 (lanes 2,5 and 6) were either directly immunoblotted with a mouse monoclonal anti-E2F-1 antibody (lanes 1 and 2) or first subjected to immunoprecipitation with the indicated antibodies and then immunoblotted with a mouse monoclonal anti-E2F-1 antibody (lanes 3-6). Antibodies used in the immunoprecipitations are: anti-Skp2 serum (lanes 3), NRS (lane 4), purified rabbit polyclonal anti-Flag (lane 5), purified rabbit IgG (lane 6).

[0375] The methodology used in this example can also be applied to identify novel substrates of any FBP, including, but not limited to, the FBPs of the invention, such as FBP1, FBP2, FBP3a, FBP3b, FBP4, FBP5, FBP6, FBP7, FBP8, FBP9, FBP10, FBP11, FBP12, FBP13, FBP14, FBP15, FBP16, FBP17, FBP18, FBP19, FBP20, FBP21, FBP22, FBP23, FBP24, and FBP25.

[0376] The invention is not to be limited in scope by the specific embodiments described which are intended as single illustrations of individual aspects of the invention, and

functionally equivalent methods and components are within the scope of the invention. Indeed various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

[0377] All references cited herein are incorporated herein by reference for all purposes.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 90

<210> SEQ ID NO 1

<211> LENGTH: 2151

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

tgcgttgct gcgccctggc accaaagggg cgccccggc ggagagcga ccagtggcc	60
tcggcgatta tggaccggc cgaggcggg ctgcaagaga aggcactcaa gtttatgaat	120
tcctcagaga gagaagactg taataatggc gaaccccta ggaagataat accagagaag	180
aattcactta gacagacata caacagctgt gccagactct gcttaacca agaaacagta	240
tgtttagcaa gcaactgctat gaagactgag aattgtgtgg ccaaaacaaa acttgccaat	300
ggcacttcca gtatgattgt gccaagcaa cggaaactct cagcaagcta tgaaaaggaa	360
aaggaaactgt gtgtcaaata ctttgagcag tggtcagagt cagatcaagt ggaatttgtg	420
gaacatctta tatcccaaat gtgtcattac caacatggc acataaactc gtatcttaaa	480
cctatgttgc agagagattt cataactgct ctgccagctc ggggattgga tcatatcgct	540
gagaacattc tgtcatacct ggatgcaaaa tcaactatgt ctgctgaact tgtgtgcaag	600
gaatggtacc gactgacctc tgatggcatg ctgtggaaga agcttatcga gagaatggtc	660
aggacagatt ctctgtggag aggcctggca gaacgaagag gatggggaca gtatttatc	720
aaaaacaac ctctgacgg gaatgtcct ccaactctt ttatagagc actttatcct	780
aaaattatac aagacattga gacaatagaa tctaattgga gatgtggaag acatagtta	840
cagagaattc actgccgaag tgaacaagc aaaggagttt actgtttaca gtatgatgat	900
cagaaaaatag taagcggcct tcgagacaac acaatcaaga tctgggataa aaacacattg	960
gaatgcaagc gaattctcac aggccatata ggttcagtcc tctgtctcca gtatgatgag	1020
agagtgatca taacaggatc atcggattcc acggtcagag tgtgggatgt aaatacaggt	1080
gaaatgctaa acacgttgat tcaccattgt gaagcagttc tgcacttgcg tttcaataat	1140
ggcatgatgg tgacctgctc caaagatcgt tcattgctg tatgggatat ggcctcccca	1200
actgacatta ccctccggag ggtgctggtc ggacaccgag ctgctgtcaa tgtttagtagc	1260
tttgatgaca agtacattgt ttctgcatct ggggatagaa ctataaaggt atggaacaca	1320
agtacttggt aatttgtaag gaccttaaat ggacacaaac gaggcattgc ctgtttgcag	1380
tacagggaca ggctggtagt gactggctca tctgacaaca ctatcagatt atgggacata	1440
gaatgtgggt catgtttacg agtgtagaa ggccatgagg aattggtgcg ttgtattcga	1500

-continued

```

tttgataaca agaggatagt cagtggggcc tatgatggaa aaattaaagt gtgggatctt 1560
gtggctgctt tggacccccg tgctcctgca gggacactct gtctacggac ccttgtggag 1620
cattccggaa gagtttttctg actacagttt gatgaattcc agattgtcag tagttcacat 1680
gatgacacaa tcctcatctg ggacttccta aatgatccag ctgccaagc tgaaccccc 1740
cgttccccctt ctgaacata cacctacatc tccagataaa taaccataca ctgacctcat 1800
acttgcccag gacccattaa agttgcggtt tttaacgtat ctgccaatac caggatgagc 1860
aacaacagta acaatcaaac tactgccag tttccctgga ctagccgagg agcagggctt 1920
tgagactcct gttgggacac agttgggtctg cagtcggccc aggacggctc actcagcaca 1980
actgactgct tcagtgtctg tatcagaaga tgtcttctat caattgtgaa tgattggaac 2040
ttttaaacct cccctcctct cctcctttca cctctgcacc tagttttttc ccattggttc 2100
cagacaaagg tgacttataa atatatttag tgttttgcca gaaaaaaaa a 2151

```

<210> SEQ ID NO 2

<211> LENGTH: 569

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

Met Asp Pro Ala Glu Ala Val Leu Gln Glu Lys Ala Leu Lys Phe Met
  1             5             10            15

Asn Ser Ser Glu Arg Glu Asp Cys Asn Asn Gly Glu Pro Pro Arg Lys
      20             25            30

Ile Ile Pro Glu Lys Asn Ser Leu Arg Gln Thr Tyr Asn Ser Cys Ala
      35             40            45

Arg Leu Cys Leu Asn Gln Glu Thr Val Cys Leu Ala Ser Thr Ala Met
      50             55            60

Lys Thr Glu Asn Cys Val Ala Lys Thr Lys Leu Ala Asn Gly Thr Ser
      65             70            75            80

Ser Met Ile Val Pro Lys Gln Arg Lys Leu Ser Ala Ser Tyr Glu Lys
      85             90            95

Glu Lys Glu Leu Cys Val Lys Tyr Phe Glu Gln Trp Ser Glu Ser Asp
      100            105            110

Gln Val Glu Phe Val Glu His Leu Ile Ser Gln Met Cys His Tyr Gln
      115            120            125

His Gly His Ile Asn Ser Tyr Leu Lys Pro Met Leu Gln Arg Asp Phe
      130            135            140

Ile Thr Ala Leu Pro Ala Arg Gly Leu Asp His Ile Ala Glu Asn Ile
      145            150            155            160

Leu Ser Tyr Leu Asp Ala Lys Ser Leu Cys Ala Ala Glu Leu Val Cys
      165            170            175

Lys Glu Trp Tyr Arg Val Thr Ser Asp Gly Met Leu Trp Lys Lys Leu
      180            185            190

Ile Glu Arg Met Val Arg Thr Asp Ser Leu Trp Arg Gly Leu Ala Glu
      195            200            205

Arg Arg Gly Trp Gly Gln Tyr Leu Phe Lys Asn Lys Pro Pro Asp Gly
      210            215            220

Asn Ala Pro Pro Asn Ser Phe Tyr Arg Ala Leu Tyr Pro Lys Ile Ile
      225            230            235            240

Gln Asp Ile Glu Thr Ile Glu Ser Asn Trp Arg Cys Gly Arg His Ser

```

-continued

245					250					255					
Leu	Gln	Arg	Ile	His	Cys	Arg	Ser	Glu	Thr	Ser	Lys	Gly	Val	Tyr	Cys
			260					265					270		
Leu	Gln	Tyr	Asp	Asp	Gln	Lys	Ile	Val	Ser	Gly	Leu	Arg	Asp	Asn	Thr
		275					280					285			
Ile	Lys	Ile	Trp	Asp	Lys	Asn	Thr	Leu	Glu	Cys	Lys	Arg	Ile	Leu	Thr
	290					295					300				
Gly	His	Thr	Gly	Ser	Val	Leu	Cys	Leu	Gln	Tyr	Asp	Glu	Arg	Val	Ile
305					310					315					320
Ile	Thr	Gly	Ser	Ser	Asp	Ser	Thr	Val	Arg	Val	Trp	Asp	Val	Asn	Thr
			325					330						335	
Gly	Glu	Met	Leu	Asn	Thr	Leu	Ile	His	His	Cys	Glu	Ala	Val	Leu	His
			340				345						350		
Leu	Arg	Phe	Asn	Asn	Gly	Met	Met	Val	Thr	Cys	Ser	Lys	Asp	Arg	Ser
		355					360					365			
Ile	Ala	Val	Trp	Asp	Met	Ala	Ser	Pro	Thr	Asp	Ile	Thr	Leu	Arg	Arg
	370					375					380				
Val	Leu	Val	Gly	His	Arg	Ala	Ala	Val	Asn	Val	Val	Asp	Phe	Asp	Asp
385					390					395					400
Lys	Tyr	Ile	Val	Ser	Ala	Ser	Gly	Asp	Arg	Thr	Ile	Lys	Val	Trp	Asn
			405					410					415		
Thr	Ser	Thr	Cys	Glu	Phe	Val	Arg	Thr	Leu	Asn	Gly	His	Lys	Arg	Gly
			420					425					430		
Ile	Ala	Cys	Leu	Gln	Tyr	Arg	Asp	Arg	Leu	Val	Val	Ser	Gly	Ser	Ser
		435					440					445			
Asp	Asn	Thr	Ile	Arg	Leu	Trp	Asp	Ile	Glu	Cys	Gly	Ala	Cys	Leu	Arg
	450					455					460				
Val	Leu	Glu	Gly	His	Glu	Glu	Leu	Val	Arg	Cys	Ile	Arg	Phe	Asp	Asn
465					470					475					480
Lys	Arg	Ile	Val	Ser	Gly	Ala	Tyr	Asp	Gly	Lys	Ile	Lys	Val	Trp	Asp
			485					490					495		
Leu	Val	Ala	Ala	Leu	Asp	Pro	Arg	Ala	Pro	Ala	Gly	Thr	Leu	Cys	Leu
		500					505						510		
Arg	Thr	Leu	Val	Glu	His	Ser	Gly	Arg	Val	Phe	Arg	Leu	Gln	Phe	Asp
		515					520					525			
Glu	Phe	Gln	Ile	Val	Ser	Ser	Ser	His	Asp	Asp	Thr	Ile	Leu	Ile	Trp
	530					535					540				
Asp	Phe	Leu	Asn	Asp	Pro	Ala	Ala	Gln	Ala	Glu	Pro	Pro	Arg	Ser	Pro
545					550					555					560
Ser	Arg	Thr	Tyr	Thr	Tyr	Ile	Ser	Arg							
			565												

<210> SEQ ID NO 3

<211> LENGTH: 1476

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

atggagagaa aggactttga gacatggctt gataacattt ctgttacatt tctttctctg	60
acggacttgc agaaaaatga aactctggat cacctgatta gtctgagtgg ggcagtccag	120
ctcaggcatc tctccaataa cctagagact ctcctcaagc gggacttccct caaactcctt	180

-continued

```

ccccctggagc tcagttttta ttgttataaa tggctcgatc ctcagacttt actcacatgc 240
tgccctcgtct ctaaacacgtg gaataaggtg ataagtcctt gtacagaggt gtggcagact 300
gcatgtaaaa atttgggctg gcagatagat gattctgttc aggacgcttt gcactggaag 360
aagggtttatt tgaaggctat ttgtagaatg aagcaactgg aggaccatga agcctttgaa 420
acctcgatcat taattggaca cagtgccaga gtgtatgcac tttactacaa agatggactt 480
ctctgtacag ggtcagatga cttgtctgca aagctgtggg atgtgagcac agggcagtgc 540
gtttatggca tccagaccca cacttgtgca gcggtgaagt ttgatgaaca gaagcttgtg 600
acaggctcct ttgacaacac tgtggcttgc tgggaatgga gttccggagc caggacccag 660
cactttcggg gccacacggg gccggtatct agcgtggact acaatgatga actggatatc 720
ttggtgagcg gctctgcaga cttcactgtg aaagtatggg ctttatctgc tgggacatgc 780
ctgaacacac tcaccgggca cacggaatgg gtcaccaagg tagttttgca gaagtgcata 840
gtcaagtctc tcttgacacg tcctggagac tacatcctct taagtgcaga caaatatgag 900
attaagattt ggccaattgg gagagaaatc aactgtaagt gcttaaagac attgtctgtc 960
tctgaggata gaagtatctg cctgcagcca agacttcatt ttgatggcaa atacattgtc 1020
tgtagtctag cacttggctc ctaccagtgg gactttgcca gttatgatat tctcagggtc 1080
atcaagactc ctgagatagc aaacttggcc ttgcttggct ttggagatat ctttgccttg 1140
ctgtttgaca accgctacct gtacatcatg gacttgcgga cagagagcct gattagtgcg 1200
tggcctctgc cagagtacag ggaatcaaag agaggctcaa gcttcctggc aggcgaacat 1260
cctggctgaa tggactggat gggcacaaatg acacgggctt ggtctttgcc accagcatgc 1320
ctgaccacag tattcacctg gtgttgtgga aggagcacgg ctgacaccat gagccaccac 1380
cgctgactga ctttgggtgc cggggctgcg ggttttgggt gcacctctgc ggcacgcgac 1440
tgcatagaac aaagtcttca cctaattgta tcatca 1476

```

<210> SEQ ID NO 4

<211> LENGTH: 422

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

```

Met Glu Arg Lys Asp Phe Glu Thr Trp Leu Asp Asn Ile Ser Val Thr
 1             5             10            15
Phe Leu Ser Leu Thr Asp Leu Gln Lys Asn Glu Thr Leu Asp His Leu
20            25            30
Ile Ser Leu Ser Gly Ala Val Gln Leu Arg His Leu Ser Asn Asn Leu
35            40            45
Glu Thr Leu Leu Lys Arg Asp Phe Leu Lys Leu Leu Pro Leu Glu Leu
50            55            60
Ser Phe Tyr Leu Leu Lys Trp Leu Asp Pro Gln Thr Leu Leu Thr Cys
65            70            75            80
Cys Leu Val Ser Lys Gln Trp Asn Lys Val Ile Ser Ala Cys Thr Glu
85            90            95
Val Trp Gln Thr Ala Cys Lys Asn Leu Gly Trp Gln Ile Asp Asp Ser
100           105           110
Val Gln Asp Ala Leu His Trp Lys Lys Val Tyr Leu Lys Ala Ile Leu
115           120           125

```

-continued

Arg Met Lys Gln Leu Glu Asp His Glu Ala Phe Glu Thr Ser Ser Leu
 130 135 140
 Ile Gly His Ser Ala Arg Val Tyr Ala Leu Tyr Tyr Lys Asp Gly Leu
 145 150 155 160
 Leu Cys Thr Gly Ser Asp Asp Leu Ser Ala Lys Leu Trp Asp Val Ser
 165 170 175
 Thr Gly Gln Cys Val Tyr Gly Ile Gln Thr His Thr Cys Ala Ala Val
 180 185 190
 Lys Phe Asp Glu Gln Lys Leu Val Thr Gly Ser Phe Asp Asn Thr Val
 195 200 205
 Ala Cys Trp Glu Trp Ser Ser Gly Ala Arg Thr Gln His Phe Arg Gly
 210 215 220
 His Thr Gly Ala Val Phe Ser Val Asp Tyr Asn Asp Glu Leu Asp Ile
 225 230 235 240
 Leu Val Ser Gly Ser Ala Asp Phe Thr Val Lys Val Trp Ala Leu Ser
 245 250 255
 Ala Gly Thr Cys Leu Asn Thr Leu Thr Gly His Thr Glu Trp Val Thr
 260 265 270
 Lys Val Val Leu Gln Lys Cys Lys Val Lys Ser Leu Leu His Ser Pro
 275 280 285
 Gly Asp Tyr Ile Leu Leu Ser Ala Asp Lys Tyr Glu Ile Lys Ile Trp
 290 295 300
 Pro Ile Gly Arg Glu Ile Asn Cys Lys Cys Leu Lys Thr Leu Ser Val
 305 310 315 320
 Ser Glu Asp Arg Ser Ile Cys Leu Gln Pro Arg Leu His Phe Asp Gly
 325 330 335
 Lys Tyr Ile Val Cys Ser Ser Ala Leu Gly Leu Tyr Gln Trp Asp Phe
 340 345 350
 Ala Ser Tyr Asp Ile Leu Arg Val Ile Lys Thr Pro Glu Ile Ala Asn
 355 360 365
 Leu Ala Leu Leu Gly Phe Gly Asp Ile Phe Ala Leu Leu Phe Asp Asn
 370 375 380
 Arg Tyr Leu Tyr Ile Met Asp Leu Arg Thr Glu Ser Leu Ile Ser Arg
 385 390 395 400
 Trp Pro Leu Pro Glu Tyr Arg Glu Ser Lys Arg Gly Ser Ser Phe Leu
 405 410 415
 Ala Gly Glu His Pro Gly
 420

<210> SEQ ID NO 5

<211> LENGTH: 1407

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

```

cggggtggtg tgtgggggaa gccgccccg gcagcaggat gaaacgagga ggaagagata    60
gtgaccgtaa ttcacagaa gaaggaactg cagagaaatc caagaaactg aggactacaa    120
atgagcattc tcagacttgt gattggggtg atctccttca ggacattatt ctccaagtat    180
ttaaatattt gcctcttctt gaccgggctc atgcttcaca agtttgccgc aactggaacc    240
aggatatttca catgcctgac ttgtggagat gttttgaatt tgaactgaat cagccagcta    300
catcttattt gaaagctacc catccagagc tgatcaaaca gattattaaa agacattcaa    360

```

-continued

```

accatctaca atatgtcagc ttcaaggtag acagcagcaa ggaatcagct gaagcagctt 420
gtgatatact atcgcaactt gtgaattgct ctttaaaaac acttggaactt atttcaactg 480
ctcgaccaag ctttatggat ttaccaaagt ctcactttat ctctgcactg acagttgtgt 540
tcgtaaaactc caaatccctg tcttcgctta agatagatga tactccagta gatgatccat 600
ctctcaaagt actagtggcc aacaatagtg atacactcaa gctgttgaaa atgagcagct 660
gtcctcatgt ctctccagca ggtatccttt gtgtggctga tcagtgtcac ggcttaagag 720
aactagccct gaactaccac ttattgagtg atgagttggt acttgcaattg tcttctgaaa 780
aacatgttcg attagaacat ttgcgcattg atgtagtcag tgagaatcct ggacagacac 840
acttccatac tattcagaag agtagctggg atgctttcat cagacattca ccaaagtga 900
acttagtgat gtattttttt ttatatgaag aagaatttga ccccttcttt cgctatgaaa 960
tacctgccac ccattctgtac ttggggagat cagtaagcaa agatgtgctt ggcctgtgtg 1020
gaatgacatg ccctagactg gttgaactag tagtgtgtgc aaatggatta cgccacttg 1080
atgaagagtt aattcgcatg gcagaacgtt gcaaaaattt gtcagctatt ggactagggg 1140
aatgtgaagt ctcatgtagt gcctttgttg agtttgtgaa gatgtgtggt ggccgcctat 1200
ctcaattatc cattatggaa gaagtactaa ttcctgacca aaagtatagt ttggagcaga 1260
ttcactggga agtgtccaag catcttggtt ggggtgtggt tcccgacatg atgccactt 1320
ggtaaaaact gcatgatgaa tagcacctta atttcaagca aatgtattat aattaaagtt 1380
ttatttgctg taaaaaaaaa aaaaaaa 1407

```

<210> SEQ ID NO 6

<211> LENGTH: 428

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

```

Met Lys Arg Gly Gly Arg Asp Ser Asp Arg Asn Ser Ser Glu Glu Gly
  1             5             10            15

Thr Ala Glu Lys Ser Lys Lys Leu Arg Thr Thr Asn Glu His Ser Gln
      20            25            30

Thr Cys Asp Trp Gly Asn Leu Leu Gln Asp Ile Ile Leu Gln Val Phe
      35            40            45

Lys Tyr Leu Pro Leu Leu Asp Arg Ala His Ala Ser Gln Val Cys Arg
      50            55            60

Asn Trp Asn Gln Val Phe His Met Pro Asp Leu Trp Arg Cys Phe Glu
      65            70            75            80

Phe Glu Leu Asn Gln Pro Ala Thr Ser Tyr Leu Lys Ala Thr His Pro
      85            90            95

Glu Leu Ile Lys Gln Ile Ile Lys Arg His Ser Asn His Leu Gln Tyr
      100           105           110

Val Ser Phe Lys Val Asp Ser Ser Lys Glu Ser Ala Glu Ala Ala Cys
      115           120           125

Asp Ile Leu Ser Gln Leu Val Asn Cys Ser Leu Lys Thr Leu Gly Leu
      130           135           140

Ile Ser Thr Ala Arg Pro Ser Phe Met Asp Leu Pro Lys Ser His Phe
      145           150           155           160

Ile Ser Ala Leu Thr Val Val Phe Val Asn Ser Lys Ser Leu Ser Ser

```


-continued

165					170					175					
Leu	Lys	Ile	Asp	Asp	Thr	Pro	Val	Asp	Asp	Pro	Ser	Leu	Lys	Val	Leu
			180					185					190		
Val	Ala	Asn	Asn	Ser	Asp	Thr	Leu	Lys	Leu	Leu	Lys	Met	Ser	Ser	Cys
		195					200					205			
Pro	His	Val	Ser	Pro	Ala	Gly	Ile	Leu	Cys	Val	Ala	Asp	Gln	Cys	His
	210					215					220				
Gly	Leu	Arg	Glu	Leu	Ala	Leu	Asn	Tyr	His	Leu	Leu	Ser	Asp	Glu	Leu
225					230					235					240
Leu	Leu	Ala	Leu	Ser	Ser	Glu	Lys	His	Val	Arg	Leu	Glu	His	Leu	Arg
			245						250					255	
Ile	Asp	Val	Val	Ser	Glu	Asn	Pro	Gly	Gln	Thr	His	Phe	His	Thr	Ile
		260						265					270		
Gln	Lys	Ser	Ser	Trp	Asp	Ala	Phe	Ile	Arg	His	Ser	Pro	Lys	Val	Asn
		275					280					285			
Leu	Val	Met	Tyr	Phe	Phe	Leu	Tyr	Glu	Glu	Glu	Phe	Asp	Pro	Phe	Phe
	290					295					300				
Arg	Tyr	Glu	Ile	Pro	Ala	Thr	His	Leu	Tyr	Phe	Gly	Arg	Ser	Val	Ser
305					310					315					320
Lys	Asp	Val	Leu	Gly	Arg	Val	Gly	Met	Thr	Cys	Pro	Arg	Leu	Val	Glu
			325					330						335	
Leu	Val	Val	Cys	Ala	Asn	Gly	Leu	Arg	Pro	Leu	Asp	Glu	Glu	Leu	Ile
		340					345						350		
Arg	Ile	Ala	Glu	Arg	Cys	Lys	Asn	Leu	Ser	Ala	Ile	Gly	Leu	Gly	Glu
	355						360					365			
Cys	Glu	Val	Ser	Cys	Ser	Ala	Phe	Val	Glu	Phe	Val	Lys	Met	Cys	Gly
	370					375					380				
Gly	Arg	Leu	Ser	Gln	Leu	Ser	Ile	Met	Glu	Glu	Val	Leu	Ile	Pro	Asp
385					390				395						400
Gln	Lys	Tyr	Ser	Leu	Glu	Gln	Ile	His	Trp	Glu	Val	Ser	Lys	His	Leu
			405					410					415		
Gly	Arg	Val	Trp	Phe	Pro	Asp	Met	Met	Pro	Thr	Trp				
	420						425								

<210> SEQ ID NO 7

<211> LENGTH: 1444

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

atggcgaggaa gcgagccgcg cagcggaaca aattcgccgc cgccgccctt cagcgactgg	60
ggccgcctgg aggcggccat cctcagcggc tggaagacct tctggcagtc agtgagcaag	120
gatagggtgg cggtacgac ctcccgggag gaggtggatg aggcggccag caccctgacg	180
cgggtgccga ttgatgtaca gctatatatt ttgtcctttc tttcacctca tgatctgtgt	240
cagttgggaa gtacaaatca ttattggaat gaaactgtaa gaaatccaat tctgtggaga	300
tactttttgt tgagggatct tccttcttgg tcttctgttg actggaagtc tcttccatat	360
ctacaaatct taaaaaagcc tatatctgag gtctctgatg gtgcattttt tgactacatg	420
gcagtctatc taatgtgctg tccatacaca agaagagctt caaaatccag ccgtcctatg	480
tatggagctg tcacttcttt ttacactcc ctgatcattc ccaatgaacc tcgatttgct	540

-continued

```

ctgtttggac cacgtttgga acaattgaat acctctttgg tgttgagctt gctgtcttca 600
gaggaacttt gcccaacagc tggtttgcc cagaggcaga ttgatggat tggtacagga 660
gtcaattttc agttgaacaa ccaacataaa ttcaacattc taatcttata ttcaactacc 720
agaaaggaaa gagatagagc aagggaagag catacaagt cagttaacaa gatgttcagt 780
cgacacaatg aagggtgatg tcgaccagga agccgggtaca gtgtgattcc acagattcaa 840
aaactgtgtg aagttgtaga tgggttcac tatgttgcaa atgctgaagc tcataaaaga 900
catgaatggc aagatgaatt ttctcatatt atggcaatga cagatccagc ctttgggtct 960
tcgggaagac cattgttgtt tttatcttgt atttctcaag gggatgtaaa aagaatgcc 1020
tgtttttatt tggctcatga gctgcatctg aatcttctaa atcaccatg gctgggtccag 1080
gatacagagg ctgaaactct gactgggttt ttgaatggca ttgagtggat tcttgaagaa 1140
gtggaatcta agcgtgcaag atgattctct tttcagatct tgggaactga aaccatttga 1200
aatttattac taagtgctg atgtgaatat ttgctcagtc agcccacctt gtccctgcctt 1260
tttgagata ggctttcatt tggacagcta taactgctgt gttttttata ttatttttac 1320
tttttaccat aaatcaatta caagaaaaga gtttcagtcc tagtatttag ccccaaatg 1380
aacctttaa catttttttg gtaattttta tttttctgt ctttttaaaa atattaaatt 1440
ttgg 1444

```

<210> SEQ ID NO: 8

<211> LENGTH: 472

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

```

Met Ala Gly Ser Glu Pro Arg Ser Gly Thr Asn Ser Pro Pro Pro Pro
 1           5           10          15
Phe Ser Asp Trp Gly Arg Leu Glu Ala Ala Ile Leu Ser Gly Trp Lys
 20          25          30
Thr Phe Trp Gln Ser Val Ser Lys Asp Arg Val Ala Arg Thr Thr Ser
 35          40          45
Arg Glu Glu Val Asp Glu Ala Ala Ser Thr Leu Thr Arg Leu Pro Ile
 50          55          60
Asp Val Gln Leu Tyr Ile Leu Ser Phe Leu Ser Pro His Asp Leu Cys
 65          70          75          80
Gln Leu Gly Ser Thr Asn His Tyr Trp Asn Glu Thr Val Arg Asn Pro
 85          90          95
Ile Leu Trp Arg Tyr Phe Leu Leu Arg Asp Leu Pro Ser Trp Ser Ser
100         105         110
Val Asp Trp Lys Ser Leu Pro Tyr Leu Gln Ile Leu Lys Lys Pro Ile
115         120         125
Ser Glu Val Ser Asp Gly Ala Phe Phe Asp Tyr Met Ala Val Tyr Leu
130         135         140
Met Cys Cys Pro Tyr Thr Arg Arg Ala Ser Lys Ser Ser Arg Pro Met
145         150         155         160
Tyr Gly Ala Val Thr Ser Phe Leu His Ser Leu Ile Ile Pro Asn Glu
165         170         175
Pro Arg Phe Ala Leu Phe Gly Pro Arg Leu Glu Gln Leu Asn Thr Ser
180         185         190

```

-continued

Leu Val Leu Ser Leu Leu Ser Ser Glu Glu Leu Cys Pro Thr Ala Gly
 195 200 205
 Leu Pro Gln Arg Gln Ile Asp Gly Ile Gly Ser Gly Val Asn Phe Gln
 210 215 220
 Leu Asn Asn Gln His Lys Phe Asn Ile Leu Ile Leu Tyr Ser Thr Thr
 225 230 235 240
 Arg Lys Glu Arg Asp Arg Ala Arg Glu Glu His Thr Ser Ala Val Asn
 245 250 255
 Lys Met Phe Ser Arg His Asn Glu Gly Asp Asp Arg Pro Gly Ser Arg
 260 265 270
 Tyr Ser Val Ile Pro Gln Ile Gln Lys Leu Cys Glu Val Val Asp Gly
 275 280 285
 Phe Ile Tyr Val Ala Asn Ala Glu Ala His Lys Arg His Glu Trp Gln
 290 295 300
 Asp Glu Phe Ser His Ile Met Ala Met Thr Asp Pro Ala Phe Gly Ser
 305 310 315 320
 Ser Gly Arg Pro Leu Leu Val Leu Ser Cys Ile Ser Gln Gly Asp Val
 325 330 335
 Lys Arg Met Pro Cys Phe Tyr Leu Ala His Glu Leu His Leu Asn Leu
 340 345 350
 Leu Asn His Pro Trp Leu Val Gln Asp Thr Glu Ala Glu Thr Leu Thr
 355 360 365
 Gly Phe Leu Asn Gly Ile Glu Trp Ile Leu Glu Glu Val Glu Ser Lys
 370 375 380
 Arg Ala Arg Phe Ser Phe Gln Ile Leu Gly Thr Glu Thr Ile Asn Leu
 385 390 395 400
 Leu Leu Arg Ser Cys Glu Tyr Leu Leu Ser Gln Pro Thr Leu Ser Cys
 405 410 415
 Leu Phe Ala Asp Arg Leu Ser Phe Gly Gln Leu Leu Leu Cys Phe Leu
 420 425 430
 Tyr Tyr Phe Tyr Phe Leu Pro Ile Asn Tyr Lys Lys Arg Val Ser Val
 435 440 445
 Leu Val Phe Ser Pro Lys Met Asn Leu Thr Phe Phe Trp Phe Leu Tyr
 450 455 460
 Phe Leu Ser Phe Lys Tyr Ile Leu
 465 470

<210> SEQ ID NO 9

<211> LENGTH: 2076

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

```

aggttgctca gctgcccccg gagcggttcc tccacctgag gcagacacca cctcggttg 60
catgagccgg cgccctgca gctgcgcct acggccaccc cgctgctoct gcagcgccag 120
ccccagcgca gtgacagccg ccggggcgccc tcgaccctcg gatagttgta aagaagaaag 180
ttctaccctt tctgtcaaaa tgaagtgtga ttttaattgt aaccatgttc attccggact 240
taaactggta aaacctgatg acattggaag actagtttcc tacaccctcg catatctgga 300
aggttcctgt aaagactgca ttaaagacta tgaaaggctg tcattgtattg ggtcaccgat 360
tgtgagccct aggattgtac aacttgaaac tgaaagcaag cgcttgcata acaaggaaaa 420

```

-continued

```

tcaacatgtg caacagacac ttaatagtag aaatgaaata gaagcactag agaccagtag 480
actttatgaa gacagtggct attcctcatt ttctctacaa agtggcctca gtgaacatga 540
agaaggtagc ctccctggagg agaatttcgg tgacagtcta caatcctgcc tgctacaaat 600
acaaagccca gaccaatata ccaacaaaaa cttgctgccca gttcttcatt ttgaaaaagt 660
ggtttgttca acattaaaaa agaatgcaaa acgaaatcct aaagtagatc gggagatgct 720
gaaggaaatt atagccagag gaaattttag actgcagaat ataattggca gaaaaatggg 780
cctagaatgt gtagatattc tcagcgaact ctttcgaagg ggactcagac atgtcttagc 840
aactatttta gcacaactca gtgacatgga cttaatcaat gtgtctaaag tgagcacaac 900
ttggaagaag atcctagaag atgataaggg ggcattccag ttgtacagta aagcaataca 960
aagagttacc gaaaacaaca ataaattttc acctcatgct tcaaccagag aatatgttat 1020
gttcagaacc ccactggcct ctgttcagaa atcagcagcc cagactttct tcaaaaaaga 1080
tgctcaaacc aagttatcca atcaagggtg tcagaaaggt tctacttata gtcgacacaa 1140
tgaattctct gaggttgcca agacattgaa aaagaacgaa agcctcaaag cctgtattcg 1200
ctgtaattca cctgcaaaat atgattgcta tttacaacgg gcaacctgca aacgagaagg 1260
ctgtggattt gattattgta cgaagtgtct ctgtaattat catactacta aagactgttc 1320
agatggcaag ctccctcaaag ccagttgtaa aataggtccc ctgcctggta caaagaaaag 1380
caaaaagaat ttacgaagat tgtgatctct tattaatat attgttactg atcatgaatg 1440
ttagttagaa aatgttaggt tttaacttaa aaaaaattgt attgtgattt tcaattttat 1500
gttgaatcgt gtgtagtata ctgaggtttt tttcccccca gaagataaag aggatagaca 1560
acctcttaaa atattttttac aatttaatga gaaaaagttt aaaattctca atacaaatca 1620
aacaatttaa atattttaag aaaaaggaa aagtagatag tgatactgag ggtaaaaaaa 1680
aaattgattc aattttatgg taaaggaaac ccatgcaatt ttacctagac agtcttaaat 1740
atgtctgggt ttccatctgt tagcatttca gacattttat gttcctctta ctcaattgat 1800
accaacagaa atatcaactt ctggagtcta ttaaatgtgt tgtcaccttt ctaaagcttt 1860
ttttcattgt gtgtatttcc caagaaagta tcctttgtaa aaacttgctt gttttcctta 1920
tttctgaaat ctgttttaat atttttgtat acatgtaaat atttctgtat tttttatatg 1980
tcaaagaata tgtctcttgt atgtacatat aaaaataaat tttgctcaat aaaattgtaa 2040
gcttaaaaaa aaaaaaaaaa aactcgagac tagtgc 2076

```

<210> SEQ ID NO 10

<211> LENGTH: 447

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

```

Met Ser Arg Arg Pro Cys Ser Cys Ala Leu Arg Pro Pro Arg Cys Ser
 1             5             10             15
Cys Ser Ala Ser Pro Ser Ala Val Thr Ala Ala Gly Arg Pro Arg Pro
          20             25             30
Ser Asp Ser Cys Lys Glu Glu Ser Ser Thr Leu Ser Val Lys Met Lys
          35             40             45
Cys Asp Phe Asn Cys Asn His Val His Ser Gly Leu Lys Leu Val Lys
          50             55             60

```

-continued

Pro	Asp	Asp	Ile	Gly	Arg	Leu	Val	Ser	Tyr	Thr	Pro	Ala	Tyr	Leu	Glu	65	70	75	80
Gly	Ser	Cys	Lys	Asp	Cys	Ile	Lys	Asp	Tyr	Glu	Arg	Leu	Ser	Cys	Ile	85	90	95	
Gly	Ser	Pro	Ile	Val	Ser	Pro	Arg	Ile	Val	Gln	Leu	Glu	Thr	Glu	Ser	100	105	110	
Lys	Arg	Leu	His	Asn	Lys	Glu	Asn	Gln	His	Val	Gln	Gln	Thr	Leu	Asn	115	120	125	
Ser	Thr	Asn	Glu	Ile	Glu	Ala	Leu	Glu	Thr	Ser	Arg	Leu	Tyr	Glu	Asp	130	135	140	
Ser	Gly	Tyr	Ser	Ser	Phe	Ser	Leu	Gln	Ser	Gly	Leu	Ser	Glu	His	Glu	145	150	155	160
Glu	Gly	Ser	Leu	Leu	Glu	Glu	Asn	Phe	Gly	Asp	Ser	Leu	Gln	Ser	Cys	165	170	175	
Leu	Leu	Gln	Ile	Gln	Ser	Pro	Asp	Gln	Tyr	Pro	Asn	Lys	Asn	Leu	Leu	180	185	190	
Pro	Val	Leu	His	Phe	Glu	Lys	Val	Val	Cys	Ser	Thr	Leu	Lys	Lys	Asn	195	200	205	
Ala	Lys	Arg	Asn	Pro	Lys	Val	Asp	Arg	Glu	Met	Leu	Lys	Glu	Ile	Ile	210	215	220	
Ala	Arg	Gly	Asn	Phe	Arg	Leu	Gln	Asn	Ile	Ile	Gly	Arg	Lys	Met	Gly	225	230	235	240
Leu	Glu	Cys	Val	Asp	Ile	Leu	Ser	Glu	Leu	Phe	Arg	Arg	Gly	Leu	Arg	245	250	255	
His	Val	Leu	Ala	Thr	Ile	Leu	Ala	Gln	Leu	Ser	Asp	Met	Asp	Leu	Ile	260	265	270	
Asn	Val	Ser	Lys	Val	Ser	Thr	Thr	Trp	Lys	Lys	Ile	Leu	Glu	Asp	Asp	275	280	285	
Lys	Gly	Ala	Phe	Gln	Leu	Tyr	Ser	Lys	Ala	Ile	Gln	Arg	Val	Thr	Glu	290	295	300	
Asn	Asn	Asn	Lys	Phe	Ser	Pro	His	Ala	Ser	Thr	Arg	Glu	Tyr	Val	Met	305	310	315	320
Phe	Arg	Thr	Pro	Leu	Ala	Ser	Val	Gln	Lys	Ser	Ala	Ala	Gln	Thr	Ser	325	330	335	
Leu	Lys	Lys	Asp	Ala	Gln	Thr	Lys	Leu	Ser	Asn	Gln	Gly	Asp	Gln	Lys	340	345	350	
Gly	Ser	Thr	Tyr	Ser	Arg	His	Asn	Glu	Phe	Ser	Glu	Val	Ala	Lys	Thr	355	360	365	
Leu	Lys	Lys	Asn	Glu	Ser	Leu	Lys	Ala	Cys	Ile	Arg	Cys	Asn	Ser	Pro	370	375	380	
Ala	Lys	Tyr	Asp	Cys	Tyr	Leu	Gln	Arg	Ala	Thr	Cys	Lys	Arg	Glu	Gly	385	390	395	400
Cys	Gly	Phe	Asp	Tyr	Cys	Thr	Lys	Cys	Leu	Cys	Asn	Tyr	His	Thr	Thr	405	410	415	
Lys	Asp	Cys	Ser	Asp	Gly	Lys	Leu	Leu	Lys	Ala	Ser	Cys	Lys	Ile	Gly	420	425	430	
Pro	Leu	Pro	Gly	Thr	Lys	Lys	Ser	Lys	Lys	Asn	Leu	Arg	Arg	Leu		435	440	445	

<210> SEQ ID NO 11

<211> LENGTH: 1535

<212> TYPE: DNA

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

```

gcgcgttcgg gagcttcggc cctgcgtagg aggcgggtgc aggtgtgggt gctgagccgc   60
ccgccgcctg gagggggaga cagcttcagg acacgcaggc cgcagcgagg gcccgggccc   120
gggggatccc aggccatgga cgctccccac tccaaagcag ccctggacag cattaacgag   180
ctgcccgata acatcctgct ggagctgttc acgcacgtgc ccgccgccca gctgctgctg   240
aactgccgcc tgggtctgca cctctggcgg gacctcatcg acctcctgac cctctggaaa   300
cgcaagtgcc tgcgaaaggg cttcatcacc aaggactggg accagcccggt ggccgactgg   360
aaaaatctct acttcctacg gagcctgcat aggaacctcc tgcgcaaccc gtgtgtgtaa   420
aacgatatgt ttgcattgga aattgatttc aatgggtggg accgctggaa ggtggatagc   480
ctccctggag cccacgggac agaatttcct gaccccaaag tcaagaagtc ttttgtcaca   540
tcctacgaac tgtgcctcaa gtgggagctg gtggaccttc tagccgaccg ctactgggag   600
gagctactag acacattccg gccggacatc gtggttaagg actggtttgc tgccagagcc   660
gactgtggct gcacctacca actcaaagtg cagctggcct cggctgacta ctctgtgttg   720
gcctccttcg agccccacc tgtgaccatc caacagtgga acaatgccac atggacagag   780
gtctcctaca ctttctcaga ctacccccgg ggtgtccgct acatcctctt ccagcatggg   840
ggcagggaca cccagtactg ggcaggctgg tatgggcccc gagtccacaa cagcagcatt   900
gtcgtcagcc ccaagatgac caggaaccag gcctcgtccg aggtctagcc tgggcagaag   960
catggacagg agggaggtgc ccaatcgccc tacggagctg ttgtccagat tttctgacag  1020
ctgtccatcc tgtgtctggg tcagccagag gttcctccag gcaggagctg agcatggggg  1080
gggcagttag gtccctgtac cagcgactcc tgccccggtt caaccctacc agcttgtggg  1140
aacttactgt cacatagctc tgacgttttg ttgtaataaa tgttttcagg ccgggcactg  1200
tggtctcacg ctgtaatccc agcacttttg gagaccgagg caggtggatc acgaggtcag  1260
gagacagaga ccatcctggc caacacgggtg aaaccctgtc tctactaaaa atacaaaaaa  1320
ttagccgggc gtggtggcgg gcgcctgtag tcccagctac tcgggagggt gatgcagaag  1380
aatggcgtga acccggaagg cagagcttgc agtgagccga gatcacgcca ctgcactcca  1440
gcctgggtga cagagcgaga ctctggctca taaaataata ataataataa ataaataaaa  1500
aataaatggt tttcagtaaa aaaaaaaaaa aaaaaa  1535

```

<210> SEQ ID NO 12

<211> LENGTH: 338

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

```

Ala Arg Ser Gly Ala Ser Ala Leu Arg Arg Arg Arg Val Gln Val Trp
 1             5             10             15
Val Leu Ser Arg Pro Pro Pro Gly Gly Gly Asp Ser Phe Arg Thr Arg
          20             25             30
Arg Pro Gln Arg Gly Pro Gly Pro Gly Gly Ser Gln Ala Met Asp Ala
          35             40             45
Pro His Ser Lys Ala Ala Leu Asp Ser Ile Asn Glu Leu Pro Asp Asn
          50             55             60

```

-continued

Ile	Leu	Leu	Glu	Leu	Phe	Thr	His	Val	Pro	Ala	Arg	Gln	Leu	Leu	Leu	65	70	75	80
Asn	Cys	Arg	Leu	Val	Cys	Ser	Leu	Trp	Arg	Asp	Leu	Ile	Asp	Leu	Leu	85	90	95	
Thr	Leu	Trp	Lys	Arg	Lys	Cys	Leu	Arg	Lys	Gly	Phe	Ile	Thr	Lys	Asp	100	105	110	
Trp	Asp	Gln	Pro	Val	Ala	Asp	Trp	Lys	Ile	Phe	Tyr	Phe	Leu	Arg	Ser	115	120	125	
Leu	His	Arg	Asn	Leu	Leu	Arg	Asn	Pro	Cys	Ala	Glu	Asn	Asp	Met	Phe	130	135	140	
Ala	Trp	Gln	Ile	Asp	Phe	Asn	Gly	Gly	Asp	Arg	Trp	Lys	Val	Asp	Ser	145	150	155	160
Leu	Pro	Gly	Ala	His	Gly	Thr	Glu	Phe	Pro	Asp	Pro	Lys	Val	Lys	Lys	165	170	175	
Ser	Phe	Val	Thr	Ser	Tyr	Glu	Leu	Cys	Leu	Lys	Trp	Glu	Leu	Val	Asp	180	185	190	
Leu	Leu	Ala	Asp	Arg	Tyr	Trp	Glu	Glu	Leu	Leu	Asp	Thr	Phe	Arg	Pro	195	200	205	
Asp	Ile	Val	Val	Lys	Asp	Trp	Phe	Ala	Ala	Arg	Ala	Asp	Cys	Gly	Cys	210	215	220	
Thr	Tyr	Gln	Leu	Lys	Val	Gln	Leu	Ala	Ser	Ala	Asp	Tyr	Phe	Val	Leu	225	230	235	240
Ala	Ser	Phe	Glu	Pro	Pro	Pro	Val	Thr	Ile	Gln	Gln	Trp	Asn	Asn	Ala	245	250	255	
Thr	Trp	Thr	Glu	Val	Ser	Tyr	Thr	Phe	Ser	Asp	Tyr	Pro	Arg	Gly	Val	260	265	270	
Arg	Tyr	Ile	Leu	Phe	Gln	His	Gly	Gly	Arg	Asp	Thr	Gln	Tyr	Trp	Ala	275	280	285	
Gly	Trp	Tyr	Gly	Pro	Arg	Val	Thr	Asn	Ser	Ser	Ile	Val	Val	Ser	Pro	290	295	300	
Lys	Met	Thr	Arg	Asn	Gln	Ala	Ser	Ser	Glu	Ala	Gln	Pro	Gly	Gln	Lys	305	310	315	320
His	Gly	Gln	Glu	Glu	Ala	Ala	Gln	Ser	Pro	Tyr	Gly	Ala	Val	Val	Gln	325	330	335	

Ile Phe

<210> SEQ ID NO 13

<211> LENGTH: 1763

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

tggaattccc atggaccatg tctaataccc gatttacaat tacattgaac tacaaggatc	60
ccctcactgg agatgaagag accttggtt catatgggat tgtttctggg gacttgatat	120
gtttgattct tcacgatgac attccaccgc ctaatatacc ttcattccaca gattcagagc	180
attttcact ccagaacaat gagcaaccct ctttgccac cagctccaat cagactagca	240
tacaggatga acaaccaagt gattcattcc aaggacaggc agcccagtct ggtgtttgga	300
atgacgacag tatgttaggg ctagtcaaa attttgaagc tgagtcaatt caagataatg	360
cgcataatgg agagggcaca ggtttctatc cctcagaacc cctgctctgt agtgaatcgg	420
tggaagggca agtgccacat tcattagaga ccttgatatc atcagctgac tgttctgatg	480

-continued

```

ccaatgatgc gttgatagtg ttgatacatc ttctcatggt ggagtcaggt tacatacctc 540
agggcaccga agccaaagca ctgtccctgc cggagaagtg gaagttgagc ggggtgtata 600
agctgcagta catgcatcat ctctgcgagg gcagctccgc tactctcacc tgtgtgcctt 660
tgggaaacct gattgttgta aatgctacac taaaaatcaa caatgagatt agaagtgtga 720
aaagattgca gctgctacca gaatctttta ttgcaaaga gaaactaggg gaaaatgtag 780
ccaacatata caaagatctt cagaaactct ctgcctctt taaagaccag ctggtgtatc 840
ctcttctggc ttttaccgca caagcactga acctaccaa tgtatttggg ttggtcgtcc 900
tcccattgga actgaaacta cggatcttcc gacttctgga tgttcgttcc gtcttgtctt 960
tgtctgcggt ttgtcgtgac ctctttactg cttcaaatga cccactcctg tggaggtttt 1020
tatatctgcy tgatttttca gacaatactg tcagagttca agacacagat tggaaagaac 1080
tgtacaggaa gaggcacata caaagaaaag aatccccgaa agggcgggtt gtgctgctcc 1140
tgccatcgtc aaccacacc attccattct atcccaacc cttgcacctt aggccatttc 1200
ctagctcccc cttctctcca ggaattatcg ggggtgaata tgaccaaaga ccaacacttc 1260
cctatgttgg agacccaatc agttcactca ttcttggtcc tggggagacg cccagccagt 1320
tacctccact gagaccacgc ttgatccag ttggccact tccaggacct aacccatct 1380
tgccagggcy aggcggcccc aatgacagat ttccctttag acccagcagc ggtcgccaa 1440
ctgatggccg cctgtcattc atgtgattga ttgttaattt catttctgga gctccatttg 1500
ttttgtttc taaactacag atgtcactcc ttggggtgct gatctcgagt gttattttct 1560
gattgtggtg ttgagagttg cactcccaga aaccttttaa gagatacatt tatagcccta 1620
ggggtggtat gacccaaagc ttctctgtg acaagggttg ccttggaat agttggctgc 1680
caatctccct gctcttggtt ctctctaga ttgaagtttg tttctgatg ctgttcttac 1740
cagattaaaa aaaagtgtaa att 1763

```

<210> SEQ ID NO 14

<211> LENGTH: 482

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

```

Met Ser Asn Thr Arg Phe Thr Ile Thr Leu Asn Tyr Lys Asp Pro Leu
  1             5             10             15
Thr Gly Asp Glu Glu Thr Leu Ala Ser Tyr Gly Ile Val Ser Gly Asp
          20             25             30
Leu Ile Cys Leu Ile Leu His Asp Asp Ile Pro Pro Pro Asn Ile Pro
          35             40             45
Ser Ser Thr Asp Ser Glu His Ser Ser Leu Gln Asn Asn Glu Gln Pro
          50             55             60
Ser Leu Ala Thr Ser Ser Asn Gln Thr Ser Ile Gln Asp Glu Gln Pro
          65             70             75             80
Ser Asp Ser Phe Gln Gly Gln Ala Ala Gln Ser Gly Val Trp Asn Asp
          85             90             95
Asp Ser Met Leu Gly Pro Ser Gln Asn Phe Glu Ala Glu Ser Ile Gln
          100            105            110
Asp Asn Ala His Met Ala Glu Gly Thr Gly Phe Tyr Pro Ser Glu Pro
          115            120            125

```


-continued

```

Leu Leu Cys Ser Glu Ser Val Glu Gly Gln Val Pro His Ser Leu Glu
 130          135          140

Thr Leu Tyr Gln Ser Ala Asp Cys Ser Asp Ala Asn Asp Ala Leu Ile
145          150          155          160

Val Leu Ile His Leu Leu Met Leu Glu Ser Gly Tyr Ile Pro Gln Gly
          165          170          175

Thr Glu Ala Lys Ala Leu Ser Leu Pro Glu Lys Trp Lys Leu Ser Gly
          180          185          190

Val Tyr Lys Leu Gln Tyr Met His His Leu Cys Glu Gly Ser Ser Ala
          195          200          205

Thr Leu Thr Cys Val Pro Leu Gly Asn Leu Ile Val Val Asn Ala Thr
          210          215          220

Leu Lys Ile Asn Asn Glu Ile Arg Ser Val Lys Arg Leu Gln Leu Leu
225          230          235          240

Pro Glu Ser Phe Ile Cys Lys Glu Lys Leu Gly Glu Asn Val Ala Asn
          245          250          255

Ile Tyr Lys Asp Leu Gln Lys Leu Ser Arg Leu Phe Lys Asp Gln Leu
          260          265          270

Val Tyr Pro Leu Leu Ala Phe Thr Arg Gln Ala Leu Asn Leu Pro Asn
          275          280          285

Val Phe Gly Leu Val Val Leu Pro Leu Glu Leu Lys Leu Arg Ile Phe
          290          295          300

Arg Leu Leu Asp Val Arg Ser Val Leu Ser Leu Ser Ala Val Cys Arg
305          310          315          320

Asp Leu Phe Thr Ala Ser Asn Asp Pro Leu Leu Trp Arg Phe Leu Tyr
          325          330          335

Leu Arg Asp Phe Arg Asp Asn Thr Val Arg Val Gln Asp Thr Asp Trp
          340          345          350

Lys Glu Leu Tyr Arg Lys Arg His Ile Gln Arg Lys Glu Ser Pro Lys
          355          360          365

Gly Arg Phe Val Leu Leu Leu Pro Ser Ser Thr His Thr Ile Pro Phe
          370          375          380

Tyr Pro Asn Pro Leu His Pro Arg Pro Phe Pro Ser Ser Arg Leu Pro
385          390          395          400

Pro Gly Ile Ile Gly Gly Glu Tyr Asp Gln Arg Pro Thr Leu Pro Tyr
          405          410          415

Val Gly Asp Pro Ile Ser Ser Leu Ile Pro Gly Pro Gly Glu Thr Pro
          420          425          430

Ser Gln Leu Pro Pro Leu Arg Pro Arg Phe Asp Pro Val Gly Pro Leu
          435          440          445

Pro Gly Pro Asn Pro Ile Leu Pro Gly Arg Gly Gly Pro Asn Asp Arg
          450          455          460

Phe Pro Phe Arg Pro Ser Arg Gly Arg Pro Thr Asp Gly Arg Leu Ser
465          470          475          480

Phe Met

```

<210> SEQ ID NO 15

<211> LENGTH: 43

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

-continued

Leu Pro Ala Arg Gly Leu Asp His Ile Ala Glu Asn Ile Leu Ser Tyr
1 5 10 15
Leu Asp Ala Lys Ser Leu Cys Ala Ala Glu Leu Val Cys Lys Glu Trp
20 25 30
Tyr Arg Val Thr Ser Asp Gly Met Leu Trp Lys
35 40

<210> SEQ ID NO 16
<211> LENGTH: 40
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

Leu Pro Leu Glu Leu Ser Phe Tyr Leu Leu Lys Trp Leu Asp Pro Gln
1 5 10 15
Thr Leu Leu Thr Cys Cys Leu Val Ser Lys Gln Trp Asn Lys Val Ile
20 25 30
Ser Ala Cys Thr Glu Val Trp Gln
35 40

<210> SEQ ID NO 17
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

Leu Leu Gln Asp Ile Ile Leu Gln Val Phe Lys Tyr Leu Pro Leu Leu
1 5 10 15
Asp Arg Ala His Ala Ser Gln Val Cys Arg Asn Trp Asn Gln Val Phe
20 25 30
His Met Pro Asp Leu Trp Arg
35

<210> SEQ ID NO 18
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

Leu Pro Ile Asp Val Gln Leu Tyr Ile Leu Ser Phe Leu Ser Pro His
1 5 10 15
Asp Leu Cys Gln Leu Gly Ser Thr Asn His Tyr Trp Asn Glu Thr Val
20 25 30
Arg Asn Pro Ile Leu Trp Arg
35

<210> SEQ ID NO 19
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

Leu Arg His Val Leu Ala Thr Ile Leu Ala Gln Leu Ser Asp Met Asp
1 5 10 15
Leu Ile Asn Val Ser Lys Val Ser Thr Thr Trp Lys Lys Ile Leu Glu
20 25 30
Asp Asp Lys Gly Ala Phe Gln

-continued

35

<210> SEQ ID NO 20
<211> LENGTH: 40
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

Leu Pro Asp Asn Ile Leu Leu Glu Leu Phe Thr His Val Pro Ala Arg
1 5 10 15
Gln Leu Leu Leu Asn Cys Arg Leu Val Cys Ser Leu Trp Arg Asp Leu
20 25 30
Ile Asp Leu Leu Thr Leu Trp Lys
35 40

<210> SEQ ID NO 21
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

Leu Pro Leu Glu Leu Lys Leu Arg Ile Phe Arg Leu Leu Asp Val Arg
1 5 10 15
Ser Val Leu Ser Leu Ser Ala Val Cys Arg Asp Leu Phe Thr Ala Ser
20 25 30
Asn Asp Pro Leu Leu Trp Arg
35

<210> SEQ ID NO 22
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22

Leu Pro Asp Glu Leu Leu Leu Gly Ile Phe Ser Cys Leu Cys Leu Pro
1 5 10 15
Glu Leu Leu Lys Val Ser Gly Val Cys Lys Arg Trp Tyr Arg Leu Ala
20 25 30
Ser Asp Glu Ser Leu Trp Gln
35

<210> SEQ ID NO 23
<211> LENGTH: 1323
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 23

acatttttcta atgtttacag aatgaagagg aacagtttat ctgttgagaa taaaattgtc 60
cagttgtcag gagcagcgaa acagccaaaa gttgggttct actcttctct caaccagact 120
catacacaca cggttcttct agactggggg agtttgccct accatgtagt attacaaatt 180
tttcagtatc ttcttttact agatcggggc tgtgcatctt ctgtatgtag gaggtggaat 240
gaagtttttc atatttctga cctttggaga aagtttgaat ttgaactgaa ccagtcagct 300
acttcattctt ttaagtccac tcatcctgat ctcatcagc agatcattaa aaagcatttt 360
gctcatcttc agtatgtcag cttaaggtt gacagtagcg ctgagtcagc agaagctgcc 420
tgtgatatac tctctcagct ggtaaattgt tccatccaga ccttgggctt gatttcaaca 480

-continued

```

gccaaagccaa gtttcatgaa tgtgtcggag tctcattttg tgtcagcact tacagttggt 540
tttatcaact caaaatcatt atcatcaatc aaaattgaag atacaccagt ggatgatcct 600
tcattgaaga ttcttgtggc caataatagt gacactctaa gactcccaa gatgagtagc 660
tgtcctcatg tttcatctga tggaattctt tgtgtagctg accgttgtca aggccttaga 720
gaactggcgt tgaattatta catcctaact gatgaacttt tccttgcaact ctcaagcgag 780
actcatgtta accttgaaca tcttcgaatt gatgttgtga gtgaaaatcc tggacagatt 840
aaatttcag ctgttaaaaa acacagttgg gatgcactta ttaaacattc ccctagagtt 900
aatgttggtta tgcacttctt tctatatgaa gaggaattcg agacgttctt caaagaagaa 960
acccctgtta ctcaccttta ttttggtcgt tcagtcagca aagtggtttt aggacgggta 1020
ggtctcaact gtcctcgact gattgagttta gtggtgtgtg ctaatgatct tcagcctctt 1080
gataatgaac ttatttgtat tgctgaacac tgtacaaacc taacagcctt gggcctcagc 1140
aaatgtgaag ttagctgcag tgccttcac aggtttgtta gactgtgtga gagaaggta 1200
acacagctct ctgtaatgga ggaagttttg atccctgatg aggattatag cctagatgaa 1260
attcacactg aagtctccaa atacctggga agagtatggt tccctgatgt gatgcctctc 1320
tgg 1323

```

<210> SEQ ID NO 24

<211> LENGTH: 434

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

```

Met Lys Arg Asn Ser Leu Ser Val Glu Asn Lys Ile Val Gln Leu Ser
 1             5             10             15
Gly Ala Ala Lys Gln Pro Lys Val Gly Phe Tyr Ser Ser Leu Asn Gln
          20             25             30
Thr His Thr His Thr Val Leu Leu Asp Trp Gly Ser Leu Pro His His
          35             40             45
Val Val Leu Gln Ile Phe Gln Tyr Leu Pro Leu Leu Asp Arg Ala Cys
          50             55             60
Ala Ser Ser Val Cys Arg Arg Trp Asn Glu Val Phe His Ile Ser Asp
          65             70             75             80
Leu Trp Arg Lys Phe Glu Phe Glu Leu Asn Gln Ser Ala Thr Ser Ser
          85             90             95
Phe Lys Ser Thr His Pro Asp Leu Ile Gln Gln Ile Ile Lys Lys His
          100            105            110
Phe Ala His Leu Gln Tyr Val Ser Phe Lys Val Asp Ser Ser Ala Glu
          115            120            125
Ser Ala Glu Ala Ala Cys Asp Ile Leu Ser Gln Leu Val Asn Cys Ser
          130            135            140
Ile Gln Thr Leu Gly Leu Ile Ser Thr Ala Lys Pro Ser Phe Met Asn
          145            150            155            160
Val Ser Glu Ser His Phe Val Ser Ala Leu Thr Val Val Phe Ile Asn
          165            170            175
Ser Lys Ser Leu Ser Ser Ile Lys Ile Glu Asp Thr Pro Val Asp Asp
          180            185            190
Pro Ser Leu Lys Ile Leu Val Ala Asn Asn Ser Asp Thr Leu Arg Leu

```

-continued

195	200	205
Pro Lys Met Ser Ser Cys	Pro His Val Ser Ser	Asp Gly Ile Leu Cys
210	215	220
Val Ala Asp Arg Cys	Gln Gly Leu Arg Glu Leu	Ala Leu Asn Tyr Tyr
225	230	235 240
Ile Leu Thr Asp Glu Leu Phe Leu Ala Leu Ser Ser Glu Thr His Val		
	245	250 255
Asn Leu Glu His Leu Arg Ile Asp Val Val Ser Glu Asn Pro Gly Gln		
	260	265 270
Ile Lys Phe His Ala Val Lys Lys His Ser Trp Asp Ala Leu Ile Lys		
	275	280 285
His Ser Pro Arg Val Asn Val Val Met His Phe Phe Leu Tyr Glu Glu		
	290	295 300
Glu Phe Glu Thr Phe Phe Lys Glu Glu Thr Pro Val Thr His Leu Tyr		
	305	310 315 320
Phe Gly Arg Ser Val Ser Lys Val Val Leu Gly Arg Val Gly Leu Asn		
	325	330 335
Cys Pro Arg Leu Ile Glu Leu Val Val Cys Ala Asn Asp Leu Gln Pro		
	340	345 350
Leu Asp Asn Glu Leu Ile Cys Ile Ala Glu His Cys Thr Asn Leu Thr		
	355	360 365
Ala Leu Gly Leu Ser Lys Cys Glu Val Ser Cys Ser Ala Phe Ile Arg		
	370	375 380
Phe Val Arg Leu Cys Glu Arg Arg Leu Thr Gln Leu Ser Val Met Glu		
	385	390 395 400
Glu Val Leu Ile Pro Asp Glu Asp Tyr Ser Leu Asp Glu Ile His Thr		
	405	410 415
Glu Val Ser Lys Tyr Leu Gly Arg Val Trp Phe Pro Asp Val Met Pro		
	420	425 430
Leu Trp		

<210> SEQ ID NO 25

<211> LENGTH: 1970

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

```

ggaaacgtca aaattgggat agtcggcagt tctggcccct gcagctggag gtaccctgag      60
ttctgagggg cgtagtgtctg tttctggtat tctcatcgcg gtcaccteta ccggtgtgga    120
caagtaaagt ttgaatcagc ttctccatgg cctgggcacc agttcccggc tgagccattt    180
tcctttttggc taaaagtccc cgcccagagg ccaattcgtc gcggcggcgg tggagatcgc    240
aggtcgctca ggcttcgaga tgggtcaagg gttgtggaga gtggtcagaa accagcagct    300
gcaacaagaa ggctacagtg agcaaggcta cctcaccaga gagcagagca ggagaatggc    360
tgcgagcaac atttctaaca ccaatcatcg taaacaagtc caaggaggca ttgacatata    420
tcattctttt aaggcaagga aatcgaaaaga acaggaagga ttcattaatt tggaaatgtt    480
gcctcctgag ctaagcttta ccatcttctc ctacotgaat gcaactgacc tttgcttgge    540
ttcatgtgtt tggcaggacc ttgcgaatga tgaacttctc tggcaagggt tgtgcaaadc    600
cacttggggt cactgttcca tatacaataa gaaccacact ttaggatattt cttttagaaa    660

```

-continued

```

aktgtatatg cagctggatg aaggcagcct cacctttaat gccaacccag atgagggagt 720
gaactacttt atgtccaagg gtatcctgga tgattcgcca aaggaaatag caaagtttat 780
cttctgtaca agaacactaa attggaaaaa actgagaatc tatcttgatg aaaggagaga 840
tgtcttggat gaccttgtaa cattgcataa ttttagaaat cagttcttgc caaatgcact 900
gagagaattt tttcgtcata tccatgcccc tgaagagcgt ggagagtatc ttgaaactct 960
tataacaaag ttctcacata gattctgtgc ttgcaaccct gatttaatgc gagaacttgg 1020
ccttagtcct gatgtgtct atgtactgtg ctactctttg attctacttt ccattgacct 1080
cactagccct catgtgaaga ataaatgtc aaaaaggga tttattcgaa ataccgcgtc 1140
cgctgctcaa aatatttagt aagattttgt agggcatctt tatgacaata tctaccttat 1200
tggccatgtg gctgcataaa aagcaccaatt gctaggactt cagtttttac ttcagactaa 1260
agctacccaa ggacttagca gatatggggg ttacatcagt gctggtcatt gtagcctgag 1320
tatacaatca agcttcagt tgcaaccttt ttttcttttg ccattttcta ttttagtaat 1380
ttccttgggg aactaaataa ttttgcagaa ttttctctaa ttttgtttat caggttttgc 1440
acaaagcaga gccactgtct aacacagctg ttaacgaatg ataaactgac attatactct 1500
aaaagatggt gtatttgtgc attagatttg cctgaaaaac tttatccatt tccattcttt 1560
atacaaatca catgtaatgt gtacatattt aactaaagag atttatagtc ataattattt 1620
tattgtaaa attttaacta aagtttttcc ttttctctca aactgagttc tgaattttat 1680
ttgattctga tctgaaacta ttgtctycgt aaaagttaga tctgacttca grcagaaacc 1740
aataccagct tccttttctt ttaaactttg aagagtgttg atttgttact atattactat 1800
gcaaaactgg cagttatttt tataatataa atttataatt tgatttttta ttttaaaaac 1860
tgggttaatc aagtctcggt aagtccttta aaccatttag gatttttaaa acatcaaaat 1920
ttatgattta cattcatagg aataaaataa aatatyatta gaactctggt 1970

```

```

<210> SEQ ID NO 26
<211> LENGTH: 634
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: SITE
<222> LOCATION: all Xaa positions
<223> OTHER INFORMATION: Xaa=unknown amino acid residue

```

```

<400> SEQUENCE: 26

```

```

Glu Thr Ser Lys Leu Gly Ser Ala Val Leu Ala Pro Ala Ala Gly Gly
 1             5             10            15
Thr Leu Ser Ser Glu Gly Arg Ser Ala Val Ser Gly Ile Leu Ile Ala
      20            25            30
Val Thr Ser Thr Gly Val Asp Lys Ser Leu Asn Gln Leu Leu His Gly
      35            40            45
Leu Gly Thr Ser Ser Arg Leu Ser His Phe Pro Phe Gly Lys Ser Pro
      50            55            60
Pro Arg Gly Gln Phe Val Ala Ala Ala Val Glu Ile Ala Gly Arg Ser
      65            70            75            80
Gly Leu Gln Met Gly Gln Gly Leu Trp Arg Val Val Arg Asn Gln Gln
      85            90            95
Leu Gln Gln Glu Gly Tyr Ser Glu Gln Gly Tyr Leu Thr Arg Glu Gln
      100           105           110

```

-continued

Ser	Arg	Arg	Met	Ala	Ala	Ser	Asn	Ile	Ser	Asn	Thr	Asn	His	Arg	Lys
	115						120					125			
Gln	Val	Gln	Gly	Gly	Ile	Asp	Ile	Tyr	His	Leu	Leu	Lys	Ala	Arg	Lys
	130					135					140				
Ser	Lys	Glu	Gln	Glu	Gly	Phe	Ile	Asn	Leu	Glu	Met	Leu	Pro	Pro	Glu
145					150					155					160
Leu	Ser	Phe	Thr	Ile	Leu	Ser	Tyr	Leu	Asn	Ala	Thr	Asp	Leu	Cys	Leu
				165					170					175	
Ala	Ser	Cys	Val	Trp	Gln	Asp	Leu	Ala	Asn	Asp	Glu	Leu	Leu	Trp	Gln
			180					185					190		
Gly	Leu	Cys	Lys	Ser	Thr	Trp	Gly	His	Cys	Ser	Ile	Tyr	Asn	Lys	Asn
	195						200					205			
Pro	Pro	Leu	Gly	Phe	Ser	Phe	Arg	Lys	Xaa	Tyr	Met	Gln	Leu	Asp	Glu
	210					215					220				
Gly	Ser	Leu	Thr	Phe	Asn	Ala	Asn	Pro	Asp	Glu	Gly	Val	Asn	Tyr	Phe
225					230					235					240
Met	Ser	Lys	Gly	Ile	Leu	Asp	Asp	Ser	Pro	Lys	Glu	Ile	Ala	Lys	Phe
				245					250					255	
Ile	Phe	Cys	Thr	Arg	Thr	Leu	Asn	Trp	Lys	Lys	Leu	Arg	Ile	Tyr	Leu
			260					265					270		
Asp	Glu	Arg	Arg	Asp	Val	Leu	Asp	Asp	Leu	Val	Thr	Leu	His	Asn	Phe
	275						280					285			
Arg	Asn	Gln	Phe	Leu	Pro	Asn	Ala	Leu	Arg	Glu	Phe	Phe	Arg	His	Ile
	290					295					300				
His	Ala	Pro	Glu	Glu	Arg	Gly	Glu	Tyr	Leu	Glu	Thr	Leu	Ile	Thr	Lys
305					310					315					320
Phe	Ser	His	Arg	Phe	Cys	Ala	Cys	Asn	Pro	Asp	Leu	Met	Arg	Glu	Leu
				325					330					335	
Gly	Leu	Ser	Pro	Asp	Ala	Val	Tyr	Val	Leu	Cys	Tyr	Ser	Leu	Ile	Leu
			340					345					350		
Leu	Ser	Ile	Asp	Leu	Thr	Ser	Pro	His	Val	Lys	Asn	Lys	Met	Ser	Lys
	355						360					365			
Arg	Glu	Phe	Ile	Arg	Asn	Thr	Arg	Arg	Ala	Ala	Gln	Asn	Ile	Ser	Glu
	370					375					380				
Asp	Phe	Val	Gly	His	Leu	Tyr	Asp	Asn	Ile	Tyr	Leu	Ile	Gly	His	Val
385					390					395					400
Ala	Ala	Lys	Ala	Gln	Leu	Leu	Gly	Leu	Gln	Phe	Leu	Leu	Gln	Thr	Lys
				405				410						415	
Ala	Thr	Gln	Gly	Leu	Ser	Arg	Tyr	Gly	Gly	Tyr	Ile	Ser	Ala	Gly	His
			420					425					430		
Cys	Ser	Leu	Ser	Ile	Gln	Ser	Ser	Phe	Ser	Val	Gln	Pro	Phe	Phe	Leu
		435					440					445			
Leu	Pro	Phe	Ser	Ile	Leu	Val	Ile	Ser	Leu	Gly	Asn	Ile	Ile	Leu	Gln
	450					455					460				
Asn	Phe	Ser	Phe	Cys	Leu	Ser	Arg	Phe	Ala	Gln	Ser	Arg	Ala	Thr	Val
465					470					475					480
His	Ser	Cys	Arg	Met	Ile	Asn	His	Tyr	Thr	Leu	Lys	Asp	Gly	Val	Phe
				485					490					495	
Val	His	Ile	Cys	Leu	Lys	Asn	Phe	Ile	His	Phe	His	Ser	Leu	Tyr	Lys
			500					505						510	

-continued

Tyr His Val Met Cys Thr Tyr Leu Thr Lys Glu Ile Tyr Ser His Asn
515 520 525

Tyr Phe Ile Val Lys Ile Leu Thr Lys Val Phe Pro Phe Leu Ser Asn
530 535 540

Val Leu Lys Phe Ile Phe Ser Glu Thr Ile Val Xaa Val Lys Val Arg
545 550 555 560

Ser Asp Phe Arg Gln Lys Pro Ile Pro Ala Ser Phe Ser Phe Lys Leu
565 570 575

Arg Val Leu Ile Cys Tyr Tyr Ile Thr Met Gln Asn Trp Gln Leu Phe
580 585 590

Leu Tyr Lys Phe Ile Ile Phe Phe Ile Leu Lys Thr Gly Leu Ile Lys
595 600 605

Ser Arg Val Leu Thr Ile Asp Phe Asn Ile Lys Ile Tyr Asp Leu His
610 615 620

Ser Glu Asn Lys Ile Xaa Leu Glu Leu Trp
625 630

<210> SEQ ID NO 27

<211> LENGTH: 4168

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

```

gatggcggcg gcagcagtcg acagcgcgat ggaggtggtg ccggcgctgg cggaggaggc    60
cgcgccggag gtagcggggc tcagctgcct cgtcaacctg ccgggtgagg tgctggagta    120
cctcctgtgc tgcggctcgc tgacggccgc cgacatcggc cgtgtctcca gcacctgccg    180
gcggctgcgc gagctgtgcc agagcagcgg gaaggtgtgg aaggagcagt tccgggtgag    240
gtggccttcc cttatgaaac actacagccc caccgactac gtcaattggt tggaaagata    300
taaagttcgg caaaaagctg ggtagaagc gcggaagatt gtagcctcgt tctcaaagag    360
gttcttttca gagcacgttc cttgtaattg cttcagtgac attgagaacc ttgaaggacc    420
agagattttt tttaggatg aactgggtgt taccctaaat atggaaggaa gaaaagcttt    480
gacctggaaa tactacgcaa aaaaaattct ttactacctg cggcaacaga agatcttaaa    540
taatcttaag gcctttcttc agcagccaga tgactatgag tcgtatcttg aagggtgctgt    600
atatattgac cagtactgca atcctctctc cgacatcagc ctcaaagaca tccaggccca    660
aattgacagc atcgtggagc ttgtttgcaa aacccttcgg ggcataaaca gtcgccaccc    720
cagcttggcc ttcaaggcag gtgaatcatc catgataatg gaaatagaac tccagagcca    780
ggtgctggat gccatgaact atgtccttta cgaccaactg aagttcaagg ggaatcgaat    840
ggattactat aatgccctca acttatatat gcatacaggt ttgattcgca gaacaggaat    900
cccaatcagc atgtctctgc tctatttgac aattgctcgg cagttgggag tcccactgga    960
gcctgtcaac ttcccaagtc acttcttatt aagggtgtgc caaggcgag aagggcgagc   1020
cctggacatc ttgactaca tctacataga tgcttttggg aaaggcaagc agctgacagt   1080
gaaagaatgc gagtacttga tcggccagca cgtgactgca gactgtatg ggggtgtcaa   1140
tgtcaagaag gtgttacaga gaatgggtgg aaacctgtta agcctgggga agcgggaagg   1200
catcgaccag tcataccagc tcctgagaga ctgctgggat ctctatcttg caatgtaccc   1260
ggaccagggt cagcttctcc tcctccaagc caggctttac ttccacctgg gaatctggcc   1320

```


-continued

agagaaggtg cttgacatcc tccagcacat ccaaacccta gaccgggggc agcacggggc	1380
ggtgggctac ctggtgcagc acactctaga gcacattgag cgcaaaaagg aggaggtggg	1440
cgtagaggtg aagctgcgct ccgatgagaa gcacagagat gtctgctact ccatcgggct	1500
cattatgaag cataagaggt atggctataa ctgtgtgatac tacggctggg accccacctg	1560
catgatggga cagcagtgga tccggaacat gaacgtccac agcctgccgc acggccacca	1620
ccagcctttc tataacgtgc tgggtggagga cggctcctgt cgatacgag cccaagaaaa	1680
cttggaaatat aacgtggagc ctcaagaaat ctcacaccct gacgtgggac gctattttctc	1740
agagtttact ggcaactcact acatcccaaa cgcagagctg gagatccggg atccagaaga	1800
tctggagttt gtctatgaaa cgggtgcagaa tatttacagt gcaagaaaag agaacaataga	1860
tgagtaaagt ctagagagga cattgcacct ttgctgctgc tgctatcttc caagagaacg	1920
ggactccgga agaagacgtc tccacggagc cctcgggacc tgctgcacca ggaaagccac	1980
tccaccagta gtgctggttg cctcctacta agtttaaata ccgtgtgctc tccccagct	2040
gcaaaagaaa tgtgtctctc cgcctacact agtgaattaa tctgaaaggc actgtgtcag	2100
tggcatggct tgtatgcttg tcctgtggtg acagtttgtg acattctgtc ttcagaggt	2160
ctcacagtcg acgctcctgt aatcattctt tgtattcact ccattcccct gtctgtctgc	2220
atttgtctca gaacatttcc ttggctggac agatgggggtt atgcatttgc aataatttcc	2280
ttctgatttc tctgtggaac gtgttcggtc ccgagttagg actgtgtgtc tttttaccct	2340
gaagttagtt gcatattcag aggtaaagt gtgtgctatc ttggcagcat cttagagatg	2400
gagacattaa caagctaagt gtaattagaa tcatttgaat ttattttttt ctaatatgtg	2460
aaacacagat ttcaagtgtt ttatcttttt tttttaaat taaatgggaa tataacacag	2520
ttttcccttc catattcctc tcttgagttt atgcacatct ctataaatca ttagttttct	2580
attttattac ataaaattct tttagaaaa gcaaatagtg aactttgtga atggattttt	2640
ccatactcat ctacaattcc tccattttta atgactactt ttatttttta atttaaaaa	2700
tctacttcag tatcatgagt aggtcttaca tcagtgatgg gttctttttg tagtgagaca	2760
tacaaatctg atgttaatgt ttgctcttag aagtcatact ccattggtctt caaagaccaa	2820
aaaatgaggt ttgctcttg taatcaggaa aaaaaaaaa taatgaacct taaaaaaaaa	2880
aaaaaagggt ttgaaggga aaaaagtggt ttcacacctc ttgttattcc ttagagtcac	2940
ttcaaggcct gtttgaatgt ggcagggttag aaagagagag aatgtctttc atttgaagag	3000
tggttgacct gtgtgaaagg agatgtgcgt gttggaatct gcttttccaa gccgccaggg	3060
tcctgacggc agcagacga agcctgttgt ggcgtcttct gggaaagcct gaccgtgtgt	3120
tcggacggca ctggctcctt tccgaagttc tcagtaactg agcccagagt aactgcacgc	3180
ctttgtgcag ctctggagct ccaccaactc tcggcctgcc agttctcaag cgagctaatac	3240
ttgtcattaa tcgatagaag ctaacttcg aagttaggac ctagttaact tgctctcaac	3300
atttaaaata atgcagttgc tctagtgaat ggggcgttag gggcctgtct ctgcacctgt	3360
ctgtccatct gcatgcagta ttctcaccca tgttgaatgc ctgctgcttg tttacccttt	3420
ggaaaccctg gggtagccaa ggtttggaaa gccacctgag accacttcac agcaaggga	3480
ggctttaagc agttactaga aagagatggg gatttggccc ctggctcctc cagcctgaat	3540
gagctattta atccactgtc catgttcctc atcagtcaaa tccaaagtca aaggatttga	3600

-continued

```

acctgcatct ggaaacgtaa ccactcacag cacctggccc gccaaaggttg ggaggattgt 3660
acactacttt cattaagagg ggaaagtttg ataatacgga attaattaat atgaatgaga 3720
tgcattaata agaacctgag catgctgaga gttgcaattg ttggttttct ggtttgattg 3780
atttcctttt ttcttagaca catcaaagtc aagaaagatg gttttacctt tactgaccca 3840
gctgtacata tgtatctaga ctgtttttta atgtctttct tcatgaatgc ttcattggggc 3900
tccaggaagc ctgtatcacc tgtgtaagtt ggtatttggg cactttatat ttttctaaaa 3960
acgtgttttg gatcctgtac tctaataaat cataagtttc tttttaaaaa ttttccaaaa 4020
cttttctcca ttttaaaaag ccctgttata aacgttgaac tttcacaatg ttaaaatgtt 4080
aaatatttgg atatagcaac ttcttttctc ttcaaataaa tgccaagatt tttttgtaca 4140
atgattaata aatggaactt atccagag 4168

```

```

<210> SEQ ID NO 28
<211> LENGTH: 621
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 28

```

```

Met Ala Ala Ala Val Asp Ser Ala Met Glu Val Val Pro Ala Leu
 1           5           10          15
Ala Glu Glu Ala Ala Pro Glu Val Ala Gly Leu Ser Cys Leu Val Asn
          20          25          30
Leu Pro Gly Glu Val Leu Glu Tyr Ile Leu Cys Cys Gly Ser Leu Thr
          35          40          45
Ala Ala Asp Ile Gly Arg Val Ser Ser Thr Cys Arg Arg Leu Arg Glu
          50          55          60
Leu Cys Gln Ser Ser Gly Lys Val Trp Lys Glu Gln Phe Arg Val Arg
          65          70          75          80
Trp Pro Ser Leu Met Lys His Tyr Ser Pro Thr Asp Tyr Val Asn Trp
          85          90          95
Leu Glu Glu Tyr Lys Val Arg Gln Lys Ala Gly Leu Glu Ala Arg Lys
          100         105         110
Ile Val Ala Ser Phe Ser Lys Arg Phe Phe Ser Glu His Val Pro Cys
          115         120         125
Asn Gly Phe Ser Asp Ile Glu Asn Leu Glu Gly Pro Glu Ile Phe Phe
          130         135         140
Glu Asp Glu Leu Val Cys Ile Leu Asn Met Glu Gly Arg Lys Ala Leu
          145         150         155         160
Thr Trp Lys Tyr Tyr Ala Lys Lys Ile Leu Tyr Tyr Leu Arg Gln Gln
          165         170         175
Lys Ile Leu Asn Asn Leu Lys Ala Phe Leu Gln Gln Pro Asp Asp Tyr
          180         185         190
Glu Ser Tyr Leu Glu Gly Ala Val Tyr Ile Asp Gln Tyr Cys Asn Pro
          195         200         205
Leu Ser Asp Ile Ser Leu Lys Asp Ile Gln Ala Gln Ile Asp Ser Ile
          210         215         220
Val Glu Leu Val Cys Lys Thr Leu Arg Gly Ile Asn Ser Arg His Pro
          225         230         235         240
Ser Leu Ala Phe Lys Ala Gly Glu Ser Ser Met Ile Met Glu Ile Glu
          245         250         255

```

-continued

```

Leu Gln Ser Gln Val Leu Asp Ala Met Asn Tyr Val Leu Tyr Asp Gln
      260                      265                      270

Leu Lys Phe Lys Gly Asn Arg Met Asp Tyr Tyr Asn Ala Leu Asn Leu
      275                      280                      285

Tyr Met His Gln Val Leu Ile Arg Arg Thr Gly Ile Pro Ile Ser Met
      290                      295                      300

Ser Leu Leu Tyr Leu Thr Ile Ala Arg Gln Leu Gly Val Pro Leu Glu
      305                      310                      315                      320

Pro Val Asn Phe Pro Ser His Phe Leu Leu Arg Trp Cys Gln Gly Ala
      325                      330                      335

Glu Gly Ala Thr Leu Asp Ile Phe Asp Tyr Ile Tyr Ile Asp Ala Phe
      340                      345                      350

Gly Lys Gly Lys Gln Leu Thr Val Lys Glu Cys Glu Tyr Leu Ile Gly
      355                      360                      365

Gln His Val Thr Ala Ala Leu Tyr Gly Val Val Asn Val Lys Lys Val
      370                      375                      380

Leu Gln Arg Met Val Gly Asn Leu Leu Ser Leu Gly Lys Arg Glu Gly
      385                      390                      395                      400

Ile Asp Gln Ser Tyr Gln Leu Leu Arg Asp Ser Leu Asp Leu Tyr Leu
      405                      410                      415

Ala Met Tyr Pro Asp Gln Val Gln Leu Leu Leu Leu Gln Ala Arg Leu
      420                      425                      430

Tyr Phe His Leu Gly Ile Trp Pro Glu Lys Val Leu Asp Ile Leu Gln
      435                      440                      445

His Ile Gln Thr Leu Asp Pro Gly Gln His Gly Ala Val Gly Tyr Leu
      450                      455                      460

Val Gln His Thr Leu Glu His Ile Glu Arg Lys Lys Glu Glu Val Gly
      465                      470                      475                      480

Val Glu Val Lys Leu Arg Ser Asp Glu Lys His Arg Asp Val Cys Tyr
      485                      490                      495

Ser Ile Gly Leu Ile Met Lys His Lys Arg Tyr Gly Tyr Asn Cys Val
      500                      505                      510

Ile Tyr Gly Trp Asp Pro Thr Cys Met Met Gly His Glu Trp Ile Arg
      515                      520                      525

Asn Met Asn Val His Ser Leu Pro His Gly His His Gln Pro Phe Tyr
      530                      535                      540

Asn Val Leu Val Glu Asp Gly Ser Cys Arg Tyr Ala Ala Gln Glu Asn
      545                      550                      555                      560

Leu Glu Tyr Asn Val Glu Pro Gln Glu Ile Ser His Pro Asp Val Gly
      565                      570                      575

Arg Tyr Phe Ser Glu Phe Thr Gly Thr His Tyr Ile Pro Asn Ala Glu
      580                      585                      590

Leu Glu Ile Arg Tyr Pro Glu Asp Leu Glu Phe Val Tyr Glu Thr Val
      595                      600                      605

Gln Asn Ile Tyr Ser Ala Lys Lys Glu Asn Ile Asp Glu
      610                      615                      620

```

<210> SEQ ID NO 29

<211> LENGTH: 278

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: modified_base

-continued

<222> LOCATION: all n positions
 <223> OTHER INFORMATION: n=a, c, g or t

<400> SEQUENCE: 29

```
ccgtagtact ggnntccggc gggctggtga ggaatggagc cggtagntgc ttgcggcgag    60
tcccgggntc ctccgtagac ccgcgganac cttcgtgttg agtaacctgg cggagggtggt    120
ggagcgtgtg ctcaccttcc tgcccgccaa ggcgttgctg cgggtggcct gcgtgtgccg    180
cttatggagg gagtgtgtgc gcagagtatt gcggacccat cggagcgtaa cctggatctc    240
cgcaggcctg gcggaggccg gccacctggn ggggcatt                                278
```

<210> SEQ ID NO 30
 <211> LENGTH: 91
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: SITE
 <222> LOCATION: all Xaa positions
 <223> OTHER INFORMATION: Xaa=unknown amino acid residue

<400> SEQUENCE: 30

```
Arg Ser Thr Gly Phe Arg Arg Ala Gly Glu Glu Trp Ser Arg Xaa Leu
  1             5             10            15
Ala Ala Ser Pro Gly Xaa Leu Arg Arg Pro Ala Xaa Thr Phe Val Leu
          20            25            30
Ser Asn Leu Ala Glu Val Val Glu Arg Val Leu Thr Phe Leu Pro Ala
          35            40            45
Lys Ala Leu Leu Arg Val Ala Cys Val Cys Arg Leu Trp Arg Glu Cys
          50            55            60
Val Arg Arg Val Leu Arg Thr His Arg Ser Val Thr Trp Ile Ser Ala
          65            70            75            80
Gly Leu Ala Glu Ala Gly His Leu Xaa Gly His
          85            90
```

<210> SEQ ID NO 31
 <211> LENGTH: 592
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31

```
gcggccgcgc ccggtgcagc aacagcagca gcagcccccg cagcagccgc cgccgcagcc    60
gccccagcag cagccgcccc agcagcagcc tccgccgcgc ccgcagcagc agcagcagca    120
gcagcctccg ccgccgccac cgccgctccc gccgctgcct caggagcggg acaacgtcgg    180
cgagcgggat gatgatgtgc ctgcagatat gggtgcagaa gaatcaggtc ctggtgcaca    240
aaatagtcca taccaacttc gtagaaaaac tcttttgccg aaaagaacag cgtgtccccc    300
aaagaacagt atggaggggc cctcaacttc aactacagaa aactttggtc atcgtgcaaa    360
acgtgcaaga gtgtctggaa aatcacaaga tctatcagca gcacctgctg aacagtatct    420
tcaggagaaa ctgccagatg aagtggttct aaaaatcttc tcttacttgc tggaacagga    480
tctttgtaga gcagcttggt tatgtaaacg cttcagttaa cttgctaata atcccaattt    540
gtggaaacga ttatatatgg aagtatttga atatactcgc cctatgatgc at                                592
```

<210> SEQ ID NO 32
 <211> LENGTH: 197

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 32

Arg Pro Arg Pro Val Gln Gln Gln Gln Gln Gln Pro Pro Gln Gln Pro
 1             5             10             15

Pro Pro Gln Pro Pro Gln Gln Gln Pro Pro Gln Gln Gln Pro Pro Pro
      20             25             30

Pro Pro Gln Gln Gln Gln Gln Gln Gln Pro Pro Pro Pro Pro Pro Pro
 35             40             45

Pro Pro Pro Leu Pro Gln Glu Arg Asn Asn Val Gly Glu Arg Asp Asp
 50             55             60

Asp Val Pro Ala Asp Met Val Ala Glu Glu Ser Gly Pro Gly Ala Gln
 65             70             75             80

Asn Ser Pro Tyr Gln Leu Arg Arg Lys Thr Leu Leu Pro Lys Arg Thr
      85             90             95

Ala Cys Pro Thr Lys Asn Ser Met Glu Gly Ala Ser Thr Ser Thr Thr
      100            105            110

Glu Asn Phe Gly His Arg Ala Lys Arg Ala Arg Val Ser Gly Lys Ser
      115            120            125

Gln Asp Leu Ser Ala Ala Pro Ala Glu Gln Tyr Leu Gln Glu Lys Leu
      130            135            140

Pro Asp Glu Val Val Leu Lys Ile Phe Ser Tyr Leu Leu Glu Gln Asp
      145            150            155            160

Leu Cys Arg Ala Ala Cys Val Cys Lys Arg Phe Ser Glu Leu Ala Asn
      165            170            175

Asp Pro Asn Leu Trp Lys Arg Leu Tyr Met Glu Val Phe Glu Tyr Thr
      180            185            190

Arg Pro Met Met His
      195

```

```

<210> SEQ ID NO 33
<211> LENGTH: 537
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 33

gcggccgcgg ccgggactcc gcggtgggcg agcgccctgt gaggtgacca tggaggctgg      60
tggcctcccc ttggagctgt ggcgcctgat cttagcctac ttgcaccttc ccgacctggg      120
ccgctgcagc ctggtatgca gggcctggta tgaactgata ctcagtctcg acagcaccgc      180
ctggcggcag ctgtgtcttg gttgcaccga gtgccgccat cccaattggc ccaaccagcc      240
agatgtggag cctgagctct ggagagaagc cttcaagcag cattaccttg catccaagac      300
atggaccaag aatgccttgg acttggagtc ttccatctgc ttttctctat tccgccggag      360
gagggaaacga cgtacctga gtgttgggcc aggcctgtag tttgacagcc tgggcagtgc      420
cttgcccatg gccagcctgt atgaccgaat tgtgctcttc ccagggtgtg acgaagagca      480
aggtgaaatc atcttgaagg tgcctgtgga gattgtaggg caggggaagt tgggtga      537

```

```

<210> SEQ ID NO 34
<211> LENGTH: 178
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

-continued

<400> SEQUENCE: 34

Arg Pro Arg Pro Gly Leu Arg Gly Gly Arg Ala Pro Cys Glu Val Thr
 1 5 10 15
 Met Glu Ala Gly Gly Leu Pro Leu Glu Leu Trp Arg Met Ile Leu Ala
 20 25 30
 Tyr Leu His Leu Pro Asp Leu Gly Arg Cys Ser Leu Val Cys Arg Ala
 35 40 45
 Trp Tyr Glu Leu Ile Leu Ser Leu Asp Ser Thr Arg Trp Arg Gln Leu
 50 55 60
 Cys Leu Gly Cys Thr Glu Cys Arg His Pro Asn Trp Pro Asn Gln Pro
 65 70 75 80
 Asp Val Glu Pro Glu Ser Trp Arg Glu Ala Phe Lys Gln His Tyr Leu
 85 90 95
 Ala Ser Lys Thr Trp Thr Lys Asn Ala Leu Asp Leu Glu Ser Ser Ile
 100 105 110
 Cys Phe Ser Leu Phe Arg Arg Arg Arg Glu Arg Arg Thr Leu Ser Val
 115 120 125
 Gly Pro Gly Arg Glu Phe Asp Ser Leu Gly Ser Ala Leu Ala Met Ala
 130 135 140
 Ser Leu Tyr Asp Arg Ile Val Leu Phe Pro Gly Val Tyr Glu Glu Gln
 145 150 155 160
 Gly Glu Ile Ile Leu Lys Val Pro Val Glu Ile Val Gly Gln Gly Lys
 165 170 175
 Leu Gly

<210> SEQ ID NO 35

<211> LENGTH: 751

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 35

gagaccgaga cggcgccgct gaccctagag tcgctgccc a ccgatcccct gctcctcatc 60
 ttatcctttt tggactatcg ggatctaac aactgttggt atgtcagtcg aagattaagc 120
 cagctatcaa gtcatgatcc gctgtggaga agacattgca aaaaatactg gctgatatct 180
 gaggaagaga aaacacagaa gaatcagtgt tggaaatctc tcttcataga tacttactct 240
 gatgtaggaa gatacattga ccattatgct gctattaaaa aggcctcggg aatgatctca 300
 agaaatattt ggagcccagg tgcctcggg tgggttttat ctctgaaaga ggggtgctcg 360
 agaggaagac ctcgatgctg tggaagcgca gattgggctg caagtctcct ggacgattat 420
 cgatgttcat accgaattca caatggacag aagttagttg gttcctgggg ttattgggaa 480
 gcatggcact gtctaatac tctcgttctg aagatttggt agacgtcgat acagctgccg 540
 gagattccag cagagacagg gactgaaata ctgtctccct ttaacttttg catacatact 600
 ggtttgagtc agtacatagc agtggagcgt gcagagggtt gaaacaaaaa tgaagttttc 660
 taccaatgtc agacagtaga acgtgtgttt aaatatggca ttaagatgtg ttctgatggt 720
 tgtataaatg gcatgcatta ggtattttca g 751

<210> SEQ ID NO 36

<211> LENGTH: 247

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 36

Glu Thr Glu Thr Ala Pro Leu Thr Leu Glu Ser Leu Pro Thr Asp Pro
 1 5 10 15
 Leu Leu Leu Ile Leu Ser Phe Leu Asp Tyr Arg Asp Leu Ile Asn Cys
 20 25 30
 Cys Tyr Val Ser Arg Arg Leu Ser Gln Leu Ser Ser His Asp Pro Leu
 35 40 45
 Trp Arg Arg His Cys Lys Lys Tyr Trp Leu Ile Ser Glu Glu Glu Lys
 50 55 60
 Thr Gln Lys Asn Gln Cys Trp Lys Ser Leu Phe Ile Asp Thr Tyr Ser
 65 70 75 80
 Asp Val Gly Arg Tyr Ile Asp His Tyr Ala Ala Ile Lys Lys Ala Ser
 85 90 95
 Gly Met Ile Ser Arg Asn Ile Trp Ser Pro Gly Val Leu Gly Trp Val
 100 105 110
 Leu Ser Leu Lys Glu Gly Cys Ser Arg Gly Arg Pro Arg Cys Cys Gly
 115 120 125
 Ser Ala Asp Trp Ala Ala Ser Phe Leu Asp Asp Tyr Arg Cys Ser Tyr
 130 135 140
 Arg Ile His Asn Gly Gln Lys Leu Val Gly Ser Trp Gly Tyr Trp Glu
 145 150 155 160
 Ala Trp His Cys Leu Ile Thr Ile Val Leu Lys Ile Cys Thr Ser Ile
 165 170 175
 Gln Leu Pro Glu Ile Pro Ala Glu Thr Gly Thr Glu Ile Leu Ser Pro
 180 185 190
 Phe Asn Phe Cys Ile His Thr Gly Leu Ser Gln Tyr Ile Ala Val Glu
 195 200 205
 Ala Ala Glu Gly Asn Lys Asn Glu Val Phe Tyr Gln Cys Gln Thr Val
 210 215 220
 Glu Arg Val Phe Lys Tyr Gly Ile Lys Met Cys Ser Asp Gly Cys Ile
 225 230 235 240
 Asn Gly Met His Val Phe Ser
 245

<210> SEQ ID NO 37

<211> LENGTH: 368

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: all n positions

<223> OTHER INFORMATION: n=a, c, g or t

<400> SEQUENCE: 37

ggctccgggt tccgggccgg cgggtggccg ctcacccatgc ccgnaagca ccagcatttc 60
 caggaacctg aggtcggctg ctgcgggaaa tacttcctgt ttggcttcaa cattgtcttc 120
 tgggtgctgg gagccctgtt cctggctatc ggcctctggg cctggggtga gaagggcggt 180
 ctctgaaca tctcagcgt gacagatctg ggaggccttg accccgtgtg gcttgtttgt 240
 ggtagttgga ggcgtcatgt cgggtgctgg ctttgctggg ctgcaattgg ggcctccgg 300
 gagaacacct tcctgctcaa gttttctcnc gngttcctcg gtctcatott ctctctggag 360
 ctggcaac 368

-continued

<210> SEQ ID NO 38
 <211> LENGTH: 122
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: SITE
 <222> LOCATION: all Xaa positions
 <223> OTHER INFORMATION: Xaa=unknown amino acid residue

<400> SEQUENCE: 38

```

Gly Ser Gly Phe Arg Ala Gly Gly Trp Pro Leu Thr Met Pro Gly Lys
 1             5             10             15
His Gln His Phe Gln Glu Pro Glu Val Gly Cys Cys Gly Lys Tyr Phe
      20             25             30
Leu Phe Gly Phe Asn Ile Val Phe Trp Val Leu Gly Ala Leu Phe Leu
      35             40             45
Ala Ile Gly Leu Trp Ala Trp Gly Glu Lys Gly Val Leu Ser Asn Ile
      50             55             60
Ser Ala Leu Thr Asp Leu Gly Gly Leu Asp Pro Val Trp Leu Val Cys
      65             70             75             80
Gly Ser Trp Arg Arg His Val Gly Ala Gly Leu Cys Trp Ala Ala Ile
      85             90             95
Gly Ala Leu Arg Glu Asn Thr Phe Leu Leu Lys Phe Phe Xaa Xaa Phe
      100            105            110
Leu Gly Leu Ile Phe Phe Leu Glu Leu Ala
      115            120
  
```

<210> SEQ ID NO 39
 <211> LENGTH: 774
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 39

```

gcggcgccgc ccgccgcgta cctggacgag ctgcccgagc cgctgctgct gcgcgtgctg      60
gccgcactgc cggcccgcca gctgggtgcag gcctgccgcc tgggtgtgct gcgctggaag      120
gagctggtgg acggcgcccc gctgtggctg ctcaagtgcc agcaggaggg gctgggtgcc      180
gagggcgccg tggaggagga gcgcgaccac tggcagcagt tctacttctt gagcaagcgg      240
cgccgcaacc ttctgcgtaa cccgtgtggg gaagaggact tggaaggctg gtgtgacgtg      300
gagcatgggt gggacggctg gaggggtggag gagctgcctg gagacagtgg ggtggagttc      360
accacgatg agagcgtaaa gaagtacttc gctcctctct ttgagtgggt tcgcaaagca      420
caggtcattg acctgcaggc tgagggttac tgggaggagc tgctggacac gactcagccg      480
gccatcgtgg tgaaggactg gtactcgggc cgcagcgacg ctggttgccct ctacgagctc      540
accgttaagc tactgtccga gcacgagaac gtgctggctg agttcagcag cgggcagggt      600
gcagtgcccc aagacagtga cggcgggggc tggatggaga tctccacac cttcaccgac      660
tacgggccgg gcgtccgctt cgtccgcttc gagcacgggg ggcaggggctc cgtctactgg      720
aagggtggtt tcggggcccc ggtgaccaac agcagcgtgt gggtagaacc ctga      774
  
```

<210> SEQ ID NO 40
 <211> LENGTH: 257
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 40

Ala Ala Ala Ala Ala Tyr Leu Asp Glu Leu Pro Glu Pro Leu Leu
 1 5 10 15
 Leu Arg Val Leu Ala Ala Leu Pro Ala Ala Glu Leu Val Gln Ala Cys
 20 25 30
 Arg Leu Val Cys Leu Arg Trp Lys Glu Leu Val Asp Gly Ala Pro Leu
 35 40 45
 Trp Leu Leu Lys Cys Gln Gln Glu Gly Leu Val Pro Glu Gly Gly Val
 50 55 60
 Glu Glu Glu Arg Asp His Trp Gln Gln Phe Tyr Phe Leu Ser Lys Arg
 65 70 75 80
 Arg Arg Asn Leu Leu Arg Asn Pro Cys Gly Glu Glu Asp Leu Glu Gly
 85 90 95
 Trp Cys Asp Val Glu His Gly Gly Asp Gly Trp Arg Val Glu Glu Leu
 100 105 110
 Pro Gly Asp Ser Gly Val Glu Phe Thr His Asp Glu Ser Val Lys Lys
 115 120 125
 Tyr Phe Ala Ser Ser Phe Glu Trp Cys Arg Lys Ala Gln Val Ile Asp
 130 135 140
 Leu Gln Ala Glu Gly Tyr Trp Glu Glu Leu Leu Asp Thr Thr Gln Pro
 145 150 155 160
 Ala Ile Val Val Lys Asp Trp Tyr Ser Gly Arg Ser Asp Ala Gly Cys
 165 170 175
 Leu Tyr Glu Leu Thr Val Lys Leu Leu Ser Glu His Glu Asn Val Leu
 180 185 190
 Ala Glu Phe Ser Ser Gly Gln Val Ala Val Pro Gln Asp Ser Asp Gly
 195 200 205
 Gly Gly Trp Met Glu Ile Ser His Thr Phe Thr Asp Tyr Gly Pro Gly
 210 215 220
 Val Arg Phe Val Arg Phe Glu His Gly Gly Gln Gly Ser Val Tyr Trp
 225 230 235 240
 Lys Gly Trp Phe Gly Ala Arg Val Thr Asn Ser Ser Val Trp Val Glu
 245 250 255
 Pro

<210> SEQ ID NO 41

<211> LENGTH: 957

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 41

atgggcgaga aggcggtccc ttgctaagg aggaggcggg tgaagagaag ctgcccttct 60
 tgtggtcgcg agcttggggg tgaagagaag aggggggaaag gaaatccgat ttccatccag 120
 ttgttcccc cagagctggt ggagcatatc atctcattcc tcccagtcag agaccttggt 180
 gccctcgcc agacctgcc ctacttcac gaagtgtgcg atggggaagg cgtgtggaga 240
 cgcattctgc gcagactcag tccgcgcctc caagatcagg acacgaaggg cctgtatttc 300
 caggcatttg gaggcgcgcg ccgatgtctc agcaagagcg tggccccctt gctagcccac 360
 ggctaccgcc gcttcttgcc caccaaggat cacgtcttca ttcttgacta cgtggggacc 420
 ctcttcttcc tcaaaaatgc cctggtctcc accctcggcc agatgcagtg gaagcggggc 480

-continued

```

tgtcgctatg ttgtgttggtg tcgtggagcc aaggattttg cctcggaccc aaggtgtgac 540
acagtttacc gtaaatacct ctacgtcttg gccactcggg agccgcagga agtgggtgggt 600
accaccagca gccgggcctg tgactgtgtt gaggtctatc tgcagtctag tgggcagcgg 660
gtcttcaaga tgacattcca ccaactcaatg accttcaagc agatcgtgct ggttggtcag 720
gagacccagc gggctctact gctcctcaca gaggaaggaa agatctactc ttggttagtg 780
aatgagaccc agcttgacca gccacgctcc tacacgggtc agctggccct gaggaagggtg 840
tcccactacc tgccctacct gcgcgtggcc tgcagtactt ccaaccagag cagcaccctc 900
tacgtcacag atcctattct gtgctcttgg ctacaaccac cttggcctgg tggatga 957

```

<210> SEQ ID NO 42

<211> LENGTH: 318

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 42

```

Met Gly Glu Lys Ala Val Pro Leu Leu Arg Arg Arg Val Lys Arg
 1           5           10          15
Ser Cys Pro Ser Cys Gly Ser Glu Leu Gly Val Glu Glu Lys Arg Gly
 20          25          30
Lys Gly Asn Pro Ile Ser Ile Gln Leu Phe Pro Pro Glu Leu Val Glu
 35          40          45
His Ile Ile Ser Phe Leu Pro Val Arg Asp Leu Val Ala Leu Gly Gln
 50          55          60
Thr Cys Arg Tyr Phe His Glu Val Cys Asp Gly Glu Gly Val Trp Arg
 65          70          75          80
Arg Ile Cys Arg Arg Leu Ser Pro Arg Leu Gln Asp Gln Asp Thr Lys
 85          90          95
Gly Leu Tyr Phe Gln Ala Phe Gly Gly Arg Arg Arg Cys Leu Ser Lys
100         105         110
Ser Val Ala Pro Leu Leu Ala His Gly Tyr Arg Arg Phe Leu Pro Thr
115         120         125
Lys Asp His Val Phe Ile Leu Asp Tyr Val Gly Thr Leu Phe Phe Leu
130         135         140
Lys Asn Ala Leu Val Ser Thr Leu Gly Gln Met Gln Trp Lys Arg Ala
145         150         155         160
Cys Arg Tyr Val Val Leu Cys Arg Gly Ala Lys Asp Phe Ala Ser Asp
165         170         175
Pro Arg Cys Asp Thr Val Tyr Arg Lys Tyr Leu Tyr Val Leu Ala Thr
180         185         190
Arg Glu Pro Gln Glu Val Val Gly Thr Thr Ser Ser Arg Ala Cys Asp
195         200         205
Cys Val Glu Val Tyr Leu Gln Ser Ser Gly Gln Arg Val Phe Lys Met
210         215         220
Thr Phe His His Ser Met Thr Phe Lys Gln Ile Val Leu Val Gly Gln
225         230         235         240
Glu Thr Gln Arg Ala Leu Leu Leu Thr Glu Glu Gly Lys Ile Tyr
245         250         255
Ser Leu Val Val Asn Glu Thr Gln Leu Asp Gln Pro Arg Ser Tyr Thr
260         265         270

```

-continued

Val Gln Leu Ala Leu Arg Lys Val Ser His Tyr Leu Pro His Leu Arg
275 280 285

Val Ala Cys Met Thr Ser Asn Gln Ser Ser Thr Leu Tyr Val Thr Asp
290 295 300

Pro Ile Leu Cys Ser Trp Leu Gln Pro Pro Trp Pro Gly Gly
305 310 315

<210> SEQ ID NO 43

<211> LENGTH: 1590

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 43

```

cgagggggaa gcaaggaag ggaagagga agggaaaagc gagcgagagg ggcaaggcgg      60
aagaggaagc agggcggaag ggaagcccgg gccgcagacg gcaaggagg cagcggggccg    120
ggggctgagg cgggagcgag gacacgcccc agagaggaag cagagggagg cggaagcgtg    180
gaggaagggg cgagaggcat catcaaagga gatgagggga gcgtaggggc cgggaaagag    240
gcacaaggaa gaaagtatgg gaaggaggaa tggaggggtca gggctaggcg gcgggagggc    300
gccaggccgg gaagagtaca aggacaagga ggtcaggttt gggcctacat cccggggaca    360
ggggcgggcca tggcgggcggc agccaggagg gaggaggagg aggcgggctcg ggagtcagcc  420
gcctgcccgg ctgcgggggc agcgctctgg cgctgcccgg aagtgtctgt gctgcacatg    480
tgctcctacc tcgacatgcg ggccctcggc cgctggggcc aggtgtaccg ctggctgtgg    540
cacttcacca actgcgacct gctccggcgc cagatagcct gggcctcgct caactccggc    600
ttcacgcggc tcggcaccaa cctgatgacc agtgtcccag tgaagggtgc tcagaactgg    660
atagtggggg gctgccgaga ggggattctg ctgaagtgga gatgcagtca gatgccctgg    720
atgcagctag aggatgatgc tttgtacata tcccaggcta atttcactct ggcctaccag    780
ttccgtccag atggtgccag ctgtaaccgt cagcctctgg gagtctctgc tgggcatgat    840
gaggacgttt gccactttgt gctggccacc tcgcatattg tcagtgcagg aggagatggg    900
aagattggcc ttggtaagat tcacagcacc ttcgctgcca agtactgggc tcatgaacag    960
gaggtgaact gtgtggattg caaagggggc atcatatcat ttggctccag ggacaggacg   1020
gccaaggtgt gcccttttgc ctcaggccag ctggggcagt gtttatacac catccagact   1080
gaagacaaaa tctggtctgt tgctatcagg ccattactca gctcttttgt gacagggacg   1140
gcttgttgtg ggcacttctc acccctgaaa atctgggacc tcaacagtgg gcagctgatg   1200
acacacttgg acagagactt tccccaaagg gctgggggtgc tggatgtcat atatgagtcc   1260
cctttcgcac tgctctcctg tggctatgac acctatgttc gctactggga ctgccgcacc   1320
agtgtccgga aatgtgtcat ggagtgggag gagccccaca acagcacccct gtactgcctg   1380
cagacagatg gcaaccactt gcttgccaca ggctcctcct tctatagcgt tgtacggctg   1440
tgggaccggc accaaagggc ctgccgcac accttcccgc tgacgtcgac ccgcctcggc   1500
agccctgtgt actgctgca tctcaccacc aagcatctct atgctgcgct gtcttacaac   1560
ctccacgtcc tggatattca aaaccogtga                                     1590

```

<210> SEQ ID NO 44

<211> LENGTH: 529

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 44

```

Arg Gly Gly Ser Glu Gly Arg Gly Arg Gly Arg Glu Lys Arg Ala Arg
 1           5           10           15
Gly Ala Arg Arg Lys Arg Lys Gln Gly Gly Arg Glu Ala Arg Ala Ala
 20           25           30
Asp Gly Glu Gly Gly Ser Gly Pro Gly Ala Glu Ala Gly Ala Arg Thr
 35           40           45
Arg Pro Arg Glu Glu Ala Glu Gly Gly Gly Ser Val Glu Glu Gly Ala
 50           55           60
Arg Gly Ile Ile Lys Gly Asp Glu Gly Ser Val Gly Ala Gly Lys Glu
 65           70           75           80
Ala Gln Gly Arg Lys Tyr Gly Lys Glu Glu Trp Arg Val Arg Ala Arg
 85           90           95
Arg Arg Glu Gly Ala Arg Pro Gly Arg Val Gln Gly Gln Gly Gly Gln
100          105          110
Val Trp Ala Tyr Ile Pro Gly Thr Gly Ala Ala Met Ala Ala Ala Ala
115          120          125
Arg Glu Glu Glu Glu Glu Ala Ala Arg Glu Ser Ala Ala Cys Pro Ala
130          135          140
Ala Gly Pro Ala Leu Trp Arg Leu Pro Glu Val Leu Leu Leu His Met
145          150          155          160
Cys Ser Tyr Leu Asp Met Arg Ala Leu Gly Arg Leu Ala Gln Val Tyr
165          170          175
Arg Trp Leu Trp His Phe Thr Asn Cys Asp Leu Leu Arg Arg Gln Ile
180          185          190
Ala Trp Ala Ser Leu Asn Ser Gly Phe Thr Arg Leu Gly Thr Asn Leu
195          200          205
Met Thr Ser Val Pro Val Lys Val Ser Gln Asn Trp Ile Val Gly Cys
210          215          220
Cys Arg Glu Gly Ile Leu Leu Lys Trp Arg Cys Ser Gln Met Pro Trp
225          230          235          240
Met Gln Leu Glu Asp Asp Ala Leu Tyr Ile Ser Gln Ala Asn Phe Ile
245          250          255
Leu Ala Tyr Gln Phe Arg Pro Asp Gly Ala Ser Leu Asn Arg Gln Pro
260          265          270
Leu Gly Val Ser Ala Gly His Asp Glu Asp Val Cys His Phe Val Leu
275          280          285
Ala Thr Ser His Ile Val Ser Ala Gly Gly Asp Gly Lys Ile Gly Leu
290          295          300
Gly Lys Ile His Ser Thr Phe Ala Ala Lys Tyr Trp Ala His Glu Gln
305          310          315          320
Glu Val Asn Cys Val Asp Cys Lys Gly Gly Ile Ile Ser Phe Gly Ser
325          330          335
Arg Asp Arg Thr Ala Lys Val Trp Pro Leu Ala Ser Gly Gln Leu Gly
340          345          350
Gln Cys Leu Tyr Thr Ile Gln Thr Glu Asp Gln Ile Trp Ser Val Ala
355          360          365
Ile Arg Pro Leu Leu Ser Ser Phe Val Thr Gly Thr Ala Cys Cys Gly
370          375          380
His Phe Ser Pro Leu Lys Ile Trp Asp Leu Asn Ser Gly Gln Leu Met

```

-continued

385	390	395	400
Thr His Leu Asp Arg Asp Phe Pro Pro Arg Ala Gly Val Leu Asp Val	405	410	415
Ile Tyr Glu Ser Pro Phe Ala Leu Leu Ser Cys Gly Tyr Asp Thr Tyr	420	425	430
Val Arg Tyr Trp Asp Cys Arg Thr Ser Val Arg Lys Cys Val Met Glu	435	440	445
Trp Glu Glu Pro His Asn Ser Thr Leu Tyr Cys Leu Gln Thr Asp Gly	450	455	460
Asn His Leu Leu Ala Thr Gly Ser Ser Phe Tyr Ser Val Val Arg Leu	465	470	475
Trp Asp Arg His Gln Arg Ala Cys Pro His Thr Phe Pro Leu Thr Ser	485	490	495
Thr Arg Leu Gly Ser Pro Val Tyr Cys Leu His Leu Thr Thr Lys His	500	505	510
Leu Tyr Ala Ala Leu Ser Tyr Asn Leu His Val Leu Asp Ile Gln Asn	515	520	525

Pro

<210> SEQ ID NO 45

<211> LENGTH: 1214

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 45

```

gcattgctat aattttacta tactctcatc taaatctaaa atcagtcttc aaaataaaaa    60
caaattgtcc ttgcccataa atttttttaa tcgcacaatt aattgacatt aactgccaat    120
tctttttggc taattgacta attttaactt ctgtgttgct tttccagagg catggctatt    180
gcaccttggg agaagccttt aatcggttag acttctcaag tgcaattcaa gatatccgaa    240
cgttcaatta tgtggtcaaa ctggtgcagc taattgcaa atcccagta acttcattga    300
gtggcgtggc acagaagaat tacttcaaca ttttgataa aatcgttcaa aaggttcttg    360
atgaccacca caatcctcgc ttaatcaaa atcttctgca agacctaa gc tctaccctct    420
gcattcttat tagaggagta gggaagtctg tattagtggg aaacatcaat atttgattt    480
gccgattaga aactattctc gcctggcaac aacagctaca ggatcttcag atgactaagc    540
aagtgaacaa tggcctcacc ctcaagtacc ttcctctgca catgctgaac aacatcctat    600
accggttctc agacggatgg gacatcatca ccttaggcca ggtgaccccc acgttgtata    660
tgcttagtga agacagacag ctgtggaaga agctttgtca gtaccatttt gctgaaaagc    720
agttttgtag acatttgatc ctttcagaaa aaggtcatat tgaatggaag ttgatgtact    780
ttgcacttca gaaacattac ccagcgaagg agcagtacgg agacacactg catttctgtc    840
ggcactgcag cattctcttt tggaaggact caggacaccc ctgcacggcg gccgacctg    900
acagctgctt cagcctgtg tctccgcagc acttcatcga cctcttcaag ttttaagggc    960
tgcccctgcc atccctattg gagattgtga atcctgctgt ctgtgcaggg ctcatagtga   1020
gtgttctgtg aggtgggtgg agactcctcg gaagccccctg cttccagaaa gcctgggaag   1080
aactgccctt ctgcaagggg gggactgcat ggttgcatth tcatcactga aagtcagagg   1140
ccaaggaaat catttctact tctttaaaaa ctccttctaa gcatattaaa atgtgaaatt   1200

```

-continued

ttgcgtactc tctc

1214

<210> SEQ ID NO 46

<211> LENGTH: 272

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 46

Leu Ile Leu Thr Ser Val Leu Leu Phe Gln Arg His Gly Tyr Cys Thr
 1 5 10 15

Leu Gly Glu Ala Phe Asn Arg Leu Asp Phe Ser Ser Ala Ile Gln Asp
 20 25 30

Ile Arg Thr Phe Asn Tyr Val Val Lys Leu Leu Gln Leu Ile Ala Lys
 35 40 45

Ser Gln Leu Thr Ser Leu Ser Gly Val Ala Gln Lys Asn Tyr Phe Asn
 50 55 60

Ile Leu Asp Lys Ile Val Gln Lys Val Leu Asp His His Asn Pro
 65 70 75 80

Arg Leu Ile Lys Asp Leu Leu Gln Asp Leu Ser Ser Thr Leu Cys Ile
 85 90 95

Leu Ile Arg Gly Val Gly Lys Ser Val Leu Val Gly Asn Ile Asn Ile
 100 105 110

Trp Ile Cys Arg Leu Glu Thr Ile Leu Ala Trp Gln Gln Gln Leu Gln
 115 120 125

Asp Leu Gln Met Thr Lys Gln Val Asn Asn Gly Leu Thr Leu Ser Asp
 130 135 140

Leu Pro Leu His Met Leu Asn Asn Ile Leu Tyr Arg Phe Ser Asp Gly
 145 150 155 160

Trp Asp Ile Ile Thr Leu Gly Gln Val Thr Pro Thr Leu Tyr Met Leu
 165 170 175

Ser Glu Asp Arg Gln Leu Trp Lys Lys Leu Cys Gln Tyr His Phe Ala
 180 185 190

Glu Lys Gln Phe Cys Arg His Leu Ile Leu Ser Glu Lys Gly His Ile
 195 200 205

Glu Trp Lys Leu Met Tyr Phe Ala Leu Gln Lys His Tyr Pro Ala Lys
 210 215 220

Glu Gln Tyr Gly Asp Thr Leu His Phe Cys Arg His Cys Ser Ile Leu
 225 230 235 240

Phe Trp Lys Asp Ser Gly His Pro Cys Thr Ala Ala Asp Pro Asp Ser
 245 250 255

Cys Phe Thr Pro Val Ser Pro Gln His Phe Ile Asp Leu Phe Lys Phe
 260 265 270

<210> SEQ ID NO 47

<211> LENGTH: 4059

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 47

agtacggcag tgaggggcaaa ggcagctcga gcatctcatc tgacgtgagt tcaagtacag 60

atcacacgcc cactaaagcc cagaagaatg tggctaccag cgaagactcc gacctgagca 120

tgcgcacact gagcacgccc agcccagccc tgatatgtcc accgaatctc ccaggatttc 180

agaatggaag gggctcgtcc acctcctcgt cctccatcac cggggagacg gtggccatgg 240

-continued

tgcaactcccc gcccccgacc cgcctcacac acccgctcat ccggtctgcc tccagacccc	300
agaaggagca ggccagcata gaccgggtcc cggaccactc catggtgcag atctttctct	360
tcctgcccac caaccagctg tgccgctcgc cgcgagtgtg ccgccgctgg tacaacctgg	420
cctgggaccc gcggctctgg aggactatcc gcctgacggg cgagaccatc aacgtggacc	480
gcgccctcaa ggtgtgacc cgcagactct gccaggacac ccccaactgt tgtctcatgc	540
tggaaacctg aactgtcagt ggctgcaggc ggctcacaga ccgagggtgt tacaccatcg	600
cccagtgtgt ccccgaactg aggcgactgg aagtctcagg ctgttacaat atctccaacg	660
aggcgtcttt tgatgtgtgt tccctctgcc ctaatctgga gcacctggat gtgtcaggat	720
gctccaaagt gacctgcac agcttgacct gggaggcctc cattaaactg tcacccttgc	780
atggcaaaca gatctccatc cgctacctgg acatgacgga ctgcttcgtg ctggaggacg	840
aaggcctgca caccatcgcg gcgcactgca cgcagctcac ccacctctac ctgcgccgct	900
gcgtccgcct gaccgacgaa ggcctgcgct acctggtgat ctactgcgcc tccatcaagg	960
agctgagcgt cagcgactgc cgcttcgtca gcgacttcgg cctgcgggag atcgccaagc	1020
tgaggtcccc cctgcggtac ctgagcatcg cgcactgcgg ccgggtcacc gacgtgggca	1080
tccgctacgt ggccaagtac tgcagcaagc tgcgctacct caacgcgagg ggctgcgagg	1140
gcatcacgga ccacggtgtg gagtacctcg ccaagaactg caccaaactc aaatccctgg	1200
atatcggaac atgccctttg gtatccgaca cgggcctgga gtgcctggcc ctgaactgct	1260
tcaacctcaa gcggctcagc ctcaagtcct gcgagagcat caccggccag ggcttgacga	1320
tcgtggccgc caactgcttt gacctccaga cgctgaatgt ccaggactgc gaggtctccg	1380
tgaggccctt gcgctttgtc aaacgccact gcaagcgtgt cgtcatcgag cacaccaacc	1440
cggctttctt ctgaagggac agagttcatc cggcgttgta ttcacacaaa cctgaacaaa	1500
gcaaattttt ttaaaagcag cgtatgtaag caccgacacc cactcaaac agctctttct	1560
tccgggaagg ttattaggaa tctggccttt atttttcttc atttctcatg ggcaacagag	1620
gccaaagaaa cgaagcaaga caaacagcaa acaggcattt tggtcaggtc atttgtaggc	1680
agtttctctt ctacacaaa agtgaactaa gcaggctgat cgctgttctt tgagcaaggc	1740
gcttactctc ctccgctcag gcccccaagg ccgccctttc cctcgcacac aggcccccac	1800
ccccagttc caccgcccc cccaaggcc acaccctccc tccctagagc agcagcgagg	1860
atccatcatc agaatacacag tgctctccag acctcctctc taaactgctt cattgacctt	1920
agtcactctc ttcaatccca ccccatgga cattcttctc aactcaatac catagcactt	1980
tgcataggca aaatactttt caggcctttt taaaaaatc attacagcaa acagctgggg	2040
aaggacatgc agtcctcccc cagctctgtc aatgactatg accttgggca aagcacttca	2100
ctgctctggg ctgcagcttc cagcactgaa tcagaggcca cacagcccaa agattagctt	2160
catgtccatt atagcattga gggagcagag ataccatac acagaagcac cttggcatag	2220
agcaccagc catcgacctc ttccaggaga actgattctg tggatggatg tgatttcagg	2280
agattgtgca gtgccagcat cagtgcataa agggctcctgt atgtcctttg gctgcaaatc	2340
acccacttcc ctgtgtttca gtgggagaat ttcctctccc acctcctcac atcctctttt	2400
gccaggctgg atgctgtcgt ctctgtacac aaatactttc tgcattcccc cctccacacc	2460
atcctagcga ggcaccagca cacctaatac cagcaaagcc cagatccccc catcagttgc	2520

-continued

```

ttttactcag tgttttcaaa taggagtaaa ggcccttgca atttttaatt aacaagcaag 2580
gccccaggga acacatgtcc tcaaaagttt ttctgatccc tcgccttgca cacctggcat 2640
gcatcaggga catctgtcct acagctggga gagacagatg cctcggttct ttgtcattca 2700
gattgcattt gacctcttct catctattta tttctttata catccagact tcatcacatg 2760
aagcctattg gggttaagtt tgtaagtgtt taattgtgca aattgccacc ctgtgtacct 2820
cctccatgtc tgtctgctg tttccacca aagaatgcaa agcagacttc cagggtgtta 2880
aattctgttc actcaacaat gccagatgaa tggaagaggg aacacactga gatgacttag 2940
actctgggcc accaaccaga cccttgaaa ggaatactaa aatcattaca aggtatggat 3000
tttaaatgga tgaacttca aattatctta tttggataga agtctatatt ctagcctcat 3060
ttgcatgaag tcagatagcc agaagaaatt ccattgctgg ttttcacgaa attcacttgt 3120
cttttgctaa taaacacatg gccctttccc agattattct ctagccaagc cccacctttg 3180
ttacgttgaa atccctcatt tattttcttc tcaaatgcc cattatccaa atgcagaacc 3240
tctgcatctc caagccagtt atgctgaatt tgtcaaatc agacaccctt gacaactgca 3300
ctcctactgt aggtcctgt gcatactgtc gtcttctgtg ggggatggag aggttagtgt 3360
gatgaggtgg tgtctgccc ggaggtttct ttcaaaccatc atggcctccc atccaatcaa 3420
catcatcaaa ttacatgtgt aatcaaggct ctgtgccatg ggggaaatga atcatttagc 3480
taggccagga tctagtgaat gccacagagt ttaaaacat gaaagaagtt gaaggcagca 3540
ttcctcagct ctgtgacttg tgacctatt tgaagtttca ggatttgggt gtcacaaagg 3600
attgtcccta atccttggcc ctggggtctt ccgagtgagc tggtttaata ctctgagaat 3660
gagcagggag atccagagaa tgaatccctg accgcatcac ctaaactgtc ttccaaacat 3720
gagacaaagc tgactgttca cactgattgc ccagcacata ccgtcttgcc agtttcttct 3780
ttctcccg tctcctgttc atccattctg ttctcccttg gggtggaat ctatgatgga 3840
ggttactggg gaaacagctc agcagatttt tggagaccaa accaaaggtc tcaactaggaa 3900
atttatctgt tttaaaacat tgcttcttc ctggtctgtc taaattgaat gctcattgtt 3960
tgttgttgtt gtttttaaat tctaattgtc aaatcactgc gtgctgtatg aatctagaaa 4020
gccttaattt actaccaaga aataaagcaa tatgttcgt 4059

```

<210> SEQ ID NO 48

<211> LENGTH: 483

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 48

```

Tyr Gly Ser Glu Gly Lys Gly Ser Ser Ser Ile Ser Ser Asp Val Ser
 1             5             10             15
Ser Ser Thr Asp His Thr Pro Thr Lys Ala Gln Lys Asn Val Ala Thr
 20             25             30
Ser Glu Asp Ser Asp Leu Ser Met Arg Thr Leu Ser Thr Pro Ser Pro
 35             40             45
Ala Leu Ile Cys Pro Pro Asn Leu Pro Gly Phe Gln Asn Gly Arg Gly
 50             55             60
Ser Ser Thr Ser Ser Ser Ile Thr Gly Glu Thr Val Ala Met Val
 65             70             75             80

```


His	Ser	Pro	Pro	Pro	Thr	Arg	Leu	Thr	His	Pro	Leu	Ile	Arg	Leu	Ala	
				85					90				95			
Ser	Arg	Pro	Gln	Lys	Glu	Gln	Ala	Ser	Ile	Asp	Arg	Leu	Pro	Asp	His	
				100				105					110			
Ser	Met	Val	Gln	Ile	Phe	Ser	Phe	Leu	Pro	Thr	Asn	Gln	Leu	Cys	Arg	
				115			120					125				
Cys	Ala	Arg	Val	Cys	Arg	Arg	Trp	Tyr	Asn	Leu	Ala	Trp	Asp	Pro	Arg	
				130		135					140					
Leu	Trp	Arg	Thr	Ile	Arg	Leu	Thr	Gly	Glu	Thr	Ile	Asn	Val	Asp	Arg	
145					150					155						160
Ala	Leu	Lys	Val	Leu	Thr	Arg	Arg	Leu	Cys	Gln	Asp	Thr	Pro	Asn	Val	
				165					170						175	
Cys	Leu	Met	Leu	Glu	Thr	Val	Thr	Val	Ser	Gly	Cys	Arg	Arg	Leu	Thr	
				180					185						190	
Asp	Arg	Gly	Leu	Tyr	Thr	Ile	Ala	Gln	Cys	Cys	Pro	Glu	Leu	Arg	Arg	
				195					200						205	
Leu	Glu	Val	Ser	Gly	Cys	Tyr	Asn	Ile	Ser	Asn	Glu	Ala	Val	Phe	Asp	
				210					215						220	
Val	Val	Ser	Leu	Cys	Pro	Asn	Leu	Glu	His	Leu	Asp	Val	Ser	Gly	Cys	
225					230					235						240
Ser	Lys	Val	Thr	Cys	Ile	Ser	Leu	Thr	Arg	Glu	Ala	Ser	Ile	Lys	Leu	
				245					250						255	
Ser	Pro	Leu	His	Gly	Lys	Gln	Ile	Ser	Ile	Arg	Tyr	Leu	Asp	Met	Thr	
				260					265						270	
Asp	Cys	Phe	Val	Leu	Glu	Asp	Glu	Gly	Leu	His	Thr	Ile	Ala	Ala	His	
				275					280						285	
Cys	Thr	Gln	Leu	Thr	His	Leu	Tyr	Leu	Arg	Arg	Cys	Val	Arg	Leu	Thr	
				290					295						300	
Asp	Glu	Gly	Leu	Arg	Tyr	Leu	Val	Ile	Tyr	Cys	Ala	Ser	Ile	Lys	Glu	
305					310					315						320
Leu	Ser	Val	Ser	Asp	Cys	Arg	Phe	Val	Ser	Asp	Phe	Gly	Leu	Arg	Glu	
				325					330						335	
Ile	Ala	Lys	Leu	Glu	Ser	Arg	Leu	Arg	Tyr	Leu	Ser	Ile	Ala	His	Cys	
				340					345						350	
Gly	Arg	Val	Thr	Asp	Val	Gly	Ile	Arg	Tyr	Val	Ala	Lys	Tyr	Cys	Ser	
				355					360						365	
Lys	Leu	Arg	Tyr	Leu	Asn	Ala	Arg	Gly	Cys	Glu	Gly	Ile	Thr	Asp	His	
				370					375						380	
Gly	Val	Glu	Tyr	Leu	Ala	Lys	Asn	Cys	Thr	Lys	Leu	Lys	Ser	Leu	Asp	
385					390					395						400
Ile	Gly	Lys	Cys	Pro	Leu	Val	Ser	Asp	Thr	Gly	Leu	Glu	Cys	Leu	Ala	
				405					410						415	
Leu	Asn	Cys	Phe	Asn	Leu	Lys	Arg	Leu	Ser	Leu	Lys	Ser	Cys	Glu	Ser	
				420					425						430	
Ile	Thr	Gly	Gln	Gly	Leu	Gln	Ile	Val	Ala	Ala	Asn	Cys	Phe	Asp	Leu	
				435					440						445	
Gln	Thr	Leu	Asn	Val	Gln	Asp	Cys	Glu	Val	Ser	Val	Glu	Ala	Leu	Arg	
				450					455						460	
Phe	Val	Lys	Arg	His	Cys	Lys	Arg	Cys	Val	Ile	Glu	His	Thr	Asn	Pro	
465					470					475						480
Ala	Phe	Phe														

-continued

<210> SEQ ID NO 49
<211> LENGTH: 850
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 49

```

tgcgccgcgc cccgcaccgc caccggcacc cagccccacg cccgaggaag ggccccgacgc      60
gggctgggga gaccgcattc ccttggaat cctggcgag attttcgggt tgttggtggc      120
ggcgacggc cccatgccct tcctgggcag ggctgcgcgc gtgtgccgc gctggcagga      180
ggccgcttcc caaccgcgc tctggcacac cgtgacctg tcgtccccgc tggtcggccg      240
gcctgccaa ggcgggggtca aggcggagaa gaagctcctt gcttccttg agtggcttat      300
gcccacatcg ttttcacagc tccagaggct gacctcatc cactggaagt ctacagtaca      360
ccccgtgttg aagctggtag gtgagtgtg tcctcggtc actttcctca agctctccgg      420
ctgccacggt gtgactgctg acgctctggt catgctagcc aaagcctgct gccagctcca      480
tagcctggac ctacagcact ccatggtgga gtccacagct gtggtgagct tcttgagga      540
ggcaggggtc cgaatgcgca agttgtggct gacctacagc tcccagacga cagccatcct      600
gggcgcattg ctgggcagct gctgccccca gctccaggtc ctggaggtga gcaccggcat      660
caaccgtaat agcattcccc ttcagctgcc tgtcgaggct ctgcagaaag gctgccctca      720
gtccagagtg ctgcggtgtg tgaacctgat gtggctgccc aagcctccgg gacgaggggt      780
ggctcccga ccaggcttcc ctagcctaga ggagctctgc ctggcgagct caacctgcaa      840
ctttgtgagc                                     850

```

<210> SEQ ID NO 50
<211> LENGTH: 283
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 50

```

Ala Ala Ala Pro Ala Pro Ala Pro Ala Pro Thr Pro Thr Pro Glu Glu
  1             5             10             15
Gly Pro Asp Ala Gly Trp Gly Asp Arg Ile Pro Leu Glu Ile Leu Val
             20             25             30
Gln Ile Phe Gly Leu Leu Val Ala Ala Asp Gly Pro Met Pro Phe Leu
             35             40             45
Gly Arg Ala Ala Arg Val Cys Arg Arg Trp Gln Glu Ala Ala Ser Gln
             50             55             60
Pro Ala Leu Trp His Thr Val Thr Leu Ser Ser Pro Leu Val Gly Arg
             65             70             75             80
Pro Ala Lys Gly Gly Val Lys Ala Glu Lys Lys Leu Leu Ala Ser Leu
             85             90             95
Glu Trp Leu Met Pro Asn Arg Phe Ser Gln Leu Gln Arg Leu Thr Leu
             100            105            110
Ile His Trp Lys Ser Gln Val His Pro Val Leu Lys Leu Val Gly Glu
             115            120            125
Cys Cys Pro Arg Leu Thr Phe Leu Lys Leu Ser Gly Cys His Gly Val
             130            135            140
Thr Ala Asp Ala Leu Val Met Leu Ala Lys Ala Cys Cys Gln Leu His
             145            150            155            160

```

-continued

Ser Leu Asp Leu Gln His Ser Met Val Glu Ser Thr Ala Val Val Ser
165 170 175

Phe Leu Glu Glu Ala Gly Ser Arg Met Arg Lys Leu Trp Leu Thr Tyr
180 185 190

Ser Ser Gln Thr Thr Ala Ile Leu Gly Ala Leu Leu Gly Ser Cys Cys
195 200 205

Pro Gln Leu Gln Val Leu Glu Val Ser Thr Gly Ile Asn Arg Asn Ser
210 215 220

Ile Pro Leu Gln Leu Pro Val Glu Ala Leu Gln Lys Gly Cys Pro Gln
225 230 235 240

Leu Gln Val Leu Arg Leu Leu Asn Leu Met Trp Leu Pro Lys Pro Pro
245 250 255

Gly Arg Gly Val Ala Pro Gly Pro Gly Phe Pro Ser Leu Glu Glu Leu
260 265 270

Cys Leu Ala Ser Ser Thr Cys Asn Phe Val Ser
275 280

<210> SEQ ID NO 51
<211> LENGTH: 1777
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: all n positions
<223> OTHER INFORMATION: n=a, c, g or t

<400> SEQUENCE: 51

```

acaacactgc tctcagaagg atactgcaga actccttaga ggtcttagcc tatggaatca    60
tgctgaagag cgacagaart tttttaataa ttccgtggat gaaaagtcag ataaagaagc    120
agaagtgtca gaacactcca caggtataac ccatcttcct cctgaggtaa tgctgtcaat    180
tttcagctat cttaatcctc aagagttatg tcgatgcagt caagtaagca tgaaatggtc    240
tcagctgaca aaaacgggat cgctttggaa acatctttac cctgttcatt gggccagagg    300
tgactgggtat agtgggtccc caactgaact tgatactgaa cctgatgatg aatgggtgaa    360
aaataggaaa gatgaaagtc gtgcttttca tgagtgggat gaagatgctg acattgatga    420
atctgaagag tctgcggagg aatcaattgc tatcagcatt gcacaaatgg aaaaacgttt    480
actccatggc ttaattcata acgtttctacc atatgttggt acttctgtaa aaaccttagt    540
attagcatac agctctgcag tttccagcaa aatggttagg cagattttag agctttgtcc    600
taacctggag catctggatc ttaccagac tgacatttca gattctgcat ttgacagttg    660
gtcttggtct ggttgctgcc agagtcttcg gcactctgat ctgtctggtt gtgagaaaaa    720
cacagatgtg gccctagaga agatttccag agctcttgga attctgacat ctcatcaaag    780
tggttttttg aaaacatcta caagcaaaat tacttcaact gcgtggaaaa ataaagacat    840
taccatgcag tccaccaagc agtatgcctg tttgcacgat ttaactaaca agggcattgg    900
agaagaaata gataatgaac acccctggac taagcctggt tcttctgaga atttcacttc    960
tccttatgtg tggatgtag atgctgaaga tttggctgat attgaagata ctgtggaatg   1020
gagacataga aatgttgaaa gtctttgtgt aatggaaaca gcaccaact ttagttgttc   1080
cacctctggt tgttttagta aggacattgt tggactaagg actagtgtct gttggcagca   1140
gcattgtgct tctccagcct ttgcgtattg tggtcactca ttttgttgta caggaacagc   1200

```

-continued

```

tttaagaact atgtcatcac tcccagaatc ttctgcaatg tgtagaaaag cagcaaggac 1260
tagattgcct aggggaaaag acttaattta ctttgggagt gaaaaatctg atcaagagac 1320
tggacgtgta cttctgtttc tcagtttata tggatgttat cagatcacag accatggtct 1380
cagggttttg actctgggag gagggctgcc ttatttgag caccttaac tctctggttg 1440
tcttactata actggtgcag gcctgcagga ttgggtttca gcatgtcctt ccttgaatga 1500
tgaatacttt tactactgtg acaacattaa cggctcctcat gctgataccg ccagtggatg 1560
ccagaatttg cagtgtggtt ttcgagcctg ctgccgctct ggcgaaatgac ccttgacttc 1620
tgatctttgt ctacttcatt tagctgagca ggctttcttt catgcacttt actcatagca 1680
catttcttgt gtaaccatc cctttttgag cgtgacttgt tttgggcca ttnyttacaa 1740
cttcagaaat ctttaattacc agtgrattgt aatgttg 1777

```

```

<210> SEQ ID NO 52
<211> LENGTH: 590
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: SITE
<222> LOCATION: all Xaa positions
<223> OTHER INFORMATION: Xaa=unknown amino acid residue

```

```

<400> SEQUENCE: 52

```

```

Gln His Cys Ser Gln Lys Asp Thr Ala Glu Leu Leu Arg Gly Leu Ser
  1             5             10             15
Leu Trp Asn His Ala Glu Glu Arg Gln Lys Phe Phe Lys Tyr Ser Val
          20             25             30
Asp Glu Lys Ser Asp Lys Glu Ala Glu Val Ser Glu His Ser Thr Gly
          35             40             45
Ile Thr His Leu Pro Pro Glu Val Met Leu Ser Ile Phe Ser Tyr Leu
          50             55             60
Asn Pro Gln Glu Leu Cys Arg Cys Ser Gln Val Ser Met Lys Trp Ser
          65             70             75             80
Gln Leu Thr Lys Thr Gly Ser Leu Trp Lys His Leu Tyr Pro Val His
          85             90             95
Trp Ala Arg Gly Asp Trp Tyr Ser Gly Pro Ala Thr Glu Leu Asp Thr
          100            105            110
Glu Pro Asp Asp Glu Trp Val Lys Asn Arg Lys Asp Glu Ser Arg Ala
          115            120            125
Phe His Glu Trp Asp Glu Asp Ala Asp Ile Asp Glu Ser Glu Glu Ser
          130            135            140
Ala Glu Glu Ser Ile Ala Ile Ser Ile Ala Gln Met Glu Lys Arg Leu
          145            150            155            160
Leu His Gly Leu Ile His Asn Val Leu Pro Tyr Val Gly Thr Ser Val
          165            170            175
Lys Thr Leu Val Leu Ala Tyr Ser Ser Ala Val Ser Ser Lys Met Val
          180            185            190
Arg Gln Ile Leu Glu Leu Cys Pro Asn Leu Glu His Leu Asp Leu Thr
          195            200            205
Gln Thr Asp Ile Ser Asp Ser Ala Phe Asp Ser Trp Ser Trp Leu Gly
          210            215            220
Cys Cys Gln Ser Leu Arg His Leu Asp Leu Ser Gly Cys Glu Lys Ile

```

-continued

225	230	235	240
Thr Asp Val Ala	Leu Glu Lys Ile Ser Arg	Ala Leu Gly Ile Leu Thr	
	245	250	255
Ser His Gln Ser	Gly Phe Leu Lys Thr Ser Thr Ser Lys Ile Thr Ser		
	260	265	270
Thr Ala Trp Lys	Asn Lys Asp Ile Thr Met Gln Ser Thr Lys Gln Tyr		
	275	280	285
Ala Cys Leu His	Asp Leu Thr Asn Lys Gly Ile Gly Glu Glu Ile Asp		
	290	295	300
Asn Glu His Pro	Trp Thr Lys Pro Val Ser Ser Glu Asn Phe Thr Ser		
	305	310	315
Pro Tyr Val Trp	Met Leu Asp Ala Glu Asp Leu Ala Asp Ile Glu Asp		
	325	330	335
Thr Val Glu Trp	Arg His Arg Asn Val Glu Ser Leu Cys Val Met Glu		
	340	345	350
Thr Ala Ser Asn	Phe Ser Cys Ser Thr Ser Gly Cys Phe Ser Lys Asp		
	355	360	365
Ile Val Gly Leu	Arg Thr Ser Val Cys Trp Gln Gln His Cys Ala Ser		
	370	375	380
Pro Ala Phe Ala	Tyr Cys Gly His Ser Phe Cys Cys Thr Gly Thr Ala		
	385	390	395
Leu Arg Thr Met	Ser Ser Leu Pro Glu Ser Ser Ala Met Cys Arg Lys		
	405	410	415
Ala Ala Arg Thr	Arg Leu Pro Arg Gly Lys Asp Leu Ile Tyr Phe Gly		
	420	425	430
Ser Glu Lys Ser	Asp Gln Glu Thr Gly Arg Val Leu Leu Phe Leu Ser		
	435	440	445
Leu Ser Gly Cys	Tyr Gln Ile Thr Asp His Gly Leu Arg Val Leu Thr		
	450	455	460
Leu Gly Gly Gly	Leu Pro Tyr Leu Glu His Leu Asn Leu Ser Gly Cys		
	465	470	475
Leu Thr Ile Thr	Gly Ala Gly Leu Gln Asp Leu Val Ser Ala Cys Pro		
	485	490	495
Ser Leu Asn Asp	Glu Tyr Phe Tyr Tyr Cys Asp Asn Ile Asn Gly Pro		
	500	505	510
His Ala Asp Thr	Ala Ser Gly Cys Gln Asn Leu Gln Cys Gly Phe Arg		
	515	520	525
Ala Cys Cys Arg	Ser Gly Glu Pro Leu Thr Ser Asp Leu Cys Leu Leu		
	530	535	540
His Leu Ala Glu	Gln Ala Phe Phe His Ala Leu Tyr Ser His Ile Ser		
	545	550	555
Cys Val Asn His	Pro Phe Leu Ser Val Thr Cys Phe Gly Pro Ile Xaa		
	565	570	575
Tyr Asn Phe Arg	Asn Leu Asn Tyr Gln Xaa Ile Val Met Leu		
	580	585	590

<210> SEQ ID NO 53
 <211> LENGTH: 1681
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: all n positions

-continued

<223> OTHER INFORMATION: n=a, c, g or t

<400> SEQUENCE: 53

```

ttttactgta cacagttgat gtattttgat gctgggcctg tctgggtctgt cttgaggatt    60
attaaccttt agaggatatca gagaagcaaa tgggtactgg tgaggctgct cattagggaa    120
gagggcaaaa ggagcactag ctaggtcaga gccatgtttc aggtcacaaat gtgatgtcag    180
atgttgctta taaatccttt cttgtcttcg ccattcttaa atcttgatag gtgcctgttg    240
ggaaactgta aatgcctttc ccaatggaga atcaacagat tgggtgatgg tggagtcggt    300
caggaagact caggtcttct agaggaaagg atgcctcatc accccttngg cccaggcagc    360
tgctgtcaga gaatgacaca gcacctgcac agtcgctgtc cacttcctgc cactgctgtc    420
ggtgggggtga cgggagcaaa gtaggcgtgg actttgacat gagggagctg agcccgcatc    480
cgcttgatgc ctgcacgggt aacctgctgg cagtcgtaca gctcgaggcg ctccaggcct    540
cggcagttct ctaggtgttc cagggccaca tcagtgatga ggaggcagtt gtccaactcc    600
agtacccgca gcctctcatg gccacaggta ctgttgctca ggtgcaggat cccatcatct    660
gkgatgagtt cacagtggga caggctcagg gcttgcatgt taggacagtg aatggagagc    720
tggatgagtg tgctgtcgtt tatcaggatg cawtcttcaa gatccatctt ctccaattcg    780
tggaattcc gagctaaaag tgtaaacctt gcgtcagtca aatgggagca tcgggcagcc    840
tccaaaattt gcagtcgcgg acagttcaaa cccagggtcg taagagaggc atctgtgagg    900
ttgtgcaac ccgaaaggca gagagcctgt agccggtgac agcccctgca tatctgcacc    960
acaccttcat ccgtgatacg tgagcaggac tgcaagttga ggctcacaag ctcatggcag   1020
taattctgaa tgtgtttcag agcttcatct tctaactgtg tgcagccctc caggagcagg   1080
gttttcaggc ctgcacaacc tcgcaccagt gcctcgatgc catccttcgt gatctgatca   1140
caccaagaga ggttcaggta ctccaggttt cggcagccct cactgatccc ctccaaggag   1200
ctgtttgtaa tagacacaca ggaggtcaga wccagatggt tcagcttgga acagaatctg   1260
ctaaggctat aacacgtgct gtcagtgatt tttgtgcac cattgagggt caaatgttca   1320
atgtttcggc agttctgtgc aaaggtcttc aaggaggaat ccccaacacc aatgcagcct   1380
cgcaagctga gtttcctcag gaatccaacg catcgcttcg agatattttc caccactcga   1440
ccctctacat ctatttgaaa gttaaaaaga tctattcttt gccagttgct tccatccagg   1500
gctaagatgt tccaagcctt ggaaatctgt gcacatcggc acaaagttac tatatccaag   1560
aaggaaaata ttcttaacag aagttctttg ggtaactttt tgtaataaag gccttcatca   1620
ttgtttgaga aaaccatggc cgaagagcgg cgagcgagcc cacagcccga agtcacacgg   1680
c                                                                    1681

```

<210> SEQ ID NO 54

<211> LENGTH: 437

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: SITE

<222> LOCATION: all Xaa positions

<223> OTHER INFORMATION: Xaa=unknown amino acid residue

<400> SEQUENCE: 54

```

Arg Val Thr Ser Gly Cys Gly Leu Ala Arg Gly Ser Ser Ala Met Val
  1             5             10             15

```

-continued

Phe	Ser	Asn	Asn	Asp	Glu	Gly	Leu	Ile	Asn	Lys	Lys	Leu	Pro	Lys	Glu
			20					25					30		
Leu	Leu	Leu	Arg	Ile	Phe	Ser	Phe	Leu	Asp	Ile	Val	Thr	Leu	Cys	Arg
		35					40					45			
Cys	Ala	Gln	Ile	Ser	Lys	Ala	Trp	Asn	Ile	Leu	Ala	Leu	Asp	Gly	Ser
	50					55					60				
Asn	Trp	Gln	Arg	Ile	Asp	Leu	Phe	Asn	Phe	Gln	Ile	Asp	Val	Glu	Gly
	65				70					75					80
Arg	Val	Val	Glu	Asn	Ile	Ser	Lys	Arg	Cys	Val	Gly	Phe	Leu	Arg	Lys
				85					90					95	
Leu	Ser	Leu	Arg	Gly	Cys	Ile	Gly	Val	Gly	Asp	Ser	Ser	Leu	Lys	Thr
		100						105					110		
Phe	Ala	Gln	Asn	Cys	Arg	Asn	Ile	Glu	His	Leu	Asn	Leu	Asn	Gly	Cys
		115					120					125			
Thr	Lys	Ile	Thr	Asp	Ser	Thr	Cys	Tyr	Ser	Leu	Ser	Arg	Phe	Cys	Ser
	130					135						140			
Lys	Leu	Lys	His	Leu	Xaa	Leu	Thr	Ser	Cys	Val	Ser	Ile	Thr	Asn	Ser
	145				150					155					160
Ser	Leu	Lys	Gly	Ile	Ser	Glu	Gly	Cys	Arg	Asn	Leu	Glu	Tyr	Leu	Asn
			165					170						175	
Leu	Ser	Trp	Cys	Asp	Gln	Ile	Thr	Lys	Asp	Gly	Ile	Glu	Ala	Leu	Val
		180						185					190		
Arg	Gly	Cys	Arg	Gly	Leu	Lys	Ala	Leu	Leu	Leu	Arg	Gly	Cys	Thr	Gln
		195				200						205			
Leu	Glu	Asp	Glu	Ala	Leu	Lys	His	Ile	Gln	Asn	Tyr	Cys	His	Glu	Leu
	210					215					220				
Val	Ser	Leu	Asn	Leu	Gln	Ser	Cys	Ser	Arg	Ile	Thr	Asp	Glu	Gly	Val
	225				230					235					240
Val	Gln	Ile	Cys	Arg	Gly	Cys	His	Arg	Leu	Gln	Ala	Leu	Cys	Leu	Ser
			245						250					255	
Gly	Cys	Ser	Asn	Leu	Thr	Asp	Ala	Ser	Leu	Thr	Ala	Leu	Gly	Leu	Asn
			260					265					270		
Cys	Pro	Arg	Leu	Gln	Ile	Leu	Glu	Ala	Ala	Arg	Cys	Ser	His	Leu	Thr
		275				280						285			
Asp	Ala	Gly	Phe	Thr	Leu	Leu	Ala	Arg	Asn	Cys	His	Glu	Leu	Glu	Lys
	290					295					300				
Met	Asp	Leu	Glu	Xaa	Cys	Ile	Leu	Ile	Thr	Asp	Ser	Thr	Leu	Ile	Gln
	305				310					315					320
Leu	Ser	Ile	His	Cys	Pro	Lys	Leu	Gln	Ala	Leu	Ser	Leu	Ser	His	Cys
			325						330					335	
Glu	Leu	Ile	Xaa	Asp	Asp	Gly	Ile	Leu	His	Leu	Ser	Asn	Ser	Thr	Cys
		340						345					350		
Gly	His	Glu	Arg	Leu	Arg	Val	Leu	Glu	Leu	Asp	Asn	Cys	Leu	Leu	Ile
		355				360						365			
Thr	Asp	Val	Ala	Leu	Xaa	His	Leu	Glu	Asn	Cys	Arg	Gly	Leu	Glu	Arg
	370					375					380				
Leu	Glu	Leu	Tyr	Asp	Cys	Gln	Gln	Val	Thr	Arg	Ala	Gly	Ile	Lys	Arg
	385				390					395					400
Met	Arg	Ala	Gln	Leu	Pro	His	Val	Lys	Val	His	Ala	Tyr	Phe	Ala	Pro
			405						410					415	

-continued

Val Thr Pro Pro Thr Ala Val Ala Gly Ser Gly Gln Arg Leu Cys Arg
 420 425 430
Cys Cys Val Ile Leu
 435

<210> SEQ ID NO 55

<211> LENGTH: 1866

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 55

```

atgtcaccgg tctttcccat gttaacagtt ctgaccatgt tttattatat atgccttcgg      60
cgccgagcca ggacagctac aagaggagaa atgatgaaca cccatagagc tatagaatca    120
aacagccaga cttccctctt caatgcagag gtagtccagt atgccaaaga agtagtggt      180
ttcagttccc attatggaag tgagaatagt atgtcctata ctatgtggaa tttggctggt    240
gtaccaaagt tattcccaag ttctggtgac tttactcaga cagctgtggt tcgaacttat    300
gggacatggt gggatcagtg tcctagtgtc tccttgccat tcaagaggac gccacctaatt   360
tttcagagcc aggactatgt ggaacttact tttgaacaac aggtgtatcc tacagctgta    420
catgttctag aaacctatca tcccggagca gtcattagaa ttctcgcttg ttctgcaaatt   480
ccttattccc caaatccacc agctgaagta agatgggaga ttctttggtc agagagacct    540
acgaaggtga atgcttccca agctcgccag tttaaacctt gtattaagca gataaatttc    600
ccccaaaatc ttatacgact ggaagtaaat agttctcttc tggaatatta cactgaatta    660
gatgcagttg tgctacatgg tgtgaaggac aagccagtgc tttctctcaa gacttcactt    720
attgacatga atgatataga agatgatgcc tatgcagaaa aggatggttg tggaatggac    780
agtcttaaca aaaagtttag cagtgtgtgc ctcggggaag ggccaaataa tgggtatttt    840
gataaactac cttatgagct tattcagctg attctgaatc atcttacct accagacctg    900
tgtagattag cacagacttg caaactactg agccagcatt gctgtgatcc tctgcaatac    960
atccacctca atctgcaacc atactgggca aaactagatg acacttctct ggaatttcta   1020
cagtctcgct gcaacttgtt ccagtggcct aatttatctt ggactggcaa tagaggcttc   1080
atctctgttg caggatttag caggtttctg aaggtttgtg gatccgaatt agtacgcctt   1140
gaattgtctt gcagccactt tcttaatgaa acttgcttag aagttatttc tgagatgtgt   1200
ccaaatctac aggccttaaa tctctcctcc tgtgataagc taccacctca agctttcaac   1260
cacattgcca agttatgcag ccttaaacga cttgttctct atcgaacaaa agtagagcaa   1320
acagcactgc tcagcatttt gaacttctgt tcagagcttc agcacctcag tttaggcagt   1380
tgtgtcatga ttgaagacta tgatgtgata gctagcatga taggagccaa gtgtaaaaaa   1440
ctccggaccc tggatctgtg gagatgtaag aatattactg agaatggaat agcagaactg   1500
gttcttgggt gtccactact ggaggagctt gaccttggct ggtgcccaac tctgcagagc   1560
agcaccgggt gcttcaccag actggcacac cagctcccaa acttgcaaaa actctttctt   1620
acagctaata gatctgtgtg tgacacagac attgatgaat tggcatgtaa ttgtaccagg   1680
ttacagcagc tggacatatt aggaacaaga atggtaagtc cggcatcctt aagaaaactc   1740
ctggaatctt gtaaagatct ttctttactt gatgtgtcct tctgttcgca gattgataac   1800
agagctgtgc tagaactgaa tgcaagcttt ccaaaagtgt tcataaaaaa gagctttact   1860

```


-continued

cagtga

1866

<210> SEQ ID NO 56

<211> LENGTH: 621

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 56

```

Met Ser Pro Val Phe Pro Met Leu Thr Val Leu Thr Met Phe Tyr Tyr
 1           5           10           15

Ile Cys Leu Arg Arg Ala Arg Thr Ala Thr Arg Gly Glu Met Met
 20           25           30

Asn Thr His Arg Ala Ile Glu Ser Asn Ser Gln Thr Ser Pro Leu Asn
 35           40           45

Ala Glu Val Val Gln Tyr Ala Lys Glu Val Val Asp Phe Ser Ser His
 50           55           60

Tyr Gly Ser Glu Asn Ser Met Ser Tyr Thr Met Trp Asn Leu Ala Gly
 65           70           75           80

Val Pro Asn Val Phe Pro Ser Ser Gly Asp Phe Thr Gln Thr Ala Val
 85           90           95

Phe Arg Thr Tyr Gly Thr Trp Trp Asp Gln Cys Pro Ser Ala Ser Leu
100          105          110

Pro Phe Lys Arg Thr Pro Pro Asn Phe Gln Ser Gln Asp Tyr Val Glu
115          120          125

Leu Thr Phe Glu Gln Gln Val Tyr Pro Thr Ala Val His Val Leu Glu
130          135          140

Thr Tyr His Pro Gly Ala Val Ile Arg Ile Leu Ala Cys Ser Ala Asn
145          150          155          160

Pro Tyr Ser Pro Asn Pro Pro Ala Glu Val Arg Trp Glu Ile Leu Trp
165          170          175

Ser Glu Arg Pro Thr Lys Val Asn Ala Ser Gln Ala Arg Gln Phe Lys
180          185          190

Pro Cys Ile Lys Gln Ile Asn Phe Pro Thr Asn Leu Ile Arg Leu Glu
195          200          205

Val Asn Ser Ser Leu Leu Glu Tyr Tyr Thr Glu Leu Asp Ala Val Val
210          215          220

Leu His Gly Val Lys Asp Lys Pro Val Leu Ser Leu Lys Thr Ser Leu
225          230          235          240

Ile Asp Met Asn Asp Ile Glu Asp Asp Ala Tyr Ala Glu Lys Asp Gly
245          250          255

Cys Gly Met Asp Ser Leu Asn Lys Lys Phe Ser Ser Ala Val Leu Gly
260          265          270

Glu Gly Pro Asn Asn Gly Tyr Phe Asp Lys Leu Pro Tyr Glu Leu Ile
275          280          285

Gln Leu Ile Leu Asn His Leu Thr Leu Pro Asp Leu Cys Arg Leu Ala
290          295          300

Gln Thr Cys Lys Leu Leu Ser Gln His Cys Cys Asp Pro Leu Gln Tyr
305          310          315          320

Ile His Leu Asn Leu Gln Pro Tyr Trp Ala Lys Leu Asp Asp Thr Ser
325          330          335

Leu Glu Phe Leu Gln Ser Arg Cys Thr Leu Val Gln Trp Leu Asn Leu
340          345          350

```

-continued

Ser Trp Thr Gly Asn Arg Gly Phe Ile Ser Val Ala Gly Phe Ser Arg
 355 360 365
 Phe Leu Lys Val Cys Gly Ser Glu Leu Val Arg Leu Glu Leu Ser Cys
 370 375 380
 Ser His Phe Leu Asn Glu Thr Cys Leu Glu Val Ile Ser Glu Met Cys
 385 390 395 400
 Pro Asn Leu Gln Ala Leu Asn Leu Ser Ser Cys Asp Lys Leu Pro Pro
 405 410 415
 Gln Ala Phe Asn His Ile Ala Lys Leu Cys Ser Leu Lys Arg Leu Val
 420 425 430
 Leu Tyr Arg Thr Lys Val Glu Gln Thr Ala Leu Leu Ser Ile Leu Asn
 435 440 445
 Phe Cys Ser Glu Leu Gln His Leu Ser Leu Gly Ser Cys Val Met Ile
 450 455 460
 Glu Asp Tyr Asp Val Ile Ala Ser Met Ile Gly Ala Lys Cys Lys Lys
 465 470 475 480
 Leu Arg Thr Leu Asp Leu Trp Arg Cys Lys Asn Ile Thr Glu Asn Gly
 485 490 495
 Ile Ala Glu Leu Ala Ser Gly Cys Pro Leu Leu Glu Glu Leu Asp Leu
 500 505 510
 Gly Trp Cys Pro Thr Leu Gln Ser Ser Thr Gly Cys Phe Thr Arg Leu
 515 520 525
 Ala His Gln Leu Pro Asn Leu Gln Lys Leu Phe Leu Thr Ala Asn Arg
 530 535 540
 Ser Val Cys Asp Thr Asp Ile Asp Glu Leu Ala Cys Asn Cys Thr Arg
 545 550 555 560
 Leu Gln Gln Leu Asp Ile Leu Gly Thr Arg Met Val Ser Pro Ala Ser
 565 570 575
 Leu Arg Lys Leu Leu Glu Ser Cys Lys Asp Leu Ser Leu Leu Asp Val
 580 585 590
 Ser Phe Cys Ser Gln Ile Asp Asn Arg Ala Val Leu Glu Leu Asn Ala
 595 600 605
 Ser Phe Pro Lys Val Phe Ile Lys Lys Ser Phe Thr Gln
 610 615 620

<210> SEQ ID NO 57

<211> LENGTH: 984

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 57

```

atgcaacttg tacctgatat agagttcaag attacttata cccggtctcc agatggtgat      60
ggcgttgga acagctacat tgaagataat gatgatgaca gcaaaatggc agatctcttg      120
tcctacttcc agcagcaact cacatttcag gagtctgtgc ttaaactgtg tcagcctgag      180
cttgagagca gtcagattca catatcagtg ctgccaatgg aggtcctgat gtacatcttc      240
cgatgggtgg tgtctagtga cttggacctc agatcattgg agcagttgtc gctggtgtgc      300
agaggattct acatctgtgc cagagacctt gaaatatggc gtctggcctg cttgaaagtt      360
tggggcagaa gctgtattaa acttggtccg tacacgtcct ggagagagat gtttttagaa      420
cggcctcgtg ttcggtttga tggcgtgtat atcagtaaaa ccacatatat tcgtcaaggg      480
gaacagtctc ttgatggttt ctatagagcc tggcaccaag tggaatatta caggtacata      540

```

-continued

```

agattctttc ctgatggcca tgtgatgatg ttgacaaccc ctgaagagcc tcagtccatt    600
gttccacggt taagaactag gaataccagg actgatgcaa ttctactggg tcactatcgc    660
ttgtcacaag acacagacaa tcagacacaa gtatttgctg taataactaa gaaaaaagaa    720
gaaaaaacac ttgactataa atacagatat ttctgctcgtg tccctgtaca agaagcagat    780
cagagttttc atgtggggct acagctatgt tccagtggtc accagagggt caacaaactc    840
atctggatac atcattcttg tcacattact tacaatcaa ctggtgagac tgcagtcagt    900
gcttttgaga ttgacaagat gtacaccccc ttgttcttcg ccagagtaag gagctacaca    960
gctttctcag aaaggcctct gtag                                           984

```

<210> SEQ ID NO 58

<211> LENGTH: 327

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 58

```

Met Gln Leu Val Pro Asp Ile Glu Phe Lys Ile Thr Tyr Thr Arg Ser
 1             5             10             15
Pro Asp Gly Asp Gly Val Gly Asn Ser Tyr Ile Glu Asp Asn Asp Asp
      20             25             30
Asp Ser Lys Met Ala Asp Leu Leu Ser Tyr Phe Gln Gln Gln Leu Thr
      35             40             45
Phe Gln Glu Ser Val Leu Lys Leu Cys Gln Pro Glu Leu Glu Ser Ser
      50             55             60
Gln Ile His Ile Ser Val Leu Pro Met Glu Val Leu Met Tyr Ile Phe
      65             70             75             80
Arg Trp Val Val Ser Ser Asp Leu Asp Leu Arg Ser Leu Glu Gln Leu
      85             90             95
Ser Leu Val Cys Arg Gly Phe Tyr Ile Cys Ala Arg Asp Pro Glu Ile
      100            105            110
Trp Arg Leu Ala Cys Leu Lys Val Trp Gly Arg Ser Cys Ile Lys Leu
      115            120            125
Val Pro Tyr Thr Ser Trp Arg Glu Met Phe Leu Glu Arg Pro Arg Val
      130            135            140
Arg Phe Asp Gly Val Tyr Ile Ser Lys Thr Thr Tyr Ile Arg Gln Gly
      145            150            155            160
Glu Gln Ser Leu Asp Gly Phe Tyr Arg Ala Trp His Gln Val Glu Tyr
      165            170            175
Tyr Arg Tyr Ile Arg Phe Phe Pro Asp Gly His Val Met Met Leu Thr
      180            185            190
Thr Pro Glu Glu Pro Gln Ser Ile Val Pro Arg Leu Arg Thr Arg Asn
      195            200            205
Thr Arg Thr Asp Ala Ile Leu Leu Gly His Tyr Arg Leu Ser Gln Asp
      210            215            220
Thr Asp Asn Gln Thr Lys Val Phe Ala Val Ile Thr Lys Lys Lys Glu
      225            230            235            240
Glu Lys Pro Leu Asp Tyr Lys Tyr Arg Tyr Phe Arg Arg Val Pro Val
      245            250            255
Gln Glu Ala Asp Gln Ser Phe His Val Gly Leu Gln Leu Cys Ser Ser
      260            265            270

```

-continued

Gly His Gln Arg Phe Asn Lys Leu Ile Trp Ile His His Ser Cys His
 275 280 285

Ile Thr Tyr Lys Ser Thr Gly Glu Thr Ala Val Ser Ala Phe Glu Ile
 290 295 300

Asp Lys Met Tyr Thr Pro Leu Phe Phe Ala Arg Val Arg Ser Tyr Thr
 305 310 315 320

Ala Phe Ser Glu Arg Pro Leu
 325

<210> SEQ ID NO 59
 <211> LENGTH: 765
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: all n positions
 <223> OTHER INFORMATION: n=a, c, g or t

<400> SEQUENCE: 59

```
gcagccctgg atcctgactt agagaatgat gatttctttg tcagaaagac tggggctttc    60
catgcaaadc catatgttct ccgagctttt gaagacttta gaaagttctc tgagcaagat    120
gattctgtag agcgagatat aattttacag tgtagagaag gtgaacttgt acttccggat    180
ttggaaaaag atgatatgat tgttcgccga atcccagcac agaagaaaga agtgccgctg    240
tctggggccc cagatagata ccacccagtc ccttttcccg aaccctggac tcttcctcca    300
gaaattcaag caaaatttct ctgtgtactt gaaaggacat gcccatccaa agaaaaaagt    360
aatagctgta gaatattagt tccttcatat cggcagaaga aagatgacat gctgacacgt    420
aagattcagt cctggaaact gggaactacc gtgcctccca tcagtttcac ncctggcccc    480
tgcaagtgagg ctgacttgaa gagatgggag gccatccggg aggccagcag actcaggcac    540
aagaaaaaggc tgatggtgga gagactcttt caaaagattt atggtgagaa tgggagtaag    600
tccatgagtg atgtcagcgc agaagatggt caaaacttgc gtcagctgcg ttacgaggag    660
atgcagaaaa taaaatcaca attaaaagaa caagatcaga aatggcagga tgaccttgca    720
aaatggaag atcgtcgaaa aagttacact tcagatctgc agaag                    765
```

<210> SEQ ID NO 60
 <211> LENGTH: 255
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 60

Ala Ala Leu Asp Pro Asp Leu Glu Asn Asp Asp Phe Phe Val Arg Lys
 1 5 10 15

Thr Gly Ala Phe His Ala Asn Pro Tyr Val Leu Arg Ala Phe Glu Asp
 20 25 30

Phe Arg Lys Phe Ser Glu Gln Asp Asp Ser Val Glu Arg Asp Ile Ile
 35 40 45

Leu Gln Cys Arg Glu Gly Glu Leu Val Leu Pro Asp Leu Glu Lys Asp
 50 55 60

Asp Met Ile Val Arg Arg Ile Pro Ala Gln Lys Lys Glu Val Pro Leu
 65 70 75 80

Ser Gly Ala Pro Asp Arg Tyr His Pro Val Pro Phe Pro Glu Pro Trp
 85 90 95

-continued

Thr Leu Pro Pro Glu Ile Gln Ala Lys Phe Leu Cys Val Leu Glu Arg
 100 105 110
 Thr Cys Pro Ser Lys Glu Lys Ser Asn Ser Cys Arg Ile Leu Val Pro
 115 120 125
 Ser Tyr Arg Gln Lys Lys Asp Asp Met Leu Thr Arg Lys Ile Gln Ser
 130 135 140
 Trp Lys Leu Gly Thr Thr Val Pro Pro Ile Ser Phe Thr Pro Gly Pro
 145 150 155 160
 Cys Ser Glu Ala Asp Leu Lys Arg Trp Glu Ala Ile Arg Glu Ala Ser
 165 170 175
 Arg Leu Arg His Lys Lys Arg Leu Met Val Glu Arg Leu Phe Gln Lys
 180 185 190
 Ile Tyr Gly Glu Asn Gly Ser Lys Ser Met Ser Asp Val Ser Ala Glu
 195 200 205
 Asp Val Gln Asn Leu Arg Gln Leu Arg Tyr Glu Glu Met Gln Lys Ile
 210 215 220
 Lys Ser Gln Leu Lys Glu Gln Asp Gln Lys Trp Gln Asp Asp Leu Ala
 225 230 235 240
 Lys Trp Lys Asp Arg Arg Lys Ser Tyr Thr Ser Asp Leu Gln Lys
 245 250 255

<210> SEQ ID NO 61
 <211> LENGTH: 36
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 61

Leu Pro Pro Glu Leu Ser Phe Thr Ile Leu Ser Tyr Leu Asn Ala Thr
 1 5 10 15
 Asp Leu Cys Leu Ala Ser Cys Val Trp Gln Asp Leu Ala Asn Asp Glu
 20 25 30
 Leu Leu Trp Gln
 35

<210> SEQ ID NO 62
 <211> LENGTH: 42
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 62

Leu Pro Gly Glu Val Leu Glu Tyr Ile Leu Cys Cys Gly Ser Leu Thr
 1 5 10 15
 Ala Ala Asp Ile Gly Arg Val Ser Ser Thr Cys Arg Arg Leu Arg Glu
 20 25 30
 Leu Cys Gln Ser Ser Gly Lys Val Trp Lys
 35 40

<210> SEQ ID NO 63
 <211> LENGTH: 44
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 63

Leu Ala Glu Val Val Glu Arg Val Leu Thr Phe Leu Pro Ala Lys Ala
 1 5 10 15
 Leu Leu Arg Val Ala Cys Val Cys Arg Leu Trp Arg Glu Cys Val Arg

-continued

	20		25		30
Arg	Val	Leu	Arg	Thr	His
	35			40	
Arg	Ser	Val	Thr	Trp	Ile

<210> SEQ ID NO 64
 <211> LENGTH: 39
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 64

Leu	Pro	Asp	Glu	Val	Val	Leu	Lys	Ile	Phe	Ser	Tyr	Leu	Leu	Glu	Gln
1				5				10					15		
Asp	Leu	Cys	Arg	Ala	Ala	Cys	Val	Cys	Lys	Arg	Phe	Ser	Glu	Leu	Ala
	20					25							30		

Asn	Asp	Pro	Asn	Leu	Trp	Lys
	35					

<210> SEQ ID NO 65
 <211> LENGTH: 41
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 65

Leu	Pro	Leu	Glu	Leu	Trp	Arg	Met	Ile	Leu	Ala	Tyr	Leu	His	Leu	Pro
1				5				10					15		
Asp	Leu	Gly	Arg	Cys	Ser	Leu	Val	Cys	Arg	Ala	Trp	Tyr	Glu	Leu	Ile
	20					25							30		

Leu	Ser	Leu	Asp	Ser	Thr	Arg	Trp	Arg
	35					40		

<210> SEQ ID NO 66
 <211> LENGTH: 39
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 66

Leu	Pro	Thr	Asp	Pro	Leu	Leu	Leu	Ile	Leu	Ser	Phe	Leu	Asp	Tyr	Arg
1				5				10					15		
Asp	Leu	Ile	Asn	Cys	Cys	Tyr	Val	Ser	Arg	Arg	Leu	Ser	Gln	Leu	Ser
	20					25							30		

Ser	His	Asp	Pro	Leu	Trp	Arg
	35					

<210> SEQ ID NO 67
 <211> LENGTH: 40
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 67

Leu	Pro	Glu	Pro	Leu	Leu	Leu	Arg	Val	Leu	Ala	Ala	Leu	Pro	Ala	Ala
1				5				10					15		
Glu	Leu	Val	Gln	Ala	Cys	Arg	Leu	Val	Cys	Leu	Arg	Trp	Lys	Glu	Leu
	20					25							30		

Val	Asp	Gly	Ala	Pro	Leu	Trp	Leu
	35					40	

<210> SEQ ID NO 68
 <211> LENGTH: 40

-continued

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 68
Leu Phe Pro Pro Glu Leu Val Glu His Ile Ile Ser Phe Leu Pro Val
1 5 10 15
Arg Asp Leu Val Ala Leu Gly Gln Thr Cys Arg Tyr Phe His Glu Val
20 25 30
Cys Asp Gly Glu Gly Val Trp Arg
35 40

<210> SEQ ID NO 69
<211> LENGTH: 44
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 69
Leu Pro Glu Val Leu Leu His Met Cys Ser Tyr Leu Asp Met Arg
1 5 10 15
Ala Leu Gly Arg Leu Ala Gln Val Tyr Arg Trp Leu Trp His Phe Thr
20 25 30
Asn Cys Asp Leu Leu Arg Arg Gln Ile Ala Trp Ala
35 40

<210> SEQ ID NO 70
<211> LENGTH: 40
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 70
Leu Pro Leu His Met Leu Asn Asn Ile Leu Tyr Arg Phe Ser Asp Gly
1 5 10 15
Trp Asp Ile Ile Thr Leu Gly Gln Val Thr Pro Thr Leu Tyr Met Leu
20 25 30
Ser Glu Asp Arg Gln Leu Trp Lys
35 40

<210> SEQ ID NO 71
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 71
Leu Pro Asp His Ser Met Val Gln Ile Phe Ser Phe Leu Pro Thr Asn
1 5 10 15
Gln Leu Cys Arg Cys Ala Arg Val Cys Arg Arg Trp Tyr Asn Leu Ala
20 25 30
Trp Asp Pro Arg Leu Trp Arg
35

<210> SEQ ID NO 72
<211> LENGTH: 44
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 72
Ile Pro Leu Glu Ile Leu Val Gln Ile Phe Gly Leu Leu Val Ala Ala
1 5 10 15

-continued

Asp Gly Pro Met Pro Phe Leu Gly Arg Ala Ala Arg Val Cys Arg Arg
20 25 30

Trp Gln Glu Ala Ala Ser Gln Pro Ala Leu Trp His
35 40

<210> SEQ ID NO 73
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 73

Leu Pro Pro Glu Val Met Leu Ser Ile Phe Ser Tyr Leu Asn Pro Gln
1 5 10 15

Glu Leu Cys Arg Cys Ser Gln Val Ser Met Lys Trp Ser Gln Leu Thr
20 25 30

Lys Thr Gly Ser Leu Trp Lys
35

<210> SEQ ID NO 74
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 74

Leu Pro Lys Glu Leu Leu Arg Ile Phe Ser Phe Leu Asp Ile Val
1 5 10 15

Thr Leu Cys Arg Cys Ala Gln Ile Ser Lys Ala Trp Asn Ile Leu Ala
20 25 30

Leu Asp Gly Ser Asn Trp Gln
35

<210> SEQ ID NO 75
<211> LENGTH: 48
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 75

Leu Pro Tyr Glu Leu Ile Gln Leu Ile Leu Asn His Leu Thr Leu Pro
1 5 10 15

Asp Leu Cys Arg Leu Ala Gln Thr Cys Lys Leu Leu Ser Gln His Cys
20 25 30

Cys Asp Pro Leu Gln Tyr Ile His Leu Asn Leu Gln Pro Tyr Trp Ala
35 40 45

<210> SEQ ID NO 76
<211> LENGTH: 44
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 76

Leu Pro Met Glu Val Leu Met Tyr Ile Phe Arg Trp Val Val Ser Ser
1 5 10 15

Asp Leu Asp Leu Arg Ser Leu Glu Gln Leu Ser Leu Val Cys Arg Gly
20 25 30

Phe Tyr Ile Cys Ala Arg Asp Pro Glu Ile Trp Arg
35 40

-continued

<210> SEQ ID NO 77
<211> LENGTH: 49
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 77

Leu Pro Pro Glu Ile Gln Ala Lys Phe Leu Cys Val Leu Glu Arg Thr
1 5 10 15
Cys Pro Ser Lys Glu Lys Ser Asn Ser Cys Arg Ile Leu Val Pro Ser
20 25 30
Tyr Arg Gln Lys Lys Asp Asp Met Leu Thr Arg Lys Ile Gln Ser Trp
35 40 45

Lys

<210> SEQ ID NO 78
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 78

Leu Pro His His Val Val Leu Gln Ile Phe Gln Tyr Leu Pro Leu Leu
1 5 10 15
Asp Arg Ala Cys Ala Ser Ser Val Cys Arg Arg Trp Asn Glu Val Phe
20 25 30
His Ile Ser Asp Leu Trp Arg
35

<210> SEQ ID NO 79
<211> LENGTH: 43
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 79

Leu Trp Ala Trp Gly Glu Lys Gly Val Leu Ser Asn Ile Ser Ala Leu
1 5 10 15
Thr Asp Leu Gly Gly Leu Asp Pro Val Trp Leu Val Cys Gly Ser Trp
20 25 30
Arg Arg His Val Gly Ala Gly Leu Cys Trp Ala
35 40

<210> SEQ ID NO 80
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide

<400> SEQUENCE: 80

agtagtaaca aaggtaaa acagttgact gtatcgctcga ggatgccttc aattaagtt 59

<210> SEQ ID NO 81
<211> LENGTH: 58
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide

<400> SEQUENCE: 81

-continued

gcgggttactt acttagagct cgacgtctta ctacttagc tcacttctct tcacacca 58

<210> SEQ ID NO 82
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 82

Cys Asp Gly Glu Lys Asp Thr Tyr Ser Tyr Leu Ala
1 5 10

<210> SEQ ID NO 83
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 83

Cys Glu Ser Ser Phe Ser Leu Asn Met Asn Phe Ser Ser Lys Arg Thr
1 5 10 15

Lys Phe Lys Ile Thr Thr Ser Met Gln
20 25

<210> SEQ ID NO 84
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 84

Cys Glu Glu Ala Gln Val Arg Lys Glu Asn Gln Trp
1 5 10

<210> SEQ ID NO 85
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: Phosphorylation
<222> LOCATION: 8
<223> OTHER INFORMATION: Phosothreonine

<400> SEQUENCE: 85

Asn Ala Gly Ser Val Glu Gln Thr Pro Lys Lys Pro Gly Leu Arg Arg
1 5 10 15

Arg Gln Thr

<210> SEQ ID NO 86
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide

<400> SEQUENCE: 86

cctgggggat gttctca

17

<210> SEQ ID NO 87
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
Oligonucleotide

-continued

```

<400> SEQUENCE: 87
ggcttccggg cathtag                17

<210> SEQ ID NO 88
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
      Oligonucleotide

<400> SEQUENCE: 88
catctggcac gattcca                17

<210> SEQ ID NO 89
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
      Oligonucleotide

<400> SEQUENCE: 89
ccgctcatcg tatgaca                17

<210> SEQ ID NO 90
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: Phosphorylation
<222> LOCATION: 8
<223> OTHER INFORMATION: Phosotyrosine

<400> SEQUENCE: 90
Ala Glu Ile Gly Val Gly Ala Tyr Gly Thr Val Tyr Lys Ala Arg Asp
  1           5           10          15
Pro His Ser

```

1-49. (canceled)

50. An isolated nucleic acid molecule comprising a nucleotide sequence which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO: 10.

51. An isolated nucleic acid molecule which encodes an F-box polypeptide, or a fragment thereof, said nucleic acid molecule comprising a nucleotide sequence of SEQ ID NO: 9.

52. An isolated nucleic acid sequence derived from a mammalian genome that:

- a) hybridizes under highly stringent conditions to the nucleotide sequence of SEQ ID NO: 9; and
- b) encodes a gene product which contains an F-box motif and binds to Skp 1.

53. A nucleotide vector containing the nucleotide sequence of claim 50, 51 or **52**.

54. An expression vector containing the nucleotide sequence of claim 50, 51 or **52**, in operative association with a nucleotide regulatory sequence that controls expression of the nucleotide sequence in a host cell.

55. A genetically engineered host cell that contains the nucleotide sequence of claim 50, 51 or **52**, in operative association with a nucleotide regulatory sequence that controls expression of the nucleotide sequence in the host cell.

56. A transgenic animal having cells which harbor a transgene comprising the nucleic acid of claim 50, 51 or **52**.

57. An animal inactivated in the locus comprising the nucleotide sequence of claim 50, 51 or **52**.

58. An isolated F-box polypeptide having the amino acid sequence of SEQ ID NO: 10.

59. An antibody that immunospecifically binds the polypeptide of claim 58.

60. A method of diagnosing proliferative and differentiative related disorders comprising measuring FBP5 gene expression in a patient sample.

61. A method for screening compounds useful for the treatment of proliferative and differentiative disorders comprising contacting a compound with a cell expressing an F-box polypeptide having the amino acid sequence of SEQ ID NO: 10, or a fragment thereof, and its substrate, and detecting a change in the F-box polypeptide activity.

62. The method of claim 61 wherein the change in the F-box polypeptide activity is detected by detecting a change in the interaction of the F-box polypeptide with one or more polypeptides.

63. The method of claim 62 in which one of the one or more polypeptides is the substrate of the F-box polypeptide.

64. The method of claim 62 in which at least one of the one or more polypeptides is a component of the ubiquitin pathway.

65. The method of claim 62 in which one of the one or more polypeptides is Skp1.

66. The method of claim 61 wherein the change in the F-box polypeptide activity is detected by detecting a change in the ubiquitination or degradation of the substrate.

67. A method for screening compounds useful for the treatment of proliferative and differentiative disorders comprising contacting a compound with a cell or a cell extract expressing Fbp5 and one or more Fbp5 substrates, and detecting a change in the activity of Fbp5.

68. The method of claim 67 wherein the change in the activity of Fbp5 is detected by detecting a change in the

interaction of Fbp5 with one of the one or more Fbp5 substrates.

69. The method of claim 67 wherein the change in the activity of Fbp5 is detected by detecting a change in the ubiquitination or degradation of one of the one or more Fbp5 substrates.

70. A method for treating a proliferative or differentiative disorder in a mammal comprising administering to the mammal a compound to the mammal that modulates the synthesis, expression or activity of the FBP5 gene or gene product so that symptoms of the disorder are ameliorated.

71. The method of claim 70 in which the disorder is breast cancer.

72. The method of claim 70 in which the disorder is ovarian cancer.

73. The method of claim 70 in which the disorder is prostate cancer.

74. The method of claim 70 in which the disorder is small cell lung carcinoma.

* * * * *