



AFRICAN REGIONAL INDUSTRIAL PROPERTY
ORGANISATION (ARIPO)

886

(11) (A)

<p>(21) Application Number: AP/P/98/01195</p> <p>(22) Filing Date: 19960718</p> <p>(24) Date of Grant & 20001103 (45) Publication</p>	<p>(73) Applicant(s): RHONE-POULENC AGROCHIMIE 14-20 rue Pierre Baizet F-69009 Lyon France</p>
<p>(30) Priority Data</p> <p>(33) Country: FR</p> <p>(31) Number: 95/08.979</p> <p>(32) Date: 19950719</p>	<p>(72) Inventors: MICHEL LEBRUN 224, Rue De Saint-Cyr F-69009 Lyon France</p> <p>(See Overleaf)</p>
<p>(84) Designated States:</p> <p>SD KE</p>	<p>(74) Representative GALLOWAY & COMPANY P O BOX WGT 28 WESTGATE HARARE</p>

- (51) **International Patent Classification (Int.Cl.7):** C12N 15/54; A01H 1/06
- (54) **Title:** Mutated 5-Enol Pyruvylshikimate-3-Phosphate Synthase, Gene Coding For Said Protein And Transformed Plants Containing Said Gene
- (57) **Abstract:**

A mutated glyphosate resistance gene of 5-enol pyruvylshikimate-3-phosphate synthase (EPSPS) including at least one substitution of threonine 102 by isoleucine, and useful for producing glyphosate-resistant transformed plants, is disclosed.

Inventors Continued

2. ALAIN SAILLAND
38, rue Ernest-Fabreque
F69009 Lyon
FRANCE

3. GEORGES FREYSSINET
29, rue de Nervieux
F-69540 Saint-cyr-au-Mont-D'or
France

4. ERIC DEGRYSE
4 rue des Allsiers
F-67100
Strasbourg
France

Mutated 5-enolpyruvylshikimate-3-phosphate synthase,
gene coding for this protein and transformed plants
containing this gene

The present invention relates to a new
5 5-enolpyruvylshikimate-3-phosphate synthase (or EPSPS)
which displays increased tolerance with respect to
herbicides which are competitive inhibitors with
respect to phosphoenolpyruvate (PEP) of EPSPS
activity. This more tolerant EPSP synthase possesses at
10 least one "threonine by isoleucine" substitution. The
invention also relates to a gene coding for such a
protein, to plant cells transformed by chimeric gene
constructions containing this gene, to the plants
regenerated from these cells and also to the plants
15 originating from crossing using these transformed
plants.

Glyphosate, sulfosate or fosametine are
broad-spectrum systemic herbicides of the
phosphonomethylglycine family. They act essentially as
20 competitive inhibitors of 5-enolpyruvylshikimate-3-
phosphate synthase (EC 2.5.1.19) or EPSPS with respect
to the PEP (phosphoenolpyruvate). After their
application to the plant, they are translocated in the
plant where they accumulate in the rapidly growing
25 parts, in particular the cauline and root apices,
causing damage to the point of destruction of sensitive
plants.

AP/P/98/01195

Plastid EPSPS, the main target of these products, is an enzyme of the pathway of biosynthesis of aromatic amino acids, which is encoded by one or more nuclear genes and synthesized in the form of a cytoplasmic precursor, then imported into the plastids where it accumulates in its mature form.

The tolerance of plants to glyphosate and to products of the family is obtained by stable introduction into their genome of an EPSPS gene, of plant or bacterial origin, which is mutated or otherwise in respect of the characteristics of inhibition by glyphosate of the product of this gene. In view of the mode of action of glyphosate and the degree of tolerance to glyphosate of the product of the genes which are used, it is advantageous to be able to express the product of the translation of this gene so as enable it to be accumulated in substantial amounts in the plastids.

It is known, for example from US Patent 4,535,060, to confer on a plant a tolerance to a herbicide of the above type, especially N-phosphonomethylglycine or glyphosate, by introducing into the genome of plants a gene coding for an EPSPS carrying at least one mutation that makes this enzyme more resistant to its competitive inhibitor (glyphosate) after localization of the enzyme in the plastid compartment. These techniques, however, need to be improved in order to obtain greater reliability in the use of these plants under agricultural conditions.

In the present description, "plant" is understood to mean any differentiated multicellular organism capable of photosynthesis, and "plant cell" is understood to mean any cell originating from a plant and capable of constituting undifferentiated tissues such as calluses or differentiated tissues such as embryos or plant parts or seeds.

The subject of the present invention is the production of transformed plants having increased tolerance to herbicides having EPSP synthase as their target, e.g. of the phosphonomethylglycine family, e.g. glyphosate or a glyphosate precursor, by regeneration of cells transformed by means of new chimeric genes containing a gene for tolerance to these herbicides.

The invention accordingly provides a DNA sequence coding for a mutated 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), characterized in that the EPSPS is from maize and comprises a first mutation consisting in the threonine 102 → isoleucine substitution and a second mutation consisting in a substitution of proline 106 by serine; the invention also provides a DNA sequence coding for a mutated 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), characterised in that it comprises the coding region of the DNA sequence represented in SEQ ID No. 2, comprising a first mutation consisting in the threonine 102 isoleucine substitution and a second mutation consisting in a substitution of proline 106 by serine; and a DNA sequence coding for a mutated 5-enolpyruvylshikimate -3-phosphate synthase (EPSPS), characterised in that it comprises the coding sequence of SEQ ID No. 4.

According to a further feature of the invention there is provided a 5-enolpyruvylshikimate -3-phosphate synthase (EPSPS), characterized in that the EPSPS is from maize and comprises a first mutation consisting in the substitution of threonine 102 by isoleucine and a second mutation consisting in a substitution of proline 106 by serine.

The invention also provides a chimeric gene comprising a coding sequence as well as regulatory elements at position 5' and 3' which are heterologous and capable of functioning in plants, characterized in that it comprises as coding sequence at least one sequence according to the invention.

AP 0 0 0 8 8 6

3a

The chimeric gene of the invention for conferring on plants increased tolerance with respect to a herbicide having EPSPS as its target, generally comprises, in the direction of transcription: a promoter region, optionally a transit peptide region, a sequence of a gene coding for a glyphosate tolerance enzyme and an
5 untranslated polyadenylation signal region at the 3' end, characterized in that the glyphosate tolerance gene contains, relative to the gene from which it is derived, a "threonine 102 by isoleucine" substitution in the "aroA" (EPSPS) region and, in the same region, a "proline 106 by serine" substitution. These substitutions can be introduced or be present in an EPSPS sequence of any

origin, in particular of plant, bacterial, algal or fungal origin.

The transit peptides which can be used in the transit peptide region can be, known per se, of plant origin, for example originating from maize, sunflower, pea, tobacco or the like. The first and the second transit peptide can be identical, similar or different. They can, in addition, each comprise one or more transit peptide units according to European Patent Application EP 0 508 909. It is the role of this characteristic region to permit the release of a mature and native protein, and especially the above mutated EPSPS, with maximum efficacy in the plasmid compartment.

The promoter region of the chimeric gene according to the invention may comprise a plant virus promoter; the region may be advantageously composed of at least one gene promoter or promoter fragment which is expressed naturally in plants (e.g. tubulin, generally α -tubulin, introns actin, histone).

The untranslated transcription termination signal region at the 3' end of the chimeric gene may be of any origin, for example of bacterial origin, such as that of the nopaline synthase gene, or of plant origin, such as that of the *Arabidopsis thaliana* histone H4A748 gene according to the European Patent Application (European Application 633 317).

The chimeric gene according to the invention can comprise, in addition to the essential portions

above, at least one untranslated intermediate (linker) region, which can be located between the different transcribed regions described above. This intermediate region can be of any origin, for example of bacterial, viral or plant origin.

5 The method of the invention for the production of plants with improved tolerance to a herbicide having EPSP synthase activity as its target, characterised in that plant cells or protoplasts are transformed with a gene according to the invention and in that the transformed cells are subjected to a regeneration. The plants obtained may be used to prepare hybrid lines.

10 Isolation of a cDNA coding for a maize EPSPS:

The different steps which led to the obtaining of maize EPSPS cDNA, which served as substrate for the introduction of the two mutations, are described below. All the operations described below are given by way of example, and correspond to a choice made from among the different methods available for arriving
15 at the same result. This choice has no effect on the quality of the result, and consequently any suitable method may be used by a person skilled in the art to arrive at the same result. Most of the methods of engineering of DNA fragments are described in "Current Protocols in Molecular Biology" Volumes 1 and 2, Ausubel
F.M. et al., published by Greene Publishing Associates and Wiley-Interscience
20 (1989) (hereinafter, references to protocols described in this work will be designated "ref. CPMB"). The operations relating to DNA which were performed according to the protocols described in this work are especially the following: ligation of DNA fragments, treatment with Klenow DNA polymerase and T4 DNA polymerase, preparation of plasmid and of bacteriophage λ DNA, either as a minipreparation or as
25 a maxipreparation, and DNA and

minipreparation or as a maxipreparation, and DNA and RNA analyses according to the Southern and Northern techniques, respectively. Other methods described in this work were followed, and only significant
5 modifications or additions to these protocols have been described below.

Example 1:

1. Obtaining of an *Arabidopsis thaliana* EPSPS fragment

10 a) Two 20-mer oligonucleotides of respective sequences:

5'-GCTCTGCTCATGTCTGCTCC-3'

5'-GCCCGCCCTTGACAAAGAAA-3'

were synthesized from the sequence of an *Arabidopsis thaliana* EPSPS gene (Klee H.J. et al. (1987) Mol. Gen. Genet., 210, 437-442). These two oligonucleotides are
15 at positions 1523 to 1543 and 1737 to 1717, respectively, of the published sequence, and in opposite orientations.

20 b) *Arabidopsis thaliana* (var. *columbia*) total DNA was obtained from Clontech (catalogue reference: 6970-1).

c) 50 nanograms (ng) of DNA are mixed with 300 ng of each of the oligonucleotides and subjected to
25 35 amplification cycles with a Perkin-Elmer 9600 apparatus, under the conditions of standard medium for amplification which are recommended by the supplier.

The resulting 204-bp fragment constitutes the *Arabidopsis thaliana* EPSPS fragment.

2. Construction of a library of a cDNA from a BMS maize cell line

5 a) 5 g of filtered cells are ground in liquid nitrogen, and the total nucleic acids are extracted according to the method described by Shure et al. with the following modifications:

- 10 - the pH of the lysis buffer is adjusted to PH 9.0;
- after precipitation with isopropanol, the pellet is taken up in water and, after dissolution, adjusted to 2.5 M LiCl. After incubation for 12 h at °C, 15 the pellet from centrifugation for 15 min at 30,000 g at 4°C is resolubilized. The LiCl precipitation step is then repeated. The resolubilized pellet constitutes the RNA fraction of the 20 total nucleic acids.

b) The poly(A)⁺ RNA fraction of the RNA fraction is obtained by chromatography on an oligo(dT)-cellulose column as described in "Current Protocols in Molecular Biology".

25 c) Synthesis of double-stranded cDNA having a synthetic EcoRI end: this is carried out according to the protocol of the supplier of the different reagents needed for this synthesis penny form of a kit: the

"copy kit" from the company In Vitrogen.

Two single-stranded and partially complementary oligonucleotides of respective sequences:

5'-AATTCCTGGG-3'

5
5'-CCCGGG-3' (the latter being phosphorylated)

are ligated with the blunt-ended double-stranded cDNAs.

This ligation of the adaptors results in the creation of SmaI sites attached to the double-stranded cDNAs and EcoRI sites in cohesive form at each end of the double-stranded cDNAs.

d) Creation of the library:

The cDNAs possessing the artificial cohesive EcoRI sites at their ends are ligated with bacteriophage λ gt10 cDNA which has been cut with EcoRI and dephosphorylated according to the protocol of the supplier New England Biolabs.

An aliquot of the ligation reaction was encapsidated in vitro with encapsidation extracts, namely Gigapack Gold, according to the supplier's instructions; this library was titrated using the bacterium *E. coli* C600hfl. The library thereby obtained is amplified and stored according to the instructions of the same supplier, and constitutes the BMS maize cell suspension cDNA library.

3. Screening of the BMS maize cell suspension cDNA library with the *Arabidopsis thaliana* EPSP probe

The protocol followed is that of "Current

Protocols in Molecular Biology" Volumes 1 and 2,
Ausubel F.M. et al., published by Greene Publishing
Associates and Wiley-Interscience (1989) (CPMB).
Briefly, approximately 10^6 recombinant phages are
5 plated out on LB dishes at an average density of 100
phages/cm². The lytic plaques are replicated in
duplicate on Amersham Hybond N membranes.

h) The DNA was fixed to the filters by 1600kJ
UV treatment (Stratagene Stratalinker). The filters
10 were prehybridized in 6×SSC/0.1%SDS/0.25 skimmed milk
for 2 h at 65°C. The *Arabidopsis thaliana* EPSPS probe
was labelled with [³²P]dCTP by random priming according
to the supplier's instructions (Pharmacia Ready to Go
kit). The specific activity obtained is of the order of
15 10⁸ cpm per µg of fragment. After denaturation for 5
min at 100°C, the probe is added to the
prehybridization medium and hybridization is continued
for 14 hours at 55°C. The filters are fluorographed for
48 h at -80°C with Kodak XAR5 film and Amersham
20 Hyperscreen RPN enhancing screens. Alignment of the
positive spots on the filter with the dishes from which
they originate enables zones corresponding to the
phages displaying a positive hybridization response
with the *Arabidopsis thaliana* EPSPS probe to be picked
25 out from the dish. This step of plating out, transfer,
hybridization and recovery is repeated until all the
spots in the dish of the successively purified phages
prove 100% positive in hybridization. A plaque of

AP/98/01195

independent phage lysis is then picked out in diluent λ medium (Tris-Cl pH 7.5; 10mM MgSO₄; 0.1M NaCl; 0.1% gelatin); these phages in solution constitute the EPSP-positive clones of the BMS maize cell suspension.

5 4. Preparation and analysis of the DNA of the EPSP clones of the BMS maize cell suspension

Approximately 5×10^8 phages are added to 20 ml of C600hfl bacteria at an OD_{600nm} value of 2/ml and incubated for 15 minutes at 37°C. This suspension is then diluted in 200 ml of bacterial growth medium in a 1-1 Erlenmeyer and stirred in a rotary stirrer at 250 rpm. Lysis is noted when the medium clarifies, corresponding to the lysis of the turbid bacteria, and takes place after approximately 4 h of stirring. This supernatant is then treated as described in "Current Protocols in Molecular Biology". The DNA obtained corresponds to the EPSP clones of the BMS maize cell suspension.

One to two μ g of this DNA are cut with EcoRI and separated on 0.8% LGTA/TBE agarose gel (ref. CPMB). A final verification consists in checking that the purified DNA does indeed display a hybridization signal with the *Arabidopsis thaliana* EPSPS probe. After electrophoresis, the DNA fragments are transferred onto Amersham Hybond N membranes according to the protocol of Southern described in "Current Protocols in Molecular Biology". The filter is hybridized with the *Arabidopsis thaliana* EPSPS probe according to the

conditions described in section 3 above. The clone displaying a hybridization signal with the *Arabidopsis thaliana* EPSPS probe and containing the longest EcoRI fragment has a size estimated on gel as approximately
5 1.7 kbp.

5. Obtaining of the clone pRPA-ML-711

Ten μg of the phage clone containing the 1.7-kbp insert are digested with EcoRI and separated on 0.8% LGTA/TBE agarose gel (ref. CPMB). The gel fragment
10 containing the 1.7-kbp insert is excised from the gel by BET staining, and the fragment is treated with β -agarse according to the protocol of the supplier, New England Biolabs. The purified DNA of the 1.7-kbp
15 fragment is ligated at 12°C for 14 h with the DNA of plasmid pUC 19 (New England Biolabs) cut with EcoRI according to the ligation protocol described in "Current Protocols in Molecular Biology". Two μl of the above ligation mixture are used for the transformation
20 of an aliquot of electrocompetent *E. coli* DH10B; transformation is accomplished by electroporation using the following conditions: the mixture of competent bacteria and and ligation medium is introduced into an electroporation cell of thickness 0.2 cm (Biorad) previously cooled to 0°C. The physical conditions of
25 the electroporation using an electroporator made by Biorad are 2500 volts, 25 μF and 200 Ω . Under these conditions, the mean discharge time of the condenser is of the order of 4.2 milliseconds. The bacteria are then.

taken up in 1 ml of SOC medium (ref. CPMB) and stirred for 1 hour at 200 rpm on a rotary stirrer in 15-ml Corning tubes. After plating out on LB/agar medium supplemented with 100 µg/ml of carbenicillin, minipreparations of the bacterial clones which have grown after one night at 37°C is produced according to the protocol described in "Current Protocols in Molecular Biology". After digestion of the DNA with EcoRI and separation by electrophoresis on 0.8% LGTA/TBE agarose gel (ref. CPMB), the clones possessing a 1.7-kbp insert are retained. A final verification consists in checking that the purified DNA does indeed display a hybridization signal with the *Arabidopsis thaliana* EPSPS probe. After electrophoresis, the DNA fragments are transferred onto Amersham Hybond N membranes according to the protocol of Southern described in "Current Protocols in Molecular Biology". The filter is hybridized with the *Arabidopsis thaliana* EPSPS probe according to the conditions described in section 3 above. The plasmid clone possessing a 1.7-kbp insert and hybridizing with the *Arabidopsis thaliana* EPSPS probe was prepared on a larger scale, and the DNA resulting from the lysis of the bacteria was purified on a CsCl gradient as described in "Current Protocols in Molecular Biology". The purified DNA was partially sequenced with a Pharmacia kit according to the supplier's instructions and using as primers the M13 direct and reverse universal primers ordered from the

AP/P/98/01195

same supplier. The partial sequence produced covers approximately 0.5 kbp. The derived amino acid sequence in the region of the mature protein (approximately 50 amino acid residues) displays 100% identity with the
5 corresponding amino sequence of mature maize EPSPS described in American Patent USP 4,971,908). This clone, corresponding to a 1.7-kbp EcoRI fragment of the EPSP DNA of the BMS maize cell suspension, was designated pRPA-ML-711. The complete sequence of this
10 clone was determined on both strands using the protocol of the Pharmacia kit and synthesizing complementary oligonucleotides and those of the opposite orientation every 250 bp approximately. The complete sequence obtained of this 1713-bp clone is presented in SEQ ID
15 No. 1.

6. Obtaining of the clone pRPA-ML-715

Analysis of the sequence of the clone pRPA-ML-711, and especially comparison of the derived amino acid sequence with that of maize, shows a sequence
20 extension of 92 bp upstream of the GCG codon coding for the NH₂-terminal alanine of the mature portion of maize EPSPS (American Patent USP 4,971,908). Similarly, an extension of 288 bp downstream of the AAT codon coding for the COOH-terminal asparagine of the mature portion
25 of maize EPSPS (American Patent USP 4,971,908) is observed. These two portions could correspond, in the case of the NH₂-terminal extension to a portion of the sequence of a transit peptide for plastid localization,

and, in the case of the COOH-terminal extension, to the untranslated 3' region of the cDNA.

In order to obtain a cDNA coding for the mature portion of the maize EPSPS cDNA, as described in USP 4,971,908, the following operations were carried out:

a) Removal of the untranslated 3' region: construction of pRPA-ML-712:

The clone pRPA-ML-711 was cut with the restriction enzyme AseI, and the ends resulting from this cleavage were rendered blunt by treatment with the Klenow fragment of DNA polymerase I according to the protocol described in CPMB. A cleavage with the restriction enzyme SacII was then performed. The DNA resulting from these operations was separated by electrophoresis on 1% LGTA/TBE agarose gel (ref. CPMB).

The gel fragment containing the 0.4-kbp "AseI-blunt ends/SacII" insert was excised from the gel and purified according to the protocol described in section 5 above. The DNA of the clone pRPA-ML-711 was cut with the restriction enzyme HindIII [lacuna] located in the polylinker of the cloning vector pUC19, and the ends resulting from this cleavage were rendered blunt by treatment with the Klenow fragment of DNA polymerase I. A cleavage with the restriction enzyme SacII was then performed. The DNA resulting from these manipulations was separated by electrophoresis on 0.7% LGTA/TBE agarose gel (ref. CPMB).

AP/P/98/01195

The gel fragment containing the approximately 3.7-kbp HindIII-blunt ends/SacII insert was excised from the gel and purified according to the protocol described in section 5 above.

5 The two inserts were ligated, and 2 μ l of the ligation mixture were used to transform *E. coli* DH10B as described above in section 5.

The plasmid DNA content of different clones was analysed according to the procedure described for
10 pRPA-ML-711. One of the plasmid clones selected contains an approximately 1.45-kbp EcoRI-HindIII insert. The sequence of the terminal ends of this clone reveals that the 5' end of the insert corresponds exactly to the corresponding end of pRPA-ML-711, and
15 that the 3'-terminal end possesses the following sequence:

"5'-...AATTAAGCTCTAGAGTCGACCTGCAGGCATGCAAGCTT-3'".

The underlined sequence corresponds to the codon of the COOH-terminal amino acid asparagine, the
20 next codon corresponding to the translation stop codon. The nucleotides downstream correspond to sequence elements of the pUC19 polylinker. This clone comprising the pRPA-ML-711 sequence up to the translation
25 termination site of mature maize EPSPS and followed by sequences of the pUC 19 polylinker up to the HindIII site was designated pRPA-ML-712.

b) Modification of the 5' end of pRPA-ML-712:
construction of pRPA-ML-715:

The clone pRPA-ML-712 was cut with the restriction enzymes PstI and HindIII. The DNA resulting from these manipulations was separated by electrophoresis on 0.8% LGTA/TBE agarose gel (ref. CPMB). The gel fragment containing the 1.3-kbp PstI-EcoRI insert was excised from the gel and purified according to the protocol described in section 5 above. This insert was ligated in the presence of an equimolecular amount of each of the two partially complementary oligonucleotides of sequence:

Oligo 1: 5'-GAGCCGAGCTCCATGGCCGGCGCCGAGGAGATCGTGCTGCA-3'

Oligo 2: 5'-GCACGATCTCCTCGGCGCCGGCCATGGAGCTCGGCTC-3'

as well as in the presence of plasmid pUC19 DNA digested with the restriction enzymes BamHI and HindIII.

Two μ l of the ligation mixture were used to transform *E. coli* DH10B as described above in section 5. After analysis of the plasmid DNA content of different clones according to the procedure described above in section 5, one of the clones possessing an approximately 1.3-kbp insert was retained for subsequent analyses. The sequence of the 5'-terminal end of the selected clone reveals that the DNA sequence in this region is the following: sequence of the pUC19 polylinker from the EcoRI to the BamHI sites, followed by the sequence of the oligonucleotides used in the cloning, followed by the remainder of the sequence present in pRPA-ML-712. This clone was designated pRPA-

ML-713. This clone possesses a methionine ATG codon included in an NcoI site upstream of the N-terminal alanine codon of mature EPSP synthase. Furthermore, the alanine and glycine codons of the N-terminal end have
5 been preserved, but modified on the third variable base: initial GCGGGT gives modified GCCGGC.

The clone pRPA-ML-713 was cut with the restriction enzyme HindIII, and the ends of this cleavage were rendered blunt by treatment with the
10 Klenow fragment of DNA polymerase I. A cleavage with restriction enzyme SacI was then performed. The DNA resulting from these manipulations was separated by electrophoresis on 0.8% LGTA/TBE agarose gel (ref. CPMB). The gel fragment containing the 1.3-kbp
15 "HindIII-blunt ends/SacI" insert was excised from the gel and purified according to the protocol described in section 5 above. This insert was ligated in the presence of plasmid pUC19 DNA digested with restriction enzyme XbaI, and the ends of this cleavage were
20 rendered blunt by treatment with the Klenow fragment of DNA polymerase I. A cleavage with the restriction enzyme SacI was then performed. Two μ l of the ligation mixture were used to transform *E. coli* DH10B as described above in section 5. After analysis of the
25 plasmid DNA content of different clones according to the procedure described above in section 5, one of the clones possessing an approximately 1.3-kbp insert was retained for subsequent analyses. The sequence of the

terminal ends of the selected clone reveals that the DNA sequence is the following: sequence of the pUC19 polylinker from the EcoRI to SacI sites, followed by the sequence of the oligonucleotides used in the cloning from which the 4 bp GATCC of the oligonucleotide 1 described above have been deleted, followed by the remainder of the sequence present in pRPA-ML-712 up to the HindIII site and sequence of the pUC19 polylinker from XbaI to HindIII. This clone was designated pRPA-ML-715.

7. Obtaining of a cDNA coding for a mutated maize EPSPS

All the mutagenesis steps were carried out with the Pharmacia U.S.E. mutagenesis kit according to the supplier's instructions. The principle of this mutagenesis system is as follows: plasmid DNA is denatured by heat and reassociated in the presence of a molar excess of, on the one hand the mutagenesis oligonucleotide, and on the other hand an oligonucleotide enabling a unique restriction enzyme site present in the polylinker to be eliminated. After the reassociation step, synthesis of the complementary strand is carried out by the action of T4 DNA polymerase in the presence of T4 DNA ligase and gene 32 protein in a suitable buffer which is supplied. The synthesis product is incubated in the presence of the restriction enzyme for which the site is assumed to have disappeared by mutagenesis. The *E. coli* strain

possessing, in particular, the mutS mutation is used as host for the transformation of this DNA. After growth in liquid medium, the total plasmid DNA is prepared and incubated in the presence of the restriction enzyme used before. After these treatments, *E. coli* strain DH10B is used as host for the transformation. The plasmid DNA of the clones isolated is prepared, and the presence of the mutation introduced is verified by sequencing.

10 A)- modification of sites or sequences without in principle affecting the EPSPS-resistance character of maize to products which are competitive inhibitors of EPSP synthase activity: elimination of an internal NcoI site from pRPA-ML-715.

15 The pRPA-ML-715 sequence is numbered arbitrarily by placing the first base of the N-terminal alanine codon GCC at position 1. This sequence possesses an NcoI site at position 1217. The site-modification oligonucleotide possesses the sequence:

20 5'-CCACAGGATGGCGATGGCCTTCTCC-3'.

 After sequencing according to the references given above, the sequence read after mutagenesis corresponds to that of the oligonucleotide used. The NcoI site has indeed been eliminated, and the translation into amino acids in this region preserves the initial sequence present in pRPA-ML-715.

 This clone was designated pRPA-ML-716.

 The 1340-bp sequence of this clone is

presented in SEQ ID No. 2 and SEQ ID No. 3.

B)- sequence modifications enabling the EPSPS-resistance character of maize to products which are competitive inhibitors of EPSP synthase activity to be increased.

The following oligonucleotides were used:

a) mutation Thr 102 → Ile.

5'-GAATGCTGGAATCGCAATGCGGCCATTGACAGC-3'

b) mutation Pro 106 → Ser.

10 5'-GAATGCTGGAAGTCAATGCGGTCCTTGACAGC-3'

c) mutations Gly 101 → Ala and Thr 102 → Ile.

5'-CTTGGGGAATGCTGCCATCGCAATGCGGCCATTG-3'

d) mutations Thr 102 → Ile and Pro 106 → Ser.

5'-GGGGAATGCTGGAATCGCAATGCGGTCCTTGACAGC-3'

15 After sequencing, the sequence read after mutagenesis on the three mutated fragments is identical to the parent pRPA-ML-716 DNA sequence, with the exception of the mutagenized region which corresponds to that of the mutagenesis oligonucleotides used. These clones were designated: pRPA-ML-717 for the mutation Thr 102 → Ile, pRPA-ML-718 for the mutation Pro 106 → Ser, pRPA-ML-719 for the mutations Gly 101 → Ala and Thr 102 → Ile and pRPA-ML-720 for the mutations Thr 102 → Ile and Pro 106 → Ser.

The 1340-bp sequence of pRPA-ML-720 is presented in SEQ ID No. 4 and SEQ ID No. 5.

The 1395-bp NcoI-HindIII insert is the basis of all the constructions used for the transformation of plants for the introduction of resistance to herbicides which are competitive inhibitors of EPSPS, and especially glyphosate resistance. This insert will be designated in the remainder of the description "the maize EPSPS double mutant".

10 Example 2:

Glyphosate tolerance of the different mutants in vitro

2.a: Extraction of EPSP synthase

The different EPSP synthase genes are introduced in the form of an NcoI-HindIII cassette into the plasmid vector pTrc99a (Pharmacia, ref: 27-5007-01) cut with NcoI and HindIII. Recombinant *E. coli* DH10B bacteria overexpressing the different EPSP synthases are sonicated in 40 ml of buffer per 10 g of pelleted cells, and washed with this same buffer (200 mM Tris-HCl pH 7.8, 50 mM mercaptoethanol, 5 mM EDTA and 1 mM PMSF), to which 1 g of polyvinylpyrrolidone is added. The suspension is stirred for 15 minutes at 4°C and then centrifuged for 20 minutes at 27,000 g and 4°C.

25 Ammonium sulphate is added to the supernatant to bring the solution to 40% saturation with respect to ammonium sulphate. The mixture is centrifuged for 20 minutes at 27,000 g and 4°C. Ammonium sulphate is

added to the new supernatant to bring the solution to 70% saturation with respect to ammonium sulphate. The mixture is centrifuged for 30 minutes at 27,000 g and 4°C. The EPSP synthase present in this protein pellet
5 is taken up in 1 ml of buffer (20 mM Tris-HCl pH 7.8 and 50 mM mercaptoethanol). This solution is dialysed overnight against two litres of this same buffer at 4°C.

2.b: Enzyme activity

10 The activity of each enzyme, as well as its glyphosate resistance, is measured in vitro over 10 minutes at 37°C in the following reaction mixture: 100 mM maleic acid pH 5.6, 1 mM phosphoenolpyruvate, 3 mM shikimate 3-phosphate (prepared according to
15 Knowles P.F. and Sprinson D.B. 1970. Methods in Enzymol 17A, 351-352 from *Aerobacter aerogenes* strain ATCC 25597) and 10 mM potassium fluoride. The enzyme extract is added at the last moment after the addition of glyphosate, the final concentration of which varies
20 from 0 to 20 mM.

The activity is measured by assaying the phosphate liberated according to the technique of Tausky H.A. and Shorr E. 1953. J. Biol. Chem. 202, 675-685.

25 Under these conditions, the wild-type (WT) enzyme is already 85% inhibited at a glyphosate concentration of 0.12 mM. At this concentration, the mutant enzyme known as Ser106 is only 50% inhibited,

and the other three mutants, Ile102, Ile102/Ser106 and Ala101/Ile102, show little or no inhibition.

The glyphosate concentration has to be multiplied by ten, that is to say 1.2 mM, in order to produce a 50% inhibition of the mutant enzyme Ile102, the mutants Ile102/Ser106, Ala/Ile and Ala still not being inhibited.

It should be noted that the activity of the mutants Ala/Ile and Ala is not inhibited up to glyphosate concentrations of 10mM, and that that of the mutant Ile102/Ser106 is not reduced even if the glyphosate concentration is multiplied by 2, that is to say 20 mM.

Example 3:

Resistance of transformed tobacco plants

1-1- Transformation

The vector pRPA-RD-173 is introduced into *Agrobacterium tumefaciens* strain EHA101 (Hood et al., 1987) carrying the cosmid pTVK291 (Komari et al., 1986). The transformation technique is based on the procedure of Horsh et al. (1985).

1-2- Regeneration

The regeneration of PBD6 tobacco (source SEITA France) from leaf explants is carried out on a Murashige and Skoog (MS) basal medium comprising 30 g/l of sucrose as well as 200 µg/ml of kanamycin. The leaf explants are removed from plants cultivated in the greenhouse or *in vitro* and transformed according to the

leaf disc technique (Science, 1985, Vol. 227, pp. 1229-1231) in three successive steps: the first comprises the induction of shoots on a medium supplemented with 30 g/l of sucrose containing 0.05 mg/l of naphthylacetic acid (NAA) and 2 mg/l of benzylaminopurine (BAP) for 15 days. The shoots formed during this step are then developed for 10 days by culturing on an MS medium supplemented with 30 g/l of sucrose but not containing any hormone. Shoots which have developed are then removed and cultured on an MS rooting medium having half the content of salts, vitamins and sugar and not containing any hormone. After approximately 15 days, the rooted shoots are transferred to soil.

15 1-3- Glyphosate resistance

Twenty transformed plants were regenerated and transferred to the greenhouse for the construction of pRPA-RD-173. These plants were treated in the greenhouse at the 5-leaf stage with an aqueous suspension of RoundUp corresponding to 0.8 kg of glyphosate active substance per hectare.

The results correspond to the observation of phytotoxicity indices recorded 3 weeks after treatment. Under these conditions, it is found that the plants transformed with the construction pRPA-RD-173 display very good tolerance, whereas the untransformed control plants are completely destroyed.

These results show clearly the improvement

brought about by the use of a chimeric gene according to the invention for the same gene coding for glyphosate tolerance.

Example 4:

5 Transformation and selection of maize cells
BMS (Black Mexican Sweet) maize cells in an exponential growth phase are bombarded with the construction pRPA-RD-130 according to the principle and the protocol described by Klein et al. 1987 (Klein
10 T.M., Wolf E.D., Wu R. and Sandford J.C. (1987): High velocity microprojectiles for delivering nucleic acids into living cells, NATURE Vol. 327 pp. 70-73).

Two days after bombardment, the cells are transferred to the same medium containing 2 mM
15 N-(phosphomethyl)glycine.

After 8 weeks of selection on this medium, calluses which develop are selected, then amplified and analysed by PCR, and reveal clearly the presence of the chimeric OTP-EPSPS gene.

20 Cells not bombarded and grown on the same medium containing 2 mM N-(phosphomethyl)glycine are blocked by the herbicide and do not develop.

The transformed plants according to the invention may be used as parents for obtaining lines
25 and hybrids having the phenotypic character corresponding to the expression of the chimeric gene introduced.

Description of the constructions of the
plasmids

pRPA-RD-124: Addition of a "nos"
polyadenylation signal to pRPA-ML-720 with creation of
5 a cloning cassette containing the maize double mutant
EPSPS gene (Thr 102 → Ile and Pro 106 → Ser). pRPA-ML-
720 is digested with HindIII and treated with the
Klenow fragment of *E. coli* DNA polymerase I to produce
a blunt end. A second digestion is performed with NcoI,
10 and the EPSPS fragment is purified. The EPSPS gene is
then ligated with purified pRPA-RD-12 (a cloning
cassette containing the polyadenylation signal of
nopaline synthase) to give pRPA-RD-124. To obtain the
useful purified vector pRPA-RD-12, it was necessary for
15 the latter to be digested beforehand with SalI, treated
with Klenow DNA polymerase and then digested a second
time with NcoI.

pRPA-RD-125: Addition of an optimized transit
peptide (OTP) to pRPA-RD-124 with creation of a cloning
20 cassette containing the EPSPS gene targeted on the
plasmids. pRPA-RD-7 (European Patent Application
EP 652 286) is digested with SphI, treated with T4 DNA
polymerase and then digested with SpeI, and the OTP
fragment is purified. This OTP fragment is cloned into
25 pRPA-RD-124 which has previously been digested with
NcoI, treated with Klenow DNA polymerase to remove the
protruding 3' portion and then digested with SpeI. This
clone is then sequenced in order to ensure correct

translational fusion between the OTP and the EPSPS gene. pRPA-RD-125 is then obtained.

pRPA-RD-130: Addition of the H3C4 maize histone promoter and of *adh1* intron 1 sequences of
5 pRPA-RD-123 (Patent Application EP 507 698) to pRPA-RD-125 with creation of a cassette for expression in plants for the expression of the double mutant EPSPS gene in the tissues of monocotyledons. pRPA-RD-123 (a cassette containing the H3C4 maize histone promoter
10 fused with the *adh1* intron 1) is digested with *NcoI* and *SacI*. The DNA fragment containing the promoter derived from pRPA-RD-123 is then purified and ligated with pRPA-RD-125 which has previously been digested with *NcoI* and *SacI*.

15 pRPA-RD-159: Addition of the H4A748 *Arabidopsis* histone double promoter (Patent Application EP 507 698) to pRPA-RD-125 with creation of a cassette for expression in plants for the expression of the "OTP-double mutant EPSPS gene" gene in the tissues of
20 dicotyledons. pRPA-RD-132 (a cassette containing the H4A748 double promoter (Patent Application EP 507 698)) is digested with *NcoI* and *SacI*. The purified promoter fragment is then cloned into [lacuna] which has been digested with *EcoI* and *SacI*.

25 pRPA-RD-173: Addition of the "H4A748 promoter-OTP-double mutant EPSPS gene" gene of pRPA-RD-159 to plasmid pRPA-BL-150A (European Patent Application 508 909) with creation of an *Agrobacterium*

AP000886

28

tumefaciens transformation vector. pRPA-RD-159 is digested with NotI and treated with Klenow polymerase. This fragment is then cloned into pRPA-BL-150A with SmaI.

AP/P/98/01195

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Lebrun, Michel
 De Rube, Richard T
 Sailland, Alain

(ii) TITLE OF INVENTION: 5-enol pyruvylshikimate-3-phosphate
 synthase mutee, gene codant pour cette proteine et plantes
 transformees contenant ce gene

(iii) NUMBER OF SEQUENCES: 5

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Francois Chretien
 (B) STREET: 1420 rue Pierre Baizet
 (C) CITY: Lyon Cedex 09
 (E) COUNTRY: France
 (F) ZIP: 69263

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
 (B) FILING DATE:
 (C) CLASSIFICATION:

(vii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Chretien, Francois

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (33)72-29-26-46
 (B) TELEFAX: (33)72-29-28-43

(2) INFORMATION FOR SEQ ID NO:1:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1713 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Zea mays
 (B) STRAIN: Black Mexican Sweet
 (F) TISSUE TYPE: Callus

(vii) IMMEDIATE SOURCE:

(A) LIBRARY: lambda gt10
 (B) CLONE: pRPA-HL-711

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AATCAATTTC ACACAGGAAA CAGCTATGAC CATGATTACG AATTGGGGCC CCGGGCGGTG	60
ATCGGGGGCC GGCAGCGGGC GCGGGCGGTG AGCGGGGTGC CGAGGAGATC GTGCTGCGAG	120
CGATCAAGGA GATCTCGGGC ACGTCAAGC TCGCGGGTGC CAAGTGGCTT TCCACCGGA	180
TCTCTCTACT CGCGCGCGTG TCGAGGGGA CAACAGTGGT TGATMCGTG CTGACAGTG	240
AGGATGTCCA CTACATGCTC GCGGCGTTGA GGACTCTTGG TCTCTCTGTC GAAGCGGACA	300
AAGCTGCCAA AAGAGCTGTA GTTGTGGCT GTGGTGGAAA GTTCCGAGTT GAGGATGCCA	360
AAGAGGAAGT GCAGCTCTTC TGGGGAAAG CTGGAACTGC AATCGGGCCA TTGACAGCAG	420
CTGTTACTGC TCTGTGTGGA AATGCCAAGT ACGTGGTTGA TGGATTACCA AGAATGAGGG	480

AP/P/98/01195

AATGACCCAT TGGGACTTG GTTGTGGAT TGAAGCAKCT TGGTTCAGAT GTTGATTGT 510
 TCTTTTACAC TCACTTCTCA CTTTCTGTTG TCAATGGAAT CCGAGGACTA CCGTGTGGCA 600
 AATTCAGGT GTCTTCTTCC ATCAGCATTG ACTTACTTGG TGGCTTCTTG ATGGCTGCTC 640
 GTTGGCTCT TGGTATGTTG GAGATTGAAA TCATTGATAA ATTAATCTCC ATTCCTTAGG 720
 TGTAAATGAC ATTGAGATTG ATGGAGGTTT TTGGTGTGAA AGCAGAGCAT TGTGATAGCT 780
 GGGACAGATT CTACATTAA GAGGTCAAA AATACAAGTC CCTAATAAAT GCCTATGTTG 840
 AAGGTGATGC CTCAGGCCA AGCTATTCT TGGCTGGTGC TGCATTACT GGAGGACTG 900
 TGACTGTGGA AGGTGTGGC ACCACCATT TGCAGGTGA TGTGAAGTTT GCTGAGTAC 960
 TGGAGATGAT GGGAGCGAAG GTTACATGGA CCGAGACTAG CSTAAGTGT ACTGGCCDAC 1020
 CCGGGAGCC ATTTGGGAGG AAACACTCA AGGCGATTGA TGTCAAGATG AACAGATGC 1080
 CTGATGTGC CATGACTCTT GCTGTGGTTG CCTCTTTGC CGATGGCCCG ACAGCCATCA 1140
 GAGAGTGGC TTCTGGAGA GTAAAGGAGA CCGAGAGGAT GGTTCGATC CCGACGGAGC 1200
 TAACCAAGCT GGGAGCTCT GTTGAGGAG GCGCGACTA CTGCATCATC ACCCGCCGG 1260
 AGAAGGTGAA CGTGAAGCCG ATCGACAGT AGGAGSACA CAGGATGGCC ATGGCTTCT 1320
 CCTTGGCCG CTGTCCGAG GTCCCGTCA CCTCCGGGA CCTGGGTGC ACCCGAAGA 1380
 CCTCCCGCA CTACTTGGAT GTGCTGAGCA CTTTGTGCA GAATTAATAA AGGTGCGAT 1440
 ACTACCAGC AGCTTGATTG AAGTGAJAG CTGTGTCTGA GGAATACAT TTCTTTTGT 1500
 CTGTTTTCT CTTTACGGG ATTAAGTTT GAGTCTGAA CTTTGTGT TTGZAGCAG 1560
 TTTCTATTG GGATCTAAG TTTGTGCACT GTAAGCCAAA TTTCAATTCA AGAGTGGTC 1620
 GTTGGATAA TAAGAATAA AAATTAGCT TCAGTGAAA AAAAAAAAAA AAAAAAAAAA 1680
 AAAAAAAAAA AAAAAAAAAA AACCCGGAA TTC 1713

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1340 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Zea mays
 - (B) STRAIN: Black Mexican Sweet

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: pRPA-ML-716

- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 6..1337

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CCATG GCC GGC GGC GAG GAG ATC GTG CTG CAG CCC ATC AAG GAG ATC 47
 Ala Gly Ala Glu Glu Ile Val Leu Gln Pro Ile Lys Glu Ile
 1 5 10
 TCC GGC ACC GTC AAG CTG CCG GGG TCC AAG TCG CTT TCC AAC CGG ATC 95
 Ser Gly Thr Val Lys Leu Pro Gly Ser Lys Ser Leu Ser Asn Arg Ile
 15 20 25 30
 CTC CTA CTC GGC GGC CTG TCC GAG GGG ACA ACA GTG GTT GAT AAC CTG 143
 Leu Leu Leu Ala Ala Leu Ser Glu Gly Thr Thr Val Val Asp Asn Leu
 35 40 45
 CTG AAC AGT GAG GAT GTC CAC TAC ATG CTC GGG GCC TTG AGG ACT CTT 191
 Leu Asn Ser Glu Asp Val His Tyr Met Leu Gly Ala Leu Arg Thr Leu

AP/P/98/01195

	50	55	60		
GAT CTC TCT CTC GAA GGT GAC AAA GCT GCC AAA AGA GCT GTA GTT GTT					239
Gly Leu Ser Val Glu Ala Asp Lys Ala Ala Lys Arg Ala Val Val Val	65	70	75		
GAT TGT GTT GGA AAG TTC CCA GTT GAG GAT GCT AAA GAG GAA GTG CAG					287
Gly Cys Gly Gly Lys Phe Pro Val Glu Asp Ala Lys Glu Glu Val Gln	80	85	90		
CTC TTC TTG GGG AAT GCT GGA ACT GCA ATG CCG CCA TTG ACA GCA GCT					335
Leu Phe Leu Gly Asn Ala Gly Thr Ala Met Arg Pro Leu Thr Ala Ala	95	100	105	110	
GTT ACT GCT GCT GGT GGA AAT GCA ACT TAC GTG CTT GAT GGA GTA CCA					383
Val Thr Ala Ala Gly Gly Asn Ala Thr Tyr Val Leu Asp Gly Val Pro	115	120	125		
AGA ATG AGG GAG AGA CCC ATT GGC GAC TTG GTT GTC GGA TTG AAG CAG					431
Arg Met Arg Glu Arg Pro Ile Gly XHP Leu Val Val Gly Leu Lys Gln	130	135	140		
CTT GGT GCA GAT GTT GAT TGT TTC CTT GGC ACT GAC TGC CCA CCT GTT					479
Leu Gly Ala Asp Val Asp Cys Phe Leu Gly Thr Asp Cys Pro Pro Val	145	150	155		
GCT GTC AAT GGA ATC GGA GGG CTA CCT GGT GGC AAG GTC AAG CTG TCT					527
Arg Val Asn Gly Ile Gly Gly Leu Pro Gly Gly Lys Val Lys Leu Ser	160	165	170		
GGC TCC ATC AGC AGT CAG TAC TTG AGT GCC TTG CTG ATG GCT GCT CCT					575
Gly Ser Ile Ser Ser Gln Tyr Leu Ser Ala Leu Leu Met Ala Ala Pro	175	180	185	190	
TTG GCT CTT GGG GAT GTG GAG ACT GAA ATC ATT GAT AAA TTA ATC TCC					623
Leu Ala Leu Gly Asp Val Glu Ile Glu Ile Ile Asp Lys Leu Ile Ser	195	200	205		
ATT CCG TAC GTC GAA ATG ACA TTG AGA TTG ATG GAG CGT TTT GGT GTG					671
Ile Pro Tyr Val Glu Met Thr Leu Arg Leu Met Glu Arg Phe Gly Val	210	215	220		
AAA GCA GAG CAT TCT GAT AGC TGG GAC AGA TTC TAC ATT AAG GGA GGT					719
Lys Ala Glu His Ser Asp Ser Trp Asp Arg Phe Tyr Ile Lys Gly Gly	225	230	235		
CAA AAA TAC AAG TCC CCT AAA AAT GCC TAT GTT GAA GGT GAT GCC TCA					767
Gln Lys Tyr Lys Ser Pro Lys Asn Ala Tyr Val Glu Gly Asp Ala Ser	240	245	250		
AGC GCA AGC TAT TTC TTG GCT GGT GCT GCA ATT ACT GGA GGG ACT GTG					815
Ser Ala Ser Tyr Phe Leu Ala Gly Ala Ala Ile Thr Gly Gly Thr Val	255	260	265	270	
ACT GTG GAA GGT TGT GGC ACC ACC AGT TTG CAG GGT GAT GTG AAG TTT					863
Thr Val Glu Gly Cys Gly Thr Thr Ser Leu Gln Gly Asp Val Lys Phe	275	280	285		
GCT GAG GTA CTG GAG ATG ATG GGA GCG AAG GTT ACA TGG ACC GAG ACT					911
Ala Glu Val Leu Glu Met Met Gly Ala Lys Val Thr Trp Thr Glu Thr	290	295	300		
AGC GTA ACT GTT ACT GGC CCA CCG CCG GAG CCA TTT GGG AGG AAA CAC					959
Ser Val Thr Val Thr Gly Pro Pro Arg Glu Pro Phe Gly Arg Lys His	305	310	315		
CTC AAG GCG ATT GAT GTC AAC ATG AAC AAG ATG CCT GAT GTC GCC ATG					1007
Leu Lys Ala Ile Asp Val Asn Met Asn Lys Met Pro Asp Val Ala Met	320	325	330		
ACT CTT GCT GTG GTT GCC CTC TTT GCC GAT GGC CCG ACA GCC ATC AGA					1055
Thr Leu Ala Val Val Ala Leu Phe Ala Asp Gly Pro Thr Ala Ile Arg	335	340	345	350	
GAC GTG GCT TCC TGG AGA GTA AAG GAG ACC GAG AGG ATG GTT GCG ATC					1103
Asp Val Ala Ser Trp Arg Val Lys Glu Thr Glu Arg Met Val Ala Ile	355	360	365		
CGG ACC GAG CTA ACC AAG CTG GGA GCA TCT GTT GAG GAA GGG CCG GAC					1151
Arg Thr Glu Leu Thr Lys Leu Gly Ala Ser Val Glu Glu Gly Pro Asp					

AP/P/98/01195

AP000886

32

370	375	380	
TAC TCC ATC ATC ACC CGT CGT GAG AAG CTG AAC GTG ACC GGT ATC GAC			1199
Tyr Cys Ile Ile Thr Pro Pro Glu Lys Leu Asn Val Thr Ala Ile Asp	390	395	
ACT TAC GAC GAC CAC ACG ATG GCG ATG GCG TTC TCC CTC GCG GCG TGT			1217
Thr Tyr Asp Asp His Arg Met Ala Met Ala Phe Ser Leu Ala Ala Cys	405	410	
GCC GAG GTC CCC GTC ACC ATC CGG GAC CCT GGG TGC ACC CGG AAG ACC			1295
Ala Glu Val Pro Val Thr Ile Arg Asp Pro Gly Cys Thr Arg Lys Thr	420	425	430
TTC CCC GAC TAC TTC GAT GTG CTG AGC ACT TTC GTC AAG AAT			1337
Phe Pro Asp Tyr Phe Asp Val Leu Ser Thr Phe Val Lys Asn	435	440	
TAA			1340

(2) INFORMATION FOR SEQ ID NO:3:

- (1) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 444 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(11) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

Ala Gly Ala Glu Glu Ile Val Leu Gln Pro Ile Lys Glu Ile Ser Gly
 1           5           10           15
Thr Val Lys Leu Pro Gly Ser Lys Ser Leu Ser Asn Arg Ile Leu Leu
          20           25           30
Leu Ala Ala Leu Ser Glu Gly Thr Thr Val Val Asp Asn Leu Leu Asn
          35           40           45
Ser Glu Asp Val His Tyr Met Leu Gly Ala Leu Arg Thr Leu Gly Leu
          50           55           60
Ser Val Glu Ala Asp Lys Ala Ala Lys Arg Ala Val Val Val Gly Cys
          65           70           75           80
Gly Gly Lys Phe Pro Val Glu Asp Ala Lys Glu Glu Val Gln Leu Phe
          85           90           95
Leu Gly Asn Ala Gly Thr Ala Met Arg Pro Leu Thr Ala Ala Val Thr
          100          105          110
Ala Ala Gly Gly Asn Ala Thr Tyr Val Leu Asp Gly Val Pro Arg Met
          115          120          125
Arg Glu Arg Pro Ile Gly Asp Leu Val Val Gly Leu Lys Gln Leu Gly
          130          135          140
Ala Asp Val Asp Cys Phe Leu Gly Thr Asp Cys Pro Pro Val Arg Val
          145          150          155          160
Asn Gly Ile Gly Gly Leu Pro Gly Gly Lys Val Lys Leu Ser Gly Ser
          165          170          175
Ile Ser Ser Gln Tyr Leu Ser Ala Leu Leu Met Ala Ala Pro Leu Ala
          180          185          190
Leu Gly Asp Val Glu Ile Glu Ile Ile Asp Lys Leu Ile Ser Ile Pro
          195          200          205
Tyr Val Glu Met Thr Leu Arg Leu Met Glu Arg Phe Gly Val Lys Ala
          210          215          220
Glu His Ser Asp Ser Trp Asp Arg Phe Tyr Ile Lys Gly Gly Gln Lys
          225          230          235          240
Tyr Lys Ser Pro Lys Asn Ala Tyr Val Glu Gly Asp Ala Ser Ser Ala
          245          250          255
    
```

AP/P/98/01195

Ser Tyr Phe Leu Ala Gly Ala Ala Ile Thr Gly Gly Thr Val Thr Val
 260 265 270
 Glu Gly Cys Gly Thr Thr Ser Leu Gln Gly Asp Val Lys Phe Ala Glu
 275 280 285
 Val Leu Glu Met Met Gly Ala Lys Val Thr Trp Thr Glu Thr Ser Val
 290 295 300
 Thr Val Thr Gly Pro Pro Arg Glu Pro Phe Gly Arg Lys His Leu Lys
 305 310 315 320
 Ala Ile Asp Val Asn Met Asn Lys Met Pro Asp Val Ala Met Thr Leu
 325 330 335
 Ala Val Val Ala Leu Phe Ala Asp Gly Pro Thr Ala Ile Arg Asp Val
 340 345 350
 Ala Ser Trp Arg Val Lys Glu Thr Glu Arg Met Val Ala Ile Arg Thr
 355 360 365
 Glu Leu Thr Lys Leu Gly Ala Ser Val Glu Glu Gly Pro Asp Tyr Cys
 370 375 380
 Ile Ile Thr Pro Pro Glu Lys Leu Asn Val Thr Ala Ile Asp Thr Tyr
 385 390 395 400
 Asp Asp His Arg Met Ala Met Ala Phe Ser Leu Ala Ala Cys Ala Glu
 405 410 415
 Val Pro Val Thr Ile Arg Asp Pro Gly Cys Thr Arg Lys Thr Phe Pro
 420 425 430
 Asp Tyr Phe Asp Val Leu Ser Thr Phe Val Lys Asn
 435 440

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1340 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Zea mays
- (B) STRAIN: Black Mexican Sweet

(vii) IMMEDIATE SOURCE:

- (B) CLONE: pRPA-HL-720

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 6..1337

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCATG GGC GGC GGC GAG GAG ATC GTG CTG CAG CCC ATC AAG GAG ATC	47
Ala Gly Ala Glu Glu Ile Val Leu Gln Pro Ile Lys Glu Ile	
1 5 10	
TCC GGC ACC GTC AAG CTG CCG GGG TCC AAG TCG CTT TCC AAC CGG ATC	95
Ser Gly Thr Val Lys Leu Pro Gly Ser Lys Ser Leu Ser Asn Arg Ile	
15 20 25 30	
CTC CTA CTC GGC GGC CTG TCC GAG GGG ACA ACA GTG GTT GAT AAC CTG	143
Leu Leu Leu Ala Ala Leu Ser Glu Gly Thr Thr Val Val Asp Asn Leu	
35 40 45	
CTG AAC AGT GAG GAT GTC CAC TAC ATG CTC GGG GCC TTG AGG ACT CTT	191
Leu Asn Ser Glu Asp Val His Tyr Met Leu Gly Ala Leu Arg Thr Leu	
50 55 60	
GGT CTC TCT GTC GAA GCG GAC AAA GCT GCC AAA AGA GCT GEA GTT GTT	239
Gly Leu Ser Val Glu Ala Asp Lys Ala Ala Lys Arg Ala Val Val Val	
65 70 75	

AP/P/98/01195

AP000886

34

GAC	TGT	GTT	GGA	AAG	TTC	CCA	GTT	GAG	GAT	GCT	AAA	GAG	GAA	GTG	CAG	287
Gly	Cys	Gly	Gly	Lys	Phe	Pro	Val	Glu	Asp	Ala	Lys	Glu	Glu	Val	Gln	
	40					45					90					
GTC	TTC	TTG	GGG	AAT	GCT	GGA	ATC	GCA	ATG	GGT	TCC	TTG	ACA	GCA	GCT	335
Leu	Phe	Leu	Gly	Asn	Ala	Gly	Ile	Ala	Met	Arg	Ser	Leu	Thr	Ala	Ala	
	55			100					105						110	
GTT	ACT	GCT	GCT	GGT	GGA	AAT	GCA	ACT	TAC	GTG	CTT	GAT	GGA	GTA	CCA	383
Val	Thr	Ala	Ala	Gly	Asn	Ala	Thr	Tyr		Val	Leu	Asp	Gly	Val	Pro	
				115				120						125		
AGA	ATG	AGG	GAG	AGA	CCC	ATT	GGC	GAC	TTG	GTT	GTC	GGA	TTG	AAG	CAG	431
Arg	Met	Arg	Glu	Arg	Pro	Ile	Gly	Asp	Leu	Val	Val	Gly	Leu	Lys	Gln	
			130					135					140			
CTT	GGT	GCA	GAT	GTT	GAT	TGT	TTC	CTT	GGC	ACT	GAC	TGC	CCA	CCT	GTT	479
Leu	Gly	Ala	Asp	Val	Asp	Cys	Phe	Leu	Gly	Thr	Asp	Cys	Pro	Pro	Val	
			145				150					155				
CGT	GTC	AAT	GGA	ATC	GGA	GGG	GTA	CCT	GGT	GGC	AAG	GTC	AAG	CTG	TCT	527
Arg	Val	Asn	Gly	Ile	Gly	Gly	Leu	Pro	Gly	Gly	Lys	Val	Lys	Leu	Ser	
	160					165					170					
GGC	TCC	ATC	AGC	AGT	CAG	TAC	TTG	AGT	GCC	TTG	CTG	ATG	GCT	GCT	CCT	575
Gly	Ser	Ile	Ser	Ser	Gln	Tyr	Leu	Ser	Ala	Leu	Leu	Met	Ala	Ala	Pro	
	175				180					185					190	
TTG	GCT	CTT	GGG	GAT	GTG	GAG	ATT	GAA	ATC	ATT	GAT	AAA	TTA	ATC	TCC	623
Leu	Ala	Leu	Gly	Asp	Val	Glu	Ile	Glu	Ile	Ile	Asp	Lys	Leu	Ile	Ser	
				195				200						205		
ATT	CCG	TAC	GTC	GAA	ATG	ACA	TTG	AGA	TTG	ATG	GAG	CGT	TTT	GGT	GTG	671
Ile	Pro	Tyr	Val	Glu	Met	Thr	Leu	Arg	Leu	Met	Glu	Arg	Phe	Gly	Val	
			210				215						220			
AAA	GCA	GAG	CAT	TCT	GAT	AGC	TGG	GAC	AGA	TTC	TAC	ATT	AAG	GGA	GGT	719
Lys	Ala	Glu	His	Ser	Asp	Ser	Trp	Asp	Arg	Phe	Tyr	Ile	Lys	Gly	Gly	
			225				230					235				
CAA	AAA	TAC	AAG	TCC	CCT	AAA	AAT	GCC	TAT	GTT	GAA	GGT	GAT	GCC	TCA	767
Gln	Lys	Tyr	Lys	Ser	Pro	Lys	Asn	Ala	Tyr	Val	Glu	Gly	Asp	Ala	Ser	
	240					245					250					
AGC	GCA	AGC	TAT	TTC	TTG	GCT	GGT	GCT	GCA	ATT	ACT	GGA	GGG	ACT	GTG	815
Ser	Ala	Ser	Tyr	Phe	Leu	Ala	Gly	Ala	Ala	Ile	Thr	Gly	Gly	Thr	Val	
				260					265						270	
ACT	GTG	GAA	GGT	TGT	GGC	ACC	ACC	AGT	TTG	CAG	GGT	GAT	GTG	AAG	TTT	863
Thr	Val	Glu	Gly	Cys	Gly	Thr	Thr	Ser	Leu	Gln	Gly	Asp	Val	Lys	Phe	
				275				280						285		
GCT	GAG	GTA	CTG	GAG	ATG	ATG	GGA	GCG	AAG	GTT	ACA	TGG	ACC	GAG	ACT	911
Ala	Glu	Val	Leu	Glu	Met	Met	Gly	Ala	Lys	Val	Thr	Trp	Thr	Glu	Thr	
			290					295					300			
AGC	GTA	ACT	GTT	ACT	GGC	CCA	CCG	CGG	GAG	CCA	TTT	GGG	AGG	AAA	CAC	959
Ser	Val	Thr	Val	Thr	Gly	Pro	Pro	Arg	Glu	Pro	Phe	Gly	Arg	Lys	His	
			305			310						315				
CTC	AAG	GCG	ATT	GAT	GTC	AAC	ATG	AAC	AAG	ATG	CCT	GAT	GTC	GCC	ATG	1007
Leu	Lys	Ala	Ile	Asp	Val	Asn	Met	Asn	Lys	Met	Pro	Asp	Val	Ala	Met	
	320					325					330					
ACT	CTT	GCT	GTG	GTT	GCC	CTC	TTT	GCC	GAT	GGC	CCG	ACA	GCC	ATC	AGA	1055
Thr	Leu	Ala	Val	Val	Ala	Leu	Phe	Ala	Asp	Gly	Pro	Thr	Ala	Ile	Arg	
	335				340				345						350	
GAC	GTG	GCT	TCC	TGG	AGA	GTA	AAG	GAG	ACC	GAG	AGG	ATG	GTT	GCG	ATC	1103
Asp	Val	Ala	Ser	Arg	Val	Lys	Glu	Thr	Glu	Arg	Met	Val	Val	Ala	Ile	
			355					360						365		
CGG	ACG	GAG	CTA	ACC	AAG	CTG	GGA	GCA	TCT	GTT	GAG	GAA	GGG	CCG	GAC	1151
Arg	Thr	Glu	Leu	Thr	Lys	Leu	Gly	Ala	Ser	Val	Glu	Glu	Gly	Pro	Asp	
			370					375					380			
TAC	TGC	ATC	ATC	ACG	CCG	CCG	GAG	AAG	CTG	AAC	GTG	ACG	GCG	ATC	GAC	1199
Tyr	Cys	Ile	Ile	Thr	Pro	Pro	Glu	Lys	Leu	Asn	Val	Thr	Ala	Ile	Asp	
			385				390						395			

AP/P/98/01195

ACT TAC GAC GAC CAC ARG ATG GGT ATG GCG TTC TCC CTT GCG GCG TGT 1247
 The Tyr Asp Asp His Arg Met Ala Met Ala Phe Ser Leu Ala Ala Cys
 400 405 410

GCT GAG GTC CCC GTC ACC ATC GGG GAC CCT GAG TGC ACC GGG AAG ACC 1295
 Ala Glu Val Pro Val Thr Ile Arg Asp Pro Gly Cys Thr Arg Lys Thr
 415 420 425 430

TTC CCC GAC TAC TTC GAT GTG CTG AGC ACT TTC GTC AAG AAT 1337
 Phe Pro Asp Tyr Phe Asp Val Leu Ser Thr Phe Val Lys Asn
 435 440

TAA 1340

(2) INFORMATION FOR SEQ ID NO:5:

- (1) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 444 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(11) MOLECULE TYPE: protein

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Ala Gly Ala Glu Glu Ile Val Leu Gln Pro Ile Lys Glu Ile Ser Gly
 1 5 10 15

Thr Val Lys Leu Pro Gly Ser Lys Ser Leu Ser Asn Arg Ile Leu Leu
 20 25 30

Leu Ala Ala Leu Ser Glu Gly Thr Thr Val Val Asp Asn Leu Leu Asn
 35 40 45

Ser Glu Asp Val His Tyr Met Leu Gly Ala Leu Arg Thr Leu Gly Leu
 50 55 60

Ser Val Glu Ala Asp Lys Ala Ala Lys Arg Ala Val Val Val Gly Cys
 65 70 75 80

Gly Gly Lys Phe Pro Val Glu Asp Ala Lys Glu Glu Val Gln Leu Phe
 85 90 95

Leu Gly Asn Ala Gly Ile Ala Met Arg Ser Leu Thr Ala Ala Val Thr
 100 105 110

Ala Ala Gly Gly Asn Ala Thr Tyr Val Leu Asp Gly Val Pro Arg Met
 115 120 125

Arg Glu Arg Pro Ile Gly Asp Leu Val Val Gly Leu Lys Gln Leu Gly
 130 135 140

Ala Asp Val Asp Cys Phe Leu Gly Thr Asp Cys Pro Pro Val Arg Val
 145 150 155 160

Asn Gly Ile Gly Gly Leu Pro Gly Gly Lys Val Lys Leu Ser Gly Ser
 165 170 175

Ile Ser Ser Gln Tyr Leu Ser Ala Leu Leu Met Ala Ala Pro Leu Ala
 180 185 190

Leu Gly Asp Val Glu Ile Glu Ile Ile Asp Lys Leu Ile Ser Ile Pro
 195 200 205

Tyr Val Glu Met Thr Leu Arg Leu Met Glu Arg Phe Gly Val Lys Ala
 210 215 220

Glu His Ser Asp Ser Trp Asp Arg Phe Tyr Ile Lys Gly Gly Gln Lys
 225 230 235 240

Tyr Lys Ser Pro Lys Asn Ala Tyr Val Glu Gly Asp Ala Ser Ser Ala
 245 250 255

Ser Tyr Phe Leu Ala Gly Ala Ala Ile Thr Gly Gly Thr Val Thr Val
 260 265 270

Glu Gly Cys Gly Thr Thr Ser Leu Gln Gly Asp Val Lys Phe Ala Glu
 275 280 285

AP/P/98/01195

AP000886

36

Val Leu Glu Met Met Gly Ala Lys Val Thr Trp Thr Glu Thr Ser Val
270 295

Thr Val Thr Gly Pro Pro Arg Glu Pro Phe Gly Arg Lys His Leu Lys
305 310 315

Ala Ile Asp Val Asn Met Asn Lys Met Pro Asp Val Ala Met Thr Leu
325 330 335

Ala Val Val Ala Leu Phe Ala Asp Gly Pro Thr Ala Ile Arg Asp Val
340 345 350

Ala Ser Trp Arg Val Lys Glu Thr Glu Arg Met Val Ala Ile Arg Thr
355 360 365

Glu Leu Thr Lys Leu Gly Ala Ser Val Glu Glu Gly Pro Asp Tyr Cys
370 375 380

Ile Ile Thr Pro Pro Glu Lys Leu Asn Val Thr Ala Ile Asp Thr Tyr
385 390 395

Asp Asp His Arg Met Ala Met Ala Phe Ser Leu Ala Ala Cys Ala Glu
405 410 415

Val Pro Val Thr Ile Arg Asp Pro Gly Cys Thr Arg Lys Thr Phe Pro
420 425 430

Asp Tyr Phe Asp Val Leu Ser Thr Phe Val Lys Asn
435 440

AP/P/98/01195

CLAIMS

- 5 1. A DNA sequence coding for a mutated 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), characterized in that the EPSPS is from maize and comprises a first mutation consisting in the threonine 102 → isoleucine substitution and a second mutation consisting in a substitution of proline 106 by serine.
- 10 2. DNA sequence coding for a mutated 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), characterised in that it comprises the coding region of the DNA sequence represented in SEQ ID No. 2, comprising a first mutation consisting in the threonine 102 isoleucine substitution and a second mutation consisting in a substitution of proline 106 by serine.
- 15 3. DNA sequence coding for a mutated 5-enolpyruvylshikimate -3-phosphate synthase (EPSPS), characterised in that it comprises the coding sequence of SEQ ID No. 4.
- 20 4. A mutated 5-enolpyruvylshikimate -3-phosphate synthase (EPSPS), characterized in that the EPSPS is maize and comprises a first mutation consisting in the substitution of threonine 102 by isoleucine and a second mutation consisting in a substitution of proline 106 by serine.
- 25 5. A mutated 5-enolpyruvylshikimate -3-phosphate synthase (EPSPS), characterized in that it comprises the peptide sequence of SEQ ID No. 5.
- 30 6. Chimeric gene comprising a coding sequence as well as regulatory elements at position 5' and 3' which are heterologous and capable of functioning in plants, characterized in that it comprises as coding sequence at least one sequence according to any one of claims 1 to 3.

7. Chimeric gene according to claim 6, characterized in that it comprises a plant virus promoter.

5 8. Chimeric gene according to claim 6, characterized in that it comprises a plant promoter.

9. Chimeric gene according to claim 8 in which the plant promoter is α -tubulin, histone, intron or actin.

10 10. Chimeric gene according to any one of claims 6 to 9, characterised in that it comprises a zone coding for a transit peptide.

11. Chimeric gene according to claim 10, characterized in that it comprises one or more transit peptide units.

15 12. Vector for the transformation of plants, characterized in that it comprises at least one gene according to any one of claims 6 to 11.

20 13. Plant cell, characterized in that it comprises at least one gene according to any one of claims 6 to 11.

25 14. Method for the production of plants with improved tolerance to a herbicide having EPSP synthase as its target, characterised in that plant cells or protoplasts are transformed with a gene according to any one of claims 6 to 11, and in that the transformed cells are subjected to a regeneration.

30 15. Method as claimed in claim 14, characterised in that the plants with improved tolerance are used for the preparation of hybrid lines.

AP 0 0 0 8 8 6 / 0 1 1 9 5

16. Method of treatment of plants with a herbicide having EPSPS as its target, characterized in that the herbicide is applied to plants comprising plant cells according to Claim 13.

17. Method according to claim 16, characterized in that glyphosate or a glyphosate precursor is applied.

AP/98/01195