



US00PP20253P2

(12) **United States Plant Patent**
Handley

(10) **Patent No.:** **US PP20,253 P2**

(45) **Date of Patent:** **Sep. 1, 2009**

(54) **ERYSIMUM PLANT NAMED 'JENNY BROOK'**

(58) **Field of Classification Search** Plt./263.1
See application file for complete search history.

(50) Latin Name: *Erysimum linifolium*
Varietal Denomination: **Jenny Brook**

(56) **References Cited**

(76) Inventor: **Roger Handley**, Caerolweg, Lon
Glanfred, Llandre, Bow Street,
Ceredigion (GB), SY24 5BY

PUBLICATIONS

Upov Plant Variety Database Mar. 2008 search for PBR
20060957 for cultivar Jenny Brook. p. 1. 2008.*

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

* cited by examiner

Primary Examiner—Annette H Para
(74) *Attorney, Agent, or Firm*—Jondle & Associates, P.C.

(21) Appl. No.: **12/079,984**

(57) **ABSTRACT**

(22) Filed: **Mar. 31, 2008**

A new *Erysimum* plant particularly distinguished by flowers
which change color through the season starting out as
purple-mauve changing to peach and by its upright, long
flower stems that are held above green linear foliage is dis-
closed.

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./263.1**

1 Drawing Sheet

1

2

Genus and species: *Erysimum linifolium*
Variety denomination: 'Jenny Brook'.

the colors shown are as true as can be reasonably obtained by
conventional photographic procedures.

BACKGROUND OF THE NEW PLANT

DESCRIPTION OF THE NEW CULTIVAR

The present invention comprises a new and distinct culti-
var of *Erysimum*, botanically known as *Erysimum linifolium*
and hereinafter referred to by the cultivar name 'Jenny
Brook'. The new cultivar was discovered in a cultivated gar-
den in Cerdigion, Wales in 2001. Both the male and female
parents are unknown individual plants of *Erysimum*. 'Jenny
Brook' was first propagated in 2001 in Cerdigion, Wales.

5 The following detailed descriptions set forth the distinc-
tive characteristics of 'Jenny Brook'. The data which define
these characteristics were collected from asexual reproduc-
tions carried out in Lancaster, Pa. The plant history was
taken on one-year-old plants grown outdoors. Color readings
10 were taken outdoors under natural light. Color references
primarily to the R.H.S. Colour Chart of The Royal Horticul-
tural Society of London (R.H.S.) (2001 edition).

The new cultivar was discovered in Cerdigion, Wales in
2001 and has been asexually reproduced repeatedly by veg-
etative cuttings over a 7-year period in Cerdigion, Wales and
Lancaster, Pa. The present invention has been found to retain
its distinctive characteristics through successive asexual
propagations.

DESCRIPTION OF THE NEW PLANT

Plant Breeder's Rights for this cultivar were applied for in
the European Union on Apr. 19, 2006. 'Jenny Brook' has not
been made publicly available more than one year prior to
filing of this application.

Classification:

Family.—Brassicaceae.
Botanical.—*Erysimum linifolium*.
Common name.—Perennial wallflower.

SUMMARY OF THE INVENTION

The following are the most outstanding and distinguishing
characteristics of this new cultivar when grown under nor-
mal horticultural practices in Cerdigion, Wales and
Lancaster, Pa.

Parentage:

Female parent.—Unknown.
Male parent.—Unknown.

1. Flowers change color through the season starting out as
purple-mauve changing to peach; and
2. Upright, long flower stems that are held above green
linear foliage.

Plant:

Type.—Perennial.
25 *Habit*.—Woody base, branching, numerous linear
shaped leaves.
Height (from top of soil).—60.0 cm.
Width (horizontal plant diameter).—60.0 cm.
Propagation.—Tip cuttings, tissue culture.
30 *Time to produce a finished flowering plant*.—8 weeks.
Time to initiate and develop roots.—10 days.
Root description.—White, fibrous.

DESCRIPTION OF THE PHOTOGRAPH

This new *Erysimum* plant is illustrated by the accompany-
ing photograph which shows a close-up of the inflorescence;

Stems:

Average number.—8.
35 *Length of stems/branches (from the base of the stem to
the tip)*.—20.0 cm.
Internode length.—0.5 cm.
Diameter of branches (from midpoint).—0.3 cm.
Stem color.—RHS 137A (Green).

- Anthocyanin*.—RHS 59A to RHS 59B (Red-Purple).
Texture.—Rough.
- Leaves:
Arrangement.—Alternate.
Size.—Length: 5.0 cm to 11.0 cm. Width: 0.5 cm.
Shape.—Linear.
Apex.—Acute.
Base.—Cuneate.
Margin.—Undulate.
Texture.—Upper surface: Rough. Lower surface: Smooth.
Immature leaf.—Color: Upper surface: RHS 137B (Green). Lower surface: RHS 137C (Green).
Mature leaf.—Color: Upper surface: RHS 137A (Green). Lower surface: RHS 137D (Green).
Venation pattern.—Pinnate, very faint.
Venation color.—Upper surface: RHS 137A (Green). Lower surface: RHS 137D (Green).
Petioles.—Absent.
- Flower bud:
Shape.—Elliptic.
Size.—Length: 0.5 cm. Diameter: 0.3 cm.
Color at tight bud.—RHS 59A (Red-Purple).
Texture.—Smooth.
- Inflorescence:
Blooming habit (flowering season).—May to July and August.
Inflorescence type.—Raceme.
Average number of flowering stems.—20.
Average number of flowers and buds per stem.—30.
Lastingness of individual blooms on the plant.—10 days.
Fragrance.—Slight.
Inflorescence length.—30.0 cm to 40.0 cm.
Inflorescence diameter.—5.0 cm.
- Flowers:
Type.—Single flower, four petals, borne in dense elongating corymb like racemes.
Shape.—Cruciform.
Size.—Length or depth: 2.0 cm. Width or diameter: 3.0 cm.
Petals.—Number per flower: 4. Length: 1.0 cm. Width: 1.0 cm. Shape: Obocordate. Apex: Retuse. Base: Attenuate. Margin: Entire. Texture: Upper surface: Smooth. Lower surface: Smooth. Color: Upper sur-

- face: RHS 35C (Orange-Red). Lower surface: RHS 35D (Orange-Red).
- Calyx arrangement*.—Tubular, deeply cut to form 4 segments.
- Sepals*.—Quantity per flower: 4. Length: 1.0 cm. Width: 0.3 cm. Shape: Tubular. Apex: Acute. Base: Cuneate. Margin: Entire. Texture: Upper surface: Slightly rough. Lower surface: Smooth. Color: Upper surface: RHS 59A (Red-Purple). Lower surface: RHS 59D (Red-Purple).
- Peduncles:
Length.—30.0 cm to 40.0 cm.
Diameter.—0.5 cm.
Aspect.—Erect.
Texture.—Rough.
Color.—RHS 137A (Green).
- Pedicels:
Length.—1.0 cm.
Diameter.—0.2 cm.
Texture.—Slightly rough.
Color.—RHS 137A (Green).
- Reproductive organs:
Stamens.—Quantity per flower: 5. Filament length: 0.8 cm. Filament diameter: 0.1 cm. Anthers: Shape: Ligulate Length: 0.3 cm. Diameter: 0.1 cm. Color: RHS 152A (Yellow-Green). Pollen amount: Very sparse. Pollen color: RHS 153A (Yellow-Green).
Pistils.—Number per flower: 1. Length: 1.0 cm. Diameter: 0.1 cm. Stigma: Length: 0.01 cm. Diameter: 0.1 cm. Shape: Rounded. Color: RHS 150C (Yellow-Green). Style: Length: 0.9 cm. Diameter: 0.1 cm. Shape: Tubular. Color: RHS 150C (Yellow-Green).
 Fruit and seed set: Has not been observed.
 Disease and insect resistance: Typical of *Erysimum*.

COMPARISON WITH COMMERCIAL
 CULTIVARS

‘Jenny Brook’ differs from the commercial cultivar ‘Bowles Mauve’ (unpatented) in that ‘Jenny Brook’ has flowers that change color through the flowering season from purple-mauve to peach, while ‘Bowles Mauve’ has flowers that are lilac and pink.

I claim:

1. A new and distinct cultivar of *Erysimum* plant as shown and described herein.

* * * * *

