



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 341 071 B1**

EUROPEAN PATENT SPECIFICATION

- (45) Date of publication of patent specification: **03.08.94** (51) Int. Cl.⁵: **C11D 1/83, C11D 1/94, C11D 3/32**
- (21) Application number: **89304522.9**
- (22) Date of filing: **05.05.89**

(54) Detergent compositions.

- | | |
|---|--|
| <p>(30) Priority: 06.05.88 GB 8810820</p> <p>(43) Date of publication of application:
08.11.89 Bulletin 89/45</p> <p>(45) Publication of the grant of the patent:
03.08.94 Bulletin 94/31</p> <p>(84) Designated Contracting States:
CH DE ES FR GB IT LI NL SE</p> <p>(56) References cited:
EP-A- 0 070 076
EP-A- 0 155 737
EP-A- 0 216 301
US-A- 4 668 422</p> <p>(73) Proprietor: UNILEVER PLC
Unilever House
Blackfriars
P.O. Box 68
London EC4P 4BO(GB)</p> <p>(84) Designated Contracting States:
GB</p> <p>(73) Proprietor: UNILEVER N.V.
Weena 455
NL-3013 AL Rotterdam(NL)</p> | <p>(84) Designated Contracting States:
CH DE ES FR IT LI NL SE</p> <p>(72) Inventor: Naik, Appaya Raghunath
8 Kilmalcolm Close
Oxton
Birkenhead Merseyside, L43 9OT(GB)</p> <p>(74) Representative: Elliott, Peter William et al
Unilever plc
Patent Division
Colworth House
Sharnbrook
Bedford MK44 1LO (GB)</p> |
|---|--|

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

EP 0 341 071 B1

DescriptionTECHNICAL FIELD

5 The present invention relates to novel detergent compositions and especially, but not exclusively, to foaming liquid compositions useful for light-duty applications such as manual dishwashing. The compositions may also have use for fabric washing, cleaning carpets and as shampoos.

DESCRIPTION OF THE PRIOR ART

10

Light-duty liquid detergent compositions such as are suitable for use in washing dishes are well known. Many commercially-available formulations are based on a sulphate - or sulphonate-type anionic detergent, especially alkyl sulphate, alkyl benzene sulphonate and alkane sulphonate, in conjunction with an alkyl polyethoxy sulphate (alkyl ether sulphate). The sulphonate-type detergent generally predominates.

15

The use of conventional dishwashing liquids based on alkyl benzene sulphonate/alkyl ether sulphate, or alkyl sulphate/alkyl ether sulphate has been found to have a deleterious effect on the hand condition of users. Mildness in detergent products, particularly in washing-up liquids, has hence become a desirable quality.

20

Alkyl mono- and poly-glycosides are known and have been disclosed in a number of prior art patents and literature publications. For example, Boettner (US3219656) discusses the utility of alkyl mono-glycosides as foam stabilisers for anionic and nonionic surfactants. Lew (US 3772269) discloses that decyl glucoside containing less than 1.6 glucose units acts as a good foamer. Several Procter and Gamble disclosures (eg EP 0 070 074, EP 0 070 075 and EP 0 070 076) discuss the utility of alkylpolyglucosides having greater than 1.5 glucose units in detergent compositions. Henkel (EP 216 301 A) disclose the beneficial use of alkyl glycosides having from 1 to 1.4 glucose units which are incorporated in dishwashing liquids comprising ether sulphate and ethanolamide.

25

SUMMARY OF THE INVENTION

30

The present invention relates to safe, mild liquid detergents with a good foam stability and a good cleansing ability based upon alkyl polyglycosides. The invention aims to provide novel, high-foaming nonionic-based liquid detergents comprising an alkyl polyglycoside. It is a further aim to provide a mild, ecologically safe liquid detergent.

Accordingly, the present invention relates to an aqueous liquid detergent composition comprising:

35

- (a) a C₈-C₁₈ alkyl polyglycoside surfactant having an average degree of polymerisation of from 1 to 3;
- (b) an anionic surfactant;
- (c) a surface-active betaine and/or amine oxide; and optionally
- (d) an ethanolamide.

40

wherein the amount of anionic surfactant is not greater than 1.5 times (on a molar basis) the level of betaine and/or amine oxide and wherein the level of betaine and/or amine oxide and, if present, ethanolamide is from 12 to 30 wt% of the total active.

DESCRIPTION OF PREFERRED EMBODIMENTS

45

Alkyl Polyglycoside

50

Alkyl polyglycosides, though nonionic in nature, differ from conventional ethoxylated nonionics such as alkyl ethoxylates or alkylphenol ethoxylates in having a different type of hydrophilic group and in consequence exhibit different physical, chemical and hence functional properties compared to conventional ethoxylated nonionics. For example, ethoxylates are low melting solids or liquids at room temperature, are completely miscible in non-polar solvents and their aqueous solutions have cloud points. In comparison, alkyl polyglycosides are high melting solids, sparingly soluble in non-polar solvents and their aqueous solutions do not show cloud points. Alkyl polyglycosides also generally give lower surface tensions and give a more stable foam compared to comparable, conventional nonionics. Alkyl polyglycosides are also prepared by different routes and can be produced entirely based on natural raw materials. Hence the compounds and compositions made therewith can be expected to be biologically safer and more acceptable than conventional nonionics and compositions containing them. Impurities present will also differ from conventional nonionics.

55

Alkyl polyglycosides may be manufactured by the Fischer glycoxidation of a detergent alcohol, eg C₈-C₁₈



5

Specification of the ROG_x product depends on control of the ratios of the reactants. A product will be a ratio of, eg ROG₁:ROG₂:ROG₃.

The alkyl polyglycosides used in the present invention have a hydrophobic group containing 8 to 20 carbon atoms, preferably about 10 to about 16, most preferably from 12 to 14, and a polysaccharide hydrophilic group containing from 1 to 3, preferably 1 to 1.5, most preferably 1.2 to 1.4 saccharide units on average. The saccharide unit may be, for example, a galactoside, glucoside, fructoside or glycosyl. Mixtures thereof may be used.

Preferred alkyl polyglycosides are APG 500 and APG 550 from Horizon (APG is a trademark). APG 500 has an average degree of polymerisation of 1.4, APG 550 of 1.8. US 4 599 188 (Llenado) gives further description and characterisation of alkyl polyglycosides.

The alkyl polyglycoside is preferably the major active component and may, for example, be present at from 30 to 60 wt% of the total active.

Anionic Active

20

The co-active anionic surfactant used according to the present invention may be of the sulphate or sulphonate type, or a mixture thereof. The sulphates may be conventional anionic synthetic detergents, and may be sulphuric acid half esters of either alkanols (ie where n = 0) or ethoxylated alkanols (ie, where n = 1-40). Suitable sulphates are from the linear, branched, primary or secondary alcohols, with or without ethoxylation. Preferred alcohols are:

25

(i) the linear primary alcohols having 10-14 carbon atoms, which, if ethoxylated, have up to 12 mols ethylene oxide per mol of alcohol;

(ii) the mainly straight-chain primary alcohols having 10-15 carbon atoms and about 25% 2-methyl branching, which, if ethoxylated, have up to 12 mols ethylene oxide per mol of alcohol;

30

(iii) the random secondary alcohols having 11-15 carbon atoms and 7-30 mols ethylene oxide per mol of alcohol.

Primary alcohols with substantial branching may be used.

The preferred cations which neutralise the anionic surfactant are sodium and ammonium, and the latter sulphates can be an aid to hydrotropeing of the liquid compositions of the invention.

35

The sulphonates may be those conventionally used, including linear alkylbenzene sulphonate, secondary alkyl sulphonate and alpha olefin sulphonate.

Amine Oxide/Betaine


The third essential feature of the present invention is the presence of an amine oxide or a betaine as a lather booster.

It is preferred to avoid using substantial amounts of these, especially amine oxides, for the sake of economy and consequent cost effectiveness. Preferably then the amount of amine oxide is not more than 10% by weight of the active detergent mixture. Preferably the amount of betaine is not more than 30% by weight of this mixture. The total amount of amine oxide and betaine, and, if present, ethanolamide is from 12 to 30% by weight of the active detergent mixture. The lather booster is preferably present at from 2 to 6% by weight of the total composition. It is preferred to use betaines alone.

45

Suitable betaines include simple betaines of formula:


50




55

EP 0 341 071 B1

and amido betaines of formula:




In both formulae R is a C₈ to C₁₈ straight or branched alkyl group. It may be a lauryl group or a middle cut coconut alkyl group. R₆ and R₇ are each C₁ to C₃ alkyl or C₁ to C₃ hydroxyalkyl. Examples of sulphobetaines have the above formulae with -CH₂CO⁻₂ replaced by:



A suitable simple betaine is Empigen BB from Albright & Wilson. It has the formula quoted above in which R is C₁₂ to C₁₄ alkyl, derived from coconut, and R₆ and R₇ are both methyl. Also preferred is Tego L7 from Goldschmidt, which has a whole coconut alkyl group.

Suitable amine oxides have the formula:



wherein R is a straight- or branched-chain C₈ to C₁₈ alkyl group and R₆ and R₇ are each C₁ to C₃ alkyl, or C₁ to C₃ hydroxyalkyl. A suitable amine oxide is Empigen OB from Albright & Wilson. In it R is middle-cut coconut alkyl and R₆ and R₇ are both methyl.

Additional Ingredients

The compositions according to the present invention may include other conventional ingredients such as minor amounts of builders, urea, sodium or ammonium chloride, corrosion inhibitors, colourings and perfumes. Conventional foam boosters such as ethanolamides may be included.

The presence of ethanolamide is optional. If present, the total level of betaine and/or amine oxide and ethanolamide is from 12 to 30 wt% of the total active. A preferred ethanolamide is Empilan LME (Albright & Wilson).

The compositions according to the present invention are of particular use as light duty liquid detergent compositions for manual dishwashing.

The term "dishes" as used herein means any utensils involved in food preparation or consumption which may be required to be washed to free them from food particles and other food residues, greases, proteins, starches, gums dyes, oils and burnt organic residues.

Typical compositions for manual dishwashing comprise from about 5 to about 60 wt%, preferably about 10 to about 40 wt% of the mixture of active components hereinbefore described, the balance comprising water and solvents.

The compositions according to the present invention are mild and environmentally safe. They exhibit good foam characteristics.

The invention will now be described in more detail by the following Examples.

EXAMPLES

The foaming performance and mildness of various aqueous formulations were compared. Those formulations falling within the scope of the present application were compared with comparable compositions outside the scope.

Foaming performance was assessed by means of a modified Schlachter-Dierkes test based on the principle described in Fette und Seifen 1951, 53, 207. A 100 ml aqueous solution of each material tested, having a concentration of 0.04% active detergent in 24°H water (French hardness) at 45° C was rapidly oscillated using a vertically oscillating perforated disc within a graduated cylinder. After the initial generation of foam, increments (0.2 g) of soil (9.5 parts commercial cooking fat, 0.25 parts oleic acid, 0.25 parts stearic acid and 10 parts wheat starch in 120 parts water) were added at 15 second intervals (10 seconds' mild agitation and 5 seconds' rest) until the foam collapsed. The result was recorded as the number of soil increments (NSI score): a score difference of 6 or less is generally regarded as insignificant. Each result was typically the average of 3 or 4 runs.

Several in-vitro and in-vivo methods for evaluating protein denaturation potency of surfactants and their mixtures have been reported (see Miyazowa et al. Int J Cos Sci 6 33-46 1984, and the references cited therein). One such method is the study of interaction of detergents with acid phosphatase enzyme either from skin (Prottey et al, Int J Cos Sci 6 263-273 1984) or from Wheatgerm (Tanaka et al, Anal Biochem 66 489-498 1975).

In vivo mildness of formulations can be assessed using a flex wash test. In this test neat products were rubbed on the forearm of panellists and rinsed. The process was repeated four times a day for five days and the level of erythema developed was assessed by trained assessors.

A comparison of wheatgerm acid phosphatase (WGAP) test and flex wash test results indicated that formulations giving less than 50% enzyme inhibition under the test conditions are substantially mild; any mildness differences between products giving <40% inhibition do not show any detectable mildness differences in flex wash test, indicating that the enzyme test is very sensitive and that in a real life situation there is a threshold level of protein denaturation below which all actives and products are indistinguishably mild. The results of the WGAP test are expressed as percentage inhibition (ie, 100% minus percentage activity remaining). Water gave no inhibition at all, ie 100% of activity remained.

Materials Used

- APG : APG 500 or APG 550 from Horizon Chemicals. Both have C₁₂-C₁₄ alkyl groups. Average degree of polymerisation of 1.4 and 1.8 respectively.
- PAS : Dobanol 23-S (Shell)
- Betaine : Empigen BB (Albright & Wilson)
- Amine Oxide : Empigen OB (Albright & Wilson)
- LME : Emplilan LME (Albright & Wilson)

The varying amounts of ingredients used in the compositions tested are shown below as a percentage of the total weight of the composition.

5
10
15
20
25
30
35
40
45
50
55

<u>Example Number</u>	<u>Product</u>	<u>Plunger Test</u> 24°H, 45°C, 0.04% AD
1	19/6/4/5	40
2	22/6/6	35
3	22/6/6	37
4	18/8/8	37
5	18/8/8	39
6	22/8/6/4	39
8	9/9/8/8	35
9	18/8/8	29
A	22/12	15
B	18/8/8	31
C	22/8/4	29
D	22/12	27

The foam tests show that compositions falling within the scope of the invention perform better than those outside (A, B and D). Furthermore, APG 500 performs better than APG 550.
The following enzyme inhibition tests were performed;

	Product	% Enzyme Inhibition
19/6/4/5	Dobanol 91-8EO/PAS/Betaine/LME	4
19/6/4/5	APG 500/PAS/Betaine/LME	4

In the WGAP test, APG and nonionic formulations give similar low enzyme inhibition, and hence both have similar mildness.

Claims

1. An aqueous liquid detergent composition comprising:
 - (a) a C₈-C₁₈ alkyl polyglycoside surfactant having an average degree of polymerisation of from 1 to 3;
 - (b) an anionic surfactant;
 - (c) a surface-active betaine and/or amine oxide; and optionally;
 - (d) an ethanolamide;

wherein the amount of anionic surfactant is not greater than 1.5 times (on a molar basis) the level of betaine and/or amine oxide and wherein the level of betaine and/or amine oxide and, if present, ethanolamide is from 12 to 30 wt % of the total active.
2. A composition as claimed in claim 1, wherein the alkyl polyglycoside has an average degree of polymerisation of from 1 to 1.4.
3. A composition as claimed in claim 1 or claim 2, wherein the anionic surfactant is a primary alkyl sulphate.
4. A composition as claimed in any one of claims 1 to 3, wherein a betaine alone is used as a lather booster.
5. A composition according to any one of claims 1 to 4 wherein a nonionic active is also present.
6. An aqueous liquid detergent composition as claimed in claim 1 comprising, based on the total amount of active material:
 - (a) from 50 to 60 wt% of an alkyl polyglycoside having an average degree of polymerisation of from 1 to 1.4;
 - (b) from 15 to 20 wt% of a primary alkyl sulphate;
 - (c) from 10 to 15 wt% of a betaine;
 - (d) from 10 to 15% of an ethanolamide.

Patentansprüche

1. Wässrige flüssige Waschmittelzusammensetzung, umfassend:
 - (a) ein C₈-C₁₈-Alkylpolyglycosid-Tensid mit einem durchschnittlichen Polymerisationsgrad von 1 bis 3;
 - (b) ein anionisches Tensid;
 - (c) ein oberflächenaktives Betain und/oder Aminoxid; und gegebenenfalls
 - (d) ein Ethanolamid,

wobei die Menge an anionischem Tensid nicht größer als das 1,5-fache (auf molarer Basis) des Anteils an Betain und/oder Aminoxid ist und wobei der Anteil an Betain und/oder Aminoxid und, falls vorliegend, Ethanolamid 12 bis 30 Gew.-% des gesamten Aktivstoffs ausmacht.
2. Mittel nach Anspruch 1, wobei das Alkylpolyglycosid einen durchschnittlichen Polymerisationsgrad von 1 bis 1,4 aufweist.
3. Mittel nach Anspruch 1 oder 2, wobei das anionische Tensid ein primäres Alkylsulfat ist.
4. Mittel nach einem der Ansprüche 1 bis 3, wobei nur ein Betain als Schaumverstärker verwendet wird.

5. Mittel nach einem der Ansprüche 1 bis 4, wobei ebenfalls ein nichtionischer Aktivstoff vorliegt.
6. Wässrige flüssige Waschmittelzusammensetzung nach Anspruch 1, umfassend auf der Grundlage der Gesamtmenge an aktivem Stoff:
- 5 (a) 50 bis 60 Gew.-% eines Alkylpolyglycosids mit einem durchschnittlichen Polymerisationsgrad von 1 bis 1,4;
(b) 15 bis 20 Gew.-% eines primären Alkylsulfats;
(c) 10 bis 15 Gew.-% eines Betains;
(d) 10 bis 15 % eines Ethanolamids.

10

Revendications

1. Composition détergente liquide aqueuse comprenant :
- 15 (a) un alkylpolyglycoside tensioactif en C₈-C₁₈ présentant un degré moyen de polymérisation de 1 à 3 ;
(b) un tensioactif anionique ;
(c) une bétaine et/ou un oxyde d'amine tensioactif ; et, facultativement :
(d) un éthanolamide,
- 20 dans laquelle la quantité du tensioactif anionique n'est pas supérieure à 1,5 fois (sur une base molaire) le niveau de bétaine et/ou d'oxyde d'amine et dans laquelle le niveau de bétaine et/ou d'oxyde d'amine et, éventuellement, d'éthanolamide représente de 12 à 30% en poids du total des ingrédients actifs.
2. Composition selon la revendication 1, dans laquelle l'alkylpolyglycoside présente un degré moyen de polymérisation de 1 à 1,4.
- 25 3. Composition selon la revendication 1 ou 2, dans laquelle le tensioactif anionique est un alkylsulfate primaire.
4. Composition selon l'une quelconque des revendications 1 à 3, dans laquelle on utilise la seule bétaine comme agent de renforcement du moussage.
- 30 5. Composition selon l'une quelconque des revendications 1 à 4, qui contient également un tensioactif non ionique.
- 35 6. Composition détergente liquide aqueuse selon la revendication 1, qui comprend par rapport au total des matières actives :
- (a) de 50 à 60% en poids d'un alkylpolyglycoside ayant un degré moyen de polymérisation de 1 à 1,4 ;
(b) de 15 à 20% en poids d'un alkylsulfate primaire ;
40 (c) de 10 à 15% en poids d'une bétaine ;
(d) de 10 à 15% d'un éthanolamide.

45

50

55