

(72) **Inventeurs(suite)/Inventors(continued):** ECKSTEIN, HELLMUT, DE; KIRSCHBAUM, THOMAS, DE;
BOMMARIUS NEE RIEBEL, BETTINA, US

(57) **Abrégé(suite)/Abstract(continued):**

used. The invention additionally concerns a process for inserting corresponding nucleic acid sequences into a vector for expression in methylotrophic yeast strains and it concerns appropriate vectors and host strains.

Abstract

The invention concerns a process for the recombinant production and expression of eukaryotic alkaline phosphatase in yeast cells in which in particular a DNA coding for a eukaryotic highly active alkaline phosphatase having a specific activity above 3000 U/mg is used. The invention additionally concerns a process for inserting corresponding nucleic acid sequences into a vector for expression in methylotrophic yeast strains and it concerns appropriate vectors and host strains.

Roche Diagnostics GmbH

5387/00/

Expression of alkaline phosphatase in yeast

The invention concerns a process for the recombinant production and expression of eukaryotic alkaline phosphatase. The invention additionally concerns a codon-optimized DNA which codes for a eukaryotic highly active alkaline phosphatase having a specific activity of more than 3000 U/mg. Furthermore the invention concerns a process for inserting the DNA into a vector for expression in yeast cells.

Alkaline phosphatases (AP) are dimeric, zinc-containing, non-specific phosphomonoesterases which occur in prokaryotic as well as in eukaryotic organisms e.g. in *E. coli* and mammals (McComb et al., 1979 *Alkaline Phosphatases Plenum Press, New York*). Comparison of the primary structures of various alkaline phosphatases showed a high degree of homology (25-30 % homology between *E. coli* and mammalian AP; Millàn, 1988 *Anticancer Res.* 8, 995-1004; Harris, 1989 *Clin. Chim. Acta* 186, 133-150).

In humans and higher animals the AP family comprises four members that are located in different gene loci (Millan, 1988 *Anticancer Res.* 8, 995-1004; Harris 1989 *Clin. Chim. Acta* 186, 133-150). The alkaline phosphatase family includes the tissue-specific APs (placental AP (PLAP), germ cell AP (GCAP) and intestinal AP (IAP)) and the non-tissue specific APs (TnAP) which are primarily located in the liver, kidney and bones.

A decisive property of the previously known APs is the large variability of the catalytic activity of mammalian APs which have a 10-100-fold higher k_{cat} s value than *E. coli* AP. Among the mammalian APs, the APs from the bovine intestine (bIAP) exhibit the highest specific activities. This property makes the bIAPs attractive for biochemical applications such as the use of corresponding enzyme conjugates as a diagnostic reagent, or to dephosphorylate DNA. The existence of various alkaline phosphatases from the bovine intestine having specific activities of varying magnitudes is described in EP 0 955 369 and Manes et al. (1998), *J. Biol. Chem.* **273** No. 36, 23353-23360. Up to now recombinant expression of eukaryotic alkaline phosphatases of low activity (up to 3000 U/mg) has been described in various eukaryotic cell lines such as CHO cells (bIAP I/WO 93/18139; Weissig et al. 1993, *Biochem. J.* 260, 503-508), COS cells (human placental AP/Berger et al., 1987, *Biochemistry* **84**, 4885-4889) or baculovirus expression systems (human placental AP/Davis et al. 1992, *Biotechnology* **10**, 1149-1150). The expression of more active APs (specific activity > 3000 U/mg) from the bovine intestine in CHO cells has also been described (bIAP II, III and IV/Manes et al. 1998, *J. Biol. Chem.* **273** No. 36, 23353-23360). However, a disadvantage of expressing alkaline phosphatases in these expression systems is the low expression output which makes the recombinant production especially of a highly active AP uneconomic.

Although it is in principle possible to express eukaryotic alkaline phosphatases in prokaryotic expression hosts such as *E. coli* (human placental AP/Beck and Burtscher, 1994 *Protein Expression and Purification* **5**, 192-197), the alkaline phosphatases

expressed in prokaryotes are not glycosylated which is essential especially for preparing enzyme conjugates.

Consequently the object of the present invention is to develop a robust and stable expression process for the production of glycosylated eukaryotic alkaline phosphatase having a high specific activity, which, due to the high expression output, allows an economic production of such an alkaline phosphatase and, moreover, yields an enzyme whose properties are comparable to native alkaline phosphatase of high activity or low activity (commercially available for example from Roche Diagnostics GmbH, Biozyme, Oriental Yeast) with regard to for example specific activity and thermostability.

The object is achieved according to the invention by a process for the production of a eukaryotic alkaline phosphatase having a high specific activity in yeast and especially in a methylotrophic yeast comprising the steps:

- a) cloning a gene sequence into different vectors
- b) transformation of the yeast,
- c) expression and
- d) purification of the alkaline phosphatase,
characterized in that
 - (i) a first vector has a resistance gene for a first selection marker
 - (ii) transformants which have integrated the resistance gene and the gene sequence into the genome are selected by growth on nutrient medium containing a low concentration of a first selection marker,

- (iii) gene copy number is increased by multiple transformation in which multiple transformants are selected by growth on a nutrient medium at an increased selection pressure,
- (iv) a second vector is added which has a resistance gene for a second selection marker in addition to the gene sequence,
- (v) the gene copy number is increased by multiple transformation with the second vector in which multiple transformants are selected by growth on nutrient medium at an increased selection pressure and
- (vi) the clones are selected which have integrated several copies of the gene sequence and the selection marker resistance genes into the genome in a stable manner.

A preferred gene sequence is a DNA sequence which codes for a eukaryotic alkaline phosphatase that has a specific activity of more than 3000 U/mg and in special cases of more than 7000 U/mg to about 10,000 U/mg. For example a DNA sequence according to SEQ ID NO:1 has proven to be suitable according to the invention. A codon-optimized DNA sequence which corresponds to the gene sequence SEQ ID NO:1 at the amino acid level is particularly preferred. Codon-optimization means that each codon for example of SEQ ID NO:1 has been optimized by silent mutations i.e. changes at the DNA level which, however, have no effect at the amino acid level in order to increase the translation according to the requirements of the selected expression host which results for example in the gene sequence according to SEQ ID NO:5. It is, however, also possible to incorporate other sequences than SEQ ID NO:1 into the vector which code for alkaline phosphatases and are optionally codon-optimized such as bIAP1, III, IV (DE

198 19 962 and EP 0 955 369). It is particularly preferable for the process according to the invention to use a codon-optimized gene sequence according to SEQ ID NO:5. The corresponding gene sequence is then cloned into one or several vector(s) which is or are selected depending on the host to be transformed.

Methylotrophic yeasts e.g. *Pichia pastoris* yeast, *Hansenula polymorpha* and also other yeasts such as *Saccharomyces cerevisiae*, *Yarrowia lipolytica* or *Schizosaccharomyces pombe* are particularly suitable as the yeast host. Suitable vectors are known to a person skilled in the art such as pPICZαA, pIIC9K, Yes vectors, pTEF1/Zeo, pYDI (e.g. Invitrogen). The expression vector that is formed in this manner is preferably transformed into various strains of *Pichia pastoris* and integrated into the genome in a stable manner. Stable integration into the yeast genome has the advantage that selection pressure is not required during the subsequent production of the, for example, eukaryotic, highly active alkaline phosphatase in large volume ferments. Stable integration into the genome means that the expression vector is incorporated into the genome of for example *Pichia pastoris* by means of homologous recombination and is thus transmitted by heredity as a permanent component from generation to generation (Cregg, J.M. et al., Mol. Cell. Biol. 5 (1985), 3376-3385).

The gene copy number was increased in the methylotrophic yeast by multiple transformation while at the same time increasing the selection pressure with a suitable selection marker e.g. an antibiotic such as Zeocin[®] or Geneticin (G418) or an auxotrophy marker after which only those clones can survive which have integrated

several copies of the expression vector into the genome in a stable manner. In order to be resistant to higher concentrations of the antibiotic used as the selection marker, it is necessary that the clones produce an increasing amount of resistance protein. This can for example be achieved by multiple integration of the expression vector which contains the resistance gene for the antibiotic used as the selection marker in addition to the expression cassette for the highly active alkaline phosphatase for example.

The object of producing eukaryotic alkaline phosphatase economically in a robust and stable expression process with a high expression output could not be achieved until measures (i) to (vi) were combined. Thus for example transformation of a *Pichia pastoris* strain X-33 with an expression vector which contains the bIAP^{II} gene according to SEQ ID NO:1 did not lead to the desired result without these measures (see examples 1 and 2). Although the process enabled a considerable increase in the expression output compared to expression of bIAP^{II} in CHO cells (Manes et al., 1998, J. Biol. Chem. **273** No.36, 23353-23360) the process does not allow a recombinant alkaline phosphatase to be produced economically.

One of the necessary measures for the process according to the invention is the synthesis of a codon-optimized gene sequence. A complete de novo synthesis of the ca. 1,5 kbp long gene which codes for the eukaryotic highly active alkaline phosphatase was necessary in order to optimize each codon for expression in yeast. It was possible to optimize each codon, as required, by retranslation of the amino acid sequence of the eukaryotic highly active alkaline phosphatase according

to SEQ ID NO:4 (bIAP-II) and by utilizing the degenerate code. For this purpose the gene was divided into 28 oligonucleotides having a length of 54 to 82 nucleotides. The oligonucleotides were designed as an alternating sequence of sense strand and antistrand fragments the 5' and 3' ends of which each overlapped in a complementary manner with the neighbouring oligonucleotides. The overlapping region was in each case selected in such a manner that unspecific binding was largely prevented during the annealing reaction in the subsequent PCR reaction. The oligonucleotides at the 5' and 3' ends of the gene were provided with recognition sites for restriction endonucleases upstream and downstream of the coding region which could be used for a later insertion of the synthetic gene according to SEQ ID NO:5 into expression vectors. Hence a recognition site for the restriction endonuclease EcoRI was incorporated upstream and a recognition site for the restriction endonuclease Asp718 was incorporated downstream. The sequences of the oligonucleotides are shown in SEQ ID NO:6 to 33.

The gene synthesis was carried out by means of a PCR reaction. For this purpose the coding region was firstly divided into three segments (oligonucleotides 6 to 15, 16 to 23, 24 to 33) and these segments were generated in separate PCR reactions. During the gene synthesis by means of PCR reaction using overlapping complementary oligonucleotides the gene fragment is elongated stepwise to form the full length product which is then amplified in subsequent cycles. The annealing temperature in this process depends on the overlapping region having the lowest melting temperature.

The three segments were subsequently analysed by agarose

gel electrophoresis, the products having the expected length were isolated from the gel by means of the QIAquickTM gel extraction kit (Qiagen) and synthesized in a further PCR reaction to form the complete gene product. In this process the PCR reaction was carried out in the first 5 cycles without adding the primers at the 5' end and at the 3' end of the total gene so that only a few fragments of the gene product having the expected length were initially formed from the three segments. The annealing temperature depends on the overlapping region having the lowest melting temperature. Subsequently the terminal primers were added and the annealing temperature was increased to correspond with the annealing temperature of the primer having the lowest melting temperature. The gene fragment having the expected length was amplified to a high degree in a further 25 cycles.

The PCR mixture was analysed by agarose gel electrophoresis and the gene fragment having the expected size was isolated (QIAquickTM gel extraction kit/Qiagen).

The cloning of such a PCR fragment, transformation in *Pichia pastoris* and the expression is described in example 3.

The codon-optimized gene for the highly active alkaline phosphatase enabled the expression output to be increased three-fold compared to the first experiments with the wild-type gene.

However, these clones did not provide an economic process for producing the highly active alkaline

phosphatase.

One measure which can increase the expression output of heterologous and homologous proteins in *Pichia pastoris* is to increase the gene copy number in the cell by multiple transformation. This measure can increase the transcription product i.e. the mRNA of the target gene. The gene copy number is increased by multiple transformation of a clone containing the expression vector while simultaneously increasing the selection pressure during the subsequent growth of the transformants on nutrient plates containing increased concentrations of the antibiotic used as the selection marker. In this process an expression clone which has already taken up at least one copy of the expression vector from the first transformation cycle is again made competent (see example 1) and is again transformed with the expression vector. Transformants are selected which have integrated several copies of the expression vector into the genome by plating out on nutrient plates with a higher selection pressure i.e. plates containing a higher concentration of the antibiotic (e.g. Zeocin[®]) used as the selection marker than during the first transformation cycle. For this the highest concentration of the antibiotic used as the selection marker at which the clone from the first transformation cycle can still grow is determined and the concentration of the antibiotic used as the selection marker is increased accordingly above the determined threshold value in the YPDS agar plates after the additional transformation. Increasing the copy number of the expression vector also increases the copy number of the resistance gene which is a component of the expression vector and hence also increases resistance to higher concentrations of the antibiotic used as the selection marker. It is also

possible to select clones containing different copy numbers of the expression vector in the genome by varying the concentration of the antibiotic used as the selection marker in the nutrient plates (ca. 100 to 2000 $\mu\text{g/ml}$ / see example 4).

A further measure which can be used to increase the expression output of heterologous and homologous proteins in yeast such as *Pichia pastoris* is to increase the gene copy number by multiple selection. In order to achieve this an expression clone that has been already optimized by multiple transformation with an expression vector which contains the expression cassette containing the target gene (e.g. the gene which codes for the highly active alkaline phosphatase according to SEQ ID NO:5) and a resistance gene for the first antibiotic used as the selection marker (e.g. Zeocin[®]) is transformed with a second expression vector which contains the target gene (e.g. the gene which codes for the highly active alkaline phosphatase according to SEQ ID NO:5) and a resistance gene for the second antibiotic (e.g. Geneticin (G418)) used as the selection marker. When the transformants are subsequently plated out on nutrient plates which contain the second antibiotic as the selection marker, only those clones are selected which have also taken up at least one copy of the expression vector containing the resistance gene for the second antibiotic used as the selection marker in addition to copies of the expression vector containing the resistance gene for the first antibiotic used as the selection marker. These expression clones can now be in turn subjected to a further multiple transformation with the expression vector containing the resistance gene for the second antibiotic used as the selection marker (see example 5).

By combining the measures of multiple transformation and double selection it was possible to increase the expression output four-fold compared to the expression output of clones from the first transformation cycle containing the codon-optimized gene.

The recombinant alkaline phosphatase can be extracted from the biomass by extraction methods that are in principle known to a person skilled in the art e.g. Protein Purification, Springer Verlag, editor Robert Scopes (1982). A pure band product having a specific activity of more than 7000 U/mg is obtained by chromatographic separation methods and especially those using hydrophobic column materials and a cation exchanger.

The purified product was subjected to N-terminal sequencing in order to characterize the recombinant highly active alkaline phosphatase.

The dominant sequence EAEAEFLIPA (SEQ ID NO:36) was determined. The sequence unequivocally correlates with the N-terminal sequence of the AP"LIPA" (SEQ ID NO:37) and with the linker peptide of the construct EAEAEF (SEQ ID NO:38) which is formed by the strategy of cloning the gene sequence into the vector and by cleavage of the α -factor signal peptide by a Kex2 signal peptidase (e.g. Invitrogen).

The stability of the recombinant alkaline phosphatase product was examined in comparison with the naturally occurring alkaline phosphatase. The samples yielded comparable results when the solutions were subjected to a thermal stress (55°C).

Hence the present invention describes for the first time a process which enables an economic production of a recombinant alkaline phosphatase from mammalian cells such as bovine intestine which has properties that are comparable to the native highly active alkaline phosphatase from bovine intestine and is glycosylated.

The present invention also concerns a DNA sequence according to SEQ ID NO:5 as a codon-optimized gene sequence for the expression of the gene for the highly active alkaline phosphatase in *Pichia pastoris*.

A further subject matter of the invention is a vector containing SEQ ID NO:5 and particularly preferably the vector pHAP10-3 according to fig. 2. pHAP10-3 is the vector pPICZ α A which is commercially available (Invitrogen) and contains the inventive gene according to SEQ ID NO:5 which is under the control of the AOX 1 promoter.

A further subject matter of the invention is a host strain which has been transformed with the vectors according to the invention. The *Pichia pastoris* X-33 strain transformed with the vector pHAP10-3 is particularly preferred.

Another preferred vector is a vector which contains the entire expression cassette from pHAP10-3 which essentially comprises the AOX 1 promoter, the signal peptide of the α -factor from *Saccharomyces cerevisiae* which is cloned in the correct reading frame behind the signal peptide, the codon-optimized target gene according to SEQ ID NO:5 which codes for the highly active alkaline phosphatase and the AOX 1 transcription

termination region (see fig.3). The vector pHAP 10-3/9K is particularly preferred which comprises the commercially available vector pPIC9K (Invitrogen) and the expression cassette from pHAP 10-3 including the synthetic gene according to SEQ ID NO:5.

The vectors pHAP 10-3 and pHAP 10-3/9K are equally relevant since the final production clone contains copies of both vectors.

A further subject matter of the invention is a host strain which has been transformed with the pHAP10-3/9K vector. However, other vectors and strains known to a person skilled in the art are also suitable in the sense of the present invention such as YES vectors, pYD1, pTEF1/ZEO (Invitrogen) and *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Hansenula polymorpha*, *Yarrowia lipolytica* and in particular *Pichia pastoris* X-33. The *Pichia pastoris* X-33 strain transformed with the vector pHAP10-3/9K is especially preferable for the invention.

Hence a further subject matter of the invention is a process for producing a eukaryotic highly active alkaline phosphatase by expressing the protein in a host strain which has been transformed with one or several vectors according to the invention and especially with the pHAP 10-3 vector or the pHAP 10-3/9K vector. *Pichia pastoris* strains which have been transformed with the inventive vectors are particularly preferred for the inventive process. The strain *Pichia pastoris* X-33 which has been transformed with a pHAP 10-3 and a pHAP 10-3/9K vector is especially preferred in this connection.

Figures

Figure 1

Plasmid map of the expression vector pHAP-1 containing the bIAP^{II} gene in pICZ α A (Invitrogen).

Figure 2

Plasmid map of the expression vector pHAP 10-3 containing the synthetic gene in pPIC9K (Invitrogen).

Figure 3

Plasmid map of the expression vector pHAP 10-3/9K containing the synthetic gene in pPIC9K (Invitrogen).

Abbreviations

YPD: yeast peptone dextrose
YPDS: yeast peptone dextrose sorbitol
BMGY: buffered glycerol complex medium
BMMY: buffered methanol complex medium

Example 1:Cloning the bIAP^{II} gene

The bIAP^{II} gene according to SEQ ID NO:1 (EP 0 955 369; Manes et al., 1998, *J. Biol. Chem.* **273** No. 36, 23353-23360) was firstly provided upstream and downstream with restriction endonuclease cleavage sites suitable for cloning into expression vectors for *Pichia pastoris* by means of PCR and selection of suitable primers according to SEQ ID NO:2 and 3. Hence the restriction endonuclease cleavage site for EcoRI was attached upstream and the restriction endonuclease cleavage site for Asp718 I was attached downstream.

The PCR fragment was recleaved with EcoRI and Asp718 I (Roche Diagnostics GmbH), isolated again (QIAquickTM gel extraction kit/Qiagen) and subsequently ligated into a vector fragment of the expression vector pPICZ α A (Invitrogen) that had been linearized with EcoRI and Asp718 I (Roche Diagnostics GmbH) and isolated (QIAquickTM gel extraction kit/Qiagen). In this vector the bIAP^{II} gene is under the control of the AOX 1 promoter (promoter for alcohol oxidase 1 from *Pichia pastoris* and inducible with methanol) and is cloned in the correct reading frame behind the signal peptide of the α -factor from *Saccharomyces cerevisiae*. It was then examined whether the gene fragment inserted in this manner was free of errors by means of restriction analysis and sequencing. The expression vector formed in this manner which contains the bIAP^{II} gene which codes for the eukaryotic highly active alkaline phosphatase was named pHAP-1 (see fig. 1).

Transformation of pHAP-1 in Pichia pastoris

For the transformation of pHAP-1 in *Pichia pastoris* X-33 and subsequent integration into the genome, the vector was firstly linearized with SacI (Roche Diagnostics GmbH). The transformation was carried out by means of electroporation using a Gene Pulser II (Biorad).

For this 5 ml YPD medium (Invitrogen) was inoculated with a colony of *Pichia pastoris* wild-type strain and incubated at 30°C overnight while shaking. 200 ml fresh YPD medium (Invitrogen) was subsequently transfer inoculated 1:2000 with the overnight culture and incubated overnight at 30°C while shaking until the OD₆₀₀ reached 1.3 - 1.5. The cells were centrifuged (1500 x g/5 minutes) and the pellet was resuspended in 200 ml ice-cold sterile water (0°C). The cells were again centrifuged (1500 x g/5 minutes) and resuspended in 100 ml ice-cold, sterile water (0°C). The cells were again centrifuged and resuspended in 10 ml ice-cold (0°C) 1 M sorbitol (ICN). The cells were again centrifuged and resuspended in 0.5 ml ice-cold (0°C) 1 M sorbitol (ICN). The cells obtained in this manner were kept on ice and used immediately for transformation.

About 1 µg linearized pHAP-1 vector DNA was added to 80 µl of the cells and the entire mixture was transferred to an ice-cold (0°C) electroporation cuvette and incubated for a further 5 minutes on ice. Subsequently the cuvette was transferred to the Gene Pulser IITM (Biorad) and the transformation was carried out at 1 kV, 1 kΩ and 25 µF. After electroporation, 1 ml 1 M sorbitol (ICN) was added to the mixture and subsequently 100 to 150 µl was plated out on a YPDS agar plate (Invitrogen) containing 100 µg/ml Zeocin[®] (Invitrogen). The plates were subsequently incubated at 30°C for 2-4 days.

Raster MD (=minimal dextrose) plates were inoculated with the clones and they were analysed further. Growing clones were picked out, resuspended in 20 μ l sterile water, lysed with 17.5 U lyticase (Roche Diagnostics GmbH) (1 hour, 37°C) and examined directly for the correct integration of the bIAPII expression cassette by means of PCR.

Clones which had integrated the complete expression cassette during transformation into the genome were then used in expression experiments.

Expression of the highly active alkaline phosphatase


3 ml BMGY medium (Invitrogen) was inoculated with positive clones and incubated overnight at 30°C while shaking. Subsequently the OD was determined at 600 nm and 10 ml BMMY medium (Invitrogen) was transfer inoculated in such a manner that an OD₆₀₀ of 1 was obtained. The BMMY medium (Invitrogen) contains methanol (Mallinckrodt Baker B.V.) which induces the expression of the highly active alkaline phosphatase via the AOX 1 promoter.

The shaking flasks were incubated at 30°C while shaking, samples were taken every 24 hours, the OD₆₀₀ was determined, an activity test for the expression of the highly active alkaline phosphatase was carried out and 0.5 % methanol (Mallinckrodt Baker B.V.) was added for the further induction. The expression experiments were carried out for 96 hours.

Example 2:Activity test for the highly active alkaline phosphatase

500 μ l of the expression culture of example 1 was removed, the OD₆₀₀ was determined and the cells were centrifuged. The supernatant was stored and the cell pellet was resuspended for the lysis in an amount of Y-PERTM (50 to 300 μ l/Pierce) corresponding to the OD₆₀₀ and shaken for 1 hour at room temperature. Subsequently the lysate was centrifuged to remove cell debris (15000 x g/5 minutes) and the supernatant was transferred to fresh reaction vessels. 5 μ l of the lysate was then used in the activity test.

The principle of the activity test is as follows:



The increase in absorbance at 405 nm is measured.

50 μ l 4-nitrophenyl phosphate solution (0.67 mol/l 4-nitrophenyl phosphate, Na salt (Roche Diagnostics GmbH)) was added to 3 ml diethanolamine buffer (1 mol/l diethanolamine (Merck) pH 9.8, 0.5 mmol/l MgCl₂ (Riedel de Haen)) and the mixture was incubated at 37°C. Subsequently the reaction was started by adding 5 μ l lysate and the change in absorbance at 37°C was determined for 3 minutes and from this the $\Delta A/\text{min}$ was calculated.

The activity was then calculated according to the following formula:

$$\text{activity} = \frac{3.10}{\epsilon \times 0.005 \times l} \times \Delta A / \text{min} \times \frac{1}{\text{factor} \times} [\text{U/ml sample solution}]$$

$$\epsilon = 18.2 [1 \times \text{mmol}^{-1} \times \text{cm}^{-1}]$$

factor x = concentration factor after cell lysis

The activity of the medium supernatant of the expression cultures was determined in a similar manner. The reaction in this case was also started with 5 μ l of the supernatant but 0.5 mM ZnCl_2 was added additionally. In this case the calculation was carried out without factor x.

Example 3:

Cloning the PCR fragment from the gene synthesis

The PCR fragment was recleaved with EcoRI and Asp718 (Roche Diagnostics GmbH), isolated again (QIAquickTM gel extraction kit/Qiagen) and subsequently ligated into a vector fragment of the expression vector pPICZ α A (Invitrogen) that had been linearized with EcoRI and Asp718 (Roche Diagnostics GmbH) and isolated (QIAquickTM gel extraction kit/Qiagen). In this vector the synthetic gene is under the control of the AOX 1 promoter (promoter for alcohol oxidase 1 from *Pichia pastoris*, inducible with methanol (Mallinckrodt Baker B.V.) and is cloned in the correct reading frame behind the signal peptide of the α -factor from *Saccharomyces cerevisiae*. It was then examined whether the gene fragment inserted in this manner was free of errors by means of restriction analysis and sequencing. The expression vector formed in this manner which contains a synthetic

gene which codes for the eukaryotic highly active alkaline phosphatase was named pHAP10-3 (see fig. 2).

Transformation of pHAP10-3 in *Pichia pastoris*

For the transformation of pHAP10-3 in *Pichia pastoris* X-33 and subsequent integration into the genome, the vector was firstly linearized with SacI (Roche Diagnostics GmbH). The transformation was carried out by means of electroporation using a Gene Pulser II (Biorad). For this 5 ml YPD medium (Invitrogen) was inoculated with a colony of *Pichia pastoris* and incubated at 30°C overnight while shaking. 200 ml fresh YPD medium (Invitrogen) was subsequently transfer inoculated 1:2000 with the overnight culture and incubated overnight at 30°C while shaking until the OD₆₀₀ reached 1.3 - 1.5. The cells were centrifuged (1500 x g/5 minutes) and the pellet was resuspended in 200 ml ice-cold sterile water (0°C). The cells were again centrifuged (1500 x g/5 minutes) and resuspended in 100 ml ice-cold, sterile water (0°C). The cells were again centrifuged and resuspended in 10 ml ice-cold (0°C) 1 M sorbitol (ICN). The cells were again centrifuged and resuspended in 0.5 ml ice-cold (0°C) 1 M sorbitol (ICN). The cells obtained in this manner were kept on ice and used immediately for transformation.

About 1 µg linearized pHAP10-3 vector DNA was added to 80 µl of the cells and the entire mixture was transferred to an ice-cold (0°C) electroporation cuvette and incubated for a further 5 minutes on ice.

Subsequently the cuvette was transferred to the Gene Pulser II (Biorad) and the transformation was carried out at 1 kV, 1 kΩ and 25 µF. After electroporation 1 ml 1 M sorbitol (ICN) was added to the mixture and subsequently 100 to 150 µl was plated out on a YPDS agar

plate (Invitrogen) containing 100 $\mu\text{g/ml}$ Zeocin[®] (Invitrogen). The plates were subsequently incubated at 30°C for 2-4 days.

Raster MD (=minimal dextrose) plates were inoculated with the clones and they were analysed further. Growing clones were picked out, resuspended in 20 μl sterile water, lysed with 17.5 U lyticase (Roche Diagnostics GmbH) (1 hour, 37°C) and examined directly for the correct integration of the synthetic AP expression cassette by means of PCR.

Clones which had integrated the complete expression cassette during transformation into the genome were then used in expression experiments.

Expression of the highly active alkaline phosphatase

3 ml BMGY medium (Invitrogen) was inoculated with positive clones and incubated overnight at 30°C while shaking. Subsequently the OD was determined at 600 nm and 10 ml BMMY medium (Invitrogen) was transfer inoculated in such a manner that an OD₆₀₀ of 1 was obtained. The BMMY medium (Invitrogen) contains methanol (Mallinckrodt Baker B.V.) which induces the expression of the highly active alkaline phosphatase via the AOX 1 promoter.

The shaking flasks were incubated at 30°C while shaking, samples were taken every 24 hours, the OD₆₀₀ was determined, an activity test for the expression of the highly active alkaline phosphatase was carried out and 0.5 % methanol (Mallinckrodt Baker B.V.) was added for the further induction. The expression experiments were carried out for 96 hours.

Activity test for the highly active alkaline phosphatase

500 μ l of the expression culture was removed, the OD₆₀₀ was determined and the cells were centrifuged. The supernatant was stored and the cell pellet was resuspended for lysis in an amount of Y-PER™ (50 to 300 μ l/Pierce) corresponding to the OD₆₀₀ and shaken for 1 hour at room temperature. Subsequently the lysate was centrifuged to remove cell debris (15000 x g/5 minutes) and the supernatant was transferred to fresh reaction vessels. 5 μ l of the lysate was then used in the activity test.

The activity test was carried out as described above.

Example 4:Increasing the expression output by multiple transformation

The best clones from the expression experiments were in turn prepared for extrapolation as described above and again transformed with 1 μ g linearized pHAP10-3 vector DNA and the transformation mixture was plated out on YPDS agar plates (Invitrogen) containing 1000 to 2000 μ g/ml Zeocin® (Invitrogen). As a result the selection pressure was increased to such an extent that only clones could grow that had integrated several copies of the expression vector pHAP10-3 and thus also several copies of the respective resistance gene (in this case Zeocin®) into the genome. The Zeocin® resistance protein is the product of the bleomycin gene of *Streptoalloteichus hindustanus* (Chalmers, T. et al., Curr. Genet. 20 (1991), 309-314; Drocourt, D. et al., Nucleic Acid Research 18 (1990), 4009) which binds Zeocin® in a stoichiometric concentration ratio and thus makes the cell resistant to Zeocin®. The higher the concentration of Zeocin® in the

YPDS agar plates, the more resistance protein the cell has to generate in order to quantitatively bind the Zeocin® and thus enable growth. This is possible for example when multiple copies of the resistance gene have been integrated into the genome. As described above raster MD plates were transfer inoculated with the clones and they were again checked as described above by PCR analysis for the correct integration of the haAP expression cassette. Subsequently these clones were in turn tested for haAP activity as described above.

Example 5:

Increasing the expression output by using a second selection pressure

Increasing the Zeocin® concentration above 2000 µg/ml did not lead to an improvement in the expression output of the highly active alkaline phosphatase. In order to further increase the gene copy number in the expression clones of the gene according to SEQ ID NO:5 which codes for the highly active alkaline phosphatase and which is codon-optimized for expression in yeast, the integration of additional expression vectors into the genome of the expression clone derived from examples 3 and 4 that had the highest expression output was selected by means of a second selection pressure, preferably G418 (Roche Diagnostics GmbH). For this purpose the entire expression cassette from pHAP10-3 comprising AOX1 promoter, signal peptide of the α -factor from *Saccharomyces cerevisiae*, codon-optimized gene for the highly active alkaline phosphatase according to SEQ ID NO:5 and AOX 1 transcription termination region isolated by means of PCR using appropriately selected primers as described below, was cloned into the vector pIC9K the integration of which into the genome of *Pichia pastoris*

was selected by means of G418 (Roche Diagnostics GmbH). The primers are shown in SEQ ID NO:34 and 35.

The PCR mixture was analysed by agarose gel electrophoresis, the gene fragment having the expected size was isolated (QIAquickTM gel extraction kit/Qiagen), recleaved with SacI and NotI (Roche Diagnostics GmbH), subsequently isolated again from the agarose gel (QIAquickTM gel extraction kit/Qiagen) and ligated into an isolated vector fragment from pPIC9K that had also been linearized with SacI/NotI (Roche Diagnostics GmbH). This ensures that the entire expression cassette from pHAP10-3 is present in an identical form in pPIC9K. The inserted fragment was checked by means of restriction analysis and sequencing with the flanking regions. The expression vector formed in this manner was named pHAP10-3/9K (see fig.3).

The clones with the highest haAP expression output from the multiple transformation with pHAP10-3 (Zeocin resistance) were prepared for electroporation as described above and transformed as described above with 1 μ g of the vector fragment from pHAP10-3/9K linearized with SacI (Roche Diagnostics GmbH). The transformation mixture was subsequently stored for 1 to 3 days at 4°C in 1 M sorbitol (ICN) (to form the G418 resistance) and 100 to 200 μ l was plated out on YPD plates (Invitrogen) containing 1, 2 and 4 mg/ml G418 (Roche Diagnostics GmbH) and incubated for 3 to 5 days at 30°C. The resulting clones were again examined by means of the activity test for an increased expression of the eukaryotic highly active alkaline phosphatase as described above.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: Roche Diagnostics GmbH
(B) STREET: Sandhofer Strasse 116
(C) CITY: D-68305
(D) STATE/PROVINCE: Mannheim
(E) COUNTRY: Germany
(F) POSTAL CODE/ZIP:
(G) TELEPHONE: (617) 667-8000
(I) TELEFAX: (617) 632-7098

(ii) TITLE OF INVENTION: Expression of alkaline phosphatase in yeast

(iii) NUMBER OF SEQUENCES: 38

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Borden Ladner Gervais LLP
(B) STREET: 60 Queen Street
(C) CITY: Ottawa
(D) PROVINCE: Ontario
(E) COUNTRY: CANADA
(F) POSTAL CODE: K1P 5Y7

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Ver. 2.1

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: 2,352,609
(B) FILING DATE: 23-JUL-2001

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: DE 100 36 491.8
(B) FILING DATE: 25-JUL-2000

(viii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Fritz, Joachim T.
- (B) REGISTRATION NUMBER: 4173
- (C) REFERENCE/DOCKET NUMBER: PAT 49675-1

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: (613) 237-5160
- (B) TELEFAX: (613) 787-3558

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1476 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Bovine
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

gaattcctca tcccagctga ggaggaaaac cccgccttct ggaaccgcca ggcagcccag 60
gcccttgatg tagccaagaa gttgcagccg atccagacag ctgccaagaa tgtcatcctc 120
ttcttggggg atgggatggg ggtgcctacg gtgacagcca ctcgatcct aaaggggcag 180
atgaatggca aactgggacc tgagacaccc ctggccatgg accagttccc atacgtggct 240
ctgtccaaga catacaacgt ggacagacag gtgccagaca ggcagggcac tgccactgcc 300
tacctgtgtg ggggtcaagg caactacaga accatcggtg taagtgcagc cgcccgtctac 360
aatcagtgca acacgacacg tgggaatgag gtcacgtctg tgatcaaccg ggccaagaaa 420
gcaggggaagg ccgtgggagt ggtgaccacc accaggggtgc agcatgcctc cccagccggg 480
gcctacgcgc acacggtgaa ccgaaactgg tactcagacg ccgacctgcc tgctgatgca 540
cagaagaatg gctgccagga catcgccgca cagctgggtct acaacatgga tattgacgtg 600

27

atcctgggtg gaggccgaat gtacatgttt cctgagggga cccagaccc tgaataacca 660
 gatgatgcca gtgtgaatgg agtccggaag gacaagcaga acctgggtgca ggaatggcag 720
 gccaagcacc agggagccca gtatgtgtgg aaccgcactg cgctccttca ggcggccgat 780
 gactccagtg taacacacct catgggcctc tttgagccgg cagacatgaa gtataatgtt 840
 cagcaagacc acaccaagga cccgaccctg gcggagatga cggaggcggc cctgcaagtg 900
 ctgagcagga acccccgggg cttctacctc ttctgtggagg gaggccgcat tgaccacggt 960
 caccatgacg gcaaagctta tatggcactg actgaggcga tcatgtttga caatgccatc 1020
 gccaaggcta acgagctcac tagcgaactg gacacgctga tccttggtcac tgcagaccac 1080
 tcccatgtct tctcttttgg tggctacaca ctgcgtggga cctccatttt cggctctggcc 1140
 cccggcaagg ccttagacag caagtcctac acctccatcc tctatggcaa tggcccaggc 1200
 tatgcgcttg gcgggggctc gaggcccgat gttaatggca gcacaagcga ggaaccctca 1260
 taccggcagc aggcggccgt gcccctggct agcgagaccc acgggggcca agacgtggcg 1320
 gtgttcgcgc gaggcccga ggcgcacctg gtgcacggcg tgcaggagga gaccttcgtg 1380
 gcgcacatca tggcctttgc gggctgcgtg gagccctaca ccgactgcaa tctgccagcc 1440
 cccgccaccg ccaccagcat ccccgactag ggtacc 1476

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 40 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

gcgcgaattc ctcatcccag ctgaggagga aaaccccgcc

40

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

cgcggtacc ctagtcgggg atgctggtgg cggtgg

36

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 487 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: PRT

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Leu	Ile	Pro	Ala	Glu	Glu	Glu	Asn	Pro	Ala	Phe	Trp	Asn	Arg	Gln	Ala	
1				5					10					15		
Ala	Gln	Ala	Leu	Asp	Val	Ala	Lys	Lys	Leu	Gln	Pro	Ile	Gln	Thr	Ala	
			20					25					30			
Ala	Lys	Asn	Val	Ile	Leu	Phe	Leu	Gly	Asp	Gly	Met	Gly	Val	Pro	Thr	
		35					40				45					
Val	Thr	Ala	Thr	Arg	Ile	Leu	Lys	Gly	Gln	Met	Asn	Gly	Lys	Leu	Gly	
	50					55					60					

29

Pro	Glu	Thr	Pro	Leu	Ala	Met	Asp	Gln	Phe	Pro	Tyr	Val	Ala	Leu	Ser	65	70	75	80
Lys	Thr	Tyr	Asn	Val	Asp	Arg	Gln	Val	Pro	Asp	Ser	Ala	Gly	Thr	Ala	85	90	95	
Thr	Ala	Tyr	Leu	Cys	Gly	Val	Lys	Gly	Asn	Tyr	Arg	Thr	Ile	Gly	Val	100	105	110	
Ser	Ala	Ala	Ala	Arg	Tyr	Asn	Gln	Cys	Asn	Thr	Thr	Arg	Gly	Asn	Glu	115	120	125	
Val	Thr	Ser	Val	Ile	Asn	Arg	Ala	Lys	Lys	Ala	Gly	Lys	Ala	Val	Gly	130	135	140	
Val	Val	Thr	Thr	Thr	Arg	Val	Gln	His	Ala	Ser	Pro	Ala	Gly	Ala	Tyr	145	150	155	160
Ala	His	Thr	Val	Asn	Arg	Asn	Trp	Tyr	Ser	Asp	Ala	Asp	Leu	Pro	Ala	165	170	175	
Asp	Ala	Gln	Lys	Asn	Gly	Cys	Gln	Asp	Ile	Ala	Ala	Gln	Leu	Val	Tyr	180	185	190	
Asn	Met	Asp	Ile	Asp	Val	Ile	Leu	Gly	Gly	Gly	Arg	Met	Tyr	Met	Phe	195	200	205	
Pro	Glu	Gly	Thr	Pro	Asp	Pro	Glu	Tyr	Pro	Asp	Asp	Ala	Ser	Val	Asn	210	215	220	
Gly	Val	Arg	Lys	Asp	Lys	Gln	Asn	Leu	Val	Gln	Glu	Trp	Gln	Ala	Lys	225	230	235	240
His	Gln	Gly	Ala	Gln	Tyr	Val	Trp	Asn	Arg	Thr	Ala	Leu	Leu	Gln	Ala	245	250	255	
Ala	Asp	Asp	Ser	Ser	Val	Thr	His	Leu	Met	Gly	Leu	Phe	Glu	Pro	Ala	260	265	270	
Asp	Met	Lys	Tyr	Asn	Val	Gln	Gln	Asp	His	Thr	Lys	Asp	Pro	Thr	Leu	275	280	285	
Ala	Glu	Met	Thr	Glu	Ala	Ala	Leu	Gln	Val	Leu	Ser	Arg	Asn	Pro	Arg	290	295	300	
Gly	Phe	Tyr	Leu	Phe	Val	Glu	Gly	Gly	Arg	Ile	Asp	His	Gly	His	His	305	310	315	320

30

Asp Gly Lys Ala Tyr Met Ala Leu Thr Glu Ala Ile Met Phe Asp Asn
 325 330 335

Ala Ile Ala Lys Ala Asn Glu Leu Thr Ser Glu Leu Asp Thr Leu Ile
 340 345 350

Leu Val Thr Ala Asp His Ser His Val Phe Ser Phe Gly Gly Tyr Thr
 355 360 365

Leu Arg Gly Thr Ser Ile Phe Gly Leu Ala Pro Gly Lys Ala Leu Asp
 370 375 380

Ser Lys Ser Tyr Thr Ser Ile Leu Tyr Gly Asn Gly Pro Gly Tyr Ala
 385 390 395 400

Leu Gly Gly Gly Ser Arg Pro Asp Val Asn Gly Ser Thr Ser Glu Glu
 405 410 415

Pro Ser Tyr Arg Gln Gln Ala Ala Val Pro Leu Ala Ser Glu Thr His
 420 425 430

Gly Gly Glu Asp Val Ala Val Phe Ala Arg Gly Pro Gln Ala His Leu
 435 440 445

Val His Gly Val Gln Glu Glu Thr Phe Val Ala His Ile Met Ala Phe
 450 455 460

Ala Gly Cys Val Glu Pro Tyr Thr Asp Cys Asn Leu Pro Ala Pro Ala
 465 470 475 480

Thr Ala Thr Ser Ile Pro Asp
 485

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1476 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

gaattcttga ttccagctga agaagaaaat ccagcttttt ggaatagaca agctgctcaa 60
 gctttggatg ttgctaagaa gttgcaacca attcaaactg ctgctaagaa tgttattttg 120
 tttttgggtg atggtatggg tgttccaact gttactgcta ctagaatttt gaaggggtcaa 180
 atgaatggta agttgggtcc agaaactcca ttggctatgg atcaatttcc atacgttgct 240
 ttgtctaaga cttacaatgt tgatagacaa gttccagatt ctgctgggtac tgctactgct 300
 tacttggtg gtggttaaggg taattacaga actattgggtg tttctgctgc tgctagatac 360
 aatcaatgta atactactag aggtaatgaa gttacttctg ttattaatag agctaagaag 420
 gctggtaagg ctgttggtgt tgttactact actagagttc aacatgcttc tccagctgggt 480
 gcttacgctc atactgttaa tagaaattgg tactctgatg ctgatttgcc agctgatgct 540
 caaaagaatg gttgtcaaga tattgctgct caattggttt acaatatgga tattgatggt 600
 attttgggtg gtggtagaat gtacatgttt ccagaaggta ctccagatcc agaataccca 660
 gatgatgctt ctgttaatgg tgttagaaag gataagcaaa atttggttca agaatggcaa 720
 gctaagcatc aagggtgctca atatgtttgg aatagaactg ctttggttgca agctgctgat 780
 gattctagtg ttactcattt gatgggtttg tttgaaccag ctgatatgaa gtataatggt 840
 caacaagatc atactaagga tccaactttg gctgaaatga ctgaagctgc tttgcaagtt 900
 ttgtctagaa atccaagagg tttttacttg tttggtgaag gtggtagaat tgatcatggt 960
 catcatgatg gtaaggctta tatggctttg actgaagcta ttatgtttga taatgctatt 1020
 gctaaggcta atgaattgac ttctgaattg gatactttga ttttgggttac tgctgatcat 1080
 agtcatgttt tttctttttg tggttacact ttgagaggta cttctatttt tggtttggct 1140
 ccaggtaagg ctttggatag taagtcttac acttctattt tgtatggtaa tgggtccagg 1200
 tatgctttgg gtgggtgggtc tagaccagat gttaatggta gtactagtga agaaccatct 1260
 tacagacaac aagctgctgt tccattggct agtgaaactc atgggtgggtga agatgttgct 1320
 gtttttgcta gaggtccaca agctcatttg gttcatgggtg ttcaagaaga aacttttggt 1380
 gctcatatta tggcttttgc tgggttggtt gaaccataca ctgattgtaa tttgccagct 1440

ccagctactg ctactagtat tccagattaa ggtacc

1476

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 78 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

gcgcgaattc ttgattccag ctgaagaaga aaatccagct ttttgggaata gacaagctgc 60

tcaagctttg gatgttgc

78

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 70 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ccaaaaacaa aataacattc ttagcagcag tttgaattgg ttgcaacttc ttagcaacat 60

ccaaagcttg

70

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 69 bases

33

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

gaatggttatt ttgttttttgg gtgatgggtat ggggtgttcca actggttactg ctactagaat 60
tttgaaggg 69

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 70 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ggaaattgat ccatagccaa tggagtttct ggaccaact taccattcat ttgacccttc 60
aaaattctag 70

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 71 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

34

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

gctatggatc aatttccata cgttgctttg tctaagactt acaatggtga tagacaagtt 60
ccagattctg c 71

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 71 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ccaatagttc tgtaattacc cttaacacca cacaagtaag cagtagcagt accagcagaa 60
tctggaactt g 71

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 72 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

35

gtaattacag aactattggt gtttctgctg ctgctagata caatcaatgt aatactacta 60
gaggtaatga ag 72

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 74 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

agtaacaaca ccaacagcct taccagcctt cttagctcta ttaataacag aagtaacttc 60
attacctcta gtag 74

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 74 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

gctgttggtg ttgttactac tactagagtt caacatgctt ctccagctgg tgcttacgct 60
catactgtta atag 74

36

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 68 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

caaccattct tttgagcatc agctggcaaa tcagcatcag agtaccaatt tctattaaca 60
gtatgagc 68

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

gatgctcaaa agaatggttg tcaagatatt gctgctcaat tggtttaciaa tatgg 55

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 72 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

37

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(A) NAME/KEY: Artificial Sequence

(B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

ccttctggaa acatgtacat tctaccacca cccaaaataa catcaatatc catattgtaa 60
accaattgag ca 72

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 71 bases

(B) TYPE: nucleic acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(A) NAME/KEY: Artificial Sequence

(B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

gtacatgttt ccagaaggta ctccagatcc agaataccca gatgatgctt ctgttaatgg 60
tgttagaaag g 71

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 73 bases

(B) TYPE: nucleic acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(A) NAME/KEY: Artificial Sequence

(B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

38

catattgagc accttgatgc ttagcttgcc attcttgaac caaattttgc ttatcctttc 60
taacaccatt aac 73

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 71 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

gcatcaaggc gctcaatatg tttggaatag aactgctttg ttgcaagctg ctgatgattc 60
tagtggttact c 71

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 54 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

cttcatatca gctgggttcaa acaaaccat caaatgagta acactagaat catc 54

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 59 bases
- (B) TYPE: nucleic acid

39

- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

gaaccagctg atatgaagta taatgttcaa caagatcata ctaaggatcc aactttggc 59

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 67 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

cctcttggat ttctagacaa aacttgcaaa gcagcttcag tcatttcagc caaagttgga 60
tccttag 67

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 69 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

gtctagaaat ccaagagggt tttacttggt tggtgaaggt ggtagaattg atcatgggtca 60
tcatgatgg 69

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 73 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

ccttagcaat agcattatca aacataatag cttcagtcaa agccatataa gccttaccat 60
catgatgacc atg 73

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 74 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

gataatgcta ttgctaaggc taatgaattg acttctgaat tggatacttt gattttgggtt 60
actgctgatc atag 74

41

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 73 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

ccaaaccaa aatagaagta cctctcaaag tgtaaccacc aaaagaaaa acatgactat 60
gatcagcagt aac 73

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 73 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

cttctatattt tggtttggct ccaggtaagg ctttggatag taagtcttac acttctattt 60
tgtatggtaa tgg 73

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 76 bases

42

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

ctagtactac cattaacatc tgggtctagaa ccaccaccca aagcataacc tggaccatta 60
ccatacaaaa tagaag 76

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 77 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

gatgttaatg gtagtactag tgaagaacca tcttacagac aacaagctgc tgttccattg 60
gctagtgaaa ctcattgg 77

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 73 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

43

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(A) NAME/KEY: Artificial Sequence

(B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

caccatgaac caaatgagct tgtggacctc tagcaaaaac agcaacatct tcaccaccat 60
gagtttcact agc 73

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 74 bases

(B) TYPE: nucleic acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(A) NAME/KEY: Artificial Sequence

(B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

gctcatttgg ttcattggtgt tcaagaagaa actttttgttg ctcatattat ggcttttgct 60
ggttgtgttg aacc 74

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 82 bases

(B) TYPE: nucleic acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(A) NAME/KEY: Artificial Sequence

(B) LOCATION:

44

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

gcgcggtacc ttaatctgga atactagtag cagtagctgg agctggcaaa ttacaatcag 60
tgtatgggttc aacacaacca gc 82

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

gcgcgcctag gagatctaac atccaaagac g 31

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: Artificial Sequence
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

cgcgcgctag cggatccgca caaacgaag 29

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: PRT

(ix) FEATURE:

- (A) NAME/KEY: *Saccharomyces cerevisiae*
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Glu Ala Glu Ala Glu Phe Leu Ile Pro Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: PRT

(ix) FEATURE:

- (A) NAME/KEY: *Saccharomyces cerevisiae*
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Leu Ile Pro Ala
1

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

46

(ii) MOLECULE TYPE: PRT

(ix) FEATURE:

(A) NAME/KEY: *Saccharomyces cerevisiae*

(B) LOCATION:


(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

Glu	Ala	Glu	Ala	Glu	Phe
1				5	

CLAIMS:


1. Process for the production of a eukaryotic alkaline phosphatase in yeast cells comprising the steps:
 - a) cloning a cDNA sequence codon-optimized for expression in yeast, which codes for a eukaryotic alkaline phosphatase into different vectors
 - b) transformation of the yeast,
 - c) expression and
 - d) purification of the alkaline phosphatase, wherein
 - a first vector has a resistance gene for a first selection marker
 - transformants which have integrated the resistance gene and the cDNA sequence into the genome are selected by growth on nutrient medium containing a low concentration of a first selection marker,
 - the gene copy number is increased by multiple transformation in which multiple transformants are selected by growth on nutrient medium at an increased selection pressure,
 - a second vector is added which has a resistance gene for a second selection marker,
 - the gene copy number is increased by multiple transformation with the second vector in which multiple transformants are selected by growth on nutrient medium at an increased selection pressure and
 - those clones are selected which have integrated several copies of the cDNA sequence and the selection marker resistance genes in a stable manner,
 - wherein the cDNA sequence corresponds to SEQ ID NO:5.
2. Process as claimed in claim 1, wherein methylotrophic yeast cells are used.
3. Process as claimed in claim 2, wherein *Pichia pastoris* or *Hansenula polymorpha* is used as the yeast strain.

4. DNA according to SEQ ID NO:5.
5. Vector containing SEQ ID NO:5.
6. Vector as claimed in claim 5, which corresponds to pHAP10-3.
7. Vector containing the entire expression cassette from pHAP10-3 which comprises SEQ ID NO:5.
8. Vector as claimed in claim 7, which corresponds to pHAP10-3/9K.
9. Host strain transformed with a vector as claimed in any one of claims 5 to 8.
10. Host strain as claimed in claim 9, wherein *Pichia pastoris* or *Hansenula polymorpha* is used as the host strain.
11. *Pichia pastori* X-33 strain transformed with a vector as claimed in any one of claims 6 to 8.
12. Process for producing a eukaryotic highly active alkaline phosphatase, wherein the enzyme is expressed in a host strain as claimed in any one of claims 9 to 11.




bIAPII gene for highly active AP via EcoRI/Asp718 in pICZalpha

Fig.1



Synthetic gene for highly active AP via EcoRI/Asp718 in pPICZalpha

Fig. 2




PCR product hAP10-3 (JM105)/PIC5a3a via SacI/NotI in pPIC9K

Mismatch: 0 MinCuts = 1 MaxCuts: 2

With 198 enzymes: *

Fig. 3



Synthetic gene for highly active AP via EcoRI/Asp718 in pPICZalpha