

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3354727 T3**

(12) Oversættelse af
europæisk patentskrift


-
- (51) Int.Cl.: **C 12 N 9/10 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2020-11-16**
- (80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2020-10-07**
- (86) Europæisk ansøgning nr.: **17208538.3**
- (86) Europæisk indleveringsdag: **2010-01-08**
- (87) Den europæiske ansøgnings publiceringsdag: **2018-08-01**
- (30) Prioritet: **2009-01-08 US 143401 P**
- (62) Stamansøgningsnr: **10729606.3**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC
MK MT NL NO PL PT RO SE SI SK SM TR**
- (73) Patenthaver: **Codexis, Inc., 200 Penobscot Drive, Redwood City, CA 94063, USA**
- (72) Opfinder: **MILLER, Gregory, 200 Penobscot Drive, Redwood City, CA 94063, USA
ZHANG, Xiyun, 200 Penobscot Drive, Redwood City, CA 94063, USA
DHAWAN, Ish, Kumar, 200 Penobscot Drive, Redwood City, CA 94063, USA**
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **TRANSAMINASEPOLYPEPTIDER**
- (56) Fremdragne publikationer:
**WO-A2-2006/063336
WO-A2-2007/130518**

DESCRIPTION

3. BACKGROUND

[0001] Aminotransferases, also known as transaminases (E.C. 2.6.1) catalyze the transfer of an amino group, a pair of electrons, and a proton from a primary amine of an amino donor substrate to the carbonyl group of an amino acceptor molecule. Omega-transaminases (ω -transaminases) transfer amine groups which are separated from a carboxyl group by at least one methylene insertion.

[0002] A general transaminase reaction is shown in Reaction I, below. In this reaction, an amino acceptor (I, keto, or ketone), which is the precursor of the desired amino acid product, is reacted with an amino donor (II). The transaminase enzyme exchanges the amino group of the amino donor with the keto group of the amino acceptor. The reaction therefore results in the desired chiral amine product (III) and a new amino acceptor (keto) compound (IV), which is a by-product.




[0003] Various ω -transaminases have been isolated from microorganisms, including, but not limited to, *Alcaligenes denitrificans*, *Bordetella bronchiseptica*, *Bordetella parapertussis*, *Brucella melitensis*, *Burkholderia malle*, *Burkholderia pseudomallei*, *Chromobacterium violaceum*, *Oceanicola granulosus* HTCC2516, *Oceanobacter* sp. RED65, *Oceanospirillum* sp. MED92, *Pseudomonas putida*, *Ralstonia solanacearum*, *Rhizobium meliloti*, *Rhizobium* sp. (strain NGR234), *Vibrio fluvialis*, *Bacillus thuringensis*, and *Klebsiella pneumoniae* (Shin et al., 2001, Biosci. Biotechnol. Biochem. 65:1782-1788).

[0004] Several aminotransferase gene and enzyme sequences have also been reported, e.g., *Ralstonia solanacearum* (Genbank Acc. No. YP_002257813.1, GI:207739420), *Burkholderia pseudomallei* 1710b (Genbank Acc. No. ABA47738.1, GI:76578263), and *Bordetella petrii* (Genbank Acc. No. AM902716.1, GI:163258032). Two transaminases, EC 2.6.1.18 and EC 2.6.1-19, have been crystallized and characterized (see Yonaha et al., 1983, Agric. Biol. Chem. 47 (10):2257-2265).

[0005] The enzyme ω -amino acid:pyruvate transaminase (ω -APT, E.C. 2.6.1.18) from *Vibrio fluvialis* JS17 carries out the following reaction





using pyridoxal-5'-phosphate as a cofactor. The transaminase from *Vibrio fluvialis* has been reported to show catalytic activity toward aliphatic amines not bearing a carboxyl group.

[0006] Transaminase enzymes have potential industrial use for stereoselective synthesis of optically pure chiral amines and the enantiomeric enrichment of chiral amines and amino acids (Shin et al., 2001, Biosci. Biotechnol. Biochem. 65:1782-1788; Iwasaki et al., 2003, Biotech. Lett. 25:1843-1846; Iwasaki et al., 2004, Appl. Microb. Biotech. 69:499-505, Yun et al., 2004, Appl. Environ. Microbiol. 70:2529-2534; and Hwang et al., 2004, Enzyme Microbiol. Technol. 34:429-426). Chiral amines play an important role in the pharmaceutical, agrochemical and chemical industries and are frequently used as intermediates or synthons for the preparation of various pharmaceuticals, such as cephalosporine or pyrrolidine derivatives. Examples of the use of aminotransferases to generate useful chemical compounds include: preparation of intermediates and precursors of pregabalin (e.g., WO 2008/127646); the stereospecific synthesis and enantiomeric enrichment of β -amino acids (e.g., WO 2005/005633); the enantiomeric enrichment of amines (e.g., US Patent No. US 4,950,606; US Patent No. 5,300,437; and US Patent No. 5,169,780); and the production of amino acids and derivatives (e.g., US Patent No. 5,316,943; US Patent No. 4,518,692; US Patent No. 4,826,766; US Patent No. 6,197,558; and US Patent No. 4,600,692).

[0007] In a great number of the various applications of chiral amines, only one particular optically active form, either the (*R*) or the (*S*) enantiomer is physiologically active. Hence, transaminases are useful for the enantiomeric enrichment and stereoselective synthesis of chiral amines.

[0008] However, transaminases used to mediate transamination reactions can have undesirable properties, such as instability and narrow substrate recognition profiles, thus making them undesirable for commercial applications. Thus, there is a need for other types of transaminases that can be used in processes for preparing chiral amines in an optically active form.

4. SUMMARY


[0009] The present invention provides a transaminase enzyme, comprising a polypeptide having transaminase activity with increased thermostability at 50°C as compared to the polypeptide of SEQ ID NO: 2, wherein the polypeptide comprises an amino acid sequence that is at least 80% identical to SEQ ID NO: 18 and has one or more residue differences as compared to SEQ ID NO: 2 of X9 is T; X86 is Y and X294 is V.

The present invention also provides a polynucleotide encoding a polypeptide of the present invention.

The present invention also provides an expression vector comprising a polynucleotide of the present invention.

The present invention also provides a host cell comprising a polynucleotide of the present invention.

The present invention also provides a method for the conversion of a substrate of Formula I to the chiral amine product of Formula III in stereomeric excess,



wherein the chiral carbon atom of the chiral amine product is marked with an *; each of R^1 , R^2 , R^3 and R^4 , when taken independently, is an unsubstituted or substituted alkyl, alkylaryl, or aryl group, wherein R^1 is different from R^2 in structure or chirality, or R^1 and R^2 , taken together, is a hydrocarbon chain of 4 or more carbon atoms containing a center of chirality, the method comprising contacting the substrate of Formula I with a transaminase polypeptide of the invention in presence of an amino donor under suitable reaction conditions for the conversion of the substrate to the product of Formula III in stereomeric excess

The present disclosure provides transaminase polypeptides having the ability to catalyze the transfer of an amino group from a donor amine to an amine acceptor molecule, polynucleotides encoding such polypeptides, and methods for using the polypeptides. Generally, transaminase polypeptides are useful for the enantiomeric enrichment and stereoselective synthesis of chiral amines.

[0010] In one aspect, the transaminase polypeptides described herein have altered properties as compared to a reference enzyme, such as the naturally occurring transaminase obtained with *Vibrio fluvialis* (e.g., SEQ ID NO:2) or an engineered transaminase, such as the polypeptide of SEQ ID NO:18. Changes to enzyme properties can include, among others, improvements in enzymatic activity, stereoselectivity, stereospecificity, thermostability, solvent stability, and/or reduced substrate or product inhibition. The altered properties are based on engineered transaminase polypeptides having residue differences at specific residue positions as compared to a reference sequence of a naturally occurring *Vibrio fluvialis* transaminase or a reference engineered transaminase, such as the sequence of SEQ ID NO:18. In some aspects of the disclosure, the residue differences are present at one or more of the following residue positions: X4, X6, X9, X12, X21, X30, X31, X44, X45, X, 56, X57, X81, X82, X85, X86, X95, X112, X113, X127, X147, X153, X157, X166, X177, X181, X208, X211, X228, X233, X253, X272, X294, X297, X302, X311, X314, X316, X317, X318, X319, X320, X321, X324, X385, X391, X398, X407, X408, X409, X415, X417, X418, X420, X431, X434, X438, X444, and X446. Amino acid residues that can be present at the specified residue positions and the associated changes to enzyme properties are provided in the detailed description.

[0011] The transaminase polypeptides of the disclosure may be characterized by increased thermostability as compared to the wild-type polypeptide under the same reaction conditions.

Thus, the polypeptides are capable of mediating transamination reactions (e.g., reaction schemes I, II or III), as indicated by the continued formation of products, at higher temperatures and for longer times than the wild-type polypeptide. In some embodiments, the transaminase polypeptides of the invention are improved in being able to retain activity under higher concentrations of the donor amine, such as 2 M concentration of isopropylamine. In some aspects of the disclosure, the engineered transaminases with increased thermostability and/or increased isopropylamine stability comprises an amino acid sequence that is at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the reference sequence of SEQ ID NO: 10, 16 or 18.

[0012] In some embodiments, the engineered transaminase polypeptide can have increased enzymatic activity as compared to the wild type transaminase enzyme or a reference engineered transaminase for transforming the substrates to the products. The amount of the improvement can range from more than 1.1 times (or fold) the enzymatic activity of the corresponding wild-type or reference transaminase enzyme, to as much as 2 times, 5 times, 10 times, 20 times, or more. In some embodiments, the engineered transaminase enzyme exhibits improved enzymatic activity that is at least 1.1-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold, 25-fold, 50-fold, 100-fold, 500-fold, or 1000-fold greater than that of the wild-type transaminase of SEQ ID NO:2 or a reference transaminase enzyme of SEQ ID NO:18. Improvements in enzyme activity may also have associated increases in stereoselectivity, stereospecificity, thermostability, solvent stability, and/or substrate binding, or reduced substrate and/or product inhibition.

[0013] In some embodiments, the engineered transaminases are characterized by activity on a variety of structurally different amine acceptor substrates. In some embodiments, the engineered transaminases are capable of stereoselectively converting substrates 3,4-dihydronaphthalen-1(2*H*)-one, 1-phenylbutan-2-one, 3,3-dimethylbutan-2-one, octan-2-one, 1-(4-bromophenyl)ethanone, 4-phenylbutan-2-one, ethyl 3-oxobutanoate, 1-(6-methoxynaphthalen-2-yl)ethanone, 1-(2,3-dihydrobenzo[*b*][1,4]dioxin-6-yl)ethanone, 1-(4-phenoxyphenyl)ethanone, (*R*)-4-oxotetrahydro-2*H*-pyran-3-yl-benzoate and/or (*R*)-3-(benzyloxy)dihydro-2*H*-pyran-4(3*H*)-one to the corresponding amine product at a rate that is greater than the polypeptides of SEQ ID NO:2 and/or SEQ ID NO:18.

[0014] In some aspects of the disclosure, the improved transaminase polypeptide comprises an amino acid sequence corresponding to SEQ ID NO: 4, 6, 8, 10, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 176, 178, 180, 182, 184, 186, 188, 190, 192, 194, 196, or 198.

[0015] The engineered transaminase enzymes described herein can be obtained by mutagenizing a gene encoding a naturally-occurring wild-type transaminase enzyme that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%,

95%, 96%, 97%, 98%, 99% or more identical to the amino acid sequence of *Vibrio fluvialis* transaminase (SEQ ID NO:2).

[0016] In another aspect, the present disclosure provides polynucleotides encoding the engineered transaminases described herein or polynucleotides that hybridize to such polynucleotides under highly stringent conditions. The polynucleotide can include promoters and other regulatory elements useful for expression of the encoded engineered transaminase, and can utilize codons optimized for specific desired expression systems.

[0017] In some aspects of the disclosure, the polynucleotide encoding the improved transaminases comprises a sequence corresponding to SEQ ID NO: 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 7, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 159, 161, 163, 165, 167, 169, 171, 173, 175, 177, 179, 181, 183, 185, 187, 189, 191, 193, 195, or 197.

[0018] In another aspect, the present disclosure provides host cells comprising the polynucleotides and/or expression vectors described herein. The host cells can be *Vibrio fluvialis* or they may be a different organism, such as *E. coli*. The host cells can be used for the expression and isolation of the engineered transaminase enzymes described herein, or, alternatively, they can be used directly for the conversion of the substrate to the stereoisomeric product.

[0019] In a further aspect, also provided are methods for carrying out reaction Schemes I, II or III (below) using any of the engineered transaminases described herein, which method comprises contacting or incubating the amino acceptor substrate with a transaminase polypeptide of the disclosure in presence of an amino donor under reaction conditions suitable for the conversion of the substrate to the amine product, thereby transforming the substrates to the product compounds. Whether carrying out the method with whole cells, cell extracts or purified transaminase enzymes, a single transaminase enzyme can be used or, alternatively, mixtures of two or more transaminase enzymes can be used.

5. BRIEF DESCRIPTION OF THE FIGURES

[0020] FIG. 1 illustrates the role of transaminases in the conversion of an amino acceptor (ketone) substrate of general formula I and an amino donor of general formula II to the chiral amine (III) product and an amino acceptor (ketone) byproduct (IV). This reaction uses a transaminase described herein and a co-factor such as pyridoxal-5'-phosphate.

6. DETAILED DESCRIPTION

6.1 Abbreviations

[0021] The abbreviations used for the genetically encoded amino acids are conventional and are as follows:

Amino Acid	Three-Letter Abbreviation	One-Letter Abbreviation
Alanine	Ala	A
Arginine	Arg	R
Asparagine	Asn	N
Aspartate	Asp	D
Cysteine	Cys	C
Glutamate	Glu	E
Glutamine	Gln	Q
Glycine	Gly	G
Histidine	His	H
Isoleucine	Ile	I
Leucine	Leu	L
Lysine	Lys	K
Methionine	Met	M
Phenylalanine	Phe	F
Proline	Pro	P
Serine	Ser	S
Threonine	Thr	T
Tryptophan	Trp	W
Tyrosine	Tyr	Y
Valine	Val	V

[0022] When the three-letter abbreviations are used, unless specifically preceded by an "L" or a "D" or clear from the context in which the abbreviation is used, the amino acid may be in either the L- or D-configuration about α -carbon (C_α). For example, whereas "Ala" designates alanine without specifying the configuration about the α -carbon, "D-Ala" and "L-Ala" designate D-alanine and L-alanine, respectively. When the one-letter abbreviations are used, upper case letters designate amino acids in the L-configuration about the α -carbon and lower case letters designate amino acids in the D-configuration about the α -carbon. For example, "A" designates L-alanine and "a" designates D-alanine. When polypeptide sequences are presented as a string of one-letter or three-letter abbreviations (or mixtures thereof), the sequences are presented in the amino (N) to carboxy (C) direction in accordance with common convention.

[0023] The abbreviations used for the genetically encoding nucleosides are conventional and are as follows: adenosine (A); guanosine (G); cytidine (C); thymidine (T); and uridine (U). Unless specifically delineated, the abbreviated nucleotides may be either ribonucleosides or 2'-deoxyribonucleosides. The nucleosides may be specified as being either ribonucleosides or 2'-deoxyribonucleosides on an individual basis or on an aggregate basis. When nucleic acid sequences are presented as a string of one-letter abbreviations, the sequences are presented in the 5' to 3' direction in accordance with common convention, and the phosphates are not indicated.


6.2 Definitions

[0024] As used herein, the following terms are intended to have the following meanings.

[0025] "Protein", "polypeptide," and "peptide" are used interchangeably herein to denote a polymer of at least two amino acids covalently linked by an amide bond, regardless of length or post-translational modification (e.g., glycosylation, phosphorylation, lipidation, myristilation, ubiquitination, etc.). Included within this definition are D- and L-amino acids, and mixtures of D- and L-amino acids.

[0026] "Aminotransferase" and "transaminase" are used interchangeably herein to refer to a polypeptide having an enzymatic capability of transferring an amino group (NH_2), a pair of electrons, and a proton from a primary amine to a carbonyl group ($\text{C}=\text{O}$) of an acceptor molecule. More specifically, the transaminase polypeptides are capable of stereoselectively catalyzing the process of Scheme 1 (below). Transaminases as used herein include naturally occurring (wild type) transaminase as well as non-naturally occurring engineered polypeptides generated by human manipulation.


[0027] "Amino acceptor" and "amine acceptor", "keto substrate", "keto" and "ketone" are used interchangeably herein to refer to a carbonyl (keto, or ketone) compound which accepts an amino group from a donor amine. Amino acceptors are molecules of general Formula I,



in which each of R^1 , R^2 , when taken independently, is an alkyl, an alkylaryl group, or aryl group which is unsubstituted or substituted with one or more enzymatically acceptable groups. R^1 may be the same or different from R^2 in structure or chirality. R^1 and R^2 , taken together, may form a ring that is unsubstituted, substituted, or fused to other rings. Amino acceptors include keto carboxylic acids and alkanones (ketones). Typical keto carboxylic acids are α -keto carboxylic acids such as glyoxalic acid, pyruvic acid, oxaloacetic acid, and the like, as well as salts of these acids. Amino acceptors also include substances which are converted to an amino acceptor by other enzymes or whole cell processes, such as fumaric acid (which can be

converted to oxaloacetic acid), glucose (which can be converted to pyruvate), lactate, maleic acid, and others. Amino acceptors that can be used include, by way of example and not limitation, 3,4-dihydronaphthalen-1(2*H*)-one, 1-phenylbutan-2-one, 3,3-dimethylbutan-2-one, octan-2-one, ethyl 3-oxobutanoate, 4-phenylbutan-2-one, 1-(4-bromophenyl)ethanone, 2-methyl-cyclohexanone, 7-methoxy-2-tetralone, 1-hydroxybutan-2-one, pyruvic acid, acetophenone, 2-methoxy-5-fluoroacetophenone, levulinic acid, 1-phenylpropan-1-one, 1-(4-bromophenyl)propan-1-one, 1-(4-nitrophenyl)propan-1-one, 1-phenylpropan-2-one, 2-oxo-3-methylbutanoic acid, 1-(3-trifluoromethylphenyl)propan-1-one, hydroxypropanone, methoxyoxypropanone, 1-phenylbutan-1-one, 1-(2,5-dimethoxy-4-methylphenyl)butan-2-one, 1-(4-hydroxyphenyl)butan-3-one, 2-acetylnaphthalene, phenylpyruvic acid, 2-ketoglutaric acid, and 2-ketosuccinic acid, including both (*R*) and (*S*) single isomers where possible.

[0028] "Amino donor" refers to an amino compound which donates an amino group to the amino acceptor, thereby becoming a carbonyl species. Amino donors are molecules of general Formula II,



in which each of R^3 , R^4 , when taken independently, is an alkyl, an alkylaryl group, or aryl group which is unsubstituted or substituted with one or more enzymatically non-inhibiting groups. R^3 can be the same or different from R^4 in structure or chirality. R^3 and R^4 , taken together, may form a ring that is unsubstituted, substituted, or fused to other rings. Typical amino donors that can be used with the invention include chiral and achiral amino acids, and chiral and achiral amines. Amino donors that can be used with the invention include, by way of example and not limitation, isopropylamine (also termed 2-aminopropane), α -phenethylamine (also termed 1-phenylethanamine), and its enantiomers (*S*)-1-phenylethanamine and (*R*)-1-phenylethanamine, 2-amino-4-phenylbutane, glycine, L-glutamic acid, L-glutamate, monosodium glutamate, L-alanine, D-alanine, D,L-alanine, L-aspartic acid, L-lysine, L-ornithine, β -alanine, taurine, n-octylamine, cyclohexylamine, 1,4-butanediamine, 1,6-hexanediamine, 6-aminohexanoic acid, 4-aminobutyric acid, tyramine, and benzyl amine, 2-aminobutane, 2-amino-1-butanol, 1-amino-1-phenylethane, 1-amino-1-(2-methoxy-5-fluorophenyl)ethane, 1-amino-1-phenylpropane, 1-amino-1-(4-hydroxyphenyl)propane, 1-amino-1-(4-bromophenyl)propane, 1-amino-1-(4-nitrophenyl)propane, 1-phenyl-2-aminopropane, 1-(3-trifluoromethylphenyl)-2-aminopropane, 2-aminopropanol, 1-amino-1-phenylbutane, 1-phenyl-2-aminobutane, 1-(2,5-dimethoxy-4-methylphenyl)-2-aminobutane, 1-phenyl-3-aminobutane, 1-(4-hydroxyphenyl)-3-aminobutane, 1-amino-2-methylcyclopentane, 1-amino-3-methylcyclopentane, 1-amino-2-methylcyclohexane, 1-amino-1-(2-naphthyl)ethane, 3-methylcyclopentylamine, 2-methylcyclopentylamine, 2-ethylcyclopentylamine, 2-methylcyclohexylamine, 3-methylcyclohexylamine, 1-aminotetralin, 2-aminotetralin, 2-amino-5-methoxytetralin, and 1-aminoindan, including both (*R*) and (*S*) single isomers where possible and including all possible salts of the amines.

[0029] "Chiral amine" refers to amines of general formula $R^1\text{-CH(NH}_2\text{)-R}^2$ and is employed herein in its broadest sense, including a wide variety of aliphatic and alicyclic compounds of different, and mixed, functional types, characterized by the presence of a primary amino group bound to a secondary carbon atom which, in addition to a hydrogen atom, carries either (i) a divalent group forming a chiral cyclic structure, or (ii) two substituents (other than hydrogen) differing from each other in structure or chirality. Divalent groups forming a chiral cyclic structure include, for example, 2-methylbutane-1,4-diyl, pentane-1,4-diyl, hexane-1,4-diyl, hexane-1,5-diyl, 2-methylpentane-1,5-diyl. The two different substituents on the secondary carbon atom (R^1 and R^2 above) also can vary widely and include alkyl, aralkyl, aryl, halo, hydroxy, lower alkyl, lower alkoxy, lower alkylthio, cycloalkyl, carboxy, carbalkoxy, carbamoyl, mono- and di-(lower alkyl) substituted carbamoyl, trifluoromethyl, phenyl, nitro, amino, mono- and di-(lower alkyl) substituted amino, alkylsulfonyl, arylsulfonyl, alkylcarboxamido, arylcarboxamido, etc., as well as alkyl, aralkyl, or aryl substituted by the foregoing.

[0030] "Pyridoxal-phosphate", "PLP", "pyridoxal-5'-phosphate", "PYP", and "P5P" are used interchangeably herein to refer to the compound that acts as a coenzyme in transaminase reactions. In some embodiments, pyridoxal phosphate is defined by the structure 1-(4'-formyl-3'-hydroxy-2'-methyl-5'-pyridyl)methoxyphosphonic acid, CAS number [54-47-7]. Pyridoxal-5'-phosphate can be produced *in vivo* by phosphorylation and oxidation of pyridoxol (also known as Vitamin B₆). In transamination reactions using transaminase enzymes, the amine group of the amino donor is transferred to the coenzyme to produce a keto byproduct, while pyridoxal-5'-phosphate is converted to pyridoxamine phosphate. Pyridoxal-5'-phosphate is regenerated by reaction with a different keto compound (the amino acceptor). The transfer of the amine group from pyridoxamine phosphate to the amino acceptor produces a chiral amine and regenerates the coenzyme. In some embodiments, the pyridoxal-5'-phosphate can be replaced by other members of the vitamin B₆ family, including pyridoxine (PN), pyridoxal (PL), pyridoxamine (PM), and their phosphorylated counterparts; pyridoxine phosphate (PNP), and pyridoxamine phosphate (PMP).

[0031] "Coding sequence" refers to that portion of a nucleic acid (e.g., a gene) that encodes an amino acid sequence of a protein.

[0032] "Naturally-occurring" or "wild-type" refers to the form found in nature. For example, a naturally occurring or wild-type polypeptide or polynucleotide sequence is a sequence present in an organism that can be isolated from a source in nature and which has not been intentionally modified by human manipulation.

[0033] "Recombinant" or "engineered" or "non-naturally occurring" when used with reference to, e.g., a cell, nucleic acid, or polypeptide, refers to a material, or a material corresponding to the natural or native form of the material, that has been modified in a manner that would not otherwise exist in nature, or is identical thereto but produced or derived from synthetic materials and/or by manipulation using recombinant techniques. Non-limiting examples include, among others, recombinant cells expressing genes that are not found within the native

(non-recombinant) form of the cell or express native genes that are otherwise expressed at a different level.

[0034] "Percentage of sequence identity" and "percentage homology" are used interchangeably herein to refer to comparisons among polynucleotides and polypeptides, and are determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence for optimal alignment of the two sequences. The percentage may be calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Alternatively, the percentage may be calculated by determining the number of positions at which either the identical nucleic acid base or amino acid residue occurs in both sequences or a nucleic acid base or amino acid residue is aligned with a gap to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Those of skill in the art appreciate that there are many established algorithms available to align two sequences. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith and Waterman, 1981, *Adv. Appl. Math.* 2:482, by the homology alignment algorithm of Needleman and Wunsch, 1970, *J. Mol. Biol.* 48:443, by the search for similarity method of Pearson and Lipman, 1988, *Proc. Natl. Acad. Sci. USA* 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the GCG Wisconsin Software Package), or by visual inspection (see generally, *Current Protocols in Molecular Biology*, F. M. Ausubel et al., eds., Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (1995 Supplement) (Ausubel)). Examples of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al., 1990, *J. Mol. Biol.* 215: 403-410 and Altschul et al., 1977, *Nucleic Acids Res.* 3389-3402, respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information website. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as, the neighborhood word score threshold (Altschul et al, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-

scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff, 1989, Proc Natl Acad Sci USA 89:10915). Exemplary determination of sequence alignment and % sequence identity can employ the BESTFIT or GAP programs in the GCG Wisconsin Software package (Accelrys, Madison WI), using default parameters provided.

[0035] "Reference sequence" refers to a defined sequence used as a basis for a sequence comparison. A reference sequence may be a subset of a larger sequence, for example, a segment of a full-length gene or polypeptide sequence. Generally, a reference sequence is at least 20 nucleotide or amino acid residues in length, at least 25 residues in length, at least 50 residues in length, or the full length of the nucleic acid or polypeptide. Since two polynucleotides or polypeptides may each (1) comprise a sequence (i.e., a portion of the complete sequence) that is similar between the two sequences, and (2) may further comprise a sequence that is divergent between the two sequences, sequence comparisons between two (or more) polynucleotides or polypeptide are typically performed by comparing sequences of the two polynucleotides or polypeptides over a "comparison window" to identify and compare local regions of sequence similarity.

[0036] In some embodiments, a "reference sequence" can be based on a primary amino acid sequence, where the reference sequence is a sequence that can have one or more changes in the primary sequence. For instance, a "reference sequence based on SEQ ID NO:2 having at the residue corresponding to X85 an alanine or valine" refers to a reference sequence in which the corresponding residue at X85 in SEQ ID NO:2, which is a phenylalanine, has been changed to an alanine or valine.

[0037] "Comparison window" refers to a conceptual segment of at least about 20 contiguous nucleotide positions or amino acids residues wherein a sequence may be compared to a reference sequence of at least 20 contiguous nucleotides or amino acids and wherein the portion of the sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The comparison window can be longer than 20 contiguous residues, and includes, optionally 30, 40, 50, 100, or longer windows.

[0038] "Substantial identity" refers to a polynucleotide or polypeptide sequence that has at least 80 percent sequence identity, at least 85 percent identity and 89 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison window of at least 20 residue positions, frequently over a window of at least 30-50 residues, wherein the percentage of sequence identity is calculated by comparing the reference sequence to a sequence that includes deletions or additions which

total 20 percent or less of the reference sequence over the window of comparison. In specific embodiments applied to polypeptides, the term "substantial identity" means that two polypeptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 80 percent sequence identity, preferably at least 89 percent sequence identity, at least 95 percent sequence identity or more (e.g., 99 percent sequence identity). Preferably, residue positions which are not identical differ by conservative amino acid substitutions.

[0039] "Corresponding to", "reference to" or "relative to" when used in the context of the numbering of a given amino acid or polynucleotide sequence refers to the numbering of the residues of a specified reference sequence when the given amino acid or polynucleotide sequence is compared to the reference sequence. In other words, the residue number or residue position of a given polymer is designated with respect to the reference sequence rather than by the actual numerical position of the residue within the given amino acid or polynucleotide sequence. For example, a given amino acid sequence, such as that of an engineered transaminase, can be aligned to a reference sequence by introducing gaps to optimize residue matches between the two sequences. In these cases, although the gaps are present, the numbering of the residue in the given amino acid or polynucleotide sequence is made with respect to the reference sequence to which it has been aligned.

[0040] "Stereoselectivity" refers to the preferential formation in a chemical or enzymatic reaction of one stereoisomer over another. Stereoselectivity can be partial, where the formation of one stereoisomer is favored over the other, or it may be complete where only one stereoisomer is formed. When the stereoisomers are enantiomers, the stereoselectivity is referred to as enantioselectivity, the fraction (typically reported as a percentage) of one enantiomer in the sum of both. It is commonly alternatively reported in the art (typically as a percentage) as the enantiomeric excess (e.e.) calculated therefrom according to the formula $[\text{major enantiomer} - \text{minor enantiomer}] / [\text{major enantiomer} + \text{minor enantiomer}]$. Where the stereoisomers are diastereoisomers, the stereoselectivity is referred to as diastereoselectivity, the fraction (typically reported as a percentage) of one diastereomer in a mixture of two diastereomers, commonly alternatively reported as the diastereomeric excess (d.e.). Enantiomeric excess and diastereomeric excess are types of stereomeric excess.

[0041] "Highly stereoselective" refers to a transaminase polypeptide that is capable of converting the substrate to the corresponding chiral amine product with at least about 85% stereomeric excess.

[0042] "Stereospecificity" refers to the preferential conversion in a chemical or enzymatic reaction of one stereoisomer over another. Stereospecificity can be partial, where the conversion of one stereoisomer is favored over the other, or it may be complete where only one stereoisomer is converted.

[0043] "Chemoselectivity" refers to the preferential formation in a chemical or enzymatic reaction of one product over another.

[0044] "Improved enzyme property" refers to a transaminase polypeptide that exhibits an improvement in any enzyme property as compared to a reference transaminase. For the engineered transaminase polypeptides described herein, the comparison is generally made to the wild-type transaminase enzyme, although in some embodiments, the reference transaminase can be another improved engineered transaminase. Enzyme properties for which improvement is desirable include, but are not limited to, enzymatic activity (which can be expressed in terms of percent conversion of the substrate), thermo stability, solvent stability, pH activity profile, cofactor requirements, refractoriness to inhibitors (e.g., substrate or product inhibition), stereospecificity, and stereoselectivity (including enantioselectivity).

[0045] "Increased enzymatic activity" refers to an improved property of the engineered transaminase polypeptides, which can be represented by an increase in specific activity (e.g., product produced/time/weight protein) or an increase in percent conversion of the substrate to the product (e.g., percent conversion of starting amount of substrate to product in a specified time period using a specified amount of transaminase) as compared to the reference transaminase enzyme. Exemplary methods to determine enzyme activity are provided in the Examples. Any property relating to enzyme activity may be affected, including the classical enzyme properties of K_m , V_{max} or k_{cat} , changes of which can lead to increased enzymatic activity. Improvements in enzyme activity can be from about 1.1 times the enzymatic activity of the corresponding wild-type transaminase enzyme, to as much as 2 times, 5 times, 10 times, 20 times, 25 times, 50 times, 75 times, 100 times, or more enzymatic activity than the naturally occurring transaminase or another engineered transaminase from which the transaminase polypeptides were derived. In specific embodiments, the engineered transaminase enzyme exhibits improved enzymatic activity in the range of 1.5 to 50 times, 1.5 to 100 times greater than that of the parent transaminase enzyme. It is understood by the skilled artisan that the activity of any enzyme is diffusion limited such that the catalytic turnover rate cannot exceed the diffusion rate of the substrate, including any required cofactors. The theoretical maximum of the diffusion limit, or k_{cat}/K_m , is generally about 10^8 to 10^9 ($M^{-1} s^{-1}$). Hence, any improvements in the enzyme activity of the transaminase will have an upper limit related to the diffusion rate of the substrates acted on by the transaminase enzyme. Transaminase activity can be measured by any one of standard assays, such as by monitoring changes in spectrophotometric properties of reactants or products. In some embodiments, the amount of products produced can be measured by High-Performance Liquid Chromatography (HPLC) separation combined with UV absorbance or fluorescent detection following o-phthalaldehyde (OPA) derivitization. Comparisons of enzyme activities are made using a defined preparation of enzyme, a defined assay under a set condition, and one or more defined substrates, as further described in detail herein. Generally, when lysates are compared, the numbers of cells and the amount of protein assayed are determined as well as use of identical expression systems and identical host cells to minimize variations in amount of enzyme produced by the host cells and present in the lysates.

[0046] "Conversion" refers to the enzymatic conversion of the substrate(s) to the corresponding product(s). "Percent conversion" refers to the percent of the substrate that is

converted to the product within a period of time under specified conditions. Thus, the "enzymatic activity" or "activity" of a transaminase polypeptide can be expressed as "percent conversion" of the substrate to the product.

[0047] "Thermostable" refers to a transaminase polypeptide that maintains similar activity (more than 60% to 80% for example) after exposure to elevated temperatures (e.g., 40-80°C) for a period of time (e.g., 0.5-24 hrs) compared to the wild-type enzyme.

[0048] "Solvent stable" refers to a transaminase polypeptide that maintains similar activity (more than e.g., 60% to 80%) after exposure to varying concentrations (e.g., 5-99%) of solvent (ethanol, isopropyl alcohol, dimethylsulfoxide (DMSO), tetrahydrofuran, 2-methyltetrahydrofuran, acetone, toluene, butyl acetate, methyl tert-butyl ether, etc.) for a period of time (e.g., 0.5-24 hrs) compared to the wild-type enzyme.

[0049] "pH stable" refers to a transaminase polypeptide that maintains similar activity (more than e.g., 60% to 80%) after exposure to high or low pH (e.g. 4.5-6 or 8 to 12) for a period of time (e.g. 0.5-24 hrs) compared to the untreated enzyme.

[0050] "Thermo- and solvent stable" refers to a transaminase polypeptide that is both thermostable and solvent stable.

[0051] "Derived from" as used herein in the context of engineered transaminase enzymes, identifies the originating transaminase enzyme, and/or the gene encoding such transaminase enzyme, upon which the engineering was based. For example, the engineered transaminase enzyme of SEQ ID NO:18 was obtained by artificially evolving, over multiple generations the gene encoding the *Vibrio fluvialis* transaminase enzyme of SEQ ID NO:2. Thus, this engineered transaminase enzyme is "derived from" the wild-type transaminase of SEQ ID NO:2.

[0052] "Hydrophilic Amino Acid or Residue" refers to an amino acid or residue having a side chain exhibiting a hydrophobicity of less than zero according to the normalized consensus hydrophobicity scale of Eisenberg et al., 1984, J. Mol. Biol. 179:125-142. Genetically encoded hydrophilic amino acids include L-Thr (T), L-Ser (S), L-His (H), L-Glu (E), L-Asn (N), L-Gln (Q), L-Asp (D), L-Lys (K) and L-Arg (R).

[0053] "Acidic Amino Acid or Residue" refers to a hydrophilic amino acid or residue having a side chain exhibiting a pK value of less than about 6 when the amino acid is included in a peptide or polypeptide. Acidic amino acids typically have negatively charged side chains at physiological pH due to loss of a hydrogen ion. Genetically encoded acidic amino acids include L-Glu (E) and L-Asp (D).

[0054] "Basic Amino Acid or Residue" refers to a hydrophilic amino acid or residue having a side chain exhibiting a pK value of greater than about 6 when the amino acid is included in a peptide or polypeptide. Basic amino acids typically have positively charged side chains at

physiological pH due to association with hydronium ion. Genetically encoded basic amino acids include L-Arg (R) and L-Lys (K).

[0055] "Polar Amino Acid or Residue" refers to a hydrophilic amino acid or residue having a side chain that is uncharged at physiological pH, but which has at least one bond in which the pair of electrons shared in common by two atoms is held more closely by one of the atoms. Genetically encoded polar amino acids include L-Asn (N), L-Gln (Q), L-Ser (S) and L-Thr (T).

[0056] "Hydrophobic Amino Acid or Residue" refers to an amino acid or residue having a side chain exhibiting a hydrophobicity of greater than zero according to the normalized consensus hydrophobicity scale of Eisenberg et al., 1984, J. Mol. Biol. 179:125-142. Genetically encoded hydrophobic amino acids include L-Pro (P), L-Ile (I), L-Phe (F), L-Val (V), L-Leu (L), L-Trp (W), L-Met (M), L-Ala (A) and L-Tyr (Y).

[0057] "Aromatic Amino Acid or Residue" refers to a hydrophilic or hydrophobic amino acid or residue having a side chain that includes at least one aromatic or heteroaromatic ring. Genetically encoded aromatic amino acids include L-Phe (F), L-Tyr (Y) and L-Trp (W). Although owing to the pKa of its heteroaromatic nitrogen atom L-His (H) it is sometimes classified as a basic residue, or as an aromatic residue as its side chain includes a heteroaromatic ring, herein histidine is classified as a hydrophilic residue or as a "constrained residue" (see below).

[0058] "Constrained Amino Acid or Residue" refers to an amino acid or residue that has a constrained geometry. Herein, constrained residues include L-pro (P) and L-his (H). Histidine has a constrained geometry because it has a relatively small imidazole ring. Proline has a constrained geometry because it also has a five membered ring.

[0059] "Non-polar Amino Acid or Residue" refers to a hydrophobic amino acid or residue having a side chain that is uncharged at physiological pH and which has bonds in which the pair of electrons shared in common by two atoms is generally held equally by each of the two atoms (i.e., the side chain is not polar). Genetically encoded non-polar amino acids include L-Gly (G), L-Leu (L), L-Val (V), L-Ile (I), L-Met (M) and L-Ala (A).

[0060] "Aliphatic Amino Acid or Residue" refers to a hydrophobic amino acid or residue having an aliphatic hydrocarbon side chain. Genetically encoded aliphatic amino acids include L-Ala (A), L-Val (V), L-Leu (L) and L-Ile (I).

[0061] "Cysteine" represented by L-Cys (C) is unusual in that it can form disulfide bridges with other L-Cys (C) amino acids or other sulfanyl- or sulfhydryl-containing amino acids. The "cysteine-like residues" include cysteine and other amino acids that contain sulfhydryl moieties that are available for formation of disulfide bridges. The ability of L-Cys (C) (and other amino acids with -SH containing side chains) to exist in a peptide in either the reduced free -SH or oxidized disulfide-bridged form affects whether L-Cys (C) contributes net hydrophobic or hydrophilic character to a peptide. While L-Cys (C) exhibits a hydrophobicity of 0.29 according

to the normalized consensus scale of Eisenberg (Eisenberg *et al.*, 1984, *supra*), it is to be understood that for purposes of the present disclosure L-Cys (C) is categorized into its own unique group.

[0062] "Small Amino Acid or Residue" refers to an amino acid or residue having a side chain that is composed of a total three or fewer carbon and/or heteroatoms (excluding the α -carbon and hydrogens). The small amino acids or residues may be further categorized as aliphatic, non-polar, polar or acidic small amino acids or residues, in accordance with the above definitions. Genetically-encoded small amino acids include L-Ala (A), L-Val (V), L-Cys (C), L-Asn (N), L-Ser (S), L-Thr (T) and L-Asp (D).

[0063] "Hydroxyl-containing Amino Acid or Residue" refers to an amino acid containing a hydroxyl (-OH) moiety. Genetically-encoded hydroxyl-containing amino acids include L-Ser (S) L-Thr (T) and L-Tyr (Y).

[0064] "Amino acid difference" or "residue difference" refers to a change in the residue at a specified position of a polypeptide sequence when compared to a reference sequence. For example, a residue difference at position X9, where the reference sequence has an alanine, refers to a change of the residue at position X9 to any residue other than alanine. As disclosed herein, an enzyme can include one or more residue differences relative to a reference sequence, where multiple residue differences typically are indicated by a list of the specified positions where changes are made relative to the reference sequence.

[0065] "Conservative" amino acid substitutions or mutations refer to the interchangeability of residues having similar side chains, and thus typically involves substitution of the amino acid in the polypeptide with amino acids within the same or similar defined class of amino acids. However, in some embodiments, conservative mutations as used herein do not include substitutions from a hydrophilic to hydrophilic, hydrophobic to hydrophobic, hydroxyl-containing to hydroxyl-containing, or small to small residue, if the conservative mutation can instead be a substitution from an aliphatic to an aliphatic, non-polar to non-polar, polar to polar, acidic to acidic, basic to basic, aromatic to aromatic, or constrained to constrained residue. Further, as used herein, A, V, L, or I can be conservatively mutated to either another aliphatic residue or to another non-polar residue. The table below shows exemplary conservative substitutions.

Residue	Possible Conservative Mutations
A, L, V, I	Other aliphatic (A, L, V, I) Other non-polar (A, L, V, I, G, M)
G, M	Other non-polar (A, L, V, I, G, M)
D, E	Other acidic (D, E)
K, R	Other basic (K, R)
P, H	Other constrained (P, H)
N, Q, S, T	Other polar
Y, W, F	Other aromatic (Y, W, F)
C	None

[0066] "Non-conservative substitution" refers to substitution or mutation of an amino acid in the polypeptide with an amino acid with significantly differing side chain properties. Non-conservative substitutions may use amino acids between, rather than within, the defined groups listed above. In one embodiment, a non-conservative mutation affects (a) the structure of the peptide backbone in the area of the substitution (e.g., proline for glycine) (b) the charge or hydrophobicity, or (c) the bulk of the side chain.

[0067] "Deletion" refers to modification to the polypeptide by removal of one or more amino acids from the reference polypeptide. Deletions can comprise removal of 1 or more amino acids, 2 or more amino acids, 5 or more amino acids, 10 or more amino acids, 15 or more amino acids, or 20 or more amino acids, up to 10% of the total number of amino acids, or up to 20% of the total number of amino acids making up the reference enzyme while retaining enzymatic activity and/or retaining the improved properties of an engineered transaminase enzyme. Deletions can be directed to the internal portions and/or terminal portions of the polypeptide. In various embodiments, the deletion can comprise a continuous segment or can be discontinuous.

[0068] "Insertion" refers to modification to the polypeptide by addition of one or more amino acids from the reference polypeptide. In some embodiments, the improved engineered transaminase enzymes comprise insertions of one or more amino acids to the naturally occurring transaminase polypeptide as well as insertions of one or more amino acids to other improved transaminase polypeptides. Insertions can be in the internal portions of the polypeptide, or to the carboxy or amino terminus. Insertions as used herein include fusion proteins as is known in the art. The insertion can be a contiguous segment of amino acids or separated by one or more of the amino acids in the naturally occurring polypeptide.

[0069] "Fragment" as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion, but where the remaining amino acid sequence is identical to the corresponding positions in the sequence. Fragments can be at least 14 amino acids long, at least 20 amino acids long, at least 50 amino acids long or longer, and up to 70%, 80%, 90%, 95%, 98%, and 99% of the full-length transaminase polypeptide, for example the polypeptide of SEQ ID NO:2 or engineered transaminase of SEQ ID NO:10.

[0070] "Isolated polypeptide" refers to a polypeptide which is substantially separated from other contaminants that naturally accompany it, e.g., protein, lipids, and polynucleotides. The term embraces polypeptides which have been removed or purified from their naturally-occurring environment or expression system (e.g., host cell or *in vitro* synthesis). The improved transaminase enzymes may be present within a cell, present in the cellular medium, or prepared in various forms, such as lysates or isolated preparations. As such, in some embodiments, the improved transaminase enzyme can be an isolated polypeptide.

[0071] "Substantially pure polypeptide" refers to a composition in which the polypeptide

species is the predominant species present (i.e., on a molar or weight basis it is more abundant than any other individual macromolecular species in the composition), and is generally a substantially purified composition when the object species comprises at least about 50 percent of the macromolecular species present by mole or % weight. Generally, a substantially pure transaminase composition will comprise about 60 % or more, about 70% or more, about 80% or more, about 90% or more, about 95% or more, and about 98% or more of all macromolecular species by mole or % weight present in the composition. In some embodiments, the object species is purified to essential homogeneity (i.e., contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species. Solvent species, small molecules (<500 Daltons), and elemental ion species are not considered macromolecular species. In some embodiments, the isolated improved transaminases polypeptide is a substantially pure polypeptide composition.

[0072] "Stringent hybridization" is used herein to refer to conditions under which nucleic acid hybrids are stable. As known to those of skill in the art, the stability of hybrids is reflected in the melting temperature (T_m) of the hybrids. In general, the stability of a hybrid is a function of ion strength, temperature, G/C content, and the presence of chaotropic agents. The T_m values for polynucleotides can be calculated using known methods for predicting melting temperatures (see, e.g., Baldino et al., Methods Enzymology 168:761-777; Bolton et al., 1962, Proc. Natl. Acad. Sci. USA 48:1390; Bresslauer et al., 1986, Proc. Natl. Acad. Sci USA 83:8893-8897; Freier et al., 1986, Proc. Natl. Acad. Sci USA 83:9373-9377; Kierzek et al., Biochemistry 25:7840-7846; Rychlik et al., 1990, Nucleic Acids Res 18:6409-6412 (erratum, 1991, Nucleic Acids Res 19:698); Sambrook et al., supra); Suggs et al., 1981, In Developmental Biology Using Purified Genes (Brown et al., eds.), pp. 683-693, Academic Press; and Wetmur, 1991, Crit Rev Biochem Mol Biol 26:227-259). In some aspects, the polynucleotide encodes the polypeptide disclosed herein and hybridizes under defined conditions, such as moderately stringent or highly stringent conditions, to the complement of a sequence encoding an engineered transaminase enzyme of the present disclosure.

[0073] "Hybridization stringency" relates to hybridization conditions, such as washing conditions, in the hybridization of nucleic acids. Generally, hybridization reactions are performed under conditions of lower stringency, followed by washes of varying but higher stringency. The term "moderately stringent hybridization" refers to conditions that permit target-DNA to bind a complementary nucleic acid that has about 60% identity, preferably about 75% identity, about 85% identity to the target DNA, with greater than about 90% identity to target-polynucleotide. Exemplary moderately stringent conditions are conditions equivalent to hybridization in 50% formamide, 5× Denhart's solution, 5×SSPE, 0.2% SDS at 42°C, followed by washing in 0.2×SSPE, 0.2% SDS, at 42°C. "High stringency hybridization" refers generally to conditions that are about 10°C or less from the thermal melting temperature T_m as determined under the solution condition for a defined polynucleotide sequence. In some embodiments, a high stringency condition refers to conditions that permit hybridization of only those nucleic acid sequences that form stable hybrids in 0.018M NaCl at 65°C (i.e., if a hybrid is not stable in 0.018M NaCl at 65°C, it will not be stable under high stringency conditions, as contemplated

herein). High stringency conditions can be provided, for example, by hybridization in conditions equivalent to 50% formamide, 5× Denhart's solution, 5×SSPE, 0.2% SDS at 42°C, followed by washing in 0.1×SSPE, and 0.1% SDS at 65°C. Another high stringency condition is hybridizing in conditions equivalent to hybridizing in 5X SSC containing 0.1% (w:v) SDS at 65°C and washing in 0.1x SSC containing 0.1% SDS at 65°C. Other high stringency hybridization conditions, as well as moderately stringent conditions, are described in the references cited above.

[0074] "Heterologous" polynucleotide refers to any polynucleotide that is introduced into a host cell by laboratory techniques, and includes polynucleotides that are removed from a host cell, subjected to laboratory manipulation, and then reintroduced into a host cell.

[0075] "Codon optimized" refers to changes in the codons of the polynucleotide encoding a protein to those preferentially used in a particular organism such that the encoded protein is efficiently expressed in the organism of interest. Although the genetic code is degenerate in that most amino acids are represented by several codons, called "synonyms" or "synonymous" codons, it is well known that codon usage by particular organisms is nonrandom and biased towards particular codon triplets. This codon usage bias may be higher in reference to a given gene, genes of common function or ancestral origin, highly expressed proteins versus low copy number proteins, and the aggregate protein coding regions of an organism's genome. The polynucleotides encoding the transaminases enzymes may be codon optimized for optimal production from the host organism selected for expression.

[0076] "Preferred, optimal, high codon usage bias codons" refers interchangeably to codons that are used at higher frequency in the protein coding regions than other codons that code for the same amino acid. The preferred codons may be determined in relation to codon usage in a single gene, a set of genes of common function or origin, highly expressed genes, the codon frequency in the aggregate protein coding regions of the whole organism, codon frequency in the aggregate protein coding regions of related organisms, or combinations thereof. Codons whose frequency increases with the level of gene expression are typically optimal codons for expression. A variety of methods are known for determining the codon frequency (e.g., codon usage, relative synonymous codon usage) and codon preference in specific organisms, including multivariate analysis, for example, using cluster analysis or correspondence analysis, and the effective number of codons used in a gene (see GCG CodonPreference, Genetics Computer Group Wisconsin Package; CodonW, John Peden, University of Nottingham; McInerney, J. O, 1998, *Bioinformatics* 14:372-73; Stenico et al., 1994, *Nucleic Acids Res.* 22:437-46; Wright, F., 1990, *Gene* 87:23-29). Codon usage tables are available for a growing list of organisms (see for example, Wada et al., 1992, *Nucleic Acids Res.* 20:2111-2118; Nakamura et al., 2000, *Nucl. Acids Res.* 28:292; Duret, et al., *supra*; Henaut and Danchin, "Escherichia coli and Salmonella," 1996, Neidhardt, et al. Eds., ASM Press, Washington D.C., p. 2047-2066. The data source for obtaining codon usage may rely on any available nucleotide sequence capable of coding for a protein. These data sets include nucleic acid sequences actually known to encode expressed proteins (e.g., complete protein coding sequences-CDS), expressed sequence tags (ESTS), or predicted coding regions of genomic sequences (see for

example, Mount, D., Bioinformatics: Sequence and Genome Analysis, Chapter 8, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 2001; Uberbacher, E. C., 1996, Methods Enzymol. 266:259-281; Tiwari et al., 1997, Comput. Appl. Biosci. 13:263-270).

[0077] "Control sequence" is defined herein to include all components, which are necessary or advantageous for the expression of a polynucleotide and/or polypeptide of the present disclosure. Each control sequence may be native or foreign to the nucleic acid sequence encoding the polypeptide. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the nucleic acid sequence encoding a polypeptide.

[0078] "Operably linked" is defined herein as a configuration in which a control sequence is appropriately placed (i.e., in a functional relationship) at a position relative to a polynucleotide of interest such that the control sequence directs or regulates the expression of the polynucleotide and/or polypeptide of interest.

[0079] "Promoter sequence" refers to a nucleic acid sequence that is recognized by a host cell for expression of a polynucleotide of interest, such as a coding sequence. The promoter sequence contains transcriptional control sequences, which mediate the expression of a polynucleotide of interest. The promoter may be any nucleic acid sequence which shows transcriptional activity in the host cell of choice including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

[0080] "Alkyl" by itself or as part of another substituent refers to a saturated or unsaturated branched, straight-chain or cyclic monovalent hydrocarbon radical having the stated number of carbon atoms (e.g., C₁-C₆ means one to six carbon atoms) that is derived by the removal of one hydrogen atom from a single carbon atom of a parent alkane, alkene or alkyne. The term "alkyl" is specifically intended to include groups having any degree or level of saturation, i.e., groups having exclusively single carbon-carbon bonds, groups having one or more double carbon-carbon bonds, groups having one or more triple carbon-carbon bonds and groups having mixtures of single, double and triple carbon-carbon bonds. Where a specific level of saturation is intended, the expressions "alkanyl," "alkenyl," and "alkynyl" are used. The expression "lower alkyl" refers to alkyl groups composed of from 1 to 6 carbon atoms, preferably 1-4 carbon atoms.

[0081] "Alkanyl" by itself or as part of another substituent refers to a saturated branched, straight-chain or cyclic alkyl derived by the removal of one hydrogen atom from a single carbon atom of a parent alkane. Alkanyl groups include, but are not limited to, methanyl; ethanyl; propanyls such as propan-1-yl, propan-2-yl (isopropyl), cyclopropan-1-yl, etc.; butanyls such as butan-1-yl, butan-2-yl (sec-butyl), 2-methyl-propan-1-yl (isobutyl), 2-methyl-propan-2-yl (*t*-

butyl), cyclobutan-1-yl, etc.; and the like. In some embodiments, the alkanyl groups are (C₁-C₆) alkyl.

[0082] "Alkenyl" by itself or as part of another substituent refers to an unsaturated branched, straight-chain or cyclic alkyl having at least one carbon-carbon double bond derived by the removal of one hydrogen atom from a single carbon atom of a parent alkene. The group may be in either the *cis* or *trans* conformation about the double bond(s). Alkenyl groups include, but are not limited to, ethenyl; propenyls such as prop-1-en-1-yl, prop-1-en-2-yl, prop-2-en-1-yl, prop-2-en-2-yl, cycloprop-1-en-1-yl; cycloprop-2-en-1-yl ; butenyls such as but-1-en-1-yl, but-1-en-2-yl, 2-methyl-prop-1-en-1-yl, but-2-en-1-yl, but-2-en-2-yl, buta-1,3-dien-1-yl, buta-1,3-dien-2-yl, cyclobut-1-en-1-yl, cyclobut-1-en-3-yl, cyclobuta-1,3-dien-1-yl, etc.; and the like. In some embodiments, the alkenyl group is (C₂-C₆) alkenyl.

[0083] "Alkynyl" by itself or as part of another substituent refers to an unsaturated branched, straight-chain or cyclic alkyl having at least one carbon-carbon triple bond derived by the removal of one hydrogen atom from a single carbon atom of a parent alkyne. Typical alkynyl groups include, but are not limited to, ethynyl; propynyls such as prop-1-yn-1-yl , prop-2-yn-1-yl, etc.; butynyls such as but-1-yn-1-yl, but-1-yn-3-yl, but-3-yn-1-yl , etc.; and the like. In some embodiments, the alkynyl group is (C₂-C₆) alkynyl.

[0084] "Cycloalkyl" and "Heterocycloalkyl" by themselves or as part of another substituent refer to cyclic versions of "alkyl" and "heteroalkyl" groups, respectively. For heteroalkyl groups, a heteroatom can occupy the position that is attached to the remainder of the molecule. Typical cycloalkyl groups include, but are not limited to, cyclopropyl; cyclobutyls such as cyclobutanyl and cyclobutenyl; cyclopentyls such as cyclopentanyl and cyclopentenyl; cyclohexyls such as cyclohexanyl and cyclohexenyl; and the like. Typical heterocycloalkyl groups include, but are not limited to, tetrahydrofuranyl (e.g., tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, etc.), piperidinyl (e.g., piperidin-1-yl, piperidin-2-yl, etc.), morpholinyl (e.g., morpholin-3-yl, morpholin-4-yl, etc.), piperazinyl (e.g., piperazin-1-yl, piperazin-2-yl, etc.), and the like.

[0085] "Aryl" by itself or as part of another substituent refers to a monovalent aromatic hydrocarbon group having the stated number of carbon atoms (e.g., C₅-C₁₅ means from 5 to 15 carbon atoms) derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. In some embodiments, the aryl group is (C₅-C₁₅) aryl, with (C₅-C₁₀) being even more preferred. In some embodiments, the aryls are cyclopentadienyl, phenyl and naphthyl.

[0086] "Heteroaryl" by itself or as part of another substituent refers to a monovalent heteroaromatic group having the stated number of ring atoms (e.g., "5-14 membered" means from 5 to 14 ring atoms) derived by the removal of one hydrogen atom from a single atom of a parent heteroaromatic ring system. In some embodiments, the heteroaryl group is a 5-14 membered heteroaryl. In some embodiments, the heteroaryl group is a 5-10 membered heteroaryl.

[0087] "Alkoxy" by itself or as part of another substituent refers to $-OR^7$, where R^7 represents an alkyl or cycloalkyl group as defined herein. Typical alkoxy groups include, but are not limited to, methoxy, ethoxy, propoxy, butoxy, cyclohexyloxy, and the like.

[0088] "Alkylcarbonyl" by itself or as part of another substituent refers to $-C(O)-R^8$, where R^8 is an alkyl, as defined above. Typical alkoxycarbonyl include, but are not limited to, acetyl, ethylcarbonyl, n-propylcarbonyl, and the like.

[0089] "Alkoxycarbonyl" by itself or as part of another substituent, refers to $C(O)OR^9$ where R^9 represents an alkyl or cycloalkyl group as defined herein. Typical alkoxycarbonyl groups include, but are not limited to, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, cyclohexyloxycarbonyl, and the like.

[0090] "Amino" by itself or as part of another substituent refers to the group $-NH_2$. Substituted amino refers to the group $-NHR^{10}$, $NR^{10}R^{10}$, and $NR^{10}R^{10}R^{10}$, where each R^{10} is independently selected from substituted or unsubstituted alkyl, cycloalkyl, cycloheteroalkyl, alkoxy, aryl, heteroaryl, heteroarylalkyl, acyl, alkoxycarbonyl, sulfanyl, sulfinyl, sulfonyl, and the like. Typical amino groups include, but are limited to, dimethylamino, diethylamino, trimethylamino, triethylamino, methylsulfonylamino, furanyloxy-sulfamino, and the like.

[0091] "Substituted alkyl, aryl, arylalkyl, heteroaryl or heteroarylalkyl" refers to an alkyl, aryl, arylalkyl, heteroaryl or heteroarylalkyl group in which one or more hydrogen atoms is replaced with another substituent group. Exemplary substituent groups include, but are not limited to, $-OR^{11}$, $-SR^{11}$, $-NR^{11}R^{11}$, $-NO_2$, $-NO$, $-CN$, $-CF_3$, halogen (e.g., $-F$, $-Cl$, $-Br$ and $-I$), $-C(O)R^{11}$, $-C(O)OR^{11}$, $-C(O)NR^{11}$, $-S(O)_2R^{11}$, $-S(O)_2NR^{11}R^{11}$, where each R^{10} is independently selected from the group consisting of hydrogen and (C_1-C_6) alkyl.


[0092] "Substituted" as used herein means one or more hydrogen atoms (e.g., 1, 2, 3, 4, 5, or 6 hydrogen atoms) of the group is replaced with a substituent atom or group commonly used in pharmaceutical chemistry. Each substituent can be the same or different. Examples of suitable substituents include, but are not limited to, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, aralkyl, cycloheteroalkyl, heteroaryl, OR^{12} (e.g., hydroxyl, alkoxy (e.g., methoxy, ethoxy, and propoxy), aryloxy, heteroaryloxy, aralkyloxy, ether, ester, carbamate, etc.), hydroxyalkyl, alkoxycarbonyl, alkoxyalkoxy, perhaloalkyl, perfluoroalkyl (e.g., CF_3 , CF_2 , CF_3), perfluoroalkoxy (e.g., OCF_3 , OCF_2CF_3), alkoxyalkyl, SR^{12} (e.g., thiol, alkylthio, arylthio, heteroarylthio, aralkylthio, etc.), $S^+R^{11}_2$, $S(O)R^{12}$, SO_2R^{12} , $NR^{12}R^{12}$ (e.g., primary amine (i.e., NH_2), secondary amine, tertiary amine, amide, carbamate, urea, etc.), hydrazide, halide, nitrile, nitro, sulfide, sulfoxide, sulfone, sulfonamide, thiol, carboxy, aldehyde, keto, carboxylic acid, ester, amide, imine, and imide, including seleno and thio derivatives thereof, wherein each of the substituents can be optionally further substituted. In embodiments in which a functional group with an aromatic

carbon ring is substituted, such substitutions will typically number about 1 to 5, with about 1 or 2 substitutions being preferred.

6.3 Transaminase Enzymes

[0093] The present disclosure provides transaminase polypeptides having the ability to catalyze the transfer of an amino group from a donor amine to an amine acceptor molecule, polynucleotides encoding such polypeptides, and methods for using the polypeptides. Transaminase polypeptides are useful for the enantiomeric enrichment and stereoselective synthesis of chiral amines.


[0094] The transaminase enzymes described herein are capable of catalyzing the transfer of an amino group from an amino donor of general Formula II to an amino acceptor (ketone substrate) of general Formula I. The products of this reaction are a chiral amine (III) having a stereogenic carbon atom indicated by the * and an amino acceptor (ketone) byproduct (IV), as shown in Scheme 1:



Scheme 1

where the chiral carbon to which the amine is bonded has is predominantly (S) or (R) stereochemistry. As such, the chiral amine product is produced in stereomeric excess.


[0095] In some embodiments, the transaminase enzymes described herein are capable of catalyzing the transfer of an amino group from an amino donor of general Formula II to an amino acceptor (ketone substrate) of general Formula I and stereoselectively forming a chiral amine (III_S) in which the chiral carbon to which the amine is bonded has predominately (S) stereochemistry, and an amino acceptor (ketone) byproduct (IV), as shown in Scheme 2:



Scheme 2

[0096] In some embodiments, the transaminase enzymes described herein are capable of catalyzing the transfer of an amino group from an amino donor of general Formula II to an amino acceptor (ketone) of general Formula I, in which the products of the reaction are a chiral amine (III_R) having predominately (R) stereochemistry, and an amino acceptor (ketone)

byproduct (IV), as shown in Scheme 3:




Scheme 3

[0097] Each of R^1 , R^2 , R^3 and R^4 , when taken independently, can be an alkyl, an alkylaryl group, or aryl group which is unsubstituted or substituted with one or more enzymatically non-inhibiting groups, where R^1 is different from R^2 in structure or chirality, or R^1 and R^2 , taken together, are a hydrocarbon chain of 4 or more carbon atoms containing a center of chirality. In some embodiments, the alkyl group can be a substituted or unsubstituted branched or straight chain alkyl. R^3 may be the same or different from R^4 in structure or chirality. In some embodiments, R^3 and R^4 , taken together, may form a ring that is unsubstituted, substituted, or fused to other rings. In some embodiments, the rings can be substituted or unsubstituted cycloalkyl or heterocycloalkyl. In some embodiments for R^3 and R^4 , the alkyl, either alone or as a substituted or unsubstituted alkylaryl is a lower alkyl.


[0098] In general, the enzymatic process operates on only one chiral form of an amine, or operates on one chiral form to a far greater extent than the other. Thus, the transaminase polypeptides described herein are capable of performing stereoselective synthesis, in which one of the chiral amine products (III_S or III_R) illustrated above is produced in an amount substantially greater than the other. "Substantially greater" as used herein refers to a percentage of the combined amount of both chiral forms that is at least about 51% (e.g., at least 51, 55, 60, 65, 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99%). As such, in some embodiments, the product of the transaminase can be in a stereomeric excess of at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, or 99% or greater.

[0099] Reactions shown in Schemes 1 to 3 use an amino acceptor substrate and an amino donor substrate. Amino acceptors are compounds of general Formula I:



in which each of R^1 , R^2 , when taken independently, can be an alkyl, an alkylaryl group, or aryl group which is unsubstituted or substituted with one or more enzymatically non-inhibiting groups. As noted above, R^1 may be the same or different from R^2 in structure or chirality. In some embodiments, R^1 and R^2 , taken together, may form a ring that is unsubstituted, substituted, or fused to other rings. In some embodiments, the amino acceptor is any compound according to Formula 1, above.

[0100] Amino donors are compounds of general Formula II:



in which each of R^3 , R^4 , when taken independently, is an alkyl, an alkylaryl group, or aryl group which is unsubstituted or substituted with one or more enzymatically non-inhibiting groups. R^3 may be the same or different from R^4 in structure or chirality. R^3 and R^4 , taken together, may form a ring that is unsubstituted, substituted, or fused to other rings. In some embodiments for R^3 and R^4 , the alkyl, either alone or as a substituted or unsubstituted alkylaryl is a lower alkyl. The amino donors can include all salt forms of the compounds of Formula II.

[0101] In some embodiments, the amino donor is any compound according to Formula II, above and all possible salt forms. Exemplary amino donors include, among others, isopropylamine (2-amino propane), D-alanine, L-alanine, or D,L-alanine, and all possible salts of these amino donors.

[0102] As illustrated, the reaction can proceed in either the forward or reverse direction. In some embodiments, either the forward or reverse reactions can be favored by adding or removing starting materials or reaction products. For example, when one stereoselectively synthesizes one chiral form of an amine (III_S or III_R), as shown in Reactions I and II, additional amino acceptor (ketone) or amino donor can be added (up to saturation) and/or the chiral amine or ketone by product formed can be removed.

[0103] For performing the transamination reactions, the present disclosure provides engineered transaminase enzymes that are capable of stereoselectively converting an amino acceptor to the corresponding chiral amino product and which have an improved property when compared with the naturally-occurring enzyme obtained from *Vibrio fluvialis* (SEQ ID NO:2) or when compared with other engineered transaminase enzymes (e.g., SEQ ID NO:18).

[0104] In the characterizations of the transaminases herein, the polypeptides can be described in reference to the amino acid sequence of a naturally occurring transaminase of *Vibrio fluvialis* or another engineered transaminase. As such, the amino acid residue position is determined in the transaminases beginning from the initiating methionine (M) residue (i.e., M represents residue position 1), although it will be understood by the skilled artisan that this initiating methionine residue may be removed by biological processing machinery, such as in a host cell or in vitro translation system, to generate a mature protein lacking the initiating methionine residue. The amino acid residue position at which a particular amino acid or amino acid change is present in an amino acid sequence is sometimes describe herein in terms "Xn" , or "position n", where n refers to the residue position. A substitution mutation, which is a replacement of an amino acid residue in a reference sequence with a different amino acid residue may be denoted by the symbol "→", or by conventional notations used in the art, for example, Y(number)Z, where Y is the amino acid residue in the reference sequence, the

"number" refers to the residue position, and Z is the amino residue substitution.

[0105] The polynucleotide sequence encoding the naturally occurring transaminase of *Vibrio fluvialis* JS17 is described in Shin et al., 2003, "Purification, characterization, and molecular cloning of a novel amine:pyruvate transaminase from *Vibrio fluvialis* JS17" Appl Microbiol Biotechnol. 61(5-6):463-471). The polynucleotide and amino acid sequence of the *Vibrio fluvialis* transaminase is also presented herein as SEQ ID NO:1 and SEQ ID NO:2, respectively.

[0106] As noted above, the transaminases of the disclosure are characterized by an improved enzyme property as compared to the naturally occurring parent enzyme or another engineered transaminase polypeptide. In the embodiments herein, the improved property of the transaminase can be with respect to, among others, enzyme activity, stability (e.g., solvent- and/or thermostability), stereoselectivity, stereospecificity, inhibitor resistance, and substrate recognition. In some embodiments, the transaminase polypeptides can have more than one improved property, such as increased stability and substrate recognition.

[0107] The engineered transaminases disclosed herein have various residue differences as compared to a reference sequence (e.g., naturally-occurring polypeptide of SEQ ID NO:2 or an engineered polypeptide of SEQ ID NO:18) that result in changes to the enzyme properties. The residue differences can be characterized as "modifications" of the reference sequence, where the modifications include amino acid substitutions, deletions, and insertions. Any one or a combination of modifications can be present to generate the improved engineered transaminases. These residue differences can also be described as particular features of the improved engineered polypeptides at specified residue positions.

[0108] The number of residue differences from the reference transaminase polypeptide that provides an improved transaminase property can comprise differences at one or more residue positions, 2 or more residue positions, 3 or more residue positions, 5 or more residue positions, 10 or more residue positions, or 20 or more residue positions, up to 10% of the total number of amino acids, up to 20% of the total number of amino acids, or up to 30% of the total number of amino acids of the reference enzyme sequence. The number of residue differences as compared to the reference sequence can be about 1-2, 1-3, 1-4, 1-5, 1-6, 1-7, 1-8, 1-9, 1-10, 1-15, 1-20, 1-21, 1-22, 1-23, 1-24, 1-25, 1-30, 1-35, 1-40, 1-45, 1-50, 1-55, or 1-60 residue positions. The number of differences as compared to the reference sequence can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 30, 35, 40, 45, 50, 55, or 60 residue positions.

[0109] The residue differences may comprise substitutions as compared to the reference sequence. In some aspects of the disclosure, the substitutions as compared to the reference sequence can be at one or more residue positions, 2 or more residue positions, 3 or more residue positions, 5 or more residue positions, 10 or more residue positions, or 20 or more residue positions, up to 10% of the total number of amino acids, up to 20% of the total number of amino acids, or up to 30% of the total number of amino acids of the reference enzyme sequence. The number of substitutions as compared to the reference sequence can be

substitutions about 1-2, 1-3, 1-4, 1-5, 1-6, 1-7, 1-8, 1-9, 1-10, 1-15, 1-20, 1-21, 1-22, 1-23, 1-24, 1-25, 1-30, 1-35, 1-40, 1-45, 1-50, 1-55, or 1-60 residue positions. The number of substitutions as compared to the reference sequence can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 30, 35, 40, 45, 50, 55, or 60 residue positions.

[0110] In some aspects of the disclosure, the residue differences of the engineered transaminases of the disclosure can occur at one or more of the following residue positions: X4, X9, X12, X21, X30, X31, X44, X45, X, 56, X57, X81, X82, X85, X86, X95, X112, X113, X127, X147, X153, X157, X166, X177, X181, X208, X211, X228, X233, X253, X272, X294, X297, X302, X311, X314, X316, X317, X318, X319, X320, X321, X324, X385, X391, X398, X407, X408, X409, X415, X417, X418, X420, X431, X434, X438, X444, and X446. As described herein, the presence of particular amino acid residues at these residue positions is associated with changes to properties of the transaminases. In some aspects of the disclosure, the number of residue differences can be at least 2, 3, 4, 5, or 6 or more of the specified residue positions.

[0111] In some aspects of the disclosure, the choice of amino acid residues for the specified residue positions can be based on the following features. X4 is a basic residue; X9 is a polar residue; X12 a non-polar residue; X21 is a polar residue; X30 is an aliphatic or non-polar residue; X31 is an aliphatic or non-polar residue; X44 is an aliphatic or non-polar residue; X45 is a constrained residue; X56 is a non-polar or aliphatic residue; X57 is an aromatic, aliphatic, non-polar, polar or cysteine residue; X81 is an acidic residue; X82 is an aromatic or constrained residue; X85 is an aliphatic, non-polar, polar, or cysteine residue; X86 is a constrained, aromatic, or polar residue; X95 is a polar residue; X112 is a non-polar or aliphatic residue; X113 is a cysteine, aromatic or constrained residue; X127 is an aliphatic or non-polar residue; X147 is a non-polar residue; X153 is an aliphatic, non-polar, or polar residue; X157 is a polar residue; X166 is a polar residue; X177 is a non-polar or aliphatic residue; X181 is a basic residue; X208 is a non-polar or aliphatic residue; X211 is an acidic residue; X228 is a non-polar residue; X233 is a polar or aliphatic residue; X253 is a non-polar residue; X272 is a non-polar or aliphatic residue; X294 is a non-polar or aliphatic residue; X297 is an aliphatic residue; X302 is an acidic residue; X311 is an aliphatic residue; X314 is an aliphatic or polar residue; X316 is an acidic residue; X317 is an aromatic, non-polar or aliphatic residue; X318 is an aromatic, basic or non-polar residue; X319 is a non-polar, aliphatic, or polar residue; X320 is an aliphatic residue; X321 is an aliphatic residue; X324 is a non-polar residue; X385 is a basic residue; X398 is a basic residue; X407 is a polar residue; X408 is an aliphatic residue; X409 is non-polar residue; X415 is a non-polar residue; X417 is a non-polar, aliphatic, polar, acidic, or cysteine residue; X418 is a non-polar or aliphatic residue; X420 is a polar residue; X431 is an acidic residue; X434 is an aliphatic residue; X438 is an aliphatic residue; X444 is a non-polar or aliphatic residue; and X446 is a non-polar or aliphatic residue. In some aspects of the disclosure, where the amino acid residue at the corresponding residue position of the reference sequence are encompassed within the category of amino acids described for the specified position, a different amino acid within that amino acid category can be used in light of the guidance provided herein.

[0112] In some aspects, the transaminase polypeptides can have additionally one or more residue differences at residue positions not specified by an X above as compared to a reference sequence, for example SEQ ID NO:2. In some aspects, the differences can be 1-2, 1-3, 1-4, 1-5, 1-6, 1-7, 1-8, 1-9, 1-10, 1-15, 1-20, 1-21, 1-22, 1-23, 1-24, 1-25, 1-30, 1-35, 1-40, 1-45, 1-50, 1-55, or 1-60 residue differences at other amino acid residue positions not defined by X above. In some aspects, the number of differences can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 30, 35, 40, 45, 50, 55, or 60 residue differences at other amino acid residue positions. In some aspects, the differences comprise conservative mutations.

[0113] In some aspects of the disclosure, the amino acid residues at the specified positions can be selected from one or more of the following features: X4 is R; X9 is T; X12 is G; X21 is N; X30 is A; X31 is A; X44 is A; X45 is H; X56 is V; X57 is L, I, F, A, S, or C; X81 is D; X82 is H; X85 is A, S, V, T, N, or C; X86 is S, H, N, F, G, or A; X95 is T; X112 is I; X113 is H or C; X127 is L; X147 is G; X153 is A, S, T, N, G, Q, or C; X157 is T; X166 is S; X177 is L; X181 is R; X208 is I; X211 is K; X228 is G; X233 is T, S or L; X253 is M; X272 is A; X294 is V; X297 is A; X302 is K; X311 is V; X314 is V or T; X316 is K; X317 is L, Y or M; X318 is G, R, or F; X319 is V or Q; X320 is A; X321 is L; X324 is G; X385 is R; X391 is A; X398 is R; X407 is S; X408 is A; X409 is G; X415 is M; X417 is F, A, I, C, T, S, or C; X418 is V; X420 is N; X431 is D; X434 is V; X438 is L; X444 is V; and X446 is V. In some aspects, the transaminase polypeptides can have additionally one or more residue differences at residue positions not specified by an X above as compared to a reference sequence, for example SEQ ID NO:2. In some aspects, the differences can be 1-2, 1-3, 1-4, 1-5, 1-6, 1-7, 1-8, 1-9, 1-10, 1-15, 1-20, 1-21, 1-22, 1-23, 1-24, 1-25, 1-30, 1-35, 1-40, 1-45, 1-50, 1-55, or 1-60 residue differences at other amino acid residue positions not defined by X above. In some aspects, the number of differences can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 30, 35, 40, 45, 50, 55, or 60 residue differences at other amino acid residue positions. In some aspects, the differences comprise conservative mutations. In some aspects of the disclosure, the engineered transaminases can comprise an amino acid sequence that is at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more identical to a reference based on SEQ ID NO:2 having one or more of the preceding features, and is characterized one or more of the improved enzyme properties described herein.

[0114] In some aspects of the disclosure, the improved transaminase polypeptides comprise an amino acid sequence having at least one or more residue differences as compared to the reference sequence, e.g., naturally occurring sequence of SEQ ID NO:2 or an engineered transaminase sequence of SEQ ID NO:18, at the following residue positions: X57, X85, X86, X153, X228, X233, X317, X318, and X417. In some aspects of the disclosure, the residues at these residue positions can be selected from the following features: X57 is an aromatic, aliphatic, non-polar, polar or cysteine residue; X85 is an aliphatic, non-polar, polar, or cysteine residue; X86 is a constrained, aromatic, or polar residue; X153 is an aliphatic, non-polar, or polar residue; X228 is a non-polar residue; X233 is a polar or aliphatic residue; X317 is an aromatic, non-polar or aliphatic residue; X318 is an aromatic, basic or non-polar residue; and X417 is a non-polar, aliphatic, polar, acidic, or cysteine residue. In some aspects of the

disclosure, the residues at these residue positions can be selected from the following features: X57 is L, I, F, A, S, or C; X85 is A, S, V, T, N, or C; X86 is S, H, N, F, G, or A; X153 is A, S, T, N, G, Q, or C; X228 is G; X233 is T, S or L; X317 is L, Y or M; X318 is G, R, or F; and X417 is F, A, I, C, T, S, or C. In some aspects of the disclosure, the engineered transaminases can comprise an amino acid sequence that is at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more identical to a reference based on SEQ ID NO:2 having one or more of the preceding features, and is characterized one or more of the improved enzyme properties described herein.

[0115] In some aspects of the disclosure, the improved transaminase polypeptide can comprise an amino acid sequence that is at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more identical to a sequence corresponding to SEQ ID NO: 4, 6, 8, 10, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 176, 178, 180, 182, 184, 186, 188, 190, 192, 194, 196, or 198. In some of these aspects of the disclosure, the transaminase polypeptides can have one or more of the improved properties described herein.

[0116] In some aspects of the disclosure, the improved transaminase polypeptides comprises an amino acid sequence corresponding to the sequence of SEQ ID NO: 4, 6, 8, 10, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 176, 178, 180, 182, 184, 186, 188, 190, 192, 194, 196, or 198.

[0117] In some aspects of the disclosure, the improved property of the transaminase polypeptide is with respect to its stability, such as thermostability and/or solvent stability. It is advantageous for transaminase polypeptides to be able to retain enzymatic activity at elevated temperatures and solvent conditions in order to drive the reaction to completion. For example, the transaminase polypeptide of SEQ ID NO:2 may lose activity at elevated temperature (e.g., 40-45°C) or in presence of reaction components, e.g., amine donor isopropylamine. The thermostable polypeptides of the disclosure may be characterized by higher retention of activity when treated for about 0.5 to 24 hours at about 35°C to 50°C, at a pH of about 7.5 to 9.0, as compared to transaminase of SEQ ID NO:2. In some embodiments, the amount of enzymatic activity remaining, i.e., residual activity, following treatment at the defined elevated temperature can be at least 5%, 10%, 20%, 30%, 40%, 50% or more of the activity of an untreated polypeptide as assayed at a defined condition. In some embodiments, the elevated temperature condition is 50°C for 23 hrs. Under the latter condition, the naturally-occurring transaminase of SEQ ID NO:2 has no residual activity following treatment at the defined condition while the engineered transaminases of the disclosure retains significant activity. For example, the transaminase of SEQ ID NO:16 or 18 following treatment at 50°C for 23 hrs is

capable of converting 10% and 23%, respectively, of the substrate pyruvate to product alanine in presence of amino donor isopropylamine, under the assay condition described in the Examples.

[0118] In some aspects of the disclosure, the engineered transaminase polypeptides are characterized by resistance against inactivation by a solvent or reaction component, e.g., isopropylamine. In some aspects of the disclosure, the engineered transaminase enzymes are resistant to inactivation by up to 1 M or by up to 2 M isopropylamine. In some embodiments, the amount of enzymatic activity remaining, i.e., residual activity, following treatment with solvent or reaction component (e.g., isopropylamine) can be at least 5%, 10%, 20%, 30%, 40%, 50% or more of the activity of an untreated polypeptide.

[0119] In some aspects of the disclosure, the improved transaminase polypeptide characterized by increased thermo- and/or solvent stability comprises an amino acid sequence that is at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the sequence of SEQ ID NO: 10, 16, or 18. In some embodiments, the transaminase polypeptide characterized by increased thermostability corresponds to the sequence of SEQ ID NO: 10, 16, or 18.

[0120] In some aspects of the disclosure, the transaminase polypeptides having improved thermostability (as compared to SEQ ID NO:2) can have residue differences as compared to SEQ ID NO:2 at one or more of the following residue positions: X4, X9, X12, X21, X45, X86, X157, X177, X208, X211, X253, X272, X294, X302, X316, X324, X391, X398, X418, X420, X431, X444, and X446. In some aspects of the disclosure, the number of residue differences can be at least 2, 3, 4, 5, or 6 or more of the specified residue positions associated with increased thermostability.

[0121] In some aspects of the disclosure, the transaminase polypeptides improved as compared to SEQ ID NO:2 with respect to their thermostability have one or more of the following features: X4 is R, Q, or L; X9 is T; X12 is K; X21 is N; X45 is H; X86 is Y; X157 is T; X177 is L; X208 is I; X211 is K; X253 is M; X272 is A; X294 is V; X302 is K; X316 is K; X324 is G; X391 is A; X398 is R; X418 is V; X420 is N; X431 is D; X444 is V; and X446 is V.

[0122] In some aspects of the disclosure, the improved transaminase polypeptide characterized by increased thermostability includes, but are not limited to, an amino acid sequence having one of the following feature or set of features: X45 is H, X86 is Y, X177 is L, X211 is K, X294 is V, X324 is G, X391 is A, X398 is R, and X420 is N; X9 is T, X45 is H, X86 is Y, X177 is L, X211 is K, X294 is V, X324 is G, and X391 is A; X21 is N, X45 is H, X177 is L, X208 is I, X211 is K, X324 is G, X391 is A; and X9 is T, X21 is N, X86 is Y, X208 is I, and X294 is V.

[0123] In some aspects of the disclosure, the transaminase polypeptide is improved with respect to, among other properties, enzyme activity, substrate recognition, stereospecificity and/or stereoselectivity. These characteristics can be present in the context of a thermo-

and/or solvent stable engineered transaminase, such as SEQ ID NO: 10, 16 or 18. For example, the thermo/solvent stable transaminases can be modified to have improvements in other enzyme properties while retaining its thermo/solvent stable characteristics. A significant advantage of having a thermo- and/or solvent stable transaminase as a basis for obtaining improvements in other enzyme properties, e.g., substrate recognition, enzyme activity, stereoselectivity, stereospecificity, is that the engineered transaminase can be used under a process condition that is unsuitable for the naturally occurring enzyme, for example, among others, (a) screening for activity on various substrates at conditions of elevated temperature and/or high solvent concentrations; (b) scale-up conditions where transamination reactions may be carried out for longer time periods, and/or (c) under high substrate loading conditions. In some embodiments, screening with substrates at the elevated temperature and/or higher solvent concentration maintains the thermo- and/or solvent stable features of the transaminase polypeptide while allowing for identification of improvements in other enzyme properties.

[0124] However, in some aspects of the disclosure, the improvements in other enzyme properties can be in the context of the naturally occurring transaminase polypeptide, such as the sequence of SEQ ID NO:2. Accordingly, in some aspects of the disclosure, the engineered transaminase polypeptides characterized by improvements in enzyme activity, substrate recognition or stereoselectivity can comprise an amino acid sequence that is at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the sequence of SEQ ID NO:2 and having the indicated improved property.

[0125] In some aspects, the improved property of the transaminase polypeptide is with respect to an increase in its rate of conversion of the substrate to the product. This improvement in enzymatic activity can be manifested by the ability to use less of the improved polypeptide as compared to the wild-type (SEQ ID NO:2), or other reference sequence (e.g., SEQ ID NO:18), to produce the same amount of product in a given amount of time. In some aspects, the improvement in enzymatic activity can be manifested by the ability to convert a greater percentage of the substrate to product in a defined time with the same amount of enzyme as compared to the wild type or reference enzyme. In some aspects, the transaminase polypeptides are capable of converting the substrate to products at a rate that is at least 1.1-fold, 1.5 fold, 2 fold, 3 fold, 4 fold, 5-fold, 10-fold, 25-fold, 50-fold, 75-fold, 100-fold, 150-fold, 200-fold, 250-fold, 300-fold, 500-fold, 1000-fold or more over the rate of SEQ ID NO:2 or SEQ ID NO:18.

[0126] In some aspects of the disclosure, the transaminase polypeptides characterized by improved enzymatic activity, i.e., greater rate of converting the substrates to products as compared to SEQ ID NO:2 and/or SEQ ID NO:18, can comprise an amino acid sequence having one or more of the following features: X30 is A; X31 is A; X44 is A; X45 is N; X56 is V; X57 is, A, C, F, I, L, or S; X81 is D; X82 is H; X85 is A, C, S, N, T, G, or V; X86 is F, S, N, A, G, or H; X95 is T; X112 is I; X113 is C or H; X127 is L; X147 is G; X153 is A, C, G, N, M, Q, S, or T; X166 is S; X177 is V; X181 is R; X208 is I; X211 is R; X228 is G or T; X233 is L, S, I, V, N, G, or T; X294 is M; V297 is A, S, T, I, M, Q, C, or G; X311 is V; X314 is T or V; X317 is L, M, or Y; X318 is G, F, C, K, W, or R; X319 is Q, G, M, N, or V; X320 is A or K; X321 is L, M, or I; X324 is

S; X385 is R; X398 is R; X407 is S; X408 is A; X409 is G; X415 is M; X417 is A, C, E, F, I, N, Q, Y, S, T, or V; X420 is N; X434 is V; and X438 is L.

[0127] In some aspects of the disclosure, transaminase polypeptides that are capable of an increased enzymatic activity can comprise an amino acid sequence having one or more of the following features: X30 is A; X31 is A; X56 is V; X81 is D; X82 is H; X95 is T; X113 is C or H.

[0128] In some aspects of the disclosure, the transaminase polypeptides that are improved as compared to SEQ ID NO:2, and/or SEQ ID NO:18 with respect to their rate of enzymatic activity, e.g., their rate of converting the substrates to products can comprise an amino acid sequence having one of the following set of features: X12 is G and X434 is V; X44 is A and X166 is S; X57 is I and X153 is S; X81 is D and X86 is H; X82 is H and X417 is F; X85 is A and X317 is L; X85 is S and X153 is A; X85 is A and X153 is A; X85 is S and X153 is S. X86 is S and X153 is S; X86 is H and X153 is A; X112 is I and X317 is L; or X113 is H and X407 is S; X153 is S and X233 is S; X311 is V and X314 is T; X314 is V and X409 is G; and X318 is G and X408 is A.

[0129] In some aspects of the disclosure, the transaminase polypeptides that are improved as compared to SEQ ID NO:2, and/or SEQ ID NO:18 with respect to their rate of enzymatic activity, i.e., their rate of converting the substrates to products can comprise an amino acid sequence having at least one of the following set of features: X57 is A, X153 is S and X318 is G; X57 is F, X86 is H and X153 is Q; X57 is I, X86 is F and X320 is A; X57 is F, X86 is F and X153 is Q; X57 is A, X153 is C and X321 is L; X57 is C, X86 is S and X417 is T; X57 is C, X86 is A and X317 is L; X57 is F, X318 is F and X417 is S; X57 is L, X86 is S and X153 is A; X57 is F, X318 is G and X417 is I; X57 is L, X86 is F and X318 is F; X57 is L, X417 is C and X438 is L; X57 is F, X127 is L and X417 is C; X57 is S, X233 is L and X417 is V; X57 is S, X86 is G and X417 is C; X85 is A, X147 is G and X153 is A; X85 is S, X153 is A and X233 is T; X85 is A, X153 is S and X417 is S; X86 is H, X153 is A and X417 is C; X86 is H, X153 is S and X417 is C; X86 is F, X153 is C and X297 is A; X86 is S, X153 is T and X297 is A; X86 is H, X233 is L and X417 is A; X86 is F, X318 is R and X417 is A; X86 is N, X228 is G and X317 is L; X95 is T, X153 is A and X417 is C; X113 is C, X385 is R and X417 is C; X153 is A, X317 is L and X318 is G; X153 is A, X233 is T and X417 is C; X153 is S, X318 is R and X417 is E; X153 is C, X233 is L and X318 is R; X153 is T, X228 is G and X321 is L; X153 is S, X317 is L and X417 is C; X153 is T, X319 is V and X417 is I; X153 is S, X228 is G and X417 is V; X153 is C, X317 is Y and X319 is Q; X153 is T, X228 is G and X417 is A; X228 is G, X317 is L and X417 is C; X228 is G, X318 is G and X417 is C; X233 is L, X321 is L and X417 is I; and X317 is L, X318 is R and X417 is T.

[0130] In some aspects of the disclosure, the transaminase polypeptides that are improved as compared to SEQ ID NO:2 and/or SEQ ID NO:18 with respect to their rate of enzymatic activity, i.e., their rate of converting the substrates to products, can comprise an amino acid sequence having one the following set of features: X57 is L and X86 is F and X153 is S and X233 is T and X417 is T; X86 is H and X153 is A and XA228 is G and X417 is I; and X86 is H and X153 is S and X181 is R and X417 is T.

[0131] In some aspects, the transaminase polypeptides described herein are capable of stereoselectively converting amino acceptor substrates to the corresponding chiral amine products, at a rate that is greater than the wild-type transaminase of *Vibrio fluvialis* (SEQ ID NO:2). Polypeptides that are capable of the above reaction include, but are not limited to, a polypeptide comprising an amino acid sequence corresponding to SEQ ID NO:18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 176, 178, 180, 182, 184, 186, 188, 190, or 192. Activities of the engineered transaminases with respect to specific substrates are described in further detail below.

[0132] In some aspects, the improved property of the transaminase polypeptide is with respect to an increase of its stereoselectivity for converting the amino acceptor substrate to a chiral amino product. In some aspects, the improved property of the transaminase is with respect to an increase in stereoselectivity, i.e., herein, an increase in the stereomeric excess of the product. In some aspects, the improved property of the transaminase property is with respect to producing the *S* enantiomers of chiral amines in stereomeric excess. In some aspects, the improved property of the transaminase property is with respect to producing the *R* enantiomers of chiral amines in stereomeric excess.

[0133] In some aspects, the transaminase polypeptides are characterized by the ability to convert amino acceptor substrates to the chiral amine products of the (*R*) configuration in stereomeric excess greater than the polypeptide of SEQ ID NO:18. In some aspects of the disclosure, the *R*-selective transaminase polypeptides comprise an amino acid sequence having one or more of the following features: X57 is L; X81 is D; X82 is H; X85 is A, C, N, T, or V; X86 is S, F, or H; X95 is T; X112 is I; X147 is G; X153 is A, S, N, G, or T; X233 is S or T; X317 is L, X318 is G or R; and X319 is V.

[0134] In some aspects of the disclosure, the transaminase polypeptides capable of converting amino acceptor substrates to the chiral amine products of predominately the (*R*) configuration at a rate that is greater than the thermostable polypeptide of SEQ ID NO:18 can comprise an amino acid sequence having one of the following features: X85 is A; X85 is A and X153 is A; and X85 is S and X153 is A.

[0135] In some embodiments, the transaminase polypeptide capable of converting amino acceptor substrates to the chiral amine products of predominately the (*R*) configuration at a rate that is greater than the polypeptide of SEQ ID NO:18 comprises an amino acid sequence, corresponding to SEQ ID NO: 22, 30, 32, 34, 36, 40, 44, 48, 56, 58, 60, 62, 64, 66, 68, 70, 72, 76, 84, 88, 90, 92, 104, 108, 110, 120, 122, 124, 126, 128, 132, 160, or 176.


[0136] In some embodiments, the transaminase polypeptide capable of converting amino acceptor substrates to the chiral amine products of predominately the (*R*) configuration at a rate that is greater than the thermostable polypeptide of SEQ ID NO:18 comprises an amino

acid sequence corresponding to SEQ ID NO: 32, 36, 40, 56, 58, 60, 62, 64, 66, 122, 124, 126, 128, or 132

[0137] In some embodiments, the transaminase polypeptide capable of converting amino acceptor substrates to the chiral amine products of predominately the (*R*) configuration, at a rate that is greater than the thermostable polypeptide of SEQ ID NO:18 comprises an amino acid sequence corresponding to SEQ ID NO: 122, 124, 126, 128, 130, or 132.

[0138] The engineered transaminase polypeptides may be capable of binding and converting structurally diverse substrates. In some instances, the conversion rates for structurally different substrates for the naturally occurring *V fluvialis* enzyme may be insignificant. Accordingly, the engineered transaminase polypeptides may be characterized by increased conversion rates for substrates that are not converted at significant rates by the polypeptide of SEQ ID NO:2 or another engineered transaminase, such as SEQ ID NO:18.

[0139] In some embodiments, the transaminase polypeptides are characterized by the capability of stereoselectively converting the substrate 3,4-dihydronaphthalen-1(2*H*)-one, to the product (*S*)-1,2,3,4-tetrahydronaphthalen-1-amine, as below,



at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate 3,4-dihydronaphthalen-1(2*H*)-one include one or more of the following: X30, X44, X57, X82, X85, X113, X153, X166, X233, X314, X311, X317, X318, X319, X320; X407, X409, and X417.


[0140] In some aspects of the disclosure, the transaminase polypeptides capable of converting 3,4-dihydronaphthalen-1(2*H*)-one to (*S*)-1,2,3,4-tetrahydronaphthalen-1-amine at a rate greater than SEQ ID NO:2 or SEQ ID NO:18 comprises an amino acid sequence having at least one or more of the following features or set of features: X30 is A; X31 is A; X57 is I; X82 is H; X85 is V; X85 is C; X153 is S; X317 is M; X317 is Y; X417 is A; X319 is Q; X320 is A; X417 is I; X113 is H and X407 is S; X44 is A and X166 is S; X314 is V and X409 is G; X153 is S and X233 is S; X311 is V and X314 is T; X153 is S, X318 is R and X417 is E; X153 is C, X233 is L and X318 is R; X153 is S, X317 is L and X417 is C; X233 is L, X321 is L, and X417 is I; and X153 is C, X317 is Y and X319 is Q.

[0141] In some embodiments, the transaminase polypeptides capable of converting 3,4-dihydronaphthalen-1(2*H*)-one to (*S*)-1,2,3,4-tetrahydronaphthalen-1-amine at a rate greater than SEQ ID NO:2 and/or SEQ ID NO:18 comprises an amino acid sequence corresponding to SEQ ID NO: 38, 42, 44, 52, 56, 66, 74, 76, 78, 80, 82, 84, 94, 96, 98, 100, 104, 118, 160, 164, 172, 184, or 188.

[0142] In some embodiments, the transaminase polypeptide is capable of stereoselectively converting 3,4-dihydronaphthalen-1(2*H*)-one to (S)-1,2,3,4-tetrahydronaphthalen-1-amine at a rate that is at least 5-fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Polypeptides that are capable of the above reaction include, but are not limited to, a polypeptide comprising an amino acid sequence corresponding to SEQ ID NO: 44, 104, 164, 172, 184, or 188.

[0143] In some embodiments, the transaminase polypeptide is capable of stereoselectively converting 3,4-dihydronaphthalen-1(2*H*)-one to (S)-1,2,3,4-tetrahydronaphthalen-1-amine at a rate that is at least 10 fold greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Polypeptides that are capable of the above reaction include, but are not limited to, a polypeptide comprising an amino acid sequence corresponding to SEQ ID NO: 164 or 188.

[0144] In some embodiments, the transaminase polypeptide is capable of stereoselectively converting the substrate 1-phenylbutan-2-one to the product (S)-1-phenylbutan-2-amine, as below



at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate 1-phenylbutan-2-one include one or more of the following: X57; X81, X86; X153; X181; X228; X233; X317; X318; X319; X321; and X417.


[0145] In some aspects of the disclosure, the transaminase polypeptides capable of converting 1-phenylbutan-2-one, to the product (S)-1-phenylbutan-2-amine at a rate that is greater than the wild-type *Vibrio fluvialis* (SEQ ID NO:2) comprise an amino acid sequence having at least one of the following features or set of features: X81 is D and X86 is H; X86 is H and X153 is A; X86 is H, X153 is A and X417 is C; X86 is H, X153 is S and X417 is C; X86 is H, X153 is A, X228 is G and X417 is I; X86 is H, X153 is S, X181 is R and X417 is T; X228 is G, X317 is L and X417 is C; X57 is A, X153 is S and X318 is G; X57 is F, X86 is H, and X153 is Q; X57 is A, X153 is C and X321 is L; X86 is F, X318 is R and X417 is A; X228 is G, X318 is G and X417 is C; X153 is S, X318 is R and X417 is E; X153 is C, X233 is L and X318 is R; X86 is N, X228 is G and X317 is L; X153 is T, X228 is G and X321 is L; X153 is S, X317 is L, X417 is C; X57 is L, X86 is S and X153 is A; X153 is S, X228 is G and X417 is V; X153 is C, X317 is Y and X319 is Q; and X153 is T, X228 is G and X417 is A.

[0146] In some embodiments, the transaminase polypeptide capable of stereoselectively converting 1-phenylbutan-2-one, to the (S)-1-phenylbutan-2-amine comprises an amino acid sequence corresponding to SEQ ID NO: 22, 88, 90, 92, 106, 110, 138, 140, 142, 148, 150, 152, 160, 164, 168, 170, 172, 178, 182, 188, or 190.

[0147] In some embodiments, the transaminase polypeptide is capable of stereoselectively

converting 1-phenylbutan-2-one, to (S)-1-phenylbutan-2-amine at a rate that is at least 5-fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. In some embodiments, the transaminase polypeptide capable of stereoselectively converting 1-phenylbutan-2-one to the (S)-1-phenylbutan-2-amine at a rate at least 5 times greater than the polypeptide of SEQ ID NO:2 or SEQ ID NO:8 comprises an amino acid sequence corresponding to SEQ ID NO: 22, 88, 90, 92, 106 or 110.

[0148] In some embodiments, the transaminase polypeptide described herein is capable of stereoselectively converting the substrate 3,3-dimethylbutan-2-one, to the product (S)-3,3-dimethylbutan-2-amine, as below:




at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate 3,3-dimethylbutan-2-one include one or more of the following: X12, X30, X31, X44, X57, X81, X82, X85, X86, X95, X112, X113, X127, X153, X166, X181, X228, X233, X297, X311, X314, X317, X318, X319, X320, X321, X385, X407, X408, X409, X417, X434, and X438.

[0149] In some aspects of the disclosure, the transaminase polypeptides capable of converting 3,3-dimethylbutan-2-one, to the product (S)-3,3-dimethylbutan-2-amine at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18 comprises an amino acid sequence having at least one of the following features or set of features: X30 is A; X31 is A; X57 is L; X57 is I; X82 is H; X85 is V; X85 is S; X85 is T; X85 is A; X85 is N; X85 is C; X86 is S; X86 is N; X86 is F; X153 is T; X153 is N; X153 is G; X153 is S; X317 is M; X317 is Y; X319 is Q; X320 is A; X417 is A; X417 is I; X417 is C; X12 is G and X434 is V; X44 is A and X166 is S; X57 is I and X153 is S; X81 is D and X86 is H; X82 is H and X417 is F; X85 is S and X153 is A; X85 is A and X153 is A; X85 is A and X317 is L; X86 is H and X153 is A; X86 is S and X153 is S; X112 is I and X317 is L; X113 is H and X407 is S; X153 is S and X233 is S; X311 is V and X314 is T; X314 is V and X409 is G; X318 is G and X408 is A; X57 is L, X417 is C and X438 is L; X57 is F, X127 is L and X417 is C; X57 is S, X233 is L and X417 is V; X57 is A, X153 is S and X318 is G; X57 is F, X86 is H and X153 is Q; X57 is I, X86 is F and X320 is A; X57 is F, X86 is F and X153 is Q; X57 is C, X86 is S and X417 is T; X57 is F, X318 is F and X417 is S; X57 is L, X86 is S and X153 is A; X57 is L, X86 is F and X318 is F; X57 is F, X318 is G and X417 is I; X86 is H, X153 is S and X417 is C; X86 is H, X153 is A and X417 is C; X86 is F, X153 is C and X297 is A; X86 is H, X233 is L and X417 is A; X86 is N, X228 is G and X317 is L; X95 is T, X153 is A, and X417 is C; X113 is C, X385 is R and X417 is C; X153 is A, X317 is L and X318 is G; X153 is A, X233 is T and X417 is C; X153 is C, X233 is L and X318 is R; X153 is T, X228 is G, and X321 is L; X153 is S, X317 is L, and X417 is C; X153 is T, X319 is V and X417 is I; X153 is S, X228 is G and X417 is V; X153 is C, X317 is Y, and X319 is Q; X153 is T, X228 is G and X417 is A; X233 is L, X321 is L and X417 is I; X228 is G, X318 is G and X417 is C; X317 is L, X318 is R and X417 is T; X86 is H, X153 is A, X228 is G and X417 is I; and X86 is H, X153 is S, X181 is R and X417 is T.

[0150] In some embodiments, the transaminase polypeptide capable of stereoselectively converting 3,3-dimethylbutan-2-one to (S)-3,3-dimethylbutan-2-amine comprises an amino acid sequence corresponding to SEQ ID NO: 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 134, 140, 142, 144, 146, 152, 154, 156, 158, 162, 164, 168, 170, 172, 174, 176, 178, 180, 182, 184, 186, 188, 190, or 192.

[0151] In some embodiments, the transaminase polypeptide is capable of stereoselectively converting 3,3-dimethylbutan-2-one to (S)-3,3-dimethylbutan-2-amine, at a rate that is at least 1.1 fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. In some embodiments, the transaminase polypeptide capable of stereoselectively converting 3,3-dimethylbutan-2-one to (S)-3,3-dimethylbutan-2-amine at a rate that is at least 1.1 fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18 comprises an amino acid sequence corresponding to SEQ ID NO: 30, 34, 38, 44, 52, 54, 70, 72, 74, 76, 80, 82, 96, 98, 100, 104, 118, 120, 164, 178, or 188.

[0152] In some embodiments, the transaminase polypeptide described herein is capable of stereoselectively converting the substrate octan-2-one, to the product (S)-octan-2-amine, as below:



at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate octan-2-one include one or more of the following: X12, X30, X31, X44, X57, X81, X82, X85, X86, X95, X112, X113, X127, X153, X166, X181, X228, X233, X297, X311, X314, X317, X318, X319, X320, X321, X385, X407, X408, X409, X417, X434, and X438.

[0153] In some aspects of the disclosure, the transaminase polypeptides capable of converting octan-2-one, to the product (S)-octan-2-amine comprises an amino acid sequence having at least one of the following features or set of features: X30 is A; X31 is A; X57 is I; X57 is L; X82 is H; X85 is V; X85 is S; X85 is T; X85 is A; X85 is N; X85 is C; X86 is S; X86 is N; X86 is F; X153 is S; X153 is T; X153 is N; X153 is G; X317 is M; X317 is Y; X417 is A; X319 is Q; X320 is A; X417 is I; X417 is C; X12 is G and X434 is V; X44 is A and X166 is S; X57 is I and X153 is S; X81 is D and X86 is H; X82 is H and X417 is F; X85 is A and X317 is L; X85 is S and X153 is A; X85 is A and X153 is A; X86 is H and X153 is A; X86 is S and X153 is S; X112 is I and X317 is L; X113 is H and X407 is S; X153 is S and X233 is S; X311 is V and X314 is T; X314 is V and X409 is G; X318 is G and X408 is A; X57 is L, X417 is C and X438 is L; X57 is F, X127 is L and X417 is C; X57 is S, X233 is L and X417 is V; X57 is S, X86 is G and X417 is C; X57 is A, X153 is S and X318 is G; X57 is F, X86 is H and X153 is Q; X57 is I, X86 is F and X320 is A; X57 is F, X86 is F and X153 is Q; X57 is A, X153 is C and X321 is L; X57 is C, X86 is S and X417 is T; X57 is C, X86 is A and X317 is L; X57 is F, X318 is F and X417 is S; X57 is L, X86 is S and


X153 is A; X57 is L, X86 is F and X318 is F; X57 is F, X318 is G and X417 is I; X86 is F, X318 is R and X417 is A; X86 is F, X153 is C and X297 is A; X86 is S, X153 is T and X297 is A; X86 is H, X233 is L and X417 is A; X86 is N, X228 is G, and X317 is L; X86 is H, X153 is A and X417 is C; X86 is H, X153 is S and X417 is C; X95 is T, X153 is A and X417 is C; X113 is C, X385 is R and X417 is C; X153 is A, X233 is T and X417 is C; X153 is S, X318 is R and X417 is E; X153 is C, X233 is L and X318 is R; X153 is S, X228 is G and X417 is V; X153 is C, X317 is Y and X319 is Q; X153 is T, X228 is G, and X417 is A; X153 is T, X319 is V and X417 is I; X153 is T, X228 is G and X321 is L; X153 is S, X317 is L and X417 is C; X153 is A, X317 is L and X318 is G; X228 is G, X318 is G and X417 is C; X228 is G, X317 is L and X417 is C; X233 is L, X321 is L and X417 is I; X317 is L, X318 is R and X417 is T; X86 is H, X153 is A, X228 is G and X417 is I; and X86 is H, X153 is S, X181 is R and X417 is T.

[0154] In some embodiments, the transaminase polypeptide capable of stereoselectively converting octan-2-one to (S)-octan-2-amine comprises an amino acid sequence corresponding to SEQ ID NO: 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 134, 136, 138, 140, 142, 144, 146, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 178, 180, 182, 184, 186, 188, 190, or 192.

[0155] In some embodiments, the transaminase polypeptide is capable of stereoselectively converting octan-2-one to (S)-octan-2-amine at a rate that is at least 1.1 fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. In some embodiments, the transaminase polypeptide capable of stereoselectively converting octan-2-one to (S)-octan-2-amine at a rate that is at least 1.1 fold greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18 comprises an amino acid sequence corresponding to SEQ ID NO: 20, 22, 24, 26, 28, 30, 34, 38, 42, 44, 46, 48, 52, 54, 70, 72, 74, 76, 80, 82, 84, 86, 92, 96, 98, 102, 104, 108, 110, 112, 114, 116, 134, 136, 138, 140, 142, 144, 146, 152, 154, 156, 164, 166, 168, 170, 172, 174, 178, 180, 182, 184, 188, 190, or 192.

[0156] In some embodiments, the transaminase polypeptide is capable of stereoselectively converting octan-2-one to (S)-octan-2-amine at a rate that is at least 5 fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Exemplary polypeptides that are capable of stereoselectively converting octan-2-one (S)-octan-2-amine at a rate that is at least 5 fold greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18 comprises an amino acid sequence corresponding to SEQ ID NO: 22 or 192.

[0157] In some embodiments, the transaminase polypeptides described herein are capable of stereoselectively converting the substrate 1-(4-bromophenyl)ethanone, to the product (S)-1-(4-bromophenyl)ethanamine, as below



at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate 1-(4-bromophenyl)ethanone include one or more of the following: X12, X30, X31, X44, X57, X81, X82, X85, X86, X95, X112, X113, X127, X153, X153, X166, X181, X228, X233, X297, X311, X314, X317, X318, X319, X320, X321, X385, X407, X408, X409, X417, X434, and X438.

[0158] In some aspects of the disclosure, the transaminase polypeptide capable of stereoselectively converting 1-(4-bromophenyl)ethanone, to (S)-1-(4-bromophenyl)ethanamine comprises an amino acid sequence having at least one of the following features or set of features: X30 is A; X57 is L; X57 is I; X82 is H; X85 is V; X85 is T; X85 is A; X85 is N; X86 is N; X86 is F; X86 is S; X153 is S; X153 is T; X153 is N; X153 is G; X317 is M; X317 is Y; X319 is Q; X320 is A; X417 is A; X417 is I; X417 is C; X12 is G and X434 is V; X44 is A and X166 is S; X57 is I and X153 is S; X81 is D and X86 is H; X82 is H and X417 is F; X85 is S and X153 is A; X85 is A and X153 is A; X86 is S and X153 is S; X86 is H and X153 is A; X112 is I and X317 is L; X113 is H and X407 is S; X153 is S and X233 is S; X314 is V and X409 is G; X318 is G and X408 is A; X31 is A, X311 is V and X314 is T; X57 is L, X417 is C and X438 is L; X57 is F, X127 is L and X417 is C; X57 is I, X86 is F and X320 is A; X57 is S, X86 is G and X417 is C; X57 is C, X86 is S and X417 is T; X57 is C, X86 is A and X317 is L; X57 is F, X318 is F and X417 is S; X57 is L, X86 is S and X153 is A; X57 is F, X318 is G and X417 is I; X86 is H, X153 is A and X417 is C; X86 is H, X153 is S and X417 is C; X86 is F, X318 is R and X417 is A; X86 is F, X153 is C and X297 is A; X86 is S, X153 is T and X297 is I; X86 is N, X228 is G and X317 is L; X86 is H, X233 is L and X417 is A; X95 is T, X153 is A and X417 is C; X113 is C, X385 is R and X417 is C; X153 is A, X233 is T and X417 is C; X153 is A, X317 is L and X318 is G; X153 is S, X318 is R and X417 is E; X153 is T, X228 is G and X321 is L; X153 is S, X317 is L and X417 is C; X153 is C, X233 is L and X318 is R; X153 is S, X228 is G and X417 is V; X153 is T, X319 is V and X417 is I; X153 is C, X317 is Y and X319 is Q; X153 is T, X228 is G and X417 is A; X228 is G, X317 is L and X417 is C; X228 is G, X318 is G and X417 is C; X233 is L, X321 is L and X417 is I; X317 is L, X318 is R and X417 is T; X86 is H, X153 is A, X228 is G and X417 is I; and X86 is H, X153 is S, X181 is R and X417 is T.


[0159] In some embodiments, the transaminase polypeptide capable of converting 1-(4-bromophenyl)ethanone to (S)-1-(4-bromophenyl)ethanamine comprises an amino acid sequence corresponding to SEQ ID NO: 20, 22, 24, 26, 28, 30, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 60, 62, 64, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 136, 138, 144, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 176, 178, 182, 184, 186, 188, 190, or 192.

[0160] In some embodiments, the transaminase polypeptide is capable of converting 1-(4-bromophenyl)ethanone to (S)-1-(4-bromophenyl)ethanamine at a rate that is at least 1.1 fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. In some embodiments, the transaminase polypeptide capable of converting 1-(4-bromophenyl)ethanone to (S)-1-(4-bromophenyl)ethanamine at a rate that is at least 1.1 fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18 comprises an amino acid sequence corresponding to SEQ ID NO: 20, 22, 24, 26, 28, 34, 38, 42, 44, 46, 48, 50, 52, 54, 68, 70, 72, 74, 76, 78, 80, 82, 84,

86, 88, 92, 94, 96, 98, 100, 102, 104, 108, 110, 112, 114, 116, 118, 136, 150, 152, 154, 156, 160, 164, 166, 170, 172, 174, 182, 184, 186, 188, or 192.

[0161] In some embodiments, the transaminase polypeptide is capable of converting 1-(4-bromophenyl)ethanone to the product (S)-1-(4-bromophenyl)ethanamine at a rate that is at least 5 fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. In some embodiments, the transaminase polypeptide capable of converting 1-(4-bromophenyl)ethanone to (S)-1-(4-bromophenyl)ethanamine at a rate that is at least 5-fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18 comprises an amino acid sequence corresponding to SEQ ID NO: 84, 102, 108, 120, 122, 124, 128, 132, 172, 184, or 186.

[0162] In some embodiments, the transaminase polypeptide described herein are capable of stereoselectively converting the substrate 4-phenylbutan-2-one, to the product (R)-4-phenylbutan-2-amine, as below:



at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate 4-phenylbutan-2-one include one or more of the following: X30, X31, X44, X57, X81, X82, X85, X86, X113, X153, X166, X181, X228, X233, X297, X317, X318, X319, X320, X321, X385, X407, X408, X417, and X438.

[0163] In some aspects of the disclosure, the transaminase polypeptide capable of stereoselectively converting 4-phenylbutan-2-one to (R)-4-phenylbutan-2-amine comprises an amino acid sequence having at least one of the following features or set of features: X30 is A; X31 is A; X57 is L; X57 is I; X82 is H; X85 is V; X85 is S; X85 is T; X85 is A; X86 is S; X86 is N; X86 is F; X153 is S; X153 is T; X153 is N; X153 is G; X319 is Q; X317 is M; X317 is Y; X320 is A; X417 is A; X417 is I; X417 is C; X44 is A and X166 is S; X57 is I and X153 is S; X81 is D and X86 is H; X82 is H and X417 is F; X85 is A and X153 is A; X85 is A and X317 is L; X85 is S and X153 is A; X86 is S and X153 is S; X86 is H and X153 is A; X113 is H and X407 is S; X318 is G and X408 is A; X153 is S and X233 is S; X57 is A, X153 is S and X318 is G; X57 is F, X86 is H and X153 is Q; X57 is I, X86 is F and X320 is A; X57 is F, X86 is F, and X153 is Q; X57 is A, X153 is C and X321 is L; X57 is L, X417 is C and X438 is L; X57 is S, X233 is L and X417 is V; X57 is C, X86 is S, and X417 is T; X57 is S, X86 is G and X417 is C; X57 is C; X86 is A and X317 is L; X57 is F, X318 is F and X417 is S; X57 is L, X86 is S and X153 is A; X57 is L, X86 is F and X318 is F; X57 is F, X318 is G and X417 is I; X86 is H, X153 is S and X417 is C; X86 is H, X153 is A and X417 is C; X86 is F, X318 is R and X417 is A; X86 is F, X153 is C and X297 is A; X86 is S, X153 is T and X297 is A; X86 is H, X233 is L and X417 is A; X86 is N, X228 is G and X317 is L; X113 is C, X385 is R and X417 is C; X153 is A, X233 is T and X417 is C X153 is A, X317 is L and X318 is G; X153 is S, X318 is R and X417 is E; X153 is C, X233 is L and X318 is R; X153 is T, X228 is G and X321 is L; X153 is S, X317 is L and X417 is C; X153 is S, X228


is G and X417 is V; X153 is C, X317 is Y and X319 is Q; X153 is T, X228 is G and X417 is A; X228 is G, X317 is L and X417 is C; X228 is G, X318 is G and X417 is C; X233 is L, X321 is L and X417 is I; X317 is L, X318 is R and X417 is T; X86 is H, X153 is A, X228 is G and X417 is I; and X86 is H, X153 is S, X181 is R and X417 is T.

[0164] In some embodiments, the transaminase polypeptide capable of stereoselectively converting 4-phenylbutan-2-one to (*R*)-4-phenylbutan-2-amine comprises an amino acid sequence corresponding to SEQ ID NO: 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 178, 182, 184, 186, 188, 190, or 192.

[0165] In some embodiments, the transaminase polypeptide is capable of stereoselectively converting 4-phenylbutan-2-one to (*R*)-4-phenylbutan-2-amine at a rate that is at least 1.1 fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. In some embodiments, the transaminase polypeptide capable of stereoselectively converting 4-phenylbutan-2-one to (*R*)-4-phenylbutan-2-amine at a rate that is at least 1.1 fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18 comprises an amino acid sequence corresponding to SEQ ID NO: 20, 22, 24, 26, 28, 30, 34, 42, 44, 46, 52, 68, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 102, 104, 110, 112, 114, 134, 136, 138, 140, 142, 144, 146, 148, 150, 154, 156, 158, 162, 164, 166, 168, 172, 174, 178, 182, 184, 186, 188, 190, or 192.

[0166] In some embodiments, the transaminase polypeptide is capable of stereoselectively converting 4-phenylbutan-2-one, to (*R*)-4-phenylbutan-2-amine at a rate that is at least 5-fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. In some embodiments, the transaminase polypeptide capable of stereoselectively converting 4-phenylbutan-2-one to (*R*)-4-phenylbutan-2-amine at a rate that is at least 5-fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18, comprises an amino acid sequence corresponding to SEQ ID NO: 154, 172, or 178.

[0167] In some embodiments, the transaminase polypeptides described herein are capable of stereoselectively converting the substrate ethyl 3-oxobutanoate, to the product (*R*)-ethyl 3-aminobutanoate, as below



at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate ethyl 3-oxobutanoate include one or more of the following: X57, X85, X86, X147, X153, X233, X317, X318, and X417.


[0168] In some aspects of the disclosure, the transaminase polypeptide capable of stereoselectively converting ethyl 3-oxobutanoate to (*R*)-ethyl 3-aminobutanoate comprises an amino acid sequence having at least one of the following features or set of features: X85 is V;

X85 is S; X85 is T; X85 is A; X85 is N; X85 is C; X153 is T; X153 is N; X153 is G; X153 is S; X417 is C; X85 is S and X153 is A; X85 is A and X153 is A; X85 is A and X317 is L; X86 is H and X153 is A; X85 is A, X153 is S and X417 is S; X85 is S, X153 is A and X233 is T; X85 is A, X147 is G and X153 is A; X86 is H, X153 is A and X417 is C; X86 is H, X153 is S and X417 is C; X153 is A, X317 is L and X318 is G; X57 is I, X85 is A, X86 is H and X417 is C; and X57 is L, X86 is F, X153 is S, X233 is T and X417 is T.

[0169] In some embodiments, the transaminase polypeptide capable of stereoselectively converting ethyl 3-oxobutanoate to (*R*)-ethyl 3-aminobutanoate comprises an amino acid sequence corresponding to SEQ ID NO: 32, 34, 36, 40, 44, 56, 58, 60, 62, 64, 66, 68, 70, 72, 86, 88, 90, 92, 122, 126, 128, 130, or 132.

[0170] In some embodiments, the transaminase polypeptide is capable of stereoselectively converting ethyl 3-oxobutanoate to (*R*)-ethyl 3-aminobutanoate at a rate that is at least 5-fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Polypeptides that are capable of the above reaction include, but are not limited to, a polypeptide comprising an amino acid sequence corresponding to SEQ ID NO: 32, 36, 40, 56, 58, 60, 62, 64, 66, 126, 128, or 130.

[0171] In some embodiments, the transaminase polypeptides described herein are capable of stereoselectively converting the substrate ethyl 3-oxobutanoate, to the product (*S*)-ethyl 3-aminobutanoate, as below:



at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate ethyl 3-oxobutanoate include one or more of the following: X12, X30, X31, X44, X57, X81, X82, X85, X86, X112, X113, X127, X153, X166, X228, X233, X297, X311, X314, X317, X318, X319, X320, X321, X385, X407, X408, X409, X417, X434, and X438.

[0172] In some aspects of the disclosure, the transaminase polypeptide capable of converting ethyl 3-oxobutanoate to (*S*)-ethyl 3-aminobutanoate comprises an amino acid sequence having at least one of the following features or set of features: X30 is A; X31 is A; X57 is I; X57 is L; X82 is H; X86 is S; X86 is N; X86 is F; X317 is M; X417 is A; X319 is Q; X320 is A; X12 is G and X434 is V; X44 is A and X166 is S; X57 is I and X153 is S; X81 is D and X86 is H; X82 is H and X417 is F; X85 is S and X153 is S; X86 is S and X153 is S; X112 is I and X317 is L; X113 is H and X407 is S; X318 is G and X408 is A; X314 is V and X409 is G; X311 is V and X314 is T; X57 is L, X417 is C and X438 is L; X57 is F, X127 is L and X417 is C; X57 is S, X233 is L and X417 is V; X57 is S, X86 is G and X417 is C; X57 is A, X153 is S and X318 is G; X57 is F, X86 is H and X153 is Q; X57 is I, X86 is F and X320 is A; X57 is F, X86 is F and X153 is Q; X57 is A, X153 is C and X321 is L; X57 is C, X86 is A and X317 is L; X57 is F, X318 is F and X417 is S; X57 is L, X86 is S and X153 is A; X57 is L, X86 is F and X318 is F; X57 is C, X86 is S and


X417 is T; X57 is F, X318 is G and X417 is I; X86 is F, X318 is R and X417 is A; X86 is N, X228 is G and X317 is L; X86 is F, X153 is C and X297 is A; X86 is S, X153 is T and X297 is A; X86 is H, X233 is L and X417 is A; X113 is C, X385 is R and X417 is C; X153 is A, X233 is T and X417 is C; X153 is C, X233 is L and X318 is R; X153 is T, X228 is G and X321 is L; X153 is S, X317 is L and X417 is C; X153 is S, X228 is G and X417 is V; X153 is C, X317 is Y and X319 is Q; X153 is T, X228 is G and X417 is A; X228 is G, X317 is L and X417 is C; X228 is G, X318 is G and X417 is C; X233 is L, X321 is L and X417 is I; X317 is L, X318 is R and X417 is T; and X86 is H, X153 is A, X228 is G and X417 is I.

[0173] In some embodiments, the transaminase polypeptide capable of stereoselectively converting ethyl 3-oxobutanoate to (S)-ethyl 3-aminobutanoate comprises an amino acid sequence corresponding to SEQ ID NO: 20, 22, 24, 26, 28, 30, 38, 42, 46, 48, 50, 52, 54, 74, 78, 80, 82, 94, 96, 98, 100, 102, 106, 112, 114, 116, 118, 120, 124, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 162, 164, 166, 168, 170, 172, 174, 178, 180, 182, 184, 186, 188, 190 or 192.

[0174] In some embodiments, the transaminase polypeptide is capable of stereoselectively converting ethyl 3-oxobutanoate to (S)-ethyl 3-aminobutanoate at a rate that is at least 1.1 fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. In some embodiments, the transaminase polypeptide capable of converting ethyl 3-oxobutanoate to (S)-ethyl 3-aminobutanoate at a rate that is at least 1.1 fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18 comprises an amino acid sequence corresponding to SEQ ID NO: 22, 26, 28, 38, 46, 56, 60, 64, 78, 84, 86, 90, 100, 102, 106, 112, 118, 136, 144, 146, 154, 156, 158, 160, 164, 166, 168, 170, 172, 174, 178, 182, 188, or 190.

[0175] In some embodiments, the transaminase polypeptide is capable of stereoselectively converting ethyl 3-oxobutanoate to (S)-ethyl 3-aminobutanoate at a rate that is at least 5-fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. In some embodiments, the transaminase polypeptide capable of stereoselectively converting ethyl 3-oxobutanoate to (S)-ethyl 3-aminobutanoate at a rate that is at least 5-fold or greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18 comprises an amino acid sequence corresponding to SEQ ID NO: 22, 26, 28, 38, 46, 56, 60, 64, 78, 84, 86, 90, 100, 102, 106, 112, 118, 136, 144, 154, 166, 170, 174, 178, or 188.

[0176] In some embodiments, the transaminase polypeptides described herein are capable of stereoselectively converting the substrate 1-(6-methoxynaphthalen-2-yl)ethanone, to the product (S)-1-(6-methoxynaphthalen-2-yl)ethanamine, as below:




at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate 1-(6-methoxynaphthalen-2-yl)ethanone include one or more of the following: X57, X86, X153, X233, X317, X318, X320,

X321, X417, and X438.

[0177] In some aspects of the disclosure, the transaminase polypeptide capable of converting 1-(6-methoxynaphthalen-2-yl)ethanone to (S)-1-(6-methoxynaphthalen-2-yl)ethanamine comprises an amino acid sequence having at least one of the following features or set of features: X57 is L; X86 is F or N; X417 is C, I or A; X57 is L, X86 is F, X153 is S, X233 is T and X417 is T; X57 is C, X86 is S and X417 is T; X57 is S, X233 is L, and X417 is V; X57 is A, X153 is S and X318 is G; X57 is A, X153 is C, and X321 is L; X57 is L, X86 is S, and X153 is A; X57 is S, X86 is G and X417 is C; X57 is L, X86 is F, X318 is F; X57 is F, X86 is F, and X153 is Q; X57 is I; X86 is F and X320 is A; X57 is C, X86 is A and X317 is L; X57 is L, X417 is C and X438 is L; X86 is F, X318 is R and X417 is A; X57 is F, X86 is H and X153 is Q; X57 is F, X318 is F and X417 is S; X86 is H, X153 is S, X181 is R and X417 is T; X153 is S, X317 is L and X417 is C; X153 is T, X228 is G and X417 is A; X57 is F, X127 is L and X417 is C; X228 is G, X317 is L and X417 is C; X233 is L, X321 is L and X417 is I; X228 is G, X318 is G and X417 is C; X86 is H, X153 is S and X417 is C; X86 is H, X233 is L and X417 is A; X86 is S and X153 is S; X86 is F, X153 is C and X297 is A; X153 is S, X228 is G and X417 is V; X57 is F, X318 is G and X417 is I; X153 is T, X228 is G and X321 is L; X317 is L, X318 is R and X417 is T; X153 is C, X317 is Y, and X319 is Q; X57 is I and X153 is S; X86 is H, X153 is A and X417 is C; X153 is A, X233 is T and X417 is C; X95 is T, X153 is A and X417 is C; X86 is N, X228 is G and X317 is L; and X86 is S, X153 is T, X297 is A.

[0178] In some embodiments, the transaminase polypeptide capable of stereoselectively converting 1-(6-methoxynaphthalen-2-yl)ethanone to (S)-1-(6-methoxynaphthalen-2-yl)ethanamine comprises an amino acid sequence corresponding to SEQ ID NO: 24, 26, 28, 46, 48, 78, 84, 86, 88, 92, 102, 108, 110, 114, 116, 126, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 162, 166, 168, 170, 172, 174, 178, 180, 182, 184, 186, 188, 190, or 192.

[0179] In some embodiments, the transaminase polypeptides described herein are capable of stereoselectively converting the substrate 1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanone, to the product (S)-1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanamine, as below:




at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate 1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanone include one or more of the following: X57, X86, X153, X228, X233, X297, X318, X320, X417, and X438.

[0180] In some aspects of the disclosure, the transaminase polypeptide capable of converting 1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanone, to the product (S)-1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanamine comprises an amino acid sequence having at least one of the

following features or set of features: X31 is A; X57 is C, I or L; X86 is S or F; X153 is G or S; X417 is T or I; X57 is F, X86 is F, and X153 is Q; X57 is I and X153 is S; X57 is L, X86 is F, X153 is S, X233 is T and X417 is T; X86 is F, X153 is C, and X297 is A; X57 is L, X417 is C, and X438 is L; X57 is S, X233 is L, and X417 is V; X57 is L, X86 is S, X153 is A; X57 is S, X86 is G, and X417 is C; X57 is A, X153 is S, X318 is G; X57 is I, X86 is F and X320 is A; X57 is L, X86 is F, X318 is F; X153 is S, X228 is G, and X417 is V; X57 is C, X86 is A, and X317 is L; X153 is S and X233 is S; X57 is A, X153 is C and X321 is L; X86 is S and X153 is S; X86 is H, X153 is S, X181 is R, and X417 is T; X21 is N, X45 is N, X177 is V, X208 is I, X211 is R, X324 is S, and X391 is T; X86 is H, X153 is A, X228 is G, and X417 is I; and X153 is C, X233 is L and X318 is R.

[0181] In some embodiments, the transaminase polypeptide capable of stereoselectively converting 1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanone, to the product (S)-1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanamine comprises an amino acid sequence corresponding to SEQ ID NO: 16, 26, 28, 42, 44, 46, 48, 72, 84, 98, 104, 106, 110, 114, 126, 134, 136, 140, 144, 146, 148, 152, 156, 164, 166, 178, 180, or 182.

[0182] In some embodiments, the transaminase polypeptides described herein are capable of stereoselectively converting the substrate 1-(4-phenoxyphenyl)ethanone, to the product (S)-1-(4-phenoxyphenyl)ethanamine, as below:




at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate 1-(4-phenoxyphenyl)ethanone include one or more of the following: X57, X86, X153, X181, X233, X297, X318, X417, and X438.

[0183] In some aspects of the disclosure, the transaminase polypeptide capable of converting 1-(4-phenoxyphenyl)ethanone, to the product (S)-1-(4-phenoxyphenyl)ethanamine comprises an amino acid sequence having at least one of the following features or set of features: X31 is A; X86 is S or F; X57 is L or I; X153 is S or G; X417 is A or I; X57 is I and X153 is S; X86 is S and X153 is S; X153 is S and X233 is S; X86 is H and X153 is A; X86 is H; X153 is A and X417 is C; X95 is T, X153 is A and X417 is C; X86 is H, X153 is S and X417 is C; X57 is L, X417 is C and X438 is L; X57 is S, X233 is L and X417 is V; X57 is S, X86 is G and X417 is C; X57 is A, X153 is S, X318 is G; X57 is F, X86 is H and X153 is Q; X57 is I, X86 is F and X320 is A; X57 is F, X86 is F and X153 is Q; X57 is C, X86 is S and X417 is T; X86 is F, X153 is C and X297 is A; X86 is H, X233 is L and X417 is A; X153 is C, X233 is L and X318 is R; X57 is C, X86 is A and X317 is L; X57 is L, X86 is S and X153 is A; X57 is L, X86 is F and X318 is F; X153 is S, X228 is G and X417 is V; X153 is T, X228 is G and X417 is A; X86 is H, X153 is A, X228 is G and X417 is I; X86 is H, X153 is S, X181 is R, and X417 is T; and X57 is L, X86 is F, X153 is S, X233 is T and X417 is T.

[0184] In some embodiments, the transaminase polypeptide capable of stereoselectively converting 1-(4-phenoxyphenyl)ethanone to the product (S)-1-(4-phenoxyphenyl)ethanamine comprises an amino acid sequence corresponding to SEQ ID NO: 16, 20, 26, 28, 42, 44, 46, 48, 72, 78, 84, 88, 90, 92, 98, 104, 106, 108, 110, 114, 126, 134, 136, 140, 142, 144, 146, 154, 156, 162, 164, 166, 178, 180, 182, or 190.

[0185] In some embodiments, the transaminase polypeptides described herein are capable of stereoselectively converting the substrate (*R*)-4-oxotetrahydro-2*H*-pyran-3-yl-benzoate to the product (3*S*,4*S*)-3-aminotetrahydro-2*H*-pyran-3-yl benzoate, as below:




at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate (*R*)-4-oxotetrahydro-2*H*-pyran-3-yl-benzoate include one or more of the following: X57, X86, X153, X233, X317, X318, X320, X321, and X417.

[0186] In some aspects of the disclosure, the transaminase polypeptide capable of converting (*R*)-4-oxotetrahydro-2*H*-pyran-3-yl-benzoate to the product (3*S*,4*S*)-3-aminotetrahydro-2*H*-pyran-3-yl benzoate comprises an amino acid sequence having at least one of the following features or set of features: X57 is I and X153 is S; X57 is A, X153 is S and X318 is G; X57 is A, X153 is C and X321 is L; X57 is L, X86 is S and X153 is A; X57 is I, X86 is F, X320 is A; X57 is S, X86 is G, and X417 is C; X57 is C, X86 is S and X417 is T; X57 is C, X86 is A and X317 is L; X57 is L, X86 is F, X153 is S. X233 is T and X417 is T.

[0187] In some embodiments, the transaminase polypeptide capable of stereoselectively converting (*R*)-4-oxotetrahydro-2*H*-pyran-3-yl-benzoate, to the product (3*S*,4*S*)-3-benzyloxytetrahydro-2*H*-pyran-4-amine comprises an amino acid sequence corresponding to SEQ ID NO: 46, 126, 136, 140, 144, 148, 154, 166, or 178.

[0188] In some embodiments, the transaminase polypeptides described herein are capable of stereoselectively converting the substrate (*R*)-3-(benzyloxy)dihydro-2*H*-pyran-4(3*H*)-one, to the product (3*S*,4*S*)-3-(benzyloxy)tetrahydro-2*H*-pyran-4-amine, as below:



at a rate that is greater than the polypeptide of SEQ ID NO:2 and/or SEQ ID NO:18. Residue positions associated with increase in activity on substrate (*R*)-3-(benzyloxy)dihydro-2*H*-pyran-4(3*H*)-one include one or more of the following: X57, X86, X153, X233, X317, X318, X320, X321, and X417.

[0189] In some aspects of the disclosure, the transaminase polypeptide capable of converting (*R*)-3-(benzyloxy)dihydro-2*H*-pyran-4(3*H*)-one, to (3*S*,4*S*)-3-(benzyloxy)tetrahydro-2*H*-pyran-4-amine comprises an amino acid sequence having at least one of the following features or set of features: X30 is A; X31 is A; X57 is I or L; X82 is H; X85 is S, T, A; X86 is S, F; X153 is N, G, S, T; X317 is M or Y; X319 is Q; X320 is A; X417 is A, I or C; X57 is I and X153 is S; X85 is S and X153 is S; X81 is D and X86 is H; X86 is H and X153 is A; X86 is S and X153 is S; X85 is S and X153 is A; X85 is A and X153 is A; X314 is V and X409 is G; X311 is V and X314 is T; X82 is H and X417 is F; X85 is A and X317 is L; X318 is G and X408 is A; X153 is S and X233 is S; X113 is H and X407 is S; X12 is G and X434 is V; X112 is I and X317 is L; X44 is A and X166 is S; X86 is H, X153 is S and X417 is C; X57 is F, X127 is L and X417 is C; X57 is S, X86 is G and X417 is C; X228 is G, X317 is L, X417 is C; X57 is F, X86 is H and X153 is Q; X86 is F, X318 is R and X417 is A; X86 is F, X153 is C and X297 is A; X86 is N, X228 is G and X317 is L; X57 is L, X86 is F and X318 is F; X153 is S, X228 is G and X417 is V; X57 is F, X86 is F and X153 is Q; X57 is L, X417 is C and X438 is L; X57 is A, X153 is C and X321 is L; X57 is S, X233 is L and X417 is V; X86 is S, X153 is T and X297 is A; X57 is F, X318 is F and X417 is S; X153 is T, X319 is V and X417 is I; X57 is S, X233 is L and X417 is V; X57 is I, X86 is F and X320 is A; X57 is F, X318 is G and X417 is I; X153 is C, X317 is Y and X319 is Q; X57 is L, X86 is S and X153 is A; X233 is L, X321 is L and X417 is I; X95 is T, X153 is A; and X417 is C; X86 is H, X153 is A and X417 is C; X228 is G, X318 is G and X417 is C; X85 is S, X153 is A and X233 is T; X57 is A, X153 is S and X318 is G; X153 is S, X318 is R and X417 is E; X153 is C, X233 is L and X318 is R; X153 is A; X233 is T and X417 is C; X85 is A, X147 is G and X153 is A; X153 is S, X317 is L and X417 is C; X317 is L, X318 is R and X417 is T; X153 is T, X228 is G and X417 is A; X57 is C, X86 is A and X317 is L; X153 is A, X317 is L and X318 is G; X57 is C, X86 is S and X417 is T; X86 is H, X233 is L and X417 is A; X153 is T, X228 is G, X321 is L; X113 is C, X385 is R and X417 is C; X86 is H, X153 is S, X181 is R and X417 is T; X86 is H, X153 is A, X228 is G and X417 is I; X57 is I, X85 is A, X86 is H and X417 is C; and X57 is L, X86 is F, X153 is S, X233 is T and X417 is T.

[0190] In some embodiments, the transaminase polypeptide capable of stereoselectively converting (*R*)-3-(benzyloxy)dihydro-2*H*-pyran-4(3*H*)-one to (3*S*,4*S*)-3-(benzyloxy)tetrahydro-2*H*-pyran-4-amine comprises an amino acid sequence corresponding to SEQ ID NO: 20, 22, 26, 28, 30, 32, 34, 36, 40, 42, 44, 46, 48, 50, 52, 58, 60, 68, 70, 72, 78, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 114, 116, 118, 122, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172; 174, 176, 178, 180, 182, 184, 186, 188, 190, or 192.

[0191] The expression of the transaminase polypeptides of the invention may be improved as compared to the expression of the polypeptide of SEQ ID NO:2 encoded by the polynucleotide of SEQ ID NO:1 in a host cell, particularly an *E. coli* host cell. In aspects of the disclosure, the

transaminase polypeptide can have an amino acid sequence having at least one or more of the following features: X4 is R; X6 is R or I or N; X9 is T or G; X6 is R and X133 is T; X12 is K and X302 is K; and X9 is T, X86 is Y and X294 is V. In some aspects of the disclosure, the transaminase polypeptide with increased expression can be encoded by a polynucleotide sequence having one or more of the following residue differences as compared to the polynucleotide sequence of SEQ ID NO:1 or SEQ ID NO:17: X852 is T and X861 is A; X18 is T and X286 is T; X891 is C; X12 is A and X15 is A and X18 is T; X309 is T; and X561 is C.

[0192] Exemplary transaminase polypeptides that are capable of increased expression include, but are not limited to, polypeptides comprising an amino acid sequence corresponding to SEQ ID NO: 4, 6, 8, 12, 14, 194, 196, or 198.

[0193] Because the reference polypeptide having the amino acid sequence of SEQ ID NO:18 is capable of converting the substrate to the product at a rate (for example, 100% conversion in 20 hours of 1 g/L substrate with about 10 g/L of the transaminase, in 50% isopropylamine at pH 8) and with a stereoselectivity that is improved over wild-type (SEQ ID NO:2), the polypeptides herein that are improved over SEQ ID NO:18 are also improved over wild-type.

[0194] Table 2 below provides a list of SEQ ID NOs having increased thermo- and/or solvent stability as compared to the wildtype transaminase of SEQ ID NO:2. All sequences are derived from the wild-type *Vibrio fluvialis* transaminase sequence (SEQ ID NO:2). In Table 2 below, each row lists two SEQ ID NOs, where the odd numbers in the column labeled "Nuc ID" refer to the nucleotide sequence that codes for the amino acid sequence provided by the even numbers in the column labeled "Pep ID". The amino acid substitutions as compared to the thermostable *Vibrio fluvialis* amino acid sequence of SEQ ID NO:2 are listed in the column "Active Amino Acid Mutations."

Table 2: List of Sequences and Corresponding Thermostability Improvements

Nuc ID	Pep ID	Active Amino Acid Mutations (as compared to SEQ ID NO:2)	% Residual Activity 50°C 23 h ^a	% Activity at 50°C rel. to 35°C, 23 h ^b
1	2	-	ND ¹	ND ¹
9	10	A9T; S398R; S420N	6.0	ND ¹
11	12	A9T; F86Y; M294V	4.6	0
15	16	A9T; D21N; F86Y; T208I; M294V		18%
17	18	A9T; N45H; F86Y; V177L; R211K; M294V; S324G; T391A		53%

¹ND = not detectable
^a % activity measured as % conversion at 50°C of 1.0M pyruvate with 1.0M isopropylamine at pH 9.
^b % activity measured as % conversion of pyruvate after 23 h at 50°C compared to

same reaction at 35°C.

[0195] **Table 3** below provides a list of SEQ ID NOs with their associated activities of transaminases for converting seven different amino acceptor substrates to the corresponding S-enantiomers of the amino products. Substrates 1 to 7 are (Sub 1) 3,4-dihydronaphthalen-1(2*H*)-one; (Sub 2) 1-phenylbutan-2-one; (Sub 3) 3,3-dimethylbutan-2-one; (Sub 4) octan-2-one; (Sub 5) ethyl 3-oxobutanoate; (Sub 6) 4-phenylbutan-2-one; and (Sub 7) 1-(4-bromophenyl)ethanone. The amino acid substitutions as compared to the thermostable *Vibrio fluvialis* amino acid sequence of SEQ ID NO:18 are listed in the column "Active Amino Acid Mutations." In the activity columns, one or more plus "+" signs indicate polypeptides having an improved ability to convert the amino acceptor substrate to the chiral amine product as compared to the wild-type amino acid sequence of SEQ ID NO:2. Two plus signs "++" indicates that the polypeptide is about 1.1 to 5-fold improved as compared to SEQ ID NO:18. Three plus signs "+++" indicates that the polypeptide is about 5 to 10-fold improved as compared to SEQ ID NO:18. Four plus signs "++++" indicates that the polypeptide is more than 10-fold improved as compared to SEQ ID NO:18. A blank indicates that data are unavailable. Substrates 3-7 show increased activity with most of the mutated polypeptides. Substrates 1 and 2 show increased activity with fewer polypeptides.

Table 3: List of Sequences and Corresponding Activity Improvement

Nuc ID	Pep ID	Active Amino Acid Mutations	Sub 1	Sub 2	Sub 3	Sub 4	Sub 5	Sub 6	Sub 7
19	20	Y86S			+	++	++	++	++
21	22	G81D; Y86H		++++	+	+++	+	++	++
23	24	Y86N			+	++	++	++	++
25	26	Y86F			+	++	+	++	++
27	28	W57L			+	++	+	++	++
29	30	Y82H; L417F			++	++	++	++	+
31	32	F85A; F317L;			+	+	+	+	
33	34	V153A; F317L; P318G			++	++	++	++	++
35	36	F85S;V153A			+	+	+	+	+
37	38	Y113H;C407S	+		++	++	+	+	++
39	40	F85A; V153A			+	+	+	+	+
41	42	W57I	+		+	++	+	++	++
43	44	V153S	++		++	++	++	++	++
45	46	W57I; V153S			+	++	+	++	++
47	48	Y86S; V153S			+	++	++	+	++
49	50	P318G; T408A			+	+	++	+	++
51	52	Y82H	+		++	++	++	++	++
53	54	E12G; M434V			++	++	++	+	++

Nuc ID	Pep ID	Active Amino Acid Mutations	Sub 1	Sub 2	Sub 3	Sub 4	Sub 5	Sub 6	Sub 7
55	56	F85V	+		+	+	+	+	+
57	58	F85S			+	+	+	+	
59	60	F85T			+	+	+	+	+
61	62	F85A			+	+	+	+	+
63	64	F85N			+	+	+	+	
65	66	F85C			+	+	+	+	
67	68	V153T	+		++	++	++	++	++
69	70	V153N			++	++	++	+	++
71	72	V153G			++	++	++	++	++
73	74	F317M	+		++	++	++	++	++
75	76	F317Y	+		++	++	++	++	++
77	78	L417A	+		+	+	+	++	++
79	80	H319Q	+		++	++	++	++	++
81	82	G320A	+		++	++	++	++	++
83	84	L417I	+		+	++	+	++	+++
85	86	L417C			+	++	+	++	++
87	88	Y86H; V153A; L417C		+++	+	+	++	++	++
89	90	Y86H;V153A		+++	+	+	+	++	+
91	92	Y86H; V153S; L417C		+++	+	++	++	++	++
93	94	T30A	+		+	+	++	++	++
95	96	V44A; N166S	+		++	++	++	++	++
97	98	V31A	+		++	++	++	++	++
99	100	I314V; D409G	+		++	+	+	+	++
101	102	V153A; P233T; L417C			+	++	+	++	+++
103	104	V153S; P233S	++		++	++	++	++	++
105	106	Y86H;V153A; A228G; L417I		+++	+	+	+	+	+
107	108	M95T; V153A; L417C			+	++	++	+	+++
109	110	Y86H;V153S; C181R; L417T		+++	+	++	++	++	++
111	112	Y113C; K385R; L417C			+	++	+	++	++
113	114	W57L; L417C; F438L			+	++	+	++	++
115	116	W57F;M127L; L417C			+	++	+	+	++
117	118	I311V; I314T	+		++	+	+	+	++

Nuc ID	Pep ID	Active Amino Acid Mutations	Sub 1	Sub 2	Sub 3	Sub 4	Sub 5	Sub 6	Sub 7
119	120	F112I; F317L			++	+	++		+
133	134	W57S;P233L; L417V			+	++	+	++	
135	136	W57S;Y86G; L417C				++	+	++	++
137	138	A228G;F317L; L417C		+		++	++	++	+
139	140	W57A;V153S; P318G		+	+	++	++	++	
141	142	W57F;Y86H; V153Q		++	+	++	++	++	
143	144	W57I;Y86F; G320A			+	++	+	++	+
145	146	W57F;Y86F; V153Q			+	++	+++	++	
147	148	W57A;V153C; F321L		+		++	++	++	
149	150	Y86F;P318R; L417A		+		+	++	++	++
151	152	A228G;P318G; L417C		+	+	++	++	+	++
153	154	W57C;Y86S; L417T			+	++	+	+++	++
155	156	Y86F;V153C; V297A			+	++	+++	++	++
157	158	Y86S;V153T; V297A			+	+	+++	++	+
159	160	V153S;P318R; L417E	+	+		+	+++	+	++
161	162	Y86H;P233L; L417A			+	+	++	++	+
163	164	V153C;P233L; P318R	++++	+	++	++	+++	++	++
165	166	W57C;Y86A; F317L				++	+	++	++
167	168	Y86N;A228G; F317L		+	+	++	+++	++	+
169	170	V153T; A228G F321L;		++	+	++	++++	+	++
171	172	V153S; F317L; L417C	++	++	+	++	+++	+++	+++
173	174	W57F; P318F; L417S			+	++	+	++	++
175	176	V153T; H319V; L417I			+		++	+	+
177	178	W57L; Y86S; V153A		+	++	++	+	+++	+
179	180	W57L; Y86F; P318F			+	++	++	++	
181	182	V153S; A228G; L417V;		++	+	++	+++	++	++
183	184	P233L; F321L; L417I	++		+	++	++	++	+++
185	186	F317L; P318R; L417T			+	+	++	++	+++
187	188	V153C; F317Y; H319Q	+++	+	++	++	++++	++	++
189	190	V153T; A228G; L417A		++	+	++	+++	++	+
191	192	W57F; P318G; L417I			+	+++	++	++	++

[0196] Table 4 below provides a list of SEQ ID NOs with their associated activities for

converting two different amino acceptor substrates to the corresponding *R*-enantiomers of the amino products. Substrates 5 and 6 are (Sub 5) ethyl 3-oxobutanoate and (Sub 6) 4-phenylbutan-2-one. All sequences below are derived from the wild-type *Vibrio fluvialis* transaminase sequence (SEQ ID NO:2). In Table 4 below, the odd numbered SEQ ID NOs in the column labeled "Nuc ID" refer to the nucleotide sequence that codes for the amino acid sequence provided by the even numbered SEQ ID NOs in the column labeled "Pep ID". The amino acid substitutions as compared to the amino acid sequence of SEQ ID NO:18 are listed in the column "Active Amino Acid Mutations." In the activity columns, one or more plus "+" signs indicate polypeptides having an improved ability to convert the amino acceptor substrate to the chiral amine product as compared to the wild-type amino acid sequence of SEQ ID NO:2. Two plus signs "++" indicates that the polypeptide is about 1.1 to 5-fold improved as compared to SEQ ID NO:18. Three plus signs "+++" indicates that the polypeptide is about 5 to 10-fold improved as compared to SEQ ID NO:18. Four plus signs "++++" indicates that the polypeptide is more than 10-fold improved as compared to SEQ ID NO:18. A blank indicates that data are unavailable. Substrate 5 shows increased activity with most of the mutated polypeptides. Substrate 6 shows increased activity with fewer polypeptides.

Table 4: List of Sequences and Corresponding Activity Improvement

Nuc ID	Pep ID	Active Amino Acid Mutations (as compared to SEQ ID NO:18)	Sub 5	Sub 6
21	22	G81D; Y86H	+	
29	30	Y82H; L417F	+	
31	32	F85A; F317L	++++	
33	34	V153A; F317L; P318G	++	
35	36	F85S; V153A	++++	
39	40	F85A; V153A	++++	
43	44	V153S	++	
47	48	Y86S; V153S	++	
55	56	F85V	++++	
57	58	F85S	++++	
59	60	F85T	++++	
61	62	F85A	++++	
63	64	F85N	++++	
65	66	F85C	++++	
67	68	V153T	++	
69	70	V153N	++	
71	72	V153G	++	
75	76	F317Y	++	
83	84	L417I	+	
85	86	L417C	+	

Nuc ID	Pep ID	Active Amino Acid Mutations (as compared to SEQ ID NO:18)	Sub 5	Sub 6
87	88	Y86H; V153A; L417C	++	
89	90	Y86H; V153A	+	
91	92	Y86H; V153S; L417C	++	
103	104	V153S; P233S	++	
107	108	M95T; V153A; L417C	+	
109	110	Y86H; V153S; C181R; L417T	++	
119	120	F112I; F317L		+
121	122	F85A; W147G; V153A	++	++++
123	124	F85S; V153S		++++
125	126	W57L; Y86F; V153S; P233T; L417T	+++	
127	128	F85S; V153A; P233T	+++	++++
129	130	W57I; F85A; Y86H; L417C	+++	
131	132	F85A; V153S; L417S	++	++++
159	160	V153S; P318R; L417E	++	
175	176	V153T; H319V; L417I	++	

[0197] In some aspects of the disclosure, an improved transaminase comprises an amino acid sequence that is at least about 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to SEQ ID NO:2 or SEQ ID NO:18, wherein the amino acid sequence comprises any one of the set of mutations contained in any one of the polypeptide sequences listed in **Table 3** or **Table 4** as compared to a reference sequence corresponding to SEQ ID NO:2 or SEQ ID NO:18. In some aspects of the disclosure, this transaminase polypeptide comprises any one of the set of mutations listed in

Table 3 or **Table 4**. In some aspects, the engineered transaminases can have additionally about 1-2, 1-3, 1-4, 1-5, 1-6, 1-7, 1-8, 1-9, 1-10, 1-15, 1-20, 1-21, 1-22, 1-23, 1-24, 1-25, 1-30, 1-35, 1-40, 1-45, 1-50, 1-55, or 1-60 residue differences at other residue positions. In some aspects, the number of differences as compared to the reference sequence can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 30, 35, 40, 45, 50, 55, or 60 residue positions. In some aspects, the residue differences comprise conservative mutations.

[0198] In some aspects of the disclosure, an improved transaminase polypeptide comprises an amino acid sequence that is at least about 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to a reference sequence corresponding to SEQ ID NO: 4, 6, 8, 10, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172,

174, 176, 178, 180, 182, 184, 186, 188, 190, 192, 194, 196, or 198, wherein the engineered transaminase amino acid sequence comprises any one of the set of mutations contained in any one of the polypeptide sequences listed in **Table 3** or **Table 4** as compared to a reference sequence corresponding to SEQ ID NO:2 or SEQ ID NO:18. In some aspects of the disclosure, this transaminase polypeptide comprises any one of the set of mutations listed in **Table 3** or **Table 4**. In some aspects, these engineered transaminases can have additionally about 1-2, 1-3, 1-4, 1-5, 1-6, 1-7, 1-8, 1-9, 1-10, 1-15, 1-20, 1-21, 1-22, 1-23, 1-24, 1-25, 1-30, 1-35, 1-40, 1-45, 1-50, 1-55, or 1-60 residue differences at other residue positions. In some aspects, the number of differences as compared to the reference sequence can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 30, 35, 40, 45, 50, 55, or 60 residue positions. In some aspects, the residue differences comprise conservative mutations.

[0199] In some aspects of the disclosure, the amino acid sequences of the polypeptides of the current disclosure specifically exclude those that differ from the wild-type sequence of *Vibrio fluvialis* by only one of any of the following single mutations: residue X57 is G; residue X147 is G; residue X231 is G; residue X233 is L; residue X265 is L; residue X285 is A; residue X297 is A; and residue X415 is L.

[0200] The engineered transaminase enzymes described herein can be obtained by mutagenizing a gene encoding a naturally-occurring wild-type transaminase enzyme that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence of *Vibrio fluvialis* transaminase (SEQ ID NO:2) utilizing standard laboratory techniques, including various mutagenesis and recombination techniques.

[0201] As will be appreciated by those of skill in the art, some of the above-defined categories of amino acid residues used to describe the engineered transaminases herein, unless otherwise specified, are not mutually exclusive. Thus, amino acids having side chains exhibiting two or more physico-chemical properties can be included in multiple categories. The appropriate classification of any amino acid or residue will be apparent to those of skill in the art, especially in light of the detailed disclosure provided herein.

[0202] In some aspects, the improved engineered transaminase polypeptides comprise deletions of the engineered transaminase polypeptides described herein. Thus, for each and every transaminase polypeptides of the disclosure, the deletions can comprise one or more amino acids, 2 or more amino acids, 3 or more amino acids, 4 or more amino acids, 5 or more amino acids, 6 or more amino acids, 8 or more amino acids, 10 or more amino acids, 15 or more amino acids, or 20 or more amino acids, up to 10% of the total number of amino acids, up to 10% of the total number of amino acids, up to 20% of the total number of amino acids, or up to 30% of the total number of amino acids of the transaminase polypeptides, as long as the functional activity and/or improved properties of the transaminase is maintained. In some aspects, the deletions can comprise, 1-2, 1-3, 1-4, 1-5, 1-6, 1-7, 1-8, 1-9, 1-10, 1-15, 1-20, 1-21, 1-22, 1-23, 1-24, 1-25, 1-30, 1-35, 1-40, 1-45, 1-50, 1-55, or 1-60 amino acid residues. In some aspects, the number of deletions can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 16, 17, 18, 19,

20, 21, 22, 23, 24, 25, 30, 30, 35, 40, 45, 50, 55, or 60 amino acids. In some aspects, the deletions can comprise deletions of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, 20, 21, 22, 23, 24, 25 or 30 amino acid residues.

[0203] The improved engineered transaminase polypeptides may comprise fragments of the engineered transaminase enzymes described herein. In some aspects, the polypeptide fragments can be 70%, 80%, 90%, 95%, 98%, or 99% of the full-length transaminase polypeptide, such as the transaminase of SEQ ID NO:18.

[0204] The polypeptides described herein are not restricted to the genetically encoded amino acids. In addition to the genetically encoded amino acids, the polypeptides described herein may be comprised, either in whole or in part, of naturally-occurring and/or synthetic non-encoded amino acids. Certain commonly encountered non-encoded amino acids of which the polypeptides described herein may be comprised include, but are not limited to: the D-stereomers of the genetically-encoded amino acids; 2,3-diaminopropionic acid (Dpr); α -aminoisobutyric acid (Aib); ϵ -aminohexanoic acid (Aha); δ -aminovateric acid (Ava); N-methylglycine or sarcosine (MeGly or Sar); ornithine (Orn); citrulline (Cit); t-butylalanine (Bua); t-butylglycine (Bug); N-methylisoleucine (Melle); phenylglycine (Phg); cyclohexylalanine (Cha); norleucine (Nle); naphthylalanine (Nal); 2-chlorophenylalanine (Ocf); 3-chlorophenylalanine (Mcf); 4-chlorophenylalanine (Pcf); 2-fluorophenylalanine (Off); 3-fluorophenylalanine (Mff); 4-fluorophenylalanine (Pff); 2-bromophenylalanine (Obf); 3-bromophenylalanine (Mbf); 4-bromophenylalanine (Pbf); 2-methylphenylalanine (Omf); 3-methylphenylalanine (Mmf); 4-methylphenylalanine (Pmf); 2-nitrophenylalanine (Onf); 3-nitrophenylalanine (Mnf); 4-nitrophenylalanine (Pnf); 2-cyanophenylalanine (Ocf); 3-cyanophenylalanine (Mcf); 4-cyanophenylalanine (Pcf); 2-trifluoromethylphenylalanine (Otf); 3-trifluoromethylphenylalanine (Mtf); 4-trifluoromethylphenylalanine (Ptf); 4-aminophenylalanine (Paf); 4-iodophenylalanine (Pif); 4-aminomethylphenylalanine (Pamf); 2,4-dichlorophenylalanine (Opof); 3,4-dichlorophenylalanine (Mpcf); 2,4-difluorophenylalanine (Opff); 3,4-difluorophenylalanine (Mpff); pyrid-2-ylalanine (2pAla); pyrid-3-ylalanine (3pAla); pyrid-4-ylalanine (4pAla); naphth-1-ylalanine (1nAla); naphth-2-ylalanine (2nAla); thiazolylalanine (taAla); benzothienylalanine (bAla); thienylalanine (tAla); furylalanine (fAla); homophenylalanine (hPhe); homotyrosine (hTyr); homotryptophan (hTrp); pentafluorophenylalanine (5ff); styrylalanine (sAla); authrylalanine (aAla); 3,3-diphenylalanine (Dfa); 3-amino-5-phenypentanoic acid (Afp); penicillamine (Pen); 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid (Tic); β -2-thienylalanine (Thi); methionine sulfoxide (Mso); N(w)-nitroarginine (nArg); homolysine (hLys); phosphonomethylphenylalanine (pmPhe); phosphoserine (pSer); phosphothreonine (pThr); homoaspartic acid (hAsp); homoglutamic acid (hGlu); 1-aminocyclopent-(2 or 3)-ene-4 carboxylic acid; pipercolic acid (PA), azetidine-3-carboxylic acid (ACA); 1-aminocyclopentane-3-carboxylic acid; allylglycine (aOly); propargylglycine (pgGly); homoalanine (hAla); norvaline (nVal); homoleucine (hLeu), homovaline (hVal); homoisoleucine (hIle); homoarginine (hArg); N-acetyl lysine (AcLys); 2,4-diaminobutyric acid (Dbu); 2,3-diaminobutyric acid (Dab); N-methylvaline (MeVal); homocysteine (hCys); homoserine (hSer); hydroxyproline (Hyp) and homoproline (hPro). Additional non-encoded amino acids of which the polypeptides described herein may be comprised will be apparent to those of skill in the art (see, e.g., the various

amino acids provided in Fasman, 1989, CRC Practical Handbook of Biochemistry and Molecular Biology, CRC Press, Boca Raton, FL, at pp. 3-70 and the references cited therein). These amino acids may be in either the L- or D-configuration.

[0205] Those of skill in the art will recognize that amino acids or residues bearing side chain protecting groups may also comprise the polypeptides described herein. Non-limiting examples of such protected amino acids, which in this case belong to the aromatic category, include (protecting groups listed in parentheses), but are not limited to: Arg(tos), Cys(methylbenzyl), Cys (nitropyridinesulphenyl), Glu(δ -benzylester), Gln(xanthyl), Asn(N- δ -xanthyl), His(bom), His(benzyl), His(tos), Lys(fmoc), Lys(tos), Ser(O-benzyl), Thr (O-benzyl) and Tyr(O-benzyl).

[0206] Non-encoding amino acids that are conformationally constrained of which the polypeptides described herein may be composed include, but are not limited to, N-methyl amino acids (L-configuration); 1-aminocyclopent-(2 or 3)-ene-4-carboxylic acid; pipercolic acid; azetidine-3-carboxylic acid; homoproline (hPro); and 1-aminocyclopentane-3-carboxylic acid.

[0207] As described above the various modifications introduced into the naturally occurring polypeptide to generate an engineered transaminase enzyme can be targeted to a specific property of the enzyme.

[0208] In some embodiments, the polypeptides of the disclosure can be in the form of fusion polypeptides in which the engineered polypeptides are fused to other polypeptides, such as, by way of example and not limitation, antibody tags (e.g., myc epitope), purification sequences (e.g., His tags for binding to metals), and cell localization signals (e.g., secretion signals). Thus, the engineered polypeptides described herein can be used with or without fusions to other polypeptides.

[0209] In some embodiments, the polypeptide described herein can be provided in the form of kits. The enzymes in the kits may be present individually or as a plurality of enzymes. The kits can further include reagents for carrying out the enzymatic reactions, substrates for assessing the activity of enzymes, as well as reagents for detecting the products. The kits can also include reagent dispensers and instructions for use of the kits.

[0210] In some embodiments, the polypeptides can be provided on a physical substrate. In some embodiments, the polypeptides can be provided in the form of an array in which the polypeptides are arranged in positionally distinct locations. The array can be used to test a variety of substrate compounds for conversion by the polypeptides. "Substrate," "support," "solid support," "solid carrier," or "resin" in the context of arrays refer to any solid phase material. Substrate also encompasses terms such as "solid phase," "surface," and/or "membrane." A solid support can be composed of organic polymers such as polystyrene, polyethylene, polypropylene, polyfluoroethylene, polyethyleneoxy, and polyacrylamide, as well as co-polymers and grafts thereof. A solid support can also be inorganic, such as glass, silica, controlled pore glass (CPG), reverse phase silica or metal, such as gold or platinum. The configuration of a substrate can be in the form of beads, spheres, particles, granules, a gel, a

membrane or a surface. Surfaces can be planar, substantially planar, or non-planar. Solid supports can be porous or non-porous, and can have swelling or non-swelling characteristics. A solid support can be configured in the form of a well, depression, or other container, vessel, feature, or location. A plurality of supports can be configured on an array at various locations, addressable for robotic delivery of reagents, or by detection methods and/or instruments.

[0211] In certain embodiments, the kits of the present disclosure include arrays comprising a plurality of different transaminase polypeptides at different addressable position, wherein the different polypeptides are different variants of a reference sequence each having at least one different improved enzyme property. Such arrays comprising a plurality of engineered polypeptides and methods of their use are described in, e.g., WO2009/008908A2.

[0212] In some embodiments, the transaminase polypeptides can be bound on a physical substrate. The transaminase polypeptide can be bound non-covalently or covalently. Various methods for conjugation to substrates, e.g., membranes, beads, glass, etc. are described in, among others, Hermanson, G.T., *Bioconjugate Techniques*, Second Edition, Academic Press; (2008), and *Bioconjugation Protocols: Strategies and Methods*, In *Methods in Molecular Biology*, C.M. Niemeyer ed., Humana Press (2004).

6.4 Polynucleotides Encoding Engineered Transaminases

[0213] In another aspect, the present disclosure provides polynucleotides encoding the engineered transaminase enzymes. The polynucleotides may be operatively linked to one or more heterologous regulatory sequences that control gene expression to create a recombinant polynucleotide capable of expressing the polypeptide. Expression constructs containing a heterologous polynucleotide encoding the engineered transaminase can be introduced into appropriate host cells to express the corresponding transaminase polypeptide.

[0214] Because of the knowledge of the codons corresponding to the various amino acids, availability of a protein sequence provides a description of all the polynucleotides capable of encoding the subject. The degeneracy of the genetic code, where the same amino acids are encoded by alternative or synonymous codons allows an extremely large number of nucleic acids to be made, all of which encode the improved transaminase enzymes disclosed herein. Thus, having identified a particular amino acid sequence, those skilled in the art could make any number of different nucleic acids by simply modifying the sequence of one or more codons in a way which does not change the amino acid sequence of the protein. In this regard, the present disclosure specifically contemplates each and every possible variation of polynucleotides that could be made by selecting combinations based on the possible codon choices, and all such variations are to be considered specifically disclosed for any polypeptide disclosed herein, including the amino acid sequences presented in **Table 3** or **Table 4**.

[0215] In various embodiments, the codons are preferably selected to fit the host cell in which the protein is being produced. For example, preferred codons used in bacteria are used to

express the gene in bacteria; preferred codons used in yeast are used for expression in yeast; and preferred codons used in mammals are used for expression in mammalian cells.

[0216] In certain embodiments, all codons need not be replaced to optimize the codon usage of the transaminases since the natural sequence will comprise preferred codons and because use of preferred codons may not be required for all amino acid residues. Consequently, codon optimized polynucleotides encoding the transaminase enzymes may contain preferred codons at about 40%, 50%, 60%, 70%, 80%, or greater than 90% of codon positions of the full length coding region.

[0217] In some aspects of the disclosure, the polynucleotide comprises a nucleotide sequence encoding a transaminase polypeptide with an amino acid sequence that has at least about 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to any of the reference engineered transaminase polypeptides described herein. Accordingly, in some aspects of the disclosure, the polynucleotide encodes an amino acid sequence that is at least about 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to SEQ ID NO: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 176, 178, 180, 182, 184, 186, 188, 190, 192, 194, 196, or 198, where the polypeptide has transaminase activity and one or more of the improved properties described herein.

[0218] In some aspects of the disclosure, the polynucleotide comprises a nucleotide sequence encoding an engineered transaminase polypeptide with an amino acid sequence that has at least about 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the polypeptide comprising an amino acid corresponding to SEQ ID NO: 10, where the polypeptide has transaminase activity and one or more of the improved properties described herein.

[0219] In some aspects of the disclosure, the polynucleotide comprises a nucleotide sequence encoding an engineered transaminase polypeptide with an amino acid sequence that has at least about 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the polypeptide comprising an amino acid corresponding to SEQ ID NO: 16, where the polypeptide has transaminase activity and one or more of the improved properties described herein.

[0220] In some aspects of the disclosure, the polynucleotide comprises a nucleotide sequence encoding an engineered transaminase polypeptide with an amino acid sequence that has at least about 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the polypeptide comprising an amino acid corresponding to SEQ ID NO:18, where the polypeptide has transaminase activity and one or more of the improved properties described herein.

[0221] In some aspects of the disclosure, the polynucleotides encoding the engineered transaminases are selected from SEQ ID NO: 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 159, 161, 163, 165, 167, 169, 171, 173, 175, 177, 179, 181, 183, 185, 187, 189, 191, 193, 195, and 197.

[0222] In some aspects of the disclosure, the polynucleotides are capable of hybridizing under highly stringent conditions to a polynucleotide corresponding to SEQ ID NO: 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 159, 161, 163, 165, 167, 169, 171, 173, 175, 177, 179, 181, 183, 185, 187, 189, 191, 193, 195, or 197, or a complement thereof, and encodes a polypeptide having transaminase activity with one or more of the improved properties described herein.

[0223] In some aspects of the disclosure, the polynucleotides encode the polypeptides described herein but have about 80% or more sequence identity, about 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity at the nucleotide level to a reference polynucleotide encoding the engineered transaminase. In some aspects of the disclosure, the reference polynucleotide sequence is selected from SEQ ID NO: 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 159, 161, 163, 165, 167, 169, 171, 173, 175, 177, 179, 181, 183, 185, 187, 189, 191, 193, 195, and 197.

[0224] An isolated polynucleotide encoding an improved transaminase polypeptide may be manipulated in a variety of ways to provide for expression of the polypeptide. Manipulation of the isolated polynucleotide prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotides and nucleic acid sequences utilizing recombinant DNA methods are well known in the art. Guidance is provided in Sambrook et al., 2001, *Molecular Cloning: A Laboratory Manual*, 3rd Ed., Cold Spring Harbor Laboratory Press; and *Current Protocols in Molecular Biology*, Ausubel, F. ed., Greene Pub. Associates, 1998, updates to 2006.

[0225] For bacterial host cells, suitable promoters for directing transcription of the nucleic acid constructs of the present disclosure, include the promoters obtained from the *E. coli* lac operon, *Streptomyces coelicolor* agarase gene (dagA), *Bacillus subtilis* levansucrase gene (sacB), *Bacillus licheniformis* alpha-amylase gene (amyL), *Bacillus stearothermophilus* maltogenic amylase gene (amyM), *Bacillus amyloliquefaciens* alpha-amylase gene (amyQ), *Bacillus licheniformis* penicillinase gene (penP), *Bacillus subtilis* xylA and xylB genes, and

prokaryotic beta-lactamase gene (Villa-Kamaroff et al., 1978, Proc. Natl Acad. Sci. USA 75: 3727-3731), as well as the *tac* promoter (DeBoer et al., 1983, Proc. Natl Acad. Sci. USA 80: 21-25).

[0226] For filamentous fungal host cells, suitable promoters for directing the transcription of the nucleic acid constructs of the present disclosure include promoters obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (glaA), *Rhizomucor miehei* lipase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Aspergillus nidulans* acetamidase, and *Fusarium oxysporum* trypsin-like protease (WO 96/00787), as well as the NA2-tpi promoter (a hybrid of the promoters from the genes for *Aspergillus niger* neutral alpha-amylase and *Aspergillus oryzae* triose phosphate isomerase), and mutant, truncated, and hybrid promoters thereof.

[0227] In a yeast host, useful promoters can be from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* galactokinase (GAL1), *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP), and *Saccharomyces cerevisiae* 3-phosphoglycerate kinase. Other useful promoters for yeast host cells are described by Romanos et al., 1992, Yeast 8:423-488.

[0228] The control sequence may also be a suitable transcription terminator sequence, a sequence recognized by a host cell to terminate transcription. The terminator sequence is operably linked to the 3' terminus of the nucleic acid sequence encoding the polypeptide. Any terminator which is functional in the host cell of choice may be used in the present invention.

[0229] For example, exemplary transcription terminators for filamentous fungal host cells can be obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* alpha-glucosidase, and *Fusarium oxysporum* trypsin-like protease.

[0230] Exemplary terminators for yeast host cells can be obtained from the genes for *Saccharomyces cerevisiae* enolase, *Saccharomyces cerevisiae* cytochrome C (CYC1), and *Saccharomyces cerevisiae* glyceraldehyde-3-phosphate dehydrogenase. Other useful terminators for yeast host cells are described by Romanos et al., 1992, *supra*.

[0231] The control sequence may also be a suitable leader sequence, a nontranslated region of an mRNA that is important for translation by the host cell. The leader sequence is operably linked to the 5' terminus of the nucleic acid sequence encoding the polypeptide. Any leader sequence that is functional in the host cell of choice may be used. Exemplary leaders for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase and *Aspergillus nidulans* triose phosphate isomerase. Suitable leaders for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* 3-phosphoglycerate kinase, *Saccharomyces cerevisiae* alpha-factor, and

Saccharomyces cerevisiae alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP).

[0232] The control sequence may also be a polyadenylation sequence, a sequence operably linked to the 3' terminus of the nucleic acid sequence and which, when transcribed, is recognized by the host cell as a signal to add polyadenosine residues to transcribed mRNA. Any polyadenylation sequence which is functional in the host cell of choice may be used in the present invention. Exemplary polyadenylation sequences for filamentous fungal host cells can be from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Fusarium oxysporum* trypsin-like protease, and *Aspergillus niger* alpha-glucosidase. Useful polyadenylation sequences for yeast host cells are described by Guo and Sherman, 1995, Mol Cell Bio 15:5983-5990.

[0233] The control sequence may also be a signal peptide coding region that codes for an amino acid sequence linked to the amino terminus of a polypeptide and directs the encoded polypeptide into the cell's secretory pathway. The 5' end of the coding sequence of the nucleic acid sequence may inherently contain a signal peptide coding region naturally linked in translation reading frame with the segment of the coding region that encodes the secreted polypeptide. Alternatively, the 5' end of the coding sequence may contain a signal peptide coding region that is foreign to the coding sequence. The foreign signal peptide coding region may be required where the coding sequence does not naturally contain a signal peptide coding region.

[0234] Alternatively, the foreign signal peptide coding region may simply replace the natural signal peptide coding region in order to enhance secretion of the polypeptide. However, any signal peptide coding region which directs the expressed polypeptide into the secretory pathway of a host cell of choice may be used in the present invention.

[0235] Effective signal peptide coding regions for bacterial host cells are the signal peptide coding regions obtained from the genes for *Bacillus* NCIB 11837 maltogenic amylase, *Bacillus stearothermophilus* alpha-amylase, *Bacillus licheniformis* subtilisin, *Bacillus licheniformis* beta-lactamase, *Bacillus stearothermophilus* neutral proteases (nprT, nprS, nprM), and *Bacillus subtilis* prsA. Further signal peptides are described by Simonen and Palva, 1993, Microbiol Rev 57: 109-137.

[0236] Effective signal peptide coding regions for filamentous fungal host cells can be the signal peptide coding regions obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* neutral amylase, *Aspergillus niger* glucoamylase, *Rhizomucor miehei* aspartic proteinase, *Humicola insolens* cellulase, and *Humicola lanuginosa* lipase.

[0237] Useful signal peptides for yeast host cells can be from the genes for *Saccharomyces cerevisiae* alpha-factor and *Saccharomyces cerevisiae* invertase. Other useful signal peptide coding regions are described by Romanos et al., 1992, *supra*.

[0238] The control sequence may also be a propeptide coding region that codes for an amino acid sequence positioned at the amino terminus of a polypeptide. The resultant polypeptide is known as a proenzyme or propolypeptide (or a zymogen in some cases). A propolypeptide can be converted to a mature active polypeptide by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide. The propeptide coding region may be obtained from the genes for *Bacillus subtilis* alkaline protease (aprE), *Bacillus subtilis* neutral protease (nprT), *Saccharomyces cerevisiae* alpha-factor, *Rhizomucor miehei* aspartic proteinase, and *Myceliophthora thermophila* lactase (WO 95/33836).

[0239] Where both signal peptide and propeptide regions are present at the amino terminus of a polypeptide, the propeptide region is positioned next to the amino terminus of a polypeptide and the signal peptide region is positioned next to the amino terminus of the propeptide region.

[0240] It may also be desirable to add regulatory sequences, which allow the regulation of the expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those which cause the expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. In prokaryotic host cells, suitable regulatory sequences include the lac, tac, and trp operator systems. In yeast host cells, suitable regulatory systems include, as examples, the ADH2 system or GAL1 system. In filamentous fungi, suitable regulatory sequences include the TAKA alpha-amylase promoter, *Aspergillus niger* glucoamylase promoter, and *Aspergillus oryzae* glucoamylase promoter.

[0241] Other examples of regulatory sequences are those which allow for gene amplification. In eukaryotic systems, these include the dihydrofolate reductase gene, which is amplified in the presence of methotrexate, and the metallothionein genes, which are amplified with heavy metals. In these cases, the nucleic acid sequence encoding the polypeptide of the present invention would be operably linked with the regulatory sequence.

[0242] Thus, in another aspect, the present disclosure is also directed to a recombinant expression vector comprising a polynucleotide encoding an engineered transaminase polypeptide or a variant thereof, and one or more expression regulating regions such as a promoter and a terminator, a replication origin, etc., depending on the type of hosts into which they are to be introduced. The various nucleic acid and control sequences described above may be joined together to produce a recombinant expression vector which may include one or more convenient restriction sites to allow for insertion or substitution of the nucleic acid sequence encoding the polypeptide at such sites. Alternatively, the nucleic acid sequence of the present disclosure may be expressed by inserting the nucleic acid sequence or a nucleic acid construct comprising the sequence into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression.

[0243] The recombinant expression vector may be any vector (e.g., a plasmid or virus), which can be conveniently subjected to recombinant DNA procedures and can bring about the

expression of the polynucleotide sequence. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vectors may be linear or closed circular plasmids.

[0244] The expression vector may be an autonomously replicating vector, i.e., a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one which, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. Furthermore, a single vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the host cell, or a transposon may be used.

[0245] The expression vector of the present invention preferably contains one or more selectable markers, which permit easy selection of transformed cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like. Examples of bacterial selectable markers are the *dal* genes from *Bacillus subtilis* or *Bacillus licheniformis*, or markers, which confer antibiotic resistance such as ampicillin, kanamycin, chloramphenicol (Example 1) or tetracycline resistance. Suitable markers for yeast host cells are ADE2, HIS3, LEU2, LYS2, MET3, TRP1, and URA3.

[0246] Selectable markers for use in a filamentous fungal host cell include, but are not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hph* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase), *sC* (sulfate adenylyltransferase), and *trpC* (anthranilate synthase), as well as equivalents thereof. Embodiments for use in an *Aspergillus* cell include the *amdS* and *pyrG* genes of *Aspergillus nidulans* or *Aspergillus oryzae* and the *bar* gene of *Streptomyces hygroscopicus*.

[0247] The expression vectors of the present invention can contain an element(s) that permits integration of the vector into the host cell's genome or autonomous replication of the vector in the cell independent of the genome. For integration into the host cell genome, the vector may rely on the nucleic acid sequence encoding the polypeptide or any other element of the vector for integration of the vector into the genome by homologous or nonhomologous recombination.

[0248] Alternatively, the expression vector may contain additional nucleic acid sequences for directing integration by homologous recombination into the genome of the host cell. The additional nucleic acid sequences enable the vector to be integrated into the host cell genome at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements should preferably contain a sufficient number of nucleic acids, such as 100 to 10,000 base pairs, preferably 400 to 10,000 base pairs, and most preferably 800 to 10,000 base pairs, which are highly homologous with the corresponding target sequence to enhance the probability of homologous recombination. The integrational

elements may be any sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding nucleic acid sequences. On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination.

[0249] For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. Examples of bacterial origins of replication are P15A ori (as shown in the plasmid of FIG. 5) or the origins of replication of plasmids pBR322, pUC19, pACYCI77 (which plasmid has the P15A ori), or pACYC184 permitting replication in *E. coli*, and pUB110, pE194, pTA1060, or pAM β 1 permitting replication in *Bacillus*. Examples of origins of replication for use in a yeast host cell are the 2 micron origin of replication, ARS1, ARS4, the combination of ARS1 and CEN3, and the combination of ARS4 and CEN6. The origin of replication may be one having a mutation which makes its functioning temperature-sensitive in the host cell (see, e.g., Ehrlich, 1978, Proc Natl Acad Sci. USA 75:1433).

[0250] More than one copy of a nucleic acid sequence of the present invention may be inserted into the host cell to increase production of the gene product. An increase in the copy number of the nucleic acid sequence can be obtained by integrating at least one additional copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the nucleic acid sequence where cells containing amplified copies of the selectable marker gene, and thereby additional copies of the nucleic acid sequence, can be selected for by cultivating the cells in the presence of the appropriate selectable agent.

[0251] Many of the expression vectors for use in the present invention are commercially available. Suitable commercial expression vectors include p3xFLAGTM™ expression vectors from Sigma-Aldrich Chemicals, St. Louis MO., which includes a CMV promoter and hGH polyadenylation site for expression in mammalian host cells and a pBR322 origin of replication and ampicillin resistance markers for amplification in *E. coli*. Other suitable expression vectors are pBluescriptII SK(-) and pBK-CMV, which are commercially available from Stratagene, LaJolla CA, and plasmids which are derived from pBR322 (Gibco BRL), pUC (Gibco BRL), pREP4, pCEP4 (Invitrogen) or pPoly (Lathe et al., 1987, Gene 57:193-201).

6.5 Host Cells for Expression of Transaminase Polypeptides

[0252] In another aspect, the present disclosure provides a host cell comprising a polynucleotide encoding an improved transaminase polypeptide of the present disclosure, the polynucleotide being operatively linked to one or more control sequences for expression of the transaminase enzyme in the host cell. Host cells for use in expressing the polypeptides encoded by the expression vectors of the present invention are well known in the art and include but are not limited to, bacterial cells, such as *E. coli*, *Vibrio fluvialis*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells (e.g., *Saccharomyces cerevisiae* or *Pichia pastoris* (ATCC Accession No. 201178)); insect cells such as *Drosophila* S2 and

Spodoptera Sf9 cells; animal cells such as CHO, COS, BHK, 293, and Bowes melanoma cells; and plant cells. Appropriate culture mediums and growth conditions for the above-described host cells are well known in the art.

[0253] Polynucleotides for expression of the transaminase may be introduced into cells by various methods known in the art. Techniques include among others, electroporation, biolistic particle bombardment, liposome mediated transfection, calcium chloride transfection, and protoplast fusion. Various methods for introducing polynucleotides into cells will be apparent to the skilled artisan.

[0254] An exemplary host cell is *Escherichia coli* W3110. The expression vector was created by operatively linking a polynucleotide encoding an improved transaminase into the plasmid pCK110900 operatively linked to the *lac* promoter under control of the *lacI* repressor. The expression vector also contained the P15a origin of replication and the chloramphenicol resistance gene.

6.6 Methods of Generating Engineered Transaminase Polypeptides.

[0255] To make the improved polynucleotides and polypeptides of the present disclosure, the naturally-occurring transaminase enzyme that catalyzes the transamination reaction can be obtained (or derived) from *Vibrio fluvialis*. The parent polynucleotide sequence may be codon optimized to enhance expression of the transaminase in a specified host cell. As an illustration, the parental polynucleotide sequence encoding the wild-type polypeptide of *Vibrio fluvialis* was constructed from oligonucleotides prepared based upon the known polypeptide sequence of *Vibrio fluvialis* sequence (Shin et al., 2003, "Purification, characterization, and molecular cloning of a novel amine:pyruvate transaminase from *Vibrio fluvialis* JS17" Appl Microbiol Biotechnol. 61(5-6):463-471). The parental polynucleotide sequence, designated as SEQ ID NO: 1, was codon optimized for expression in *E. coli* and the codon-optimized polynucleotide cloned into an expression vector, placing the expression of the transaminase gene under the control of the *lac* promoter and *lacI* repressor gene. Clones expressing the active transaminase in *E. coli* were identified and the genes sequenced to confirm their identity. The sequence designated SEQ ID NO: 17 was the parent sequence utilized as the starting point for most experiments and library construction of engineered transaminases evolved from the *Vibrio fluvialis* transaminase.

[0256] The engineered transaminases can be obtained by subjecting the polynucleotide encoding the naturally occurring transaminase to mutagenesis and/or directed evolution methods, as discussed above. An exemplary directed evolution technique is mutagenesis and/or DNA shuffling as described in Stemmer, 1994, Proc Natl Acad Sci USA 91:10747-10751; WO 95/22625; WO 97/0078; WO 97/35966; WO 98/27230; WO 00/42651; WO 01/75767 and U.S. Pat. 6,537,746. Other directed evolution procedures that can be used include, among others, staggered extension process (StEP), in vitro recombination (Zhao et al., 1998, Nat. Biotechnol. 16:258-261), mutagenic PCR (Caldwell et al., 1994, PCR Methods

Appl. 3:S136-S140), and cassette mutagenesis (Black et al., 1996, Proc Natl Acad Sci USA 93:3525-3529). Mutagenesis and directed evolution techniques useful for the purposes herein are also described in the following references: Ling, et al., 1997, Anal. Biochem. 254(2):157-78; Dale et al., 1996, "Oligonucleotide-directed random mutagenesis using the phosphorothioate method," In Methods Mol. Biol. 57:369-74; Smith, 1985, Ann. Rev. Genet. 19:423-462; Botstein et al., 1985, Science 229:1193-1201; Carter, 1986, Biochem. J. 237:1-7; Kramer et al., 1984, Cell, 38:879-887; Wells et al., 1985, Gene 34:315-323; Minshull et al., 1999, Curr Opin Chem Biol 3:284-290; Christians et al., 1999, Nature Biotech 17:259-264; Crameri et al., 1998, Nature 391:288-291; Crameri et al., 1997, Nature Biotech 15:436-438; Zhang et al., 1997, Proc Natl Acad Sci USA 94:4504-4509; Crameri et al., 1996, Nature Biotech 14:315-319; Stemmer, 1994, Nature 370:389-391; Stemmer, 1994, Proc Natl Acad Sci USA 91:10747-10751; WO 95/22625; WO 97/0078; WO 97/35966; WO 98/27230; WO 00/42651; WO 01/75767 and U.S. Pat. 6,537,746.

[0257] The clones obtained following mutagenesis treatment can be screened for engineered transaminases having a desired improved enzyme property. Measuring enzyme activity from the expression libraries can be performed using the standard biochemistry techniques, such as HPLC analysis following OPA derivitization of the product amine.

[0258] Where the improved enzyme property desired is thermostability, enzyme activity may be measured after subjecting the enzyme preparations to a defined temperature and measuring the amount of enzyme activity remaining after heat treatments. Clones containing a polynucleotide encoding a transaminase are then isolated, sequenced to identify the nucleotide sequence changes (if any), and used to express the enzyme in a host cell.

[0259] Where the sequence of the engineered polypeptide is known, the polynucleotides encoding the enzyme can be prepared by standard solid-phase methods, according to known synthetic methods. Fragments of up to about 100 bases can be individually synthesized, then joined (e.g., by enzymatic or chemical ligation methods, or polymerase mediated methods) to form any desired continuous sequence. For example, polynucleotides and oligonucleotides of the invention can be prepared by chemical synthesis using, e.g., the classical phosphoramidite method described by Beaucage et al., 1981, Tet Lett 22:1859-69, or the method described by Matthes et al., 1984, EMBO J. 3:801-05, e.g., as it is typically practiced in automated synthetic methods. According to the phosphoramidite method, oligonucleotides are synthesized, e.g., in an automatic DNA synthesizer, purified, annealed, ligated and cloned in appropriate vectors. In addition, essentially any nucleic acid can be obtained from any of a variety of commercial sources.


[0260] Engineered transaminase enzymes expressed in a host cell can be recovered from the cells and or the culture medium using any one or more of the well known techniques for protein purification, including, among others, lysozyme treatment, sonication, filtration, salting-out, ultra-centrifugation, and chromatography. Suitable solutions for lysing and the high efficiency extraction of proteins from bacteria, such as *E. coli*, are commercially available under the trade name CellLytic B™ from Sigma-Aldrich of St. Louis MO.

[0261] Chromatographic techniques for isolation of the transaminase polypeptide include, among others, reverse phase chromatography high performance liquid chromatography, ion exchange chromatography, gel electrophoresis, and affinity chromatography. Conditions for purifying a particular enzyme will depend, in part, on factors such as net charge, hydrophobicity, hydrophilicity, molecular weight, molecular shape, etc., and will be apparent to those having skill in the art.

[0262] Affinity techniques may be used to isolate the improved transaminase enzymes. For affinity chromatography purification, any antibody which specifically binds the transaminase polypeptide may be used. For the production of antibodies, various host animals, including but not limited to rabbits, mice, rats, etc., may be immunized by injection with a transaminase polypeptide, or a fragment thereof. The transaminase polypeptide or fragment may be attached to a suitable carrier, such as BSA, by means of a side chain functional group or linkers attached to a side chain functional group. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvants such as BCG (bacilli Calmette Guerin) and Corynebacterium parvum.

6.7 Methods of Using the Engineered Transaminase Enzymes and Compounds Prepared Therewith

[0263] In some embodiments, the transaminases described herein can be used in a method for performing processes shown in Schemes 1 to 3, *supra*, in which an amino group from an amino donor of general Formula II is transferred to an amino acceptor (ketone substrate) of general Formula I to produce a chiral amine. The reaction can produce the *R* chiral amine or *S* chiral amine in stereomeric excess. Generally, the method for performing the transamination reaction can comprise contacting or incubating the amino donor of Formula II and an amino acceptor of Formula I,



with an engineered transaminase polypeptide of the disclosure under reaction conditions suitable for converting the amine acceptor to the (*S*) or (*R*) chiral amine in stereomeric excess. Suitable groups for R^1 , R^2 , R^3 and R^4 are described above.

[0264] In some embodiments of the method, the amino acceptors can be selected from pyruvate, dihydronaphthalen-1(2*H*)-one, 1-phenylbutan-2-one, 3,3-dimethylbutan-2-one,

octan-2-one, ethyl 3-oxobutanoate, 4-phenylbutan-2-one, 1-(4-bromophenyl)ethanone, and 2-methyl-cyclohexanone and 7-methoxy-2-tetralone, 1-(6-methoxynaphthalen-2-yl)ethanone, 1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanone, 1-(4-phenoxyphenyl)ethanone, (*R*)-4-oxotetrahydro-2*H*-pyran-3-yl-benzoate, and (*R*)-3-(benzyloxy)dihydro-2*H*-pyran-4(3*H*)-one.

[0265] Accordingly, in some embodiments, the method comprises contacting dihydronaphthalen-1(2*H*)-one with a transaminase polypeptide of the disclosure in presence of an amino donor under suitable reaction conditions for the conversion of dihydronaphthalen-1(2*H*)-one to (*S*)-1,2,3,4-tetrahydronaphthalen-1-amine in enantiomeric excess. In some embodiments of the process, the product (*S*)-1,2,3,4-tetrahydronaphthalen-1-amine can be formed in at least 10% 20% 30% 40% 50% 60%, 70%, 80%, 90%, or 95% or more stereomeric excess.

[0266] In some embodiments, the method comprises contacting 1-phenylbutan-2-one with a transaminase polypeptide of the disclosure in presence of an amino donor under suitable reaction conditions for the conversion of 1-phenylbutan-2-one to (*S*)-1-phenylbutan-2-amine in enantiomeric excess. In some embodiments of the process, the product (*S*)-1-phenylbutan-2-amine can be formed in at least 10% 20% 30% 40% 50% 60%, 70%, 80%, 90%, or 95% or more stereomeric excess.

[0267] In some embodiments, the method comprises contacting 3,3-dimethylbutan-2-one with a transaminase polypeptide of the disclosure in presence of an amino donor under suitable reaction conditions for the conversion of 3,3-dimethylbutan-2-one to (*S*)-3,3-dimethylbutan-2-amine in enantiomeric excess. In some embodiments of the process, the product (*S*)-3,3-dimethylbutan-2-amine can be formed in at least 10% 20% 30% 40% 50% 60%, 70%, 80%, 90%, or 95% or more stereomeric excess.

[0268] In some embodiments, the method comprises contacting octan-2-one with a transaminase polypeptide of the disclosure in presence of an amino donor under suitable reaction conditions for the conversion of octan-2-one to (*S*)-octan-2-amine in enantiomeric excess. In some embodiments of the process, the product (*S*)-octan-2-amine can be formed in at least 10% 20% 30% 40% 50% 60%, 70%, 80%, 90%, or 95% or more stereomeric excess.

[0269] In some embodiments, the method comprises contacting 1-(4-bromophenyl)ethanone with a transaminase polypeptide of the disclosure in presence of an amino donor under suitable reaction conditions for the conversion of 1-(4-bromophenyl)ethanone to (*S*)-1-(4-bromophenyl)ethanamine in enantiomeric excess. In some embodiments of the process, the product (*S*)-1-(4-bromophenyl)ethanamine can be formed in at least 10% 20% 30% 40% 50% 60%, 70%, 80%, 90%, or 95% or more stereomeric excess.

[0270] In some embodiments, the method comprises contacting 4-phenylbutan-2-one with a transaminase polypeptide of the disclosure in presence of an amino donor under suitable reaction conditions for the conversion of 4-phenylbutan-2-one to (*R*)-4-phenylbutan-2-amine in enantiomeric excess. In some embodiments of the process, the product (*R*)-4-phenylbutan-2-

amine can be formed in at least 10% 20% 30% 40% 50% 60%, 70%, 80%, 90%, or 95% or more stereomeric excess.

[0271] In some embodiments, the method comprises contacting ethyl 3-oxobutanoate with a transaminase polypeptide of the disclosure in presence of an amino donor under suitable reaction conditions for the conversion of ethyl 3-oxobutanoate to (*R*)-ethyl 3-aminobutanoate in enantiomeric excess. In some embodiments of the process, the product (*R*)-ethyl 3-aminobutanoate can be formed in at least 10% 20% 30% 40% 50% 60%, 70%, 80%, 90%, or 95% or more stereomeric excess.

[0272] In some embodiments, the method comprises contacting ethyl 3-oxobutanoate with a transaminase polypeptide of the disclosure in presence of an amino donor under suitable reaction conditions for the conversion of ethyl 3-oxobutanoate to (*S*)-ethyl 3-aminobutanoate in enantiomeric excess. In some embodiments of the process, the product (*S*)-ethyl 3-aminobutanoate can be formed in at least 10% 20% 30% 40% 50% 60%, 70%, 80%, 90%, or 95% or more stereomeric excess.

[0273] In some embodiments, the method comprises contacting 1-(6-methoxynaphthalen-2-yl)ethanone with a transaminase polypeptide of the disclosure in presence of an amino donor under suitable reaction conditions for the conversion of 1-(6-methoxynaphthalen-2-yl)ethanone to (*S*)-1-(6-methoxynaphthalen-2-yl)ethanamine in stereomeric excess.

[0274] In some embodiments, the method comprises contacting 1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanone with a transaminase polypeptide of the disclosure in presence of an amino donor under suitable reaction conditions for the conversion of 1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanone to (*S*)-1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanamine in stereomeric excess.

[0275] In some embodiments, the method comprises contacting 1-(4-phenoxyphenyl)ethanone with a transaminase polypeptide of the disclosure in presence of an amino donor under suitable reaction conditions for the conversion of 1-(4-phenoxyphenyl)ethanone to (*S*)-1-(4-phenoxyphenyl)ethanamine in stereomeric excess.

[0276] In some embodiments, the method comprises contacting (*R*)-3-(benzyloxy)dihydro-2H-pyran-4(3H)-one with a transaminase polypeptide of the disclosure in presence of an amino donor under suitable reaction conditions for the conversion of (*R*)-3-(benzyloxy)dihydro-2H-pyran-4(3H)-one, to (3*S*,4*S*)-3-(benzyloxy)tetrahydro-2H-pyran-4-amine in stereomeric excess.

[0277] In some embodiments of the method, the amino donor comprises a compound of Formula II, as described herein. In some embodiments of the method, the amino donor is selected from isopropylamine (2-amino propane), α -phenethylamine, D-alanine, L-alanine, or D,L-alanine, particularly isopropylamine.

[0278] In the methods, any of the engineered transaminase described herein can be used to

convert the amine acceptor in presence of an amino donor to produce the chiral amine. In some aspects of the disclosure, the engineered transaminases used in the method comprises an amino acid sequence that is at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the sequence of SEQ ID NO: 10, 16, or 18. These engineered transaminases can have one or more of improved properties, including among others, thermostability, solvent stability, isopropylamine resistance, increased enzymatic activity, stereoselectivity, and/or substrate recognition/binding.


[0279] In some aspects of the disclosure of the method, the engineered transaminases can have one or more residue differences as compared to the naturally occurring transaminase of SEQ ID NO:2 or an a reference engineered transaminase, such as SEQ ID NO:18, at the following residue positions: X4, X9, X12, X21, X30, X31, X44, X45, X, 56, X57, X81, X82, X85, X86, X95, X112, X113, X127, X147, X153, X157, X166, X177, X181, X208, X211, X228, X233, X253, X272, X294, X297, X302, X311, X314, X316, X317, X318, X319, X320, X321, X324, X385, X391, X398, X407, X408, X409, X415, X417, X418, X420, X431, X434, X438, X444, and X446.

[0280] Guidance in choosing an engineered transaminase for a specific amine acceptor substrate is provided in the descriptions herein, such as in **Table 3** and **Table 4**, which shows the activities of various engineered transaminases for structurally different amino acceptor substrates. In some aspects of the disclosure the method usesan engineered transaminase comprising an amino acid sequence corresponding to SEQ ID NO: 4, 6, 8, 10, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88. 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170. 172, 174, 176, 178, 180, 182, 184, 186, 188, 190, 192, 194, 196, or 198.

[0281] In some embodiments of the method for enantiomeric enrichment of a chiral amine, additional quantities of the amino acceptor can be added (up to saturation) and/or the amino acceptor (ketone) formed can be continuously removed from the reaction mixture. Conversely, when the undesired chiral form of the amine is converted to the amino acceptor (ketone) byproduct and the desired chiral form is not, the latter can be readily isolated by conventional techniques. A partial separation can be effected by acidification, extraction with a hydrocarbon such as heptane to remove the ketone, rendering the aqueous phase basic, and re-extraction with a hydrocarbon such as heptane. When, on the other hand, both chiral forms of the amine are desired, the form which is converted to the ketone can be removed from the reaction mixture (or from the aqueous phase in a two phase mixture) and independently subjected to the action of an omega-transaminase in the presence of an amino donor to generate the same chiral form which was initially converted to the ketone.

[0282] As noted herein, the transaminases of the disclosure can be used to mediate the reverse reaction in Scheme 1, i.e., the conversion of the chiral amine of Formula III to the ketone of Formula I along with conversion of the ketone of Formula IV to the amine of Formula

II. Stereospecific conversion of either the (*R*) or (*S*) chiral amine can be used for the chiral resolution of mixtures of (*R*) and (*S*) amines. Accordingly, in some embodiments, the process for chiral resolution can comprise contacting a mixture of (*R*) and (*S*) chiral amine of formula IIIa (e.g., a racemic mixture) with a stereospecific transaminase of the disclosure in presence of ketone of Formula IV,



under suitable reaction conditions for formation of the ketone of Formula I and the amine of Formula II, thereby generating a mixture having a stereomeric excess of the chiral amine of Formula IIIb.

[0283] As is known by those of skill in the art, transamination reactions typically require a cofactor. Reactions catalyzed by the engineered transaminase enzymes described herein also typically require a cofactor, although many embodiments of the engineered transaminases require far less cofactor than reactions catalyzed with wild-type transaminase enzymes. As used herein, the term "cofactor" refers to a non-protein compound that operates in combination with a transaminase enzyme. Cofactors suitable for use with the engineered transaminase enzymes described herein include, but are not limited to, pyridoxal-5'-phosphate (also known as pyridoxal-phosphate, PLP, P5P). In some embodiments, the PLP cofactor is provided in the cell extract and does not need to be added. In some embodiments utilizing highly purified transaminase enzyme, the cofactor is added to the reaction mixture either at the beginning of the reaction or additional cofactor may be added during the reaction. In some embodiments, a different member of the vitamin B₆ family, such as pyridoxine, is used in place of PLP.

[0284] In some embodiments, the transaminase reactions can be carried in presence of reduced cofactor, nicotinamide adenine dinucleotide (NADH), which can limit the inactivation of the transaminase enzyme (see van Ophem et al., 1998, Biochemistry 37(9):2879-88). In some embodiments, a cofactor regeneration system, such as glucose dehydrogenase (GDH) and glucose or formate dehydrogenase and formate can be used to regenerate NADH in the reaction medium.

[0285] The transamination reactions described herein are generally carried out in a solvent. Suitable solvents include water, organic solvents (e.g., ethanol, dimethyl sulfoxide (DMSO), ethyl acetate, butyl acetate, 1-octanol, heptane, octane, methyl t-butyl ether (MTBE), toluene, and the like), ionic liquids (e.g., 1-ethyl 4-methylimidazolium tetrafluoroborate, 1-butyl-3-methylimidazolium tetrafluoroborate, 1-butyl-3-methylimidazolium hexafluorophosphate, and the like). In some embodiments, aqueous solvents, including water and aqueous co-solvent systems, are used.

[0286] Exemplary aqueous co-solvent systems comprises water and one or more organic

solvent. In general, an organic solvent component of an aqueous co-solvent system is selected such that it does not completely inactivate the transaminase enzyme. Appropriate co-solvent systems can be readily identified by measuring the enzymatic activity of the specified engineered transaminase enzyme with a defined substrate of interest in the candidate solvent system, utilizing an enzyme activity assay, such as those described herein.

[0287] The organic solvent component of an aqueous co-solvent system may be miscible with the aqueous component, providing a single liquid phase, or may be partly miscible or immiscible with the aqueous component, providing two liquid phases. Generally, when an aqueous co-solvent system is employed, it is selected to be biphasic, with water dispersed in an organic solvent, or vice-versa. Generally, when an aqueous co-solvent system is utilized, it is desirable to select an organic solvent that can be readily separated from the aqueous phase. In general, the ratio of water to organic solvent in the co-solvent system is typically in the range of from about 1% to about 99% (v/v) organic solvent to water. The co-solvent system may be pre-formed prior to addition to the reaction mixture, or it may be formed *in situ* in the reaction vessel.

[0288] The aqueous solvent (water or aqueous co-solvent system) may be pH-buffered or unbuffered. Generally, the transaminase reaction can be carried out at a pH of about 10 or below, usually in the range of from about 5 to about 10. In some embodiments, the transaminase reaction is carried out at a pH of about 9 or below, usually in the range of from about 5 to about 9. In some embodiments, the transaminase reaction is carried out at a pH of about 8 or below, often in the range of from about 5 to about 8, and usually in the range of from about 6 to about 8. The transaminase reaction may also be carried out at a pH of about 7.8 or below, or 7.5 or below. Alternatively, the transaminase reaction may be carried out at a neutral pH, i.e., about 7.

[0289] During the course of the transamination reactions, the pH of the reaction mixture may change. The pH of the reaction mixture may be maintained at a desired pH or within a desired pH range by the addition of an acid or a base during the course of the reaction. Alternatively, the pH may be controlled by using an aqueous solvent that comprises a buffer. Suitable buffers to maintain desired pH ranges are known in the art and include, for example, phosphate buffer, triethanolamine buffer, and the like. Combinations of buffering and acid or base addition may also be used.

[0290] In carrying out the transamination reactions described herein, the engineered transaminase enzyme may be added to the reaction mixture in the form of the purified enzymes, whole cells transformed with gene(s) encoding the enzymes, and/or cell extracts and/or lysates of such cells. The gene(s) encoding the engineered transaminase enzymes can be transformed into host cells separately or together into the same host cell. For example, in some embodiments one set of host cells can be transformed with gene(s) encoding one engineered transaminase enzyme and another set can be transformed with gene(s) encoding another engineered transaminase. Both sets of transformed cells can be utilized together in the reaction mixture in the form of whole cells, or in the form of lysates or extracts derived

therefrom. In other embodiments, a host cell can be transformed with gene(s) encoding multiple engineered transaminase enzymes. In some embodiments the engineered polypeptides can be expressed in the form of secreted polypeptides and the culture medium containing the secreted polypeptides can be used for the transaminase reaction.

[0291] Whole cells transformed with gene(s) encoding the engineered transaminase enzyme or cell extracts, lysates thereof, and isolated enzymes may be employed in a variety of different forms, including solid (e.g., lyophilized, spray-dried, and the like) or semisolid (e.g., a crude paste).

[0292] The cell extracts or cell lysates may be partially purified by precipitation (ammonium sulfate, polyethyleneimine, heat treatment or the like, followed by a desalting procedure prior to lyophilization (e.g., ultrafiltration, dialysis, and the like). Any of the cell preparations may be stabilized by crosslinking using known crosslinking agents, such as, for example, glutaraldehyde or immobilization to a solid phase (e.g., Eupergit C, and the like).

[0293] The solid reactants (e.g., enzyme, salts, etc.) may be provided to the reaction in a variety of different forms, including powder (e.g., lyophilized, spray dried, and the like), solution, emulsion, suspension, and the like. The reactants can be readily lyophilized or spray dried using methods and equipment that are known to those having ordinary skill in the art. For example, the protein solution can be frozen at -80°C in small aliquots, then added to a prechilled lyophilization chamber, followed by the application of a vacuum.

[0294] The quantities of reactants used in the transamination reaction will generally vary depending on the quantities of product desired, and concomitantly the amount of transaminase substrate employed. Those having ordinary skill in the art will readily understand how to vary these quantities to tailor them to the desired level of productivity and scale of production. In general, the transaminase substrates are kept at levels that achieve essentially complete or near complete conversion of the substrates into products.

[0295] The order of addition of reactants is not critical. The reactants may be added together at the same time to a solvent (e.g., monophasic solvent, biphasic aqueous co-solvent system, and the like), or alternatively, some of the reactants may be added separately, and some together at different time points. For example, the cofactor, transaminase, and transaminase substrate may be added first to the solvent.

[0296] For improved mixing efficiency when an aqueous co-solvent system is used, the transaminase, and cofactor may be added and mixed into the aqueous phase first. The organic phase may then be added and mixed in, followed by addition of the transaminase substrate. Alternatively, the transaminase substrate may be premixed in the organic phase, prior to addition to the aqueous phase

[0297] Suitable conditions for carrying out the transaminase-catalyzed transamination reactions described herein include a wide variety of conditions which can be readily optimized

by routine experimentation that includes, but is not limited to, contacting the engineered transaminase enzyme and substrate at an experimental pH and temperature and detecting product, for example, using the methods described in the Examples provided herein.

[0298] The transaminase catalyzed reaction is typically carried out at a temperature in the range of from about 15°C to about 60°C. For some embodiments, the reaction is carried out at a temperature in the range of from about 20°C to about 55°C. For some embodiments, the reaction is carried out at a temperature in the range of from about 35°C to about 50°C. In still other embodiments, it is carried out at a temperature in the range of from about 40°C to about 50°C. The reaction may also be carried out under ambient conditions.

[0299] The transamination reaction is generally allowed to proceed until essentially complete, or near complete, transformation of substrate is obtained. Transformation of substrate to product can be monitored using known methods by detecting substrate and/or product. Suitable methods include gas chromatography, HPLC, and the like. Conversion yields of the chiral amine product generated in the reaction mixture are generally greater than about 50%, may also be greater than about 60%, may also be greater than about 70%, may also be greater than about 80%, may also be greater than 90%, and are often greater than about 97%.

7. EXAMPLES

[0300] Various features and embodiments of the disclosure are illustrated in the following representative examples, which are intended to be illustrative, and not limiting.

Example 1: Screen for thermostable transaminase variants using isopropylamine as amine-donor.

[0301] The following method was used to screen transaminase libraries for a thermostable backbone from which most polypeptide variants were evolved. Biotransformations were performed in 96-well plates and contained the following: 1.0 M isopropylamine (pH 7.5) and 1.0 M pyruvate in 100 mM triethanolamine hydrochloride buffer (pH 7.5). The amine product (alanine) was then converted to its corresponding isoindole product using o-phthalaldehyde/N-acetylcysteine reagent (OPA), separated by HPLC, and quantified using external standards.

Reagents, per well:

Component	Volume/Well	Final conc.
Pyruvate, pH 7.5 (2.5M)	80 μ L	1.0 M
IPM (5.0M, pH 7.5)	40 $\square$ μ L	1.0 M
1.0M TEA, pH 7.5	20 $\square$ μ L	100 mM
Water	20 $\square$ μ L	

Component	Volume/Well	Final conc.
Lysate	40 μ L	
200 μ L/well final volume.		

[0302] Biotransformations. Lysate containing transaminase variants was added to 96-well round bottom plates (40 μ L/well). The plates were heat-sealed and incubated at 50°C for 24 h. After 24 h at 50°C, reactions were initiated by addition of a 160 μ L/well of a 50°C pre-heated solution of 1.25 M pyruvate, 1.25 M isopropylamine, and 125 mM triethanolamine (pH 7.5). The reactions were briefly agitated, heat sealed, and returned to the 50°C incubator. After 2 h, 10 μ L/well biotransformation was transferred to a 96-well plate containing 190 μ L/well aqueous 0.5 mM hydroxylamine. This hydroxylamine-quenched plate was heat-sealed and stored in a 4°C refrigerator for later analysis. The reaction plate was heat-sealed and returned to the 50°C incubator. After 24 h total reaction time, another 10 μ L/well aliquot of biotransformation was added to a new 96-well plate containing 190 μ L/well 0.5 mM hydroxylamine. The quenched biotransformations (both the 2 h and the 24 h timepoints) were then OPA derivatized and analyzed by HPLC. Quantification of alanine was performed as follows:

1. 20 μ L/well quenched biotransformation was transferred to an empty 96-well plate.
2. Six standards containing mixtures of known amounts of isopropylamine and alanine (50 mM total amine concentration) were prepared and 20 μ L of each standard was added to a second 96-well plate.
3. Both the quenched biotransformation plate and the standards plate were transferred to an HPLC equipped with an autosampler.
4. On-line OPA derivatization of each sample to the corresponding isoindole product was performed immediately prior to HPLC analysis by automated addition of 80 μ L OPA derivatization reagent to each 20 μ L aliquot.
5. The isoindole products were separated by HPLC and detected by absorbance at 320 nm.
6. Peak areas of the standards were used to generate a calibration curve.
7. Alanine product concentrations were calculated from the calibration curve and accounted for dilution steps.

Example 2: Generation of highly expressed *Vibrio fluvialis* transaminase variants.

[0303] A gene encoding the wild-type omega transaminase from *Vibrio fluvialis* SEQ ID NO:1 based on the reported amino acid sequence (Kim and/or Shin) was synthesized and cloned into an in-house pCK110900 vector system (United States Patent Application Publication 20060195947) and subsequently expressed in *E. coli* W3110*fhuA*. Two random mutagenesis libraries were generated, with individual colonies grown in 96-well plates. Crude lysate from the


cells was then used to screen for increased activity. Screening conditions were as follows: 1.0 M isopropylamine and 110 mM pyruvate in 100 mM triethanolamine buffer (pH 9) at 50°C for 4-4.5 h. Reactions were quenched by addition of 0.5 mM aqueous hydroxylamine hydrochloride and then analyzed by HPLC. Variants that produced significantly higher yields of the product (L-alanine) were sequenced and analyzed by SDS-PAGE, revealing that N-terminal mutations, both coding and non-coding (up to and including the 12th residue), are capable of greatly increasing expression levels.

Table 5

Nuc ID	Amino Acid Sequence Changes	Silent mutations	Expression Level
193	E12K; E302K	g309t	+++++
3		c18t; c286t	+++++
5	A9T	c852t; g861a	+++
7	P4R	g891c	+++++
11	A9T; F86Y; M294V;	c852t; g861a	+++++
13		g12a; g15a; c18t	+++++
195	S6R; A133T; Y184F; P252Q; I314F	t561c	+++
197	S6I		+++++
+++ : 1-5x increase over wt ++++ : 5-10x increase over wt +++++ : 10-20x increase over wt			

Example 3: Amination of ethyl 3-oxobutanoate using isopropylamine as amine-donor

[0304] The following example describes the conditions for performing the following reaction:



[0305] A high-throughput method was used to screen transaminase libraries for polypeptide variants using isopropylamine as the amine-donor. Biotransformations were performed in 96-well plates and contained the following: 50 mM isopropylamine (pH 7.0) and 50 mM ethyl 3-oxobutanoate in 100 mM triethanolamine hydrochloride buffer (pH 7.0) containing 5% ethanol. Amine products were converted to their corresponding isoindole products using o-phthalaldehyde/N-acetylcysteine reagent (OPA), separated by HPLC, and quantified using external standards.

[0306] Biotransformations were initiated by addition of 70 μL lysate containing 100 μM PLP to 30 μL "reaction mixture." Reaction mixture consisted of 167 mM isopropylamine (pH 7.0) and 167 mM ethyl 3-oxobutanoate in 333 mM triethanolamine buffer (pH 7.0) containing 16.7% ethanol. Biotransformations were agitated at room temperature for 17 h and then quenched by addition of 0.1 volume of 1.5 mM aqueous hydroxylamine hydrochloride.


[0307] Quantification of ethyl 3-aminobutanoate was performed as follows:

1. 1. 20 μL /well quenched biotransformation was transferred to an empty 96-well plate.
2. 2. Six standards containing mixtures of known amounts of isopropylamine and ethyl 3-aminobutanoate (50 mM total amine concentration) were prepared and 20 μL of each standard was added to a second 96-well plate.
3. 3. Both the quenched biotransformation plate and the standards plate were transferred to an HPLC equipped with an autosampler.
4. 4. On-line OPA derivatization of each sample to the corresponding isoindole product was performed immediately prior to HPLC analysis by automated addition of 80 μL OPA derivatization reagent to each 20 μL aliquot.
5. 5. The isoindole products were separated by HPLC and detected by absorbance at 320 nm.
6. 6. Peak areas of the standards were used to generate a calibration curve.
7. 7. ethyl 3-aminobutanoate product concentrations were calculated from the calibration curve and accounted for dilution steps.

The procedure in this example was also used to screen transaminase libraries with 3,4-dihydronaphthalen-1(2*H*)-one, 1-phenylbutan-2-one, 2-methylcyclohexanone, and 1-(4-bromophenyl)ethanone as the amino acceptor.

Example 4: Amination of octan-2-one using isopropylamine as amine-donor

[0308] The following example describes the conditions for performing the following reaction:



[0309] The following high-throughput method was used to screen transaminase libraries for variants using isopropylamine as the amine-donor. Biotransformations were performed in 96-well plates and contained the following: 200 mM isopropylamine (pH 7.0) and 100 mM octan-2-one in 100 mM triethanolamine hydrochloride buffer (pH 7.0) containing 5% ethanol. Amine products were converted to their corresponding isoindole products using o-phthalaldehyde/N-acetylcysteine reagent (OPA), separated by HPLC, and quantified using external standards.


[0310] Biotransformations were initiated by addition of 80 μL lysate containing 100 μM PLP to 20 μL "reaction mixture." Reaction mixture consisted of 1.0 M isopropylamine (pH 7.0) and 500 mM octan-2-one in 500 mM triethanolamine buffer (pH 7.0) containing 25% ethanol. Biotransformations were agitated at room temperature for 17 h and then quenched by addition of 3 volumes of 0.5 mM aqueous hydroxylamine hydrochloride.

[0311] Quantification of octan-2-amine was performed as follows:

1. 1. 20 μL /well quenched biotransformation was transferred to an empty 96-well plate.
2. 2. Six standards containing mixtures of known amounts of isopropylamine and octan-2-amine (50 mM total amine concentration) were prepared and 20 μL of each standard was added to a second 96-well plate.
3. 3. Both the quenched biotransformation plate and the standards plate were transferred to an HPLC equipped with an autosampler.
4. 4. On-line OPA derivatization of each sample to the corresponding isoindole product was performed immediately prior to HPLC analysis by automated addition of 80 μL OPA derivatization reagent to each 20 μL aliquot.
5. 5. The isoindole products were separated by HPLC and detected by absorbance at 320 nm.
6. 6. Peak areas of the standards were used to generate a calibration curve.
7. 7. Octan-2-amine product concentrations were calculated from the calibration curve and accounted for dilution steps.

Example 5: Amination of 3,3-dimethylbutan-2-one using isopropylamine as amine-donor

[0312] The following example describes the conditions for performing the following reaction,



[0313] The following high-throughput method was used to screen transaminase libraries for polypeptide variants using isopropylamine as the amine-donor. Biotransformations were performed in 96-well plates and contained the following: 200 mM isopropylamine (pH 7.0) and 100 mM 3,3-dimethylbutan-2-one in 100 mM triethanolamine hydrochloride buffer (pH 7.0) containing 5% ethanol. Amine products were converted to their corresponding isoindole products using o-phthalaldehyde/N-acetylcysteine reagent (OPA), separated by HPLC, and quantified using external standards.

[0314] Biotransformations were initiated by addition of 40 μL lysate containing 100 μM PLP to 60 μL "reaction mixture." Reaction mixture consisted of 333 mM isopropylamine (pH 7.0) and 167 mM 3,3-dimethylbutan-2-one in 167 mM triethanolamine buffer (pH 7.0) containing 8.3% ethanol. Biotransformations were agitated at room temperature for 17 h and then quenched by addition of 3 volumes of 0.5 mM aqueous hydroxylamine hydrochloride.


[0315] Quantification of 3,3-dimethylbutan-2-amine was performed as follows:

1. 1. 20 μL /well quenched biotransformation was transferred to an empty 96-well plate.
2. 2. Six standards containing mixtures of known amounts of isopropylamine and 3,3-dimethylbutan-2-amine (50 mM total amine concentration) were prepared and 20 μL of each standard was added to a second 96-well plate.
3. 3. Both the quenched biotransformation plate and the standards plate were transferred to an HPLC equipped with an autosampler.
4. 4. On-line OPA derivatization of each sample to the corresponding isoindole product was performed immediately prior to HPLC analysis by automated addition of 80 μL OPA derivatization reagent to each 20 μL aliquot.
5. 5. The isoindole products were separated by HPLC and detected by absorbance at 320 nm.
6. 6. Peak areas of the standards were used to generate a calibration curve.
7. 7. 3,3-dimethylbutan-2-amine product concentrations were calculated from the calibration curve and accounted for dilution steps.

[0316] The procedure in this example was also used to screen transaminase libraries with 4-phenylbutan-2-one as the amino acceptor.

Example 6: Amination of 1-(6-methoxynaphthalene-2-yl)ethanone to its corresponding S-amine.

[0317]



[0318] Reaction Conditions and Analysis: Transaminase variants were assayed for conversion of the ketone substrate 1-(6-methoxynaphthalene-2-yl)ethanone to its corresponding amine product (S)-1-(6-methoxynaphthalene-2-yl)ethanamine according to the following reaction conditions: 1 mg/mL (or 6.25 mg/mL) ketone substrate, 0.5 mg/mL PLP, 0.5 mg/mL NAD, 0.1 mg/mL lactate dehydrogenase (LDH), 1 mg/mL formate dehydrogenase (FDH), 90 mg/mL L-

alanine, 40 mg/mL formate, 5 vol% DMSO, buffer pH 7.5. The reaction mixture was allowed to react for 24 h with shaking at 30°C. Alternatively, GDH and glucose can replace FDH and formate for recycling NAD co-factor. Typical concentrations are as follows: 1 mg/mL GDH, 90 mg/mL glucose.

[0319] Additionally, a subset of variants were screened under the same conditions except using isopropylamine (iPr-NH₂) rather than L-alanine as the amine donor for the reaction. according to the following conditions: 1 mg/mL (or 5 mg/mL) ketone substrate, 0.5 mg/mL PLP, 1 M iPr-NH₂, 5-40 vol% DMSO in 100 mM triethanolamine buffer, pH 7.5 or 8.5. The reaction mixture was allowed to react for 24 h at 30°C.

[0320] Following the reaction, the mixture is extracted into an organic solvent phase (e.g., ethyl acetate, propyl acetate or methyl t-butyl ether) and analyzed by HPLC. Percent conversion is calculated based on peak areas as the ratio of amine product peak area to (ketone substrate peak area + amine product peak area).

[0321] As shown by the results listed in **Table 6**, reactions using alanine as amine donor resulted in substantially higher conversion than reactions using isopropylamine as donor. Reactions with alanine as donor and 1.0 mg/mL substrate loading were able to convert up to 21.3% of the substrate -(6-methoxynaphthalene-2-yl)ethanone to its corresponding amine product.

[0322] At least the following residue differences are associated with the ability to convert at least 2.0% substrate to its corresponding amine product: W57A/C/F/I/L/S, Y86A/F/G/H/S, V153A/C/Q/S, P233L/T, F317L, P318F/G/R, G320A, F321L, L417A/SAT7V, and F438L.

Table 6


SEQ ID NO:	Residue differences (as compared to SEQ ID NO:18)	iPr-NH ₂ donor	1.0 mg/mL substrate (alanine donor)	6.25 mg/mL substrate (alanine donor)
126	W57L; Y86F; V153S; P233T; L417T	+++	+++++	++++
154	W57C; Y86S; L417T	+++	+++++	++++
134	W57S; P233L; L417V		+++++	++++
140	W57A; V153S; P318G	++	+++++	+++++
148	W57A; V153C; F321L		+++++	++++
178	W57L; Y86S; V153A	++	+++++	+++
136	W57S; Y86G; L417C		+++++	++++
180	W57L; Y86F; P318F	++	+++++	++++
146	W57F; Y86F; V153Q	++	++++	++++
144	W57I; Y86F; G320A	+	++++	+++
166	W57C; Y86A; F317L		++++	+++

SEQ ID NO:	Residue differences (as compared to SEQ ID NO:18)	iPr-NH ₂ donor	1.0 mg/mL substrate (alanine donor)	6.25 mg/mL substrate (alanine donor)
114	W57L; L417C; F438L	++	+++	+++
150	Y86F; P318R; L417A		+++	+++
142	W57F; Y86H; V153Q		+++	+++
174	W57F; P318F; L417S		+++	++
110	Y86H; V153S; C181R; L417T	++	+++	++
172	V153S; F317L; L417C	+	+++	++
190	V153T; A228G; L417A		+++	++
116	W57F; M127L; L417C		+++	++
138	A228G; F317L; L417C		+++	++
184	P233L; F321L; L417I		+++	
152	A228G; P318G; L417C		+++	++
28	W57L	++	+++	+++
86	L417C	+	+++	+
92	Y86H; V153S; L417C	++	+++	+
162	Y86H; P233L; L417A		+++	++
48	Y86S; V153S		++	+
156	Y86F; V153C; V297A	++	++	++
182	V153S; A228G; L417V		++	
192	W57F; P318G; L417I		++	
78	L417A		++	+
26	Y86F		++	
84	L417I		++	
170	V153T; A228G; F321L		++	
186	F317L; P318R; L417T		++	
188	V153C; F317Y; H319Q		++	
24	Y86N		+	
46	W57I; V153S	+	+	+
88	Y86H; V153A; L417C		+	
102	V153A; P233T; L417C		+	
108	M95T; V153A; L417C		+	
168	Y86N; A228G; F317L		+	

SEQ ID NO:	Residue differences (as compared to SEQ ID NO:18)	iPr-NH ₂ donor	1.0 mg/mL substrate (alanine donor)	6.25 mg/mL substrate (alanine donor)
158	Y86S; V153T; V297A		+	
+ ≥ 0.2% ++ ≥ 0.5% +++ ≥ 1.0% ++++ ≥ 5.0% +++++ ≥ 10.0%				

Example 7: Transaminase variants capable of converting ketone substrate 1-(2,3-dihydrobenzo[b][1.4]dioxin-6-yl)ethanone to its corresponding S-amine product.

[0323]



[0324] Reaction conditions and analysis: Transaminase variants were assayed for conversion of the ketone substrate 1-(2,3-dihydrobenzo[b][1.4]dioxin-6-yl)ethanone to its corresponding amine product (S)-1-(2,3-dihydrobenzo[b][1.4]dioxin-6-yl)ethanamine according to the following reaction conditions: 1 mg/mL ketone substrate, 0.5 mg/mL PLP, 0.5 mg/mL NAD, 0.1 mg/mL LDH, 1 mg/mL FDH, 90 mg/mL L-alanine, 40 mg/mL formate, 5 vol% DMSO, buffer pH 7.5. The reaction mixture was allowed to react for 24 h at 30°C.

[0325] Following the reaction, the mixture is extracted into an organic solvent phase (e.g., ethyl acetate, propyl acetate or methyl t-butyl ether) and analyzed by HPLC. Percent conversion is calculated based on peak areas as the ratio of amine product peak area to (ketone substrate peak area + amine product peak area).


[0326] As shown by the results listed in **Table 7**, variants were able to convert up to 9.3% of ketone substrate 1-(2,3-dihydrobenzo[b][1.4]dioxin-6-yl)ethanone to its corresponding S-amine product. The positive control fluvialis TA of SEQ ID NO:18 exhibited only 0.7% conversion.

[0327] At least the following residue differences are associated with the ability to convert at least 2.0% substrate of formula (2) to its corresponding amine product: W57A/C/F/I/L/S, Y86F/G/S, V153A/C/Q/S, A228G, P233L/T, V297A, P318F/G, G320A, L417C/T/V, and F438L.

Table 7

SEQ ID NO:	Residue Differences (as compared to SEQ ID NO:18)	Conv. (%)
152	W57C; Y86S; L417T	+++
146	W57F; Y86F; V153Q	+++
126	W57L; Y86F; V153S; P233T; L417T	+++
156	Y86F; V153C; V297A	++
28	W57L	++
114	W57L; L417C; F438L	++
134	W57S; P233L; L417V	++
178	W57L; Y86S; V153A	++
136	W57S; Y86G; L417C	++
140	W57A; V153S; P318G	++
42	W57I	++
46	W57I; V153S	++
144	W57I; Y86F; G320A	++
180	W57L; Y86F; P318F	++
182	V153S; A228G; L417V	++
26	Y86F	++
166	W57C; Y86A; F317L	+
104	V153S; P233S	+
148	W57A; V153C; F321L	+
72	V153G	+
48	Y86S; V153S	+
84	L417I	+
110	Y86H; V153S; C181R; L417T	+
16	D21N; H45N; L177V; T208I; K211R; G324S; A391T	+
44	V153S	+
98	V31A	+
106	Y86H; V153A; A228G; L417I	+
164	V153C; P233L; P318R	+
+ ≥ 0.8% ++ ≥ 2.0% +++ ≥ 5.0%		

Example 8: Transaminase variants capable of converting ketone substrate 1-(4-phenoxyphenyl)ethanone to its corresponding S-amine product.



[0329] Reaction conditions and analysis for conversion of ketone substrate: Transaminase variants were assayed for conversion of the ketone substrate 1-(4-phenoxyphenyl)ethanone to its corresponding amine product (S)-1-(4-phenoxyphenyl)ethanamine according to the following reaction conditions: 1 mg/mL ketone substrate, 0.5 mg/mL PLP, 0.5 mg/mL NAD, 0.1 mg/mL LDH, 1 mg/mL FDH, 90 mg/mL L-alanine, 40 mg/mL formate, 5 vol% DMSO, buffer pH 7.5. The reaction mixture was allowed to react for 24 h at 30°C.

[0330] Following the reaction, the mixture is extracted into an organic solvent phase (e.g., ethyl acetate, propyl acetate or methyl t-butyl ether) and analyzed by HPLC. Percent conversion is calculated based on peak areas as the ratio of amine product peak area to (ketone substrate peak area + amine product peak area).

[0331] As shown by the results listed in **Table 8**, variants were able to convert up to 2.0% of ketone substrate 1-(4-phenoxyphenyl)ethanone to its corresponding S-amine product. The positive control fluvialis TA of SEQ ID NO:18 exhibited no detectable conversion under the assay conditions.

[0332] At least the following residue differences are associated with the ability to convert at least 0.5% substrate of formula (3) to its corresponding amine product: W57C/F/I/L, Y86H/F/S, V153A/C/Q/S, C181R, P233T, V297A, P318F, L417C/T, and F438L.


Table 8

SEQ ID NO:	Residue Differences (as compared to SEQ ID NO:18)	Conv. (%)
16	D21N; H45N; L177V; T208I; K211R; G324S; A391T	+
20	Y86S	+
26	Y86F	+
28	W57L	++
42	W57I	+
44	V153S	+
46	W57I; V153S	+
48	Y86S; V153S	+
72	V153G	+

SEQ ID NO:	Residue Differences (as compared to SEQ ID NO:18)	Conv. (%)
78	L417A	+
84	L417I	+
88	Y86H; V153A; L417C	+
90	Y86H; V153A	+
92	Y86H; V153S; L417C	+
98	V31A	+
104	V153S; P233S	+
106	Y86H; V153A; A228G; L417I	+
108	M95T; V153A; L417C	+
110	Y86H; V153S; C181R; L417T	+
114	W57L; L417C; F438L	+
126	W57L; Y86F; V153S; P233T; L417T	++
134	W57S; P233L; L417V	+
136	W57S; Y86G; L417C	+
140	W57A; V153S; P318G	+
142	W57F; Y86H; V153Q	+
144	W57I; Y86F; G320A	+
146	W57F; Y86F; V153Q	++
154	W57C; Y86S; L417T	++
156	Y86F; V153C; V297A	++
162	Y86H; P233L; L417A	+
164	V153C; P233L; P318R	+
166	W57C; Y86A; F317L	+
178	W57L; Y86S; V153A	++
180	W57L; Y86F; P318F	+
182	V153S; A228G; L417V	+
190	V153T; A228G; L417A	+
+ ≥ 0.2%		
++ ≥ 1.0%		

Example 9: Transaminase variants capable of converting ketone substrate (R)-4-oxotetrahydro-2H-pyran-3-yl benzoate to its corresponding S-amine product.

[0333]



[0334] Reaction conditions and analysis: Transaminase variants were assayed for conversion of the ketone substrate (*R*)-4-oxotetrahydro-2*H*-pyran-3-yl benzoate to its corresponding amine product (3*S*,4*S*)-4-aminotetrahydro-2*H*-pyran-3-yl benzoate according to the following reaction conditions: 1 mg/mL ketone substrate, 0.5 mg/mL PLP, 0.5 mg/mL NAD, 0.1 mg/mL LDH, 1 mg/mL FDH, 90 mg/mL L-alanine, 40 mg/mL formate, 5 vol% DMSO, buffer pH 7.5. The reaction mixture was allowed to react for 24 h at 30°C.

[0335] Following the reaction, the mixture is extracted into an organic solvent phase (e.g., ethyl acetate, propyl acetate or methyl t-butyl ether) and analyzed by HPLC. Percent conversion is calculated based on peak areas as the ratio of amine product peak area to (ketone substrate peak area + amine product peak area).

[0336] As shown by the results listed in **Table 9**, variants were able to convert up to 57% of ketone substrate (*R*)-4-oxotetrahydro-2*H*-pyran-3-yl benzoate to its corresponding S-amine product. The positive control fluvialis TA of SEQ ID NO:18 exhibited no detectable conversion.

[0337] At least the following residue differences are associated with the ability to convert at least 1% substrate (*R*)-4-oxotetrahydro-2*H*-pyran-3-yl benzoate to its corresponding S-amine product: W57A/C/I/L/S, Y86A/F/G/S, V153C/S, P233T, F317L, P318G, G320A, F321L, and L417T.


Table 9

SEQ ID NO:	Residue Differences (as compared to SEQ ID NO:18)	% Conversion
140	W57A; V153S; P318G	+++
148	W57A;V153C;F321L	+++
178	W57L;Y86S;V153A	++
144	W57I; Y86F; G320A	++
136	W57S; Y86G; L417C	++
126	W57L; Y86F; V153S; P233T; L417T	++
154	W57C; Y86S; L417T	++
166	W57C; Y86A; F317L	++
46	W57I; V153S	+

+ ≥ 1%
++ ≥ 5%
+++ ≥ 50%

Example 10: Transaminase variants capable of converting ketone substrate (*R*)-3-(benzyloxy)dihydro-2H-pyran-4(3H)-one to its corresponding S-amine product.

[0338]



[0339] Reaction conditions and analysis: Transaminase variants were assayed for conversion of the ketone substrate (*R*)-3-(benzyloxy)dihydro-2H-pyran-4(3H)-one to its corresponding amine product (3*S*,4*S*)-3-(benzyloxy)tetrahydro-2H-pyran-4-amine according to the following reaction conditions: 1 mg/mL ketone substrate, 0.5 mg/mL PLP, 0.5 mg/mL NAD, 0.1 mg/mL LDH, 1 mg/mL FDH, 90 mg/mL L-alanine, 40 mg/mL formate, 5 vol% DMSO, buffer pH 7.5. The reaction mixture was allowed to react for 24 h at 30°C.

[0340] Following the reaction, the mixture is extracted into an organic solvent phase (e.g., ethyl acetate, propyl acetate or methyl t-butyl ether) and analyzed by HPLC. Percent conversion is calculated based on peak areas as the ratio of amine product peak area to (ketone substrate peak area + amine product peak area).

[0341] As shown by the results listed in **Table 10**, variants were able to convert up to 100% of ketone substrate (*R*)-3-(benzyloxy)dihydro-2H-pyran-4(3H)-one to its corresponding amine product. The positive control fluvialis TA of SEQ ID NO:18 exhibited only 9% conversion.

[0342] At least the following residue differences are associated with the ability to convert at least 90% substrate (*R*)-3-(benzyloxy)dihydro-2H-pyran-4(3H)-one to its corresponding S-amine product: W57F/I/S; G81D; F85S; Y86F/G/H/N; M127L; V153C/Q/S; A228G; V297A; F317L; P318F/R; L417A/C/V; and F438L.

Table 10

SEQ ID NO:	Residue Differences (as compared to SEQ ID NO:18)	% Conversion
42	W57I	++++

SEQ ID NO:	Residue Differences (as compared to SEQ ID NO:18)	% Conversion
46	W57I; V153S	++++
92	Y86H; V153S; L417C	++++
116	W57F; M127L; L417C	++++
126	W57L; Y86F; V153S; P233T; L417T	++++
134	W57S; P233L; L417V	++++
136	W57S; Y86G; L417C	++++
140	W57A; V153S; P318G	++++
148	W57A; V153C; F321L	++++
154	W57C; Y86S; L417T	++++
166	W57C; Y86A; F317L	++++
178	W57L; Y86S; V153A	++++
180	W57L; Y86F; P318F	++++
28	W57L	++++
144	W57I; Y86F; G320A	++++
114	W57L; L417C; F438L	++++
22	G81D; Y86H	++++
146	W57F; Y86F; V153Q	+++
192	W57F; P318G; L417I	+++
110	Y86H; V153S; C181R; L417T	+++
106	Y86H; V153A; A228G; L417I	+++
156	Y86F; V153C; V297A	+++
172	V153S; F317L; L417C	+++
174	W57F; P318F; L417S	+++
142	W57F; Y86H; V153Q	+++
190	V153T; A228G; L417A	+++
186	F317L; P318R; L417T	+++
90	Y86H; V153A	+++
176	V153T; H319V; L417I	+++
182	V153S; A228G; L417V	+++
48	Y86S; V153S	+++
108	M95T; V153A; L417C	+++
88	Y86H; V153A; L417C	+++
128	F85S; V153A; P233T	+++
150	Y86F; P318R; L417A	+++

SEQ ID NO:	Residue Differences (as compared to SEQ ID NO:18)	% Conversion
98	V31A	+++
130	W57I; F85A; Y86H; L417C	+++
138	A228G; F317L; L417C	+++
36	F85S; V153A	+++
26	Y86F	+++
158	Y86S; V153T; V297A	+++
162	Y86H; P233L; L417A	++
102	V153A; P233T; L417C	++
122	F85A; W147G; V153A	++
40	F85A; V153A	++
86	L417C	++
100	I314V; D409G	++
132	F85A; V153S; L417S	++
170	V153T; A228G; F321L	++
50	P318G; T408A	++
184	P233L; F321L; L417I	++
52	Y82H	++
78	L417A	++
188	V153C; F317Y; H319Q	++
94	T30A	++
164	V153C; P233L; P318R	++
34	V153A; F317L; P318G	++
96	V44A; N166S	++
152	A228G; P318G; L417C	++
44	V153S	++
72	V153G	++
104	V153S; P233S	++
84	L417I	++
70	V153N	++
68	V153T	+
160	V153S; P318R; L417E	+
168	Y86N; A228G; F317L	+
60	F85T	+
20	Y86S	+

SEQ ID NO:	Residue Differences (as compared to SEQ ID NO:18)	% Conversion
118	I311V; I314T	+
30	Y82H; L417F	+
32	F85A; F317L	+
58	F85S	+
82	G320A	+
62	F85A	+
80	H319Q	+
74	F317M	+
76	F317Y	+
112	Y113C; K385R; L417C	+
38	Y113H; C407S	+
54	E12G; M434V	+
120	F112I; F317L	+
+ ≥ 10% ++ ≥ 20% +++ ≥ 50% ++++ ≥ 90%		

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2008127646A \[0006\]](#)
- [WO2005005633A \[0006\]](#)
- [US4950606A \[0006\]](#)
- [US5300437A \[0006\]](#)
- [US5169780A \[0006\]](#)
- [US5316943A \[0006\]](#)
- [US4518692A \[0006\]](#)

- [US4826766A \[0006\]](#)
- [US6197558B \[0006\]](#)
- [US4600692A \[0006\]](#)
- [WO2009008908A2 \[0211\]](#)
- [WO9600787A \[0226\]](#)
- [WO9533836A \[0238\]](#)
- [WO9522625A \[0256\] \[0256\]](#)
- [WO970078A \[0256\] \[0256\]](#)
- [WO9735966A \[0256\] \[0256\]](#)
- [WO9827230A \[0256\] \[0256\]](#)
- [WO0042651A \[0256\] \[0256\]](#)
- [WO0175767A \[0256\] \[0256\]](#)
- [US6537746B \[0256\] \[0256\]](#)
- [US20060195947A \[0303\]](#)

Non-patent literature cited in the description

- SHIN et al. Biosci. Biotechnol. Biochem., 2001, vol. 65, 1782-1788 [\[0003\]](#)
- YONAHARA et al. Agric. Biol. Chem., 1983, vol. 47, 102257-2265 [\[0004\]](#)
- SHIN et al. Biosci. Biotechnol. Biochem., 2001, vol. 65, 1782-1788 [\[0006\]](#)
- IWASAKI et al. Biotech. Lett., 2003, vol. 25, 1843-1846 [\[0006\]](#)
- IWASAKI et al. Appl. Microb. Biotech., 2004, vol. 69, 499-505 [\[0006\]](#)
- YUN et al. Appl. Environ. Microbiol., 2004, vol. 70, 2529-2534 [\[0006\]](#)
- HWANG et al. Enzyme Microbiol. Technol., 2004, vol. 34, 429-426 [\[0006\]](#)
- CHEMICAL ABSTRACTS, 54-47-7 [\[0030\]](#)
- SMITHWATERMAN Adv. Appl. Math., 1981, vol. 2, 482- [\[0034\]](#)
- NEEDLEMAN WUNSCH J. Mol. Biol., 1970, vol. 48, 443- [\[0034\]](#)
- PEARSON LIPMAN Proc. Natl. Acad. Sci. USA, 1988, vol. 85, 2444- [\[0034\]](#)
- Current Protocols, a joint venture between Current Protocols in Molecular Biology Greene Publishing Associates, Inc. 19950000 [\[0034\]](#)
- ALTSCHUL et al. J. Mol. Biol., 1990, vol. 215, 403-410 [\[0034\]](#)
- ALTSCHUL et al. Nucleic Acids Res., 1977, 3389-3402 [\[0034\]](#)
- HENIKOFF HENIKOFF Proc Natl Acad Sci USA, 1989, vol. 89, 10915- [\[0034\]](#)
- EISENBERG et al. J. Mol. Biol., 1984, vol. 179, 125-142 [\[0052\]](#) [\[0056\]](#)
- BALDINO et al. Methods Enzymology, vol. 168, 761-777 [\[0072\]](#)
- BOLTON et al. Proc. Natl. Acad. Sci. USA, 1962, vol. 48, 1390- [\[0072\]](#)
- BRESSLAUER et al. Proc. Natl. Acad. Sci USA, 1986, vol. 83, 8893-8897 [\[0072\]](#)
- FREIER et al. Proc. Natl. Acad. Sci USA, 1986, vol. 83, 9373-9377 [\[0072\]](#)
- KIERZEK et al. Biochemistry, vol. 25, 7840-7846 [\[0072\]](#)
- RYCHLIK et al. Nucleic Acids Res, 1990, vol. 18, 6409-6412 [\[0072\]](#)

- Nucleic Acids Res, 1991, vol. 19, 698- [\[0072\]](#)
- **SUGGS et al.** Developmental Biology Using Purified Genes Academic Press 1981 0000683-693 [\[0072\]](#)
- **WETMUR** Crit Rev Biochem Mol Biol, 1991, vol. 26, 227-259 [\[0072\]](#)
- **CODONWJOHN PEDENGCG** CodonPreferenceGenetics Computer Group Wisconsin Package [\[0076\]](#)
- **MCINERNEY, J. O** Bioinformatics, 1998, vol. 14, 372-73 [\[0076\]](#)
- **STENICO et al.** Nucleic Acids Res., 1994, 222437-46 [\[0076\]](#)
- **WRIGHT, F.** Gene, 1990, vol. 87, 23-29 [\[0076\]](#)
- **WADA et al.** Nucleic Acids Res., 1992, vol. 20, 2111-2118 [\[0076\]](#)
- **NAKAMURA et al.** Nucl. Acids Res., 2000, vol. 28, 292- [\[0076\]](#)
- **HENAUTDANCHIN et al.** Escherichia coli and Salmonella ASM Press 1996 00002047-2066 [\[0076\]](#)
- **MOUNT, D.** Bioinformatics: Sequence and Genome Analysis Cold Spring Harbor Laboratory Press 2001 0000 [\[0076\]](#)
- **UBERBACHER, E. C.** Methods Enzymol., 1996, vol. 266, 259-281 [\[0076\]](#)
- **TIWARI et al.** Comput. Appl. Biosci., 1997, vol. 13, 263-270 [\[0076\]](#)
- **SHIN et al.** Purification, characterization, and molecular cloning of a novel amine:pyruvate transaminase from Vibrio fluvialis JS17 Appl Microbiol Biotechnol., 2003, vol. 61, 5-6463-471 [\[0105\]](#) [\[0255\]](#)
- **FASMAN** CRC Practical Handbook of Biochemistry and Molecular Biology CRC Press 1989 00003-70 [\[0204\]](#)
- **HERMANSON, G.T.** Bioconjugate Techniques Academic Press 2008 0000 [\[0212\]](#)
- Bioconjugation Protocols: Strategies and Methods Methods in Molecular Biology Humana Press 2004 0000 [\[0212\]](#)
- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual Cold Spring Harbor Laboratory Press 2001 0000 [\[0224\]](#)
- Current Protocols in Molecular Biology Greene Pub. Associates 1998 0000 [\[0224\]](#)
- **VILLA-KAMAROFF et al.** Proc. Natl Acad. Sci. USA, 1978, vol. 75, 3727-3731 [\[0225\]](#)
- **DEBOER et al.** Proc. Natl Acad. Sci. USA, 1983, vol. 80, 21-25 [\[0225\]](#)
- **ROMANOS et al.** Yeast, 1992, vol. 8, 423-488 [\[0227\]](#)
- **GUOSHHERMAN** Mol Cell Bio, 1995, vol. 15, 5983-5990 [\[0232\]](#)
- **SIMONENPALVA** Microbiol Rev, 1993, vol. 57, 109-137 [\[0235\]](#)
- **EHRlich** Proc Natl Acad Sci. USA, 1978, vol. 75, 1433- [\[0249\]](#)
- **LATHE et al.** Gene, 1987, vol. 57, 193-201 [\[0251\]](#)
- **STEMMER** Proc Natl Acad Sci USA, 1994, vol. 91, 10747-10751 [\[0256\]](#) [\[0256\]](#)
- **ZHAO et al.** Nat. Biotechnol., 1998, vol. 16, 258-261 [\[0256\]](#)
- **CALDWELL et al.** PCR Methods Appl., 1994, vol. 3, S136-S140 [\[0256\]](#)
- **BLACK et al.** Proc Natl Acad Sci USA, 1996, vol. 93, 3525-3529 [\[0256\]](#)
- **LING et al.** Anal. Biochem., 1997, vol. 254, 2157-78 [\[0256\]](#)
- **DALE et al.** Oligonucleotide-directed random mutagenesis using the phosphorothioate method Methods Mol. Biol., 1996, vol. 57, 369-74 [\[0256\]](#)
- **SMITH** Ann. Rev. Genet., 1985, vol. 19, 423-462 [\[0256\]](#)
- **BOTSTEIN et al.** Science, 1985, vol. 229, 1193-1201 [\[0256\]](#)

- CARTER Biochem. J., 1986, vol. 237, 1-7 [0256]
- KRAMER et al. Cell, 1984, vol. 38, 879-887 [0256]
- WELLS et al. Gene, 1985, vol. 34, 315-323 [0256]
- MINSHULL et al. Curr Opin Chem Biol, 1999, vol. 3, 284-290 [0256]
- CHRISTIANS et al. Nature Biotech, 1999, vol. 17, 259-264 [0256]
- CRAMERI et al. Nature, 1998, vol. 391, 288-291 [0256]
- CRAMERI et al. Nature Biotech, 1997, vol. 15, 436-438 [0256]
- ZHANG et al. Proc Natl Acad Sci USA, 1997, vol. 94, 45-44509- [0256]
- CRAMERI et al. Nature Biotech, 1996, vol. 14, 315-319 [0256]
- STEMMER Nature, 1994, vol. 370, 389-391 [0256]
- BEAUCAGE et al. Tet Lett, 1981, vol. 22, 1859-69 [0259]
- MATTHES et al. EMBO J., 1984, vol. 3, 801-05 [0259]
- VAN OPHEM et al. Biochemistry, 1998, vol. 37, 92879-88 [0284]

Patentkrav**1.** Transaminaseenzym, omfattende:

et polypeptid med transaminaseaktivitet med forøget varmemestabilitet ved 50°C

- 5 sammenlignet med polypeptidet af SEQ ID NO: 2, hvor polypeptidet omfatter en aminosyresekvens, som er mindst 80% identisk med SEQ ID NO: 18 og har en eller flere restforskelle sammenlignet med SEQ ID NO: 2, hvor X9 er T; X86 er Y og X294 er V.

- 10 **2.** Transaminaseenzymet ifølge krav 1, hvor polypeptidet har mindst ca. 10% restaktivitet i omdannelsen af pyruvat til L-alanin under tilstedeværelsen af aminodonorisopropylamin efter behandling af polypeptidet ved 50°C i 23 timer.

- 3.** Transaminaseenzymet ifølge krav 1 eller 2, som omfatter en aminosyresekvens
15 svarende til SEQ ID NO: 10, 16 eller 18.

4. Transaminaseenzymet ifølge krav 1 eller 2:

- i) yderligere med forøget transaminaseaktivitet sammenlignet med SEQ ID
20 NO: 2 til omdannelse af et aminoacceptorsubstrat til det tilsvarende chirale aminoprodukt;

ii) hvor polypeptidet er i stand til at omdanne aminoacceptorsubstratet til det tilsvarende aminoprodukt med forbedret *R*-enantioselektivitet sammenlignet med SEQ ID NO: 18;

- iii) som er i stand til at omdanne aminoacceptorsubstratet valgt fra 3,4-dihydronaphthalen-1(2*H*)-on, 1-phenylbutan-2-on, 3,3-dimethylbutan-2-on, octan-2-on, ethyl-3-oxobutanoat, 4-phenylbutan-2-on og 1-(4-bromphenyl)ethanon til det tilsvarende aminoprodukt ved en rate, som er større end polypeptidet af SEQ ID NO: 2 eller 18;

- iv) som er i stand til stereoselektivt at omdanne 1-(6-methoxynaphthalen-2-yl)ethanon, til produktet (*S*)-1-(6-methoxynaphthalen-2-yl)ethanamin
30 ved en rate, som er større end polypeptidet af SEQ ID NO: 2 eller 18;

v) som er i stand til stereoselektivt at omdanne 1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanon, til produktet (S)-1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanamin ved en rate som er større end polypeptidet af SEQ ID NO: 2 eller 18;

5 vi) som er i stand til stereoselektivt at omdanne 1-(4-phenoxyphenyl)ethanon, til produktet (S)-1-(4-phenoxyphenyl)ethanamin ved en rate som er større end polypeptidet af SEQ ID NO: 2 eller 18;

vii) som er i stand til stereoselektivt at omdanne (*R*)-4-oxotetrahydro-2*H*-pyran-3-yl-benzoat til produktet (3*S*,4*S*)-3-aminotetrahydro-2*H*-pyran-3-yl-benzoat ved en rate, som er større end polypeptidet af SEQ ID NO: 2
10 eller 18; eller

viii) som er i stand til stereoselektivt at omdanne (*R*)-3-benzyloxydihydro-2*H*-pyran-4(3*H*)-on, til produktet (3*S*,4*S*)-3-benzyloxytetrahydro-2*H*-pyran-4-amin ved en rate, som er større end polypeptidet af SEQ ID NO: 2
15 eller 18.

5. Transaminaseenzymet ifølge krav 1 eller 2:

i) yderligere med forøget transaminaseaktivitet sammenlignet med SEQ ID NO: 2 til omdannelse af et aminoacceptorsubstrat til det tilsvarende chirale aminoprodukt, og som omfatter en aminosyresekvens svarende til SEQ ID
20 NO: 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 176, 178,
25 180, 182, 184, 186, 188, 190 eller 192;

ii) hvor polypeptidet er i stand til at omdanne aminoacceptorsubstratet til det tilsvarende aminoprodukt med forbedret *R*-enantioselektivitet sammenlignet med SEQ ID NO: 18, og som omfatter en aminosyresekvens svarende til SEQ ID NO: 22, 30, 32, 34, 36, 40, 44, 48, 56, 58, 60, 62, 64,
30 66, 68, 70, 72, 76, 84, 88, 90, 92, 104, 108, 110, 120, 122, 124, 126, 128, 132, 160 eller 176;

- iii) som er i stand til at omdanne aminoacceptorsubstratet valgt fra 3,4-dihydronaphthalen-1(2*H*)-on, 1-phenylbutan-2-on, 3,3-dimethylbutan-2-on, octan-2-on, ethyl 3-oxobutanoat, 4-phenylbutan-2-on og 1-(4-bromphenyl)ethanon til det tilsvarende aminprodukt ved en rate, som er større end polypeptidet af SEQ ID NO: 2 eller 18, og
- 5
- a) som omfatter en aminosyresekvens svarende til SEQ ID NO: 38, 42, 44, 52, 56, 66, 74, 76, 78, 80, 82, 84, 94, 96, 98, 100, 104, 118, 160, 164, 172, 184 eller 188;
- b) hvor polypeptidet er i stand til at omdanne
- 10 aminoacceptorsubstratet 1-phenylbutan-2-on til aminoproduktet (S)-1-phenylbutan-2-amin med forbedret S-enantioselektivitet sammenlignet med SEQ ID NO: 2 eller 18;
- c) hvor polypeptidet er i stand til at omdanne
- 15 aminoacceptorsubstratet 3,3-dimethylbutan-2-on til aminoproduktet (S)-3,3-dimethylbutan-2-amin med forbedret S-enantioselektivitet sammenlignet med SEQ ID NO: 2 eller 18;
- d) hvor polypeptidet er i stand til at omdanne
- 20 aminoacceptorsubstratet octan-2-on til aminoproduktet (S)-octan-2-amin med forbedret S-enantioselektivitet sammenlignet med SEQ ID NO: 2 eller 18;
- e) hvor polypeptidet er i stand til at omdanne
- 25 aminoacceptorsubstratet 1-(4-bromphenyl)ethanon til aminoproduktet (S)-1-(4-bromphenyl)ethanamin med forbedret S-enantioselektivitet sammenlignet med SEQ ID NO: 2 eller 18; eller
- f) som er i stand til stereoselektivt at omdanne 4-phenylbutan-2-on til (R)-4-phenylbutan-2-amin ved en rate, som er større end polypeptidet af SEQ ID NO: 2 eller 18;
- iv) som er i stand til stereoselektivt at omdanne 1-(6-methoxynaphthalen-2-yl)ethanon til produktet (S)-1-(6-methoxynaphthalen-2-yl)ethanamin
- 30 ved en rate, som er større end polypeptidet af SEQ ID NO: 2 eller 18, og

som omfatter en aminosyresekvens svarende til SEQ ID NO: 24, 26, 28, 46, 48, 78, 84, 86, 88, 92, 102, 108, 110, 114, 116, 126, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 162, 166, 168, 170, 172, 174, 178, 180, 182, 184, 186, 188, 190 eller 192;

5 v) som er i stand til stereoselektivt at omdanne 1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanon til produktet (S)-1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanamin ved en rate, som er større end polypeptidet af SEQ ID NO: 2 eller 18, og som omfatter en aminosyresekvens svarende til SEQ ID NO: 16, 26, 28, 42, 44, 46, 48, 72, 84, 98,
10 104, 106, 110, 114, 126, 134, 136, 140, 144, 146, 148, 152, 156, 164, 166, 178, 180 eller 182;

vi) som er i stand til stereoselektivt at omdanne 1-(4-phenoxyphenyl)ethanon til produktet (S)-1-(4-phenoxyphenyl)ethanamin ved en rate, som er større end polypeptidet af SEQ ID NO: 2 eller 18, og
15 som omfatter en aminosyresekvens svarende til SEQ ID NO: 16, 20, 26, 28, 42, 44, 46, 48, 72, 78, 84, 88, 90, 92, 98, 104, 106, 108, 110, 114, 126, 134, 136, 140, 142, 144, 146, 154, 156, 162, 164, 166, 178, 180, 182 eller 190;

vii) som er i stand til stereoselektivt at omdanne (R)-4-oxotetrahydro-2H-pyran-3-yl-benzoat til produktet (3S,4S)-3-aminotetrahydro-2H-pyran-3-yl benzoat ved en rate, som er større end polypeptidet af SEQ ID NO: 2 eller 18, og som omfatter en aminosyresekvens svarende til SEQ ID NO: 46,
20 126, 136, 140, 144, 148, 154, 166 og 178; eller

viii) som er i stand til stereoselektivt at omdanne (R)3-benzyloxydihydro-2H-pyran-4(3H)-on til produktet (3S,4S)-3-benzyloxytetrahydro-2H-pyran-4-amin ved en rate, som er større end polypeptidet af SEQ ID NO: 2 eller 18, og som omfatter en aminosyresekvens svarende til SEQ ID NO: 20, 22, 26, 28, 30, 32, 34, 36, 40, 42, 44, 46, 48, 50, 52, 58, 60, 68, 70, 72, 78, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 114,
25 116, 118, 122, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172; 174, 176, 178, 180, 182, 184, 186, 188, 190 eller 192.

6. Transaminaseenzymet af ifølge:

- i) krav 5(iii)(b) som omfatter en aminosyresekvens svarende til SEQ ID NO: 22, 88, 90, 92, 106, 110, 138, 140, 142, 148, 150, 152, 160, 164, 168, 170, 172, 178, 182, 188 eller 190;
- 5 ii) krav 5(iii)(c) som omfatter en aminosyresekvens svarende to SEQ ID NO: 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 134, 140, 142, 144, 146, 152, 154, 156, 158, 162, 164, 168, 170, 172, 174, 176, 10 178, 180, 182, 184, 186, 188, 190 eller 192;
- iii) krav 5(iii)(d) som omfatter en aminosyresekvens svarende til SEQ ID NO: 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 134, 136, 15 138, 140, 142, 144, 146, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 178, 180, 182, 184, 186, 188, 190 eller 192;
- iv) krav 5(iii)(e) som omfatter en aminosyresekvens svarende til SEQ ID NO: 20, 22, 24, 26, 28, 30, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 60, 62, 64, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 20 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 136, 138, 144, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 176, 178, 182, 184, 186, 188, 190 og 192; eller
- v) krav 5(iii)(f) som omfatter en aminosyresekvens svarende til SEQ ID NO: 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 25 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 178, 182, 184, 186, 188, 190 eller 192;

7. Transaminaseenzymet ifølge et hvilket som helst af kravene 1-6, hvor
 30 polypeptidet har forbedret ekspresion i en værtselle sammenlignet med SEQ ID

NO: 2.

8. Polynukleotid som koder for polypeptidet ifølge et hvilket som helst af kravene 1-7.

5


9. Polynukleotidet ifølge krav 8, som omfatter en polynukleotidsekvens svarende til SEQ ID NO: 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 159, 161, 163, 165, 167, 169, 171, 173, 175, 177, 179, 181, 183, 185, 187, 189, 191, 193, 195 eller 197.

10. Ekspressionsvektor omfattende polynukleotidet ifølge krav 8 eller krav 9.

15

11. Værtselle omfattende polynukleotidet ifølge krav 8 eller 9.

12. Fremgangsmåde til omdannelsen af et substrat med formel I til det chirale aminprodukt med formel III i stereomert overskud,



20

hvor

det chirale carbonatom af det chirale aminprodukt er markeret med en *;

hver af R^1 , R^2 , R^3 og R^4 , når taget uafhængigt, er en ikke-substitueret eller substitueret alkyl-, alkylaryl- eller arylgruppe, hvor R^1 er forskellig fra R^2 i struktur eller chiralitet, eller R^1 og R^2 tilsammen er en carbonhydridkæde med 4 eller flere carbonatomer indeholdende et chiralitetscenter,

25

5 hvor fremgangsmåden omfatter at bringe substratet med formel I i kontakt med et transaminasepolypeptid ifølge et hvilket som helst af kravene 1 til 7 i tilstedeværelsen af en aminodonor under egnede reaktionsbetingelser for omdannelsen af substratet til produktet med formel III i stereomert overskud.

13. Fremgangsmåden ifølge krav 12, hvor:

- i) substratet med formel I er 3,4-dihydronaphthalen-1(2*H*)-on, som omdannes til produktet (S)-1,2,3,4-tetrahydronaphthalen-1-amin i stereomert overskud;
- 10 ii) substratet med formel I er 1-phenylbutan-2-on, som omdannes til produktet (S)-1-phenylbutan-2-amin i stereomert overskud;
- iii) substratet med formel I er 3,3-dimethylbutan-2-on, som omdannes til produktet (S)-3,3-dimethylbutan-2-amin i stereomert overskud;
- 15 iv) substratet med formel I er octan-2-on, som omdannes til produktet (S)-octan-2-amin i stereomert overskud;
- v) substratet med formel I er 1-(4-bromphenyl)ethanon, som omdannes til produktet (S)-1-(4-bromphenyl)ethanamin i stereomert overskud;
- vi) substratet med formel I er 4-phenylbutan-2-on, som omdannes til produktet (R)-4-phenylbutan-2-amin i stereomert overskud;
- 20 vii) substratet med formel I er ethyl-3-oxobutanoat, som omdannes til produktet (R)-ethyl 3-aminobutanoat i stereomert overskud;
- viii) substratet med formel I er ethyl-3-oxobutanoat, som omdannes til produktet (S)-ethyl 3-aminobutanoat i stereomert overskud;
- 25 ix) substratet med formel I er 1-(6-methoxynaphthalen-2-yl)ethanon, som omdannes til produktet (S)-1-(6-methoxynaphthalen-2-yl)ethanamin i stereomert overskud;

x) substratet med formel I er 1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanon, som omdannes til produktet (S)-1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanamin i stereomert overskud;

5 xi) substratet med formel I er 1-(4-phenoxyphenyl)ethanon, som omdannes til produktet (S)-1-(4-phenoxyphenyl)ethanamin i stereomert overskud;

xii) substratet med formel I er (R)-4-oxotetrahydro-2H-pyran-3-yl-benzoat, som omdannes til produktet (3S,4S)-3-aminotetrahydro-2H-pyran-3-yl benzoat i stereomert overskud; eller

10 xiii) substratet med formel I er (R)-3-benzyloxydihydro-2H-pyran-4(3H)-on, som omdannes til produktet (3S,4S)-3-benzyloxytetrahydro-2H-pyran-4-amin i stereomert overskud.

14. Fremgangsmåden ifølge krav 12, hvor:

15 i) substratet med formel I er 3,4-dihydronaphthalen-1(2H)-on, som omdannes til produktet (S)-1,2,3,4-tetrahydronaphthalen-1-amin i stereomert overskud, hvor transaminasepolypeptidet omfatter en aminosyresekvens svarende til SEQ ID NO: 38, 42, 44, 52, 56, 66, 74, 76, 78, 80, 82, 84, 94, 96, 98, 100, 104, 118, 160, 164, 172, 184 eller 188;

20 ii) substratet med formel I er 1-phenylbutan-2-on, som omdannes til produktet (S)-1-phenylbutan-2-amin i stereomert overskud, hvor transaminasepolypeptidet omfatter en aminosyresekvens svarende til SEQ ID NO: 22, 88, 90, 92, 106, 110, 138, 140, 142, 148, 150, 152, 160, 164, 168, 170, 172, 178, 182, 188 eller 190;

25 iii) substratet med formel I er 3,3-dimethylbutan-2-on, som omdannes til produktet (S)-3,3-dimethylbutan-2-amin i stereomert overskud, hvor transaminasepolypeptidet omfatter en aminosyresekvens svarende til SEQ ID NO: 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 134,

140, 142, 144, 146, 152, 154, 156, 158, 162, 164, 168, 170, 172, 174, 176, 178, 180, 182, 184, 186, 188, 190 eller 192;

iv) substratet med formel I er octan-2-on, som omdannes til produktet (S)-octan-2-amin i stereomert overskud, hvor transaminasepolypeptidet
 5 omfatter en aminosyresekvens svarende til SEQ ID NO: 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 134, 136, 138, 140, 142, 144, 146, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 178,
 10 180, 182, 184, 186, 188, 190 eller 192;

v) substratet med formel I er 1-(4-bromphenyl)ethanon, som omdannes til produktet (S)-1-(4-bromphenyl)ethanamin i stereomert overskud, hvor transaminasepolypeptidet omfatter en aminosyresekvens svarende til SEQ
 ID NO: 20, 22, 24, 26, 28, 30, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54,
 15 56, 60, 62, 64, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 136, 138, 144, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 176, 178, 182, 184, 186, 188, 190 eller 192;

vi) substratet med formel I er 4-phenylbutan-2-on, som omdannes til
 20 produktet (R)-4-phenylbutan-2-amin i stereomert overskud, hvor transaminasepolypeptidet omfatter en aminosyresekvens svarende til SEQ ID NO: 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 134,
 25 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 178, 182, 184, 186, 188, 190 eller 192;

vii) substratet med formel I er ethyl-3-oxobutanoat, som omdannes til produktet (R)-ethyl 3-aminobutanoat i stereomert overskud, hvor transaminasepolypeptidet omfatter en aminosyresekvens svarende til SEQ
 ID NO: 32, 34, 36, 40, 44, 56, 58, 60, 62, 64, 66, 68, 70, 72, 86, 88, 90,
 30 92, 122, 126, 128, 130 eller 132;

- viii) substratet med formel I er ethyl-3-oxobutanoat, som omdannes til produktet (S)-ethyl-3-aminobutanoat i stereomert overskud, hvor transaminasepolypeptidet omfatter en aminosyresekvens svarende til SEQ ID NO: 20, 22, 24, 26, 28, 30, 38, 42, 46, 48, 50, 52, 54, 74, 78, 80, 82, 94, 96, 98, 100, 102, 106, 112, 114, 116, 118, 120, 124, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 162, 164, 166, 168, 170, 172, 174, 178, 180, 182, 184, 186, 188, 190 eller 192;
- ix) substratet med formel I er 1-(6-methoxynaphthalen-2-yl)ethanon, som omdannes til produktet (S)-1-(6-methoxynaphthalen-2-yl)ethanamin i stereomert overskud, transaminasepolypeptidet omfatter en aminosyresekvens svarende til SEQ ID NO: 24, 26, 28, 46, 48, 78, 84, 86, 88, 92, 102, 108, 110, 114, 116, 126, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 162, 166, 168, 170, 172, 174, 178, 180, 182, 184, 186, 188, 190 eller 192;
- x) substratet med formel I er 1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanon, som omdannes til produktet (S)-1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)ethanamin i stereomert overskud, hvor transaminasepolypeptidet omfatter en aminosyresekvens svarende til SEQ ID NO: 16, 26, 28, 42, 44, 46, 48, 72, 84, 98, 104, 106, 110, 114, 126, 134, 136, 140, 144, 146, 148, 152, 156, 164, 166, 178, 180 eller 182;
- xi) substratet med formel I er 1-(4-phenoxyphenyl)ethanon, som omdannes til produktet (S)-1-(4-phenoxyphenyl)ethanamin i stereomert overskud, transaminasepolypeptidet omfatter en aminosyresekvens svarende til SEQ ID NO: 16, 20, 26, 28, 42, 44, 46, 48, 72, 78, 84, 88, 90, 92, 98, 104, 106, 108, 110, 114, 126, 134, 136, 140, 142, 144, 146, 154, 156, 162, 164, 166, 178, 180, 182 eller 190;
- xii) substratet med formel I er (R)-4-oxotetrahydro-2H-pyran-3-yl-benzoat, som omdannes til produktet (3S,4S)-3-aminotetrahydro-2H-pyran-3-yl benzoat i stereomert overskud, hvor transaminasepolypeptidet omfatter en aminosyresekvens svarende til SEQ ID NO: 46, 126, 136, 140, 144, 148, 154, 166 eller 178; eller

xiii) substratet med formel I er (*R*)-3-benzyloxydihydro-2*H*-pyran-4(3*H*)-on, som omdannes til produktet (3*S*,4*S*)-3-benzyloxytetrahydro-2*H*-pyran-4-amin i stereomert overskud, hvor transaminasepolypeptidet omfatter en aminosyresekvens svarende til SEQ ID NO: 20, 22, 26, 28, 30, 32, 34, 36, 40, 42, 44, 46, 48, 50, 52, 58, 60, 68, 70, 72, 78, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 114, 116, 118, 122, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172; 174, 176, 178, 180, 182, 184, 186, 188, 190 eller 192.

10 **15.** Fremgangsmåden ifølge krav 12:

(i) hvor aminodonoren er valgt fra gruppen bestående af isopropylamin, α -phenethylamin, D-alanin eller L-alanin;

(ii) hvor reaktionsbetingelsen er:

i) 20°C til 55°C; eller

15 ii) 35°C til 50°C;

(iii) hvor aminoacceptor-biproduktet fjernes; eller

(iv) som udføres under tilstedeværelsen af reduceret nikotinamidadenin-dinukleotid (NADH).

DRAWINGS




FIG. 1