

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 March 2005 (10.03.2005)

PCT

(10) International Publication Number
WO 2005/021779 A1

(51) International Patent Classification⁷: C12Q 1/00, (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US (patent), UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number: PCT/US2004/028238

(22) International Filing Date: 30 August 2004 (30.08.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 10/652,732 29 August 2003 (29.08.2003) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application: US 10/652,732 (CIP)
Filed on 29 August 2003 (29.08.2003)

(71) Applicant (for all designated States except US): PERKINELMER LAS, INC. [US/US]; 549 Albany Street, Boston, MA 02118 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): CERDA, Blas [MX/US]; 9 Eben Street, Milford, MA 01757 (US).

(74) Agent: GUY, Pamela, M.; 549 Albany Street, Boston, MA 02118 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2005/021779 A1

(54) Title: MASS SPECTROMETRY METHODS FOR SIMULTANEOUS DETECTION OF METABOLIC ENZYME ACTIVITY AND METABOLITE LEVELS

(57) **Abstract:** The present invention provides a method for detecting a metabolic disorder in an individual. The method involves (a) contacting a sample containing (i) one or more metabolically indicative enzymes and (ii) one or more metabolic analytes, with one or more substrates for said one or more enzymes to produce a reaction admixture, under conditions in which at least one of said enzymes is capable of acting on a corresponding substrate to generate at least one product; (b) contacting said reaction admixture with a reagent that inhibits the ability of said one or more enzymes to act on a corresponding substrate, wherein said one or more metabolic analytes and said at least one product are soluble in said reagent; to produce a test sample and (c) determining the presence or amount of said one or more metabolic analytes and said at least one product contained in said test sample using mass spectrometry, wherein a determined presence or amount of said one or more metabolic analytes and said at least one product correlates with presence or absence of said metabolic disorder.

MASS SPECTROMETRY METHODS FOR SIMULTANEOUS DETECTION OF
METABOLIC ENZYME ACTIVITY AND METABOLITE LEVELS

5

BACKGROUND OF THE INVENTION

The present invention relates generally to diagnostic medicine and more specifically to methods for diagnosing metabolic disorders.

10 Metabolism is the process of building the body's molecular structures from nutrients (anabolism) and breaking them down for energy (catabolism). Metabolic processes result in growth, produce energy, eliminate wastes, and control distribution of nutrients in the body. Homeostasis, or a steady state, in the body is a result of normal metabolism.

15 Enzymes play an important role in metabolism because they catalyze conversion of one molecule to another during metabolic processes. When a metabolic enzyme is faulty or present in an abnormal amount in an individual, a metabolic disorder can result. Metabolic disorders are frequently due to genetic inheritance that leads to absence or overproduction of an enzyme or production of a faulty enzyme.

20 As an example, deficiencies produced by an inactive gene can prevent the body from making an enzyme (or enzymes) needed to break down certain amino acids or types of fats. Such an enzyme deficiency typically prevents normal metabolism of a nutrient, causing the nutrient or its metabolite to build up in the body to toxic levels. An enzyme deficiency can also cause nutrient deficiencies when an enzyme is unable to make a normal end-product of a metabolic process. Nutrient deficiencies resulting from metabolic disorders can lead to impaired growth and development, and other 25 severe health problems.

30 Many inherited metabolic disorders are fatal in the first weeks or months of postnatal life, for example, severe defects in the conversion of pyruvate to acetyl

coenzyme A (CoA), some urea cycle defects, and severe defects in the processing of fructose. Infants and children with a treatable metabolic disorder are often identified by newborn screening in developed countries. For example, testing for phenylketonuria is routine practice in developed countries. Phenylketonuria is an 5 example of a relatively common inherited metabolic disorder that occurs in about one out of 16,000 live births in the United States. Individuals having phenylketonuria do not produce the enzyme necessary to break down phenylalanine (an amino acid). Fortunately, when recognized, this metabolic disorder can be successfully treated by dietary restriction.

10

Unfortunately, newborns are not typically screened for other metabolic disorders such as homocysteinuria, maple syrup urine disease, organic acid disorders, and disorders of fatty acid oxidation. As a result, these disorders are often detected in 15 infants and children after damage has occurred and effects such as developmental delay and mental retardation become apparent. However, such effects often can be reduced or avoided with early detection and sustained dietary restriction. Such early detection involves examining a blood sample for an enzyme activity or a metabolic marker (a metabolite).

20

Wide-spread newborn testing for multiple inherited metabolic disorders has been unavailable due, in part, to cost associated with each test. One approach for reducing costs is to examine multiple enzyme activities and metabolites and in a single test. However, development of multi-enzyme/metabolite testing has been hindered by the different conditions needed for enzyme tests in comparison to metabolite tests. 25 Currently, enzyme and metabolite tests cannot be carried out simultaneously from a single patient sample.

30 Thus, there exists a need for methods for efficiently diagnosing a metabolic disorder in an individual. The present invention satisfies this need and provides related advantages as well.

SUMMARY OF THE INVENTION

The present invention provides a method for detecting a metabolic disorder in an individual. The method involves (a) contacting a sample containing (i) one or 5 more metabolically indicative enzymes and (ii) one or more metabolic analytes, with one or more substrates for said one or more enzymes to produce a reaction admixture, under conditions in which at least one of said enzymes is capable of acting on a corresponding substrate to generate at least one product; (b) contacting said reaction admixture with a reagent that inhibits the ability of said one or more enzymes 10 to act on a corresponding substrate, wherein said one or more metabolic analytes and said at least one product are soluble in said reagent; to produce a test sample and (c) determining the presence or amount of said one or more metabolic analytes and said at least one product contained in said test sample using mass spectrometry, wherein a determined presence or amount of said one or more metabolic analytes and said at 15 least one product correlates with presence or absence of said metabolic disorder. A sample used in a method of the invention can be, for example, a body fluid sample, such as blood, and can be obtained from a neonate, newborn, child or adult. In one embodiment, the method can include, in step (a) contacting the sample with one or more reference substrates. In another embodiment, the method can include, in step 20 (b), contacting the sample with one or more reference products. In a further embodiment, the method can include, in step (d), prior to determining, adding one or more reference products corresponding to the at least one product. In yet another embodiment, the method can include, in step (d), prior to determining, adding one or more reference analytes corresponding to the one or more metabolic analytes 25 contained in the sample.

The invention provides another method for detecting a metabolic disorder in an individual. The method involves (a) contacting a vessel containing one or more substrates in dried form with a sample from the individual, the sample containing (i) one or more metabolically indicative enzymes and (ii) one or more metabolic analytes, 30 in aqueous solution; under conditions wherein at least one of the enzymes is capable of acting on a corresponding substrate to generate at least one product; (b) adding a reagent that inhibits the ability of the one or more enzymes to act on a corresponding

substrate, and wherein the one or more metabolic analytes and at least one product are soluble in the reagent; (c) determining the presence or amount of the one or more metabolic analytes and the at least one product in the resultant mixture using mass spectrometry, wherein a determined presence or amount of said one or more metabolic analytes and said at least one product correlates with presence or absence of said metabolic disorder. A sample used in a method of the invention can be, for example, a body fluid sample, such as blood, and can be obtained from a neonate, newborn, child or adult. In an embodiment, the vessel further can contain one or more reference substrates, one or more metabolic analytes, one or more references products, or a combination of such references.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows simultaneous detection of biotinidase activity and metabolic analytes (amino acids and carnitines) using a method of the invention.

DETAILED DESCRIPTION OF THE INVENTION

This invention relates to the discovery of metabolic disorder diagnostic tests that allow detection of at least two relevant markers simultaneously and from a single sample. In particular, the inventor has discovered that one sample can be used to simultaneously detect a metabolic analyte, as well as activity of a metabolic enzyme, using mass spectrometry.

The present invention involves detecting at least two markers or indicia of one or more metabolic disorders from a single sample. At least one index corresponds to the activity of a metabolically indicative enzyme in the sample, and at least one index corresponds to the presence or amount of a metabolic analyte endogenously contained in the sample. Previous clinical methods for genetic screening using mass spectrometry detected only metabolic analytes in a single sample. In particular, Chace (US 6,258,605 and US 6,455,321) describes preparing a patient sample (dried blood spot) by extraction in methanol prior to analysis. Under such conditions,

enzymes are inactive such that measuring enzyme activity is unfeasible, moreover, uncontemplated.

Conversely, well-known methods for determining enzymatic activity do not 5 include detection of metabolic analytes but instead measure a tag on the enzymatic product, such as a fluorescent tag. Modes of detection for tagged products cannot simultaneously detect metabolic analytes present in a patient sample. Another disadvantage of fluorimetric methods with respect to enzymatic assays is that artificial substrates tagged with a fluorescent group are different enough from their natural 10 counterparts that enzyme activity toward such substrates is inherently reduced and thus sensitivity of the assays is reduced. The methods of the invention advantageously can involve detection of metabolically indicative enzyme activity toward natural substrate.

15 As described in Example I, the activity of metabolically indicative enzyme biotinidase, as well as the concentration of alpha amino acids and carnitines, were simultaneously determined from a single sample. As is shown in Figure I, in spite of its multiplex character, the sensitivity of the method for detecting biotinidase (Fig. 1) is comparable to a well-known fluorimetric enzyme assay method. Similarly, as shown 20 in Tables 3 and 4, a method of the invention has sensitivity comparable to well-known mass spectrometry methods that measure analytes without regard for enzyme activities. Based on this discovery of simultaneous detection of metabolic analytes and enzyme activity from a single sample, the invention provides methods for detecting metabolic disorders.

25

Detecting of a metabolic disorder can be used in diagnosing or predicting susceptibility to a metabolic disorder because biochemical indicators examined using the methods of the invention are indicative of a metabolic disorder, whether or not physiologic or behavioral symptoms of the disorder have become apparent.

30

The methods of the invention also are useful for monitoring the metabolism of an individual, such as one undergoing treatment for a metabolic disorder. As a non-limiting example, the methods can be used for determining therapeutic efficacy of a

particular treatment. Based on this determination, the individual can be offered additional or alternative therapeutic options. The methods of the invention also can be useful for assessing patient compliance with a particular treatment modality, such as dietary restriction. Therefore, the invention is applicable to screening, diagnosis, 5 prognosis, monitoring therapy and compliance, and any other application in which determining the presence or amount of one or more metabolic analytes and at least one metabolically indicative enzyme activity is useful.

The methods of the invention involve detecting two or more analytes and 10 metabolically indicative enzyme activities simultaneously. By providing a multiplexed format, the invention advantageously reduces technical performance errors because the number of manipulations is fewer than tests performed separately. Moreover, assay costs associated with simultaneous detection of multiple analytes and enzyme 15 activities is lower than those associated with individual tests because fewer materials (assay vessels, liquid transfer materials, and the like) are used and less operator time is required.

As used herein, the term "metabolic disorder" means any abnormal condition in 20 the body that causes loss of metabolic control of the body's steady state. Metabolism is the physiological and biochemical process by which food is converted in the body to forms that provide energy for bodily activities. Metabolism includes creation of specific molecules that the body uses for its myriad activities. Such molecules include hormones, neurotransmitters, proteins such as enzymes, and membrane constituents. Metabolism also includes degradation processes that enable cells to excrete waste 25 products. Exemplary metabolic processes include processes for absorption and modification of vitamins and minerals; for degrading molecules to provide energy or to be excreted; processes for making acetyl-coenzyme A, nonessential amino acids, cholesterol, long-chain fatty acids, prostaglandins, complex lipids and proteins; and for neutralizing toxins. Therefore, a metabolic disorder is a condition that interferes with 30 normal creation or destruction of biological molecules that regulate health. A metabolic disorder arising early in life and resulting from genetic inheritance is generally referred to as an "inborn error of metabolism", whereas a metabolic disorder

that arises as a result of disease or lifestyle is generally referred to as an "acquired metabolic disorder", although genetic predisposition generally exists.

Elevated amino acids, free carnitine and acylcarnitine levels are examples of
5 metabolites that can be indicative of one or more of several metabolic disorders. Free carnitine and acylcarnitines are markers for disorders that are classified as fatty acid oxidation (FAO) disorders and organic aciduria disorders (OAD). Similarly, amino acids are used as markers for several metabolic disorders collectively known as amino acidopathies.

10

In Fatty Acid Oxidation (FAO) Disorders disorders, enzymes necessary for fatty acid breakdown are unavailable or have reduced activity. Breakdown, or oxidation, of fatty acids is necessary for energy production when glucose levels, the body's main source of energy, are low. Without this energy supply some individuals may have
15 recurring incidences of low blood sugar levels. In cases of fasting, often caused by illnesses such as ear infections or flu, there may be metabolic crisis. Affected individuals may show vomiting, diarrhea, lethargy, seizures and coma. Failure to diagnose FAO disorders may result in excessive fat buildup in the liver, heart and kidneys. This buildup can cause a variety of symptoms, ranging from hepatic failure,
20 encephalopathy, and heart and eye complications to general problems with muscle development. Many of these clinical symptoms can lead to death. Many deaths due to FAO disorders have been misdiagnosed as SIDS or Reye's Syndrome.

In Organic Aciduria (OA) Disorders, the metabolic pathways of organic acids
25 are disrupted in OA disorders and thus accumulation of the acids in blood and urine alters the acid-base balance of the body. Resulting modifications or adaptations to intermediary metabolic pathways may cause numerous clinical symptoms, including metabolic acidosis, ketosis, hyperammonemia, failure to thrive, sepsis or coma.

30

In disorders of amino acid metabolism (amino acidopathies), enzymes necessary for the metabolism of certain amino acids are unavailable or have reduced activity. As a result, the concentration of the affected amino acids and alternate metabolites increases in the infant's body. These excesses can have deleterious

effects on the infant's health including death. Some commonly studied amino acidopathies are:

Phenylketonuria (PKU) is a disorder of aromatic amino acid metabolism in which phenylalanine cannot be converted to tyrosine. If untreated, PKU leads to various degrees of mental retardation. Hyperphenylalaninemia leads to mental retardation and muscular rigidity. Homocystinuria leads to vascular occlusive disease, osteoporosis, accumulation of homocystine and methionine, and variable developmental delays. Maple Syrup Urine Disease (MSUD) is caused by a disorder of branched-chain amino acid metabolism resulting in elevated levels of leucine, isoleucine and valine in the blood. If untreated, lethargy progressive to coma, developmental delay, and convulsions will develop. Tyrosinemia type I (hereditary tyrosinemia), leads to acute hepatic failure or chronic cirrhosis and hepatocellular carcinoma. Citrullinemia leads to convulsions, anorexia, vomiting and lethargy, followed rapidly by potentially lethal coma.

As used herein, the term "metabolically indicative" when used in reference to an enzyme means that the amount or activity of the enzyme is altered as a result of a metabolic disorder in an individual. Therefore, the amount or activity of a metabolically indicative enzyme can be increased or decreased in an individual having a metabolic disorder in comparison to an individual free of the metabolic disorder. Although the activity of a metabolically indicative enzyme can be altered in an individual having a metabolic disorder, it is understood that the individual can be free of, or can have outward signs or symptoms of, the disorder.

25

As used herein, the term "metabolic analyte" means a substance that is altered in amount, activity, or physical characteristic in the body of an individual experiencing a metabolic disorder in comparison to an individual not experiencing a metabolic disorder. Exemplary metabolic analytes include an amino acid, fatty acid and organic acid, and other small biological molecules, as well as a macromolecule such as a nucleic acid, polypeptide or carbohydrate, and the like.

The invention provides methods for detecting a metabolic disorder in an individual. One method involves (a) contacting a sample containing (i) one or more metabolically indicative enzymes and (ii) one or more metabolic analytes, with one or more substrates for the one or more enzymes to produce a reaction admixture, under 5 conditions in which at least one of the enzymes is capable of acting on a corresponding substrate to generate at least one product; (b) contacting the reaction admixture with a reagent that inhibits the ability of the one or more enzymes to act on a corresponding substrate, wherein the one or more metabolic analytes and the at least one product are soluble in the reagent; to produce a test sample and (c) 10 determining the presence or amount of the one or more metabolic analytes and the at least one product contained in the test sample using mass spectrometry.

The invention provides another method for detecting a metabolic disorder. The method involves (a) contacting a vessel containing one or more substrates in dried 15 form with a sample from the individual, the sample containing (i) one or more metabolically indicative enzymes and (ii) one or more metabolic analytes, in aqueous solution; under conditions wherein at least one of the enzymes is capable of acting on a corresponding substrate to generate at least one product; (b) adding a reagent that inhibits the ability of the one or more enzymes to act on a corresponding substrate, 20 wherein the one or more metabolic analytes and at least one product are soluble in the reagent; (c) determining the presence or amount of the one or more metabolic analytes and the at least one product in the resultant mixture using mass spectrometry wherein a determined presence or amount of the one or more metabolic analytes and the at least one product correlates with presence or absence of the metabolic disorder.

25

The invention provides a further method for detecting a metabolic disorder in an individual. The method involves (a) contacting a sample comprising: (i) one or more metabolically indicative enzymes and (ii) one or more metabolic analytes, with one or more substrates for the one or more enzymes to produce a 30 reaction admixture, under conditions in which at least one of the enzymes is capable of acting on a corresponding substrate to generate at least one product; (b) contacting the reaction admixture with a reagent that inhibits the ability of the one or more enzymes to act on a corresponding substrate, wherein the one or more

metabolic analytes and the at least one product are soluble in the reagent, to produce a test sample; (c) contacting the reaction admixture with one or more reference products and one or more reference analytes, and (d) determining the presence or amount of the one or more metabolic analytes and the at least one product contained 5 in the test sample, with respect to the one or more reference products and one or more reference analytes, using mass spectrometry, wherein a determined presence or amount of the one or more metabolic analytes and the at least one product correlates with presence or absence of the metabolic disorder.

10 Another method for detecting a metabolic disorder in an individual involves (a) contacting a vessel containing one or more substrates in dried form with a sample from the individual, the sample containing (i) one or more metabolically indicative enzymes and (ii) one or more metabolic analytes, (b) contacting the vessel with a solution to generate conditions wherein at least one of the enzymes is capable of 15 acting on a corresponding substrate to generate at least one product; (c) adding a reagent that inhibits the ability of the one or more enzymes to act on a corresponding substrate, wherein the one or more metabolic analytes and at least one product are soluble in the reagent; (d) determining the presence or amount of the one or more metabolic analytes and the at least one product in the resultant mixture using 20 mass spectrometry, wherein a determined presence or amount of the one or more metabolic analytes and the at least one product correlates with presence or absence of the metabolic disorder.

Further, the invention provides a method for detecting a metabolic disorder in 25 an individual that involves (a) separating a sample into two sample portions, the sample comprising: (i) one or more metabolically indicative enzymes and (ii) one or more metabolic analytes; (b) contacting each sample portion with one or more substrates for the one or more enzymes to produce two reaction admixtures, under conditions in which at least one of the enzymes in each sample portion is 30 capable of acting on a corresponding substrate to generate at least one product; (c) contacting each reaction admixture with a reagent that inhibits the ability of the one or more enzymes to act on a corresponding substrate, wherein the reagent can be the same or different for each reaction admixture, and wherein the one or more

metabolic analytes and the at least one product are soluble in the reagent or reagents; to produce a test sample and (d) combining the two reaction admixtures to produce a test sample, and (e) determining the presence or amount of the one or more metabolic analytes and the at least one product contained in the test sample using mass spectrometry, wherein a determined presence or amount of the one or more metabolic analytes and the at least one product correlates with presence or absence of the metabolic disorder.

Yet another method for detecting a metabolic disorder in an individual provided by the invention involves (a) separating a sample into two sample portions, the sample comprising: (i) one or more metabolically indicative enzymes and (ii) one or more metabolic analytes; (b) contacting each sample portion with individual vessels, each vessel containing one or more substrates for the one or more enzymes in dried form, (c) contacting each vessel with a solution to produce two reaction admixtures, under conditions in which at least one of the enzymes in each sample portion is capable of acting on a corresponding substrate to generate at least one product; (d) contacting each reaction admixture with a reagent that inhibits the ability of the one or more enzymes to act on a corresponding substrate, wherein the reagent can be the same or different for each reaction admixture, and wherein the one or more metabolic analytes and the at least one product are soluble in the reagent or reagents; to produce a test sample and (e) combining the two reaction admixtures to produce a test sample, and (f) determining the presence or amount of the one or more metabolic analytes and the at least one product contained in the test sample using mass spectrometry, wherein a determined presence or amount of the one or more metabolic analytes and the at least one product correlates with presence or absence of the metabolic disorder.

In any of the above methods for detecting a metabolic disorder, one or more reference products and/or reference analytes can be added to a test sample prior to mass spectrometry. Similarly, one or more reference substrates can be added to a reaction admixture or solution before an enzymatic reaction is allowed to proceed. Such additions can be useful when the methods are carried out in a quantitative format, as is described in more detail herein below.

The methods of the invention can be carried out in a variety of formats. As an example of a typical procedure, a sample from a normal individual or one suspected of having a metabolic disorder is placed in a vessel. The type of the sample (for 5 example, whole blood, serum, tissue) is chosen based on the particular metabolic disorder, and any sample pre-treatment to release enzymes is performed based on the sample type. An enzyme that correlates with the test subject metabolic disorder is known, as is the enzyme substrate. At least one metabolic analyte known to correlate with the metabolic disorder, or a related or different metabolic disorder, is also known.

10

To the sample is added an aqueous solution containing a substrate for the enzyme indicative of the metabolic disease. The enzymatic reaction is allowed to proceed for a time period sufficient for the enzyme to produce a detectable or desired amount of product. The reaction is then stopped by adding a stop reagent that 15 includes a reference analyte to be detected. The reagent denatures the enzyme, but maintains the product and reference form of metabolic analyte in solution (reference analyte). The sample is subjected to tandem mass spectrometry to detect the product, metabolic analyte, and reference analyte. The amount of metabolic analyte is determined based on the reference analyte, and the presence of the product is 20 detected.

As another example, a sample from a normal individual or one suspected of having a metabolic disorder is added to a vessel containing the metabolically indicative enzyme substrate. If the sample is contained in a suitable solution, the 25 enzymatic reaction is allowed to proceed. If the sample is in dried form, such as on membrane or paper, a suitable solution is applied to the sample when in the vessel. After a time period sufficient to solubilize the enzyme or allow enzyme to be released from the material on which it is dried (if present), the enzymatic reaction is allowed to proceed. As above, the enzymatic reaction is allowed to proceed for a time period 30 sufficient for the enzyme to produce a detectable or desired amount of product. The reaction is then stopped by adding a stop reagent that includes a reference analyte and a reference product. The amounts of metabolic analyte and enzymatic product are determined based on the reference analyte and product.

As a further example of a procedure for performing a method of the invention, after addition of the stop reagent, the test sample can be processed to separate soluble from insoluble material. Separation can be performed, for example, by 5 centrifugation, filtration and the like. The resultant clarified test sample can then be subjected to mass spectrometry. Another optional step is to add a detergent or other solubilizing agent to the test sample, for example in the case that the reaction admixture and reagent are immiscible.

10 Although not required, the test sample can be derivatized prior to mass spectrometry to enhance detection of the amount of metabolic analyte, enzymatic product and/or reference substance. An exemplary well-known derivatization method is reaction of sample with acid and butanol.

15 A method of the invention can be used to simultaneously determine presence or amount of one or more metabolic analytes and at least one enzymatic product. As such, the methods are applicable to simultaneously determining presence or amount of two or more metabolic analytes and at least one enzymatic product. three or more metabolic analytes and at least one enzymatic product, four or more metabolic 20 analytes and at least one enzymatic product, including at least five metabolic analytes, at least five metabolic analytes, at least ten metabolic analytes and at least twenty metabolic analytes, together with a at least one enzymatic product. Similarly, the methods are applicable to simultaneously determining presence of amount of one or more metabolic analytes and two or more enzymatic products, one or more metabolic 25 analytes and three or more enzymatic products, one or more metabolic analytes and four or more enzymatic products, one or more metabolic analytes and five more enzymatic products, as well as one or more metabolic analytes and more than five enzymatic products.

30 The methods of the invention involve detecting a metabolic disorder by testing a "sample." As used herein, the term "sample" means any biological fluid, cell, tissue, or fraction thereof, that includes one or more enzymes having an endogenous enzyme indicative of a metabolic state, and one or more endogenous metabolic analytes. A

sample can be, for example, a specimen obtained from an individual or can be derived from such a specimen. For example, a sample can be a tissue section obtained by biopsy, or cells that are placed in or adapted to tissue culture. Exemplary samples therefore include cultured fibroblasts, cultured amniotic fluid cells, and chorionic villus sample. A sample can also be a biological fluid specimen such as urine, blood, plasma, serum, saliva, semen, sputum, cerebral spinal fluid, tears, mucus, and the like. A sample can be further fractionated, if desired, to a fraction containing particular cell types. For example, a blood sample can be fractionated into serum or into fractions containing particular types of blood cells such as red blood cells or white blood cells (leukocytes). If desired, a sample can be a combination of samples from an individual such as a combination of a tissue and fluid sample, and the like.

For used in the methods of the invention, a sample contains one or more metabolically indicative enzymes and one or more endogenous metabolic analytes. To allow enzyme activity to occur in solution while practicing the methods of the invention, the sample can be suitably processed to release metabolically indicative enzyme(s) and/or endogenous metabolic analytes into solution. Well known methods for releasing or solubilizing enzymes in samples containing cells include treatment with detergent, freeze/thaw treatment, drying and rehydrating, and the like.

20

The origin of a sample used in a method of the invention will depend upon the metabolically indicative enzyme(s) and metabolic analyte(s) to be detected. The body fluids and tissues in which metabolically indicative enzymes and metabolic analytes are contained are well known to those skilled in the art. As non-limiting examples, a whole blood sample can be used when biotinidase is the metabolically indicative enzyme; a liver biopsy can be used when carbamylphosphate synthetase is the metabolically indicative enzyme; urine can be used when branched-chain ketoacid dehydrogenase is the metabolically indicative enzyme and a skin biopsy can be used when alpha-L-fucosidase is the metabolically indicative enzyme.

25

Methods for obtaining samples that preserve the activity or integrity of molecules in the sample are well known to those skilled in the art. Such methods include the use of appropriate buffers and/or inhibitors, including nuclease, protease

and phosphatase inhibitors, that preserve or minimize changes in the molecules in the sample. Such inhibitors include, for example, chelators such as ethylenediamine tetraacetic acid (EDTA), ethylene glycol bis(P-aminoethyl ether)N,N,N1,N1-tetraacetic acid (EGTA), protease inhibitors such as phenylmethylsulfonyl fluoride (PMSF), aprotinin, leupeptin, antipain and the like, and phosphatase inhibitors such as phosphate, sodium fluoride, vanadate and the like. Appropriate buffers and conditions for isolating molecules are well known to those skilled in the art and can be varied depending, for example, on the type of molecule in the sample to be characterized (see, for example, Ausubel et al. *Current Protocols in Molecular Biology* (Supplement 47), John Wiley & Sons, New York (1999); Harlow and Lane, *Antibodies: A Laboratory Manual* (Cold Spring Harbor Laboratory Press (1988); Harlow and Lane, *Using Antibodies: A Laboratory Manual*, Cold Spring Harbor Press (1999); Tietz *Textbook of Clinical Chemistry*, 3rd ed. Burtis and Ashwood, eds. W.B. Saunders, Philadelphia, (1999)). A sample also can be processed to eliminate or minimize the presence of interfering substances.

Various methods to fractionate a fluid sample or cell extract are well known to those skilled in the art, including subcellular fractionation or chromatographic techniques such as ion exchange, hydrophobic and reverse phase, size exclusion, affinity, hydrophobic charge-induction chromatography, and the like (Ausubel et al. *supra*, 1999; Scopes, *Protein Purification: Principles and Practice*, third edition, Springer-Verlag, New York (1993); Burton and Harding, *J. Chromatoqr. A* 814:71-81 (1998)). As an example, a serum sample from an individual containing leukocytes can be fractionated to isolate leukocytes, if desired, or subfractionated, for example, into macrophages, T cells, B cells, granulocytes, monocytes, neutrophils, eosinophils, basophils, mast cells, and the like.

For use in a method of the invention, a sample can be in a variety of physical states. For example, a sample can be a liquid or solid, can be dissolved or suspended in a liquid, can be in an emulsion or gel, and can be absorbed onto a material. As a non-limited example, a sample can be a liquid blood sample, liquid serum sample, liquid white blood cell sample, dried blood, serum, or white cell sample, or such a

sample absorbed onto a paper or polymer substrate.

Example 1 describes use of a dried blood sample in a method of the invention.

A sample used in a method of the invention can be separated prior to use.

5 Separation can be desirable if particular metabolically indicative enzymatic activities present in the sample require distinct assay conditions, or if particular metabolic analytes endogenously present in the sample are soluble in different solution environments. Separation also can be desirable to reduce the presence of substances that interfere with detection of a metabolically indicative enzyme or

10 metabolic analyte in a sample, and to retain that interfering portion of the sample in order to detect one or more distinct metabolically indicative enzymes or metabolic analytes in the portion.

Individuals of all ages can be affected by a metabolic disorder. Therefore, a

15 sample used in a method of the invention can be obtained from an individual of any age, including a neonate, newborn, baby, child, and adult. Exemplary metabolic disorders that can be characterized by adult onset include Schindler disease (lysosomal enzyme alpha-N-acetylgalactosaminidase deficiency), urea cycle disorders and sphingolipidoses. The methods of the invention are applicable to human and non-

20 human individuals. As such, the methods can be used on a sample obtained, for example, from any veterinary or research subject for which a metabolically indicative enzyme and endogenous metabolic analyte are known. Exemplary non-human animals include a horse, dog, cat, rabbit, rat, mouse, fish, turtle and lizard. The methods of the invention also are applicable to lower organisms, such as yeast,

25 archebacteria and bacteria, and plants, so long as a sample containing an endogenous metabolically indicative enzyme and endogenous metabolic analyte can be obtained.

The methods of the invention involve detecting the enzymatic product of at least

30 one "metabolically indicative enzyme." Any metabolically indicative enzyme can be assayed using the methods of the invention, so long as a substrate is known for the enzyme, which substrate is soluble in a selected "reagent," and which substrate can be modified to a form distinctly detectable by mass spectrometry, if desired, for use as

an internal standard. Metabolically indicative enzymes are represented in every enzyme classe, including oxidoreductases, hydrolases, lyases, transferases, ligases and isomerases. As a non-limiting example, well-known metabolically indicative enzymes and their associated metabolic disorders are shown below in Table 1. A 5 specific example of a metabolically indicative enzyme is biotinidase.

Table 1:

Metabolic disorder	Metabolically indicative enzyme	Exemplary metabolic analyte
3-methylglutaconic aciduria	3-methylglutaconic hydratase	3-methylglutaconic acid
Smith-Lemli-Opitz Syndrome		7-dehydrocholesterol
4-hydroxylbutyric aciduria	Succinic semialdehyde dehydrogenase	
Adenylosuccinate lyase deficiency	Adenylosuccinate lyase	succinylAICA riboside and succinyladenosine
Beta-Ketothiolase Deficiency	3-oxoacid CoA transferase and beta-ketothiolase.	
Biotinidase deficiency	Biotinidase	Biotin
Carbamylphosphate synthetase deficiency	Carbamylphosphate synthetase	
Duarte Galactosemia	Gal-1-P-uridyltransferase	Gal-1-P
Ehlers-Danlos Syndrome, Type VI	Lysyl hydroxylase	pyridinoline (Pyr), and deoxypyridinoline, (Dpyr), and their ratio
Fucosidosis	alpha-L-fucosidase	
Guanidinoacetate Methyltransferase Deficiency	Guanidinoacetate methyltransferase	Guanidinoacetate
Krabbe Disease	Beta-Galactocerebrosidase	
Long Chain Acyl-CoA Dehydrogenase Deficiency	Acyl-CoA Dehydrogenases: Short-, Medium-, and Long Chain	
Maple Syrup Urine Disease	Branched-Chain Ketoacid Dehydrogenase	branched chain amino acids (valine, isoleucine and leucine)

Methyltetrahydrofolate Reductase Deficiency	Methyltetrahydrofolate Reductase	
Phenylketonurea	phenylalanine hydroxylase	Biopterin, Pteridine, Phenylalanine
Tay-Sachs Disease	Beta-N-Acetylglucosaminidase, Hexosaminidase (Beta-Hexosaminidase-A)	Gangliosides
Urea Cycle Disorder	Argininase, Argininosuccinate Lyase, Carbamyl Phosphate Synthetase	Orotic acid
Zellweger Syndrome	Dihydroxyacetonephosphate acyltransferase, Phytanic Acid Oxidase	Pipecolic Acid, Very Long Chain Fatty Acids
Classic galactosemia	Galactose-1-phosphate uridyl transferase (GALT)	
Galactosemia	Galactokinase	
Galactosemia	Uridine diphosphate galactose-4-epimerase	
G6PD deficiency	Glucose-6-phosphate dehydrogenase (G6PD)	
Sialidosis type II	Neuraminidase	
Hurler disease	α -Iduronidase	
Hunter disease	Iduronate sulfatase	
Sanfilipo disease type A (MPS IIIA)	Heparan sulphamidase	
Sanfilipo disease type B (MPS IIIB)	α -N-acetyl glucosaminidase	
Sanfilipo disease type C (MPS IIIC)	Glucosamine N-acetyl transferase	
Morquio disease type A (MPS IVA)	N-acetyl galactosamine-6-sulphatase	
Morquio disease type B (MPS IVB)	β -Galactosidase	
Maroteaux-Lamy disease (MPS VI)	Arylsulphatase B	
Sly disease (MPS VII)	β -Glucuronidase	
α -Mannosidosis	α -Mannosidase	

β -Mannosidosis	β -Mannosidase	
α -Fucosidosis	α -Fucosidase	
Sialidosis	α -N-acetylneuraminidase	
Galactosialidosis	α -N-acetylneuraminidase, β -Galactosidase	
Aspartylglucosaminuria	Aspartylglucosaminidase	
Schidler disease	α -N-acetylgalactosaminidase,	
Pompe disease	α -Glucosidase	
Tay-Sachs	α -Hexosaminidase	
Sandhoff disease	α -Hexosaminidase	
GM-1 gangliosidosis	β -Galactosidase	
Gaucher disease	β -Glucosidase, Chitotriosidase	
Krabbe disease	Galactocerebrosidase	
Metachromatic leucodystrophy	Arylsulphatase A	
Fabry disease	α -Galactosidase A	
Farber disease	Ceramidase	
Niemann-Pick disease	Sphingomyelinase	

Substrates for metabolically indicative enzymes are well known to those skilled in the art. As used herein, the term "substrate" means a molecule or complex acted upon by a metabolically indicative enzyme. In particular, a substrate has a physiological affinity for and must be able to be acted upon by a corresponding enzyme. A substrate useful in the methods of the invention can be native or modified. Modified substrates useful in the invention have a mass different from the native substrate and retain the ability to be acted upon by the corresponding enzyme. A variety of chemical modification reactions useful for increasing the mass of a substrate are well known to those skilled in the art. Exemplary modifications suitable for substrates include isotopic labeling and any chemical modification that alters the mass or mass-to-charge ratio of the substrate. Once a substrate is modified, activity can be confirmed using routine methods, such as comparison of enzyme activity using native substrate vs. modified substrate, or by quantitation of the product via internal standardization using one or more reference products of known concentration. As a non-limiting example, when the metabolically indicative enzyme is biotinidase, a substrate can be native biocytin or modified biocytin.

As used herein, the term "product" when used in reference to enzymatic activity means a molecular entity that is produced as a result of enzyme acting on substrate. Therefore, the presence or amount of a product in a sample correlates with the 5 presence or amount of enzymatic activity in the sample. As such, an amount of product determined using a method of the invention can be expressed as a level of enzymatic activity using routine calculations, if desired, for example, when assessing differences between test sample and reference level enzymatic activity.

10 The methods of the invention involve use of a reagent that inhibits the ability of one or more metabolically indicative enzymes to act on a corresponding substrate, As used herein, the term "inhibit" means to reduce or block enzyme activity such that product formation is undetectable or insignificant.

15 As used herein, a "reagent" for use in the methods of the invention is or contains a chemical agent that specifically or non-specifically inhibits the ability of a metabolically indicative enzyme to act on a corresponding substrate, and does not substantially interfere with mass spectrometry analysis. Exemplary non-specific reagents that inhibit enzymatic reactions are chaotropes, which are chemical agents 20 that denature proteins, and solutions containing chaotropes. Non-limiting examples of chaotropes are alcohols, such as methanol, ethanol and isobutyl alcohol; urea and guanidine salts. Exemplary specific reagents that inhibit enzymatic reactions include enzyme inhibitors effective for a selected enzyme. A large number of enzyme inhibitors are widely available and well-known to those skilled in the art. A reagent for 25 use in the methods of the invention can be selected based on, for example, suitability for inhibiting activity of metabolically indicative enzyme(s), and ability to maintain solubility of metabolic analyte(s) and enzymatic product(s). A selected reagent can be selected to have a minimal number of components in order to minimize background for mass spectrometry. A selected reagent can be miscible or immiscible with a particular 30 reaction admixture or solution, with rapid mixing being one method for ensuring that enzymatic reaction is inhibited. Examples of reagents useful in the invention include organic solvents and solutions thereof, organic acids and solutions thereof, alcohols and solutions thereof, reagents that are non-polar relative to the reaction admixture or

solution, and reagents that have a pH distinct from the reaction admixture or solution, the pH creating an unsuitable environment for enzymatic activity, and the like. Non-limiting specific reagents include methanol and solutions thereof; trifluoroacetic acid and solutions thereof; and acetonitrile and solutions thereof. Those skilled in the art of 5 mass spectrometry will be able to identify reagents that do not interfere with mass spectrometry determinations.

The ability of a selected reagent to inhibit an enzymatic reaction can be determined by routine methods. For example, the effectiveness of a candidate 10 "reagent" can be tested by adding the reagent or a known inhibitor of the enzyme to an enzymatic reaction in progress. The amount of enzymatic activity under each condition can then be compared to confirm if a selected reagent is suitable for use in the invention methods.

15 The methods of the invention involve detecting at least one "metabolic analyte." A variety of metabolic analytes can be detected using the methods of the invention, so long as the analyte is soluble in a selected reagent, has a known mass, and can be modified, if desired, for use as an internal standard. Metabolic analytes associated with particular metabolic disorders are well known to those skilled in the art. As non- 20 limiting examples, well-known metabolic analytes and their associated metabolic disorders are shown above in Table 1. Amino acid and carnitine metabolic analytes are shown in Table 2, and can correspond with common metabolic disorders shown in Table 3. Amino acid disorders and corresponding amino acid metabolic analytes are also shown in Table 4. Fatty acid disorders and corresponding fatty acid metabolic 25 analytes are shown in Table 5; additional fatty acid disorders shown in Table 3.

Table 2

ANALYTE NAME	ABBREVIATION
Amino Acids	
Alanine	Ala
Arginine	Arg
Citruline	Cit
Glycine	Gly
Leucine	Leu

Methionine	Met
Ornithine	Orn
5-Oxoproline	5-Oxo Pro
Phenylalanine	Phe
Tyrosine	Tyr
Valine	Val
Proline	Pro
Carnitines	
Free carnitine	C0
Acetylcarnitine	C2
Propionylcarnitine	C3
Malonylcarnitine	C3DC
Butyrylcarnitine	C4
3-Hydroxy-butyrylcarnitine	C4OH
Isovalerylcarnitine	C5
Tiglylcarnitine	C5:1
Glutarylcarnitine	C5DC
3-Hydroxy-isovalerylcarnitine	C5OH
Hexanoylcarnitine	C6
Adipylcarnitine	C6DC
Octanoylcarnitine	C8
Octenoylcarnitine	C8:1
Decanoylcarnitine	C10
Decenoylcarnitine	C10:1
ANALYTE NAME	
ABBREVIATION	
Carnitines	
Decadienoylcarnitine	C10:2
Dodecanoylcarnitine	C12
Dodecenoylcarnitine	C12:1
Tetradecanoylcarnitine (Myristoylcarnitine)	C14
Tetradecenoylcarnitine	C14:1
Tetradecadienoylcarnitine	C14:2
3-Hydroxy-tetradecanoylcarnitine	C14OH
Hexadecanoylcarnitine (palmitoylcarnitine)	C16
Hexadecenoylcarnitine	C16:1
3-Hydroxy-hexadecanoylcarnitine	C16OH
3-Hydroxy-hexadecenoylcarnitine	C16:1OH
Octadecanoylcarnitine (Stearoylcarnitine)	C18
Octadecenoylcarnitine (Oleylcarnitine)	C18:1
Octadecadienoylcarnitine (Linoleylcarnitine)	C18:2
3-Hydroxy-octadecanoylcarnitine	C18OH
3-Hydroxy-octadecenoylcarnitine	C18:1OH

Table 3

Amino Acid Disorders	
ARGD	Argininemia
ASA, ASL, ALD	Argininosuccinic aciduria (Argininosuccinate lyase deficiency, Argininosuccinase deficiency)
ASD, ASS	Citrullinemia (Argininosuccinic acid synthetase deficiency, Argininosuccinate synthetase deficiency)
HCU, HCYS	Homocystinuria (Cystathione synthase deficiency)
Hmet	Hypermethioninemia
HHH	Hyperornithinemia, Hyperammonemia, Hyperhomocitrullinuria syndrome (Ornithine translocase deficiency)
PRO	Hyperprolinemia
MSUD, BCKA	Maple Syrup Urine Disease (Branched chain ketoaciduria)
NKG, NKHI	Nonketotic hyperglycinemia
PKU	Phenylketonuria
PYP/PIP	Pyroglutamic/pipecolic acidemia Tyrosenemia, Type I Tyrosenemia, Type II 5-oxoprolinuria (Pyroglutamic aciduria)
Fatty Acid and Organic Acid Disorders	
2-MBCD	2-methylbutyryl CoA dehydrogenase deficiency 2,4-Dienoyl-CoA reductase deficiency
HMG	3 -hydroxy-3-methylglutaryl CoA lyase deficiency (hydroxymethylglutaric acidemia)
3MCC, 3-MMC	3-methylcrotonyl CoA carboxylase deficiency (3-methylcrotonylglycinemia)
CPT I	Carnitine palmitoyltransferase, type I deficiency
CPT II	Carnitine palmitoyltransferase, type II deficiency
CTD	Carnitine transporter defect
CATR, CACT	Carnitine/acylcarnitine translocase defect
EMA	Ethylmalonic acidemia
GA I	Glutaric acidemia, type I (Glutaryl CoA dehydrogenase deficiency)
IBCD	Isobutyryl CoA dehydrogenase deficiency
IVA	Isovaleric acidemia
LCAD	Long-chain acyl-CoA dehydrogenase deficiency
LCHAD	Long-chain hydroxyacyl-CoA dehydrogenase deficiency
MA	Malonic aciduria
MCAD	Medium-chain acyl-CoA dehydrogenase deficiency
MMA	Methylmalonic acidemia
BKT	Mitochondrial acetoacetyl CoA thiolase deficiency (Beta-Ketothiolase deficiency)
MADD, GA II	Multiple acyl-CoA dehydrogenase deficiency (Glutaric acidemia, type II)
MCD	Multiple Co-A carboxylase deficiency (Holocarboxylase synthetase deficiency)
PA, PPA	Propionic acidemia
SCAD	Short-chain acyl-CoA dehydrogenase deficiency

SCHAD	Short-chain hydroxyacyl-CoA dehydrogenase deficiency
TFP	Trifunctional protein deficiency
VLCAD	Very-long-chain acyl-CoA dehydrogenase deficiency

Table 4

Aminoacidopathies	
Phenylketonuria	
Phe ^{2,4,6}	
Phe, Phe/Tyr ⁷	
Phe, Tyr ^{1,5}	
Maple syrup urine disease	
Leu+Ile ¹	
Leu+Ile, Val ^{2,4,6}	
Leu, Val ⁵	
Leu, Leu/Phe ⁷	
Homocystinuria	
Met ^{1,2,4,6}	
Met, Met/Phe, Met/Leu ⁵	
Met, Met/Phe ⁷	
Hypermethioninemia	
Met ²	
Met, Met/Phe ⁷	
Citrullinemia	
Cit ^{1,2,6}	
Cit/Phe, Cit/Tyr ⁴	
Cit, Orn/Cit, Cit/Arg ⁵	
Cit, Cit/Arg ⁷	
Nonketotic hyperglycinemia	
Gly ^{2,4,5}	
Tyrosinemia Type I	
Met, Tyr ^{1,4}	
Tyr ^{2,6}	
Tyr, Tyr/Phe ⁷	
Tyrosinemia II	
Tyr ²	
Tyr ⁷	
Argininemia	
Arg ²	
Arg, Arg/Orn ²	
PRO Hyperprolinemia	
Pro/Phe ⁴	
HHH	
Orn, HomoCit ²	
Orn, Orn/Cit ⁷	
5-oxoprolinuria	
5-Oxopro ²	
pyroGlu/Phe ⁴	
Argininosuccinic aciduria	
Cit ^{1,6}	
Cit/Tyr ⁴	
Asa ⁵	
Cit, Cit/Arg ⁷	

Table 5

Fatty Acid Oxidation Disorders	
SCAD C4, C6 ¹ C4 ^{2,5,7} C4, C5, C4/C2, C5/C2 ⁴ C4, C4/C2, C4/C3 ⁶	LCAD ⁴ C14:2, C14:1, C14:1/C16
MCAD C8, C10:1 ¹ C8, C10, C10:1, C6 ² C6, C8, C10, C10:1, C8/C10 ³ C6, C8, C10:1 ^{4,7} C6, C8, C10:1, C10, C8/C2, C8/C10, C8/C12 ⁵ C6, C8, C10:1, C8/C10 ⁶	VLCAD ¹ C14, C14:1, C16, C18 ² C14:1, C14, C16 ³ C14:1, C14, C16:1, C16, C18:1, C18, C14:1/C12:1 ⁴ C14:2, C14:1, C14:1/C16 ⁵ C14:1 or C14 ⁶ C14, C14:1, C14:2, C16:1, C14:1/C16 ⁷ C14:1
Glutaric acidemia Type II C5DC ¹ C4, C5, C8:1, C8, C12, C14, C16, C5DC ² C4, C5DC, C5, C6, C8, C10 ³ C6, C8, C10, C5/C2 ⁴ C4, C5, C6, C8, C10 ⁶ C8, C10, C5DC ⁷	LCHAD ¹ C14, C14:1, C16, C18, C14OH, C14:1OH, C16OH, C18OH ² C16OH, C18:1OH, C18OH ³ C16OH, C18:1OH, C18OH, C16OH/C16 ⁴ C14OH, C16:1OH, C16OH, C18:1OH, C18:OH, C14:1, C14 ⁶ C16OH, C18:1, C18:1OH, C18:2, C18:2OH ⁷ C16OH
Carnitine transporter defect C0, C2 ⁵	Trifunctional protein deficiency ² C16OH, C18:1OH, C18OH ⁶ C16OH, C18:1OH, C18OH, C16OH/C16 ⁵ C14OH, C16:1OH, C16OH, C18:1OH, C18:OH, C14:1, C14
CPT I C16, C18:1, C18 ² C0, C16, C18, C0/(C16+C18) ⁵	Carnitine/acylcarnitine translocase deficiency ² C16, C18:1, C18 ⁵ C0, C16, C18, C0/(C16+C18) ⁷ C16
Ethylmalonic acidemia C4, C5, C4/C2, C5/C2 ⁴	CPT II ¹ C14, C14:1, C16, C16:1 ³ C16, C18, C18:1, C16/C14:1 ⁶ C16, C18:1, C18:2 ⁷ C16
2,4 Dienoyl CoA reductase deficiency ⁶ C10:2	

In the methods of the invention, the presence or amount of one or more metabolic analytes can be determined. An amount of a metabolic analyte refers to an absolute amount of a molecule in a sample or to a relative amount of the molecule, 5 including amounts determined under steady-state or non-steady-state conditions. The amount of a metabolic analyte can be determined relative to a reference molecule in a sample.

The methods of the invention involve determining the presence or amount of at 10 least one metabolic analyte and/or product, where the presence or amount of each metabolic analyte and/or product correlates the presence or absence of a metabolic disorder. To quantitate the amount of a metabolic analyte or product in a test sample, an internal standard can be used, as is described below. The methods of the invention can be used quantitatively, if desired, to allow comparison of test sample 15 results with known or pre-determined standard amounts of particular analyte and/or product amounts. The methods of the invention can be used qualitatively when the presence of the metabolic analyte or enzyme in the sample is indicative of a metabolic disorder, for example, when the metabolic analyte detected is the result of abnormal metabolic processes, or when enzyme activity is not detected in a normal sample. 20 The methods can also be used qualitatively when a test sample is compared with a reference sample, which can be either a normal reference or metabolic disorder reference. In this format, the relative amount of analyte or product can be indicative of a metabolic disorder.

25 An internal standard, or reference analyte, for a metabolic analyte useful in a method of the invention can be any modification or analog of the analyte that is detectable by mass spectrometry. A reference analyte is separately detectable from the metabolic analyte based on a unique physical characteristic, such as a unique mass or charge-to-mass ratio. In addition, a suitable generic reference standard can 30 be used. Such an internal standard will, for example, co-elute with the analyte or product if a separation method such as chromatography is used prior to mass spectrometric analysis. A commonly used internal standard for mass spectrometry is a stable isotopically labeled form or chemical derivative of the analyte or product.

Non-limiting examples of internal standards for metabolic analytes that are amino acids include isotopically labeled amino acids such as $^{15}\text{N},2\text{-}^{13}\text{C}$ -Glycine, $^{2}\text{H4}$ -Alanine, ^{2}H 8-Valine, ^{2}H 3-Leucine, ^{2}H 3-Methionine, ^{2}H 5-Phenylalanine, ^{2}H H4-Tyrosine, ^{2}H 3-Aspartate, ^{2}H 3-Glutamate, $^{2}\text{H2}$ -Ornithine $^{2}\text{H Cl}$, $^{2}\text{H2}$ -Citrulline, and ^{2}H 4-13C-Arginine·HCl.

Non-limiting examples of internal standards for metabolic analytes that are carnitines include isotopically labeled carnitines such as $^{2}\text{H9}$ -carnitine, ^{2}H 3-acetylcarnitine, ^{2}H 3-propionylcarnitine, ^{2}H 3-butyrylcarnitine, ^{2}H 9-isovalerylcarnitine, ^{2}H 3-octanoylcarnitine, ^{2}H 9-myristoylcarnitine, and ^{2}H 3-palmitoylcarnitine. An internal standard for an enzymatic product, or reference product, can be any modification or analog of the product that is identifiable by mass spectrometry. A reference product is separately detectable from the product based on a unique physical characteristic, such as a unique mass or charge-to-mass ratio. A non-limiting example of an internal standard for an enzymatic product is an isotopically labeled form of the product, such as a deuterated form of the product.

15

Generally, a cut-off value for each metabolic analyte or enzyme activity will be known in the art for commonly tested analytes and enzymes. A cut-off value is typically an analyte concentration or enzyme activity, or ratio thereof, above or below which is considered indicative of a metabolic disorder or cause for retest. Thus, in accordance with the present invention a reference level of a metabolic analyte or enzymatic activity (as indicated by presence or amount of enzymatic product) in a particular sample type is identified as a cut-off value, above which there is a significant correlation between the presence of the analyte or enzymatic activity and presence or absence of a metabolic disorder. Exemplary cut-off values for amino acid and carnitine/acylcarnitine butyl esters are described in US 6,455,321. Also described are procedures for determining significance of an elevated amount of a metabolic analyte in a blood sample from a newborn. This method can be applied to a variety of types of samples (for example, serum and tissue biopsy) from a subject of any age.

30

Those of skill in the art will recognize that some cut-off values are not absolute in that clinical correlations are still significant over a range of values on either side of the cutoff; however, it is possible to select an optimal cut-off value (e.g. varying H-scores, and the like) of analyte, enzyme activity for a particular sample types. Cut-off

values determined for use in a method of the invention generally are compared with published ranges but can be individualized to the methodology used and patient population. It is understood that improvements in optimal cut-off values could be determined depending on the sophistication of statistical methods used and on the 5 number and source of samples used to determine reference level values for the different analyte, enzyme activity and sample types. Therefore, established cut-off values can be adjusted up or down, on the basis of periodic re-evaluations or changes in methodology or population distribution. In addition, instrument-specific cut-off values can be used, if desired, for example such as when inter-instrument performance 10 comparability is <10%.

The reference level can be determined by a plurality of methods, provided that the resulting reference level accurately provides an amount of metabolic analyte or enzyme activity above which exists a first group of individuals having a different 15 probability of metabolic disorder than that of a second group of individuals having metabolic analyte or enzyme activity amount below the reference level. The reference level can be determined by comparison of metabolic analyte or enzyme activity amount in populations of patients having the same metabolic disorder. This can be accomplished, for example, by histogram analysis, in which an entire cohort of 20 patients are graphically presented, wherein a first axis represents the amount of metabolic analyte or enzyme activity and a second axis represents the number of individuals in the cohort whose sample contain metabolic analyte or enzyme activity at a given amount. Two or more separate groups of individuals can be determined by identification of subsets populations of the cohort which have the same or similar 25 levels of metabolic analyte or enzyme activity. Determination of the reference level can then be made based on an amount which best distinguishes these separate groups. The reference level can be a single number, equally applicable to every individual, or the reference level can vary, according to specific subpopulations of individuals. For example, older individuals might have a different reference level than younger 30 individuals for the same metabolic disorder.

The methods of the invention use mass spectrometry for determining the amount or presence of one more metabolic analytes and at least one metabolically

indicative enzyme product. A variety of configurations of mass spectrometers can be used in a method of the invention. Several types of mass spectrometers are available or can be produced with various configurations. In general, a mass spectrometer has the following major components: a sample inlet, an ion source, a mass analyzer, a 5 detector, a vacuum system, and instrument-control system, and a data system. Difference in the sample inlet, ion source, and mass analyzer generally define the type of instrument and its capabilities. For example, an inlet can be a capillary-column liquid chromatography source or can be a direct probe or stage such as used in matrix-assisted laser desorption. Common ion sources are, for example, electrospray, 10 including nanospray and microspray or matrix-assisted laser desorption. Common mass analyzers include a quadrupole mass filter, ion trap mass analyzer and time-of-flight mass analyzer.

The ion formation process is a starting point for mass spectrum analysis.

15 Several ionization methods are available and the choice of ionization method depends on the sample to be analyzed. For example, for the analysis of amino acids a relatively gentle ionization procedure such as electrospray ionization (ESI) can be desirable. For ESI, a solution containing the sample is passed through a fine needle at high potential which creates a strong electrical field resulting in a fine spray of highly charged 20 droplets that is directed into the mass spectrometer. Other ionization procedures include, for example, fast-atom bombardment (FAB) which uses a high-energy beam of neutral atoms to strike a solid sample causing desorption and ionization. Matrix-assisted laser desorption ionization (MALDI) is a method in which a laser pulse is used to strike a sample that has been crystallized in an UV-absorbing compound matrix.

25 Other ionization procedures known in the art include, for example, plasma and glow discharge, plasma desorption ionization, resonance ionization, and secondary ionization.

Electrospray ionization (ESI) has several properties that are useful for the 30 invention described herein. For example, ESI can be used for biological molecules such as amino acids and carnitines that are difficult to ionize or vaporize. In addition, the efficiency of ESI can be very high which provides the basis for highly sensitive measurements. Furthermore, ESI produces charged molecules from solution, which is

convenient for analyzing metabolic analytes, products and reference molecules that are in solution. In contrast, ionization procedures such as MALDI require crystallization of the sample prior to ionization.

5 Since ESI can produce charged molecules directly from solution, it is compatible with samples from liquid chromatography systems. For example, a mass spectrometer can have an inlet for a liquid chromatography system, such as an HPLC, so that fractions flow from the chromatography column into the mass spectrometer. This in-line arrangement of a liquid chromatography system and mass spectrometer is
10 sometimes referred to as LC- MS. A LC-MS system can be used, for example, to separate metabolic analytes, products and reference molecules from complex mixtures before mass spectrometry analysis. In addition, chromatography can be used to remove salts or other buffer components from the sample before mass spectrometry analysis. For example, desalting of a sample using a reversed-phase
15 HPLC column, in-line or off-line, can be used to increase the efficiency of the ionization process and thus improve sensitivity of detection by mass spectrometry.

20 A variety of mass analyzers are available that can be paired with different ion sources. Different mass analyzers have different advantages as known to one skilled in the art and as described herein. The mass spectrometer and methods chosen for detection depends on the particular assay, for example, a more sensitive mass analyzer can be used when a small amount of ions are generated for detection. Several types of mass analyzers and mass spectrometry methods are described below.

25

Quadrupole mass spectrometry utilizes a quadrupole mass filter or analyzer. This type of mass analyzer is composed of four rods arranged as two sets of two electrically connected rods. A combination of rf and dc voltages are applied to each pair of rods which produces fields that cause an oscillating movement of the ions as
30 they move from the beginning of the mass filter to the end. The result of these fields is the production of a high-pass mass filter in one pair of rods and a low-pass filter in the other pair of rods. Overlap between the high-pass and low- pass filter leaves a defined m/z that can pass both filters and traverse the length of the quadrupole. This m/z is

selected and remains stable in the quadrupole mass filter while all other m/z have unstable trajectories and do not remain in the mass filter. A mass spectrum results by ramping the applied fields such that an increasing m/z is selected to pass through the mass filter and reach the detector. In addition, quadrupoles can also be set up to 5 contain and transmit ions of all m/z by applying a rf-only field. This allows quadrupoles to function as a lens or focusing system in regions of the mass spectrometer where ion transmission is needed without mass filtering. This will be of use in tandem mass spectrometry as described further below.

10 A quadrupole mass analyzer, as well as the other mass analyzers described herein, can be programmed to analyze a defined m/z or mass range. This property of mass spectrometers is useful for the invention described herein. Since the mass range of metabolic analytes, enzymatic products and reference molecules will be known prior to an assay, a mass spectrometer can be programmed to transmit ions of the 15 projected correct mass range while excluding ions of a higher or lower mass range. The ability to select a mass range can decrease the background noise in the assay and thus increase the signal- to-noise ratio as well as increasing the specificity of the assay. Therefore, the mass spectrometer can accomplish an inherent separation step as well as detection and identification of metabolic analytes, enzymatic products and 20 reference molecules.

Ion trap mass spectrometry utilizes an ion trap mass analyzer. In these mass analyzers, fields are applied so that ions of all m/z are initially trapped and oscillate in the mass analyzer. Ions enter the ion trap from the ion source through a focusing 25 device such as an octapole lens system. Ion trapping takes place in the trapping region before excitation and ejection through an electrode to the detector. Mass analysis is accomplished by sequentially applying voltages that increase the amplitude of the oscillations in a way that ejects ions of increasing m/z out of the trap and into the detector. In contrast to quadrupole mass spectrometry, all ions are retained in the 30 fields of the mass analyzer except those with the selected m/z. One advantage to ion traps is that they have very high sensitivity, as long as one is careful to limit the number of ions being trapped at one time. Control of the number of ions can be accomplished by varying the time over which ions are injected into the trap. The mass

resolution of ion traps is similar to that of quadrupole mass filters, although ion traps do have low m/z limitations.

Time-of-flight mass spectrometry utilizes a time-of-flight mass analyzer. For this 5 method of m/z analysis, an ion is first given a fixed amount of kinetic energy by acceleration in an electric field (generated by high voltage). Following acceleration, the ion enters a field-free or "drift" region where it travels at a velocity that is inversely proportional to its m/z. Therefore, ions with low m/z travel more rapidly than ions with high m/z. The time required for ions to travel the length of the field-free region is 10 measured and used to calculate the m/z of the ion.

One consideration in this type of mass analysis is that the set of ions being 15 studied be introduced into the analyzer at the same time. For example, this type of mass analysis is well suited to ionization techniques like MALDI which produces ions in short well-defined pulses. Another consideration is to control velocity spread produced by ions that have variations in their amounts of kinetic energy. The use of longer flight tubes, ion reflectors, or higher accelerating voltages can help minimize the 20 effects of velocity spread. Time-of-flight mass analyzers have a high level of sensitivity and a wider m/z range than quadrupole or ion trap mass analyzers. Also data can be acquired quickly with this type of mass analyzer because no scanning of the mass analyzer is necessary.

Tandem mass spectrometry can utilize combinations of the mass analyzers 25 described above. Tandem mass spectrometers can use a first mass analyzer to separate ions according to their m/z in order to isolate an ion of interest for further analysis. The isolated ion of interest is then broken into fragment ions (called collisionally activated dissociation or collisionally induced dissociation) and the fragment ions are analyzed by the second mass analyzer. These types of tandem mass spectrometer systems are called tandem in space systems because the two mass 30 analyzers are separated in space, usually by a collision cell. Tandem mass spectrometer systems also include tandem in time systems where one mass analyzer is used, however the mass analyzer is used sequentially to isolate an ion, induce fragmentation, and then perform mass analysis.

Mass spectrometers in the tandem in space category have more than one mass analyzer. For example, a tandem quadrupole mass spectrometer system can have a 5 first quadrupole mass filter, followed by a collision cell, followed by a second quadrupole mass filter and then the detector. Another arrangement is to use a quadrupole mass filter for the first mass analyzer and a time-of-flight mass analyzer for the second mass analyzer with a collision cell separating the two mass analyzers. Other tandem systems are known in the art including reflectron-time-of-flight, tandem 10 sector and sector-quadrupole mass spectrometry.

Mass spectrometers in the tandem in time category have one mass analyzer that performs different functions at different times. For example, an ion trap mass spectrometer can be used to trap ions of all m/z. A series of rf scan functions are 15 applied which ejects ions of all m/z from the trap except the m/z of ions of interest. After the m/z of interest has been isolated, an rf pulse is applied to produce collisions with gas molecules in the trap to induce fragmentation of the ions. Then the m/z values of the fragmented ions are measured by the mass analyzer. Ion cyclotron resonance instruments, also known as Fourier transform mass spectrometers, are an 20 example of tandem-in-time systems.

Several types of tandem mass spectrometry experiments can be performed by controlling the ions that are selected in each stage of the experiment. The different types of experiments utilize different modes of operation, sometimes called "scans," of 25 the mass analyzers. In a first example, called a mass spectrum scan, the first mass analyzer and the collision cell transmit all ions for mass analysis into the second mass analyzer. In a second example, called a product ion scan, the ions of interest are mass-selected in the first mass analyzer and then fragmented in the collision cell. The ions formed are then mass analyzed by scanning the second mass analyzer. In a third 30 example, called a precursor ion scan, the first mass analyzer is scanned to sequentially transmit the mass analyzed ions into the collision cell for fragmentation. The second mass analyzer mass-selects the product ion of interest for transmission to the detector. Therefore, the detector signal is the result of all precursor ions that can

be fragmented into a common product ion. Other experimental formats include neutral loss scans where a constant mass difference is accounted for in the mass scans. The use of these different tandem mass spectrometry scan procedures can be advantageous when large sets of analytes are measured in a single experiment as

5 with multiplex experiments.

In view of the above, those skilled in the art recognize that different mass spectrometry methods, for example, quadrupole mass spectrometry, ion trap mass spectrometry, time-of-flight mass spectrometry and tandem mass spectrometry, can

10 use various combinations of ion sources and mass analyzers which allows for flexibility in designing customized detection protocols. In addition, mass spectrometers can be programmed to transmit all ions from the ion source into the mass spectrometer either sequentially or at the same time. Furthermore, a mass spectrometer can be programmed to select ions of a particular mass for transmission

15 into the mass spectrometer while blocking other ions. The ability to precisely control the movement of ions in a mass spectrometer allows for greater options in detection protocols which can be advantageous when a large number of analytes, for example, from a multiplex experiment, are being analyzed.

20 Different mass spectrometers have different levels of resolution, that is, the ability to resolve peaks between ions closely related in mass. The resolution is defined as $R=m/\Delta m$, where m is the ion mass and Δm is the difference in mass between two peaks in a mass spectrum. For example, a mass spectrometer with a resolution of 1000 can resolve an ion with a m/z of 100.0 from an ion with a m/z of

25 100.1. Those skilled in the art will therefore select a mass spectrometer having a resolution appropriate for the analyte(s) to be detected.

Mass spectrometers can resolve ions with small mass differences and measure the mass of ions with a high degree of accuracy. Therefore, analytes of similar masses

30 can be used together in the same experiment since the mass spectrometer can differentiate the mass of even closely related molecules. The high degree of resolution and mass accuracy achieved using mass spectrometry methods allows the use of large sets of analytes because they can be distinguished from each other.

Additional mass spectrometry methods are well known in the art (see Burlingame et al. Anal. Chem. 70:647R-716R (1998); Kinter and Sherman, New York (2000)). Exemplary descriptions of mass spectrometry methods for detecting 5 metabolic analytes include Chace DH, Hillman SL, Van Hove JLK, Naylor EW. Clin Chem 1997;43:210613; Rashed MS, Bucknall MP, Little D, et al. Clin Chem 1997;43:112941; Matern D, Strauss AW, Hillman SL, Mayatepek E, Millington DS, Trefz FK. Pediatr Res 1999;46:459, and Millington DS, Kodo N, Terada N, Roe D, Chace DH. International Journal of Mass Spectrometry and Ion Processes 10 1991;111:21128.

One or more steps of a method of the invention can be automated, if desired. Those skilled in the art will know how to use autosamplers, liquid handling devices, robots, computers and other routine laboratory instruments for automation.

15

All journal articles, reference, and patent citations provided above, in parentheses or otherwise, whether previously stated or not, are incorporated herein by reference.

20

It is understood that modifications that do not substantially affect the function of the various embodiments of this invention are also included within the definition of the invention provided herein. Accordingly, the following examples are intended to illustrate but not limit the present invention.

EXAMPLE 1

This example describes simultaneous determination of the amount of a
5 metabolic analyte and an metabolically indicative enzyme product from a patient
sample.

The enzyme biotinidase is a known metabolically indicative enzyme for the
metabolic disorder termed biotinidase deficiency. In the untreated state, profound
10 biotinidase deficiency in newborns typically is characterized by neurological and skin-
related symptoms, such as seizures, hypotonia, and rash, often accompanied by
hyperventilation, laryngeal stridor, and apnea. Older children may also have alopecia,
ataxia, developmental delay, neurosensory hearing loss, optic atrophy, and recurrent
15 infections. Individuals with partial biotinidase deficiency may have hypotonia, skin
rash, and hair loss, particularly during times of stress. Once detected, biotinidase
deficiency is often treatable by administration of biotin.

It has been previously established that individuals with profound biotinidase
deficiency have <10% of mean normal serum biotinidase activity. Individuals with
20 partial biotinidase deficiency have 10-30% of mean normal serum biotinidase activity.
Therefore, it is understood that a decrease in biotinidase activity compared to normal
reference level is indicative of biotinidase deficiency.

Acylcarnitines and α -amino acid also were assayed. Different patterns of the
25 plasma acylcarnitine profile can indicate the diagnosis of fatty acid oxidation disorders
(MCAD, VLCAD, SCAD, MAD, LCHAD, and CPTII), as well as some organic
acidemias (propionic acidemia, methylmalonic acidemia, isovaleric acidemia, glutaric
acidemia type I, 3-methylcrotonyl CoA carboxylase deficiency, B-ketothiolase
30 deficiency, etc.). Plasma amino acids analysis will detect specific amino acid
disorders such as phenylketonuria (PKU), Maple Syrup Urine Disease (MSUD), urea
cycle defects, non-ketotic hyperglycinemia and homocystinuria. This analysis is also
indicated in patients with general failure to thrive as an initial screen. For assessment

of a metabolic disorder, plasma amino acids are compared to age-matched normal values

Activity of the enzyme biotinidase and the amount of the α -amino acid and 5 acylcarnitine analytes were measured simultaneously as follows:

Dried blood samples on paper were each punched into a well of a microwell plate. Each sample was reconstituted with 50 microliter of buffer solution containing isotopically labeled biocytin (the substrate for biotinidase), and the mixture is 10 incubated for 17 hours at 42°C. 140 microliter of methanol containing isotopically labeled internal standards for α -amino acids and acylcarnitines and for the product of the biotinidase reaction were then added, and the plate was shaken for 30 minutes at 27°C. The plate was then covered with aluminum foil and placed on an autosampler. Each processed sample was analyzed by tandem mass spectrometry, and 15 concentrations of biotinidase product, endogenous α -amino acids and endogenous acylcarnitines were quantified based on internal standards.

Tables 6 and 7 show data obtained using the above method, as compared to data obtained using well known mass spectrometry screening methods for metabolic 20 analytes without simultaneously measuring enzyme activity. A slope of one indicates identical results between the compared methods.

Table 6: α -Amino acid (AA) comparison:

25

AA	SLOPE	R2
ALA	1.04	0.9916
ARG	1.11	0.9975
CIT	1.13	0.9912
GLY	1.36	0.9836
LEU	1.14	0.9987
MET	1.00	0.9989
ORN	1.02	0.9951
PHE	0.99	0.9949
TYR	1.10	0.9917
VAL	1.04	0.9836
MEAN	1.09	0.9927

30

Table 7: Acylcarnitine comparison:

AC	SLOPE	R2
C0	1.05	0.9994
C2	1.12	0.9902
C3	1.11	0.9915
C4	1.10	0.9946
C5DC	0.89	0.9641
C6	0.99	0.9973
C8	1.00	0.9987
C10	1.05	0.9987
C12	1.01	0.9968
C14	1.06	0.9952
C16	1.21	0.9837
C18	1.16	0.9844
MEAN	1.06	0.9912

These results show that the above method produces data that correlate with well known mass spectrometry screening methods in which α -amino acids and acylcarnitines are extracted and measured without measuring enzyme activity simultaneously. These results also show that the above method produces data that correlate with known enzyme activity screening methods that measure fluorescence of a tagged product without measuring simultaneously concentrations of α -amino acids and acylcarnitines.

What is claimed is:

1. A method for detecting a metabolic disorder in an individual, comprising:
 - 5 (a) contacting a sample comprising:
 - (i) one or more metabolically indicative enzymes and
 - (ii) one or more metabolic analytes,
with one or more substrates for said one or more enzymes to produce a reaction admixture, under conditions in which at least one of said enzymes is capable
 - 10 of acting on a corresponding substrate to generate at least one product;
 - (b) contacting said reaction admixture with a reagent that inhibits the ability of said one or more enzymes to act on a corresponding substrate, wherein said one or more metabolic analytes and said at least one product are soluble in said reagent; to produce a test sample and
 - 15 (c) determining the presence or amount of said one or more metabolic analytes and said at least one product contained in said test sample using mass spectrometry,
wherein a determined presence or amount of said one or more metabolic analytes and said at least one product correlates with presence or absence of said
 - 20 metabolic disorder.
2. The method of claim 1, wherein said sample is a body fluid sample.
3. The method of claim 2, wherein said body fluid sample is blood.
- 25 4. The method of claim 1, wherein said sample is dried.
5. The method of claim 1, wherein said individual is a human suspected of having a metabolic disorder.
- 30 6. The method of claim 1, wherein said individual is a neonate.
7. The method of claim 1, wherein said individual is a newborn.

8. The method of claim 1, wherein said individual is a child.
9. The method of claim 1, wherein said individual is an adult.
- 5 10. The method of claim 1, wherein said metabolic disorder is an inborn error of metabolism.
- 10 11. The method of claim 1, wherein said metabolic disorder is an acquired metabolic disorder.
12. The method of claim 1, wherein said metabolically indicative enzyme is selected from the group consisting of oxidoreductase, hydrolase, lyase, transferase, ligase and isomerase.
- 15 13. The method of claim 12, wherein the enzyme is a hydrolase.
14. The method of claim 13, wherein the enzyme is a biotinidase.
- 20 15. The method of claim 14, wherein said substrate is biocytin.
16. The method of claim 1, wherein said metabolic analyte is one or more amino acids.
- 25 17. The method of claim 1, wherein said metabolic analyte is an acylcarnitine or plurality of acylcarnitines.
18. The method of claim 1, wherein step (a) further comprises contacting said sample with one or more reference substrates.
- 30 19. The method of claim 1, wherein step (b) further comprises contacting said sample with one or more reference products.

20. The method of claim 1, wherein step (d) further comprises, prior to determining, adding one or more reference products corresponding to the at least one product .

5

21. The method of claim 1, wherein step (d) further comprises, prior to determining, adding one or more reference analytes corresponding to the one or more metabolic analytes contained in said sample.

10

22. The method of claim 1, wherein said reaction admixture is aqueous.

23. The method of claim 22, wherein said reagent is non-aqueous.

15

24. The method of claim 23, wherein said reagent comprises an organic solvent.

25. The method of claim 23, wherein said reagent comprises an alcohol.

26. The method of claim 25, wherein said reagent is methanol.

20

27. The method of claim 1, wherein said mass spectrometry is tandem mass spectrometry.

28. A method for detecting a metabolic disorder in an individual, comprising:

25

(a) contacting a vessel containing one or more substrates in dried form with a sample from said individual, the sample containing

- (i) one or more metabolically indicative enzymes and
- (ii) one or more metabolic analytes,

in aqueous solution;

30

under conditions wherein at least one of said enzymes is capable of acting on a corresponding substrate to generate at least one product;

(b) adding a reagent that inhibits the ability of said one or more enzymes to act on a corresponding substrate, wherein said one or more metabolic analytes and at least one product are soluble in said reagent;

5 (c) determining the presence or amount of said one or more metabolic analytes and said at least one product in the resultant mixture using mass spectrometry,

wherein a determined presence or amount of said one or more metabolic analytes and said at least one product correlates with presence or absence of said metabolic disorder.

10 .

29. The method of claim 28, wherein said sample is a body fluid sample.

30. The method of claim 29, wherein said body fluid sample is blood.

15

31. The method of claim 30, wherein said sample is dried.

32. The method of claim 28, wherein said individual is a human suspected of having a metabolic disorder.

20

33. The method of claim 28, wherein said individual is a neonate.

34. The method of claim 28, wherein said individual is a newborn.

35. The method of claim 28, wherein said individual is a child.

25

36. The method of claim 28, wherein said individual is an adult.

37. The method of claim 28, wherein said metabolic disorder is an inborn error of metabolism.

30

38. The method of claim 28, wherein said metabolic disorder is an acquired metabolic disorder.

39. The method of claim 28, wherein said metabolically indicative enzyme is selected from the group consisting of oxidoreductase, hydrolase, lyase, transferase, ligase and isomerase.

5 40. The method of claim 39, wherein the enzyme is a hydrolase.

41. The method of claim 40, wherein the enzyme is a biotinidase.

42. The method of claim 41, wherein said substrate is biocytin.

10 43. The method of claim 28, wherein said metabolic analyte is one or more amino acids.

15 44. The method of claim 28, wherein said metabolic analyte is an acylcarnitine or plurality of acylcarnitines.

45. The method of claim 28, wherein said vessel further contains one or more reference substrates.

20 46. The method of claim 28, wherein said vessel further contains one or more reference analytes corresponding to the one or more metabolic analytes in said sample.

25 47. The method of claim 28, wherein said vessel further contains one or more reference products.

48. The method of claim 28, wherein said reagent is non-aqueous.

49. The method of claim 48, wherein said reagent is an organic solvent.

30 50. The method of claim 28, wherein said reagent contains alcohol.

51. The method of claim 50, wherein said alcohol is methanol.

52. The method of claim 28, wherein said mass spectrometry is tandem mass spectrometry.

5 53. The method of claim 28, wherein said vessel is a well of a multi-well plate.

54. A method for detecting a metabolic disorder in an individual, comprising:

(a) contacting a sample comprising:

(i) one or more metabolically indicative enzymes and

10 (ii) one or more metabolic analytes,

with one or more substrates for said one or more enzymes to produce a reaction admixture, under conditions in which at least one of said enzymes is capable of acting on a corresponding substrate to generate at least one product;

(b) contacting said reaction admixture with a reagent that inhibits the ability

15 of said one or more enzymes to act on a corresponding substrate, wherein said one or more metabolic analytes and said at least one product are soluble in said reagent, to produce a test sample;

(c) contacting said reaction admixture with one or more reference products and one or more reference analytes, and

20 (d) determining the presence or amount of said one or more metabolic analytes and said at least one product contained in said test sample, with respect to said one or more reference products and one or more reference analytes, using mass spectrometry,

wherein a determined presence or amount of said one or more metabolic analytes and said at least one product correlates with presence or absence of said metabolic disorder.

55. A method for detecting a metabolic disorder in an individual, comprising:

(a) separating a sample into two sample portions, the sample comprising:

30 (i) one or more metabolically indicative enzymes and

(ii) one or more metabolic analytes;

(b) contacting each sample portion with one or more substrates for said one or more enzymes to produce two reaction admixtures, under conditions in which at

least one of said enzymes in each sample portion is capable of acting on a corresponding substrate to generate at least one product;

(c) contacting each reaction admixture with a reagent that inhibits the ability of said one or more enzymes to act on a corresponding substrate, wherein said

5 reagent can be the same or different for each reaction admixture, and wherein said one or more metabolic analytes and said at least one product are soluble in said reagent or reagents; to produce a test sample and

(d) combining the two reaction admixtures to produce a test sample, and

10 (e) determining the presence or amount of said one or more metabolic analytes and said at least one product contained in said test sample using mass spectrometry,

wherein a determined presence or amount of said one or more metabolic analytes and said at least one product correlates with presence or absence of said metabolic disorder.

15

56. A method for detecting a metabolic disorder in an individual, comprising:

(a) separating a sample into two sample portions, the sample comprising:

(i) one or more metabolically indicative enzymes and

(ii) one or more metabolic analytes;

20 (b) contacting each sample portion with individual vessels, each vessel containing one or more substrates for said one or more enzymes in dried form,

(c) contacting each vessel with a solution to produce two reaction admixtures, under conditions in which at least one of said enzymes in each sample portion is capable of acting on a corresponding substrate to generate at least one product;

(d) contacting each reaction admixture with a reagent that inhibits the ability of said one or more enzymes to act on a corresponding substrate, wherein said reagent can be the same or different for each reaction admixture, and wherein said one or more metabolic analytes and said at least one product are soluble in said reagent or reagents; to produce a test sample and

30 (e) combining the two reaction admixtures to produce a test sample, and

(f) determining the presence or amount of said one or more metabolic analytes and said at least one product contained in said test sample using mass spectrometry,

wherein a determined presence or amount of said one or more metabolic analytes and said at least one product correlates with presence or absence of said metabolic disorder.

57. A method for detecting a metabolic disorder in an individual, comprising:

5 (a) contacting a vessel containing one or more substrates in dried form with
10 a sample from said individual, the sample containing


- (i) one or more metabolically indicative enzymes and
- (ii) one or more metabolic analytes,

15 (b) contacting said vessel with a solution to generate conditions wherein at least one of said enzymes is capable of acting on a corresponding substrate to generate at least one product;

(c) adding a reagent that inhibits the ability of said one or more enzymes to act on a corresponding substrate, wherein said one or more metabolic analytes and at least one product are soluble in said reagent;

20 (d) determining the presence or amount of said one or more metabolic analytes and said at least one product in the resultant mixture using mass spectrometry,

wherein a determined presence or amount of said one or more metabolic analytes and said at least one product correlates with presence or absence of said metabolic disorder.



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US04/28238

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12Q 1/00, 1/34; G01N 33/53
US CL : 435/4, 7.71, 7.72, 18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/4, 7.71, 7.72, 18

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 6,258,605 B1 (CHACE) 10 July 2001 (10.07.2001), column 1, lines 36-55 and column 5, line 60-column 8, line 49	1-57
Y	US 5,719,035 (ROSENTHAL et al) 17 February 1998 (17.02.1998), column 4, lines 12-63	1-57
Y	IM, W.B. et al. Bacterial Degradation of Biotin. J. Biol. Chem. Volo. 248, No.22, p.7798-7805, especially p. 7798, 7799 and 7801.	1-57
Y	US 6,455,321 B1 (CHACE) 24 September 2002 (24.09.2002), column 6, line 9-column 10, line 57	1-57
A	US 5,629,210 (HERCULES et al) 13 May 1997 (13.05.1997), column 1, lines 15-42.	1-57
A, P	US 6,670,194 B1 (AEBERSOLD et al) 30 December 2003 (30.12.2003), entire document.	1-57

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

14 December 2004 (14.12.2004)

Date of mailing of the international search report

13 JAN 2005

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Faxsimile No. (703) 305-3230

Authorized officer

Lora E Barnhart

Telephone No. 571-272-1600

**DEBORAH A. THOMAS
PARALEGAL SPECIALIST**

GROUP 1000 Dct