



- (51) International Patent Classification:
C07K 14/47 (2006.01) *C12N 15/113* (2010.01)
- (21) International Application Number:
PCT/IB2013/001990
- (22) International Filing Date:
5 August 2013 (05.08.2013)
- (25) Filing Language: English
- (26) Publication Language: English
- (71) Applicants: **SOCIÉTÉ DE DÉVELOPPEMENT ET DE RECHERCHE INDUSTRIELLE** [FR/FR]; 42 Rue de Longvic, F-21300 Chenove (FR). **CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE - CNRS -** [FR/FR]; 3, rue Michel Ange, F-75794 Paris Cedex 16 (FR). **UNIVERSITÉ PARIS DIDEROT - PARIS 7** [FR/FR]; 5, rue Thomas Mann, F-75013 Paris (FR). **INSERM (INSTITUT NATIONAL DE LA SANTÉ ET DE LA RECHERCHE MÉDICALE)** [FR/FR]; 101, rue de Tolbiac, F-75013 Paris (FR). **ÉCOLE NORMALE SUPÉRIEURE** [FR/FR]; 45 rue d'Ulm, F-75005 Paris (FR).
- (72) Inventors: **DUGAST-DARZACQ, Claire**; 24 avenue Raymond Aron, F-92160 Antony (FR). **NOIZET, Maïté**; 76-78 Avenue de Paris, Bâtiment Le Vau, F-78000 Versailles (FR). **DARZACQ, Xavier**; 24 avenue Raymond Aron, F-92160 Antony (FR).
- (74) Agents: **BERNARDI, Céline** et al.; Cabinet Plasseraud, 52 rue de la Victoire, F-75440 Paris Cedex 09 (FR).
- (81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- Published:
- with international search report (Art. 21(3))
 - with sequence listing part of description (Rule 5.2(a))

(54) Title: MOLECULAR TARGETS FOR THE TREATMENT OF WOUNDS, IN PARTICULAR CHRONIC WOUNDS

(57) Abstract: The present invention relates to a therapeutic compound comprising: - an agent that inhibits the activity of at least one gene selected from the group consisting of MEOX2, MAF, SIX2 and homologues thereof having at least 50% identity with said genes and/or - an agent that enhances the activity of at least one gene selected from the group consisting of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, FOXS1, GLI1, SOX9 and homologues thereof having at least 50% identity with said gene for use in the treatment of wounds, preferably chronic wounds.



Molecular targets for the treatment of wounds, in particular chronic wounds

Field of the invention

The present invention relates to at least one molecular target for healing or treating wounds, in particular chronic wounds and more preferentially non healing chronic wounds. Further, the invention concerns a novel therapeutic for treating wounds and a novel gene therapy approach, involving said molecular target for treating wounds. Additionally, the invention concerns a method for treating wounds using said therapeutic or said gene therapy.

Background of the invention

The natural wound healing is divided into three sequential phases; each phase is characterized by specific cellular activities: the inflammatory phase, the proliferative phase and the remodeling phase.

The first phase, called the inflammatory phase, begins minutes after injury. The blood vessels rupture induces the clot formation, composed mainly of fibrin and fibronectin. The clot fills partially the lesion and allows the migration of the inflammatory cells within the lesion. The inflammatory cells are recruited to debride the wound. Platelets secrete factors, such as growth factors or cytokines, which induce the recruitment of cells implicated in the wound healing (inflammatory cells such as neutrophils and macrophages, fibroblasts and endothelial cells).

The second phase is called the proliferative phase and corresponds to the development of the granulation tissue. Fibroblasts migrate into the wound area, proliferate and form a new provisional extracellular matrix by secreting extracellular matrix (ECM) proteins. Then endothelial cells migrate to promote the neovascularization or angiogenesis of the lesion. Inside the granulation tissue, fibroblasts activate and differentiate into myofibroblasts, presenting contractile properties thanks to their expression of alpha-smooth muscle actin (similar to that in smooth muscle cells). Myofibroblasts have a key role in wound healing as they provide the contraction of the wound. Finally, keratinocytes migrate from the wound edge, proliferate and differentiate to reconstitute the epidermis.

The last phase of the wound healing process appears after the wound closure. It corresponds to the remodeling of the granulation tissue. The granulation tissue is reorganized, type III collagen is replaced by type I collagen, as normal dermis is principally composed of type I

collagen. During this phase, myofibroblasts in excess are eliminated by apoptosis. The last phase of the wound healing is long. One year after injury, the scar is remodeled; it gets less red and thinner.

However, this process is not only complex but fragile; it is susceptible to interruption or failure leading to the formation of chronic or non-healing wounds or formation of abnormal scars. Factors which may contribute to this include diseases (such as diabetes, venous or arterial disease), age, infection or tissue localization.

Chronic or non-healing wounds

Chronic wounds are a worldwide health problem, in part due to a lack of adequate methods of treatment. In 2010, more than 7 million people worldwide suffered from chronic wounds, and the projected annual increase is at least ten percent.

Chronic wounds are sometimes non-healing wounds. Common types of chronic wounds include, but are not limited to, venous leg ulcers, diabetic foot ulcers, decubitus ulcers, arterial leg ulcers, those of mixed etiology (venous and arterial) or those with no known etiology. We can also find acute wounds that become chronic as they do not heal correctly.

Non-healing wounds or chronic wounds are a challenge for the patient, the health care professional, and the health care system. They significantly impair the quality of life for millions of people and impart burden on society in terms of lost productivity and health care money.

Wound healing is a dynamic pathway that leads to the restoration of tissue integrity and functions. A chronic wound or non-healing wound develops when the normal reparative process is disturbed. By understanding the biology of wound healing, the physician can optimize the wound healing by choosing the adequate treatment.

In chronic or non-healing wounds, the natural healing process is altered, and thus healing is prolonged, incomplete and sometimes wounds never close. A chronic wound occurs when some factor causes the disruption of the normal inflammatory and proliferative phases. By enhancing or manipulating factors involved in wound healing it may therefore be possible to correct the process, thereby reducing the likely occurrence of a chronic wound or accelerate its subsequent repair.

Role of fibroblasts in wound healing

Fibroblasts are implicated in the process of wound healing, this involves several steps of differentiation from a quiescent fibroblast to a mobilized fibroblast that will transform into a myofibroblast and finally enter apoptosis. In chronic or non-healing wounds, this process is misregulated and fibroblasts fail to undertake the myofibroblast differentiation and are found in the wound as unfunctional fibroblasts, called pseudo senescent fibroblasts (Telgenhoff D, Shroot B (2005) Cellular senescence mechanisms in chronic wound healing. *Cell Death Differ* 12: 695-698). The aim of the present invention is to map, at the whole genome scale, the different genes that needs to be activated or deactivated during this process, and thus facilitates the healing process of the wound.

Human fibroblasts have the ability to enter into a physiological process named senescence, which permits a limited replicative cell cycle and thus avoids loss of genetic information. It usually occurs when a cell has already conducted several rounds of replication (called replicative senescence and dependent from telomere length), but can also occur in response to environmental stress (Muller M (2009) Cellular senescence: molecular mechanisms, in vivo significance, and redox considerations. *Antioxid Redox Signal* 11: 59-98). Senescence cells are arrested in cell cycle but maintain metabolic activity (Telgenhoff D, Shroot B (2005) Cellular senescence mechanisms in chronic wound healing. *Cell Death Differ* 12: 695-698).

Fibroblasts of chronic wounds lose some of their functionalities, and more particularly, they lose part or all of their replicative function (Telgenhoff D, Shroot B (2005) Cellular senescence mechanisms in chronic wound healing. *Cell Death Differ* 12: 695-698). In a wound, human fibroblasts are also associated with an up-regulation of APA-1, a protein which induces matrix remodeling, demonstrating that pseudo-senescence fibroblast phenotype was not induced by telomere attrition (Benanti JA, Williams DK, Robinson KL, Ozer HL, Galloway DA (2002) Induction of extracellular matrix-remodeling genes by the senescence-associated protein APA-1. *Mol Cell Biol* 22: 7385-7397). Thus, senescent fibroblasts in chronic wounds would appear more particularly due to a chronic inflammation than to a telomere shortening (Telgenhoff D, Shroot B (2005) Cellular senescence mechanisms in chronic wound healing. *Cell Death Differ* 12: 695-698).

The present invention can improve the quality of a patient's life by treating wounds in a way that actively promotes healing.

SUMMARY OF THE INVENTION

The present invention relates to a therapeutic compound comprising :

- 5 - an agent that inhibits the activity of at least one gene selected from the group consisting of MEOX2, MAF, SIX2 and homologues thereof having at least 50% identity with said genes and/or
- an agent that enhances the activity of at least one gene selected from the group consisting of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, FOXS1, GLI1, SOX9,
- 10 preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 and homologues thereof having at least 50% identity with said gene
- for use in the treatment of wounds, preferably chronic wounds.

The invention also relates to a pharmaceutical composition comprising a therapeutic compound as defined above, together with a pharmaceutically acceptable carrier.

- 15 In one aspect, the invention relates to a method for preparing a pharmaceutical composition as defined above comprising bringing said therapeutic compound in conjunction or association with a pharmaceutically or veterinary acceptable carrier or vehicle.

In another aspect, the invention relates to a method for treating a mammalian wound, wherein said method comprises: administering to said wound a therapeutic compound comprising:

- 20 - an agent that inhibits the activity of at least one gene selected from the group consisting of MEOX2, MAF, SIX2 and homologues thereof having at least 50% identity with said genes and/or
- an agent that enhances the activity of at least one gene selected from the group consisting of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, FOXS1, GLI1, SOX9,
- 25 preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 and homologues thereof having at least 50% identity with said gene

The present invention also relates to a kit for treating a wound, preferably a chronic wound, wherein said kit comprises:

- (a) at least one therapeutic compound or pharmaceutical composition as defined above; and

(b) at least one dressing for applying to said wound.

In yet another aspect, the invention relates to a combination therapeutic for treating a wound, preferably a chronic wound, comprising:

- 5 a) an agent that inhibits the activity of at least one gene selected from the group consisting of MEOX2, MAF, SIX2 and homologues thereof having at least 50% identity with said genes and/or
- an agent that enhances the activity of at least one gene selected from the group consisting of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, FOXS1, GLI1, SOX9,
- 10 preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 and homologues thereof having at least 50% identity with said gene
- and
- b) at least one further therapeutic.

- 15 The present invention also relates to the use of
- an agent that inhibits the activity of at least one gene selected from the group consisting of MEOX2, MAF, SIX2 and homologues thereof having at least 50% identity with said genes and/or
- an agent that enhances the activity of at least one gene selected from the group consisting of
- 20 CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, FOXS1, GLI1, SOX9, preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 and homologues thereof having at least 50% identity with said gene
- for treating a wound, preferably a chronic wound, wherein said agent modulates fibroblast and myofibroblast differentiation and/or activity.

25

DETAILED DESCRIPTION OF THE INVENTION

Accordingly, in one aspect of the invention there is provided a therapeutic compound comprising:

- an agent that inhibits the activity of at least one gene selected from the group consisting of MEOX2, MAF, SIX2 and homologues thereof having at least 50% identity with said genes and/or

- an agent that enhances the activity of at least one gene selected from the group consisting of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, FOXS1, GLI1, SOX9, preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 and homologues thereof having at least 50% identity with said gene

for use in the treatment of wounds, preferably chronic wounds.

In a preferred embodiment of the invention, said therapeutic composition also comprises at least:

(a) - an agent that inhibits the activity of PPARG or homologues thereof having at least 50% identity with PPARG and/or

(b) - an agent that enhances the activity of at least one gene selected from the group consisting of SMAD3, SMAD4 and homologues thereof having at least 50% identity with said genes.

Indeed, the inventors have found that these genes were involved in the healing process and that their down-regulation or up-regulation would be useful for the treatment of wounds, in particular chronic wounds.

Thus, the invention involves a novel gene therapy approach and/or a novel protein therapy approach.

As used herein, the expression “agent that inhibits the activity of a gene” or “inhibitor” refers to an agent than can downregulate said gene. It encompasses agents that act at the level of the gene expression, as well as agents that act at the level of the protein, either by decreasing the amount of protein present in a given cell, or by inhibiting the protein’s activity.

As used herein, the expression “agent that enhances the activity of a gene” or “enhancer” refers to an agent than can upregulate said gene. It encompasses agents that act at the level of the gene expression, as well as agents that act at the level of the protein, either by increasing the amount of protein present in a given cell, or by increasing the protein’s activity. It also encompasses agents that act upstream or downstream of said gene or protein in a signaling pathway.

In one embodiment of the invention, the novel therapeutic comprises an inhibitor or enhancer of the gene expression, this inhibitor or enhancer can be either an anti-sense DNA or RNA, siRNA, shRNA, cDNA, TALENS or ribozymes, either naked or in the form of plasmid and viral vectors or a drug.

In a preferred embodiment, said therapeutic compound is a siRNA selected from the group consisting of siRNA having the sequence as set forth in SEQ ID No:1 to SEQ ID No:8 and mixtures thereof.

Typically, the expression of the MAF gene can be inhibited by the siRNAs having the sequences as set forth in SEQ ID No:1 and/or SEQ ID No:2.

SEQ ID NO:1: 5'-UUCGAUCUGAUGAAGUUUGAA

SEQ ID NO:2: 5'-CGGCAGGAGAAUGGCAUCAGA

Typically, the expression of the MEOX2 gene can be inhibited by the siRNAs having the sequences as set forth in SEQ ID No:3 and/or SEQ ID No:4.

SEQ ID NO:3: 5'-AGCAUGCGCACUUAUGAUUA

SEQ ID NO:4: 5'-CCCGCCCGUCCUGUGCUCCAA

Typically, the expression of the PPARG gene can be inhibited by the siRNAs having the sequences as set forth in SEQ ID No:5 and/or SEQ ID No:6.

SEQ ID NO:5: 5'-UCCGGAGAACAAUCAGAUUGA

SEQ ID NO:6: 5'-GAGGGCGAUCUUGACAGGAAA

Typically, the expression of the SIX2 gene can be inhibited by the siRNAs having the sequences as set forth in SEQ ID No:7 and/or SEQ ID No:8.

SEQ ID NO:7: 5'-CAACGAGAACUCCAAUUCUAA

SEQ ID NO:8: 5'-CCCGCUGAAUGGCAGCGGCAA

In one embodiment, the agent that enhances the activity of a gene is a cDNA.

Typically, the expression of the CREB5 gene can be enhanced by administering the CREB5 cDNA.

The same applies mutatis mutandis for all the other genes listed.

- 5 The cDNA can be administered or delivered to the wound to be treated in any suitable form known to the skilled person in art. It can be delivered as naked DNA, using plasmid vectors, viral vectors, or any other suitable means.

In another embodiment, the novel therapeutic comprises an inhibitor or enhancer of the protein encoded by the gene function, this inhibitor or enhancer can be either a binding agent
10 that binds, either reversibly or irreversibly, to inhibit or enhance protein function such as an antibody or a known, or synthesized, protein agonist or antagonist; or an agent that works upstream or downstream of the protein signaling mechanism to inhibit or enhance the protein signaling and so negate or enhance the effects of expression of the protein in wound tissue.

Said agent (inhibitor or enhancer) can be any agent known in the art to act upon a given
15 molecular target.

Typically, the enhancer of PPARG can be selected in the group consisting of thiazolidinediones, such as rosiglitazone and pioglitazone (Curr Drug Targets Cardiovasc Haematol Disord. 2005 Oct;5(5):377-86. Role of PPAR- gamma agonist thiazolidinediones in treatment of pre-diabetic and diabetic individuals: a cardiovascular perspective. Dumasia R,
20 Eagle KA, Kline-Rogers E, May N, Cho L, Mukherjee D).

Typically, the inhibitor of PPARG can be G3335 (CAS 36099-95-3) (Chembiochem. 2006 Jan;7(1):74-82. The dipeptide H-Trp-Glu-OH shows highly antagonistic activity against PPARGgamma: bioassay with molecular modeling simulation. Ye F, Zhang ZS, Luo HB, Shen JH, Chen KX, Shen X, Jiang HL.).

25

The therapeutic of the invention is for use in treating wounds, more ideally in chronic wounds, preferentially in non-healing wounds. These wounds are preferentially mammalian wounds, more preferentially human wounds. Chronic wounds that are preferably treated using the invention are leg ulcers, diabetic foot ulcers, pressure ulcers.

An antibody for use in the invention is most ideally a monoclonal antibody or a humanized antibody.

In the above aspects and embodiments of the invention the therapeutic is formulated for topical application, but it can also be formulated for oral, cutaneous, transcutaneous,
5 transdermal, intravenous or any application known.

Alternatively, in the above aspects of the invention the therapeutic is formulated for application into a dressing or impregnation of a dressing.

The therapeutic of the invention may be administered with another active agent. Such active agent may be an antibiotic or antibacterial agent, an antiseptic, an antiviral, an antifungal, an
10 analgesic, an anti-inflammatory agent, a wound healing agent, a keratolytic agent, an anesthetic agent. Such actives are well known to skilled practitioners.

In another aspect of the invention, there is provided a pharmaceutical composition for use in treating wounds comprising a therapeutic of the invention together with a pharmaceutical acceptable carrier.

15 Other active materials may also be present in the pharmaceutical composition, as may be considered appropriate or advisable for the wound being treated.

The carrier, or, if more than one be present, each of the carriers, must be acceptable in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient.

20 The formulations include those suitable for topical, oral, rectal, nasal or any administration known and may be prepared by any methods well known in the art of pharmacy.

The composition may be prepared by bringing into association the therapeutic of the invention and the carrier. In general, the formulations are prepared by uniformly and intimately bringing into association the active agent with liquid carriers or finely bringing into
25 association the active with liquid carriers or finely divided solid carriers or both, and then if necessary shaping the product. The invention extends to methods for preparing a pharmaceutical composition comprising bringing a therapeutic of the invention in conjugation or association with a pharmaceutically acceptable carrier or vehicle.

For topical application to the skin, compounds of conventional use may be made up into a cream, ointment, gel, jelly, solution or suspension etc. Cream or ointment formulations that may be used for the composition are conventional formulations well known in the art.

Formulations for oral administration in the present invention may be presented as: capsules, sachets or tablets each containing a predetermined amount of the active agent; as a powder or granules; as a solution or a suspension of the active agent in an aqueous liquid or a nonaqueous liquid; or as an oil-in-water liquid emulsion or a water in oil liquid emulsion; or as a bolus etc.

For compositions for oral administration (e.g. tablets and capsules), the term "acceptable carrier" includes vehicles such as common excipients e.g. binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, polyvinylpyrrolidone (Povidone), methylcellulose, ethylcellulose, sodium carboxymethylcellulose, hydroxypropylmethylcellulose, sucrose and starch; fillers and carriers, for example corn starch, gelatin, lactose, sucrose, microcrystalline cellulose, kaolin, mannitol, dicalcium phosphate, sodium chloride and alginic acid; and lubricants such as magnesium stearate, sodium stearate and other metallic stearates, glycerol stearate stearic acid, silicone fluid, talc waxes, oils and colloidal silica. Flavouring agents such as peppermint, oil of wintergreen, cherry flavouring and the like can also be used. It may be desirable to add a colouring agent to make the dosage form readily identifiable. Tablets may also be coated by methods well known in the art.

Additionally, or alternatively, the further aspect of the invention also, or alternatively, comprises a novel method for treating a mammalian wound, typically a chronic wound, which method comprises administering to said wound

- an agent that inhibits the activity of at least one gene selected from the group consisting of MEOX2, MAF, SIX2 and homologues thereof having at least 50% identity with said genes and/or

- an agent that enhances the activity of at least one gene selected from the group consisting of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, FOXS1, GLI1, SOX9, preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 and homologues thereof having at least 50% identity with said gene

The invention also comprises a novel method for treating a mammalian wound, typically a chronic wound, which method further comprises administering to said wound

- an agent that inhibits the activity of PPARG or homologues thereof having at least 50% identity with PPARG

and/or

- an agent that enhances the activity of at least one gene selected from the group consisting of SMAD3, SMAD4 and homologues thereof having at least 50% identity with said genes.

According to yet a further aspect of the invention there is provided a kit for treating a wound, preferably a chronic wound, wherein said kit comprises:

(a) at least one therapeutic as above described; and

(b) at least one medical device for applying to said wound

The term “medical device” includes an instrument, apparatus, implant, in vitro reagent, or similar or related article that is used to diagnose, prevent, or treat disease or other conditions, and does not achieve its purposes through chemical action within or on the body (which would make it a drug). Whereas medicinal products (also called pharmaceuticals) achieve their principal action by pharmacological, metabolic or immunological means, medical devices act by other means like physical, mechanical, or thermal means.

According to a yet further aspect of the invention there is provided a combination therapeutic for treating a wound comprising an inhibitor of MEOX2, MAF, SIX2 or enhancer of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, SOX9, GLI1, FOXS1 preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 gene expression and an inhibitor of MEOX2, MAF, SIX2 or enhancer of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, SOX9, GLI1, FOXS1, preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 protein activity.

The invention also comprises a combination therapeutic for treating a wound, which combination therapeutic further comprises an inhibitor of PPARG gene expression or enhancer of SMAD3, SMAD4 gene expression and an inhibitor of PPARG or enhancer of SMAD3, SMAD4 protein activity.

According to a further aspect of the invention there is provided a therapeutic for treating a wound comprising an inhibitor of MEOX2, MAF, PPARG, SIX2 protein, or a homologue thereof or an enhancer of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SMAD3, SMAD4,

SRF, STAT4, TCF4, SOX9, GLI1, FOXS1, preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 protein, or a homologue thereof.

The invention also comprises a therapeutic for treating a wound further comprising an inhibitor of PPARG protein, or a homologue thereof or an enhancer of SMAD3, SMAD4 protein, or a homologue thereof.

According to a further aspect of the invention there is provided use of an inhibitor of MEOX2, MAF, SIX2 protein, or a homologue thereof, in the manufacture of a medicament for treating a wound. According to a further aspect of the invention there is provided use of an inhibitor of MEOX2, MAF, SIX2, or a homologue thereof, for treating a wound.

The invention further provides use of an inhibitor of PPARG protein, or a homologue thereof, in the manufacture of a medicament for treating a wound. According to a further aspect of the invention there is provided use of an inhibitor of PPARG or a homologue thereof, for treating a wound.

The term "homologue" as used herein refers to amino acid sequences which have a sequence at least 50% homologous to the amino acid sequence of MEOX2, MAF, PPARG, SIX2 and which retain the biological activity of the MEOX2, MAF, PPARG, SIX2 sequence. It is preferred that homologues are at least 75% homologous to the MEOX2,ETS1, MAF, PPARG, SIX2 peptide sequence and, in increasing order of preference, at least 80%, 85%, 90%, 95% or 99% homologous to the MEOX2, MAF, PPARG, SIX2 peptide sequence.

According to a further aspect of the invention there is provided use of an enhancer of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, SOX9, GLI1 , FOXS1, preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 protein, or a homologue thereof, in the manufacture of a medicament for treating a wound.

According to a further aspect of the invention there is provided use of an enhancer of SMAD3, SMAD4, or a homologue thereof, for treating a wound.

The term "homologue" as used herein refers to amino acid sequences which have a sequence at least 50% homologous to the amino acid sequence of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SMAD3, SMAD4, SRF, STAT4, TCF4, SOX9, GLI1, FOXS1 and which retain the biological activity of the CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SMAD3,

SMAD4, SRF, STAT4, TCF4, SOX9, GLI1, FOXS1 sequence. It is preferred that homologues are at least 75% homologous to the CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SMAD3, SMAD4, SRF, STAT4, TCF4, SOX9, GLI1, FOXS1 peptide sequence and, in increasing order of preference, at least 80%, 85%, 90%, 95% or 99% homologous to the
5 CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SMAD3, SMAD4, SRF, STAT4, TCF4 SOX9 GLI1, FOXS1 peptide sequence.

Treatment of a wound described herein includes reference to human or veterinary use.

In the claims which follow and in the preceding description of the invention, except where the context requires otherwise due to express language or necessary implication, the word
10 "comprises", or variations such as "comprises" or "comprising" is used in an inclusive sense i.e. to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.

Preferred features of each aspect of the invention may be as described in connection with any of the other aspects.

15 Other features of the present invention will become apparent from the following examples. Generally speaking, the invention extends to any novel one, or any novel combination, of the features disclosed in this specification (including the accompanying claims and drawings). Thus, features, integers, characteristics, compounds or chemical moieties described in conjunction with a particular aspect, embodiment or example of the invention are to be
20 understood to be applicable to any other aspect, embodiment or example described herein, unless incompatible therewith.

Moreover, unless stated otherwise, any feature disclosed herein may be replaced by an alternative feature serving the same or a similar purpose.

The full identity of the genes according to the invention is available on the NCBI database
25 (<http://www.ncbi.nlm.nih.gov/>), or is well known to those skilled in the art.

FIGURE LEGENDS

The legends of the figures are the following:

Figure 1a: Schematic representation of the different treatments applied to the Normal Human Dermal Fibroblast (NHDF)

Figure 1b: Graphic representation of the α SMA mRNA levels as assessed by RT-qPCR with the different treatments

Figure 2a: Graphic representation of the α SMA mRNA levels as assessed by RT-qPCR. NHDFs were treated either with mock siRNA or siRNA directed against different mRNA (EGR2, SRF, HIC1, STAT4, TCF4, GLI1, JUN, IRF7, E2F1, MYC, CREB5, FOXS1, SOX9 or STAT1) and concomitantly subjected or not to TGF β 1 treatment. The RTqPCR were normalized with TUBB and the mock siRNA treated (T-E-) condition was set to one. The treatments of NHDFs with siRNA against SRF, HIC1 or STAT4 lead to extensive cell death (*): no analysis was possible.

Figure 2b: Graphic representation of the percentage of differentiated cells as assessed by the percentage of alpha SMA positive cells after treatment of NHDFs as described in a). The treatments of NHDFs with siRNA against SRF, HIC1 or STAT4 lead to extensive cell death (*): no analysis was possible.

Figure 2c: Graphic representation of the α SMA mRNA levels as assessed by RT-qPCR. Primary human dermal fibroblasts were treated either with mock siRNA or siRNA directed against different mRNA (PPARG, MAF, MEOX2, SIX2, STAT1 or USF2). The RTqPCR were normalized with TUBB and the mock siRNA treated (T-E-) condition was set to 1.

Figure 2d: Graphic representation of the α SMA mRNA levels as assessed by RT-qPCR. NHDFs were treated either with mock siRNA or siRNA directed against different mRNA (PPARG, MAF, MEOX2, SIX2, STAT1 or USF2) and concomitantly treated either with CWF1 alone or in combination with TGF β 1. The RT-qPCR were normalized with TUBB and to the mock siRNA treated (T-E-) condition (set to 1 but not represented on this graph). To help in analyzing the effect of the different siRNA, the condition where the cells were treated with mock siRNA and TGF β 1 is added and represents the normal differentiation (light grey bar in the mock histogram).

Figure 2e: Graphic representation of the α SMA mRNA levels as assessed by RT-qPCR. NHDFs were treated as depicted in d) except for the exudate treatment which was performed with a different exudate from a different patient suffering from chronic wound (CWF2).

Figure 3: short and long timing after TGF β 1 treatment for EGR2, FOXS1, SOX9, SRF, STAT4, TCF4, MYC, JUN, IRF7, E2F1, CREB5 and GLI1. For each Factor, graphic

representation of the mRNA levels after increasing time of treatment of the NHDF with TGF β as assessed by RT-qPCR.

Figure 4: Key transcription factors in fibroblast to myofibroblast differentiation

a) Graphical explanation of the in silico gene network analysis

- 5 b) Table representing the different transcription factors identified by bioinformatical and network analysis.

Figure 5: short and long timing after TGF β 1 treatment for PPARG, MAF, MEOX2 and SIX2. For each Factor, graphic representation of the mRNA levels after increasing time of treatment of the NHDF with TGF β as assessed by RT-qPCR.

- 10 **Figure 6:** Graphic representation of the TCF4 (a), EGR2 (b), SOX9 (c), STAT4 (d), FOXS1 (e), PPARG (f), MAF (g), MEOX2 (h), and SIX2 (i) mRNA levels after NHDF (donor A) treatment for 48h (light grey) or 72h (dark grey) either with mock siRNA or siRNA directed against different TF mRNA (SOX9, EGR2, TCF4, or FOXS1) and concomitantly treated with TGF- β . For all graphs, the mock siRNA treated with TGF- β (T+E) condition was set to 100%
15 for each time of treatment (48h and 72h).

Table 1: genes to be increased or decreased in order to treat wounds, non- healing or chronic wounds

Table 2: siRNA sequences for each target gene.

EXAMPLE

- 20 In response to a lesion, fibroblasts migrate into the wound where they differentiate into contractile myofibroblasts that will finally enter into apoptosis during the remodeling phase. This differentiation process can be studied ex-vivo in environmentally controlled tissue culture conditions, and therefore the timely controlled succession of different gene expression patterns can be addressed.

25 **Materials and methods**

Establishment of an ex vivo model of chronic wounds

Normal Dermal Fibroblast Cell Culture and exudate collection

NHDF, isolated from human explants, were purchased from Promocell. NHDF were cultivated in DMEM-F12 (Invitrogen), supplemented with 10% FCS (Invitrogen, 5µg/mL of insulin and 1ng/mL of b-FGF (PromoKine)).

To collect exudates, two patients with mixed ulcers were recruited. For patient selection, it was decided to exclude any other comorbidity factor potentially involved in wound etiology: diabetes, peripheral arterial diseases, malnutrition. Exudates were collected from negative pressure therapy. All the exudates were centrifuged at 1,500 x g for 3 minutes to remove cell debris. The supernatant was filtered and stored at -80°C until use. Aliquots were used to determine protein concentration according to BCA method (Sigma).

10 Establishment of an ex vivo model of chronic wounds

For experiments, cells were deprived of insulin and b-FGF during 48 hours. Then, the cells were cultivated on collagen coated culture plates in DMEM-F12, supplemented with 10% FCS, 10ng/mL of TGF-β1 (Promocell) for 4 days. Four points were tested in order to appreciate the effect of exudate on fibroblast differentiation: untreated cells (T-E-), fibroblasts treated with TGF-β1 (T+E-), cells treated with exudate (T-E+) and finally fibroblasts treated with TGF-β1 and exudate at the same time (T+E+).

The efficiency of fibroblast differentiation was estimated by analyzing the expression of the myofibroblast marker alpha smooth muscle actin (αSMA).

Western Blotting assay

Total proteins were extracted by scratching the cells with lysis buffer (TRIS, NaCl, NP40, EDTA, IMDTT) and incubated 30 min in ice. To remove cell debris, the samples were centrifuged at 13,000 x g for 10 min at 4°C and stored at -20°C until use. Protein concentration was determined according to BCA method (Sigma). Equal amounts of total protein (20µg) were loaded to NuPAGE 10% BIS-Tris gel (Invitrogen), separated by migration at 150 V, and transferred to nitrocellulose membrane (Whatman) 1 hour at 30 V. Then, membranes were stained for αSMA (Abcam) and tubulin (Abcam). Incubations were followed by secondary antibodies goat anti-rabbit IgG and goat anti-mouse IgG, respectively, conjugated with horseradish-peroxidase (HRP) (Promega). Signals were detected by ECL chemiluminescence using UptiLight HS WB Substrate (Uptima, Interchim). Bands were digitized with a scanner and the ratio between all bands density of the same blot was

calculated by software (ImageJ 1.43u, 64-bit). Relative α SMA expression was normalized to the respective value for tubulin.

Total RNA Sample Preparation

After four days of experiment, treated fibroblasts were lysed with TRIzol Reagent (Invitrogen) and stored at -80°C . Then RNA was purified using chloroform and precipitated by isopropanol. Total RNA was quantified on the NanoDrop 2000c Spectrophotometer (Thermo Scientific). Reverse transcription of 500 ng total RNA to cDNA was done with oligot dT (Invitrogen) using SuperScript III RT (Invitrogen) and RNase OUT (Invitrogen). The cDNA was store at -20°C .

10 Quantitative real-time RT-PCR

Quantitative real-time PCR (RT-qPCR) was done using 5 μL of 1:20 diluted cDNA on the LightCycler480 system (Roche) using Maxima SYBR Green qPCR Master Mix (Fermentas). Forward and reverse primers for α SMA were designed by Eurofins (MWG, α SMA forward:CTGTTTTCCCATCCATTGTG (SEQ ID NO:9), α SMA reverse: 15 CCATGTTCTATCGGGTACTT (SEQ ID NO:10)) and a 100 μM stock was stored at -20°C . Forward and reverse primer pairs were used for each RT-qPCR reaction. The cycling conditions were as follows : an initial 95°C for 10 minutes, followed by 45 cycles of 95°C for 15 sec, 58°C for 30 sec, 72°C for 20 sec. LightCycler 480 SW 1.5 was used to evaluate the T_{M} curves, to determine the C_{p} and to approximate the relative concentration for each 20 amplification reaction.

siRNA treatment

α -Smooth Muscle Actin Immunofluorescence

Cells grown in collagen coated culture dishes, and treated as previously described, were fixed with 4% paraformaldehyde (PFA) in PBS for 15 minutes and permeabilized with 2.5% Triton X-100 (Euromedex, 2000-B) in PBS for 3 minutes. After saturation with 5% BSA in PBS, 25 cells were stained for α -SMA (Abcam, ab5694) and for DNA (DAPI). As secondary antibody, CyTM3 conjugated anti rabbit (GE Healthcare, PA43004) was used. Samples were observed with an oil immersion objective (Plan Fluor 40X/1.30 Oil, Nikon) on a Nikon ECLIPSE Ti (Nikon). Digital images were taken with a digital camera (Cool SNAP HQ², Photometrics) 30 and software (MetaMorf 7.5.4.0). To estimate the percentage of fibroblast differentiation due

to the different treatments, the total number of cells per field was determined by the DAPI, and myofibroblasts, differentiated fibroblasts, were counted using the α -SMA staining. Then, STUDENT (t-) and χ^2 tests were realized to evaluate the differential expression of α SMA between the untreated fibroblasts (T-E-) and the treated ones.

5 mRNA Expression Analysis

Total RNA Sample Preparation and cDNA Synthesis

After four days of experiment, treated fibroblasts were detached with TRIzol Reagent (Invitrogen (Life Technologies), 15596-018) and stored at -80°C. Then RNA was purified using chloroform and precipitate by isopropanol. Total RNA was quantified on the NanoDrop
 10 2000c Spectrophotometer (Thermo Scientific) and their quality was evaluated on the RNA Nano Chips (Agilent 2100 bioanalyzer, Agilent, 5067-1511). Reverse transcription of 500 ng total RNA to cDNA was done with oligot dT (Invitrogen (Life technologies), 18418-020) using SuperScript III RT (Invitrogen (Life technologies), 18080-085) and RNase OUT (Invitrogen (Life technologies), 10777-019). The cDNA was stored at -20°C. Only samples
 15 with a good bioanalyzer profile were used for qPCR analysis.

Network analysis

In order to enlighten master regulators of fibroblast fate after each different treatment, we performed a gene network analysis treating gene expression lists determined after mRNA seq deep sequencing analysis of the gene profile of T-E- with the gene profile of T+E-, T+E+ and
 20 T-E+ 1. In these analysis and based on the assumption that the decrease or increase of interconnected genes is of stronger significance than a significant Log FC, we have used lists of genes selected only based on their P value and not on the value of their Log FC. We have performed two types of analysis: an ingenuity “upstream regulator analysis” and a DIRE (<http://dire.dcode.org/>) analysis. The Ingenuity “upstream regulator analysis”, given the
 25 particular profile of genes expression between two conditions, consists in selecting potential upstream regulators. The DIRE analysis is based on the selection of potential common regulatory elements between genes based on these elements conservation during evolution. From these identified elements, DIRE is able to provide a list of master regulators for a list of co-regulated genes. From those two analyses, and for each list analyzed, we have selected
 30 Transcription Factors (TFs) expressed in at least one of the two conditions considered in the concerned list (i.e. number of sequencing his superior to twenty in at least one of the two

conditions). Then, we have deeply compared the two sets of analysis and decided to keep in the “key regulators lists” transcription factors belonging to both analyses. Because of possible bias in these two analyses we also decided to rescue transcription factors belonging only to one analysis and not the other but presenting very interesting target genes pattern in one list or the other. Altogether, these genes networks analysis allowed us to propose a list of TFs being key regulators in one or the other fibroblast fate.

For the chronic or non-healing wound model, chronic wound exudates were added to cell cultures (500µg/mL of total proteins of exudate). The experiments that were performed are depicted in Figure 1a): cells were either not treated (T-E-), either treated with TGFβ alone (T+E-), exudate alone (T-E+) or TGF-β and exudate (T+E+) for 4 days. The assays described previously were used to assess the level of differentiation. Chronic wound exudates decrease the expression of αSMA (mRNA, figure 1b). This indicated that chronic wound exudates clearly inhibit fibroblast differentiation. This is correlated to the fact that in chronic wounds, non-functional fibroblasts, also called pseudo-senescent fibroblasts, are present.

Gene expression route upon fibroblast to myofibroblast differentiation

Identification of the main molecular targets implicated in fibroblast differentiation of human primary fibroblasts under normal and pathological conditions

We performed an in silico gene network analysis to enlighten putative upstream regulators of the different gene expression routes defined previously. This approach was original in the sense that we used global gene network analysis to identify potential key regulators and we did not take into account a change in these factors expression to select them. For example, we used the DIRE program (<http://dire.dcode.org/>) to identify evolutionary conserved potential regulatory elements in the different genes lists which allowed us to enlighten transcription factors that could potentially bind to these elements and thus regulate these sets of genes.

Twenty-three transcription factors were selected out from this analysis.

To prioritize the extensive study of the different Transcription Factors (TFs), we performed a time response study of TFs after the different fibroblast treatments. We did a short-term (between 30mn and 8 hours) and a long-term (between 8 hours and 96 hours) analysis of their changes in expression after the different treatments.

We performed an exhaustive siRNA-based approach to study on one hand the role of these different factors in normal fibroblast to myofibroblast differentiation pathway and on the other hand, in the chronic-exudate-dependent non-differentiation cell fate.

5 The siRNA knock-down of fourteen of the potential key transcription factors identified therein inhibited the fibroblast to myofibroblast differentiation pathway as assessed by analyzing the α SMA expression from TGF β - and siRNA-treated NHDFs :GLI1, HIC1, TCF4, SOX9, STAT4 MYC, CREB5, IRF7, JUN,E2F1, EGR2, SRF, SMAD3, FOXS1 as their knockdown inhibitor myofibroblast differentiation to various extents (Fig. 2a-b). Very interestingly, except for SOX9, FOXS1 and EGR2, whose expression is strongly and rapidly
10 up-regulated upon TGF β treatment, the mRNA levels of the other factors is constant during the first day or so after TGF β treatment and overall unchanged during the four days of differentiation (Fig 3). This indicates that the maintenance of their expression but not their over-expression is necessary for fibroblast to myofibroblast differentiation.

The siRNA knock-down of four other potential key transcription factors identified by the in
15 silico analysis (MAF, SIX2, MEOX2 and PPARG) seemed to induce the fibroblast to myofibroblast differentiation in absence of TGF β to the same extend as the one obtained with mock transfected cells treated with TGF β . Even more interestingly the induction of myofibroblast differentiation was old true even in presence of chronic wound exudates and was very similar to the differentiation obtained in presence of TGF β -only for mock
20 transfected cells. In absence of these factors, the fibroblasts seemed able to bypass the exudate dominant action in order to differentiate into myofibroblasts. We can also enlighten that, except for FOXS1, all these factors are slightly too strongly down regulated upon TGF β treatment (Fig. 5) supporting that their down regulation could be necessary for differentiation.

Altogether these results showed that with knocking-down approaches we were able either to
25 reduce fibroblast to myofibroblast differentiation or to induce this differentiation even in a chronic wound context (table 1).

An in silico gene network analysis allowed us to identify potential key regulators of fibroblast cell fate either during differentiation into myofibroblast or in a context of chronic wounds. By
30 knocking down approaches, we found a strong effect on differentiation for nineteen factors. The strength of our analysis comes from the unbiased approach to look for evolutionary

conserved common DNA element in the different lists of genes (via the DIRE program) and from there to identify putative upstream transcription regulators, rather than to look at differences in the expression levels of the transcription factors. It is very likely that we would have missed those cell fate regulators by looking at their differential expression only.

- 5 By knocking-down approaches we identified transcription factors required for normal fibroblast to myofibroblast differentiation (Fig.2). We used the α SMA to follow myofibroblast differentiation as it is their best ex-vivo marker.

10 We have also identified factors which seemed to play a role but maybe not as strongly as the ones described in the paragraph before as their knockdown leads to consistent but mild decrease of α SMA expression. These factors are MYC, JUN, E2F1, IRF7 and CREB5.

Very interestingly, we showed that the inactivation of some transcription factors leads to an increase of fibroblast differentiation per se. The invalidation of some of these factors could even increase fibroblast to myofibroblast differentiation in a chronic wound context.

- 15 The knocking down of PPARG mRNA leads to an increase of fibroblast to myofibroblast differentiation (Fig.2c-d) even in the presence of chronic exudate.

FOXS1 belongs to the forkhead family of transcription factor often involved in developmental processes such as morphogenesis and differentiation. It has been shown that FOXS1 is of primary importance in the development of testicular vasculature. Moreover, FOXS1 was
20 described as an early sensory neuronal marker. Here we show that inactivation of FOXS1 leads to an increase of myofibroblast differentiation in absence of TGF β .

MEOX2 has already been described as implicated in TGF β pathway as it was identified as an important factor in cleft palate development in TGF β 3 knockout mice. Experiments in C2C12 myoblast cells showed that MEOX2 is also important for skeletal muscle development and
25 differentiation. Here, we showed that siRNA directed against MEOX2 lead to a bypass of the exudate effect by fibroblasts to be able to differentiate into myofibroblasts.

In T cells, it has been shown that MAF was responsible for inhibition of IL22 expression by neutralizing TGF β . TGF β and MAF have antagonist/opposite effects on IL21 expression in CD4(+) T cells. In the same connection, in this study, we implicated MAF as an inhibitor of
30 fibroblast to myofibroblast differentiation in absence or presence of exudate as its inactivation

by siRNA leads to an increase of myofibroblast differentiation (Fig.2c-d). On the contrary, during chondrocyte differentiation, a long form of MAF interacts and cooperates with SOX9 to activate downstream targets. This is another example of the differences between myofibroblast and chondrocyte differentiations.

- 5 SIX2 has been involved in maintaining pluripotency in kidney: in embryonal renal mesenchyme cells it is able to suppress differentiation and during kidney development it maintains the progenitor pool. Here, in dermal fibroblast, the invalidation of SIX2 leads to a bypass of the dominant exudate effect on TGF β signaling (Fig.2c-d).

Regulatory interactions between key transcription factors during fibroblast to myofibroblast differentiation

- 10 The expression of TCF4 mRNA was not modified after SOX9, EGR2 or FOXS1 knockdown (Fig 6) placing it at the top of the regulatory interaction network between these factors. On the contrary EGR2 expression was largely inhibited upon siRNA treatment against TCF4 and SOX9 but remained unchanged upon treatment against FOXS1 (Fig 6) placing it after TCF4 and SOX9 but before FOXS1 in the network. SOX9 mRNA remained largely unchanged upon TCF4 and FOXS1 knockdown. We placed FOXS1 as a downstream target of SOX9. On the contrary SOX9 is upregulated upon EGR2 knockdown (Fig 6) The STAT4 mRNA is downregulated by TCF4, SOX9 and EGR2 knockdown (Fig 6) placing is as a downstream target of those TF whereas it remained unchanged upon FOXS1 siRNA treatment placing it beforehand. Consistently, FOXS1 mRNA is downregulated upon TCF4, SOX9 and EGR2 siRNA treatment (Fig 6) placing it at the end of this cascade. Globally, the PPARG, MAF and MEOX2 mRNA were upregulated upon siRNA treatment against TCF4, SOX9, EGR2 and FOXS1 (Fig 6) consistent with their role as antagonist in fibroblast differentiation. SIX2 mRNA level was unchanged upon EGR2 and FOXS1 knockdown (Fig 6) but upregulated in the same manner by TCF4 and SOX9 consistent with a close interconnexion between these two TF and suggesting the existence of a balanced signal between the differentiation agonists SOX9 and TCF4 and the antagonist SIX2.

- 30 The identification of transcription factors able to bypass chronic wound exudates effect is of major importance in the chronic wound field as it gives new insight into targets that can be used in the treatment of chronic wounds such as leg ulcers. In this study, by focusing on

fibroblast, by no mean we tried to dissimulate the importance of other cells like neutrophils and macrophages in the skin healing process but we willingly simplified the biological context to draw a clearer picture of the situation.

Claims

1. A therapeutic compound comprising :
- an agent that inhibits the activity of at least one gene selected from the group consisting of
5 MEOX2, MAF, SIX2, preferentially MEOX2, MAF, SIX2 and homologues thereof having at least 50% identity with said genes
and/or
 - an agent that enhances the activity of at least one gene selected from the group consisting of
10 CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, FOXS1, GLI1, SOX9, preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 and homologues thereof having at least 50% identity with said gene
for use in the treatment of wounds, preferably chronic wounds.
2. A therapeutic compound according to claim 1 further comprising:
- 15 - an agent that inhibits the activity of PPARG or homologues thereof having at least 50% identity with PPARG
and/or
 - an agent that enhances the activity of at least one gene selected from the group consisting of SMAD3, SMAD4 and homologues thereof having at least 50% identity with said genes.
- 20 3. A therapeutic compound according to claim 1 or 2 wherein said agent is selected from the group consisting of: anti-sense DNA or RNA, siRNA, shRNA, cDNA, TALENS or ribozymes, either naked or in the form of plasmid or viral vectors.
4. A therapeutic compound according to claim 1 or 2 wherein said agent is an inhibitor or enhancer of protein function.
- 25 5. A therapeutic compound according to claim 4 wherein said agent is selected from the group consisting of: a binding agent that binds, either reversibly or irreversibly, to inhibit protein function such as an antibody or a synthetic antagonist; or an agent that works upstream or downstream of the protein signaling mechanism to inhibit protein function.
6. A therapeutic compound according to any preceding claim wherein the therapeutic
30 compound is for use in treating mammalian wounds.

7. A therapeutic compound according to any preceding claim wherein the therapeutic compound is for use in treating chronic wounds such as venous ulcers, diabetic ulcers, or pressure ulcers.
8. A therapeutic compound according to any preceding claim wherein the therapeutic is for use in treating human wounds.
9. A therapeutic compound according to any preceding claim wherein the therapeutic compound is for topical application.
10. A therapeutic compound according to any preceding claim wherein the therapeutic compound is for application to a medical device or impregnation of a medical device.
11. A pharmaceutical composition comprising a therapeutic compound according to any preceding claim together with a pharmaceutically acceptable carrier.
12. A method for preparing a pharmaceutical composition according to claim 11 comprising bringing said therapeutic compound in conjunction or association with a pharmaceutically or veterinary acceptable carrier or vehicle.
13. A method for treating a mammalian wound, wherein said method comprises: administering to said wound a therapeutic compound comprising:
- an agent that inhibits the activity of at least one gene selected from the group consisting of MEOX2, MAF, SIX2 and homologues thereof having at least 50% identity with said genes and/or
 - an agent that enhances the activity of at least one gene selected from the group consisting of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, FOXS1, GLI1, SOX9 , preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 and homologues thereof having at least 50% identity with said gene
14. A kit for treating a wound, preferably a chronic wound, wherein said kit comprises:
- (a) at least one therapeutic compound according to any of claims 1 -10 or a composition according to claim 11; and
 - (b) at least one dressing for applying to said wound.
15. A combination therapeutic for treating a wound, preferably a chronic wound, comprising:

a) an agent that inhibits the activity of at least one gene selected from the group consisting of MEOX2, MAF, SIX2 and homologues thereof having at least 50% identity with said genes and/or

5 - an agent that enhances the activity of at least one gene selected from the group consisting of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, FOXS1, GLI1, SOX9, preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 and homologues thereof having at least 50% identity with said gene

and

b) at least one further therapeutic.

10 16. Use of an

an agent that inhibits the activity of at least one gene selected from the group consisting of MEOX2, MAF, SIX2 and homologues thereof having at least 50% identity with said genes and/or


15 - an agent that enhances the activity of at least one gene selected from the group consisting of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, FOXS1, GLI1, SOX9, preferentially FOXS1, EGR2, SOX9, TCF4, STAT4 and homologues thereof having at least 50% identity with said gene

for treating a wound,

wherein said agent modulates fibroblast and myofibroblast differentiation and/or activity.

20

a)



b)






Figure 1



Figures 2a) and b)



Figures 2c) 2d) and 2e)





Figure 3

5/9

a)



b)

Symbol	Entrez Gene Name
CREB5	cAMP responsive element binding protein 5
E2F1	E2F transcription factor 1
EGR2	early growth response 2
ETV1	v-ets erythroblastosis virus E26 oncogene homolog 1 (avian)
FOXS1	forkhead box S1
GLI1	GLI family zinc finger 1
HC1	hypermethylated in cancer 1
IRF7	interferon regulatory factor 7
JUN	jun oncogene
MAF	v-maf musculoaponeurotic fibrosarcoma oncogene homolog (avian)
MEOX2	mesenchyme homeobox 2
MYC	v-myc myelocytomatosis viral oncogene homolog (avian)
POU2F2	POU class 2 homeobox 2
PPARG	peroxisome proliferator-activated receptor gamma
SIX2	SIX homeobox 2
SMAD3	SMAD family member 3
SMAD4	SMAD family member 4
SOX9	SRY (sex determining region Y)-box 9
SRF	serum response factor (c-fos serum response element-binding transcription factor)
STAT1	signal transducer and activator of transcription 1 91kDa
STAT4	signal transducer and activator of transcription 4
TCF4	transcription factor 4
USF2	upstream transcription factor 2 c-fos interacting

Figure 4




Figure 5




Figure 6

Symbol	Wound
CREB5	To increase
E2F1	To increase
EGR2	To increase
FOXS1	To increase
GLI1	To increase
HIC1	To increase
IRF7	To increase
JUN	To increase
MAF	To decrease
MEOX2	To decrease
MYC	To increase
PPARG	To decrease
SIX2	To decrease
SOX9	To increase
SMAD3	To increase
SMAD4	To increase
SRF	To increase
STAT4	To increase
TCF4	To increase

Table 1

CREB5	CTGGCGAAGGGUGGAGACGA	Q. SIB4157174	11
	AACAGUAIUUCUGUAGGALICUA	Q. SIB4141893	12
E2F1	UCGGAGAAUUCUAGAUU		13
	GAGAAGUACCGCUAUGAGA	T. L-003259-00	14
	GAGCAGAUUGUUAUGUGA		15
	GAACAGGGCCACUAGACUCU		16
EGR2	GAAGGCALAAUCAUAUUG		17
	CUACUGUGGCCCGAAAGUUI	T. L-006527-00	18
	GAACCCAGACCUUCACUUA		19
	GAGAAGAGUUCGUUGGAUC		20
FOX51	AGGCGCAAUAAAGCCAUUGUA	Q. SIB4152540	21
	CUAGCUUAGGACCCUGAAGAA	Q. SIB0387527	22
GLI1	GCAAAUAGGECUUCACAU		23
	AGGCUACAGCUUGUGUGUAA	T. L-003836-00	24
	GGACGAGGGACCUUGCAU		25
	CAGCUAGAGUCCAGAGGUU		26
HIC1	GCACAGCAACGCCAACUCA		27
	GAGCUUCGGUGACAAACCU		28
	UGAUCAUUGUGUGUCAGAA		29
	GACCAUCGACCGUUCUCU	T. L-006532-00	30
IRF7	GCACAAGGUGUACGCGCUC		31
	CAACUUCGCGUGGCGACUG	T. L-011810-00	32
	CGCGGCAUUCUACAAGGCCU		33
	CAGGCAAGGACACAGACUA		34
JUN	GAGCGGACCUUAUGGCUAC		35
	GAACAGGUGGCCACAGCUA	T. L-003268-00	36
	GAACAGACCUUCUAUAGCG		37
	UGAAAGCUCAGAAUUGGGA		38
MAF	UUCGAUCUGAUGAAGUUGAA	Q. SIB0076049	1
	CGCGAGGAGAAUUGCAUCAGA	Q. SIB03086063	2
MEOX2	AGCAUCCGCAUUAUUAUUA	Q. SIB4299421	3
	CCCGCCGUGUCUGGCUCAA	Q. SIB4262174	4
MYC	AGGGAACUCUUGUGGUA		39
	GAACACACAACGUCUUGGA	T. L-003282-00	40
	AACGUUAGCUUCACCAACA		41
	CGAUGUUGUUCUUGUGGAA		42
PPAR γ	UCGGGAGAAACAUCAGAUUA	Q. SIB3115266	5
	GAGGCGCAUCUUGACAGGAAA	Q. SIB0071673	6
SIX2	CAACGAGAACUCCAAUUCUAA	Q. SIB0719313	7
	CCCGCCUGAAUGGACGCGCAA	Q. SIB4141095	8
SMAD3	CAACAGGAAUGCAACAGUG		43
	GAGUUCGCCUUCAAUUA	T. L-020067-00	44
	GGACGGAGGUUCUCCAAC		45
	UUAGAGACAUCAGUAUGG		46
SMAD4	GCAAUUGAAAGUUGGUAA		47
	CCGACAAACUUAAGACUGA	T. L-003902-00	48
	GAUCCAUUAUCUACGAA		49
	GUACAGAGUUAUACUAG		50
SOX9	GGAAACAACCGUCUACACA		51
	GAACAAGCCCGACGUCAAG	T. L-021507-00	52
	GACCUUCGAUGUACACGAG		53
	GGAAGUCGUGAAGAAACGG		54
SRF	UAGAGACGGCCAUUGUAU		55
	GGACUGUGCUGAAGAGUAC	T. L-009800-00	56
	GCACCAAGAGUGAAGAUUC		57
	GCACCAUGUCUGCUAGUG		58
STAT1	GCACGAGGGGCUAGCUU		59
	CUACGAACAUGACCCUAUC	T. L-003543-00	60
	GAACCUAGACUUCACGCG		61
	AGAAAGAGCUUGACAGUAA		62
STAT4	GAACUAAACUAUCAGGUAA		63
	GCAUGUAGCUUGGUUAU	T. L-011784-00	64
	CAAUUAGCUACAGUGAUG		65
	CUCCGAGACUACAAGUUA		66
TCF4	GCACUUGGCUUGAUUUAU		67
	GACAAACAGCGAGUGAUU	T. L-004594-00	68
	GCACAGCUGUUGGUUAG		69
	CAACGGACAGACAGUUA		70
USF2	GCAAGCGGGAGCGAGUAA		71
	GGAGCGGACAAGAUCAACAA	T. L-003618-00	72
	GAAGAGCCAGCACAACGA		73
	CAAAAUCCUUCAGCAUUG		74

Table 2

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2013/001990

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07K14/47 C12N15/113
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MARTIN P ET AL: "Growth factors and cutaneous wound repair", PROGRESS IN GROWTH FACTOR RESEARCH, PERGAMON PRESS, GB, vol. 4, no. 1, 1 January 1992 (1992-01-01) , pages 25-44, XP026324179, ISSN: 0955-2235, DOI: 10.1016/0955-2235(92)90003-Z [retrieved on 1992-01-01] page 34 second full paragraph -page 36 first full paragraph. ----- -/--	1-16



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

8 April 2014

Date of mailing of the international search report

08/07/2014

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Kools, Patrick

INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2013/001990

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>P. LIU ET AL: "Cross Talk Among Smad, MAPK, and Integrin Signaling Pathways Enhances Adventitial Fibroblast Functions Activated by Transforming Growth Factor- 1 and Inhibited by Gax", ARTERIOSCLEROSIS, THROMBOSIS, AND VASCULAR BIOLOGY, vol. 28, no. 4, 1 April 2008 (2008-04-01), pages 725-731, XP055111941, ISSN: 1079-5642, DOI: 10.1161/ATVBAHA.107.159889 the whole document</p>	1-16
A	<p>DANIEL NOWINSKI ET AL: "Analysis of Gene Expression in Fibroblasts in Response to Keratinocyte-Derived Factors In Vitro: Potential Implications for the Wound Healing Process1", JOURNAL OF INVESTIGATIVE DERMATOLOGY, vol. 122, no. 1, 1 January 2004 (2004-01-01), pages 216-221, XP055111564, ISSN: 0022-202X, DOI: 10.1046/j.0022-202X.2003.22112.x the whole document</p>	1-16
A	<p>WO 2012/120269 A1 (UNIV CARDIFF [GB]; JIANG WENGUO [GB]; HARDING KEITH [GB]) 13 September 2012 (2012-09-13) whole document, especially the claims.</p>	1-16
A	<p>DUSTIN M BERMUDEZ ET AL: "Inhibition of stromal cell-derived factor-1 further impairs diabetic wound healing", JOURNAL OF VASCULAR SURGERY, C.V. MOSBY CO., ST. LOUIS, MO, US, vol. 53, no. 3, 5 October 2010 (2010-10-05), pages 774-784, XP028147676, ISSN: 0741-5214, DOI: 10.1016/J.JVS.2010.10.056 [retrieved on 2010-10-22] abstract</p>	1-16
A	<p>DAVID H GORSKI ET AL: "Inhibition of endothelial cell activation by the homeobox gene Gax", JOURNAL OF SURGICAL RESEARCH, vol. 111, no. 1, 1 May 2003 (2003-05-01), pages 91-99, XP055111506, ISSN: 0022-4804, DOI: 10.1016/S0022-4804(03)00042-8 the whole document</p>	1-16

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB2013/001990

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-16(partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-16(partially)

Therapeutic compound comprising an agent inhibiting the activity of ME0X2 or a gene having 50% identity to said gene for use in the treatment of wounds. Pharmaceutical composition comprising said therapeutic compound. Method for preparing the pharmaceutical composition. Method for treating a mammalian wound by administration of the therapeutic compound. Kit for treating a wound wherein the kit comprises the therapeutic compound. A combination therapeutic for treating a wound comprising the therapeutic compound and a further therapeutic. Use of the therapeutic for treating a wound, wherein said therapeutic modulates fibroblast and myofibroblast differentiation and/or activity.

2. claims: 1-16(partially)

As for subject 1 now for agents inhibiting the expression of MAF.

3. claims: 1-16(partially)

As for subject 1 now for agents inhibiting the expression of SIX2.

4-16. claims: 1-16(partially)

As for subject 1 now for agents increasing the expression of CREB5, E2F1, EGR2, HIC1, IRF7, JUN, MYC, SRF, STAT4, TCF4, FOXS1, GLI1, or SOX9, and homologues thereof having at least 50% identity with said gene.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2013/001990

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2012120269 A1	13-09-2012	AU 2012226593 A1	12-09-2013
		CA 2829010 A1	13-09-2012
		CN 103443276 A	11-12-2013
		EP 2683823 A1	15-01-2014
		GB 2502017 A	13-11-2013
		JP 2014511392 A	15-05-2014
		WO 2012120269 A1	13-09-2012
