

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
30 September 2004 (30.09.2004)

PCT

(10) International Publication Number
WO 2004/083447 A2

(51) International Patent Classification⁷: **C12Q**
(21) International Application Number:
PCT/US2004/007634
(22) International Filing Date: 12 March 2004 (12.03.2004)
(25) Filing Language: English
(26) Publication Language: English
(30) Priority Data:
60/454,436 13 March 2003 (13.03.2003) US

(74) Agents: SHAYESTEH, Laleh et al.; Exelixis, Inc., 170 Harbor Way, P.O. Box 511, South San Francisco, CA 94083-0511 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(71) Applicant (*for all designated States except US*): EX-ELIXIS, INC. [US/US]; 170 Harbor Way, P.O. Box 511, South San Francisco, CA 94083-0511 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): COSTA, Michael, R. [US/US]; 18 Hazelwood Avenue, San Francisco, CA 94112 (US). GENDREAU, Steven, Brian [US/US]; 2801 Turk Street, #103, San Francisco, CA 94118 (US). DORA, Emery, G., III [US/US]; 1847 32nd Avenue, San Francisco, CA 94122 (US). KADYK, Lisa, C. [US/US]; 431 Mangels Avenue, San Francisco, CA 94127 (US). HUBER, Jennifer, Lynn [US/US]; 477 Reina Del Mar, Pacifica, CA 94044 (US). LICKTEIG, Kim [US/US]; 1921 Grove Street, San Francisco, CA 94117 (US).

(54) Title: MBCATS AS MODIFIERS OF THE BETA-CATENIN PATHWAY AND METHODS OF USE

(57) Abstract: Human MBCAT genes are identified as modulators of the beta-catenin pathway, and thus are therapeutic targets for disorders associated with defective beta-catenin function. Methods for identifying modulators of beta-catenin, comprising screening for agents that modulate the activity of MBCAT are provided.

WO 2004/083447 A2

MBCATS AS MODIFIERS OF THE BETA-CATENIN PATHWAY AND METHODS OF USE

REFERENCE TO RELATED APPLICATIONS

5 This application claims priority to U.S. provisional patent application 60/454,436 filed 3/13/2003. The contents of the prior application are hereby incorporated in their entirety.

BACKGROUND OF THE INVENTION

10 Beta-catenin is an adherens junction protein. Adherens junctions (AJs; also called the zonula adherens) are critical for the establishment and maintenance of epithelial layers, such as those lining organ surfaces. AJs mediate adhesion between cells, communicate a signal that neighboring cells are present, and anchor the actin cytoskeleton. In serving these roles, AJs regulate normal cell growth and behavior. At several stages of
15 embryogenesis, wound healing, and tumor cell metastasis, cells form and leave epithelia. This process, which involves the disruption and reestablishment of epithelial cell-cell contacts, may be regulated by the disassembly and assembly of AJs. AJs may also function in the transmission of the 'contact inhibition' signal, which instructs cells to stop dividing once an epithelial sheet is complete.

20 The AJ is a multiprotein complex assembled around calcium-regulated cell adhesion molecules called cadherins (Peifer, M.(1993) Science 262: 1667-1668). Cadherins are transmembrane proteins: the extracellular domain mediates homotypic adhesion with cadherins on neighboring cells, and the intracellular domain interacts with cytoplasmic proteins that transmit the adhesion signal and anchor the AJ to the actin
25 cytoskeleton. These cytoplasmic proteins include the alpha-, beta-, and gamma-catenins. The beta-catenin protein shares 70% amino acid identity with both plakoglobin, which is found in desmosomes (another type of intracellular junction), and the product of the Drosophila segment polarity gene 'armadillo'. Armadillo is part of a multiprotein AJ complex in Drosophila that also includes some homologs of alpha-catenin and cadherin,
30 and genetic studies indicate that it is required for cell adhesion and cytoskeletal integrity.

 Beta-catenin, in addition to its role as a cell adhesion component, also functions as a transcriptional co-activator in the Wnt signaling pathway through its interactions with the family of Tcf and Lef transcription factors (for a review see Polakis, (1999) Current Opinion in Genetics & Development, 9:15–21 and Gat U., et al., (1998) Cell 95:605-614).

The APC gene, which is mutant in adenomatous polyposis of the colon, is a negative regulator of beta-catenin signaling (Korinek, V. et al., (1997) Science 275: 1784-1787; Morin, P. J., et al., (1997) Science 275: 1787-1790). The APC protein normally binds to beta-catenin and, in combination with other proteins (including glycogen synthase kinase-3b and axin, is required for the efficient degradation of b-catenin. The regulation of beta-catenin is critical to the tumor suppressive effect of APC and that this regulation can be circumvented by mutations in either APC or beta-catenin.

While mammals contain only a single beta-catenin gene, *C. elegans* contains three (Korswagen HC, et al., (2000) Nature 406:527-32). Each worm beta-catenin appears to carry out unique functions (Korswagen HC, et al., (2000) Nature 406:527-32, Nartarajan L et al. (2001) Genetics 159: 159-72). Because of the divergence of function in *C. elegans*, it is possible to specifically study beta-catenin role in cell adhesion, which is mediated by the *C. elegans* beta-catenin HMP-2.

The ability to manipulate the genomes of model organisms such as *C. elegans* provides a powerful means to analyze biochemical processes that, due to significant evolutionary conservation, have direct relevance to more complex vertebrate organisms. Due to a high level of gene and pathway conservation, the strong similarity of cellular processes, and the functional conservation of genes between these model organisms and mammals, identification of the involvement of novel genes in particular pathways and their functions in such model organisms can directly contribute to the understanding of the correlative pathways and methods of modulating them in mammals (see, for example, Dulubova I, et al, J Neurochem 2001 Apr;77(1):229-38; Cai T, et al., Diabetologia 2001 Jan;44(1):81-8; Pasquinelli AE, et al., Nature. 2000 Nov 2;408(6808):37-8; Ivanov IP, et al., EMBO J 2000 Apr 17;19(8):1907-17; Vajo Z et al., Mamm Genome 1999 Oct;10(10):1000-4). For example, a genetic screen can be carried out in an invertebrate model organism having underexpression (e.g. knockout) or overexpression of a gene (referred to as a "genetic entry point") that yields a visible phenotype. Additional genes are mutated in a random or targeted manner. When a gene mutation changes the original phenotype caused by the mutation in the genetic entry point, the gene is identified as a "modifier" involved in the same or overlapping pathway as the genetic entry point. When the genetic entry point is an ortholog of a human gene implicated in a disease pathway, such as beta-catenin, modifier genes can be identified that may be attractive candidate targets for novel therapeutics.

All references cited herein, including patents, patent applications, publications, and sequence information in referenced Genbank identifier numbers, are incorporated herein in their entireties.

5

SUMMARY OF THE INVENTION

We have discovered genes that modify the beta-catenin pathway in *C. elegans*, and identified their human orthologs, hereinafter referred to as Modifier of beta catenin (MBCAT). The invention provides methods for utilizing these beta-catenin modifier genes and polypeptides to identify MBCAT-modulating agents that are candidate
10 therapeutic agents that can be used in the treatment of disorders associated with defective or impaired beta-catenin function and/or MBCAT function. Preferred MBCAT-modulating agents specifically bind to MBCAT polypeptides and restore beta-catenin function. Other preferred MBCAT-modulating agents are nucleic acid modulators such as antisense oligomers and RNAi that repress MBCAT gene expression or product activity
15 by, for example, binding to and inhibiting the respective nucleic acid (i.e. DNA or mRNA).

MBCAT modulating agents may be evaluated by any convenient *in vitro* or *in vivo* assay for molecular interaction with an MBCAT polypeptide or nucleic acid. In one embodiment, candidate MBCAT modulating agents are tested with an assay system
20 comprising a MBCAT polypeptide or nucleic acid. Agents that produce a change in the activity of the assay system relative to controls are identified as candidate beta-catenin modulating agents. The assay system may be cell-based or cell-free. MBCAT-modulating agents include MBCAT related proteins (e.g. dominant negative mutants, and biotherapeutics); MBCAT -specific antibodies; MBCAT -specific antisense oligomers and
25 other nucleic acid modulators; and chemical agents that specifically bind to or interact with MBCAT or compete with MBCAT binding partner (e.g. by binding to an MBCAT binding partner). In one specific embodiment, a small molecule modulator is identified using a binding assay. In specific embodiments, the screening assay system is selected from an apoptosis assay, a cell proliferation assay, an angiogenesis assay, and a hypoxic
30 induction assay.

In another embodiment, candidate beta-catenin pathway modulating agents are further tested using a second assay system that detects changes in the beta-catenin pathway, such as angiogenic, apoptotic, or cell proliferation changes produced by the originally identified candidate agent or an agent derived from the original agent. The

second assay system may use cultured cells or non-human animals. In specific embodiments, the secondary assay system uses non-human animals, including animals predetermined to have a disease or disorder implicating the beta-catenin pathway, such as an angiogenic, apoptotic, or cell proliferation disorder (e.g. cancer).

5 The invention further provides methods for modulating the MBCAT function and/or the beta-catenin pathway in a mammalian cell by contacting the mammalian cell with an agent that specifically binds a MBCAT polypeptide or nucleic acid. The agent may be a small molecule modulator, a nucleic acid modulator, or an antibody and may be administered to a mammalian animal predetermined to have a pathology associated with
10 the beta-catenin pathway.

DETAILED DESCRIPTION OF THE INVENTION

Genetic screens were designed to identify modifiers of the beta-catenin pathway in *C. elegans*. A weak allele of beta-catenin was used in our screen (a homozygous viable
15 mutant of beta-catenin, allele qm39). The hmp-2 (qm-39) strain produces larval worms with a highly penetrant lumpy body phenotype in first stage larval worms (L1s). Various specific genes were silenced by RNA inhibition (RNAi). Methods for using RNAi to silence genes in *C. elegans* are known in the art (Fire A, et al., 1998 Nature 391:806-811; Fire, A. Trends Genet. 15, 358-363 (1999); WO9932619). Genes causing altered
20 phenotypes in the worms were identified as modifiers of the beta-catenin pathway. Modifiers were identified followed by the identification of their orthologs. Accordingly, vertebrate orthologs of these modifiers, and preferably the human orthologs, MBCAT genes (i.e., nucleic acids and polypeptides) are attractive drug targets for the treatment of pathologies associated with a defective beta-catenin signaling pathway, such as cancer.
25 Table 1 (Example II) lists the modifiers and their orthologs.

In vitro and in vivo methods of assessing MBCAT function are provided herein. Modulation of the MBCAT or their respective binding partners is useful for understanding the association of the beta-catenin pathway and its members in normal and disease conditions and for developing diagnostics and therapeutic modalities for beta-catenin
30 related pathologies. MBCAT-modulating agents that act by inhibiting or enhancing MBCAT expression, directly or indirectly, for example, by affecting an MBCAT function such as enzymatic (e.g., catalytic) or binding activity, can be identified using methods provided herein. MBCAT modulating agents are useful in diagnosis, therapy and pharmaceutical development.

Nucleic acids and polypeptides of the invention

Sequences related to MBCAT nucleic acids and polypeptides that can be used in the invention are disclosed in Genbank (referenced by Genbank identifier (GI) or RefSeq number), shown in Table 1 and in the appended sequence listing.

5 The term "MBCAT polypeptide" refers to a full-length MBCAT protein or a functionally active fragment or derivative thereof. A "functionally active" MBCAT fragment or derivative exhibits one or more functional activities associated with a full-length, wild-type MBCAT protein, such as antigenic or immunogenic activity, enzymatic activity, ability to bind natural cellular substrates, etc. The functional activity of MBCAT
10 proteins, derivatives and fragments can be assayed by various methods known to one skilled in the art (Current Protocols in Protein Science (1998) Coligan *et al.*, eds., John Wiley & Sons, Inc., Somerset, New Jersey) and as further discussed below. In one embodiment, a functionally active MBCAT polypeptide is an MBCAT derivative capable of rescuing defective endogenous MBCAT activity, such as in cell based or animal assays;
15 the rescuing derivative may be from the same or a different species. For purposes herein, functionally active fragments also include those fragments that comprise one or more structural domains of an MBCAT, such as a kinase domain or a binding domain. Protein domains can be identified using the PFAM program (Bateman A., et al., Nucleic Acids Res, 1999, 27:260-2). Methods for obtaining MBCAT polypeptides are also further
20 described below. In some embodiments, preferred fragments are functionally active, domain-containing fragments comprising at least 25 contiguous amino acids, preferably at least 50, more preferably 75, and most preferably at least 100 contiguous amino acids of an MBCAT. In further preferred embodiments, the fragment comprises the entire functionally active domain.

25 The term "MBCAT nucleic acid" refers to a DNA or RNA molecule that encodes a MBCAT polypeptide. Preferably, the MBCAT polypeptide or nucleic acid or fragment thereof is from a human, but can also be an ortholog, or derivative thereof with at least 70% sequence identity, preferably at least 80%, more preferably 85%, still more preferably 90%, and most preferably at least 95% sequence identity with human MBCAT.
30 Methods of identifying orthologs are known in the art. Normally, orthologs in different species retain the same function, due to presence of one or more protein motifs and/or 3-dimensional structures. Orthologs are generally identified by sequence homology analysis, such as BLAST analysis, usually using protein bait sequences. Sequences are assigned as a potential ortholog if the best hit sequence from the forward BLAST result

retrieves the original query sequence in the reverse BLAST (Huynen MA and Bork P, Proc Natl Acad Sci (1998) 95:5849-5856; Huynen MA *et al.*, Genome Research (2000) 10:1204-1210). Programs for multiple sequence alignment, such as CLUSTAL (Thompson JD *et al.*, 1994, Nucleic Acids Res 22:4673-4680) may be used to highlight
5 conserved regions and/or residues of orthologous proteins and to generate phylogenetic trees. In a phylogenetic tree representing multiple homologous sequences from diverse species (e.g., retrieved through BLAST analysis), orthologous sequences from two species generally appear closest on the tree with respect to all other sequences from these two species. Structural threading or other analysis of protein folding (e.g., using software by
10 ProCeryon, Biosciences, Salzburg, Austria) may also identify potential orthologs. In evolution, when a gene duplication event follows speciation, a single gene in one species, such as *C.elegans*, may correspond to multiple genes (paralogs) in another, such as human. As used herein, the term "orthologs" encompasses paralogs. As used herein, "percent (%) sequence identity" with respect to a subject sequence, or a specified portion of a subject
15 sequence, is defined as the percentage of nucleotides or amino acids in the candidate derivative sequence identical with the nucleotides or amino acids in the subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1997) 215:403-410) with all the search
20 parameters set to default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A % identity value is determined by the number of matching identical nucleotides or amino acids divided by the sequence length for which the percent identity is
25 being reported. "Percent (%) amino acid sequence similarity" is determined by doing the same calculation as for determining % amino acid sequence identity, but including conservative amino acid substitutions in addition to identical amino acids in the computation.

A conservative amino acid substitution is one in which an amino acid is substituted
30 for another amino acid having similar properties such that the folding or activity of the protein is not significantly affected. Aromatic amino acids that can be substituted for each other are phenylalanine, tryptophan, and tyrosine; interchangeable hydrophobic amino acids are leucine, isoleucine, methionine, and valine; interchangeable polar amino acids are glutamine and asparagine; interchangeable basic amino acids are arginine, lysine and

histidine; interchangeable acidic amino acids are aspartic acid and glutamic acid; and interchangeable small amino acids are alanine, serine, threonine, cysteine and glycine.

Alternatively, an alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman (Smith and Waterman, 1981, *Advances in Applied Mathematics* 2:482-489; database: European Bioinformatics Institute; Smith and Waterman, 1981, *J. of Molec.Biol.*, 147:195-197; Nicholas et al., 1998, "A Tutorial on Searching Sequence Databases and Sequence Scoring Methods" (www.psc.edu) and references cited therein.; W.R. Pearson, 1991, *Genomics* 11:635-650). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff (Dayhoff: *Atlas of Protein Sequences and Structure*, M. O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA), and normalized by Gribskov (Gribskov 1986 *Nucl. Acids Res.* 14(6):6745-6763). The Smith-Waterman algorithm may be employed where default parameters are used for scoring (for example, gap open penalty of 12, gap extension penalty of two). From the data generated, the "Match" value reflects "sequence identity."

Derivative nucleic acid molecules of the subject nucleic acid molecules include sequences that hybridize to the nucleic acid sequence of an MBCAT. The stringency of hybridization can be controlled by temperature, ionic strength, pH, and the presence of denaturing agents such as formamide during hybridization and washing. Conditions routinely used are set out in readily available procedure texts (*e.g.*, *Current Protocol in Molecular Biology*, Vol. 1, Chap. 2.10, John Wiley & Sons, Publishers (1994); Sambrook *et al.*, *Molecular Cloning*, Cold Spring Harbor (1989)). In some embodiments, a nucleic acid molecule of the invention is capable of hybridizing to a nucleic acid molecule containing the nucleotide sequence of an MBCAT under high stringency hybridization conditions that are: prehybridization of filters containing nucleic acid for 8 hours to overnight at 65° C in a solution comprising 6X single strength citrate (SSC) (1X SSC is 0.15 M NaCl, 0.015 M Na citrate; pH 7.0), 5X Denhardt's solution, 0.05% sodium pyrophosphate and 100 µg/ml herring sperm DNA; hybridization for 18-20 hours at 65° C in a solution containing 6X SSC, 1X Denhardt's solution, 100 µg/ml yeast tRNA and 0.05% sodium pyrophosphate; and washing of filters at 65° C for 1h in a solution containing 0.1X SSC and 0.1% SDS (sodium dodecyl sulfate).

In other embodiments, moderately stringent hybridization conditions are used that are: pretreatment of filters containing nucleic acid for 6 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.1% PVP, 0.1%

Ficoll, 1% BSA, and 500 $\mu\text{g/ml}$ denatured salmon sperm DNA; hybridization for 18-20h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 $\mu\text{g/ml}$ salmon sperm DNA, and 10% (wt/vol) dextran sulfate; followed by washing twice for 1 hour at 55° C in a solution containing 2X SSC and 0.1% SDS.

Alternatively, low stringency conditions can be used that are: incubation for 8 hours to overnight at 37° C in a solution comprising 20% formamide, 5 x SSC, 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 $\mu\text{g/ml}$ denatured sheared salmon sperm DNA; hybridization in the same buffer for 18 to 20 hours; and washing of filters in 1 x SSC at about 37° C for 1 hour.

Isolation, Production, Expression, and Mis-expression of MBCAT Nucleic Acids and Polypeptides

MBCAT nucleic acids and polypeptides are useful for identifying and testing agents that modulate MBCAT function and for other applications related to the involvement of MBCAT in the beta-catenin pathway. MBCAT nucleic acids and derivatives and orthologs thereof may be obtained using any available method. For instance, techniques for isolating cDNA or genomic DNA sequences of interest by screening DNA libraries or by using polymerase chain reaction (PCR) are well known in the art. In general, the particular use for the protein will dictate the particulars of expression, production, and purification methods. For instance, production of proteins for use in screening for modulating agents may require methods that preserve specific biological activities of these proteins, whereas production of proteins for antibody generation may require structural integrity of particular epitopes. Expression of proteins to be purified for screening or antibody production may require the addition of specific tags (*e.g.*, generation of fusion proteins). Overexpression of an MBCAT protein for assays used to assess MBCAT function, such as involvement in cell cycle regulation or hypoxic response, may require expression in eukaryotic cell lines capable of these cellular activities. Techniques for the expression, production, and purification of proteins are well known in the art; any suitable means therefore may be used (*e.g.*, Higgins SJ and Hames BD (eds.) Protein Expression: A Practical Approach, Oxford University Press Inc., New York 1999; Stanbury PF et al., Principles of Fermentation Technology, 2nd edition, Elsevier Science, New York, 1995; Doonan S (ed.) Protein Purification Protocols, Humana Press, New Jersey, 1996; Coligan JE et al, Current Protocols in Protein Science

(eds.), 1999, John Wiley & Sons, New York). In particular embodiments, recombinant MBCAT is expressed in a cell line known to have defective beta-catenin function. The recombinant cells are used in cell-based screening assay systems of the invention, as described further below.

5 The nucleotide sequence encoding an MBCAT polypeptide can be inserted into any appropriate expression vector. The necessary transcriptional and translational signals, including promoter/enhancer element, can derive from the native MBCAT gene and/or its flanking regions or can be heterologous. A variety of host-vector expression systems may be utilized, such as mammalian cell systems infected with virus (*e.g.* vaccinia virus,
10 adenovirus, *etc.*); insect cell systems infected with virus (*e.g.* baculovirus); microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage, plasmid, or cosmid DNA. An isolated host cell strain that modulates the expression of, modifies, and/or specifically processes the gene product may be used.

 To detect expression of the MBCAT gene product, the expression vector can
15 comprise a promoter operably linked to an MBCAT gene nucleic acid, one or more origins of replication, and, one or more selectable markers (*e.g.* thymidine kinase activity, resistance to antibiotics, *etc.*). Alternatively, recombinant expression vectors can be identified by assaying for the expression of the MBCAT gene product based on the physical or functional properties of the MBCAT protein in *in vitro* assay systems (*e.g.*
20 immunoassays).

 The MBCAT protein, fragment, or derivative may be optionally expressed as a fusion, or chimeric protein product (i.e. it is joined via a peptide bond to a heterologous protein sequence of a different protein), for example to facilitate purification or detection. A chimeric product can be made by ligating the appropriate nucleic acid sequences
25 encoding the desired amino acid sequences to each other using standard methods and expressing the chimeric product. A chimeric product may also be made by protein synthetic techniques, *e.g.* by use of a peptide synthesizer (Hunkapiller *et al.*, Nature (1984) 310:105-111).

 Once a recombinant cell that expresses the MBCAT gene sequence is identified,
30 the gene product can be isolated and purified using standard methods (*e.g.* ion exchange, affinity, and gel exclusion chromatography; centrifugation; differential solubility; electrophoresis). Alternatively, native MBCAT proteins can be purified from natural sources, by standard methods (*e.g.* immunoaffinity purification). Once a protein is obtained, it may be quantified and its activity measured by appropriate methods, such as

immunoassay, bioassay, or other measurements of physical properties, such as crystallography.

The methods of this invention may also use cells that have been engineered for altered expression (mis-expression) of MBCAT or other genes associated with the beta-catenin pathway. As used herein, mis-expression encompasses ectopic expression, over-expression, under-expression, and non-expression (*e.g.* by gene knock-out or blocking expression that would otherwise normally occur).

Genetically modified animals

Animal models that have been genetically modified to alter MBCAT expression may be used in *in vivo* assays to test for activity of a candidate beta-catenin modulating agent, or to further assess the role of MBCAT in a beta-catenin pathway process such as apoptosis or cell proliferation. Preferably, the altered MBCAT expression results in a detectable phenotype, such as decreased or increased levels of cell proliferation, angiogenesis, or apoptosis compared to control animals having normal MBCAT expression. The genetically modified animal may additionally have altered beta-catenin expression (*e.g.* beta-catenin knockout). Preferred genetically modified animals are mammals such as primates, rodents (preferably mice or rats), among others. Preferred non-mammalian species include zebrafish, *C. elegans*, and *Drosophila*. Preferred genetically modified animals are transgenic animals having a heterologous nucleic acid sequence present as an extrachromosomal element in a portion of its cells, *i.e.* mosaic animals (see, for example, techniques described by Jakobovits, 1994, *Curr. Biol.* 4:761-763.) or stably integrated into its germ line DNA (*i.e.*, in the genomic sequence of most or all of its cells). Heterologous nucleic acid is introduced into the germ line of such transgenic animals by genetic manipulation of, for example, embryos or embryonic stem cells of the host animal.

Methods of making transgenic animals are well-known in the art (for transgenic mice see Brinster *et al.*, *Proc. Nat. Acad. Sci. USA* 82: 4438-4442 (1985), U.S. Pat. Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Pat. No. 4,873,191 by Wagner *et al.*, and Hogan, B., *Manipulating the Mouse Embryo*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1986); for particle bombardment see U.S. Pat. No., 4,945,050, by Sandford *et al.*; for transgenic *Drosophila* see Rubin and Spradling, *Science* (1982) 218:348-53 and U.S. Pat. No. 4,670,388; for transgenic insects see Berghammer A.J. *et al.*, *A Universal Marker for Transgenic Insects* (1999) *Nature* 402:370-371; for transgenic

Zebrafish see Lin S., Transgenic Zebrafish, Methods Mol Biol. (2000);136:375-3830); for microinjection procedures for fish, amphibian eggs and birds see Houdebine and Chourrout, Experientia (1991) 47:897-905; for transgenic rats see Hammer *et al.*, Cell (1990) 63:1099-1112; and for culturing of embryonic stem (ES) cells and the subsequent production of transgenic animals by the introduction of DNA into ES cells using methods such as electroporation, calcium phosphate/DNA precipitation and direct injection see, e.g., Teratocarcinomas and Embryonic Stem Cells, A Practical Approach, E. J. Robertson, ed., IRL Press (1987)). Clones of the nonhuman transgenic animals can be produced according to available methods (see Wilmut, I. *et al.* (1997) Nature 385:810-813; and PCT International Publication Nos. WO 97/07668 and WO 97/07669).

In one embodiment, the transgenic animal is a "knock-out" animal having a heterozygous or homozygous alteration in the sequence of an endogenous MBCAT gene that results in a decrease of MBCAT function, preferably such that MBCAT expression is undetectable or insignificant. Knock-out animals are typically generated by homologous recombination with a vector comprising a transgene having at least a portion of the gene to be knocked out. Typically a deletion, addition or substitution has been introduced into the transgene to functionally disrupt it. The transgene can be a human gene (e.g., from a human genomic clone) but more preferably is an ortholog of the human gene derived from the transgenic host species. For example, a mouse MBCAT gene is used to construct a homologous recombination vector suitable for altering an endogenous MBCAT gene in the mouse genome. Detailed methodologies for homologous recombination in mice are available (see Capecchi, Science (1989) 244:1288-1292; Joyner *et al.*, Nature (1989) 338:153-156). Procedures for the production of non-rodent transgenic mammals and other animals are also available (Houdebine and Chourrout, *supra*; Pursel *et al.*, Science (1989) 244:1281-1288; Simms *et al.*, Bio/Technology (1988) 6:179-183). In a preferred embodiment, knock-out animals, such as mice harboring a knockout of a specific gene, may be used to produce antibodies against the human counterpart of the gene that has been knocked out (Claesson MH *et al.*, (1994) Scan J Immunol 40:257-264; Declerck PJ *et al.*, (1995) J Biol Chem. 270:8397-400).

In another embodiment, the transgenic animal is a "knock-in" animal having an alteration in its genome that results in altered expression (e.g., increased (including ectopic) or decreased expression) of the MBCAT gene, e.g., by introduction of additional copies of MBCAT, or by operatively inserting a regulatory sequence that provides for altered expression of an endogenous copy of the MBCAT gene. Such regulatory

sequences include inducible, tissue-specific, and constitutive promoters and enhancer elements. The knock-in can be homozygous or heterozygous.

Transgenic nonhuman animals can also be produced that contain selected systems allowing for regulated expression of the transgene. One example of such a system that
5 may be produced is the cre/loxP recombinase system of bacteriophage P1 (Lakso *et al.*, PNAS (1992) 89:6232-6236; U.S. Pat. No. 4,959,317). If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two
10 transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman *et al.* (1991) Science 251:1351-1355; U.S. Pat. No. 5,654,182). In a preferred embodiment, both Cre-LoxP and Flp-Frt are used in the same system to regulate expression of the
15 transgene, and for sequential deletion of vector sequences in the same cell (Sun X *et al.* (2000) Nat Genet 25:83-6).

The genetically modified animals can be used in genetic studies to further elucidate the beta-catenin pathway, as animal models of disease and disorders implicating defective beta-catenin function, and for *in vivo* testing of candidate therapeutic agents, such as those
20 identified in screens described below. The candidate therapeutic agents are administered to a genetically modified animal having altered MBCAT function and phenotypic changes are compared with appropriate control animals such as genetically modified animals that receive placebo treatment, and/or animals with unaltered MBCAT expression that receive candidate therapeutic agent.

25 In addition to the above-described genetically modified animals having altered MBCAT function, animal models having defective beta-catenin function (and otherwise normal MBCAT function), can be used in the methods of the present invention. For example, a beta-catenin knockout mouse can be used to assess, *in vivo*, the activity of a candidate beta-catenin modulating agent identified in one of the *in vitro* assays described
30 below. Preferably, the candidate beta-catenin modulating agent when administered to a model system with cells defective in beta-catenin function, produces a detectable phenotypic change in the model system indicating that the beta-catenin function is restored, i.e., the cells exhibit normal cell cycle progression.

Modulating Agents

The invention provides methods to identify agents that interact with and/or modulate the function of MBCAT and/or the beta-catenin pathway. Modulating agents identified by the methods are also part of the invention. Such agents are useful in a variety of diagnostic and therapeutic applications associated with the beta-catenin pathway, as well as in further analysis of the MBCAT protein and its contribution to the beta-catenin pathway. Accordingly, the invention also provides methods for modulating the beta-catenin pathway comprising the step of specifically modulating MBCAT activity by administering a MBCAT-interacting or -modulating agent.

As used herein, an "MBCAT-modulating agent" is any agent that modulates MBCAT function, for example, an agent that interacts with MBCAT to inhibit or enhance MBCAT activity or otherwise affect normal MBCAT function. MBCAT function can be affected at any level, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In a preferred embodiment, the MBCAT - modulating agent specifically modulates the function of the MBCAT. The phrases "specific modulating agent", "specifically modulates", etc., are used herein to refer to modulating agents that directly bind to the MBCAT polypeptide or nucleic acid, and preferably inhibit, enhance, or otherwise alter, the function of the MBCAT. These phrases also encompass modulating agents that alter the interaction of the MBCAT with a binding partner, substrate, or cofactor (e.g. by binding to a binding partner of an MBCAT, or to a protein/binding partner complex, and altering MBCAT function). In a further preferred embodiment, the MBCAT- modulating agent is a modulator of the beta-catenin pathway (e.g. it restores and/or upregulates beta-catenin function) and thus is also a beta-catenin-modulating agent.

Preferred MBCAT-modulating agents include small molecule compounds; MBCAT-interacting proteins, including antibodies and other biotherapeutics; and nucleic acid modulators such as antisense and RNA inhibitors. The modulating agents may be formulated in pharmaceutical compositions, for example, as compositions that may comprise other active ingredients, as in combination therapy, and/or suitable carriers or excipients. Techniques for formulation and administration of the compounds may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton, PA, 19th edition.

Small molecule modulators

Small molecules are often preferred to modulate function of proteins with enzymatic function, and/or containing protein interaction domains. Chemical agents, referred to in the art as "small molecule" compounds are typically organic, non-peptide molecules, having a molecular weight up to 10,000, preferably up to 5,000, more preferably up to 1,000, and most preferably up to 500 daltons. This class of modulators includes chemically synthesized molecules, for instance, compounds from combinatorial chemical libraries. Synthetic compounds may be rationally designed or identified based on known or inferred properties of the MBCAT protein or may be identified by screening compound libraries. Alternative appropriate modulators of this class are natural products, particularly secondary metabolites from organisms such as plants or fungi, which can also be identified by screening compound libraries for MBCAT-modulating activity. Methods for generating and obtaining compounds are well known in the art (Schreiber SL, Science (2000) 151: 1964-1969; Radmann J and Gunther J, Science (2000) 151:1947-1948).

Small molecule modulators identified from screening assays, as described below, can be used as lead compounds from which candidate clinical compounds may be designed, optimized, and synthesized. Such clinical compounds may have utility in treating pathologies associated with the beta-catenin pathway. The activity of candidate small molecule modulating agents may be improved several-fold through iterative secondary functional validation, as further described below, structure determination, and candidate modulator modification and testing. Additionally, candidate clinical compounds are generated with specific regard to clinical and pharmacological properties. For example, the reagents may be derivatized and re-screened using *in vitro* and *in vivo* assays to optimize activity and minimize toxicity for pharmaceutical development.

Protein Modulators

Specific MBCAT-interacting proteins are useful in a variety of diagnostic and therapeutic applications related to the beta-catenin pathway and related disorders, as well as in validation assays for other MBCAT-modulating agents. In a preferred embodiment, MBCAT-interacting proteins affect normal MBCAT function, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In another embodiment, MBCAT-interacting proteins are useful in detecting and providing information about the function of MBCAT proteins, as is relevant to beta-catenin related disorders, such as cancer (e.g., for diagnostic means).

An MBCAT-interacting protein may be endogenous, i.e. one that naturally interacts genetically or biochemically with an MBCAT, such as a member of the MBCAT pathway that modulates MBCAT expression, localization, and/or activity. MBCAT-modulators include dominant negative forms of MBCAT-interacting proteins and of MBCAT proteins themselves. Yeast two-hybrid and variant screens offer preferred methods for identifying endogenous MBCAT-interacting proteins (Finley, R. L. et al. (1996) in DNA Cloning-Expression Systems: A Practical Approach, eds. Glover D. & Hames B. D (Oxford University Press, Oxford, England), pp. 169-203; Fashema SF et al., Gene (2000) 250:1-14; Drees BL Curr Opin Chem Biol (1999) 3:64-70; Vidal M and Legrain P Nucleic Acids Res (1999) 27:919-29; and U.S. Pat. No. 5,928,868). Mass spectrometry is an alternative preferred method for the elucidation of protein complexes (reviewed in, e.g., Pandley A and Mann M, Nature (2000) 405:837-846; Yates JR 3rd, Trends Genet (2000) 16:5-8).

An MBCAT-interacting protein may be an exogenous protein, such as an MBCAT-specific antibody or a T-cell antigen receptor (see, e.g., Harlow and Lane (1988) Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory; Harlow and Lane (1999) Using antibodies: a laboratory manual. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press). MBCAT antibodies are further discussed below.

In preferred embodiments, an MBCAT-interacting protein specifically binds an MBCAT protein. In alternative preferred embodiments, an MBCAT-modulating agent binds an MBCAT substrate, binding partner, or cofactor.

Antibodies

In another embodiment, the protein modulator is an MBCAT specific antibody agonist or antagonist. The antibodies have therapeutic and diagnostic utilities, and can be used in screening assays to identify MBCAT modulators. The antibodies can also be used in dissecting the portions of the MBCAT pathway responsible for various cellular responses and in the general processing and maturation of the MBCAT.

Antibodies that specifically bind MBCAT polypeptides can be generated using known methods. Preferably the antibody is specific to a mammalian ortholog of MBCAT polypeptide, and more preferably, to human MBCAT. Antibodies may be polyclonal, monoclonal (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab').sub.2 fragments, fragments produced by a FAb expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

Epitopes of MBCAT which are particularly antigenic can be selected, for example, by routine screening of MBCAT polypeptides for antigenicity or by applying a theoretical method for selecting antigenic regions of a protein (Hopp and Wood (1981), Proc. Natl. Acad. Sci. U.S.A. 78:3824-28; Hopp and Wood, (1983) Mol. Immunol. 20:483-89; 5 Sutcliffe et al., (1983) Science 219:660-66) to the amino acid sequence of an MBCAT. Monoclonal antibodies with affinities of 10^8 M^{-1} preferably 10^9 M^{-1} to 10^{10} M^{-1} , or stronger can be made by standard procedures as described (Harlow and Lane, *supra*; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed) Academic Press, New York; and U.S. Pat. Nos. 4,381,292; 4,451,570; and 4,618,577). Antibodies may be 10 generated against crude cell extracts of MBCAT or substantially purified fragments thereof. If MBCAT fragments are used, they preferably comprise at least 10, and more preferably, at least 20 contiguous amino acids of an MBCAT protein. In a particular embodiment, MBCAT-specific antigens and/or immunogens are coupled to carrier proteins that stimulate the immune response. For example, the subject polypeptides are 15 covalently coupled to the keyhole limpet hemocyanin (KLH) carrier, and the conjugate is emulsified in Freund's complete adjuvant, which enhances the immune response. An appropriate immune system such as a laboratory rabbit or mouse is immunized according to conventional protocols.

The presence of MBCAT-specific antibodies is assayed by an appropriate assay 20 such as a solid phase enzyme-linked immunosorbant assay (ELISA) using immobilized corresponding MBCAT polypeptides. Other assays, such as radioimmunoassays or fluorescent assays might also be used.

Chimeric antibodies specific to MBCAT polypeptides can be made that contain different portions from different animal species. For instance, a human immunoglobulin 25 constant region may be linked to a variable region of a murine mAb, such that the antibody derives its biological activity from the human antibody, and its binding specificity from the murine fragment. Chimeric antibodies are produced by splicing together genes that encode the appropriate regions from each species (Morrison et al., Proc. Natl. Acad. Sci. (1984) 81:6851-6855; Neuberger et al., Nature (1984) 312:604-608; 30 Takeda et al., Nature (1985) 31:452-454). Humanized antibodies, which are a form of chimeric antibodies, can be generated by grafting complementary-determining regions (CDRs) (Carlos, T. M., J. M. Harlan. 1994. Blood 84:2068-2101) of mouse antibodies into a background of human framework regions and constant regions by recombinant DNA technology (Riechmann LM, et al., 1988 Nature 323: 323-327). Humanized

antibodies contain ~10% murine sequences and ~90% human sequences, and thus further reduce or eliminate immunogenicity, while retaining the antibody specificities (Co MS, and Queen C. 1991 Nature 351: 501-501; Morrison SL. 1992 Ann. Rev. Immun. 10:239-265). Humanized antibodies and methods of their production are well-known in the art (U.S. Pat. Nos. 5,530,101, 5,585,089, 5,693,762, and 6,180,370).

MBCAT-specific single chain antibodies which are recombinant, single chain polypeptides formed by linking the heavy and light chain fragments of the Fv regions via an amino acid bridge, can be produced by methods known in the art (U.S. Pat. No. 4,946,778; Bird, Science (1988) 242:423-426; Huston et al., Proc. Natl. Acad. Sci. USA (1988) 85:5879-5883; and Ward et al., Nature (1989) 334:544-546).

Other suitable techniques for antibody production involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors (Huse et al., Science (1989) 246:1275-1281). As used herein, T-cell antigen receptors are included within the scope of antibody modulators (Harlow and Lane, 1988, *supra*).

The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, antibodies will be labeled by joining, either covalently or non-covalently, a substance that provides for a detectable signal, or that is toxic to cells that express the targeted protein (Menard S, et al., Int J. Biol Markers (1989) 4:131-134). A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, fluorescent emitting lanthanide metals, chemiluminescent moieties, bioluminescent moieties, magnetic particles, and the like (U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241). Also, recombinant immunoglobulins may be produced (U.S. Pat. No. 4,816,567). Antibodies to cytoplasmic polypeptides may be delivered and reach their targets by conjugation with membrane-penetrating toxin proteins (U.S. Pat. No. 6,086,900).

When used therapeutically in a patient, the antibodies of the subject invention are typically administered parenterally, when possible at the target site, or intravenously. The therapeutically effective dose and dosage regimen is determined by clinical studies. Typically, the amount of antibody administered is in the range of about 0.1 mg/kg –to about 10 mg/kg of patient weight. For parenteral administration, the antibodies are formulated in a unit dosage injectable form (e.g., solution, suspension, emulsion) in

association with a pharmaceutically acceptable vehicle. Such vehicles are inherently nontoxic and non-therapeutic. Examples are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Nonaqueous vehicles such as fixed oils, ethyl oleate, or liposome carriers may also be used. The vehicle may contain minor amounts of additives, such as buffers and preservatives, which enhance isotonicity and chemical stability or otherwise enhance therapeutic potential. The antibodies' concentrations in such vehicles are typically in the range of about 1 mg/ml to about 10 mg/ml.

Immunotherapeutic methods are further described in the literature (US Pat. No. 5,859,206; WO0073469).

Nucleic Acid Modulators

Other preferred MBCAT-modulating agents comprise nucleic acid molecules, such as antisense oligomers or double stranded RNA (dsRNA), which generally inhibit MBCAT activity. Preferred nucleic acid modulators interfere with the function of the MBCAT nucleic acid such as DNA replication, transcription, translocation of the MBCAT RNA to the site of protein translation, translation of protein from the MBCAT RNA, splicing of the MBCAT RNA to yield one or more mRNA species, or catalytic activity which may be engaged in or facilitated by the MBCAT RNA.

In one embodiment, the antisense oligomer is an oligonucleotide that is sufficiently complementary to an MBCAT mRNA to bind to and prevent translation, preferably by binding to the 5' untranslated region. MBCAT-specific antisense oligonucleotides, preferably range from at least 6 to about 200 nucleotides. In some embodiments the oligonucleotide is preferably at least 10, 15, or 20 nucleotides in length. In other embodiments, the oligonucleotide is preferably less than 50, 40, or 30 nucleotides in length. The oligonucleotide can be DNA or RNA or a chimeric mixture or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide may include other appending groups such as peptides, agents that facilitate transport across the cell membrane, hybridization-triggered cleavage agents, and intercalating agents.

In another embodiment, the antisense oligomer is a phosphothioate morpholino oligomer (PMO). PMOs are assembled from four different morpholino subunits, each of which contain one of four genetic bases (A, C, G, or T) linked to a six-membered morpholine ring. Polymers of these subunits are joined by non-ionic phosphodiamidate

intersubunit linkages. Details of how to make and use PMOs and other antisense oligomers are well known in the art (e.g. see WO99/18193; Probst JC, Antisense Oligodeoxynucleotide and Ribozyme Design, Methods. (2000) 22(3):271-281; Summerton J, and Weller D. 1997 Antisense Nucleic Acid Drug Dev. :7:187-95; US Pat. No.

5 5,235,033; and US Pat No. 5,378,841).

Alternative preferred MBCAT nucleic acid modulators are double-stranded RNA species mediating RNA interference (RNAi). RNAi is the process of sequence-specific, post-transcriptional gene silencing in animals and plants, initiated by double-stranded RNA (dsRNA) that is homologous in sequence to the silenced gene. Methods relating to
10 the use of RNAi to silence genes in *C. elegans*, *Drosophila*, plants, and humans are known in the art (Fire A, et al., 1998 Nature 391:806-811; Fire, A. Trends Genet. 15, 358-363 (1999); Sharp, P. A. RNA interference 2001. Genes Dev. 15, 485-490 (2001); Hammond, S. M., et al., Nature Rev. Genet. 2, 110-1119 (2001); Tuschl, T. Chem. Biochem. 2, 239-245 (2001); Hamilton, A. et al., Science 286, 950-952 (1999); Hammond, S. M., et al.,
15 Nature 404, 293-296 (2000); Zamore, P. D., et al., Cell 101, 25-33 (2000); Bernstein, E., et al., Nature 409, 363-366 (2001); Elbashir, S. M., et al., Genes Dev. 15, 188-200 (2001); WO0129058; WO9932619; Elbashir SM, et al., 2001 Nature 411:494-498).

Nucleic acid modulators are commonly used as research reagents, diagnostics, and therapeutics. For example, antisense oligonucleotides, which are able to inhibit gene
20 expression with exquisite specificity, are often used to elucidate the function of particular genes (see, for example, U.S. Pat. No. 6,165,790). Nucleic acid modulators are also used, for example, to distinguish between functions of various members of a biological pathway. For example, antisense oligomers have been employed as therapeutic moieties in the treatment of disease states in animals and man and have been demonstrated in numerous
25 clinical trials to be safe and effective (Milligan JF, *et al*, Current Concepts in Antisense Drug Design, J Med Chem. (1993) 36:1923-1937; Tonkinson JL *et al.*, Antisense Oligodeoxynucleotides as Clinical Therapeutic Agents, Cancer Invest. (1996) 14:54-65). Accordingly, in one aspect of the invention, an MBCAT-specific nucleic acid modulator is used in an assay to further elucidate the role of the MBCAT in the beta-catenin pathway,
30 and/or its relationship to other members of the pathway. In another aspect of the invention, an MBCAT-specific antisense oligomer is used as a therapeutic agent for treatment of beta-catenin-related disease states.

Assay Systems

The invention provides assay systems and screening methods for identifying specific modulators of MBCAT activity. As used herein, an "assay system" encompasses all the components required for performing and analyzing results of an assay that detects and/or measures a particular event. In general, primary assays are used to identify or confirm a modulator's specific biochemical or molecular effect with respect to the MBCAT nucleic acid or protein. In general, secondary assays further assess the activity of a MBCAT modulating agent identified by a primary assay and may confirm that the modulating agent affects MBCAT in a manner relevant to the beta-catenin pathway. In some cases, MBCAT modulators will be directly tested in a secondary assay.

In a preferred embodiment, the screening method comprises contacting a suitable assay system comprising an MBCAT polypeptide or nucleic acid with a candidate agent under conditions whereby, but for the presence of the agent, the system provides a reference activity (e.g. binding activity), which is based on the particular molecular event the screening method detects. A statistically significant difference between the agent-biased activity and the reference activity indicates that the candidate agent modulates MBCAT activity, and hence the beta-catenin pathway. The MBCAT polypeptide or nucleic acid used in the assay may comprise any of the nucleic acids or polypeptides described above.

Primary Assays

The type of modulator tested generally determines the type of primary assay.

Primary assays for small molecule modulators

For small molecule modulators, screening assays are used to identify candidate modulators. Screening assays may be cell-based or may use a cell-free system that recreates or retains the relevant biochemical reaction of the target protein (reviewed in Sittampalam GS *et al.*, Curr Opin Chem Biol (1997) 1:384-91 and accompanying references). As used herein the term "cell-based" refers to assays using live cells, dead cells, or a particular cellular fraction, such as a membrane, endoplasmic reticulum, or mitochondrial fraction. The term "cell free" encompasses assays using substantially purified protein (either endogenous or recombinantly produced), partially purified or crude cellular extracts. Screening assays may detect a variety of molecular events, including protein-DNA interactions, protein-protein interactions (*e.g.*, receptor-ligand binding),

transcriptional activity (*e.g.*, using a reporter gene), enzymatic activity (*e.g.*, via a property of the substrate), activity of second messengers, immunogenicity and changes in cellular morphology or other cellular characteristics. Appropriate screening assays may use a wide range of detection methods including fluorescent, radioactive, colorimetric,

5 spectrophotometric, and amperometric methods, to provide a read-out for the particular molecular event detected.

Cell-based screening assays usually require systems for recombinant expression of MBCAT and any auxiliary proteins demanded by the particular assay. Appropriate methods for generating recombinant proteins produce sufficient quantities of proteins that
10 retain their relevant biological activities and are of sufficient purity to optimize activity and assure assay reproducibility. Yeast two-hybrid and variant screens, and mass spectrometry provide preferred methods for determining protein-protein interactions and elucidation of protein complexes. In certain applications, when MBCAT-interacting proteins are used in screens to identify small molecule modulators, the binding specificity
15 of the interacting protein to the MBCAT protein may be assayed by various known methods such as substrate processing (*e.g.* ability of the candidate MBCAT-specific binding agents to function as negative effectors in MBCAT-expressing cells), binding equilibrium constants (usually at least about 10^7 M^{-1} , preferably at least about 10^8 M^{-1} , more preferably at least about 10^9 M^{-1}), and immunogenicity (*e.g.* ability to elicit MBCAT
20 specific antibody in a heterologous host such as a mouse, rat, goat or rabbit). For enzymes and receptors, binding may be assayed by, respectively, substrate and ligand processing.

The screening assay may measure a candidate agent's ability to specifically bind to or modulate activity of a MBCAT polypeptide, a fusion protein thereof, or to cells or membranes bearing the polypeptide or fusion protein. The MBCAT polypeptide can be
25 full length or a fragment thereof that retains functional MBCAT activity. The MBCAT polypeptide may be fused to another polypeptide, such as a peptide tag for detection or anchoring, or to another tag. The MBCAT polypeptide is preferably human MBCAT, or is an ortholog or derivative thereof as described above. In a preferred embodiment, the screening assay detects candidate agent-based modulation of MBCAT interaction with a
30 binding target, such as an endogenous or exogenous protein or other substrate that has MBCAT-specific binding activity, and can be used to assess normal MBCAT gene function.

Suitable assay formats that may be adapted to screen for MBCAT modulators are known in the art. Preferred screening assays are high throughput or ultra high throughput

and thus provide automated, cost-effective means of screening compound libraries for lead compounds (Fernandes PB, Curr Opin Chem Biol (1998) 2:597-603; Sundberg SA, Curr Opin Biotechnol 2000, 11:47-53). In one preferred embodiment, screening assays uses fluorescence technologies, including fluorescence polarization, time-resolved
5 fluorescence, and fluorescence resonance energy transfer. These systems offer means to monitor protein-protein or DNA-protein interactions in which the intensity of the signal emitted from dye-labeled molecules depends upon their interactions with partner molecules (*e.g.*, Selvin PR, Nat Struct Biol (2000) 7:730-4; Fernandes PB, *supra*; Hertzberg RP and Pope AJ, Curr Opin Chem Biol (2000) 4:445-451).

10 A variety of suitable assay systems may be used to identify candidate MBCAT and beta-catenin pathway modulators (*e.g.* U.S. Pat. No. 6,165,992 (kinase assays); U.S. Pat. Nos. 5,550,019 and 6,133,437 (apoptosis assays); and U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434 (angiogenesis assays), among others). Specific preferred assays are described in more detail below.

15 **Kinase assays.** In some preferred embodiments the screening assay detects the ability of the test agent to modulate the kinase activity of an MBCAT polypeptide. In further embodiments, a cell-free kinase assay system is used to identify a candidate beta-catenin modulating agent, and a secondary, cell-based assay, such as an apoptosis or
20 hypoxic induction assay (described below), may be used to further characterize the candidate beta-catenin modulating agent. Many different assays for kinases have been reported in the literature and are well known to those skilled in the art (*e.g.* U.S. Pat. No. 6,165,992; Zhu et al., Nature Genetics (2000) 26:283-289; and WO0073469).
Radioassays, which monitor the transfer of a gamma phosphate are frequently used. For
25 instance, a scintillation assay for p56 (lck) kinase activity monitors the transfer of the gamma phosphate from gamma -³³P ATP to a biotinylated peptide substrate; the substrate is captured on a streptavidin coated bead that transmits the signal (Beveridge M *et al.*, J Biomol Screen (2000) 5:205-212). This assay uses the scintillation proximity assay (SPA), in which only radio-ligand bound to receptors tethered to the surface of an SPA
30 bead are detected by the scintillant immobilized within it, allowing binding to be measured without separation of bound from free ligand.

Other assays for protein kinase activity may use antibodies that specifically recognize phosphorylated substrates. For instance, the kinase receptor activation (KIRA) assay measures receptor tyrosine kinase activity by ligand stimulating the intact receptor

in cultured cells, then capturing solubilized receptor with specific antibodies and quantifying phosphorylation via phosphotyrosine ELISA (Sadick MD, Dev Biol Stand (1999) 97:121-133).

Another example of antibody based assays for protein kinase activity is TRF (time-resolved fluorometry). This method utilizes europium chelate-labeled anti-phosphotyrosine antibodies to detect phosphate transfer to a polymeric substrate coated onto microtiter plate wells. The amount of phosphorylation is then detected using time-resolved, dissociation-enhanced fluorescence (Braunwalder AF, et al., Anal Biochem 1996 Jul 1;238(2):159-64).

10

Apoptosis assays. Assays for apoptosis may be performed by terminal deoxynucleotidyl transferase-mediated digoxigenin-11-dUTP nick end labeling (TUNEL) assay. The TUNEL assay is used to measure nuclear DNA fragmentation characteristic of apoptosis (Lazebnik *et al.*, 1994, Nature 371, 346), by following the incorporation of fluorescein-dUTP (Yonehara *et al.*, 1989, J. Exp. Med. 169, 1747). Apoptosis may further be assayed by acridine orange staining of tissue culture cells (Lucas, R., et al., 1998, Blood 15:4730-41). Other cell-based apoptosis assays include the caspase-3/7 assay and the cell death nucleosome ELISA assay. The caspase 3/7 assay is based on the activation of the caspase cleavage activity as part of a cascade of events that occur during programmed cell death in many apoptotic pathways. In the caspase 3/7 assay (commercially available Apo-ONE™ Homogeneous Caspase-3/7 assay from Promega, cat# 67790), lysis buffer and caspase substrate are mixed and added to cells. The caspase substrate becomes fluorescent when cleaved by active caspase 3/7. The nucleosome ELISA assay is a general cell death assay known to those skilled in the art, and available commercially (Roche, Cat# 1774425). This assay is a quantitative sandwich-enzyme-immunoassay which uses monoclonal antibodies directed against DNA and histones respectively, thus specifically determining amount of mono- and oligonucleosomes in the cytoplasmic fraction of cell lysates. Mono and oligonucleosomes are enriched in the cytoplasm during apoptosis due to the fact that DNA fragmentation occurs several hours before the plasma membrane breaks down, allowing for accumulation in the cytoplasm. Nucleosomes are not present in the cytoplasmic fraction of cells that are not undergoing apoptosis. An apoptosis assay system may comprise a cell that expresses an MBCAT, and that optionally has defective beta-catenin function (e.g. beta-catenin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the apoptosis assay system and changes in

30

induction of apoptosis relative to controls where no test agent is added, identify candidate beta-catenin modulating agents. In some embodiments of the invention, an apoptosis assay may be used as a secondary assay to test a candidate beta-catenin modulating agents that is initially identified using a cell-free assay system. An apoptosis assay may also be used to test whether MBCAT function plays a direct role in apoptosis. For example, an apoptosis assay may be performed on cells that over- or under-express MBCAT relative to wild type cells. Differences in apoptotic response compared to wild type cells suggests that the MBCAT plays a direct role in the apoptotic response. Apoptosis assays are described further in US Pat. No. 6,133,437.

10

Cell proliferation and cell cycle assays. Cell proliferation may be assayed via bromodeoxyuridine (BRDU) incorporation. This assay identifies a cell population undergoing DNA synthesis by incorporation of BRDU into newly-synthesized DNA. Newly-synthesized DNA may then be detected using an anti-BRDU antibody (Hoshino *et al.*, 1986, *Int. J. Cancer* 38, 369; Campana *et al.*, 1988, *J. Immunol. Meth.* 107, 79), or by other means.

Cell proliferation is also assayed via phospho-histone H3 staining, which identifies a cell population undergoing mitosis by phosphorylation of histone H3. Phosphorylation of histone H3 at serine 10 is detected using an antibody specific to the phosphorylated form of the serine 10 residue of histone H3. (Chadlee, D.N. 1995, *J. Biol. Chem* 270:20098-105). Cell Proliferation may also be examined using [³H]-thymidine incorporation (Chen, J., 1996, *Oncogene* 13:1395-403; Jeoung, J., 1995, *J. Biol. Chem.* 270:18367-73). This assay allows for quantitative characterization of S-phase DNA syntheses. In this assay, cells synthesizing DNA will incorporate [³H]-thymidine into newly synthesized DNA. Incorporation can then be measured by standard techniques such as by counting of radioisotope in a scintillation counter (e.g., Beckman LS 3800 Liquid Scintillation Counter). Another proliferation assay uses the dye Alamar Blue (available from Biosource International), which fluoresces when reduced in living cells and provides an indirect measurement of cell number (Voytik-Harbin SL *et al.*, 1998, *In Vitro Cell Dev Biol Anim* 34:239-46). Yet another proliferation assay, the MTS assay, is based on in vitro cytotoxicity assessment of industrial chemicals, and uses the soluble tetrazolium salt, MTS. MTS assays are commercially available, for example, the Promega CellTiter 96[®] Aqueous Non-Radioactive Cell Proliferation Assay (Cat.# G5421).

Cell proliferation may also be assayed by colony formation in soft agar (Sambrook et al., Molecular Cloning, Cold Spring Harbor (1989)). For example, cells transformed with MBCAT are seeded in soft agar plates, and colonies are measured and counted after two weeks incubation.

5 Cell proliferation may also be assayed by measuring ATP levels as indicator of metabolically active cells. Such assays are commercially available, for example Cell Titer-Glo™, which is a luminescent homogeneous assay available from Promega.

Involvement of a gene in the cell cycle may be assayed by flow cytometry (Gray JW et al. (1986) Int J Radiat Biol Relat Stud Phys Chem Med 49:237-55). Cells
10 transfected with an MBCAT may be stained with propidium iodide and evaluated in a flow cytometer (available from Becton Dickinson), which indicates accumulation of cells in different stages of the cell cycle.

Accordingly, a cell proliferation or cell cycle assay system may comprise a cell that expresses an MBCAT, and that optionally has defective beta-catenin function (e.g.
15 beta-catenin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the assay system and changes in cell proliferation or cell cycle relative to controls where no test agent is added, identify candidate beta-catenin modulating agents. In some embodiments of the invention, the cell proliferation or cell cycle assay may be used as a secondary assay to test a candidate beta-catenin modulating agents that is
20 initially identified using another assay system such as a cell-free assay system. A cell proliferation assay may also be used to test whether MBCAT function plays a direct role in cell proliferation or cell cycle. For example, a cell proliferation or cell cycle assay may be performed on cells that over- or under-express MBCAT relative to wild type cells. Differences in proliferation or cell cycle compared to wild type cells suggests that the
25 MBCAT plays a direct role in cell proliferation or cell cycle.

Angiogenesis. Angiogenesis may be assayed using various human endothelial cell systems, such as umbilical vein, coronary artery, or dermal cells. Suitable assays include Alamar Blue based assays (available from Biosource International) to measure
30 proliferation; migration assays using fluorescent molecules, such as the use of Becton Dickinson Falcon HTS FluoroBlock cell culture inserts to measure migration of cells through membranes in presence or absence of angiogenesis enhancer or suppressors; and tubule formation assays based on the formation of tubular structures by endothelial cells on Matrigel® (Becton Dickinson). Accordingly, an angiogenesis assay system may

comprise a cell that expresses an MBCAT, and that optionally has defective beta-catenin function (e.g. beta-catenin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the angiogenesis assay system and changes in angiogenesis relative to controls where no test agent is added, identify candidate beta-catenin modulating agents. In some embodiments of the invention, the angiogenesis assay may be used as a secondary assay to test a candidate beta-catenin modulating agents that is initially identified using another assay system. An angiogenesis assay may also be used to test whether MBCAT function plays a direct role in cell proliferation. For example, an angiogenesis assay may be performed on cells that over- or under-express MBCAT relative to wild type cells. Differences in angiogenesis compared to wild type cells suggests that the MBCAT plays a direct role in angiogenesis. U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434, among others, describe various angiogenesis assays.

Hypoxic induction. The alpha subunit of the transcription factor, hypoxia inducible factor-1 (HIF-1), is upregulated in tumor cells following exposure to hypoxia in vitro. Under hypoxic conditions, HIF-1 stimulates the expression of genes known to be important in tumour cell survival, such as those encoding glycolytic enzymes and VEGF. Induction of such genes by hypoxic conditions may be assayed by growing cells transfected with MBCAT in hypoxic conditions (such as with 0.1% O₂, 5% CO₂, and balance N₂, generated in a Napco 7001 incubator (Precision Scientific)) and normoxic conditions, followed by assessment of gene activity or expression by Taqman®. For example, a hypoxic induction assay system may comprise a cell that expresses an MBCAT, and that optionally has defective beta-catenin function (e.g. beta-catenin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the hypoxic induction assay system and changes in hypoxic response relative to controls where no test agent is added, identify candidate beta-catenin modulating agents. In some embodiments of the invention, the hypoxic induction assay may be used as a secondary assay to test a candidate beta-catenin modulating agents that is initially identified using another assay system. A hypoxic induction assay may also be used to test whether MBCAT function plays a direct role in the hypoxic response. For example, a hypoxic induction assay may be performed on cells that over- or under-express MBCAT relative to wild type cells. Differences in hypoxic response compared to wild type cells suggests that the MBCAT plays a direct role in hypoxic induction.

Cell adhesion. Cell adhesion assays measure adhesion of cells to purified adhesion proteins, or adhesion of cells to each other, in presence or absence of candidate modulating agents. Cell-protein adhesion assays measure the ability of agents to modulate the adhesion of cells to purified proteins. For example, recombinant proteins are produced, diluted to 2.5g/mL in PBS, and used to coat the wells of a microtiter plate. The wells used for negative control are not coated. Coated wells are then washed, blocked with 1% BSA, and washed again. Compounds are diluted to 2× final test concentration and added to the blocked, coated wells. Cells are then added to the wells, and the unbound cells are washed off. Retained cells are labeled directly on the plate by adding a membrane-permeable fluorescent dye, such as calcein-AM, and the signal is quantified in a fluorescent microplate reader.

Cell-cell adhesion assays measure the ability of agents to modulate binding of cell adhesion proteins with their native ligands. These assays use cells that naturally or recombinantly express the adhesion protein of choice. In an exemplary assay, cells expressing the cell adhesion protein are plated in wells of a multiwell plate. Cells expressing the ligand are labeled with a membrane-permeable fluorescent dye, such as BCECF, and allowed to adhere to the monolayers in the presence of candidate agents. Unbound cells are washed off, and bound cells are detected using a fluorescence plate reader.

High-throughput cell adhesion assays have also been described. In one such assay, small molecule ligands and peptides are bound to the surface of microscope slides using a microarray spotter, intact cells are then contacted with the slides, and unbound cells are washed off. In this assay, not only the binding specificity of the peptides and modulators against cell lines are determined, but also the functional cell signaling of attached cells using immunofluorescence techniques in situ on the microchip is measured (Falsey JR et al., Bioconjug Chem. 2001 May-Jun;12(3):346-53).

Tubulogenesis. Tubulogenesis assays monitor the ability of cultured cells, generally endothelial cells, to form tubular structures on a matrix substrate, which generally simulates the environment of the extracellular matrix. Exemplary substrates include MatrigelTM (Becton Dickinson), an extract of basement membrane proteins containing laminin, collagen IV, and heparin sulfate proteoglycan, which is liquid at 4° C and forms a solid gel at 37° C. Other suitable matrices comprise extracellular components such as collagen, fibronectin, and/or fibrin. Cells are stimulated with a pro-angiogenic

stimulant, and their ability to form tubules is detected by imaging. Tubules can generally be detected after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Tube formation assays are well known in the art (e.g., Jones MK et al., 1999, Nature Medicine 5:1418-1423). These assays have traditionally involved

5 stimulation with serum or with the growth factors FGF or VEGF. Serum represents an undefined source of growth factors. In a preferred embodiment, the assay is performed with cells cultured in serum free medium, in order to control which process or pathway a candidate agent modulates. Moreover, we have found that different target genes respond differently to stimulation with different pro-angiogenic agents, including inflammatory

10 angiogenic factors such as TNF- α . Thus, in a further preferred embodiment, a tubulogenesis assay system comprises testing an MBCAT's response to a variety of factors, such as FGF, VEGF, phorbol myristate acetate (PMA), TNF- α , ephrin, etc.

Cell Migration. An invasion/migration assay (also called a migration assay) tests

15 the ability of cells to overcome a physical barrier and to migrate towards pro-angiogenic signals. Migration assays are known in the art (e.g., Paik JH et al., 2001, J Biol Chem 276:11830-11837). In a typical experimental set-up, cultured endothelial cells are seeded onto a matrix-coated porous lamina, with pore sizes generally smaller than typical cell size. The matrix generally simulates the environment of the extracellular matrix, as

20 described above. The lamina is typically a membrane, such as the transwell polycarbonate membrane (Corning Costar Corporation, Cambridge, MA), and is generally part of an upper chamber that is in fluid contact with a lower chamber containing pro-angiogenic stimuli. Migration is generally assayed after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Migration is assessed as the number of

25 cells that crossed the lamina, and may be detected by staining cells with hemotoxylin solution (VWR Scientific, South San Francisco, CA), or by any other method for determining cell number. In another exemplary set up, cells are fluorescently labeled and migration is detected using fluorescent readings, for instance using the Falcon HTS FluoroBlok (Becton Dickinson). While some migration is observed in the absence of

30 stimulus, migration is greatly increased in response to pro-angiogenic factors. As described above, a preferred assay system for migration/invasion assays comprises testing an MBCAT's response to a variety of pro-angiogenic factors, including tumor angiogenic and inflammatory angiogenic agents, and culturing the cells in serum free medium.

Sprouting assay. A sprouting assay is a three-dimensional *in vitro* angiogenesis assay that uses a cell-number defined spheroid aggregation of endothelial cells ("spheroid"), embedded in a collagen gel-based matrix. The spheroid can serve as a starting point for the sprouting of capillary-like structures by invasion into the extracellular matrix (termed "cell sprouting") and the subsequent formation of complex anastomosing networks (Korff and Augustin, 1999, J Cell Sci 112:3249-58). In an exemplary experimental set-up, spheroids are prepared by pipetting 400 human umbilical vein endothelial cells into individual wells of a nonadhesive 96-well plates to allow overnight spheroidal aggregation (Korff and Augustin: J Cell Biol 143: 1341-52, 1998). Spheroids are harvested and seeded in 900 μ l of methocel-collagen solution and pipetted into individual wells of a 24 well plate to allow collagen gel polymerization. Test agents are added after 30 min by pipetting 100 μ l of 10-fold concentrated working dilution of the test substances on top of the gel. Plates are incubated at 37°C for 24h. Dishes are fixed at the end of the experimental incubation period by addition of paraformaldehyde. Sprouting intensity of endothelial cells can be quantitated by an automated image analysis system to determine the cumulative sprout length per spheroid.

Primary assays for antibody modulators

For antibody modulators, appropriate primary assays test is a binding assay that tests the antibody's affinity to and specificity for the MBCAT protein. Methods for testing antibody affinity and specificity are well known in the art (Harlow and Lane, 1988, 1999, *supra*). The enzyme-linked immunosorbant assay (ELISA) is a preferred method for detecting MBCAT-specific antibodies; others include FACS assays, radioimmunoassays, and fluorescent assays.

In some cases, screening assays described for small molecule modulators may also be used to test antibody modulators.

Primary assays for nucleic acid modulators

For nucleic acid modulators, primary assays may test the ability of the nucleic acid modulator to inhibit or enhance MBCAT gene expression, preferably mRNA expression. In general, expression analysis comprises comparing MBCAT expression in like populations of cells (*e.g.*, two pools of cells that endogenously or recombinantly express MBCAT) in the presence and absence of the nucleic acid modulator. Methods for analyzing mRNA and protein expression are well known in the art. For instance, Northern

blotting, slot blotting, ribonuclease protection, quantitative RT-PCR (*e.g.*, using the TaqMan®, PE Applied Biosystems), or microarray analysis may be used to confirm that MBCAT mRNA expression is reduced in cells treated with the nucleic acid modulator (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm DH and Guiseppi-Elie, A Curr Opin Biotechnol 2001, 12:41-47). Protein expression may also be monitored. Proteins are most commonly detected with specific antibodies or antisera directed against either the MBCAT protein or specific peptides. A variety of means including Western blotting, ELISA, or in situ detection, are available (Harlow E and Lane D, 1988 and 1999, *supra*).

In some cases, screening assays described for small molecule modulators, particularly in assay systems that involve MBCAT mRNA expression, may also be used to test nucleic acid modulators.

15 **Secondary Assays**

Secondary assays may be used to further assess the activity of MBCAT-modulating agent identified by any of the above methods to confirm that the modulating agent affects MBCAT in a manner relevant to the beta-catenin pathway. As used herein, MBCAT-modulating agents encompass candidate clinical compounds or other agents derived from previously identified modulating agent. Secondary assays can also be used to test the activity of a modulating agent on a particular genetic or biochemical pathway or to test the specificity of the modulating agent's interaction with MBCAT.

Secondary assays generally compare like populations of cells or animals (*e.g.*, two pools of cells or animals that endogenously or recombinantly express MBCAT) in the presence and absence of the candidate modulator. In general, such assays test whether treatment of cells or animals with a candidate MBCAT-modulating agent results in changes in the beta-catenin pathway in comparison to untreated (or mock- or placebo-treated) cells or animals. Certain assays use "sensitized genetic backgrounds", which, as used herein, describe cells or animals engineered for altered expression of genes in the beta-catenin or interacting pathways.

Cell-based assays

Cell based assays may detect endogenous beta-catenin pathway activity or may rely on recombinant expression of beta-catenin pathway components. Any of the

aforementioned assays may be used in this cell-based format. Candidate modulators are typically added to the cell media but may also be injected into cells or delivered by any other efficacious means.

5 *Animal Assays*

A variety of non-human animal models of normal or defective beta-catenin pathway may be used to test candidate MBCAT modulators. Models for defective beta-catenin pathway typically use genetically modified animals that have been engineered to mis-express (*e.g.*, over-express or lack expression in) genes involved in the beta-catenin
10 pathway. Assays generally require systemic delivery of the candidate modulators, such as by oral administration, injection, etc.

In a preferred embodiment, beta-catenin pathway activity is assessed by monitoring neovascularization and angiogenesis. Animal models with defective and normal beta-catenin are used to test the candidate modulator's affect on MBCAT in
15 Matrigel® assays. Matrigel® is an extract of basement membrane proteins, and is composed primarily of laminin, collagen IV, and heparin sulfate proteoglycan. It is provided as a sterile liquid at 4° C, but rapidly forms a solid gel at 37° C. Liquid Matrigel® is mixed with various angiogenic agents, such as bFGF and VEGF, or with human tumor cells which over-express the MBCAT. The mixture is then injected
20 subcutaneously(SC) into female athymic nude mice (Taconic, Germantown, NY) to support an intense vascular response. Mice with Matrigel® pellets may be dosed via oral (PO), intraperitoneal (IP), or intravenous (IV) routes with the candidate modulator. Mice are euthanized 5 - 12 days post-injection, and the Matrigel® pellet is harvested for hemoglobin analysis (Sigma plasma hemoglobin kit). Hemoglobin content of the gel is
25 found to correlate the degree of neovascularization in the gel.

In another preferred embodiment, the effect of the candidate modulator on MBCAT is assessed via tumorigenicity assays. Tumor xenograft assays are known in the art (see, *e.g.*, Ogawa K et al., 2000, *Oncogene* 19:6043-6052). Xenografts are typically implanted SC into female athymic mice, 6-7 week old, as single cell suspensions either
30 from a pre-existing tumor or from *in vitro* culture. The tumors which express the MBCAT endogenously are injected in the flank, 1×10^5 to 1×10^7 cells per mouse in a volume of 100 μ L using a 27gauge needle. Mice are then ear tagged and tumors are measured twice weekly. Candidate modulator treatment is initiated on the day the mean tumor weight reaches 100 mg. Candidate modulator is delivered IV, SC, IP, or PO by bolus

administration. Depending upon the pharmacokinetics of each unique candidate modulator, dosing can be performed multiple times per day. The tumor weight is assessed by measuring perpendicular diameters with a caliper and calculated by multiplying the measurements of diameters in two dimensions. At the end of the experiment, the excised
5 tumors maybe utilized for biomarker identification or further analyses. For immunohistochemistry staining, xenograft tumors are fixed in 4% paraformaldehyde, 0.1M phosphate, pH 7.2, for 6 hours at 4°C, immersed in 30% sucrose in PBS, and rapidly frozen in isopentane cooled with liquid nitrogen.

In another preferred embodiment, tumorigenicity is monitored using a hollow fiber
10 assay, which is described in U.S. Pat No. US 5,698,413. Briefly, the method comprises implanting into a laboratory animal a biocompatible, semi-permeable encapsulation device containing target cells, treating the laboratory animal with a candidate modulating agent, and evaluating the target cells for reaction to the candidate modulator. Implanted cells are generally human cells from a pre-existing tumor or a tumor cell line. After an appropriate
15 period of time, generally around six days, the implanted samples are harvested for evaluation of the candidate modulator. Tumorigenicity and modulator efficacy may be evaluated by assaying the quantity of viable cells present in the macrocapsule, which can be determined by tests known in the art, for example, MTT dye conversion assay, neutral red dye uptake, trypan blue staining, viable cell counts, the number of colonies formed in
20 soft agar, the capacity of the cells to recover and replicate in vitro, etc.

In another preferred embodiment, a tumorigenicity assay use a transgenic animal, usually a mouse, carrying a dominant oncogene or tumor suppressor gene knockout under the control of tissue specific regulatory sequences; these assays are generally referred to as transgenic tumor assays. In a preferred application, tumor development in the transgenic
25 model is well characterized or is controlled. In an exemplary model, the "RIP1-Tag2" transgene, comprising the SV40 large T-antigen oncogene under control of the insulin gene regulatory regions is expressed in pancreatic beta cells and results in islet cell carcinomas (Hanahan D, 1985, Nature 315:115-122; Parangi S et al, 1996, Proc Natl Acad Sci USA 93: 2002-2007; Bergers G et al, 1999, Science 284:808-812). An "angiogenic
30 switch," occurs at approximately five weeks, as normally quiescent capillaries in a subset of hyperproliferative islets become angiogenic. The RIP1-TAG2 mice die by age 14 weeks. Candidate modulators may be administered at a variety of stages, including just prior to the angiogenic switch (e.g., for a model of tumor prevention), during the growth of small tumors (e.g., for a model of intervention), or during the growth of large and/or

invasive tumors (e.g., for a model of regression). Tumorigenicity and modulator efficacy can be evaluating life-span extension and/or tumor characteristics, including number of tumors, tumor size, tumor morphology, vessel density, apoptotic index, etc.

5 **Diagnostic and therapeutic uses**

Specific MBCAT-modulating agents are useful in a variety of diagnostic and therapeutic applications where disease or disease prognosis is related to defects in the beta-catenin pathway, such as angiogenic, apoptotic, or cell proliferation disorders. Accordingly, the invention also provides methods for modulating the beta-catenin pathway
10 in a cell, preferably a cell pre-determined to have defective or impaired beta-catenin function (e.g. due to overexpression, underexpression, or misexpression of beta-catenin, or due to gene mutations), comprising the step of administering an agent to the cell that specifically modulates MBCAT activity. Preferably, the modulating agent produces a detectable phenotypic change in the cell indicating that the beta-catenin function is
15 restored. The phrase "function is restored", and equivalents, as used herein, means that the desired phenotype is achieved, or is brought closer to normal compared to untreated cells. For example, with restored beta-catenin function, cell proliferation and/or progression through cell cycle may normalize, or be brought closer to normal relative to untreated cells. The invention also provides methods for treating disorders or disease
20 associated with impaired beta-catenin function by administering a therapeutically effective amount of an MBCAT -modulating agent that modulates the beta-catenin pathway. The invention further provides methods for modulating MBCAT function in a cell, preferably a cell pre-determined to have defective or impaired MBCAT function, by administering an MBCAT -modulating agent. Additionally, the invention provides a method for treating
25 disorders or disease associated with impaired MBCAT function by administering a therapeutically effective amount of an MBCAT -modulating agent.

The discovery that MBCAT is implicated in beta-catenin pathway provides for a variety of methods that can be employed for the diagnostic and prognostic evaluation of diseases and disorders involving defects in the beta-catenin pathway and for the
30 identification of subjects having a predisposition to such diseases and disorders.

Various expression analysis methods can be used to diagnose whether MBCAT expression occurs in a particular sample, including Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR, and microarray analysis. (e.g., Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc.,

chapter 4; Freeman WM *et al.*, *Biotechniques* (1999) 26:112-125; Kallioniemi OP, *Ann Med* 2001, 33:142-147; Blohm and Guiseppi-Elie, *Curr Opin Biotechnol* 2001, 12:41-47).

Tissues having a disease or disorder implicating defective beta-catenin signaling that express an MBCAT, are identified as amenable to treatment with an MBCAT modulating agent. In a preferred application, the beta-catenin defective tissue overexpresses an MBCAT relative to normal tissue. For example, a Northern blot analysis of mRNA from tumor and normal cell lines, or from tumor and matching normal tissue samples from the same patient, using full or partial MBCAT cDNA sequences as probes, can determine whether particular tumors express or overexpress MBCAT. Alternatively, the TaqMan® is used for quantitative RT-PCR analysis of MBCAT expression in cell lines, normal tissues and tumor samples (PE Applied Biosystems).

Various other diagnostic methods may be performed, for example, utilizing reagents such as the MBCAT oligonucleotides, and antibodies directed against an MBCAT, as described above for: (1) the detection of the presence of MBCAT gene mutations, or the detection of either over- or under-expression of MBCAT mRNA relative to the non-disorder state; (2) the detection of either an over- or an under-abundance of MBCAT gene product relative to the non-disorder state; and (3) the detection of perturbations or abnormalities in the signal transduction pathway mediated by MBCAT.

Kits for detecting expression of MBCAT in various samples, comprising at least one antibody specific to MBCAT, all reagents and/or devices suitable for the detection of antibodies, the immobilization of antibodies, and the like, and instructions for using such kits in diagnosis or therapy are also provided.

Thus, in a specific embodiment, the invention is drawn to a method for diagnosing a disease or disorder in a patient that is associated with alterations in MBCAT expression, the method comprising: a) obtaining a biological sample from the patient; b) contacting the sample with a probe for MBCAT expression; c) comparing results from step (b) with a control; and d) determining whether step (c) indicates a likelihood of the disease or disorder. Preferably, the disease is cancer. The probe may be either DNA or protein, including an antibody.

EXAMPLES

The following experimental section and examples are offered by way of illustration and not by way of limitation.

I. C. elegans beta-catenin screen

The identification of mutants that suppress the cell adhesion defect of beta-catenin may lead to unique therapeutic targets that inhibit cell migration or metastasis. *hmp-2* was initially identified in an EMS screen for defects in body elongation during embryonic morphogenesis (see Costa *et al.*, (1998) The Journal of Cell Biology 1998, 141: 297-308). The loss of function allele *hmp-2 (zu364)* exhibits 99% embryonic lethality, with mutant embryos arresting during elongation and abnormal bulges forming on the dorsal side. About 1% of these embryos hatch to form viable lumpy larvae. The reduction of function allele *hmp-2 (qm39)* yields viable larvae with a characteristic lumpy appearance. When grown at 15°C, approximately 92% (SD 3.9) of the L1 larvae show this lumpy phenotype, with the penetrance of the phenotype decreasing as the animals molt and move through successive larval stages. For this screen, *hmp-2 (qm39)* worms were soaked at 15°C in double stranded RNA (dsRNA) at the L4 larval stage and the progeny were scored as L1 larvae for modification of the adhesion defect. The screen protocol is described below.

15

1) *hmp-2 (qm39)* animals were bleached and hatched on peptone free agarose plates to produce a synchronous population. Starved L1s were transferred to 10x peptone plates seeded with 750 µl OP50 (25% w/v in TB) and allowed to develop to the L4 larval stage.

20

2) dsRNA was dispensed in 6 µl aliquots into 96 well round bottom plates (Nunc #262162). L4 animals were collected by suspension in M9 buffer, washed 2x with M9 to remove any excess OP50, and dispensed in 2 µl aliquots into the RNA to a total worm density of 75-100 worms per well. As a control, multiple wells contained only RNA resuspension buffer (1x IM buffer).

25

3) Animals were soaked in dsRNA at 15°C for 24 hours.

4) Following dsRNA soaking, the animals were fed in the wells by addition of 25µl liquid NGM + 3% OP50. The animals were kept at 15°C and allowed to become gravid and lay progeny in the wells, which took approximately 72 hours. Food levels were monitored visually during maturation and more was added as needed.

30

5) Following maturation, animals from each well were plated onto individual 6cm peptone free agarose plates and placed at 15°C overnight.

6) Animals on each plate were scored visually under the dissecting microscope for modification of the lumpy phenotype. Scoring was performed

qualitatively, with an increase in dead embryos scored as enhancement and an increase in wild type appearing animals scored as suppression of the defect.

7) Retests of interesting suppressor candidates followed the same protocol as the primary screen with certain modifications: several retests were performed for each suppressor, retested candidates were encoded so that they could be scored blindly, and retested candidates were scored quantitatively. Each plate was scored by counting 100 total objects. An object was defined as either an embryo or an L1 stage larva. Each object was scored as one of the following: a wildtype appearing animal, a lumpy appearing animal, or an unhatched embryo. Scores were represented as the percentage of wildtype appearing animals relative to all objects scored. Wildtype animals were defined as L1 larvae with smooth cuticles that did not have any sort of lumpy body morphology.

A confirmed suppressor was one that was ≥ 2 standard deviations away from the mean of the controls for at least 3 of the four retest experiments.

II. Analysis of Table 1

BLAST analysis (Altschul et al., *supra*) was employed to identify orthologs of *C. elegans* modifiers. The columns "MBCAT symbol", and "MBCAT name aliases" provide a symbol and the known name abbreviations for the Targets, where available, from Genbank. "MBCAT RefSeq_NA or GI_NA", "MBCAT GI_AA", "MBCAT NAME", and "MBCAT Description" provide the reference DNA sequences for the MBCATs as available from National Center for Biology Information (NCBI), MBCAT protein Genbank identifier number (GI#), MBCAT name, and MBCAT description, all available from Genbank, respectively. The length of each amino acid is in the "MBCAT Protein Length" column.

Names and Protein sequences of *C. elegans* modifiers of beta-catenin from screen (Example I), are represented in the "Modifier Name" and "Modifier GI_AA" column by GI#, respectively.

Table 1

MBCAT symbol	MBCAT name aliases	MBCAT RefSeq_N A or GI_NA	NA SEQ ID NO	MBCAT GI_AA or RefSeq_A A	AA SEQ ID NO	MBCAT name	MBCA T descript ion	MBCAT protein length	Modifi er name	Modifier GI_AA
LOC130600	LOC130600 similar to ribosomal protein L23, cytosolic [validated] - rat na	XM_0657 96	1	17444877	8	similar to ribosomal protein L23, cytosolic [validated] - rat	na	113	rpl-23	17554750
LOC139436	LOC139436 similar to ribosomal protein L23, cytosolic [validated] - rat na	XM_0666 98	2	17485849	9	similar to ribosomal protein L23, cytosolic [validated] - rat	na	200	rpl-23	17554750
LOC222901	LOC222901 similar to ribosomal protein L23, cytosolic [validated] - rat na	XM_1672 75	3	20539732	10	similar to ribosomal protein L23, cytosolic [validated] - rat	na	107	rpl-23	17554750
LOC285850	LOC285850 similar to ribosomal protein L23, cytosolic [validated] - rat na	XM_2083 58	4	27498549	11	similar to ribosomal protein L23, cytosolic [validated] - rat	na	79	rpl-23	17554750
RPL23	RPL23 rpL17 60S ribosomal protein L23 ribosomal protein L23	NM_0009 78	5	4506605	12	ribosomal protein L23	structur al constitu ent of ribosom e	140	rpl-23	17554750

STK3	KRS1 MST2 serine/threonine kinase 3 (Ste20, yeast homolog) serine/threonine kinase 3 (STE20 homolog, yeast) STK3	NM_006281	6	5454094	13	serine/threonine kinase 3 (STE20 homolog, yeast)	protein kinase; protein serine/threonine kinase; protein serine/threonine kinase	491	XE961	8489865
STK4	KRS2 MST1 serine/threonine kinase 4 (Ste20, yeast homolog) serine/threonine kinase 4 STK4	NM_006282	7	5454096	14	serine/threonine kinase 4	protein serine/threonine kinase; protein serine/threonine kinase	487	XE961	8489865

III. High-Throughput In Vitro Fluorescence Polarization Assay

Fluorescently-labeled MBCAT peptide/substrate are added to each well of a 96-well microtiter plate, along with a test agent in a test buffer (10 mM HEPES, 10 mM NaCl, 6 mM magnesium chloride, pH 7.6). Changes in fluorescence polarization, determined by using a Fluorolite FPM-2 Fluorescence Polarization Microtiter System (Dynatech Laboratories, Inc), relative to control values indicates the test compound is a candidate modifier of MBCAT activity.

IV. High-Throughput In Vitro Binding Assay.

³³P-labeled MBCAT peptide is added in an assay buffer (100 mM KCl, 20 mM HEPES pH 7.6, 1 mM MgCl₂, 1% glycerol, 0.5% NP-40, 50 mM beta-mercaptoethanol, 1 mg/ml BSA, cocktail of protease inhibitors) along with a test agent to the wells of a Neutralite-avidin coated assay plate and incubated at 25°C for 1 hour. Biotinylated substrate is then added to each well and incubated for 1 hour. Reactions are stopped by washing with PBS, and counted in a scintillation counter. Test agents that cause a

difference in activity relative to control without test agent are identified as candidate beta-catenin modulating agents.

V. Immunoprecipitations and Immunoblotting

5 For coprecipitation of transfected proteins, 3×10^6 appropriate recombinant cells containing the MBCAT proteins are plated on 10-cm dishes and transfected on the following day with expression constructs. The total amount of DNA is kept constant in each transfection by adding empty vector. After 24 h, cells are collected, washed once with phosphate-buffered saline and lysed for 20 min on ice in 1 ml of lysis buffer
10 containing 50 mM Hepes, pH 7.9, 250 mM NaCl, 20 mM -glycerophosphate, 1 mM sodium orthovanadate, 5 mM p-nitrophenyl phosphate, 2 mM dithiothreitol, protease inhibitors (complete, Roche Molecular Biochemicals), and 1% Nonidet P-40. Cellular debris is removed by centrifugation twice at $15,000 \times g$ for 15 min. The cell lysate is incubated with 25 μ l of M2 beads (Sigma) for 2 h at 4 °C with gentle rocking.

15 After extensive washing with lysis buffer, proteins bound to the beads are solubilized by boiling in SDS sample buffer, fractionated by SDS-polyacrylamide gel electrophoresis, transferred to polyvinylidene difluoride membrane and blotted with the indicated antibodies. The reactive bands are visualized with horseradish peroxidase coupled to the appropriate secondary antibodies and the enhanced chemiluminescence
20 (ECL) Western blotting detection system (Amersham Pharmacia Biotech).

VI. Kinase assay

A purified or partially purified MBCAT is diluted in a suitable reaction buffer, e.g., 50 mM Hepes, pH 7.5, containing magnesium chloride or manganese chloride (1-20 mM)
25 and a peptide or polypeptide substrate, such as myelin basic protein or casein (1-10 μ g/ml). The final concentration of the kinase is 1-20 nM. The enzyme reaction is conducted in microtiter plates to facilitate optimization of reaction conditions by increasing assay throughput. A 96-well microtiter plate is employed using a final volume 30-100 μ l. The reaction is initiated by the addition of ^{33}P -gamma-ATP (0.5 μ Ci/ml) and
30 incubated for 0.5 to 3 hours at room temperature. Negative controls are provided by the addition of EDTA, which chelates the divalent cation (Mg^{2+} or Mn^{2+}) required for enzymatic activity. Following the incubation, the enzyme reaction is quenched using EDTA. Samples of the reaction are transferred to a 96-well glass fiber filter plate (MultiScreen, Millipore). The filters are subsequently washed with phosphate-buffered

saline, dilute phosphoric acid (0.5%) or other suitable medium to remove excess radiolabeled ATP. Scintillation cocktail is added to the filter plate and the incorporated radioactivity is quantitated by scintillation counting (Wallac/Perkin Elmer). Activity is defined by the amount of radioactivity detected following subtraction of the negative control reaction value (EDTA quench).

VII. Expression analysis

All cell lines used in the following experiments are NCI (National Cancer Institute) lines, and are available from ATCC (American Type Culture Collection, Manassas, VA 20110-2209). Normal and tumor tissues are obtained from Impath, UC Davis, Clontech, Stratagene, Ardaïs, Genome Collaborative, and Ambion.

TaqMan® analysis is used to assess expression levels of the disclosed genes in various samples.

RNA is extracted from each tissue sample using Qiagen (Valencia, CA) RNeasy kits, following manufacturer's protocols, to a final concentration of 50ng/μl. Single stranded cDNA is then synthesized by reverse transcribing the RNA samples using random hexamers and 500ng of total RNA per reaction, following protocol 4304965 of Applied Biosystems (Foster City, CA).

Primers for expression analysis using TaqMan® assay (Applied Biosystems, Foster City, CA) are prepared according to the TaqMan® protocols, and the following criteria: a) primer pairs are designed to span introns to eliminate genomic contamination, and b) each primer pair produced only one product. Expression analysis is performed using a 7900HT instrument.

TaqMan® reactions are carried out following manufacturer's protocols, in 25 μl total volume for 96-well plates and 10 μl total volume for 384-well plates, using 300nM primer and 250 nM probe, and approximately 25ng of cDNA. The standard curve for result analysis is prepared using a universal pool of human cDNA samples, which is a mixture of cDNAs from a wide variety of tissues so that the chance that a target will be present in appreciable amounts is good. The raw data are normalized using 18S rRNA (universally expressed in all tissues and cells).

For each expression analysis, tumor tissue samples are compared with matched normal tissues from the same patient. A gene is considered overexpressed in a tumor when the level of expression of the gene is 2 fold or higher in the tumor compared with its matched normal sample. In cases where normal tissue is not available, a universal pool of

cDNA samples is used instead. In these cases, a gene is considered overexpressed in a tumor sample when the difference of expression levels between a tumor sample and the average of all normal samples from the same tissue type is greater than 2 times the standard deviation of all normal samples (i.e., $\text{Tumor} - \text{average}(\text{all normal samples}) > 2 \times \text{STDEV}(\text{all normal samples})$).

A modulator identified by an assay described herein can be further validated for therapeutic effect by administration to a tumor in which the gene is overexpressed. A decrease in tumor growth confirms therapeutic utility of the modulator. Prior to treating a patient with the modulator, the likelihood that the patient will respond to treatment can be diagnosed by obtaining a tumor sample from the patient, and assaying for expression of the gene targeted by the modulator. The expression data for the gene(s) can also be used as a diagnostic marker for disease progression. The assay can be performed by expression analysis as described above, by antibody directed to the gene target, or by any other available detection method.

15

WHAT IS CLAIMED IS:

1. A method of identifying a candidate beta-catenin pathway modulating agent, said method comprising the steps of:
 - 5 (a) providing an assay system comprising a MBCAT polypeptide or nucleic acid;
 - (b) contacting the assay system with a test agent under conditions whereby, but for the presence of the test agent, the system provides a reference activity; and
 - (c) detecting a test agent-biased activity of the assay system, wherein a difference between the test agent-biased activity and the reference activity identifies the test agent as
 - 10 a candidate beta-catenin pathway modulating agent.
2. The method of Claim 1 wherein the assay system comprises cultured cells that express the MBCAT polypeptide.
- 15 3. The method of Claim 2 wherein the cultured cells additionally have defective beta-catenin function.
4. The method of Claim 1 wherein the assay system includes a screening assay comprising a MBCAT polypeptide, and the candidate test agent is a small molecule
- 20 modulator.
5. The method of Claim 4 wherein the assay is a binding assay.
6. The method of Claim 1 wherein the assay system is selected from the group consisting
- 25 of an apoptosis assay system, a cell proliferation assay system, an angiogenesis assay system, and a hypoxic induction assay system.
7. The method of Claim 1 wherein the assay system includes a binding assay comprising a MBCAT polypeptide and the candidate test agent is an antibody.
- 30 8. The method of Claim 1 wherein the assay system includes an expression assay comprising a MBCAT nucleic acid and the candidate test agent is a nucleic acid modulator.

9. The method of Claim 8 wherein the nucleic acid modulator is an antisense oligomer.
10. The method of Claim 8 wherein the nucleic acid modulator is a PMO.
- 5 11. The method of Claim 1 additionally comprising:
(d) administering the candidate beta-catenin pathway modulating agent identified in (c) to a model system comprising cells defective in beta-catenin function and, detecting a phenotypic change in the model system that indicates that the beta-catenin function is restored.
- 10 12. The method of Claim 11 wherein the model system is a mouse model with defective beta-catenin function.
13. A method for modulating a beta-catenin pathway of a cell comprising contacting a
15 cell defective in beta-catenin function with a candidate modulator that specifically binds to a MBCAT polypeptide, whereby beta-catenin function is restored.
14. The method of Claim 13 wherein the candidate modulator is administered to a
vertebrate animal predetermined to have a disease or disorder resulting from a defect in
20 beta-catenin function.
15. The method of Claim 13 wherein the candidate modulator is selected from the group consisting of an antibody and a small molecule.
- 25 16. The method of Claim 1, comprising the additional steps of:
(e) providing a secondary assay system comprising cultured cells or a non-human animal expressing MBCAT ,
(f) contacting the secondary assay system with the test agent of (b) or an agent derived therefrom under conditions whereby, but for the presence of the test agent or agent
30 derived therefrom, the system provides a reference activity; and
(g) detecting an agent-biased activity of the second assay system,
wherein a difference between the agent-biased activity and the reference activity of the second assay system confirms the test agent or agent derived therefrom as a candidate beta-catenin pathway modulating agent,

and wherein the second assay detects an agent-biased change in the beta-catenin pathway.

5 17. The method of Claim 16 wherein the secondary assay system comprises cultured cells.

18. The method of Claim 16 wherein the secondary assay system comprises a non-human animal.

10 19. The method of Claim 18 wherein the non-human animal mis-expresses a beta-catenin pathway gene.

15 20. A method of modulating beta-catenin pathway in a mammalian cell comprising contacting the cell with an agent that specifically binds a MBCAT polypeptide or nucleic acid.

21. The method of Claim 20 wherein the agent is administered to a mammalian animal predetermined to have a pathology associated with the beta-catenin pathway.

20 22. The method of Claim 20 wherein the agent is a small molecule modulator, a nucleic acid modulator, or an antibody.

23. A method for diagnosing a disease in a patient comprising:

- 25 (a) obtaining a biological sample from the patient;
(b) contacting the sample with a probe for MBCAT expression;
(c) comparing results from step (b) with a control;
(d) determining whether step (c) indicates a likelihood of disease.

30 24. The method of claim 23 wherein said disease is cancer.

SEQUENCE LISTING

<110> EXELIXIS, INC

<120> MBCATS AS MODIFIERS OF THE BETA-CATENIN PATHWAY AND METHODS OF USE

<130> EX04-015C-PC

<150> US 60/454,436

<151> 2003-03-13

<160> 14

<170> PatentIn version 3.2

<210> 1

<211> 987

<212> DNA

<213> Homo sapiens

<400> 1

```

atggccttgg gcaggggaacc agttgtggat ttcccggggg caggggttggg ggcagtcctt      60
ggaaccgggc ctagcccgtc tgggagcggg ggggtagaag tgaaaggctt aagtgtgggt      120
gcgggggactt gcagccgcac agacaaggca gaccgggact gggaagctgc ccggagcggc      180
tgctgcgact ctccgtccg acctagccgg ggcagccctc cctgccccgg gagaagacgg      240
cgagcaggag gacccgcaga gcccgcgccg gctgctccgg ccccgggggc actggcacca      300
tcgggctgtg gtcttttccc gatgcccgag ccgtggacgc ggccaccta gagcgggtccg      360
gcgaaggcag cgcagccggt gtcccaggga ccgagcaggg ccgagttttc ctggggagcc      420
tgggtccccg cggccgggcg gccgctggga ggtggaggag cctctccagc cccggccgca      480
gctcatcgtg agccaaccgc gccacctgcc ggccagatgc gggagtgtgc ggccccggag      540
gaagggcgag cggacaaaga ccgcaacccg caatccgcaa cccgcagag aggtgatgtt      600
gacgcgacgc gaggacgtgg tgggacctct ggtgcaaat tccgaatgtc cttgagtttc      660
ccggtaggag ctgcggtcaa ctgtgctgac aacaccggag ccaaaagcct gtatatcatc      720
tccacgaagg ggatcaaggg acggctgaac agacttcccg ctgctgatgt gggtgacatg      780
gtgatgacca cagtcaagaa aggcaaacca gagctcagaa aaagggtagc ttcagcagtg      840
gtcatttcac aacagaagtc atactggaga aaaggtggca tgtttcctta ttttgaagat      900
aatgcagggg tcatagcaaa caataaaggt gagatgaaag gttctgccat tacaggacca      960
gtagcaaaga agcgtgtaga cttgtga                                     987

```

<210> 2

<211> 603

<212> DNA

<213> Homo sapiens

<400> 2
 atgggtcatg agttgattac tgttgaagta ggtggcaagt acatgaacag tacaaatgca 60
 ctgggaagac tggggaagcc ccagatttca caagagacac tcacaagtct agcctctcac 120
 tccacatctg cttgcatggt aattgggcca tttcagggag ccaaagagga gttagggaag 180
 gcatggtcac caaaacttgc cttagaggac cacactttta agatattgaa gggaggacgt 240
 gatgggtctt ccagtatgaa attctggatt tccctgggtc ttctgttagg agtggaatca 300
 actgtgctga caacacagga gctgaaaaat ctgtgtatct tctcagtga ggggaatcaag 360
 ggatggctga acagactctc cgctgctggt gtgggtgatg tggatgatggc tacagttaag 420
 aaaggcaaac cacagcttag aaagaagata aaagatgaaa tgcttcttta ttttgaagat 480
 aatgtggaag ttataggaat taataaagggt aagataaaaa tttctgctat cacaggacag 540
 attgtgaagg aggggtgcaga attgtgcccc agtattagat acagctctgg tagcattgca 600
 tga 603

<210> 3
 <211> 633
 <212> DNA
 <213> Homo sapiens

<400> 3
 tttccagcgt tcaagatgtg gaagcgagga cgtgggtgggt cctctgggtg gaacttccgg 60
 atttctgtgg gtcttccggt aggagctgtg atcaactgtg ctgacaacac aggagccaaa 120
 aacctgtata tcatctccgt gaaggggatc aagggacggc tgaacagact tcccgtgct 180
 ggtgtgggtg acatggggat ggctacagtc aagaaaggga aactagagct cagaaaaaag 240
 tgctcaggaa ctctgggtcaa aggaacacca ctagaaaaca ctgaaggagg catgtcaggt 300
 ggttctgctt ctccatggct ccctacgatg aatgcctaga gagggatttg acatggcaac 360
 ctgatgccag cacatctgtc cacagctgcc attgcaagga tctaccttgg tgccctctaa 420
 agaaattctg aattttctga gaacaagagg tctagtccca tatgtggaag tcttttctgt 480
 atgtctctga gtagaaagtg agaattgata tgccaaggga tgagtaatga gcctcatctt 540
 gatgcagagt cccaagaac caacttttac tgaggagcat gcaggcaact ttatacccca 600
 tttgacaaca ttaaacccta aagaagaaaa tca 633

<210> 4
 <211> 482
 <212> DNA
 <213> Homo sapiens

<400> 4
 ctttttctt ctttccggct ttcaagatgt tgaagcgagg acgtgggtggg tcctctgggtg 60
 tgaaattcca gatttccttg ggtcttctgg taggagtgtg atcagttgtg ctgacaacac 120

```

aggagccaaa aacctgtatg tcatctccgt gaaaggggat caagggatgg ctgaacagac 180
ttcccactgc tgggtgtgggt gacacgggtga tggccacagt caagaaaggc aaaccagagc 240
tcagaaaaaa ggtacatcca gcagtgggta ttcgacaacg aaagtcatac cgcagaaaag 300
atggcggtgtt tctttatttt gaagataatg cagggggtcac agtgaacaat aaaggcgaga 360
tgaaagggttc tgccattaca ggaccagtag caaaggagtg tgcagacttg tggccctgga 420
ttgcggtccag tgctggcagc attgcatgat tctccagtat atttgtaaaa aataaaaaaa 480
aa 482

```

<210> 5
 <211> 493
 <212> DNA
 <213> Homo sapiens

```

<400> 5
cttttttctt ttttccggcg ttcaagatgt cgaagcgagg acgtgggtggg tcctctgggtg 60
cgaaattccg gatttccttg ggtcttccgg taggagctgt aatcaattgt gctgacaaca 120
caggagccaa aaacctgtat atcatctccg tgaaggggat caagggacgg ctgaacagac 180
ttcccgtctgc tgggtgtgggt gacatgggtga tggccacagt caagaaaggc aaaccagagc 240
tcagaaaaaa ggtacatcca gcagtgggtca ttcgacaacg aaagtcatac cgtagaaaag 300
atggcggtgtt tctttatttt gaagataatg caggagtcac agtgaacaat aaaggcgaga 360
tgaaagggttc tgccattaca ggaccagtag caaaggagtg tgcagacttg tggccccgga 420
ttgcatccaa tgctggcagc attgcatgat tctccagtat atttgtaaaa aataaaaaaa 480
aactaaaccc att 493

```

<210> 6
 <211> 2820
 <212> DNA
 <213> Homo sapiens

```

<400> 6
ccgcggagtt acgggaaagt tgggtccgagt tcccagagtt tccctctgtg gtgccctagg 60
cttcggccccg gtgccccggc tcctttcctc ctttcggcct tcgccgtcca ccagggtccct 120
ctctctgtcc cggccgccat ggagcagccg ccggcgcccta agagtaaact aaaaaagctg 180
agtgaagaca gtttgactaa gcagcctgaa gaagtttttg atgtattaga gaagcttgga 240
gaagggtctt atggaagtgt atttaaagca atacacaagg aatccgggtca agttgtcgca 300
attaaacaag tacctgttga atcagatctt caggaaataa tcaaagaaat ttccataatg 360
cagcaatgtg acagcccata tggtgtaaag tactatggca gttattttta gaatacagac 420
ctctggattg ttatggagta ctgtggcgct ggctctgtct cagacataat tagattacga 480

```

aacaagacat	taatagaaga	tgaaattgca	accattctta	aatctacatt	gaaaggacta	540
gaatatttgc	actttatgag	aaaaatacac	agagatataa	aagctggaaa	tattctcctc	600
aatacagaag	gacatgcaaa	attggcagat	tttggagtgg	ctggtcagtt	aacagataca	660
atggcaaaac	gcaatactgt	aataggaact	ccattttgga	tggctcctga	ggtgattcaa	720
gaaataggct	ataactgtgt	ggccgacatc	tggtcacctg	gcattacttc	tatagaaatg	780
gctgaaggaa	aacctcctta	tgctgatata	catccaatga	gggctatttt	tatgattccc	840
acaaatccac	caccaacatt	cagaaaagcca	gaactttggt	ccgatgatgt	caccgatgtt	900
gttaaaaagt	gtttggtgaa	gaatcctgag	cagagagcta	ctgcaacaca	acttttacag	960
catcctttta	tcaagaatgc	caaacctgta	tcaatattaa	gagacctgat	cacagaagct	1020
atggagatca	aagctaaaag	acatgacgaa	cagcaacgag	aattggaaga	ggaagaagaa	1080
aattcggatg	aagatgagct	ggattcccac	accatgggtga	agactagtgt	gggagagtgt	1140
ggcaccatgc	gggccacaag	cacgatgagt	gaaggggccc	agaccatgat	tgaacataat	1200
agcacgatgt	tggaatccga	cttggggacc	atggtgataa	acagtgagga	tgaggaagaa	1260
gaagatggaa	ctatgaaaag	aaatgcaacc	tcaccacaag	tacaaagacc	atctttcatg	1320
gactactttg	ataagcaaga	cttcaagaat	aagagtcacg	aaaactgtaa	tcagaacatg	1380
catgaaccct	tccctatgtc	caaaaacggt	tttcctgata	actggaaagt	tcctcaagat	1440
ggagactttg	actttttgaa	aaatctaagt	ttagaagaac	tacagatgcg	gttaaaagca	1500
ctggacccca	tgatggaacg	ggagatagaa	gaacttcgtc	agagatacac	tgcgaaaaga	1560
cagcccattc	tggtatgcat	ggatgcaaag	aaaagaaggc	agcaaaactt	ttgagtctaa	1620
tttcctctct	gtttttaact	attctggaga	ccaagaaacc	actaggaatt	gaaggaatat	1680
ttggatattt	ttaatcctaa	gattttgccc	tacaattagg	cagagggtcaa	aaagtgacaa	1740
tggtacatgc	ccaggtaaat	tcccaaaagg	cagaattgac	agttgtatct	gctgtgcatt	1800
cactctaaga	tgaggagaac	aaaagaagtg	tattctcttg	ttctgtcagc	tgcataccag	1860
taataaaaact	gttatgaaat	ggattttcaa	ggtctctaaa	ccttgaaaat	ccaaaggcta	1920
ttgttgcatc	gtacagcact	gaaagggctt	tatgttacaa	tattctttat	tcctatctag	1980
tatactaggc	tatttattgt	ccccttaggt	aaacttattt	atztatgcta	ttttggcttt	2040
gtttcatttt	ttaaggacaa	gatcaggata	gctttggtga	aggtaggggc	atattaatat	2100
gatgataatg	tgcaaccaat	ttatactttc	tgcagggagc	tatgggggtac	attccttgat	2160
ttccaggata	gtttttcaaa	taggaaagca	ataatggcag	tagttctcaa	atgggctagg	2220
ccttttttat	attgaagcaa	taattccatt	tttacccttt	gaaattttgt	ttttttgatt	2280
tttgatgttt	ggtacaaaata	gaactatata	tatttaggta	aaatagatct	atcgtgttta	2340

aaaccaaaga aatcaatgga acccttgac	aaaaaagtgt gataaatatt tttaaataaa	2400
aacttaatac aaatgtaatt tgtaataatt	gtttcatggt ttatgtgtag atctaatagc	2460
tgaactgatt caaactgtaa taagctcatc	aatttcattt ctatgaaaat gtgctctggt	2520
gtcacaggat gtttctgttg attttattca	tttcctggga attggtaaac atcatgttcc	2580
tgatgataac ccagtagcaa aaacatttgt	actgagtggg acaagccttg gggactgaaa	2640
aaaaaaaaag attaaaacca ttaaaaagaa	actcattttt acgctgaatg aacatttata	2700
tgattgcatt gggaccagtc atttcctaag	ctacatatgg ccatcttgac agtggttttt	2760
cttttgtgtg ttttaattatt atgtgtaa	atcataagaca aataaatttc actgtgccac	2820

<210> 7

<211> 6344

<212> DNA

<213> Homo sapiens

<400> 7

gcggaagtgt gggaggtct gcggggcggg	ctcaggaggt ccgcgggagg atggagcagt	60
gagcgggtct gggcggtgc tggcagcgcc	atggagacgg tacagctgag gaaccgcgcg	120
cgccggcagc tgaaaaagt ggatgaagat	agtttaacca aacaaccaga agaagtattt	180
gatgtcttag agaaacttgg agaaggtcc	tatggcagcg tatacaaagc tattcataaa	240
gagaccggcc agattgttgc tattaagcaa	gttcctgtgg aatcagacct ccaggagata	300
atcaaagaaa tctctataat gcagcaatgt	gacagccctc atgtagtcaa atattatggc	360
agttatttta agaacacaga cttatggatc	gttatggagt actgtggggc tgggtctgta	420
tctgatatca ttcgattacg aaataaaacg	ttaacagaag atgaaatagc tacaatatta	480
caatcaactc ttaagggact tgaatacctt	cattttatga gaaaaataca ccgagatata	540
aaggcaggaa atattttgct aaatacagaa	ggacatgcaa aacttgcaga ttttggggta	600
gcagggtcaac ttacagatac catggccaag	cggaaatacag tgataggaac accattttgg	660
atggctccag aagtgattca ggaaattgga	tacaactgtg tagcagacat ctggtccctg	720
ggaataactg ccatagaaat ggctgaagga	aagccccctt atgctgatat ccatccaatg	780
agggcaatct tcatgattcc tacaatcct	cctcccacat tccgaaaacc agagctatgg	840
tcagataact ttacagattt tgtgaaacag	tgtcttgtaa agagccctga gcagagggcc	900
acagccactc agctcctgca gcacccattt	gtcaggagtg ccaaaggagt gtcaatactg	960
cgagacttaa ttaatgaagc catggatgtg	aaactgaaac gccaggaatc ccagcagcgg	1020
gaagtggacc aggacgatga agaaaactca	gaagaggatg aaatggattc tggcacgatg	1080
gttcgagcag tgggtgatga gatgggcact	gtccgagtag ccagcaccat gactgatgga	1140

gccaaacta	tgattgagca	cgatgacacg	ttgccatcac	aactgggcac	catggtgatc	1200
aatgcagagg	atgaggaaga	ggaaggaact	atgaaaagaa	gggatgagac	catgcagcct	1260
gcgaaaccat	cctttcttga	atattttgaa	caaaaagaaa	aggaaaacca	gatcaacagc	1320
tttggcaaga	gtgtacctgg	tccactgaaa	aattcttcag	attggaaaat	accacaggat	1380
ggagactacg	agtttcttaa	gagttggaca	gtggaggacc	ttcagaagag	gctcttggcc	1440
ctggacccca	tgatggagca	ggagattgaa	gagatccggc	agaagtacca	gtccaagcgg	1500
cagcccatcc	tggatgccat	agaggctaag	aagagacggc	aacaaaactt	ctgagcaagg	1560
ccaggctgtg	agggccccag	ctccaccacg	gctttgggtg	aattctggat	ggcttgcctc	1620
atgtttgtta	gccagcactt	ctgctctgtc	gtctctccac	agcacctttg	tgaactcagg	1680
aatgtgcgcc	agtgggaagg	gctctcttga	cagtcagcgt	gccatcttga	tgtgtgtatg	1740
tacattggtc	aggtatatta	tctcaaagga	tttatattgg	cgcttttaac	tcagagtttt	1800
aaaccccgag	aacagagact	cctagttgag	tgatagctgg	gaaagtttta	cattgtctgt	1860
ttttcttctc	ccaatagctt	tcaattgttc	tttctggaag	acttttataa	aaatataaat	1920
atgcatatat	atatataaat	tataaataga	ttccccacgc	agtgtggtgg	catctctgta	1980
caggtacagt	tttaaacggt	ttgcctcttt	tctgtaagat	tatggtactg	tggaacatga	2040
gggcagagga	caccgggagg	ctgttagggg	gtcactgaat	cccaggagcc	aacctcccc	2100
tttgcagggc	tgcatttaaa	aattagggtt	gggacagttc	ttgtaccgtg	gtttcagcct	2160
tgtgtggtca	tactggctt	ctggagctat	tggatgatgc	caagggaaag	ctttgagagt	2220
ttatgtttac	tctttgagtc	ccaggagaag	cctggcaccc	tctttgcaaa	ttggcctttg	2280
ctctttcaat	gcctttcatc	catctccact	ctctcaactg	cctaaagtca	cagcacagat	2340
actgcccagt	gccttaagag	gagacatgat	ctctaccagg	gactctcagc	aaacacggga	2400
ctgtgttcag	tccacaaagg	aaaagcggtt	ttgaagctct	cattgttcat	gtaaaaatca	2460
tacacgtggc	atgttgctcc	acattcctta	cacacagggg	tagaggggat	tgcttttgtg	2520
accacagttc	aaatatgtga	ctgttttctt	ttctctttta	ctgctaagca	gcctggaaag	2580
gataaatgaa	tattagacta	agatttgttt	tccaggaggc	tcaatctgaa	cacacagaat	2640
gtcagagctg	gaagggacta	tagagatcat	ctgatctgat	cctcttgtag	ggatgatcgc	2700
aaaactgagg	tgtagagagg	ggaatggcca	aaatcacaaa	gcaagttagc	gttaagagct	2760
gagactagaa	ttcagggtcc	tactcccag	gccaccgaac	catgcagccc	cttctttggg	2820
ggaagagacc	tgtgtcagtc	ttggttaatt	gttccaggga	accttgctaa	cagaaacttg	2880
ctcttgcctt	ggctcttcag	tagatgacct	ggctgtaaag	agattccctg	gacgagccag	2940
atcattcagt	ttcagcgagt	ccttgagctc	cacaacatct	accagatata	gcagacaagc	3000

acccatggag	gcaggtttcg	ggcctgaagc	agatcagagg	gctttgcaaa	agacagcata	3060
gagccatctt	cctgcaactt	tacctctttc	cctcagatgg	ggagccatga	ctggggttgca	3120
cctcaggata	ctgtaatttg	actccataat	tgcttttgct	cctgaaacct	gggaatcaat	3180
ggaaaggcag	ggaatgtgcc	tcttctgtgg	ccagattctg	ttatttgcaa	ttaaagcaag	3240
tttttaaaaa	atgcaagagg	cagttgttag	tcttcagggc	ttggcaactg	aaatagctat	3300
gtggcgggata	cggaaaacag	aggacaattt	gaggatcttg	ctggaataat	aaatgacagc	3360
taccatttgt	tgagcaccta	ttatatatca	ggcactgagc	tggttaggct	ctaaacttca	3420
caataaccct	gtgacttaac	tactttatct	ccattttgta	gttgaagaaa	taagttcaga	3480
gagaaagatt	ccttcccaag	gtcatgcagc	tagtaaataga	tagaatcagg	attcatagca	3540
tcactatagg	gggtcaatat	ttacacaaaa	aaggaaagtc	acaagcctgt	ttaaaatgaa	3600
gtgaccacct	tttcttgcat	agactaaata	actcgaactg	gcatttttag	gttggaagaa	3660
cagctgaatt	agtagttaag	tctgatagcc	aagtaagttt	taaaaaccaa	agcatccagg	3720
atgcacaccc	ctgcaccatt	tgctgtgcga	attaatagtt	ctgtctctct	ctctctttct	3780
tttttctttt	tattctttga	gatggatttt	cgctcttgtc	gccaggctg	gagtacaatg	3840
gcacgatctt	ggctcactgc	aacctccgcc	tcccgggttc	aagcgattct	tctgctggga	3900
ttacagcata	tgccaccatg	cccagattat	ttttttgtat	ttgtagtaga	gacgggggtt	3960
caccatgtca	gtcaggctgg	tcttgaactc	ctgacctcag	gtgatccacc	cgcctcagcc	4020
tcccacactg	ctgggattac	aggcatgagc	caccgctcct	ggcctctctt	tcttttttaa	4080
acaaagaact	ttgcacttgg	ccagagagga	ggagaaagcc	cattttctcc	cttcctaagc	4140
tagatccaaa	taaaagaaag	ttcagttttc	ccccataact	attcttgggt	catgaacttt	4200
gatctggagt	ttgttttggt	tcaggaatgt	gtgcaccacg	cttctctgat	caacaaagtc	4260
tattgcttac	cagtctagct	tgatgaagcc	ttttggccag	aagtcaattt	gttttggatc	4320
agagaaattt	cctgacaagg	tataattgtt	ttctagtgc	agaaaggcaa	aggaacaagt	4380
cctagtgtgt	gttggtgttg	ttgaatacta	aatttaagat	atgtcagctt	gctttcaatg	4440
agccttgggc	ttctgttatt	gcttgagcat	ttggaactcg	agcttccaga	gaaatttgag	4500
gtcctcgctt	gttctctgcc	ttcaagaaac	aatgacctga	ttctgtcttt	aaaaaaaaaa	4560
atctcagaat	tctttttttg	tttgtgtttt	tttttttttt	tgagacagag	tctcactctg	4620
ttgcccaggc	tggagtgcag	tggcgccatc	tgggctcact	gcaacctccg	cctcccaggt	4680
tcaagcaatt	ctcctgcctc	agcctcccag	gtagctgcca	ctacagggtg	tgaccacca	4740
cggccggcta	atttttgtat	tttttagtaga	gacagggttt	caccatatta	gccagggtgg	4800
tcttgaactc	ctgaccttgt	gatccaccgc	cctcggcctc	ccaaagtgt	gggattacag	4860

```

gcgtgagcca ccttgccctgg ccaaaaaatct cagaattctt taagactggt ttaattgctc 4920
catcagtaat tttgaagcac tttccttttt tttttttttt cccctttttg tccctttccc 4980
caagccacca attggatgga tgaatgtttg acggggaaga ggaagggtag gaggatgcat 5040
ggatgagtgg atgagtggat cgatggatgt attgataaat agatagaacc agtcatctga 5100
agcaacttaa gaattgtagc cttgactcct tgagactgta gatttcgatc caggaaacat 5160
ttatttagca cctgccagat gccagaaatt tataccattt aaaactcagt aagtctttta 5220
aatatcagga aggagagaag cgacatcatg atacatccta tgggtattaa aaagccaata 5280
gaatattatg aataatttta tgctaataaa tttaacaact tcaacatcat aaacaaattc 5340
cttgaaaaat aaaaagtacc aaaattcatt caagaagaaa tagataccag cctgagcaac 5400
atggcaaaat cccatctcta caaaacatca aaaaaaaaaa aaattagtcg ggcatggtgg 5460
tgcacacctg taatcccagc ttgtcaggag gctgaagtgg gaggatcacc tgagcccagg 5520
gagggtcaagg atgcagttag ccatggtctc accactgcac tctagcctgg gtgacagaat 5580
gagaccccgt ctcaaaaaaa aagaagaagt agataatctg aatagcccta tatctataga 5640
aacttaatag tgctgggaga tataggtatt attatcctca ttttacagat gtgaaaattg 5700
aggctcagag aagtaaagtc tattgctcaa ggtcatgtgg ctagaatatg gcagagccat 5760
gattcagatc caggctctct gattcttatt ccagtgtcct ttctagcata ccatgttgcc 5820
tctaaagatt gcagctcctt atttactaga aaattgttcc tgcccaatct acatctccac 5880
ctcaccccat cttttcttaa gcactatggt tgtgttttta tcagtattat attcattgtc 5940
tttggaatac atgttcttgt ttgtgtttgg aaaaaaaatc tcttttacca gcttgacctc 6000
ggaccaactt ggaaaaaaa aagcttaaat gtttttgcta tgtacagttt aaaaatgtga 6060
agtttgtagc ttttaacttt tgtaagaaaa tctaataaca ctggcttaag tgctgacttg 6120
aaatgctatt ttgtaagggt tggatgtaag taatcaattg aggtcagcag tttgtatgag 6180
acatagcttc ctccattgcc cccactcctt ttttcttttt taagtttgag atgcttcctg 6240
tgtttttatg ttagaattgt tgttctcctt cttttcttct tcctatacct catcacgttt 6300
gttttaaata aactgtcctt tggaccacaa aaaaaaaaaa aaaa 6344

```

<210> 8
 <211> 113
 <212> PRT
 <213> Homo sapiens

<400> 8

Met Ser Leu Ser Phe Pro Val Gly Ala Ala Val Asn Cys Ala Asp Asn
 1 5 10 15

Thr Gly Ala Lys Ser Leu Tyr Ile Ile Ser Thr Lys Gly Ile Lys Gly
20 25 30

Arg Leu Asn Arg Leu Pro Ala Ala Asp Val Gly Asp Met Val Met Thr
35 40 45

Thr Val Lys Lys Gly Lys Pro Glu Leu Arg Lys Arg Val Pro Ser Ala
50 55 60

Val Val Ile Ser Gln Gln Lys Ser Tyr Trp Arg Lys Gly Gly Met Phe
65 70 75 80

Pro Tyr Phe Glu Asp Asn Ala Gly Val Ile Ala Asn Asn Lys Gly Glu
85 90 95

Met Lys Gly Ser Ala Ile Thr Gly Pro Val Ala Lys Lys Arg Val Asp
100 105 110

Leu

<210> 9
<211> 200
<212> PRT
<213> Homo sapiens

<400> 9

Met Gly His Glu Leu Ile Thr Val Glu Val Gly Gly Lys Tyr Met Asn
1 5 10 15

Ser Thr Asn Ala Leu Gly Arg Leu Gly Lys Pro Gln Ile Ser Gln Glu
20 25 30

Thr Leu Thr Ser Leu Ala Ser His Ser Thr Ser Ala Cys Met Val Ile
35 40 45

Gly Pro Phe Gln Gly Ala Lys Glu Glu Leu Gly Lys Ala Trp Ser Pro
50 55 60

Lys Leu Ala Leu Glu Asp His Thr Phe Lys Ile Leu Lys Gly Gly Arg
65 70 75 80

Asp Gly Ser Ser Ser Met Lys Phe Trp Ile Ser Leu Gly Leu Leu Leu
85 90 95

Gly Val Glu Ser Thr Val Leu Thr Thr Gln Glu Leu Lys Asn Leu Cys
100 105 110

Ile Phe Ser Val Lys Gly Ile Lys Gly Trp Leu Asn Arg Leu Ser Ala
 115 120 125

Ala Gly Val Gly Asp Val Val Met Ala Thr Val Lys Lys Gly Lys Pro
 130 135 140

Gln Leu Arg Lys Lys Ile Lys Asp Glu Met Leu Leu Tyr Phe Glu Asp
 145 150 155 160

Asn Val Glu Val Ile Gly Asn Asn Lys Gly Lys Ile Lys Ile Ser Ala
 165 170 175

Ile Thr Gly Gln Ile Val Lys Glu Gly Ala Glu Leu Cys Pro Ser Ile
 180 185 190

Arg Tyr Ser Ser Gly Ser Ile Ala
 195 200

<210> 10
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 10

Met Trp Lys Arg Gly Arg Gly Gly Ser Ser Gly Val Asn Phe Arg Ile
 1 5 10 15

Ser Val Gly Leu Pro Val Gly Ala Val Ile Asn Cys Ala Asp Asn Thr
 20 25 30

Gly Ala Lys Asn Leu Tyr Ile Ile Ser Val Lys Gly Ile Lys Gly Arg
 35 40 45

Leu Asn Arg Leu Pro Ala Ala Gly Val Gly Asp Met Gly Met Ala Thr
 50 55 60

Val Lys Lys Gly Lys Leu Glu Leu Arg Lys Lys Cys Ser Gly Thr Leu
 65 70 75 80

Val Lys Gly Thr Pro Leu Glu Asn Thr Glu Gly Gly Met Ser Gly Gly
 85 90 95

Ser Ala Ser Pro Trp Leu Pro Thr Met Asn Ala
 100 105

<210> 11
 <211> 79

<212> PRT

<213> Homo sapiens

<400> 11

Met Ala Thr Val Lys Lys Gly Lys Pro Glu Leu Arg Lys Lys Val His
 1 5 10 15

Pro Ala Val Val Ile Arg Gln Arg Lys Ser Tyr Arg Arg Lys Asp Gly
 20 25 30

Val Phe Leu Tyr Phe Glu Asp Asn Ala Gly Val Thr Val Asn Asn Lys
 35 40 45

Gly Glu Met Lys Gly Ser Ala Ile Thr Gly Pro Val Ala Lys Glu Cys
 50 55 60

Ala Asp Leu Trp Pro Trp Ile Ala Ser Ser Ala Gly Ser Ile Ala
 65 70 75

<210> 12

<211> 140

<212> PRT

<213> Homo sapiens

<400> 12

Met Ser Lys Arg Gly Arg Gly Gly Ser Ser Gly Ala Lys Phe Arg Ile
 1 5 10 15

Ser Leu Gly Leu Pro Val Gly Ala Val Ile Asn Cys Ala Asp Asn Thr
 20 25 30

Gly Ala Lys Asn Leu Tyr Ile Ile Ser Val Lys Gly Ile Lys Gly Arg
 35 40 45

Leu Asn Arg Leu Pro Ala Ala Gly Val Gly Asp Met Val Met Ala Thr
 50 55 60

Val Lys Lys Gly Lys Pro Glu Leu Arg Lys Lys Val His Pro Ala Val
 65 70 75 80

Val Ile Arg Gln Arg Lys Ser Tyr Arg Arg Lys Asp Gly Val Phe Leu
 85 90 95

Tyr Phe Glu Asp Asn Ala Gly Val Ile Val Asn Asn Lys Gly Glu Met
 100 105 110

Lys Gly Ser Ala Ile Thr Gly Pro Val Ala Lys Glu Cys Ala Asp Leu
 115 120 125

Trp Pro Arg Ile Ala Ser Asn Ala Gly Ser Ile Ala
 130 135 140

<210> 13

<211> 491

<212> PRT

<213> Homo sapiens

<400> 13

Met Glu Gln Pro Pro Ala Pro Lys Ser Lys Leu Lys Lys Leu Ser Glu
 1 5 10 15

Asp Ser Leu Thr Lys Gln Pro Glu Glu Val Phe Asp Val Leu Glu Lys
 20 25 30

Leu Gly Glu Gly Ser Tyr Gly Ser Val Phe Lys Ala Ile His Lys Glu
 35 40 45

Ser Gly Gln Val Val Ala Ile Lys Gln Val Pro Val Glu Ser Asp Leu
 50 55 60

Gln Glu Ile Ile Lys Glu Ile Ser Ile Met Gln Gln Cys Asp Ser Pro
 65 70 75 80

Tyr Val Val Lys Tyr Tyr Gly Ser Tyr Phe Lys Asn Thr Asp Leu Trp
 85 90 95

Ile Val Met Glu Tyr Cys Gly Ala Gly Ser Val Ser Asp Ile Ile Arg
 100 105 110

Leu Arg Asn Lys Thr Leu Ile Glu Asp Glu Ile Ala Thr Ile Leu Lys
 115 120 125

Ser Thr Leu Lys Gly Leu Glu Tyr Leu His Phe Met Arg Lys Ile His
 130 135 140

Arg Asp Ile Lys Ala Gly Asn Ile Leu Leu Asn Thr Glu Gly His Ala
 145 150 155 160

Lys Leu Ala Asp Phe Gly Val Ala Gly Gln Leu Thr Asp Thr Met Ala
 165 170 175

Lys Arg Asn Thr Val Ile Gly Thr Pro Phe Trp Met Ala Pro Glu Val
 180 185 190

Ile Gln Glu Ile Gly Tyr Asn Cys Val Ala Asp Ile Trp Ser Leu Gly

195					200					205						
Ile	Thr	Ser	Ile	Glu	Met	Ala	Glu	Gly	Lys	Pro	Pro	Tyr	Ala	Asp	Ile	
210					215					220						
His	Pro	Met	Arg	Ala	Ile	Phe	Met	Ile	Pro	Thr	Asn	Pro	Pro	Pro	Thr	
225					230					235					240	
Phe	Arg	Lys	Pro	Glu	Leu	Trp	Ser	Asp	Asp	Phe	Thr	Asp	Phe	Val	Lys	
245					250					255						
Lys	Cys	Leu	Val	Lys	Asn	Pro	Glu	Gln	Arg	Ala	Thr	Ala	Thr	Gln	Leu	
260					265					270						
Leu	Gln	His	Pro	Phe	Ile	Lys	Asn	Ala	Lys	Pro	Val	Ser	Ile	Leu	Arg	
275					280					285						
Asp	Leu	Ile	Thr	Glu	Ala	Met	Glu	Ile	Lys	Ala	Lys	Arg	His	Asp	Glu	
290					295					300						
Gln	Gln	Arg	Glu	Leu	Glu	Glu	Glu	Glu	Glu	Asn	Ser	Asp	Glu	Asp	Glu	
305					310					315					320	
Leu	Asp	Ser	His	Thr	Met	Val	Lys	Thr	Ser	Val	Gly	Glu	Cys	Gly	Thr	
325					330					335						
Met	Arg	Ala	Thr	Ser	Thr	Met	Ser	Glu	Gly	Ala	Gln	Thr	Met	Ile	Glu	
340					345					350						
His	Asn	Ser	Thr	Met	Leu	Glu	Ser	Asp	Leu	Gly	Thr	Met	Val	Ile	Asn	
355					360					365						
Ser	Glu	Asp	Glu	Glu	Glu	Glu	Asp	Gly	Thr	Met	Lys	Arg	Asn	Ala	Thr	
370					375					380						
Ser	Pro	Gln	Val	Gln	Arg	Pro	Ser	Phe	Met	Asp	Tyr	Phe	Asp	Lys	Gln	
385					390					395					400	
Asp	Phe	Lys	Asn	Lys	Ser	His	Glu	Asn	Cys	Asn	Gln	Asn	Met	His	Glu	
405					410					415						
Pro	Phe	Pro	Met	Ser	Lys	Asn	Val	Phe	Pro	Asp	Asn	Trp	Lys	Val	Pro	
420					425					430						
Gln	Asp	Gly	Asp	Phe	Asp	Phe	Leu	Lys	Asn	Leu	Ser	Leu	Glu	Glu	Leu	
435					440					445						

Gln Met Arg Leu Lys Ala Leu Asp Pro Met Met Glu Arg Glu Ile Glu
 450 455 460

Glu Leu Arg Gln Arg Tyr Thr Ala Lys Arg Gln Pro Ile Leu Asp Ala
 465 470 475 480

Met Asp Ala Lys Lys Arg Arg Gln Gln Asn Phe
 485 490

<210> 14
 <211> 487
 <212> PRT
 <213> Homo sapiens

<400> 14

Met Glu Thr Val Gln Leu Arg Asn Pro Pro Arg Arg Gln Leu Lys Lys
 1 5 10 15

Leu Asp Glu Asp Ser Leu Thr Lys Gln Pro Glu Glu Val Phe Asp Val
 20 25 30

Leu Glu Lys Leu Gly Glu Gly Ser Tyr Gly Ser Val Tyr Lys Ala Ile
 35 40 45

His Lys Glu Thr Gly Gln Ile Val Ala Ile Lys Gln Val Pro Val Glu
 50 55 60

Ser Asp Leu Gln Glu Ile Ile Lys Glu Ile Ser Ile Met Gln Gln Cys
 65 70 75 80

Asp Ser Pro His Val Val Lys Tyr Tyr Gly Ser Tyr Phe Lys Asn Thr
 85 90 95

Asp Leu Trp Ile Val Met Glu Tyr Cys Gly Ala Gly Ser Val Ser Asp
 100 105 110

Ile Ile Arg Leu Arg Asn Lys Thr Leu Thr Glu Asp Glu Ile Ala Thr
 115 120 125

Ile Leu Gln Ser Thr Leu Lys Gly Leu Glu Tyr Leu His Phe Met Arg
 130 135 140

Lys Ile His Arg Asp Ile Lys Ala Gly Asn Ile Leu Leu Asn Thr Glu
 145 150 155 160

Gly His Ala Lys Leu Ala Asp Phe Gly Val Ala Gly Gln Leu Thr Asp
 165 170 175

Thr Met Ala Lys Arg Asn Thr Val Ile Gly Thr Pro Phe Trp Met Ala
 180 185 190

Pro Glu Val Ile Gln Glu Ile Gly Tyr Asn Cys Val Ala Asp Ile Trp
 195 200 205

Ser Leu Gly Ile Thr Ala Ile Glu Met Ala Glu Gly Lys Pro Pro Tyr
 210 215 220

Ala Asp Ile His Pro Met Arg Ala Ile Phe Met Ile Pro Thr Asn Pro
 225 230 235 240

Pro Pro Thr Phe Arg Lys Pro Glu Leu Trp Ser Asp Asn Phe Thr Asp
 245 250 255

Phe Val Lys Gln Cys Leu Val Lys Ser Pro Glu Gln Arg Ala Thr Ala
 260 265 270

Thr Gln Leu Leu Gln His Pro Phe Val Arg Ser Ala Lys Gly Val Ser
 275 280 285

Ile Leu Arg Asp Leu Ile Asn Glu Ala Met Asp Val Lys Leu Lys Arg
 290 295 300

Gln Glu Ser Gln Gln Arg Glu Val Asp Gln Asp Asp Glu Glu Asn Ser
 305 310 315 320

Glu Glu Asp Glu Met Asp Ser Gly Thr Met Val Arg Ala Val Gly Asp
 325 330 335

Glu Met Gly Thr Val Arg Val Ala Ser Thr Met Thr Asp Gly Ala Asn
 340 345 350

Thr Met Ile Glu His Asp Asp Thr Leu Pro Ser Gln Leu Gly Thr Met
 355 360 365

Val Ile Asn Ala Glu Asp Glu Glu Glu Glu Gly Thr Met Lys Arg Arg
 370 375 380

Asp Glu Thr Met Gln Pro Ala Lys Pro Ser Phe Leu Glu Tyr Phe Glu
 385 390 395 400

Gln Lys Glu Lys Glu Asn Gln Ile Asn Ser Phe Gly Lys Ser Val Pro
 405 410 415

Gly Pro Leu Lys Asn Ser Ser Asp Trp Lys Ile Pro Gln Asp Gly Asp
420 425 430

Tyr Glu Phe Leu Lys Ser Trp Thr Val Glu Asp Leu Gln Lys Arg Leu
435 440 445

Leu Ala Leu Asp Pro Met Met Glu Gln Glu Ile Glu Glu Ile Arg Gln
450 455 460

Lys Tyr Gln Ser Lys Arg Gln Pro Ile Leu Asp Ala Ile Glu Ala Lys
465 470 475 480

Lys Arg Arg Gln Gln Asn Phe
485