

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199962382 B2
(10) Patent No. 759399

(54) Title
Compounds and methods for detecting biomolecules

(51)⁷ International Patent Classification(s)
C12Q 001/68

(21) Application No: **199962382**

(22) Application Date: **1999.06.02**

(87) WIPO No: **WO99/67619**

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/087641	1998.06.02	US
09/323424	1999.06.01	US

(43) Publication Date : **2000.01.10**

(43) Publication Journal Date : **2000.03.23**

(44) Accepted Journal Date : **2003.04.17**

(71) Applicant(s)
Ambergen, Inc.

(72) Inventor(s)
Kenneth J. Rothschild; Jerzy Olejnik


(74) Agent/Attorney
SPRUSON and FERGUSON,GPO Box 3898,SYDNEY NSW 2001

62382/99

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N	A2	(11) International Publication Number: WO 99/67619 (43) International Publication Date: 29 December 1999 (29.12.99)
(21) International Application Number: PCT/US99/12328 (22) International Filing Date: 2 June 1999 (02.06.99) (30) Priority Data: 60/087,641 2 June 1998 (02.06.98) US NOT FURNISHED 1 June 1999 (01.06.99) US 09/323424 (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US Not furnished (CIP) Filed on Not furnished (71) Applicant (for all designated States except US): AMBERGEN, INC. [US/US]; 1106 Commonwealth Avenue, Boston, MA 02215 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ROTHSCHILD, Kenneth, J. [US/US]; 397 Dorcar Road, Newton, MA 02459 (US). OLEJNIK, Jerzy [PL/US]; Apartment 3, 99 Boylston Street, Brookline, MA 02146 (US).		(74) Agents: CARROLL, Peter, G. et al.; Medlen & Carroll, LLP, Suite 2200, 220 Montgomery Street, San Francisco, CA 94104 (US). (81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: COMPOUNDS AND METHODS FOR DETECTING BIOMOLECULES (57) Abstract Methods are described for synthesizing compounds useful for detecting nucleic acids, and in particular, the hybridization of nucleic acids. Photocleavable agents are described, including but not limited to photocleavable nucleotides and photocleavable phosphoramidites.		



Compounds and Methods for Detecting Biomolecules

Field of the Invention

The present invention relates to compounds and methods for detecting biomolecules, and in particular, the detection of nucleic acids and nucleic acid hybridisation using photocleavable phosphoramidites.

Background

Numerous methods have been developed for the incorporation of ligands into synthetic oligonucleotides using phosphoramidite chemistry. See S.L. Beaucage and P.I. Radhakrishnan, *Tetrahedron* 49:1925 (1993). One approach is based on the incorporation of an aliphatic amino group into synthetic oligonucleotides by use of amino-modifier phosphoramidites or modified CPGs. See S. Agrawal *et al.*, *Nucleic Acids Res.*, 14:6227 (1986). See P.S. Nelson *et al.*, *Nucleic Acids Res.* 20:6253 (1992); P.S. Nelson *et al.*, *Nucleic Acids Res.* 17:7179 (1989); P.S. Nelson *et al.*, *Nucleic Acids Res.* 17:7187 (1989). Such amino groups allow the post-synthetic addition of a variety of molecules including fluorescent markers, biotin, haptens or enzymes which can be used as hybridisation probes for non-radioactive detection of specific target DNA/RNA or as sequencing primers. Amino-modified oligonucleotides have also been used for preparation of affinity matrices and as immobilised PCR primers.

One disadvantage of covalent labelling and immobilisation of oligonucleotides is that the covalent bond is not selectively cleavable, and, therefore does not allow for the recovery/release of oligonucleotide or the removal of the label. In order to circumvent this problem, several approaches have been proposed based on chemical cleavage. These include the use of cleavable disulfide bonds or the introduction of acid cleavable linkers. See W Bannwarth and J. Wippler, *Helv. Chim. Acta* 73:1139 (1990); B.D. Gildea *et al.*, *Tetrahedron Lett.* 31:7095 (1990); E. Leikauf *et al.*, *Tetrahedron* 51:3793 (1995); J. Olejnik *et al.*, *Nucleic Acids Res.* 24:361 (1996). These approaches, however, require the application of chemical reagents that are not suitable for many applications. They also necessitate removal of the excess of cleaving reagents and result in residual modification of the oligonucleotide.

Summary of the Invention

The present invention relates to compounds and methods for detecting biomolecules, and in particular, the detection of nucleic acid and nucleic acid hybridisation using photocleavable agents, including photocleavable detectable moieties, photocleavable nucleotides (*e.g.* photocleavable NTPs) and photocleavable phosphoramidites. It is not intended that the present invention be limited to the type of biomolecule detected. Both



nucleic acids and proteins can be detected using the photocleavable agents of the present invention. The details regarding detection of various biomolecules (including proteins) are set forth in Attachments A, B, C, D and E hereto, all of which constitute the disclosure and specification of the present application.

5 In one embodiment, the present invention contemplates a photocleavable nucleotide. In a specific embodiment, the photocleavable nucleotide is a photocleavable dideoxynucleotide. The latter is useful in a variety of formats and assays, including but not limited to, nucleic acid sequencing.

10 In one embodiment, the present invention contemplates a composition, comprising a photocleavable aminotag phosphoramidite selected from the group consisting of compounds 6 and 7 of Figure 2. From Figure 2, it should be clear that the present invention contemplates particular protecting groups ("PG") suitable for such compounds, including but not limited to, 9-fluorenylmethoxycarbonyl (Fmoc). In a preferred embodiment, the photocleavable aminotag phosphoramidite is incorporated into an
15 oligonucleotide during synthesis at the 5' end of an oligonucleotide.

Thus, according to one embodiment of the invention, there is provided a composition, comprising a photocleavable aminotag phosphoramidite selected from the group of compounds A and B of Figure 8 and the compound of Figure 10.

20 According to another embodiment of the invention, there is provided a composition, comprising a photocleavable aminotag phosphoramidite selected from the group of compounds 6 and 7 of Figure 2 and compound 9 of Figure 9.

25 In another embodiment, the present invention contemplates a composition, comprising a photocleavable sulfhydryltag phosphoramidite such as that shown in Figure 4. In a preferred embodiment, the photocleavable sulfhydryltag phosphoramidite is incorporated at the 5' end of an oligonucleotide.

Thus, according to another embodiment of the invention, there is provided a composition, comprising the photocleavable sulfhydryltag phosphoramidite set forth in Figure 4.

30 The present invention also contemplates methods of synthesis of photocleavable phosphoramidites and methods of synthesis of oligonucleotides containing such photocleavable phosphoramidites. Moreover, the present invention contemplates methods of incorporating detectable moieties or markers ("M") to a phosphoramidite and photocleaving so as to release said marker.

Thus, according to another embodiment of the invention, there is provided a method, comprising:



a) providing:

i) a compound comprising a photocleavable phosphoramidite and a protected tag linked to said photocleavable phosphoramidite, wherein said protected tag is selected from the group consisting of an amino tag, a sulfhydryl tag, a nucleotide tag and
5 a mass tag,

ii) oligonucleotide synthesis means; and

b) synthesising an oligonucleotide with said synthesis means and said photocleavable phosphoramidite under conditions such that an oligonucleotide having a 3' end and a 5' end is produced wherein said photocleavable phosphoramidite is
10 incorporated at said 5' end.

According to another embodiment of the invention, there is provided a method, comprising:

a) providing:

i) a compound comprising a photocleavable phosphoramidite and a
15 protected tag linked to said photocleavable phosphoramidite, wherein said protected tag is selected from the group consisting of an amino tag, a sulfhydryl tag, nucleotide tag and a mass tag,

ii) oligonucleotide synthesis means,

iii) a marker molecule,

20 iv) a source of electromagnetic radiation;

b) synthesising an oligonucleotide with said synthesis means and said photocleavable phosphoramidite under conditions such that an oligonucleotide having a 3' end and a 5' end is produced wherein said photocleavable phosphoramidite is incorporated at said 5' end;

25 c) linking said marker molecule to said synthesised oligonucleotide of step (b) so as to produce a photocleavable marker/oligonucleotide complex; and

d) exposing said complex to electromagnetic radiation from said source of electromagnetic radiation under conditions such that said oligonucleotide is released from said marker.

30 In a further embodiment, the present invention contemplates oligonucleotides linked to a solid support or surface via a photocleavable linkage. In such an embodiment, the oligonucleotide can be released from the solid support (e.g. a bead) or a surface (e.g. a gold surface) by photocleavage.

Accordingly, in another embodiment of the invention, there is provided a method, comprising:



a) providing:

i) a compound comprising photocleavable phosphoramidite, and a protected tag linked to said photocleavable phosphoramidite, wherein said protected tag is selected from the group consisting of an amino tag, a sulfhydryltag, nucleotide tag and a mass tag;

ii) oligonucleotide synthesis means,

iii) a surface,

iv) a source of electromagnetic radiation;

b) synthesising an oligonucleotide with said synthesis means and said photocleavable phosphoramidite under conditions such that an oligonucleotide having a 3' end and a 5' end is produced wherein said photocleavable phosphoramidite is incorporated at said 5' end;

c) attaching said synthesised oligonucleotide of step (b) to a surface to create an immobilised oligonucleotide; and

d) exposing said immobilised oligonucleotide to electromagnetic radiation from said source of electromagnetic radiation under conditions such that said oligonucleotide is released from said surface.

Description of Drawings

Figure 1 schematically shows a generic structure for the photocleavable phosphoramidites of the present invention.

Figure 2 schematically shows one embodiment of a synthesis scheme for a variety of aminotag phosphoramidites of the present invention.

Figure 3 schematically shows one embodiment of linking a marker ("M") to an aminotag phosphoramidite that has been incorporated into an oligonucleotide, followed by photocleavage.



Figure 4 shows the structure of one embodiment of the sulfhydryltag phosphoramidites of the present invention.

Figure 5 schematically shows one embodiment of a synthesis scheme for an oligonucleotide conjugate comprising a photocleavable aminotag phosphoramidite and a peptide marker.

Figure 6 schematically shows the detection of a peptide marker of an oligonucleotide conjugate by mass spectrometry.

Figure 7 shows one embodiment of a photocleavable nucleotide of the present invention.

Figure 8 shows two compounds of Figure 7 where the protecting group is indicated



generically as "PG".

Figure 9 schematically shows one embodiment of a synthesis scheme for a novel photocleavable aminotag phosphoramidite.

Figure 10 shows one compound of Figure 9 where the protecting group is indicated generically as "PG".

DEFINITIONS

To facilitate understanding of the invention, a number of terms are defined below.

The terms "marker" and "marker molecule" refer to a detectable moiety, i.e. a chemical group, structure of compound that possesses a specifically identifiable physical property which can be distinguished from the physical properties of other chemicals present in the heterologous mixture. It is not intended that the present invention be limited to a particular detectable moiety. For example, the present invention contemplates detectable moieties having the properties of fluorescence, phosphorescence and luminescence. A variety of molecules have such properties as set forth in U.S. Application Ser. No. 08/345,807, hereby incorporated by reference. Preferred detectable moieties are those that are detectable by spectrometry, and in particular, by mass spectrometry.

"Nucleic acid sequence" and "nucleotide sequence" as used herein refer to an oligonucleotide or polynucleotide, and fragments or portions thereof, and to DNA or RNA of genomic or synthetic origin which may be single- or double-stranded, and represent the sense or antisense strand.

The photocleavable agents of the present invention can be "linked" or "incorporated" into proteins and nucleic acids. By use of the term "linked" or "incorporated" it is not meant that the entire photocleavable agent need be part of the final molecule. Some photocleavable agents of the present invention have reactive groups (i.e. the agent is an "activated agent") and leaving groups such that the photocleavable agent upon incorporation or linkage loses one or more groups.

The term "portion" when used in reference to a nucleotide sequence refers to fragments of that nucleotide sequence. The fragments may range in size from 1

nucleotide residues (and more preferably 5 or greater) to the entire nucleotide sequence minus one nucleic acid residue. The present invention contemplates, for example, that the photocleavable agents may be incorporated into just a portion of a nucleic acid.

As used herein, the terms "complementary" or "complementarity" are used in reference to "polynucleotides" and "oligonucleotides" (which are interchangeable terms that refer to a sequence of nucleotides) related by the base-pairing rules. For example, for the sequence 5'-AGT-3' is complementary to the sequence 5'-ACT-3'. Complementarity can be "partial" or "total." "Partial" complementarity is where one or more nucleic acid bases is not matched according to the base pairing rules. "Total" or "complete" complementarity between nucleic acids is where each and every nucleic acid base is matched with another base under the base pairing rules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. The present invention contemplates hybridization (e.g. complete, partial, low and high stringency) using oligonucleotides having photocleavable agents incorporated therein.

The terms "homology" and "homologous" as used herein in reference to nucleotide sequences refer to a degree of complementarity with other nucleotide sequences. There may be partial homology or complete homology (*i.e.*, identity). A nucleotide sequence which is partially complementary, *i.e.*, "substantially homologous," to a nucleic acid sequence is one that at least partially inhibits a completely complementary sequence from hybridizing to a target nucleic acid sequence. The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or Northern blot, solution hybridization and the like) under conditions of low stringency. A substantially homologous sequence or probe will compete for and inhibit the binding (*i.e.*, the hybridization) of a completely homologous sequence to a target sequence under conditions of low stringency. This is not to say that conditions of low stringency are such that non-specific binding is permitted; low stringency conditions require that the binding of two sequences to one another be a specific (*i.e.*, selective) interaction. The absence of non-specific binding may be tested by the use of

a second target sequence which lacks even a partial degree of complementarity (*e.g.*, less than about 30% identity); in the absence of non-specific binding the probe will not hybridize to the second non-complementary target.

Low stringency conditions comprise conditions equivalent to binding or hybridization at 42°C in a solution consisting of 5X SSPE (43.8 g/l NaCl, 6.9 g/l NaH₂PO₄•H₂O and 1.85 g/l EDTA, pH adjusted to 7.4 with NaOH), 0.1% SDS, 5X Denhardt's reagent [50X Denhardt's contains per 500 ml: 5 g Ficoll (Type 400, Pharmacia), 5 g BSA (Fraction V; Sigma)] and 100 µg/ml denatured salmon sperm DNA followed by washing in a solution comprising 5X SSPE, 0.1% SDS at 42°C when a probe of about 500 nucleotides in length is employed.

The art knows well that numerous equivalent conditions may be employed to comprise low stringency conditions; factors such as the length and nature (DNA, RNA, base composition) of the probe and nature of the target (DNA, RNA, base composition, present in solution or immobilized, *etc.*) and the concentration of the salts and other components (*e.g.*, the presence or absence of formamide, dextran sulfate, polyethylene glycol), as well as components of the hybridization solution may be varied to generate conditions of low stringency hybridization different from, but equivalent to, the above listed conditions. In addition, the art knows conditions which promote hybridization under conditions of high stringency (*e.g.*, increasing the temperature of the hybridization and/or wash steps, the use of formamide in the hybridization solution, *etc.*).

When used in reference to a double-stranded nucleic acid sequence such as a cDNA or genomic clone, the term "substantially homologous" refers to any probe which can hybridize to either or both strands of the double-stranded nucleic acid sequence under conditions of low stringency as described above.

When used in reference to a single-stranded nucleic acid sequence, the term "substantially homologous" refers to any probe which can hybridize (*i.e.*, it is the complement of) to the single-stranded nucleic acid sequence under conditions of low stringency as described above.

As used herein, the term "hybridization" is used in reference to the pairing of

complementary nucleic acids using any process by which a strand of nucleic acid joins with a complementary strand through base pairing to form a hybridization complex. Hybridization and the strength of hybridization (*i.e.*, the strength of the association between the nucleic acids) is impacted by such factors as the degree of complementarity
5 between the nucleic acids, stringency of the conditions involved, the T_m of the formed hybrid, and the G:C ratio within the nucleic acids.

As used herein the term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bounds between complementary G and C bases and between complementary A and T bases; these
10 hydrogen bonds may be further stabilized by base stacking interactions. The two complementary nucleic acid sequences hydrogen bond in an antiparallel configuration. A hybridization complex may be formed in solution (as determined, *e.g.*, by C_0t or R_0t analysis) or between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized to a solid support [*e.g.*, a nylon membrane or a nitrocellulose
15 filter as employed in Southern and Northern blotting, dot blotting or a glass slide as employed in *in situ* hybridization, including FISH (fluorescent *in situ* hybridization)].

"Stringency" when used in reference to nucleic acid hybridization typically occurs in a range from about $T_m - 5^\circ\text{C}$ (5°C below the T_m of the probe) to about 20°C to 25°C below T_m . The term " T_m " is used in reference to the "melting temperature." The melting
20 temperature is the temperature at which a population of double-stranded nucleic acid molecules becomes half dissociated into single strands. As will be understood by those of skill in the art, a stringent hybridization can be used to identify or detect identical polynucleotide sequences or to identify or detect similar or related polynucleotide sequences. Under "stringent conditions" a sequence or fragments thereof will hybridize
25 to the sequence's exact complement and closely related sequences.

DNA molecules are said to have "5' ends" and "3' ends" because mononucleotides are reacted to make oligonucleotides in a manner such that the 5' phosphate of one mononucleotide pentose ring is attached to the 3' oxygen of its neighbor in one direction via a phosphodiester linkage. Therefore, an end of an oligonucleotide is referred to as the
30 "5' end" if its 5' phosphate is not linked to the 3' oxygen of a mononucleotide pentose

ring. An end of an oligonucleotide is referred to as the "3' end" if its 3' oxygen is not linked to a 5' phosphate of another mononucleotide pentose ring. As used herein, a nucleic acid sequence, even if internal to a larger oligonucleotide, also may be said to have 5' and 3' ends. In either a linear or circular DNA molecule, discrete elements are referred to as being "upstream" or 5' of the "downstream" or 3' elements. This terminology reflects the fact that transcription proceeds in a 5' to 3' fashion.

DESCRIPTION OF THE INVENTION

The present invention relates to compounds and methods for detecting biomolecules, and in particular, the detection of nucleic acid and nucleic acid hybridization using photocleavable agents, including photocleavable detectable moieties, photocleavable nucleotides and photocleavable phosphoramidites. Figure 1 schematically shows a generic structure for the photocleavable phosphoramidites of the present invention. In one embodiment, the present invention contemplates non-nucleosidic photocleavable phosphoramidites. These reagents introduce a photocleavable amino group on the 5'-terminal phosphate of synthetic oligonucleotides and are fully compatible with automated solid support synthesis. The 5' photocleavable amino group enables introduction of a variety of amine-reactive markers onto synthetic oligonucleotides as well as immobilization on activated solid supports. The photocleavable ("PC") bond on the 5'-phosphate can then be selectively cleaved by near-UV illumination thereby enabling release of the marker or detachment of the oligonucleotide from a solid support.

It is not intended that the present invention be limited to the particular marker linked to the 5' terminal phosphate of the oligonucleotide. The preparation of photocleavable conjugates with polypeptides, biotin, digoxigenin, and tetramethylrhodamine are contemplated.

It is specifically contemplated that the 5'-PC-amino group can be used as an affinity tag for photocleavage-mediated affinity purification and phosphorylation of synthetic oligonucleotides in conjunction with activated supports. Upon irradiation with 300-350 nm light, the 5'-PC-amino group is cleaved with high efficiency, resulting in rapid release of 5'-phosphorylated oligonucleotides into solution. Such 5'-PC-amino

labeled oligonucleotides are useful in a variety of applications in molecular biology including multiple non-radioactive probing of DNA/RNA blots, affinity isolation and purification of nucleic acids binding proteins, diagnostic assays requiring release of the probe-target complex or specific marker, cassette mutagenesis and PCR. They enable the spatially-addressable photorelease of the probe-target complexes or marker molecules for diagnostic purposes.

A. Photocleavable AminoTag Phosphoramidites

A variety of aminotag phosphoramidites are contemplated. In one embodiment (Figure 2), the PC-aminotag phosphoramidite (compound 6) comprises a protected amino group linked through a methylene group to a photocleavable 1-(2-nitrophenyl)ethyl linker derivatized with N,N'-diisopropyl-2-cyanoethyl phosphoramidite. In another embodiment, the present invention contemplates a "PC-X-aminotag" (Figure 2, compound 7) which comprises an additional spacer (*e.g.* 6-aminocaproic acid, referred to as "X") between the amino group and the photocleavable linker. The properties of modified oligonucleotides prepared using these PC-aminotag phosphoramidites and their conjugates are such that the present invention provides advantages over other cleavable systems (discussed above).

Unlike biotin, which can interact only with avidin/streptavidin coated surfaces, PC-aminotag-phosphoramidite introduces a photocleavable amino group onto the 5' end of an oligonucleotide. As such it can be used to react with a wide variety of activated surfaces and molecules, thereby facilitating the formation of many different PC-linkers and PC-markers.

B. Synthesis of Oligonucleotides With AminoTag Phosphoramidites

The photocleavable-aminotag phosphoramidites of the present invention can be readily used in any automated DNA/RNA synthesizer employing standard phosphoramidite chemistry (*i.e.* an oligonucleotide synthesis means). The selective reaction of PC-aminotag phosphoramidites with the free 5'-OH group of a growing oligonucleotide chain, followed by cleavage from the support and deprotection, results in the introduction of a phosphodiester group linked to a primary aliphatic amino group through a photocleavable

linker. This amino group can then be used to introduce a variety of photocleavable markers through a postsynthetic modification reaction (Figure 3) with amine reactive reagents. It can also be used to attach synthetic oligonucleotides to activated solid supports for a variety of applications requiring immobilization and release of DNA molecules. Upon irradiation with near-UV light, the phosphodiester bond between the photocleavable linker and the phosphate is cleaved, resulting in the formation of a 5'-monophosphate on the released oligonucleotide. The 1-(2-nitrophenyl)ethyl moiety is converted into a 2-nitrosoacetophenone derivative and is released together with the attached marker molecule M.

C. Photocleavable SulfhydrylTag Phosphoramidites

The present invention also contemplates sulfhydryltag phosphoramidites. The design of such compounds (Figure 4) is similar to that for the aminotag phosphoramidites such that the photocleavable sulfhydryl group can be incorporated at the 5'-end of an oligonucleotide. This phosphoramidite allows for reactions with a variety of thiol-activated supports and biomolecules. A trityl protective group is also incorporated for protection purposes.

D. Photocleavable Phosphoramidites Comprising Peptide Markers

The present invention contemplates photocleavable phosphoramidites comprising peptide markers. Peptides as markers have a number of desirable features when used as mass markers, including a large set of distinct masses which are easily resolved by MALDI-MS. When attached through a photocleavable linker to a specific oligonucleotide sequence, the resulting conjugate can be used in a wide variety of applications in DNA-analysis by mass spectrometry.

The compounds can be made using a conventional solid phase nucleic acid synthesizer (e.g. a synthesis means). The DNA probe is first assembled on an automated DNA synthesizer in the conventional fashion (Figure 5). The aminotag phosphoramidite is incorporated at the 5' end of the sequence. At the end of the cycle, the support (CPG) is treated with base, which selectively removes the Fmoc protective group from the linker.

In the next step, an N-Fmoc protected peptide marker is introduced and an activating mixture is added, resulting in the formation of the peptide bond between the peptide marker and photocleavable linker DNA. Conventional cleavage and deprotection with ammonia yields the desired conjugate with a free amino group on the peptide marker.

E. Detection Of Peptide Markers

The present invention contemplates the detection of markers, including but not limited to peptide markers. In one embodiment, the present invention contemplates detection by mass spectrometry (e.g. MALDI-MS). In one embodiment, the DNA moiety of the mass marker is hybridized to an immobilized target DNA single strand. The bond between the hybridization probe and the mass tag is then cleaved by the UV-MALDI desorption laser pulse (Figure 6) and the released mass tag (e.g. an octapeptide mass tag) is detected in the mass spectrum indicating the presence of the complementary sequence of its hybridization probe. This method potentially offers both higher sensitivity compared to the direct detection of the DNA hybridization probe and is less prone to signal degradation due to metastable fragmentation and cation adduct formation. Furthermore, it allows for simultaneous detection of several target sequences, since peptide mass markers can be readily synthesized with easily resolvable mass differences.

F. Photocleavable Nucleotides

The present invention also contemplates photocleavable nucleotides (PC-nucleotides). Using these PC-nucleotides, photocleavable tags (PC-tags) can be enzymatically incorporated into an oligonucleotide through a wide range of enzyme based methodologies for the synthesis and modification of DNA/RNA. These include PCR, terminal deoxynucleotidyl transferase (single labeling or tailing at the 3'-end) and nick translation reaction or replacement synthesis using T4 DNA polymerase (multiple label incorporation). For RNA labeling, appropriate polymerases, such as SP6, T7 and T3 polymerases can be used. Alternatively, T4 RNA ligase can be used to join 3'-labeled (deoxy)nucleotide 5'-monophosphate to target RNA. Dideoxy-PC-nucleotides can be used to end-label the products of polymerase based sequencing reactions thereby facilitating

MALDI-MS based sequencing.

EXPERIMENTAL

The following examples serve to illustrate certain preferred embodiments and aspects of the present invention and are not to be construed as limiting the scope thereof.

5 In the experimental disclosure which follows, the following abbreviations apply:
eq (equivalents); μ (micron); M (Molar); μ M (micromolar); mM (millimolar); N
(Normal); mol (moles); mmol (millimoles); μ mol (micromoles); nmol (nanomoles); g
(grams); mg (milligrams); μ g (micrograms); ng (nanograms); L (liters); ml (milliliters);
10 μ l (microliters); cm (centimeters); mm (millimeters); μ m (micrometers); nM
(nanomolar); $^{\circ}$ C (degrees Centigrade); PBS (phosphate buffered saline); TLC (thin layer
chromatography); HPTLC (high performance thin layer chromatography); HPLC (high
pressure liquid chromatography); U (units); d(days).

All chemicals used in the synthesis were purchased from Aldrich Chemical Co.
(Milwaukee, WI), unless otherwise indicated. Biotinamidohexanoic acid NHS ester was
15 purchased from Sigma (St. Louis, MO), digoxigenin-X-NHS ester (digoxigenin-3-O-
methylcarbonyl-(-aminocaproic acid N-hydroxysuccinimide ester) was obtained from
Boehringer-Mannheim (Indianapolis, IN) and tetramethylrhodamine-X-NHS ester (6-
tetramethylrhodamine-5-(and-6)-carboxamidohexanoic acid succinimidyl ester, TAMRA-
X-NHS) was from Molecular Probes (Eugene, OR). Fmoc-NHS and N-Fmoc-6-
20 aminocaproic acid were from Novabiochem (La Jolla, CA). 1 H NMR spectra were
recorded in $CDCl_3$ on a Varian Unity Plus (Palo Alto, CA) spectrometer at 400 MHz with
chemical shifts (δ , ppm) reported relative to a tetramethylsilane internal standard.
Oligonucleotide synthesis was performed on an Applied Biosystem DNA/RNA
synthesizer, model 392 (Foster City, CA). Samples were irradiated with a Blak Ray XX-
25 15 UV lamp (Ultraviolet Products Inc., San Gabriel, CA) at a distance of 15 cm (emission
peak 365 nm; 300 nm cut-off; intensity 2.3 mW/cm 2 @365 nm). UV-VIS spectra were
recorded on a Shimadzu 2101PC spectrophotometer (Columbia, MD). HPLC analysis was
performed on a Waters (Milford, MA) system consisting of a U6K injector, 600
Controller, Novapak C18 (3.9 x 150 mm) column and a 996 photodiode array detector.

Buffer A: 0.1 N triethylamine acetate, pH = 7.0; buffer B: acetonitrile. Elution was performed using a linear gradient (8 - 50%) of buffer B in buffer A over 45 minutes at a 1 ml/min. flow. Preparative purifications were achieved on a Waters Novapak C18 RCM cartridge (8 x 100 mm) using conditions as specified above, except for flow rate, which was increased to 2 ml/min. Fractions were then analyzed by analytical HPLC, fractions pooled and freeze-dried. No special precautions were necessary to protect the reagents and the labeled oligonucleotides from light.

EXAMPLE 1

This example describes the synthesis of 5-N-Fmoc(aminomethyl)-1-(2-nitrophenyl)-ethanol (Figure 2, compound 4). As shown in Figure 2, this compound can be obtained from 5-bromomethyl-2-nitroacetophenone (compound 1) through a quaternization reaction with hexamethylenetetramine and decomposition of the resulting quaternary salt (compound 2) under acidic conditions. Crude 5-aminomethyl-2-nitroacetophenone hydrochloride (2.5 mmole) was resuspended in 12 ml of methanol and cooled in an ice bath. 48 mg (1.26 mmole, 2.0 eq.) of sodium borohydride was added in small portions with continuous stirring during 10 minutes. Subsequent portions of borohydride (30 mg each) were added to reaction mixture until TLC showed complete conversion (Figure 2, compound 3). The reaction was quenched by addition of 10 ml acetone. 10 ml water was added and the reaction mixture was concentrated under vacuum to approximately 10 mL. 10 ml of acetonitrile was added and the pH of the resulting solution was adjusted to 8.5 using 1N HCl. Fmoc-NHS (1.03 g; 3 mmole, 1.2 eq. in acetonitrile (10 ml) was added slowly during 15 minutes, the pH being adjusted to 8.5 with triethylamine. The reaction mixture was stirred for an additional 1 hour, then it was acidified to pH = 3 and extracted with chloroform. The organic layer was dried over anhydrous sodium sulphate and evaporated, yielding 1.4 g of compound 4 as yellow oil, which was purified on a silica gel column using a step gradient (0-2%) of methanol in chloroform. The product was isolated at 1% of MeOH. Yield - 720 mg (69 %) ¹H NMR (CDCl₃ + CD₃OD): 1.57-1.52 (d, 3H), 2.39 (s, 1H), 4.17-4.19 (m, 1H), 4.39-4.47 (d-d, 2H), 5.34-5.36 (q, 1H), 5.83 (s-br, 1H), 7.17-7.19 (d, 1H), 7.21-7.30 (t, 2H), 7.36-7.40

(t, 2H), 7.55-7.57 (d, 2H), 7.69 (s, 1H), 7.73-7.75 (d, 2H), 7.83-7.85 (d, 1H).

EXAMPLE 2

This example describes the synthesis of compound 5 (Figure 2). This reaction was carried out as for compound 4 (See Example 1), except N-Fmoc-6-aminocaproic acid NHS ester was used instead of the Fmoc-NHS. To the solution of compound 3 (5.3 mmol in 40 ml acetonitrile:H₂O, 1:1, pH = 8.5) a solution of N-Fmoc-6-aminohexanoic acid NHS ester (2.9 g; 1.2 eq.) in 20 ml of dioxane was added during 15 minutes. After work-up, the crude product was purified on a silica gel column using step gradient of methanol in chloroform (0-3%), product (compound 5) being eluted at 2% methanol. Yield - 2.03 g (72%). R_f = (CHCl₃ : MeOH, 95:5, v:v) 1H NMR (CDCl₃ + CD₃OD): 7.79-7.77(d, 1H), 7.72-7.67(m, 3H), 7.60-7.50(d, 2H), 7.35-7.31(t, 2H), 7.27-7.19(m, 3H), 6.42-6.37(app. t, br, 1H), 5.37-5.33(q, 1H), 5.16-5.08(t, br, 1H), 4.46-4.41(m, 2H), 4.33-4.30(d, 2H), 4.15-4.10(t, 1H), 3.14-3.09(q, 2H), 2.23-2.18(t, 2H), 1.67-1.59(m, 2H), 1.48-1.42(m, 3H), 1.35-1.28(m, 2H).

EXAMPLE 3

This example describes the synthesis of yet another photocleavable aminotag phosphoramidite of the present invention, namely 5-N-Fmoc-(aminomethyl)-1-(2-nitrophenyl)ethyl-2-cyanoethyl-N,N-diisopropylamino-phosphoramidite (compound 6, Figure 2). For this reaction, compound 4 (200mg, 0.52 mmole) was dried in vacuo over P₂O₅ for 5 hrs. 3 ml of anhydrous acetonitrile was injected through septum, followed by N,N-diisopropylethylamine (350 μ l, 2.08 mmole, 4 eq.) and finally cyanoethoxy-N,N-diisopropylchlorophosphine (0.52 mmole, 117 μ l, 1 eq.). The reaction mixture was stirred at RT for 1 hr then another 0.5 eq. of phosphine was added. After 2 hours the reaction mixture was treated with 1 ml of ethyl acetate and worked-up between aq. NaCl and dichloromethane. The organic layer was washed with water, dried over sodium sulphate and evaporated. Product was purified on silica gel column using a step gradient of ethyl acetate in hexane (0-30%) containing 0.2% of triethylamine. Product was isolated at 22%

of ethyl acetate, yielding 130 mg of compound 6 (42%). R_f (hexane:ethyl acetate:Et₃N, 45:45:10, v:v:v) 0.80 and 0.86 (two diastereoisomers). ESI-MS M⁺ = 619.3 (+H) (618.26 calculated). Elemental analysis: CHN%(calculated); C64.03(64.07); H6.23(6.35); N8.76(9.06). ¹H NMR (CDCl₃): 7.88-7.84(m, 1H), 7.77-7.27(m, 3H), 7.60-7.55 (m, 2H), 7.40-7.34 (m, 2H), 7.30-7.24(m, 3H), 5.67(t, 1H), 5.55-5.49(q, 1H), 4.45-4.37(m, 4H), 4.24-4.16(m, 1H), 3.87-3.79(m, 1H), 3.65-3.58(m, 2H), 3.48-3.42(m, 1H), 2.62-2.40(m, 2H), 1.55-1.53(d, J = 6.8Hz + s, 4H), 1.19-1.08(d+d+d, J = 6.8Hz, 12H), 0.89-0.85(d, J = 6.8Hz, 3H).

EXAMPLE 4

This example describes the synthesis of an oligonucleotide comprising a photocleavable aminotag phosphoramidite, namely Compound 7 (Figure 2). This compound was prepared from compound 5 (1.0 g; 1.92 mmol) N,N-diisopropylethylamine (1.3 ml; 4 eq.) and cyanoethoxy-N,N-diisopropylchlorophosphine (1.5 eq.) in 12 ml acetonitrile using a procedure similar to that described for compound 6. Product was purified on a silica gel column using step gradient of ethyl acetate in hexane (10-70%, v/v) containing 0.8 % triethylamine. Product was isolated at 50% ethyl acetate as two diastereoisomers (R_f = 0.69, 0.61; AcOEt : hexane : Et₃N, 8:1:1, v/v). Yield - 580 mg (41%). ESI-MS, M⁺ = 731.2 (731.3, calculated); elemental analysis C 65.34(64.01) H 6.65(6.89) N 10.08(9.57) (calculated) ¹H NMR: 7.84-7.82(d, J = 8.4, 1H), 7.72-7.69 (m, 3H), 7.57-7.52 (m, 2H), 7.37-7.32(m, 2H), 7.29-7.25(m, 3H), 6.73-6.65(t, J = 6Hz, 1H), 5.49-5.44(q, 1H), 5.10-5.00(t, 1H), 4.48-4.30(m, 4H), 4.20-4.05(m, 2H), 3.50-3.45(m, 2H), 2.66-2.64(m, 2H), 2.22-2.16(t, 2H), 1.54-1.48(d, J = 6.0Hz, 4H), 1.24-1.20(m, 6H), 1.16-1.13(d, J = 6.4 Hz, 4H), 1.09-1.05(d, J = 6.8 Hz, 4H)

EXAMPLE 5

This example describes the syntheses of 5'-PC-amino-oligonucleotides. The reaction was carried out using a commercially available synthesizer in the Trityl-On mode. Coupling efficiency for PC-aminotag phosphoramidites was determined by HPLC analysis of several sequences and found to be in the range 95-98%. Standard cleavage and

deprotection procedures were used. The composition of 5'-PC-amino-(dT)₇ and 5'-PC-amino-X-(dT)₇ was verified before and after photocleavage reaction by MALDI-TOF: 5'-PC-amino-(dT)₇ – 2,326.3 (uncleaved) 2,148.4 (cleaved); 5'-PC-amino-X-(dT)₇ 2,438.4 (uncleaved) 2,147.2 (cleaved). Control 5'-phosphorylated sequences were synthesised using the chemical phosphorylation reagent Phosphalink" (Applied Biosystem, Foster City, CA) according to the manufacture's instructions.

Example 6

This example describes the reaction of the oligonucleotides of Example 5 with marker molecules. Crude 5'-PC-amino-d(T)₇ or 5'-PC-amino-X-d(T)₇ (5.0 OD) was dissolved in 200 µl of 0.1 N NaHCO₃ (pH = 8.5). To this solution was added 100 µl of the NHS ester (biotin, digoxigenin or rhodamine) solution in anhydrous DMF (10 mg/ml). The solution was incubated overnight at room temperature, then it was purified by gel filtration (Sephadex G-25, NAP-10 column, Pharmacia, Piscataway, NJ), freeze dried and purified to homogeneity on RP HPLC. For photocleavage studies, a solution of 0.2 OD of oligonucleotides in 50 µl of H₂O was irradiated for 5 minutes in clear polypropylene microcentrifuge tube and a 20 µl aliquot was then analysed by HPLC.

Example 7

This example describes reactions with activated supports. Two oligonucleotides, 44-mer(5'-H₂N-PC-GGC-CGC-TAC-CTC-TCA-(CCA)6-CTG-ATG-AGC-T-3' - SEQ ID NO:1) and 35-mer (5'-H₂N-PC-CAT-CAG-(TGG)6-TGA-GAG-GTA-GC-3' - SEQ ID NO:2) were synthesised at 0.2 µmol scale with 5'-PC-aminotag modifier. These were cleaved and deprotected using standard procedures (28% NH₃, 1 hr RT, 18 hrs 55°C) and then freeze-dried. 6-aminohexanoic acid-NHS agarose (100mg) (Sigma, St Louis, MO) was suspended in 300 µl of water, vortexed briefly and spin-filtered using a 0.22 µm Ultrafree MC filter (Millipore, Bedford, MA), then resuspended in 300 µl of water. Crude 5'-PC-amino-oligonucleotide (10.0 OD) was added to a suspension of activated agarose, followed by 40 µl of 1M NaHCO₃. The mixture was vortexed at room temperature for 1 hour, spin-filtered and washed with 100 µl of wash buffer (2M NaCl, EDTA, pH = 8.0). Finally, the



beads were resuspended in 300 μ l of wash buffer and irradiated with near UV light for 5 minutes. After irradiation, the suspension was spin-filtered and the beads were washed with wash buffer (100 μ l). The combined filtrates were then desalted on a NAP-10 column, analysed by UV absorption spectroscopy and by denaturing PAGE.

Example 8

This example describes the detection of a peptide mass marker. All DNA syntheses were performed on an Expedite 8909 synthesiser (PerSeptive Biosystems, Framingham, MA) using phosphoramidite chemistry at either 0.2 or 1.0 μ mol scale. For the preparation of photocleavable peptide-DNA mass markers, DNA was assembled in the synthesiser at 1.0 μ mol scale and modified on the 5'-end with PC-Aminotag phosphoramidite. The Fmoc protective group was then removed by brief treatment with tetramethylguanidine, the support was washed with acetonitrile, dried and coupled with activated, Fmoc protected peptide for 2 h. The support was washed with DMF and acetonitrile, dried and then deprotected with ammonia and freeze-dried. The conjugate was purified to homogeneity using RP-HPLC. For this presentation 3 different mass markers were synthesised (see Table 1).

Table 1

#	Composition of Mass Marker	Mass Tag (M+H) ⁺	Intact Mass Marker (M+H) ⁺
1	H ₂ N-GRALGVFV-PC-5'-CAC-GTA-CAG-GAT-GTA-CAG-3' (SEQ ID NO:3)	1095.4 Da	6707.0 Da
2	H ₂ N-GRALGVFL-PC-5'-TCG-AGA-GGT-ACA-TCG-TG-3' (SEQ ID NO:4)	1109.4 Da	6438.8 Da
3	H ₂ N-GRALGVFF-PC-5'-AAG-CGG-TAC-GAG-TAG-CA-3' (SEQ ID NO:5)	1143.4 Da	6490.9 Da

Target DNA was synthesised on DMT-C27 beads (Penninsula Laboratories, CA). An aliquot containing approximately 4 nmol of target DNA was taken from the bead suspension and mixed with 10 pmol of the PC mass markers. Hybridisation was carried out in hybridisation buffer (25 mM NaCl, 20 mM Tris-HCl, 10 mM MgCl₂, pH = 7.5) under the following conditions: denaturation: 72°C, 10 min., annealing: 72-37°C over 30 min. Samples were incubated for an additional 30 min. at 37°C, centrifuged, washed with ice-cold water (3 x 100 μ l) and resuspended in 50 μ l water.



MALDI-MS measurements on immobilized DNA target sequences were carried out on a Vision 2000 reflectron-TOF mass spectrometer: ($U_{acc} = 5$ kV, eq. flight path length 1.7 m, N_2 laser, $\lambda = 337$ nm, $\tau = 3$ ns, SEM detector). Solution spectra were obtained on a laboratory built linear TOF-MS instrument equipped with delayed ion extraction ($U_{acc} = 25$ kV, equivalent flight path length 12.1 m, Nd:YAG laser, $\lambda = 355$ nm, $\tau = 5$ ns, additional Er:YAG laser, $\lambda = 2.94$ μ m, $\tau = 80$ ns, MCP detector) 3-hydroxypicolinic acid and 2,5-dihydroxybenzoic acid were used as matrices for UV-MALDI measurements; for IR-MALDI measurements, succinic acid was used. For measurements on immobilized DNA, a 0.5 - 1 μ l aliquot of the bead suspension was codeposited on the target with 1-2 μ l of matrix solution (dried droplet preparation).

In UV-MALDI mass spectra obtained from solutions of each of the mass markers, signals corresponding to the released peptide mass tag, the cleaved DNA and the intact conjugate were observed. In control experiments with IR-MALDI, only a signal corresponding to the intact mass marker and a very weak signal corresponding to the cleaved DNA sequence are detected, while signals of the peptide mass tag are absent. This proves the postulated photocleavage mechanism.

Signals corresponding to mass tags released from mass markers hybridized to immobilized DNA target sequences were readily obtained in UV-MALDI. In a competition experiment, an equimolar mixture of the 3 mass markers was incubated with beads presenting a DNA sequence complementary to the hybridization probe regions of one of them. Only the mass tag corresponding to the target sequence was detected in UV-MALDI mass spectra obtained from these beads, demonstrating the selectivity of hybridization.

In a different experiment evaluating the possibility of multiplex detection of several target sequences an immobilized strand containing the three target sequences with two nucleotides in between was incubated with an equimolar solution of the 3 mass markers. Signals of all three mass markers were detected in a UV-MALDI mass spectrum obtained from these beads. Relative signal intensities were found not to vary with the arrangement of the target sequences within the immobilized strand.

EXAMPLE 9

Figure 9 schematically shows one embodiment of a synthesis scheme for a novel photocleavable aminotag phosphoramidite. The present invention contemplates utilizing such novel phosphoramidites in the synthesis of oligonucleotides containing such photocleavable phosphoramidites.

Briefly, acetovanillone (1) (4.1 g; 24.7 mmol) was dissolved in 20 ml of anhydrous DMF. To this solution was added methyl 4-bromobutyrate (5 g; 28 mmol) and finely powdered anhydrous K_2CO_3 (5.13 g 37 mmol). The resulting slurry was stirred at room temperature overnight. The reaction mixture was partitioned between brine and EtOAc, and acidified with 1N HCl. The organic layer was dried and evaporated to give (2) as yellowish oil (~6.8g).

Compound 2 (6.8 g) was dissolved in 22.5 ml of 2:1 pyridine:H₂O (v/v) and to the resulting solution hydroxylamine hydrochloride (2.14 g; 31 mmol) was added. The solution was stirred at room temperature overnight and then partitioned between saturated NaCl and EtOAc. Organic phase was dried and evaporated to give the oxime (3) as a creamy solid (7.1 g).

The oxime (3) (8.45 g; 30.04 mmol) was dissolved in glacial acetic acid (48 ml) and to this solution was added 10% Pd/C (300 mg). The slurry was placed under vacuum for 5 min. and then connected to the hydrogen tank with ~1 atm (15 psi) pressure and stirred overnight. Another 300 mg of 10% Pd/C was added and the mixture stirred under hydrogen for another 24 hrs. The Pd/C was filtered off and the filtrate acidified with 1.2 eq. of 1N HCl, followed by extraction with $CHCl_3$. Filtrate was discarded and the aqueous layer evaporated to dryness under reduced pressure to give amine (4) as hydrochloride salt.

The amine (4) (3.66 g; 13.6 mmol) was dissolved in 28 ml of pyridine and cooled in an ice bath. To this solution trifluoroacetic anhydride (2.9 ml; 20.5 mmol) was added gradually over a period of 5 minutes. The reaction mixture was stirred for additional 30 min., and then partitioned between saline and EtOAc. Organic layer was dried and evaporated to dryness to give the trifluoroacetamide (5) (4.64 g; 96.5% yield).

Compound 5 (4.58 g) was added to 100 ml of 70% HNO_3 at 0°C and the resulting

solution was stirred at °C for 2 hrs, then poured onto 1 kg of ice. The precipitate formed was filtered off, washed with ice-cold water and dried to give 3.50 g of compound 6.

Compound 6 (1.84 g; 4.5 mmol) was suspended in 50 ml MeOH, and sodium borohydride was added in several portions (total 4.7 g) over 5 hrs. 100 ml of water was added and the resulting solution was stirred overnight, then acidified to pH = 1.0 and extracted with chloroform. Organic layer was discarded, aqueous phase was basified with 1N NaOH and extracted again with chloroform. Organic layer was dried and evaporated to give 410 mg of amine (7) (32% yield).

Compound 7 (410 mg, 1.44 mmol) was dissolved in 1:1 acetonitrile:water (v/v; 10 ml) and then 1 eq. (199 ul) of triethylamine was added.. Then a solution of Fmoc-NHS (584 mg; 1.2 eq.) in 3.5 ml acetonitrile was added during 5 minutes. The reaction mixture was stirred for additional 20 min., the precipitate formed was filtered off, washed with 1 N HCl, water and dried in vacuum to give Fmoc derivative 8. ¹H NMR (DMSO-d₆) 8.02, 8.00 (d); 7.86, 7.84 (d); 7.63, 7.61 (d); 7.46 (s); 7.40, 7.38, 7.36 (t), 7.29, 7.27, 7.25 (t); 7.23 (s); 5.21, 5.19, 5.17 (t); 4.40, 4.39, 4.37(t); 4.27 (m), 4.17 (m), 4.05 (m), 3.85 (s), 3.59-3.43 (q); 1.75 (t); 1.54 (t); 1.41 (d).

Compound 8, (800 mg; 1.58 mmol) was dissolved in an anhydrous acetonitrile (7.5 ml) and to this solution N,N-diisopropylethylamine (1.07 ml; 6.33 mmol) was added. To this solution 2-cyanoethyl-N,N-diisopropyl-chlorophosphine (390 ul x 3, 3.3 equiv. total) was added and the reaction stirred for 2 hrs. Then 3 ml of EtOAc was added and the reaction mixture was partitioned between saline and EtOAc. Organic layer was dried, evaporated to dryness and purified on a silica gel using step gradient of hexane/EtOAc/triethylamine to give 160 mg of the final product (compound 9). ¹H NMR (DMSO-d₆) 8.02, 8.00 (d); 7.86, 7.84 (d); 7.63, 7.61 (d); 7.46 (s); 7.40, 7.38, 7.36 (t), 7.29, 7.27, 7.25 (t); 7.23 (s); 5.21, 5.19, 5.17 (t); 4.40, 4.39, 4.37(t); 4.26 (m), 4.08 (m); 4.17 (m), 4.05 (m), 3.84 (s), 3.59-3.40 (m); 2.88-2.56 (m) 1.75 (m); 1.54 (t); 1.41 (d); 1.23-1.08 (m).

It is not intended that the above synthesis scheme be limited to a particular protecting group. Figure 10 shows compound 9 of Figure 9 where the protecting group is indicated generically as "PG".

EDITORIAL NOTE - NO 62382/99

Sequence listing pages 1-2 is part of the description.

The claims are to follow.

SEQUENCE LISTING

<110> Rothschild, Kenneth J.
Olejniak, Jerzy

<120> Compounds and Methods for Detecting Biomolecules

<130> AMBER-03888

<140> 09/323,424

<141> 1999-06-01

<150> 60/087,641

<151> 1998-06-02

<160> 6

<170> PatentIn Ver. 2.0

<210> 1

<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 1
cgccgctac ctctcaccac caccaccacc accactgatg agct 44

<210> 2
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic

<400> 2
ccatcagtggt tggtggtggt ggtggtgaga ggtagc 36

<210> 3
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic

<400> 3
cacgtacagg atgtacag 18

<210> 4
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic

<400> 4
tcgagaggta catcgtg 17



<210> 5
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic

<400> 5
aagcggtagc agtagca



17

<210> 6
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic



<400> 6
tcgagaggta catcgtg

17



The claims defining the invention are as follows:

1. A composition, comprising a photocleavable aminotag phosphoramidite selected from the group of compounds A and B of Figure 8 and the compound of Figure 10.
- 5 2. The composition of claim 1, wherein said photocleavable aminotag phosphoramidite is incorporated at the 5' end of an oligonucleotide.
3. The composition of claim 2, wherein said oligonucleotide is immobilized.
4. The composition of claim 3, wherein said oligonucleotide is immobilized via said aminotag.
- 10 5. The composition of claim 3, wherein said oligonucleotide is immobilized to an activated surface.
6. The composition of claim 5, wherein said surface is the surface of a bead.
7. The composition of claim 2, wherein a marker molecule is linked to said aminotag phosphoramidite.
- 15 8. The composition of claim 7, wherein said marker molecule is selected from the group consisting of activated polypeptides, activated biotin, activated digoxigenin, and activated tetramethylrhodamine.
9. The composition of claim 7, wherein said marker is detectable by spectrometry.
- 20 10. The composition of claim 9, wherein said marker is detectable by mass spectrometry.
11. A composition, comprising a photocleavable aminotag phosphoramidite selected from the group of compounds 6 and 7 of Figure 2 and compound 9 of Figure 9.
12. The composition of claim 11, wherein said photocleavable aminotag
25 phosphoramidite is incorporated at the 5' end of an oligonucleotide.
13. The composition of claim 12, wherein said oligonucleotide is immobilized.
14. The composition of claim 13, wherein said oligonucleotide is immobilized via said aminotag.
15. The composition of claim 13, wherein said oligonucleotide is immobilized to
30 an activated surface.
16. The composition of claim 15, wherein said surface is the surface of a bead.
17. The composition of claim 12, wherein a ~~marker~~ molecule is linked to said aminotag phosphoramidite.



18. The composition of claim 17, wherein said marker molecule is selected from the group consisting of activated polypeptides, activated biotin, activated digoxigenin, and activated tetramethylrhodamine.

19. The composition of claim 17, wherein said marker is detectable by spectrometry.

20. The composition of claim 19, wherein said marker is detectable by mass spectrometry.

21. A composition, comprising the photocleavable sulfhydryl phosphoramidite set forth in Figure 4.

22. The composition of claim 21, wherein said photocleavable sulfhydryltag phosphoramidite is incorporated at the 5' end of an oligonucleotide.

23. The composition of claim 22, wherein said oligonucleotide is immobilized.

24. The composition of claim 23, wherein said oligonucleotide is immobilized via said sulfhydryltag.

25. The composition of claim 23, wherein said oligonucleotide is immobilized to a surface.

26. The composition of claim 25, wherein said surface is a gold surface.

27. The composition of claim 22, wherein a marker molecule is linked to said sulfhydryltag phosphoramidite.

28. The composition of claim 27, wherein said marker molecule is selected from the group consisting of polypeptides, biotin, digoxigenin, and tetramethylrhodamine.

29. The composition of claim 27, wherein said marker is detectable by spectrometry.

30. The composition of claim 29, wherein said marker is detectable by mass spectrometry.




31. A method, comprising:

a) providing:

i) a compound comprising a photocleavable phosphoramidite and a protected tag linked to said photocleavable phosphoramidite, wherein said protected tag is selected from the group consisting of an amino tag, a sulfhydryl tag, a nucleotide tag and a mass tag,

ii) oligonucleotide synthesis means; and

b) synthesising an oligonucleotide with said synthesis means and said photocleavable phosphoramidite under conditions such that an oligonucleotide having a 3'



end and a 5' end is produced wherein said photocleavable phosphoramidite is incorporated at said 5' end.

32. The method of claim 31, wherein said photocleavable phosphoramidite is an aminotag phosphoramidite protected by an 9-fluorenylmethoxycarbonyl group.

5 33. The method of claim 32, wherein said photocleavable aminotag phosphoramidite is selected from compounds 6 and 7 of Figure 2 and compound 9 of Figure 9.

34. The method of claim 31, wherein said photocleavable phosphoramidite is a sulfhydryltag phosphoramidite protected by a trityl group.

10 35. The method of claim 34, wherein said sulfhydryl tag phosphoramidite is set forth in Figure 4.

36. A method of preparing a composition comprising a photocleavable phosphoramidite, substantially as hereinbefore described with reference to any one of the examples.

15 37. A composition comprising a photocleavable phosphoramidite, prepared by a method of any one of claims 31 to 36.

38. A composition comprising a photocleavable phosphoramidite, substantially as hereinbefore described with reference to any one of the examples.

39. A method, comprising:

20 a) providing:

i) a compound comprising a photocleavable phosphoramidite and a protected tag linked to said photocleavable phosphoramidite, wherein said protected tag is selected from the group consisting of an amino tag, a sulfhydryl tag, nucleotide tag and a mass tag,

25 ii) oligonucleotide synthesis means,

iii) a marker molecule,

iv) a source of electromagnetic radiation;

b) synthesising an oligonucleotide with said synthesis means and said photocleavable phosphoramidite under conditions such that an oligonucleotide having a 3' end and a 5' end is produced wherein said photocleavable phosphoramidite is incorporated at said 5' end;

c) linking said marker molecule to said synthesised oligonucleotide of step (b) so as to produce a photocleavable marker/oligonucleotide complex; and

d) exposing said complex to electromagnetic radiation from said source of electromagnetic radiation under conditions such that said oligonucleotide is released from said marker.

40. The method of claim 39, wherein said photocleavable phosphoramidite is an aminotag phosphoramidite protected by a 9-fluorenylmethoxy carbonyl group.

41. The method of claim 40, wherein said photocleavable aminotag phosphoramidite is selected from compounds 6 and 7 of Figure 2 and compound 9 of Figure 9.

42. The method of claim 39, wherein said photocleavable phosphoramidite is a sulfhydryltag phosphoramidite protected by a trityl group.

43. The method of claim 42, wherein said sulfhydryltag phosphoramidite is set forth in Figure 4.

44. A method as defined in claim 39, substantially as hereinbefore described with reference to any one of the examples.

45. An oligonucleotide obtained by a method according to any one of claims 39 to 44.

46. A method, comprising:

a) providing:

i) a compound comprising photocleavable phosphoramidite, and a protected tag linked to said photocleavable phosphoramidite, wherein said protected tag is selected from the group consisting of an amino tag, a sulfhydryltag, nucleotide tag and a mass tag;

ii) oligonucleotide synthesis means,



iii) a surface,

iv) a source of electromagnetic radiation;

b) synthesising an oligonucleotide with said synthesis means and said photocleavable phosphoramidite under conditions such that an oligonucleotide having a 3' end and a 5' end is produced wherein said photocleavable phosphoramidite is incorporated at said 5' end;

c) attaching said synthesised oligonucleotide of step (b) to a surface to create an immobilised oligonucleotide; and

d) exposing said immobilised oligonucleotide to electromagnetic radiation from said source of electromagnetic radiation under conditions such that said oligonucleotide is released from said surface.



47. The method of claim 46, wherein said photocleavable phosphoramidite is an aminotag phosphoramidite protected by a 9-fluorenylmethoxycarbonyl group.

48. The method of claim 47, wherein said photocleavable aminotag phosphoramidite is selected from compounds 6 and 7 of Figure 2 and compound 9 of Figure 9.

49. The method of claim 46, wherein said photocleavable phosphoramidite is a sulfhydryltag phosphoramidite protected by a trityl group.

50. The method of claim 49, wherein said sulfhydryltag phosphoramidite is set forth in Figure 4.

51. A method as defined in claim 46, substantially as hereinbefore described with reference to any one of the examples.

52. An oligonucleotide obtained by a method according to any one of claims 46 to 51.

Dated 10 February, 2003

Ambergen, Inc.

Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON





Figure 1

Tag = biotin, NH_2 , SH

Figure 2




Figure 3





Figure 4



201200 62382

Figure 5




Figure 6







Figure 7





A



B

Figure 8

Figure 9




Figure 10

