

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Substances for preventing and treating autoimmune diseases

(51)⁶ International Patent Classification(s)
C12N 5/00 101BHAU
(2006.01) PCT/US2003/02462
C12N 5/00 5
20060101AFI20060

(21) Application No: **2003296892** (22) Application Date: **2003.08.06**

(87) WIPO No: **WO04/034966**

(30) Priority Data

(31) Number (32) Date (33) Country
60/401,652 2002.08.06 US

(43) Publication Date : **2004.05.04**
(43) Publication Journal Date : **2004.06.10**

(71) Applicant(s)
Loma Linda University

(72) Inventor(s)
Escher, Alan P.; Li, Fengchun

(74) Agent/Attorney
**Freehills Patent & Trade Mark Attorneys, Level 43 101 Collins Street,
Melbourne, VIC, 3000**

(56) Related Art
Rabinovitch et al. Diabetes 1999 June; 48:1223-1229

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 April 2004 (29.04.2004)

PCT

(10) International Publication Number
WO 2004/034966 A3

- (51) International Patent Classification⁷: C12N 5/00, 15/00, 15/87, A61K 48/00
- (21) International Application Number: PCT/US2003/024625
- (22) International Filing Date: 6 August 2003 (06.08.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/401,652 6 August 2002 (06.08.2002) US
- (71) Applicant (for all designated States except US): LOMA LINDA UNIVERSITY [US/US]; 2488 Prospect Street, Loma Linda, CA 92354 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): ESCHIER, Alan, P. [US/US]; 463 Jefferson Street, Redlands, CA 92374 (US). LI, Fengchun [CN/US]; 24530 University Avenue, Apt. 10, Loma Linda, CA 92354-2732 (US).
- (74) Agents: FARAH, David, A. et al.; Sheldon & Mak, PC, 225 South Lake Avenue, 9th Floor, Pasadena, CA 91101 (US).
- (81) Designated States (national): AL, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
- with international search report
 - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- (88) Date of publication of the international search report: 8 July 2004
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2004/034966 A3

(54) Title: SUBSTANCES FOR PREVENTING AND TREATING AUTOIMMUNE DISEASES

(57) Abstract: A substance for preventing, delaying the onset of, or treating one or more than one autoimmune disease, the substance comprising a polynucleotide construct comprising a polynucleotide sequence encoding the pro-apoptotic protein BAX and encoding one or more than one autoantigen for the autoimmune disease. A method for preventing, delaying the onset of or treating an autoimmune disease in a patient comprising selecting a patient who is susceptible to developing the autoimmune disease, who is developing the autoimmune disease or who has the autoimmune disease and administering to the patient one or more than one dose of a polynucleotide construct comprising a polynucleotide sequence encoding the pro-apoptotic protein BAX and encoding one or more than one autoantigen for an autoimmune disease, or comprising a polynucleotide sequence encoding the adenoviral protein E3-GP19k, or comprising a polynucleotide sequence encoding Δ BCL-2.

SUBSTANCES FOR PREVENTING AND TREATING AUTOIMMUNE DISEASES**STATEMENT REGARDING FEDERALLY SPONSORED
RESEARCH OR DEVELOPMENT**

This invention was made with United States Government support under Cooperative Agreement Number DAMD-17-97-2-7016 with the National Medical Technology Testbed, Inc., United States Department of the Army. The United States Government has certain rights in this invention.

CROSS-REFERENCE TO RELATED APPLICATION

The present Application claims the benefit of United States provisional patent application 60/401,652, titled "Method and Substances for the Suppression of Diabetes," filed August 6, 2002, the contents of which are incorporated in this disclosure by reference in their entirety.

BACKGROUND

Autoimmune diseases cause significant human morbidity and mortality. These diseases include approximately 80 diseases, such as rheumatoid arthritis, systemic lupus and multiple sclerosis, and affect approximately 5% of the population of the United States. One autoimmune disease, type 1 diabetes, is the most frequent chronic disease in children, and has a steadily increasing worldwide incidence.

Generally, the onset of type 1 diabetes begins with the display by antigen presenting cells (APCs) of autoantigens synthesized by pancreatic beta cells. This display results in the immune system destruction of pancreatic beta cells mediated mostly by T helper 1 (Th1) and cytotoxic T lymphocytes and, thereby, to the loss of insulin production.

Many prophylactic and therapeutic approaches for type 1 diabetes attempt to prevent the destruction of beta cells by inducing the immune system to delete, inactivate or suppress pathogenic self-reactive lymphocytes, such as by administering vaccines that solely deliver autoantigen, or by administering substances are direct effectors of the immune system, such as cytokines. However, currently available DNA-based vaccines are not completely efficient in preventing the disease, and the use of some of these vaccines are associated with inducing or enhancing autoimmunity rather than preventing the disease. Additionally, the use of cytokines is associated with significant morbidity.

Therefore, there is a need for a new method for preventing, delaying the onset of, or

treating autoimmune diseases using vaccines that are not associated with these disadvantages. Further, there is a need for a new method for preventing, delaying the onset of, or treating type 1 diabetes using vaccines that are not associated with these disadvantages.

SUMMARY

5 According to one embodiment of the present invention, there is provided a substance for preventing, delaying the onset of or treating one or more than one autoimmune disease. The substance comprises a polynucleotide construct comprising a polynucleotide sequence encoding the pro-apoptotic protein BAX and encoding one or more than one autoantigen for the autoimmune disease.

10 According to another embodiment of the present invention, there is provided a use of a polynucleotide construct comprising a polynucleotide sequence encoding the pro-apoptotic protein BAX and encoding one or more than one autoantigen for an autoimmune disease for the manufacture of a medicament for preventing, delaying the onset of or treating the one or more than one autoimmune disease.

15 According to another embodiment of the present invention, there is provided a use of a polynucleotide construct comprising a polynucleotide sequence encoding the adenoviral protein E3-GP19k for the manufacture of a medicament for preventing, delaying the onset of or treating one or more than one autoimmune disease.

20 According to another embodiment of the present invention, there is provided a use of a polynucleotide construct comprising a polynucleotide sequence encoding Δ BCL-2 for the manufacture of a medicament for preventing, delaying the onset of or treating one or more than one autoimmune disease.

25 In one embodiment, the medicament is manufactured in dosage units of between about 0.5 mg to about 5 mg. In another embodiment, the medicament is manufactured in dosage units of between about 1 mg to about 4 mg. In another embodiment, the medicament is manufactured in dosage units of between about 2.5 mg to about 3 mg. In another embodiment, the medicament is manufactured in a form suitable for intramuscular administration. In another embodiment, the medicament is manufactured in a form suitable for intravenous administration.

30 According to another embodiment of the present invention, there is provided a method for preventing, delaying the onset of or treating an autoimmune disease in a patient. The method comprises selecting a patient who is susceptible to developing the autoimmune

disease, who is developing the autoimmune disease or who has the autoimmune disease; and administering to the patient one or more than one dose of a polynucleotide construct comprising a polynucleotide sequence encoding the pro-apoptotic protein BAX and encoding one or more than one autoantigen for the autoimmune disease, or a polynucleotide construct comprising a polynucleotide sequence encoding the adenoviral protein E3-GP19k, or a polynucleotide construct comprising a polynucleotide sequence encoding Δ BCL-2, or a combination of the preceding polynucleotide constructs.

In one embodiment, the autoimmune disease is type I diabetes. In another embodiment, selecting the patient comprises identifying in the patient the presence of anti-insulin or anti-GAD autoantibodies or both anti-insulin and anti-GAD autoantibodies. In another embodiment, selecting the patient comprises identifying in the patient the presence of increasing hyperglycemia. In another embodiment, selecting the patient comprises identifying in the patient the presence of glycosuria. In another embodiment, selecting the patient comprises identifying in the patient the presence of a genetic predisposition to the autoimmune disease.

In another embodiment, the one or more than one dose is a plurality of doses. In another embodiment, administering to the patient one or more than one dose comprises injecting the patient intramuscularly with the one or more than one dose. In another embodiment, the method further comprises, after administering, monitoring the patient for the development of the autoimmune disease.

FIGURES

These and other features, aspects and advantages of the present invention will become better understood with regard to the following description, appended claims, and accompanying figures where:

25 Figure 1 are schematic depictions of three substances according to the present invention; and

Figure 2 are schematic depictions of the fifteen plasmids that were tested for their efficiency in preventing, delaying the onset of or treating an autoimmune disease in accordance with a method of the present invention.

30 **DESCRIPTION**

According to one embodiment of the present invention, there are provided substances for preventing, delaying the onset of or treating one or more than one autoimmune disease.

According to another embodiment of the present invention, there is provided a method of preventing, delaying the onset of or treating one or more than one autoimmune disease. In one embodiment, the autoimmune disease is type 1 diabetes. In a preferred embodiment, the method comprising using a substance according to the present invention is a vaccine. The substances and method of the present invention do not use solely the delivery of autoantigen, and do not use molecules that are direct effectors of the immune system as in prior methods. Instead, the present invention uses a vaccine to prevent apoptosis of one or more than one type of cell capable of suppressing the autoimmune disease. Because these one or more than one type of cell capable of suppressing the autoimmune disease are still be subject to physiological and immune regulation, the risk of inducing or enhancing autoimmunity is greatly reduced by the present method as compared to some prior art methods. Further, because the present invention does not involve administering substances that are direct effectors of the immune system, such as cytokines, the present invention does not pose the risk side effects associated with such direct effectors of the immune system. Further advantageously, a genetic vaccine comprising primarily plasmid DNA can be produced in large quantities at relatively low cost and does not require a "cold chain" for storage. Therefore, the substances and methods according to the present invention are both economical and practical for use to prevent, delay the onset of or treat an autoimmune disease. Further, a genetic vaccine according to the present invention modifies the genetic material of an organism directly which means that native epitopes will be processed by the organism's immune system unlike protein-based vaccines. The substances and method of the present invention will now be disclosed in detail.

As used in this disclosure, the term "autoimmune disease" comprises both diseases due in part or in total to destruction of normal cells or tissues by the organism's own immune system, and also comprises destruction of cells or tissues that were transplanted into the organism to take the place of defective or absent cells or tissues, such as islet cell transplants, or partial or whole organ transplants, by the organism's own immune system.

As used in this disclosure, the term "comprise" and variations of the term, such as "comprising" and "comprises," are not intended to exclude other additives, components, integers or steps.

In one embodiment, the present invention includes three substances that can be used either individually, sequentially or simultaneously to prevent, delay the onset of or treat one

or more than one autoimmune disease. One of the three substances is a DNA construct comprising a polynucleotide sequence, SEQ ID NO:1, encoding the pro-apoptotic protein BAX, and encoding one or more than one autoantigen for the autoimmune disease. Another of the three substances is a DNA construct comprising a polynucleotide sequence, SEQ ID NO:2, encoding the adenoviral protein E3-GP19k, which prevents presentation of an antigen on MHC-I molecules in the endoplasmic reticulum. Another of the three substances is a DNA construct comprising a polynucleotide sequence, SEQ ID NO:3, encoding a truncated form of BCL-2 designated Δ BCL-2 in this disclosure.

As will be understood by those with skill in the art with reference to this disclosure, though specific sequences are given for the polynucleotide sequences as disclosed in this disclosure, such as the polynucleotide sequences encoding the pro-apoptotic protein BAX, the adenoviral protein E3-GP19k and Δ BCL-2, the present invention includes any other sequence that does not cause a change in the translated amino acid sequence, as well as any sequence that does cause a change in the translated amino acid sequence but where the change does not substantially affect the function of the translated amino acid sequence so as to make it unsuitable for the uses contemplated in this disclosure.

Referring now to Figure 1, there are shown schematic depictions of three substances according to the present invention. As can be seen, each substance comprises a plasmid DNA construct. Substance A comprises a plasmid construct comprising a polynucleotide encoding an autoantigen for the autoimmune disease, such as secreted glutamic acid decarboxylase that is an autoantigen for type 1 diabetes, followed by a polynucleotide, SEQ ID NO:1, encoding BAX. Substance B comprises a plasmid construct comprising a polynucleotide, SEQ ID NO:2, encoding E3-GP19k without a polynucleotide encoding an autoantigen for the autoimmune disease. Substance C comprises a plasmid construct comprising a polynucleotide, SEQ ID NO:3, encoding a truncated form of the anti-apoptotic protein BCL-2 without a polynucleotide encoding an autoantigen for the autoimmune disease. As used in the Figures, "CMV" represents the cytomegalovirus promoter element, "pA" represents a polyadenylation site, and "IRES" represents an internal ribosome binding site from the EMCV virus, SEQ ID NO:4.

In order to demonstrate the advantages of the present invention, fifteen plasmids were constructed and used as vaccines. Each construct was cloned into the vector pND2. Referring now to Figure 2, there are shown schematic depictions of the fifteen plasmids that

were tested for their efficiency in preventing, delaying the onset of or treating an autoimmune disease. As can be seen, each plasmid was under the plasmid transcriptional control of the same promoter (CMVp) to ensure expression of both open reading frames in each transfected cells. During construction of these plasmids containing the cDNA encoding BCL-2, it was found that plasmid deletions occurred due to the large size of the cDNA. Therefore, a truncated version of *bcl-2* designated $\Delta bcl-2$ was used to construct the plasmids. As shown in Figure 2, the plasmids comprised cDNA encoding cytoplasmic GAD, SEQ ID NO:5, (plasmid 1); secreted GAD (SGAD), SEQ ID NO:6, (plasmid 2); a control secreted luciferase, SEQ ID NO:7, (plasmid 3); truncated human anti-apoptotic protein BCL-2 ($\Delta BCL-2$), SEQ ID NO:3, (plasmid 4); anti-apoptotic protein BAX, SEQ ID NO:1, (plasmid 5); E3-GP19k, SEQ ID NO:2, (plasmid 6); $\Delta BCL-2$, SEQ ID NO:3, in combination with cytoplasmic GAD, SEQ ID NO:5, secreted GAD, SEQ ID NO:6, and secreted luciferase, SEQ ID NO:7, (plasmids 7-9, respectively), BAX, SEQ ID NO:1, in combination with cytoplasmic GAD, SEQ ID NO:5, secreted GAD, SEQ ID NO:6, and secreted luciferase, SEQ ID NO:7, (plasmids 10-12, respectively); and E3-GP19k, SEQ ID NO:2, in combination with cytoplasmic GAD, SEQ ID NO:5, secreted GAD, SEQ ID NO:6, and secreted luciferase, SEQ ID NO:7, (plasmids 13-15, respectively).

All plasmids were generated, the open reading frame amplified using PCR, and the amplification products were inspected after DNA sequencing and found to be without mutations. Each construct was then used to transfect simian COS-7 cells transiently for immunoblot analysis of cell lysates, which confirmed that a gene product of the correct size was encoded (data not shown).

Next, the effects of the 15 plasmids on non-obese diabetic (NOD) mice were determined as follows. First, plasmid DNA was isolated using Qiagen Endofree kits (Qiagen Inc., Chatsworth, CA, US), and 300 ug of each of the 15 plasmid DNAs was injected intramuscularly into groups of fifteen 4-5-week-old female NOD mice. The 300 ug dose was selected as a dose relevant to the human clinical setting based on organism weight. The onset of diabetes was monitored until the age of 35 weeks, using urine and blood glucose analysis. The mice were considered diabetic after testing positive for high levels of glycosuria, with blood glucose levels greater than 300 mg/dl on two consecutive days.

The results of these experiments demonstrated the following. The percentage of diabetic animals at 35 weeks of age ranged from 73-93% for mice vaccinated with plasmids

- 1-3; 60-67% for mice vaccinated with plasmids 4 or 7-9; 47-85% for mice vaccinated with plasmids 5 and 10-12; and 53-73% for mice vaccinated with plasmids 6 and 13-15. Control animals (those not vaccinated) had an incidence of diabetes of about 93%. Therefore, administration of 300 ug of plasmid vector alone or of 300 ug of plasmid vector encoding
- 5 antigens alone, plasmids 1-3, did not result in significant diabetes suppression. Mice vaccinated with plasmids 6-9 and 11 showed statistically significant suppression of diabetes when compared to untreated mice ($P < 0.05$ for plasmid 7, and $P < 0.02$ for plasmid 9). In addition, mice receiving pND2-E3-GP19k, plasmid 6 or pND2-SGAD55-BAX, plasmid 11 showed a significantly decreased incidence of diabetes at 35 weeks when compared to mice
- 10 receiving plasmid pND2-GAD65, plasmid 1 or pND2-GAD65-BAX, plasmid 10 ($P < 0.04$), and mice receiving pND2-GAD65- Δ BCL2, plasmid 7 or pND2-SGAD55- Δ BCL2, plasmid 8 showed significantly decreased diabetes when compared to mice receiving pND2-GAD65, plasmid 1 ($P < 0.05$). Suppression of diabetes was associated with decreased islet inflammation (data not shown). These results will be disclosed now in greater detail.
- 15 Mice that were vaccinated with plasmids comprising Δ bcl-2, plasmids 4 and 7-9, showed a 4-5 weeks delay in diabetes onset regardless of the co-expressed antigen, and a decrease in the incidence of diabetes at 35 weeks of age (60-67% compared to about 93% for the unvaccinated control mice) regardless of the co-expressed antigen. Therefore, co-expression of GAD autoantigen did not suppress the effect.
- 20 Mice that were vaccinated with plasmids comprising *bax*, plasmids 5 and 10-12, did not show diabetes suppression, with the exception of *sgad55-bax*, plasmid 11. While mice vaccinated with plasmid 11 started to develop diabetes at a time similar to other mice vaccinated with a plasmid comprising only *bax*, plasmid 5, the incidence of diabetes in mice vaccinated with plasmid 11 at 35 weeks of age was only 47% compared with a 93%
- 25 incidence for the unvaccinated control mice ($p < 0.05$).
- Mice that were vaccinated with plasmids comprising *E3-gp19k*, plasmids 6 and 13-15 showed wide differences in diabetes onset, depending on the antigen that was co-expressed. Mice that were vaccinated with the plasmid comprising *E3-gp19k* without autoantigen, plasmid 6 started to develop diabetes with a 4-5 week delay, and showed decreased diabetes
- 30 at 35 weeks of age (53% vs 93% for the unvaccinated control mice for control) ($p < 0.05$). Mice that were vaccinated with the plasmids comprising *E3-gp19k* with autoantigen, plasmids 13-15, suppressed the effect, both with respect to the delay in the onset of diabetes and with

respect to the incidence of diabetic animals at 35 weeks.

Next, immune responses were characterized using a GAD-specific ELISpot assay and ELISA of serum anti-GAD IgG isotypes to determine whether diabetes suppression by the administration of the substances of the present invention was associated with suppression of inflammatory Th1-like activity, and up-regulation of anti-inflammatory Th2 like response.

The ELISpot assay was conducted as follows. Splenocytes were isolated from the mice at time of diabetes onset, or at the end of the observation period for non-diabetic animals. The cells were then stimulated with recombinant GAD protein, and the number of cells secreting IFN-gamma (for Th1-like activity), and IL-4 (for Th2-like activity) were counted, following a standard manufacturer's protocol. The number of cells secreting the cytokines in the absence of GAD stimulation was then subtracted, and results analyzed. For IFN-gamma the data clearly indicated that suppression of diabetes by plasmid 6, encoding E3-GP19k alone, or by plasmids 4 and 7-9, encoding Δ BCL-2 alone or together with an antigen, were associated with a suppression of GAD-specific activity. Therefore, E3-19k and Δ BCL-2 could induce an immune response that was able to suppress autoractivity against beta cells. Surprisingly, the SGAD55-BAX combination did not appear to significantly suppress Th1-like activity. Further, SGAD55 alone, which did not suppress diabetes, did suppress GAD-specific Th1-like response.

With respect to IL-4, the data indicated an increase in GAD-specific activity for mice that received plasmid 6 encoding E3-GP19k alone (diabetes suppression), plasmid 13 encoding SGAD55 and E3-19k (no diabetes suppression), and plasmid 8 SGAD55 and Δ BCL-2(diabetes suppression). Thus, increased Th2-like activity was not always associated with decreased Th1-like activity or disease suppression.

The ELISA was conducted as follows. Animal sera were used for ELISA of anti-GAD IgG2a,b and IgG1 isotypes, which indicate a Th1-like and Th2-like activity, respectively. ELISA of anti-GAD IgG2a,b indicated that three of the plasmid DNAs coding for Δ BCL-2, plasmids 4, 8 and 9, showed a significant reduction in Th1-like activity, when compared to plasmid 5 coding for BAX, but not with the unvaccinated control mice. ELISA of anti-GAD IgG1 indicated that all plasmid DNAs encoding BAX, plasmids 5 and 10-12, resulted in decreased Th2-like activity.

These data taken together indicate that, first, *bax*, a plasmid cDNA coding for a pro-apoptotic protein, can be used as a molecular adjuvant for genetic vaccines to prevent

autoimmune disease, such as a vaccine comprising a polynucleotide encoding a secreted form of an autoantigen. Second, a plasmid cDNA encoding E3-GP19k or encoding a truncated BCL-2 alone could suppress autoimmune disease, though a plasmid cDNA encoding E3-GP19k or encoding a truncated BCL-2 combined with an autoantigen was less effective.

5 In one embodiment of the present invention, there is provided a method of preventing, delaying the onset of or treating an autoimmune disease. The method comprises, first, selecting a patient who is susceptible to developing the autoimmune disease, who is developing the autoimmune disease or who has the autoimmune disease. The selection can be made using standard methods as will be understood by those with skill in the art with
10 reference to this disclosure. For example, if the autoimmune disease is diabetes, the selection can be made by identifying in the patient the presence of anti-insulin or anti-GAD autoantibodies or both anti-insulin and/or anti-GAD autoantibodies, the presence of increasing hyperglycemia, the presence of glycosuria, the presence of a genetic predisposition to diabetes or more than one of these.

15 Next, the patient is administered one or more than one dose of a plasmid construct according to the present invention. That is, a plasmid construct comprising a polynucleotide encoding an autoantigen for the autoimmune disease and encoding BAX, or a plasmid construct comprising a polynucleotide encoding E3-GP19k but without a polynucleotide encoding an autoantigen for the autoimmune disease, or a plasmid construct comprising a
20 polynucleotide encoding a truncated form of the anti-apoptotic protein BCL-2 but without a polynucleotide encoding an autoantigen for the autoimmune disease. In a preferred embodiment, the organism is administered two plasmid constructs according to the present invention. In a particularly preferred embodiment, the organism is administered all three plasmid constructs according to the present invention.

25 In a preferred embodiment, the plasmid construct is administered in a plurality of doses. In another preferred embodiment, the dose is between about 0.001 mg/Kg and about 10 mg/Kg. In another preferred embodiment, the dose is between about 0.01 mg/Kg and about 1 mg/Kg. In another preferred embodiment, the dose is about 0.05 mg/Kg. In a preferred embodiment, a suitable dose for a human adult is between about 0.5 mg and 5 mg.
30 In a preferred embodiment, a suitable dose for a human adult is between about 1 mg and 4 mg. In a preferred embodiment, a suitable dose for a human adult is between about 2.5 mg and 3 mg. In another preferred embodiment, the dose is administered weekly between about

2003296892 21 Feb 2005

WO 2004/034966

PCT/US2003/024625

10

2 and about 10 times. In a particularly preferred embodiment, the dose is administered weekly 4 times. In another particularly preferred embodiment, the dose is administered only once.

Administration can be by a suitable route. In a preferred embodiment, the route is intramuscular or intravenous.

Additionally, the method can comprise, after administering, monitoring the patient for the development of the autoimmune disease.

EXAMPLE I

PREVENTION OF DIABETES

According to the present invention, the onset of diabetes in a patient is delayed or prevented, for example, as follows. First, the patient is selected based on the presence of circulating anti-insulin and anti-GAD autoantibodies. Next, the patient is injected intramuscularly with 0.05 mg/Kg of a plasmid construct comprising a polynucleotide sequence, SEQ ID NO:1, encoding the pro-apoptotic protein BAX and encoding SGAD, SEQ ID NO:6, or comprising a polynucleotide sequence, SEQ ID NO:2, encoding the adenoviral protein E3-GP19k, or comprising a polynucleotide sequence, SEQ ID NO:3, encoding Δ BCL-2. The injection is repeated weekly for 3 weeks while the level of circulating anti-insulin and anti-GAD autoantibodies is monitored. The treatment is ended when the level of circulating anti-insulin and anti-GAD autoantibodies has returned to normal.

Although the present invention has been discussed in considerable detail with reference to certain preferred embodiments, other embodiments are possible. Therefore, the scope of the appended claims should not be limited to the description of preferred embodiments contained in this disclosure.

Reference to any prior art in the specification is not, and should not be taken as, an acknowledgment or any form of suggestion that this prior art forms part of the common general knowledge in Australia or any other jurisdiction or that this prior art could reasonably be expected to be ascertained, understood and regarded as relevant by a person skilled in the art.

2003296892 01 Aug 2005

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A substance for preventing, delaying the onset of or treating one or more than one autoimmune disease, the substance comprising a polynucleotide construct comprising a polynucleotide sequence encoding the pro-apoptotic protein BAX and encoding one or more than one autoantigen for the autoimmune disease.
2. Use of a polynucleotide construct comprising a polynucleotide sequence encoding the pro-apoptotic protein BAX and encoding one or more than one autoantigen for an autoimmune disease for the manufacture of a medicament for preventing, delaying the onset of or treating the one or more than one autoimmune disease.
3. Use according to claim 2, where the medicament is manufactured in dosage units of between about 0.5 mg to about 5 mg.
4. Use according to claim 2, where the medicament is manufactured in dosage units of between about 1 mg to about 4 mg.
5. Use according to claim 2, where the medicament is manufactured in dosage units of between about 2.5 mg to about 3 mg.
6. Use according to any one of claims 2-5, where the medicament is manufactured in a form suitable for intramuscular administration.
7. Use according to any one of claims 2-5, where the medicament is manufactured in a form suitable for intravenous administration.
- 20 8. A method for preventing, delaying the onset of or treating an autoimmune disease in a patient comprising:
 - a) selecting a patient who is susceptible to developing the autoimmune disease, who is developing the autoimmune disease or who has the autoimmune disease; and
 - 25 b) administering to the patient one or more than one dose of a polynucleotide construct comprising a polynucleotide sequence encoding the pro-apoptotic protein BAX and encoding one or more than one autoantigen for the autoimmune disease.

2003296892 05 Dec 2005

9. The method of claim 8, where the autoimmune disease is type I diabetes.
10. The method of claim 9, where selecting the patient comprises identifying in the patient the presence of anti-insulin or anti-GAD autoantibodies or both anti-insulin and anti-GAD autoantibodies.
11. The method of claim 9, where selecting the patient comprises identifying in the patient the presence of increasing hyperglycemia.
12. The method of claim 9, where selecting the patient comprises identifying in the patient the presence of glycosuria.
13. The method of claim 8, where selecting the patient comprises identifying in the patient the presence of a genetic predisposition to the autoimmune disease.
14. The method according to any one of claims 8-13, where the one or more than one dose is a plurality of doses.
15. The method according to any one of claims 8-13, where administering to the patient one or more than one dose comprises injecting the patient intramuscularly with the one or more than one dose.
16. The method according to any one of claims 8-15, further comprising, after administering, monitoring the patient for the development of the autoimmune disease.
17. A method for preventing, delaying the onset of or treating an autoimmune disease in a patient comprising:

- 20 a) selecting a patient who is susceptible to developing the autoimmune disease, who is developing the autoimmune disease or who has the autoimmune disease; and
- b) administering to the patient one or more than one dose of a polynucleotide construct comprising a polynucleotide sequence encoding the pro-apoptotic protein BAX and one or more than one autoantigen for the autoimmune disease,
- 25 together with at least one protein selected from the adenoviral protein E3-GP19k, and the protein encoded by SEQ ID: 3.

2003296892 05 Dec 2005

20

25

18. The method of claim 17, where the autoimmune disease is type I diabetes.
19. The method of claim 18, where selecting the patient comprises identifying in the patient the presence of anti-insulin or anti-GAD autoantibodies or both anti-insulin and anti-GAD autoantibodies.
20. The method of claim 18, where selecting the patient comprises identifying in the patient the presence of increasing hyperglycemia.
21. The method of claim 18, where selecting the patient comprises identifying in the patient the presence of glycosuria.
22. The method of claim 17, where selecting the patient comprises identifying in the patient the presence of a genetic predisposition to the autoimmune disease.
23. The method according to any one of claims 17-22, where the one or more than one dose is a plurality of doses.
24. The method according to any one of claims 17-22, where administering to the patient one or more than one dose comprises injecting the patient intramuscularly with the one or more than one dose.
25. The method according to any one of claims 17-23, further comprising, after administering, monitoring the patient for the development of the autoimmune disease.
26. A substance for preventing, delaying the onset of or treating one or more than one autoimmune disease substantially as hereinbefore described with reference to the example.
27. Use of a polynucleotide construct comprising a polynucleotide sequence encoding the pro-apoptotic protein BAX substantially as hereinbefore described with reference to the example.
28. A method for preventing, delaying the onset of or treating an autoimmune disease in a patient substantially as hereinbefore described with reference to the example.

2003296892 05 Dec 2005

Dated : 5 December 2005

Freehills Patent & Trade Mark Attorneys

Patent & Trade Mark Attorneys for the Applicant:

Loma Linda University

FIG.1





FIG. 2



SEQUENCE LISTING

<110> Loma Linda University
 ESCHER, Alan P.
 LI, Fengchun

<120> Substances for Preventing and Treating Autoimmune Diseases

<130> 14102-1PCT

<140> to be assigned
 <141> 2003-08-06

<150> US 60/401,652
 <151> 2002-08-06

<160> 7

<170> PatentIn version 3.2

<210> 1
 <211> 579
 <212> DNA
 <213> Homo sapiens

<400> 1
 atggacgggt cgggggagca gccagaggc gggggggcca ccagctctga gcagatcatg 60
 aagacagggg cccttttget tcagggttcc atccaggatc gagcaggggc aatggggggg 120
 gaggcaccgc agctggccct ggaccgggtg cctcaggatg cgtccacca gaagctgagc 180
 gagtgtctca agcgcatcgg ggacgaactg gacagtaaca tggagctgca gaggatgatt 240
 gccgcggtgg acacagactc ccccgagag gtctttttcc gagtggcagc tgacatgttt 300
 tctgacggca acttcaactg gggcggggtt gtgcgccctt tctactttgc cagcaaaactg 360
 gtgtctcaagg cctgtgtcac caagggtccg gaactgatca gaaccatcat gggctggaca 420
 ttggacttcc tccgggagcg gctgttgggc tggatccaag accaggggtg ttgggacggc 480
 ctctctctct actttgggac gccacgtgg cagacogtga ccatctttgt ggcgggagtg 540
 ctcaccgct cgtcaccat ctggaagaag atgggctga 579

<210> 2
 <211> 492
 <212> DNA
 <213> Human adenovirus type 2

<400> 2
 atgagggtaca tgatttttagg cttgctcgcc cttgcggcag tctgcagcgc tgccaaaaag 60
 gttgagttaa aggaaccagc ttgcaatggt acattttaa atcagaagctaa tgaatgcact 120
 actcttataa aatgcaccac agaacatgaa aagcttatta ttgccacaa agacaaaatt 180
 ggcaagtatg ctgtatatgc tatttggcag ccaggtgaca ctaacgacta taatgtcaca 240
 gtotccaag gtgaaaatcg taaaactttt atgtataaat ttccatttta tgaatgtgc 300
 gatattacca tgtacatgag caaacagtac aagttgtggc cccacaaaa gtgttttagag 360
 aacactggca ccttttgttc caccgctctg cttattacag cgtttgcttt ggtatgtacc 420
 ttactttatc tcaaatataa aagcagacgc agttttattg atgaaaagaa aatgccttga 480
 ttttcgctt gc 492

<210> 3
 <211> 599
 <212> DNA
 <213> Homo sapiens

<400> 3
 atggcgacag ctgggagaag tggttacgat aaccgggaga tagtgatgaa gtacatccat 60
 tataagctgt cgcagagggg ctacgagtgg gatgctacg cggctgccgc ggggcctgcg 120
 ctacagcccg tgccacctgt ggtccacctg accctccgac aggcgcgcga cgaattctcc 180
 cgcgcctacc gcgcgactt cgcgcgagatg tccagccagc tgcacctgac gcccttcacc 240
 gcgcggggat gctttgccac ggtggtggag gagctcttca gggacggggg gaactggggg 300
 aggattgtgg ccttctttga gttcgggtgg gtcattgtgtg tggagagcgt caaccgggag 360
 atgtcgcgcc tgggtggaca catcgccctg tggatgactg agtaacctgaa ccggcacctg 420
 cacacctgga tccaggataa cggaggctgg gatgcctttg tggaaactgta cggccccagc 480

atgogggccto tgtttgattt ctcttggtg tctctgaaga ctctgctcag tttggccctg 540

gtgggagctt gcatcacctt ggggtgcctat ctggggcaca agtgaagtca acatgcctg 599

<210> 4

<211> 619

<212> DNA

<213> Encephalomyocarditis virus

<400> 4

tctagataat acgactcact atagggcgaa ttccccctct cctccccccc cccctaacgt 60

tactggccga agccgcttgg aataaggcgg gtgtggttt gtctatatgt tattttccac 120

catattgcgg tcttttgcca atgtgagggc cgggaaacct ggccctgtct tcttgacgag 180

cattcctagg ggtctttccc ctctcgccaa aggaatgcaa ggtctgttga atgtcgtgaa 240

ggaagcagtt cctctggaag cttcttgaag acaacaacg tctgtagcga ccttttgcag 300

gcagoggaa cccccactg ggcacagggt cctctggggc caaaagccag gtgtataaga 360

tacacctgca aaggcggcac aacccagtg ccacgttgtg agtggaata gttgtggaaa 420

gagtcaaatg gctctcctca agcgtattca acaaggggct gaaggatgcc cagaaggtag 480

cccatgtat gggatctgat ctggggcctc ggtgcacatg ctttacctgt gtttagtcga 540

ggttaaaaaa cgtctaggcc cccaaccac ggggacgtgg ttttcctttg aaaaacacga 600

ttattatatt gcctctaga 619

<210> 5

<211> 1868

<212> DNA

<213> Homo sapiens

<400> 5

gagctccacc ggggtggcgg ccgctctaga ccaccatggc atctccgggc tctggctttt 60

ggtcttttcgg gtgggaagat ggtcttgggg attccgagaa tcccggcaca ggcgcagcct 120

ggtgccaaat ggtctcagaag ttcacgggcg gcatcggaaa caaactgtgc gccctgctct 180

acggagacgc cgagaagccg gcggagagcg gggggagcca accccgcgg gccgcgcgcc 240

```

ggaaggccgc ctgcgcctgc gaccagaagc cctgcagctg ctccaaagt gatgtcaact 300
acgcgctttot ccatgcaaca gacctgctgc cggcgtgtga tggagaaagg cccactttgg 360
cgttttctgca agatgttatg aacattttac ttcagtatgt ggtgaaaagt ttcgatagat 420
caaccaaagt gattgatttc cattatccta atgagcttct ccaagaatat aattgggaat 480
tggcagacca accacaaaat ttggaggaaa ttttgatgca ttgccaaaca actctaaaat 540
atgcaattaa aacagggcat cctagatact tcaatcaact ttctactggt ttggatatgg 600
ttggattagc agcagactgg ctgacatcaa cagcaaatac taacatgttc acctatgaaa 660
ttgtccagt atttgtctt ttggaatag tccactaaa gaaatgaga gaaatcattg 720
gttgccagg gggctctggc gatgggatat tttctccgg tggcgccata tctaactgt 780
atgccatgat gatgcacgc tttaagatgt tcccagaagt caaggagaaa ggaatggctg 840
ctcttccag gctcattgcc ttcacgtctg aacatagtca tttttctctc aagaaggag 900
ctgcagcctt agggattgga agagacagcg tgattctgat taaatgtgat gagagaggga 960
aatgatttc atctgatctt gaaagaagga ttctgaagc caaacagaaa gggtttgttc 1020
ctttctctgt gagtgcaca gctggaacca cgtgtacgg agcatttgac cccctcttag 1080
ctgtcgtga catttgcaaa aagtataaga tctggatgca tgtggatgca gcttggggtg 1140
gggattact gatgtccga aaacacaagt ggaaactgag tggcgtggag agggccaact 1200
ctgtgacgtg gaatccacac aagatgatgg gagtcccttt gcagtgtct gctctcttg 1260
ttaagagaag gggattgatg cagaattgca accaaatgca tgctctctac ctctttcagc 1320
aagataaaca ttatgacctg tcctatgaca ctggagacaa ggccttacag tgcggacgcc 1380
acgttgatgt ttttaacta tggctgatgt ggagggcaa ggggactacc gsgtttgaag 1440
cgcatgttga taaatgtttg gagtggcag agtatattata caacatcata aaaaaccgag 1500
aaggatatga gatggtgttt gatgggaagc ctgaggacac aaatgtctgc ttctggtaca 1560
ttctccaag cttcgtact ctggaagaca atgaagagag aatgagtcgc ctctcgaagg 1620

```

tggctccagt gattaaagcc agaatgatgg agtatggaac cacaatggtc agctaccaac 1680
 ccttgggaga caaggtcaat ttcttcogca tggctcatctc aaacccagcg goaactcacc 1740
 aagacattga ctctctgatt gaagaaatag aacgccttgg acaagattta taataacctt 1800
 gctcaccaag ctgttccact tctctaggta gcgacctcga gcggccgctc gagggggggc 1860
 ccggtacc . 1868

<210> 6
 <211> 1638
 <212> DNA
 <213> Artificial

<220>
 <223> secreted form of human GAD

<400> 6
 atgtacagga tgcaactcct gtcttgcatc gcactaagtc ttgcacttgt cacaacacgt 60
 gcacctactt acgcgtttct ccatgcaaca gacctgctgc cggcgtgtga tggagaaagg 120
 cccacttttg cgtttctgca agatgttatg aacattttac ttcagtatgt ggtgaaaagt 180
 ttctagatag caaccaaagt gattgatttc cattatccta atgagcttct ccaagaatat 240
 aattgggaat tggcagacca accacaaaat ttggaggaaa ttttgatgca ttgocaaaca 300
 actctaaaat atgcaattaa aacagggcat cctagatact tcaatcaact ttctacttgt 360
 ttggatatgg ttggattagc agcagactgg ctgacatcaa cagcaaatat taacatgttc 420
 acctatgaaa ttgctccagt atttgtgctt ttggaatatg tcacactaaa gaaatgaga 480
 gaaatcattg gctggccagg gggctctggc gatgggatat tttctcccg tggcgccata 540
 tctaacatgt atgccatgat gatcgacgc tttaagatgt tcccagaagt caaggagaaa 600
 ggaatggctg ctcttccag gctcattgcc ttcacgtctg aacatagtca tttttctctc 660
 aagaagggag ctgcagcctt agggattgga agagacagcg tgattctgat taaatgtgat 720
 gagagagggg aatgattcc atctgatctt gaaagaagga ttcttgaagc caaacagaaa 780
 gggtttgttc ctctctcgt gagtgcaca gctggaacca ccgtgtacgg agcatttgac 840

```

ccccctcttag ctgtcgctga catttgcaaa aagtataaga tctggatgca tgtggatgca 900
gcttgggggtg ggggattact gatgtccga aaacacaagt ggaaactgag tggcgtggag 960
agggccaact ctgtgacgtg gaatccacac aagatgatgg gagtcccttt gcagtgggtct 1020
gctctcctgg ttagagaaga gggattgatg cagaattgca accaaatgca tgcctcctac 1080
ctctttcagc aagataaaca ttatgacctg tcctatgaca ctggagacaa ggccttacag 1140
tgcggacgcc acgttgatgt ttttaacta tggctgatgt ggagggcaaa ggggactacc 1200
gggtttgaag cgcattgtga taaatgtttg gagttggcag agtatttata caacatcata 1260
aaaaaccgag aaggatatga gatggtgttt gatgggaagc ctgaggacac aaatgtctgc 1320
ttctggtaca ttctccaag cttgcgtact ctggaagaca atgaagagag aatgagtcgc 1380
ctctogaagg tggctccagt gattaaagcc agaatgatgg agtatggaac cacaatggtc 1440
agctaccaac ccttgggaga caaggtcaat ttcttcgca tggctatctc aaaccagcg 1500
gcaactcacc aagacattga cttctgatt gaagaaatag aacgccttg acaagattta 1560
taataacctt gctcaccaag ctgttcact tctctaggta ggcacctga gggccgctc 1620
gagggggggc ccggtacc 1638

```

<210> 7
 <211> 1271
 <212> DNA
 <213> Artificial

<220>
 <223> secreted form of Renilla luciferase

```

<400> 7
atgtacagga tgcaactcct gtcttgcatg gcaactaagtc ttgcacttgt cacaacagtc 60
gcacctactg aattcagctt aaagatgact tcgaaagttt atgatccaga acaaaggaaa 120
cggatgataa ctggtccgca gtggtgggcc agatgtaaac aaatgaatgt tcttgattca 180
tttattaatt attatgattc agaaaaacat gcagaaaatg ctgttatattt ttacatggt 240

```

aacgcggcct cttcttattt atggcgacat gttgtgccac atattgagcc agtagcgcgg	300
tgtattatac cagatcttat tggatatggc aaatcaggca aatctggtaa tggttcttat	360
aggttacttg atcattacaa atatcttact gcatggtttg aacttcttaa tttaccaaag	420
aagatcattt ttgtcggcca tgattggggt gctgcttttg catttcatta tagctatgag	480
catcaagata agatcaaagc aatagttcac gctgaaagt tagtagatgt gattgaatca	540
tgggatgaat ggctgatat tgaagaagat attgcgttga tcaaatctga agaaggagaa	600
aaaatggttt tggagaataa cttcttctgt gaaaccatgt tgccatcaaa aatcatgaga	660
aagttagaac cagaagaatt tgcagcatat cttgaacat tcaaagagaa aggtgaagtt	720
cgctgtccaa cattatcatg gcctcgtgaa atcccgtag taaaaggtag taaacctgac	780
gttgtacaaa ttgttaggaa ttataatgct tatctacgtg caagtgatga tttacaaaaa	840
atgtttattg aatcggatcc aggattcttt tccaatgcta ttgttgaagg cgccaagaag	900
tttcttaata ctgaatttgt caaagtataa ggtcttcatt tttcgcaaga agatgcacct	960
gatgaaatgg gaaaatataa caaatcgttc gttgagcgag ttctcaaaaa tgaacaataa	1020
ttacttttgt tttttattta ctttttccc gggtttaata atataaatgt ctttttcaac	1080
aattttattt taactgaata tttcacagg aacattcata tatgttgatt aatttagctc	1140
gaactttact ctgtcatatc attttggaat attacctctt tcaatgaac tttataaaca	1200
gtggttcaat taattaatat atattataat tacatttggt atgtaataaa ctgggtttta	1260
ttataaaaaa a	1271