

(19) DANMARK

(10) DK/EP 2280021 T3



(12)

Oversættelse af
europæisk patent

Patent- og
Varemærkestyrelsen

(51)	Int.Cl.:	C 07 K 1/36 (2006.01) A 61 K 9/16 (2006.01) A 61 K 47/48 (2006.01) C 07 K 1/30 (2006.01) C 07 K 14/62 (2006.01)	A 61 K 9/00 (2006.01) A 61 K 38/28 (2006.01) B 82 Y 5/00 (2011.01) C 07 K 1/32 (2006.01)	A 61 K 9/14 (2006.01) A 61 K 47/22 (2006.01) C 07 D 241/08 (2006.01) C 07 K 14/60 (2006.01)
------	----------	--	---	--

(45) Oversættelsen bekendtgjort den: **2016-05-02**

(80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2016-02-17**

(86) Europæisk ansøgning nr.: **10075475.3**

(86) Europæisk indleveringsdag: **2000-06-29**

(87) Den europæiske ansøgnings publiceringsdag: **2011-02-02**

(30) Prioritet: **1999-06-29 US 141433 P**

(62) Stamansøgningsnr: **07075344.7**

(84) Designerede stater: **AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE**

(73) Patenthaver: **MannKind Corporation, 25134 Rye Canyon Loop, Suite 300, Valencia, CA 91355, USA**

(72) Opfinder: **Steiner, Solomon S., 24 Old Wagon Road, Mount Kisco, NY 10549, USA**
Sulner, Joseph W., 330 Bullard Avenue, Paramus, NJ 07652, USA
Woods, Rodney J., 395 Candlewood Lake Road North, New Milford, CT 06776, USA

(74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**

(54) Benævnelse: **Farmaceutiske formuleringer omfattende insulin kompleksbundet til et diketopiperazin**

(56) Fremdragne publikationer:
WO-A1-93/18754
WO-A1-96/09813
WO-A1-96/10396

DK/EP 2280021 T3

DESCRIPTION

Background Of The Invention

[0001] The present invention is generally in the field of pharmaceutical formulations, and more particularly related to methods and compositions for purifying and stabilizing insulin, for use in pharmaceutical applications.

[0002] In a normal person, the β -cells of the pancreatic islets of Langerhans produce insulin, required by the body for glucose metabolism, in response to an increase in blood glucose concentration. The insulin metabolizes incoming glucose and temporarily stops the liver's conversion of glycogen and lipids to glucose thereby allowing the body to support metabolic activity between meals. The Type I diabetic, however, has a reduced ability or absolute inability to produce insulin due to β -cell destruction and needs to replace the insulin via daily injections or an insulin pump. More common than Type I diabetes, though, is Type II diabetes, which is characterized by insulin resistance and increasingly impaired pancreatic β -cell function. Type II diabetics may still produce insulin, but they may also require insulin replacement therapy.

[0003] Type II diabetics typically exhibit a delayed response to increases in blood glucose levels. While normal persons usually release insulin within 2-3 minutes following the consumption of food, Type II diabetics may not secrete endogenous insulin for several hours after consumption. As a result, endogenous glucose production continues after consumption (Pfeiffer, Am. J. Med., 70:579-88 (1981)), and the patient experiences hyperglycemia due to elevated blood glucose levels.

[0004] Loss of glucose-induced insulin secretion is one of the earliest disturbances of β -cell function (Cerasi et al., Diabetes, 21:224-34 (1972); Polonsky et al., N. Engl. J Med., 318:1231-39 (1988)), but the causes and degree of β cell dysfunction are unknown in most cases. While genetic factors play an important role, (Leahy, Curr. Opin. Endocrinol. Diabetes, 2:300-06 (1995)), some insulin secretory disturbances seem to be acquired and may be at least partially reversible through optimal glucose control. Optimal glucose control via insulin therapy after a meal can lead to a significant improvement in natural glucose-induced insulin release by requiring both normal tissue responsiveness to administered insulin and an abrupt increase in serum insulin concentrations. Therefore, the challenge presented in the treatment of early stage Type II diabetics, those who do not have excessive loss of β -cell function, is to restore the release of insulin following meals.

[0005] Most early stage Type II diabetics currently are treated with oral agents, but with little success. Subcutaneous injections of insulin are also rarely effective in providing insulin to Type II diabetics and may actually worsen insulin action because of delayed, variable, and shallow onset of action. It has been shown, however, that if insulin is administered intravenously with a meal, early stage Type II diabetics experience the shutdown of hepatic gluconeogenesis and exhibit increased physiological glucose control. In addition, their free fatty acids levels fall at a faster rate than without insulin therapy. While possibly effective in treating Type II diabetes, intravenous administration of insulin, is not a reasonable solution, as it is not safe or feasible for patients to intravenously administer insulin at every meal.

[0006] Insulin, a polypeptide with a nominal molecular weight of 6,000 Daltons, traditionally has been produced by processing pig and cow pancreas to isolate the natural product. More recently, however, recombinant technology has been used to produce human insulin *in vitro*. Natural and recombinant human insulin in aqueous solution is in a hexameric configuration, that is, six molecules of recombinant insulin are noncovalently associated in a hexameric complex when dissolved in water in the presence of zinc ions. Hexameric insulin is not rapidly absorbed. In order for recombinant human insulin to be absorbed into a patient's circulation, the hexameric form must first associate into dimeric and/or monomeric forms before the material can move into the blood stream. The delay in absorption requires that the recombinant human insulin be administered approximately one half hour prior to meal time in order to produce therapeutic insulin blood level, which can be burdensome to patients who are required to accurately anticipate the times they will be eating. To overcome this delay, analogs of recombinant human insulin, such as HUMALOG™, have been developed, which rapidly disassociate into a virtually entirely monomeric form following subcutaneous administration. Clinical studies have demonstrated that HUMALOG™ is absorbed quantitatively faster than recombinant human insulin after subcutaneous administration. See, for example, U.S. Patent No. 5,547,929 to Anderson Jr., et al.

[0007] In an effort to avoid the disadvantages associated with delivery by injection and to speed absorption, administration of monomeric analogs of insulin via the pulmonary route has been developed. For example, U.S. Patent No. 5,888,477 to Gonda, et al. discloses having a patient inhale an aerosolized formulation of monomeric insulin to deposit particles of insulin on the patient's lung tissue. However, the monomeric formulation is unstable and rapidly loses activity, while the rate of uptake remains unaltered.

[0008] While it would be desirable to produce rapidly absorbable insulin derived from natural sources, transformation of the

hexameric form into the monomeric form, such as by removing the zinc from the complex, yields an insulin that is unstable and has an undesirably short shelf life. It therefore would be desirable to provide monomeric forms of insulin, while maintaining its stability in the absence of zinc. It also would be advantageous to provide diabetic patients with monomeric insulin compositions that are suitable for pulmonary administration, provide rapid absorption, and which can be produced in ready-to-use formulations that have a commercially useful shelf-life.

[0009] These problems with impurities, metal ions that affect stability or bioavailability, occur with many other proteins and peptides.

[0010] U.S. Patent No. 6,071,497 to Steiner, et al. discloses microparticle drug delivery systems in which the drug is encapsulated in diketopiperazine microparticles which are stable at a pH of 6.4 or less and unstable at pH of greater than 6.4, or which are stable at both acidic and basic pH, but which are unstable at pH between about 6.4 and 8. The patent does not describe monomeric insulin compositions that are suitable for pulmonary administration, provide rapid absorption, and which can be produced in ready-to-use formulations that have a commercially useful shelf-life.

[0011] It would therefore be advantageous to develop alternative insulin delivery compositions for Type II diabetics that provide more rapid elevation of insulin blood levels and are easily administered to ensure patient compliance. It also would be desirable to apply the delivery compositions and methods to other biologically active agents.

[0012] The present application describes improved methods for purifying peptides and proteins, especially in the preparation of compositions suitable for pulmonary administration.

[0013] The present application describes stable monomeric peptide compositions suitable for pulmonary delivery.

[0014] The present application describes methods and compositions for the facilitated transport of insulin and other biologically active agents across biological membranes.

[0015] The present application describes methods and compositions for the improved absorption of insulin or other biologically active agents in the bloodstream.

[0016] The present application describes methods and compositions for the improved absorption of insulin or other biologically active agents in the bloodstream characterized by ease of administration.

Summary Of The Invention

[0017] The present invention provides a composition comprising microparticles, which comprise insulin complexed to a diketopiperazine, for use in medicine; wherein the composition releases monomeric or dimeric insulin upon administration; and wherein the microparticles are obtainable through a process comprising -

1. (i) providing the diketopiperazine as pre-formed microparticles in a suspension;
2. (ii) combining the microparticles in suspension with a solution of insulin; and
3. (iii) lyophilizing or freeze drying said suspension;

thereby to yield diketopiperazine microparticles having an insulin coating.

[0018] The diketopiperazine may, for example, be 2,5-diketo-3,6-di(4-X-aminobutyl)piperazine, wherein X represents a succinyl, glutaryl, maleyl or fumaryl group; and may preferably be 2,5-diketo-3,6-di(4-fumaryl-aminobutyl)piperazine.

[0019] The insulin complexed to a diketopiperazine may be monomeric insulin. The insulin complexed to a diketopiperazine may be predominately in the dimeric and monomeric state, as opposed to the hexameric state. The insulin may, for example, be human insulin.

[0020] In one option, the microparticles are obtainable by the process as defined above, and wherein that process further comprises the step of removing zinc ions by washing the microparticles with an appropriate solvent.

[0021] The composition of the present invention may comprise a stabilized form of insulin which has been purified to remove zinc.

[0022] The present invention also provides for the use of a composition comprising microparticles, which comprise insulin complexed to a diketopiperazine in the manufacture of a medicament for treating diabetes, such as type II diabetes; wherein the composition releases monomeric or dimeric insulin upon administration; and wherein the microparticles are obtainable through a process comprising providing the diketopiperazine as pre-formed microparticles in a suspension; adding a solution of insulin to said suspension; and then lyophilizing or freeze drying said suspension to form diketopiperazine microparticles having an insulin coating.

[0023] Methods are also described herein for purifying peptides and proteins by incorporating the peptide or protein into a diketopiperazine or competitive complexing agent to facilitate removal one or more impurities, i.e. undesirable components, from the peptide or protein. In a preferred embodiment, a peptide, such as insulin, containing one or more impurities, e.g., zinc ions, is entrapped in diketopiperazine to form a precipitate of peptide/diketopiperazine/impurity, which is then washed with a solvent for the impurity to be removed, which is a nonsolvent for the diketopiperazine and a nonsolvent for the peptide. Alternatively, the impurity can be removed by using complexing agents to selectively complex with and displace the impurities, for example, such as by dialysis.

[0024] Formulations and methods also are described for the improved transport of active agents across biological membranes, resulting, for example, in a rapid increase in blood agent concentration. The formulations include microparticles formed of (i) an active agent, which may be charged or neutral, and (ii) a transport enhancer that masks the charge of the agent and/or that forms hydrogen bonds with the target biological membrane in order to facilitate transport.

[0025] The present invention provides a composition comprising microparticles, which comprise insulin complexed to a diketopiperazine, for use in medicine; wherein the composition releases monomeric or dimeric insulin upon administration; and wherein the microparticles are obtainable through a process comprising -

1. (i) providing the diketopiperazine as pre-formed microparticles in a suspension;
2. (ii) combining the microparticles in suspension with a solution of insulin; and
3. (iii) lyophilizing or freeze drying said suspension;

thereby to yield diketopiperazine microparticles having an insulin coating.

[0026] In a preferred embodiment, the insulin is administered via pulmonary delivery of microparticles comprising fumaryl diketopiperazine and insulin in its biologically active form. The charge on the insulin molecule is masked by hydrogen bonding it to the diketopiperazine, thereby enabling the insulin to pass through the target membrane. This method of delivering insulin results in a rapid increase in blood insulin concentration that is comparable to the increase resulting from intravenous delivery.

Brief Description of the Drawings

[0027]

Figure 1a is a graph of mean blood glucose values over time (minutes). Figure 1b is a graph of mean C-peptide concentrations during experiments comparing levels of C-peptide (ng/ml) over time (minutes) when insulin was administered intravenously, subcutaneously, and by inhalation.

Figure 2a is a graph of glucose infusion rate (mg/kg/min) over time (minutes) comparing insulin administered intravenously, subcutaneously, and by inhalation. Figure 2b is a graph of mean insulin concentrations (μ U/ml) over time (minutes) comparing insulin administered intravenously, subcutaneously, and by inhalation.

Detailed Description Of The Invention

[0028] As described herein, encapsulation or entrapment of large polymers, such as proteins and peptides, in diketopiperazines can be used to remove impurities or contaminants such as metal ions or other small molecules. The present invention relies on the formation of a complex between insulin and diketopiperazine, wherein the composition releases monomeric or dimeric insulin upon administration, and wherein the microparticles are obtainable through a process comprising -

1. (i) providing the diketopiperazine as pre-formed microparticles in a suspension;

2. (ii) combining the microparticles in suspension with a solution of insulin; and
3. (iii) lyophilizing or freeze drying said suspension;

thereby to yield diketopiperazine microparticles having an insulin coating. The diketopiperazines also serve both to stabilize and enhance delivery of the complexed insulin. Formulations also have been developed for the enhanced transport of insulin across biological membranes. These formulations include microparticles formed of a diketopiperazine and insulin coated thereon. The diketopiperazine can mask the charge of the insulin and/or can form hydrogen bonds with the membrane. The formulations can provide rapid increases in the concentration of the insulin in the blood following administration of the formulations.

[0029] For example, it was discovered that hexameric insulin can be delivered to the lung in fumaryl diketopiperazine formulation, reaching peak blood concentrations within 3-10 minutes. In contrast, insulin administered by the pulmonary route without fumaryl diketopiperazine typically takes between 25-60 minutes to reach peak blood concentrations, while hexameric insulin takes 30-90 minutes to reach peak blood level when administered by subcutaneous injection. This feat has been successfully replicated several times and in several species, including humans.

[0030] Removing zinc from insulin typically produces unstable insulin with an undesirably short shelf life. Purification to remove zinc, stabilization and enhanced delivery of insulin is demonstrated by the examples. Formulations of insulin trapped in fumaryl diketopiperazine were found to be stable and have an acceptable shelf life. Measurement of the zinc levels demonstrated that the zinc had been largely removed during the entrapment process, yielding monomeric insulin in a stable delivery formulation.

[0031] Rapid absorption of a number of other peptides, including salmon calcitonin, parathyroid hormone 1-34, octreotide, leuprolide and RSV peptide, has been observed when the peptide is pulmonarily delivered in fumaryl diketopiperazineproviding peak blood concentrations within 3-10 minutes after pulmonary delivery.

I. Materials

A. Agent to be Delivered

[0032] In accordance with the disclosure of the present application, the agent to be delivered is referred to herein as the active agent, or molecule to be encapsulated or entrapped. For the purposes of the presently-claimed invention, the active agent is insulin complexed to a diketopiperazine, in the form of microparticles obtainable through a process comprising -

1. (i) providing the diketopiperazine as pre-formed microparticles in a suspension;
2. (ii) combining the microparticles in suspension with a solution of insulin; and
3. (iii) lyophilizing or freeze drying said suspension;

thereby to yield diketopiperazine microparticles having an insulin entrapped thereon as a coating. The insulin may or may not be a charged species.

[0033] The exact mechanism by which the diketopiperazines form a complex with the materials to be delivered, such as insulin, is not known, but it is believed that the diketopiperazines form a complex with the material to be purified. This process is referred to herein interchangeably as entrapment or complexation.

[0034] In a preferred embodiment, the active agent is monomeric insulin or a stabilized form of insulin which has been purified to remove zinc

[0035] Impurities which can be removed from the insulin include metal ions such as zinc, and other di- or multi-valent ions, and small inorganic molecules and solvent residuals.


B. Diketopiperazines

[0036] Diketopiperazines useful in the present compositions and methods are described, for example, in U.S. Patent No. 6,071,497.

(i). General Formula

[0037] The diketopiperazines are rigid planar rings with at least six ring atoms containing heteroatoms and unbonded electron pairs. Also disclosed are diketomorpholine and diketodioxane, respectively. Although it is possible to replace a nitrogen with a sulfur atom, this does not yield a stable structure.

[0038] The general formula for diketopiperazine and its analogs is shown below.



[0039] Wherein R3 and R6 are side chains defined as Q-T-Q-U or Q-U, Q is, independently, a C₁₋₂₀ straight, branched or cyclic alkyl, aralkyl, alkaryl, alkenyl, alkynyl, heteroalkyl, heterocyclic, alkyl-heterocyclic, or heterocyclic-alkyl; T is -C(O)O, -OC(O), -C(O)NH, -NH,-NQ, -OQO, -O, -NHC(O), -OP(O), -P(O)O, -OP(O)₂, -P(O)₂O, -OS(O)₂, or -S(O)₃; U is an acid group, such as a carboxylic acid, phosphoric acid, phosphonic acid and sulfonic acid, or a basic group, such as primary, secondary and tertiary amines, quaternary ammonium salts, guanidine, aniline, heterocyclic derivatives, such as pyridine and morpholine, or a zwitterionic C₁₋₂₀ chain containing at least one acidic group and at least one basic group, for example, those described above, wherein the side chains can be further functionalized with an alkene or alkyne group at any position, one or more of the carbons on the side chain can be replaced with an oxygen, for example, to provide short polyethylene glycol chains, one or more of the carbons can be functionalized with an acidic or basic group, as described above, and wherein the ring atoms X at positions 1 and 4 are N.

[0040] Examples of acidic side chains include, but are not limited to, cis and trans -CH=CH-CO₂H, -CH(CH₃)=CH(CH₃)-CO₂H, -(CH₂)₃-CO₂H, -CH₂CH(CH₃)-CO₂H, -CH(CH₂CO₂H)=CH₂, -(tetrafluoro)benzoic acid, -benzoic acid and -CH(NHC)(O)CF₃)-CH₂-CO₂H.

[0041] Examples of basic side chains include, but are not limited to, -aniline, -phenyl-C(NH)NH₂, -phenyl-C(NH)NH(alkyl), -phenyl-C(NH)N(alkyl)₂ and -(CH₂)₄NHC(O)CH(NH₂)CH(NH₂)CO₂H.

[0042] Examples of zwitterionic side chains include, but are not limited to, -CH(NH₂)-CH₂-CO₂H and -NH(CH₂)₁₋₂₀CO₂H.

[0043] The term aralkyl refers to an aryl group with an alkyl substituent.

[0044] The term heterocyclic-alkyl refers to a heterocyclic group with an alkyl substituent.

[0045] The term alkaryl refers to an alkyl group that has an aryl substituent.

[0046] The term alkyl-heterocyclic refers to an alkyl group that has a heterocyclic substituent.

[0047] The term alkene, as referred to herein, and unless otherwise specified, refers to an alkene group of C₂ to C₁₀, and specifically includes vinyl and allyl.

[0048] The term alkyne, as referred to herein, and unless otherwise specified, refers to an alkyne group of C₂ to C₁₀.

[0049] As used herein, "diketopiperazines" includes diketopiperazines and derivatives and modifications thereof falling within the scope of the above-general formula.

[0050] Fumaryl diketopiperazine is most preferred for pulmonary applications.

(ii). Synthesis

[0051] Diketopiperazines can be formed by cyclodimerization of amino acid ester derivatives, as described by Katchalski, et al., J. Amer. Chem. Soc. 68:879-80 (1946), by cyclization of dipeptide ester derivatives, or by thermal dehydration of amino acid derivatives in high-boiling solvents, as described by Kopple, et al., J. Org. Chem. 33(2):862-64 (1968). 2,5-diketo-3,6-di(aminobutyl)piperazine (Katchalski *et al.* refer to this as lysine anhydride) was prepared via cyclodimerization of N-epsilon-P-L-lysine in molten phenol, similar to the Kopple method in *J. Org. Chem.*, followed by removal of the blocking (P)-groups with 4.3 M HBr in acetic acid. This route is preferred because it uses a commercially available starting material, it involves reaction conditions that are reported to preserve stereochemistry of the starting materials in the product and all steps can be easily scaled up for manufacture.

[0052] Diketomorpholine and diketooxetane derivatives can be prepared by stepwise cyclization in a manner similar to that disclosed in Katchalski, et al., J. Amer. Chem. Soc. 68:879-80 (1946).

[0053] Diketopiperazines can be radiolabelled. Means for attaching radiolabels are known to those skilled in the art. Radiolabelled diketopiperazines can be prepared, for example, by reacting tritium gas with those compounds listed above that contain a double or triple bond. A carbon-14 radiolabelled carbon can be incorporated into the side chain by using ¹⁴C labeled precursors which are readily available. These radiolabelled diketopiperazines can be detected *in vivo* after the resulting microparticles are administered to a subject.

(a) Synthesis of Symmetrical

Diketopiperazine Derivatives

[0054] The diketopiperazine derivatives are symmetrical when both side chains are identical. The side chains can contain acidic groups, basic groups, or combinations thereof.

[0055] One example of a symmetrical diketopiperazine derivative is 2,5-diketo-3,6-di(4-succinylaminobutyl)piperazine. 2,5-diketo-3,6-di(aminobutyl) piperazine is exhaustively succinylated with succinic anhydride in mildly alkaline aqueous solution to yield a product which is readily soluble in weakly alkaline aqueous solution, but which is quite insoluble in acidic aqueous solutions. When concentrated solutions of the compound in weakly alkaline media are rapidly acidified under appropriate conditions, the material separates from the solution as microparticles.

[0056] Other preferred compounds can be obtained by replacing the succinyl group(s) in the above compound with glutaryl, maleyl or fumaryl groups.

(b) Synthesis of Asymmetrical

Diketopiperazine Derivatives

[0057] One method for preparing unsymmetrical diketopiperazine derivatives is to protect functional groups on the side chain, selectively deprotect one of the side chains, react the deprotected functional group to form a first side chain, deprotect the second functional group, and react the deprotected functional group to form a second side chain.

[0058] Diketopiperazine derivatives with protected acidic side chains, such as cyclo-Lys(P)Lys(P), wherein P is a benzyloxycarbonyl group, or other protecting group known to those skilled in the art, can be selectively deprotected. The protecting groups can be selectively cleaved by using limiting reagents, such as HBr in the case of the benzyloxycarbonyl group, or fluoride ion in the case of silicon protecting groups, and by using controlled time intervals. In this manner, reaction mixtures which contain unprotected, monoprotected and di-protected diketopiperazine derivatives can be obtained. These compounds have different solubilities in various solvents and pH ranges, and can be separated by selective precipitation and removal. An appropriate solvent, for example, ether, can then be added to such reaction mixtures to precipitate all of these materials together. This can stop the deprotection reaction before completion by removing the diketopiperazines from the reactants used to deprotect the protecting groups. By stirring the mixed precipitate with water, both the partially and completely reacted species can

be dissolved as salts in the aqueous medium. The unreacted starting material can be removed by centrifugation or filtration. By adjusting the pH of the aqueous solution to a weakly alkaline condition, the asymmetric monoprotected product containing a single protecting group precipitates from the solution, leaving the completely deprotected material in solution.

[0059] In the case of diketopiperazine derivatives with basic side chains, the basic groups can also be selectively deprotected. As described above, the deprotection step can be stopped before completion, for example, by adding a suitable solvent to the reaction. By carefully adjusting the solution pH, the deprotected derivative can be removed by filtration, leaving the partially and totally deprotected derivatives in solution. By adjusting the pH of the solution to a slightly acidic condition, the monoprotected derivative precipitates out of solution and can be isolated.

[0060] Zwitterionic diketopiperazine derivatives can also be selectively deprotected, as described above. In the last step, adjusting the pH to a slightly acidic condition precipitates the monoprotected compound with a free acidic group. Adjusting the pH to a slightly basic condition precipitates the monoprotected compound with a free basic group.

[0061] Limited removal of protecting groups by other mechanisms, including but not limited to cleaving protecting groups that are cleaved by hydrogenation by using a limited amount of hydrogen gas in the presence of palladium catalysts. The resulting product is also an asymmetric partially deprotected diketopiperazine derivative. These derivatives can be isolated essentially as described above.

[0062] The monoprotected diketopiperazine is reacted to produce a diketopiperazine with one sidechain and protecting group. Removal of protecting groups and coupling with other side chains yields unsymmetrically substituted diketopiperazines with a mix of acidic, basic, and zwitterionic sidechains.

[0063] Other materials that exhibit this response to pH can be obtained by functionalizing the amide ring nitrogens of the diketopiperazine ring.

C. Transport Enhancers

[0064] The insulin is complexed with a diketopiperazine, which preferably acts as a transport enhancer which is degradable and capable of forming hydrogen bonds with the target biological membrane in order to facilitate transport of the agent across the membrane. The transport enhancer also is capable of forming hydrogen bonds with the insulin, if charged, in order to mask the charge and facilitate transport of the insulin across the membrane.

[0065] The diketopiperazine transport enhancer preferably is biodegradable and may provide linear, pulsed or bulk release of the insulin.

[0066] A preferred transport enhancer is fumaryl diketopiperazine. Other diketopiperazines which may be useful as a transport enhancer are described above.

[0067] Like most proteins and peptides, insulin is a charged molecule, which impedes its ability to cross charged biological membranes. It has been found that when insulin hydrogen bonds to fumaryl diketopiperazine, the charge of the peptide is masked, thereby facilitating or enhancing the passage of insulin across the membranes, such as mucosal membranes, and into the blood.

II. Methods

A. Encapsulation

[0068] It is disclosed herein that an active agent may (a) be encapsulated within microparticles by dissolving a diketopiperazine with acidic side chains in bicarbonate or other basic solution, adding the active agent in solution or suspension, and then precipitating the microparticle by adding acid, such as 1 M citric acid.

[0069] Alternatively, as disclosed herein, an active agent may (b) be encapsulated within microparticles by dissolving a diketopiperazine with basic side chains in an acidic solution, such as 1 M citric acid, adding the active agent in solution or

suspension, and then precipitating the microparticle by adding bicarbonate or another basic solution.

[0070] Alternatively, as also disclosed herein, an active agent may (c) be encapsulated within microparticles by dissolving a diketopiperazine with both acidic and basic side chains in an acidic or basic solution, adding the active agent in solution or suspension to be encapsulated, then precipitating the microparticle by neutralizing the solution.

[0071] The microparticles can be stored in the dried state and suspended for administration to a patient. In option (a), the reconstituted microparticles maintain their stability in an acidic medium and dissociate as the medium approaches physiological pH in the range of between 6 and 14. In option (b) embodiment, suspended microparticles maintain their stability in a basic medium and dissociate at a pH of between 0 and 6. In option (c), the reconstituted microparticles maintain their stability in an acidic or basic medium and dissociate as the medium approaches physiological pH in the range of pH between 6 and 8.

[0072] The impurities typically are removed when the microparticles are precipitated. However, impurities also can be removed by washing the particles to dissolve the impurities. A preferred wash solution is water or an aqueous buffer. Solvents other than water also can be used to wash the microspheres or precipitate the diketopiperazines, in order to remove impurities that are not water soluble. Any solvent in which neither the cargo nor the fumaryl diketopiperazine is soluble are suitable. Examples include acetic acid, ethanol, and toluene.

B. Entrapment/Complexation

[0073] The presently-claimed invention provides an alternative to the encapsulation process discussed in section A. More specifically, the presently-claimed invention provides a composition comprising microparticles of diketopiperazine complexed to insulin, wherein the composition releases monomeric or dimeric insulin upon administration; and wherein the microparticles are obtainable through a process comprising (i) providing the diketopiperazine as pre-formed microparticles in a suspension; (ii) combining the microparticles in suspension with a solution of insulin; and (iii) lyophilizing or freeze drying said suspension; thereby to yield diketopiperazine microparticles having an insulin coating.

[0074] Accordingly, microparticles of diketopiperazine are prepared and provided in a suspension, typically an aqueous suspension, to which a solution of the insulin then is added. The suspension is then lyophilized or freeze dried to yield diketopiperazine microparticles having a coating of insulin. In one embodiment, the insulin is in a hexameric form. Zinc ions can then be removed by washing the microparticles with an appropriate solvent.

[0075] As used herein, the term "entrapped" with reference to insulin in/with a diketopiperazine refers to coating of the insulin onto microparticles of the diketopiperazine.

[0076] The diketopiperazine microparticles have been found to have a higher affinity for insulin than does zinc. Insulin has been found to be stabilized within an ordered lattice array of fumaryl diketopiperazine. In this state, in the sufficient absence of zinc ions, the insulin is predominately dimeric and monomeric, as opposed to its hexameric state. The insulin therefore more readily dissociates to its monomeric state, which is the state in which insulin exerts its biological activity.

C. Administration

[0077] The compositions of the present invention, as described herein, can be administered to patients in need of insulin. The compositions preferably are administered in the form of microparticles, which can be in a dry powder form for pulmonary administration or suspended in an appropriate pharmaceutical carrier, such as saline.

[0078] The microparticles preferably are stored in dry or lyophilized form until immediately before administration. The microparticles then can be administered directly as a dry powder, such as by inhalation using, for example, dry powder inhalers known in the art. Alternatively, the microparticles can be suspended in a sufficient volume of pharmaceutical carrier, for example, as an aqueous solution for administration as an aerosol.

[0079] The microparticles also can be administered via oral, subcutaneous, and intravenous routes.

[0080] The compositions can be administered to any targeted biological membrane, preferably a mucosal membrane of a patient. In a preferred embodiment, the patient is a human suffering from Type II diabetes. In a preferred embodiment, the

composition delivers insulin in biologically active form to the patient, which provides a spike of serum insulin concentration which simulates the normal response to eating.

[0081] In a preferred embodiment, hexameric insulin is entrapped in fumaryl diketopiperazine to form a solid precipitate of monomeric insulin in the fumaryl diketopiperazine, which then is washed with aqueous solution to remove the free zinc. This formulation demonstrates blood uptake following pulmonary administration at a rate 2.5 times the rate of insulin uptake following subcutaneous injection, with peak blood levels occurring at between 7.5 and 10 minutes after administration.

[0082] The range of loading of the drug to be delivered is typically between about 0.01% and 90%, depending on the form and size of the drug to be delivered and the target tissue.

[0083] The preferred range is from 0.1% to 50% loading by weight of drug. The appropriate dosage can be determined, for example, by the amount of complexed/entrapped insulin, the rate of its release from the microparticles, and, in a preferred embodiment, the patient's blood glucose level.

[0084] The compositions and methods described herein are further described by the following non-limiting examples.

Example 1: Removal of Zinc From U.S.P. Injectable Insulin

[0085] Insulin trapped in fumaryl diketopiperazine was analyzed to assess whether zinc was removed during the entrapment process. The insulin used as the starting material met U.S.P. standards for injectable insulin, and according to the certificate of analysis, the insulin contained a considerable quantity of zinc: 0.41 %. This insulin was then entrapped in fumaryl diketopiperazine to form a solid fumaryl diketopiperazine/insulin mixture, as described above.

[0086] Following entrapment of the insulin in fumaryl diketopiperazine, the amount of zinc theoretically should be present in the same proportion as it existed in the neat insulin. Using the certificate of analysis value, it was calculated that one should expect to find 697 parts per million (ppm) of zinc per gram in the solid yield of fumaryl diketopiperazine/insulin. Surprisingly, the quantity of zinc present the solid fumaryl diketopiperazine/insulin was measured to be only 6 ppm. The "missing" zinc was presumably eliminated with the water used to wash the insulin/fumaryl diketopiperazine precipitate.

Example 2: Bioavailability of Insulin in Diketopiperazine Pulmonary Formulation

Subjects and Methods

[0087] The study was reviewed and approved by the ethical review committee of the Heinrich-Heine-University, Düsseldorf, and conducted according to local regulations, the Declaration of Helsinki and the rules of Good Clinical Practice.

[0088] The study was conducted with 5 healthy male volunteers. Inclusion criteria were good health, as judged by physical examination, age: 18 to 40 years, body mass index: 18 to 26 kg/m², capability to reach peak inspiratory flow of 4 l/sec measured by a computer assisted spirometry and a FEV₁ equal to or greater than 80% of predicted normal (FEV₁ = forced expiratory volume in one second). Exclusion criteria were Diabetes mellitus type 1 or 2, prevalence of human insulin antibodies, history of hypersensitivity to the study medication or to drugs with similar chemical structures, history or severe or multiple allergies, treatment with any other investigational drug in the last 3 months before study entry, progressive fatal disease, history of drug or alcohol abuse, current drug therapy with other drugs, history significant cardiovascular, respiratory, gastrointestinal, hepatic, renal, neurological, psychiatric and/or hematological disease, ongoing respiratory tract infection or subjects defined as being smokers with evidence or history of tobacco or nicotine use.

Conduct of the Study

[0089] On the morning of the study days, the subjects came to the hospital (fasting, except for water, from midnight onward) at 7:30 a.m. The subjects were restricted from excessive physical activities and an intake of alcohol for 24 hours before each treatment day. They were randomly assigned to one of the three treatment arms. The subjects received a constant intravenous

regular human insulin infusion, which was kept at $0.15 \text{ mU min}^{-1} \text{ kg}^{-1}$ so that serum insulin concentrations were established at 10-15 U/ml during a period of 2 hours before time point 0. This low-dose infusion was continued throughout the test to suppress endogenous insulin secretion. Blood glucose was kept constant at a level of 90 mg/dl throughout the glucose clamp by a glucose controlled infusion system (BIOSTATOR™). The glucose clamp algorithm was based on the actual measured blood glucose concentration and the grade of variability in the minutes before to calculate the glucose infusion rates for keeping the blood glucose concentration constant. The insulin application (5 U i.v. or 10 U s.c. injection or three deep breaths inhalation per capsule (2 capsules with 50 U each) applied with a commercial inhalation device (Boehringer Ingelheim)) had to be finished immediately before time point 0. The duration of the clamp experiment was 6 hours from time point 0. Glucose infusion rates, blood glucose, serum-insulin and C-peptide were measured.

Bioefficacy and Bioavailability

[0090] To determine bioefficacy, the areas under the curve of the glucose infusion rates were calculated for the first 3 hours (AUC_{0-180}) after the administration and for the overall observation period of six hours after the administration (AUC_{0-360}) and were correlated to the amount of insulin applied. To determine bioavailability, the areas under the curve of the insulin concentrations were calculated for the first 3 hours (AUC_{0-180}) after the administration and for the overall observation period of six hours after the administration (AUC_{0-360}) and correlated to the amount of insulin applied.

[0091] In this clamp study, inhalation of 100 U of TECHNOSPHERE™/Insulin was well tolerated and was demonstrated to have a substantial blood glucose lowering effect with a relative bioavailability of 25.8% for the first three hours as calculated from the achieved serum insulin concentrations. TECHNOSPHERES™ are microparticles (also referred to herein as microspheres) formed of diketopiperazine that self-assembles into an ordered lattice array at particular pHs, typically a low pH. They typically are produced to have a mean diameter between about 1 and about 5 μm .

Results

[0092] The pharmacokinetic results are illustrated in Figures 1 and 2 and in Table 1.

Efficacy Results

[0093] Inhalation of 100 U of TECHNOSPHERE™/Insulin (inhalation of 100 U) revealed a peak of insulin concentration after 13 min (intravenous (i.v.) (5U): 5 min, subcutaneous (s.c.) (10 U): 121 min) and a return of the insulin levels to baseline after 180 min (i.v.: 60 min, s.c. 360 min). Biological action as measured by glucose infusion rate peaked after 39 min (i.v. 14 min, s.c.: 163 min) and lasted for more than 360 min (i.v.: 240 min, s.c.: > 360 min). Absolute bioavailability (comparison to i.v. application) was $14.6 \pm 5.1\%$ for the first 3 hours and $15.5 \pm 5.6\%$ for the first 6 hours. Relative bioavailability (comparison to s.c. application) was $25.8 \pm 11.7\%$ for the first 3 hours and $16.4 \pm 7.9\%$ for the first 6 hours.

Table 1: Pharmacokinetic Parameters

<u>Parameter Calculated on Glucose Infusion Rate</u>	<u>Intravenous Administration</u>	<u>Inhaled</u>	<u>Subcutaneous Administration</u>
T50%*	9 min	13 min	60 min
Tmax	14 min	39 min	163 min
T-50%**	82 min	240 min	240 min
T to baseline	240 min	>360 min	>360 min
<u>Parameter Calculated on Insulin Levels</u>	<u>Intravenous Administration</u>	<u>Inhaled</u>	<u>Subcutaneous Administration</u>
T50%*	2 min	2.5 min	27 min
Tmax	5 min	13 min	121 min
T-50% **	6 min	35 min	250 min
T to baseline	60 min	180 min	360 min

* time from baseline to half-maximal values
 ** time from baseline to half-maximal after passing Tmax

Safety Results

[0094] TECHNOSPHERE™/Insulin was shown to be safe in all patients. One patient was coughing during the inhalation without any further symptoms or signs of deterioration of the breathing system.

Conclusions

[0095] Inhalation of 100 U of TECHNOSPHERE™/Insulin was well tolerated and was demonstrated to have a substantial blood glucose lowering effect with a relative bioavailability of 25.8% for the first 3 hours as calculated from the achieved serum insulin concentrations.

Summary

[0096] In this study, the inhalation of TECHNOSPHERE™/Insulin (the formulation of example 1) was demonstrated in healthy human subjects to have a time-action profile with a rapid peak of insulin concentration (Tmax: 13 min) and rapid onset of action (Tmax: 39 min) and a sustained action over more than 6 hours. The total metabolic effect measured after inhalation of 100 U of TECHNOSPHERE™/Insulin was larger than after subcutaneous injection of 10U of insulin. The relative bioefficacy of TECHNOSPHERE™/Insulin was calculated to be 19.0%, while the relative bioavailability was determined to be 25.8% in the first three hours.

[0097] The data also show that inhalation of TECHNOSPHERE™/Insulin resulted in a much more rapid onset of action than s.c. insulin injection that was close to the onset of action of i.v. insulin injection, while duration of action of TECHNOSPHERE™/Insulin was comparable to that of s.c. insulin injection.

[0098] The drug was well tolerated and no serious adverse events were reported during the entire trial.

Example 3: Removal of Impurity From Proprietary Peptide

[0099] A proprietary peptide containing an impurity was trapped in fumaryl diketopiperazine, forming a peptide/fumaryl diketopiperazine precipitate. The precipitate was washed with water to remove the impurity. The peptide is rather unstable and trapping it in fumaryl diketopiperazine markedly improves its stability; both as a dry powder and in aqueous suspension for injection.

Example 4: Stabilized Glucagon Formulations

Formulation

[0100] Glucagon was formulated under sterile conditions, into a stabilized complex by precipitation in acidic solution with fumaryl diketopiperazine (3,6-bis[N-fumaryl-N-(n-butyl)amino]-2,5-diketopiperazine). The complex was washed and lyophilized, yielding a sterile dry powder formulation of diketopiperazine/glucagon (hereinafter referred to as "TG") containing from 1.2 to 8.2% glucagon by weight, depending upon the formulation parameters desired (allowing physicians to increase dose yet keep the volume constant). The TG powder was suspended in an appropriate media suitable for subcutaneous delivery in a MiniMed 507C infusion pump.

Stability Protocol

[0101] Glucagon and TG were suspended in infusion media and incubated at 40 °C in a water bath for varying amounts of time up to 150 hours.

Glucagon HPLC Analysis

[0102] An adaptation of USP method for glucagon analysis was employed. A Waters Symmetry Shield RP8 column (5 μ m, 3.9 x 150 mm) and guard RP8 column (5 μ m, 3.9 x 20 mm) were used at a flow rate of 1 mL/min. and a detection wavelength of 214 nm. The gradient method consisted of mobile phase A: 9.8 g NaH₂PP4 (0.0816 M) and 170 mg L-cysteine (1.4 mM) per liter HPLC grade water, adjusted pH to 2.6 with phosphoric acid; and B: acetonitrile. Glucagon solutions were diluted as needed with water and injected. TG samples were prepared by adding 1/10th volume 1 M Tris pH 10.0 to sample to solubilize the fumaryl diketopiperazine.

Rat Study Protocol

[0103] Sprague Dawley rats 200-250 g were fasted overnight and given subcutaneous injection of glucagon or TG (0.75 mg/kg) in an appropriate media that had been held at 25 °C for 0, 24, or 48 hours. Blood samples were taken at -10, -5, 0, 5, 10, 15, 20, 30, 45, and 60 minutes post dose and analyzed for blood glucose (HemCue B-glucose analyzer, Hemocue AB, Angelholm Sweden). Mean baseline was determined (pre-dose measurements) and was subtracted from the subsequent data and plotted vs. time. This was done to assure that the TG formulation, which appeared to not degrade significantly, showed appropriate pharmacological activity.

Results

[0104] Following 40 °C incubation, HPLC analysis showed an increase in breakdown products in the glucagon preparation. By contrast, TG has only one minor degradation peak (RT=6) which correlated with the slightly less active oxidative form of glucagon. Glucagon without diketopiperazine (i.e. without TECHNOSPHERES™) had many degradation peaks, some of which contributed to an enhanced effect and others that reduced the potency of glucagon.

[0105] The TG sterile lyophilized powder was shipped frozen to a hospital, where it was re-suspended in sterile media. The material re-suspended well and each vial was continuously infused over a 72 hour period.

Conclusion

[0106] Standard preparations of glucagon are not suitable for regulation of blood glucose by continuous subcutaneous infusion. Administration of such preparations containing variable amounts of the deamidated and hydrolysed forms resulted in highly variable blood glucose levels. Suspensions of TECHNOSPHERES™/glucagon, which is stabilized, does not aggregate and contains clinically irrelevant amounts of breakdown products. As such TG can be and has been used as a therapy for hyperinsulinemia, providing consistent, elevated glucose levels when administered subcutaneously over time.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US5547929A [0006]
- US5888477A [0007]
- US6071497A [0010] [0036]

Non-patent literature cited in the description

- **PFEIFFER**Am. J. Med., 1981, vol. 70, 579-88 [0003]
- **CERASI et al.**Diabetes, 1972, vol. 21, 224-34 [0004]
- **POLONSKY et al.**N. Engl. J Med., 1988, vol. 318, 1231-39 [0004]
- **LEAHY**Curr. Opin. Endocrinol. Diabetes, 1995, vol. 2, 300-06 [0004]
- **KATCHALSKI et al.**J. Amer. Chem. Soc., 1946, vol. 68, 879-80 [0051] [0052]
- **KOPPLE et al.**J. Org. Chem., 1968, vol. 33, 2862-64 [0051]

Patentkrav

1. Sammensætning, som omfatter mikropartikler, som omfatter insulin, som er kompleksbundet til et diketopiperazin, til anvendelse inden for medicin;5
hvor sammensætningen frigiver monomer- eller dimerinsulin ved indgivelse; og hvor mikropartiklerne kan opnås ved en fremgangsmåde, omfattende -
(i) at tilvejebringe diketopiperazinet som præ-formede mikropartikler i en suspen-
sion;
(ii) at kombinere mikropartiklerne i en suspension med en oplosning af insu-
lin; og10
(iii) at lyofilisere eller frysetørre suspensionen;
hvorved der frembringes diketopiperazinmikropartikler, som har en insulin-
coating.
2. Sammensætning til anvendelse ifølge krav 1, hvor diketopiperazinet er
15 2,5-diketo-3,6-di(4-X-aminobutyl)piperazin, hvor X repræsenterer en succi-
nyl-, glutaryl-, maleyl- eller fumarylgruppe.
3. Sammensætning til anvendelse ifølge krav 1 eller 2, hvor diketopiperazinet
er 2,5-diketo-3,6-di(4-fumaryl-aminobutyl)piperazin.
20
4. Sammensætning til anvendelse ifølge krav 1, hvor insulinen, som er kom-
pleksbundet til et diketopiperazin, er monomerinsulin.
- 25 5. Sammensætning til anvendelse ifølge krav 1, hvor insulinen, som er kom-
pleksbundet til et diketopiperazin, hovedsageligt befinder sig i dimer- og mo-
nomertilstanden i stedet for hexamertilstanden.

6. Sammensætning til anvendelse ifølge krav 1, hvor insulinen er human insulin.

5 7. Sammensætning til anvendelse ifølge krav 1, hvor mikropartiklerne kan opnås ved fremgangsmåden ifølge krav 1, og hvor fremgangsmåden desuden omfatter trinnet, hvor der fjernes zinkioner, ved at mikropartiklerne vaskes med et egnet opløsningsmiddel.

10 8. Sammensætning til anvendelse ifølge krav 1, hvor insulinen er en stabiliseret form for insulin, som er blevet renset for at fjerne zink.

15 9. Sammensætning til anvendelse ifølge et af kravene 1 til 8, hvor sammensætningen anvendes til behandling af diabetes.

20 10. Sammensætning til anvendelse ifølge et af kravene 1 til 8, hvor sammensætningen anvendes til behandling af type II diabetes.

25 11. Anvendelse af en sammensætning, som omfatter mikropartikler, som omfatter insulin, som er kompleksbundet til et diketopiperazin, inden for fremstillingen af et medikament til behandling af diabetes; hvor sammensætningen frigiver monomer- eller dimerinsulin ved indgivelse; og hvor mikropartiklerne kan opnås ved en fremgangsmåde, som omfatter at tilvejebringe diketopiperazinet som præ-formede mikropartikler i en suspension; tilsætte en oplosning af insulin til suspensionen; og derefter lyofilisere eller frysetørre suspensionen for at danne diketopiperazinmikropartikler, som har en insulin-coating.

12. Anvendelse ifølge krav 11, hvor diabetes er type II diabetes.

DRAWINGS




FIG. 1A




FIG. 1B




FIG. 2A




FIG. 2B