LV 11631

(19) LATVIJAS REPUBLIKAS
PATENTU VALDE

(11) LV 11631 B

(51) Int.Cl. [6] C12N15/13
C07K16/28
C07K16/46
A61K39/395

Latvijas patents uz izgudrojumu
1995.g. 30.marta Latvijas Republikas likums

(12) Īsziņas

(21) Pieteikuma numurs: P-96-218

(22) Pieteikuma datums: 04.07.1996

(41) Pieteikuma publikācijas datums: 20.12.1996

(45) Patenta publikācijas datums: 20.04.1997

(30) Prioritāte:
9325182.5 08.12.1993 GB

(86) PCT pieteikums:
PCT/IB94/00387, 21.11.1994

(87) PCT publikācija:
WO 95/16038, 15.06.1995

(73) Īpašnieks(i):
T CELL SCIENCES, INC.;
115 Fourth Avenue, Needham MA 02194, US

(72) Izgudrotājs(i):
Augustine, Yee-Tharn LIN (US)

(74) Pilnvarotais vai pārstāvis:
Ābrams FOGELS,
Patentu birojs "ALFA-PATENTS",
Mārstaļu iela 2/4, Rīga LV-1050, LV

(54) Virsraksts: **Cilvēkam piemērota antiviela un tās pielietojums**

(57) Kopsavilkums: Aprakstīta cilvēkam piemērota antiviela vai tajā ietilpstoša antigēnu saistoša olbaltumviela ar specifisku saistību pret tādu T šūnu subpopulāciju, kurai raksturīgas mainīgas uzbūves beta ķēdes, sevišķi pret tādu subpopulāciju, kas ekspresē cilvēka Vb 5.2 un/vai 5.3, kā arī Vb 8.1. Izgudrojums attiecas arī uz šādu antivielu iegūšanu, tās saturošām farmaceitiskām kompozīcijām un to pielietojumu ārstniecībā, sevišķi autoimūno slimību ārstēšanai.

Izgudrojuma formula

1. Cilvēkam piemērota antiviela vai tajā ietilpstošā antigēnu saistošā olbaltumviela ar tādu aminoskābju secību, kas ietver visu vai daļu no tās aminoskābju secības, kura pamatā ietilpst komplimentaritāti nosakošos apgabalos monoklonālajā antivielā ar specifisku saistību pret noteiktu T šūnu subpopulāciju, kas ekspresē mainīgos T šūnu antigēna receptora beta-ķēdes apgabalus Vβ5.2 vai Vβ5.3, saistībā ar dažādiem izmeklētiem olbaltumvielas fragmentiem no minētās monoklonālās antivielas, pie kam antiviela vai tajā ietilpstošā antigēnu saistošā olbaltumviela spēj saistīties ar minētajām noteiktajām T šūnu subpopulācijām vismaz ar tikpat lielu afinitāti, kā minētā monoklonālā antiviela, pie kam

cilvēkam piemērotā antiviela vai tajā ietilpstošā antigēnu saistošā olbaltumviela satur aminoskābju secību TM27 V_K:

1	**DIQMTQSPSSLSASVGDRVTITCSASQGISNYLNWYQQTPGKAPKLLIYY**	50
51	**TSSLHSGVPSRFSGSGSGTDYTFTISSLQPEDIATYYCQQYSKLPRTFGQ**	100
101	**GTKLQIT**	107

un papildus tai satur aminoskābju secību, kas ņemta no šādas secību grupas: secība TM27 V_H

1	**QVQLQESGPGLVRPSQTLSLTCTVSGFSLTAYGVNWVROPPGRGLEWLGM**	50
51	**IWGDGNTDYNSALKSRVTMLKDTSKNQKSLRLSSVTAADTAVYYCARDRV**	100
101	**TATLYAMDYWGQGSLVTVSS**	120

secība TM27 V_H, kurā 48. aminoskābes atlikums aizvietots ar izoleicīna atlikumu (I);

secība TM27 V_H, kurā 78. un 79. aminoskābju atlikumi ir valīns (V) un fenilalanīns (F);

secība TM27 V_H, kurā aminoskābju atlikumi no 67. līdz 70. ar secību VTML ir attiecīgi aizvietoti ar secību LSIS, bet 73. aminoskābes atlikums ir asparagīns (N);

secība TM27V_H, kurā 92. aminoskābes atlikums ir arginīns (R).

2. Cilvēkam piemērota antiviela vai tajā ietilpstošā antigēnu saistošā olbaltumviela ar tādu aminoskābju secību, kas ietver visu vai daļu no tās aminoskābju secības, kura pamatā ietilpst komplimentaritāti nosakošos apgabalos monoklonālajā antivielā ar specifisku saistību pret noteiktu T šūnu subpopulāciju, kas ekspresē mainīgo T šūnu antigēna receptora beta-ķēdes apgabalu Vβ8.1, saistībā ar dažādiem izmeklētiem olbaltumvielas fragmentiem no minētās monoklonālās antivielas, pie kam antiviela vai tajā ietilpstošā antigēnu saistošā olbaltumviela spēj saistīties ar minētajām noteiktajām T šūnu subpopulācijām vismaz ar tikpat lielu afinitāti, kā minētā monoklonālā antiviela, pie kam

cilvēkam piemērotā antiviela vai tajā ietilpstošā antigēnu saistošā olbaltumviela satur aminoskābju secību TM29 V_K:

1 **DIQMTQSPSSLSASVGDRVTITCRASSSVNYIYWYQQTPGKAPKLLIYYT** 50
51 **SNLAPGVPSRFSGSGSGTDYTFTISSLQPEDIATYYCQQQFTSSPFTFGQGT** 100
101 **KLQIT** 106.

un papildus tai satur aminoskābju secību, kas ņemta no šādas secību grupas:
secība TM29 V_H

1 **EVQLVESGGGVVQPGRSLRLSCSSSGFTFSNFGMHWVRQAPGKGLEWVAY** 50
51 **ISSGSSTIYYADTLKGRFTISRDNSKNTLFLQMDSLRPEDTGVYFCARRG** 100
101 **EGAMDYWGQGTPVTVSS** 117.

secība TM29 V_H, kurā 23. un 24. aminoskābju atlikumi abi aizvietoti ar alanīna atlikumu (A);

secība TM29 V_H, kurā 75. aminoskābes atlikums ir aizvietots ar prolīna atlikumu (P);

secība TM29 V_H, kurā aminoskābju atlikumi 92. un 93. aizvietoti, attiecīgi, ar alanīna (A) un metionīna (M) atlikumu, bet 95. aminoskābes atlikums ir aizvietots ar tirozīna (Y) atlikumu.

3. Cilvēkam piemērota antiviela pēc 1. vai 2. punkta, kas veido izotipa IgG2a secību.

4. Cilvēkam piemērota antiviela pēc 3. punkta, kurā izmainītas viena vai vairākas aminoskābes konstantajos Ig domēnos ar nolūku izmainīt minētā konstantā domēna efektora funkciju.

5. Cilvēkam piemērota antiviela vai tajā ietilpstošā antigēnu saistošā olbaltumviela pēc 1. vai 2. punkta, kas iegūta ar rekombinatās DNS tehnoloģiju.

6. Paņēmiens cilvēkam piemērotas antivielas vai tajā ietilpstošās antigēnu saistošās olbaltumvielas pēc jebkura no iepriekšējiem punktiem iegūšanai, kas paredz:

a) izolēt pirmo DNS secību, kas pamatā kodē vismaz vienu komplimentaritāti nosakošo apgabalu peles monoklonālajā antivielā anti-Vβ5.2/Vβ5.3 vai anti-Vβ8.1 ar specifisku saistību pret noteiktu T šūnu subpopulāciju;

b) klonēt minēto pirmo DNS secību piemērotā vektorā:

c) ar mutaģenēzi ievadīt izmeklētas cilvēka olbaltumvielas aminoskābes, nolūkā izveidot konstrukciju, kurā iesaistīti komplimentaritāti nosakošie apgabali;

d) transformēt saimniekšūnu ar posmā c) iegūto konstrukciju;

e) kultivēt minēto transformēto saimniekšūnu, lai producētu minēto cilvēkam piemēroto antivielu vai tajā ietilpstošo antigēnu saistošo olbaltumvielu.

7. Paņēmiens pēc 6. punkta, kurā saimniekšūna ir CHO šūna.


8. Cilvēkam piemērotā antiviela vai tajā ietilpstošā antigēnu saistošā olbaltumviela pēc 1. vai 2. punkta pielietojumam ārstniecībā.

9. Farmaceitiskā kompozīcija, kas satur cilvēkam piemēroto antivielu vai tajā ietilpstošo antigēnu saistošo olbaltumvielu pēc 1. vai 2. punkta kopā ar farmaceitiski pieņemamu atšķaidītāju.

10. Cilvēkam piemērotās antivielas vai tajā ietilpstošās antigēnu saistošās olbaltumvielas pēc 1. vai 2. punkta ***pielietojums*** tāda medikamenta ražošanai, kas paredzēts Krona (Crohn) slimības ārstēšanai.

11. Cilvēkam piemērotās antivielas vai tajā ietilpstošās antigēnu saistošās olbaltumvielas pēc 1. vai 2. punkta ***pielietojums*** tāda medikamenta ražošanai, kas paredzēts multiplās sklerozes ārstēšanai.

FIG.3




PELES 4H11	y = 393.85 - 5.0535x R^2 = 0.985
TM27L/NS0	y = 415.77 - 2.3218x R^2 = 0.939
TM27L/CHO	y = 415.64 - 2.1880x R^2 = 0.987

UG/ML	BEZ IEZĪMES [%]	PELES 4H11	TM27L/NS0	TM27L/CHO
5.000	50.000	149.687	303.633	299.073
3.750	42.900	181.330	296.867	328.207
2.500	33.300	219.803	349.850	342.667
1.875	27.300	251.783	367.640	358.120
1.250	20.000	288.993	360.657	374.003
0.938	15.800	308.113	380.870	378.773
0.625	11.100	333.177	388.950	395.623
0.312	5.900	349.383	405.483	401.297
0.156	3.030	380.860	400.113	402.853
0.000	0.000	417.797	417.797	417.792

FIG.4

TM27 SKEČERDA ANALĪZE AR HPB ŠŪNĀM



	Kd (nM)
TM27 CH0	12.3
TM27 NS0	8.2
4H11	4.5

Humanized Antibodies and Uses Thereof

The present invention relates to humanized antibodies and binding proteins thereof, capable of binding to T cells exhibiting particular variable beta chains, and particularly those subpopulations expressing human Vβ 5.2 and/or 5.3, and Vβ 8.1. The present invention also relates to the preparation of such antibodies, to pharmaceutical compositions containing them, and therapeutic utilization of the antibodies, specifically in autoimmune diseases.

T cells play a pivotal role in the differentiation and regulation of effector mechanisms within the immune system (Paul et al., (1987) Science 195:1293-1300). The co-recognition of antigen and major histocompatibility molecules by a T cell must be specific and precisely regulated, since improper immune regulation fosters autoimmunity. Several laboratories have studied diseases in which there appears to be improper immune regulation, such as autoimmunity, and some forms of immunodeficiency, and have implicated T cells in the pathogenesis of such diseases.

Several situations exist where there have been reported a clonal or oligoclonal expansion of a particular T call receptor composition. The most obvious examples are in conditions of malignancy which have resulted in a T cell leukemia or lymphoma. In situations of T cell leukemias or lymphomas, the T cell receptor acts as a unique tumor marker since the T cell receptor is stably rearranged and presented on the surface of the cell. Another situation where a particular T cell receptor composition is implicated is in the recipient of an organ graft whose T lymphocytes have T cell receptors making them aggressive against the MHC molecules of the donor individual, as for instance the donor of a bone marrow graft.

More importantly, several groups have reported selective T cell antigen receptor V region gene usage in certain autoimmune situations. For instance, Grunwald et al. have noted a

preferential expression of the Vα 2.3 gene product in CD4+ T cells in the broncheoalveolar lavage when compared to peripheral blood lymphocytes of patients with Sarcoidosis (Grunwald et al., (1992) Eur. J. Immunol. 22:129). In Kawasaki disease, the preferential expansion of Vβ 2 and Vβ 8 T cells was noted at the onset of disease (Abe et al., (1992) Proc. Natl. Acad. Sci. USA 89:4066).

Rheumatoid arthritis has also been extensively studied in this regard. Several investigators have noted preferential expansion of subsets of T cells, as for instance: DerSimonian et al., (1993) J. Exp. Med. 177:1623 (preferential expansion of Vα 12.1 bearing T cells in CD8+ peripheral blood T lymphocytes); Stamenkovic et al., (1988) Proc. Natl. Acad. Sci. USA 85:1179 (synovial membrane-infiltrating T cells grown in IL2 were oligoclonal by southern blot analysis); Paliard et al., (1991) Science 253:34 (hypothesized that a Superantigen activated Vβ 14+ T cells, including autoreactive T cells which expand clonally and migrate to the synovial fluid of rheumatoid arthritis patients); Howell et al., (1991) Proc. Natl. Acad. Sci. USA 88: 10921 (noted in particular Vβ 3, 14, and 17 T cell V region gene usage in IL 2R+ cells from synovial fluid of rheumatoid arthritis patients); Uematsu et al., Proc. Natl. Acad. Sci. USA 88:8534 (showed oligoclonal T cell V region gene usage in synovial fluid T cells of a single RA individual); and International Patent Application No. WO90/06758 (implicating Vβ 3, 9, and 10 in RA).

Inflammatory bowel disease has also been extensively studied. several groups have noted expanded T cell populations or preferential T cell receptor V region gene usage as for instance: Posnett et al., (1990) J. Clin. Invest. 85:1770; Spencer et al., (1991) J. Clin. Pathol. 44:915; Trejjdosiewicz et al., (1991) Clin. Exp. Immunol. 84:440; and Van Kerckhove et al., (1992) J. Exp. Med. 175:57. Still others have reported preferential T cell V gene usage in *Mycobacterium leprae* (van Shooten et al., (1992) Proc. Natl. Acad. Sci. USA 89:11244; Wang et al., (1993) Proc. Natl. Acad. Sci. USA 90:188.

In humans, expansion of Vβ 8.1 T cells in inflammatory tissue has been found in association with several autoimmune diseases, including Crohn's Disease (Posnett et al., (1990) J. Clin. Invest. 85:1770-1776), Kawasaki Disease (Abe et al., (1992) Proc. Natl. Acad. Sci. USA 89:4066-4070 and Abe et al., (1993) J. Exp. Med. 177:791-796) and rheumatoid arthritis (Brennan et al., (1988) Clin. Exp. Immunol. 73:417-423).

Multiple Sclerosis (MS) is another autoimmune disease that has been intensively studied. MS is an immune mediated disease characterized by central nervous system

mononuclear cell infiltration and demyelination. Although the pathogenesis of MS is unknown, both genetic and environmental factors have been implicated in the disease process. Major elements of the genetic predisposition include an association of disease with particular class II major histocompatibility complex (MHC) haplotypes, in particular HLA-DR21 and DQW1 (Terasaid et al., (1976) Science 1933:1245-1247; Ho et al., (1982) Immuno-genetics 15:509-517; Spielman et al., (1982) Epidemiol. Rev. 4:45-65; Francis et al., (1986) Lancet 1:211; Elian et al., (1987) Disease Markers 5:89-99; Urban et al., (1988) Cell 54:577-592, Vandenbark et al., (1989) Nature 341:541-544: Howell et al., (1989) Science 246:668-670).

It has been shown that T cells isolated from the cerebrospinal fluid of patients suffering from MS utilize a limited set of V region genes. The demonstration of in vivo activated myelin basic protein specific T cells in MS patients implicates MBP reactive T cells in the pathogenesis of the disease (Wucherpfennig et al., (1990) Science 248:1016-1019). When the TCR Vβ usage of MBP reactive T cell lines is determined via polymerase chain reaction (PCR) amplification of cDNA with T cell receptor Vβ primers, preferential usage of a limited number of Vβ genes has been found (Wucherpfennig et al., (1990) (supra). Vβ 17 and to a lesser extent Vβ 12 are frequently used in the recognition of the immunodominant region of the human autoantigen MBP); Oksenberg et al., (1993) Nature 362:68. The results of some studies also imply that there is a limited T cell receptor Vα gene expression in MS brain lesions (Oksenberg et al., (1990) Nature 345:344-346). T cell repertoire analysis using quantitative PCR and monoclonal antibody (mAb) staining has shown that Vβ 5.2 and/or 5.3 are predominantly utilised by the myelin basic protein (MBP) specific T cells isolated from MS patients compared with control (Oksenberg et al., (1993) (supra) (rearranged Vβ 5.2 genes were detected in the brains of patients with a certain HLA phenotype) and Kotzin et al. (1991) Proc. Natl. Acad. Sci. USA 88:9196 (a bias for use of the β chain variable region 5.2 and to a lesser extent Vβ 6.1 was seen among MBP specific clones from patients with MS).

Currently, no effective treatment for MS is known. (*Harrison's Principles of Internal Medicine*, 12th ed. Wilson et al., McGraw Hill, Inc. 1991). Therapeutic efforts are directed toward amelioration of the acute episode, prevention of relapses or progression of the disease, and relief of symptoms.

However, expression of the mouse Vβ 8.2 variable region gene has been found to correlate with experimental allergic encephalomyelitis (EAE), a mouse model of human MS. It has been demonstrated that treatment with mouse Vβ 8.2 specific mAb can both prevent and

reduce the disease (Acha-Orbea et al., (1988) Cell 54:263 and Urban et al., (1988) Cell 54:577). Thus, there is a great need to develop an antibody or "antibody-like" molecule, suitable for treatment of this disease.

Antibodies typically comprise two heavy chains linked together by disulfide bonds and a light chain associated with the N-terminal end of each heavy chain. Each heavy chain has at its N-terminal end a variable domain followed by a constant domain at its other end. Each light chain also has at its N-terminal end a variable domain followed by a constant domain. The variable domains of each pair of light and heavy chains form the antigen binding site. The variable domains on the light and heavy chain have the same general structure and each domain comprises a framework of four regions, whose sequences are relatively conserved, connected by three complementarity determining regions (CDRs). The four framework regions adopt somewhat of a beta-sheet conformation and the CDRs form loops connecting the beta-sheet construction. The CDRs are held in close proximity by the framework regions and contribute to the formation of the antigen binding site. CDRs and framework regions of antibodies may be determined by reference to the Kabat numbering system (Kabat et al., (1987) "Sequences of Proteins of Immunological Interest", US Dept. of Health and Human Services, US Government Printing Office) in conjunction with x-ray crystallography, as set forth in WO91/09967.

In order to produce an antibody which can target a specific antigen, the procedure of Kohler and Milstein (Kohler et al., (1976) Nature 256:495-497) is usually used. This generally involves immunising a mouse with the antigen, fusing spleen cells from the immunised mouse with mouse myeloma cells and selecting from the hybridomas thus produced one or more hybridomas which secrete a monoclonal antibody specific for the target antigen.

It would be desirable to use such mAbs in therapy. However, such mAbs are essentially of murine origin and are therefore themselves antigenic in humans. If such mAbs are administered repeatedly to a human, the human's immune system mounts a response to the mouse mAb, thus rendering it ineffective.

It has therefore been proposed, originally by Winter and his co-workers (see, for instance, Reichmann et al., (1988) Nature 332:323-327 and Verhoeyen et al., (1988) Science 239:1534-1536) that the CDRs of a mouse mAb should be grafted onto a human framework to produce a CDR-grafted antibody having the binding properties of the donor mouse mAb and the human compatibility of the acceptor human framework.

However, the function of an antibody molecule is dependent on its three dimensional structure, which in turn is dependent on its primary amino acid sequence. Changing the amino acid sequence of an antibody may adversely affect its activity. Likewise, fragments of the antibody may not retain the appropriate three dimensional structure necessary to foster binding activity. Moreover, a change in the DNA sequence coding for the antibody may affect the ability of the cell containing the DNA sequence to express, secrete or assemble the antibody. The exact residues comprising CDRs are difficult to define and do not necessarily correspond to all the residues in the hypervariable regions, as defined by the Kabat numbering system. There are also critical framework residues which are important in positioning the CDRs for interaction with antigen or which are involved in interactions between the heavy and light chains. It may be necessary to alter certain framework residues so that they correspond to the donor residues at certain positions, rendering the CDR-grafted antibody less "human" in character.

There have been published various proposals for identifying the CDR and framework residues which need to be changed to donor residues in order to produce a useful CDR-grafted antibody (see, for instance, Queen et al., WO90/07861; Kurrle et al., EP-A-0 403 156; Adair et al., WO91/09967; Queen et al., WO92/11018; and Bendig et al., WO92/01568). It emerges from these documents that the production of a useful CDR-grafted antibody in any particular case is not straightforward.

Despite the problems which are inherent in attempting to produce a specific CDR-grafted antibody, in a preferred embodiment, the present inventors have succeeded in producing a CDR-grafted antibody based on human framework regions and having an antigen binding site specific for variable beta chain regions peculiar to certain subpopulations of T cells. Surprisingly, the humanized antibodies of the invention demonstrate as good or better binding affinity to target T cells when compared to their murine prototypes. The currently claimed antibodies or parts thereof offer the further advantage of being less immunogenic than their murine prototypes, thus diminishing adverse patient reactions when used in therapy.

SUMMARY OF THE INVENTION

The present invention is directed to humanized antibodies having a selective binding specificity for certain T cell subpopulations, which antibodies are highly sensitive in their binding to these subpopulations, and demonstrate specificity and affinity in that binding,

comparable with, if not better than, that of the prototype mAbs from which a substantial portion of their CDR regions are derived. The present invention is also directed to methods of preparation of such humanized antibodies, utilizing novel cDNA encoding certain hypervariable and framework residues and grafting these into human heavy chain and human light chain frameworks. Further embodiments include pharmaceutical compositions and therapeutic methods of using these antibodies.

In one particularly preferred embodiment, the mouse mAb, 16G8, which recognizes human Vβ 8.1, was humanized by CDR grafting certain CDR and select framework residues from the murine mAb into KOL heavy chain and REI light chain frameworks. The cDNAs encoding humanized heavy (IgG1) and light (K) chains in mammalian cell expression vectors with Neo and DHFR selection markers, respectively, were transfected into a DHFR- Chinese hamster fibroblast (CHO) cell line followed by selection and amplification. The humanized mAb secreted, designated "TM29", maintains the specificity for human TCR Vβ 8.1 with affinity comparable to that of the prototype mouse mAb (16G8). The antibodies of the invention have demonstrated that humanization of anti-TCR antibodies can be accomplished while maintaining the correct subset specificity and affinity.

Other embodiments of the invention include the use of these humanized antibodies as a therapeutic agent for treatment of human autoimmune disease, and particularly Crohn's disease.

In another particularly preferred embodiment of the present invention, a mouse mAb, TM23, which recognizes human Vβ 5.2 and 5.3, was humanized by CDR grafting into NEWM heavy chain and REI light chain frameworks. The cDNAs encoding humanized heavy (IgG1) and light (K) chains in mammalian cell expression vectors with Neo and DHFR selection markers, respectively, were transfected into a DHFR-Chinese hamster fibroblast (CHO) cell line followed by selection and amplification. The humanized mAb that was secreted was designated "M27", and maintains specificity for human TCR Vβ 5.2 and 5.3 with affinity comparable to, if not better than, that of the prototype mouse mAb (TM23). Other embodiments of the invention include the use of TM27 as a therapeutic agent for treatment of human MS.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 depicts various variable light chain sequences and heavy chain sequences for TM27:

Figure 2 depicts various variable light chain sequences and heavy chain sequences for TM29;

Figure 3 depicts the results of a competition assay comparing TM27L/NSO and TM27L/CHO with 4H11; and

Figure 4 depicts a Scatchard plot of the experiment shown in Figure 3.

DETAILED DESCRIPTION OF THE INVENTION

Certain definitions used in the context of the description of the invention are set forth as follows.

In the present application, the term "antibody" is used to describe Igs or any fragments thereof, light chain or heavy chain monomers or dimers, and single chain antibodies, such as a single chain Fvs in which the heavy and light chain variable domains are joined by a peptide linker, whether natural or produced by recombinant DNA technology or otherwise, provided that the antibody includes at least one antigen binding site. The remainder of the antibody need not comprise only Ig-derived protein sequences. For instance, a gene may be constructed in which a DNA sequence encoding part of a human Ig chain is fused to a DNA sequence encoding the amino acid sequence of a polypeptide effector or reporter molecule. Thus "antibody" encompasses hybrid antibodies (see below).

The term "binding protein" as used herein refers to a construct having a particular amino acid sequence derived from an antibody or combination of antibodies, said sequence sterically configured in such a manner as to form at least one antigen binding site.

The abbreviation "mAb" is used to indicate a monoclonal antibody as produced by a hybridoma or derivative cell line.

The term "recombinant antibody" is used to describe an antibody produced by a process involving the use of recombinant DNA technology.

The term "CDR" or "complementarity determining region" refers to those portions of an antibody's heavy chain or light chain variable regions juxtapositioned in three-dimensional space to form an antigen binding surface.

The term "chimeric antibody" is used to describe an antibody in which the variable domains as a whole are derived from an antibody from a first mammalian species and have been fused onto at least one constant domain from an antibody from a different mammalian species.

The term "CDR-grafted" is used to describe an antibody or portion thereof in which the CDRs are substantially derived from an antibody from a first mammalian species, grafted into a variable framework regions substantially derived from a second mammalian species. Certain

select amino acids in said framework regions may also be derived from said first mammalian species.

The term "hybrid antibody" is used to describe a protein comprising at least the binding portion of an Ig attached by peptide linkage to at least part of another protein. It will be appreciated that certain skilled workers may also use the word "chimeric" to describe such constructs, but in the present specification such constructs are referred to as hybrid antibodies and the term chimeric antibodies is used in the sense defined above.

The term "humanized antibody" is used to describe an antibody having at least one, and preferably two or three, of its CDRs in one or both of the variable domains derived from an antibody from a first species, it being understood that this may involve certain select framework amino acids in conjunction with certain hypervariable amino acid sequences. The remaining Ig-derived parts of the antibody being derived from one or more different antibodies. The variable domains may be made by use of recombinant DNA technology or by peptide synthesis.

"Expression vector" includes vectors which are capable of expressing DNA sequences contained therein, i.e. the coding sequences are operably linked to other sequences capable of effecting their expression. A useful, but not always necessary (i.e. insect cells), element of an effective expression vector is a marker encoding sequence, i.e. a sequence encoding a vector sequence which results in a phenotypic property (e.g. neomycin resistance, methionine sulfoximine resistance or tryptophan prototrophy) of the cells containing the protein which permits those cells to be readily identified. In sum, "expression vector" is given a functional definition and any DNA sequence which is capable of effecting expression of a specified contained DNA code is included in this term as it is applied to the specified sequence. As at present, such vectors are frequently in the form of plasmids. Thus "plasmid" and "expression vector" are often used interchangeably. However, the invention is intended to include such other forms of expression vectors which serve equivalent functions and which may, from time to time, become known in the art, including retroviruses, adenoviruses, in vitro systems (Baranov et al., (1989) Gene 84:2:463) and the like.

As stated previously, the DNA sequences will be expressed in host cells after the sequences have been operably linked to (i.e. positioned to ensure the functioning of) an expression control sequence. These expression vectors are typically replicable in the host organisms either as episomes or as an integral part of the host chromosomal DNA.

"Recombinant host cells" refers to cells which have been transformed with vectors constructed using recombinant DNA techniques. By virtue of this transformation, the host cell is able to produce the desired product in useful quantities, rather than in lesser amounts, or more commonly, in less than detectable amounts, as one would expect to be produced by the untransformed host. The antibody and binding proteins of the present invention may be produced by a recombinant host cell in quantities useful to carry out additional experimentation or in commercially acceptable quantities, such as about 100 grams or more.

In descriptions of processes for isolation of antibodies or binding proteins thereof from recombinant hosts, the terms "cell" and "cell culture" are used interchangeably to denote the source of antibody unless it is clearly specified otherwise. In other words, recovery from the "cells" may mean either from spun down whole cells, or from the cell culture containing both the medium and the suspended cells, or, additionally, as is possible in the case of myeloma cell lines, from ascites culture.

The humanized antibodies or binding proteins thereof of the present invention have amino acid sequences that comprise all or a portion of its CDRs substantially derived from a monoclonal antibody having a specificity for a select subpopulation of T cells, and particularly the Vβ 8.1 subpopulation or the Vβ 5.2/5.3 subpopulation. In the preferred embodiments, the monoclonal antibody is murine in origin. The framework amino acid sequence of the variable domains of the antibodies or portions thereof is substantially human in origin in the preferred embodiments, hence the term "humanized antibody". This "humanization" is believed to be useful in reducing the immunogenicity of the antibody when administered therapeutically to human patients. Preferred for use herein are the NEWM or KOL human framework regions for the heavy chain, and REI human framework regions for the light chain. Certain select framework residues are maintained as murine, rather than human. This is believed necessary to achieve the appropriate three-dimensional structuring of the molecule, and to thereby increase binding specificity and affinity for the specific T cell subset.

Any portion of the humanized antibodies (hence, the broad definition of the term antibody) made in accordance with the present teachings is within the contemplation of the present invention, as long as binding specificity and affinity for select subpopulations of T cells is maintained. Thus, binding proteins derived from said antibodies are clearly within the scope of the invention, as are other fragments that maintain these capabilities, at least to a degree suitable for therapeutic use, as described below.

One skilled in the art may test the binding parameters of various constructs made following the teachings herein through the use of conventional binding assays conducted with specific T cells or T cell receptor proteins exhibiting the select variable region that is targeted. For example, competition binding assays with prototype mAbs, binding to various cell lines, Scatchard analyses and the like may be performed and compared to results obtained with prototype mAbs. Alternatively, one may resort to various animal model systems, such as those described below, or clinical trial studies to assess the therapeutic effectiveness of the constructs made in accordance with the invention, it being understood that a construct may be a very suitable therapeutic agent even when its binding pattern to target antigen differs from that of the prototype mAb.

In particularly preferred embodiments, the amino acid sequence of the humanized antibody or binding protein thereof comprises all or a portion of a variable light chain selected from the following group:

TM 29 light chain

1 DIQMTQSPSSLSASVGDRVTITCRASSSVNYIYWYQQTPGKAPKLLIYYT 50

51 SNLAPGVPSRFSGSGSGTDYTFTISSLQPEDIATYYCOOFTSSPFTFGQG 100

101 TKLQIT 106

TM 27 light chain

1 DIQMTQSPSSLSASVGDRVTITCSASQGISNYLNWYQQTPGKAPKLLIYY 50

51 TSSLHSGVPSRFSGSGSGTDYTFTISSLQPEDIATYYCQQYSKLPRTFGQ 100

101 GTKLQIT 107

alone or in combination with all or a portion of a variable heavy chain selected from the following group:

TM 29 heavy chain

1 EVQLVESGGGVVQPGRSLRLSCSSSGFTFSNFGMHWVRQAPGKGLEWVAY 50

51 ISSGSSTIYYADTLKGRFTISRDNSKNTLFLQMDSLRPEDTGVYFCARRG 100

101 EGAMDYWGQGTPVTVSS 117

TM 27 heavy chain

1 QVQLQESGPGLVRPSQTLSLTCTVSGFSLTAYGVNWVRQPPGRGLEWVGM 50

51 IWGDGNTDYNSALKSRVTMLKDTSKNQFSLRLSSVTAADTAVYYCARDRV 100

101 TATLYAMDYWGQGSLVTVSS 120

Select framework residues in the vicinity of the individual CDRs are important to the support of the three-dimensional conformation of the CDRs (Foote and Winter, (1992) J. Mol. Biol. 224:487-499). Thus, in certain preferred embodiments, modified heavy chains were designed. In the first at residues 66 to 69 inclusive and 73, the murine sequence Leu-Ser-Ile-Ser (66-69) and Asp (73) are substituted for the human counterparts.

In other preferred embodiments, murine residues valine and phenylalanine are provided at positions 78 and 79, respectively, or murine arginine is provided at position 92.

Figure 1 exemplifies these substitutions, showing the CDRs in bold and underlining. These preferred antibody constructs (including binding proteins thereof) bind to Vβ 5.2/5.3.

Other preferred antibodies and binding proteins thereof are provided herein by modification of heavy chains at residues 23 and 24. At these positions, the murine counterpart sequence Ala- is substituted. Other constructs demonstrate that residue 75 may be the murine proline residue, or that residues 88, 89 and 91 may be alanine, methionine or tyrosine respectively. These antibody constructs and binding proteins thereof bind Vβ 8.1 T cells and are more fully depicted in Figure 2.

Particularly preferred for use herein are CDR-grafted antibodies consisting essentially of the TM27 light chain in conjunction with the TM27 heavy chain, wherein residue 48 is either murine leucine or isoleucine. Also particularly preferred are CDR-grafted antibodies consisting essentially of the TM29 light chain in conjunction with the TM29 heavy chain.

The humanized antibodies or binding proteins thereof of the present invention may be produced by a variety of techniques, with expression in transfected cells, such as yeast, insect, CHO or myeloma cells, being preferred. Most preferably, the host cell is a CHO host cell.

To design a CDR-grafted antibody or a binding protein thereof in accordance with the present invention, it is first necessary to ascertain the variable domain sequence of an antibody having the desired binding properties. Suitable source cells for such DNA sequences include avian, mammalian or other vertebrate sources such as chickens, mice, rats and rabbits, and preferably mice. The variable domain sequences ($V_H$and V_L) may be determined from heavy

and light chain cDNA, synthesized from the respective mRNA by techniques generally known to the art. The hypervariable regions may then be determined using the Kabat method (supra). The CDRs may be determined by structural analysis using X -ray crystallography or molecular modelling techniques. Composite CDRs may then be constructed containing all the residues corresponding to the hypervariable regions, along with certain select residues from the framework regions. The resulting composite CDRs may then be transferred as the "antigen binding site", while the remainder of the antibody, if it is desired to construct a complete Ig, will include the heavy and light chain constant domains and remaining framework residues. The latter portions may be based on human antibodies of different classes.

Constant domains may be selected to have desired effector functions appropriate to the intended use of the antibody so constructed. For example, human IgG isotypes, IgG_1 and IgG_3 are effective for complement fixation and cell mediated lysis. For other purposes, other isotypes, such as IgG_2 and IgG_4, or other classes, such as IgM and IgE, may be more suitable.

For human therapy, it is particularly desirable to use human isotypes, to minimize antiglobulin responses during therapy. Human constant domain DNA sequences, preferably in conjunction with their variable domain framework regions can be prepared in accordance with wellknown procedures. An example of this is CAMPATH 1H available from Burroughs Wellcome Ltd.

In accordance with preferred embodiments of the present invention, certain CDR-grafted antibodies are provided which contain select alterations to the human-like framework region (in other words, outside of the CDRs of the variable domains), resulting in a CDR-grafted antibody with satisfactory binding affinity. Such binding affinity is preferably from about $10^{5}.M^{-1}$ to about $10^{12}.M.1$ and is more preferably at least about $10^{8}.M^{-1}$. Most preferably the binding affinity is about equal to or greater than that of the prototype antibody (in preferred instances, this prototype antibody would be murine).

In constructing the CDR-grafted antibodies of the present invention, the V_H and/or V_L gene segments may be altered by mutagenesis. One skilled in the art will also understand that various other nucleotides coding for amino acid residues or sequences contained in the Fc portion or other areas of the antibody may be altered in like manner (see, for example, PCT/US89/00297).

Exemplary techniques include the addition, deletion or nonconservative substitution of a limited number of various nucleotides or the conservative substitution of many nucleotides, provided that the proper reading frame is maintained.

Substitutions, deletions, insertions or any subcombination may be used to arrive at a final construct. Since there are 64 possible codon sequences but only twenty known amino acids, the genetic code is degenerate in the sense that different codons may yield the same amino acid. However, the code is precise for each amino acid. Thus, there is at least one codon for each amino acid, i.e. each codon yields a single amino acid and no other. It will be apparent that, during translation, the proper reading frame must be maintained in order to obtain the proper amino acid sequence in the polypeptide ultimately produced.

Techniques for additions, deletions or substitutions at predetermined amino acid sites having a known sequence are well known. Exemplary techniques include oligonucleotide-mediated site-directed mutagenesis and the PCR.

Oligonucleotide site-directed mutagenesis in essence involves hybridizing an oligonucleotide coding for a desired mutation with a single strand of DNA containing the region to be mutated and using the single strand as a template for extension of the oligonucleotide to produce a strand containing the mutation. This technique, in various forms, is described in Zoller and Smith, (1982) Nuc. Acids Res. 10:6487, Norris et al., (1983) Nuc. Acids Res. 11:5103, Zoller and Smith, (1984) DNA 3:479, Kramer et al ., (1982) Nuc. Acids Res. 10:6475.

PCR in essence involves exponentially amplifying DNA in vitro using sequence-specific oligonucleotides. The oligonucleotides can incorporate sequence alterations if desired. The PCR technique is described in Mullis and Foloona, (1987), Meth. Enz. 155:335. Examples of mutagenesis using PCR are described in Higuchi et al., (1988) Nuc. Acids Res. 16:7351, Ho et al., (1989) Gene, 77:51, Ho et al., Engineering Hybridization Restriction Genes without the Use of Restriction Enzymes: Gene Splicing by Overlap Extension; Horton et al., (1989) Gene 77:61.

The nucleotide sequences of the present invention, capable of ultimately expressing the desired antibodies, or binding proteins thereof, can be formed from a variety of different polynucleotides (genomic DNA, cDNA, RNA or synthetic oligonucleotides). At present, it is preferred that the polynucleotide sequence comprises a fusion of cDNA and genomic DNA. The polynucleotide sequence may encode various Ig components (e.g. V, J, D, and C domains). They may be constructed by a variety of different techniques. Joining appropriate genomic and

cDNA sequences is presently the most common method of production, but cDNA sequences may also be utilized (see EP-A-0 239 400 and Reichmann et al., (1988) Nature 332:323.)

Certain suitable expression vectors and host cells are described in US-A-4 816 567.

The vectors and methods disclosed herein are suitable for use in host cells over a wide range of prokaryotic and eukaryotic organisms.

In general, of course, prokaryotes are preferred for cloning of DNA sequences for constructing the vectors useful in the invention. For example, various *E. coli* plasmid cloning vectors are particularly useful, such as the Bluescript M13 vectors available from Stratagene, San Diego, California; pUC19c, Genbank Accession # VB0026, pBR322 (described below) and the like. These examples are, of course, intended to be illustrative rather than limiting.

Prokaryotes may also be used for expression. The aforementioned *E. coli* strains, bacilli such as *Bacillus subtilis*, and other enterobacteriaceae, such as *Salmonella typhimurium* or *Serratia marcescens* and various *Pseudomonas* species may be used.

In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication site as well as marking sequences which are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is typically transformed using one of the many derivatives of pBR322, a plasmid derived from an *E. coli* species (Bolivar et al., (1977) Gene 2:95.) pBR322 contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR322 plasmid, its descendants or other microbial plasmids may also contain, or be modified to contain, promoters which can be used by the microbial organism for the expression of recombinant proteins. Those promoters commonly used in recombinant DNA construction include lactose promoter systems (Chang et al., (1978) Nature 275:615; Itakura et al., (1978) Science 198:1056; Goedell et al., (1979) Nature 281:544) and tryptophan (trp) promoter systems (Goedell et al., (1980) Nuc. Acids Res. 8:4057) and EP-A-0 036 776. While these are the most commonly used, other microbial promoters have been discovered and utilized, and details concerning their nucleotide sequences have been published, enabling a skilled worker to ligate them functionally into plasmid vectors (Siebenlist et al., (1980) Cell 20:269).

In addition to prokaryotes, eukaryotic microbes, such as yeast cultures, may also be used. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among eukaryotic microorganisms, although a number of other strains are commonly available. For

expression in *Saccharomyces*, the plasmid YRp7, for example, (Stinchcomb et al., (1979) Nature 282:39; Kingsman et al., (1979) Gene 7:141; Tschemper et al., (1980) Gene 19:157) is commonly used. This plasmid already contains the trpl gene which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example ATCC No. 44076 or PEP4-1 (Jones, (1977) Genetics 8:85:12). The presence of the trpl lesion as a characteristic of the yeast host cell genome than provides an effective environment for detecting transformation by growth in the absence of tryptophan.

Suitable promoting sequences in yeast vectors include the promoters for 3-phosphoglycerate kinase or other glycolytic enzymes, such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase and glucokinase (Hess et al., (1968) J. Adv. Enzyme Reg. 7:149; Holland et al., (1978) Biochemistry 17:4900.) In constructing suitable expression plasmids, the termination sequences associated with these genes are also ligated into the expression vector 3' of the sequence desired to be expressed to provide polyadenylation of the mRNA and termination. Other promoters, which have the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism and the aforementioned glyceraldehyde-3-phosphate dehydrogenase, an enzyme responsible for maltose and galactose utilization. Any plasmid vector containing a yeast compatible promoter, origin of replication and termination sequences is suitable.

In addition to microorganisms, cultures of cells derived from multicellular organisms may also be used as hosts. In principle, any such cell culture is workable, whether from a vertebrate or an invertebrate organism. However, to date, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure in recent years (Kruse and Patterson, (1973), Tissue Culture, Academic Press). Examples of such useful host cell lines are VERO, HeLa, Chinese hamster ovary (CHO), W138, BHK, COS-7, MDCK and myeloma and gs myeloma (available from Celltech, Slough, U.K.) cell lines. Expression vectors for such cells may include (if necessary) an appropriate origin of replication, as well as a promoter located in front of the gene to be expressed, along with any necessary ribosome binding sites, RNA splice sites, polyadenylation sites and transcriptional terminator sequences.

For use in mammalian calls, the control functions on the expression vectors are often provided by viral material. For example, commonly used promoters are derived from human cytomegalovirus (HCNV), polyoma virus, adenovirus 2 and, most frequently, simian virus 40 (SV40). The early and late promoters of SV40 virus are particularly useful because both are obtained easily from the virus as a fragment which also contains the SV40 viral origin of replication (Fiers et al., (1978) Nature 273:113). Further, it is also possible, and often desirable, to utilize promoter or control sequences normally associated with the desired gene sequence, provided such control sequences are compatible with the host cell system.

An origin of replication may be provided either by construction of the vector to include an erogenous origin, such as may be derived from SV40 or other viral (e.g. polyoma virus, adenovirus, VSV or BPV) source, or may be provided by the host cell chromosomal replication mechanism. If the vector is integrated into the host cell chromosome, the latter is often sufficient.

The vectors containing the DNA segments of interest (e.g. the heavy and light chain encoding sequences and expression control sequences) can be transferred into the host call by well-known methods, which vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment, lipofection or electroporation may be used for other cellular hosts (Maniatis et al., (1990) Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor Press).

Once expressed, the constructs of the present invention can be purified according to standard procedures of the art, including ammonium sulfate precipitation, affinity columns, column chromatography and gel electrophoresis (Scopes, (1982) Protein Purification, Springer Verlag, NY). Binding affinities of the constructs so expressed may be ascertained by techniques known to the art, as more fully exemplified in the example section of this specification.

Substantially pure CDR-grafted antibodies or binding proteins thereof of at least 90 to 95% homogeneity are preferred, and 98 to 99% or more homogeneity is most preferred for pharmaceutical uses. Once purified, partially or to homogeneity as desired, the CDR-grafted antibodies may then be used diagnostically or therapeutically (including extracorporeally) or in developing and performing assay procedures, immunofluorescent stainings and the like (Lefkovite and Pernis, (1979 and 1981) Immunological Methods, Volumes I and II, Academic Press, NY).

The CDR-grafted antibodies or binding proteins thereof of the present invention will typically find use in treating T cell mediated disorders. For example, typical disease states suitable for treatment include autoimmune diseases, such as Type I diabetes, multiple sclerosis, rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease and myasthenia gravis.

Therapeutic use of the T cell antibodies of the invention is premised upon the correlation between a specific immune related disease and the preferential expression of a particular T cell antigen receptor V region gene product or the expanded usage of a particular T cell antigen receptor V gene. The T cell antigen receptor V regions are useful in part because it is possible to regulate the immune response in an individual by specific therapeutic intervention utilizing the T cell antigen receptor. Specifically, the presence or expression of a particular variable region locus has been shown to correlate with particular immune-related disorders. By determining the particular V region loci associated with a particular immune disorder, one can treat the individual by inhibiting the attack by the T cells carrying the particular V region.

The term treatment in the instant invention is meant to include the situations of "prevention", "suppression" or "treatment" of the disease. "Prevention" involves administration of the protective composition prior to the induction of the disease. Thus, for example, in the animal model, EAE, successful administration of a protective composition prior to injection of the encephalitogen that induces the disease results in "prevention" of the disease.

"Suppression" involves administration of the composition after the inductive event but prior to the clinical appearance of the disease. Again, using the EAE example, successful administration of a protective composition after injection of the encephalitogen, but prior to the appearance of neurological symptoms comprises "suppression" of the disease.

"Treatment" involves administration of the protective composition after the appearance of the disease. In the EAE example, successful administration of a protective composition after injection of the encephalitogen and after clinical signs have developed comprises "treatment" of the disease.

Animal model systems which can be used to screen the effectiveness of the antibodies or binding proteins thereof in protecting against or treating the disease are available. Systemic lupus erythematosus (SLE) is tested in susceptible mice as disclosed by Knight et al., (1978) J.Exp. Med. 147:1653 and Reinersten et al., (1978) New Eng. J. Med. 299:515. Myasthenia Gravis (MG) is tested in SJL/J female mice by inducing the disease with soluble AChR protein from another species as described in Lindstrom et al., (1988) Adv. Immunol. 42:233-284.

Arthritis is induced in a susceptible strain of mice by injection of Type II collagen as described by Stuart et al., (1984) Ann. Rev. Immunol. 42:233-284. Adjuvant arthritis is induced in susceptible rats by injection of mycobacterial heat shock protein as described by Van Eden et al., (1988) Nature 331:171-173. Thyroiditis is induced in mice by administration of thyroglobulin as described by Maron et al., (1980) J. Exp. Med. 152:1115-1120. Insulin dependent diabetes mellitus (IDDM) occurs naturally or can be induced in certain strains of mice such as those described by Kanasawa et al., (1984) Diabetologia 27:113. EAE in mouse and rat serves as a model for MS in human. In this model, the demyelinating disease is induced by administration of myelin basic protein as described by Paterson, (1986) Textbook of Immunopathology (Mischer et al., eds.) Grune and Stratton, New York, pp 179-213; McFarlin et al., (1973) Science 179:478-480: and Satoh et al., (1987) J.Immunol. 138:179-184.

The CDR-grafted antibodies or binding proteins thereof of the present invention may also be used in combination with other antibodies, particularly mAbs reactive with other markers on human cells responsible for the diseases. For example, suitable T cell markers can include those grouped into the so-called "Clusters of Differentiation," as named by the First International Leukocyte Differentiation Workshop (Bernhard et al., (1984) Leukocyte Typing, Springer Verlag, NY.

Generally, the present CDR-grafted antibodies or binding proteins will be utilized in purified form together with pharmacologically appropriate carriers. Typically, these carriers include aqueous or alcoholiclaqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride and lactated Ringer's. Suitable physiologically acceptable adjuvants, if necessary to keep the complex in suspension, may be chosen from thickeners such as carboxymethylcellulose, polyvinylpyrrolidone, gelatin and alginates.

Intravenous vehicles include fluid and nutrient replenishers and electrolyte replenishers, such as those based on Ringer's dextrose. Preservatives and other additives, such as antimicrobials, antioxidants, chelating agents and inert gases, can also be present (Mack, (1982) Remington's Pharmaceutical Sciences, 16th Edition).

The constructs of the present invention may be used as separately administered compositions or in conjunction with other agents. These can include various immunotherapeutic drugs, such as cylcosporine, methotrexate, adriamycin or cisplatinum, and immunotoxins. Pharmaceutical compositions can include "cocktails" of various cytotoxic or other agents in

conjunction with the CDR-grafted antibodies or binding proteins thereof of the present invention, or even combinations of constructs according to the present invention and CDR-grafted antibodies having different specificities.

The route of administration of pharmaceutical compositions according to the invention may be any of those commonly known to those of ordinary skill in the art. For therapy, including without limitation immunotherapy, the CDR-grafted antibodies or binding proteins thereof of the invention can be administered to any patient in accordance with standard techniques. The administration can be by any appropriate mode, including parenterally, intravenously, intramuscularly, intraperitoneally, or also, appropriately, by direct infusion with a catheter. The dosage and frequency of administration will depend on the age, sex and condition of the patient, concurrent administration of other drugs, counter indications and other parameters to be taken into account by the clinician.

The constructs of this invention can be lyophilized for storage and reconstituted in a suitable carrier prior to use. This technique has been shown to be effective with conventional immunoglobulins and art-known lyophilization and reconstitution techniques can be employed. It will be appreciated by those skilled in the art that lyophilization and reconstitution can load to varying degrees of antibody activity loss (e.g. with conventional immunoglobulins, IgM antibodies tend to have greater activity loss than IgG antibodies) and that use levels may have to be adjusted to compensate.

The compositions containing the present CDRgrafted antibodies or a cocktail thereof can be administered for prophylactic and/or therapeutic treatments. In certain therapeutic applications, an adequate amount to accomplish at least partial inhibition, suppression, modulation, killing, or some other measurable parameter, of a population of selected cells is defined as a "therapeutically-effective dose". Amounts needed to achieve this dosage will depend upon the severity of the disease and the general state of the patient's own immune system, but generally range from 0.005 to 5.0 mg of CDR-grafted antibody or binding protein thereof per kilogram of body weight, with doses of 0.05 to 2.0 mg/kg/dose being more commonly used. For prophylactic applications, compositions containing the present constructs or cocktails thereof may also be administered in similar or slightly lower dosages.

A composition containing a construct according to the present invention may be utilized in prophylactic and therapeutic settings to aid in the alteration, inactivation, killing or removal of a select T cell target population in a mammal.

In another embodiment, the constructs described herein may be used extracorporeally or in vitro selectively to kill, deplete or otherwise effectively remove the target cell population from a heterogenous collection of cells. Blood from the mammal may be combined extracorporeally with the CDR-grafted antibodies or binding proteins thereof whereby the undesired cells are killed or otherwise removed from the blood for return to the mammal in accordance with standard techniques.

In addition to the therapeutic uses, the constructs will find use in other settings. In one embodiment, the present antibodies or binding proteins are useful in an investigational setting. The T cell receptor binding proteins or antibodies of the invention are also useful in structural and functional studies regarding the T cell receptor. Besides serving as substrates or binding domains for specific forms of T cell receptors, these antibody constructs may serve as tools for the investigation of conformational studies to approximate the native configurations of various portions of the T cell receptor. In yet another embodiment, the T cell antibodies or binding proteins thereof can be used as diagnostic probes. In one specific embodiment, the constructs of the present invention may be used in determining the amount or the presence of a certain T cell V region subfamily in a biological sample. One such assay is described in Rittershaus - WO92/08981, published May 26, 1992 entitled "Therapeutic and Diagnostic Methods Using Total Leukocyte Surface Antigens". As such, the present invention provides for methods for diagnosing an immune related disease, such as MS, based on detecting the specific subset of T cell dntigen receptor in a biological sample.

The CDR-grafted antibodies or their binding proteins may be labelled in accordance with techniques known to the art. The constructs are also suitable for other in vlvo purposes. For example, they can be used for selective cell treatment of peripheral blood cells where it is desired to eliminate only target T lymphocytes or similarly in cell culture to eliminate unwanted T lymphocytes.

The following Examples are intended to more specifically describe certain aspects of the present invention but are not to be considered limitative thereof.

Example 1

TX27 - SYNOPSIS OF WORK

1. BALB/c mice were immunized with the T cell leukemia cell line HPB-All (once intraperitoneally and twice intravenously).

2. Spleen cells were harvested from these mice and fused with mouse myeloma immortalizing cell line P3XS3AG8.653.

3. Selection of appropriate clones was accomplished in HAT media.

4. Screening of the clones so selected for appropriate binding was accomplished by immunoprecipitation of radiolabelled HPB-ALL.

5. Subcloning was accomplished by limiting dilution.

6. Specificity was checked by PBL stimulation and CD3 modulation studies.

7. A Hybridoma secreting appropriate antibody was designated 4H11.

8. RNA was obtained from this clone and mRNA isolated. cDNA to the mRNA was prepared.

9. The cDNA was sequenced and checked for accuracy.

10. The isolated cDNAs (H and K) were cloned into M13 vectors.

11. Select human framework regions were introduced through mutagenesis, as exemplified in Example 2.

12. Humanized antibodies (-1 mg each) were purified from NS0 transfectants for preliminary evaluation.

Five different TM27's have been made:

TM27L: with Leucine at position 48 of heavy chain.

TM27I: with isoleucine at position 48 of heavy chain.

TM27.1: with FS to VF (78-79) changes.

TM27.2: with VTML/T to LSIS/N (67-70/73) changes.

TM27.3: with V to R (92) changes.

DETAILED EXPERIMENTAL DESCRIPTION

SUMMARY

The aim of this project was to humanize a murine anti-human TCR (T cell receptor) Vβ 5.2/5.3 monoclonal antibody (TM23) and to generate a humanized antibody expressing cell line suitable for process development. The humanized anti-TCR Vβ 5.2/5.3 monoclonal antibody is TM27.

Murine Antibody

The 4H11 hybridoma was developed by fusion of a mouse myeloma cell line, P3X63AGS.653, to spleen cells from Balb/c mice immunized with the T cell leukemia cell line HPB-ALL. The hybridoma expresses the monoclonal antibody TM23 with specificity to the human T cell receptor (TCR) Vβ 5.2 and 5.3. The isotype is IgG2a/kappa.

Humanization

The cDNA encoding the heavy and light chains of TM23 was isolated from the 4H11 cDNA library constructed using mouse IgG2a and kappa specific 3' end primers. The cDNA clones isolated were confirmed by DNA sequencing and internal amino acid sequencing. The CDRs (complementarity determining regions) of the heavy and light chains were identified according to the alignment of conserved mouse framework sequences. The L, V, (D), J (L, leader; V, variable; D, diversity; J, joining) DNA fragments were isolated using PCR and cloned into CHO cell expression vectors, pTCSLCxDHFR and pTCSLCg1NeoAp, which carried human kappa and IgG1 constant region cDNA, respectively, for mouse V/human C chimeric antibody expression.

Based on the comparison of amino acid sequences deduced from cDNA clones, the TM23 heavy chain was found to belong to Kabat subgroup IB. The V_K sequence was found to belong to Kabat V_KV. Therefore, the human NEWM and REI frameworks were chosen for humanization of heavy and light chain respectively. The rearranged V(D)J fragments (except part of 5'V and 3'J) were cloned, into M13 vectors which carried leader and intron sequences in genomic configuration, to generate complete genomic L-V(D)J constructs.

Myeloma Expression

After construction of CDR-grafted V(D)J regions by mutagenesis, the DNA fragments were cloned into myeloma expression vectors, which carried the constant region of human kappa and IgG1, respectively. Five humanized heavy chains with various mutations were constructed. Each of the five heavy chain constructs was co-transfected with a humanized light chain into NSO myeloma cell line. Humanized antibodies were purified from culture supernatants of the expanded human IgG1/κ positive transfectants for evaluation. The results indicated that TM27L, with the least mouse amino acid sequences, stained Vβ 5.3 T cells as well as the other humanized versions, if not better. TM27L was therefore chosen for further development and was designated as TM27.

CHO Cell Expression

The heavy and light chain leader and V regions of TM27 were isolated from myeloma constructs by PCR for cloning in cDNA configuration into CHO cell expression systems, pTCSLCg1NeoAp* (heavy) and pTCSLCκDHFR (light) as described above. The CHO cell transfectants were selected and human IgG1/κ positive cells were cloned. One of the highest producers, TM27L-662-35, was expanded for cell banking and small scale humanized Ab preparation. The CHO expressed TM27 antibody has been characterized throughout the progress of the project and is comparable with that from myeloma cell lines and the parental TM23. Concurrently, CHO cell clones were subjected to amplification and cloning to increase antibody expression and demonstrate suitability of the strain for productivity improvement.

DETAILED DESCRIPTION

Immunization of BALB/c Mice with EPB-ALL

The β F1 (an anti-human β chain T cell receptor constant region monoclonal antibody) immunoprecipitate of human T cell leukemia HPB-ALL cells (3×10^7) obtained from Dr. Michael Brenner (Harvard Medical School) was mixed with complete Freund's adjuvant and injected intraperitoneally (i.p.) into a BALB/c mouse. After 5 weeks, the mouse was boosted with 3×10^7cells in saline on 3 consecutive days, first i.p. and then twice intravenously (i.v.).

Fusion of Spleen Calls with P3X63AG8.653

The fusion was done by established techniques three days after the final i.v. boost. The spleen cells, in 5-fold excess, were fused with P3X63Ag8.653 cells (a mouse myeloma, non-Ig secreting line derived from BALB/c mouse, ATCC CRL 1580) using 50% polyethylene glycol 1500 (BDH, Dorset, England).

RAT Selection of Fused Cells

The fused cells were plated in 96-well plates with $2x10^5$ BALB/c thymocytes per well as feeders and selected in RPMI-1640 media containing 15% FBS (fetal bovine serum) and HAT (6 mM hypoxanthine, 50 mM aminopterin and 2 mM thymidine). The cells were incubated in humidified air with 5% CO_2 at 37 °C.

Screening of Hybridoma

After 13 - 14 days, the wells with growth were screened by both ELISA and immunoprecipitation of ^{125}I-radio-labeled HPB-ALL cells. Under the conditions of immunoprecipitation, CD3 molecules are dissociated from the TCR α and β chains. The hybridomas which reacted with HPB-ALL were expanded into T-75 flasks for further cloning.

Cloning of Hybridomas of Interest

Hybridomas with the specificity for HPB-ALL TCAR (T cell antigen receptor) were cloned by limiting dilution. Cells were plated out on 96 well culture plates with 2 fold serial dilutions of cell density ranging from 2 to 0.015 cell/well. The best growing colony with the highest productivity was chosen for repeating the above procedure twice. One of the final colonies after these two additional rounds of selection, 4H11, was identified. The mAb expressed by clone 4H11 immunoprecipated molecules identical to the TCR α and β chains precipitated with the β F1 mAb, which reacts with constant region of TCR β chain. The result suggested that this mAb had the specificity to TCR β chain expressed on the HPB-ALL T cells. The antibody was called TM23 for convenience.

Characterization of Monoclonal Antibody from Hybridoma 4H11, TM23

Isotyping of TM23 mAbs was performed using a commercially available kit (Zymed, South San Francisco, CA). The results indicated that it is a mouse IgG2a/κ isotype.

To confirm the anti-TCR reactivity of the TM23 mAb, it was screened for its ability to comodulate the CD3 molecule from the surface of HPB-ALL cells. Previous studies have shown that mAbs to either the a or b chain of TCR can comodulate CD3 from the cell surface when cells are preincubated with the anti-TCR mAbs. When HPB-ALL cells were preincubated with TM23, a significant decrease in reactivity with the anti-CD3 mAb Leu-4 was seen when compared to the Leu-4 reactivity seen in untreated cells.

Normal human PBL (peripheral blood lymphocyte) was tested for reactivity with TM23. From 1.6% to 3.0% of the PBL T cells studied reacted with TM23. This result is typical of results obtained with several other mAbs which have been shown to react with specific V region determinants.

To determine if TM23 binds to the same epitope as 1C1, a mAb which reacts with human Vβ 5.2 and 5.3 subfamilies (Boylston, et al., 1986 J. Immunol. 137:741), the ability of one mAb to block the binding of the other was studied. When HPB-ALL cells were first incubated with TM23, the binding of fluorescein-conjugated 1C1 was inhibited. Also, when cells were incubated first with 1C1, the binding to TM23 was inhibited. This result suggests that TM23 and 1C1 bind to the same or closely adjacent epitopes.

To further assess whether TM23 and 1c1 have similar reactivities, PBL lines were derived by stimulating with the other mAb. Each line was then examined for reactivity with each mAb. When a PBL line was established by stimulation with TM23, both 1C1 and 4H11 reacted with a similar percentage of cells. The same is true of a 1C1 stimulated PBL line.

Growing of 4H11 Cell Culture

The 4H11 clone TM23.1 7-31-91 was grown and expanded in DMEM/F12 media (1:1) plus 1% fetal calf serum, 100 m/ml Penicillin, 100 m/ml Streptomycin and 12.5 mM glutamine.

Preparation of cDNA Encoding TH23 Heavy and Light Chains

About 10^8 4H11 cells were isolated by centrifugation and washed once with ice cold PBS. Total RNA was prepared using Promega RNAgents Total RNA Isolation Kit (Promega, Cat # Z5110). Poly $(A)^+$ RNA was prepared using Promega PolyATract mRNA Isolation System (Promega Cat # Z5200). The mouse IgG2a and κ specific cDNA library was prepared using the BRL SuperScript Kit (BRL Cat # 82YBSA) with the IgG2a primer

5' ATATGCSGCCGCTCATTTACCCGGAGTCCGGGAGAAGCTCTTAGT 3'

and κ primer

5' ATATGCGGCCGCTTAACACTCATTCCTGTTGAAGCTCTTGACAAT 3'.

The two primers contain Xho I restriction enzyme sites and sequences complementary to the 3' ends of mouse IgG2a and κ coding regions, respectively. The cDNA was cloned into plasmid pSPORT, supplied in the Kit, and transformed into *E. coli* DH5α competent cells by electroporation. Plasmid clones carrying the right size inserts were selected for sequencing.

Identification of CDNA Clones Encoding TM23 Heavy and Light chains

The mini-prep plasmid DNA with inserts of the right size (light chain clones # 1, 6, 19, 20, 25 and 26; heavy chain clones # 4, 9, 11, 15, 22 and 34) were subjected to double stranded DNA sequencing using the Sequencing Kit (USB Cat # 70770) which has a T7 primer 5' to the DNA inserts. The sequences of light chain clones # 6, 20, and 25 and heavy chain clone # 34 indicated that they were mouse Ig clones. Light chain clones # 6, 20 and 25 had identical sequences and # 6 had a point mutation and was a nonfunctional clone. Clones # 20 and 34 were sequenced using κ

5' GACATTGATGTCTTTGGGGTAGAAGTTGTT 3'

and IgG2a

5' GGTCACTGGCTCAGGGAAATAACCCTTGAC 3'

constant region specific primers. Again, the results indicated that clone # 6 was a non-functional clone and clones # 20 and 34 were the right clones for light and heavy chain respectively. The cDNA and amino acid sequences deduced from the two clones are shown in Fig. 1.

Due to N-terminal blocking, micro-amino acid sequencing was performed from CNBr cleaved peptide fragments isolated from polyacrylamide gels. The internal V region amino acid sequences obtained from this process were consistent with that deduced from cDNA sequences. The results confirmed that the cDNA clones isolated were the ones encoding TM23 heavy and light chain.

The CDRs of the heavy and light chains were identified according to the method of Kabat and Wu (supra) and are highlighted and underlined in Figure 1.

PCR Isolation of V Region DNA of H and L Chains and DNA sequencing

PRODUCTION OF CHIMERIC TM23 ANTIBODY

Isolation of Murine VH DNA and NK DNA

The TM23 heavy and light chain variable regions were adapted for insertion into expression vectors. The DNA sequence coding for the TM23 murine VH was amplified from TM23MuVH.MuIgG2a by PCR using oligonucleotide primers VH1BACK
(5' AGGTSMARCTGCAGSAGTCWGG 3')
which contains a site for the restriction enzyme PstI and VH1FOR
5' TGAGGAGACGGTGACCGTGGTCCCTTGGCCCCAG 3'
which contains likewise a BstEII site. M stands for C or A, S stands for C or G, R stands for A or G and W stands for A or T. These sites were thereby introduced at the 5' and 3' ends respectively of TM23VH DNA. The amplified DNA was cut with PstI and BstEII and ligated into the PstI and BstEII sites of double-stranded replicative (RF) form DNA of the vector M13 VHPCR1, cut with the same enzymes.

M13 VHPCRI is an M13 phage vector derived from M13-HuVHNP (Jones et al., (1986) Nature 321:522-525) and contains an immunoglobulin promoter, signal sequence and appropriate splice sites (Orlandi et al., (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837).

The ligated DNA was transformed into competent *E.coli* strain TG1 (K12, Δ(lac-pro), supE, thi, hsd D5 [F' tra D36, pro^{A+B+}, $lacO^{3}$, lacZ M15]. Single-stranded DNA was prepared from the recombinant phage plaques obtained.

The DNA sequence coding for the TM23 murine VK was amplified from TM23MuVK.MuCK by PCR using oligonucleotide primers VK1BACK
5' GACATCCAGCTGACCCAGTCTCCA 3'
which contains a PvuII site and VK1FOR
5' GTTAGATCTCCAGCTTGGTCCC 3'
which contains a BglII site in conjunction with two overlapping internal primers designed to remove the HindIII site in CDR3, oligo 943 5' CCGAGGAAGTTTACTATACTG 3',
and oligo 944 5' CAGTATAGTAAACTTCCTCGG 3'

(Horton et al, (1990). PvuII and BglII sites were thereby introduced at the 5' and 3' ends of TM23 VK DNA. The amplified DNA was digested with PvuII and BglII and ligated in PvuII and BclI cut M13 VKPCR1. The enzymes BglII and BclI cut DNA to produce compatible cohesive ends which can be ligated together.

M13VKPCR1 contains the same promoter, signal sequence and splice sites as M13 VHPCR1 (Orlandi et al, (1989).

The ligated DNA was transformed into *E-coli* TG1 and single-stranded DNA prepared from the recombinant phage plaques obtained.

PCR products of the correct size, 359 base pairs (bp) for VH and 288 bp for VK were obtained. The VH DNA was inserted into the vector M13VHPCR1 to give M13 TM23MuVH. The VK DNA was inserted into the vector M13VKPCR1 to give M13 TM23MuVK.

Confirmation of DNA Sequences

It was confirmed that the correct DNA sequences had been cloned and that no spurious mutations had been introduced by the PCR. The sequence of single-stranded M13 phage DNA of the entire 823 base pair (bp) VH region including the promoter, signal sequence, splice sites and 5' and 3' untranslated sequences was obtained by the dideoxy method (Sanger et al, 1977) using T7 DNA polymerase or Sequenase.

The sequence of single-stranded M13 phage DNA of the entire 630 base pair (bp) VK region including the signal sequence, splice sites and 5' and 3' untranslated sequences was obtained by the dideoxy method, as above.

DNA clones with the desired sequence for TM23 VH adapted for insertion in the expression vector were identified. One of these, #2, was selected for further work. The introduction of the requisite restriction sites necessary for insertion of VH into the expression vector resulted in 1 amino acid change at the N-terminus; lysine (K) at.position 5 to glutamine (G) and one change at the C-terminus, serine (S) at position 108 to threonine (T).

DNA clones with the desired sequence for TM23 VK adapted for insertion in the expression vector were identified. One of these, #4, was selected for further work. The amino acid squence of CDR3 was not altered by the removal of the Hind III site, but the introduction of the requisite restriction sites necessary for insertion of VK into the expression vector resulted in three amino acid changes at the N-terminus; methionine (M) at position 4 to leucine (L),

threonine (T) at position 7 to serine (S) and threonine (T) at position 8 to proline (P), and one change at the C-terminus, valine (V) at position 105 to glutamine (E), due to the primers used.

Cloning of V region DNA Fragments into M13 Vector

Transfer of the cloned murine variable region to expression vectors containing human constant region genes for subsequent expression of the antibody in mammalian cells was accomplished.

Selection of IgG Isotype for Humanization

Human IgG1 isotype was chosen for TM27 for the following reasons:

1. IgGl has the capability of mediating both ADCC (antibody dependent cell cytotoxicity) and CDC (complement dependent cytotoxicity). Therefore, it may be more effective in down-regulating Vb 5.2/5.3 T cells; and
2. IgG1 has been used more often in Ab humanization.

Human Framework Selection for CDR Graft

PRODUCTION OF HUMANISED TH23 ANTIBODY

Design of Humanised TH23 Variable Region Sequences

Selection of human framework sequences to receive the TM23 CDRs and determination of essential murine residues that must be substituted in the human framework was carried out. Computer assisted comparisons of the amino acid sequences of TM23 VH and VK were made both with the known sequences of other murine antibodies and human antibody VH and VK sequences.

The program ALLIGN from DNASTAR Ltd, London W13 9BL, UK, was used, run on a Compaq 386 processor. AALIGN aligns two proteins sequences using the Lipman-Pearson and Needleman-Wunsch methods. First, the Lipman-Pearson method determines the best regions of homology. Then the Needleman-Wunsch method provides the final alignment of the sequences.

The alignments produced were then examined paying particular attention to factors such as the length of the CDRs and residues believed to be critical in supporting the conformation of

the CDRS, for example positions 71 and 94 and residues close to the CDRs (Tramontano et al, (1990); Foote and Winter, (1992)).

Comparison with consensus sequences for the Kabat subgroups assigned the TM23 murine VH to Kabat subgroup VK V. There are no amino acid sequences which are unsual for these subgroups in the TM23 frameworks.

The human antibodies NEWM and REI provided the framework sequences for VH and VK respectively. For VH, in addition to the CDRS, the murine residues at 27 to 30 inclusive and 71 were used. These framework residues are thought to be important in supporting the conformation of the CDRs (Foote and Winter, (1992)). Residue 71 may fix the relative dispositions of CDR1 and CDR2, according to whether there is a bulky side chain (lysine or arginine) or a smaller side chain (valine or alanine) present (Tramontano et al., (1990). Therefore valine 71 in NEWM VH was replaced with lysine as in TM23 VH. Residues 27 to 30 form part of the structural, rather than hypervariable, antigen-binding loop as defined by Chothia and Lesk (1987). Framework residues 47 to 49 may also be important for antibody structure (Foote and Winter, (1992). Therefore two alternative humanised heavy chains were constructed: one with isoleucine (I) at position 48 as in NEWM; and one with leucine (L) at position 48 so that residues 47 to 49 from the murine VH were preserved. A comparison of the murine VH sequence with the humanised version is shown in Example 2.

No alterations were made to the human REI framework chosen for TM23 VK. A comparison of the murine VK sequence with the humanised version is also shown in Example 2.

Description of mamalian Expression Vector pTCSNeo

The plasmid pTC5Neo contains a unique Xho I cloning site into which genes of interest are inserted. Transcription of the inserted gene is driven by the regulatory sequences consisting of a SV40 enhancer upstream of the adenovirus 2 major late promoter (Ad2 MLP). The Ad2 MLP is itself located upstream of the adenovirus tripartite leader. The inserted gene is flanked by a polyadenylation signal from murine immunoglobulin kappa (Igκ) at the 31 end. Mammalian selection is conferred by the neomycin resistance gene (Neo^r) . The plasmid also provides the bacterial origin of replication and b-lactamase gene (Amp^r).

Construction of Expression Vector pTCSLNeo

Two complementary oligonucleotides with the sequences: anti-sense,

5' CGACATCGATGATATCGAATTCGCGGCCGCCAGCTGGGGCCCTCTAGAC 3'

and sense,

5' CGAGTCTAGAGGGCCCCAGCTGGCGGCCGCGAATTCGATATCATCGATG 3'

were reannealed and ligated to pTCSNeo which had been digested with XhoI and filled in with dT. The resulting plasmid, pTCSLNeo, with the polylinker consisting of multiple restriction enzymatic sites in the orientation of

5' XhoI / XbaI / ApaI / PvuII / NotI / EcoRI / EcoRV / ClaI 3',

is convenient for inserting the gene of interest for expression. The neo^r gene can be changed to other selection marker genes as well.

Construction of Expression Vector pTCSLDHPR-

The neo^r gene in pTCSLNeo was removed by digestion with the restriction enzyme HindIII and BamHI and religated with the murine mutant DHFR (dihydofolate reductase) (DHFR-) gene fragment generated by HindIII and BamHI digestion of plasmid pSV2-DHFR- obtained from Dr. Paul Berg (Department of Biochemistry, Stanford University Medical School).

Description of Plasmid pSV184 H-Neo/DNS-VκCκ

The plasmid pSV184 ΔH-Neo/DNS-VκCκ (obtained from Dr. Lee Herzenberg, Stanford University Medical School - see Thesis of Jeffery Dangl, page 48-67) is a mouse V and human C chimeric construct in which the rearranged κ chain V-J region gene segment isolated from a Dansyl Chloride specific murine hybridoma was fused to a human genomic Cκ fragment (pSV184ΔH-Neo-Cκ.

Isolation of the human kappa chain constant region by the PCR method.

Using the undigested plasmid pSV184ΔH Neo/DNS-VκCκ (pSVHuk) which contains the human kappa chain constant region, PCR was performed with two primers (one forward, one reverse) to isolate the constant region. The oligomers (primers) were obtained from operon. The PCR reaction reagents were from the GeneAmp DNA amplification reagent kit with AmpliTaq (Perkin-Elmer Cetus) The PCR reactions were set up in three different concentrations

of Mg^{2+}. Aliquots of the PCR products were analyzed on a 1% agarose gel. The PCR product from the first sample of Mg^{2+} (concentration 1.5 mM) was chosen. The primers, which flank the C region, were designed to contain restriction enzyme sites for cloning purposes. Thus, the amplified DNA was digested with EcoRI (at the 3' end). The site at the 5' end (PvuII) was not digested. The digested DNA fragment was then run on a 1% agarose gel and the 300 bp band excised. The DNA was extracted from the gel and a small aliquot was analyzed on a 1% agarose gel to determine quality and to estimate quantity. The 300 bp band appeared to be clean and the total amount of DNA estimated to be ~ 140 ng.

Human kappa chain constant region cloned into the pBS $KS^{\pm}$ vector and sequenced.

The EcoRI-digested DNA fragment of the human kappa chain constant region was inserted into the vector pBS KS^{+}, which had been digested with Sma I and EcoR I, using T4 DNA ligase. This ligation mixture was used to transform *E. coli* DH5α competent cells. Single colonies were picked and used to inoculate 24 liquid cultures. DNA was prepared from each of the cultures. Each DNA sample was digested with Pvu II (5' end of Cκ) and EcoR I (3' end of Cκ), and analyzed on a 2% agarose gel. Due to difficulty in separating the lower range bands (300 bp), the DNA samples were digested with PvuII only. After analysis on a 2% agarose gel, 19 out of 24 were correct for C_k insert. Samples #1 - 5 (of the 19 correct samples) were chosen to be sequenced, using T7 and T3 primers (sequences in the vector flanking the insert) . Other sequencing reagents were from the Sequenase Version 2.0 kit (USB). Sample #5 proved to have correct sequence for the human kappa chain constant region.

Human kappa chain constant region cloned into the mammalian expression vector pTCSLDRPR* to become pTCSLCκDRFR*.

The human kappa chain constant region was isolated from the pBS KS vector (Sample #5) using PvuII and EcoRI restriction enzymes. However, due to the previous problem of band separation, a second digest was performed with the construct pBS/Human Cκ. The second digest involved EcoRI and XbaI (located 5' to the PvuII site) yielding a fragment ~ 300 bp. This fragment was then gel purified and subjected to a PvuII digest, to remove the remaining small fragment of vector at the 5' end of the Cκ. This final fragment was gel purified and then analyzed on a 1% agarose gel for quality and to estimate quantity. The Cκ DNA appeared to be a clean band at ~ 280 bp and the total quantity estimated to be ~ 135 ng.

Due to the existence of three EcoRI sites in the expression vector, pTCSLDHFR*, it was determined that cloning into the PvuII site only would be best. Since the Cκ DNA was prepared by digestion with PvuII - ECORI, the EcoRI end was repaired (to form blunt end) using Klenow fragment. The repaired, blunt-ended Cκ DNA was ligated to the pTCSLDHFR* vector which had been PvuII digested and phosphatase-treated (using calf intestinal phosphatase). This ligation mixture was used to transform *E. coli* DH5α competent cells. single colonies were picked and used to inoculate 24 liquid cultures. DNA was prepared from each. All DNA samples were digested with PvuII and HindIII, and analyzed on a it agarose gel. It was determined that five samples, out of 24, contained the Cκ insert in the correct orientation. Sample #7 was chosen for continuation.

Replacement of the mutant DHPR* gene with the wild type DEPR gene.

For the expression of the vector pTCSLCκDHFR* in CHO(dhfr⁻)DUX B11 cells, the DHFR* gene was replaced with the DHFR gene. DNA from pTCSLCκDHFR* sample #7 as well as pSV2-DHFR (plasmid obtained from Dr. Paul Berg, Department of Biochemistry, Stanford University Medical School) was digested with restriction enzymes HindIII and BglII to isolate the respective gene fragments. Upon analysis, it was determined that an additional BglII site in the pTCSLCκDHFR* vector removed a portion of the polyA site also, which would interfere with the vector function. Therefore, the digests were repeated using HindIII and BamHI. The (DHFR*)pTCSLCκ portion of the vector was treated with phosphatase. Then both the isolated fragments (pTCSLCκ and DHFR) were gel purified and analyzed on a 1% agarose gel for quality and to estimate quantity. Both bands appeared to be clean and estimated to be of similar concentration, ~ 40-50 ng/ml. The DHFR gene was ligated to the pTCSLCκ vector using T4 DNA ligase. The ligation mixture was used to transform *E. coli* DH5α competent cells. Single colonies were picked and used to inoculate 24 liquid cultures. DNA was prepared from each. The DNA was analyzed by restriction enzyme digestion with HindIII and BamHI.

From 24 cultures, 23 had the correct bands indicating the insertion of the DHFR gene into the pTCSLCκ vector. Sample #22 was selected for further use.

in vitro Mutagenesis to Construct CDR Grafted V Regions with Human Framework

Construction of Humanized Variable Genes

A humanized VH gene was constructed by site-directed mutagenesis of M13 VHPCR1 template, which comprises a synthetic VH gene with the framework regions of the human myeloma protein NEWM (see Jones et al, 1986). single stranded template DNA was prepared from M13 VHPCR1 grown in a dut$^-$, ung$^-$ *E.coli* strain RZ1032 so that the template DNA contained uracil in place of thymine. Mutagenic oligonucleotide primers were synthesised to encode the TM23 CDRs and approximately 15 bp of perfectly matched flanking sequence on either side. If the TM23 CDR sequence matched the template sequence shorter oligonucleotide primers were used. The primers used were

CDR1

5' TGGCTGTCTCACCCAGTTTACACCATAGGCGGTTAATGAGAAGCCAGACACGG 3',

CDR2

5'

GCTGGTGTCCTTTCAGCATTGTCACTCTGGATTTGAGAGCTGAATTATAGTCTGTA-

TTTCCATCACCCCATATCATTCCAAT1 CCACTCAAGACC 3',

and CDR3

5' CCAGTAGTCCATAGCATAGAGGGTAGCCGTAACCCTATCTCTTGCACAATAATAG 3'.

The oligonucleotide primer for CDR was extended at the 3' end to include the required residues at positions 27 to 30. A mixed oligonucleotide was made for CDR2, extended at the 3' end of the CDR with either G or T at base number 402 of the VH providing 2 alternative codons such that position 48 of the VH became either L or I. The oligonucleotides (10 pmol of each) were phosphorylated with 5 units of T4 polynucleotide kinase in a total volume of 50 µl at 37 °C for 1 hour. 1 pmol of each mutagenic primer and 1 pmol of the M13 sequencing primer 3' to the V gene were annealed to 0.5 µg of template in a total volume of 20 µl by heating to 90 °C for 30 seconds, rapidly cooling to 70 °C and slowly to 37 °C. The annealed primers were then extended and ligated with the addition of T7 DNA polymerase and T4 DNA ligase, incubated at room temperature for 1 hour. The uracil-containing template DNA was removed by treating the DNA with 1 unit of uracil DNA glycosylase at 37 °C for 1 hour, leaving the mutated strand intact. This DNA was amplified by PCR then analysed by agarose gel electrophoresis to check

that DNA of the correct size had been produced. The DNA was cut with HindIII and BamHI and ligated into the HindIII and BamHI sites of M13mp19.

Similarly, the reshaped TM23 VK gene (TM23HuVK) was constructed by mutagenesis of an M13 phage template which comprises the DNA sequence of a VK gene with the framework regions of the human Bence-Jones protein REI. This was derived from M13 VKPCR1 which was used in the construction of the chimeric light chain, with the PvuII and BclI restriction enzyme sites used for cloning removed. The primers used were

CDR1

5' TCTGCTTGGTACCAGTTTAAATAATTGCTAATGCCCTGACTTGCACTACAGGTGATGGTC 3',

CDR2

5' TCTGCTTGGCACACCTGAGTGTAAACTTGATGTGTAGTAGATCAGCAGCTT 3',

and CDR3

5' CCCTTGGCCGAACGTCCGAGGAAGTTTACTATACTGCTGGCAGTAGTAG 3'.

PCR products of the expected size, 823 bp for both VHs and 620 bp for VK were obtained and cloned into the M14mp19 vector.

Verification of Humanized Variable-Region DNA Sequences

It was confirmed that the required CDRs had been inserted, that no spurious changes had occurred and that the correct humanized VH and VK genes had been constructed. The complete variable regions were sequenced by the dideoxy method using T7 DNA polymerase or Sequenase. The sequences obtained were compared to the designed sequences. Clones giving the expected sequence for the two VHs were selected. These clones were named M13 TM23NMVH and M13 TM23NMVHL. A clone giving the expected sequence for the reshaped VK, #11, was selected and named M13 TM23HuVK.

Cloning of Rearranged V Regions into Myeloma Expression Vectors

(Cloning of Humanized TM23 Variable Regions into Expression Vectors)

The cloned humanized variable regions were transferred to expression vectors containing human constant region genes for subsequent expression of the antibody in mammlian cells. The two humanized VH genes were isolated from M13 TM23NMVH and M13 TM23NMVHL as HindIII to BamHl fragments and ligated into the HindIII and BamHI sites of the heavy chain expression vector, pSVgpt.HuIgG1.

The humanized VK gene was isolated from M13 TM23HuVX as a HindIII to BamHI fragment and ligated into the HindIII and BamHI sites of the light chain expression vector pSVhvg.HuCK.

Restriction enzyme analysis showed that the two alternative humanized VHs were correctly inserted into the heavy chain vector to give the plasmids pSVgptTM23NMVH.HuIgG1 and pSVgptNMVHL.HuIgG1.

Likewise, the humanized VK was correctly inserted into the light chain vector to give the plasmid pSVhygTM23HuVK.HuCK.

Co-transfection of H and L Chains Constructs into NS0 Cells

Cotransfection of Humanized Heavy and Light Chain Genes

Humanized expression plasmids were transferred to the mammalian cell lines, NS0 and P3-8.653F8879.

The humanized expression plasmids were cotransfected into NS0 and P3-8.653F8879 cells using the same procedure as for the chimeric expression plasmids. The two humanized heavy chain plasmids, pSVgptTM23NMVH.HuIgG1 and pSVgptNMVHL.HuIgG1 were each cotransfected with the humanized light chain plasmid, pSVhygTM23HuVK.HuCK to produce cell lines expressing two alternative humanized antibodies, TM23HuVH/HuVK and TM23HuVHL/HuVK. These expression plasmids were also cotransfected in combination with the chimeric expression plasmids to produce cell lines expressing the hybrid antibodies TM23HuVH/MuVK, TM23HuVHL/MuVK and TM23MuVH/HuVK. These hybrid antibodies provide important tools to locate the cause of any binding defect to either the heavy chain or the light chain in the humanized antibody.

Fourteen days after the transfection colonies of NS0 cells were visible for all the various plasmid combinations. Only one single colony of P3-8.653F8879 cells was obtained from all the plates seeded and these were therefore discarded.

Selection of Human IgG1/kappa Positive Clones

Selection for Cells Producing Human Antibody

NS0 cell lines producing humanized and hybrid chimeric/humanized antibodies were isolated. Samples of supernatant from wells containing cell colonies were screened by ELISA.

TM23HuVH/HuVK:-NSO cells from 4 wells were selected in which the culture supernatant gave the highest readings in the ELISA; D5, D10, C6 and E11. Well D5 contained 2 visible colonies, the other wells one. However, the cell lines produced from these are not necessarily clonal. Cell lines from these wells were cultured and frozen in liquid nitrogen as TM23HuVH/HuVK D5, D10, C6 and E11.

TM23HuVHL/HuVK:- NSO cells from 4 wells were selected in which the culture supernatant gave the highest readings in the ELISA; B2, B7, D8 and E9. Well E9 contained 2 visible colonies, the other wells one. However, the cell lines produced from these are not necessarily clonal. Cell lines from these wells were cultured and frozen in liquid nitrogen as TM23HuVHL/HuVK B2, B7, D8 and E9.

TM23HuVH/MuVK:- NSO cells from 4 wells were selected in which the culture supernatant gave the highest readings in the ELISA; D9, E6, F10 and G10. Wells F19 and G10 contained 2 visible colonies, the other wells one. However, the cell lines produced from these are not necessarily clonal. Cell lines from these wells were cultured and frozen in liquid nitrogen as TM23HuVH/MuVK D9, E6, F10 and G10.

TM23HuVHL/MuVKL:- NSO cells from 4 wells were selected in which the culture supernatant gave the highest readings in the ELISA; B8, C6, D2 and D8. Wells C6 and D8 contained 2 visible colonies, the other wells one. However, the cell lines produced from these are not necessarily clonal. Cell lines from these wells were cultured and frozen in liquid nitrogen as TM23HuVHL/MuVK B8, C6, D2 and D8.

TM23MuVH/HuVK:- NSO cells from 4 wells were selected in which the culture supernatant gave the highest readings in the ELISA; B9, G4, G6 and G11. All wells contained at least 4 visible colonies. The cell lines produced from these are not necessarily clonal. Cell lines from these wells were cultured and frozen in liquid nitrogen as TM23MuVH/HuVK B9, G4, G6 and G11.

First Strand cDNA Synthesis for CHO Cell Expression

Preparation of Cytoplasmic RNA

RNA was prepared for reverse transcription. A vial of NS0 TM23 HuVHL/HuVK cells was recovered from liquid nitrogen by thawing rapidly at 37 °C. The cells were gradually diluted with 10 ml of Dulbecco's Modified Eagle's Medium/Ham's F12 (DMEM/F12) supplemented with antibiotics and 5% gamma globulin free fetal bovine serum (GG-free FBS). The cells were centrifuged at 1500 rpm for 5 minutes, the medium removed and resuspended in 20 ml of fresh medium in a 75 cm^3 flask. The culture was expanded to 30 ml in a 175 cm^3 flask. Actively growing cells were harvested by centrifugation and washed 3 times with cold phosphate buffered saline (PBS). Cytoplasmic RNA was prepared from the cells by the method of Favoloro et al, (1980). A total of 350 mg of cytoplasmic RNA was obtained.

cDNA Synthesis

First strand cDNA for was prepared VH and VK regions. First strand variable region cDNAs were synthesised in reactions containing 5 μg of RNA, 250 μM deoxynucleotide triphosphates (dNTPs), 15 units of ribonuclease inhibitor (Pharmacia RNAguard) and 25 pM of oligonucleotide primer. The primers were oligo 1109 for VH

5' GGGGAAGACCGATGGGCCCTTGGTGGAGGCTGAGGAGACGGTGACC 3' and oligo 1095 for VK

5' AGATTTCAGCTGCTCATCAGATGGCGGGAAGATGAAGACAGATGG 3'.

The reactions were heated to 70 °C for 10 minutes then coolled slowly to 37 °C. 100 units of Moloney Murine Leukemia Virus Reverse Transcriptase was added and the reactions incubated at 37 °C for 30 minutes.

The cDNAs were amplified by PCR using the primers above in conjunction with oligo 1096 for VH

5' GCCGCTCGAGCCTCACCATGGGATGGAGCTGTATCATCCTCTTCTTGGTAG 3' and oligo 1094 for VK

5' GCCGCTCGAGCCTCACCATGGGATGGAGCTGTATCATCCTCTTCTTGGTAGCAAC-AGCTACAGGTGTCCACTCCGACATCCAGATGACCCAGAG 3'.

The PCR products were checked by agarose gel electrophoresis and were of the expected size (approximately 450 base pairs) and the DNA bands were purified from the gel.

Restriction Enzyme Digestions

The PCR products were digested for cloning into CHO vectors. The VH PCR product was digested with XhoI and ApaI. The VK PCR product was digested with XhoI and PvuII. Samples of the digested DNAs were electrophoresed on a 0.8% agarose gel. Bands of the expected size, 450 base pairs, were seen.

Isolate V regions (V_H [TCS and S] and $V\kappa$ [S]) from cDNA by PCR amplification V_H [TCS]

First strand cDNA samples of TM27 Vκ and TM27 V_H, as well as oligonucleotide primers used for amplification of these molecules by PCR, were obtained. PCR using the TM27 V_H cDNA with appropriate primers was performed using Taq polymerase. The product was run on an agarose gel and purified using the Geneclean II kit (Bio101) . Recovery was estimated at 1.5 ng/ml.

Transfection of COS Cells with pSR1Neo4H11 (light) and pBJ14H11 (Heavy) for C Region cDNA Isolation

The mammalian expression vector pSR1Neo and PBJ1 are derivatives of pcDL-SRa296 (Takebe, et al., (1988) Mol. Call. Biol. 8:466) as described in Lin, et al., (1990) Science 249:677 in which a XhoI site located between the SRa promoter and the SV40 polyadenylation site has been replaced for convenient gene cloning by a polylinker that contains the following restriction sites: 5' XhoI / XbaI / SfiI / NotI / EcoRI / EcoRV / HindIII / ClaI 3'. In the pSR1Neo, a neomycin resistance gene was inserted between the ampicillin resistance gene and the SRa promoter. The cDNA encoding leader and V-D-J region of TM23 heavy chain was first fused to the genomic heavy chain constant region followed by insertion into the XhoI and NotI cut pBJ1 vector to generate pBJ14H11 (heavy). Similarly, the light chain cDNA encoding leader and V-J was first fused to genomic κ fragment followed by insertion into XhoI / NotI cut pSR1Neo to generate pSR1Neo4H11 (light). The two plasrnids, pBJ14H11 (heavy) and pSR1Neo4H11 (light) were then co-transfected into COS-7 cells (a SV40 transformed African green monkey kidney cell line, ATCC CRL 1651) for transient expression.

Isolation of Human IgG1 Constant Region

Human IgG1 constant region cDNA was prepared from poly(A)$^+$ RNA isolated from transfected COS cells by PCR using IgG1 constant region primers: forward primer 5' GCGTGACAAGGGCCCATCGGTCTTCCCCCTGGCACCCTC 3'
and reverse primer
5' GCGTGACAAGAATTCTCATTTACCCGGAGACAGGGAGAGGCTCTT 3'.
The PCR constant region DNA fragment with EcoRI at 3' end (flanking the coding region) and ApaI at the 5' end (in the coding region) was cloned into pBS KS$^+$ plasmid for subcloning and sequencing. The entire constant region was sequenced using T3 and T7 primers, complemented with constant region internal primers. The clone with the correct sequence was then subcloned into plasmid pTCSLNeo to generate pTCSLCg1Neo.

Construction of Plasmid pTCSLCg1Neo

The IgG1Cg1 clone with the correct sequence in pBS KS$^+$ was cut out using the restriction enzymes EcoRI and ApaI and ligated into pTCSNeo digested with the same enzymes. The resulting plasmid, pTCSLCg1Neo, was used for further modification to generate pTCSLCgNeoApa$^-$.

Elimination of ApaI site in pTCSLCκ1Neo to generate pTCSLCgNeoApa$^-$

The pTCSLCκ1Neo CHO expression vector contains two sites for the restriction enzyme ApaI: one site located within the polylinker region used for insertion of cloned fragments; the other site located just 3' to the neo^r gene. In order to simplify subsequent cloning steps, the ApaI site near neo^r was eliminated. pTCSLC$_k$1Neo was partially digested with ApaI; a small aliquot was run on an agarose gel to check for presence of linear molecules (i.e. plasmids cut at only one of the two ApaI sites. The digested material was then treated with T4 DNA polymerase, which selectively removes the single-strand extensions left by ApaI without digesting double-strand DNA. Plasmids were then recircularized by ligation and transformed into HB101 competent *E. coli* cells. Thirty transformants were picked and used to inoculate liquid cultures. DNA was prepared from each culture, digested with ApaI and run on agarose gels. Unfortunately, plasmid DNA was degraded in these digests. ApaI digests were repeated on 21 of the 30 samples. Two isolates (# 4 and 12) appeared to be linearized by ApaI while all others were cut at two sites. The two candidate clones were then digested with ApaI and HindIII

to determine which ApaI site was present. In this test, the ApaI site 3' to neo^r in pTCSLCg1NeoApa• isolate #12 was eliminated, as desired. Additional restriction digests were performed which confirmed the structure of this isolate.

Cloning the PcR isolated human kappa and heavy chain V regions into the CHO Vectors, pTCSLCκDHFR and pTCSLC$_g$1NeoApa•, Respectively.

Clone the TH27 Vκ fragment into pTCSLCκDHPR

The cDNA for the kappa chain V region for TM27 had been gel purified and digested with the appropriate restriction enzymes. A small aliquot of the DNA sample was analyzed on a 1% agarose gel to estimate quantity and to verify the quality. An extra (contaminating) band was observed at ~ 800 bp the concentration was estimated at 1.25 ng/ml. The V region was ligated to the pTCSLCκDHFR vector which had been digested with restriction enzymes XhoI and PvuII and phosphatase-treated. The ligation mixture was used to transform *E. coli* DH5α competent cells. Yield of transformant colonies was very low. Single colonies were picked and used to inoculate twelve liquid cultures. DNA was prepared from each, digested with XhoI and PvuII and analyzed on a it agarose gel. There were two positive samples, #3 and #4. Each of the positive samples was sequenced.

Clone TH27 V_H fragment into pTCSLC$_g$1NeoApa•

The PCR product corresponding to the V_H region of TM27 was gel-purified and digested with the appropriate restriction enzymes (XhoI and ApaI). However, since it had been noted that the ApaI digestion may not have been complete, this digestion was repeated. A small aliquot was analyzed on an agarose gel to estimate quantity and to verify the quality. The concentration of the band corresponding to the V_H region was estimated at 0.4 ng/ml. As in the case of the Vκ fragment, an extra band was observed at ~ 800 bp, suggesting that the purity of this material was also not as expected. The V_H region was ligated to the pTCSLC$_g$1NeoApa• vector, which had been digested with the restriction enzymes XhoI and ApaI and phosphatase-treated. The ligation mixture was used to transform *E. coli* DH5α competent cells. Very few transformants were recovered. Six colonies were picked and used to inoculate liquid cultures. Miniprep DNA was prepared from each, digested with XhoI and EcoRI, and analyzed on agarose gels. One candidate clone was recovered from the first ligation, pTCSLHuV_H4HC$_g$1NeoApa• (isolate #6). In order to obtain more candidate clones (in case the

single isolate #6 failed to contain the correct sequence in subsequent tests) additional ligations were set up using 1) the V_H PCR product, repurified on low melting point agarose to remove the high-molecular-weight contaminating band and 2) a V_H PCR product synthesized using cDNA and primers. From these ligation reactions, eight additional transformants were analyzed and five candidate clones recovered: isolates #9, 10, 11, 12, and 14.

Sequence cloned V regions on both strands to confirm expected DNA sequence

Primers designed to facilitate DNA sequencing analysis of V regions inserted in the pTCSLC$_g$1NeoApa$^-$ and pTCSLCκDHFR vectors were obtained from Operon Research. The forward primer was located within pTCS vector sequences and was used for sequencing V inserts in both pTCSLC$_g$1NeoApa$^-$ and pTCSLCκDHFR vectors. Reverse primers were situated within the Cκ or C$_g$1 sequences, and used for sequencing Vκ and V_H inserts, respectively.

Double strand sequencing of the human kappa chain V region

The two samples from TM23 (#3 and #4) were sequenced for both strands. The reverse strand was sequenced entirely, through the restriction enzyme sites (3' - 5'; PVUII - XhoI) using the primer HUCK5PR. The forward strand sequence was determined using two different primers, PTCSFOR and PTCSFWD. Full length sequence through the restriction enzyme sites (5' - 3'; XhoI - PvuII) was obtained by combining the sequences yielded from each of the primers. A small area of compression was seen to be present in the forward strands which was not resolved with either of the primers or by using a sequencing method that alleviates such compressions. However, the reverse strand proved to have correct sequence in this area. Both samples #3 and #4 appeared to have correct sequence, with sample #4 chosen to be used for mammalian cell transfection.

Double strand sequencing of the TH27 gamma chain V region.

The V_H fragments of pTCSLHuV$_H$4HC$_g$1NeoApa$^-$ (isolates #6 and 9) were sequenced completely on both strands. The humanized V_H fragment of isolate #6 matched the expected sequence through the coding region. There was a single base pair substitution (C to T) at position -2 (relative to initiating ATG), which was not expected to affect expression. The V_H fragment of isolate #9 contained two substitutions in the coding region. The four additional candidate clones were not sequenced.

Co-transfect COS and CHO cells with TH27 and CONTROL plasmids

Recipient cells were CHO dhfr-DUX B11 and COS-1. The CHO DUX B11 cell line was obtained from Biogen Inc. For CHO transfection, five micrograms of the TM27 plasmids pTCSLHuVκ4HCκDHFR (isolate #4), pTCSLHuV$_H$4HC$_g$1NeoApa- (isolate #6), and the vectors pTCSLDHFRApa- and pTCSLNeoApa- was linearized by digestion with the restriction enzyme AatII. The TM27-containing plasmids (labelled "TM27L") were co-precipitated in ethanol and resuspended in H_2O. Similarly, the vector plasmids (labelled "CONTROL") were co-precipitated and resuspended in H_2O. Each of the co-precipitated plasmids were transfected into 10^7 CHO cells by electroporation, using a BioRad Gene Pulser at 250 V and 960 mF. Transfected cells were then grown in α Minimal Essential Medium (αMEM) supplemented with thymidine, adenosine and deoxyadenosine (non-selective media) for two days prior to selection.

For COS transfection, five micrograms of uncut TM27L plasmids and vector plasmids were co-precipitated in ethanol and resuspended in H_2O. Plasmids were transfected into $4.1x10^6$ COS cells by electroporation. Transfected cells were grown in Dulbecco's Modified Eagle Medium (DMEM) for three days, then supernatants harvested for assays.

Analyze supernatants of COS transfactants by human IgG1 ELISA and staining

Production of human IgG by COS cells was measured by human IgG1 ELISA on cell culture supernatants. Microtiter plates were coated with mouse anti-human IgG1 Fc antibody in PBS and blocked in blocking buffer (1% BSA in PBS). Supernatants were added in 100 µl samples; appropriate dilutions in blocking buffer were assayed as necessary. Captured IgG was detected with horseradish peroxidase-conjugated goat anti-human $Ig_κ$ antibody and OPD (o-phenylenediamine). Supernatants from TM27L-transfected COS cells showed levels of 95 ng/ml human IgG. Human IgG from CONTROL-transfected COS cells was undetectable. Flow cytometry was performed on supernatants from TM27L-transfected COS cells, concentrated ~6-fold. Intensity of staining on HPB cells, as indicated by mean channel fluoresence, was about 9-fold lower than that obtained with purified TM27L (purified from NS0 cells) at ~ 2.5 mg/ml. On Jurkat cells, staining was at background level.

Selection of CHO transfectants

CHO transfectants were placed under selection in αMEM media lacking nucleotide supplements (to select DHFR$^-$ on pTCSLHuVκ4HC$_g$DHFR) and containing Geneticin (G418: to select neo^r on pTCSLHuV$_H$4HC$_g$1NeoApa$^-$). Cells were plated at 2.0x10^5 cells/ml in 24-well plates.

Analyze supernatants of CHO transfectants by human IgG1 ELISA

After 12 days in selective media, supernatants from confluent wells were harvested and assayed by human IgG1/κ ELISA. Concentrations in TM27L supernatants ranged from 182 - 236 ng/ml. Most were >200 ng/ml. Human IgG1/κ was undetectable in CONTROL supernatants. Cells from twelve high-producing TM27 wells were expanded into T25 flasks. Single vials were frozen as backup cultures and stored under liquid nitrogen. Cells from two CONTROL wells were also expanded to T25 flasks and frozen.

First limiting-dilution plating of CHO transfectants

The four highest-producing wells were selected for limiting-dilution plating in 96-well plates: wells #1B1 (232.6 ng/ml), 1D6 (233.3 ng/ml), 2B2 (235.5 ng/ml) and 2C1 (236.0 ng/ml). For each well, three plates were established in selective media: one at 1 cell/well and two at 0.5 cell/well.

Supernatants from the first round of cloning were screened by human IgG1/κ ELISA. Cells from 90 high-producing wells were expanded to 24-well plates and allowed to grow to confluency. Supernatants of 12 of these cultures were assayed by human IgG1/κ ELISA and by flow geometry. Concentrations in the human IgG1/κ ELISA ranged from 211 - 1048 ng/ml. All samples stained HPB cells (mean channel fluoresence ranged from 46.5 to 169.9) and did not stain Jurkat cells. The six highest-producing clones were expanded and frozen (as backups) in liquid nitrogen. Of these, the two highest-producing clones (1B1-C7 and 2B2-H9) were selected to continue into subsequent rounds of cloning.

Two primary cultures of CONTROL constructs were also placed through one round of limiting-dilution plating in 96-well plates. One clone of each was selected and expanded. Supernatants were assayed and shown to give no detectable signal on the human IgG1/κ ELISA (page 640-83). Cells were frozen and are stored in liquid nitrogen.

Expansion of uncloned cultures to roller bottle scale for antibody purification

Two primary cultures (1B1 and 2B2) were expanded in order to produce sufficient material for TM27 antibody purification. The cells were expanded to 2L roller bottles and adapted to lower serum concentration (via T25 in 10% serum → T75 in 10% serum → 3xT75 in 5% serum → 3x2L bottles in 1% serum) (page 640-24). The 2L bottles were established on day 1, and fed (by replacing 1/2 volume with fresh media with 1% serum) on days 2, 5 and 8. First harvest was on day 10, second on day 12 and third on day 15. Harvested supernatants were spun and stored at -20 °C until purification.

Purification of antibody on Protein A column

Supernatants from the 2B2 cultures were pooled (total volume of 2.7 liters) and filtered through a 0.22 micron nylon membrane. Human IgG1 ELISA indicated that the total amount of starting material was 1090 mg. Antibody was purified on a Prosep-A (Protein A) column, eluted with 0.1M citrate at pH 5.0, 4.0 and 3.0. Eluates were dialyzed into PBS. Amounts of purified antibody in the eluted fractions were measured by human IgG1 ELISA. The pH 3.0 fraction contained most of the purified antibody, while a small amount was found in the pH 4.0 fraction. The human IgG1 ELISA was repeated to obtain more precise concentration measurements. Total yield of TM27 (pH 3.0 plus pH 4.0 fractions) was 960 mg.

Protein-A Purification of Antibodies from NS0 Transfectant Culture Supernatant

The antibodies from NS0 transfectant culture supernatants were purified on a Protein A column substantially as described above.

Characterization of TM 27 Antibody from NSO Myeloma Cell Lines

The TM27 mAb purified from the culture supernatants of NSO transfectants using Protein A column chromatography was used to stain HPB-ALL cells (Vβ 5.2) and Jurkat (Vβ 8.1) cells as a negative control in comparison with mouse mAb, TM23, and chimeric TM27. TM27 stained positively on HPB-ALL but not Jurkat.

To determine the specificity of TM27, normal PBL T cells were stained with TM27 in comparison with TM23 mAb. Both the TM27 and TM23 mAbs stained ~ 3.0 % of total human PBL T cells. A competition assay was set up in which a fixed concentration of FITC labeled TM23 mAb was added to various concentrations of unlabeled 4H11 or TM27 mAb. The

antibody mixtures were then used to stain HBP-ALL cells. Both the unlabeled TM23 and TM27 blocked staining by the FITC labeled TM23. To determine whether TM27 can comodulate with CD3 antigen, HPB-ALL cells were incubated with various concentration of TM27 or TM23 mAb overnight and then stained with anti-CD3 mAb. The results indicated that TM27 as well as TM23 could cause the endocytosis of TCR/CD3 complex.

A Scatchard analysis was performed to measure the binding affinity of TM27 on HPB-ALL cells. In both the Scatchard and competition (with TM23 mAb) assay, the results indicated that TM27 maintained approximately the same affinity (Kd = 2.0 x 10^{-8} M^{-1}), and could compete well with 4H11 mAb.

Comparison of TM27L/NS0 and TM27L/CHO in a competition assay with 4H11 was also performed and the results are depicted in Figure 3. A Scatchard analysis was performed and the results are depicted in Figure 4.

Second limiting-dilution plating of CHO transfectants

A limiting-dilution plating was set up as above. Wells were screened on the human IgG1 ELISA. Cells from six wells of each (1B1-C7 and 2B2-H9) were expanded to 24-well plates and assayed by human IgG1 ELISA. The two highest-producing clones from each were expanded and frozen in liquid nitrogen. The highest-producing clones from each were selected for the final round of subcloning.

Third limiting-dilution plating of CHO transfectants

A limiting-dilution plating was set up as above. Wells were screened on the human IgG1 ELISA. Cells from six wells of each (1B1-C7 and 2B2-H9) were expanded to 24-well plates and assayed by human IgG1 ELISA. The three highest-producing clones from each were expanded to T25 flasks and frozen in liquid nitrogen. A single clone from each was expanded to a T75 flask. At this time, Geneticin was dropped from the media, thus removing seletion for neo^r. Selection for DHFR was retained. Supernatants were assayed by human IgG1 ELISA. The 1B1-C7 clone produced 4.69 mg/10^6 cells/day. The 2B2-H9 clone produced 2.65 mg/10^6 cells/day. The C7 isolate was therefore chosen as the final clone and the H9 isolate was designated as an alternate clone.

Preparation of frozen cell stocks

The final clone chosen above was expanded to five T225 flasks to provide a sufficient number of cells to establish a bank of frozen cell stocks. Cells were harvested from these flasks and pooled. The pooled suspension contained 98.3% viable cells. The total viable cell count was 1.58 X 10^8, sufficient to prepare the desired number of 72 vials at 2 x 10^6 cells/vial. Cells were frozen and labelled "TM27L-662-35". In addition, the alternate H9 clone was expanded to one T225 flask to establish a small frozen cell bank. Cells harvested from this flask were 97.3% viable. The suspension contained sufficient cells to prepare 9.7 vials at 2 x 10^6 cells/vial (page 662-89). Cells were frozen in liquid nitrogen and labelled as "TM27L-662-89".

Testing of cells revived from frozen stock.

Viability and Mycoplasma Testing

One vial of frozen cell stock TM27L-662-35 was thawed and recovered in αMEM. Cell viability was determined to be 88% by trypan blue staining. A confluent culture grown from this vial was tested for mycoplasma contamination using a Bionique Testing Laboratories kit. The test detected no contamination. In addition, one vial of frozen cell stock TM27L-662-89 was thawed. Cell viability was determined to be 93.8% by trypan blue staining. A confluent culture grown from this vial was tested for mycoplasma contamination as above. The test detected no contamination. Both cultures were expanded to T75 flasks and cell culture supernatants assayed by human IgG1 ELISA. The TM27L-662-35 clone produced 3.24 μg/10^6 cells/day, and the TM27L-662-89 clone produced 2.52 μg/10^6 cells/day. In a second assay, TM27L-662-35 produced 3.03 μg/10^6, cells/day, and TM27L-662-89 produced 2.74 μg/10^6cells/day.

Expression of TM27L-662-35 to roller bottle scale for antibody purification

The TM27L-662-35 culture was expanded in order to produce sufficient material for TM27L antibody purification. The cells were expanded to 2L roller bottles and adapted to lower serum concentration (via T25 in 10% serum to T75 in 10% serum to 3 x T75 in 5% serum to 3 x 2L bottles in 1% serum). The 2L bottles were established on day 1 and fed (by replacing 1/2 volume with fresh media with 1% serum) on days 2, 5, 7 and 9. First harvest was on day 12, second on day 14 and third on day 16. Harvested supernatants were spun, filtered and stored at -70 °C until purification.

Purification of antibody on Protein A column

Supernatants from the TM27L-662-35 cultures were pooled for a total volume of 5.1 liters. Antibody was bound to a Prosep-A (Protein A) column and eluted with 0.1 M citrate at pH 5.0 and 3.0. The pH 3.0 eluate was dialyzed into PBS. Amounts of purified antibody in the starting material and eluted fractions were measured by human IgG1/kappa ELISA. The pH 3.0 fraction contained approximately 3.9 mg of antibody in 25 ml. The material was concentrated on a Centriprep-30 concentrator (Amicon) and the human IgG1/ kappa ELISA was repeated. Total yield of TM27L was -3.2 mg at 1.6 mg/ml.

Primary characterization of purified antibody

Flow cytometry was performed on ~2.5 mg of material from each of the pH 3.0 and pH 4.0 fractions and TM27 purified from NS0 cells. Intensity of staining on HPB cells, as indicated by mean channel fluorescence, was measured as follows: 283.46 for the pH 3.0 fraction; 321.98 for the pH 4.0 fraction; 506.79 for NS0-derived TM27. On Jurkat cells, staining was at background level for all samples.

Example 2

Sequences of CDR Grafted TM27 Antibodies Compared to Human Framework Cassettes

TM27 Vκ

```
1     DIQMTQSPSSLSASVGDRVTITCSASQGISNYLNWYQQTPGKAPKLLIYY 50
REI   ----------------------------------------Q------D-IK----------------------------------E
TM23  ----------TT----------L----------S----------------------------K--DGTV----------
WAL   ----------------------------------------R------S----------.----------K----------------A

51    TSSLHSGVPSRFSGSGSGTDYTFTISSLQPEDIATYYCQQYSKLPRTFGQ 100
REI   A-N--QA----------------------------------------------------------------QS---Y-------
TM23  ----------------------------------------SL-----N--E----------------------------------G
WAL   A-----Q-----T----------F-L----------S----------SYSTLI-------

101   GTKLQIT   107
REI   ----------
TM23  ------E-K
WAL   -----R-E-K
```

TM27 VH

```
1*    QVQLQESGPGLVRPSQTLSLTCTVSGFSLTAYGVNWVRQPPGRGLEWIGM  50
NEWM --------EQ----------------------------------STFSNDYYT----------------------I--Y
TM23 --------K----------A-------S-----I-------------------------------------------K---------.--
HIL  ----K---VQA-G--V--Q--GRS--R-S---IA------TFSN----MH---------A-----K--------VAV

51    WGDGNTDYNSALKSRVTMLKDTSKNQKSLRLSSVTAADTAVYYCARDRV  100
NEW  VFYH-TS---DTTPLR---------.-V-------------------------------------------------NLI
TM23 ------------------------------LSIS----N---S---VF-KMN--LQTD----R-------------
HIL  ----YN-S---Y--GDSV-G--F---ISR--N---RTLYMZMN--LRTE-----V----------PD
          R

101   TATLYAMDYWGQGSLVTYSS  120
NEWM AGCIDV-------------------------.--
TM23 ------------------------------TS-----.--
HIL  IL--AFSF----------------VL-----.--
```

* HUVHL has L AT #48

TM27L	with Leucine at position 48 of heavy chain.
TM27I	with Isoleucine at position 48 of heavy chain.
TM27.1	with FS to VF (78-79) changes.
TM27.2	with VTML/T to LSIS/N (67-70/73) changes.
TM27.3	with V to R (92) changes.

Example 3

Scatchard Analysis of TM23

Exp	Kd	Receptors/cell
617:096	3.27e-8 M	3.04e5
617:115	2.30e-8 M	2.43e5
617:119	2.46e-8 M	2.46e5

Mean

Kd = 2.52 (±0.67)e-8 M

Receptors/cell = 2.64e5

Scatchard Analysis of TM27

Exp	Kd	Receptors/cell
613:78	1.48e-8 M	3.31e4
613:82	1.14e-8 M	2.47e4

Mean

Kd = 1.31(±0.24)e-8 M

Receptors/cell = 2.89e4

Example 4

Summary of Work on TM27I

TH27I (48) in COS/CHO:

* cDNA was prepared from the CDR-grafted V regions in HS0 producing line, amplified by PCR, sequenced.
* DNA fragments were cloned into pTCSLNeo in cDNA configuration.
* The plasmid was co-transfected with pTCSLDHFRTM27κ into COS and CHO.
* Positive ELISA results of human IgG1/κ expression from COS cell supernatant indicated that the plasmids constructs were good.
* CHO cell transfectants were selected, expanded and cloned. The cells were frozen in LN_2 after one cloning.

Example 5

AMINO ACID SEQUENCE DATA DERIVED FROM 16G8 cDNA

HEAVY CHAIN

```
          10         20         30         40         50
           *          *          *          *          *
TTRAP RSSHS VISTE HRPLT MDSRL NLVFL VLILK GVQCD VQLVE SGGGL

          60         70         80         90         100
           *          *          *          *          *
VQPGG SRKLS CAASG FTFSN FGMHW VRQAP DKGLE WVAYI SSGSS TIYYA

         110        120        130        140        150
           *          *          *          *          *
DTLKG RFTIS RDNPK NTLFL QMTSL RSEDT AMYYC ARRGE GAMDY WGQGT

         160        170
           *          *
SVTVS SAKTT PPSVY PLAPG
```

LIGHT CHAIN

```
          10         20         30         40         50
           *          *          *          *          *
ISQGT KFKYT MDFQV QIFSF LLISI SVVMS RGENV LTQSP AIMSA SLGEK

          60         70         80         90         100
           *          *          *          *          *
VTMSC RASSS VNYIY WYQQK SDASP KLWIY YTSNL APGVP TRFSG SGSGN

         110        120        130        140        150  153
           *          *          *          *          *    *
SYSLT ISSME GEDAA TYYCQ QFTSS PFTFG SGTKL EIKRA DAAPT VSIFP PSS
```

PROTEIN SEQUENCING DATA:

H = DVQLVE?GGGLVQPG

L = ENVLTQ

Example 6

SUMMARY OF WORK ON TM29

* Hybridomas were prepared substantially as in Example 1 and cDNA (depicted in Example 5) was isolated.
* Isolated cDNAs (H and κ) (from the 16G8 hybridoma) were cloned into M13 vectors.
* Human framework ($VH_{KOL}/V\kappa_{REI}$) was introduced through mutagenesis.
* CDR-grafted V regions were cloned into myeloma expression vectors.
* Humanized antibodies (~ 1 mg each) were purified from NS0 transfectants for preliminary evaluation.

Four different TM29's have been made:

TM 29

TM29.1 with SS to AA (23-24) changes.

TM29.2 with S to P (75) change.

TM29.3 with GV/F to AM/Y (92-93/95) changes.

DETAILED EXPERIMENTATION - TM29

Isolation of the human kappa chain constant region by the PCR method.

Using the undigested plasmid pSV184DH-Neo/DNS-VκCκ (pSVHuk which contains the human kappa chain constant region) PCR was performed with two primers (one forward, one reverse) to isolate the constant region. The oligomers (primers) were obtained from Operon. The PCR reaction reagents were from the GeneAmp DNA amplification reagent kit with AmpliTaq (Perkin-Elmer Cetus). The PCR reactions were set up in three different concentrations of Mg^{2+}. Aliquots of the PCR products were analyzed on a 1% agarose gel. The PCR product from the first sample of Mg^{2+} (concentration 1.5 mM) was chosen. The primers, which flank the C region, were designed containing restriction enzyme sites for cloning purposes. Thus, the amplified DNA was digested with EcoRI (at the 3' end). The site at the 5' end (PvuII) was not digested. The digested DNA fragment was then run on a 1% agarose gel

and the 300 bp band excised. The DNA was extracted from the gel and a small aliquot was analyzed on a 1% agarose gel to determine quality and to estimate quantity. The 300 bp band appeared to be clean and the total amount of DNA estimated to be ~ 140 ng.

Human kappa chain constant region cloned into the pBS KS$^{\pm}$ vector and sequenced.

The EcoR I-digested DNA fragment of the human kappa chain constant region was inserted into the vector pBS KS^{+}, which had been digested with Sma I and EcoR I, using T4 DNA ligase. This ligation mixture was used to transform *E. coli* DH5α competent cells. Single colonies were picked and used to inoculate 24 liquid cultures. DNA was prepared from each of the cultures. Each DNA sample was digested with PvuII (5' end of Cκ) and ECoRI (3' end of Cκ), and analyzed on a 2% agarose gel. Due to difficulty in separating the lower range bands (300 bp), the DNA samples were digested with PvuII only. After analysis on a 2% agarose gel, 19 out of 24 were correct for Cκ insert. Samples #1 - 5 (of the 19 correct samples) were chosen to be sequenced, using T7 and T3 primers (sequences in the vector flanking the insert). Other sequencing reagents were from the Sequenase Version 2.0 kit (USB). Sample #5 proved to have correct sequence for the human kappa chain constant region.

Human kappa chain constant region cloned into the mammalian expression vector - pTCSLDHFR* to become pTCSLCκDHFR*

The human kappa chain constant region was isolated from the pBS KS^{+} vector (Sample #5) using PvuII and EcoRI restriction enzymes. However, due to the previous problem of band separation, a second digest was performed with the construct pBS/Human Cκ. The second digest involved EcoRI and XbaI (located 5' to PvuII site) yielding a fragment ~ 300 bp. This fragment was then gel purified and subjected to a PvuII digest, to remove the remaining small fragment of vector at the 5' end of the Cκ. This final fragment was gel purified and then analyzed on a 11 agarose gel for quality and to estimate quantity. The Cκ DNA appeared to be a clean band at ~ 280 bp and the total quantity estimated to be ~135 ng.

Due to the existence of three EcoRI sites in the expression vector, pTCSLDHFR*, it was determined that cloning into the PvuII site only would be best. Since the Cκ DNA was prepared PvuII - EcoRI, the EcoRI end was repaired (to form blunt end) using Klenow fragment. The repaired, blunt-ended Cκ DNA was ligated to the pTCSLDHFR* vector which had been PvuII digested and phosphatase treated (using calf intestinal phosphatase). This

ligation mixture was used to transform *E. coli* DH5α competent cells. Single colonies were picked and used to inoculate 24 liquid cultures. DNA was prepared from each. All DNA samples were digested with PvuII and HindIII, and analyzed on a 1% agarose gel. It was determined that five samples, out of 24, contained the Cκ insert in the correct orientation. Sample #7 was chosen for continuation.

Replacement of the mutant DHFR* gene with the wild type DHFR gene.

For the expression of the vector pTCSLCκDHFR* in CHO(dhfr$^-$)DUX B11 cells, the DHFR* gene was replaced with the DHFR gene. DNA from pTCSLCκDHFR* sample #7 as well as pSV2-DHFR (plasmid obtained from Dr. Paul Berg, Department of Biochemistry, Stanford University Medical School) was digested with restriction enzymes HindIII and BglII, to isolate the respective gene fragments. Upon analysis, it was determined that an additional BglII site in the pTCSLCκDHFR* vector removed a portion of the polyA site also, which would interfere with the vector function. Therefore, the digests were repeated using HindIII and BamHI. The (DHFR*)pTCSLCκ portion of the vector was treated with phosphatase. Then both the isolated fragments (pTCSLCκ and DHFR) were gel purified, and analyzed on a 1% agarose gel for quality and to estimate quantity. Both bands appeared to be clean and estimated to be of similar concentration, ~ 40-50 ng/ml. The DHFR gene was ligated to the pTCSLCκ vector using T4 DNA ligase. The ligation mixture was used to transform *E. coli* DH5α competent cells. Single colonies were picked and used to inoculate 24 liquid cultures. DNA was prepared from each. The DNA was analyzed by restriction enzyme digestion with HindIII and BamHI. From 24 cultures, 23 had the correct bands indicating the insertion of the DHFR gene into the pTCSLCκ vector. Sample #22 was selected for further use.

Cloning the PCR isolated human kappa chain V region into the CHO vector pTCSLCκDHFR.

The cDNA for the kappa chain V region for 16G8 had been gel purified and digested with the appropriate restriction enzymes. A small aliquot of the DNA sample was analyzed on a 1% agarose gel to estimate quantity and to verify the quality. An extra (contaminating) band was observed at ~ 800 bp and the concentration was estimated at 1.25 ng/ml. The V region was ligated to the pTCSLCκDHFR vector, which had been digested with restriction enzymes XhoI and PvuII and phosphatase-treated. The ligation mixture was used to transform *E. coli* DH5α

competent cells. Yield of transformant colonies was very low. Single colonies were picked and used to inoculate twelve liquid cultures. DNA was prepared from each, digested with XhoI and PvuII and analyzed on a 1% agarose gel. There were four positive samples: #1, #4, #7 and #10. Each of the positive samples was sequenced. The PCR isolated TM29 gamma chain V region was cloned into the CHO vector pTCSLC$_g$1NeoApa*.

Double strand sequencing of the human kappa chain V region.

The four samples from 16G8 (#1, #4, #7 and #10) were sequenced for both strands. The reverse strand was sequenced entirely, through the restriction enzyme sites (3' - 5'; PvuII - XhoI) using the primer HUCK5PR. Both samples #l and #4 appeared to have correct sequence, sample #7 did not sequence, and sample #10 had the wrong sequence. For samples #1 and #4, the forward strand sequence was determined using two different primers, PTCSFOR and PTCSFWD. Full length sequence through the restriction enzyme sites (5' - 3'; XhoI - PvuII) was obtained by combining the sequences yielded from each of the primers. A small area of compression was seen to be present in the forward strands which was not resolved with either of the primers or by using a sequencing method that alleviates such compressions. However the reverse strands proved to have correct sequence in this area. Both samples #l and # 4 appeared to have correct sequence, with sample #4 chosen to be used for mammalian cell transfection.

Co-transfection of TH29 light and heavy chain plasmids into COS and CHO cells.

Both Cos-1 and CHO(dhfr$^-$)DUX B11 cells were used in the co-transfection of TM29 light and heavy chain plasmids. The CHO DUX B11 cell line was obtained from Biogen, Inc. and originally was derived by Dr. Lawrence Chaisin (Columbia University).

For the Cos-1 cell transfection, 5 mg of each undigested TM29 plasmid (pTCSLHuVκ16GCκDHFR and pTCSLHuV$_H$16GC$_g$1NeoApa*; TM29-4 and TM29-26, respectively) Was precipitated in ethanol and resuspended in H_2O at a 1 mg/ml concentration. The plasmids were then co-transfected into 3.8 x 10^6 cells by electroporation, using a BioRad Gene Pulser at 250V, 960 mFD. The transfected cells were resuspended in Dulbecco's Modified Eagle's medium (DMEM) and placed in culture for three days. Control vector plasmids (pTCSLDHFRApa* and pTCSLNeoApa*) were prepared and co-transfected similarly to the TM29 plasmids. The subsequent transfected cells were resuspended and cultured similarly to the TM29 transfected Cos cells.

For the CHO DUX B11 cell transfection, 5 mg of each TM29 plasmid (as above) was digested with the restriction enzyme AatII to yield a linear fragment of DNA, precipitated in ethanol and resuspended in H_2O at a 1 mg/ml concentration. The plasmids were then co-transfected into 5 x 10^6 cells by electroporation, as above. The transfected cells were resuspended in α Minimal Essential medium (αMEM) supplemented with thymidine, adenosine and deoxyadenosine (non-selective medium) and placed in culture for three days. control vector plasmids (as above) were prepared and co-transfected similarly to the TM29 plasmids. The subsequent transfected cells were resuspended and cultured similarly to the TM29 transfected CHO cells.

Analysis of TH29 transfacted Cos cell supernatant by human $IgG_{1\kappa}$ ELISA.

After three days in culture, the TM29 transfected Cos cells appeared to be confluent. Therefore, the supernatant was collected and the cells were discarded. Human IgG production by the cells was determined from the culture supernatant using the human $IgG_{1\kappa}$ ELISA. The supernatant IgG was captured by a mouse anti-human IG_1 Fc antibody, which had been coated on a 96-well plate, then detected by a horseradish peroxidase-conjugated goat anti-human Igκ antibody and color development with o-phenylenediamine (OPD). The supernatant IgG concentrations were determined by comparing optical density measurements to a curve generated by a purified human $IgG_{1\kappa}$ antibody of known concentration. The TM29 transfected Cos cell supernatant yielded a human IgG concentration of 201.3 ng/ml. The control DNA transfected Cos cell supernatant yielded a human IgG concentration of 0 ng/ml.

Selection of TH29 transfected CHO cells.

After three days in culture, the TM29 transfected CHO cells appeared to be confluent. Therefore, the cells were treated with 0.25% trypsin (in Hank's Balanced Salt solution) to dissociate them from the culture flasks. The cells were collected, centrifuged and the supernatant removed. The cell pellet was resuspended in αMEM selection media at 2 x 10^3 cells/ml and plated at 1 ml/well in a 24-well plate. The αMEM selection media had a dual selection capability, lacking nucleotide supplements and replacing FBS with dialyzed FBS to select for $DHFR^+$ on pTCSLHuVκ16GCκDHFR, and adding Geneticin (G-418) to select for neo^r on $pTCSLHuV_H16GC_g$NeoApa*.

Analysis of TX29 transfected CHO cell supernatant by human $IgG_{1\kappa}$ ELISA and by flow cytometry.

After 14 days in selection media, TM29 transfected CHO cell supernatants were collected from each well and assayed using the human $IgG_{1\kappa}$ ELISA. The human IgG concentration in the TM29 transfected CHO cell supernatants ranged 144.6 ng/ml - 291.9 ng/ml, with an average concentration of 240.5 ng/ml. The human IgG concentration in the control DNA transfected CHO cell supernatants was 0 ng/ml.

Supernatants from the three highest producing wells for the TM29 transfected CHO cells (2C3, 2C4, 2B6) and one well for the control DNA transfected CHO cells (3D4) were also assayed by flow cytometry. The three TM29 samples, 2C3, 2C4 and 2B6, stained Jurkat cells positively (mean channel fluorescence ranged 186.61 - 290.01) and did not stain HPB cells positively (mean channel fluorescence ranged 11.77 - 13.82). The control sample, 3D4, did not stain Jurkat cells positively (mean channel fluorescence 11.96).

Cloning of TH29 transfected CHO cells by limiting dilution.

The three highest producing wells for the TM29 transfected CHO cells (2C3, 291.9 ng/ml; 2C4, 289.7 ng/ml; 2B6, 289.5 ng/ml) were chosen for cloning by limiting dilution. For each of the wells, the cell concentrations were diluted in αMEM selection media and plated at 2 cells/well, 1 cell/well, and 0.5 cell/well in 96-well plates. Ten days after cloning, colony growth was observed in numerous wells on each of the 0.5 cell/well plates for 2C3, 2C4 and 2B6. Therefore, supernatants from all wells from each of the 0.5 cell/well plates were collected and assayed using the human $IgG_{1\kappa}$ ELISA. All wells that exhibited cell growth were positive in the ELISA. A total of 24 of the highest producing wells (10 from 2C3, 14 from 2B6) were chosen for expansion into a 24-well plate and grown to confluency. Supernatants were collected from each well and assayed using the human $IgG_{1\kappa}$ ELISA. The human IgG concentration ranged 0.7 mg/ml - 4.8 mg/ml, with an average concentration of 2.9 mg/ml. Supernatants from the six highest producing wells for the TM29 transfected CHO clones (1C7, 1D9, 1G2, 2G1, 2G10, 2H2a) were then assayed by flow cytometry. All six samples stained Jurkat cells positively (mean channel fluorescence ranged 313.68 - 356.87) and did not stain HPB cells positively (mean channel fluorescence ranged 7.59 - 8.16). Each of the six samples was expanded to a 25

cm^3 flask, frozen and stored under liquid nitrogen. The three highest producing wells (2C3-1G2, 2C3-2G10, 2C3-2H2a) were chosen for subcloning.

The one well for control DNA transfected CHO cells (3D4) was cloned by limiting dilution similar to the TM29 wells. By diluting the cell concentration 10-fold lower than desired, no colony growth was observed ten days after cloning. Therefore, a fresh culture was started from a frozen vial (3D4). Cloning by limiting dilution was repeated using this new culture. Sixteen days after cloning, six of the most confluent wells from the 0.5 cell/well plates (1BS, 1C11, 1D6, 2E7, 2F3, 2F9) were chosen for expansion into a 24-well plate. Each well was subsequently expanded to a 25 cm^3 flask and then to a 75 cm^3 flask and grown to confluency. Supernatants were collected from each flask and assayed using the human $IgG_{1\kappa}$ ELISA. The human IgG concentration was 0 ng/ml. Each flask was then frozen and stored under liquid nitrogen.

Enansion of uncloned TM29 transfacted CHO cell cultures to roller bottle scale.

The three highest producing wells for the TM29 transfected CHO cells (2C3, 291.9 ng/ml; 2C4, 289.7 ng/ml; 2B6, 289.5 ng/ml) were also chosen for expansion into 25 cm^3 flasks as uncloned cultures. At this time, one vial of each of the TM29 cultures (as well as one vial of the control DNA transfected CHO cell culture) was frozen and stored under liquid nitrogen. Each of the uncloned cultures was then expanded to a 75 cm^3 flask. From one 75 cm^3 flask, each culture was expanded to 3 75 cm^3 flasks while decreasing the serum concentration in the αMEM medium from 10% to 5%. Each of the 75 cm^3 flasks for both 2C3 and 2B6 was then expanded to a 2L roller bottle. All 3 75 cm^3 flasks for 2C4 were frozen and stored under liquid nitrogen. Once the cultures were in the roller bottles, the serum concentration in the αMEM medium was decreased again, from 5% to 1%. After nine days in culture, the 2B6 bottles appeared to be confluent while the 2C3 bottles appeared to be ~ 25% confluent. The supernatants from all 2B6 and 2C3 bottles were collected on Day 9, Day 14, and Day 16, after which the cells were discarded. The collected supernatants were centrifuged and stored at -20 °C. An aliquot of each of the three harvested supernatants for both 2B6 and 2C3 was assayed using the human $IgG_{1\kappa}$ ELISA. The total quantity of human IgG for 2B6 was determined to be 820 mg; for 2C3, 1923 mg.

Subcloning of TH29 transfected CHO cell clones by limiting dilution.

The three highest producing wells for the TM29 transfected CHO cell clones (2C3-1G2, 4.8 mg/ml; 2C3-2H2a, 4.1 mg/ml; 2C3-2G10, 4.0 mg/ml) were chosen for subcloning by limiting dilution. Fourteen days after subcloning, colony growth was observed in numerous wells on each of the 0.5 cell/well plates for 1G2, 2G10 and 2H2a. Therefore, supernatants from those wells possessing colony growth were collected and assayed using the human $IgG_{1\kappa}$ ELISA. All wells exhibiting cell growth were positive in the ELISA, yielding a cloning efficiency of 100%. A total of 24 of the highest producing wells (2 from 2H2a, 5 from 2G10, 17 from 1G2) were chosen for expansion into a 24-well plate and grown to confluency. Supernatants were collected from each well and assayed using the human $IgG_{1\kappa}$ ELISA. The human IgG concentration ranged 1.2 mg/ml ~ 9.2 mg/ml, with an average concentration of 4.2 mg/ml. Supernatants from the four highest producing wells for the TM29 transfected CHO subclones (2G5, 2H12, 1H6, 2E9) were also assayed by flow cytometry. All four samples stained Jurkat cells positively (mean channel fluorescence ranged 211.51 - 292.45) and did not stain HPB cells positively (mean channel fluorescence ranged 8.36 - 8.80). Each of the four samples was expanded to a 25 cm^3 flask.

Second subcloning of TM29 transfected CHO cell clones by limiting dilution.

The two best TM29 transfected CHO cell clones (2C3-1G2-2G5 and 2C3-1G2-1H6), based on the combined results of the ELISA and flow cytometry data, were chosen for a second subcloning by limiting dilution. At this time, one vial of each of the clones (as well as one vial of each of 2C3-1G2-2H12 and 2C3-1G2-2E9) was frozen and stored under liquid nitrogen. Twelve days after subcloning, colony growth was observed in numerous wells on each of the 0.5 cell/well plates for 2G5 and 1H6. Therefore, supernatants from those walls possessing colony growth were collected and assayed using the human $IgG_{1\kappa}$ ELISA. All wells exhibiting cell growth were positive in the ELISA, yielding a cloning efficiency of 100%. A total of 24 of the highest producing wells (20 from 2G5, 4 from 1H6) were chosen for expansion into a 24-well plate (#646-101) and grown to confluency. Supernatants were collected from each well and assayed using the human $IgG_{1\kappa}$ ELISA. Due to lost activity with the human $IgG_{1\kappa}$ standard used in the assay, the concentrations for the unknown samples could not be determined. Therefore, the highest producing wells were determined by optical density. The six highest

producing wells (5 from 2G5, 1 from 1H6) were chosen for expansion into 25 cm^3 flasks, then into 75 cm^3 flasks and grown to confluency. Supernatants were collected from each flask and assayed using the human IgG_{1K} ELISA. The human IgG concentration ranged 14.6 mg/ml - 21.2 mg/ml, with an average concentration of 18.6 mg/ml. Supernatants from the six flasks for the TM29 transfected CHO subclones (1BS, 1D1, 1F11, 2B10, 2F5, 2F9) were then assayed by flow cytometry (#646-122). All six samples stained Jurkat Cells positively (mean channel fluorescence ranged 304.51 - 383.00) and did not stain HPB cell positively (mean channel fluorescence ranged 8.67 - 10.25).

Preparation of a frozen cell stock for the TH29 transfacted CHO cell clone.

The TM29 transfected CHO cell clone chosen to prepare a frozen cell stock was 2C3-1G2-1H6-2F5. This clone was expanded from one 75 cm^3 flask to five 225 cm^3 flasks in order to provide a sufficient number of cells to establish a frozen cell bank for research purposes. The frozen cell bank was prepared following the SOP TMB1-001.00. The cells were treated with trypsin, collected from each flask and pooled. The pooled cell suspension was found to contain 3.2×10^8 cells, with a 99% viability. Seventy-two vials were prepared, each containing 2×10^6 cells in a 1 ml volume. All vials for the clone have been labelled TM29-646-132.

Example 7

Sequences of CDR Grafted TM29 Antibodies Compared to Other Framework Regions

TM29 Vk

1	DIQMTQSPSSLSASVGDRVTITCRASSSVNYIYWYQQTPGKAPKLLIYYT	50
REI	----------Q---QDIIKYLN----------EA	
16G8	ENVL--------AIM------L--EK---MS---------- ----------KSDAS------W-------	
HIC	E--VL--------GT --L---P--ERA--LS--------Q---SSYLA---------K----Q-----R------GA	
51	SNLAPGVPSRPSSGSGTDYTFTISSLQPEDIATYYCQQFTSSPFTFGQGT	100
REI	------QA----------YQ--L-Y----------	
16G8	----------T----------NS---SL-------MEG----A----------S---	
HIC	-SR----T--I--D----------F---L-----R-E- ----F-V---------YG----W----------	

```
101   KLQIT      106
REI   --------
16G8  ----E-K
HIC   --VE-K
```

TM29 VH

```
1     EVOLVESGGGVVOPGRSLRLSCSSSGFTFSNFGMHWVROAPGKGLEWVAY
KOL   Q----------------------------------------------I --SYA---Y----------I
16G8  D----------L---------G-RK------AA--------  ----------D----------
WEA   Q--------D---------L--E-----G-----------A-----------AND---N----------LSF

51    ISSGSSTIYYADTLKGRFTISRDNSKNTLFLOMDSLRPEDTGVYFCARRG
KOL   -WDDG----DQH------SV-------------------------------------------------------D-
16G8  ----------------------------------------------P----------T------S-------AM-Y---------
WEA   -GGSG----------SV----------NB-------S-Y--------S------A-------A--YY---------

101   EGAM----------DYWGOGTPVTVSS
KOL   GHGFCSSASCFGP----------
16G8  ----          ----------S---------
WEA   --WLL----------N----------L---------
```

TM29 with the sequences of lst version

TM29.1 with SS to AA (23-24) changes

TM29.2 With S to P (75) change

TM29.3 with GV/F to AM/Y (92-93/95) changes

Example 8

Scatchard analysis of TM29 mAb

Scatchard analyses of TM29 and 16G8 antibodies were performed in three independent experiments. The table below shows the Kd determined for each of the experiments.

Exp.	TM29 (Kd M)	16G8 (Kd M)
1	1.45×10^{-9}	2.86×10^{-9}
2	1.65×10^{-9}	2.65×10^{-9}
3	1.15×10^{-9}	2.44×10^{-9}
Mean	1.42×10^{-9}	2.65×10^{-9}

Example 9

Summary of Work on Chimeric TM29

Chimerics in COS and CHO:

* V(D)J region cDNAs were cloned into CHO cell expression vectors, pTCSLNeo (H) and pTCSLDHFR (k) in cDNA configuration.
* Expression plasmids were co-transfected into COS and CHO.
* Ab concentration in supernatant from COS cells ~ 35 ng/ml.
* CHO cell transfectants were analyzed (by FACS and ELISA), cloned once, and were amplified with MTX (20 nM, 100 nM and 500 nM).
* The antibody concentration from unamplified CHO cell clones was ~ 3 μg/ml; and stained Jurkat cells positively. The unamplified clones were frozen in LN_2.

Example 10

Expression of Chimeric 16G8 in COS and CHO Cells 16G8 Chimeric

Light chain and heavy chain constructs contained complete mouse V region with human constant region in the same expression vectors used for CDR-grafted constructs: pTCSDHFR link pTCSNeo link - the constructs had double selection capability.

COS cell transfectants - ~ 35 nglml

CHO cell transfectants - 16 positive wells/54 wells

	ng/ml		corrected for volume
1A1	151.6		
1A6	166.6		
1B3	107.2		
1B6	179.6		
1D2	101.0		
1D5	215.1	*	2.2 μg/ml
2A4	199.8		
2B1	184.5		
2B2	213.8	*	2.1 μg/ml
2B4	130.6		

2C2	119.6		
2C6	214.4	*	2.1 μg/ml
2D3	216.6	*	2.2 μg/ml
2D6	139.7		
3A1	198.7		
3A2	172.2		

all expanded to 25 cm^3 flasks

* Flow cytometry performed.

Example 11

Summary of Work on TM29

TM29

TM29 in COS/CHO:

* cDNA was prepared from the CDR-grafted V regions in NS0 producing line, amplified by PRCR, sequenced.
* DNA fragments were cloned into pTCSLNeo (H) and pTCSLDHFR (k) in cDNA configuration.
* Plasmids were co-transfected into COS and CHO.
* Supernatant from COS cells contains ~ 2 μg/ml Ab.
* CHO cell transfectants secreted ~ 3 μg/ml Ab before cloning and amplification, supernatant stained Jurkat cells positively. Cell bank from the best unamplified Ab producer was prepared after three times of cloning.
* The CHO cell lines were amplified under various concentration of MTX.

Example 12

Comparison of TM27L, TM27I and TM23 in 4H11-FITC Competition Assay

Exp 1:

$$\frac{\text{slope TM23}}{\text{slope TM27L}} = 1.56$$

$$\frac{\text{slope TM23}}{\text{slope TM27I}} = 2.04$$

Exp 2:

$$\frac{\text{slope TM23}}{\text{slope TM27L}} = 1.82$$

$$\frac{\text{slope TM23}}{\text{slope TM27I}} = 2.44$$

Compared with TM23, the values from both TM27L and TM27I are all <3 fold.

Example 13

TM27 Methotrexate Amplification

Productivity (micrograms/10^6 cells/day)

		Methotrexate Level (nM)				
		0	20	100	10-200	20-200
C7	*7/30	27	nd	nd	nd	265.8
	*7/22	3.5	nd	19.2	14.7	34.2
	7/6	2.6	nd	nd	9.1	nd
	6/25	2	4.8	4.5	nd	nd
	6/18	3	8.9	nd	nd	nd
H9	*7/30	22.7	nd	nd	nd	181.9
	&7/22	3.9	nd	9.4	11.6	34.4
	6/25	1.5	1.8	4.7	3.6	nd
	6/18	1.8	1.5	nd	nd	nd

* No interassay comparison.

Intra-assay relative value

		0	20	100	20-100	20-200
C7	*7/30	1	nd	nd	nd	9.8
	*7/22	1	nd	5.5	4.2	9.8
	7/6	1	nd	nd	3.5	nd
	6/25	1	2.4	2.3	nd	nd
	6/18	1	3.0	nd	nd	nd
H9	*7/30	1	nd	nd	nd	8.0
	*7/22	1	nd	2.4	3.0	8.8
	6/25	1	1.2	3.1	2.4	nd
	6/18	1	0.8	nd	nd	nd

Status

		0	20	100	20-100	20-200
C7	8/11	fm	fm,s,sf4	fm	fm	fm,s*
H9	8/11	fm	discard	fm,s,f2	fm	fm,s*

Key fm = froze 1 tube mixed culture

s = subculture by one limiting dilution

s* = first limiting dilution in progress

f# = froze one vial of each of # wells

CLAIMS

1. A humanized antibody or binding protein thereof having an amino acid sequence that comprises all or a portion of CDRs substantially derived from a monoclonal antibody having a specificity for a select subpopulation of T cells expressing the Vβ5.2 or Vβ5.3 variable region of the T cell antigen receptor beta chain, in conjunction with select variable framework residues also derived from said monoclonal antibody, said humanized antibody or binding protein thereof capable of binding to said select subpopulations of T cells with a binding affinity at least comparable to that exhibited by said monoclonal antibody, wherein

the humanized antibody or binding protein thereof comprises the TM27 Vκ amino acid sequence:

1 DIQMTQSPSSLSASVGDRVTITCSASQGISNYLNWYQQTPGKAPKLLIYY 50

51 TSSLHSGVPSRFSGSGSGTDYTFTISSLQPEDIATYYCQQYSKLPRTFGQ 100

101 GTKLQIT 107

and further comprises an amino acid sequence selected from the group of the TM27 V_H amino acid sequence:

1 QVQLQESGPGLVRPSQTLSLTCTVSGFSLTAYGVNWVRQPPGRGLEWLGM 50

51 IWGDGNTDYNSALKSRVTMLKDTSKNQKSLRLSSVTAADTAVYYCARDRV 100

101 TATLYAMDYWGQGSLVTVSS 120,

the TM27 V_H sequence wherein amino acid residue 48 is replaced with isoleucine (I),

the TM27 V_H sequence wherein amino acid residues 78 and 79 are valine (V) and phenylalanine (F),

the TM27 V_H sequence wherein amino acid residues 67 to 70 having the sequence VTML are replaced with LSIS, respectively, and amino acid residue 73 is asparagine (N), and the TM27 V_H sequence in which amino acid residue 92 is an arginine (R).

2. A humanized antibody or binding protein thereof having an amino acid sequence that comprises all or a portion of CDRs substantially derived from a monoclonal antibody having a specificity for a select subpopulation of T cells expressing the Vβ8.1 variable region of the T cell antigen receptor beta chain, in conjunction with select variable framework residues also derived from said monoclonal antibody, said humanized antibody or binding protein thereof capable of binding to said select subpopulations of T cells with a binding affinity at least comparable to that exhibited by said monoclonal antibody, wherein:

the humanized antibody or binding protein thereof comprises the TM29 Vκ amino acid sequence:

1 DIQMTQSPSSLSASVGDRVTITCRASSSVNYIYWYQQTPGKAPKLLIYYT 50

51 SNLAPGVPSRFSGSGSGTDYTFTISSLQPEDIATYYCQQFTSSPFTFGQGT 100

101 KLQIT 106,

and further comprises an amino acid sequence selected from the group consisting of the TM29 V_H sequence:

1 EVQLVESGGGVVQPGRSLRLSCSSSGFTF SNFGMHWVRQAPGKGLEWVAY 50

51 ISSGSSTIYYADTLKGRFTISRDNSKNTLFLQMDSLRPEDTGVYFCARRG 100

101 EGAMDYWGQGTPVTVSS 117,

the TM29 V_H sequence wherein amino acid residues 23 and 24 are both replaced with alanine (A) residues,

the TM29 V_H sequence wherein amino acid residue 75 is replaced with proline (P), and

the TM29 V_H sequence wherein amino acid residues 92 and 93 are replaced with alanine (A) and methionine (M), respectively, and amino acid residue 95 is replaced with tyrosine (Y).

3. The humanized antibody of claim 1 or 2 which is of isotype IgG2a.

4. The humanized antibody of claim 3 wherein one or more residues in the constant domains of the Ig has been altered in order to alter the effector function of said constant domain.

5. The humanized antibody or binding portion thereof of any one of claims 1 or 2 produced by recombinant DNA technology.

6. A method for producing a humanized antibody or binding protein thereof of any one of claims 1 or 2, which method comprises:

a) isolating a first DNA sequence encoding at least one CDR substantially derived from an anti-Vβ5.2/Vβ5.3 or anti-Vb8.1 murine monoclonal antibody having a specificity for a select subpopulation of T cells;

b) cloning said first DNA sequence into a suitable vector;

c) introducing select human framework amino acids through mutagenesis, thereby forming a CDR grafted construct;

d) transforming a host cell with said construct of step c); and

e) culturing said host cell so transformed to produce said humanized antibody or binding protein thereof.

7. The method of claim 6 wherein the host cell is a CHO cell.

8. A humanized antibody or binding protein thereof in accordance with any one of claims 1 or 2 for use in therapy.

9. A pharmaceutical composition comprising a humanized antibody or binding protein thereof in accordance with any one of claims 1or 2 in combination with a pharmaceutically acceptable excipient.

10. The use of a humanized antibody or binding protein thereof according to any one of claims 1 or 2 in the manufacture of a medicament for treatment of Crohn's disease.

11. The use of a humanized antibody or binding protein thereof according to any one of claims 1 or 2 in the manufacture of a medicament for treatment of multiple sclerosis.



10 20 30 40 50
TM23 V_L DIQMTQTTSSLSASLGDRVTISCSASQGISNYLNWYQQKPDGTVKLLIYY
TM27 V_L DIQMTQSPSSLSASVGDRVTITCSASQGISNYLNWYQQTPGKAPKLLIYY
CDRI
60 70 80 90 100
TSSLHSGVPSRFSGSGSGTDYSLTISNLEPEDIATYYCQQYSKLPRTFGG
TSSLHSGVPSRFSGSGSGTDYTFTISSLQPEDIATYYCQQYSKLPRTFGQ
CDR2
CDR3
107
GTKVEIK
GTKLQIT
10 20 30 40 50
TM23 V_H QVQLKESGPGLVAPSQSLSITCTVSGFSLTAYGVNWVRQPPGKGLEWLGM
TM27L V_H QVQLQESGPGLVRPSQTLSLTCTVSGFSLTAYGVNWVRQPPGRGLEWLGM
TM27I V_H ==========AYGVN==========I=M
TM27.1V_H ==========AYGVN==========L=M
TM27.2V_H ==========AYGVN==========L=M
TM27.3V_H ==========AYGVN==========L=M
CDRI

FIG.1



60 70 80 90 100
IWGDGNTDYNSALKSRLSISKDNSKSQVFLKMNSLQTDDTARYYCARDRV
IWGDGNTDYNSALKSRVTMLKDTSKNQFSLRLSSVTAADTAVYYCARDRV
IWGDGNTDYNSALK==========DRV
IWGDGNTDYNSALK==========VF==========DRV
IWGDGNTDYNSALK==LSIS==N==========DRV
IWGDGNTDYNSALK==========R=====DRV
CDR2
110 120
TATLYAMDYWGQGTSVTVSS
TATLYAMDYWGQGSLVTVSS
TATLYAMDY==========
TATLYAMDY==========
TATLYAMDY==========
TATLYAMDY==========
CDR3

FIG.1(contd.)



10 20 30 40 50
16G8 V_L ENVLTQSPAIMSASLGEKVTMSCRASSSVNYIYWYQQKSDASPKLWIYYT
TM29 V_L DIQMTQSPSSLSASVGDRVTITCRASSSVNYIYWYQQTPGKAPKLLIYYT
CDRI
60 70 80 90 100
SNLAPGVPTRFSGSGSGNSYSLTISSMEGEDAATYYCQQFTSSPFTFGSG
SNLAPGVPSRFSGSGSGTDYTFTISSLQPEDIATYYCQQFTSSPFTFGQG
CDR2
CDR3
106
TKLEIK
TKLQIT
10 20 30 40 50
16G8 V_H DVQLVESGGGLVQPGGSRKLSCAASGFTFSNFGMHWVRQAPDKGLEWVAY
TM29 V_H EVQLVESGGGVVQPGRSLRLSCSSSGFTFSNFGMHWVRQAPGKGLEWVAY
TM29.1V_H ==========AA======NFGMH==========Y
TM29.2V_H ==========NFGMH==========Y
TM29.3V_H ==========NFGMH==========Y
CDRI

FIG.2





FIG.2(contd.)

LV 11631

FIG.3




MURINE4H11	y = 393.85 - 5.0535x	R^2 = 0.985
TM27L/NS0	y = 415.77 - 2.3218x	R^2 = 0.939
TM27L/CHO	y = 415.64 - 2.1880x	R^2 = 0.987

UG/ML	%UNLABELED	MURINE4H11	TM27L/NS0	TM27L/CHO
5.000	50.000	149.687	303.633	299.073
3.750	42.900	181.330	296.867	328.207
2.500	33.300	219.803	349.850	342.667
1.875	27.300	251.783	367.640	358.120
1.250	20.000	288.993	360.657	374.003
0.938	15.800	308.113	380.870	378.773
0.625	11.100	333.177	388.950	395.623
0.312	5.900	349.383	405.483	401.297
0.156	3.030	380.860	400.113	402.853
0.000	0.000	417.797	417.797	417.792

FIG.4

TM27 SCATCHARD ANALYSIS - HPB CELLS



	Kd (nM)
● TM27 CH0	12.3
◆ TM27 NS0	8.2
■ 4H11	4.5