

PCTWORLD INTELLECT
Inte

INTERNATIONAL APPLICATION PUBLISHED

WO 9604301A2

(51) International Patent Classification ⁶ : C07K 14/18, C12N 15/51, A61K 39/29		A2	(11) International Publication Number: WO 96/04301
			(43) International Publication Date: 15 February 1996 (15.02.96)
(21) International Application Number: PCT/US95/10035			(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).
(22) International Filing Date: 26 July 1995 (26.07.95)			
(30) Priority Data:			
08/282,959 29 July 1994 (29.07.94) US			
Not furnished 25 July 1995 (25.07.95) US			
(71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street, Emeryville, CA 94608 (US).			Published <i>Without international search report and to be republished upon receipt of that report.</i>
(72) Inventors: SELBY, Mark; 136 Galewood Circle, San Francisco, CA 94131 (US). HOUGHTON, Michael; 53 Rosemead Court, Danville, CA 94526 (US).			
(74) Agents: HARBIN, Alisa, A. et al.; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US).			

(54) Title: NOVEL HEPATITIS C E1 AND E2 TRUNCATED POLYPEPTIDES AND METHODS OF OBTAINING THE SAME**(57) Abstract**

Novel Hepatitis C E1 and E2 truncated polypeptides and complexes comprising these polypeptides, are disclosed. The polypeptides are C-terminally truncated to remove all or a portion of their membrane spanning domains. Hence, the polypeptides are capable of secretion when expressed recombinantly.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

NOVEL HEPATITIS C E1 AND E2 TRUNCATED POLYPEPTIDES
AND METHODS OF OBTAINING THE SAME

Background of the Invention

5

Technical Field

The present invention pertains generally to viral proteins. In particular, the invention relates to truncated, secreted forms of hepatitis C virus E1 and E2 proteins and the isolation and recombinant production of the same.

10

Background of the Invention

Hepatitis C Virus (HCV) is the principal cause of parenteral non-A, non-B hepatitis which is transmitted largely through blood transfusion and sexual contact. The virus is present in between 0.4 to 2.0% of blood donors. Chronic hepatitis develops in approximately 50% of infections and of these, approximately 20% of infected individuals develop liver cirrhosis which sometimes leads to hepatocellular carcinoma. Accordingly, the study and control of the disease is of medical importance.

15

The viral genomic sequence of HCV is known, as are methods for obtaining the sequence. See, e.g., International Publication Nos. WO 89/04669; WO 90/11089; and WO 90/14436. In particular, HCV has a 9.5 kb positive-sense, single-stranded RNA genome and is a member of the Flaviridae family of viruses. Currently, there are 6 distinct, but related genotypes of HCV based on phylogenetic analyses (Simmonds et al., *J. Gen. Virol.* (1993) 74:2391-2399). The virus encodes a single polyprotein having more than 3000 amino acid residues (Choo et al., *Science* (1989) 244:359-362; Choo et al., *Proc. Natl. Acad. Sci. USA* (1991) 88:2451-2455; Han et al., *Proc. Natl. Acad. Sci. USA* (1991) 88:1711-1715). The polyprotein is processed co- and post-translationally into both structural and non-structural (NS) proteins.

20

In particular, there are three putative structural proteins, consisting of the N-terminal nucleocapsid protein (termed "core") and two envelope glycoproteins, "E1"

(also known as E) and "E2" (also known as E2/NS1). (See, Houghton et al., *Hepatology* (1991) 14:381-388, for a discussion of HCV proteins, including E1 and E2.) E1 is detected as a 32-35 kDa species and is converted into a single endo H-sensitive band of approximately 18 kDa. By contrast, E2 displays a complex pattern upon
5 immunoprecipitation consistent with the generation of multiple species (Grakoui et al., *J. Virol.* (1993) 67:1385-1395; Tomei et al., *J. Virol.* (1993) 67:4017-4026.). The HCV envelope glycoproteins E1 and E2 form a stable complex that is co-immunoprecipitable (Grakoui et al., *J. Virol.* (1993) 67:1385-1395; Lanford et al., *Virology* (1993) 197:225-235; Ralston et al., *J. Virol.* (1993) 67:6753-6761). The HCV
10 E1 and E2 glycoproteins are of considerable interest because they have been shown to be protective in primate studies. (Choo et al., *Proc. Natl. Acad. Sci. USA* (1994) 91:1294-1298).

The envelope of the HCV virion remains uncharacterized. Thus, expression studies using recombinant cDNA templates are the only means currently available to
15 study envelope biosynthesis. E1 and E2 are retained within cells and lack complex carbohydrate when expressed stably or in a transient Vaccinia virus system (Spaete et al., *Virology* (1992) 188:819-830; Ralston et al., *J. Virol.* (1993) 67:6753-6761). Since the E1 and E2 proteins are normally membrane-bound in these expression systems, it would be desirable to produce secreted forms to facilitate purification of the proteins for
20 further use.

It has been found that removal of the transmembrane domain of the viral cell surface glycoproteins of influenza virus (Sveda, et al., *Cell* (1982) 30:649-656; Gething and Sambrook, *Nature* (1982) 300:598-603) and vesicular stomatitis virus (Rose and Bergmann, *Cell* (1982) 30:753-762) results in secretion of the truncated glycoprotein
25 from mammalian host cells. See also EPO Publication No. 139,417. Similarly, truncated cytomegalovirus gH is secreted when expressed in baculovirus cells (International Publication No. WO 92/02628, published 20 February 1992). A C-terminally truncated HCV E2 molecule, capable of secretion from mammalian cells, has been described. Spaete et al., *Virology* (1992) 188:819-830. However, the
30 transmembrane anchor region of E1 has not heretofore been elucidated and hence the production of truncated forms of HCV E1 for secretion has not been previously

disclosed. Furthermore, complexes of truncated, secreted E1 and E2 polypeptides have not been previously described.

Summary of the Invention

5 The present invention is based on the elucidation of sequences of E1 and E2 important for anchoring the proteins to the endoplasmic reticulum (ER) and for co-precipitation of E2 with E1. Thus, the elimination of these sequences serves to produce truncated forms of the glycoproteins which are secreted rather than retained in the ER membrane. Furthermore, truncation facilitates purification of the E2 protein
10 without associated E1, and vice versa. The truncated E1 and E2 proteins, when expressed together or combined after expression, are capable of forming a complex.

 Accordingly, in one embodiment, the subject invention is directed to an HCV E1 polypeptide, lacking all or a portion of its membrane spanning domain such that the polypeptide is capable of secretion into growth medium when expressed recombinantly
15 in a host cell. In preferred embodiments, the HCV E1 polypeptide lacks at least a portion of its C-terminus beginning at or near about amino acid 370 or about amino acid 360, numbered with reference to the HCV1 E1 amino acid sequence.

 In another embodiment, the invention is directed to an HCV E2 polypeptide lacking at least a portion of its membrane spanning domain such that the polypeptide is
20 capable of secretion into growth medium when expressed recombinantly in a host cell, wherein the polypeptide lacks at least a portion of its C-terminus beginning at or near about amino acid 730 but not extending beyond about amino acid 699, numbered with reference to the HCV1 E2 amino acid sequence. In particularly preferred embodiments, the HCV E2 polypeptide lacks at least a portion of its C-terminus beginning at about
25 amino acid 725, numbered with reference to the HCV1 E2 amino acid sequence.

 Other embodiments of the subject invention pertain to polynucleotides encoding the above polypeptides, vectors comprising these polynucleotides, host cells transformed with the vectors and methods of recombinantly producing the polypeptides.

 In yet another embodiment, the subject invention is directed to a secreted
30 E1/secreted E2 complex comprising:

(a) an HCV E1 polypeptide, lacking all or a portion of its membrane spanning domain such that the E1 polypeptide is capable of secretion into growth medium when expressed recombinantly in a host cell; and

5 (b) an HCV E2 polypeptide, lacking all or a portion of its membrane spanning domain such that said E2 polypeptide is capable of secretion into growth medium when expressed recombinantly in a host cell.

In preferred embodiments, the secreted E1/secreted E2 complex includes an E1 polypeptide which lacks at least a portion of its C-terminus beginning at about amino acid 360 or 370, numbered with reference to the HCV1 E1 amino acid sequence, and
10 said E2 polypeptide lacks at least a portion of its C-terminus beginning at about amino acid 725 or 730, numbered with reference to the HCV1 E2 amino acid sequence.

In still further embodiments, the subject invention is directed to vaccine compositions comprising the truncated HCV E1 and/or E2 polypeptides and/or complexes of the E1 and E2 polypeptides.

15 These and other embodiments of the present invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

Brief Description of the Figures

Figure 1 is a schematic of the E1 region of HCV1.

20 Figure 2 shows the nucleotide sequence and corresponding amino acid sequence for HCV1 E1, including the N-terminal signal sequence and the C-terminal membrane anchor domain.

Figure 3 is a schematic of the E2 region of HCV1.

Figures 4A-4C show the nucleotide sequence and the corresponding amino acid
25 sequence for the HCV1 E2/NS2 region, including the N-terminal signal sequence for E2 and the C-terminal membrane anchor domain for E2.

Figure 5 depicts the HCV E1 cDNA templates used for transfection in the Examples. The core through NS2 region is shown on the top and is drawn to scale; the distal NS3 through NS5 is not drawn to scale. The E1 region has been expanded to
30 better display the templates used. The numbers to the right refer to the amino acid endpoint used in each template.

Figure 6 depicts the HCV E2 cDNA templates used for transfection in the Examples. The core through NS2 region is shown on the top and is drawn to scale; the distal NS3 through NS5 is not drawn to scale. The E2/NS2 region has been expanded to better display the templates used. The column to the left refers to the amino acid endpoint used in each template.

Figure 7 depicts plasmid pMHE2-715, containing a gene encoding for a truncated E2 protein having amino acids 383-715 of HCV1. Also present in the vector is the SV40 origin of replication, the mouse cytomegalovirus immediate early promoter (MCMV ie), a tpa leader, the SV40 polyadenylation signal and DHFR cDNA for selection.

Detailed Description of the Invention

The practice of the present invention will employ, unless otherwise indicated, conventional methods of virology, microbiology, molecular biology and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, et al. *Molecular Cloning: A Laboratory Manual* (2nd Edition, 1989); *DNA Cloning: A Practical Approach*, vol. I & II (D. Glover, ed.); *Oligonucleotide Synthesis* (N. Gait, ed., 1984); *Nucleic Acid Hybridization* (B. Hames & S. Higgins, eds., 1985); *Transcription and Translation* (B. Hames & S. Higgins, eds., 1984); *Animal Cell Culture* (R. Freshney, ed., 1986); Perbal, *A Practical Guide to Molecular Cloning* (1984); *Fundamental Virology*, 2nd Edition, vol. I & II (B.N. Fields and D.M. Knipe, eds.)

All publications, patents and patent applications cited herein, whether *supra* or *infra*, are hereby incorporated by reference in their entirety.

As used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural references unless the content clearly dictates otherwise.

I. Definitions

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

By an "E1 polypeptide" is meant a molecule derived from an HCV E1 region. Such a molecule can be physically derived from the region or produced recombinantly or synthetically, based on the known sequence. The mature E1 region of HCV1 begins at approximately amino acid 192 of the polyprotein and continues to approximately amino acid 383 (see Figures 1 and 2). Amino acids at around 173 through approximately 191 serve as a signal sequence for E1. Thus, by an "E1 polypeptide" is meant either a precursor E1 protein, including the signal sequence, or a mature E1 polypeptide which lacks this sequence, or even an E1 polypeptide with a heterologous signal sequence. Furthermore, as elucidated herein, the E1 polypeptide includes a C-terminal membrane anchor sequence which occurs at approximately amino acid positions 360-383.

By an "E2 polypeptide" is meant a molecule derived from an HCV E2 region. Such a molecule can be physically derived from the region or produced recombinantly or synthetically, based on the known sequence. The mature E2 region of HCV1 is believed to begin at approximately amino acid 384-385 (see Figures 3 and 4A-4C). A signal peptide begins at approximately amino acid 364 of the polyprotein. Thus, by an "E2 polypeptide" is meant either a precursor E2 protein, including the signal sequence, or a mature E2 polypeptide which lacks this sequence, or even an E1 polypeptide with a heterologous signal sequence. Furthermore, as elucidated herein, the E2 polypeptide includes a C-terminal membrane anchor sequence which occurs at approximately amino acid positions 715-730 and may extend as far as approximately amino acid residue 746 (see, Lin et al., *J. Virol.* (1994) 68:5063-5073).

Representative E1 and E2 regions from HCV1 are shown in Figures 2 and 4A-4C, respectively. For purposes of the present invention, the E1 and E2 regions are defined with respect to the amino acid number of the polyprotein encoded by the genome of HCV1, with the initiator methionine being designated position 1. However, it should be noted that the term an "E1 polypeptide" or an "E2 polypeptide" as used herein is not limited to the HCV1 sequence. In this regard, the corresponding E1 or E2 regions in another HCV isolate can be readily determined by aligning sequences from the two isolates in a manner that brings the sequences into maximum alignment. This can be performed with any of a number of computer software packages, such as ALIGN

1.0, available from the University of Virginia, Department of Biochemistry (Attn: Dr. William R. Pearson). See, Pearson et al., *Proc. Natl. Acad. Sci. USA* (1988) 85:2444-2448.

Furthermore, an "E1 polypeptide" or an "E2 polypeptide" as defined herein is
5 not limited to a polypeptide having the exact sequence depicted in the Figures. Indeed, the HCV genome is in a state of constant flux and contains several variable domains which exhibit relatively high degrees of variability between isolates. As will become evident herein, all that is important is that the region which serves to anchor the polypeptide to the endoplasmic reticulum be identified such that the polypeptide can be
10 modified to remove all or part of this sequence for secretion. It is readily apparent that the terms encompass E1 and E2 polypeptides from any of the various HCV isolates including isolates having any of the 6 genotypes of HCV described in Simmonds et al., *J. Gen. Virol.* (1993) 74:2391-2399). Furthermore, the term encompasses any such E1 or E2 protein regardless of the method of production, including those proteins
15 recombinantly and synthetically produced.

Additionally, the terms "E1 polypeptide" and "E2 polypeptide" encompass proteins which include additional modifications to the native sequence, such as additional internal deletions, additions and substitutions (generally conservative in nature). These modifications may be deliberate, as through site-directed mutagenesis, or
20 may be accidental, such as through naturally occurring mutational events. All of these modifications are encompassed in the present invention so long as the modified E1 and E2 polypeptides function for their intended purpose. Thus, for example, if the E1 and/or E2 polypeptides are to be used in vaccine compositions, the modifications must be such that immunological activity (i.e., the ability to elicit an antibody response to the
25 polypeptide) is not lost. Similarly, if the polypeptides are to be used for diagnostic purposes, such capability must be retained.

An E1 or E2 polypeptide "lacking all or a portion of its membrane spanning domain" is an E1 or E2 polypeptide, respectively, as defined above, which has been C-terminally truncated to delete all or a part of the membrane anchor sequence which
30 functions to associate the polypeptide to the endoplasmic reticulum. Such a polypeptide is therefore capable of secretion into growth medium in which an organism expressing

the protein is cultured. The truncated polypeptide need only lack as much of the membrane anchor sequence as necessary in order to effect secretion. Secretion into growth media is readily determined using a number of detection techniques, including, e.g., polyacrylamide gel electrophoresis and the like and immunological techniques such as immunoprecipitation assays as described in the examples. With E1, generally polypeptides terminating with about amino acid position 370 and higher (based on the numbering of HCV1 E1) will be retained by the ER and hence not secreted into growth media. With E2, polypeptides terminating with about amino acid position 731 and higher (also based on the numbering of the HCV1 E2 sequence) will be retained by the ER and not secreted. It should be noted that these amino acid positions are not absolute and may vary to some degree.

Although not all possible C-terminal truncations have been exemplified herein, it is to be understood that intervening truncations, such as e.g., E1 polypeptides ending in amino acids 351, 352, 353 and so on, or E2 polypeptides ending in for example amino acids 716, 717, 718 and so on, are also encompassed by the present invention. Hence, all E1 polypeptides, terminating at about amino acids 369 and lower, and all E2 polypeptides, terminating at about amino acids 730 and lower, which are capable of secretion into growth medium when expressed recombinantly, are intended to be captured herein.

Furthermore, the C-terminal truncation can extend beyond the transmembrane spanning domain towards the N-terminus. Thus, for example, E1 truncations occurring at positions lower than, e.g., 360 and E2 truncations occurring at positions lower than, e.g., 715, are also encompassed by the present invention. All that is necessary is that the truncated E1 and E2 polypeptides be secreted and remain functional for their intended purpose. However, particularly preferred E2 constructs will be those with C-terminal truncations that do not extend beyond amino acid position 699.

A "secreted E1/secreted E2 complex" refers to a complex of the E1 and E2 proteins, each of which lacks all or a portion of the membrane spanning domain, as described above. The mode of association of E1 and E2 in such a complex is immaterial. Indeed, such a complex may form spontaneously simply by mixing secreted E1 and E2 proteins which have been produced individually. Similarly, when co-

expressed, the secreted E1 and secreted E2 proteins can form a complex spontaneously in the media. Formation of a "secreted E1/secreted E2 complex" is readily determined using standard protein detection techniques such as polyacrylamide gel electrophoresis and immunological techniques such as immunoprecipitation.

5 Two polynucleotides or protein molecules are "substantially homologous" when at least about 40-50%, preferably at least about 70-80%, and most preferably at least about 85-95%, of the nucleotides or amino acids from the molecules match over a defined length of the molecule. As used herein, substantially homologous also refers to molecules having sequences which show identity to the specified nucleic acid or protein
10 molecule. Nucleic acid molecules that are substantially homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions, as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook et al., *supra*; *DNA Cloning*, vols I & II, *supra*; *Nucleic Acid Hybridization*, *supra*.

15 An "isolated" protein or polypeptide is a protein which is separate and discrete from a whole organism with which the protein is normally associated in nature. Thus, a protein contained in a cell free extract would constitute an "isolated" protein, as would a protein synthetically or recombinantly produced. Likewise, an "isolated" polynucleotide is a nucleic acid molecule separate and discrete from the whole organism with which the
20 sequence is found in nature; or a sequence devoid, in whole or part, of sequences normally associated with it in nature; or a sequence, as it exists in nature, but having heterologous sequences (as defined below) in association therewith.

A "coding sequence" or a sequence which "encodes" a selected protein, is a nucleic acid sequence which is transcribed (in the case of DNA) and translated (in the
25 case of mRNA) into a polypeptide *in vitro* or *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A coding sequence can include, but is not limited to cDNA from viral nucleotide sequences as well as synthetic DNA sequences. A transcription
30 termination sequence may be located 3' to the coding sequence.

A "polynucleotide" can include, but is not limited to, viral sequences, procaryotic sequences, viral RNA, eucaryotic mRNA, cDNA from eucaryotic mRNA, genomic DNA sequences from eucaryotic (e.g., mammalian) DNA, and even synthetic DNA sequences. The term also captures sequences that include any of the known base
5 analogs of DNA and RNA such as, but not limited to 4-acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinylcytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl)uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyluracil, dihydrouracil, inosine, N6-isopentenyladenine, 1-methyladenine, 1-methylpseudouracil, 1-methylguanine, 1-methylinosine, 2,2-
10 dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine,
15 pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

"Control elements" refers collectively to promoter sequences, ribosome binding sites, polyadenylation signals, transcription termination sequences, upstream regulatory
20 domains, enhancers, and the like, which collectively provide for the transcription and translation of a coding sequence in a host cell. Not all of these control elements need always be present in a recombinant vector so long as the desired gene is capable of being transcribed and translated.

A control element "directs the transcription" of a coding sequence in a cell when
25 RNA polymerase will bind the promoter sequence and transcribe the coding sequence into mRNA, which is then translated into the polypeptide encoded by the coding sequence.

"Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, control
30 elements operably linked to a coding sequence are capable of effecting the expression of the coding sequence when RNA polymerase is present. The control elements need not

be contiguous with the coding sequence, so long as they functions to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between, e.g., a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

"Recombinant" as used herein to describe a nucleic acid molecule means a polynucleotide of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation: (1) is not associated with all or a portion of the polynucleotide with which it is associated in nature; and/or (2) is linked to a polynucleotide other than that to which it is linked in nature. The term "recombinant" as used with respect to a protein or polypeptide means a polypeptide produced by expression of a recombinant polynucleotide. "Recombinant host cells," "host cells," "cells," "cell lines," "cell cultures," and other such terms denoting procaryotic microorganisms or eucaryotic cell lines cultured as unicellular entities, are used interchangeably, and refer to cells which can be, or have been, used as recipients for recombinant vectors or other transfer DNA, and include the progeny of the original cell which has been transfected. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement to the original parent, due to accidental or deliberate mutation. Progeny of the parental cell which are sufficiently similar to the parent to be characterized by the relevant property, such as the presence of a nucleotide sequence encoding a desired peptide, are included in the progeny intended by this definition, and are covered by the above terms.

A "vector" is a replicon in which a heterologous polynucleotide segment is attached, so as to bring about the replication and/or expression of the attached segment, such as a plasmid, transposon, phage, etc.

By "vertebrate subject" is meant any member of the subphylum cordata, including, without limitation, humans and other primates, including non-human primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs; birds, including domestic,

wild and game birds such as chickens, turkeys and other gallinaceous birds, ducks, geese, and the like. The term does not denote a particular age. Thus, both adult and newborn individuals are intended to be covered. The system described above is intended for use in any of the above vertebrate species, since the immune systems of all
5 of these vertebrates operate similarly.

II. Modes of Carrying Out the Invention

The present invention is based on the discovery of novel E1 and E2 polypeptides which are C-terminally truncated such that they are capable of secretion into growth
10 medium when produced recombinantly in a host cell. The secreted polypeptides are also surprisingly able to form complexes with one another. This interaction is unexpected since, as shown herein, the ability of E1 and E2 to co-precipitate is lost upon elimination of the membrane spanning domain.

In particular, analysis of transient transfections of serially extended templates
15 covering the E2/NS2 region provided evidence for three E2 species with distinct C-termini. One form was E2 terminating at amino acid 729 while the larger two species represented fusions with the downstream NS2A and NS2A/NS2B proteins terminating at amino acids 809 and 1026, respectively. Using the same E2 templates, a region of E2 important for co-immunoprecipitation of E1 has been defined which also
20 prevents E2 secretion. Similarly, a membrane spanning domain of E1 was identified.

More specifically, a membrane spanning domain which presumably serves to anchor E1 to the ER membrane, has been identified at about amino acid positions 360-383. A number of C-terminally truncated E1 polypeptides, lacking portions of this membrane spanning domain, have been constructed. (See Figure 5). It has been found
25 that E1 polypeptides, ending in amino acid positions 370 and higher, are not secreted into growth media when recombinantly expressed.

Similarly, a series of E2 molecules which include C-terminal truncations as depicted in Figure 6, have been recombinantly expressed and culture media from transformed cells tested for the presence of the E2 polypeptides to determine which
30 truncated constructs are capable of secretion. It has been found that molecules ending at amino acid positions higher than 730, are not secreted into growth media and

presumably are retained in the ER membrane. (It should be noted that a small amount of secretion is observed with constructs terminating at amino acid position 730). An inverse relationship between E2 secretion and its association with E1 has also been found. Hence, the E2 polypeptides which are secretable do not co-precipitate with E1
5 from lysates whereas those that are not secreted, do.

However, surprisingly, when secreted forms of E1 and E2 polypeptides are co-expressed, the secreted polypeptides are able to form a complex, detectable using antibodies to either of E1 or E2. Such complex formation is significant as it demonstrates that the regions of E1 and E2 which are important for their interaction are
10 retained despite the elimination of C-terminal membrane anchors.

These discoveries provide efficient methods for purifying E1, E2 and secreted E1/E2 complexes, for future use. In particular, secreted proteins are more easily purified than intracellularly expressed proteins. Similarly, since as described above, the native E1 and E2 proteins are known to form a complex, the invention herein described
15 provides a method for obtaining either of E1 or E2 free from the other protein. Additionally, should an E1/E2 complex be desired, the secreted proteins can either be co-expressed or mixed together (either in culture media or in purified or semipurified form), for spontaneous complex formation.

The truncated E1 and E2 polypeptides can be produced using a variety of
20 techniques. For example, the polypeptides can be generated using recombinant techniques, well known in the art. In this regard, oligonucleotide probes can be devised based on the known sequences of the HCV genome and used to probe genomic or cDNA libraries for E1 and E2 genes. The genes can then be further isolated using standard techniques and, e.g., restriction enzymes employed to truncate the gene at
25 desired portions of the full-length sequence. Similarly, the E1 and E2 genes can be isolated directly from cells and tissues containing the same, using known techniques, such as phenol extraction and the sequence further manipulated to produce the desired truncations. *See, e.g.,* Sambrook et al., *supra*, for a description of techniques used to obtain and isolate DNA. Finally, the genes encoding the truncated E1 and E2
30 polypeptides can be produced synthetically, based on the known sequences. The nucleotide sequence can be designed with the appropriate codons for the particular

amino acid sequence desired. In general, one will select preferred codons for the intended host in which the sequence will be expressed. The complete sequence is generally assembled from overlapping oligonucleotides prepared by standard methods and assembled into a complete coding sequence. See, e.g., Edge (1981) *Nature* 292:756; Nambair *et al.* (1984) *Science* 223:1299; Jay *et al.* (1984) *J. Biol. Chem.* 259:6311.

Once coding sequences for the desired proteins have been isolated or synthesized, they can be cloned into any suitable vector or replicon for expression. Numerous cloning vectors are known to those of skill in the art, and the selection of an appropriate cloning vector is a matter of choice. Examples of recombinant DNA vectors for cloning and host cells which they can transform include the bacteriophage λ (*E. coli*), pBR322 (*E. coli*), pACYC177 (*E. coli*), pKT230 (gram-negative bacteria), pGV1106 (gram-negative bacteria), pLAFR1 (gram-negative bacteria), pME290 (non-*E. coli* gram-negative bacteria), pHV14 (*E. coli* and *Bacillus subtilis*), pBD9 (*Bacillus*), pIJ61 (*Streptomyces*), pUC6 (*Streptomyces*), YIp5 (*Saccharomyces*), YCp19 (*Saccharomyces*) and bovine papilloma virus (mammalian cells). See, generally, *DNA Cloning*: Vols. I & II, *supra*; Sambrook *et al.*, *supra*; B. Perbal, *supra*.

Insect cell expression systems, such as baculovirus systems, can also be used and are known to those of skill in the art and described in, e.g., Summers and Smith, *Texas Agricultural Experiment Station Bulletin No. 1555* (1987). Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, *inter alia*, Invitrogen, San Diego CA ("MaxBac" kit).

Viral systems, such as a vaccinia based infection/transfection system, as described in Tomei *et al.*, *J. Virol.* (1993) 67:4017-4026 and Selby *et al.*, *J. Gen. Virol.* (1993) 74:1103-1113, will also find use with the present invention. In this system, cells are first transfected *in vitro* with a vaccinia virus recombinant that encodes the bacteriophage T7 RNA polymerase. This polymerase displays exquisite specificity in that it only transcribes templates bearing T7 promoters. Following infection, cells are transfected with the DNA of interest, driven by a T7 promoter. The polymerase expressed in the cytoplasm from the vaccinia virus recombinant transcribes the transfected DNA into RNA which is then translated into protein by the host translational

machinery. The method provides for high level, transient, cytoplasmic production of large quantities of RNA and its translation product(s).

The gene can be placed under the control of a promoter, ribosome binding site (for bacterial expression) and, optionally, an operator (collectively referred to herein as "control" elements), so that the DNA sequence encoding the desired E1 or E2 polypeptide is transcribed into RNA in the host cell transformed by a vector containing this expression construction. The coding sequence may or may not contain a signal peptide or leader sequence. With the present invention, both the naturally occurring signal peptides or heterologous sequences can be used. Leader sequences can be removed by the host in post-translational processing. *See, e.g.*, U.S. Patent Nos. 4,431,739; 4,425,437; 4,338,397.

Other regulatory sequences may also be desirable which allow for regulation of expression of the protein sequences relative to the growth of the host cell. Such regulatory sequences are known to those of skill in the art, and examples include those which cause the expression of a gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Other types of regulatory elements may also be present in the vector, for example, enhancer sequences.

The control sequences and other regulatory sequences may be ligated to the coding sequence prior to insertion into a vector, such as the cloning vectors described above. Alternatively, the coding sequence can be cloned directly into an expression vector which already contains the control sequences and an appropriate restriction site.

In some cases it may be necessary to modify the coding sequence so that it may be attached to the control sequences with the appropriate orientation; *i.e.*, to maintain the proper reading frame. It may also be desirable to produce mutants or analogs of the E1 or E2 protein. Mutants or analogs may be prepared by the deletion of a portion of the sequence encoding the protein, by insertion of a sequence, and/or by substitution of one or more nucleotides within the sequence. Techniques for modifying nucleotide sequences, such as site-directed mutagenesis, are well known to those skilled in the art. *See, e.g.*, Sambrook *et al.*, *supra*; *DNA Cloning*, Vols. I and II, *supra*; *Nucleic Acid Hybridization*, *supra*.

The expression vector is then used to transform an appropriate host cell. A number of mammalian cell lines are known in the art and include immortalized cell lines available from the American Type Culture Collection (ATCC), such as, but not limited to, Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, 5 monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), Madin-Darby bovine kidney ("MDBK") cells, as well as others. Similarly, bacterial hosts such as *E. coli*, *Bacillus subtilis*, and *Streptococcus spp.*, will find use with the present expression constructs. Yeast hosts useful in the present invention include *inter alia*, *Saccharomyces cerevisiae*, *Candida albicans*, *Candida maltosa*, *Hansenula* 10 *polymorpha*, *Kluyveromyces fragilis*, *Kluyveromyces lactis*, *Pichia guilliermondii*, *Pichia pastoris*, *Schizosaccharomyces pombe* and *Yarrowia lipolytica*. Insect cells for use with baculovirus expression vectors include, *inter alia*, *Aedes aegypti*, *Autographa californica*, *Bombyx mori*, *Drosophila melanogaster*, *Spodoptera frugiperda*, and *Trichoplusia ni*. Depending on the expression system and host selected, the proteins 15 of the present invention are produced by growing host cells transformed by an expression vector described above under conditions whereby the protein of interest is expressed. The protein is then isolated from the host cells and purified. Since the present invention provides for secretion of the E1 and E2 polypeptides, the proteins can be purified directly from the media. The selection of the appropriate growth conditions 20 and recovery methods are within the skill of the art.

The E1 and E2 polypeptides of the present invention can also be produced using conventional methods of protein synthesis, based on the known amino acid sequences. In general, these methods employ the sequential addition of one or more amino acids to a growing peptide chain. Normally, either the amino or carboxyl group of the first 25 amino acid is protected by a suitable protecting group. The protected or derivatized amino acid can then be either attached to an inert solid support or utilized in solution by adding the next amino acid in the sequence having the complementary (amino or carboxyl) group suitably protected, under conditions that allow for the formation of an amide linkage. The protecting group is then removed from the newly added amino acid 30 residue and the next amino acid (suitably protected) is then added, and so forth. After the desired amino acids have been linked in the proper sequence, any remaining

protecting groups (and any solid support, if solid phase synthesis techniques are used) are removed sequentially or concurrently, to render the final polypeptide. By simple modification of this general procedure, it is possible to add more than one amino acid at a time to a growing chain, for example, by coupling (under conditions which do not
5 racemize chiral centers) a protected tripeptide with a properly protected dipeptide to form, after deprotection, a pentapeptide. *See, e.g.,* J. M. Stewart and J. D. Young, *Solid Phase Peptide Synthesis*, 2nd Ed., Pierce Chemical Co., Rockford, IL (1984) and G. Barany and R. B. Merrifield, *The Peptides: Analysis, Synthesis, Biology*, editors E. Gross and J. Meienhofer, Vol. 2, Academic Press, New York, (1980), pp. 3-254, for
10 solid phase peptide synthesis techniques; and M. Bodansky, *Principles of Peptide Synthesis*, Springer-Verlag, Berlin (1984) and E. Gross and J. Meienhofer, Eds., *The Peptides: Analysis, Synthesis, Biology, supra*, Vol. 1, for classical solution synthesis.

As explained above, the present invention also provides a method for producing secreted E1/secreted E2 complexes. Such complexes are readily produced by e.g., co-
15 transfecting host cells with constructs encoding for the E1 and E2 truncated proteins. Co-transfection can be accomplished either in *trans* or *cis*, i.e., by using separate vectors or by using a single vector which bears both of the E1 and E2 genes. If done using a single vector, both genes can be driven by a single set of control elements or, alternatively, the genes can be present on the vector in individual expression cassettes,
20 driven by individual control elements. Following expression, the secreted E1 and E2 proteins will spontaneously associate. Alternatively, the complexes can be formed by mixing the individual proteins together which have been produced separately, either in purified or semi-purified form, or even by mixing culture media in which host cells expressing the proteins, have been cultured.

25 The novel, secreted E1 and E2 polypeptides of the present invention, complexes thereof, or the polynucleotides coding therefor, can be used for a number of diagnostic and therapeutic purposes. For example, the proteins and polynucleotides can be used in a variety of assays, to determine the presence of E1 and E2 proteins in a biological sample to aid in the diagnosis of HCV disease. The E1 and E2 polypeptides and
30 polynucleotides encoding the polypeptides can also be used in vaccine compositions, individually or in combination, in e.g., prophylactic (i.e., to prevent infection) or

therapeutic (to treat HCV following infection) vaccines. Indeed, such secreted envelope glycoproteins are particularly useful in nucleic acid immunization where secretable molecules may be more effective than the corresponding intracellular proteins in generating an immune response. The vaccines can comprise mixtures of one or more
5 of the E1 and E2 proteins (or nucleotide sequences encoding the proteins), such as E1 and E2 proteins derived from more than one viral isolate. The vaccine may also be administered in conjunction with other antigens and immunoregulatory agents, for example, immunoglobulins, cytokines, lymphokines, and chemokines, including but not limited to IL-2, modified IL-2 (cys125→ser125), GM-CSF, IL-12, γ -interferon, IP-10,
10 MIP1 β and RANTES.

The vaccines will generally include one or more "pharmaceutically acceptable excipients or vehicles" such as water, saline, glycerol, ethanol, etc. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles.

15 A carrier is optionally present which is a molecule that does not itself induce the production of antibodies harmful to the individual receiving the composition. Suitable carriers are typically large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycollic acids, polymeric amino acids, amino acid copolymers, lipid aggregates (such as oil droplets or liposomes), and inactive virus
20 particles. Such carriers are well known to those of ordinary skill in the art. Furthermore, the HCV polypeptide may be conjugated to a bacterial toxoid, such as toxoid from diphtheria, tetanus, cholera, etc.

Adjuvants may also be used to enhance the effectiveness of the vaccines. Such adjuvants include, but are not limited to: (1) aluminum salts (alum), such as aluminum
25 hydroxide, aluminum phosphate, aluminum sulfate, etc.; (2) oil-in-water emulsion formulations (with or without other specific immunostimulating agents such as muramyl peptides (see below) or bacterial cell wall components), such as for example (a) MF59 (International Publication No. WO 90/14837), containing 5% Squalene, 0.5% Tween 80, and 0.5% Span 85 (optionally containing various amounts of MTP-PE (see below),
30 although not required) formulated into submicron particles using a microfluidizer such as Model 110Y microfluidizer (Microfluidics, Newton, MA), (b) SAF, containing 10%

Squalane, 0.4% Tween 80, 5% pluronic-blocked polymer L121, and thr-MDP (see below) either microfluidized into a submicron emulsion or vortexed to generate a larger particle size emulsion, and (c) Ribit[™] adjuvant system (RAS), (Ribi Immunochem, Hamilton, MT) containing 2% Squalene, 0.2% Tween 80, and one or more bacterial cell wall components from the group consisting of monophosphoryl lipid A (MPL), trehalose dimycolate (TDM), and cell wall skeleton (CWS), preferably MPL + CWS (Detox[™]); (3) saponin adjuvants, such as Stimulon[™] (Cambridge Bioscience, Worcester, MA) may be used or particle generated therefrom such as ISCOMs (immunostimulating complexes); (4) Complete Freund's Adjuvant (CFA) and Incomplete Freund's Adjuvant (IFA); (5) cytokines, such as interleukins (IL-1, IL-2, etc.), macrophage colony stimulating factor (M-CSF), tumor necrosis factor (TNF), etc.; and (6) other substances that act as immunostimulating agents to enhance the effectiveness of the composition. Alum and MF59 are preferred.

Muramyl peptides include, but are not limited to, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-normuramyl-L-alanyl-D-isoglutamate (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE), etc.

Typically, the vaccine compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared. The preparation also may be emulsified or encapsulated in liposomes for enhanced adjuvant effect, as discussed above.

The vaccines will comprise a therapeutically effective amount of the E1 and/or E2 truncated proteins, or complexes of the proteins, or nucleotide sequences encoding the same, and any other of the above-mentioned components, as needed. By "therapeutically effective amount" is meant an amount of an E1 and/or E2 truncated protein which will induce an immunological response in the individual to which it is administered. Such a response will generally result in the development in the subject of a secretory, cellular and/or antibody-mediated immune response to the vaccine. Usually, such a response includes but is not limited to one or more of the following effects; the production of antibodies from any of the immunological classes, such as

immunoglobulins A, D, E, G or M; the proliferation of B and T lymphocytes; the provision of activation, growth and differentiation signals to immunological cells; expansion of helper T cell, suppressor T cell, and/or cytotoxic T cell and/or $\gamma\delta$ T cell populations.

5 Preferably, the effective amount is sufficient to bring about treatment or prevention of disease symptoms. The exact amount necessary will vary depending on the subject being treated; the age and general condition of the individual to be treated; the capacity of the individual's immune system to synthesize antibodies; the degree of protection desired; the severity of the condition being treated; the particular HCV
10 polypeptide selected and its mode of administration, among other factors. An appropriate effective amount can be readily determined by one of skill in the art. A "therapeutically effective amount" will fall in a relatively broad range that can be determined through routine trials.

 Once formulated, the vaccines are conventionally administered parenterally, e.g.,
15 by injection, either subcutaneously or intramuscularly. Additional formulations suitable for other modes of administration include oral and pulmonary formulations, suppositories, and transdermal applications. Dosage treatment may be a single dose schedule or a multiple dose schedule. The vaccine may be administered in conjunction with other immunoregulatory agents.

20 As explained above, vaccines containing polynucleotides encoding for the truncated, secreted proteins can be used for nucleic acid immunization. Such a method comprises the introduction of a polynucleotide encoding one or more of the E1 and/or E2 polypeptides into a host cell, for the *in vivo* expression of the proteins. The polynucleotide can be introduced directly into the recipient subject, such as by
25 injection, inhalation or the like, or can be introduced *ex vivo*, into cells which have been removed from the host. In the latter case, the transformed cells are reintroduced into the subject where an immune response can be mounted against the protein encoded by the polynucleotide. Methods of nucleic acid immunization are known in the art and disclosed in e.g., International Publication No. WO 93/14778 (published 5 August
30 1993); International Publication No. WO 90/11092 (published 4 October 1990); Wang et

al. *Proc. Natl. Acad. Sci. USA* (1993) 90:4156-4160; Tang et al. *Nature* (1992) 356:152-154; and Ulmer et al. *Science* (1993) 259:1745-1749.

Generally, the polynucleotide is administered as a vector which has been encapsulated in a liposome and formulated into a vaccine composition as described above.

5

III. Experimental

Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

10 Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

Materials and Methods

15

Cells and transient expression

BSC40 cells and chimpanzee fibroblasts F503 (Perot et al., *J. Gen. Virol.* (1992) 73:3281-3284) were used in infection/transfection experiments as previously described (Selby et al., *J. Gen. Virol.* (1993) 74:1103-1113). Briefly, subconfluent monolayers of
20 cells in 60 mm dishes were infected with VV_{T7} (moi = 10) in serum free DME. After 30-60 minutes, the inoculum was removed and replaced with the appropriate cDNA templates cloned into the pTM1 vector (Elroy-Stein and Moss, *Proc. Natl. Acad. Sci. USA* (1990) 87:6743-6747), as described below. Vector DNA was complexed with either Lipofectin (BRL) or Lipofectamine (BRL). After transfection for 2.5-3 hours, the
25 DNA was removed and the cells were starved for 30 minutes in met/cys-deficient DME. Approximately 100-200 μ Ci of ³⁵S-Express Label (NEN) was added to cells for 3-4 hours. The cells were lysed in 1X lysis buffer (100 mM NaCl, 20 mM Tris-HCL pH 7.5, 1 mM EDTA, 0.5% NP40, 0.5% deoxycholate and 100 mM PMSF, 0.5 ug/ml leupeptin and 2 mg/ml aprotinin), stored on ice for 10 minutes and cleared by
30 centrifugation (15,000 x g for 5 minutes). Lysates were immunoprecipitated with the designated antibody immobilized on Protein A Sepharose (BioRad).

HCV templates

All E1 and E2 templates were generated by PCR and confirmed by sequencing. The appropriate 5' primer containing a methionine residue and an *NcoI* site was used along with 3' primers that had a termination codon following the designated envelope endpoint and finally, for E1, a *BamHI* site. Both oligos had non-specific sequences on the ends to facilitate more efficient digestions by *NcoI* and *BamHI* enzymes. Digested PCR fragments were ligated into *NcoI/BamHI*-digested pTM1 (Elroy-Stein and Moss, *Proc. Natl. Acad. Sci. USA* (1990) 87:6743-6747). The pTM1 vector contains the T7 promoter and the EMC leader proximal to the *NcoI* cloning site which corresponds to the first met residue encoded by the designated DNA. E2 templates were digested with *NcoI* and *AscI* and cloned into *NcoI* (partial)/*AscI*-pTM1-CE2 (Selby et al., *J. Gen. Virol.* (1993) 74:1103-1113) to generate the H clones where translations began at amino acid 1 and encode core, E1 and the designated E2 regions. For the truncated E1 polypeptides, coding templates began with a methionine residue, followed by isoleucine and then amino acid 172. For the truncated E2 constructs, the methionine at position 364 was used as the N-terminus in the constructions. After identifying possible clones and amplifying the DNA, all were sequenced. All the E1 clones were shown to be correct by sequencing. Most of the E2 clones showed the correct sequence with the exception of the loss of a single leucine residue within E2. This loss did not influence the conclusions as it was not in the vicinity of the C-terminus.

Immunoprecipitations

Immunoprecipitations were performed on media from transfected cells as described in Selby et al., *J. Gen. Virol.* (1993) 74:1103-1113, except that the precipitates were washed at least once with lysis buffer containing 500 mM NaCl instead of 100 mM. The precipitates were resuspended in Laemmli buffer, boiled and analyzed on 12.5% or 15% acrylamide gels. Gels were enhanced (Amplify, Amersham), dried and exposed to film at -80°C. The antisera used were previously described in Selby et al., *J. Gen. Virol.* (1993) 74:1103-1113. Endo H treatments of immunoprecipitates were performed according to the manufacturer's specifications (Oxford Glycosystems).

For the immunoprecipitation of secreted E1 and E2 proteins, media collected from transfected cells was microcentrifuged and immunoprecipitated in tubes containing either monoclonal anti-E1 antibody or rabbit polyclonal anti-E2 antibody immobilized on protein G sepharose (Pharmacia) or protein A sepharose (Sigma), respectively. After an
5 overnight incubation at 4°C, the sepharose Ab-Ag complexes were washed with lysis buffer twice, once with lysis buffer containing 500 mM NaCl and finally with 120 mM Tris, 8.0. After all the liquid was aspirated, approximately 30 µl of Laemmli sample buffer was added, the samples were boiled and then loaded onto a 12.5% acrylamide gel. Following electrophoresis, the gels were fixed, amplified and dried onto 3MM.
10 The dried gels were exposed to film with an intensifying screen. Transfection controls were done with a template for the β-galactosidase cDNA (pTM1-β-gal).

NS2B sequencing

BSC40 cells were transfected with pE2₁₀₀₆, labeled with ³⁵S-met (NEN) and lysed
15 with lysis buffer containing protease inhibitors. Cleared lysates were precipitated with rabbit anti-E2 overnight, washed and analyzed on a 15% acrylamide gel. The gel was then transferred to PVDF in 20% methanol/1x running buffer for 2 hours at 50 volts. The region containing NS2B was identified by autoradiography and cut out. NS2B was sequenced via sequential Edmann degradation on a Applied Biosystems 470A gas-phase
20 sequencer (Speicher, (1989). Microsequencing with PVDF membranes: Efficient electroblotting, direct protein adsorption and sequencer program modifications. in *Techniques in Protein Chemistry*. Ed. T.E. Hugli. Academic Press, San Diego, CA. pp 24-35.). Butyl chloride-containing fractions were evaporated and counted with scintillation fluid.

25

Example 1

Secretion/Retention of E1

As explained above, a series of E1 templates were generated in pTM1, using PCR (Figure 5). In particular, coding templates beginning with a methionine residue,
30 followed by isoleucine and then amino acid 172 of the HCV polyprotein and continuing to amino acid 330, and clones of 10 amino acid increments through amino acid 380,

were generated. Amino acids 173 through 191 correspond to the C-terminus of core which apparently serves a role as a signal sequence. Mature E1 is thought to begin at amino acid 192 of the polyprotein following signal sequence cleavage.

Anti-E1 immunoreactive material was recovered in the media of cells transfected with templates through amino acid 360. No E1 was detected in the media of cells transfected with clones terminating at amino acids 370 and 380.

Example 2

Secretion/Retention of E2

A series of E2 templates were also generated in pTM1, using PCR (Figure 6). In particular, the first coding amino acid of E2 corresponds to methionine at position 364 and this was used as the N-terminus in the constructions. Amino acid 364 corresponds to the approximate start of the E2 signal peptide (Hijikata et al., *Proc. Natl. Acad. Sci. USA* (1991) 88:5547-5551; Ralston et al., *J. Virol.* (1993) 67:6753-6761). Mature E2 is thought to begin with amino acid 385. The staggered C-termini ranged from amino acid 661 through 1006. In particular, the clones terminated at amino acids 661, 699, 710, 715, 720, 725, 730, 760, 780, 807, 837, 906 and 1006.

Anti-E2 immunoreactive material was recovered in the media of cells transfected with templates through 725. Small amounts of E2 were detected in media transfected with clones terminating at amino acid 730. Little or no E2 was detected in the media of cells transfected with clones terminating at amino acids beyond 730. The sequence just before 730 is quite hydrophobic, reminiscent of a membrane anchor sequence. Thus, it appears that sequences between 715 and 730 and extending as far as approximately amino acid residue 746 (see, Lin et al., *J. Virol.* (1994) 68:5063-5073), serve to anchor E2 to the ER, preventing secretion.

An additional secreted E2 molecule was made which included amino acids 383_{aa} through 715_{lys}. The E2 molecule was expressed using a Chinese hamster ovary cell/dihydrofolate reductase (CHO/DHFR) expression system as follows. A DNA fragment of HCV E2 from amino acid 383 to amino acid 715 of HCV1 was generated by PCR and then ligated into the pMH vector which includes the strong mouse

cytomegalovirus immediate early (MCMV ie) promoter and the selectable marker DHFR, to render plasmid pMHE2-715 (see Figure 7). This plasmid was then stably transfected into the CHO cell line, Dg44, as follows. 100 μ g of DNA were combined with lipofectin (Gibco-BRL) for transfection into 1×10^7 cells on 10 cm dishes. The cells
5 were transfected and incubated for 24 hours in non-selective medium (Ham's nutrient mixture F-12, JRH Biosciences series no. 51, proline, glutamine and sodium pyruvate) then allowed to recover for 24 hours in non-selective medium supplemented with fetal bovine serum. The non-selective medium was then replaced with selective medium (Ham's nutrient mixture F-12, JRH Biosciences series no. 52, proline, glutamine, and
10 sodium pyruvate) supplemented with dialyzed fetal bovine serum. The medium was changed every 3-4 days until colonies began to form.

The DG44 cell line lacks endogenous DHFR activity. Thus, only those cells transfected with the pMHE2-715 plasmid having the DHFR gene can grow and form colonies in the selective medium which lacks hypoxanthine, aminopterin and thymidine.
15 Approximately 2200 colonies were picked and grown up in 96-well plates. When the clones reached confluency 5 days later, the media was assayed for secreted E2 by ELISA using monoclonal antibody 3E5-1 that was raised against a linear determinant of E2 as the capture antibody and E3 expressed in CHO cells and purified from a monoclonal antibody (5E5H7 reactive to a conformational determinant of E2) affinity
20 column as the standard. The top 83 expressing clones were expanded to 24-well plates. Once confluency was reached in the 24-well plates, the media was assayed and then the top 41 clones were expanded to 6-well plates. When the clones reached confluency, the media was assayed and the top 21 clones were expanded to 75 cm₂ flasks. At this point expression was confirmed by the ELISA and by fluorescent immunostaining using the
25 monoclonal antibody 3E5-1. 5 clones had 100% of their cells fluorescing. On the basis of both assays, the top 3 clones were chosen to be purified by limiting dilution.

The top 7 clones were pooled and plated out in 10 cm dishes for growth in a range of methotrexate (MTX) concentrations. MTX is an inhibitor of DHFR. Among a population of cultured cells, variant clones that are overexpressing DHFR are resistant
30 to the toxic MTX. It has been observed that the other genes on a plasmid transfected into CHO cells can be co-amplified with the DHFR gene. The cells expressing HCV

E2 were amplified in selective media with final concentrations of 10, 20, 50, 100, 200 nM MTX. Colonies grew up in the 10 and 20 nM MTX media only. These 384 clones were picked, expanded up and checked for expression as done before in the initial selection. In adherent cultures, the top amplified clone had an expression level that was
5 31% higher than the top unamplified clone and was fluorescing 100%.

Example 3

E2:E1 Co-immunoprecipitation

E1 immunoprecipitates with E2 using monoreactive antisera to E2 or E1
10 (Grakoui et al., *J. Virol.* (1993) 67:1385-1395; Lanford et al., *Virology* (1993) 197:225-235; Ralston et al., *J. Virol.* (1993) 67:6753-6761). This interaction is very strong as it resists disruption by 0.5% SDS (Grakoui et al., *J. Virol.* (1993) 67:1385-1395) or by high salt/non-ionic detergent (Ralston et al., *J. Virol.* (1993) 67:6753-6761); Glazer and Selby, unpublished observations). To define the region of
15 E2 important for this interaction, BSC40 cells with H templates (core through E2) were transfected and radiolabeled lysates immunoprecipitated with rabbit polyclonal anti-E2 antiserum. E2 species encoded by templates that terminated at amino acid 730, 760 and 780 associated with E1. There was no quantitative difference in E1 co-precipitation with all templates between 730 and 1006. In contrast, E2 species encoded by templates
20 661, 699, 710, 715, 720 and 725 failed to significantly co-precipitate E1. As a control, the same lysates were immunoprecipitated with patient antiserum LL. E1 proteins from all templates were clearly precipitated with LL; E2 proteins were precipitated efficiently with the exception of template 661H. Relatively poor detection of E2₆₆₁ has consistently been observed, possibly owing to a different structure of this protein compared to longer
25 versions.

Thus, the sequences between amino acids 715 and 730 are important for efficient E1 association. Removal of this C-terminal anchor seems to preclude the association that normally appears between E1 and E2. However, when both envelopes are secreted, it appears that some association occurs. This result is opposite of the secretion data
30 with respect to E2 presented above, thereby establishing an inverse relationship between secretion of E2 and interaction of E2 with E1.

Example 4

Formation of Truncated E1 and E2 Complexes

To test whether truncated forms of E1 and E2, as produced above, were capable of associating with one another, the construct coding for an E1 polypeptide, C-terminally truncated at amino acid 360, was co-transfected with constructs coding for an E2 polypeptide, C-terminally truncated at amino acid 715 or an E2 polypeptide, C-terminally truncated at amino acid 725. Cells were transformed as described above and media collected. E1/E2 complexes were observed clearly with anti-E2 monoclonal antibody where E1 was co-precipitated. The converse precipitation with monoclonal anti-E1 yielded much less E1 compared to the former precipitation. A similar observation was made in Ralston et al., *J. Virol.* (1993) 67:6753-6761, with respect to non-truncated constructs. Such complex formation is significant because it demonstrates that the regions of E1 and E2 important for their interaction are retained, despite elimination of all or a part of the membrane spanning domains.

Example 5

Multiple E2 Species: Endo H Studies

Three E2 bands resulted from endo F treatment of E2, suggesting that more than one E2 species exists although the endpoints remain undefined (Grakoui et al., *J. Virol.* (1993) 67:1385-1395). Furthermore, a high molecular weight band reactive with both E2 and NS2 antisera led to the speculation of a possible E2-NS2 species (Grakoui et al., *J. Virol.* (1993) 67:1385-1395; Tomei et al., *J. Virol.* (1993) 67:4017-4026). Endo H treatment was used to define these three E2 glycoproteins. Endo H is a suitable deglycosidase for these experiments as only immature, high mannose glycoproteins are observed in transient expression systems (Spaete et. al., *Virology* (1992) 188:819-830; Ralston et al., *J. Virol.* (1993) 67:6753-6761). Endo H treatment of E2 proteins encoded by templates E2₆₆₁ through E2₇₃₀ yielded single bands that increased in size concomitantly with the elongated templates. Endo H-treated E2 proteins encoded by templates E2₇₆₀, E2₇₈₀ and E2₈₀₇ showed an additional band that stopped increasing in size at amino acid 807 when compared to longer templates. These data suggest that one form of E2 terminates near amino acid 730. Recent data suggest that E2 may terminate

at about amino acid 746 (see, Lin et al., *J. Virol.* (1994) 68:5063-5073). A second form of E2 appears to terminate near amino acid position 807. On templates E2₉₀₆ and E2₁₀₀₆ a third band was note which migrated higher than the previous doublet. The sizes of the bands from these two templates were consistent with E2 molecules that were
5 encoded by the entire templates. The mobility of E2-NS2 fusion in pTM1-HCV was slightly decreased relative to E2₁₀₀₆. These observations are consistent with the third form of E2 ending co-incidentally with the NS2/NS3 junction at amino acid 1026 (Grakoui et al., *Proc. Natl. Acad. Sci. USA* (1993) 90:10583-10587).

NS2 has been identified preceding NS3 (Grakoui et al., *J. Virol.* (1993) 67:1385-1395; Hijikata et al., *Proc. Natl. Acad. Sci. USA* (1993) 90:10773-10777;
10 Selby et al., *J. Gen. Virol.* (1993) 74:1103-1113). The results presented here and other recent reports (Lin et al., *J. Virol.* (1994) 68:5063-5073) support differential precursor processing whereby extensions of E2 beyond amino acids 730-746 represent E2 fusions to NS2. We tentatively suggest that the small protein coding region approximately
15 between amino acids 730-746 and 807 predicted by the endo H experiments corresponds to NS2A while the downstream protein is NS2B. This nomenclature is based on the similar organization of proteins amongst the Flaviviridae. It is possible that NS2A is not independently cleaved from the polyprotein as is NS2B and it remains to be determined if NS2A is in fact a non-structural protein.

20

Example 6

NS2B: E2: Co-immunoprecipitation and the

N-terminus of NS2B

Patient serum LL detected the E2 glycoproteins and two other species of 14 kDa
25 and 21 kDa from transfections with templates E2₉₀₆ and E2₁₀₀₆. These latter species correspond to NS2B because the difference in size correlates perfectly with the difference in the template lengths and a 23 kDa NS2B was detected from the full-length HCV template, pTM1-HCV (Selby et al., *J. Gen. Virol.* (1993) 74:1103-1113). When an anti-E2 antibody was used to immunoprecipitate the same lysates, NS2B
30 co-precipitated with E2. Under these conditions E1 also co-immunoprecipitates. Additionally, higher anti-E2 reactive species were also seen in 906, 1006 and

pTM1-HCV precipitations, which correspond to E2 fusions with NS2A and different lengths of NS2B.

Using the various E2 templates, we defined one region of E2 important for co-precipitation of NS2B. E2 templates ending at amino acids 699, 730, 760, 780 and
5 807 were co-transfected with pTM1-NS25, which encodes amino acids 730 through 3011 (Selby et al., *J. Gen. Virol.* (1993) 74:1103-1113). pTM1-NS25 was used since this template was previously determined to encode NS2 and an independent template was required for NS2B expression to define sequences important for the E2:NS2B interaction. Anti-E2 antibody immunoprecipitated E2 proteins and also co-precipitated
10 NS2B (23 kDa) and probable NS4B (27 kDa) from all lysates except from that transfected with the E2₆₉₉ template. As a positive control, truncated NS2B from the E2₁₀₀₆ template was co-precipitated with E2. These data demonstrate that amino acids 699 through 730 are important for NS2B and NS4B association with E2.

The anti-E2 antibody that co-precipitated NS2B was made against an SOD-E2
15 fusion that terminated at amino acid 662. This excludes the possibility of cross-reactivity to NS2B as a cause for NS2B detection. This NS2B band was not glycosylated as evidenced by its insensitivity to endo H treatment. Because co-precipitation of NS2B was with little background and efficient, this association property was used to radio-sequence the N-terminus of NS2B. The immunoprecipitate
20 of the ³⁵S-methionine-labeled lysate from an E2₁₀₀₆ transfection was analyzed on a 15% acrylamide gel and transferred to PVDF for sequencing. Using a methionine residue 18 amino acids away from residue 810 as a reference, the amino terminus of NS2B was identified as amino acid 810. This assignment agrees with the predicted E2 C-terminus based on the endo H pattern from the E2 deletion templates and confirms the cleavage
25 sites recently reported by Grakoui et al., *Proc. Natl. Acad. Sci. USA* (1993) 90:10583-10587 and Mizushima et al., *J. Virol.* (1994) 68:2731-2734. The presumptive NS2A termini is predicted to reside as amino acids 730 and 809 if its exists as an independent protein.

Thus, novel secreted E1 and E2 forms are disclosed. Although preferred
30 embodiments of the subject invention have been described in some detail, it is

understood that obvious variations can be made without departing from the spirit and the scope of the invention as defined by the appended claims.

CLAIMS

1. A hepatitis C virus (HCV) E1 polypeptide, lacking all or a portion of its membrane spanning domain such that said polypeptide is capable of secretion into growth medium when expressed recombinantly in a host cell.

2. The HCV polypeptide of claim 1, wherein said polypeptide lacks at least a portion of its C-terminus beginning at about amino acid 370, numbered with reference to the HCV1 E1 amino acid sequence.

3. The HCV polypeptide of claim 2, wherein said polypeptide lacks at least a portion of its C-terminus beginning at about amino acid 360, numbered with reference to the HCV1 E1 amino acid sequence.

4. A hepatitis C virus (HCV) E2 polypeptide lacking at least a portion of its membrane spanning domain such that said polypeptide is capable of secretion into growth medium when expressed recombinantly in a host cell, wherein said polypeptide lacks at least a portion of its C-terminus beginning at about amino acid 730 but not extending beyond about amino acid 699, numbered with reference to the HCV1 E2 amino acid sequence.

5. The HCV polypeptide of claim 4, wherein said polypeptide lacks at least a portion of its C-terminus beginning at about amino acid 725, numbered with reference to the HCV1 E2 amino acid sequence.

6. The HCV polypeptide of claim 4, wherein said polypeptide lacks a portion of its C-terminus beginning at about amino acid 715, numbered with reference to the HCV1 E2 amino acid sequence.

7. A polynucleotide comprising a coding sequence for an HCV polypeptide according to claim 1.

8. A polynucleotide comprising a coding sequence for an HCV polypeptide according to claim 2.

5 9. A polynucleotide comprising a coding sequence for an HCV polypeptide according to claim 3.

10. A polynucleotide comprising a coding sequence for an HCV polypeptide according to claim 4.

10 11. A polynucleotide comprising a coding sequence for an HCV polypeptide according to claim 5.

12. A polynucleotide comprising a coding sequence for an HCV polypeptide according to claim 6.

15

13. A recombinant vector comprising:
(a) a polynucleotide according to claim 7; and
(b) control elements that are operably linked to said coding sequence whereby said coding sequence can be transcribed and translated in a host cell.

20

14. A recombinant vector comprising:
(a) a polynucleotide according to claim 8; and
(b) control elements that are operably linked to said coding sequence whereby said coding sequence can be transcribed and translated in a host cell.

25

15. A recombinant vector comprising:
(a) a polynucleotide according to claim 9; and
(b) control elements that are operably linked to said coding sequence whereby said coding sequence can be transcribed and translated in a host cell.

30

16. A recombinant vector comprising:

- (a) a polynucleotide according to claim 10; and
- (b) control elements that are operably linked to said coding sequence whereby said coding sequence can be transcribed and translated in a host cell.

5 17. A recombinant vector comprising:

- (a) a polynucleotide according to claim 11; and
- (b) control elements that are operably linked to said coding sequence whereby said coding sequence can be transcribed and translated in a host cell.

10 18. A recombinant vector comprising:

- (a) a polynucleotide according to claim 12; and
- (b) control elements that are operably linked to said coding sequence whereby said coding sequence can be transcribed and translated in a host cell.

15 19. A host cell transformed with the recombinant vector of claim 13.

20 20. A host cell transformed with the recombinant vector of claim 14.

21. A host cell transformed with the recombinant vector of claim 15.

22. A host cell transformed with the recombinant vector of claim 16.

23. A host cell transformed with the recombinant vector of claim 17.

25 24. A host cell transformed with the recombinant vector of claim 18.

25. A method of producing a secreted HCV polypeptide, said method comprising:

- (a) providing a population of host cells according to claim 19; and
- 30 (b) culturing said population of cells under conditions whereby said HCV polypeptide is expressed and secreted.

26. A method of producing a secreted HCV polypeptide, said method comprising:

- (a) providing a population of host cells according to claim 20; and
- (b) culturing said population of cells under conditions whereby said HCV polypeptide is expressed and secreted.

27. A method of producing a secreted HCV polypeptide, said method comprising:

- (a) providing a population of host cells according to claim 21; and
- (b) culturing said population of cells under conditions whereby said HCV polypeptide is expressed and secreted.

28. A method of producing a secreted HCV polypeptide, said method comprising:

- (a) providing a population of host cells according to claim 22; and
- (b) culturing said population of cells under conditions whereby said HCV polypeptide is expressed and secreted.

29. A method of producing a secreted HCV polypeptide, said method comprising:

- (a) providing a population of host cells according to claim 23; and
- (b) culturing said population of cells under conditions whereby said HCV polypeptide is expressed and secreted.

30. A method of producing a secreted HCV polypeptide, said method comprising:

- (a) providing a population of host cells according to claim 24; and
- (b) culturing said population of cells under conditions whereby said HCV polypeptide is expressed and secreted.

31. A secreted E1/secreted E2 complex comprising:

(a) a hepatitis C virus (HCV) E1 polypeptide, lacking all or a portion of its membrane spanning domain such that said E1 polypeptide is capable of secretion into growth medium when expressed recombinantly in a host cell; and

5 (b) a hepatitis C virus (HCV) E2 polypeptide, lacking all or a portion of its membrane spanning domain such that said E2 polypeptide is capable of secretion into growth medium when expressed recombinantly in a host cell.

32. The secreted E1/secreted E2 complex of claim 31, wherein said E1 polypeptide lacks at least a portion of its C-terminus beginning at about amino acid 370,
10 numbered with reference to the HCV1 E1 amino acid sequence and said E2 polypeptide lacks at least a portion of its C-terminus beginning at about amino acid 730, numbered with reference to the HCV1 E2 amino acid sequence.


33. The secreted E1/secreted E2 complex of claim 31, wherein said E1
15 polypeptide lacks at least a portion of its C-terminus beginning at about amino acid 360, numbered with reference to the HCV1 E1 amino acid sequence and said E2 polypeptide lacks at least a portion of its C-terminus beginning at about amino acid 725, numbered with reference to the HCV1 E2 amino acid sequence.

20 34. A vaccine composition comprising a pharmaceutically acceptable excipient and an HCV E1 polypeptide according to claim 1.

35. A vaccine composition comprising a pharmaceutically acceptable excipient and an HCV E2 polypeptide according to claim 4.

25

36. A vaccine composition comprising a pharmaceutically acceptable excipient and an HCV secreted E1/secreted E2 complex according to claim 31.



= presumptive signal sequence for E1, derived from the C-terminus of the core coding region



= region of the HCV polyprotein which contributes to E1 retention within cells

Figure 1

170 MetIleCysSerPheSerIlePheLeuLeuAlaLeuLeuSerCysLeuThrValProAla
 ATGATTGCTCTTTCTCTATCTTCCTTCTGGCCCTGCTCTCTTGCTTGACTGTGCCCGCT
 TACTAAACGAGAAAGAGATAGAAGGANGACCGGGACGAGAGAAOGAACTGACACGGGCGA

Mature E1

190 SerAlaTyrGlnValArgAsnSerThrGlyLeuTyrHisValThrAsnAspCysProAsn
 TCGGCTACCAAGTGGCGCAACTCCACGGGGCTCTACCAAGTCACCAATGATTGCCCTAAC
 AGCGGATGGTTACGCGTTGAGGTGCCCGAGATGGTGAGTGGTTACTAACGGGATTG

210 SerSerIleValTyrGluAlaAlaAspAlaIleLeuHisThrProGlyCysValProCys
 TCGAGTATTGTGTACGAGGCGGCGATGCCATCTGCACACTCGGGGTGGTCCCTTGC
 AGCTCATACACATGCTCGCGCGCTACGGTAGGAAGTGTGAGGCCCAAGCAGGGAACG

230 ValArgGluGlyAsnAlaSerArgCysTrpValAlaMetThrProThrValAlaThrArg
 GTTCGTAGGGCAACGCTCGAGGTGTGGGTGGCGATGACCCCTACGGTGGCCACCAGG
 CAAGCACTCCCGTTGCGGAGCTCCACAACCCACCGCTACTGGGGATGCCACGGTGTGCC

250 AspGlyLysLeuProAlaThrGlnLeuArgArgHisIleAspLeuLeuValGlySerAla
 GATGGCAAACTCCCGCGACGCGAGCTTCGACGTCACATCGATCTGCTTGTGGGAGCGCC
 CTACCGTTTGGGGGCGCTGCGTGAAGCTGCAGTGTAGCTAGACGAACAGCCCTCGCGG

270 ThrLeuCysSerAlaLeuTyrValGlyAspLeuCysGlySerValPheLeuValGlyGln
 ACCCTCTGTTCGGCCCTCTACGTGGGGGACCTCTGGGGTCTGTCTTTCTGTGGCCAA
 TGGGAGACAAGCGGGAGATGCACCCCTGGAGACGCCAGACAGAAAGAACAGCCGGTT

290 LeuPheThrPheSerProArgArgHisTrpThrThrGlnGlyCysAsnCysSerIleTyr
 CTGTTTACCTTCTCTCCAGGCGCCACTGGACGACGCAAGGTTGCAATTGCTCTATCTAT
 GACAAATGGAAGAGAGGGTCCGCGGTGACCTGCTGCGTTCCACGTTAACGAGATAGATA

310 ProGlyHisIleThrGlyHisArgMetAlaTrpAspMetMetMetAsnTrpSerProThr
 CCGGGCCATATAACGGGTCAACGCGATGGCATGGGATATGATGATGAAGTGGTCCCTACG
 GGGCGGTATATTGCCAGTGGCGTACCGTACCCTATACTACTACTTGACCAGGGGATGC


330 ThrAlaLeuValMetAlaGlnLeuLeuArgIleProGlnAlaIleLeuAspMetIleAla
 ACGGCGTTGGTAATGGCTCAGCTGCTCGGATCCCAAGCCATCTTGGACATGATCGCT
 TGCGCAACCATTACCGAGTCGACGAGGCCTAGGGTGTTOGGTAGAACCTGTACTAGCGA

C-terminal Anchor

350 GlyAlaHisTrpGlyValLeuAlaGlyIleAlaTyrPheSerMetValGlyAsnTrpAla
 GGTGCTCACTGGGGAGTCTGGCGGGCATAGGTATTTCTCCATGGTGGGGAACTGGGGG
 CCACGAGTGACCCCTCAGGACCGCCCGTATCGCATAAAGAGGTACCACCCCTTGACCCG

370 LysValLeuValValLeuLeuLeuPheAlaGlyOP
 AAGGTCCTGGTAGTGTCTGCTATTTGCCGGCTGA
 TTCCAGGACCATCAAGACGACGATAACGGCGGACT

Figure 2



= presumptive signal sequence for E2, derived from the C-terminus of the E1 coding region



= region of HCV polypeptide which contributes E2 retention within cells

Figure 3

364 MetValGlyAsnTrpAlaLysValLeuValValLeuLeuLeuPheAlaGlyValAspAla
ATGGTGGGGAAC TGGCGAAGGTCTGCTAGTGTCTGCTATTGTCGGGCGTCCGACGG
TACCACCCCTTGACCGCTTCCAGGACCATCACGACGACGATAAACGGCGCGAGCTGCGC

Mature E2

384 GluThrHisValThrGlyGlySerAlaGlyHisThrValSerGlyPheValSerLeuLeu
GAAACCCACGTCACCGGGGAAGTGGCGGCCACACTGTGTCTGGATTGTTAGCCTCCTC
CTTTGGGTGCAGTGGCCCCCTTCACGGCCGGTGTGACACAGACCTAAACAATCGGAGGAG

404 AlaProGlyAlaLysGlnAsnValGlnLeuIleAsnThrAsnGlySerTrpHisLeuAsn
GCACCAGGCGCAAGCAGAACGTCCAGCTGATCAACACCAACGGCAGTTGGCACCTCAAT
CGTGGTCCGGGTTCTGTCTGCAGGTGACTAGTTGTGGTTGCGTCAACCGTGGAGTTA

424 SerThrAlaLeuAsnCysAsnAspSerLeuAsnThrGlyTrpLeuAlaGlyLeuPheTyr
AGCACGGCCCTGAAGTCAATGATAGCTCAACACCGGCTGGTGGCAGGGCTTTTCTAT
TGTGTCGGGACTTGACCTTACTATOGGAGTTGTGCGCGACCAACCGTCCGAAAAGATA

444 HisHisLysPheAsnSerSerGlyCysProGluArgLeuAlaSerCysArgProLeuThr
CACCACAAGTCAACTCTTCAGGCTGTCTGAGAGGCTAGCCAGCTGCGACCCCTTACC
GTGGTGTTCAGTTGAGAAGTCCGACAGGACTCTCGATCGGTGACGGCTGGGAATGG

464 AspPheAspGlnGlyTrpGlyProIleSerTyrAlaAsnGlySerGlyProAspGlnArg
GATTTTGACGAGGCTGGGGCCCTATCAGTTATGCCAAGGAGCGGCCCGACGAGCGC
CTAAACTGGTCCCGACCCCGGATAGTCAATACGGTTGCCTTCGCGGGGCTGGTTCGGC

484 ProTyrCysTrpHisTyrProProLysProCysGlyIleValProAlaLysSerValCys
CCCTACTGCTGGCACTACCCCCAAAACCTTGCGGTATTGTGCCCGGAAGAGTGTGTGT
GGGATGACGACGTGATGGGGGTTTTTGAACGCCATAACACGGGCGCTTCTCACACACA

504 GlyProValTyrCysPheThrProSerProValValValGlyThrThrAspArgSerGly
GGTCCGGTATATTGCTTCACTCCCAGCCCCGTGGTGGTGGGAACGACGACAGGTGGGGC
CCAGGCCATATAACGAAGTGAGGGTGGGGCACCACCACCTTGTGGCTGTCCAGCCCCG

524 AlaProThrTyrSerTrpGlyGluAsnAspThrAspValPheValLeuAsnAsnThrArg
GCGCCACCTACAGCTGGGGTGAAAATGATACGGACGTCTTGTCTTAAACAATACCAGG
CGCGGGTGGATGTCGACCCCACTTTTACTATGCCTGCAGAAGCAGGAATTGTTATGGTCC

544 ProProLeuGlyAsnTrpPheGlyCysThrTrpMetAsnSerThrGlyPheThrLysVal
CCACCGCTGGGCAATTGGTTTCGGTTGTACCTGGATGAACCTCAACTGGATTACCAAAGTG
GGTGGCGACCCGTTAACCAAGCCAACATGACCTACTTGAGTTGACCTAAGTGGTTTCAC

564 CysGlyAlaProProCysValIleGlyGlyAlaGlyAsnAsnThrLeuHisCysProThr
TGCGGAGCGCCTCCTTGTGTATCGGAGGGGCGGGCAACAACACCTGCACTGCCCCACT
ACGCTCGCGGAGGAACACAGTAGCCTCCCGCCCGTTGTTGTGGGACGTGACGGGGTGA

584 AspCysPheArgLysHisProAspAlaThrTyrSerArgCysGlySerGlyProTrpIle
GATTGCTTCGCAAGCATCCGGACGCCACATACTCTCGGTGCGGCTCGGTCCCTGGATC
CTAACAAGGCGTTCGTAGGCTGCGGTGTATGAGAGCCACGCGAGGCCAGGGACCTAG

604 ThrProArgCysLeuValAspTyrProTyrArgLeuTrpHisTyrProCysThrIleAsn
ACACCCAGGTGCCGTGGTGGTACTACCGTATAGGCTTTGGCATTATCCTGTACCATCAAC
TGTGGGTCCAAGGACCGAGCTGATGGGCATATCCGAAACCGTAATAGGAACATGGTAGTTG

624 TyrThrIlePheLysIleArgMetTyrValGlyGlyValGluHisArgLeuGluAlaAla
TACACCATATTTAAATCAGGATGTACGTGGGAGGGGTGGAACACAGGCTGGAAGCTGCC
ATGTGGTATAAATTTAGTCTACATGCACCTCCCGAGCTTGTGTCCGACCTTCGACGG

644 CysAsnTrpThrArgGlyGluArgCysAspLeuGluAspArgAspArgSerGluLeuSer
TGCAACTGGAAGCGGGGGAACGTTGCGATCTGGAAGATAGGGACAGGTCCGAGCTCAGC
ACGTTGACCTGCGCCCGCTTGCAACGCTAGACCTTCTATCCCTGTCCAGGCTCGAGTGC

Figure 4A

664 ProLeuLeuLeuThrThrThrGlnTrpGlnValLeuProCysSerPheThrThrLeuPro
COGTTACTGCTGACCACTACACAGTGGCAGGTCTCCCGTGTTCCTTCACAACCCCTGCCA
GGCAATGACGACTGGTGATGTGTACCGTCCAGGAGGGCACAAGGAAGTGTGGGACGGT

684 AlaLeuSerThrGlyLeuIleHisLeuHisGlnAsnIleValAspValGlnTyrLeuTyr
GCCTGTGCCACGGCCTCATCCACCTCCACCAGAACATTGTGGACGTGCAGTACTTGTAC
CGGAACAGGTGGCCGGAGTAGGTGGAGTGGTCTTGTAACACCTGCACGTCAATGAACATG

704 GlyValGlySerSerIleAlaSerTrpAlaIleLysTrpGluTyrValValLeuLeuPhe
GGGGTGGGGTCAAGCATCGCTCCTGGGCCATTAGTGGGAGTACGTCTCTCTCTGTTC
CCCCACCCAGTTCTGTAGCGCAGGACCGGTAATTCACCTCATGCAGCAGGAGGACAAG

C-terminal anchor

724 LeuLeuLeuAlaAspAlaArgValCysSerCysLeuTrpMetMetLeuLeuIleSerGln
CTTCTGCTTGCAGACGCGCGCTCTGCTCTCTGCTGTGGATGATCTACTCATATCCCAA
GAAGACGAACGCTCTGCGCGCAGACGAGGACGAACACCTACTACGATGAGTATAGGGTT

NS2A

744 AlaGluAlaAlaLeuGluAsnLeuValIleLeuAsnAlaAlaSerLeuAlaGlyThrHis
GCGAAGCGGCTTGGAGAACCTCGTAATCTTAATGCAGCATCCCTGGCGGGACGCAC
CGCCTTCGCGAAACCTCTTGGAGCATTATGAATTACGTCGTAGGACCGGCCCTGCGTG

764 GlyLeuValSerPheLeuValPhePheCysPheAlaTrpTyrLeuLysGlyLysTrpVal
GGTCTGTATCTCTCTCTGTCTCTCTGCTTGTGATGGTATCTGAAGGGTAAGTGGGTG
CCAGAACATAGGAAGGAGCACAAGAAGACGAAACGTACCATAGACTTCCCATTCACCCAC

784 ProGlyAlaValTyrThrPheTyrGlyMetTrpProLeuLeuLeuLeuLeuAlaLeu
CCGGAGGGGTCTACACCTTCTACGGGATGTGGCCTCTCTCTCTCTCTGTGGCGTTG
GGGCTCGCCAGATGTGGAAGATGCCCTACACGGAGAGGAGGAAGAGGACAACCGCAAC

NS2B

804 ProGlnArgAlaTyrAlaLeuAspThrGluValAlaAlaSerCysGlyGlyValValLeu
CCCCAGCGGGCTACGGCTGGACACGGAGGTGGCGCGCTCGTGTGGCGGTGTTGTTCTC
GGGTCGCGCGCATGCGCGACCTGTGCCTCCACGGCGCAGCACACCGCCACAACAAGAG

824 ValGlyLeuMetAlaLeuThrLeuSerProTyrTyrLysArgTyrIleSerTrpCysLeu
GTGGGTGATGCGGCTAACTCTGTACCATATTACAAGCGCTATATCAGCTGGTGCTTG
CAGCCCAACTACCGGATTGAGACAGTGGTATAATGTTGCGATATAGTCGACCAGAAC

844 TrpTrpLeuGlnTyrPheLeuThrArgValGluAlaGlnLeuHisValTrpIleProPro
TGGTGGCTTCAGTATTTTCTGACAGAGTGAAGCGCACTGCACGTGTGGATTCCCCC
ACCACCGAAGTCATAAAGACTGGTCTCACCTTCGGTTGACGTGCACACCTAAGGGGGG

864 LeuAsnValArgGlyGlyArgAspAlaValIleLeuLeuMetCysAlaValHisProThr
CTCAACGTCCGAGGGGGGCGACGCGCTCATCTTACTCATGTGTGCTGTACACCGACT
GAGTTGCAGGCTCCCCCGCGCTGCGGCAGTAGAATGAGTACACACGACATGTGGGCTGA

884 LeuValPheAspIleThrLysLeuLeuLeuAlaValPheGlyProLeuTrpIleLeuGln
CTGGTATTTGACATCACCAATTGCTGCTGGCGCTCTTCGGACCCCTTGGATTCTTCAA
GACCATAAACTGTAGTGGTTTAAACGACGCGCAGAACCTGGGGAAACCTAAGAAGTT

904 AlaSerLeuLeuLysValProTyrPheValArgValGlnGlyLeuLeuArgPheCysAla
GCCAGTTTGCTTAAAGTACCCTACTTTGTGCGCGTCCAAGGCCTTCTCCGGTTCTGCGCG
CGGTCAAACGAATTTTCATGGGATGAAACACGCGCAGGTTCCGGAAGAGGCCAAGACGCGC

924 LeuAlaArgLysMetIleGlyGlyHisTyrValGlnMetValIleIleLysLeuGlyAla
TTAGCGCGAAGATGATCGGAGGCCATTACGTGCAATGGTCATCATTAAGTTAGGGGCG
AATCGCGCTTCTACTAGCTCGGTAATGCACGTTTACCAGTAGTAATTCAATCCCCGC

944 LeuThrGlyThrTyrValTyrAsnHisLeuThrProLeuArgAspTrpAlaHisAsnGly
CTTACTGGCACCTATGTTTATAACCATCTCACTCTCTCTCGGGACTGGGCGCACACGGC
GAATGACCGTGATACAAATATTGGTAGAGTGAGGAGAAGCCCTGACCGCGTGTGCGG

Figure 4B

964 LeuArgAspLeuAlaValAlaValGluProValValPheSerGlnMetGluThrLysLeu
TTGCGAGATCTGGCCGTGGCTGTAGAGCCAGTCGTCTTCTCCCAAATGGAGACCAAGCTC
AACGCTCTAGACCGGCACCGACATCTCGGTCAGCAGAAGAGGGTTTACCTCTGGTTGAG

984 IleThrTrpGlyAlaAspThrAlaAlaCysGlyAspIleIleAsnGlyLeuProValSer
ATCAGTGGGGGGCAGATACCGCCGCGTGGGTGACATCATCAACGGCTTGCCTGTTCC
TAGTGCACCCCCGTCTATGGCGGCGCACGCCACTGTAGTAGTTGCCGAACGGACAAAGG

1004 AlaArgArgGlyArgGluIleLeuLeuGlyProAlaAspGlyMetValSerLysGlyTrp
GCCCGCAGGGGCGGGAGATACTGCTCGGGCCAGCCGATGGAATGGTCTCCAAGGGTTGG
CGGGCGTCCCGGCCCTCTATGACGAGCCCGGTGGGTACCTTACCAGAGGTTCCCAACC

1024 ArgLeuLeu
AGGTGCTG
TCCAACGAC

Figure 4C




Figure 5




Figure 6




FIG. 7