

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
09 November 2017 (09.11.2017)



(10) International Publication Number
WO 2017/191114 A1

(51) International Patent Classification:

C07D 403/14 (2006.01) *A61K 31/426* (2006.01)
C07D 417/14 (2006.01) *A61P 9/00* (2006.01)
A61K 31/435 (2006.01) *A61P 13/00* (2006.01)
A61K 31/495 (2006.01)

GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/EP2017/060383

(22) International Filing Date:

02 May 2017 (02.05.2017)

Declarations under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

(25) Filing Language:

English

(26) Publication Language:

English

Published:

— *with international search report (Art. 21(3))*

(30) Priority Data:

16168172.1 03 May 2016 (03.05.2016) EP

(71) Applicants: **BAYER AKTIENGESELLSCHAFT** [DE/DE]; Kaiser-Wilhelm-Allee 1, 51373 Leverkusen (DE). **BAYER PHARMA AKTIENGESELLSCHAFT** [DE/DE]; Müllerstr. 178, 13353 Berlin (DE).

(72) Inventors: **COLLIN, Marie-Pierre**; Katemberger Schulweg 52, 42113 Wuppertal (DE). **KOLKHOF, Peter**; Falkenberg 121, 42113 Wuppertal (DE). **NEUBAUER, Thomas**; Kaiser-Wilhelm-Allee 12, 42117 Wuppertal (DE). **FÜRSTNER, Chantal**; Arnoldstr. 33, 45478 Mülheim/Ruhr (DE). **POOK, Elisabeth**; Im Lehmbruch 24, 42109 Wuppertal (DE). **WITWERT, Matthias, Beat**; Grünwalder Berg 41, 42105 Wuppertal (DE). **SCHMECK, Carsten**; Carl-Friedrich-Goerdeler-Straße 24, 45472 Mülheim (DE). **WASNAIRE, Pierre**; Am Botanischen Garten 41, 40225 Düsseldorf (DE). **TINEL, Hanna**; In der Beek 16, 42113 Wuppertal (DE). **SCHIRMER, Heiko**; Dingshauser Straße 51a, 42655 Solingen (DE).

(74) Agent: **BIP PATENTS**; Alfred-Nobel-Str. 10, 40789 Monheim am Rhein NRW (DE).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH,

(54) Title: HYDROXYALKYL-SUBSTITUTED HETEROARYLTRIAZOLE DERIVATIVES AND USES THEREOF

(57) Abstract: The present invention relates to novel 5-(hydroxyalkyl)-1-heteroaryl-1,2,4-triazole derivatives, to processes for the preparation of such compounds, to pharmaceutical compositions containing such compounds, and to the use of such compounds or compositions for the treatment and/or prevention of diseases, in particular for the treatment and/or prevention of renal and cardiovascular diseases.



WO 2017/191114 A1

HYDROXYALKYL-SUBSTITUTED HETEROARYLTRIAZOLE DERIVATIVES AND USES THEREOF

The present invention relates to novel 5-(hydroxyalkyl)-1-heteroaryl-1,2,4-triazole derivatives, to processes for the preparation of such compounds, to pharmaceutical compositions containing such
5 compounds, and to the use of such compounds or compositions for the treatment and/or prevention of diseases, in particular for the treatment and/or prevention of renal and cardiovascular diseases.

Vasopressin is a neurohormone which basically regulates water homeostasis and vascular tone. It is produced in specialized endocrine neurons in the *Nucleus supraopticus* and *N. paraventricularis* in the wall of the third ventricle (hypothalamus) and is transported from there along the neural
10 processes into the posterior lobes of the hypophysis (neurohypophysis). There, the hormone is released into the bloodstream in response to different physiological and pathophysiological stimuli. A disturbed neurohormonal regulation essentially manifests itself in an elevation of the sympathetic tone and inappropriate activation of the renin-angiotensin-aldosterone system (RAAS). While the inhibition of these components by beta-receptor blockers on the one hand by ACE inhibitors or
15 angiotensin-receptor blockers on the other is now an inherent part of the pharmacological treatment of cardiovascular diseases, the inappropriate elevation of vasopressin secretion is at present still not adequately treatable.

Vasopressin exerts its action mainly via binding to three receptors, which are classified as V1a, V1b and V2 receptors and which belong to the family of G protein-coupled receptors.

20 V2 receptors are located in the distal tubular epithelium and the epithelium of the collecting tubules in the kidney. Their activation renders these epithelia permeable to water. This phenomenon is due to the incorporation of aquaporins (special water channels) in the luminal membrane of the epithelial cells. Consequently, pharmacological inhibition of the action of vasopressin on the V2 receptor results in increased urine excretion. Hence, drugs with V2 antagonistic activity appear
25 particularly suitable for the treatment of all disease conditions which are associated with an overloading of the body with water.

V1b receptors (also named V3 receptors) are mainly detectable in the central nervous system. Together with corticotropin-releasing hormone (CRH), vasopressin regulates the basal and stress-induced secretion of adrenocorticotrophic hormone (ACTH) via the V1b receptor.

30 V1a receptors are mainly located on vascular smooth muscle cells (VSMC) but also on cardiomyocytes, fibroblasts and specialized renal cells like glomerular mesangial cells or cells of the macula densa which control the release of renin [Wasilewski MA, Myers VD, Recchia FA, Feldman AM, Tilley DG, *Cell Signal.*, 28(3), 224-233, (2016)]. The activation of VSMC V1a receptor by vasopressin gives rise to intracellular calcium release and according vasoconstriction.
35 Therefore, stimulation of VSMC V1a receptors causes increased vascular resistance and increased

cardiac afterload. Cardiac output is adversely affected by V1a-mediated vasoconstriction. The increase in afterload and direct stimulation of V1a receptors on cardiomyocytes can lead to cardiac hypertrophy and remodeling including fibrosis. Mice with cardiac-specific overexpression of V1a receptor develop cardiac hypertrophy leading to dilation and left ventricular dysfunction, suggesting an essential role for V1a receptor in the development of heart failure [Li X, Chan TO, Myers V, Chowdhury I, Zhang XQ, Song J, Zhang J, Andrel J, Funakoshi H, Robbins J, Koch WJ, Hyslop T, Cheung JY, Feldman AM, *Circulation.*; 124, 572-581 (2011)].

V1a receptor is also expressed in the renal cortical and medullary vasculature, where it mediates vasoconstriction of renal vessels and affecting overall renal blood flow. Thus, the activation of V1a receptor can decrease renal medullary blood flow inducing further pathological processes as tissue hypoxia, reduced oxygen and accordingly energy supply for tubular transport processes as well as direct damages of mesangial and macula densa cells. It has been demonstrated that mesangial V1a receptor activation mediates TGF β signaling and causes an increase in production of collagen IV. While this signaling contributes to extracellular matrix accumulation and remodeling in the kidney, similar signaling pathways are believed to occur in cardiac cells especially after myocardial infarction, which emphasizes the central role of V1a receptor in the development of hypertrophic and fibrotic processes in response to pathophysiological elevated vasopressin levels [Wasilewski MA, Myers VD, Recchia FA, Feldman AM, Tilley DG. *Arginine vasopressin receptor signaling and functional outcomes in heart failure. Cell Signal.*, 28(3), 224-233 (2016)].

Since V1a receptors are mainly expressed on VSMCs and thus participating in vascular function, a link to vascular diseases as peripheral arterial disease (PAD) including claudication and critical limb ischemia as well as coronary microvascular dysfunction (CMD) is conceivable.

Apart from this, V1a receptors are also expressed on human platelets and in the liver. The meaning of platelet V1a receptors is not fully understood although vasopressin induces aggregation of human platelets via V1a receptor at high concentrations *ex vivo*. Therefore, inhibition of vasopressin-induced platelet aggregation by V1a receptor antagonists is a useful pharmacological *ex vivo* assay making use of human tissue endogenously expressing the V1a receptor [Thibonnier M, Roberts JM, *J Clin Invest.*; 76:1857-1864, (1985)].

Vasopressin stimulates gluconeogenesis and glycogenolysis via activation of the hepatic V1a receptor. Animal studies have shown that vasopressin impairs glucose tolerance which could be inhibited by a V1a receptor antagonist thereby providing a link of vasopressin receptor V1a to diabetes mellitus. [Taveau C, Chollet C, Waeckel L, Desposito D, Bichet DG, Arthus MF, Magnan C, Philippe E, Paradis V, Fougelle F, Hainault I, Enhorning S, Velho G, Roussel R, Bankir L, Melander O, Bouby N. Vasopressin and hydration play a major role in the development of glucose intolerance and hepatic steatosis in obese rats. *Diabetologia*, 58(5), 1081-1090, (2015)].

Vasopressin was shown to contribute to the development of albuminuria and to diabetes-induced nephropathy in animal models which is consistent with epidemiological findings in humans.

It was found recently that vasopressin also seems to play a causal role in the development of preeclampsia. Chronic infusion of vasopressin during pregnancy in mice is sufficient to induce all
5 of the major maternal and fetal phenotypes associated with human preeclampsia, including pregnancy-specific hypertension [Santillan MK, Santillan DA, Scroggins SM, Min JY, Sandgren JA, Pearson NA, Leslie KK, Hunter SK, Zamba GK, Gibson-Corley KN, Grobe JL. Vasopressin in preeclampsia: a novel very early human pregnancy biomarker and clinically relevant mouse model. Hypertension. 64(4), 852-859, (2014)].

10 Vasopressin levels can be elevated in women with dysmenorrhoea (a gynecological disorder which is characterised by cyclical cramping pelvic pain) during menstruation, which appear to increase myometrial smooth muscle contraction. It was found recently that a selective vasopressin V1a receptor antagonist (relcovaptan/SR-49059) can reduce intrauterine contractions elicited by vasopressin.

15 For these reasons, agents which inhibit the action of vasopressin on the V1a receptor appear suitable for the treatment of several cardiovascular diseases. In particular, agents which inhibit the action of vasopressin selectively on the V1a receptor offer an especially ideal profile for the treatment of otherwise normovolemic patients, i.e. those which are not eligible for decongestion by e.g. high doses of loop diuretics or V2 antagonists, and where induced aquaresis via V2 inhibition
20 may be undesired.

Certain 4-phenyl-1,2,4-triazol-3-yl derivatives have been described in WO 2005/063754-A1 and WO 2005/105779-A1 to act as vasopressin V1a receptor antagonists that are useful for the treatment of gynecological disorders, notably menstrual disorders such as dysmenorrhea.

In WO 2011/104322-A1, a particular group of bis-aryl-bonded 1,2,4-triazol-3-ones, including
25 phenyl-1,2,4-triazol-3-yl and 1-phenyl-1,2,3-triazol-4-yl derivatives thereof, has been disclosed as antagonists of vasopressin V2 and/or V1a receptors being useful for the treatment and/or prevention of cardiovascular diseases. The described compounds, however, do not show sufficient selectivity towards the V1a receptor and mostly show combined activity on both vasopressin V1a and V2 receptors. Yet, as outlined above, a high affinity as well as selectivity for the V1a receptor
30 is a desirable prerequisite for the treatment of disease conditions where a decongestion is not desired and may lead to a dysregulated body fluid homeostasis including decreased blood plasma osmolality in otherwise normovolemic individuals.

In WO 2016/071212-A1 certain 5-(hydroxyalkyl)-1-phenyl-1,2,4-triazole derivatives have been disclosed, which act as potent antagonists of both vasopressin V1a and V2 receptors and, in
35 addition, exhibit significantly enhanced aquaretic potency in vivo after oral application. The

compounds are described to be useful for the treatment and/or prevention of cardiovascular and renal diseases. Yet, as outlined above, a high affinity as well as selectivity for the V1a receptor is a desirable prerequisite for the treatment of disease conditions where a decongestion is not desired and may lead to a dysregulated body fluid homeostasis including decreased blood plasma osmolality in otherwise normovolemic individuals.


An activity profile with a high selectivity for the V1a receptor has a low potential to cause unwanted off-target related side effects and would also help towards reducing the amount of substance which is going to be required to achieve and maintain the desired therapeutic effect, thus limiting the potential for unacceptable side effects and/or unwanted drug-drug interactions during the treatment of patients which might already be at high risk, such as, for example, in acute or chronic heart and kidney diseases.

The technical problem to be solved according to the present invention may therefore be seen in identifying and providing new compounds that act as potent antagonists of the vasopressin V1a receptor. A further object of the invention is to identify and provide new compounds with a high affinity and selectivity vis-à-vis the vasopressin V1a receptor. The compounds are intended to avoid inducing aquaresis via V2 inhibition. The compounds are further intended to have a similar or improved therapeutic profile compared to the compounds known from the prior art, for example with respect to their in vivo properties, for example their pharmacokinetic and pharmacodynamic characteristics and/or their metabolic profile and/or their dose-activity relationship.

Surprisingly, it has now been found that certain 5-(hydroxyalkyl)-1-heteroaryl-1,2,4-triazole derivatives represent highly potent and selective antagonists of the V1a receptor. This specific profile renders the compounds of the present invention useful for the treatment and/or prevention of diseases, which are associated with V1a receptor activation. The compounds of the present invention are particularly useful for the treatment and/or prevention of renal and cardiovascular diseases in subjects which do not suffer from fluid overload and who therefore should not be decongested.

The compounds of the present invention have valuable pharmacological properties and can be used for the prevention and/or treatment of various diseases and disease-induced states in humans and other mammals.

In one aspect, the present invention relates to 5-(hydroxyalkyl)-1-heteroaryl-1,2,4-triazole derivatives of the general formula (I)



R¹ represents a group selected from a hydrogen atom and methyl,

Ar represents a 5- or 6-membered heteroaryl group attached via a ring carbon atom having one or two ring heteroatoms selected from a nitrogen atom and a sulfur atom,

wherein any 5- or 6-membered heteroaryl group are each optionally substituted, identically or differently, with one or two groups selected from a halogen atom, nitro, cyano, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-alkylsulfanyl, (C₁-C₄)-alkoxycarbonyl, aminocarbonyl and -S(=O)₂NH₂,

wherein said (C₁-C₄)-alkyl group, said (C₁-C₄)-alkoxy group and said (C₁-C₄)-alkylsulfanyl group are each optionally substituted with up to three fluorine atoms,

The compounds according to this invention can also be present in the form of their salts, solvates and/or solvates of the salts.

The term “substituted” means that one or more hydrogen atoms on the designated atom or group are replaced with a selection from the indicated group, provided that the designated atom's normal valency under the existing circumstances is not exceeded. Combinations of substituents and/or variables are permissible.

The term “optionally substituted” means that the number of substituents can be equal to or different from zero. Unless otherwise indicated, it is possible that optionally substituted groups are substituted with as many optional substituents as can be accommodated by replacing a hydrogen atom with a non-hydrogen substituent on any available carbon atom or heteroatom.

When groups in the compounds according to the invention are substituted, it is possible for said groups to be mono-substituted or poly-substituted with substituent(s), unless otherwise specified. Within the scope of the present invention, the meanings of all groups which occur repeatedly are independent from one another. It is possible that groups in the compounds according to the invention are substituted with one, two or three identical or different substituents.

The term “comprising” when used in the specification includes “consisting of”.

If within the present text any item is referred to as “as mentioned herein”, it means that it may be mentioned anywhere in the present text.

The terms as mentioned in the present text have the following meanings:

The term “5- to 6-membered heteroaryl” means a monovalent, monocyclic aromatic ring having 5
5 or 6 ring atoms, which contains at least one ring heteroatom and optionally one, two or three further ring heteroatoms from a nitrogen atom and a sulfur atom, and which is bound via a ring carbon atom or optionally via a ring nitrogen atom (if allowed by valency).

Said heteroaryl group can be a 5-membered heteroaryl group, such as, for example, thienyl, furanyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl,
10 triazolyl, thiadiazolyl or tetrazolyl; or a 6-membered heteroaryl group, such as, for example, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl or triazinyl; or a tricyclic heteroaryl group, such as, for example, carbazolyl, acridinyl or phenazinyl.

In general, and unless otherwise mentioned, the heteroaryl or heteroarylene groups include all possible isomeric forms thereof, *e.g.*: tautomers and positional isomers with respect to the point of
15 linkage to the rest of the molecule. Thus, for some illustrative non-restricting examples, the term pyridinyl includes pyridin-2-yl, pyridin-3-yl and pyridin-4-yl; or the term thienyl includes thien-2-yl and thien-3-yl.

Particularly, the heteroaryl group in substituent Ar in the general formula (I), *supra*, is a pyridinyl, a pyrazinyl, a pyridazinyl, an imidazolyl or a thiazolyl group.

20 The term “halogen atom” means a fluorine, chlorine, bromine or iodine atom, particularly a fluorine, or chlorine atom.

The term “C₁-C₄”, as used in the present text, *e.g.* in the context of the definition of “C₁-C₄-alkyl”, “C₁-C₄-alkoxy”, “ ” or “C₁-C₄-alkylsulfanyl”, means an alkyl group having a finite number of carbon atoms of 1 to 4, *i.e.* 1, 2, 3, or 4 carbon atoms.

25 The term “C₁-C₄-alkyl” means a linear or branched, saturated, monovalent hydrocarbon group having 1, 2, 3, or 4 carbon atoms, *e.g.* a methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl, isobutyl, *tert*-butyl, or an isomer thereof. Particularly, said group has 1, 2, 3 or 4 carbon atoms (“C₁-C₄-alkyl”), *e.g.* a methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl isobutyl, or *tert*-butyl group, more particularly 1, 2 or 3 carbon atoms (“C₁-C₃-alkyl”), *e.g.* a methyl, ethyl, *n*-propyl or
30 isopropyl group, even more particularly a methyl group.

The term “C₁-C₄-alkylsulfanyl” means a linear or branched, saturated, monovalent group of formula (C₁-C₄-alkyl)-S-, in which the term “C₁-C₄-alkyl” is as defined *supra*, *e.g.* a methylsulfanyl, ethylsulfanyl, propylsulfanyl, isopropylsulfanyl, butylsulfanyl, *sec*-butylsulfanyl, isobutylsulfanyl, *tert*-butylsulfanyl group.

The term “C₁-C₄-alkoxy” means a linear or branched, saturated, monovalent group of formula (C₁-C₄-alkyl)-O-, in which the term “C₁-C₄-alkyl” is as defined *supra*, e.g. a methoxy, ethoxy, *n*-propoxy, isopropoxy, *n*-butoxy, *sec*-butoxy, isobutoxy, *tert*-butoxy, or an isomer thereof.

It is possible for the compounds of general formula (I) to exist as isotopic variants. The invention therefore includes one or more isotopic variant(s) of the compounds of general formula (I), particularly deuterium-containing compounds of general formula (I).

The term “Isotopic variant” of a compound or a reagent is defined as a compound exhibiting an unnatural proportion of one or more of the isotopes that constitute such a compound.

The term “Isotopic variant of the compound of general formula (I)” is defined as a compound of general formula (I) exhibiting an unnatural proportion of one or more of the isotopes that constitute such a compound.

The expression “unnatural proportion” means a proportion of such isotope which is higher than its natural abundance. The natural abundances of isotopes to be applied in this context are described in “Isotopic Compositions of the Elements 1997”, Pure Appl. Chem., 70(1), 217-235, 1998.

Examples of such isotopes include stable and radioactive isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, sulfur, fluorine, chlorine, bromine and iodine, such as ²H (deuterium), ³H (tritium), ¹¹C, ¹³C, ¹⁴C, ¹⁵N, ¹⁷O, ¹⁸O, ³²P, ³³P, ³³S, ³⁴S, ³⁵S, ³⁶S, ¹⁸F, ³⁶Cl, ⁸²Br, ¹²³I, ¹²⁴I, ¹²⁵I, ¹²⁹I and ¹³¹I, respectively.

With respect to the treatment and/or prevention of the disorders specified herein the isotopic variant(s) of the compounds of general formula (I) preferably contain deuterium (“deuterium-containing compounds of general formula (I)”). Isotopic variants of the compounds of general formula (I) in which one or more radioactive isotopes, such as ³H or ¹⁴C, are incorporated are useful e.g. in drug and/or substrate tissue distribution studies. These isotopes are particularly preferred for the ease of their incorporation and detectability. Positron emitting isotopes such as ¹⁸F or ¹¹C may be incorporated into a compound of general formula (I). These isotopic variants of the compounds of general formula (I) are useful for in vivo imaging applications. Deuterium-containing and ¹³C-containing compounds of general formula (I) can be used in mass spectrometry analyses in the context of preclinical or clinical studies.

Isotopic variants of the compounds of general formula (I) can generally be prepared by methods known to a person skilled in the art, such as those described in the schemes and/or examples herein, by substituting a reagent for an isotopic variant of said reagent, preferably for a deuterium-containing reagent. Depending on the desired sites of deuteration, in some cases deuterium from D₂O can be incorporated either directly into the compounds or into reagents that are useful for

synthesizing such compounds. Deuterium gas is also a useful reagent for incorporating deuterium into molecules. Catalytic deuteration of olefinic bonds and acetylenic bonds is a direct route for incorporation of deuterium. Metal catalysts (i.e. Pd, Pt, and Rh) in the presence of deuterium gas can be used to directly exchange deuterium for hydrogen in functional groups containing hydrocarbons. A variety of deuterated reagents and synthetic building blocks are commercially available from companies such as for example C/D/N Isotopes, Quebec, Canada; Cambridge Isotope Laboratories Inc., Andover, MA, USA; and CombiPhos Catalysts, Inc., Princeton, NJ, USA.

The term “deuterium-containing compound of general formula (I)” is defined as a compound of general formula (I), in which one or more hydrogen atom(s) is/are replaced by one or more deuterium atom(s) and in which the abundance of deuterium at each deuterated position of the compound of general formula (I) is higher than the natural abundance of deuterium, which is about 0.015%. Particularly, in a deuterium-containing compound of general formula (I) the abundance of deuterium at each deuterated position of the compound of general formula (I) is higher than 10%, 20%, 30%, 40%, 50%, 60%, 70% or 80%, preferably higher than 90%, 95%, 96% or 97%, even more preferably higher than 98% or 99% at said position(s). It is understood that the abundance of deuterium at each deuterated position is independent of the abundance of deuterium at other deuterated position(s).

The selective incorporation of one or more deuterium atom(s) into a compound of general formula (I) may alter the physicochemical properties (such as for example acidity [C. L. Perrin, et al., J. Am. Chem. Soc., 2007, 129, 4490], basicity [C. L. Perrin et al., J. Am. Chem. Soc., 2005, 127, 9641], lipophilicity [B. Testa et al., Int. J. Pharm., 1984, 19(3), 271]) and/or the metabolic profile of the molecule and may result in changes in the ratio of parent compound to metabolites or in the amounts of metabolites formed. Such changes may result in certain therapeutic advantages and hence may be preferred in some circumstances. Reduced rates of metabolism and metabolic switching, where the ratio of metabolites is changed, have been reported (A. E. Mutlib et al., Toxicol. Appl. Pharmacol., 2000, 169, 102). These changes in the exposure to parent drug and metabolites can have important consequences with respect to the pharmacodynamics, tolerability and efficacy of a deuterium-containing compound of general formula (I). In some cases deuterium substitution reduces or eliminates the formation of an undesired or toxic metabolite and enhances the formation of a desired metabolite (e.g. Nevirapine: A. M. Sharma et al., Chem. Res. Toxicol., 2013, 26, 410; Efavirenz: A. E. Mutlib et al., Toxicol. Appl. Pharmacol., 2000, 169, 102). In other cases the major effect of deuteration is to reduce the rate of systemic clearance. As a result, the biological half-life of the compound is increased. The potential clinical benefits would include the ability to maintain similar systemic exposure with decreased peak levels and increased trough levels. This could result in lower side effects and enhanced efficacy, depending on the particular

compound's pharmacokinetic/ pharmacodynamic relationship. ML-337 (C. J. Wenthur et al., J. Med. Chem., 2013, 56, 5208) and Odanacatib (K. Kassahun et al., WO2012/112363) are examples for this deuterium effect. Still other cases have been reported in which reduced rates of metabolism result in an increase in exposure of the drug without changing the rate of systemic clearance (e.g. Rofecoxib: F. Schneider et al., *Arzneim. Forsch. / Drug. Res.*, 2006, 56, 295; Telaprevir: F. Maltais et al., J. Med. Chem., 2009, 52, 7993). Deuterated drugs showing this effect may have reduced dosing requirements (e.g. lower number of doses or lower dosage to achieve the desired effect) and/or may produce lower metabolite loads.

A compound of general formula (I) may have multiple potential sites of attack for metabolism. To optimize the above-described effects on physicochemical properties and metabolic profile, deuterium-containing compounds of general formula (I) having a certain pattern of one or more deuterium-hydrogen exchange(s) can be selected. Particularly, the deuterium atom(s) of deuterium-containing compound(s) of general formula (I) is/are attached to a carbon atom and/or is/are located at those positions of the compound of general formula (I), which are sites of attack for metabolizing enzymes such as e.g. cytochrome P₄₅₀.

Where the plural form of the word compounds, salts, polymorphs, hydrates, solvates and the like, is used herein, this is taken to mean also a single compound, salt, polymorph, isomer, hydrate, solvate or the like.

By "stable compound" or "stable structure" is meant a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

The compounds of the present invention optionally contain one asymmetric centre, depending upon the location and nature of the various substituents desired. It is possible that one asymmetric carbon atom is present in the (R) or (S) configuration, which can result in racemic mixtures. In certain instances, it is possible that asymmetry also be present due to restricted rotation about a given bond, for example, the central bond adjoining two substituted aromatic rings of the specified compounds. Preferred compounds are those which produce the more desirable biological activity. Separated, pure or partially purified isomers and stereoisomers or racemic mixtures of the compounds of the present invention are also included within the scope of the present invention. The purification and the separation of such materials can be accomplished by standard techniques known in the art.

The optical isomers can be obtained by resolution of the racemic mixtures according to conventional processes, for example, by the formation of diastereoisomeric salts using an optically active acid or base or formation of covalent diastereomers. Examples of appropriate acids are

tartaric, diacetyltartaric, ditoluoyltartaric and camphorsulfonic acid. Mixtures of diastereoisomers can be separated into their individual diastereomers on the basis of their physical and/or chemical differences by methods known in the art, for example, by chromatography or fractional crystallisation. The optically active bases or acids are then liberated from the separated diastereomeric salts. A different process for separation of optical isomers involves the use of chiral chromatography (*e.g.*, HPLC columns using a chiral phase), with or without conventional derivatisation, optimally chosen to maximise the separation of the enantiomers. Suitable HPLC columns using a chiral phase are commercially available, such as those manufactured by Daicel, *e.g.*, Chiracel OD and Chiracel OJ, for example, among many others, which are all routinely selectable. Enzymatic separations, with or without derivatisation, are also useful. The optically active compounds of the present invention can likewise be obtained by chiral syntheses utilizing optically active starting materials. In order to distinguish different types of isomers from each other reference is made to IUPAC Rules Section E (Pure Appl Chem 45, 11-30, 1976).

The present invention includes all possible stereoisomers of the compounds of the present invention as single stereoisomers, or as any mixture of said stereoisomers, *e.g.* (R)- or (S)- isomers, in any ratio. Isolation of a single stereoisomer, *e.g.* a single enantiomer or a single diastereomer, of a compound of the present invention is achieved by any suitable state of the art method, such as chromatography, especially chiral chromatography, for example.

Further, it is possible for the compounds of the present invention to exist as tautomers. The present invention includes all possible tautomers of the compounds of the present invention as single tautomers, or as any mixture of said tautomers, in any ratio.

Further, the compounds of the present invention can exist as N-oxides, which are defined in that at least one nitrogen of the compounds of the present invention is oxidised. The present invention includes all such possible N-oxides.

The present invention also covers useful forms of the compounds of the present invention, such as metabolites, hydrates, solvates, salts, in particular pharmaceutically acceptable salts, and/or co-precipitates.

The compounds of the present invention can exist as a hydrate, or as a solvate, wherein the compounds of the present invention contain polar solvents, in particular water, methanol or ethanol for example, as structural element of the crystal lattice of the compounds. It is possible for the amount of polar solvents, in particular water, to exist in a stoichiometric or non-stoichiometric ratio. In the case of stoichiometric solvates, *e.g.* a hydrate, hemi-, (semi-), mono-, sesqui-, di-, tri-, tetra-, penta- *etc.* solvates or hydrates, respectively, are possible. The present invention includes all such hydrates or solvates. Hydrates are preferred solvates in the context of the present invention.

Further, it is possible for the compounds of the present invention to exist in free form, *e.g.* as a free base, or as a free acid, or as a zwitterion, or to exist in the form of a salt. Said salt may be any salt, either an organic or inorganic addition salt, particularly any pharmaceutically acceptable organic or inorganic addition salt, which is customarily used in pharmacy, or which is used, for example, for
5 isolating or purifying the compounds of the present invention.

The term "pharmaceutically acceptable salt" refers to an inorganic or organic acid addition salt of a compound of the present invention. For example, see S. M. Berge, *et al.* "Pharmaceutical Salts," J. Pharm. Sci. 1977, 66, 1-19.

A suitable pharmaceutically acceptable salt of the compounds of the present invention may be, for
10 example, an acid-addition salt of a compound of the present invention bearing a nitrogen atom, in a chain or in a ring, for example, which is sufficiently basic, such as an acid-addition salt with an inorganic acid, or "mineral acid", such as hydrochloric, hydrobromic, hydroiodic, sulfuric, sulfamic, bisulfuric, phosphoric, or nitric acid, for example, or with an organic acid, such as formic, acetic, acetoacetic, pyruvic, trifluoroacetic, propionic, butyric, hexanoic, heptanoic, undecanoic,
15 lauric, benzoic, salicylic, 2-(4-hydroxybenzoyl)-benzoic, camphoric, cinnamic, cyclopentanepropionic, digluconic, 3-hydroxy-2-naphthoic, nicotinic, pamoic, pectinic, 3-phenylpropionic, pivalic, 2-hydroxyethanesulfonic, itaconic, trifluoromethanesulfonic, dodecylsulfuric, ethanesulfonic, benzenesulfonic, para-toluenesulfonic, methanesulfonic, 2-naphthalenesulfonic, naphthalenedisulfonic, camphorsulfonic acid, citric, tartaric, stearic, lactic,
20 oxalic, malonic, succinic, malic, adipic, alginic, maleic, fumaric, D-gluconic, mandelic, ascorbic, glucoheptanoic, glycerophosphoric, aspartic, sulfosalicylic, or thiocyanic acid, for example.

Further, another suitably pharmaceutically acceptable salt of a compound of the present invention which is sufficiently acidic, is an alkali metal salt, for example a sodium or potassium salt, an
25 alkaline earth metal salt, for example a calcium, magnesium or strontium salt, or an aluminium or a zinc salt, or an ammonium salt derived from ammonia or from an organic primary, secondary or tertiary amine having 1 to 20 carbon atoms, such as ethylamine, diethylamine, triethylamine, ethyldiisopropylamine, monoethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, dimethylaminoethanol, diethylaminoethanol, tris(hydroxymethyl)aminomethane, procaine,
30 dibenzylamine, *N*-methylmorpholine, arginine, lysine, 1,2-ethylenediamine, *N*-methylpiperidine, *N*-methyl-glucamine, *N,N*-dimethyl-glucamine, *N*-ethyl-glucamine, 1,6-hexanediamine, glucosamine, sarcosine, serinol, 2-amino-1,3-propanediol, 3-amino-1,2-propanediol, 4-amino-1,2,3-butanetriol, or a salt with a quarternary ammonium ion having 1 to 20 carbon atoms, such as tetramethylammonium, tetraethylammonium, tetra(*n*-propyl)ammonium, tetra(*n*-butyl)ammonium,
35 *N*-benzyl-*N,N,N*-trimethylammonium, choline or benzalkonium.

Those skilled in the art will further recognise that it is possible for acid addition salts of the claimed compounds to be prepared by reaction of the compounds with the appropriate inorganic or organic acid via any of a number of known methods. Alternatively, alkali and alkaline earth metal salts of acidic compounds of the present invention are prepared by reacting the compounds of the present invention with the appropriate base via a variety of known methods.

The present invention includes all possible salts of the compounds of the present invention as single salts, or as any mixture of said salts, in any ratio.

In the present text, in particular in the Experimental Section, for the synthesis of intermediates and of examples of the present invention, when a compound is mentioned as a salt form with the corresponding base or acid, the exact stoichiometric composition of said salt form, as obtained by the respective preparation and/or purification process, is, in most cases, unknown.

Unless specified otherwise, suffixes to chemical names or structural formulae relating to salts, such as "hydrochloride", "trifluoroacetate", "sodium salt", or "x HCl", "x CF₃COOH", "x Na⁺", for example, mean a salt form, the stoichiometry of which salt form not being specified.

This applies analogously to cases in which synthesis intermediates or example compounds or salts thereof have been obtained, by the preparation and/or purification processes described, as solvates, such as hydrates, with (if defined) unknown stoichiometric composition.

Furthermore, the present invention includes all possible crystalline forms, or polymorphs, of the compounds of the present invention, either as single polymorph, or as a mixture of more than one polymorph, in any ratio.

In a distinct embodiment, the present invention relates to compounds of formula (I), *supra*, wherein

R¹ represents a methyl group,

Ar represents a group selected from pyridine, pyrazine, pyridazine, 1-methylimidazole, 1,3-thiazole,

wherein any pyridine group is each optionally substituted, identically or differently, with one or two groups selected from a fluorine atom, a chlorine atom, methoxy and trifluoromethoxy,

wherein any pyrazine group is optionally substituted with methoxy,

or a pharmaceutically acceptable salt, hydrate and/or solvate thereof.

In a distinct embodiment, the present invention relates to compounds of formula (I), *supra*, wherein

R¹ represents a methyl group,

Ar represents a group selected from pyridine, pyrazine, pyridazine, 1-methylimidazole, 1,3-thiazole,

wherein any pyridine group is each optionally substituted, identically or differently, with one or two groups selected from a fluorine atom, a chlorine atom, methoxy, trifluoromethyl and trifluoromethoxy,


wherein any pyrazine group is optionally substituted with methoxy,

5 or a pharmaceutically acceptable salt, hydrate and/or solvate thereof.

In a preferred embodiment, the present invention relates to compounds according to formula (I), *supra*, wherein

R¹ represents a methyl group,

Ar represents a group of the formula



in which

#¹ represents the point of attachment to the nitrogen atom,

R^{2A} represents a group selected from a hydrogen atom, a chlorine atom and trifluoromethoxy,

15 R^{2B} represents a group selected from a hydrogen atom and a chlorine atom


R^{2C} represents a chlorine atom,

or a pharmaceutically acceptable salt, hydrate and/or solvate thereof.

In another preferred embodiment, the present invention relates to compounds according to formula (I), *supra*, wherein

20 R¹ represents a methyl group,

Ar represents a group of the formula



in which

#¹ represents the point of attachment to the nitrogen atom,

25 R^{2A} represents a group selected from a hydrogen atom, a chlorine atom, trifluoromethyl and trifluoromethoxy,


R^{2B} represents a group selected from a hydrogen atom and a chlorine atom

R^{2C} represents a chlorine atom,

or a pharmaceutically acceptable salt, hydrate and/or solvate thereof.

In accordance with a further preferred embodiment, the present invention covers compounds of
5 general formula (I), *supra*, wherein

R¹ represents a (2*S*)-3,3,3-trifluoro-2-hydroxypropyl group of the formula




in which

#¹ represents the point of attachment to the nitrogen atom,

10 or a pharmaceutically acceptable salt, hydrate and/or solvate thereof.

R¹ represents a (2*R*)-3,3,3-trifluoro-2-hydroxypropyl group of the formula



in which

#¹ represents the point of attachment to the nitrogen atom,

15 or a pharmaceutically acceptable salt, hydrate and/or solvate thereof.

In a particular further embodiment of the first aspect, the present invention covers combinations of two or more of the above mentioned embodiments under the heading “further embodiments of the first aspect of the present invention”.


The present invention covers any sub-combination within any embodiment or aspect of the present
20 invention of compounds of general formula (I), *supra*.

The present invention covers any sub-combination within any embodiment or aspect of the present invention of intermediate compounds of general formula (II), (III), (IV), (V), (VI), (VII), (XV).

The present invention covers the compounds of general formula (I) which are disclosed in the Example Section of this text, *infra*.

25 In accordance with a second aspect, the present invention covers methods of preparing compounds of general formula (I) as defined *supra*, said methods comprising the step


[A] of allowing an intermediate compound of formula (II):



in which

R³ represents a (C₁-C₄)-alkyl group, in particular a methyl group,

to react in a first step in the presence of an at least stoichiometric amount of a base with a
 5 compound of general formula (III):




in which

R¹ is as defined for the compound of general formula (I) as defined supra,

PG represents a suitable alcohol protecting group, such as acetate,


10 to give an intermediate compound, which is then allowed to react in a second step with a hydrazine compound of general formula (IV) or a respective salt thereof



in which

Ar is as defined for the compound of general formula (I) as defined supra,

15 thereby giving an intermediate compound of general formula (V):



in which

R¹ and Ar is as defined for the compound of general formula (I) as defined supra,


PG represents a suitable alcohol protecting group, such as acetyl

- 5 followed by alcohol deprotection of (V) with a suitable deprotection agent thereby giving a compound of general formula (I) in which

R¹ and Ar is as defined for the compound of general formula (I) as defined supra,

or

[B] allowing an intermediate compound of formula (VI):




10

in which,

R¹ is as defined for the compound of general formula (I) as defined supra,

and/or a tautomer thereof to react with a compound of general formula (VII):



15

in which

Ar is as defined for the compound of general formula (I) as defined supra,

R^{XA} and R^{XB} represent, independently from each other, a hydrogen atom, or

R^{XA} and R^{XB} together form a $-CH_2CH_2-$ or $-C(CH_3)_2C(CH_3)_2-$ bridge,

in the presence of a copper catalyst and an amine base thereby giving a compound of general formula (I) in which

R^1 and Ar are as defined for the compound of general formula (I) as defined supra,

5 each [A] and [B] optionally followed, where appropriate, by (i) separating the compounds of formula (I) thus obtained into their respective diastereomers, and/or (ii) converting the compounds of formula (I) into their respective hydrates, solvates, salts and/or hydrates or solvates of the salts by treatment with the corresponding solvents and/or acids or bases.

The present invention covers methods of preparing compounds of the present invention of general
10 formula (I), said methods comprising the steps as described in the Experimental Section herein.

The schemes and procedures described below illustrate synthetic routes to the compounds of general formula (I) of the invention and are not intended to be limiting. It is clear to the person skilled in the art that the order of transformations as exemplified in schemes 1, 2, 3, 4, 5, 6, 7 and 8 can be modified in various ways. The order of transformations exemplified in these schemes is
15 therefore not intended to be limiting. In addition, interconversion of any of the substituents R^1 , R^2 , PG, R^{XA} , R^{XB} , X and Ar can be achieved before and/or after the exemplified transformations. These modifications can be such as the introduction of protecting groups, cleavage of protecting groups, reduction or oxidation of functional groups, halogenation, metallation, substitution or other reactions known to the person skilled in the art. These transformations include those which
20 introduce a functionality which allows for further interconversion of substituents. Appropriate protecting groups and their introduction and cleavage are well-known to the person skilled in the art (see for example T.W. Greene and P.G.M. Wuts in *Protective Groups in Organic Synthesis*, 3rd edition, Wiley 1999). Specific examples are described in the subsequent paragraphs.

The multicomponent cyclization (II) \rightarrow (V) is carried out by first reacting imidate of formula (II)
25 with an acid chloride of formula (III) in the presence of a base to form an intermediate which is in a subsequent step reacted with the hydrazine compound of formula (IV). Typically the formed intermediate is not isolated and the reaction over the two steps is performed in one-pot. The hydrazine compound of formula (IV) may also be used in form of its salts, such as a hydrochloride salt or a tosylate salt. Under the alkaline reaction conditions, the hydrazine salt will be reconverted
30 into the free base form. The amount of base added may then be adjusted in this respect. It may be beneficial in the second step to add a copper or zinc salt, such as copper(II) sulfate, copper(II) chloride, zinc(II) sulfate and zinc(II) chloride. Typically and preferably copper(II)sulfate is used.


Suitable bases for both steps of the cyclization are typically tertiary amine bases, such as *N,N*-diisopropylethylamine (DIPEA), triethylamine, triisopropylamine, *N*-methylimidazole, *N*-methylmorpholine,
35 pyridine and 4-(*N,N*-dimethylamino)pyridine. Preferably, *N,N*-

diisopropylethylamine (DIPEA) is used as base. The reaction is performed in an inert organic solvent, such as dichloromethane, 1,2-dichloroethane, methyl *tert*-butyl ether, tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane, toluene, pyridine, ethyl acetate, acetonitrile or *N,N*-dimethylformamide, or in a mixture of these solvents. Preferably tetrahydrofuran or dioxane or a mixture thereof are used as solvents. The first step is generally carried out at a temperature in the range of -10°C to +120°C, preferably at 0°C. The second step is generally carried out at a temperature in the range of +20°C to +120°C, preferably at room temperature. Concomitant microwave irradiation may have a beneficial effect in this reaction as well at a temperature in the range of +60°C to +150°C, preferably at +120°C.

Introduction and removal of the protective group (PG) (V) → (I) is carried out by customary methods known from the literature [see, for example, T.W. Greene and P.G.M. Wuts, *Protective Groups in Organic Synthesis*, Wiley, New York, 1999]. Thus, the acetyl group is preferably removed with the aid of a base, such as an aqueous sodium hydroxide solution.

The coupling reaction (VI) + (VII) → (I) is typically carried out with the aid of a copper catalyst and an amine base ["Chan-Lam coupling" conditions; see, for instance, D. M. T. Chan et al., *Tetrahedron Lett.* **44** (19), 3863-3865 (2003); J. X. Qiao and P. Y. S. Lam, *Synthesis*, 829-856 (2011); K. S. Rao and T.-S. Wu, *Tetrahedron* **68**, 7735-7754 (2012)]. Copper catalysts suitable for this process are in particular copper(II) salts, such as copper(II) acetate, copper(II) trifluoromethanesulfonate or copper(II) bromide. Practical amine bases include, for example, triethylamine, *N,N*-diisopropylethylamine, pyridine and 4-(*N,N*-dimethylamino)pyridine. The reaction is performed in an inert organic solvent, such as dichloromethane, 1,2-dichloroethane, methyl *tert*-butyl ether, tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane, toluene, pyridine, ethyl acetate, acetonitrile or *N,N*-dimethylformamide, or in a mixture of these solvents. Preferably, pyridine is used both as solvent and base. The coupling is generally carried out at a temperature in the range of +20°C to +120°C, preferably at +20°C to +70°C. Concomitant microwave irradiation may have a beneficial effect in this reaction as well.

Compounds of formula (I), wherein R¹ represents methyl, can also be obtained in diastereomerically pure form by employing the appropriate enantiomer of acyl chloride (III) [R¹ = methyl], *i.e.* (III-A) or (III-B)




30


or a salt thereof, in the condensation reaction described above.

Compounds of general formula (II) as defined *supra*, can be prepared by a method comprising the step

[a] of allowing a compound of formula (VIII):




5 to react with a nitrile compound of general formula (IX),



in which

X represents a leaving group, such as chlorine, bromine, iodine, mesylate or tosylate, in particular chlorine or bromine,


10 thereby giving a compound of general formula (X)



followed by a subsequent step

[b] of allowing the compound of formula (X) obtained in step [a] to react with a basic alcoholate, preferably sodium methanolate, thereby giving a compound of general formula (II),

15



in which


R^3 represents a (C₁-C₄)-alkyl group, in particular a methyl group.

The *N*-alkylation reaction (VIII) + (IX) → (X) (step [a]) is typically carried out in the presence of a base. Typical and exemplary bases include sodium carbonate, potassium carbonate, cesium carbonate, *N,N*-diisopropylethylamine, triethylamine, sodium *tert*-butylate or potassium *tert*-butylate in acetonitrile, methylisobutylketone, dioxane, dimethylformamide, dimethylacetamide, *N*-methylpyrrolidinone, dimethylsulfoxide and sulfolane, preference is given to potassium carbonate in methylisobutylketone or acetonitrile. The reaction may optionally be carried out in an advantageous manner with addition of an alkylation catalyst such as, for example, lithium bromide, sodium iodide, lithium iodide, tetra-*n*-butylammoniumbromide, tetra-*n*-butylammoniumiodide or benzyltriethylammoniumchloride. The reactions are generally carried out in a temperature range of from +40°C to +120°C, preferably at from +60°C to +80°C. The reactions can be carried out at atmospheric, at elevated or at reduced pressure (for example at from 0.5 to 5 bar); in general, the reactions are carried out at atmospheric pressure. It may be advantageous to slowly perform the addition of the alkylation agent (IX) over a longer time span.

Conversion to the imidates of general formula (II) can be achieved via standard reaction protocols known to the person skilled in the art (step [b]: (X) → (II)). The reaction is typically carried out under basic reactions conditions by reacting with a basic alcoholate. Typical bases, which may be used are sodium methanolate, sodium ethanolate, sodium propanolate, sodium isopropoxide, sodium *tert*-butylate or potassium *tert*-butylate in methanol, ethanol, *n*-propanol, isopropanol, *n*-butanol, isobutanol and *tert*-butanol. Preference is given to sodium methanolate in methanol. The reactions are generally carried out in a temperature range of from +20°C to +80°C, preferably at from +20°C to +60 °C.

Alternatively, the nitrile compounds of general formula (X) may optionally also be prepared as shown in the synthetic scheme 1 below:

Scheme 1:



TFAA = trifluoroacetic acid anhydride

The amide coupling (XI) → (XII) can be carried out directly with the help of a condensing agent or activating agent in the presence of a base or over two steps via an acyl chloride or carboxylic acid imidazolide. Typical condensation and activating agents for the amide formation in process steps (XI) → (XII) include, for example, carbodiimides such as *N,N'*-diethyl-, *N, N'*-dipropyl-, *N,N'*-diisopropyl-*N,N'*-dicyclohexylcarbodiimide (DCC) or *N*-(3-dimethylaminoisopropyl)-*N'*-ethylcarbodiimide hydrochloride (EDC), phosgene derivatives such as *N, N'*-carbonyldiimidazole (CDI), 1,2-oxazolium compounds such as 2-ethyl-5-phenyl-1,2-oxazolium-3-sulphate or 2-tert-butyl-5-methyl-isoxazolium perchlorate, acylamino compounds such as 2-ethoxy-1-ethoxy-carbonyl-1,2-dihydroquinoline, or isobutyl chloroformate, propanephosphonic anhydride, diethyl cyanophosphonate, bis(2-oxo-3-oxazolidinyl)phosphoryl chloride, benzotriazol-1-yloxytris-(dimethylamino)phosphonium hexafluorophosphate, benzotriazol-1-yloxytris(pyrrolidino)-phosphonium hexafluorophosphate (PyBOP), *O*-(benzotriazol-1-yl)-*N, N, N', N'*-tetramethyluronium tetrafluoroborate (TBTU), *O*-(benzotriazol-1-yl)-*N, N, N', N'*-tetramethyluronium hexafluorophosphate (HBTU), 2-(2-oxo-1-(2H)-pyridyl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TPTU), *O*-(7-azabenzotriazol-1-yl)-*N, N, N', N'*-tetramethyluroniumhexafluorophosphate (HATU) or *O*-(1H-6-chloro-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TCTU), optionally in combination with other additives such as 1-hydroxybenzotriazole (HOBt) or *N*-hydroxysuccinimide (HOSu). The acyl chlorides are typically prepared with thionyl chloride or oxalyl chloride in an inert solvent like dichloromethane or *N,N*-dimethylformamide. It is also possible to use mixtures of the solvents mentioned

Typical and exemplary bases include sodium carbonate, potassium carbonate, cesium carbonate, *N,N*-diisopropylethylamine, triethylamine, sodium *tert*-butylate or potassium *tert*-butylate in acetonitrile, methylisobutylketone, dioxane, dimethylformamide, dimethylacetamide, *N*-methylpyrrolidinone, dimethylsulfoxide and sulfolane, preference is given to potassium carbonate in methylisobutylketone or acetonitrile. The reaction may optionally be carried out in an advantageous manner with addition of an alkylation catalyst such as, for example, lithium bromide, sodium iodide, lithium iodide, tetra-*n*-butylammoniumbromide, tetra-*n*-butylammoniumiodide or benzyltriethylammoniumchloride. The conversion to the nitrile (XII) → (X) can be carried out with the help of dehydrating agent. Typical dehydrating agents include, for example trifluoroacetic acid


anhydride, phosphorous pentoxide (P₄O₁₀), phosphoryl chloride (POCl₃), phosphorous pentachloride (PCl₅), CCl₄-PPh₃ (Appel reagent), hexamethylphosphoramide (HMPA); methyl *N*-(triethylammoniumsulfonyl)carbamate (Burgess reagent), (Chloromethylene)dimethyliminium chloride (Vilsmeier reagent), oxalyl chloride/DMSO and thionylchloride (SOCl₂).

- 5 Typical and exemplary solvents for both steps (XI) → (XII) and (XII) → (X) include for example, ethers such as diethyl ether, dioxane, tetrahydrofuran, glycol dimethyl ether or diethylene glycol dimethyl ether, hydrocarbons such as benzene, toluene, xylene, hexane, cyclohexane or mineral oil, fractions, halogenated hydrocarbons such as dichloromethane, trichloromethane, carbon tetrachloride, 1,2-dichloroethane, trichloroethylene or chlorobenzene, or other solvents such as
 10 acetone, ethyl acetate, acetonitrile, pyridine, dimethyl sulphoxide, N,N-dimethylformamide, N,N'-dimethylpropyleneurea (DMPU) or N-methylpyrrolidone (NMP). It is also possible to use mixtures of the solvents mentioned.


In a typical and preferred procedure the carboxylic acid (XI) is first reacted with pivaloyl chloride in the presence of pyridine to form an intermediate which is in a subsequent step reacted with
 15 ammonia. Typically the formed intermediate is not isolated and the reaction over the two steps is performed in one-pot. Suitable as bases for the first step are preferably, pyridine, 4-(N,N-dimethylamino)pyridine or N,N-diisopropylethylamine (DIPEA). The conversion of carboxamide (XII) into nitrile (X) is then typically performed by reaction with trifluoroacetic anhydride. Both reactions are conducted in an inert organic solvent, preferably tetrahydrofuran.

- 20 Compounds of general formula (VI) as defined *supra*, can be prepared by a method comprising the step


[a] of allowing a compound of formula (XIII):



to first react with hydrazine to give the hydrazine of formula (XIV)



which is then being condensed with an amidine of formula (XV)




or a salt thereof,

5 in which

R¹ is as defined for the compound of general formula (I) as defined supra,

in the presence of a base to give a 1,2,4-triazole derivative of formula (VI)



and/or a tautomer thereof,

10 in which


R¹ is as defined for the compound of general formula (I) as defined supra,

The transformation (XIII) → (XIV) is carried out in the usual way by treating methyl ester (II) with hydrazine or hydrazine hydrate in an alcoholic solvent, such as methanol, ethanol, *n*-propanol, isopropanol or *n*-butanol, at a temperature in the range of +20°C to +100°C.

15 The condensation reaction (XIV) + (XV) → (VI) is usually carried out in an inert dipolar-aprotic solvent, such as *N,N*-dimethylformamide (DMF), *N,N*-dimethylacetamide (DMA), dimethylsulfoxide (DMSO), *N*-methylpyrrolidinone (NMP) or *N,N'*-dimethylpropylene urea (DMPU), in the


presence of a sufficiently strong base, such as sodium hydride or a sodium or potassium alkoxide, for example sodium or potassium methoxide, sodium or potassium ethoxide, or sodium or potassium *tert*-butoxide. The amidine (XV) may be employed as such in this reaction or in salt form, *e.g.* as the hydrochloride salt. In the latter case, a proportional excess of base is used. The reaction is generally performed at a temperature between +80°C and +150°C. Heating by means of a microwave reactor device may have a beneficial effect for this condensation reaction.

The 1,2,4-triazole derivative of formula (VI) produced by this reaction may also be present in other tautomeric forms, such as (VI-A) or (VI-B)



or as a mixture of tautomers.

Compounds of formula (I), wherein R¹ represents methyl, can also be obtained in diastereomerically pure form by employing the appropriate enantiomer of amidine (XV) [R¹ = methyl], *i.e.* (XV-A) or (XV-B)




or a salt thereof, in the condensation reaction described above.

The compounds of formula (VIII), (XI) and (XIII) can be synthesized by the procedures described in Int. Pat. Appl. WO 2010/105770 and WO 2011/104322 (see also synthesis schemes 2a, 2b and 3 below).


The compounds of the formula III, IV, VII, IX and XV are either commercially available, known from the literature, or can be prepared from readily available starting materials by adaptation of standard methods described in the literature. Detailed procedures and literature references for preparing the starting materials can also be found in the Experimental Part in the section on the preparation of the starting materials and intermediates.

The preparation of the compounds of the invention may be illustrated by means of the following synthesis schemes:

Scheme 2


5 [cf. Int. Pat. Appl. WO 2011/104322-A1].

Scheme 3



[cf. Int. Pat. Appl. WO 2011/104322-A1].


Scheme 4a




5

TFAA = trifluoroacetic acid anhydride


Scheme 4b




Scheme 5




Scheme 6




5 Scheme 7a



Scheme 7b



Scheme 8



The compounds of general formula (I) of the present invention can be converted to any salt, preferably pharmaceutically acceptable salts, as described herein, by any method which is known to the person skilled in the art. Similarly, any salt of a compound of general formula (I) of the present invention can be converted into the free compound, by any method which is known to the person skilled in the art.

The compounds of the present invention have valuable pharmacological properties and can be used for the prevention and/or treatment of various diseases and disease-induced states in humans and other mammals. Compounds of general formula (I) of the present invention demonstrate a valuable pharmacological spectrum of action and pharmacokinetic profile, both of which could not have been predicted. Compounds of the present invention have surprisingly been found to effectively inhibit the vasopressin V1a receptor and it is possible therefore that said compounds be used for the

treatment and/or prevention of diseases, preferably renal and cardiovascular diseases in humans and animals.

In the context of the present invention, the term "treatment" or "treating" includes inhibiting, delaying, relieving, mitigating, arresting, reducing, or causing the regression of a disease, disorder, condition, or state, the development and/or progression thereof, and/or the symptoms thereof. The term "prevention" or "preventing" includes reducing the risk of having, contracting, or experiencing a disease, disorder, condition, or state, the development and/or progression thereof, and/or the symptoms thereof. The term prevention includes prophylaxis. Treatment or prevention of a disorder, disease, condition, or state may be partial or complete.

Throughout this document, for the sake of simplicity, the use of singular language is given preference over plural language, but is generally meant to include the plural language if not otherwise stated. For example, the expression "A method of treating a disease in a patient, comprising administering to a patient an effective amount of a compound of formula (I)" is meant to include the simultaneous treatment of more than one disease as well as the administration of more than one compound of formula (I).

The compounds of the present invention are highly potent and in particular selective antagonists of the vasopressin V1a receptor. The compounds of the invention are therefore expected to be highly valuable as therapeutic agents for the treatment and/or prevention of diseases, in particular for the treatment and/or prevention of renal and cardiovascular diseases.

As used herein, the term "vasopressin V1a receptor antagonist" refers to a compound that functions by inhibiting (partially or completely) or blocking the vasopressin V1a receptor, thereby preventing activation of the receptor by vasopressin.

In one embodiment, compounds described herein are active at the V1a receptor. In another embodiment compounds described herein exhibit inhibition of the V1a receptor according to the study in B-1 with an $IC_{50} < 100$ nM. In another embodiment compounds described herein exhibit inhibition of the V1a receptor according to the study in B-1 with an $IC_{50} < 20$ nM. In another embodiment compounds described herein exhibit inhibition of the V1a receptor according to the study in B-1 with an $IC_{50} < 10$ nM. In another embodiment compounds described herein exhibit inhibition of the V1a receptor according to the study in B-1 with an $IC_{50} < 5$ nM. In another embodiment compounds described herein exhibit inhibition of the V1a receptor according to the study in B-1 with an $IC_{50} < 1$ nM.

In a further embodiment, compounds described herein are selectively active at the V1a receptor, and are less active, substantially less active, and/or inactive at other vasopressin receptors, such as the V1b and/or V2 subtypes. In another embodiment, compounds described herein are at least 10-fold selective for the V1a receptor compared to the V2 receptor as determined according to the

study in B-1. In another embodiment, compounds described herein are at least 15-fold selective for the V1a receptor compared to the V2 receptor as determined according to the study in B-1. In another embodiment, compounds described herein are at least 20-fold selective for the V1a receptor compared to the V2 receptor as determined according to the study in B-1. In another
5 embodiment, compounds described herein are at least 30-fold selective for the V1a receptor compared to the V2 receptor as determined according to the study in B-1.

The compounds according to the invention are suitable for the treatment and/or prevention of renal diseases, in particular of acute and chronic kidney diseases, diabetic kidney diseases, and of acute and chronic renal failure. The general terms 'renal disease' or 'kidney disease' describe a class of
10 conditions in which the kidneys fail to filter and remove waste products from the blood. There are two major forms of kidney disease: acute kidney disease (acute kidney injury, AKI) and chronic kidney disease (CKD). The compounds according to the invention may further be used for the treatment and/or prevention of sequelae of acute kidney injury arising from multiple insults such as ischemia-reperfusion injury, radiocontrast administration, cardiopulmonary bypass surgery, shock
15 and sepsis. In the sense of the present invention, the term renal failure or renal insufficiency comprises both acute and chronic manifestations of renal insufficiency, as well as underlying or related kidney diseases such as renal hypoperfusion, intradialytic hypotension, obstructive uropathy, glomerulopathies, IgA nephropathy, glomerulonephritis, acute glomerulonephritis, glomerulosclerosis, tubulointerstitial diseases, nephropathic diseases such as primary and
20 congenital kidney disease, nephritis, Alport syndrome, kidney inflammation, immunological kidney diseases such as kidney transplant rejection, immune complex-induced kidney diseases, nephropathy induced by toxic substances, contrast medium-induced nephropathy; minimal change glomerulonephritis (lipoid); Membranous glomerulonephritis; focal segmental glomerulosclerosis (FSGS); hemolytic uremic syndrome (HUS), amyloidosis, Goodpasture's syndrome, Wegener's
25 granulomatosis, Purpura Schönlein-Henoch, diabetic and non-diabetic nephropathy, pyelonephritis, renal cysts, nephrosclerosis, hypertensive nephrosclerosis and nephrotic syndrome, which can be characterized diagnostically, for example, by abnormally reduced creatinine and/or water excretion, abnormally increased blood concentrations of urea, nitrogen, potassium and/or creatinine, altered activity of renal enzymes such as, for example, glutamyl synthetase, altered urine osmolarity or
30 urine volume, increased microalbuminuria, macroalbuminuria, lesions of glomeruli and arterioles, tubular dilatation, hyperphosphataemia and/or the need for dialysis. The present invention also comprises the use of the compounds according to the invention for the treatment and/or prevention of sequelae of renal insufficiency, for example pulmonary edema, heart failure, uraemia, anaemia, electrolyte disturbances (*e.g.* hyperkalaemia, hyponatraemia) and disturbances in bone and
35 carbohydrate metabolism. The compounds according to the invention are also suitable for the treatment and/or prevention of polycystic kidney disease (PCKD) and of the syndrome of inadequate ADH secretion (SIADH).

Cardiovascular diseases in this context that may be treated and/or prevented with the compounds of the invention include, but are not limited to, the following: acute and chronic heart failure including worsening chronic heart failure (or hospitalization for heart failure) and including congestive heart failure, arterial hypertension, resistant hypertension, arterial pulmonary hypertension, coronary heart disease, stable and unstable angina pectoris, atrial and ventricular arrhythmias, disturbances of atrial and ventricular rhythm and conduction disturbances, for example atrioventricular blocks of degree I-III (AVB I-III), supraventricular tachyarrhythmia, atrial fibrillation, atrial flutter, ventricular fibrillation, ventricular flutter, ventricular tachyarrhythmia, torsade-de-pointes tachycardia, atrial and ventricular extrasystoles, AV-junction extrasystoles, sick-sinus syndrome, syncope, AV-node re-entry tachycardia and Wolff-Parkinson-White syndrome, acute coronary syndrome (ACS), autoimmune heart diseases (pericarditis, endocarditis, valvulitis, aortitis, cardiomyopathies), shock such as cardiogenic shock, septic shock and anaphylactic shock, aneurysms, Boxer cardiomyopathy (premature ventricular contraction), furthermore thromboembolic diseases and ischaemias such as peripheral perfusion disturbances, reperfusion injury, arterial and venous thromboses, myocardial insufficiency, endothelial dysfunction, micro- and macrovascular damage (vasculitis) and for preventing restenoses such as after thrombolysis therapies, percutaneous transluminal angioplasty (PTA), percutaneous transluminal coronary angioplasty (PTCA), heart transplantation and bypass operations, arteriosclerosis, disturbances of lipid metabolism, hypolipoproteinaemias, dyslipidemias, hypertriglyceridemias, hyperlipidemias and combined hyperlipidemias, hypercholesterolaemias, abetalipoproteinaemia, sitosterolemia, xanthomatosis, Tangier disease, adipositas, obesity, metabolic syndrome, transitory and ischemic attacks, stroke, inflammatory cardiovascular diseases, peripheral and cardiac vascular diseases, peripheral circulation disorders, spasms of the coronary arteries and peripheral arteries, and edema such as, for example, pulmonary edema, cerebral edema, renal edema and heart failure-related edema.

In the sense of the present invention, the term heart failure also includes more specific or related disease forms such as right heart failure, left heart failure, global insufficiency, ischemic cardiomyopathy, dilatative cardiomyopathy, congenital heart defects, heart valve defects, heart failure with heart valve defects, mitral valve stenosis, mitral valve insufficiency, aortic valve stenosis, aortic valve insufficiency, tricuspidal stenosis, tricuspidal insufficiency, pulmonary valve stenosis, pulmonary valve insufficiency, combined heart valve defects, heart muscle inflammation (myocarditis), chronic myocarditis, acute myocarditis, viral myocarditis, diabetic heart failure, alcoholic cardiomyopathy, cardiac storage diseases, heart failure with preserved ejection fraction (HFpEF or diastolic heart failure), and heart failure with reduced ejection fraction (HFrEF or systolic heart failure).

The compounds of the present invention may be particularly useful for the treatment and/or prevention of the cardiorenal syndrome (CRS) and its various subtypes. This term embraces certain dis-

orders of the heart and kidneys whereby acute or chronic dysfunction in one organ may induce acute or chronic dysfunction of the other.

Moreover, the compounds according to the invention may be used for the treatment and/or prevention of peripheral arterial disease (PAD) including claudication and including critical limb ischemia as well as coronary microvascular dysfunction (CMD) including CMD type 1-4, primary and secondary Raynaud's phenomenon, microcirculation disturbances, peripheral and autonomic neuropathies, diabetic microangiopathies, diabetic retinopathy, diabetic limb ulcers, gangrene, CREST syndrome, erythematous disorders, rheumatic diseases and for promoting wound healing.

Furthermore, the compounds of the invention are suitable for treating urological diseases and diseases of the male and female urogenital system such as, for example, benign prostatic syndrome (BPS), benign prostatic hyperplasia (BPH), benign prostatic enlargement (BPE), bladder outlet obstruction (BOO), lower urinary tract syndromes (LUTS), neurogenic overactive bladder (OAB), interstitial cystitis (IC), urinary incontinence (UI) such as for example mixed, urge, stress and overflow incontinence (MUI, UUI, SUI, OUI), pelvic pains, erectile dysfunction, dysmenorrhea and endometriosis.

The compounds according to the invention may also be used for the treatment and/or prevention of inflammatory diseases, asthmatic diseases, chronic obstructive pulmonary disease (COPD), acute respiratory distress syndrome (ARDS), acute lung injury (ALI), alpha-1-antitrypsin deficiency (AATD), pulmonary fibrosis, pulmonary emphysema (*e.g.* smoking-induced pulmonary emphysema) and cystic fibrosis (CF). In addition, the compounds of the invention may be used for the treatment and/or prevention of pulmonary arterial hypertension (PAH) and other forms of pulmonary hypertension (PH), including pulmonary hypertension associated with left ventricular disease, HIV infection, sickle cell anaemia, thromboembolism (CTEPH), sarcoidosis, chronic obstructive pulmonary disease (COPD) or pulmonary fibrosis.

Additionally, the compounds according to the invention may be used for the treatment and/or prevention of liver cirrhosis, ascites, diabetes mellitus and diabetic complications such as, for example, neuropathy and nephropathy.

Further, the compounds of the invention are suitable for the treatment and/or prevention of central nervous disorders such as anxiety states, depression, glaucoma, cancer, in particular pulmonary tumors, and circadian rhythm misalignment such as jet lag and shift work.

Furthermore, the compounds according to the invention may be useful for the treatment and/or prevention of pain conditions, diseases of the adrenals such as, for example, pheochromocytoma and adrenal apoplexy, diseases of the intestine such as, for example, Crohn's disease and diarrhea, menstrual disorders such as, for example, dysmenorrhea, endometriosis, preterm labor and tocolysis.

Due to their activity and selectivity profile, the compounds of the present invention are believed to be particularly suitable for the treatment and/or prevention of acute and chronic kidney diseases including diabetic nephropathy, acute and chronic heart failure, preeclampsia, peripheral arterial disease (PAD), coronary microvascular dysfunction (CMD), Raynaud's syndrome, and
5 dysmenorrhea.

The diseases mentioned above have been well characterized in humans, but also exist with a comparable etiology in other mammals, and may be treated in those with the compounds and methods of the present invention.

Thus, the present invention further relates to the use of the compounds according to the invention
10 for the treatment and/or prevention of diseases, especially of the aforementioned diseases.

The present invention further relates to the use of the compounds according to the invention for preparing a pharmaceutical composition for the treatment and/or prevention of diseases, especially of the aforementioned diseases.

The present invention further relates to the use of the compounds according to the invention in a
15 method for the treatment and/or prevention of diseases, especially of the aforementioned diseases.

The present invention further relates to a method for the treatment and/or prevention of diseases, especially of the aforementioned diseases, by using an effective amount of at least one of the compounds according to the invention.

In accordance with another aspect, the present invention covers pharmaceutical combinations, in
20 particular medicaments, comprising at least one compound of general formula (I) of the present invention and at least one or more further active ingredients, in particular for the treatment and/or prevention of diseases, especially of the aforementioned diseases.

Particularly, the present invention covers a pharmaceutical combination, which comprises:

- one or more first active ingredients, in particular compounds of general formula (I) as
25 defined *supra*, and
- one or more further active ingredients, in particular for the treatment and/or prevention of diseases, especially of the aforementioned diseases.

The term "combination" in the present invention is used as known to persons skilled in the art, it being possible for said combination to be a fixed combination, a non-fixed combination or a kit-of-
30 parts.

A "fixed combination" in the present invention is used as known to persons skilled in the art and is defined as a combination wherein, for example, a first active ingredient, such as one or more compounds of general formula (I) of the present invention, and a further active ingredient are present together in one unit dosage or in one single entity. One example of a "fixed combination" is
35 a pharmaceutical composition wherein a first active ingredient and a further active ingredient are

present in admixture for simultaneous administration, such as in a formulation. Another example of a “fixed combination” is a pharmaceutical combination wherein a first active ingredient and a further active ingredient are present in one unit without being in admixture.

5 A non-fixed combination or “kit-of-parts” in the present invention is used as known to persons skilled in the art and is defined as a combination wherein a first active ingredient and a further active ingredient are present in more than one unit. One example of a non-fixed combination or kit-of-parts is a combination wherein the first active ingredient and the further active ingredient are present separately. It is possible for the components of the non-fixed combination or kit-of-parts to be administered separately, sequentially, simultaneously, concurrently or chronologically
10 staggered.

The compounds of the present invention can be administered as the sole pharmaceutical agent or in combination with one or more other pharmaceutically active ingredients where the combination causes no unacceptable adverse effects. The present invention also covers such pharmaceutical combinations. For example, the compounds of the present invention can be combined with known
15 agents for the treatment and/or prevention of diseases, especially of the aforementioned diseases.

In particular, the compounds of the present invention may be used in fixed or separate combination with

- antithrombotic agents, for example and preferably from the group of platelet aggregation inhibitors, anticoagulants and profibrinolytic substances;
- 20 • blood pressure lowering agents, for example and preferably from the group of calcium antagonists, angiotensin AII antagonists, ACE inhibitors, NEP inhibitors, vasopeptidase inhibitors, endothelin antagonists, renin inhibitors, alpha-blockers, beta-blockers, mineralocorticoid receptor antagonists and diuretics;
- antidiabetic agents (hypoglycemic or antihyperglycemic agents), such as for example and
25 preferably insulin and derivatives, sulfonylureas, biguanides, thiazolidinediones, acarbose, DPP4 inhibitors, GLP-1 analogues, or SGLT inhibitors (gliflozins).
- organic nitrates and NO-donors, for example sodium nitroprusside, nitroglycerin, isosorbide mononitrate, isosorbide dinitrate, molsidomine or SIN-1, and inhalational NO;
- compounds that inhibit the degradation of cyclic guanosine monophosphate (cGMP), for
30 example inhibitors of phosphodiesterases (PDE) 1, 2, 5 and/or 9, in particular PDE-5 inhibitors such as sildenafil, vardenafil, tadalafil, udenafil, dasantafil, avanafil, mirodenafil, lodenafil, CTP-499 or PF-00489791;

- positive-inotropic agents, such as for example cardiac glycosides (digoxin) and beta-adrenergic and dopaminergic agonists such as isoproterenol, adrenalin, noradrenalin, dopamine or dobutamine;
- natriuretic peptides, such as for example atrial natriuretic peptide (ANP, anaritide), B-type natriuretic peptide or brain natriuretic peptide (BNP, nesiritide), C-type natriuretic peptide (CNP) or urodilatin;
- calcium sensitizers, such as for example and preferably levosimendan;
- NO- and heme-independent activators of soluble guanylate cyclase (sGC), for example and with preference the compounds described in WO 01/19355, WO 01/19776, WO 01/19778, WO 01/19780, WO 02/070462 and WO 02/070510;
- NO-independent, but heme-dependent stimulators of guanylate cyclase (sGC), for example and with preference the compounds described in WO 00/06568, WO 00/06569, WO 02/42301, WO 03/095451, WO 2011/147809, WO 2012/004258, WO 2012/028647 and WO 2012/059549;
- agents, that stimulates the synthesis of cGMP, for example and with preference sGC modulators, for example and with preference riociguat, cinaciguat, vericiguat or BAY 1101042;
- inhibitors of human neutrophil elastase (HNE), such as for example sivelestat or DX-890 (reltran);
- compounds inhibiting the signal transduction cascade, in particular tyrosine and/or serine/threonine kinase inhibitors, such as for example nintedanib, dasatinib, nilotinib, bosutinib, regorafenib, sorafenib, sunitinib, cediranib, axitinib, telatinib, imatinib, brivanib, pazopanib, vatalanib, gefitinib, erlotinib, lapatinib, canertinib, lestaurtinib, pelitinib, semaxanib or tandutinib;
 - compounds influencing the energy metabolism of the heart, such as for example and preferably etomoxir, dichloroacetate, ranolazine or trimetazidine, or full or partial adenosine A1 receptor agonists as GS-9667 (previously known as CVT-3619), capadenoson and neladenoson bialanate (BAY 1067197);;
- compounds influencing the heart rate, such as for example and preferably ivabradine;
- cardiac myosin activators, such as for example and preferably omecamtiv mecarbil (CK-1827452);
- anti-inflammatory drugs such as non-steroidal anti-inflammatory drugs (NSAIDs) including acetylsalicylic acid (aspirin), ibuprofen and naproxen, glucocorticoids, , such as for example and preferably prednison, prednisolon, methylprednisolon, triamcinolon, dexamethason, beclomethason, betamethason, flunisolid, budesonid or fluticason, 5-aminosalicylic acid

derivatives, leukotriene antagonists, TNF-alpha inhibitors and chemokine receptor antagonists such as CCR1, 2 and/or 5 inhibitors;

- fat metabolism altering agents, for example and preferably from the group of thyroid receptor agonists, cholesterol synthesis inhibitors, such as for example and preferably HMG-CoA-reductase or squalene synthesis inhibitors, ACAT inhibitors, CETP inhibitors, MTP inhibitors, PPAR-alpha, PPAR-gamma and/or PPAR-delta agonists, cholesterol absorption inhibitors, lipase inhibitors, polymeric bile acid adsorbers, bile acid reabsorption inhibitors and lipoprotein(a) antagonists.

Antithrombotic agents are preferably to be understood as compounds from the group of platelet aggregation inhibitors, anticoagulants and profibrinolytic substances.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a platelet aggregation inhibitor, for example and preferably aspirin, clopidogrel, ticlopidine or dipyridamole.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a thrombin inhibitor, for example and preferably ximelagatran, dabigatran, melagatran, bivalirudin or enoxaparin.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a GPIIb/IIIa antagonist, for example and preferably tirofiban or abciximab.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a factor Xa inhibitor, for example and preferably rivaroxaban, apixaban, otamixaban, fidexaban, razaxaban, fondaparinux, idraparinux, DU-176b, PMD-3112, YM-150, KFA-1982, EMD-503982, MCM-17, MLN-1021, DX 9065a, DPC 906, JTV 803, SSR-126512 or SSR-128428.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with heparin or a low molecular weight (LMW) heparin derivative.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a vitamin K antagonist, for example and preferably coumarin.

Blood pressure lowering agents are preferably to be understood as compounds from the group of calcium antagonists, angiotensin AII antagonists, ACE inhibitors, NEP inhibitors, vasopeptidase inhibitors, endothelin antagonists, renin inhibitors, alpha-blockers, beta-blockers, mineralocorticoid receptor antagonists and diuretics.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a calcium antagonist, for example and preferably nifedipine, amlodipine, verapamil or diltiazem.

5 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with an alpha-1-receptor blocker, for example and preferably prazosin or tamsulosin.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a beta-blocker, for example and preferably propranolol, atenolol, timolol, pindolol, alprenolol, oxprenolol, penbutolol, bupranolol, metipranolol, nadolol, mepindolol, 10 carazolol, sotalol, metoprolol, betaxolol, celiprolol, bisoprolol, carteolol, esmolol, labetalol, carvedilol, adaprolol, landiolol, nebivolol, epanolol or bucindolol.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with an angiotensin AII receptor antagonist, for example and preferably losartan, candesartan, valsartan, telmisartan, irbesartan, olmesartan, eprosartan, embursartan or 15 azilsartan.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a vasopeptidase inhibitor or inhibitor of neutral endopeptidase (NEP), such as for example and preferably sacubitril, omapatrilat or AVE-7688.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a dual angiotensin AII receptor antagonist/NEP inhibitor (ARNI), for 20 example and preferably LCZ696.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with an ACE inhibitor, for example and preferably enalapril, captopril, lisinopril, ramipril, delapril, fosinopril, quinopril, perindopril, benazepril ortrandopril.

25 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with an endothelin antagonist, for example and preferably bosentan, darusentan, ambrisentan, tezosentan, sitaxsentan, avosentan, macitentan or atrasentan.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a renin inhibitor, for example and preferably aliskiren, SPP-600 or SPP- 30 800.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a mineralocorticoid receptor antagonist, for example and preferably finerenone, esaxerenone (CS-3150), or apararenone (MT-3995), spironolactone, canrenone, potassium canrenoate, eplerenone, CS-3150, or MT-3995.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a diuretic, such as for example and preferably furosemide, bumetanide, piretanide, torsemide, bendroflumethiazide, chlorothiazide, hydrochlorothiazide, xipamide, indapamide, hydroflumethiazide, methyclothiazide, polythiazide, trichloromethiazide, chlorothalidone, metolazone, quinethazone, acetazolamide, dichlorophenamide, methazolamide, glycerine, isosorbide, mannitol, amiloride or triamterene.

Fat metabolism altering agents are preferably to be understood as compounds from the group of CETP inhibitors, thyroid receptor agonists, cholesterol synthesis inhibitors such as HMG-CoA-reductase or squalene synthesis inhibitors, ACAT inhibitors, MTP inhibitors, PPAR-alpha, PPAR-gamma and/or PPAR-delta agonists, cholesterol absorption inhibitors, polymeric bile acid adsorbers, bile acid reabsorption inhibitors, lipase inhibitors and lipoprotein(a) antagonists.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a CETP inhibitor, for example and preferably dalcetrapib, anacetrapib, BAY 60-5521 or CETP-vaccine (Avant).

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a thyroid receptor agonist, for example and preferably D-thyroxin, 3,5,3'-triiodothyronin (T3), CGS 23425 or axitirome (CGS 26214).

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with an HMG-CoA-reductase inhibitor from the class of statins, for example and preferably lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin, rosuvastatin or pitavastatin.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a squalene synthesis inhibitor, for example and preferably BMS-188494 or TAK-475.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with an ACAT inhibitor, for example and preferably avasimibe, melinamide, pactimibe, eflucimibe or SMP-797.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with an MTP inhibitor, for example and preferably implitapide, R-103757, BMS-201038 or JTT-130.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a PPAR-gamma agonist, for example and preferably pioglitazone or rosiglitazone.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a PPAR-delta agonist, for example and preferably GW 501516 or BAY 68-5042.

5 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a cholesterol absorption inhibitor, for example and preferably ezetimibe, tiqueside or pamaqueside.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a lipase inhibitor, for example and preferably orlistat.

10 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a polymeric bile acid adsorber, for example and preferably cholestyramine, colestipol, colesolvam, CholestaGel or colestimide.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a bile acid reabsorption inhibitor, for example and preferably ASBT (= IBAT) inhibitors such as AZD-7806, S-8921, AK-105, BARI-1741, SC-435 or SC-635.

15 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a lipoprotein(a) antagonist, for example and preferably gemcabene calcium (CI-1027) or nicotinic acid.

20 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a TGFbeta antagonist, by way of example and with preference pirfenidone or fresolimumab.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with HIF-PH inhibitors, by way of example and with preference molidustat or roxadustat.

25 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a CCR2 antagonist, by way of example and with preference CCX-140.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a TNFalpha antagonist, by way of example and with preference adalimumab.

30 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a galectin-3 inhibitor, by way of example and with preference GCS-100.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a BMP-7 agonist, by way of example and with preference THR-184.

5 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a NOX1/4 inhibitor, by way of example and with preference GKT-137831.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a medicament which affects the vitamin D metabolism, by way of example and with preference cholecalciferol or paracalcitol.

10 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a cytostatic agent, by way of example and with preference cyclophosphamide.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with an immunosuppressive agent, by way of example and with
15 preference ciclosporin.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a phosphate binder, by way of example and with preference sevelamer or lanthanum carbonate.

In a preferred embodiment of the invention, the compounds according to the invention are
20 administered in combination with a calcimimetic for therapy of hyperparathyroidism.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with agents for iron deficit therapy, by way of example and with preference iron products.

In a preferred embodiment of the invention, the compounds according to the invention are
25 administered in combination with agents for the therapy of hyperurikaemia, by way of example and with preference allopurinol or rasburicase.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with glycoprotein hormone for the therapy of anaemia, by way of example and with preference erythropoietin.

30 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with biologics for immune therapy, by way of example and with preference abatacept, rituximab, eculizumab or belimumab.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with Jak inhibitors, by way of example and with preference

ruxolitinib, tofacitinib, baricitinib, CYT387, GSK2586184, lestaurtinib, pacritinib (SB1518) or TG101348.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with prostacyclin analogs for therapy of microthrombi.

- 5 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with an alkali therapy, by way of example and with preference sodium bicarbonate.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with an mTOR inhibitor, by way of example and with preference
10 everolimus or rapamycin.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with an NHE3 inhibitor, by way of example and with preference AZD1722.

In a preferred embodiment of the invention, the compounds according to the invention are
15 administered in combination with an eNOS modulator, by way of example and with preference sapropterin.

In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with a CTGF inhibitor, by way of example and with preference FG-3019.

- 20 In a preferred embodiment of the invention, the compounds according to the invention are administered in combination with antidiabetics (hypoglycemic or antihyperglycemic agents), such as for example and preferably insulin and derivatives, sulfonylureas such as tolbutamide, carbutamide, acetohexamide, chlorpropamide, glipizide, gliclazide, glibenclamide, glyburide, glibornuride, gliquidone, glisoxepide, glyclopyramide, glimepiride, JB253 and JB558, meglitinides
25 such as repaglinide and nateglinide, biguanides such as metformin and buformin, thiazolidinediones such as rosiglitazone and pioglitazone, alpha-glucosidase inhibitors such as miglitol, acarbose and voglibose, DPP4 inhibitors such as vildagliptin, sitagliptin, saxagliptin, linagliptin, alogliptin, septagliptin and teneligliptin, GLP-1 analogues such as exenatide (also exendin-4, liraglutide, lixisenatide and taspoglutide, or SGLT inhibitors (gliflozins) such as
30 canagliflozin, dapagliflozin and empagliflozin.

In a particularly preferred embodiment, the compounds of the present invention are administered in combination with one or more additional therapeutic agents selected from the group consisting of diuretics, angiotensin AII antagonists, ACE inhibitors, beta-receptor blockers, mineralocorticoid receptor antagonists, antidiabetics, organic nitrates and NO donors, activators and stimulators of
35 the soluble guanylate cyclase (sGC), and positive-inotropic agents.

In a further particularly preferred embodiment, the compounds of the present invention are administered in combination with one or more additional therapeutic agents selected from the group consisting of diuretics, angiotensin AII antagonists, ACE inhibitors, beta-receptor blockers, mineralocorticoid receptor antagonists, antidiabetics, organic nitrates and NO donors, activators and stimulators of the soluble guanylate cyclase (sGC), antiinflammatory agents, immunosuppressive agents, phosphate binders and/or compounds which modulate vitamin D metabolism.

Thus, in a further embodiment, the present invention relates to pharmaceutical compositions comprising at least one of the compounds according to the invention and one or more additional therapeutic agents for the treatment and/or prevention of diseases, especially of the aforementioned diseases.

Furthermore, the compounds of the present invention may be utilized, as such or in compositions, in research and diagnostics, or as analytical reference standards and the like, which are well known in the art.

When the compounds of the present invention are administered as pharmaceuticals, to humans and other mammals, they can be given *per se* or as a pharmaceutical composition containing, for example, 0.1% to 99.5% (more preferably, 0.5% to 90%) of active ingredient in combination with one or more pharmaceutically acceptable excipients.

Thus, in another aspect, the present invention relates to pharmaceutical compositions comprising at least one of the compounds according to the invention, conventionally together with one or more inert, non-toxic, pharmaceutically acceptable excipients, and to the use thereof for the treatment and/or prevention of diseases, especially of the aforementioned diseases.

It is possible for the compounds according to the invention to have systemic and/or local activity. For this purpose, they can be administered in a suitable manner, such as, for example, via the oral, parenteral, pulmonary, nasal, sublingual, lingual, buccal, rectal, vaginal, dermal, transdermal, conjunctival, otic route or as an implant or stent.

For these administration routes, it is possible for the compounds according to the invention to be administered in suitable administration forms.

For oral administration, it is possible to formulate the compounds according to the invention to dosage forms known in the art that deliver the compounds of the invention rapidly and/or in a modified manner, such as, for example, tablets (uncoated or coated tablets, for example with enteric or controlled release coatings that dissolve with a delay or are insoluble), orally-disintegrating tablets, films/wafers, films/lyophilisates, capsules (for example hard or soft gelatine capsules), sugar-coated tablets, granules, pellets, powders, emulsions, suspensions, aerosols or

solutions. It is possible to incorporate the compounds according to the invention in crystalline and/or amorphised and/or dissolved form into said dosage forms.

Parenteral administration can be effected with avoidance of an absorption step (for example intravenous, intraarterial, intracardial, intraspinal or intralumbal) or with inclusion of absorption
5 (for example intramuscular, subcutaneous, intracutaneous, percutaneous or intraperitoneal). Administration forms which are suitable for parenteral administration are, inter alia, preparations for injection and infusion in the form of solutions, suspensions, emulsions, lyophilisates or sterile powders.

Examples which are suitable for other administration routes are pharmaceutical forms for
10 inhalation [inter alia powder inhalers, nebulizers], nasal drops, nasal solutions, nasal sprays; tablets/films/wafers/capsules for lingual, sublingual or buccal administration; suppositories; eye drops, eye ointments, eye baths, ocular inserts, ear drops, ear sprays, ear powders, ear-rinses, ear tampons; vaginal capsules, aqueous suspensions (lotions, mixturae agitandae), lipophilic suspensions, emulsions, ointments, creams, transdermal therapeutic systems (such as, for example,
15 patches), milk, pastes, foams, dusting powders, implants or stents.

The compounds according to the invention can be incorporated into the stated administration forms. This can be effected in a manner known per se by mixing with pharmaceutically suitable excipients. Pharmaceutically suitable excipients include, inter alia,

- 20 • fillers and carriers (for example cellulose, microcrystalline cellulose (such as, for example, Avicel[®]), lactose, mannitol, starch, calcium phosphate (such as, for example, Di-Cafos[®])),
- ointment bases (for example petroleum jelly, paraffins, triglycerides, waxes, wool wax, wool wax alcohols, lanolin, hydrophilic ointment, polyethylene glycols),
- bases for suppositories (for example polyethylene glycols, cacao butter, hard fat),
- 25 • solvents (for example water, ethanol, isopropanol, glycerol, propylene glycol, medium chain-length triglycerides fatty oils, liquid polyethylene glycols, paraffins),
- surfactants, emulsifiers, dispersants or wetters (for example sodium dodecyl sulfate, lecithin, phospholipids, fatty alcohols (such as, for example, Lanette[®]), sorbitan fatty acid esters (such as, for example, Span[®]), polyoxyethylene sorbitan fatty acid esters (such as, for example, Tween[®]), polyoxyethylene fatty acid glycerides (such as, for example, Cremophor[®]), polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers,
30 glycerol fatty acid esters, poloxamers (such as, for example, Pluronic[®]),
- buffers, acids and bases (for example phosphates, carbonates, citric acid, acetic acid, hydrochloric acid, sodium hydroxide solution, ammonium carbonate, trometamol, triethanolamine),
- 35 • isotonicity agents (for example glucose, sodium chloride),
- adsorbents (for example highly-disperse silicas),
- viscosity-increasing agents, gel formers, thickeners and/or binders (for example polyvinylpyrrolidone, methylcellulose, hydroxypropylmethylcellulose, hydroxypropyl-

cellulose, carboxymethylcellulose-sodium, starch, carbomers, polyacrylic acids (such as, for example, Carbopol[®]); alginates, gelatine),

- disintegrants (for example modified starch, carboxymethylcellulose-sodium, sodium starch glycolate (such as, for example, Explotab[®]), cross- linked polyvinylpyrrolidone, croscarmellose-sodium (such as, for example, AcDiSol[®])),
- flow regulators, lubricants, glidants and mould release agents (for example magnesium stearate, stearic acid, talc, highly-disperse silicas (such as, for example, Aerosil[®])),
- coating materials (for example sugar, shellac) and film formers for films or diffusion membranes which dissolve rapidly or in a modified manner (for example polyvinylpyrrolidones (such as, for example, Kollidon[®]), polyvinyl alcohol, hydroxypropylmethylcellulose, hydroxypropylcellulose, ethylcellulose, hydroxypropylmethylcellulose phthalate, cellulose acetate, cellulose acetate phthalate, polyacrylates, polymethacrylates such as, for example, Eudragit[®])),
- capsule materials (for example gelatine, hydroxypropylmethylcellulose),
- synthetic polymers (for example polylactides, polyglycolides, polyacrylates, polymethacrylates (such as, for example, Eudragit[®]), polyvinylpyrrolidones (such as, for example, Kollidon[®]), polyvinyl alcohols, polyvinyl acetates, polyethylene oxides, polyethylene glycols and their copolymers and blockcopolymers),
- plasticizers (for example polyethylene glycols, propylene glycol, glycerol, triacetine, triacetyl citrate, dibutyl phthalate),
- penetration enhancers,
- stabilisers (for example antioxidants such as, for example, ascorbic acid, ascorbyl palmitate, sodium ascorbate, butylhydroxyanisole, butylhydroxytoluene, propyl gallate),
- preservatives (for example parabens, sorbic acid, thiomersal, benzalkonium chloride, chlorhexidine acetate, sodium benzoate),
- colourants (for example inorganic pigments such as, for example, iron oxides, titanium dioxide),
- flavourings, sweeteners, flavour- and/or odour-masking agents.

The present invention furthermore relates to a pharmaceutical composition which comprise at least one compound according to the invention, conventionally together with one or more pharmaceutically suitable excipient(s), and to their use according to the present invention.

Based upon standard laboratory techniques known to evaluate compounds useful for the treatment of cardiovascular and renal disorders, by standard toxicity tests and by standard pharmacological assays for the determination of treatment of the conditions identified above in mammals, and by comparison of these results with the results of known active ingredients or medicaments that are used to treat these conditions, the effective dosage of the compounds of the present invention can readily be determined for treatment of each desired indication. The amount of the active ingredient to be administered in the treatment of one of these conditions can vary widely according to such considerations as the particular compound and dosage unit employed, the mode of administration, the period of treatment, the age and sex of the patient treated, and the nature and extent of the condition treated.

The total amount of the active ingredient to be administered will generally range from about 0.001 mg/kg to about 200 mg/kg body weight per day, and preferably from about 0.01 mg/kg to about 20 mg/kg body weight per day. Clinically useful dosing schedules will range from one to three times a day dosing to once every four weeks dosing. In addition, it is possible for "drug holidays", in which
5 a patient is not dosed with a drug for a certain period of time, to be beneficial to the overall balance between pharmacological effect and tolerability. It is possible for a unit dosage to contain from about 0.5 mg to about 1500 mg of active ingredient, and can be administered one or more times per day or less than once a day. The average daily dosage for administration by injection, including intravenous, intramuscular, subcutaneous and parenteral injections, and use of infusion techniques
10 will preferably be from 0.01 to 200 mg/kg of total body weight. Illustratively, the compound of the present invention may be administered parenterally at a dose of about 0.001 mg/kg to about 10 mg/kg, preferably of about 0.01 mg/kg to about 1 mg/kg of body weight. In oral administration, an exemplary dose range is about 0.01 to 100 mg/kg, preferably about 0.01 to 20 mg/kg, and more preferably about 0.1 to 10 mg/kg of body weight. Ranges intermediate to the above-recited values
15 are also intended to be part of the invention.

Of course the specific initial and continuing dosage regimen for each patient will vary according to the nature and severity of the condition as determined by the attending diagnostician, the activity of the specific compound employed, the age and general condition of the patient, time of administration, route of administration, rate of excretion of the drug, drug combinations, and the
20 like. The desired mode of treatment and number of doses of a compound of the present invention or a pharmaceutically acceptable salt or ester or composition thereof can be ascertained by those skilled in the art using conventional treatment tests.

The following exemplary embodiments illustrate the invention. The invention is not restricted to the examples.

25 The percentages in the following tests and examples are, unless stated otherwise, by weight; parts are by weight. Solvent ratios, dilution ratios and concentrations reported for liquid/liquid solutions are each based on volume.

EXPERIMENTAL SECTION

NMR peak forms are stated as they appear in the spectra, possible higher order effects have not been considered.

Chemical names were generated using the ACD/Name software from ACD/Labs. In some cases
5 generally accepted names of commercially available reagents were used in place of ACD/Name generated names.

The following table 1 lists the abbreviations used in this paragraph and in the Examples section as far as they are not explained within the text body. Other abbreviations have their meanings customary *per se* to the skilled person.

10 Table 1: Abbreviations

The following table lists the abbreviations used herein.

Abbreviation	Meaning
br	broad (¹ H-NMR signal)
cat.	catalytic
CI	chemical ionisation
d	doublet
DAD	diode array detector
DCM	dichloromethane
dd	double-doublet
DMF	<i>N,N</i> -dimethylformamide
DMSO	dimethylsulfoxide
dt	double-triplet
ELSD	Evaporative Light Scattering Detector
ESI	electrospray (ES) ionisation
h	hour(s)
HCl	hydrochloric acid
HPLC	high performance liquid chromatography
LC-MS	liquid chromatography mass spectrometry
m	multiplet

Abbreviation	Meaning
min	minute(s)
MS	mass spectrometry
NMR	nuclear magnetic resonance spectroscopy: chemical shifts (δ) are given in ppm. The chemical shifts were corrected by setting the DMSO signal to 2.50 ppm unless otherwise stated.
PDA	Photo Diode Array
q	quartet
quint.	quintuplet
Rt	retention time (as measured either with HPLC or UPLC) in minutes
s	singlet
SQD	Single-Quadrupole-Detector
t	triplet
td	triple-doublet
TFA	trifluoroacetic acid
THF	tetrahydrofuran
UPLC	ultra performance liquid chromatography

The various aspects of the invention described in this application are illustrated by the following examples which are not meant to limit the invention in any way.

The example testing experiments described herein serve to illustrate the present invention and the invention is not limited to the examples given.

EXPERIMENTAL SECTION - GENERAL PART

All reagents, for which the synthesis is not described in the experimental part, are either commercially available, or are known compounds or may be formed from known compounds by known methods by a person skilled in the art.

10 The compounds and intermediates produced according to the methods of the invention may require purification. Purification of organic compounds is well known to the person skilled in the art and there may be several ways of purifying the same compound. In some cases, no purification may be necessary. In some cases, the compounds may be purified by crystallization. In some cases,

impurities may be stirred out using a suitable solvent. In some cases, the compounds may be purified by chromatography, particularly flash column chromatography, using for example prepacked silica gel cartridges, e.g. Biotage SNAP cartridges KP-Sil[®] or KP-NH[®] in combination with a Biotage autopurifier system (SP4[®] or Isolera One[®]) and eluents such as gradients of hexane/ethyl acetate or DCM/methanol. In some cases, the compounds may be purified by preparative HPLC using for example a Waters autopurifier equipped with a diode array detector and/or on-line electrospray ionization mass spectrometer in combination with a suitable prepacked reverse phase column and eluents such as gradients of water and acetonitrile which may contain additives such as trifluoroacetic acid, formic acid or aqueous ammonia.

In some cases, purification methods as described above can provide those compounds of the present invention which possess a sufficiently basic or acidic functionality in the form of a salt, such as, in the case of a compound of the present invention which is sufficiently basic, a trifluoroacetate or formate salt for example, or, in the case of a compound of the present invention which is sufficiently acidic, an ammonium salt for example. A salt of this type can either be transformed into its free base or free acid form, respectively, by various methods known to the person skilled in the art, or be used as salts in subsequent biological assays. It is to be understood that the specific form (e.g. salt, free base etc.) of a compound of the present invention as isolated and as described herein is not necessarily the only form in which said compound can be applied to a biological assay in order to quantify the specific biological activity.

UPLC-MS Standard Procedures

Method 1 (LC/MS):

Instrument: Waters ACQUITY SQD UPLC System; Column: Waters Acquity UPLC HSS T3 1.8 μ 50 x 1 mm; eluent A: 1 L water + 0.25 ml 99% formic acid, eluent B: 1 L acetonitrile + 0.25 ml 99% formic acid; gradient: 0.0 min 90% A \rightarrow 1.2 min 5% A \rightarrow 2.0 min 5% A; oven: 50°C; flow rate: 0.40 ml/min; UV detection: 208 – 400 nm.

Method 2 (LC/MS) :

Instrument: Agilent MS Quad 6150; HPLC: Agilent 1290; Column: Waters Acquity UPLC HSS T3 1.8 μ 50 x 2.1 mm; eluent A: 1 L water + 0.25 ml 99% formic acid , eluent B: 1 L acetonitrile + 0.25 ml 99% formic acid; gradient: 0.0 min 90% A \rightarrow 0.3 min 90% A \rightarrow 1.7 min 5% A \rightarrow 3.0 min 5% A; oven: 50°C; flow rate: 1,20 ml/min; UV detection: 205 – 305 nm.

Method 3:

Instrument: Waters ACQUITY SQD UPLC System; Column: Waters Acquity UPLC HSS T3 1.8 μ 50 x 1 mm; eluent A: 1 L water + 0.25 ml 99% formic acid, eluent B: 1 L acetonitrile + 0.25 ml

99% formic acid; gradient: 0.0 min 95% A → 6.0 min 5% A → 7.5 min 5% A; oven: 50°C; flow rate: 0.35 ml/min; UV detection: 210 – 400 nm.

Method 4:

Column: Chromatorex or Reprosil; C18; 10 μm; 125 x 30 mm; Eluent A: 10 L Water + 20 ml 50%
5 Formic acid, Eluent B: Acetonitrile; Gradient: 0-5.50 min 10% B : 90% A; Sample injection: at 3.00 min; 5.50 min 90% A : 10% B → 17.65 min 5% A : 95% B; 17.65-19.48 min 5% A : 95% B; Flow: 75 ml/min; UV- Detection: 210 nm.


Method 5 (LC/MS):

Instrument MS: Thermo Scientific FT-MS; Instrument type UHPLC+: Thermo Scientific UltiMate
10 3000; Column: Waters, HSST3, 2.1 x 75 mm, C18 1.8 μm; eluent A: 1 L water + 0.01% formic acid; eluent B: 1 L acetonitrile + 0.01% formic acid; gradient: 0.0 min 10% B → 2.5 min 95% B → 3.5 min 95% B; oven: 50°C; flow rate: 0.90 ml/min; UV detection: 210 nm/ optimum integration path 210-300 nm.

15 **EXPERIMENTAL SECTION – STARTING MATERIALS AND INTERMEDIATES**

Example 1A

{3-(4-Chlorophenyl)-5-oxo-4-[(2*S*)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1*H*-1,2,4-triazol-1-yl} acetonitrile




20 In a 2 L reaction vessel, 100 g (273 mmol) of {3-(4-chlorophenyl)-5-oxo-4-[(2*S*)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1*H*-1,2,4-triazol-1-yl} acetic acid (synthesis described as Example 8A in WO 2010/105770-A1), 43.3 g (547 mmol) of pyridine and 33 mg (0.3 mmol) of 4-dimethylaminopyridine were dissolved in 300 ml THF. The resulting solution was treated at 5 °C with 52.8 g (438 mmol) of 2,2-dimethylpropanoylchloride over 15 minutes and the resulting
25 mixture was stirred at room temperature for 2.5 hours. After cooling to 0 °C, 183 ml of 28% aqueous ammonia solution was added over 1 h while the solution temperature was kept between 10 °C and 20 °C and at the resulting mixture then stirred at 5 °C for an additional time period of 1 h.

500 ml methyl tert-butylether and 300 ml 20% aqueous citric acid were then added while keeping the internal temperature between 10 °C and 20 °C. The phases were separated and the organic phase was washed with 300 ml of 20% aqueous citric acid followed by 300 ml saturated aqueous sodium hydrogencarbonate solution and finally with 300 ml of 10% aqueous sodium chloride solution. The organic phase was evaporated at 60 °C under reduced pressure until an oily residue was obtained. 300 ml THF was then added and the solution was evaporated again until an oily solution was obtained. This operation was repeated a second time. The oil residue was retaken in 360 ml THF and treated with 172 g (820 mmol) trifluoroacetic acid anhydride over 20 min at a temperature between 10 °C and 20 °C. The resulting solution was then stirred at room temperature for 1 h. 720 ml 4-methyl-2-pentanone and 650 ml 7.5% aqueous sodium hydroxide solution were added at a temperature between 10 °C and 20 °C. Finally the pH-value was adjusted to pH = 9.5 using 7.5% aqueous sodium hydroxide solution. After phase separation, the organic phase was washed twice with 450 ml 10% aqueous sodium chloride solution. The organic phase was evaporated at a temperature of 80 °C under reduced pressure while 1200 ml n-heptane was added. The formed suspension was cooled to 20 °C and a solid formed which was filtered off and washed with 200 ml n-heptane and then dried under reduced pressure (50°C, 30 mbar) affording 88 g (93 % of th.) of {3-(4-chlorophenyl)-5-oxo-4-[(2*S*)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1*H*-1,2,4-triazol-1-yl}acetonitrile as a solid.

¹H-NMR (400 MHz, DMSO-d₆): δ [ppm] = 7.78 (d, 2H), 7.55 (d, 2H), 6.91 (d, 1H), 5.17 (s, 2 H), 4.34-4.23 (m, 1 H), 3.98 (dd, 1H), 3.81 (dd, 1H).

Example 2A

Methyl-2-{3-(4-chlorophenyl)-5-oxo-4-[(2*S*)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1*H*-1,2,4-triazol-1-yl}ethanimidate




25 In a 4 L reaction vessel, 200 g (576.9 mmol) of {3-(4-chlorophenyl)-5-oxo-4-[(2*S*)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1*H*-1,2,4-triazol-1-yl}acetonitrile (Example 1A) in 1600 ml methanol was treated with 5.2 g (28 mmol) sodium methanolate (30% in methanol) and the resulting mixture was stirred at 50 °C for 2.5 hours. The solution was then evaporated at 50 °C under reduced pressure until an oily solution was obtained. 2000 ml methyl tert-butylether was

added and the solution was concentrated until a volume of 800 ml was achieved. 3000 ml n-heptane was then added and a suspension was formed. After cooling at 20 °C, the solid was filtered and washed with 500 ml n-heptane and then dried under reduced pressure (50°C, 30 mbar) affording 175 g (80 % of th.) of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate as a solid.

¹H-NMR (400 MHz, DMSO-d₆): δ [ppm] = 8.01 (s, 1H), 7.78 (d, 2H), 7.62 (d, 2H), 6.93 (br. s, 1H), 4.50 (s, 2 H), 4.35-4.23 (m, 1 H), 3.96 (dd, 1H), 3.81 (dd, 1H), 3.67 (s, 3 H).

Example 3A

(1R)-1-[3-(3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl)methyl)-1-(4-chloropyridin-3-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 200 mg, 528 μmol) in THF (2.0 ml) was treated at 0 °C with N,N-diisopropylethylamine (280 μl, 1.6 mmol) and (2R)-1-chloro-1-oxopropan-2-yl acetate (79 μl, 580 μmol) and stirred at 0°C for 30 min. 4-chloro-3-hydrazinylpyridine (83.4 mg, 581 μmol) was then added and the resulting mixture was stirred 2 h at RT, followed by 3 h at 150°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 178 mg (58 % of Th.) of the title compound.

LC-MS (Method 1): R_t = 0.99 min; MS (ESIpos): m/z = 586 [M+H]⁺

¹H-NMR (500 MHz, DMSO-d₆) δ [ppm]: 8.94-8.68 (m, 2H), 8.02-7.52 (m, 5H), 6.89 (d, 1H), 5.88-5.61 (m, 1H), 5.13 (m, 2H), 4.48-4.17 (m, 1H), 4.08-3.77 (m, 2H), 1.79 (s, 3H), 1.55 (d, 3H).

Example 4A

(1S)-1-[3-(3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl)methyl)-1-(4-chloropyridin-3-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




Under argon, a solution of methyl 2-{{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} ethanimidate (Example 2A, 150 mg, 396 μ mol) in THF (1.5 ml) was treated at 0°C with N,N-diisopropylethylamine (210 μ l, 1.2 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μ l, 440 μ mol) and stirred at 0°C for 30 min. 4-chloro-3-hydrazinylpyridine hydrochloride (78.4 mg, 436 μ mol) was then added and the resulting mixture was stirred 3 h at room temperature, followed by 3 h at 150°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 118 mg (51 % of Th.) of the title compound.

10 LC-MS (Method 1): $R_t = 0.98$ min; MS (ESIpos): $m/z = 586$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-d₆) δ [ppm]: 8.97-8.65 (m, 2H), 8.01-7.46 (m, 5H), 6.89 (d, 1H), 5.75 (m, 1H), 5.12 (m, 2H), 4.36-4.20 (m, 1H), 4.14-3.76 (m, 2H), 1.79 (s, 3H), 1.55 (d, 3H).

Example 5A

15 (1S)-1-[3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(3-chloropyridin-2-yl)-1H-1,2,4-triazol-5-yl] ethyl acetate




Under argon, a solution of methyl 2-{{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} ethanimidate (Example 2A, 150 mg, 396 μ mol) in THF (1.5 ml) was treated at 0°C with N,N-diisopropylethylamine (210 μ l, 1.2 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μ l, 440 μ mol) and stirred at 0°C for 30 min. 3-chloro-2-hydrazinylpyridine (62.5 mg, 436 μ mol) was then added and the resulting mixture was stirred overnight at room temperature and evaporated. The residue was purified by preparative HPLC (Method 4) affording 78.1 mg (34 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 1.03$ min; MS (ESIpos): $m/z = 586$ $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ [ppm]: 8.59 (dd, 1H), 8.32 (dd, 1H), 7.83-7.48 (m, 6H), 6.89 (d, 1H), 5.87 (q, 1H), 5.12 (m, 2H), 4.42-4.21 (m, 1H), 3.98 (d, 1H), 3.85 (dd, 1H), 1.75 (s, 3H), 1.56 (d, 3H).

5 **Example 6A**

(1R)-1-[3-(3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl)methyl)-1-(3,5-dichloropyridin-4-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 200 mg, 528 μmol) in THF (2.0 ml) was treated at 0°C with N,N-diisopropylethylamine (280 μl , 1.6 mmol) and (2R)-1-chloro-1-oxopropan-2-yl acetate (79 μl , 580 μmol) and stirred at 0°C for 30 min. 3,5-dichloro-4-hydrazinylpyridine (103 mg, 581 μmol) was then added and the resulting mixture was stirred 2 h at room temperature, followed by 3 h at 150°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 108 mg (31 % of Th.) of the title compound as a mixture of rotamers.

LC-MS (Method 1): $R_t = 1.06$ min; MS (ESIpos): $m/z = 620$ $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ [ppm] = 8.99 (d, 1H), 7.84-7.51 (m, 5H), 6.89 (m, 1H), 5.81 (m, 1H), 5.16 (m, 2H), 4.37-4.14 (m, 1H), 4.09-3.69 (m, 2H), 1.77 (s, 3H), 1.57 (d, 3H).

20 **Example 7A**

(1R)-1-[3-(3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl)methyl)-1-(3-chloropyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 500 mg, 1.32 mmol) in (5.0 ml) THF was treated at 0°C with N,N-diisopropylethylamine (690 µl, 4.0 mmol) and (2R)-1-chloro-1-oxopropan-2-yl acetate (55 µl, 1.5 mmol) and stirred at 0°C for 30 min. A solution of 3-chloro-2-hydrazinylpyridine (208 mg, 1.45 mmol) in (2.0 µl) THF and (1.0 µl) dioxane was then added and the resulting mixture was stirred overnight at room temperature, followed by 2 h at 150°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 85.2 mg (11 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 1.02$ min; MS (ESIpos): $m/z = 586$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-d₆) δ [ppm]: 8.59 (dd, 1H), 8.32 (dd, 1H), 7.86-7.49 (m, 5H), 6.89 (d, 1H), 5.87 (q, 1H), 5.12 (m, 2H), 4.42-4.18 (m, 1H), 4.07-3.75 (m, 2H), 1.75 (s, 3H), 1.56 (d, 3H).

Example 8A

(1R)-1-[3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-(3-chloropyridin-4-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate



15

Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 200 mg, 528 µmol) in THF (2.0 ml, 25 mmol) was treated at 0°C with N,N-diisopropylethylamine (280 µl, 1.6 mmol) and (2R)-1-chloro-1-oxopropan-2-yl acetate (55 µl, 580 µmol) and stirred at 0°C for 30 min. 3-chloro-4-hydrazinylpyridine (83.4 mg, 581 µmol) was then added and the resulting mixture was stirred overnight at room temperature, followed by 3 h at 150°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 152 mg (49 % of Th.) of the title compound.


LC-MS (Method 1): $R_t = 1.00$ min; MS (ESIpos): $m/z = 586$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-d₆) δ [ppm]: 8.96 (s, 1H), 8.78 (d, 1H), 7.89-7.54 (m, 5H), 6.89 (d, 1H), 5.79 (q, 1H), 5.13 (m, 2H), 4.29 (m, 1H), 4.09-3.75 (m, 2H), 1.76 (s, 3H), 1.56 (d, 3H).

25

Example 9A

(1S)-1-[3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-(3-chloropyridin-4-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 150 mg, 396 μmol) in THF (1.5 ml) was treated at 0°C with N,N-diisopropylethylamine (210 μl , 1.2 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μl , 440 μmol) and stirred at 0°C for 30 min. 3-chloro-4-hydrazinylpyridine (62.5 mg, 436 μmol) was then added and the resulting mixture was stirred 3 h at room temperature, followed by 3 h at 150°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 133 mg (57 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.99$ min; MS (ESIpos): $m/z = 586$ $[\text{M}+\text{H}]^+$

$^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ [ppm]: 8.97 (s, 1H), 8.78 (d, 1H), 7.84-7.58 (m, 5H), 6.89 (d, 1H), 5.88-5.72 (m, 1H), 5.13 (m, 2H), 4.40-4.20 (m, 1H), 4.09-3.79 (m, 2H), 1.76 (s, 3H), 1.56 (d, 3H).

15 **Example 10A**

(1S)-1-[3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-(3-methoxypyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate



Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 100 mg, 264 μmol) in THF (2.0 ml) was treated at 0°C with N,N-diisopropylethylamine (180 μl , 1.1 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (37 μl , 290 μmol) and stirred at 0°C for 30 min. 2-hydrazinyl-3-methoxypyridine (404 mg, 10 % purity, 290 μmol) was then added and the resulting mixture was


stirred overnight at room temperature and evaporated. The residue was purified by preparative HPLC (Method 4) affording 54.2 mg (35 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.93$ min; MS (ESIpos): $m/z = 582$ $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ [ppm]: 8.15 (dd, 1H), 7.85-7.56 (m, 6H), 6.90 (d, 1H), 5.79 (q, 5
1H), 5.09 (m, 2H), 4.38-4.21 (m, 1H), 3.82 (m, 5H), 1.71 (s, 3H), 1.52 (d, 3H).

Example 11A

(1S)-1-[3-(3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl)methyl)-1-(3,5-dichloropyridin-4-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




10 Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 150 mg, 396 μmol) in THF (1.5 ml) was treated at 0°C with N,N-diisopropylethylamine (210 μl , 1.2 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μl , 440 μmol) and stirred at 0°C for 30 min. 3,5-dichloro-4-hydrazinylpyridine (77.6 mg, 436 μmol) was then added and the resulting mixture was stirred 3 h
15 at room temperature, followed by 3 h at 150°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 89.6 mg (33 % of Th.) of the title compound as a mixture of rotamers.

LC-MS (Method 1): $R_t = 1.07$ min; MS (ESIpos): $m/z = 620$ $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ [ppm]: 8.99 (d, 1H), 7.80-7.51 (m, 5H), 6.88 (m, 1H), 5.81 (q, 20
1H), 5.15 (m, 2H), 4.43-4.21 (m, 1H), 4.11-3.74 (m, 2H), 1.77 (s, 3H), 1.57 (d, 3H).

Example 12A

(1S)-1-[3-(3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl)methyl)-1-(1,3-thiazol-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




Under argon, a solution of methyl 2-{{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} ethanimidate (Example 2A, 150 mg, 396 μmol) in THF (1.5 ml) was treated at 0°C with N,N-diisopropylethylamine (210 μl , 1.2 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μl , 440 μmol) and stirred at 0°C for 30 min. 2-hydrazinyl-1,3-thiazole hydrochloride (1:1) (66.1 mg, 436 μmol) was then added and the resulting mixture was stirred 2 h at room temperature, followed by 3 h at 150°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 61.3 mg (28 % of Th.) of the title compound.

10 LC-MS (Method 1): $R_t = 1.02$ min; MS (ESIpos): $m/z = 558$ $[\text{M}+\text{H}]^+$

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ [ppm] = 7.85-7.52 (m, 6H), 6.90 (d, 1H), 6.50 (q, 1H), 5.11 (m, 2H), 4.29 (m, 1H), 4.08-3.78 (m, 2H), 1.96 (s, 3H), 1.61 (d, 3H).

Example 13A

15 (1S)-1-[3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(3-fluoropyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate



Under argon, a solution of methyl 2-{{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} ethanimidate (Example 2A, 150 mg, 396 μmol) in THF (1.5 ml) was treated at 0°C with N,N-diisopropylethylamine (210 μl , 1.2 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μl , 440 μmol) and stirred at 0°C for 30 min. 3-fluoro-2-hydrazinylpyridine (55.4 mg, 436 μmol) was then added and the resulting mixture was stirred overnight at room temperature and evaporated. The residue was purified by preparative HPLC (Method 4) affording 152 mg (67 % of Th.) of the title compound.


LC-MS (Method 1): $R_t = 1.01$ min; MS (ESIpos): $m/z = 570$ $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ [ppm] = 8.46 (br. d, 1H), 8.23-8.05 (m, 1H), 7.88-7.53 (m, 5H), 6.89 (d, 1H), 5.93 (q, 1H), 5.12 (m, 2H), 4.30 (m, 1H), 4.08-3.71 (m, 2H), 1.79 (s, 3H), 1.59 (d, 3H).

5

Example 14A

(1S)-1-{3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-[3-(trifluoromethoxy)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl acetate



- 10 Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 250 mg, 660 μmol) in THF (5.0 ml) was treated at 0°C with N,N-diisopropylethylamine (460 μl , 2.6 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (92 μl , 730 μmol) and stirred at 0°C for 30 min. 2-hydrazinyl-3-(trifluoromethoxy)pyridine 4-methylbenzenesulfonate (1:1) (265 mg, 726 μmol) was then added
- 15 and the resulting mixture was stirred overnight at room temperature, followed by 2 h at 150°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 188 mg (45 % of Th.) of the title compound.


A

LC-MS (Method 1): $R_t = 1.08$ min; MS (ESIpos): $m/z = 636$ $[M+H]^+$

- 20 $^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ [ppm] = 8.62 (dd, 1H), 8.27 (dt, 1H), 7.94-7.50 (m, 5H), 6.89 (d, 1H), 5.99 (q, 1H), 5.13 (m, 2H), 4.43-3.72 (m, 3H), 1.74 (s, 3H), 1.58 (d, 3H).

Example 15A

(1R)-1-[3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-(3-fluoropyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




Under argon, a solution of methyl 2-{{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} ethanimidate (Example 2A, 200 mg, 528 μmol) in THF (2.0 ml) was treated at 0°C with N,N-diisopropylethylamine (280 μl , 1.6 mmol) and (2R)-1-chloro-1-oxopropan-2-yl acetate (73 μl , 580 μmol) and stirred at 0°C for 30 min. 3-fluoro-2-hydrazinylpyridine (73.8 mg, 581 μmol) was then added and the resulting mixture was stirred overnight at room temperature and evaporated. The residue was purified by preparative HPLC (Method 4) affording 195 mg (65 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 1.01$ min; MS (ESIpos): $m/z = 570$ $[\text{M}+\text{H}]^+$

¹H-NMR (400 MHz, DMSO- d_6) δ [ppm]: 8.46 (br. d, 1H), 8.14 (m, 1H), 7.86-7.53 (m, 5H), 6.89 (d, 1H), 5.93 (q, 1H), 5.12 (m, 2H), 4.30 (m, 1H), 4.11-3.74 (m, 2H), 1.79 (s, 3H), 1.59 (d, 3H).

Example 16A

(1S)-1-[3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(pyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate



15

Under argon, a solution of methyl 2-{{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} ethanimidate (Example 2A, 200 mg, 528 μmol) in THF (2.0 ml) was treated at 0°C with N,N-diisopropylethylamine (280 μl , 1.6 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (79 μl , 580 μmol) and stirred at 0°C for 30 min. 2-hydrazinylpyridine (63.4 mg, 581 μmol) was then added and the resulting mixture was stirred overnight at room temperature and evaporated. The residue was purified by preparative HPLC (Method 4) affording 199 mg (68 % yield) of the title compound.


LC-MS (Method 1): $R_t = 1.05$ min; MS (ESIpos): $m/z = 552$ $[\text{M}+\text{H}]^+$

20

¹H-NMR (500 MHz, DMSO-d₆) δ [ppm]: 8.55-8.47 (m, 1H), 8.19-8.02 (m, 1H), 7.83-7.44 (m, 6H), 6.90 (d, 1H), 6.51 (q, 1H), 5.17-5.05 (m, 2H), 4.31 (m, 1H), 4.06-3.81 (m, 2H), 1.89 (s, 3H), 1.62 (d, 3H).

Example 17A

- 5 (1S)-1-[3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-(5-chloropyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 150 mg, 396 μmol) in THF (1.5 ml) was treated at 0°C with N,N-diisopropylethylamine (210 μl, 1.2 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μl, 440 μmol) and stirred at 0°C for 30 min. 5-chloro-2-hydrazinylpyridine (62.5 mg, 436 μmol) was then added and the resulting mixture was stirred overnight at room temperature and evaporated. The residue was purified by preparative HPLC (Method 4) affording 165 mg (71 % of Th.) of the title compound.

- 15 LC-MS (Method 1): R_t = 1.14 min; MS (ESIpos): m/z = 586 [M+H]⁺

¹H-NMR (400 MHz, DMSO-d₆) δ [ppm]: 8.58 (d, 1H), 8.21 (dd, 1H), 7.89-7.54 (m, 5H), 6.90 (d, 1H), 6.46 (q, 1H), 5.23-5.00 (m, 2H), 4.31 (m, 1H), 4.12-3.76 (m, 2H), 1.92 (s, 3H), 1.61 (d, 3H).

Example 18A

- 20 (1R)-1-{3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-[3-(trifluoromethoxy)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl acetate




Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} ethanimidate (Example 2A, 250 mg, 660 μmol) in THF (5.0 ml) was treated at 0°C with N,N-diisopropylethylamine (460 μl , 2.6 mmol) and (2R)-1-chloro-1-oxopropan-2-yl acetate (92 μl , 730 μmol) and stirred at 0°C for 30 min. 2-hydrazinyl-3-(trifluoromethoxy)pyridine 4-methylbenzenesulfonate (1:1) (265 mg, 726 μmol) was then added and the resulting mixture was stirred 5 h at room temperature, followed by 3 h at 120°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 199 mg (47 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 1.08$ min; MS (ESIpos): $m/z = 636$ $[\text{M}+\text{H}]^+$

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ [ppm] 8.62 (dd, 1H), 8.27 (d, 1H), 7.99-7.55 (m, 5H), 6.89 (d, 1H), 5.99 (q, 1H), 5.25-5.01 (m, 2H), 4.41-4.18 (m, 1H), 4.05-3.78 (m, 2H), 1.74 (s, 3H), 1.58 (d, 3H).

Example 19A

(1S)-1-[3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(6-fluoropyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} ethanimidate (Example 2A, 150 mg, 396 μmol) in THF (1.5 ml) was treated at 0°C with N,N-diisopropylethylamine (210 μl , 1.2 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μl , 440 μmol) and stirred at 0°C for 30 min. 2-fluoro-6-hydrazinylpyridine (55.4 mg, 436 μmol) was then added and the resulting mixture was stirred 3 h at room temperature, followed by 3 h at 150°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 168 mg (74 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 1.04$ min; MS (ESIpos): $m/z = 570$ $[\text{M}+\text{H}]^+$

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ [ppm]: 8.26 (q, 1H), 7.80-7.55 (m, 5H), 7.32 (dd, 1H), 6.90 (d, 1H), 6.43 (q, 1H), 5.12 (m, 2H), 4.31 (m, 1H), 4.11-3.94 (m, 1H), 3.86 (dd, 1H), 1.93 (s, 3H), 1.62 (d, 3H).

Example 20A

(1S)-1-[3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-(pyridin-4-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 150 mg, 396 μmol) in THF (1.5 ml) was treated at 0°C with N,N-diisopropylethylamine (210 μl , 1.2 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μl , 440 μmol) and stirred at 0°C for 30 min. 4-hydrazinylpyridine (47.5 mg, 436 μmol) was then added and the resulting mixture was stirred overnight at room temperature, followed by 1 h at 100°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 141 mg (63 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.93$ min; MS (ESIpos): $m/z = 552$ $[\text{M}+\text{H}]^+$

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ [ppm]: 8.78 (d, 2H), 7.81-7.51 (m, 6H), 6.89 (d, 1H), 6.02 (q, 1H), 5.11 (m, 2H), 4.30 (m, 1H), 4.10-3.77 (m, 2H), 1.81 (s, 3H), 1.60 (d, 3H).

15 **Example 21A**

(1S)-1-[3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-(pyrazin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate



Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 150 mg, 396 μmol) in THF (1.5 ml) was treated at 0°C with N,N-diisopropylethylamine (210 μl , 1.2 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μl , 440 μmol) and stirred at 0°C for 30 min. 2-hydrazinylpyrazine (48.0 mg, 436 μmol) was then added and the resulting mixture was stirred


overnight at room temperature and evaporated. The residue was purified by preparative HPLC (Method 4) affording 150 mg (68 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.95$ min; MS (ESIpos): $m/z = 553$ $[M+H]^+$

1H -NMR (400 MHz, DMSO- d_6) δ [ppm] = 9.09 (d, 1H), 8.77 (d, 1H), 8.61 (m, 1H), 7.82-7.52 (m, 4H), 6.91 (d, 1H), 6.44 (q, 1H), 5.15 (m, 2H), 4.31 (m, 1H), 4.10-3.78 (m, 2H), 1.92 (s, 3H), 1.62 (d, 3H).

Example 22A

(1S)-1-[3-(3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl)methyl)-1-(pyridazin-3-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate



10

Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 150 mg, 396 μ mol) in THF (1.5 ml) was treated at 0°C with N,N-diisopropylethylamine (210 μ l, 1.2 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μ l, 440 μ mol) and stirred at 0°C for 30 min. 3-hydrazinylpyridazine (48.0 mg, 436 μ mol) was then added and the resulting mixture was stirred overnight at room temperature and evaporated. The residue was purified by preparative HPLC (Method 4) affording 85.6 mg (39 % of Th.) of the title compound.

15


LC-MS (Method 1): $R_t = 0.90$ min; MS (ESIpos): $m/z = 553$ $[M+H]^+$

1H -NMR (400 MHz, DMSO- d_6): δ [ppm] = 9.35 (dd, 1H), 8.19-7.92 (m, 2H), 7.86-7.54 (m, 4H), 6.90 (d, 1H), 6.47 (q, 1H), 5.15 (m, 2H), 4.31 (m, 1H), 4.14-3.79 (m, 2H), 1.90 (s, 3H), 1.66 (d, 3H).

20

Example 23A

(1S)-1-[3-(3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl)methyl)-1-(3-methoxy-pyridazin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




Under argon, a solution of methyl 2-{{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 150 mg, 396 μmol) in THF (3.0 ml) was treated at 0°C with N,N-diisopropylethylamine (280 μl , 1.6 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μl , 440 μmol) and stirred at 0°C for 30 min. 2-hydrazinyl-3-methoxypyrazine dihydrochloride (92.8 mg, 436 μmol) was then added and the resulting mixture was stirred 5 h at room temperature, followed by 3 h at 120°C in the microwave and evaporated. The residue was purified by preparative HPLC (Method 4) affording 68.5 mg (30 % of Th.) of the title compound.

10 LC-MS (Method 1): R_t = 1.01 min; MS (ESIpos): m/z = 583 [M+H]⁺

¹H-NMR (400 MHz, DMSO-d₆) δ [ppm]: 8.53 (d, 1H), 8.29 (d, 1H), 7.81-7.57 (m, 4H), 6.89 (d, 1H), 5.85 (q, 1H), 5.11 (m, 2H), 4.37-4.23 (m, 1H), 4.08-3.77 (m, 5H), 1.74 (s, 3H), 1.56 (d, 3H).

Example 24A

15 (1S)-1-[3-({3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-(1-methyl-1H-imidazol-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




20 Under argon, a solution of methyl 2-{{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 150 mg, 396 μmol) in THF (1.5 ml) was treated at 0°C with N,N-diisopropylethylamine (280 μl , 1.6 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (55 μl , 440 μmol) and stirred at 0°C for 30 min. 2-hydrazinyl-1-methyl-1H-imidazole dihydrochloride (80.6 mg, 436 μmol) was then added and the resulting mixture was stirred overnight at room temperature and evaporated. The residue was purified by preparative HPLC (Method 4) affording 171 mg (77 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.94$ min; MS (ESIpos): $m/z = 555$ $[M+H]^+$

1H -NMR (400 MHz, DMSO- d_6) δ [ppm] = 7.79-7.55 (m, 4H), 7.38 (d, 1H), 7.03-6.80 (m, 2H), 5.81-5.75 (m, 1H), 5.12 (m, 2H), 4.29 (m, 1H), 4.07-3.76 (m, 2H), 3.48 (s, 3H), 1.89 (s, 3H), 1.56 (d, 3H).

5 **Example 25A**

(1R)-1-[3-(3-(4-Chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl)methyl)-1-(pyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate




Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} ethanimidate (Example 2A, 200 mg, 528 μ mol) in THF (2.0 ml) was treated at 0°C with N,N-diisopropylethylamine (280 μ l, 1.6 mmol) and (2R)-1-chloro-1-oxopropan-2-yl acetate (55 μ l, 580 μ mol) and stirred at 0°C for 30 min. 2-hydrazinylpyridine (63.4 mg, 581 μ mol) was then added and the resulting mixture was stirred overnight at room temperature and evaporated. The residue was purified by preparative HPLC (Method 4) affording 214 mg (73 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 1.05$ min; MS (ESIpos): $m/z = 552$ $[M+H]^+$

1H -NMR (400 MHz, DMSO- d_6): δ [ppm] = 8.51 (dd, 1H), 8.08 (m, 1H), 7.84-7.45 (m, 6H), 6.90 (d, 1H), 6.51 (q, 1H), 5.11 (m, 2H), 4.31 (m, 1H), 4.08-3.80 (m, 2H), 1.89 (s, 3H), 1.62 (d, 3H).

Example 26A

20 Methyl {3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}acetate




Under argon, potassium *t*-butoxide (9.1 g, 81.3 mmol) was added portionwise at room temperature to a solution of 5-(4-chlorophenyl)-4-[(2*S*)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3*H*-1,2,4-triazol-3-one (Example 5A in WO 2011/104322-A1; 20 g, 65.01 mmol) in THF (40 ml). To this solution was added methyl bromoacetate (10.9 g, 71.5 mmol), and the mixture was stirred at room temperature overnight. The reaction mixture was then diluted with water and extracted with ethyl acetate. The combined organic phases were dried over sodium sulfate, filtered, and concentrated *in vacuo* affording 16.4 g (46 % of th.) of the title compound.

LC/MS (Method 1): $R_t = 0.90$ min; MS (ESIpos): $m/z = 380$ [M+H]⁺

¹H NMR (400 MHz, DMSO-*d*₆): δ [ppm] = 7.73-7.81 (m, 2H), 7.60-7.69 (m, 2H), 6.92 (d, 1H), 4.72 (s, 2H), 4.19-4.33 (m, 1H), 4.00 (dd, 1H), 3.85 (dd, 1H), 3.70 (s, 3H).

Example 27A

2-{3-(4-Chlorophenyl)-5-oxo-4-[(2*S*)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1*H*-1,2,4-triazol-1-yl}acetohydrazine




7.2 g (19.0 mmol) of methyl {3-(4-chlorophenyl)-5-oxo-4-[(2*S*)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1*H*-1,2,4-triazol-1-yl}acetate (Example 26A) were dissolved in absolute ethanol (60 ml). To this solution were added 2.09 g (41.7 mmol) hydrazine hydrate, and the mixture was stirred under reflux for 5 h and then at room temperature overnight. The resulting mixture was partially concentrated *in vacuo* and then diluted with water and extracted with ethyl acetate. The combined organic phases were dried over sodium sulfate, filtered, and concentrated *in vacuo*. The residue was dissolved in dichloromethane, and after crystallization the white solid was filtered off and dried under high vacuum affording 7.0 g (97% of th.) of the title compound.

LC/MS (Method 1): $R_t = 0.73$ min; MS (ESIpos): $m/z = 380$ [M+H]⁺

¹H NMR (400 MHz, DMSO-*d*₆): δ [ppm] = 9.23 (t, 1H), 9.23 (t, 1H), 7.73-7.78 (m, 2H), 7.61-7.66 (m, 2H), 6.90 (d, 1H), 4.38 (d, 2H), 4.24-4.34 (m, 3H), 3.96 (dd, 1H), 3.82 (dd, 1H).

Example 28A

5-(4-Chlorophenyl)-2-({5-[1-hydroxyethyl]-4H-1,2,4-triazol-3-yl}methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one (diastereomeric mixture)




Under argon, sodium ethoxide (1.53 g, 21.69 mmol) was added portionwise at room temperature to a solution of 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}acetohydrazine (Example 27A, 4.1 g, 10.8 mmol) and 2-hydroxypropanimidamide hydrochloride (1.48 g, 11.9 mmol) in DMF (110 ml). The reaction mixture was stirred at 120°C for 4.5 h. After cooling, the reaction mixture was partially concentrated *in vacuo* and then diluted with ethyl acetate. The resulting mixture was washed with water, and after phase separation, the aqueous phase was extracted twice with ethyl acetate. The combined organic phases were dried over sodium sulfate, filtered, and concentrated under reduced pressure. The resulting solid was dried under high vacuum affording 4.9 g (92% of th.) of the title compound as a mixture of diastereomers.

LC/MS (Method 1): $R_t = 0.82$ min; MS (ESIpos): $m/z = 433$ [M+H]⁺

¹H NMR (400 MHz, DMSO-*d*₆): δ [ppm] = 13.68 (s, 1H), 7.70-7.77 (d, 2H), 7.58-7.65 (m, 2H), 6.93 (d, 1H), 5.73 (d, 1H), 4.89-5.00 (m, 2H), 4.80 (quin, 1H), 4.24-4.36 (m, 1H), 3.93-4.02 (m, 1H), 3.79-3.88 (m, 1H), 1.39 (d, 3H).

Example 29A

(1S)-1-{3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-[3-(trifluoromethyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl acetate



Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}ethanimidate (Example 2A, 250 mg, 660 μ mol)


in tetrahydrofuran (2.5 ml) was treated at 0°C with N,N-diisopropylethylamine (340 μ l, 2.0 mmol) and (2S)-1-chloro-1-oxopropan-2-yl acetate (92 μ l, 730 μ mol) and stirred at 0°C for 30 min. 2-hydrazinyl-3-(trifluoromethyl)pyridine (76 μ l, 730 μ mol) and anhydrous zinc sulfate (107 mg, 660 μ mol) were added and the resulting mixture was stirred 1 h at 45°C. The reaction mixture was diluted with ethyl acetate and washed with water (3 times) followed by brine. The organic phase was dried over magnesium sulfate and evaporated. The residue was purified by preparative HPLC (Method 4). The product containing fraction were evaporated and the residue purified by preparative HPLC (Method 4) affording 103.5 mg (22% of th.) of the title compound.

LC-MS (Method 5): $R_t = 1.91$ min; MS (ESIpos): $m/z = 620.1$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-d₆) δ [ppm]: 8.93-8.86 (m, 1H), 8.57 (dd, 1H), 7.92 (dd, 1H), 7.80-7.56 (m, 4H), 6.88 (d, 1H), 5.96 (q, 1H), 5.20-5.01 (m, 2H), 4.41-4.21 (m, 1H), 4.05-3.78 (m, 2H), 1.72 (s, 3H), 1.58 (d, 3H).

Example 30A

(1R)-1-{3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-[3-(trifluoromethyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl} ethyl acetate



Under argon, a solution of methyl 2-{3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} ethanimidate (Example 2A, 250 mg, 660 μ mol) in tetrahydrofuran (2.5 ml) was treated at 0°C with N,N-diisopropylethylamine (340 μ l, 2.0 mmol) and (2R)-1-chloro-1-oxopropan-2-yl acetate (109 mg, 726 μ mol) and stirred at 0°C for 30 min. 2-hydrazinyl-3-(trifluoromethyl)pyridine (76 μ l, 730 μ mol) and anhydrous zinc sulfate (107 mg, 660 μ mol) were added and the resulting mixture was stirred 1 h at 45°C. The reaction mixture was diluted with ethyl acetate and washed with water (3 times) followed by brine. The organic phase was dried over magnesium sulfate and evaporated. The residue was purified by preparative HPLC (Method 4). The product containing fraction were evaporated and the residue purified by preparative HPLC (Method 4) affording 101 mg (24 % of th.) of the title compound.


LC-MS (Method 5): $R_t = 1.94$ min; MS (ESIpos): $m/z = 620.1$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-d₆) δ [ppm]: 8.90 (d, 1H), 8.63-8.52 (m, 1H), 7.98-7.88 (m, 1H), 7.78-7.56 (m, 4H), 6.89 (d, 1H), 5.96 (q, 1H), 5.21-5.02 (m, 2H), 4.41-4.22 (m, 1H), 4.08-3.76 (m, 2H), 1.72 (s, 3H), 1.58 (d, 3H).

5 EXPERIMENTAL SECTION – EXAMPLES

Example 1

5-(4-Chlorophenyl)-2-({1-(4-chloropyridin-3-yl)-5-[(1R)-1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




- 10 A solution of (1R)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(4-chloropyridin-3-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 3A, 168 mg, 287 μmol) in methanol (1.7 ml) was treated with an aqueous sodium hydroxide solution (290 μl, 1.0 M, 290 μmol) and stirred for 1.5 h at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture
- 15 was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 103 mg (65 % of Th.) of the title compound.

LC-MS (Method 1): R_t = 0.87 min; MS (ESIpos): m/z = 544 [M+H]⁺

¹H-NMR (400 MHz, DMSO-d₆) δ [ppm] = 8.82-8.64 (m, 2H), 7.92-7.56 (m, 5H), 7.01-6.78 (m, 1H), 5.50 (br. s, 1H), 5.08 (m, 2H), 4.72 (q, 1H), 4.30 (m, 1H), 4.08-3.76 (m, 2H), 1.41 (d, 3H).

20 Example 2

5-(4-Chlorophenyl)-2-({1-(4-chloropyridin-3-yl)-5-[(1S)-1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




At 0°C, a solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(4-chloropyridin-3-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 4A, 182 mg, 310 μmol) in methanol (1.8 ml) was treated with an aqueous sodium hydroxide solution (310 μl, 1.0 M, 310 μmol) and stirred for 1.5 h at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 103 mg (59 % of Th.) of the title compound.

10 LC-MS (Method 1): $R_t = 0.87$ min; MS (ESIpos): $m/z = 544$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 8.82-8.61 (m, 2H), 7.89-7.57 (m, 5H), 6.96-6.83 (m, 1H), 5.50 (br. s, 1H), 5.08 (m, 2H), 4.72 (q, 1H), 4.29 (m, 1H), 4.08-3.74 (m, 2H), 1.41 (d, 3H).

Example 3

15 5-(4-Chlorophenyl)-2-({1-(3-chloropyridin-2-yl)-5-[(1S)-1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




A solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(3-chloropyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 5A, 72.0 mg, 123 μmol) in methanol (730 μl) was treated with an aqueous sodium hydroxide solution (120 μl, 1.0 M, 120 μmol) and stirred 30 min at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 63.3 mg (89 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.88$ min; MS (ESIpos): $m/z = 544$ $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$): δ [ppm] = 8.57 (dd, 1H), 8.25 (dd, 1H), 7.78-7.58 (m, 5H), 6.90 (d, 1H), 5.50 (d, 1H), 5.07 (m, 2H), 4.83 (quin, 1H), 4.30 (m, 1H), 4.10-3.78 (m, 2H), 1.43 (d, 3H).

Example 4

- 5 5-(4-Chlorophenyl)-2-({1-(3,5-dichloropyridin-4-yl)-5-[(1R)-1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




- At 0°C , a solution of (1R)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(3,5-dichloropyridin-4-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 6A, 100 mg, 161 μmol) in methanol (960 μl) was treated with an aqueous sodium hydroxide solution (160 μl , 1.0 M, 160 μmol) and stirred for 1.5 h at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol and the filtrate was evaporated. The residue was purified via preparative HPLC (Chromatorex C18, 10 μm , 125 x 30 mm. water/acetonitrile-gradient 0.1% formic acid). affording 21.0 mg (23 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.96$ min; MS (ESIpos): $m/z = 578$ $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ [ppm] = 8.90 (s, 2H), 7.82-7.52 (m, 4H), 6.89 (d, 1H), 5.69 (d, 1H), 5.18-4.97 (m, 2H), 4.79 (quin, 1H), 4.28 (m, 1H), 4.11-3.74 (m, 2H), 1.44 (d, 3H).

20 **Example 5**

- 5-(4-Chlorophenyl)-2-({1-(3-chloropyridin-2-yl)-5-[(1R)-1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




At 0°C, a solution of (1R)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(3-chloropyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 7A, 253 mg, 431 μmol) in methanol (2.6 ml) was treated with an aqueous sodium hydroxide solution (430 μl, 1.0 M, 430 μmol) and stirred 30 min at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 234 mg (84 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.90$ min; MS (ESIpos): $m/z = 544$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 8.57 (dd, 1H), 8.25 (dd, 1H), 7.80-7.57 (m, 5H), 6.90 (d, 1H), 5.50 (d, 1H), 5.07 (m, 2H), 4.83 (quin, 1H), 4.30 (m, 1H), 4.09-3.78 (m, 2H), 1.43 (d, 3H).

Example 6

5-(4-Chlorophenyl)-2-({1-(3-chloropyridin-4-yl)-5-[(1R)-1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one



15

At 0°C, a solution of (1R)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(3-chloropyridin-4-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 8A, 141 mg, 240 μmol) in methanol (1.4 ml) was treated with an aqueous sodium hydroxide solution (240 μl, 1.0 M, 240 μmol) and stirred 30 min at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 121 mg (93 % of Th.) of the title compound.


LC-MS (Method 1): $R_t = 0.89$ min; MS (ESIpos): $m/z = 544$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 8.91 (s, 1H), 8.73 (d, 1H), 7.80-7.56 (m, 5H), 6.89 (d, 1H), 5.58 (d, 1H), 5.15-4.97 (m, 2H), 4.77 (quin, 1H), 4.29 (m, 1H), 4.07-3.75 (m, 2H), 1.42 (d, 3H).

Example 7

25

5-(4-Chlorophenyl)-2-({1-(3-chloropyridin-4-yl)-5-[(1S)-1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




At 0°C, a solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(3-chloropyridin-4-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 9A, 122 mg, 208 μmol) in methanol (1.2 ml) was treated with an aqueous sodium hydroxide solution (210 μl, 1.0 M, 210 μmol) and stirred 2.5 h at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 98.9 mg (87 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.89$ min; MS (ESIpos): $m/z = 544$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 8.91 (s, 1H), 8.73 (d, 1H), 7.79-7.56 (m, 5H), 6.89 (d, 1H), 5.59 (d, 1H), 5.08 (m, 2H), 4.77 (quin, 1H), 4.36-4.22 (m, 1H), 4.08-3.78 (m, 2H), 1.42 (d, 3H).

Example 8

5-(4-Chlorophenyl)-2-({5-[(1S)-1-hydroxyethyl]-1-(3-methoxypyridin-2-yl)-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one



A solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(3-methoxypyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 10A, 47.0 mg, 80.8 μmol) in (160 μl, 1.0 M, 160 μmol) was treated with an aqueous sodium hydroxide solution (1.0 ml) and stirred 1 h at room temperature. 50% aqueous


formic acid (12 μ l) was added and the resulting mixture was purified via preparative HPLC (Reprosil; C18; 10 μ m; 125 x 30 mm; water/acetonitrile- gradient 0,1% formic acid) affording 32.6 mg (75 % of Th.) of the title compound.

LC-MS (Method 1): R_t = 0.84 min; MS (ESIpos): m/z = 540 $[M+H]^+$

- 5 $^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ [ppm] = 8.14 (dd, 1H), 7.83-7.54 (m, 6H), 6.91 (d, 1H), 5.39 (d, 1H), 5.05 (m, 2H), 4.68 (quin, 1H), 4.38-4.21 (m, 1H), 4.12-3.75 (m, 5H), 1.38 (d, 3H).

Example 9

5-(4-Chlorophenyl)-2-({1-(3,5-dichloropyridin-4-yl)-5-[(1S)-1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one



10

A solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(3,5-dichloropyridin-4-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 11A, 90.0 mg, 145 μ mol) in (4.1 ml, 7.0 M, 29 mmol) was treated with an aqueous sodium hydroxide solution and stirred 30 min at room temperature. Formic acid (3 ml) was added, the resulting mixture concentrated and purified via preparative HPLC (Chromatorex; C18; 10 μ m; 125 x 30 mm; water/acetonitrile- gradient 0,1% formic acid) affording 22.3 mg (27 % of Th.) of the title compound.


15

LC-MS (Method 1): R_t = 0.92 min; MS (ESIpos): m/z = 578 $[M+H]^+$

- 20 $^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ [ppm] = 8.90 (s, 2H), 7.82-7.55 (m, 4H), 6.89 (d, 1H), 5.69 (d, 1H), 5.09 (m, 2H), 4.80 (quin, 1H), 4.28 (m, 1H), 4.07-3.78 (m, 2H), 1.44 (d, 3H).

Example 10

5-(4-Chlorophenyl)-2-({5-[(1S)-1-hydroxyethyl]-1-(1,3-thiazol-2-yl)-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




At 0°C, a solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(1,3-thiazol-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 12A, 53.0 mg, 95.0 μmol) in methanol (560 μl) was treated with an aqueous sodium hydroxide solution (95 μl, 1.0 M, 95 μmol) and stirred 2.5 h at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 48.6 mg (quant.) of the title compound.

LC-MS (Method 1): $R_t = 0.93$ min; MS (ESIpos): $m/z = 516$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 7.82-7.54 (m, 6H), 6.90 (d, 1H), 5.71-5.48 (m, 2H), 5.19-5.01 (m, 2H), 4.38-4.18 (m, 1H), 4.08-3.80 (m, 2H), 1.48 (d, 3H).

Example 11

5-(4-Chlorophenyl)-2-({1-(3-fluoropyridin-2-yl)-5-[(1S)-1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one



15

A solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(3-fluoropyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 13A, 135 mg, 237 μmol) in methanol (1.4 ml) was treated with an aqueous sodium hydroxide solution (240 μl, 1.0 M, 240 μmol) and stirred 30 min at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 94.8 mg (76 % of Th.) of the title compound.


LC-MS (Method 1): $R_t = 0.90$ min; MS (ESIpos): $m/z = 528$ [M+H]⁺

20

¹H-NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 8.45 (d, 1H), 8.08 (m, 1H), 7.84-7.51 (m, 5H), 6.90 (d, 1H), 5.53 (br d, 1H), 5.18-5.02 (m, 2H), 4.98-4.85 (m, 1H), 4.30 (m, 1H), 4.08-3.74 (m, 2H), 1.45 (d, 3H).

Example 12

- 5 5-(4-Chlorophenyl)-2-({5-[(1S)-1-hydroxyethyl]-1-[3-(trifluoromethoxy)pyridin-2-yl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




- A solution of (1S)-1-{{3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-[3-(trifluoromethoxy)pyridin-2-yl]-1H-1,2,4-triazol-5-yl} ethyl acetate (Example 14A, 80.5 mg, 127 μmol) in (250 μl, 1.0 M, 250 μmol) was treated with an aqueous sodium hydroxide solution methanol (2.0 ml) and stirred 2 h at room temperature. 50% aqueous formic acid (29 μL) was added and the resulting mixture was purified via preparative HPLC (Reprosil; C18; 10 μm; 125 x 30 mm; water/acetonitrile-gradient 0,1% HCOOH) affording 68.5 mg (87 % of Th.) of the title compound.

- 15 LC-MS (Method 3): $R_t = 2.94$ min; MS (ESIpos): $m/z = 594$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 8.61 (dd, 1H), 8.19 (dt, 1H), 7.83-7.57 (m, 5H), 6.90 (d, 1H), 5.49 (d, 1H), 5.08 (m, 2H), 4.92 (quin, 1H), 4.36-4.24 (m, 1H), 4.11-3.75 (m, 2H), 1.44 (d, 3H).

Example 13

- 20 5-(4-Chlorophenyl)-2-({1-(3-fluoropyridin-2-yl)-5-[(1R)-1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




At 0°C, a solution of (1R)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(3-fluoropyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 15A, 184 mg, 323 μmol) in methanol (1.9 ml) was treated with an aqueous sodium hydroxide solution (320 μl, 1.0 M, 320 μmol) and stirred 30 min at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 167 mg (98 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.90$ min; MS (ESIpos): $m/z = 528$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 8.45 (d, 1H), 8.08 (ddd, 1H), 7.81-7.54 (m, 5H), 6.90 (d, 1H), 5.52 (br d, 1H), 5.08 (s, 2H), 4.98-4.87 (m, 1H), 4.30 (m, 1H), 4.09-3.76 (m, 2H), 1.45 (d, 3H).

Example 14

5-(4-Chlorophenyl)-2-({5-[(1S)-1-hydroxyethyl]-1-(pyridin-2-yl)-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




At 0°C, a solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(pyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 16A, 194 mg, 352 μmol) in methanol (2.1 ml) was treated with an aqueous sodium hydroxide solution (350 μl, 1.0 M, 350 μmol) and stirred for 1.5 h at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 157 mg (88 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.94$ min; MS (ESIpos): $m/z = 510$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 8.61-8.54 (m, 1H), 8.14-8.03 (m, 1H), 7.85-7.48 (m, 6H), 6.90 (d, 1H), 5.6 (br. s, 1H), 5.41 (q, 1H), 5.17-4.99 (m, 2H), 4.31 (m, 1H), 4.10-3.78 (m, 2H), 1.49 (d, 3H).

Example 15

5-(4-Chlorophenyl)-2-({1-(5-chloropyridin-2-yl)-5-[(1S)-1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




At 0°C, a solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(5-chloropyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 17A, 154 mg, 263 μmol) in methanol (1.6 ml) was treated with an aqueous sodium hydroxide solution (260 μl, 1.0 M, 260 μmol) and stirred 2.5 h at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 141 mg (99 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 1.04$ min; MS (ESIpos): $m/z = 544$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 8.65 (d, 1H), 8.19 (dd, 1H), 7.84-7.55 (m, 5H), 6.90 (d, 1H), 5.54-5.35 (m, 2H), 5.20-5.02 (m, 2H), 4.31 (m, 1H), 4.07-3.79 (m, 2H), 1.48 (d, 3H).

15 Example 16

5-(4-Chlorophenyl)-2-({5-[(1R)-1-hydroxyethyl]-1-[3-(trifluoromethoxy)pyridin-2-yl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one



A solution of (1R)-1-{3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-[3-(trifluoromethoxy)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl acetate (Example 18A, 190 mg, 299 μmol) in (600 μl, 1.0 M, 600 μmol) was treated with an aqueous sodium hydroxide solution methanol (4.6 ml) and stirred 1 h at room temperature. 50%


aqueous formic acid (68 μ l) was added and the resulting mixture was purified via preparative HPLC (Reprosil; C18; 10 μ m; 125 x 30 mm; water/acetonitrile-gradient 0.1% formic acid) affording 162 mg (87 % of Th.) of the title compound.

LC-MS (Method 2): R_t = 1.23 min; MS (ESIpos): m/z = 594 $[M+H]^+$

- 5 $^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ [ppm] = 8.61 (dd, 1H), 8.19 (dt, 1H), 7.83-7.56 (m, 5H), 6.90 (d, 1H), 5.49 (d, 1H), 5.14-5.01 (m, 2H), 4.92 (quin, 1H), 4.30 (br d, 1H), 4.04-3.78 (m, 2H), 1.44 (d, 3H).

Example 17

- 5-(4-Chlorophenyl)-2-({1-(6-fluoropyridin-2-yl)-5-[(1S)-1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one
- 10




- At 0°C, a solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(6-fluoropyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 19A, 122 mg, 214 μ mol) in methanol (1.3 ml) was treated with an aqueous sodium hydroxide solution (210 μ l, 1.0 M, 210 μ mol) and stirred 2.5 h at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 130 mg (quant.) of the title compound.
- 15

- 20 LC-MS (Method 1): R_t = 0.97 min; MS (ESIpos): m/z = 528 $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ [ppm] = 8.24 (q, 1H), 7.79-7.57 (m, 5H), 7.32 (dd, 1H), 6.90 (d, 1H), 5.54-5.40 (m, 2H), 5.15-5.02 (m, 2H), 4.31 (m, 1H), 4.09-3.78 (m, 2H), 1.49 (d, 3H).

Example 18

- 5-(4-Chlorophenyl)-2-({5-[(1S)-1-hydroxyethyl]-1-(pyridin-4-yl)-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one
- 25




At 0°C, a solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-(pyridin-4-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 20A, 133 mg, 241 μmol) in methanol (1.4 ml) was treated with an aqueous sodium hydroxide solution (240 μl, 1.0 M, 240 μmol) and stirred 30 min at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated and purified via preparative HPLC (Chromatorex; C18; 10 μm; 250 x 20 mm; water/acetonitrile-gradient 0.1% formic acid) 55.3 mg (45 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.83$ min; MS (ESIpos): $m/z = 510$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 8.80-8.69 (m, 2H), 7.81-7.54 (m, 6H), 6.89 (d, 1H), 5.88 (d, 1H), 5.18-4.86 (m, 3H), 4.30 (m, 1H), 4.07-3.77 (m, 2H), 1.51 (d, 3H).

Example 19

15 5-(4-Chlorophenyl)-2-({5-[(1S)-1-hydroxyethyl]-1-(pyrazin-2-yl)-1H-1,2,4-triazol-3-yl}methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one



At 0°C, a solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-(pyrazin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 21A, 140 mg, 253 μmol) in methanol (1.5 ml) was treated with an aqueous sodium hydroxide solution (250 μl, 1.0 M, 250 μmol) and stirred 2.5 h at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and


washed with methanol. The filtrate was evaporated affording 120 mg (93 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.85$ min; MS (ESIpos): $m/z = 511$ $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ [ppm] = 9.05 (d, 1H), 8.84-8.61 (m, 2H), 7.84-7.53 (m, 4H),
5 6.91 (d, 1H), 5.59-5.33 (m, 2H), 5.12 (m, 2H), 4.39-4.20 (m, 1H), 4.10-3.78 (m, 2H), 1.50 (d, 3H).

Example 20

5-(4-Chlorophenyl)-2-({5-[(1S)-1-hydroxyethyl]-1-(pyridazin-3-yl)-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




10 At 0°C , a solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(pyridazin-3-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 22A, 80.0 mg, 145 μmol) in methanol (860 μl) was treated with an aqueous sodium hydroxide solution (140 μl , 1.0 M, 140 μmol) and stirred 30 min at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and
15 the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 58.8 mg (80 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.82$ min; MS (ESIpos): $m/z = 511$ $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ [ppm] = 9.35 (dd, 1H), 8.11-7.96 (m, 2H), 7.82-7.56 (m, 4H),
20 6.90 (d, 1H), 5.67-5.38 (m, 2H), 5.22-5.03 (m, 2H), 4.31 (m, 1H), 4.08-3.77 (m, 2H), 1.52 (d, 3H).

Example 21

5-(4-Chlorophenyl)-2-({5-[(1S)-1-hydroxyethyl]-1-(3-methoxypyrazin-2-yl)-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




A solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(3-methoxypyrazin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 23A, 60.0 mg, 103 μmol) in methanol (100 μl , 1.0 M, 100 μmol) was treated with an aqueous sodium hydroxide solution (1.0 ml) and stirred 30 min at room temperature. 50% aqueous formic acid (8 μl) and the resulting mixture was purified via preparative HPLC (Reprosil; C18; 10 μm ; 125 x 30 mm, water/acetonitrile-gradient 0.1% formic acid) affording 26.3 mg (46 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.90$ min; MS (ESIpos): $m/z = 541$ $[\text{M}+\text{H}]^+$

$^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$): δ [ppm] = 8.45 (d, 1H), 8.24 (d, 1H), 7.80-7.58 (m, 4H), 6.90 (d, 1H), 5.49 (d, 1H), 5.06 (s, 2H), 4.85 (quin, 1H), 4.30 (m, 1H), 4.05-3.76 (m, 5H), 1.43 (d, 3H).

Example 22

5-(4-Chlorophenyl)-2-({5-[(1S)-1-hydroxyethyl]-1-(1-methyl-1H-imidazol-2-yl)-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one



15

At 0°C , a solution of (1S)-1-[3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl} methyl)-1-(1-methyl-1H-imidazol-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 24A, 162 mg, 292 μmol) in methanol (1.7 ml) was treated with an aqueous sodium hydroxide solution (290 μl , 1.0 M, 290 μmol) and stirred 30 min at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 50.2 mg (32 % of Th.) of the title compound.


20

LC-MS (Method 1): $R_t = 0.82$ min; MS (ESIpos): $m/z = 513$ $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ [ppm] = 7.79-7.59 (m, 4H), 7.35 (d, 1H), 7.02-6.86 (m, 2H), 5.65 (d, 1H), 5.23-5.03 (m, 2H), 4.84 (quin, 1H), 4.29 (m, 1H), 4.07-3.77 (m, 2H), 3.42 (s, 3H), 1.42 (d, 3H).

5 **Example 23**

5-(4-Chlorophenyl)-2-(5-[(1R)-1-hydroxyethyl]-1-(pyridin-2-yl)-1H-1,2,4-triazol-3-yl)methyl-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




A solution of (1R)-1-[3-(3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl)methyl]-1-(pyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl acetate (Example 25A, 203 mg, 368 μmol) in methanol (2.2 ml) was treated with an aqueous sodium hydroxide solution (370 μl , 1.0 M, 370 μmol) and stirred 30 min at room temperature. 0.5 g of activated ion exchange resin (Dowex 50WX8, 200-400 mesh) were added and the resulting mixture was stirred at room temperature for 15 min. The resin was filtered off and washed with methanol. The filtrate was evaporated affording 180 mg (96 % of Th.) of the title compound.

LC-MS (Method 1): $R_t = 0.95$ min; MS (ESIpos): $m/z = 510$ $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ [ppm] = 8.58 (dd, 1H), 8.08 (td, 1H), 7.82-7.47 (m, 6H), 6.91 (d, 1H), 5.61 (br s, 1H), 5.41 (q, 1H), 5.09 (s, 2H), 4.31 (m, 1H), 4.08-3.79 (m, 2H), 1.49 (d, 3H).

Example 24

5-(4-Chlorophenyl)-2-(5-[(1-hydroxyethyl]-1-(pyridin-3-yl)-1H-1,2,4-triazol-3-yl)methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one (diastereomeric mixture)



To a solution of 5-(4-chlorophenyl)-2-{{[5-(1-hydroxyethyl)-1H-1,2,4-triazol-3-yl]methyl}}-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one (Example 28A, 300 mg, 0.69 mmol) in pyridine (6 ml) were added pyridine-3-boronic acid (170.4 mg, 1.39 mmol) and
 5 Cu(OAc)₂ (257.4 mg, 1.39 mmol). The reaction mixture was heated to 60 °C for 2 h and then stirred for 5 days at room temperature, after which boronic acid (54.5 mg, 0.35 mmol) was added. After 4 h at 60 °C, the reaction was concentrated *in vacuo*, then diluted with ethyl acetate and quenched with aqueous HCl (1 M). After phase separation, the aqueous phase was extracted twice with ethyl acetate. The combined organic phases were dried over sodium sulfate, filtered, and
 10 concentrated *in vacuo*. The crude was purified by preparative HPLC (Chromatorex C18 10µm 125x30mm, water-acetonitrile-gradient, 0.05% trifluoroacetic acid) affording 32 mg (9% of th.) of the title compound as a mixture of diastereoisomers.

LC/MS (Method 3): R_t = 2.38 min; MS [ESIpos]: m/z = 510 [M+H]⁺

¹H NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 8.84 (d, 1 H), 8.70 (dd, 1 H), 8.05 - 8.10 (m, 1 H), 7.72 -
 15 7.78 (m, 2 H), 7.58 - 7.65 (m, 3 H), 6.89 (br. s., 1 H), 5.03 - 5.13 (m, 2 H), 4.83 (q, 1 H), 4.30 (br. s., 1 H), 4.01 (dd, 1 H), 3.85 (dd, 1 H), 1.47 (d, 3 H).

The two diastereomers were separated by preparative chiral HPLC [sample preparation: 28 mg dissolved in 1 ml 2-propanol; injection volume: 0.5 ml; column: Daicel Chiralcel[®] OX-H 5 µm, 250 x 20 mm; eluent: isohexane/isopropanol 50:50; flow rate: 15 ml/min; temperature: 40°C; UV
 20 detection: 220 nm]. After separation, 8 mg of diastereomer 1 (Example 25), which eluted first, and 10 mg of diastereomer 2 (Example 26), which eluted later, were isolated.

Example 25

5-(4-Chlorophenyl)-2-({5-(1-hydroxyethyl)-1-(pyridin-3-yl)-1H-1,2,4-triazol-3-yl}methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one (diastereomer 1)

25 LC/MS (Method 3): R_t = 2.38 min; MS [ESIpos]: m/z = 510 [M+H]⁺

Analytical chiral HPLC: R_t = 9.05 min, d.e. = 100% [column: Chiralcel OX-H, 5 µm, 250 x 4.6 mm; eluent: 50% isohexane/ 30% 2-propanol + 0.2% TFA+1% H₂O; flow rate: 1 ml/min; temperature: 40°C; UV detection: 220 nm].

¹H NMR (500 MHz, DMSO-*d*₆) δ [ppm] = 8.83 (d, 1 H), 8.69 (dd, 1 H), 8.05 - 8.09 (m, 1 H), 7.73 - 7.78 (m, 2 H), 7.60 - 7.64 (m, 3 H), 6.90 (d, 1 H), 5.74 (d, 1 H), 5.08 (s, 2 H), 4.83 (quin, 1 H), 4.26 - 4.35 (m, 1 H), 4.01 (dd, 1 H), 3.85 (dd, 1 H), 1.47 (d, 3 H).

Example 26

- 5 5-(4-Chlorophenyl)-2-({5-(1-hydroxyethyl)-1-(pyridin-3-yl)-1H-1,2,4-triazol-3-yl}methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one (diastereomer 2)


LC/MS (Method 3): R_t = 2.39 min; MS [ESIpos]: m/z = 510 [M+H]⁺

- Analytical chiral HPLC: R_t = 11.39 min, d.e. = 99% [column: Chiralcel OX-H, 5 μm, 250 x 4.6 mm; eluent: 50% isohexane/ 30% 2-propanol + 0.2 TFA+1% H₂O; flow rate: 1 ml/min; temperature: 40°C; UV detection: 220 nm].

¹H NMR (500 MHz, DMSO-*d*₆) δ [ppm] = 8.83 (d, 1 H), 8.70 (dd, 1 H), 8.05 - 8.09 (m, 1 H), 7.73 - 7.78 (m, 2 H), 7.59 - 7.65 (m, 3 H), 6.90 (d, 1 H), 5.74 (d, 1 H), 5.04 - 5.13 (m, 2 H), 4.83 (quin, 1 H), 4.25 - 4.34 (m, 1 H), 4.01 (dd, 1 H), 3.85 (dd, 1 H), 1.47 (d, 3 H).

Example 27

- 15 5-(4-Chlorophenyl)-2-({1-(5-chloropyridin-3-yl)-5-(1-hydroxyethyl)-1H-1,2,4-triazol-3-yl}methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one (diastereomeric mixture)



- To a solution of 5-(4-chlorophenyl)-2-{{5-(1-hydroxyethyl)-1H-1,2,4-triazol-3-yl}methyl}-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one (Example 28A, 300 mg, 0.69 mmol) in pyridine (6 ml) were added (5-chloropyridin-3-yl)boronic acid (218 mg, 1.39 mmol) and copper(II) acetate (257.4 mg, 1.39 mmol). The reaction mixture was heated to 60 °C for 2 h and then stirred for 5 days at room temperature, after which boronic acid (54.5 mg, 0.35 mmol) was added. After 4 h at 60 °C, the reaction was concentrated *in vacuo*, then diluted with ethyl acetate and quenched with aqueous HCl (1 M). After phase separation, the aqueous phase was extracted twice with ethyl acetate. The combined organic phases were dried over sodium sulfate, filtered, and concentrated *in vacuo*. The crude was purified by preparative HPLC (Chromatorex C18 10μm

125x30mm, water-acetonitrile-gradient, 0.05% trifluoroacetic acid) affording 82.6 mg (21%) of the title compound as a mixture of diastereoisomers.

LC/MS (Method 3): $R_t = 2.77$ min; MS [ESIpos]: $m/z = 544$ [M+H]⁺

¹H-NMR (400 MHz, Hz, DMSO-*d*₆) δ [ppm] = 8.82 (d, 1 H), 8.78 (d, 1 H), 8.31 (t, 1 H), 7.72 - 7.78 (m, 2 H), 7.59 - 7.65 (m, 2 H), 6.89 (br. d, 1 H), 5.79 (br. s., 1 H), 5.03 - 5.14 (m, 2 H), 4.90 (q, 1 H), 4.30 (br. s., 1 H), 4.01 (dd, 1 H), 3.85 (dd, 1 H), 1.48 (1.48 (d, 3 H), d, 3 H).

The two diastereoisomers were separated by preparative chiral HPLC [sample preparation: 79 mg dissolved in 2 ml ethanol; injection volume: 0.7 ml; column: Daicel Chiralcel® OX-H 5 μ m, 250 x 20 mm; eluent: isohexane/ethanol 60:40; flow rate: 15 ml/min; temperature: 35°C; UV detection: 220 nm]. After separation, 32 mg of diastereomer 1 (Example 28), which eluted first, and 32 mg of diastereomer 2 (Example 29), which eluted later, were isolated.

Example 28

5-(4-Chlorophenyl)-2-({1-(5-chloropyridin-3-yl)-5-[1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one (diastereomer 1)

LC/MS (Method 3): $R_t = 2.80$ min; MS [ESIpos]: $m/z = 544$ [M+H]⁺

Analytical chiral HPLC: $R_t = 9.45$ min, d.e. = 99.2% [column: Chiralcel OX-H, 5 μ m, 250 x 4.6 mm; eluent: 70 % isohexane/ 30% ethanol + 0.2% TFA+1% H₂O; flow rate: 1 ml/min; temperature: 35°C; UV detection: 220 nm].

¹H NMR (400 MHz, DMSO-*d*₆) δ [ppm] = 8.82 (d, 1 H), 8.78 (d, 1 H), 8.31 (t, 1 H), 7.72 - 7.78 (m, 2 H), 7.60 - 7.65 (m, 2 H), 6.90 (d, 1 H), 5.79 (d, 1 H), 5.09 (s, 2 H), 4.90 (quin, 1 H), 4.24 - 4.35 (m, 1 H), 4.01 (dd, 1 H), 3.85 (dd, 1 H), 1.48 (d, 3 H).

Example 29

5-(4-Chlorophenyl)-2-({1-(5-chloropyridin-3-yl)-5-[1-hydroxyethyl]-1H-1,2,4-triazol-3-yl} methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one (diastereomer 2)


LC/MS (Method 3): $R_t = 2.80$ min; MS [ESIpos]: $m/z = 544$ [M+H]⁺

Analytical chiral HPLC: $R_t = 11.53$ min, d.e. = 100% [column: Chiralcel OX-H, 5 μ m, 250 x 4.6 mm; eluent: 70 % isohexane/ 30% ethanol + 0.2% TFA+1% H₂O; flow rate: 1 ml/min; temperature: 35°C; UV detection: 220 nm].

¹H NMR (500 MHz, DMSO-*d*₆) δ [ppm] = 8.82 (d, 1 H), 8.78 (d, 1 H), 8.30 (t, 1 H), 7.73 - 7.77 (m, 2 H), 7.60 - 7.65 (m, 2 H), 6.89 (d, 1 H), 5.80 (d, 1 H), 5.04 - 5.14 (m, 2 H), 4.90 (quin, 1 H), 4.25 - 4.34 (m, 1 H), 4.01 (dd, 1 H), 3.85 (dd, 1 H), 1.48 (d, 3 H).

Example 30

5-(4-chlorophenyl)-2-({5-[(1S)-1-hydroxyethyl]-1-[3-(trifluoromethyl)pyridin-2-yl]-1H-1,2,4-triazol-3-yl}methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one




- 5 At 0°C, (1S)-1-{{3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-[3-(trifluoromethyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl acetate (Example 29A; 103 mg, 166 μmol) was treated with a solution of ammonia in methanol (2.0 ml, 7.0 M, 14 mmol) and stirred for overnight at 0°C. The reaction mixture was acidified with formic acid and purified by preparative HPLC (Method 4) affording 63.4 mg (66 % of th.) of the title compound.

LC-MS (Method 1): $R_t = 0.94$ min; MS (ESIpos): $m/z = 578.0$ [M+H]⁺

¹H-NMR (400 MHz, DMSO-d₆) δ [ppm]: 8.87 (d, 1H), 8.50 (d, 1H), 7.92-7.56 (m, 4H), 6.95-6.84 (m, 1H), 5.45 (d, 1H), 5.06 (d, 2H), 4.90 (quin, 1H), 4.29 (br d, 1H), 4.06-3.76 (m, 2H), 1.43 (d, 3H).

15 **Example 31**

5-(4-chlorophenyl)-2-({5-[(1R)-1-hydroxyethyl]-1-[3-(trifluoromethyl)pyridin-2-yl]-1H-1,2,4-triazol-3-yl}methyl)-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-2,4-dihydro-3H-1,2,4-triazol-3-one



- 20 At 0°C, (1R)-1-{{3-({3-(4-chlorophenyl)-5-oxo-4-[(2S)-3,3,3-trifluoro-2-hydroxypropyl]-4,5-dihydro-1H-1,2,4-triazol-1-yl}methyl)-1-[3-(trifluoromethyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl acetate (Example 30A, 107 mg, 173 μmol) was treated with an solution of ammonia in methanol (2.9 ml, 7.0 M, 21 mmol), stirred for 1.5 h at 0°C and 30 min at room temperature. The

reaction mixture was acidified with formic acid and purified by preparative HPLC (Method 4) affording 68.4 mg (69 % of th.) of the title compound.

LC-MS (Method 5): $R_t = 1.70$ min; MS (ESIpos): $m/z = 578.1$ $[M+H]^+$

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ [ppm]: 8.87 (d, 1H), 8.54-8.46 (m, 1H), 7.87 (dd, 1H), 7.78-7.57 (m, 4H), 6.94-6.86 (m, 1H), 5.45 (d, 1H), 5.15-4.99 (m, 2H), 4.90 (quin, 1H), 4.39 – 4.20 (br m, 1H), 4.05-3.76 (m, 2H), 1.43 (d, 3H).

EXPERIMENTAL SECTION – BIOLOGICAL ASSAYS

Abbreviations and Acronyms:

Acc. No.	accession number
AVP	arginine vasopressin
B_{\max}	maximal ligand binding capacity
BSA	bovine serum albumin
cAMP	cyclic adenosine monophosphate
Cat. No.	catalogue number
cDNA	complementary deoxyribonucleic acid
CHO	chinese hamster ovary
CRE	cAMP response element
Ct	cycle threshold
DMEM/F12	Dulbecco's modified Eagle's medium / Ham's F12 medium (1:1)
DNA	deoxyribonucleic acid
DMSO	dimethylsulfoxide
DTT	dithiothreitol
EC_{50}	half-maximal effective concentration
EDTA	ethylenediamine-tetraacetic acid
FAM	carboxyfluorescein succinimidyl ester
f.c.	final concentration
FCS	fetal calf serum
HEPES	4-(2-hydroxyethyl)piperazine-1-ethanesulfonic acid

IC ₅₀	half-maximal inhibitory concentration
K _d	dissociation constant
K _i	dissociation constant of an inhibitor
mRNA	messenger ribonucleic acid
PBS	phosphate buffered saline
PEG	polyethylene glycol
p.o.	<i>per os</i> , peroral
RNA	ribonucleic acid
RTPCR	real-time polymerase chain reaction
SPA	scintillation proximity assay
TAMRA	carboxytetramethylrhodamine
TRIS; Tris	2-amino-2-hydroxymethylpropane-1,3-diol

Demonstration of the activity of the compounds of the present invention may be accomplished through *in vitro*, *ex vivo*, and *in vivo* assays that are well known in the art. For example, to demonstrate the activity of the compounds of the present invention, the following assays may be used.

5 B-1. Cellular *in vitro* assay for determining vasopressin receptor activity

The identification of agonists and antagonists of the V1a and V2 vasopressin receptors from humans, rats and dogs as well as the quantification of the activity of the compounds of the invention is carried out using recombinant cell lines. These cell lines originally derive from a hamster's ovary epithelial cell (Chinese Hamster Ovary, CHO K1, ATCC: American Type Culture Collection, Manassas, VA 20108, USA). The test cell lines constitutively express the human, rat or dog V1a or V2 receptors. In case of the G_{αq}-coupled V1a receptors, cells are also stably transfected with a modified form of the calcium-sensitive photoproteins aequorin (human and rat V1a) or obelin (dog V1a), which, after reconstitution with the cofactor coelenterazine, emit light when there are increases in free calcium concentrations [Rizzuto R, Simpson AW, Brini M, Pozzan T, *Nature* 358, 325-327 (1992); Illarionov BA, Bondar VS, Illarionova VA, Vysotski ES, *Gene* 153 (2), 273-274 (1995)]. The resulting vasopressin receptor cells react to stimulation of the recombinantly expressed V1a receptors by intracellular release of calcium ions, which can be quantified by the resulting photoprotein luminescence. The G_s-coupled V2 receptors are stably transfected into cell lines expressing the gene for firefly luciferase under control of a CRE-responsible promoter. Activation of V2 receptors induces the activation of the CRE-responsive promoter *via* cAMP increase,

thereby inducing the expression of firefly luciferase. The light emitted by photoproteins of V1a cell lines as well as the light emitted by firefly luciferase of V2 cell lines corresponds to the activation or inhibition of the respective vasopressin receptor. The bioluminescence of the cell lines is detected using a suitable luminometer [Milligan G, Marshall F, Rees S, *Trends in Pharmacological Sciences* 17, 235-237 (1996)].

Test procedure:

Vasopressin V1a receptor cell lines:

On the day before the assay, the cells are plated out in culture medium (DMEM/F12, 2% FCS, 2 mM glutamine, 10 mM HEPES, 5 µg/ml coelenterazine) in 384-well microtiter plates and kept in a cell incubator (96% humidity, 5% v/v CO₂, 37°C). On the day of the assay, test compounds in various concentrations are placed for 10 minutes in the wells of the microtiter plate before the agonist [Arg⁸]-vasopressin at EC₅₀ concentration is added. The resulting light signal is measured immediately in a luminometer.

Vasopressin V2 receptor cell lines:

On the day before the assay, the cells are plated out in culture medium (DMEM/F12, 2% FCS, 2 mM glutamine, 10 mM HEPES) in 384-well microtiter plates and kept in a cell incubator (96% humidity, 5% v/v CO₂, 37°C). On the day of the assay, test compounds in various concentrations and the agonist [Arg⁸]-vasopressin at EC₅₀ concentration are added together to the wells, and plates are incubated for 3 hours in a cell incubator. Upon addition of the cell lysis reagent Triton™ and the substrate luciferin, luminescence of firefly luciferase is measured in a luminometer.

Table 1A below lists individual IC₅₀ values for the compounds of the invention (including racemic mixtures as well as separated enantiomers) that were obtained from cell lines transfected with the human V1a or V2 receptor:

Table 1A

Example No.	IC₅₀ hV1a [µM]	IC₅₀ hV2 [µM]	ratio IC₅₀ hV2/hV1a
1	0.00049	0.01595	32.6
2	0.00053	0.00565	10.7
3	0.00057	0.00807	14.2
4	0.00130	0.00865	6.7
5	0.00135	0.02700	20.0
6	0.00235	0.04033	17.2
7	0.00300	0.01600	5.3

Example No.	IC₅₀ hV1a [μM]	IC₅₀ hV2 [μM]	ratio IC₅₀ hV2/hV1a
8	0.00365	0.00940	2.6
9	0.00413	0.02450	5.9
10	0.00425	0.10500	24.7
11	0.00560	0.02270	4.1
12	0.00570	0.06467	11.3
13	0.00605	0.03867	6.4
14	0.00645	0.06550	10.2
15	0.00825	0.01175	1.4
16	0.01300	0.38667	29.7
17	0.01450	0.03950	2.7
18	0.02860	0.19900	7.0
19	0.02950	0.32000	10.8
20	0.04760	0.23450	4.9
21	0.04900	0.04600	0.9
22	0.06808	0.13793	2.0
23	0.08650	0.10267	1.2
24	0.04910	0.77350	15.8
25	0.09075	0.24000	2.6
26	0.03430	0.32500	9.5
27	0.01230	0.01875	1.5
28	0.02055	0.05175	2.5
29	0.00771	0.01520	2.0
30	0.00330	0.04850	14.9
31	0.00490	0.17000	35.1

The IC₅₀ data listed in Table 1A demonstrate that the compounds of the present invention are acting as selective and potent vasopressin V1a receptor antagonists.

B-2. Radioactive binding assay

IC₅₀ and K_i values can be determined in radioactive binding assays using membrane fractions of recombinant human embryonic kidney cell line 293 (HEK293) or CHO-K1 cell lines expressing the respective human vasopressin V1a and V2 receptors.

5 Human recombinant vasopressin V1a receptors expressed in HEK293 cells are used in 50 mM Tris-HCl buffer, pH 7.4, 5 mM MgCl₂, 0.1% BSA using standard techniques. Aliquots of prepared membranes are incubated with test compounds in various concentrations in duplicates and 0.03nM [¹²⁵I]Phenylacetyl-D-Tyr(Me)-Phe-Gln-Asn-Arg-Pro-Arg-Tyr-NH₂ for 120 minutes at 25°C. Non-specific binding is estimated in the presence of 1 μM [Arg⁸]Vasopressin. Receptors are filtered and
10 washed, the filters are then counted to determine [¹²⁵I]Phenylacetyl-D-Tyr(Me)-Phe-Gln-Asn-Arg-Pro-Arg-Tyr-NH₂ specifically bound.

CHO-K1 cells stably transfected with a plasmid encoding human vasopressin V2 receptor are used to prepare membranes in 50 mM Tris-HCl buffer, pH 7.4, 10 mM MgCl₂, 0.1% BSA using standard techniques. Aliquots of prepared membrane are incubated with test compounds in various
15 concentrations in duplicates and 4 nM [³H](Arg⁸)-Vasopressin for 120 minutes at 25°C. Non-specific binding is estimated in the presence of 1 mM (Arg⁸)-vasopressin. Membranes are filtered and washed 3 times and the filters are counted to determine [³H](Arg⁸)-Vasopressin specifically bound.

IC₅₀ values are determined by a non-linear, least squares regression analysis using MathIQTTM (ID
20 Business Solutions Ltd., UK). The inhibition constant K_i is calculated using the equation of Cheng and Prusoff (Cheng, Y., Prusoff, W.H., Biochem. Pharmacol. 22:3099-3108, 1973).

B-3. Cellular in vitro assay for detecting the action of vasopressin V1a receptor antagonists on the regulation of pro-fibrotic genes

The cell line H9C2 (American Type Culture Collection ATCC No. CRL-1446), described as a
25 cardiomyocyte type isolated from rat cardiac tissue, endogenously expresses the vasopressin V1a receptor AVPR1A in high copy number, whereas AVPR2 expression cannot be detected. Likewise, the cell line NRK49F (ATCC No. CRL1570) isolated from rat kidney tissue, shows similar expression pattern of high AVPR1A mRNA expression and diminishing AVPR2 expression. For cell assays detecting the inhibition of AVPR1A receptor-dependent regulation of gene expression
30 by receptor antagonists, the procedure is as follows:

H9C2 cells or NRK49F cells are seeded in 6-well microtiter plates for cell culture at a cell density of 50 000 cells/ well in 2.0 ml of Opti-MEM medium (Invitrogen Corp., Carlsbad, CA, USA, Cat. No. 11058-021) and held in a cell incubator (96% humidity, 8% v/v CO₂, 37°C). After 24 hours, sets of three wells (triplicate) are charged with vehicle solution (negative control) and vasopressin
35 solution ([Arg⁸]-vasopressin acetate, Sigma, Cat. No. V9879), or test compound (dissolved in

vehicle: water with 20% v/v ethanol) and vasopressin solution. In the cell culture, the final vasopressin concentration is 1 nM. The test compound solution is added to the cell culture in small volumes, so that a final concentration of 0.03% of ethanol in the cell assay is not exceeded. After an incubation time of 5 hours, the culture supernatant is drawn off under suction, the adherent cells
5 are lysed in 350 μ l of RLT buffer (Qiagen, Cat. No. 79216), and the RNA is isolated from the lysate using the RNeasy kit (Qiagen, Cat. No. 74104). This is followed by DNase digestion (Invitrogen, Cat. No. 18068-015), cDNA synthesis (Promega, ImProm-II Reverse Transcription System, Cat. No. A3800) and Reverse Transcription Polymerase Chain Reaction (RT-PCR) (pPCR MasterMix RT-QP2X-03-075, Eurogentec, Seraing, Belgium). All procedures take place in
10 accordance with the working protocols of the test reagents' manufacturers. The primer sets for the RT-PCR are selected on the basis of the mRNA gene sequences (NCBI GenBank Entrez Nucleotide Data Base) using the Primer3Plus program with 6-FAM TAMRA-labelled probes. The RT-PCR for determining the relative mRNA expression in the cells of the various assay batches is carried out using the Applied Biosystems ABI Prism 7700 Sequence Detector in 384-well microtiter plate
15 format in accordance with the instrument operating instructions. The relative gene expression is represented by the delta-delta Ct value [Applied Biosystems, User Bulletin No. 2 ABI Prism 7700 SDS, December 11, 1997 (updated 10/2001)] with reference to the level of expression of the ribosomal protein L-32 gene (GenBank Acc. No. NM_013226) and the threshold Ct value of Ct = 35.

20 **B-4. Inhibition of vasopressin induced aggregation of human platelets**

Human platelets endogenously express the V1a receptor. It was found that relatively high vasopressin concentrations (ca. 50-100 nM) stimulate platelet aggregation *ex vivo*. Therefore, platelets enriched from human blood may serve as a V1a expressing tissue for pharmacological studies with corresponding high concentrations of vasopressin antagonists.

25 Human blood is collected in a 10 mM trisodium citrate solution by venous puncture from nonsmoking healthy volunteers (n=4-8) who were drug free for at least 1 week. Platelet-rich plasma (PRP) is obtained by centrifuging the blood sample at 140 g for 20 min at 4°C. The resulting pellet is further centrifuged (15.000 rpm, 2 min) to produce platelet-poor plasma (PPP). Platelet aggregation is measured turbidimetrically using an aggregometer (APACT 4). The reaction
30 is followed by monitoring changes in light transmission on 178 μ L PRP aliquots, under continuous stirring at 37°C, against PPP control. Various concentrations of vasopressin antagonists (in 2 μ L) are added to PRP 5 min before the addition of 20 μ L Arg-vasopressin (final concentration 100 nM). The inhibitory effects of the compounds are determined by measuring the height of the aggregation wave from the bottom of the shape change compared with the control response. IC50 values are
35 calculated a dose-response inhibition curve by an iterative nonlinear regression program

B-5. Effects on the contraction of isolated rat vessel rings**Isolated aorta**

Test compounds can be investigated on isolated aortic rings from male Wistar rats endogenously expressing the V1a receptor. Male Wistar rats are euthanized using carbon dioxide. The aorta is removed and placed in ice-cold Krebs-Henseleit buffer of following composition (in mmol/l): NaCl 112, KCl 5.9, CaCl₂ 2.0, MgCl₂ 1.2, NaH₂PO₄ 1.2, NaHCO₃ 25, glucose 11.5. The aorta is cut into 3 mm rings and transferred to 20 ml organ baths containing Krebs-Henseleit solution equilibrated with 95% O₂, 5% CO₂ at 37°C. For recording of isometric tension the rings are mounted between two hooks. The resting tension is adjusted to 3 g. After an equilibration period, each experiment is started by exposing the preparation to K⁺ (50 mM) Krebs-Henseleit solution. The aortic rings are then pre-contracted using 1 nmol/l Arg-vasopressin. After a stable contraction is established, a cumulative dose response curve of the test compound is constructed. The stabilized contraction induced by Arg-vasopressin is defined as 100% tension. The relaxation is expressed as percentage tension.

15 Isolated A. renalis

Male Wistar rats (200-250 g) are euthanized using carbon dioxide. The A. renalis is removed and placed in ice-cold Krebs-Henseleit buffer of following composition (in mmol/l): NaCl 112, KCl 5.9, CaCl₂ 2.0, MgCl₂ 1.2, NaH₂PO₄ 1.2, NaHCO₃ 25, glucose 11.5. For measurement of isometric tension, ring segments, 2 mm in length, are mounted in a small vessel chamber myograph (Danish Myo Technology A/S, Denmark) using two tungsten wires fixed to mounting jaws. One mounting jaw is attached to a micrometer, allowing control of vessel circumference. The other mounting jaw is attached to a force transducer for measurement of tension development. The whole preparation is kept in a chamber with physiological salt solution at 37°C, bubbled with oxygen. After a 30 min equilibration period, the vessels are stretched to their optimal lumen diameter for active tension development which is determined based on the internal circumference-wall tension ratio. The internal circumference is set to 90% of what the vessels would have if they are exposed to a passive tension equivalent to that produced by a transmural pressure of 100 mmHg.

Afterwards, the vessels are washed three times with Krebs-Henseleit buffer and left to equilibrate for 30 min. The contractility is then tested by a twofold exposure to a high K⁺ solution (50 mmol/l KCl). After washing with Krebs-Henseleit buffer the vessels are then pre-contracted using 1 nmol/l Arg-vasopressin. After a stable contraction is established, a cumulative dose response curve of the test compound is constructed. The stabilized contraction induced by Arg-vasopressin is defined as 100% tension. The relaxation is expressed as percentage tension.

B-6. In vivo assay for detecting cardiovascular effects: blood pressure measurement in anaesthetized rats (vasopressin 'challenge' model)

Male Sprague-Dawley rats (250-350 g body weight) are used under ketamine/ xylazine/ pentobarbital injection anaesthesia. Polyethylene tubes (PE-50, Intramedic®), prefilled with heparin-containing (500 IU/ml) isotonic sodium chloride solution, are introduced into the jugular vein and the femoral vein and then tied in. Arg-vasopressin (SIGMA) is injected *via* one venous access, with the aid of a syringe; the test substance is administered *via* the second venous access. For determination of the systolic blood pressure, a pressure catheter (Millar SPR-320 2F) is tied into the carotid artery. The arterial catheter is connected to a pressure transducer which feeds its signals to a recording computer equipped with suitable recording software. In a typical experiment, the experimental animal is administered 3-4 successive bolus injections at intervals of 10-15 min with a defined amount of Arg-vasopressin (30 ng/kg) in isotonic sodium chloride solution. When the blood pressure has reached initial levels again, the test substance is administered as a bolus, with subsequent continuous infusion, in a suitable solvent. After this, at defined intervals (10-15 min), the same amount of Arg-vasopressin as at the start is administered again. On the basis of the blood pressure values, a determination is made of the extent to which the test substance counteracts the hypertensive effect of Arg-vasopressin. Control animals only receive solvent instead of the test substance.

Following intravenous administration, the compounds of the invention, in comparison to the solvent controls, bring about an inhibition of the blood pressure increase caused by Arg-vasopressin.

B-7. In vivo assay for detecting protective renal effects: Acute ischemia/reperfusion injury model in rodents

Laboratory bred male C57Bl/6J mice 6-8 weeks old are obtained from Taconic Biosciences, male 6-8 weeks old Sprague Dawley® rat are obtained from Charles River. Both rats and mice are maintained under standard laboratory conditions, 12 hour light-dark cycles with access to normal chow and drinking water at libitum. For the ischemia reperfusion injury model a total of 10-12 rats or mice is used in each control and experimental group.

Animals are anesthetized with continuous inhaled isoflurane. A right nephrectomy is performed through a right flank incision 7 days before the ischemic procedures in the contralateral kidneys. For renal ischemia a left flank incision is made. Renal vessels are exposed by dissection of the left renal pedicle. Non-traumatic vascular clamps are used to stop blood flow (artery and vein) during 45 min (rats) or 25 min (mice) of ischemia. Reperfusion is established by removing the clamps. The abdominal wall (muscular layer and skin) is closed with 5.0 polypropylene sutures. Temgesic® (Buprenorphin, 0.025 mg/kg s.c.) is applied as an analgesic.

Urine of each animal is collected in metabolic cages over night before sacrifice at 24h post ischemia. Upon sacrifice, blood samples are obtained under terminal anesthesia. After centrifugation of the blood samples, serum is isolated. Both serum creatinine and serum urea are measured via clinical biochemistry analyzer (Pentra 400). For the assessment of serum and urinary kidney injury biomarkers (Neutrophil gelatinase-associated lipocalin [NGAL], kidney injury molecule- 1 [KIM-1] and Osteopontin) ELISA's are performed according to the manufacturers protocol. Both urinary creatinine and albumin are measured to determine the albumin/creatinine ratio.

Total RNA is isolated from kidneys. Left kidneys are snap-frozen in liquid nitrogen at sacrifice. Kidney tissue is then homogenized and RNA is obtained. Total RNA is transcribed to cDNA. Using TaqMan real-time PCR renal NGAL, Osteopontin, KIM-1, Nephhrin and Podocin mRNA expression is analyzed in whole kidney tissue.

Differences between groups are analyzed by one-way ANOVA with Dunnett's corrections for multiple comparisons. Statistical significance is defined as $p < 0.05$. All statistical analyses are done using GraphPad Prism 6.

B-8. In vivo assay for detecting cardiovascular effects: hemodynamic investigations in anaesthetized dogs

Male beagle dogs (Beagle, Marshall BioResources, USA) with a weight of between 10 and 15 kg are anesthetized with pentobarbital (30 mg/kg iv, Narcoren®, Merial, Germany) for the surgical interventions and the hemodynamic and functional investigation termini. Pancuroniumbromide (Pancuronium Inresa, Inresa, Germany, 2-4 mg/animal i.v.) serves additionally as a muscle relaxant. The dogs are intubated and ventilated with an oxygen/ambient air mixture (30/70%), about 2,5-4 L/min. Ventilation takes place using a ventilator from GE Healthcare (Avance, Germany) and is monitored using a carbon dioxide analyzer (-Datex Ohmeda). The anesthesia is maintained by continual infusion of pentobarbital (50 µg/kg/min); fentanyl is used as an analgesic (10 µg/kg/h).

In preparatory interventions, the dogs are fitted with a cardiac pacemaker. At start of experiment, a cardiac pacemaker from Biotronik (Logos®, Germany) is implanted into a subcutaneous skin pocket and is contacted with the heart via a pacemaker electrode (Siello S60®, Biotronik, Germany) which is advanced through the external jugular vein, with illumination, into the right ventricle.

Thereafter accesses are removed and the dog wakes spontaneously from the anesthesia. After a further 7 days, the above-described pacemaker is activated and the heart is stimulated at a frequency of 220 beats per minute.

The actual drug testing experiments take place 28 days after the beginning of pacemaker stimulation, using the following instrumentation:

- Introduction of a bladder catheter for bladder relief and for measuring the flow of urine
- Attachment of electrocardiography (ECG) leads to the extremities for ECG measurement
- 5 • Introduction of a sheath introducer filled with sodium chloride solution into the femoral artery. This tube is connected to a pressure sensor (Braun Melsungen, Melsungen, Germany) for measuring the systemic blood pressure
- Introduction of a Millar Tip catheter (type 350 PC, Millar Instruments, Houston, USA) through a port secured in the carotid artery, for measuring cardiac hemodynamics .
- 10 • Introduction of a Swan-Ganz catheter (CCOmbo 7.5F, Edwards, Irvine, USA) via the jugular vein into the pulmonary artery, for measuring the cardiac output, oxygen saturation, pulmonary arterial pressures and central venous pressure
- Siting of a venous catheter in the cephalic vein, for infusing pentobarbital, for liquid replacement and for blood sampling (determination of the plasma levels of substance or other clinical blood values)
- 15 • Siting of a venous catheter in the saphenous vein, for infusing fentanyl and for administration of substance
- Infusion of vasopressin (Sigma) in increasing dosage, up to a dose of 4 mU/kg/min. The pharmacological substances are then tested with this dosage.
- 20 The primary signals are amplified if necessary (ACQ7700, Data Sciences International, USA or Edwards-Vigilance-Monitor, Edwards, Irvine, USA) and subsequently fed into the Ponemah system (Data Sciences International, USA) for evaluation. The signals are recorded continuously throughout the experimental period, and are further processed digitally by said software, and averaged over 30 seconds.

25 **C) Working examples of pharmaceutical compositions**

The substances according to the invention can be converted to pharmaceutical preparations as follows:

Tablet:

Composition:

- 30 100 mg of the compound of Example 1, 50 mg of lactose (monohydrate), 50 mg of maize starch, 10 mg of polyvinylpyrrolidone (PVP 25) (from BASF, Germany) and 2 mg of magnesium stearate.

Tablet weight 212 mg. Diameter 8 mm, radius of curvature 12 mm.

Production:

The mixture of the compound of Example 1, lactose and starch is granulated with a 5% strength solution (m/m) of the PVP in water. After drying, the granules are mixed with the magnesium stearate for 5 min. This mixture is compressed in a conventional tableting press (see above for format of the tablet).

Oral suspension:

Composition:

1000 mg of the compound of Example 1, 1000 mg of ethanol (96%), 400 mg of Rhodigel (xanthan gum) (from FMC, USA) and 99 g of water.

10 ml of oral suspension correspond to a single dose of 100 mg of the compound of the invention.

Production:

The Rhodigel is suspended in ethanol, and the compound of Example 1 is added to the suspension. The water is added while stirring. The mixture is stirred for about 6 h until swelling of the Rhodigel is complete.


Sterile i.v. solution:

The compound according to the invention is dissolved at a concentration below saturation solubility in a physiologically acceptable solvent (for example isotonic sodium chloride solution, glucose solution 5% and/or PEG 400 solution 30%). The solution is sterilized by filtration and filled into sterile and pyrogen-free injection containers.

Although the invention has been disclosed with reference to specific embodiments, it is apparent that other embodiments and variations of the invention may be devised by others skilled in the art without departing from the true spirit and scope of the invention. The claims are intended to be construed to include all such embodiments and equivalent variations.

CLAIMS:


1. A compound of general formula (I):



in which

- 5 R^1 represents a group selected from a hydrogen atom and methyl,
- Ar represents a 5- or 6-membered heteroaryl group attached via a ring carbon atom having one or two ring heteroatoms selected from a nitrogen atom and a sulfur atom,
- wherein any 5- or 6-membered heteroaryl group are each optionally substituted, identically or differently, with one or two groups selected from a halogen atom, nitro, cyano, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-alkylsulfanyl, (C₁-C₄)-alkoxycarbonyl, aminocarbonyl and -S(=O)₂NH₂,
- 10 wherein said (C₁-C₄)-alkyl group, said (C₁-C₄)-alkoxy group and said (C₁-C₄)-alkylsulfanyl group are each optionally substituted with up to three fluorine atoms,
- 15 or a pharmaceutically acceptable salt, hydrate and/or solvate thereof.
2. A compound of general formula (I) according to Claim 1, wherein
- R^1 represents a methyl group,
- Ar represents a group selected from pyridine, pyrazine, pyridazine, 1-methylimidazole, 1,3-thiazole,
- 20 wherein any pyridine group is each optionally substituted, identically or differently, with one or two groups selected from a fluorine atom, a chlorine atom, methoxy, trifluoromethyl and trifluoromethoxy,
- wherein any pyrazine group is optionally substituted with methoxy,
- or a pharmaceutically acceptable salt, hydrate and/or solvate thereof.
- 25 3. A compound of general formula (I) according to Claim 1 or 2, wherein
- R^1 represents a methyl group,

Ar represents a group of the formula



in which

#¹ represents the point of attachment to the nitrogen atom,

5 R^{2A} represents a group selected from a hydrogen atom, a chlorine atom, trifluoromethyl and trifluoromethoxy,


R^{2B} represents a group selected from a hydrogen atom and a chlorine atom

R^{2C} represents a chlorine atom,

or a pharmaceutically acceptable salt, hydrate and/or solvate thereof.

10 4. A method of preparing a compound of general formula (I) according to any one of claims 1 to 3 said method comprising the step


[A] of allowing an intermediate compound of formula (II):



in which

15 R³ represents a (C₁-C₄)-alkyl group, in particular a methyl group,

to react in a first step in the presence of an at least stoichiometric amount of a base with a compound of general formula (III):




in which

R¹ is as defined for the compound of general formula (I) according to any one of claims 1 to 3

PG represents a suitable alcohol protecting group, such as acetate,


to give an intermediate compound, which is then allowed to react in a second step with a hydrazine compound of general formula (IV) or a respective salt thereof



in which

Ar is as defined for the compound of general formula (I) according to any one of claims 1 to 3,

thereby giving an intermediate compound of general formula (V):



in which

R¹ and Ar are as defined for the compound of general formula (I) according to any one of claims 1 to 3


PG represents a suitable alcohol protecting group, such as acetyl,

followed by alcohol deprotection of (V) with a suitable deprotection agent thereby giving a compound of general formula (I) in which

R¹ and Ar are as defined for the compound of general formula (I) according to any one of claims 1 to 3,

or


[B] allowing an intermediate compound of formula (VI):



in which,

R¹ is as defined for the compound of general formula (I) according to any one of claims 1 to 3

5 and/or a tautomer thereof to react with a compound of general formula (VII):



in which

Ar is as defined for the compound of general formula (I) according to any one of claims 1 to 3, and

10 R^{XA} and R^{XB} represent, independently from each other, a hydrogen atom, or

R^{XA} and R^{XB} together form a -CH₂CH₂- or -C(CH₃)₂-C(CH₃)₂- bridge,

in the presence of a copper catalyst and an amine base thereby giving a compound of general formula (I) in which

15 R¹ and Ar are as defined for the compound of general formula (I) according to any one of claims 1 to 3,

each [A] and [B] optionally followed, where appropriate, by (i) separating the compounds of formula (I) thus obtained into their respective diastereomers, and/or (ii) converting the compounds of formula (I) into their respective hydrates, solvates, salts and/or hydrates or solvates of the salts by treatment with the corresponding solvents and/or acids or bases.

20 5. Compound for use as defined in any of Claims 1 to 3 for the treatment and/or prevention of diseases.

6. Compound as defined in any of Claims 1 to 3 for use in a method for the treatment and/or prevention of acute and chronic kidney diseases including diabetic nephropathy, acute and

chronic heart failure, preeclampsia, peripheral arterial disease (PAD), coronary microvascular dysfunction (CMD), Raynaud's syndrome and dysmenorrhea.

7. Use of a compound as defined in any of Claims 1 to 3 for the manufacture of a pharmaceutical composition for the treatment and/or prevention of acute and chronic kidney diseases including diabetic nephropathy, acute and chronic heart failure, preeclampsia, peripheral arterial disease (PAD), coronary microvascular dysfunction (CMD), Raynaud's syndrome and dysmenorrhea.
8. Pharmaceutical composition comprising a compound as defined in any of Claims 1 to 3 and one or more pharmaceutically acceptable excipients.
9. Pharmaceutical composition of Claim 8 comprising comprising one or more first active ingredients, in particular compounds of general formula (I) according to any one of claims 1 to 4, and one or more further active ingredients, in particular one or more additional therapeutic agents selected from the group consisting of diuretics, angiotensin AII antagonists, ACE inhibitors, beta-receptor blockers, mineralocorticoid receptor antagonists, antidiabetics, organic nitrates and NO donors, activators and stimulators of the soluble guanylate cyclase (sGC), antiinflammatory agents, immunosuppressive agents, phosphate binders and/or compounds which modulate vitamin D metabolism.
10. The pharmaceutical composition as defined in Claim 8 or 9 for the treatment and/or prevention of acute and chronic kidney diseases including diabetic nephropathy, acute and chronic heart failure, preeclampsia, peripheral arterial disease (PAD), coronary microvascular dysfunction (CMD), Raynaud's syndrome and dysmenorrhea.
11. Method for the treatment and/or prevention of acute and chronic kidney diseases including diabetic nephropathy, acute and chronic heart failure, preeclampsia, peripheral arterial disease (PAD), coronary microvascular dysfunction (CMD), Raynaud's syndrome and dysmenorrhea in a human or other mammal, comprising administering to a human or other mammal in need thereof a therapeutically effective amount of one or more compounds as defined in any of Claims 1 to 3, or of a pharmaceutical composition as defined in any of Claims 8 to 10.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2017/060383

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C07D403/14 C07D417/14 A61K31/435 A61K31/495 A61K31/426
 A61P9/00 A61P13/00
 ADD.
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2011/104322 A1 (BAYER PHARMA AG [DE]; FUERSTNER CHANTAL [DE]; KELDENICH JOERG [DE]; DE) 1 September 2011 (2011-09-01) cited in the application example 61; claim 1	1-11
X,P	WO 2016/071212 A1 (BAYER PHARMA AG [DE]) 12 May 2016 (2016-05-12) cited in the application claim 1	1-11

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search 8 June 2017	Date of mailing of the international search report 21/06/2017
---	---

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Wolf, Claudia
--	--

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/EP2017/060383

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2011104322	A1	01-09-2011	AU 2011219746 A1 27-09-2012 BR 112012021498 A2 28-06-2016 CA 2791100 A1 01-09-2011 CL 2012002356 A1 17-05-2013 CN 103189363 A 03-07-2013 CU 20120122 A7 30-01-2013 EA 201290836 A1 29-03-2013 EC SP12012125 A 28-09-2012 EP 2539326 A1 02-01-2013 GT 201200250 A 13-11-2014 HK 1187049 A1 08-04-2016 IL 221511 A 30-04-2015 JP 5826773 B2 02-12-2015 JP 2013520470 A 06-06-2013 KR 20130004316 A 09-01-2013 MA 34019 B1 01-02-2013 NZ 602018 A 31-10-2014 PE 06832013 A1 20-06-2013 SG 183439 A1 27-09-2012 UA 109650 C2 25-09-2015 US 2013190330 A1 25-07-2013 US 2016051518 A1 25-02-2016 WO 2011104322 A1 01-09-2011
WO 2016071212	A1	12-05-2016	AR 102487 A1 01-03-2017 AU 2015342017 A1 27-04-2017 TW 201629044 A 16-08-2016 US 2016122325 A1 05-05-2016 UY 36383 A 01-06-2016 WO 2016071212 A1 12-05-2016