



(51) International Patent Classification:

A01N 59/00 (2006.01) A01N 33/12 (2006.01)
A01P 1/00 (2006.01) A01N 31/02 (2006.01)

(21) International Application Number:

PCT/GB2012/051118

(22) International Filing Date:

18 May 2012 (18.05.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/491,974 1 June 2011 (01.06.2011) US

(71) Applicant (for all designated States except MN, US):

RECKITT BENCKISER LLC [US/US]; Morris Corporate Center IV, 399 Interpace Parkway, Parsippany, New Jersey 07054-0225 (US).

(71) Applicant (for MN only): RECKITT & COLMAN

(OVERSEAS) LIMITED [GB/GB]; 103-105 Bath Road, Slough Berkshire SL1 3UH (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): IJAZ, Khalid

[US/US]; 27 Bulger Avenue, Apt#B, New Milford, New Jersey 07646 (US). RUBINO, Joseph [US/US]; 20 Per-

cypeny Lane, Parsippany, New Jersey 07054 (US). ZHU, Yun-Peng [US/US]; 7-17 Oak Street, Fairlawn, New Jersey 07410 (US).

(74) Agents: BOWERS, Craig et al.; Reckitt Benckiser Corporate Services Limited-, Legal Department - Patents Group, Dansom Lane, Hull Humberside HU8 7DS (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available):


AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available):

ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

[Continued on next page]

(54) Title: AQUEOUS ALCOHOLIC MICROBICIDAL COMPOSITIONS COMPRISING COPPER IONS



(57) Abstract: Liquid treatment compositions comprise (or in certain preferred embodiments may consist essentially of, or may consist of): a copper source material which releases copper ions into the treatment composition, at least one alcohol which independently of other constituents present exhibits a microbicidal effect, at least one quaternary ammonium compound which provides a microbicidal benefit, optionally but very preferably also at least one detergent surfactant, further optionally one or more further constituents which impart one or more advantageous technical or aesthetic benefits to the compositions, including one or more detergent surfactants, and water, wherein the compositions are at a pH such that the surface treatment compositions, exhibit a microbicidal or antimicrobial effect on treated inanimate surfaces or when used to treat an airspace, e.g. ambient air, characterized in exhibiting a microbicidal benefit when tested against one or more challenge microorganisms, preferably against Poliovirus type 1 Sabin ("PV1"), according to one or more of the following standardized test protocols: ASTM E1052 Standard Test Method for Efficacy of Antimicrobial Agents against Viruses in Suspension, or ASTM E1053 Standard Test Method to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces, or European Standard Surface Test, EN-13697, or AOAC Germicidal Spray Products as Disinfectant Test Method, AOAC Index, 17th Ed. (2000).



TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG). **Published:**

— with international search report (Art. 21(3))

AQUEOUS ALCOHOLIC MICROBICIDAL COMPOSITIONS COMPRISING COPPER IONS

5 The present invention relates to aqueous alcoholic compositions which comprise copper ions which compositions exhibit a microbicidal benefit when applied to inanimate surfaces or when used to treat the air. The aqueous alcoholic compositions provide a surprisingly high degree of microbicidal activity against various undesirable microorganisms (sometimes referred to as ‘pathogens’) including various bacteria, mycobacteria, viruses, and fungi.

10 While ethanol and other monohydric alcohols are known to the art as having a beneficial microbicidal benefit, at the same time it is a volatile organic compound (“VOC”) and there is a substantial interest in regulating the use of ethanol (as well as other volatile organic compounds) in products wherein the ethanol or other VOC is exposed to the environment. Such regulatory interests are, however, completely contrary to the technical benefits provided by ethanol and
15 other monohydric alcohols, and in particular ethanol, as a microbicidal agent, as increased levels of ethanol in a composition have long been known to find increased microbicidal benefits against undesirable microorganisms.

 The technical art has proposed several compositions which are lauded to provide some degree of microbicidal / germicidal / antimicrobial efficacy, at the same time to comprise
20 reduced amounts of ethanol and other monohydric alcohols while still providing an appreciable microbicidal benefit. However, these compositions are not wholly successful in providing microbicidal / germicidal / antimicrobial benefit against a broad range of undesirable microorganisms, and in particular in providing effective microbicidal benefit against particularly difficult to eradicate microorganisms including non-enveloped viruses, and in particular
25 polioviruses [e.g., poliovirus type 1 (Sabin)]. As is recognized in the art, demonstrated eradication of poliovirus is highly advantageous as such compositions would not only be effective in controlling this dangerous microorganism but at the same time such a high level of efficacy would also be recognized against relatively easier to eradicate microorganisms including but not limited to bacteria, mycobacteria, other non-enveloped and enveloped virus strains
30 including fungi.

The prior art discloses various compositions which are cited to provide a microbicidal effect. For example, in US 5180749 are described largely aqueous compositions comprising about 65 – 88%wt. water and which include as further essential constituents both about 10 – 30%wt. ethanol with about 2 – 5%wt. benzyl alcohol. However, the use of water soluble metal salts is not disclosed nor is the pH of the compositions disclosed. The compositions were tested
5 against *Staphylococcus aureus*, *Salmonella choleraesuis*, *Pseudomonas aeruginosa*, rhinovirus type 39, herpes simplex 1, herpes simplex 2, adenovirus type 2, respiratory syncytial, influenza A2, influenza B, human rotavirus, *Mycobacterium tuberculosis* var. bovis, as well as fungi of types *Aspergillus niger* and *Trichopython mentgrophytes*. In that patent, when contrasting the
10 data from Table B to the data from Table A, the necessary inclusion of benzyl alcohol in conjunction with ethanol in order to achieve increased microbicidal efficacy is shown. The poor microbicidal efficacy of compositions comprising 30%wt. ethanol and water and where benzyl alcohol is absent is demonstrated on Table B. In US 3992146 are disclosed germicidal and antifungal compositions which are based on aqueous solutions of a copper compound and a
15 surfactant. The surfactants disclosed are primarily anionic surfactants based on sulfate or sulfonated organic compounds. The use of ethanol or of specific pH ranges are not clearly disclosed or demonstrated.

US 5728404 discloses certain virucidal disinfectant compositions which are described as including one or more C₁-C₄ aliphatic alcohols, 0.1 – 1%wt. of a hydrolyzed metal ion, and
20 water. Compositions comprising ethyl alcohol and isopropyl alcohol and ratios of 8:1 to 1:1 are noted to be particularly effective and preferred. While the document alleges that the amount of the aliphatic alcohol may be in the range of 40% - 90%wt., such is not demonstrated as in the four examples provided the amount of the aliphatic alcohols are respectively 80%wt., 70%wt., 80%wt. and 80%wt. Furthermore, when formed as described in that document, the composition
25 according to Example 1 of this patent document exhibited a pH of 5.48, the composition of Example 2 exhibited a pH of 5.63, and the composition of Example 3 exhibited a pH of 5.63, which indicates that the foregoing compositions consistently demonstrated an acidic pH.

US 6034043 and US 6017861 disclose liquid skin cleaning compositions comprising (1) a so-called mild surfactant system of which at least 10%wt. of which (and which preferably at
30 least 25%wt. of which), is an anionic surfactant, (2) 0.1 – 10%wt. of a polyvalent cation or cations selected from zinc, copper, tin, aluminum, cobalt, nickel, chromium, titanium, and/or

manganese and mixtures thereof, and (3) 1 – 99%wt. water wherein the cations provide antimicrobial activity. These patents suggest that microbiocidal activity of the liquid skin cleaning compositions was due to the combination of the mild surfactant system with the polyvalent cation or cations whereas the polyvalent cation or cations themselves did not provide a microbiocidal benefit. Further, none of the demonstrated compositions include lower alkyl monohydric alcohols.

US 2004/0213750 discloses aqueous alcoholic compositions which comprise 40%wt. – 70%wt. of a lower alkanol, optionally a quaternary ammonium cationic compound which itself provides germicidal properties, water and a pH adjusting agent to provide a final pH of between 7 and 13. The compositions are shown to be effective against various microorganisms including gram-positive and gram-negative types of pathogenic bacteria, as well as Poliovirus (Type 1) at a 10 minute contact time. The reference however makes no mention of the use of copper ions in the compositions.

US 2007/0184013 discloses compositions which are cited to be effective against non-enveloped virus particles. The compositions comprise a C₁-C₆ alcohol and an efficacy-enhancing amount of one or more of: cationic oligomers and polymers, proton donors, chaotropic agents, and mixtures thereof with the proviso that when the compositions include a proton donor that a cationic oligomer or polymer is also present. The cationic oligomers and polymers disclosed are defined to include cationic polyalkylene imines, cationic ethoxy polyalkylene imines, cationic poly[N-[3-(dialkylammonio)alkyl]N'[3-(alkyleneoxyalkylene dialkylammonio)alkyl]urea dichloride], vinyl caprolactam/VP/dialkylaminoalkyl alkylate copolymers and polyquaternium copolymers. The example compositions disclosed in the reference demonstrate compositions having 62%wt. and even greater amounts of the C₁-C₆ alcohol as being present.

US 2008/0045491 discloses certain surface sanitizer compositions which are described as comprising 50%-90% wt. of a water miscible alcohol component and an acid component to maintain the pH below about 5, a multivalent cation and the balance being water. The multivalent cation is described as including polymers having at least two positive charges such as polyamines, chitosan, polylysine, metal ions in metal compounds.

The treatment of biofilms by compositions which include certain heavy metals are known from US 2008/0118573. The treatment steps require that the biofilms be contacted with the said

compositions for 4 hours or more. The biofilms are defined to be conglomerates of microbial organisms embedded in highly hydrated matrices of exopolymers, typically polysaccharides, and other macromolecules.

5 US 2009/0226494 discloses certain antibacterial formulations which comprise a water-soluble copper compound, a water-soluble ammonium agent, and a water-soluble acid when the composition necessarily has an acidic pH.

10 US 2010/0233098 discloses methods and compositions for disinfecting hard surfaces which are aqueous compositions which comprise 40%wt.-70%wt. of an alcohol constituent selected from the group consisting of methanol, ethanol, n-propanol, isopropanol, n-butanol, benzyl alcohol, and mixtures thereof and a pH in the range of from about 7.0 – 14.0. The compositions may include further optional constituents, including ancillary antimicrobial agents, and surfactants, but the use of water soluble metal salts is not disclosed.

15 US 2008/0045491 discloses certain surface sanitizer compositions which are recited to include 50-90%wt. of an alcohol component, 10 – 50%wt. of water, an acid component to maintain the pH of the composition between 2 – 5, and 0.05 – 5%wt. of a multivalent cation constituent. The multivalent cation constituent may be a one of a selected list of polymers, a metal ion or a metal compound. The compositions may further optionally include one or more further constituents including oxidative agents, plant derived alkenes or essential oils, emollients, humectants, lubricants and one or more antimicrobial compounds, e.g., quaternary ammonium
20 compounds. The single example of US 2008/0045491 tested demonstrates that a composition having 78%wt. ethanol exhibits efficacy against *Candida albicans*, *Aspergillus niger*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus* and adenovirus type 5. Further examples disclosed in US 2008/0045491 are not disclosed to have been tested against any microorganisms.

25 Notwithstanding these various known art compositions, there is still an urgent need in the art to produce treatment compositions, adapted for the control or eradication of undesired microorganisms where such treatment compositions comprise reduced amounts of VOC, and in particular aliphatic alcohols which provide a microbicidal effect such as ethanol, yet which compositions are highly effective against particularly difficult to eradicate undesired
30 microorganisms, especially poliovirus, particularly where the treatment compositions are applied to an inanimate surface or are used to treat an airspace.

In a broad aspect, the compositions of the present invention are generally directed to liquid inanimate surface treatment compositions which impart an antimicrobial or microbicidal benefit to treated surfaces which compositions comprise (or in certain preferred embodiments may consist essentially of, or may consist of): a copper source material which releases copper ions into the treatment composition, at least one quaternary ammonium compound which provides a microbicidal benefit, at least one alcohol which independently of other constituents present exhibits a microbicidal effect, and optionally but very preferably, also at least one further detergic surfactant, and water. These compositions may optionally additionally include one or more further constituents which impart one or more advantageous technical or aesthetic benefits to the compositions. These compositions are at a pH such that the surface treatment compositions exhibit a microbicidal or antimicrobial or germicidal effect on treated surfaces or when used to treat an airspace, e.g. ambient air. Preferably these liquid inanimate surface treatment compositions are characterized in exhibiting a microbicidal benefit when tested against one or more challenge organisms according to one or more of the following standardized test protocols: ASTM E1052 Standard Test Method for Efficacy of Antimicrobial Agents against Viruses in Suspension, or ASTM E1053 Standard Test Method to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces, or European Standard Surface Test, EN13697, or AOAC Germicidal Spray Products as Disinfectant Test Method, AOAC Index, 17th Ed. (2000). Preferably the compositions are liquid, or sprayable liquid compositions (e.g. pumpable but are not aerosol compositions) and exhibit a viscosity of not more than about 100 cPs, preferably not more than 50 cPs, more preferably not more than 10 cPs at 20°C, when tested according to conventional quantitative methods (e.g., Brookfield Viscometer) and are pourable, readily flowable liquids. Such may be provided in any other apparatus or device wherein the liquid composition may be poured or sprayed onto a surface or into the air. The inventive compositions provide a high degree of microbicidal activity against various undesirable microorganisms (sometimes referred to as 'pathogens') including various bacteria, mycobacteria, viruses, and fungi.

Within this broad aspect, the present inventors have surprisingly observed that there may be formed compositions which exhibit a synergistic improvement in microbicidal effect when there are added to aqueous alcoholic liquid compositions containing at least one quaternary ammonium compound which provides a microbicidal benefit at specific pH ranges (especially

preferably at alkaline pH ranges), small but effective amounts of a material which provides a copper ion to the aqueous alcoholic liquid compositions, and (optionally but in most cases), especially wherein at least one further surfactant is also present. Such an effect is surprising and also particularly technically advantageous as improved microbicidal efficacy has been observed
5 against particularly difficult to control (or eradicate) microorganisms and in particular the poliovirus, while at the same time achieving these effects in aqueous alcoholic liquid compositions having a reduced VOC content. As is known to the art, non-enveloped viruses including poliovirus is particularly difficult to control or eradicate and demonstrated
10 microbicidal efficacy against poliovirus is expected to be indicative of microbicidal efficacy against other non-enveloped viruses and microorganisms which are less difficult to control or eradicate.

In a first aspect the present invention provides liquid, inanimate surface treatment compositions which impart a microbicidal benefit to such treated surfaces which compositions comprise (or in certain preferred embodiments may consist essentially of, or may consist of):
15 a copper source material which releases copper ions into the treatment composition, preferably a source of Cu(I) and/or Cu(II) ions;
at least 20%wt of at least one lower alkyl aliphatic monohydric alcohol, and preferably at least about 30%wt. of a lower alkyl aliphatic monohydric alcohol;
at least one quaternary ammonium compound which provides a microbicidal benefit;
20 water;
optionally, one or more further constituents which impart one or more advantageous technical or aesthetic benefits to the compositions, including one or more deterative surfactants;
wherein the composition has a pH of at least 5,
wherein the surface treatment compositions are characterized in exhibiting a microbicidal
25 benefit when tested against one or more challenge microorganisms according to one or more of the following standardized test protocols: ASTM E1052 Standard Test Method for Efficacy of Antimicrobial Agents against Viruses in Suspension, or ASTM E1053 Standard Test Method to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces, or European Standard Surface Test, EN1369, or AOAC Germicidal
30 Spray Products as Disinfectant Test Method, AOAC Index, 17th Ed. (2000.) against one or more challenge microorganisms, especially preferably against poliovirus type 1 (Sabin) ("PV1").

In a second aspect the present invention provides liquid, inanimate surface treatment compositions which impart a microbicidal benefit to such treated surfaces which compositions comprise (or in certain preferred embodiments may consist essentially of, or may consist of):

a copper source material which releases copper ions into the treatment composition,
5 preferably a source of Cu(I) and/or Cu(II) ions;

at least 20%wt of at least one lower alkyl aliphatic monohydric alcohol, and preferably at least about 30%wt. of a lower alkyl aliphatic monohydric alcohol;

at least one quaternary ammonium compound which provides a microbicidal benefit;

at least one deterative surfactant, and preferably at least one nonionic surfactant;

10 water;

optionally, one or more further constituents which impart one or more advantageous technical or aesthetic benefits to the compositions, including one or more deterative surfactants;

wherein the composition has a pH of at least 5,

wherein the surface treatment compositions are characterized in exhibiting a microbicidal
15 benefit when tested against one or more challenge microorganisms according to one or more of the following standardized test protocols: ASTM E1052 Standard Test Method for Efficacy of Antimicrobial Agents against Viruses in Suspension, or ASTM E1053 Standard Test Method to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces, or European Standard Surface Test, EN1369, or AOAC Germicidal
20 Spray Products as Disinfectant Test Method, AOAC Index, 17th Ed. (2000).) against one or more challenge microorganisms, especially preferably against poliovirus type 1 (Sabin) ("PV1").

Preferably according to this second aspect the at least one further deterative surfactant, is a nonionic surfactant which provides a microbicidal benefit, as compared to where such at least
25 one further deterative surfactant (which is preferably a nonionic surfactant) is absent.

According to a third aspect of the invention there are provided compositions according to the first aspect of the invention wherein the amount of the of at least one alcohol, of a lower alkyl aliphatic monohydric alcohol ranges from at least 20%wt. to 40%wt., preferably to an amount of up to, but excluding 40%wt.

30 In a further aspect of the invention there is provided a microbicidal control system of constituents which are in and of themselves effective in providing effective control of poliovirus

independently of further and optional constituents. This first microbicidal control system of constituents comprises (or consists essentially of, or consists of): water, a copper source material which releases copper ions into the treatment composition, at least 20%wt. of one or more one or more C₁-C₄ aliphatic alcohols and especially preferably wherein ethanol is the predominant or sole C₁-C₄ aliphatic alcohols present, a cationic quaternary ammonium compound which provides a microbicidal benefit and, where necessary a buffer or pH adjusting agent to impart an alkaline pH, preferably an alkaline pH of 7.5 or greater. This first microbicidal control system of constituents may thereafter optionally include further constituents which may or may not provide a further microbicidal benefit. Preferably the microbicidal control system is characterized in exhibiting a microbicidal benefit when tested against one or more challenge microorganisms according to one or more of the following standardized test protocols: ASTM E1052 Standard Test Method for Efficacy of Antimicrobial Agents against Viruses in Suspension, or ASTM E1053 Standard Test Method to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces, or European Standard Surface Test, EN1369, or AOAC Germicidal Spray Products as Disinfectant Test Method, AOAC Index, 17th Ed. (2000), especially preferably against poliovirus type 1 (Sabin) ("PV1").

In a further aspect of the invention there is provided a further microbicidal control system of constituents which are in and of themselves effective in providing effective control of poliovirus independently of further and optional constituents. This second microbicidal control system of constituents comprises (or consists essentially of, or consists of) water, a copper source material which releases copper ions into the treatment composition, at least 20%wt. of one or more one or more C₁-C₄ aliphatic alcohols and especially preferably wherein ethanol is the predominant or sole C₁-C₄ aliphatic alcohol present, a cationic quaternary ammonium compound which provides a microbicidal benefit, a surfactant constituent which is preferably at least one nonionic surfactant, and where necessary a buffer or pH adjusting agent to impart an alkaline pH, and preferably an alkaline pH of 7.5 or greater. Preferably the surfactant constituent is one or both of a nonionic surfactant and/or a cationic surfactant. This second microbicidal control system of constituents may thereafter optionally include further constituents which may or may not provide a further microbicidal benefit. Preferably this second microbicidal control system is characterized in exhibiting a microbicidal benefit when tested against one or more challenge microorganisms according to one or more of the following

standardized test protocols: ASTM E1052 Standard Test Method for Efficacy of Antimicrobial Agents against Viruses in Suspension, or ASTM E1053 Standard Test Method to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces, or European Standard Surface Test, EN1369, or AOAC Germicidal Spray Products as Disinfectant Test Method, AOAC Index, 17th Ed. (2000), especially preferably against poliovirus type 1 (Sabin) (“PV1”).

Either of the foregoing microbicidal control system of constituents may be included in a liquid, inanimate surface treatment composition.

It is to be understood that in each of the foregoing aspects the inanimate surface treatment compositions may instead or also be used as air treatment compositions for a microbicidal benefit to treated air, particularly in a volume of air or headspace, e.g, in a closed room or the interior of a vehicle.

In a further aspect the present invention provides surface treatment compositions according to any foregoing aspects of the invention which compositions exhibit a pH of at least about 5, preferably however exhibit a pH of about 6 to about 12.

In a still further aspect the present invention provides a method of controlling the incidence of undesired microorganisms on an inanimate surface, the method comprising the step of: contacting an inanimate surface which is in need of treatment or upon which the presence of one or more undesirable microorganisms are suspected or are known to be present, with an effective amount of a liquid, inanimate surface treatment composition as described herein to provide a surface treatment benefit thereto, preferably to provide a microbicidal benefit to the contacted surface.

According to a further aspect of the present invention there is provided a method of controlling the incidence of undesired microorganisms in air, or in a headspace such as the ambient air within a closed volume such as a room or the interior of a vehicle, the method comprising the step of: delivering and dispersing within an airspace an effective amount of the liquid inanimate surface treatment composition as described herein to provide a microbicidal benefit to the treated air, preferably to provide a microbicidal benefit to the treated air.

In an additional aspect the present invention provides a method for the manufacture of a vendible product which comprises a treatment composition as described herein.

These and further aspects of the invention will become more apparent from a reading of the following specification.

Figure 1 is a graph of the respective efficacy against Poliovirus type 1 (Sabin) ("PV1") of compositions later described on Table C and Table 1, at an approximate pH range of 8 – 9.

5 Figure 2 is a graph of the respective efficacy against Poliovirus type 1 (Sabin) ("PV1") of compositions later described on Table C and Table 1, at an approximate pH range of 9 - 10.

Figure 3 is a graph of the respective efficacy against Poliovirus type 1 (Sabin) ("PV1") of compositions later described on Table C and Table 1, at an approximate pH range of 10 - 11.

A first essential constituent of the invention is a copper source material which releases
10 copper ions into the treatment composition, preferably a source of Cu(I) and/or Cu(II) ions. The copper ions should be dispersible, miscible or soluble in the surface treatment compositions. Any material or compound which may function as a source of copper ions, e.g., Cu(I) and/or Cu(II) ions, to deliver or provide such copper ions into largely alcoholic liquid compositions, such as those described in this patent specification and particularly with reference to the
15 examples, may be used in the present inventive compositions. Non-limiting examples of such materials or compounds include copper sulfate, copper chloride, copper nitrate, copper oxychloride, $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$, $\text{Cu}(\text{AcO})_2 \cdot \text{H}_2\text{O}$, Cu D-gluconate, $\text{Cu}(\text{I})\text{Cl} \cdot \text{H}_2\text{O}$ or any other chemical compound or chemical species which may be used to provide Cu(I) and especially Cu(II) ions into a largely alcoholic liquid composition. Such are to be expressly understood as non-limiting
20 examples and that other materials which may function to provide copper ions may be used, e.g., further copper containing salts of organic or inorganic compounds or materials. The copper ions need not be fully soluble within the largely alcoholic liquid compositions and may, for example, be dispersions. The copper source material may be present in the treatment compositions in any effective amount but advantageously is at least about 0.001%wt. to about 2.0%wt, preferably
25 from about 0.01%wt to about 1%wt., and particularly preferably from about 0.01%wt. to about 0.5%wt. of the copper source material. Alternately, the copper source material may be present in the treatment compositions in a sufficient amount such that the copper source material releases copper ions into the treatment composition so to provide between about 1 ppm to about 10,000 ppm of Cu(I) and/or Cu(II) ions, preferably between about 20 ppm and about 5000 ppm of Cu(I)
30 and/or Cu(II) ions, yet more preferably between about 50 ppm to about 1000 ppm, and

particularly preferably between about 50 ppm to about 500 ppm of Cu(I) and/or Cu(II) ions within the inventive compositions taught herein.

Exclusive of counterions of surfactant compounds or counterions of other materials described herein which might be present, most preferably the copper source material is the sole material present in the composition which releases available metal ions to the treatment
5 compositions taught herein.

A further essential constituent of the inventive compositions is at least one lower alkyl aliphatic monohydric alcohol. Preferably this at least one of a lower alkyl aliphatic monohydric alcohol also exhibits a biocidal effect against microorganisms independently of the other
10 constituents which may be present in the compositions. Exemplary and preferred are C₁-C₆ monohydric alcohols, especially methanol, ethanol, n-propanol, isopropanol, and all isomers of butanol. Of these, C₁-C₃ monohydric alcohols and especially C₁-C₃ monohydric alcohols are preferred, especially ethanol. A single such alcohol, or mixture of two or more such alcohols, may be present. In certain embodiments when a plurality of alcohols are present, ethanol is the
15 predominant alcohol present, and especially preferably comprises at least 50.1%wt., and especially preferably and in order of increasing preference, at least 51%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99%, 99.5% and 100% by weight of the at least one lower alkyl aliphatic monohydric alcohol present. The at least one lower alkyl aliphatic monohydric alcohol comprises at least 20%wt. of the treatment composition of which it forms a
20 part. Preferably the at least one lower alkyl aliphatic monohydric alcohol constituent is present in the treatment composition in an amount of at least about 21%wt., and in order of increasing preference comprises at least 22%, 22.5%, 23%, 23.5%, 24%, 24.5%, 25%, 25.5 %, 26%, 26.5%, 27%, 27.5% 28%, 28.5%, 29%, 29.5%, 30%, 30.5%, 31%, 31.5%, 32%, 32.5%, 33%, 33.5%, 34%, 34.5%, 35%, 35.5%, 36%, 36.5%, 37%, 37.5%, 38%, 38.5%, 39%, 39.5%, 40%, 40.5%,
25 41%, 41.5%, 42%, 42.5%, 43%, 43.5%, 44%, 44.5%, 45%, 45.5%, 46%, 46.5%, 47%, 47.5%, 48%, 48.5%, 49%, 49.5%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60% 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69% and 70% by weight. Concurrently and preferably the at least one lower alkyl aliphatic monohydric alcohol constituent is present in the treatment composition in an amount of up to about 95%wt., and in order of increasing preference is present
30 in an amount up to 90%, 85%, 80%, 75%, 70%, 69%, 68%, 67%, 66%, 65%, 64%, 63%, 62%, 61%, 60%, 59%, 58%, 57%, 56%, 55%, 54%, 53%, 52%, 51%, 50%, 49.5%, 49%, 48.5%, 48%,

47.5%, 47%, 46.5%, 46%, 45.5%, 45%, 44.5% 44%, 43.5%, 43%, 42.5%, 42%, 41.5%, 41%, 40.5%, 40%, 39.5%, 39%, 38.5%, 38.5, 37.5%, 37%, 36.5%, 36%, 35.5%, 35%, 34.5%, 34%, 33.5%, 33%, 32.5%, 32%, 31.5%, 31%, 30.5%, 30%, 29.5%, 29%, 28.5%, 28%, 27.5%, 27%, 26.5%, 26%, 25.5%, 25%, 24.5%, 24%, 23.5%, 23%, 22.5%, 22% , 21.5% and 21% by weight of the treatment composition of which it forms a part.


As noted previously, in certain preferred embodiments the compositions comprise from 20%wt. to 40%wt. of at least one lower alkyl aliphatic monohydric alcohol, and preferably in an amount up to but excluding 40%wt.

Advantageously the at least one lower alkyl aliphatic monohydric alcohol constituent exhibits a microbicidal effect against one or more pathogens even in the absence of the further constituents of the treatment compositions taught herein. For this reason, C₁-C₄ monohydric aliphatic alcohols, e.g., methanol, ethanol and the various isomers of propanol are particularly preferred whether used singly or in mixtures of two or more selected C₁-C₄ monohydric aliphatic alcohols. In certain embodiments a single C₁-C₄ monohydric aliphatic alcohol is present as the second essential constituent. In certain embodiments, ethanol is the sole constituent of the lower alkyl aliphatic monohydric alcohol constituent.

A third further essential constituent is at least one quaternary ammonium compound which provides a microbicidal benefit. For purposes of the present invention described herein, such quaternary ammonium compounds are to be understood as being outside of the scope of the defined deterative surfactants as such materials are primarily provided to impart a microbicidal effect and not to provide an appreciable deterative benefit. Any cationic surfactant which satisfies these requirements may be used and is considered to be within the scope of the present invention. Mixtures of two or more cationic surface active agents, viz., cationic surfactants, may also be used. Cationic surfactants are well known and useful cationic surfactants may be one or more of those described for example in *McCutcheon's Functional Materials, Vol.2, 1998*; *Kirk-Othmer, Encyclopedia of Chemical Technology, 4th Ed., Vol. 23, pp. 481-541 (1997)*, the contents of which are herein incorporated by reference. These are also described in the respective product specifications and literature available from the suppliers of these cationic surfactants.

Examples of preferred cationic surfactant compositions useful in the practice of the instant invention are those which provide a microbicidal or germicidal effect to the compositions,


and especially preferred are quaternary ammonium compounds and salts thereof, which may be characterized by the general structural formula:



where at least one of R₁, R₂, R₃ and R₄ is a alkyl, aryl or alkylaryl substituent of from 6 to 26 carbon atoms, and the entire cation portion of the molecule has a molecular weight of at least 165. The alkyl substituents may be long-chain alkyl, long-chain alkoxyaryl, long-chain alkylaryl, halogen-substituted long-chain alkylaryl, long-chain alkylphenoxyalkyl, arylalkyl, etc. The remaining substituents on the nitrogen atoms other than the abovementioned alkyl substituents are hydrocarbons usually containing no more than 12 carbon atoms. The substituents R₁, R₂, R₃ and R₄ may be straight-chained or may be branched, but are preferably straight-chained, and may include one or more amide, ether or ester linkages. The counterion X may be any salt-forming anion which permits for the solubility of the quaternary ammonium complex within the treatment composition.

Exemplary quaternary ammonium salts within the above description include the alkyl ammonium halides such as cetyl trimethyl ammonium bromide, alkyl aryl ammonium halides such as octadecyl dimethyl benzyl ammonium bromide, N-alkyl pyridinium halides such as N-cetyl pyridinium bromide, and the like. Other suitable types of quaternary ammonium salts include those in which the molecule contains either amide, ether or ester linkages such as octyl phenoxy ethoxy ethyl dimethyl benzyl ammonium chloride, N-(laurylcocoaminoformylmethyl)-pyridinium chloride, and the like. Other very effective types of quaternary ammonium compounds which are useful as germicides include those in which the hydrophobic radical is characterized by a substituted aromatic nucleus as in the case of lauryloxyphenyltrimethyl ammonium chloride, cetylaminophenyltrimethyl ammonium methosulfate, dodecylphenyltrimethyl ammonium methosulfate, dodecylbenzyltrimethyl ammonium chloride, chlorinated dodecylbenzyltrimethyl ammonium chloride, and the like.

Preferred quaternary ammonium compounds which act as germicides and which are useful in the practice of the present invention include those which have the structural formula:



wherein R₂ and R₃ are the same or different C₈-C₁₂alkyl, or R₂ is C₁₂₋₁₆alkyl, C₈₋₁₈alkylethoxy, C₈₋₁₈alkylphenoethoxy and R₃ is benzyl, and X is a halide, for example chloride, bromide or iodide, a saccharinate counterion or is a methosulfate anion. The alkyl groups recited in R₂ and R₃ may be straight-chained or branched, but are preferably substantially linear.

Particularly useful quaternary ammonium compounds include compositions which include a single quaternary compound, as well as mixtures of two or more different quaternary compounds. Such useful quaternary compounds are available under the BARDAC®, BARQUAT®, HYAMINE®, LONZABAC®, and ONYXIDE® trademarks, which are more fully described in, for example, *McCutcheon's Functional Materials* (Vol. 2), North American Edition, 1998, as well as the respective product literature from the suppliers identified below. Such include, for example, BARDAC® 205M which is described to be a liquid containing alkyl dimethyl benzyl ammonium chloride, octyl decyl dimethyl ammonium chloride; didecyl dimethyl ammonium chloride, and dioctyl dimethyl ammonium chloride (50% active) (also available as 80% active (BARDAC® 208M)); BARDAC® 2050 which is described to be a combination of octyl decyl dimethyl ammonium chloride/didecyl dimethyl ammonium chloride, and dioctyl dimethyl ammonium chloride (50% active) (also available as 80% active (BARDAC® 2080)); BARDAC® 2250 which is described to be didecyl dimethyl ammonium chloride (50% active); BARDAC® LF (or BARDAC® LF-80), described as being based on dioctyl dimethyl ammonium chloride (BARQUAT® MB-50, MX-50, OJ-50 (each 50% liquid) and MB-80 or MX-80 (each 80% liquid) are each described as an alkyl dimethyl benzyl ammonium chloride; BARDAC® 4250 and BARQUAT® 4250Z (each 50% active) or BARQUAT® 4280 and BARQUAT® 4280Z (each 80% active) are each described as alkyl dimethyl benzyl ammonium chloride/alkyl dimethyl ethyl benzyl ammonium chloride. Also, HYAMINE® 1622, described as diisobutyl phenoxy ethoxy ethyl dimethyl benzyl ammonium chloride (50% solution); HYAMINE® 3500 (50% actives), described as alkyl dimethyl benzyl ammonium chloride (also available as 80% active (HYAMINE® 3500-80)); and HYMAINE® 2389 described as being based on methyl dodecyl benzyl ammonium chloride and/or

methyldodecylxylene-bis-trimethyl ammonium chloride. (BARDAC® , BARQUAT® and HYAMINE® are presently commercially available from Lonza, Inc., Fairlawn, New Jersey). BTC® 50 NF (or BTC® 65 NF) is described to be alkyl dimethyl benzyl ammonium chloride (50% active); BTC® 99 is described as didecyl dimethyl ammonium chloride (50% active);

5 BTC® 776 is described to be myrisalkonium chloride (50% active); BTC® 818 is described as being octyl decyl dimethyl ammonium chloride, didecyl dimethyl ammonium chloride, and dioctyl dimethyl ammonium chloride (50% active) (available also as 80% active (BTC® 818-80%)); BTC® 824 and BTC® 835 are each described as being of alkyl dimethyl benzyl ammonium chloride (each 50% active); BTC® 885 is described as a combination of BTC® 835

10 and BTC® 818 (50% active) (available also as 80% active (BTC® 888)); BTC® 1010 is described as didecyl dimethyl ammonium chloride (50% active) (also available as 80% active (BTC® 1010-80)); BTC® 2125 (or BTC® 2125 M) is described as alkyl dimethyl benzyl ammonium chloride and alkyl dimethyl ethylbenzyl ammonium chloride (each 50% active) (also available as 80% active (BTC® 2125 80 or BTC® 2125 M)); BTC® 2565 is described as alkyl

15 dimethyl benzyl ammonium chlorides (50% active) (also available as 80% active (BTC® 2568)); BTC® 8248 (or BTC® 8358) is described as alkyl dimethyl benzyl ammonium chloride (80% active) (also available as 90% active (BTC® 8249)); ONYXIDE® 3300 is described as n-alkyl dimethyl benzyl ammonium saccharinate (95% active). (BTC® and ONYXIDE® are presently commercially available from Stepan Company, Northfield, Illinois.) Polymeric quaternary

20 ammonium salts based on these monomeric structures are also considered desirable for the present invention. One example is POLYQUAT®, described as being a 2-butenyldimethyl ammonium chloride polymer.

When present, the quaternary ammonium compound(s) may be present in any effective amount, but generally need not be present in amounts in excess of about 10%wt. based on the

25 total weight of the composition. Preferably the germicidal quaternary ammonium compounds may be present in the inventive compositions in amounts of from about 0.001 %wt. to up to about 10%wt., very preferably about 0.01-8%wt., more preferably in amounts of between about 0.01-2%wt., and most preferably from about 0.01 - 1%wt.. It is particularly advantageous that the preferred germicidal cationic surfactant(s) are present in amounts of at least about 200 parts

30 per million (ppm), preferably in amounts of from about 1 ppm to 10,000 ppm, preferably from about 50 ppm to 2000 ppm, more preferably in amounts of from about 100 ppm to 1,000 ppm.

Particularly preferred amounts of one or more quaternary ammonium compound(s) and preferred amounts are identified with reference to the examples.

While not wishing to be bound by the following, the present inventors have surprisingly found that by careful selection of both: (1) the nature and amounts of the copper source material which releases copper ions into the treatment composition, and especially preferably wherein the copper source material is a source of Cu(I) and/or Cu(II) ions, and (2) the at least one lower alkyl aliphatic monohydric alcohol which preferably also exhibits an independent microbicidal effect, and wherein the composition is at a pH in excess of 5, preferably in excess of 8, therein is provided what appears to be a synergistic increase in the activity of the at least one lower alkyl aliphatic monohydric alcohol, especially preferably when these constituents are concurrently present with one or more further surfactant compounds e.g., one or more deterative surfactant compounds, especially where one or more nonionic surfactant compounds are present in addition to at least one quaternary ammonium compound. The resultant compositions provide unexpectedly superior microbicidal efficacy against a range of undesirable microorganisms including certain non-enveloped viruses, mycobacteria, bacteria and certain fungi, which has heretofore not been expected from compositions which have the reduced amounts of the alcohol constituent as provided in the inventive compositions. Such an effect has been observed even when a very limited amount of the copper source material is present, and wherein the amount of the one lower alkyl aliphatic monohydric alcohol is also present in reduced amounts, e.g, in an amount of between 20%wt. to about 50%wt, or even lesser amounts of up to but excluding 40%wt. Reference is made to the various Examples provided in this specification which demonstrates this effect, particularly as against comparative formulations which omit one or more of the copper source material, the at least one lower alkyl aliphatic monohydric alcohol, the at least one quaternary ammonium compound or which exhibits a pH level outside a preferred range. Although the literature describe the biochemical mechanisms of the separate microbicidal actives (the first, the second and the third essential constituents) when these microbicidal actives are used singly, inanimate surface treatment compositions as now disclosed by the inventors are believed to be unknown, particularly wherein such treatment compositions exhibit what is believed to be a synergistic benefit.

As the inventive compositions are, in part, aqueous, water is added in order to provide to 100% by weight of the compositions of the invention and is thus a further essential constituent.

The water may be tap water, but is preferably distilled and is most preferably deionized water. If the water is tap water, it is preferably substantially free of any undesirable impurities such as organics or inorganics, especially minerals salts which are present in hard water and which may undesirably interfere with the operation of the constituents present in the aqueous compositions according to the invention. In preferred embodiments the total amount of water and the one lower alkyl aliphatic monohydric alcohol(s) present comprise at least 80%wt., yet more preferably and in order of increasing preference comprise at least 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, and 99% by weight of the compositions of which they form a part.

In certain preferred embodiments the treatment compositions necessarily include at least at least one further deterative surfactant, (which is preferably a nonionic surfactant) other than a germicidally effective quaternary ammonium compound, which least one further deterative surfactant provides a further microbicidal benefit within the treatment composition of which it forms a part, as compared to where such at least one such further deterative surfactant is absent from said composition.

In certain embodiments the treatment compositions necessarily include at least one further, deterative surfactant, although such may be considered an optional constituent according to other embodiments of the invention.

Non-limiting examples of the major surfactant types that can be used as deterative surfactants of the present invention include those which are known as anionic, nonionic, amphoteric, and zwitterionic surfactants as well as further cationic surfactants which are not primarily present to provide a microbicidal or germicidal benefit. Such include, e.g.: sulfates and sulfonates of oils and fatty acids, sulfates and sulfonates, ethoxylated alkylphenols, sulfates of alcohols, sulfates of ethoxylated alcohols, sulfates of fatty esters, sulfonates of benzene, cumene, toluene and xylene, sulfonates of condensed naphthalenes, sulfonates of dodecyl and tridecylbenzenes, sulfonates of naphthalene and alkyl naphthalene, sulfonates of petroleum, sulfosuccinamates, sulfosuccinates and derivatives, soaps, taurates, thio and mercapto derivatives, tridecyl and dodecyl benzene sulfonic acids, alkanolamides, alkanolamines, alkylaryl sulfonates, alkylaryl sulfonic acids, alkylbenzenes, amine acetates, amine oxides, amines, sulfonated amines and amides, betaine derivatives, block polymers, carboxylated alcohol or alkylphenol ethoxylates, carboxylic acids and fatty acids, ethoxylated alcohols, ethoxylated

alkylphenols, ethoxylated amines and/or amides, ethoxylated fatty acids, ethoxylated fatty esters and oils, fatty esters, fluorocarbon-based surfactants, glycerol esters, glycol esters, heterocyclic-type products, imidazolines and imidazoline derivatives, isethionates, lanolin-based derivatives, lecithin and lecithin derivatives, lignin and lignin derivatives, maleic or succinic anhydrides, methyl esters, monoglycerides and derivatives, olefin sulfonates, phosphate esters, phosphorous organic derivatives, polyethylene glycols, polymeric (polysaccharides, acrylic acid, and acrylamide) surfactants, propoxylated and ethoxylated fatty acid alcohols or alkyl phenols, protein-based surfactants, sarcosine derivatives, silicone-based surfactants, sorbitan derivatives, sucrose and glucose esters and derivatives, as well as further surfactants known to the art but not elucidated here.

Additional non-limiting examples of surfactants that can be used to carry out the present invention include one or more nonionic surfactants, especially one or more compounds based on the condensation products of alkylene oxide groups with an organic hydrophobic compound, such as an aliphatic compound or with an alkyl aromatic compound. The nonionic synthetic organic detergents generally are the condensation products of an organic aliphatic or alkyl aromatic hydrophobic compound and hydrophilic ethylene oxide groups. Practically any hydrophobic compound having a carboxy, hydroxy, amido, or amino group with a free hydrogen attached to the nitrogen can be condensed with ethylene oxide or with the polyhydration product thereof, polyethylene glycol, to form a water soluble nonionic detergent. Further, the length of the polyethenoxy hydrophobic and hydrophilic elements may be varied to adjust these properties. Illustrative examples of such a nonionic surfactant include the condensation product of one mole of an alkyl phenol having an alkyl group containing from 6 to 12 carbon atoms with from about 5 to 25 moles of an alkylene oxide. Another example of such a nonionic surfactant is the condensation product of one mole of an aliphatic alcohol which may be a primary, secondary or tertiary alcohol having from 6 to 18 carbon atoms with from 1 to about 10 moles of alkylene oxide. Preferred alkylene oxides are ethylene oxides or propylene oxides which may be present singly, or may be both present.

Non-limiting, illustrative examples of nonionic surfactants include primary and secondary linear and branched alcohol ethoxylates, such as those based on C₆-C₁₈ alcohols which further include an average of from 2 to 80 moles of ethoxylation per mol of alcohol. Examples include the Genapol® series of linear alcohol ethoxylates from Clariant Corp., Charlotte, NC.


The 26-L series is based on the formula $RO(CH_2CH_2O)_nH$ wherein R is a mixture of linear, even carbon-number hydrocarbon chains ranging from $C_{12}H_{25}$ to $C_{16}H_{33}$ and n represents the number of repeating units and is a number of from 1 to about 12, such as 26-L-1, 26-L-1.6, 26-L-2, 26-L-3, 26-L-5, 26-L-45, 26-L-50, 26-L-60, 26-L-60N, 26-L-75, 26-L-80, 26-L-98N, and the 24-L series, derived from synthetic sources and typically contain about 55% C_{12} and 45% C_{14} alcohols, such as 24-L-3, 24-L-45, 24-L-50, 24-L-60, 24-L-60N, 24-L-75, 24-L-92, and 24-L-98N. From product literature, the single number following the "L" corresponds to the average degree of ethoxylation (numbers between 1 and 5) and the two digit number following the letter "L" corresponds to the cloud point in °C of a 1.0 wt.% solution in water.

Further examples of useful nonionic surfactants include secondary C_{12} - C_{15} alcohol ethoxylates, including those which have from about 3 to about 10 moles of ethoxylation. Such are available in the Tergitol® series of nonionic surfactants (Dow Chemical, Midland, MI), particularly those in the Tergitol® "15-S-" series. Further exemplary nonionic surfactants include linear primary C_{11} - C_{15} alcohol ethoxylates, including those which have from about 3 to about 10 moles of ethoxylation. Such are available in the Tomadol® series of nonionic surfactants under the following tradenames: Tomadol 1-3 (linear C_{11} alcohol with 3 moles (average) of ethylene oxide); Tomadol 1-5 (linear C_{11} alcohol with 5 moles (average) of ethylene oxide); Tomadol 1-7 (linear C_{11} alcohol with 7 moles (average) of ethylene oxide); Tomadol 1-9 (linear C_{11} alcohol with 9 moles (average) of ethylene oxide); Tomadol 23-1 (linear C_{12-13} alcohol with 1 mole (average) of ethylene oxide); Tomadol 23-3 (linear C_{12-13} alcohol with 3 moles (average) of ethylene oxide); Tomadol 23-5 (linear C_{12-13} alcohol with 5 moles (average) of ethylene oxide); Tomadol 23-6.5 (linear C_{12-13} alcohol with 6.6 moles (average) of ethylene oxide); Tomadol 25-12 (linear C_{12-15} alcohol with 11.9 moles (average) of ethylene oxide); Tomadol 25-3 (linear C_{12-15} alcohol with 2.8 moles (average) of ethylene oxide); Tomadol 25-7 (linear C_{12-15} alcohol with 7.3 moles (average) of ethylene oxide); Tomadol 25-9 (linear C_{12-15} alcohol with 8.9 moles (average) of ethylene oxide); Tomadol 45-13 (linear C_{14-15} alcohol with 12.9 moles (average) of ethylene oxide); Tomadol 45-2.25 (linear C_{14-15} alcohol with 2.23 moles (average) of ethylene oxide); Tomadol 45-7 (linear C_{14-15} alcohol with 7 moles (average) of ethylene oxide); Tomadol 91-2.5 (linear C_{9-11} alcohol with 2.7 moles (average) of ethylene oxide); Tomadol 91-6 (linear C_{9-11} alcohol with 6 moles (average) of ethylene oxide); Tomadol


91-8 (linear C₉₋₁₁ alcohol with 8.3 moles (average) of ethylene oxide) (Tomah Products, Inc., Milton, WI).

Further examples of useful nonionic surfactants include C₆-C₁₅ straight chain alcohols ethoxylated with about 1 to 13 moles of ethylene oxide, particularly those which include about 3 to about 6 moles of ethylene oxide. Examples of such nonionic surfactants include Alfonic® 810-4.5, which is described as having an average molecular weight of 356, an ethylene oxide content of about 4.85 moles and an HLB of about 12; Alfonic® 810-2, which is described as having an average molecular weight of 242, an ethylene oxide content of about 2.1 moles and an HLB of about 12; and Alfonic® 610-3.5, which is described as having an average molecular weight of 276, an ethylene oxide content of about 3.1 moles, and an HLB of 10.

A further class of nonionic surfactants which may find use in the present inventive compositions include ethoxylated octyl and nonyl phenols include those having one of the following general structural formulas:



15 or,



in which the C₉H₁₉ group in the latter formula is a mixture of branched chained isomers, and x indicates an average number of ethoxy units in the side chain. Particularly suitable non-ionic ethoxylated octyl and nonyl phenols include those having from about 7 to about 13 ethoxy groups. Such compounds are commercially available under the trade name Triton® X (Dow Chemical, Midland, MI), as well as under the tradename Igepal® (Rhodia, Princeton, NJ). One exemplary and particularly preferred nonylphenol ethoxylate is Igepal® CO-630.

Still further examples of suitable nonionic surfactants which may be advantageously included in the inventive compositions are alkoxy block copolymers, and in particular, compounds based on ethoxy/propoxy block copolymers. Polymeric alkylene oxide block copolymers include nonionic surfactants in which the major portion of the molecule is made up

of block polymeric C2-C4 alkylene oxides. Such nonionic surfactants, while preferably built up from an alkylene oxide chain starting group, can have as a starting nucleus almost any active hydrogen containing group including, without limitation, amides, phenols, thiols and secondary alcohols.

5 One group of such useful nonionic surfactants containing the characteristic alkylene oxide blocks are those which may be generally represented by the formula (A):



where EO represents ethylene oxide,

PO represents propylene oxide,

10 y equals at least 15,

(EO)_{x+z} equals 20 to 50% of the total weight of said compounds, and,

the total molecular weight is preferably in the range of about 2000 to 15,000.


Another group of nonionic surfactants for use in the new inventive compositions can be represented by the formula (B):



wherein R is an alkyl, aryl or aralkyl group, where the R group contains 1 to 20 carbon atoms, the weight percent of EO is within the range of 0 to 45% in one of the blocks a, b, and within the range of 60 to 100% in the other of the blocks a, b, and the total number of moles of combined EO and PO is in the range of 6 to 125 moles, with 1 to 50 moles in the PO rich block and 5 to
20 100 moles in the EO rich block.

Further nonionic surfactants which in general are encompassed by formula (B) include butoxy derivatives of propylene oxide/ethylene oxide block polymers having molecular weights within the range of about 2000-5000.

25 Still further useful nonionic surfactants containing polymeric butoxy (BO) groups can be represented by formula (C) as follows:



wherein R is an alkyl group containing 1 to 20 carbon atoms,

n is about 5-15 and x is about 5-15.

30 Also useful as the nonionic block copolymer surfactants, which also include polymeric butoxy groups, are those which may be represented by the following formula (D):




wherein n is about 5-15, preferably about 15,

x is about 5-15, preferably about 15, and

y is about 5-15, preferably about 15.

5 Still further useful nonionic surfactants include ethoxylated derivatives of propoxylated ethylene diamine, which may be represented by the following formula:



where (EO) represents ethoxy,

(PO) represents propoxy,

10 the amount of $(\text{PO})_x$ is such as to provide a molecular weight prior to ethoxylation of about 300 to 7500, and the amount of $(\text{EO})_y$ is such as to provide about 20% to 90% of the total weight of said compound.

Further examples of useful nonionic surfactants are one or more amine oxides.

Exemplary amine oxides include:

15 A) Alkyl di (lower alkyl) amine oxides in which the alkyl group has about 10-20, and preferably 12-16 carbon atoms, and can be straight or branched chain, saturated or unsaturated.


The lower alkyl groups include between 1 and 7 carbon atoms. Examples include lauryl dimethyl amine oxide, myristyl dimethyl amine oxide, and those in which the alkyl group is a mixture of different amine oxide, dimethyl cocoamine oxide, dimethyl (hydrogenated tallow) amine oxide, and myristyl/palmityl dimethyl amine oxide;

20 B) Alkyl di (hydroxy lower alkyl) amine oxides in which the alkyl group has about 10-20, and preferably 12-16 carbon atoms, and can be straight or branched chain, saturated or unsaturated. Examples are bis(2-hydroxyethyl) cocoamine oxide, bis(2-hydroxyethyl) tallowamine oxide; and bis(2-hydroxyethyl) stearylamine oxide;

25 C) Alkylamidopropyl di(lower alkyl) amine oxides in which the alkyl group has about 10-20, and preferably 12-16 carbon atoms, and can be straight or branched chain, saturated or unsaturated. Examples are cocoamidopropyl dimethyl amine oxide and tallowamidopropyl dimethyl amine oxide; and

D) Alkylmorpholine oxides in which the alkyl group has about 10-20, and preferably 12-16 carbon atoms, and can be straight or branched chain, saturated or unsaturated.

Preferably the amine oxide constituent is an alkyl di (lower alkyl) amine oxide as denoted above and which may be represented by the following structure:



5

wherein each:

R_1 is a straight chained C_1 - C_4 alkyl group, preferably both R_1 are methyl groups; and,

R_2 is a straight chained C_8 - C_{18} alkyl group, preferably is C_{10} - C_{14} alkyl group, most preferably is a C_{12} alkyl group.

10

Each of the alkyl groups may be linear or branched, but most preferably are linear.


Technical grade mixtures of two or more amine oxides may be used, wherein amine oxides of varying chains of the R_2 group are present. Preferably, the amine oxides used in the present invention include R_2 groups which comprise at least 50%wt., preferably at least 60%wt. of C_{12} alkyl groups and at least 25%wt. of C_{14} alkyl groups, with not more than 15%wt. of C_{16} , C_{18} or

15

higher alkyl groups as the R_2 group.

Further specific examples of useful nonionic surfactants are alkanolamide surfactant compounds. Exemplary useful alkanolamides include one or more monoethanol amides, and diethanol amides of fatty acids having an acyl moiety which contains from about 8 to about 18 carbon atoms, and which may be represented in accordance with the formula:

20




where R_1 represents a saturated or unsaturated aliphatic hydrocarbon radical of from about 7 to 21 carbon atoms, but preferably from about 11 to 17 carbon atoms; R_2 represents a $-\text{CH}_2-$ or $-\text{CH}_2\text{CH}_2-$, and m is an integer from 1 to 3, but is preferably 1. Preferably, R_1 is a saturated or unsaturated aliphatic hydrocarbon radical comprising from about 11 to 17 carbon atoms, and m is 1. Specific examples of such compounds include mono-ethanol amine coconut fatty acid amide and diethanol amine dodecyl fatty acid amide. An exemplary useful and particularly preferred fatty acid amides include cocomonethanol amide or cocodiethanolamide, which are

25

presently commercially available under the Monamid® tradename. Further exemplary useful alkanolamides which provide such functions include *inter alia*: cocamide MEA, cocamide DEA, soyamide DEA, lauramide DEA, oleamide MIPA, stearamide MEA, myristamide MEA, lauramide MEA, capramide DEA, ricinoleamide DEA, myristamide DEA, stearamide DEA, 5 oleylamide DEA, tallowamide DEA, lauramide MIPA, tallowamide MEA, isostearamide DEA, isostearamide MEA, and mixtures thereof. Further useful alkanolamide surfactant compounds include alkanolamides, particularly fatty monoalkanolamides and fatty dialkanolamides, including one or more of those marketed under the Ninol® tradename. Further exemplary alkanolamide surfactant compounds include monoethanol amides and diethanol amides include 10 those marketed under the trade names Alakamide® and Cyclomide® by Rhône-Poulenc Co., (Cranbury, NJ) e.g., Cyclomide® CDD-518 described to be a nonionic surfactant based on coconut diethanolamide; Cyclomide® C212 described to be a nonionic surfactant based on coconut monoethanolamide; Cyclomide® DC212/SE described to be a nonionic surfactant based on 1:1 fatty acid diethanolamide; Cyclomide® DIN 100 described to be a nonionic surfactant 15 based on lauric/linoleic diethanolamide; Cyclomide® DIN-295/S described to be a nonionic surfactant based on 1:1 linoleic diethanolamide; Cyclomide® DL203 described to be a nonionic surfactant based on 2:1 lauric diethanolamide.

Further specific examples of useful nonionic surfactants include alkyl polyglycosides. The alkyl polyglycosides which can be used as nonionic surfactants in the composition are 20 generally represented by the formula:



wherein R₁ is a monovalent organic radical having from about 6 to about 30 carbon atoms; R₂ is a divalent alkylene radical having from 2 to 4 carbon atoms; Z is a saccharide residue having 5 or 6 carbon atoms; b is a number having a value from 0 to about 12; a is a number having a value 25 from 1 to about 6. Preferred alkyl polyglycosides have the formula I wherein Z is a glucose residue and b is zero. Such alkyl polyglycosides are commercially available, for example, as APG®, GLUCOPON®, or PLANTAREN® surfactants from Cogis Corp. Specific examples of such surfactants include but are not limited to: APG® 225, described to be an alkyl polyglycoside in which the alkyl group contains 8 to 10 carbon atoms and having an average 30 degree of polymerization of 1.7; GLUCOPON® 425, described to be an alkyl polyglycoside in which the alkyl group contains 8 to 16 carbon atoms and having an average degree of

polymerization of 1.48.; GLUCOPON® 625, described to be an alkyl polyglycoside in which the alkyl group contains 12 to 16 carbon atoms and having an average degree of polymerization of 1.6; APG® 325, described to be an alkyl polyglycoside in which the alkyl group contains 9 to 11 carbon atoms and having an average degree of polymerization of 1.5; GLUCOPON® 600, 5 described to be an alkyl polyglycoside in which the alkyl group contains 12 to 16 carbon atoms and having an average degree of polymerization of 1.4; PLANTAREN® 2000, described to be an alkyl polyglycoside in which the alkyl group contains 8 to 16 carbon atoms and having an average degree of polymerization of 1.4; and, PLANTAREN® 1300, described to be an alkyl polyglycoside in which the alkyl group contains 12 to 16 carbon atoms and having an average 10 degree of polymerization of 1.6. Other examples include alkyl polyglycoside surfactant compositions which are comprised of mixtures of compounds of formula I wherein Z represents a moiety derived from a reducing saccharide containing 5 or 6 carbon atoms; a is a number having a value from 1 to about 6; b is zero; and R.sub.1 is an alkyl radical having from 8 to 20 carbon atoms. The compositions are characterized in that they have increased surfactant 15 properties and an HLB in the range of about 10 to about 16 and a non-Flory distribution of glycosides, which is comprised of a mixture of an alkyl monoglycoside and a mixture of alkyl polyglycosides having varying degrees of polymerization of 2 and higher in progressively decreasing amounts, in which the amount by weight of polyglycoside having a degree of polymerization of 2, or mixtures thereof with the polyglycoside having a degree of 20 polymerization of 3, predominate in relation to the amount of monoglycoside, said composition having an average degree of polymerization of about 1.8 to about 3. Such compositions, also known as peaked alkyl polyglycosides, can be prepared by separation of the monoglycoside from the original reaction mixture of alkyl monoglycoside and alkyl polyglycosides after removal of the alcohol. This separation may be carried out by molecular distillation and normally results in 25 the removal of about 70-95% by weight of the alkyl monoglycosides. After removal of the alkyl monoglycosides, the relative distribution of the various components, mono- and poly-glycosides, in the resulting product changes and the concentration in the product of the polyglycosides relative to the monoglycoside increases as well as the concentration of individual polyglycosides to the total, i.e. DP2 and DP3 fractions in relation to the sum of all DP fractions. Such 30 compositions are disclosed in U.S. Pat. No. 5,266,690, the entire contents of which are incorporated herein by reference.

Other alkyl polyglycosides which can be used in the compositions according to the invention are those in which the alkyl moiety contains from 6 to 18 carbon atoms in which and the average carbon chain length of the composition is from about 9 to about 14 comprising a mixture of two or more of at least binary components of alkylpolyglycosides, wherein each
5 binary component is present in the mixture in relation to its average carbon chain length in an amount effective to provide the surfactant composition with the average carbon chain length of about 9 to about 14 and wherein at least one, or both binary components, comprise a Flory distribution of polyglycosides derived from an acid-catalyzed reaction of an alcohol containing 6-20 carbon atoms and a suitable saccharide from which excess alcohol has been separated.

10 Also useful as nonionic surfactants are ethylene oxides condensed with sorbitan fatty acid esters. Such materials are presently commercially available under the tradename TWEEN (ex. ICI) and/or CRILL (ex. Croda) which include polyoxyethylene sorbitan monolaurate, polyoxyethylene sorbitan monopalmitate, polyoxyethylene sorbitan monostearate, polyoxyethylene sorbitan tristearate, polyoxyethylene sorbitan monooleate, polyoxyethylene
15 sorbitan trioleates which are available in a variety of grades, and with differing amounts of polyoxyethylene groups per molecule.

The inventive compositions most desirably, although not always essentially, include at least one nonionic surfactant. An example of an especially preferred nonionic surfactant is at least one alcohol ethoxylate based nonionic surfactant in an amount of from about 0.01 – 10%wt.
20 In order of increasing preference, when present, the at least one nonionic surfactant comprises in %wt. at least 0.025, 0.05, 0.075, 0.1, 0.2, 0.25, 0.3, 0.4, 0.5, 0.6, 0.75, 0.8, 0.9, 1, 1.1, 1.2, 1.3, 1.4 and 1.5%wt. and similarly in order of increasing preference the at least one nonionic surfactant comprises, in %wt., not more than 10, 9, 8, 7.5, 7, 6, 5, 4.75, 4.5, 4, 3.75, 3.5, 3.25, 3, 2.75 and 2%wt. based on the total weight of a treatment composition of which they form a part.

25 Especially preferred nonionic surfactants and the amounts in which they are preferably present are disclosed with reference to one or more of the Examples. In certain embodiments at least one nonionic surfactant is necessarily present and is considered as a further essential constituent of the invention.

Non-limiting examples of further surfactants which may be included in the treatment
30 compositions of the invention include zwitterionic and amphoteric surfactants. Zwitterionic surfactants may also be present either by themselves or in admixture with another ionic


surfactant providing there are no troublesome interactions. Typical examples of amphoteric or zwitterionic surfactants are alkyl betaines, alkyl amidobetaines, aminopropionates, aminoglycinates, imidazolium betaines and sulfobetaines. Within this group, alkyl betaines and alkyl amidobetaines are particularly preferred. Alkyl betaines are known surfactants which are
5 mainly produced by carboxyalkylation, preferably carboxymethylation of aminic compounds. Typical examples are the carboxymethylation products of hexyl methyl amine, hexyl dimethyl amine, octyl dimethyl amine, decyl dimethyl amine, dodecyl methyl amine, dodecyl dimethyl amine, dodecyl ethyl methyl amine, C12/14 cocoalkyl dimethyl amine, myristyl dimethyl amine, cetyl dimethyl amine, stearyl dimethyl amine, stearyl ethyl methyl amine, oleyl dimethyl amine,
10 C16/18 tallow alkyl dimethyl amine and technical mixtures thereof.

Alkyl amidobetaines which represent carboxyalkylation products of amidoamines are also suitable. Typical examples are reaction products of fatty acids containing 6 to 22 carbon atoms, namely caproic acid, caprylic acid, capric acid, lauric acid, myristic acid, palmitic acid, palmitoleic acid, stearic acid, isostearic acid, oleic acid, elaidic acid, petroselic acid, linoleic
15 acid, linolenic acid, elaeostearic acid, arachic acid, gadoleic acid, behenic acid and erucic acid and technical mixtures thereof, with N,N-dimethylaminoethyl amine, N,N-dimethylaminopropyl amine, N,N-diethylaminoethyl amine and N,N-diethylaminopropyl amine which are condensed with sodium chloroacetate. The condensation product of C8/18 cocofatty acid-N,N-dimethylaminopropyl amide with sodium chloroacetate is preferably used.

Further specific examples of particular amphoteric surfactants which may be used in the treatment compositions of the invention include one or more amphoteric surfactants. Exemplary amphoteric surfactants include alkylampho(mono)acetates, alkylampho(di)acetates, alkylampho(mono)propionates, and alkylampho(di)propionates. Examples of these amphoteric surfactants can be found under the tradename Miranol from Rhodia (Cranbury, NJ). Some
25 examples include Miranol C2M-Conc. NP, described to be disodium cocoamphodiacetate; Miranol FA-NP, described to be sodium cocoamphodiacetate; Miranol DM, described to be sodium steroamphodiacetate; Miranol HMA, described to be sodium lauroamphodiacetate; Miranol C2M, described to be cocoamphodipropionic acid; Miranol C2M-SF, described to be disodium cocoamphodipropionate; Miranol CM-SF Conc., described as being cocoamphopropionate;
30 Mirataine H2C-HA, described as sodium lauriminodipropionate; Miranol Ultra L-32, described as sodium lauroamphodiacetate; and Miranol Ultra C-37, described as sodium cocoamphodiacetate.

Other amphoteric surfactants are also available under the tradename Amphoterger from Lonza (Fair Lawn, NJ) such as Amphoterger K described to sodium cocoamphopropionate; Amphoterger K-2, described as disodium cocoamphodipropionate; Amphoterger W, described to be sodium cocoamphoacetate; and Amphoterger W-2, described to be disodium
 5 cocoamphodiacetate.

Further useful amphoteric surfactants include those which may be represented by the following general formula



in which, R represents a C₄ to C₂₄ alkyl group, and is preferably a C₁₀ to C₁₆ alkyl group, R₁
 10 and R₂ independently represent a C₁ to C₈ alkyl group, is preferably -CH₂CH₂- or -CH₂CH₂CH₂-, and M may be any salt-forming anion which permits water solubility or water miscibility of the compound, e.g., chloride, bromide, methosulfate, ethosulfate, lactate, saccharinate, acetate or phosphate. Such compounds are presently commercially available, such
 15 as those marketed in the Tomamine Amphoteric series of amphoteric surfactants, ex. Air Products Inc.

While the one or more deterative surfactants may be present in any effective amount which may be observed to improve the microbicidal efficacy of the system of the essential constituents, these one or more surfactants, when present, are advantageously present in an amount of from about 0.001 – 15%wt., preferably from about 0.01 – 10%wt. and particularly
 20 preferably from about 0.05 – 5%wt., based on the total weight of the treatment composition within which they are present.

In certain preferred embodiments, at least one deterative surfactant, preferably at least one nonionic surfactant, is a necessary constituent of the inventive compositions. While not wishing to be bound by the following, it is suspected that the presence of at least one surfactant and
 25 especially at least one nonionic surfactant aids in the penetration of organic soils and/or the penetration of one or more undesired microorganisms and hastens the activity of the essential constituents, viz. the copper ions provided by the source of copper ions, and/or the lower alkyl aliphatic monohydric alcohol in reducing, deactivating or destroying these undesired

microorganisms and thus may aid in both providing an improved speed and/or degree of control, reduction or elimination of the one or more undesired microorganisms being treated with the treatment compositions taught herein.

In certain embodiments, the sole surfactants present in the compositions are one or more of the cationic surfactants described above, and/or one or more of the nonionic surfactants described above. In certain embodiments the sole surfactant(s) present is one or more cationic surfactants. In certain embodiments the sole surfactant(s) present is one or more nonionic surfactants.

The pH of the treatment compositions is preferably established and thereafter maintained at a desired pH or within a bounded pH range. As is better understood from a consideration of the example compositions, the inventors have also found that the pH of the treatment compositions plays a significant role in establishing the overall efficacy of a treatment composition in reducing, deactivating or destroying undesired microorganisms. It was generally observed that compositions having a higher, more alkaline pH but concurrently including lesser amounts of alcohol (specifically ethanol) provided similar microbicidal performance to other compositions having a lower pH but which included increased amounts of ethanol. Thus, a reasonable degree of flexibility in formulating compositions of the invention is provided by judicious control of the pH and the amount of the lower alkyl monohydric alcohol present. Specific reference is made to the example formulations described hereinafter which demonstrate this effect. The pH of the inventive compositions is at least 5, but is preferably greater and in certain particularly preferred embodiments is substantially alkaline. While the pH of the composition may be 5 or greater, preferably the pH of the compositions is at least about 6, and more preferably is in the range of from about 7 – 14, especially in the range of about 9 – 12. Thus in preferred embodiments the pH of the treatment compositions (and/or microbicidal control system) is at least 5, and in order of increasing preference is at least 6, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, 7, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9, 8, 8.1, 8.2, 8.3, 8.4, 8.5, 8.6, 8.7, 8.8, 8.9, 9, 9.1, 9.2, 9.3, 9.4, 9.5, 9.6, 9.7, 9.8, 9.9, 10, 10.1, 10.2, 10.3, 10.4, 10.5, 10.6, 10.7, 10.8, 10.9, 11, 11.1, 11.2, 11.3, 11.4, 11.5, 11.6, 11.7, 11.8, 11.9, 12, 12.1, 12.2, 12.3, 12.4, 12.5. In preferred embodiments, and in order of increasing preference the pH of the treatment compositions (and/or microbicidal control system) is not in excess of: 12.5, 12.4, 12.3, 12.2, 12.1, 12, 11.9, 11.8, 11.7, 11.6, 11.5, 11.4, 11.3, 11.2, 11.1, 11, 10.9, 10.8, 10.7, 10.6, 10.5, 10.4,

10.3, 10.2, 10.1, 10, 9.9, 9.8, 9.7, 9.6, 9.5. It is expected that compositions of the invention may have lower pHs, in the range of 1 – 14 if desired; however preferred pHs are indicated in the foregoing ranges and are demonstrated by the Examples. The pH of the surface treatment compositions may be established, adjusted and/or maintained by the addition of an effective amount of a pH adjustment constituent.

Optionally but preferably the treatment compositions of the invention include a pH adjusting constituent which may be used to establish and/or maintain, viz., buffer, a treatment composition at a desired pH or within a bounded pH range. Essentially any material which may increase or decrease the pH of the treatment composition is suitable as a pH adjusting constituent. Suitable pH adjusting constituents are one or more acids and/or bases whether such be based on organic and/or inorganic compounds or materials. By way of non-limiting example, pH adjusting agents include phosphorus containing compounds, monovalent and polyvalent salts such as of silicates, carbonates, and borates, certain acids and bases, tartrates and certain acetates. Further exemplary pH adjusting agents include mineral acids, basic compositions, and organic acids, which are typically required in only minor amounts. By way of further non-limiting example, pH buffering compositions include the alkali metal phosphates, polyphosphates, pyrophosphates, triphosphates, tetraphosphates, silicates, metasilicates, polysilicates, carbonates, hydroxides, and mixtures of the same. Certain salts, such as the alkaline earth phosphates, carbonates, and hydroxides, can also function as buffers. It may also be suitable to use as buffers such materials as aluminosilicates (zeolites), borates, aluminates and certain organic materials such as gluconates, succinates, maleates, citrates, and their alkali metal salts. Particularly useful and preferred is citric acid and metal salts thereof such as sodium citrate which are widely available and which are effective in providing these pH adjustment and buffering effects. Further exemplary and useful pH adjusting constituents include monoalkanolamines, dialkanolamines, trialkanolamines, and alkylalkanolamines such as alkyl-dialkanolamines, and dialkyl-monoalkanolamines. Such may also function as detergent surfactants. The alkanol and alkyl groups are generally short to medium chain length, that is, from 1 to 7 carbons in length. For di- and trialkanolamines and dialkyl-monoalkanolamines, these groups can be combined on the same amine to produce for example, methylethylhydroxypropylhydroxylamine. One of ordinary skill in the art can readily ascertain other members of this group. Preferred alkanolamines include monoethanolamine.

When present, the one or more pH adjusting constituents are included in amounts which are effective in establishing and/or maintaining the pH of a treatment composition at the desired pH value or within a range of pH values. Advantageously the one or more pH adjusting constituents comprise from about 0.001 – 2.5%wt., preferably from about 0.01 – 1.5%wt. of the treatment composition of which the one or more pH adjusting constituents form a part. Preferred pH adjusting constituents include those demonstrated in or more of the Examples. In certain preferred embodiments, one or more pH adjusting constituents are necessarily present and are to be understood as essential constituents of the treatment compositions.


The liquid inanimate surface treatment compositions of the invention may include one or more further optional constituents or materials which impart a desired technical and/or aesthetic benefit to the inventive compositions. However in certain embodiments one or more of such further optional constituents or materials may be expressly excluded from the treatment compositions. Non-limiting examples of such one or more further optional constituents are hereinafter described.

The treatment compositions of the invention may include one or more acids, which include not only organic and inorganic acids but also acid salts of organic acids. Preferred examples of the organic acid to be used in the present invention include linear aliphatic acids such as formic acid, acetic acid, propionic acid, butyric acid and valeric acid; dicarboxylic acids such as oxalic acid, malonic acid, succinic acid, glutaric acid, adipic acid, pimelic acid, fumaric acid and maleic acid; acidic amino acids such as glutamic acid and aspartic acid; and hydroxy acids such as glycolic acid, lactic acid, hydroxyacrylic acid, alpha-hydroxybutyric acid, glyceric acid, tartronic acid, malic acid, tartaric acid and citric acid, as well as acid salts of these organic acids. Exemplary inorganic acids include phosphoric acid, potassium dihydrogenphosphate, sodium dihydrogenphosphate, sodium sulfite, potassium sulfite, sodium pyrosulfite (sodium metabisulfite), potassium pyrosulfite (potassium metabisulfite), acid sodium hexametaphosphate, acid potassium hexametaphosphate, acid sodium pyrophosphate, acid potassium pyrophosphate and sulfamic acid. These acids can be used singly or as a mixture of two or more inorganic and/or organic acids. Such one or more acids may be used to adjust the pH of the inventive compositions, and/or buffer the pH of the treatment compositions. When present, these may be included in effective amounts.

The treatment compositions of the invention may also include one or more further compounds, constituents or materials which provide an ancillary microbicidal benefit or effect. These are distinguished from the essential constituents of the invention described above. When present, they may be included in amounts which are effective in order to provide an ancillary microbicidal benefit. Non-limiting examples of such materials include non-cationic microbicidal agents which are particularly useful in the present invention: pyrithiones (especially zinc pyrithione which is also known as ZPT), dimethyldimethylol hydantoin (Glydant), methylchloroisothiazolinone/methylisothiazolinone (Kathon CG), sodium sulfite, sodium bisulfite, imidazolidinyl urea (Germall 115), diazolidinyl urea (Germaill II), benzyl alcohol, 2-bromo-2-nitropropane-1,3-diol (Bronopol), formalin (formaldehyde), iodopropenyl butylcarbamate (Polyphase P100), chloroacetamide, methanamine, methyldibromonitrile glutaronitrile (1,2-Dibromo-2,4-dicyanobutane or Tektamer), glutaraldehyde, 5-bromo-5-nitro-1,3-dioxane (Bronidox), phenethyl alcohol, o-phenylphenol/sodium o-phenylphenol, sodium hydroxymethylglycinate (Suttocide A), polymethoxy bicyclic oxazolidine (Nuosept C), dimethoxane, thimersal dichlorobenzyl alcohol, captan, chlorphenenesin, dichlorophene, chlorbutanol, glyceryl laurate, halogenated diphenyl ethers like 2,4,4-trichloro-2-hydroxy-diphenyl ether (Triclosan or TCS), 2,2-dihydroxy-5,5-dibromo-diphenyl ether, phenolic compounds like phenol, 2-methyl phenol, 3-methyl phenol, 4-methyl phenol, 4-ethyl phenol, 2,4-dimethyl phenol, 2,5-dimethyl phenol, 3,4-dimethyl phenol, 2,6-dimethyl phenol, 4-n-propyl phenol, 4-n-butyl phenol, 4-n-amyl phenol, 4-tert-amyl phenol, 4-n-hexyl phenol, 4-n-heptyl phenol, mono- and poly-alkyl and aromatic halophenols such as p-chlorophenol, methyl p-chlorophenol, ethyl p-chlorophenol, n-propyl p-chlorophenol, n-butyl p-chlorophenol, n-amyl p-chlorophenol, sec-amyl p-chlorophenol, n-hexyl p-chlorophenol, cyclohexyl p-chlorophenol, n-heptyl p-chlorophenol, n-octyl p-chlorophenol, o-chlorophenol, methyl o-chlorophenol, ethyl o-chlorophenol, n-propyl o-chlorophenol, n-butyl o-chlorophenol, n-amyl o-chlorophenol, tert-amyl o-chlorophenol, n-hexyl o-chlorophenol, n-heptyl o-chlorophenol, o-benzyl p-chlorophenol, o-benzyl-m-methyl p-chlorophenol, o-benzyl-m, m-dimethyl p-chlorophenol, o-phenylethyl p-chlorophenol, o-phenylethyl-m-methyl p-chlorophenol, 3-methyl p-chlorophenol, 3,5-dimethyl p-chlorophenol, 6-ethyl-3-methyl p-chlorophenol, 6-n-propyl-3-methyl p-chlorophenol, 6-iso-propyl-3-methyl p-chlorophenol, 2-ethyl-3,5-dimethyl p-chlorophenol, 6-sec-butyl-3-methyl p-chlorophenol, 2-iso-propyl-3,5-dimethyl p-chlorophenol, 6-diethylmethyl-


3-methyl p-chlorophenol, 6-iso-propyl-2-ethyl-3-methyl p-chlorophenol, 2-sec-amyl-3,5-dimethyl p-chlorophenol 2-diethylmethyl-3,5-dimethyl p-chlorophenol, 6-sec-octyl-3-methyl p-chlorophenol, p-chloro-m-cresol, p-bromophenol, methyl p-bromophenol, ethyl p-bromophenol, n-propyl p-bromophenol, n-butyl p-bromophenol, n-amyl p-bromophenol, sec-amyl p-bromophenol, n-hexyl p-bromophenol, cyclohexyl p-bromophenol, o-bromophenol, tert-amyl o-bromophenol, n-hexyl o-bromophenol, n-propyl-m,m-dimethyl o-bromophenol, 2-phenyl phenol, 4-chloro-2-methyl phenol, 4-chloro-3-methyl phenol, 4-chloro-3,5-dimethyl phenol, 2,4-dichloro-3,5-dimethylphenol, 3,4,5,6-terabromo-2-methylphenol, 5-methyl-2-pentylphenol, 4-isopropyl-3-methylphenol, para-chloro-meta-xyleneol, dichloro meta xylenol, chlorothymol, 5-chloro-2-hydroxydiphenylmethane, resorcinol and its derivatives including methyl resorcinol, ethyl resorcinol, n-propyl resorcinol, n-butyl resorcinol, n-amyl resorcinol, n-hexyl resorcinol, n-heptyl resorcinol, n-octyl resorcinol, n-nonyl resorcinol, phenyl resorcinol, benzyl resorcinol, phenylethyl resorcinol, phenylpropyl resorcinol, p-chlorobenzyl resorcinol, 5-chloro 2,4-dihydroxydiphenyl methane, 4-chloro 2,4-dihydroxydiphenyl methane, 5-bromo 2,4-dihydroxydiphenyl methane, and 4-bromo 2,4-dihydroxydiphenyl methane, bisphenolic compounds like 2,2-methylene bis (4-chlorophenol), 2,2-methylene bis (3,4,6-trichlorophenol), 2,2-methylene bis (4-chloro-6-bromophenol), bis (2-hydroxy-3,5-dichlorophenyl) sulphide, and bis (2-hydroxy-5-chlorobenzyl)sulphide, benzoic esters (parabens) like methylparaben, propylparaben, butylparaben, ethylparaben, isopropylparaben, isobutylparaben, benzylparaben, sodium methylparaben, and sodium propylparaben, halogenated carbanilides (e.g., 3,4,4-trichlorocarbanilides (Triclocarban or TCC), 3-trifluoromethyl-4,4-dichlorocarbanilide, 3,3,4-trichlorocarbanilide, etc.).

Of these, preferred are phenol based non-cationic microbicides, especially those based on one or more phenolic compounds, particularly 2-hydroxydiphenyl compounds which may be exemplified by the following classes of compounds:




wherein Y is chlorine or bromine, Z is SO₂ H, NO₂, or C₁-C₄ alkyl, r is 0 to 3, o is 0 to 3, p is 0 or 1, m is 0 or 1, and n is 0 or 1. In preferred embodiments, Y is chlorine or bromine, m is 0, n is 0 or 1, o is 1 or 2, r is 1 or 2, and p is 0, and according to especially preferred embodiments, Y is chlorine, m is 0, n is 0, o is 1, r is 2, and p is 0.

5 Particularly useful 2-hydroxydiphenyl compounds include those which may be represented by the structure:



which is commonly referred to as "TRICLOSAN" and which is presently commercially available from Ciba Specialty Chemicals Corp., as well as halogenated carbanilides, e.g., TCC.


10 Further exemplary useful phenolic based disinfecting agents include 2,2'-hydroxy-5,5'-dibromo-diphenyl ether which may be represented by the structure:



wherein R₁ is hydro, hydroxy, C₁-C₄ alkyl, chloro, nitro, phenyl, or benzyl; R₂ is hydro, hydroxy, C₁-C₆ alkyl, or halo; R₃ is hydro, C₁-C₆ alkyl, hydroxy, chloro, nitro, or a sulfur in the form of an alkali metal salt or ammonium salt; R₄ is hydro or methyl, and R₅ is hydro or nitro. Halo is bromo or, preferably, chloro.

Specific examples of phenol derivatives include, but are not limited to, chlorophenols (o-, m-, p-), 2,4-dichlorophenol, p-nitrophenol, picric acid, xylenol, p-chloro-m-xyleneol, cresols (o-, m-, p-), p-chloro-m-cresol, pyrocatechol, resorcinol, 4-n-hexylresorcinol, pyrogallol, phloroglucin, carvacrol, thymol, p-chlorothymol, o-phenylphenol, o-benzylphenol, p-chloro-o-benzylphenol, phenol, 4-ethylphenol, and 4-phenolsulfonic acid.

Still further useful phenol derivatives include those which may be represented by the structure:



- 5 wherein X is sulfur or a methylene group, R₁ and R'₁ are hydroxy, and R₂, R'₂, R₃, R'₃, R₄, R'₄, R₅, and R'₅, independent of one another, are hydro or halo. Specific, nonlimiting examples of diphenyl compounds are hexachlorophene, tetrachlorophene, dichlorophene, 2,3-dihydroxy-5,5'-dichlorodiphenyl sulfide, 2,2'-dihydroxy-3,3',5,5'-tetrachlorodiphenyl sulfide, 2,2'-dihydroxy-3,5',5,5', 6,6'-hexachlorodiphenyl sulfide, and 3,3'-dibromo-5,5'-dichloro-2,2'-
- 10 dihydroxydiphenylamine. Of the foregoing, a particularly useful phenol derivative is commonly referred to as triclocarban, or 3,4,4'-trichlorocarbanilide as well as derivatives thereto. When present, one or more such further compounds, constituents or materials which provide an ancillary microbicidal benefit or effect may be present in effective amounts, e.g., in amounts of up to about 5%wt., although depending upon the efficacy of one or more selected such further
- 15 compounds, constituents or materials are usually effective in reduced amounts, e.g., 0.001 – 2%wt. of the treatment composition.

The treatment compositions of the invention may optionally include a fragrance constituent, which may be based on natural and/or synthetic fragrances and most commonly are mixtures or blends of a plurality of such fragrances, optionally in conjunction with a carrier such

20 as an organic solvent or a mixture of organic solvents in which the fragrances are dissolved, suspended or dispersed. Such may be natural fragrances, e.g, natural extracts of plants, fruits, roots, stems, leaves, wood extracts, e.g. terpineols, resins, balsams, animal raw materials, e.g., civet and beaver, as well as typical synthetic perfume compounds which are frequently products of the ester, ether, aldehyde, ketone, alcohol and hydrocarbon type, e.g., benzyl acetate, linalyl

25 acetate, citral, citronellal, methyl cedryl ketone, eugenol, isoeugenol, geraniol, linalool, and Typically it is preferred to use mixtures of different perfume compounds which, together, produce an agreeable fragrance. Other suitable perfume oils are essential oils of relatively low

volatility which are mostly used as aroma components. Examples are sage oil, camomile oil, clove oil, melissa oil, mint oil, cinnamon leaf oil, lime-blossom oil, juniper berry oil, vetiver oil, olibanum oil, galbanum oil, labolanum oil and lavandin oil. When present in a treatment composition, in accordance with certain of the preferred embodiments, the fragrance constituent may be present in any effective amount such that it can be discerned by a consumer of the composition, however such is advantageously present in amounts of up to about 1%wt., preferably in amounts of from about 0.00001%wt. to about 0.5%wt., and most preferably in an amount of from about 0.0001%wt. to 0.5%wt. based on the total weight of the treatment composition of which it forms a part.

10 A further optional constituent of the treatment compositions of the invention include colorant, such as dyes and pigments which may be used to impart a color to the compositions of which they form a part.

The treatment compositions of the invention may also optionally include a preservative constituent which is used to control undesired microorganisms within the treatment composition particularly when the treatment composition is in long-term storage and at elevated temperatures. While these are normally not present due to the microbicidal efficacy of the compositions as taught herein, such ancillary preservative constituents may be included in minor but effective amounts. Nonlimiting examples include one or more of parabens, including methyl parabens and ethyl parabens, glutaraldehyde, formaldehyde, 2-bromo-2-nitropropoane-1,3-diol, 5-chloro-2-methyl-4-isothiazolin-3-one, 2-methyl-4-isothiazoline-3-one, and mixtures thereof. One exemplary composition is a combination 5-chloro-2-methyl-4-isothiazolin-3-one and 2-methyl-4-isothiazolin-3-one where the amount of either component may be present in the mixture anywhere from 0.001 to 99.99 weight percent, based on the total amount of the preservative. Further exemplary useful preservatives include those which are commercially including a mixture of 5-chloro-2-methyl-4-isothiazolin-3-one and 2-methyl-4-isothiazolin-3-one marketed under the trademark KATHON® CG/ICP as a preservative composition presently commercially available from Rohm and Haas (Philadelphia, PA).

25 A further optional constituent in the inventive treatment compositions is one or more chelating agents. Exemplary useful chelating agents include those known to the art, including by way of non-limiting example; aminopolycarboxylic acids and salts thereof wherein the amino nitrogen has attached thereto two or more substituent groups. Preferred chelating agents include

acids and salts, especially the sodium and potassium salts of ethylenediaminetetraacetic acid, diethylenetriaminepentaacetic acid, N-hydroxyethylethylenediaminetriacetic acid, and of which the sodium salts of ethylenediaminetetraacetic acid may be particularly advantageously used. Such chelating agents may be omitted, or they may be included in generally minor amounts such as from about 0.001 - 0.5 %wt. based on the weight of the chelating agents and/or salt forms thereof. When present, advantageously, such chelating agents are included in the present inventive composition in amounts from about 0.001 - 5%wt., but are most desirably present in reduced weight percentages from about 0.01 - 0.5%wt.

The amounts or presence of chelating agents should be carefully controlled and may in some preferred embodiments be excluded from the treatment compositions. This is due the fact that the presence of chelating agents may undesirably form insoluble complexes with the Cu(I) and/or Cu(II) ions present in the compositions, and thus reduce or remove their beneficial effects in the treatment compositions. Desirably when such are present they are included in only limited amounts so as not to deleteriously affect the improved microbicidal efficacy of the inventive compositions, but in particularly preferred embodiments such chelating agents are desirably excluded from the compositions of the invention

The treatment compositions of the invention may include one or more further organic solvents, which are differentiated from the essential alkyl aliphatic monohydric alcohol constituent. Such further optional organic solvents may include one or more of: alcohols other than the essential lower alkyl aliphatic monohydric alcohol described previously, glycols, acetates, ether acetates, glycerols, as well as polyethylene glycols and glycol ethers. Mixtures of these further optional organic solvents can also be used. Typically such further one or more organic solvents are ones which have no appreciable microbicidal effect and are thus differentiated from the essential alkyl aliphatic monohydric alcohol constituent. Non-limiting examples of useful glycol ethers and examples include those glycol ethers having the general structure $R_a-O-[CH_2-CH(R)-(CH_2)-O]_n-H$, wherein R_a is C_{1-20} alkyl or alkenyl, or a cyclic alkane group of at least 6 carbon atoms, which may be fully or partially unsaturated or aromatic; n is an integer from 1 to 10, preferably from 1 to 5; each R is selected from H or CH_3 ; and a is the integer 0 or 1. Specific and preferred solvents are selected from propylene glycol methyl ether, dipropylene glycol methyl ether, tripropylene glycol methyl ether, propylene glycol n-propyl ether, ethylene glycol n-butyl ether, diethylene glycol n-butyl ether, diethylene glycol

methyl ether, propylene glycol, ethylene glycol, diethylene glycol monoethyl ether acetate and the like. When present such further optional one or more organic solvents may be present in any effective amount, preferably in amounts of between about 0.001 – 10%wt., and preferably between about 0.01 – 5%wt. based on the total weight of the treatment composition of which they form a part.

When one or more such further optional constituents are present in the treatment compositions, preferably their cumulative amount does not exceed about 25%wt. and preferably does not exceed about 20%wt., of the treatment composition of which they form a part.

The inventive compositions are preferably liquids which have a viscosity in the range of about 200 centipoise (“cP”) or less, preferably and in order of increasing preference, viscosities of 150cP, 100 cP, 75 cP, 50 cP, 25 cP, 20 cP, 15 cP, 10 cP, 5 cP, 3 cP, 2 cP, and 1 cP, when measured using conventional quantitative method, e.g., as measured at 20°C or 25°C by a Brookfield Type LVT or Type RVT viscometer using a standard spindle provided by that manufacturer and measuring the samples at room temperature (20 - 25°C).

As the treatment compositions taught herein are used to treat inanimate surfaces including porous and nonporous surfaces and are not provided as a topical skin treatment composition or personal care composition or for that matter as a wound dressing or a preparation for use in wound dressings, the treatment compositions most preferably exclude (unless already described previously) as constituents known-art certain additives and adjuvants which are conventional in the cosmetic, pharmaceutical or dermatological field, specifically hydrophilic or lipophilic gelling agents, hydrophilic or lipophilic active agents, humectants, opacifiers, light stabilizers including UV absorbers, and Polyquaternium type polymers.

The treatment compositions most preferably exclude (unless already described previously) thickener components especially one or more of polysaccharide thickeners such as cellulose, alkyl celluloses, alkoxy celluloses, hydroxy alkyl celluloses, naturally occurring polysaccharide polymers such as xanthan gum, guar gum, locust bean gum, tragacanth gum, or derivatives thereof, polycarboxylate polymers, polyacrylamides, clays, and mixtures thereof.

The treatment compositions of the invention are not provided with an aerosol propellant gas or constituent, and are not packaged or sold as vendible articles in pressured containers, e.g., aerosol canisters. The surface treatment compositions are pourable and pumpable and may be dispensed by pumping the composition through a manually operated or a power driven (e.g.,

motor driven, pressure driven) dispensing device, such as a sprayer, viz “trigger” sprayer or spray pump affixed to a container containing a quantity of the surface treatment composition. The surface treatment composition may also be a pourable composition which may be dispensed from the open end of a suitable flask, bottle or other container, or may be dispensed via a suitable nozzle or spout, e.g., which may be operated by either inversion of the container, and optionally compressing some or part of the container, so to expel it from the container to a surface to be treated. Between such dispensing operations; however, the contents of such a container which includes the surface treatment composition are not pressurized. Thus a further aspect of the invention provides a closed container containing the inventive composition as described herein.

The treatment compositions of the invention may also be supplied within a water dispersible, water miscible or water soluble sachet or pouch or water-soluble package; such may be formed from a water soluble material, such as a water soluble or water dispersible polymeric film (e.g. polyvinyl alcohol), or alternately may be formed from a water insoluble material, such as a water insoluble polymeric film. Additionally the sachet, pouch or package may be formed in a manner where only part of the sachet is physically breachable or only part of the sachet, pouch or package is water soluble or dispersible. Thus a further aspect of the invention provides a closed, a water dispersible, a water miscible or a water soluble sachet or pouch containing the inventive composition as described herein.

The treatment compositions can also be applied to a hard surface by using a wet wipe. The wipe can be of a woven or non-woven nature. Fabric substrates can include nonwoven or woven pouches, sponges, or in the form of abrasive or non-abrasive cleaning pads. Such fabrics are known commercially in this field and are often referred to as wipes. Such substrates can be resin bonded, hydroentangled, thermally bonded, meltblown, needlepunched or any combination of the former.

The nonwoven fabrics may be a combination of wood pulp fibers and textile length synthetic fibers formed by well known dry-form or wet-lay processes. Synthetic fibers such as rayon, nylon, orlon and polyester as well as blends thereof can be employed. The wood pulp fibers should comprise about 30 to about 60 percent by weight of the nonwoven fabric, preferably about 55 to about 60 percent by weight, the remainder being synthetic fibers. The wood pulp fibers provide for absorbency, abrasion and soil retention whereas the synthetic fibers provide for substrate strength and resiliency.

The substrate of the wipe may also be a film forming material such as a water soluble polymer. Such self-supporting film substrates may be sandwiched between layers of fabric substrates and heat sealed to form a useful substrate. The free standing films can be extruded utilizing standard equipment to devolatilize the blend. Casting technology can be used to form and dry films, or a liquid blend can be saturated into a carrier and then dried in a variety of known methods.

The treatment compositions of the present invention are absorbed onto the wipe to form a saturated wipe and sold as a vendible product. The wipe can then be sealed individually in a pouch which can then be opened when needed or a multitude of wipes can be placed in a container for use on an as-needed basis. The container, when closed, is sufficiently sealed to prevent evaporation of any components from the compositions. Thus a further aspect of the invention provides a closed container containing one or more wipes which include the treatment composition as described herein.

The treatment compositions of the invention may be used to provide or impart a microbicidal effect on treated inanimate surfaces. Preferably the surface treatment compositions are characterized in exhibiting a microbicidal benefit when tested against one or more challenge microorganisms according to one or more of the following standardized test protocols: ASTM E1052 Standard Test Method for Efficacy of Antimicrobial Agents against Viruses in Suspension, or ASTM E1053 Standard Test Method to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces, or European Standard Surface Test, EN13697 or AOAC Germicidal Spray Products as Disinfectant Test Method, AOAC Index, 17th Ed. (2000). In particularly preferred embodiments the treatment compositions exhibit a high degree of microbicidal efficacy against various undesirable microorganisms (sometimes referred to as 'pathogens') including various bacteria, viruses, and fungi. In particularly preferred embodiments treatment compositions of the invention exhibit a high degree of microbicidal efficacy against poliovirus type 1 (Sabin) ("PV1").

The surface treatment compositions may be applied to inanimate surfaces in order to impart a cleaning effect thereto, but preferably are applied to impart a microbicidal benefit thereto. Inanimate surfaces include hard surfaces, which are typically nonporous hard surfaces. By way of example, hard surfaces include surfaces composed of refractory materials such as: glazed and unglazed tile, brick, porcelain, ceramics as well as stone including marble, granite,

and other stones surfaces; glass; metals; plastics e.g. polyester, vinyl; fiberglass, Formica®, Corian® and other hard surfaces known to the industry. Hard surfaces which are to be particularly denoted are lavatory fixtures, lavatory appliances (toilets, bidets, shower stalls, bathtubs and bathing appliances), wall and flooring surfaces especially those which include refractory materials and the like. Further hard surfaces which are particularly denoted are those associated with kitchen environments and other environments associated with food preparation. Hard surfaces which are those associated with hospital environments, medical laboratories and medical treatment environments. Inanimate surfaces which may be treated by the surface treatment compositions of the invention include soft surfaces, non-limiting examples of which include: carpets, rugs, upholstery, curtains and drapes, fabrics, textiles, garments, and the like

The treatment compositions described herein may also be used to provide an air treatment benefit if they are sprayed or dispersed into the air, particularly if the surface treatment composition is provided as comminuted particles, viz., droplets within an airspace, such that the treatment composition contacts said airspace and provides a technical benefit thereto, e.g., fragrancng, odor masking, odor elimination, malodour neutralization, air sanitization and the like. The method may be practiced within an open airspace, e.g., a larger volume such as a room, public space within the interior of a building, a cabin or compartment within a vehicle, as well as within a closed container or other relatively smaller space, e.g., the interior of a storage cabinet, a closet, a shower stall, a garbage container or refuse bin, and the like. When used to provide an air treatment benefit, it is preferred that the surface treatment composition comprise at least about 30%wt. of the lower alkyl aliphatic monohydric alcohol constituent, in addition to the further essential constituents heretofore described.

The treatment compositions may also be dispensed, e.g. to a surface, or delivered to an airspace, by means of a mist generator means. Such a mist generator means typically includes an element or member which operates to comminute the unpressurized liquid treatment composition into small particles which form a mist, e.g. nebulize or atomize the unpressurized liquid treatment composition. Such a mist generator means may also be considered an aerosol delivery system which is however not generated from a device wherein the treatment composition also includes a propellant constituent. The mist generator means may comprise a vibrating member which includes a metal or ceramic plate; the plate may be solid or porous, or micropierced in the form of a grid or in the form of one or more segments or slots passing through the vibrating

member, and a piezoelectric actuator which, when operated, causes vibratory motion in the vibrating member. Alternately, the mist generator means may be an electrostatic spray device. Alternately the mist generator means may be an ultrasonic nozzle device. Such devices are known to the art. Non-limiting examples of such mist generators and devices which include such
5 mist generator means include those disclosed in one or more of: US Patent 5743251, US Patent 6234167, US 6491233, US 6501052, US 6516796, US 6568390, US 6640050, US 6681998, US 6766220, US 6772757, US 6804458, US 6883516, US 7229029, US 2007/0011940, US 2007/0169775, US 2007/0235555, US 2008/00419272, US 2009/0121043, US 2009/0272818, the entire contents of each of which are herein incorporated by reference thereto.

10 The mist generator means may be an ultrasonic nozzle device. Such ultrasonic nozzle devices may be obtained from commercial sources, e.g., Sono-Tek, Inc. (Milton, NY, USA) as well as Sonaer Inc., (Farmingdale, N.Y., USA) as well as being disclosed in published patent applications, US 2009/0254020, and US 2009/0224066, the contents of which are herein incorporated by reference.

15 The treatment compositions may also be dispensed, e.g. to a surface, or delivered to an airspace, by means of evaporation of the unpressurized liquid treatment composition particularly to an airspace. For example, the unpressurized liquid treatment composition may be provided in a container for containing the same, and a wick inserted into the unpressurized liquid treatment composition which wick also extends outwardly from the container and from whence the
20 unpressurized liquid treatment composition may evaporate or be otherwise delivered to a surface or to an airspace. Nonlimiting examples of such devices include those disclosed in one or more of: US 7168631, US 6699432, US 6580875, US 4898328, the entire contents of each of which are herein incorporated by reference thereto.

25 Thus a further aspect of the invention provides a closed container containing the inventive composition as described herein.

As certain embodiments of the invention there are provided processes for the treatment of surfaces, or air, including inanimate hard surfaces and inanimate soft surfaces which method includes the step of: contacting such a surface which is in need of treatment or upon which the presence of one or more undesirable microorganisms are suspected or are known to be present,
30 with an effective amount of a surface treatment composition as described herein to provide a surface treatment benefit thereto, preferably to provide a microbicidal benefit to the surface,

particularly against various undesirable microorganisms (sometimes referred to as ‘pathogens’) including various bacteria, mycobacteria, viruses, and fungi, and particularly preferably against poliovirus type 1 (Sabin) (“PV1”). Such methods require the application of an effective amount of a treatment composition as taught herein to such surfaces, so that the desired microbicidal benefit is imparted to the treated surface. Desirably such an effective amount is a sufficient amount of a treatment composition which will provide at least a 1 log₁₀ reduction, more preferably at least, and in order of increasing preference, a 1.25, 1.5, 1.75, 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, 4, 4.25, 4.5, 4.75, and 5 log₁₀ reduction against one or more challenge microorganisms, and especially preferably against poliovirus type 1 (Sabin) (“PV1”) in accordance with one or more of the testing protocols described hereinafter, and/or degrees of microbicidal efficacy of poliovirus type 1 (Sabin) or other challenge microorganism, as are demonstrated with reference to one or more of the Examples.

With reference to the Examples described later, and with reference to Tables C and 1, the disclosed compositions were subjected to one or more of the following test protocols in order to evaluate their microbicidal efficacy against one or more of the other challenge microorganisms which are identified on Table B. As is known in the art, amongst the most difficult to control or eradicate are non-enveloped viruses, such as poliovirus type 1 (Sabin,) and while microbicidal efficacy against the poliovirus type 1 (Sabin) presumptively demonstrates that the same composition would be expected to be effective against the bacteria and the other non-enveloped viruses disclosed on Table B, however, the converse is not expected to be true by a skilled artisan. Thus, for example, while a composition which exhibits good microbicidal efficacy against a Gram-positive or Gram-negative bacteria, such would not be expected to be particularly effective against the poliovirus type 1 (Sabin), while the converse would be expected to be true. Further, even demonstrated efficacy of a composition against a relatively easier to control or eradicate non-enveloped viruses, such as the rotaviruses, or human adenovirus would not necessarily be expected by a skilled artisan to be particularly effective against the poliovirus type 1 (Sabin), while the converse would be expected to be true.

Preferred treatment compositions of the invention demonstrate a microbicidal benefit when tested according to the standardized protocol outlined in ASTM E1052 Standard Test Method for Efficacy of Antimicrobial Agents against Viruses in Suspension. This test is per se, known to the art. Preferred treatment compositions exhibit a microbicidal benefit against one,

but preferably against two or more of the following challenge organisms, specifically “challenge” non-enveloped viruses: hepatitis A (supplied as hepatitis A virus, strain MH-175 ex. Dr. Mark Sobsey, University of North Carolina, Chapel Hill, NC ; human adenovirus type 5 (supplied as ATCC VR-5); feline calicivirus strain F-9 (supplied as ATCC VR-782); herpes simplex type 1 (supplied as ATCC VR-1493); human rhinovirus type 14 strain 1059 (supplied as ATCC VR-284), and especially preferably against poliovirus type 1 (Sabin) (supplied by U.S. Centers for Disease Control and Prevention (CDC)), in accordance with this test protocol. As is known to the skilled artisan, of these forgoing challenge viruses, the most resistant to control or eradicate is the poliovirus type 1 (Sabin) and is it commonly presumed that any composition which shows an effective degree of control or eradication against the poliovirus type 1 (Sabin) virus will exhibit an even greater degree of control or eradication of the further prior listed viruses. As is known from the literature, e.g., *Hierarchy of susceptibility of viruses to environmental surface disinfectants: a predictor of activity against new and emerging viral pathogens*. J. AOAC International 90:1655-1658, Sattar, S.A. (2007), the efficacy of a composition in controlling or eradicating poliovirus type 1 (Sabin) provides an excellent prediction of the composition’s efficacy against further challenge non-enveloped viruses.

Preferred treatment compositions of the invention demonstrate a microbicidal benefit when tested according to the standardized protocol outlined in ASTM E1053 Standard Test Method to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces. This test is also, per se, known to the art. Preferred treatment compositions exhibit a microbicidal benefit against poliovirus type 1 (Sabin) (supplied by U.S. Centers for Disease Control and Prevention (CDC)), in accordance with this test protocol. Again, as is known to the skilled artisan, of these forgoing challenge viruses the most resistant to control or eradication is the poliovirus type 1 (Sabin) and is it commonly presumed that any composition which shows an effective degree of control or eradication against the poliovirus type 1 (Sabin) will exhibit an even greater degree of control or eradication of many other viruses, including hepatitis A virus, feline calicivirus strain F-9, Herpes simplex type 1 and human rhinovirus type 14 strain 1059 as identified above.

Preferred treatment compositions of the invention demonstrate a microbicidal benefit when tested according to the standardized protocol outlined in European Standard Surface Test, EN13697. This test too is, per se, known to the art. Preferred treatment compositions exhibit a

microbicidal benefit against one or more of the following bacteria or fungi: *Staphylococcus aureus* (supplied as ATCC 6538); *Escherichia coli* (supplied as ATCC 10536); *Pseudomonas aeruginosa* (supplied as ATCC 15442); *Enterococcus hirae* (supplied as ATCC 10541) and/or the fungus, *Trichophyton mentagrophytes* (supplied as ATCC 9533) in accordance with the protocols of the test.

Certain preferred treatment compositions of the invention also demonstrate a microbicidal benefit when tested according to the standardized protocol identified as the AOAC Germicidal Spray Products as Disinfectant Test Method, AOAC Index, 17th Ed. (2000). This test is also, per se, known to the art. Preferred treatment compositions exhibit a microbicidal benefit against one or more of the following bacteria or fungi: *Staphylococcus aureus* (“S. aureus” or “Sa”) (supplied as ATCC 6538); *Escherichia coli* (“E. coli”, or “Ec”) (supplied as ATCC 10536), in accordance with the protocols of the test.

The following examples below illustrate exemplary formulations as well as preferred embodiments of the invention. It is to be understood that these examples are provided by way of illustration only and that further useful formulations falling within the scope of the present invention and the claims may be readily produced by one of ordinary skill in the art without deviating from the scope and spirit of the invention.

Examples

A number of treatment compositions as well as a number of comparative compositions were produced and are described on the following Tables. In these identified compositions, the constituents were used “as supplied” from their respective suppliers and may constitute less than 100%wt. “actives”, or may have been supplied as constituting 100%wt. “active” of the named compound, as indicated below. Treatment compositions which are considered to fall within the scope of the present invention are identified by a digit prepended with the letter “E” which indicates this to be an “example” composition, while compositions provided only for the purposes of comparison are identified by a digit prepended with the letter “C”, which indicates this to be a comparative composition and falling outside of the scope of the present invention. In certain of the treatment compositions, one or more constituents, e.g, a pH adjusting agent, or deionized water was added in “quantum sufficient” “q.s.” in order to provide a desired pH or to

provide a sufficient mass in order to provide 100%wt. of each composition. The example compositions disclosed hereinafter include certain presently preferred embodiments of the invention. The comparative compositions are presented on Table C, while treatment compositions of the invention are identified on one or more of the further tables, e.g, Table 1.

5 The compositions disclosed on the following tables were produced by simple mixing, under stirring, of the identified constituents, generally in accordance with the following protocol. To a suitably sized laboratory beaker outfitted with a mechanical stirrer or a magnetic stirrer, was first supplied a major proportion of the deionized water. All of the constituents, as well as the laboratory beaker were at room temperature (approx. 20°C) and as the beaker was open, mixing
 10 was at normal atmospheric pressure. Thereafter under stirring conditions (approx. 300 rpm) was added the source of copper ions, and mixing continued until this material was dissolved. Subsequently while stirring continued, was next added the quaternary ammonium compound(s), and next any pH adjusting agents when such were included. Stirring continued for a further 15 – 30 minutes to ensure a homogenous mixture, to which was next added the alcohol constituent.
 15 Subsequently were added any remaining constituents including any further quantity of pH adjusting constituents (if present) in order to establish the desired pH of the surface treatment composition. Constituents identified as being added “q.s.” were added in order to adjust the pH of the formed composition or to bring the weight of the formed composition to 100%wt. Stirring continued for a further 1 – 15 minutes to ensure the formation of a homogenous mixture, after
 20 which the surface treatment composition was withdrawn from the beaker and used or tested.

These compositions as identified below were formed using the constituents identified on the following Table A which identifies the specific constituents used.

Table A	Constituents
CuSO ₄ .5H ₂ O	CuSO ₄ .5H ₂ O, technical grade (100%wt. actives)
CuCl ₂ .2H ₂ O	CuCl ₂ .2H ₂ O, technical grade (100%wt. actives)
Cu(AcO) ₂ .H ₂ O	Cu(AcO) ₂ .H ₂ O, technical grade (100%wt. actives)
Cu D-gluconate	Cu D-gluconate, technical grade (100%wt. actives)
Cu(I)Cl. H ₂ O	Cu(I)Cl. H ₂ O, technical grade (100%wt. actives)
ethanol (100%)	ethanol, technical grade, 100%wt. actives
ethanol (95%)	ethanol, technical grade, 95%wt. actives

isopropanol	isopropanol, technical grade, 100%wt. actives
n-propanol	n-propanol, technical grade, 100%wt. actives
BTC-65 (50%)	C ₁₂ -C ₁₆ alkyl dimethyl benzyl ammonium chloride provided in an aqueous alcoholic carrier (50%wt. actives) (ex. Stepan)
Onyxide 3300 (33%)	quaternary ammonium complex with saccharinate counterion, (33%wt. actives, balance ethanol) (ex. Stepan Co.)
Neodol 91-6	nonionic surfactant, C ₉ -C ₁₁ linear primary alcohol ethoxylate, avg. 6 mols. ethoxylation, 100%wt. actives (ex. Shell Chemical)
Steol CS370	anionic surfactant, sodium laureth sulfate, 3 mols EO (70%wt. actives) (ex. Stepan Co.)
Glucopon 425N	alkylpolyglycoside (50%wt. actives) (ex. Cognis)
Plurafac SL62	nonionic surfactant, linear alcohol ethoxylate (100%wt. active) (ex. BASF)
Tween 20	sorbitan based surfactant
Silwet Hydrostable 212	silylated/organic surfactant blend (used as supplied) (ex. Momentive Performance Materials, Inc.).
triethanolamine	triethanolamine, technical grade (100%wt. active) (ex. Dow Chemical Company)
ethanolamine	ethanolamine, technical grade (100%wt. active) (ex. Huntsman)
NH ₄ OH (29.86%)	aqueous solution of NH ₄ OH (29.6%wt. active)
NaOH (10%)	aqueous solution of sodium hydroxide, 10%wt. active
NaOH (50%)	aqueous solution of sodium hydroxide, 50%wt. active
Citrosol 502(50%)	aqueous solution of citric acid (50%wt. active) (ex. ADM)
Citrosol 502(50%)/6	aqueous solution of citric acid (50%wt. active) (ex. ADM) further diluted 1:5 in deionized water
citric acid (1.56%)	aqueous solution of anhydrous laboratory grade citric acid (1.56%wt. active)
ammonium acetate	anhydrous solid, reagent grade (ex. JT Baker) (100%wt. actives)
Monacor BE	borate ester blend, used as supplied comprising MEA-borate ester and MIPA-borate ester (ex. Croda, tradename Crodacor BE)
IL2976	corrosion inhibitor comprising mixed acid esters, a proprietary composition believed to be borate esters, used 'as supplied'

Trilon BX	aqueous solution of tetrasodium ethylene diamine tetraacetic acid 38.9%wt. actives) (ex. BASF)
di H ₂ O	deionized water, (100%wt. active)

Further, wherein a specific composition was evaluated for microbicidal efficacy against a challenge microorganism according to one or more of the test protocols identified above, the results of these tests are reported as well. Wherein multiple challenge microorganisms were evaluated in any one test, multiple results are reported.

In the following tables, the tested microorganisms and their identity as reported on the tables are as indentified on Table B:

Table B	Microorganisms
Identifier	Type / Challenge microorganism
“PV1”	Virus / Poliovirus type 1 Sabin, ex. supplied by U.S. Centers for Disease Control and Prevention (CDC)
“HAV”	Virus / Hepatitis A virus, strain MH-175 ex. Dr. Mark Sobsey, University of North Carolina, Chapel Hill, NC
“HAdV”	Virus / Human adenovirus type 5, supplied as ATCC VR-5
“FCV”	Virus / Feline calicivirus strain F-9, supplied as ATCC VR-782 (The use of the feline calicivirus strain F-9 was as a surrogate virus for Norovirus)
“IV-A”	Virus / Influenza A virus, A/California/04/2009 (H1N1), supplied as Biodefence and Emerging Infections Research Resources Repository (BEI Resource) NR-13658
“HSV”	Virus / Herpes simplex type 1, supplied as ATCC VR-1493
“HRV”	Virus / Human rhinovirus type 14 strain 1059, supplied as ATCC VR-284
“Rota”	Virus / Rotavirus Wa strain, supplied as ATCC VR-2018
“ <i>M. bovis</i> ” or “Mb”	Bacteria / <i>Mycobacterium bovis</i> - BCG, supplied by Organon Teknika, Durham, NC to ATS Labs (used as a surrogate for testing efficacy against Tuberculosis)
“ <i>S.aureus</i> ” or “Sa”	Bacteria / <i>Staphylococcus aureus</i> , supplied as ATCC 6538

"E.coli" or "Ec"	Bacteria / <i>Escherichia coli</i> , supplied as ATCC 10536
"P.aeruginosa" or "Pa"	Bacteria / <i>Pseudomonas aeruginosa</i> ("P. aeruginosa") (supplied as ATCC 15442);
"E.hirae", or "Eh"	Bacteria / <i>Enterococcus hirae</i> , supplied as ATCC 10541
"T. mentagropytes" or "T.ment"	Fungus / <i>Trichophyton mentagropytes</i> , supplied as ATCC 9533

In the following tables, Table C describes various "comparative" examples, (which may be identified by the prepended letter "C") while subsequent Table 1 describes various examples of compositions according to the invention, (which may also be identified by the prepended letter "E") as well as the observed physical properties and the results of microbiodical testing according to one or more of the following standardized test protocols:

- A) ASTM E1052 - 96(2002) Standard Test Method for Efficacy of Antimicrobial Agents against Viruses in Suspension, (for a 5 minute contact time, unless specified otherwise) identified on Table 1 as "ASTM E 1052 (log10 reduction)";
- B) ASTM E 1053 - 11 Standard Test Method to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces, (for a 10 minute contact time, unless specified otherwise) identified on Table 1 as "ASTM 1053 (log10 reduction)";
- C) European Standard Surface Test, EN13697, identified on Table 1 as "EN 13697 (log10 reduction)";
- D) AOAC Germicidal Spray Products as Disinfectant Test Method, AOAC Index, 17th Ed. (2000), identified on Table 1 as "AOAC Germicidal Spray". In this test, a result of "0/60" or "1/60" is equivalent to a result of "pass" according to that test's protocols. Results of "2" excess thereof for "/60" tested plates/samples are considered as being equivalent to a "fail" according to that test's protocols.
- E) The European Standard Surface Test, EN13697 protocol was used for testing antifungal efficacy against *Trichophyton mentagrophytes* (supplied as ATCC 9533); the results reported on Tables C and 1 are the log10 reduction of the fungus.

It is noted that each tested composition was not necessarily tested according to all of the foregoing protocols as test results of microbicidal efficacy against Poliovirus type 1 Sabin supports the presumption of efficacy against easier to control or eradicate microorganisms.

In the following tables the amount of the copper ions present are also indicated in parts per million (ppm) and this number is based on the empirical calculation of the available metal ions present in the indicated composition and 100% disassociation of the copper ion from the copper ion source is presumed for this empirical calculation.

5 In the following tables the appearance of the test compositions both of Table 1 and C are indicated; all of the liquid compositions were transparent unless indicated otherwise, e.g. “slight haze” and many were bluish in tint of a lesser or greater substantivity.

All of the compositions of both Table 1 and C were liquids which were readily pourable and pumpable and had a “water-thin” viscosity.

10

Table C (Comparative Examples)			
	C1	C2	C5
CuSO ₄ ·5H ₂ O	--	--	0.1
ethanol (100%)	--	45	--
ethanol (95%)	--	--	--
BTC-65 (50%)	0.2	0.2	--
Onyde 3300 (33%)	--	--	--
Neodol 91-6	0.1	0.1	0.1
triethanolamine	0.022	0.04	0.39
di H ₂ O	99.67	54.66	99.4
pH	9	9.09	8.08
copper ion content (ppm)	0	0	254
appearance	colorless	colorless	light blue
ASTM E 1052 (log10 reduction)	PV1 =< 3.00 IV-A > 5.00 HSV > 6.00 HAdV < 2.0	PV1 < 2.67 PV1 < 2.17 PV1 < 1.50 HSV > 6.00 HAdV > 4.5 HRV = <3.83	PV1 = <2.5 IV-A >6.00 IV-A >5.17 HSV >7.5 HSV >7.0 HAdV <1.0
ASTM 1053 (log10 reduction)	--	--	--
AOAC Germicidal Spray	--	--	Sa =57/60 Pa = 27/60
EN 13697 (log10 reduction)	Sa =4.31 EC >5.50 Pa >4.52 Eh =5.48	--	Sa <1.17 Ec =2.12 Pa =2.84 Eh <1.3
<i>T. ment</i> (log10 reduction)	--	--	--

Table C						
	C6	C7	C8	C9	C10	C11
CuSO ₄ .5H ₂ O	0.1	0.1	--	0.1	--	0.1
ethanol (100%)	--	5	5	10	10	50
ethanol (95%)	--	--	--	--	--	--
BTC-65 (50%)	--	0.2	0.2	0.2	0.2	0.2
Onyde 3300 (33%)	--	--	--	--	--	--
Neodol 91-6	--	0.1	0.1	0.1	0.1	0.1
Plurafac SL62	--	--	--	--	--	--
triethanolamine	0.30	0.31	--	0.31	--	--
citric acid (1.56%)	--	--	--	--	--	--
di H ₂ O	99.56	94.3	94.7	89.3	89.70	49.60
pH	8.08	8.04	8.44	7.97	8.35	4.74
copper ion content (ppm)	254	254	0	254	0	--
appearance	medium blue	medium blue	colorless	light blue	colorless	--
ASTM E 1052 (log10 reduction)	PV1 = <1.5 IV-A <1.33 IV-A <0.33 HSV <2.17 HAdV <1.73	PV1 = < 0.23 IV-A >5.00 HSV >6.00	PV1 <0.23	PV1 <0.9	PV1 <0.23	--
ASTM 1053 (log10 reduction)	--	--	--	--	--	HRV =1.50
AOAC Germicidal Spray	Sa = 60/60 Pa = 23/60	Sa = 0/30 Pa = 0/30	--	--	--	--
EN 13697 (log10 reduction)	Sa <1.97 Ec =0.44 Pa =0.78	Sa =3.48 Ec >6.58 Pa >5.16 Eh =3.43	Sa >6.40 Ec >6.58 Pa >5.16 Eh >6.74	--	--	--
<i>T. ment</i> (log10 reduction)	0.06	--	--	--	--	--

Table C			
	C12	C13	C14
CuSO ₄ ·5H ₂ O	--	--	--
ethanol (100%)	50	50	50
ethanol (95%)	--	--	--
BTC-65 (50%)	0.2	--	--
Onyxide 3300 (33%)	--	--	--
Neodol 91-6	0.1	0.1	--
Plurafac SL62	--	--	--
triethanolamine	0.012	0.014	0.015
di H ₂ O	49.68	49.89	49.985
pH	8.64	8.87	8.98
copper ion content (ppm)	0	0	0
appearance	colorless	colorless	--
ASTM E 1052 (log10 reduction)	PV1 <1.25 HAdv >4.50 HRV >6.00 FCV >4.33 FCV >6.00	PV1 = <2.0	PV1 =<1.5
ASTM 1053 (log10 reduction)	--	--	--
AOAC Germicidal Spray	--	--	--
EN 13697 (log10 reduction)	Sa >6.12 Ec >5.66 Pa >3.80 Eh >6.50	--	--
<i>T.ment</i> (log10 reduction)	> 5.74	--	--

Table C				
	C15	C16	C17	C18
CuSO ₄ .5H ₂ O	--	--	--	--
ethanol (100%)	35	35	35	--
ethanol (95%)	--	--	--	--
BTC-65 (50%)	0.2	--	--	0.2
Onyxide 3300 (33%)	--	--	--	--
Neodol 91-6	0.1	0.1	--	--
triethanolamine	0.016	0.02	0.025	--
di H ₂ O	64.68	64.88	64.97	99.8
pH	8.89	9.08	9.23	8.58
copper ion content (ppm)	0	0	0	0
appearance	clear	clear	clear	clear
ASTM E 1052 (log10 reduction)	PV1 =<2.17	PV1 =<1.17	PV1 =<1.83 FCV >6.00	PV1 =<1.83 IV-A >5.00 HSV >6.00 HAdv <1.5
ASTM 1053 (log10 reduction)	--	--	--	--
AOAC Gemicidal Spray	--	--	--	Sa =1/60 PA =1/60
EN 13697 (log10 reduction)	Sa >6.35, Ec >5.35, Pa >3.85	Sa >6.55 Ec >5.50 Pa >4.52 Eh >6.68	Sa =4.60, Ec =4.19, Pa >4.52 Eh >6.68	Sa =4.47, Ec =4.28, Pa =>4.52 Eh =3.83
<i>T.ment</i> (log10 reduction)	--	--	4.84	2.73

Table 1 (Examples)								
	E1	E2	E3	E4	E5	E6	E7	E8
CuSO ₄ .5H ₂ O	0.1	0.1	0.199	0.1	0.1	0.1	0.1	0.1
ethanol (100%)	25	45	45	75	35	35	50	50
ethanol (95%)	--	--	--	--	--	--	--	--
BTC-65 (50%)	0.2	0.2	--	0.2	0.2	0.2	0.2	0.2
Onyxide 3300 (33%)	--	--	0.30	--	--	--	--	--
Neodol 91-6	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
triethanolamine	0.42	0.19	0.3	0.302	0.443	--	--	0.37
ethanolamine	--	--	--	--	--	--	--	--
NH ₄ OH (29.8%)	--	0.1	0.09	--	--	--	--	--
NaOH (10%)	--	--	--	--	--	--	--	--
Citrosol 502(50%)	--	--	--	--	--	--	--	--
ammonium acetate	--	--	--	--	--	2.0	2.0	--
Monacor BE	--	0.1	0.1	--	--	--	--	--
Trilon BX	--	--	--	--	--	--	--	--
di H ₂ O	74.18	54.21	53.907	24.298	64.157	62.6	47.6	49.22
pH	8.09	9.5	8.85	8.23	8.06	6.82	7.08	8.06
copper ion content (ppm)	254	254	506	254	254	254	254	254
appearance	colorless	light blue	light blue	light blue	light blue	light blue	light blue	light blue
ASTM E 1052 (log ₁₀ reduction)	PV1 >3.67	PV1 >5.27 PV1 >5.88 HAV >6.00 FCV >6.00	PV1 >5.67	PV1 >5.67	PV1 >4.87 FCV >6.00	--	--	PV1 >5.89 HAdV >5.00 HRV >6.00 HAV >6.00 FCV >6.00
ASTM 1053 (log ₁₀ reduction)	--	1 minute: PV1 >4.50 5 minutes: PV1 >4.50	--	--	--	PV1 >4.38 HRV >2.75 Rota >2.5	PV1 >3.50	PV1 >3.50 PV1 >4.50
AOAC Germicidal Spray Test	--	Sa = 0/60 Pa = 0/60 Mb = 0/30 (hard surface test) and ≥ 5.32 (suspension)	--	--	--	--	--	Sa = 0/30 Pa = 0/30

		(test)						
EN 13697 (log10 reduction)	--	--	--	--	--	--	--	Sa >6.29, Ec >5.90, Pa >4.23, En > 5.67
<i>T. ment</i> (log10 reduction)	--	**	--	--	--	--	--	>5.74

Table 1					
	E9	E10	E11	E12	E13
CuSO ₄ .5H ₂ O	0.1	0.1	0.1	0.1	0.1
ethanol (100%)	40	40	40	40	40
ethanol (95%)	--	--	--	--	--
BTC-65 (50%)	--	--	--	--	--
Onyxide 3300 (33%)	0.303	0.303	0.303	0.303	0.303
Neodol 91-6	0.1	0.1	0.1	0.1	0.1
triethanolamine	0.13	0.13	0.13	0.13	0.13
ethanolamine	0.05	0.05	0.05	0.05	0.05
NH ₄ OH (29.8%)	0.12	0.12	0.12	0.12	0.12
NaOH (10%)	--	--	--	--	--
Citrosol 502(50%)	--	--	--	--	--
ammonium acetate	--	--	--	--	--
Monacor BE	--	--	0.20	--	--
Trilon BX	--	--	--	--	--
IL2976	0.060	0.20	--	1.22	0.11
di H ₂ O	58.6	59.0	59.0	58.0	59.1
pH	9.10	9.10	9.59	9.50	9.10
copper ion content (ppm)	254	254	254	254	254
appearance	light blue	light blue	light blue	light blue	light blue
ASTM E 1052 (log10 reduction)	PV1 =<3.44	PV1 =<3.54	PV1 =<5.10	PV1 =<3.10	PV1 =<2.44
ASTM 1053 (log10 reduction)	--	--	--	--	--
AOAC Gemicidal Spray Test	--	--	--	--	--
EN 13697 (log10 reduction)	--	--	--	--	--
<i>T.ment</i> (log10 reduction)	--	--	--	--	--

Table 1					
	E14	E15	E16	E17	E18
CuSO ₄ .5H ₂ O	0.1	0.1	0.1	0.1	0.1
ethanol (100%)	50	50	50	50	50
ethanol (95%)	--	--	--	--	--
BTC-65 (50%)	0.2	0.2	0.2	0.2	0.2
Onyxide 3300 (33%)	--	--	--	--	--
Neodol 91-6	0.1	--	0.1	0.1	0.1
triethanolamine	0.48	0.37	0.41	--	0.48
ethanolamine	--	--	--	--	--
NH ₄ OH (29.8%)	--	--	--	--	--
NaOH (50%)	--	--	0.054	0.03	--
Citrosol 502(50%)	--	--	q.s.	--	--
sodium citrate	--	--	--	0.1	--
ammonium acetate	--	--	--	--	--
Monacor BE	--	--	--	--	0.06
di H ₂ O	49.12	49.32	49.45	49.57	49.06
pH	8.06	8.10	8.24	8.45	8.52
copper ion content (ppm)	254	254	254	254	254
appearance	light blue	light blue	cloudy	colorless	--
ASTM E 1052 (log10 reduction)	PV1 >5.00	PV1 =<4.73	PV1 =<5.00	PV1 >5.67	--
ASTM 1053 (log10 reduction)	PV1 >3.50 PV1 >4.50	--	--	--	--
AOAC Germicidal Spray Test	--	--	--	--	--
EN 13697 (log10 reduction)	--	--	--	--	--

	E19	E20	E21	E22	E23
CuSO ₄ .5H ₂ O	0.1	0.1	0.1	0.1	0.1
ethanol (100%)	--	45	45	25	45
ethanol (95%)	50	--	--	--	--
isopropanol	--	--	--	20	--
BTC-65 (50%)	0.1	--	--	0.2	0.2
Onyxide 3300 (33%)	--	0.3	0.3	--	--
Neodol 91-6	0.1	0.1	0.1	0.1	0.1
Steol CS370	--	--	--	--	0.143
triethanolamine	0.31	0.48	0.18	0.4	0.43
ethanolamine	--	--	--	--	--
NH ₄ OH (29.8%)	--	--	0.11	--	--
NaOH (10%)	--	--	--	--	--
Citrosol 502(50%)	--	--	--	--	--
Citrosol 502(50%)/6	--	--	--	--	--
ammonium acetate	--	--	--	--	--
Monacor BE	--	--	--	--	--
Trilon BX	--	--	--	--	--
di H ₂ O	49.39	54.02	54.21	54.2	54.027
pH	8.1	8.2	9.17	8.03	8.08
copper ion content (ppm)	254	254	254	254	254
appearance	light blue	light blue	light blue	light blue	dark blue, slight haze
ASTM E 1052 (log10 reduction)	--	PV1 >5.27 HRV >4.83 HSV >6.00	PV1 >4.50	PV1 = 4.83	PV1 >6.00
ASTM 1053 (log10 reduction)	--	--	PV1 >4.50 (at 1 min), PV1 >4.50 (at 5 min)	--	--
AOAC Germicidal Spray Test	--	--	Sa = 0/60 Pa = 0/60	--	--
EN 13697 (log10 reduction)	--	--	--	--	--

Table 1						
	E24	E25	E26	E27	E28	E29
CuSO ₄ .5H ₂ O	0.1	0.1	0.1	0.1	0.1	0.1
ethanol (100%)	45	--	45	45	45	45
ethanol (95%)	--	--	--	--	--	--
isopropanol	--	--	--	--	--	--
n-propanol	--	45	--	--	--	--
BTC-65 (50%)	0.2	0.2	0.2	--	--	0.2
Onyxide 3300	--	--	--	--	--	--
BTC-2125M	--	--	--	0.2	--	--
BTC-1010	--	--	--	--	0.2	--
Neodol 91-6	0.1	0.1	--	0.1	0.1	--
Steol CS370	--	--	0.143	--	--	--
Glucopon 425N	--	--	--	--	--	0.2
triethanolamine	0.41	0.4	0.41	0.42	0.46	0.42
ethanolamine	--	--	--	--	--	--
NH ₄ OH (29.8%)	--	--	--	--	--	--
NaOH (10%)	--	--	--	--	--	--
Citrosol 502(50%)	--	--	--	--	--	--
ammonium acetate	--	--	--	--	--	--
Monacor BE	--	--	--	--	--	--
Trilon BX	--	--	--	--	--	--
di H ₂ O	54.19	54.2	54.14	54.18	54.14	54.08
pH	8.09	8.09	8.11	8.12	8.12	8.13
copper ion content (ppm)	254	254	254	254	254	254
appearance	light blue	--	dark blue, slight	light blue	light blue	dark blue
ASTM E 1052 (log ₁₀ reduction)	PV1 >6.33 HRV >6.00	PV1 <3.00	PV1 >6.00	PV1 >6.00	PV1 >6.00	PV1 >6.00
ASTM 1053 (log ₁₀ reduction)	--	--	--	--	--	--
AOAC Germicidal Spray Test	Sa = 0/30 Pa = 0/30	--	--	--	--	--
EN 13697 (log ₁₀)	--	--	--	--	--	--




Table 1	E30	E31	E32	E33	E34	E35	E36	E37
CuSO ₄ ·5H ₂ O	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
ethanol (100%)	45	45	40	38	45	38	38	38
ethanol (95%)	--	--	--	--	--	--	--	--
isopropanol	--	--	--	--	--	--	--	--
n-propanol	--	--	--	--	--	--	--	--
BTC-65 (50%)	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Onyxde 3300 (33%)	--	--	--	--	--	--	--	--
BTC-2125M	--	--	--	--	--	--	--	--
Neodol 91-6	--	--	0.1	0.1	0.1	0.1	0.1	0.1
Steol CS370	--	--	--	--	--	--	--	--
Ammonyx LO	0.33	--	--	--	--	--	--	--
Plurafac SL62	--	0.1	--	--	--	--	--	--
triethanolamine	0.48	0.43	0.42	0.015	0.0102	0.42	0.08	--
ethanolamine	--	--	--	--	--	--	--	--
NH ₄ OH (29.8%)	--	--	--	--	--	--	0.11	0.11
NaOH (10%)	--	--	--	--	--	--	--	--
Citrosol 502(50%)	--	--	--	0.08	0.066	--	--	--
ammonium acetate	--	--	--	--	--	--	--	--
Monacor BE	--	--	--	--	--	--	--	--
Trilon BX	--	--	--	--	--	--	--	--
di H ₂ O	53.88	54.17	59.18	61.58	54.4	61.18	61.41	61.49
pH	8.18	8.14	8.10	5.0	5.04	8.08	9.20	9.38
copper ion content (ppm)	254	254	254	254	254	254	254	254
appearance	dark blue	light blue	light blue	light blue	light blue	light blue	light blue	--
ASTM E 1052 (log10 reduction)	PV1 >6.00	PV1 >6.00	PV1 >5.69 HRV >6.00	PV1 >5.62 PV1 >4.73 HAV >=6.00 HAdV >4.50 FCV > 3.33	PV1 >=4.73	PV1 = 4.54 HRV >6.00	PV1 >5.65 HAV >6.00 FCV >6.00	--
ASTM 1053 (log10 reduction)	--	--	--	--	PV1 >=3 (5 min, aluminum Can, complete inactivation)	--	--	--

					PV1=3.5 (5min, glass jar pressured, completed inactivation)			
ADAC Germicidal Spray Test	-	-	-	-	-	-	Sa = 0/30 Pa = 0/30	-
EN 13697 (log10 reduction)	-	-	-	Sa >6.27 Ec >6.25 Pa >4.97 Eh >6.56	-	-	-	-

Table 1	E38	E39	E40	E41	E42	E43
CuSO ₄ ·5H ₂ O	0.1	0.1	0.1	0.1	0.1	0.1
ethanol (100%)	35	35	35	30	25	20
ethanol (95%)	--	--	--	--	--	--
BTC-65 (50%)	0.2	0.2	0.2	0.2	--	--
Oryxide 3300 (33%)	--	--	--	--	0.30	0.30
Neodol 91-6	--	0.1	0.1	0.1	0.1	0.1
triethanolamine	0.41	0.4	0.1	0.45	0.1	0.28
ethanolamine	--	--	0.1	--	--	--
NH ₄ OH (29.8%)	--	--	0.17	0.21	0.10	0.18
NaOH (10%)	--	--	0.25	--	--	--
Citrosol 502(50%)	--	--	--	--	--	--
ammonium acetate	--	--	--	--	0.06	--
Monacor BE	--	--	--	--	--	--
di H ₂ O	64.29	64.2	63.98	68.94	74.2	79.0
pH	8.1	8.1	11.03	9.51	9.21	9.16
copper ion content (ppm)	254	254	254	254	254	254
appearance	light blue	light blue	light blue	light blue	light blue	light blue
ASTM E 1052 (log10 reduction)	PV1 <3.33	PV1 = 3.17, FCV >6.0	PV1 >5.88	PV1 >5.17, HAV ≤2.00, HAdV >4.50, HSV-1 >6.0	PV1 ≤3.23	PV1 <3.00, FCV >6.00
ASTM 1053 (log10 reduction)	--	--	--	--	--	--
AOAC Gericidal Spray Test	--	--	--	--	--	--
EN 13697 (log10 reduction)	--	Sa >6.12 Ec >5.66 Pa >3.80 Eh > 6.50	--	Sa >6.62 Ec >6.63 Pa >6.69 Eh >6.74	--	--

Table 1	E44	E45	E46	E47
CuSO ₄ ·5H ₂ O	0.099	0.099	0.05	0.025
ethanol (100%)	45	45	45	45
ethanol (95%)	--	--	--	--
BTC-65 (50%)	--	--	--	--
Onyxide 3300 (33%)	0.151	0.076	0.303	0.303
Neodal 91-6	0.1	0.1	0.1	0.1
triethanolamine	0.35	0.28	0.2	0.1
ethanolamine	--	--	--	--
NH ₄ OH (29.8%)	0.09	0.09	0.13	0.08
NaOH (10%)	--	--	--	--
Citrosol 502(50%)	--	--	--	--
ammonium acetate	--	--	--	--
Monacor BE	--	--	0.06	0.06
di H ₂ O	54.21	54.35	54.15	54.33
pH	8.97	9.05	9.84	10.06
copper ion content (ppm)	254	254	127	64
appearance	light blue	light blue	light blue	light blue
ASTM E 1052 (log ₁₀ reduction)	PV1 >5.67	PV1 >5.67	PV1 >5.67	PV1 >5.67
ASTM 1053 (log ₁₀ reduction)	--	--	--	--
AOAC Germicidal Spray Test	--	--	--	--
EN 13697 (log ₁₀ reduction)	--	--	--	--

	E48	E49	E50	E51
CuSO ₄ ·5H ₂ O	--	--	--	--
CuCl ₂ ·2H ₂ O	0.1	--	--	--
Cu(AcO) ₂ ·H ₂ O	--	0.1	--	--
Cu D-gluconate	--	--	0.1	--
Cu(I)Cl·H ₂ O	--	--	--	0.1
ethanol (100%)	45	45	45	45
ethanol (95%)	--	--	--	--
BTC-65 (50%)	0.2	0.2	0.2	--
Onyxide 3300 (33%)	--	--	--	0.3
Neodol 91-6	0.1	0.1	0.1	0.1
triethanolamine	0.6	0.45	0.45	0.45
ethanolamine	--	--	--	--
NH ₄ OH (29.8%)	--	--	--	--
NaOH (10%)	--	--	--	--
Citrosol 502(50%)	--	--	--	--
di H ₂ O	54.0	54.15	54.35	54.05
pH	8.1	8.03	8.03	8.14
copper ion content (ppm)	372	318	140	641
appearance	light blue	light blue	--	light blue, slight haze
ASTM E 1052 (log10 reduction)	PV1 >6.00	PV1 >6.00	PV1 =<5.50	PV1 >6.00 FCV =>6.00
ASTM 1053 (log10 reduction)	--	--	--	--
EN 13697 (log10 reduction)	--	--	--	--

Several of the compositions of Table 1 were also tested for product stability and performance after being stored for an extended period of time (4 weeks or more) at room temperature (approx. 20 - 22°C) to determine the microbicidal efficacy of the compositions following storage. The results in the following Table 1A below demonstrate the continued microbicidal efficacy of compositions according to the invention.

	E8	E2	E21
pH	8.06	9.5	9.17
ASTM E 1052, (log10 reduction)	at 4 months: PV1 =>4.73 at 6 months: PV1 =>4.56 FCV >6.00	at 3 months: PV1 >5.50	at 3 months: PV1 >4.27

As can be seen from the foregoing results, the compositions of the invention exhibited excellent microbicidal efficacy as demonstrated by the various test results, even wherein reduced levels of ethanol (e.g, less than 50%wt, especially 45%wt.) or less ethanol) was present as a constituent.

Although this invention has been shown and described with respect to the detailed embodiments thereof, it will be understood by those of ordinary skill in the art that various changes may be made and equivalents may be substituted for elements thereof without departing from the scope of the invention. In addition, modifications may be made to adapt a particular situation or material to the teachings of the invention without departing from the essential scope thereof. Therefore, it is intended that the invention not be limited to the particular embodiments disclosed in the above detailed description, but that the invention will include all embodiments falling within the scope of the appended claims.

C:\ANPCMB\102792\1210\PCTAppn.doc

Claims:

1. A liquid, inanimate surface treatment composition which imparts a microbicidal benefit
5 to such treated surfaces which composition comprises:
a copper source material which releases copper ions into the treatment
composition, preferably a source of Cu(I) and/or Cu(II) ions;
at least 20%wt. of at least one lower alkyl aliphatic monohydric alcohol,
at least one quaternary ammonium compound which provides a microbicidal
10 benefit;
water;
optionally, one or more further constituents which impart one or more
advantageous technical or aesthetic benefits to the compositions, including one or more
detergent surfactants;
15 wherein the composition has a pH of at least 5,
wherein the surface treatment compositions are characterized in exhibiting a
microbicidal benefit when tested against one or more challenge microorganisms,
according to one or more of the following standardized test protocols: ASTM E1052
Standard Test Method for Efficacy of Antimicrobial Agents against Viruses in
20 Suspension, or ASTM E1053 Standard Test Method to Assess Virucidal Activity of
Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces,
or European Standard Surface Test, EN13697, or AOAC Germicidal Spray Products as
Disinfectant Test Method, AOAC Index, 17th Ed. (2000) and, especially preferably
wherein the challenge microorganism is poliovirus type 1 (Sabin) ("PVI").
25
2. A composition according to claim 1 which comprises at least 30%wt. of at least one
lower alkyl aliphatic monohydric alcohol.
3. A composition according to claim 1 which comprises wherein ethanol is the predominant
30 or sole lower alkyl aliphatic monohydric alcohol present in the composition.

4. A composition according to any of claims 1 – 3, which further comprises at least one nonionic surfactant constituent.
5. A composition according to any of claims 1 – 4, wherein the compositions have a pH of at least 8.
6. A composition according to claim 1, which composition comprises:
a copper source material which releases copper ions into the treatment composition, preferably a source of Cu(I) and/or Cu(II) ions;
at least 40%wt. of at least one lower alkyl aliphatic monohydric alcohol,
at least one quaternary ammonium compound which provides a microbicidal benefit;
water;
optionally, one or more further constituents which impart one or more advantageous technical or aesthetic benefits to the compositions, including one or more deterative surfactants;
wherein the composition has a pH of at least 8.5,
wherein the surface treatment compositions are characterized in exhibiting at least a 3 log₁₀ reduction of human Rhinovirus when tested according to the standardized test protocols of ASTM E1053 Standard Test Method to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces.
7. A composition according to claim 1, which composition comprises:
a copper source material which releases copper ions into the treatment composition, preferably a source of Cu(I) and/or Cu(II) ions;
at least 40%wt. of at least one lower alkyl aliphatic monohydric alcohol,
at least one quaternary ammonium compound which provides a microbicidal benefit;
water;

optionally, one or more further constituents which impart one or more advantageous technical or aesthetic benefits to the compositions, including one or more deterative surfactants;

wherein the composition has a pH of at least 8.5,

5 wherein the surface treatment compositions are characterized in exhibiting at least a 3 log₁₀ reduction of poliovirus type 1 Sabin when tested according to the standardized test protocols of ASTM E1053 Standard Test Method to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces.

10 8. A microbicidal control system of constituents which are in and of themselves effective in providing effective control of poliovirus independently of further and optional constituents, which microbicidal control system of constituents comprises (or consists essentially of, or consists of): as a first constituent, water, as a next constituent at least 20%wt. of one or more one or more C₁-C₄ aliphatic alcohols and especially preferably
15 wherein ethanol is the predominant or sole C₁-C₄ aliphatic alcohols present, a cationic quaternary ammonium compound which provides a microbicidal benefit and, where necessary, a buffer or pH adjusting agent to impart an alkaline pH, preferably an alkaline pH of 7.5 or greater.

20 9. A microbicidal control system of constituents according to claim 8, which further comprises at least one nonionic surfactant constituent.

10. A microbicidal control system of constituents which are in and of themselves effective in providing effective control of poliovirus independently of further and optional
25 constituents which microbicidal control system of constituents comprises (or consists essentially of, or consists of) as a first constituent, water, as a next constituent at least 20%wt. of one or more one or more C₁-C₄ aliphatic alcohols and especially preferably wherein ethanol is the predominant or sole C₁-C₄ aliphatic alcohol present, a cationic quaternary ammonium compound which provides a microbicidal benefit, a surfactant
30 constituent which is preferably at least one nonionic surfactant, and where necessary a

buffer or pH adjusting agent to impart an alkaline pH, and preferably an alkaline pH of 7.5 or greater.

5 11. A microbicidal control system of constituents according to claim 9, which comprises at least one nonionic surfactant constituent.

12. A method of controlling the incidence of undesired microorganisms on an inanimate surface, the method comprising the step of:

10 contacting an inanimate surface which is in need of treatment or upon which the presence of one or more undesirable microorganisms are suspected or are known to be present, with an effective amount of a liquid, inanimate surface treatment composition according to any of claims 1 - 7 or with an effective amount of a microbicidal control system according to any of claims 8 - 11 to provide a microbicidal benefit to the contacted surface.

15

13. An air treatment composition effective in controlling the incidence of undesired microorganisms in air which comprises:

a copper source material which releases copper ions into the treatment composition, preferably a source of Cu(I) and/or Cu(II) ions;

20

at least 20%wt. of at least one lower alkyl aliphatic monohydric alcohol,

at least one quaternary ammonium compound which provides a microbicidal benefit;

water;

25

optionally, one or more further constituents which impart one or more advantageous technical or aesthetic benefits to the compositions, including one or more deterative surfactants;

wherein the composition has a pH of at least 5,

30

wherein the surface treatment compositions are characterized in exhibiting a microbicidal benefit when tested against one or more challenge microorganisms, according to one or more of the following standardized test protocols: ASTM E1052 Standard Test Method for Efficacy of Antimicrobial Agents against Viruses in

5 Suspension, or ASTM E1053 Standard Test Method to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces, or European Standard Surface Test, EN1369, or AOAC Germicidal Spray Products as Disinfectant Test Method, AOAC Index, 17th Ed. (2000) against one or more challenge
microorganisms, especially preferably wherein the challenge microorganism is poliovirus
type 1 (Sabin) ("PVI").

10 14. A composition according to claim 13 which comprises wherein ethanol is the predominant or sole lower alkyl aliphatic monohydric alcohol present in the composition.

15 15. A composition according to claim 13, which further comprises at least one nonionic surfactant constituent.

20 16. A method of controlling the incidence of undesired microorganisms in air, or in a headspace such as the ambient air within a closed volume such as a room or the interior of a vehicle, the method comprising the step of: delivering and dispersing within an airspace an effective amount of an air treatment composition according to any of claims 13 - 15, to provide a microbicidal benefit to the treated air.

Fig. 1






Figure 2
pH 9 - 10





INTERNATIONAL SEARCH REPORT

International application No

PCT/GB2012/051118

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>Anonymous: "Disinfectant", Wikipedia, the free encyclopedia</p> <p>9 October 2010 (2010-10-09), XP002680035, Retrieved from the Internet: URL: http://web.archive.org/web/20101009063613/http://en.wikipedia.org/wiki/Disinfectant [retrieved on 2012-07-16] sections 2.1 and 2.5</p> <p>-----</p>	1-16
Y	<p>US 5 373 025 A (GAY WALTER A [US]) 13 December 1994 (1994-12-13) claims 1-5 column 9, line 4 - line 21</p> <p>-----</p>	1-16
Y	<p>US 2009/136581 A1 (GUTIERREZ MARTINEZ ALFONSO [MX] ET AL) 28 May 2009 (2009-05-28) claims 1, 7</p> <p>-----</p>	1-16
Y	<p>EP 1 767 495 A1 (HISHIDA IWA0 [JP]) 28 March 2007 (2007-03-28) abstract claim 1</p> <p>-----</p>	1-16
Y	<p>WO 03/073857 A2 (SPIESS URANIA CHEMICALS GMBH [DE]; FRANKE FRIEDRICH [DE]; GOEBEL GERHA) 12 September 2003 (2003-09-12) abstract claims 1-11</p> <p>-----</p>	1-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/GB2012/051118

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2009130608 A2	29-10-2009	US 2010015245 A1 WO 2009130608 A2	21-01-2010 29-10-2009

US 2010209460 A1	19-08-2010	NONE	

US 5373025 A	13-12-1994	AU 7046094 A US 5373025 A WO 9429266 A1	03-01-1995 13-12-1994 22-12-1994

US 2009136581 A1	28-05-2009	NONE	

EP 1767495 A1	28-03-2007	EP 1767495 A1 JP 4561558 B2 JP 2007084493 A US 2007065519 A1	28-03-2007 13-10-2010 05-04-2007 22-03-2007

WO 03073857 A2	12-09-2003	AU 2003213997 A1 BR 0308182 A DE 10209600 A1 EP 1480518 A2 US 2005159482 A1 WO 03073857 A2	16-09-2003 11-01-2005 18-09-2003 01-12-2004 21-07-2005 12-09-2003
