



(51) International Patent Classification:

C07D 231/14 (2006.01) A61P 11/00 (2006.01)
A61K 31/415 (2006.01)

(21) International Application Number:

PCT/US20 12/07 1232

(22) International Filing Date:

21 December 2012 (21.12.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/580,495 27 December 2011 (27.12.2011) US

(71) Applicant: ALLERGAN, INC. [US/US]; 2525 Dupont Drive, Irvine, California 92612 (US).

(72) Inventors; and

(71) Applicants : CARLING, William R. [GB/GB]; 28 Yeomans Close, Bishop's Stortford Hertfordshire CM23 4EU (GB). MARTOS, Jose L. [ES/GB]; 28 Newberry Side, Basildon Essex SSI 5 5XB (GB). KANGASMETSA, Jussi J. [SE/GB]; 9 Railway Cottages, Wendens Ambo, Saffron Essex CB11 4LA (GB). WANG, Jenny W. [US/US]; 2 Giverny, Newport Coast, California 92657 (US). WOODWARD, David F. [US/US]; 22736 Islamare Lane, Lake Forest, California 92630 (US).


(74) Agents: FORRESTAL, Kevin et al; Allergan, Inc., 2525 Dupont Drive, Irvine, California 92612 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on nextpage]

(54) Title: COMPOUNDS ACTING AT MULTIPLE PROSTAGLANDIN RECEPTORS GIVING A GENERAL ANTI-INFLAMMATORY RESPONSE



(57) Abstract: The present invention provides a compound, that is wherein Y, W, Z, R1, R2, R4, R5 and R6 are as defined in the specification. The compounds may be administered to treat DP, FP, EP1, TP and/or EP4 receptor-mediated diseases or conditions.

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

**COMPOUNDS ACTING AT MULTIPLE PROSTAGLANDIN RECEPTORS GIVING
A GENERAL ANTI-INFLAMMATORY RESPONSE**

BY INVENTORS

**WILLIAM R. CARLING, JOSE L. MARTOS, JUSSI J. KANGASMETSA,
JENNY W. WANG AND DAVID F. WOODWARD**

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Patent Application No. 61/580,495,
filed on December 27, 2011, the disclosure of each of which is incorporated herein in its
entirety by this specific reference.

BACKGROUND OF THE INVENTION

This invention relates to compounds, to processes for their preparation, to pharmaceutical
compositions containing them and to their use in medicine, in particular their use in the
treatment of conditions mediated by the action of ligands for the DP₁, FP, TP, EP₁ and EP₄
prostaglandin (PG) receptors. The present compounds have the general structure shown
below and act at different prostaglandin receptors to thereby provide a general anti-
inflammatory response.

SUMMARY OF THE RELATED ART

The EP₁ receptor is a 7-transmembrane receptor and its natural ligand is the prostaglandin
PGE₂. PGE₂ also has affinity for the other EP receptors (types EP₂, EP₃ and EP₄). The EP₁
receptor is associated with smooth muscle contraction, pain (in particular inflammatory,
neuropathic and visceral), inflammation, allergic activities, renal regulation and gastric or
enteric mucus secretion.

Prostaglandin E₂ (PGE₂) exerts allodynia through the EP₁ receptor subtype and hyperalgesia
through EP₂ and EP₃ receptors in the mouse spinal cord. Furthermore, it has been shown that
in the EP₁ knock-out mouse pain-sensitivity responses are reduced by approximately 50%.
EP₁ receptor antagonist (ONO-8711) reduces hyperalgesia and allodynia in a rat model of
chronic constriction injury and inhibits mechanical hyperalgesia in a rodent model of post-
operative pain. The efficacy of EP₁ receptor antagonists in the treatment of visceral pain in a

human model of hypersensitivity has been demonstrated. Thus, selective prostaglandin ligands, agonists or antagonists, depending on which prostaglandin E receptor subtype is being considered, have anti-inflammatory, antipyretic and analgesic properties similar to a conventional non-steroidal anti-inflammatory drug, and in addition, inhibit hormone-induced uterine contractions and have anti-cancer effects. These compounds have a diminished ability to induce some of the mechanism-based side effects of NSAIDs which are indiscriminate cyclooxygenase inhibitors. In particular, the compounds have a reduced potential for gastrointestinal toxicity, a reduced potential for renal side effects, a reduced effect on bleeding times and a lessened ability to induce asthma attacks in aspirin-sensitive asthmatic subjects. Moreover, as a result of sparing potentially beneficial prostaglandin pathways, these agents may have enhanced efficacy and safety over NSAIDs and/or COX-2 inhibitors. EP₄ receptors have also been implicated in pain, hyperalgesia, allodynia, and inflammation. (See Pub. No. US 2005/0065200 for other diseases that may be treated by EP₄ receptor antagonists.)

The TP (also known as TxA₂) receptor is a prostanoid receptor subtype stimulated by the endogenous mediator thromboxane. Activation of this receptor results in various physiological actions primarily incurred by its platelet aggregatory and smooth muscle constricting effects, thus opposing those of prostacyclin receptor activation.

TP receptors have been identified in human kidneys in the glomerulus and extraglomerular vascular tissue. Activation of TP receptors constricts glomerular capillaries and suppresses glomerular filtration rates indicating that TP receptor antagonists could be useful for renal dysfunction in glomerulonephritis, diabetes mellitus and sepsis.

Activation of TP receptors induces bronchoconstriction, an increase in microvascular permeability, formation of mucosal edema and mucus secretion, which are typical characteristic features of bronchial asthma. TP antagonists have been investigated as potential asthma treatments resulting in, for example, orally active Seratrodist (AA-2414). Ramatroban is another TP receptor antagonist currently undergoing phase III clinical trials as an anti-asthmatic compound.

Since DPi receptor stimulation may trigger an asthmatic response in certain individuals, compounds that have DPi antagonist properties may be useful as anti-asthmatic drugs. (See Pub. No. 2004/0162323 for the disclosure of other diseases and conditions that may be treated with DP antagonists.)

5

Finally, the FP receptor modulates intraocular pressure and mediates smooth muscle contraction of the sphincter muscles in the gastrointestinal tract and the uterus. Thus, antagonists of the FP receptor are useful for treating reproductive disorders. (See US Patent No. 6,511,999 for other diseases and conditions that may be treated with FP receptor antagonists.)

10

As further background for the present invention, see US Published Patent Application 2007/0060596.

15 BRIEF SUMMARY OF THE INVENTION

This invention provides novel compounds, that are 1-[(5-halo or alkyl or fluoroalkyl or alkoxy-2-[(cycloalkyl)oxy]phenyl)methyl]-(5-alkyl or fluoroalkyl)-1H-pyrazole-3-(carboxylic acid or methylene carboxylic acids) and alkyl or aryl ester or sulfonamides thereof

20 Preferably the ester or sulfonamide is an alkyl ester or sulfonamide. Preferably said compound is a 1-[(5-halo-2-[(cycloalkyl)oxy]phenyl)methyl]-(5-alkyl)-1H-pyrazole-3-(carboxylic acid) or an alkyl or aryl ester or sulfonamide thereof. Preferably, said halo is chloro or bromo and said cycloalkylalkyl is cyclopentyl

25 The invention further relates to pharmaceutical compositions containing the above compounds in combination with a pharmaceutically-acceptable excipient and to their use in medicine, in particular their use in the treatment of conditions mediated by the action of ligands for the DPi, FP, EPi and EP₄ prostaglandin (PG) receptors. The compounds of this invention are also useful for treating conditions mediated by the action of ligands for the
30 thromboxane (TP) receptor.

Some embodiments of the present invention include:

1. A compound, that is a 1-[(5-halo or alkyl or fluoroalkyl or alkoxy-2-
 {(cycloalkyl)oxy}phenyl)methyl]-(5-alkyl or fluoroalkyl)-1H-pyrazole-3-(carboxylic acid or
 methylene carboxylic acid) or an alkyl or aryl ester or sulfonamide thereof.


2. A compound according to paragraph 1 wherein said compound is a 1-({5-halo-2-[(2-
 cycloalkyl)oxy]phenyl}methyl)-3-carboxylic acid or ester or sulfonamide thereof.

3. A compound according to paragraph 1 wherein said compound is a 1-({5-halo-2-[(2-
 cycloalkyl)oxy]phenyl}methyl)-3-carboxylic acid.

4. A compound according to paragraph 2 wherein said ester or sulfonamide is an alkyl
 ester or sulfonamide.

5. A compound according to paragraph 3, wherein said halo is selected from the group
 consisting of chloro and bromo.

6. A compound represented by the following formula



wherein Y is $(CH_2)_m$ wherein m is 0 or an integer of from 1 to 3;

Z is selected from the group consisting of O, S, SO, SO_2 and $(CH_2)_p$, wherein p is 0 or an
 integer of from 1 to 3;

W is substituted phenyl ring, alkyl, substituted alkyl, cycloalkyl or substituted cycloalkyl:

R_i is selected from the group consisting of OR₇, N(R₇)₂, and N(R₇)SO₂R₇;

R₂ is H;

5

R₄ is selected from the group consisting of alkyl, halogen-substituted alkyl and amino,

R₅ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxyl, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy;

10

R_e is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxyl, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy, wherein at least one of R₅ and R₆ is halogen; and

15

R₇ is selected from the group consisting of H, hydrocarbyl and substituted hydrocarbyl.

8. The compound of paragraph 7 wherein R_i is OH.

20

9. The compound of paragraph 8 wherein R₇ is selected from the group consisting of carbocyclic aryl and alkyl

10. The compound of paragraph 9 wherein W is cycloalkyl.

25

11. The compound of paragraph 10 wherein R₄ is selected from the group consisting of alkyl and chloro and bromo-substituted alkyl.

12. The compound of paragraph 10 wherein R₄ is H.

30

13. The compound of paragraph 10 wherein Y is absent, i.e. n is 0.

14. The compound of paragraph 10 wherein Z is O.

15. The compound of paragraph 10 wherein W is cyclopentyl


16. The compound of paragraph 10 that is selected from the group consisting of

1-(5-Chloro-2-cyclopentylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid,

1-(5-Bromo-2-cyclopentylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid and

1-(5-Bromo-2-cyclobutylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid

17. A method comprising administering a compound represented by the following formula



wherein Y is $(\text{CH}_2)_m$ wherein m is 0 or an integer of from 1 to 3;

Z is selected from the group consisting of O, S, SO, SO_2 and $(\text{CH}_2)_p$, wherein p is 0 or an integer of from 1 to 3;

W is substituted phenyl ring, alkyl, substituted alkyl, cycloalkyl or substituted cycloalkyl:

R_i is selected from the group consisting of OR₇, N(R₇)₂, and N(R₇)SO₂R₇;

R₂ is H ;

R₄ is selected from the group consisting of alkyl, halogen-substituted alkyl and amino;

R₅ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxyl, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy;

- 5 R₆ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxy, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy, wherein at least one of R₅ and R₆ are halogen;and

R₇ is selected from the group consisting of H, hydrocarbyl and substituted hydrocarbyl.

10

18. The method of paragraph 17 wherein R_i is OH.

19. The method compound of paragraph 18 wherein R₇ is selected from the group consisting of carbocyclic aryl and alkyl

15

20. The method of paragraph 19 wherein W is cycloalkyl.

21. The method of paragraph 20 wherein R₄ is selected from the group consisting of alkyl and chloro and bromo-substituted alkyl.

20

22. The method of paragraph 20 wherein R₄ is H.

23. The method of paragraph 20 wherein Y is absent, i.e. n

25 24. The method of paragraph 20 wherein Z is O. is 0.

25. The compound of paragraph 20 wherein W is cyclopentyl

26. The compound of paragraph 20 that is selected from the group consisting of

30

1-(5-Chloro-2-cyclopentylmethoxy-benzyl)-5 -methyl-1H-pyrazole-3-carboxylic acid,

1-(5-Bromo-2-cyclopentylmethoxy-benzyl)-5 -methyl-1H-pyrazole-3-carboxylic acid and

35 1-(5-Bromo-2-cyclobutylmethoxy-benzyl)-5 -methyl-1H-pyrazole-3-carboxylic acid

27. The method of paragraph 26 wherein said compound is administered to treat DPI, FP, EP1, TP and/or EP4 receptor mediated diseases or conditions.

5

28. The method of paragraph 27 wherein said condition or disease is related to inflammation.

29. The method of paragraph 27 wherein said DPI , FP, EP1, TP and/or EP4 receptor mediated condition or disease is selected from the group consisting of allergic conditions, asthma, allergic asthma, allergic rhinitis, uveitis and related disorders , atherosclerosis, blood coagulation disorders, bone disorders, cancer, cellular neoplastic transformations, chronic obstructive pulmonary diseases and other forms of lung inflammation, congestive heart failure, diabetic retinopathy, diseases or conditions requiring a treatment of anti-coagulation, diseases requiring control of bone formation and resorption, endometriosis , fertility disorders, , gangrene, glaucoma, hyperpyrexia, immune and autoimmune diseases, inflammatory conditions, metastatic tumor growth, migraine, mucus secretion disorders, nasal congestion, nasal inflammation, occlusive vascular diseases, ocular hypertension, ocular hypotension, osteoporosis, pre-term labor rheumatoid arthritis , pain, perennial rhinitis, pulmonary congestion, pulmonary hypotension, Raynaud's disease, rejection in organ transplant and by-pass surgery, respiratory conditions, hirsutism , rhinorrhea, shock, sleep disorders, and sleep-wake cycle disorders.

30. The method of paragraph 27 wherein said compound is administered as a surgical adjunct in ophthalmology for cataract removal and artificial lens insertion, ocular implant procedures, photorefractive radial keratotomy and other ophthalmological laser procedures.

31. The method of paragraph 27 wherein said compound is administered as a surgical adjunct in a procedure involving skin incisions, relief of pain and inflammation and scar formation/keloids post-surgery, for treating sports injuries and general aches and pains in muscles and joints.

32. The method of paragraph 27 wherein said DPi, FP, EPi, TP , and/or EP₄ receptor mediated condition or disease is an EPi and/or EP₄ receptor mediated condition or disease.

33. The method of paragraph 27 wherein said DPi, FP, EPi, TP and/or EP₄ receptor mediated condition or disease is an allergic condition.

34. The method of paragraph 33 wherein said condition is dermatological allergy.

35. The method of paragraph 27 wherein said condition is an ocular allergy.

36. The method of paragraph 27 wherein said condition is a respiratory allergy.

37. The method of paragraph 27 wherein said condition or disease is selected from the group consisting of nasal congestion, rhinitis, and asthma.

38. The method of paragraph 27 wherein said condition or disease is related to pain.

39. The method of paragraph 27 wherein said condition or disease is selected from the group consisting of arthritis, migraine, and headache.

40. The method of paragraph 27 wherein said condition or disease is associated with the gastrointestinal tract.

41. The method of paragraph 27 wherein said condition or disease is selected from the group consisting of peptic ulcer, heartburn, reflux esophagitis, erosive esophagitis, non-ulcer dyspepsia, infection by *Helicobacter pylori*, alrinitis, and irritable bowel syndrome.

42. The method of paragraph 27 wherein said condition or disease is selected from the group consisting of hyperalgesia and allodynia.

43. The method of paragraph 27 wherein said condition or disease is related to mucus secretion.

44. The method of paragraph 27 wherein said mucus secretion is gastrointestinal.

45. The method of paragraph 27 wherein said mucus secretion occurs in the nose, sinuses, throat, or lungs.

5

46. The method of paragraph 27 wherein said condition or disease is related to abdominal cramping.

47. The method of paragraph 27 wherein said condition or disease is irritable bowel syndrome.

10

48. The method of paragraph 27 wherein said condition or disease is a bleeding disorder.

49. The method of paragraph 27 wherein said condition or disease is a sleep disorder.

15

50. The method of paragraph 27 wherein said condition or disease is mastocytosis.

51. The method of paragraph 27 wherein said condition or disease is associated with elevated body temperature.

20

52. The method of paragraph 27 wherein said condition or disease is associated with ocular hypertension and glaucoma.

53. The method of paragraph 27 wherein said condition or disease is associated with ocular hypotension.

25

54. The method of paragraph 27 wherein said condition relates to surgical procedures to treat pain, inflammation and other unwanted sequelae wherein said surgical procedure includes incision , laser surgery or implantation.


30

55. The method of paragraph 27 where said condition is related to pain and inflammation and post-surgical scar and keloid formation.

56. The method of paragraph 27 where said condition is related to diseases of female reproduction, associated with menstrual cramping, endometriosis, and pre-term labor

57 A pharmaceutical product comprising a compound having the following formula

5



Y is $(CH_2)_m$ wherein m is 0 or an integer of from 1 to 3;

10

Z is selected from the group consisting of O, S, SO, SO₂ and $(CH_2)_p$, wherein p is 0 or an integer of from 1 to 3;

W is substituted phenyl ring, alkyl, substituted alkyl, cycloalkyl or substituted cycloalkyl :

15

R_i is selected from the group consisting of OR₇, N(R₇)₂, and N(R₇)SO₂R₇;

R₂ is H ;

20 R₄ is selected from the group consisting of alkyl, halogen-substituted alkyl and amino ;

R₅ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxy, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy;


25

Re is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxy, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy, wherein at least one of R_5 and R_6 are halogen; and

- 5 R_7 is selected from the group consisting of H, hydrocarbyl and substituted hydrocarbyl.

or a pharmaceutically acceptable salt or a prodrug thereof, wherein said product is packaged and labeled for the treatment or prevention of a disease or condition selected from the group consisting of uveitis, allergic conditions, asthma, allergic asthma, allergic rhinitis, atherosclerosis, blood coagulation disorders, bone disorders, cancer, cellular neoplastic transformations, chronic obstructive pulmonary diseases and other forms of lung inflammation, congestive heart failure, diabetic retinopathy, diseases or conditions requiring a treatment of anti-coagulation, diseases requiring control of bone formation and resorption, endometriosis fertility disorders, hyperpyrexia, gangrene, glaucoma, hypothermia, immune and autoimmune diseases, inflammatory conditions, menstrual cramping, metastatic tumor growth, migraine, mucus secretion disorders, nasal congestion, nasal inflammation, occlusive vascular diseases, ocular hypertension, ocular hypotension, osteoporosis, pain, perennial rhinitis, pre-term labor pulmonary congestion, pulmonary hypotension, Raynaud's disease, rejection in organ transplant and by-pass surgery, respiratory conditions, rheumatoid arthritis, rhinorrhea, shock, sleep disorders, sleep-wake cycle disorders, sports injuries, muscle aches and pains, and surgical adjunct for minimizing pain, inflammation and scar/keloid formation.

58. A pharmaceutical composition comprising a compound having the following formula



Y is $(\text{CH}_2)_m$ wherein m is 0 or an integer of from 1 to 3;

5 Z is selected from the group consisting of O, S, SO, SO_2 and $(\text{CH}_2)_p$, wherein p is 0 or an integer of from 1 to 3;

W is substituted phenyl ring, alkyl, substituted alkyl, cycloalkyl or substituted cycloalkyl;

10 Ri is selected from the group consisting of OR_7 , $\text{N}(\text{R}_7)_2$, and $\text{N}(\text{R}_7)\text{SO}_2\text{R}_7$;

R_2 is H ;

R_4 is selected from the group consisting of alkyl, halogen-substituted alkyl and amino;

15 R_5 is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxy, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy;

20 R_6 is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxy, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy, wherein at least one of R_5 and R_6 are halogen;and

DETAILED DESCRIPTION OF THE INVENTION

25 The following terms are used to define the disclosed invention.

"Hydrocarbyl" refers to a hydrocarbon radical having only carbon and hydrogen atoms. Preferably, the hydrocarbyl radical has from 1 to 20 carbon atoms, more preferably from 1 to 12 carbon atoms and most preferably from 1 to 7 carbon atoms.

30

"Substituted hydrocarbyl" refers to a hydrocarbyl radical wherein one or more, but not all, of the hydrogen and/or the carbon atoms are replaced by a halogen, nitrogen, oxygen, sulfur or

phosphorus atom or a radical including a halogen, nitrogen, oxygen, sulfur or phosphorus atom, e.g. fluoro, chloro, cyano, nitro, hydroxyl, phosphate, thiol, etc."

"Alkyl" refers to a straight-chain, branched or cyclic saturated aliphatic hydrocarbon.

- 5 Preferably, the alkyl group has 1 to 12 carbons. More preferably, it is an alkyl of from 4 to 10 carbons, most preferably 4 to 8 carbons. Typical alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tertiary butyl, pentyl, hexyl and the like. The alkyl group may be optionally substituted with one or more substituents selected from the group consisting of hydroxyl, cyano, alkoxy, =O, =S, NO₂, halogen, dimethyl amino, and SH.

10

"Cycloalkyl" refers to a cyclic saturated aliphatic hydrocarbon group. Preferably, the cycloalkyl group has 3 to 12 carbons. More preferably, it has from 4 to 7 carbons, most preferably 5 or 6 carbons.

- 15 "Substituted cycloalkyl" refers to a cycloalkyl radical wherein one or more, but not all, of the hydrogen and/or the carbon atoms are replaced by a halogen, nitrogen, oxygen, sulfur or phosphorus atom or a radical including a halogen, nitrogen, oxygen, sulfur or phosphorus atom, e.g. fluoro, chloro, cyano, nitro, hydroxyl, phosphate, thiol, etc. Specific substituted cycloalkyls, i.e. cyclopentyl, may be referred to as substituted cyclopentyl with the
20 understanding that the substituents are the same."

"Aryl" refers to an aromatic group which has at least one ring having a conjugated pi electron system and includes carbocyclic aryl, heterocyclic aryl and biaryl groups. The aryl group may be optionally substituted with one or more substituents selected from the group
25 consisting of alkyl, hydroxyl, halogen, COOR⁶, NO₂, CF₃, N(R⁶)₂, CON(R⁶)₂, SR⁶, sulfoxy, sulfone, CN and OR⁶, wherein R⁶ is alkyl.


"Carbocyclic aryl" refers to an aryl group wherein the ring atoms are carbon.

- 30 "Heteroaryl or heterocyclic aryl" refers to an aryl group having from 1 to 3 heteroatoms as ring atoms, the remainder of the ring atoms being carbon. Heteroatoms include oxygen, sulfur, and nitrogen. Thus, heterocyclic aryl groups include furanyl, thienyl, pyridyl, pyrrolyl, N-lower alkyl pyrrolo, pyrimidyl, pyrazinyl, imidazolyl and the like. Preferably, the

heteroaryl group has from 2 to 10 carbons. More preferably, it has from 3 to 10 carbons, most preferably 3 carbons.

The present invention provides compounds having the general formula I:

5



Y is $(CH_2)_m$ wherein m is 0 or an integer of from 1 to 3;

10 Z is selected from the group consisting of O, S, SO, SO₂ and $(CH_2)_p$, wherein p is 0 or an integer of from 1 to 3;

W is substituted phenyl ring, alkyl, substituted alkyl, cycloalkyl or substituted cycloalkyl:

15 Ri is selected from the group consisting of OR₇, N(R₇)₂, and N(R₇)SO₂R₇;

R₂ is H ;

R₄ is selected from the group consisting of alkyl, halogen-substituted alkyl and amino;

20

R₅ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxyl, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy;

25 R₆ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxyl, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy, wherein at least one of R₅ and R₆ are halogen;and

Preferably, Ri is OH.

Preferably, Y is absent, i.e. n is 0.

5 Preferably, Z is O.

Preferably, W is selected from the group consisting of cyclopentyl, cyclobutyl, cyclohexyl and substituted cyclopentyl, cyclobutyl and cyclohexyl.

10 More preferably W is cyclopentyl and cyclobutyl.

The most preferred compounds of the present invention are selected from the group consisting of

15

The compounds of the present invention may be prepared by the methods disclosed in the Examples.

The following examples are intended to illustrate the present invention.

20

The reagents and conditions used in the Examples may be abbreviated as follows:

Ac is acetyl;

25 DCM is dichloromethane;

DTAD is Di-tert-butyl azodicarboxylate;

TFA is trifluoroacetic acid;

30

RT is room temperature;

Ph is phenyl;

DiBAL-H is diisobutylaluminumhydride;

DMF is dimethylformamide;

5 Et is ethyl;

THF is tetrahydrofuran;

DMAP is 4-dimethylaminopyridine;


10

HEPES is 4- (2-hydroxyethyl)-1 -piperazineethanesulfonic acid).

EXAMPLE 1

15


1-(5-CHLORO-2-CYCLOPENTYLMETHOXY-BENZ YL)-5-METHYL- 1H-PYRAZOLE-3-CARBOXYLIC ACID, 4



STEP 1

20

N'-(5-chloro-2-hydroxy-benzyl)-hydrazinecarboxylic acid tert-butyl ester, **1**




A solution of 5-chloro-2-hydroxybenzaldehyde (1.5 g 9.3 mmol), tert-butylcarbazate (1.25 g, 9.3 mmol) and acetic acid (0.54 mL, 9.3 mmol) in CH_2Cl_2 (50 mL) was stirred under a nitrogen atmosphere for 30 min at RT. Then sodium triacetoxyborohydride (6.20 g, 27.9 mmol) was added portion wise and the resulting mixture was stirred at RT overnight. The reaction was thoroughly quenched with 2 M HCl (15 mL) and stirred at RT for 1 h. The reaction mixture was partitioned between water (50 mL) and CH_2Cl_2 (25 mL). The aqueous layer was extracted with CH_2Cl_2 (25 mL). The combined organic layers were washed with water (2 x 75 mL), dried (Na_2SO_4) and evaporated to dryness to give hydrazine **1** as a white solid.

$^1\text{H-NMR}$ (CDCl_3 , 300 MHz) 57.17 (dd, 1H, ArH), 7.03 (d, 1H, ArH), 6.83 (d, 1H, ArH), 1.49 (s, 9H, $-\text{C}(\text{CH}_3)_3$).
LC-MS: m/z 273 $\text{M} + \text{H}^+$

STEP 2

1-(5-chloro-2-hydroxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **2**




A suspension of N'-(5-chloro-2-hydroxy-benzyl)-hydrazinecarboxylic acid tert-butyl ester **1** (2.5g, 9.3 mmol) in CH_2Cl_2 was treated with TFA (20 mL) and stirred at RT overnight. The volatiles were removed *in vacuo*. The residue was dissolved in AcOH (20 mL) and slowly added to a solution of ethyl-2,4-dioxopentanoate in AcOH (10 mL). The resulting mixture was refluxed for 1 h and let cool down and stirred at RT for 16 h. Precipitated 1-(5-Chloro-2-hydroxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester **2** was filtered and washed with ether. The white solid was dried overnight in a dessicator yielding 1-(5-chloro-2-hydroxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester **2**.

$^1\text{H-NMR}$ (CDCl_3 , 300 MHz) 510.16 (s, 1H, ArOH), 7.17 (dd, 1H, ArH), 6.86 (d, 1H, ArH), 6.65 (d, 1H, ArH), 6.58 (s, 1H, ArH), 5.24 (s, 2H, ArCH_2), 4.24 (q, 2H, $-\text{CH}_2\text{CH}_3$), 2.28 (s, 3H, CH_3), 1.26 (t, 3H, $-\text{CH}_2\text{CH}_3$). LC-MS: m/z 295 $\text{M} + \text{H}^+$

STEP 3

1-(5-chloro-2-cyclopentylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **3**.




A solution of 1-(5-chloro-2-hydroxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **2**, (0.15g, 0.51 mmol), polymer supported triphenylphosphine (0.2g, 0.56 mmol), diethylazodicarboxylate (O.lg, 0.56 mmol) and cyclopentanemethanol (0.056 ml, 0.56 mmol) in a mixture of THF (5 mL) was refluxed for 48 hours. Solid support reagent was removed by filtration and the volatiles were removed *in vacuo*. The crude product was purified on silica to yield 1-(5-chloro-2-hydroxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **3a** pale yellow oil, 0.086 g.

¹H-NMR (CDCl₃, 300 MHz) δ 7.18 (dd, 1H, ArH) , 6.79 (d, 1H, ArH), 6.67 (s, 1H, ArH), 6.53 (dd, 1H, ArH) , 5.37 (s, 2H, ArCH₂) , 4.43 (q, 2H, -CH₂CH₃) , 3.88 (dd, 2H, O-CH₂) , 2.40 (m, 1H OCH₂CH) , 2.22 (s, 3H, CH₃) , 1.94-1.22 (m, 8H, CH₂) , 1.42 (t, 3H, -CH₂CH₃) .
LC-MS: m/z 377 M + H⁺

STEP 4

1-(5-chloro-2-cyclopentylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid, **4**.



To a solution of ester **3** (0.086g, 0.25 mmol) in MeOH (10 mL) was added a solution of 2 M NaOH 2 mL (0.40 mmol) and the resulting mixture was stirred at RT for 4 hours. The


volatiles were removed in vacuo. The residue was diluted with water (5 mL) and acidified to pH 1 with 2 M HCl. The acid 4 was isolated by filtration as a white solid and washed with water and dried overnight over KOH in a dessicator to yield 1-(5-chloro-2-cyclopentylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid, 4, as a white solid.

¹H-NMR(CDC1₃, 300 MHz) δ 7.20 (dd, 1H, ArH), 6.81 (d, 1H, ArH), 6.71 (s, 1H, ArH), 6.63 (dd, 1H, ArH), 5.36 (s, 2H, ArCH₂), 3.88 (dd, 2H, O-CH₂), 2.40 (m, 1H OCH₂CH), 2.26 (s, 3H, CH₃), 1.94-1.22 (m, 8H, CH₂).

LC-MS: m/z 349 M + H⁺


EXAMPLE 2

1-(5-BROMO-2-CYCLOPENTYLMETHOXY-BENZYL)-5-METHYL-1H-PYRAZOLE-3-CARBOXYLIC ACID, 8.



STEP 1

N'-(5-bromo-2-hydroxy-benzyl)-hydrazinecarboxylic acid tert-butyl ester, 5.




The title compound was prepared from 5-bromosalicylaldehyde following the method in Example 1, Step 1.

¹H-NMR(CDC1₃, 300 MHz) 57.31 (dd, 1H, ArH), 7.17 (d, 1H, ArH), 6.79 (d, 1H, ArH), 1.49 (s, 9H, -C(CH₃)₃).

LC-MS: m/z 318 M + H⁺

STEP 2

1-(5-bromo-2-hydroxy-benzyl)-5-methyl-1h-pyrazole-3-carboxylic acid ethyl ester, **6**



5


The title compound was prepared from compound **5** following the method in Example 1, Step 2.

10 $^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ 9.51 (s, 1H, ArOH), 7.33 (dd, 1H, ArH), 7.22 (d, 1H, ArH), 6.87 (d, 1H, ArH), 6.59 (s, 1H, ArH), 5.20 (s, 2H, ArCH_2), 4.38 (q, 2H, $-\text{CH}_2\text{CH}_3$), 2.42 (s, 3H, CH_3), 1.40 (t, 3H, $-\text{CH}_2\text{CH}_3$). LC-MS: m/z 340 $\text{M} + \text{H}^+$

15

STEP 3

1-(5-bromo-2-cyclopentylmethoxy-benzyl)-5-methyl-1h-pyrazole-3-carboxylic acid ethyl ester, **7**.




20 The title compound was prepared from compound **6** following the method in Example 1, Step 3.

25 $^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ 7.24 (dd, 1H, ArH), 6.66 (d, 1H, ArH), 6.61 (dd, 1H, ArH), 6.57 (s, 1H, ArH), 5.23 (s, 2H, ArCH_2), 4.34 (q, 2H, $-\text{CH}_2\text{CH}_3$), 3.79 (dd, 2H, $\text{O}-\text{CH}_2$), 2.31 (m, 1H, OCH_2CH_2), 2.13 (s, 3H, CH_3), 1.83-1.11 (m, 8H, CH_2), 1.33 (t, 3H, CH_2CH_3). LC-MS: m/z 422 $\text{M} + \text{H}^+$

30

STEP 4

1-(5-bromo-2-cyclopentylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid, **8**.



5

The title compound was prepared from compound **7** following the method in Example 1, Step 4.


¹H-NMR (CDCl₃, 300 MHz) δ 7.35 (dd, 1H, **ArH**), 6.80 (d, 1H, **ArH**), 6.77 (dd, 1H, **ArH**), 6.70 (s, 1H, **ArH**), 5.36 (s, 2H, **ArCH₂**), 3.88 (dd, 2H, **O-CH₂**), 2.40 (m, 1H **OCH₂CH**), 2.27 (s, 3H, **CH₃**), 1.94-1.10 (m, 8H, **CH₂**).
LC-MS: m/z 394 M + H⁺

15

EXAMPLE 3

1-(2-CYCLOPENTYLMETHOXY-5-TRIFLUOROMETHYL-BENZYL)-5-METHYL-1H-PYRAZOLE-3-CARBOXYLIC ACID, 12.


20



25

STEP 1

N'-(2-hydroxy-5-trifluoromethyl-benzyl)-hydrazinecarboxylic acid tert-butyl ester, **9**.




The title compound was prepared from 5-trifluoromethylaldehyde following the method in Example 1, Step 1.

¹H-NMR(CDC1₃, 300 MHz) 5.971 (broad s, 1H, OH), 7.49 (dd, 1H, ArH), 7.32 (s, 1H, ArH), 6.97 (d, 1H, ArH), 6.15 (broad s, 1H, NH), 4.43 (broad s, 1H, NH), 1.50 (s, 9H, -C(CH₃)₃).
LC-MS: m/z 307 M + H⁺

STEP 2

1-(2-hydroxy-5-(trifluoromethyl)benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **10**.




The title compound was prepared from compound **9** following the method in Example 1, Step 2.

¹H-NMR(CDC1₃, 300 MHz) δ 7.52 (dd, 1H, ArH), 7.40 (d, 1H, ArH), 7.08 (d, 1H, ArH), 6.60 (s, 1H, ArH), 5.27 (s, 2H, ArCH₂), 4.39 (q, 2H, -CH₂CH₃), 2.44 (s, 3H, CH₃), 1.40 (t, 3H, -CH₂CH₃). LC-MS: m/z 329 M + H⁺

STEP 3

1-(2-cyclopentylmethoxy-5-trifluoromethyl-benzyl)-5-methyl-1h-pyrazole-3-carboxylic acid ethyl ester, **11**.




The title compound was prepared from compound **10** following the method in Example 1, Step 3.

$^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ 7.51 (dd, 1H, **ArH**), 6.94 (dd, 1H, **ArH**), 6.85 (d, 1H, **ArH**), 6.67 (s, 1H, **ArH**), 5.42 (s, 2H, **ArCH₂**), 4.43 (q, 2H, **-CH₂CH₃**), 3.96 (dd, 2H, **O-CH₂**), 2.43 (m, 1H, **OCH₂CH**), 2.23 (s, 3H, **CH₃**), 1.97-1.05 (m, 8H, **CH₂**), 1.42 (t, 3H, **CH₂CH₃**).
LC-MS: m/z 411 $M + H^+$

STEP 4

1-(2-cyclopentylmethoxy-5-trifluoromethyl-benzyl)-5-methyl-1h-pyrazole-3-carboxylic acid, **12**.




The title compound was prepared from compound **11** following the method in Example 1, Step 4.

¹H-NMR(CDC1₃, 300 MHz) δ 7.53 (dd, 1H, ArH) , 6.96 (dd, 1H, ArH), 6.94 (d 1H, ArH), 6.72 (s, 1H, ArH), 5.40 (s, 2H, ArCH₂), 3.96 (dd, 2H, O-CH₂), 2.43 (m, 1H OCH₂CH), 2.27 (s, 3H, CH₃), 1.94-1.21 (m, 8H, CH₂).

5 LC-MS: m/z 383 M + H⁺

EXAMPLE 4

10 N-{1-[5-CHLORO-2-(4-METHOXY-BENZYLOXY)-BENZYL]-5-METHYL-1H-PYRAZOLE-3-CARBONYL}-BENZENESULFONAMIDE, 16.




15

STEP 1

1-[5-chloro-2-(4-methoxy-benzyloxy)-benzyl]-5-methyl-1h-pyrazole-3-carboxylic acid ethyl ester, 13.

20




To a solution of 1-(5-chloro-2-hydroxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester **2** 0.2g (0.6 mmol) in DMF (5 mL) were added potassium carbonate (0.23 g, 1.7 mmol), potassium iodide (0.1 g, 0.6 mmol) and 4-methoxybenzyl chloride (0.1 ml, 0.7 mmol). The resulting mixture was heated at 100 °C in an Emrys microwave reactor for 30 minutes. The volatiles were removed *in vacuo*. The crude product was purified on silica to yield 1-[5-chloro-2-(4-methoxy-benzyloxy)-benzyl]-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester **13** as a white solid.

¹H-NMR(CDCl₃, 300 MHz) δ 7.32 (d, 2H, ArH), 7.20 (dd, 1H, ArH), 6.94 (d, 2H, ArH), 6.88 (d, 1H, ArH), 6.64 (s, 1H, ArH), 6.60 (d, 1H, ArH), 5.35 (s, 2H, ArCH₂), 5.03 (s, 2H, ArCH₂), 4.42 (q, 2H, -CH₂CH₃), 3.85 (3H, OCH₃), 2.15 (s, 3H, CH₃), 1.41 (t, 3H, -CH₂CH₃). LC-MS: m/z 415 M + H⁺

STEP 2

1-[5-chloro-2-(4-methoxy-benzyloxy)-benzyl]-5-methyl-1H-pyrazole-3-carboxylic acid, **14**.




To a solution of 1-[5-chloro-2-(4-methoxy-benzyloxy)-benzyl]-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **13**, 0.28g (0.6 mmol) in EtOH (2 mL) was added a solution of LiOH (0.15g in 0.5 ml H₂O). The resulting mixture was heated at 100 °C on microwave for 10 minutes. The mixture was poured into water (20 mL) and extracted with EtOAc (3 x 15 mL). The organic layers were combined, washed with brine (30 mL), dried (MgSO₄) and the volatiles removed *in vacuo* to yield 1-[5-chloro-2-(4-methoxy-benzyloxy)-benzyl]-5-methyl-1H-pyrazole-3-carboxylic acid, **14** as a white solid.

¹H-NMR(CDCl₃, 300 MHz) δ 7.32 (d, 2H, ArH), 7.20 (dd, 1H, ArH), 6.94 (d, 2H, ArH), 6.88 (d, 1H, ArH), 6.62 (s, 1H, ArH), 6.60 (d, 1H, ArH), 5.35 (s, 2H, ArCH₂), 5.03 (s, 2H, ArCH₂), 3.85 (3H, OCH₃), 2.16 (s, 3H, CH₃). LC-MS: m/z 387 M + H⁺

STEP 3

1-[5-chloro-2-(4-methoxy-benzyloxy)-benzyl]-5-methyl-1H-pyrazole-3-carbonyl fluoride, **15**.



5

To a solution of acid 5-chloro-2-(4-methoxy-benzyloxy)-benzyl]-5-methyl-1H-pyrazole-3-carboxylic acid, **14**, 0.2g (0.52 mmol) in dry THF under N₂ atm was added 500 μL of pyridine (6.2 mmol) and 440 μL (5.2 mmol) of cyanuric fluoride. The mixture was refluxed


10 After drying over MgSCl solvents were removed in vacuo to yield the crude acid fluoride, **15** which was used in sequential step without further purification.

¹H-NMR (CDCl₃, 300 MHz) δ 7.31 (d, 2H, ArH), 7.24 (dd, 1H, ArH), 6.94 (d, 2H, ArH), 6.91 (d, 1H, ArH), 6.76 (d, 1H, ArH), 6.70 (s, 1H, ArH), 5.35 (s, 2H, ArCH₂), 5.02 (s, 2H, ArCH₂), 3.85 (3H, OCH₃), 2.17 (s, 3H, CH₃).

¹⁹F-NMR (CDCl₃, 300 MHz) δ +21.8.

STEP 4

5 n-{1-[5-chloro-2-(4-methoxy-benzyloxy)-benzyl]-5-methyl-1H-pyrazole-3-carbonyl} - benzenesulfonamide, **16**.




10 To a solution of 1-[5-chloro-2-(4-methoxy-benzyloxy)-benzyl]-5-methyl-1H-pyrazole-3-carbonyl fluoride, **15**, 0.06g (0.18 mmol) and K_2CO_3 0.100g (0.72 mmol) in dry DCM (2 mL), benzenesulfonamide 0.055g (0.35mmol) was added. The mixture was stirred under a nitrogen atmosphere for 16 hours before diluting with EtOAc. The organic phase was washed with 2M HCl, followed by brine, dried over $MgSO_4$ and evaporated to dryness. The crude
 15 acyl sulphonamide was purified on silica to yield N-{1-[5-chloro-2-(4-methoxy-benzyloxy)-benzyl]-5-methyl-1H-pyrazole-3-carbonyl} - benzenesulfonamide, **16**.

1H -NMR ($CDCl_3$, 300 MHz) δ 8.17 (dd, 2H, ArH,), 7.96 (dd, 2H, ArH,), 7.70-7.49 (m, 3H, ArH), 7.27 (m, 2H, ArH,), 6.92 (m, 2H, ArH), 6.73 (d, 1H, ArH), 6.54 (s, 1H, ArH), 5.19 (s, 2H, ArCH₂), 5.00 (s, 2H, ArCH₂), 4.81 (broad s, 1H, NH), 3.85 (3H, OCH₃), 2.10 (s, 3H, CH₃).
 20 LC-MS : m/z 527 M + H⁺

25

EXAMPLE 5

N-{1-[5-CHLORO-2-(4-CHLORO-BENZYLOXY)-BENZYL]-5-METHYL-1H-PYRAZOLE-3-CARBONYL}-BENZENESULFONAMIDE, 17.




The title compound was prepared following the method described in example 4 but using the appropriate reagents.

¹H-NMR (DMSO, 300 MHz) δ 8.17 (dd, 2H, ArH), 7.96 (dd, 2H, ArH), 7.70-7.49 (m, 3H, ArH), 7.27 (m, 2H, ArH), 6.92 (m, 2H, ArH), 6.73 (d, 1H, ArH), 6.54 (s, 1H, ArH), 5.22 (s, 2H, ArCH₂), 5.19 (s, 2H, ArCH₂), 2.08 (s, 3H, CH₃).

LC-MS: m/z 530 M + H⁺


EXAMPLE 6

1-[5-BROMO-2-(2-ETHYL-BUTOXY)-BENZYL]-5-METHYL-1H-PYRAZOLE-3-CARBOXYLIC ACID, 19



STEP 1

l-[5-bromo-2-(2-ethyl-butoxy)-benzyl]-5-methyl-lh-pyrazole-3-carboxylic acid ethyl ester, **18**.




The title compound was prepared from compound **6** following the method in Example 1, Step 3.

^1H -NMR (CDCl_3 , 300 MHz) δ 7.34 (dd, 1H, ArH), 6.77 (d, 1H, ArH), 6.67 (d, 1H, ArH), 6.64 (d, 1H, ArH), 5.38 (s, 2H, ArCH₂), 4.43 (q, 2H, -CH₂CH₃), 3.91 (dd, 2H, O-CH₂), 2.21 (s, 3H, CH₃), 1.72 (m, 1H, CH₂CH), 1.58-1.43 (m, 4H, CH₂), 1.42 (t, 3H, CH₂CH₃), 0.96 (t, 6H, CH₂CH₃).

LC-MS: m/z 424 M + H⁺

STEP 2

l-[5-bromo-2-(2-ethyl-butoxy)-benzyl]-5-methyl-lh-pyrazole-3-carboxylic acid, **19**.




The title compound was prepared from compound **18** following the method in Example 4, Step 2.

$^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ 7.36 (dd, 1H, ArH), 6.78 (dd, 1H, ArH), 6.74 (d, 1H, ArH), 6.71 (d, 1H, ArH), 5.53 (s, 2H, ArCH_2), 3.91 (dd, 2H, O-CH_2), 2.25 (s, 3H, CH_3), 1.72 (m, 1H, CH_2CH), 1.57-1.42 (m, 4H, CH_2), 0.96 (t, 6H, CH_2CH_3).

LC-MS: m/z 396 $\text{M} + \text{H}^+$


EXAMPLE 7

1-[5-BROMO-2-(2-ETHYL-2-METHYL-BUTOXY)-BENZYL]-5-METHYL-1H-PYRAZOLE-3-CARBOXYLIC ACID, 23.



STEP 1

2-ethyl-2-methyl-butyric acid ethyl ester, 20.



Diisopropylamine (1.13 mL, 8.05mmol) was dissolved in THF (1ml) under nitrogen. The mixture was cooled to -78°C and $n\text{-Buli}$ (3.08 mL, 7.7mmol) was added dropwise to form LDA. A solution of 2-ethyl-butyric acid ethyl ester (0.57 mL, 3.5 mmol) in THF (2ml) was added dropwise to this solution. The resulting mixture was stirred at -78°C under nitrogen for 30 minutes before iodomethane (0.5 mL, 8.05mmol) was added dropwise and the mixture was first stirred for 1 hour at -78°C , and then 30 minutes at 0°C , and finally 30 minutes at room temperature. The reaction mixture was quenched with H_2O (5ml). The mixture was extracted twice with Et_2O (5ml), washed with 2M HCl (5ml) then dried over MgSO_4 .


$^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ 4.14 (q, 2H, CH_2CH_3), 1.78-1.60 (m, 2H, CH_2CH_3), 1.54-1.36 (m, 2H, CH_2CH_3), 1.27 (t, 3H, CH_2CH_3), 1.11 (s, 3H, CCH_3), 0.84 (t, 6H, CH_2CH_3).

LC-MS : m/z 159 M + H⁺

STEP 2

5

2-ethyl-2-methyl-1-butan-1-ol, **21**.




10 To a solution of 2-ethyl-2-methyl-butyric acid ethyl ester **20** (0.23 mL, 1.26mmol) in dry toluene (5 ml) under nitrogen was added DIBAL-H (12.6 mL, 14.5mmol) dropwise. The resulting mixture was stirred at room temperature for 3.5 hours. Aqueous Rochelle salts solution (10%, 15ml) was added to the reaction mixture at 0°C. After 10 minutes at 0°C it was stirred at room temperature for 40 minutes. The mixture was diluted with EtOAc (15ml)
 15 and washed with 2M HCl (15ml), saturated ammonium chloride (15ml) and brine (15ml) then dried over MgSC[^]. The volatiles were removed *in vacuo* to yield 2-ethyl-2-methyl-butan-1-ol **21**, which was used in the next step without further purification.

20 ¹H-NMR (CDCl₃, 300 MHz) δ 3.38 (broad s, 2H, CH₂OH), 1.39-1.22 (m, 4H, CH₂CH₃), 1.18 (broad s, CH₂OH), 0.87 (s, 3H, CCH₃), 0.83 (t, 6H, CH₂CH₃).
 LC-MS : m/z 117 M + H⁺

STEP 3

25

1-[5-bromo-2-(2-ethyl-2-methyl-butoxy)-benzyl]-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **22**.



30

A solution of 1-(5-bromo-2-hydroxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **6** (0.20g, 0.60 mmol), triphenylphosphine (0.31g, 1.2 mmol), di-tert-butylazodicarboxylate (0.27g, 1.2 mmol) and 2-ethyl-2-methyl-butan-1-ol **20** (0.13g, 1.2 mmol) in a mixture of THF (8 mL) was heated at 100 °C in an Emrys microwave reactor for

20 minutes. The volatiles were removed in vacuo and the crude product was purified on silica to yield 1-[5-bromo-2-(2-ethyl-2-methyl-butoxy)-benzyl]-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **22**.


¹H-NMR (CDCl₃, 300 MHz) δ 7.32 (dd, 1H, ArH), 6.77 (d, 1H, ArH), 6.67 (s, 1H, ArH), 6.57 (d, 1H, ArH), 5.39 (s, 2H, ArCH₂), 4.42 (q, 2H, -CH₂CH₃), 3.68 (s, 2H, OCH₂), 2.20 (s, 3H, CH₃), 1.50-1.37 (m, 4H, CCH₂), 1.42 (s, 3H, CCH₃), 0.86 (t, 6H, CH₂CH₃).
LC-MS: m/z 438 M + H⁺

10

STEP 4

1-[5-bromo-2-(2-ethyl-2-methyl-butoxy)-benzyl]-5-methyl-1H-pyrazole-3-carboxylic acid, **23**.

15



To a solution of 1-[5-bromo-2-(2-ethyl-2-methyl-butoxy)-benzyl]-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **22**, (0.17g, 0.38mmol) in a mixture of MeOH (3 mL) and THF (6 ml) was added a solution of LiOH in 3 ml of H₂O. The resulting mixture was heated at 100 °C in an Emrys microwave reactor for 20 minutes. The mixture was poured into water (20 mL) and extracted with EtOAc (3 x 15 mL). The organic layers were combined, washed with brine (30 mL), dried (**MgSO₄**) and the volatiles were removed in vacuo to yield 1-[5-bromo-2-(2-ethyl-2-methyl-butoxy)-benzyl]-5-methyl-1H-pyrazole-3-carboxylic acid, **23**.

20


25

¹H-NMR (CDCl₃, 300 MHz) δ 7.34 (dd, 1H, ArH), 6.78 (d, 1H, ArH), 6.73 (s, 1H, ArH), 6.65 (d, 1H, ArH), 5.41 (s, 2H, ArCH₂), 3.69 (s, 2H, OCH₂), 2.22 (s, 3H, CH₃), 1.45 (q, 4H, CCH₂), 0.97 (s, 3H, CCH₃), 0.86 (t, 6H, CH₂CH₃).
LC-MS: m/z 410 M + H⁺


30

EXAMPLE 8**1-(5-BROMO-2-CYCLOBUTYLMETHOXY-BENZYL)-5-METHYL-1H-PYRAZOLE-3-CARBOXYLIC ACID, 25**

5

**STEP 1**

10 1-(5-bromo-2-cyclobutylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, 24



15 A solution of 1-(5-bromo-2-hydroxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **6** (O.lg, 0.3 mmol), triphenylphosphine (0.155g, 0.6 mmol), di-tert-butylazodicarboxylate (0.14g, 0.6 mmol) and cyclobutanemethanol (0.06 ml, 0.6 mmol) in a mixture of THF (8 mL) was heated at 100 °C in an Emrys microwave reactor for 20 minutes. The volatiles were removed in vacuo and the crude product was purified on silica to yield 1-


20 [5-bromo-2-(2-ethyl-2-methyl-butoxy)-benzyl]-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **24**.

¹H-NMR (CDCl₃, 300 MHz) δ 7.32 (m, 1H, ArH), 6.70 (d, 2H, ArH), 6.67 (s, 1H, ArH), 5.39 (s, 2H, ArCH₂), 4.42 (q, 2H, -CH₂CH₃), 4.00 (s, 2H, OCH₂), 2.80 (m, 1H, CH), 2.20 (s, 3H, CH₃), 2.10-1.80 (m, 6H, CCH₂) 1.40 (t, 3H, OCH₂CH₃).

25

STEP 2

1-(5-bromo-2-cyclobutylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid, **25**



5

To a solution of 1-(5-bromo-2-cyclobutylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid ethyl ester, **24** (0.12g, 0.29mmol) in a mixture of MeOH (2 mL) and THF (4 mL) was added a solution of NaOH (0.061g, 1.45 mmol) in 2 mL of H₂O. The resulting mixture was heated at 100 °C in an Emrys microwave reactor for 20 minutes. The mixture was poured into water (20 mL) and extracted with EtOAc (3 x 15 mL). The organic layers were combined, washed with brine (30 mL), dried (MgSO₄) and the volatiles were removed in vacuum to yield 1-(5-bromo-2-cyclobutylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid, **25**


¹H-NMR (CDCl₃, 300 MHz) δ 7.32 (d, 1H, ArH), 6.70 (m, 2H, ArH), 6.67 (s, 1H, ArH), 5.32 (s, 2H, ArCH₂), 3.97 (d, 2H, OCH₂), 2.75 (m, 1H, CH), 2.20 (s, 3H, CH₃), 2.10-1.80 (m, 6H, CCH₂)
LC-MS: m/z 380 M + H⁺

20

EXAMPLE 9

1-[5-BROMO-2-(1-TRIFLUOROMETHYL-CYCLOBUTYLMETHOXY)-BENZYL]-5-METHYL-1H-PYRAZOLE-3-CARBOXYLIC ACID, 26

25



The title compound was prepared following the methods described in example 8 but using the appropriate reagents.

¹H-NMR (CDCl₃, 300 MHz) δ 7.39 (m, 1H, ArH), 6.80 (s, 1H, ArH), 6.70 (m, 2H, ArH), 5.39 (s, 2H, ArCH₂), 4.19 (s, 2H, OCH₂), 2.45 (m, 2H, -CH₂CH₂-), 2.20 (s, 3H, CH₃), 2.15 (m, 4H, CCH₂)

LC-MS: m/z 448 M + H⁺

The above compounds were tested for PG antagonist activity as follows using human recombinant prostanoid receptor (DPi, EPi₄, FP, IP and TP) stable cell lines:

In order to measure the response of G_s and G_i coupled prostanoid receptors as a Ca²⁺ signal, chimeric G protein cDNAs were used. Stable cell lines over-expressing human prostanoid DPi, EPi₄, FP, IP, and TP receptors were established as follows:

Briefly, human prostanoid DPi, EP₂, and EP₄ receptor cDNAs were co-transfected with chimeric G_{qs} cDNA containing a haemagglutinin (HA) epitope; human prostanoid EP₃ receptors were co-transfected with chimeric G_{qi}-HA; human EPi, FP, IP, and TP receptor cDNAs were expressed with no exogenous G-proteins. G_{qs} and G_{qi} chimeric cDNAs (Molecular Devices, Sunnyvale, CA, U.S.A.), as well as cDNAs of prostanoid receptors, were cloned into a pCEP₄ vector with a hygromycin B selection marker. Transfection into HEK-293 EBNA (Epstein-Barr virus nuclear antigen) cells was achieved by the FuGENE 6 transfection Reagent (Roche Applied Science, Indianapolis, IN, USA). Stable transfectants were selected according to hygromycin resistance. Because G_{qs} and G_{qi} contained an HA epitope, G-protein expression was detected by Western blotting analysis using anti-mouse HA monoclonal antibody and horseradish peroxidase (HRP)-conjugated secondary antibody, while functional expression of prostanoid receptors was detected by FLIPR screening (Matias *et al.*, 2004). These stable cell lines were validated using previously published antagonists at 10 μM against serial dilutions of standard agonists by FLIPR functional assays for Ca²⁺ Signaling (as described below).

Ca²⁺ signaling studies were performed using a FLIPR TETRA system (Molecular Devices, Sunnyvale, CA, USA) in the 384-format. This is a high-throughput instrument for cell-based assays to monitor Ca²⁺ signaling associated with GPCRs and ion channels. Cells were seeded at a density of 5 x 10⁴ cells/well in BioCoat poly-D-lysine coated, black wall, clear bottom 384-well plates (BD Biosciences, Franklin lakes, NJ, USA) and allowed to attach overnight

in an incubator at 37°C. The cells were then washed twice with HBSS-HEPES buffer (Hanks' balanced salt solution without bicarbonate and phenol red, 20mM HEPES, pH 7.4) using an ELx405 Select CW Microplate Washer (BioTek, Winooski, VT, USA). After 60 min of dye-loading in the dark using the Ca^{2+} -sensitive dye Fluo-4AM (Invitrogen, Carlsbad, CA, USA), at a final concentration of $2 \times 10^{-6}\text{M}$, the plates were washed 4 times with HBSS-HEPES buffer to remove excess dye and leaving 50 μl of buffer in each well. The plates were then placed in the FLIPR TETRA instrument and allowed to equilibrate at 37°C. Compounds were added in a 25 μl volume to each well to give final concentrations of 0.1 μM , 0.3 μM , 1 μM , 3 μM , 10 μM , and 30 μM ; or 0.067 μM , 0.1 μM , 0.2 μM , 0.3 μM , 0.67 μM , and 1 μM for cells over-expressing TP receptors. After 4.5 minutes, a 7-point serial dilution of the standard agonist for the corresponding receptor, in a 25 μl volume was injected at the final concentrations from 10^{-11}M to 10^{-5}M in 10-fold serial dilution increments for cells expressing human recombinant **DPI**, **EPI**, **EP₂**, **EP₃**, **EP₄**, **FP**, and **IP** receptors. The dose range for the standard agonist for human recombinant TP receptors was from 10^{-12}M to 10^{-6}M . HBSS-HEPES buffer was used as the negative control for the standard agonists. Cells were excited with LED (light emitting diode) excitation at 470-495nm and emission was measured through an emission filter at 515-575nm. Assay plates were read for 3.5 minutes using the **FLIPR^{TE}TRA**. The peak increase in fluorescence intensity was recorded for each well. On each plate, negative controls, dose response of positive controls, and co-treatments of antagonist-agonist for each dose were in triplicates. Standard agonists were as follows: **DP** =BW 245C, **EPI-EP₄** =**PGE₂**, **FP** =17-phenyl-**PGF_{2 α}** , **IP** =Cicaprost, and **TP** =U-46619. The peak fluorescence change in each well containing drug was expressed relative to vehicle controls with the standard agonist at 10^{-6}M (the positive control). To obtain concentration-response curves, compounds were tested in triplicate in each plate over the desired concentration range.

DATA PROCESSING

All plates were subjected to appropriate baseline corrections. Maximum fluorescence values were exported. The raw data of $n=1$ was first processed by Activity Base using nonlinear regression curve fit to calculate the percentage activity of each data point relative to the positive control ($=10^{-6}\text{M}$ of the standard agonist). Then $n=3$ of this data were exported to GraphPad Prism 4 to calculate the average EC_{50} of the standard agonist, and the IC_{50} (the concentration of the antagonist required to inhibit half the standard agonist activity) were

calculated using nonlinear regression curve fit, with constraints of bottom constant equal to 0 and top constant equal to 100. Calculation of $K_b = [\text{Antagonist Concentration}] / (IC_{50}/EC_{50} - 1)$. When no antagonism was detected or when $K_b > 10,000\text{nM}$, the antagonist is defined as not active (NA).

5

The results of the above testing are reported in TABLE 1, below.

Compound No.	FP	DP	EP1	EP2	EP3	EP4	IP	TP
Example 1	150	800	30	NA	NA	800	NA	<1
Example 2	20	400	10	NA	NA	400	1300	<1
Example 3	80	700	5	8900	NA	700	4000	1
Example 4	400	800	10	4900	2500	140	900	2
Example 5	1700	2600	10	NA	1500	75	2000	10
Example 6	35	150	10	6200	NA	800	500	<1
Example 7	80	150	19	3600	NA	2070	1700	11
Example 8	8	130	10	NA	4100	700	5600	8
Example 9	30	190	16	3300	2600	600	NA	10

10 As shown in TABLE 1, the preferred compounds of this invention are pan antagonists which have activity at the FP, DPi, EPi, EP₄ and TP receptors, but are generally inactive at the IP, EP₂ and EP₃ receptors. Thus, these compounds have a biological selectivity profile making them useful in treating diseases and conditions which are ameliorated by the IP/EP₂ and/or EP₃ receptor stimulation, without the symptoms side effects mediated by the FP, DP, EPi, EP₄ and TP receptors. Also, based on the data generated for this TABLE 1, it appears that the cyclobutyl and cyclopentyl compounds are more active at the FP receptor than the corresponding phenyl compounds. Therefore, cycloalkyl compounds are preferred over carbocyclic aryl compounds. Further evidence for the preference of the cycloalkyl compounds of this invention over the corresponding carbocyclic aryl compounds is that the cycloalkyl compounds are essentially inactive at EP₂ and EP₃ receptors. Thus, these compounds have a biological selectivity profile making them useful in treating diseases and conditions which are ameliorated by the IP/EP₂ and/or EP₃ receptor stimulation, without the side effects mediated by the FP, DP, EP₁, EP₄ and TP receptors.

20

Thus, the compounds of this invention compound may be administered to treat DPi, FP, EPi, TP and/or EP₄ receptor mediated diseases or conditions.

5 For example, said condition or disease may be related to inflammation, or said , FP, EPi, TP and/or EP₄ receptor mediated condition or disease may be selected from the group consisting of allergic conditions, asthma, allergic asthma, allergic rhinitis, uveitis and related disorders , atherosclerosis, blood coagulation disorders, bone disorders, cancer, cellular neoplastic transformations, chronic obstructive pulmonary diseases and other forms of lung
10 inflammation, congestive heart failure, diabetic retinopathy, diseases or conditions requiring a treatment of anti-coagulation, diseases requiring control of bone formation and resorption, endometriosis, fertility disorders, gangrene, glaucoma, hyperpyrexia, immune and autoimmune diseases, inflammatory conditions, metastatic tumor growth, migraine, mucus secretion disorders, nasal congestion, nasal inflammation, occlusive vascular diseases, ocular
15 hypertension, ocular hypotension, osteoporosis, rheumatoid arthritis , pain, perennial rhinitis, pre-term labor, pulmonary congestion, pulmonary hypotension, Raynaud's disease, rejection in organ transplant and by-pass surgery, respiratory conditions, hirsutism, rhinorrhea, shock, sleep disorders, and sleep-wake cycle disorders.

20 Said compound may be administered as a surgical adjunct in ophthalmology for cataract removal and artificial lens insertion , ocular implant procedures, photorefractive radial keratotomy and other ophthalmological laser procedures or as a surgical adjunct in a procedure involving skin incisions, relief of pain and inflammation and scar formation/keloids post-surgery, for treating sports injuries and general aches and pains in muscles and joints.

25 Preferably, said DPi, FP, EPi, TP , and/or EP₄ receptor mediated condition or disease is an EPi and/or EP₄ receptor mediated condition or disease.

Preferably, saidDPi, FP, EPi , TP and/or EP₄ receptor mediated condition or disease is an
30 allergic condition, e.g. an dermatological allergy, or an ocular allergy, or a respiratory allergy, e.g. nasal congestion, rhinitis, and asthma.

Said condition or disease may be related to pain.

Said condition or disease may be selected from the group consisting of arthritis, migraine, and headache.

- 5 Said condition or disease may be associated with the gastrointestinal tract, wherein said condition or disease may be peptic ulcer, heartburn, reflux esophagitis, erosive esophagitis, non-ulcer dyspepsia, infection by *Helicobacter pylori*, alrnyitis, and irritable bowel syndrome.

- 10 Said condition or disease may be selected from the group consisting of hyperalgesia and allodynia, or said condition or disease may be related to mucus secretion, wherein said mucus secretion is gastrointestinal, or occurs in the nose, sinuses, throat, or lungs.

Said condition or disease is related to abdominal cramping, e.g. said condition, menstrual cramping or disease may be irritable bowel syndrome.

15

Said condition or disease may be a bleeding disorder, or a sleep disorder, or mastocytosis.

Said condition or disease may be associated with elevated body temperature, or ocular hypertension and glaucoma, or ocular hypotension.

20

Said condition may relate to surgical procedures to treat pain, inflammation and other unwanted sequelae wherein said surgical procedure includes incision, laser surgery or implantation.

- 25 The present invention also relates to a method of treating inflammation resulting from inflammatory diseases characterized by monocytic infiltration caused by the secretion of cytokines and/or chemokines by administration, to a patient in need of said treatment, of a pharmaceutical composition comprising a compound of the present invention

- 30 The current finding that the compounds of this invention are effective in attenuating the production of TNF family cytokines (TNFa), and the classical interleukin-1 (IL-1) family cytokines is especially important. These cytokines exert a broad spectrum of biological and pathological effects. They play key roles in inflammation and RA pathogenesis by

stimulating the release of multiple proinflammatory cytokines, including themselves, through the NFkB signaling pathway. Although alleviating the symptoms of RA in 50-65% of patients, a TNFa antibody is very expensive to use compared to chemically synthesized small molecules, inconvenient to administer usually requiring injections, and has been linked to tuberculosis, lymphoma, and other adverse effects. Unlike a TNFa antibody that totally eliminates all circulating TNFa in the system; the compounds of this invention only attenuate the production of TNFa by inhibiting proinflammatory PG receptors. Therefore, the adverse effects associated with a TNFa antibody in elevating infectious and cancerous tendency is less likely.

Proinflammatory elements TNF, RANTES, and MCP-1 are involved in the cascade of events in the early and late stages of atherosclerosis. Plasma MCP-1 levels have been linked to cardiovascular disease risk factors in clinical studies. Platelet activation leads to the release of MIP-1a, RANTES, and IL-8, which attract leukocytes and further activate other platelets. These evidences provide a direct linkage between homeostasis, infection, and inflammation and the development of atherosclerosis. The compounds of this invention are able to target multiple biomarkers of inflammation, thrombosis, and atherothrombosis simultaneously, which may confer pharmaceutical potential on the compounds of this invention in treating atherosclerosis and atherothrombosis. As a result, the compounds of this invention are unlikely to be associated with cardiovascular liability as in the case of the COXIBs, conversely it may even have a beneficial effect on cardiovascular function.

In summary, because of their ability to suppress the synthesis of some key proinflammatory cytokines/chemokines IL-8, MCP-1, MDC, RANTES, and TNFa, the compounds of the present invention are believed to be, not only at least as effective as COXIBs and NSAIDs in RA treatment, but also are a safer therapy in RA treatment. They are also a potential therapy for cardiovascular diseases.

The compounds of this invention are believed to treat or prevent inflammation at least in part by the decreasing the amount of the secretion of certain cytokines and/ or chemokines that result from the exposure of the patient to a stimulant.

In particular, the secretion of VEGF, MIP-1 β , IL-8, MCP-1, MDC, and RANTES may be reduced in those instances where said secretions are triggered by lipopolysaccharides (LPS) and or TNFa.

- 5 Interleukin-8 (IL-8): functions as a potent chemoattractant and activator of neutrophils, IL-8 is produced in response to stimulation with either IL-1 or TNFa. IL-8 not only accounts for a significant proportion of the chemotactic activity for neutrophils in rheumatoid arthritis (RA) synovial fluids, but also is a potent angiogenic factor in the RA synovium.
- 10 Monocyte chemoattractant protein- 1 (MCP-1, or CCL-2): is not only believed to play a role in inflammatory diseases characterized by monocytic infiltration, such as RA rheumatoid arthritis, psoriasis, and atherosclerosis, but is also implicated in other diseases, such as atopic dermatitis, renal disease, pleurisy, allergy and asthma, colitis, endometriosis, polymyositis and dermatomyositis, uveitis, restenosis, brain inflammation and obesity. MCP-1 also
- 15 controls leukocyte trafficking in vascular cells involved in diabetes and diabetes-induced atherosclerosis. MCP-1 antibodies are potential therapeutic agents for treating MCP-1/CCR2-mediated multiple inflammatory diseases.

- Tumor necrosis factor α (TNFa): mainly secreted by macrophages and recognized for its
- 20 importance in activating the cytokine cascade. TNFa stimulates the production of proinflammatory cytokines/chemokines, collagenases, metalloproteinases, and other inflammatory mediators; activates endothelial cells and neutrophils; promotes T- and B-cell growth, as well as stimulating bone resorption. The TNFa antibody infliximab not only decreases the production of local and systemic proinflammatory cytokines/chemokines, but
 - 25 also reduces serum MMP-3 production, nitric oxide synthase activity, VEGF release, and angiogenesis in inflamed joints.

- Macrophage-derived chemokine (MDC) induces chemotaxis for monocyte-derived dendritic cells, activated T cells and natural killer (NK) cells (Ho *et al.*, 2003). Highly expressed by
- 30 the three major cell types involved in allergic inflammation: eosinophils, basophils, and Th2 lymphocytes (Garcia *et al.*, 2005), as well as highly expressed in atopic dermatitis (Pivarcsi *et al.*, 2005), MDC plays a role in inflammatory diseases such as allergic asthma and atopic dermatitis (Ho *et al.*, 2003). Significantly enhanced in keratinocytes of patients with atopic

dermatitis, MDC could be a candidate therapeutic target for inflammatory skin disease such as atopic dermatitis (Qi *et al*, 2009). MDC is also implicated in disease activity of RA. After combination treatment with the disease-modifying anti-rheumatic drugs leflunomide and methotrexate in RA patients, plasma MCP-1 and MDC concentrations were significantly lower, and so was the recruitment of inflammatory cells into the sites of inflammation (Ho *et al*, 2003). Moreover, MDC also amplify platelet activation and has been associated with the pathogenesis of atherosclerotic disease including thrombosis (Gleissner *et al*, 2008).

Regulated on Activation, Normal T Cell Expressed and Secreted (RANTES) is a chemoattractant for blood monocytes, memory T-helper cells and eosinophils, and plays an active role in recruiting leukocytes into inflammatory sites. It also stimulates the release of histamine from basophils, activates eosinophils and causes hypodense eosinophils, which are associated with diseases such as asthma and allergic rhinitis. RANTES receptor CCR5 is also expressed on cells involved in atherosclerosis (e.g. monocytes/macrophages, T lymphocytes, or Th1-type cells), and is specialized in mediating RANTES-triggered atherosclerotic plaque formation (Zernecke *et al*, 2008). Like MCP-1, stimulation with RANTES enhances production of IL-6 and IL-8 in RA fibroblast-like synovial cells; elevated MMP-3 production by chondrocytes, and inhibited proteoglycan synthesis and enhanced proteoglycan release from the chondrocytes (Iwamoto *et al*, 2008). Both MCP-1 and RANTES were found to play an important role in allergic lung inflammation, lung leukocyte infiltration, bronchial hyper-responsiveness, and the recruitment of eosinophils in the pathogenesis of asthma (Conti *et al*, 2001). Similar to MCP-1, RANTES also enhances the inflammatory response within the nervous system, which plays an apparent role in the pathogenesis of multiple sclerosis (Conti *et al*, 2001). Inhibitors for RANTES may provide clinical benefits in treating inflammation, CNS disorders, parasitic disease, cancer, autoimmune and heart diseases (Castellani *et al*, 2007).

While the use of the compounds of this invention are believed to decrease the secretion of the above cytokines, it is also believed that the compounds of this invention are effective to decrease the secretion of ENA-7, PAI-1, CD-10, G-CSF, GM-CSF, IL-1 α and IL-18, as well.

The compounds of this invention may be also tested for efficacy in treating uveitis as described below.

ARACHIDONATE INDUCED UVEITIS

5 The rational for this protocol is to use arachidonate to directly produce ocular anterior segment uveitis, as opposed to using lipopolysaccharide (LPS) to indirectly release arachidonic acid.

Induction of uveitis:

10 Conscious male or female Dutch-belted pigmented rabbits weighing 2.5 - 3 kg are used for all in vivo slit lamp studies. Four animals are employed per test group. The right eye of each animal receiving 35 μ l of topically administered test and the contralateral left eye of each animal receiving 35 μ l of topically administered vehicle (t = 0 minutes), followed 30 minutes later by treatment with 35 μ l of 0.5% sodium arachidonate onto the surface of both eyes (t = 30 minutes). Both eyes are examined by slit lamp 60 minutes following sodium arachdionate challenge (t = 90 minutes) at 16x magnification under both white light and blue light
15 illumination at an approximate angle of 45° through 1 mm and 5 mm slit widths.

Measurement of anterior chamber leukocyte infiltration:


Anterior chamber leukocyte infiltration is measured using a numerical scoring system to estimate cell number per field defined by a 5 mm slit width: 0 = no cells per field (no response); 1 = 1 - 10 cells per field (mild); 2 = 11 - 20 cells per field (moderate); 3 = 26 - 50
20 cells per field (severe); 4 = >50 cells per filed (florid). Results are reported as the mean score value $\pm$ S.E.M.

The compounds of this invention may be tested according to the method described in "Characterization of Receptor Subtypes Involved in Prostanoid-Induced Conjunctival Pruritis and Their Role in Mediating Conjunctival Itching", Vol. 279, No.1,(JPET)279, 137-
25 142' 1996 for their efficacy in alleviating itch to thereby indicate that the compounds of this invention are useful in treating allergic conjunctivitis..

While the use of the compounds of this invention are believed to decrease the secretion of the above cytokines, it also is believed that the compounds of this invention are effective to decrease the secretion of ENA-7, PAI-1, CD-10, G-CSF, GM-CSF, IL-1 α and IL-18, as well.

- 5 Finally, said condition that may be treated with the compounds of this invention may be related to pain and inflammation and post-surgical scar and keloid formation.

In view of the various diseases and conditions that may be treated with the compositions of this invention there is provided a pharmaceutical product comprising a compound having the
10 following formula



wherein Y is $(CH_2)_m$ wherein m is 0 or an integer of from 1 to 3;

15

Z is selected from the group consisting of O, S, SO, SO₂ and $(CH_2)_p$, wherein p is 0 or an integer of from 1 to 3;

W is substituted aryl, alkyl, substituted alkyl, cycloalkyl or substituted cycloalkyl:

20

R₁ is selected from the group consisting of OR₇, N(R₇)₂, and N(R₇)SO₂R₇;

R₂ is H ;

25 R₄ is selected from the group consisting of alkyl, halogen-substituted alkyl and amino,

R₅ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxyl, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy;

- 5 R₆ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxyl, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy, wherein at least one of R₅ and R₆ is halogen;and

R₇ is selected from the group consisting of H, hydrocarbyl and substituted hydrocarbyl and/or
10 a pharmaceutically acceptable salt or a prodrug thereof, wherein said product is packaged and labeled for the treatment or prevention of a disease or condition selected from the group consisting of uveitis , allergic conditions, asthma, allergic asthma, allergic rhinitis, atherosclerosis, blood coagulation disorders, bone disorders, cancer, cellular neoplastic transformations, chronic obstructive pulmonary diseases and other forms of lung
15 inflammation, congestive heart failure, diabetic retinopathy, diseases or conditions requiring a treatment of anti-coagulation, diseases requiring control of bone formation and resorption, fertility disorders, hyperpyrexia , endometriosis gangrene, glaucoma, hypothermia, immune and autoimmune diseases, inflammatory conditions, metastatic tumor growth, migraine, mucus secretion disorders, nasal congestion, nasal inflammation, occlusive vascular diseases, ocular
20 hypertension, ocular hypotension, osteoporosis, pain, perennial rhinitis, pre-term labor pulmonary congestion, pulmonary hypotension, Raynaud's disease, rejection in organ transplant and by-pass surgery, respiratory conditions, rheumatoid arthritis, rhinorrhea, shock, sleep disorders, sleep-wake cycle disorders, sports injuries , muscle aches and pains , and surgical adjunct for minimizing pain , inflammation and scar/keloid formation.

25

Those skilled in the art will readily understand that for administration the compounds disclosed herein can be admixed with pharmaceutically acceptable excipients which, per se, are well known in the art. Specifically, a drug to be administered systemically, it may be formulated as a powder, pill, tablet or the like, or as a solution, emulsion, suspension, aerosol,
30 syrup or elixir suitable for oral or parenteral administration or inhalation.

For solid dosage forms, non-toxic solid carriers include, but are not limited to, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin,

the polyalkylene glycols, talcum, cellulose, glucose, sucrose and magnesium carbonate. The solid dosage forms may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the technique described in the U.S. Pat. Nos. 4,256,108; 4,166,452; and 4,265,874 to form osmotic therapeutic tablets for control release. Liquid pharmaceutically administrable dosage forms can, for example, comprise a solution or suspension of one or more of the compounds of the present invention and optional pharmaceutical adjuncts in a carrier, such as for example, water, saline, aqueous dextrose, glycerol, ethanol and the like, to thereby form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of nontoxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like. Typical examples of such auxiliary agents are sodium acetate, sorbitan monolaurate, triethanolamine, sodium acetate, triethanolamine oleate, etc. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., 16th Edition, 1980. The composition of the formulation to be administered, in any event, contains a quantity of one or more of the presently useful compounds in an amount effective to provide the desired therapeutic effect.

Parenteral administration is generally characterized by injection, either subcutaneously, intramuscularly or intravenously. Injectable formulations can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol and the like. In addition, if desired, the injectable pharmaceutical compositions to be administered may also contain minor amounts of nontoxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like.

The amount of the presently useful compound or compounds of the present invention administered is, of course, dependent on the therapeutic effect or effects desired, on the specific mammal being treated, on the severity and nature of the mammal's condition, on the manner of administration, on the potency and pharmacodynamics of the particular compound

or compounds employed, and on the judgment of the prescribing physician. The therapeutically effective dosage of the presently useful compound or compounds is preferably in the range of about 0.5 ng/kg/day or about 1 ng/kg/day to about 100 mg/kg/day.

- 5 For ophthalmic application, solutions are often prepared using a physiological saline solution as a major vehicle. Ophthalmic solutions should preferably be maintained at a comfortable pH with an appropriate buffer system. The formulations may also contain conventional, pharmaceutically acceptable preservatives, stabilizers and surfactants.
- 10 Preservatives that may be used in the pharmaceutical compositions of the present invention include, but are not limited to, benzalkonium chloride, chlorobutanol, thimerosal, phenylmercuric acetate and phenylmercuric nitrate. A useful surfactant is, for example, Tween 80. Likewise, various useful vehicles may be used in the ophthalmic preparations of the present invention. These vehicles include, but are not limited to, polyvinyl alcohol,
- 15 povidone, hydroxypropyl methyl cellulose, poloxamers, carboxymethyl cellulose, hydroxyethyl cellulose and purified water.

Tonicity adjustors may be added as needed or convenient. They include, but are not limited to, salts, particularly sodium chloride, potassium chloride, mannitol and glycerin, or any other

20 suitable ophthalmically acceptable tonicity adjustor.

Various buffers and means for adjusting pH may be used so long as the resulting preparation is ophthalmically acceptable. Accordingly, buffers include acetate buffers, citrate buffers, phosphate buffers and borate buffers. Acids or bases may be used to adjust the pH of these

25 formulations as needed.

Similarly, an ophthalmically acceptable antioxidant for use in the present invention includes, but is not limited to, sodium metabisulfite, sodium thiosulfate, acetylcysteine, butylated hydroxyanisole and butylated hydroxytoluene.

30 Other excipient components which may be included in the ophthalmic preparations are chelating agents. A useful chelating agent is edentate disodium, although other chelating agents may also be used in place or in conjunction with it.


For topical use, creams, ointments, gels, solutions or suspensions, etc., containing the compound of the present invention are employed. Topical formulations may generally be comprised of a pharmaceutical carrier, cosolvent, emulsifier, penetration enhancer, preservative system, and emollient.

The actual dose of the compounds of the present invention depends on the specific compound, and on the condition to be treated; the selection of the appropriate dose is well within the knowledge of the skilled artisan.

The present invention is not to be limited in scope by the exemplified embodiments, which are only intended as illustrations of specific aspects of the invention. Various modifications of the invention, in addition to those disclosed herein, will be apparent to those skilled in the art by a careful reading of the specification, including the claims, as originally filed. It is intended that all such modifications will fall within the scope of the appended claims.

We claim:

1. A compound, that is a 1-[(5-halo or alkyl or fluoroalkyl or alkoxy-2-[(cycloalkyl or substituted cycloalkyl)oxy]phenyl)methyl]-(5-alkyl or fluoroalkyl)-1H-pyrazole-3-(carboxylic acid or methylene carboxylic acid) or an alkyl or aryl ester or sulfonamide thereof.
2. A compound, that is a 1-[(5-halo or alkyl or fluoroalkyl or alkoxy-2-[(cycloalkyl)oxy]phenyl)methyl]-(5-alkyl or fluoroalkyl)-1H-pyrazole-3-(carboxylic acid or methylene carboxylic acid) or an alkyl or aryl ester or sulfonamide thereof.
3. A compound according to claim 1 wherein said compound is a 1-[(5-halo-2-[(2-cycloalkyl)oxy]phenyl)methyl]-3-carboxylic acid or ester or sulfonamide thereof.
4. A compound according to claim 1 wherein said compound is a 1-[(5-halo-2-[(2-cycloalkyl)oxy]phenyl)methyl]-3-carboxylic acid.
5. A compound according to claim 3 wherein said ester or sulfonamide is an alkyl ester or sulfonamide.
6. A compound according to claim 3, wherein said halo is selected from the group consisting of chloro and bromo.
7. A compound represented by the following formula



wherein Y is $(\text{CH}_2)_m$ wherein m is 0 or an integer of from 1 to 3;

Z is selected from the group consisting of O, S, SO, SO_2 and $(\text{CH}_2)_p$, wherein p is 0 or an integer of from 1 to 3;

W is substituted aryl, alkyl, substituted alkyl, cycloalkyl or substituted cycloalkyl:

R_i is selected from the group consisting of OR₇, N(R₇)₂, and N(R₇)SO₂R₇;

R₂ is H ;

R₄ is selected from the group consisting of alkyl, halogen-substituted alkyl and amino

R₅ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxyl, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy;

R₆ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxy, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy, wherein at least one of R₅ and R₆ is halogen;and


R₇ is selected from the group consisting of H, hydrocarbyl and substituted hydrocarbyl.

8. The compound of claim 7 wherein R₇ is selected from the group consisting of carbocyclic aryl and alkyl.

9. The compound of claim 7 wherein R_i is OH.

10. The compound of claim 9 wherein R₄ is selected from the group consisting of alkyl and chloro and bromo-substituted alkyl.

11. The compound of claim 9 wherein R_4 is H.
12. The compound of claim 9 wherein Y is absent.
13. The compound of claim 9 wherein Z is O.
14. The compound of claim 9 wherein W is cyclopentyl
15. The compound of claim 9 that is selected from the group consisting of
 1-(5-Chloro-2-cyclopentylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid,
 1-(5-Bromo-2-cyclopentylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid and
 1-(5-bromo-2-cyclobutylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid
16. A method comprising administering a compound represented by the following formula



wherein Y is $(CH_2)_m$ wherein m is 0 or an integer of from 1 to 3;

Z is selected from the group consisting of O, S, SO, SO_2 and $(CH_2)_p$, wherein p is 0 or an integer of from 1 to 3;

W is substituted aryl, alkyl, substituted alkyl, cycloalkyl or substituted cycloalkyl:

R_i is selected from the group consisting of OR₇, N(R₇)₂, and N(R₇)SO₂R₇;

R₂ is H;

R₄ is selected from the group consisting of alkyl, halogen-substituted alkyl and amino;

R₅ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxy, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy;

R_e is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxy, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy, wherein at least one of R₅ and R₆ are halogen; and

R₇ is selected from the group consisting of H, hydrocarbyl and substituted hydrocarbyl.

17. The method of claim 16 wherein R_i is OH.

18. The method compound of claim 17 wherein R₇ is selected from the group consisting of carbocyclic aryl and alkyl

19. The method of claim 18 wherein W is cycloalkyl.

20. The method of claim 19 wherein R₄ is selected from the group consisting of alkyl and chloro and bromo-substituted alkyl.

21. The method of claim 19 wherein R₄ is H.

22. The method of claim 19 wherein Y is absent, i.e. n.

23. The method of claim 19 wherein Z is O. is 0.
24. The compound of claim 19 wherein W is cyclopentyl.
25. The compound of claim 19 that is selected from the group consisting of
1-(5-Chloro-2-cyclopentylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid,
1-(5-Bromo-2-cyclopentylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid and
1-(5-bromo-2-cyclobutylmethoxy-benzyl)-5-methyl-1H-pyrazole-3-carboxylic acid
26. The method of claim 25 wherein said compound is administered to treat DPI, FP, EP1, TP and/or EP4 receptor mediated diseases or conditions.
27. The method of claim 26 wherein said condition or disease is related to inflammation.
28. The method of claim 26 wherein said DPI, FP, EP1, TP and/or EP4 receptor mediated condition or disease is selected from the group consisting of allergic conditions, asthma, allergic asthma, allergic rhinitis, uveitis and related disorders, atherosclerosis, blood coagulation disorders, bone disorders, cancer, cellular neoplastic transformations, chronic obstructive pulmonary diseases and other forms of lung inflammation, congestive heart failure, diabetic retinopathy, diseases or conditions requiring a treatment of anti-coagulation, diseases requiring control of bone formation and resorption, endometriosis, fertility disorders, gangrene, glaucoma, hyperpyrexia, immune and autoimmune diseases, inflammatory conditions, metastatic tumor growth, migraine, mucus secretion disorders, nasal congestion, nasal inflammation, occlusive vascular diseases, ocular hypertension, ocular hypotension, osteoporosis, pre-term labor, rheumatoid arthritis, pain, perennial rhinitis, pulmonary congestion, pulmonary hypotension, Raynaud's disease, rejection in organ transplant and by-pass surgery, respiratory conditions, hirsutism, rhinorrhea, shock, sleep disorders, and sleep-wake cycle disorders.

29. The method of claim 26 wherein said compound is administered as a surgical adjunct in ophthalmology for cataract removal and artificial lens insertion, ocular implant procedures, photorefractive radial keratotomy and other ophthalmological laser procedures.
30. The method of claim 26 wherein said compound is administered as a surgical adjunct in a procedure involving skin incisions, relief of pain and inflammation and scar formation/keloids post-surgery, for treating sports injuries and general aches and pains in muscles and joints.
31. The method of claim 26 wherein said DPi, FP, EPi, TP , and/or EP₄ receptor mediated condition or disease is an EPi and/or EP₄ receptor mediated condition or disease.
32. The method of claim 26 wherein said DPi, FP, EPi, TP and/or EP₄ receptor mediated condition or disease is an allergic condition.
33. The method of claim 32 wherein said condition is dermatological allergy.
34. The method of claim 26 wherein said condition is an ocular allergy.
35. The method of claim 26 wherein said condition is a respiratory allergy.
36. The method of claim 26 wherein said condition or disease is selected from the group consisting of nasal congestion, rhinitis, and asthma.
37. The method of claim 26 wherein said condition or disease is related to pain.
38. The method of claim 26 wherein said condition or disease is selected from the group consisting of arthritis, migraine, and headache.
39. The method of claim 26 wherein said condition or disease is associated with the gastrointestinal tract.

19043 PCT (AP)

40. The method of claim 26 wherein said condition or disease is selected from the group consisting of peptic ulcer, heartburn, reflux esophagitis, erosive esophagitis, non-ulcer dyspepsia, infection by *Helicobacter pylori*, alrnyitis, and irritable bowel syndrome.

41. The method of claim 26 wherein said condition or disease is selected from the group consisting of hyperalgesia and allodynia.

42. The method of claim 26 wherein said condition or disease is related to mucus secretion.

43. The method of claim 26 wherein said mucus secretion is gastrointestinal.

44. The method of claim 26 wherein said mucus secretion occurs in the nose, sinuses, throat, or lungs.

45. The method of claim 26 wherein said condition or disease is related to abdominal cramping.

46. The method of claim 26 wherein said condition or disease is irritable bowel syndrome.

47. The method of claim 26 wherein said condition or disease is a bleeding disorder.

48. The method of claim 26 wherein said condition or disease is a sleep disorder.

49. The method of claim 26 wherein said condition or disease is mastocytosis.

50. The method of claim 26 wherein said condition or disease is associated with elevated body temperature.

51. The method of claim 26 wherein said condition or disease is associated with ocular hypertension and glaucoma.

19043 PCT (AP)


52. The method of claim 26 wherein said condition or disease is associated with ocular hypotension.

53. The method of claim 26 wherein said condition relates to surgical procedures to treat pain, inflammation and other unwanted sequelae wherein said surgical procedure includes incision, laser surgery or implantation.

54. The method of claim 26 where said condition is related to pain and inflammation and post-surgical scar and keloid formation.

55. The method of claim 26 where said condition is related to diseases of female reproduction, associated with menstrual cramping, endometriosis, and pre-term labor

56. A pharmaceutical product comprising a compound having the following formula



wherein Y is $(CH_2)_m$ wherein m is 0 or an integer of from 1 to 3;

Z is selected from the group consisting of O, S, SO, SO_2 and $(CH_2)_p$, wherein p is 0 or an integer of from 1 to 3;

W is substituted aryl, alkyl, substituted alkyl, cycloalkyl or substituted cycloalkyl:

R_7 is selected from the group consisting of OR_7 , $N(R_7)_2$, and $N(R_7)SO_2R_7$;

R₂ is H ;

R₄ is selected from the group consisting of alkyl, halogen-substituted alkyl and amino ;


R₅ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxy, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy;

R₆ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxy, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy, wherein at least one of R₅ and R₆ are halogen;and

R₇ is selected from the group consisting of H, hydrocarbyl and substituted hydrocarbyl.

or a pharmaceutically acceptable salt or a prodrug thereof, wherein said product is packaged and labeled for the treatment or prevention of a disease or condition selected from the group consisting of uveitis, allergic conditions, asthma, allergic asthma, allergic rhinitis, atherosclerosis, blood coagulation disorders, bone disorders, cancer, cellular neoplastic transformations, chronic obstructive pulmonary diseases and other forms of lung inflammation, congestive heart failure, diabetic retinopathy, diseases or conditions requiring a treatment of anti-coagulation, diseases requiring control of bone formation and resorption, endometriosis fertility disorders, hyperpyrexia , gangrene, glaucoma, hypothermia, immune and autoimmune diseases, inflammatory conditions, menstrual cramping ,metastatic tumor growth, migraine, mucus secretion disorders, nasal congestion, nasal inflammation, occlusive vascular diseases, ocular hypertension, ocular hypotension, osteoporosis, pain, perennial rhinitis, pre-term labor pulmonary congestion, pulmonary hypotension, Raynaud's disease, rejection in organ transplant and by-pass surgery, respiratory conditions, rheumatoid arthritis, rhinorrhea, shock, sleep disorders, sleep-wake cycle disorders, sports injuries , muscle aches and pains, and surgical adjunct for minimizing pain, inflammation and scar/keloid formation.

57. A pharmaceutical composition comprising a compound having the following formula



Wherein Y is $(CH_2)_m$ wherein m is 0 or an integer of from 1 to 3;

Z is selected from the group consisting of O, S, SO, SO_2 and $(CH_2)_p$, wherein p is 0 or an integer of from 1 to 3;

W is cycloalkyl;

R₁ is selected from the group consisting of OR_7 , $N(R_7)_2$, and $N(R_7)SO_{2R_7}$;

R₂ is H ;

R₄ is selected from the group consisting of alkyl, halogen-substituted alkyl and amino;

R₅ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxy, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy;

R₆ is selected from the group consisting of H, hydroxyl, alkyl, aryl, alkoxy, aryloxy, halogen, nitro, amino, cyano and hydroxyl, halogen, nitro, amino and cyano-substituted alkyl, aryl, alkoxy or aryloxy, wherein at least one of R₅ and R₆ are halogen; and

R₇ is selected from the group consisting of H, hydrocarbyl and substituted hydrocarbyl, or a pharmaceutically acceptable salt or a prodrug thereof, and a pharmaceutically acceptable excipient.

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2012/071232

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C07D231/14 A61K31/415 A61P11/00
 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal , CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	wo 2005/040128 Al (GLAXO GROUP LTD [GB] ; GIBLIN GERARD MARTIN PAUL [GB] ; HALL ADRIAN [GB]) 6 May 2005 (2005-05-06) cited in the applicati on page 1, line 1 - page 1, line 7 page 3, line 12 - page 5, line 9 -----	1-6
A	wo 2006/114313 Al (GLAXO GROUP LTD [GB] ; CONWAY ELIZABETH ANN [GB] ; GIBLIN GERARD MARTIN) 2 November 2006 (2006-11-02) page 1, line 1 - page 1, line 5 page 3, line 10 - page 4, line 30 -----	1-6
A, P	wo 2012/003414 Al (ALLERGAN INC [US] ; WOODWARD DAVID F [US] ; MARTOS JOSE L [GB] ; WANG JEN) 5 January 2012 (2012-01-05) page 1, line 15 - page 1, line 21 claims 1-11 -----	1-6



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) on which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 February 2013

Date of mailing of the international search report

02/05/2013

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040,
 Fax: (+31-70) 340-3016

Authorized officer

Usuel 1i , Ambrogio o

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2012/071232

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos. :

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos. :

1-6

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-6

Compounds of claim 1

2. claims: 7-57

Compounds of claim 7

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2012/071232

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
wo 2005040128 AI	06-05-2005	AT 386725 T DE 602004011966 T2 EP 1675832 AI ES 2298832 T3 JP 2007509104 A US 2007060596 AI wo 2005040128 AI	15-03-2008 12-02-2009 05-07-2006 16-05-2008 12-04-2007 15-03-2007 06-05-2005
wo 2006114313 AI	02-11-2006	EP 1874303 AI JP 2008539185 A US 2009239845 AI wo 2006114313 AI	09-01-2008 13-11-2008 24-09-2009 02-11-2006
wo 2012003414 AI	05-01-2012	AR 082073 AI AU 2011272716 AI CA 2804194 AI CN 103038219 A EP 2588459 AI SG 186886 AI TW 201206891 A US 2012165382 AI wo 2012003414 AI	07-11-2012 07-02-2013 05-01-2012 10-04-2013 08-05-2013 28-02-2013 16-02-2012 28-06-2012 05-01-2012