



(86) Date de dépôt PCT/PCT Filing Date: 2005/12/17
(87) Date publication PCT/PCT Publication Date: 2006/07/20
(45) Date de délivrance/Issue Date: 2014/02/18
(85) Entrée phase nationale/National Entry: 2007/07/12
(86) N° demande PCT/PCT Application No.: EP 2005/013634
(87) N° publication PCT/PCT Publication No.: 2006/074789
(30) Priorité/Priority: 2005/01/14 (FR0500422)


(51) Cl.Int./Int.Cl. *C07D 209/42* (2006.01),
A61K 31/404 (2006.01), *A61K 31/4439* (2006.01),
A61P 3/06 (2006.01), *A61P 3/10* (2006.01),
A61P 9/10 (2006.01), *C07D 401/04* (2006.01)

(72) Inventeurs/Inventors:
ADJE, NATHALIE, FR;
ROCHE, DIDIER, FR;
YVON, STEPHANE, FR

(73) Propriétaire/Owner:
MERCK PATENT GESELLSCHAFT MIT
BESCHRAENKTER HAFTUNG, DE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : DERIVES D'ACIDE 1H-INDOLE-3-CARBOXYLIQUE ET UTILISATION EN TANT QU'AGONISTES DE PPAR
(54) Title: 1H-INDOLE-3-CARBOXYLIC ACID DERIVATIVES AND THEIR USE AS PPAR AGONISTS



(57) Abrégé/Abstract:

A compound of the formula (1): (see formula 1), or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof, in which: R¹ is -O-R'¹ or -NR'¹R''¹, wherein R'¹ and R''¹ are each H or optionally substituted alkyl,


(57) Abrégé(suite)/Abstract(continued):

alkenyl, alkynyl, cycloalkyl, aryl, or heteroaryl; R^2 is: optionally substituted alkyl, alkenyl or alkynyl radical; an aryl radical, optionally substituted and/or optionally fused to a monocyclic or polycyclic, saturated or unsaturated 5- to 8-membered nucleus optionally containing one or more hetero atoms chosen from O, N and S, the said nucleus itself being optionally substituted, or a saturated, unsaturated or aromatic, optionally substituted 5- to 8-membered monocyclic heterocyclic radical; R^3 is H or optionally substituted alkyl; and A is a linear or branched C_{1-6} alkylene chain; the processes for the preparation of these compounds, the uses thereof for the treatment of dyslipidaemia, atherosclerosis and diabetes, and the pharmaceutical compositions comprising them.

26474-1095

Abstract

A compound of the formula (1):



or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable
 5 addition salt thereof, in which: R^1 is $-O-R'^1$ or $-NR'^1R''^1$, wherein R'^1 and R''^1 are each
 H or optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, or heteroaryl; R^2 is:
 optionally substituted alkyl, alkenyl or alkynyl radical; an aryl radical, optionally
 substituted and/or optionally fused to a monocyclic or polycyclic, saturated or
 10 unsaturated 5- to 8-membered nucleus optionally containing one or more hetero
 atoms chosen from O, N and S, the said nucleus itself being optionally substituted, or
 a saturated, unsaturated or aromatic, optionally substituted 5- to 8-membered
 monocyclic heterocyclic radical; R^3 is H or optionally substituted alkyl; and A is a
 linear or branched C_{1-6} alkylene chain; the processes for the preparation of these
 compounds, the uses thereof for the treatment of dyslipidaemia, atherosclerosis and
 15 diabetes, and the pharmaceutical compositions comprising them.

1H-INDOLE-3-CARBOXYLIC ACID DERIVATIVES AND THEIR USE AS PPAR AGONISTS

5 **[0001]** The present invention relates to 1*H*-indole-3-carboxylic acid derivatives that can be used in the treatment of dyslipidaemia, atherosclerosis and diabetes. The invention also relates to pharmaceutical compositions comprising them and to processes for the preparation of these compounds.

10 **[0002]** In addition, the invention relates to the use of these compounds for the production of medicaments for the treatment of dyslipidaemia, atherosclerosis and diabetes.

15 **[0003]** The chronic effect of a calorie imbalance has resulted in an epidemic increase in the incidence of metabolic diseases in modern society. As a result, the World Health Organization has estimated that the global incidence of type 2 diabetes will exceed 300 million in 2030. Although several therapeutic options exist, none of them reverses the progress of this plague.

20 **[0004]** Although the control of glycated haemoglobin and plasmatic glycaemia in the fasted state are still considered as the primary objectives of antidiabetic treatments, acknowledgement of the fact that the diabetic state encompasses a range of metabolic disorders has broadened scope and expectations of future therapies. In the course of the last decade, hyperglycaemia has been shown to be not the only component of a series of anomalies affecting type-2 diabetic patients. Concurrent diseases, including insulin resistance, obesity, hypertension and dyslipidaemia, which, if they are present together or in part, constitutes what has been described as metabolic syndrome or syndrome X. This array of metabolic disorders forms the bases of a substantial increase in the incidence of cardiovascular disease in these patients.

25 **[0005]** In the search for novel and improved treatment options for diabetic patients, the family of receptors activated by the peroxisome proliferators ("peroxisome proliferator-activated receptor": PPAR) appears potentially to

30

be an ideal target. This family of ligand-activated transcription factors modulates numerous aspects of lipid and carbohydrate metabolism, thus having the possibility of attacking several facets of the diabetic phenotype. There are three types of PPAR: PPAR alpha, gamma and delta (PPAR α , PPAR γ and PPAR δ , respectively).

[0006] PPAR α is involved in stimulating the β -oxidation of fatty acids. In rodents, a change transmitted by a PPAR α in the expression of genes involved in fatty acid metabolism is the basis of the phenomenon of peroxisome proliferation, a pleiotropic cellular response, mainly limited to the liver and the kidneys, which can lead to hepatocarcinogenesis in rodents. The phenomenon of peroxisome proliferation is not encountered in man. In addition to its role in peroxisome proliferation in rodents, PPAR α is also involved in controlling the levels of HDL cholesterol in rodents and humans. This effect is at least partially based on a transcription regulation transmitted by a PPAR α of the major HDL apolipoproteins, apo A-I and apo A-II. The hypotriglyceridaemic action of fibrates and fatty acids also involves PPAR α and can be summarised as follows: (i) increased lipolysis and clearance of the remaining particles, due to changes in the levels of lipoprotein lipase and of apo C-III, (ii) stimulation of fatty acid uptake by the cell and its subsequent conversion into acyl-CoA derivatives by induction of a protein for binding fatty acids and acyl-CoA synthase, (iii) induction of the β -oxidation pathways of fatty acids, (iv) reduction in the synthesis of fatty acids and triglycerides, and finally (v) reduction in the production of VLDL. As a result, both the improved catabolism of the triglyceride-rich particles and the reduced secretion of VLDL particles constitute mechanisms that contribute towards the hypolipidaemic effect of fibrates.

[0007] Fibric acid derivatives, such as clofibrate, fenofibrate, benzafibrate, ciprofibrate, beclofibrate and etofibrate, and also gemfibrozil, each of which are PPAR α ligands and/or activators, produce a substantial reduction in plasmatic triglycerides and also a certain increase in HDLs. The effects on LDL cholesterol are contradictory and may depend on the compound and/or

the dyslipidaemic phenotype. For these reasons, this class of compounds was first used for the treatment of hypertriglyceridaemia (i.e. Fredrickson Type IV and V) and/or mixed hyperlipidaemia.

[0008] The activation of a PPAR δ was initially reported as not being
5 involved in the modulation of the levels of glucose or of triglycerides (Berger et al., *J. Biol. Chem.*, (1999), Vol. 274, pp. 6718-6725). Later, it was shown that the activation of PPAR δ leads to higher levels of HDL cholesterol in dbldb mice (Leibowitz et al., *FEBS Letters*, (2000), **473**, 333-336). Furthermore, a PPAR δ agonist, during its administration to obese adult insulin-
10 resistant rhesus monkeys, caused a dramatic dose-dependent increase in HDL cholesterol in the serum, while at the same time reducing the levels of low-density LDLs, by depleting the triglycerides and the insulin (Oliver et al., *PNAS*, (2001), **98**, 5306-5311). The same publication also showed that the activation of PPAR δ increased the AI cassette binding the ATP inverse
15 transporter of cholesterol and induced a flow of cholesterol specific for apolipoprotein A1. Taken together, these observations suggest that the activation of PPAR δ is useful for the treatment of and preventing diseases and cardiovascular states comprising atherosclerosis, hypertriglyceridaemia and mixed dyslipidaemia (PCT publication WO 01/00603 (Chao et al.)).

[0009] The subtypes of PPAR γ receptor are involved in the activation of
20 the programme of adipocyte differentiation and are not involved in the stimulation of peroxisome proliferation in the liver. There are two known isoforms of PPAR γ protein: PPAR γ 1 and PPAR γ 2, which differ only in the fact that PPAR γ 2 contains 28 additional amino acids at the amino end. The DNA
25 sequences for the human isotypes are described by Elbrecht et al., *BBRC*, **224**, (1996), 431-437. In mice, PPAR γ 2 is specifically expressed in the fat cells. Tontonoz et al., *Cell*, **79**, (1994), 1147-1156, provide proof showing that one physiological role of PPAR γ 2 is to induce adipocyte differentiation. As with other members of the superfamily of nuclear hormone receptors,
30 PPAR γ 2 regulates the expression of genes via an interaction with other proteins and binding to hormone response elements, for example in the 5'

lateral regions of the response genes. An example of a PPAR γ 2 response gene is the tissue-specific P2 adipocyte gene. Although peroxisome proliferators, comprising fibrates and fatty acids, activate the transcriptional activity of PPAR receptors, only prostaglandin J₂ derivatives have been identified as potential natural ligands of the PPAR γ subtype, which also binds antidiabetic thiazolidinedione agents with high affinity.

[0010] It is generally thought that glitazones exert their effects by binding to receptors of the family of peroxisome proliferator-activated receptors (PPAR), by controlling certain transcription elements in relation with the biological species listed above. See Hulin et al., *Current Pharm. Design*, (1996), 2, 85-102. In particular, PPAR γ has been imputed as a major molecular target for the glitazone class of insulin sensitisers.

[0011] Many compounds of glitazone type, which are PPAR agonists, have been approved for use in the treatment of diabetes. These are troglitazone, rosiglitazone and pioglitazone, which are all primary or exclusive agonists of PPAR γ .

[0012] This indicates that the search for compounds having varying degrees of PPAR α , PPAR γ and PPAR δ activation might lead to the discovery of medicaments that efficiently reduce triglycerides and/or cholesterol and/or glucose, presenting great potential in the treatment of diseases, such as type 2 diabetes, dyslipidaemia, syndrome X (comprising metabolic syndrome, i.e. reduced glucose tolerance, insulin resistance, hypertriglyceridaemia and/or obesity), cardiovascular diseases (comprising atherosclerosis) and hypercholesterolaemia.

[0013] The combinations of the PPAR activities that have been studied the most extensively are the PPAR alpha plus gamma combination (dual agonists) with, especially, tesaglitazar, and also the alpha, gamma plus delta triple combination (PPARpan agonists).

[0014] Although glitazones are beneficial in the treatment of NIDDM, a number of serious unfavourable side effects associated with the use of these compounds have been found. The most serious of these was toxicity to the

liver, which has resulted in a certain number of deaths. The most serious problems arose in the use of troglitazone, which has recently been removed from the market for toxicity reasons.


[0015] Besides the potential hepatic toxicity of glitazones, other deleterious effects have been associated with PPAR gamma full agonists, for instance weight gain, anaemia and oedema, which limit their use (rosiglitazone, pioglitazone).

[0016] On account of the problems that have been encountered with glitazones, researchers in many laboratories have studied classes of PPAR agonists that are not glitazones and do not contain 1,3-thiazolidinedione species, but which modulate the three known subtypes of PPAR, together or separately, to variable degrees (measured by intrinsic power, maximum breadth of functional response or spectrum of changes in gene expression).

[0017] Thus, recent studies (cf. WO 01/30343 and WO 02/08188) have revealed that certain compounds have PPAR agonist or partial agonist properties, which are useful in the treatment of type 2 diabetes with reduced side effects with respect to the heart weight and body weight.

[0018] The inventors have now discovered a novel class of compounds that are partial or full agonists of PPAR γ , with differing degrees of PPAR α and/or PPAR δ activity.

[0019] More specifically, the invention relates to compounds derived from the 1*H*-indole-3-carboxylic acid of the formula (1) below:



in which:

R¹ represents -O-R'¹ or -NR'¹R''¹, with R'¹ and R''¹, which may be identical or different, being chosen from a hydrogen atom, an alkyl radical, an

26474-1095

6

alkenyl radical, an alkynyl radical, a cycloalkyl radical, an aryl radical and a heteroaryl radical;

R^2 is chosen from:

- an alkyl, alkenyl or alkynyl radical;
- 5 • an aryl radical, optionally substituted and/or optionally fused to a monocyclic or polycyclic, saturated or unsaturated 5- to 8-membered nucleus optionally containing one or more hetero atoms chosen from O, N and S, the said nucleus itself being optionally substituted, and
- 10 • a saturated, unsaturated or aromatic, optionally substituted 5- to 8-membered monocyclic heterocyclic radical containing one or more hetero atoms chosen from O, N and S;

R^3 is chosen from a hydrogen atom and an alkyl radical; and


A represents a linear or branched alkylene chain containing from 1 to
15 6 carbon atoms;

the possible optical isomers, oxide forms and solvates thereof, and also pharmaceutically acceptable addition salts thereof with acids or bases.

26474-1095

6a

[0019a] The invention may also relate to a compound of formula (1):



or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof, in which: R^1 is $-O-R'^1$ or $-NR'^1R''^1$; R'^1 and R''^1 are each

5 independently H, optionally substituted alkyl having 1 to 10 carbon atoms, optionally substituted alkenyl having 2 to 10 carbon atoms, optionally substituted alkynyl having 2 to 10 carbon atoms, optionally substituted cycloalkyl having 4 to 9 carbon atoms, optionally substituted aryl having 6 to 18 carbon atoms, or optionally substituted heteroaryl; R^2 is optionally substituted alkyl having 1 to 10 carbon atoms, optionally

10 substituted alkenyl having 2 to 10 carbon atoms, or optionally substituted alkynyl having 2 to 10 carbon atoms, aryl having 6 to 18 carbon atoms which is optionally substituted and/or optionally fused to a monocyclic or polycyclic, saturated or unsaturated 5- to 8-membered nucleus optionally containing one or more hetero atoms which are O, N or S, and

15 said nucleus is optionally substituted, or a saturated, unsaturated or aromatic, optionally substituted 5- to 8-membered monocyclic heterocyclic radical containing one or more hetero atoms which are O, N or S; R^3 is H or optionally substituted alkyl having 1 to 10 carbon atoms; and A is a linear or branched alkylene chain containing from 1 to 6 carbon atoms; wherein optionally substituted alkyl groups listed above are

20 each optionally substituted one or more times by halogen, $-O$ -alkyl having 1 to 10 carbon atoms, aryl having 6 to 18 carbon atoms and which is optionally substituted by halogen, methyl, ethyl, methoxy, phenyl, trifluoromethyl or

26474-1095

6b

trifluoromethoxy, cycloalkyl having 4 to 9 carbon atoms, or a heterocyclic group; wherein optionally substituted alkenyl groups listed above are each unsubstituted or substituted one or more times by halogen, trifluoromethyl, trifluoromethoxy, hydroxyl, alkoxy, alkoxycarbonyl, carboxyl or oxo; wherein optionally substituted alkynyl groups

5 listed above are each unsubstituted or substituted one or more times by halogen, trifluoromethyl, trifluoromethoxy, hydroxyl, alkoxy, alkoxycarbonyl, carboxyl or oxo; wherein optionally substituted aryl groups listed above are each unsubstituted or substituted one or more times by halogen, methyl, ethyl, methoxy, phenyl, trifluoromethyl or trifluoromethoxy; wherein optionally substituted heterocyclic groups

10 listed above are each unsubstituted or substituted one or more times by halogen, methyl, ethyl, methoxy, phenyl, trifluoromethyl or trifluoromethoxy; wherein aromatic heterocyclic groups or heteroaryl groups are selected from pyridyl, pyrimidinyl, triazolyl, thiadiazolyl, oxazolyl, thiazolyl, thienyl, quinolyl, benzothiazolyl, acridinyl, phenazinyl, carbazolyl, and benzothiophenyl; wherein saturated or unsaturated

15 heterocyclic groups are selected morpholinyl, piperidinyl, thiazolidinyl, oxazolidinyl, tetrahydrothienyl, tetrahydrofuryl, pyrrolidinyl, isoxazolidinyl, imidazolidinyl, and pyrazolidinyl; and wherein optionally substituted cycloalkyl is unsubstituted or substituted one or more times by halogen atoms, alkyl, alkenyl, alkynyl, trifluoromethyl, trifluoromethoxy, hydroxyl, alkoxy, alkoxycarbonyl, carboxyl, or oxo.

20 **[0020]** The acids that can be used for the formation of salts of compounds of the formula (1) are mineral or organic acids. The resulting salts are, for example, the hydrochlorides, hydrobromides, sulfates, hydrogen sulfates, dihydrogen phosphates, citrates, maleates, fumarates, trifluoroacetates, 2-naphthalenesulfonates and *para*-toluenesulfonates.

25 **[0021]** The bases that can be used for the formation of salts of compounds of the formula (1) are organic or mineral bases. The resulting salts are, for example, the salts formed with metals and especially alkali metals, alkaline-earth metals and transition metals (such as sodium, potassium, calcium, magnesium or aluminium) or

26474-1095

6c

with bases, for instance ammonia or secondary or tertiary amines (such as diethylamine, triethylamine, piperidine, piperazine or morpholine) or with basic amino acids, or with osamines (such as meglumine) or with amino alcohols (such as 3-aminobutanol and 2-aminoethanol).

[0022] The invention especially encompasses the pharmaceutically acceptable salts, but also salts that allow a suitable separation or crystallisation of the compounds of the formula (1), such as the salts obtained with chiral amines or chiral acids.

5 **[0023]** Examples of chiral amines that can be used include quinine, brucine, (S)-1-(benzyloxymethyl)propylamine (III), (-)-ephedrine, (4S,5R)-(+)-1,2,3,4-tetramethyl-5-phenyl-1,3-oxazolidine, (R)-1-phenyl-2-*p*-tolylethylamine, (S)-phenylglycinol, (-)-N-methylephedrine, (+)-(2S,3R)-4-dimethylamino-3-methyl-1,2-diphenyl-2-butanol, (S)-phenylglycinol and (S)- α -methylbenzylamine, or a mixture of two or more thereof.

10 **[0024]** Examples of chiral acids that can be used include (+)-d-di-O-benzoyltartaric acid, (-)-l-di-O-benzoyltartaric acid, (-)-di-O,O'-*p*-toluyl-l-tartaric acid, (+)-di-O,O'-*p*-toluyl-d-tartaric acid, (R)-(+)-malic acid, (S)-(-)-malic acid, (+)-camphanic acid, (-)-camphanic acid, *R*-(-)-1,1'-binaphthalene-2,2'-diyl hydrogen phosphate, (S)-(+)-1,1'-binaphthalene-2,2'-diyl hydrogen phosphate, (+)-camphoric acid, (-)-camphoric acid, (S)-(+)-2-phenylpropionic acid, (R)-(-)-2-phenylpropionic acid, d-(-)-mandelic acid, l-(+)-mandelic acid, d-tartaric acid and l-tartaric acid, or a mixture of two or more thereof.

15 **[0025]** The chiral acid is preferably chosen from (-)-di-O,O'-*p*-toluyl-l-tartaric acid, (+)-di-O,O'-*p*-toluyl-d-tartaric acid, (R)-(-)-1,1'-binaphthalene-2,2'-diyl hydrogen phosphate, (S)-(+)-1,1'-binaphthalene-2,2'-diyl hydrogen phosphate, d-tartaric acid and L-tartaric acid, or a mixture of two or more thereof.

20 **[0026]** The invention also encompasses the possible optical isomers, in particular stereoisomers and diastereoisomers, where appropriate, of the compounds of the formula (1), and also mixtures of the optical isomers in any proportions, including racemic mixtures.

25 **[0027]** Depending on the nature of the substituents, the compounds of the formula (1) may also be in various tautomeric forms, which are also included in the present invention, alone or as mixtures of two or more thereof, in all proportions.

30

26474-1095

8

[0028] Prodrugs of the compounds of the formula (1) above are also possible.

[0029] The term "prodrugs" means compounds which, once administered to the patient, are chemically and/or biologically converted by the living body,
5 into compounds of the formula (1).

[0030] In the compounds of the formula (1) defined above, the term "alkyl radical" means a linear or branched hydrocarbon-based chain containing from 1 to 10 carbon atoms and better still from 1 to 6 carbon atoms, for example from 1 to 4 carbon atoms.

10 **[0031]** Examples of alkyl radicals are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, isopentyl, neopentyl, 2-methylbutyl, 1-ethylpropyl, hexyl, isohexyl, neohexyl, 1-methylpentyl, 3-methylpentyl, 1,1-dimethylbutyl, 1,3-dimethylbutyl, 1-ethylbutyl, 1-methyl-1-ethylpropyl, heptyl, 1-methylhexyl, 1-propylbutyl, 4,4-dimethylpentyl, octyl, 1-methylheptyl, 2-
15 methylhexyl, 5,5-dimethylhexyl, nonyl, decyl, 1-methylnonyl, 3,7-dimethyloctyl and 7,7-dimethyloctyl.

[0032] The alkyl radicals present as substituents of the compounds of the formula (1) according to the present invention may be optionally substituted by one or more chemical species chosen from:

- 20
- halogen atom;
 - -O-alkyl radical;
 - aryl radical;
 - cycloalkyl radical; and
 - heterocyclic radical.

25 **[0033]** The term "alkylene chain" means a divalent radical of linear or branched aliphatic hydrocarbon-based type derived from the alkyl groups defined above by abstraction of a hydrogen atom. Preferred examples of alkylenediyl chains are $-(CH_2)_k-$ chains in which k represents an integer chosen from 1, 2, 3, 4, 5 and 6, and the chains $>CH(CH_3)$, $>C(CH_3)_2$,
30 $-CH_2-CH(CH_3)-CH_2-$ and $-CH_2-C(CH_3)_2-CH_2-$.

[0034] The term "alkenyl radical" means a linear or branched hydrocarbon-based chain containing from 2 to 10 carbon atoms, preferably from 2 to 8 carbon atoms and advantageously from 2 to 6 carbon atoms, containing one, two or more unsaturations in the form of a double bond, the said chain
5 being optionally substituted by one or more substituents, which may be identical or different, chosen from halogen atoms and trifluoromethyl, trifluoromethoxy, hydroxyl, alkoxy, alkoxycarbonyl, carboxyl and oxo radicals.

[0035] Examples of alkenyl radicals that may be mentioned include the ethylenyl radical, the propenyl radical, the isopropenyl radical, the but-2-enyl
10 radical, pentenyl radicals and hexenyl radicals.

[0036] The term "alkynyl radical" means a linear or branched hydrocarbon-based chain containing from 2 to 10 carbon atoms, preferably from 2 to 8 carbon atoms and advantageously from 2 to 6 carbon atoms, containing one, two or more unsaturations in the form of a triple bond, the said chain
15 being optionally substituted by one or more substituents, which may be identical or different, chosen from halogen atoms and trifluoromethyl, trifluoromethoxy, hydroxyl, alkoxy, alkoxycarbonyl, carboxyl and oxo radicals.

[0037] Examples of alkynyl radicals that may be mentioned include the ethynyl radical, the propynyl radical, the but-2-ynyl radical, pentynyl radicals
20 and hexynyl radicals.

[0038] According to the invention, the term "aryl radical" means a monocyclic or polycyclic carbocyclic aromatic radical containing from 6 to 18 carbon atoms and preferably from 6 to 10 carbon atoms. Aryl radicals that may be mentioned include phenyl, naphthyl, anthryl and phenanthryl radicals.

[0039] In the present invention, the cycloalkyl radical is taken to mean a cyclic hydrocarbon-based radical containing from 4 to 9 carbon atoms, preferably 5, 6 or 7 carbon atoms and advantageously 5 or 6 carbon atoms, optionally containing one or more unsaturations in the form of double and/or triple bonds, the said cycloalkyl radical being optionally substituted by one or
25
30 more substituents, which may be identical or different, chosen from halogen

atoms and alkyl, alkenyl, alkynyl, trifluoromethyl, trifluoromethoxy, hydroxyl, alkoxy, alkoxy carbonyl, carboxyl and oxo radicals.

[0040] Preferred examples of cycloalkyl radicals are cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cyclohexadienyl, cycloheptyl, cycloheptenyl and cycloheptadienyl.

[0041] The cycloalkyl radicals are generally monocyclic radicals, but may also be polycyclic, especially bicyclic or tricyclic, optionally containing one or more unsaturations in the form of double bonds.

[0042] The polycyclic cycloalkyl radicals are, for example, tetrahydronaphthyl, perhydronaphthyl, indanyl, bicyclooctyl, bicyclononyl and bicyclodecyl radicals.

[0043] Unless otherwise indicated, the heterocyclic radicals are monocyclic, bicyclic or tricyclic radicals containing one or more hetero atoms generally chosen from O, S and N, optionally in oxidised form (in the case of S and N), and optionally one or more unsaturations in the form of double bonds. If they are totally saturated, the heterocyclic radicals are said to be aromatic or heteroaryl radicals.

[0044] Preferably, at least one of the monocycles constituting the heterocycle contains from 1 to 4 endocyclic hetero atoms and better still from 1 to 3 hetero atoms.

[0045] Preferably, the heterocycle consists of one or more monocycles, each of which is 5- to 8-membered.

[0046] Examples of 5- to 8-membered monocyclic aromatic heterocyclic radicals are the heteroaryl radicals derived, by abstraction of a hydrogen atom, from aromatic heterocycles, such as pyridine, furan, thiophene, pyrrole, imidazole, thiazole, isoxazole, isothiazole, furazane, pyridazine, pyrimidine, pyrazine, thiazines, oxazole, pyrazole, oxadiazole, triazole and thiadiazole.

[0047] Preferred aromatic heterocyclic radicals that may be mentioned include pyridyl, pyrimidinyl, triazolyl, thiadiazolyl, oxazolyl, thiazolyl and thienyl radicals.

[0048] Examples of bicyclic heteroaryls in which each monocycle is 5- to 8-membered are chosen from indolizine, indole, isoindole, benzofuran, benzothiophene, indazole, benzimidazole, benzothiazole, benzofurazane, benzothiofurazane, purine, quinoline, isoquinoline, cinnoline, phthalazine, quinazoline, quinoxaline, naphthyridines, pyrazolotriazines (such as pyrazolo-1,3,4-triazine), pyrazolopyrimidine and pteridine.

[0049] Preferred heteroaryl radicals that may be mentioned include the quinolyl, pyridyl, benzothiazolyl and triazolyl radicals.

[0050] The tricyclic heteroaryls in which each monocycle is 5- to 8-membered are chosen, for example, from acridine, phenazine and carbazole.

[0051] Saturated or unsaturated, 5- to 8-membered monocyclic heterocycles are the saturated or, respectively, unsaturated derivatives of the aromatic heterocycles mentioned above.

[0052] More particularly, mention may be made of morpholine, piperidine, thiazolidine, oxazolidine, tetrahydrothienyl, tetrahydrofuryl, pyrrolidine, isoxazolidine, imidazolidine and pyrazolidine.

[0053] The aryl and heterocyclic radicals are optionally substituted by one or more of the following radicals G:

trifluoromethyl; trifluoromethoxy; styryl; halogen atom; monocyclic, bicyclic or tricyclic aromatic heterocyclic radical containing one or more hetero atoms chosen from O, N and S; and optionally substituted by one or more radicals T as defined below; group Het-CO- in which Het represents an aromatic heterocyclic radical as defined above, optionally substituted by one or more radicals T; a C₁-C₆ alkylene chain; a C₁-C₆ alkylenedioxy chain; nitro; cyano; (C₁-C₁₀)alkyl; (C₁-C₁₀)alkylcarbonyl; (C₁-C₁₀)alkoxycarbonyl-A- in which A represents (C₁-C₆)alkylene, (C₂-C₆)alkenylene or a bond; (C₃-C₁₀)cycloalkyl; trifluoromethoxy; di(C₁-C₁₀)alkylamino; (C₁-C₁₀)alkoxy(C₁-C₁₀)alkyl; (C₁-C₁₀)alkoxy; (C₆-C₁₈)aryl optionally substituted by one or more radicals T; (C₆-C₁₈)aryl(C₁-C₁₀)alkoxy(CO)_n- in which n is 0 or 1 and aryl is optionally substituted by one or more radicals T; (C₆-C₁₈)aryloxy-(CO)_n- in which n is 0 or 1 and aryl is optionally substituted by one or more radicals T;

(C₆-C₁₈)arylthio in which aryl is optionally substituted by one or more radicals T; (C₆-C₁₈)aryloxy(C₁-C₁₀)alkyl(CO)_n- in which n is 0 or 1 and aryl is optionally substituted by one or more radicals T; a saturated or unsaturated, 5- to 8-membered monocyclic heterocycle containing one or more hetero atoms
 5 chosen from O, N and S, optionally substituted by one or more radicals T; (C₆-C₁₈)arylcarbonyl optionally substituted by one or more radicals T; (C₆-C₁₈)arylcarbonyl-B-(CO)_n- in which n is 0 or 1; B represents (C₁-C₆)-alkylene or (C₂-C₆)alkenylene and aryl is optionally substituted by one or more radicals T; (C₆-C₁₈)aryl-C-(CO)_n- in which n is 0 or 1, C represents
 10 (C₁-C₆)alkylene or (C₂-C₆)alkenylene and aryl is optionally substituted by one or more radicals T; (C₆-C₁₈)aryl fused with a saturated or unsaturated heterocycle as defined above, optionally substituted by one or more radicals T; and (C₂-C₁₀)alkynyl.

[0054] T is chosen from a halogen atom; (C₆-C₁₈)aryl; (C₁-C₆)alkyl;
 15 (C₁-C₆)alkoxy; (C₁-C₆)alkoxy(C₆-C₁₈)aryl; nitro; carboxyl; (C₁-C₆)alkoxy-carboxyl; and T may represent oxo if it substitutes a saturated or unsaturated heterocycle; or alternatively T represents (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkyl; or (C₁-C₆)alkylcarbonyl((C₁-C₆)alkyl)_n- in which n is 0 or 1.

[0055] The term "halogen atom" means a chlorine, bromine, iodine or
 20 fluorine atom, preferably fluorine or chlorine.

[0056] Among the compounds of the formula (1), the ones that are preferred are those for which R¹ represents -O-R'¹ and most particularly those for which R¹ represents -O-R'¹, R'¹ being a hydrogen atom or an alkyl radical.

[0057] A first preferred group of compounds of the invention consists of
 25 compounds having one or more of the following characteristics, taken separately or as a combination of one, several or all of them:

R¹ represents -O-R'¹, R'¹ being chosen from a hydrogen atom and an alkyl radical;

R² represents an alkyl radical optionally substituted by a radical -O-alkyl, aryl or cycloalkyl, or alternatively represents an optionally substituted
 30 aryl radical, or alternatively an optionally substituted heterocyclic radical;

R^3 is chosen from a hydrogen atom and an alkyl radical optionally substituted by a radical -O-alkyl, aryl or cycloalkyl; and

A represents a linear or branched alkylene chain containing from 1 to 6 carbon atoms;

5 the possible optical isomers, oxide forms and solvates thereof, and also pharmaceutically acceptable addition salts thereof with acids or bases.

[0058] Another even more preferred group of compounds of the invention consists of compounds having one or more of the following characteristics, taken separately or as a combination of one, several or all of them:

10 R^1 represents -O- R'^1 , R'^1 being chosen from a hydrogen atom and an alkyl radical containing from 1 to 6 carbon atoms;

R^2 represents an alkyl radical containing from 1 to 6 carbon atoms and optionally substituted by a radical -O-alkyl containing from 1 to 6 carbon atoms, or substituted by a phenyl radical or a 5- or 6-membered cycloalkyl radical, or alternatively represents an optionally substituted phenyl radical, or
15 alternatively an optionally substituted heterocyclic radical;

R^3 is chosen from a hydrogen atom and an alkyl radical containing from 1 to 6 carbon atoms and optionally substituted by a radical -O-alkyl containing from 1 to 6 carbon atoms, or substituted by a phenyl radical or a
20 5- or 6-membered cycloalkyl radical; and

A represents a linear or branched alkylene chain containing from 1 to 6 carbon atoms;

the possible optical isomers, oxide forms and solvates thereof, and also pharmaceutically acceptable addition salts thereof with acids or bases.

25 **[0059]** Another preferred group of compounds of the invention consists of compounds having one or more of the following characteristics, taken separately or as a combination of one, several or all of them:

R^1 represents -O- R'^1 , R'^1 being chosen from a hydrogen atom, a methyl radical and an ethyl radical;

30 R^2 represents an alkyl radical containing from 1 to 6 carbon atoms and optionally substituted by a methoxy or ethoxy radical, or substituted by a

substituted phenyl radical or a cyclopentyl or cyclohexyl radical, or alternatively represents an optionally substituted phenyl radical, or alternatively an optionally substituted aromatic heterocyclic radical containing at least one nitrogen atom;

5 R^3 is chosen from a hydrogen atom and an alkyl radical containing from 1 to 6 carbon atoms and optionally substituted by a methoxy or ethoxy radical, or substituted by a substituted phenyl radical or a cyclopentyl or cyclohexyl radical; and

A represents an alkylene chain of the formula $-(CH_2)_k-$, in which k
10 represents an integer between 1 and 6, limits inclusive, or a $-C(CH_3)_2-$ chain;

the possible optical isomers, oxide forms and solvates thereof, and also pharmaceutically acceptable addition salts thereof with acids or bases.

[0060] Another even more preferred group of compounds of the invention consists of compounds having one or more of the following characteristics,
15 taken separately or as a combination of one, several or all of them:

R^1 represents a hydrogen atom;

R^2 is chosen from methyl, ethyl, propyl and n-hexyl radicals, optionally substituted by a methoxy or ethoxy radical, or substituted by a substituted phenyl radical, or with a cyclopentyl radical, or alternatively represents
20 an optionally substituted phenyl radical, or alternatively an optionally substituted pyridyl radical;

R^3 is chosen from a hydrogen atom, a methyl radical, an ethyl radical, a propyl radical, an isopropyl radical, an isobutyl radical and an isopentyl radical, each of these radicals being optionally substituted by a methoxy or
25 ethoxy radical, or with a substituted phenyl radical; and

A represents an alkylene chain of the formula $-(CH_2)_k-$, in which k represents 1, 2 or 3, or a $-C(CH_3)_2-$ chain;

the possible optical isomers, oxide forms and solvates thereof, and also pharmaceutically acceptable addition salts thereof with acids or bases.

[0061] The substituents on the aryl and heterocyclic radicals are preferably chosen from halogen atoms, preferably fluorine and/or chlorine, and methyl, ethyl, methoxy, phenyl, trifluoromethyl and trifluoromethoxy radicals.

[0062] The heterocyclic radicals are preferentially chosen from thienyl, benzothiophenyl, pyridyl and oxazolyl radicals.

[0063] More particularly, the preferred compounds of the formula (1) are those chosen from:

- ethyl 1-benzyl-5-(3-carboxypropoxy)-2-methyl-6-pyrid-3-yl-1*H*-indole-3-carboxylate;
- 10 • ethyl 5-(3-carboxypropoxy)-1-(4-chlorobenzyl)-2-methyl-6-pyrid-3-yl-1*H*-indole-3-carboxylate;
- ethyl 5-(1-carboxy-1-methylethoxy)-2-methyl-1-(3-methylbutyl)-6-pyrid-4-yl-1*H*-indole-3-carboxylate;
- ethyl 5-carboxymethoxy-6-hexyl-1-isobutyl-2-methyl-1*H*-indole-3-carboxylate;
- 15 • ethyl 5-carboxymethoxy-6-(3-ethoxypropyl)-1-isobutyl-2-methyl-1*H*-indole-3-carboxylate;
- ethyl 5-carboxymethoxy-6-(3-cyclopentylpropyl)-1-(2-methoxyethyl)-2-methyl-1*H*-indole-3-carboxylate;
- 20 • ethyl 5-carboxymethoxy-6-[2-(4-fluorophenyl)ethyl]-1-(2-methoxyethyl)-2-methyl-1*H*-indole-3-carboxylate;
- ethyl 5-(3-carboxypropoxy)-6-hexyl-1-(2-methoxyethyl)-2-methyl-1*H*-indole-3-carboxylate;
- ethyl 5-(3-carboxypropoxy)-6-(3-ethoxypropyl)-1-(2-methoxyethyl)-2-methyl-1*H*-indole-3-carboxylate; and
- 25 • ethyl 5-(3-carboxypropoxy)-6-(3-cyclopentylpropyl)-1-(2-methoxyethyl)-2-methyl-1*H*-indole-3-carboxylate;

and the possible optical isomers, oxide forms and solvates thereof, and also pharmaceutically acceptable addition salts of these compounds with acids or bases.

[0064] The invention also relates to pharmaceutical compositions comprising a pharmaceutically effective amount of at least one compound of the formula (1) as defined above in combination with one or more pharmaceutically acceptable vehicles.

5 **[0065]** These compositions can be administered orally in the form of tablets, gel capsules or granules with immediate release or controlled release, intravenously in the form of an injectable solution, transdermally in the form of an adhesive transdermal device, or locally in the form of a solution, cream or gel.

10 **[0066]** A solid composition for oral administration is prepared by adding to the active principle a filler and, where appropriate, a binder, a disintegrant, a lubricant, a dye or a flavour enhancer, and by forming the mixture into a tablet, a coated tablet, a granule, a powder or a capsule.

[0067] Examples of fillers include lactose, corn starch, sucrose, glucose, sorbitol, crystalline cellulose and silicon dioxide, and examples of binders include poly(vinyl alcohol), poly(vinyl ether), ethylcellulose, methylcellulose, acacia, gum tragacanth, gelatine, shellac, hydroxypropylcellulose, hydroxypropylmethylcellulose, calcium citrate, dextrin and pectin. Examples of lubricants include magnesium stearate, talc, polyethylene glycol, silica and hardened plant oils. The dye can be any dye permitted for use in medicaments. Examples of flavour enhancers include cocoa powder, mint in herb form, aromatic powder, mint in oil form, borneol and cinnamon powder. Needless to say, the tablet or granule may be appropriately coated with sugar, gelatine or the like.

25 **[0068]** An injectable form comprising the compound of the present invention as active principle is prepared, where appropriate, by mixing the said compound with a pH regulator, a buffer, a suspending agent, a solubilising agent, a stabiliser, a tonicity agent and/or a preserving agent, and by converting the mixture into a form for intravenous, subcutaneous or intramuscular injection according to a standard process. Where appropriate, the
30 injectable form obtained can be freeze-dried via a standard process.

[0069] Examples of suspending agents include methylcellulose, polysorbate 80, hydroxyethylcellulose, acacia, powdered gum tragacanth, sodium carboxymethyl cellulose and polyethoxylated sorbitan monolaurate.

[0070] Examples of solubilising agents include castor oil solidified with polyoxyethylene, polysorbate 80, nicotinamide, polyethoxylated sorbitan monolaurate and the ethyl ester of castor oil fatty acid.

[0071] In addition, the stabiliser encompasses sodium sulfite, sodium metasilicate and ether, while the preserving agent encompasses methyl p-hydroxybenzoate, ethyl p-hydroxybenzoate, sorbic acid, phenol, cresol and chlorocresol.

[0072] The present invention also relates to the use of a compound of the formula (1) of the invention for the preparation of a medicament for the prevention or treatment dyslipidaemia, atherosclerosis and diabetes.

[0073] The effective administration doses and posologies of the compounds of the invention, intended for the prevention or treatment of a disease, condition or state caused by or associated with modulation of the activity of the PPARs, depends on a large number of factors, for example on the nature of the modulator, the size of the patient, the desired aim of the treatment, the nature of the pathology to be treated, the specific pharmaceutical composition used and the observations and conclusions of the treating doctor.

[0074] For example, in the case of an oral administration, for example, a tablet or a gel capsule, a possible suitable dosage of the compounds of the formula (1) is between about 0.1 mg/kg and about 100 mg/kg of body weight per day, preferably between about 0.5 mg/kg and about 50 mg/kg of body weight per day, more preferentially between about 1 mg/kg and about 10 mg/kg of body weight per day and more preferably between about 2 mg/kg and about 5 mg/kg of body weight per day of active material.

[0075] If representative body weights of 10 kg and 100 kg are considered in order to illustrate the daily oral dosage range that can be used and as described above, suitable dosages of the compounds of the formula (1) will

be between about 1-10 mg and 1000-10 000 mg per day, preferably between about 5-50 mg and 500-5000 mg per day, more preferably between about 10.0-100.0 mg and 100.0-1000.0 mg per day and even more preferentially between about 20.0-200.0 mg and about 50.0-500.0 mg per day of active
5 material comprising a preferred compound.

[0076] These dosage ranges represent total amounts of active material per day for a given patient. The number of administrations per day at which a dose is administered may vary within wide proportions as a function of pharmacokinetic and pharmacological factors, such as the half-life of the
10 active material, which reflects its rate of catabolism and of clearance, and also the minimum and optimum levels of the said active material reached in the blood plasma or other bodily fluids of the patient and which are required for therapeutic efficacy.

[0077] Many other factors should also be considered in deciding upon the
15 number of daily administrations and the amount of active material that should be administered at a time. Among these other factors, and not the least of which, is the individual response of the patient to be treated.


[0078] The present invention also relates to a general process for the preparation of the compounds of the formula (1), starting with ethyl 6-bromo-
20 5-hydroxy-2-methyl-1*H*-indole-3-carboxylate, the hydroxyl function of which is subjected to the action of a compound of the formula (2):



in which A is as defined above for the compounds of the formula (1) and R represents a protecting group for the acid function, for example an
25 alkyl radical, such as methyl or ethyl,


in the presence of a base, such as an alkali metal or alkaline-earth metal hydroxide or carbonate, for example sodium hydroxide or potassium carbonate (K_2CO_3), in polar aprotic medium, for example in dimethylformamide (DMF) solvent,

30 to give the compound of the formula (3):



in which A and R are as defined above,


of which compound of the formula (3) the nitrogen atom may optionally be substituted, under the action of a bromide R³-Br, in which R³ is as defined for the compounds of the formula (1), under conditions similar to those outlined above (for example NaOH or K₂CO₃/DMF), so as to obtain the compound of the formula (4):



in which A, R³ and R are as defined above,


and then subjected to a Heck reaction (R. F. Heck et al., *J. Org. Chem.*, (1972), **37**, 2320 sqq.) in which a compound of the formula R'-CH=CH-R'' (precursor of the radical R²-, radical R²- possibly being represented by the radical R'-CH₂-CH-R'') is treated with a borane, for example 9-borabicyclo[3.3.1]nonane, to give an R²-borane, in which R² is as defined for the compounds of the formula (1),

which is then coupled with the compound of the formula (3), in the presence of a palladium catalyst, such as 1,1'-bis(diphenylphosphino)ferrocenedichloropalladium II (PdCl₂dppf), in basic medium, for example in potassium phosphate, in a polar aprotic solvent, for instance tetrahydrofuran, to give the compound of the formula (1_R):



in which A, R², R³ and R are as defined above,

which compound of the formula (1_R) is then converted, according to standard techniques known to those skilled in the art, into the corresponding acid of the formula (1_{OH}):



which is a special case of the compounds of the formula (1) in which R¹ represents a hydroxyl group,

and the acid is optionally esterified, or converted into the corresponding amide, also according to standard techniques, to form the set of compounds of the formula (1), with R¹ other than a hydroxyl group.


[0079] This synthetic method applies to all the compounds of the formula (1) according to the present invention, and is described in greater detail in the synthesis of the compounds of Examples 1, 2, 12 and 13 below.

[0080] It should be understood that the compounds of the formula (1_R) above, if R represents an alkyl radical, form part of the compounds of the formula (1) according to the present invention.

[0081] If such compounds are desired, the steps of deprotection of the acid function and then of esterification are superfluous.


[0082] According to one variant, the compounds of the formula (1) can also be prepared from ethyl 6-bromo-5-hydroxy-2-methyl-1*H*-indole-3-car-

boxylate, the hydroxyl function of which is protected in a conventional manner, for example by reaction with acetic anhydride, in the presence of pyridine, and the nitrogen atom is then optionally substituted, under the action of a bromide R^3 -Br, as indicated for the production of compound (4) defined above, and the hydroxyl function is then deprotected, in the presence of a base and in an alcohol, for example sodium hydroxide in methanol, so as to obtain the compound of the formula (5):




in which R^3 is as defined above,

for which compound of the formula (5) the synthesis upto the compound of the formula (1) is continued by means of a technique of synthesis on resin, for example of grafted Wang type and corresponding to formula (6):




in which A is as defined for the compounds of the formula (1) and represents the resin support,

which, when placed in contact with compound (5), in basic medium and a polar aprotic solvent, for example K_2CO_3 /DMF, in the presence of potassium iodide, gives the compound of the formula (7):




in which A, R³ and  are as defined above,
the bromine atom of which is replaced with the substituent R², under the
same operating conditions as those described above for the production of the
compound of the formula (1_R), thus leading to the compound of the formula
5 (8):



in which A, R², R³ and  are as defined above,
which is then detached from the resin support according to the usual condi-
tions, for example using trifluoroacetic acid, so as to give the compound of
the formula (1_{OH}) described above, which is then, where appropriate, esteri-
fied or converted into the corresponding amide, to form the set of compounds
10 of the formula (1) with R¹ other than a hydroxyl group.

[0083] This method is more particularly detailed in the preparation of the
compounds of Examples 3, 14-17 and 29-31.

15 **[0084]** According to one alternative, the attachment to the grafted resin
(6) can be performed directly on ethyl 6-bromo-5-hydroxy-2-methyl-1H-
indole-3-carboxylate, so as to obtain the bromo derivative of the formula (9):



in which A and  are as defined above,

20 in which derivative of the formula (9) the nitrogen atom may be
optionally substituted by the radical R³, and the bromine atom replaced with

the radical R^2 (in any order), according to the techniques outlined above for the preparations of the compounds of the formulae (5) and (8), respectively.


[0085] This method is more particularly detailed in the preparation of the compounds of Examples 4-11 and 18-28.

5 **[0086]** The compounds of the formula (1) in which R^1 represents H can advantageously be obtained by saponification of the corresponding compounds of the formula (1) in which R^1 represents an alkyl radical, or alternatively starting with the compounds of the formula (1_R), in which R represents an alkyl radical. The saponification can be performed via the action of a base,
10 such as a mineral base chosen from lithium hydroxide, potassium hydroxide, sodium hydroxide, sodium hydrogen carbonate, potassium hydrogen carbonate, sodium carbonate and potassium carbonate. The molar amount of base to be used generally ranges from 1 to 20 equivalents and preferably from 1 to 12 equivalents depending on the strength of the selected base.

15 **[0087]** The reaction is preferably performed in a solvent of polar protic type and more preferably in a mixture of a lower (C₁-C₄) alkanol and water, such as a mixture of ethanol and water or methanol and water.

[0088] The reaction temperature advantageously ranges between room temperature and 120°C and better still between 20°C and 100°C, for example
20 between 20°C and reflux.

[0089] The compound of the formula (6) can readily be obtained by coupling a resin of Wang type, of the formula (10), with an acid of the formula (11), according to the following reaction scheme:



25 in which A is as defined for the compounds of the formula (1) and  represents the resin support,

according to standard techniques known to those skilled in the art, for example in the presence of 4-dimethylaminopyridine (DMAP) and N,N'-diiso-

propylcarbodiimide (DIC), in apolar organic medium, for example in methylene chloride.

[0090] This grafting of the Wang resin is described in greater detail in Example 3, step 4.

5 **[0091]** In the processes described above, it should be understood that the operating conditions may vary substantially as a function of the various substituents present in the compounds of the formula (1) that it is desired to prepare. Such variations and adaptations are readily accessible to those skilled in the art, for example from scientific reviews, the patent literature, Chemical
10 Abstracts, and computer databases, including the Internet. Similarly, the starting materials are either commercially available or accessible via syntheses that a person skilled in the art can readily find, for example in the various publications and databases described above.

[0092] The optical isomers of the compounds of the formula (1) can be
15 obtained on the one hand via standard techniques for separating and/or purifying isomers known to those skilled in the art, starting with the racemic mixture of the compound of the formula (1). The optical isomers can also be obtained directly via stereoselective synthesis of an optically active starting compound, or via separation or recrystallisation of the optically active salts of
20 the compounds of the formula (1), the salts being obtained with chiral amines or chiral acids.

[0093] The examples that follow illustrate the present invention without limiting it in any way. In these examples and the proton nuclear magnetic resonance data (300 MHz NMR), the following abbreviations have been
25 used: s for singlet, d for doublet, t for triplet, q for quartet, o for octet and m for complex multiplet. The chemical shifts δ are expressed in ppm.

EXAMPLES

Example 1: Ethyl 5-ethoxycarbonylmethoxy-6-hexyl-1-isobutyl-2-methyl-1*H*-
30 indole-3-carboxylate

Step 1

[0094] A mixture of ethyl 6-bromo-5-hydroxy-2-methyl-1*H*-indole-3-carboxylate (5 g; 16.77 mmol), ethyl 2-bromoacetate (4.2 g; 25.16 mmol) and potassium carbonate (2.3 g; 16.89 mmol) in dimethylformamide (DMF) (20 ml) is heated at 60°C for 1 hour 30 minutes. The reaction medium is poured into a mixture of ice and ethyl acetate. The aqueous phase is extracted twice with ethyl acetate. The combined organic phases are washed with water, dried over sodium sulfate and then concentrated. The residue obtained (6.85 g) is purified by chromatography on silica (3.63 g; 57%).

¹H NMR (300 MHz, chloroform-D), δ ppm: 1.2 (t, $J=6.9$ Hz, 3 H); 1.4 (t, $J=7.2$ Hz, 3 H); 2.6 (s, 3 H); 4.2 (m, 4 H); 4.7 (s, 2 H); 7.2 (s, 1 H); 7.3 (s, 1 H); 7.5 (s, 1 H).

Step 2

[0095] A mixture of the compound obtained in step 1 (3.63 g; 9.45 mmol), isobutyl bromide (2.6 g; 18.98 mmol) and potassium carbonate (2.68 g; 19.68 mmol) in DMF (20 ml) is heated for 48 hours. A large excess of the halide, of potassium carbonate and of potassium iodide (0.615 mg) is added and the mixture is heated for a further 48 hours. The reaction medium is then poured into a mixture of ice and ethyl acetate. The aqueous phase is extracted twice with ethyl acetate. The combined organic phases are washed with water, dried over sodium sulfate and then concentrated. The residue obtained (3.73 g) is purified by chromatography on silica (gradient: 0-30% ethyl acetate in heptane). 1.73 g (42%) of the expected product are obtained.

¹H NMR (300 MHz, chloroform-D), δ ppm: 0.9 (d, $J=6.7$ Hz, 6 H); 1.3 (t, $J=7.2$ Hz, 3 H); 1.4 (t, $J=7.2$ Hz, 3 H); 2.2 (m, 1 H); 2.7 (s, 3 H); 3.8 (d, $J=7.8$ Hz, 2 H); 4.3 (q, $J=7.2$ Hz, 2 H); 4.4 (q, $J=7.2$ Hz, 2 H); 4.7 (s, 2 H); 7.5 (s, 1 H); 7.6 (s, 1 H).

Step 3

[0096] Preparation of the organoborane solution: a solution of hexene

(0.42 g; 5 mmol) in tetrahydrofuran (THF) (2 ml) is cooled under nitrogen to 3°C. A commercial 0.5 N solution of 9-borabicyclo[3.3.1]nonane (9-BBN) in THF (10 ml, 5 mmol) is then added dropwise so as to keep the temperature below 5°C. The colourless solution obtained is then stirred for 3 hours at
5 room temperature.

[0097] A mixture of the compound obtained in step 2 (0.208 g; 0.47 mmol), PdCl₂dppf (12.3 mg; 15.1 µmol) and potassium phosphate (237 mg; 0.89 mmol) in THF (2 ml) is refluxed and the organoborane solution (2.3 ml; 0.96 mmol) is then added in a single portion. The medium, which
10 quickly turns black, is refluxed for 1 hour 30 minutes. The reaction medium is then poured into a mixture of water and ethyl ether. The aqueous phase is extracted. The combined organic phases are washed with water, dried over sodium sulfate and then concentrated. The oily residue obtained (0.5 g) is purified by chromatography on silica (4/1 heptane/ethyl acetate) and then
15 dispersed in heptane (82.0 mg; 39%).

¹H NMR (300 MHz, chloroform-D), δ ppm: 0.9 (t, *J*=6.7 Hz, 3 H); 0.9 (d, *J*=6.7 Hz, 6 H); 1.4 (m, 6 H); 1.3 (t, *J*=7.2 Hz, 3 H); 1.4 (t, *J*=7.2 Hz, 3 H); 1.7 (m, 2 H); 2.2 (s, 1 H); 2.8 (m, 2 H); 2.7 (s, 3 H); 3.9 (d, *J*=7.4 Hz, 2 H); 4.3 (q, *J*=7.2 Hz, 2 H); 4.4 (q, *J*=7.2 Hz, 2 H); 4.7 (s, 2 H); 7.0 (s, 1 H); 7.5 (s,
20 1 H).

Example 2: Ethyl 5-carboxymethoxy-6-hexyl-1-isobutyl-2-methyl-1*H*-indole-3-carboxylate

[0098] A mixture of the compound obtained in Example 1 (82.0 mg; 0.184 mmol), methanol (2 ml) and aqueous 1 N sodium hydroxide (0.265 ml; 0.265 mmol) is stirred overnight at room temperature. The solvents are evaporated off. The residue is dissolved in water and then treated with concentrated hydrochloric acid. After extraction with ethyl ether and drying over sodium sulfate, evaporation gives a white solid (71 mg; 92%).

¹H NMR (300 MHz, chloroform-D), δ ppm: 0.9 (m, 3 H); 0.9 (d, *J*=6.7 Hz, 6 H); 1.3 (m, 7 H); 1.4 (t, *J*=7.2 Hz, 3 H); 1.6 (m, 2 H); 2.2 (m, 1 H); 2.8
30

(m, 2 H); 2.7 (s, 3 H); 3.9 (d, $J=7.6$ Hz, 2 H); 4.4 (q, $J=7.2$ Hz, 2 H) . 4.7 (s, 2 H); 7.0 (s, 1 H); 7.5 (s, 1 H).

Example 3: Ethyl 5-(3-carboxypropoxy)-6-hexyl-1-(2-methoxyethyl)-2-methyl-1*H*-indole-3-carboxylate

Step 1

[0099] A mixture composed of ethyl 6-bromo-5-hydroxy-2-methyl-1*H*-indole-3-carboxylate (14.91 g; 50 mmol), acetic anhydride (100 ml; 1.06 mol) and pyridine (12.08 ml; 0.15 mol) is refluxed for one hour. The cooled reaction medium is then poured into saturated sodium bicarbonate solution. After extraction with ethyl acetate, the combined organic phases are washed with water, dried over sodium sulfate and then concentrated. The residue obtained (14.0 g) is dispersed in methylene chloride (8.3 g). The mother liquors, concentrated and then chromatographed on a column of alumina (methylene chloride), give a further amount of the expected product (2.9 g). The two batches (8.3 g and 2.9 g) are combined and dispersed in ethyl ether. 9.6 g of pure product are obtained.

Melting point: 200°C

^1H NMR (300 MHz, chloroform- D), δ ppm: 1.4 (t, $J=7.1$ Hz, 3 H); 2.4 (s, 3 H); 2.6 (s, 3 H); 4.4 (q, $J=7.1$ Hz, 2 H); 7.3 (s, 1 H); 7.7 (s, 1 H); 8.6 (s, 1 H).

Step 2

[0100] A mixture composed of the derivative obtained in step 1 (9.05 g; 26.60 mmol), sodium hydroxide (NaOH) (1.17 g; 29.26 mmol) and 2-methoxyethyl bromide (7.50 ml; 79.81 mmol) in dry DMF is stirred at 40°C for 4 hours. The cooled reaction medium is then poured into ice-cold dilute hydrochloric acid. After extraction with ethyl acetate, the combined organic phases are washed with water and dried over sodium sulfate. The evaporation residue is dispersed in pentane (7.84 g).

¹H NMR (300 MHz, chloroform-D), δ ppm: 1.4 (t, $J=7.1$ Hz, 3 H); 2.4 (s, 3 H); 2.8 (s, 3 H); 3.3 (s, 3 H); 3.6 (t, $J=5.5$ Hz, 2 H); 4.2 (t, $J=5.5$ Hz, 2 H); 4.4 (q, $J=7.1$ Hz, 2 H); 7.5 (s, 1 H); 7.8 (s, 1 H).

5 Step 3

[0101] A mixture composed of the derivative obtained in step 2 (7.81 g; 19.61 mmol) and aqueous 1 N sodium hydroxide (23.5 ml; 23.5 mmol) in methanol (80 ml) is stirred at room temperature for 1 hour. The reaction medium is then poured into dilute hydrochloric acid. The precipitate formed is
10 filtered off, washed and drained by suction (6.65 g; 90%).

Melting point: 195°C

¹H NMR (300 MHz, chloroform-D), δ ppm: 1.4 (t, $J=7.2$ Hz, 3 H); 2.8 (s, 3 H); 3.3 (s, 3 H); 3.6 (t, $J=5.6$ Hz, 2 H); 4.2 (t, $J=5.6$ Hz, 2 H); 4.4 (q, $J=7.2$ Hz, 2 H); 5.3 (s, 1 H); 7.4 (s, 1 H); 7.7 (s, 1 H).

15

Step 4

[0102] Grafting of 4-bromobutyric acid onto the Wang resin:

[0103] A mixture composed of Wang resin (8.24 g; 0.91 mM/g), 4-bromobutyric acid (5.51 g; 33 mmol), 4-dimethylaminopyridine (DMAP) (0.183 g; 1.5 mM) and N,N'-diisopropylcarbodiimide (DIC) (5.14 g; 33 mmol) in methylene chloride (130 ml) is stirred with an orbital stirrer for 20 hours in a 250 ml flask. The resin is washed three times with methylene chloride and then three times with methanol. After drying under vacuum at room temperature, 9.16 g of grafted resin are obtained (theory: 9.36 g).

25 [0104] 2.63 g of the above resin are taken and poured into dry DMF (14 ml). The 5-hydroxy-6-bromoindole derivative obtained in step 3 (1.5 g; 4.21 mmol), potassium iodide (349.5 mg; 2.1 mmol) and potassium carbonate (291 mg; 2.1 mmol) are then successively added. The mixture is heated at 80°C overnight. The resin is filtered off and washed three times with DMF
30 (5 ml), then three times with a 1/1 tetrahydrofuran/water mixture (5 ml), then

three times with tetrahydrofuran (5 ml) and then three times with methanol (5 ml) and finally dried under vacuum (3.14 g, theory: 3.21 g).

Step 5

5 [0105] Preparation of the organoborane solution: a 0.5 N solution of 9-BBN in THF (1.2 ml; 0.60 mmol) is cooled to 0°C and hexene (75 µl; 0.60 mmol) is then added. The colourless solution obtained is then stirred for three hours at room temperature.

[0106] The organoborane solution (0.957 ml; 0.45 mmol) is added to a
10 mixture of resin obtained in step 4 (227.27 mg), Pd(PPh₃)₄ (8.67 mg; 75.0 µmol) and aqueous 2 M sodium carbonate (94 µl; 188 µmol) in DMF (2 ml). The medium is heated at 80°C overnight. The resin is filtered off and washed three times with DMF, then three times with a 1/1 tetrahydrofuran/water mixture, three times with methanol and then three times with
15 methylene chloride and finally dried under vacuum.

Step 6

[0107] The resin obtained in step 5 is treated for 2 hours at room temperature with an 8/2 mixture of methylene chloride/trifluoroacetic acid
20 (2 ml).

[0108] The medium is filtered and the resin is then washed with methylene chloride. The filtrate is concentrated to dryness (10 mg). The resin is then treated again for 2 hours with a 1/1 mixture of methylene chloride/trifluoroacetic acid.

25 [0109] After filtration and washing, evaporation of the filtrate gives an additional 18 mg of product (total 28 mg).

¹H NMR (300 MHz, chloroform-D), δ ppm: 0.9 (m, 3 H); 1.3 (m, 6 H); 1.4 (t, J=7.2 Hz, 3 H); 1.6 (m, 2 H); 2.2 (m, 2 H); 2.7 (m, 7 H); 3.3 (s, 3 H); 3.6 (t, J=5.7 Hz, 2 H); 4.1 (t, J=5.8 Hz, 2 H); 4.3 (t, J=5.6 Hz, 2 H); 4.4 (m, 2 H);
30 5.3 (s, 1 H) 7.0 (s, 1 H); 7.6 (s, 1 H).

LC/MS: ES+ 448.5 ES- 446.4

Example 4: Ethyl 1-benzyl-5-(3-carboxypropoxy)-2-methyl-6-pyrid-3-yl-1*H*-indole-3-carboxylate

Step 1

5 [0110] 2.85 g of grafted Wang resin (loading 1.09 mM/g) are taken up with 4-bromobutyric acid (preparation identical to that described in step 4 of Example 3, but starting with a 1.3 mM/g Wang resin) and poured into dry DMF (58 ml). Next, ethyl 6-bromo-5-hydroxy-2-methyl-1*H*-indole-3-carboxy-
10 late (3.71 g; 12.44 mmol), potassium iodide (0.52 g; 3.13 mmol) and potassium carbonate (0.43 g; 3.11 mmol) are successively added. The mixture is heated at 80°C for 16 hours. The resin is filtered off and washed twice with DMF, three times with a 1/1 tetrahydrofuran/water mixture, three times with tetrahydrofuran (THF) and three times with methanol and finally dried under vacuum (3.37 g, theory: 3.52 g).

15

Step 2

[0111] 602 mg of the above resin (loading 0.88 mM/g) in DMF (5 ml) are treated with sodium hydride (NaH) at 60% in oil (63.3 mg; 1.58 mmol) for 15 minutes, and potassium iodide (88 mg; 0.53 mmol) and benzyl bromide
20 (362.6 mg; 2.11 mmol) are then added. The mixture is stirred for 20 hours at room temperature under nitrogen.

[0112] Ethyl acetate (0.5 ml) is added and the resin is then filtered off and washed three times with DMF (10 ml), three times with 1/1 THF/H₂O, three times with THF and three times with methanol and finally dried under vac-
25 uum.

Step 3

[0113] A mixture of the above resin (146.3 mg; loading 0.82 mM/g), PdP(Ph₃)₄ (27.7 mg; 24 µmol), aqueous 2 M sodium carbonate (120 µl; 240 mmol) and 3-pyridylboronic acid (59 mg; 480 µmol) in DMF (2 ml) is
30

heated at 120°C for 12 hours. The mixture is stirred for 20 hours at room temperature under nitrogen.

[0114] The resin is filtered off and washed three times with DMF (2 ml), three times with 1/1 THF/H₂O, three times with THF, three times with methanol and three times with methylene chloride and finally dried under vacuum.

[0115] The resin obtained is treated with an 8/2 mixture of methylene chloride/trifluoroacetic acid (1.5 ml) for 1.5 hours at room temperature.

[0116] The medium is filtered and the resin is then washed with methylene chloride. The filtrate is concentrated to dryness (39 mg).

10 LC/MS: ES+ 473.2

Example 5: Ethyl 5-(3-carboxypropoxy)-2-methyl-6-(3,4-dichlorophenyl)-1*H*-indole-3-carboxylate

15 [0117] A mixture of the resin obtained in step 1 of Example 4 (144 mg; loading 0.94 mmol/g), a 0.025 M solution of PdP(Ph₃)₄ in DMF (1 ml; 25 µmol), aqueous 2M sodium carbonate (135 µl; 270 mmol) and 3,4-dichlorophenylboronic acid (52 mg; 270 µmol) in DMF (2 ml) is heated at 120°C for 16 hours. The resin is filtered off and washed six times with DMF
20 (3 ml), six times with dimethyl sulfoxide (DMSO) (3 ml), three times with water (3 ml), three times with methanol (3 ml) and four times with methylene chloride (3 ml).

[0118] The resin suspended in methylene chloride (1 ml) is then treated with a 6/4 mixture of methylene chloride/trifluoroacetic acid (2 ml) for 1 hour
25 at room temperature.

[0119] The medium is filtered and the resin is then washed with methylene chloride. The filtrate is concentrated to dryness (20 mg).

LC/MS: ES+ 450.3 452.3 454.2. 2 chlorine atoms.

30 [0120] Compounds 6 to 31 were prepared according to protocols similar to those described for the preparation of the compounds of Examples 1 to 5


above.

[0121] The structures of compounds 6 to 31 are collated in Table 1 below, in which "Method" specifies the method number (1, 2 or 3 described above) used for the preparation of each compound:

















5













-- TABLE 1 --

Structures of compounds 6 to 31



Ex. No.	R ¹	A	R ²	R ³	Method
6	-H	-(CH ₂) ₃ -			3
7	-H	-(CH ₂) ₃ -		-H	3
8	-H	-C(CH ₃) ₂ -		-H	3
9	-H	-C(CH ₃) ₂ -		-H	3
10	-H	-C(CH ₃) ₂ -			3
11	-H	-C(CH ₃) ₂ -			3

Ex. No.	R^1	A	R^2	R^3	Method
12	-H	-CH ₂ -			1
13	-H	-CH ₂ -			1
14	-H	-CH ₂ -			2
15	-H	-CH ₂ -	-(CH ₂) ₅ -CH ₃		2
16	-H	-CH ₂ -			2
17	-H	-CH ₂ -			2
18	-H	-(CH ₂) ₃ -		-H	3
19	-H	-(CH ₂) ₃ -		-H	3
20	-H	-(CH ₂) ₃ -		-H	3
21	-H	-(CH ₂) ₃ -		-H	3
22	-H	-C(CH ₃) ₂ -		-H	3

Ex. No.	R^1	A	R^2	R^3	Method
23	-H	-C(CH ₃) ₂ -		-H	3
24	-H	-C(CH ₃) ₂ -		-H	3
25	-H	-C(CH ₃) ₂ -		-H	3
26	-H	-C(CH ₃) ₂ -		-H	3
27	-H	-C(CH ₃) ₂ -		-H	3
28	-H	-C(CH ₃) ₂ -		-H	3
29	-H	-(CH ₂) ₃ -			2
30	-H	-(CH ₂) ₃ -			2
31	-H	-(CH ₂) ₃ -			2

[0122] The results of the analyses of the synthesised products 6 to 31 are given in Table 2 below, in which table:

- M represents the theoretical molar mass of the compound;
- 5 - LC/MS indicates the result of the analysis by mass spectrometry coupled to liquid-phase chromatography; and

- NMR indicates the chemical shifts δ (in ppm) of the proton by magnetic resonance at 300 MHz.

-- TABLE 2 --

5

Example No	M	LC/MS	NMR
6	506.98	ES+ 507.2/509.2 1 Cl	
7	382.41	ES+ 383.3	
8	395.45	ES- 394.2	
9	450.32	ES- 448.1 / 450.1 2 Cl	
10	521.53	ES+ 522.4	
11	452.55	ES+ 453.4	
12	443.58		¹ H NMR (300 MHz, DMSO-D6), δ ppm: 0.9 (d, J=6.5 Hz, 6 H); 1.0 (m, 2 H); 1.5 (m, 10 H); 2.1 (m, 1 H); 2.7 (m, 5 H); 3.3 (m, 4 H); 4.0 (d, J=7.2 Hz, 2 H); 4.2 (q, J=7.1 Hz, 2 H); 4.7 (s, 2 H); 7.3 (s, 1 H); 7.3 (s, 1 H); 12.9 (s, 1 H).
13	419.51	ES- 418.5	
14	445.55	ES- 444.4	¹ H NMR (300 MHz, chloroform-D), δ ppm: 1.1 (d, J=6.3 Hz, 2 H); 1.6 (m, 14 H); 2.8 (m, 5 H); 3.3 (s, 3 H); 3.7 (t, J=5.6 Hz, 2 H); 4.3 (t, J=5.6 Hz, 2 H); 4.4 (q, J=7.1 Hz, 2 H); 4.8 (s, 2 H); 7.1 (s, 1 H); 7.5 (s, 1 H).
15	419.51	ES+ 420.3	
16	457.50	ES- 456.3	
17	421.49	ES+ 422.3	¹ H NMR (300 MHz, chloroform-D), δ ppm: 1.2 (t, J=7.0 Hz, 3 H); 1.4 (t, J=7.1 Hz, 3 H); 1.9 (m, 2 H); 2.7 (s, 3 H); 2.9 (t, J=7.2 Hz, 2 H); 3.3 (s, 3 H); 3.6 (m, 6 H); 4.3 (t, J=5.6 Hz, 2 H); 4.4 (q, J=6.9 Hz, 2 H); 4.7 (s, 2 H); 6.8 (s, 1 H); 7.1 (s, 1 H); 7.5 (s, 1 H).
18	449.42	ES+ 450.3	

Example No	M	LC/MS	NMR
19	449.42	ES- 448.4	
20	465.42	ES- 464.4	
21	415.87	ES+ 416.3	
22	425.43	ES- 424.3	
23	449.42	ES+ 450.4	
24	449.42	ES+ 450.3	
25	465.42	ES+ 466.3	
26	411.45	ES+ 412.4	
27	411.45	ES+ 412.4	
28	415.87	ES- 414.3 /416.3 1 Cl	
29	449.54	ES+ 450.4	¹ H NMR (300 MHz, chloroform-D), δ ppm: 1.3 (t, J=6.8 Hz, 3 H); 1.4 (t, J=7.0 Hz, 3 H); 1.9 (m, 2 H); 2.2 (m, 2 H); 2.6 (t, J=6.6 Hz, 2 H); 2.8 (m, 5 H); 3.3 (s, 3 H); 3.6 (m, 6 H); 4.1 (t, J=5.5 Hz, 2 H); 4.3 (t, J=5.7 Hz, 2 H); 4.4 (q, J=7 Hz, 2 H); 7.0 (s, 1 H); 7.6 (s, 1 H).
30	485.55	ES+ 486.4	¹ H NMR (300 MHz, chloroform-D), δ ppm: 1.4 (t, J=7.1 Hz, 2 H); 2.2 (m, 2 H); 2.7 (m, 6 H); 2.9 (m, 4 H); 3.3 (s, 3 H); 3.5 (t, J=5.7 Hz, 2 H); 4.2 (m, 4 H); 4.4 (q, J=7.1 Hz, 2 H); 6.0 (s, 1 H); 6.8 (s, 1 H); 6.9 (m, , 2 H); 7.1 (m, 2 H); 7.6 (s, 1 H).
31	473.61	ES+ 474.5	¹ H NMR (300 MHz, chloroform-D), δ ppm: 1.5 (m, 16 H); 2.2 (m, 2 H); 2.7 (m, 7 H); 3.3 (s, 3 H); 3.7 (t, J=5.6 Hz, 2 H); 4.1 (m, 2 H); 4.3 (t, J=5.7 Hz, 2 H); 4.4 (m, 2 H); 7.0 (s, 1 H); 7.6 (s, 1 H).

RESULTS

[0123] The measurement of the PPAR activation was performed accord-
5 ing to a technique described by Lehmann et al. (*J. Biol. Chem.*, **270**, (1995),
12953-12956).

[0124] CV-1 cells (monkey kidney cells) are cotransfected with an expression vector for the chimeric protein PPAR γ -Gal4 and with a "reporter" plasmid that allows expression of the luciferase gene placed under the control of a promoter comprising Gal4 response elements.

5 **[0125]** The cells are seeded in 96-well microplates and cotransfected using a commercial reagent with the reporter plasmid (pG5-tk-pGL3) and the expression vector for the chimeric protein (PPAR γ -Gal4). After incubation for 4 hours, whole culture medium (comprising 10% foetal calf serum) is added to the wells. After 24 hours, the medium is removed and replaced with whole
10 medium comprising the test products. The products are left in contact with the cells for 18 hours. The cells are then lysed and the luciferase activity is measured using a luminometer. A PPAR γ activation factor can then be calculated by means of the activation of the expression of the reporter gene induced by the product (relative to the control cells that have received no
15 product).

[0126] In the absence of the PPAR γ ligand binding domain (vector expressing Gal4 alone), the luciferase activity measured in the presence of an agonist is zero.

[0127] The following transactivation result was obtained with a concentration of 50 μ M on PPAR γ .
20

<i>Ex.</i>	<i>Concentration</i>	<i>Activation factor of the chimeric protein PPARγ-Gal4</i>
4	50 μ M	17
Without agonist (Control)	-	1

Example of biological activities of partial agonists

25 Transactivation test

[0128] The transactivation test using the expression of a chimeric protein Gal-4-PPAR γ makes it possible to determine also whether an agonist func-

tions as a "full" agonist or as a "partial" agonist in this system.

[0129] An agonist is "partial" in this system if it induces a weaker response, i.e. it has lower efficacy, than rosiglitazone, which is a "full" agonist. In concrete terms, in our system, the transactivation obtained at the plateau
5 with a partial agonist will be between 20% and 50% of the maximum response (efficacy) at the plateau of rosiglitazone.


Ex.	Maximum stimulation of the PPAR γ chimeric protein obtained with rosiglitazone	Concentration to reach the maximum stimulation of the PPAR γ chimeric protein
31	24%	1 μ M

26474-1095

39

CLAIMS:

1. A compound of formula (1):



- or an optical isomer thereof, an oxide form thereof, or a
5 pharmaceutically acceptable addition salt thereof,

in which:

R^1 is $-O-R^{1'}$ or $-NR^{1'}R^{''1}$;

- $R^{1'}$ and $R^{''1}$ are each independently H, optionally substituted alkyl
having 1 to 10 carbon atoms, optionally substituted alkenyl having 2 to 10 carbon
10 atoms, optionally substituted alkynyl having 2 to 10 carbon atoms, optionally
substituted cycloalkyl having 4 to 9 carbon atoms, optionally substituted aryl having 6
to 18 carbon atoms, or optionally substituted heteroaryl;

- R^2 is optionally substituted alkyl having 1 to 10 carbon atoms, optionally
substituted alkenyl having 2 to 10 carbon atoms, or optionally substituted alkynyl
15 having 2 to 10 carbon atoms, aryl having 6 to 18 carbon atoms which is optionally
substituted and/or optionally fused to a monocyclic or polycyclic, saturated or
unsaturated 5- to 8-membered nucleus optionally containing one or more hetero
atoms which are O, N or S, and said nucleus is optionally substituted, or a saturated,
unsaturated or aromatic, optionally substituted 5- to 8-membered monocyclic
20 heterocyclic radical containing one or more hetero atoms which are O, N or S;

R^3 is H or optionally substituted alkyl having 1 to 10 carbon atoms; and

26474-1095

40

A is a linear or branched alkylene chain containing from 1 to 6 carbon atoms;

wherein optionally substituted alkyl groups listed above are each optionally substituted one or more times by halogen, -O-alkyl having 1 to 10 carbon atoms, aryl having 6 to 18 carbon atoms and which is optionally substituted by halogen, methyl, ethyl, methoxy, phenyl, trifluoromethyl or trifluoromethoxy, cycloalkyl having 4 to 9 carbon atoms, or a heterocyclic group;

wherein optionally substituted alkenyl groups listed above are each unsubstituted or substituted one or more times by halogen, trifluoromethyl, trifluoromethoxy, hydroxyl, alkoxy, alkoxycarbonyl, carboxyl or oxo;

wherein optionally substituted alkynyl groups listed above are each unsubstituted or substituted one or more times by halogen, trifluoromethyl, trifluoromethoxy, hydroxyl, alkoxy, alkoxycarbonyl, carboxyl or oxo;

wherein optionally substituted aryl groups listed above are each unsubstituted or substituted one or more times by halogen, methyl, ethyl, methoxy, phenyl, trifluoromethyl or trifluoromethoxy;

wherein optionally substituted heterocyclic groups listed above are each unsubstituted or substituted one or more times by halogen, methyl, ethyl, methoxy, phenyl, trifluoromethyl or trifluoromethoxy;

wherein aromatic heterocyclic groups or heteroaryl groups are selected from pyridyl, pyrimidinyl, triazolyl, thiadiazolyl, oxazolyl, thiazolyl, thienyl, quinolyl, benzothiazolyl, acridinyl, phenazinyl, carbazolyl, and benzothiophenyl;

wherein saturated or unsaturated heterocyclic groups are selected morpholinyl, piperidinyl, thiazolidinyl, oxazolidinyl, tetrahydrothienyl, tetrahydrofuryl, pyrrolidinyl, isoxazolidinyl, imidazolidinyl, and pyrazolidinyl; and

26474-1095

41

wherein optionally substituted cycloalkyl is unsubstituted or substituted one or more times by halogen atoms, alkyl, alkenyl, alkynyl, trifluoromethyl, trifluoromethoxy, hydroxyl, alkoxy, alkoxycarbonyl, carboxyl, or oxo.

2. A compound according to claim 1, or an optical isomer thereof, an oxide
5 form thereof, or a pharmaceutically acceptable addition salt thereof, wherein:

R^1 is -O- R'^1 , and R'^1 is H or alkyl;

R^2 is alkyl optionally substituted by -O-alkyl, aryl or cycloalkyl, or R^2 is optionally substituted aryl, or R^2 is an optionally substituted heterocyclic radical;

R^3 is H, or is alkyl optionally substituted by -O-alkyl, aryl or cycloalkyl;

10 and

A is a linear or branched alkylene chain containing from 1 to 6 carbon.

3. A compound according to claim 1, or an optical isomer thereof, an oxide
form thereof, or a pharmaceutically acceptable addition salt thereof, wherein:

R^1 is -O- R'^1 , and R'^1 is H or alkyl containing from 1 to 6 carbon atoms;

15 R^2 is alkyl containing from 1 to 6 carbon atoms which is optionally substituted by -O-alkyl containing from 1 to 6 carbon atoms, phenyl, or 5- or 6-membered cycloalkyl, or R^2 is optionally substituted phenyl, or R^2 is an optionally substituted heterocyclic radical;

20 R^3 is H or alkyl containing from 1 to 6 carbon atoms which is optionally substituted by -O-alkyl containing from 1 to 6 carbon atoms, phenyl, or 5- or 6-membered cycloalkyl; and

A is a linear or branched alkylene chain containing from 1 to 6 carbon atoms.

26474-1095

42

4. A compound according to claim 1, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof, wherein:

R^1 is $-O-R'^1$, and R'^1 is H, methyl, or ethyl;

5 R^2 is alkyl containing from 1 to 6 carbon atoms optionally substituted by methoxy, ethoxy, substituted phenyl, cyclopentyl, or cyclohexyl, or R^2 is optionally substituted phenyl, or R^2 is an optionally substituted aromatic heterocyclic radical containing at least one nitrogen atom;

10 R^3 is H, or is alkyl containing from 1 to 6 carbon atoms which is optionally substituted by methoxy, ethoxy, substituted phenyl, cyclopentyl, or cyclohexyl; and

A is an alkylene chain of the formula $-(CH_2)_k-$, in which k is 1, 2, 3, 4, 5, or 6, or A is $-C(CH_3)_2-$.

5. A compound according to claim 1, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof, wherein:

15 R^1 is OH;

R^2 is methyl, ethyl, propyl, or n-hexyl, in each case optionally substituted by a methoxy, ethoxy, substituted phenyl, or cyclopentyl, or R^2 is optionally substituted phenyl, or R^2 is optionally substituted pyridyl;

20 R^3 is H, or is methyl, ethyl, propyl, isopropyl, isobutyl, or isopentyl, which in each is optionally substituted by methoxy, ethoxy, or substituted phenyl; and

A is an alkylene chain of the formula $-(CH_2)_k-$, in which k is 1, 2, or 3, or A is $-C(CH_3)_2-$.

6. A compound according to claim 1, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof, wherein

26474-1095

43

substituents on the aryl and heterocyclic radicals are each halogens, methyl, ethyl, methoxy, phenyl, trifluoromethyl, or trifluoromethoxy.

7. A compound according to claim 1, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof, wherein said
5 heterocyclic radicals are each thienyl, benzothiophenyl, pyridyl, or oxazolyl.
8. A compound according to claim 1, which is ethyl 1-benzyl-5-(3-carboxypropoxy)-2-methyl-6-pyrid-3-yl-1H-indole-3-carboxylate, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof.
9. A compound according to claim 1, which is ethyl 5-(3-carboxypropoxy)-
10 1-(4-chlorobenzyl)-2-methyl-6-pyrid-3-yl-1H-indole-3-carboxylate, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof.
10. A compound according to claim 1, which is ethyl 5-(1-carboxy-1-methylethoxy)-2-methyl-1-(3-methylbutyl)-6-pyrid-4-yl-1H-indole-3-carboxylate, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable
15 addition salt thereof.
11. A compound according to claim 1, which is ethyl 5-carboxymethoxy-6-hexyl-1-isobutyl-2-methyl-1H-indole-3-carboxylate, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof.
12. A compound according to claim 1, which is ethyl 5-carboxymethoxy-6-
20 (3-ethoxypropyl)-1-isobutyl-2-methyl-1H-indole-3-carboxylate, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof.
13. A compound according to claim 1, which is ethyl 5-carboxymethoxy-6-(3-cyclopentylpropyl)-1-(2-methoxyethyl)-2-methyl-1H-indole-3-carboxylate, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable
25 addition salt thereof.

26474-1095

44

14. A compound according to claim 1, which is ethyl 5-carboxymethoxy-6-[2-(4-fluorophenyl)ethyl]-1-(2-methoxyethyl)-2-methyl-1H-indole-3-carboxylate, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof.

5 15. A compound according to claim 1, which is ethyl 5-(3-carboxypropoxy)-6-hexyl-1-(2-methoxyethyl)-2-methyl-1H-indole-3-carboxylate, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof.

16. A compound according to claim 1, which is ethyl 5-(3-carboxypropoxy)-6-(3-ethoxypropyl)-1-(2-methoxyethyl)-2-methyl-1H-indole-3-carboxylate, or an
10 optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof.

17. A compound according to claim 1, which is ethyl 5-(3-carboxypropoxy)-6-(3-cyclopentylpropyl)-1-(2-methoxyethyl)-2-methyl-1H-indole-3-carboxylate, or an
15 optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof.

18. Pharmaceutical composition comprising at least one compound as defined in any one of claims 1 to 17, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof, in combination with one or more pharmaceutically acceptable vehicles.

20 19. Pharmaceutical composition according to claim 18 for use in the prevention or treatment of dyslipidaemia, atherosclerosis or diabetes.

20. Use of a compound as defined in any one of claims 1 to 17, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof, for the preparation of a medicament for the prevention or
25 treatment of dyslipidaemia, atherosclerosis or diabetes.

26474-1095

45

21. Use of a compound as defined in any one of claims 1 to 17, or an optical isomer thereof, an oxide form thereof, or a pharmaceutically acceptable addition salt thereof, for the prevention or treatment of dyslipidaemia, atherosclerosis or diabetes.

