



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 38/49, A61P 7/02, 7/10 // (A61K 38/49, 38:17)	A2	(11) International Publication Number: WO 00/59535 (43) International Publication Date: 12 October 2000 (12.10.00)
(21) International Application Number: PCT/IL00/00208 (22) International Filing Date: 3 April 2000 (03.04.00) (30) Priority Data: 129332 5 April 1999 (05.04.99) IL (71) Applicant (for all designated States except US): HADASIT MEDICAL RESEARCH SERVICES & DEVELOPMENT LTD. [IL/IL]; P.O. Box 12000, 91120 Jerusalem (IL). (72) Inventor; and (75) Inventor/Applicant (for US only): HIGAZI, Abd Al-Roof [IL/IL]; Neve Shalom, 99761 D.N. Shimshon (IL). (74) Agents: LUZZATTO, Kfir et al.; Luzzatto & Luzzatto, P.O. Box 5352, 84152 Beer-Sheva (IL).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: suPAR STIMULATING ACTIVITY OF tcuPA-MEDIATED FIBRINOLYSIS AND DIFFERENT USES THEREOF (57) Abstract The present invention relates to a complex (tcuPA/suPAR) of the two chain urokinase plasminogen activator (tcuPA) with the soluble urokinase plasminogen activator (suPAR), to pharmaceutical compositions comprising this complex and to different uses of the compositions in the treatment and/or prevention of thrombotic events, particularly those associated with the formation of fibrin clots. In the complex (tcuPA/suPAR), the suPAR stimulates fibrinolytic activity mediated by tcuPA under physiological conditions. The complex acts preferably on freshly-formed clots and is specific to such clots.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

suPAR STIMULATING ACTIVITY OF tcuPA-MEDIATED FIBRINOLYSIS AND DIFFERENT USES THEREOF

FIELD OF THE INVENTION

The present invention relates a complex of the two chain urokinase plasminogen activator (tcuPA) with the soluble urokinase plasminogen activator (suPAR) or with fragments thereof, pharmaceutical compositions comprising said complex as active ingredient and to the different uses of said compositions in the treatment and/or prevention of thrombotic events.

BACKGROUND OF THE INVENTION

Acute myocardial infraction, stroke, pulmonary emboli and other thrombotic events are the major causes of death in developed countries. Tremendous efforts have been made to develop drugs that can prevent or improve acute thromboembolic events. Amongst these drugs are thrombolytic agents that are used widely in the acute setting. However, the utility of currently available thrombolytics is limited by the high-risk of bleeding which accompanies their use. This danger results from the high doses that are required to lyse clots, which cause plasmin to accumulate in the circulation, resulting in systemic fibrinolysis and fibrinogenolysis. Intracarnial hemorrhage occurs in approximately 1% of patients receiving thrombolytic agents and other major hemorrhagic complications occur with a comparable incidence. These side effects limit the use of thrombolytic agents to the relief of ischemia only in the most severe cases.

Other limitations of available thrombolytic treatment include the length of time required for clot lysis to be achieved and blood vessel potency to be reestablished. In addition, available thrombolytic agents do not distinguish between newly formed clots, responsible for the acute event, and "aged" clots which may serve a physiological purpose.

Based upon these considerations, it would be advantageous to develop a thrombolytic agent that would act only in the presence of fibrin, would have a rapid onset of action and be specific for recently formed clots. These properties would make it possible to lyse clots more rapidly, with lower concentrations of the agent, thereby reducing the prevalence of systemic bleeding. The ability of such an agent to distinguish between recently formed and older clots would reduce the probability of further bleeding.

Plasminogen activators are widely used in the treatment of thromboembolic diseases. One of these activators is urokinase plasminogen activator (uPA), known to be synthesized as a proenzyme consisting of a single-chain protein (scuPA) [Pannell, R. & Gurewich V. *Blood* **69**:22-28 (1987)] and to be involved in several important biological processes including angiogenesis, wound healing, inflammation, ovulation and placental development, atherosclerosis, aneurysm and neointima formation and the formation of tumor metastases.

Limited proteolysis of scuPA results in the formation of two chains (tcuPA), considered to be the active form of the enzyme [Kasai, S. *et al.* *J. Biol. Chem.* **260**:12382-12389 (1985)]. One of the most important regulators of uPA is the plasminogen activator inhibitor-1 (PAI-1). This regulator interacts with tcuPA in a very rapid two-step reaction, leading to the formation of an inactive, SDS-stable 1/1 complex [Lindahl, T. L. *et al.* *Biochem. J.* **265**:109-113 (1990)]. The binding of tcuPA to uPAR only slightly reduces the susceptibility of tcuPA to the inhibitory effect of PAI-1 [Ellis, V. *et al.* *J. Biol. Chem.* **265**:9904-9908 (1990)].

Over the last years, the inventors have been engaged in developing and characterizing novel plasminogen activators that fulfill the above specified criteria.

It has now been surprisingly found that soluble human urokinase receptor (suPAR) stimulates the activity of the commonly used thrombolytic agent tcuPA, under specific conditions *in vitro*. As will be shown in the following Examples, the activity of the tcuPA/suPAR complex which is formed between the said two components, is fibrin- dependent and requires the presence of specific plasma or serum components(s).

Further, as will also be shown in the following Examples, under *in vivo* conditions, suPAR, by complexing with tcuPA, significantly enhanced lysis of pulmonary emboli, compared to lysis by tcuPA in the absence of the receptor.

SUMMARY OF THE INVENTION

The present invention relates to a complex of tcuPA with suPAR, or fragment/s of said suPAR (said complex also referred to herein as tcuPA/suPAR), in which the suPAR, or said fragment/s thereof, stimulates fibrinolytic activity mediated by tcuPA under physiological conditions.

The complex according to the invention acts preferably on freshly-formed clots and therefore is specific thereto.

In a second aspect, the invention relates to a pharmaceutical composition comprising as active ingredient a therapeutically effective amount of the suPAR/tcuPA complex of the invention. The compositions according to the invention may optionally further comprise pharmaceutically acceptable carriers, diluents, adjuvants and preserving agents.

In a preferred embodiment, the pharmaceutical compositions of the invention are for the treatment and/or prevention of thrombotic events associated with the formation of fibrin clots.

The invention is also concerned with the use of suPAR as a stimulator of the fibrinolytic activity of tCuPA, particularly for the treatment of thrombotic events.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 *Effect of suPAR on tCuPA-mediated clot lysis*

Lysis of clots prepared from human plasma. Clots were prepared by adding thrombin (0.4 NIH U/ml final concentration). TcuPA or equimolar concentrations of suPAR and tCuPA (tcuPA/suPAR) were added for 2-3 hrs, at 37°C. Lanes 1 and 4 show the fibrinolytic activity mediated by tCuPA alone. Lane 2 shows the fibrinolytic activity mediated by tCuPA/suPAR at a concentration ratio of 1:1. Lane 3 shows the activity mediated by tCuPA/suPAR at a concentration ratio of 1:10. The sizes of the lytic areas generated by tCuPA/suPAR are observed.

Figure 2 *suPAR stimulates tCuPA-mediated fibrinolysis*

Plasma clots were supplemented with ^{125}I -fibrin and incubated with plasminogen and tCuPA in the presence of the indicated concentrations of suPAR. The released radioactivity (RelRad) was determined after 1 hr., at 37°C and the radioactivity released into the supernatant was measured. The mean $\pm$ SEM of three repeated experiments is shown. The concentration of HMW- or LMW-tCuPA was 25nM. The values indicated are means of three experiments.

Figure 3 *The effect of suPAR on HMW-tCuPA mediated fibrinolysis*

HMW-tCuPA was incubated with freshly prepared clots, in the presence and absence of suPAR. The values indicated are means $\pm$ SEM of three repeated experiments. T designates time (in minutes).

Figure 4 *suPAR stimulates tcuPA-mediated fibrinolysis in vivo*

uPA^{-/-} mice (uPA-depleted mice) and their syngeneic wild type littermates received tcuPA, tcuPA/suPAR, suPAR or PBS, via continuous IV infusion. One hour after injection of ¹²⁵I-microemboli, the radioactivity in the lungs was measured. The data is expressed as the difference between spontaneous clot lysis in wild type mice and the obtained in uPA^{-/-} mice infused with PBS, tcuPA, tcuPA/suPAR or suPAR. The mean $\pm$ SEM of 2 experiments is presented. ID designates injected dose. For cases designated by ***, $p = <0.001$ compared to uPA^{-/-} mice infused with PBS.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a complex of tcuPA with suPAR or with fragments of suPAR, which complex is believed to directly or indirectly induce fibrinolysis of plasma-derived clots, by suPAR stimulation of the fibrinolytic activity mediated by tcuPA. In particular, the fibrinolytic activity ascribed to suPAR/tcuPA is essentially specific to fresh fibrin clots.

It should be clear that the term suPAR refers to the receptor *per se*, or to functional fragments or derivatives thereof. By the term functional fragment or derivative is meant any fragment or derivative of suPAR which exhibits the biological activity of suPAR, particularly the stimulation of fibrinolytic activity mediated by tcuPA.

The terms 'fibrin clots' or 'plasma-derived clots' mean clots formed by the conversion of fibrinogen to fibrin, whereby red blood cells and other formed elements are entrapped within the coagulated plasma. The enzyme catalyzing this conversion is known to be thrombin, which is formed in shed

blood and converts fibrinogen into fibrin, thereby producing fibrin clots by a hydrolyzing action.

The fibrinolytic activity of the scuPA/suPAR complex, *in vivo*, under physiological conditions, as compared to the activity of tcuPA alone or in combination with suPAR, was described in WO98/25641. The results presented therein clearly show that suPAR has only marginal effect on tcuPA-mediated fibrinolysis, when applied to a fibrin clot at a 1:1 stoichiometry.

However, the inventors have now found that despite the consensus that pro-uPA's, e.g. scuPA or tcuPA, bind to their receptor, uPAR, in a 1:1 ratio [A. Nykjaer *et al* J. Biol. Chem. **269**(41):25668-25676 (1994)], when employing an excess amount of suPAR (for example, suPAR/tcuPA 10:1), the receptor or at least a functional fragment thereof, has an unexpected stimulatory effect on tcuPA-mediated fibrinolysis. This novel finding may indicate a different and novel mechanism of action of tcuPA, being activated only in the presence of an excess amount of suPAR.

In a second aspect, the invention concerns a pharmaceutical composition comprising as active ingredient a therapeutically effective amount of the tcuPA/suPAR complex of the invention.

The 'therapeutically effective amount', for purposes herein, is determined by such considerations as are known in the art. The amount must be effective to achieve improvement including but not limited to improved survival rate or more rapid recovery, or improvement or elimination of symptoms and other indicators as are selected as appropriate measures by those skilled in the art. The doses may be single doses or multiple doses over a period of several days, but single doses are preferred.

The tcuPA/suPAR complex or the pharmaceutical composition of the invention can be administered in various ways and may comprise, in addition to the active ingredient, pharmaceutically acceptable carriers, diluents, adjuvants, preserving agents and vehicles. According to one embodiment, the pharmaceutical composition of the invention is in a dosage unit form.

The pharmaceutical compositions can be administered subcutaneously or parenterally including intravenous, intraarterial, intramuscular, and intraperitoneal administration, as well as intrathecal techniques. Implants of the pharmaceutical preparations may also be useful. The pharmaceutically acceptable carriers, diluents, adjuvants and vehicles as well as implant carriers generally refer to inert, non-toxic solid or liquid fillers, diluents, or encapsulating material not reacting with the active ingredients of the invention.

When administering the complex or the pharmaceutical composition of the invention parenterally, it will generally be formulated in a unit dosage injectable form (solution, suspension, emulsion). The pharmaceutical formulations suitable for injection include sterile aqueous solutions and sterile powders for reconstitution into sterile injectable solutions. The carrier can be any physiologically acceptable suitable carrier, for example, water, or aqueous buffer solutions.

Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Non-aqueous vehicles such as cottonseed oil, sesame oil, olive oil, soybean oil, corn oil, sunflower oil or peanut oil and esters, such as isopropyl myristate, may also be used as solvent systems for compound compositions.

In addition, various additives which enhance the stability, sterility and isotonicity of the compositions, including antimicrobial preservatives, antioxidants, chelating agents and buffers can be added. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid and the like. In many cases it will be desirable to include isotonic agents, for example, sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin. According to the present invention, any vehicle, diluent, or additive used would have to be compatible with the compositions.

Sterile injectable solutions can be prepared by incorporating the compositions utilized in practicing the present invention in the required amount of the appropriate solvent with various of the other ingredients, as desired.

Nonetheless, the composition disclosed herein in detail can be administered orally to the patient. Conventional forms such as administering the composition as tablets, suspensions, solutions, emulsions, capsules, powders, syrups and the like are usable. Known techniques which deliver it orally or intravenously and retain the biological activity are preferred.

For delivery within the CNS intrathecal delivery can be used with for example an Ommaya reservoir. United States Patent 5,455,044 provides for use of a dispersion system for CNS delivery or see United States Patent 5,558,852 for a discussion of CNS delivery. In addition, pharmacological formulations that cross the blood brain barrier can be administered. Such formulations can take advantage of methods now available to produce chimeric structures in which the complex of the invention, alone or in combination with human IgG or said IgG-derived peptide/s are coupled to a

brain transport vector thus allowing transportation across the barrier. Further, in appropriate cases blood brain barrier disruption can be utilized.

The pharmaceutical composition according to the invention is intended for the treatment of thrombotic events associated with formation of fibrin clots. The term thrombotic event is well known to the man of the art, and may include *inter alia*, acute myocardial infraction, stroke, pulmonary emboli, cerebro-vascular events, disseminated intravascular coagulation (DIC), or deep vein thrombosis, but by no means should be limited thereto.

Evidently, any other disorder associated with the damaging formation of fibrin clots may also be treated or prevented by use of suPAR or the complex of the present invention.

In a further aspect, the invention relates to the use of suPAR as a stimulating agent for the fibrinolytic activity of tcuPA.

Further, the invention relates to the use of a complex of tcuPA/suPAR, in the preparation of a pharmaceutical composition, the composition being particularly for the treatment of thrombotic events associated with the formation of fibrin clots. As indicated above, the thrombotic events against which suPAR or suPAR/tcuPA may be used may include acute myocardial infraction, stroke, pulmonary emboli, cerebro-vascular events or deep vein thrombosis.

Yet further, the invention relates to a method for treating and/or preventing a thromboembolic disorder associated with the formation of fibrin clots in a patient in need of such treatment by administering to said patient a therapeutically effective amount of scPAR or of a complex comprising suPAR/tcuPA. The method according to the invention may be utilized for the treatment of acute myocardial infraction, stroke, pulmonary emboli,

cerebro-vascular events or deep vein thrombosis, however, should not be limited to these disorders.

The invention will now be described in more detail on hand of the following Examples, which are illustrative only and do not limit the scope of the invention, which is defined by the appended claims.

EXAMPLES

Materials and methods

SuPAR was a gift of Dr. J. Henkin and Dr. A. Mazar, Abbott Laboratories, Abbott Park, IL. Human fibrinogen and human thrombin were purchased from Sigma, St. Louis, MO. TcuPA was purchased from American Diagnostics, Greenwich, CT. Plasminogen was prepared as described by Deutsch and Mertz [Deutsch D. & Mertz E.T., Science **170**:1095-1096 (1970)]. Plasma was obtained from the Hadassah Hospital blood bank. Blood used to obtain plasma was drawn from healthy volunteers. Blood (450 ml) was collected in bags produced by Travenol Laboratories, Ashdod, Israel, containing 63 ml of CPD solution (containing 1.66 g sodium citrate (hydrous), 61 g dextrose, 206 mg citric acid and 140 mg moon basic sodium phosphate). Plasma was separated by centrifugation.

Radiolabeled fibrinogen

Plasminogen-depleted human fibrinogen (Fib 1-1340, American Diagnostics, Greenwich, CT) was trace radiolabeled with ^{125}I (NEN Life Sciences). ^{125}I -fibrinogen ($\sim 40 \times 10^6$ cpm) was added to 1 ml unlabeled fibrinogen (35 mg/ml) before the micro-emboli were prepared.

Preparation of micro-emboli

Human blood from healthy volunteers was collected in citrate (final concentration 0.32%). Plasma was isolated from the human blood by centrifugation at 1200xg.

Clots were formed by first adding 1 ml trace-labeled human ^{125}I -fibrinogen to 2.5 ml plasma. CaCl_2 and human thrombin (Sigma) were then added at a final concentration of 20 mM and 0.2 U/ml, respectively, followed by incubation for 1 hr at room temperature and further incubated overnight at 4°C. From this point all steps were performed at 4°C. The fibrin clots were decanted onto the lids of culture dishes, and then cut into small pieces and re-suspended in 2 ml of PBS buffer. The micro emboli were then suspended in 13 ml PBS buffer containing 3 mg/ml BSA. Immediately before injection, the preparations were sedimented for 5 min. in order to remove any larger aggregates which may have been formed. The supernatant was aliquoted into 0.2 ml doses for injection. Random aliquots of micro-emboli were selected to characterize the size distribution (characterized previously as 10-100 μm) using a Coulter Counter.

Mice

uPA depleted mice (uPA-/-), on a 25% Swiss/75% C57 blank background, and littermate controls were kindly provided by Dr. P. Carmeliet (Leuven, Belgium). All mice weighed 20-30 g at the time of the study.

In Vitro Experiments

Assessment of fibrinolysis by reduction in clot size

Human fibrinogen [Sigma, St. Louis MO] was reconstituted in phosphate buffered saline (PBS, pH 7.4) or in plasma to a concentration of 9 mg/ml, after which human thrombin was added (0.4NIH U/ml) to form clots. The mixture was decanted onto culture dish lids and incubated for 60 min, at room temperature. After clot formation, aliquots of phosphate-buffered saline (PBS), pH 7.4, containing tcuPA (10pmole in 10 μl PBS) was added to the surface of the clot in the presence or absence of 0-100 pmol suPAR. As a control, no tcuPA was added to several lids containing formed clots. The clots, in the presence or absence of suPAR, were then incubated for a further

period of 2-3 hr, at 37°C, during which the appearance of digestion areas was observed. At that stage, the clots were washed several times with PBS and incubated overnight with 0.2% trypan blue. On the following day the clots were rinsed four times with PBS and photographed. The size of the lytic zones was calculated by the NIH image program (Fig. 1).

Assessment of fibrinolysis by measuring the release of radioactivity

Human fibrinogen (American Diagnostics USA) was radiolabeled with ^{125}I [Higazi, AA-R *et al.* J. Biol. Chem. **270**:9472-9477 (1995)] and re-suspended in either PBS solution (pH 7.4) or in plasma at a fibrinogen concentration of 3 mg/ml. The final radioactivity of these preparations was ~30,000 cpm/ml. Clots were formed in 16 mm tissue culture wells (Costar, Cambridge, MA) by the addition of thrombin (0.4 NIH U/ml) to each well. The plasminogen activator (400 μl of HMW-tcuPA, 25 nM), was added directly to the center of each formed clot and after incubation for specific periods the wells were washed with PBS and the level of radioactivity released into the lavage solution was determined by a gamma counter. Figure 2 illustrates the plasminogen activation by the complex suPAR/tcuPA after 1 hr of incubation at 37°C.

Fibrinolysis was also measured as previously described [Higazi, AA-R., *et al.* Biochem. J. **300**:251-255 (1994)], by addition of 0.4 ml serum or PBS containing plasminogen activator, tcuPA (25nM) in the presence or absence of suPAR (50nM). The plates were incubated at 37°C for various time periods in which aliquots of 25 μl were removed for counting the solubilized label in a gamma counter. Fibrinolytic activity was determined by following the release from the ^{125}I -labeled fibrin clot of labeled soluble degradation products of fibrin and is illustrated in Figure 3.

Results

Plasma derived clots were used to examine the effect of suPAR on the activity of tcuPA. The activity of tcuPA was substantially stimulated at

molar concentrations of suPAR greater than 1:1 (Figs. 1 and 2). This stimulation could be suppressed by amino terminal fragment of urokinase (ATF) (which competes with tcuPA for receptor binding). In the presence of LMW-tcuPA, lacking the receptor-binding determinant, no stimulation was observed (data not shown). Blocking of the interaction of tcuPA with suPAR by ATF, caused almost complete inhibition of the stimulatory effect of suPAR on tcuPA-mediated fibrinolysis (Figs. 1 and 2).

Figure 3 show that in the presence of tcuPA, suPAR-mediated cleavage rate of the plasma clots was faster and to a greater extent than do equimolar concentrations of tcuPA.

To exclude the possibility that the stimulatory effect of tcuPA was due to contaminating scuPA, SDS-PAGE was performed, which confirmed the absence of scuPA in the reaction mixture (data not shown).

When clots formed from purified fibrinogen were employed, no stimulation by suPAR of tcuPA-mediated fibrinolysis was observed.

The data presented herein indicate that all of the components necessary for suPAR to stimulate the activity of tcuPA are present in freshly isolated serum. Nonetheless, the effect of serum was lost when kept overnight at 30°C, however, no loss of stimulatory activity was observed with plasma maintained under the same conditions, even for several days. These results suggest that the stimulatory factor(s) in serum underwent activation during storage. The differences between plasma and serum may indicate that after coagulation is initiated, the half-life of the stimulatory factor is limited. Further, these results indicate that freshly-formed clots will be lysed more efficiently than aged clots.

***In vivo* Experiments**

The effect of suPAR on the ability of tcuPA to cleave plasma clots in vivo

Radiolabeled micro-emboli were injected into the tail vein of mice and the plasminogen activator was injected into the jugular vein. The fibrinolytic activity of the various agents, i.e. tcuPA, tcuPA/suPAR or suPAR alone, was compared to a spontaneous clot lysis which served as a control, by measuring the rate in which radioactivity was cleared from the lungs of the mice. The results presented in Figure 4 clearly indicate that while suPAR alone has no fibrinolytic activity, in the presence of tcuPA, the fibrinolytic action mediated by the latter is substantially enhanced. Therefore, it is evident that suPAR stimulates tcuPA mediated fibrinolysis.

Distribution of micro-emboli

Freshly prepared micro-emboli (15-30,000 cpm/0.2 ml) were re-suspended and injected into the tail vein of mice. At 10 min, 1, 3, and 5 hr post-injection, the mice were anesthetized with metophane and 0.1 ml of blood were withdrawn from them into a heparinized capillary tube by retro-orbital puncture. The mice were then sacrificed by cervical dislocation and their major organs were immediately harvested, rinsed in saline, dried on filter paper and weighed. The radioactivity level in each tissue was determined from which the exact dose (cpm) injected into each animal was calculated by subtracting the residual radioactivity in the tube and syringe after the injection, from the original amount.

The radioactivity associated with the tail of each mouse was determined in order to verify that injection into the vein was complete. Pilot studies indicated that >50% of the total injected dose of microemboli were distributed in a homologous pattern throughout the lung on autoradiography and light microscopy. In contrast, <5% of ¹²⁵I-fibrinogen were found in association with the lungs.

Lysis of pulmonary micro-emboli

uPA^{-/-} mice and their syngeneic wild type littermates were anesthetized by intraperitoneal injection on Nembutal. The jugular vein was then cannulated using a siliconized polyethylene tube and PBS or tcuPA (alone, with soluble suPAR or with a combination of tcuPA/suPAR) were infused (IV infusion) via a PHD 2000 multi-syringe pump at a rate of 0.5 mg/kg/hr or 0.25 mg/kg/h, for 60 min. Five minutes after the onset of the infusion, ¹²⁵I-micro-emboli were injected into the tail's vein, during which the mice remained anesthetized. At the end of the infusion, the mice were sacrificed, the tissues were collected and their radioactivity was determined as described above.

Results

Figure 3 shows that while uPA^{-/-} mice have impaired lysis of pulmonary micro-emboli compared to that with syngeneic wild type mice, in the presence of tcuPA lysis was enhanced and in the presence of tcuPA/suPAR the rate of lysis of such pulmonary micro-emboli was substantially elevated. This finding clearly teaches the stimulatory effect of suPAR on the thrombolytic activity of tcuPA.

CLAIMS

1. A complex comprising tcuPA and suPAR, or at least one functional fragment of suPAR, in which the suPAR, or said fragment/s thereof, stimulates fibrinolytic activity mediated by tcuPA under physiological conditions.
2. The complex as claimed in claim 1, wherein said fibrinolytic activity is specific to substantially fresh clots.
3. A pharmaceutical composition comprising as active ingredient a therapeutically effective amount of the complex tcuPA/suPAR, in which the said suPAR stimulates fibrinolytic activity mediated by tcuPA.
4. The pharmaceutical composition as claimed in claim 3, further comprising pharmaceutically acceptable carriers, diluents, adjuvants and preserving agents.
5. The pharmaceutical composition as claimed in claim 4, for the treatment of thrombotic events associated with the formation of fibrin clots.
6. The pharmaceutical composition as claimed in claim 5, wherein said thrombotic events include acute myocardial infraction, stroke, pulmonary emboli, cerebro-vascular events, disseminated intravascular coagulation (DIC) or deep vein thrombosis.
7. The pharmaceutical composition as claimed in any one of claims 3 to 6, in a dosage unit form.
8. Use of suPAR as a stimulator of the fibrinolytic activity of tcuPA.
9. Use of tcuPA/suPAR complex in the preparation of a pharmaceutical composition for the treatment of thrombotic events associated with the formation of fibrin clots.
10. Use as claimed in claim 9, wherein said thrombotic events include acute myocardial infraction, stroke, pulmonary emboli, cerebro-vascular events disseminated intravascular coagulation (DIC) or deep vein thrombosis.

11. A method for treating and/or preventing a thromboembolic disorder associated with the formation of fibrin clots in a patient in need of such treatment by administering to said patient a therapeutically effective amount of a complex comprising tcnPA/suPAR.
12. The method as claimed in claim 11 for the treatment of acute myocardial infraction, stroke, pulmonary emboli, cerebro-vascular events or deep vein thrombosis.

1/3




Fig. 1

2/3




Fig. 2




Fig. 3

3/3




Fig. 4