

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
9 December 2004 (09.12.2004)

PCT

(10) International Publication Number
WO 2004/106366 A1

(51) International Patent Classification⁷: **C07K 7/56**, 1/04,
16/20, A61K 38/12, 39/015

(74) Agent: **NOBBE, Matthias**; Viering, Jentschura & Partner,
Centroallee 263, 46047 Oberhausen (DE).

(21) International Application Number:
PCT/EP2004/005952

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(22) International Filing Date: 2 June 2004 (02.06.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03012520.7 2 June 2003 (02.06.2003) EP

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **PEVION
BIOTECH LTD.** [CH/CH]; Rehhagstrasse 79, CH-3018
Bern (CH).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **SWISS TROPICAL
INSTITUTE** [CH/CH]; Socinstrasse 57, CH-4002 Basel
(CH). **UNIVERSITÄT ZÜRICH** [CH/CH]; Winterthur-
erstrasse 190, CH-8092 Zürich (CH). **PLUSCHKE, Gerd**
[DE/DE]; Im Vogelsang 3, 79189 Bad Kronzing (DE).
KIENZL, Ursula [CH/CH]; Affolternstr. 146, CH-8050
Zürich (CH). **ROBINSON, John** [CH/CH]; Tobelstr. 24,
CH-8615 Wermatswil (CH). **ZURBRIGGEN, Rinaldo**
[CH/CH]; Rainstrasse 40, CH-3185 Schmitten (CH).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: METHODS FOR SYNTHESIZING CONFORMATIONALLY CONSTRAINED PEPTIDES, PEPTIDOMETICS AND
THE USE THEREOF AS SYNTHETIC VACCINES

(57) Abstract: The present invention relates to methods for synthesizing conformationally constrained peptides and cyclic pep-
tidomimetics obtainable by these methods which are conformationally constrained due an internal cross-link. This cross-link is
formed between the side chain of an amino acid residue or analog and a (2S, 4S)-4- functionalized proline residue. The invention
further relates to the use of (2S, 4S)- 4-functionalized proline residues as building units in the synthesis of such peptidomimetics
and to the use thereof as antigens, alone or in combination with suitable immunopotentiating delivery systems, for example im-
munopotentiating reconstituted influenza virosomes to elicit an immune response in a mammal. Moreover, the invention also relates
to pharmaceutical compositions containing the same.



WO 2004/106366 A1

Methods for synthesizing conformationally constrained peptides, peptidomimetics and the use thereof as synthetic vaccines

- 5 The present invention relates to methods for synthesizing conformationally constrained peptides and cyclic peptidomimetics obtainable by these methods which are conformationally constrained due an internal cross-link. This cross-link is formed between the side chain of an amino acid
10 residue or analog and a (2S,4S)-4-functionalized proline residue. The invention further relates to the use of (2S,4S)-4-functionalized proline residues as building units in the synthesis of such peptidomimetics and to the use thereof as antigens, alone or in combination with suitable
15 immunopotentiating delivery systems, for example immunopotentiating reconstituted influenza virosomes to elicit an immune response in a mammal. Moreover, the invention also relates to compositions containing the same.
- 20 New scientific knowledge and technologies allow to identify and localize essential neutralizing epitopes of microorganisms causing infections and other diseases. These antigen structures can often be mimicked using small linear peptides or polypeptides which are generally designated as
25 peptidomimetics. Such mimetics appear to be promising candidates for vaccination strategies and in the design of novel synthetic vaccines.

However, conformation plays a key role in the ability of
30 peptides to elicit antibody responses against folded proteins. Linear peptides often elicit antibodies that bind well to denatured proteins, but less frequently recognize conformational epitopes in native protein structures. In

general, these peptides are more conformationally mobile and therefore unlikely to adopt a stable secondary structure in aqueous solution which corresponds to the three-dimensional epitope on the surface of a protein.

5

To overcome these limitations, conformational constraints can be incorporated into the primary peptide chain which will reduce the degree of structural flexibility. If the "frozen" conformation of such a peptide imitates the corresponding secondary structure of the target epitope, the peptide antigen can be used to raise antibodies which potentially cross-react with protein structures bearing such epitopes.

There are several approaches known in the prior art to introduce conformational constraints into a linear peptide or polypeptide chain. For example, bridging between two neighbouring amino acids in a peptide leads to a local conformational modification, the flexibility of which is limited in comparison with that of regular peptides. Some possibilities for forming such bridges include incorporation of lactams and piperazinones (for review see Giannis & Kolter, *Angew. Chem. Int. Ed.*, 1993, 32: 1244).

Global restrictions in the conformation of a peptide are possible by limiting the flexibility of the peptide strand through cyclization (Hruby et al., *Biochem. J.*, 268:249, 1990). The common modes of cyclization are the same found in naturally occurring cyclic peptides. These include side chain to side chain cyclization or side chain to end-group cyclization. Another common cyclization is the end-to-end cyclization.

Another conceptual approach to the conformational constraint of peptides was introduced by Gilon, et al., (Biopolymers, 31:745, 1991) who proposed backbone to backbone cyclization of peptides. The theoretical advantages of this strategy
5 include the ability to effect cyclization via the carbons or nitrogens of the peptide backbone without interfering with side chains that may be crucial for interaction with the antibody.

10 Yet another approach in the design of conformationally constrained peptidomimetics, which is described in US10/114918, is to attach a short amino acid sequence of interest to a template, to generate a cyclic constrained peptidomimetic. Such cyclic peptidomimetics are not only
15 structurally stabilized by their templates, and thereby offer three-dimensional conformations that may imitate conformational epitopes on viruses and parasites, but they are also more resistant than linear peptides to proteolytic degradation in serum.

20

US10/114918 further discloses the synthesis of conformationally constrained cross-linked peptidomimetics by preparation of synthetic amino acids for backbone coupling to appropriately positioned amino acids in order to stabilize
25 the supersecondary structure of mimetics. Cross-linking can be achieved by amide coupling of the primary amino group of an orthogonally protected (2S,3R)-3-aminoproline residue to a suitably positioned side chain carboxyl group of glutamate. This approach has been followed in the preparation of
30 conformationally constraint tetrapeptide repeats of the CS protein wherein at least one proline has been replaced by (2S,3R)-3-aminoproline and, in order to introduce a side chain

carboxyl group, glutamate has been incorporated as a replacement for alanine.

There are several approaches in the prior art to use modified
5 proline analogs for incorporating conformational constraints
into peptides. For example, Zhang et al., J. Med Chem., 1996,
39: 2738-2744, describes synthesis and biological testing of
several cyclic analogs of angiotensin II being cyclized
between positions 5 and 7 to study the conformational
10 features of molecular recognition of angiotensin II.
Cyclization has been achieved by forming amide bonds between
4-amino-trans-proline and side chain carboxyl groups of
aspartate and glutamate on one hand and on the other hand, by
disulfide bridges between homocysteine residues and 4-
15 mercapto-trans/cis-prolines.

Furthermore, (2S,3R,4R)-diaminoproline as a diketopiperazine
was incorporated by solid-phase peptide synthesis into a
protein loop mimetic. This was shown by NMR analysis to adopt
20 a stable beta-hairpin conformation in DMSO (Pfeifer &
Robinson, Chem. Comm., 1998, 1977-1978).

Vaccination is arguably the most successful medical invention
and global vaccination programs have yielded impressive
25 results. However, the appearance of novel infectious diseases
of global threat, like AIDS and the comeback of infectious
diseases that seemed to be mastered by chemotherapy, like
tuberculosis and malaria causes a great need to develop new
sophisticated vaccines.

30

In particular, malaria is an increasing health problem
throughout the third world. Several hundred million people
suffer from the disease and the most acute form, caused by

the protozoan parasite *Plasmodium falciparum*, kills over a million children each year in Africa alone.

There is presently no effective vaccine against the parasite,
5 and older established drugs like chloroquine are rapidly loosing their effectiveness due to resistance. On the other hand, ongoing research has provided many new antigens as potential malaria vaccine candidates.

10 Several antibody targets on the malaria parasite have been identified, one of which is the circumsporozoite (CS) protein present on the surface of early sporozoites (Potocnjak et al., J. Exp. Med., 1980, 151: 1504-1513). The central portion of the CS protein ($M_r = 44$ kDa contains 41 tandem repeats of
15 a tetrapeptide, 37 of which are Asn-Ala-Asn-Pro and four of which are Asn-Val-Asp-Pro (Godson et al., Nature, 1983, 305: 29-33; Dame et al., Science, 1984, 225: 593-599).

It was shown that linear tandemly repeated NANP peptides can
20 elicit antibodies in mice and rabbits that recognize the native CS protein and block sporozoite invasion of hepatocytes (Dame et al., Science, 1984, 225: 593-599; Ballou et al., Science, 1985, 228: 996-999; Young et al., Science, 1985, 228: 958-962; and Zavala et al., Science, 1985, 228:
25 1436-1440). These results were a prelude to vaccination studies in humans with synthetic (Asn-Ala-Asn-Pro)₃ peptides conjugated to tetanus toxin. However, the immune response was not strong enough for these conjugates to be useful as malaria vaccine (Herrington et al., Nature, 1987, 328: 257).
30 Subsequently, a number of studies were initiated to optimise the immune response to (Asn-Ala-Asn-Pro)_n peptides (Etlinger et al., Eur. J. Immunol., 1991, 21: 1505-1511; Tam et al., J. Exp. Med., 1990, 171: 299-306; Pessi et al., Eur. J.

Immunol., 1991, 21: 2273-2276; and deOliviera et al., Vaccine, 1994, 12: 1012-1017).

All of these efforts should be seen in view of the fact that the conformations of the (Asn-Ala-Asn-Pro) repeats in the CS protein were not known and so could not be taken into account in the design process. It seemed most likely that short linear (Asn-Ala-Asn-Pro)₃ peptides would be largely unstructured in aqueous solution, and be susceptible to rapid proteolytic degradation in serum. A later study also suggested that a significant part of the immune response against a linear (Asn-Ala-Asn-Pro)₃ peptide is directed against the chain termini, which of course are not present in the native CS protein (Etlinger & Trzeczak, Phil. Trans. Roy. Soc. Lond. B, 1993, 340: 69-72). Although the structure of the (Asn-Ala-Asn-Pro)-repeat region in the CS protein is still unknown, modelling suggests that the (Asn-Pro Asn-Ala-Asn)-motif may adopt a helical beta-turn, which is tandemly repeated in the CS protein to generate a novel supersecondary structure. Cyclic peptidomimetics of this (Asn-Pro Asn-Ala-Asn)-motif were synthesized and shown by NMR to adopt helical turns in aqueous solution (Pfeifer et al., Chimia, 2001, 55(4): 334-339).

Using the approach to incorporate conformational constraints by means of internal cross-linking, US10/114918 describes the preparation of a conformationally constrained peptidomimetic corresponding to the afore-mentioned tetrapeptide repeat region of the CS protein. In a five-fold repeat, one proline residue has been replaced by (2S,3R)-3-aminoproline and further, one alanine residue has been replaced by glutamate. Internal cross-linking has then been achieved by amide

coupling between the primary amino group of the modified proline unit and the side chain carboxyl group of glutamate.

As outlined above, peptide and protein mimetics are potentially of great value in synthetic vaccine design. The mimetics should function by stimulating the immune system to produce antibodies that recognize the intact parasite. However, the difficulty of presenting in a mimetic the conformational epitopes found on the native antigenic that are required for protective antibody responses are not yet fully overcome.

There is still a need for new routes to incorporate conformational constraints into a linear peptide chain to freeze a specific conformation. Since it is impossible to exactly predict the final conformation of the resulting peptide, it is generally required to create a whole set of constrained peptides of the same primary sequence, which are then analysed for their ability to mimic an epitope of the antigenic protein. For example, such an analysis can be carried out by testing the individual constrained peptides for their cross-reactivity towards known antibodies of said antigenic protein. It is therefore necessary to have a variety of different modes for incorporating a conformational constraint into a precursor peptide at hand.

Surprisingly, it has now been found that conformationally constrained peptides which are cyclized through a 4-substituted proline residue, having a specific stereochemistry, provide for a distinct and unique three-dimensional structure of a given peptide. The conformational constraint is achieved by an internal cross-link between a modified proline residue carrying a functional group at the

4-position and a spatially adjacent side chain functional group of a second residue. The modified proline residue is characterised by a (2S,4S) stereochemistry.

5 Cyclic peptidomimetics cyclized via (2S,4S)-4-substituted proline residues provide for a new class of conformationally constrained peptides which appear to be more stable in aqueous solution and consequently less prone to degradation in serum when, for example, administered as vaccine.

10

Synthesis of peptidomimetics containing (2S,4S)-4-substituted proline residue(s) are much easier accessible as for example their 3-substituted counterparts due to a shortened synthesis route comprising less individual steps. Furthermore, starting materials to synthesize the (2S,4S)-4-substituted proline unit are available in large amounts as they are standard substances which can directly be purchased from several suppliers, for example from Neosystem, 7 rue de Boulogne, 67100 Strasbourg, France.

20

The incorporation of a (2S,4S)-4-substituted proline residue instead of, for example a 3-substituted proline unit achieves a conformationally constrained peptide which has a substantially altered three-dimensional peptide structure. It is surprising that this minor change of the chemical structure has such a strong impact on the conformation and immunogenicity of the resulting peptide of the peptide.

25 Amino acids and amino acid residues described herein may be referred to according to the accepted one or three letter code referenced in text books well known to those of skill in the art, such as Stryer, Biochemistry, 4th Ed., Freeman and

30

Co., New York, 1995 and Creighton, Proteins, 2nd Ed. Freeman and Co. New York, 1993.

As used herein, the terms "peptide" and "polypeptide" are
5 used synonymously and in their broadest sense to refer to a compound of two or more amino acid residues, or amino acid analogs. The amino acid residues may be linked by peptide bonds, or alternatively by other bonds, e.g. ester, ether etc. As used herein, the term "amino acid" or "amino acid
10 residue" refers to either natural and/or unnatural or synthetic amino acids, including both the D or L enantiomeric forms, and amino acid analogs.

The term "epitope" or "B cell epitope" as used herein,
15 designates the structural component of a molecule that is responsible for specific interactions with corresponding antibody (immunoglobulin) molecules elicited by the same or related antigen. More generally, the term refers to a peptide having the same or similar immunoreactive properties, such as
20 specific antibody binding affinity, as the antigenic protein or peptide used to generate the antibody. An epitope that is formed by a specific peptide sequence generally refers to any peptide which is reactive with antibodies against the specific sequence.

25

The term "antigen" as used herein, means a molecule which is used to induce production of antibodies. The term is alternatively used to denote a molecule which is reactive with a specific antibody.

30

The term "immunogen" as used herein, describes an entity that induces antibody production in a host animal. In some

instances the antigen and the immunogen are the same entity, while in other instances the two entities are different.

5 The term "immunopotentiating" is used herein to refer to an enhancing effect on immune functions which may occur through stimulation of immune effector cells and may lead to increased resistance to infectious or parasitic agents.

10 The term "synthetic" as used herein relates to peptides produced by a chemical method as described above, for example.

15 The term "peptidomimetic" is used herein to denote a peptide or peptide analog that biologically mimics active determinants on parasites, viruses, or other bio-molecules.

20 The term "conformation" as used herein denotes the various nonsuperimposable three-dimensional arrangements of atoms that are interconvertible without breaking covalent bonds.

In a first embodiment, the present invention relates to a method for synthesizing a conformationally constrained peptide which comprises one or more regions of general formula (I):

25 (N-terminus) ...-[X]_m-Y-[X]_n-Z-[X]_o-... (C-terminus) (I)

30 wherein X = an amino acid residue or an amino acid analog and can be the same or different if n > 1;
m, o ≥ 0 and n ≥ 1, preferably ≥ 2;
Y, Z = A or B and Y ≠ Z, wherein

A = a 4-FgA-proline residue, wherein FgA is a functional group at the 4-position of the proline residue;

B = an amino acid residue or an amino acid analog having a side chain functional group FgB,

wherein the functional groups FgA and FgB are capable of forming an internal link by coupling the functional group FgA at the 4-position of A and the side chain functional group FgB of B;

and comprises the following steps of (a) providing amino acid residues Y and Z having appropriate functional groups FgA and FgB, said functional groups being optionally protected, (b) synthesizing a linear peptide comprising the amino acid residues Y and Z, and (c) optionally deprotecting, if said functional groups are optionally protected, and reacting the functional groups FgA and FgB for converting the linear peptide into the cross-linked form by coupling the functional group FgA at the 4-position of the proline residue (A) and the side chain functional group FgB of B.

Preferably, A is a (2S,4S)-4-FgA-proline residue, more preferably a (2S,4S)-4-aminoproline residue. Preferably, B is an amino acid residue or an amino acid analog having a side chain carboxyl group, preferably, glutamate or aspartate, wherein an internal cross-link is formed between A and B by amide coupling the amino group at the 4-position of the (2S,4S)-4-aminoproline residue and the carboxyl group of B.

Preferably, said conformationally constrained peptide is capable of eliciting a pathogen-specific immune response in a mammal.


- 5 It is preferred that the linear peptide of step (b) comprises one or more portions of the malaria circumsporozoite (CS) protein of a Plasmodium species, preferably Plasmodium falciparum. More preferably, this sequence comprises one or more tetrapeptides, which are selected from the group
10 consisting of Asn-Pro-Asn-Ala, Asn-Pro-Asn-Val, Asp-Pro-Asn-Ala and Asp-Pro-Asn-Val.

Preferably, the method for synthesizing conformationally constrained peptides according to the present invention is
15 carried out using solid phase synthesis techniques in the assembling step (c). The linear peptide can be assembled using Fmoc-chemistry. Cleavage from the resin and removal of side-chain protecting groups can proceed in one step and the introduction of the internal cross-link between the (2S,4S)-
20 4-substituted proline unit, preferably (2S,4S)-4-aminoproline, and a spatially adjacent side chain functional group can be achieved by cyclization in dimethylformamide (DMF) with a coupling reagent such as O-(7-Azabenzotriazole-1-yl)-N,N,N',N-tetramethyluronium-hexafluorophosphate (HATU).

25 Optionally, this method further includes the step of attaching the cyclized peptide to a phospholipid moiety (e.g. PE). The phospholipid anchor is preferably attached via a linker, preferably a dicarboxylate linker, more preferably a
30 succinate linker.

In another embodiment, this invention relates to conformationally constrained peptides which are obtainable by

the method(s) according to the invention. These peptides comprise one or more regions of general formula (I):



5

wherein X = an amino acid residue or an amino acid analog and can be the same or different if $n > 1$;

$m, o \geq 0$ and $n \geq 1$, preferably ≥ 2 ;

10

$Y, Z = A$ or B and $Y \neq Z$, wherein

A = a 4-FgA-proline residue, wherein FgA is a functional group at the 4-position of the proline residue;

15

B = an amino acid residue or an amino acid analog having a side chain functional group FgB,

20

wherein the functional groups FgA and FgB are forming an internal link by coupling the functional group FgA at the 4-position of the proline residue and the side chain functional group FgB of B.

Preferably, A is a (2S,4S)-4-FgA-proline residue, more preferably a (2S,4S)-4-aminoproline residue.

25

Preferably, the functional groups FgA and FgB are chosen to form a cyclic ester or cyclic amide bond. Preferably, B is an amino acid residue or an amino acid analog having a side chain carboxyl group, preferably glutamate or aspartate, wherein an internal cross-link is formed between A and B by amide coupling the amino group at the 4-position of the (2S,4S)-4-aminoproline residue and the carboxyl group of B.

30

More preferably, the functional group at the 4-position of the modified proline unit is an amino group and cyclization is achieved through amide coupling of said 4-amino of (2S,4S)-4-aminoproline (Apro) to a side chain carboxyl group
5 FgB of an spatially adjacent residue.

More preferred are peptides wherein the internal cross-link forming residues A and B are separated by more than one amino acid residues or analogs, i.e. $n \geq 2$.

10

Another embodiment of the invention relates to peptides which are conformationally constrained due to an internal cross-link between a (2S,4S)-4-functionalized proline unit and a suitable side chain functional group of a second residue and
15 which are mimicking the three-dimensional structure of, for example, an epitope on the surface of an antigenic protein. These peptides, designated hereinafter as peptidomimetics, are characterized in that the above-described internal cross-link stabilizes a supersecondary 3D structure of the peptide.

20

To form this internal cross-link between the modified proline unit and the side chain of another spatially adjacent amino acid residue or analog, said side chain must provide a suitable functional group to allow the formation of a stable
25 chemical bond between both residues. Preferably, the amino acid residue or analog B is selected from the group consisting of glutamate and aspartate and the afore-mentioned internal cross-link is formed through amide coupling with (2S,4S)-4-aminoproline. More preferably, B is glutamate.

30

Therefore, the present invention also refers to the use of a 4-Fg-proline, preferably a (2S,4S)-4-Fg-proline for synthesizing conformationally constrained peptides, wherein

Fg is a functional group, selected from amino, hydroxy, sulfhydryl, halogen, sulfonyl, carboxy, thiocarboxy or substituted derivatives thereof, and being preferably selected from amino, hydroxy, halogen, carboxy, or substituted derivatives thereof. The invention furthermore refers to such conformationally constrained peptides in which at least one 4-Fg-proline, preferably a (2S,4S)-4-Fg-proline is incorporated.

10 The conformationally constrained peptides according to the present invention closely resemble the three-dimensional conformations found on an intact pathogenic protein, thus providing improved epitopes for the generation of pathogen-specific antibodies that efficiently cross-react with pathogens.

In yet another embodiment, the inventive peptide comprises one or more portions of the malaria circumsporozoite (CS) protein of a Plasmodium species, preferably of Plasmodium falciparum.

Protection of mammals, including man, against infection by the etiologic agent of malaria, Plasmodium can be achieved by eliciting an immune response directed against the circumsporozoite (CS) protein. Four species of Plasmodium are known to infect man. These are *P. falciparum*, *P. vivax*, *P. ovale* and *P. malariae*. The CS protein of *P. falciparum* comprises about 412 amino acids with an approximate molecular weight of 44,000. It comprises 41 tandem repeats of a tetrapeptide. Synthetic peptides of a length of 5-20 residues derived from the repeat region of the CS protein of *P. falciparum* are preferred.

Although, the three-dimensional structure of the tetrapeptide repeat region in the CS protein is still unknown, theoretical studies suggest that it is likely to adopt a stable and repetitious conformation, possibly based on beta-helical turns or similar structures. The present invention provides peptides for the molecular mimicry of the conformational epitopes of the native malaria CS protein which are structurally optimized in order to elicit cross-reactive antibodies with higher efficiency.

In a preferred embodiment of the invention, the peptide comprises one or more tetrapeptides selected from the group consisting of Asn-Pro-Asn-Ala, Asn-Pro-Asn-Val, Asp-Pro-Asn-Ala and Asp-Pro-Asn-Val. More preferably, these peptides comprise 3 to 10, and most preferably 4 to 6 of such tetrapeptide units.

The present invention relates to peptidomimetics for the molecular mimicry of the conformational epitopes of the CS protein of Plasmodium species, preferably *P. falciparum*. More preferably, the conformationally constrained peptide comprises (Asn-Pro-Asn-Ala)_n wherein n is 2, 3, 4 or 5.

Models of these conformationally constrained peptidomimetics may be assessed for the stability and adoption of supersecondary structure in molecular dynamics (MD) simulations in solvent. Adoption of a supersecondary structure by these model peptidomimetics may be evidence that their structures are close to the preferred conformation of the tetrapeptide-repeat region in the native CS protein. Furthermore, 2D NOESY (nuclear Overhauser enhancement spectroscopy) spectra of conformationally constrained peptidomimetics may be examined for connectivities between

the peptide NH groups in the tandemly repeated helical turns. Such connectivities provide evidence for the relatively stable helical turn formation in the context of a supersecondary structure conformation.

5

In a more preferred embodiment, the invention relates to peptides wherein at least one of said Asn-Pro-Asn-Ala tetrapeptides is replaced by Asn-Pro-Asn-Glu and wherein the glutamate residue of Asn-Pro-Asn-Glu forms an internal cross-
 10 link through amide coupling with (2S,4S)-4-aminoproline.

Preferably, the peptide comprises one or more units of formula (II):

15 Formula (II):

(N-terminus)...Asn-Pro-Asn-Ala-Asn-Apro-(Asn-Ala-Asn-Pro)₂-Asn-Glu-Asn-Pro-Asn-Ala...(C-terminus)




wherein Apro is (2S,4S)-4-aminoproline.

It is more preferred that the peptide essentially comprises a five-fold tandem repeat of (Asn-Pro-Asn-Ala) of the CS
 20 protein of *P. falciparum*, wherein Pro of the second repeat is replaced by a (2S,4S)-4-aminoproline unit and alanine of the fourth repeat is replaced by glutamate (formula III, hereinafter denoted as UK39).

25 Formula (III):

H₂N-Asn-Pro-Asn-Ala-Asn-Apro-Asn-Ala-Asn-Pro-Asn-Ala-Asn-Pro-Asn-Glu-Asn-Pro-Asn-Ala-COOH



wherein Apro is (2S,4S)-4-aminoproline.

Through amide coupling, an internal cross-link is formed
5 which stabilizes a structure mimicking the native
conformation of CS-protein tandem repeat epitopes.

As described in more detail in the following example,
modeling studies concerning the peptidomimetic UK39 (formula
10 V) have revealed a unique and distinct conformation compared
to the isomeric mimetic BP66 which merely differs to UK39 in
that the cyclization has been achieved through (2S,3R)-3-
aminoproline. Due to the easily accessible (2S,4S)-4-
aminoproline, UK39 (in contrast to BP66) can be obtained by a
15 short and standardized synthesis route. Since UK39 requires
less individual synthesis steps and the starting material are
cataloged standard substances and are therefore available in
large quantities, high amounts of UK39 can easily be obtained
in higher amounts if compared to the corresponding synthesis
20 of BP66.

Furthermore, UK39 is less prone to be degraded in aqueous
solution than its isomeric counterpart BP66. This is
apparently due to a gain of stability of the conformation.
25


Moreover, UK39 shows an excellent antigenicity and parasite
clearance in vivo. If compared with BP66, it is a better
mimetic. With regard to cross reactivity to antibodies which
were raised against *P. falciparum*, more antibodies bind to
30 UK39 than to BP66.

The skilled artisan will appreciate that the afore-mentioned
beneficial features are not limited to amide bond

cyclizations. That is to say, UK39 derivatives wherein the internal cross-link is formed by a disulfide bridge or by an ester bond, i.e. wherein the modified proline unit is (2S,4S)-4-mercaptoproline or (2S,4S)-4-hydroxyproline, respectively, and wherein in the first case alanine is replaced by cysteine instead of glutamate are also encompassed by the present invention. The specific conformation behind the afore-mentioned beneficial features is essentially determined by the stereochemistry and the position of the cross-linking functional group at the modified proline ring.

The skilled artisan will appreciate that the afore-mentioned antigenic polypeptide molecules may be administered with one or more adjuvants in order to enhance the immunological response. For example, depending on the host species, adjuvants which may be used include, but are not limited to: mineral salts or mineral gels such as aluminum hydroxide, aluminum phosphate, and calcium phosphate; surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, and dinitrophenol; immunostimulatory molecules, such as cytokines, saponins, muramyl dipeptides and tripeptide derivatives, CpG dinucleotides, CpG oligonucleotides, monophosphoryl Lipid A, and polyphosphazenes; particulate and microparticulate adjuvant, such as emulsions, liposomes, virosomes, cochleates; or an immune stimulating complex mucosal adjuvants, Freund's (complete and incomplete, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum.)

In another embodiment, the peptides according to the invention are coupled to a phospholipid. Preferably, the N-



terminus of these peptides is coupled via a linker, preferably a dicarboxylate linker, more preferably a succinate linker to a fatty acid derivative of phosphaditylethanolamine, preferably 1-palmitoyl-3-oleoyl-phosphatidylethanolamine (PE, formula IV). Said fatty acid derivative is preferably a mono- to di-ester of glycerol with one or two C₁₀ to C₃₀ fatty acids which may have one or more double bonds and may be same or different.

Formula (IV):

The peptides according to the invention can be used in synthetic vaccine design. Peptidomimetics function by stimulating the immune system to produce antibodies that recognize the intact parasite. Preferably, such mimetics are presented to the immune system in a way that leads to a more efficient antibody production. For example, cyclic peptidomimetics can be presented on immunopotentiating reconstituted influenza virosomes (IRIVs) or liposomes, a form of antigen delivery that is practised already in human clinical use.

The attachment of a phospholipid anchor to the N-terminus of peptidomimetics functioning as antigen allows to combine the peptide with immunopotentiating delivery systems. According to a further aspect of the invention, peptides are thus

preferred which are combined with an immunopotentiating delivery system. Preferred immunopotentiating delivery systems are selected from the group consisting of liposomes, multiple-antigen peptides and immunopotentiating
5 reconstituted virosomes.


For example, immunopotentiating reconstituted influenza virosomes (IRIV) as human compatible immunopotentiating delivery agents are capable of presenting the
10 conformationally constrained peptidomimetics in multiple copies to the immune system and therefore, further improves the generation of efficient pathogen cross-reactive antibody responses. IRIVs are spherical, unilamellar vesicles, prepared from a mixture of phospholipids and influenza virus
15 surface glycoproteins. The hemagglutinin membrane glycoprotein of influenza virus plays a key role in the mode of action of IRIVs. This major antigen of influenza is a fusion-inducing component, which facilitates antigen delivery to immunocompetent cells.

20

In addition, peptides according to the invention (T cell epitopes) can be encapsulated into virosomes in order to be protected from enzymatic degradation by the body fluids and will be presented to the immune system via the MHC I class
25 pathway.

Particularly preferred is a peptide of five (Asn-Pro-Asn-Ala) tandem repeats that is internally cross-linked as described above and which further comprises a PE moiety at the N-
30 terminus which provides for the attachment to an immunopotentiating reconstituted influenza virosome (formula V).

Formula (V):



For in vivo experiments peptides according to the invention
 5 can be combined with adjuvants in order to enhance the immunological response. For example, depending on the host species, adjuvants which may be used include, but are not limited to: mineral salts or mineral gels such as aluminum hydroxide, aluminum phosphate, and calcium phosphate; surface
 10 active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, and dinitrophenol; immunostimulatory molecules, such as cytokines, saponins, muramyl dipeptides and tripeptide derivatives, CpG dinucleotides, CpG
 15 oligonucleotides, monophosphoryl Lipid A, and polyphosphazenes; particulate and microparticulate adjuvant, such as emulsions, liposomes, virosomes, cochleates; or an immune stimulating complex mucosal adjuvants, Freund's (complete and incomplete, and potentially useful human
 20 adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum.) However, it is noted that strong adjuvants, e.g. Freund's adjuvants can cause severe undesirable side effects that they are not accepted by regulatory authorities for human use.

Yet another embodiment concerns the use of (2S,4S)-4-substituted proline for synthesizing conformationally constrained peptides. Preferably, this covalently modified proline is used in one of the above described method for
5 synthesizing conformationally constrained peptides which have been cyclized through coupling of said modified proline unit, preferably (2S,4S)-4-aminoproline, and a second residue of the peptide chain. In this method, (2S,4S)-4-aminoproline is incorporated during the assembling step (b).

10

The cross-linked peptidomimetic can be prepared by solid phase synthesis methods well-known in the art. The linear peptide can be assembled using Fmoc-chemistry. Cleavage from the resin and removal of side-chain protecting groups can
15 proceed in one step and the key backbone coupling of the modified proline residue and glutamate can be achieved by cyclization in DMF with a coupling reagent such as HATU.

In a further embodiment, the invention relates to the use of
20 the inventive peptides for the manufacture of a vaccine for the treatment of malaria. Preferably, conformationally constrained peptidomimetics of the central (Asn-Pro-Asn-Ala) repeat region of the CS protein of the malaria parasite *P. falciparum* can be used to mimick the surface structure of CS
25 protein and thereby elicit a humoral immune response. Antibodies raised against such a mimetic are capable to cross-react with the native CS protein on *P. falciparum* sporozoites. Therefore, peptidomimetics according to the invention can be widely used in the design of molecularly
30 defined combined synthetic vaccines, including those targeted against multiple antigens and development stages of *P. falciparum*, and against other infectious agents.

In a further embodiment, the invention concerns a method for producing antibodies in a host against Plasmodium species, preferably against Plasmodium falciparum comprising the step of administering an above-described peptide to said host.

5

Another embodiment of the present invention relates to an in vitro method for detecting Plasmodium species in a sample comprising the steps (a) contacting said sample with an antibody according to the invention under conditions such
10 that binding to CS protein epitopes occurs if CS protein is present; and (b) detecting the presence of said antibody bound to an CS protein epitope.

In detail, the method comprises incubating a test sample with
15 one or more antibodies of the present invention and assaying whether the antibody binds to the test sample. The presence of CS protein may indicate malaria disease.

Conditions for incubating an antibody with a test sample
20 vary. Incubation conditions depend on the format employed in the assay, the detection methods employed and the type and nature of the antibody used in the art. Examples of such assays can be found in Tijssen, "Practice and theory of enzyme immunoassays: Laboratory Techniques in Biochemistry
25 and Molecular Biology," Elsevier Science Publishers, NL (1985).

Yet another embodiment of the present invention relates to a kit for detecting the presence of Plasmodium species in a
30 sample, wherein said kit comprises: (i) a first container means containing an antibody according to the invention, and (ii) second container means containing a conjugate comprising a binding partner of the antibody and a label. In another

preferred embodiment, the kit further comprises one or more other containers comprising one or more of the following: wash reagents and reagents capable of detecting the presence of bound antibodies.

5

Examples of detection reagents include, but are not limited to, labeled secondary antibodies, or in the alternative, if the primary antibody is labeled, the chromophoric, enzymatic, or antibody binding reagents which are capable of reacting with the labeled antibody.

10

Having now generally described the present invention, the same may be more readily understood by reference to the following example in connecting with the accompanying Figures 1-2.

15

FIG. 1 schematically illustrates the route of synthesis to the conformationally constrained peptidomimetic Asn-Pro-Asn-Ala-Asn-(Apro-Asn-Ala-Asn-Pro-Asn-Ala-Asn-Pro-Asn-Glu)_{cyclo}-Asn-Pro-Asn-Ala attached to a PE moiety.

20

FIG. 2 shows the grade of purification of the inventive peptidomimetic according to formula (IV) in an HPLC chromatogram. After completing the synthesis, the solvent was removed and the resulting residue purified using a C4 reverse phase HPLC column (Vydac 214 TP 1010, 25cmx10mm) using a gradient starting with 50 % ethanol in water to 100 % ethanol (+0.1% TFA) over 15 minutes. UK39 appears as a broad peak at about 90 % ethanol. m/z 1427 (M+2H)²⁺.

25

30

FIG. 3 shows a comparison of average NMR structures in aqueous solution of the prior art peptide BP66 and the peptide UK39 according to the present invention. BP66 only differs to UK39 in that the modified proline unit (2S,4S)-4-aminoproline has been replaced by the (2S,3R)-3-aminoproline isomer. Due to the change of the position of the cross-link from the 3- to the 4-position, the conformation of the entire macrocyclic portion of the peptide has been changed. The figure shows average NMR structures deduced in water by NMR and dynamic simulated annealing (SA). The figure was prepared using MOLMOL (Konradi R. et al., J. Mol. Graph. 1996, 14, 51-55).

It should be understood that the following example is for illustrative purposes only and should not be construed as limiting this invention in any way to the specific embodiment recited therein.

Unless otherwise specified, general chemical and peptide synthesis procedures, such as those set forth in Voet, Biochemistry, Wiley, 1990; Stryer; Peptide Chemistry. A Practical Textbook, 2nd ed., Miklos Bodanszky, Springer-Verlag, Berlin, 1993; Principles of Peptide Synthesis, 2nd ed., Miklos Bodanszky, Springer-Verlag, Berlin, 1993; Chemical Approaches to the Synthesis of Peptides and Proteins, P. Lloyd-Williams, F. Albericio, E. Giralt, CRC Press, Boca Raton, 1997; Bioorganic Chemistry: Peptides and Proteins, S. M. Hecht, Ed., Oxford Press, Oxford, 1998, are used.

Examples

Synthesis and purification of the inventive peptidomimetic according to formula (V)

The following demonstrates the synthesis of the conformationally constrained peptidomimetic Asn-Pro-Asn-Ala-Asn-(Apro-Asn-Ala-Asn-Pro-Asn-Ala-Asn-Pro-Asn-Glu)_{cyclo}-Asn-Pro-Asn-Ala attached to a PE moiety. The synthesis of the linear peptide precursor was performed on Rink Amide MBHA resin (0.73 mM/g) (Novabiochem) on an Applied Biosystems ABI433A peptide synthesizer. The peptide was synthesized on a 0.25 mmol scale using 4 eq of each Fmoc-protected amino acid each activated with HBTU/HOBt (4 eq.). The amino acids used were: Fmoc-Asn(Mtt)-OH, Fmoc-Pro-OH, Fmoc-Ala-OH, Fmoc-Glu(tBu)-OH and Fmoc-(4S,2S)-4-aminoproline(Boc)-OH.

The cleavage of the linear peptide from the resin was carried out using TFA containing 2.5% TIS and 2.5% water over 3h at room temp. After removal of most of the TFA in vacuo, the peptide was precipitated using diisopropyl ether, washed with iPr₂O, and dried in vacuo for 1h. The product can be analyzed by reverse phase HPLC on a C18 column using a gradient of MeCN/H₂O (+0.1% TFA; 5->95% MeCN ; t_R = 10 min). m/z: 2294 (M+H)⁺.

For cyclization, the crude linear peptide from above was stirred overnight at room temp. together with 4 eq HATU, 4 eq HOAt in DMF and 1 % v/v DIEA (2 mg/ml peptide). The solvent was removed and the peptide dissolved in 20 % piperidine / DMF and stirred for 15 min. at room temp. to remove the Fmoc group. The solvent was evaporated and peptide was precipitated using diisopropyl ether and dried in vacuo. The product can be analyzed by reverse phase HPLC on a C18 column using a gradient of MeCN/H₂O (+0.1% TFA; 5->95% MeCN ; t_R = 11 min). m/z: 2276 (M+H)⁺.

The foregoing product (40 mg) in DMF (5 ml) with 4 eq each of HATU/HOAt was treated with a solution of PE-CO-(CH₂)₂-COOH (PE-succinate; 4 eq.) in DCM (5 ml) together with 1 % of DIEA and stirred overnight at room temp. The solvent was removed and the resulting residue purified using a C4 reverse phase HPLC column (Vydac 214 TP 1010, 25cmx10mm) using a gradient starting with 50 % ethanol in water to 100 % ethanol (+0.1% TFA) over 15 minutes. UK39 appears as a broad peak at about 90 % ethanol. m/z 1427 (M+2H)²⁺ (see below).

Cross-reactivity to antibodies raised against the CS protein of *P. falciparum*

The following demonstrates the cross-reactivity of both UK39 and the isomeric form BP66 to antibodies raised against the CS protein of *P. falciparum*. BP66 merely differs to UK39 in that the modified proline unit (2S,4S)-4-aminoproline (formula VI) of UK39 has been replaced by the (2S,3R)-3-aminoproline isomer.

20

Tab. 1 shows that more monoclonal antibodies raised against the CS protein bind to UK39 than to BP66. Peptidomimetic UK39 comprising (2S,4S)-4-aminoproline thus provide for an improved mimicking of the tandem repeat region of the CS protein

25

The benefits in immunological reactivity arise from the use of (2S,4S)-4-aminoproline rather than the earlier (2S,3R)-3-aminoproline. So the improvement arises by moving the site of cross-linking from the 3- to the 4-position of the proline ring. The immunological response generated by the novel mimetic is stronger and different. This difference is seen in the fact that a monoclonal antibody could be isolated from

30

the immunization with the mimetic containing (2S,4S)-4-aminoproline that binds the parasite but does not bind the mimetic containing 3-aminoproline. This indicates that the new molecule mimics parts of the parasite surface protein that are not represented at all in the earlier mimetic containing (2S,3R)-3-aminoproline (the antibody binds the parasite and the mimetic with (2S,4S)-4-aminoproline but not the mimetic with 2S,3R)-3-aminoproline). This can be easily rationalized, since by changing the position of the cross-link from the 3-position to the 4-position the inventors inevitably changed the conformation (shape) of the macrocyclic portion of the molecule. This is a crucial part for recognition by antibodies. Molecular modelling studies with both mimetics supports the notion that changing the position of the cross-link also changes the conformation of the backbone. The change in conformation may be small. But even small changes in conformation may lead to changes in the way the mimetic is recognized by antibodies, and hence change the ability of the molecule to mimic epitopes on the surface of the parasite (Table 1).

Tab. 1: Immunological cross-reactivity of monoclonal antibodies raised against *P. falciparum* sporozoites with the closely related mimotopes UK39 and BP66.

Mab	BP 66	UK 39
Sp4-5F2	+	+
Sp4-2H1	+	+
Sp3-E6	+	+
Sp3-C6	+	+
Sp3-E9	+	+
Sp4-4B6	+	+
Sp4-7C2	+	+

Sp4-7E4	+	+
Sp4-7H1	-	+
Sp4-4D7	+	+
Sp3-B4-C12	+	+
Sp4-1B4	-	+

Modelling of the conformation of the conformationally constrained peptidomimetics UK39 and BP66

To determine which conformation the constrained peptide antigen will adopt in aqueous solution, the peptidomimetic can be studied by NMR and MD methods in aqueous solution in analogy to previous studies (Bisang, C. et al., J. Am. Chem. Soc. 1998, 120, 7439-7449). Average solution structures for the conformationally constrained peptidomimetics are calculated using NOE-derived distance restraints by dynamic simulated annealing (SA) and molecular dynamics (MD) simulations, using methods described earlier (Bisang, C. et al., J. Am. Chem. Soc. 1998, 120, 7439-7449)

Modeling studies concerning UK39 and BP66 show that cyclization through the (2S,4S)-4-aminoproline residue stabilizes the conformation of UK39. Moreover, modeling reveals that BP66 adopts a different structure than UK39. Therefore, it can be concluded that the change of the cyclization from position 3 to 4 together with the specific stereochemistry of the primary amino group at the modified proline residue creates a new conformation in the NPNA motifs.

Preparation of mimetic-loaded virosomes

For the preparation of PE-mimetic-IRIV, a solution of 4 mg purified Influenza A/Singapore hemagglutinin is centrifuged for 30 min at 100,000g and the pellet is dissolved in 1.33 ml.

of PBS containing 100mM OEG (PBS-OEG). 32 mg phosphatidylcholine (Lipoid, Ludwigshafen, Germany), 6 mg phosphatidylethanolamine and the PE-mimetics are dissolved in a total volume of 2.66 ml PBS-OEG. The phospholipids and the
5 hemagglutinin solution are mixed and sonicated for 1 min. This solution is then centrifuged for 1 hour at 100,000g and the supernatant is sterile filtered (0.22 µm). Virosomes are then formed by detergent removal using BioRad SM BioBeads (BioRad, Glattbrugg, Switzerland).

10

Immunogenicity studies for the UK39-peptidomimetic-IRIV

Antibody responses elicited by IRIVs loaded with the conformationally constrained peptidomimetic UK39 are studied in BALB/c mice. Preimmunization is achieved with the
15 influenza vaccine Inflexal Berna TM (Berna-Products, Bern, Switzerland). Immunization is achieved with several doses of IRIV-peptidomimetic.

BALB/c mice are preimmunized intramuscularly with commercial
20 whole virus influenza vaccine (0.1 ml; Inflexal Berna, Berna Products, Bern, Switzerland) on day 21. Starting on day 0, they received at three-weekly intervals three doses of conformationally constrained UK39 peptidomimetic linked to IRIV intramuscularly at doses of 50 µg of mimetic. Blood is
25 collected two weeks after the third immunization and analyzed by ELISA and IFA.

Enzyme-Linked Immunosorbent Assays (ELISA)

ELISA microtiter plates (Immunolon 413, Dynatech, Embrach,
30 Switzerland) are coated at 4°C overnight with 50 µl of a 5 mg/ml solution of peptidomimetic constructs in PBS, pH 7.2. Wells are then blocked with 5% milk powder in PBS for 1 h at 37°C followed by three washings with PBS containing 0.05%

Tween-20. Plates are then incubated with twofold serial dilutions of mouse serum or hybridoma cell supernatants in PBS containing 0.05% Tween-20 and 0.5% milk powder for 2 h at 37°C. After washing, the plates are incubated with alkaline phosphatase-conjugated goat anti mouse IgG (g-chain specific) antibodies (Sigma, St. Louis, MO) for 1 h at 37°C and then washed. Phosphatase substrate (1 mg/ml p-nitrophenyl phosphate, Sigma) in buffer (0.14% Na₂CO₃, 0.3% NaHCO₃, 0.02% MgCl₂, pH 9.6) is added and incubated at room temperature. The optical density (OD) of the reaction product is recorded after appropriate time at 405 nm using a microplate reader (Titertek Multiscan MCC/340, Labsystems, Finland). Titration curves are registered and analyzed using GENESIS LITE 2.16 software (Life Sciences Ltd., Basingstoke, UK). Effective dose 20% values (ED20%) are calculated for each curve and the corresponding titers are set as endpoint titers.


Immunofluorescence assays

Immunofluorescence assays are performed to assess cross-reactivity of the antibodies obtained. Air-dried unfixed *P. falciparum* salivary gland sporozoites (strain NF54) attached to microscope glass slides are incubated in a moist chamber for 20 min at 37 °C with serum diluted in PBS. The slides are then washed five times with PBS containing 0.1% bovine serum albumin (PBS-BSA) and dried. FITC-labelled goat anti-mouse IgG (Fab-specific) antibodies (Sigma), optimally diluted in PBS containing 0.1 g/l Evans43blue (Merck, Germany), are added. After incubation for 20 min at 37 °C the slides are again washed five times with PBS-BSA, dried, mounted with glycerol, and covered with a cover slide. A Leitz Dialux 20 microscope using 12.5/18 ocular and a 40x/1.30 oil fluorescence 160/0.17 objective is used to detect

fluorescence staining at 495 nm excitation and 525 nm emission wavelengths.

Claims

1. A method for synthesizing a conformationally constrained peptide comprising one or more regions of general
5 formula (I):



10 wherein X = an amino acid residue or an amino acid analog and can be the same or different if $n > 1$;

$m, o \geq 0$ and $n \geq 1$, preferably ≥ 2 ;

$Y, Z = A$ or B and $Y \neq Z$, wherein

15 A = a (2S,4S)-4-FgA-proline residue, wherein FgA is a functional group at the 4-position of the proline residue;

B = an amino acid residue or an amino acid analog having a side chain functional group FgB,

20 wherein the functional groups FgA and FgB are capable of forming an internal link by coupling the functional group FgA at the 4-position of the proline residue and the side chain functional group FgB of B;

25 said method comprising the following steps of

- (a) providing amino acid residues Y and Z having appropriate functional groups FgA and FgB, said functional groups being optionally protected,
- (b) synthesizing a linear peptide comprising the amino
30 acid residues Y and Z, and
- (c) deprotecting, if said functional groups are optionally protected, and reacting the functional groups FgA and FgB for converting the linear peptide

into the cross-linked form by coupling the functional group FgA at the 4-position of the proline residue and the side chain functional group FgB of B.

5 2. The method of claim 1 wherein said modified 4-FgA-proline residue is a modified (2S,4S)-4-amino-proline residue.

10 3. The method of claims 1 or 2 further comprising the step (d) of attaching the cross-linked peptide to a phospholipid moiety.

15 4. The method of claim 3 wherein said phospholipid anchor is attached via a linker, preferably a dicarboxylate linker, more preferably a succinate linker.

5. The method of any of claims 1 to 4,
wherein A = an (2S,4S)-4-aminoproline residue;

20 B = an amino acid residue or an amino acid analog having a side chain carboxyl group, wherein an internal cross-link is formed between A and B by amide coupling the amino group at the 4-position of the (2S,4S)-4-aminoproline residue and the carboxyl group of B.

25

6. The method of any of claims 1 to 5 wherein $n \geq 2$.

30 7. The method of any of claims 1 to 6 wherein the amino acid residue or an amino acid analog B is selected from the group consisting of glutamate and aspartate.

8. The method of claim 7 wherein the amino acid residue is glutamate.

9. The method of any of claims 1 to 8 wherein said linear peptide prepared in step (b) comprises one or more portions of the malaria circumsporozoite (CS) protein of a Plasmodium species, preferably Plasmodium falciparum.

10. The method of claim 9 wherein said linear peptide prepared in step (b) comprises one or more tetrapeptides selected from the group consisting of Asn-Pro-Asn-Ala, Asn-Pro-Asn-Val, Asp-Pro-Asn-Ala and Asp-Pro-Asn-Val.

11. The method of claim 10 wherein said linear peptide prepared in step (b) comprises 3 to 10, preferably 4 to 6 of said tetrapeptides.

12. The method of claims 10 or 11 wherein at least one of said Asn-Pro-Asn-Ala tetrapeptides is replaced by Asn-Pro-Asn-Glu and wherein the glutamate residue of said Asn-Pro-Asn-Glu is said B of general formula (I).

13. The method of any of claims 1 to 12, wherein solid phase synthesis techniques are used in the synthesizing step (b).

14. A peptide obtainable by the method of any of claims 1 to 13.


15. The peptide of claim 14 wherein the internal cross-link stabilizes a supersecondary 3D structure of the peptide.

16. The peptide of claim 14 or 15 comprising one or more units of formula (II):

(N-terminus)...-Asn-Pro-Asn-Ala-Asn-Pro-(Asn-Ala-Asn-Pro)₂-Asn-Glu-Asn-Pro-Asn-Ala...(C-terminus)

wherein Apro is (2S,4S)-4-aminoproline.

17. The peptide of claim 14 or 15 having the formula
5 (III):



wherein Apro is (2S,4S)-4-aminoproline.

18. The peptide of any of claims 14 to 17 wherein said
peptide is coupled to a phospholipid.

10

19. The peptide of claim 18, wherein the N-terminus of
said peptide is coupled via a linker, preferably a
dicarboxylate linker, more preferably a succinate linker to a
fatty acid derivative of phosphaditylethanolamine, preferably
15 1-palmitoyl-3-oleoyl-phosphatidylethanolamine (PE).

20. The peptide of any of claims 14 to 19, wherein said
peptide is delivered with an immunopotentiating delivery
system which is selected from the group consisting of
20 liposomes, multiple-antigen peptides and immunopotentiating
reconstituted virosomes.

21. The peptide of claim 20, wherein the internally
cross-linked peptide of formula (III) comprising a PE moiety
25 at the N-terminus is attached to an immunopotentiating
reconstituted influenza virosome.

22. A composition comprising one or more peptides of any
of claims 14 to 21.

23. The composition of claim 22 further comprising one or more pharmaceutically acceptable adjuvants.

5 24. Use of a 4-Fg-proline, preferably a (2S,4S)-4-Fg-proline for synthesizing conformationally constrained peptides, wherein Fg is a functional group, selected from amino, hydroxy, sulfhydryl, halogen, sulfonyl, carboxy, thiocarboxy or substituted derivatives thereof.

10 25. Use of claim 24, wherein Fg is amino or C1-C12-alkylsubstituted amino.

15 26. Use of (2S,4S)-4-aminoproline in the method of claims 1 to 13, wherein (2S,4S)-4-aminoproline is incorporated during the assembling step (b).

20 27. Use of the peptide of any of claims 14 to 21 for the manufacture of a vaccine for treating malaria.

25 28. A method for producing an antibody in a host against Plasmodium species, preferably against Plasmodium falciparum comprising the step of administering a peptide of any of claims 14 to 21 to said host.

29. Antibody obtainable by the method of claim 28.

30 30. An in vitro method for detecting Plasmodium species in a sample comprising the steps (a) contacting said sample with an antibody of claim 29 under conditions such that binding to CS protein epitopes occurs if CS protein is present; and (b) detecting the presence of said antibody bound to an CS protein epitope.

31. Kit for detecting the presence of Plasmodium species
in a sample, wherein said kit comprises: (i) a first
container means containing an antibody of claims 28, and (ii)
5 second container means containing a conjugate comprising a
binding partner of the antibody and a label.

Fig. 1/3







Fig. 2/3

Fig. 3/3



SEQUENCE LISTING

<110> Pevion Biotech Ltd.
 University Zürich
 Swiss Tropical Institute

<120> Methods for synthesizing conformationally constrained peptides,
 peptidomimetics and the use thereof as synthetic vaccines

<130> P200925

<150> EP 03012520.7
 <151> 2003-06-02

<160> 10

<170> PatentIn version 3.2

<210> 1
 <211> 4
 <212> PRT
 <213> Artificial

<220>
 <223> Tandem repeat sequence of the central portion of CS-protein

<400> 1
 Asn Ala Asn Pro
 1

<210> 2
 <211> 4
 <212> PRT
 <213> Artificial

<220>
 <223> Tandem repeat sequence of the central portion of CS-protein

<400> 2
 Asn Val Asp Pro
 1

<210> 3
 <211> 12
 <212> PRT
 <213> Artificial

<220>
 <223> Tandem repeat sequence of the central portion of CS-protein

<400> 3
 Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro
 1 5 10

<210> 4
 <211> 5
 <212> PRT
 <213> Artificial

<220>
 <223> Tandem repeat sequence of the central portion of CS-protein

<400> 4

Asn Pro Asn Ala Asn
1 5

<210> 5
<211> 4
<212> PRT
<213> Artificial

<220>
<223> Portion of the malaria circumsporozoite (CS) protein of a Plasmodium species

<400> 5

Asn Pro Asn Ala
1

<210> 6
<211> 4
<212> PRT
<213> Artificial

<220>
<223> Portion of the malaria circumsporozoite (CS) protein of a Plasmodium species

<400> 6

Asn Pro Asn Val
1

<210> 7
<211> 4
<212> PRT
<213> Artificial

<220>
<223> Portion of the malaria circumsporozoite (CS) protein of a Plasmodium species

<400> 7

Asp Pro Asn Ala
1

<210> 8
<211> 4
<212> PRT
<213> Artificial

<220>
<223> Portion of the malaria circumsporozoite (CS) protein of a Plasmodium species

<400> 8

Asp Pro Asn Val
1

<210> 9
<211> 4

<212> PRT
 <213> Artificial

<220>
 <223> Modified protein sequence of a tandem repeat sequence of the
 central portion of CS-protein

<400> 9

Asn Pro Asn Glu
 1

<210> 10
 <211> 20
 <212> PRT
 <213> Artificial

<220>
 <223> conformational constrained peptidomimetic

<220>
 <221> misc_feature
 <222> (6)..(6)
 <223> Amide coupling of (2S,4S)-4-aminoproline in pos. 6 with glutamate
 residue in pos. 16

<400> 10

Asn Pro Asn Ala Asn Xaa Asn Ala Asn Pro Asn Ala Asn Pro Asn Glu
 1 5 10 15

Asn Pro Asn Ala
 20

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/005952

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K7/56 C07K1/04 C07K16/20 A61K38/12 A61K39/015

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ZHANG W-J ET AL: "Novel cyclic analogs of angiotensin II with cyclization between positions 5 and 7: Conformational and biological implications" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 39, no. 14, 1996, pages 2738-2744, XP002234087 ISSN: 0022-2623 cited in the application	1,2, 13-15, 22,24
Y	the whole document	1-31
X	EP 0 341 603 A (ABBOTT LAB) 15 November 1989 (1989-11-15) the whole document	1,2,6, 13-15, 22,24
	----- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* & * document member of the same patent family

Date of the actual completion of the international search

24 August 2004

Date of mailing of the international search report

03/09/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Vogt, T

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2004/005952

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PFEIFFER ET AL.: "A Virosome-mimotope approach to synthetic vaccine design and optimization: Synthesis, conformation, and immune recognition of a potential malaria-vaccine candidate." ANGEW. CHEM. INT. ED., vol. 42, 30 May 2003 (2003-05-30), pages 2368-2371, XP002254701	29,30
Y	the whole document	1-31
X	MORENO R ET AL: "Exploiting conformationally constrained peptidomimetics and an efficient human-compatible delivery system in synthetic vaccine design." CHEMBIOCHEM: A EUROPEAN JOURNAL OF CHEMICAL BIOLOGY. GERMANY 5 NOV 2001, vol. 2, no. 11, 5 November 2001 (2001-11-05), pages 838-843, XP002254702 ISSN: 1439-4227	29,30
Y	the whole document	1-31

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/005952

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
EP 0341603	A	15-11-1989	EP	0341603 A2	15-11-1989
			WO	8910935 A1	16-11-1989