

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 May 2001 (25.05.2001)

PCT

(10) International Publication Number  
WO 01/36631 A1

(51) International Patent Classification<sup>7</sup>: C12N 15/12,  
C07K 14/47, 16/18

(21) International Application Number: PCT/GB00/04345

(22) International Filing Date:  
14 November 2000 (14.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
9926995.3 15 November 1999 (15.11.1999) GB  
0001550.3 24 January 2000 (24.01.2000) GB

(71) Applicant: SMITHKLINE BEECHAM P.L.C.  
[GB/GB]; New Horizons Court, Brentford, Middle-  
sex TW8 9EP (GB).

(72) Inventors: MICHALOVICH, David; SmithKline  
Beecham Pharmaceuticals, New Frontiers Science Park  
South, Third Avenue, Harlow, Essex CM19 5AW (GB).  
PRINJHA, Rabinder; SmithKline Beecham Pharmaceu-  
ticals, New Frontiers Science Park South, Third Avenue,  
Harlow, Essex CM19 5AW (GB).

(74) Agent: CONNELL, Anthony, Christopher; Corporate  
Intellectual Property, SmithKline Beecham, Two New  
Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(81) Designated State (national): JP.

(84) Designated States (regional): European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE, TR).

Published:

— With international search report.

[Continued on next page]

(54) Title: HUMAN NOGO-C POLYNUCLEOTIDE AND POLYPEPTIDE AND THEIR USES

A

MEDLDQSPVSSSDSPRPQPAFKYQVREPEDEEEEEEEDEDEDELEEVLERKPA60  
AGLSAAPVPTAPAGAPLMDFGNDFVPPAPRGPLPAAPPVAPERQPSWDPSPVSTVPAP120  
SPLSAAAVSPSKLPEDDEPPARPPPPASVSPQAEFVWTPAPAPAAPFSTPAAPKRRG180  
SSGSVDLTFALPAASEPVIRSSAENMDLKEQPGNTISAGQEDFPSVLLTAASLPSLSP240  
LSAASFKEHEYLGNLSLTPTEGLQENVSEASKVSEKAKTLLIDRLTFSELEYSEM300  
GSSFSVSPKAESAIVANFREEIIVKNKDEEKLVSNNILHMQELPTALTKLVEDEVV360  
SSEKAKDSFNEKRVAVEAPMRSEYADFKPFRVNEVKOSKEDSDMLAAGKTESNLESKV420  
DKKCFADSLQTNHEKDSSESSNDTSFSTPBGIKDRPGAYITCAPFNPATESIATNIF480  
PLLGDPTSSENKTDKKEKKAQIVTEKNTSTKTSNFFLVAAQDSRTDVTNLTNLT540  
EVVANMPEGLTFDLVQACSESELNEVTGKIAETKMDLVQTSEVMQESLYPAAQLCPSF600  
ESEATPSPVLPDIVMEAPLNSAVPSAGASVIQPSSSPLEASSVNYSEIKHEPENPPYE660  
EAMSVSLKRVSGIKKEIKPENINAALQTEAPYISLACDLIKETKLSAEPADPDFDYSE720  
MAKVBPQVPDHSSELVEDSSPDSEFVDFSDSIPDPVQKQDETVMVLKESLTETSFESMI780  
EVENKRLSALPPEGGKPYLESFKSLDNTKDTLLPDEVSTLSKKEKIPLOMEKSLTAVY840  
SNDLFLISKEAQIRETETFSDDSPIETIDEFPTLISKTDSPSKLAREYTDLEVSHKSEI900  
ANAPDGAGSLPCTELPHDLSLKNIQPKVEEKISFSDDFSKNGSATSKVLLLPDVSALAT960  
QAEIESIVKPKVLVKEARKKLPSTTEKEDRSFSAELSKTSVVDLLYWRDLKKTGVV1020  
FGASLPLLSLTVFSIVSVTAYIALALLSVTISFRIYKGVIAIQKSDGHPFRAYLESE1080  
VAISEELVQKYSNSALGHVNCTIKELRRLFLVDDLVDLSLKFVLMWVTVYVGLFNGLTL1140  
LILALISLSFVSVPIYERHQAQIDHYLGLANKNVKDAMAKIQAKIPGLKRAE

1192

B

MEDLDQSPVSSSDSPRPQPAFKYQVREPEDEEEEEEEDEDEDELEEVLERKPA60  
AGLSAAPVPTAPAGAPLMDFGNDFVPPAPRGPLPAAPPVAPERQPSWDPSPVSTVPAP120  
SPLSAAAVSPSKLPEDDEPPARPPPPASVSPQAEFVWTPAPAPAAPFSTPAAPKRRG180  
SSGSVVDLLYWRDLKKTGVVFGASLPLLSLTVFSIVSVTAYIALALLSVTISFRIYK240  
VIQAIQKSDGHPFRAYLESEVAISEELVQKYSNSALGHVNCTIKELRRLFLVDDLVDLSL300  
KFVLMWVTVYVGLFNGLTLILALISLSFVSVPIYERHQAQIDHYLGLANKNVKDAMAK360  
IQAKIPGLKRAE

373

C

MDGQKKNWKKVVDLLYWRDLKKTGVVFGASLPLLSLTVFSIVSVTAYIALALLSVTIS60  
FRIYKGVIAIQKSDGHPFRAYLESEVAISEELVQKYSNSALGHVNCTIKELRRLFLVD120  
DLVDLSLKFVLMWVTVYVGLFNGLTLILALISLSFVSVPIYERHQAQIDHYLGLANKNV180  
KDAMAKIQAKIPGLKRAE



WO 01/36631 A1



— *Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## HUMAN NOGO-C POLYNUCLEOTIDE AND POLYPEPTIDE AND THEIR USES

**Field of the Invention**

5 This invention relates to newly identified polypeptides and polynucleotides encoding such polypeptides, to their use in diagnosis and in identifying compounds that may be agonists, antagonists that are potentially useful in therapy, and to production of such polypeptides and polynucleotides.

**Background of the Invention**

10 The drug discovery process is currently undergoing a fundamental revolution as it embraces "functional genomics", that is, high throughput genome- or gene-based biology. This approach as a means to identify genes and gene products as therapeutic targets is rapidly superceding earlier approaches based on "positional cloning". A phenotype, that is a biological function or genetic disease, would be identified and this would then be tracked back to the responsible gene, based on its  
15 genetic map position.

Functional genomics relies heavily on high-throughput DNA sequencing technologies and the various tools of bioinformatics to identify gene sequences of potential interest from the many molecular biology databases now available. There is a continuing need to identify and characterise further genes and their related polypeptides/proteins, as targets for drug discovery.

20

**Summary of the Invention**

The present invention relates to NOGO-C, in particular NOGO-C polypeptides and NOGO-C polynucleotides, recombinant materials and methods for their production. Such polypeptides and polynucleotides are of interest in relation to methods of treatment of certain  
25 diseases, including, but not limited to, neuropathies, spinal injury, brain injury, stroke, neuronal degeneration for example Alzheimer's disease and Parkinson's disease, neuromuscular disorders, psychiatric disorders and developmental disorders, hereinafter referred to as " diseases of the invention". In a further aspect, the invention relates to methods for identifying agonists and antagonists (*e.g.*, inhibitors) using the materials provided by the invention, and treating conditions  
30 associated with NOGO-C imbalance with the identified compounds. In a still further aspect, the invention relates to diagnostic assays for detecting diseases associated with inappropriate NOGO-C activity or levels.

**Description of the Invention**

35 In a first aspect, the present invention relates to NOGO-C polypeptides. Such polypeptides include (a) an isolated polypeptide encoded by a polynucleotide comprising the sequence of SEQ ID NO:1;

- (b) an isolated polypeptide encoded by a polynucleotide having the sequence of SEQ ID NO:1;
- (c) the polypeptide having the amino acid sequence of SEQ ID NO:2; and
- (d) fragments and variants of such polypeptides in (a) to (c).

Polypeptides of the present invention are believed to be members of the reticulon family of

- 5 polypeptides. They are therefore of interest because members of this family have been shown to display prominent but not exclusive expression in cells of the nervous system. Expression of one isoform of these polypeptides, NSP-C (J. Hens et al. *Cell Tissue Res.* 292:229-237, 1998) has been shown to correlate with neuronal differentiation. Alternative splicing of the genes of this family of polypeptides is known to generate differentially expressed isoforms with overlapping
- 10 and distinct functions in different tissues (J.G. Geisler et al. *Mamm. Genome* 9:164-173, 1998). Amino acid similarity between members of this family of polypeptides with fragments of a high-molecular weight protein purified from bovine spinal cord (A.A. Spillmann et al. *J. Biol. Chem.* 273:15487-15493, 1998) indicates a potential role in axonal growth inhibition role for these proteins. Similarly, expression of NSP-A in specific cancerous cells (N. Senden et al. *Histochem. Cell Biol.* 108:155-165, 1997) may indicate a potential use of these polypeptides in the diagnosis
- 15 and or treatment of cancers.

- The detection of polynucleotides comprising portions of NOGO in human fetal brain and human adult spinal cord cDNA and an abundant >5kb mRNA isoform in human adult brain together with different transcripts potentially arising by alternative splicing in heart, lung, liver,
- 20 kidney and skeletal muscle suggests that NOGO isoforms may similarly serve overlapping and distinct functions in these tissues. By analogy with the semaphorin family of neurite-modulatory polypeptides it might be postulated that these different isoforms would function in each of these tissues to control local innervation by distinct neuronal populations. Aberrant expression of specific isoforms within, for example, skeletal muscle would be predicted to alter motor and
- 25 sensory neuron innervation in diseases such as ALS.

The presence of characteristic signature polypeptides and highly hydrophobic regions in the polypeptide of the present invention suggests that its expression in regions of the nervous system and in tissues forming boundaries for growth may modulate growth and pathfinding both during development and following pathological or injurious processes.

- 30 Expression of the polypeptide of the invention, fragments thereof or alternatively spliced variants of the polypeptide on the surface of cells either naturally, as a secreted protein or following release by cellular damage may act on other cells either through specific receptors or pathologically through non-specific interactions to modulate cell attachment, spreading, migration or growth. This inhibition might be expected to be either reversible or permanent possibly
- 35 resulting in cell death. .

The biological properties of the NOGO-C are hereinafter referred to as "biological activity of NOGO-C" or "NOGO-C activity". Preferably, a polypeptide of the present invention exhibits at least one biological activity of NOGO-C.

Polypeptides of the present invention also includes variants of the aforementioned polypeptides, including all allelic forms and splice variants. Such polypeptides vary from the reference polypeptide by insertions, deletions, and substitutions that may be conservative or non-conservative, or any combination thereof. Particularly preferred variants are those in which several, for instance from 50 to 30, from 30 to 20, from 20 to 10, from 10 to 5, from 5 to 3, from 3 to 2, from 2 to 1 or 1 amino acids are inserted, substituted, or deleted, in any combination.

Preferred fragments of polypeptides of the present invention include an isolated polypeptide comprising an amino acid sequence having at least 30, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID NO: 2, or an isolated polypeptide comprising an amino acid sequence having at least 30, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO: 2. Preferred fragments are biologically active fragments that mediate the biological activity of NOGO-C, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also preferred are those fragments that are antigenic or immunogenic in an animal, especially in a human.

Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention. The polypeptides of the present invention may be in the form of the "mature" protein or may be a part of a larger protein such as a precursor or a fusion protein. It is often advantageous to include an additional amino acid sequence that contains secretory or leader sequences, pro-sequences, sequences that aid in purification, for instance multiple histidine residues, or an additional sequence for stability during recombinant production.

Polypeptides of the present invention can be prepared in any suitable manner, for instance by isolation from naturally occurring sources, from genetically engineered host cells comprising expression systems (*vide infra*) or by chemical synthesis, using for instance automated peptide synthesizers, or a combination of such methods.. Means for preparing such polypeptides are well understood in the art.

In a further aspect, the present invention relates to NOGO-C polynucleotides. Such polynucleotides include:

- (a) an isolated polynucleotide encoding the polypeptide of SEQ ID NO:2;
- (b) an isolated polynucleotide having the polynucleotide sequence of SEQ ID NO:1; and

polynucleotides that are fragments and variants of the above mentioned polynucleotides or that are complementary to above mentioned polynucleotides, over the entire length thereof.

Preferred fragments of polynucleotides of the present invention include an isolated polynucleotide having at least 15, 30, 50 or 100 contiguous nucleotides from the sequence of SEQ ID NO: 1 or an isolated polynucleotide having at least 30, 50 or 100 contiguous nucleotides truncated or deleted from the sequence of SEQ ID NO: 1.

Preferred variants of polynucleotides of the present invention include splice variants, allelic variants, and polymorphisms, including polynucleotides having one or more single nucleotide polymorphisms (SNPs).

In a further aspect, the present invention provides polynucleotides that are RNA transcripts of the DNA sequences of the present invention. Accordingly, there is provided an RNA polynucleotide encoding the polypeptide of SEQ ID NO:2 and an RNA transcript of the DNA sequence of SEQ ID NO:1 and RNA polynucleotides that are complementary thereto.

The polynucleotide sequence of SEQ ID NO:1 is a splice variant of the human NOGO gene which maps to human chromosome 2p21. Two splice variants of human NOGO have already been disclosed in WO00/05364 (SmithKline Beecham). The polynucleotide sequence of SEQ ID NO:1 is a cDNA sequence that encodes the polypeptide of SEQ ID NO:2. The polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence of SEQ ID NO:1 or it may be a sequence other than SEQ ID NO:1, which, as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2. The polypeptide of the SEQ ID NO:2 is related to other proteins of the reticulon family, having homology and/or structural similarity with Human neuroendocrine-specific protein C (A.J. Roebroek et al., J. Biol. Chem. 268: 13439-13447, 1993). The amino acid sequence of SEQ ID NO:2 has been deposited in GenBank with accession number AJ251385.

WO00/05364 (SmithKline Beecham) discloses NOGO-A and NOGO-B splice-variant polynucleotide and polypeptide sequences. SEQ ID NO:3 shows the cDNA sequence of NOGO-A which encodes the polypeptide of SEQ ID NO:4 (deposited in GenBank as AJ251383). SEQ ID NO:5 shows the cDNA sequence of NOGO-B which encodes the polypeptide of SEQ ID NO:6 (deposited in GenBank as AJ251384).

Preferred polypeptides and polynucleotides of the present invention are expected to have, *inter alia*, similar biological functions/properties to their homologous polypeptides and polynucleotides. Furthermore, preferred polypeptides and polynucleotides of the present invention have at least one NOGO-C activity.

Polynucleotides of the present invention may be obtained using standard cloning and screening techniques from a cDNA library derived from mRNA in cells of human fetal brain and

spinal cord, (see for instance, Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

5 When polynucleotides of the present invention are used for the recombinant production of polypeptides of the present invention, the polynucleotide may include the coding sequence for the mature polypeptide, by itself, or the coding sequence for the mature polypeptide in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence that  
10 facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites  
15 and sequences that stabilize mRNA.

Polynucleotides that are identical, or have sufficient identity to a polynucleotide sequence of SEQ ID NO:1, may be used as hybridization probes for cDNA and genomic DNA or as primers for a nucleic acid amplification reaction (for instance, PCR). Such probes and primers may be used to isolate full-length cDNAs and genomic clones encoding polypeptides of the present invention and to  
20 isolate cDNA and genomic clones of other genes (including genes encoding paralogs from human sources and orthologs and paralogs from species other than human) that have a high sequence similarity to SEQ ID NO:1, typically at least 95% identity. Preferred probes and primers will generally comprise at least 15 nucleotides, preferably, at least 30 nucleotides and may have at least 50, if not at least 100 nucleotides. Particularly preferred probes will have between 30 and 50  
25 nucleotides. Particularly preferred primers will have between 20 and 25 nucleotides.

A polynucleotide encoding a polypeptide of the present invention, including homologs from species other than human, may be obtained by a process comprising the steps of screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 or a fragment thereof, preferably of at least 15 nucleotides; and isolating full-length cDNA and  
30 genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to the skilled artisan. Preferred stringent hybridization conditions include overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA; followed by washing the filters in  
35 0.1x SSC at about 65°C. Thus the present invention also includes isolated polynucleotides, preferably with a nucleotide sequence of at least 100, obtained by screening a library under

stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO.1 or a fragment thereof, preferably of at least 15 nucleotides.

The skilled artisan will appreciate that, in many cases, an isolated cDNA sequence will be incomplete, in that the region coding for the polypeptide does not extend all the way through to the 5' terminus. This is a consequence of reverse transcriptase, an enzyme with inherently low "processivity" (a measure of the ability of the enzyme to remain attached to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during first strand cDNA synthesis.

There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman et al., Proc Nat Acad Sci USA 85, 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon (trade mark) technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon (trade mark) technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the cDNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using 'nested' primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the known gene sequence). The products of this reaction can then be analysed by DNA sequencing and a full-length cDNA constructed either by joining the product directly to the existing cDNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

Recombinant polypeptides of the present invention may be prepared by processes well known in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems comprising a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Polynucleotides may be introduced into host cells by methods described in many standard laboratory manuals, such as Davis et al., Basic Methods in Molecular Biology (1986) and Sambrook *et al. (ibid)*. Preferred

methods of introducing polynucleotides into host cells include, for instance, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

5           Representative examples of appropriate hosts include bacterial cells, such as *Streptococci*, *Staphylococci*, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

10           A great variety of expression systems can be used, for instance, chromosomal, episomal and virus-derived systems, *e.g.*, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and  
15           phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector that is able to maintain, propagate or express a polynucleotide to produce a polypeptide in a host may be used. The appropriate polynucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, (*ibid*). Appropriate secretion  
20           signals may be incorporated into the desired polypeptide to allow secretion of the translated protein into the lumen of the endoplasmic reticulum, the periplasmic space or the extracellular environment. These signals may be endogenous to the polypeptide or they may be heterologous signals.

          If a polypeptide of the present invention is to be expressed for use in screening assays, it is generally preferred that the polypeptide be produced at the surface of the cell. In this event, the  
25           cells may be harvested prior to use in the screening assay. If the polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide. If produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

          Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid  
30           extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and/or  
35           purification.

Polynucleotides of the present invention may be used as diagnostic reagents, through detecting mutations in the associated gene. Detection of a mutated form of the gene characterised by the polynucleotide of SEQ ID NO:1 in the cDNA or genomic sequence and which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or altered spatial or temporal expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques well known in the art.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or it may be amplified enzymatically by using PCR, preferably RT-PCR, or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled NOGO-C nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence difference may also be detected by alterations in the electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing (see, for instance, Myers *et al.*, Science (1985) 230:1242). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton *et al.*, Proc Natl Acad Sci USA (1985) 85: 4397-4401).

An array of oligonucleotides probes comprising NOGO-C polynucleotide sequence or fragments thereof can be constructed to conduct efficient screening of *e.g.*, genetic mutations. Such arrays are preferably high density arrays or grids. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability, see, for example, M.Chee *et al.*, Science, 274, 610-613 (1996) and other references cited therein.

Detection of abnormally decreased or increased levels of polypeptide or mRNA expression may also be used for diagnosing or determining susceptibility of a subject to a disease of the invention. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit comprising:

- (a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO: 1, or a fragment or an RNA transcript thereof;
- (b) a nucleotide sequence complementary to that of (a);
- (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or a fragment thereof; or
- (d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, particularly diseases of the invention, amongst others.

The polynucleotide sequences of the present invention are also valuable tools for tissue expression studies. Such studies allow the determination of expression patterns of polynucleotides of the present invention which may give an indication as to the expression patterns of the encoded polypeptides in tissues, by detecting the mRNAs that encode them. The techniques used are well known in the art and include in situ hybridisation techniques to clones arrayed on a grid, such as cDNA microarray hybridisation (Schena *et al*, Science, 270, 467-470, 1995 and Shalon *et al*, Genome Res, 6, 639-645, 1996) and nucleotide amplification techniques such as PCR. A preferred method uses the TAQMAN (Trade mark) technology available from Perkin Elmer. Results from these studies can provide an indication of the normal function of the polypeptide in the organism. In addition, comparative studies of the normal expression pattern of mRNAs with that of mRNAs encoded by an alternative form of the same gene (for example, one having an alteration in polypeptide coding potential or a regulatory mutation) can provide valuable insights into the role of the polypeptides of the present invention, or that of inappropriate expression thereof in disease. Such inappropriate expression may be of a temporal, spatial or simply quantitative nature.

The polypeptides of the present invention are expressed in brain, heart, liver, skeletal muscle, pancreas and kidney; based on Northern blot data provided in figure 1.

A further aspect of the present invention relates to antibodies. The polypeptides of the invention or their fragments, or cells expressing them, can be used as immunogens to produce antibodies that are immunospecific for polypeptides of the present invention. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against polypeptides of the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, or cells to an animal, preferably a non-human animal, using routine protocols. For preparation of monoclonal antibodies, any technique

which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983) 4:72) and the EBV-hybridoma technique (Cole *et al.*, *Monoclonal Antibodies and Cancer Therapy*, 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies, such as those described in U.S. Patent No. 4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms, including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against polypeptides of the present invention may also be employed to treat diseases of the invention, amongst others.

Polypeptides and polynucleotides of the present invention may also be used as vaccines. Accordingly, in a further aspect, the present invention relates to a method for inducing an immunological response in a mammal that comprises inoculating the mammal with a polypeptide of the present invention, adequate to produce antibody and/or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said animal from disease whether that disease is already established within the individual or not. An immunological response in a mammal may also be induced by a method comprises delivering a polypeptide of the present invention *via* a vector directing expression of the polynucleotide and coding for the polypeptide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases of the invention. One way of administering the vector is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a modified nucleic acid, or a DNA/RNA hybrid. For use a vaccine, a polypeptide or a nucleic acid vector will be normally provided as a vaccine formulation (composition). The formulation may further comprise a suitable carrier. Since a polypeptide may be broken down in the stomach, it is preferably administered parenterally (for instance, subcutaneous, intramuscular, intravenous, or intradermal injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions that may contain anti-oxidants, buffers, bacteriostats and solutes that render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions that may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The

vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

5

Polypeptides of the present invention have one or more biological functions that are of relevance in one or more disease states, in particular the diseases of the invention hereinbefore mentioned. It is therefore useful to to identify compounds that stimulate or inhibit the function or level of the polypeptide. Accordingly, in a further aspect, the present invention provides for a method of screening compounds to identify those that stimulate or inhibit the function or level of the polypeptide. Such methods identify agonists or antagonists that may be employed for therapeutic and prophylactic purposes for such diseases of the invention as hereinbefore mentioned. Compounds may be identified from a variety of sources, for example, cells, cell-free preparations, chemical libraries, collections of chemical compounds, and natural product mixtures. Such agonists or antagonists so-identified may be natural or modified substrates, ligands, receptors, enzymes, etc., as the case may be, of the polypeptide; a structural or functional mimetic thereof (see Coliganet *al.*, Current Protocols in Immunology 1(2):Chapter 5 (1991)) or a small molecule.

The screening method may simply measure the binding of a candidate compound to the polypeptide, or to cells or membranes bearing the polypeptide, or a fusion protein thereof, by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve measuring or detecting (qualitatively or quantitatively) the competitive binding of a candidate compound to the polypeptide against a labeled competitor (*e.g.* agonist or antagonist). Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells bearing the polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide of the present invention, to form a mixture, measuring a NOGO-C activity in the mixture, and comparing the NOGO-C activity of the mixture to a control mixture which contains no candidate compound.

Polypeptides of the present invention may be employed in conventional low capacity screening methods and also in high-throughput screening (HTS) formats. Such HTS formats include not only the well-established use of 96- and, more recently, 384-well micotiter plates but also emerging methods such as the nanowell method described by Schullek et al, Anal Biochem., 246, 20-29, (1997).

Fusion proteins, such as those made from Fc portion and NOGO-C polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists for the polypeptide of the present invention (see D. Bennett *et al.*, J Mol Recognition, 8:52-58 (1995); and K. Johanson *et al.*, J Biol Chem, 270(16):9459-9471 (1995)).

5       The polynucleotides, polypeptides and antibodies to the polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to  
10       discover agents that may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

A polypeptide of the present invention may be used to identify membrane bound or soluble receptors, if any, through standard receptor binding techniques known in the art. These include, but are not limited to, ligand binding and crosslinking assays in which the polypeptide is  
15       labeled with a radioactive isotope (for instance,  $^{125}\text{I}$ ), chemically modified (for instance, biotinylated), or fused to a peptide sequence suitable for detection or purification, and incubated with a source of the putative receptor (cells, cell membranes, cell supernatants, tissue extracts, bodily fluids). Other methods include biophysical techniques such as surface plasmon resonance and spectroscopy. These screening methods may also be used to identify agonists and antagonists  
20       of the polypeptide that compete with the binding of the polypeptide to its receptors, if any. Standard methods for conducting such assays are well understood in the art.

Examples of antagonists of polypeptides of the present invention include antibodies or, in some cases, oligonucleotides or proteins that are closely related to the ligands, substrates, receptors, enzymes, etc., as the case may be, of the polypeptide, *e.g.*, a fragment of the ligands, substrates,  
25       receptors, enzymes, etc.; or a small molecule that bind to the polypeptide of the present invention but do not elicit a response, so that the activity of the polypeptide is prevented.

Screening methods may also involve the use of transgenic technology and NOGO-C gene. The art of constructing transgenic animals is well established. For example, the NOGO-C gene may be introduced through microinjection into the male pronucleus of fertilized oocytes, retroviral transfer into pre- or post-implantation embryos, or injection of genetically modified,  
30       such as by electroporation, embryonic stem cells into host blastocysts. Particularly useful transgenic animals are so-called "knock-in" animals in which an animal gene is replaced by the human equivalent within the genome of that animal. Knock-in transgenic animals are useful in the drug discovery process, for target validation, where the compound is specific for the human target. Other useful transgenic animals are so-called "knock-out" animals in which the expression  
35       of the animal ortholog of a polypeptide of the present invention and encoded by an endogenous

DNA sequence in a cell is partially or completely annulled. The gene knock-out may be targeted to specific cells or tissues, may occur only in certain cells or tissues as a consequence of the limitations of the technology, or may occur in all, or substantially all, cells in the animal.

Transgenic animal technology also offers a whole animal expression-cloning system in which introduced genes are expressed to give large amounts of polypeptides of the present invention

Screening kits for use in the above described methods form a further aspect of the present invention. Such screening kits comprise:

- (a) a polypeptide of the present invention;
  - (b) a recombinant cell expressing a polypeptide of the present invention;
  - 10 (c) a cell membrane expressing a polypeptide of the present invention; or
  - (d) an antibody to a polypeptide of the present invention;
- which polypeptide is preferably that of SEQ ID NO:2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

15

### Glossary

The following definitions are provided to facilitate understanding of certain terms used frequently hereinbefore.

“Antibodies” as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

“Isolated” means altered “by the hand of man” from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not “isolated,” but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is “isolated”, as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is “isolated” even if it is still present in said organism, which organism may be living or non-living.

30 “Polynucleotide” generally refers to any polyribonucleotide (RNA) or polydeoxribonucleotide (DNA), which may be unmodified or modified RNA or DNA.

“Polynucleotides” include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, “polynucleotide” refers to triple-stranded regions

35

comprising RNA or DNA or both RNA and DNA. The term "polynucleotide" also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any polypeptide comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as post-translational processing, or by chemical modification techniques that are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications may occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present to the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from post-translation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, biotinylation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination (see, for instance, *Proteins - Structure and Molecular Properties*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993; Wold, F., *Post-translational Protein Modifications: Perspectives and Prospects*, 1-12, in *Post-translational Covalent Modification of Proteins*, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and

nonprotein cofactors", Meth Enzymol, 182, 626-646, 1990, and Rattan *et al.*, "Protein Synthesis: Post-translational Modifications and Aging", Ann NY Acad Sci, 663, 48-62, 1992).

"Fragment" of a polypeptide sequence refers to a polypeptide sequence that is shorter than the reference sequence but that retains essentially the same biological function or activity as the reference polypeptide. "Fragment" of a polynucleotide sequence refers to a polynucleotide sequence that is shorter than the reference sequence of SEQ ID NO:1.

"Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains the essential properties thereof. A typical variant of a polynucleotide differs in nucleotide sequence from the reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from the reference polypeptide. Generally, alterations are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, insertions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. Typical conservative substitutions include Gly, Ala; Val, Ile, Leu; Asp, Glu; Asn, Gln; Ser, Thr; Lys, Arg; and Phe and Tyr. A variant of a polynucleotide or polypeptide may be naturally occurring such as an allele, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis. Also included as variants are polypeptides having one or more post-translational modifications, for instance glycosylation, phosphorylation, methylation, ADP ribosylation and the like. Embodiments include methylation of the N-terminal amino acid, phosphorylations of serines and threonines and modification of C-terminal glycines.

"Allele" refers to one of two or more alternative forms of a gene occurring at a given locus in the genome.

"Polymorphism" refers to a variation in nucleotide sequence (and encoded polypeptide sequence, if relevant) at a given position in the genome within a population.

"Single Nucleotide Polymorphism" (SNP) refers to the occurrence of nucleotide variability at a single nucleotide position in the genome, within a population. An SNP may occur within a gene or within intergenic regions of the genome. SNPs can be assayed using Allele Specific Amplification (ASA). For the process at least 3 primers are required. A common primer is used in reverse complement to the polymorphism being assayed. This common primer can be between 50 and 1500 bps from the polymorphic base. The other two (or more) primers are

identical to each other except that the final 3' base wobbles to match one of the two (or more) alleles that make up the polymorphism. Two (or more) PCR reactions are then conducted on sample DNA, each using the common primer and one of the Allele Specific Primers.

"Splice Variant" as used herein refers to cDNA molecules produced from RNA molecules initially transcribed from the same genomic DNA sequence but which have undergone alternative RNA splicing. Alternative RNA splicing occurs when a primary RNA transcript undergoes splicing, generally for the removal of introns, which results in the production of more than one mRNA molecule each of that may encode different amino acid sequences. The term splice variant also refers to the proteins encoded by the above cDNA molecules.

"Identity" reflects a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, determined by comparing the sequences. In general, identity refers to an exact nucleotide to nucleotide or amino acid to amino acid correspondence of the two polynucleotide or two polypeptide sequences, respectively, over the length of the sequences being compared.

"% Identity" - For sequences where there is not an exact correspondence, a "% identity" may be determined. In general, the two sequences to be compared are aligned to give a maximum correlation between the sequences. This may include inserting "gaps" in either one or both sequences, to enhance the degree of alignment. A % identity may be determined over the whole length of each of the sequences being compared (so-called global alignment), that is particularly suitable for sequences of the same or very similar length, or over shorter, defined lengths (so-called local alignment), that is more suitable for sequences of unequal length.

"Similarity" is a further, more sophisticated measure of the relationship between two polypeptide sequences. In general, "similarity" means a comparison between the amino acids of two polypeptide chains, on a residue by residue basis, taking into account not only exact correspondences between a between pairs of residues, one from each of the sequences being compared (as for identity) but also, where there is not an exact correspondence, whether, on an evolutionary basis, one residue is a likely substitute for the other. This likelihood has an associated "score" from which the "% similarity" of the two sequences can then be determined.

Methods for comparing the identity and similarity of two or more sequences are well known in the art. Thus for instance, programs available in the Wisconsin Sequence Analysis Package, version 9.1 (Devereux J et al, Nucleic Acids Res, 12, 387-395, 1984, available from Genetics Computer Group, Madison, Wisconsin, USA), for example the programs BESTFIT and GAP, may be used to determine the % identity between two polynucleotides and the % identity and the % similarity between two polypeptide sequences. BESTFIT uses the "local homology" algorithm of Smith and Waterman (J Mol Biol, 147,195-197, 1981, Advances in Applied Mathematics, 2, 482-489, 1981) and finds the best single region of similarity between two

sequences. BESTFIT is more suited to comparing two polynucleotide or two polypeptide sequences that are dissimilar in length, the program assuming that the shorter sequence represents a portion of the longer. In comparison, GAP aligns two sequences, finding a "maximum similarity", according to the algorithm of Neddleman and Wunsch (J Mol Biol, 48, 443-453, 1970). GAP is more suited to comparing sequences that are approximately the same length and an alignment is expected over the entire length. Preferably, the parameters "Gap Weight" and "Length Weight" used in each program are 50 and 3, for polynucleotide sequences and 12 and 4 for polypeptide sequences, respectively. Preferably, % identities and similarities are determined when the two sequences being compared are optimally aligned.

Other programs for determining identity and/or similarity between sequences are also known in the art, for instance the BLAST family of programs (Altschul S F et al, J Mol Biol, 215, 403-410, 1990, Altschul S F et al, Nucleic Acids Res., 25:389-3402, 1997, available from the National Center for Biotechnology Information (NCBI), Bethesda, Maryland, USA and accessible through the home page of the NCBI at [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)) and FASTA (Pearson W R, Methods in Enzymology, 183, 63-99, 1990; Pearson W R and Lipman D J, Proc Nat Acad Sci USA, 85, 2444-2448, 1988, available as part of the Wisconsin Sequence Analysis Package).

Preferably, the BLOSUM62 amino acid substitution matrix (Henikoff S and Henikoff J G, Proc. Nat. Acad Sci. USA, 89, 10915-10919, 1992) is used in polypeptide sequence comparisons including where nucleotide sequences are first translated into amino acid sequences before comparison.

Preferably, the program BESTFIT is used to determine the % identity of a query polynucleotide or a polypeptide sequence with respect to a reference polynucleotide or a polypeptide sequence, the query and the reference sequence being optimally aligned and the parameters of the program set at the default value, as hereinbefore described.

"Identity Index" is a measure of sequence relatedness which may be used to compare a candidate sequence (polynucleotide or polypeptide) and a reference sequence. Thus, for instance, a candidate polynucleotide sequence having, for example, an Identity Index of 0.95 compared to a reference polynucleotide sequence is identical to the reference sequence except that the candidate polynucleotide sequence may include on average up to five differences per each 100 nucleotides of the reference sequence. Such differences are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion. These differences may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between these terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. In other words, to obtain a polynucleotide sequence having an Identity Index of 0.95 compared to a reference polynucleotide sequence, an average of up to 5 in every 100 of the

nucleotides of the in the reference sequence may be deleted, substituted or inserted, or any combination thereof, as hereinbefore described. The same applies *mutatis mutandis* for other values of the Identity Index, for instance 0.96, 0.97, 0.98 and 0.99.

Similarly, for a polypeptide, a candidate polypeptide sequence having, for example, an Identity Index of 0.95 compared to a reference polypeptide sequence is identical to the reference sequence except that the polypeptide sequence may include an average of up to five differences per each 100 amino acids of the reference sequence. Such differences are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion. These differences may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between these terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. In other words, to obtain a polypeptide sequence having an Identity Index of 0.95 compared to a reference polypeptide sequence, an average of up to 5 in every 100 of the amino acids in the reference sequence may be deleted, substituted or inserted, or any combination thereof, as hereinbefore described. The same applies *mutatis mutandis* for other values of the Identity Index, for instance 0.96, 0.97, 0.98 and 0.99.

The relationship between the number of nucleotide or amino acid differences and the Identity Index may be expressed in the following equation:

$$n_a \leq x_a - (x_a \bullet I),$$

in which:

$n_a$  is the number of nucleotide or amino acid differences,

$x_a$  is the total number of nucleotides or amino acids in SEQ ID NO:1 or SEQ ID NO:2, respectively,

$I$  is the Identity Index ,

$\bullet$  is the symbol for the multiplication operator, and

in which any non-integer product of  $x_a$  and  $I$  is rounded down to the nearest integer prior to subtracting it from  $x_a$ .

"Homolog" is a generic term used in the art to indicate a polynucleotide or polypeptide sequence possessing a high degree of sequence relatedness to a reference sequence. Such relatedness may be quantified by determining the degree of identity and/or similarity between the two sequences as hereinbefore defined. Falling within this generic term are the terms "ortholog", and "paralog". "Ortholog" refers to a polynucleotide or polypeptide that is the functional equivalent of the polynucleotide or polypeptide in another species. "Paralog" refers to a polynucleotide or polypeptide that within the same species which is functionally similar.

"Fusion protein" refers to a protein encoded by two, often unrelated, fused genes or fragments thereof. In one example, EP-A-0 464 \*\*\* discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, employing an immunoglobulin Fc region as a part of a fusion  
5 protein is advantageous for use in therapy and diagnosis resulting in, for example, improved pharmacokinetic properties [see, *e.g.*, EP-A 0232 262]. On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified.

10 All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is  
15 also incorporated by reference herein in its entirety in the manner described above for publications and references.

## Examples

### Example 1 – Cloning of NOGO cDNAs

Primers designed against ESTs containing peptides reported for a bovine IN-1 reactive inhibitory protein (Spillmann, A.A., et al (1998) *J Biol Chem.* **273**, 19283-19293) were used in standard PCR amplifications from human fetal brain or human spinal cord cDNA (Clontech). These products were cloned into pGEMTeasy (Promega), sequenced and this data used to search for new ESTs extending the known sequence until multiple sequences containing in-frame upstream stop codons were identified. IMAGE clones (NOGO-B IMAGE 590987, NOGO-C IMAGE 561867) were obtained from Research Genetics, and fully sequenced on both strands. The full ORF for NOGO-A was amplified from human spinal cord cDNA (Clontech) using the sense primer (GTGCTCGAGC CAGCCATGGAAGACCTGGACCAGTCTCCTCTGG) and the antisense primer (GAATCTAGACTAATTATTTTGGGCGTTTTTCATTTCAGCTTTGC). Epitope tagged versions of NOGO-A cDNA were made in pcDNA3 (Invitrogen) using standard PCR techniques (N terminal FLAG tag or C terminal myc tag).

### Example 2 - Tissue distribution of NOGO

A radioactively labelled DNA probe corresponding to the C-terminal 2kb of the NOGO open reading frame was hybridised with a Clontech MTN-1 filter containing equivalent loadings of mRNA from each of the indicated tissues and then washed to high stringency to only detect NOGO transcripts. At least three bands are visible (>5kb, 2.4kb and <2kb). The NOGO-A transcript is present in adult brain and to a lesser degree in heart and skeletal muscle. The NOGO-B transcript is present at essentially similar levels in all tissues while the NOGO-C transcript is more specifically expressed, most abundantly in skeletal muscle, brain and kidney with low levels detectable in the pancreas.

### Example 3 – Demonstration of neurite inhibitory activity

In order to assess whether NOGO possessed neurite inhibitory activity a soluble version of the 220 kD isoform (NOGO-A) was prepared. The extracellular region of the 220 kD protein was prepared as a bivalent Fc chimeric protein using a CD33 signal-sequence and human IgG-Fc sequences and the purified protein was placed in model systems to assess neurite inhibitory activity.

### Fc Preparation

The entire putative extracellular domain (ECD) of NOGO-A (residues 1-1024) was amplified by PCR and cloned into signal pIg-plus (R&D Systems) between the CD33 signal sequence and

human IgG sequence using engineered XhoI-XbaI sites to produce a secreted chimeric fusion protein with a carboxy terminal Fc portion. NOGO fusion protein was purified from COS-7 conditioned medium, following DEAE-Dextran transfections, using protein A sepharose (Sigma). Dissociated cerebellar granule neurons (CGC) were allowed to adhere to substrate and then

5 treated with control Fc protein (SIRP-Fc), MAG-Fc, or NOGO-Fc. The control protein SIRP-Fc had no effect on neurite-outgrowth while the inhibitory effects of MAG and NOGO were dose dependent and were equally potent in this model. This experiment shows that recombinant soluble NOGO produced as a bivalent Fc is a potent neurite outgrowth inhibitory molecule comparable to MAG.

10

#### **Example 4 – Purified GST-NOGO fusion proteins inhibit hippocampal neurite outgrowth**

This example investigated the effect of purified GST-NOGO fusion proteins (GST: glutathione-S-transferase) on cultured hippocampal neurite outgrowth. The fusion proteins created for examples 4 and 5 consist of:

15

- i) GST alone;
- ii) GST-Nogo "splice" encoding Nogo-A amino acids 185-1004 (which represent the entire splice region specific to Nogo-A); and
- iii) GST-Nogo "loop" encoding amino acids 1055-1120 (representing the intertransmembrane loop present in all Nogo isoforms).

20

Cultured E18 embryonic hippocampal neurons in polylysine coated 96 well plates at a density of 20,000 cells per well in Eagle's basal medium supplemented to contain 25mM KCl, 10% FCS and 50ug/ml gentamycin were supplemented with purified fusion proteins in triplicate wells then incubated for 48 hours. Cells were fixed with 4% paraformaldehyde then stained using coomassie blue. Neurite length was determined using a KS300 Image analysis system. Results are expressed

25 as a percentage of the length of neurites of cells treated with PBS alone and are shown in Table 1.

Both NOGO fusion proteins clearly inhibit hippocampal neurite-outgrowth confirming the presence of two independent neurite inhibitory domains, one present in the NOGO-A specific splice region and one in the NOGO-A, B and C common intertransmembrane loop region.

30

Table 1.

| Treatment         | Neurite Length (% of untreated control) |
|-------------------|-----------------------------------------|
| GST 20nM          | 93 ± 5.1                                |
| GST 160nM         | 95 ± 9                                  |
| GST-Nogo loop 1nM | 78 ± 3.6                                |

|                       |          |
|-----------------------|----------|
| GST-Nogo loop 20nM    | 60 ± 2.3 |
| GST-Nogo loop 160nM   | 66 ± 1.8 |
| GST-Nogo splice 1nM   | 98 ± 3.1 |
| GST-Nogo splice 20nM  | 95 ± 3.8 |
| GST-Nogo splice 160nM | 62 ± 3.9 |

#### **Example 5 - Purified GST-NOGO fusion proteins inhibit differentiated PC12 neurite outgrowth**

5 NGF differentiated PC12 cells were cultured in 96 well plates and treated with purified GST and GST-Nogo fusion proteins (described in example 4) at the indicated concentrations, incubated for a further 72hours, fixed and stained with coomassie blue. Neurite-length was determined using a KS300 Image analysis system. Results are expressed as the length of neurites in  $\mu\text{m}$  and are shown in Table 2.

10

GST-NOGO loop and GST-NOGO splice both significantly affect neurite outgrowth from PC12 cells at the concentrations tested. Both Nogo fusion proteins clearly inhibit PC12 neurite-outgrowth confirming the presence of two independent neurite inhibitory domains, one present in the NOGO-A specific splice region and one in the NOGO-A, B and C common

15 intertransmembrane loop region.

Table 2.

| Treatment                        | Mean Neurite-length ( $\mu\text{m}$ ) | Standard deviation |
|----------------------------------|---------------------------------------|--------------------|
| No Protein                       | 32.09                                 |                    |
| GST 6 $\mu\text{M}$              | 27.95                                 | 2.31               |
| GSTNogoLoop 2.6 $\mu\text{M}$    | 24.7                                  | 3.43               |
| GSTNogoloop 8 $\mu\text{M}$      | 20.59                                 | 1.99               |
| GSTNogo splice 3.8 $\mu\text{M}$ | 29.49                                 | 5.25               |
| GSTNogo splice 6.4 $\mu\text{M}$ | 20.2                                  | 4.65               |

#### **Example 6 – Nogo expression in pMCAO rat model of stroke**

20 TAQMAN™ analysis of cDNA preparations made from samples of tissues taken from either the ipsilateral or contralateral side of permanent MCAO treated rats was performed to analyse changes in NOGO-A expression following the ischaemic insult. The results are shown in Fig 3.

NOGO-A expression was significantly increased in the MCAO treated samples within three hours and remained high to the end of the trial period at 24 hours. Similar changes were not observed in samples from the contralateral side or in sham operated animal samples.

- 5 These data suggest very strongly that increases in NOGO-A expression correlate with the onset of ischaemic damage and may be a causative factor.

Antagonists of NOGO-A activity could therefore be expected to protect from ischaemic damage as occurs in stroke patients.

**In the figures:**

Figure 1 depicts a Northern blot showing tissue distribution of human NOGO. Where lane 1 contains human adult heart RNA, lane 2 contains human adult brain RNA, lane 3 contains human placental RNA, lane 4 contains human adult lung RNA, lane 5 contains human adult liver RNA, lane 6 contains human adult skeletal muscle RNA, lane 7 contains human adult kidney RNA and lane 8 contains human adult pancreas RNA.

Figure 2 Amino acids matching all six published peptide sequences from purified bNI-220 are shown with a thick underline. The location of putative transmembrane domains is shown double underlined. An acid box motif is marked with a dotted underline. Amino acids absent in the shorter splice variants of NOGO are shown in italics (residues 186-1004).

A. Predicted amino acid sequence of NOGO-A (GenBank Accession number AJ251383).

B. Predicted amino acid sequence of NOGO-B (GenBank Accession number AJ251384).

C. Predicted amino acid sequence of NOGO-C (GenBank Accession number AJ251385).

Figure 3. Nogo-A expression levels in pMCAO rat stroke model. NOGO-A levels are shown as % GAPDH (a standard housekeeping gene). The data is shown for ipsilateral (L) and contralateral (R) in naïve animals and at post-operative time points of 3 hours, 6 hours, 12 hours and 24 hours. Data is also shown for sham operated (white bars) animals.

## SEQUENCE INFORMATION

## SEQ ID NO:1

ATGGACGGTCAGAAGAAAAATTGGAAGGACAAGGTTGTTGACCTCCTGTACTGGAGAGAC  
 ATTAAGAAGACTGGAGTGGTGTGTTGGTGCCAGCCTATTCTGCTGCTTTCATTGACAGTA  
 5 TTCAGCATTGTGAGCGTAACAGCCTACATTGCCTTGGCCCTGCTCTCTGTGACCATCAGC  
 TTTAGGATATACAAGGGTGTGATCCAAGCTATCCAGAAATCAGATGAAGGCCACCCATTC  
 AGGGCATATCTGGAATCTGAAGTTGCTATATCTGAGGAGTTGGTTCAGAAGTACAGTAAT  
 TCTGCTCTTGGTTCATGTGAACTGCACGATAAAGGAACTCAGGCGCCTCTTCTTAGTTGAT  
 GATTTAGTTGATTCTCTGAAGTTTGCAGTGTGATGTGGGTATTTACCTATGTTGGTGCC  
 10 TTGTTTAATGGTCTGACACTACTGATTTTGGCTCTCATTTCACTCTTCAGTGTTCCTGTT  
 ATTTATGAACGGCATCAGGCACAGATAGATCATTATCTAGGACTTGCAAATAAGAATGTT  
 AAAGATGCTATGGCTAAAATCCAAGCAAAAATCCCTGGATTGAAGCGCAAAGCTGAATGA

## SEQ ID NO:2

15 MDGQKKNWKDKVVDLLYWRDIKKTGVVFGASLFLLLSLTVFSIVSVTAYIALALLSVTIS  
 FRIYKGVQIAIQKSDEGHPFRAYLESEVAISEELVQKYSNSALGHVNCTIKELRRLFLVD  
 DLVDSLKFAVLMWVFTYVGALFNGLTLLILALISLFSVPVIYERHQAQIDHYLGLANKNV  
 KDAMAKIQAKIPGLKRKAE

## SEQ ID NO:3

20 ATGGAAGACCTGGACCAGTCTCCTCTGGTCTCGTCCTCGGACAGCCACCCCGGCCGAG  
 CCCGCGTTCAAGTACCAGTTCGTGAGGGAGCCGAGGACGAGGAGGAAGAAGAGGAGGAG  
 GAAGAGGAGGACGAGGACGAAGACCTGGAGGAGCTGGAGGTGCTGGAGAGGAAGCCCGCC  
 GCCGGGCTGTCCGCGGCCCCAGTGCCACCGCCCCCTGCCGCGGGCGCGCCCTGATGGAC  
 25 TTCGGAAATGACTTCGTGCCGCCGGCGCCCCGGGGACCCCTGCCGGCCGCTCCCCCGTC  
 GCCCCGGAGCGGCAGCCGTCTTGGGACCCGAGCCCGGTGTCTGTCGACCGTGCCCGCGCCA  
 TCCCCGCTGTCTGTGCCGAGTCTCGCCCTCCAAGCTCCCTGAGGACGACGAGCCTCCG  
 GCCCCGCTCCCCCTCCTCCCCCGGCCAGCGTGAGCCCCCAGGCAGAGCCCGTGTGGACC  
 CCGCCAGCCCCGGCTCCCCCGCGCCCCCTCCACCCCGGCCGCGCCCAAGCGCAGGGGC  
 30 TCCTCGGGCTCAGTGGATGAGACCCCTTTTTGCTCTTCTGCTGCATCTGAGCCTGTGATA  
 CGTCTCTGCGAGAAAATATGGACTTGAAGGAGCAGCCAGGTAACACTATTTTCGGCTGGT  
 CAAGAGGATTTCCCATCTGTCTCTGCTTGAACTGCTGCTTCTCTTCTTCTGTCTCCT  
 CTCTCAGCCGCTTCTTTCAAAGAACATGAATACCTTGGTAATTTGTCAACAGTATTACCC  
 ACTGAAGGAACACTTCAAGAAAATGTGAGTGAAGCTTCTAAAGAGGTCTCAGAGAAGGCA  
 35 AAAACTCTACTCATAGATAGAGATTTAACAGAGTTTTTCAAGATTAGAATACTCAGAAATG  
 GGATCATCGTTCAGTGTCTCTCCAAAAGCAGAATCTGCCGTAATAGTAGCAAATCCTAGG  
 GAAGAAAATAATCGTGAAAAATAAAGATGAAGAAGAGAAGTTAGTTAGTAATAACATCCTT  
 CATAATCAACAAGAGTTACCTACAGCTCTTACTAAATTGGTTAAAGAGGATGAAGTTGTG  
 TCTTCAGAAAAAGCAAAAGACAGTTTTAATGAAAAGAGAGTTGCAGTGGAAGCTCCTATG  
 40 AGGGAGGAATATGCAGACTTCAAACATTTGAGCGAGTATGGGAAGTGAAAGATAGTAAG  
 GAAGATAGTGATATGTTGGCTGCTGGAGGTAAATCGAGAGCAACTTGGAAGTAAAGTG  
 GATAAAAAATGTTTTGCAGATAGCCTTGAGCAAACTAATCACGAAAAAGATAGTGAGAGT  
 AGTAATGATGATACTTCTTTCCCCAGTACGCCAGAAGGTATAAAGGATCGTCCAGGAGCA

TATATCACATGTGCTCCCTTTAACCCAGCAGCAACTGAGAGCATTGCAACAAACATTTTT  
CCTTTGTTAGGAGATCCTACTTCAGAAAATAAGACCGATGAAAAAAAAATAGAAGAAAAG  
AAGGCCCAAATAGTAACAGAGAAGAATACTAGCACCAAACATCAAACCCCTTTCTTGTA  
GCAGCACAGGATTCTGAGACAGATTATGTACACAACAGATAATTTAACAAAGGTGACTGAG  
5 GAAGTCGTGGCAAACATGCCTGAAGGCCTGACTCCAGATTTAGTACAGGAAGCATGTGAA  
AGTGAATTGAATGAAGTTACTGGTACAAAGATTGCTTATGAAACAAAATGGACTTGGTT  
CAAACATCAGAAGTTATGCAAGAGTCACTCTATCCTGCAGCACAGCTTTGCCCATCATT  
GAAGAGTCAGAAGCTACTCCTTCACCAGTTTTGCCTGACATTGTTATGGAAGCACCATTG  
AATTCTGCAGTTCCTAGTGCTGGTGCTTCCGTGATACAGCCCAGCTCATCACCATTAGAA  
10 GCTTCTTCAGTTAATTATGAAAGCATAAAACATGAGCCTGAAAACCCCCCACCATATGAA  
GAGGCCATGAGTGTATCACTAAAAAAGTATCAGGAATAAAGGAAGAAATTAAAGAGCCT  
GAAAATATTAATGCAGCTCTTCAAGAAACAGAAGCTCCTTATATATCTATTGCATGTGAT  
TTAATTAAAGAAACAAAGCTTTCTGCTGAACCAGCTCCGGATTTCTCTGATTATTCAGAA  
ATGGCAAAGTTGAACAGCCAGTGCCTGATCATTCTGAGCTAGTTGAAGATTCCTCACCT  
15 GATTCTGAACCAGTTGACTTATTTAGTGATGATTCAATACCTGACGTTCCACAAAAACAA  
GATGAAACTGTGATGCTTGTGAAAGAAAGTCTCACTGAGACTTCATTTGAGTCAATGATA  
GAATATGAAAATAAGGAAAAACTCAGTGCTTTGCCACCTGAGGGAGGAAAGCCATATTTG  
GAATCTTTTAAGCTCAGTTTAGATAACACAAAAGATACCCTGTTACCTGATGAAGTTTCA  
ACATTGAGCAAAAAGGAGAAAATTCCTTTGCAGATGGAGGAGCTCAGTACTGCAGTTTAT  
20 TCAAATGATGACTTATTTATTTCTAAGGAAGCACAGATAAGAGAACTGAAACGTTTTCA  
GATTCTATCTCCAATTGAAATTATAGATGAGTTCCTACATTGATCAGTTCTAAAAGTAT  
TCATTTTCTAAATTAGCCAGGGAATATACTGACCTAGAAGTATCCACAAAAGTGAAATT  
GCTAATGCCCCGGATGGAGCTGGGTTCATTGCCTTGACAGAAATTGCCCCATGACCTTTCT  
TTGAAGAACATACAACCCAAAGTTGAAGAGAAAATCAGTTTCTCAGATGACTTTTCTAAA  
25 AATGGGTCTGCTACATCAAAGGTGCTCTTATTGCCTCCAGATGTTTCTGCTTTGGCCACT  
CAAGCAGAGATAGAGAGCATAGTTAAACCCAAAGTTCTTGTAAGAAGCTGAGAAAAAA  
CTTCCTTCCGATACAGAAAAAGAGGACAGATCACCATCTGCTATATTTTCAGCAGAGCTG  
AGTAAAACTTCAGTTGTTGACCTCCTGTACTGGAGAGACATTAAGAAGACTGGAGTGGTG  
TTTGGTGCCAGCCTATTCTGCTGCTTTTCATTGACAGTATTGAGCATTGTGAGCGTAACA  
30 GCCTACATTGCCTTGGCCCTGCTCTCTGTGACCATCAGCTTTAGGATATACAAGGGTGTG  
ATCCAAGCTATCCAGAAATCAGATGAAGGCCACCCATTGAGGCATATCTGGAATCTGAA  
GTTGCTATATCTGAGGAGTTGGTTTCAAGTACAGTAATTCTGCTCTTGGTCATGTGAAC  
TGCACGATAAAGGAACTCAGGCGCCTCTTCTTAGTTGATGATTTAGTTGATTCTCTGAAG  
TTTGCAGTGTGATGTGGGTATTTACCTATGTTGGTGCCTTGTTTAATGGTCTGACACTA  
35 CTGATTTTGGCTCTCATTTCACTCTTCAGTGTTTCTGTTATTTATGAACGGCATCAGGCG  
CAGATAGATCATTATCTAGGACTTGCAAATAAGAATGTTAAAGATGCTATGGCTAAAATC  
CAAGCAAAAATCCCTGGATTGAAGCGCAAAGCTGAATGA

**SEQ ID NO:4**

40 MEDLDQSPVLVSSSDSPRPQPAFKYQFVREPEDEEEEEEEEEDEDEDLEELEVLERKPA  
AGLSAAPVPTAPAAGAPLMDFGNDFVPPAPRGPLPAAPPVAPERQPSWDSPVSSTVPAP  
SPLSAAAVSPSKLPEDDEPPARPPPPPPASVSPQAEPVWTPPAPAPAAPPSTPAAPKRRG  
SSGSVDETLFALPAASEPVIRSSAENMDLKEQPGNTISAGQEDFPSVLLLETAASLPSLP

LSAASFKEHEYLGNLSTVLPTEGTLQENVSEASKEVSEKAKTLLIDRDLTEFSELEYSEM  
 GSSFSVSPKAESAVIVANPREEIIIVKNKDEEEKLVSNILHNQQELPTALTCLKVKEDEVV  
 SSEKAKDSFNEKRVAEAPMREEYADFKPFERVWEVKDSKEDSDMLAAGGKIESNLESKV  
 DKKCFADSLEQTNHEKDSSESNDDTSFPSTPEGIKDRPGAYITCAPFNPAATESIATNIF  
 5 PLLGDPTSENKTDEKKIEEKKQIVTEKNTSTKTSNPFLVAAQDSETDYVTTDNLTKVTE  
 EVVANMPEGLTPDLVQEACESELNEVTGTKIAYETKMDLVQTSEVMQESLYPAAQLCPSF  
 EESEATPSPVLPDIVMEAPLNSAVPSAGASVIQPSSSPLEASSVNYESIKHEPENPPPYE  
 EAMSVSLKKVSGIKEEIKEPENINAALQETEAPYISIACDLIKETKLSAEPAPDFSDYSE  
 MAKVEQPVPDHSSELVEDSSPDSEPVDFSDDSIPDVPQKQDETVMVLVKESLTETSFESMI  
 10 EYENKEKLSALPPEGKPYLESFKLSLDNTKDTLLPDEVSTLSKKEKIPLQMEELSTAVY  
 SNDDLFISKEAQIRETETFSDSSPIEIIDEFPTLISSKTDSSFKLAREYTDLEVSHKSEI  
 ANAPDGAGSLPCTELPHDLSLKNIQPKVEEKISFSDDFSKNGSATSQVLLLPDVSALAT  
 QAEIESIVKPKVLVKEAEKLPSTDEKEDRSPSAIFSAELSKTSVVDLLYWRDIKKTGVV  
 FGASLFLLLSLTVFSIVSVTAYIALALLSVTISFRIYKQVIAIQKSDEGHPFRAYLESE  
 15 VAISEELVQKYSNSALGHVNCTIKELRRLFLVDDLVDLSLKFAVLMWVFTYVGALFNGLTLL  
 LILALISLFSVPVIYERHQAQIDHYLGLANKNVKDAMAKIQAKIPGLKRKAE

**SEQ ID NO:5**

ATGGAAGACCTGGACCAGTCTCCTCTGGTCTCGTCCCTCGGACAGCCACCCCGGCCGAG  
 20 CCCGCGTTCAAGTACCAGTTCGTGAGGGAGCCCGAGGACGAGGAGGAAGAAGAGGAGGAG  
 GAAGAGGAGGACGAGGACGAAGACCTGGAGGAGCTGGAGGTGCTGGAGAGGAAGCCCGCC  
 GCCGGGCTGTCCGCGGCCCCAGTGCCACCGCCCCCTGCCGCCGGCGCGCCCCCTGATGGAC  
 TTCGGAATGACTTCGTGCCGCCGGCGCCCCGGGACCCCTGCCGCCCGCTCCCCCGTC  
 GCCCCGGAGCGGCAGCCGTCTTGGGACCCGAGCCCGGTGTCGTCGACCGTGCCCCGCGCA  
 25 TCCCCGCTGTCTGCTGCCGAGTCTCGCCCTCCAAGCTCCCTGAGGACGACGAGCCTCCG  
 GCCCCGCCTCCCCCTCCTCCCCGGCCAGCGTGAGCCCCCAGGCAGAGCCCGTGTGGACC  
 CCGCCAGCCCCGGCTCCCGCCGCGCCCCCTCCACCCCGCGCGCCCAAGCGCAGGGGC  
 TCCTCGGGCTCAGTGTTGTTGACCTCCTGTACTGGAGAGACATTAAGAAGACTGGAGTG  
 GTGTTTGGTGCCAGCCTATTCTGCTGCTTTTATTGACAGTATTAGCATTGTGAGCGTA  
 30 ACAGCCTACATTGCCTTGGCCCTGCTCTGTGACCATCAGCTTTAGGATATACAAGGGT  
 GTGATCCAAGCTATCCAGAAATCAGATGAAGGCCACCCATTAGGGCATATCTGGAATCT  
 GAAGTTGCTATATCTGAGGAGTTGGTTCAGAAGTACAGTAATTCTGCTCTTGGTCATGTG  
 AACTGCACGATAAAGGAACTCAGGCGCCTCTTCTTAGTTGATGATTTAGTTGATTCTCTG  
 AAGTTTGCAGTGTTGATGTGGGTATTTACCTATGTTGGTGCCCTGTTTAAATGGTCTGACA  
 35 CTACTGATTTTGGCTCTCATTTCACTCTTCAGTGTTTCTGTTATTTATGAACGGCATCAG  
 GCACAGATAGATCATTATCTAGGACTTGCAAATAAGAATGTTAAAGATGCTATGGCTAAA  
 ATCCAAGCAAAAATCCCTGGATTGAAGCGCAAAGCTGAATGA

**SEQ ID NO:6**

40 MEDLDQSPLVSSSDSPRPQPAFKYQFVREPEDEEEEEEEEEDEDEDLEELEVLERKPA  
 AGLSAAVPPTAPAAGAPLMDFGNDFVPPAPRGPLPAAPPVAPERQPSWDPSPVSSVTPAP  
 SPLSAAAVSPSKLPEDDEPPARPPPPPPASVSPQAEPVWTPAPAPAAPSTPAAPKRRG  
 SSGSVVVDLLYWRDIKKTGVVFGASLFLLLSLTVFSIVSVTAYIALALLSVTISFRIYKG

VIQAIQKSDEGHPFRAYLESEVAISEELVQKYSNSALGHVNCTIKELRRLFLVDDLVDL  
KFAVLMWVFYVVGALFNGLTLLILALISLFSVPVIYERHQAQIDHYLGLANKNVKDAMAK  
IQAKIPGLKRKAE

5

**Claims**

1. An isolated polypeptide encoded by a polynucleotide comprising the sequence of SEQ ID NO:1.
2. An isolated polypeptide according to claim 1 encoded by a polynucleotide having the sequence of  
5 SEQ ID NO:1.
3. An isolated polypeptide according to claim 1 or 2 having the amino acid sequence of SEQ ID NO:2.
- 10 4. An isolated polypeptide which is a fragment or variant of an isolated polypeptide of any one of claims 1 to 3.
5. An isolated polynucleotide selected from the group consisting of:  
(a) an isolated polynucleotide encoding the polypeptide of SEQ ID NO:2;  
15 (b) an isolated polynucleotide having the polynucleotide sequence of SEQ ID NO:1; and  
(c) an isolated polynucleotide that is a fragment or variant of the polynucleotides of (a) or (b); or polynucleotides that are complementary to such polynucleotides, over the entire length thereof.
6. An expression system comprising a polynucleotide capable of producing a polypeptide of claim  
20 1 when said expression vector is present in a compatible host cell.
7. A recombinant host cell comprising the expression vector of claim 6 or a membrane thereof expressing the polypeptide of claim 1.
- 25 8. A process for producing a polypeptide of claim 1 comprising the step of culturing a host cell as defined in claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture medium.
9. An antibody immunospecific for the polypeptide of any one of claims 1 to 3.
- 30 10. A method for screening to identify compounds that stimulate or inhibit the function or level of the polypeptide of claim 1 comprising a method selected from the group consisting of:  
(a) measuring or, detecting, quantitatively or qualitatively, the binding of a candidate compound to the polypeptide (or to the cells or membranes expressing the polypeptide) or a fusion protein  
35 thereof by means of a label directly or indirectly associated with the candidate compound;

- (b) measuring the competition of binding of a candidate compound to the polypeptide (or to the cells or membranes expressing the polypeptide) or a fusion protein thereof in the presence of a labeled competitor;
- (c) testing whether the candidate compound results in a signal generated by activation or
- 5 inhibition of the polypeptide, using detection systems appropriate to the cells or cell membranes expressing the polypeptide;
- (d) mixing a candidate compound with a solution containing a polypeptide of claim 1, to form a mixture, measuring activity of the polypeptide in the mixture, and comparing the activity of the mixture to a control mixture which contains no candidate compound; or
- 10 (e) detecting the effect of a candidate compound on the production of mRNA encoding said polypeptide or said polypeptide in cells, using for instance, an ELISA assay.

Figure 1




Figure 2

**A**

MEDLDQSPLVSSSDSPRPQPAFKYQFVREPEDEEEEEEEEEDEDEDLEELEVLERKPA 60  
 AGLSAAPVPTAPAAGAPLMDFGNDFVPPAPRGPLPAAPPVAPERQPSWDPSVVSSTVPAP 120  
 SPLSAAAVSPSKLPEDDEPPARPPPPPPASVSPQAEPVWTPPAPAPAAPPSTPAAPKRRG 180  
 SSGSVDETFLFALPAASEPVIRSSAENMDLKEQPGNTISAGQEDFPSVLLETAASLPSLSP 240  
 LSAASFKEHEYLGNLSTVLPTEGTLQENVSEASKEVSEKAKTLLIDRDLTEFSELEYSEM 300  
 GSSFSVSPKAESAVIVANPREEIIVKNKDEEEKLVSNILHNQQELPTALTKLVKEDEVV 360  
 SSEKAKDSFNEKRVAVEAPMREEYADFKPFERVWEVKDSKEDSDMLAAGGKIESNLESKV 420  
 DKKCFADSLEQTNHEKDSSESSNDDTSFPSTPEGIKDRPGAYITCAPFNPAATESIATNIF 480  
 PLLGDPTSENKTDEKKIEEKKAIQIVTEKNTSTKTSNPFLVAAQDSETDYVTTDNLTKVTE 540  
 EVVANMPEGLTPDLVQEACESELNEVTGTKIAYETKMDLVQTSEVMQESLYPAAQLCPSF 600  
 EESEATPSPVLPDIVMEAPLNSAVPSAGASVIQSSSPLEASSVNYESIKHEPENPPPYE 660  
 EAMSVSLKKVSGIKEEIKEPENINAALQETEAPYISIACDLIKETKLSAEPAPDFSDYSE 720  
 MAKVEQPVPDHSSELVEDSSPDSEPVDLFSDDSIQDVPQKQDETVMVLKESLTETSFESMI 780  
 EYENKEKLSALPPEGGKPYLESFKLSLDNTKDTLLPDEVSTLSKKEKIPLQMEELSTAVY 840  
 SNDDLFIKSAQIRETETFSDDSSPIEIIIDEFPTLISSKTDSFSKLAREYTDLEVSHKSEI 900  
 ANAPDGAGSLPCTELPHDLSLKNIQPKVEEKISFSDDFSKNGSATSKVLLLPDVSALAT 960  
 QAEIESIVKPKVLVKEAEKKLPSDTEKEDRSPSAIFSAELSKTSVVDLLYWRDIKKTGVV 1020  
 FGASLFLLLSLTVFSIVSVTAYIALALLSVTISFRIYKGVIAIQKSDEGHPFRAYLESE 1080  
 VAISEELVQKYSNSALGHVNCTIKELRRLFLVDDLVDLSLKFAVLMWVFTYVGALFNGLTL 1140  
 LILALISLFSVPVIYERHQAQIDHYLGLANKNVKDAMAKIQAKIPGLKRKAE 1192

**B**


MEDLDQSPLVSSSDSPRPQPAFKYQFVREPEDEEEEEEEEEDEDEDLEELEVLERKPA 60  
 AGLSAAPVPTAPAAGAPLMDFGNDFVPPAPRGPLPAAPPVAPERQPSWDPSVVSSTVPAP 120  
 SPLSAAAVSPSKLPEDDEPPARPPPPPPASVSPQAEPVWTPPAPAPAAPPSTPAAPKRRG 180  
 SSGSVVVDLLYWRDIKKTGVVFGASLFLLLSLTVFSIVSVTAYIALALLSVTISFRIYKG 240  
 VIQAIQKSDEGHPFRAYLESEVAISEELVQKYSNSALGHVNCTIKELRRLFLVDDLVDLSL 300  
 KFAVLMWVFTYVGALFNGLTLLILALISLFSVPVIYERHQAQIDHYLGLANKNVKDAMAK 360  
 IQAKIPGLKRKAE 373

Figure 2 (continued)

**C**

MDGQKKNWKDKVVDLLYWRDIKKTGVVFGASLFLLLSLTVFSIVSVTAYIALALLSVTIS 60  
FRIYKGVIAIQKSDEGHPFRAYLESEVAISEELVQKYSNSALGHVNCTIKELRRLFLVD 120  
DLVDSLKFAVLMWVFTYVGALFNGLTLLILALISLFSVPVIYERHQAQIDHYLGLANKNV 180  
KDAMAKIQAKIPGLKRKAE

Figure 3



## SEQUENCE LISTING

5           <110> SmithKline Beecham plc  
           <120> Novel Compounds  
           <130> GP30165A  
 10           <160> 6  
           <170> FastSEQ for Windows Version 3.0  
 15           <210> 1  
           <211> 600  
           <212> DNA  
           <213> Homo sapiens  
 20           <400> 1  
 atggacggtc agaagaaaaa ttggaaggac aaggttgttg acctcctgta ctggagagac       60  
 attaagaaga ctggagtggg gtttggtgcc agcctattcc tgctgctttc attgacagta       120  
 ttcagcattg tgagcgtaac agcctacatt gccttggccc tgctctctgt gaccatcagc       180  
 tttaggatat acaaggggtg gatccaagct atccagaaat cagatgaagg ccacccattc       240  
 25 agggcatatc tggaatctga agttgctata tctgaggagt tggttcagaa gtacagtaat       300  
 tctgctcttg gtcattgtga ctgcacgata aaggaactca ggcgcctctt cttagttgat       360  
 gatttagttg attctctgaa gtttgctgag ttgatgtggg tatttaccta tgttgggtgcc       420  
 ttgtttaatg gtctgacact actgattttg gctctcattt cactcttcag tgttcctggt       480  
 atttatgaac ggcatcaggc acagatagat cattatctag gacttgcaaa taagaatggt       540  
 30 aaagatgcta tggctaaaat ccaagcaaaa atccctggat tgaagcgcaa agctgaatga       600  
           <210> 2  
           <211> 199  
           <212> PRT  
 35           <213> Homo sapiens  
           <400> 2  
 Met Asp Gly Gln Lys Lys Asn Trp Lys Asp Lys Val Val Asp Leu Leu  
           1                   5                   10                   15  
 40 Tyr Trp Arg Asp Ile Lys Lys Thr Gly Val Val Phe Gly Ala Ser Leu  
                   20                   25                   30  
 Phe Leu Leu Leu Ser Leu Thr Val Phe Ser Ile Val Ser Val Thr Ala

35 40 45  
 Tyr Ile Ala Leu Ala Leu Leu Ser Val Thr Ile Ser Phe Arg Ile Tyr  
 50 55 60  
 Lys Gly Val Ile Gln Ala Ile Gln Lys Ser Asp Glu Gly His Pro Phe  
 5 65 70 75 80  
 Arg Ala Tyr Leu Glu Ser Glu Val Ala Ile Ser Glu Glu Leu Val Gln  
 85 90 95  
 Lys Tyr Ser Asn Ser Ala Leu Gly His Val Asn Cys Thr Ile Lys Glu  
 100 105 110  
 10 Leu Arg Arg Leu Phe Leu Val Asp Asp Leu Val Asp Ser Leu Lys Phe  
 115 120 125  
 Ala Val Leu Met Trp Val Phe Thr Tyr Val Gly Ala Leu Phe Asn Gly  
 130 135 140  
 Leu Thr Leu Leu Ile Leu Ala Leu Ile Ser Leu Phe Ser Val Pro Val  
 15 145 150 155 160  
 Ile Tyr Glu Arg His Gln Ala Gln Ile Asp His Tyr Leu Gly Leu Ala  
 165 170 175  
 Asn Lys Asn Val Lys Asp Ala Met Ala Lys Ile Gln Ala Lys Ile Pro  
 180 185 190  
 20 Gly Leu Lys Arg Lys Ala Glu  
 195

&lt;210&gt; 3

&lt;211&gt; 3579

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 3

atggaagacc tggaccagtc tctctctggtc tcgtcctcgg acagcccacc ccggccgcag 60  
 30 cccgcgttca agtaccagtt cgtgagggag cccgaggacg aggaggaaga agaggaggag 120  
 gaagaggagg acgaggacga agacctggag gagctggagg tgctggagag gaagcccgcc 180  
 gccgggctgt ccgcggcccc agtgcccacc gcccctgccg ccggcgcgcc cctgatggac 240  
 ttcggaatatg acttcgtgcc gccggcgccc cggggacccc tgccggccgc tcccccgctc 300  
 gccccggagc ggcagccgtc ttgggacccg agcccgggtgt cgtcgaccgt gcccgcgcca 360  
 35 tccccgctgt ctgctgccgc agtctcgccc tccaagctcc ctgaggacga cgagcctccg 420  
 gcccggcctc cccctcctcc cccggccagc gtgagccccc aggcagagcc cgtgtggacc 480  
 ccgccagccc cggctcccg cgcgcccccc tccaccccgg ccgcgccc aa ggcagggggc 540  
 tctcgggct cagtggatga gacccttttt gctcttctctg ctgcatctga gcctgtgata 600  
 cgctcctctg cagaaaaatat ggacttgaag gagcagccag gtaacactat ttcggctggt 660  
 40 caagaggatt tcccatctgt cctgcttgaa actgctgctt ctcttccttc tctgtctcct 720  
 ctctcagccg cttcttttaa agaacatgaa taccttggtg atttgtcaac agtattaccc 780  
 actgaaggaa cacttcaaga aaatgtcagt gaagcttcta aagaggtctc agagaaggca 840

|    |                                                                    |      |
|----|--------------------------------------------------------------------|------|
|    | aaaactctac tcatagatag agatttaaca gagttttcag aattagaata ctcagaaatg  | 900  |
|    | ggatcatcgt tcagtgtctc tccaaaagca gaatctgccg taatagtagc aaatcctagg  | 960  |
|    | gaagaaataa tcgtgaaaaa taaagatgaa gaagagaagt tagttagtaa taacatcctt  | 1020 |
|    | cataatcaac aagagttacc tacagctctt actaaattgg ttaaagagga tgaagttgtg  | 1080 |
| 5  | tcttcagaaa aagcaaaaga cagttttaat gaaaagagag ttgcagtgga agctcctatg  | 1140 |
|    | agggaggaat atgcagactt caaaccattt gagcgagtat gggaagtga agatagtaag   | 1200 |
|    | gaagatagtg atatgttggc tgctggaggt aaaatcgaga gcaacttgga aagtaaagtg  | 1260 |
|    | gataaaaaat gttttgcaga tagccttgag caaactaatc acgaaaaaga tagtgagagt  | 1320 |
|    | agtaatgatg atacttcttt cccagtagc ccagaaggta taaaggatcg tccaggagca   | 1380 |
| 10 | tatatcacat gtgtccctt taaccagca gcaactgaga gcattgcaac aaacattttt    | 1440 |
|    | cctttgttag gagatcctac ttcagaaaat aagaccgatg aaaaaaaat agaagaaaag   | 1500 |
|    | aaggcccaa tagtaacaga gaagaatact agcaccaaaa catcaaacc ttttcttgta    | 1560 |
|    | gcagcacagg attctgagac agattatgtc acaacagata atttaacaaa ggtgactgag  | 1620 |
|    | gaagtcgtgg caaacatgcc tgaaggcctg actccagatt tagtacagga agcatgtgaa  | 1680 |
| 15 | agtgaattga atgaagttac tggtaaaaag attgcttatg aaacaaaaat ggacttgggt  | 1740 |
|    | caaacatcag aagttagtca agagtcactc tatctcgag cacagctttg cccatcattt   | 1800 |
|    | gaagagtcag aagctactcc ttcaccagtt ttgcctgaca ttgttatgga agcaccattg  | 1860 |
|    | aattctgcag ttcttagtgc tgggtgcttcc gtgatacagc ccagctcatc accattagaa | 1920 |
|    | gcttcttcag ttaattatga aagcataaaa catgagcctg aaaaccccc accatatgaa   | 1980 |
| 20 | gaggccatga gtgtatcact aaaaaaagta tcaggaataa aggaagaaat taaagagcct  | 2040 |
|    | gaaaatatta atgcagctct tcaagaaaca gaagtcctt atatatctat tgcagtgtat   | 2100 |
|    | ttaattaaag aaacaaagct ttctgctgaa ccagctccgg atttctctga ttattcagaa  | 2160 |
|    | atggcaaaag ttgaacagcc agtgcctgat cattctgagc tagttgaaga ttcctcacct  | 2220 |
|    | gattctgaac cagttgactt atttagtgat gattcaatac ctgacgttcc acaaaaacaa  | 2280 |
| 25 | gatgaaactg tgatgcttgt gaaagaaagt ctactgaga cttcatttga gtcaatgata   | 2340 |
|    | gaatatgaaa ataaggaaaa actcagtgtc ttgccacctg agggaggaaa gccatatttg  | 2400 |
|    | gaatctttta agctcagttt agataacaca aaagataccc tgttacctga tgaagtttca  | 2460 |
|    | acattgagca aaaaggagaa aattcctttg cagatggagg agctcagtac tgcagtttat  | 2520 |
|    | tcaaatgatg acttatttat ttctaaggaa gcacagataa gagaaactga aacgttttca  | 2580 |
| 30 | gattcatctc caattgaaat tatagatgag ttccctacat tgatcagttc taaaactgat  | 2640 |
|    | tcattttcta aattagccag ggaatatact gacctagaag tatcccacaa aagtgaattt  | 2700 |
|    | gctaattgcc cggtggagc tgggtcattg ccttgccacag aattgcccc tgacctttct   | 2760 |
|    | ttgaagaaca tacaacccaa agttgaagag aaaatcagtt tctcagatga cttttctaaa  | 2820 |
|    | aatgggtctg ctacatcaaa ggtgctctta ttgcctccag atgtttctgc tttggccact  | 2880 |
| 35 | caagcagaga tagagagcat agttaaaccc aaagtctctg tgaaagaagc tgagaaaaaa  | 2940 |
|    | cttccttcg atacagaaaa agaggacaga tcaccatctg ctatattttc agcagagctg   | 3000 |
|    | agtaaaactt cagttgttga cctcctgtac tggagagaca ttaagaagac tggagtgggtg | 3060 |
|    | tttggtgcca gcctattcct gctgctttca ttgacagtat tcagcattgt gagcgtaaca  | 3120 |
|    | gcctacattg ccttgccct gctctctgtg accatcagct ttaggatata caagggtgtg   | 3180 |
| 40 | atccaagcta tccagaaatc agatgaaggc caccatttca gggcatatct ggaatctgaa  | 3240 |
|    | gttgctatat ctgaggagtt ggttcagaag tacagtaatt ctgctcttgg tcatgtgaac  | 3300 |
|    | tgcacgataa aggaactcag gcgcctcttc ttagttgatg atttagttga ttctctgaag  | 3360 |

```

tttgcagtgt tgatgtgggt atttacctat gttggtgcct tgtttaatgg tctgacacta 3420
ctgatttttg ctctcatttc actcttcagt gttcctgtta tttatgaacg gcatcaggcg 3480
cagatagatc attatctagg acttgcaaat aagaatgtta aagatgctat ggctaaaatc 3540
caagcaaaaa tccttggatt gaagcgcaaa gctgaatga 3579

```

5

```

<210> 4
<211> 1192
<212> PRT
<213> Homo sapiens

```

10

```

<400> 4

```

15

```

Met Glu Asp Leu Asp Gln Ser Pro Leu Val Ser Ser Ser Asp Ser Pro
 1          5          10          15
Pro Arg Pro Gln Pro Ala Phe Lys Tyr Gln Phe Val Arg Glu Pro Glu
20          25          30
Asp Glu Glu Glu Glu Glu Glu Glu Glu Glu Asp Glu Asp Glu Asp
35          40          45
Leu Glu Glu Leu Glu Val Leu Glu Arg Lys Pro Ala Ala Gly Leu Ser
50          55          60
20 Ala Ala Pro Val Pro Thr Ala Pro Ala Ala Gly Ala Pro Leu Met Asp
65          70          75          80
Phe Gly Asn Asp Phe Val Pro Pro Ala Pro Arg Gly Pro Leu Pro Ala
85          90          95
Ala Pro Pro Val Ala Pro Glu Arg Gln Pro Ser Trp Asp Pro Ser Pro
25          100          105          110
Val Ser Ser Thr Val Pro Ala Pro Ser Pro Leu Ser Ala Ala Ala Val
115          120          125
Ser Pro Ser Lys Leu Pro Glu Asp Asp Glu Pro Pro Ala Arg Pro Pro
130          135          140
30 Pro Pro Pro Pro Ala Ser Val Ser Pro Gln Ala Glu Pro Val Trp Thr
145          150          155          160
Pro Pro Ala Pro Ala Pro Ala Ala Pro Pro Ser Thr Pro Ala Ala Pro
165          170          175
Lys Arg Arg Gly Ser Ser Gly Ser Val Asp Glu Thr Leu Phe Ala Leu
35          180          185          190
Pro Ala Ala Ser Glu Pro Val Ile Arg Ser Ser Ala Glu Asn Met Asp
195          200          205
Leu Lys Glu Gln Pro Gly Asn Thr Ile Ser Ala Gly Gln Glu Asp Phe
210          215          220
40 Pro Ser Val Leu Leu Glu Thr Ala Ala Ser Leu Pro Ser Leu Ser Pro
225          230          235          240
Leu Ser Ala Ala Ser Phe Lys Glu His Glu Tyr Leu Gly Asn Leu Ser

```

|    |                                                                 |     |     |
|----|-----------------------------------------------------------------|-----|-----|
|    | 245                                                             | 250 | 255 |
|    | Thr Val Leu Pro Thr Glu Gly Thr Leu Gln Glu Asn Val Ser Glu Ala |     |     |
|    | 260                                                             | 265 | 270 |
|    | Ser Lys Glu Val Ser Glu Lys Ala Lys Thr Leu Leu Ile Asp Arg Asp |     |     |
| 5  | 275                                                             | 280 | 285 |
|    | Leu Thr Glu Phe Ser Glu Leu Glu Tyr Ser Glu Met Gly Ser Ser Phe |     |     |
|    | 290                                                             | 295 | 300 |
|    | Ser Val Ser Pro Lys Ala Glu Ser Ala Val Ile Val Ala Asn Pro Arg |     |     |
|    | 305                                                             | 310 | 315 |
| 10 | Glu Glu Ile Ile Val Lys Asn Lys Asp Glu Glu Glu Lys Leu Val Ser |     |     |
|    | 325                                                             | 330 | 335 |
|    | Asn Asn Ile Leu His Asn Gln Gln Glu Leu Pro Thr Ala Leu Thr Lys |     |     |
|    | 340                                                             | 345 | 350 |
|    | Leu Val Lys Glu Asp Glu Val Val Ser Ser Glu Lys Ala Lys Asp Ser |     |     |
| 15 | 355                                                             | 360 | 365 |
|    | Phe Asn Glu Lys Arg Val Ala Val Glu Ala Pro Met Arg Glu Glu Tyr |     |     |
|    | 370                                                             | 375 | 380 |
|    | Ala Asp Phe Lys Pro Phe Glu Arg Val Trp Glu Val Lys Asp Ser Lys |     |     |
|    | 385                                                             | 390 | 395 |
| 20 | Glu Asp Ser Asp Met Leu Ala Ala Gly Gly Lys Ile Glu Ser Asn Leu |     |     |
|    | 405                                                             | 410 | 415 |
|    | Glu Ser Lys Val Asp Lys Lys Cys Phe Ala Asp Ser Leu Glu Gln Thr |     |     |
|    | 420                                                             | 425 | 430 |
|    | Asn His Glu Lys Asp Ser Glu Ser Ser Asn Asp Asp Thr Ser Phe Pro |     |     |
| 25 | 435                                                             | 440 | 445 |
|    | Ser Thr Pro Glu Gly Ile Lys Asp Arg Pro Gly Ala Tyr Ile Thr Cys |     |     |
|    | 450                                                             | 455 | 460 |
|    | Ala Pro Phe Asn Pro Ala Ala Thr Glu Ser Ile Ala Thr Asn Ile Phe |     |     |
|    | 465                                                             | 470 | 475 |
| 30 | Pro Leu Leu Gly Asp Pro Thr Ser Glu Asn Lys Thr Asp Glu Lys Lys |     |     |
|    | 485                                                             | 490 | 495 |
|    | Ile Glu Glu Lys Lys Ala Gln Ile Val Thr Glu Lys Asn Thr Ser Thr |     |     |
|    | 500                                                             | 505 | 510 |
|    | Lys Thr Ser Asn Pro Phe Leu Val Ala Ala Gln Asp Ser Glu Thr Asp |     |     |
| 35 | 515                                                             | 520 | 525 |
|    | Tyr Val Thr Thr Asp Asn Leu Thr Lys Val Thr Glu Glu Val Val Ala |     |     |
|    | 530                                                             | 535 | 540 |
|    | Asn Met Pro Glu Gly Leu Thr Pro Asp Leu Val Gln Glu Ala Cys Glu |     |     |
|    | 545                                                             | 550 | 555 |
| 40 | Ser Glu Leu Asn Glu Val Thr Gly Thr Lys Ile Ala Tyr Glu Thr Lys |     |     |
|    | 565                                                             | 570 | 575 |
|    | Met Asp Leu Val Gln Thr Ser Glu Val Met Gln Glu Ser Leu Tyr Pro |     |     |

|    |                                                                 |     |     |
|----|-----------------------------------------------------------------|-----|-----|
|    | 580                                                             | 585 | 590 |
|    | Ala Ala Gln Leu Cys Pro Ser Phe Glu Glu Ser Glu Ala Thr Pro Ser |     |     |
|    | 595                                                             | 600 | 605 |
|    | Pro Val Leu Pro Asp Ile Val Met Glu Ala Pro Leu Asn Ser Ala Val |     |     |
| 5  | 610                                                             | 615 | 620 |
|    | Pro Ser Ala Gly Ala Ser Val Ile Gln Pro Ser Ser Ser Pro Leu Glu |     |     |
|    | 625                                                             | 630 | 635 |
|    | Ala Ser Ser Val Asn Tyr Glu Ser Ile Lys His Glu Pro Glu Asn Pro |     |     |
|    | 645                                                             | 650 | 655 |
| 10 | Pro Pro Tyr Glu Glu Ala Met Ser Val Ser Leu Lys Lys Val Ser Gly |     |     |
|    | 660                                                             | 665 | 670 |
|    | Ile Lys Glu Glu Ile Lys Glu Pro Glu Asn Ile Asn Ala Ala Leu Gln |     |     |
|    | 675                                                             | 680 | 685 |
|    | Glu Thr Glu Ala Pro Tyr Ile Ser Ile Ala Cys Asp Leu Ile Lys Glu |     |     |
| 15 | 690                                                             | 695 | 700 |
|    | Thr Lys Leu Ser Ala Glu Pro Ala Pro Asp Phe Ser Asp Tyr Ser Glu |     |     |
|    | 705                                                             | 710 | 715 |
|    | Met Ala Lys Val Glu Gln Pro Val Pro Asp His Ser Glu Leu Val Glu |     |     |
|    | 725                                                             | 730 | 735 |
| 20 | Asp Ser Ser Pro Asp Ser Glu Pro Val Asp Leu Phe Ser Asp Asp Ser |     |     |
|    | 740                                                             | 745 | 750 |
|    | Ile Pro Asp Val Pro Gln Lys Gln Asp Glu Thr Val Met Leu Val Lys |     |     |
|    | 755                                                             | 760 | 765 |
|    | Glu Ser Leu Thr Glu Thr Ser Phe Glu Ser Met Ile Glu Tyr Glu Asn |     |     |
| 25 | 770                                                             | 775 | 780 |
|    | Lys Glu Lys Leu Ser Ala Leu Pro Pro Glu Gly Gly Lys Pro Tyr Leu |     |     |
|    | 785                                                             | 790 | 795 |
|    | Glu Ser Phe Lys Leu Ser Leu Asp Asn Thr Lys Asp Thr Leu Leu Pro |     |     |
|    | 805                                                             | 810 | 815 |
| 30 | Asp Glu Val Ser Thr Leu Ser Lys Lys Glu Lys Ile Pro Leu Gln Met |     |     |
|    | 820                                                             | 825 | 830 |
|    | Glu Glu Leu Ser Thr Ala Val Tyr Ser Asn Asp Asp Leu Phe Ile Ser |     |     |
|    | 835                                                             | 840 | 845 |
|    | Lys Glu Ala Gln Ile Arg Glu Thr Glu Thr Phe Ser Asp Ser Ser Pro |     |     |
| 35 | 850                                                             | 855 | 860 |
|    | Ile Glu Ile Ile Asp Glu Phe Pro Thr Leu Ile Ser Ser Lys Thr Asp |     |     |
|    | 865                                                             | 870 | 875 |
|    | Ser Phe Ser Lys Leu Ala Arg Glu Tyr Thr Asp Leu Glu Val Ser His |     |     |
|    | 885                                                             | 890 | 895 |
| 40 | Lys Ser Glu Ile Ala Asn Ala Pro Asp Gly Ala Gly Ser Leu Pro Cys |     |     |
|    | 900                                                             | 905 | 910 |
|    | Thr Glu Leu Pro His Asp Leu Ser Leu Lys Asn Ile Gln Pro Lys Val |     |     |

915                      920                      925  
 Glu Glu Lys Ile Ser Phe Ser Asp Asp Phe Ser Lys Asn Gly Ser Ala  
 930                      935                      940  
 Thr Ser Lys Val Leu Leu Leu Pro Pro Asp Val Ser Ala Leu Ala Thr  
 5 945                      950                      955                      960  
 Gln Ala Glu Ile Glu Ser Ile Val Lys Pro Lys Val Leu Val Lys Glu  
 965                      970                      975  
 Ala Glu Lys Lys Leu Pro Ser Asp Thr Glu Lys Glu Asp Arg Ser Pro  
 980                      985                      990  
 10 Ser Ala Ile Phe Ser Ala Glu Leu Ser Lys Thr Ser Val Val Asp Leu  
 995                      1000                      1005  
 Leu Tyr Trp Arg Asp Ile Lys Lys Thr Gly Val Val Phe Gly Ala Ser  
 1010                      1015                      1020  
 Leu Phe Leu Leu Leu Ser Leu Thr Val Phe Ser Ile Val Ser Val Thr  
 15 1025                      1030                      1035                      104  
 Ala Tyr Ile Ala Leu Ala Leu Leu Ser Val Thr Ile Ser Phe Arg Ile  
 1045                      1050                      1055  
 Tyr Lys Gly Val Ile Gln Ala Ile Gln Lys Ser Asp Glu Gly His Pro  
 1060                      1065                      1070  
 20 Phe Arg Ala Tyr Leu Glu Ser Glu Val Ala Ile Ser Glu Glu Leu Val  
 1075                      1080                      1085  
 Gln Lys Tyr Ser Asn Ser Ala Leu Gly His Val Asn Cys Thr Ile Lys  
 1090                      1095                      1100  
 Glu Leu Arg Arg Leu Phe Leu Val Asp Asp Leu Val Asp Ser Leu Lys  
 25 1105                      1110                      1115                      112  
 Phe Ala Val Leu Met Trp Val Phe Thr Tyr Val Gly Ala Leu Phe Asn  
 1125                      1130                      1135  
 Gly Leu Thr Leu Leu Ile Leu Ala Leu Ile Ser Leu Phe Ser Val Pro  
 1140                      1145                      1150  
 30 Val Ile Tyr Glu Arg His Gln Ala Gln Ile Asp His Tyr Leu Gly Leu  
 1155                      1160                      1165  
 Ala Asn Lys Asn Val Lys Asp Ala Met Ala Lys Ile Gln Ala Lys Ile  
 1170                      1175                      1180  
 Pro Gly Leu Lys Arg Lys Ala Glu  
 35 1185                      1190

&lt;210&gt; 5

&lt;211&gt; 1122

&lt;212&gt; DNA

40 &lt;213&gt; Homo sapiens

&lt;400&gt; 5

atggaagacc tggaccagtc tcctctggtc tcgtcctcgg acagcccacc ccggccgcag 60  
 cccgcgttca agtaccagtt cgtgagggag cccgaggacg aggaggaaga agaggaggag 120  
 gaagaggagg acgaggacga agacctggag gagctggagg tgctggagag gaagcccgcc 180  
 gccgggctgt ccgcggcccc agtgcccacc gcccctgccg ccggcgcgcc cctgatggac 240  
 5 ttcggaatatg acttcgtgcc gccggcgccc cggggacccc tgccggccgc tcccccgtc 300  
 gccccggagc ggcagccgtc ttgggacccg agcccgggtg cgtcgaccgt gcccgcgcca 360  
 tccccgctgt ctgctgcgcg agtctcgccc tccaagctcc ctgaggacga cgagcctccg 420  
 gcccggcctc cccctcctcc cccggccagc gtgagccccc aggcagagcc cgtgtggacc 480  
 ccgccagccc cggctcccg cgcgcccccc tccaccccg cgcgcccga ggcaggggc 540  
 10 tcctcgggct cagtgggtgt tgacctcctg tactggagag acattaagaa gactggagtg 600  
 gtgtttggtg ccagcctatt cctgctgctt tcattgacag tattcagcat tgtgagcgta 660  
 acagcctaca ttgccttggc cctgctctct gtgaccatca gctttaggat atacaagggc 720  
 gtgatccaag ctatccagaa atcagatgaa ggccacccat tcagggcata tctggaatct 780  
 gaagtgtcta tatctgagga gttggttcag aagtacagta attctgctct tggcatgtg 840  
 15 aactgcacga taaaggaact caggcgctc ttcttagttg atgatttagt tgattctctg 900  
 aagtttgagc tgttgatgtg ggtatttacc tatgttggtg ccttgtttaa tggctgaca 960  
 ctactgattt tggctctcat ttcactcttc agtgttcctg ttatttatga acggcatcag 1020  
 gcacagatag atcattatct aggacttgca aataagaatg ttaaagatgc tatggctaaa 1080  
 atccaagcaa aaatccctgg attgaagcgc aaagctgaat ga 1122

20

&lt;210&gt; 6

&lt;211&gt; 373

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

25

&lt;400&gt; 6

Met Glu Asp Leu Asp Gln Ser Pro Leu Val Ser Ser Ser Asp Ser Pro  
 1 5 10 15  
 Pro Arg Pro Gln Pro Ala Phe Lys Tyr Gln Phe Val Arg Glu Pro Glu  
 20 25 30  
 Asp Glu Glu Glu Glu Glu Glu Glu Glu Asp Glu Asp Glu Asp  
 35 40 45  
 Leu Glu Glu Leu Glu Val Leu Glu Arg Lys Pro Ala Ala Gly Leu Ser  
 50 55 60  
 35 Ala Ala Pro Val Pro Thr Ala Pro Ala Ala Gly Ala Pro Leu Met Asp  
 65 70 75 80  
 Phe Gly Asn Asp Phe Val Pro Pro Ala Pro Arg Gly Pro Leu Pro Ala  
 85 90 95  
 Ala Pro Pro Val Ala Pro Glu Arg Gln Pro Ser Trp Asp Pro Ser Pro  
 100 105 110  
 40 Val Ser Ser Thr Val Pro Ala Pro Ser Pro Leu Ser Ala Ala Val  
 115 120 125

Ser Pro Ser Lys Leu Pro Glu Asp Asp Glu Pro Pro Ala Arg Pro Pro  
 130 135 140  
 Pro Pro Pro Pro Ala Ser Val Ser Pro Gln Ala Glu Pro Val Trp Thr  
 145 150 155 160  
 5 Pro Pro Ala Pro Ala Pro Ala Ala Pro Pro Ser Thr Pro Ala Ala Pro  
 165 170 175  
 Lys Arg Arg Gly Ser Ser Gly Ser Val Val Val Asp Leu Leu Tyr Trp  
 180 185 190  
 Arg Asp Ile Lys Lys Thr Gly Val Val Phe Gly Ala Ser Leu Phe Leu  
 10 195 200 205  
 Leu Leu Ser Leu Thr Val Phe Ser Ile Val Ser Val Thr Ala Tyr Ile  
 210 215 220  
 Ala Leu Ala Leu Leu Ser Val Thr Ile Ser Phe Arg Ile Tyr Lys Gly  
 225 230 235 240  
 15 Val Ile Gln Ala Ile Gln Lys Ser Asp Glu Gly His Pro Phe Arg Ala  
 245 250 255  
 Tyr Leu Glu Ser Glu Val Ala Ile Ser Glu Glu Leu Val Gln Lys Tyr  
 260 265 270  
 Ser Asn Ser Ala Leu Gly His Val Asn Cys Thr Ile Lys Glu Leu Arg  
 20 275 280 285  
 Arg Leu Phe Leu Val Asp Asp Leu Val Asp Ser Leu Lys Phe Ala Val  
 290 295 300  
 Leu Met Trp Val Phe Thr Tyr Val Gly Ala Leu Phe Asn Gly Leu Thr  
 305 310 315 320  
 25 Leu Leu Ile Leu Ala Leu Ile Ser Leu Phe Ser Val Pro Val Ile Tyr  
 325 330 335  
 Glu Arg His Gln Ala Gln Ile Asp His Tyr Leu Gly Leu Ala Asn Lys  
 340 345 350  
 Asn Val Lys Asp Ala Met Ala Lys Ile Gln Ala Lys Ile Pro Gly Leu  
 30 355 360 365  
 Lys Arg Lys Ala Glu  
 370

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/04345

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 C07K16/18

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, STRAND

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category ° | Citation of document, with indication, where appropriate, of the relevant passages                                                                                                                                                                                    | Relevant to claim No. |
|------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| X          | <p>WO 98 06841 A (INCYTE PHARMACEUTICALS, INC.) 19 February 1998 (1998-02-19)<br/> abstract<br/> page 1, line 1 -page 2, line 41<br/> page 5, line 7 -page 28, line 8<br/> SEQ ID NOS: 1 and 2<br/> page 38 -page 39<br/> page 46; claims 1-14<br/> ---<br/> -/--</p> | 1-10                  |



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

° Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

14 March 2001

Date of mailing of the international search report

30/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fuchs, U

# INTERNATIONAL SEARCH REPORT

Intern      1al Application No

PCT/GB 00/04345

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category ° | Citation of document, with indication, where appropriate, of the relevant passages                                                                                                                                                                                                                                                                                                                                                                                                                        | Relevant to claim No. |
|------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| X          | <p>WO 98 56804 A (HUMAN GENOME SCIENCES, INC.) 17 December 1998 (1998-12-17)<br/> abstract<br/> page 2, line 1 - line 8<br/> gene no.: 69<br/> page 62, line 1 - line 32<br/> page 82; table 1<br/> page 85, line 1 -page 105, line 7<br/> page 111, line 6 -page 112, line 16<br/> page 113, line 1 -page 120, line 8<br/> SEQ ID NO: 79<br/> page 235 -page 236<br/> SEQ ID NO: 301<br/> page 354 -page 355<br/> page 370 -page 373; claims 1-23<br/> ---</p>                                           | 1-10                  |
| X          | <p>EMBL database, Heidelberg, FRG<br/> Emhum1 accession number AF077050<br/> 27 April 1999<br/> SONG, H. ET AL.: "Homo sapiens<br/> neuroendocrine-specific protein C homolog<br/> mRNA, complete cds."<br/> XP002162897<br/> the whole document<br/> ---&amp; EMBL database, Heidelberg, FRG<br/> Trembl accession number Q9Y293<br/> 1 November 1999<br/> SONG, H. ET AL.: "Human<br/> neuroendocrine-specific protein C (NSP)<br/> homolg gene."<br/> XP002162898<br/> the whole document<br/> ---</p> | 1-8                   |
| X          | <p>---&amp; EMBL database, Heidelberg, FRG<br/> Trembl accession number Q9Y293<br/> 1 November 1999<br/> SONG, H. ET AL.: "Human<br/> neuroendocrine-specific protein C (NSP)<br/> homolg gene."<br/> XP002162898<br/> the whole document<br/> ---</p>                                                                                                                                                                                                                                                    | 1-8                   |
| A          | <p>SPILLMANN, A.A. ET AL: "Identification<br/> and Characterization of a Bovine Neurite<br/> Growth Inhibitor (bNI-220)"<br/> JOURNAL OF BIOLOGICAL CHEMISTRY,<br/> vol. 273, no. 30,<br/> 24 July 1998 (1998-07-24), pages<br/> 19283-19293, XP002162896<br/> cited in the application<br/> see especially<br/> * page 19288, line 40 - line 56 *<br/> * page 19290; table 2 *<br/> the whole document<br/> ---</p>                                                                                      | 1-10                  |
|            | -/--                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |                       |

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/04345

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages                                                                                                                                                                                                                                                                    | Relevant to claim No. |
|------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| A          | ROEBROEK, A.J. ET AL.: "Cloning and Expression of Alternative Transcripts of a Novel Neuroendocrine-specific Gene and Identification of Its 135-kDa Translational Product"<br>JOURNAL OF BIOLOGICAL CHEMISTRY,<br>vol. 268, no. 18,<br>25 June 1993 (1993-06-25), pages<br>13439-13447, XP002045108<br>cited in the application<br>the whole document | 1-10                  |
| P,X        | PRINJHA, R. ET AL.: "Inhibitor of neurite outgrowth in humans"<br>NATURE,<br>vol. 403, no. 6768,<br>27 January 2000 (2000-01-27), pages<br>383-384, XP002144397<br>the whole document                                                                                                                                                                 | 1-8                   |
| P,X        | -& EMBL database, Heidelberg, FRG<br>Emhum4 accession number AJ251385<br>22 July 2000<br>MICHALOVICH, D.: "Homo sapiens mRNA for Nogo-C protein (Nogo gene)"<br>XP002162899<br>the whole document                                                                                                                                                     | 1-8                   |

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 00/04345

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s)                                                                                                                   | Publication<br>date                                                                                                        |
|-------------------------------------------|---------------------|----------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|
| WO 9806841 A                              | 19-02-1998          | US 5858708 A<br>AU 3902297 A<br>EP 0918856 A<br>JP 2000517173 T                                                                              | 12-01-1999<br>06-03-1998<br>02-06-1999<br>26-12-2000                                                                       |
| WO 9856804 A                              | 17-12-1998          | AU 8066798 A<br>EP 1042346 A<br>AU 6241698 A<br>EP 0988385 A<br>US 6046031 A<br>WO 9831818 A<br>AU 8474398 A<br>EP 1000084 A<br>WO 9902546 A | 30-12-1998<br>11-10-2000<br>07-08-1998<br>29-03-2000<br>04-04-2000<br>23-07-1998<br>08-02-1999<br>17-05-2000<br>21-01-1999 |