

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
22 January 2009 (22.01.2009)

PCT

(10) International Publication Number  
WO 2009/012467 A2

## (51) International Patent Classification:

*C12N 15/11* (2006.01) *A01H 5/00* (2006.01)  
*C12N 15/82* (2006.01) *A01H 1/00* (2006.01)

(74) Agents: **BASTIAN, Kevin L.** et al.; Townsend and Townsend and Crew LLP, Two Embarcadero Center, 8th Floor, San Francisco, California 94111 (US).

## (21) International Application Number:

PCT/US2008/070507

(22) International Filing Date: 18 July 2008 (18.07.2008)

(25) Filing Language: English

(26) Publication Language: English


(30) Priority Data:  
60/950,853 19 July 2007 (19.07.2007) US(71) Applicant (for all designated States except US): **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, 12th Floor, Oakland, California 94607 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL,

[Continued on next page]

## (54) Title: INCREASING GRAIN YIELD THROUGH TARGETED REDUCTION IN ETHYLENE SIGNALING



(57) Abstract: The present invention is directed to plant genetic engineering. In particular, it is directed to producing green leaves and increasing productivity through inhibition of ethylene. The compositions and methods of the invention involve dominant negative ethylene receptors that interfere with ethylene signaling.

WO 2009/012467 A2

Figure 1



NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— *without international search report and to be republished upon receipt of that report*

## INCREASING GRAIN YIELD THROUGH TARGETED REDUCTION IN ETHYLENE SIGNALING

### BACKGROUND OF THE INVENTION

5 [0001] Ethylene is known to be a regulator of programmed cell death (PCD) during plant development (Campbell and Drew, *Planta* 157:350-357 (1983); Drew *et al.*, *Planta* 147:83-88 (1979); He *et al.*, *Plant Physiol.* 112:1679-1685 (1996)) and plays a role in orchestrating PCD in developing cereal endosperm: exogenous ethylene can accelerate the onset of the cell death program in developing endosperm whereas inhibitors of ethylene biosynthesis or 10 perception delay the program (Young *et al.*, *Plant Physiol.* 119:737-751 (1997); Young and Gallie, *Plant Mol. Biol.* 39:915-926 (1999); Young and Gallie, *Plant Mol. Biol.* 42:397-414 (2000)). Ethylene controls many aspects of plant growth and development such as fruit development, root and leaf growth and seed germination.

15 [0002] Ethylene perception involves membrane-localized receptors that, in *Arabidopsis*, include ETR1, ERS1, ETR2, ERS2 and EIN4 (Chang *et al.*, *Science* 262:539-544 (1993); Hua *et al.*, *Science* 269:1712-1714 (1995), Hua *et al.*, *Plant Cell* 10:1321-1332 (1998), Sakai *et al.*, *Proc. Natl. Acad. Sci. USA* 95:5812-5817 (1998)). ETR1, ETR2 and EIN4 are composed of three domains, an N-terminal transmembrane ethylene binding domain (Schaller and Bleeker, *Science* 270:1809-1811 (1995)), a histidine protein kinase domain, and a C-terminal 20 receiver domain. ERS1 and ERS2 lack the receiver domain. These genes have been grouped into two subfamilies based on homology, where ETR1 and ERS1 comprise one subfamily and ETR2, ERS2, and EIN4 comprise the other (Hua *et al.*, *Plant Cell* 10:1321-1332 (1998)). ETR1 and ERS1 both contain a functional histidine kinase domain and autophosphorylate histidine residues, whereas other family members lack particular residues required for 25 histidine kinase activity. However, *in vitro* studies indicate that ETR2, ERS2, and EIN4 are capable of autophosphorylation on serine and threonine residues (Moussatche and Klee (2004), *J. Biol. Chem.*, 279:48734).

30 [0003] Only two ethylene receptor genes have been identified in maize (*i.e.*, ZmETR2 and ZmERS1) in contrast to the five types identified in *Arabidopsis* (Gallie and Young (2004) *Mol Genet Genomics* 271: 267-281). ZmETR2 has two variants, named ZmETR9 and ZmETR40. Members of both ethylene receptor families are expressed to substantially higher

levels in the developing embryo relative to the endosperm. This explains why, despite the fact that the endosperm and embryo each contribute to the synthesis of ethylene, the developing endosperm exhibits a low threshold to ethylene-induced cell death while the embryo is protected.

5 [0004] The endosperm of cereals serves as the major storage organ for grain but undergoes cell death during mid to late seed development. Ethylene regulates the timing of the onset of cell death in the developing endosperm. Because ethylene is a gas that can pass freely through membranes, all organs of the developing kernel might be expected to be exposed to ethylene generated by a specific organ and diluted only by their distance from the generating 10 source. The ability to limit cell death to specific organs within the developing kernel suggests tight control of the expression of the ethylene biosynthetic and perception machinery.

15 [0005] The role of ethylene in photosynthesis is unclear. Photosynthesis is often inhibited during conditions of stress, e.g., drought, ozone exposure, or chilling (Flexas and Medrano (2002), *Annals Bot.* 89: 183-189; Chaves *et al.* (2002), *Annals Bot.* 89: 907-916; Ramachandra *et al.* (2004), *J Plant Physiol.* 161: 1189-1202). Photosynthetic capacity increases during leaf expansion and declines with leaf age until it reaches low levels prior to the onset of leaf senescence (Gay and Thomas (1995), *New Phytol.* 130: 159-168). The rate of initiation and execution of a senescence program significantly impacts the ultimate 20 contribution that a leaf can make to a plant. This is of particular relevance to those crops, such as cereals, where yield potential is reduced by adverse environmental conditions that induce premature leaf senescence.

25 [0006] The effect of ethylene on photosynthesis has been controversial with reports suggesting either no effect or an inhibitory effect (Pallaghy and Raschke (1972), *Plant Physiol.* 49: 275-276; Kays and Pallas (1980), *Nature* 285: 51-52; Pallas and Kays (1982), *Plant Physiol.* 70: 598-601; Squier *et al.* (1985), *Environ Sci Technol* 19: 432-437; Taylor and Gunderson (1986), *Merr. Plant Physiol.* 86: 85-92; Woodrow *et al.* (1988), *Mill. J Exp Bot.* 39: 667-684). The difference in species, growth conditions, intact versus excised leaves, and concentration of exogenous ethylene used may have contributed to the variation in 30 observed effects. A mutant approach was employed to examine the effect of ethylene on photosynthetic activity and grain development in maize (Young *et al.* (2004), *Plant J* 40: 813-825). The authors in this study found that maize mutants with defective ethylene

production had increased amounts of chlorophyll and rates of CO<sub>2</sub> assimilation relative to wild type plants.

[0007] Because ethylene plays such a large role in plant growth and development, the identification of genes involved in the ethylene response pathway is useful for creating plants 5 with phenotypes associated with an altered ethylene-related process, such as plants having staygreen traits. Accordingly, a need exists for the identification of genes involved in cereal ethylene signal transduction pathways.

### BRIEF SUMMARY OF THE INVENTION

10 [0008] The present invention provides compositions and methods to affect the ethylene signal transduction pathway in plants.

[0009] The invention provides for an isolated nucleic acid which can encode a dominant negative ethylene receptor polypeptide, for example, a dominant negative ZmERS1, ZmETR9, or ZmETR40 (represented by SEQ ID NOs:2, 4 and 6, respectively). In addition, 15 the invention includes the polynucleotide sequences encoding each of these polypeptides (e.g., SEQ ID NOs:1, 3 and 5). The polynucleotides of the invention include sequences having at least 90%, and more typically at least 95%, 99%, or 100% identity to SEQ ID NOs:1, 3, and 5. In some embodiments of the invention, the isolated nucleic acid encodes a polypeptide with at least 90%, more typically 95%, 99% or 100% identity to SEQ ID NO:2, 20 wherein amino acid 65 is not a cysteine. In some embodiments, amino acid 65 of SEQ ID NO:2 is a tyrosine. In some embodiments, the isolated nucleic acid encodes a polypeptide with at least 90%, more typically 95%, 99%, or 100% identity to SEQ ID NO:4, wherein amino acid 102 is not a cysteine. In some embodiments, amino acid 102 of SEQ ID NO:4 is a tyrosine. In some embodiments, the isolated nucleic acid encodes a polypeptide with at 25 least 90%, more typically 95%, 99%, or 100% identity to SEQ ID NO:6, wherein amino acid 102 is not a cysteine. In some embodiments, amino acid 102 of SEQ ID NO:6 is a tyrosine.

[0010] In some embodiments, the nucleic acids encode truncation mutants of ZmERS1 and ZmETR2 (for example, comprising amino acids 1-110 of SEQ ID NO:2 or amino acids 1-147 SEQ ID NOs:4 and 6). In such embodiments, the truncation mutant may be slightly longer, 30 e.g., 1-150, 1-200, 1-250, 1-300, 1-325, or 1-350 of SEQ ID NO:2 or amino acids 1-200, 1-250, 1-300, 1-350, or 1-375 of SEQ ID NO:4 or 6.

[0011] In some embodiments, the invention provides a recombinant expression cassette comprising a promoter sequence operably linked to a nucleic acid sequence encoding a dominant negative ethylene receptor polypeptide, for example, a dominant negative ZmERS1, ZmETR9, or ZmETR40 (represented by SEQ ID NOs:2, 4, and 6, respectively). In 5 some embodiments, the nucleic acids encode truncation mutants of ZmERS1 and ZmETR2 (for example, comprising amino acids 1-110 of SEQ ID NO:2 or amino acids 1-147 SEQ ID NO:4 or 6). In addition, the invention includes the polynucleotide sequences of SEQ ID NOs:1, 3, and 5, and nucleic acid sequences having at least 90%, and more typically at least 95%, 99%, or 100% identity to SEQ ID NOs:1, 3, and 5.

10 [0012] In some embodiments, the invention provides for a transgenic plant comprising a recombinant expression cassette comprising a promoter sequence operably linked to a nucleic acid sequence encoding a dominant negative ethylene receptor polypeptide of the invention. In some embodiments, the nucleic acids encode ZmERS1, ZmETR9, or ZmETR40 (represented by SEQ ID NOs:2, 4, and 6, respectively). In other embodiments, the dominant 15 negative ethylene receptor polypeptide comprises amino acids 1-110 of SEQ ID NO:2 or amino acids 1-147 SEQ ID NO:4 or 6. In some embodiments, the nucleic acid sequence is SEQ ID NO:1, 3, or 5, and nucleic acid sequences at least 90%, and more typically at least 95%, 99%, or 100% identical to SEQ ID NOs:1, 3, and 5. In some embodiments, the transgenic plant is a cereal plant, such as wheat, rice, barley, rye, millet, sorghum, or oat. In 20 some embodiments, the transgenic plant is maize.

[0013] In further embodiments, the invention provides a method of reducing ethylene sensitivity in a plant comprising the steps of (a) introducing a construct comprising a promoter operably linked to a nucleic acid sequence encoding a dominant negative ethylene receptor polypeptide and (b) selecting a plant with reduced ethylene sensitivity. Preferred 25 dominant negative ethylene receptor polypeptides include ZmERS1, ZmETR9, or ZmETR40 (represented by SEQ ID NOs:2, 4, and 6, respectively). In other embodiments, the dominant negative ethylene receptor polypeptide comprises amino acids 1-110 of SEQ ID NO:2 or amino acids 1-147 SEQ ID NO:4 or 6. In some embodiments, the nucleic acid sequences are SEQ ID NOs:1, 3, or 5, and nucleic acid sequences at least 90%, and more typically at least 30 95%, 99%, or 100% identical to SEQ ID NOs:1, 3, or 5.

[0014] In another aspect, the invention provides a method of producing a staygreen phenotype in a plant. The method comprises the steps of (a) introducing a construct

comprising a promoter operably linked to a nucleic acid sequence encoding a dominant negative ethylene receptor polypeptide and (b) selecting a plant with staygreen characteristics. In some embodiments, the ethylene receptor polypeptides include ZmERS1, ZmETR9, or ZmETR40 (represented by SEQ ID NOs:2, 4, and 6, respectively). In other 5 embodiments, the dominant negative ethylene receptor polypeptide comprises amino acids 1-110 of SEQ ID NO:2 or amino acids 1-147 SEQ ID NO:4 or 6. In some embodiments, the nucleic acid sequence is SEQ ID NOs:1, 3, or 5, and nucleic acid sequences at least 90%, and more typically at least 95%, 99%, or 100% identical to SEQ ID NOs:1, 3, or 5. In some 10 embodiments, the plant is selected for delayed senescence, increased photosynthetic capacity or for multiple embryos in a single seed.

**[0015]** Other objects, advantages and embodiments of the invention will be apparent from review of the Detailed Description that follows.

#### BRIEF DESCRIPTION OF THE DRAWINGS

15 **[0016]** Figure 1: Quantitation of the ethylene insensitivity conferred by the expression of mutant ZmERS1 or ZmETR2 in *Arabidopsis* over a range of ACC concentrations. Seeds expressing mutant ZmERS1 (MS1-11) or ZmETR2 (MT2-9) were germinated on ACC at the concentrations indicated and grown in the dark for 5 days. Wild type seeds (WT) were included as a control.

20

#### DETAILED DESCRIPTION OF THE INVENTION

##### A. INTRODUCTION

25 **[0017]** The present invention provides new methods of delaying senescence in a maize plant by inhibiting ethylene responses in the plant. The delay in senescence can be achieved by mutation of ethylene receptor proteins, as well as overexpression of wild-type or mutant ethylene receptor proteins. The present invention also provides methods for selecting for a cereal plant with a delayed senescence pattern or characteristic. A delayed senescence pattern will result in a cereal plant with an altered phenotype as compared to a wild type plant. An altered phenotype includes, but is not limited to, staygreen traits, *e.g.*, leaves that 30 remain green late in the growing season, increased photosynthetic capacity, improved drought tolerance, improved silage, increased grain yield, and increased tolerance to planting

at higher densities, and kernels with multiple embryos. Accordingly, a plant with increased biomass and/or yield can be identified by inhibiting ethylene response, through the production of mutated ethylene receptor proteins alone or in combination with other methods.

## B. DEFINITIONS

5 [0018] The phrase "nucleic acid" or "polynucleotide sequence" refers to a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Nucleic acids may also include modified nucleotides that permit correct read through by a polymerase and do not alter the expression of a polypeptide encoded by that nucleic acid.

10 [0019] The phrase "nucleic acid sequence encoding" refers to a nucleic acid which directs the expression of a specific protein or peptide. The nucleic acid sequences include both the DNA strand sequence that is transcribed into RNA and the RNA sequence that is translated into protein. The nucleic acid sequences include both the full length nucleic acid sequences as well as non-full length sequences derived from the full length sequences. It should be further understood that the sequence includes the degenerate codons of the native sequence or 15 sequences which may be introduced to provide codon preference in a specific host cell.

20 [0020] The term "promoter" refers to a region or sequence determinants located upstream or downstream from the start of transcription and which are involved in recognition and binding of RNA polymerase and other proteins to initiate transcription. A "plant promoter" is a promoter capable of initiating transcription in plant cells. Such promoters need not be of plant origin, for example, promoters derived from plant viruses, such as the CaMV35S promoter, can be used in the present invention.

25 [0021] The term "plant" includes whole plants, shoot vegetative organs/structures (*e.g.* leaves, stems and tubers), roots, flowers and floral organs/structures (*e.g.* bracts, sepals, petals, stamens, carpels, anthers and ovules), seeds (including embryo, endosperm, and seed coat) and fruit (the mature ovary), plant tissue (*e.g.* vascular tissue, ground tissue, and the like) and cells (*e.g.* guard cells, egg cells, trichomes and the like), and progeny of same. The class of plants that can be used in the method of the invention is generally as broad as the class of higher and lower plants amenable to transformation techniques, including angiosperms (monocotyledonous and dicotyledonous plants), gymnosperms, ferns, 30 bryophytes, and multicellular algae. It includes plants of a variety of ploidy levels, including aneuploid, polyploid, diploid, haploid and hemizygous.

**[0022]** The phrase "host cell" refers to a cell from any organism. Preferred host cells are derived from plants, bacteria, yeast, fungi, insects or other animals. Methods for introducing polynucleotide sequences into various types of host cells are well known in the art.

5 **[0023]** A polynucleotide sequence is "heterologous to" a second polynucleotide sequence if it originates from a foreign species, or, if from the same species, is modified by human action from its original form. For example, a promoter operably linked to a heterologous coding sequence refers to a coding sequence from a species different from that from which the promoter was derived, or, if from the same species, a coding sequence which is different from any naturally occurring allelic variants.

10 **[0024]** A polynucleotide "exogenous to" an individual plant is a polynucleotide which is introduced into the plant, or a predecessor generation of the plant, by any means other than by a sexual cross. Examples of means by which this can be accomplished are described below, and include Agrobacterium-mediated transformation, biolistic methods, electroporation, in planta techniques, and the like.

15 **[0025]** "Increased or enhanced expression or activity" of a particular polypeptide or nucleic acid of the invention refers to an augmented change in activity of the polypeptide. Examples of such increased activity or expression include the following: Activity of the polypeptide or expression of the gene encoding the polypeptide is increased above the level (or is present for a longer period of time) of that in wild-type, non-transgenic control plants. Activity of a 20 polypeptide or expression of a gene is present in an organ, tissue or cell where it is not normally detected in wild-type, non-transgenic control plants (*i.e.* spatial distribution of a polypeptide or expression of the gene encoding the polypeptide is altered).

25 **[0026]** "Decreased expression or activity" of a polypeptide or nucleic acid of the invention refers to a decrease in activity of the polypeptide. Examples of such decreased activity or expression include the following: Activity of the polypeptide or expression of the gene is decreased below the level of that in a wild-type, non-transgenic control plant.

**[0027]** The term "reproductive structures" or "reproductive tissues" as used herein includes fruit, ovules, seeds, pollen, flowers, or flower parts such as pistils, stamens, anthers, sepals, petals, carpels, or any embryonic tissue.

30 **[0028]** The term "vegetative structures" or "vegetative tissues" as used herein includes leaves, stems, tubers, roots, vascular tissue, or root and shoot meristem.

**[0029]** An "expression cassette" refers to a nucleic acid construct, which when introduced into a host cell, results in transcription and/or translation of a RNA or polypeptide, respectively. Antisense or sense constructs that are not or cannot be translated are expressly included by this definition.

5    **[0030]** In the case of both expression of transgenes and inhibition of endogenous genes (e.g., by antisense, or sense suppression) one of skill will recognize that the inserted polynucleotide sequence need not be identical and may be "substantially identical" to a sequence of the gene from which it was derived. As explained below, these variants are specifically covered by this term.

10   **[0031]** In the case where the inserted polynucleotide sequence is transcribed and translated to produce a functional polypeptide, one of skill will recognize that because of codon degeneracy, a number of polynucleotide sequences will encode the same polypeptide. These variants are specifically covered by the term "polynucleotide sequence from a gene of the invention". In addition, the term specifically includes sequences (e.g., full length sequences) 15 substantially identical (determined as described below) with a gene sequence encoding a polypeptide of the invention, and that encode proteins that retain the function of a polypeptide of the invention.

20   **[0032]** In the case of polynucleotides used to inhibit expression of an endogenous gene, the introduced sequence need not be perfectly identical to a sequence of the target endogenous gene. The introduced polynucleotide sequence will typically be at least substantially 25 identical (as determined below) to the target endogenous sequence.

**[0033]** Two nucleic acid sequences or polypeptides are said to be "identical" if the sequence of nucleotides or amino acid residues, respectively, in the two sequences is the same when aligned for maximum correspondence as described below. The term "complementary to" is used herein to mean that the sequence is complementary to all or a portion of a reference polynucleotide sequence.

**[0034]** Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman, *Add. APL. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48:443 (1970), by 30 the search for similarity method of Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT,

BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

[0035] "Percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

[0036] The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 85% sequence identity to a dominant negative ethylene receptor polynucleotide (*e.g.*, SEQ ID NO:1, 3, or 5). Alternatively, percent identity can be any integer from 85% to 100%. Most embodiments include at least: 85%, 90%, 95%, or 99% compared to a reference sequence using the programs described herein; preferably BLAST using standard parameters, as described below. Accordingly, sequences encoding a polypeptide used in the methods of the present invention include nucleic acid sequences that have substantial identity to the sequences disclosed here. One of skill will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like.

[0037] Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 90% to a dominant negative ethylene receptor polypeptide (*e.g.*, SEQ ID NO:2, 4, or 6). Percent identity of polypeptides can be any integer from 90% to 100%, for example, 90%, 95%, or 99%. Polypeptides that are "substantially similar" share sequences as noted above except that residue positions which are not identical may differ by conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of

amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Conservative amino acids substitution groups include: valine-leucine-isoleucine, phenylalanine-tyrosine, 5 lysine-arginine, alanine-valine, aspartic acid-glutamic acid, and asparagine-glutamine.

[0038] Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other, or a third nucleic acid, under stringent conditions. Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting 10 point (Tm) for the specific sequence at a defined ionic strength and pH. The Tm is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is about 0.02 molar at pH 7 and the temperature is at least about 60°C or 65°C.

15 [0039] For the purposes of this disclosure, stringent conditions for hybridizations are those which include at least one wash in 0.2X SSC at 63°C for 20 minutes, or equivalent conditions. Moderately stringent conditions include at least one wash (usually 2) in 0.2X SSC at a temperature of at least about 50°C, usually about 55°C, for 20 minutes, or equivalent conditions.

20 [0040] The phrase "phenotype associated with an ethylene-related process" refers to a phenotype that is modulated by ethylene. Exemplary phenotypes include, but are not limited to, staygreen traits, such as improved drought tolerance, improved silage, leaves that stay green later in the season, and increased tolerance to planting at higher densities. Modulation of ethylene-related processes can result from, *e.g.*, overproduction of ethylene, 25 underproduction of ethylene, increased sensitivity to ethylene in a cell or decreased sensitivity to ethylene in a cell.

[0041] The term "ethylene receptor" or "ethylene receptor protein," refers to ethylene receptors from any of the ethylene receptor families present in a plant. By way of example, in *Arabidopsis*, ethylene receptor proteins include ETR1, ERS1, ETR2, ERS2, and EIN4. 30 *Zea mays* ethylene receptor proteins include ZmERS1 and ZmETR2 (including the ZmETR2 variants ZmETR9 and ZmETR40). As used herein, "ETR2", "ZmETR2", "ZmETR9" and

"ZmETR40" can be used synonymously. Similarly, "ERS1" and "ZmERS1" can be used synonymously.

[0042] Ethylene receptors of the invention may be isolated from any species of plant, and include species homologs of the exemplary ethylene receptors. Ethylene receptors also 5 include proteins with naturally-occurring and induced mutations, including insertion, deletion, and point mutations.

[0043] "Dominant negative" refers to a gene product that adversely affects the function of the normal, wild-type gene product within the same cell even when the cell is heterozygous (wild-type and dominant negative). Expression of the dominant negative mutant generally 10 results in a decrease in the normal function. A "dominant negative ethylene receptor" may be formed by preserving the dimerization domain, but disrupting or truncating other functional domains of the protein. For example, an ethylene receptor mutant that does not bind ethylene (called a "non-ethylene binding ethylene receptor" herein) acts in a dominant negative fashion. The non-ethylene binding ethylene receptor of the invention include the 15 polypeptides of SEQ ID NOs:2, 4, and 6, and substantially identical variants thereof, so long as residue 65 of SEQ ID NO:2 or residue 102 of SEQ ID NOs:4 or 6 is not a cysteine.

[0044] Ethylene receptor polypeptides dimerize via a disulfide link at the N terminus of the polypeptides (see, e.g., Schaller *et al.* (1995) *J. Biol. Chem.* 270:12526-30). For the purposes of an ethylene receptor, the "dimerization domain" refers to at least one of the N terminal 20 cysteine residues that mediate this dimerization. The cysteines involved in dimerization are distinct from the cysteine residue(s) that mediate ethylene binding. The dimerization domain in ZmERS1 is found at amino acids 4 and 6 of SEQ ID NO:2. The dimerization domains of ZmETR9 and ZmETR40 are located at residues 37 and 40 of SEQ ID NO:4, and amino acids 38 and 40 of SEQ ID NO:6, respectively.

[0045] The term "ethylene binding domain" roughly corresponds to the transmembrane 25 domain, found at the N terminus of the ethylene receptor polypeptides. The residues involved in ethylene binding in ZmERS1 span amino acids 20-119 of SEQ ID NO:2. The residues involved in ethylene binding in ZmETR2 span amino acids 53-156 of SEQ ID NOs:4 and 6. The ethylene binding ability of the receptor can be tested using the methods 30 described herein.

[0046] The term "transmembrane domain" or "intermembrane region" of an ethylene receptor refers to region spanning the three or four membrane-spanning helices of the

ethylene receptor polypeptide. Like most transmembrane domains, the ethylene receptor transmembrane domain is largely hydrophobic. In most ethylene receptor polypeptides, the transmembrane domain roughly corresponds to the "ethylene binding domain." By way of example, the ZmERS1 transmembrane domain spans approximately amino acids 20-107 of SEQ ID NO:2. The transmembrane domain of ZmETR2 spans approximately 5-144 of SEQ ID NOs:4 and 6.

[0047] The "histidine kinase domain" or "kinase domain" of an ethylene receptor refers to a region that is involved in signal transduction pathways. ETR1 and ERS1 receptors autophosphorylate at a conserved histidine residue in the region in response to ethylene. The other ethylene receptor family proteins (ETR2, ERS2, and EIN4) lack residues that are required for histidine kinase activity. However, these proteins are still capable of conferring ethylene responsiveness on a plant cell.

[0048] The "C terminal receiver domain" is found only in the ETR1, ETR2 and EIN4 ethylene receptors. The receiver domain plays a role in regulating responses in combination with the kinase domain. The kinase domain and receiver domains interact with CTR1 to modulate ethylene responsiveness. By way of example, the C terminal receiver domain in ETR2 spans approximately amino acids 612-767 of SEQ ID NOs:4 and 6.

[0049] The term "staygreen" refers to the ability of a hybrid plant to maintain plant health later into the growing season as compared to a wild type plant. Staygreen traits have been associated with increased grain yield, increased photosynthetic capacity, improved drought tolerance, improved silage and an increase in tolerance to planting at higher densities. The staygreen phenotype can be conveniently assayed using standard assays. For example, dark-induced senescence assays can be used. Such assays typically involve sheathing leaves while still attached to the plant for one week. The lack of light induces leaf senescence, which can be delayed, as compared to controls, in plants of the invention.

[0050] The term "photosynthetic capacity" refers to the ability of a plant to convert light energy into chemical energy. Photosynthetic capacity may be determined, for example, by measuring the amount of chlorophyll in a plant structure or the amount of CO<sub>2</sub> assimilation. Such techniques are known in the art.

[0051] A "biomarker" is any detectable marker that indicates the presence of a trait or gene of interest. In some cases, a biomarker is a readily detectable phenotype that is linked to or cosegregates with the trait or gene of interest. For example, some biomarkers confer

resistance to chemical or nutrient stress. In some embodiments, the biomarker is a readily detectable protein, *e.g.*, a fluorophore or similar reporter construct. In some embodiments, the biomarker is a nucleotide sequence, such as a unique restriction site or a readily amplified sequence. Biomarkers can be used to indicate the presence of a transgene or an endogenous 5 gene, *e.g.*, a particular allele. Use of biomarkers is known in the art, and the technique can be modified to suit the particular situation.

### C. INHIBITING ACTIVITY OF THE POLYPEPTIDES OF THE INVENTION

[0052] The invention provides methods of modulating ethylene-related processes by 10 inhibiting the activity of an ethylene receptor protein in a plant. In some embodiments, the invention provides a method of inhibiting ethylene receptor activity in a plant, the method comprising introducing a construct comprising a promoter operably linked to a nucleic acid sequence encoding an ethylene receptor, for example, the sequence of SEQ ID NO:1, 3, or 5, variants, or truncation mutants thereof.

[0053] In some embodiments, non-functional or dominant negative mutations of 15 polypeptides of the invention can be prepared by expressing a transgene that encodes a mutated dominant negative polypeptide. As explained above, dominant negative mutant polypeptides adversely affect the function of the normal, wild-type gene product within the same cell even when the cell is heterozygous (wild-type and dominant negative). Without 20 wanting to be bound by theory, the dominant negative gene product is thought to function and interact with at least some of the same elements as the wild-type gene product, but blocks some aspect of the wild-type function.

[0054] An example of a dominant negative mutation is a protein that is functional as a 25 dimer. A mutation that removes the functional domain but retains the dimerization domain causes a dominant negative phenotype because some fraction of the wild-type protein is bound by non-functional protein, resulting in a non-functional dimer. Ethylene receptor proteins are amenable to this model, because they function as homodimers in the cell membrane. Thus, as an example, a dominant negative ethylene receptor may be formed by preserving the dimerization domain but disrupting or truncating other functional domains of 30 the protein.

[0055] Non-ethylene binding ethylene receptors act in a dominant negative fashion. Thus, disruption of at least one residue in the ethylene binding domain of ZmERS1 or ZmETR2 can result in ethylene insensitivity. For example, we have found that disruption of Cys65 in ZmERS1, and Cys102 in ZmETR2 results in ethylene insensitivity in plants. Disruption of 5 other residues in the ethylene binding domain can also interfere with ethylene binding, e.g., any of residues 20-119, 25-36, or 61-91 of ZmERS1, or residues 53-156, 58-67, or 90-120 of ZmETR2.

[0056] Use of dominant negative mutants to produce inactive target genes in transgenic plants is described in Mizukami *et al.*, *Plant Cell* 8:831-845 (1996). As noted above, this 10 approach can be used to decrease ethylene sensitivity in plants by introducing dominant negative mutants of ethylene receptors into plants. For example, an altered *Arabidopsis* ERS gene can be used to confer dominant ethylene insensitivity (Hua *et al.*, *Science* 269:1712-4 (1995)). *Arabidopsis* ETR1 mutants have also been used (Wilkinson *et al.*, *Nat Biotechnol.*, 15 :444-7 (1997) and Chang *et al.*, *Science*, 262:539-44 (1993)).

15 [0057] Another strategy is to inhibit the ability of a polypeptide of the invention to interact with itself or with other molecules. This can be achieved, for instance, using specific antibodies. For example, cell-specific expression of antibodies can be used to inactivate functional domains through antibody:antigen recognition (see, Hupp *et al.*, *Cell* 83:237-245 (1995)).

20 [0058] In some embodiments, the invention includes an ethylene receptor with reduced ethylene responsiveness. In the absence of ethylene, ethylene receptors negatively regulate ethylene response pathways. Signaling downstream of ethylene receptor proteins is mediated by CTR1, a negative regulator of ethylene responses that is related to mammalian RAF-type serine/threonine kinases (Kieber *et al.*, *Cell*, 72:427-441 (1993)). Ethylene binding results in 25 decreased activity of CTR-1, and consequently, an increase in EIN2 activity (which acts downstream of CTR-1) that ultimately leads to an increase in ethylene responsiveness (Bleeker and Schaller, *Plant Physiol.*, 111:653-660 (1996); Hua and Meyerowitz, *Cell*, 72:427-441 (1998)). Thus, ethylene binding inhibits negative receptor signaling.

[0059] Accordingly, dominant negative ethylene insensitivity may be conferred by a 30 mutation that prevents ethylene binding to the receptor or one that causes constitutive signaling through CTR-1. A particular example of a dominant negative ethylene insensitive mutant is the *etr1-1* mutant of the *Arabidopsis* ETR1 gene. It has a Cys to Tyr mutation in

the second transmembrane region that prevents ethylene from binding to the receptor. Additional ethylene receptor mutants that interfere with ethylene binding or that allow CTR-1 signaling even in the presence of ethylene are known in the art (see, e.g., Wang *et al.* (2006), *Plant Cell*, 18:3429-3442).

5 [0060] In some embodiments, a truncation mutant of the ethylene receptor protein is used to confer ethylene insensitivity. The truncation may be on the N-terminal or C-terminal end of the ethylene receptor, or both. An example is an ethylene receptor that lacks the histidine protein kinase and C-terminal receiver domains. Another example is a truncation mutant that lacks the intracellular domain after the transmembrane region, for example, after

10 approximately residue 110 of SEQ ID NO:2 or residue 147 of SEQ ID NO:4 or 6. These truncation mutants can be used alone or in combination with other mutations to reduce ethylene responsiveness. In some embodiments, the truncation mutant can be slightly longer, e.g., 1-150, 1-200, 1-250, 1-300, 1-325, or 1-350 of SEQ ID NO:2 or amino acids 1-200, 1-250, 1-300, 1-350, or 1-375 of SEQ ID NO:4 or 6.

15 [0061] Site directed mutagenesis techniques may be employed with the nucleotide sequences of the invention. For example, restriction endonuclease digestion of DNA followed by ligation may be used to generate deletion variants of ethylene receptors, as described in section 15.3 of Sambrook *et al.* (*Molecular Cloning: A Laboratory Manual*, 2nd Ed., 1989 Cold Spring Harbor Laboratory Press, New York, N.Y.). A similar strategy may be used to

20 construct insertion variants, as described in section 15.3 of Sambrook *et al.*, *supra*. More recently Zhu *et al.* (1999), *Proc. Natl. Acad. Sci. USA* 96:8768-73, have devised a method of targeting mutations to plant genes *in vivo* using chimeric RNA/DNA oligonucleotides.

25 [0062] Oligonucleotide-directed mutagenesis may also be employed for preparing substitution variants of this invention. It may also be used to conveniently prepare the deletion and insertion variants of this invention. This technique is well known in the art as described by Adelman *et al.* (1983) *DNA* 2:183; Sambrook *et al.*, *supra*; "Current Protocols in Molecular Biology", 1991, Wiley (NY), F. T. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. D. Seidman, J. A. Smith and K. Struhl, eds.

30 [0063] PCR techniques may also be used to introduce mutations into the target nucleotide sequence. For example, PCR primers may be designed to introduce an insertion, deletion, or point mutation into the targeted sequence. Such methods are well understood in the art.

[0064] The above techniques may be combined with other methods to reduce ethylene responsiveness in a plant. For example, using specified promoters, the skilled practitioner can direct the expression of an exogenous ethylene receptor (e.g., ERS1, ERS2, ETR1, ETR2, or EIN4) and create a plant with desirable phenotypic characteristics, e.g., staygreen traits.

5 The skilled practitioner can choose from a variety of known promoters, whether constitutive, inducible, tissue specific. In a preferred embodiment, ethylene sensitivity is reduced using methods to increase expression of a functional ethylene receptor. In a further preferred embodiment, ethylene sensitivity is reduced using methods to decrease expression of endogenous ethylene receptors. These methods are described in detail in U.S. Patent

10 Application No. 10/876,086, which is incorporated by reference in its entirety.

#### D. ISOLATION OF NUCLEIC ACIDS USED IN THE PRESENT INVENTION

[0065] In some embodiments the invention provides for an isolated nucleic acid which encodes an ethylene receptor protein or a mutated version thereof. The isolated nucleic acid is at least 90% identical to the wild type polynucleotide sequence of the ethylene receptor gene. In one embodiment, the isolated nucleic acid comprises a sequence derived from a *Zea mays* ethylene receptor. In an exemplary embodiment, the polynucleotide sequence is at least 90%, and more typically 95%, 99%, or 100% identical to SEQ ID NO:1, 3, or 5. In other embodiments truncation mutants comprising less than the full length ethylene receptor proteins can be used. In some embodiments, the truncation mutant comprises the N terminal transmembrane domain. In a preferred embodiment, the truncation mutant comprises approximately nucleotides 1-330 of SEQ ID NO:1 (encoding amino acids 1-110 of SEQ ID NO:2). In another preferred embodiment, the truncation mutant comprises 1-441 of SEQ ID NO:3 or 5 (encoding amino acids 1-147 of SEQ ID NOs:4 and 6, respectively).

25 [0066] The isolation of nucleic acids used in the present invention may be accomplished by a number of techniques. For instance, oligonucleotide probes based on the known sequences for ethylene receptor genes in another plant species can be used to identify the desired gene in a cDNA or genomic DNA library from a desired plant species. To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation, e.g. using restriction endonucleases, and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. To prepare a library of embryo-specific cDNAs,

mRNA is isolated from embryos and a cDNA library that contains the gene transcripts is prepared from the mRNA.

**[0067]** The cDNA or genomic library can then be screened using a probe based upon the sequence of a cloned ethylene receptor gene. Probes may be used to hybridize with genomic

5 DNA or cDNA sequences to isolate homologous genes in the same or different plant species.

**[0068]** Alternatively, the nucleic acids of interest can be amplified from nucleic acid

samples using amplification techniques. For instance, polymerase chain reaction (PCR) technology can be used to amplify the sequences of the genes directly from mRNA, from cDNA, from genomic libraries or cDNA libraries. PCR and other *in vitro* amplification

10 methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes.

**[0069]** Appropriate primers and probes for identifying genes encoding polypeptides of the invention from plant tissues are generated from comparisons of the sequences of known

15 ethylene receptor proteins. For a general overview of PCR, see PCR Protocols: A Guide to Methods and Applications. (Innis, M., Gelfand, D., Sninsky, J. and White, T., eds.), Academic Press, San Diego (1990). The amplification conditions depend on many factors, including the length of the primers, the degree of identity between the primer and target sequences and the length of the target sequence to be amplified. Typical reaction conditions 20 are as follows. Reaction components: 10 mM Tris HCl, pH 8.3, 50 mM potassium chloride, 1.5 mM magnesium chloride, 0.001% gelatin, 200  $\mu$ M dATP, 200  $\mu$ M dCTP, 200  $\mu$ M dGTP, 200  $\mu$ M dTTP, 0.4  $\mu$ M primers, and 100 units per mL Taq polymerase. Program: 96°C for 3 min., 30 cycles of 96°C for 45 sec., 50°C for 60 sec., 72°C for 60 sec., followed by 72°C for 5 min.

25 **[0070]** Polynucleotides may also be synthesized by well-known techniques as described in the technical literature. See, *e.g.*, Carruthers *et al.*, *Cold Spring Harbor Symp. Quant. Biol.*

47:411-418 (1982), and Adams *et al.*, *J. Am. Chem. Soc.* 105:661 (1983). Double stranded DNA fragments may then be obtained either by synthesizing the complementary strand and annealing the strands together under appropriate conditions, or by adding the complementary

30 strand using DNA polymerase with an appropriate primer sequence.

[0071] The genus of sequences of the present invention include genes and gene products identified and characterized by analysis using the nucleic acid sequences, including SEQ ID NOs:1, 3, and 5, and protein sequences, including SEQ ID NOs:2, 4, and 6. Sequences encoding the polypeptides used in the present invention include nucleic acid sequences 5 having substantial identity to SEQ ID NOs: 1, 3, and 5. Sequences encoding the polypeptides used in the present invention include those encoding polypeptide sequences having substantial identity to SEQ ID NOs:2, 4, and 6.

[0072] Once a nucleic acid is isolated using the method described above, standard methods can be used to determine if the nucleic acid encodes an ethylene receptor protein. A nucleic 10 acid that encodes a polypeptide of the invention can be used to create a transgenic plant having staygreen traits. A transgenic plant having enhanced or increased expression of, for example, an ethylene receptor polypeptide identical or substantially identical to SEQ ID NOs: 2, 4, or 6, or a truncation mutant thereof will display a phenotype associated with an altered ethylene process within the plant, *e.g.*, delayed senescence.

15 [0073] Using standard methods, the skilled practitioner can compare the sequence of a putative nucleic acid sequence thought to encode, for example, an ethylene receptor polypeptide to a nucleic acid sequence encoding an ethylene receptor polypeptide to determine if the putative nucleic acid encodes an actual ethylene receptor polypeptide. A nucleic acid that encodes an ethylene receptor polypeptide, *e.g.*, nucleic acids comprising 20 sequences identical or substantially identical to SEQ ID NOs:1, 3, or 5, or truncation mutants thereof can be used in the methods of the present invention.

#### E. PREPARATION OF RECOMBINANT VECTORS

[0074] The invention provides a recombinant expression cassette comprising a promoter sequence operably linked to a nucleic acid sequence encoding an ethylene receptor 25 polypeptide sequence. In some embodiments, the ethylene receptor is ERS, for example, the ERS1 nucleotides represented by SEQ ID NO:1 or a truncation mutant of SEQ ID NO:1. In some embodiments, the ethylene receptor is ETR, for example the ETR2 nucleotides represented by SEQ ID NO:3 or 5, or a truncation mutant of SEQ ID NO:3 or 5.

[0075] To use isolated sequences in the above techniques, recombinant DNA vectors 30 suitable for transformation of plant cells are prepared. Techniques for transforming a wide variety of higher plant species are well known and described in the technical and scientific

literature, *e.g.*, Weising *et al.*, *Ann. Rev. Genet.* 22:421-477 (1988). A DNA sequence coding for the desired polypeptide, for example a cDNA sequence encoding a full length protein, will preferably be combined with transcriptional and translational initiation regulatory sequences which will direct the transcription of the sequence from the gene in the intended 5 tissues of the transformed plant.

[0076] For example, for overexpression, a plant promoter fragment may be employed which will direct expression of the gene in all tissues of a regenerated plant. Such promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation.

10 [0077] Alternatively, the plant promoter may direct expression of the polynucleotide of the invention in a specific tissue (tissue-specific promoters, organ-specific promoters) or specific environmental condition (inducible promoters).

15 [0078] If proper polypeptide expression is desired, a polyadenylation region at the 3'-end of the coding region should be included. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA.

20 [0079] The vector comprising the sequences (*e.g.*, promoters or coding regions) from genes of the invention will typically comprise a marker gene that confers a selectable phenotype on plant cells. For example, the marker may encode biocide resistance, particularly antibiotic resistance, such as resistance to kanamycin, G418, bleomycin, hygromycin, or herbicide resistance, such as resistance to glyphosate, chlorosulfuron or glufosinate.

25 [0080] Nucleic acid sequences of the invention, *e.g.*, nucleic acid sequences that encode ethylene receptor proteins, are expressed recombinantly in plant cells to enhance and increase levels of endogenous plant transcription factors. For example, ethylene receptor nucleic acid sequences of the invention are expressed recombinantly in plant cells to enhance and increase levels of endogenous ethylene receptor polypeptides. A variety of different expression constructs, such as expression cassettes and vectors suitable for transformation of plant cells can be prepared. Techniques for transforming a wide variety of higher plant species are well known and described in the technical and scientific literature, *e.g.*, Weising *et al.*, *Ann. Rev. Genet.* 22:421-477 (1988). A DNA sequence coding for a polypeptide described in the 30 present invention, *e.g.*, a cDNA sequence encoding a full length ethylene receptor protein, can be combined with cis-acting (promoter and enhancer) transcriptional regulatory

sequences to direct the timing, tissue type and levels of transcription in the intended tissues of the transformed plant. Translational control elements can also be used.

[0081] The invention provides a nucleic acid encoding an ethylene receptor polypeptide operably linked to a promoter which, in some embodiments, is capable of driving the transcription of the coding sequence in plants. The promoter can be, *e.g.*, derived from plant or viral sources. The promoter can be, *e.g.*, constitutively active, inducible, or tissue specific. In construction of recombinant expression cassettes, vectors, transgenics, of the invention, different promoters can be chosen and employed to differentially direct gene expression, *e.g.*, in some or all tissues of a plant or animal. Typically, as discussed above, desired promoters are identified by analyzing the 5' sequences of a genomic clone corresponding to the embryo-specific genes described here.

### 1. Constitutive Promoters

[0082] A promoter fragment can be employed which will direct expression of a nucleic acid encoding an ethylene receptor protein in all transformed cells or tissues, *e.g.* as those of a regenerated plant. Such promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include those from viruses which infect plants, such as the cauliflower mosaic virus (CaMV) 35S transcription initiation region (see, *e.g.*, Dagless, *Arch. Virol.* 142:183-191 (1997)); the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens* (see, *e.g.*, Mengiste (1997) *supra*; O'Grady, *Plant Mol. Biol.* 29:99-108 (1995); the promoter of the tobacco mosaic virus; the promoter of Figwort mosaic virus (see, *e.g.*, Maiti, *Transgenic Res.* 6:143-156 (1997)); actin promoters, such as the *Arabidopsis* actin gene promoter (see, *e.g.*, Huang, *Plant Mol. Biol.* 33:125-139 (1997)); alcohol dehydrogenase (Adh) gene promoters (see, *e.g.*, Millar, *Plant Mol. Biol.* 31:897-904 (1996)); ACT11 from *Arabidopsis* (Huang *et al.*, *Plant Mol. Biol.* 33:125-139 (1996)), Cat3 from *Arabidopsis* (GenBank No. U43147, Zhong *et al.*, *Mol. Gen. Genet.* 251:196-203 (1996)), the gene encoding stearoyl-acyl carrier protein desaturase from *Brassica napus* (Genbank No. X74782, Solocombe *et al.*, *Plant Physiol.* 104:1167-1176 (1994)), GPc1 from maize (GenBank No. X15596, Martinez *et al.*, *J. Mol. Biol.* 208:551-565 (1989)), Gpc2 from maize (GenBank No. U45855, Manjunath *et al.*, *Plant Mol. Biol.* 33:97-112 (1997)), other transcription initiation regions from various plant genes known to those of skill. See also Holtorf *et al.*, *Plant Mol. Biol.* 29:637-646 (1995).

## 2. Inducible Promoters

**[0083]** Alternatively, a plant promoter may direct expression of the nucleic acids described in the present invention, *e.g.*, nucleic acids encoding an ethylene receptor protein, under the influence of changing environmental conditions or developmental conditions. Examples of 5 environmental conditions that may effect transcription by inducible promoters include anaerobic conditions, elevated temperature, drought, or the presence of light. Examples of developmental conditions that may effect transcription by inducible promoters include senescence. Such promoters are referred to herein as "inducible" promoters. For example, the invention incorporates the drought-inducible promoter of maize (Busk (1997) *supra*); the 10 cold, drought, and high salt inducible promoter from potato (Kirch *et al.*, *Plant Mol. Biol.* 33:897-909 (1997)). Examples of developmental conditions include cell aging, and embryogenesis. For example, the invention incorporates the senescence inducible promoter of Arabidopsis, SAG 12, (Gan and Amasino, *Science* 270:1986-1988 (1995)) and the embryogenesis related promoters of LEC1 (Lotan *et al.*, *Cell*, 93:1195-1205 (1998)), LEC2 15 (Stone *et al.*, *Proc. Natl. Acad. Sci. USA* 98:11806-11811 (2001)), FUS3 (Luerssen, *Plant J.* 15:755-764 (1998)), AtSERK1 (Hecht *et al.*, *Plant Physiol.* 127:803-816 (2001)), and AGL15 (Heck *et al.*, *Plant Cell* 7:1271-1282 (1995)).

**[0084]** Alternatively, plant promoters which are inducible upon exposure to plant hormones, such as auxins or cytokinins, are used to express the nucleic acids of the invention. 20 For example, the invention can use the auxin response elements E1 promoter fragment (AuxREs) in the soybean (*Glycine max L.*) (Liu, *Plant Physiol.* 115:397-407 (1997)); the auxin-responsive Arabidopsis GST6 promoter (also responsive to salicylic acid and hydrogen peroxide) (Chen, *Plant J.* 10: 955-966 (1996)); the auxin-inducible parC promoter from tobacco (Sakai, *Plant Cell Physiol.* 37:906-913 (1996)); a plant biotin response element 25 (Streit, *Mol. Plant Microbe Interact.* 10:933-937 (1997)); and, the promoter responsive to the stress hormone abscisic acid (Sheen, *Science* 274:1900-1902 (1996)). The invention can also use the cytokinin inducible promoters of ARR5 (Brandstatter and Kieber, *Plant Cell* 10:1009-1019 (1998)), ARR6 (Brandstatter and Kieber, *Plant Cell* 10:1009-1019 (1998)), ARR2 30 (Hwang and Sheen, *Nature* 413:383-389 (2001)), the ethylene responsive promoter of ERF1 (Solano *et al.*, *Genes Dev.* 12:3703-3714 (1998)), and the  $\beta$ -estradiol inducible promoter of XVE (Zuo *et al.*, *Plant J.* 24:265-273 (2000)).

**[0085]** Plant promoters which are inducible upon exposure to chemicals reagents which can be applied to the plant, such as herbicides or antibiotics, are also used to express the nucleic

acids of the invention. For example, the maize In2 2 promoter, activated by benzenesulfonamide herbicide safeners, can be used (De Veylder, *Plant Cell Physiol.* 38:568-577 (1997)) as well as the promoter of the glucocorticoid receptor protein fusion inducible by dexamethasone application (Aoyama, *Plant J.* 11:605-612 (1997)); application of different 5 herbicide safeners induces distinct gene expression patterns, including expression in the root, hydathodes, and the shoot apical meristem. The coding sequence of the described nucleic acids can also be under the control of, e.g., a tetracycline inducible promoter, e.g., as described with transgenic tobacco plants containing the *Avena sativa* L. (oat) arginine 10 decarboxylase gene (Masgrau, *Plant J.* 11:465-473 (1997)); or, a salicylic acid responsive element (Stange, *Plant J.* 11:1315-1324 (1997)).

### 3. Tissue-Specific Promoters

[0086] Alternatively, the plant promoter may direct expression of the polynucleotide of the invention in a specific tissue (tissue-specific promoters). Tissue specific promoters are transcriptional control elements that are only active in particular cells or tissues at specific 15 times during plant development, such as in vegetative tissues or reproductive tissues.

[0087] Examples of tissue-specific promoters under developmental control include 20 promoters that initiate transcription only (or primarily only) in certain tissues, such as vegetative tissues, e.g., roots, leaves or stems, or reproductive tissues, such as fruit, ovules, seeds, pollen, pistils, flowers, anthers, or any embryonic tissue. Reproductive tissue-specific promoters may be, e.g., ovule-specific, embryo-specific, endosperm-specific, integument-specific, seed and seed coat-specific, pollen-specific, petal-specific, sepal-specific, anther-specific or some combination thereof.

[0088] Suitable seed-specific promoters are derived from the following genes: MAC1 from maize (Sheridan, *Genetics* 142:1009-1020 (1996)); Cat3 from maize (GenBank No. L05934, 25 Abler Plant Mol. Biol. 22:10131-10138 (1993)); viviparous-1 from *Arabidopsis* (Genbank No. U93215); atmycl from *Arabidopsis* (Urao, *Plant Mol. Biol.* 32:571-576 (1996); Conceicao Plant 5:493-505 (1994)); napA and BnCysP1 from *Brassica napus* (GenBank No. J02798, Josefsson, *JBL* 26:12196-12201 (1987), Wan *et al.*, *Plant J.* 30:1-10 (2002)); and the napin 30 gene family from *Brassica napus* (Sjodahl, *Planta* 197:264-271 (1995)). Fruit specific promoters include the promoter from the CYP78A9 gene (Ito and Meyerowitz, *Plant Cell* 12:1541-1550 (2000)).

[0089] The ovule-specific BEL1 gene described in Reiser, *Cell* 83:735-742 (1995), GenBank No. U39944, can also be used. See also Ray, *Proc. Natl. Acad. Sci. USA* 91:5761-5765 (1994). The egg and central cell specific FIE1 promoter is also a useful reproductive tissue-specific promoter.

5 [0090] Sepal and petal specific promoters are also used to express nucleic acids of the invention in a reproductive tissue-specific manner. For example, the *Arabidopsis* floral homeotic gene APETALA1 (AP1) encodes a putative transcription factor that is expressed in young flower primordia, and later becomes localized to sepals and petals (see, e.g., Gustafson Brown, *Cell* 76:131-143 (1994); Mandel, *Nature* 360:273-277(1992)). A related promoter, 10 for AP2, a floral homeotic gene that is necessary for the normal development of sepals and petals in floral whorls, is also useful (see, e.g., Drews, *Cell* 65:991-1002 (1991); Bowman, *Plant Cell* 3:749-758 (1991)). Another useful promoter is that controlling the expression of the unusual floral organs (ufo) gene of *Arabidopsis*, whose expression is restricted to the junction between sepal and petal primordia (Bossinger, *Development* 122:1093-1102 (1996)).

15 [0091] A pollen specific promoter has been identified in maize (Guerrero, *Mol. Gen. Genet.* 224:161-168 (1990)). Other genes specifically expressed in pollen are described, e.g., by Wakeley, *Plant Mol. Biol.* 37:187-192 (1998); Ficker, *Mol. Gen. Genet.* 257:132-142 (1998); Kulikauskas, *Plant Mol. Biol.* 34:809-814 (1997); Treacy, *Plant Mol. Biol.* 34:603-611 (1997).

20 [0092] Promoters specific for pistil and siliques valves, inflorescence meristems, caulin leaves, and the vasculature of stem and floral pedicels include promoters from the FUL gene Mandel and Yanofsky, *Plant Cell*, 7:1763-1771 (1995). Promoters specific for developing carpels, placenta, septum, and ovules are also used to express LEC2 nucleic acids in a tissue-specific manner. They include promoters from the SHP1 and SHP2 genes (Flanagan *et al.* 25 *Plant J* 10:343-353 (1996), Savidge *et al.*, *Plant Cell* 7(6):721-733 (1995)). Promoters specific for the anther tapetum may be derived from the TA29 gene (Goldberg *et al.*, *Philos Trans. R. Soc. Lond. B. Biol. Sci.* 350:5-17 (1995)).

[0093] Other suitable promoters include those from genes encoding embryonic storage proteins. For example, the gene encoding the 2S storage protein from *Brassica napus*, 30 Dasgupta, *Gene* 133:301-302 (1993); the 2s seed storage protein gene family from *Arabidopsis*; the gene encoding oleosin 20kD from *Brassica napus*, GenBank No. M63985; the genes encoding oleosin A, Genbank No. U09118, and, oleosin B, Genbank No. U09119,

from soybean; the gene encoding oleosin from *Arabidopsis*, Genbank No. Z17657; the gene encoding oleosin 18kD from maize, GenBank No. J05212, Lee, *Plant Mol. Biol.* 26:1981-1987 (1994); and, the gene encoding low molecular weight sulphur rich protein from soybean, Choi, *Mol Gen Genet.* 246:266-268 (1995), can be used. The tissue specific E8

5 promoter from tomato is particularly useful for directing gene expression so that a desired gene product is located in fruits. Suitable promoters may also include those from genes expressed in vascular tissue, such as the ATHB-8, AtPIN1, AtP5K1 or TED3 genes (Baima *et al.*, *Plant Physiol.* 126:643-655 (2001), Galawieiler *et al.*, *Science* 282:2226-2230 (1998), Elge *et al.*, *Plant J.* 26:561-571 (2001), Igarashi *et al.*, *Plant Mol. Biol.* 36:917-927 (1998)).

10 [0094] A tomato promoter active during fruit ripening, senescence and abscission of leaves and, to a lesser extent, of flowers can be used (Blume, *Plant J.* 12:731-746 (1997)). Other exemplary promoters include the pistil specific promoter in the potato (*Solanum tuberosum L.*) SK2 gene, encoding a pistil specific basic endochitinase (Ficker, *Plant Mol. Biol.* 35:425-431 (1997)); the Blec4 gene from pea (*Pisum sativum* cv. Alaska), active in epidermal tissue 15 of vegetative and floral shoot apices of transgenic alfalfa. This makes it a useful tool to target the expression of foreign genes to the epidermal layer of actively growing shoots.

15 [0095] A variety of promoters specifically active in vegetative tissues, such as leaves, stems, roots and tubers, can also be used to express the nucleic acids used in the methods of the invention. For example, promoters controlling patatin, the major storage protein of the 20 potato tuber, can be used, *e.g.*, Kim, *Plant Mol. Biol.* 26:603-615 (1994); Martin, *Plant J.* 11:53-62 (1997). The ORF13 promoter from *Agrobacterium rhizogenes* which exhibits high activity in roots can also be used (Hansen, *Mol. Gen. Genet.* 254:337-343 (1997)). Other useful vegetative tissue-specific promoters include: the tarin promoter of the gene encoding a globulin from a major taro (*Colocasia esculenta* L. Schott) corm protein family, tarin 25 (Bezerra, *Plant Mol. Biol.* 28:137-144 (1995)); the curculin promoter active during taro corm development (de Castro, *Plant Cell* 4:1549-1559 (1992)) and the promoter for the tobacco root specific gene TobRB7, whose expression is localized to root meristem and immature central cylinder regions (Yamamoto, *Plant Cell* 3:371-382 (1991)).

30 [0096] Leaf-specific promoters, such as the ribulose biphosphate carboxylase (RBCS) promoters can be used. For example, the tomato RBCS1, RBCS2 and RBCS3A genes are expressed in leaves and light grown seedlings, only RBCS1 and RBCS2 are expressed in developing tomato fruits (Meier, *FEBS Lett.* 415:91-95 (1997)). A ribulose bisphosphate

carboxylase promoters expressed almost exclusively in mesophyll cells in leaf blades and leaf sheaths at high levels, described by Matsuoka, *Plant J.* 6:311-319 (1994), can be used.

Another leaf-specific promoter is the light harvesting chlorophyll a/b binding protein gene promoter, see, e.g., Shiina, *Plant Physiol.* 115:477-483 (1997); Casal, *Plant Physiol.*

5 116:1533-1538 (1998). The *Arabidopsis thaliana* myb-related gene promoter (Atmyb5) described by Li, *FEBS Lett.* 379:117-121 (1996), is leaf-specific. The Atmyb5 promoter is expressed in developing leaf trichomes, stipules, and epidermal cells on the margins of young rosette and cauline leaves, and in immature seeds. Atmyb5 mRNA appears between fertilization and the 16-cell stage of embryo development and persists beyond the heart stage.

10 10 A leaf promoter identified in maize by Busk, *Plant J.* 11:1285-1295 (1997), can also be used.

**[0097]** Another class of useful vegetative tissue-specific promoters are meristematic (root tip and shoot apex) promoters. For example, the “SHOOTMERISTEMLESS” and “SCARECROW” promoters, which are active in the developing shoot or root apical meristems, described by Di Laurenzio, *Cell* 86:423-433 (1996); and, Long, *Nature* 379:66-69 (1996); can be used. Another useful promoter is that which controls the expression of 3 hydroxy 3 methylglutaryl coenzyme A reductase HMG2 gene, whose expression is restricted to meristematic and floral (secretory zone of the stigma, mature pollen grains, gynoecium vascular tissue, and fertilized ovules) tissues (see, e.g., Enjuto, *Plant Cell* 7:517-527 (1995)).

15 Also useful are kn1 related genes from maize and other species which show meristem specific expression, see, e.g., Granger, *Plant Mol. Biol.* 31:373-378 (1996); Kerstetter, *Plant Cell* 6:1877-1887 (1994); Hake, *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 350:45-51 (1995). For example, the *Arabidopsis thaliana* KNAT1 or KNAT2 promoters. In the shoot apex, KNAT1 transcript is localized primarily to the shoot apical meristem; the expression of KNAT1 in the shoot meristem decreases during the floral transition and is restricted to the cortex of the inflorescence stem (see, e.g., Lincoln, *Plant Cell* 6:1859-1876 (1994)).

**[0098]** One of skill will recognize that a tissue-specific promoter may drive expression of operably linked sequences in tissues other than the target tissue. Thus, as used herein a tissue-specific promoter is one that drives expression preferentially in the target tissue, but may also lead to some expression in other tissues as well.

30 **[0099]** In another embodiment, a nucleic acid described in the present invention is expressed through a transposable element. This allows for constitutive, yet periodic and infrequent expression of the constitutively active polypeptide. The invention also provides

for use of tissue-specific promoters derived from viruses which can include, *e.g.*, the tobamovirus subgenomic promoter (Kumagai, *Proc. Natl. Acad. Sci. USA* 92:1679-1683 (1995)) the rice tungro bacilliform virus (RTBV), which replicates only in phloem cells in infected rice plants, with its promoter which drives strong phloem specific reporter gene expression; the cassava vein mosaic virus (CVMV) promoter, with highest activity in vascular elements, in leaf mesophyll cells, and in root tips (Verdaguer, *Plant Mol. Biol.* 31:1129-1139 (1996)).

#### F. PRODUCTION OF TRANSGENIC PLANTS

[0100] In a further aspect, the invention provides a transgenic plant comprising a recombinant expression cassette comprising a promoter sequence operably linked to a nucleic acid sequence encoding an ethylene receptor, such as ERS (represented, for example, by SEQ ID NO:1), or ETR (represented, for example, by SEQ ID NO:3 or 5), wherein the isolated nucleic acid is at least 90% identical to the polynucleotide sequence. Alternatively, the isolated nucleic acid is a truncation mutant of SEQ ID NO:1, 3, or 5.

[0101] DNA constructs of the invention may be introduced into the genome of the desired plant host by a variety of conventional techniques. For example, the DNA construct may be introduced directly into the genomic DNA of the plant cell using techniques such as electroporation and microinjection of plant cell protoplasts, or the DNA constructs can be introduced directly to plant tissue using biotics, *e.g.*, DNA particle bombardment.

[0102] Microinjection techniques are known in the art and well described in the scientific and patent literature. The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski *et al. Embo J.* 3:2717-2722 (1984). Electroporation techniques are described in Fromm *et al. Proc. Natl. Acad. Sci. USA* 82:5824 (1985). Biotic transformation techniques are described in Klein *et al. Nature* 327:70-73 (1987).

[0103] Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is infected by the bacteria. *Agrobacterium tumefaciens*-mediated transformation techniques, including disarming and use of binary vectors, are well described in the scientific literature. See, for example Horsch *et al. Science* 233:496-498 (1984), and Fraley *et al. Proc. Natl. Acad. Sci.*

USA 80:4803 (1983) and *Gene Transfer to Plants*, Potrykus, ed. (Springer-Verlag, Berlin 1995).

[0104] Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed

5 genotype and thus the desired phenotype such as decreased farnesyltransferase activity. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker that has been introduced together with the desired nucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans *et al.*, *Protoplasts Isolation and Culture, Handbook of Plant*  
10 *Cell Culture*, pp. 124-176, MacMillan Publishing Company, New York, 1983; and Binding, *Regeneration of Plants, Plant Protoplasts*, pp. 21-73, CRC Press, Boca Raton, 1985.

Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee *et al.*, *Ann. Rev. of Plant Phys.* 38:467-486 (1987).

15 [0105] The nucleic acids of the invention can be used to confer desired traits on essentially any plant, including maize. Thus, the invention has use over a broad range of plants, including species from the genera *Anacardium*, *Arachis*, *Asparagus*, *Atropa*, *Avena*, *Brassica*, *Chlamydomonas*, *Chlorella*, *Citrus*, *Citrullus*, *Capsicum*, *Carthamus*, *Cocos*, *Coffea*, *Cucumis*, *Cucurbita*, *Cyrtomium*, *Daucus*, *Elaeis*, *Fragaria*, *Glycine*, *Gossypium*, *Helianthus*,  
20 *Heterocallis*, *Hordeum*, *Hyoscyamus*, *Lactuca*, *Laminaria*, *Linum*, *Lolium*, *Lupinus*, *Lycopersicon*, *Macrocytis*, *Malus*, *Manihot*, *Majorana*, *Medicago*, *Nereocystis*, *Nicotiana*, *Olea*, *Oryza*, *Osmunda*, *Panieum*, *Pannisetum*, *Persea*, *Phaseolus*, *Pistachia*, *Pisum*, *Pyrus*, *Polypodium*, *Prunus*, *Pteridium*, *Raphanus*, *Ricinus*, *Secale*, *Senecio*, *Sinapis*, *Solanum*, *Sorghum*, *Theobromus*, *Trigonella*, *Triticum*, *Vicia*, *Vitis*, *Vigna*, and *Zea*.

## 25 G. DETECTION OF THE TRANSGENIC PLANTS OF THE INVENTION

[0106] The invention provides a method of modulating ethylene receptor activity in a plant.

In some embodiments, the method further comprises selecting a plant with a phenotype of delayed senescence in its reproductive plant structure. In some embodiments, the

reproductive structure is a seed. In some embodiments, the phenotype is multiple embryos in  
30 a single seed. In some embodiments, the construct is introduced by a sexual cross.

[0107] In some embodiments, screening further comprises detecting a plant having a desirable phenotype. For example, leaf color can be examined to determine if the photosynthetic life-span of the plant has been affected. Plants with extended photosynthetic life cycles are characterized by leaves that stay green for a longer duration of time as compared to wild type plants. In addition, chlorophyll levels and CO<sub>2</sub> assimilation can be measured using well known techniques. The size of plant vegetative and reproductive structures can be examined to determine if they are larger or smaller than those of a wild type plants. Transgenic plants of the present invention may possess larger fruit, ovules, seeds, pollen, embryonic tissue, flowers, flower parts such as pistils, stamens, sepals, petals, carpels, leaves, stems, tubers, roots, vascular tissue, provascular tissue or root or stem meristems. The resultant transgenic plants can be assayed for increased drought tolerance. Methods for assaying for increased drought tolerance are known and include measuring transpiration rate of transgenic plants, stomatal conductance, rate of water loss in a detached leaf assay or examining leaf turgor. Transgenic plants with decreased transpiration rates, for example, have increased drought tolerance.

[0108] Plants with reduced ethylene sensitivity can also be selected by testing the ability of the plant to grow at higher density. Plants with this ethylene resistant phenotype are advantageously planted at high density, regardless of selection process. Similarly, plants with an ethylene resistant phenotype are advantageously planted in drought conditions, regardless of the selection process.

[0109] Plants with reduced ethylene sensitivity can also be selected by detecting the presence of a biomarker, *e.g.*, a marker gene. A marker gene generally gives rise to a readily detectable phenotype that cosegregates with a staygreen phenotype, which may not be as immediately detectable.

[0110] Ethylene sensitivity can also be observed by the “triple response” phenotype, which involves radial expansion of the hypocotyl, inhibition of root and hypocotyl elongation, and exaggeration of the apical hook (Neljubow (1901), *Pflanzen Beih. Bot. Zentralbl.*, 10:128-139). Mutants that cause constitutive ethylene signaling are characterized by a reduced stature compared to wild type plants (Guzman and Ecker (1990), *Plant Cell*, 2:513-523; Kieber *et al.* (1993) *Cell* 72:427-441).

[0111] Conversely, ethylene insensitivity may be observed by delayed dark-induced loss of chlorophyll and/ or an increase in plant size. Ethylene insensitive plants may be larger, even

when compared to wild type plants grown under optimal conditions, *i.e.*, in the absence of trace levels of ethylene that may accumulate in enclosed growth facilities. However, it will be recognized by those with skill in the art that ethylene responses vary according to plant species and environmental conditions. The staygreen phenotype can be conveniently assayed 5 using standard assays. For example, dark-induced senescence assays can be used. Such assays typically involve sheathing leaves while still attached to the plant for one week. The lack of light induces leaf senescence, which can be delayed, as compared to controls, in plants of the invention.

[0112] Biochemical means for detecting transgenic plants are also well known, *e.g.*, 10 detecting expression of a biomarker such as a reporter gene inserted on the transgenic construct, or the transgenic nucleic acid or protein sequences in the plant. Means for detecting and quantifying mRNA and proteins are well known in the art, *e.g.*, Northern Blots, PCR, Western Blots, and activity assays. For example, after introduction of the expression cassette into a plant, the plants are screened for the presence of the transgene and crossed to 15 an inbred or hybrid line. Progeny plants are then screened for the presence of the transgene and self-pollinated. Progeny from the self-pollinated plants are grown. The resultant transgenic plants can be examined for any of the phenotypic characteristics associated with altered ethylene-related processes, *e.g.*, characteristics associated with staygreen traits or delayed senescence. For example, using the methods of the present invention, inhibition of 20 the nucleic acids or proteins described in the present invention may delay senescence in cells of a vegetative or reproductive plant structure.

[0113] It is understood that the examples and embodiments described herein are for 25 illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety for all purposes.

## EXAMPLES

[0114] Standard methods were used in the methods described briefly below.

**Example 1: Expression of *Arabidopsis etr1-1* in *Zea mays* results in a delayed ethylene response**

*Construct design and plant transformation*

[0115] The *Arabidopsis etr1-1* mutant has been described previously. It is a dominant negative mutant ethylene receptor that results in ethylene insensitivity. Transgenic maize containing the wheat *Arabidopsis etr1-1* under the control of the maize ubiquitin (Ub) promoter in pACH18 (Christensen and Quail (1996), *Transgenic Res.*, 5: 213-218) were generated by particle bombardment of embryogenic A188 X B73 (Hill) callus as described (Gordon-Kamm *et al.* (1990), *Plant Cell*, 2: 603-618). Co-transformation with the *bar* gene provided bialaphos selection for the transformed callus (De Block *et al.* (1987) *EMBO J.* 6: 2513-2518). Regenerants containing the Ub- *etr1-1* constructs were identified using PCR.

*Results*

[0116] To determine whether introducing a state of ethylene insensitivity can alter ethylene-mediated events in maize, the transgenic maize lines expressing *Arabidopsis etr1-1* under the control of the maize ubiquitin (Ub) promoter were grown under greenhouse conditions. Leaves of T<sub>0</sub> progeny confirmed by PCR to harbor the *etr1-1* transgene exhibited altered growth phenotypes. To examine whether the expression from the *etr1-1* transgene resulted in an alteration in leaf senescence or a staygreen phenotype, leaf 6 and leaf 7 from adult T<sub>0</sub> generation plants were sheathed while still attached to the plant to prevent light from reaching the leaves for one week. The lack of light normally induces leaf senescence, referred to as dark-induced senescence, and is often used as a measure of staygreen potential.

[0117] Transgenic leaves expressing *etr1-1* exhibited delayed leaf senescence, suggesting that a state of ethylene insensitivity in maize results in a staygreen phenotype. These results demonstrate that engineering a state of ethylene insensitivity in maize through the expression of dominant negative mutant ethylene receptor, *e.g.*, following a Cys to Tyr mutation in the second transmembrane or other mutations with similar effect can alter ethylene-mediated processes, *e.g.*, leaf aging and senescence.

**Example 2: Expression of dominant negative ZmERS1 and ZmETR2 in *Arabidopsis* results in ethylene insensitivity**

[0118] We next set out to determine whether expression of a mutant *Zea mays* ethylene receptor would result in ethylene insensitivity in *Arabidopsis* in the same manner.

*Construct design and plant transformation*

[0119] *Arabidopsis* were transformed with a pBI121 construct, according to the method described in Example 1. The constructs comprised a 35S promoter operably linked to one of the following coding sequences: *ein2-5* (an ethylene insensitive *Arabidopsis* mutation described in Alonso *et. al.* (1999) *Science* 284:2148-52); mutant ZmERS1; or mutant ZmETR2 (ZmETR40). The ZmERS1 and ZmETR2 mutants are dominant negative ethylene receptors (non-ethylene binding), and the sequences are described herein. Wild type plants were used as an ethylene-sensitive control (designated WT). Germination on ACC, the precursor to ethylene, inhibits seedling growth. Ethylene insensitive transformants were screened by germinating the transformants in the dark on medium containing 20  $\mu$ M ACC. Three independent lines of ZmERS1 transformants (designated MS1-11, MS2-12, and MS1-15) and three independent lines of ZmETR2 transformants (designated MT2-4, MT2-5, or MT2-9) were selected for further study.

*Results*

[0120] To show that mutant ZmETR2 and ZmERS1 ethylene receptors confer a state of ethylene insensitivity, the mutant lines were germinated on medium containing 20  $\mu$ M ACC or on medium containing 20  $\mu$ M Ag, which chemically confers a state of ethylene insensitivity in *Arabidopsis*. The data show that seed expressing either mutant ZmERS1 or ZmETR2 are ethylene insensitive in that they do not exhibit the classic "triple response" to ethylene but rather exhibit long, thin hypocotyls with no apical hook. Similar results were observed for the ethylene insensitive mutant, *ein2-5*. In contrast, wild type seedlings exhibit the classic response to ethylene. In the presence of Ag, WT seedlings are also ethylene insensitive. Quantitative measurements for 5 day old seedlings are shown in Table 1 ( $p < 0.001$  unless otherwise stated). These data show that mutant ZmERS1 and ZmETR2 ethylene receptors can confer ethylene insensitivity in *Arabidopsis*.

*Table 1: Expression of mutant ZmETR2 and ZmERS1 receptors confers ethylene insensitivity in Arabidopsis*

	20mM ACC		5mM Ag	
	Hypocotyl length (mm)	Root length (mm)	Hypocotyl length (mm)	Root length (mm)
WT	5.0 $\pm$ 0.71	3.00 $\pm$ 1.03	14.7 $\pm$ 1.12	6.69 $\pm$ 1.41
<i>ein2-5</i>	13.9 $\pm$ 1.71	7.43 $\pm$ 2.63	11.0 $\pm$ 3.13	6.27 $\pm$ 1.93 $p = 0.371$

MT2-4	14.2 ± 1.61	6.05 ± 1.36	13.0 ± 3.20	7.33 ± 1.81 p = 0.122
MT2-5	14.4 ± 1.97	4.87 ± 1.16	11.5 ± 2.87	5.39 ± 1.27
MT2-9	13.8 ± 2.36	6.27 ± 1.76	12.1 ± 2.40	7.07 ± 1.58 p = 0.354
WT	6.6 ± 133	3.10 ± 0.92	16.9 ± 2.41	5.48 ± 0.94
<i>ein2-5</i>	17.0 ± 2.98	6.51 ± 1.56	14.7 ± 3.78	5.97 ± 1.56 p = 0.349
MS1-11	17.6 ± 3.10	4.60 ± 0.88	10.9 ± 3.68	6.35 ± 1.87 p = 0.086
MS1-15	18.0 ± 1.38	5.88 ± 1.40	11.7 ± 5.01	5.51 ± 1.45 p = 0.921
MS1-12	8.8 ± 1.37	3.33 ± 0.97 p = 0.406	17.3 ± 1.95 p = 0.561	7.14 ± 1.97

[0121] To examine the level of ethylene insensitivity conferred over a range of ACC concentrations, seeds from one ZmERS1 line (MS1-11) and one ZmETR2 line (MT2-9) were germinated on medium containing various levels of ACC (from 1.0 to 20 µM). Expression of 5 the mutant ethylene receptors conferred ethylene insensitivity across the range of ACC concentrations. There was some decrease in the level of ethylene insensitivity achieved at higher ACC concentrations but it remained substantially greater than in WT seedlings. Quantitative measurements are shown in Figure 1.

[0122] Ethylene insensitivity could also be shown when seedlings were germinated on 20 µM ACC in the light. WT seedlings exhibited non-expanded cotyledons whereas lines 10 expressing either mutant ZmERS1 or ZmETR2 were similar to the ethylene insensitive mutant *ein2-5*. Subsequent growth on 10 µM ACC in the light showed that WT seedlings were substantially smaller than those grown on Ag, whereas lines expressing either mutant ZmERS1 or ZmETR2 were similar to WT seedlings grown on Ag or to the *ein2-5* ethylene 15 insensitive mutant.

[0123] Plants expressing either mutant ZmERS1 or ZmETR2 also had a larger leaf size, similar to the *ein2-5* and *etr1-1* ethylene insensitive mutants. At 7 weeks of age, plants 20 expressing either mutant ZmERS1 or ZmETR2 showed delayed leaf senescence and a staygreen phenotype, similar to that observed for *ein2-5*. When assayed at 4 weeks, little difference in chlorophyll content was observed (Table 2). However, plants expressing either mutant ZmERS1 or ZmETR2 showed some delay in flowering and consequently, an increase in leaf number.

Table 2: Phenotypes of *Arabidopsis* expressing mutant *Zea mays* ethylene receptors

	Flowering time (days)	Leaf number	Chlorophyll a (ng/mg FW)	Chlorophyll b (ng/mg FW)	a/b Ratio
WT	22.0	11.1 ± 1.7	960 ± 92	281 ± 21	3.42
<i>ein2-5</i>	22.5	12.1 ± 2.0	923 ± 67	296 ± 21	3.11
MT2-4	22.5	12.9 ± 1.9	988 ± 86	298 ± 27	3.32
MT2-5	22.5	13.8 ± 2.0	903 ± 93	271 ± 16	3.33
MT2-9	28.0	16.2 ± 1.2	965 ± 104	304 ± 39	3.18
MS1-11	24.0	12.5 ± 1.7	1016 ± 144	312 ± 33	3.25

[0124] We next sought to determine if expression of the mutant ZmERS1 is fully dominant in conferring a state of ethylene insensitivity when present in a hemizygous versus 5 homozygous state. Crosses between ZmERS1 mutant (MS1-11) homozygotes and wild type seeds were performed to obtain seeds hemizygous for the ZmERS1 mutant. These were tested against seed that were homozygous for mutant ZmERS1. The plants were grown in 20 µM ACC for 5 days in the dark. As shown in Table 3, the hemizygous seedlings are as 10 insensitive to ethylene as homozygous seedlings in this triple response assay ( $p < 0.001$ ). The hemizygous and homozygous mutants were also equally insensitive when grown in the light on 20 µM ACC.

Table 3: ZmERS1 mutants are dominant following their hemizygous expression in *Arabidopsis*

	Hypocotyl length (mm)	Root length (mm)
WT	8.52 ± 0.65	2.78 ± 0.83
Hemizygous MS1-11	16.5 ± 4.78	4.78 ± 1.60
Homozygous MS1-11	17.3 ± 4.77	4.77 ± 0.92

## SEQUENCE LISTING

## SEQ ID NO:1

## Zea mays ERS1-like coding sequence (Cys → Tyr 65)

ATGGACGGATGCGATTGCATAGAGCCACTATGGCCTACCGATGATCTTCTCGTCA  
 5 AGTATCAGTACATCTCAGACTTCTCATGCCCTGCGTACTTCTGATTCCATTG  
 GAGCTCATATATTTGTGAAGAAGTCGCTTCTCCCACAGATGGGTCTGA  
 TCCAGTTGGTGCCTTATAGTTCTTATGGGCAACCCATCTGATAAACCTGTGG  
 ACGTTCACCAACATACAAAGACCGTTGCGATGGTCATGACCATAGCGAAGATT  
 CTACAGCAGTCGTGTCTGTGCAACTGCTTGTGCTCGTTATATCATTCCGAC  
 10 TTGTTGAGCGTGAAGAACTAGGGAGTTGTTCTGAAGAATAAAGCTGAGGAGCTT  
 GATAGAGAGATGGGACTTATAAGGACGCAAGAGGGAGACTGGTAGACATGTTAGG  
 ATGCTTACACATGAAATCAGAAGTACTCTTGATAGACATACAATTTGAAGACTA  
 CTCTCGTTGAGCTAGGAAGGACCTTGGGCTTGGAGAAATGTGCATTGTGGATGCC  
 ATCTCGAAGTGGCTCAAGCCTTCAGCTTCTCATACTTGCGCCACCAGATTACTG  
 15 TTGGATCATCGGTGCCAATGAATCTTCTGTCGTCATCAAGTGTCAAGTCAAGTGGAGCT  
 CCGGGCAATCATAATACCCACACATCTTGGCGGGGTTCGACCTCTGCA  
 GGGCGATATGTTCCACCAGAAGTGGCCGAGTCCGTGTACCTCTACATCTT  
 CAAACTTCAAATAATGATTGGCCTGAGCTCAGCAAAAGCTTGCAATCAT  
 GGTTTGATGCTTCCATCTGATAGTGTAGAAAATTGCATGTGCATGAATTGGAG  
 20 CTGGTTGAGGTCGTTGCTGATCAGGTAGCAGTTGCACTATCTCATGCAGCTATT  
 TCGAAGAGTCATGCGGGCACGTGATTACTAATGGAGCAGAAATTGCGCTGG  
 ATTTAGCTCGAAGAGAGGGCTGAGATGGCTATCCGTGCTCGCAATGATTCCCTAGC  
 TGTTATGAATCACGAAATGAGAACACCCATGAATGCAATAATAGCCCTTCCTCC  
 TTGCTTTGGAAACTGAGCTTACTCCTGAGCAGCGTCTAATGGTGGAAACAGTAC  
 25 TGAAAAGCAGCAATTGTTAGCAACACTCATCAATGATGTTCTGGATCTTCAA  
 ACTCGAGGATGGAAGCCTGAACTGGAGATTAAAGCATTCAATCTCATGCTGTT  
 TTCAAAGAAGTAATGGGTTTCACTAAACCAATTGCATCTACAGAGGCTATCTG  
 TATCGGTTATGTTGGCACCAGATCTGCCGTTATGTCAATTGGTGTGAAAAGAG  
 ACTCATGCAAACATTCTGAACATCTCTGGCAATGCTGTAAAGTTACCAAGGGAG  
 30 GGACACATCACGCTTGTAGCTTCCATTGTGAAGGCTGACTCTTGAGAGAGTTCA  
 GAACCCAGAATTCTACCAACTGCAAGTGTGAAACATTCTATTGAAAGTTCA  
 GGTAAAAGATACAGGCTGTGGAGTTAGCCTCAGGATCTACCTCATGTATTCA  
 AAGTTGCTCATCCTCAAAGTGGAGGAACCGAGGGTTAATGGTAGTGGTCTTG  
 GCCTGCCATATGCAAGAGGTTGTTAGTCTTATGGGAGGGCACATCTGGATCGA  
 35 CAGCGAAGGAACCGGAAGAGGTTGCACCGCAACATTGTCATCAAGCTCGCGT  
 GTGTGACAACACAAACACCTACCAAAAGCAGCTGGTCTCTAATCTGGCAAG  
 CAGTGCAGACTCCAATTGTCGCTCCGAAAGTGTGCTGCCGACGGGAGAGGATCT  
 GTTCCCTGAAATCTCGTACCAAGAAGCGTA

## 40 SEQ ID NO:2

## Zea mays ERS1-like protein sequence (Cys → Tyr 65)

MDGCDCIEPLWPTDDLLVKYQYISDFIALAYFSIPLIELYFVKKSSFPYRWVLIQFG  
 AFIVLYGATHLINLWTFTHTKTVAMVMTIAKVSTAVVSCATMLVHIIIPDLSVK  
 TRELFLKNKAELDREMGLIRTQEETGRHVRMLTHEIRSTLDRHMLKTTLVELGRTL  
 45 GLEECALWMPSRSGSSLQLSHTLHHQITVGSSVPINLPVINQVFSSNRAIIIPHTSPLARI  
 RPLTGRYVPPEVAARVPLLHLSNFQINDWPELSAKSFAIMVLMPLSDSARKWHVHE  
 LELVEVVADQVAVALSHAIALEESMRARDLLMEQNVALDLARREAEMAIRARNDL

5 AVMNHEMRTPMNIIALSSLLTELTELPEQRLMVETVLKSSNLLATLINDVLDLSKLE  
 DGSLELEIKAFNLHAVFKEVMGFIKPIASIKRLSVVMLAPDLPLCAIGDEKRLMQTIL  
 NISGNAVKFTKEGHITLVASIVKADSLREFRTPEFHPTASDDHFYLKVQVKDTGCGIG  
 PQDLPHVFTKFAHPQSGGNRGFNGSGLGLAICKRFVSLMGGHIWIDSEGTGRGCTAT  
 FVVKLGVCDNTNTYQQQLIPLVWPSSADSDLRAPKPLPDGRGSTPLKSRYQRSV

## SEQ ID NO:3

**Zea mays ETR2-like (ZmETR 9) coding sequence (Cys → Tyr 102)**

10 ATGGTGGTGGGAACGGCACTGCTGCGCGGGTTTCCCTCCGCGTGGATCCTCCTGT  
 TCCTCTCCTCCCTGCTCCTCTCGCCGTAGCGGCGTCTGCGATTCCGGCCACTGC  
 GCGCGCTGCGACGACGCCGACGACGGCGCCCTCTCCAGCACCTATAACATCCTG  
 CAATGCCAGAAGGTAGCGACTTCCTCATCGCCGCGCCTACTTCTCCATCCCCTG  
 TCGAGCTGCTCTACTTCGCCACCTGCTCCGACCTCTTCCCCCTAAATGGATCGTG  
 CTGCAGTTCGCGCCTTCATCGTGCCTACGGCCTCACGCACCTCATCACTGTGTT  
 15 CACCTACGAGCCGCACTCCTCACCTCGTACTCGCCCTTACCGTCGCCAAGTTCC  
 TGACGGCACTGGTCTCCTCGCGACGGCCATCACCTGCTGACGCTGATACCACA  
 GCTCCTGAGGGTGAAGGTAGGGAAAACCTCCTGATGAACAAGGCACGCTGGATGTG  
 GGACCGGGAGGTGGGGAGGATGAAAAGGAAGAAGAGGGCAGCTGGCATGTGC  
 GCATGCTCACACAGGAGATCCGCAAGTCGCTCGACAGACATACCATCTTGTACAC  
 20 CACCATGGTTGAGCTCTCGAAGGCACTGGAACACTCGAGAATTGTGCTGTGGATg  
 CCTGATGAGACCAGGAGCACGATGATCTAACACATCAGCTGAGGGAAAGGGAT  
 ATAATGGACCCACAGAAACACTCGATTCCATTGATGATCCGGATGTTCAAGAAA  
 TAAAGGCAACCAAGGATGCAAAGTTCTGGCCAGATTGGCGCTAGGGGTT  
 CTAGCCGAAGCAAGCATGAAGCAGGGCCTGTGGCTGCAATAAGGATGCCGATGT  
 25 TAAGGGTGTCAAATTCAAAGGAGGGACTCCGGAAGTGTGATGCAGACGAGCTATG  
 CTATCTGGTTCTGGTTTGCTTAATGATGGTTATTAGGGTGGGTCGAAGAGA  
 GTTGGAGATTGTTGAGGTAGTTGCTGACCAAGTTGCTGAGCTGCTCTGTACATGCT  
 GCACTCCTAGAGGAGTCTCAGCTGATGCGAGAGAAGCTGCGAGCAGCATAGG  
 GACTTGCTGCAGGCAAAGGATGAAGCCATGAGGGCAGGGGACGCTAGGAATTCC  
 30 TTCCAGACTGCAATGTACGATGGAATGCGAAGGCCAATGCACTCAATCCTGGTC  
 TCGTCTCAATGATGCAACAGGAGAGCATGAATCCAGAGCAAAGGCTTGATGG  
 ATGCCATTGCCAAGACAAGCAGTGTGCACTTGTCTGGTCAGGAGGCCGTTAACCT  
 AAACATCGACAATGAACTGTGAGCACTTGTCTGGTCAGATGTCTAACGGTTGCAAG  
 35 GGTGTGGAGTTGAGTTCAAGTGGAGAATTCTTGCCAGAAAGGATATTGGTG  
 ATGAGAAGAGAGTCTCCATTGTCCTGCACATGGTAGGCACCTAACAGACCG  
 ATGTAATGCTGGCTGTATCTCATTATGTAATGTCCATAATGAGGTTGAAGAT  
 AGGCATAATCATGACTGGATGCTGCGAAGAGCAAACCTCTGGGGCTATGTAT  
 GTGTGAAATTGAGATTAGGATTAGAAAATCAAAGGGCTATCTGTTGAGTTCATC  
 40 AAGCAGTCAGATAAGTCAGGGATCCAACCCAAACAATTCTGAGATGGGCTTAG  
 CTTCAATATGTCAAGAAGATTGTGAGATGATGAATGCAATTGGTCAGTA  
 TCAGATTCTAAAGCATGGAGAAACTATCATGCTAGTCCTCCAGTTCCAGTTGG  
 AACCTGTGACTCCGGTCTGGAGCGTCTCAGATTGTACAGATCATCCGCAAT  
 TCCCAACTTTAATGGGCTCAGAGTCCTCTGCGGACAGCGACTGCACCAACCGA  
 45 GCTGTAACTCACAGGCTCTAGAGAAGCTTGGTGCAGTCCTTCCAGCTGGTGGT  
 CTGGCGTCCAATGCATCAGCTCCTCGCTGCGGAGTCGTCCTCCAGCTGGTGGT  
 TCTTGATCTTGACATGCAGACGATGGATGGATTGCAAGTAGCCCGCGATCAGG  
 AAGTTCAGTAGCAATAGTTGGCTGCCGTTGATTATTGCCCTAGCAGCAAGAATCG

ACGACAAACATCCGGGATCGTGCAGAGGTCAGGAGTAAATGGCCTGATCCAGA  
AACCGGTACATTAGCCGCGCTGGGAGATGAACCTGTATAGTCCTTCAGAACAA  
AT

5 **SEQ ID NO:4****Zea mays ETR2-like (ZmETR 9) protein sequence (Cys → Tyr 102)**

MVVGTLRGSWILLFLSSLLSPSAASVDFGHCGGCDDADDGALSSTYNILQC  
QKVSDFLIAAYFSIPELLYFATCSDLFPLKWIVLQFGAFIVLYGLTHLITVFTYEPHS  
FHLVLALTVAKFALTVSFATAITLLTIPQLLRVKVRENFLMNKARELDREVGRMK  
10 RKEEASWHVRMLTQEIRKSLDRHTILYTTMVELSKALELQNCAVWMPDETSTMIL  
THQLRERDIMDPQKHSIPIDDPDVQEIKATKDAKVLGPD SALGVSSRSKHEAGPVAII  
RMPMLRVSNFKGGTPEVMQTSYAILVLVLPNDGSLGWRRELEIVEVADQVAVAL  
SHAALLEESQLMREKLAEQHRDLLQAKDEAMRAGDARNFSQTAMYDGMRRPMHSI  
LGLVSMQQESMNPEQRLVMDAIAKTSSVASTLMNDVMQTSTMCEHLSLVRRPF  
15 NLHSFIKEVVGVVRCLTGCKGVEFQVENSLPERIIGDEKRVFHIVLHMVGTLTDRC  
NAGCISLYVNNEVEDRHNHDWMLRRANFSGGYVCVKFEIRIRSKGYLLSSSSQ  
ISQGSKPNNSEMGLSFNMCKKIVQMMNGNIWSVSDSKSIGETIMLVLFQLEPVTPVS  
GASSDLYRSSAIPNFNGLRVLLADSDCTNRAVTHRLLKGCRVLSVASGVQCISSFA  
AESSFQLVVLDLDMQTMDGFEVARAIRKFSSNSWLPLIIALAARIDDNIRDRCQRSGV  
20 NGLIQKPVTLAALGDELYRVLQNN

**SEQ ID NO:5****Zea mays ETR2-like (ZmETR 40) coding sequence (Cys → Tyr 102)**

ATGGTGGTGGGAACGGCGCCGTGCGGGGTCTCCGTCTCCCTCCGTGGATCCTCC  
25 TGCTCCTTCCCTCCCTGCTCCTCTCGCCGTGGCGCGTCCGTGATTCCGGCAC  
TGCAGCTGCGACGACGCCGACGACGGCGCCCTCTCGAGCACCTACAACATCCTG  
CAATGCCAGAAGGTCAAGCAGCTTCCATCGCCGCGCCTACTTCTCCATCCCAC  
TCGAGCTGCTCTACTTCGCCACCTGCTCCGACCTTTCCCCCTCAAATGGATCGT  
CTGCAGTTGGCGCCTTCATCGTGCCTACGGCCTCACGCACCTCATCACCGTGT  
30 CACCTACGACCCGCACTCCTCACCTCGTGCCTGCCCTCACCGTCGCCAAGTTC  
ATGACGGCACTAGTCTCCTCGCCACAGCCATCACGCTGCTGACACTGATACCGC  
AGCTCCTGAGGGTGAAGGTCAAGGGAAAATTCCTGGTGAACAAGGCACGTGAGC  
TGGACCGGGAGGTGGGGATGATGAAATGAAAGAAGAGGGCAGCTGGCATGTG  
CGTATGCTCACACAGGAGATCCGCAAGTCGCTCGACAGGCACACCACCTGTACA  
35 CCACCATGGTTGAGCTCTCGAAAGCGCTGGAACTGCAGAATTGTGCTGTGGAT  
GCCCGATGAAACCAGGGAGCGAGATGATCTTAACATCAGCCAAGGGAAAGGGA  
TATAATGGACCAAGCAGAACTGCTCGATTCTATTGATGATCCAGATGTTCAAGAA  
ATAAAGGCTACCAAGGACGCAAAAGTTCTGGGCCAGATTGGCACTAGGGGTT  
GCTACCCGCAAGCTTGACGTGGGCCTGTGGCTGCAATAAGGATGCCGATGTTA  
40 AGGGTGTCAAATTCAAAGGAGGGACTCCAGAAGTGAATGCAGACGAGCTATGCT  
ATCTGGTTCTGGTTTGCCTAATGATGGTTCATGGGGTGGGGTAGAAGAGAGT  
TGGAGATTGTTGAAGTAGTTGCTGACCAAGTTGCGGGTGGCTTGTACATGCTGC  
ACTCCTAGAGGGAGTCTCAGCTGATGCGAGAGAAACTTGCTGAGCAGTATAAGGGA  
CTTGCTGCAGGCAAAGCATGAAGCCATGAGGGCAGGGGAAGCTCGGAATTCTT  
45 CCAGACTGCAATGTACGACGGAATGCGAAGGCCAATGCACTCAATCCTGGTCTT  
GTCTCAATGATGCAACAGGAGAGCATGAATCCAGAGCAAAGGGTTGTGATGGAT  
GCCATTGCCAAGACAAGCAGTGTGCGTCCACACTGATGAATGATGTGATGCAA

5 ACATCGACAATGAACTGTGAGCACTTGTCTTGGTGAGGAGGCCGTTCAATCTTC  
 ATTCTTTATTAAAGAAGCTGTTGGAGTGGTCAGATGTCTAATGGTTGCAAGGG  
 TGAGAGAGTTGAGTTCAAGTGGATAATTCTTGCACAGAAAGGATCATTGGTGAT  
 GAGAAGAGAGTCTCACATTGTCCTGCACATGGTAGGCACCCCTAATAAACCGAT  
 10 5 GTAATGTCGGCTGTATCTCGTTATATGTCAATGGTATAATGAGGTTGAAGAGAG  
 GCATAATCATGACTGGATGCTGCGGAGAACAAACTCTCTGGGGCTATGTTGT  
 GTGAAATTGAGATTAGGATTAGAAAATCCAAGGACTATCTTGAGTTCAAACG  
 GTCAGATAAGTCATGGGTCCAACCAAACAATTCTGAGATGGGCTTAGCTCA  
 ATATGTGCAAGAAGATTGTGAGATGATGAACGGCAACATTGGTCAGTATCAG  
 15 10 ATTCTAAAAGCGTTGGAGAAACCATCATGCTGGCCTCCAGTTCCAGCTGCAGCC  
 TCTGACTGCGGTCTCCTCCGCGCGTCTCAGACTTGAGCCGATCGTCCGCAATC  
 CCCAACTTCAACGGGCTCAGAGTCCTCTGGCGGACAGCGACGACACCAACAGA  
 GCAGTAACACACAGGCTCTGGAGAAGCTCGGCTGCCGGCTTCGGTCGCCT  
 20 15 CCGGTGTCCAATGCACGAGCTCCTCGCCGCCAGCCGCTTCAGCTGGTGGT  
 CCTGGACCTCGCCTTGCAGAGGACGGACGGAGCCGCTCGAAGTGGCCCGCGATCAG  
 GAAGTTCACTAGCAATAGCTGGCTGCCGCTGATCGTCGCCCTAGCTGCGAGGATC  
 GATGACAAGGTCCGAGACGGATGCCAGAGGTGGGATAAGCGGCCTGATCCAG  
 AAACCGGCCACGTTAGCTGCGCTGGAGATGAGCTGTATAGGGCCTTCAGAAC  
 AGT

**SEQ ID NO:6**

**Zea mays ETR2-like (ZmETR 40) protein sequence (Cys → Tyr 102)**

25 MVVGTAPCGVSVSSWILLLSSLLSPSAASVDFGHCGCDDADDGALSSTYNILQC  
 QKVSDFLIAAYFSIPELLYFATCSDLFPLKWIVLQFGAFIVLYGLTHLITVFTYDPHS  
 FHLVLALTVAKFMTALVSFATAITLLTLIPQLLRVKVRENFLVNKARELDREVGMMK  
 MKEEASWHVRMLTQEIRKSLDRHTILYTTMVELSKALELQNCAVWMPDETRSEML  
 THQPRERDIMDQQNCSIPDDPDVQEIKATKDAKVLGPDSALGVATRKLDVGPVAII  
 RMPMLRVSNFKGGTPEVMQTSYAILVLVLPNDGSLGWGRRELEIVEVADQVAVAL  
 30 30 SHAALLEESQLMREKLAEQYRDLQAKHEAMRAGEARNNSFQTAMYDGMRRPMHSI  
 LGLVSMMQQESMNPEQRVVMDAIAKTSSVASTLMNDVMQTSTMCEHLSLVRPF  
 NLHSFIKEAVGVVRCLTGCKGVEFQVDNSLPERIIGDEKRVFHIVLHMVGTLINRC  
 NVGCISLYVNNGHNEVEERHNHDWMLRTNFSGGYVCVKEIRIRKSODYLLSSNGQI  
 SHGSKPNNSEMGLSFNMCKKIVQMMNGNIWSVSDSKSVGETIMVLQFQLQPLTAV  
 35 35 SSAASSDLRSSAIPNFNGLRVLLADSDDTNRAVTHRLEKLGCRVLSVASGVQCTSS  
 FAAEPSFQLVVLSDLALQRTDGLEVARAIRKFSSNSWLPLIVALAARIIDDKVRDGCQR  
 SGISGLIQQPATLAALGDELYRVLQNS

WHAT IS CLAIMED IS:

1                   1. An isolated polynucleotide encoding a non-ethylene binding ethylene  
2 receptor polypeptide, wherein:

3                   (a) the non-ethylene binding ethylene receptor polypeptide comprises a  
4 sequence at least 95% identical to a sequence selected from the group consisting of: SEQ ID  
5 NO:2, SEQ ID NO:4, and SEQ ID NO:6; and

6                   (b) expression of the non-ethylene binding ethylene receptor polypeptide  
7 results in a staygreen phenotype in a plant.

1                   2. The polynucleotide of claim 1, wherein the polypeptide sequence is  
2 SEQ ID NO:2.

1                   3. The polynucleotide of claim 1, wherein the polypeptide sequence is  
2 SEQ ID NO:4.

1                   4. The polynucleotide of claim 1, wherein the polypeptide sequence is  
2 SEQ ID NO:6.

1                   5. The polynucleotide of claim 1, wherein the polynucleotide comprises  
2 the sequence of SEQ ID NO:1.

1                   6. The polynucleotide of claim 1, wherein the polynucleotide comprises  
2 the sequence of SEQ ID NO:3.

1                   7. The polynucleotide of claim 1, wherein the polynucleotide comprises  
2 the sequence of SEQ ID NO:5.

1                   8. A recombinant expression cassette comprising a promoter sequence  
2 operably linked to the polynucleotide of claim 1.

1                   9. A transgenic plant comprising the expression cassette of claim 8,  
2 wherein the plant has a staygreen phenotype.

1                   10. The transgenic plant of claim 9, wherein the plant is a cereal plant.

1                   11. The transgenic plant of claim 9, wherein the plant is maize.

1           12. A method of reducing ethylene sensitivity in a plant comprising the  
2 steps of:

3           (a) introducing an expression cassette comprising the isolated polynucleotide  
4 of claim 1 operably linked to a promoter; and  
5           (b) selecting a plant with reduced ethylene sensitivity.

1           13. A method of producing a staygreen phenotype in a plant, comprising  
2 the steps of:

3           (a) introducing an expression cassette comprising the isolated polynucleotide  
4 of claim 1 operably linked to a promoter; and  
5           (b) selecting a plant with staygreen characteristics.

1           14. The method of claim 13, wherein the plant is selected based on  
2 increased photosynthetic capacity.

1           15. The method of claim 13, wherein the plant is selected based on the  
2 phenotype of multiple embryos in a single seed.

1           16. The method of claim 13, wherein the plant is selected based on delayed  
2 senescence.

1           17. The method of claim 13, wherein the plant is selected based on the  
2 detection of a biomarker.

1           18. The method of claim 13, wherein the plant is a cereal plant.

19. The method of claim 13, wherein the plant is maize.

1/1




Figure 11

1/1




Figure 11