



US 20020123616A1

(19) **United States**

(12) **Patent Application Publication**

(10) **Pub. No.: US 2002/0123616 A1**

Lilja et al.

(43) **Pub. Date: Sep. 5, 2002**

(54) **PROSTATE SPECIFIC ANTIGEN (PSA)
-PROTEINASE INHIBITORS COMPLEXES**

(30) **Foreign Application Priority Data**

Jul. 23, 1990 (SE) 9002480-3

(76) Inventors: **Hans Lilja**, Skannor (SE); **Ulf-Hakan Stenman**, Grankulla (FI)

Publication Classification

Correspondence Address:

Stephen A. Bent
FOLEY & LARDNER
Washington Harbour
3000 K Street, N.W., Suite 500
Washington, DC 20007-5109 (US)

(51) **Int. Cl.⁷** **G01N 33/574; C07K 16/30**

(52) **U.S. Cl.** **530/387.7; 435/7.23**

(57) **ABSTRACT**

(21) Appl. No.: **09/905,998**

(22) Filed: **Jul. 17, 2001**

Related U.S. Application Data

(60) Continuation of application No. 09/264,176, filed on Mar. 5, 1999, now abandoned, which is a continuation of application No. 08/446,526, filed on May 19, 1995, now patented, which is a division of application No. 07/956,509, filed on Mar. 19, 1993, now patented, filed as 371 of international application No. PCT/FI91/00223, filed on Jul. 22, 1991.

According to the method of the invention, immunoassays are applied to measure free PSA as well as a proteinase inhibitor complex. Free PSA and PSA complex are according to the invention measured by a non-competitive immunoassay employing at least two different monoclonal antibodies. The invention is further characterized by that the PSA-proteinase inhibitor complex of interest is formed either with α_1 -antichymotrypsin, α_1 -protease inhibitor (API) or α_2 -macroglobulin. Moreover, the invention is characterized by that free PSA, the PSA-proteinase inhibitor complex and their ratio are applied in the diagnosis of patients with prostate cancer.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11

PROSTATE SPECIFIC ANTIGEN (PSA) -PROTEINASE INHIBITORS COMPLEXES

[0001] The present invention relates to an immunoassay of prostate-specific antigen (PSA), in which specific reagent materials (antibodies) are used that allow the measurement of free PSA as well as the PSA proteinase inhibitor complex.

[0002] It also relates to the use of free PSA and the PSA proteinase inhibitor complex and their ratio as a useful marker in diagnosis of patients with prostate cancer.

BACKGROUND OF THE INVENTION

[0003] The prostate specific antigen (PSA) was first purified from prostatic tissue (Wang et al. Invest Urol 1979), but the same protein was almost simultaneously and independently characterized in the seminal plasma (Hara et al. J Lab Clin Med 1989; Graves et al. N Engl J Med 1985). PSA is now known to be a 33-kDa glycosylated single chain serine protease (Lilja, J Clin Invest 1985; Watt et al. Proc Natl Acad Sci (USA) 1986). The 237 amino-acid polypeptide backbone has extensive similarities with that of the glandular kallikreins (Lundwall et al. FEBS Lett 1987; Schaller et al. Eur J Biochem 1987). Unlike the trypsin-like glandular kallikreins, which display Arg-restricted substrate specificity (MacDonald et al. Biochem J 1988), PSA displays chymotrypsin-like substrate specificity (Akiyama et al. FEBS Lett 1987; Christensson et al. Manuscript 1990; Lilja et al. J Biol Chem 1989). PSA has been predicted to be produced as a presumably inactive zymogen (Lundwall et al. FEBS Lett 1987). Active PSA is secreted into the seminal plasma (Lilja, J Clin Invest 1985) where it is one of the most abundant proteins of the prostate (Lilja et al. The Prostate 1988; Dubé et al. J Androl 1987). The biological activity of PSA in semen relates to its limited proteolytic fragmentation of the predominant proteins secreted by the seminal vesicles (Lilja, J Clin Invest 1985; Lilja et al. J Clin Invest 1987; McGee et al. Biol Reprod 1988).

[0004] Secondary to the release from the prostate epithelium PSA may also be detected in the circulation (Papsidero et al. Cancer Res 1980). Measurements of the serum concentration of PSA have now found widespread use in monitoring of patients with prostate cancer, although increased serum concentrations of PSA have also been reported in benign prostatic hyperplasia and secondary to surgical trauma of the prostate (Duffy, Ann Clin Biochem 1989; Brawer et al. Urology suppl 1989). However, it is presently unknown whether the immunoreactivity in serum represents the PSA-zymogen, the active PSA or PSA inactivated by extracellular proteinase inhibitors and contradictory results have been reported on the molecular mass of this immunoreactivity. Papsidero reported in 1980 the PSA-immunoreactivity to elute as a single 90 to 100 kDa peak (Papsidero et al. Cancer Res 1980), whereas Alfthan and Stenman reported the predominant part of this immunoreactivity to elute as a 30-kDa protein (Alfthan et al. Clin Chem 1988) when subjected to gel filtration chromatography.

[0005] In the proceeding invention we showed that PSA has the ability to form complexes with proteinase inhibitors that occur in high concentration in the human extracellular fluids and that PSA occurs in these fluids both in a free and complexed form. In addition, the invention proved to be very useful in diagnosis of prostate cancer patients.

SUMMARY OF THE INVENTION

[0006] According to the method of the invention, immunoassays are applied to measure free PSA as well as PSA as a proteinase inhibitor complex. Free PSA and the PSA complex are according to the invention measured by a non-competitive immunoassay employing at least two different monoclonal antibodies. The invention is further characterized by that the PSA proteinase inhibitor complex of interest is formed either with α_1 -antichymotrypsin, α_1 -protease inhibitor (API) or α_2 -macroglobulin. Moreover, the invention is characterized by that free PSA, the PSA-proteinase inhibitor complex and their ratio are applied in the diagnosis of patients with prostate cancer.

DESCRIPTION OF THE DRAWINGS

[0007] FIG. 1 presents a polyclonal antibody and three monoclonal antibodies used to probe proteins blotted onto PVDF-membranes after agarose gel electrophoresis. In lane 1 is 1 μ g PSA, in lane 2 is 1 μ g PSA incubated at 37° C. for 30 min with 6 μ g of α_1 -antichymotrypsin, and in lane 3 is 6 μ g of α_1 -antichymotrypsin.

[0008] FIG. 2 presents a polyclonal and three monoclonal antibodies used to probe proteins blotted onto PVDF-membranes after SDS-PAGE. In lane 1 is 1 μ g PSA, in lane 2 is 1 μ g PSA incubated at 37° C. for 30 min with 6 μ g of antichymotrypsin, and in lane 3 is 6 μ g of α_1 -antichymotrypsin.

[0009] FIG. 3 presents the specificity of the three assay versions.

[0010] FIG. 4 presents a correlation of PSA immunoreactivity in serum samples from 65 individual patients when analyzed with assay versions A and C.

[0011] FIG. 5 presents a correlation of PSA immunoreactivity in serum samples from 65 individual patients when analyzed with assay versions A and B.

[0012] FIG. 6 presents a gel filtration of a patient sample B on a TSK 250 HPLC column. The PSA immunoreactivity of the eluted fractions were analyzed by assay versions A, B and C.

[0013] FIG. 7 presents the a filtration of a patient sample C on a TSK 250 HPLC column. The PSA immunoreactivity of the eluted fractions were analyzed by assay versions A, B and C.

[0014] FIG. 8 presents a gel filtration of a patient sample D on a TSK 250 HPLC column. The PSA immunoreactivity of the eluted fractions were analyzed by assay versions A, B and C.

[0015] FIG. 9 presents a characterization of PSA immunoreactivity in a serum sample with a PSA level of 10000 μ g/l by gel filtration. PSA and the PSA-ACT complex were measured by IFMA.

[0016] FIG. 10 presents the proportion of the PSA-ACT complex of the total PSA immunoreactivity in sera of patients with prostatic cancer as a function of the PSA concentration. The level of PSA was measured by IRMA and that of PSA-ACT by IFMA.

[0017] FIG. 11 presents the concentration of the PSA-API complex measured by IFMA as a function of the PSA concentration measured by IRMA in sera from patients with prostatic cancer.

DETAILED DESCRIPTION

[0018] 1. Production and Characterization of Monoclonal Antibodies

[0019] Production of anti-PSA specific monoclonal antibodies Balb/c mice were immunized by intraperitoneous injections with 70 μg of PSA emulsified in equal volymes with Freund's complete adjuvant. The immunization was repeated after 3, 6 and 9 weeks with 50 μg of PSA emulsified with Freund's incomplete adjuvant. Three weeks later the mice were given a final booster with 40 μg of PSA and the mice were killed four days later. Lymphoid cells of the spleen were prepared and mixed in a 1:1 ratio with plasmacytoma cells (NS-1). The cells were fused and harvested in microtiter wells in KC-2000 (Hazleton Biologics Inc., Lenexa, USA) containing 200 g/L foetal calf serum and HAT-supplement H-0262 (1:50, Sigma) (Matikainen et al. J Gen Microbiol 1983).

[0020] Anti-PSA specific antibody production by the master clones was assayed with well strip plates coated with rabbit anti-mouse IgG (Lövgren et al. Talanta 1984). The strips were incubated with either the hybridoma supernatants or the standard (monoclonal antibody against PSA; 0812 Hybritech), washed, incubated with Eu-labelled PSA (50 ng per well), and the amount of bound Eu-labelled PSA was determined.

[0021] Cloning of the master clones by limited dilutions was performed as described (Staszewski and Yale, J Biol Med 1984). The desired clones were expanded intraperitoneally in Balb/c mice; the ascitic fluid being collected in 10 days. The IgG-fraction-of the ascitic fluid was purified by chromatography on protein A-Sepharose following the protocol recommended by the manufacturer.

[0022] Solid phase bound PSA was used to test if one unlabelled monoclonal antibody could block the binding of another Eu-labelled anti-PSA MAb to the solid-phase bound PSA. The solid-phase bound PSA was obtained by the incubation of 25 μL aliquots of purified PSA (25 $\mu\text{g}/\text{L}$) and 200 μL of Assay-buffer DELFIA[®] (50 mmol/L Tris, pH 7.75, 0.15 mol/L NaCl, 0.5 g/L BSA, and 0.5 g/L NaN_3) for 2 h in well strip plates coated with the 2E9 or the 5A10 anti-PSA MAb. The strips were washed and then incubated for 1 h with 200 μL of one unlabelled anti-PSA MAb (0.005-50 $\mu\text{g}/\text{L}$). Again, the strips were washed, incubated for 1 h with another Eu-labelled anti-PSA MAb and the amount of bound Eu-labelled anti-PSA was determined.

[0023] Partial Characterization of the Epitope Specificity of Three Monoclonal Antibodies Against PSA

[0024] Several clones produced monoclonal antibodies against PSA as indicated by fluorometric assay. Three of these (designated 2E9, 2H11 and 5A10) were expanded and the antibodies isolated from the ascitic fluid. The three monoclonal antibodies against PSA were used to probe proteins blotted onto PVDF-membranes after agarose gel electrophoresis (FIG. 1) or SDS-PAGE (FIG. 2) of 1 μg of PSA (lane 1); 1 μg of PSA incubated at 37° C. for 30 min with 6 μg of α_1 -antichymotrypsin (lane 2); and 6 μg of α_1 -antichymotrypsin (lane 3). PSA blotted to PVDF-membranes from the agarose gels was identified by all three monoclonal antibodies whereas the PSA complexed to α_1 -antichymotrypsin was identified by the 2E9 and the 2H11 antibodies but not by the 5A10 antibody. The 2E9 antibody

was the only anti-PSA MY that readily identified PSA and PSA complexed to α_1 -antichymotrypsin when these proteins were blotted onto PVDF-membranes after SDS-PAGE. However, a minute reaction was also obtained with the PSA (but not with the PEA complexed to α_1 -antichymotrypsin) when the 2H11 and the 5A10 antibodies were used to probe these proteins blotted onto PVDF-membranes after SDS-PAGE.

[0025] The epitope specificity of the three monoclonal antibodies was also characterized using three different sets of solid-phase sandwich assays. Thereby assay (A), where the 2E9 antibody was used as solid-phase catcher and Eu-labelled 2H11 was used as detecting antibody, displayed an almost identical dose-response for PSA as compared with PSA complexed to α_1 -antichymotrypsin (Table 1; FIG. 3). This contrasts with both assay (B), where the 5A10 antibody was used as catcher and Eu-labelled 2H11 was used as detecting antibody, which preferentially recognized PSA but only poorly recognized PSA complexed to α_1 -antichymotrypsin, and with assay (C), where the 2E9 antibody was used as catcher and Eu-labelled antibody against α_1 -antichymotrypsin was used as detecting antibody, which only recognized PSA complexed to α_1 -antichymotrypsin (Table 1; FIG. 3).

[0026] Solid-phase bound PSA was used to further characterize the epitope specificity of the three monoclonal antibodies against PSA; the solid-phase binding of PSA having been achieved by the use of well strip plates coated with the 2E9 or the 5A10 antibody. It was thereby found that none of the anti-PSA MAb's 2E9, 2H11 or the 5A10 significantly blocked the binding of each other when we tested the ability of one anti-PSA MAb to block the binding of another Eu-labelled anti-PSA MAb to the solid-phase bound PSA.

[0027] 2. The Occurrence of PSA-Proteinase Inhibitor Complexes in Human Serum**[0028]** Analysis of PSA in Human Serum

[0029] Serum from individual patient samples (n=65) were analyzed with the three different sets of assays (A, B and C). Regression analysis of the results obtained with assay A and assay C gave $y=0.89x+6.55$, $r=0.97$ (FIG. 4); and the regression analysis between assay A and assay B gave $y=0.10x+9.56$, $r=0.82$ (FIG. 5).

[0030] The total recovery of the immunoreactivity from the gel filtration experiments of patient samples on the TSK 250 HPLC column was equally high (82 to 107%) with all three assay procedures used (A, B and C). The gel filtration experiments of patient samples on the TSK 250 HPLC column showed that the predominant peak of PSA-immunoreactivity, when analyzed with assay A, was identified in fractions eluting at a position corresponding to a molecular mass of 80 to 90 kDa while a minor peak of this immunoreactivity was found in fractions eluting at a position corresponding to a molecular mass of 25 to 40 kDa (FIGS. 6-8). In much the same way, the analysis of the fractions eluted with assay C (specific for PSA complexed to α_1 -antichymotrypsin) identified one predominant immunoreactive peak in the range 80 to 90 kDa (FIGS. 6-8). However, when assay B was used to analyze the fractions eluted from the gel filtration experiments the predominant immunoreactive peak eluted at a position corresponding to mass of 25 to 40 kDa.

The elution position of this peak corresponded to the minor immunoreactive peak Identified with assay A (FIGS. 6-8).

[0031] When serum samples from men with various levels of PSA (10-10000 $\mu\text{g/L}$) were fractionated by gel filtration, two components corresponding to PSA and PSA-ACT were also observed. In samples with high PSA-levels the PSA-ACT complex dominated (FIG. 9). In female sera these components were not seen (not shown). The proportion of PSA-ACT of total PSA immunoreactivity increased with increasing PSA levels (FIG. 10). In sera from healthy males with PSA levels below 2.8 $\mu\text{g/L}$ the proportion of PSA-ACT was 23-47%, in samples with PSA levels of 2.8-10 $\mu\text{g/L}$ the proportion was 26-86% and in samples with higher levels the proportion increased further being 70-100% at PSA levels over 1000 $\mu\text{g/L}$ (FIG. 10).

[0032] On the basis of the concentrations of the complexes expressed in arbitrary units the main complex of PSA in sera was PSA-ACT complex. In samples with low PSA levels the concentration of both PSA-API and PSA-ACT were close to the detection limit. Therefore it was not possible to calculate the proportion of these complexes in normal samples. Clearly elevated levels of PSA-API complex occurred in samples with PSA levels over 40 $\mu\text{g/L}$ and the levels tended to increase with increasing levels of PSA (FIG. 11).

[0033] 3. PSA and PSA- α_1 -antichymotrypsin Complexes in the Diagnosis of Patients With Prostate Cancer

[0034] The three assay versions referred to under section "Characterization of the epitope specificity of three monoclonal antibodies against PSA" were used to test 144 patients with benign prostatic hyperplasia (BPH) and 122 patients with different stages of prostate cancer (CAP). The ratios between A: PSA complexed with α_1 -antichymotrypsin/PSA total and B: PSA free non-complexed/PSA total were calculated as well as the clinical sensitivity and specificity for the measurement of total PSA and PSA α_1 -antichymotrypsin alone (Table 2). It is obvious from the presented data that increased clinical specificity is achieved by measuring the PSA α_1 -antichymotrypsin complex and that the ratios between PSA free/PSA total and PSA free/PSA complexed with α_1 -antichymotrypsin are significantly different between BPH and CAP patients.

[0035] Table 1

[0036] The table 1 presents a dose-response of purified PSA and PSA complexed to α_1 -antichymotrypsin when analyzed by three different sets of assays.

[0037] The assay A is 2E9 anti-PSA MAb as solid phase catcher and Eu-labelled 2H11 anti-PSA MAB as detecting antibody.

[0038] The assay B is 5A10 anti-PSA MAB as solid phase catcher and Eu-labelled 2H11 anti-PSA MAB as detecting antibody.

[0039] The assay C is 2E9 anti-PSA MAB as solid phase catcher and Eu-labelled rabbit antibody against α_1 -antichymotrypsin as detecting antibody.

[0040] The columns 1 indicate the purified PSA and columns 2 indicate the PSA complexed to α_1 -antichymotrypsin.

[0041] Tables 2a and 2b

[0042] The tables 2a and 2b present the results of the testing of the patient samples with three assay versions for free, complexed and total PSA. In the table BPH indicates benign prostatic hyperplasia, CAP indicates prostate cancer, G indicates the differentiation grade and T indicates the grade. The table 2b presents the sensitivity and the specificity.

TABLE 1

PSA $\mu\text{g/L}$	PSA assay					
	A		B		C	
	1	2	1	2	1	2
1	6664	5250	6733	2119	435	4208
5	26897	23535	31487	3179	487	15662
10	53452	41064	65146	4573	559	30283
100	534860	460464	600057	33006	2105	267223
500	2231640	1826790	2631640	156712	12073	726596

1. Purified PSA;

2. PSA complexed to α_1 -antichymotrypsin.

[0043]

TABLE 2a

		Correlation coefficient	Ratio mean
BPH (n = 144)	A. PSA c/PSA tot	0.932	0.970
	B. PSA f/PSA tot	0.853	0.302
CAP (n = 122)	A.	0.994	1.219
	B.	0.784	0.191
CAP, G1 (n = 31)	A.	0.994	1.628
	B.	0.922	0.190
CAP, G2 (n = 47)	A.	0.972	1.141
	B.	0.956	0.169
CAP, G3 (n = 43)	A.	0.996	1.014
	B.	0.818	0.218
CAP T1-2 (n = 56)	A.	0.985	1.044
	B.	0.868	0.178
CAP T3-4 (n = 65)	A.	0.993	1.372
	B.	0.770	0.204
CAP T4 (n = 25)	A.	0.997	1.174
(not treated)	B.	0.825	0.188
BPH (n = 84) PSA \leq 5	A.	0.879	1.059
	B.	0.850	0.301
BPH (n = 60) PSA > 5	A.	0.888	0.846
	B.	0.735	0.303
CAP (n = 26) PSA \leq 5	A.	0.913	1.773
	B.	0.826	0.202
CAP (n = 94) PSA > 5	A.	0.993	1.065
	B.	0.778	0.188
CAP (n = 25)	A.	0.919	1.025
PSA > 5 \leq 20	B.	0.502	0.187
CAP (n = 69)	A.	0.993	1.080
PSA > 20	B.	0.770	0.184

[0044]

TABLE 2b

Sensitivity and specificity				
PSA tot	Sensitivity	PSA tot	>5	95/121 = 0.785
			>10	80/121 = 0.661
	Specificity	PSA tot	<5	84/144 = 0.583
			<10	116/144 = 0.806

TABLE 2b-continued

Sensitivity and specificity				
PSA c	Sensitivity	PSA c	≥5	93/121 = 0.769
			>10	81/121 = 0.669
	Specificity	PSA c	<5	92/144 = 0.639
			<10	124/144 = 0.861

1. An immunoassay of prostate-specific antigen (PSA) characterized by that in the immunoassay of prostate-specific antigen (PSA)

an amount of prostate-specific antigen complexed with a proteinase inhibitor and/or

an amount of free non-complexed prostate-specific antigen and/or

an amount of the total prostate-specific antigen are measured:

2. An immunoassay according to claim 1 characterized by that

the PSA-proteinase inhibitor complex (complexed PSA) and/or

the free non-complexed prostate-specific antigen (free PSA) and/or

the total prostate-specific antigen (total PSA) are measured by a non-competitive immunoassay.

3. An immunoassay according to claim 2 characterized by that

the PSA-proteinase inhibitor complex (complexed PSA) and/or

the free non-complexed prostate-specific antigen (free PSA) and/or

the total prostate-specific antigen (total PSA) are measured by a non-competitive immunoassay employing at least two different monoclonal antibodies.

4. An immunoassay according to claim 3 characterized by that the monoclonal antibodies used for the recognition of the PSA-proteinase inhibitor complex (complexed PSA) bind either to

prostate-specific antigen (PSA), or to

the PSA-proteinase inhibitor complex (complexed PSA) or to

the proteinase inhibitor.

5. An immunoassay according to claim 4 characterised by that the proteinase inhibitor is a₁-antichymotrypsin.

6. An immunoassay according to claim 4 characterized by that the proteinase inhibitor is o-proteinase inhibitor.

7. An immunoassay according to claim 4 characterized by that the proteinase inhibitor is macroglobulin.

8. An immunoassay according to claim 2 characterized by that the ratio between the free non-complexed prostate-specific antigen (free PSA) and the PSA-proteinase inhibitor complex (complexed PSA) is determined.

9. An immunoassay according to claim 2 characterized by that the ratio between the free non-complexed prostate-specific antigen (free PSA) and the total prostate-specific antigen (total PSA) is determined.

10. An immunoassay according to claim 2 characterized by that the ratio between the PSA-proteinase inhibitor complex (complexed PSA) and the total prostate-specific antigen (total PSA) is determined.

* * * * *