

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 June 2002 (20.06.2002)

PCT

(10) International Publication Number
WO 02/48298 A1

(51) International Patent Classification⁷: **C11D 3/00**,
3/20, 3/48

(21) International Application Number: PCT/EP01/14356

(22) International Filing Date: 6 December 2001 (06.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
00811192.4 14 December 2000 (14.12.2000) EP

(71) Applicant (for all designated States except US): **CIBA
SPECIALTY CHEMICALS HOLDING INC.** [CH/CH];
Klybeckstrasse 141, CH-4057 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **OCHS, Dietmar**
[DE/DE]; Webergasse 11g, 79650 Schopfheim (DE).
BRUGGER, François [FR/FR]; 15B, ru de l'Avenir,
F-68510 Waltenheim (FR). **SCHNYDER, Marcel**
[CH/CH]; Schulstrasse 11/4, CH-4127 Birsfelden (CH).

(74) Common Representative: **CIBA SPECIALTY CHEMI-
ICALS HOLDING INC.**; Patentabteilung, Klybeckstrasse
141, CH-4057 Basel (CH).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,
YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: SURFACE-ACTIVE COMPOSITIONS

(57) Abstract: A surface-active composition is described which comprises (a) from 0.01 to 90 % by weight of a mixture of a mi-
crobicidal active ingredient of (a₁) a diphenyl ether compound and (a₂) a phenol derivative, (b) from 0 to 50 % by weight of one
or more hydrotropic agents, (c) from 0 to 80 % by weight of one or more synthetic detergents or of a soap or of combinations of
the mentioned substances and/or of a salt of a saturated and/or unsaturated C₈-C₂₂ fatty acid, (d) from 0 to 50 % by weight of an
alcohol, (e) from 0 to 50 % by weight of typical ingredients for cleaning and disinfectant compositions and optionally (f) tap water
or deionised water ad 10 %. The compositions are used for the antimicrobial treatment of human skin, and hair, of hard surfaces and
of textile fibre materials.



WO 02/48298 A1

Surface-active compositions

The present invention relates to surface-active compositions and to the use of such compositions for the antimicrobial treatment of human skin and hair and for the treatment of hard surfaces and textile fibre materials.

Cleaning and disinfectant compositions comprising antimicrobial active ingredients, e.g. personal care preparations, hand and machine dishwashing formulations, cleaning and disinfecting formulations for hard surfaces and liquid and solid textile washing formulations, are becoming ever more widespread. Phenol derivatives and diphenyl ether compounds are known as antibacterial active ingredients.

It has now been found, surprisingly, that a combination of diphenyl ether compounds and phenol derivatives exhibits strong bactericidal effects.

The present invention accordingly relates to a surface-active composition comprising


- (a) from 0.01 to 90 % by weight of a mixture of a microbicidal active ingredient of
 - (a₁) a diphenyl ether compound and
 - (a₂) a phenol derivative,
- (b) from 0 to 50 % by weight of one or more hydrotropic agents,
- (c) from 0 to 80 % by weight of one or more synthetic detergents or of a soap or of combinations of the mentioned substances and/or of a salt of a saturated and/or unsaturated C₈-C₂₂ fatty acid,
- (d) from 0 to 50 % by weight of an alcohol,
- (e) from 0 to 50 % by weight of typical ingredients for cleaning and disinfectant compositions and optionally
- (f) tap water or deionised water ad 100 %.

Preferably, the present invention accordingly relates to a surface-active composition comprising

- (a) from 0.01 to 10 % by weight of a mixture of a microbicidal active ingredient of

- (a₁) a diphenyl ether compound and
- (a₂) a phenol derivative,
- (b) from 0 to 50 % by weight of one or more hydrotropic agents,
- (c) from 5 to 80 % by weight of one or more synthetic detergents or of a soap or of combinations of the mentioned substances and/or of a salt of a saturated and/or unsaturated C₈-C₂₂ fatty acid,
- (d) from 0 to 50 % by weight of an alcohol, and optionally
- (f) tap water or deionised water ad 100 %.

The composition according to the invention preferably comprises as component (a₁) a hydroxy-diphenyl ether of formula



Y is chlorine or bromine,

Z is SO₂H, NO₂ or C₁-C₄alkyl,

r is from 0 to 3,


o is from 0 to 3,

p is 0, 1 or 2,

m is 1 or 2 and

n is 0 or 1,

and more especially a compound of




Y is chlorine and


r is 1 or 2.

Very special preference is given to a compound of formula

- 3 -



As component (a₁) there can also be used a non-halogenated hydroxydiphenyl ether of formula



U₁ and U₂ are each independently of the other hydrogen, hydroxy, unsubstituted or hydroxy-substituted C₁-C₂₀alkyl, C₅-C₇cycloalkyl, C₁-C₆alkylcarbonyl, C₁-C₂₀alkoxy, phenyl or phenyl-C₁-C₃alkyl;

U₃ is hydrogen, C₁-C₂₀alkyl, C₁-C₂₀alkoxy or C₁-C₆alkylcarbonyl; and

U₄ is hydrogen, unsubstituted or hydroxy-substituted C₁-C₂₀alkyl, C₅-C₇cycloalkyl, hydroxy, formyl, acetonyl, C₁-C₆alkylcarbonyl, C₂-C₂₀alkenyl, carboxy, carboxy-C₁-C₃alkyl, C₁-C₃alkylcarbonyl-C₁-C₃alkyl or carboxyallyl.

U₁, U₂, U₃ and U₄ having the meaning of C₁-C₂₀alkyl are straight-chain or branched alkyl radicals, for example methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, isopentyl, tert-pentyl, hexyl, cyclohexyl, heptyl, octyl, isooctyl, nonyl, decyl and the like.

U₁, U₂ and U₃ as C₁-C₂₀alkoxy are straight-chain or branched alkoxy radicals, for example methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-butoxy, pentyloxy, isopentyloxy, tert-pentyloxy, heptyloxy, octyloxy, isooctyloxy, nonyloxy, decyloxy and the like.

U₁, U₂, U₃ and U₄ having the meaning of C₁-C₆alkylcarbonyl are straight-chain or branched carbonyl radicals, for example acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl and the like.

U_1 , U_2 and U_4 having the meaning of hydroxy-substituted C_1 - C_{20} alkyl are, for example, hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, hydroxypentyl, hydroxyhexyl, hydroxyheptyl, hydroxyoctyl, hydroxynonyl, hydroxydecyl and the like.


According to the invention, preference is given to the use of compounds of formula (1') wherein OH is in the meta- or para-position relative to the ether bond.

Preferably, U_1 and U_2 are each independently of the other hydrogen, C_1 - C_{20} alkyl, C_1 - C_6 alkylcarbonyl or C_1 - C_{20} alkoxy.


U_3 is preferably hydrogen, C_1 - C_{20} alkyl or C_1 - C_{20} alkoxy.

U_4 is preferably hydrogen, C_1 - C_{20} alkyl, hydroxy, formyl, acetonyl, allyl, carboxymethyl, carboxyallyl, hydroxy-substituted C_1 - C_{20} alkyl or C_1 - C_6 alkylcarbonyl.


Compounds of formula (1') that are of special interest are:




Other compounds of formula (1') that are of special interest are:




U_1 is C_1 - C_3 alkyl, for example the compound of formula




Also of interest are compounds of formula




U_4 is C_1 - C_3 alkyl, for example the compound of formula




The following compounds are of special interest:





The compounds of formula (1') are known or can be prepared using methods analogous to those known.

Compounds suitable as component (a₂) are preferably those selected from phenol derivatives of formula



- R₁ is hydrogen, hydroxy, C₁-C₂₀alkyl, chloro, phenyl, benzyl or nitro,
- R₂ is hydrogen, hydroxy, C₁-C₂₀alkyl or halogen,
- R₃ is hydrogen, C₁-C₂₀alkyl, hydroxy or chloro,
- R₄ is hydrogen or methyl, and
- R₅ is hydrogen or nitro.

Exemplary compounds are chlorophenols (o-, m-, p-chlorophenols), 2,4-dichlorophenol, p-nitrophenol, xlenol, p-chloro-m-xlenol, cresols (o-, m-, p-cresols), p-chloro-m-cresol, pyrocatechol, resorcinol, orcinol, 4-n-hexylresorcinol, pyrogallol, phloroglucinol, carvacrol, thymol, p-chlorothymol, o-phenylphenol, o-benzylphenol and p-chloro-o-benzylphenol.

Further exemplary representatives of component (a₂) are chlorhexidines, for example 1,1'-hexamethylene-bis(5-(p-chlorophenyl)biguanide), together with organic and inorganic acids and chlorhexidine derivatives, such as their diacetates, digluconates or dihydrochloride compounds.

Further exemplary phenol derivatives are 1-phenoxypropan-2-ol and 3-(4-chlorophenoxy)-1,2-propanediol.

Very special preference is given to the use of o-phenylphenol as component (a₂).

In the composition according to the invention, the combination of (a₁) the compound of formula (2) or (3) and (a₂) o-phenylphenol is especially used.

The following compounds are suitable as component (b):

- sulfonates of terpenoids, or of mono- or di-nuclear aromatic compounds, for example sulfonates of camphor, toluene, xylene, cumene or of naphthol;
- saturated or unsaturated C₃-C₁₂ di- or poly-carboxylic acids, for example malonic, succinic, glutaric, adipic, pimelic, suberic, azelaic and sebacic acid, undecane- and dodecane-dioic acid, fumaric, maleic, tartaric and malic acid, and citric and aconitic acid;
- aminocarboxylic acids, such as ethylenediaminetetraacetic acid, hydroxyethylethylenediaminetetraacetic acid and nitrilotriacetic acid;
- cycloaliphatic carboxylic acids, such as camphoric acid;
- aromatic carboxylic acids, such as benzoic, phenylacetic, phenoxyacetic and cinnamic acid, 2-, 3- and 4-hydroxybenzoic acid, anilic acid, and o-, m- and p-chlorophenylacetic acid and o-, m- and p-chlorophenoxyacetic acid;
- isethionic acid;
- tannic acid;

- acid amides of formula
$$R_1-CO-N \begin{array}{l} \nearrow R_2 \\ \searrow R_3 \end{array}$$
 wherein

R₁ is hydrogen or C₁-C₁₂alkyl and

R₂ and R₃ are each independently of the other hydrogen, C₁-C₁₂alkyl, C₂-C₁₂alkenyl, C₂-C₁₂hydroxyalkenyl, C₂-C₁₂hydroxyalkyl or a polyglycol ether chain having from 1 to 30 -CH₂-CH₂-O- or -CHY₁-CHY₂-O- groupings, wherein one of the radicals Y₁ and Y₂ is hydrogen and the other is methyl, such as N-methylacetamide;

- urea derivatives of formula
$$\begin{array}{c} R_1 \\ \diagdown \\ N-CO-N \\ \diagup \quad \diagdown \\ R_2 \quad R_3 \\ \quad \quad R_4 \end{array}$$
 wherein

R_1 , R_2 , R_3 and R_4 are each independently of the others hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_1 - C_8 hydroxyalkyl or C_2 - C_8 hydroxyalkenyl.

All the organic acids mentioned under (b) can also be in the form of their water-soluble salts, such as the alkali metal salts, especially the sodium or potassium salts, or the amine ($NR_1R_2R_3$) salts wherein

R_1 , R_2 and R_3 are each independently of the others hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_1 - C_8 -hydroxyalkyl, C_5 - C_8 cycloalkyl or polyalkyleneoxy- C_1 - C_{18} alkyl or

R_1 , R_2 and R_3 , together with the nitrogen atom to which they are bonded, are unsubstituted or C_1 - C_4 alkyl-substituted morpholino.

Component (b) can consist of a single compound or a plurality of different compounds.

Very special preference is given to a combination of cumenesulfonate and citric acid monohydrate.

As component (c), anionic, nonionic, or zwitterionic and amphoteric synthetic detergents are suitable.

Suitable anionic detergents are

- sulfates, for example fatty alcohol sulfates, the alkyl chain of which has from 8 to 18 carbon atoms, for example sulfated lauryl alcohol;
- fatty alcohol ether sulfates, for example the acid esters or salts thereof of a polyaddition product of from 2 to 30 mol of ethylene oxide and 1 mol of a C_8 - C_{22} fatty alcohol;
- the alkali metal, ammonium or amine salts, referred to as soaps, of C_8 - C_{20} fatty acids, for example coconut fatty acid;
- alkylamide sulfates;
- alkylamine sulfates, for example monoethanolamine lauryl sulfate;

- alkylamide ether sulfates;
- alkylaryl polyether sulfates;
- monoglyceride sulfates;
- alkanesulfonates, the alkyl chain of which contains from 8 to 20 carbon atoms, for example dodecyl sulfonate;
- alkylamide sulfonates;
- alkylaryl sulfonates;
- α -olefin sulfonates;
- sulfosuccinic acid derivatives, for example alkyl sulfosuccinates, alkyl ether sulfosuccinates or alkylsulfosuccinamide derivatives;

- N-[alkylamidoalkyl]amino acids of formula $\text{CH}_3(\text{CH}_2)_n\text{-CO-N} \begin{array}{c} \text{Y} \\ \text{CH-Z-COO}^-\text{M}^+ \\ \text{X} \end{array}$ wherein

X is hydrogen, $\text{C}_1\text{-C}_4$ alkyl or $\text{-COO}^-\text{M}^+$,

Y is hydrogen or $\text{C}_1\text{-C}_4$ alkyl,


Z is $\text{-(CH}_2\text{)}_{m_1-1}$

m_1 is from 1 to 5,

n_1 is an integer from 6 to 18 and

M is an alkali metal cation or amine cation,

- alkyl and alkylaryl ether carboxylates of formula (13) $\text{CH}_3\text{-X-Y-A}$ wherein

X is a radical of formula $\text{-(CH}_2\text{)}_{5-19}\text{-O-}$, $\text{-(CH}_2\text{)}_{5-11}\text{-}$  -O- or $\text{-(CH}_2\text{)}_{5-19}\text{-N} \begin{array}{c} \text{R} \\ \text{ } \end{array}$,

R is hydrogen or $\text{C}_1\text{-C}_4$ alkyl,

Y is -(CHCHO)_{1-50} ,

A is $\text{(CH}_2\text{)}_{m_2-1}\text{-COO}^-\text{M}^+$ or $\begin{array}{c} \text{O} \\ \parallel \\ \text{---P---O}^-\text{M}^+ \\ | \\ \text{O}^-\text{M}^+ \end{array}$

m_2 is from 1 to 6 and

M is an alkali metal cation or amine cation.

Also used as anionic surfactants are fatty acid methyl taurides, alkyl isothionates, fatty acid polypeptide condensation products and fatty alcohol phosphoric acid esters. The alkyl radicals occurring in those compounds preferably have from 8 to 24 carbon atoms.

The anionic surfactants are generally in the form of their water-soluble salts, such as the alkali metal, ammonium or amine salts. Examples of such salts include lithium, sodium, potassium, ammonium, triethylamine, ethanolamine, diethanolamine and triethanolamine salts. The sodium, potassium or ammonium ($\text{NR}_1\text{R}_2\text{R}_3$) salts, especially, are used, with R_1 , R_2 and R_3 each independently of the others being hydrogen, C_1 - C_4 alkyl or C_1 - C_4 hydroxyalkyl.

Especially preferred anionic surfactants in the composition according to the invention are monoethanolamine lauryl sulfate or the alkali metal salts of fatty alcohol sulfates, especially sodium lauryl sulfate and the reaction product of from 2 to 4 mol of ethylene oxide and sodium lauryl ether sulfate.

Suitable zwitterionic and amphoteric surfactants include C_8 - C_{18} betaines, C_8 - C_{18} sulfobetaines, C_8 - C_{24} alkylamido- C_1 - C_4 alkylenebetaines, imidazoline carboxylates, alkylamphocarboxy-carboxylic acids, alkylamphocarboxylic acids (e.g. lauroamphoglycinate) and N-alkyl- β -aminopropionates or -iminodipropionates, with preference being given to C_{10} - C_{20} alkylamido- C_1 - C_4 alkylenebetaines and especially to coconut fatty acid amide propylbetaine.

Nonionic surfactants that may be mentioned include, for example, derivatives of the adducts of propylene oxide/ethylene oxide having a molecular weight of from 1000 to 15 000, fatty alcohol ethoxylates (1-50 EO), alkylphenol polyglycol ethers (1-50 EO), polyglucosides, ethoxylated hydrocarbons, fatty acid glycol partial esters, for example diethylene glycol monostearate, fatty acid alkanolamides and dialkanolamides, fatty acid alkanolamide ethoxylates and fatty amine oxides.

As component (c) there may also be used the salts of saturated and unsaturated C_8 - C_{22} fatty acids either alone or in the form of a mixture with one another or in the form of a mixture with other detergents mentioned as component (c). Examples of such fatty acids include, for example, capric, lauric, myristic, palmitic, stearic, arachidic, behenic, caproic, dodecenoic, tetradecenoic, octadecenoic, oleic, eicosenoic and erucic acid, and the commercial mixtures of such acids, such as, for example, coconut fatty acid. Such acids are present in the form of salts, there coming into consideration as cations alkali metal cations, such as sodium and potassium cations, metal atoms, such as zinc and aluminium atoms, and nitrogen-containing organic compounds of sufficient alkalinity, such as amines and ethoxylated amines. Such salts may also be prepared *in situ*.

As component (d) there come into consideration as dihydric alcohols especially those compounds having from 2 to 6 carbon atoms in the alkylene moiety, such as ethylene glycol, 1,2- or 1,3-propanediol, 1,3-, 1,4- or 2,3-butanediol, 1,5-pentanediol and 1,6-hexanediol.

Preference is given to 1,2-propanediol (propylene glycol).

Preferred monohydric alcohols are ethanol, n-propanol and isopropanol and mixtures of those alcohols.

The composition according to the invention comprises, as component (e), builders (zeolites/layered silicates), bleaching agents or bleaching systems (perborate/percarbonate plus TAED), fluorescent whitening agents and enzymes.

Furthermore, the washing composition can comprise enzymes, enzyme stabilisers, thickeners, sequestering agents, for example EDTA or phosphoric acid salts, corrosion inhibitors, colourants, perfumes, fluorescent whitening agents, buffer compounds or the like.

Compositions according to the invention can be prepared by mixing components (a) and optionally (b), (c), (d) and (e) in any desired order with the requisite amount of deionised water and stirring the batch until homogeneous. The composition is made up to 100 % with tap water or deionised water. The procedure is purely physical. No chemical reaction takes place between the individual components.

Cleaning and disinfecting formulations according to the present invention may further comprise thickening agents, sequestering agents, antioxidants, UV absorbers, dyes, perfumes, buffer compounds, vitamins, moisturizers, body care substances, solids like waxes etc..

The formulations according to the invention exhibit strong bactericidal activity in two respects:

- rapid destruction of germs present.

This can be demonstrated, for example, by a suspension test, e.g. according to test method EN 1276.

- long-term bactericidal activity on the treated surface, as a result of which recolonisation is prevented effectively.

This can be demonstrated, for example, by the AATCC 100-1993 method.

They are therefore suitable for disinfecting and cleaning human skin and hands, hard articles and textile fibre materials and can be applied thereto in dilute or undiluted form, an amount of at least 2 ml, preferably in the undiluted form, being suitable for disinfection of the hands.

The compositions according to the invention are very especially used in washing and cleaning formulations, for example in household washing formulations, powder washing formulations, washing pastes, fabric softeners, solid soaps, dishwashing formulations, all-purpose cleaners, especially in liquid washing formulations for textile fibre materials.

The invention accordingly relates also to a method for the antimicrobial treatment of textile fibre materials in washing liquor, which method comprises treating the textile fibre materials in the washing liquor with a composition comprising

- (a) a microbicidal active ingredient of
 - (a₁) from 0 to 5 % by weight of a diphenyl ether compound and
 - (a₂) from 0.1 to 5 % by weight of a phenol derivative,
- (b) from 0 to 50 % by weight of one or more hydrotropic agents,
- (c) from 5 to 80 % by weight of one or more synthetic detergents or of a soap or of combinations of the mentioned substances and/or of a salt of a saturated and/or unsaturated C₈-C₂₂ fatty acid,
- (f) from 0 to 50 % by weight of an alcohol, and optionally
- (e) from 0 to 50 % by weight of typical ingredients for cleaning and disinfectant compositions and optionally
- (f) tap water or deionised water ad 100 %.

In the method according to the invention preference is given to a washing liquor that is free of diphenyl ether compounds, that is to say contains no component (a₁).

The invention relates also to a method for imparting antimicrobial properties to textile fibre materials, which method comprises treating the textile fibre materials in the washing liquor with a composition comprising

- (a) from 0.01 to 10 % by weight of a mixture of a microbicidal active ingredient of
 - (a₁) a diphenyl ether compound and
 - (a₂) a phenol derivative,
- (b) from 0 to 50 % by weight of one or more hydrotropic agents,
- (c) from 5 to 80 % by weight of one or more synthetic detergents or of a soap or of combinations of the mentioned substances and/or of a salt of a saturated and/or unsaturated C₈-C₂₂ fatty acid,
- (g) from 0 to 50 % by weight of an alcohol,
- (h) from 0 to 50 % by weight of typical ingredients for cleaning and disinfectant compositions and optionally and optionally

- (i) tap water or deionised water ad 100 %, at least a fraction of the antimicrobial active ingredient remaining on the textile fibre material.

The textile materials that can be treated in accordance with the invention are undyed or dyed or printed, natural or synthetic fibre materials, for example of silk, wool, polyamide or polyurethanes, and especially cellulosic fibre materials of all kinds. Such fibre materials are, for example, natural cellulose fibres, such as cotton, linen, jute and hemp, as well as cellulose and regenerated cellulose. Preferred suitable textile fibre materials are of cotton.

Using the composition according to the invention it is possible to destroy bacteria present on the washing material in the dilute liquor during the washing procedure. At the same time, antimicrobial properties are imparted to the washed textile material, that is to say bacteria that get on the textile material while it is being worn are destroyed.

The following Examples illustrate the invention. Percentages and parts are percentages by weight and parts by weight, respectively.

Examples 1: Preparation of a liquid washing formulations (1) – (5)

Liquid formulations having the following compositions are prepared:

Formulation	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>
combination of 30 % of the compound of formula (3) and 70 % of propylene glycol	0.6	0.6	0.6	-	-
o-phenylphenol	0.5	1	1	1	2
sodium dodecylbenzenesulfonate	6	6	6	6	6
sodium lauryl sulfate	8	8	8	8	8
Pareth 45-7 (Dobanol 45-7)	4	4	4	4	4
ethanol	9	9	9	9	9
sodium cumenesulfonate	5	-	5	5	-
soap noodles (Mettler)	5	7	7	5	7
trisodium citrate dihydrate	2	2	2	2	2
triethanolamine	5	5	5	5	5
fluorescent whitening agents	0.3	0.3	0.3	0.3	0.3
water to	100	100	100	100	100

Example 2: Determination of the bactericidal efficacy of formulations (1) to (5) in accordance with EN 1276 (concentration 80 %, contact time 5 minutes) in log reduction

Test principle:

1.0 ml of a bacterial suspension is added to 8.0 ml of the formulation in question (the test concentration is multiplied by a factor of 1.25) and to 1.0 ml of a suspension of 0.3 % (factor 10) bovine albumin and mixed vigorously. After the contact time (see above) at 21°C (+/- 1°C), a 0.1 ml sample is removed and added to 50 ml of TSB + inactivator (= test neutralisation mixture, 10⁰). 500 µl of the neutralisation mixture are added to 9 ml of TSB + inactivator to give a 10⁻² dilution. Each test neutralisation mixture and the dilutions are filtered over a membrane and washed with 150 ml of distilled water. The membranes are

incubated for 48 hours on the surface of agar plates. After incubation, the colonies are counted and listed in a Table, and the log reduction is calculated.

The results are given in Table 1.

<u>Table 1</u>					
Formulation	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>
S.a. ATCC 6538	>5	> 5	4.6	4.5	> 5
E.c. ATCC 10536	>5	> 5	4.5	3.8	> 5
E.h. ATCC 10541	>5	> 5	>5	>5	> 5
Ps.a. ATCC 15442	3	4.3	4	3.9	> 5

The results in Table 1 show that good bactericidal effects can be achieved on the textile material using the formulations according to the invention.

Example 3: Determination of the bactericidal effects, during wear (from 0 to 24 hours), on textile material washed under standard conditions

Textile material: cotton

washing formulation: 2.3 g

water: 300 ml

liquor ratio: 1:10

duration of treatment: 10 min

temperature: 40°C

Test principle:

Round cotton textile patches which have been washed under standard conditions (2.3 g detergent in a 300 ml liquor; 30 g textile; washing period: 10 minutes at 40°C) are placed in sterile Petri dishes (diameter: 55 mm).

All the samples are then inoculated with 0.25 ml of a bacterial suspension (approx. $\sim 10^5$ cfu/sample) and placed in a humidity chamber at 37°C.

Directly after inoculation and after 8 and 24 hours at 37°C, the inoculated textile patches are placed in 50 ml of 0.07 molar phosphate buffer (pH 7.4, containing 1 % Tween 80 and 0.3 % lecithin) and shaken for 1 minute. After shaking, a dilution series in sterile distilled water, down to a concentration of 10^{-2} , is prepared. 100 μ l samples of the undiluted solution and of the 10^{-1} and 10^{-2} dilutions are applied to the plates using a spiralometer. After incubation, the surviving colonies are counted, calculated as cfu/sample and given in Table 2 herein below.

Formulation of the detergents	<u>6</u>	<u>7</u>	<u>8</u>
combination of 30 % of the compound of formula (3) and 70 % of propylene glycol	-	-	0.6
o-phenylphenol	-	2	0.5
sodium dodecylbenzenesulfonate	20	20	20
Pareth 45-7 (Dobanol 45-7)	14	14	14
ethanol	9	9	9
soap noodles (Mettler)	10	10	10
trisodium citrate dihydrate	4	4	4
triethanolamine	5	5	5
water	ad 100 %	ad 100 %	ad 100 %
pH "as is"	10.5	10.3	10.3
appearance	clear, yellowish	clear, yellowish	clear, yellowish

Table 2				
AATCC 100-1993 cotton washed with a washing formulation under standard conditions (2.3 g of det. / 300 ml of water/ washing liquor 1:10, 10 min., 40°C)				
Formulation		<u>6</u>	<u>7</u>	<u>8</u>
S. aureus ATCC 6538	0h	4.4×10^5		
			3.6×10^5	4.1×10^5
	8h		3.9×10^5	
		8.2×10^5		
	24h	1.0×10^7	4.1×10^5	< 100
Klebsiella pneumoniae ATCC 4352	0h	1.9×10^5		
			1.8×10^5	1.8×10^5
	8h	5.7×10^8	4.3×10^8	3.1×10^2
				1×10^2
	24h	2.7×10^9	1.6×10^9	< 100

Only detergent compositions comprising the compound of formula (3) show a distinct antimicrobial activity on the textile material.

Example 4: Preparation of further liquid washing formulations

	<u>Formulation</u>										
<u>Components</u>	<u>9</u>	<u>10</u>	<u>11</u>	<u>12</u>	<u>13</u>	<u>14</u>	<u>15</u>	<u>16</u>	<u>17</u>	<u>18</u>	<u>19</u>
combination of 30 % of the compound of formula (3) and 70 % of propylene glycol	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
o-phenylphenol	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9
dodecylbenzenesulfonic acid	7.5							8.5			
sodium dodecylbenzenesulfonate		27	23.6	10	28				20	24	6
sodium laureth sulfate 3 EO	17	10									
sodium lauryl sulfate						6					8
coconut acid	12.5			10	4	4	10			10	
C ₁₂₋₁₃ Pareth-7	10							26.9	27.8	25	4
PEG-7 C ₁₃ oxoalcohol				20	9	14.5	12	29	26		
PEG-8 C ₁₃₋₁₅ fatty alcohol							10				
alkyl polyglucoside			5			1	2				
laureth-10		5									
PPG				2	3	8					
sodium carbonate			2								
sodium tripolyphosphate			20								
potassium tripolyphosphate 50 %		22									
sodium cumenesulfonate 40 %		25									
trisodium citrate	5.5					2					2
lauryltrimonium chloride	0.7										
polycarboxylate				13	18	15	10	23	16.2		

	<u>Formulation</u>										
<u>Components</u>	<u>9</u>	<u>10</u>	<u>11</u>	<u>12</u>	<u>13</u>	<u>14</u>	<u>15</u>	<u>16</u>	<u>17</u>	<u>18</u>	<u>19</u>
2-propanol	6			7	3		4	9.5	8		
ethanol	6										9
glycerol										20	
propylene glycol										6	
NaOH	3.2			2	1	2.3	1.8	1.1		1.8	4
fluorescent whitening agent Tinopal CBS-x	0.1	0.1	0.1	0.1	0.1	0.1	0.1				0.1
fluorescent whitening agent Tinopal CBS-CL								0.1	0.1	0.1	
Soap											7
water to	100	100	100	100	100	100	100	100	100	100	

Example 5: Preparation of different formulations:

	<u>formulation</u>				
<u>components</u>	<u>20</u>	<u>21</u>	<u>22</u>	<u>23</u>	<u>24</u>
combination of 30 % of the compound of formula (3) and 70 % of propylene glycol	0.13	0.13	0.6	0.6	0.3
o-phenylphenol	0.2	0.2	0.9	0.9	0.45
sodium laureth sulfate	9.0	15	1.2		
cocamidopropyl betaine	3.0	4.5	1		
decyl glucoside	3.0				
citric acid	0.1	0.1		3	
polyquaternium-7		0.4			
lauramine oxide			1		
sodium Citrate			4		
sodium carbonate			3		

	<u>formulation</u>				
<u>components</u>	<u>20</u>	<u>21</u>	<u>22</u>	<u>23</u>	<u>24</u>
ethanol			3		
sodium C ₁₄₋₁₇ alkyl sec. sulfonate				16.6	
sodium laurylsulfate				20	
Laureth-09				3	
sodium cumolsulfonate				5	
sodium chloride				3	
Quaternium 18 and iospropylalcohol					4
Pareth-25-7					0.5
water to	100	100	100	100	100

Formulation 20: shower gel

Formulation 21: shampoo


Formulation 22: all purpose cleaner

Formulation 23: dish washing detergent

Formulation 24: softener detergent

What is claimed is:

1. A surface-active composition according to claim 1 comprising
 - (a) from 0.01 to 90 % by weight of a mixture of a microbicidal active ingredient of
 - (a₁) a diphenyl ether compound and
 - (a₂) a phenol derivative,
 - (b) from 0 to 50 % by weight of one or more hydrotropic agents,
 - (c) from 0 to 80 % by weight of one or more synthetic detergents or of a soap or of combinations of the mentioned substances and/or of a salt of a saturated and/or unsaturated C₈-C₂₂ fatty acid,
 - (d) from 0 to 50 % by weight of an alcohol,
 - (e) from 0 to 50 % by weight of typical ingredients for cleaning and disinfectant compositions and optionally
 - (f) tap water or deionised water ad 100 %.
2. Composition according to claim 1 comprising
 - (a) from 0.01 to 10 % by weight of a mixture of a microbicidal active ingredient of
 - (a₁) a diphenyl ether compound and
 - (a₂) a phenol derivative,
 - (b) from 0 to 50 % by weight of one or more hydrotropic agents,
 - (c) from 5 to 80 % by weight of one or more synthetic detergents or of a soap or of combinations of the mentioned substances and/or of a salt of a saturated and/or unsaturated C₈-C₂₂ fatty acid,
 - (d) from 0 to 50 % by weight of an alcohol, and optionally
 - (e) tap water or deionised water ad 100 %.
3. A composition according to claim 1 or 2, wherein there is used as component (a₁) a 2-hydroxy-diphenyl ether of formula




Y is chlorine or bromine,

- 23 -


Z is SO₂H, NO₂ or C₁-C₄alkyl,
 r is from 0 to 3,
 o is from 0 to 3,
 p is 0, 1 or 2,
 m is 1 or 2 and
 n is 0 or 1.

4. A composition according to claim 3, wherein there is used a compound of formula




Y is chlorine and
 r is 1 or 2.

5. A composition according to any one of claims 1 to 4, wherein there is used a 2-hydroxy-diphenyl ether of formula



6. A composition according to claim 1 or 2, wherein there is used as component (a₁) a diphenyl ether compound of formula




U₁ and U₂ are each independently of the other hydrogen, hydroxy, unsubstituted or hydroxy-substituted C₁-C₂₀alkyl; C₃-C₇cycloalkyl, C₁-C₆alkylcarbonyl, C₁-C₂₀alkoxy, phenyl or phenyl-C₁-C₃alkyl;

U_3 is hydrogen, C_1 - C_{20} alkyl, C_1 - C_{20} alkoxy or C_1 - C_6 alkylcarbonyl; and

U_4 is hydrogen, unsubstituted or hydroxy-substituted C_1 - C_{20} alkyl; C_5 - C_7 cycloalkyl, hydroxy, formyl, acetyl, C_1 - C_6 alkylcarbonyl, C_2 - C_{20} alkenyl, carboxy, carboxy- C_1 - C_3 alkyl, C_1 - C_3 alkylcarbonyl- C_1 - C_3 alkyl or carboxyallyl.

7. A composition according to any one of claims 1 to 6, wherein there is used as component (a₂) a compound of formula



R_1 is hydrogen, hydroxy, C_1 - C_{20} alkyl, chloro, phenyl, benzyl or nitro,

R_2 is hydrogen, hydroxy, C_1 - C_{20} alkyl or halogen,

R_3 is hydrogen, C_1 - C_{20} alkyl, hydroxy or chloro,

R_4 is hydrogen or methyl, and

R_5 is hydrogen or nitro.

8. A composition according to claim 7, wherein o-phenylphenol is used as component (a₂).

9. A composition according to any one of claims 1 to 8, wherein the compound of formula (2) or (3) is used as component (a₁) and o-phenylphenol is used as component (a₂).

10. A composition according to any one of claims 1 to 9, wherein a sulfonate of a terpenoid or of a mono- or di-nuclear aromatic compound is used as component (b).

11. A composition according to claim 9, wherein a mono- or di-nuclear aromatic compound the sulfonate of camphor, toluene, xylene, cumene or of naphthol is used as component (b).

12. A composition according to any one of claims 1 to 9, wherein a saturated or unsaturated C_3 - C_{12} di- or poly-carboxylic acid is used as component (b).

13. A composition according to either claim 10 or claim 11, wherein a combination of cumenesulfonate and citric acid monohydrate is used as component (b).
14. A composition according to any one of claims 1 to 121, wherein a C₁₀-C₂₀alkylamido-C₁-C₄alkylenebetaine is used as component (b).
15. A composition according to any one of claims 1 to 13, wherein a salt of lauric, myristic, palmitic, stearic, arachidic, behenic, caproic, dodecenoic, tetradecenoic, octadecenoic, oleic, eicosenoic or erucic acid is used as component (c).
16. A composition according to any one of claims 1 to 154, wherein propylene glycol is used as component (d).
17. A composition according to any one of claims 1 to 15, wherein ethanol, propanol, isopropanol or a mixture of those alcohols is used as component (d).
18. Use of the composition according to any one of claims 1 to 17 for the antimicrobial treatment of human skin and hair.
19. Use of the composition according to any one of claims 1 to 17 for the antimicrobial treatment of hard surfaces.
20. Use according to claim 19, wherein the composition is used in dishwashing formulations.
21. Use according to claim 19, wherein the composition is used in all-purpose cleaners.
22. Use of the composition according to any one of claims 1 to 17 for the antimicrobial treatment of textile fibre materials.

23. Use according to claim 22, wherein the composition is used in powder washing formulations, washing pastes, liquid washing formulations, fabric softeners or solid soaps.

24. A method for the antimicrobial treatment of textile fibre materials in washing liquor, which method comprises treating the textile fibre materials in the washing liquor with a composition comprising

- (a) a microbicidal active ingredient of
 - (a₁) from 0 to 5 % by weight of a diphenyl ether compound and
 - (a₂) from 0.1 to 5 % by weight of a phenol derivative,
- (b) from 0 to 50 % by weight of one or more hydrotropic agents,
- (c) from 5 to 80 % by weight of one or more synthetic detergents or of a soap or of combinations of the mentioned substances and/or of a salt of a saturated and/or unsaturated C₈-C₂₂ fatty acid,
- (f) from 0 to 50 % by weight of an alcohol, and optionally
- (e) from 0 to 50 % by weight of typical ingredients for cleaning and disinfectant compositions and optionally
- (f) tap water or deionised water ad 100 %.

25. A method according to claim 24, wherein the washing liquor contains no component (a₁).

26. A method for imparting antimicrobial properties to textile fibre materials, which method comprises treating the textile fibre materials in the washing liquor with a composition comprising

- (a) from 0.01 to 10 % by weight of a mixture of a microbicidal active ingredient of
 - (a₁) a diphenyl ether compound and
 - (a₂) a phenol derivative,
- (b) from 0 to 50 % by weight of one or more hydrotropic agents,
- (c) from 5 to 80 % by weight of one or more synthetic detergents or of a soap or of combinations of the mentioned substances and/or of a salt of a saturated and/or unsaturated C₈-C₂₂ fatty acid,

- (d) from 0 to 50 % by weight of an alcohol,
 - (e) from 0 to 50 % by weight of typical ingredients for cleaning and disinfectant compositions and optionally and optionally
 - (f) tap water or deionised water ad 100 %,
- at least a fraction of the antimicrobial active ingredient remaining on the textile fibre material.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/14356

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C11D3/00 C11D3/20 C11D3/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	US 6 136 771 A (FOX PRISCILLA S ET AL) 24 October 2000 (2000-10-24) claims 1-8,12,21,40,41 ---	1,3-11, 17-19 2,20-23
X	WO 98 55092 A (PROCTER & GAMBLE) 10 December 1998 (1998-12-10) page 23, line 1 - line 8 claims 1,10,13 ---	1,2,7-9, 18
X	WO 00 13656 A (AHMED FAHIM Y) 16 March 2000 (2000-03-16) table 14 ---	1-7,14, 16,18
X	US 6 107 261 A (FOX PRISCILLA S ET AL) 22 August 2000 (2000-08-22) claims 1-36 --- -/--	1-7,10, 11,18-21

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* & * document member of the same patent family

Date of the actual completion of the international search

30 May 2002

Date of mailing of the international search report

05/06/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Richards, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/14356

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 855 439 A (PROCTER & GAMBLE) 29 July 1998 (1998-07-29) example 7 ----	1-11, 19-21
A	WO 96 06152 A (MOLDOVANYI LASZLO ;CIBA GEIGY AG (CH)) 29 February 1996 (1996-02-29) claims 1-20 ----	1-21
A	US 3 503 885 A (WEDELL HANS) 31 March 1970 (1970-03-31) claims 1-3,7,8 ----	1,2, 8-11,17, 19-21
A	EP 0 903 401 A (CIBA GEIGY AG) 24 March 1999 (1999-03-24) claims 1-13 ----	1-5
A	US 5 955 408 A (KAISER NANCY E ET AL) 21 September 1999 (1999-09-21) claim 1 ----	1-5, 16-18
E	EP 1 167 503 A (CIBA SC HOLDING AG) 2 January 2002 (2002-01-02) claims 1-20 -----	24,25

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/14356

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6136771	A	24-10-2000	US 6107261 A AU 5472000 A BR 0011860 A EP 1191843 A2 WO 0078275 A2 AU 5471000 A WO 0078141 A1 US 6204230 B1	22-08-2000 09-01-2001 30-04-2002 03-04-2002 28-12-2000 09-01-2001 28-12-2000 20-03-2001
WO 9855092	A	10-12-1998	AU 7803598 A CN 1265582 T EP 0996422 A1 JP 2001518940 T WO 9855092 A1 US 6294186 B1 ZA 9804761 A	21-12-1998 06-09-2000 03-05-2000 16-10-2001 10-12-1998 25-09-2001 04-12-1998
WO 0013656	A	16-03-2000	AU 5804399 A WO 0013656 A1	27-03-2000 16-03-2000
US 6107261	A	22-08-2000	AU 5471000 A AU 5472000 A BR 0011860 A EP 1191843 A2 WO 0078141 A1 WO 0078275 A2 US 6136771 A US 6204230 B1	09-01-2001 09-01-2001 30-04-2002 03-04-2002 28-12-2000 28-12-2000 24-10-2000 20-03-2001
EP 0855439	A	29-07-1998	EP 0855439 A1 AU 5917198 A BR 9807005 A CN 1255156 T EP 0855440 A1 EP 0971995 A1 JP 2002511110 T PL 336995 A1 TR 9901727 T2 WO 9832821 A1 US 6152152 A	29-07-1998 18-08-1998 14-03-2000 31-05-2000 29-07-1998 19-01-2000 09-04-2002 31-07-2000 21-05-2001 30-07-1998 28-11-2000
WO 9606152	A	29-02-1996	AU 3345195 A BG 101306 A BR 9508767 A CZ 9700555 A3 WO 9606152 A2 EP 0777716 A2 FI 970741 A HU 77706 A2 JP 10504591 T SK 24597 A3	14-03-1996 30-09-1997 11-11-1997 11-06-1997 29-02-1996 11-06-1997 21-02-1997 28-07-1998 06-05-1998 06-08-1997
US 3503885	A	31-03-1970	BE 690213 A CH 519571 A CH 484268 A DE 1467619 A1 FR 1501612 A NL 6615294 A	25-05-1967 29-02-1972 15-01-1970 13-02-1969 10-11-1967 29-05-1967

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/14356

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3503885	A		SE 317765 B	24-11-1969
EP 0903401	A	24-03-1999	EP 0903401 A1	24-03-1999
			AU 737803 B2	30-08-2001
			AU 8520298 A	01-04-1999
			BR 9803462 A	23-11-1999
			JP 11166197 A	22-06-1999
			PL 328483 A1	29-03-1999
			TR 9801841 A2	21-10-1999
			US 6365563 B1	02-04-2002
			ZA 9808453 A	17-03-1999
US 5955408	A	21-09-1999	AU 726543 B2	09-11-2000
			AU 3663997 A	02-02-1998
			CA 2260126 A1	15-01-1998
			EP 0939618 A1	08-09-1999
			WO 9801110 A1	15-01-1998
			US 6090772 A	18-07-2000
EP 1167503	A	02-01-2002	EP 1167503 A1	02-01-2002
			US 2002032241 A1	14-03-2002