

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
20 April 2006 (20.04.2006)

PCT

(10) International Publication Number
WO 2006/040182 A1

(51) International Patent Classification:

C07D 207/09 (2006.01) C07D 205/04 (2006.01)
C07D 401/12 (2006.01) C07D 413/10 (2006.01)
C07D 405/12 (2006.01) C07D 413/14 (2006.01)
C07D 409/12 (2006.01) C07D 211/28 (2006.01)
C07D 401/04 (2006.01) A61K 31/397 (2006.01)
C07D 403/04 (2006.01) A61K 31/445 (2006.01)
C07D 413/12 (2006.01) A61P 25/16 (2006.01)

(21) International Application Number:

PCT/EP2005/011106

(22) International Filing Date: 14 October 2005 (14.10.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/618,878 14 October 2004 (14.10.2004) US

(71) Applicant (for all designated States except US): **ABBOTT GMBH & CO. KG** [DE/DE]; Max-Planck-Ring 2, 65205 Wiesbaden (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DRESCHER, Karla** [DE/DE]; Unteres Bieth 10, 69221 Dossenheim (DE). **HAUPT, Andreas** [DE/DE]; Schaelzigweg 52, 68723 Schwetzingen (DE). **UNGER, Liliane** [DE/DE]; Wollstrasse 129, 67065 Ludwigshafen (DE). **TURNER, Sean, C.** [GB/DE]; Lameystr. 21, 68165 Mannheim (DE). **BRAJE, Wilfried** [DE/DE]; Meerfeldstr. 52, 68163 Mannheim (DE). **GRANDEL, Roland** [DE/DE]; Birkenweg 49, 69221 Dossenheim (DE). **HENRY, Christophe** [FR/DE]; Gartenstrasse 1, 68723 Schwetzingen (DE). **BACKFISCH, Gisela** [DE/DE]; Talstrasse 43, 69221

Dossenheim (DE). **BEYERBACH, Armin** [DE/DE]; Portugieser Strasse 5, 67098 Bad Dürkheim (DE). **LU-BISCH, Wilfried** [DE/DE]; Dreikreuzweg 58, 69151 Neckargemünd (DE).

(74) Agent: **REITSTÖTTER-KINZEBACH**; Reitstötter Kinzebach & Partner (GbR), Patentanwälte, Ludwigsplatz 4, 67059 Ludwigshafen (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, NZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HETEROCYCLIC COMPOUNDS SUITABLE FOR TREATING DISORDERS THAT RESPOND TO MODULATION OF THE DOPAMINE D₃ RECEPTOR



(57) Abstract: The invention relates to compounds of the formula (I) wherein n is 0, 1 or 2; G is CH₂ or CHR³; R¹ is H, C₁-C₆-alkyl, C₁-C₆-alkyl substituted by C₃-C₆-cycloalkyl, C₁-C₆-hydroxyalkyl, fluorinated C₁-C₆-alkyl, C₃-C₆-cycloalkyl, fluorinated C₃-C₆-cycloalkyl, C₃-C₆ alkenyl, fluorinated C₃-C₆-alkenyl, formyl, acetyl or propionyl; R², R³ and R⁴ are, independently of each other, H, methyl, fluoromethyl, difluoromethyl, or trifluoromethyl; A is phenylene, pyridylene, pyrimidylene, pyrazinylene, pyridazinylene or thiophenylene, which can be substituted by one or more substituents selected from halogen, methyl, methoxy and CF₃; E is NR⁵ or

CH₂, wherein R⁵ is H or C₁-C₃-alkyl; Ar is a cyclic radical selected from the group consisting of phenyl, a 5- or 6- membered heteroaromatic radical comprising as ring members 1, 2 or 3 heteroatoms selected from N, O and S and a phenyl ring fused to a saturated or unsaturated 5- or 6-membered carbocyclic or heterocyclic ring, where the heterocyclic ring comprises as ring members 1, 2 or 3 heteroatoms selected from N, O and S and/or 1, 2 or 3 heteroatom-containing groups each independently selected from NR⁶, where R⁶ is H, C₁-C₄-alkyl, fluorinated C₁-C₄-alkyl, C₁-C₄-alkylcarbonyl or fluorinated C₁-C₄-alkylcarbonyl, and where the cyclic radical Ar may carry 1, 2 or 3 substituents R^a, wherein the variable R^a has the meanings given in the claims and in the description; and physiologically tolerated acid addition salts thereof. The invention also relates to the use of a compound of the formula I or a pharmaceutically acceptable salt thereof for preparing a pharmaceutical composition for the treatment of a medical disorder susceptible to treatment with a dopamine D₃ receptor ligand.

HETEROCYCLIC COMPOUNDS SUITABLE FOR TREATING DISORDERS THAT RESPOND TO MODULATION OF THE DOPAMINE D₃ RECEPTOR

Background Of The Invention

5

The present invention relates to novel heterocyclic compounds. The compounds possess valuable therapeutic properties and are suitable, in particular, for treating diseases that respond to modulation of the dopamine D₃ receptor.

10 Neurons obtain their information by way of G protein-coupled receptors, inter alia. A large number of substances exert their effect by way of these receptors. One of them is dopamine. Confirmed findings exist with regard to the presence of dopamine and its physiological function as a neurotransmitter. Disorders in the dopaminergic transmitter system result in diseases of the central nervous system which include, for example,
15 schizophrenia, depression and Parkinson's disease. These diseases, and others, are treated with drugs which interact with the dopamine receptors.

Up until 1990, two subtypes of dopamine receptor had been clearly defined pharmacologically, namely the D₁ and D₂ receptors. More recently, a third subtype was found,
20 namely the D₃ receptor which appears to mediate some effects of antipsychotics and antiparkinsonians (J.C. Schwartz et al., The Dopamine D₃ Receptor as a Target for Antipsychotics, in Novel Antipsychotic Drugs, H.Y. Meltzer, Ed. Raven Press, New York 1992, pages 135-144; M. Dooley et al., Drugs and Aging 1998, 12, 495-514, J.N. Joyce, Pharmacology and Therapeutics 2001, 90, pp. 231-59 "The Dopamine D₃ Receptor as a Therapeutic Target for Antipsychotic and Antiparkinsonian Drugs").
25


Since then, the dopamine receptors have been divided into two families. On the one hand, there is the D₂ group, consisting of D₂, D₃ and D₄ receptors, and, on the other hand, the D₁ group, consisting of D₁ and D₅ receptors. Whereas D₁ and D₂ receptors
30 are widely distributed, D₃ receptors appear to be expressed regioselectively. Thus, these receptors are preferentially to be found in the limbic system and the projection regions of the mesolimbic dopamine system, especially in the nucleus accumbens, but also in other regions, such as the amygdala. Because of this comparatively regioselective expression, D₃ receptors are regarded as being a target having few side-effects
35 and it is assumed that while a selective D₃ ligand would have the properties of known antipsychotics, it would not have their dopamine D₂ receptor-mediated neurological side-effects (P. Sokoloff et al., Localization and Function of the D₃ Dopamine Receptor,

Arzneim. Forsch./Drug Res. 42(1), 224 (1992); P. Sokoloff et al. Molecular Cloning and Characterization of a Novel Dopamine Receptor (D₃) as a Target for Neuroleptics, Nature, 347, 146 (1990)).

- 5 Compounds having an affinity for the dopamine D₃ receptor have been described in the prior art on various occasions, e.g. in WO 95/04713, WO 96/23760, WO 97/45503, WO98/27081 and WO 99/58499. Some of these compounds possess moderate affinities and or selectivities for the dopamine D₃ receptor. They have therefore been proposed as being suitable for treating diseases of the central nervous system. Some of
- 10 the compounds described in these publications possess a pyrrolidinyphenyl structure. Unfortunately their affinity and selectivity towards the D₃ receptor or their pharmacological profile are not satisfactory. Consequently there is an ongoing need to provide new compounds, which either have an high affinity and an improved selectivity. The compounds should also have good pharmacological profile, e.g. a high brain plasma
- 15 ratio, a high bioavailability, good metabolic stability, or a decreased inhibition of the mitochondrial respiration.

Summary Of The Invention

- 20 The invention is based on the object of providing compounds which act as highly selective dopamine D₃ receptor ligands. This object is surprisingly achieved by means of compounds of the formula I



25

wherein

n is 0, 1 or 2;

30 G is CH₂ or CHR³;

3

- R^1 is H, C₁-C₆-alkyl, C₁-C₆-alkyl substituted by C₃-C₆-cycloalkyl, C₁-C₆-hydroxyalkyl, fluorinated C₁-C₆-alkyl, C₃-C₆-cycloalkyl, fluorinated C₃-C₆-cycloalkyl, C₃-C₆-alkenyl, fluorinated C₃-C₆-alkenyl, formyl, acetyl or propionyl;
- 5 R^2 , R^3 and R^4 are, independently of each other, H, methyl, fluoromethyl, difluoromethyl, or trifluoromethyl;
- A is phenylene, pyridylene, pyrimidylene, pyrazinylene, pyridazinylene or thio-phenylene, which can be substituted by one or more substituents selected from
10 halogen, methyl, methoxy and CF₃;
- E is NR⁵ or CH₂, wherein R⁵ is H or C₁-C₃-alkyl;
- Ar is a cyclic radical selected from the group consisting of phenyl, a 5- or 6-
15 membered heteroaromatic radical comprising as ring members 1, 2 or 3 heteroatoms selected from N, O and S and a phenyl ring fused to a saturated or unsaturated 5- or 6-membered carbocyclic or heterocyclic ring, where the heterocyclic ring comprises as ring members 1, 2 or 3 heteroatoms selected from N, O and S and/or 1, 2 or 3 heteroatom-containing groups each independently selected from
20 NR⁸, where R⁸ is H, C₁-C₄-alkyl, fluorinated C₁-C₄-alkyl, C₁-C₄-alkylcarbonyl or fluorinated C₁-C₄-alkylcarbonyl, and where the cyclic radical Ar may carry 1, 2 or 3 substituents R^a;
- R^a is halogen, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₁-C₆-alkoxy-
25 C₁-C₆-alkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₃-C₆-cycloalkyl, fluorinated C₃-C₆-cycloalkyl, C₁-C₆-alkoxy, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, phenylsulfonyl, benzyloxy, phenoxy, where the phenyl radical in the
30 3 last-mentioned radicals may be unsubstituted or may carry 1 to 3 substituents selected from C₁-C₄-alkyl, fluorinated C₁-C₄-alkyl and halogen, CN, nitro, C₁-C₆-alkylcarbonyl, fluorinated C₁-C₆-alkylcarbonyl, C₁-C₆-alkylcarbonylamino, fluorinated C₁-C₆-alkylcarbonylamino, carboxy, NH-C(O)-NR⁶R⁷, NR⁶R⁷, NR⁶R⁷-C₁-C₆-alkylene, O-NR⁶R⁷, where R⁶ and R⁷ are, independently of each other, H, C₁-C₄-
35 alkyl, fluorinated C₁-C₄-alkyl or C₁-C₄-alkoxy or may form, together with N, a 4-, 5- or 6-membered saturated or unsaturated ring, or is a saturated or unsaturated 3- to 7-membered heterocyclic ring comprising as ring members 1, 2, 3 or 4 het-

eroatoms selected from N, O and S and/or 1, 2 or 3 heteroatom-containing groups selected from NR^9 , where R^9 has one of the meanings given for R^8 , SO , SO_2 and CO , and where the heterocyclic ring may carry 1, 2 or 3 substituents selected from hydroxy, halogen, $\text{C}_1\text{-C}_6\text{-alkyl}$, fluorinated $\text{C}_1\text{-C}_6\text{-alkyl}$ and $\text{C}_1\text{-C}_6\text{-alkoxy}$;

and physiologically tolerated acid addition salts thereof.

The present invention therefore relates to compounds of the general formula I and to their physiologically tolerated acid addition salts.

The present invention also relates to a pharmaceutical composition which comprises at least one compound of the formula I and/or at least one physiologically tolerated acid addition salt of I, where appropriate together with physiologically acceptable carriers and/or auxiliary substances.

The present invention also relates to a method for treating disorders which respond to influencing by dopamine D_3 receptor antagonists or dopamine D_3 agonists, said method comprising administering an effective amount of at least one compound of the formula I and/or at least one physiologically tolerated acid addition salt of I to a subject in need thereof.

Detailed Description Of The Invention

The diseases which respond to the influence of dopamine D_3 receptor antagonists or agonists include, in particular, disorders and diseases of the central nervous system, in particular affective disturbances, neurotic disturbances, stress disturbances and somatoform disturbances and psychoses, especially schizophrenia and depression and, in addition, disturbances of kidney function, in particular kidney function disturbances which are caused by diabetes mellitus (see WO 00/67847).

According to the invention, at least one compound of the general formula I having the meanings mentioned at the outset is used for treating the above mentioned indications. Provided the compounds of the formula I of a given constitution may exist in different spatial arrangements, for example if they possess one or more centers of asymmetry, polysubstituted rings or double bonds, or as different tautomers, it is also possible to use enantiomeric mixtures, in particular racemates, diastereomeric mixtures and

5

tautomeric mixtures, preferably, however, the respective essentially pure enantiomers, diastereomers and tautomers of the compounds of formula I and/or of their salts.

Particularly, the carbon atom of the nitrogen-containing ring carrying the group A may
5 have (S) or (R) configuration. However, the (S) configuration is preferred.

Moreover, the radical A may be in a cis or trans position to either of the substituents R², R³ or R⁴ (if at least one of those is not hydrogen). However, the cis position is preferred.

10

It is likewise possible to use physiologically tolerated salts of the compounds of the formula I, especially acid addition salts with physiologically tolerated acids. Examples of suitable physiologically tolerated organic and inorganic acids are hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, C₁-C₄-alkylsulfonic acids, such as
15 methanesulfonic acid, aromatic sulfonic acids, such as benzenesulfonic acid and toluenesulfonic acid, oxalic acid, maleic acid, fumaric acid, lactic acid, tartaric acid, adipic acid and benzoic acid. Other utilizable acids are described in Fortschritte der Arzneimittelforschung [Advances in drug research], Volume 10, pages 224 ff., Birkhäuser Verlag, Basel and Stuttgart, 1966.

20

The organic moieties mentioned in the above definitions of the variables are - like the term halogen - collective terms for individual listings of the individual group members. The prefix C_n-C_m indicates in each case the possible number of carbon atoms in the group.

25

The term halogen denotes in each case fluorine, bromine, chlorine or iodine, in particular fluorine, chlorine or bromine.

C₁-C₄ Alkyl is a straight-chain or branched alkyl group having from 1 to 4 carbon atoms. Examples of an alkyl group are methyl, ethyl, n-propyl, iso-propyl, n-butyl, 2-butyl, iso-butyl or *tert*-butyl. C₁-C₂ Alkyl is methyl or ethyl, C₁-C₃ alkyl is additionally n-propyl or isopropyl.

C₁-C₆ Alkyl is a straight-chain or branched alkyl group having from 1 to 6 carbon atoms. Examples include C₁-C₄ alkyl as mentioned above and also pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl,

35

6

4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl.

5

Fluorinated C₁-C₆ alkyl is a straight-chain or branched alkyl group having from 1 to 6, especially 1 to 4 carbon atoms, in particular 1 to 3 carbon atoms, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by a fluorine atom such as in fluoromethyl, difluoromethyl, trifluoromethyl, (R)-1-fluoroethyl, (S)-1-fluoroethyl, 2-fluoroethyl, 1,1-difluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, (R)-1-fluoropropyl, (S)-1-fluoropropyl, 2-fluoropropyl, 3-fluoropropyl, 1,1-difluoropropyl, 2,2-difluoropropyl, 3,3-difluoropropyl, 3,3,3-trifluoropropyl, (R)-2-fluoro-1-methylethyl, (S)-2-fluoro-1-methylethyl, (R)-2,2-difluoro-1-methylethyl, (S)-2,2-difluoro-1-methylethyl, (R)-1,2-difluoro-1-methylethyl, (S)-1,2-difluoro-1-methylethyl, (R)-2,2,2-trifluoro-1-methylethyl, (S)-2,2,2-trifluoro-1-methylethyl, 2-fluoro-1-(fluoromethyl)ethyl, 1-(difluoromethyl)-2,2-difluoroethyl, 1-(trifluoromethyl)-2,2,2-trifluoroethyl, 1-(trifluoromethyl)-1,2,2,2-tetrafluoroethyl, (R)-1-fluorobutyl, (S)-1-fluorobutyl, 2-fluorobutyl, 3-fluorobutyl, 4-fluorobutyl, 1,1-difluorobutyl, 2,2-difluorobutyl, 3,3-difluorobutyl, 4,4-difluorobutyl, 4,4,4-trifluorobutyl, and the like.

20

Branched C₃-C₆ alkyl is alkyl having 3 to 6 carbon atoms at least one being a secondary or tertiary carbon atom. Examples are isopropyl, tert.-butyl, 2-butyl, isobutyl, 2-pentyl, 2-hexyl, 3-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1-methyl-1-ethylpropyl.

25

Fluorinated branched C₃-C₆ alkyl is alkyl having 3 to 6 carbon atoms at least one being a secondary or tertiary carbon atom wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by a fluorine atom.

30

C₁-C₆ Alkoxy is a straight-chain or branched alkyl group having from 1 to 6, in particular 1 to 4 carbon atoms, which is bound to the remainder of the molecule via an oxygen atom. Examples include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, 2-butoxy, iso-butoxy, tert.-butoxy, pentyloxy, 1-methylbutoxy, 2-methylbutoxy, 3-methylbutoxy, 2,2-dimethylpropoxy, 1-ethylpropoxy, hexyloxy, 1,1-dimethylpropoxy, 1,2-dimethylpropoxy, 1-methylpentyloxy, 2-methylpentyloxy, 3-methylpentyloxy, 4-methylpentyloxy, 1,1-dimethylbutyloxy, 1,2-dimethylbutyloxy, 1,3-dimethylbutyloxy, 2,2-dimethylbutyloxy, 2,3-dimethylbutyloxy, 3,3-dimethylbutyloxy, 1-ethylbutyloxy, 2-

35

ethylbutyloxy, 1,1,2-trimethylpropoxy, 1,2,2-trimethylpropoxy, 1-ethyl-1-methylpropoxy and 1-ethyl-2-methylpropoxy.

Fluorinated C₁-C₆ alkoxy is a straight-chain or branched alkoxy group having from 1 to 6, in particular 1 to 4 carbon atoms, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by a fluorine atoms such as in fluoromethoxy, difluoromethoxy, trifluoromethoxy, (R)-1-fluoroethoxy, (S)-1-fluoroethoxy, 2-fluoroethoxy, 1,1-difluoroethoxy, 2,2-difluoroethoxy, 2,2,2-trifluoroethoxy, (R)-1-fluoropropoxy, (S)-1-fluoropropoxy, (R)-2-fluoropropoxy, (S)-2-fluoropropoxy, 3-fluoropropoxy, 1,1-difluoropropoxy, 2,2-difluoropropoxy, 3,3-difluoropropoxy, 3,3,3-trifluoropropoxy, (R)-2-fluoro-1-methylethoxy, (S)-2-fluoro-1-methylethoxy, (R)-2,2-difluoro-1-methylethoxy, (S)-2,2-difluoro-1-methylethoxy, (R)-1,2-difluoro-1-methylethoxy, (S)-1,2-difluoro-1-methylethoxy, (R)-2,2,2-trifluoro-1-methylethoxy, (S)-2,2,2-trifluoro-1-methylethoxy, 2-fluoro-1-(fluoromethyl)ethoxy, 1-(difluoromethyl)-2,2-difluoroethoxy, (R)-1-fluorobutoxy, (S)-1-fluorobutoxy, 2-fluorobutoxy, 3-fluorobutoxy, 4-fluorobutoxy, 1,1-difluorobutoxy, 2,2-difluorobutoxy, 3,3-difluorobutoxy, 4,4-difluorobutoxy, 4,4,4-trifluorobutoxy, and the like.

C₁-C₆ Hydroxyalkyl is a straight-chain or branched alkyl group having from 1 to 6, especially 1 to 4 carbon atoms, in particular 1 to 3 carbon atoms, wherein one of the hydrogen atoms is replaced by a hydroxy group, such as in 2-hydroxyethyl or 3-hydroxypropyl.

C₁-C₆-Alkoxy-C₁-C₆-alkyl is a straight-chain or branched alkyl group having from 1 to 6, especially 1 to 4 carbon atoms, in particular 1 to 3 carbon atoms, wherein one of the hydrogen atoms is replaced by a C₁-C₆-alkoxy group, such as in methoxymethyl, 2-methoxyethyl, ethoxymethyl, 3-methoxypropyl, 3-ethoxypropyl and the like.

C₁-C₆-Alkoxy-C₁-C₆-alkoxy is a straight-chain or branched alkyl group having from 1 to 6, especially 1 to 4 carbon atoms, in particular 1 to 3 carbon atoms, wherein one of the hydrogen atoms is replaced by a C₁-C₆-alkoxy group, such as in 2-methoxyethoxy, ethoxymethoxy, 2-ethoxyethoxy, 3-methoxypropoxy, 3-ethoxypropoxy and the like.

C₁-C₆-Alkylcarbonyl is a straight-chain or branched alkyl group having from 1 to 6, especially 1 to 4 carbon atoms, in particular 1 to 3 carbon atoms, wherein one of the hydrogen atoms is replaced by a carbonyl group (CO), such as in acetyl and propionyl.

Fluorinated C₁-C₆-alkylcarbonyl is a straight-chain or branched alkyl group having from 1 to 6, especially 1 to 4 carbon atoms, in particular 1 to 3 carbon atoms, wherein one of the hydrogen atoms is replaced by a carbonyl group (CO) and wherein at least one of the remaining hydrogen atoms, e.g. 1, 2, 3, or 4 of the hydrogen atoms are replaced by a fluorine atom, such as in trifluoroacetyl and 3,3,3-trifluoropropionyl.

C₁-C₆-Alkylcarbonylamino is a straight-chain or branched alkyl group having from 1 to 6, especially 1 to 4 carbon atoms, in particular 1 to 3 carbon atoms, wherein one of the hydrogen atoms is replaced by a carbonylamino group (CO-NH-), such as in acetamido (CH₃CONH-) and propionamido (CH₃CH₂CONH-).

Fluorinated C₁-C₆-alkylcarbonylamino is a straight-chain or branched alkyl group having from 1 to 6, especially 1 to 4 carbon atoms, in particular 1 to 3 carbon atoms, wherein one of the hydrogen atoms is replaced by a carbonylamino group (CO-NH-) and wherein at least one of the remaining hydrogen atoms, e.g. 1, 2, 3, or 4 of the hydrogen atoms are replaced by a fluorine atom, such as in trifluoroacetyl and 3,3,3-trifluoropropionyl.

C₁-C₆ Alkylthio (also termed as C₁-C₆-alkylsulfanyl) (or C₁-C₆-alkylsulfinyl or C₁-C₆-alkylsulfonyl, respectively) refer to straight-chain or branched alkyl groups having 1 to 6 carbon atoms, e.g. 1 to 4 carbon atoms, which are bound to the remainder of the molecule via a sulfur atom (or S(O)O in case of alkylsulfinyl or S(O)₂O in case of alkylsulfonyl, respectively), at any bond in the alkyl group. Examples for C₁-C₄-alkylthio include methylthio, ethylthio, propylthio, isopropylthio, and n-butylthio. Examples for C₁-C₄-alkylsulfinyl include methylsulfinyl, ethylsulfinyl, propylsulfinyl, isopropylsulfinyl, and n-butylsulfinyl. Examples for C₁-C₄-alkylsulfonyl include methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, and n-butylsulfonyl.

Fluorinated C₁-C₆ alkylthio (also termed fluorinated C₁-C₆-alkylsulfanyl) is a straight-chain or branched alkylthio group having from 1 to 6, in particular 1 to 4 carbon atoms, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by fluorine atoms. Fluorinated C₁-C₆ alkylsulfinyl is a straight-chain or branched alkylsulfinyl group having from 1 to 6, in particular 1 to 4 carbon atoms, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by fluorine atoms. Fluorinated C₁-C₆ alkylsulfonyl is a straight-chain or branched alkylsulfonyl group having from 1 to 6, in particular 1 to 4 carbon atoms, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by fluorine atoms.

C₃-C₆ Cycloalkyl is a cycloaliphatic radical having from 3 to 6 C atoms, such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. The cycloalkyl radical may be unsubstituted or may carry 1, 2, 3 or 4 C₁-C₄ alkyl radicals, preferably a methyl radical. One
 5 alkyl radical is preferably located in the 1-position of the cycloalkyl radical, such as in 1-methylcyclopropyl or 1-methylcyclobutyl.

Fluorinated C₃-C₆ cycloalkyl is a cycloaliphatic radical having from 3 to 6 C atoms, such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl, wherein at least one, e.g. 1, 2,
 10 3, 4 or all of the hydrogen atoms are replaced by a fluorine atoms such as in 1-fluorocyclopropyl, 2-fluorocyclopropyl, (S)- and (R)-2,2-difluorocyclopropyl, 1,2-difluorocyclopropyl, 2,3-difluorocyclopropyl, pentafluorocyclopropyl, 1-fluorocyclobutyl, 2-fluorocyclobutyl, 3-fluorocyclobutyl, 2,2-difluorocyclobutyl, 3,3-difluorocyclobutyl, 1,2-difluorocyclobutyl, 1,3-difluorocyclobutyl, 2,3-difluorocyclobutyl, 2,4-difluorocyclobutyl,
 15 or 1,2,2-trifluorocyclobutyl.

C₂-C₆-Alkenyl is a singly unsaturated hydrocarbon radical having 2, 3, 4, 5 or 6 C-atoms, e.g. vinyl, allyl (2-propen-1-yl), 1-propen-1-yl, 2-propen-2-yl, methallyl (2-methylprop-2-en-1-yl) and the like. C₃-C₆-Alkenyl is, in particular, allyl, 1-methylprop-2-en-1-yl, 2-buten-1-yl, 3-buten-1-yl, methallyl, 2-penten-1-yl, 3-penten-1-yl, 4-penten-1-yl, 1-methylbut-2-en-1-yl or 2-ethylprop-2-en-1-yl.
 20

Fluorinated C₂-C₆-alkenyl is a singly unsaturated hydrocarbon radical having 2, 3, 4, 5 or 6 C-atoms, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are
 25 replaced by a fluorine atoms such as in 1-fluorovinyl, 2-fluorovinyl, 2,2-fluorovinyl, 3,3,3-fluoropropenyl, 1,1-difluoro-2-propenyl 1-fluoro-2-propenyl and the like.

C₁-C₆-Alkylene is a hydrocarbon bridging group having 1, 2, 3, 4, 5 or 6 carbon atoms, like methylene, ethylene, 1,2- and 1,3-propylene, 1,4-butylene and the like.
 30

Examples of 5- or 6-membered heteroaromatic radicals comprise 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, pyrazinyl, 3- or 4-pyridazinyl, 2- or 3-thienyl, 2- or 3-furyl, 2- or 3-pyrrolyl, 2-, 3- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 3- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-imidazolyl, 2- or 5-[1,3,4]oxadiazolyl, 4- or
 35 5-[1,2,3]oxadiazolyl, 3- or 5-[1,2,4]oxadiazolyl, 2- or 5-[1,3,4]thiadiazolyl, 2- or 5-[1,3,4]thiadiazolyl, 4- or 5-[1,2,3]thiadiazolyl, 3- or 5-[1,2,4]thiadiazolyl 1*H*-, 2*H*- or 3*H*- 1,2,3-triazol-4-yl, 2*H*-triazol-3-yl, 1*H*-, 2*H*-, or 4*H*-1,2,4-triazolyl and 1*H*- or 2*H*-

tetrazolyl, which may be unsubstituted or which may carry 1, 2 or 3 of the aforementioned radicals R^a .

5 Examples of a phenyl ring fused to a saturated or unsaturated 5- or 6-membered carbocyclic or heterocyclic ring comprise indenyl, indanyl, naphthyl, 1,2- or 2,3-dihydronaphthyl, tetralin, benzofuranyl, 2,3-dihydrobenzofuranyl, benzothienyl, indolyl, indazolyl, benzimidazolyl, benzoxathiazolyl, benzoxadiazolyl, benzothiadiazolyl, benzoxazinyl, dihydrobenzoxazinyl, chinolinyl, isochinolinyl, tetrahydroisochinolinyl, chromenyl, chromanyl and the like, which may be unsubstituted or which may carry 1,
10 2 or 3 of the aforementioned radicals R^a . This fused system may be bonded to the remainder of the molecule (more precisely to the sulfonyl group) via carbon atoms of the phenyl moiety or via ring atoms (C- or N-atoms) of the ring fused to phenyl.

15 Examples for saturated or unsaturated 3- to 7-membered heterocyclic rings (as radicals R^a) comprise saturated or unsaturated, aromatic or non-aromatic heterocyclic rings. Examples therefore include, apart from the above-defined 5- or 6-membered heteroaromatic radicals, aziridyl, diaziridinyl, oxiranyl, azetidiny, azetiny, di- and tetrahydrofuranyl, pyrrolinyl, pyrrolidinyl, oxopyrrolidinyl, pyrazolinyl, pyrazolidinyl, imidazoliny, imidazolidinyl, oxazoliny, oxazolidinyl, oxo-oxazolidinyl, isoxazoliny, isoxazolidinyl,
20 piperidinyl, piperazinyl, morpholinyl and the like.

If R^6 and R^7 form together with N a 4-, 5- or 6-membered ring, examples for this type of radical comprise, apart from the above-defined 5- or 6-membered heteroaromatic radicals containig at least one N atom as ring member, azetidiny, azetiny, pyrrolinyl, pyrrolidinyl, pyrazolinyl, pyrazolidinyl, imidazoliny, imidazolidinyl, oxazoliny, oxazolidinyl, piperidinyl, piperazinyl, morpholinyl and the like.

In a specific embodiment,

30 R^1 is H, C_1 - C_6 -alkyl which may be substituted by C_3 - C_6 -cycloalkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_3 - C_6 -alkenyl, fluorinated C_3 - C_6 -alkenyl, formyl, acetyl or propionyl; and

35 Ar is a cyclic radical selected from the group consisting of phenyl, a 5- or 6-membered heteroaromatic radical comprising as ring members 1, 2 or 3 heteroatoms selected from N, O and S and a phenyl ring fused to a saturated or unsaturated 5- or 6-membered carbocyclic or heterocyclic ring, where the heterocyclic ring comprises as ring members 1, 2 or 3 heteroatoms selected from N, O and S,

11

and where the cyclic radical may carry 1, 2 or 3 substituents R^a selected from the group consisting of halogen, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, CN, acetyl, carboxy, NR^6R^7 , NR^6R^7 - C_1 - C_6 -alkylene, where R^6 and R^7 are, independently of each other, H, C_1 - C_4 -alkyl or fluorinated C_1 - C_4 -alkyl or may form, together with N, a 4-, 5- or 6-membered saturated or unsaturated ring, and a saturated or unsaturated 5- or 6-membered heterocyclic ring comprising as ring members 1, 2 or 3 heteroatoms selected from N, O and S.

- 10 In compounds of formula I, n preferably is 0 or 1; i.e. the nitrogen-containing ring is an azetidiny group or a pyrrolidiny group; and particularly, n is 1, which means that in a particularly preferred embodiment, the nitrogen-containing ring is a pyrrolidiny ring.

Preferably, the radical R^1 is selected from H, C_1 - C_4 -alkyl, C_1 - C_4 -alkyl which is substituted by C_3 - C_6 -cycloalkyl or hydroxy, fluorinated C_1 - C_4 -alkyl and C_2 - C_4 -alkenyl. More preference is given to H, propyl, cyclopropylmethylene, fluorinated ethyl, e.g. 2-fluoroethyl, fluorinated propyl, e.g. 3-fluoropropyl, hydroxypropyl, e.g. 3-hydroxypropyl, propionyl and allyl. More preferably, R^1 is selected from H, propyl, cyclopropylmethylene, 2-fluoroethyl, 3-fluoropropyl, 3-hydroxypropyl, and allyl. In a particularly preferred embodiment, R^1 is n-propyl or allyl and especially n-propyl.

In an alternative embodiment, R^1 is preferably selected from H, C_1 - C_4 -alkyl, C_1 - C_4 -alkyl which is substituted by C_3 - C_6 -cycloalkyl, fluorinated C_1 - C_4 -alkyl and C_2 - C_4 -alkenyl. More preference is given to H, propyl, cyclopropylmethylene, fluorinated ethyl, e.g. 2-fluoroethyl, fluorinated propyl, e.g. 3-fluoropropyl, and allyl. In a particularly preferred embodiment, R^1 is n-propyl or allyl and especially n-propyl.

Preferably, R^2 , R^3 and R^4 are H.

- 30 The group A is preferably phenylene, pyridylene or pyrimidylene. In a more preferred embodiment, A is 1,4-phenylene, 1,2-phenylene, 2,5-pyridylene 3,6-pyridylene or 2,5-pyrimidylene, where A may be substituted as described above. In an even more preferred embodiment, A is 1,4-phenylene, 1,2-phenylene, 3,6-pyridylene or 2,5-pyrimidylene. If A is substituted, preferred substituents are selected from halogen, in particular fluorine, and methoxy. Examples include 2-fluoro-1,4-phenylene, 3-fluoro-1,4-phenylene, 2-methoxy-1,4-phenylene, and 3-methoxy-1,4-phenylene. In a specific embodiment, A is not substituted. Particularly, A is 1,4-phenylene.

The group E is preferably NR^5 , more preferably NH or NCH_3 and in particular NH.

Preferred cyclic radicals of the group Ar are phenyl, 2- or 3-thienyl, in particular 3-thienyl, imidazolyl, in particular 4-imidazolyl, isoxazolyl, in particular 4-isoxazolyl, thiazolyl, in particular 2-thiazolyl, triazolyl, in particular 3-[1,2,4]triazolyl, thiadiazolyl, in particular 3- and 5-[1,2,4]thiadiazolyl and 2-[1,3,4]thiadiazolyl, 2-, 3- or 4-pyridyl, 2- and 5-pyrimidinyl, 1-, 2-, 3-, 4- or 5-indanyl, 2-, 3-, 4- or 5-benzofuranyl, quinolinyl, in particular 8-quinolinyl, isoquinolinyl, in particular 5-isoquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, in particular 7-1,2,3,4-tetrahydroisoquinolin-7-yl, benzothienyl, in particular 2-benzothienyl, benzothiazolyl, in particular 6-benzothiazolyl, benzoxadiazolyl, in particular 4-[2,1,3]benzoxadiazolyl, benzothiadiazolyl, in particular 4-[2,1,3]benzothiadiazolyl, benzoxazin and dihydrobenzoxazin. The numbers indicate the position at which Ar is bound to the sulfonyl group. More preferred radicals Ar are phenyl, 2-thienyl, 3-pyridyl, 5-pyridyl, 5-indanyl, 2-benzofuranyl, and 2,3-dihydrobenzofuran-2-yl. Even more preferred radicals Ar are phenyl, 2-thienyl, 5-indanyl, benzofuran-2-yl and 2,3-dihydrobenzofuran-2-yl. Specifically, Ar is phenyl.

In an alternative embodiment, preferred cyclic radicals of the group Ar are phenyl, 2- or 3-thienyl, 2-, 3- or 4-pyridyl, 1-, 2-, 3-, 4- or 5-indanyl, 2-, 3-, 4- or 5-benzofuranyl, in particular 2-thienyl, 2- or 3-pyridinyl, 5-indanyl, 5-benzofuranyl, and especially phenyl.

Preferably, R^a selected from the group consisting of halogen, $\text{C}_1\text{-C}_6$ -alkyl, fluorinated $\text{C}_1\text{-C}_6$ -alkyl, $\text{C}_1\text{-C}_6$ -hydroxyalkyl, $\text{C}_1\text{-C}_6$ -alkoxy- $\text{C}_1\text{-C}_6$ -alkyl, $\text{C}_2\text{-C}_6$ -alkenyl, $\text{C}_3\text{-C}_6$ -cycloalkyl, fluorinated $\text{C}_3\text{-C}_6$ -cycloalkyl, $\text{C}_1\text{-C}_6$ -alkoxy, $\text{C}_1\text{-C}_6$ -alkoxy- $\text{C}_1\text{-C}_6$ -alkoxy, fluorinated $\text{C}_1\text{-C}_6$ -alkoxy, fluorinated $\text{C}_1\text{-C}_6$ -alkylthio, $\text{C}_1\text{-C}_6$ -alkylsulfonyl, phenylsulfonyl, benzyloxy, phenoxy, CN, nitro, acetyl, trifluoroacetyl, acetamido, carboxy, NH-C(O)-NH_2 , NR^6R^7 , $\text{NR}^6\text{R}^7\text{-C}_1\text{-C}_6$ -alkylene, $\text{O-NR}^6\text{R}^7$, where R^6 and R^7 are, independently of each other, H, $\text{C}_1\text{-C}_4$ -alkyl, fluorinated $\text{C}_1\text{-C}_4$ -alkyl or $\text{C}_1\text{-C}_4$ -alkoxy, and a saturated or unsaturated 3- to 7-membered heterocyclic ring comprising as ring members 1, 2, 3 or 4 heteroatoms selected from N, O and S and/or 1, 2 or 3 heteroatom-containing groups selected from NR^9 , where R^9 has one of the meanings given for R^8 , SO , SO_2 and CO , and where the 3- to 7-membered heterocyclic ring may carry 1, 2 or 3 substituents selected from hydroxy, halogen, $\text{C}_1\text{-C}_6$ -alkyl, fluorinated $\text{C}_1\text{-C}_6$ -alkyl and $\text{C}_1\text{-C}_6$ -alkoxy.

13

Preferably, the saturated or unsaturated 3- to 7-membered heterocyclic ring is selected from azetidin-1-yl, 2-methylazetidiny, 3-methoxyazetidiny, 3-hydroxyazetidiny, 3-fluoroazetidiny, pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, 2- and 3-fluoropyrrolidin-1-yl, 2,2-difluoropyrrolidin-1-yl, 3,3-difluoropyrrolidin-1-yl, 2- and 3-methylpyrrolidin-1-yl, 1-methylpyrrolidin-2-yl, 2,2-dimethylpyrrolidin-1-yl, 3,3-dimethylpyrrolidin-1-yl, 2-oxo-pyrrolidin-1-yl, 2- and 3-trifluoromethylpyrrolidin-1-yl, 2-oxo-oxazolidin-1-yl, piperidin-1-yl, 2-methylpiperidin-1-yl, piperazin-1-yl, 4-methylpiperazin-1-yl, morpholin-4-yl, thiomorpholin-4-yl, 1-oxothiomorpholin-4-yl, 1,1-dioxothiomorpholin-4-yl, pyrrol-1-yl, pyrrol-2-yl, pyrrol-3-yl, 1-methylpyrrol-2-yl, 1-methylpyrrol-3-yl, furan-2-yl, furan-3-yl, thiophen-2-yl, thiophen-3-yl, 5-propylthiophen-2-yl, pyrazol-1-yl, pyrazol-3-yl, pyrazol-4-yl, 1-methylpyrazol-4-yl, imidazol-1-yl, imidazol-2-yl, 1-methylimidazol-2-yl, oxazol-2-yl, oxazol-4-yl, oxazol-5-yl, isoxazol-3-yl, isoxazol-4-yl, isoxazol-5-yl, [1,2,3]triazol-1-yl, [1,2,4]triazol-1-yl, [1,2,3]triazol-2-yl, [1,2,4]triazol-3-yl, [1,2,4]triazol-4-yl, 4-methyl-[1,2,4]triazol-3-yl, 2-methyl-[1,2,3]triazol-4-yl, [1,3,4]-oxadiazol-2-yl, [1,2,4]-oxadiazol-3-yl, [1,2,4]-oxadiazol-5-yl, [1,2,3]-oxadiazol-5-yl, [1,2,3]thiadiazol-4-yl, tetrazol-1-yl, tetrazol-5-yl, 2-methyltetrazol-5-yl, 1-methyltetrazol-5-yl, furazan-3-yl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, pyrimidin-2-yl, pyrimidin-4-yl and pyrimidin-5-yl. More preferably, the saturated or unsaturated 3- to 7-membered heterocyclic ring is selected from azetidin-1-yl, pyrrolidin-1-yl, 3,3-difluoropyrrolidin-1-yl, 2-oxo-pyrrolidin-1-yl, 2-oxo-oxazolidin-1-yl, morpholin-4-yl, 2-furanyl, 5-propylthien-2-yl, pyrrol-1-yl, pyrazol-1-yl, 1-methylpyrazol-4-yl, 1-ethylpyrazol-4-yl, oxazol-5-yl, isoxazol-3-yl, isoxazol-5-yl, 4-[1,2,3]thiadiazolyl. Even more preferably, the saturated or unsaturated 3- to 7-membered heterocyclic ring is selected from azetidin-1-yl, pyrrolidin-1-yl, 3,3-difluoropyrrolidin-1-yl, 2-oxo-pyrrolidin-1-yl, morpholin-4-yl, pyrrol-1-yl, furan-2-yl, pyrazol-1-yl, 1-methylpyrazol-4-yl, oxazol-5-yl, isoxazol-5-yl, 4-[1,2,3]thiadiazolyl.

In a preferred embodiment, the cyclic radical Ar is unsubstituted or substituted by 1, 2 or 3 substituents R^a selected from the group consisting of halogen, C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, fluorinated C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, fluorinated C_1 - C_4 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxy, C_2 - C_4 -alkenyl, fluorinated C_2 - C_4 -alkenyl, NR^6R^7 , ONR^6R^7 , C_1 - C_6 -alkylene- NR^6R^7 , where R^6 and R^7 are independently of each other H, C_1 - C_4 -alkyl or or C_1 - C_4 -alkoxy, ureido ($NHCONH_2$) C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, acetyl, carboxyl, hydroxy, cyano, nitro, benzoxy, methylsulfanyl, fluoromethylsulfanyl, difluoromethylsulfanyl, trifluoromethylsulfanyl, methylsulfonyl and one of the above-mentioned saturated or unsaturated 3- to 7-membered heterocyclic rings. In a more preferred embodiment, R^a is selected from the group consisting of halogen, C_1 - C_6 -alkyl, fluorinated C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, OCF_3 , $OCHF_2$, OCH_2F , C_2 - C_4 -alkenyl, C_3 - C_6 -

cycloalkyl, fluorinated C₃-C₆- cycloalkyl, ureido, acetyl, carboxyl, hydroxy, cyano, benzoxy, trifluoromethylsulfanyl, methylsulfonyl, azetidin-1-yl, pyrrolidin-1-yl, 3,3-difluoropyrrolidin-1-yl, 2-oxo-pyrrolidin-1-yl, 2-oxo-oxazolidin-1-yl, morpholin-4-yl, 2-furanyl, 5-propylthien-2-yl, pyrrol-1-yl, pyrazol-1-yl, 1-methylpyrazol-4-yl, 1-ethylpyrazol-4-yl, oxazol-5-yl, isoxazol-3-yl, isoxazol-5-yl, and 4-[1,2,3]thiadiazolyl.

Even more preferably, R^a is selected from the group consisting of halogen, C₁-C₆-alkyl, fluorinated C₁-C₄-alkyl, C₁-C₄-alkoxy, OCF₃, OCHF₂, C₂-C₄-alkenyl, C₃-C₆-cycloalkyl, fluorinated C₃-C₆- cycloalkyl, ureido, acetyl, carboxyl, hydroxy, benzoxy, trifluoromethylsulfanyl, azetidin-1-yl, pyrrolidin-1-yl, 3,3-difluoropyrrolidin-1-yl, 2-oxo-pyrrolidin-1-yl, morpholin-4-yl, pyrrol-1-yl, furan-2-yl, pyrazol-1-yl, 1-methylpyrazol-4-yl, oxazol-5-yl, isoxazol-5-yl, and 4-[1,2,3]thiadiazolyl.

If Ar is a heteroaromatic ring, R^a in this case is in particular selected from halogen, C₂-C₄-alkenyl, oxazolyl and isoxazolyl. If Ar is a fused system, it is preferentially not substituted.

In an alternative embodiment, the cyclic radical is substituted by 1, 2 or 3 substituents R^a selected from the group consisting of halogen, C₁-C₆-alkyl, fluorinated C₁-C₄-alkyl, C₁-C₄-alkoxy, fluorinated C₁-C₄-alkoxy, C₂-C₄-alkenyl, fluorinated C₂-C₄-alkenyl, NR⁶R⁷, C₁-C₆-alkylene-NR⁶R⁷, where R⁶ and R⁷ are independently of each other H, C₁-C₄-alkyl or fluorinated C₁-C₄-alkyl or may form, together with N, a 4-, 5- or 6-membered saturated or unsaturated ring, like CH₂N(CH₃)₂, C₃-C₆-cycloalkyl optionally substituted by halogen, acetyl or carboxyl. In a more preferred embodiment, Ar is phenyl which is substituted by 1, 2 or 3 substituents selected from the group consisting of halogen, C₁-C₆-alkyl, fluorinated C₁-C₄-alkyl, C₁-C₄-alkoxy, OCF₃, OCHF₂, OCH₂F, C₂-C₄-alkenyl, fluorinated C₂-C₄-alkenyl, CH₂N(CH₃)₂, NR⁶R⁷, where R⁶ and R⁷ are independently of each other H, C₁-C₄-alkyl or fluorinated C₁-C₄-alkyl, C₃-C₆-cycloalkyl optionally substituted by halogen, acetyl or carboxyl, or Ar is thienyl, pyridyl, benzofuranyl or indanyl, which are optionally substituted by halogen, C₁-C₄-alkyl or C₁-C₄-alkenyl. More preferably, Ar is phenyl which is substituted by 1, 2 or 3 substituents R^a selected from fluorine or bromine, C₁-C₆-alkyl, especially methyl, ethyl, n-propyl, isopropyl, sec-butyl, isobutyl, dimethylpropyl, and particularly isopropyl, fluorinated C₁-C₄-alkyl, especially CF₃ or fluorinated isopropyl, C₁-C₄-alkoxy, especially methoxy, ethoxy, propoxy, isopropoxy or butoxy, OCF₃, OCHF₂, OCH₂F, isopropenyl, CH₂N(CH₃)₂, NR⁶R⁷, where R⁶ and R⁷ are independently of each other H, C₁-C₄-alkyl or fluorinated C₁-C₄-alkyl, C₃-C₆-cycloalkyl, especially cyclopentyl, fluorinated C₃-C₆-cycloalkyl, especially 2,2-difluorocyclopropyl, acetyl or carboxyl. Alternatively, Ar is thienyl or pyridyl which carry

15


1, 2 or 3 substituents selected from halogen, especially chlorine, and C₁-C₄-alkenyl, especially isopropenyl, or Ar is benzofuranyl or indanyl.

In the aforementioned 5-membered heteroaromatic radicals, Ar preferably carries one radical R^a in the 5-position (related to the 2-position of the SO₂-radical) and optionally one or two further radicals selected from halogen, in particular fluorine or chlorine.

Phenyl and the aforementioned 6-membered heteroaromatic radicals Ar preferably carry one radical R^a in the 4-position (related to the 1-position of the SO₂-radical) and optionally one or two further radicals selected from halogen, in particular fluorine or bromine.

In a very preferred embodiment of the invention Ar is phenyl that carries one radical R^a in the 4-position of the phenyl ring and optionally 1 or 2 halogen atoms, which are selected from halogen, in particular from fluorine or chlorine.

In a even more preferred embodiment, Ar preferably carries one radical R^a, which has the formula R^{a'}



wherein

Y is N, CH or CF,

R^{a1} and R^{a2} are independently of each other selected from C₁-C₂-alkyl, in particular methyl, fluorinated C₁-C₂-alkyl, in particular fluoromethyl, difluoromethyl or trifluoromethyl, provided for Y being CH or CF one of the radicals R^{a1} or R^{a2} may also be hydrogen or fluorine, or

R^{a1} and R^{a2} form a radical (CH₂)_m wherein 1 or 2 of the hydrogen atoms may be replaced by fluorine and wherein m is 2, 3 or 4, in particular CH₂-CH₂, CHF-CH₂, CF₂-CH₂, CH₂-CH₂-CH₂, CHF-CH₂-CH₂, CF₂-CH₂-CH₂, CH₂-CHF-CH₂, CH₂-CF₂-CH₂.

In case R^{a1} and R^{a2} are different from each other, the radical of the aforementioned formula Ra may have either (R)- or (S)-configuration with regard to the Y-moiety.

16

Examples for preferred radicals of the formula R^a comprise isopropyl, (R)-1-fluoroethyl, (S)-1-fluoroethyl, 2-fluoroethyl, 1,1-difluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, (R)-1-fluoropropyl, (S)-1-fluoropropyl, 2-fluoropropyl, 3-fluoropropyl, 1,1-difluoropropyl, 2,2-difluoropropyl, 3,3-difluoropropyl, 3,3,3-trifluoropropyl, (R)-2-fluoro-1-methylethyl, (S)-2-fluoro-1-methylethyl, (R)-2,2-difluoro-1-methylethyl, (S)-2,2-difluoro-1-methylethyl, (R)-1,2-difluoro-1-methylethyl, (S)-1,2-difluoro-1-methylethyl, (R)-2,2,2-trifluoro-1-methylethyl, (S)-2,2,2-trifluoro-1-methylethyl, 2-fluoro-1-(fluoromethyl)ethyl, 1-(difluoromethyl)-2,2-difluoroethyl, 1-fluoro-1-methylethyl cyclopropyl, cyclobutyl, 1-fluorocyclopropyl, (S)- and (R)-2,2-difluorocyclopropyl and 2-fluorocyclopropyl.

10

Amongst the radicals of the formula R^a those are preferred which carry 1, 2, 3 or 4, in particular 1, 2 or 3 fluorine atoms.

15

In a particularly preferred embodiment, radical R^a is in the 4-position of the phenyl ring.


Preferred examples for Ar are in particular the following: 4-methylphenyl, 4-ethylphenyl, 4-propylphenyl, 4-isopropylphenyl, 4-sec-butylphenyl, 4-isobutylphenyl, 4-(1,1-dimethylpropyl)-phenyl, 4-vinylphenyl, 4-isopropenylphenyl, 4-fluorophenyl, 4-chlorophenyl, 4-bromophenyl, 4-(fluoromethyl)phenyl, 3-(fluoromethyl)phenyl, 2-(fluoromethyl)phenyl, 4-(difluoromethyl)phenyl, 3-(difluoromethyl)phenyl, 2-(difluoromethyl)phenyl, 4-(trifluoromethyl)phenyl, 3-(trifluoromethyl)phenyl, 2-(trifluoromethyl)phenyl, 4-(1-fluoroethyl)-phenyl, 4-((S)-1-fluoroethyl)-phenyl, 4-((R)-1-fluoroethyl)-phenyl, 4-(2-fluoroethyl)-phenyl, 4-(1,1-difluoroethyl)-phenyl, 4-(2,2-difluoroethyl)-phenyl, 4-(2,2,2-trifluoroethyl)-phenyl, 4-(3-fluoropropyl)-phenyl, 4-(2-fluoropropyl)-phenyl, 4-((S)-2-fluoropropyl)-phenyl, 4-((R)-2-fluoropropyl)-phenyl, 4-(3,3-difluoropropyl)-phenyl, 4-(3,3,3-trifluoropropyl)-phenyl, 4-(1-fluoro-1-methylethyl)-phenyl, 4-(2-fluoro-1-methylethyl)-phenyl, 4-((S)-2-fluoro-1-methylethyl)-phenyl, 4-((R)-2-fluoro-1-methylethyl)-phenyl, 4-(2,2-difluoro-1-methylethyl)-phenyl, 4-((S)-2,2-difluoro-1-methylethyl)-phenyl, 4-((R)-2,2-difluoro-1-methylethyl)-phenyl, 4-(2,2,2-trifluoro-1-methylethyl)-phenyl, 4-((S)-2,2,2-trifluoro-1-methylethyl)-phenyl, 4-((R)-2,2,2-trifluoro-1-methylethyl)-phenyl, 4-(2-fluoro-1-fluoromethylethyl)-phenyl, 4-(1-difluoromethyl-2,2-difluoroethyl)-phenyl, 4-(1,1-dimethyl-2-fluoroethyl)-phenyl, 4-methoxyphenyl, 4-ethoxyphenyl, 4-propoxyphenyl, 4-isopropoxyphenyl, 4-butoxyphenyl, 4-(fluoromethoxy)-phenyl, 4-(difluoromethoxy)-phenyl, 4-(trifluoromethoxy)-phenyl, 3-(trifluoromethoxy)-phenyl, 4-(2-fluoroethoxy)-phenyl, 4-(2,2-difluoroethoxy)-phenyl, 4-(2,2,2-trifluoroethoxy)-phenyl, 4-(1,1,2,2-tetrafluoroethoxy)-phenyl, 4-cyclopropylphenyl, 4-cyclobutylphenyl, 4-

35


cyclopentylphenyl, 4-(2,2-difluorocyclopropyl)-phenyl, 3,4-difluorophenyl, 4-bromo-3-fluorophenyl, 4-bromo-2-fluorophenyl, 4-bromo-2,5-difluorophenyl, 2-fluoro-4-isopropylphenyl, 3-fluoro-4-isopropylphenyl, 4-(1-hydroxy-1-methylethyl)-phenyl, 4-(2-hydroxy-2-methylpropyl)-phenyl, 4-acetylphenyl, 4-carboxyphenyl, 4-cyanophenyl, 4-hydroxyphenyl, 4-(O-benzyl)-phenyl, 4-(2-methoxyethoxy)-phenyl, 4-(CH₂-N(CH₃)₂)-phenyl, 4-(NH-CO-NH₂)-phenyl, 4-(methylsulfanyl)-phenyl, 4-(fluoromethylsulfanyl)-phenyl, 4-(difluoromethylsulfanyl)-phenyl, 4-(trifluoromethylsulfanyl)-phenyl, 4-(methylsulfonyl)-phenyl, 4-(N-methoxy-N-methyl-amino)-phenyl, 4-(methoxyamino)-phenyl, 4-(ethoxyamino)-phenyl, 4-(N-methylaminooxy)-phenyl, 4-(N,N-dimethylaminooxy)-phenyl, 4-(azetidin-1-yl)-phenyl, 4-(2-methylazetidin-1-yl)-phenyl, 4-((S)-2-methylazetidin-1-yl)-phenyl, 4-((R)-2-methylazetidin-1-yl)-phenyl, 4-(3-fluoroazetidin-1-yl)-phenyl, 4-(3-methoxyazetidin-1-yl)-phenyl, 4-(3-hydroxyazetidin-1-yl)-phenyl, 4-(pyrrolidin-1-yl)-phenyl, 4-(pyrrolidin-2-yl)-phenyl, 4-((S)-pyrrolidin-2-yl)-phenyl, 4-((R)-pyrrolidin-2-yl)-phenyl, 4-(pyrrolidin-3-yl)-phenyl, 4-((S)-pyrrolidin-3-yl)-phenyl, 4-((R)-pyrrolidin-3-yl)-phenyl, 4-(2-fluoropyrrolidin-1-yl)-phenyl, 4-((S)-2-fluoropyrrolidin-1-yl)-phenyl, 4-((R)-2-fluoropyrrolidin-1-yl)-phenyl, 4-(3-fluoropyrrolidin-1-yl)-phenyl, 4-((S)-3-fluoropyrrolidin-1-yl)-phenyl, 4-((R)-3-fluoropyrrolidin-1-yl)-phenyl, 4-(2,2-difluoropyrrolidin-1-yl)-phenyl, 4-(3,3-difluoropyrrolidin-1-yl)-phenyl, 4-(2-methylpyrrolidin-1-yl)-phenyl, 4-((S)-2-methylpyrrolidin-1-yl)-phenyl, 4-((R)-2-methylpyrrolidin-1-yl)-phenyl, 4-(3-methylpyrrolidin-1-yl)-phenyl, 4-((S)-3-methylpyrrolidin-1-yl)-phenyl, 4-((R)-3-methylpyrrolidin-1-yl)-phenyl, 4-(1-methylpyrrolidin-2-yl)-phenyl, 4-((S)-1-methylpyrrolidin-2-yl)-phenyl, 4-((R)-1-methylpyrrolidin-2-yl)-phenyl, 4-(1-methylpyrrolidin-3-yl)-phenyl, 4-((S)-1-methylpyrrolidin-3-yl)-phenyl, 4-((R)-1-methylpyrrolidin-3-yl)-phenyl, 4-(2,2-dimethylpyrrolidin-1-yl)-phenyl, 4-(3,3-dimethylpyrrolidin-1-yl)-phenyl, 4-(2-trifluoromethylpyrrolidin-1-yl)-phenyl, 4-((S)-2-trifluoromethylpyrrolidin-1-yl)-phenyl, 4-((R)-2-trifluoromethylpyrrolidin-1-yl)-phenyl, 4-(3-trifluoromethylpyrrolidin-1-yl)-phenyl, 4-((S)-3-trifluoromethylpyrrolidin-1-yl)-phenyl, 4-((R)-3-trifluoromethylpyrrolidin-1-yl)-phenyl, 4-(2-oxopyrrolidin-1-yl)-phenyl, 4-(2-oxo-oxazolidin-3-yl)-phenyl, 4-(piperidin-1-yl)-phenyl, 4-(2-methylpiperidin-1-yl)-phenyl, 4-((S)-2-methylpiperidin-1-yl)-phenyl, 4-((R)-2-methylpiperidin-1-yl)-phenyl, 4-(piperazin-1-yl)-phenyl, 4-(4-methylpiperazin-1-yl)-phenyl, 4-(morpholin-4-yl)-phenyl, 4-(thiomorpholin-4-yl)-phenyl, 4-(1-oxo-thiomorpholin-4-yl)-phenyl, 4-(1,1-dioxo-thiomorpholin-4-yl)-phenyl, 4-(pyrrol-1-yl)-phenyl, 4-(pyrrol-2-yl)-phenyl, 4-(pyrrol-3-yl)-phenyl, 4-(1-methylpyrrol-2-yl)-phenyl, 4-(1-methylpyrrol-3-yl)-phenyl, 4-(furan-2-yl)-phenyl, 4-(furan-3-yl)-phenyl, 4-(thiophen-2-yl)-phenyl, 4-(thiophen-3-yl)-phenyl, 4-(5-propylthien-2-yl)-phenyl, 4-(pyrazol-1-yl)-phenyl, 4-(pyrazol-3-yl)-phenyl, 4-(pyrazol-4-yl)-phenyl, 4-(1-methyl-1H-pyrazol-4-yl)-phenyl,

phenyl, 4-(1-ethyl-1H-pyrazol-4-yl)-phenyl, 4-(1-methyl-1H-pyrazol-5-yl)-phenyl, 4-(1H-imidazol-2-yl)-phenyl, 4-(imidazol-1-yl)-phenyl, 4-(1-methylimidazol-2-yl)-phenyl, 4-(oxazol-2-yl)-phenyl, 4-(oxazol-4-yl)-phenyl, 4-(oxazol-5-yl)-phenyl, 4-(isoxazol-3-yl)-phenyl, 4-(isoxazol-4-yl)-phenyl, 4-(isoxazol-5-yl)-phenyl, 4-([1,2,3]-triazol-1-yl)-phenyl, 4-([1,2,4]-triazol-1-yl)-phenyl, 4-([1,2,3]-triazol-2-yl)-phenyl, 4-(4H-[1,2,4]-triazol-3-yl)-phenyl, 4-([1,2,4]-triazol-4-yl)-phenyl, 4-(2H-[1,2,3]-triazol-4-yl)-phenyl, 4-(4-methyl-4H-[1,2,4]-triazol-3-yl)-phenyl, 4-(2-methyl-2H-[1,2,3]-triazol-4-yl)-phenyl, 4-([1,3,4]-oxadiazol-2-yl)-phenyl, 4-([1,2,4]-oxadiazol-3-yl)-phenyl, 4-([1,2,4]-oxadiazol-5-yl)-phenyl, 4-([1,2,3]-oxadiazol-4-yl)-phenyl, 4-([1,2,3]-oxadiazol-5-yl)-phenyl, 4-([1,2,3]-thiadiazol-4-yl)-phenyl, 4-(1H-tetrazol-5-yl)-phenyl, 4-(tetrazol-1-yl)-phenyl, 4-(2-methyl-2H-tetrazol-5-yl)-phenyl, 4-(1-methyl-1H-tetrazol-5-yl)-phenyl, 4-furazan-3-yl-phenyl, 4-(pyrid-2-yl)-phenyl, 4-(pyrid-3-yl)-phenyl, 4-(pyrid-4-yl)-phenyl, 4-(pyrimidin-2-yl)-phenyl, 4-(pyrimidin-4-yl)-phenyl, 4-(pyrimidin-5-yl)-phenyl, 5-isopropylthiophen-2-yl, 2-chlorothiophen-5-yl, 2,5-dichlorothiophen-4-yl, 2,3-dichlorothiophen-5-yl, 2-chloro-3-nitrothiophen-5-yl, 2-(phenylsulfonyl)-thiophen-5-yl, 2-(pyridin-2-yl)thiophen-5-yl, 2-(5-(trifluoromethyl)isoxazol-3-yl)-thiophen-5-yl, 2-(2-methylthiazol-4-yl)-thiophen-5-yl, 1-methyl-1H-imidazol-4-yl, 1,2-dimethyl-1H-imidazol-4-yl, 3,5-dimethylisoxazol-4-yl, thiazol-2-yl, 4-methylthiazol-2-yl, 4-isopropylthiazol-2-yl, 4-trifluoromethylthiazol-2-yl, 5-methylthiazol-2-yl, 5-isopropylthiazol-2-yl, 5-trifluoromethylthiazol-2-yl, 2,4-dimethylthiazol-5-yl, 2-acetamido-4-methylthiazol-5-yl, 4H-[1,2,4]triazol-3-yl, 5-methyl-4H-[1,2,4]triazol-3-yl, 4-methyl-4H-[1,2,4]triazol-3-yl, 5-isopropyl-4H-[1,2,4]triazol-3-yl, 5-trifluoromethyl-4H-[1,2,4]triazol-3-yl, 4,5-dimethyl-4H-[1,2,4]triazol-3-yl, 5-isopropyl-4-methyl-4H-[1,2,4]triazol-3-yl, 5-trifluoromethyl-4-methyl-4H-[1,2,4]triazol-3-yl, [1,3,4]thiadiazol-2-yl, 5-methyl-[1,3,4]thiadiazol-2-yl, 5-isopropyl-[1,3,4]thiadiazol-2-yl, 5-trifluoromethyl-[1,3,4]thiadiazol-2-yl, 3-bromo-2-chloropyrid-5-yl, 2-(4-morpholino)-pyrid-5-yl, 2-phenoxy-pyrid-5-yl, (2-isopropyl)-pyrimidin-5-yl, (5-isopropyl)-pyrimidin-2-yl, 8-quinolyl, 5-isoquinolyl, 2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl, 5-chloro-3-methylbenzothiophen-2-yl, 3,4-dihydro-4-methyl-2H-benzo[b][1,4]oxazinyl, benzothiazol-6-yl, benzo[2,1,3]oxadiazol-4-yl, 5-chlorobenzo[2,1,3]oxadiazol-4-yl, 7-chlorobenzo[2,1,3]oxadiazol-4-yl and benzo[2,1,3]thiadiazol-4-yl.


Particularly preferred compounds I are those of formulae I.a, I.b, I.c, I.d, I.e, I.f, I.g, I.h, I.i, I.k, I.l, I.m, I.n, I.o, I.p, I.q, I.r, I.s and I.t, wherein R¹ and Ar have the above-defined meanings. Preferred meanings of R¹ and Ar are as defined above.




I.a




I.b




I.c




I.d




I.e




I.f




I.g




I.h




I.i




I.k




I.l




I.m




I.n




I.o




I.p




I.q



I.r



I.s



I.t

21

Examples of preferred compounds which are represented by the formulae I.a, I.b, I.c, I.d, I.e, I.f, I.g, I.h, I.i, I.k, I.l, I.m, I.n, I.o, I.p, I.q, I.r, I.s and I.t are the individual compounds listed above, where the variables Ar and R¹ have the meanings given in one row of table A.

5

Table A

No.	R ¹	Ar
1.	propyl	4-methylphenyl
2.	propyl	4-ethylphenyl
3.	propyl	4-propylphenyl
4.	propyl	4-isopropylphenyl
5.	propyl	4-sec-butylphenyl
6.	propyl	4-isobutylphenyl
7.	propyl	4-(1,1-dimethylpropyl)-phenyl
8.	propyl	4-vinylphenyl
9.	propyl	4-isopropenylphenyl
10.	propyl	4-fluorophenyl
11.	propyl	4-chlorophenyl
12.	propyl	4-bromophenyl
13.	propyl	4-(fluoromethyl)phenyl
14.	propyl	3-(fluoromethyl)phenyl
15.	propyl	2-(fluoromethyl)phenyl
16.	propyl	4-(difluoromethyl)phenyl
17.	propyl	3-(difluoromethyl)phenyl
18.	propyl	2-(difluoromethyl)phenyl
19.	propyl	4-(trifluoromethyl)phenyl
20.	propyl	3-(trifluoromethyl)phenyl
21.	propyl	2-(trifluoromethyl)phenyl
22.	propyl	4-(1-fluoroethyl)-phenyl
23.	propyl	4-((S)-1-fluoroethyl)-phenyl
24.	propyl	4-((R)-1-fluoroethyl)-phenyl
25.	propyl	4-(2-fluoroethyl)-phenyl
26.	propyl	4-(1,1-difluoroethyl)-phenyl
27.	propyl	4-(2,2-difluoroethyl)-phenyl
28.	propyl	4-(2,2,2-trifluoroethyl)-phenyl

No.	R ¹	Ar
29.	propyl	4-(3-fluoropropyl)-phenyl
30.	propyl	4-(2-fluoropropyl)-phenyl
31.	propyl	4-((S)-2-fluoropropyl)-phenyl
32.	propyl	4-((R)-2-fluoropropyl)-phenyl
33.	propyl	4-(3,3-difluoropropyl)-phenyl
34.	propyl	4-(3,3,3-trifluoropropyl)-phenyl
35.	propyl	4-(1-fluoro-1-methylethyl)-phenyl
36.	propyl	4-(2-fluoro-1-methylethyl)-phenyl
37.	propyl	4-((S)-2-fluoro-1-methylethyl)-phenyl
38.	propyl	4-((R)-2-fluoro-1-methylethyl)-phenyl
39.	propyl	4-(2,2-difluoro-1-methylethyl)-phenyl
40.	propyl	4-((S)-2,2-difluoro-1-methylethyl)-phenyl
41.	propyl	4-((R)-2,2-difluoro-1-methylethyl)-phenyl
42.	propyl	4-(2,2,2-trifluoro-1-methylethyl)-phenyl
43.	propyl	4-((S)-2,2,2-trifluoro-1-methylethyl)-phenyl
44.	propyl	4-((R)-2,2,2-trifluoro-1-methylethyl)-phenyl
45.	propyl	4-(2-fluoro-1-fluoromethylethyl)-phenyl
46.	propyl	4-(1-difluoromethyl-2,2-difluoroethyl)-phenyl
47.	propyl	4-(1,1-dimethyl-2-fluoroethyl)-phenyl
48.	propyl	4-methoxyphenyl
49.	propyl	4-ethoxyphenyl
50.	propyl	4-propoxyphenyl
51.	propyl	4-isopropoxyphenyl
52.	propyl	4-butoxyphenyl
53.	propyl	4-(fluoromethoxy)-phenyl
54.	propyl	4-(difluoromethoxy)-phenyl
55.	propyl	4-(trifluoromethoxy)-phenyl
56.	propyl	3-(trifluoromethoxy)-phenyl
57.	propyl	4-(2-fluoroethoxy)-phenyl
58.	propyl	4-(2,2-difluoroethoxy)-phenyl
59.	propyl	4-(2,2,2-trifluoroethoxy)-phenyl
60.	propyl	4-(1,1,2,2-tetrafluoroethoxy)-phenyl
61.	propyl	4-cyclopropylphenyl
62.	propyl	4-cyclobutylphenyl
63.	propyl	4-cyclopentylphenyl

23

No.	R ¹	Ar
64.	propyl	4-(2,2-difluorocyclopropyl)-phenyl
65.	propyl	3,4-difluorophenyl
66.	propyl	4-bromo-3-fluorophenyl
67.	propyl	4-bromo-2-fluorophenyl
68.	propyl	4-bromo-2,5-difluorophenyl
69.	propyl	2-fluoro-4-isopropylphenyl
70.	propyl	3-fluoro-4-isopropylphenyl
71.	propyl	4-(1-hydroxy-1-methylethyl)-phenyl
72.	propyl	4-(2-hydroxy-2-methylpropyl)-phenyl
73.	propyl	4-acetylphenyl
74.	propyl	4-carboxyphenyl
75.	propyl	4-cyanophenyl
76.	propyl	4-hydroxyphenyl
77.	propyl	4-(O-benzyl)-phenyl
78.	propyl	4-(2-methoxyethoxy)-phenyl
79.	propyl	4-(CH ₂ -N(CH ₃) ₂)-phenyl
80.	propyl	4-(NH-CO-NH ₂)-phenyl
81.	propyl	4-(methylsulfanyl)-phenyl
82.	propyl	4-(fluoromethylsulfanyl)-phenyl
83.	propyl	4-(difluoromethylsulfanyl)-phenyl
84.	propyl	4-(trifluoromethylsulfanyl)-phenyl
85.	propyl	4-(methylsulfonyl)-phenyl
86.	propyl	4-(N-methoxy-N-methyl-amino)-phenyl
87.	propyl	4-(methoxyamino)-phenyl
88.	propyl	4-(ethoxyamino)-phenyl
89.	propyl	4-(N-methylaminooxy)-phenyl
90.	propyl	4-(N,N-dimethylaminooxy)-phenyl
91.	propyl	4-(azetidin-1-yl)-phenyl
92.	propyl	4-(2-methylazetidin-1-yl)-phenyl
93.	propyl	4-((S)-2-methylazetidin-1-yl)-phenyl
94.	propyl	4-((R)-2-methylazetidin-1-yl)-phenyl
95.	propyl	4-(3-fluoroazetidin-1-yl)-phenyl
96.	propyl	4-(3-methoxyazetidin-1-yl)-phenyl
97.	propyl	4-(3-hydroxyazetidin-1-yl)-phenyl
98.	propyl	4-(pyrrolidin-1-yl)-phenyl

No.	R ¹	Ar
99.	propyl	4-(pyrrolidin-2-yl)-phenyl
100.	propyl	4-((S)-pyrrolidin-2-yl)-phenyl
101.	propyl	4-((R)-pyrrolidin-2-yl)-phenyl
102.	propyl	4-(pyrrolidin-3-yl)-phenyl
103.	propyl	4-((S)-pyrrolidin-3-yl)-phenyl
104.	propyl	4-((R)-pyrrolidin-3-yl)-phenyl
105.	propyl	4-(2-fluoropyrrolidin-1-yl)-phenyl
106.	propyl	4-((S)-2-fluoropyrrolidin-1-yl)-phenyl
107.	propyl	4-((R)-2-fluoropyrrolidin-1-yl)-phenyl
108.	propyl	4-(3-fluoropyrrolidin-1-yl)-phenyl
109.	propyl	4-((S)-3-fluoropyrrolidin-1-yl)-phenyl
110.	propyl	4-((R)-3-fluoropyrrolidin-1-yl)-phenyl
111.	propyl	4-(2,2-difluoropyrrolidin-1-yl)-phenyl
112.	propyl	4-(3,3-difluoropyrrolidin-1-yl)-phenyl
113.	propyl	4-(2-methylpyrrolidin-1-yl)-phenyl
114.	propyl	4-((S)-2-methylpyrrolidin-1-yl)-phenyl
115.	propyl	4-((R)-2-methylpyrrolidin-1-yl)-phenyl
116.	propyl	4-(3-methylpyrrolidin-1-yl)-phenyl
117.	propyl	4-((S)-3-methylpyrrolidin-1-yl)-phenyl
118.	propyl	4-((R)-3-methylpyrrolidin-1-yl)-phenyl
119.	propyl	4-(1-methylpyrrolidin-2-yl)-phenyl
120.	propyl	4-((S)-1-methylpyrrolidin-2-yl)-phenyl
121.	propyl	4-((R)-1-methylpyrrolidin-2-yl)-phenyl
122.	propyl	4-(1-methylpyrrolidin-3-yl)-phenyl
123.	propyl	4-((S)-1-methylpyrrolidin-3-yl)-phenyl
124.	propyl	4-((R)-1-methylpyrrolidin-3-yl)-phenyl
125.	propyl	4-(2,2-dimethylpyrrolidin-1-yl)-phenyl
126.	propyl	4-(3,3-dimethylpyrrolidin-1-yl)-phenyl
127.	propyl	4-(2-trifluoromethylpyrrolidin-1-yl)-phenyl
128.	propyl	4-((S)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
129.	propyl	4-((R)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
130.	propyl	4-(3-trifluoromethylpyrrolidin-1-yl)-phenyl
131.	propyl	4-((S)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
132.	propyl	4-((R)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
133.	propyl	4-(2-oxopyrrolidin-1-yl)-phenyl

No.	R ¹	Ar
134.	propyl	4-(2-oxo-oxazolidin-3-yl)-phenyl
135.	propyl	4-(piperidin-1-yl)-phenyl
136.	propyl	4-(2-methylpiperidin-1-yl)-phenyl
137.	propyl	4-((S)-2-methylpiperidin-1-yl)-phenyl
138.	propyl	4-((R)-2-methylpiperidin-1-yl)-phenyl
139.	propyl	4-(piperazin-1-yl)-phenyl
140.	propyl	4-(4-methylpiperazin-1-yl)-phenyl
141.	propyl	4-(morpholin-4-yl)-phenyl
142.	propyl	4-(thiomorpholin-4-yl)-phenyl
143.	propyl	4-(1-oxo-thiomorpholin-4-yl)-phenyl
144.	propyl	4-(1,1-dioxo-thiomorpholin-4-yl)-phenyl
145.	propyl	4-(pyrrol-1-yl)-phenyl
146.	propyl	4-(pyrrol-2-yl)-phenyl
147.	propyl	4-(pyrrol-3-yl)-phenyl
148.	propyl	4-(1-methylpyrrol-2-yl)-phenyl
149.	propyl	4-(1-methylpyrrol-3-yl)-phenyl
150.	propyl	4-(furan-2-yl)-phenyl
151.	propyl	4-(furan-3-yl)-phenyl
152.	propyl	4-(thiophen-2-yl)-phenyl
153.	propyl	4-(thiophen-3-yl)-phenyl
154.	propyl	4-(5-propylthien-2-yl)-phenyl
155.	propyl	4-(pyrazol-1-yl)-phenyl
156.	propyl	4-(pyrazol-3-yl)-phenyl
157.	propyl	4-(pyrazol-4-yl)-phenyl
158.	propyl	4-(1-methyl-1H-pyrazol-4-yl)-phenyl
159.	propyl	4-(1-ethyl-1H-pyrazol-4-yl)-phenyl
160.	propyl	4-(1-methyl-1H-pyrazol-5-yl)-phenyl
161.	propyl	4-(1H-imidazol-2-yl)-phenyl
162.	propyl	4-(imidazol-1-yl)-phenyl
163.	propyl	4-(1-methylimidazol-2-yl)-phenyl
164.	propyl	4-(oxazol-2-yl)-phenyl
165.	propyl	4-(oxazol-4-yl)-phenyl
166.	propyl	4-(oxazol-5-yl)-phenyl
167.	propyl	4-(isoxazol-3-yl)-phenyl
168.	propyl	4-(isoxazol-4-yl)-phenyl

No.	R ¹	Ar
169.	propyl	4-(isoxazol-5-yl)-phenyl
170.	propyl	4-([1,2,3]-triazol-1-yl)-phenyl
171.	propyl	4-([1,2,4]-triazol-1-yl)-phenyl
172.	propyl	4-([1,2,3]-triazol-2-yl)-phenyl
173.	propyl	4-(4H-[1,2,4]-triazol-3-yl)-phenyl
174.	propyl	4-([1,2,4]-triazol-4-yl)-phenyl
175.	propyl	4-(2H-[1,2,3]-triazol-4-yl)-phenyl
176.	propyl	4-(4-methyl-4H-[1,2,4]-triazol-3-yl)-phenyl
177.	propyl	4-(2-methyl-2H-[1,2,3]-triazol-4-yl)-phenyl
178.	propyl	4-([1,3,4]-oxadiazol-2-yl)-phenyl
179.	propyl	4-([1,2,4]-oxadiazol-3-yl)-phenyl
180.	propyl	4-([1,2,4]-oxadiazol-5-yl)-phenyl
181.	propyl	4-([1,2,3]-oxadiazol-4-yl)-phenyl
182.	propyl	4-([1,2,3]-oxadiazol-5-yl)-phenyl
183.	propyl	4-([1,2,3]-thiadiazol-4-yl)-phenyl
184.	propyl	4-(1H-tetrazol-5-yl)-phenyl
185.	propyl	4-(tetrazol-1-yl)-phenyl
186.	propyl	4-(2-methyl-2H-tetrazol-5-yl)-phenyl
187.	propyl	4-(1-methyl-1H-tetrazol-5-yl)-phenyl
188.	propyl	4-furazan-3-yl-phenyl
189.	propyl	4-(pyrid-2-yl)-phenyl
190.	propyl	4-(pyrid-3-yl)-phenyl
191.	propyl	4-(pyrid-4-yl)-phenyl
192.	propyl	4-(pyrimidin-2-yl)-phenyl
193.	propyl	4-(pyrimidin-4-yl)-phenyl
194.	propyl	4-(pyrimidin-5-yl)-phenyl
195.	propyl	5-isopropylthiophen-2-yl
196.	propyl	2-chlorothiophen-5-yl
197.	propyl	2,5-dichlorothiophen-4-yl
198.	propyl	2,3-dichlorothiophen-5-yl
199.	propyl	2-chloro-3-nitrothiophen-5-yl
200.	propyl	2-(phenylsulfonyl)-thiophen-5-yl
201.	propyl	2-(pyridin-2-yl)thiophen-5-yl
202.	propyl	2-(5-(trifluoromethyl)isoxazol-3-yl)-thiophen-5-yl
203.	propyl	2-(2-methylthiazol-4-yl)-thiophen-5-yl

No.	R ¹	Ar
204.	propyl	1-methyl-1H-imidazol-4-yl
205.	propyl	1,2-dimethyl-1H-imidazol-4-yl
206.	propyl	3,5-dimethylisoxazol-4-yl
207.	propyl	thiazol-2-yl
208.	propyl	4-methylthiazol-2-yl
209.	propyl	4-isopropylthiazol-2-yl
210.	propyl	4-trifluoromethylthiazol-2-yl
211.	propyl	5-methylthiazol-2-yl
212.	propyl	5-isopropylthiazol-2-yl
213.	propyl	5-trifluoromethylthiazol-2-yl
214.	propyl	2,4-dimethylthiazol-5-yl
215.	propyl	2-acetamido-4-methylthiazol-5-yl
216.	propyl	4H-[1,2,4]triazol-3-yl
217.	propyl	5-methyl-4H-[1,2,4]triazol-3-yl
218.	propyl	4-methyl-4H-[1,2,4]triazol-3-yl
219.	propyl	5-isopropyl-4H-[1,2,4]triazol-3-yl
220.	propyl	5-trifluoromethyl-4H-[1,2,4]triazol-3-yl
221.	propyl	4,5-dimethyl-4H-[1,2,4]triazol-3-yl
222.	propyl	5-isopropyl-4-methyl-4H-[1,2,4]triazol-3-yl
223.	propyl	5-trifluoromethyl-4-methyl-4H-[1,2,4]triazol-3-yl
224.	propyl	[1,3,4]thiadiazol-2-yl
225.	propyl	5-methyl-[1,3,4]thiadiazol-2-yl
226.	propyl	5-isopropyl-[1,3,4]thiadiazol-2-yl
227.	propyl	5-trifluoromethyl-[1,3,4]thiadiazol-2-yl
228.	propyl	3-bromo-2-chloropyrid-5-yl
229.	propyl	2-(4-morpholino)-pyrid-5-yl
230.	propyl	2-phenoxy-pyrid-5-yl
231.	propyl	(2-isopropyl)-pyrimidin-5-yl
232.	propyl	(5-isopropyl)-pyrimidin-2-yl
233.	propyl	8-quinolyl
234.	propyl	5-isoquinolyl
235.	propyl	2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl
236.	propyl	5-chloro-3-methylbenzothiophen-2-yl
237.	propyl	3,4-dihydro-4-methyl-2H-benzo[b][1,4]oxazinyl

No.	R ¹	Ar
238.	propyl	benzothiazol-6-yl
239.	propyl	benzo[2,1,3]oxadiazol-4-yl
240.	propyl	5-chlorobenzo[2,1,3]oxadiazol-4-yl
241.	propyl	7-chlorobenzo[2,1,3]oxadiazol-4-yl
242.	propyl	benzo[2,1,3]thiadiazol-4-yl
243.	ethyl	4-methylphenyl
244.	ethyl	4-ethylphenyl
245.	ethyl	4-propylphenyl
246.	ethyl	4-isopropylphenyl
247.	ethyl	4-sec-butylphenyl
248.	ethyl	4-isobutylphenyl
249.	ethyl	4-(1,1-dimethylpropyl)-phenyl
250.	ethyl	4-vinylphenyl
251.	ethyl	4-isopropenylphenyl
252.	ethyl	4-fluorophenyl
253.	ethyl	4-chlorophenyl
254.	ethyl	4-bromophenyl
255.	ethyl	4-(fluoromethyl)phenyl
256.	ethyl	3-(fluoromethyl)phenyl
257.	ethyl	2-(fluoromethyl)phenyl
258.	ethyl	4-(difluoromethyl)phenyl
259.	ethyl	3-(difluoromethyl)phenyl
260.	ethyl	2-(difluoromethyl)phenyl
261.	ethyl	4-(trifluoromethyl)phenyl
262.	ethyl	3-(trifluoromethyl)phenyl
263.	ethyl	2-(trifluoromethyl)phenyl
264.	ethyl	4-(1-fluoroethyl)-phenyl
265.	ethyl	4-((S)-1-fluoroethyl)-phenyl
266.	ethyl	4-((R)-1-fluoroethyl)-phenyl
267.	ethyl	4-(2-fluoroethyl)-phenyl
268.	ethyl	4-(1,1-difluoroethyl)-phenyl
269.	ethyl	4-(2,2-difluoroethyl)-phenyl
270.	ethyl	4-(2,2,2-trifluoroethyl)-phenyl
271.	ethyl	4-(3-fluoropropyl)-phenyl
272.	ethyl	4-(2-fluoropropyl)-phenyl

No.	R ¹	Ar
273.	ethyl	4-((S)-2-fluoropropyl)-phenyl
274.	ethyl	4-((R)-2-fluoropropyl)-phenyl
275.	ethyl	4-(3,3-difluoropropyl)-phenyl
276.	ethyl	4-(3,3,3-trifluoropropyl)-phenyl
277.	ethyl	4-(1-fluoro-1-methylethyl)-phenyl
278.	ethyl	4-(2-fluoro-1-methylethyl)-phenyl
279.	ethyl	4-((S)-2-fluoro-1-methylethyl)-phenyl
280.	ethyl	4-((R)-2-fluoro-1-methylethyl)-phenyl
281.	ethyl	4-(2,2-difluoro-1-methylethyl)-phenyl
282.	ethyl	4-((S)-2,2-difluoro-1-methylethyl)-phenyl
283.	ethyl	4-((R)-2,2-difluoro-1-methylethyl)-phenyl
284.	ethyl	4-(2,2,2-trifluoro-1-methylethyl)-phenyl
285.	ethyl	4-((S)-2,2,2-trifluoro-1-methylethyl)-phenyl
286.	ethyl	4-((R)-2,2,2-trifluoro-1-methylethyl)-phenyl
287.	ethyl	4-(2-fluoro-1-fluoromethylethyl)-phenyl
288.	ethyl	4-(1-difluoromethyl-2,2-difluoroethyl)-phenyl
289.	ethyl	4-(1,1-dimethyl-2-fluoroethyl)-phenyl
290.	ethyl	4-methoxyphenyl
291.	ethyl	4-ethoxyphenyl
292.	ethyl	4-propoxyphenyl
293.	ethyl	4-isopropoxyphenyl
294.	ethyl	4-butoxyphenyl
295.	ethyl	4-(fluoromethoxy)-phenyl
296.	ethyl	4-(difluoromethoxy)-phenyl
297.	ethyl	4-(trifluoromethoxy)-phenyl
298.	ethyl	3-(trifluoromethoxy)-phenyl
299.	ethyl	4-(2-fluoroethoxy)-phenyl
300.	ethyl	4-(2,2-difluoroethoxy)-phenyl
301.	ethyl	4-(2,2,2-trifluoroethoxy)-phenyl
302.	ethyl	4-(1,1,2,2-tetrafluoroethoxy)-phenyl
303.	ethyl	4-cyclopropylphenyl
304.	ethyl	4-cyclobutylphenyl
305.	ethyl	4-cyclopentylphenyl
306.	ethyl	4-(2,2-difluorocyclopropyl)-phenyl
307.	ethyl	3,4-difluorophenyl

No.	R ¹	Ar
308.	ethyl	4-bromo-3-fluorophenyl
309.	ethyl	4-bromo-2-fluorophenyl
310.	ethyl	4-bromo-2,5-difluorophenyl
311.	ethyl	2-fluoro-4-isopropylphenyl
312.	ethyl	3-fluoro-4-isopropylphenyl
313.	ethyl	4-(1-hydroxy-1-methylethyl)-phenyl
314.	ethyl	4-(2-hydroxy-2-methylpropyl)-phenyl
315.	ethyl	4-acetylphenyl
316.	ethyl	4-carboxyphenyl
317.	ethyl	4-cyanophenyl
318.	ethyl	4-hydroxyphenyl
319.	ethyl	4-(O-benzyl)-phenyl
320.	ethyl	4-(2-methoxyethoxy)-phenyl
321.	ethyl	4-(CH ₂ -N(CH ₃) ₂)-phenyl
322.	ethyl	4-(NH-CO-NH ₂)-phenyl
323.	ethyl	4-(methylsulfanyl)-phenyl
324.	ethyl	4-(fluoromethylsulfanyl)-phenyl
325.	ethyl	4-(difluoromethylsulfanyl)-phenyl
326.	ethyl	4-(trifluoromethylsulfanyl)-phenyl
327.	ethyl	4-(methylsulfonyl)-phenyl
328.	ethyl	4-(N-methoxy-N-methyl-amino)-phenyl
329.	ethyl	4-(methoxyamino)-phenyl
330.	ethyl	4-(ethoxyamino)-phenyl
331.	ethyl	4-(N-methylaminooxy)-phenyl
332.	ethyl	4-(N,N-dimethylaminooxy)-phenyl
333.	ethyl	4-(azetidin-1-yl)-phenyl
334.	ethyl	4-(2-methylazetidin-1-yl)-phenyl
335.	ethyl	4-((S)-2-methylazetidin-1-yl)-phenyl
336.	ethyl	4-((R)-2-methylazetidin-1-yl)-phenyl
337.	ethyl	4-(3-fluoroazetidin-1-yl)-phenyl
338.	ethyl	4-(3-methoxyazetidin-1-yl)-phenyl
339.	ethyl	4-(3-hydroxyazetidin-1-yl)-phenyl
340.	ethyl	4-(pyrrolidin-1-yl)-phenyl
341.	ethyl	4-(pyrrolidin-2-yl)-phenyl
342.	ethyl	4-((S)-pyrrolidin-2-yl)-phenyl

No.	R ¹	Ar
343.	ethyl	4-((R)-pyrrolidin-2-yl)-phenyl
344.	ethyl	4-(pyrrolidin-3-yl)-phenyl
345.	ethyl	4-((S)-pyrrolidin-3-yl)-phenyl
346.	ethyl	4-((R)-pyrrolidin-3-yl)-phenyl
347.	ethyl	4-(2-fluoropyrrolidin-1-yl)-phenyl
348.	ethyl	4-((S)-2-fluoropyrrolidin-1-yl)-phenyl
349.	ethyl	4-((R)-2-fluoropyrrolidin-1-yl)-phenyl
350.	ethyl	4-(3-fluoropyrrolidin-1-yl)-phenyl
351.	ethyl	4-((S)-3-fluoropyrrolidin-1-yl)-phenyl
352.	ethyl	4-((R)-3-fluoropyrrolidin-1-yl)-phenyl
353.	ethyl	4-(2,2-difluoropyrrolidin-1-yl)-phenyl
354.	ethyl	4-(3,3-difluoropyrrolidin-1-yl)-phenyl
355.	ethyl	4-(2-methylpyrrolidin-1-yl)-phenyl
356.	ethyl	4-((S)-2-methylpyrrolidin-1-yl)-phenyl
357.	ethyl	4-((R)-2-methylpyrrolidin-1-yl)-phenyl
358.	ethyl	4-(3-methylpyrrolidin-1-yl)-phenyl
359.	ethyl	4-((S)-3-methylpyrrolidin-1-yl)-phenyl
360.	ethyl	4-((R)-3-methylpyrrolidin-1-yl)-phenyl
361.	ethyl	4-(1-methylpyrrolidin-2-yl)-phenyl
362.	ethyl	4-((S)-1-methylpyrrolidin-2-yl)-phenyl
363.	ethyl	4-((R)-1-methylpyrrolidin-2-yl)-phenyl
364.	ethyl	4-(1-methylpyrrolidin-3-yl)-phenyl
365.	ethyl	4-((S)-1-methylpyrrolidin-3-yl)-phenyl
366.	ethyl	4-((R)-1-methylpyrrolidin-3-yl)-phenyl
367.	ethyl	4-(2,2-dimethylpyrrolidin-1-yl)-phenyl
368.	ethyl	4-(3,3-dimethylpyrrolidin-1-yl)-phenyl
369.	ethyl	4-(2-trifluoromethylpyrrolidin-1-yl)-phenyl
370.	ethyl	4-((S)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
371.	ethyl	4-((R)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
372.	ethyl	4-(3-trifluoromethylpyrrolidin-1-yl)-phenyl
373.	ethyl	4-((S)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
374.	ethyl	4-((R)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
375.	ethyl	4-(2-oxopyrrolidin-1-yl)-phenyl
376.	ethyl	4-(2-oxo-oxazolidin-3-yl)-phenyl
377.	ethyl	4-(piperidin-1-yl)-phenyl

No.	R ¹	Ar
378.	ethyl	4-(2-methylpiperidin-1-yl)-phenyl
379.	ethyl	4-((S)-2-methylpiperidin-1-yl)-phenyl
380.	ethyl	4-((R)-2-methylpiperidin-1-yl)-phenyl
381.	ethyl	4-(piperazin-1-yl)-phenyl
382.	ethyl	4-(4-methylpiperazin-1-yl)-phenyl
383.	ethyl	4-(morpholin-4-yl)-phenyl
384.	ethyl	4-(thiomorpholin-4-yl)-phenyl
385.	ethyl	4-(1-oxo-thiomorpholin-4-yl)-phenyl
386.	ethyl	4-(1,1-dioxo-thiomorpholin-4-yl)-phenyl
387.	ethyl	4-(pyrrol-1-yl)-phenyl
388.	ethyl	4-(pyrrol-2-yl)-phenyl
389.	ethyl	4-(pyrrol-3-yl)-phenyl
390.	ethyl	4-(1-methylpyrrol-2-yl)-phenyl
391.	ethyl	4-(1-methylpyrrol-3-yl)-phenyl
392.	ethyl	4-(furan-2-yl)-phenyl
393.	ethyl	4-(furan-3-yl)-phenyl
394.	ethyl	4-(thiophen-2-yl)-phenyl
395.	ethyl	4-(thiophen-3-yl)-phenyl
396.	ethyl	4-(5-propylthien-2-yl)-phenyl
397.	ethyl	4-(pyrazol-1-yl)-phenyl
398.	ethyl	4-(pyrazol-3-yl)-phenyl
399.	ethyl	4-(pyrazol-4-yl)-phenyl
400.	ethyl	4-(1-methyl-1H-pyrazol-4-yl)-phenyl
401.	ethyl	4-(1-ethyl-1H-pyrazol-4-yl)-phenyl
402.	ethyl	4-(1-methyl-1H-pyrazol-5-yl)-phenyl
403.	ethyl	4-(1H-imidazol-2-yl)-phenyl
404.	ethyl	4-(imidazol-1-yl)-phenyl
405.	ethyl	4-(1-methylimidazol-2-yl)-phenyl
406.	ethyl	4-(oxazol-2-yl)-phenyl
407.	ethyl	4-(oxazol-4-yl)-phenyl
408.	ethyl	4-(oxazol-5-yl)-phenyl
409.	ethyl	4-(isoxazol-3-yl)-phenyl
410.	ethyl	4-(isoxazol-4-yl)-phenyl
411.	ethyl	4-(isoxazol-5-yl)-phenyl
412.	ethyl	4-([1,2,3]-triazol-1-yl)-phenyl

No.	R ¹	Ar
413.	ethyl	4-([1,2,4]-triazol-1-yl)-phenyl
414.	ethyl	4-([1,2,3]-triazol-2-yl)-phenyl
415.	ethyl	4-(4H-[1,2,4]-triazol-3-yl)-phenyl
416.	ethyl	4-([1,2,4]-triazol-4-yl)-phenyl
417.	ethyl	4-(2H-[1,2,3]-triazol-4-yl)-phenyl
418.	ethyl	4-(4-methyl-4H-[1,2,4]-triazol-3-yl)-phenyl
419.	ethyl	4-(2-methyl-2H-[1,2,3]-triazol-4-yl)-phenyl
420.	ethyl	4-([1,3,4]-oxadiazol-2-yl)-phenyl
421.	ethyl	4-([1,2,4]-oxadiazol-3-yl)-phenyl
422.	ethyl	4-([1,2,4]-oxadiazol-5-yl)-phenyl
423.	ethyl	4-([1,2,3]-oxadiazol-4-yl)-phenyl
424.	ethyl	4-([1,2,3]-oxadiazol-5-yl)-phenyl
425.	ethyl	4-([1,2,3]-thiadiazol-4-yl)-phenyl
426.	ethyl	4-(1H-tetrazol-5-yl)-phenyl
427.	ethyl	4-(tetrazol-1-yl)-phenyl
428.	ethyl	4-(2-methyl-2H-tetrazol-5-yl)-phenyl
429.	ethyl	4-(1-methyl-1H-tetrazol-5-yl)-phenyl
430.	ethyl	4-furazan-3-yl-phenyl
431.	ethyl	4-(pyrid-2-yl)-phenyl
432.	ethyl	4-(pyrid-3-yl)-phenyl
433.	ethyl	4-(pyrid-4-yl)-phenyl
434.	ethyl	4-(pyrimidin-2-yl)-phenyl
435.	ethyl	4-(pyrimidin-4-yl)-phenyl
436.	ethyl	4-(pyrimidin-5-yl)-phenyl
437.	ethyl	5-isopropylthiophen-2-yl
438.	ethyl	2-chlorothiophen-5-yl
439.	ethyl	2,5-dichlorothiophen-4-yl
440.	ethyl	2,3-dichlorothiophen-5-yl
441.	ethyl	2-chloro-3-nitrothiophen-5-yl
442.	ethyl	2-(phenylsulfonyl)-thiophen-5-yl
443.	ethyl	2-(pyridin-2-yl)thiophen-5-yl
444.	ethyl	2-(5-(trifluoromethyl)isoxazol-3-yl)-thiophen-5-yl
445.	ethyl	2-(2-methylthiazol-4-yl)-thiophen-5-yl
446.	ethyl	1-methyl-1H-imidazol-4-yl
447.	ethyl	1,2-dimethyl-1H-imidazol-4-yl

No.	R ¹	Ar
448.	ethyl	3,5-dimethylisoxazol-4-yl
449.	ethyl	thiazol-2-yl
450.	ethyl	4-methylthiazol-2-yl
451.	ethyl	4-isopropylthiazol-2-yl
452.	ethyl	4-trifluoromethylthiazol-2-yl
453.	ethyl	5-methylthiazol-2-yl
454.	ethyl	5-isopropylthiazol-2-yl
455.	ethyl	5-trifluoromethylthiazol-2-yl
456.	ethyl	2,4-dimethylthiazol-5-yl
457.	ethyl	2-acetamido-4-methylthiazol-5-yl
458.	ethyl	4H-[1,2,4]triazol-3-yl
459.	ethyl	5-methyl-4H-[1,2,4]triazol-3-yl
460.	ethyl	4-methyl-4H-[1,2,4]triazol-3-yl
461.	ethyl	5-isopropyl-4H-[1,2,4]triazol-3-yl
462.	ethyl	5-trifluoromethyl-4H-[1,2,4]triazol-3-yl
463.	ethyl	4,5-dimethyl-4H-[1,2,4]triazol-3-yl
464.	ethyl	5-isopropyl-4-methyl-4H-[1,2,4]triazol-3-yl
465.	ethyl	5-trifluoromethyl-4-methyl-4H-[1,2,4]triazol-3-yl
466.	ethyl	[1,3,4]thiadiazol-2-yl
467.	ethyl	5-methyl-[1,3,4]thiadiazol-2-yl
468.	ethyl	5-isopropyl-[1,3,4]thiadiazol-2-yl
469.	ethyl	5-trifluoromethyl-[1,3,4]thiadiazol-2-yl
470.	ethyl	3-bromo-2-chloropyrid-5-yl
471.	ethyl	2-(4-morpholino)-pyrid-5-yl
472.	ethyl	2-phenoxy-pyrid-5-yl
473.	ethyl	(2-isopropyl)-pyrimidin-5-yl
474.	ethyl	(5-isopropyl)-pyrimidin-2-yl
475.	ethyl	8-quinolyl
476.	ethyl	5-isoquinolyl
477.	ethyl	2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl
478.	ethyl	5-chloro-3-methylbenzothiophen-2-yl
479.	ethyl	3,4-dihydro-4-methyl-2H-benzo[b][1,4]oxazinyl
480.	ethyl	benzothiazol-6-yl
481.	ethyl	benzo[2,1,3]oxadiazol-4-yl

No.	R ¹	Ar
482.	ethyl	5-chlorobenzo[2,1,3]oxadiazol-4-yl
483.	ethyl	7-chlorobenzo[2,1,3]oxadiazol-4-yl
484.	ethyl	benzo[2,1,3]thiadiazol-4-yl
485.	methyl	4-methylphenyl
486.	methyl	4-ethylphenyl
487.	methyl	4-propylphenyl
488.	methyl	4-isopropylphenyl
489.	methyl	4-sec-butylphenyl
490.	methyl	4-isobutylphenyl
491.	methyl	4-(1,1-dimethylpropyl)-phenyl
492.	methyl	4-vinylphenyl
493.	methyl	4-isopropenylphenyl
494.	methyl	4-fluorophenyl
495.	methyl	4-chlorophenyl
496.	methyl	4-bromophenyl
497.	methyl	4-(fluoromethyl)phenyl
498.	methyl	3-(fluoromethyl)phenyl
499.	methyl	2-(fluoromethyl)phenyl
500.	methyl	4-(difluoromethyl)phenyl
501.	methyl	3-(difluoromethyl)phenyl
502.	methyl	2-(difluoromethyl)phenyl
503.	methyl	4-(trifluoromethyl)phenyl
504.	methyl	3-(trifluoromethyl)phenyl
505.	methyl	2-(trifluoromethyl)phenyl
506.	methyl	4-(1-fluoroethyl)-phenyl
507.	methyl	4-((S)-1-fluoroethyl)-phenyl
508.	methyl	4-((R)-1-fluoroethyl)-phenyl
509.	methyl	4-(2-fluoroethyl)-phenyl
510.	methyl	4-(1,1-difluoroethyl)-phenyl
511.	methyl	4-(2,2-difluoroethyl)-phenyl
512.	methyl	4-(2,2,2-trifluoroethyl)-phenyl
513.	methyl	4-(3-fluoropropyl)-phenyl
514.	methyl	4-(2-fluoropropyl)-phenyl
515.	methyl	4-((S)-2-fluoropropyl)-phenyl
516.	methyl	4-((R)-2-fluoropropyl)-phenyl

No.	R ¹	Ar
517.	methyl	4-(3,3-difluoropropyl)-phenyl
518.	methyl	4-(3,3,3-trifluoropropyl)-phenyl
519.	methyl	4-(1-fluoro-1-methylethyl)-phenyl
520.	methyl	4-(2-fluoro-1-methylethyl)-phenyl
521.	methyl	4-((S)-2-fluoro-1-methylethyl)-phenyl
522.	methyl	4-((R)-2-fluoro-1-methylethyl)-phenyl
523.	methyl	4-(2,2-difluoro-1-methylethyl)-phenyl
524.	methyl	4-((S)-2,2-difluoro-1-methylethyl)-phenyl
525.	methyl	4-((R)-2,2-difluoro-1-methylethyl)-phenyl
526.	methyl	4-(2,2,2-trifluoro-1-methylethyl)-phenyl
527.	methyl	4-((S)-2,2,2-trifluoro-1-methylethyl)-phenyl
528.	methyl	4-((R)-2,2,2-trifluoro-1-methylethyl)-phenyl
529.	methyl	4-(2-fluoro-1-fluoromethylethyl)-phenyl
530.	methyl	4-(1-difluoromethyl-2,2-difluoroethyl)-phenyl
531.	methyl	4-(1,1-dimethyl-2-fluoroethyl)-phenyl
532.	methyl	4-methoxyphenyl
533.	methyl	4-ethoxyphenyl
534.	methyl	4-propoxyphenyl
535.	methyl	4-isopropoxyphenyl
536.	methyl	4-butoxyphenyl
537.	methyl	4-(fluoromethoxy)-phenyl
538.	methyl	4-(difluoromethoxy)-phenyl
539.	methyl	4-(trifluoromethoxy)-phenyl
540.	methyl	3-(trifluoromethoxy)-phenyl
541.	methyl	4-(2-fluoroethoxy)-phenyl
542.	methyl	4-(2,2-difluoroethoxy)-phenyl
543.	methyl	4-(2,2,2-trifluoroethoxy)-phenyl
544.	methyl	4-(1,1,2,2-tetrafluoroethoxy)-phenyl
545.	methyl	4-cyclopropylphenyl
546.	methyl	4-cyclobutylphenyl
547.	methyl	4-cyclopentylphenyl
548.	methyl	4-(2,2-difluorocyclopropyl)-phenyl
549.	methyl	3,4-difluorophenyl
550.	methyl	4-bromo-3-fluorophenyl
551.	methyl	4-bromo-2-fluorophenyl

No.	R ¹	Ar
552.	methyl	4-bromo-2,5-difluorophenyl
553.	methyl	2-fluoro-4-isopropylphenyl
554.	methyl	3-fluoro-4-isopropylphenyl
555.	methyl	4-(1-hydroxy-1-methylethyl)-phenyl
556.	methyl	4-(2-hydroxy-2-methylpropyl)-phenyl
557.	methyl	4-acetylphenyl
558.	methyl	4-carboxyphenyl
559.	methyl	4-cyanophenyl
560.	methyl	4-hydroxyphenyl
561.	methyl	4-(O-benzyl)-phenyl
562.	methyl	4-(2-methoxyethoxy)-phenyl
563.	methyl	4-(CH ₂ -N(CH ₃) ₂)-phenyl
564.	methyl	4-(NH-CO-NH ₂)-phenyl
565.	methyl	4-(methylsulfanyl)-phenyl
566.	methyl	4-(fluoromethylsulfanyl)-phenyl
567.	methyl	4-(difluoromethylsulfanyl)-phenyl
568.	methyl	4-(trifluoromethylsulfanyl)-phenyl
569.	methyl	4-(methylsulfonyl)-phenyl
570.	methyl	4-(N-methoxy-N-methyl-amino)-phenyl
571.	methyl	4-(methoxyamino)-phenyl
572.	methyl	4-(ethoxyamino)-phenyl
573.	methyl	4-(N-methylaminooxy)-phenyl
574.	methyl	4-(N,N-dimethylaminooxy)-phenyl
575.	methyl	4-(azetidin-1-yl)-phenyl
576.	methyl	4-(2-methylazetidin-1-yl)-phenyl
577.	methyl	4-((S)-2-methylazetidin-1-yl)-phenyl
578.	methyl	4-((R)-2-methylazetidin-1-yl)-phenyl
579.	methyl	4-(3-fluoroazetidin-1-yl)-phenyl
580.	methyl	4-(3-methoxyazetidin-1-yl)-phenyl
581.	methyl	4-(3-hydroxyazetidin-1-yl)-phenyl
582.	methyl	4-(pyrrolidin-1-yl)-phenyl
583.	methyl	4-(pyrrolidin-2-yl)-phenyl
584.	methyl	4-((S)-pyrrolidin-2-yl)-phenyl
585.	methyl	4-((R)-pyrrolidin-2-yl)-phenyl
586.	methyl	4-(pyrrolidin-3-yl)-phenyl

No.	R ¹	Ar
587.	methyl	4-((S)-pyrrolidin-3-yl)-phenyl
588.	methyl	4-((R)-pyrrolidin-3-yl)-phenyl
589.	methyl	4-(2-fluoropyrrolidin-1-yl)-phenyl
590.	methyl	4-((S)-2-fluoropyrrolidin-1-yl)-phenyl
591.	methyl	4-((R)-2-fluoropyrrolidin-1-yl)-phenyl
592.	methyl	4-(3-fluoropyrrolidin-1-yl)-phenyl
593.	methyl	4-((S)-3-fluoropyrrolidin-1-yl)-phenyl
594.	methyl	4-((R)-3-fluoropyrrolidin-1-yl)-phenyl
595.	methyl	4-(2,2-difluoropyrrolidin-1-yl)-phenyl
596.	methyl	4-(3,3-difluoropyrrolidin-1-yl)-phenyl
597.	methyl	4-(2-methylpyrrolidin-1-yl)-phenyl
598.	methyl	4-((S)-2-methylpyrrolidin-1-yl)-phenyl
599.	methyl	4-((R)-2-methylpyrrolidin-1-yl)-phenyl
600.	methyl	4-(3-methylpyrrolidin-1-yl)-phenyl
601.	methyl	4-((S)-3-methylpyrrolidin-1-yl)-phenyl
602.	methyl	4-((R)-3-methylpyrrolidin-1-yl)-phenyl
603.	methyl	4-(1-methylpyrrolidin-2-yl)-phenyl
604.	methyl	4-((S)-1-methylpyrrolidin-2-yl)-phenyl
605.	methyl	4-((R)-1-methylpyrrolidin-2-yl)-phenyl
606.	methyl	4-(1-methylpyrrolidin-3-yl)-phenyl
607.	methyl	4-((S)-1-methylpyrrolidin-3-yl)-phenyl
608.	methyl	4-((R)-1-methylpyrrolidin-3-yl)-phenyl
609.	methyl	4-(2,2-dimethylpyrrolidin-1-yl)-phenyl
610.	methyl	4-(3,3-dimethylpyrrolidin-1-yl)-phenyl
611.	methyl	4-(2-trifluoromethylpyrrolidin-1-yl)-phenyl
612.	methyl	4-((S)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
613.	methyl	4-((R)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
614.	methyl	4-(3-trifluoromethylpyrrolidin-1-yl)-phenyl
615.	methyl	4-((S)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
616.	methyl	4-((R)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
617.	methyl	4-(2-oxopyrrolidin-1-yl)-phenyl
618.	methyl	4-(2-oxo-oxazolidin-3-yl)-phenyl
619.	methyl	4-(piperidin-1-yl)-phenyl
620.	methyl	4-(2-methylpiperidin-1-yl)-phenyl
621.	methyl	4-((S)-2-methylpiperidin-1-yl)-phenyl

No.	R ¹	Ar
622.	methyl	4-((R)-2-methylpiperidin-1-yl)-phenyl
623.	methyl	4-(piperazin-1-yl)-phenyl
624.	methyl	4-(4-methylpiperazin-1-yl)-phenyl
625.	methyl	4-(morpholin-4-yl)-phenyl
626.	methyl	4-(thiomorpholin-4-yl)-phenyl
627.	methyl	4-(1-oxo-thiomorpholin-4-yl)-phenyl
628.	methyl	4-(1,1-dioxo-thiomorpholin-4-yl)-phenyl
629.	methyl	4-(pyrrol-1-yl)-phenyl
630.	methyl	4-(pyrrol-2-yl)-phenyl
631.	methyl	4-(pyrrol-3-yl)-phenyl
632.	methyl	4-(1-methylpyrrol-2-yl)-phenyl
633.	methyl	4-(1-methylpyrrol-3-yl)-phenyl
634.	methyl	4-(furan-2-yl)-phenyl
635.	methyl	4-(furan-3-yl)-phenyl
636.	methyl	4-(thiophen-2-yl)-phenyl
637.	methyl	4-(thiophen-3-yl)-phenyl
638.	methyl	4-(5-propylthien-2-yl)-phenyl
639.	methyl	4-(pyrazol-1-yl)-phenyl
640.	methyl	4-(pyrazol-3-yl)-phenyl
641.	methyl	4-(pyrazol-4-yl)-phenyl
642.	methyl	4-(1-methyl-1H-pyrazol-4-yl)-phenyl
643.	methyl	4-(1-ethyl-1H-pyrazol-4-yl)-phenyl
644.	methyl	4-(1-methyl-1H-pyrazol-5-yl)-phenyl
645.	methyl	4-(1H-imidazol-2-yl)-phenyl
646.	methyl	4-(imidazol-1-yl)-phenyl
647.	methyl	4-(1-methylimidazol-2-yl)-phenyl
648.	methyl	4-(oxazol-2-yl)-phenyl
649.	methyl	4-(oxazol-4-yl)-phenyl
650.	methyl	4-(oxazol-5-yl)-phenyl
651.	methyl	4-(isoxazol-3-yl)-phenyl
652.	methyl	4-(isoxazol-4-yl)-phenyl
653.	methyl	4-(isoxazol-5-yl)-phenyl
654.	methyl	4-([1,2,3]-triazol-1-yl)-phenyl
655.	methyl	4-([1,2,4]-triazol-1-yl)-phenyl
656.	methyl	4-([1,2,3]-triazol-2-yl)-phenyl

No.	R ¹	Ar
657.	methyl	4-(4H-[1,2,4]-triazol-3-yl)-phenyl
658.	methyl	4-([1,2,4]-triazol-4-yl)-phenyl
659.	methyl	4-(2H-[1,2,3]-triazol-4-yl)-phenyl
660.	methyl	4-(4-methyl-4H-[1,2,4]-triazol-3-yl)-phenyl
661.	methyl	4-(2-methyl-2H-[1,2,3]-triazol-4-yl)-phenyl
662.	methyl	4-([1,3,4]-oxadiazol-2-yl)-phenyl
663.	methyl	4-([1,2,4]-oxadiazol-3-yl)-phenyl
664.	methyl	4-([1,2,4]-oxadiazol-5-yl)-phenyl
665.	methyl	4-([1,2,3]-oxadiazol-4-yl)-phenyl
666.	methyl	4-([1,2,3]-oxadiazol-5-yl)-phenyl
667.	methyl	4-([1,2,3]-thiadiazol-4-yl)-phenyl
668.	methyl	4-(1H-tetrazol-5-yl)-phenyl
669.	methyl	4-(tetrazol-1-yl)-phenyl
670.	methyl	4-(2-methyl-2H-tetrazol-5-yl)-phenyl
671.	methyl	4-(1-methyl-1H-tetrazol-5-yl)-phenyl
672.	methyl	4-furazan-3-yl-phenyl
673.	methyl	4-(pyrid-2-yl)-phenyl
674.	methyl	4-(pyrid-3-yl)-phenyl
675.	methyl	4-(pyrid-4-yl)-phenyl
676.	methyl	4-(pyrimidin-2-yl)-phenyl
677.	methyl	4-(pyrimidin-4-yl)-phenyl
678.	methyl	4-(pyrimidin-5-yl)-phenyl
679.	methyl	5-isopropylthiophen-2-yl
680.	methyl	2-chlorothiophen-5-yl
681.	methyl	2,5-dichlorothiophen-4-yl
682.	methyl	2,3-dichlorothiophen-5-yl
683.	methyl	2-chloro-3-nitrothiophen-5-yl
684.	methyl	2-(phenylsulfonyl)-thiophen-5-yl
685.	methyl	2-(pyridin-2-yl)thiophen-5-yl
686.	methyl	2-(5-(trifluoromethyl)isoxazol-3-yl)-thiophen-5-yl
687.	methyl	2-(2-methylthiazol-4-yl)-thiophen-5-yl
688.	methyl	1-methyl-1H-imidazol-4-yl
689.	methyl	1,2-dimethyl-1H-imidazol-4-yl
690.	methyl	3,5-dimethylisoxazol-4-yl
691.	methyl	thiazol-2-yl

41

No.	R ¹	Ar
692.	methyl	4-methylthiazol-2-yl
693.	methyl	4-isopropylthiazol-2-yl
694.	methyl	4-trifluoromethylthiazol-2-yl
695.	methyl	5-methylthiazol-2-yl
696.	methyl	5-isopropylthiazol-2-yl
697.	methyl	5-trifluoromethylthiazol-2-yl
698.	methyl	2,4-dimethylthiazol-5-yl
699.	methyl	2-acetamido-4-methylthiazol-5-yl
700.	methyl	4H-[1,2,4]triazol-3-yl
701.	methyl	5-methyl-4H-[1,2,4]triazol-3-yl
702.	methyl	4-methyl-4H-[1,2,4]triazol-3-yl
703.	methyl	5-isopropyl-4H-[1,2,4]triazol-3-yl
704.	methyl	5-trifluoromethyl-4H-[1,2,4]triazol-3-yl
705.	methyl	4,5-dimethyl-4H-[1,2,4]triazol-3-yl
706.	methyl	5-isopropyl-4-methyl-4H-[1,2,4]triazol-3-yl
707.	methyl	5-trifluoromethyl-4-methyl-4H-[1,2,4]triazol-3-yl
708.	methyl	[1,3,4]thiadiazol-2-yl
709.	methyl	5-methyl-[1,3,4]thiadiazol-2-yl
710.	methyl	5-isopropyl-[1,3,4]thiadiazol-2-yl
711.	methyl	5-trifluoromethyl-[1,3,4]thiadiazol-2-yl
712.	methyl	3-bromo-2-chloropyrid-5-yl
713.	methyl	2-(4-morpholino)-pyrid-5-yl
714.	methyl	2-phenoxy-pyrid-5-yl
715.	methyl	(2-isopropyl)-pyrimidin-5-yl
716.	methyl	(5-isopropyl)-pyrimidin-2-yl
717.	methyl	8-quinolyl
718.	methyl	5-isoquinolyl
719.	methyl	2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl
720.	methyl	5-chloro-3-methylbenzothiophen-2-yl
721.	methyl	3,4-dihydro-4-methyl-2H-benzo[b][1,4]oxazinyl
722.	methyl	benzothiazol-6-yl
723.	methyl	benzo[2,1,3]oxadiazol-4-yl
724.	methyl	5-chlorobenzo[2,1,3]oxadiazol-4-yl
725.	methyl	7-chlorobenzo[2,1,3]oxadiazol-4-yl

No.	R ¹	Ar
726.	methyl	benzo[2,1,3]thiadiazol-4-yl
727.	H	4-methylphenyl
728.	H	4-ethylphenyl
729.	H	4-propylphenyl
730.	H	4-isopropylphenyl
731.	H	4-sec-butylphenyl
732.	H	4-isobutylphenyl
733.	H	4-(1,1-dimethylpropyl)-phenyl
734.	H	4-vinylphenyl
735.	H	4-isopropenylphenyl
736.	H	4-fluorophenyl
737.	H	4-chlorophenyl
738.	H	4-bromophenyl
739.	H	4-(fluoromethyl)phenyl
740.	H	3-(fluoromethyl)phenyl
741.	H	2-(fluoromethyl)phenyl
742.	H	4-(difluoromethyl)phenyl
743.	H	3-(difluoromethyl)phenyl
744.	H	2-(difluoromethyl)phenyl
745.	H	4-(trifluoromethyl)phenyl
746.	H	3-(trifluoromethyl)phenyl
747.	H	2-(trifluoromethyl)phenyl
748.	H	4-(1-fluoroethyl)-phenyl
749.	H	4-((S)-1-fluoroethyl)-phenyl
750.	H	4-((R)-1-fluoroethyl)-phenyl
751.	H	4-(2-fluoroethyl)-phenyl
752.	H	4-(1,1-difluoroethyl)-phenyl
753.	H	4-(2,2-difluoroethyl)-phenyl
754.	H	4-(2,2,2-trifluoroethyl)-phenyl
755.	H	4-(3-fluoropropyl)-phenyl
756.	H	4-(2-fluoropropyl)-phenyl
757.	H	4-((S)-2-fluoropropyl)-phenyl
758.	H	4-((R)-2-fluoropropyl)-phenyl
759.	H	4-(3,3-difluoropropyl)-phenyl
760.	H	4-(3,3,3-trifluoropropyl)-phenyl

No.	R ¹	Ar
761.	H	4-(1-fluoro-1-methylethyl)-phenyl
762.	H	4-(2-fluoro-1-methylethyl)-phenyl
763.	H	4-((S)-2-fluoro-1-methylethyl)-phenyl
764.	H	4-((R)-2-fluoro-1-methylethyl)-phenyl
765.	H	4-(2,2-difluoro-1-methylethyl)-phenyl
766.	H	4-((S)-2,2-difluoro-1-methylethyl)-phenyl
767.	H	4-((R)-2,2-difluoro-1-methylethyl)-phenyl
768.	H	4-(2,2,2-trifluoro-1-methylethyl)-phenyl
769.	H	4-((S)-2,2,2-trifluoro-1-methylethyl)-phenyl
770.	H	4-((R)-2,2,2-trifluoro-1-methylethyl)-phenyl
771.	H	4-(2-fluoro-1-fluoromethylethyl)-phenyl
772.	H	4-(1-difluoromethyl-2,2-difluoroethyl)-phenyl
773.	H	4-(1,1-dimethyl-2-fluoroethyl)-phenyl
774.	H	4-methoxyphenyl
775.	H	4-ethoxyphenyl
776.	H	4-propoxyphenyl
777.	H	4-isopropoxyphenyl
778.	H	4-butoxyphenyl
779.	H	4-(fluoromethoxy)-phenyl
780.	H	4-(difluoromethoxy)-phenyl
781.	H	4-(trifluoromethoxy)-phenyl
782.	H	3-(trifluoromethoxy)-phenyl
783.	H	4-(2-fluoroethoxy)-phenyl
784.	H	4-(2,2-difluoroethoxy)-phenyl
785.	H	4-(2,2,2-trifluoroethoxy)-phenyl
786.	H	4-(1,1,2,2-tetrafluoroethoxy)-phenyl
787.	H	4-cyclopropylphenyl
788.	H	4-cyclobutylphenyl
789.	H	4-cyclopentylphenyl
790.	H	4-(2,2-difluorocyclopropyl)-phenyl
791.	H	3,4-difluorophenyl
792.	H	4-bromo-3-fluorophenyl
793.	H	4-bromo-2-fluorophenyl
794.	H	4-bromo-2,5-difluorophenyl
795.	H	2-fluoro-4-isopropylphenyl

No.	R ¹	Ar
796.	H	3-fluoro-4-isopropylphenyl
797.	H	4-(1-hydroxy-1-methylethyl)-phenyl
798.	H	4-(2-hydroxy-2-methylpropyl)-phenyl
799.	H	4-acetylphenyl
800.	H	4-carboxyphenyl
801.	H	4-cyanophenyl
802.	H	4-hydroxyphenyl
803.	H	4-(O-benzyl)-phenyl
804.	H	4-(2-methoxyethoxy)-phenyl
805.	H	4-(CH ₂ -N(CH ₃) ₂)-phenyl
806.	H	4-(NH-CO-NH ₂)-phenyl
807.	H	4-(methylsulfanyl)-phenyl
808.	H	4-(fluoromethylsulfanyl)-phenyl
809.	H	4-(difluoromethylsulfanyl)-phenyl
810.	H	4-(trifluoromethylsulfanyl)-phenyl
811.	H	4-(methylsulfonyl)-phenyl
812.	H	4-(N-methoxy-N-methyl-amino)-phenyl
813.	H	4-(methoxyamino)-phenyl
814.	H	4-(ethoxyamino)-phenyl
815.	H	4-(N-methylaminooxy)-phenyl
816.	H	4-(N,N-dimethylaminooxy)-phenyl
817.	H	4-(azetidin-1-yl)-phenyl
818.	H	4-(2-methylazetidin-1-yl)-phenyl
819.	H	4-((S)-2-methylazetidin-1-yl)-phenyl
820.	H	4-((R)-2-methylazetidin-1-yl)-phenyl
821.	H	4-(3-fluoroazetidin-1-yl)-phenyl
822.	H	4-(3-methoxyazetidin-1-yl)-phenyl
823.	H	4-(3-hydroxyazetidin-1-yl)-phenyl
824.	H	4-(pyrrolidin-1-yl)-phenyl
825.	H	4-(pyrrolidin-2-yl)-phenyl
826.	H	4-((S)-pyrrolidin-2-yl)-phenyl
827.	H	4-((R)-pyrrolidin-2-yl)-phenyl
828.	H	4-(pyrrolidin-3-yl)-phenyl
829.	H	4-((S)-pyrrolidin-3-yl)-phenyl
830.	H	4-((R)-pyrrolidin-3-yl)-phenyl

No.	R ¹	Ar
831.	H	4-(2-fluoropyrrolidin-1-yl)-phenyl
832.	H	4-((S)-2-fluoropyrrolidin-1-yl)-phenyl
833.	H	4-((R)-2-fluoropyrrolidin-1-yl)-phenyl
834.	H	4-(3-fluoropyrrolidin-1-yl)-phenyl
835.	H	4-((S)-3-fluoropyrrolidin-1-yl)-phenyl
836.	H	4-((R)-3-fluoropyrrolidin-1-yl)-phenyl
837.	H	4-(2,2-difluoropyrrolidin-1-yl)-phenyl
838.	H	4-(3,3-difluoropyrrolidin-1-yl)-phenyl
839.	H	4-(2-methylpyrrolidin-1-yl)-phenyl
840.	H	4-((S)-2-methylpyrrolidin-1-yl)-phenyl
841.	H	4-((R)-2-methylpyrrolidin-1-yl)-phenyl
842.	H	4-(3-methylpyrrolidin-1-yl)-phenyl
843.	H	4-((S)-3-methylpyrrolidin-1-yl)-phenyl
844.	H	4-((R)-3-methylpyrrolidin-1-yl)-phenyl
845.	H	4-(1-methylpyrrolidin-2-yl)-phenyl
846.	H	4-((S)-1-methylpyrrolidin-2-yl)-phenyl
847.	H	4-((R)-1-methylpyrrolidin-2-yl)-phenyl
848.	H	4-(1-methylpyrrolidin-3-yl)-phenyl
849.	H	4-((S)-1-methylpyrrolidin-3-yl)-phenyl
850.	H	4-((R)-1-methylpyrrolidin-3-yl)-phenyl
851.	H	4-(2,2-dimethylpyrrolidin-1-yl)-phenyl
852.	H	4-(3,3-dimethylpyrrolidin-1-yl)-phenyl
853.	H	4-(2-trifluoromethylpyrrolidin-1-yl)-phenyl
854.	H	4-((S)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
855.	H	4-((R)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
856.	H	4-(3-trifluoromethylpyrrolidin-1-yl)-phenyl
857.	H	4-((S)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
858.	H	4-((R)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
859.	H	4-(2-oxopyrrolidin-1-yl)-phenyl
860.	H	4-(2-oxo-oxazolidin-3-yl)-phenyl
861.	H	4-(piperidin-1-yl)-phenyl
862.	H	4-(2-methylpiperidin-1-yl)-phenyl
863.	H	4-((S)-2-methylpiperidin-1-yl)-phenyl
864.	H	4-((R)-2-methylpiperidin-1-yl)-phenyl
865.	H	4-(piperazin-1-yl)-phenyl

No.	R ¹	Ar
866.	H	4-(4-methylpiperazin-1-yl)-phenyl
867.	H	4-(morpholin-4-yl)-phenyl
868.	H	4-(thiomorpholin-4-yl)-phenyl
869.	H	4-(1-oxo-thiomorpholin-4-yl)-phenyl
870.	H	4-(1,1-dioxo-thiomorpholin-4-yl)-phenyl
871.	H	4-(pyrrol-1-yl)-phenyl
872.	H	4-(pyrrol-2-yl)-phenyl
873.	H	4-(pyrrol-3-yl)-phenyl
874.	H	4-(1-methylpyrrol-2-yl)-phenyl
875.	H	4-(1-methylpyrrol-3-yl)-phenyl
876.	H	4-(furan-2-yl)-phenyl
877.	H	4-(furan-3-yl)-phenyl
878.	H	4-(thiophen-2-yl)-phenyl
879.	H	4-(thiophen-3-yl)-phenyl
880.	H	4-(5-propylthien-2-yl)-phenyl
881.	H	4-(pyrazol-1-yl)-phenyl
882.	H	4-(pyrazol-3-yl)-phenyl
883.	H	4-(pyrazol-4-yl)-phenyl
884.	H	4-(1-methyl-1H-pyrazol-4-yl)-phenyl
885.	H	4-(1-ethyl-1H-pyrazol-4-yl)-phenyl
886.	H	4-(1-methyl-1H-pyrazol-5-yl)-phenyl
887.	H	4-(1H-imidazol-2-yl)-phenyl
888.	H	4-(imidazol-1-yl)-phenyl
889.	H	4-(1-methylimidazol-2-yl)-phenyl
890.	H	4-(oxazol-2-yl)-phenyl
891.	H	4-(oxazol-4-yl)-phenyl
892.	H	4-(oxazol-5-yl)-phenyl
893.	H	4-(isoxazol-3-yl)-phenyl
894.	H	4-(isoxazol-4-yl)-phenyl
895.	H	4-(isoxazol-5-yl)-phenyl
896.	H	4-([1,2,3]-triazol-1-yl)-phenyl
897.	H	4-([1,2,4]-triazol-1-yl)-phenyl
898.	H	4-([1,2,3]-triazol-2-yl)-phenyl
899.	H	4-(4H-[1,2,4]-triazol-3-yl)-phenyl
900.	H	4-([1,2,4]-triazol-4-yl)-phenyl

No.	R ¹	Ar
901.	H	4-(2H-[1,2,3]-triazol-4-yl)-phenyl
902.	H	4-(4-methyl-4H-[1,2,4]-triazol-3-yl)-phenyl
903.	H	4-(2-methyl-2H-[1,2,3]-triazol-4-yl)-phenyl
904.	H	4-([1,3,4]-oxadiazol-2-yl)-phenyl
905.	H	4-([1,2,4]-oxadiazol-3-yl)-phenyl
906.	H	4-([1,2,4]-oxadiazol-5-yl)-phenyl
907.	H	4-([1,2,3]-oxadiazol-4-yl)-phenyl
908.	H	4-([1,2,3]-oxadiazol-5-yl)-phenyl
909.	H	4-([1,2,3]-thiadiazol-4-yl)-phenyl
910.	H	4-(1H-tetrazol-5-yl)-phenyl
911.	H	4-(tetrazol-1-yl)-phenyl
912.	H	4-(2-methyl-2H-tetrazol-5-yl)-phenyl
913.	H	4-(1-methyl-1H-tetrazol-5-yl)-phenyl
914.	H	4-furazan-3-yl-phenyl
915.	H	4-(pyrid-2-yl)-phenyl
916.	H	4-(pyrid-3-yl)-phenyl
917.	H	4-(pyrid-4-yl)-phenyl
918.	H	4-(pyrimidin-2-yl)-phenyl
919.	H	4-(pyrimidin-4-yl)-phenyl
920.	H	4-(pyrimidin-5-yl)-phenyl
921.	H	5-isopropylthiophen-2-yl
922.	H	2-chlorothiophen-5-yl
923.	H	2,5-dichlorothiophen-4-yl
924.	H	2,3-dichlorothiophen-5-yl
925.	H	2-chloro-3-nitrothiophen-5-yl
926.	H	2-(phenylsulfonyl)-thiophen-5-yl
927.	H	2-(pyridin-2-yl)thiophen-5-yl
928.	H	2-(5-(trifluoromethyl)isoxazol-3-yl)-thiophen-5-yl
929.	H	2-(2-methylthiazol-4-yl)-thiophen-5-yl
930.	H	1-methyl-1H-imidazol-4-yl
931.	H	1,2-dimethyl-1H-imidazol-4-yl
932.	H	3,5-dimethylisoxazol-4-yl
933.	H	thiazol-2-yl
934.	H	4-methylthiazol-2-yl
935.	H	4-isopropylthiazol-2-yl

No.	R ¹	Ar
936.	H	4-trifluoromethylthiazol-2-yl
937.	H	5-methylthiazol-2-yl
938.	H	5-isopropylthiazol-2-yl
939.	H	5-trifluoromethylthiazol-2-yl
940.	H	2,4-dimethylthiazol-5-yl
941.	H	2-acetamido-4-methylthiazol-5-yl
942.	H	4H-[1,2,4]triazol-3-yl
943.	H	5-methyl-4H-[1,2,4]triazol-3-yl
944.	H	4-methyl-4H-[1,2,4]triazol-3-yl
945.	H	5-isopropyl-4H-[1,2,4]triazol-3-yl
946.	H	5-trifluoromethyl-4H-[1,2,4]triazol-3-yl
947.	H	4,5-dimethyl-4H-[1,2,4]triazol-3-yl
948.	H	5-isopropyl-4-methyl-4H-[1,2,4]triazol-3-yl
949.	H	5-trifluoromethyl-4-methyl-4H-[1,2,4]triazol-3-yl
950.	H	[1,3,4]thiadiazol-2-yl
951.	H	5-methyl-[1,3,4]thiadiazol-2-yl
952.	H	5-isopropyl-[1,3,4]thiadiazol-2-yl
953.	H	5-trifluoromethyl-[1,3,4]thiadiazol-2-yl
954.	H	3-bromo-2-chloropyrid-5-yl
955.	H	2-(4-morpholino)-pyrid-5-yl
956.	H	2-phenoxy-pyrid-5-yl
957.	H	(2-isopropyl)-pyrimidin-5-yl
958.	H	(5-isopropyl)-pyrimidin-2-yl
959.	H	8-quinolyl
960.	H	5-isoquinolyl
961.	H	2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl
962.	H	5-chloro-3-methylbenzothiophen-2-yl
963.	H	3,4-dihydro-4-methyl-2H-benzo[b][1,4]oxazinyl
964.	H	benzothiazol-6-yl
965.	H	benzo[2,1,3]oxadiazol-4-yl
966.	H	5-chlorobenzo[2,1,3]oxadiazol-4-yl
967.	H	7-chlorobenzo[2,1,3]oxadiazol-4-yl
968.	H	benzo[2,1,3]thiadiazol-4-yl
969.	3-fluoropropyl	4-methylphenyl

No.	R ¹	Ar
970.	3-fluoropropyl	4-ethylphenyl
971.	3-fluoropropyl	4-propylphenyl
972.	3-fluoropropyl	4-isopropylphenyl
973.	3-fluoropropyl	4-sec-butylphenyl
974.	3-fluoropropyl	4-isobutylphenyl
975.	3-fluoropropyl	4-(1,1-dimethylpropyl)-phenyl
976.	3-fluoropropyl	4-vinylphenyl
977.	3-fluoropropyl	4-isopropenylphenyl
978.	3-fluoropropyl	4-fluorophenyl
979.	3-fluoropropyl	4-chlorophenyl
980.	3-fluoropropyl	4-bromophenyl
981.	3-fluoropropyl	4-(fluoromethyl)phenyl
982.	3-fluoropropyl	3-(fluoromethyl)phenyl
983.	3-fluoropropyl	2-(fluoromethyl)phenyl
984.	3-fluoropropyl	4-(difluoromethyl)phenyl
985.	3-fluoropropyl	3-(difluoromethyl)phenyl
986.	3-fluoropropyl	2-(difluoromethyl)phenyl
987.	3-fluoropropyl	4-(trifluoromethyl)phenyl
988.	3-fluoropropyl	3-(trifluoromethyl)phenyl
989.	3-fluoropropyl	2-(trifluoromethyl)phenyl
990.	3-fluoropropyl	4-(1-fluoroethyl)-phenyl
991.	3-fluoropropyl	4-((S)-1-fluoroethyl)-phenyl
992.	3-fluoropropyl	4-((R)-1-fluoroethyl)-phenyl
993.	3-fluoropropyl	4-(2-fluoroethyl)-phenyl
994.	3-fluoropropyl	4-(1,1-difluoroethyl)-phenyl
995.	3-fluoropropyl	4-(2,2-difluoroethyl)-phenyl
996.	3-fluoropropyl	4-(2,2,2-trifluoroethyl)-phenyl
997.	3-fluoropropyl	4-(3-fluoropropyl)-phenyl
998.	3-fluoropropyl	4-(2-fluoropropyl)-phenyl
999.	3-fluoropropyl	4-((S)-2-fluoropropyl)-phenyl
1000.	3-fluoropropyl	4-((R)-2-fluoropropyl)-phenyl
1001.	3-fluoropropyl	4-(3,3-difluoropropyl)-phenyl
1002.	3-fluoropropyl	4-(3,3,3-trifluoropropyl)-phenyl
1003.	3-fluoropropyl	4-(1-fluoro-1-methylethyl)-phenyl
1004.	3-fluoropropyl	4-(2-fluoro-1-methylethyl)-phenyl

No.	R ¹	Ar
1005.	3-fluoropropyl	4-((S)-2-fluoro-1-methylethyl)-phenyl
1006.	3-fluoropropyl	4-((R)-2-fluoro-1-methylethyl)-phenyl
1007.	3-fluoropropyl	4-(2,2-difluoro-1-methylethyl)-phenyl
1008.	3-fluoropropyl	4-((S)-2,2-difluoro-1-methylethyl)-phenyl
1009.	3-fluoropropyl	4-((R)-2,2-difluoro-1-methylethyl)-phenyl
1010.	3-fluoropropyl	4-(2,2,2-trifluoro-1-methylethyl)-phenyl
1011.	3-fluoropropyl	4-((S)-2,2,2-trifluoro-1-methylethyl)-phenyl
1012.	3-fluoropropyl	4-((R)-2,2,2-trifluoro-1-methylethyl)-phenyl
1013.	3-fluoropropyl	4-(2-fluoro-1-fluoromethylethyl)-phenyl
1014.	3-fluoropropyl	4-(1-difluoromethyl-2,2-difluoroethyl)-phenyl
1015.	3-fluoropropyl	4-(1,1-dimethyl-2-fluoroethyl)-phenyl
1016.	3-fluoropropyl	4-methoxyphenyl
1017.	3-fluoropropyl	4-ethoxyphenyl
1018.	3-fluoropropyl	4-propoxyphenyl
1019.	3-fluoropropyl	4-isopropoxyphenyl
1020.	3-fluoropropyl	4-butoxyphenyl
1021.	3-fluoropropyl	4-(fluoromethoxy)-phenyl
1022.	3-fluoropropyl	4-(difluoromethoxy)-phenyl
1023.	3-fluoropropyl	4-(trifluoromethoxy)-phenyl
1024.	3-fluoropropyl	3-(trifluoromethoxy)-phenyl
1025.	3-fluoropropyl	4-(2-fluoroethoxy)-phenyl
1026.	3-fluoropropyl	4-(2,2-difluoroethoxy)-phenyl
1027.	3-fluoropropyl	4-(2,2,2-trifluoroethoxy)-phenyl
1028.	3-fluoropropyl	4-(1,1,2,2-tetrafluoroethoxy)-phenyl
1029.	3-fluoropropyl	4-cyclopropylphenyl
1030.	3-fluoropropyl	4-cyclobutylphenyl
1031.	3-fluoropropyl	4-cyclopentylphenyl
1032.	3-fluoropropyl	4-(2,2-difluorocyclopropyl)-phenyl
1033.	3-fluoropropyl	3,4-difluorophenyl
1034.	3-fluoropropyl	4-bromo-3-fluorophenyl
1035.	3-fluoropropyl	4-bromo-2-fluorophenyl
1036.	3-fluoropropyl	4-bromo-2,5-difluorophenyl
1037.	3-fluoropropyl	2-fluoro-4-isopropylphenyl
1038.	3-fluoropropyl	3-fluoro-4-isopropylphenyl
1039.	3-fluoropropyl	4-(1-hydroxy-1-methylethyl)-phenyl

No.	R ¹	Ar
1040.	3-fluoropropyl	4-(2-hydroxy-2-methylpropyl)-phenyl
1041.	3-fluoropropyl	4-acetylphenyl
1042.	3-fluoropropyl	4-carboxyphenyl
1043.	3-fluoropropyl	4-cyanophenyl
1044.	3-fluoropropyl	4-hydroxyphenyl
1045.	3-fluoropropyl	4-(O-benzyl)-phenyl
1046.	3-fluoropropyl	4-(2-methoxyethoxy)-phenyl
1047.	3-fluoropropyl	4-(CH ₂ -N(CH ₃) ₂)-phenyl
1048.	3-fluoropropyl	4-(NH-CO-NH ₂)-phenyl
1049.	3-fluoropropyl	4-(methylsulfanyl)-phenyl
1050.	3-fluoropropyl	4-(fluoromethylsulfanyl)-phenyl
1051.	3-fluoropropyl	4-(difluoromethylsulfanyl)-phenyl
1052.	3-fluoropropyl	4-(trifluoromethylsulfanyl)-phenyl
1053.	3-fluoropropyl	4-(methylsulfonyl)-phenyl
1054.	3-fluoropropyl	4-(N-methoxy-N-methyl-amino)-phenyl
1055.	3-fluoropropyl	4-(methoxyamino)-phenyl
1056.	3-fluoropropyl	4-(ethoxyamino)-phenyl
1057.	3-fluoropropyl	4-(N-methylaminooxy)-phenyl
1058.	3-fluoropropyl	4-(N,N-dimethylaminooxy)-phenyl
1059.	3-fluoropropyl	4-(azetidin-1-yl)-phenyl
1060.	3-fluoropropyl	4-(2-methylazetidin-1-yl)-phenyl
1061.	3-fluoropropyl	4-((S)-2-methylazetidin-1-yl)-phenyl
1062.	3-fluoropropyl	4-((R)-2-methylazetidin-1-yl)-phenyl
1063.	3-fluoropropyl	4-(3-fluoroazetidin-1-yl)-phenyl
1064.	3-fluoropropyl	4-(3-methoxyazetidin-1-yl)-phenyl
1065.	3-fluoropropyl	4-(3-hydroxyazetidin-1-yl)-phenyl
1066.	3-fluoropropyl	4-(pyrrolidin-1-yl)-phenyl
1067.	3-fluoropropyl	4-(pyrrolidin-2-yl)-phenyl
1068.	3-fluoropropyl	4-((S)-pyrrolidin-2-yl)-phenyl
1069.	3-fluoropropyl	4-((R)-pyrrolidin-2-yl)-phenyl
1070.	3-fluoropropyl	4-(pyrrolidin-3-yl)-phenyl
1071.	3-fluoropropyl	4-((S)-pyrrolidin-3-yl)-phenyl
1072.	3-fluoropropyl	4-((R)-pyrrolidin-3-yl)-phenyl
1073.	3-fluoropropyl	4-(2-fluoropyrrolidin-1-yl)-phenyl
1074.	3-fluoropropyl	4-((S)-2-fluoropyrrolidin-1-yl)-phenyl

No.	R ¹	Ar
1075.	3-fluoropropyl	4-((R)-2-fluoropyrrolidin-1-yl)-phenyl
1076.	3-fluoropropyl	4-(3-fluoropyrrolidin-1-yl)-phenyl
1077.	3-fluoropropyl	4-((S)-3-fluoropyrrolidin-1-yl)-phenyl
1078.	3-fluoropropyl	4-((R)-3-fluoropyrrolidin-1-yl)-phenyl
1079.	3-fluoropropyl	4-(2,2-difluoropyrrolidin-1-yl)-phenyl
1080.	3-fluoropropyl	4-(3,3-difluoropyrrolidin-1-yl)-phenyl
1081.	3-fluoropropyl	4-(2-methylpyrrolidin-1-yl)-phenyl
1082.	3-fluoropropyl	4-((S)-2-methylpyrrolidin-1-yl)-phenyl
1083.	3-fluoropropyl	4-((R)-2-methylpyrrolidin-1-yl)-phenyl
1084.	3-fluoropropyl	4-(3-methylpyrrolidin-1-yl)-phenyl
1085.	3-fluoropropyl	4-((S)-3-methylpyrrolidin-1-yl)-phenyl
1086.	3-fluoropropyl	4-((R)-3-methylpyrrolidin-1-yl)-phenyl
1087.	3-fluoropropyl	4-(1-methylpyrrolidin-2-yl)-phenyl
1088.	3-fluoropropyl	4-((S)-1-methylpyrrolidin-2-yl)-phenyl
1089.	3-fluoropropyl	4-((R)-1-methylpyrrolidin-2-yl)-phenyl
1090.	3-fluoropropyl	4-(1-methylpyrrolidin-3-yl)-phenyl
1091.	3-fluoropropyl	4-((S)-1-methylpyrrolidin-3-yl)-phenyl
1092.	3-fluoropropyl	4-((R)-1-methylpyrrolidin-3-yl)-phenyl
1093.	3-fluoropropyl	4-(2,2-dimethylpyrrolidin-1-yl)-phenyl
1094.	3-fluoropropyl	4-(3,3-dimethylpyrrolidin-1-yl)-phenyl
1095.	3-fluoropropyl	4-(2-trifluoromethylpyrrolidin-1-yl)-phenyl
1096.	3-fluoropropyl	4-((S)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
1097.	3-fluoropropyl	4-((R)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
1098.	3-fluoropropyl	4-(3-trifluoromethylpyrrolidin-1-yl)-phenyl
1099.	3-fluoropropyl	4-((S)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
1100.	3-fluoropropyl	4-((R)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
1101.	3-fluoropropyl	4-(2-oxopyrrolidin-1-yl)-phenyl
1102.	3-fluoropropyl	4-(2-oxo-oxazolidin-3-yl)-phenyl
1103.	3-fluoropropyl	4-(piperidin-1-yl)-phenyl
1104.	3-fluoropropyl	4-(2-methylpiperidin-1-yl)-phenyl
1105.	3-fluoropropyl	4-((S)-2-methylpiperidin-1-yl)-phenyl
1106.	3-fluoropropyl	4-((R)-2-methylpiperidin-1-yl)-phenyl
1107.	3-fluoropropyl	4-(piperazin-1-yl)-phenyl
1108.	3-fluoropropyl	4-(4-methylpiperazin-1-yl)-phenyl
1109.	3-fluoropropyl	4-(morpholin-4-yl)-phenyl

No.	R ¹	Ar
1110.	3-fluoropropyl	4-(thiomorpholin-4-yl)-phenyl
1111.	3-fluoropropyl	4-(1-oxo-thiomorpholin-4-yl)-phenyl
1112.	3-fluoropropyl	4-(1,1-dioxo-thiomorpholin-4-yl)-phenyl
1113.	3-fluoropropyl	4-(pyrrol-1-yl)-phenyl
1114.	3-fluoropropyl	4-(pyrrol-2-yl)-phenyl
1115.	3-fluoropropyl	4-(pyrrol-3-yl)-phenyl
1116.	3-fluoropropyl	4-(1-methylpyrrol-2-yl)-phenyl
1117.	3-fluoropropyl	4-(1-methylpyrrol-3-yl)-phenyl
1118.	3-fluoropropyl	4-(furan-2-yl)-phenyl
1119.	3-fluoropropyl	4-(furan-3-yl)-phenyl
1120.	3-fluoropropyl	4-(thiophen-2-yl)-phenyl
1121.	3-fluoropropyl	4-(thiophen-3-yl)-phenyl
1122.	3-fluoropropyl	4-(5-propylthien-2-yl)-phenyl
1123.	3-fluoropropyl	4-(pyrazol-1-yl)-phenyl
1124.	3-fluoropropyl	4-(pyrazol-3-yl)-phenyl
1125.	3-fluoropropyl	4-(pyrazol-4-yl)-phenyl
1126.	3-fluoropropyl	4-(1-methyl-1H-pyrazol-4-yl)-phenyl
1127.	3-fluoropropyl	4-(1-ethyl-1H-pyrazol-4-yl)-phenyl
1128.	3-fluoropropyl	4-(1-methyl-1H-pyrazol-5-yl)-phenyl
1129.	3-fluoropropyl	4-(1H-imidazol-2-yl)-phenyl
1130.	3-fluoropropyl	4-(imidazol-1-yl)-phenyl
1131.	3-fluoropropyl	4-(1-methylimidazol-2-yl)-phenyl
1132.	3-fluoropropyl	4-(oxazol-2-yl)-phenyl
1133.	3-fluoropropyl	4-(oxazol-4-yl)-phenyl
1134.	3-fluoropropyl	4-(oxazol-5-yl)-phenyl
1135.	3-fluoropropyl	4-(isoxazol-3-yl)-phenyl
1136.	3-fluoropropyl	4-(isoxazol-4-yl)-phenyl
1137.	3-fluoropropyl	4-(isoxazol-5-yl)-phenyl
1138.	3-fluoropropyl	4-([1,2,3]-triazol-1-yl)-phenyl
1139.	3-fluoropropyl	4-([1,2,4]-triazol-1-yl)-phenyl
1140.	3-fluoropropyl	4-([1,2,3]-triazol-2-yl)-phenyl
1141.	3-fluoropropyl	4-(4H-[1,2,4]-triazol-3-yl)-phenyl
1142.	3-fluoropropyl	4-([1,2,4]-triazol-4-yl)-phenyl
1143.	3-fluoropropyl	4-(2H-[1,2,3]-triazol-4-yl)-phenyl
1144.	3-fluoropropyl	4-(4-methyl-4H-[1,2,4]-triazol-3-yl)-phenyl

No.	R ¹	Ar
1145.	3-fluoropropyl	4-(2-methyl-2H-[1,2,3]-triazol-4-yl)-phenyl
1146.	3-fluoropropyl	4-([1,3,4]-oxadiazol-2-yl)-phenyl
1147.	3-fluoropropyl	4-([1,2,4]-oxadiazol-3-yl)-phenyl
1148.	3-fluoropropyl	4-([1,2,4]-oxadiazol-5-yl)-phenyl
1149.	3-fluoropropyl	4-([1,2,3]-oxadiazol-4-yl)-phenyl
1150.	3-fluoropropyl	4-([1,2,3]-oxadiazol-5-yl)-phenyl
1151.	3-fluoropropyl	4-([1,2,3]-thiadiazol-4-yl)-phenyl
1152.	3-fluoropropyl	4-(1H-tetrazol-5-yl)-phenyl
1153.	3-fluoropropyl	4-(tetrazol-1-yl)-phenyl
1154.	3-fluoropropyl	4-(2-methyl-2H-tetrazol-5-yl)-phenyl
1155.	3-fluoropropyl	4-(1-methyl-1H-tetrazol-5-yl)-phenyl
1156.	3-fluoropropyl	4-furazan-3-yl-phenyl
1157.	3-fluoropropyl	4-(pyrid-2-yl)-phenyl
1158.	3-fluoropropyl	4-(pyrid-3-yl)-phenyl
1159.	3-fluoropropyl	4-(pyrid-4-yl)-phenyl
1160.	3-fluoropropyl	4-(pyrimidin-2-yl)-phenyl
1161.	3-fluoropropyl	4-(pyrimidin-4-yl)-phenyl
1162.	3-fluoropropyl	4-(pyrimidin-5-yl)-phenyl
1163.	3-fluoropropyl	5-isopropylthiophen-2-yl
1164.	3-fluoropropyl	2-chlorothiophen-5-yl
1165.	3-fluoropropyl	2,5-dichlorothiophen-4-yl
1166.	3-fluoropropyl	2,3-dichlorothiophen-5-yl
1167.	3-fluoropropyl	2-chloro-3-nitrothiophen-5-yl
1168.	3-fluoropropyl	2-(phenylsulfonyl)-thiophen-5-yl
1169.	3-fluoropropyl	2-(pyridin-2-yl)thiophen-5-yl
1170.	3-fluoropropyl	2-(5-(trifluoromethyl)isoxazol-3-yl)-thiophen-5-yl
1171.	3-fluoropropyl	2-(2-methylthiazol-4-yl)-thiophen-5-yl
1172.	3-fluoropropyl	1-methyl-1H-imidazol-4-yl
1173.	3-fluoropropyl	1,2-dimethyl-1H-imidazol-4-yl
1174.	3-fluoropropyl	3,5-dimethylisoxazol-4-yl
1175.	3-fluoropropyl	thiazol-2-yl
1176.	3-fluoropropyl	4-methylthiazol-2-yl
1177.	3-fluoropropyl	4-isopropylthiazol-2-yl
1178.	3-fluoropropyl	4-trifluoromethylthiazol-2-yl
1179.	3-fluoropropyl	5-methylthiazol-2-yl

55

No.	R ¹	Ar
1180.	3-fluoropropyl	5-isopropylthiazol-2-yl
1181.	3-fluoropropyl	5-trifluoromethylthiazol-2-yl
1182.	3-fluoropropyl	2,4-dimethylthiazol-5-yl
1183.	3-fluoropropyl	2-acetamido-4-methylthiazol-5-yl
1184.	3-fluoropropyl	4H-[1,2,4]triazol-3-yl
1185.	3-fluoropropyl	5-methyl-4H-[1,2,4]triazol-3-yl
1186.	3-fluoropropyl	4-methyl-4H-[1,2,4]triazol-3-yl
1187.	3-fluoropropyl	5-isopropyl-4H-[1,2,4]triazol-3-yl
1188.	3-fluoropropyl	5-trifluoromethyl-4H-[1,2,4]triazol-3-yl
1189.	3-fluoropropyl	4,5-dimethyl-4H-[1,2,4]triazol-3-yl
1190.	3-fluoropropyl	5-isopropyl-4-methyl-4H-[1,2,4]triazol-3-yl
1191.	3-fluoropropyl	5-trifluoromethyl-4-methyl-4H-[1,2,4]triazol-3-yl
1192.	3-fluoropropyl	[1,3,4]thiadiazol-2-yl
1193.	3-fluoropropyl	5-methyl-[1,3,4]thiadiazol-2-yl
1194.	3-fluoropropyl	5-isopropyl-[1,3,4]thiadiazol-2-yl
1195.	3-fluoropropyl	5-trifluoromethyl-[1,3,4]thiadiazol-2-yl
1196.	3-fluoropropyl	3-bromo-2-chloropyrid-5-yl
1197.	3-fluoropropyl	2-(4-morpholino)-pyrid-5-yl
1198.	3-fluoropropyl	2-phenoxy-pyrid-5-yl
1199.	3-fluoropropyl	(2-isopropyl)-pyrimidin-5-yl
1200.	3-fluoropropyl	(5-isopropyl)-pyrimidin-2-yl
1201.	3-fluoropropyl	8-quinolyl
1202.	3-fluoropropyl	5-isoquinolyl
1203.	3-fluoropropyl	2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl
1204.	3-fluoropropyl	5-chloro-3-methylbenzothiophen-2-yl
1205.	3-fluoropropyl	3,4-dihydro-4-methyl-2H-benzo[b][1,4]oxazinyl
1206.	3-fluoropropyl	benzothiazol-6-yl
1207.	3-fluoropropyl	benzo[2,1,3]oxadiazol-4-yl
1208.	3-fluoropropyl	5-chlorobenzo[2,1,3]oxadiazol-4-yl
1209.	3-fluoropropyl	7-chlorobenzo[2,1,3]oxadiazol-4-yl
1210.	3-fluoropropyl	benzo[2,1,3]thiadiazol-4-yl
1211.	2-fluoroethyl	4-methylphenyl
1212.	2-fluoroethyl	4-ethylphenyl
1213.	2-fluoroethyl	4-propylphenyl

No.	R ¹	Ar
1214.	2-fluoroethyl	4-isopropylphenyl
1215.	2-fluoroethyl	4-sec-butylphenyl
1216.	2-fluoroethyl	4-isobutylphenyl
1217.	2-fluoroethyl	4-(1,1-dimethylpropyl)-phenyl
1218.	2-fluoroethyl	4-vinylphenyl
1219.	2-fluoroethyl	4-isopropenylphenyl
1220.	2-fluoroethyl	4-fluorophenyl
1221.	2-fluoroethyl	4-chlorophenyl
1222.	2-fluoroethyl	4-bromophenyl
1223.	2-fluoroethyl	4-(fluoromethyl)phenyl
1224.	2-fluoroethyl	3-(fluoromethyl)phenyl
1225.	2-fluoroethyl	2-(fluoromethyl)phenyl
1226.	2-fluoroethyl	4-(difluoromethyl)phenyl
1227.	2-fluoroethyl	3-(difluoromethyl)phenyl
1228.	2-fluoroethyl	2-(difluoromethyl)phenyl
1229.	2-fluoroethyl	4-(trifluoromethyl)phenyl
1230.	2-fluoroethyl	3-(trifluoromethyl)phenyl
1231.	2-fluoroethyl	2-(trifluoromethyl)phenyl
1232.	2-fluoroethyl	4-(1-fluoroethyl)-phenyl
1233.	2-fluoroethyl	4-((S)-1-fluoroethyl)-phenyl
1234.	2-fluoroethyl	4-((R)-1-fluoroethyl)-phenyl
1235.	2-fluoroethyl	4-(2-fluoroethyl)-phenyl
1236.	2-fluoroethyl	4-(1,1-difluoroethyl)-phenyl
1237.	2-fluoroethyl	4-(2,2-difluoroethyl)-phenyl
1238.	2-fluoroethyl	4-(2,2,2-trifluoroethyl)-phenyl
1239.	2-fluoroethyl	4-(3-fluoropropyl)-phenyl
1240.	2-fluoroethyl	4-(2-fluoropropyl)-phenyl
1241.	2-fluoroethyl	4-((S)-2-fluoropropyl)-phenyl
1242.	2-fluoroethyl	4-((R)-2-fluoropropyl)-phenyl
1243.	2-fluoroethyl	4-(3,3-difluoropropyl)-phenyl
1244.	2-fluoroethyl	4-(3,3,3-trifluoropropyl)-phenyl
1245.	2-fluoroethyl	4-(1-fluoro-1-methylethyl)-phenyl
1246.	2-fluoroethyl	4-(2-fluoro-1-methylethyl)-phenyl
1247.	2-fluoroethyl	4-((S)-2-fluoro-1-methylethyl)-phenyl
1248.	2-fluoroethyl	4-((R)-2-fluoro-1-methylethyl)-phenyl

57

No.	R ¹	Ar
1249.	2-fluoroethyl	4-(2,2-difluoro-1-methylethyl)-phenyl
1250.	2-fluoroethyl	4-((S)-2,2-difluoro-1-methylethyl)-phenyl
1251.	2-fluoroethyl	4-((R)-2,2-difluoro-1-methylethyl)-phenyl
1252.	2-fluoroethyl	4-(2,2,2-trifluoro-1-methylethyl)-phenyl
1253.	2-fluoroethyl	4-((S)-2,2,2-trifluoro-1-methylethyl)-phenyl
1254.	2-fluoroethyl	4-((R)-2,2,2-trifluoro-1-methylethyl)-phenyl
1255.	2-fluoroethyl	4-(2-fluoro-1-fluoromethylethyl)-phenyl
1256.	2-fluoroethyl	4-(1-difluoromethyl-2,2-difluoroethyl)-phenyl
1257.	2-fluoroethyl	4-(1,1-dimethyl-2-fluoroethyl)-phenyl
1258.	2-fluoroethyl	4-methoxyphenyl
1259.	2-fluoroethyl	4-ethoxyphenyl
1260.	2-fluoroethyl	4-propoxyphenyl
1261.	2-fluoroethyl	4-isopropoxyphenyl
1262.	2-fluoroethyl	4-butoxyphenyl
1263.	2-fluoroethyl	4-(fluoromethoxy)-phenyl
1264.	2-fluoroethyl	4-(difluoromethoxy)-phenyl
1265.	2-fluoroethyl	4-(trifluoromethoxy)-phenyl
1266.	2-fluoroethyl	3-(trifluoromethoxy)-phenyl
1267.	2-fluoroethyl	4-(2-fluoroethoxy)-phenyl
1268.	2-fluoroethyl	4-(2,2-difluoroethoxy)-phenyl
1269.	2-fluoroethyl	4-(2,2,2-trifluoroethoxy)-phenyl
1270.	2-fluoroethyl	4-(1,1,2,2-tetrafluoroethoxy)-phenyl
1271.	2-fluoroethyl	4-cyclopropylphenyl
1272.	2-fluoroethyl	4-cyclobutylphenyl
1273.	2-fluoroethyl	4-cyclopentylphenyl
1274.	2-fluoroethyl	4-(2,2-difluorocyclopropyl)-phenyl
1275.	2-fluoroethyl	3,4-difluorophenyl
1276.	2-fluoroethyl	4-bromo-3-fluorophenyl
1277.	2-fluoroethyl	4-bromo-2-fluorophenyl
1278.	2-fluoroethyl	4-bromo-2,5-difluorophenyl
1279.	2-fluoroethyl	2-fluoro-4-isopropylphenyl
1280.	2-fluoroethyl	3-fluoro-4-isopropylphenyl
1281.	2-fluoroethyl	4-(1-hydroxy-1-methylethyl)-phenyl
1282.	2-fluoroethyl	4-(2-hydroxy-2-methylpropyl)-phenyl
1283.	2-fluoroethyl	4-acetylphenyl

No.	R ¹	Ar
1284.	2-fluoroethyl	4-carboxyphenyl
1285.	2-fluoroethyl	4-cyanophenyl
1286.	2-fluoroethyl	4-hydroxyphenyl
1287.	2-fluoroethyl	4-(O-benzyl)-phenyl
1288.	2-fluoroethyl	4-(2-methoxyethoxy)-phenyl
1289.	2-fluoroethyl	4-(CH ₂ -N(CH ₃) ₂)-phenyl
1290.	2-fluoroethyl	4-(NH-CO-NH ₂)-phenyl
1291.	2-fluoroethyl	4-(methylsulfanyl)-phenyl
1292.	2-fluoroethyl	4-(fluoromethylsulfanyl)-phenyl
1293.	2-fluoroethyl	4-(difluoromethylsulfanyl)-phenyl
1294.	2-fluoroethyl	4-(trifluoromethylsulfanyl)-phenyl
1295.	2-fluoroethyl	4-(methylsulfonyl)-phenyl
1296.	2-fluoroethyl	4-(N-methoxy-N-methyl-amino)-phenyl
1297.	2-fluoroethyl	4-(methoxyamino)-phenyl
1298.	2-fluoroethyl	4-(ethoxyamino)-phenyl
1299.	2-fluoroethyl	4-(N-methylaminooxy)-phenyl
1300.	2-fluoroethyl	4-(N,N-dimethylaminooxy)-phenyl
1301.	2-fluoroethyl	4-(azetidin-1-yl)-phenyl
1302.	2-fluoroethyl	4-(2-methylazetidin-1-yl)-phenyl
1303.	2-fluoroethyl	4-((S)-2-methylazetidin-1-yl)-phenyl
1304.	2-fluoroethyl	4-((R)-2-methylazetidin-1-yl)-phenyl
1305.	2-fluoroethyl	4-(3-fluoroazetidin-1-yl)-phenyl
1306.	2-fluoroethyl	4-(3-methoxyazetidin-1-yl)-phenyl
1307.	2-fluoroethyl	4-(3-hydroxyazetidin-1-yl)-phenyl
1308.	2-fluoroethyl	4-(pyrrolidin-1-yl)-phenyl
1309.	2-fluoroethyl	4-(pyrrolidin-2-yl)-phenyl
1310.	2-fluoroethyl	4-((S)-pyrrolidin-2-yl)-phenyl
1311.	2-fluoroethyl	4-((R)-pyrrolidin-2-yl)-phenyl
1312.	2-fluoroethyl	4-(pyrrolidin-3-yl)-phenyl
1313.	2-fluoroethyl	4-((S)-pyrrolidin-3-yl)-phenyl
1314.	2-fluoroethyl	4-((R)-pyrrolidin-3-yl)-phenyl
1315.	2-fluoroethyl	4-(2-fluoropyrrolidin-1-yl)-phenyl
1316.	2-fluoroethyl	4-((S)-2-fluoropyrrolidin-1-yl)-phenyl
1317.	2-fluoroethyl	4-((R)-2-fluoropyrrolidin-1-yl)-phenyl
1318.	2-fluoroethyl	4-(3-fluoropyrrolidin-1-yl)-phenyl

No.	R ¹	Ar
1319.	2-fluoroethyl	4-((S)-3-fluoropyrrolidin-1-yl)-phenyl
1320.	2-fluoroethyl	4-((R)-3-fluoropyrrolidin-1-yl)-phenyl
1321.	2-fluoroethyl	4-(2,2-difluoropyrrolidin-1-yl)-phenyl
1322.	2-fluoroethyl	4-(3,3-difluoropyrrolidin-1-yl)-phenyl
1323.	2-fluoroethyl	4-(2-methylpyrrolidin-1-yl)-phenyl
1324.	2-fluoroethyl	4-((S)-2-methylpyrrolidin-1-yl)-phenyl
1325.	2-fluoroethyl	4-((R)-2-methylpyrrolidin-1-yl)-phenyl
1326.	2-fluoroethyl	4-(3-methylpyrrolidin-1-yl)-phenyl
1327.	2-fluoroethyl	4-((S)-3-methylpyrrolidin-1-yl)-phenyl
1328.	2-fluoroethyl	4-((R)-3-methylpyrrolidin-1-yl)-phenyl
1329.	2-fluoroethyl	4-(1-methylpyrrolidin-2-yl)-phenyl
1330.	2-fluoroethyl	4-((S)-1-methylpyrrolidin-2-yl)-phenyl
1331.	2-fluoroethyl	4-((R)-1-methylpyrrolidin-2-yl)-phenyl
1332.	2-fluoroethyl	4-(1-methylpyrrolidin-3-yl)-phenyl
1333.	2-fluoroethyl	4-((S)-1-methylpyrrolidin-3-yl)-phenyl
1334.	2-fluoroethyl	4-((R)-1-methylpyrrolidin-3-yl)-phenyl
1335.	2-fluoroethyl	4-(2,2-dimethylpyrrolidin-1-yl)-phenyl
1336.	2-fluoroethyl	4-(3,3-dimethylpyrrolidin-1-yl)-phenyl
1337.	2-fluoroethyl	4-(2-trifluoromethylpyrrolidin-1-yl)-phenyl
1338.	2-fluoroethyl	4-((S)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
1339.	2-fluoroethyl	4-((R)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
1340.	2-fluoroethyl	4-(3-trifluoromethylpyrrolidin-1-yl)-phenyl
1341.	2-fluoroethyl	4-((S)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
1342.	2-fluoroethyl	4-((R)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
1343.	2-fluoroethyl	4-(2-oxopyrrolidin-1-yl)-phenyl
1344.	2-fluoroethyl	4-(2-oxo-oxazolidin-3-yl)-phenyl
1345.	2-fluoroethyl	4-(piperidin-1-yl)-phenyl
1346.	2-fluoroethyl	4-(2-methylpiperidin-1-yl)-phenyl
1347.	2-fluoroethyl	4-((S)-2-methylpiperidin-1-yl)-phenyl
1348.	2-fluoroethyl	4-((R)-2-methylpiperidin-1-yl)-phenyl
1349.	2-fluoroethyl	4-(piperazin-1-yl)-phenyl
1350.	2-fluoroethyl	4-(4-methylpiperazin-1-yl)-phenyl
1351.	2-fluoroethyl	4-(morpholin-4-yl)-phenyl
1352.	2-fluoroethyl	4-(thiomorpholin-4-yl)-phenyl
1353.	2-fluoroethyl	4-(1-oxo-thiomorpholin-4-yl)-phenyl

No.	R ¹	Ar
1354.	2-fluoroethyl	4-(1,1-dioxo-thiomorpholin-4-yl)-phenyl
1355.	2-fluoroethyl	4-(pyrrol-1-yl)-phenyl
1356.	2-fluoroethyl	4-(pyrrol-2-yl)-phenyl
1357.	2-fluoroethyl	4-(pyrrol-3-yl)-phenyl
1358.	2-fluoroethyl	4-(1-methylpyrrol-2-yl)-phenyl
1359.	2-fluoroethyl	4-(1-methylpyrrol-3-yl)-phenyl
1360.	2-fluoroethyl	4-(furan-2-yl)-phenyl
1361.	2-fluoroethyl	4-(furan-3-yl)-phenyl
1362.	2-fluoroethyl	4-(thiophen-2-yl)-phenyl
1363.	2-fluoroethyl	4-(thiophen-3-yl)-phenyl
1364.	2-fluoroethyl	4-(5-propylthien-2-yl)-phenyl
1365.	2-fluoroethyl	4-(pyrazol-1-yl)-phenyl
1366.	2-fluoroethyl	4-(pyrazol-3-yl)-phenyl
1367.	2-fluoroethyl	4-(pyrazol-4-yl)-phenyl
1368.	2-fluoroethyl	4-(1-methyl-1H-pyrazol-4-yl)-phenyl
1369.	2-fluoroethyl	4-(1-ethyl-1H-pyrazol-4-yl)-phenyl
1370.	2-fluoroethyl	4-(1-methyl-1H-pyrazol-5-yl)-phenyl
1371.	2-fluoroethyl	4-(1H-imidazol-2-yl)-phenyl
1372.	2-fluoroethyl	4-(imidazol-1-yl)-phenyl
1373.	2-fluoroethyl	4-(1-methylimidazol-2-yl)-phenyl
1374.	2-fluoroethyl	4-(oxazol-2-yl)-phenyl
1375.	2-fluoroethyl	4-(oxazol-4-yl)-phenyl
1376.	2-fluoroethyl	4-(oxazol-5-yl)-phenyl
1377.	2-fluoroethyl	4-(isoxazol-3-yl)-phenyl
1378.	2-fluoroethyl	4-(isoxazol-4-yl)-phenyl
1379.	2-fluoroethyl	4-(isoxazol-5-yl)-phenyl
1380.	2-fluoroethyl	4-([1,2,3]-triazol-1-yl)-phenyl
1381.	2-fluoroethyl	4-([1,2,4]-triazol-1-yl)-phenyl
1382.	2-fluoroethyl	4-([1,2,3]-triazol-2-yl)-phenyl
1383.	2-fluoroethyl	4-(4H-[1,2,4]-triazol-3-yl)-phenyl
1384.	2-fluoroethyl	4-([1,2,4]-triazol-4-yl)-phenyl
1385.	2-fluoroethyl	4-(2H-[1,2,3]-triazol-4-yl)-phenyl
1386.	2-fluoroethyl	4-(4-methyl-4H-[1,2,4]-triazol-3-yl)-phenyl
1387.	2-fluoroethyl	4-(2-methyl-2H-[1,2,3]-triazol-4-yl)-phenyl
1388.	2-fluoroethyl	4-([1,3,4]-oxadiazol-2-yl)-phenyl

61

No.	R ¹	Ar
1389.	2-fluoroethyl	4-([1,2,4]-oxadiazol-3-yl)-phenyl
1390.	2-fluoroethyl	4-([1,2,4]-oxadiazol-5-yl)-phenyl
1391.	2-fluoroethyl	4-([1,2,3]-oxadiazol-4-yl)-phenyl
1392.	2-fluoroethyl	4-([1,2,3]-oxadiazol-5-yl)-phenyl
1393.	2-fluoroethyl	4-([1,2,3]-thiadiazol-4-yl)-phenyl
1394.	2-fluoroethyl	4-(1H-tetrazol-5-yl)-phenyl
1395.	2-fluoroethyl	4-(tetrazol-1-yl)-phenyl
1396.	2-fluoroethyl	4-(2-methyl-2H-tetrazol-5-yl)-phenyl
1397.	2-fluoroethyl	4-(1-methyl-1H-tetrazol-5-yl)-phenyl
1398.	2-fluoroethyl	4-furazan-3-yl-phenyl
1399.	2-fluoroethyl	4-(pyrid-2-yl)-phenyl
1400.	2-fluoroethyl	4-(pyrid-3-yl)-phenyl
1401.	2-fluoroethyl	4-(pyrid-4-yl)-phenyl
1402.	2-fluoroethyl	4-(pyrimidin-2-yl)-phenyl
1403.	2-fluoroethyl	4-(pyrimidin-4-yl)-phenyl
1404.	2-fluoroethyl	4-(pyrimidin-5-yl)-phenyl
1405.	2-fluoroethyl	5-isopropylthiophen-2-yl
1406.	2-fluoroethyl	2-chlorothiophen-5-yl
1407.	2-fluoroethyl	2,5-dichlorothiophen-4-yl
1408.	2-fluoroethyl	2,3-dichlorothiophen-5-yl
1409.	2-fluoroethyl	2-chloro-3-nitrothiophen-5-yl
1410.	2-fluoroethyl	2-(phenylsulfonyl)-thiophen-5-yl
1411.	2-fluoroethyl	2-(pyridin-2-yl)thiophen-5-yl
1412.	2-fluoroethyl	2-(5-(trifluoromethyl)isoxazol-3-yl)-thiophen-5-yl
1413.	2-fluoroethyl	2-(2-methylthiazol-4-yl)-thiophen-5-yl
1414.	2-fluoroethyl	1-methyl-1H-imidazol-4-yl
1415.	2-fluoroethyl	1,2-dimethyl-1H-imidazol-4-yl
1416.	2-fluoroethyl	3,5-dimethylisoxazol-4-yl
1417.	2-fluoroethyl	thiazol-2-yl
1418.	2-fluoroethyl	4-methylthiazol-2-yl
1419.	2-fluoroethyl	4-isopropylthiazol-2-yl
1420.	2-fluoroethyl	4-trifluoromethylthiazol-2-yl
1421.	2-fluoroethyl	5-methylthiazol-2-yl
1422.	2-fluoroethyl	5-isopropylthiazol-2-yl
1423.	2-fluoroethyl	5-trifluoromethylthiazol-2-yl

62

No.	R ¹	Ar
1424.	2-fluoroethyl	2,4-dimethylthiazol-5-yl
1425.	2-fluoroethyl	2-acetamido-4-methylthiazol-5-yl
1426.	2-fluoroethyl	4H-[1,2,4]triazol-3-yl
1427.	2-fluoroethyl	5-methyl-4H-[1,2,4]triazol-3-yl
1428.	2-fluoroethyl	4-methyl-4H-[1,2,4]triazol-3-yl
1429.	2-fluoroethyl	5-isopropyl-4H-[1,2,4]triazol-3-yl
1430.	2-fluoroethyl	5-trifluoromethyl-4H-[1,2,4]triazol-3-yl
1431.	2-fluoroethyl	4,5-dimethyl-4H-[1,2,4]triazol-3-yl
1432.	2-fluoroethyl	5-isopropyl-4-methyl-4H-[1,2,4]triazol-3-yl
1433.	2-fluoroethyl	5-trifluoromethyl-4-methyl-4H-[1,2,4]triazol-3-yl
1434.	2-fluoroethyl	[1,3,4]thiadiazol-2-yl
1435.	2-fluoroethyl	5-methyl-[1,3,4]thiadiazol-2-yl
1436.	2-fluoroethyl	5-isopropyl-[1,3,4]thiadiazol-2-yl
1437.	2-fluoroethyl	5-trifluoromethyl-[1,3,4]thiadiazol-2-yl
1438.	2-fluoroethyl	3-bromo-2-chloropyrid-5-yl
1439.	2-fluoroethyl	2-(4-morpholino)-pyrid-5-yl
1440.	2-fluoroethyl	2-phenoxy-pyrid-5-yl
1441.	2-fluoroethyl	(2-isopropyl)-pyrimidin-5-yl
1442.	2-fluoroethyl	(5-isopropyl)-pyrimidin-2-yl
1443.	2-fluoroethyl	8-quinolyl
1444.	2-fluoroethyl	5-isoquinolyl
1445.	2-fluoroethyl	2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl
1446.	2-fluoroethyl	5-chloro-3-methylbenzothiophen-2-yl
1447.	2-fluoroethyl	3,4-dihydro-4-methyl-2H-benzo[b][1,4]oxazinyl
1448.	2-fluoroethyl	benzothiazol-6-yl
1449.	2-fluoroethyl	benzo[2,1,3]oxadiazol-4-yl
1450.	2-fluoroethyl	5-chlorobenzo[2,1,3]oxadiazol-4-yl
1451.	2-fluoroethyl	7-chlorobenzo[2,1,3]oxadiazol-4-yl
1452.	2-fluoroethyl	benzo[2,1,3]thiadiazol-4-yl
1453.	cyclopropylmethyl	4-methylphenyl
1454.	cyclopropylmethyl	4-ethylphenyl
1455.	cyclopropylmethyl	4-propylphenyl
1456.	cyclopropylmethyl	4-isopropylphenyl
1457.	cyclopropylmethyl	4-sec-butylphenyl

No.	R ¹	Ar
1458.	cyclopropylmethyl	4-isobutylphenyl
1459.	cyclopropylmethyl	4-(1,1-dimethylpropyl)-phenyl
1460.	cyclopropylmethyl	4-vinylphenyl
1461.	cyclopropylmethyl	4-isopropenylphenyl
1462.	cyclopropylmethyl	4-fluorophenyl
1463.	cyclopropylmethyl	4-chlorophenyl
1464.	cyclopropylmethyl	4-bromophenyl
1465.	cyclopropylmethyl	4-(fluoromethyl)phenyl
1466.	cyclopropylmethyl	3-(fluoromethyl)phenyl
1467.	cyclopropylmethyl	2-(fluoromethyl)phenyl
1468.	cyclopropylmethyl	4-(difluoromethyl)phenyl
1469.	cyclopropylmethyl	3-(difluoromethyl)phenyl
1470.	cyclopropylmethyl	2-(difluoromethyl)phenyl
1471.	cyclopropylmethyl	4-(trifluoromethyl)phenyl
1472.	cyclopropylmethyl	3-(trifluoromethyl)phenyl
1473.	cyclopropylmethyl	2-(trifluoromethyl)phenyl
1474.	cyclopropylmethyl	4-(1-fluoroethyl)-phenyl
1475.	cyclopropylmethyl	4-((S)-1-fluoroethyl)-phenyl
1476.	cyclopropylmethyl	4-((R)-1-fluoroethyl)-phenyl
1477.	cyclopropylmethyl	4-(2-fluoroethyl)-phenyl
1478.	cyclopropylmethyl	4-(1,1-difluoroethyl)-phenyl
1479.	cyclopropylmethyl	4-(2,2-difluoroethyl)-phenyl
1480.	cyclopropylmethyl	4-(2,2,2-trifluoroethyl)-phenyl
1481.	cyclopropylmethyl	4-(3-fluoropropyl)-phenyl
1482.	cyclopropylmethyl	4-(2-fluoropropyl)-phenyl
1483.	cyclopropylmethyl	4-((S)-2-fluoropropyl)-phenyl
1484.	cyclopropylmethyl	4-((R)-2-fluoropropyl)-phenyl
1485.	cyclopropylmethyl	4-(3,3-difluoropropyl)-phenyl
1486.	cyclopropylmethyl	4-(3,3,3-trifluoropropyl)-phenyl
1487.	cyclopropylmethyl	4-(1-fluoro-1-methylethyl)-phenyl
1488.	cyclopropylmethyl	4-(2-fluoro-1-methylethyl)-phenyl
1489.	cyclopropylmethyl	4-((S)-2-fluoro-1-methylethyl)-phenyl
1490.	cyclopropylmethyl	4-((R)-2-fluoro-1-methylethyl)-phenyl
1491.	cyclopropylmethyl	4-(2,2-difluoro-1-methylethyl)-phenyl
1492.	cyclopropylmethyl	4-((S)-2,2-difluoro-1-methylethyl)-phenyl

No.	R ¹	Ar
1493.	cyclopropylmethyl	4-((R)-2,2-difluoro-1-methylethyl)-phenyl
1494.	cyclopropylmethyl	4-(2,2,2-trifluoro-1-methylethyl)-phenyl
1495.	cyclopropylmethyl	4-((S)-2,2,2-trifluoro-1-methylethyl)-phenyl
1496.	cyclopropylmethyl	4-((R)-2,2,2-trifluoro-1-methylethyl)-phenyl
1497.	cyclopropylmethyl	4-(2-fluoro-1-fluoromethylethyl)-phenyl
1498.	cyclopropylmethyl	4-(1-difluoromethyl-2,2-difluoroethyl)-phenyl
1499.	cyclopropylmethyl	4-(1,1-dimethyl-2-fluoroethyl)-phenyl
1500.	cyclopropylmethyl	4-methoxyphenyl
1501.	cyclopropylmethyl	4-ethoxyphenyl
1502.	cyclopropylmethyl	4-propoxyphenyl
1503.	cyclopropylmethyl	4-isopropoxyphenyl
1504.	cyclopropylmethyl	4-butoxyphenyl
1505.	cyclopropylmethyl	4-(fluoromethoxy)-phenyl
1506.	cyclopropylmethyl	4-(difluoromethoxy)-phenyl
1507.	cyclopropylmethyl	4-(trifluoromethoxy)-phenyl
1508.	cyclopropylmethyl	3-(trifluoromethoxy)-phenyl
1509.	cyclopropylmethyl	4-(2-fluoroethoxy)-phenyl
1510.	cyclopropylmethyl	4-(2,2-difluoroethoxy)-phenyl
1511.	cyclopropylmethyl	4-(2,2,2-trifluoroethoxy)-phenyl
1512.	cyclopropylmethyl	4-(1,1,2,2-tetrafluoroethoxy)-phenyl
1513.	cyclopropylmethyl	4-cyclopropylphenyl
1514.	cyclopropylmethyl	4-cyclobutylphenyl
1515.	cyclopropylmethyl	4-cyclopentylphenyl
1516.	cyclopropylmethyl	4-(2,2-difluorocyclopropyl)-phenyl
1517.	cyclopropylmethyl	3,4-difluorophenyl
1518.	cyclopropylmethyl	4-bromo-3-fluorophenyl
1519.	cyclopropylmethyl	4-bromo-2-fluorophenyl
1520.	cyclopropylmethyl	4-bromo-2,5-difluorophenyl
1521.	cyclopropylmethyl	2-fluoro-4-isopropylphenyl
1522.	cyclopropylmethyl	3-fluoro-4-isopropylphenyl
1523.	cyclopropylmethyl	4-(1-hydroxy-1-methylethyl)-phenyl
1524.	cyclopropylmethyl	4-(2-hydroxy-2-methylpropyl)-phenyl
1525.	cyclopropylmethyl	4-acetylphenyl
1526.	cyclopropylmethyl	4-carboxyphenyl
1527.	cyclopropylmethyl	4-cyanophenyl

65

No.	R ¹	Ar
1528.	cyclopropylmethyl	4-hydroxyphenyl
1529.	cyclopropylmethyl	4-(O-benzyl)-phenyl
1530.	cyclopropylmethyl	4-(2-methoxyethoxy)-phenyl
1531.	cyclopropylmethyl	4-(CH ₂ -N(CH ₃) ₂)-phenyl
1532.	cyclopropylmethyl	4-(NH-CO-NH ₂)-phenyl
1533.	cyclopropylmethyl	4-(methylsulfanyl)-phenyl
1534.	cyclopropylmethyl	4-(fluoromethylsulfanyl)-phenyl
1535.	cyclopropylmethyl	4-(difluoromethylsulfanyl)-phenyl
1536.	cyclopropylmethyl	4-(trifluoromethylsulfanyl)-phenyl
1537.	cyclopropylmethyl	4-(methylsulfonyl)-phenyl
1538.	cyclopropylmethyl	4-(N-methoxy-N-methyl-amino)-phenyl
1539.	cyclopropylmethyl	4-(methoxyamino)-phenyl
1540.	cyclopropylmethyl	4-(ethoxyamino)-phenyl
1541.	cyclopropylmethyl	4-(N-methylaminooxy)-phenyl
1542.	cyclopropylmethyl	4-(N,N-dimethylaminooxy)-phenyl
1543.	cyclopropylmethyl	4-(azetidin-1-yl)-phenyl
1544.	cyclopropylmethyl	4-(2-methylazetidin-1-yl)-phenyl
1545.	cyclopropylmethyl	4-((S)-2-methylazetidin-1-yl)-phenyl
1546.	cyclopropylmethyl	4-((R)-2-methylazetidin-1-yl)-phenyl
1547.	cyclopropylmethyl	4-(3-fluoroazetidin-1-yl)-phenyl
1548.	cyclopropylmethyl	4-(3-methoxyazetidin-1-yl)-phenyl
1549.	cyclopropylmethyl	4-(3-hydroxyazetidin-1-yl)-phenyl
1550.	cyclopropylmethyl	4-(pyrrolidin-1-yl)-phenyl
1551.	cyclopropylmethyl	4-(pyrrolidin-2-yl)-phenyl
1552.	cyclopropylmethyl	4-((S)-pyrrolidin-2-yl)-phenyl
1553.	cyclopropylmethyl	4-((R)-pyrrolidin-2-yl)-phenyl
1554.	cyclopropylmethyl	4-(pyrrolidin-3-yl)-phenyl
1555.	cyclopropylmethyl	4-((S)-pyrrolidin-3-yl)-phenyl
1556.	cyclopropylmethyl	4-((R)-pyrrolidin-3-yl)-phenyl
1557.	cyclopropylmethyl	4-(2-fluoropyrrolidin-1-yl)-phenyl
1558.	cyclopropylmethyl	4-((S)-2-fluoropyrrolidin-1-yl)-phenyl
1559.	cyclopropylmethyl	4-((R)-2-fluoropyrrolidin-1-yl)-phenyl
1560.	cyclopropylmethyl	4-(3-fluoropyrrolidin-1-yl)-phenyl
1561.	cyclopropylmethyl	4-((S)-3-fluoropyrrolidin-1-yl)-phenyl
1562.	cyclopropylmethyl	4-((R)-3-fluoropyrrolidin-1-yl)-phenyl

No.	R ¹	Ar
1563.	cyclopropylmethyl	4-(2,2-difluoropyrrolidin-1-yl)-phenyl
1564.	cyclopropylmethyl	4-(3,3-difluoropyrrolidin-1-yl)-phenyl
1565.	cyclopropylmethyl	4-(2-methylpyrrolidin-1-yl)-phenyl
1566.	cyclopropylmethyl	4-((S)-2-methylpyrrolidin-1-yl)-phenyl
1567.	cyclopropylmethyl	4-((R)-2-methylpyrrolidin-1-yl)-phenyl
1568.	cyclopropylmethyl	4-(3-methylpyrrolidin-1-yl)-phenyl
1569.	cyclopropylmethyl	4-((S)-3-methylpyrrolidin-1-yl)-phenyl
1570.	cyclopropylmethyl	4-((R)-3-methylpyrrolidin-1-yl)-phenyl
1571.	cyclopropylmethyl	4-(1-methylpyrrolidin-2-yl)-phenyl
1572.	cyclopropylmethyl	4-((S)-1-methylpyrrolidin-2-yl)-phenyl
1573.	cyclopropylmethyl	4-((R)-1-methylpyrrolidin-2-yl)-phenyl
1574.	cyclopropylmethyl	4-(1-methylpyrrolidin-3-yl)-phenyl
1575.	cyclopropylmethyl	4-((S)-1-methylpyrrolidin-3-yl)-phenyl
1576.	cyclopropylmethyl	4-((R)-1-methylpyrrolidin-3-yl)-phenyl
1577.	cyclopropylmethyl	4-(2,2-dimethylpyrrolidin-1-yl)-phenyl
1578.	cyclopropylmethyl	4-(3,3-dimethylpyrrolidin-1-yl)-phenyl
1579.	cyclopropylmethyl	4-(2-trifluoromethylpyrrolidin-1-yl)-phenyl
1580.	cyclopropylmethyl	4-((S)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
1581.	cyclopropylmethyl	4-((R)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
1582.	cyclopropylmethyl	4-(3-trifluoromethylpyrrolidin-1-yl)-phenyl
1583.	cyclopropylmethyl	4-((S)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
1584.	cyclopropylmethyl	4-((R)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
1585.	cyclopropylmethyl	4-(2-oxopyrrolidin-1-yl)-phenyl
1586.	cyclopropylmethyl	4-(2-oxo-oxazolidin-3-yl)-phenyl
1587.	cyclopropylmethyl	4-(piperidin-1-yl)-phenyl
1588.	cyclopropylmethyl	4-(2-methylpiperidin-1-yl)-phenyl
1589.	cyclopropylmethyl	4-((S)-2-methylpiperidin-1-yl)-phenyl
1590.	cyclopropylmethyl	4-((R)-2-methylpiperidin-1-yl)-phenyl
1591.	cyclopropylmethyl	4-(piperazin-1-yl)-phenyl
1592.	cyclopropylmethyl	4-(4-methylpiperazin-1-yl)-phenyl
1593.	cyclopropylmethyl	4-(morpholin-4-yl)-phenyl
1594.	cyclopropylmethyl	4-(thiomorpholin-4-yl)-phenyl
1595.	cyclopropylmethyl	4-(1-oxo-thiomorpholin-4-yl)-phenyl
1596.	cyclopropylmethyl	4-(1,1-dioxo-thiomorpholin-4-yl)-phenyl
1597.	cyclopropylmethyl	4-(pyrrol-1-yl)-phenyl

No.	R ¹	Ar
1598.	cyclopropylmethyl	4-(pyrrol-2-yl)-phenyl
1599.	cyclopropylmethyl	4-(pyrrol-3-yl)-phenyl
1600.	cyclopropylmethyl	4-(1-methylpyrrol-2-yl)-phenyl
1601.	cyclopropylmethyl	4-(1-methylpyrrol-3-yl)-phenyl
1602.	cyclopropylmethyl	4-(furan-2-yl)-phenyl
1603.	cyclopropylmethyl	4-(furan-3-yl)-phenyl
1604.	cyclopropylmethyl	4-(thiophen-2-yl)-phenyl
1605.	cyclopropylmethyl	4-(thiophen-3-yl)-phenyl
1606.	cyclopropylmethyl	4-(5-propylthien-2-yl)-phenyl
1607.	cyclopropylmethyl	4-(pyrazol-1-yl)-phenyl
1608.	cyclopropylmethyl	4-(pyrazol-3-yl)-phenyl
1609.	cyclopropylmethyl	4-(pyrazol-4-yl)-phenyl
1610.	cyclopropylmethyl	4-(1-methyl-1H-pyrazol-4-yl)-phenyl
1611.	cyclopropylmethyl	4-(1-ethyl-1H-pyrazol-4-yl)-phenyl
1612.	cyclopropylmethyl	4-(1-methyl-1H-pyrazol-5-yl)-phenyl
1613.	cyclopropylmethyl	4-(1H-imidazol-2-yl)-phenyl
1614.	cyclopropylmethyl	4-(imidazol-1-yl)-phenyl
1615.	cyclopropylmethyl	4-(1-methylimidazol-2-yl)-phenyl
1616.	cyclopropylmethyl	4-(oxazol-2-yl)-phenyl
1617.	cyclopropylmethyl	4-(oxazol-4-yl)-phenyl
1618.	cyclopropylmethyl	4-(oxazol-5-yl)-phenyl
1619.	cyclopropylmethyl	4-(isoxazol-3-yl)-phenyl
1620.	cyclopropylmethyl	4-(isoxazol-4-yl)-phenyl
1621.	cyclopropylmethyl	4-(isoxazol-5-yl)-phenyl
1622.	cyclopropylmethyl	4-([1,2,3]-triazol-1-yl)-phenyl
1623.	cyclopropylmethyl	4-([1,2,4]-triazol-1-yl)-phenyl
1624.	cyclopropylmethyl	4-([1,2,3]-triazol-2-yl)-phenyl
1625.	cyclopropylmethyl	4-(4H-[1,2,4]-triazol-3-yl)-phenyl
1626.	cyclopropylmethyl	4-([1,2,4]-triazol-4-yl)-phenyl
1627.	cyclopropylmethyl	4-(2H-[1,2,3]-triazol-4-yl)-phenyl
1628.	cyclopropylmethyl	4-(4-methyl-4H-[1,2,4]-triazol-3-yl)-phenyl
1629.	cyclopropylmethyl	4-(2-methyl-2H-[1,2,3]-triazol-4-yl)-phenyl
1630.	cyclopropylmethyl	4-([1,3,4]-oxadiazol-2-yl)-phenyl
1631.	cyclopropylmethyl	4-([1,2,4]-oxadiazol-3-yl)-phenyl
1632.	cyclopropylmethyl	4-([1,2,4]-oxadiazol-5-yl)-phenyl

No.	R ¹	Ar
1633.	cyclopropylmethyl	4-([1,2,3]-oxadiazol-4-yl)-phenyl
1634.	cyclopropylmethyl	4-([1,2,3]-oxadiazol-5-yl)-phenyl
1635.	cyclopropylmethyl	4-([1,2,3]-thiadiazol-4-yl)-phenyl
1636.	cyclopropylmethyl	4-(1H-tetrazol-5-yl)-phenyl
1637.	cyclopropylmethyl	4-(tetrazol-1-yl)-phenyl
1638.	cyclopropylmethyl	4-(2-methyl-2H-tetrazol-5-yl)-phenyl
1639.	cyclopropylmethyl	4-(1-methyl-1H-tetrazol-5-yl)-phenyl
1640.	cyclopropylmethyl	4-furazan-3-yl-phenyl
1641.	cyclopropylmethyl	4-(pyrid-2-yl)-phenyl
1642.	cyclopropylmethyl	4-(pyrid-3-yl)-phenyl
1643.	cyclopropylmethyl	4-(pyrid-4-yl)-phenyl
1644.	cyclopropylmethyl	4-(pyrimidin-2-yl)-phenyl
1645.	cyclopropylmethyl	4-(pyrimidin-4-yl)-phenyl
1646.	cyclopropylmethyl	4-(pyrimidin-5-yl)-phenyl
1647.	cyclopropylmethyl	5-isopropylthiophen-2-yl
1648.	cyclopropylmethyl	2-chlorothiophen-5-yl
1649.	cyclopropylmethyl	2,5-dichlorothiophen-4-yl
1650.	cyclopropylmethyl	2,3-dichlorothiophen-5-yl
1651.	cyclopropylmethyl	2-chloro-3-nitrothiophen-5-yl
1652.	cyclopropylmethyl	2-(phenylsulfonyl)-thiophen-5-yl
1653.	cyclopropylmethyl	2-(pyridin-2-yl)thiophen-5-yl
1654.	cyclopropylmethyl	2-(5-(trifluoromethyl)isoxazol-3-yl)-thiophen-5-yl
1655.	cyclopropylmethyl	2-(2-methylthiazol-4-yl)-thiophen-5-yl
1656.	cyclopropylmethyl	1-methyl-1H-imidazol-4-yl
1657.	cyclopropylmethyl	1,2-dimethyl-1H-imidazol-4-yl
1658.	cyclopropylmethyl	3,5-dimethylisoxazol-4-yl
1659.	cyclopropylmethyl	thiazol-2-yl
1660.	cyclopropylmethyl	4-methylthiazol-2-yl
1661.	cyclopropylmethyl	4-isopropylthiazol-2-yl
1662.	cyclopropylmethyl	4-trifluoromethylthiazol-2-yl
1663.	cyclopropylmethyl	5-methylthiazol-2-yl
1664.	cyclopropylmethyl	5-isopropylthiazol-2-yl
1665.	cyclopropylmethyl	5-trifluoromethylthiazol-2-yl
1666.	cyclopropylmethyl	2,4-dimethylthiazol-5-yl
1667.	cyclopropylmethyl	2-acetamido-4-methylthiazol-5-yl

No.	R ¹	Ar
1668.	cyclopropylmethyl	4H-[1,2,4]triazol-3-yl
1669.	cyclopropylmethyl	5-methyl-4H-[1,2,4]triazol-3-yl
1670.	cyclopropylmethyl	4-methyl-4H-[1,2,4]triazol-3-yl
1671.	cyclopropylmethyl	5-isopropyl-4H-[1,2,4]triazol-3-yl
1672.	cyclopropylmethyl	5-trifluoromethyl-4H-[1,2,4]triazol-3-yl
1673.	cyclopropylmethyl	4,5-dimethyl-4H-[1,2,4]triazol-3-yl
1674.	cyclopropylmethyl	5-isopropyl-4-methyl-4H-[1,2,4]triazol-3-yl
1675.	cyclopropylmethyl	5-trifluoromethyl-4-methyl-4H-[1,2,4]triazol-3-yl
1676.	cyclopropylmethyl	[1,3,4]thiadiazol-2-yl
1677.	cyclopropylmethyl	5-methyl-[1,3,4]thiadiazol-2-yl
1678.	cyclopropylmethyl	5-isopropyl-[1,3,4]thiadiazol-2-yl
1679.	cyclopropylmethyl	5-trifluoromethyl-[1,3,4]thiadiazol-2-yl
1680.	cyclopropylmethyl	3-bromo-2-chloropyrid-5-yl
1681.	cyclopropylmethyl	2-(4-morpholino)-pyrid-5-yl
1682.	cyclopropylmethyl	2-phenoxy-pyrid-5-yl
1683.	cyclopropylmethyl	(2-isopropyl)-pyrimidin-5-yl
1684.	cyclopropylmethyl	(5-isopropyl)-pyrimidin-2-yl
1685.	cyclopropylmethyl	8-quinolyl
1686.	cyclopropylmethyl	5-isoquinolyl
1687.	cyclopropylmethyl	2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl
1688.	cyclopropylmethyl	5-chloro-3-methylbenzothiophen-2-yl
1689.	cyclopropylmethyl	3,4-dihydro-4-methyl-2H-benzo[b][1,4]oxazinyl
1690.	cyclopropylmethyl	benzothiazol-6-yl
1691.	cyclopropylmethyl	benzo[2,1,3]oxadiazol-4-yl
1692.	cyclopropylmethyl	5-chlorobenzo[2,1,3]oxadiazol-4-yl
1693.	cyclopropylmethyl	7-chlorobenzo[2,1,3]oxadiazol-4-yl
1694.	cyclopropylmethyl	benzo[2,1,3]thiadiazol-4-yl
1695.	allyl	4-methylphenyl
1696.	allyl	4-ethylphenyl
1697.	allyl	4-propylphenyl
1698.	allyl	4-isopropylphenyl
1699.	allyl	4-sec-butylphenyl
1700.	allyl	4-isobutylphenyl
1701.	allyl	4-(1,1-dimethylpropyl)-phenyl

No.	R ¹	Ar
1702.	allyl	4-vinylphenyl
1703.	allyl	4-isopropenylphenyl
1704.	allyl	4-fluorophenyl
1705.	allyl	4-chlorophenyl
1706.	allyl	4-bromophenyl
1707.	allyl	4-(fluoromethyl)phenyl
1708.	allyl	3-(fluoromethyl)phenyl
1709.	allyl	2-(fluoromethyl)phenyl
1710.	allyl	4-(difluoromethyl)phenyl
1711.	allyl	3-(difluoromethyl)phenyl
1712.	allyl	2-(difluoromethyl)phenyl
1713.	allyl	4-(trifluoromethyl)phenyl
1714.	allyl	3-(trifluoromethyl)phenyl
1715.	allyl	2-(trifluoromethyl)phenyl
1716.	allyl	4-(1-fluoroethyl)-phenyl
1717.	allyl	4-((S)-1-fluoroethyl)-phenyl
1718.	allyl	4-((R)-1-fluoroethyl)-phenyl
1719.	allyl	4-(2-fluoroethyl)-phenyl
1720.	allyl	4-(1,1-difluoroethyl)-phenyl
1721.	allyl	4-(2,2-difluoroethyl)-phenyl
1722.	allyl	4-(2,2,2-trifluoroethyl)-phenyl
1723.	allyl	4-(3-fluoropropyl)-phenyl
1724.	allyl	4-(2-fluoropropyl)-phenyl
1725.	allyl	4-((S)-2-fluoropropyl)-phenyl
1726.	allyl	4-((R)-2-fluoropropyl)-phenyl
1727.	allyl	4-(3,3-difluoropropyl)-phenyl
1728.	allyl	4-(3,3,3-trifluoropropyl)-phenyl
1729.	allyl	4-(1-fluoro-1-methylethyl)-phenyl
1730.	allyl	4-(2-fluoro-1-methylethyl)-phenyl
1731.	allyl	4-((S)-2-fluoro-1-methylethyl)-phenyl
1732.	allyl	4-((R)-2-fluoro-1-methylethyl)-phenyl
1733.	allyl	4-(2,2-difluoro-1-methylethyl)-phenyl
1734.	allyl	4-((S)-2,2-difluoro-1-methylethyl)-phenyl
1735.	allyl	4-((R)-2,2-difluoro-1-methylethyl)-phenyl
1736.	allyl	4-(2,2,2-trifluoro-1-methylethyl)-phenyl

No.	R ¹	Ar
1737.	allyl	4-((S)-2,2,2-trifluoro-1-methylethyl)-phenyl
1738.	allyl	4-((R)-2,2,2-trifluoro-1-methylethyl)-phenyl
1739.	allyl	4-(2-fluoro-1-fluoromethylethyl)-phenyl
1740.	allyl	4-(1-difluoromethyl-2,2-difluoroethyl)-phenyl
1741.	allyl	4-(1,1-dimethyl-2-fluoroethyl)-phenyl
1742.	allyl	4-methoxyphenyl
1743.	allyl	4-ethoxyphenyl
1744.	allyl	4-propoxyphenyl
1745.	allyl	4-isopropoxyphenyl
1746.	allyl	4-butoxyphenyl
1747.	allyl	4-(fluoromethoxy)-phenyl
1748.	allyl	4-(difluoromethoxy)-phenyl
1749.	allyl	4-(trifluoromethoxy)-phenyl
1750.	allyl	3-(trifluoromethoxy)-phenyl
1751.	allyl	4-(2-fluoroethoxy)-phenyl
1752.	allyl	4-(2,2-difluoroethoxy)-phenyl
1753.	allyl	4-(2,2,2-trifluoroethoxy)-phenyl
1754.	allyl	4-(1,1,2,2-tetrafluoroethoxy)-phenyl
1755.	allyl	4-cyclopropylphenyl
1756.	allyl	4-cyclobutylphenyl
1757.	allyl	4-cyclopentylphenyl
1758.	allyl	4-(2,2-difluorocyclopropyl)-phenyl
1759.	allyl	3,4-difluorophenyl
1760.	allyl	4-bromo-3-fluorophenyl
1761.	allyl	4-bromo-2-fluorophenyl
1762.	allyl	4-bromo-2,5-difluorophenyl
1763.	allyl	2-fluoro-4-isopropylphenyl
1764.	allyl	3-fluoro-4-isopropylphenyl
1765.	allyl	4-(1-hydroxy-1-methylethyl)-phenyl
1766.	allyl	4-(2-hydroxy-2-methylpropyl)-phenyl
1767.	allyl	4-acetylphenyl
1768.	allyl	4-carboxyphenyl
1769.	allyl	4-cyanophenyl
1770.	allyl	4-hydroxyphenyl
1771.	allyl	4-(O-benzyl)-phenyl

No.	R ¹	Ar
1772.	allyl	4-(2-methoxyethoxy)-phenyl
1773.	allyl	4-(CH ₂ -N(CH ₃) ₂)-phenyl
1774.	allyl	4-(NH-CO-NH ₂)-phenyl
1775.	allyl	4-(methylsulfanyl)-phenyl
1776.	allyl	4-(fluoromethylsulfanyl)-phenyl
1777.	allyl	4-(difluoromethylsulfanyl)-phenyl
1778.	allyl	4-(trifluoromethylsulfanyl)-phenyl
1779.	allyl	4-(methylsulfonyl)-phenyl
1780.	allyl	4-(N-methoxy-N-methyl-amino)-phenyl
1781.	allyl	4-(methoxyamino)-phenyl
1782.	allyl	4-(ethoxyamino)-phenyl
1783.	allyl	4-(N-methylaminooxy)-phenyl
1784.	allyl	4-(N,N-dimethylaminooxy)-phenyl
1785.	allyl	4-(azetidin-1-yl)-phenyl
1786.	allyl	4-(2-methylazetidin-1-yl)-phenyl
1787.	allyl	4-((S)-2-methylazetidin-1-yl)-phenyl
1788.	allyl	4-((R)-2-methylazetidin-1-yl)-phenyl
1789.	allyl	4-(3-fluoroazetidin-1-yl)-phenyl
1790.	allyl	4-(3-methoxyazetidin-1-yl)-phenyl
1791.	allyl	4-(3-hydroxyazetidin-1-yl)-phenyl
1792.	allyl	4-(pyrrolidin-1-yl)-phenyl
1793.	allyl	4-(pyrrolidin-2-yl)-phenyl
1794.	allyl	4-((S)-pyrrolidin-2-yl)-phenyl
1795.	allyl	4-((R)-pyrrolidin-2-yl)-phenyl
1796.	allyl	4-(pyrrolidin-3-yl)-phenyl
1797.	allyl	4-((S)-pyrrolidin-3-yl)-phenyl
1798.	allyl	4-((R)-pyrrolidin-3-yl)-phenyl
1799.	allyl	4-(2-fluoropyrrolidin-1-yl)-phenyl
1800.	allyl	4-((S)-2-fluoropyrrolidin-1-yl)-phenyl
1801.	allyl	4-((R)-2-fluoropyrrolidin-1-yl)-phenyl
1802.	allyl	4-(3-fluoropyrrolidin-1-yl)-phenyl
1803.	allyl	4-((S)-3-fluoropyrrolidin-1-yl)-phenyl
1804.	allyl	4-((R)-3-fluoropyrrolidin-1-yl)-phenyl
1805.	allyl	4-(2,2-difluoropyrrolidin-1-yl)-phenyl
1806.	allyl	4-(3,3-difluoropyrrolidin-1-yl)-phenyl

No.	R ¹	Ar
1807.	allyl	4-(2-methylpyrrolidin-1-yl)-phenyl
1808.	allyl	4-((S)-2-methylpyrrolidin-1-yl)-phenyl
1809.	allyl	4-((R)-2-methylpyrrolidin-1-yl)-phenyl
1810.	allyl	4-(3-methylpyrrolidin-1-yl)-phenyl
1811.	allyl	4-((S)-3-methylpyrrolidin-1-yl)-phenyl
1812.	allyl	4-((R)-3-methylpyrrolidin-1-yl)-phenyl
1813.	allyl	4-(1-methylpyrrolidin-2-yl)-phenyl
1814.	allyl	4-((S)-1-methylpyrrolidin-2-yl)-phenyl
1815.	allyl	4-((R)-1-methylpyrrolidin-2-yl)-phenyl
1816.	allyl	4-(1-methylpyrrolidin-3-yl)-phenyl
1817.	allyl	4-((S)-1-methylpyrrolidin-3-yl)-phenyl
1818.	allyl	4-((R)-1-methylpyrrolidin-3-yl)-phenyl
1819.	allyl	4-(2,2-dimethylpyrrolidin-1-yl)-phenyl
1820.	allyl	4-(3,3-dimethylpyrrolidin-1-yl)-phenyl
1821.	allyl	4-(2-trifluoromethylpyrrolidin-1-yl)-phenyl
1822.	allyl	4-((S)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
1823.	allyl	4-((R)-2-trifluoromethylpyrrolidin-1-yl)-phenyl
1824.	allyl	4-(3-trifluoromethylpyrrolidin-1-yl)-phenyl
1825.	allyl	4-((S)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
1826.	allyl	4-((R)-3-trifluoromethylpyrrolidin-1-yl)-phenyl
1827.	allyl	4-(2-oxopyrrolidin-1-yl)-phenyl
1828.	allyl	4-(2-oxo-oxazolidin-3-yl)-phenyl
1829.	allyl	4-(piperidin-1-yl)-phenyl
1830.	allyl	4-(2-methylpiperidin-1-yl)-phenyl
1831.	allyl	4-((S)-2-methylpiperidin-1-yl)-phenyl
1832.	allyl	4-((R)-2-methylpiperidin-1-yl)-phenyl
1833.	allyl	4-(piperazin-1-yl)-phenyl
1834.	allyl	4-(4-methylpiperazin-1-yl)-phenyl
1835.	allyl	4-(morpholin-4-yl)-phenyl
1836.	allyl	4-(thiomorpholin-4-yl)-phenyl
1837.	allyl	4-(1-oxo-thiomorpholin-4-yl)-phenyl
1838.	allyl	4-(1,1-dioxo-thiomorpholin-4-yl)-phenyl
1839.	allyl	4-(pyrrol-1-yl)-phenyl
1840.	allyl	4-(pyrrol-2-yl)-phenyl
1841.	allyl	4-(pyrrol-3-yl)-phenyl

No.	R ¹	Ar
1842.	allyl	4-(1-methylpyrrol-2-yl)-phenyl
1843.	allyl	4-(1-methylpyrrol-3-yl)-phenyl
1844.	allyl	4-(furan-2-yl)-phenyl
1845.	allyl	4-(furan-3-yl)-phenyl
1846.	allyl	4-(thiophen-2-yl)-phenyl
1847.	allyl	4-(thiophen-3-yl)-phenyl
1848.	allyl	4-(5-propylthien-2-yl)-phenyl
1849.	allyl	4-(pyrazol-1-yl)-phenyl
1850.	allyl	4-(pyrazol-3-yl)-phenyl
1851.	allyl	4-(pyrazol-4-yl)-phenyl
1852.	allyl	4-(1-methyl-1H-pyrazol-4-yl)-phenyl
1853.	allyl	4-(1-ethyl-1H-pyrazol-4-yl)-phenyl
1854.	allyl	4-(1-methyl-1H-pyrazol-5-yl)-phenyl
1855.	allyl	4-(1H-imidazol-2-yl)-phenyl
1856.	allyl	4-(imidazol-1-yl)-phenyl
1857.	allyl	4-(1-methylimidazol-2-yl)-phenyl
1858.	allyl	4-(oxazol-2-yl)-phenyl
1859.	allyl	4-(oxazol-4-yl)-phenyl
1860.	allyl	4-(oxazol-5-yl)-phenyl
1861.	allyl	4-(isoxazol-3-yl)-phenyl
1862.	allyl	4-(isoxazol-4-yl)-phenyl
1863.	allyl	4-(isoxazol-5-yl)-phenyl
1864.	allyl	4-([1,2,3]-triazol-1-yl)-phenyl
1865.	allyl	4-([1,2,4]-triazol-1-yl)-phenyl
1866.	allyl	4-([1,2,3]-triazol-2-yl)-phenyl
1867.	allyl	4-(4H-[1,2,4]-triazol-3-yl)-phenyl
1868.	allyl	4-([1,2,4]-triazol-4-yl)-phenyl
1869.	allyl	4-(2H-[1,2,3]-triazol-4-yl)-phenyl
1870.	allyl	4-(4-methyl-4H-[1,2,4]-triazol-3-yl)-phenyl
1871.	allyl	4-(2-methyl-2H-[1,2,3]-triazol-4-yl)-phenyl
1872.	allyl	4-([1,3,4]-oxadiazol-2-yl)-phenyl
1873.	allyl	4-([1,2,4]-oxadiazol-3-yl)-phenyl
1874.	allyl	4-([1,2,4]-oxadiazol-5-yl)-phenyl
1875.	allyl	4-([1,2,3]-oxadiazol-4-yl)-phenyl
1876.	allyl	4-([1,2,3]-oxadiazol-5-yl)-phenyl


75

No.	R ¹	Ar
1877.	allyl	4-([1,2,3]-thiadiazol-4-yl)-phenyl
1878.	allyl	4-(1H-tetrazol-5-yl)-phenyl
1879.	allyl	4-(tetrazol-1-yl)-phenyl
1880.	allyl	4-(2-methyl-2H-tetrazol-5-yl)-phenyl
1881.	allyl	4-(1-methyl-1H-tetrazol-5-yl)-phenyl
1882.	allyl	4-furazan-3-yl-phenyl
1883.	allyl	4-(pyrid-2-yl)-phenyl
1884.	allyl	4-(pyrid-3-yl)-phenyl
1885.	allyl	4-(pyrid-4-yl)-phenyl
1886.	allyl	4-(pyrimidin-2-yl)-phenyl
1887.	allyl	4-(pyrimidin-4-yl)-phenyl
1888.	allyl	4-(pyrimidin-5-yl)-phenyl
1889.	allyl	5-isopropylthiophen-2-yl
1890.	allyl	2-chlorothiophen-5-yl
1891.	allyl	2,5-dichlorothiophen-4-yl
1892.	allyl	2,3-dichlorothiophen-5-yl
1893.	allyl	2-chloro-3-nitrothiophen-5-yl
1894.	allyl	2-(phenylsulfonyl)-thiophen-5-yl
1895.	allyl	2-(pyridin-2-yl)thiophen-5-yl
1896.	allyl	2-(5-(trifluoromethyl)isoxazol-3-yl)-thiophen-5-yl
1897.	allyl	2-(2-methylthiazol-4-yl)-thiophen-5-yl
1898.	allyl	1-methyl-1H-imidazol-4-yl
1899.	allyl	1,2-dimethyl-1H-imidazol-4-yl
1900.	allyl	3,5-dimethylisoxazol-4-yl
1901.	allyl	thiazol-2-yl
1902.	allyl	4-methylthiazol-2-yl
1903.	allyl	4-isopropylthiazol-2-yl
1904.	allyl	4-trifluoromethylthiazol-2-yl
1905.	allyl	5-methylthiazol-2-yl
1906.	allyl	5-isopropylthiazol-2-yl
1907.	allyl	5-trifluoromethylthiazol-2-yl
1908.	allyl	2,4-dimethylthiazol-5-yl
1909.	allyl	2-acetamido-4-methylthiazol-5-yl
1910.	allyl	4H-[1,2,4]triazol-3-yl
1911.	allyl	5-methyl-4H-[1,2,4]triazol-3-yl

76

No.	R ¹	Ar
1912.	allyl	4-methyl-4H-[1,2,4]triazol-3-yl
1913.	allyl	5-isopropyl-4H-[1,2,4]triazol-3-yl
1914.	allyl	5-trifluoromethyl-4H-[1,2,4]triazol-3-yl
1915.	allyl	4,5-dimethyl-4H-[1,2,4]triazol-3-yl
1916.	allyl	5-isopropyl-4-methyl-4H-[1,2,4]triazol-3-yl
1917.	allyl	5-trifluoromethyl-4-methyl-4H-[1,2,4]triazol-3-yl
1918.	allyl	[1,3,4]thiadiazol-2-yl
1919.	allyl	5-methyl-[1,3,4]thiadiazol-2-yl
1920.	allyl	5-isopropyl-[1,3,4]thiadiazol-2-yl
1921.	allyl	5-trifluoromethyl-[1,3,4]thiadiazol-2-yl
1922.	allyl	3-bromo-2-chloropyrid-5-yl
1923.	allyl	2-(4-morpholino)-pyrid-5-yl
1924.	allyl	2-phenoxy-pyrid-5-yl
1925.	allyl	(2-isopropyl)-pyrimidin-5-yl
1926.	allyl	(5-isopropyl)-pyrimidin-2-yl
1927.	allyl	8-quinolyl
1928.	allyl	5-isoquinolyl
1929.	allyl	2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl
1930.	allyl	5-chloro-3-methylbenzothiophen-2-yl
1931.	allyl	3,4-dihydro-4-methyl-2H-benzo[b][1,4]oxazinyl
1932.	allyl	benzothiazol-6-yl
1933.	allyl	benzo[2,1,3]oxadiazol-4-yl
1934.	allyl	5-chlorobenzo[2,1,3]oxadiazol-4-yl
1935.	allyl	7-chlorobenzo[2,1,3]oxadiazol-4-yl
1936.	allyl	benzo[2,1,3]thiadiazol-4-yl
1937.	allyl	6-chloroimidazo[2,1-b]thiazolyl

Compounds I of the present invention can be synthesized as outlined in the synthetic routes A, B and C below.



Scheme 1:

- 5 In scheme 1, A, Ar, G, n, R², and R⁴ are as defined above. R' is either R¹ or is a precursor of R¹.

Route A

- 10 In route A, the amino compound (II-1) is reacted with a suitable sulfonic acid derivative to give the sulfonamide (I-1) (E = NH). A suitable sulfonic acid derivative is e.g. the sulfonyl chloride Ar-SO₂Cl. The sulfonation reaction is preferably carried out in the presence of a base, according to standard procedures in the art. In the reaction depicted in the above scheme 1, the sulfonation takes place under the reaction conditions
- 15 which are customary for preparing arylsulfonamide compounds or arylsulfonic esters, respectively, and which are described, for example, in J. March, Advanced Organic Chemistry, 3rd edition, John Wiley & Sons, New York, 1985 page 444ff and the literature cited therein, European J. Org. Chem. 2002 (13), pp. 2094-2108, Tetrahedron 2001, 57 (27) pp. 5885-5895, Bioorganic and Medicinal Chemistry Letters, 2000, 10(8),
- 20 pp. 835-838 and Synthesis 2000 (1), pp. 103-108. The reaction customarily takes place in an inert solvent, for example in an ether, such as diethyl ether, diisopropyl ether, methyl tert-butyl ether or tetrahydrofuran, a halohydrocarbon, such as dichloromethane, an aliphatic or cycloaliphatic hydrocarbon, such as pentane, hexane or cyclohexane, or an aromatic hydrocarbon, such as toluene, xylene, cumene and the
- 25 like, or in a mixture of the abovementioned solvents. The reaction with Cl-SO₂-Ar is

customarily carried out in the presence of an auxiliary base. Suitable bases are inorganic bases, such as sodium carbonate or potassium carbonate, or sodium hydrogencarbonate or potassium hydrogencarbonate, and organic bases, for example trialkylamines, such as triethylamine, or pyridine compounds, such as pyridine, lutidine and the like.

- 5 The latter compounds can at the same time serve as solvents. The auxiliary base is customarily employed in at least equimolar quantities, based on the amine compound (II-1).

- 10 Prior to the sulfonation reaction, the radical NH_2 can be converted into a $\text{NR}^{5'}$ group, in which $\text{R}^{5'}$ has the meanings different from hydrogen which are specified for R^5 (not shown in scheme 1).

- 15 If in the resulting sulfonamide (I'-1) R' is not the desired radical R^1 but a precursor thereof, the compound can be modified as outlined below to obtain the desired substituent R^1 . A precursor is a radical which can be easily removed and replaced by the desired group R^1 or which can be modified to give R^1 . The precursor can also be an N-protective group.

- 20 If R' is allyl, the allyl group can be cleaved to obtain a compound wherein R' is hydrogen. The cleavage of the allyl group is achieved, for example, by reacting compound (I'-1) [$\text{R}' = \text{allyl}$] with an allyl trapping agent, such as mercaptobenzoic acid or 1,3-dimethylbarbituric acid, in the presence of catalytic quantities of palladium (0) compounds or palladium compounds which are able to form a palladium(0) compound under reaction conditions, e.g. palladium dichloride, tetrakis(triphenylphosphine)-
- 25 palladium(0) or tris(dibenzylideneacetone)dipalladium(0), advantageously in combination with phosphine ligands, e.g. triarylphosphines, such as triphenylphosphine, trialkylphosphines, such as tributylphosphine, and cycloalkylphosphines, such as tricyclohexylphosphine, and especially with phosphine chelate ligands, such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl or 1,4-bis(diphenylphosphino)butane, using
- 30 methods known from the literature (with regard to eliminating N-allyl in the presence of mercaptobenzoic acid, see WO 94/24088; with regard to eliminating in the presence of 1,3-dimethylbarbituric acid, see J. Am. Chem. Soc. 2001, 123 (28), pp. 6801-6808 and J. Org. Chem 2002, 67(11) pp. 3718-3723). Alternatively, the cleavage of N-allyl can also be effected by reacting in the presence of rhodium compounds, such as
- 35 tris(triphenylphosphine)chlororhodium(I), using methods known from the literature (see J. Chem. Soc., Perkin Transaction I: Organic and Bio-Organic Chemistry 1999 (21) pp. 3089-3104 and Tetrahedron Asymmetry 1997, 8(20), pp. 3387 - 3391).

If R' is benzyl, this substituent may also be cleaved to obtain a compound (I'-1) wherein R' is H. The reaction conditions for the cleavage are known in the art. Typically, the benzyl group is removed by a hydrogenation reaction in the presence of a suitable Pd catalyst, such as Pd on carbon or palladium hydroxide.

R' can also be a protective group. The protective group may be removed to yield a compound (I'-1) wherein R' is H. Suitable protective groups are known in the art and are, for example, selected from tert-butoxycarbonyl (boc), benzyloxycarbonyl (Cbz), 9-fluorenylmethoxycarbonyl (Fmoc), triphenylmethyl (Trt) and nitrobenzenesulfonyl (Nps). A preferred protective group is boc. The protective groups can be removed by known methods, such as treatment of the protected amine with an acid, e.g. halogen acid, such as HCl or HBr, or trifluoroacetic acid, or by hydrogenation, optionally in the presence of a Pd catalyst.

The resulting compound, wherein R' is H, can then be reacted, in a known manner, in the sense of an alkylation, with a compound R¹-X. In this compound, R¹ is C₁-C₄-alkyl, C₃-C₆-cycloalkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl or C₃-C₆-cycloalkyl-C₁-C₄-alkyl and X is a nucleophilically displaceable leaving group, e.g. halogen, trifluoroacetate, alkylsulfonate, arylsulfonate, alkyl sulfate and the like. The reaction conditions which are required for the alkylation have been adequately disclosed, e.g. in *Bioorganic and Medicinal Chemistry Lett.* 2002, 12(7), pp. 2443-2446 and also 2002, 12(5), pp. 1917-1919.

The alkylation can also be achieved, in the sense of a reductive amination, by reacting the compound (I'-1), wherein R' = H, with a suitable ketone or aldehyde in the presence of a reducing agent, e.g. in the presence of a borohydride such as sodium borohydride, sodium cyanoborohydride or sodium triacetoxyborohydride. The skilled person is familiar with the reaction conditions which are required for a reductive amination, e.g. from *Bioorganic and Medicinal Chemistry Lett.* 2002, 12(5), pp. 795-798 and 12(7) pp. 1269-1273.

In case R' is hydrogen, the resulting sulfonamide (I'-1) can further be reacted with an acyl halide to obtain a compound of the formula I wherein R¹ is C₁-C₃-alkylcarbonyl. The carbonyl group in these compounds can be reduced with diborane to obtain compounds of the general formula I, wherein R¹ is C₂-C₄-alkyl. The carbonyl group can also be reacted with a fluorinating agent to obtain a compound I wherein R¹ is 1,1-

difluoroalkyl. Acylation and reduction can be achieved by standard methods, which are discussed in Jerry March, *Advanced Organic Chemistry*, 3rd ed. J. Wiley & Sons, New York 1985, p.370 and 373 (acylation) and p. 1099 f. and in the literature cited in this publication (with regard to acylation, see also *Synth. Commun.* 1986, 16, p. 267, and
5 with regard to reduction, see also *J. Heterocycl. Chem.* 1979, 16, p. 1525).

Route B

In route B, the bromo substituted compound (II-2) is reacted with an appropriate sulfonamide $\text{ArSO}_2\text{NHR}^5$ to give the sulfonamide (I'-1). The reaction is generally carried out under activating conditions, e.g. under microwave conditions. Pd, especially Pd(0), or Cu catalysts may also be used for coupling (see, e.g. *Org. Lett.* 2000, 2, 1101; *J. Am. Chem. Soc.* 2002, 124, 6043; *Org. Lett.* 2003, 5, 4373; *Tetrahedron Lett.* 2003, 44, 3385). Examples for suitable Pd (0) catalysts are tetrakis(triphenylphosphine)-
10 palladium(0) and $\text{Pd}_2(\text{dba})_3$ (tris(dibenzylideneacetone) -dipalladium(0)), which are customarily used in the presence of a tri(substituted)phosphine, e.g. a triarylphosphine such as triphenylphosphine, tritolyphosphine or xantphos, tri(cyclo)alkylphosphine such as tris-n-butylphosphine, tris(tert.-butyl)phosphine or tris(cyclohexylphosphine). This route is especially useful in cases where the corresponding sulfonyl chloride is not
15 available.

Alternatively, the bromo substituent may be replaced by an amino substituent, e.g. by reacting with a benzophenone imine or with lithium bis(trimethylsilyl)amide in the presence of a palladium(0) compound such as tris(dibenzylideneacetone)dipalladium(0) in
20 the presence of a tri(substituted)phosphine, e.g. a triarylphosphine such as triphenylphosphine or tritolyphosphine, tri(cyclo)alkylphosphine such as tris-n-butylphosphine, tris(tert.-butyl)phosphine or tris(cyclohexylphosphine), preferably in the presence of a base such as sodium hydride according to the method described in (see, e.g. *J. Org. Chem.*, 68 (1993) pp 8274-8276, *J. Org. Chem.* 2000, 65, 2612). The resulting amino
25 compound may then be subjected to the sulfonation reaction of route A.

Route C


In route C, compound (II-3) is reacted with a mercapto compound HS-Ar in the presence of a base, such as sodium hydride or sodium alkoxide or with an alkali metal salt thereof thereby yielding a thioether compound. The thioether moiety is then oxidized to a sulfone moiety, e.g. by oxone, to yield the sulfone (I'-2).
30

The substituent Ar can be varied by either using different sulfonyl chlorides or by modifying the substituents of the cyclic group Ar after the formation of the sulfonamide (I'-1) by known methods. For example, a bromine substituent of the Ar group may be replaced by an N-bound pyrrolidinyl group according to the procedure described in Tetrahedron Asym. 1999, 10, 1831. A bromine substituent of the Ar group may be replaced by an isopropenyl group according to a Stille coupling where the bromo compound is reacted with an alkenyl tributyl stannate in the presence of an appropriate Pd coupling catalyst, e.g. tetrakis(triphenylphosphine) palladium(0) (see, e.g. Tetrahedron, 2003, 59(34), 6545 and Bioorg. Med. Chem. 1999, 7(5), 665). The isopropenyl group may then be converted into the isopropyl group by known hydrogenation methods.

Compounds of formula (II) (II-1, II-2 and II-3) can be synthesized by as shown below.

1. Synthesis of compounds (II-1)

Scheme 2



In scheme 2, A, G, n and R' are as defined above.

20

The conversion of the acid (III) into its methyl ester (IV) is performed by standard techniques, e.g. as described in Jerry March, Advanced Organic Chemistry, John Wiley, 3rd edition, page 348ff. For instance, the acid is transformed into the corresponding acid chloride, e.g. by reacting it with SOCl₂. The chloride is then converted into the ester by reaction with methanol.

25

The reduction in step (ii) is suitably carried out under standard conditions for the conversion of carboxylic esters into alcohols. Appropriate reaction conditions and reducing agents are described, e.g. in Jerry March, *Advanced Organic Chemistry*, John Wiley, 3rd edition, page 1093ff. Typical reducing agents are metal hydrides and complex hydrides. Examples of suitable metal hydrides include BH_3 , 9-BBN, AlH_3 and $\text{AlH}(\text{i-Bu})_2$ (DIBAL-H), suitably in the presence of complexing solvents, such as tetrahydrofuran and diethylether. Complex hydrides are e.g. NaBH_4 , LiAlH_4 and $\text{LiAlH}(\text{OR})_3$, where R is C_1 - C_4 -alkyl, such as methyl, ethyl, isobutyl or tert-butyl. A preferred reducing agent is LiAlH_4 . The reduction is suitably carried out in complexing solvents, such as open-chain and cyclic ethers, e.g. tetrahydrofuran, diethyl ether, dipropyl ether, diisopropyl ether, dibutyl ether and methylbutyl ether. A preferred solvent is tetrahydrofuran.

In the mesylation step (iii), the alcohol functionality is converted into a better leaving group. The mesylation is performed under standard conditions, e.g. by reacting the alcohol with methanesulfonyl chloride in the presence of a base. Suitable bases are, among others, alkyl amines, such as diethyl amine, triethyl amine and ethyldiisopropyl amine. In this step, other functionalities representing good leaving groups, such as trifluoroacetate, other alkylsulfonates, arylsulfonates, e.g. tosylates, alkyl sulfates and the like tosylate, may be introduced instead of the methanesulfonyl group.

In the cyclisation step (iv), compound (VI) or a suitable derivative thereof is reacted with a primary amine $\text{NH}_2\text{R}'$. In case the primary amine is a liquid, it may also be used as solvent, no further solvent being necessary. If the amine is viscous or a solid, the reaction is advantageously carried out in a suitable solvent.

The reaction of step (v) takes place under the reaction conditions which are customary for a nitration reaction on an aromatic radical and which are described, for example, in Jerry March, *Advanced Organic Chemistry*, John Wiley, 3rd edition, page 468ff, *Tetrahedron* 1999, 55(33), pp. 10243-10252, *J. Med. Chem.* 1997, 40(22), pp. 3679-3686 and *Synthetic Communications*, 1993, 23(5), pp. 591-599. For example, compound (VII) is reacted with concentrated nitric acid or a nitrate, such as potassium or sodium nitrate, in the presence of concentrated sulfuric acid. The resulting product (VIII) may in the form of different regioisomers (e.g. ortho, meta or para, if A is phenyl or a 6-membered hetaryl). In the case of A being phenyl or a 6-membered hetaryl, the para-nitro compound generally predominates. However, some ortho product may also be obtained, whereas the meta product is not produced at all or only in neglectable


amounts. By separating ortho and para products, compounds of formula I, wherein A is 1,4-bound aryl or hetaryl as well as compounds I, wherein A is 1,2-bound aryl or hetaryl, are accessible via the reaction path shown in scheme 2.

- 5 In step (vi), the nitro group in (VIII) is reduced to an NH_2 group. Subsequently, the NH_2 group can be converted into a $-\text{NR}^{5'}$ group, in which $\text{R}^{5'}$ has the meanings different from hydrogen which are specified for R^5 . The reaction conditions which are required for step (vi) correspond to the customary conditions for reducing aromatic nitro groups which have been described extensively in the literature (see, for example, J. March,
- 10 Advanced Organic Chemistry, 3rd ed., J. Wiley & Sons, New-York, 1985, p. 1183 and the literature cited in this reference). The reduction is achieved, for example, by reacting the nitro compound VII with a metal such as iron, zinc or tin under acidic reaction conditions, i.e. using nascent hydrogen, or using a complex hydride such as lithium aluminum hydride or sodium borohydride, preferably in the presence of transition metal
- 15 compounds of nickel or cobalt such as $\text{NiCl}_2(\text{P}(\text{phenyl})_3)_2$, or CoCl_2 , (see Ono et al. Chem. Ind. (London), 1983 p.480), or using NaBH_2S_3 (see Lalancette et al. Can. J. Chem. 49, 1971, p. 2990), with it being possible to carry out these reductions, depending on the given reagent, in substance or in a solvent or diluent. Alternatively, the reduction can be carried out with hydrogen in the presence of a transition metal catalyst,
- 20 e.g. using hydrogen in the presence of catalysts based on platinum, palladium, nickel, ruthenium or rhodium. The catalysts can contain the transition metal in elemental form or in the form of a complex compound, of a salt or of an oxide of the transition metal, with it being possible, for the purpose of modifying the activity, to use customary ligands, e.g. organic phosphine compounds, such as triphenylphosphine, tricyclohexylphosphine or tri-n-butylphosphines or phosphites. The catalyst is customarily employed in quantities of from 0.001 to 1 mol per mol of nitro compound, calculated as catalyst metal. In a preferred variant, the reduction is effected using tin(II) chloride in analogy with the methods described in Bioorganic and Medicinal Chemistry Letters, 2002,
- 25 12(15), pp. 1917-1919 and J. Med. Chem. 2002, 45(21), pp. 4679-4688. The reaction of VII with tin(II) chloride is preferably carried out in an inert organic solvent, preferably an alcohol such as methanol, ethanol, isopropanol or butanol.
- 30

- For compounds, wherein n is 1 and A is phenylene (i.e. (I) is N-(pyrrolidin-3-yl)-phenyl-sulfonamide), starting compound (III) is e.g. commercially available (S) or (R) phenylsuccinic acid or a racemic mixture thereof. By starting from enantiomerically pure (S)- or (R)-compound (III), pure (S)- or (R)- may be obtained:
- 35

84

a) (S) isomer




- 5 In step (i), commercially available (S)-phenylsuccinic acid (II-S) is converted into the the methyl ester (III); this is reduced to the alcohol (IV) which is reacted with methylsulfonylchloride. Cyclisation with a primary amine gives the phenyl pyrrolidine (VI). The phenyl group is first nitrated, then the nitro group is reduced to an amino function which is reacted with a sulfonyl chloride to give the desired sulfonyl amide (I'-S).

10

a) (R) isomer

The (R)-isomer can be obtained in an analogous way starting from commercially available (R)-phenylsuccinic acid (III-R):

15



c) Isomeric mixtures

20

Isomeric mixtures of compound I'-S and I'-R can be obtained by starting from racemic III or from a mixture of III-S and III-R.

The skilled person will appreciate that the synthesis described in scheme 2 is also suitable for the preparation of compounds (II) and consequently for compounds (I), wherein R^2 , R^3 and R^4 are different from H, e.g. by starting from the correspondingly substituted compound (III). The same applies to the synthesis of enantiomerically pure (I) which can be synthesized by starting from the corresponding enantiomer (III).

2. Synthesis of compounds (II-2)

Compounds of formula (II-2) can be synthesized by carrying out in step (v) of scheme 2 a halogenation instead of a nitration. Halogenation reactions of aryl and hetaryl groups are widespread standard methods and are, e.g., discussed in Jerry March, Advanced Organic Chemistry, John Wiley, 3rd edition page 476 ff.

3. Synthesis of compounds (II-3)


The synthesis of these compounds also belongs to standard reaction methods and can be performed by monohalogenating the methyl group of a methyl-substituted aryl or hetaryl compound.

4. Specific syntheses

4.1 Synthesis of compounds, wherein n is 5 (pyrrolidiny) sulfone derivatives)

4.1.1

Scheme 3:



In scheme 3, A and R^3 are as defined above.

The pyrrolidine ring is also available by a [3+2] dipolar cycloaddition of a non-stabilized azomethine ylid to a 1-alkenylaryl or hetaryl derivative (IX) (e.g. a vinyl benzene, $R^3 = H$). This procedure is generally described in J. Org. Chem 1987, 52, 235. The precursor of the ylid, the amine $N(CH_2R^b)(CH_2SiMe_3)(CH_2OCH_3)$ (X), is commercially available or can be synthesized from $NH_2(CH_2R^b)$, Me_3SiCH_2Cl and $HCHO$ in the presence of methanol.

The 1-alkenyl-(hetero)aromatic compound (IX) can be synthesized e.g. by a Stille coupling of a halogeno benzene, e.g. a bromo benzene, with the corresponding alkenyl tributyl stannate, such as vinyl or isobutenyl tributyl stannate, in the presence of an appropriate Pd coupling catalyst, e.g. tetrakis(triphenylphosphine) palladium(0) (see, e.g. Tetrahedron, 2003, 59(34), 6545 and Bioorg. Med. Chem. 1999, 7(5), 665). By choosing a special Stille isomer (e.g. cis- or trans-isobutenyl tributyl stannate), the corresponding cis- or trans alkyl phenyl pyrrolidine can be prepared selectively.

Alternatively, the 1-alkenyl-(hetero)aromatic compound (IX) can be synthesized by a Wittig reaction of aryl aldehyde with a Wittig reagent such as $PPh_3=CHR$ (R is H, or C_1 - C_3 -alkyl). Conditions for the Wittig reaction are well known in the art and are, e.g. discussed in Jerry March, Advanced Organic Chemistry, John Wiley, 3rd edition, page 845 ff.

Advantageously, the 1-(hetero)alkenyl-aromatic compound (IX) further carries a nitro group or another halogeno substituent ($X = NO_2$ or halogen). In this case, the subsequent reaction steps can be carried out as outlined in route A or B. If $X = H$, the A ring may be first nitrated as described in scheme 2, step (v) and then subjected to the reaction of scheme 2, step (vi) and scheme 1, route A; or ring A may be halogenated and then subjected to the procedure of route B.

The group CH_2R^b of the precursor amine advantageously corresponds either to the desired group R^1 of the final compound I or is alternatively a cleavable group, such as benzyl, which can be removed to give the N-unsubstituted pyrrolidine. The latter can subsequently be functionalized as described above (see route A).

The synthesis of hetarylpyrrolidines is e.g. described in Chem. Pharm. Bull., 1985, 33, 2762-66; J. Heterocyclic Chemistry, 1996, 1995-2005; J. Heterocyclic Chemistry, 2001, 38, 1039-1044; Tetrahedron Letters, 1992, 33, 44, 6607-10; Heterocycles, 1998, 48,


87

12, 2535-2541 for A being pyridylene. The synthesis of vinyl-substituted thiophene and thiazol is e.g. described in Bioorg. Med. Chem. 1999, 7(5), 665.

4.1.2

5

Scheme 4:



- Phenylpyrrolidines can also be prepared by a [3+2] dipolar cycloaddition of a non-stabilized azomethine ylid to a 1-alkynylbenzene (XII) (see, e.g., Tetrahedron 1996, 52, 59). The resulting pyrroline (XIII) or the final product (I') is then hydrogenated to the corresponding pyrrolidine (XI). If the hydrogenation is carried out under chiral conditions, e.g. by using chiral catalysts, the enantiomerically pure phenylpyrrolidine compounds can be obtained. Chiral hydrogenation catalysts are well known in the art.
- The subsequent conversion to the desired sulfonamide can be carried out as described in route A or B.

4.1.3

- Alternatively, hetarylpyrrolidinyl compounds can be prepared from hetaryl halides which are subjected to a Pd-mediated cross coupling with an organozinc pyrrolidine compound. This process is described in further detail below in route F. In this alternative, too, the hetaryl halide advantageously carries a nitro group. In this case, the conversion to the desired sulfonamides can be carried out as described in route A. Alternatively, the hetaryl halide carries a halogen atom. In this case, the conversion to the desired sulfonamides can be achieved as described in route B.

4.1.4

- Compounds I, wherein n is 1, G is CH_2 , A is 1,3-bound arylene or hetarylene and E is NH, can be prepared in a similar way compared to the 1,4-bound compound from a 3-aminoaryl or hetaryl-pyrrolidine which is reacted with an appropriate sulfonyl chloride. Advantageously, the N-atom of the pyrrolidine ring is protected by a urathane-based

protective group, like benzyloxycarbonyl (cbz) and tert-butyloxycarbonyl (boc). This group may be replaced by the desired substituent R¹ by treating the compound with an acid, such as hydrochloric acid, thus removing the acid group, and then introducing the desired substituent as described in route A.

5

The 3-aminoaryl or hetaryl-pyrrolidine may be prepared by way of a Heck reaction where a protected pyrroline is reacted with 1-iodo-3-nitrobenzene under typical Heck conditions. Catalytic hydrogenation of the pyrroline double bond and reduction of the nitro group according to the procedure described in scheme 2 yields the desired product.


10

4.2 Synthesis of N-(azetidin-3-yl)-sulfonamides

Compounds I, wherein n is 0 (azetidine compounds) can be synthesized as follows:

15

Scheme 5:



In scheme 5, Ar and R1 are as defined above. X and Y are, independently of each other, CH or N.

Starting from 1-benzhydryl-azetidin-3-ol, Pd-mediated deprotection of the amine (Tetrahedron 2002, 58, 9865-9870), carbamate formation and subsequent halogenation generate an intermediate that undergoes Zn insertion (Tetrahedron 1987, 43, 2203-2212; J. Org. Chem. 1988, 53, 2390-2392). The thus obtained organozinc species can react with an appropriate 2-halo-nitro-ring (Synlett 1998, 4, 379-380; J. Am. Chem. Soc. 2003, 125, 12527-12530) to give the nitro-aryl-azetidine core. If one utilizes a 2-halo-halo-ring, there is also the possibility to realize the direct coupling between the aryl-azetidine halide and the appropriate sulfonamides (Org. Lett. 2000, 2, 1101-1104; J. Am. Chem. Soc. 2002, 124, 6043-6048; Org. Lett. 2003, 5, 4373-4376; Tetrahedron Lett. 2003, 44, 3385-3386). The amine may be regenerated by cleavage of the carbamate (e.g. with trifluoroacetic acid in the case of a Boc carbamate) and subsequently converted into an amide by reaction with the appropriate acyl chloride. The nitro group can be reduced to the amine via tin chloride or catalytic hydrogenation (e.g. Pd-C) and then converted to the desired sulfonamide by reaction with the appropriate sulfonyl chloride in the presence of a base such as pyridine. Ultimate reduction of the amide via hydroboration furnishes the final compounds.

Of course, the reaction also applies to compounds wherein the (hetero)aromatic ring bound to the azetidine group is a 5-membered heterocyclic radical, e.g. thienyl.

4.3 Synthesis of N-(piperidin-3-yl)-sulfonamides

Further to the above-described syntheses (routes A, B and C), compounds I, wherein n is 2 and E is NR⁵ (piperidin-3-yl sulfonamides) may be prepared by starting from commercially available 3-aryl or 3-hetaryl piperidines. These starting compounds may then be converted into the amino-substituted or halogenated derivative and then be subjected to the synthetic path of route A or B.

A skilled person will readily appreciate that compounds of the formula I can also be obtained from structurally similar compounds by functional group interconversion. In particular N-bound radicals R^a can be introduced into compounds of the formula I by reacting the corresponding halogen compound, i.e. a compound of the formula I, which instead of R^a carries a halogen atom, in particular a bromine or iodine atom, with a pri-

mary or secondary amine in the presence of a base, preferably also in the presence of a palladium catalyst in terms of a Buchwald-Hartwig reaction.


If not indicated otherwise, the above-described reactions are generally carried out in a solvent at temperatures between room temperature and the boiling temperature of the solvent employed. Alternatively, the activation energy which is required for the reaction can be introduced into the reaction mixture using microwaves, something which has proved to be of value, in particular, in the case of the reactions catalyzed by transition metals (with regard to reactions using microwaves, see Tetrahedron 2001, 57, p. 9199 ff. p. 9225 ff. and also, in a general manner, "Microwaves in Organic Synthesis", André Loupy (Ed.), Wiley-VCH 2002.

The sulfonylchlorides $\text{Cl-SO}_2\text{-Ar}$ are either commercially available or can be prepared according to standard synthetic methods. Sulfonylchlorides containing a fluorinated radical R^a may be prepared by different synthetic routes, e.g. by reacting suitable hydroxy or oxo precursor (e.g. a compound $\text{Cl-SO}_2\text{-Ar}$, carrying a hydroxy or oxo substituted radical) with fluorinating reagents like DAST (diethylaminosulfurtrifluoride), morpholine-DAST, deoxo-fluor (bis(2-methoxyethyl)aminosulfur trifluoride), Ishikawa's reagent (N,N-diethyl-(1,1,2,3,3,3-hexafluoropropyl)amine; Journal of Fluorine Chemistry, 1989, 43, 371-377). More conventionally, the hydroxy group of an aromatic compound which carries a hydroxy substituted radical but not a chlorosulfonyl group, is transformed into a leaving group which is then replaced by a fluoride ion (J. Org. Chem., 1994, 59, 2898-22901; Tetrahedron Letters, 1998, 7305-6; J. Org. Chem., 1998, 63, 9587-9589, Synthesis, 1987, 920-21)). Subsequent direct chlorosulfonylation with chlorosulfonic acid (Heterocycles, 2001, 55, 9, 1789-1803; J. Org. Chem., 2000, 65, 1399-1406) or a two step process preparing first the sulfonic acid derivatives which are then transformed to the sulfonylchlorides with e.g. chlorosulfonic acid, phosphorous pentachloride (Eur. J. Med. Chem., 2002, 36, 809-828) and the like, yields the desired sulfonylchloride (Tetrahedron Letters, 1991, 33, 50 7787-7788)) Sulfonylchlorides may also be prepared by diazotation of suitable amine precursor Ar-NH_2 with sodium nitrite under acidic conditions and reaction with sulfur dioxide in acetic acid (scheme (iii); J. Org. Chem., 1960, 25, 1824-26); by oxidation of suitable heteroaryl-thiols HS-Ar or heteroaryl-benzyl-thioethers $\text{C}_6\text{H}_5\text{-CH}_2\text{-S-Ar}$ with chlorine (Synthesis, 1998, 36-38; J. Am. Chem. Soc., 1950, 74, 4890-92;) directly to the corresponding sulfonyl chlorides. The further are known in the art or may be prepared by standard methods. E.g. mercapto-pyrimidines or pyrimidinyl-benzylthioether precursors can e.g. be prepared according to literature (Chemische Berichte, 1960, 1208-11; Chemische Berichte, 1960,

95, 230-235; Collection Czechoslow. Chem. Comm., 1959, 24, 1667-1671; Austr. J. Chem., 1966, 19, 2321-30; Chemiker-Zeitung, 101, 6, 1977, 305-7; Tetrahedron, 2002, 58, 887-890; Synthesis, 1983, 641-645).

- 5 In the following schemes 6 to 8 several routes are shown which are suitable to prepare benzenesulfonyl chlorides carrying a fluorinated propyl radical.

Scheme 6:




10

- The 4-(1,1-difluoro-2-propyl)benzene-1-sulfonyl chloride intermediate can be prepared from the commercially available 2-phenylpropanoic acid. In the first step a) the 2-phenylpropanoic acid is converted to the alkyl ester by esterification with an alcohol (e.g. methanol or ethanol) under acid catalysis (e.g. HCl, SO₂Cl₂). The ester can be reduced to the corresponding 2-phenylpropanal by a reducing agent such as DIBAL (diisobutylaluminium hydride). The aldehyde is converted to the 1,1-difluoro-2-propyl derivative by reaction with a suitable fluorinating reagent like DAST (diethylaminosulfur trifluoride), morpholine-DAST, deoxo-fluor (bis(2-methoxyethyl)aminosulfur trifluoride), Ishikawa's reagent (N,N-diethyl-(1,1,2,3,3,3-hexafluoropropyl)amine; Journal of Fluorine Chemistry, 1989, 43, 371-377) (step b). The thus obtained 1,1-difluoro-2-phenylpropane can be converted into 4-(1,1-difluoro-2-propyl)benzenesulfonyl chloride by either direct chlorosulfonylation with chlorosulfonic acid (Heterocycles, 2001, 55, 9, 1789-1803; J. Org. Chem., 2000, 65, 1399-1406) (step c) or by a two step process preparing first the sulfonic acid derivatives (step d) which are then transformed to the sulfonyl chlorides (step e) by reaction with e.g. chlorosulfonic acid, phosphorous pentachloride (Eur. J. Med. Chem., 2002, 36, 809-828); through diazotisation of suitable amine precursors with sodium nitrite under acidic conditions and reaction with sulfur dioxide in acetic acid (J. Org. Chem., 1960, 25, 1824-26); oxidation of suitable
- 15
- 20
- 25

heteroaryl-thiols or heteroaryl-benzyl-thioethers with chlorine (Synthesis, 1998, 36-38; J. Am. Chem. Soc., 1950, 74, 4890-92) directly to the corresponding sulfonyl chlorides.

5 The synthesis shown in scheme 6 can also be performed using (R)-2-phenylpropanoic acid and (S)-2-phenylpropanoic acid respectively to give the corresponding chiral 4-(1,1-difluoropropan-2-yl)benzene-1-sulfonyl chlorides.

Scheme 7:



10

4-(1,1,1-Trifluoropropan-2-yl)benzene-1-sulfonyl chloride intermediate can be prepared from the commercially available 2,2,2-trifluoro-1-phenylethanone by a synthetic route shown in scheme 7. The ketone can be converted to the 3,3,3-trifluoro-2-phenylpropene by a Wittig reaction with a suitable ylide such as methylene-triphenylphosphane (prepared by reaction of methyltriphenylphosphonium halide and a suitable base such as lithium diisopropylamide or potassium tert-butoxide) or according to a Horner-Emmons reaction by reacting the ketone with a suitable phosphonate such as diethyl methylphosphonate and a suitable suitable base such as lithium diisopropylamide or potassium tert-butoxide. The thus obtained 3,3,3-trifluoro-2-phenylpropene can then be reduced to the saturated alkane by catalytic hydrogenation (eg Pd-C) followed by conversion to the sulfonyl chloride by the methods described in scheme 6.


20

The synthesis of scheme 7 can also be performed using a chiral catalyst for the alkene hydrogenation to allow the preparation of the corresponding chiral 4-(1,1,1-trifluoropropan-2-yl)benzene-1-sulfonyl chlorides.

25

Scheme 8:

93



The 4-(1,1,1-trifluoropropan-2-yl)benzene-1-sulfonyl chloride can be also prepared from the commercially available 1-phenyl-ethanone by a four step procedure as shown in scheme 8. The ketone can be converted to the trifluoromethyl hydroxyl intermediate by reaction with trimethyl-trifluoromethyl-silane (Journal of Organic Chemistry, 2000, 65, 8848-8856; Journal of Fluorine Chemistry, 2003, 122, 243-246) which can then be converted to the trifluoromethyl bromide (Journal of the American Chemical Society, 1987, 109, 2435-4). Dehalogenation by catalytic hydrogenation (eg Pd-C) can then be followed by conversion to the sulfonyl chloride by the methods discussed above.

Examples of solvents which can be used are ethers, such as diethyl ether, diisopropyl ether, methyl *tert*-butyl ether or tetrahydrofuran, aprotic polar solvent, such as dimethyl-formamide, dimethyl sulfoxide, dimethoxyethane, and acetonitrile, aromatic hydrocarbons, such as toluene and xylene, ketones, such as acetone or methyl ethyl ketone, halohydrocarbons, such as dichloromethane, trichloromethane and dichloroethane, esters, such as ethyl acetate and methyl butyrate, carboxylic acids, such as acetic acid or propionic acid, and alcohols, such as methanol, ethanol, n-propanol, isopropanol, n-butanol, isobutanol, 2-butanol and *tert*-butanol.

If desired, it is possible for a base to be present in order to neutralize protons which are released in the reactions. Suitable bases include inorganic bases, such as sodium carbonate, potassium carbonate, sodium hydrogen carbonate or potassium hydrogen carbonate, and, in addition, alkoxides, such as sodium methoxide or sodium ethoxide, alkali metal hydrides, such as sodium hydride, and also organometallic compounds, such as butyllithium compounds or alkylmagnesium compounds, or organic nitrogen bases, such as triethylamine or pyridine. The latter compounds can at the same time serve as solvents.

The crude product is isolated in a customary manner, for example by filtering, distilling off the solvent or extracting from the reaction mixture, etc. The resulting compounds

can be purified in a customary manner, for example by means of recrystallizing from a solvent, by means of chromatography or by means of converting into an acid addition salt.

5 The acid addition salts are prepared in a customary manner by mixing the free base with a corresponding acid, where appropriate in solution in an organic solvent, for example a lower alcohol, such as methanol, ethanol or propanol, an ether, such as methyl *tert*-butyl ether or diisopropyl ether, a ketone, such as acetone or methyl ethyl ketone, or an ester, such as ethyl acetate.

10

The compounds according to the invention of the formula I are surprisingly highly selective dopamine D₃ receptor ligands which, because of their low affinity for other receptors such as D₁ receptors, D₄ receptors, α1-adrenergic and/or α2-adrenergic receptors, muscarinergic receptors, histamine receptors, opiate receptors and, in particular,
15 dopamine D₂ receptors, give rise to fewer side-effects than do the classic neuroleptics, which are D₂ receptor antagonists. A compound of the invention can be a dopamine D₃ receptor agonist, including partial agonistic activity, or a dopamine D₃ receptor antagonist, including partial antagonistic activity.

20

The high affinity of the compounds according to the invention for D₃ receptors is reflected in very low in-vitro receptor binding constants ($K_i(D_3)$ values) of as a rule less than 50 nM (nmol/l), preferably of less than 10 nM and, in particular of less than 5 nM. The displacement of [¹²⁵I]-iodosulpride can, for example, be used in receptor binding studies for determining binding affinities for D₃ receptors.

25

The selectivity of the compounds according to the invention, i.e. the ratio $K_i(D_2)/K_i(D_3)$ of the receptor binding constants, is as a rule at least 50, preferably at least 100, even better at least 150. The displacement of [³H]SCH23390, [¹²⁵I] idosulpride or [¹²⁵I] spiperone can be used, for example, for carrying out receptor binding studies on D₁, D₂
30 and D₄ receptors.

35

Because of their binding profile, the compounds can be used for treating diseases which respond to dopamine D₃ receptor ligands (or which are susceptible to treatment with a dopamine D₃ receptor ligand, respectively), i.e. they are effective for treating
those medical disorders or diseases in which exerting an influence on (modulating) the dopamine D₃ receptors leads to an improvement in the clinical picture or to the disease

being cured. Examples of these diseases are disorders or diseases of the central nervous system.

Disorders or diseases of the central nervous system are understood as meaning disorders which affect the spinal chord and, in particular, the brain. Within the meaning of the invention, the term "disorder" denotes disturbances and/or anomalies which are as a rule regarded as being pathological conditions or functions and which can manifest themselves in the form of particular signs, symptoms and/or malfunctions. While the treatment according to the invention can be directed toward individual disorders, i.e. anomalies or pathological conditions, it is also possible for several anomalies, which may be causatively linked to each other, to be combined into patterns, i.e. syndromes, which can be treated in accordance with the invention.

The disorders which can be treated in accordance with the invention are, in particular, psychiatric and neurological disturbances. These disturbances include, in particular, organic disturbances, including symptomatic disturbances, such as psychoses of the acute exogenous reaction type or attendant psychoses of organic or exogenous cause, e.g., in association with metabolic disturbances, infections and endocrinopathologies; endogenous psychoses, such as schizophrenia and schizotypal and delusional disturbances; affective disturbances, such as depressions, mania and/or manic-depressive conditions; and also mixed forms of the above-described disturbances; neurotic and somatoform disturbances and also disturbances in association with stress; dissociative disturbances, e.g. loss of consciousness, clouding of consciousness, double consciousness and personality disturbances; disturbances in attention and waking/sleeping behavior, such as behavioral disturbances and emotional disturbances whose onset lies in childhood and youth, e.g. hyperactivity in children, intellectual deficits, in particular attention disturbances (attention deficit disorders), memory disturbances and cognitive disturbances, e.g. impaired learning and memory (impaired cognitive function), dementia, narcolepsy and sleep disturbances, e.g. restless legs syndrome; development disturbances; anxiety states, delirium; sexual disturbances, e.g. impotence in men; eating disturbances, e.g. anorexia or bulimia; addiction; and other unspecified psychiatric disturbances.

The disorders which can be treated in accordance with the invention also include Parkinson's disease and epilepsy and, in particular, the affective disturbances connected thereto.

The addiction diseases include psychic disorders and behavioral disturbances which are caused by the abuse of psychotropic substances, such as pharmaceuticals or narcotics, and also other addiction diseases, such as addiction to gaming (impulse control disorders not elsewhere classified). Examples of addictive substances are: opioids (e.g. morphine, heroin and codeine), cocaine; nicotine; alcohol; substances which interact with the GABA chloride channel complex, sedatives, hypnotics and tranquilizers, for example benzodiazepines; LSD; cannabinoids; psychomotor stimulants, such as 3,4-methylenedioxy-N-methylamphetamine (ecstasy); amphetamine and amphetamine-like substances such as methylphenidate and other stimulants including caffeine. Addictive substances which come particularly into consideration are opioids, cocaine, amphetamine or amphetamine-like substances, nicotine and alcohol.

With regard to the treatment of addiction diseases, particular preference is given to those compounds according to the invention of the formula I which themselves do not possess any psychotropic effect. This can also be observed in a test using rats, which, after having been administered compounds which can be used in accordance with the invention, reduce their self administration of psychotropic substances, for example cocaine.

According to another aspect of the present invention, the compounds according to the invention are suitable for treating disorders whose causes can at least partially be attributed to an anomalous activity of dopamine D₃ receptors.

According to another aspect of the present invention, the treatment is directed, in particular, toward those disorders which can be influenced, within the sense of an expedient medicinal treatment, by the binding of preferably exogeneously administered binding partners (ligands) to dopamine D₃ receptors.

The diseases which can be treated with the compounds according to the invention are frequently characterized by progressive development, i.e. the above-described conditions change over the course of time; as a rule, the severity increases and conditions may possibly merge into each other or other conditions may appear in addition to those which already exist.

The compounds according to the invention can be used to treat a large number of signs, symptoms and/or malfunctions which are connected with the disorders of the central nervous system and, in particular, the abovementioned conditions. These signs,

symptoms and/or malfunctions include, for example, a disturbed relationship to reality, lack of insight and ability to meet customary social norms or the demands made by life, changes in temperament, changes in individual drives, such as hunger, sleep, thirst, etc., and in mood, disturbances in the ability to observe and combine, changes in personality, in particular emotional lability, hallucinations, ego-disturbances, distracted-
5 ness, ambivalence, autism, depersonalization and false perceptions, delusional ideas, chanting speech, lack of synkinesia, short-step gait, flexed posture of trunk and limbs, tremor, poverty of facial expression, monotonous speech, depressions, apathy, impeded spontaneity and decisiveness, impoverished association ability, anxiety, nervous
10 agitation, stammering, social phobia, panic disturbances, withdrawal symptoms in association with dependency, manifold syndromes, states of excitation and confusion, dysphoria, dyskinetic syndromes and tic disorders, e.g. Huntington's chorea and Gilles-de-la-Tourette's syndrome, vertigo syndromes, e.g. peripheral positional, rotational and oscillatory vertigo, melancholia, hysteria, hypochondria and the like.

15 Within the meaning of the invention, a treatment also includes a preventive treatment (prophylaxis), in particular as relapse prophylaxis or phase prophylaxis, as well as the treatment of acute or chronic signs, symptoms and/or malfunctions. The treatment can be orientated symptomatically, for example as the suppression of symptoms. It can be
20 effected over a short period, be orientated over the medium term or can be a long-term treatment, for example within the context of a maintenance therapy.

Therefore the compounds according to the invention are preferentially suitable for treating diseases of the central nervous system, in particular for treating affective disorders;
25 neurotic disturbances, stress disturbances and somatoform disturbances and psychoses, and, in particular, for treating schizophrenia and depression. Because of their high selectivity with regard to the D₃ receptor, the compounds I according to the invention are also suitable for treating disturbances of kidney function, in particular disturbances of kidney function which are caused by diabetes mellitus (see WO 00/67847) and, especially, diabetic nephropathy.
30

Within the context of the treatment, the use according to the invention of the described compounds involves a method. In this method, an effective quantity of one or more compounds, as a rule formulated in accordance with pharmaceutical and veterinary
35 practice, is administered to the individual to be treated, preferably a mammal, in particular a human being, productive animal or domestic animal. Whether such a treatment is indicated, and in which form it is to take place, depends on the individual case

and is subject to medical assessment (diagnosis) which takes into consideration signs, symptoms and/or malfunctions which are present, the risks of developing particular signs, symptoms and/or malfunctions, and other factors.

- 5 As a rule, the treatment is effected by means of single or repeated daily administration, where appropriate together, or alternating, with other active compounds or active compound-containing preparations such that a daily dose of preferably from about 0.1 to 1000 mg/kg of bodyweight, in the case of oral administration, or of from about 0.1 to 100 mg/kg of bodyweight, in the case of parenteral administration, is supplied to an individual to be treated.

The invention also relates to the production of pharmaceutical compositions for treating an individual, preferably a mammal, in particular a human being, productive animal or domestic animal. Thus, the ligands are customarily administered in the form of pharmaceutical compositions which comprise a pharmaceutically acceptable excipient together with at least one compound according to the invention and, where appropriate, other active compounds. These compositions can, for example, be administered orally, rectally, transdermally, subcutaneously, intravenously, intramuscularly or intranasally.

20 Examples of suitable pharmaceutical formulations are solid medicinal forms, such as powders, granules, tablets, in particular film tablets, lozenges, sachets, cachets, sugar-coated tablets, capsules, such as hard gelatin capsules and soft gelatin capsules, suppositories or vaginal medicinal forms, semisolid medicinal forms, such as ointments, creams, hydrogels, pastes or plasters, and also liquid medicinal forms, such as solutions, emulsions, in particular oil-in-water emulsions, suspensions, for example lotions, injection preparations and infusion preparations, and eyedrops and eardrops. Implanted release devices can also be used for administering inhibitors according to the invention. In addition, it is also possible to use liposomes or microspheres.

30 When producing the compositions, the compounds according to the invention are optionally mixed or diluted with one or more excipients. Excipients can be solid, semisolid or liquid materials which serve as vehicles, carriers or medium for the active compound.

35 Suitable excipients are listed in the specialist medicinal monographs. In addition, the formulations can comprise pharmaceutically acceptable carriers or customary auxiliary substances, such as glidants; wetting agents; emulsifying and suspending agents; pre-

servatives; antioxidants; antiirritants; chelating agents; coating auxiliaries; emulsion stabilizers; film formers; gel formers; odor masking agents; taste corrigents; resin; hydrocolloids; solvents; solubilizers; neutralizing agents; diffusion accelerators; pigments; quaternary ammonium compounds; refatting and overfatting agents; raw materials for ointments, creams or oils; silicone derivatives; spreading auxiliaries; stabilizers; sterilants; suppository bases; tablet auxiliaries, such as binders, fillers, glidants, disintegrants or coatings; propellants; drying agents; opacifiers; thickeners; waxes; plasticizers and white mineral oils. A formulation in this regard is based on specialist knowledge as described, for example, in Fiedler, H.P., Lexikon der Hilfsstoffe für Pharmazie, Kosmetik und angrenzende Gebiete [Encyclopedia of auxiliary substances for pharmacy, cosmetics and related fields], 4th edition, Aulendorf: ECV-Editio-Kantor-Verlag, 1996.

The following examples serve to explain the invention without limiting it.

- 15 The compounds were either characterized via proton-NMR in d₆-dimethylsulfoxid or d-chloroform, if not stated otherwise, on a 400 MHz or 500 MHz NMR instrument (Bruker AVANCE), or by mass spectrometry, generally recorded via HPLC-MS in a fast gradient on C18-material (electrospray-ionisation (ESI) mode), or melting point.
- 20 The magnetic nuclear resonance spectral properties (NMR) refer to the chemical shifts (δ) expressed in parts per million (ppm). The relative area of the shifts in the ¹H NMR spectrum corresponds to the number of hydrogen atoms for a particular functional type in the molecule. The nature of the shift, as regards multiplicity, is indicated as singlet (s), broad singlet (s. br.), doublet (d), broad doublet (d br.), triplet (t), broad triplet (t br.), quartet (q), quintet (quint.) and multiplet (m).

Preparation Examples:

- I. Preparation of the intermediates
- 30 a. Preparation of sulfonyl chlorides
- a.1 4-((S)-2-Fluoro-1-methyl-ethyl)-benzenesulfonyl chloride
- 35 a.1.1 Toluene-4-sulfonic acid (S)-2-phenyl-propyl ester

100

To a solution of 20 g of (S)-(-)-2-phenyl-1-propanol in 240 ml of dichloromethane was added in portions 28 g of p-toluenesulfonyl chloride (146.8 mmol). After stirring for 18 h at room temperature, the organic phase was washed with 100 ml of water, dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to yield 43 g of the title compound.

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.65 (d, 2H), 7.15-7.3 (m, 5H), 7.1 (d, 2H), 4.0-4.1 (m, 2H), 3.1 (m, 1H), 2.4 (s, 3H), 1.3 (d, 3H).

10 a.1.2 ((S)-2-Fluoro-1-methyl-ethyl)-benzene

9.62 g of toluene-4-sulfonic acid (S)-2-phenyl-propyl ester (33.13 mmol) were dissolved in 80 ml of polyethylenglycol 400. 9.62 g of potassium fluoride (165.6 mmol) were added and the reaction mixture was stirred at 50°C for 3 days and another 2 days at 55-70°C. The reaction was treated with 150 ml of saturated aqueous sodium chloride solution, extracted three times with diethyl ether, and the combined organic layers were dried over magnesium sulfate, filtered, and the solvent was evaporated under reduced pressure. The crude product was purified via silica gel chromatography using cyclohexane/ethyl acetate 15% as eluent. 2.85 g of the desired product were isolated, containing ~ 25% of the elimination side product.

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.2-7.4 (m, 5H), 4.3-4.6 (several m, 2H), 3.15 (m, 1H), 1.3 (m, 3H).

25

a.1.3 4-((S)-2-Fluoro-1-methyl-ethyl)-benzenesulfonyl chloride

3.5 g of ((S)-2-fluoro-1-methyl-ethyl)-benzene (25.32 mmol) were dissolved in 80 ml of dichloromethane. At 0-5°C, 11.81 g of chlorosulfonic acid (101.31 mmol), dissolved in 20 ml of dichloromethane, were added dropwise. The reaction mixture was stirred for 30 min at room temperature and 2 h at 30°C. The solvent was evaporated. 150 ml of diethyl ether were added to the residue, washed once with 150 ml of water, and the organic layer was dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure. The crude product was purified via silica gel chromatography with n-heptane-dichloromethane (6:4) as eluent to give 1.5 g of the title compound.

35

101

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.0 (d, 2H), 7.5 (d, 2H), 4.5 (dd, 2H), 3.25 (m, 1H), 1.4 (d, 3H).

a.2 4-((R)-2-Fluoro-1-methyl-ethyl)-benzenesulfonyl chloride

5

a.2.1 Toluene-4-sulfonic acid (R)-2-phenyl-propyl ester

Following the procedure analogous to that used for the synthesis of toluene-4-sulfonic acid (S)-2-phenyl-propyl ester, but using (R)-2-phenyl-1-propanol, the title compound was prepared

10

a.2.2 ((R)-2-Fluoro-1-methyl-ethyl)-benzene

The title compound was prepared as described above for the synthesis of ((S)-2-fluoro-1-methyl-ethyl)-benzene, but using toluene-4-sulfonic acid (R)-2-phenyl-propyl ester instead of toluene-4-sulfonic acid (S)-2-phenyl-propyl ester.

15

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.2-7.4 (m, 5H), 4.3-4.6 (several m, 2H), 3.15 (m, 1H). 1.3 (m, 3H).

20

a.2.3 4-((R)-2-Fluoro-1-methyl-ethyl)-benzenesulfonyl chloride

1.3 g of ((R)-2-fluoro-1-methyl-ethyl)-benzene (9.4 mmol) were dissolved in 50 ml of dichloromethane. At 0-5°C, 1.1 g of chlorosulfonic acid (9.4 mmol), dissolved in 10 ml of dichloromethane, were added dropwise. The reaction mixture was stirred for 20 min at 0-5°C and then added to a solution of 2.15 g of phosphorous pentachloride dissolved in 40 ml of dichloromethane. The reaction mixture was stirred for 30 min at 0-5°C and 1 h at room temperature. The solvent was evaporated, 100 ml of diethyl ether were added, the mixture was washed once with 150 ml of water, and the organic layer dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure. The crude product was purified via silica gel chromatography with n-heptane-dichloromethane (1:1) as eluent to give 0.261 g of the title compound.

25

30

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.0 (d, 2H), 7.5 (d, 2H), 4.5 (dd, 2H), 3.25 (m, 1H), 1.4 (d, 3H).

35

102

a.3 4-(2-Fluoro-1-methyl-ethyl)-benzenesulfonyl chloride

Following the procedures analogous to that used for the preparation of 4-((S)-2-fluoro-1-methyl-ethyl)-benzenesulfonyl chloride, but starting with 2-phenyl-1-propanol in step a.3.1, the title compound was prepared.

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.0 (d, 2H), 7.5 (d, 2H), 4.5 (dd, 2H), 3.25 (m, 1H), 1.4 (d, 3H).

a.4 4-(2-Fluoro-1-fluoromethyl-ethyl)-benzenesulfonyl chloride

a.4.1 (2-Fluoro-1-fluoromethyl-ethyl)-benzene

4 g of 3-phenylglutaric acid (19.21 mmol) were suspended in 350 ml of dichloromethane. At room temperature, 6.5 g of xenon difluoride (38.42 mmol) were added and the reaction mixture was stirred at room temperature for 18 h. The organic phase was washed once with 975 ml of 6% aqueous sodium hydrogencarbonate, dried over magnesium sulfate, filtered, and the solvent evaporated. The remaining residue was distilled at a bath temperature of 123°C at 21 mm to yield 0.78 g of the title compound that contained ~ 50% of 4-(2-Fluoro-1-methyl-ethyl)-benzene.

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.2-7.4 (m, 5H), 4.6-4.8 (dd, 4H), 3.3 (m, 1H).

a.4.2 4-(2-Fluoro-1-fluoromethyl-ethyl)-benzenesulfonyl chloride

Following the procedures analogous to that used for the preparation of 4-((S)-2-fluoro-1-methyl-ethyl)-benzenesulfonyl chloride, but using 5 equivalents of chlorosulfonic acid, 0,12 g of the title compound were obtained.

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.05 (d, 2H), 7.55 (d, 2H), 4.75 (dd, 4H), 3.4 (m, 1H).

a.5 4-(3,3,3-Trifluoropropyl)-benzenesulfonyl chloride

103

2.9 g were obtained from commercially available (3,3,3-trifluoropropyl)-benzene following the procedure used for the synthesis of 4-((S)-2-fluoro-1-methyl-ethyl)-benzenesulfonyl chloride described above.

5 ¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.0 (d, 2H), 7.45 (d, 2H), 3.0 (t, 2H), 2.45 (m, 2H).

a.6 4-(2,2,2-Trifluoroethyl)-benzenesulfonyl chloride

10 The product was obtained from commercially available (2,2,2-trifluoroethyl)-benzene following the procedure as described in J. Org. Chem., 1960, 25, 1824-26.

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.05 (d, 2H), 7.55 (d, 2H), 3.5 (q, 2H).

15

a.7 4-(3-Fluoropropyl)-benzenesulfonyl chloride

a.7.1 (3-Fluoropropyl)-benzene

20 15.6 g of diethylaminosulfurtrifluoride (DAST, 96.91 mmol) were dissolved in 18 ml of dichloromethane. At 0-5°C, 12 g of 3-phenyl-1-propanol (88.1 mmol) dissolved in 30 ml of dichloromethane, were added dropwise. The reaction mixture was stirred for 18 h, and, after addition of 30 ml of dichloromethane, poured onto 100 ml of ice water. The organic layer was separated, dried over magnesium sulfate, filtered, and the solvent evaporated. The crude product was purified by distillation at a bath temperature of 106°C at 20 mm to yield 7.4 g of the title compound.

25

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.1-7.3 (m, 5H), 4.4 (dt, 2H), 2.7 (m, 2H). 2.0 (m, 2H).

30

a.7.2 4-(3-Fluoropropyl)-benzenesulfonyl chloride

35 4.1 g of (3-fluoro-propyl)-benzene (29.67 mmol) were dissolved in 40 ml of dichloromethane. At 0-5°C, 6.91 g of chlorosulfonic acid (59.34 mmol), dissolved in 10 ml of dichloromethane, were added dropwise. The reaction mixture was stirred for 45 min at 0-5°C and then added to a solution of 6.8 g of phosphorous

104

pentachloride (32.63 mmol) dissolved in 50 ml of dichloromethane. The reaction mixture was stirred for 1 h at 5-10°C. The solvent was evaporated, 150 ml of diethyl ether added, washed once with 150 ml of ice water, and the organic layer dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure. The crude product was purified via silica gel chromatography with n-heptane-dichloromethane (11:9) as eluent to give 5.5 g of the title compound.

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.95 (d, 2H), 7.45 (d, 2H), 4.5 (dt, 2H), 2.9 (t, 2H), 2.05 (m, 2H).

a.8 4-(2,2-Difluoro-cyclopropyl)-benzenesulfonyl chloride

2.07 g of were obtained from commercially available (2,2-difluorocyclopropyl)-benzene following the procedure used for the synthesis of (3-fluoropropyl)-benzenesulfonyl chloride with the exception that only 1.1 equivalents of phosphorous pentachloride were used.

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.0 (d, 2H), 7.45 (d, 2H), 2.85 (m, 1H), 2.0 (m, 1H), 1.75 (m, 1H).

a.9 3-Bromo-4-trifluoromethoxy-benzenesulfonyl chloride

2.0 g of 1-bromo-2-(trifluoro-methoxy)benzene (8.3 mmol) were dissolved in 30 ml of dichloromethane. At 0-5°C, 1.06 g of chlorosulfonic acid (9.13 mmol), dissolved in 3 ml of dichloromethane, were added dropwise. The reaction mixture was stirred for 30 min at room temperature. Additional 5.5 equivalents of chlorosulfonic in dichloromethane were added to drive the reaction to completion. Standard work-up was followed and silica gel chromatography with n-heptane-dichloromethane (6:4) as eluent gave 2.19 g of the title compound.

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.3 (d, 1H), 8.05 (dd, 1H), 7.5 (dd, 1H).

a.10 4-(2-Fluoroethyl)-benzenesulfonyl chloride

a.10.1 (2-Fluoroethyl)-benzene

105

6.8 g of the title compound were obtained from commercially available 2-phenyl-ethanol following the procedure used for the synthesis of (3-fluoropropyl)-benzene.

5 $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ [ppm] 7.1-7.3 (m, 5H), 4.6 (m, 1H), 4.45 (m, 1H), 2.95 (m, 1H), 2.9 (m, 1H).

a.10.2 4-(2-Fluoroethyl)-benzenesulfonyl chloride

10

3.55 g were obtained following the procedure used for the synthesis of 4-((R)-2-fluoro-1-methyl-ethyl)-benzenesulfonyl chloride.

15 $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ [ppm] 8.0 (d, 2H), 7.5 (d, 2H), 4.7 (dt, 2H), 3.05-3.2 (dt, 2H).

a.11 5-Propylthiophene-2-sulfonyl chloride

20

Following the procedures analogous to that used for the preparation of (3-fluoropropyl)-benzenesulfonyl chloride, but using only 1 equivalent of phosphorous pentachloride, the title compound was prepared.

25 $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ [ppm] 7.7 (d, 1H), 6.85 (d, 1H), 2.9 (t, 2H), 1.75 (m, 2H), 1.0 (t, 3H).

a.12 4-(1-Methyl-1H-pyrazol-4-yl)-benzenesulfonyl chloride

a.12.1 1-Methyl-4-phenyl-1H-pyrazole

30 1 g of 2-phenylmalonaldehyde (6.75 mmol) were dissolved in 25 ml of ethanol. 0.36 ml of N-methyl-hydrazine (6.75 mmol) were added, the reaction mixture was stirred under reflux for 4 h, the solvent evaporated under reduced pressure to yield 1.09 g of the product.

35 ESI-MS: 159.1 $[\text{M}+\text{H}]^+$

106

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.75 (s, 1H), 7.6 (s, 1H), 7.45 (d, 2H), 7.35 (t, 2H), 7.2 (t, 1H), 3.9 (s, 3H)

a.12.2 4-(1-Methyl-1H-pyrazol-4-yl)-benzenesulfonyl chloride

5

0.5 g of 1-methyl-4-phenyl-1H-pyrazole (3.16 mmol) were dissolved in 20ml of dichloromethane. At 0°C, 0.232 ml of chlorosulfonic acid were added and the reaction mixture was stirred for 1 h under ice cooling. Additional 0.7 ml of chlorosulfonic acid were added, the mixture was stirred at 0°C for 30 minutes and then 90 minutes at 50°C. The two phases were separated and the lower layer put on ice, extracted twice with diethyl ether, dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to yield 0.496 g of the product.

10

15

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.0 (d, 2H), 7.85 (s, 1H), 7.75 (s, 1H), 7.65 (d, 2H), 4.0 (s, 3H).

a.13 4-(1,1,1-Trifluoropropan-2-yl)benzenesulfonyl chloride and
2-(1,1,1-trifluoropropan-2-yl)benzenesulfonyl chloride

20

Prepared on a 14 g scale following the procedure outlined in Scheme 7. 2-(1,1,1-Trifluoropropan-2-yl)benzenesulfonyl chloride is a by-product of the reaction.

25

4-(1,1,1-Trifluoropropan-2-yl)benzenesulfonyl chloride:

MS (ESI) m/z: 273.1 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 7.62 (d, 2H), 7.33 (d, 2H), 3.81 (m, 1H), 1.42 (d, 3H).

30

2-(1,1,1-Trifluoropropan-2-yl)benzenesulfonyl chloride:

MS (ESI) m/z: 273.1 [M+H]⁺

a.14 4-(1,1-Difluoropropan-2-yl)benzenesulfonyl chloride and
2-(1,1-Difluoropropan-2-yl)benzene-1-sulfonyl chloride

Prepared on an 11 g scale following the procedure outlined in Scheme 6. 2-(1,1-Difluoropropan-2-yl)benzene-1-sulfonyl chloride is a by-product of the reaction.

35

4-(1,1-Difluoropropan-2-yl)benzenesulfonyl chloride:

MS (ESI) m/z : 255.0 $[M+H]^+$

$^1\text{H-NMR}$ (DMSO- d_6): δ [ppm] 8.03 (d, 2H), 7.55 (d, 2H), 5.88 (dt, 1H), 3.34 (m, 1H), 1.47 (d, 3H).

$^{13}\text{C-NMR}$ (DMSO- d_6): δ [ppm] 146.43, 143.54, 129.77, 127.28, 117.06 (t), 43.76, 13.78.

2-(1,1-difluoropropan-2-yl)benzene-1-sulfonyl chloride:

Isolated by chromatography on 110 mg scale.

MS (ESI) m/z : 255.0 $[M+H]^+$

$^1\text{H-NMR}$ (DMSO- d_6): δ [ppm] 8.15 (d, 1H), 7.77 (t, 1H), 7.70 (d, 1H), 7.54 (t, 1H), 5.99 (dt, 1H), 4.43 (m, 1H), 1.51 (d, 3H).

$^{13}\text{C-NMR}$ (DMSO- d_6): δ [ppm] 143.45, 138.63, 135.53, 130.93, 129.04, 128.17, 116.61 (t), 38.38, 13.68.

b. Preparation of toluene-4-sulfonic acid 3-fluoro-propyl ester

5 g of 3-fluoro-propanol (64.03 mmol) and 18 ml of triethylamine (129.32 mmol) were dissolved in 50 ml of dichloromethane. At 0-5°C, 12.9 g toluene-4-sulfonylchloride (67.66 mmol) were added and the reaction mixture stirred at room temperature for 18 h. Standard work-up yielded 13.7 g of toluene-4-sulfonic acid 3-fluoro-propyl ester.

ESI-MS: 233.1 $[M+H]^+$

II. Preparation of the compounds I

EXAMPLE 1

4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

1.1 (S)-2-Phenyl-succinic acid dimethyl ester

5 g of (S)-2-phenyl succinic acid (25.75 mmol) were dissolved in 50 ml of methanol. At 4°C, 4.7 ml of thionyl chloride (64.37 mmol) were added dropwise. The reaction mixture was stirred at room temperature for 2h, the solvents were evaporated under reduced pressure. The residue that remained was dissolved in diethyl ether, washed once with saturated aqueous NaHCO_3 solution, reextracted with diethyl ether, and the combined

organic layers dried over magnesium sulfate, filtered, and evaporated to dryness to yield 5.8 g of the desired product.

ESI-MS: 223.1 $[M+H]^+$

5 1.2 (S)-2-Phenyl-butane-1,4-diol

2.54 g of lithium aluminium hydride (66.95 mmol) were suspended under ice cooling in 25 ml of tetrahydrofuran. 5.8 g of (S)-2-phenyl succinic acid dimethyl ester (25.75 mmol) dissolved in 25 ml of tetrahydrofuran were added slowly at 5-10°C. Stirring was continued for 15 minutes and then 15 ml of tetrahydrofuran/water (1:1) were added dropwise. The suspension was adjusted to pH 3-4 with conc. hydrochloric acid, filtered and the filter washed with dichloromethane. The filtrate was evaporated to dryness, taken up in diethylether, washed with saturated sodium hydrogen carbonate solution, reextracted with diethylether, and the combined organic layers dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to yield 4.2 g of the diol.

ESI-MS: 189.1 $[M+Na]^+$

$^1\text{H-NMR}$ (CDCl_3): δ [ppm] 7.25-7.4 (m, 2H), 7.15-7.3 (m, 3H), 4.2-4.35 (m, 2H), 3.2 (m, 1H), 3.1 (m, 1H), 2.1-2.3 (m, 3H).

20

1.3 Methanesulfonic acid (S)-4-methanesulfonyloxy-3-phenyl-butyl ester

4.19 g of (S)-2-phenyl-butane-1,4-diol (25.21 mmol) were dissolved in 50 ml of dichloromethane. 10.53 ml of triethylamine (75.6 mmol) were added, and, under ice cooling, 5 ml of methansulfonyl chloride (64.34 mmol). Stirring was continued for 15 minutes and then 40 ml of water were added. The organic phase was separated, and the aqueous phase extracted with dichloromethane. The combined organic layers were dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to yield 8.37 g of the product.

30 1.4 (S)-3-phenyl-1-propyl-pyrrolidine

2.0 g of methanesulfonic acid (S)-4-methanesulfonyloxy-3-phenyl-butyl ester (5.51 mmol) were dissolved in 5 ml of n-propylamine (60.82 mmol). The reaction mixture was stirred for 15 h at room temperature, diethyl ether added, the organic phase washed twice with water. The aqueous phase was reextracted once with diethylether, the organic layers combined, dried over magnesium sulfate, filtered, and the solvent evapo-

rated under reduced pressure to yield 1.09 g of the product.

ESI-

MS: 190.1 [M+H]⁺

1.5 (S)-3-(4-Nitro-phenyl)-1-propyl-pyrrolidine

5

0.3 g of (S)-3-phenyl-1-propyl-pyrrolidine (1.48 mmol) was dissolved in 2 ml of conc. sulphuric acid under argon and ice cooling. 165.16 mg of potassium nitrate (1.63 mmol) were added in small portions. The reaction mixture was stirred for 15 minutes under ice cooling, for 15 h at room temperature, and poured onto crushed ice. The aqueous solution was made alkaline with 25 % sodium hydroxide, extracted three times with diethyl ether, the aqueous phase reextracted once with diethylether, the organic layers combined, dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to yield 0.326 g of a brownish oil. A second reaction yielded another 0.919 g of the desired product.

10

15 ESI-MS: 235.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 8.15 (d, 2H), 7.45 (d, 2H), 3.4-3.5 (m, 1H), 2.9-3.0 (m, 1H), 2.75 (m, 1H), 2.3-2.6 (m, 4H), 1.8-1.9 (m, 1H), 1.5-1.65 (m, 3H), 0.95 (m, 3H).

1.6 (S)-3-(4-Amino-phenyl)-1-propyl-pyrrolidine

20

0.907 g of (S)-3-(4-nitro-phenyl)-1-propyl-pyrrolidine (3.59 mmol) were dissolved in 20 ml of methanol, 7.0 g of tin dichloride (31.02 mmol) added, and the reaction mixture stirred under reflux for 1 h. The methanol was evaporated, 60 ml of 1 N sodium hydroxide and dichloromethane added, and the phases separated after extensive stirring. The aqueous phase was extracted twice with dichloromethane, the organic layers combined, dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to yield 0.744 g of the crude amino compound.

25

ESI-MS:

205.2 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 6.9 (d, 2H), 6.45 (d, 2H), 4.7 (s, broad, 2H), 3.1 (m, 1H), 2.85 (m, 1H), 2.65 (m, 1H), 2.55 (m, 1H), 2.25-2.45 (m, 3H), 2.1 (m, 1H), 1.65 (m, 1H), 1.4-1.5 (m, 2H), 0.85 (m, 3H).

30

1.7 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

35

110

0.4 g of (S)-3-(4-amino-phenyl)-1-propyl-pyrrolidine (1.96 mmol) and 0.407 mg of 4-isopropyl-phenylsulfonyl chloride (1.86 mmol) were dissolved in 15 ml of tetrahydrofuran. 0.82 ml of triethylamine (5.87 mmol) were added and the reaction mixture stirred for 15 h at room temperature. The solvents were evaporated under reduced pressure, the residue treated with water and adjusted to an alkaline pH with sodium hydroxide solution. The aqueous layer was extracted three times with diethyl ether, the organic layers combined, dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure. The crude product was purified with silica gel chromatography with ethyl acetate/methanol (2.5-3%) as eluent, yielding 0.225 g of the purified product. This material was dissolved in 15 ml of diethyl ether and 1 ml of dichloromethane, 0.61 ml of 1 N HCl in diethyl ether added, and after formation of a precipitate, the suspension evaporated under reduced pressure to yield 0.235 g of a white precipitate.

ESI-MS: 387.2 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 11.3 and 11.1 (2 s, broad, 1H), 10.35 (m, 1H), 7.7 (d, 2H), 7.4 (d, 2H), 7.15-7.3 (m, 2H), 7.1 (m, 2H), 3.2-3.8 (several m, 4H), 2.85-3.15 (several m, 4H), 2.3 (m, 1H), 1.8-2.0 (m, 1H), 1.6-1.75 (m, 2H), 1.15 (d, 6H), 0.9 (m, 3H).

EXAMPLE 2

4-(1,1-Dimethyl-propyl)-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

0.219 g of the desired product were obtained following the same synthetic procedure as described for 4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-(1,1-dimethyl-propyl)benzenesulfonylchloride.

ESI-MS: 415.5 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 11.3 and 11.1 (2 s, broad, 1H), 10.3 (m, 1H), 7.7 (d, 2H), 7.5 (d, 2H), 7.1-7.3 (m, 2H), 7.1 (m, 2H), 3.15-3.8 (several m, 4H), 2.85-3.15 (several m, 3H), 2.3 (m, 1H), 1.8-2.0 (m, 1H), 1.5-1.75 (several m, 4H), 1.2 (s, 6H), 0.9 (m, 3H), 0.55 (m, 3H).

EXAMPLE 3

4-(Isopropyl)-N-[4-((S)-1-allyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

3.1 (S)-1-Allyl-3-phenyl-pyrrolidine

1.3 g of the desired product were obtained following the same synthetic procedure as described for (S)-3-phenyl-1-propyl-pyrrolidine, using allylamine.

ESI-MS: 188.2 [M+H]⁺

5

3.2 (S)-1-Allyl-3-(4-nitro-phenyl)-pyrrolidine

1.27 g of the desired product were obtained following the same synthetic procedure as described for (S)-3-(4-nitro-phenyl)-1-propyl-pyrrolidine.

10 ESI-MS: 233.3 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 8.15 (d, 2H), 7.4 (d, 2H), 5.85-6.0 (m, 1H), 5.2 (m, 1H), 5.1 (m, 1H), 3.4-3.5 (m, 1H), 3.05-3.2 (m, 2H), 3.0 (m, 1H), 2.75 (m, 2H), 2.6 (m, 1H), 2.3-2.4 (m, 1H), 1.8-1.9 (m, 1H).

15 3.3 (S)-1-Allyl-3-(4-amino-phenyl)-pyrrolidine

1.01 g of the desired product were obtained following the same synthetic procedure as described for (S)-3-(4-nitro-phenyl)-1-propyl-pyrrolidine.

ESI-MS: 203.1 [M+H]⁺

20

3.4 4-(Isopropyl)-N-[4-((S)-1-allyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

0.184 g of the desired product were obtained following the same synthetic procedure as described for 4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride.

25

ESI-MS: 385.1 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 11.8 and 11.5 (2 s, broad, 1H), 10.45 (m, 1H), 7.7 (d, 2H), 7.4 (d, 2H), 7.2-7.3 (m, 2H), 7.1 (m, 2H), 6.0 (m, 1H), 5.4-5.55 (m, 2H), 3.8 m (2H), 3.3-3.7 (several m, 3H), 3.2 (m, 1H), 2.9-3.1 (several m, 2H), 2.3 (m, 1H), 1.85-2.05 (m, 1H), 1.15 (s, 6H).

30

EXAMPLE 4

4-(Isopropyl)-N-[4-((R)-1-allyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

35

4.1 (R)-1-Allyl-3-(4-nitro-phenyl)-pyrrolidine

2.1 g of the desired product were obtained following the same synthetic procedure as described for (S)-3-(4-nitro-phenyl)-1-propyl-pyrrolidine.

ESI-MS: 233.1 [M+H]⁺

5 ¹H-NMR (CDCl₃): δ [ppm] 8.15 (d, 2H), 7.4 (d, 2H), 5.85-6.0 (m, 1H), 5.2 (m, 1H), 5.1 (m, 1H), 3.4-3.5 (m, 1H), 3.05-3.2 (m, 2H), 3.0 (m, 1H), 2.75 (m, 2H), 2.6 (m, 1H), 2.3-2.4 (m, 1H), 1.8-1.9 (m, 1H).

4.2 (R)-1-Allyl-3-(4-amino-phenyl)-pyrrolidine

10

1.12 g of the desired product were obtained following the same synthetic procedure as described for (S)-1-allyl-3-(4-amino-phenyl)-1-propyl-pyrrolidine.

ESI-MS: 203.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm]

15

4.3 4-(Isopropyl)-N-[4-((R)-1-allyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

20 0.138 g of the desired product were obtained following the same synthetic procedure as described for 4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride.

ESI-MS: 385.1 [M+H]⁺

25 ¹H-NMR (DMSO-d₆): δ [ppm] 11.65 and 11.45 (2 s, broad, 1H), 10.35 (m, 1H), 7.7 (d, 2H), 7.4 (d, 2H), 7.2-7.3 (m, 2H), 7.1 (m, 2H), 6.0 (m, 1H), 5.4-5.55 (m, 2H), 3.8 m (2H), 3.3-3.7 (several m, 3H), 3.2 (m, 1H), 2.9-3.1 (several m, 2H), 2.3 (m, 1H), 1.85-2.05 (m, 1H), 1.15 (s, 6H).

EXAMPLE 5

30 4-(Isopropyl)-N-[4-((R)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

0.525 g of 4-(isopropyl)-N-[4-((R)-1-allyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide (1.37 mmol) were dissolved in 15 ml ethanol, 0.075 g Pd/C (10%) added, warmed to 80°C and 0.861 g ammonium formate (13.65 mmol) dissolved in 7.5 ml of water, 35 added. After stirring for 15 h at room temperature, the catalyst was filtered off, washed with water and dichloromethane. The aqueous phase was adjusted to pH 9 with 1 N

113

aq. sodium hydroxide, extracted twice with dichloromethane, and the combined organic layers washed with saturated sodium chloride solution, dried over magnesium sulfate, filtered, and the solvent evaporated to dryness to yield 0.55 g of a mixture of the N-propyl- and N-dealkylated products. This mixture was dissolved in 25 ml of dichloromethane, 0.116 ml of propionic aldehyde (1.6 mmol), 0.14 ml acetic acid (2.39 mmol) and 0.508 g sodium triacetoxyborohydride (2.39 mmol) added. After stirring for 1 h at room temperature, the mixture was evaporated to dryness, water added and the pH adjusted to pH 9 with 1 N aq. sodium hydroxide. The aqueous layer was extracted three times with diethyl ether, the organic layers combined, dried over magnesium sulfate, filtered, and evaporated to dryness. The crude product was purified via silica gel chromatography with ethyl acetate / methanol (17.5 %). Fractions containing the product were combined, evaporated under reduced pressure, and the residue partitioned between alkaline (pH 9-10) aqueous and diethyl ether, followed by a second extraction with ethyl ether and one with ethyl acetate. The combined organic layers were dried over magnesium sulfate, filtered, and evaporated to dryness to yield 0.21 g of the purified product. The material was dissolved in 10 ml of diethyl ether, and 0.285 ml of 2 N HCl in diethyl ether added. The forming suspension was evaporated to dryness under reduced pressure to yield 0.201 g of the product as hydrochloride.

ESI-MS: 387.2 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 11.4 and 11.3 (2 s, broad, 1H), 10.35 (m, 1H), 7.7 (d, 2H), 7.4 (d, 2H), 7.15-7.3 (m, 2H), 7.1 (m, 2H), 3.2-3.8 (several m, 4H), 2.85-3.15 (several m, 4H), 2.3 (m, 1H), 1.8-2.0 (m, 1H), 1.6-1.75 (m, 2H), 1.15 (d, 6H), 0.9 (m, 3H).

25 EXAMPLE 6

4-Ethyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

0.096 g of the desired product were obtained following the same synthetic procedure as described for 4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-ethyl-benzenesulfonylchloride.

ESI-MS: 373.3 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 11.35 and 11.15 (2 s, broad, 1H), 10.3 (m, 1H), 7.7 (d, 2H), 7.4 (d, 2H), 7.15-7.3 (m, 2H), 7.05 (m, 2H), 2.9-3.8 (several m, 7H), 2.6-2.7 (m, 2H), 2.3 (m, 1H), 1.8-2.0 (m, 1H), 1.6-1.75 (m, 2H), 1.2 (m, 3H), 0.9 (m, 3H).

EXAMPLE 7

4-Trifluoromethoxy-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

- 5 0.067 g of the desired product were obtained following the same synthetic procedure as described for 4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-trifluoromethoxy-benzenesulfonylchloride.

ESI-MS: 429.1 [M+H]⁺

- 10 ¹H-NMR (DMSO-d₆): δ [ppm] 11.25 and 11.05 (2 s, broad, 1H), 10.65 (m, 1H), 8.0 (m, broad, 4H), 7.2-7.35 (m, 2H), 7.1 (m, 2H), 2.9-3.8 (several m, 7H), 2.3 (m, 1H), 1.8-2.05 (m, 1H), 1.6-1.75 (m, 2H), 0.9 (m, 3H).

EXAMPLE 8

- 15 4-Trifluoromethyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

- 0.12 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-trifluoromethyl-benzenesulfonylchloride.

ESI-MS: 413.1 [M+H]⁺

- 20 ¹H-NMR (DMSO-d₆): δ [ppm] 11.3 and 11.1 (2 s, broad, 1H), 10.55 (m, 1H), 7.9 (d, 2H), 7.55 (d, 2H), 7.2-7.35 (m, 2H), 7.1 (m, 2H), 2.9-3.8 (several m, 7H), 2.3 (m, 25 1H), 1.8-2.05 (m, 1H), 1.6-1.75 (m, 2H), 0.9 (m, 3H).

EXAMPLE 9

4-Difluoromethoxy-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

30

0.125 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-difluoromethoxy-benzenesulfonylchloride.

- 35 ESI-MS: 411.1 [M+H]⁺

115

¹H-NMR (CDCl₃): δ [ppm] 7.8 (d, 2H), 7.0-7.2 (m, 6H), 7.1 (m, 2H), 6.55 (t, 1H), 3.25 (m, 1H), 3.0 (m, 1H), 2.85 (m, 1H), 2.65 (m, 1H), 2.4-2.55 (m, 3H), 2.25 (m, 1H), 1.8 (m, 1H), 1.5 (m, 2H), 0.9 (t, 3H).

5 EXAMPLE 10

4-Methyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

0.31 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-methyl-benzenesulfonylchloride.

ESI-MS: 359.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 7.7 (m, 1H), 7.2 (m, 1H), 7.05 (m, 1H), 3.3 (m, 1H), 3.0 (m, 1H), 2.85 (m, 1H), 2.65 (m, 1H), 2.2-2.6 (several m, 4H), 2.35 (s, 3H), 1.8 (m, 1H), 1.55 (m, 2H), 0.9 (m, 3H).

EXAMPLE 11

6-Chloro-pyridine-3-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide

0.163 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 6-chloro-pyridine-sulfonylchloride.

ESI-MS: 380.1 [M+H]⁺

EXAMPLE 12

4-Methoxy-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

0.154 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-methoxy-benzene-sulfonylchloride.

ESI-MS: 375.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 7.7 (m, 1H), 7.05 (m, 1H), 6.85 (m, 1H), 3.8 (s, 3H), 3.3 (m, 1H), 3.1 (m, 1H), 2.9 (m, 1H), 2.7 (m, 1H), 2.4-2.6 (several m, 3H), 2.3 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 0.9 (m, 3H).

EXAMPLE 13

4-Chloro-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

- 5 0.175 g of the desired product were obtained following the same synthetic procedure as described for 4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-chloro-benzene-sulfonylchloride.

ESI-MS: 379.05 [M+H]⁺

- 10 ¹H-NMR (CDCl₃): δ [ppm] 7.7 (d, 2H), 7.35 (d, 2H), 7.0-7.2 (m, 4H), 7.1 (m, 2H), 3.3 (m, 1H), 3.05 (m, 1H), 2.85 (m, 1H), 2.65 (m, 1H), 2.4-2.55 (m, 3H), 2.25 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 0.9 (m, 3H).

EXAMPLE 14

- 15 2,3-Dihydro-benzofuran-5-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide

0.207 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 2,3-Dihydro-

- 20 benzofuran-5-sulfonylchloride.

ESI-MS: 387.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 7.5-7.6 (m, 2H), 7.15 (m, 2H), 6.95 (m, 2H), 6.7 (m, 1H), 4.65 (m, 2H), 3.35 (m, 1H), 3.2 (m, 2H), 3.0 (m, 1H), 2.8 (m, 1H), 2.6 (m, 1H), 2.3-2.5 (m, 3H), 2.2-2.3 (m, 1H), 1.8 (m, 1H), 1.5 (m, 2H), 0.9 (m, 3H).

25

EXAMPLE 15

4-Bromo-3-fluoro-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

- 30 0.289 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-bromo-3-fluoro-benzene-sulfonylchloride.

ESI-MS: 441.0 / 443.0 [M+H]⁺

- 35 ¹H-NMR (CDCl₃): δ [ppm] 7.65 (m, 1H), 7.5 (m, 1H), 7.4 (m, 1H), 7.15 (d, 2H), 7.0 (d, 2H), 3.3 (m, 1H), 3.0 (m, 1H), 2.8 (m, 1H), 2.65 (m, 1H), 2.35-2.5 (m, 3H), 2.3 (m, 1H), 1.8 (m, 1H), 1.5 (m, 2H), 0.9 (m, 3H).

EXAMPLE 16

4-Bromo-3-fluoro-N-[2-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

- 5 0.02 g of the ortho product could be separated during purification of the major para product.

ESI-MS: 441.0 / 443.0 [M+H]⁺

- ¹H-NMR (CDCl₃): δ [ppm] 7.5-7.7 (m, 3H), 7.1-7.2 (m, 3H), 7.0 (m, 1H), 3.5 (m, 1H), 3.3 (m, 1H), 3.1-3.2 (m, 2H), 2.7-2.9 (m, 3H), 2.3-2.4 (m, 1H), 1.9-2.1 (m, 2H), 1.6-1.8 (m, 2H), 0.9 (m, 3H).
- 10

EXAMPLE 17

4-Bromo-2-fluoro-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

- 15 0.387 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-bromo-2-fluoro-benzene-sulfonylchloride.

ESI-MS: 441.0 / 443.0 [M+H]⁺

- 20 ¹H-NMR (CDCl₃): δ [ppm] 7.65 (m, 1H), 7.3-7.4 (m, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 3.3 (m, 1H), 3.0 (m, 1H), 2.8 (m, 1H), 2.65 (m, 1H), 2.35-2.55 (m, 3H), 2.25 (m, 1H), 1.75 (m, 1H), 1.5 (m, 2H), 0.9 (m, 3H).

EXAMPLE 18

- 25 4-Isopropyl-N-[4-((S)-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

- 0.515 g of 4-isopropyl-N-[4-((S)-1-allyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide (1.34 mmol) were dissolved in 10 ml of tetrahydrofuran and added to a solution of 0.049 g tris-(dibenzylidenacetone)-dipalladium(0) (0.05 mmol) and 0.023 g of 1,4-bis-(diphenylphosphino)-butane (0.05 mmol) in 3 ml of tetrahydrofuran under argon, followed by addition of 0.227 mg of 2-mercapto benzoic acid (1.47 mmol) in 3 ml of tetrahydrofuran. The mixture was stirred for 2 h at room temperature, the solvent evaporated under reduced pressure and water containing 1 N hydrochloric acid added to adjust the pH to acidic values. The aqueous phase was extracted three times with ethyl acetate, then adjusted to alkaline pH with 1 N sodium hydroxide, extracted twice with diethyl ether and twice with dichloromethane. The combined organic layers were dried
- 30
- 35

over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to yield 0.215 g of the secondary amine.

ESI-MS: 345.1 [M+H]⁺

5 EXAMPLE 19

2-Fluoro-4-isopropenyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

0.2 g of 2-fluoro-4-bromo-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-
benzenesulfonamide (0.45 mmol), 1.05 g of tributyl-isopropenyl-stannane (3.17 mmol)
10 and 0.026 g of tetrakis(triphenyl)phosphinepalladium(0) (0.02 mmol) were dissolved in 3
ml of tetrahydrofuran and stirred for 40 minutes at 150°C in the microwave (CEM). The
reaction mixture was filtered over celite, washed with methanol, and the filtrate evapo-
rated to dryness under reduced pressure. The residue was purified via silica gel chro-
matography with cyclohexane/ethyl acetate (20%), ethyl acetate, and ethyl ace-
15 tate/methanol (15%). Fractions containing the product were combined, the solvent
evaporated to yield 0.176 g of the title product.

ESI-MS: 403.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 7.7 (m, 1H), 7.2-7.3 (m, 2H), 7.1 (m, 2H), 7.0 (m, 2H),
5.45 (m, 1H), 5.25 (m, 1H), 3.3-3.4 (m, 1H), 2.6 (m, 1H), 2.25-2.35 (m, 1H), 2.1 (s,
20 3H), 1.8-1.9 (m, 2H), 1.55-1.7 (m, 3H), 1.2-1.4 (m, 3H), 0.9 (m, 3H).

EXAMPLE 20

3-Fluoro-4-isopropenyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

25 3-Fluoro-4-isopropenyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide
were obtained following the same synthetic procedure as described for 2-fluoro-4-
isopropenyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide starting
from 3-fluoro-4-bromo-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide
yielding 0.17 g of the desired.

30 ESI-MS: 403.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 7.3-7.5 (m, 3H), 7.15 (d, 2H), 7.0 (d, 2H), 5.3 (m, 2H),
3.35 (m, 1H), 2.6 (m, 2H), 2.3 (m, 1H), 2.1 (s, 3H), 1.8-1.9 (m, 2H), 1.55-1.7 (m, 3H),
1.2-1.4 (m, 1H), 0.9 (m, 4H).

35 EXAMPLE 21

2-Fluoro-4-isopropenyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

0.161 g of 2-fluoro-4-isopropenyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide (0.37 mmol) were dissolved in 20 ml methanol, a tip of a spatula 10% Pd/C added, and the reaction mixture hydrogenated for 3 h at 50°C. The reaction mixture was filtered over celite, washed with methanol, and the filtrate evaporated to dryness under reduced pressure. The residue was purified via silica gel chromatography on a chromabond column with ethyl acetate and ethyl acetate/methanol (10%). Fractions containing the product were combined, the solvent evaporated to yield 0.11 g of the title product.

ESI-MS: 405.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 7.7 (m, 1H), 6.9-7.2 (several m, 6H), 3.5-3.6 (m, 2H), 3.2-3.4 (m, 2H), 2.85-3.05 (m, 4H), 2.4 (m, 1H), 2.1 (m, 1H), 1.8 (m, 2H), 1.2 (d, 6H), 0.95 (m, 3H).

EXAMPLE 22

3-Fluoro-4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

0.122 g of 3-fluoro-4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide were obtained following the same synthetic procedure as described for 2-fluoro-4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide starting from 3-fluoro-4-isopropenyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide.

ESI-MS: 405.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 7.5 (m, 1H), 7.4 (m, 1H), 7.3 (m, 1H), 7.1 (d, 2H), 7.0 (d, 2H), 3.4 (m, 1H), 3.2 (m, 2H), 3.05 (m, 1H), 2.9 (m, 1H), 2.6-2.7 (m, 3H), 2.1 (m, 1H), 1.9 (m, 1H), 1.6-1.7 (m, 2H), 1.2 (d, 6H), 0.9 (m, 3H).

EXAMPLE 23

4-Isopropoxy-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

30

0.083 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-isopropoxybenzene-sulfonylchloride.

ESI-MS: 403.3 [M+H]⁺

35

120

¹H-NMR (DMSO-d₆): δ [ppm] 11.3 and 11.1 (2 s, broad, 1H), 10.2 (m, 1H), 7.7 (d, 2H), 7.25 (m, 1H), 7.2 (m, 1H), 6.95-7.1 (m, 4H), 4.7 (m, 1H), 2.9-3.8 (several m, 7H), 2.3 (m, 1H), 1.85-2.0 (m, 1H), 1.7 (m, 2H), 1.2 (d, 6H), 0.9 (m, 3H).

5 EXAMPLE 24

Indan-5-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide hydrochloride

0.15 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-

10 benzenesulfonamide hydrochloride using commercially available indan-5-sulfonylchloride.

ESI-MS: 385.1 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 11.3 and 11.1 (2 s, broad, 1H), 10.25 (m, 1H), 7.6 (s, 1H), 7.5 (d, 1H), 7.35 (d, 1H), 7.25 (m, 1H), 7.2 (m, 1H), 7.05 (m, 2H), 3.2-3.8 (several m, 4H), 2.8-3.1 (several m, 7H), 2.3 (m, 1H), 1.8-2.05 (m, 3H), 1.6-1.7 (m, 2H), 0.9 (m, 3H).

EXAMPLE 25

4-Bromo-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

20

0.07 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-

benzenesulfonamide hydrochloride using commercially available 4-bromo-benzene-sulfonylchloride.

25 ESI-MS: 425.0 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 11.3 and 11.05 (2 s, broad, 1H), 10.45 (m, 1H), 7.8 (d, 2H), 7.7 (d, 2H), 7.3 (m, 1H), 7.2 (m, 1H), 7.05 (m, 2H), 2.9-3.8 (several m, 7H), 2.3 (m, 1H), 1.85-2.05 (m, 1H), 1.6-1.8 (m, 2H), 0.9 (m, 3H).

30 EXAMPLE 26

4-Acetyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

0.159 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-

35 benzenesulfonamide using commercially available 4-acetyl-benzene-sulfonylchloride.

ESI-MS: 387.1 [M+H]⁺

121

¹H-NMR (CDCl₃): δ [ppm] 8.0 (d, 2H), 7.8 (d, 2H), 7.15 (d, 2H), 6.95 (d, 2H), 3.3 (m, 1H), 3.0 (m, 1H), 2.8 (m, 1H), 2.65 (m, 1H), 2.6 (s, 3H), 2.35-2.5 (m, 3H), 2.25 (m, 1H), 1.8 (m, 1H), 1.5 (m, 2H), 0.9 (m, 3H).

5 EXAMPLE 27

4-Cyclopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

0.3 g of 4-bromo-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide (0.71 mmol), 0.079 g of cyclopropylboronic acid (0.92 mmol), 0.526 g of potassium phosphate (2.48 mmol) and 0.02 g of tricyclohexylphosphine were dissolved in a mixture of 4 ml of toluene and 2 ml of water. After addition of 0.008 g of palladium(II)-acetate (0.04 mmol), the reaction mixture was stirred for 1 h at 100°C in the microwave (CEM). The solution was decanted, the remaining suspension containing the catalyst washed again with ethyl acetate, the combined solvent extracts evaporated to dryness, redissolved in ethyl acetate and washed with water. The aqueous phase was reextracted once with ethyl acetate, the organic layers combined, dried over magnesium sulfate, filtered and evaporated under reduced pressure. The crude product was purified via silica gel chromatography with dichloromethane, dichloromethane-methanol (5.5 %). Fractions containing the product were combined, the solvent removed, to yield 0.066 g of the title product.

ESI-MS: 385.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 7.65 (d, 2H), 6.95-7.15 (m, 6H), 3.3 (m, 1H), 3.1 (m, 1H), 2.9 (m, 1H), 2.75 (m, 1H), 2.5 (m, 3H), 2.35 (m, 1H), 2.0 (m, 1H), 1.8 (m, 2H), 1.55 (m, 2H), 1.25 (m, 1H), 1.0 (m, 1H), 0.9 (m, 3H), 0.7 (m, 1H).

25 EXAMPLE 28

5-Bromo-thiophene-2-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide hydrochloride

0.05 g of the title product were obtained following the same synthetic procedure as described for 4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 5-bromo-thiophene-2-sulfonylchloride.

ESI-MS: 431.0 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 11.2 and 11.0 (2 s, broad, 1H), 10.65 (m, 1H), 7.2-7.4 (several m, 4H), 7.1 (m, 2H), 3.0-3.8 (several m, 7H), 2.3 (m, 1H), 1.85-2.0 (m, 1H), 1.7 (m, 2H), 0.9 (m, 3H).

EXAMPLE 29

5-Isopropenyl-thiophene-2-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide

- 5 0.154 g of the title product were obtained following the same synthetic procedure as described for 2-fluoro-4-Isopropenyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide starting from 5-bromo-thiophene-2-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide.

ESI-MS: 391.1 [M+H]⁺

- 10 ¹H-NMR (CDCl₃): δ [ppm] 7.3 (m, 1H), 7.25 (m, 1H), 7.2 (d, 2H), 7.05 (d, 2H), 6.9 (m, 1H), 5.4 (s, 1H), 5.1 (s, 1H), 3.3 (m, 1H), 3.05 (m, 1H), 2.9 (m, 1H), 2.7 (m, 1H), 2.5 (m, 2H), 2.3 (m, 1H), 2.05 (s, 3H), 1.85 (m, 1H), 1.6 (m, 2H), 1.3 (m, 1H), 0.9 (m, 3H).

EXAMPLE 30

- 15 4-Propyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

0.037 g of the title product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-propyl-benzene-sulfonylchloride.

ESI-MS: 387.2 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 10.3 (m, 1H), 7.7 (d, 2H), 7.3 (d, 2H), 7.25 (m, 1H), 7.2 (m, 1H), 7.05 (m, 2H), 2.9-3.8 (several m, 7H), 2.55 (m, 2H), 2.3 (m, 1H), 1.85-2.0 (m, 1H), 1.7 (m, 2H), 1.55 (m, 2H), 0.9 (m, 3H), 0.8 (m, 3H).

25

EXAMPLE 31

N-[6-(1-propyl-pyrrolidin-3-yl)-pyridin-3-yl]-4-isopropyl-benzene-sulfonamide

31.1 5-Nitro-2-vinyl-pyridine

30

- 0.5 g of 2-chloro-5-nitro-pyridine (3.15 mmol), 1.2 g of tributyl-vinyl-stannane (3.78 mmol), 0.036 g of tetrakis(triphenylphosphine)palladium(0) (0.03 mmol), and 0.024 g of triphenylphosphine (0.09 mmol) were dissolved in 20 ml of toluene and stirred for 2 h under reflux. Upon cooling to room temperature, 10 ml of water were added, the aqueous layer was extracted with ethyl acetate, and the combined organic layers were washed with saturated sodium chloride, dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure. The crude product was purified via silica
- 35

123

gel chromatography with cyclohexane/ethyl acetate (10%). Fractions containing the product were combined, the solvent evaporated to yield 0.528 g of the desired product.

ESI-MS: 151.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 9.4 (s, 1H), 8.4 (m, 1H), 7.45 (m, 1H), 6.9 (q, 1H), 6.45 (m, 1H), 5.7 (m, 1H).

31.2 2-(1-Benzyl-pyrrolidin-3-yl)-5-nitro-pyridine

0.15 g of 5-nitro-2-vinyl-pyridine were dissolved in 2.5 ml of dichloromethane, 0.149 g of trifluoroacetic acid (1.31 mmol) were added, followed by slow addition of 0.928 g of N-benzyl-N-(methoxymethyl)-N-trimethylsilylmethylamine (3.91 mmol). Stirring was continued at room temperature for 1 h, before the reaction mixture was washed with aqueous sodium hydrogencarbonate, the aqueous layer reextracted with dichloromethane, the organic layers combined, dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure. The crude product was purified via silica gel chromatography with ethyl acetate. Fractions containing the product were combined, the solvent evaporated to yield 0.186 g of the desired product.

ESI-MS: 284.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 9.25 (m, 1H), 8.3 (m, 1H), 7.4 (m, 1H), 7.15-7.3 (m, 5H), 3.6 (d, 2H), 3.55 (m, 1H), 2.9 (m, 1H), 2.7 (m, 2H), 2.65 (m, 1H), 2.3 (m, 1H), 2.0 (m, 1H).

31.3 2-(1-Benzyl-pyrrolidin-3-yl)-5-amino-pyridine

0.181 g of 2-(1-benzyl-pyrrolidin-3-yl)-5-nitro-pyridine were dissolved in 10 ml of methanol and 1.15 g of tin dichloride SnCl₂ (5.11 mmol) were added in portions. Stirring was continued for 1 h under refluxing conditions. The solvent was evaporated, the residue treated with 1 N aqueous sodium hydroxide and ethyl acetate, and filtered. The two phases were separated, the aqueous phase was extracted twice with ethyl acetate, and the combined organic layers were dried over magnesium sulfate, filtered, and the solvent was evaporated under reduced pressure to yield 0.191 g of the crude amino product which was used in the next reaction without further purification.

ESI-MS: 254.1 [M+H]⁺

31.4 N-[6-(1-Benzyl-pyrrolidin-3-yl)-pyridin-3-yl]-4-isopropyl-benzene-sulfonamide

124

0.12 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-propyl-benzene-sulfonylchloride. The crude product was thereby purified via silica gel chromatography on a cromabond column with ethyl acetate as eluent.

ESI-MS: 436.1 [M+H]⁺

31.5 N-[6-(Pyrrolidin-3-yl)-pyridin-3-yl]-4-isopropyl-benzene-sulfonamide

0.12 g of N-[6-(1-benzyl-pyrrolidin-3-yl)-pyridin-3-yl]-4-isopropyl-benzene-sulfonamide (0.28 mmol) were dissolved in 20 ml of methanol, a tip of a spatula 10% Pd/C was added, and the reaction mixture was hydrogenated for 2 h at room temperature and 4 h at 50°C. The catalyst was removed by filtration and the filtrate evaporated to dryness to yield 0.088 g of the desired debenzylated compound.

ESI-MS: 346.1 [M+H]⁺

31.6 N-[6-(1-propyl-pyrrolidin-3-yl)-pyridin-3-yl]-4-isopropyl-benzene-sulfonamide

0.088 g of N-[6-(1-benzyl-pyrrolidin-3-yl)-pyridin-3-yl]-4-isopropyl-benzene-sulfonamide (0.25 mmol) were dissolved in 10 ml of dichloromethane. 0.022 g of propionyl aldehyde (0.38 mmol), 0.023 g of acetic acid (0.38 mmol) and 0.081 g of sodium triacetoxyborohydride (0.38 mmol) were added and stirred for 1 h at room temperature. Water was added, the pH adjusted to 10 with 1 N sodium hydroxide, extracted three times with ethyl acetate, the organic layers combined and evaporated to dryness. The crude material was purified via silica gel chromatography with dichloromethane/methanol 20%. Fractions containing the product were combined and the solvent evaporated to yield 0.026 mg of the product.

ESI-MS: 388.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 8.15 (m, 1H), 7.7 (d, 2H), 7.5 (m, 1H), 7.3 (d, 2H), 7.1 (m, 1H), 3.5 (m, 1H), 3.2 (m, 1H), 2.95 (m, 2H), 2.75 (m, 2H), 2.5-2.65 (m, 2H), 2.3 (m, 1H), 2.0 (m, 1H), 1.6 (m, 2H), 1.2 (m, 6H), 0.9 (m, 3H).

EXAMPLE 32

4-Dimethylaminomethyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

32.1 4-Formyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

125

0.584 g of the desired product were obtained following the same synthetic procedure as described for 4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide using commercially available 4-formyl-benzene-sulfonylchloride. The product was used in the subsequent step without further purification.

5 ESI-MS: 373.4 [M+H]⁺

32.2 4-Dimethylaminomethyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

10 0.3 g of 4-formyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide (0.81 mmol) were dissolved in 20 ml of dichloromethane, 0.6 ml of a 2 M solution of dimethylamine in tetrahydrofuran (1.21 mmol) added, followed by addition of 0.07 ml of acetic acid and 0.256 g sodium triacetoxyborohydride. The reaction mixture was stirred for 15 h at room temperature, water added, the pH adjusted to alkaline conditions with
15 1 N aqueous sodium hydroxide, and the aqueous layer extracted twice with diethylether. The combined ether extracts were dried over magnesium sulfate, filtered and the solvent evaporated under reduced pressure. The crude product was purified via preparative HPLC on a 40 mm Deltapak column with methanol/water/0.1% acetic acid. Fractions containing the product were combined, the solvent evaporated and the
20 product isolated after extraction of an alkaline aqueous layer. (0.051 g).
ESI-MS: 402.1 [M+H]⁺

EXAMPLE 33

4-sec-Butyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride
25 ride

0.082 g of the desired product were obtained following the same synthetic procedure as described for 4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-sec-butyl-benzenesulfonylchloride (ART-Chem).
30 ESI-MS: 401.1 [M+H]⁺

EXAMPLE 34

4-Bromo-3,6-difluoro-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide
35
0.131 g of the desired product were obtained following the same synthetic procedure as described for 4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-

126

benzenesulfonamide using commercially available 4-bromo-3,6-difluoro-benzenesulfonylchloride.

ESI-MS: 459.0 / 461.0 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 7.55 (m, 1H), 7.4 (m, 1H), 7.15 (d, 2H), 7.0 (d, 2H),
5 4.7 (s, very broad, 2H), 3.3 (m, 1H), 3.0 (m, 1H), 2.85 (m, 1H), 2.7 (m, 1H), 2.4-2.6 (m, 3H), 2.25 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 0.85 (m, 3H).

EXAMPLE 35

4-Isobutyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

10

0.164 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride using commercially available 4-isobutyl-benzenesulfonylchloride.

15 ESI-MS: 401.1 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 11.3 and 11.1 (2 s, broad, 1H), 10.3 (m, 1H), 7.7 (d, 2H), 7.3 (d, 2H), 7.25 (m, 1H), 7.2 (m, 1H), 7.05 (m, 2H), 2.9-3.8 (several m, 9H), 2.3 (m, 1H), 1.8-2.0 (m, 2H), 1.7 (m, 2H), 0.9 (m, 3H), 0.8 (m, 6H).

20 EXAMPLE 36

4-Carboxy-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

0.105 g of the desired product were obtained following the same synthetic procedure as described for 4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-
25 benzenesulfonamide using commercially available 4-carboxy-benzenesulfonylchloride. The product however was finally isolated as precipitate from the aqueous layer through filtration, stirring with methanol, again filtration, and drying.

ESI-MS: 389.1 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 10.25 (s, 1H), 9.6 (s, broad, 1H), 7.9 (d, 2H), 7.7-7.8
30 (m, 4H), 7.3 (d, 2H), 3.0-3.9 (several broad m, 7H), 2.4 (m, 1H), 2.0 (m, 1H), 1.7 (m, 2H), 0.9 (m, 3H).

EXAMPLE 37

35 4-Cyclopentyl-N-[4-(S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

0.278 g of 4-bromo-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide (0.66 mmol) were dissolved in 5 ml tetrahydrofuran. 0.027 g of 1,1'-bis(diphenylphosphino)ferrocene (dppf)PdCl₂ (0.03 mmol) and 0.008 g of CuI (0.04 mmol) were added, followed by dropwise addition of 2 ml commercially available 0.5 M cyclopentylzincbromide in tetrahydrofuran. After stirring for 15 h at room temperature and further addition of cyclopentylzincbromide, the reaction mixture was treated with ethyl acetate, washed with water, and the organic layer filtered over celite, washed with ethyl acetate, washed with water, and the organic layer dried over magnesium sulfate, filtered and the solvent removed under reduced pressure. The crude product was purified via preparative HPLC (Delta Pak column, 40 mm diameter) using methanol/water/0.1 % acetic acid as eluent. Fractions containing the product were combined, methanol removed, and, after addition of 0.2 ml of 1 N hydrochloric acid, the aqueous layer lyophilised to yield 0.0107 g of the product.

ESI-MS: 413.1 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 12.4 & 12.3 (2 s, broad, 1H), 7.7-7.9 (m, 3H), 7.25 (m, 2H), 7.0-7.2 (m, 3H), 4.0 (m, broad, 2H), 3.75 (m, broad, 1H), 2.5-3.0 (several m broad, 6H), 2.35 (m, broad, 1H), 1.5-2.1 (several m, broad, 10H), 1.0 (m, broad, 3H).

EXAMPLE 38

4-Isopropyl-N-[4-((S)-1-(3-fluoro-propyl)-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

0.215 g of 4-isopropyl-N-[4-((S)-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide (0.62 mmol) and 0.16 g of toluene-4-sulfonic acid 3-fluoro-propyl ester (0.69 mmol) were dissolved in 5 ml of dimethylformamide, 0.43 ml of triethylamine added and the reaction stirred for 1 h at 50°C. Additional toluene-4-sulfonic acid 3-fluoro-propyl ester and triethylamine were added to drive the reaction to completion. After stirring at room temperature for 15 h, the solvent was evaporated, the residue re-dissolved in dichloromethane, and the organic layer washed with 1 N aqueous sodium hydroxide. The aqueous phase was extracted twice with dichloromethane, the organic layers combined, dried over magnesium sulfate, filtered and the solvent evaporated. The crude product was purified via silica gel chromatography (chromabond column) using dichloromethane/methanol 0-3%. Fractions containing the product were collected and the solvent evaporated to yield 0.115 g of the desired product.

ESI-MS: 405.1 [M+H]⁺

128

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.7 (d, 2H), 7.3 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 4.6 (m, 1H), 4.45 (m, 1H), 3.3 (m, 1H), 2.95 (m, 2H), 2.8 (m, 1H), 2.5-2.7 (several m, 3H), 2.45 (m, 1H), 2.25 (m, 1H), 1.75-2.0 (several m, 3H), 1.2 (m, 6H).

EXAMPLE 39

5 3-Trifluoromethyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

0.11 g of the desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide using commercially available 3-trifluoromethyl-benzenesulfonylchloride.

ESI-MS: 427.2 [M+H]⁺

EXAMPLE 40

15 4-Butyloxy-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

The desired product was obtained following the same synthetic procedure as described for 4-isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide using commercially available 4-butyloxy-benzenesulfonylchloride.

ESI-MS: 417.3 [M+H]⁺

20 ¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.7 (d, 2H), 7.15 (d, 2H), 6.95 (d, 2H), 6.85 (d, 2H), 5.6 (s, broad, 1H), 3.95 (t, 2H), 3.3 (m, 1H), 3.0 (m, 1H), 2.8 (m, 1H), 2.6 (m, 1H), 2.45 (m, 1H), 2.35 (m, 2H), 2.25 (m, 1H), 1.75 (m, 3H), 1.5 (m, 4H), 0.96 (m, 6H).

EXAMPLE 41

25 4-(2,2-Difluorocyclopropyl)-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

The desired product were obtained following the same synthetic procedure as described for 4-Isopropyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide using 4-(2,2-difluoro-cyclopropyl)-benzenesulfonylchloride.

ESI-MS: 421.15 [M+H]⁺

30 ¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.75 (d, 2H), 7.25 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 5.65 (bs, 1H), 3.3 (m, 1H), 3.05 (m, 1H), 2.85 (m, 1H), 2.75 (m, 1H), 2.65 (m, 1H), 2.45 (m, 3H), 2.3 (m, 1H), 1.9 (m, 1H), 1.8 (m, 1H), 1.65 (m, 1H), 1.55 (m, 2H), 0.95 (t, 3H).

EXAMPLE 42

35 N-[3-(1-Propyl-pyrrolidin-3-yl)-phenyl]-4-trifluoromethoxy-benzenesulfonamide x HCl

42.1 3-[3-(4-Trifluoromethoxy-benzenesulfonylamino)-phenyl]-pyrrolidine-1-carboxylic acid methyl ester

- 5 To a solution of 3-(3-amino-phenyl)-pyrrolidine-1-carboxylic acid methyl ester (500 mg, 2.27 mmol) and triethylamine (500 mg, 4.94 mmol) in THF (20 ml) at 10°C 4-trifluoromethoxy-benzenesulfonylchloride (600 mg, 2.3 mmol) was added. The mixture was allowed to come to room temperature and was stirred for 16 h. The mixture was poured into water and extracted three times with ethyl acetate. The organic layer was
10 washed with water and saturated sodium chloride solution, dried over MgSO₄ and evaporated under reduced pressure to give the product as a brown oil (1 g, 99%).

42.2 N-(3-Pyrrolidin-3-yl-phenyl)-4-trifluoromethoxy-benzenesulfonamide

- 15 3-[3-(4-Trifluoromethoxy-benzenesulfonylamino)-phenyl]-pyrrolidine-1-carboxylic acid methyl ester (500 mg, 1.13 mmol) and HCl (8M, 137.82 mmol) in EtOH (10 ml) were heated under reflux for 48 h. The mixture was extracted twice with ethyl acetate and then NaOH (2M) was added. The aqueous phase was extracted twice with ethyl acetate. The combined organic layers were washed with water and saturated sodium chloride solution, dried over MgSO₄ and evaporated under reduced pressure to give the
20 product as a brown oil (200 mg, 46%).

42.3 N-[3-(1-Propyl-pyrrolidin-3-yl)-phenyl]-4-trifluoromethoxy-benzenesulfonamide x HCl

- 25 A solution of N-(3-pyrrolidin-3-yl-phenyl)-4-trifluoromethoxy-benzenesulfonamide (200 mg, 0.52 mmol), acetic acid (30 µl, 0.52 mmol) and propionic aldehyde (30 mg, 0.52 mmol) were stirred in dichloromethane (20 ml) for 30 min at room temperature before sodium triacetoxyborohydride (165 mg, 0.78 mmol) was added in portions at 15°C. The mixture was stirred for 16 h at room temperature. The mixture was poured into water/dichloromethane (1/1) and the organic layer was washed with water, dried over MgSO₄ and evaporated under reduced pressure. The residue was purified by column chromatography (CH₂Cl₂/methanol – 2%, 3%, 4%, 6%) to give a brown oil. To a solution of this oil in diethylether HCl in diethylether (1M) was added and the mixture was
30 evaporated under reduced pressure to give the product as a white foam, which was dried in vacuo (55 mg, 23%).

MS (ESI) m/z: 429.15 [M+H]⁺

130

¹H-NMR (DMSO-d₆): δ [ppm] 11.0-11.2 (m, 1H), 10.6 (s, 1H), 7.90-7.95 (m, 2H), 7.60-7.65 (m, 2H), 7.20-7.25 (m, 1H), 6.95-7.10 (m, 3H), 3.35-3.80 (m, 3H), 3.20-3.30 (m, 1H), 3.05-3.15 (m, 2H), 2.90-3.05 (m, 1H), 2.25-2.35 (m, 1H), 1.85-2.05 (m, 1H), 1.65-1.80 (m, 2H), 0.90-1.00 (m, 3H).

5

EXAMPLE 43

4-Isopropyl-N-[3-(1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide x HCl

10

43.1 3-[3-(4-Isopropyl-benzenesulfonylamino)-phenyl]-pyrrolidine-1-carboxylic acid methyl ester

15

To a solution of 3-(3-amino-phenyl)-pyrrolidine-1-carboxylic acid methyl ester (500 mg, 2.27 mmol) and triethylamine (500 mg, 4.94 mmol) in THF (20 ml) at 10°C 4-isopropyl-benzenesulfonylchloride (500 mg, 2.27 mmol) was added. The mixture was allowed to come to room temperature and was stirred for 16 h. The mixture was poured into water and extracted three times with ethyl acetate. The organic layer was washed with water and saturated sodium chloride solution, dried over MgSO₄ and evaporated under reduced pressure to give the product as a brown oil (1 g, 100%).

20

43.2 4-Isopropyl-N-(3-pyrrolidin-3-yl-phenyl)-benzenesulfonamide

25

3-[3-(4-Isopropyl-benzenesulfonylamino)-phenyl]-pyrrolidine-1-carboxylic acid methyl ester (900 mg, 1.13 mmol) and HCl (8M, 273.90 mmol) in EtOH (20 ml) were heated to reflux for 48 h. The mixture was extracted twice with ethyl acetate and then NaOH (2M) was added. The aqueous phase was extracted twice with ethyl acetate. The combined organic layers were washed with water and saturated sodium chloride solution, dried over MgSO₄ and evaporated under reduced pressure to give the product as a brown oil (100 mg, 13%).

30

43.3 4-Isopropyl-N-[3-(1-Propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide x HCl

35

A solution of 4-isopropyl-N-(3-pyrrolidin-3-yl-phenyl)-benzenesulfonamide (100 mg, 0.29 mmol), acetic acid (20 µl, 0.29 mmol) and propionic aldehyde (16.86 mg, 0.29 mmol) were stirred in dichloromethane (20 ml) for 30 min at room temperature before sodium triacetoxyborohydride (123 mg, 0.58 mmol) was added in portions at room temperature. The mixture was stirred for 16 h at room temperature. The mixture was poured into saturated aqueous NaHCO₃/dichloromethane (1/1) and the organic layer

131

was washed with water and saturated sodium chloride solution, dried over MgSO_4 and evaporated under reduced pressure. The residue was purified by column chromatography (CH_2Cl_2 /methanol – 2%, 4%, 6%). To a solution of the purified product in diethylether HCl in diethylether (1M) was added and the mixture was evaporated under reduced pressure to give the product as a white foam, which was dried in vacuo (55 mg, 45%).

MS (ESI) m/z: 387.15 $[\text{M}+\text{H}]^+$

$^1\text{H-NMR}$ (CDCl_3): δ [ppm] 8.95-9.15- (m, 1H), 7.72-7.82 (m, 2H), 6.85-7.31 (m, 6H), 3.85-4.05 (m, 1H), 3.65-3.79 (m, 1H), 3.25-3.56 (m, 1H), 3.0-3.2 (m, 2H), 2.78-3.0 (m, 2H), 2.43-2.55 (m, 1H), 1.80-2.40 (m, 5H), 1.12-1.21 (m, 6H), 0.9-1.02 (m, 3H).

EXAMPLE 44

N-[3-(1-Cyclopropylmethyl-pyrrolidin-3-yl)-phenyl]-4-isopropyl-benzenesulfonamide x HCl

A solution of 4-isopropyl-N-(3-pyrrolidin-3-yl-phenyl)-benzenesulfonamide (100 mg, 0.29 mmol), acetic acid (20 μl , 0.29 mmol) and cyclopropanecarbaldehyde (20.35 mg, 0.29 mmol) were stirred in dichloromethane (20 ml) for 30 min at room temperature before sodium triacetoxyborohydride (123 mg, 0.58 mmol) was added in portions at room temperature. The mixture was stirred for 16 h at room temperature. The mixture was poured into saturated aqueous NaHCO_3 /dichloromethane (1/1) and the organic layer was washed with water and saturated sodium chloride solution, dried over MgSO_4 and evaporated under reduced pressure. The residue was purified by column chromatography (CH_2Cl_2 /methanol – 2%, 4%, 6%). To a solution of the purified product in diethylether HCl in diethylether (1M) was added and the mixture was evaporated under reduced pressure to give the product as a brown foam, which was dried in vacuo (50 mg, 40%).

MS (ESI) m/z: 399.25 $[\text{M}+\text{H}]^+$

$^1\text{H-NMR}$ (CDCl_3): δ [ppm] 9.05-9.35- (m, 1H), 7.72-7.81 (m, 2H), 6.82-7.35 (m, 6H), 3.69-4.05 (m, 2H), 2.70-3.25 (m, 5H), 2.0-2.55 (m, 3H), 1.1-1.3 (m, 7H), 0.6-0.8 (m, 2H), 0.47-0.5 (m, 2H).

EXAMPLE 45

N-[3-(1-Allyl-pyrrolidin-3-yl)-phenyl]-4-isopropyl-benzenesulfonamide x HCl

132

A solution of 4-isopropyl-N-(3-pyrrolidin-3-yl-phenyl)-benzenesulfonamide (100 mg, 0.29 mmol), allylbromide (38 mg, 0.31 mmol), potassium carbonate (60 mg, 0.43 mmol), and potassium fluoride (0.2 mg, 0.003 mmol) in acetone (20 ml) were heated to reflux for 4 h. The mixture was evaporated under reduced pressure and the residue was partitioned between ethyl acetate and water. The organic layer was washed with saturated sodium chloride solution, dried over MgSO_4 and evaporated under reduced pressure. The residue was purified by column chromatography (CH_2Cl_2 /methanol – 2%, 4%, 6%). To a solution of the purified product in dichloromethane HCl in diethylether (1M) was added and the mixture was evaporated under reduced pressure to give the product as a brown foam, which was dried in vacuo (25 mg, 21%).

MS (ESI) m/z : 385.15 $[\text{M}+\text{H}]^+$

^1H -NMR (CDCl_3): δ [ppm] 8.79-8.95- (m, 1H), 7.72-7.81 (m, 2H), 6.85-7.30 (m, 6H), 6.05-6.20 (m, 1H), 5.43-5.56 (m, 2H), 3.10-3.95 (m, 6H), 2.80-2.95 (m, 1H), 1.90-2.55 (m, 3H), 1.15-1.25 (m, 6H).

EXAMPLE 46

4-Isopropyl-N-[6-(1-propyl-azetidin-3-yl)-pyridin-3-yl]-benzenesulfonamide

46.1 3-Hydroxy-azetidine-1-carboxylic acid tert-butyl ester

To a degassed solution of 1-benzhydryl-azetidin-3-ol (4.75 g, 19.84 mmol) in methanol (methanol) (150 ml) were added ammonium formate (8.76 g, 138.91 mmol), 10% Pd/C (450 mg) and Boc_2O (di-tert-butyl dicarbonate) (13 g, 59.56 mmol). The resulting suspension was heated to reflux under N_2 for 1h. It was then cooled down to room temperature, filtered through a short pad of celite and concentrated. The residue was dissolved in CH_2Cl_2 and washed with water. The organic layer was dried (Na_2SO_4) and evaporated. The raw substance was chromatographed on silica gel (heptane:ethyl acetate (ethyl acetate), 1:1) to afford the title compound (3.30 g, 96%) as white crystals.

MS (ESI+) m/z = 118.1 $[\text{M}-t\text{Bu}+\text{H}]^+$

^1H NMR (400 MHz, CDCl_3) : δ (ppm) 1.43 (s, 9H), 2.35 (d, J = 6.2 Hz, 1H), 3.80 (dd, J = 10.4, 4.4 Hz, 2H), 4.15 (dd, J = 9.6, 6.7 Hz, 2H), 4.58 (m, 1H).

46.2 3-Iodo-azetidine-1-carboxylic acid tert-butyl ester

133

A solution of 3-hydroxy-azetidine-1-carboxylic acid tert-butyl ester (3.35 g, 19.34 mmol) in toluene (200 ml) was treated with imidazole (3.95 g, 58.01 mmol), triphenylphosphine (10.14 g, 38.65 mmol) and I₂ (7.36 g, 28.99 mmol). The mixture was heated at 100°C for 1 h, cooled down to room temperature and next poured into saturated aqueous NaHCO₃ (30 ml). Excess triphenylphosphine was destroyed by addition of iodine until I₂ coloration persisted in organic layer. The latter was washed with 5% aqueous Na₂S₂O₃, dried over Na₂SO₄ and evaporated. Purification of the crude product by flash column chromatography (heptane:ethyl acetate, 2:1) provides the title compound (5.19 g, 95%) as a light yellow oil.

MS (ESI+) m/z = 227.9 [M-*t*Bu+H]⁺

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 1.44 (s, 9H), 4.29 (dd, *J* = 10.4, 5.4 Hz, 2H), 4.47 (m, 1H), 4.64 (dd, *J* = 9.5, 8.0 Hz, 2H).

46.3 3-(5-Nitro-pyridin-2-yl)-azetidine-1-carboxylic acid tert-butyl ester

Zn dust (520 mg, 7.95 mmol) was vigorously stirred in THF (2 ml) under nitrogen and 1,2 dibromoethane (84 μl, 0.97 mmol) was added. The suspension was then heated at 80°C for 8 min and next allowed to cool to room temperature. Trimethylsilyl chloride (115 μl, 0.92 mmol) in THF (1 ml) was then added and the mixture further stirred at room temperature for 45 min. A solution of 3-iodo-azetidine-1-carboxylic acid tert-butyl ester (1.74 g, 6.14 mmol) in THF (2 ml) was then added dropwise to the solution over a period of 15 min and the reaction mixture stirred at room temperature for 2 h. Pd₂(dba)₃ (90 mg, 0.10 mmol) and P(2-furyl)₃ (85 mg, 0.36 mmol) were then added to the mixture, followed by 2-bromo-5-nitropyridine (1.37 g, 6.74 mmol) in THF (4 ml). The mixture was then heated at 55°C for 3 h, cooled to room temperature and quenched with saturated aqueous NaCl. Extraction with CH₂Cl₂, drying (Na₂SO₄) of the organic phase, filtration and evaporation in vacuo provided the crude material, which was purified by flash column chromatography (heptane:ethyl acetate, 3:1) to give the title compound (1.22 g, 71%) as a light yellow oil.

MS (ESI+) m/z = 224.1 [M-*t*Bu+H]⁺

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 1.47 (s, 9H), 3.99 (m, 1H), 4.18 (dd, *J* = 8.2, 6.1 Hz, 2H), 4.35 (t, *J* = 8.6 Hz, 2H), 7.42 (d, *J* = 8.5 Hz, 1H), 8.45 (dd, *J* = 8.5, 2.6 Hz, 1H), 9.44 (d, *J* = 2.4 Hz, 1H).

46.4 1-[3-(5-Nitro-pyridin-2-yl)-azetidin-1-yl]-propan-1-one

A solution of 3-(5-nitro-pyridin-2-yl)-azetidine-1-carboxylic acid tert-butyl ester (1.22 g, 4.36 mmol) in CH_2Cl_2 (100 ml) was treated with trifluoroacetic acid (TFA) (15 ml) and then stirred at room temperature for 2 h. After concentration, the raw mixture was digested in CH_2Cl_2 and washed with saturated aqueous NaHCO_3 . The aqueous phase was extracted with CH_2Cl_2 (x3), the combined organic layers dried over Na_2SO_4 and evaporated. The raw material was next dissolved in THF (80 ml) and the solution cooled to 0°C . Propionyl chloride (460 μl , 5.26 mmol) and triethylamine (735 μl , 5.28 mmol) were next added, the mixture allowed to reach 20°C and stirred for a further 12 h. It was then diluted with CH_2Cl_2 and washed successively with 1N aqueous HCl, saturated aqueous NaHCO_3 and water. The organic layer was dried (Na_2SO_4) and evaporated. The residue was chromatographed on silica gel (CH_2Cl_2 :methanol, 49:1) to afford the title compound (880 mg, 85% for two steps) as a brown oil.

MS (ESI+) m/z = 236.1 $[\text{M}+\text{H}]^+$

^1H NMR (400 MHz, CDCl_3) : δ (ppm) 1.16 (t, J = 7.5 Hz, 3H), 2.18 (q, J = 7.5 Hz, 2H), 4.07 (m, 1H), 4.22 (dd, J = 9.5, 6.0 Hz, 1H), 4.44 (m, 2H), 4.52 (t, J = 8.4 Hz, 1H), 7.42 (d, J = 8.5 Hz, 1H), 8.46 (dd, J = 8.5, 2.6 Hz, 1H), 9.45 (d, J = 2.5 Hz, 1H).

46.5 4-Isopropyl-N-[6-(1-propionyl-azetidin-3-yl)-pyridin-3-yl]-benzenesulfonamide

1-[3-(5-nitro-pyridin-2-yl)-azetidin-1-yl]-propan-1-one (340 mg, 1.44 mmol) was dissolved in ethanol (EtOH) (25 ml) and $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (1.63 g, 7.22 mmol) was added. The resulting mixture was refluxed for 8 h and the solvent next removed under vacuum. The raw material was dissolved in ethyl acetate and washed successively with 2N aqueous NaOH (x2) and water. The organic layer was dried (Na_2SO_4), filtered through a pad of celite and evaporated. Half of the crude material was then dissolved in CH_2Cl_2 (40 ml) and pyridine (115 μl , 1.41 mmol) followed by 4-isopropylbenzenesulfonylchloride (190 μl , 1.05 mmol) were added dropwise. After stirring at room temperature overnight, the reaction mixture was diluted with CH_2Cl_2 and washed successively with 1N aqueous HCl, saturated aqueous NaHCO_3 and water. The organic layer was dried (Na_2SO_4) and evaporated. The residue was chromatographed on silica gel (heptane:ethyl acetate, 1:3) to afford the title compound (150 mg, 54% for two steps) as a light yellow oil.

MS (ESI+) m/z = 388.1 $[\text{M}+\text{H}]^+$

135

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 1.14 (t, *J* = 7.5 Hz, 3H), 1.24 (d, *J* = 6.9 Hz, 6H), 2.14 (q, *J* = 7.5 Hz, 2H), 2.94 (m, 1H), 3.87 (m, 1H), 4.10 (dd, *J* = 9.7, 5.9 Hz, 1H), 4.34 (m, 2H), 4.42 (t, *J* = 8.5 Hz, 1H), 7.12 (d, *J* = 8.4 Hz, 1H), 7.31 (m, 3H), 7.59 (dd, *J* = 8.4, 2.6 Hz, 1H), 7.71 (d, *J* = 8.4 Hz, 2H), 8.25 (d, *J* = 2.3 Hz, 1H).

5

46.6 4-Isopropyl-N-[6-(1-propyl-azetidin-3-yl)-pyridin-3-yl]-benzenesulfonamide

To a solution of 4-isopropyl-N-[6-(1-propionyl-azetidin-3-yl)-pyridin-3-yl]-benzenesulfonamide (150 mg, 0.38 mmol) in THF (15 ml) was added dropwise 1M BH₃.THF (3.8 ml) and the mixture was stirred at room temperature for 12 h. It was then quenched by careful addition of 1N aqueous HCl (10 ml) and the resulting solution was heated at reflux for 4 h. The solution was next cooled to room temperature, adjusted to pH~8 with 2 N NaOH solution and diluted with CH₂Cl₂. Separation of the layers, drying (Na₂SO₄) of the organic phase, filtration and evaporation in vacuo provided the crude material, which was purified by flash column chromatography (CH₂Cl₂:methanol, 95:5) to give the title compound (95 mg, 66%) as a light yellow oil.

15

MS (ESI+) *m/z* = 374.1 [M+H]⁺

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 0.93 (t, *J* = 7.4 Hz, 3H), 1.24 (d, *J* = 6.9 Hz, 6H), 1.47 (m, 2H), 2.62 (m, 2H), 2.94 (m, 1H), 3.44 (m, 2H), 3.90 (m, 3H), 7.15 (d, *J* = 8.4 Hz, 1H), 7.31 (d, *J* = 8.3 Hz, 2H), 7.52 (dd, *J* = 8.4, 2.6 Hz, 1H), 7.70 (d, *J* = 8.3 Hz, 2H), 8.22 (d, *J* = 1.8 Hz, 1H).

20

EXAMPLE 47

N-[6-(1-Propyl-azetidin-3-yl)-pyridin-3-yl]-4-trifluoromethoxy-benzenesulfonamide

25

47.1 N-[6-(1-Propionyl-azetidin-3-yl)-pyridin-3-yl]-4-trifluoromethoxy-benzenesulfonamide.

Following the same procedure as described above, the second half of the crude aminopyridine obtained in 46.5 was treated with pyridine (115 μl, 1.41 mmol) and 4-(trifluoromethoxy)benzenesulfonylchloride (180 μl, 1.06 mmol) in CH₂Cl₂ (40 ml). Purification of the crude product by flash column chromatography (heptane:ethyl acetate, 1:2) provides the title compound (130 mg, 42% for two steps) as a light yellow oil.

30

MS (ESI+) *m/z* = 430.0 [M+H]⁺

136

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 1.14 (t, *J* = 7.5 Hz, 3H), 2.16 (q, *J* = 7.5 Hz, 2H), 3.86 (m, 1H), 4.08 (dd, *J* = 9.5, 6.1 Hz, 1H), 4.34 (m, 2H), 4.43 (t, *J* = 8.4 Hz, 1H), 7.12 (d, *J* = 8.4 Hz, 1H), 7.29 (d, *J* = 8.3 Hz, 2H), 7.60 (dd, *J* = 8.4, 2.6 Hz, 1H), 7.87 (d, *J* = 8.9 Hz, 2H), 8.01 (bs, 1H), 8.28 (d, *J* = 2.0 Hz, 1H).

5

47.2 N-[6-(1-Propyl-azetidin-3-yl)-pyridin-3-yl]-4-trifluoromethoxy-benzenesulfonamide

Following the same procedure as described in Example 46.6, N-[6-(1-propionyl-azetidin-3-yl)-pyridin-3-yl]-4-trifluoromethoxy-benzenesulfonamide (130 mg, 0.30 mmol) in THF (15 ml) was treated with 1M BH₃.THF (3 ml). Purification of the crude product by flash column chromatography (CH₂Cl₂:methanol, 95:5) provides the title compound (75 mg, 60%) as a light yellow oil.

MS (ESI+) *m/z* = 416.1 [M+H]⁺

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 0.96 (t, *J* = 7.4 Hz, 3H), 1.57 (m, 2H), 2.88 (t, *J* = 7.7 Hz, 2H), 3.76 (t, *J* = 8.1 Hz, 2H), 3.98 (m, 1H), 4.19 (t, *J* = 8.3 Hz, 2H), 7.04 (d, *J* = 8.4 Hz, 1H), 7.24 (d, *J* = 8.4 Hz, 2H), 7.33 (bs, 1H), 7.50 (dd, *J* = 8.3, 2.5 Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 2H), 8.33 (d, *J* = 2.3 Hz, 1H).

Example 48

20 4-Isopropyl-N-[4-(1-propyl-azetidin-3-yl)-phenyl]-benzene sulfonamide

48.1 3-(4-Nitro-phenyl)-azetidine-1-carboxylic acid tert-butyl ester

Following the same procedure as described in 46.3, the organozinc species was prepared using 3-iodo-azetidine-1-carboxylic acid tert-butyl ester (1.74 g, 6.14 mmol), Zn dust (520 mg, 7.95 mmol), 1,2 dibromoethane (84 μl, 0.97 mmol) and trimethylsilyl chloride (115 μl, 0.92 mmol). It was then coupled with 1-bromo-4-nitrobenzene (1.24 g, 6.13 mmol) using Pd₂(dba)₃ (90 mg, 0.10 mmol) and P(2-furyl)₃ (85 mg, 0.36 mmol). Purification of the crude product by flash column chromatography (heptane:ethyl acetate, 4:1) provides the title compound (880 mg, 52%) as a light yellow oil.

30

MS (ESI+) *m/z* = 223.1 [M-*t*Bu+H]⁺.

48.2 1-[3-(4-Nitro-phenyl)-azetidin-1-yl]-propan-1-one

137

Using the same procedure as described in 46.4, 3-(4-nitro-phenyl)-azetidine-1-carboxylic acid tert-butyl ester was deprotected with TFA (15 ml) in CH₂Cl₂ (100 ml) and the resulting amine treated with propionyl chloride (290 µl, 3.31 mmol) and triethylamine (470 µl, 3.37 mmol) in THF (80 ml). Purification of the crude product by flash column chromatography (heptane:ethyl acetate, 1:1) provides the title compound (570 mg, 87% for two steps) as a brown oil.

MS (ESI+) m/z = 235.1 [M+H]⁺

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 1.17 (t, J = 7.5 Hz, 3H), 2.17 (q, J = 7.5 Hz, 2H), 3.93 (m, 1H), 4.09 (dd, J = 9.8, 5.9 Hz, 1H), 4.14 (dd, J = 8.3, 6.0 Hz, 1H), 4.47 (t, J = 9.3 Hz, 1H), 4.59 (t, J = 8.6 Hz, 1H), 7.49 (d, J = 8.7 Hz, 2H), 8.24 (d, J = 8.7 Hz, 2H).

48.3 4-Isopropyl-N-[4-(1-propionyl-azetidin-3-yl)-phenyl]-benzenesulfonamide

Following the same procedure as described in 46.5, 1-[3-(4-nitro-phenyl)-azetidin-1-yl]-propan-1-one (480mg, 2.04 mmol) in EtOH (20 ml) was treated with SnCl₂·2H₂O (2.25 g, 9.97 mmol) and half of the resulting aniline in CH₂Cl₂ (15 ml) was then treated with pyridine (140 µl, 1.71 mmol) and 4-isopropylbenzenesulfonylchloride (230 µl, 1.28 mmol). Purification of the crude product by flash column chromatography (heptane:ethyl acetate, 1:1) provided the title compound (140 mg, 36% for two steps) as a pale yellow gum.

MS (ESI+) m/z = 387.1 [M+H]⁺

48.4 4-Isopropyl-N-[4-(1-propyl-azetidin-3-yl)-phenyl]-benzene sulfonamide

Following the same procedure as described in Example 46.6, 4-isopropyl-N-[4-(1-propionyl-azetidin-3-yl)-phenyl]-benzenesulfonamide (140 mg, 0.36 mmol) in THF (18 ml) was then treated with 1M BH₃·THF (3.6 ml). The crude material was chromatographed (ethyl acetate) to afford the title compound (90 mg, 67%) as a pale yellow oil.

MS (ESI+) m/z = 373.1 [M+H]⁺

¹H NMR (400 MHz, CDCl₃) : δ (ppm) δ 0.93 (t, J = 7.4 Hz, 3H), 1.24 (d, J = 6.9 Hz, 6H), 1.38 (m, 2H), 2.41 (t, J = 7.5 Hz, 2H), 2.94 (m, 1H), 3.03 (m, 2H), 3.67 (m, 3H), 7.01 (d, J = 8.4 Hz, 2H), 7.16 (d, J = 8.4 Hz, 2H), 7.28 (d, J = 8.5 Hz, 2H), 7.68 (d, J = 8.4 Hz, 2H).

EXAMPLE 49

N-[4-(1-Propyl-azetidin-3-yl)-phenyl]-4-trifluoromethoxybenzene sulfonamide

49.1 N-[4-(1-Propionyl-azetidin-3-yl)-phenyl]-4-trifluoromethoxy-benzenesulfonamide

5

Following the same procedure as described above, the second half of the resulting aniline from 48.3 was treated with pyridine (140 μ l, 1.71 mmol) and 4-(trifluoromethoxy)benzenesulfonylchloride (215 μ l, 1.26 mmol) in CH_2Cl_2 (15 ml). Purification of the crude product by flash column chromatography (heptane:ethyl acetate, 2:1) provided the title compound (170 mg, 40% for two steps) as a pale yellow gum. MS (ESI+) m/z = 429.0 $[\text{M}+\text{H}]^+$

10

49.3 N-[4-(1-Propyl-azetidin-3-yl)-phenyl]-4-trifluoromethoxybenzene sulfonamide

Following the same procedure as described in Example 47.2, N-[4-(1-propionyl-azetidin-3-yl)-phenyl]-4-trifluoromethoxy-benzenesulfonamide (170 mg, 0.39 mmol) in THF (20 ml) was then treated with 1M $\text{BH}_3\cdot\text{THF}$ (3.9 ml). The crude material was chromatographed (ethyl acetate) to afford the title compound (86 mg, 52%) as a pale yellow oil.

MS (ESI+) m/z = 415.1 $[\text{M}+\text{H}]^+$

20

^1H NMR (400 MHz, CDCl_3) : δ (ppm) 0.90 (t, J = 7.4 Hz, 3H), 1.38 (m, 2H), 2.43 (t, J = 7.5 Hz, 2H), 3.05 (m, 2H), 3.68 (m, 3H), 7.01 (d, J = 8.4 Hz, 2H), 7.18 (d, J = 8.4 Hz, 2H), 7.25 (d, J = 8.5 Hz, 2H), 7.79 (d, J = 8.8 Hz, 2H).

25 EXAMPLE 50

4-Bromo-N-[2-(1-propyl-pyrrolidin-3-yl)-pyrimidin-5-yl]-benzenesulfonamide.

50.1 5-Bromo-2-vinyl-pyrimidine

5-Bromo-2-iodo-pyrimidine (9.15 g, 32.11 mmol) was dissolved in THF (150 ml) and $\text{Pd}(\text{PPh}_3)_4$ (18.85g, 1.60 mmol) was added, followed by tributyl-vinyl-stannane (9.38 ml, 32.11 mmol). The resulting mixture was heated at 140°C for 20 min under microwave irradiations, next filtered through a pad of celite and concentrated. The crude material was diluted with CH_2Cl_2 and washed with water. Separation of the layers, drying (Na_2SO_4) of the organic phase, filtration and evaporation in vacuo provided the crude

35

material, which was purified by flash column chromatography (CH_2Cl_2) to give the title compound (4.05 g, 68%) as a volatile yellow oil crystallizing at 4°C.

^1H NMR (400 MHz, CDCl_3) : δ (ppm) 5.77 (dd, $J = 10.5, 1.4$ Hz, 1H), 6.62 (dd, $J = 17.3, 1.4$ Hz, 1H), 6.83 (dd, $J = 17.3, 10.5$ Hz, 1H).

5

50.2 5-Bromo-2-(1-propyl-pyrrolidin-3-yl)-pyrimidine

Methoxymethyl-propyl-trimethylsilanylmethyl-amine (14.13 g, 74.61 mmol) in CH_2Cl_2 (4 ml) was added dropwise to a 0°C cooled solution of 5-bromo-2-vinyl-pyrimidine (2 g, 10.80 mmol) and TFA (210 μl , 2.72 mmol) in CH_2Cl_2 (45 ml) over a period of 20 min. The reaction mixture was then stirred at room temperature for 2 h. The crude material was diluted with CH_2Cl_2 , washed with saturated aqueous NaHCO_3 and the organic layer was dried over Na_2SO_4 and evaporated. The residue was chromatographed on silica gel (CH_2Cl_2 :methanol, 97:3) to afford the title compound (1.03 g, 34%) as a brown oil.

15

MS (ESI+) $m/z = 271.9$ $[\text{M}+\text{H}]^+$

50.3 2-(1-Propyl-pyrrolidin-3-yl)-pyrimidin-5-ylamine.

A dry flask was charged with 5-bromo-2-(1-propyl-pyrrolidin-3-yl)-pyrimidine (300 mg, 1.11 mmol), $\text{Pd}_2(\text{dba})_3$ (tris(dibenzylideneacetone) dipalladium (30 mg, 0.032 mmol)), *rac*-BINAP (2,2'-bis(diphenylphosphino)-1,1'-binaphthyl) (35 mg, 0.056 mmol), tBuONa (150 mg, 1.56 mmol), benzophenone imine (300 mg, 1.65 mmol) and toluene (4 ml). It was then evacuated, flushed with nitrogen and the mixture heated at 80°C under microwave irradiations for 3h. After filtration through a pad of celite, the solvent was removed under reduced pressure. The crude material was then dissolved in THF (10 ml) and 1 N aqueous HCl (3 ml) was added. The solution was stirred at room temperature for 45 min and the organic solvent was next removed under vacuo. The resulting aqueous phase was adjusted to pH~9 with 2 N NaOH solution, washed with heptane:ethyl acetate, 2:1 (50 ml) and concentrated to give the crude amine.

25
30

MS (ESI+) $m/z = 207.1$ $[\text{M}+\text{H}]^+$

50.4 4-Bromo-N-[2-(1-propyl-pyrrolidin-3-yl)-pyrimidin-5-yl]-benzenesulfonamide

140

The crude 2-(1-propyl-pyrrolidin-3-yl)-pyrimidin-5-ylamine (about 1.11 mmol) was dissolved in CH₂Cl₂:pyridine, 9:1 (50 ml) and 4-bromobenzenesulfonyl chloride (566 mg, 2.21 mmol) was added. After stirring at room temperature overnight, the reaction mixture was filtered and concentrated. The raw material was then dissolved in a solution of

5 EtONa (400 mg, 5.87 mmol) in EtOH (35 ml) and the resulting solution was heated under reflux for 1h. Silica gel (1 g) was next added to the mixture and the solvent removed under reduced pressure. Following flash column chromatography (CH₂Cl₂:methanol, 9:1) provides the title compound (100 mg, 21% for four steps) as a yellow gum.

10 MS (ESI+) $m/z = 426.9$ [M+H]⁺

¹H NMR (400 MHz, CDCl₃) : rotamers δ (ppm) 0.99 (t, $J = 7.3$ Hz, 3H), 1.84 (m, 2H), 2.33 (m, 1H), 2.53 (m, 1H), 3.05 (m, 1H), 3.17 (m, 1H), 3.46 (t, $J = 6.0$ Hz, 2H), 3.70 (m, 2H), 3.86 (m, 1H), 5.67 (bs, 1H), 7.53 (d, $J = 8.4$ Hz, 2H), 7.70-7.75 (2d, $J = 8.4$ Hz, 2H), 8.44 (s, 2H).

15

EXAMPLE 51

N-[2-(1-Propyl-pyrrolidin-3-yl)-pyrimidin-5-yl]-4-trifluoromethoxy-benzenesulfonamide.

Following the same procedure as described in Example 50.4, a crude 2-(1-propyl-pyrrolidin-3-yl)-pyrimidin-5-ylamine (about 0.74 mmol) obtained via the above mentioned protocol was treated with 4-(trifluoromethoxy)benzenesulfonyl chloride (250 μ l, 1.47 mmol) in CH₂Cl₂:pyridine, 9:1 (50 ml). Subsequent treatment with EtONa/EtOH and purification of the crude product by flash column chromatography (CH₂Cl₂:methanol, 9:1) provided the title compound (44 mg, 14% for four steps) as a

20 yellow gum.

25

MS (ESI+) $m/z = 431.0$ [M+H]⁺

¹H NMR (400 MHz, CDCl₃) : rotamers δ (ppm) 0.98 (t, $J = 7.3$ Hz, 3H), 1.84 (m, 2H), 2.30 (m, 1H), 2.58 (m, 1H), 3.14 (m, 2H), 3.31 (m, 1H), 3.64 (m, 2H), 3.92 (m, 2H), 7.24 (m, 2H), 7.93-7.97 (2d, $J = 8.6$ Hz, 2H), 8.44 (s, 2H).

30

EXAMPLE 52

4-Isopropyl-N-[2-(1-propyl-pyrrolidin-3-yl)-pyrimidin-5-yl]-benzenesulfonamide.

141

Following the same procedure as described in Example 50.4, a crude 2-(1-propyl-pyrrolidin-3-yl)-pyrimidin-5-ylamine (about 0.13 mmol) obtained via the above mentioned protocol was treated with 4-isopropylbenzenesulfonyl chloride (56 μ l, 0.31 mmol) in CH_2Cl_2 :pyridine, 9:1 (50 ml). Subsequent treatment with EtONa/EtOH and purification of the crude product by flash column chromatography (CH_2Cl_2 :methanol, 85:15) provided the title compound (26 mg, 50% for four steps) as a yellow gum.

MS (ESI+) m/z = 389.1 $[\text{M}+\text{H}]^+$

^1H NMR (400 MHz, CDCl_3) : rotamers δ (ppm) 0.96 (t, J = 7.4 Hz, 3H), 1.21-1.23 (2d, J = 6.9 Hz, 6H), 1.80 (m, 2H), 2.26 (m, 1H), 2.52 (m, 1H), 2.90 (m, 1H), 3.07 (m, 2H), 3.24 (m, 1H), 3.55 (m, 2H), 3.85 (m, 2H), 6.34 (bs, 1H), 7.25 (m, 2H), 7.76-7.81 (2d, J = 8.2 Hz, 2H), 8.54 (s, 2H).

EXAMPLE 53

N-{4-[(S)-1-(3-Hydroxy-propyl)-pyrrolidin-3-yl]-phenyl}-4-isopropyl-benzenesulfonamide

53.1 Acetic acid 3-{(S)-3-[4-(4-isopropyl-benzenesulfonylamino)-phenyl]-pyrrolidin-1-yl}-propyl ester

0.21 g of 4-isopropyl-N-[4-((S)-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide (0.62 mmol) were dissolved in 5 ml dimethylformamide. 0.0019 mg of sodium iodide (0.01 mmol) and 0.13 ml of triethylamine (0.94 mmol) were added, followed by addition of 0.136 mg of 3-acetoxy-1-bromo-propane (0.75 mmol). Stirring continued for 15 h at room temperature before the reaction mixture was poured onto 50 ml of crushed ice. The aqueous layer was extracted three times with ethyl acetate and the combined organic phases washed with water and saturated sodium chloride solution. The ethyl acetate phase was dried over magnesium sulfate, filtered, and evaporated to dryness under reduced pressure to yield 0.26 g of the crude product.

ESI-MS: 445.1 $[\text{M}+\text{H}]^+$

53.2 N-{4-[(S)-1-(3-Hydroxy-propyl)-pyrrolidin-3-yl]-phenyl}-4-isopropyl-benzenesulfonamide

0.26 g of acetic acid 3-{(S)-3-[4-(4-isopropyl-benzenesulfonylamino)-phenyl]-pyrrolidin-1-yl}-propyl ester (0.58 mmol) were dissolved in 4 ml of tetrahydrofuran. 0.021 g lithium hydroxide (0.88 mmol) dissolved in 4 ml water were added and the reaction mixture stirred for 15 h at room temperature. An additional equivalent of lithium hydroxide was

142

added and stirring continued for 24 h before the reaction mixture was diluted with water and extracted three times with ethyl acetate. The combined organic phases were dried over magnesium sulfate, filtered, and evaporated to dryness under reduced pressure to yield 0.157g of the product.

5 ESI-MS: 403.3 $[M+H]^+$

$^1\text{H-NMR}$ (CDCl_3): δ [ppm] 7.7 (d, 2H), 7.25 (m, 2H), 7.1 (d, 2H), 6.95 (d, 2H), 3.8 (m, 2H), 3.3 (m, 1H), 3.05 (m, 1H), 2.95 (m, 1H), 2.7-2.9 (m, 4H), 2.55 (m, 1H), 2.25 (m, 1H), 1.7-1.85 (m, 2H), 1.15-1.3 (broad, 7H).

10 EXAMPLE 54

4-Cyclobutyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

The title compound was prepared in an analogous manner to that described for Example 37.

15 ESI-MS: 399.1 $[M+H]^+$

EXAMPLE 55

4-Oxazol-5-yl-N-[4-(1-propyl-azetidin-3-yl)-phenyl]-benzenesulfonamide

20 55.1 4-(1-Propyl-azetidin-3-yl)-phenylamine

1-(3-(4-Nitrophenyl)azetidin-1-yl)propan-1-one (3.16 g, 13.49 mmol) was dissolved in ethanol (200 ml) and $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (15.20 g, 67.45 mmol) was added. The resulting mixture was refluxed for 8 h and then the solvent was removed under vacuum. The raw material was dissolved in ethyl acetate and washed successively with 2N aqueous NaOH (x2) and water. The organic layer was dried (Na_2SO_4), filtered through a pad of celite and evaporated. The crude 1-[3-(4-amino-phenyl)-azetidin-1-yl]-propan-1-one was then dissolved in tetrahydrofuran (THF) (200 ml) and 1M LiAlH_4 in tetrahydrofuran (19.5 ml, 19.5 mmol) was added dropwise at 0°C . After stirring at room temperature for 2 h, the reaction mixture was carefully quenched with THF/ H_2O 9:1 (20 ml) at 0°C , then filtered through a pad of celite and the solvents were removed under reduced pressure. The crude 4-(1-propyl-azetidin-3-yl)-phenylamine was used without any further purification for the next step.

MS (ESI+) m/z = 191.1 $[M+H]^+$

35

55.2 4-Oxazol-5-yl-N-[4-(1-propyl-azetidin-3-yl)-phenyl]-benzenesulfonamide

The crude 4-(1-propyl-azetidin-3-yl)-phenylamine (40 mg, 0.21 mmol) was dissolved in CH₂Cl₂/pyridine 9:1 (10 ml) and 4-oxazol-5-yl-benzenesulfonyl chloride (51 mg, 0.21 mmol) was added. After stirring at room temperature for 2 h, the reaction mixture was diluted with CH₂Cl₂ and washed successively with 1N aqueous HCl, saturated aqueous NaHCO₃ and water. The organic layer was dried over Na₂SO₄ and evaporated. The residue was chromatographed on silica gel (CH₂Cl₂:methanol, 95:5) to afford the title compound (30 mg, 36%) as a white amorphous solid.

MS (ESI+) m/z = 398.1 [M+H]⁺

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 1.02 (t, J = 7.4 Hz, 3H), 1.75 (m, 2H), 3.08 (m, 2H), 3.72 (m, 2H), 4.24 (m, 1H), 4.62 (m, 2H), 7.10 (m, 2H), 7.22 (d, J = 7.9 Hz, 2H), 7.45 (s, 1H), 7.69 (d, J = 8.4 Hz, 2H), 7.89 (d, J = 8.3 Hz, 2H), 7.95 (s, 1H).

EXAMPLE 56

4-(2-Fluoro-ethyl)-N-[4-(1-propyl-azetidin-3-yl)-phenyl]-benzenesulfonamide

Following the same procedure as described in example 55, 4-(1-propyl-azetidin-3-yl)-phenylamine (100 mg, 0.52 mmol) in CH₂Cl₂/pyridine 9:1 (12 ml) was treated with 4-(2-fluoro-ethyl)-benzenesulfonyl chloride (117 mg, 0.52 mmol). Purification of the crude product by chromatography on reversed phase silica gel (H₂O+0.1% acetic acid:CH₃CN+0.1% acetic acid, 75:25) provided the title compound (11 mg, 6%) as a colourless gum.

MS (ESI+) m/z = 377.1 [M+H]⁺

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 0.97 (t, J = 7.4 Hz, 3H), 1.59 (m, 2H), 2.89 (m, 2H), 3.00 (t, J = 6.0 Hz, 1H), 3.06 (t, J = 6.0 Hz, 1H), 3.76 (m, 2H), 4.05 (m, 1H), 4.29 (m, 2H), 4.57 (t, J = 6.0 Hz, 1H), 4.69 (t, J = 6.0 Hz, 1H), 7.08 (d, J = 8.5 Hz, 2H), 7.12 (d, J = 8.5 Hz, 2H), 7.32 (d, J = 8.2 Hz, 2H), 7.71 (d, J = 8.2 Hz, 2H).

EXAMPLE 57

4-(3-Fluoro-propyl)-N-[4-(1-propyl-azetidin-3-yl)-phenyl]-benzenesulfonamide

Following the same procedure as described in example 55, 4-(1-propyl-azetidin-3-yl)-phenylamine (100 mg, 0.52 mmol) in CH₂Cl₂/pyridine 1:1 (12 ml) was treated with 4-(3-fluoro-propyl)-benzenesulfonyl chloride (161 mg, 0.68 mmol). Purification of the crude product by chromatography on reversed phase silica gel (H₂O+0.1% acetic acid:CH₃CN+0.1% acetic acid, 75:25) provided the title compound (40 mg, 20%) as a colourless gum.

MS (ESI+) m/z = 391.3 [M+H]⁺

144

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 0.96 (t, *J* = 7.4 Hz, 3H), 1.66 (m, 2H), 1.97 (m, 2H), 2.76 (m, 2H), 3.08 (m, 1H), 3.25 (m, 1H), 3.73 (m, 2H), 4.21 (m, 1H), 4.34 (t, *J* = 5.7 Hz, 1H), 4.46 (t, *J* = 5.7 Hz, 1H), 4.66 (m, 2H), 7.04 (d, *J* = 8.3 Hz, 2H), 7.12 (d, *J* = 8.3 Hz, 2H), 7.25 (d, *J* = 8.1 Hz, 2H), 7.70 (d, *J* = 8.3 Hz, 2H), 8.17 (s, 1H).

5

EXAMPLE 58

4-Bromo-N-[4-(1-propyl-azetidin-3-yl)-phenyl]-benzenesulfonamide

Following the same procedure as described in example 55, 4-(1-propyl-azetidin-3-yl)-phenylamine (1 g, 5.25 mmol) in CH₂Cl₂/pyridine 9:1 (50 ml) was treated with 4-bromobenzenesulfonyl chloride (1.34 g, 5.25 mmol). Purification of the crude product by flash column chromatography (CH₂Cl₂:methanol, 95:5) provided the title compound (1.1 g, 51%) as a colourless gum.

MS (ESI+) *m/z* = 410.0 [M+H]⁺

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 0.92 (t, *J* = 7.4 Hz, 3H), 1.47 (m, 2H), 2.60 (m, 2H), 3.28 (m, 2H), 3.78 (m, 1H), 3.91 (m, 2H), 6.07 (bs, 1 H), 7.07 (d, *J* = 8.4 Hz, 2H), 7.13 (d, *J* = 8.4 Hz, 2H), 7.53 (d, *J* = 8.5 Hz, 2H), 7.64 (d, *J* = 8.5 Hz, 2H).

EXAMPLE 59

4-((R)-2-Fluoro-1-methyl-ethyl)-N-[4-(1-propyl-azetidin-3-yl)-phenyl]-benzenesulfonamide

Following the same procedure as described in example 55, 4-(1-propyl-azetidin-3-yl)-phenylamine (150 mg, 0.78 mmol) in CH₂Cl₂/pyridine 1:1 (12 ml) was treated with 4-((R)-2-fluoro-1-methyl-ethyl)-benzenesulfonyl chloride (242 mg, 1.02 mmol). Purification of the crude product by chromatography on reversed phase silica gel (H₂O+0.1% acetic acid:CH₃CN+0.1% acetic acid, 75:25) provided the title compound (15 mg, 5%) as a colourless gum.

MS (ESI+) *m/z* = 391.1 [M+H]⁺

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 0.93 (t, *J* = 7.3 Hz, 3H), 1.27 (m, 3H), 1.60 (m, 2H), 2.99 (m, 2H), 3.13 (m, 1H), 3.86 (m, 2H), 4.10 (m, 1H), 4.42 (m, 5H), 4.36 (m, 1H), 4.49 (m, 1H), 7.07 (m, 4H), 7.27 (d, *J* = 8.3 Hz, 2H), 7.74 (d, *J* = 8.3 Hz, 2H).

EXAMPLE 60

4-((S)-2-Fluoro-1-methyl-ethyl)-N-[4-(1-propyl-azetidin-3-yl)-phenyl]-benzenesulfonamide

145

Following the same procedure as described in example 55, 4-(1-propyl-azetidin-3-yl)-phenylamine (110 mg, 0.57 mmol) in CH₂Cl₂/pyridine 1:1 (12 ml) was treated with 4-((S)-2-fluoro-1-methyl-ethyl)-benzenesulfonyl chloride (178 mg, 0.75 mmol). Purification of the crude product by flash column chromatography (CH₂Cl₂:methanol, 95:5)

5 provided the title compound (50 mg, 22%) as a gum.

MS (ESI+) m/z = 391.1 [M+H]⁺

¹H NMR (400 MHz, CDCl₃) : δ (ppm) 0.93 (t, J = 7.3 Hz, 3H), 1.27 (m, 3H), 1.60 (m, 2H), 2.92 (m, 2H), 3.15 (m, 1H), 3.73 (m, 2H), 4.06 (m, 1H), 4.32 (m, 2H), 4.36 (m, 1H), 4.49 (m, 1H), 7.07 (m, 4H), 7.28 (d, J = 8.3 Hz, 2H), 7.74 (d, J = 8.3 Hz, 2H).

10

EXAMPLE 61

N-[4-(1-Propyl-azetidin-3-yl)-phenyl]-4-(3,3,3-trifluoro-propyl)-benzenesulfonamide

Following the same procedure as described in example 55, 4-(1-propyl-azetidin-3-yl)-phenylamine (100 mg, 0.52 mmol) in CH₂Cl₂/pyridine 9:1 (12 ml) was treated with 4-(3,3,3-trifluoro-propyl)-benzenesulfonyl chloride (214 mg, 0.78 mmol). Purification of the crude product by flash column chromatography (CH₂Cl₂:methanol, 95:5) provided the title compound (26 mg, 11%) as a colourless gum.

MS (ESI+) m/z = 427.1 [M+H]⁺

20 ¹H NMR (400 MHz, CDCl₃) : δ (ppm) 0.99 (t, J = 7.4 Hz, 3H), 1.73 (m, 2H), 2.36 (m, 2H), 2.87 (m, 2H), 3.10 (m, 2H), 3.70-4.20 (m, 5H), 7.15 (m, 4H), 7.24 (d, J = 8.5 Hz, 2H), 7.79 (d, J = 8.3 Hz, 2H).

EXAMPLE 62

25 N-[4-(1-Propyl-azetidin-3-yl)-phenyl]-4-(2,2,2-trifluoro-ethyl)-benzenesulfonamide

Following the same procedure as described in example 55, 4-(1-propyl-azetidin-3-yl)-phenylamine (100 mg, 0.52 mmol) in CH₂Cl₂/pyridine 9:1 (10 ml) was treated with 4-(3,3,3-trifluoro-ethyl)-benzenesulfonyl chloride (163 mg, 0.63 mmol). Purification of the crude product by flash column chromatography (CH₂Cl₂:methanol, 95:5) provided the title compound (65 mg, 30%) as a colourless gum.

MS (ESI+) m/z = 413.1 [M+H]⁺

30 ¹H NMR (400 MHz, CDCl₃) : δ (ppm) 0.98 (t, J = 7.4 Hz, 3H), 1.71 (m, 2H), 3.09 (m, 2H), 3.38 (m, 2H), 3.66-4.75 (m, 5H), 7.13 (m, 4H), 7.33 (d, J = 8.0 Hz, 2H), 7.83 (d, J = 8.1 Hz, 2H).

35

EXAMPLE 63

146

N-[4-(1-Propyl-azetidin-3-yl)-phenyl]-4-(2,2,2-trifluoro-1-methyl-ethyl)-benzenesulfonamide

Following the same procedure as described in example 55, 4-(1-propyl-azetidin-3-yl)-phenylamine (100 mg, 0.52 mmol) in CH₂Cl₂/pyridine 9:1 (15 ml) was treated with 4-(2,2,2-trifluoro-1-methyl-ethyl)-benzenesulfonyl chloride (145 mg, 0.52 mmol). Purification of the crude product by flash column chromatography (CH₂Cl₂:methanol, 95:5) provided the title compound (40 mg, 18%) as a pale yellow gum.

MS (ESI+) m/z = 427.1 [M+H]⁺

¹H-NMR (400 MHz, CDCl₃) : δ (ppm) 0.98 (t, J = 7.4 Hz, 3H), 1.49 (d, J = 7.2 Hz, 3H), 1.68 (m, 2H), 3.02-3.21 (m, 2H), 3.47 (m, 1H), 3.68-4.70 (m, 5H), 7.07 (d, J = 8.6 Hz, 2H), 7.12 (d, J = 8.5 Hz, 2H), 7.40 (d, J = 8.2 Hz, 2H), 7.76 (d, J = 8.3 Hz, 2H).

EXAMPLE 64

4-(2,2-Difluoro-1-methyl-ethyl)-N-[4-(1-propyl-azetidin-3-yl)-phenyl]-benzenesulfonamide;

Following the same procedure as described in example 55, 4-(1-propyl-azetidin-3-yl)-phenylamine (150 mg, 0.78 mmol) in CH₂Cl₂/pyridine 1:1 (12 ml) was treated with 4-(2,2-difluoro-1-methyl-ethyl)-benzenesulfonyl chloride (321 mg, 1.26 mmol). Purification of the crude product by flash column chromatography (CH₂Cl₂:methanol, 95:5) provided the title compound (40 mg, 18%) as a colourless gum.

MS (ESI+) m/z = 409.1 [M+H]⁺

¹H-NMR (400 MHz, CDCl₃) : δ (ppm) 0.97 (t, J = 7.4 Hz, 3H), 1.36 (m, 3H), 1.69 (m, 2H), 3.07 (m, 2H), 3.17 (m, 1H), 3.54-4.65 (m, 5H), 5.76 (td, J = 3.5 Hz, 1H), 7.11 (m, 4H), 7.30 (d, J = 8.1 Hz, 2H), 7.78 (d, J = 8.3 Hz, 2H).

EXAMPLE 65

N-[4-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-4-[1,2,3]thiadiazol-4-yl-benzenesulfonamide

Sulfonamide coupling was achieved by following a procedure analogous to that described in example 55. Yield: 75 mg (52%).

MS (ESI) m/z: 429.0 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 9.75 (s, 1H), 8.29 (d, 2H), 7.91 (d, 2H), 7.13 (d, 2H), 7.04 (d, 1H), 3.16 (m, 1H), 2.83 (m, 1H), 2.60 (t, 1H, J = 7.3 Hz, 2H), 2.23 (m, 3H), 2.15 (m, 1H), 1.65 (m, 1H), 1.43 (m, 2H), 0.85 (t, J = 7.3 Hz, 3H).

EXAMPLE 66

N-[2-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-4-[1,2,3]thiadiazol-4-yl-benzenesulfonamide

Sulfonamide coupling was achieved by following a procedure analogous to that described in example 55. Yield: 15 mg (10%)

MS (ESI) m/z: 429.0 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 9.73 (s, 1H), 8.30 (d, 2H), 7.88 (d, 2H), 7.12 (t, 1H), 6.98 (m, 3H), 3.16 (m, 1H), 2.81 (m, 1H), 2.60 (t, 1H, J = 7.3 Hz, 2H), 2.23 (m, 3H), 2.14 (m, 1H), 1.58 (m, 1H), 1.39 (m, 2H), 0.82 (t, J = 7.3 Hz, 3H).

EXAMPLE 67

N-[4-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-4-pyrrolidin-1-yl-benzenesulfonamide

Starting from 4-bromo-N[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]benzenesulfonamide obtained in example 25 and following a procedure in analogy to example 50; (pyrrolidine, BINAP, tert-NaOC₄H₉, Pd₂(dba)₃, 140°C MW (microwave)) the title compound was obtained. Chromatography CH₂Cl₂-methanol 9:1. Yield: 36 mg (14%).

MS (ESI) m/z: 414.1 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 9.85 (br s, 1H), 7.50 (d, 2H), 7.11 (d, 2H), 7.00 (d, 2H), 6.51 (d, 2H), 3.30 (m, 6H), 2.91 (m, 2H), 2.63 (m, 3H), 2.19 (m, 1H), 1.87 (m, 4H), 1.73 (m, 1H), 1.51 (m, 2H), 0.85 (t, J = 7.3 Hz, 3H).

EXAMPLE 68

N-[4-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-4-trifluoromethylsulfanyl-benzenesulfonamide

4-((S)-1-Propylpyrrolidin-3-yl)phenylamin (50 mg, 0.24 mmol) was dissolved in THF (5 ml) at -78°C and potassium hexamethyldisilazide (146 mg, 0.73 mmol) added. The solution was stirred at -78°C for 1 hour and then 4-(trifluoromethylthio)benzene-1-sulfonyl fluoride (64 mg, 0.24 mmol) was added and the solution was allowed to reach room temperature over night. The solvent was removed in vacuo and the residue partitioned between ethyl acetate and NaOH (2M). The organic extract was separated, dried (MgSO₄), filtered and concentrated to give the product (97 mg, yield 89%).

MS (ESI) m/z: 445.0 [M+H]⁺

¹H-NMR (DMSO-d₆): δ [ppm] 10.27 (br s, 1H), 7.87 (m, 2H), 7.68 (m, 2H), 7.19 (d, 2H), 7.03 (d, 2H), 3.30 (m, 2H), 3.16 (m, 2H), 2.82 (m, 3H), 2.26 (m, 1H), 1.82 (m, 1H), 1.53 (m, 2H), 0.88 (t, J = 7.3 Hz, 3H).

148

EXAMPLE 69

N-[4-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-4-pyrazol-1-yl-benzenesulfonamide

Sulfonamide coupling was achieved by following a procedure analogous to that described in example 55.2; 2 mg (34%) of the title compound were obtained.

MS (ESI) m/z: 411.1 [M+H]⁺

¹H-NMR (CH₃OH-d₄): δ [ppm] 8.30 (s, 1H), 7.84 (m, 4H), 7.72 (s, 1H), 7.19 (d, 2H), 7.08 (d, 2H), 6.52 (s, 1H), 3.50 (m, 2H), 3.30 (m, 2H), 2.96 (m, 2H), 2.38 (m, 1H), 2.02 (m, 1H), 1.68 (m, 2H), 0.96 (t, J = 7.3 Hz, 3H).

EXAMPLE 70

4-Oxazol-5-yl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

Sulfonamide coupling was achieved by following a procedure analogous to that described in example 55.2. 148 mg (yield 84%) of the title compound were obtained.

MS (ESI) m/z: 412.1 [M+H]⁺

¹H-NMR (CH₃OH-d₄): δ [ppm] 8.28 (s, 1H), 7.79 (m, 4H), 7.62 (s, 1H), 7.20 (d, 2H), 7.13 (d, 2H), 3.69 (m, 1H), 3.53 (m, 1H), 3.47 (m, 2H), 3.12 (m, 3H), 2.38 (m, 1H), 2.10 (m, 1H), 1.75 (m, 2H), 0.99 (t, J = 7.3 Hz, 3H).

EXAMPLE 71

4-Oxazol-5-yl-N-[2-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

Sulfonamide coupling was achieved by following a procedure analogous to that described in example 55.2. 11 mg (yield 6%) of the title compound were obtained.

MS (ESI) m/z: 412.1 [M+H]⁺

¹H-NMR (CH₃OH-d₄): δ [ppm] 8.29 (s, 1H), 7.82 (m, 4H), 7.66 (s, 1H), 7.21 (m, 1H), 7.06 (m, 3H), 3.61 (m, 1H), 3.50 (m, 1H), 3.37 (m, 2H), 3.03 (m, 3H), 2.38 (m, 1H), 2.05 (m, 1H), 1.69 (m, 2H), 0.99 (t, J = 7.3 Hz, 3H).

EXAMPLE 72

4-(2-Oxo-pyrrolidin-1-yl)-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

Sulfonamide coupling was achieved by following a procedure analogous to that described in example 55.2. 86 mg (yield 66%) of the title compound were obtained.

MS (ESI) m/z: 428.1 [M+H]⁺

149

¹H-NMR (CDCl₃): δ [ppm] 7.72 (m, 4H), 7.12 (d, 2H), 7.01 (d, 2H), 3.83 (t, 2H), 3.32 (m, 1H), 3.14 (m, 1H), 2.92 (m, 1H), 2.81 (m, 1H), 2.53 (m, 5H), 2.27 (m, 1H), 2.13 (m, 2H), 1.83 (m, 1H), 1.52 (m, 2H), 0.91 (t, J = 7.3 Hz, 3H).

5 EXAMPLE 73

4-Furan-2-yl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

Starting from 4-bromo-N[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]benzenesulfonamide obtained in example 25 and following a procedure in analogy to example 19 the title
10 compound (92 mg (yield 76%)) was obtained.

MS (ESI) m/z: 412.1 [M+H]⁺

¹H-NMR (CH₃OH-d₄): δ [ppm] 7.73 (s, 4H), 7.61 (s, 1H), 7.19 (d, 2H), 7.12 (d, 2H), 6.91 (s, 1H), 6.52 (s, 1H), 3.95-3.40 (m, 5H), 3.32 (m, 1H), 3.20 (m, 2H), 2.15 (m, 1H), 1.74 (m, 2H), 0.99 (t, J = 7.3 Hz, 3H).

15

EXAMPLE 74

N-[4-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-4-pyrrol-1-yl-benzenesulfonamide

Sulfonamide coupling was achieved by following a procedure analogous to that de-
20 scribed in example 55. 86 mg (yield 66%) of the title compound were obtained.

MS (ESI) m/z: 411.1 [M+H]⁺

¹H-NMR (CH₃OH-d₄): δ [ppm] 7.78 (d, 2H), 7.76 (s, 1H), 7.61 (d, 2H), 7.26 (s, 1H), 7.18 (d, 2H), 7.09 (d, 2H), 6.31 (s, 1H), 3.36 (m, 2H), 3.17 (m, 1H), 3.08 (m, 1H), 2.82 (m, 3H), 2.32 (m, 1H), 1.95 (m, 1H), 1.62 (m, 2H), 0.97 (t, J = 7.4 Hz, 3H).

25

EXAMPLE 75

4-Azetidin-1-yl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

The title compound was prepared in analogy to the procedure described in example 67.
30 9 mg (yield 11%) of the title compound were obtained.

MS (ESI) m/z: 400.2 [M+H]⁺

¹H-NMR (CH₃OH-d₄): δ [ppm] 7.50 (d, 2H), 7.12 (d, 2H), 7.03 (d, 2H), 6.31 (d, 2H), 3.89 (m, 4H), 3.49 (m, 2H), 3.26 (m, 2H), 2.92 (m, 3H), 2.33 (m, 3H), 2.02 (m, 1H), 1.68 (m, 2H), 0.97 (t, J = 7.3 Hz, 3H).

35

EXAMPLE 76

150

4-(2-Oxo-oxazolidin-3-yl)-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

The title compound was prepared in analogy to the procedure described in example 67.

5 7 mg (yield 10%) of the title compound were obtained.

MS (ESI) m/z: 430.2 [M+H]⁺

¹H-NMR (CDCl₃): δ [ppm] 7.74 (d, 2H), 7.68 (d, 2H), 7.14 (d, 2H), 7.07 (d, 2H), 4.50 (m, 2H), 4.10 (m, 2H), 3.50 (m, 4H), 2.96 (m, 3H), 2.37 (m, 1H), 2.01 (m, 1H), 1.68 (m, 2H), 0.98 (t, J = 7.3 Hz, 3H).

10

EXAMPLE 77

4-(2,2-Difluoro-1-methyl-ethyl)-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

15 Sulfonamide coupling was achieved by following a procedure analogous to that described in example 55.2. Purification by chromatography ethyl acetate-CH₂Cl₂ 1:1 gave 36 mg (yield 12%) of the title compound.

MS (ESI) m/z: 423.1 [M+H]⁺

20 ¹H-NMR (CDCl₃): δ [ppm] 7.74 (d, 2H), 7.52 (d, 2H), 7.14 (d, 2H), 7.03 (d, 2H), 6.14 (m, 1H), 3.30 (m, 5H), 2.82 (m, 1H), 2.65 (t, 1H, J = 7.3 Hz, 2H), 2.39 (m, 2H), 2.15 (m, 1H), 1.65 (m, 1H), 1.41 (m, 1H), 1.31 (d, 3H), 0.98 (t, J = 7.3 Hz, 3H).

EXAMPLE 78

25 N-[4-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-4-(2,2,2-trifluoro-1-methyl-ethyl)-benzenesulfonamide

Sulfonamide coupling was achieved by following a procedure analogous to that described in example 55.2. 57 mg (yield 43%) of the title compound were obtained.

MS (ESI) m/z: 441.1 [M+H]⁺

30 ¹H-NMR (CH₃OH-d₄): δ [ppm] 7.72 (d, 2H), 7.49 (d, 2H), 7.12 (d, 2H), 7.03 (d, 2H), 3.65 (m, 2H), 3.24 (m, 1H), 3.10 (m, 1H), 2.90 (m, 1H), 2.70 (m, 1H), 2.51 (m, 3H), 2.25 (m, 1H), 1.82 (m, 1H), 1.57 (m, 2H), 1.47 (d, 3H), 0.94 (t, J = 7.3 Hz, 3H).

EXAMPLE 79

35 Benzofuran-2-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide

151

Sulfonamide coupling was achieved by following a procedure analogous to that described in example 55.2. Chromatography (using ethyl acetat-methanol) 1-50% gave 52 mg (yield 24%) of the title compound.

MS (ESI) m/z : 385.1 $[M+H]^+$

5 $^1\text{H-NMR}$ (DMSO-d_6): δ [ppm] 10.50 (br s, 1H), 7.76 (d, 1H), 7.70 (d, 1H), 7.55 (m, 3H), 7.34 (m, 1H), 7.18 (d, 2H), 7.10 (d, 2H), 3.10 (m, 1H), 2.90 (m, 1H), 2.61 (m, 5H), 2.22 (m, 1H), 1.76 (m, 1H), 1.52 (m, 3H), 0.86 (t, $J = 7.3$ Hz, 3H).

EXAMPLE 80

10 5-Isoxazol-5-yl-thiophene-2-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide

4-((S)-1-Propyl-pyrrolidin-3-yl)-phenylamine (100 mg, 0.49 mmol) was dissolved in tetrahydrofuran (5 ml). Subsequently, dimethylaminopyridine (24 mg, 0.20 mmol) and 5-(5-isoxazolyl)thiophene-2-sulfonyl chloride (159 mg, 0.64 mmol) were added and the
15 reaction mixture stirred overnight at room temperature. The solvent was evaporated under reduced pressure, the residue treated with water and ethyl acetate. The organic layer was dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to give the crude product. The crude product was purified with silica gel chromatography with dichloromethane/methanol (100:0 to 96:4) as eluent, yielding
20 the purified product (67 mg, 33%).

MS (ESI) m/z : 418.1 $[M+H]^+$

$^1\text{H-NMR}$ (DMSO-d_6): δ [ppm] 8.65 (s, 1H), 7.63 (d, 1H), 7.58 (d, 1H), 7.24 (d, 2H), 7.12 (m, 3H), 7.07 (s, 1H), 4.02 (m, 1H), 3.16 (d, 2H), 3.10 (m, 3H), 2.28 (m, 1H), 1.91 (m, 1H), 1.58 (m, 2H), 0.88 (t, $J = 7.3$ Hz, 3H).

25

EXAMPLE 81

5-Isoxazol-3-yl-thiophene-2-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide

Sulfonamide coupling was achieved by following a procedure analogous to that described in example 80. The title compound was obtained in an amount of 66 mg; yield
30 32%.

MS (ESI) m/z : 418.1 $[M+H]^+$

$^1\text{H-NMR}$ (DMSO-d_6): δ [ppm] 8.72 (s, 1H), 7.68 (d, 1H), 7.62 (d, 1H), 7.24 (d, 2H), 7.14 (m, 3H), 7.09 (s, 1H), 4.08 (m, 1H), 3.16 (d, 2H), 3.10 (m, 3H), 2.33 (m, 1H), 1.97 (m,
35 1H), 1.62 (m, 2H), 0.92 (t, $J = 7.3$ Hz, 3H).

EXAMPLE 82

152

5-Oxazol-5-yl-thiophene-2-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide

Sulfonamide coupling was achieved by following a procedure analogous to that described in example 80; title compound was obtained in an amount of 110 mg (yield

5 54%).

MS (ESI) m/z : 418.1 $[M+H]^+$

1H -NMR (CH_3OH-d_4): δ [ppm] 8.70 (s, 1H), 7.68 (d, 1H), 7.62 (d, 1H), 7.36 (d, 2H), 7.14 (m, 3H), 7.09 (s, 1H), 4.05 (m, 1H), 3.18 (d, 2H), 3.04 (m, 3H), 2.32 (m, 1H), 1.93 (m, 1H), 1.59 (m, 2H), 0.91 (t, $J = 7.3$ Hz, 3H).

10

EXAMPLE 83

N-(4-Azetidin-3-yl-phenyl)-4-(2,2-difluoro-1-methyl-ethyl)-benzenesulfonamide

Step 1: Sulfonamide coupling using 4-(1,1-difluoropropan-2-yl)benzene-1-sulfonyl chloride, following a procedure in analogy to that used in example 55. 90 mg (29%) of the

15

title compound were obtained.

MS (ESI) m/z : 467.1 $[M+H]^+$

Step 2: boc deprotection, following a procedure in analogy to that used in example

20

46.4. 77 mg (100%) of the title compound were obtained.

EXAMPLE 84

4-Isopropyl-N-[4-((S)-1-ethyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

25

0.3 g of 4-Isopropyl-N-((S)-4-pyrrolidin-3-yl-phenyl)-benzenesulfonamide (0.87 mmol) were dissolved in 20 ml dichloromethane. After addition of 0.07 ml of acetic acid, 0.073 ml of acetaldehyde (1.31 mmol) and 0.277 g of sodium triacetoxyborohydride (1.31 mmol), the reaction mixture was stirred at room temperature for 30 minutes. The solvent was evaporated, the residue dissolved in water and the pH adjusted to pH 8-9 with aqueous sodium hydroxide solution. The aqueous phase was extracted three times with diethyl ether, the combined organic phases dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure. The crude product was purified via silica gel chromatography (chromabond column) with dichloromethane, dichloromethane/methanol 4% as eluent to yield 64 mg of the title compound.

30

35

ESI-MS: 373.25 $[M+H]^+$

153

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.7 (d, 2H), 7.3 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 6.1 (bs, 1H), 3.35 (m, 1H), 3.1 (m, 1H), 2.95 (m, 2H), 2.65 (m, 2H), 2.55 (m, 1H), 2.45 (m, 1H), 2.3 (m, 1H), 1.85 (m, 1H), 1.25 (d, 6H), 1.15 (t, 3H).

5 EXAMPLE 85

4-Isopropyl-N-[4-((S)-1-methyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

4-Isopropyl-N-[4-((S)-1-methyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide was prepared as described for example 84, but using aqueous formaldehyde solution as
10 source for the carbonyl reagent.

ESI-MS: 359.2 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.7 (d, 2H), 7.3 (d, 2H), 7.1 (d, 2H), 7.0 (d, 2H), 6.0 (s, broad, 1H), 3.3 (m, 1H), 3.0 (m, 1H), 2.95 (m, 1H), 2.85 (m, 1H), 2.7 (m, 1H), 2.5 (m, 1H), 2.4 (s, 3H), 2.3 (m, 1H), 1.8 (m, 1H), 1.2 (m, 6H).

15

According to the examples described above, the examples 86 to 93 were prepared. The compounds are characterised by the following physical data.

EXAMPLE 86

20 4-(2-Fluoro-ethyl)-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide, hydrochloride

ESI-MS: 391.4 [M+H]⁺

¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 11.35 (bs, 1H), 10.35 (m, 1H), 7.75 (d, 2H),
25 7.45 (d, 2H), 7.35 (m, 2H), 7.1 (m, 2H), 4.7 (m, 1H), 4.6 (m, 1H), 3.8-3.2 (m, 5H), 3.15-2.95 (m, 4H), 2.3 (m, 1H), 2.0 (m, 1H), 1.7 (m, 2H), 0.9 (t, 3H).

EXAMPLE 87

30 4-((S)-2-Fluoro-1-methyl-ethyl)-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

ESI-MS: 405.15 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.75 (d, 2H), 7.3 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H),
5.35 (bs, 1H), 4.5 (m, 1H), 4.4 (m, 1H), 3.3 (m, 1H), 3.15 (m, 1H), 3.05 (m, 1H), 2.85 (m,
35 1H), 2.65 (m, 1H), 2.5 (m, 1H), 2.45 (m, 2H), 2.3 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 1.3 (d, 3H), 0.9 (t, 3H).

EXAMPLE 88

N-[4-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-4-(3,3,3-trifluoro-propyl)-benzenesulfonamide

ESI-MS: 441.1 [M+H]⁺

- 5 ¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.75 (d, 2H), 7.25 (d, 2H), 7.1 (d, 2H), 7.0 (d, 2H),
5.7 (bs, 1H), 3.3 (m, 1H), 3.05 (m, 1H), 2.9 (m, 3H), 2.7 (m, 1H), 2.2-2.6 (several m,
6H), 1.8 (m, 1H), 1.55 (m, 2H), 0.9 (t, 3H).

EXAMPLE 89

- 10 5-Propyl-thiophene-2-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide

ESI-MS: 393.1 [M+H]⁺

- 15 ¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.3 (d, 1H), 7.2 (d, 2H), 7.05 (d, 2H), 6.65 (d, 1H),
3.35 (m, 1H), 3.1 (m, 1H), 2.9 (m, 1H), 2.75 (m, 3H), 2.5 (m, 3H), 2.3 (m, 1H), 1.85 (m,
1H), 1.5-1.7 (m, 4H), 0.9 (m, 6H).

EXAMPLE 90

N-[4-((R)-1-Allyl-pyrrolidin-3-yl)-phenyl]-4-(2,2-difluoro-cyclopropyl)-
benzenesulfonamide

20

ESI-MS: 419.1 [M+H]⁺

- 25 ¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.65 (d, 2H), 7.2 (d, 2H), 7.05 (d, 2H), 6.95 (d, 2H),
6.0 (bs, 1H), 5.8 (m, 1H), 5.15 (d, 1H), 5.05 (d, 1H), 3.25 (m, 1H), 3.1 (m, 2H), 2.95 (m,
1H), 2.8 (m, 1H), 2.7 (m, 1H), 2.6 (m, 1H), 2.4 (m, 1H), 2.2 (m, 1H), 1.8 (m, 1H), 1.7 (m,
1H), 1.55 (m, 1H).

EXAMPLE 91

5-Methyl-pyridine-2-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide

- 30 ESI-MS: 360.1 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.5 (s, 1H), 7.9 (bs, 1H), 7.8 (d, 1H), 7.6 (d, 1H),
7.15 (d, 2H), 7.1 (d, 2H), 3.25 (m, 1H), 3.0 (m, 1H), 2.85 (m, 1H), 2.6 (m, 1H), 2.5
(m, 1H), 2.4 (m, 2H), 2.4 (s, 3H), 2.2 (m, 1H), 1.75 (m, 1H), 1.5 (m, 2H), 0.9 (t, 3H).

- 35 EXAMPLE 92

4-(3-Fluoro-propyl)-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

ESI-MS: 405.1 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.7 (d, 2H), 7.25 (d, 2H), 7.1 (d, 2H), 7.0 (d, 2H), 6.3 (bs, 1H), 4.4 (dt, 2H), 3.3 (m, 1H), 3.0 (m, 1H), 2.85 (m, 1H), 2.8 (m, 2H), 2.7 (m, 1H), 2.55 (m, 1H), 2.45 (m, 2H), 2.3 (m, 1H), 2.0 (m, 2H), 1.8 (m, 1H), 1.55 (m, 2H), 0.9 (t, 3H).

EXAMPLE 93

4-((R)-2-Fluoro-1-methyl-ethyl)-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

10

ESI-MS: 405.15 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.75 (d, 2H), 7.3 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 5.5 (bs, 1H), 4.5 (m, 1H), 4.4 (m, 1H), 3.3 (m, 1H), 3.15 (m, 1H), 3.05 (m, 1H), 2.85 (m, 1H), 2.65 (m, 1H), 2.5 (m, 1H), 2.45 (m, 2H), 2.3 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 1.3 (d, 3H), 0.9 (t, 3H).

15

EXAMPLE 94

N-[4-((S)-1-Cyclopropylmethyl-pyrrolidin-3-yl)-phenyl]-4-isopropyl-benzenesulfonamide

20 N-[4-((S)-1-Cyclopropylmethyl-pyrrolidin-3-yl)-phenyl]-4-isopropyl-benzenesulfonamide was prepared as described for 4-isopropyl-N-[4-((S)-1-ethyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide in example 84, but using cyclopropylcarbaldehyde as carbonyl reagent.

ESI-MS: 399.1 [M+H]⁺

25 ¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.65 (d, 2H), 7.25 (d, 2H), 7.1 (d, 2H), 6.95 (d, 2H), 5.25 (s, 1H), 3.3 (m, 1H), 3.15 (m, 1H), 2.9 (m, 2H), 2.65 (m, 1H), 2.6 (m, 1H), 2.4 (m, 1H), 2.35 (m, 2H), 2.25 (m, 1H), 1.75 (m, 1H), 1.2 (d, 6H), 0.9 (m, 1H), 0.45 (d, 2H), 0.1 (d, 2H).

30 According to the examples described above, the examples 95 to 119 were prepared. The compounds are characterised by the following physical data.

EXAMPLE 95

35 4-(2-Fluoro-1-fluoromethyl-ethyl)-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

ESI-MS: 423.15 [M+H]⁺

156

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.75 (d, 2H), 7.35 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 5.3 (s, 1H), 4.75 (d, 2H), 4.65 (d, 2H), 3.3 (m, 1H), 3.05 (m, 1H), 2.85 (m, 2H), 2.65 (m, 1H), 2.45 (m, 4H), 2.3 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 0.9 (t, 3H).

5 EXAMPLE 96

N-{4-[(S)-1-(2-Fluoro-ethyl)-pyrrolidin-3-yl]-phenyl}-4-isopropyl-benzenesulfonamide

ESI-MS: 391.1 [M+H]⁺

10 ¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.7 (d, 2H), 7.3 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 4.63 (t, 1H), 4.5 (t, 1H), 3.3 (m, 1H), 3.1 (m, 1H), 2.7-3.0 (several m, 5H), 2.5 (m, 1H), 2.25 (m, 1H), 1.8 (m, 1H), 1.2 (d, 6H).

EXAMPLE 97

15 4-Isopropyl-N-[4-((S)-1-propionyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

ESI-MS: 401.15 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.7 (d, 2H), 7.3 (m, 3H), 7.1 (m, 3H), 4.0 (m, 0.5H), 3.8 (m, 2H), 3.65 (m, 0.5H), 3.5 (m, 1H), 3.35 (m, 2H), 2.95 (sept, 1H), 2.8 (m, 3H), 1.95 (m, 1H), 1.2 (d, 6H), 1.15 (m, 3H).

20

EXAMPLE 98

N-[4-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-3-trifluoromethoxy-benzenesulfonamide

ESI-MS: 429.15 [M+H]⁺

25 ¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.7 (d, 1H), 7.55 (s, 1H), 7.5 (t, 1H), 7.4 (d, 1H), 7.15 (d, 2H), 6.95 (d, 2H), 5.3 (bs, 1H), 3.3 (m, 1H), 3.05 (m, 1H), 2.85 (m, 1H), 2.65 (m, 1H), 2.5 (m, 1H), 2.45 (m, 2H), 2.3 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 0.9 (t, 3H).

EXAMPLE 99

30 N-[2-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-3-trifluoromethoxy-benzenesulfonamide

ESI-MS: 429.15 [M+H]⁺

EXAMPLE 100

35 N-[4-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-2-trifluoromethyl-benzenesulfonamide

ESI-MS: 413.15 [M+H]⁺

157

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.0 (d, 1H), 7.85 (d, 1H), 7.65 (t, 1H), 7.55 (t, 1H), 7.1 (d, 2H), 6.95 (d, 2H), 5.9 (bs, 1H), 3.3 (m, 1H), 3.0 (m, 1H), 2.8 (m, 1H), 2.6 (m, 1H), 2.45 (m, 1H), 2.4 (m, 2H), 2.25 (m, 1H), 1.75 (m, 1H), 1.55 (m, 2H), 0.95 (t, 3H).

5 EXAMPLE 101

N-[4-((S)-1-Allyl-pyrrolidin-3-yl)-phenyl]-4-trifluoromethoxy-benzenesulfonamide

ESI-MS: 427.1 [M+H]⁺

10 ¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 11.3 (bd, 1H), 10.45 (m, 1H), 7.9 (d, 2H), 7.55 (d, 2H), 7.25 (m, 2H), 7.1 (d, 2H), 6.0 (m, 1H), 5.5 (m, 1H), 5.45 (m, 1H), 3.8 (m, 2H), 2.9-3.75 (several m, 5H), 2.3 (m, 1H), 1.95 (m, 1H).

EXAMPLE 102

N-((S)-4-Pyrrolidin-3-yl-phenyl)-4-trifluoromethoxy-benzenesulfonamide

15

ESI-MS: 387.05 [M+H]⁺

1H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 7.85 (d, 2H), 7.45 (d, 2H), 7.05 (d, 2H), 6.95 (d, 2H), 5.2 (m, 1H), 3.3 (m, 1H), 3.1 (m, 2H), 3.05 (m, 1H), 2.7 (m, 1H), 2.15 (m, 1H), 1.7 (m, 1H)

20

EXAMPLE 103

N-[4-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-4-(2,2,2-trifluoro-ethyl)-benzenesulfonamide

25 ESI-MS: 427.2 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.8 (d, 2H), 7.4 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 5.7 (bs, 1H), 3.4 (m, 2H), 3.3 (m, 1H), 3.05 (m, 1H), 2.85 (m, 1H), 2.7 (m, 1H), 2.5 (m, 3H), 2.3 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 0.9 (t, 3H).

30 EXAMPLE 104

N-[4-((S)-1-Ethyl-pyrrolidin-3-yl)-phenyl]-4-trifluoromethoxy-benzenesulfonamide

ESI-MS: 415.1 [M+H]⁺

35 ¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.8 (d, 2H), 7.25 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 4.9 (bs, 1H), 3.3 (m, 1H), 3.05 (m, 1H), 2.85 (m, 1H), 2.5-2.7 (several m, 3H), 2.45 (m, 1H), 2.3 (m, 1H), 1.8 (m, 1H), 1.15 (t, 3H).

EXAMPLE 105

5-Chloro-thiophene-2-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide

ESI-MS: 385.0 [M+H]⁺

- 5 ¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.25 (d, 1H), 7.2 (d, 2H), 7.0 (d, 2H), 6.8 (d, 1H), 5.1 (bs, 1H), 3.3 (m, 1H), 3.05 (m, 1H), 2.85 (m, 1H), 2.7 (m, 1H), 2.4-2.6 (several m, 3H), 2.3 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 0.9 (t, 3H).

EXAMPLE 106

- 10 N-[4-((S)-1-Allyl-pyrrolidin-3-yl)-phenyl]-4-((S)-2-fluoro-1-methyl-ethyl)-benzenesulfonamide

ESI-MS: 403.4 [M+H]⁺

- 15 EXAMPLE 107

N-[4-((S)-1-Allyl-pyrrolidin-3-yl)-phenyl]-4-(3-fluoro-propyl)-benzene sulfonamide

ESI-MS: 403.15 [M+H]⁺

20

EXAMPLE 108

4-(3-Fluoro-propyl)-N-((S)-4-pyrrolidin-3-yl-phenyl)-benzenesulfonamide

ESI-MS: 363.1 [M+H]⁺

- 25 ¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 7.7 (d, 2H), 7.4 (d, 2H), 7.1 (d, 2H), 7.0 (d, 2H), 4.6 (bs), 4.5 (m, 1H), 4.35 (m, 1H), 3.15 (m, 1H), 3.0 (m, 2H), 2.9 (m, 1H), 2.7 (m, 2H), 2.6 (m, 1H), 2.05 (m, 1H), 1.95 (m, 1H), 1.9 (m, 1H), 1.6 (m, 1H).

EXAMPLE 109

- 30 4-((S)-2-Fluoro-1-methyl-ethyl)-N-((S)-4-pyrrolidin-3-yl-phenyl)-benzenesulfonamide

ESI-MS: 363.05 [M+H]⁺

- 35 ¹H-NMR (CH₃OH-d₄, 400 MHz): δ [ppm] 7.7 (d, 2H), 7.4 (d, 2H), 7.1 (d, 2H), 7.0 (d, 2H), 4.5 (m, 1H), 4.4 (m, 1H), 2.95-3.35 (several m, 6H), 2.7 (m, 1H), 2.2 (m, 1H), 1.3 (d, 3H).

EXAMPLE 110

159

4-((R)-2-Fluoro-1-methyl-ethyl)-N-((S)-4-pyrrolidin-3-yl-phenyl)-benzenesulfonamide

ESI-MS: 363.1 [M+H]⁺

¹H-NMR (acetic acid-d₄, 400 MHz): δ [ppm] 7.7 (d, 2H), 7.4 (d, 2H), 7.2 (m, 4H), 4.55 (m, 1H), 4.45 (m, 1H), 3.8 (m, 1H), 3.65 (m, 1H), 3.5 (m, 2H), 3.2 (m, 2H), 2.4 (m, 1H), 2.05 (m, 1H), 1.3 (d, 3H).

EXAMPLE 111

N-[4-((S)-1-Methyl-pyrrolidin-3-yl)-phenyl]-4-trifluoromethoxy-benzenesulfonamide

10

ESI-MS: 401.1 [M+H]⁺

¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 7.9 (d, 2H), 7.55 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 3.2 (m, 1H), 2.8 (m, 1H), 2.6 (m, 2H), 2.3 (m, 1H), 2.25 (s, 3H), 2.15 (m, 1H), 1.65 (m, 1H).

15

EXAMPLE 112

N-[4-((R)-1-Allyl-pyrrolidin-3-yl)-phenyl]-4-(2,2,2-trifluoro-1-methyl-ethyl)-benzenesulfonamide

20

ESI-MS: 439.1 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ 7.8 (d, 2H), 7.4 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 5.9 (m, 1H), 5.2 (d, 1H), 5.1 (d, 1H), 3.45 (m, 1H), 3.3 (m, 1H), 3.2 (m, 2H), 3.1 (m, 1H), 2.9 (m, 1H), 2.7 (m, 1H), 2.5 (m, 1H), 2.3 (m, 1H), 1.8 (m, 1H), 1.5 (d, 3H).

25

EXAMPLE 113

N-[4-((S)-1-Allyl-pyrrolidin-3-yl)-phenyl]-4-((R)-2-fluoro-1-methyl-ethyl)-benzenesulfonamide

ESI-MS: 403.1 [M+H]⁺

30

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.75 (d, 2H), 7.3 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 5.9 (m, 1H), 5.2 (d, 1H), 5.1 (d, 1H), 4.5 (m, 1H), 4.4 (m, 1H), 3.3 (m, 1H), 3.1-3.2 (m, 3H), 3.0 (m, 1H), 2.8 (m, 1H), 2.65 (m, 1H), 2.4 (m, 1H), 2.25 (m, 1H), 1.8 (m, 1H), 1.3 (d, 3H).

35

EXAMPLE 114

N-{4-[(S)-1-(3-Fluoro-propyl)-pyrrolidin-3-yl]-phenyl}-4-trifluoromethoxy-benzenesulfonamide

ESI-MS: 447.1 [M+H]⁺

¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 10.25 (bs, 1H), 7.85 (d, 2H), 7.55 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 4.55 (t, 1H), 4.4 (t, 1H), 3.2 (m, 1H), 2.85 (m, 1H), 2.6 (m, 2H), 2.5 (m, 2H), 2.35 (m, 1H), 2.15 (m, 1H), 1.8 (m, 1H), 1.75 (m, 1H), 1.65 (m, 1H).

EXAMPLE 115

4-Methanesulfonyl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

10 ESI-MS: 423.1 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.95 (m, 4H), 7.1 (d, 2H), 7.0 (d, 2H), 6.2 (bs, 1H), 3.3 (m, 1H), 3.1 (m, 1H), 3.05 (s, 3H), 2.9 (m, 1H), 2.75 (m, 1H), 2.5 (m, 3H), 2.3 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 0.9 (t, 3H).

15 EXAMPLE 116

N-[4-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-4-ureido-benzenesulfonamide

ESI-MS: 403.1 [M+H]⁺

20 EXAMPLE 117

4-Cyano-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

ESI-MS: 370.1 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.85 (d, 2H), 7.7 (d, 2H), 7.1 (d, 2H), 6.95 (d, 2H), 6.0 (bs, 1H), 3.3 (m, 1H), 3.1 (m, 1H), 2.85 (m, 1H), 2.75 (m, 1H), 2.5 (m, 3H), 2.3 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 0.9 (t, 3H).

EXAMPLE 118

4-(1-Methyl-1H-pyrazol-4-yl)-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

ESI-MS: 425.1 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.75 (s, 1H), 7.7 (d, 2H), 7.65 (s, 1H), 7.45 (d, 2H), 7.1 (d, 2H), 7.0 (d, 2H), 3.9 (s, 3H), 3.3 (m, 1H), 3.05 (m, 1H), 2.85 (m, 1H), 2.7 (m, 1H), 2.5 (m, 3H), 2.3 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 0.9 (t, 3H).

EXAMPLE 119

161

1-Ethyl-1H-pyrazole-4-sulfonic acid [4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-amide hydrochloride

ESI-MS: 363.1 [M+H]⁺

- 5 ¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.65 (s, 1H), 7.6 (s, 1H), 7.15 (d, 2H), 7.0 (d, 2H), 4.25 (bs, 1H), 4.1 (q, 2H), 3.3 (m, 1H), 3.0 (m, 1H), 2.8 (m, 1H), 2.65 (m, 1H), 2.4-2.55 (m, 3H), 2.3 (m, 1H), 1.8 (m, 1H), 1.55 (m, 2H), 1.4 (t, 3H), 0.9 (t, 3H).

EXAMPLE 120

- 10 4-Morpholin-4-yl-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

- A solution of 0.07 g of rac-2,2'-bis(diphenylphosphino)-1,1'-binaphthalene (BINAP, 0.11 mmol) and 0.043 g tris(dibenzylidenacetone) dipalladium(0) (0.05 mmol) in 5ml of tetrahydrofuran were added dropwise to a solution of 0.25 g of 4-bromo-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide (from example 25 (0.59 mmol)), 0.078 ml morpholine (0.9 mmol), and 0.104 g of sodium tert.-butylate (1.08 mmol) in 20 ml tetrahydrofuran. The reaction mixture was refluxed for 5 ½ h, and, after additional addition of 0.04 ml of morpholine, for another 2 h. After evaporation, the residue was treated with water, extracted twice with diethyl ether and dichloromethane each, and the combined organic layers dried over magnesium sulfate, filtered, and the solvent evaporated. The thus obtained crude product was purified via silica gel chromatography using a gradient of dichloromethane/methanol 0-12%.

- ESI-MS: 430.2 [M+H]⁺
- 25 ¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 11.25 (d broad, 1H), 10.05 (m, 1H), 9.4 (bs, 1H), 7.6 (d, 2H), 7.25 (d, 1H), 7.2 (d, 1H), 7.05 (m, 2H), 7.0 (d, 2H), 3.8 (m, 2H), 3.5-3.8 (m, 5H), 3.2 (m, 4H), 3.0-3.1 (m, 4H), 2.3 (m, 1H), 1.9 (m, 1H), 1.7 (m, 2H), 0.9 (t, 3H).

- According to the examples described above, the examples 121 to 132 were prepared.
- 30 The compounds are characterised by the following physical data.

EXAMPLE 121

4-Benzyloxy-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

- 35 ESI-MS: 451.1 [M+H]⁺

EXAMPLE 122

162

4-Hydroxy-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

ESI-MS: 361.1 [M+H]⁺

¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 10.0 (bs, 1H), 7.55 (d, 2H), 7.1 (d, 2H), 7.0 (d, 2H), 6.85 (d, 2H), 3.15 (m, 1H), 2.85 (m, 1H), 2.6 (m, 2H), 2.3-2.45 (m, 3H), 2.15 (m, 1H), 1.65 (m, 1H), 1.4 (m, 2H), 0.85 (t, 3H).

EXAMPLE 123

N-[4-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-4-vinyl-benzenesulfonamide

ESI-MS: 371.1 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.7 (d, 2H), 7.4 (d, 2H), 7.15 (d, 2H), 7.0 (d, 2H), 6.7 (q, 1H), 5.8 (d, 1H), 5.4 (d, 1H), 4.25 (bs, 1H), 3.3 (m, 1H), 3.0 (m, 1H), 2.8 (m, 1H), 2.6 (m, 1H), 2.35-2.5 (m, 3H), 2.25 (m, 1H), 1.8 (m, 1H), 1.5 (m, 2H), 0.9 (t, 3H).

EXAMPLE 124

N-[2-((S)-1-Propyl-pyrrolidin-3-yl)-phenyl]-4-vinyl-benzenesulfonamide

ESI-MS: 371.1 [M+H]⁺

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.75 (d, 2H), 7.4 (d, 2H), 7.05-7.2 (m, 3H), 6.95 (d, 1H), 6.65 (q, 1H), 5.8 (d, 1H), 5.35 (d, 1H), 5.0 (bs), 3.45 (m, 1H), 3.3 (m, 1H), 3.15 (m, 1H), 3.1 (m, 1H), 2.65-2.8 (m, 3H), 2.3 (m, 1H), 1.9 (m, 1H), 1.7 (m, 2H), 0.9 (t, 3H).

EXAMPLE 125

4-Fluoro-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

ESI-MS: 363.1 [M+H]⁺

¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 11.25 (bd, 1H), 10.4 (m, 1H), 7.8 (m, 2H), 7.4 (m, 2H), 7.3 (d, 1H), 7.25 (d, 1H), 7.1 (m, 2H), 3.2-3.8 (m, 5H), 2.9-3.1 (m, 2H), 2.3 (m, 1H), 1.95 (m, 1H), 1.7 (m, 2H), 0.9 (t, 3H).

EXAMPLE 126

3,4-Difluoro-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide hydrochloride

ESI-MS: 381.2 [M+H]⁺

163

¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 11.25 (bd, 1H), 10.55 (m, 1H), 7.85 (t, 1H), 7.65 (m, 2H), 7.3 (d, 1H), 7.25 (d, 1H), 7.1 (m, 2H), 3.2-3.8 (m, 5H), 2.9-3.15 (m, 2H), 2.3 (m, 1H), 1.95 (m, 1H), 1.7 (m, 2H), 0.9 (t, 3H).

5 EXAMPLE 127

4-(3,3-Difluoro-pyrrolidin-1-yl)-N-[4-((S)-1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide

ESI-MS: 450.1 [M+H]⁺

10

EXAMPLE 128

N-[4-(1-Benzyl-pyrrolidin-3-yl)-3-fluoro-phenyl]-4-isopropyl-benzenesulfonamide

128.1 2-Fluoro-4-nitro-1-vinyl-benzene

15

¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.0 (m, 1H), 7.95 (m, 1H), 7.65 (m, 1H), 6.9 (dd 1H), 6.0 (m, 1H), 5.6 (m, 1H).

128.2 1-Benzyl-3-(2-fluoro-4-nitro-phenyl)-pyrrolidine

20

ESI-MS: 301.1 [M+H]⁺

128.3 1-Benzyl-3-(2-fluoro-4-amino-phenyl)-pyrrolidine

25

ESI-MS: 271.1 [M+H]⁺

128.4 N-[4-(1-Benzyl-pyrrolidin-3-yl)-3-fluoro-phenyl]-4-isopropyl-benzenesulfonamide

ESI-MS: 453.15 [M+H]⁺

30

According to the examples described above, the examples 129 to 131 were prepared. The compounds are characterised by the following physical data.

EXAMPLE 12935

N-(3-Fluoro-4-pyrrolidin-3-yl-phenyl)-4-isopropyl-benzenesulfonamide

ESI-MS: 363.15 [M+H]⁺

EXAMPLE 130

N-[3-Fluoro-4-(1-propyl-pyrrolidin-3-yl)-phenyl]-4-isopropyl-benzenesulfonamide, hydrochloride

5

ESI-MS: 405.15 [M+H]⁺

¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 11.2 (bs, 1H), 10.65 (m, 1H), 7.75 (d, 2H), 7.45 (m, 3H), 6.95 (m, 2H), 3.85-3.2 (m, 4H), 3.15-2.9 (m, 4H), 2.3 (m, 1H), 2.05 (m, 1H), 1.7 (m, 2H), 1.2 (d, 6H), 0.9 (t, 3H).

10

EXAMPLES 131

(-)-N-[6-(1-Propyl-pyrrolidin-3-yl)-pyridin-3-yl]-4-isopropyl-benzene-sulfonamide

15

The racemic compound, N-[6-(1-propyl-pyrrolidin-3-yl)-pyridin-3-yl]-4-isopropyl-benzene-sulfonamide, was subjected to a chiral chromatography on a preparative CHIRACEL AD column using n-hexane/ethanol/triethylamine (85:15:1) as eluent. Fractions containing only the desired enantiomer are analysed by analytical chiral HPLC and were combined.

20

ESI-MS: 388.1 [M+H]⁺

[α]_D: -18.1°

EXAMPLE 132

25

(+)-N-[6-(1-Propyl-pyrrolidin-3-yl)-pyridin-3-yl]-4-isopropyl-benzene-sulfonamide, hydrochloride

30

The racemic compound, N-[6-(1-propyl-pyrrolidin-3-yl)-pyridin-3-yl]-4-isopropyl-benzene-sulfonamide, was subjected to a chiral chromatography on a preparative CHIRACEL AD column using n-hexane/ethanol/triethylamine (85:15:1) as eluent. Fractions containing only the desired enantiomer are analysed by analytical chiral HPLC and were combined.

35

ESI-MS: 388.1 [M+H]⁺

[α]_D: +17.2°

EXAMPLE 133

N-[4-((S)-1-Propyl-piperidin-3-yl)-phenyl]-4-trifluoromethoxy-benzenesulfonamide

133.1 (S)-3-Phenyl-1-propyl-piperidine

1 g (6.2 mmol) of commercially available (S)-3-phenylpiperidine was dissolved in di-
5 chloromethane and 0.37 ml acetic acid (6.51 mmol). After addition of 0.5 ml of propanal
(6.93 mmol) and 1.97 g of sodium triacetoxyborohydride (9.3 mmol), the reaction mix-
ture was stirred at room temperature for 18 h. Water was added, the layers separated,
and the aqueous phase extracted with dichloromethane. The combined organic layers
were dried over magnesium sulfate, filtered, and evaporated to dryness to yield 1.1 g of
10 (S)-3-phenyl-1-propyl-piperidine.
ESI-MS: 204.1 [M+H]⁺

133.2 (S)-3-(4-Nitro-phenyl)-1-propyl-piperidine

15 ESI-MS: 249.1 [M+H]⁺
¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 8.15 (d, 2H), 7.4 (d, 2H), 2.95 (m, 4H), 2.3 (m, 2H),
1.9-2.1 (m, 3H), 1.65-1.85 (m, 2H), 1.4-1.6 (m, 2H), 0.9 (t, 3H).

133.3 (S)-3-(4-Amino-phenyl)-1-propyl-piperidine

20 ESI-MS: 219.1 [M+H]⁺

133.4 N-[4-((S)-1-Propyl-piperidin-3-yl)-phenyl]-4-trifluoromethoxy-
benzenesulfonamide

25 Following a procedure analogous to that described in example 55.2 sulfonamide cou-
pling was achieved
ESI-MS: 443.1 [M+H]⁺
¹H-NMR (CDCl₃, 400 MHz): δ [ppm] 7.8 (d, 2H), 7.2 (d, 2H), 7.1 (d, 2H), 7.0 (d, 2H), 6.7
30 (bs, 1H), 3.0 (m, 2H), 2.8 (m, 1H), 2.35 (m, 2H), 1.95 (m, 2H), 1.85 (m, 1H), 1.75 (m,
2H), 1.5 (m, 2H), 1.4 (m, 1H), 0.9 (t, 3H).

EXAMPLE 134

N-[2-Fluoro-4-(1-propyl-pyrrolidin-3-yl)-phenyl]-4-isopropyl-benzenesulfonamide,
35 hydrochloride

134.1 2-Fluoro-1-nitro-4-vinyl-benzene

4-Bromo-2-fluoro-1-nitro-benzene (691 mg, 3.14 mmol), tributyl-vinyl-stannane (1.2 g, 3.77 mmol), triphenylphosphine (49 mg, 0.19 mmol) and tetrakis(triphenyl)phosphinepalladium(0) (73 mg, 0.06 mmol) were dissolved in toluene (25 ml) and stirred for 5 hours at reflux. The reaction mixture was concentrated in vacuo. The residue was partitioned between water (25 ml) and diethyl ether (50 ml). The organic phase was separated, dried over MgSO₄, filtered and concentrated in vacuo to yield an oil (1.8 g). The residue was purified via silica gel chromatography with cyclohexane/ethyl acetate (gradient 0 – 5 %). Fractions containing the product were combined and the solvent evaporated to give an oil (360 mg, 69 %).

134.2 1-Benzyl-3-(3-fluoro-4-nitro-phenyl)-pyrrolidine

2-Fluoro-1-nitro-4-vinyl-benzene (360 mg, 2.15 mmol) was dissolved in dichloromethane (2 ml), trifluoroacetic acid (70 µl, 0.88 mmol) was added, followed by slow addition of N-benzyl-N-(methoxymethyl)-N-trimethylsilylmethylamine (633 mg, 2.67 mmol). Stirring was continued for 2 h at room temperature. Another portion of N-benzyl-N-(methoxymethyl)-N-trimethylsilylmethylamine (300 mg, 1.26 mmol) was added and stirring continued for another 30 minutes. The reaction mixture was diluted with ethyl acetate (25 ml), washed with aqueous NaHCO₃ (15 ml). The organic layer was dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to yield an oil (900 mg). The crude product was purified via silica gel chromatography with cyclohexane/ethyl acetate (gradient 0 – 25 %). Fractions containing the product were combined, the solvent evaporated to yield an oil (550 mg, 85 %).

ESI-MS: 301.1 [M+H]⁺

134.3 4-(1-Benzyl-pyrrolidin-3-yl)-2-fluoro-phenylamine

1-Benzyl-3-(3-fluoro-4-nitro-phenyl)-pyrrolidine (550 mg, 1.83 mmol) was dissolved in methanol (30 ml), tin dichloride (3.125 g, 16.48 mmol) was added, and the reaction mixture stirred at reflux for 2 h. The methanol was evaporated, 1 N sodium hydroxide (60 ml) and ethyl acetate were added and stirring continued. The tin salts were filtered off, the organic phase was separated, dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to yield an oil (400 mg).

ESI-MS: 271.1 [M+H]⁺

134.4 N-[4-(1-Benzyl-pyrrolidin-3-yl)-2-fluoro-phenyl]-4-isopropyl-benzenesulfonamide

4-(1-Benzyl-pyrrolidin-3-yl)-2-fluoro-phenylamine (400 mg, 1.48 mmol) and 4-isopropyl-phenylsulfonyl chloride (324 mg, 1.48 mmol) were dissolved in tetrahydrofuran (25 ml). Triethylamine (0.62 ml, 4.44 mmol) was added and the reaction mixture stirred over
5 night at room temperature (10 % conversion) and subsequently refluxed for 4 h (30 % conversion). Small portions of 4-isopropyl-phenylsulfonyl chloride and triethylamine were added and the reaction mixture was stirred 15 min at 150 °C in the microwave (CEM). This procedure was repeated until complete conversion was observed. The solvent was evaporated under reduced pressure, the residue treated with ethyl acetate
10 (50 ml) and extracted twice with slightly acidic water. The organic layer was dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to give an oil (170 mg, 21 %).
ESI-MS: 453.1 [M+H]⁺

15 134.5 N-(2-Fluoro-4-pyrrolidin-3-yl-phenyl)-4-isopropyl-benzenesulfonamide

A mixture of N-[4-(1-benzyl-pyrrolidin-3-yl)-2-fluoro-phenyl]-4-isopropyl-benzenesulfonamide (170 mg, 0.31 mmol) and 10 % palladium on carbon (20 mg) in ethyl acetate (25 ml) and acetic acid (10 ml) was hydrogenated over night (20 % con-
20 version). The reaction mixture was irradiated with an infra-red lamp for 3 h (complete conversion). The catalyst was filtered, and the solvent was removed under vacuum to yield an oil (45 mg, 64 % purity, 26 %).
ESI-MS: 363.1 [M+H]⁺

25 134.6 N-[2-Fluoro-4-(1-propyl-pyrrolidin-3-yl)-phenyl]-4-isopropyl-benzenesulfonamide, hydrochloride

N-(2-Fluoro-4-pyrrolidin-3-yl-phenyl)-4-isopropyl-benzenesulfonamide (45 mg, 0.08 mmol) and propionaldehyde (4.7 mg, 0.08 mmol) were dissolved in tetrahydrofuran (5
30 ml). Acetic acid (10 µl, 0.12 mmol) and sodium trisacetoxyborohydride (34 mg, 0.16 mmol) were sequentially added to the reaction mixture and stirred for 30 minutes at room temperature. The reaction mixture was concentrated and the residue was dissolved in aqueous NaHCO₃ (10 ml) and extracted with diethyl ether (30 ml). The organic phase was separated, dried over magnesium sulfate, filtered, and evaporated to
35 dryness to yield an oil (45 mg). The residue was dissolved in diethyl ether (25 ml) and HCl in ether was added and stirred overnight. The solvent was decanted and to the

residue was added 3 ml of H₂O. This solution was lyophilised to yield the pure product (32 mg, 91 %).

ESI-MS: 405.1 [M+H]⁺

¹H-NMR (CH₃OH-d₄): δ [ppm] 7.7 (d, 2H), 7.4 (m, 3H), 7.0 (m, 2H), 3.4 (m, 2H), 3.2-2.9 (m, 4H), 2.8 (m, 2H), 2.4 (m, 1H), 2.0 (m, 1H), 1.6 (m, 2H), 1.2 (d, 6H), 1.0 (t, 3H).

EXAMPLE 135

4-Isopropyl-N-(2-methoxy-4-pyrrolidin-3-yl-phenyl)-benzenesulfonamide, acetate

10 135.1 4-Bromo-2-methoxy-1-nitro-benzene

To a solution of 4-bromo-2-fluoro-1-nitro-benzene (2.0 g, 9.09 mmol) in methanol (50 ml) was added sodium methanolate (30 % in methanol) (1.64 g, 9.09 mmol). The reaction mixture was stirred at room temperature overnight. The reaction mixture was concentrated and the residue was dissolved in water (30 ml) and extracted twice with ethyl acetate. The combined organic phases were washed with water. The organic phase was separated, dried over magnesium sulfate, filtered, and evaporated to dryness to yield a crystalline solid (2.1 g, 99 %).

¹H-NMR (DMSO-d₆): δ [ppm] 7.9 (d, 1H), 7.6 (s, 1H), 7.3 (d, 1H), 4.0 (s, 3H).

20

135.2 2-Methoxy-1-nitro-4-vinyl-benzene

4-Bromo-2-methoxy-1-nitro-benzene (691 mg, 3.14 mmol), tributyl-vinyl-stannane (1.2 g, 3.77 mmol), triphenylphosphine (49 mg, 0.19 mmol) and tetrakis(triphenylphosphine)ladium(0) (73 mg, 0.06 mmol) were dissolved in toluene (25 ml) and stirred for 5 hours at reflux. The reaction mixture was concentrated in vacuo. The residue was partitioned between water (25 ml) and diethyl ether (50 ml). The organic phase was separated, dried over MgSO₄, filtered and concentrated in vacuo to yield an oil (2.0 g). The residue was purified via silica gel chromatography with cyclohexane/ethyl acetate (gradient 0 – 25 %). Fractions containing the product were combined, the solvent evaporated to give an oil (435 mg, 72 %).

30

135.3 1-Benzyl-3-(3-methoxy-4-nitro-phenyl)-pyrrolidine

2-Methoxy-1-nitro-4-vinyl-benzene (435 mg, 2.43 mmol) was dissolved in dichloromethane (2 ml), trifluoro acetic acid (80 µl, 1.0 mmol) was added, followed by slow addition of N-benzyl-N-(methoxymethyl)-N-trimethylsilylmethylamine (715 mg, 3.01

35

169

mmol). Stirring was continued for 2 h at room temperature. Another portion of N-benzyl-N-(methoxymethyl)-N-trimethylsilylmethylamine (300 mg, 1.26 mmol) was added and stirring continued for another 30 minutes. The reaction mixture was diluted with ethyl acetate (25 ml), washed with aqueous NaHCO₃ (15 ml). The organic layer
5 was dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to yield an oil (970 mg). The crude product was purified via silica gel chromatography with cyclohexane/ethyl acetate (gradient 0 – 50 %). Fractions containing the product were combined, the solvent evaporated to yield an oil (320 mg, 46 %).
ESI-MS: 313.1 [M+H]⁺

10

135.4 4-(1-Benzyl-pyrrolidin-3-yl)-2-methoxy-phenylamine

15

1-Benzyl-3-(3-methoxy-4-nitro-phenyl)-pyrrolidine (350 mg, 1.12 mmol) was dissolved in methanol (20 ml), tin dichloride (1.912 g, 10.08 mmol) was added, and the reaction
mixture stirred at reflux for 2 h. The methanol was evaporated, 1 N sodium hydroxide (50 ml) and ethyl acetate were added and stirring continued. The tin salts were filtered off, the organic phase was separated, dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to yield an oil (220 mg, 70 %).
ESI-MS: 283.1 [M+H]⁺

20

135.5 N-[4-(1-Benzyl-pyrrolidin-3-yl)-2-methoxy-phenyl]-4-isopropylbenzenesulfonamide

25

4-(1-Benzyl-pyrrolidin-3-yl)-2-methoxy-phenylamine (220 mg, 0.78 mmol) and 4-isopropylphenylsulfonyl chloride (170 mg, 0.78 mmol) were dissolved in tetrahydrofuran (20 ml). Triethylamine (0.32 ml, 2.34 mmol) was added and the reaction mixture stirred over night at room temperature (10 % conversion) and subsequently refluxed for 4 h (70 % conversion). One small portion of 4-isopropylphenylsulfonyl chloride and triethylamine were added and the reaction mixture was stirred 15 min at 150 °C in the microwave (CEM). The solvent was evaporated under reduced pressure, the residue treated with ethyl acetate (50 ml) and extracted twice with slightly acidic water. The organic layer was dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to give an oil (470 mg). The crude product was purified via silica gel chromatography with cyclohexane/ethyl acetate (gradient 0 – 100 %). Fractions containing the product were combined, the solvent evaporated to yield an oil (143 mg, 40 %).

30

35

ESI-MS: 465.1 [M+H]⁺

135.6 4-Isopropyl-N-(2-methoxy-4-pyrrolidin-3-yl-phenyl)-benzenesulfonamide, acetate

- 5 A mixture of N-[4-(1-benzyl-pyrrolidin-3-yl)-2-methoxy-phenyl]-4-isopropyl-benzenesulfonamide (143 mg, 0.31 mmol) and 10 % palladium on carbon (20 mg) in ethyl acetate (20 ml) and acetic acid (20 ml) was hydrogenated for 4 h at room temperature. The catalyst was filtered, and the solvent was removed under vacuum to yield an oil (100 mg, 71 %).
- 10 ESI-MS: 375.2 [M+H]⁺
¹H-NMR (D₂O): δ [ppm] 7.6 (d, 2H), 7.4 (d, 2H), 7.3 (d, 1H), 6.9 (d, 1H), 6.8 (s, 1H), 3.7 (m, 1H), 3.6-3.4 (m, 6H), 3.2 (m, 1H), 3.0 (m, 1H), 2.4 (m, 1H), 2.1 (m, 1H), 1.2 (d, 6H).

EXAMPLE 136

- 15 4-Isopropyl-N-[2-methoxy-4-(1-propyl-pyrrolidin-3-yl)-phenyl]-benzenesulfonamide, hydrochloride

4-Isopropyl-N-(2-methoxy-4-pyrrolidin-3-yl-phenyl)-benzenesulfonamide, acetate (40 mg, 0.09 mmol) and propionaldehyde (5.1 mg, 0.09 mmol) were dissolved in tetrahydrofuran (5 ml). Acetic acid (10 µl, 0.12 mmol) and sodium triacetoxyborohydride (37 mg, 0.18 mmol) were sequentially added to the reaction mixture and stirred for 30 minutes at room temperature. The reaction mixture was concentrated and the residue was dissolved in aqueous NaHCO₃ (10 ml) and extracted with diethyl ether (30 ml). The organic phase was separated, dried over magnesium sulfate, filtered, and evaporated to dryness to yield an oil (30 mg). The residue was dissolved in diethyl ether (25 ml) and HCl in ether was added and stirred overnight. The solvent was decanted and to the residue was added 3 ml of H₂O. This solution was lyophilised to yield the pure product (23 mg, 48 %).

20

- ESI-MS: 417.1 [M+H]⁺
- 30 ¹H-NMR (CH₃OH-d₄): δ [ppm] 7.6 (d, 2H), 7.4 (d, 2H), 7.3 (d, 1H), 6.9 (d, 1H), 6.8 (s, 1H), 3.5 (m, 5H), 3.3-3.2 (m, 2H), 3.0-2.9 (m, 4H), 2.4 (m, 1H), 2.0 (m, 1H), 1.7 (m, 2H), 1.2 (d, 6H), 1.0 (t, 3H).

EXAMPLE 137

- 35 N-[4-(1-Allyl-pyrrolidin-3-yl)-2-methoxy-phenyl]-4-isopropyl-benzenesulfonamide, hydrochloride

171

To 4-isopropyl-N-(2-Methoxy-4-pyrrolidin-3-yl-phenyl)-benzenesulfonamide, acetate (40 mg, 0.09 mmol) in N,N-dimethylformamide (5 ml) was added allyl bromide (8 μ l, 0.1 mmol) and K_2CO_3 (36 mg, 0.26 mmol). The reaction mixture was stirred at room temperature for 2 h. The reaction mixture was diluted with water (35 ml) and extracted
5 twice with diethyl ether (20 ml). The organic phase was separated, dried over magnesium sulfate, filtered, and evaporated to dryness to yield an oil (36 mg). The residue was dissolved in diethyl ether (25 ml) and HCl in ether was added and stirred overnight. The solvent was decanted and to the residue was added 3 ml H_2O . This solution was lyophilised to yield the pure product (17 mg, 37 %).

10 ESI-MS: 415.1 $[M+H]^+$

1H -NMR (CH_3OH-d_4): δ [ppm] 7.6 (d, 2H), 7.3 (m, 3H), 6.8 (d, 1H), 6.7 (s, 1H), 5.9 (m, 1H), 5.3 (d, 1H), 5.2 (d, 1H), 3.4 (s, 3H), 3.2 (m, 1H), 3.0-2.9 (m, 3H), 2.6 (m, 1H), 2.3 (m, 1H), 1.9 (m, 1H), 1.2 (d, 6H).

15 EXAMPLE 138

4-Isopropyl-N-[4-(1-propyl-piperidin-3-yl)-phenyl]-benzenesulfonamide, hydrochloride

138.1 3-Phenyl-1-propyl-piperidine

20 3-Phenyl-piperidine (4.0 g, 24.81 mmol) and propionaldehyde (1.8 ml, 24.81 mmol) were dissolved in tetrahydrofuran (100 ml). Acetic acid (2.14 ml, 37.21 mmol) and sodium trisacetoxyborohydride (10.515 g, 49.61 mmol) were sequentially added to the reaction mixture and stirred for 1 hour at room temperature. The reaction mixture was concentrated and the residue was dissolved in aqueous $NaHCO_3$ solution (50 ml) and
25 diethyl ether (100 ml). The organic phases was dried over magnesium sulfate, filtered, and evaporated to dryness to yield the crude product (4.6 g, 87 % purity).

ESI-MS: 204.15 $[M+H]^+$

138.2 3-(4-Nitro-phenyl)-1-propyl-piperidine

30

To ice cooled 3-phenyl-1-propyl-piperidine (4.6 g, 19.73 mmol) and KNO_3 (2.254 g, 22.29 mmol) was added concentrated H_2SO_4 . The reaction mixture was allowed to warm to room temperature and stirred for another 30 minutes. To the reaction mixture was added cautiously ice and subsequently the pH was adjusted to 9-10. The aqueous
35 phase was extracted several times with ethyl acetate. The organic phases was dried over magnesium sulfate, filtered, and evaporated to dryness to yield the crude product (5.2 g, 81 % purity).

ESI-MS: 249.15 [M+H]⁺

138.3 4-(1-Propyl-piperidin-3-yl)-phenylamine

5 3-(4-Nitro-phenyl)-1-propyl-piperidine (5.2 g, 1.86 mmol) was dissolved in methanol (35 ml), tin dichloride (3.78 g, 16.74 mmol) was added, and the reaction mixture stirred at reflux for 2 h and overnight at room temperature. The methanol was evaporated, 1 N sodium hydroxide (50 ml) and ethyl acetate were added and stirring continued. The tin salts were filtered off, the organic phase was separated, dried over magnesium sulfate,
10 filtered, and the solvent evaporated under reduced pressure to yield an the crude product (450 mg, 90 % purity).

138.4 4-Isopropyl-N-[4-(1-propyl-piperidin-3-yl)-phenyl]-benzenesulfonamide, hydrochloride

15 4-(1-Propyl-piperidin-3-yl)-phenylamine (600 mg, 1.65 mmol) and 4-isopropyl-phenylsulfonyl chloride (397 mg, 1.81 mmol) were dissolved in tetrahydrofuran (25 ml). Triethylamine (760 µl, 5.44 mmol) was added and the reaction mixture stirred for 72 hours at room temperature. The solvent was evaporated under reduced pressure, the
20 residue treated with diethyl ether (50 ml) and three times extracted with water (3 x 30 ml). The organic phase was treated with 1 molar HCl solution. The acidic aqueous solution was made alkaline with NaOH solution to pH 9-10 and then extracted with ethyl acetate (25 ml). The organic phase was dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to give the crude product (580 mg). The
25 crude product was purified via silica gel chromatography with ethyl acetate. Fractions containing the product were combined, the solvent evaporated to yield the pure product which was then converted into the hydrochloride salt via adding a solution of 1 N HCl in diethyl ether. The precipitate was filtered off and dried in vacuo to give the pure product (283 mg, 39 %).

30 ESI-MS: 401.15 [M+H]⁺

¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 10.3 (s, 1H), 10.25 (bs, 1H), 7.7 (d, 2H), 7.45 (d, 2H), 7.15 (d, 2H), 7.1 (d, 2H), 3.45 (m, 1H), 3.35 (m, 1H), 3.1-2.8 (m, 6H), 1.9 (m, 2H), 1.85 (m, 1H), 1.75 (m, 2H), 1.55 (m, 1H), 1.2 (d, 6H), 0.9 (t, 3H).

35 Following a procedure analogous to example 138.4 the compounds of examples 139 to 142 were prepared. The compounds are characterised by the following physical data.

EXAMPLE 139

N-[4-(1-Propyl-piperidin-3-yl)-phenyl]-4-trifluoromethoxy-benzenesulfonamide, hydrochloride

5 ESI-MS: 443.15 [M+H]⁺

¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 10.5 (s, 1H), 10.15 (bs, 1H), 7.9 (d, 2H), 7.6 (d, 2H), 7.2 (d, 2H), 7.1 (d, 2H), 3.45 (m, 2H), 3.1-2.8 (m, 5H), 1.9 (m, 2H), 1.85 (m, 1H), 1.7 (m, 2H), 1.55 (m, 1H), 0.9 (t, 3H).

10 EXAMPLE 140

4-Difluoromethoxy-N-[4-(1-propyl-piperidin-3-yl)-phenyl]-benzenesulfonamide, hydrochloride

ESI-MS: 425.15 [M+H]⁺

15 ¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 10.4 (s, 1H), 10.15 (bs, 1H), 7.85 (d, 2H), 7.4 (t, J = 70 Hz, 1H), 7.35 (d, 2H), 7.15 (d, 2H), 7.1 (d, 2H), 3.45 (m, 1H), 3.4 (m, 1H), 3.1-2.8 (m, 5H), 1.9 (m, 2H), 1.85 (m, 1H), 1.75 (m, 2H), 1.55 (m, 1H), 0.9 (t, 3H).

EXAMPLE 141

20 N-[4-(1-Propyl-piperidin-3-yl)-phenyl]-4-(2,2,2-trifluoro-ethyl)-benzenesulfonamide, hydrochloride

ESI-MS: 441.15 [M+H]⁺

25 ¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 10.4 (s, 1H), 10.2 (bs, 1H), 7.8 (d, 2H), 7.55 (d, 2H), 7.15 (d, 2H), 7.1 (d, 2H), 3.8 (q, 2H), 3.45 (m, 1H), 3.4 (m, 1H), 3.1-2.8 (m, 5H), 1.9 (m, 2H), 1.85 (m, 1H), 1.7 (m, 2H), 1.55 (m, 1H), 0.9 (t, 3H).

EXAMPLE 142

30 4-(2,2-Difluoro-cyclopropyl)-N-[4-(1-propyl-piperidin-3-yl)-phenyl]-benzenesulfonamide, hydrochloride

ESI-MS: 435.15 [M+H]⁺

35 ¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 10.35 (s, 1H), 10.15 (bs, 1H), 7.75 (d, 2H), 7.45 (d, 2H), 7.15 (d, 2H), 7.1 (d, 2H), 3.45 (m, 1H), 3.4 (m, 1H), 3.15-2.8 (m, 5H), 2.05 (m, 2H), 1.9 (m, 2H), 1.8 (m, 1H), 1.7 (m, 2H), 1.55 (m, 1H), 0.9 (t, 3H).

EXAMPLE 143

174

N-(3-Piperidin-3-yl-phenyl)-4-trifluoromethoxy-benzenesulfonamide

143.1 3-[3-(4-Trifluoromethoxy-benzenesulfonylamino)-phenyl]-piperidine-1-carboxylic acid tert-butyl ester

5

3-(3-Amino-phenyl)-piperidine-1-carboxylic acid tert-butyl ester (500 mg, 1.81 mmol) and dimethylaminopyridine (30 mg, 0.25 mmol) were dissolved in tetrahydrofuran (40 ml). 4-Trifluoromethoxyphenylsulfonyl chloride (519 mg, 1.99 mmol) was added and the reaction mixture stirred for 2 hours at room temperature. Additional quantities of trifluoromethoxyphenylsulfonyl chloride was added until the 3-(3-amino-phenyl)-piperidine-1-carboxylic acid tert-butyl ester was completely consumed to give the bis-sulfonylated product. The solvent was evaporated under reduced pressure, the residue treated with water (25 ml) and diethyl ether (50 ml). The organic phases was dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure. In an inert atmosphere, the residue was dissolved in ethanol (30 ml) and small pieces of metallic sodium were added and the reaction mixture was stirred for 1 hour at room temperature. The solvent was evaporated under reduced pressure, the residue treated with water (25 ml) and diethyl ether (50 ml). The organic phases was dried over magnesium sulfate, filtered, and the solvent evaporated under reduced pressure to give the crude product (385 mg). The crude product was purified via silica gel chromatography with dichloromethane/methanol (gradient 100:0 – 97:3). Fractions containing the product were combined, the solvent evaporated to yield the pure product (340 mg, 36 %).
ESI-MS: 445.05 $[M+H-C(CH_3)_3]^+$

25 143.2 N-(3-Piperidin-3-yl-phenyl)-4-trifluoromethoxy-benzenesulfonamide

3-[3-(4-Trifluoromethoxy-benzenesulfonylamino)-phenyl]-piperidine-1-carboxylic acid tert-butyl ester (340 mg, 0.66 mmol) was dissolved in dichloromethane (30 ml). Trifluoroacetic acid (2 ml) was added and the reaction mixture was stirred at room temperature for 1 hour. The reaction mixture was evaporated to dryness. Aqueous NaHCO₃ solution (10 ml) was added and extracted twice with ethyl acetate (25 ml). The combined organic phases were dried over magnesium sulfate, filtered, and evaporated to dryness to yield the pure product (250 mg, 95 % yield).
ESI-MS: 401.05 $[M+H]^+$

35 ¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 7.85 (d, 2H), 7.5 (d, 2H), 7.1 (t, 1H), 6.9 (d, 1H), 6.85 (m, 2H), 2.95 (m, 2H), 2.5 (m, 3H), 1.75 (m, 1H), 1.65 (m, 1H), 1.45 (m, 2H).

175

EXAMPLE 144

4-Isopropyl-N-(3-piperidin-3-yl-phenyl)-benzenesulfonamide

144.1 3-[3-(4-Isopropyl-benzenesulfonylamino)-phenyl]-piperidine-1-carboxylic acid
5 tert-butyl ester

The title compound was prepared similar to a procedure as described in example 143.1
ESI-MS: 403.15 $[M+H-C(CH_3)_3]^+$

10 144.2 4-Isopropyl-N-(3-piperidin-3-yl-phenyl)-benzenesulfonamide

The title compound was prepared similar to a procedure as described in example 144.1
ESI-MS: 359.15 $[M+H]^+$

1H -NMR (DMSO- d_6 , 400 MHz): δ [ppm] 7.75 (d, 2H), 7.4 (d, 2H), 7.1 (t, 1H), 6.9 (d,
15 1H), 6.85 (m, 2H), 2.95 (m, 2H), 2.85 (m, 1H), 2.5-2.35 (m, 3H), 1.75 (m, 1H), 1.6 (m,
1H), 1.4 (m, 2H), 1.2 (d, 6H).

EXAMPLE 145

N-[3-(1-Propyl-piperidin-3-yl)-phenyl]-4-trifluoromethoxy-benzenesulfonamide, hydro-
20 chloride

N-(3-Piperidin-3-yl-phenyl)-4-trifluoromethoxy-benzenesulfonamide (60 mg, 0.15 mmol)
and propionaldehyde (11 μ l, 0.15 mmol) were dissolved in tetrahydrofuran (10 ml).
Acetic acid (14 mg, 0.22 mmol) and sodium trisacetoxyborohydride (64 mg, 0.30 mmol)
25 were sequentially added to the reaction mixture and stirred for 1 hour at room tempera-
ture. The reaction mixture was concentrated and the residue was dissolved in aqueous
NaHCO₃ solution (5 ml) and diethyl ether (25 ml). The organic phases were dried over
magnesium sulfate, filtered, and evaporated to dryness to yield the crude product (62
mg). The crude product was purified via silica gel chromatography with dichloro-
30 methane/methanol (gradient 100:0 – 75:25). Fractions containing the product were
combined, the solvent evaporated to yield the pure product (42 mg) which was con-
verted into the hydrochloride salt (45 mg, 63 %).

ESI-MS: 443.15 $[M+H]^+$

1H -NMR (DMSO- d_6 , 400 MHz): δ [ppm] 10.45 (s, 1H), 10.25 (bs, 1H), 7.9 (d, 2H), 7.55
35 (d, 2H), 7.25 (t, 1H), 7.0 (m, 3H), 3.45 (m, 1H), 3.35 (m, 1H), 3.1-2.85 (m, 6H), 1.9 (m,
2H), 1.75 (m, 3H), 1.55 (m, 1H), 0.9 (t, 3H).

EXAMPLE 146

4-Isopropyl-N-[3-(1-propyl-piperidin-3-yl)-phenyl]-benzenesulfonamide, hydrochloride

5 The title compound was prepared in a procedure analogous to that described in example 145.

ESI-MS: 401.25 [M+H]⁺

¹H-NMR (DMSO-d₆, 400 MHz): δ [ppm] 10.3 (s, 1H), 10.15 (bs, 1H), 7.7 (d, 2H), 7.45 (d, 2H), 7.2 (t, 1H), 7.0 (m, 2H), 6.95 (d, 1H), 3.45 (m, 1H), 3.35 (m, 1H), 3.05-2.85 (m, 10 6H), 1.9 (m, 2H), 1.75 (m, 3H), 1.55 (m, 1H), 1.2 (d, 6H), 0.9 (t, 3H).

20 III. Examples of galenic administration forms

A) Tablets

25 Tablets of the following composition are pressed on a tablet press in the customary manner:

40 mg of substance from Example 8
120 mg of corn starch
13.5 mg of gelatin
30 45 mg of lactose
2.25 mg of Aerosil® (chemically pure silicic acid in submicroscopically fine dispersion)
6.75 mg of potato starch (as a 6% paste)

35 B) Sugar-coated tablets

20 mg of substance from Example 8
60 mg of core composition
70 mg of saccharification composition

40 The core composition consists of 9 parts of corn starch, 3 parts of lactose and 1 part of 60:40 vinylpyrrolidone/vinyl acetate copolymer. The saccharification composition con-

sists of 5 parts of cane sugar, 2 parts of corn starch, 2 parts of calcium carbonate and 1 part of talc. The sugar-coated tablets which had been prepared in this way are subsequently provided with a gastric juice-resistant coating.

5 IV. Biological investigations

Receptor binding studies:

The substance to be tested was either dissolved in methanol/Chremophor® (BASF-AG) or in dimethyl sulfoxide and then diluted with water to the desired concentration.

10

Dopamine D₃ receptor:

The assay mixture (0.250 ml) was composed of membranes derived from ~ 10⁶ HEK-293 cells possessing stably expressed human dopamine D₃ receptors, 0.1 nM [125I]-iodosulpride and incubation buffer (total binding) or, in addition, test substance (inhibition curve) or 1 μM spiperone (nonspecific binding). Each assay mixture was run in triplicate.

15

The incubation buffer contained 50 mM tris, 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 2 mM MgCl₂ and 0.1% bovine serum albumin, 10 μM quinolone and 0.1% ascorbic acid (prepared fresh daily). The buffer was adjusted to pH 7.4 with HCl.

20

Dopamine D_{2L} receptor:

The assay mixture (1 ml) was composed of membranes from ~ 10⁶ HEK-293 cells possessing stably expressed human dopamine D_{2L} receptors (long isoform) and 0.01 nM [125I] iodospiperone and incubation buffer (total binding) or, in addition, test substance (inhibition curve) or 1 μM haloperidol (nonspecific binding). Each assay mixture was run in triplicate.

30

The incubation buffer contained 50 mM tris, 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 2 mM MgCl₂ and 0.1% bovine serum albumin. The buffer was adjusted to pH 7.4 with HCl.

35 Measurement and analysis:

178

After having been incubated at 25°C for 60 minutes, the assay mixtures were filtered through a Whatman GF/B glass fiber filter under vacuum using a cell collecting device. The filters were transferred to scintillation vials using a filter transfer system. After 4 ml of Ultima Gold® (Packard) have been added, the samples were shaken for one hour and the radioactivity was then counted in a Beta-Counter (Packard, Tricarb 2000 or 2200CA). The cpm values were converted into dpm using a standard quench series and the program belonging to the instrument.

The inhibition curves were analyzed by means of iterative nonlinear regression analysis using the Statistical Analysis System (SAS) which is similar to the "LIGAND" program described by Munson and Rodbard.

The results of the receptro binding studies are expressed as receptor binding constants $K_i(D_2)$ and $K_i(D_3)$, respectively, as herein before described, and given in table 6.

In these tests, the compounds according to the invention exhibit very good affinities for the D_3 receptor (< 50 nM, or < 10 nM, frequently < 5 nM) and bind selectively to the D_3 receptor.

The results of the binding tests are given in table 6.

Table 6:

Example	$K_i(D_3)^*$ [nM]	$K_i(D_2)^*$ [nM]	$K_i(D_2)^*/ K_i(D_3)^*$
1	0.09	7.6	89
2	0.24	2.3	9
3	0.17	13.8	91
4	3.16	403	127
5	2.9	267	93
6	0.45	20.1	45
7	3.5	212	61
8	4.1	235	57
9	2.6	129	49
10	1.9	111	60
12	3.0	131	43
13	3.1	168	54

179

Example	$K_i(D3)^*$ [nM]	$K_i(D2)^*$ [nM]	$K_i(D2)^* / K_i(D3)^*$
14	3.4	123	37
15	1.9	74	39
16	12.6	393	31
17	3.1	126	41
18	2.3	90	39
19	0.48	12.6	26
20	0.3	7.3	24
21	0.4	11	27
23	1.07	46.6	44
24	1.2	72	60
25	1.6	95	60
26	28.1	1278	46
27	0.23	7.1	30
28	3.3	133	41
29	0.67	27.3	41
30	0.28	11.7	42
31	10.8	257	24
33	0.17	5.47	33
34	5.1	174	34
35	0.3	7.4	24
37	0.15	3.25	22
38	0.29	6.3	21
40	0.46	18.2	40
41	0.44	13.5	30
46	49	1.830	37
48	2.1	137	65
49	46	1,808	39
53	0.41	5.85	14
54	0.32	8.73	27
55	7.11	1175	165
56	24.4	2661.2	109
57	7.18	880	123

180

Example	$K_i(D3)^*$ [nM]	$K_i(D2)^*$ [nM]	$K_i(D2)^*/K_i(D3)^*$
58	39.80	2940.0	74
59	6.41	1525	238
60	7.63	7780	1019
61	21.3	--	--
62	25.2	3545	141
63	5.75	647.21	113
64	2.10	344	164
65	6.79	197	29
67	0.35	3.88	11
68	3.00	142	47
69	2.51	90.3	36
70	0.84	37.3	44
71	4.79	52.3	11
72	35.1	473	13
73	0.61	6.73	11
74	0.69	10.5	15
75	0.63	12.2	19
77	1.33	50	37
78	1.40	41.9	30
79	13.5	1111	82
80	37.6	1095	29
81	31.3	1162.0	37
82	5.59	437	78
84	0.95	16.2	17
85	0.57	18.5	32
86	1.18	119	101
87	0.55	20.5	37
88	0.92	41	45
89	0.6	27	45
90	11	280	25
92	0.61	17.3	28
93	0.48	14.7	31
94	0.23	4.7	21
95	1.3	27.9	22
96	0.83	57.8	70


181

Example	$K_i(D3)^*$ [nM]	$K_i(D2)^*$ [nM]	$K_i(D2)^*/K_i(D3)^*$
101	3.7	268	73
102	52	2714	52
103	1.2	45	38
104	8.3	352	43
105	6.1	309	51
107	0.42	45.8	108
108	11.2	203.8	18
109	7.4	257.9	35
110	6.6	895	135
111	16.4	723	44
112	10.3	1196	116
113	0.49	42.2	86
114	4.2	464	111
118	36.6	1169	32
120	2.0	174	86
121	3.7	163	44
122	43.9	1342	31
123	0.45	41	91
125	19.5	1305	67
126	34	1021	30
127	0.6	64.7	112
128	2.7	12.9	5
129	3.6	106.3	30
130	0.52	8.5	16
131	5.8	430	74
133	7.3	192.7	26
134	0.76		13
135	0.85		17
136	0.27		6
137	0.36		12

* Receptor binding constants obtained according to the assays as described herein before

We claim:

1. A compound of the formula (I)



wherein

15 n is 0, 1 or 2;

G is CH₂ or CHR³;

20 R¹ is H, C₁-C₆-alkyl, C₁-C₆-alkyl substituted by C₃-C₆-cycloalkyl, C₁-C₆-hydroxyalkyl, fluorinated C₁-C₆-alkyl, C₃-C₆-cycloalkyl, fluorinated C₃-C₆-cycloalkyl, C₃-C₆-alkenyl, fluorinated C₃-C₆-alkenyl, formyl, acetyl or propionyl;

25 R², R³ and R⁴ are, independently of each other, H, methyl, fluoromethyl, di-fluoromethyl, or trifluoromethyl;

30 A is phenylene, pyridylene, pyrimidylene, pyrazinylene, pyridazinylenes or thiophenylene, which can be substituted by one or more substituents selected from halogen, methyl, methoxy and CF₃;

35 E is NR⁵ or CH₂, wherein R⁵ is H or C₁-C₃-alkyl;

40 Ar is a cyclic radical selected from the group consisting of phenyl, a 5- or 6-membered heteroaromatic radical comprising as ring members 1, 2 or 3 heteroatoms selected from N, O and S and a phenyl ring fused to a saturated or unsaturated 5- or 6-membered carbocyclic or heterocyclic ring, where the heterocyclic ring comprises as ring members 1, 2 or 3 heteroatoms selected from N, O and S and/or 1, 2 or 3 heteroatom-containing groups each independently selected from NR⁸, where R⁸ is H, C₁-C₄-alkyl, fluorinated C₁-C₄-alkyl, C₁-C₄-alkylcarbonyl or fluorinated C₁-C₄-alkylcarbonyl, and where the cyclic radical Ar may carry 1, 2 or 3 substituents R^a;


R^a is halogen, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_1 - C_6 -alkoxy, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkylthio, fluorinated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, fluorinated C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, fluorinated C_1 - C_6 -alkylsulfonyl, phenylsulfonyl, benzyloxy, phenoxy, where the phenyl radical in the 3 last-mentioned radicals may be unsubstituted or may carry 1 to 3 substituents selected from C_1 - C_4 -alkyl, fluorinated C_1 - C_4 -alkyl and halogen, CN, nitro, C_1 - C_6 -alkylcarbonyl, fluorinated C_1 - C_6 -alkylcarbonyl, C_1 - C_6 -alkylcarbonylamino, fluorinated C_1 - C_6 -alkylcarbonylamino, carboxy, $NH-C(O)-NR^6R^7$, NR^6R^7 , $NR^6R^7-C_1$ - C_6 -alkylene, $O-NR^6R^7$, where R^6 and R^7 are, independently of each other, H, C_1 - C_4 -alkyl, fluorinated C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy or may form, together with N, a 4-, 5- or 6-membered saturated or unsaturated ring, or is a saturated or unsaturated 3- to 7-membered heterocyclic ring comprising as ring members 1, 2, 3 or 4 heteroatoms selected from N, O and S and/or 1, 2 or 3 heteroatom-containing groups selected from NR^9 , where R^9 has one of the meanings given for R^8 , SO, SO_2 and CO, and where the heterocyclic ring may carry 1, 2 or 3 substituents selected from hydroxy, halogen, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl and C_1 - C_6 -alkoxy;

and physiologically tolerated acid addition salts thereof.

2. A compound as claimed in claim 1,

25

30



wherein

35

R^1 is H, C_1 - C_6 -alkyl which may be substituted by C_3 - C_6 -cycloalkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_3 - C_6 -alkenyl, fluorinated C_3 - C_6 -alkenyl, formyl, acetyl or propionyl; and


40

Ar is a cyclic radical selected from the group consisting of phenyl, a 5- or 6-membered heteroaromatic radical comprising as ring members 1, 2 or 3 heteroatoms selected from N, O and S and a phenyl ring fused to a saturated or unsaturated 5- or 6-membered carbocyclic or heterocyclic ring, where the heterocyclic ring comprises as ring members 1, 2 or 3 heteroa-

184

- 5 toms selected from N, O and S, and where the cyclic radical may carry 1, 2 or 3 substituents R^a selected from the group consisting of halogen, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, CN, acetyl, carboxy, NR^6R^7 , NR^6R^7 - C_1 - C_6 -alkylene, where R^6 and R^7 are, independently of each other, H, C_1 - C_4 -alkyl or fluorinated C_1 - C_4 -alkyl or may form, together with N, a 4-, 5- or 6-membered saturated or unsaturated ring, and a saturated or unsaturated 5- or 6-membered heterocyclic ring comprising as ring members 1, 2 or 3 heteroatoms selected from N, O and S.
- 10 3. The compound as claimed in one of claims 1 or 2, wherein n is 0 or 1.
4. The compound as claimed in any of the preceding claims, wherein R^1 is hydrogen, methyl, ethyl, n-propyl, 2-fluoroethyl, 3-fluoropropyl, 3-hydroxypropyl, cyclopropylmethyl or allyl.
- 15 5. The compound as claimed in claim 4, wherein R^1 is n-propyl or allyl.
6. The compound as claimed in any of the preceding claims, wherein R^2 , R^3 and R^4 are H.
- 20 7. The compound as claimed in any of the preceding claims, wherein A is phenylene, pyridylene or pyrimidylene.
8. The compound as claimed in any of the preceding claims, wherein E is NH.
- 25 9. The compound as claimed in any of the preceding claims, wherein Ar is phenyl, thienyl, pyridyl, pyrimidyl, imidazolyl, isoxazolyl, thiazolyl, triazolyl, thiadiazolyl, quinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, benzofuranyl, benzothiophenyl, benzoxazinyl, benzothiazolyl, benzoxadiazolyl, benzothiadiazolyl or indanyl, which may be substituted as defined in claim 1.
- 30 10. The compound as claimed in of claim 9, wherein Ar is phenyl, thienyl, pyridyl, benzofuranyl or indanyl, which may be substituted as defined in claim 2.
- 35 11. The compound as claimed in claim 9 or 10, wherein Ar is phenyl which carries 1, 2 or 3 substituents selected from the group consisting of halogen, C_1 - C_6 -alkyl, fluorinated C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, fluorinated C_1 - C_4 -alkoxy, C_2 - C_4 -alkenyl, fluorinated C_2 - C_4 -alkenyl, $CH_2N(CH_3)_2$, NR^6R^7 , where R^6 and R^7 are independently of each other H, C_1 - C_4 -alkyl or fluorinated C_1 - C_4 -alkyl, C_3 - C_6 -cycloalkyl optionally substituted by halogen, acetyl or carboxyl, or Ar is thienyl, pyridyl, benzofuranyl or indanyl, which are optionally substituted by halogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkenyl.
- 40

12. The compound as claimed in any of the preceding claims, wherein Ar carries one radical R^a of the formula R^a



5 wherein

Y is N, CH or CF,

R^{a1} and R^{a2} are independently of each other selected from C_1 - C_2 -alkyl, fluorinated C_1 - C_2 -alkyl and C_1 - C_2 -alkoxy, provided for Y being CH or CF one of the radicals R^{a1} or R^{a2} may also be hydrogen or fluorine, or

- 10 R^{a1} and R^{a2} together form a radical $(CH_2)_m$ wherein 1 or 2 of the hydrogen atoms may be replaced by fluorine, hydroxy, C_1 - C_2 -alkyl or C_1 - C_2 -alkoxy, wherein one CH_2 moiety may be replaced by O, S, SO, SO_2 or NR^c , wherein R^c is H or C_1 - C_2 -alkyl, and wherein m is 2, 3, 4, 5 or 6.

- 15 13. The compound as claimed in claim 11, wherein the radical R^a is selected from isopropyl, (R)-1-fluoroethyl, (S)-1-fluoroethyl, 2-fluoroethyl, 1,1-difluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, (R)-1-fluoropropyl, (S)-1-fluoropropyl, 2-fluoropropyl, 3-fluoropropyl, 1,1-difluoropropyl, 2,2-difluoropropyl, 3,3-difluoropropyl, 3,3,3-trifluoropropyl, (R)-2-fluoro-1-methylethyl, (S)-2-fluoro-1-methylethyl, (R)-2,2-difluoro-1-methylethyl, (S)-2,2-difluoro-1-methylethyl, (R)-1,2-difluoro-1-methylethyl, (S)-1,2-difluoro-1-methylethyl, (R)-2,2,2-trifluoro-1-methylethyl, (S)-2,2,2-trifluoro-1-methylethyl, 2-fluoro-1-(fluoromethyl)ethyl, 1-(difluoromethyl)-2,2-difluoroethyl, cyclopropyl, cyclobutyl, 1-fluorocyclopropyl, 2-fluorocyclopropyl, (S)-2,2-difluorocyclopropyl and (R)-2,2-difluorocyclopropyl.

25

14. The compounds as claimed in any of claims 12 or 13, wherein the radical R^a carries 1, 2, 3 or 4 fluorine atoms.

- 30 15. The compound as claimed in any of the preceding claims, wherein Ar is phenyl that carries a radical R^a in the 4-position of the phenyl ring.

16. The compound as claimed in any of the preceding claims, wherein the absolute configuration at the carbon atom carrying the group A is (S).

- 35 17. The compound as claimed in claim 1, 3 to 9, 15 or 16, wherein Ar is phenyl which is substituted by a 5- or 6 membered heterocyclic radical selected from azetidiny, pyrrolidiny, oxopyrrolidiny, oxo-oxazolidiny, piperidiny, piperaziny, morpholiny, thiomorpholiny, 1-oxothiomorpholiny, 1,1-dioxothiomorpholiny, pyrrolyl, furanyl, thienyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, triazolyl, oxadiazolyl, furazanyl, thiadiazolyl, and tetrazolyl, where the heterocyclic radical may be unsubstituted
- 40

or may carry 1 to 3 substituents selected from halogen, C₁-C₄-alkyl, fluorinated C₁-C₄-alkyl, C₁-C₄-alkoxy and hydroxy.

- 5 18. The compound as claimed in claim 17, wherein Ar is phenyl which is substituted by a 5- or 6 membered heterocyclic radical selected from azetidiny, pyrrolidiny, oxopyrrolidiny, oxo-oxazolidiny, morpholinyl, furanyl, thienyl, pyrazolyl, oxazolyl, isoxazolyl, and thiadiazolyl, where the heterocyclic radical may be unsubstituted or may carry 1 to 3 substituents selected from halogen and C₁-C₄-alkyl.
- 10 19. A pharmaceutical composition comprising at least one compound as claimed in any of the preceding claims, optionally together with at least one physiologically acceptable carrier or auxiliary substance.
- 15 20. A method for treating a medical disorder susceptible to treatment with a dopamine D3 receptor ligand, said method comprising administering an effective amount of at least one compound as claimed in any of claims 1 to 18 to a subject in need thereof.
- 20 21. The method as claimed in claim 20, wherein the medical disorder is a disease of the central nervous system.
- 25 22. The use of a compound as claimed in any of claims 1 to 18 for preparing a pharmaceutical composition for the treatment of a medical disorder susceptible to treatment with a dopamine D3 receptor ligand.
23. The use as claimed in claim 22, wherein the medical disorder is a disease of the central nervous system.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2005/011106

A. CLASSIFICATION OF SUBJECT MATTER

C07D207/09	C07D401/12	C07D405/12	C07D409/12	C07D401/04
C07D403/04	C07D413/12	C07D205/04	C07D413/10	C07D413/14
C07D211/28	A61K31/397	A61K31/40	A61K31/445	A61P25/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98/27081 A (SMITHKLINE BEECHAM PLC) 25 June 1998 (1998-06-25) cited in the application claims 1,14; example 131 -----	1-23
X	WO 00/05225 A (MERCK PATENT GMBH; BOETTCHER, HENNING; VAN AMSTERDAM, CHRISTOPH; HARTI) 3 February 2000 (2000-02-03) page 21, lines 24,25 -----	1-4, 6-11,15

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* & * document member of the same patent family

Date of the actual completion of the international search

15 February 2006

Date of mailing of the international search report

22/02/2006

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Johnson, C

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2005/011106

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2005/011106

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 20,21 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2005/011106

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9827081	A	25-06-1998	AP 1277 A 05-05-2004
			AT 247099 T 15-08-2003
			AU 729056 B2 25-01-2001
			AU 6090498 A 15-07-1998
			BG 103530 A 31-01-2000
			BR 9713734 A 28-03-2000
			CA 2275492 A1 25-06-1998
			CN 1246116 A 01-03-2000
			CZ 9902203 A3 17-11-1999
			DE 69724142 D1 18-09-2003
			DE 69724142 T2 03-06-2004
			EA 2351 B1 25-04-2002
			EP 0946539 A1 06-10-1999
			ES 2203831 T3 16-04-2004
			HU 0000658 A2 28-02-2001
			ID 22821 A 09-12-1999
			JP 2001506646 T 22-05-2001
			MA 24426 A1 01-07-1998
			NO 993003 A 18-06-1999
			NZ 335970 A 26-10-2001
			OA 11066 A 10-03-2003
			PL 334337 A1 28-02-2000
			SK 80899 A3 14-02-2000
			TR 9901361 T2 23-08-1999
			TW 418205 B 11-01-2001
			US 6423717 B1 23-07-2002
WO 0005225	A	03-02-2000	AU 5033899 A 14-02-2000
			BR 9912299 A 20-11-2001
			CA 2338209 A1 03-02-2000
			CN 1309654 A 22-08-2001
			HU 0102622 A2 28-12-2001
			ID 27846 A 26-04-2001
			JP 2002521377 T 16-07-2002
			NO 20010333 A 19-01-2001
			PL 345283 A1 03-12-2001
			SK 492001 A3 06-08-2001
			ZA 200101379 A 20-05-2002