



(86) Date de dépôt PCT/PCT Filing Date: 2004/11/12
(87) Date publication PCT/PCT Publication Date: 2005/05/26
(45) Date de délivrance/Issue Date: 2012/11/06
(85) Entrée phase nationale/National Entry: 2006/05/09
(86) N° demande PCT/PCT Application No.: US 2004/037578
(87) N° publication PCT/PCT Publication No.: 2005/047478
(30) Priorité/Priority: 2003/11/12 (US60/519,818)

(51) Cl.Int./Int.Cl. *C12N 15/24* (2006.01),
A61K 38/20 (2006.01), *A61K 39/395* (2006.01),
A61P 31/00 (2006.01), *A61P 35/00* (2006.01),
A61P 37/00 (2006.01), *C07K 14/54* (2006.01),
C07K 16/24 (2006.01), *C12N 1/21* (2006.01),
C12N 15/63 (2006.01), *C12N 5/10* (2006.01),
C12Q 1/02 (2006.01)

(72) Inventeurs/Inventors:
KIM, SOO-HYUN, US;
DINARELLO, CHARLES A., US;
AZAM, TANIA, US

(73) Propriétaire/Owner:

(54) Titre : COMPOSITIONS ET PROCEDES DE REGULATION DU FACTEUR-ALPHA DE NECROSE TUMORALE
(54) Title: COMPOSITIONS AND METHODS FOR REGULATION OF TUMOR NECROSIS FACTOR-ALPHA

(57) **Abrégé/Abstract:**

The present invention relates to compositions and methods relating to an interleukin-18- inducible cytokine termed tumor necrosis factor-alpha inducing factor (TAIF) or interleukin-32 (IL-32). In particular, the present invention provides compositions and methods for treating autoimmune diseases and cancer, in part by regulation of tumor necrosis factor-alpha expression.



(73) **Propriétaires(suite)/Owners(continued):**

THE REGENTS OF THE UNIVERSITY OF COLORADO, A BODY CORPORATE, US

(74) **Agent:** GOWLING LAFLEUR HENDERSON LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
26 May 2005 (26.05.2005)

PCT

(10) International Publication Number
WO 2005/047478 A2

- (51) International Patent Classification⁷: **C12N**
- (21) International Application Number:
PCT/US2004/037578
- (22) International Filing Date:
12 November 2004 (12.11.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/519,818 12 November 2003 (12.11.2003) US
- (71) Applicant (for all designated States except US): **THE REGENTS OF THE UNIVERSITY OF COLORADO, A BODY CORPORATE** [US/US]; 201 Regent Administrative Center, Boulder, CO 80309 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **KIM, Soo-Hyun** [KR/US]; 19833 East Villanova Place, Aurora, CO 80013 (US). **DINARELLO, Charles, A.** [US/US]; 333 15th Street, Boulder, CO 80302 (US). **AZAM, Tania** [US/US]; 7685 East Gunnison Place, Denver, CO 80231 (US).
- (74) Agents: **LEKUTIS, Christine, A.** et al.; Medlen & Carroll, LLP, 101 Howard Street, Suite 350, San Francisco, CA 94105 (US).

- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOSITIONS AND METHODS FOR REGULATION OF TUMOR NECROSIS FACTOR-ALPHA

(57) Abstract: The present invention relates to compositions and methods relating to an interleukin18- inducible cytokine termed tumor necrosis factor-alpha inducing factor (TAIF) or interleukin-32 (IL-32). In particular, the present invention provides compositions and methods for treating autoimmune diseases and cancer, in part by regulation of tumor necrosis factor-alpha expression.

WO 2005/047478 A2

COMPOSITIONS AND METHODS FOR REGULATION OF TUMOR NECROSIS FACTOR-ALPHA

This invention was made in part with government support under grants AI-15614
5 and HL-68743, from the National Institutes Health. As such, the United States government
has certain rights in the invention.

FIELD OF THE INVENTION

The present invention relates to compositions and methods relating to an interleukin-
10 18- inducible cytokine termed tumor necrosis factor-alpha inducing factor (TAIF) or
interleukin-32 (IL-32). In particular, the present invention provides compositions and
methods for treating autoimmune diseases and cancer, in part by regulation of tumor
necrosis factor-alpha expression.

15 BACKGROUND OF THE INVENTION

Rheumatoid arthritis (RA) is a common chronic inflammatory arthritis that affects
about 1% of adults worldwide, with a female predominance and a peak onset in the fourth
decade of life (*See*, Firestein, "Rheumatoid Arthritis," in Scientific American Medicine,
2000; and Cohen, "Systemic Autoimmunity," in Paul (ed.) Fundamental Immunology,
20 Lippincott-Raven Publishers: Philadelphia, pp. 1067-1088, 1999). Intense inflammation
occurs in synovial joints, with infiltration of the synovial membrane by mononuclear
phagocytes, lymphocytes and neutrophils, causing significant joint pain. In addition, RA
patients generally develop loss of cartilage and bone around joints, which leads to a loss of
mobility.

25 Although the cause of RA has not been precisely defined, various characteristics of
the disease are indicative of an autoimmune component to RA etiology. In particular,
macrophage and fibroblast-derived cytokines are abundantly expressed in rheumatoid joints
(Firestein *et al.*, *J Immunol*, 144:3347, 1994). Tumor necrosis factor alpha (TNF α) and
interleukin-1 (IL-1) appear to be the major pathogenic factors, in that both can induce
30 synoviocyte proliferation, collagenase production, and prostaglandin release, while
overexpression can induce arthritis in animal models (Firestein, *supra*, 2000). IL-18 is also
present in RA joints and can directly activate macrophages to produce proinflammatory
cytokines (Gracie *et al.*, *J Clin Invest*, 104:1393, 1999).

Current RA therapies are directed to analgesia, control of inflammation, and alteration of the disease course. More aggressive treatment approaches are now frequently adopted, with RA patients rapidly requiring a switch from non-steroidal anti-inflammatory drugs (NSAIDs) to a second line reagent such as methotrexate. Unfortunately, methotrexate alone does not adequately control RA in most patients, causing physicians to select either add-on therapy or a series of single agents (Firestein, *supra*, 2000), for example leflunomide, sulfasalazine, or a TNF inhibitor. TNF-inhibitors that have been used with some success to treat RA include TNF-reactive monoclonal antibodies (infliximab/REMICADE and adalimumab/HUMIRA) and a soluble TNF-receptor /immunoglobulin fusion protein (etanercept/ENBREL). However, it is desirable to provide clinicians with additional therapies to use alone or as cocktails to halt the progression of this debilitating disease.

SUMMARY OF THE INVENTION

The present invention relates to compositions and methods relating to an interleukin-18- inducible cytokine termed tumor necrosis factor-alpha inducing factor (TAIF) or interleukin-32 (IL-32). In particular, the present invention provides compositions and methods for treating autoimmune diseases and cancer, in part by regulation of tumor necrosis factor-alpha expression.

The present invention provides purified nucleic acids comprising a sequence at least 80% identical to SEQ ID NO:15, wherein the sequence encodes interleukin-32 (IL-32), and wherein the sequence comprises exon 3 and exon 4 of IL-32 in substantially contiguous association. In some preferred embodiments, the IL-32 is: an alpha isoform comprising the amino acid sequence set forth as SEQ ID NO:7; a beta isoform comprising the amino acid sequence set forth as SEQ ID NO:8; or a delta isoform comprising the amino acid sequence set forth as SEQ ID NO:10. In other embodiments, the sequence lacks intron 4 of IL-32, while in particularly preferred embodiments, the sequence is at least 90% identical to SEQ ID NO:15. Also provided are purified nucleic acids, comprising a sequence selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4 and SEQ ID NO:6. In some embodiments, the sequence is operably linked to a heterologous promoter. In preferred embodiments, the nucleic acid is contained within a vector. Moreover, host cells comprising the vector are provided.

In addition, the present invention provides purified proteins encoded by nucleic acids comprising a sequence at least 80% identical to SEQ ID NO:15, wherein the sequence

encodes interleukin-32 (IL-32), and wherein the sequence comprises exon 3 and exon 4 of IL-32 in substantially contiguous association. In some preferred embodiments, the IL-32 is: an alpha isoform comprising the amino acid sequence set forth as SEQ ID NO:7; a beta isoform comprising the amino acid sequence set forth as SEQ ID NO:8; or a delta isoform comprising the amino acid sequence set forth as SEQ ID NO:10. In other embodiments, the IL-32 is not a gamma isoform, while in preferred embodiments, the IL-32 does not comprise the amino acid sequence set forth as SEQ ID NO:14. In some preferred embodiments, the IL-32 is a recombinant protein expressed in a cell selected from the group consisting of a bacterial cell, a yeast cell, an insect cell, and a mammalian cell. In a subset of these embodiments, the recombinant protein is a fusion protein.

Also provided by the present invention are antibodies, which bind to IL-32. In some preferred embodiments the antibody is a monoclonal antibody, while in other embodiments, a Fab fragment of the monoclonal antibody is provided. In some embodiments, the monoclonal antibody (mAb) is chosen from but not limited to 32-4 and 32-9. Hybridoma cells that produce the 32-4 mAb are being deposited with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Virginia 20110-2209. Likewise, hybridoma cells that produce the 32-9 mAb are being deposited with ATCC. Additionally, in some preferred embodiments, the monoclonal antibody is chosen from but not limited to a chimeric monoclonal antibody, a humanized monoclonal antibody, and a human monoclonal antibody. In a subset of embodiments, the monoclonal antibody inhibits IL-32-induced TNF α production by a target cell, inhibits IL-32-induced I κ B degradation in a target cell, and/or inhibits rapid IL-32-induced p38 MAPK phosphorylation in a target cell.

Moreover, the present invention provides methods for inducing TNF α production, comprising contacting at least one cell with an IL-32 protein under conditions suitable for inducing TNF α production. In preferred embodiments, the IL-32 protein is selected from the group consisting of an alpha isoform, a beta isoform, a gamma isoform and a delta isoform. In some embodiments the at least one cell comprises a leukocyte, while in a subset of these embodiments, the leukocyte is selected from the group consisting of monocytes and macrophages.

Also provided by the present invention are methods of treating a subject, comprising: providing a subject and an antibody that binds IL-32; and administering the antibody to the subject. In preferred embodiments, the IL-32 is selected from the group consisting of an alpha isoform, a beta isoform, a gamma isoform and a delta isoform. In

particularly preferred embodiments, the subject has, is suspected of having, or is at risk of having an autoimmune disease. In some embodiments, the autoimmune disease is chosen from but not limited to multiple sclerosis, myasthenia gravis, autoimmune neuropathy, autoimmune uveitis, Crohn's disease, ulcerative colitis, primary biliary cirrhosis, autoimmune hepatitis, autoimmune hemolytic anemia, pernicious anemia, autoimmune thrombocytopenia, type 1 diabetes mellitus, Grave's disease, Hashimoto's thyroiditis, autoimmune oophoritis and orchitis, temporal arteritis, anti-phospholipid syndrome, Vasculitides, Behcet's disease, rheumatoid arthritis, systemic lupus erythematosus, scleroderma, polymyositis, dermatomyositis, spondyloarthropathy, Sjogren's syndrome, psoriasis, dermatitis herpetiformis, pemphigus vulgaris, and vitiligo. The present invention provides antibodies chosen from but not limited to a human monoclonal antibody and a humanized mouse monoclonal antibody. In preferred embodiments, the administering is done under conditions suitable for alleviating at least one symptom of an autoimmune disease.

Furthermore, the present invention provides methods for screening for inhibitors of IL-32, comprising: providing an IL-32 protein, and at least one drug candidate; and analyzing the effect of the drug candidate on at least one activity of the IL-32 protein. In some embodiments, the IL-32 protein is a recombinant protein selected from the group consisting of an alpha isoform, a beta isoform, a gamma isoform and a delta isoform. In preferred embodiments, the drug candidate is chosen from but not limited to an IL-32-reactive monoclonal antibody, and a dominant-negative IL-32 variant. In particularly preferred embodiments, the at least one activity of the IL-32 protein comprises upregulation of TNF α expression.

Also provided by the present invention are methods of treating a subject, comprising: providing a subject and an IL-32 protein; and administering the IL-32 protein to the subject. In preferred embodiments, the IL-32 protein is a recombinant protein selected from the group consisting of an alpha isoform, a beta isoform, a gamma isoform and a delta isoform. In particularly preferred embodiments, the subject has, is suspected of having, or is at risk of having cancer.

The present invention also provides methods and kits for measuring IL-32 concentration in sera of a subject comprising providing sera from a subject and a IL-32-reactive antibody, and screening the sera with the antibody under conditions suitable for quantifying IL-32. In some embodiments, the subject is an autoimmune disease patient, while in other embodiments, the subject is a sepsis patient. In some preferred embodiments,

WO 2005/047478

PCT/US2004/037578

the screening is accomplished by electrochemiluminescence assay, while in other embodiments, the screening is accomplished by enzyme-linked immunosorbent assay. The IL-32-reactive antibody, in some embodiments, is chosen from but not limited to a polyclonal rabbit and human-IL-32 antibody, and a monoclonal mouse anti-human IL-32 antibody.

DESCRIPTION OF THE FIGURES

Figure 1 shows the expression and activity of a functional IL-18R β chain in human A549 lung carcinoma cells. Panel A provides the results of an RT-PCR analysis of IL-18R β expression in transfected and wild type A549 cells. Panels B and C provide graphs depicting IL-6 and IL-8 secretion respectively, in response to IL-18 (50 ng/ml) stimulation in transfected but not wild type A549 cells after 16 hours (N=7). Panel D shows the induction of NK4 (IL-32) RNA expression in transfected cells, in the presence and absence of IL-18.

Figure 2 graphically depicts the induction of TNF α expression upon treatment of mouse Raw 264.7 macrophage cells with recombinant IL-32 α (TAIF) in the presence of polymyxin B. Panel A shows that the level of recombinant IL-32 α in ion exchange chromatography fractions correlated with the levels of TNF α secreted by treated cells. Each ion exchange chromatograph fraction is shown after 10% SDS-PAGE followed by Coomassie blue staining. Panel B shows that recombinant IL-32 α expressed in bacteria induced TNF α expression in a dose-dependent manner. Panel C shows that recombinant IL-32 α expressed in mammalian cells also induced expression of TNF α . The amount of recombinant IL-32 α was estimated by immunoblot.

Figure 3 provides an alignment of the DNA sequences of the open reading frames of the four IL-32 splice variants (IL-32 α disclosed as SEQ ID NO:3, IL-32 β disclosed as SEQ ID NO:4, IL-32 γ disclosed as SEQ ID NO:5, and IL-32 δ disclosed as SEQ ID NO:6). The alignment was done with the ClustalW program (available on the web site of the Swiss node of EMBnet) and manually corrected.

Figure 4 provides an alignment of the four human IL-32 splice variants in panel A (IL-32 α disclosed as SEQ ID NO:7, IL-32 β disclosed as SEQ ID NO:8, IL-32 γ disclosed as SEQ ID NO:9, and IL-32 δ disclosed as SEQ ID NO:10), as well as an alignment of IL-32 β protein sequences from several mammalian species (human sequence disclosed as SEQ ID

WO 2005/047478

PCT/US2004/037578

NO: 8, equine disclosed as SEQ ID NO:16, and bovine disclosed as SEQ ID NO:17). The alignment was done with programs available on the web site of the Swiss node of EMBnet, and manually corrected. Myr and Gly indicate potential N-myristoylation or N-glycosylation sites, respectively.

Figure 5 depicts the structure of the human IL-32 gene on chromosome 16p13.3, with exons depicted by stippled boxes. The numbers above and below the schematic of the IL-32 gene delineate the eight exons, while the sequence of this 5 kb genomic fragment is disclosed herein as SEQ ID NO:11. Splicing of the four IL-32 variants (α , β , γ and δ) is also shown.

Figure 6 illustrates that rIL-32 induces proinflammatory cytokines in both macrophage cells (Raw) and monocytes (THP-1). Panel A indicates that rIL-32 α produced in *E. coli* induces both MIP-2 and TNF α secretion in a dose-dependent manner. IL-32 concentrations are indicated in units/ml on the x-axis. Panel B shows that both IL-32 α (10 U/ml) and IL-32 β (10 U/ml) variants activate mouse Raw 264.7 macrophage cells. Panel C provides a graph of human and murine TNF α induced by various sources of IL-32 β . Raw cells and PMA-differentiated-THP-1 cells were treated with 1 U/ml of Anjou65 and Cos-7S, or 2 U/ml of Cos7-T. IL-32 α produced in *E. coli* and IL-32 β produced in mammalian cells induced TNF α expression in PMA-differentiated THP-1 cells in a dose-dependent manner. Panel E illustrates that high concentrations of *E. coli* rIL-32 α (20 U/ml) or mammalian rIL-32 β (Anjou65, 10 U/ml) induce IL-8 secretion in undifferentiated THP-1 cells.

Figure 7 illustrates that *E. coli* rIL-32 α and mammalian rIL-32 β activities are neutralized by anti-IL-32 Fab (32-4) treatment (mean $\pm$ SEM of three separate experiments). Panel A shows the dose dependent reduction in TNF α secretion by mouse Raw cells cultured in the presence of *E. coli* rIL-32 α (3 U/ml) and an anti-IL32 Fab. Panel B shows that mammalian (Cos7-S) rIL-32 β (2 U/ml) induced mTNF α secretion was inhibited by an anti-IL-32 Fab (40 ng/ml).

Figure 8 depicts the endogenous expression of IL-32 at both mRNA and protein levels. Panel A shows the expression of IL-32 mRNA in various human tissues as determined by Northern blot. The numbers on the left indicate mRNA size. Panel B shows that soluble IL-32 is present in cell culture supernatants as determined by Western blot. A stable A549-R β clone was stimulated for 48h in the presence of IL-18 (50 ng/ml) or IL-1 β (10 ng/ml), in the absence of FCS as indicated. The supernatants were harvested and probed with an affinity-purified anti-IL-32 α antibody.

Figure 9 shows the detection of IL-32 in cell lysates. IL-32 production by Wish cells (panel A), A549-R β cells (panel B) and A549-WT cells (panel C) upon stimulation with IFN γ (100 U/ml), IL-18 (50 ng/ml) or IL-1 β (10 ng/ml) are shown. The data represent one of four independent experiments. IL-32 expression upon transient transfection of Cos7
 5 cells with IL-32 α and IL-32 β cDNAs was detected in cell culture supernatants (panel D) and cell lysates (panel E) by immunoblot. Panel F provides a comparison of IL-32 concentration in supernatants and lysates as measured by ECL. The data represent the mean $\pm$ SEM of five separate experiments.

Figure 10 depicts the measurement of endogenous IL-32 by
 10 electrochemiluminescence (ECL). In panel A, IL-32 was detected by ECL assay in the same samples used for immunoblot in Figure 9B. In panel B, IL-32 was detected in the supernatant of a human NK cell line after treatment with IL-12, IL-18, or IL-12 plus IL-18. The data represent mean $\pm$ SEM of 4 separate experiments. In panel C, ConA induced-IL-32 was detected in both the supernatant and lysate of human PBMC (N = 7).

15 Figure 11 depicts IL-32 α -induced I κ B degradation (panel A), and p38 MAPK phosphorylation (panel B), after IL-32 α (20 U/ml) treatment of mouse Raw 264.7 macrophage cells. For normalization purposes, the membrane was probed with goat anti-actin or rabbit anti-p38 MAPK.

Figure 12 provides amino acid sequences of equine IL-32 α (SEQ ID NO:18) and IL-
 20 32 β (SEQ ID NO:16) in panels A and C respectively, as well as cDNA sequences of equine IL-32 α (SEQ ID NO:19) and IL-32 β (SEQ ID NO:20) in panels B and D, respectively.

Figure 13 provides amino acid sequences of bovine IL-32 β (SEQ ID NO:17) and IL-
 32 γ (SEQ ID NO:22) in panels A and C respectively, as well as cDNA sequences of bovine IL-32 β (SEQ ID NO:21) and IL-32 γ (SEQ ID NO:23) in panels B and D, respectively.

25 Figure 14 provides the amino acid (SEQ ID NO:24) and cDNA (SEQ ID NO:25) sequences of ovine IL-32 α in panels A and B respectively. Also provided are the amino acid (SEQ ID NO:26) and cDNA (SEQ ID NO:27) sequences of swine IL-32 α in panels C and D, respectively.

30 Definitions

To facilitate an understanding of the present invention, a number of terms and phrases are defined below:

The term "gene" refers to a nucleic acid (*e.g.*, DNA) sequence that comprises coding sequences necessary for the production of a polypeptide or precursor or RNA (*e.g.*, tRNA, siRNA, rRNA, etc.). The polypeptide can be encoded by a full length coding sequence or by any portion of the coding sequence so long as the desired activity or functional properties (*e.g.*, enzymatic activity, ligand binding, signal transduction, etc.) of the full-length or fragment are retained. The term also encompasses the coding region of a structural gene and the sequences located adjacent to the coding region on both the 5' and 3' ends, such that the gene corresponds to the length of the full-length mRNA. The sequences that are located 5' of the coding region and which are present on the mRNA are referred to as 5' untranslated sequences. The sequences that are located 3' or downstream of the coding region and that are present on the mRNA are referred to as 3' untranslated sequences. The term "gene" encompasses both cDNA and genomic forms of a gene. A genomic form or clone of a gene contains the coding region, which may be interrupted with non-coding sequences termed "introns" or "intervening regions" or "intervening sequences." Introns are removed or "spliced out" from the nuclear or primary transcript, and are therefore absent in the messenger RNA (mRNA) transcript. The mRNA functions during translation to specify the sequence or order of amino acids in a nascent polypeptide.

In particular, the terms "TAIF gene" and "IL-32 gene" refers to the full-length IL-32 nucleotide sequence. However, it is also intended that the term encompass fragments of the IL-32 nucleotide sequence, as well as other domains (*e.g.*, functional domains) within the full-length IL-32 nucleotide sequence. Furthermore, the terms "IL-32 gene," "IL-32 nucleotide sequence," and "IL-32 polynucleotide sequence" encompass DNA, cDNA, and RNA sequences.

The term "plasmid" as used herein, refers to a small, independently replicating, piece of DNA. Similarly, the term "naked plasmid" refers to plasmid DNA devoid of extraneous material typically used to affect transfection. As used herein, a "naked plasmid" refers to a plasmid substantially free of calcium-phosphate, DEAE-dextran, liposomes, and/or polyamines.

As used herein, the term "purified" refers to molecules (polynucleotides or polypeptides) that are removed from their natural environment, isolated or separated. "Substantially purified" molecules are at least 50% free, preferably at least 75% free, and more preferably at least 90% free from other components with which they are naturally associated.

The term "recombinant DNA" refers to a DNA molecule that is comprised of segments of DNA joined together by means of molecular biology techniques. Similarly, the term "recombinant protein" refers to a protein molecule that is expressed from recombinant DNA.

5 The term "fusion protein" as used herein refers to a protein formed by expression of a hybrid gene made by combining two gene sequences. Typically this is accomplished by cloning a cDNA into an expression vector in frame with an existing gene. The fusion partner may act as a reporter (*e.g.*, β gal), may provide a tool for isolation purposes (*e.g.*, GST) or may increase the half-life of the protein *in vivo* (*e.g.*, IgG Fc).

10 Suitable systems for production of recombinant proteins include but are not limited to prokaryotic (*e.g.*, *Escherichia coli*), yeast (*e.g.*, *Saccaromyces cerevisiae*), insect (*e.g.*, baculovirus), mammalian (*e.g.*, Chinese hamster ovary), plant (*e.g.*, safflower), and cell-free systems (*e.g.*, rabbit reticulocyte).

15 As used herein, the term "coding region" refers to the nucleotide sequences that encode the amino acid sequences found in the nascent polypeptide as a result of translation of an mRNA molecule. The coding region is bounded in eukaryotes, on the 5' side by the nucleotide triplet "ATG" that encodes the initiator methionine and on the 3' side by one of the three triplets which specify stop codons (*i.e.*, TAA, TAG, and TGA).

20 Where amino acid sequence is recited herein in reference to a naturally occurring protein molecule, the term "amino acid sequence" and like terms such as "polypeptide" or "protein" are not meant to limit the amino acid sequence to the complete, native amino acid sequence associated with the recited protein molecule.

25 The term "wild-type" refers to a gene or gene product that has the characteristics of that gene or gene product when isolated from a naturally occurring source. A wild type gene is that which is most frequently observed in a population and is thus arbitrarily designed the "normal" or "wild-type" form of the gene.

30 As used herein, the terms "mutant," "polymorphism," and "variant," in reference to a gene or gene product, refer to alterations in sequence and/or functional properties (*i.e.*, different characteristics) when compared to the wild-type gene or parental gene product. In some preferred embodiments, the term mutant refers to a gene or gene product that differs from a parental gene or gene product as a result of mutation. It is noted that naturally occurring and induced mutants can be isolated; these are identified by the fact that they have altered characteristics when compared to the wild-type gene or parental gene product. In

addition, mutant genes can be artificially (*e.g.*, site-directed mutagenesis) or synthetically produced in the laboratory.

The terms "interleukine-32," "IL-32," "TAIF," "tumor necrosis factor-alpha inducing factor," "NK4," and "natural killer cell transcript 4," as used herein refer to a human IL-32 gene (*e.g.*, *Homo sapiens* - SEQ IDNO:11), and its gene products (*e.g.*, wild type alpha, beta, gamma and delta isoforms, and variants thereof). IL-32 variants that differ from the wild type IL-32 sequences in fewer than 20% of the residues (preferably 10% or fewer, more preferably 5% or fewer and most preferably 1% or fewer), are also suitable for use in the methods and compositions of the present invention (this includes but is not limited to the gene product corresponding to the murine cDNA fragment disclosed as SEQ ID NO:12). In contrast, the terms NK4, TAIF and IL-32 as used herein, do not refer to the internal fragment of the hepatocyte growth factor (HGF), designated as HGF/NK4 (or also simply NK4, for N-terminal hairpin domain and subsequent four-kringle domains), which is a specific antagonist of HGF (Date *et al.*, *FEBS Lett*, 420:1-6, 1997).

As used herein, the term "hybridization" is used in reference to the pairing of complementary nucleic acids. Hybridization and the strength of hybridization (*i.e.*, the strength of the association between the nucleic acids) is impacted by such factors as the degree of complementarity between the nucleic acids, stringency of the conditions involved, the T_m of the formed hybrid, and the G:C ratio within the nucleic acids.

As used herein, the term " T_m " is used in reference to the "melting temperature." The melting temperature is the temperature at which a population of double-stranded nucleic acid molecules becomes half dissociated into single strands. The equation for calculating the T_m of nucleic acids is well known in the art. As indicated by standard references, a simple estimate of the T_m value may be calculated by the equation: $T_m = 81.5 + 0.41(\% G + C)$, when a nucleic acid is in aqueous solution at 1 M NaCl (*See e.g.*, Anderson and Young, "Quantitative Filter Hybridization," in Nucleic Acid Hybridization, 1985). Other references include more sophisticated computations that take structural as well as sequence characteristics into account for the calculation of T_m .

As used herein the term "stringency" is used in reference to the conditions of temperature, ionic strength, and the presence of other compounds such as organic solvents, under which nucleic acid hybridizations are conducted. Those skilled in the art will recognize that "stringency" conditions may be altered by varying the parameters just described either individually or in concert. With "high stringency" conditions, nucleic acid base pairing will occur only between nucleic acid fragments that have a high frequency of

complementary base sequences (*e.g.*, hybridization under "high stringency" conditions may occur between homologs with about 85-100% identity, preferably about 70-100% identity). With medium stringency conditions, nucleic acid base pairing will occur between nucleic acids with an intermediate frequency of complementary base sequences (*e.g.*, hybridization under "medium stringency" conditions may occur between homologs with about 50-70% identity). Thus, conditions of "weak" or "low" stringency are often required with nucleic acids that are derived from organisms that are genetically diverse, as the frequency of complementary sequences is usually less.

The terms "high stringency conditions" and "stringent conditions" when used in reference to nucleic acid hybridization comprise conditions equivalent to binding or hybridization at 42 °C in a solution consisting of 5X SSPE (43.8 g/l NaCl, 6.9 g/l NaH₂PO₄ H₂O and 1.85 g/l EDTA, pH adjusted to 7.4 with NaOH), 0.5% SDS, 5X Denhardt's reagent and 100 µg/ml denatured salmon sperm DNA followed by washing in a solution comprising 0.1X SSPE, 1.0% SDS at 42 °C when a probe of about 500 nucleotides in length is employed.

"Medium stringency conditions" when used in reference to nucleic acid hybridization comprise conditions equivalent to binding or hybridization at 42 °C in a solution consisting of 5X SSPE (43.8 g/l NaCl, 6.9 g/l NaH₂PO₄ H₂O and 1.85 g/l EDTA, pH adjusted to 7.4 with NaOH), 0.5% SDS, 5X Denhardt's reagent and 100 µg/ml denatured salmon sperm DNA followed by washing in a solution comprising 1.0X SSPE, 1.0% SDS at 42 °C when a probe of about 500 nucleotides in length is employed.

"Low stringency conditions" comprise conditions equivalent to binding or hybridization at 42 °C in a solution consisting of 5X SSPE (43.8 g/l NaCl, 6.9 g/l NaH₂PO₄ H₂O and 1.85 g/l EDTA, pH adjusted to 7.4 with NaOH), 0.1% SDS, 5X Denhardt's reagent [50X Denhardt's contains per 500 ml: 5 g Ficoll (Type 400, Pharmacia), 5 g BSA (Fraction V; Sigma)] and 100 µg/ml denatured salmon sperm DNA followed by washing in a solution comprising 5X SSPE, 0.1% SDS at 42 °C when a probe of about 500 nucleotides in length is employed.

As used herein, the term "Northern blot" refers to methods for transferring denatured RNA onto a solid support for use in a subsequent hybridization assay. Total RNA or polyA-enriched RNA is typically electrophoresed in an agarose gel, transferred to a membrane and probed with a radioactively-labeled DNA or RNA fragment to detect specific RNA sequences. Northern blots are routinely used in the art (*See, e.g.*, Thomas,

WO 2005/047478

PCT/US2004/037578

Proc Natl Acad Sci USA 77:5201-5205, 1980; and Ausubel *et al.* (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., New York, 1994).

The term "Southern blot," as used herein, refers to methods for transferring denatured DNA, which has been fractionated by agarose gel electrophoresis, onto a solid support, for use in a subsequent hybridization assay. These methods typically entail the digestion of genomic DNA with a suitable restriction enzyme prior to agarose gel electrophoresis, transfer of the DNA to a membrane and incubation with a radioactively-labeled DNA or RNA fragment for detection of specific DNA sequences. Southern blots are routinely used in the art (See, Southern, *J Mol Biol* 98:503-517, 1975; and Ausubel *et al.*, *supra*, 1994).

As used herein, the term "polymerase chain reaction (PCR)" refers to a method for increasing the concentration of a segment of a target sequence in a DNA mixture without cloning or purification (See, *e.g.*, U.S. Patent Nos. 4,683,195, 4,683,202, and 4,965,188). This process for amplifying the target sequence consists of introducing a large excess of two oligonucleotide primers to the DNA mixture containing the desired target sequence, followed by a precise sequence of thermal cycling in the presence of a DNA polymerase. The two primers are complementary to their respective strands of the double stranded target sequence. To effect amplification, the mixture is denatured and the primers then annealed to their complementary sequences within the target molecule. Following annealing, the primers are extended with a polymerase so as to form a new pair of complementary strands. The steps of denaturation, primer annealing and polymerase extension can be repeated many times (*i.e.*, denaturation, annealing and extension constitute one "cycle") to obtain a high concentration of an amplified segment of the desired target sequence. The length of the amplified segment of the desired target sequence is determined by the relative positions of the primers with respect to each other, and therefore, this length is a controllable parameter. By virtue of the repeating aspect of the process, the method is referred to as the "polymerase chain reaction" (hereinafter "PCR"). Because the desired amplified segments of the target sequence become the predominant sequences (in terms of concentration) in the mixture, they are the to be "PCR amplified." When the template is RNA, a reverse transcription (RT) step is completed prior to the amplification cycles. Thus, this variation is termed "RT-PCR."

The term "antibody" refers to polyclonal and monoclonal antibodies. Polyclonal antibodies which are formed in the animal as the result of an immunological reaction against a protein of interest or a fragment thereof, can then be readily isolated from the

blood using well-known methods and purified by column chromatography, for example. Monoclonal antibodies can also be prepared using known methods (*See, e.g.,* Winter and Milstein, *Nature*, 349, 293-299, 1991). As used herein, the term "antibody" encompasses recombinantly prepared, and modified antibodies and antigen-binding fragments thereof, such as chimeric antibodies, humanized antibodies, multifunctional antibodies, bispecific or oligo-specific antibodies, single-stranded antibodies and F(ab) or F(ab)₂ fragments. The term "reactive" in used in reference to an antibody indicates that the antibody is capable of binding an antigen of interest. For example, an IL-32-reactive antibody is an antibody that binds to IL-32 or to a fragment of IL-32.

10 The term "dominant-negative mutant" refers to molecules that lack wild type activity, but which effectively compete with wild type molecules for substrates, receptors, etc., and thereby inhibit the activity of the wild type molecule. In preferred embodiments, the term "IL-32 dominant negative mutant" refers to a IL-32 mutant protein which competes with the wild type IL-32 protein for IL-32 receptors, but which fails to induce downstream effects such as degradation of I κ B, phosphorylation of p38 MAPK, and TNF α production. Suitable dominant-negative IL-32 variants may be selected from libraries of random IL-32 mutants or may be designed rationally, as has been described in the TNF system (Steed *et al., Science*, 301:1895-1898, 2003).

20 The term "portion" when used in reference to a nucleotide sequence refers to fragments of that sequence, which range in size from 10 nucleotides to the entire nucleotide sequence minus one nucleotide.

25 As used herein, the term "biologically active" refers to a molecule having structural, regulatory and or biochemical functions of a wild type IL-32 molecule. In some instances, the biologically active molecule is a homolog of a mammalian IL-32 molecule, while in other instances the biologically active molecule is a portion of a mammalian IL-32 molecule. Other biologically active molecules that find use in the compositions and methods of the present invention include but are not limited to mutant (*e.g.,* variants with at least one deletion, insertion or substitution) mammalian IL-32 molecules. Biological activity is determined for example, by measuring TNF α -induction *in vitro* as described in the experimental examples.

30 As used herein the term "animal" refers to any member of the kingdom Animalia, which includes living things that have cells differing from plant cells with regard to the absence of a cell wall and chlorophyll and the capacity for spontaneous movement.

Preferred embodiments of the present invention are primarily directed to vertebrate (backbone or notochord) members of the animal kingdom.

The terms "patient" and "subject" refer to a mammal (human or animal) that is a candidate for receiving medical treatment.

5 The term "control" refers to subjects or samples which provide a basis for comparison for experimental subjects or samples. For instance, the use of control subjects or samples permits determinations to be made regarding the efficacy of experimental procedures. In some embodiments, the term "control subject" refers to animals that receive a mock treatment (*e.g.*, PBS alone or normal rabbit IgG in saline).

10 The terms "sample" and "specimen" are used in their broadest sense. On the one hand, they are meant to include a specimen or culture. On the other hand, they are meant to include both biological and environmental samples. These terms encompasses all types of samples obtained from humans and other animals, including but not limited to, body fluids such as urine, blood, fecal matter, cerebrospinal fluid, semen, saliva, and wound exudates,
15 as well as solid tissue. However, these examples are not to be construed as limiting the sample types applicable to the present invention.

The term "leukocyte" as used herein, refers to cells called white blood cells that help the body fight infections and other diseases, and include for instance granulocytes (*e.g.*, neutrophils, eosinophils, basophils), mononuclear phagocytes, and lymphocytes (*e.g.*, B
20 cells, T cells, natural killer cells).

As used herein, the term "monocyte" refers to a mononuclear phagocyte circulating in blood that will later emigrate into tissue and differentiate into a macrophage. The term "macrophage" refers to relatively long-lived phagocytic cells of mammalian tissues, derived from blood monocytes. Macrophages from different sites have distinctly different
25 properties. Main types are peritoneal and alveolar macrophages, tissue macrophages (histiocytes), Kupffer cells of the liver and osteoclasts. Macrophages play an important role in killing some bacteria, protozoa and tumour cells, in releasing substances that stimulate other cells of the immune system, and presenting processed antigen to T lymphocytes.

The term "inflammation" as used herein, refers to the tissue response to trauma,
30 characterized by increased blood flow and entry of leukocytes into the tissues, resulting in swelling, redness, elevated temperature and pain.

As used herein, the term "symptom" refers to any subjective evidence of disease or of a patient's condition (*e.g.*, a change in a patient's condition indicative of some bodily or mental state).

For instance, the phrase "symptoms of inflammation" in the context of inflammatory bowel disease (IBD) is herein defined to include, but is not limited to symptoms such as abdominal pain, diarrhea, rectal bleeding, weight loss, fever, loss of appetite, and other more serious complications, such as dehydration, anemia and malnutrition. A number of such symptoms are subject to quantitative analysis (*e.g.*, weight loss, fever, anemia, etc.). Some symptoms are readily determined from a blood test (*e.g.*, anemia) or a test that detects the presence of blood (*e.g.*, rectal bleeding).

Similarly, the phrase "under conditions such that the symptoms are reduced" in the context of IBD refers to any degree of qualitative or quantitative reduction in detectable symptoms of IBD, including but not limited to, a detectable impact on the rate of recovery from disease (*e.g.*, rate of weight gain), or the reduction of at least one of the following symptoms: abdominal pain, diarrhea, rectal bleeding, weight loss, fever, loss of appetite, dehydration, anemia, distention, fibrosis, inflamed intestines and malnutrition.

The term "autoimmune disease" includes but is not limited to the following diseases: multiple sclerosis, myasthenia gravis, autoimmune neuropathies (such as Guillain-Barré), autoimmune uveitis, Crohn's disease, ulcerative colitis, primary biliary cirrhosis, autoimmune hepatitis, autoimmune hemolytic anemia, pernicious anemia, autoimmune thrombocytopenia, type 1 diabetes mellitus, Grave's disease, Hashimoto's thyroiditis, autoimmune oophoritis and orchitis, temporal arteritis, anti-phospholipid syndrome, Vasculitides (such as Wegener's granulomatosis), Behcet's disease, rheumatoid arthritis, systemic lupus erythematosus, scleroderma, polymyositis, dermatomyositis, spondyloarthropathies (such as ankylosing spondylitis), Sjogren's syndrome, psoriasis, dermatitis herpetiformis, pemphigus vulgaris, and vitiligo.

As used herein, the terms "rheumatoid arthritis" and "RA" refer to a chronic inflammatory disease in which there is destruction of joints. RA is considered to be an autoimmune disorder in which immune complexes are formed in joints and excite an inflammatory response (complex mediated hypersensitivity). Cell-mediated (type IV) hypersensitivity also occurs, resulting in the accumulation of macrophages and leading to the destruction of the synovial lining.

The terms "IBD" and "inflammatory bowel disease," as used herein, are general terms that encompass several disease processes, most commonly, ulcerative colitis and Crohn's disease.

As used herein, the term "Crohn's disease" refers to an inflammatory disease of the gastrointestinal tract. Common symptoms include recurrent abdominal pains, fever, nausea, vomiting, weight loss, and diarrhea that is occasionally bloody.

The terms "compound" and "drug candidate" refers to any chemical or biological entity (*e.g.*, including pharmaceuticals, drugs, and the like) that can be used to treat or prevent a disease, illness, sickness, or disorder of bodily function. Compounds comprise both known and potential therapeutic compounds. A compound can be determined to be therapeutic by screening, *e.g.*, using the screening methods of the present invention. A "known therapeutic compound" refers to a therapeutic compound that has been shown (*e.g.*, through animal trials or prior experience with administration to humans) to be effective in such treatment or prevention.

As used herein, the term "agonist" refers to molecules or compounds that mimic the action of a "native" or "natural" compound. Agonists may be homologous to these natural compounds in respect to conformation, charge or other characteristics. Thus, agonists may be recognized by receptors expressed on cell surfaces. This recognition may result in physiologic and/or biochemical changes within the cell, such that the cell reacts to the presence of the agonist in the same manner as if the natural compound was present. Agonists may include proteins, nucleic acids, carbohydrates, or any other molecules that bind or interact with IL-32 binding protein(s).

As used herein, the terms "antagonist" and "inhibitor" refer to molecules or compounds that inhibit the action of a "native" or "natural" compound. Antagonists may or may not be homologous to these natural compounds in respect to conformation, charge or other characteristics. Thus, antagonists may be recognized by the same or different receptors that are recognized by an agonist. Antagonists may have allosteric effects, which prevent the action of an agonist (*e.g.*, prevent native IL-32 from binding to IL-32 receptors). In contrast to the agonists, antagonistic compounds do not result in physiologic and/or biochemical changes within the cell such that the cell reacts to the presence of the antagonist in the same manner as if the natural compound was present. Antagonists and inhibitors may include proteins, nucleic acids, carbohydrates, or any other molecules that bind or interact with IL-32 binding proteins or which prevent formation of functional IL-32 multimers.

As used herein, the term modulate, refers to a change or an alteration in biological activity. Modulation may be an increase or a decrease in protein activity, a change in

binding characteristics, or any other change in the biological, functional, or immunological properties associated with the activity of a protein or other structure of interest.

DESCRIPTION OF THE INVENTION

5 Interleukin-18 (IL-18) is multifunctional cytokine having roles in both the innate and adapted immune responses. IL-18 has been studied for its effect in the broad spectrum of Th1 or Th2 related autoimmune diseases (Okamura *et al.*, *Nature*, 378:88-91, 1995, Nakanishi *et al.*, *Cytokine Growth Factor Rev*, 12:53-72, 2001; and Okamura *et al.*, *Adv Immunol*, 70:281-312, 1998). IL-1 and IL-18 belong to the IL-1 family, which share
10 structural similarity, require caspase-1 for processing (Bazan *et al.*, *Nature*, 379:591, 1996; and Gu *et al.*, *Science*, 275:206-209, 1997). IL-18 also triggers similar signaling pathways including recruiting IL-1 receptor-associated kinases (IRAKs), the formation of IRAK complexes with the tumor necrosis factor (TNF) receptor-associated factor-6, and activation of the cascade of I κ B α /NF- κ B (Kojima *et al.*, *Biochem Biophys Res Commun*, 244:183-186,
15 1998; Matsumoto *et al.*, *Biochem Biophys Res Commun*, 234:454-457, 1997; and Robinson *et al.*, *Immunity*, 7:571-581, 1997). Members of the IL-1 cytokine family are thought to play pathological roles in autoimmune and inflammatory diseases, since blocking IL-1 and IL-18 activities reduces disease severity in subjects with rheumatoid arthritis and systemic inflammation (Arend, *Adv Immunol*, 54:167-227, 1993; Dinarello, *Blood*, 87:2095-2147,
20 1996; Novick *et al.*, *Immunity*, 10:127-136, 1999; and Banda *et al.*, *J Immunol*, 170:2100-2105, 2003). However, there is an obstacle to studying IL-18-inducible genes, since only a few cell lines respond to IL-18, for example human NK and KG-1 cell lines. In addition, although these cell lines respond to IL-18, they require co-stimulatory factors, such as IL-2, IL-12, or IL-15, respectively, in order to manifest IL-18 responsiveness (Ahn *et al.*, *J*
25 *Immunol*, 159:2125-2131, 1997, Ohtsuki *et al.*, *Anticancer Res*, 17:3253-3258, 1997; Lauwerys *et al.*, *J Immunol*, 165:1847-1853, 2000; and Hoshino *et al.*, *J Immunol*, 162:51-59, 1999). The requirement for a co-stimulatory factor prevents an independent assessment of IL-18 induction of gene expression.

IL-18 has two known receptor chains, a ligand binding IL-18R α chain and a signal
30 transducing IL-18R β chain. Both chains of the IL-18 receptor belong to the IL-1 receptor family and consist of three Ig-like domains in the extracellular region (Kato *et al.*, *Nat Struct Biol*, 10:966-971, 2003; and Yamamoto *et al.*, *Biochem Biophys Res Commun*, 317:181-186, 2004). An additional component involved in IL-18 regulation is the IL-18 binding protein (IL-18BP). IL-18BP is not a part of the IL-18 signaling complex but rather

antagonizes IL-18 activity (Kim *et al.*, *Proc Natl Acad Sci USA*, 97:1190-1195, 2000; and Novick *et al.*, *Immunity*, 10:127-136, 1999). IL-18BP is a secreted receptor-like molecule consisting of a single Ig-like domain. IL-18BP shares significant homology with viral proteins, and neutralizes the biological activities of human IL-18 (Xiang and Moss,
5 *Virology*, 257:297-302, 1999; Reading and Smith, *J Virol*, 77:9960-9968, 2003; Esteban and Buller, *Virology*, 323:197-207, 2004; and Esteban *et al.*, *J Gen Virol*, 85:1291-1299, 2004).

In the absence of co-stimulants, non-immune cells do not respond to IL-18, due to little or no expression of the IL-18R β chain (Thomassen *et al.*, *J Interferon Cytokine Res*,
10 18:1077-1088, 1998; and Chandrasekar *et al.*, *Biochem Biophys Res Commun*, 303:1152-1158, 2003). Therefore, it was necessary to generate a stable clone expressing the IL-18R β chain, since this receptor component is required for transmitting an IL-18 signal (Thomassen *et al.*, *supra*, 1998; and Kim *et al.*, *J Immunol*, 166:148-154, 2001). Stable expression of the IL-18R β chain was accomplished in the human lung carcinoma A549 cell
15 (A549-R β) , which made these cells responsive to IL-18 even in the absence of co-stimulation. The stable A549-R β clone was used to identify IL-18 inducible genes by microarray analysis. The microarray study described in Example 1, revealed the IL-18 induction of a cytokine-like molecule that was described 12 years ago as natural killer cell transcript 4 or simply NK4 (Dahl *et al.*, *J Immunol*, 148:597-603, 1992). It should be noted
20 that the term NK4 is presently used to be described a variant of hepatocyte growth factor (Martin *et al.*, *J Cell Physiol*, 192:268-275, 2002). There is no sequence homology between the hepatocyte growth factor variant and the NK4/IL-32 transcript. More recently, increased expression of NK4 has been reported in PBMC from patients receiving high-dose IL-12 therapy for malignant melanoma (Panelli *et al.*, *Genome Biol*, 3(7): RESEARCH0035,
25 2002), but the function of NK4 has remained unknown until development of the present invention.

As is described in detail in the experimental examples, a novel inflammatory cytokine termed interleukin-32 (IL-32) or the tumor necrosis factor alpha-inducing factor (TAIF) has been identified through a microarray analysis of A549-R β cells. In addition, the
30 gene structure, expression pattern, and function of IL-32 have been elucidated. Prior to development of the present invention, a single isoform of IL-32 (TAIF γ /NK4) had been described as a lymphocyte transcript of unknown function (Dahl *et al.*, *J Immunol*, 148:597-603, 1992), which is transcribed from human chromosome 16p13.3 within or adjacent to the Familial Mediterranean fever locus (Bernot *et al.*, *Genomics*, 50:147-160, 1998). As is

shown herein, although IL-32 lacks sequence homology to the known cytokine families, it is clearly a cytokine by virtue of its ability to induce TNF α secretion and to signal through the classical pathways of known proinflammatory cytokines. Moreover, IL-32 is contemplated to be a member of the IL-1 cytokine family (Ghayur *et al.*, *Nature*, 386:619-623, 1997;

5 Cerretti *et al.*, *Science*, 256:97-100, 1992, and Kuida *et al.*, *Science*, 267:2000-2003, 1995), due to similarities in IL-32 regulation, and to the lack of a clear signal peptide. In addition, like IL-1 and IL-18, IL-32 is secreted primarily from stimulated cells. As demonstrated for the first time herein, IL-32 triggered typical proinflammatory cytokine signal pathways, NF κ B and p38 MAPK, thereby inducing production of the proinflammatory cytokine,
10 TNF α . Similarly, IL-1 β , IL-18, and LPS induced IL-32 expression, and unlike the original description of NK4 in activated NK or T-cells, ubiquitous IL-32 expression was observed in various organs. The IL-32 expression pattern resembles that of IL-15, which is also produced by a wide variety of tissues. This is in contrast to IL-2, which is exclusively produced by activated T cells (Bamford *et al.*, *Proc Natl Acad Sci USA*, 93:2897-2902,
15 1996; and Grabstein *et al.*, *Science*, 264:965-968, 1994).

The induction of TNF α by IL-32 indicates that this cytokine has an important role in autoimmune/inflammatory disease pathology. The high level of IL-32 observed in the circulation of some patients with sepsis (as compared to that observed in the sera of healthy individuals) has also not been documented before development of the present invention.

20 Thus, blocking IL-32 activity is contemplated to be a highly effective therapeutic approach in various autoimmune diseases, similar to successful TNF α blocking strategies (Davis *et al.*, *Ann Rheum Dis*, 59Suppl1:i41-3, 2000; and Shanahan and St Clair, *Clin Immunol*, 103:231-242, 2002). It is also contemplated that IL-32-based compositions will find use in the treatment of cancer and in other diseases where induction of cell death or apoptosis
25 comprising TNF α production is beneficial.

In addition, it is contemplated that the mouse Raw 264.7 macrophage cell line expresses an IL-32 receptor, which is activated when bound by IL-32 (extracellular IL-32 function). However, it is also contemplated that IL-32 has an intracellular function, since IL-32 was also detected in cell lysates. Thus, it is contemplated that IL-32 is active as both
30 an intracellular and as an extracellular protein similar to IL-1 α (Stevenson *et al.*, *Proc Natl Acad Sci USA*, 94:508-513, 1997) and high mobility group-1 (Wang *et al.*, *Science*, 285:248-251, 1999).

The secretion of IL-32 in the absence of a clear signal peptide is characteristic of several cytokine families (Cerretti *et al.*, *Science*, 256:97-100, 1992; Kuida *et al.*, *Science*,

267:2000-2003, 1995; and Ghayur *et al.*, *Nature*, 386:619-623, 1997). Similar to IL-1 β and IL-18, IL-32 is secreted as a soluble protein in cell culture media of stimulated primary cells, as well as cell lines. IL-32 does not possess potential caspase-1 cleavage sites by primary sequence analysis, and although new splice variants have been found, none possess
5 a typical hydrophobic signal peptide at the N-terminus.

The existence of IL-32 in multiple species is evidence of an evolutionally conserved molecule that is contemplated to play an important function in regulation of inflammation. Fewer IL-32 isoforms were identified in other mammalian species, although this result may reflect the relative lack of ESTs for these species. Similarly an extensive search for a mouse
10 IL-32 homologue was unsuccessful. However, it is possible that the putative murine IL-32 gene has very low homology with the human gene, since equine and bovine IL-32 homologs share only 31.8-28.1 % sequence identity with human IL-32.

Some preferred embodiments of the present invention are described in the following sections: (I) IL-32 Polynucleotides; (II) IL-32 Polypeptides; (III) IL-32 Antibodies; (IV)
15 Pharmaceutical Compositions Comprising IL-32 Polynucleotides, Polypeptides or Antibodies; and (V) Methods for Identifying IL-32 Inhibitors.

I. IL-32 Polynucleotides

The present invention provides nucleic acids encoding IL-32 proteins, homologs,
20 variants, and mutants (*e.g.*, SEQ ID NOs: 3, 4, 6). In some embodiments, the present invention provides polynucleotide sequences that are capable of hybridizing to SEQ ID NOs: 3, 4, 6 under conditions high stringency, as long as the polynucleotide sequence capable of hybridizing encodes a protein that retains the TNF α -inducing activity of the naturally occurring IL-32 gene. In some embodiments, the protein that retains the TNF α -
25 inducing activity of naturally occurring IL-32 is 80% homologous to wild-type IL-32, preferably 90% homologous to wild-type IL-32, more preferably 95%, and most preferably 99% homologous to wild-type IL-32. In particularly preferred embodiments, the protein that retains IL-32 biological activity comprises a nucleic acid sequence encoding the contiguous amino acid sequence LKARMHQAIERFYDKMQNAESGRGQV (SEQ ID
30 NO:13), and which is 99% homologous to wild-type IL-32 (IL-32 α , IL-32 β , IL-32 γ , IL-32 δ). In some preferred embodiments, the nucleic acid sequence does not encode the amino acid sequence set forth in SEQ ID NO:14. In a subset of these embodiments, the nucleic acid sequence comprises the sequence set forth in SEQ ID NO:15 (exon 3 adjacent to exon 4). In preferred embodiments, hybridization conditions are based on the melting

temperature (T_m) of the nucleic acid binding complex and confer a defined "stringency" as explained above in the definition section.

In other embodiments of the present invention, alleles of IL-32 are provided. In preferred embodiments, alleles result from a mutation, (*i.e.*, a change in the nucleic acid sequence) and generally produce altered mRNAs or polypeptides whose structure or function may or may not be altered. Any given gene may have none, one or many allelic forms. Common mutational changes that give rise to alleles are generally ascribed to deletions, additions or substitutions of nucleic acids. Each of these types of changes may occur alone, or in combination with the others, and at the rate of one or more times in a given sequence.

In still other embodiments of the present invention, the nucleotide sequences of the present invention may be engineered in order to alter a IL-32 coding sequence for a variety of reasons, including but not limited to, alterations which modify the cloning, processing and/or expression of the gene product. For example, mutations may be introduced using techniques that are well known in the art (*e.g.*, site-directed mutagenesis to insert new restriction sites, to alter glycosylation patterns, to change codon preference, etc.).

A modified peptide can be produced in which the nucleotide sequence encoding the polypeptide has been altered, such as by substitution, deletion, or addition. In some preferred embodiments, these modifications do not significantly alter the TNF α -inducing activity of the modified IL-32 (*e.g.*, IL-32 agonists), while in other preferred embodiments, these modifications eliminate TNF α -inducing activity of the modified IL-32 (*e.g.*, IL-32 antagonists). In other words, any given construct can be evaluated in order to determine whether it is a member of the genus of modified or variant IL-32's of the present invention as defined functionally, rather than structurally. In preferred embodiments, the activity of variant or mutant IL-32 is evaluated by the methods described in Example 2 (TNF α induction).

Moreover, as described above, variant forms of IL-32 are also contemplated as being equivalent to those peptides and DNA molecules that are set forth in more detail herein. For example, it is contemplated that isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid (*i.e.*, conservative mutations) will not have a major effect on the biological activity of the resulting molecule. Accordingly, some embodiments of the present invention provide variants of IL-32 disclosed herein containing conservative replacements. Conservative replacements are those that take place within a

family of amino acids that are related in their side chains. Genetically encoded amino acids can be divided into four families: (1) acidic (aspartate, glutamate); (2) basic (lysine, arginine, histidine); (3) nonpolar (alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan); and (4) uncharged polar (glycine, asparagine, glutamine, cysteine, serine, threonine, tyrosine). Phenylalanine, tryptophan, and tyrosine are sometimes classified jointly as aromatic amino acids. In similar fashion, the amino acid repertoire can be grouped as (1) acidic (aspartate, glutamate); (2) basic (lysine, arginine, histidine), (3) aliphatic (glycine, alanine, valine, leucine, isoleucine, serine, threonine), with serine and threonine optionally be grouped separately as aliphatic-hydroxyl; (4) aromatic (phenylalanine, tyrosine, tryptophan); (5) amide (asparagine, glutamine); and (6) sulfur-containing (cysteine and methionine) (*e.g.*, Stryer ed., *Biochemistry*, pg. 17-21, 2nd ed, WH Freeman and Co., 1981). Whether a change in the amino acid sequence of a peptide results in a functional homolog can be readily determined, by assessing the ability of the variant peptide to function in a fashion similar to the wild-type protein. Peptides having more than one replacement can readily be tested in the same manner.

More rarely, a variant includes "nonconservative" changes (*e.g.*, replacement of a glycine with a tryptophan). Analogous minor variations can also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues can be substituted, inserted, or deleted without abolishing biological activity can be found using computer programs (*e.g.*, LASERGENE software, DNASTAR Inc., Madison, WI).

II. IL-32 Polypeptides

In other embodiments, the present invention provides IL-32 polynucleotide sequences that encode IL-32 polypeptide sequences. IL-32 polypeptides (*e.g.*, SEQ ID NOs: 7, 8, 10) are described in Figure 4. Other embodiments of the present invention provide fragments, fusion proteins or functional equivalents of these IL-32 proteins. In still other embodiment of the present invention, nucleic acid sequences corresponding to these various IL-32 homologs and mutants may be used to generate recombinant DNA molecules that direct the expression of the IL-32 homologs and mutants in appropriate host cells. In some embodiments of the present invention, the polypeptide may be a naturally purified product, in other embodiments it may be a product of chemical synthetic procedures, and in still other embodiments it may be produced by recombinant techniques using a prokaryotic or eukaryotic host (*e.g.*, by bacterial, yeast, higher plant, insect and mammalian cells in culture). In some embodiments, depending upon the host employed in a recombinant

production procedure, the polypeptide of the present invention may be glycosylated or may be non-glycosylated. In other embodiments, the polypeptides of the invention may also include an initial methionine amino acid residue.

5 In one embodiment of the present invention, due to the inherent degeneracy of the genetic code, DNA sequences other than the polynucleotide sequences of SEQ ID NOs: 3, 4 and 6, which encode substantially the same or a functionally equivalent amino acid sequence, may be used to clone and express IL-32.

A. Vectors for Production of IL-32

10 The polynucleotides of the present invention may be employed for producing polypeptides by recombinant techniques. Thus, for example, the polynucleotide may be included in any one of a variety of expression vectors for expressing a polypeptide. In some embodiments of the present invention, vectors include, but are not limited to, chromosomal, nonchromosomal and synthetic DNA sequences (*e.g.*, derivatives of SV40, bacterial
15 plasmids, phage DNA; baculovirus, yeast plasmids, vectors derived from combinations of plasmids and phage DNA, and viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies). It is contemplated that any vector may be used as long as it is replicable and viable in the host.

In particular, some embodiments of the present invention provide recombinant
20 constructs comprising one or more of the sequences as broadly described above (*e.g.*, SEQ ID NOs: 3, 4, 6). In some embodiments of the present invention, the constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation. In still other embodiments, the heterologous structural sequence is assembled in appropriate phase with translation initiation and
25 termination sequences. In preferred embodiments of the present invention, the appropriate DNA sequence is inserted into the vector using any of a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art.

Large numbers of suitable vectors are known to those of skill in the art, and are
30 commercially available. Such vectors include, but are not limited to, the following vectors:
1) Bacterial -- pQE70, pQE60, pQE-9 (Qiagen), pBS, pD10, phagescript, psiX174, pbluescript SK, pBSKS, pNH8A, pNH16a, pNH18A, pNH46A (Stratagene); ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia); and 2) Eukaryotic -- pWLNEO, pSV2CAT, pOG44, PXT1, pSG (Stratagene) pSVK3, pBPV, pMSG, pSVL (Pharmacia).

Any other plasmid or vector may be used as long as they are replicable and viable in the host. In some preferred embodiments of the present invention, mammalian expression vectors comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation sites, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking non-transcribed sequences. In other
5 embodiments, DNA sequences derived from the SV40 splice, and polyadenylation sites may be used to provide the required non-transcribed genetic elements.

B. Host Cells for Production of IL-32

10 In a further embodiment, the present invention provides host cells containing the above-described constructs. In some embodiments of the present invention, the host cell is a higher eukaryotic cell (*e.g.*, a mammalian or insect cell). In other embodiments of the present invention, the host cell is a lower eukaryotic cell (*e.g.*, a yeast cell). In still other
15 embodiments of the present invention, the host cell can be a prokaryotic cell (*e.g.*, a bacterial cell). Specific examples of host cells include, but are not limited to, *Escherichia coli*, *Salmonella typhimurium*, *Bacillus subtilis*, and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, as well as *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Drosophila* S2 cells, *Spodoptera* Sf9 cells, Chinese hamster ovary (CHO) cells, COS-7 lines of monkey kidney fibroblasts, C127, 3T3, 293, 293T, HeLa
20 and BHK cell lines.

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. In some embodiments, introduction of the construct into the host cell can be accomplished by calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation. Alternatively, in some
25 embodiments, the IL-32 polypeptides are produced using conventional peptide synthesizers.

C. Purification of IL-32

The present invention also provides methods for recovering and purifying IL-32 from recombinant cell cultures including, but not limited to, ammonium sulfate or ethanol
30 precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. In other embodiments of the present invention, protein-refolding steps can be used as necessary, in completing configuration of the mature protein. In still other embodiments of the present invention,

high performance liquid chromatography (HPLC) can be employed for final purification steps.

The present invention further provides polynucleotides having the coding sequence fused in frame to a marker sequence, which allows for purification of the polypeptide of the present invention. A non-limiting example of a marker sequence is a hexahistidine tag which may be supplied by a vector, preferably a pQE-9 vector, which provides for purification of the polypeptide fused to the marker in the case of a bacterial host, or, for example, the marker sequence may be a hemagglutinin (HA) tag when a mammalian host (*e.g.*, COS-7 cells) is used.

D. IL-32 Fusion Proteins

The present invention also provides fusion proteins incorporating all or part of IL-32. Accordingly, in some embodiments of the present invention, the coding sequences for the polypeptide can be incorporated as a part of a fusion gene including a nucleotide sequence encoding a different polypeptide. It is contemplated that this type of expression system will find use under conditions where it is desirable to produce an immunogenic fragment of an IL-32 protein.

In addition to utilizing fusion proteins to enhance immunogenicity, it is widely appreciated that fusion proteins can also facilitate the expression of proteins, such as the IL-32 protein of the present invention. Accordingly, in some embodiments of the present invention, IL-32 can be generated as a glutathione-S-transferase (*i.e.*, GST fusion protein). It is contemplated that such GST fusion proteins will enable easy purification of IL-32, such as by the use of glutathione-derivatized matrices. In another embodiment of the present invention, a fusion gene coding for a purification leader sequence, such as a poly-(His)/enterokinase cleavage site sequence at the N-terminus of the desired portion of IL-32, can allow purification of the expressed IL-32 fusion protein by affinity chromatography using a Ni^{2+} metal resin. In still another embodiment of the present invention, the purification leader sequence can then be subsequently removed by treatment with enterokinase.

E. IL-32 Variants

Still other embodiments of the present invention provide mutant or variant forms of IL-32. It is possible to modify the structure of a peptide having an activity of IL-32 for such purposes as enhancing therapeutic or prophylactic efficacy, or stability (*e.g.*, *ex vivo* shelf

life, and/or resistance to proteolytic degradation *in vivo*). Such modified peptides are considered functional equivalents of peptides having an activity of the subject IL-32 proteins as defined herein. A modified peptide can be produced in which the amino acid sequence has been altered, such as by amino acid substitution, deletion, or addition. In
5 some embodiments, preferred IL-32 variants include IL-32 agonists (*e.g.*, IL-32 variants that possess TNF α -inducing activity), while other preferred IL-32 variants include IL-32 antagonists (*e.g.*, IL-32 variants that do not possess TNF α -inducing activity and that inhibit the TNF α -inducing activity of wild type IL-32 proteins).

10 III. IL-32 Antibodies

Antibodies can be generated to allow for the detection of IL-32 proteins. The antibodies may be prepared using various immunogens. In one embodiment, the immunogen is a human IL-32 peptide to generate antibodies that recognize human IL-32. Such antibodies include, but are not limited to polyclonal, monoclonal, chimeric, single
15 chain, Fab fragments, and Fab expression libraries.

Various procedures known in the art may be used for the production of polyclonal antibodies directed against IL-32. For the production of antibody, various host animals can be immunized by injection with the peptide corresponding to the IL-32 epitope including but not limited to rabbits, mice, rats, sheep, goats, etc. In a preferred embodiment, the
20 peptide is conjugated to an immunogenic carrier (*e.g.*, diphtheria toxoid, bovine serum albumin, or keyhole limpet hemocyanin). Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels (*e.g.*, aluminum hydroxide), surface-active substances (*e.g.*, lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, and potentially useful human adjuvants such as Bacille Calmette-Guerin).
25

For preparation of monoclonal antibodies directed toward IL-32, it is contemplated that any technique that provides for the production of antibody molecules by continuous cell lines in culture will find use with the present invention (*See e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring
30 Harbor, NY). These include but are not limited to the hybridoma technique (Köhler and Milstein, *Nature* 256:495-497, 1975), as well as the trioma technique, the human B-cell hybridoma technique (*See e.g.*, Kozbor *et al.*, *Immunol Today*, 4:72, 1983), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole *et al.*, in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96, 1985).

WO 2005/047478

PCT/US2004/037578

In addition, it is contemplated that techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778 will find use in producing IL-32 specific single chain antibodies. An additional embodiment of the invention utilizes the techniques described for the construction of Fab expression libraries (Huse *et al.*, *Science*, 246:1275-1281, 1989) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity for IL-32.

It is contemplated that any technique suitable for producing antibody fragments will find use in generating antibody fragments that contain the idiotype (antigen binding region) of the antibody molecule. For example, such fragments include but are not limited to:

10 F(ab')₂ fragment that can be produced by pepsin digestion of the antibody molecule; Fab' fragments that can be generated by reducing the disulfide bridges of the F(ab')₂ fragment, and Fab fragments that can be generated by treating the antibody molecule with papain and a reducing agent.

In the production of antibodies, it is contemplated that screening for the desired antibody will be accomplished by techniques known in the art (*e.g.*, radioimmunoassay, ELISA (enzyme-linked immunosorbant assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitation reactions, immunodiffusion assays, *in situ* immunoassays (*e.g.*, using colloidal gold, enzyme or radioisotope labels, for example), Western blots, precipitation reactions, agglutination assays (*e.g.*, gel agglutination assays, hemagglutination assays, etc.), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc.

In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in an immunoassay and are within the scope of the present invention. As is well known in the art, the immunogenic peptide should be provided free of the carrier molecule used in any immunization protocol. For example, if the peptide was conjugated to KLH, it may be conjugated to BSA, or used directly, in a screening assay.

30 The foregoing antibodies can be used in methods known in the art relating to the localization and structure of IL-32 (*e.g.*, for Western blotting), measuring levels thereof in appropriate biological samples, etc. The antibodies can be used to detect IL-32 in a biological sample from an individual. The biological sample can be a biological fluid, such

WO 2005/047478

PCT/US2004/037578

as, but not limited to, blood, serum, plasma, interstitial fluid, urine, cerebrospinal fluid, and the like, containing cells.

The biological samples can then be tested directly for the presence of human IL-32 using an appropriate strategy (*e.g.*, ELISA or radioimmunoassay) and format (*e.g.*,
5 microwells, dipstick etc.). Alternatively, proteins in the sample can be size separated (*e.g.*, by polyacrylamide gel electrophoresis (PAGE), in the presence or not of sodium dodecyl sulfate (SDS), and the presence of IL-32 detected by immunoblotting (Western blotting). Immunoblotting techniques are generally more effective with antibodies generated against a peptide corresponding to an epitope of a protein, and hence, are particularly suited to the
10 present invention.

Particularly preferred embodiments of the present invention comprise IL-32 antibodies for the treatment of autoimmune diseases such as rheumatoid arthritis. IL-32-reactive antibodies that neutralize the TNF α -inducing activity of IL-32 are contemplated
15 relieve at least one RA disease symptom and in especially preferred embodiments are contemplated to slow RA disease progression. Additionally teaching related to the production and use of therapeutic antibodies, which is contemplated to be applicable to the IL-32 system, is found in U.S. Patent No. 6,277,969 and U.S. Patent No. 6,448,380 directed to therapeutic TNF-reactive antibodies.

20 **IV. Pharmaceutical Compositions Comprising IL-32 Nucleic Acids, Peptides, or Antibodies**

The present invention further provides pharmaceutical compositions which may comprise all or portions of IL-32 polynucleotide sequences, IL-32 polypeptides, inhibitors or antagonists of IL-32 bioactivity, including antibodies, alone or in combination with at
25 least one other agent, such as a stabilizing compound, and may be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water.

The methods of the present invention find use in treating autoimmune diseases and cancer. Peptides can be administered to the patient intravenously in a pharmaceutically
30 acceptable carrier such as physiological saline. Standard methods for intracellular delivery of peptides can be used (*e.g.*, delivery via liposome). Such methods are well known to those of ordinary skill in the art. The formulations of this invention are useful for parenteral administration, such as intravenous, subcutaneous, intramuscular, and intraperitoneal.

Therapeutic administration of a polypeptide intracellularly can also be accomplished using gene therapy.

As is well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and interaction with other drugs being concurrently administered.

Accordingly, in some embodiments of the present invention, IL-32 nucleotide and IL-32 amino acid sequences can be administered to a patient alone, or in combination with other nucleotide sequences, drugs or hormones or in pharmaceutical compositions where it is mixed with excipient(s) or other pharmaceutically acceptable carriers. In one embodiment of the present invention, the pharmaceutically acceptable carrier is pharmaceutically inert. In another embodiment of the present invention, IL-32 polynucleotide sequences or IL-32 amino acid sequences may be administered alone to individuals subject to or suffering from a disease.

In other embodiments, the pharmaceutical compositions of the present invention can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral or nasal ingestion by a patient to be treated.

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. For example, an effective amount of IL-32 may be that amount that induces production of TNF α . Determination of effective amounts is well within the capability of those skilled in the art, especially in light of the disclosure provided herein.

For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. Then, preferably, dosage can be formulated in animal models (particularly murine models) to achieve a desirable circulating concentration range that adjusts TNF α levels.

A therapeutically effective dose refers to that amount of IL-32, which ameliorates at least one symptom of the disease state (*e.g.*, cancer) or to that amount of IL-32 antibody or antagonist, which ameliorates at least one symptom of a disease state (*e.g.*, autoimmune disease such as rheumatoid arthritis). Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals (LD₅₀, the dose lethal to 50% of the population; and ED₅₀, the dose

WO 2005/047478

PCT/US2004/037578

therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index, and it can be expressed as the ratio LD_{50}/ED_{50} . Compounds that exhibit large therapeutic indices are preferred. The data obtained from these cell culture assays and additional animal studies can be used in formulating a range of dosage for human use. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration. Normal dosage amounts may vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration.

V. Drug Screening Using IL-32

The present invention provides methods and compositions for using IL-32 as a target for screening drugs that can alter proinflammatory cytokine responses (e.g., $TNF\alpha$ production).

A technique for drug screening provides high throughput screening for compounds having suitable binding affinity to IL-32 peptides and is described in detail in WO 84/03564. Briefly, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are then reacted with IL-32 peptides and washed. Bound IL-32 peptides are then detected by methods well known in the art.

Another technique uses IL-32 antibodies, generated as discussed above. Such antibodies capable of specifically binding to IL-32 peptides compete with a test compound for binding to IL-32. In this manner, the antibodies can be used to detect the presence of any peptide that shares one or more antigenic determinants of the IL-32 peptide.

The present invention contemplates many other means of screening compounds. The examples provided above are presented merely to illustrate a range of techniques available. One of ordinary skill in the art will appreciate that many other screening methods can be used.

In particular, the present invention contemplates the use of cell lines transfected with IL-32 and variants or mutants thereof for screening compounds for activity, and in particular to high throughput screening of compounds from combinatorial libraries (e.g., libraries containing greater than 10^4 compounds). The cell lines of the present invention can be used in a variety of screening methods. In some embodiments, the cells can be used in

second messenger assays that monitor signal transduction following activation of cell-surface receptors. In other embodiments, the cells can be used in reporter gene assays that monitor cellular responses at the transcription/translation level.

5 In second messenger assays, the host cells are preferably transfected as described above with vectors encoding IL-32 or variants or mutants thereof. The host cells are then treated with a compound or plurality of compounds (*e.g.*, from a combinatorial library) and assayed for the presence or absence of a response. It is contemplated that at least some of the compounds in the combinatorial library can serve as agonists, antagonists, activators, or inhibitors of the protein or proteins encoded by the vectors. It is also contemplated that at
10 least some of the compounds in the combinatorial library can serve as agonists, antagonists, activators, or inhibitors of protein acting upstream or downstream of the protein encoded by the vector in a signal transduction pathway.

The cells are also useful in reporter gene assays. Reporter gene assays involve the use of host cells transfected with vectors encoding a nucleic acid comprising transcriptional
15 control elements of a target gene (*i.e.*, a gene that controls the biological expression and function of a disease target) spliced to a coding sequence for a reporter gene. Therefore, activation of the target gene results in activation of the reporter gene product.

EXPERIMENTAL

20 The following examples are provided in order to demonstrate and further illustrate certain preferred embodiments and aspects of the present invention and are not to be construed as limiting the scope thereof.

In the experimental disclosure which follows, the following abbreviations apply: eq (equivalents); M (Molar); μ M (micromolar); N (Normal); mol (moles); mmol (millimoles);
25 μ mol (micromoles); nmol (nanomoles); g (grams); mg (milligrams); μ g (micrograms); ng (nanograms); l or L (liters); ml (milliliters); μ l (microliters); cm (centimeters); mm (millimeters); μ m (micrometers); nm (nanometers); $^{\circ}$ C (degrees Centigrade); U (units), mU (milliunits); min. (minutes); sec. (seconds); % (percent); kb (kilobase); bp (base pair); PCR (polymerase chain reaction).

30 In addition, the following cells and bioassays were employed. The human NK cell line was obtained from Dr. Hans Klingerman (Rush Medical Center, Chicago, IL) and cultured in RPMI1640 medium containing 10% FCS, IL-2 (50 pg/ml) and IL-15 (200 pg/ml) (Peprotech, Rocky Hill, NJ). Mouse macrophage Raw 264.7 cells, human A549 lung carcinoma, monkey Cos7 kidney cells, Anjou65 (subclone of the human fibroblast

293T line), human epithelial Wish cells, and the human monocyte THP-1 cell line were obtained from American Type Culture Collection (ATCC) and maintained according to supplied instructions. Bioassays were performed in 96 well plates.

Briefly, Raw cells (5×10^5 /ml, 0.1 ml/per well), A549-R β cells (2×10^5 /ml, 0.1 ml/per well), and human NK cells (5×10^5 /ml, 0.1 ml/per well) were seeded in 96 well plates and cultured until cells adhered to the plates. For Raw cell assays, spent medium was removed and the cells were stimulated with fresh medium (0.2 ml) containing various concentration of rIL-32 in the presence 5 μ g/ml polymyxin B (Bedford Laboratories, Bedford, OH). For the A549-R β cell assays, spent medium was removed and cells were stimulated with fresh medium (0.2 ml) containing IL-18 (produced as described by Kim *et al.*, *J Immunol*, 166:148-154, 2001). For the NK cell assays, cells were stimulated with IL-12 (Peprotech) or IL-18, or both cytokines. THP-1 cells (5×10^5 /ml, 0.1 ml/per well) were seeded in 96 well plates in the absence of FCS, and then stimulated with rIL-32 from various sources. For phorbol 12-myristate 13-acetate (PMA) THP-1 differentiation assays, THP-1 cells (0.25×10^5 /ml, 0.1 ml/per well) were seeded in 24 well plates and then stimulated with 100 ng/ml PMA (Sigma, St Louis, MO) for 48 hr, before washing cells in medium lacking FCS, and treating the cells with rIL-32. The plates were placed in a cell culture incubator for 16-20 hours and then the culture supernatants were collected for cytokine measurement.

Human peripheral blood mononuclear cells (PBMC) were isolated from residual leukocytes following platelet-pheresis of healthy donors using Histopaque (Sigma) that was approved by the Combined Colorado Investigational Review Board. PBMC (1.5×10^7) were seeded in 6 well plates in 3 ml of RPMI containing 10% FCS, and then stimulated with 20 μ g of Con A (Sigma). The plates were placed in a cell culture incubator for 60 hours, after which time the culture supernatant and cell lysate were collected for IL-32 measurement.

IL-32 was evaluated in terms of units of biological activity, because there were differences in activity levels between rIL-32 produced in *E. coli*, and rIL-32 produced in mammalian cells, and because there were differences in activity levels between the different batches. One unit of IL-32 is defined as the amount of IL-32 that induces a 2-fold induction of human or mouse TNF α in PMA-differentiated THP-1 cells and mouse Raw cells, respectively, under the assay conditions described above. The approximate concentration of 1 unit of *E. coli* rIL-32 amounted to 20 ng/ml, while the approximate concentration of 1 unit of mammalian rIL-32 amounted to 0.1 ng/ml (as calculated by ECL and Western blot).

EXAMPLE 1

Identification of IL-18 Inducible Genes, including NK4/IL-32

Wild type human A549 lung carcinoma cells (A549-WT) express IL-18R α but not
5 IL-18R β . Thus, in order to express a functional IL-18 receptor in A549 cells, A549 cells
were transfected with an IL-18R β chain expression vector. Briefly, human lung carcinoma
A549 cells (3×10^5 per well) were seeded in 6 well plates a day before transfection. The
cells were then washed with 2 ml of Opti-MEM (Invitrogen, Carlsbad, CA) and incubated
for 25 min in 1 ml of fresh medium. The transfection mixture solution for each well was
10 prepared by mixing 5 μ l of Lipofectamine 2000 in 100 μ l of Opti-MEM followed by a 5
min incubation in the tissue culture hood. Next, 2 μ g of plasmid DNA, pTARGET/huIL-
18R β (Kim *et al.*, *J Immunol*, 166:148-154, 2001), was added, followed by a 20 min
incubation period. The transfection mixture solution was added to the wells and the plate
was incubated for an additional 4 hours at 37°C. Transfection was terminated, by the
15 addition of 2 ml of cell culture medium containing 10% FCS to each well. The next day,
cells were trypsinized and transferred to a 15 cm plate. After cells adhered to the plate, the
culture medium was replaced with fresh medium containing 800 μ g/ml of Neomycin (G418,
Invitrogen). The selection medium was exchanged every three days with fresh medium
until individual colonies appeared (~10-14 days). Small pieces of circular 3 MM papers
20 were sterilized, wetted with trypsin, and then used for picking individual colonies. Colonies
were transferred into 24 wells plate containing 1 ml of selection medium in each well.
Individual clones were grown until 10^6 cells were obtained, and then each clone was tested
for expression of the transfected gene by RT-PCR, as well as in bioassays measuring IL-6
and IL-8 in cell culture medium after IL-18 stimulation for 24 hours. Three clones were
25 positive by both RT-PCR and bioassay. Limiting dilutions cultures were prepared in order
to obtain a single clone.

As shown in Figure 1 panel A, A549 cells transfected with an IL-18R β (A549-R β)
construct (Kim *et al.*, *J Immunol*, 166:148-154, 2001) expressed both the IL-18R α and IL-
18R β chains as determined by RT-PCR. The stable clone (A549-R β) was tested for
30 induction of IL-8 after IL-18 (100 ng/ml) stimulation for 16 hours. Upon the expression of
IL-18R β in A549 cells, the A549-R β cells responded to IL-18 in the absence of a co-
stimulus by producing IL-6 and IL-8, whereas the A549 parent cell line (A549-WT)
remained unresponsive to IL-18 (See, Figure 1, panels B and C, N=7).

WO 2005/047478

PCT/US2004/037578

The A549-R β cells were then used to study IL-18 inducible gene expression by microarray. Briefly, A549-R β cells were seeded at 2×10^6 in nine cm plates one day before the experiment. The cells were then stimulated with IL-18 (50 ng/ml) for 6 hours, at which time the treated cells and controls were harvested. Total RNA was isolated with Tri-Reagent, and purified with the RNeasy kit (Qiagen, Valencia, CA). The total RNA was then used for microarray analysis per instructions from Affymetrix. Total RNA (10 μ g/ml) was converted to first strand cDNA using Superscript II RTTM (Invitrogen). The second strand cDNA was synthesized with the use of T4 DNA polymerase I. After second strand synthesis, the reaction mixture was cleaned with a kit supplied by GeneChipTM. cRNA synthesis was performed to generate biotin-labeled cRNA using a RNA transcript-labeling kit. The biotinylated-cRNA was fragmented prior to hybridization. The samples were hybridized to the Affymetrix GeneChip HG-U133, and data was analyzed in the microarray core laboratory of the University of Colorado Health Sciences Center.

The microarray data revealed that IL-18 induced several cytokine genes including IL-6 and IL-8. Several chemokines including interferon- β 2 and IL-1 β , whose expression was previously known to be IL-18 inducible, were among the group observed to have an increase in expression of greater than 3-fold (log base 2) in response to IL-18 treatment (See, Table 1 showing data from two independent experiments). Interestingly, natural killer cell transcript 4 (NK4) was also highly induced (5.3-fold). The NK4 gene was originally described as an activated NK or T cell product (Dahl *et al.*, *J Immunol*, 148:597-603, 1992), although no functional data had been collected on this gene. Expression of NK4 mRNA was also examined by RT-PCR, using the same total RNA as was used for the microarray study. The first strand cDNA was synthesized from approximately 1 μ g total RNA, using SuperScript II from Invitrogen (Carlsbad, CA). The PCR reaction was performed at 94°C for 45 s, 70°C for 2 min, 59°C for 1 min for 30 cycles with a sense primer, 5'-CTGTCCCGAG TCTGGACTTT-3' (SEQ ID NO:1) and an antisense primer, 5'-GCAAAGGTGG TGGTCAGTAT C-3' (SEQ ID NO:2). The NK4 transcript was detected in IL-18 treated A549-R β cells, but not unstimulated A549-R β cells (See, Figure 1, panel D).

Table 1. IL-18 Inducible Genes Identified By Microarray Analysis

GenBank Accession No.	Fold Increase	Gene Name
M28130	8.6	Interleukin-8 (IL-8)
U64197	8.0	Chemokine exodus-1
X04430	7.7	Interferon- β 2 (IL-6)
U81234	5.8	Chemokine- α 3 (CKA-3)
U37518	5.3	TNF-related apoptosis inducing ligand (TRAIL)
AB007872	5.3	TNF α inducing factor (TAIF, NK4)
X54489	5.0	Melanoma growth stimulatory activity (MGSA)
M36821	4.6	GRO- γ
X03656	4.0	Granulocyte colony stimulating factor (G-CSF)
Z70276	3.7	Fibroblast growth factor 12
M36820	3.2	GRO- β
X04500	3.5	Prointerleukin-1 β
J04513	3.1	Basic fibroblast growth factor (bFGF)

NK4 gene expression was also examined in the NK92 cell line (Hoshino *et al.*, *J Immunol*, 162:51-59, 1999). The NK4 gene was constitutively and highly expressed in this cell line, when maintained in a condition medium containing both IL-2 and IL-15. The IL-2 in the condition medium is contemplated to contribute to the high level of NK4 gene expression observed in the NK92 cells line.

EXAMPLE 2

TNF α Induction by Recombinant NK4/IL-32

To determine the function of this poorly described gene product, the NK4 cDNA (See, Figure 3, IL-32 α) was cloned from NK92 cells (Dahl *et al.*, *supra*, 1992), into pGEMT-Easy (Promega) for sequencing, and then the insert was transferred to pPROEX/Hta (Invitrogen) for expression in *E. coli*, or to pTARGET (Promega) for mammalian expression. Recombinant NK4 was expressed in *E. coli* and purified with a TALON affinity column (Invitrogen) by introducing a His⁶ tag at

WO 2005/047478

PCT/US2004/037578

the N-terminus of the recombinant proteins. The TALON affinity-purified protein was subjected to size exclusion chromatography (Superdex 75, ÄKTATMFPLC), and digested with Tobacco Etch Virus (Invitrogen) for 16 hrs at 4°C to remove the His⁶ tag. The cleaved recombinant proteins were dialyzed in phosphate buffer (20 mM, pH 9). This material was later subjected to ion exchange chromatography (HiTrapTMQFF, ÄKTATMFPLC). The recombinant NK4 protein purified using three sequential steps (His-tag affinity chromatography, size exclusion chromatography and ion exchange chromatography) ran as a homogenous band of approximately 20 kDa in a 10% SDS-PAGE gel, subsequently stained with Coomassie blue (See, Figure 2, panel A).

The thrice-purified recombinant NK4 protein was then tested for biological activity by measuring TNF α secretion by mouse Raw 264.7 macrophage cells in the presence of polymyxin B (100 U/ml). Mouse Raw 264.7 macrophage cells responded to recombinant NK4 and by producing a large amount of TNF α , coinciding with the peak fractions of protein eluting from the ion exchange column (See, Figure 2, panel A). In recognition of the newly discovered biological activity of NK4, the molecule was renamed IL-32/TNF α -inducing factor (TAIF). As shown in Figure 2 panel B, IL-32 α induced significant amounts of TNF α , at IL-32 α concentrations as low as 400 pg/ml (27 picomoles based on the 14.8 kDa calculated molecular weight of IL-32 α), and increased TNF α production in a dose dependent manner. TNF α is well known to possess multiple inflammatory properties that play a causative role in numerous inflammatory and autoimmune diseases (Beutler *et al.*, *Blood Cells Mol Dis*, 24:216-230, 1998).

Recombinant IL-32 α was also produced in mammalian cells, by transient-transfection of Cos-7 cells (8×10^6) with the pTARGET/IL-32 β expression vector via the DEAE-dextran technique (Sompayrac *et al.*, *Proc Natl Acad Sci USA*, 78:7575-7578, 1981). As shown in Figure 2 panel C, recombinant IL-32 α (100 pg/ml) obtained from the concentrated supernatant of IL-32 α -transfected cells induced TNF α production, whereas the concentrated supernatant from mock-transfected cells did not. The concentration of recombinant IL-32 α in Cos-7 supernatants was estimated via ECL and immunoblot. Additionally, IL-32 α produced in mammalian cells was found to possess greater TNF α -inducing activity than did recombinant IL-32 α produced in bacterial cells.

EXAMPLE 3

Identification of IL-32 Genomic Structure and Splice Variants

The structure of the IL-32 gene and its localization within the human genome was analyzed. IL-32 was also cloned by RT-PCR from the human NK92 cell line, cultured in the presence of IL-2 (50 pg/ml), and IL-15 (200 pg/ml). The following primers were used for this purpose: sense 5'-CTGTCCCGAG TCTGGACTTT-3' (SEQ ID NO:1), and antisense 5'-GCAAAGGTGG TGGTCAGTAT C-3' (SEQ ID NO:2). As shown in Figure 3 and 5, three splice variants of IL-32 (IL-32 α disclosed as SEQ ID NO:3 and GENBANK Accession No. AY495331, IL-32 β disclosed as SEQ ID NO:4 and GENBANK Accession No. AY495332, and IL-32 δ disclosed as SEQ ID NO:5 and GENBANK Accession No. AY495333) were identified from RNA derived from NK92 cells, whereas a different splice variant (IL-32 γ disclosed as SEQ ID NO:6 and GENBANK Accession No. BK004065) had been previously reported as the NK4 transcript (GENBANK Accession No: NM_004221). Thus, IL-32 is expressed as at least four variants due to alternative mRNA splicing.

The blast program from the NCBI web site was used to determine that the IL-32 gene resides on chromosome 16p13.3. Approximately 5 kb of sequence encompassing the IL-32 gene (set forth as SEQ ID NO:11 and GENBANK Accession No. AY495334) was identified from a 180 kb stretch of human chromosome 16 sequence (GENBANK Accession No. AC108134). By comparing the sequences of the splice variants with the genomic sequence, the IL-32 gene was found to contain eight small exons, with the second and third exons possessing ATG start codons. IL-32 γ has an additional 46 amino acids at its N-terminus (SEQ ID NO:14) because of the absence splicing between exons 3 and 4. However in IL-32 δ , the second exon is absent, resulting in the use of the ATG start codon in the third exon, instead of the ATG in the second exon. IL-32 α was the most abundant cDNA clone, therefore this isoform was used for many of the experiments described herein. IL-32 α has a deletion of 57 amino acid residues at its C-terminus due to splicing between exons 7 and 8, which is in contrast to the other variants that have a single large exon encoding their C-termini. The analysis of the IL-32 amino acid sequences revealed the presence of three potential N-myristoylation sites and one potential N-glycosylation site (See, Figure 4A).

The Blast program was used to search the NCBI database for homologs of IL-32. In this way, expressed sequence tag (EST) clones of equine, bovine, ovine, and swine IL-32 were identified. The equine IL-32 β protein sequence (GENBANK Accession Nos. CD469554 and BI961524) shares the highest homology with the human sequence (31.8%

WO 2005/047478

PCT/US2004/037578

identity), followed by bovine (GENBANK Accession Nos. CK834399 and CK832489),
 ovine (GENBANK Accession No. CO202364), and swine (GENBANK Accession No.
 CB287292) sequences. The alignments of human, equine, and bovine IL-32 β protein
 sequences are shown in Figure 4B). In particular, isoform A of sheep and pigs has been
 5 identified, while both isoforms A and B have been identified in horses, and isoforms B and
 C have been identified in cows. The sequences of equine IL-32 β and bovine IL-32 γ were
 each combined from two EST clones, and the open reading frames were deduced from the
 combined sequence.

10

EXAMPLE 4**Analysis of IL-32 Expression**

The expression of IL-32 in various human tissues was examined by northern blot.
 Briefly, IL-32 α and actin cDNA inserts were excised from plasmid vectors by using
 suitable restriction enzymes, size-fractionated and purified with a GeneClean II kit (Q-BIO
 15 gene, Carlsbad, CA). The cDNAs were labeled with ³²P-dCTP (NEN Life Science, Boston,
 MA) by random priming using the Klenow enzyme (New England Biolabs, Beverly, MA).
 A membrane containing human poly(A)⁺ RNA from different tissues (human MTN Blot II)
 incubated with the cDNA probes, whose binding was detected by autoradiography. As
 shown in Figure 8A, a 1.2kb IL-32 transcript was detected in most tissues, with higher
 20 levels of IL-32 detected in immune cells as compared to other cell types. The high
 expression of IL-32 mRNA in immune tissues was not due to loading differences as shown
 by reprobing the blot with an actin fragment.

Endogenous IL-32 expression was also detected at the protein level. Briefly, A549-
 R β cells were seeded at 10⁶ cells/well in 6 well plates. After the cells had adhered to the
 25 plate, the F12K culture medium was removed and replaced with serum-free medium, which
 in some wells contained IL-18 (50 ng/ml), IL-1 β (100 ng/ml) or LPS (500 ng/ml). After 48
 hrs, the supernatants were harvested and concentrated 10 times with CentriconTM
 concentrators. As shown in Figure 8B, an approximately 30 kDa species corresponding to
 endogenous IL-32 was detected by immunoblot with an affinity-purified rabbit anti-human
 30 IL-32 α polyclonal antibody. The difference in molecular weight between rIL-32 from *E.*
coli, and endogenous IL-32 is contemplated to be due to post-translational modification of
 the endogenous molecule, since analysis of the amino acid sequence revealed the presence
 of potential N-linked glycosylation and myristoylation sites. IL-32 was secreted into the
 cell culture medium of cells treated with IL-18 (100 ng/ml), IL-1 β (100 ng/ml) or LPS (500

ng/ml), but not by unstimulated cells (control). IL-32 was detected in cell culture medium as a secreted molecule, although IL-32 does not possess a typical hydrophobic signal peptide at its N-terminus. IL-18 induced IL-32 expression to a greater extent than did IL-1 β or LPS. A 60 kDa band was also observed (not shown), which is contemplated to be a dimerized form of IL-32. Like the monomer, the 60 kDa band was found in the supernatants of treated cells, but was absent from that of unstimulated cells.

Induction of IL-32 in human epithelial Wish cells treated with IFN γ since was also examined, since Wish cells are commonly used for antiviral assays and for assessments of other biological activities of IFN γ . For time courses of IL-32 production, Wish, A549-WT, and A549-R β cells were seeded at 5×10^4 cells/well in 6 well plates, incubated overnight and then stimulated with IFN γ (100 U/ml), IL-18 (50 ng/ml) or IL-1 β (10 ng/ml). Affinity purified rabbit anti-IL-32 α was used for detecting IL-32 in cell culture medium by immunoblot. Peroxidase-conjugated secondary antibodies (Jackson ImmunoResearch Laboratories, West Grove, PA) were used to develop the blots using enhanced chemiluminescence (NEN Life Science). For IL-32 detection by electrochemiluminescence (ECL), an aliquot of the affinity purified anti-IL-32 α antibody was labeled with biotin, and another aliquot was labeled with ruthenium according to the manufacturer's instructions (Igen, Gaithersburg, MD). The biotin and ruthenium labeled antibodies were used to construct a standard curve using recombinant IL-32 α . The liquid-phase ECL method was used to measure various cytokines in cell culture media and in serum samples. The amount of electrochemiluminescence was determined using an Origen Analyzer (Igen).

As shown in Figure 9A, IL-32 was induced in cell lysates of human epithelial cells in a time dependent manner, diminishing after 46 hours, whereas IL-32 was not induced in unstimulated cells. Similarly, IL-18 and IL-1 β increased the expression of endogenous IL-32 in the lysates of A549-R β cells in a time dependent manner (Figure 9B) whereas IL-32 was only induced by IL-1 β in A549-WT cells (Figure 9C). In addition, IFN γ was also observed to induce IL-32 in the lysates of A549-WT cells (data not shown).

In order to determine whether the recombinant proteins expressed from the cDNAs of IL-32 α and β in mammalian cells were comparable to endogenous IL-32, Cos7 cells were transiently transfected with IL-32 α and β cDNAs. As shown in Figures 9D and 9E, recombinant IL-32 α and IL-32 β were present in both cell culture media and lysates of the transfected cells as determined by immunoblot. The molecular size of recombinant mammalian IL-32 in the immunoblots was identical to the molecular size of the endogenous

WO 2005/047478

PCT/US2004/037578

IL-32 shown in Figure 8B. The ECL assay revealed a similar distribution of IL-32 α and IL-32 β in the culture media and lysates, although IL-32 β appears to be more efficiently secreted as compared to IL-32 α , which was predominantly cell-associated.

The polyclonal anti-human IL-32 antibody was affinity purified over a column containing IL-32 α -immobilized on agarose beads (Affi-gel 15)TM. Peroxidase conjugated anti-rabbit immunoglobulin secondary antibodies were obtained from Jackson ImmunoResearch. Since the anti-human IL-32 antibody specifically recognizes endogenous IL-32, the IL-32 level in the sera of human subjects (healthy individuals and patients with sepsis) was measured by ECL. Although IL-32 was detected in sera of healthy individuals (5 of 42 samples), the IL-32 levels were less than 70 pg/ml. In contrast, the mean IL-32 levels in sera of patients with sepsis were 35-fold higher than that of sera from healthy individuals. For the first time during development of the present invention, IL-32 is shown to be an inflammatory cytokine produced (directly or indirectly) in response to bacterial infection. In addition, IL-32 was detected in patients with active rheumatoid arthritis indicating that it also plays a role in autoimmunity.

To examine the regulation of IL-32 production, IL-32 was measured in the cell culture medium of different cell lines, as well as human PBMC. IL-32 was detected in the cell culture medium of A549-R β cells after IL-18 or IL-1 β stimulation but not in control medium (Figure 10A). Since the NK4 (IL-32) gene was isolated from IL-2-activated NK cells, the human NK cell line was stimulated with a combination of IL-12 plus IL-18. As shown in Figure 10B, there was a significant induction of IL-32 by IL-12 or IL-18 in NK cells, and the effect of the combination of these two cytokines appeared to be additive. By comparison, when this cell line was stimulated by either IL-12 or IL-18, there is little or no induction of IFN γ , whereas IFN γ production by the combination of IL-12 plus IL-18 is highly synergistic (Kim *et al.*, *J Biol Chem*, 277:10998-11003, 2002). The induction of IL-32 (via IFN γ) was examined by stimulating NK cells with IL-12 and IL-18, in the presence of a neutralizing anti-IFN γ antibody. This combination had no effect on IL-32 induction (data not shown). Human peripheral blood mononuclear cells (PBMC) contain mostly T-cells with few numbers of monocytes and B-cells. Freshly prepared PBMC were stimulated with LPS or ConA and the supernatants and lysates were harvested and assayed for IL-32 concentration. After 60 hours, there was no detectable IL-32 in the supernatants or lysates of PBMC stimulated with LPS (data not shown). However, ConA consistently induced IL-32, which was found in the supernatants and lysates. As shown in Figure 10C, the lysates contained more IL-32 than the supernatants.

WO 2005/047478

PCT/US2004/037578

EXAMPLE 5

Identification of IL-32 Responsive Signal Transduction Pathways

This example provides details of the experiments conducted to assess the effect of the cytokine IL-32 on signal transduction molecules inhibitor of kappa B (I κ B) and p38 mitogen-activated protein kinase (MAPK). Briefly, mouse RAW 264.7 macrophage cells were stimulated with IL-32 α (50 ng/ml) in the presence of 5 μ g/ml polymyxin B (Bedford Lab, Bedford, OH) for the indicated amount of time (in minutes). Cells were lysed with kinase lysis buffer (Han *et al.*, *J Biol Chem*, 277:47167-47174, 2002). The cell contents were separated in a 10% SDS-PAGE gel and transferred to a nitrocellulose membrane, which was subsequently blocked with 3% BSA. The membrane was probed with rabbit anti-I κ B and normalized with goat anti-actin (Santa Cruz Biotechnology, Santa Cruz, CA). As shown in Figure 11A, IL-32 α induced I κ B degradation in a time dependent manner, beginning 15 min after treatment and reaching a maximal level at 45 min, followed by recovery after 90 min.

The membrane was also probed with rabbit anti-phospho-p38 MAPK and normalized with rabbit anti-p38 MAPK (Cell Signaling, Beverly, MA). As shown in Figure 10B, phosph-p38 MAPK was dramatically increased 5 min after stimulation by IL-32 α and then decreased from 15 min to 30 minutes, thereafter. Interestingly, a second increase in p38 MAPK phosphorylation was observed at 45 min, which decreased more slowly.

EXAMPLE 6

Identification of IL-32 Binding Proteins

Recombinant IL-32 α (or IL-32 β , IL-32 δ and IL-32 γ) is expressed in *E. coli* and purified using three sequential steps as described in Example 2. Approximately 5 mg of the purified IL-32 α is immobilized onto Affi-gelTM 15 agarose beads (Bio-Rad Laboratories, Hercules, CA). Various potential receptor sources are applied to the IL-32 α -affinity column (*e.g.*, human serum or lysates of cells which secrete TNF α in response to IL-32 α , such as Raw 264.7 cells). The IL-32 α affinity column is washed extensively and the IL-32 receptor is then eluted with an elution buffer (*e.g.*, 50 mM citric acid, 100 mM NaCl, pH 2.5). The eluted fractions are neutralized immediately with 2 M tris base.

The IL-32 binding proteins isolated in this way are further characterized by chemical analysis (*e.g.*, peptide sequence analysis through Edman degradation and/or mass spectroscopy) and bioassay (*e.g.*, purified or recombinant receptors are tested for their ability to block IL-32 α -induced TNF α secretion by Raw 264.7 cells). It is contemplated

WO 2005/047478

PCT/US2004/037578

that authentic IL-32 binding proteins have the ability to inhibit IL-32 biological activities (similar to what has been observed with the TNFBP). It is also contemplated that other types of authentic IL-32 binding proteins have the ability to enhance IL-32 biological activities (similar to what has been observed with the IL-6 ligand binding chain).

5

EXAMPLE 7

Therapeutic Effect of IL-32-Antibodies and IL-32 Inhibitors in a Murine Model of Arthritis

This example provides details of the experiments conducted to assess the effect of
10 IL-32-antibodies and IL-32 antagonists (*e.g.*, soluble IL-32 receptors, dominant-negative IL-32 variants, small molecule inhibitors, etc.) as therapeutics for the treatment of collagen-induced arthritis (CIA) in mice. Briefly, CIA is induced in 8 to 10 week old DBA/1J mice (Jackson Laboratories, Bar Harbor, ME) by intradermal injection of type II bovine collagen (CII) as known in the art (Banda *et al.*, *Arthritis Rheum*, 46:3065, 2002). Each mouse
15 receives 100 μ l injections containing 200 μ g CII and 200 μ g of inactivated *Mycobacterium tuberculosis* (Difco, Detroit, MI) in incomplete freund's adjuvant on days 0 and 21. Mice are treated between days 21 and 42 with one of three therapies given as IP injections every 3 days: 2 mg/mouse normal rabbit IgG; 2 mg/mouse neutralizing rabbit anti-IL-32 prepared using HiTrap Protein G HP (Amersham Pharmacia Biotech AB, Uppsala, Sweden); and 2
20 mg/mouse recombinant IL-32 antagonist (*e.g.*, IL-32 receptor-IgG1 Fc fusion protein or dominant-negative IL-32 variant). The mice (5 in each group and 5 untreated mice) are sacrificed on day 42 by anesthesia and cervical dislocation.

The clinical disease activity of CIA is assessed every other day between days 21 and 42 by blinded observers using a three-point scale for each paw; 0 = normal joint; 1 = slight
25 inflammation and redness; 2 = severe erythema and swelling affecting the entire paw with inhibition of use; and 3 = deformed paw or joint, with ankylosis, joint rigidity, and loss of function. The total score for clinical disease activity is based on all four paws, with a maximum score of 12 for each animal. After sacrifice, both forepaws and the right hand limb are surgically removed on day 42 and fixed in 10% buffered formalin with preparation
30 of tissues samples and histological analysis as known in the art (Bendele *et al.*, *Arthritis Rheum*, 43:2648, 2000). An experienced observer (blinded to the treatment) scores the histological findings in paws, ankles, and knees. The data are expressed as mean scores for inflammation, pannus, cartilage damage, and bone damage, as well as an overall score, based on scales of 0 to 5.

Using published methods (Banda *et al.*, *J Immunol*, 170:2100-2105, 2003), various immune parameters are measured, including but not limited to: CII-specific proliferation by spleen and lymph node cells, production of anticollagen antibodies, CII-induced cytokine secretion by spleen cells (*e.g.*, utilizing TNF α , IFN γ , IL-1 β , IL-1Ra and IL-10 ELISAs), and steady state cytokine mRNA levels in joints (*e.g.*, TNF α , IFN γ , IL-1 β , IL-1Ra, IL-6, IL-18, MIF, TNF β , LT β , TGF β 1, TGF β 2). Some preferred embodiments of the present invention comprise a IL-32 antibody or a IL-32 antagonist that reduces clinical disease activity scores and histological scores of joint damage, by at least 50% (more preferably by at least 75%, and most preferably by at least 90%). Other preferred embodiments comprise an IL-32 antibody or an IL-32 antagonist that reduces CII-induced lymphocyte proliferation, and/or serum levels of collagen binding IgG antibodies, by at least 50% (more preferably by at least 75%, and most preferably by at least 90%). Particularly preferred embodiments, comprise an IL-32 antibody or an IL-32 antagonist that reduces steady state mRNA levels of TNF α , IFN γ , and/or IL-1 β in isolated joints, by at least 50% (more preferably by at least 75%, and most preferably by at least 90%). Importantly, preferred embodiments comprising the IL-32 antibodies or IL-32 antagonists described herein, are contemplated to find use in the treatment of human rheumatoid arthritis patients, using methods similar to that employed for administration of TNF-reactive antibodies (infliximab/REMICADE and adalimumab/HUMIRA) and soluble TNF-receptor/immunoglobulin fusion proteins (etanercept/ENBREL).

EXAMPLE 8

Biological Activity of Recombinant IL-32

IL-32 α induced significant amounts of TNF α and MIP-2 and increased the production of both cytokines in a dose dependent manner as shown in Figure 6A. The biological activity of IL-32 α as then compared with that of IL-32 β , which has a full C-terminus. IL-32 β induced similar the levels of TNF α and MIP-2 as did IL-32 α in the mouse macrophage Raw cell line (Figure 6B). Although all the experiments were performed in the presence of polymyxin B (5 μ g/ml), the possibility of endotoxin contamination of the recombinant proteins produced in *E. coli* could not be ruled out.

Recombinant IL-32 was also produced in a mammalian system in order to avoid endotoxin contamination. The rIL-32 β was produced from three different sources of mammalian cells, Anjou65 stable clone, Cos7 stable clone (Cos7-S), and Cos7 transient transfectant (Cos7-T). The transient and stable cloned cells were cultured in 0.5% FCS for

WO 2005/047478

PCT/US2004/037578

4 days, before harvest. As the maximum yield of mammalian rIL-32 β was only 1 ng/ml and IL-32 α was below 100 pg/ml concentration, each endotoxin free mammalian rIL-32 β was purified using an affinity column prepared by immobilizing an anti-IL-32 β mAb to agarose beads (Affi-Gel Hz, Bio-Rad) in the presence of sodium azide (0.2%) to prevent microorganism contamination during purification. The purified rIL-32 was dialyzed against RPMI containing penicillin/streptomycin (10 μ g/ml) overnight at 4°C prior to use for bioassay. All three mammalian rIL-32 β preparations induced TNF α in human PMA-differentiated THP-1 cells and in mouse Raw cells, respectively (Figure 6C). *E. coli* rIL-32 α and mammalian rIL-32 β increased human TNF α product in PMA differentiated-THP-1 cells in a dose dependent manner (Figure 6D). Furthermore, the high unit *E. coli* rIL-32 α and mammalian Anjou 65 rIL-32 β preparations induced human IL-8 in undifferentiated-THP-1 cells, whereas the mock Anjou65 transfectants did not (Figure 6E).

EXAMPLE 9

15 Production of a Neutralizing Fab Fragment of an Anti-IL-32 Monoclonal Antibody

A five week old female *Balb/c* mouse was immunized with 20 μ g of rIL-32 β antigen emulsified in Freund's complete adjuvant (Sigma). On days 14 and 21, the mouse was given an intravenous and intraperitoneal injection with the antigen emulsified in Freund's incomplete adjuvant (Sigma). After three injections, the mouse was sacrificed, the spleen was aseptically harvested, and splenocytes were prepared for fusion. Briefly, 1 x 10⁷ splenocytes and 1 x 10⁶ NS-1 mouse myeloma cells (ATCC) were fused using polyethylene glycol 1500 (Roche Applied Science, Indianapolis, IN). Fused cells were resuspended at 1 x 10⁶ cells/ml in hybridoma growth media with 10% FCS and HAT, and plated in 96 well plates. After 2 weeks, the culture supernatants of hybridomas were titrated in an indirect ELISA. Monoclonal antibody classes and subclasses were determined using an IMMUNO-TYPE mouse monoclonal antibody isotyping kit (BD Bioscience, San Diego, CA) according to the manufacturer's instructions.

Approximately 5 x 10⁶ cells of two hybridomas 32-4 (IgG₁) and 32-9 (IgG₁) were intraperitoneally injected into an 8 week old female *Balb/c* mouse. After one week, ascites fluid was collected using a sterile hypodermic needle. Antibodies from the ascites' supernatant was purified by using a protein A Sepharose column (Bio-Rad), and eluted with 0.1 M glycine-HCl, pH 2.7. The eluted mAbs were dialyzed in PBS, and purified

WO 2005/047478

PCT/US2004/037578

antibodies were concentrated using Centricon concentrators (YM-50, Life Sciences, Ann Arbor, MI). The concentration of purified antibody was determined by measuring absorbance at 280 nm.

As the affinity-purified monoclonal antibody (32-4) against rIL-32 β induced a high background level of mTNF α , a Fab fragment of this monoclonal (mAb) was prepared. Briefly, Fab fragments of the purified anti-IL-32 mAb (32-4) were generated by incubating with immobilized-Pepsin (PIERCE, Rockford, IL) at 37°C for 4h. Fc fragments and residual uncleaved mAbs were removed by using protein G Sepharose (Amersham Biosciences, Uppsala, Sweden). The Fab fragment was dialyzed against PBS overnight at 4°C and subjected to SDS-PAGE in order to confirm that cleavage was complete. The background level reduced dramatically after removing the Fc fragment, permitting the assessment the IL-32 neutralization activity of the anti-rIL-32 β mAb. As shown in Figure 7A, the Fab of the 32-4 mAb (50ng/ml) inhibited the biological activity of *E. coli* rIL-32 α by more than 70%. As shown in Figure 7B, the same Fab inhibits the biological activity of the affinity purified Cos7-S rIL-32 β by greater than 65%. Higher concentrations of the Fab fragment didn't further inhibit IL-32-induced TNF α production in these assays.

EXAMPLE 10

Therapeutic Effect of IL-32-Antibodies and IL-32 Inhibitors

in a Murine Model of Inflammatory Bowel Disease

This example provides details of the experiments conducted to assess the effect of IL-32-antibodies and IL-32 antagonists (e.g., soluble IL-32 receptors, dominant-negative IL-32 variants, small molecule inhibitors, etc.) as therapeutics for the treatment of dextran sulphate sodium (DSS)-induced colitis in mice.

Briefly, colitis is induced in 8-10 week old female C57BL/6 mice (Taconic Laboratories) by administering 2% (wt/vol) dextran sodium sulphate (DSS, MW 40,000 from ICN Biochemicals) from day 0 to day 7 in the drinking water ad libitum, followed by return to normal water as known in the art (Sivakumar *et al.*, *Gut*, 50:812-820, 2002). Mice are weighed every day beginning on day 0 and weight changes are recorded until day 13. Percent weight change for each mouse is calculated as follows: percent weight change = (weight on specific day – weight on day 0)/weight on day 0 x 100. Groups of mice are treated with: a control protein (50 μ g or 500 μ g); murine anti-IL-32 mAb (50 μ g or 500 μ g); and recombinant IL-32 antagonist (e.g., IL-32 receptor-IgG1 Fc fusion protein or dominant-negative IL-32 variant, at 50 μ g or 500 μ g), each in a volume of 200 μ l per injection

(endotoxin-free PBS). Mice are injected IP from day 0 to day 7. The mice are sacrificed on days 0, 2, 4, 6, or 8 following the start of DSS treatment, by anesthesia and cervical dislocation, and tissues (*e.g.*, large intestine, lymph nodes) are removed and processed for RNA extraction, histopathology and cytokine analysis, using methods known in the art. It is contemplated that administration of IL-32 antagonists is suitable for attenuation of inflammation during DSS induced colitis in mice, and that neutralizing IL-32 activity is of benefit for ameliorating the inflammation associated with intestinal diseases.

EXAMPLE 11

Therapeutic Effect of IL-32-Antibodies and IL-32 Inhibitors in a Murine Model of Hepatitis

This example provides details of the experiments conducted to assess the effect of IL-32-antibodies and IL-32 antagonists (*e.g.*, soluble IL-32 receptors, dominant-negative IL-32 variants, small molecule inhibitors, etc.) as therapeutics for the treatment of LPS-induced liver injury in mice primed with heat-killed *Propioibacterium acnes* (model of Fas/FasL-mediated liver disease).

Briefly, hepatitis is induced in 8-10 week old female BALB/c mice (Charles River Laboratories) by administering 500 µg/mouse IV of heat-killed *P. acnes* (Ribi Immuno-Chem Research) and 12 days later they are challenged with 50 µg/kg LPS IV as known in the art (Faggioni *et al.*, *J Immunol*, 167:5913-5920, 2001). Groups of mice are treated with: a control protein (50 µg or 500 µg); murine anti-IL-32 mAb (50 µg or 500 µg); and recombinant IL-32 antagonist (*e.g.*, IL-32 receptor-IgG1 Fc fusion protein or dominant-negative IL-32 variant, at 50 µg or 500 µg), each in a volume of 200 µl per injection (endotoxin-free PBS). Mice are injected IP either at the time of *P. acnes* administration or 10 min before LPS challenge. Mice are monitored for survival or sacrificed to collect livers for histological examination, mRNA and chemokine measurements, and blood for serum IFN-γ and transaminase measurements using methods known in the art. It is contemplated that administration of IL-32 antagonists is suitable for preventing LPS-induced liver damage and IFN-γ and Fas ligand expression when administered 10 minutes before LPS challenge of *P. acnes*-primed mice. In addition, the administration of IL-32 antagonists is contemplated to be suitable for decreasing *P. acnes*-induced granuloma formation, macrophage-inflammatory protein-1α and macrophage-inflammatory protein-2 production, when given at the moment of priming with *P. acnes*. Thus it is contemplated that patients

WO 2005/047478

PCT/US2004/037578

with liver diseases such as HCV-induced hepatitis, autoimmune hepatitis, and primary biliary cirrhosis will benefit from a therapy regimen comprising an IL-32 antagonist.

EXAMPLE 12

Therapeutic Effect of IL-32-Antibodies and IL-32 Inhibitors in a Murine Model of Ischemic Disease

This example provides details of the experiments conducted to assess the effect of IL-32-antibodies and IL-32 antagonists (*e.g.*, soluble IL-32 receptors, dominant-negative IL-32 variants, small molecule inhibitors, etc.) as therapeutics for stimulating tissue neovascularization in response to ischemic injury.

Briefly, male C57BL/6J mice undergo surgery to induce unilateral hindlimb ischemia using methods known in the art (Mallat et al., *Circ Res*, 91:441-448, 2002). Animals are anesthetized by isoflurane inhalation. The ligature is performed on the right femoral artery, 0.5 cm proximal to the bifurcation of the saphenous and popliteal arteries, and then housed under conventional conditions for 3 or 28 days. Groups of mice are treated with: a control protein (50 μ g or 500 μ g); murine anti-IL-32 mAb (50 μ g or 500 μ g); and recombinant IL-32 antagonist (*e.g.*, IL-32 receptor-IgG1 Fc fusion protein or dominant-negative IL-32 variant, at 50 μ g or 500 μ g), each in a volume of 200 μ l per injection (endotoxin-free PBS). Mice are injected IP at day 0, and day 7. The degree of angiogenesis is quantified on days 3 and 10, by measuring vessel density by microangiography, and ischemia-induced changes in vascularization is monitored by laser Doppler perfusion imaging. It is contemplated that administration of IL-32 antagonists is suitable for stimulating tissue neovascularization in response to ischemic injury.

Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention, which are obvious to those skilled in molecular biology, genetics, immunology or related fields are intended to be within the scope of the claims.

SEQUENCE LISTING

<110> The Regents of the University of Colorado, a Body Corporate

<120> Compositions and Methods for Regulation of Tumor Necrosis
Factor-Alpha

<130> 08905753CA

<140> not yet known

<141> 2004-11-12

<150> 60/519,818

<151> 2003-11-12

<160> 27

<170> PatentIn version 3.2

<210> 1

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 1

ctgtccccgag tctggacttt

20

<210> 2

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 2

gcaaagggtgg tggtcagtat c

21

<210> 3

<211> 396

<212> DNA

<213> Homo sapiens

<400> 3

atgtgcttcc cgaaggctct ctctgatgac atgaagaagc tgaaggcccg aatgcaccag

60

gctatagaaa gattttatga taaaatgcaa aatgcagaat caggacgtgg acaggtgatg

120

tcgagcctgg cagagctgga ggacgacttc aaagagggct acctggagac agtggcggct

180

tattatgagg agcagcacc agagctcact cctctacttg aaaaagaaag agatggatta

240

cggtgccgag gcaacagatc ccctgtcccg gatgttgagg atcccgcaac cgaggagcct

300

ggggagagct tttgtgacaa gtcctacgga gcccacggg gggacaagga ggagctgaca

360

ccccagaagt gctctgaacc ccaatcctca aaatga

396

<210> 4
 <211> 567
 <212> DNA
 <213> Homo sapiens

<400> 4
 atgtgcttcc cgaaggtcct ctctgatgac atgaagaagc tgaaggcccg aatgcaccag 60
 gccatagaaa gattttatga taaaatgcaa aatgcagaat caggacgtgg acaggtgatg 120
 tcgagcctgg cagagctgga ggacgacttc aaagagggct acctggagac agtggcggct 180
 tattatgagg agcagcacc cagagctcact cctctacttg aaaaagaaag agatggatta 240
 cgggtgccgag gcaacagatc ccctgtcccg gatgttgagg atcccgcaac cgaggagcct 300
 ggggagagct tttgtgacaa ggtcatgaga tggttccagg ccatgctgca gcggctgcag 360
 acctggtggc acgggggttct ggcttgggtg aaggagaagg tgggtggcct ggtccatgca 420
 gtgcaggccc tctggaaaca gttccagagt ttctgctgct ctctgtcaga gctcttcatg 480
 tcctctttcc agtcctacgg agccccacgg ggggacaagg aggagctgac accccagaag 540
 tgctctgaac cccaatcctc aaaatga 567

<210> 5
 <211> 705
 <212> DNA
 <213> Homo sapiens

<400> 5
 atgtgcttcc cgaaggtcct ctctgatgac atgaagaagc tgaaggcccg aatggtaatg 60
 ctctcccta cttctgctca ggggttgggg gcctgggtct cagcgtgtga cactgaggac 120
 actgtgggac acctgggacc ctggagggac aaggatccgg ccctttgggtg ccaactctgc 180
 ctctcttcac agcaccaggc catagaaaga ttttatgata aaatgcaaaa tgcagaatca 240
 ggacgtggac aggtgatgtc gagcctggca gagctggagg acgacttcaa agagggctac 300
 ctggagacag tggcggctta ttatgaggag cagcaccacg agctcactcc tctacttgaa 360
 aaagaaagag atggattacg gtgccgaggc aacagatccc ctgtcccga tgttgaggat 420
 cccgcaaccg aggagcctgg ggagagcttt tgtgacaagg tcatgagatg gttccaggcc 480
 atgctgcagc ggctgcagac ctggtggcac ggggttctgg cctgggtgaa ggagaagggtg 540
 gtggccctgg tccatgcagt gcaggccctc tggaaacagt tccagagttt ctgctgctct 600
 ctgtcagagc tcttcatgtc ctctttccag tcctacggag cccacgggg ggacaaggag 660
 gagctgacac ccagaagtg ctctgaacc caatcctcaa aatga 705

<210> 6
 <211> 537
 <212> DNA
 <213> Homo sapiens

<400> 6
 atgaagaagc tgaaggcccg aatgcaccag gccatagaaa gattttatga taaaatgcaa 60
 aatgcagaat caggacgtgg acaggtgatg tcgagcctgg cagagctgga ggacgacttc 120
 aaagagggct acctggagac agtggcggct tattatgagg agcagcaccc agagctcact 180
 cctctacttg aaaaagaaag agatggatta cggtgccgag gcaacagatc ccctgtcccg 240
 gatgttgagg atcccgcaac cgaggagcct ggggagagct tttgtgacaa ggtcatgaga 300
 tggttccagg ccatgctgca gcggctgcag acctgggtggc acgggggttct ggcttgggtg 360
 aaggagaagg tgggtggccct ggtccatgca gtgcaggccc tctggaaaca gttccagagt 420
 ttctgctgct ctctgtcaga gctcttcatg tcctctttcc agtcctacgg agccccacgg 480
 ggggacaagg aggagctgac accccagaag tgctctgaac cccaatcctc aaaatga 537

<210> 7
 <211> 131
 <212> PRT
 <213> Homo sapiens

<400> 7

Met	Cys	Phe	Pro	Lys	Val	Leu	Ser	Asp	Asp	Met	Lys	Lys	Leu	Lys	Ala
1				5				10						15	
Arg	Met	His	Gln	Ala	Ile	Glu	Arg	Phe	Tyr	Asp	Lys	Met	Gln	Asn	Ala
			20					25						30	
Glu	Ser	Gly	Arg	Gly	Gln	Val	Met	Ser	Ser	Leu	Ala	Glu	Leu	Glu	Asp
		35					40					45			
Asp	Phe	Lys	Glu	Gly	Tyr	Leu	Glu	Thr	Val	Ala	Ala	Tyr	Tyr	Glu	Glu
	50					55					60				
Gln	His	Pro	Glu	Leu	Thr	Pro	Leu	Leu	Glu	Lys	Glu	Arg	Asp	Gly	Leu
65					70					75					80
Arg	Cys	Arg	Gly	Asn	Arg	Ser	Pro	Val	Pro	Asp	Val	Glu	Asp	Pro	Ala
				85					90					95	
Thr	Glu	Glu	Pro	Gly	Glu	Ser	Phe	Cys	Asp	Lys	Ser	Tyr	Gly	Ala	Pro
			100					105					110		
Arg	Gly	Asp	Lys	Glu	Glu	Leu	Thr	Pro	Gln	Lys	Cys	Ser	Glu	Pro	Gln
		115					120					125			
Ser	Ser	Lys													
		130													

<210> 8
 <211> 188
 <212> PRT
 <213> Homo sapiens

<400> 8

Met Cys Phe Pro Lys Val Leu Ser Asp Asp Met Lys Lys Leu Lys Ala
 1 5 10 15

Arg Met His Gln Ala Ile Glu Arg Phe Tyr Asp Lys Met Gln Asn Ala
 20 25 30

Glu Ser Gly Arg Gly Gln Val Met Ser Ser Leu Ala Glu Leu Glu Asp
 35 40 45

Asp Phe Lys Glu Gly Tyr Leu Glu Thr Val Ala Ala Tyr Tyr Glu Glu
 50 55 60

Gln His Pro Glu Leu Thr Pro Leu Leu Glu Lys Glu Arg Asp Gly Leu
 65 70 75 80

Arg Cys Arg Gly Asn Arg Ser Pro Val Pro Asp Val Glu Asp Pro Ala
 85 90 95

Thr Glu Glu Pro Gly Glu Ser Phe Cys Asp Lys Val Met Arg Trp Phe
 100 105 110

Gln Ala Met Leu Gln Arg Leu Gln Thr Trp Trp His Gly Val Leu Ala
 115 120 125

Trp Val Lys Glu Lys Val Val Ala Leu Val His Ala Val Gln Ala Leu
 130 135 140

Trp Lys Gln Phe Gln Ser Phe Cys Cys Ser Leu Ser Glu Leu Phe Met
 145 150 155 160

Ser Ser Phe Gln Ser Tyr Gly Ala Pro Arg Gly Asp Lys Glu Glu Leu
 165 170 175

Thr Pro Gln Lys Cys Ser Glu Pro Gln Ser Ser Lys
 180 185

<210> 9
 <211> 234
 <212> PRT
 <213> Homo sapiens

<400> 9

Met Cys Phe Pro Lys Val Leu Ser Asp Asp Met Lys Lys Leu Lys Ala
 1 5 10 15

Arg Met Val Met Leu Leu Pro Thr Ser Ala Gln Gly Leu Gly Ala Trp
 20 25 30

Val Ser Ala Cys Asp Thr Glu Asp Thr Val Gly His Leu Gly Pro Trp
 35 40 45

Arg Asp Lys Asp Pro Ala Leu Trp Cys Gln Leu Cys Leu Ser Ser Gln
 50 55 60

His Gln Ala Ile Glu Arg Phe Tyr Asp Lys Met Gln Asn Ala Glu Ser
 65 70 75 80

Gly Arg Gly Gln Val Met Ser Ser Leu Ala Glu Leu Glu Asp Asp Phe
 85 90 95

Lys Glu Gly Tyr Leu Glu Thr Val Ala Ala Tyr Tyr Glu Glu Gln His
 100 105 110

Pro Glu Leu Thr Pro Leu Leu Glu Lys Glu Arg Asp Gly Leu Arg Cys
 115 120 125

Arg Gly Asn Arg Ser Pro Val Pro Asp Val Glu Asp Pro Ala Thr Glu
 130 135 140

Glu Pro Gly Glu Ser Phe Cys Asp Lys Val Met Arg Trp Phe Gln Ala
 145 150 155 160

Met Leu Gln Arg Leu Gln Thr Trp Trp His Gly Val Leu Ala Trp Val
 165 170 175

Lys Glu Lys Val Val Ala Leu Val His Ala Val Gln Ala Leu Trp Lys
 180 185 190

Gln Phe Gln Ser Phe Cys Cys Ser Leu Ser Glu Leu Phe Met Ser Ser
 195 200 205

Phe Gln Ser Tyr Gly Ala Pro Arg Gly Asp Lys Glu Glu Leu Thr Pro
 210 215 220

Gln Lys Cys Ser Glu Pro Gln Ser Ser Lys
 225 230

<210> 10

<211> 178

<212> PRT

<213> Homo sapiens

<400> 10

Met	Lys	Lys	Leu	Lys	Ala	Arg	Met	His	Gln	Ala	Ile	Glu	Arg	Phe	Tyr
1				5					10					15	

Asp	Lys	Met	Gln	Asn	Ala	Glu	Ser	Gly	Arg	Gly	Gln	Val	Met	Ser	Ser
			20					25					30		

Leu	Ala	Glu	Leu	Glu	Asp	Asp	Phe	Lys	Glu	Gly	Tyr	Leu	Glu	Thr	Val
		35					40					45			

Ala	Ala	Tyr	Tyr	Glu	Glu	Gln	His	Pro	Glu	Leu	Thr	Pro	Leu	Leu	Glu
	50					55					60				

Lys	Glu	Arg	Asp	Gly	Leu	Arg	Cys	Arg	Gly	Asn	Arg	Ser	Pro	Val	Pro
65					70					75					80

Asp	Val	Glu	Asp	Pro	Ala	Thr	Glu	Glu	Pro	Gly	Glu	Ser	Phe	Cys	Asp
				85					90					95	

Lys	Val	Met	Arg	Trp	Phe	Gln	Ala	Met	Leu	Gln	Arg	Leu	Gln	Thr	Trp
			100					105					110		

Trp	His	Gly	Val	Leu	Ala	Trp	Val	Lys	Glu	Lys	Val	Val	Ala	Leu	Val
		115					120					125			

His	Ala	Val	Gln	Ala	Leu	Trp	Lys	Gln	Phe	Gln	Ser	Phe	Cys	Cys	Ser
	130					135					140				

Leu	Ser	Glu	Leu	Phe	Met	Ser	Ser	Phe	Gln	Ser	Tyr	Gly	Ala	Pro	Arg
145					150					155					160

Gly	Asp	Lys	Glu	Glu	Leu	Thr	Pro	Gln	Lys	Cys	Ser	Glu	Pro	Gln	Ser
			165						170					175	

Ser Lys

<210> 11

<211> 5000

<212> DNA

<213> Homo sapiens

<400> 11

gcctaggggtg gaccctatatt caatatgact ggtgtccttt ggaaagggga aagggggaca 60

gtcacacca ggcagaacgt gatgaagatg aagatggcca tctacaaggg caggagaaac 120

ctgaacagaa tcccagctcc gggccctcag aaggaccca cgctgccac attgaccttg 180

gacctccagc ctgcagatcg tgaggggaaga gacgtcttcg acttagggcc ccttgctcgtg	240
gtacttcctt agtttggccc caggaaacca tcccaaaggc aagggcgtgg ttgtgctcag	300
ctgggggaag ggggctgggg gccgtgagga ggaggtggga ggcccagcca ggctggaggg	360
tcagaacctg tggagctaga agagcccgtg ggggagcccc aagattgctg agaccagtga	420
ccttcggccc cagatggcct tgccttggcc cagaagggtc agaaggacct ggtcagccaa	480
gctcagacag ccggcaggat gccttcacc ctgcagaggg tcctatcttg tcccacaggt	540
agatctacat caccactagc caccctcca acgtgcacag gccctgccc tcacggcgcc	600
cctcttaggt ccggcagttc ctgcctcctt ctgatccaga agtttctctg gcctctggag	660
ccggggcaca cctcatgcaa ggacagggtc caaatcctt tgtccttgga tcccacttgg	720
ctgacgtcac cttcctgtac tcaggaggtt tcccagcca gctgtcccga gtctggactt	780
tccctctgcc cctccccact ctcaggctgg tgggggtgggg aaagcagccc attcctgggc	840
tcagagactc ccacccagc tcagagggag caggggcccc gccagggacg gacctcatt	900
cctcccaggg accccagacc tctgtctctc tcgggtaagt ctccatctct gtctgtctct	960
gtctctgtct ctgtctctgt ctgtttttca cgcactcagc aaggcctcct gccctgagag	1020
aggtccgcc cactaccccc cactttcccc ataaaaccag ctgagtattt gtgccaggaa	1080
gactgcgtgc agaagggtgac tgtctcagtg gagctgggtc atctcaggtg gggagttggg	1140
gtccccgaag gtgaggacct tctggggagg aggggtgctt tctgagacac tttcttttcc	1200
tcacacctgt tcctcgccag caggccttgg ctcttgaac ttttgccgc catgtgcttc	1260
ccgaagggtga gtgagaggct gcgtgtgctt ttgtgggcat gtctgaaaac agaccgtaag	1320
ggtgcgggtg ccctcagtat tccccaggt gcctgtgtgt cagggtcag tcaggggcac	1380
ccagcggcag gaggatagt atggggtgag agtgtcagtg gaggcgctgg aggtcatatg	1440
tgtcgggggc gctggagaac ggcaggggtg tggatgagag ggagcacctg tcccaggagc	1500
ccttcacagc ccggaaagcc cggggcaggg gtggggcagg gctctgctgg aaacgactcg	1560
gagaatgctt ctctcagagg ccggctcagc tgggtgggcc caagagcaag gcctgtgtgg	1620
gtcctgggtg ctcttcctcc tttcctgggt tccctccgac ctcccatcct ctaccactgc	1680
cccaccgcaa atgctaggcc caccacacc tccagggagc tcttcggcct gtgacaatag	1740
gggtttccat gatgtggcct ggctcaggtt caggacagtg acccgagga cacatggctc	1800
ccgcatgtcg gcacgggtgt gctttcacc tggttcctgg gaaatcaggc tagcgggatg	1860
ggaccatcgc tgcctgaaag tgtgcagaca gctgccctgc ccagaatatg tcccaggcc	1920
ctgcgcactc tgtgggtgac tgtcaccact ctatagtggg ggaaaccagg catgtcacc	1980
ccgagactag gcccttgacg tgggggctca gcggggattc tgtgggggtgc ctctggcctc	2040

tgtggatgca gccacgtgtc tgcaggcagg aatggcccgg gacctgtggg tctgcatgtt	2100
ggcagtcggg aagagtggca ggttgtaggg tggacctacc tggcacccca aatattaatc	2160
agctcatcag agaggaatgg ctgctgttac cttctcaatt gtcatgtccc taaacatttt	2220
ttccttggcc aactctcacc tgggaccata gtggttgtgg gaaaccacgc tgagccagcc	2280
tgctccagga cagtgtccat cctcccgtgt gtgtacatgg gggggtgtgt gtgtgcaggg	2340
aggacacccc ggcccacgca ggccctgctc ttgtgaggag gggtcaccta ggcccacgca	2400
ggccctgctc ttgtgaggag gggtcaccta ggcccacgca ggccctgctc ttgtgaggag	2460
gggtcaccta ggcccacgca ggccctgctc ttgtgaggag gggtcaccta ggcccacgca	2520
ggccctgctc ttgtgaggag gggtcaccta ggcccacgca ggccctgctc ttgggcctgc	2580
ccagctgagc cggctcctga gagaagcgtt ttctgagtcg ttctgaggac agccctggcc	2640
ggctcttcca ggctgtgagg ggctcctggg actgctgtct cctcttatcc tgtacctctg	2700
ccatgtgtct ctgtgtgtgt gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt ataaattatc	2760
ctggaggaaa ggttaagggtg acacatggag actgagtgct accgttatct ccgcaggtcc	2820
tctctgatga catgaagaag ctgaaggccc gaatggtaat gctcctccct acttctgctc	2880
aggggttggg ggccctgggtc tcagcgtgtg aactgagga cactgtggga cacctgggac	2940
cctggaggga caaggatccg gccctttggg gccaaactctg cctctcttca cagcaccagg	3000
ccatagaaag attttatgat aaaatgcaaa atgcagaatc aggacgtgga caggtgggtg	3060
gatttcccct caggcaccag gtcacatgtc cccgccccca ggcactccac cctgtgtggg	3120
gctcaggggtg agaaggatga agagggaccc acaggctccc tcacccctta ccgtgggcaa	3180
atgcttgcac ctgggtggca gtgagtgggc ggggtggggga tctggacgcc cggggagact	3240
gagggaggca tccaagcccc agggctcctt gaggaacaaa caggggtgcc agacgtggcc	3300
cgggcccctg gctgggcccc gttcgggggtg tgtgggagct gaggactcac tgggcttgag	3360
gactgactga tgtgggggtgc agaggaggct tgggcctgga accgagtgtt ttgttcctaa	3420
caggtgatgt cgagcctggc agagctggag gtgagccgtg gcctccccct ccaccaagct	3480
tagtccctgg gtcttaggct ccacaggaca ctgggtctgg gccccgggtc cccttgggaa	3540
tcacctggac cagtgggggc cacagtggga agggggcagg caggagcagc atgaaccccc	3600
tgtgccctcc tctccccagg acgacttcaa agagggtac ctggagacag tggcggctta	3660
ttatgaggag cagcaccacg tgagtatgac acacccatct gggcaccttg ccttccttca	3720
cctctgccct gtcttttctt tctttcttct tttttgttta tttgagacag agtctcgctc	3780
tgtcgcccag gctggagtgc agtggcatga tcttggctca ctgcaacctc caaatctcgg	3840
gtttaagtga ttctcctgcc tcagcctgac aagtagttgg gactacaggc acccgccacc	3900
actccaggct gatttttttt gtgtgtgttt ttagtagaga ccaggtttca ccatgtttgc	3960

caggctggtc ttgaactcct aaccttgtgt tccgtctgcc ttggcctccc aaagtgctga 4020
 gattacaggc atgagccacc gggcccagcc aaccctgcc ctgtcttgat gtggtgtggg 4080
 cagggtgtgc ccagcccctg agcttggggg ggagggctgg gaggacagc ctagctggga 4140
 cctgcccattg gcctcactcc tcacacagtg gcacagccct caaggcacga tgagggccct 4200
 gacctggtga ccaagcagac acacccatcc tgtcactgcc atggaggtga atgcagagga 4260
 gggggactct gggaaaagtc cctcttgccc acggggctgt ggttgggaaa ccaacacctg 4320
 tgggcctccg tctcccaggg tcaggaaaag gctgagaggc ctgggtgtgg ccagggcctg 4380
 gggctgacac cccacctac agaccctgaa tgggtgctccc attccacagg agctcactcc 4440
 tctacttgaa aaagaaagag atggattacg gtgccgaggc aacagatccc ctgtcccgga 4500
 tggtgaggat cccgcaaccg aggagcctgg ggagagcttt tgtgacaagg tcatgagatg 4560
 gttccaggcc atgctgcagc ggctgcagac ctggtggcac ggggttcttg cctgggtgaa 4620
 ggagaagggtg gtggccctgg tccatgcagt gcaggccctc tggaaacagt tccagagttt 4680
 ctgctgctct ctgtcagagc tcttcatgtc ctctttccag tcctacggag cccacgggg 4740
 ggacaaggag gagctgacac ccagaagtg ctctgaacct caatcctcaa aatgaagata 4800
 ctgacaccac ctttgccctc cccgtcaccg cgcaccacc ctgaccctc cctcagctgt 4860
 cctgtgcccc gccctctccc gcacactcag tccccctgcc tggcgttcct gccgcagctc 4920
 tgacctgggtg ctgtcgccct ggcattctaa taaaacctgc ttatacttcc ctggcagggg 4980
 agataccatg atcgcgaggg 5000

<210> 12
 <211> 285
 <212> DNA
 <213> Mus musculus

<400> 12
 accgtcacat tgcccaacac ttgcttacia atctagaact cttgttttgc aacctgccaa 60
 ccttcttttt tgettccct gttttttctc ctaggaagtg tgtaagaca gtaccctgt 120
 atacttacct ggcaggggag ataccatgat cacgaagggt gttttcccag ggcgaggctt 180
 atccattgca ctccggatgt gctgaccctc gcgatttccc caaatgcggg aaactcgact 240
 gcataatttg tggtagtggg ggactgcgtt cccgctctcc cctgg 285

<210> 13
 <211> 26
 <212> PRT
 <213> Homo sapiens

<400> 13

Leu Lys Ala Arg Met His Gln Ala Ile Glu Arg Phe Tyr Asp Lys Met

1 5 10 15

Gln Asn Ala Glu Ser Gly Arg Gly Gln Val
20 25

<210> 14
<211> 46
<212> PRT
<213> Homo sapiens

<400> 14

Val Met Leu Leu Pro Thr Ser Ala Gln Gly Leu Gly Ala Trp Val Ser
1 5 10 15

Ala Cys Asp Thr Glu Asp Thr Val Gly His Leu Gly Pro Trp Arg Asp
20 25 30

Lys Asp Pro Ala Leu Trp Cys Gln Leu Cys Leu Ser Ser Gln
35 40 45

<210> 15
<211> 108
<212> DNA
<213> Homo sapiens

<400> 15
ccgaagggtcc tctctgatga catgaagaag ctgaaggccc gaatgcacca ggctatagaa 60
agatttttatg ataaaatgca aaatgcagaa tcaggacgtg gacagggtg 108

<210> 16
<211> 179
<212> PRT
<213> Equus caballus

<400> 16

Met Gly Tyr Pro Lys Thr Ser Arg Glu Asp Asn Glu Arg Trp Lys Ile
1 5 10 15

Arg Phe His Ser Thr Leu Asp Arg Trp Leu Asp Asp Ile Glu Val Gln
20 25 30

Ser Gln Gly Glu Glu Gln Val Asp Leu Gly Leu Glu Asp Leu Glu Glu
35 40 45

Lys Phe Ser Glu Asn Ile Leu Asp Ala Val Glu Glu His His Gln Lys
50 55 60

Asn Asn Ser Glu Ser Ala Pro Leu Leu Pro Asp Val Lys Pro Arg Leu
65 70 75 80

Arg Arg Arg Ala Gln Lys Ser Ser Val Leu Asn Pro Glu Pro Glu Gly
85 90 95

Pro Gly Ile Leu Gln Val Glu Ala Leu Glu Ala Pro Glu Pro Glu Glu
100 105 110

Ser Phe Trp Val Arg Ala Trp Arg Ser Phe Met Gly Met Leu Gln Arg
115 120 125

Leu Lys Gln Arg Trp Gln Ala Val Leu Ala Trp Val Arg Glu Lys Val
130 135 140

Ala Ala Gly Trp Gln Ala Leu Cys Ser Val Ala Gln Ser Ile Asn Ser
145 150 155 160

Val Leu Glu Ser Phe Cys Ser Tyr Met Ala Gly Leu Phe Arg Tyr His
165 170 175

Ile Gln Val

<210> 17

<211> 170

<212> PRT

<213> Bos taurus

<400> 17

Met Cys Phe Ala Lys Gly Val Pro Tyr Asp Gln Ala Ser Leu Arg Ser
1 5 10 15

Ile Met His Lys Arg Val Asp Asp Phe Cys Asp Lys Met Gly Asn Glu
20 25 30

Pro Glu Glu Ala Gln Met Glu Ala Ala Leu Asp Glu Thr Glu Glu Gly
35 40 45

Leu Ser Glu Asp Ile Cys Glu Phe Ile Glu Asp His Ile Gln Glu Asn
50 55 60

Leu Pro Glu Ser Leu Gln Glu Ser Ser Pro Leu Leu Gln Glu Ala Arg
65 70 75 80

Gln Gly Val Arg Arg Arg Ile Gln Arg Pro Ser Val Ser Ala Arg Leu
85 90 95

Glu Val Gln Asn Pro Glu Glu Ser Ile Trp Ala Arg Ala Leu Gly Arg
100 105 110

Phe Gln Val Ile Leu Gln Ser Leu Gln Gln Arg Cys Trp Asp Ala Leu
 115 120 125

Thr Trp Leu Arg Glu Lys Ala Val Thr Phe Leu Glu Ala Ile Cys Ser
 130 135 140

Val Val Lys Ala Val Leu Gly Val Leu Thr Asp Phe Cys Ser Ser Val
 145 150 155 160

Gly Gln Leu Phe Gly Asn Leu Ile Gln Val
 165 170

<210> 18
 <211> 125
 <212> PRT
 <213> Equus caballus

<400> 18

Met Gly Tyr Pro Lys Thr Ser Arg Glu Asp Asn Glu Arg Trp Lys Ile
 1 5 10 15

Arg Phe His Ser Thr Leu Asp Arg Trp Leu Asp Asp Ile Glu Val Gln
 20 25 30

Ser Gln Gly Glu Glu Gln Val Cys Gln Cys Ala Pro Thr Pro Cys Ser
 35 40 45

Arg Asn Leu Gly Gly Arg Val Val Thr Met Thr Met Arg Arg Lys Asn
 50 55 60

Val Pro Pro Gln Val Asp Leu Gly Pro Leu Thr Ser Pro Phe Ser Gln
 65 70 75 80

Arg Thr Phe Arg Ser Asp Leu Cys His Leu Pro Thr Leu Asp Leu Ser
 85 90 95

Leu Thr Thr Ser Leu Thr Ser Leu Leu Cys Thr Ala Trp Pro Pro Cys
 100 105 110

Pro Pro Cys Thr Ser Cys Ser Gly Phe Leu Leu Gln Val
 115 120 125

<210> 19
 <211> 482
 <212> DNA
 <213> Equus caballus

<400> 19

gcacgagctc gtgccgtgtg ctgagaggcc cttggggcag gcacagcccc tggaatcctg

60

agctgccatg ggctacccca agacgtccag agaagacaat gaacgttgga agatccgatt 120
 tcacagcact ttagaccggt ggcttgatga tatcgaagtt caatcccaag gagaggaaca 180
 ggtgtgtcag tgtgtctcca cgccttgcct ccgtaacctc gggggtcggg tggtcacgat 240
 gacgatgagg aggaagaacg tgccacctca ggctcgattta ggccctttga cgtccccctt 300
 ttcacagaga accttcagaa gtgacctttg ccacctgcct accttgacc tgtccttgac 360
 cacctccctc acctccttgc tgtgcacagc ctggccaccc tgcccacat gcacttcctg 420
 ctcaggtttc cttctgcagg tctgacttgt ggctccagcg catatgtctt aataaagttg 480
 tg 482

<210> 20
 <211> 815
 <212> DNA
 <213> Equus caballus

<400> 20
 ctgagaggcc cttggggcag gcacagcccc tggaatcctg agctgccatg ggctacccca 60
 agacgtccag agaagacaat gaacgttgga agatccgatt tcacagcact ttagaccggt 120
 ggcttgatga tatcgaagtt caatcccaag gagaggaaca ggctcgattta ggcctagaag 180
 acctggagga aaaattcagt gaaaacattc ttgacgccgt ggaggagcac catcagaaga 240
 acaactcaga atctgcgcct ttacttcctg acgtgaagcc caggttacgt cgcagagctc 300
 agaagtcctc tgtcctcaac cctgaacctg agggctccagg gatcctgcaa gttgaggctc 360
 tagaggcacc cgagcctgaa gaaagctttt gggtcagagc atggaggctg ttcattgggga 420
 tgctacagcg actgaagcag aggtggcagg ctgtactggc ctgggtgcga gagaaggctg 480
 ctgctggctg gcaggcccta tgcagtgtgg ccagtcctat taatagtgtg cttgagagtt 540
 tctgctccta tatggctggg ttgttttaggt accacatcca ggtctagggg gccccatggg 600
 gtccaggagg ggtagccaca ccttgacgcc ctttgacgct ccccttttca cagagaacct 660
 tcagaagtga cctttgccac ctgcctaccc ttgacctgtc cttgaccacc tccttcacct 720
 ccttgctgtg cacagcctgg ccacctgcc caccatgcac ttctgctca ggtttccttc 780
 tgcaggctctg acttgtggct ccagcgcata gtctt 815

<210> 21
 <211> 724
 <212> DNA
 <213> Bos taurus

<400> 21
 cggattcccc ggatgctcag ctggagctct ggctgcagga tctcaggctc cttcgggagg 60
 accctaagcc accatgtgct tcgctaaggg agtcccatat gaccaggctt ctctgaggctc 120

```

cataatgcac aaacgggtgg atgatttctg tgataagatg ggaaatgaac cagaagaagc 180
acagatggag gcagccctag atgagacgga ggagggactc agcgaggaca tctgtgaatt 240
catagaagat cacattcaag agaaccctcc cgaatccctg caggagtcca gtcccttgct 300
tcaggaagca cggcaaggag tacgccgcag aatccagaga ccttcagtct ctgccgtct 360
ggaggtccag aatccggaag agagcatctg ggccagagcc ctggggaggt tccaagtaat 420
tctgcagagt ctccagcagc ggtgttgga tgcgctcacc tggctgcggg agaaggcgg 480
gaccttcctg gaggccatct gcagtgtggt gaaggccgtc ttgggagtgc tgacggattt 540
ctgctcctct gtggggcagc tcttcggaaa cctcatccag gtctaggagc cgcaggtggt 600
tctggaggaa ctctcctca tctaggaggc cctgcaccat ccccttccca gaaaccatct 660
tgtgaagcga cctttgcact cctgctcacc cttgacccat cctttaactg ccctcacctc 720
ctgt 724

```

```

<210> 22
<211> 218
<212> PRT
<213> Bos taurus

```

```

<400> 22

```

```

Met Cys Phe Thr Lys Arg Asp Pro Arg Val Leu Ala Ser Phe Arg Val
1          5          10          15

```

```

Leu Met Val Arg Ser Ser Phe Pro Arg Ile Ala Gly Val Arg Glu Ala
          20          25          30

```

```

Trp Val Leu Leu Gly Glu Ala Glu Asn Ile Leu Ala His Leu Gly Pro
          35          40          45

```

```

Ser Arg Glu Lys Asn Arg Asp Ser Phe Thr Gln Val His Leu Cys Ser
          50          55          60

```

```

Gln His Asn Leu Val Asp Glu Phe Phe Asp Thr Met Glu Asn Glu Pro
65          70          75          80

```

```

Glu Gly Ala Gln Met Glu Ala Val Leu Ala Glu Thr Lys Glu Lys Phe
          85          90          95

```

```

Ile Lys Asp Ala Phe Lys Val Met Asp Asn His Ile Gln Glu Asn Ser
          100          105          110

```

```

Pro Glu Thr Leu Lys Glu Ser Ser Pro Leu Leu Gln Glu Ala Arg Gln
          115          120          125

```

```

Glu Val Arg Cys Arg Ile Gln Arg Arg Ser Val Ser Thr Ser Leu Glu

```

130	135	140
Val Gln Asn Pro Glu Glu Ser Ile Trp Ala Arg Ala Leu Arg Gln Phe		
145	150	155 160
Leu Gly Ile Leu Gln Ser Phe Leu Ser Gly Cys Arg Asp Ala Leu Thr		
	165	170 175
Trp Leu Trp Glu Lys Ala Ala Ala Cys Leu Gln Ala Ile Cys Ser Ala		
	180	185 190
Val Glu Ala Leu Trp Glu Val Leu Thr Asp Phe Cys Ser Phe Val Gly		
	195	200 205
Gln Leu Leu Cys Arg Ser Leu Ile Gln Val		
	210	215

<210> 23
 <211> 863
 <212> DNA
 <213> Bos taurus

<400> 23
 cgggatctca gctggagctc tggctgcagg atctcaggtc ccagcggcag gaccctaagc 60
 caccatgtgc ttcactaaga gagaccacg tgtcctggct tctttcaggg tgttaatggt 120
 aagaagctca tttccacgta tagctggggg tctggaggcc tgggttctgc tgggtgaagc 180
 tgagaacatt ctggcccact tgggaccag cagggagaag aaccgagatt cttttactca 240
 agtccatctc tgttcacagc acaaccttgt agatgaattt ttcgatacaa tggaaaatga 300
 accagaagga gcacagatgg aggcagtcct agcagagact aaggagaaat tcatcaagga 360
 cgcctttaaa gtcattggata atcacattca agagaacagt cccgaaaccc tgaaggagtc 420
 cagtccttg cttcaggaag cacggcaaga agtacgctgc agaatccaga gacgctccgt 480
 ctccacctct ctggagggtcc agaatccgga agagagcatc tgggccagag ccctgcggca 540
 gttcttgggc attctgcaga gtttcctgtc cgggtgtcgg gatgcgctca cctggctgtg 600
 ggagaaggcc gcggcctgcc tacaggccat ctgcagtgcg gtggaggccc tctgggaagt 660
 gctcacggat ttctgctcct ttggtgggca gctcttatgc agaagcctca tccaggtcta 720
 agagcctcac atggttctgg aggagcccca cctcattcag aaggccctgt acgatgccct 780
 tcccggaaac catcttctga agcgaccttt accctcctgc tcacccttga cccatccttt 840
 aactgccctc ccctcctgtc ctg 863

<210> 24
 <211> 127
 <212> PRT

<213> Ovis aries

<400> 24

Met Cys Phe Ala Arg Gly Val Pro His Asp Gln Ala Ser Leu Arg Ser
 1 5 10 15

Met Leu His Thr Trp Val Asp His Val Cys Asp Lys Met Gly Asn Glu
 20 25 30

Pro Glu Glu Ala Gln Met Glu Ala Ala Leu Ala Glu Met Glu Glu Glu
 35 40 45

Leu Ser Lys Asp Val Cys Glu Ser Trp Lys Ile Thr Phe Lys Arg Thr
 50 55 60

Phe Pro Asn Pro Cys Arg Ser Pro Val Pro Cys Phe Arg Lys Arg Ser
 65 70 75 80

Lys Lys Tyr Ala Ala Glu Ser Arg Asp Pro Gln Ser Leu Pro Val Trp
 85 90 95

Arg Thr Arg Asn Arg Lys Arg Ala Ser Gly Pro Glu Pro Cys Gly Gly
 100 105 110

Ser Glu Val Phe Cys Gly Val Ser Gly Ser Gly Val Ala Met Tyr
 115 120 125

<210> 25

<211> 811

<212> DNA

<213> Ovis aries

<400> 25

ctgcggtacc ggtccggatt cccgggagac acagtgtctc gctggagctc tggctgcagg 60
 atctcagatc ccagccggag gaccctaata caccatgtgc ttcgctaggg gagtcccaca 120
 tgaccaggct tctctgagga gcatgctgca cacctgggtg gatcatgtct gtgataagat 180
 gggaaatgaa ccagaagaag cacagatgga ggcagcccta gcagagatgg aggaggaact 240
 cagcaaggat gtctgtgaat catggaagat cacattcaag agaaccttcc cgaatccctg 300
 caggagtcca gtcccttgct tcaggaagcg cagcaagaag tacgccgcag aatccagaga 360
 ccctcagtct ctgcctgtct ggaggaccag aaaccggaag agagcatctg ggccagagcc 420
 ctgcggcggt tccgaggttt tctgcggagt ctctggcagc ggtgttgcca tgtactgacc 480
 tggctgcagg agaaggcggc ggctgcctg gaggccgtct gcagtgcggt gaagaccatc 540
 tggggagtgc tgacggattt ctgctcctct gtggggcagc tcttcagaaa cctcatccag 600
 gtctaggagc cccaggtcgt tcttgaggaa ctgctcctca tctagaaggc cctgcacaat 660

ccccttccca gaaaccatct tctgaagcga cctttaccct cctgttcacc cttcaccaat 720
 cctttaactg ccctcacctc ctgtctgcag ggacgacacc acaacatcaa gccaggtttc 780
 ccttctccaa gtctgacccg tctgtcaggg a 811

<210> 26
 <211> 141
 <212> PRT
 <213> Sus scrofa

<400> 26

Met Arg Gly Val Ser Ala Thr Arg Thr Leu Pro Lys Ala Gly Pro Gln
 1 5 10 15

Pro Arg Ser Gly Leu Gly Leu Pro Leu Pro Arg Arg Val Pro Glu Pro
 20 25 30

Pro Pro Ile Pro Ala Glu Ser Ser Pro Leu Leu Asn Glu Val Arg Gln
 35 40 45

Gly Val Arg Ser Arg Val Arg Arg Pro Pro Gly His Asn Gln Pro His
 50 55 60

Tyr Ala Leu Ala Val Arg Glu Pro Arg Gln Ser Thr Phe Arg Arg Ile
 65 70 75 80

Leu Glu Leu Phe Glu Glu Met Leu Lys Arg Leu Gln Gln Arg Trp Arg
 85 90 95

Gly Ala Leu Ala Trp Val Gln Glu Arg Ala Ala Ala Cys Phe Arg Gly
 100 105 110

Leu Cys Arg Ala Leu Glu Ala Phe Trp Ser Leu Val Gln Ser Phe Cys
 115 120 125

Ser Ser Met Gly His Ala Phe Gly Ser Val Ile Gln Val
 130 135 140

<210> 27
 <211> 603
 <212> DNA
 <213> Sus scrofa

<400> 27

atgacttgga ggggaactga gcggccaggg ccagcccctg ggaaaagtcc tggggtctgt 60
 ggggctgttg gcaggaaagc agcctgtgtc caaggcgggg catgaggggg gtgtctgcca 120
 ccaggactct cccaaaggca gggcctcagc caaggtcagg actggggctg cctctcccca 180

ggcgggtccc tgaaccaccc cccatccctg cagaatccag tcctctgctc aacgaagtcc 240
ggcagggagt ccgttctaga gtccgaaggc ctcttgcca caaccagcca cattatgcgc 300
tagcgggtccg ggagcccagg cagagcactt tcagacgcat ccttgagctg tttgaggaaa 360
tgctgaagcg cctgcagcag aggtggaggg gtgccctggc ttgggtgcag gaaagggctg 420
ctgcctgctt ccggggcttg tgcagggccc ttgaagcttt ctggagcctg gtgcagagtt 480
tttgctcttc catggggcac gccttcggga gtgtcatcca ggtctaaggc gctccaggctg 540
aaataagagt ttctagagca caacctcccc ctgccttggc taaaaaggca gctgtaagcc 600
ttt 603


CLAIMS:


1. A purified nucleic acid encoding an IL-32 alpha isoform consisting of the amino acid sequence set forth as SEQ ID NO:7.
2. The nucleic acid of Claim 1, wherein said sequence is operably linked to a heterologous promoter.
3. The nucleic acid of Claim 2, wherein said nucleic acid is contained within a vector.
4. A host cell comprising the nucleic acid of Claim 3.
5. A purified IL-32 alpha protein encoded by the nucleic acid of Claim 1.
6. The protein of Claim 5, wherein said IL-32 alpha is a recombinant protein expressed in a cell selected from the group consisting of a bacterial cell, a yeast cell, an insect cell, and a mammalian cell.
7. The protein of Claim 6, wherein said recombinant protein is a fusion protein.
8. An antibody, which specifically binds to the IL-32 alpha protein of Claim 5.
9. The antibody of Claim 8, wherein said antibody is a monoclonal antibody.
10. A Fab fragment of the monoclonal antibody of Claim 9.
11. The monoclonal antibody of Claim 9, wherein said monoclonal antibody inhibits IL-32-induced TNF α production by a target cell.
12. The monoclonal antibody of Claim 9, wherein said monoclonal antibody inhibits IL-32-induced I κ B degradation in a target cell.

13. The monoclonal antibody of Claim 9, wherein said monoclonal antibody inhibits rapid IL-32-induced p38 MAPK phosphorylation in a target cell.
14. Use of an IL-32 alpha protein of Claim 5 in the preparation of a medicament for treating a subject for a cancer.
15. Use of an antibody of Claim 11 in the preparation of a medicament for treating a subject for an autoimmune disease.
16. A method for screening for inhibitors of IL-32, comprising:
 - a) providing the IL-32 alpha isoform protein of Claim 5, and at least one drug candidate; and
 - b) analyzing the effect of said drug candidate on the induced secretion of a TNF α or MIP-2 by said IL-32 protein.
17. The method of Claim 16, wherein said drug candidate is selected from the group consisting of a IL-32-reactive monoclonal antibody, and a dominant-negative IL-32 variant.
18. The method of Claim 16, wherein the induced secretion of TNF α comprises upregulation of TNF α expression.
19. Use of an IL-32 protein of Claim 5 for treating a subject for a cancer.
20. The use of Claim 19, wherein said subject has, is suspected of having, or is at risk of having the cancer.
21. Use of an IL-32 alpha protein antibody of Claim 8 for treating a subject for an autoimmune disease.
22. The use of Claim 21, wherein said subject has, is suspected of having, or is at risk of

having the autoimmune disease.

23. The use of Claim 21, wherein the autoimmune disease is rheumatoid arthritis or inflammatory bowel disease.

**Fig. 1**

**Fig. 2**

IL-32 α	ATGTGCTTCCCGAAGGTCCTCTCTGATGACATGAAGAAGCTGAAGGCCCGAATG-----
IL-32 β	ATGTGCTTCCCGAAGGTCCTCTCTGATGACATGAAGAAGCTGAAGGCCCGAATG-----
IL-32 γ	ATGTGCTTCCCGAAGGTCCTCTCTGATGACATGAAGAAGCTGAAGGCCCGAATGGTAATG
IL-32 δ	-----ATGAAGAAGCTGAAGGCCCGAATG-----
IL-32 α	-----
IL-32 β	-----
IL-32 γ	CTCCTCCCTACTTCTGCTCAGGGGTTGGGGGCCTGGGTCTCAGCGTGTGACACTGAGGAC
IL-32 δ	-----
IL-32 α	-----
IL-32 β	-----
IL-32 γ	ACTGTGGGACACCTGGGACCCTGGAGGGACAAGGATCCGGCCCTTTGGTGCCAACTCTGC
IL-32 δ	-----
IL-32 α	-----CACCAGGCTATAGAAAGATTTTATGATAAAATGCAAAATGCAGAATCA
IL-32 β	-----CACCAGGCCATAGAAAGATTTTATGATAAAATGCAAAATGCAGAATCA
IL-32 γ	CTCTCTTCACAGCACCAGGCCATAGAAAGATTTTATGATAAAATGCAAAATGCAGAATCA
IL-32 δ	-----CACCAGGCCATAGAAAGATTTTATGATAAAATGCAAAATGCAGAATCA
IL-32 α	GGACGTGGACAGGTGATGTCGAGCCTGGCAGAGCTGGAGGACGACTTCAAAGAGGGCTAC
IL-32 β	GGACGTGGACAGGTGATGTCGAGCCTGGCAGAGCTGGAGGACGACTTCAAAGAGGGCTAC
IL-32 γ	GGACGTGGACAGGTGATGTCGAGCCTGGCAGAGCTGGAGGACGACTTCAAAGAGGGCTAC
IL-32 δ	GGACGTGGACAGGTGATGTCGAGCCTGGCAGAGCTGGAGGACGACTTCAAAGAGGGCTAC
IL-32 α	CTGGAGACAGTGGCGGCTTATTATGAGGAGCAGCACCCAGAGCTCACTCCTCTACTTGAA
IL-32 β	CTGGAGACAGTGGCGGCTTATTATGAGGAGCAGCACCCAGAGCTCACTCCTCTACTTGAA
IL-32 γ	CTGGAGACAGTGGCGGCTTATTATGAGGAGCAGCACCCAGAGCTCACTCCTCTACTTGAA
IL-32 δ	CTGGAGACAGTGGCGGCTTATTATGAGGAGCAGCACCCAGAGCTCACTCCTCTACTTGAA

Fig. 3A

IL-32 α	AAAGAAAGAGATGGATTACGGTGCCGAGGCAACAGATCCCCTGTCCCGGATGTTGAGGAT
IL-32 β	AAAGAAAGAGATGGATTACGGTGCCGAGGCAACAGATCCCCTGTCCCGGATGTTGAGGAT
IL-32 γ	AAAGAAAGAGATGGATTACGGTGCCGAGGCAACAGATCCCCTGTCCCGGATGTTGAGGAT
IL-32 δ	AAAGAAAGAGATGGATTACGGTGCCGAGGCAACAGATCCCCTGTCCCGGATGTTGAGGAT
IL-32 α	CCCGCAACCGAGGAGCCTGGGGAGAGCTTTTGTGACAAG-----
IL-32 β	CCCGCAACCGAGGAGCCTGGGGAGAGCTTTTGTGACAAGGTCATGAGATGGTTCCAGGCC
IL-32 γ	CCCGCAACCGAGGAGCCTGGGGAGAGCTTTTGTGACAAGGTCATGAGATGGTTCCAGGCC
IL-32 δ	CCCGCAACCGAGGAGCCTGGGGAGAGCTTTTGTGACAAGGTCATGAGATGGTTCCAGGCC
IL-32 α	-----
IL-32 β	ATGCTGCAGCGGCTGCAGACCTGGTGGCACGGGGTTCTGGCCTGGGTGAAGGAGAAGGTG
IL-32 γ	ATGCTGCAGCGGCTGCAGACCTGGTGGCACGGGGTTCTGGCCTGGGTGAAGGAGAAGGTG
IL-32 δ	ATGCTGCAGCGGCTGCAGACCTGGTGGCACGGGGTTCTGGCCTGGGTGAAGGAGAAGGTG
IL-32 α	-----
IL-32 β	GTGGCCCTGGTCCATGCAGTGCAGGCCCTCTGGAAACAGTTCCAGAGTTTCTGCTGCTCT
IL-32 γ	GTGGCCCTGGTCCATGCAGTGCAGGCCCTCTGGAAACAGTTCCAGAGTTTCTGCTGCTCT
IL-32 δ	GTGGCCCTGGTCCATGCAGTGCAGGCCCTCTGGAAACAGTTCCAGAGTTTCTGCTGCTCT
IL-32 α	-----TCCTACGGAGCCCCACGGGGGGACAAGGAG
IL-32 β	CTGTCAGAGCTCTTCATGTCCTCTTTCCAGTCCTACGGAGCCCCACGGGGGGACAAGGAG
IL-32 γ	CTGTCAGAGCTCTTCATGTCCTCTTTCCAGTCCTACGGAGCCCCACGGGGGGACAAGGAG
IL-32 δ	CTGTCAGAGCTCTTCATGTCCTCTTTCCAGTCCTACGGAGCCCCACGGGGGGACAAGGAG
IL-32 α	GAGCTGACACCCCAGAAGTGCTCTGAACCCCAATCCTCAAAATGA
IL-32 β	GAGCTGACACCCCAGAAGTGCTCTGAACCCCAATCCTCAAAATGA
IL-32 γ	GAGCTGACACCCCAGAAGTGCTCTGAACCCCAATCCTCAAAATGA
IL-32 δ	GAGCTGACACCCCAGAAGTGCTCTGAACCCCAATCCTCAAAATGA

Fig. 3B


A


	1	Myr
IL-32 α	MCFPKVLSDDMKKLKARM-----	
IL-32 β	MCFPKVLSDDMKKLKARM-----	
IL-32 γ	MCFPKVLSDDMKKLKARMVMLLPTSAQGLGAWVSACDTEDTVGH LGPW RD KDPALWCQLC	
IL-32 δ	-----MKKLKARM-----	
	61	Myr
IL-32 α	----HQAIERFYDKMQNAESGRGQVMSSLAELEDDFKEGYLETVAAYYEEQHPELTPLLE	
IL-32 β	----HQAIERFYDKMQNAESGRGQVMSSLAELEDDFKEGYLETVAAYYEEQHPELTPLLE	
IL-32 γ	LSSQHQA IERFYDKMQNAESGRGQVMSSLAELEDDFKEGYLETVAAYYEEQHPELTPLLE	
IL-32 δ	----HQAIERFYDKMQNAESGRGQVMSSLAELEDDFKEGYLETVAAYYEEQHPELTPLLE	
	121	Gly
IL-32 α	KERDGLRCRGNRSPVPDVEDPATEEPGESFCDK-----	
IL-32 β	KERDGLRCRGNRSPVPDVEDPATEEPGESFCDKVMRWFQAMLQRLQTWWHGVLA W VKEKV	
IL-32 γ	KERDGLRCRGNRSPVPDVEDPATEEPGESFCDKVMRWFQAMLQRLQTWWHGVLA W VKEKV	
IL-32 δ	KERDGLRCRGNRSPVPDVEDPATEEPGESFCDKVMRWFQAMLQRLQTWWHGVLA W VKEKV	
	181	Myr
IL-32 α	-----SYGAPRGDKEELTPQKCSEPQSSK	
IL-32 β	VALVHAVQALWKQFQSFCCSLSELFMSSFQSYGAPRGDKEELTPQKCSEPQSSK	
IL-32 γ	VALVHAVQALWKQFQSFCCSLSELFMSSFQSYGAPRGDKEELTPQKCSEPQSSK	
IL-32 δ	VALVHAVQALWKQFQSFCCSLSELFMSSFQSYGAPRGDKEELTPQKCSEPQSSK	


B


	1
huIL-32 β	MCFPKVLSDDMKKLKARMHQAIERFYDKMQNAESGRGQVMSSLAELEDDFKEGYLETVA A
EqIL-32	MGYPKTSREDNERWKIRFHSTLDRWLDDIEVQSQGEEQVDLGLEDLEEKFSENILDAVEE
BoIL-32	MCFAGVVPYDQASLR SIMHKRVDDFCDKMGNEPE-EAQMEAALDETEEGLSEDICEFIED
Consensus	*-----*-----*-----*-----*-----*-----*-----
	61
huIL-32 β	YYEEQHPELTPLLEKERDGLRCRGNRSPV----PDVEDP----ATE--EPGESFCDKVMR
EqIL-32	HHQKNNSSESAPLLPDVKPRLRRRAQKSSVLNPEPEPGILQVEALEAPEPEESFWVRAWR
BoIL-32	HIQENLPES--LQESSPL-LQEARQGVRRRIQRPSV-----SARLEVQNPEESI----WA
Consensus	-----*-----*-----*-----*-----*-----*-----*
	121
huIL-32 β	WFQAMLQR-L-QTWWHGVLA W VKEKVVA-----LVHAVQALWKQFQS--FCCSLSELF
EqIL-32	SFMGMLQR-LKQRWQAVLA-WVREKVAAGWQA--LCSVAQSINSVLES--FCSY MAGLF
BoIL-32	RALGRFQVIL-QSLQQR--WDALTWLREKAVTFLEAICSVVKAVLGVLTDFCSSVGQLF
Consensus	-----*-----*-----*-----*-----*-----*-----*
	181
huIL-32 β	MSSF---QSYGAPRGDKEELTPQKCSEPQSSK
EqIL-32	RYH---IQV-----
BoIL-32	---GNLIQV-----
Consensus	-----*-----







Fig. 4


**Fig. 5**


**Fig. 6**

**Fig. 7**

**Fig. 8**

A**B****C****D****E****F****Fig. 9**

**Fig. 10**

**Fig. 11**

A**EqIL-32 alpha (SEQ ID NO:18)**

MGYPKTSREDNERWKIRFHSTLDRWLDDIEVQSQGEEQVCQCAPTPCSRNLGGRVVTMTMRRKNVPPQVD
LGPLTSPFSQRTFRSDLCHLPTLDLSLTSLTSLCTAWPPCPPCTSCSGFLLQV

B**EqIL-32 alpha (SEQ ID NO:19)**

GCACGAGCTCGTGCCGTGTGCTGAGAGGCCCTTGGGGCAGGCACAGCCCCCTGGAATCCTGAGCTGCCATG
GGCTACCCCAAGACGTCCAGAGAAGACAATGAACGTTGGAAGATCCGATTTCACAGCACTTTAGACCGGT
GGCTTGATGATATCGAAGTTCAATCCCAAGGAGAGGAACAGGTGTGTGCTCAGTGTGCTCCCACGCCCTGCTC
CCGTAACCTCGGGGGTCTGGGTGGTCACGATGACGATGAGGAGGAAGAACGTGCCACCTCAGGTCGATTTA
GGCCCTTTGACGTCCCCCTTTTCACAGAGAACCTTCAGAAGTGACCTTTGCCACCTGCCTACCCTTGACC
TGTCCTTGACCACCTCCCTCACCTCCTTGCTGTGCACAGCCTGGCCACCCTGCCACCATGCACTTCCTG
CTCAGGTTTCCTTCTGCAGGTCTGACTTGTGGCTCCAGCGCATATGTCTTAATAAAGTTGTG

C**EqIL-32 beta (SEQ ID NO:16)**

MGYPKTSREDNERWKIRFHSTLDRWLDDIEVQSQGEEQVDLGLEDLEEKFSENILDAVEEHHQKNNSESA
PLLPDVKPRRLRRRAQKSSVLNPEPEPGILQVEALEAPEPEESFWVRAWRSFMGMLQRLKQRWQAVLAWV
REKVAAGWQALCSVAQSINSVLESFCSYAGLFRYHIQV

D**EqIL-32 beta (SEQ ID NO:20)**

CTGAGAGGCCCTTGGGGCAGGCACAGCCCCCTGGAATCCTGAGCTGCCATGGGCTACCCCAAGACGTCCAG
AGAAGACAATGAACGTTGGAAGATCCGATTTCACAGCACTTTAGACCGGTGGCTTGATGATATCGAAGTT
CAATCCCAAGGAGAGGAACAGGTGATTTAGGCCTAGAAGACCTGGAGGAAAAATTCAGTGAAAACATTC
TTGACGCCGTGGAGGAGCACCATCAGAAGAACAACTCAGAATCTGCGCCTTTACTTCCTGACGTGAAGCC
CAGGTTACGTGCGCAGAGCTCAGAAGTCCTCTGTCTCAACCCTGAACCTGAGGGTCCAGGGATCCTGCAA
GTTGAGGCTCTAGAGGCACCCGAGCCTGAAGAAAGCTTTTGGGTGAGAGCATGGAGGTGTTTATGGGGA
TGCTACAGCGACTGAAGCAGAGGTGGCAGGCTGTACTGGCCTGGGTGCGAGAGAAGGTGGCTGCTGGCTG
GCAGGCCCTATGCAGTGTGGCCAGTCCATTAATAGTGTGCTTGAGAGTTTCTGCTCCTATATGGCTGGG
TTGTTTAGGTACCACATCCAGGTCTAGGGGGCCCCATGGGGTCCAGGAGGGGTAGCCACACCTTGACGCC
CTTTGACGTCCCCCTTTTCACAGAGAACCTTCAGAAGTGACCTTTGCCACCTGCCTACCCTTGACCTGTC
CTTGACCACCTCCCTCACCTCCTTGCTGTGCACAGCCTGGCCACCCTGCCACCATGCACTTCCTGCTCA
GGTTTCCTTCTGCAGGTCTGACTTGTGGCTCCAGCGCATAGTCTT

Fig. 12

A**BoIL-32 beta (SEQ ID NO:17)**

MCFAKGVPPYDQASLRSIMHKRVDDFCDKMGNEPEEAQMEAAALDETEEGLSEDI CEFIEDHIQENLPESLQ
ESSPLLQEARQGVRRIQRPSVSARLEVQNPEESIWARALGRFQVILQSLQQRCWDALTWLREKAVTFLE
AICSVVKAVLGVLTDFCSSVGQLFGNLIQV

B**BoIL-32 beta (SEQ ID NO:21)**

CGGATTCCCGGGATGCTCAGCTGGAGCTCTGGCTGCAGGATCTCAGGTCCCTTCGGGAGGACCCTAAGCC
ACCATGTGCTTCGCTAAGGGAGTCCCATATGACCAGGCTTCTCTGAGGTCCATAATGCACAAACGGGTGG
ATGATTTCTGTGATAAGATGGGAAATGAACCAGAAGAAGCACAGATGGAGGCAGCCCTAGATGAGACGGA
GGAGGGACTCAGCGAGGACATCTGTGAATTCATAGAAGATCACATTCAAGAGAACCTTCCCGAATCCCTG
CAGGAGTCCAGTCCCTTGCTTCAGGAAGCACGGCAAGGAGTACGCCGCAGAATCCAGAGACCTTCAGTCT
CTGCCCCGTCTGGAGGTCCAGAATCCGGAAGAGAGCATCTGGGCCAGAGCCCTGGGGAGGTTCCAAGTAAT
TCTGCAGAGTCTCCAGCAGCGGTGTTGGGATGCGCTCACCTGGCTGCGGGAGAAGGCGGTGACCTTCCTG
GAGGCCATCTGCAGTGTGGTGAAGGCCGTCTTGGGAGTGCTGACGGATTTCTGCTCCTCTGTGGGGCAGC
TCTTCGGAAACCTCATCCAGGTCTAGGAGCCGCAGGTGGTTCTGGAGGAACTCCTCCTCATCTAGGAGGC
CCTGCACCATCCCCTTCCCAGAAACCATCTTGTGAAGCGACCTTTGCACTCCTGCTCACCTTGACCCAT
CCTTTAACTGCCCTCACCTCCTGT

C**BoIL-32 gamma (SEQ ID NO:22)**

MCFTKRDPRLASFRVLMVRSSFPRRIAGVREAWVLLGAEENILAHLGPSREKNRDSFTQVHLC SQHNLVD
EFFDTMENEPEGAQMEAVLAETKEKFIKDAFKVMDNHIQENSPETLKESSPLLQEARQEVRCRIQRRSVS
TSLEVQNPEESIWARALRQFLGILQSFLSGCRDALTWLWEKAAACLQAICSAVEALWEVLTDFCSFVGQL
LCRS LIQV

D**BoIL-32 gamma (SEQ ID NO:23)**

CGGGATCTCAGCTGGAGCTCTGGCTGCAGGATCTCAGGTCCCAGCGGCAGGACCCTAAGCCACCATGTGC
TTCACTAAGAGAGACCCACGTGTCCTGGCTTCTTT CAGGGTGTTAATGGTAAGAAGCTCATTTCCACGTA
TAGCTGGGGTTTCGGGAGGCCTGGGTCTGCTGGGTGAAGCTGAGAACATTCTGGCCCACTTGGGACCCAG
CAGGGAGAAGAACCGAGATTCTTTTACTCAAGTCCATCTCTGTT CACAGCACAACCTTG TAGATGAATTT
TTCGATACAATGGAAAATGAACCAGAAGGAGCACAGATGGAGGCAGTCCTAGCAGAGACTAAGGAGAAAT
TCATCAAGGACGCCTTTAAAGTCATGGATAATCACATTCAAGAGAACAGTCCCGAAACCTGAAGGAGTC
CAGTCCCTTGCTTCAGGAAGCACGGCAAGAAGTACGCTGCAGAATCCAGAGACGCTCCGTCTCCACCTCT
CTGGAGGTCCAGAATCCGGAAGAGAGCATCTGGGCCAGAGCCCTGCGGCAGTTCTTGGGCATTCTGCAGA
GTTTCCTGTCCGGGTGTCGGGATGCGCTCACCTGGCTGTGGGAGAAGGCCGCGGCCTGCCTACAGGCCAT
CTGCAGTGCGGTGGAGGCCCTCTGGGAAGTGCTCACGGATTTCTGCTCCTTTGTTGGGCAGCTCTTATGC
AGAAGCCTCATCCAGGTCTAAGAGCCTCACATGGTTCTGGAGGAGCCCCACCTCATT CAGAAGGCCCTGT
ACGATGCCCTTCCCGGAAACCATCTTCTGAAGCGACCTTTACCCTCCTGCTCACCTTGACCCATCCTTT
AACTGCCCTCCCCTCCTGTCTG

Fig. 13

A**OvIL-32 alpha (SEQ ID NO:24)**

MCFARGVPHDQASLRSM LHTWVDHVCDKMGNEPEEAQMEALAEMEEELSKDVCE
SWKITFKRTFPNPCR
SPVPCFRKRSKKYAAESRDPQSLPVWRTRNRKRASGP EPCGGSEVFCGVSGSGVAMY

B**OvIL-32 alpha (SEQ IDO:25)**

CTGCGGTACCGGTCCGGATTCCCGGGCGAGACAGTGCTCAGCTGGAGCTCTGGCTGCAGGATCTCAGATC
CCAGCCGGAGGACCCTAATCCACCATGTGCTTCGCTAGGGGAGTCCCACATGACCAGGCTTCTCTGAGGA
GCATGCTGCACACCTGGGTGGATCATGTCTGTGATAAGATGGGAAATGAACCAGAAGAAGCACAGATGGA
GGCAGCCCTAGCAGAGATGGAGGAGGAACCTCAGCAAGGATGTCTGTGAATCATGGAAGATCACATTCAAG
AGAACCTTCCCGAATCCCTGCAGGAGTCCAGTCCCTTGCTTCAGGAAGCGCAGCAAGAAGTACGCCGCAG
AATCCAGAGACCCTCAGTCTCTGCCTGTCTGGAGGACCAGAAACCGGAAGAGAGCATCTGGGCCAGAGCC
CTGCGGCGGTTCGGAGGTTTTCTGCGGAGTCTCTGGCAGCGGTGTTGCGATGTACTGACCTGGCTGCAGG
AGAAGGCGGCGGCCTGCCTGGAGGCCGTCTGCAGTGCGGTGAAGACCATCTGGGGAGTGCTGACGGATTT
CTGCTCCTCTGTGGGGCAGCTCTTCAGAAACCTCATCCAGGTCTAGGAGCCCCAGGTCGTTCTTGAGGAA
CTGCTCCTCATCTAGAAGGCCCTGCACAATCCCCCTTCCCAGAAACCATCTTCTGAAGCGACCTTTACCCT
CCTGTTACCCCTTCACCAATCCTTTAACTGCCCTCACCTCCTGTCTGCAGGGACGACACCACAACATCAA
GCCAGGTTTCCCTTCTCCAAGTCTGACCCGTCTGTCAGGGA

C**SwIL-32 alpha (SEQ ID NO:26)**

MRGVSATRTL PKAGPQPRSG LGLPLPRRVPEPPPIPAESSPLLNEVRQGVRSRVRRPPGHNQPHYALAVR
EPRQSTFRRI LELFEEM LKRLQQRWRGALAWVQERAAACFRGLCRALEAFWSLVQSFCSSMGHAFGSVIQ
V

D**SwIL-32 alpha (SEQ ID NO:27)**

ATGACTTGAGGGGAACTGAGCGGCCAGGCCCAGCCCCTGGGAAAAGTCCTGGGGTCTGTGGGGCTGTTG
GCAGGAAAGCAGCCTGTGTCCAAGGCGGGGCATGAGGGGGGTGTCTGCCACCAGGACTCTCCCAAAGGCA
GGGCCTCAGCCAAGGTCAGGACTGGGGCTGCCTCTCCCCAGGCGGGTCCCTGAACCACCCCCCATCCCTG
CAGAATCCAGTCCTCTGCTCAACGAAGTCCGGCAGGGAGTCCGTTCTAGAGTCCGAAGGCCTCCTGGCCA
CAACCAGCCACATTATGCGCTAGCGGTCCGGGAGCCCAGGCAGAGCACTTTCAGACGCATCCTTGAGCTG
TTTGAGGAAATGCTGAAGCGCCTGCAGCAGAGGTGGAGGGGTGCCCTGGCTTGGGTGCAGGAAAGGGCTG
CTGCCTGCTTCCGGGGCTTGTGCAGGGGCCCTTGAAGCTTTCTGGAGCCTGGTGCAGAGTTTTTGCTCCTC
CATGGGGCACGCCTTCGGGAGTGTCATCCAGGTCTAAGGTGCTCCAGGTGAAATAAGAGTTTCTAGAGCA
CAACCTCCCCCTGCCTTGGCTAAAAAGGCAGCTGTAAGCCTTT

Fig. 14