



(86) Date de dépôt PCT/PCT Filing Date: 1999/04/30
(87) Date publication PCT/PCT Publication Date: 1999/11/11
(45) Date de délivrance/Issue Date: 2007/10/09
(85) Entrée phase nationale/National Entry: 2000/10/26
(86) N° demande PCT/PCT Application No.: JP 1999/002305
(87) N° publication PCT/PCT Publication No.: 1999/057249
(30) Priorité/Priority: 1998/04/30 (JP10/137685)

(51) Cl.Int./Int.Cl. *C12N 15/54* (2006.01),
C07K 1/36 (2006.01), *C07K 16/40* (2006.01),
C12N 5/10 (2006.01), *C12N 9/10* (2006.01),
C12P 17/10 (2006.01)

(72) Inventeurs/Inventors:
MORI, SATOSHI, JP;
HIGUCHI, KYOKO, JP;
SUZUKI, KAZUYA, JP;
NISHIZAWA, NAOKO, JP;
NAKANISHI, HIROMI, JP

(73) Propriétaire/Owner:
JAPAN SCIENCE AND TECHNOLOGY
CORPORATION, JP

(74) Agent: RIDOUT & MAYBEE LLP

(54) Titre : NICOTIANAMINE SYNTHASE ET GENE CODANT CETTE DERNIERE
(54) Title: NICOTIANAMINE SYNTHASE AND GENE ENCODING THE SAME

(57) **Abrégé/Abstract:**

A nicotianamine synthase is isolated and purified. Then the gene of this enzyme is cloned and the base sequence and amino acid sequence thereof are determined. This gene is employed in constructing plants, in particular, grass plants highly tolerant to iron deficiency. A nicotianamine synthase involved in the mugineic acid biosynthesis pathway; the amino acid sequence thereof; a gene encoding the same; a vector containing this gene; cells transformed by the vector; a process for producing nicotianamine by using the same; plants transformed by the gene encoding the nicotianamine synthase; and an antibody against the nicotianamine synthase.





PCT

特許協力条約に基づいて公開された国際出願

(51) 国際特許分類6 C12N 9/00, 15/52, C12P 13/04, C07K 16/40	A1	(11) 国際公開番号 WO99/57249 (43) 国際公開日 1999年11月11日(11.11.99)
(21) 国際出願番号 PCT/JP99/02305 (22) 国際出願日 1999年4月30日(30.04.99) (30) 優先権データ 特願平10/137685 1998年4月30日(30.04.98) JP (71) 出願人 (米国を除くすべての指定国について) 科学技術振興事業団(JAPAN SCIENCE AND TECHNOLOGY CORPORATION)[JP/JP] 〒332-0012 埼玉県川口市本町4丁目1番8号 Saitama, (JP) (72) 発明者 ; および (75) 発明者 / 出願人 (米国についてのみ) 森 敏(MORI, Satoshi)[JP/JP] 〒275-0026 千葉県習志野市谷津6-7-2-301 Chiba, (JP) 樋口恭子(HIGUCHI, Kyoko)[JP/JP] 鈴木一矢(SUZUKI, Kazuya)[JP/JP] 中西啓仁(NAKANISHI, Hiromi)[JP/JP] 〒113-0032 東京都文京区弥生1-1-1 Tokyo, (JP) 西澤直子(NISHIZAWA, Naoko)[JP/JP] 〒113-0001 東京都文京区白山1-37-9-705 Tokyo, (JP)		(74) 代理人 弁理士 佐伯憲生(SAEKI, Norio) 〒103-0027 東京都中央区日本橋三丁目15番2号 高愛ビル9階 Tokyo, (JP) (81) 指定国 AU, CA, JP, US, 欧州特許 (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE) 添付公開書類 国際調査報告書
(54)Title: NICOTIANAMINE SYNTHASE AND GENE ENCODING THE SAME (54)発明の名称 ニコチアナミン合成酵素、それをコードする遺伝子 (57) Abstract A nicotianamine synthase is isolated and purified. Then the gene of this enzyme is cloned and the base sequence and amino acid sequence thereof are determined. This gene is employed in constructing plants, in particular, grass plants highly tolerant to iron deficiency. A nicotianamine synthase involved in the mugineic acid biosynthesis pathway; the amino acid sequence thereof; a gene encoding the same; a vector containing this gene; cells transformed by the vector; a process for producing nicotianamine by using the same; plants transformed by the gene encoding the nicotianamine synthase; and an antibody against the nicotianamine synthase.		

NICOTIANAMINE SYNTHASE AND GENE ENCODING THE SAME

Technical Field

The present invention relates to a nicotianamine synthase involved in the mugineic acid biosynthetic pathway, the amino acid sequence thereof, a gene encoding the same, a vector, a process for producing nicotianamine by using the same, plants transformed by the gene encoding the nicotianamine synthase, and an antibody against the nicotianamine synthase.

Background Art

Graminaceous plants that absorb by chelating the insoluble state Fe(III) in soil using mugineic acid and adopt so called the Strategy-II mechanism of Fe acquisition secrete Fe chelators (phytosiderophores) from their roots to solubilize sparingly soluble Fe in the rhizosphere (Roemheld, 1987). The amount of the secreted phytosiderophores increases under Fe-deficiency stress. The mugineic acid family is the only examples of phytosiderophores known so far (Takagi, 1976). Tolerance to Fe deficiency in graminaceous plants is thought to depend on a quantity of mugineic acid family secreted by plants (Takagi et al. 1984, Roemheld and Marschner 1986, Marschner et al. 1987, Mori et al. 1987, Kawai et al. 1988, Mori et al. 1988, Mihashi and Mori 1989, and Shingh et al. 1993).

The biosynthetic pathway of mugineic acid in plants is shown in Fig. 1. S-adenosylmethionine is synthesized from methionine by S-adenosylmethionine synthase. Subsequently, three molecules of S-adenosylmethionine are combined to form one molecule of nicotianamine by nicotianamine synthase. The generated nicotianamine is then converted to 3"-keto acid by nicotianamine aminotransferase,

and 2'-deoxymugineic acid is synthesized by the subsequent action of a reductase. A further series of hydroxylation steps produces the other mugineic acid derivatives including mugineic acid from the deoxymugineic acid (Mori and Nishizawa 1987, Shojima et al. 1989, Shojima et al. 1990 and Ma and Nomoto 1993).

A compound in Fig. 1, a compound in the lower right, wherein R_1 and R_2 are hydrogen and R_3 is hydroxyl, is mugineic acid. A compound wherein R_1 is hydrogen and R_2 and R_3 are hydroxyl, is 3-hydroxymugineic acid. Also a compound wherein R_2 is hydrogen and R_1 and R_3 are hydroxyl, is 3-epihydroxymugineic acid.

Three S-adenosylmethionine synthase genes were isolated from barley roots, but these genes were not induced by Fe deficiency (Takizawa et al. 1996). A gene *Ids3*, which is obtained from the barley by differential screening, is suspected to be a gene, which converts deoxymugineic acid to mugineic acid by hydroxylation and is strongly induced by Fe-deficiency (Nakanishi et al. 1993). Further, nicotianamine aminotransferase was purified and isolated from Fe-deficient barley roots, and two nicotianamine aminotransferase genes, *Naat-A* and *Naat-B*, were isolated (Takahashi et al. 1997). *Naat-A* expression was induced in Fe-deficient roots.

The synthesis of nicotianamine from S-adenosylmethionine is similar to polyamine synthesis from decarboxy-S-adenosylmethionine. In contrast to polyamine synthase, however, nicotianamine synthase catalyzes the combination of three S-adenosylmethionine molecules and the azetidine ring formation at the same time (Fig. 1). Such the nicotianamine synthase is a novel type of enzyme. Previously, we reported the partial purification of nicotianamine synthase from the roots of Fe-deficient barley and expression pattern of the activity (Higuchi et al. 1994, Higuchi et al. 1995, Kanazawa et al. 1995, Higuchi et al. 1996a and Higuchi et al. 1996b). Since nicotianamine synthase is easily decomposed during extraction and purification, it has

been difficult to purify sufficient quantities for amino acid sequencing.

The present invention has an object to provide a plant, especially graminaceous plant, highly tolerant to Fe-deficiency, as a result of isolating and purifying a nicotianamine synthase, being cloned the gene of this enzyme, determining the base sequence and amino acid sequence thereof, and using said enzyme.

Disclosure of Invention

The present invention relates to a nicotianamine synthase shown in SEQ ID NO: 1 comprising amino acid sequence shown in SEQ ID NO: 1, or amino acid sequence having deletion in a part thereof, being substituted by the other amino acids or being added with the other amino acids.

The present invention relates to the gene encoding said amino acid sequence of nicotianamine synthase.

The present invention also relates to a vector comprising containing said gene, and a transformant transformed by the said vector.

The present invention relates to a process for production of nicotianamine using the said transformant.

The present invention further relates to plants, especially graminaceous plants, to which said gene is introduced, and fruits obtained by growing said plants.

The present invention relates to a process for extraction of said nicotianamine synthase in the presence of thiol protease inhibitor, preferably E-64.

Further, the present invention relates to an antibody against said nicotianamine synthase.

Brief Description of Drawing

Fig. 1 shows the biosynthetic pathway of mugineic acid family.

Fig. 2 shows a comparison of nicotianamine synthase purification from Fe-dependent and control barley roots.

Fig. 3 shows a preparative SDS-PAGE (sodium dodecyl sulfate-polyacrylamide gel electrophoresis, hereinafter designates as SDS-PAGE) around 30 - 35 kDa. The horizontal bar indicates relative enzyme activity detected from the gels.

Fig. 4 shows elution pattern of nicotianamine synthase activity from the gel-filtration column.

The large closed circles (●) indicates enzyme activity.

Fig. 5 shows a comparison with a six partial amino acid sequence determined by nicotianamine synthase originated from barley and similar sequence of graminaceous plants obtained by computer search of the database. Identical amino acid residue is shown in " : " .

Fig. 6 shows full length of HvNAS1 cDNA and amino acid sequence deduced therefrom. The underlined sequences indicate the identical partial amino acid sequences of fragments in the above Fig. 5. Numbers of the nucleotide sequence are indicated to the right of each row. Amino acid numbers are indicated on the left of each row.

Fig. 7 shows comparison of the deduced amino acid sequences of the above 7 cDNA obtained from barley. Asterisks " *" indicates identical amino acid residues in all sequences.

Fig. 8 shows results of thin layer chromatographic (TLC) analysis of nicotianamine synthase activity obtained from *E. coli* crude extract expressing a fused protein of maltose binding protein - HvNAS1.

Fig. 9 shows Northern - hybridization analysis of HvNAS1 as a probe.

Fig. 10 shows Southern - hybridization analysis of HvNAS1 as a probe.

Fig. 11 shows Western-blot analysis of crude enzyme used for detection of nicotianamine synthase activity.

Fig. 12 shows Western-blot analysis of total protein extracted by trichloroacetic acid/acetone.

Fig. 13 shows comparison of nicotianamine synthase purification from Fe-deficient barley and control barley after DEAE-SephTMarose FF.

Fig. 14 shows comparison of nicotianamine synthase purification from Fe-deficient barley and control barley after Ether Toyopearl 650MTM.

Fig. 15 shows results of thin layer chromatographic (TLC) analysis of nicotianamine synthase activity obtained from *E. coli* crude extract expressing a fused protein of maltose binding protein - OsNAS1.

Fig. 16 shows Northern - hybridization analysis of OsNAS1 as a probe.

Fig. 17 shows results of thin layer chromatographic (TLC) analysis of nicotianamine synthase activity obtained from *E. coli* crude extract expressing a fused proteins of maltose binding protein - AtNAS1, AtNAS2 or AtNAS3.

Fig. 18 shows results of RT-PCR of total RNA extracted from the aboveground parts and roots of *Arabidopsis thaliana*. Right group indicates positive control.

Best Mode for Carrying Out the Invention

We have tried to isolate nicotianamine synthase (Higuchi et al. Plant & Soil, Vol. 165, p. 173-179, 1994), and since nicotianamine synthase was easily decomposed and was difficult to isolate and purify, we were unable to obtain sufficient amounts of protein to determine its partial amino acid sequence. Subsequently, it was found that a

thiol protease inhibitor E-64 (hereinafter designates as E-64) was very effective in suppressing degradation of nicotianamine synthase (Higuchi et al. Plant & Soil, Vol. 178, p. 171 - 177, 1996 a).

In the present invention, as a result that frozen roots were crushed to a fine powder in liquid N₂ and then rapidly homogenized with buffer containing 0.1 mM thiol protease inhibitor E-64, nicotianamine synthase protein could be isolated and its gene could also be isolated.

Further, the enzyme of the present invention recovered its activity by removal of SDS after SDS-PAGE treatment, but the rate of recovery was very low (Higuchi et al. Plant & Soil, Vol. 165, p. 173-179, 1994). Consequently, degree of purification should be increased up before treatment of SDS-PAGE. Then the column chromatography procedures were further improved.

We have also found that the enzyme of the present invention is relatively hydrophobic and a buffer containing a mild surface active agent CHAPS increased the rate of recovery. Several ion-exchange chromatography carriers were tested, and DEAE-Sepharose FF and DEAE SephacelTM were found to be the most effective. In addition to TSK gel Butyl ToyopearlTM, another hydrophobic chromatography carrier, TSK gel Ether Toyopearl 650M, effectively removed impurities of the 30 - 35 kDa.

The enzyme of the present invention has been reported that it was the peptide of 30 - 35 kDa, the activity of which was recovered by removing SDS after SDS-PAGE treatment, and the activity was detected as a broad molecular weight range of 30 - 35 kDa (refer to Fig. 3). Fig. 3 shows a result of preparative SDS-PAGE in the fractions showing enzyme activity. SDS-PAGE was carried out using 11% acrylamide slab gels. A portion of the gel was stained with Coomassie brilliant blue and the rest of the gel was stained with Cu. The gel containing proteins between 30 - 35 kDa in size was cut

into seven fragments (indicated by the short lines). The thick bars in Fig. 3 indicate relative enzymatic activities detected from each gel fragment.

In order to identify nicotianamine synthase peptide from the proteins having these molecular weights, the peptides, which were contained in the nicotianamine synthase fractions, purified from Fe-deficient and control barley roots were compared using SDS-PAGE. From each barley root 200 g, the present enzyme was purified according to the method described in example 3 hereinbelow.

The enzyme activity of the control was a quarter of the Fe-deficient roots.

The peptide composition of the active enzyme fraction from each purification step of the present enzyme was analyzed and compared by SDS-PAGE, and results are shown in Fig. 2, Fig. 13 and Fig. 14. Fig. 2, Fig. 13 and Fig. 14 show comparison with the active fraction from the purification step of Fe-deficient barley roots 200 g [in the figure, shown with (-)], and the active fraction from the purification step of the control barley roots 200 g [in the figure, shown with (+)]. SDS-PAGE was carried out using 12.5% acrylamide slab gels (Laemmli, Nature Vol. 227, p. 680-685, 1970). Gels were stained with Coomassie brilliant blue. Fig. 2 shows a step before DEAE-Sepharose. The upper row shows enzyme from Fe-deficient barley roots and the lower row shows enzyme from control roots. In each lane, lanes 1, crude extract, 200 μ g of protein; lanes 2, after Butyl Toyopearl 650M, 100 μ g of protein; lanes 3, after hydroxyapatite, 20 μ g of protein; and lanes 4, after Butyl Toyopearl 650M, 15 μ g of protein, are shown.

Fig. 13 shows after DEAE-Sepharose FF, each lane, 25 μ g of protein. Fig. 14 shows after Ether Toyopearl 650M; in which left shows inactive fraction, and right shows active fraction, and 1/25 of each fraction is electrophoresed.

As a result, almost no difference was observed in both Fe-deficient and control roots before DEAE-Sepharose step (refer to Fig. 2). After the DEAE-Sepharose step it

became clear that the 30- and 31-kDa peptides were induced by Fe-deficiency (refer to Fig. 13). After the Ether Toyopearl step, the 31 kDa peptide was eliminated from the active fraction. The 32 and 33 kDa peptides were found to be newly induced by Fe-deficiency (refer to Fig. 14). Activities were detected from the 32 and 33 kDa peptides, but no activity was detected from 30 kDa peptide (refer to Fig. 3).

Molecular weight of the enzyme of the present invention was determined by gel-filtration.

Estimated molecular weight of nicotianamine synthase by gel-filtration was reported to be 40,000 - 50,000 (Higuchi et al. Plant & Soil, Vol. 165, p. 173-179, 1994). But this did not correspond with the value estimated by SDS-PAGE.

In the present study, the buffer containing CHAPS effectively increased the resolution and molecular weight of the present enzyme was estimated to be 35,000 (refer to Fig. 4). this corresponds well to the value estimated by SDS-PAGE.

Fig. 4 shows elution pattern of nicotianamine synthase from the gel-filtration column. The black circles (●) indicate the enzyme activity and the solid line indicates absorption at 280 nm. The active fraction after hydroxyapatite chromatography was applied to a Sephacryl S300HRTM (Pharmacia) column (1.5 cm × 71 cm, 125 ml), equilibrated with developing buffer (50 mM Tris, 1 mM EDTA, 0.1 M KCl, 0.05% CHAPS, 0.1 mM p-APMSF and 3 mM DTT, pH 8.0). Molecular weight markers used were thyroglobulin (Mr 670,000), γ -globulin (Mr 158,000), ovalbumin (Mr 44,000), and myoglobin (Mr 17,000). The linear flow was 10 cm/hour.

Partial amino acid sequence was determined from purified nicotianamine synthase.

The above explained 30 kDa, 32 kDa and 33 kDa peptides were purified from 1 kg of Fe-deficient barley roots by using a method in example 3 hereinbelow. These

were partially degraded using a method in example 4 hereinbelow. Although 32- and 33-kDa peptides could not be completely separated from each other, these might have similar sequence or 32 kDa peptide was presumed to be the degradation product of 33 kDa peptide, and both of them were degraded in together.

The determined partial amino acid sequences indicated that these peptides were very similar in each other (Fig. 5). Further, since the molecular weights of the 33 kDa and 32 kDa (1) fragments had almost unchanged molecular weight as compared with before degradation, this sequence might be N-terminal region of the present enzyme. As a result of computer search of the database, a gene of unknown function having very similar sequence to these sequences was found to exist in Oryza sativa and Alabidopsis thaliana. Especially, EST-cDNA clones D23792 and D24790 of Oryza sativa were very similar with 80.0% identity in a 33-amino acid overlap in the former and 68.4% identity in a 19-amino acid overlap in the latter (Fig. 5).

Fig. 5 shows a comparison with a six partial amino acid sequence determined by nicotianamine synthase originated from barley and similar sequence of graminaceous plants obtained by computer search of the database. Identical amino acid residue is shown in “:”. The part of nucleotide sequences indicated by the arrows was applied for the sequences of primer used in PCR.

Cloning and nucleotide sequences of cDNA clones encoding nicotianamine synthase were performed and determined.

PCR amplification of total cDNA prepared from Fe-deficient barley roots using degenerate primers designed from the partial amino acid sequence obtained from the method explained hereinbefore was performed, but the objective DNA could not amplified. Then the primers having single nucleotide sequence (shown by arrows in Fig. 5) from sequences of Oryza sativa, D23792 and D24790, were synthesized and

PCR amplification was performed. The 205 bp fragment was amplified by PCR using NF and NR primers and the 274 bp fragment was amplified by PCR using IF and IR primers, and these contained the objective sequences. A cDNA library prepared using poly (A) ⁺ RNA from Fe-deficient barley roots was screened and 19 positive clones using the 205 bp fragment probe and 88 positive clones using the 274 fragment bp probe were obtained.

Among the thus obtained clones, the clone designated as HvNAS1, contained a translated region of 985 bp and amino acid sequence deduced therefrom was 328 amino acids residue, with deduced molecular weight of 35,144. This corresponded well with the value estimated by SDS-PAGE and gel-filtration. The partial amino acid sequences of the 32 kDa and 33 kDa peptides were included totally in HvNAS1 (Fig. 6).

Fig. 6 shows full length of HvNAS1 cDNA and amino acid sequence deduced therefrom. The underlined sequences indicate the identical partial amino acid sequences of fragments in the above Fig. 5. Numbers of the nucleotide sequence are indicated to the right of each row. Amino acid numbers are indicated on the left of each row.

The predicted pI of 5.2 matched the value estimated by native isoelectric focusing electrophoresis well. The six clones having very similar sequence other than HvNAS1, i.e. HvNAS2, HvNAS3, HvNAS4, HvNAS5, HvNAS6 and HvNAS7, were also obtained (Table 1, Fig. 7).

Fig. 7 shows comparison of the deduced amino acid sequences of the above 7 cDNA obtained from barley. Asterisks "*" indicates identical amino acid residues in all sequences.

The nucleotide sequences of these clones are shown in SEQ ID NO: 2 (HvNAS1), SEQ ID NO: 4 (HvNAS2), SEQ ID NO: 6 (HvNAS3), SEQ ID NO: 8

(HvNAS4), SEQ ID NO: 10 (HvNAS5), SEQ ID NO: 12 (HvNAS6) and SEQ ID NO: 14 (HvNAS7), respectively. The amino acid sequences of these amino acid sequences are shown in SEQ ID NO: 1 (HvNAS1), SEQ ID NO: 3 (HvNAS2), SEQ ID NO: 5 (HvNAS3), SEQ ID NO: 7 (HvNAS4), SEQ ID NO: 9 (HvNAS5), SEQ ID NO: 11 (HvNAS6) and SEQ ID NO: 13 (HvNAS7), respectively.

Table 1 Properties of nas clones

Clone	Number of Amino Acid Residues	Molecular Weight	pI	Identity to nas 1 (%)	Identity to nas 2 (%)	Identity to nas 4 (%)
HvNAS 1	328	35144	5.20	-		
HvNAS 2	336	35839	5.07	72	-	
HvNAS 3	336	36013	5.47	72	95	
HvNAS 4	330	35396	4.91	73	89	-
HvNAS 5	283	30148	5.22	61	61	59
HvNAS 6	329	35350	5.07	74	89	88
HvNAS 7	330	35244	4.98	70	86	91

The partial amino acid sequences determined from the 30 kDa peptide were all included in HvNAS5. The 5'- and 3'-non-translated regions of these clones were not similar with each other.

D23792 and D24790 similar to nicotianamine synthase of *Oryzae sativa* were found with about 80% identity to HvNAS1. AC003114 and AB005245 of *Arabidopsis thaliana* were found with about 45% identity to HvNAS1.

The obtained HvNAS1 protein was expressed in *E. coli*.

The PCR amplification of HvNAS1 ORF was cloned with vector pMAL-c2 to express HvNAS1 fused with C-terminal of maltose binding protein. The expression of fused protein is strongly induced by IPTG.

The crude extract was obtained from the transformed *E. coli*, and nicotianamine synthase activity was assayed in the state of the fused protein. The crude extract from the strain transformed with only the vector could not be detected the activity, whereas in case of inserted with HvNAS1 ORF, the activity was detected. Result is shown in Fig. 8.

Fig. 8 shows results of thin layer chromatographic (TLC) analysis of nicotianamine synthase obtained from *E. coli* crude extract expressing a fused protein of maltose binding protein - HvNAS1. In Fig. 8, lane 1: a standard nicotianamine synthase; lane 2: *E. coli* expressing maltose binding protein (SAM); and lane 3: *E. coli* expressing maltose binding protein - HvNAS1 fused protein.

Northern hybridization analysis conducted by the method described in example 7 hereinbelow indicated that this gene was strongly induced in Fe-deficient roots (Fig. 9). This coincides with expression pattern of the present enzyme activity (Higuchi et al. 1994). Fig. 9 shows a result of Northern hybridization analysis using HvNAS1 as a probe. Total RNA was extracted from after one week of Fe-deficient treatment and control barley leaves and roots, and in each lane, 5 μ g of RNA were electrophoresed.

Southern hybridization analysis of the barley genome DNA was performed according to the method described in example 8 hereinafter mentioned. Cutting of DNA with BamHI, EcoRI or HindIII produced plurality of fragments, however none of clones obtained at present could be digested by BamHI and EcoRI, consequently nicotianamine synthase gene might exist with multiple copies in genomes of barley and rice (Fig. 10).

Fig. 10 shows Southern - hybridization analysis of HvNAS1 as a probe. Genomic DNAs from barley and rice were digested with BamHI (lanes B), EcoRI (lanes

R) and HindIII (lanes H) and 10 μ g thereof were electrophoresed in each lane.

Further, using antigen prepared by the method described in example 9 hereinbelow, Western-blot analysis was performed according to the method described in example 10. It was found that the present enzyme protein was rapidly decomposed during the operation in the crude extract prepared for detecting the present enzyme activity (Fig. 11). The staining patterns coincided with the fact that the present enzyme activity was detected on the broad range between 30 - 35 kDa after SDS-PAGE (refer to Fig. 3).

Fig. 11 shows Western-blot analysis of crude enzyme used for detection of activity. SDS-PAGE was performed using 12.5% acrylamide slab gel. Protein 100 μ g was electrophoresed.

The crude extract obtained from denatured protein according to the method described in example 10 hereinbelow was detected as almost single band with 35 - 36 kDa (Fig. 12). This value coincided with the deduced value from the amino acid sequence.

Fig. 12 shows Western-blot analysis of total protein extracted by trichloroacetic acid/acetone. SDS-PAGE was performed using 12.5% acrylamide slab gel. Protein 100 μ g was electrophoresed. Proteins 200 μ g extracted from roots and proteins 500 μ g extracted from leaves were electrophoresed.

Western-blot analysis after 2-dimension electrophoresis reveals to detect several spots. This coincided with the fact of obtaining plurality of nicotianamine synthase gene. All spots were induced in Fe-deficient roots.

As a result that cDNA library from Fe-deficient rice roots poly (A) + RNA was screened using probes prepared by cutting HvNAS1 with restriction enzymes ApaLI and XhoI, 20 clones were obtained. These clones were divided into 3 types of clones

according to their sequences, and among them, only one type contains ORF full length, which was designated as OsNAS1. Nucleotide sequence of OsNAS1 is shown in SEQ ID NO: 16 and amino acid sequence is shown in SEQ ID NO: 15.

PCR amplification of OsNAS1 ORF was cloned with a vector pMAL-c2 to express a form fused with maltose binding protein C-terminal. The fused protein is strongly induced its expression by IPTG.

Crude extract from the transformed *E. coli* with the fused protein was obtained and nicotianamine synthase activity was assayed in the state of the fused protein. The same activity with HvNAS1 was detected. Result is shown in Fig. 15. Fig. 15 shows results of thin layer chromatographic (TLC) analysis of nicotianamine synthase obtained from *E. coli* crude extract expressing a fused protein of maltose binding protein - OsNAS1. In Fig. 15, lane 1: a standard nicotianamine (NA); lane 2: an extract from *E. coli* expressing maltose binding protein - OsNAS1 fused protein; and lane 3: an extract from *E. coli* expressing maltose binding protein - HvNAS1 fused protein.

Northern hybridization analysis conducted by the method described in example 7 hereinbelow indicated that in contrast to barley, the expression was induced in rice by Fe-deficient treatment not only in roots but also in leaves (Fig. 16). Fig. 16 shows a result of Northern hybridization analysis using OsNAS1 ORF as a probe. Total RNA was extracted from after two weeks of Fe-deficient treatment and control rice leaves and roots, and in each lane, 5 μ g of RNA were electrophoresed.

Nucleotide sequence of *Arabidopsis thaliana* similar to HvNAS1 obtained by computer search of the database was used as a primer. PCR amplification for genome DNA of *Arabidopsis thaliana* resulted to obtain three nicotianamine synthase genes. These were designated as AtNAS1, AtNAS2 and AtNAS3.

Nucleotide sequence of these genes are shown in SEQ ID NO: 18 (AtNAS1), SEQ ID NO: 20 (AtNAS2) and SEQ ID NO: 22 (AtNAS3). These amino acid sequences are shown in SEQ ID NO: 17 (AtNAS1), SEQ ID NO: 19 (AtNAS2) and SEQ ID NO: 21 (AtNAS3).

AtNAS1, AtNAS2 and AtNAS3 ORF were amplified with PCR and were cloned with a vector pMAL-c2. Each of them was tried to be expressed in the form of fusing with maltose binding protein C-terminal. The expression of the fused protein was strongly induced by IPTG.

Crude extract from the transformed *E. coli* with the fused protein was obtained and nicotianamine synthase activity was assayed in the state of the fused protein. The activity was detected. Result is shown in Fig. 17. Fig. 17 shows results of TLC analysis of nicotianamine synthase activity obtained from *E. coli* crude extract expressing a fused protein of maltose binding protein - AtNAS. In Fig. 17, lanes 1: a standard nicotianamine (NA) and S-adenosylmethionine; lanes 2: an extract from *E. coli* expressing only maltose binding protein; lanes 3: an extract from *E. coli* expressing maltose binding protein - AtNAS1 fused protein; lanes 4: an extract from *E. coli* expressing maltose binding protein - AtNAS2 fused protein; and lanes 5: an extract from *E. coli* expressing maltose binding protein - AtNAS3 fused protein.

RT-PCR was conducted according to the method described in example 11 hereinbelow. It was found that AtNAS1 was expressed in the roots and the aboveground parts of *Arabidopsis thaliana*, whereas AtNAS2 was expressed neither in the roots nor in the aboveground parts, and AtNAS3 was expressed only in the roots (Fig. 18). In Fig. 18, lane M shows molecular weight marker. Gene expression was conducted in the aboveground parts, roots and positive controls. In the figure, lanes C: AtNAS1 and AtNAS2 ORF full length were amplified; lanes 1: AtNAS1 specific

amplification fragments; lanes 2: AtNAS2 specific amplification fragments; and lanes 3: AtNAS3 specific amplification fragments.

The amount of secreted mugineic acid is reported increased up to 20 mg mugineic acid/g roots dry weight/day (Takagi, 1993). Crude nicotianamine synthase activity detected by the present inventors was sufficient to fulfill it. Since the present enzyme proteins exist in more than several types and 30 kDa peptide without activity exists, it can be speculated that as a result of aggregation of these peptides, the constructed structure, which is preferable for binding with 3 molecules of S-adenosylmethionine, reveals maximum activity. The molecular weight estimated by gel-filtration was 35,000 (Fig. 4).

Increase in activity by re-aggregation of subunits has not been observed at present. Since the fused protein with maltose binding protein and subunits showed its activity, we have at present an idea that the present enzyme might be a monomer. However, the possibility that large activity can be revealed by constructing multimer, can not completely denied.

The reaction mechanism synthesizing nicotianamine from S-adenosylmethionine may be similar to methyl transfer reaction using S-adenosylmethionine as a methyl donor, and a reaction synthesizing spermidine and spermine from decarboxylated S-adenosylmethionine. The common catalytic domain of these enzymes has been discussed in relation to equivalent amino acids configuration occupying similar positions in higher-order structures (Hashimoto et al. 1998 and Schluckebier et al. 1995).

In future, catalytic domain may be elucidated as the results of comparison with nicotianamine synthase from other plant species or X-ray crystallography.

Induction of nicotianamine synthase activity by Fe-deficiency is a specific

phenomenon in graminaceous plants, and is essential for mass production of mugineic acid family. Oryza sativa is a plant, in which secretion of mugineic acid family is the least among major graminaceous plants, consequently it is very weak for Fe-deficiency in calcareous soil.

Consequently, as a result of creating transformant Oryza sativa having tolerance to Fe-deficiency by introducing nicotianamine synthase gene of the present invention into the graminaceous plants, especially Oryza sativa, and expressing large amount at the Fe-deficiency, cultivation of rice in the calcareous soil can be possible.

Heretofore, in the graminaceous plants, nicotianamine has been thought to have only a role as a precursor for synthesis of mugineic acid family. However, since the present invention has elucidated that nicotianamine synthase gene constituted the multiple gene family, it may play other important roles in the graminaceous plants.

In plants, which lack the ability to secrete mugineic acid family, except for graminaceous plants, it has been proposed that nicotianamine plays a key role as an endogenous chelator of divalent metal cations, such as Fe^{2+} , Cu^{2+} , Zn^{2+} and Mn^{2+} , and that it contributes to the homeostasis of those metals (Stephan et al. 1994). Consequently, it may play the same role in the graminaceous plants.

Nicotianamine synthase activity is not induced in dicots, and expression of gene of the present invention may not be induced by Fe-deficiency. We have cloned nicotianamine synthase genes of Arabidopsis thaliana. Composition of promoter regions in these genes can elucidate the mechanism of gene expression caused by Fe-deficiency, and the gene of the present invention may play important function not only in the graminaceous plants but also in the dicots.

SEQ ID NO: 1 shows amino acid sequence of nicotianamine synthase of the present invention.

The present invention includes nicotianamine synthase having amino acid sequence shown in SEQ ID NO: 1. However, the present invention is not limited within the above nicotianamine synthase. The nicotianamine synthase of the present invention includes, unless it loses nicotianamine synthase activity, the peptides, in which a part of the amino acid sequence of said peptide is deleted, preferably 50% or less, more preferably 30% or less, or more further preferably 10% or less in the total amino acids, or is substituted by other amino acids, or to which other amino acids are further added, or in which these deletion, substitution and addition may be combined.

Nucleotide sequence coding nicotianamine synthase of the present invention is shown in SEQ ID NO: 2.

The present invention also includes not only a gene coding nicotianamine synthase shown in SEQ ID NO: 2 but also genes coding nicotianamine synthase mentioned hereinabove.

The vector of the present invention introducing the above gene is not specifically limited, and various vectors can be introduced. Preferable vector is the expression vector.

Various cells can be transformed conventionally by using recombinant vector of the present invention. Mass production of nicotianamide can be performed by using the thus obtained transformant. These methods are well known in the person skilled in the art.

Examples of hosts for introducing the gene of the present invention are bacteria, yeasts and cells. Preferable host is plants, especially the graminaceous plant.

Method for introducing gene is not limited. It can be made by using vector or can be directly introduce in genome.

Antibody of the present invention against nicotianamine synthase can be

prepared conventionally by using nicotianamine synthase of the present invention. Antibody can be a polyclonal antibody or, if necessary, monoclonal antibody.

Further, a selective breeding of plants, preferably graminaceous plants, can be made by using gene of the present invention. Especially, the gene of the present invention can be applied for improvement of varieties, which can grow even in Fe-deficient soil.

Examples

The following examples illustrate the present invention, but are not construed as limiting the present invention.

Example 1. (Preparation of plant material)

Seeds of barley (*Hordeum vulgare* L. cv Ehimehadakamugi No. 1) were germinated on wet filter paper and transferred into the standard hydroponic culture solution (Mori and Nishizawa, 1987) in a glass house at natural temperature under natural light. The pH of the hydroponic culture solution was adjusted at 5.5 by 0.5 N HCl everyday. When the third leaves developed, the plants were transferred to the hydroponic culture solution without containing Fe. The pH was maintained at 7.0 by 0.5 N NaOH everyday. The control plants were also cultured in the standard culture solution continuously. The culture solution was renewed once in every week. Two weeks after starting Fe-deficient treatment, when severe iron chlorosis significantly appeared on the 4th and 5th leaves, roots were harvested and frozen in liquid N₂ and stored at -80°C until use.

Example 2. (Assay of nicotianamine synthase activity)

Modified assay method reported previously by the present inventors (Higuchi et al. 1996a) was used. Enzyme solutions were equilibrated with reaction buffer [50 mM Tris, 1 mM EDTA, 3 mM dithiothreitol (hereinafter designates as DTT), 10 μ M (p-amidinophenyl) methanesulfonyl fluoride (hereinafter designates as p-APMSF) and 10 μ M trans-epoxysuccinyl-leucylamido-(4-guanidino) butane (hereinafter designates as E-64), pH 8.7]. Buffer exchange was performed by using ultrafiltration unit, Ultrafree C3LGC NMWL10000 (Millipore Co.). S-adenosylmethionine labeled with ^{14} C in carboxyl group (Amersham Inc.) was added to the enzyme solution at the final concentration of 20 μ M and kept at 25°C for 15 minutes. The reaction products were separated by thin layer chromatography on silica gel LK6 (Whatman Inc.) using developer (phenol : butanol : formic acid : water = 12 : 3 : 2 : 3). Radioactivity of the reaction products was detected by image Analyzer BAS-2000 (Fuji Film Co.). The protein content was assayed by Bradford method using Protein Assay Kit (Bio Rad Inc.).

Example 3 (Purification of nicotianamine synthase)

The following operations were performed at 4°C and E-64 was added to fractions containing nicotianamine synthase at the final concentration of 10 μ M.

The frozen roots were crushed into a fine powder in liquid N₂ and homogenized in a household juicer with 200 ml of extraction buffer [0.2 M Tris, 10 mM EDTA, 5% (v/v) glycerol, 10 mM DTT, 0.1 mM E-64, 0.1 mM p-APMSF and 5% (w/v) insoluble polyvinylpyrrolidone (PVP), pH 8.0] per 100 g of roots. The homogenate was centrifuged for 30 minutes at 22,500 $\times$ g to obtain supernatant. Ammonium sulfate was added to the supernatant to yield a final concentration of 0.4 M and allowed to stand for 1 hour. Again, the mixture was centrifuged for 30 minutes at 22,500 $\times$ g to

obtain supernatant.

The supernatant was loaded onto a TSK gel Butyl Toyopearl 650M column (10 ml bed volume per 100 g of roots), equilibrated with the adsorption buffer [20 mM Tris, 1 mM EDTA, 3 mM DTT, 0.4 M $(\text{NH}_4)_2\text{SO}_4$ and 0.1 mM p-APMSF, pH 8.0] and eluted with elution buffer [10 mM Tris, 1mM EDTA, 3 mM DTT, 0.1 mM p-APMSF, 5% glycerol and 0.05% 3-[(3-chloramidopropyl) dimethyl-ammonio] propanesulfonic acid (hereinafter designates as CHAPS), pH 8.0].

KCl was added to the active fraction to give a final concentration of 0.4 M, and 1 M potassium phosphate buffer (pH 8.0) was added to a final concentration of 1 mM of KCl. A hydroxyapatite 100 - 350 mesh (Nacalai Tesque), equilibrated with the adsorption buffer (1 mM K-P, 10 mM KCl, 3 mM DTT and 0.1 mM p-APMSF, pH 8.0), was prepared at 10 ml per protein 100 mg and the fractions containing nicotianamine synthase were loaded. Nicotianamine synthase was passed through without adsorption. The passed through fraction was loaded onto TSK gel Butyl Toyopearl 650M column (1 ml bed volume per 10 mg of protein), and nicotianamine synthase was eluted in the manner described above.

The active fraction was loaded onto a DEAE-Sepharose FF column (5 ml bed volume per 25 mg of protein, Pharmacia) equilibrated with the adsorption buffer (20 mM Tris, 1 mM EDTA, 3 mM DTT, 0.1 mM p-APMSF and 0.05% CHAPS, pH 8.0) and eluted with stepwise gradient elution of potassium chloride concentration of 0.05 M, 0.1 M, 0.15 M and 0.2 M. Nicotianamine synthase was eluted at 0.15 M of KCl concentration.

The active fraction was loaded onto the Ether Toyopearl 650M column (10 ml bed volume per 100g of roots), equilibrated with adsorption buffer [20 mM Tris, 1 mM EDTA, 3 mM DTT, 1.2 M $(\text{NH}_4)_2\text{SO}_4$ and 0.1 mM p-APMSF, pH 8.0]. Nicotianamine

synthase was not adsorbed and passed through from the column. The passed through fraction was loaded onto TSK gel Butyl Toyopearl 650M column and fractions containing nicotianamine synthase was eluted. The peptides in the active fraction containing nicotianamine synthase, which was purified by the above column chromatographic treatments, were separated by sodium dodecyl sulfate - polyacrylamide gel electrophoresis (hereinafter designates as SDS-PAGE) using 11% acrylamide slab gels. After SDS-PAGE the gel was stained with 0.3 M copper chloride (Dzandu et al. 1988), and the separated bands were cut out. The gel fragments were destained with 0.25 M EDTA/0.25 M Tris (pH 9.0) and homogenized with the extraction buffer (1% SDS, 25 mM Tris and 192 mM glycine). Each homogenate was electroeluted with SDS-free buffer (25 mM Tris and 192 mM glycine) and peptide was recovered.

Example 4. (Determination of partial amino acid sequence)

The isolated nicotianamine synthase was digested chemically with cyanogen bromide (Gross 1967).

After SDS-PAGE treatment, 10-fold volume of 70% (v/v) formic acid and 1% (w/v) cyanogen bromide were added to gel fragments containing nicotianamine synthase and decomposed at 4°C for overnight. After completion of digestion, the liquid part was collected and dried in vacuo. The dried substance was dissolved in SDS-PAGE sample buffer, and allowed to stand at room temperature for overnight, then the digested product was separated by SDS-PAGE using 16.5% acrylamide gel containing Tricine (Schagger and Jagow, 1987). The peptides were transferred onto a PVDF membrane by electroblotting (Towbin et al. 1979) and stained with amido black. The stained bands were cut out and the amino acid sequence was determined from N-

terminal side of each peptide by Edman degradation in gas-phase sequencer (model 492A protein sequencer, Applied Biosystems Inc.).

Example 5. (Cloning of nicotianamine synthase genes)

PCR amplification was conducted for cDNA originated from Fe-deficient barley roots using primers, which were synthesized based on the obtained partial amino acid sequence. A pYH23 cDNA library prepared from the poly (A)⁺RNA of Fe-deficient barley roots was screened with the thus obtained DNA fragments of PCR product, which was labeled with [α -³²P]dATP using the random primer kit (Takara Shuzo Co.), as the primers. The isolated cDNA clones were sequenced by cycle sequencing kit (Shimadzu Bunko Co.) using Shimadzu DNA sequencer DSQ-2000LTM.

PCR amplification was conducted for genomic DNA of *Arabidopsis thaliana* using primers, which were synthesized based on nucleotide sequences of AC003114 and AB005245 of *Arabidopsis thaliana*. The thus obtained DNA fragments were sequenced by cycle sequencing kit (Shimadzu Bunko Co.) using Shimadzu DNA sequencer DSQ-1000L.

The determined nucleotide sequence is shown in SEQ ID NO: 2.

Example 6. (Expression of NAS1 protein in *E. coli*)

A fragment, in which EcoRI site was introduced into the upstream of the first ATG of the HvNAS1 cDNA and PstI and BamHI sites were introduced into the downstream of the stop codon of the HvNAS1 cDNA, was amplified by PCR. The thus obtained amplified product was subcloned in the pBluescriptII SK- using EcoRI site and BamHI site, and the correct nucleotide sequence was confirmed. The fragment between EcoRI site and PstI site was cloned into pMAL-c2 to make expression in the

form of fusing the HvNAS1 to the C-terminal of maltose binding protein.

A fragment, in which EcoRI site was introduced into the upstream of the first ATG of the OsNAS1 and HindIII site was introduced into the downstream of the stop codon of the OsNAS1, was amplified by PCR. The thus obtained amplified product was subcloned in the pBluescriptII SK- using EcoRI site and HindIII site, and the correct nucleotide sequence was confirmed. The fragment between EcoRI site and HindIII site was cloned into pMAL-c2 to make expression in the form of fusing the OsNAS1 to the C-terminal of maltose binding protein.

A fragment, in which EcoRI site was introduced into the upstream of the first ATG of the AtNAS1, AtNAS2 and AtNAS3 and XbaI site was introduced into the downstream of the stop codon of the AtNAS1, AtNAS2 and AtNAS3, was amplified by PCR. The thus obtained amplified products were subcloned in the pBluescriptII SK-, and the correct nucleotide sequences were confirmed. The fragment between EcoRI site and XbaI site was cloned into pMAL-c2 to make expression in the form of fusing the AtNAS1, AtNAS2 and AtNAS3 to the C-terminal of maltose binding proteins, respectively.

E. coli strain XL1-Blue was used as a host for expressing the said fused protein. pMAL-c2-HvNAS1 and pMAL-c2, respectively, were introduced into XL1-Blue. The thus obtained recombinant bacteria were cultured in LB medium containing ampicillin and tetracycline, each 50 μ g/ml, at 37°C until the OD 600 of the culture reached 0.5. Isopropyl β -D-thiogalactopyranoside (IPTG) was added to the final concentration of 0.3 mM, and continuously cultured at 37°C for 3 hours, and collected bacterial cells. Cells were suspended in 10 mM Tris buffer containing 0.2 M NaCl, 1 mM EDTA, 3 mM DTT and 0.1 mM E-64, pH 7.4 and frozen with liquid nitrogen. This was melted in ice water and ultrasonication for 15 seconds was repeated for 10 times.

Nicotianamine synthase activity of the thus obtained crude extract was assayed according to the method described in example 2 and the enzyme activity was confirmed.

Example 7. (Northern hybridization)

Northern hybridization of barley RNA was performed using DNA fragment, which was prepared by cutting HvNAS1 cDNA with HindIII and NotI and labeled with [α -³²P]dATP, as a probe. Total RNA was extracted from barley (Naito et al. 1988). The extracted RNA was separated by 1.4% agarose gel electrophoresis, and blotted onto Hybond-NTM membranes (Amersham). Northern hybridization of rice RNA was performed using OsNAS1 ORF, which was labeled with [α -³²P]dATP, as a probe. Total RNA was extracted from rice. The extracted RNA was separated by 1.4% agarose gel electrophoresis, and blotted onto Hybond-N⁺ membranes (Amersham). The membrane was hybridized with the probe in 0.5 M Church phosphate buffer (Church and Gilbert 1984), 1 mM EDTA, 7% (w/v) SDS with 100 μ g/ml salmon sperm DNA at 65°C for overnight. The membrane was washed with buffer containing 40 mM Church phosphate buffer and 1% (w/v) SDS at 65°C for 10 minutes. After the washing was repeated once again, the membrane was washed with buffer containing 0.2 $\times$ SSPE and 0.1% (w/v) SDS at 65°C for 10 minutes. Radioactivity was detected using the image analyzer BAS-2000.

Results are shown in Fig. 9 and Fig. 16.

Example 8. (Southern hybridization)

Genomic DNA was extracted from leaves of barley and rice. The extract was digested with BamHI, EcoRI or HindIII, separated on a 0.8% (w/v) agarose gel

electrophoresis, and transferred onto Hybond-N⁺ membranes (Amersham). The hybridization was performed according to the method described in example 7 and radioactivity was detected.

Result is shown in Figure 10.

Example 9 (Preparation of polyclonal antibody)

Two rats were immunized using the antigen containing about 100 μ g of isolated nicotianamine synthase. The antigen was the same sample as that determined the partial amino acid sequence. The complete Freund's adjuvant was used at the first immunization and the incomplete Freund's adjuvant was used since the second immunization. All the constituents of the blood were corrected after the rats were immunized four times, and the obtained serum was preserved at -80°C.

Example 10 (Western blotting analysis)

Total protein was extracted using trichloroacetic acid and acetone (Damerval et al. 1986). The plants were crashed in the liquid nitrogen until powder was obtained, and mixed with acetone containing 0.1% (v/v) 2-mercaptoethanol. The protein was precipitated by allowing to stand at -20°C for 1 hour, and the precipitate was collected by centrifugation at 16,000 X g for 30 minutes. The precipitate was suspended in acetone containing 0.1% (v/v) 2-mercaptoethanol and allowed to stand at -20°C for 1 hour, then collected the precipitate by centrifugation at 16,000 X g for 30 minutes. The precipitate was dried in vacuo, and dissolved in the sample buffer [9.5 M urea, 2% (w/v) Triton X-100 and 5% (v/v) 2-ME], then centrifuged at 16,000 X g for 10 minutes to obtain the supernatant. The proteins contained in the supernatant were separated by SDS-PAGE or the denaturing two-dimensional electrophoresis (O'Farrell 1975) and transferred onto PVDF membrane. Western blotting analysis was performed by applying the primary antibody containing anti-nicotianamine synthase antibody prepared in example 9 and the secondary antibody containing horse radish binding anti-mouse IgG (H + L) goat antibody (Wako Pure Chemicals Co.) on the membrane and coloring with diaminobenzidin.

Result is shown in Fig. 12. SDS-PAGE was performed using 12.5% acrylamide slab gel. Protein 100 μ g was electrophoresed. Proteins of roots 200 μ g and leaves 500 μ g were electrophoresed.

Industrial Applicability

Various cells are transformed according to the conventional method by using recombinant vectors of the present invention. Mass production of nicotianamine can be performed by using the obtained transformant. These methods can be performed according to the method known in the person skilled in the art.

Selective breeding of plants, preferably graminaceous plants can also be performed using genes of the present invention. Especially, genes of the present invention can be applied for improving varieties, which can grow on Fe-deficient soil.

F **F**

Val Gly Ala Arg Met Ser Phe His Thr Ala Asp Val Ala Asp Leu Thr
180 185 190

Gln Glu Leu Gly Ala Tyr Asp Val Val Phe Leu Ala Ala Leu Val Gly
195 200 205

Met Ala Ala Glu Glu Lys Ala Lys Val Ile Ala His Leu Gly Ala His
210 215 220

Met Val Glu Gly Ala Ser Leu Val Val Arg Ser Ala Arg Pro Arg Gly
225 230 235 240

Phe Leu Tyr Pro Ile Val Asp Pro Glu Asp Ile Arg Arg Gly Gly Phe
245 250 255

Glu Val Leu Ala Val His His Pro Glu Gly Glu Val Ile Asn Ser Val
260 265 270

Ile Val Ala Arg Lys Ala Val Glu Ala Gln Leu Ser Gly Pro Gln Asn
275 280 285

Gly Asp Ala His Ala Arg Gly Ala Val Pro Leu Val Ser Pro Pro Cys
290 295 300

Asn Phe Ser Thr Lys Met Glu Ala Ser Ala Leu Glu Lys Ser Glu Glu
305 310 315 320

Leu Thr Ala Lys Glu Leu Ala Phe
325

<210> 2
<211> 1295
<212> DNA
<213> Hordeum vulgare

<400> 2

gcgttcagag	gcttccagag	ttcttccggt	caccaagaag	catttgatca	taacatggat	60
gcccagaaca	aggaggtcgc	tgctctgata	gagaagatcg	ccggtatcca	ggccgccatc	120
gccgagctgc	cgctcgctgag	cccgtccccc	gaggtcgaca	ggctcttcac	cgacctcgtc	180
acggcctgcg	tcccgccgag	ccccgtcgac	gtgacgaagc	tcagcccgga	gcaccagagg	240
atgcgggagg	ctctcatccg	cttgtgctcc	gccgccgagg	ggaagctcga	ggcgcaactac	300
gccgacctgc	tcgccacctt	cgacaacccg	ctcgaccacc	tcggcctctt	cccgtactac	360
agcaactacg	tcaacctcag	caggctggag	tacgagctcc	tggcgcgcca	cgtgccgggc	420
atcgcgccgg	cgcgcgtcgc	cttcgtcggc	tccggcccgc	tgccgttcag	ctcgctcgtc	480
ctcgccgcgc	accacctgcc	cgagaccag	ttcgacaact	acgacctgtg	cggcgcggcc	540
aacgagcgcg	ccaggaagct	gttcggcgcg	acggcggacg	gcgtcggcgc	gcgtatgtcg	600
ttccacacgg	cggacgtcgc	cgacctcacc	caggagctcg	gcgcctacga	cgtgggtcttc	660
ctcgccgcgc	tcgtcggcat	ggcagccgag	gagaaggcca	aggtgattgc	ccacctgggc	720
gcgcacatgg	tggagggggc	gtccctggtc	gtgcggagcg	cacggcccgc	cggctttctt	780
tacccattg	tcgacccgga	ggacatcagg	cgggggtggg	tcgaggtgct	ggccgtgcac	840
cacccggaag	gtgaggtgat	caactctgtc	atcgtcgccc	gtaaggccgt	cgaagcgcag	900
ctcagtgggc	cgcagaacgg	agacgcgcac	gcacggggcg	cggtgccggt	ggtcagcccg	960
ccatgcaact	tctccaccaa	gatggaggcg	agcgcgcttg	agaagagcga	ggagctgacc	1020
gccaaagagc	tggccttttg	attgaagagt	gcgcgtggtc	attctgtcgc	ctgcgatcgt	1080
ggtaactttc	ctactcgtgt	gtgttttgat	gtttgtgcct	gtaagagtta	tgcttccggc	1140
cttgtgctgt	taatttacac	gcgttacatg	tagtacttgt	atttatacct	ggaataacgg	1200
tatgtaacat	aaatattagt	gggatttgaa	gtgtaatgct	aaataataag	aaaacttgat	1260
gcagacattc	aaaaaaaaaa	aaaaaaaaaa	aaaaa			1295

Met 1	Ala	Ala	Gln	Asn 5	Asn	Gln	Glu	Val	Asp 10	Ala	Leu	Val	Glu	Lys 15	Ile
Thr	Gly	Leu	His 20	Ala	Ala	Ile	Ala	Lys 25	Leu	Pro	Ser	Leu	Ser 30	Pro	Ser
Pro	Asp	Val 35	Asp	Ala	Leu	Phe	Thr 40	Glu	Leu	Val	Thr	Ala 45	Cys	Val	Pro
Pro	Ser 50	Pro	Val	Asp	Val	Thr 55	Lys	Leu	Gly	Pro	Glu 60	Ala	Gln	Glu	Met
Arg 65	Glu	Gly	Leu	Ile	Arg 70	Leu	Cys	Ser	Glu	Ala 75	Glu	Gly	Lys	Leu	Glu 80
Ala	His	Tyr	Ser	Asp 85	Met	Leu	Ala	Ala	Phe 90	Asp	Lys	Pro	Leu	Asp 95	His
Leu	Gly	Met	Phe 100	Pro	Tyr	Tyr	Asn	Asn 105	Tyr	Ile	Asn	Leu	Ser 110	Lys	Leu
Glu	Tyr	Glu 115	Leu	Leu	Ala	Arg	Tyr 120	Val	Pro	Gly	Gly	Tyr 125	Arg	Pro	Ala
Arg	Val 130	Ala	Phe	Ile	Gly	Ser 135	Gly	Pro	Leu	Pro	Phe 140	Ser	Ser	Phe	Val
Leu 145	Ala	Ala	Arg	His	Leu 150	Pro	Asp	Thr	Met	Phe 155	Asp	Asn	Tyr	Asp	Leu 160
Cys	Gly	Ala	Ala	Asn 165	Asp	Arg	Ala	Ser	Lys 170	Leu	Phe	Arg	Ala	Asp 175	Arg
Asp	Val	Gly	Ala 180	Arg	Met	Ser	Phe	His 185	Thr	Ala	Asp	Val	Ala 190	Asp	Leu
Ala	Gly	Glu 195	Leu	Ala	Lys	Tyr	Asp 200	Val	Val	Phe	Leu	Ala 205	Ala	Leu	Val
Gly 210	Met	Ala	Ala	Glu	Asp	Lys 215	Ala	Lys	Val	Ile	Ala 220	His	Leu	Gly	Ala
His 225	Met	Ala	Asp	Gly	Ala 230	Ala	Leu	Val	Val	Arg 235	Ser	Ala	His	Gly	Ala 240
Arg	Gly	Phe	Leu	Tyr 245	Pro	Ile	Val	Asp	Pro 250	Gln	Asp	Ile	Gly	Arg 255	Gly
Gly	Phe	Glu	Val 260	Leu	Ala	Val	Cys	His 265	Pro	Asp	Asp	Asp	Val	Val	Asn
Ser	Val	Ile 275	Ile	Ala	Gln	Lys	Ser 280	Lys	Asp	Val	His	Ala 285	Asp	Gly	Leu

Gly Ser Gly Arg Gly Ala Gly Gly Gln Tyr Ala Arg Gly Thr Val Pro
 290 295 300

Val Val Ser Pro Pro Cys Arg Phe Gly Glu Met Val Ala Asp Val Thr
 305 310 315 320

Gln Asn His Lys Arg Asp Glu Phe Ala Asn Ala Glu Val Ala Phe
 325 330 335

<210> 4

<211> 1342

<212> DNA

<213> Hordeum vulgare

<400> 4

```
ctcctgtgcc tgtcctgagg taccaagaac accagtgaac tggctgcca gaacaaccag 60
gaggtggatg ccctggtgga gaagatcacc gggctccatg ccgcaatcgc caagctgccg 120
tcgctcagcc catccccgga cgtcgacgcg ctcttcacgg agctggtcac ggcgtgcgtt 180
ccaccgagtc cagtggacgt gaccaagctc gggccggagg cgcaggagat gcgggagggc 240
ctcatccgcc tatgctccga ggccgagggg aagctggagg cgcactactc cgacatgctc 300
gccgccttcg acaagccgct ggatcacctc ggcattgttc cctactacaa caactacatc 360
aacctcagca agctcgagta cgagctcctg gcccgctacg tgccctggcg ctatcgcccg 420
gcgcgcgtcg cgttcacatcg ctccggcccc ctgccgttca gctcctttgt cctggccgcg 480
cgccacctgc ccgacaccat gttcgacaac tatgacctgt gcggtgcggc caacgatcgc 540
gccagcaagc tcttcgcgcg ggatcgcgac gtgggtgccc gcatgtcgtt ccacacggcc 600
gacgtcgcgg acctcgccgg cgagctcgcc aagtacgacg ttgtcttcct ggccgcactc 660
gtcggcatgg ccgccgagga caaggcgaag gtgatcgcgc acctcggcgc acacatggca 720
gacggggcgg ccctcgtcgt gcgcagcgca cacggagcgc gcgggttcct gtacccgatc 780
gtcgaccccc aggacatcgg ccgaggcggg ttcgaggtgc tggccgtgtg ccatcccgac 840
gacgacgtgg tgaactccgt catcatcgca cagaagtcca aggacgtgca tgccgatgga 900
cttggcagcg ggcgtggtgc cgttgacag tacgcgcggg gcacggtgcc tgttgtcagc 960
cccccgtgca ggttcggcga gatggtggcg gacgtgaccc agaaccacaa gagagacgag 1020
tttgccaacg ccgaagtggc cttttgatcg ttcgctgcga ggggtgtgcat ccatgatcca 1080
tccatacctc gttctgtgat tgcataagc ttgcaatcgt atgcatttca agtcacgtgt 1140
tgcttctatc caataatgta cgtgtggtgt ttacacgcga atgtcttgta gacctttgta 1200
tgtgtacaag tgaattttta ttcacaaagta catataatgg tcaccattga aaagatgttt 1260
agtgtgtgtt ttccaatata tgtttgtgta aggttcacat tctaataaaa tatgtttgga 1320
acccaaaaaa aaaaaaaaaa aa 1342
```

<210> 5

<211> 335

<212> PRT

<213> Hordeum vulgare

<400> 5

Met Ala Ala Gln Asn Asn Asn Lys Asp Val Ala Ala Leu Val Glu Lys
 1 5 10 15

Ile Thr Gly Leu His Ala Ala Ile Ala Lys Leu Pro Ser Leu Ser Pro
 20 25 30

Ser Pro Asp Val Asp Ala Leu Phe Thr Glu Leu Val Thr Ala Cys Val
 35 40 45

Pro Pro Ser Pro Val Asp Val Thr Lys Leu Gly Pro Glu Ala Gln Glu
 50 55 60

Met Arg Glu Gly Leu Ile Arg Leu Cys Ser Glu Ala Glu Gly Lys Leu
 65 70 75 80
 Glu Ala His Tyr Ser Asp Met Leu Ala Ala Phe Asp Asn Pro Leu Asp
 85 90 95
 His Leu Gly Ile Phe Pro Tyr Tyr Ser Asn Tyr Ile Asn Leu Ser Lys
 100 105 110
 Leu Glu Tyr Glu Leu Leu Ala Arg Tyr Val Arg Arg His Arg Pro Ala
 115 120 125
 Arg Val Ala Phe Ile Gly Ser Gly Pro Leu Pro Phe Ser Ser Phe Val
 130 135 140
 Leu Ala Ala Arg His Leu Pro Asp Thr Met Phe Asp Asn Tyr Asp Leu
 145 150 155 160
 Cys Gly Ala Ala Asn Asp Arg Ala Ser Lys Leu Phe Arg Ala Asp Thr
 165 170 175
 Asp Val Gly Ala Arg Met Ser Phe His Thr Ala Asp Val Ala Asp Leu
 180 185 190
 Ala Ser Glu Leu Ala Lys Tyr Asp Val Val Phe Leu Ala Ala Leu Val
 195 200 205
 Gly Met Ala Ala Glu Asp Lys Ala Lys Val Ile Ala His Leu Gly Ala
 210 215 220
 His Met Ala Asp Gly Ala Ala Leu Val Val Arg Ser Ala His Gly Ala
 225 230 235 240
 Arg Gly Phe Leu Tyr Pro Ile Val Asp Pro Gln Asp Ile Gly Arg Gly
 245 250 255
 Gly Phe Glu Val Leu Ala Val Cys His Pro Asp Asp Asp Val Val Asn
 260 265 270
 Ser Val Ile Ile Ala Gln Lys Ser Lys Glu Val His Ala Asp Gly Leu
 275 280 285
 Gly Ser Ala Arg Gly Ala Gly Arg Gln Tyr Ala Arg Gly Thr Val Pro
 290 295 300
 Val Val Ser Pro Pro Cys Arg Phe Gly Glu Met Val Ala Asp Val Thr
 305 310 315 320
 Gln Asn His Lys Arg Asp Glu Phe Ala Asn Ala Glu Val Ala Phe
 325 330 335

<210> 6
 <211> 1314
 <212> DNA
 <213> Hordeum vulgare

<400> 6
 ctacttcact cacactagtg cccagaaaga aggctgcaat ggctgcccag aacaacaaca 60
 aggatgtcgc tgccctgggtg gagaagatca ccgggctcca cgccgccatc gccaaagctgc 120

```

cgtcgctcag cccatccccg gacgtcgacg cgctcttcac cgagctgggc acggcgtgcg 180
ttcccccgag ccccggtggac gtgaccaagc tcggccccga ggcgcaggag atgcgggagg 240
gcctcatccg cctctgctcc gaggcgcagg ggaagctgga ggcgcactac tccgacatgc 300
tcgccgcctt cgacaacccg ctggatcacc tcggcatctt cccctactac agcaactaca 360
tcaacctcag caagctggag tacgagctcc tggcacgcta cgtccggcgg catcgcccgg 420
cccgcgtcgc gttcatcggc tccggccccg tgccgttcag ctcttttgc ctggccgcgc 480
gccacctgcc cgacaccatg tttgacaact acgacctttg cggcgcggcc aacgatcgcg 540
ccagcaagct cttccgcgcg gacacggacg tgggtgcccg catgtcgttc cacacggccg 600
acgtcgcgga cctcgccagc gagctcgcca agtacgacgt cgtcttcctg gccgcgctcg 660
tcggcatggc cgccgaggac aaggccaagg tgatcgcgca cctcggcgca cacatggcag 720
acggggcggc cctcgtcgtg cgcagcgcac acggagcgcg cgggttcctg taccgattg 780
tcgacccccg ggacatcggc cgcggcgggt tcgaggtgct ggccgtgtgc caccgccagc 840
acgacgtggt gaactccgtc atcatcgcac agaagtccaa ggaggtgcat gccgatggac 900
ttggcagcgc gcgtggtgcc ggtcgacagt acgcgcgcgg cacggtgccg gttgtcagcc 960
ccccgtgcag gttcgggtgag atggtggcgg atgtgaccca gaaccacaag agagacgagt 1020
ttgccaacgc cgaagtggcc ttttgatcga tcgtcgccaa gggacaataa atgaacgtgg 1080
atgtggtagg gtaatttgcc tacctcgctg cttgatcgct tgcaatatgt gcacattttc 1140
ctactaccgc tgcttatgca tttcaagcca tgtgatgttg gtatccaata aagtatgtgt 1200
agggtttaca cgcaaattgc tttacacctt gtacgtgtaa gtgttgacaa cgatgaattt 1260
cagttcacia ttaataaata gtataatgga ttcaaaaaaa aaaaaaaaaa aaaa 1314

```

<210> 7

<211> 329

<212> PRT

<213> Hordeum vulgare

<400> 7

```

Met Asp Gly Gln Ser Glu Glu Val Asp Ala Leu Val Gln Lys Ile Thr
  1              5              10              15

Gly Leu His Ala Ala Ile Ala Lys Leu Pro Ser Leu Ser Pro Ser Pro
          20              25              30

Asp Val Asp Ala Leu Phe Thr Asp Leu Val Thr Ala Cys Val Pro Pro
          35              40              45

Ser Pro Val Asp Val Thr Lys Leu Ala Pro Glu Ala Gln Ala Met Arg
          50              55              60

Glu Gly Leu Ile Arg Leu Cys Ser Glu Ala Glu Gly Lys Leu Glu Ala
          65              70              75              80

His Tyr Ser Asp Met Leu Ala Ala Phe Asp Asn Pro Leu Asp His Leu
          85              90              95

Gly Val Phe Pro Tyr Tyr Ser Asn Tyr Ile Asn Leu Ser Lys Leu Glu
          100             105             110

Tyr Glu Leu Leu Ala Arg Tyr Val Pro Gly Arg His Arg Pro Ala Arg
          115             120             125

Val Ala Phe Ile Gly Ser Gly Pro Leu Pro Phe Ser Ser Tyr Val Leu
          130             135             140

Ala Ala Arg His Leu Pro Asp Thr Val Phe Asp Asn Tyr Asp Leu Cys
          145             150             155             160

Gly Ala Ala Asn Asp Arg Ala Thr Arg Leu Phe Arg Ala Asp Lys Asp
          165             170             175

```

Val Gly Ala Arg Met Ser Phe His Thr Ala Asp Val Ala Asp Leu Thr
 180 185 190
 Asp Glu Leu Ala Thr Tyr Asp Val Val Phe Leu Ala Ala Leu Val Gly
 195 200 205
 Met Ala Ala Glu Asp Lys Ala Lys Val Ile Ala His Leu Gly Ala His
 210 215 220
 Met Ala Asp Gly Ala Ala Leu Val Ala Arg His Gly Ala Arg Gly Phe
 225 230 235 240
 Leu Tyr Pro Ile Val Asp Pro Gln Asp Ile Gly Arg Gly Gly Phe Glu
 245 250 255
 Val Leu Ala Val Cys His Pro Asp Asp Asp Val Val Asn Ser Val Ile
 260 265 270
 Ile Ala Gln Lys Ser Asn Asp Val His Glu Tyr Gly Leu Gly Ser Gly
 275 280 285
 Arg Gly Gly Arg Tyr Ala Arg Gly Thr Val Val Pro Val Val Ser Pro
 290 295 300
 Pro Cys Arg Phe Gly Glu Met Val Ala Asp Val Thr Gln Lys Arg Glu
 305 310 315 320
 Glu Phe Ala Asn Ala Glu Val Ala Phe
 325

<210> 8
 <211> 1249
 <212> DNA
 <213> Hordeum vulgare

<400> 8
 ccactaccga ctaccgtagt accgtgcctc agagctcatc actggtcagg taccaagaag 60
 acataaaaat ggacggccag agcgaggagg tcgacgccct tgtccagaag atcaccggcc 120
 tccacgccgc catcgccaag ctgccctcgc tcagcccgtc cccggacgtc gacgcgctct 180
 tcaccgacct ggtcaccgcg tgcgtgcccc cgagccccgt ggacgtgacc aagctcgccc 240
 cggaggcgca ggcgatgcgg gagggcctca tccgcctctg ctccgaggcc gagggcaagc 300
 tggaggcgca ctactccgac atgctcgccg ccttcgacaa cccgctcgac cacctcggcg 360
 tcttccccta ctacagcaac tacatcaacc tcagcaagct tgagtacgag ctctcgcgcg 420
 gctacgtgcc cggcaggcat cgcccggccc gcgtgcctt catcggtcc ggcccgtgcg 480
 cgttcagctc ctacgtcctc gccgcgcgcc acctgcccga caccgtgttc gacaactacg 540
 acctgtgcgg cgcggccaac gaccgcgcga ccaggctgtt ccgcgcggac aaggacgtcg 600
 gcgcccgcac gtcgttccac accgccgacg tcgcggacct caccgacgag ctcgctacgt 660
 acgacgtcgt cttcctggcc gcgctcgtgg gcatggccgc cgaggacaag gccaaggtga 720
 tcgcgcacct tggcgcgcac atggcggacg gggcgccct cgttgccgcg cacggcgcgc 780
 gtgggttcct ctaccgatc gtcgatcccc aggacatcgg tcgaggcggg ttcgaggtgc 840
 tcgccgtgtg tcaccccgac gacgacgtgg tgaactccgt catcatcgca caaaagagca 900
 acgacgtgca cgagtatgga cttggcagcg ggcgtggtgg acggtacgcg cgaggcacgg 960
 tggtgccggt ggtcagccca ccctgcaggt tcggcgagat ggtggcagac gtgaccaga 1020
 agagagagga gtttgccaac gcggaagtgg ccttctgatt gctgctgaat cgcttgtgat 1080
 cgtacgtggt aatttttcta ctactcctcc tcctaccacc acctatcacc tatgtatgca 1140
 tttcaagtcg tgtgttgttt gtatccaata atgtaagtga gatgtttaca cgcgcaaaaa 1200
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1249

<210> 9
 <211> 282
 <212> PRT
 <213> Hordeum vulgare

<400> 9
 Met Glu Ala Glu Asn Gly Glu Val Ala Ala Leu Val Glu Lys Ile Thr
 1 5 10 15
 Gly Leu His Ala Ala Ile Ser Lys Leu Pro Ala Leu Ser Pro Ser Pro
 20 25 30
 Gln Val Asp Ala Leu Phe Thr Glu Leu Val Ala Ala Cys Val Pro Ser
 35 40 45
 Ser Pro Val Asp Val Thr Lys Leu Gly Pro Glu Ala Gln Glu Met Arg
 50 55 60
 Gln Asp Leu Ile Arg Leu Cys Ser Ala Ala Glu Gly Leu Leu Glu Ala
 65 70 75 80
 His Tyr Ser Asp Met Leu Thr Ala Leu Asp Ser Pro Leu Asp His Leu
 85 90 95
 Gly Arg Phe Pro Tyr Phe Asp Asn Tyr Val Asn Leu Ser Lys Leu Glu
 100 105 110
 His Asp Leu Leu Ala Gly His Val Ala Ala Pro Ala Arg Val Ala Phe
 115 120 125
 Ile Gly Ser Gly Pro Leu Pro Phe Ser Ser Leu Phe Leu Ala Thr Tyr
 130 135 140
 His Leu Pro Asp Thr Arg Phe Asp Asn Tyr Asp Arg Cys Ser Val Ala
 145 150 155 160
 Asn Gly Arg Ala Met Lys Leu Val Gly Ala Ala Asp Glu Gly Val Arg
 165 170 175
 Ser Arg Met Ala Phe His Thr Ala Glu Val Thr Asp Leu Thr Ala Glu
 180 185 190
 Leu Gly Ala Tyr Asp Val Val Phe Leu Ala Ala Leu Val Gly Met Thr
 195 200 205
 Ser Lys Glu Lys Ala Asp Ala Ile Ala His Leu Gly Lys His Met Ala
 210 215 220
 Asp Gly Ala Val Leu Val Arg Glu Ala Leu His Gly Ala Arg Ala Phe
 225 230 235 240
 Leu Tyr Pro Val Val Glu Leu Asp Asp Val Gly Arg Gly Gly Phe Gln
 245 250 255
 Val Leu Ala Val His His Pro Ala Gly Asp Glu Val Phe Asn Ser Phe
 260 265 270
 Ile Val Ala Arg Lys Val Lys Met Ser Ala
 275 280

<210> 10
 <211> 1044
 <212> DNA
 <213> Hordeum vulgare

<400> 10
 gtgacatgga ggccgaaaac ggcgaggtgg ctgctctggt cgagaagatc accggtctcc 60
 acgccgccat ctccaagctc ccggcactaa gcccgctctcc tcaagtcgac gcgctcttca 120
 ccgagctggt tgcggcgtgc gtcccatcaa gcccggtgga cgtgaccaag ctcggcccg 180
 aggcgcagga gatgcggcag gacctcatcc gtctctgctc ggccgccgag gggctgctcg 240
 aggcgcacta ctccgacatg ctcaccgcgt tggacagccc gctcgaccac ctcggccgct 300
 tcccttactt cgacaactac gtcaacctca gcaagctcga gcacgatctt ctggcaggtc 360
 acgtggcggc cccggcccgc gtggcgttca tcgggtcggg gccactgccg ttcagctcgc 420
 tcttccttgc gacgtaccac ctgccggaca cccggttcga caactacgac cgggtgcagcg 480
 tggcgaatgg ccgggcgatg aagctggtcg gcgcggcgga cgagggcgtg cgatcacgca 540
 tggcgttcca cacggccgaa gtcacggacc tcacggctga gctcggcgct tacgacgtgg 600
 tcttcctggc cgcgctcgtg ggaatgacgt ccaaggagaa ggccgacgcc atagcgcact 660
 tggggaagca catggcagat ggggcggtgc tcgtgcgcga agcgctgcac ggggcgcgag 720
 cgttcctgta tcctgtcgtg gagctggacg atgtcgggcg tgggtgggttc caagtgcctg 780
 ccgtgcacca ccctgcaggc gatgaggtgt tcaactcatt catagttgcc cggaagggtga 840
 aatgagtgat ttaaattaag aaaagggtga gcctgtctgc ttgtgcaa at ggtgtctcac 900
 attgataata accagatgat accctgcaca ttgatggggg tactgcagta tgtttcaatg 960
 aggtctggtt gtatcaaata tgagtatttg gcttaataat atcagcgaat atgtttcgat 1020
 taaaaaaaaa aaaaaaaaaa aaaa 1044

<210> 11
 <211> 328
 <212> PRT
 <213> Hordeum vulgare

<400> 11
 Met Asp Ala Gln Asn Lys Glu Val Asp Ala Leu Val Gln Lys Ile Thr
 1 5 10 15
 Gly Leu His Ala Ala Ile Ala Lys Leu Pro Ser Leu Ser Pro Ser Pro
 20 25 30
 Asp Val Asp Ala Leu Phe Thr Asp Leu Val Thr Ala Cys Val Pro Pro
 35 40 45
 Ser Pro Val Asp Val Thr Lys Leu Gly Ser Glu Ala Gln Glu Met Arg
 50 55 60
 Glu Gly Leu Ile Arg Leu Cys Ser Glu Ala Glu Gly Lys Leu Glu Ala
 65 70 75 80
 His Tyr Ser Asp Met Leu Ala Ala Phe Asp Asn Pro Leu Asp His Leu
 85 90 95
 Gly Met Phe Pro Tyr Tyr Ser Asn Tyr Ile Asn Leu Ser Lys Leu Glu
 100 105 110
 Tyr Glu Leu Leu Ala Arg Tyr Val Pro Gly Gly Ile Ala Arg Pro Ala
 115 120 125
 Val Ala Phe Ile Gly Ser Gly Pro Leu Pro Phe Ser Ser Tyr Val Leu
 130 135 140

Ala Ala Arg His Leu Pro Asp Ala Met Phe Asp Asn Tyr Asp Leu Cys
 145 150 155 160
 Ser Ala Ala Asn Asp Arg Ala Ser Lys Leu Phe Arg Ala Asp Lys Asp
 165 170 175
 Val Gly Ala Arg Met Ser Phe His Thr Ala Asp Val Ala Asp Leu Thr
 180 185 190
 Arg Glu Leu Ala Ala Tyr Asp Val Val Phe Leu Ala Ala Leu Val Gly
 195 200 205
 Met Ala Ala Glu Asp Lys Ala Lys Val Ile Pro His Leu Gly Ala His
 210 215 220
 Met Ala Asp Gly Ala Ala Leu Val Val Arg Ser Ala Gln Ala Arg Gly
 225 230 235 240
 Phe Leu Tyr Pro Ile Val Asp Pro Gln Asp Ile Gly Arg Gly Gly Phe
 245 250 255
 Glu Val Leu Ala Val Cys His Pro Asp Asp Asp Val Val Asn Ser Val
 260 265 270
 Ile Ile Ala His Lys Ser Lys Asp Val His Ala Asn Glu Arg Pro Asn
 275 280 285
 Gly Arg Gly Gly Gln Tyr Arg Gly Ala Val Pro Val Val Ser Pro Pro
 290 295 300
 Cys Arg Phe Gly Glu Met Val Ala Asp Val Thr His Lys Arg Glu Glu
 305 310 315 320
 Phe Thr Asn Ala Glu Val Ala Phe
 325

<210> 12
 <211> 1352
 <212> DNA
 <213> Hordeum vulgare

<400> 12
 ctccacttcg ctccctgtgcc tcaggtagcc acaacataca gtattaaaat ggatgcccag 60
 aacaaggagg ttgatgccct ggtccagaag atcaccggcc tccacgccgc catcgccaag 120
 ctgccgtccc tcagcccata acccgacgtc gacgcgtctt tcaccgacct ggtcaccgcg 180
 tgcgtccccc cgagccccgt ggacgtgacc aagctcgggt cggaggcgca ggagatgcgg 240
 gagggcctca tccgcctctg ctccgaggcc gaggggaagc tggaggcgca ctactccgac 300
 atgctggccg ccttcgacaa cccgctcgac cacctcggca tgttccccta ctacagcaac 360
 tacatcaacc tcagcaagct ggagtacgag ctccctggcg gctacgtgcc gggcggcata 420
 gcccgccccg ctgtcgcggt catcggtctc ggcccgtgc cgttcagctc ctacgtcctc 480
 gccgctcgcc acctgcccga cgccatgttc gacaactacg acctgtgtag cgcggccaac 540
 gaccgtgcga gcaagctgtt ccgcgcggac aaggacgtgg ggcgccgcat gtctttccac 600
 accgccgacg tagcggacct caccgcgag ctccgccgct acgacgtcgt cttcctggcc 660
 gcgctcgtgg gcatggctgc cgaggacaag gccaaggtga ttccgcacct cggcgcgcac 720
 atggcggacg gggcgccct cgtcgtgcgc agtgcgagc cacgtgggtt cctctaccgc 780
 atcgtcgatc cccaggacat cggtcgaggc gggtttgagg tgctggccgt gtgtcacccc 840
 gacgatgacg tggatgaactc cgtcatcatc gcacacaagt ccaaggacgt gcatgccaat 900
 gaacgtccca acgggcgtgg tggacagtac cggggcgcgg taccggtggt cagccccgcc 960

```

tgcaggttcg gtgagatggg ggcggacgtg acccacaaga gagaggagtt caccaacgcg 1020
gaagtggcct tctgatcggt gcgaggggaat gaaaatgaag gtggacgtgt gtggtcagca 1080
tccatacgtg gctgcctgct tcatcgcttg caatcgctact actacctacc tatgcagttc 1140
aagtcacgtg ttgtcaatgt aagtgtgatg ttacactag tctatgaaag gcagggcaga 1200
cgagggtagt gtgccaagta acagtgtgtc attataggtg taagtgttga gaataagacc 1260
atTTTTgttc acaaatagta tgatgtaatc ggtgtcatat tcgtattgag tacatttgtc 1320
aagttggttg ctaaaaaaaaa aaaaaaaaaa aa 1352

```

<210> 13
 <211> 329
 <212> PRT
 <213> Hordeum vulgare

```

<400> 13
Met Asp Ala Gln Ser Lys Glu Val Asp Ala Leu Val Gln Lys Ile Thr
  1          5          10          15
Gly Leu His Ala Ala Ile Ala Lys Leu Pro Ser Leu Ser Pro Ser Pro
          20          25          30
Asp Val Asp Ala Leu Phe Thr Asp Leu Val Thr Ala Cys Val Pro Pro
          35          40          45
Ser Pro Val Asp Val Thr Lys Leu Ala Pro Glu Ala Gln Ala Met Arg
          50          55          60
Glu Gly Leu Ile Arg Leu Cys Ser Glu Ala Glu Gly Lys Leu Glu Ala
  65          70          75          80
His Tyr Ser Asp Met Leu Ala Ala Phe Asp Asn Pro Leu Asp His Leu
          85          90          95
Gly Val Phe Pro Tyr Tyr Ser Asn Tyr Ile Asn Leu Ser Lys Leu Glu
          100          105          110
Tyr Glu Leu Leu Ala Arg Tyr Val Pro Gly Gly Ile Ala Pro Ala Arg
          115          120          125
Val Ala Phe Ile Gly Ser Gly Pro Leu Pro Phe Ser Ser Tyr Val Leu
          130          135          140
Ala Ala Arg His Leu Pro Asp Thr Val Phe Asp Asn Tyr Val Pro Val
          145          150          155          160
Arg Ala Ala Asn Asp Arg Ala Thr Arg Leu Phe Arg Ala Asp Lys Asp
          165          170          175
Val Gly Ala Arg Met Ser Phe His Thr Ala Asp Val Ala Asp Leu Thr
          180          185          190
Asp Glu Leu Ala Thr Tyr Asp Val Val Phe Leu Ala Ala Leu Val Gly
          195          200          205
Met Ala Ala Glu Asp Lys Gly Gln Gly Asp Pro His Leu Gly Ala His
          210          215          220
Met Ala Asp Gly Ala Ala Leu Val Arg Ser Ala His Gly Ala Arg Gly
          225          230          235          240

```


20					25					30					
Glu	Val	Asp	Ala	Leu	Phe	Thr	Asp	Leu	Val	Thr	Ala	Cys	Val	Pro	Ala
		35					40					45			
Ser	Pro	Val	Asp	Val	Ala	Lys	Leu	Gly	Pro	Glu	Ala	Gln	Ala	Met	Arg
	50					55					60				
Glu	Glu	Leu	Ile	Arg	Leu	Cys	Ser	Ala	Ala	Glu	Gly	His	Leu	Glu	Ala
	65					70					75				80
His	Tyr	Ala	Asp	Met	Leu	Ala	Ala	Phe	Asp	Asn	Pro	Leu	Asp	His	Leu
				85					90					95	
Ala	Arg	Phe	Pro	Tyr	Tyr	Gly	Asn	Tyr	Val	Asn	Leu	Ser	Lys	Leu	Glu
			100					105					110		
Tyr	Asp	Leu	Leu	Val	Arg	Tyr	Val	Pro	Gly	Ile	Ala	Pro	Thr	Arg	Val
		115					120					125			
Ala	Phe	Val	Gly	Ser	Gly	Pro	Leu	Pro	Phe	Ser	Ser	Leu	Val	Leu	Ala
	130					135					140				
Ala	His	His	Leu	Pro	Asp	Ala	Val	Phe	Asp	Asn	Tyr	Asp	Arg	Cys	Gly
	145					150					155				160
Ala	Ala	Asn	Glu	Arg	Ala	Arg	Arg	Leu	Phe	Arg	Gly	Ala	Asp	Glu	Gly
				165					170					175	
Leu	Gly	Ala	Arg	Met	Ala	Phe	His	Thr	Ala	Asp	Val	Ala	Thr	Leu	Thr
			180					185					190		
Gly	Glu	Leu	Gly	Ala	Tyr	Asp	Val	Val	Phe	Leu	Ala	Ala	Leu	Val	Gly
		195					200					205			
Met	Ala	Ala	Glu	Glu	Lys	Ala	Gly	Val	Ile	Ala	His	Leu	Gly	Ala	His
	210					215					220				
Met	Ala	Asp	Gly	Ala	Ala	Leu	Val	Val	Arg	Thr	Ala	His	Gly	Ala	Arg
	225					230					235			240	
Gly	Phe	Leu	Tyr	Pro	Ile	Val	Asp	Pro	Glu	Asp	Val	Arg	Arg	Gly	Gly
				245					250					255	
Phe	Asp	Val	Leu	Ala	Val	Cys	His	Pro	Glu	Asp	Glu	Val	Ile	Asn	Ser
			260					265					270		
Val	Ile	Val	Ala	Arg	Lys	Val	Gly	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala
		275					280					285			
Arg	Arg	Asp	Glu	Leu	Ala	Asp	Ser	Arg	Gly	Val	Val	Leu	Pro	Val	Val
	290					295					300				
Gly	Pro	Pro	Ser	Thr	Cys	Cys	Lys	Val	Glu	Ala	Ser	Ala	Val	Glu	Lys
	305					310					315				320
Ala	Glu	Glu	Phe	Ala	Ala	Asn	Lys	Glu	Leu	Ser	Val				
				325					330						

<210> 16
 <211> 1372
 <212> DNA
 <213> *Oryza sativa*

<400> 16
 ctccatttgg ttgtcatttt caactataat ccaccacaac tcgtgcaaca tcagctcact 60
 cgtgttccca accgcgacaa agcttcacag atggaggctc agaaccaaga ggtcgctgcc 120
 ctggtcgaga agatcgccgg cctccacgcc gccatctcca agctgccgtc gctgagccca 180
 tccgccgagg tggacgcgct cttcacccgac ctcgtcacgg cgtgcgtccc ggcgagcccc 240
 gtcgacgtgg ccaagctcgg cccggaggcg caggcgatgc gggaggagct catccgcctc 300
 tgctccgccc ccgagggcca cctcgaggcg cactacgccg acatgctcgc cgccttcgac 360
 aaccgcgtcg accacctcgc ccgcttcccg tactacggca actacgtcaa cctgagcaag 420
 ctggagtacg acctcctcgt ccgctacgtc cccggcattg cccccaccg cgtcgccttc 480
 gtcgggtcgg gcccgctgcc gttcagctcc ctcggtcgtc ctgcgacca cctgccggac 540
 gcggtgttcg acaactacga ccggtgcggc gcggccaacg agcgggcgag gaggtgttc 600
 cgcggcgccc acgagggcct cggcgcgcg c atggcggtcc acaccgccga cgtggcgacc 660
 ctgacggggg agctcggcgc gtacgacgtc gtgttcctgg cggcgctcgt gggcatggcg 720
 gccgaggaga aggcgggggt gatcgcgcac ctgggcgcg acatggcgga cggcgcgggc 780
 ctcgtcgtgc ggacggcgca cggggcgcg gggttcctgt acccgatcgt cgatcccgag 840
 gacgtcaggc gtggcggggt cgacgttctg gcggtgtgcc acccgagga cgaggtgatc 900
 aactccgtca tcgtcgcccc caaggtcggg gccgccgcc cgcgccgcgc ggcgcgcaga 960
 gacgagctcg cggactcgcg cggcggtggt ctgccgggtg tcgggcccgc gtccacgtgc 1020
 tgcaaggtgg aggcgagcgc ggttgagaag gcagaagagt ttgccgcaa caaggagctg 1080
 tccgtctaac agccggacga tcgaaaggcg cactatatta tggcaataaa tcatttgatt 1140
 atacttatgc tgcatttgcg aagctaaggt atactatgca agccatatgt ttgtgttcgt 1200
 acgtgttggt tgggacgtac agttgtgttg ttgtacgtcg tgaagtactg aagtgttcac 1260
 agtagatcac aagttcacag caatcaatga ggaccctgta agccagtgt aacgaggaac 1320
 atgccatctg tgtatgacag tgagaaatta tataagaaaa acattttgtg ac 1372

<210> 17
 <211> 320
 <212> PRT
 <213> *Arabidopsis thaliana*

<400> 17
 Met Ala Cys Gln Asn Asn Leu Val Val Lys Gln Ile Ile Asp Leu Tyr
 1 5 10 15
 Asp Gln Ile Ser Lys Leu Lys Ser Leu Lys Pro Ser Lys Asn Val Asp
 20 25 30
 Thr Leu Phe Gly Gln Leu Val Ser Thr Cys Leu Pro Thr Asp Thr Asn
 35 40 45
 Ile Asp Val Thr Asn Met Cys Glu Glu Val Lys Asp Met Arg Ala Asn
 50 55 60
 Leu Ile Lys Leu Cys Gly Glu Ala Glu Gly Tyr Leu Glu Gln His Phe
 65 70 75 80
 Ser Thr Ile Leu Gly Ser Leu Gln Glu Asp Gln Asn Pro Leu Asp His
 85 90 95
 Leu His Ile Phe Pro Tyr Tyr Ser Asn Tyr Leu Lys Leu Gly Lys Leu
 100 105 110
 Glu Phe Asp Leu Leu Ser Gln His Ser Ser His Val Pro Thr Lys Ile
 115 120 125

Ala Phe Val Gly Ser Gly Pro Met Pro Leu Thr Ser Ile Val Leu Ala
 130 135 140
 Lys Phe His Leu Pro Asn Thr Thr Phe His Asn Phe Asp Ile Asp Ser
 145 150 155 160
 His Ala Asn Thr Leu Ala Ser Asn Leu Val Ser Arg Asp Pro Asp Leu
 165 170 175
 Ser Lys Arg Met Ile Phe His Thr Thr Asp Val Leu Asn Ala Thr Glu
 180 185 190
 Ala Leu Asp Gln Tyr Asp Val Val Phe Leu Ala Ala Leu Val Gly Met
 195 200 205
 Asp Lys Glu Ser Lys Val Lys Ala Ile Glu His Leu Glu Lys His Met
 210 215 220
 Ala Pro Gly Ala Val Leu Met Leu Arg Arg Ala His Ala Leu Arg Ala
 225 230 235 240
 Phe Leu Tyr Pro Ile Val Asp Ser Ser Asp Leu Lys Gly Phe Gln Leu
 245 250 255
 Leu Thr Ile Tyr His Pro Thr Asp Asp Val Val Asn Ser Val Val Ile
 260 265 270
 Ala Arg Lys Leu Gly Gly Pro Thr Thr Pro Gly Val Asn Gly Thr Arg
 275 280 285
 Gly Cys Met Phe Met Pro Cys Asn Cys Ser Lys Ile His Ala Ile Met
 290 295 300
 Asn Asn Arg Gly Lys Lys Asn Met Ile Glu Glu Phe Ser Thr Ile Glu
 305 310 315 320

<210> 18
 <211> 963
 <212> DNA
 <213> Arabidopsis thaliana

<400> 18
 atggcttgcc aaaacaatct cgttgtgaag caaatcatcg acttgtagca ccaaactctca 60
 aagctcaaga gcttaaaacc ttccaaaaat gtcgacactt tggtcggaca actcgtgtcc 120
 acgtgcttac ccacggatac aaacatcgat gtcacaaata tgtgtgaaga agtcaaagac 180
 atgagagcta atctcatcaa gctttgtggt gaagccgaag gttatttgga gcaacacttc 240
 tccacaattt tgggatcttt acaagaagac caaaaccac ttgaccattt acacatcttt 300
 ccttactact ccaactacct caagctagge aagctcgagt tcgatctcct gagccaacac 360
 tcaagccatg tccccaccaa gattgccttc gtgggttcgg gtccgatgcc tctcacatcc 420
 atcgtattgg ccaagtttca cctccccaac acgacgttcc acaactttga catcgactca 480
 cacgcaaaca cactcgcttc aaacctcgtc tctcgcgacc cggacctctc aaaacgcatg 540
 atcttccaca caacggacgt actaaacgca accgaagccc ttgaccaata tgacgtcggt 600
 ttcttagcgg cgctttagg gatggacaaa gagtcaaagg tcaaagccat cgagcacttg 660
 gagaaacaca tggctoctgg agctgttctt atgctaagga gggctcatgc tctcagagct 720
 ttcttatatc caatcgttga ctcgtctgat ctcaaaggct ttcaactctt gaccatctat 780

```

catccaaccg atgacgtggt taactcgggt gtgatcgac gtaagctcgg tgggccgacc 840
acgcccgggg ttaatggtac tcgtggatgc atgtttatgc cttgtaactg ctccaagatt 900
cacgcgatca tgaacaaccg tggtaagaag aatatgatcg aggagttag taccatcgag 960
taa 963

```

<210> 19
 <211> 320
 <212> PRT
 <213> Arabidopsis thaliana

<400> 19
 Met Ala Cys Gln Asn Asn Leu Val Val Lys Gln Ile Met Asp Leu Tyr
 1 5 10 15
 Asn Gln Ile Ser Asn Leu Glu Ser Leu Lys Pro Ser Lys Asn Val Asp
 20 25 30
 Thr Leu Phe Arg Gln Leu Val Ser Thr Cys Leu Pro Thr Asp Thr Asn
 35 40 45
 Ile Asp Val Thr Glu Ile His Asp Glu Lys Val Lys Asp Met Arg Ser
 50 55 60
 His Leu Ile Lys Leu Cys Gly Glu Ala Glu Gly Tyr Leu Glu Gln His
 65 70 75 80
 Phe Ser Ala Ile Leu Gly Ser Phe Glu Asp Asn Pro Leu Asn His Leu
 85 90 95
 His Ile Phe Pro Tyr Tyr Asn Asn Tyr Leu Lys Leu Gly Lys Leu Glu
 100 105 110
 Phe Asp Leu Leu Ser Gln His Thr Thr His Val Pro Thr Lys Val Ala
 115 120 125
 Phe Ile Gly Ser Gly Pro Met Pro Leu Thr Ser Ile Val Leu Ala Lys
 130 135 140
 Phe His Leu Pro Asn Thr Thr Phe His Asn Phe Asp Ile Asp Ser His
 145 150 155 160
 Ala Asn Thr Leu Ala Ser Asn Leu Val Ser Arg Asp Ser Asp Leu Ser
 165 170 175
 Lys Arg Met Ile Phe His Thr Thr Asp Val Leu Asn Ala Lys Glu Gly
 180 185 190
 Leu Asp Gln Tyr Asp Val Val Phe Leu Ala Ala Leu Val Gly Met Asp
 195 200 205
 Lys Glu Ser Lys Val Lys Ala Ile Glu His Leu Glu Lys His Met Ala
 210 215 220
 Pro Gly Ala Val Val Met Leu Arg Ser Ala His Gly Leu Arg Ala Phe
 225 230 235 240
 Leu Tyr Pro Ile Val Asp Ser Cys Asp Leu Lys Gly Phe Glu Val Leu
 245 250 255

Thr Ile Tyr His Pro Ser Asp Asp Val Val Asn Ser Val Val Ile Ala
 260 265 270

Arg Lys Leu Gly Gly Ser Asn Gly Ala Arg Gly Ser Gln Ile Gly Arg
 275 280 285

Cys Val Val Met Pro Cys Asn Cys Ser Lys Val His Ala Ile Leu Asn
 290 295 300

Asn Arg Gly Met Glu Lys Asn Leu Ile Glu Glu Phe Ser Ala Ile Glu
 305 310 315 320

<210> 20
 <211> 963
 <212> DNA
 <213> Arabidopsis thaliana

<400> 20
 atggcttgcc aaaacaatct cggtgtgaag caaatcatgg acttatacaa ccaaattctca 60
 aacctcgaga gcttaaaacc atccaagaat gtcgacactt tggtcagaca acttgtgtcc 120
 acgtgcttac caacggacac gaacatcgat gtcacagaga tacacgatga aaaagtcaaa 180
 gacatgagat ctcatctcat caagctttgt ggtgaagccg aaggttattt agagcaacac 240
 ttttcagcaa tcttaggctc ttttgaagac aaccctctaa accatttaca catcttcccc 300
 tattacaaca actatctcaa actaggcaaa ctcgaaattcg atctcctttc tcagcacaca 360
 acccatgtcc cgaccaaagt cgcctttatt ggttcgggtc cgatgccact tacttccatc 420
 gtcttggtcca agttccacct ccccaacaca acgttccaca acttcgacat cgactcacac 480
 gccaacacac tcgcttcaaa cctcgtttct cgtgattctg acctttccaa acgcatgatt 540
 ttccacacaa ctgatgtatt aaacgctaag gaggggttag accaatacga tgttggtttc 600
 ttggcagctc ttgttggtat ggataaagag tcaaagggtca aagctattga gcatttagag 660
 aagcatatgg cccctggagc tgtggtgatg ctaagaagtg ctcatggtct tagagctttc 720
 ttgtatccaa tcgttgactc ttgtgatctt aaagggtttg aggtgttaac catttatcat 780
 ccgtctgacg acgtggttaa ttcggtggtc atcgcacgta agcttggtgg ttcaaattga 840
 gctcgaggca gccagatcgg acggtgtgtg gttatgcctt gtaattgctc taagggtccac 900
 gcgatcttga acaatcgtgg tatggagaag aatttgatcg aggagtttag tgccatcgag 960
 taa 963

<210> 21
 <211> 320
 <212> PRT
 <213> Arabidopsis thaliana

<400> 21
 Met Gly Cys Gln Asp Glu Gln Leu Val Gln Thr Ile Cys Asp Leu Tyr
 1 5 10 15

Glu Lys Ile Ser Lys Leu Glu Ser Leu Lys Pro Ser Glu Asp Val Asn
 20 25 30

Ile Leu Phe Lys Gln Leu Val Ser Thr Cys Ile Pro Pro Asn Pro Asn
 35 40 45

Ile Asp Val Thr Lys Met Cys Asp Arg Val Gln Glu Ile Arg Leu Asn
 50 55 60

Leu Ile Lys Ile Cys Gly Leu Ala Glu Gly His Leu Glu Asn His Phe

65						70						75						80
Ser	Ser	Ile	Leu	Thr	Ser	Tyr	Gln	Asp	Asn	Pro	Leu	His	His	Leu	Asn			
				85					90					95				
Ile	Phe	Pro	Tyr	Tyr	Asn	Asn	Tyr	Leu	Lys	Leu	Gly	Lys	Leu	Glu	Phe			
			100					105					110					
Asp	Leu	Leu	Glu	Gln	Asn	Leu	Asn	Gly	Phe	Val	Pro	Lys	Ser	Val	Ala			
		115					120					125						
Phe	Ile	Gly	Ser	Gly	Pro	Leu	Pro	Leu	Thr	Ser	Ile	Val	Leu	Ala	Ser			
	130					135					140							
Phe	His	Leu	Lys	Asp	Thr	Ile	Phe	His	Asn	Phe	Asp	Ile	Asp	Pro	Ser			
145					150					155					160			
Ala	Asn	Ser	Leu	Ala	Ser	Leu	Leu	Val	Ser	Ser	Asp	Pro	Asp	Ile	Ser			
				165					170					175				
Gln	Arg	Met	Phe	Phe	His	Thr	Val	Asp	Ile	Met	Asp	Val	Thr	Glu	Ser			
			180					185					190					
Leu	Lys	Ser	Phe	Asp	Val	Val	Phe	Leu	Ala	Ala	Leu	Val	Gly	Met	Asn			
		195					200					205						
Lys	Glu	Glu	Lys	Val	Lys	Val	Ile	Glu	His	Leu	Gln	Lys	His	Met	Ala			
	210					215					220							
Pro	Gly	Ala	Val	Leu	Met	Leu	Arg	Ser	Ala	His	Gly	Pro	Arg	Ala	Phe			
225					230					235					240			
Leu	Tyr	Pro	Ile	Val	Glu	Pro	Cys	Asp	Leu	Gln	Gly	Phe	Glu	Val	Leu			
				245					250					255				
Ser	Ile	Tyr	His	Pro	Thr	Asp	Asp	Val	Ile	Asn	Ser	Val	Val	Ile	Ser			
			260					265					270					
Lys	Lys	His	Pro	Val	Val	Ser	Ile	Gly	Asn	Val	Gly	Gly	Pro	Asn	Ser			
		275					280					285						
Cys	Leu	Leu	Lys	Pro	Cys	Asn	Cys	Ser	Lys	Thr	His	Ala	Lys	Met	Asn			
	290					295					300							
Lys	Asn	Met	Met	Ile	Glu	Glu	Phe	Gly	Ala	Arg	Glu	Glu	Gln	Leu	Ser			
305					310					315					320			

<210> 22
 <211> 963
 <212> DNA
 <213> Arabidopsis thaliana

<400> 22
 atgggttgcc aagacgaaca attggtgcaa acaatatgcg atctctacga aaagatctca 60
 aagcttgaga gtctaaaacc atccgaagat gtcaacattc tcttcaagca gtcggtttcc 120
 acatgcatac caccaaacc taacatcgat gtcaccaaga tgtgtgacag agtccaagag 180

attcgactta	atctcatcaa	gatttgtggt	ctagccgaag	gtcacttaga	aaaccatttc	240
tcttcgatct	tgacctctta	ccaagacaac	ccacttcac	atttaaaca	tttcccttat	300
tacaacaact	atttgaaact	cggaaagctc	gagttcgacc	tcctcgaaca	aaacctaaat	360
ggctttgtcc	caaagagtgt	ggctttcatt	ggatctggtc	ctcttcctct	cacttccatc	420
gttcttgctt	cattccatct	caaagacaca	atctttcaca	actttgacat	cgacccatca	480
gcgaactcac	tcgcttctct	tctggtttcc	tctgatccag	acatctctca	acgcatgttc	540
ttccacaccg	ttgatataat	ggacgtgaca	gagagcttaa	agagctttga	tgtcgtgttt	600
ctagctgctc	ttgttggaat	gaacaaagag	gagaaagtta	aagtgatcga	gcatctgcag	660
aaacacatgg	ctcctgggtgc	tgtgctcatg	cttaggagtg	ctcatgggtcc	gagagcggtt	720
ctttatccga	tcgttgagcc	gtgtgatctt	caggggttcg	aggttttgtc	tatttatcac	780
ccaacagatg	atgttatcaa	ctccgtgggtg	atctctaaaa	agcatccagt	tgtttcaatt	840
gggaatgttg	gtggtcctaa	ttcatgcttg	ctcaagcctt	gcaactgttc	caagaccac	900
gcgaaaatga	acaagaacat	gatgatcgag	gagttcggag	ctagggagga	acagttgtct	960
taa						963

CLAIMS:

1. A protein exhibiting nicotianamine synthase activity, wherein the protein is selected from:

(a) the polypeptide having the amino acid sequence of SEQ ID NO:1;

or

(b) a polypeptide having at least 90% identity with an amino acid sequence of SEQ ID NO:1 and comprising at least one consensus sequence of SEQ ID NO:1 selected from the group consisting of:

(1) ₂₅LPXLSPSPXVDRLFTXLVXACVPXSPVDVTKL₅₆;

(2) ₆₇LIRLCSXAEGXLEAHY₈₂;

(3) ₉₂PLDHLGXFPY₁₀₁;

(4) ₁₂₈VAFXGSGPLPFSS₁₄₀;

(5) ₁₉₉DVVFLAALVGM₂₀₉; and

(6) ₂₅₃RGGFXVLAVXHP₂₆₄;

wherein X is variable amino acid residue.

2. The protein according to claim 1 wherein said protein originates from barley.

3. The protein according to claim 1 or 2, wherein the protein has amino acid sequence shown in SEQ ID NO: 1, 3, 5, 7, 9, 11 or 13.

4. The protein according to claim 1 wherein said protein originates from Arabidopsis.

5. The protein according to claim 4, wherein the protein has amino acid sequence shown in SEQ ID NO: 17, 19 or 21.

6. The protein according to claim 1 wherein said protein originates from Oryza sativa.

7. The protein according to claim 6, wherein the protein has amino acid sequence shown in SEQ ID NO: 15.
8. A gene encoding amino acid sequence of the protein according to any one of claims 1 - 7.
9. The gene according to claim 8 wherein said gene is cDNA.
10. The gene according to claim 8 or 9 wherein the gene has base sequence shown in SEQ ID NO: 2, 4, 6, 8, 10, 12 or 14.
11. The gene according to claim 8 or 9 wherein the gene has base sequence shown in SEQ ID NO: 18, 20 or 22.
12. A vector containing the gene according to any one of claims 8 - 11.
13. The vector according to claim 12 wherein said vector is an expression vector.
14. A transformant comprising a single host cell transformed by the vector according to claim 12 or 13.
15. The transformant according to claim 14 wherein the vector has the base sequence shown in SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, or 22.
16. The transformant according to claim 14 or 15 wherein the host cell is bacteria.
17. A process for production of nicotianamine comprising the steps of:
 - (a) culturing the transformant according to any one of claims 14-16 in medium; and
 - (b) purifying nicotianamine from the medium.
18. A genetically-engineered plant cell wherein the gene according to any one of claims 8-11 is introduced.
19. A genetically-engineered fruit cell obtained by growing a plant from the plant cell according to claim 18.

20. An antibody against nicotianamine synthase according to any one of claims 1 - 7.
21. The antibody according to claim 21 wherein said antibody is polyclonal antibody.
22. A method for purification of nicotianamine synthase comprising the steps of:
- (a) crushing frozen plant roots into a powder in liquid N₂ and then homogenizing with buffer containing thiol protease inhibitor E-64 to obtain a homogenate;
 - (b) obtaining supernatant from the homogenate;
 - (c) fractionating the supernatant by ion-exchange column chromatography to obtain an active fraction comprising nicotianamine synthase; and
 - (d) separating the nicotianamine synthase from the active fraction by SDS-polyacrylamide gel electrophoresis.




Fig. 1

Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6

	GCG	TTC	AGA	GGC	TTC	CAG	AGT	TCT	TCC	GGT	CAC	CAA	GAA	GCA	TTT	GAT	CAT	AAC	54
19	ATG	GAT	GCC	CAG	AAC	AAG	GAG	GTC	GCT	GCT	CTG	ATC	GAG	AAG	ATC	GCC	GGT	ATC	108
	M	D	A	Q	N	K	E	V	A	A	L	I	E	K	I	A	G	I	
		①																	
37	CAG	GCC	GCC	ATC	GCC	GAG	CTG	CCG	TCG	CTG	AGC	CCG	TCC	CCC	GAG	GTC	GAC	AGG	162
	Q	A	A	I	A	E	L	P	S	L	S	P	S	P	E	V	D	R	
55	CTC	TTC	ACC	GAC	CTC	GTC	ACG	GCC	TGC	GTC	CCG	CCG	AGC	CCC	GTC	GAC	GTG	ACG	216
	L	F	T	D	L	V	T	A	C	V	P	P	S	P	V	D	V	T	
73	AAG	CTC	AGC	CCG	GAG	CAC	CAG	AGG	ATG	CGG	GAG	GCT	CTC	ATC	CGC	TTG	TGC	TCC	270
	K	L	S	P	E	H	Q	R	M	R	E	A	L	I	R	L	C	S	
										②									
91	GCC	GCC	GAG	GGG	AAG	CTC	GAG	GCG	CAC	TAC	GCC	GAC	CTG	CTC	GCC	ACC	TTC	GAC	324
	A	A	E	G	K	L	E	A	H	Y	A	D	L	L	A	T	F	D	
109	AAC	CCG	CTC	GAC	CAC	CTC	GGC	CTC	TTC	CCG	TAC	TAC	AGC	AAC	TAC	GTC	AAC	CTC	378
	N	P	L	D	H	L	G	L	F	P	Y	Y	S	N	Y	V	N	L	
127	AGC	AGG	CTG	GAG	TAC	GAG	CTC	CTG	GCG	CGC	CAC	GTG	CCG	GGC	ATC	GCG	CCG	GCG	432
	S	R	L	E	Y	E	L	L	A	R	H	V	P	G	I	A	P	A	
145	CGC	GTC	GCC	TTC	GTC	GGC	TCC	GGC	CCG	CTG	CCG	TTC	AGC	TCG	CTC	GTC	CTC	GCC	486
	R	V	A	F	V	G	S	G	P	L	P	F	S	S	L	V	L	A	
163	GCG	CAC	CAC	CTG	CCC	GAG	ACC	CAG	TTC	GAC	AAC	TAC	GAC	CTG	TGC	GGC	GCG	GCC	540
	A	H	H	L	P	E	T	Q	F	D	N	Y	D	L	C	G	A	A	
181	AAC	GAG	CGC	GCC	AGG	AAG	CTG	TTC	GGC	GCG	ACG	GCG	GAC	GGC	GTC	GGC	GCG	CGT	594
	N	E	R	A	R	K	L	F	G	A	T	A	D	G	V	G	A	R	
199	ATG	TCG	TTC	CAC	ACG	GCG	GAC	GTC	GCC	GAC	CTC	ACC	CAG	GAG	CTC	GGC	GCC	TAC	648
	M	S	F	H	T	A	D	V	A	D	L	T	Q	E	L	G	A	Y	
		③																	
217	GAC	GTG	GTC	TTC	CTC	GCC	GCG	CTC	GTC	GGC	ATG	GCA	GCC	GAG	GAG	AAG	GCC	AAG	702
	D	V	V	F	L	A	A	L	V	G	M	A	A	E	E	K	A	K	
235	GTG	ATT	GCC	CAC	CTG	GGC	GCG	CAC	ATG	GTG	GAG	GGG	GCG	TCC	CTG	GTC	GTG	CGG	756
	V	I	A	H	L	G	A	H	M	V	E	G	A	S	L	V	V	R	
253	AGC	GCA	CGG	CCC	CGC	GGC	TTT	CTT	TAC	CCC	ATT	GTC	GAC	CCG	GAG	GAC	ATC	AGG	810
	S	A	R	P	R	G	F	L	Y	P	I	V	D	P	E	D	I	R	
														④					
271	CGG	GGT	GGG	TTC	GAG	GTG	CTG	GCC	GTG	CAC	CAC	CCG	GAA	GGT	GAG	GTG	ATC	AAC	864
	R	G	G	F	E	V	L	A	V	H	H	P	E	G	E	V	I	N	
289	TCT	GTC	ATC	GTC	GCC	CGT	AAG	GCC	GTC	GAA	GCG	CAG	CTC	AGT	GGG	CCG	CAG	AAC	918
	S	V	I	V	A	R	K	A	V	E	A	Q	L	S	G	P	Q	N	
307	GGA	GAC	GCG	CAC	GCA	CGG	GGC	GCG	GTG	CCG	TTG	GTC	AGC	CCG	CCA	TGC	AAC	TTC	972
	G	D	A	H	A	R	G	A	V	P	L	V	S	P	P	C	N	F	
325	TCC	ACC	AAG	ATG	GAG	GCG	AGC	GCG	CTT	GAG	AAG	AGC	GAG	GAG	CTG	ACC	GCC	AAA	1026
	S	T	K	M	E	A	S	A	L	E	K	S	E	E	L	T	A	K	
	GAG	CTG	GCC	TTT	TGA	TTG	AAG	AGT	GCG	CGT	GGT	CAT	TCT	GTC	GCC	TGC	GAT	CGT	1080
	E	L	A	F	*														
	GGT	AAC	TTT	CCT	ACT	CGT	GTG	TGT	TTT	GAT	GTT	TGT	GCC	TGT	AAG	AGT	TAT	GCT	1134
	TCC	GGC	CTT	GTG	CTG	TTA	ATT	TAC	ACG	CGT	TAC	ATG	TAG	TAC	TTG	TAT	TTA	TAC	1188
	CTG	GAA	TAA	CGG	TAT	GTA	ACA	TAA	ATA	TTA	GTG	GGA	TTT	GAA	GTG	TAA	TGC	TAA	1242
	ATA	ATA	AGA	AAA	CTT	GAT	GCA	GAC	ATT	CAA	AAA	AAA	AAA	AAA	AAA	AAA	AAA	AA	

Fig. 7

HvNAS4	MDGQSE--EVDALVQKITGLHAAIAKLPSLSPSPDVDALFTDLVTACVPPSPVDVTKLAP
HvNAS7	MDAQSK--EVDALVQKITGLHAAIAKLPSLSPSPDVDALFTDLVTACVPPSPVDVTKLAP
HvNAS6	MDAQNK--EVDALVQKITGLHAAIAKLPSLSPSPDVDALFTDLVTACVPPSPVDVTKLGS
HvNAS2	MAAQNN-QEVDALVEKITGLHAAIAKLPSLSPSPDVDALFTDLVTACVPPSPVDVTKLGP
HvNAS3	MAAQNNNKDVAALVEKITGLHAAIAKLPSLSPSPDVDALFTDLVTACVPPSPVDVTKLGP
HvNAS1	MDAQNK--EVAALIEKIAGIQAAIAELPSLSPSPEVDRLFTDLVTACVPPSPVDVTKLSP
HvNAS5	MEAENG--EVAALVEKITGLHAAISKLPALSPSPQVDALFTDLVAACVPSSPVDVTKLGP
	* * ** * * * * * * * * * * * * * * * *
HvNAS4	EAQAMREGLIRLCSEAEGKLEAHYSOMLAAFDNPLDHLGVFPYYSNYINLSKLEYELLAR
HvNAS7	EAQAMREGLIRLCSEAEGKLEAHYSOMLAAFDNPLDHLGVFPYYSNYINLSKLEYELLAR
HvNAS6	EAQEMREGLIRLCSEAEGKLEAHYSOMLAAFDNPLDHLGMFPYYSNYINLSKLEYELLAR
HvNAS2	EAQEMREGLIRLCSEAEGKLEAHYSOMLAAFDKPLDHLGMFPYNNYINLSKLEYELLAR
HvNAS3	EAQEMREGLIRLCSEAEGKLEAHYSOMLAAFDNPLDHLGIFPYYSNYINLSKLEYELLAR
HvNAS1	EHQRMREALIRLCSAAEGKLEAHYADLLATFDNPLDHLGLFPYYSNYVNLRSLEYELLAR
HvNAS5	EAQEMRQDLIRLCSAAEGLLEAHYSOMLTALDSPLDHLGRFPYFDNYVNLKLEHDLLAG
	* * ** * * * * * * * * * * * * * * * *
HvNAS4	YVPGRRHRPARVAFIGSGPLPFSSYVLAARHLPDTPVDNYDLCGAANDRATRLFRADKD-V
HvNAS7	YVPGGIAPARVAFIGSGPLPFSSYVLAARHLPDTPVDNYVPVRAANDRATRLFRADKD-V
HvNAS6	YVPGGIAPARVAFIGSGPLPFSSYVLAARHLPDAMFDNYDLCGAANDRASKLFRADKD-V
HvNAS2	YVPGGYRPARVAFIGSGPLPFSSFVLAARHLPDTMFDNYDLCGAANDRASKLFRADRD-V
HvNAS3	YVRR-HRPARVAFIGSGPLPFSSFVLAARHLPDTMFDNYDLCGAANDRASKLFRADTD-V
HvNAS1	HVPG-IAPARVAFVGSGLPFSSVLAARHLPETQFDNYDLCGAANERARKLFGATADGV
HvNAS5	HVAA---PARVAFIGSGPLPFSSLFLATYHLPDTRFDNYDRCSVANGRAMKLVGAADGV
	* *** * * * * * * * * * * * * * * *
HvNAS4	GARMSFHTADVADLTDELATYDVVFLAALVGMAAEDKAKVIAHLGAHMADGAALV--ARH
HvNAS7	GARMSFHTADVADLTDELATYDVVFLAALVGMAAEDKGQGDPHLGAHMADGAALVR-SAH
HvNAS6	GARMSFHTADVADLTRELAAYDVVFLAALVGMAAEDKAKVIPHLGAHMADGAALVV-RSA
HvNAS2	GARMSFHTADVADLAGELAKYDVVFLAALVGMAAEDKAKVIAHLGAHMADGAALVVRSAH
HvNAS3	GARMSFHTADVADLASELAKYDVVFLAALVGMAAEDKAKVIAHLGAHMADGAALVVRSAH
HvNAS1	GARMSFHTADVADLTQELGAYDVVFLAALVGMAAEEKAKVIAHLGAHMVEGASLVV-RSA
HvNAS5	RSRMFHTAEVTDLTAEFGAYDVVFLAALVGMTSKEKADIAHLGKHMADGAVLVREALH
	** **** * ** * * * * * * * * * * * * * * *
HvNAS4	GARGFLYPIDPQDIGRGGFEVLAVCHPD-DDVNSVIIAQKSNDVHEYGLGSGR--GGR
HvNAS7	GARGFLYPIDPQDIGRGGFEVLAVCHPD-DDVNSVIIAQKSKDMFANGPRNGC--GGR
HvNAS6	QARGFLYPIDPQDIGRGGFEVLAVCHPD-DDVNSVIIAHKSKDVHANERPNGR--GGQ
HvNAS2	GARGFLYPIDPQDIGRGGFEVLAVCHPD-DDVNSVIIAQKSKDVHADGLGSGRGAGGQ
HvNAS3	GARGFLYPIDPQDIGRGGFEVLAVCHPD-DDVNSVIIAQKSKEVHADGLGSARGAGRQ
HvNAS1	RPRGFLYPIDPEDIRGGFEVLAVHPE-GEVINSVIVARKAVEAQLSGPQNGD----A
HvNAS5	GARAFLYPVVELDDVGRGGFQVLAVHHPAGDEVFNSFIVARKVKMSA-----
	* **** * * **** * * * * * * * * * * *
HvNAS4	YARGTVPVVSPPCRFG-EMVADVTDQ--KREEFANAFAVAF
HvNAS7	YARG-TVPVSPPCRFG-EMVADVTDQ--KREEFAKAEVAF
HvNAS6	YRGA--VPVSPPCRFG-EMVADVTH--KREEFTNAFAVAF
HvNAS2	YARG-TVPVSPPCRFG-EMVADVTDQNHKRDEFANAFAVAF
HvNAS3	YARG-TVPVSPPCRFG-EMVADVTDQNHKRDEFANAFAVAF
HvNAS1	HARG-AVPLVSPPCNFSTKMEASALE--KSEELTAKELAF
	** **** * * * * * * * * * * *

2327261 (Draw)

UNSCANNABLE ITEM

RECEIVED WITH THIS APPLICATION

(ITEM ON THE 10TH FLOOR ZONE 5 IN THE FILE PREPARATION SECTION)

DOCUMENT REÇU AVEC CETTE DEMANDE

NE POUVANT ÊTRE BALAYÉ

(DOCUMENT AU 10 IÈME ÉTAGE AIRE 5 DANS LA SECTION DE LA

PRÉPARATION DES DOSSIERS)