

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 June 2009 (11.06.2009)

PCT

(10) International Publication Number
WO 2009/073756 A1

(51) International Patent Classification:

A61K 31/55 (2006.01) A23K 1/165 (2006.01)
A61K 31/57 (2006.01) A61P 43/00 (2006.01)
A23K 1/16 (2006.01)

[US/US]; P.O. Box 318, 29160 Intervet Lane, Millsboro,
DE 19966 (US).

(74) Agent: **BLACKSTONE, William, M.**; Intervet Inc., P.O.
Box 318, 29160 Intervet Lane, Millsboro, DE 19966 (US).

(21) International Application Number:

PCT/US2008/085455

(22) International Filing Date:

4 December 2008 (04.12.2008)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/992,922 6 December 2007 (06.12.2007) US

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **INTERVET INTERNATIONAL B.V.** [NL/NL]; Wim De Koverstraat 35, NL-5831 AN Boxmeer (NL).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **WRAY, Mary, Irene** [US/US]; 35500 West 91st Street, Desoto, KS 66018 (US). **BRADLEY, Damon, Edward** [US/US]; 35500 West 91st Street, Desoto, KS 66018 (US). **PETERSEN, Melissa, A.** [US/US]; 35500 West 91st Street, Desoto, KS 66018 (US). **SHARP, Auddie** [US/US]; 35500 West 91st Street, Desoto, KS 66018 (US). **SHELTON, Celia** [US/US]; P.O. Box 318, 29160 Intervet Lane, Millsboro, DE 19966 (US). **MONTGOMERY, Jayden, Lloyd**

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report

(54) Title: COMBINATION THERAPIES USING MELENGESTROL ACETATE AND ZILPATEROL OR ITS SALTS

(57) Abstract: This invention generally relates to a method for promoting growth and increasing feed utilization efficiency in animals, and, more specifically, to combinations comprising zilpaterol (and salts thereof) and melengestrol acetate. Included within this invention are treatment methods comprising the administration of such combinations to animals, compositions comprising such combinations, uses of such combinations to prepare medicaments, and kits for using such combinations.



WO 2009/073756 A1

COMBINATION THERAPIES USING MELENGESTROL ACETATE AND ZILPATEROL OR ITS SALTS


FIELD OF THE INVENTION

5 [1] This invention generally relates to a method for promoting growth and increasing feed utilization efficiency in animals, and, more specifically, to combinations comprising zilpaterol (and salts thereof) and melengestrol acetate. Included within this invention are treatment methods comprising the administration of such combinations to animals, compositions comprising such combinations, uses of
10 such combinations to prepare medicaments, and kits for using such combinations.

BACKGROUND OF THE INVENTION

 [2] In many countries, commercial livestock rearing systems have become commonplace. Commercial animal husbandry techniques have been used to rear, for
15 example, livestock, poultry, and fish. These techniques have resulted in greatly increased production of food products from such animals. Successful commercial raising of animals, particularly for food, requires maximization of the growth rate and feed utilization efficiency to reduce the unit cost of production. This has led to the development and widespread use of feed additives.

20 [3] Zilpaterol is a known adrenergic β -2 agonist corresponding in structure to **Formula (I)**:




The IUPAC name for zilpaterol is 4,5,6,7-tetrahydro-7-hydroxy-6-(isopropylamino)imidazo[4,5,1-jk]-[1]benzazepin-2(1H)-one. The Chemical Abstracts name for
25 zilpaterol is 4,5,6,7-tetrahydro-7-hydroxy-6-[(1-methyl-ethyl) amino]-imidazo [4,5,1-jk][1]benzazepin-2(1H)-one.

 [4] Zilpaterol hydrochloride is sold by Intervet Inc., a part of Schering-Plough Corporation, under the trademark ZILMAX[®]. It is approved in the United States for cattle fed in confinement for harvest at daily doses ranging from 60 to 90

mg/animal to increase the rate of weight gain, improve feed efficiency, and increase carcass leanness during the last 20 to 40 days on feed. *See* NADA No. 141-258.

[5] In U.S. Patent 4,585,770, Fréchet et al. discuss compounds, such as zilpaterol, encompassed by a genus characterized as 6-amino-7-hydroxy-4,5,6,7-tetrahydro-imidazo[4,5,1-j-k][1]-benzazepin-2-(1H)-one derivatives and acid addition salts thereof. Fréchet et al. state that such compounds may be used as an active ingredient for inducing antihypertensive and hypotensive activity in a warm-blooded animal.

[6] In U.S. Patent 4,900,735, Grandadam discusses a zootechnical composition comprising zilpaterol and acid addition salts thereof. Grandadam states that such a composition may be used to increase the weight of cattle, pigs, sheep, and poultry. Grandadam also discusses combination therapies that further comprise the administration of a steroid corresponding in structure to **Formula (II)**:




Here, X is selected from the group consisting of hydrogen, optionally unsaturated alkyl of 1 to 6 carbon atoms wherein one of the carbon atoms may be replaced by oxygen, and acyl of an organic carboxylic acid of 1 to 18 carbon atoms. Such compounds include trenbolone acetate, wherein X is -C(O)CH₃. Trenbolone acetate is approved in the United States for use, alone and in combination with other active ingredients, in various implants in cattle to increase the rate of weight gain and improve feed efficiency. *See* NADA Nos. 138-612, 140-897, 140-992, 414-043, 141-269, 200-221, 200-224, 200-346, and 200-367. Grandadam goes on to discuss combinations that further comprise the administration of zeranol or estradiol (*i.e.*, 17 β -estra-1,3,5(10)-triene-3,17-diol). Zeranol is approved in the United States for use in implants to increase the rate of weight gain and improve feed efficiency in cattle and sheep. *See* NADA Nos. 038-233 and 141-0192. Estradiol and estradiol benzoate are approved in the United States for use in combination with other active ingredients in various implants for increasing the rate of weight gain and improving feed efficiency in cattle. *See* NADA Nos. 009-576, 011-427, 110-315, 118-123, 135-906, 140-897, 140-992, 141-043, 141-269, 200-221, 200-346, and 200-367.

[7] In U.S. Patents 5,731,028 and 5,847,124, Chevremont et al. discuss crystallized anhydrous zilpaterol hydrochloride, and particularly crystallized anhydrous zilpaterol hydrochloride wherein less than 5% of the crystals have a size of less than 15 μm , and at least 95% of the crystals have a size of less than 250 μm .

5 According to Chevremont et al., such crystals may be incorporated into animal feed to increase body weight and meat quality. Chevremont et al. provide methods for making such crystals, and discuss using the crystals to make animal premixes in which the crystals are secured to a corn cob support having a greater particle size. They also discuss monohydrate and trihydrate intermediates that can be useful in, for
10 example, making the crystals.

[8] In U.S. Patent 7,207,289, Montgomery discusses methods for increasing beef production, reducing feed intake while maintaining beef production, and reducing incidences of liver abscess in cattle. These methods comprise administering a feed comprising an ionophore and macrolide antibiotic during an
15 initial period, and then administering a feed comprising zilpaterol (including zilpaterol hydrochloride) with essentially no ionophore or macrolide antibiotic

[9] Melengestrol acetate (or "17 α -acetoxy-6-methyl-16-methylenepregna-4,6-diene-3,20-dione") corresponds in structure to **Formula (III)**:



20 It is reported to be an orally-active progestogen that inhibits estrus and ovulation, and increases weight gain in heifers. It is commercially available from Pfizer Animal Health under the trade name MGA[®]. US Patent 3,359,287 discusses preparation of melengestrol acetate by dehydrogenating 17 α -hydroxy-6 α -methyl-16-methylenepregna-4-ene-3,20--dione 17-acetate with chloranil. US Patent 4,154,748
25 discusses an alternative process wherein melengestrol acetate is prepared by a process comprising acetylation of 17 α -hydroxy-6-methyl-16-methylenepregna-4,6-diene-3,20-dione.

[10] Melengestrol acetate is approved in the United States for feeding to heifers for harvest at doses ranging from 0.25 to 0.50 mg/heifer daily to increase weight gain, improve gain efficiency, and suppress estrus. See NADA 034-254, 039-402, and 200-343. It is generally fed to heifers for 90 to 150 days before harvest. It also has been approved in the United States for the following combinations:

- a) With oxytetracycline to increase the rate of weight gain, improve feed efficiency, suppress estrus, and reduce liver condemnation due to liver abscesses. See NADA 046-718 and 046-719
- b) With monensin sodium to increase the rate of weight gain, improve feed efficiency, suppress estrus, and prevent and control coccidiosis due to *Eimeria bovis* and *Eimeria zuernii*. See NADA 124-309, 125-476, and 200-422.
- c) With tylosin phosphate to increase the rate of weight gain, improve feed efficiency, suppress estrus, and reduce incidence of liver abscesses. See NADA 138-995, 139-192, and 200-427.
- d) With lasalocid to increase the rate of weight gain, improve feed efficiency, and suppress estrus. See NADA 139-876, 140-288, and 200-451.
- e) With monensin sodium and tylosin phosphate to increase the rate of weight gain, improve feed efficiency, suppress estrus, prevent and control coccidiosis due to *Eimeria bovis* and *Eimeria zuernii*, and reduce incidence of liver abscesses caused by *Fusobacterium necrophorum* and *Actinomyces (Corynebacterium) pyogenes*. See NADA 138-792, 138-870, and 200-375.
- f) With tylosin phosphate and lasalocid to increase the rate of weight gain, improve feed efficiency, suppress estrus, and reduce incidence of liver abscesses. See NADA 138-904, 138-992, and 200-430.
- g) With monensin sodium and ractopamine hydrochloride to increase the rate of weight gain, improve feed efficiency, suppress estrus, increase carcass leanness, and prevent and control coccidiosis due to *Eimeria bovis* and *Eimeria zuernii*. See NADA 141-234 and 200-448.
- h) With monensin sodium, tylosin phosphate, and ractopamine hydrochloride to increase the rate of weight gain, improve feed efficiency, suppress estrus, increase carcass leanness, prevent and

control coccidiosis due to *Eimeria bovis* and *Eimeria zuernii*, and reduce liver abscesses caused by *Fusobacterium necrophorum* and *Actinomyces (Corynebacterium) pyogenes*. See NADA 141-233 and 200-424.

- 5 US Patent No. 3, 417,182 discusses using melengestrol acetate to control estrual periods and stimulate growth of domestic birds and other animals.

[11] There still exists a need for alternative methods and compositions for increasing the rate of weight gain and improving feed efficiency in animals, particularly animals raised for food, such as livestock, poultry, and/or fish. The following disclosure describes such methods and compositions.

10

SUMMARY OF THE INVENTION

[12] Briefly, this invention is directed, in part, to a method for improving feed utilization efficiency and/or increasing the rate of weight gain in an animal, particularly an un-spayed female animal. The animal may be, for example, a livestock animal (e.g., a bovine, sheep, goat, or other ruminant; swine; etc.), bird (e.g., a chicken, turkey, etc.), or fish (e.g., a salmon, trout, catfish, etc.). The method comprises administering to the animal zilpaterol (or a salt thereof, such as zilpaterol hydrochloride) and melengestrol acetate. Other benefits of this method generally include, for example, increases in carcass leanness.

15

20

[13] This invention also directed, in part, to uses of zilpaterol (or a salt thereof) and melengestrol acetate to make a medicament. Such a medicament is generally useful for improving feed utilization efficiency or increasing the rate of weight gain in an animal. Other uses of such a medicament generally include, for example, increasing carcass leanness.

25

[14] This invention also is directed, in part, to a composition. The composition comprises zilpaterol (or a salt thereof) and melengestrol acetate.

[15] This invention also is directed, in part, to an animal feedstuff for improving the efficiency of feed utilization or increasing the rate of weight gain in an animal. The feedstuff comprises a composition, which, in turn, comprises zilpaterol (or a salt thereof) and melengestrol acetate. The feedstuff also comprises at least one carrier material.

30

[16] This invention also is directed, in part, to kits for administering zilpaterol (or a salt thereof) and melengestrol acetate to an animal.

[17] In some embodiments directed to kits, the kit comprises a zilpaterol dosage form comprising zilpaterol (or a salt thereof). In some such embodiments, the kit also comprises at least one of the following: (a) a melengestrol acetate dosage form comprising melengestrol acetate, (b) instructions for administering zilpaterol (or a salt thereof) and melengestrol acetate to the animal, and (c) instructions for mixing at least a portion of the zilpaterol dosage form with melengestrol acetate.

[18] In some embodiments directed to kits, the kit comprises a melengestrol acetate dosage form comprising melengestrol acetate. In some such embodiments, the kit also comprises at least one of the following: (a) a zilpaterol dosage form comprising zilpaterol (or a salt thereof), (b) instructions for administering melengestrol acetate and zilpaterol (or a salt thereof) to the animal, and (c) instructions for mixing at least a portion of the melengestrol acetate dosage form with zilpaterol (or a salt thereof).

[19] Further benefits of Applicants' invention will be apparent to one skilled in the art from reading this specification.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[20] This detailed description of preferred embodiments is intended only to acquaint others skilled in the art with Applicants' invention, its principles, and its practical application so that others skilled in the art may adapt and apply the invention in its numerous forms, as they may be best suited to the requirements of a particular use. This detailed description and its specific examples, while indicating preferred embodiments of this invention, are intended for purposes of illustration only. This invention, therefore, is not limited to the preferred embodiments described in this specification, and may be variously modified.

[21] It has been discovered that the combination of zilpaterol (or a salt thereof, such as zilpaterol hydrochloride) and melengestrol acetate can be particularly useful for increasing the rate of weight gain and improving feed efficiency (*i.e.*, decrease the amount of feed per amount of weight gain) in animals. Other uses include, for example, increasing carcass leanness (*i.e.*, increasing protein content in carcass soft tissue).

[22] A salt of zilpaterol may be advantageous in the combination due to one or more of the salt's physical properties, such as pharmaceutical stability in differing temperatures and humidities; crystalline properties; and/or a desirable solubility in

water, oil, or other solvent. Any such salt preferably is pharmaceutically acceptable.

Suitable zilpaterol salts generally include acid addition salts. An acid addition salt typically can be prepared by reacting free zilpaterol base with an approximately stoichiometric amount of an inorganic or organic acid. Examples of often suitable

- 5 inorganic acids include hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric, and phosphoric acid. Examples of often suitable organic acids generally include, for example, aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic, and sulfonic classes of organic acids. Specific examples of often suitable organic acids include cholate, sorbate, laurate, acetate, trifluoroacetate, formate, 10 propionate, succinate, glycolate, gluconate, digluconate, lactate, malate, tartaric acid, citrate, ascorbate, glucuronate, maleate, fumarate, pyruvate, aspartate, glutamate, aryl carboxylic acid (*e.g.*, benzoate), anthranilic acid, mesylate, stearate, salicylate, *p*-hydroxybenzoate, phenylacetate, mandelate, embonate (pamoate), alkylsulfonate (*e.g.*, ethanesulfonate), arylsulfonate (*e.g.*, benzenesulfonate), pantothenate, 2- 15 hydroxyethanesulfonate, sulfanilate, cyclohexylaminosulfonate, algenic acid, β -hydroxybutyric acid, galactarate, galacturonate, adipate, alginate, butyrate, camphorate, camphorsulfonate, cyclopentanepropionate, dodecylsulfate, glycoheptanoate, glycerophosphate, heptanoate, hexanoate, nicotinate, 2-naphthalesulfonate, oxalate, palmoate, pectinate, 3-phenylpropionate, picrate, pivalate, 20 thiocyanate, tosylate, and undecanoate. In some embodiments, the salt of zilpaterol comprises a hydrochloric acid salt.

[23] Included within the scope of the compounds and salts used in the combinations of this invention are any stereoisomers, tautomers, and mixtures thereof.

- [24] The combination of this invention may generally be used, for example, 25 to increase the rate of weight gain, improve feed efficiency, and/or increase carcass leanness in animals raised for food. These include, for example, livestock (*e.g.*, bovine and other ruminant animals, swine animals, etc.), poultry (*e.g.*, chickens, turkeys, etc.), and/or fish (*e.g.*, salmon, trout, catfish, etc.). The dosing regimen of the zilpaterol (or salt thereof) and melengestrol acetate is preferably chosen to maximize 30 one or more beneficial effects of the combination.

[25] In some embodiments, the combination of this invention is orally administered. Suitable oral dosage forms may include, for example, solid dosage forms (*e.g.*, tablets, hard or soft capsules, granules, powders, etc.), pastes, and liquid dosage forms (*e.g.*, solutions, suspensions, syrups, etc.).

[26] In some embodiments, at least one of the zilpaterol (or salt thereof) and melengestrol acetate is incorporated into the intended recipient animal's drinking water.

[27] In some embodiments, at least one (and often both) of the zilpaterol (or salt thereof) and melengestrol acetate is incorporated into the intended recipient animal's feed. The zilpaterol (or salt thereof) and/or melengestrol acetate may be mixed with the feed directly or as part of a premix. Incorporating both the zilpaterol (or salt thereof) and melengestrol acetate into the feed is often particularly preferable for cattle or swine in a feedlot.

[28] In some embodiments, the zilpaterol (or salt thereof) and melengestrol acetate are mixed together or separately with one or more carrier materials. A suitable carrier material may be, for example, a normal daily feed. Alternative (or additional) carrier materials may include one or more excipients such as, for example, sweetening agents, flavoring agents, coloring agents, preservative agents, inert diluents (*e.g.*, calcium carbonate, sodium carbonate, lactose, calcium phosphate, sodium phosphate, or kaolin), granulating and disintegrating agents (*e.g.*, corn starch or alginic acid), binding agents (*e.g.*, gelatin, acacia, or carboxymethyl cellulose), lubricating agents (*e.g.*, magnesium stearate, stearic acid, or talc), wetting or dispersing agents, etc. Liquid compositions will generally comprise at least one solvent. The solvent(s) typically has sufficient chemical properties and is in a sufficient quantity to keep the zilpaterol (or salt thereof) and/or melengestrol acetate solubilized at normal storage and usage temperatures. In some instances, a liquid composition will comprise at least one preservative. The presence of a preservative may, for example, allow for the composition to be stored over a greater amount of time.

[29] In some embodiments, the zilpaterol (or salt thereof) and/or melengestrol acetate is in the form of particles adhered to a support, which, in turn, is fed to the intended recipient animal by, for example, being introduced into the intended recipient animal's feed, either directly or as part of a premix. Contemplated supports include, for example, inert supports, such as calcium carbonate, limestone, oyster shell flour, talc, soybean hulls, soybean meal, soybean feed, soybean mill run, wheat middling, rice hulls, corn meal, corn germ meal, corn gluten, starch, sucrose, and lactose. Particularly contemplated supports include corn cob supports, such as the support discussed in U.S. Patent 5,731,028.

[30] In some embodiments in which one or more of the active ingredients are supported, the active ingredient particles adhered to the support have a particle size that is less than the size of the support. In some embodiments employing a corn cob support, for example, the size of the support is from about 300 to about 800 μm .

5 In some such embodiments, the active ingredient particles (or at least about 95% of the active agent particles) are less than about 250 μm . In some such embodiments, the size of the majority of the active ingredient particles is from about 50 to about 200 μm . To avoid generating dust when making the supported composition, it is generally preferable to avoid using extremely small active ingredient particles. In some
10 embodiments, for example, size distribution of the active ingredient particles is such that less than about 5% of the particles have a particle size of less than about 15 μm .

[31] To the extent the combination is incorporated into feed, the feed mixture will vary depending on, for example, the type (*e.g.*, species and breed), age, weight, activity, and condition of the intended recipient. With bovine and swine
15 animals, for example, various feeds are well known in the art, and often comprise cereals; sugars; grains; arachidic, tournsole, and soybean press cake; flours of animal origin, such as fish flour; amino acids; mineral salts; vitamins; antioxidants; etc. In general, the zilpaterol (or salt thereof) and melengestrol acetate can be incorporated into any feed that is available and used for the intended recipient animal.

20 [32] It is contemplated that the zilpaterol (or salt thereof) and/or melengestrol acetate may be administered via non-oral routes, such as rectally, via inhalation (*e.g.*, via a mist or aerosol), transdermally (*e.g.*, via a transdermal patch), or parenterally (*e.g.*, subcutaneous injection, intravenous injection, intramuscular injection, implanted device, partially implanted device etc.). It is contemplated, for
25 example, that at least one of the zilpaterol (or salt thereof) and melengestrol acetate may be administered via an implant, such as a subcutaneous implant. For administration to bovine or swine animals, for example, the zilpaterol (or salt thereof) and/or melengestrol acetate may be administered in the form of an implant behind the ear. If the zilpaterol (or salt thereof) and/or the melengestrol acetate is/are
30 administered parenterally via an injection, the concentration of the active agent(s) in the dosage form preferably is sufficient to provide the desired amount of the active agent(s) in a volume that is acceptable for parenteral administration.

[33] Factors affecting the preferred dosage regimen may include, for example, the type (*e.g.*, species and breed), age, size, diet, activity, and condition of

the intended recipient; the type of administration used (*e.g.*, oral via feed, oral via drinking water, subcutaneous implant, other parenteral route, etc.); pharmacological considerations, such as the activity, efficacy, pharmacokinetic, and toxicology profiles of the particular composition administered; and whether the zilpaterol (or its salt) and melengestrol acetate are being administered as part of a combination with additional active ingredients. Thus, the preferred amounts of the zilpaterol (or salt thereof) and melengestrol acetate can vary, and, therefore, can deviate from the typical dosages set forth above.

[34] It is contemplated that one or both of the zilpaterol (or salt thereof) and melengestrol acetate may be administered once. In general, however, the combination is administered over time, and the period of time over which the zilpaterol (or salt thereof) is administered overlaps (at least partially) with the period of time over which the melengestrol acetate is administered. In some embodiments, the zilpaterol (or salt thereof) and melengestrol acetate are co-administered daily over a period of time in a substantially simultaneous manner, such as, for example, (a) in a single formulation (*e.g.*, in the same tablet, granule, or powder) having a fixed ratio(s) of the zilpaterol (or salt thereof) to the melengestrol acetate, and/or (b) in separate formulations. The zilpaterol (or salt thereof) and melengestrol acetate may alternatively (or additionally) be administered at different intervals over a period of time.

[35] In some embodiments (*e.g.*, where the animal recipient is a livestock animal), the zilpaterol (or a salt thereof) is administered approximately daily for at least about 2 days, more typically daily for from about 10 to about 60 days, and still more typically, from about 14 to about 42 days. In some embodiments (*e.g.*, where the animal is a bovine animal), the zilpaterol (or a salt thereof) is administered approximately daily for from about 20 to about 40 days. In some such embodiments, for example, the composition is administered approximately daily for from about the last 20 to about the last 40 days of the finishing period. The term "finishing period" refers to the later stage of the growing period for an animal. During this period, the recipient animal is typically confined (*e.g.*, in a feedlot for livestock animals). In some embodiments where the animal is a bovine animal, this period lasts for from about 90 to about 225 days, and depends on, for example, the starting body weight of the animal. There is typically a withdrawal period following the finishing period in which no zilpaterol (or salt thereof) is administered. The length of this withdrawal period may depend on, for example, the type (*e.g.*, species and breed), age, weight,

activity, and condition of the recipient animal, as well as the maximum acceptable residue concentration in the meat of the animal.

[36] In general, the melengestrol acetate is administered to the intended animal recipient daily during at least a portion of the period (and typically the entire period) that the zilpaterol (or a salt thereof) is administered. The melengestrol acetate also is typically administered for a period before the zilpaterol (or zilpaterol salt) is administered, and, in some instances, for at least a portion of the zilpaterol (or zilpaterol salt) withdrawal period afterward. In some embodiments, for example, the melengestrol acetate is administered approximately daily during the entire time that the animal is being fed in confinement for harvest, including before, during, and after the time that the zilpaterol (or salt thereof) is administered.

[37] Although single daily doses are typically preferred, it is contemplated that shorter or longer periods between doses can be used, depending on, for example, the recipient's metabolism of the zilpaterol (or its salt) and the melengestrol acetate. It is contemplated that smaller doses may be administered two or more times per day to achieve the desired total daily dose. Such multiple doses per day may, in some instances, be used to increase the total oral daily dose, if desired.

[38] In accordance with this invention, the zilpaterol (or salt thereof) and melengestrol acetate are generally used in amounts that, when combined, produce an unexpected benefit. It is believed, for example, that this combination provides an unexpected benefit with respect to improvement of feed utilization efficiency and/or increasing the rate of weight gain. Other benefits include, for example, increases in carcass leanness.

[39] In some embodiments, the amount of the zilpaterol (or salt thereof) and the amount of melengestrol acetate are, when combined, sufficient to improve the animal's average daily feed utilization efficiency over a period of at least about 3 days (or at least about 7 days, at least about 10 days, at least about 20 days, at least about 30 days, or at least about 40 days) relative to the average daily feed utilization efficiency that would be realized by administering the amount of the zilpaterol (or salt thereof) without any melengestrol acetate over the same amount of time under otherwise substantially identical feeding conditions.

[40] In some embodiments, the amount of the zilpaterol (or salt thereof) and the amount of melengestrol acetate are, when combined, sufficient to increase the animal's average daily weight gain over a period of at least about 3 days (or at least

about 7 days, at least about 10 days, at least about 20 days, at least about 30 days, or at least about 40 days) by at least about 100% (or at least about 125%, at least about 150%, at least about 175%, or at least about 200%) the average daily weight gain that is realized during the week immediately before administration of the zilpaterol (or salt thereof) and melengestrol acetate under otherwise substantially identical feeding conditions.

[41] In some embodiments, the amount of the zilpaterol (or salt thereof) and the amount of melengestrol acetate are, when combined, sufficient to increase the animal's average daily weight gain over a period of at least about 3 days (or at least about 7 days, at least about 10 days, at least about 20 days, at least about 30 days, or at least about 40 days) relative to the average daily weight gain that would be realized by administering the amount of the zilpaterol (or salt thereof) without any melengestrol acetate over the same amount of time under otherwise substantially identical feeding conditions. In some such embodiments, for example, the amount of the zilpaterol (or salt thereof) and the amount of melengestrol acetate are, when combined, sufficient to increase the animal's average daily weight gain over a period of at least about 3 days (or at least about 7 days, at least about 10 days, at least about 20 days, at least about 30 days, or at least about 40 days) by at least about 1.2 times (or at least about 1.5 times, at least about 1.7 times, at least about 2.0 times, or at least about 2.1 times) the average daily weight gain that would be realized by administering the amount of the zilpaterol (or salt thereof) without any melengestrol acetate over the same amount of time under otherwise substantially identical feeding conditions.

[42] The total daily dose of the zilpaterol (or a salt thereof) is typically at least about 1 mg/animal, particularly in the context of, for example, livestock. In some embodiments, for example, the daily dose is less than about 900 mg/animal. In some such embodiments, the daily dose is from about 1 to about 200 mg/animal, from about 20 to about 150 mg/animal, from about 50 to about 100 mg/animal, or from about 60 to about 90 mg/animal. In some embodiments, greater than about 0.01 mg/kg (*i.e.*, milligram of zilpaterol (or salt thereof) per kilogram body weight) is fed daily during the treatment period, particularly for bovine and swine animals. In some such embodiments, the daily dose is from about 0.01 to about 15 mg/kg, from about 0.01 to about 10 mg/kg, from about 0.05 to about 2 mg/kg, from about 0.1 to about 1, or from about 0.1 to about 0.3 mg/kg. To illustrate, in some embodiments for heifers, the daily dose is from about 0.1 to about 0.2 mg/kg (*e.g.*, about 0.15 mg/kg daily).

Illustrating further, in some embodiments for swine animals, the total daily dose is from about 0.13 to about 0.27 mg/kg.

[43] In some embodiments where the zilpaterol (or salt thereof) is administered in the recipient animal's feed, the concentration of the zilpaterol (or salt thereof) in the feed is at least about 0.01 ppm (by weight). In some embodiments for bovine animals, the concentration of zilpaterol (or a salt thereof) is no greater than about 75 ppm (by weight). In some such embodiments, for example, the concentration of zilpaterol (or its salt) is no greater than about 38 ppm. In other embodiments, the concentration (or its salt) is from about 0.5 to about 20 ppm, from about 3 to about 8 ppm, or from about 3.7 to about 7.5 ppm (by weight). In some embodiments for swine animals, the concentration of zilpaterol (or a salt thereof) is no greater than about 45 ppm (by weight). In some such embodiments, for example, the concentration is no greater than about 23 ppm. In other embodiments, the concentration of zilpaterol (or its salt) is from about 0.5 to about 20 ppm, from about 2 to about 10 ppm, or from about 4 to about 8 ppm (by weight). The zilpaterol and zilpaterol salt concentrations in this paragraph are all based on a feed containing approximately 90% dry matter.

[44] In general, the total daily dose of melengestrol acetate is at least about 0.001 mg/animal, particularly in the context of, for example, livestock. In some such embodiments, for example, the daily dose is less than about 5 mg/animal. To illustrate, in some embodiments, the daily dose is from about 0.01 to about 2 mg/animal, from about 0.05 to about 1 mg/animal, from about 0.1 to about 1 mg/animal, from about 0.25 to about 0.5 mg/animal, or from about 0.25 to about 0.4 mg/animal. In some embodiments, the daily dose is greater than about 0.0001 mg/kg (*i.e.*, milligram of melengestrol acetate per kilogram body weight). In some embodiments directed to bovine animals, the daily dose is from about 0.0005 mg/kg to about 0.001 mg/kg. In some embodiments directed to swine animals, the daily dose is from about 0.001 to about 0.008 mg/kg.

[45] In some embodiments where the melengestrol acetate is administered in the recipient animal's feed, the concentration of the melengestrol acetate in the feed is at least about 0.001 ppm (by weight). In some embodiments, the melengestrol acetate concentration is from about 0.002 to about 0.6 ppm, from about 0.002 to about 0.3 ppm, from about 0.002 to about 1 ppm, from about 0.002 to about 0.6 ppm, or from about 0.01 to about 0.1 ppm (by weight). In some such embodiments,

melengestrol acetate is administered in the feed at a concentration of from about 0.02 to about 0.06 ppm (by weight). The melengestrol acetate concentrations in this paragraph are all based on a feed containing approximately 90% dry matter.


[46] In some embodiments, the zilpaterol (or salt thereof) and melengestrol acetate are administered at a mass ratio of the zilpaterol (or salt thereof) to melengestrol acetate of from about 10 to about 1000, from about 50 to about 500, from about 100 to about 400, from about 120 to about 360, or from about 130 to about 170. In some embodiments, this mass ratio preferably is maintained approximately daily for at least about 2 days, and typically from about 10 to about 60 days, or from about 14 to about 42 days. In some embodiments (*e.g.*, where the animal is a bovine animal), the mass ratio is maintained approximately daily for from about 20 to about 40 days. In some such embodiments, for example, the mass ratio is maintained approximately daily for from about the last 20 to about the last 40 days of the finishing period. In some embodiments, this mass ratio is present in a composition (*e.g.*, feed) that is administered daily to the recipient animal.

[47] In some embodiments, the combination of this invention is administered in combination with one or more further active ingredients. It is contemplated that the additional active ingredient(s) may be administered once. In general, however, the additional active ingredient(s) is administered over time. In such instances, the additional active ingredient(s) may be administered substantially simultaneously with one or both of the zilpaterol (or salt thereof) and melengestrol acetate and/or at different intervals. To the extent the administration is simultaneous, the combined actives may be part of the same dosage formulation (*e.g.*, in the same tablet, granule, or powder) and/or separate formulations.

[48] In some embodiments (*e.g.*, for livestock, such as bovine or swine), the combination of this invention is administered with one or more steroids in addition to the melengestrol acetate.


[49] In some embodiments, the combination of this invention is administered as part of a dosing scheme with a steroid corresponding in structure to

Formula (IV):



(IV).

In **Formula (IV)**, X may be, for example, hydrogen; optionally unsaturated alkyl of from 1 to about 6 carbon atoms, wherein one of the carbon atoms optionally is replaced by -O-; or an acyl of an organic carboxylic acid of from 1 to about 18 carbon atoms. In some such embodiments, for example, X is -C(O)CH₃ (*i.e.*, the steroid comprises trenbolone acetate, also known as “17β-acetoxy-Δ^{4,9,11}-estratriene-3-one”). That compound corresponds in structure to **Formula (V)**:



(V).

Trenbolone acetate is marketed by Intervet Inc., a part of Schering-Plough Corporation, under the trademarks Finaplix[®] and Revalor[®]. For livestock in particular, trenbolone acetate may generally be administered parenterally, such as via injection or a subcutaneous implant. The implant may be in the form of a single or, more typically, multiple units (*e.g.*, pellets). Suitable locations for a subcutaneous implant generally include, for example, behind the ear. In some embodiments, the implant is implanted for from about 20 days to about 8 months before harvest, or from about 2 to about 7 months before harvest. The total trenbolone acetate implant dosage is typically less than about 4000 mg/animal. In some embodiments, the total trenbolone acetate implant dosage is from about 50 to about 2000 mg/animal, or from about 80 to about 500 mg/animal. In some embodiments, the preferred daily trenbolone acetate dosage is from about 0.75 to about 1.2 mg/animal (*e.g.*, about 0.95 mg/animal daily). In some embodiments, implants are implanted sequentially to administer the desirable amount of trenbolone acetate over time. In some embodiments, immediate- and delayed-release implant units (*e.g.*, pellets) are implanted at the same time to administer a desirable amount of trenbolone acetate over time. The timing of the initial implant and any subsequent implants, as well as the amount of trenbolone

acetate in the implant, will depend, for example, the species of the recipient animal, the size of the recipient animal, and the formulation(s) of the particular implant(s) used. Further discussion related to suitable dosing regimens for trenbolone acetate may be found in, for example, US Patents 4,900,735 and 6,498,153 (now US Patent RE 39,592 E). Both those patents are incorporated by reference into this patent.

[50] In some embodiments, the combination of this invention also (or alternatively) may be administered as part of a dosing scheme with zeranol, estradiol, and/or estradiol benzoate. It is contemplated that the zeranol, estradiol, or estradiol benzoate may be administered orally in the feed. Typically, however, zeranol, estradiol, or estradiol benzoate is administered parenterally, such as via injection or a subcutaneous implant. This is particularly true for livestock. Suitable locations for a subcutaneous implant generally include, for example, behind the ear. Zeranol is marketed under the trade name Ralgro[®] by Schering-Plough Corporation. In some embodiments, the implant contains other active ingredients in addition to the zeranol, estradiol, or estradiol benzoate, such as, for example, trenbolone acetate. In some embodiments, the implant is implanted for from about 20 days to about 8 months before harvest, from about 20 days to about 4 months, or from about 1 to about 3 months before harvest. In some embodiments, the implant is implanted at least 40 days before harvest. The total zeranol implant dosage is typically less than about 1500 mg. In some embodiments, the total zeranol implant dosage is from about 10 to about 150 mg. In some such embodiments, the total zeranol implant dosage is from about 30 to about 140 mg. These embodiments may be particularly useful when the recipient animal is a bovine animal. In other embodiments, the total zeranol implant dosage is from about 10 to about 50. These embodiments may be particularly useful for smaller animal recipients, such as a sheep or recently-weaned calf. The typical implant dosage for estradiol or estradiol benzoate is typically less than about 400 mg. In some embodiments, the implant dosage for estradiol or estradiol benzoate is from about 0.05 to about 50 mg. In some embodiments, implants are implanted sequentially to administer the desirable amount of zeranol, estradiol, or estradiol benzoate over time. In some embodiments, immediate- and delayed-release implant units (e.g., pellets) are implanted at the same time to administer a desirable amount of zeranol, estradiol, or estradiol benzoate over time. The timing of the implant and any subsequent implants, as well as the amount of zeranol, estradiol, or estradiol benzoate in the implant, will depend on, for example, the species of the recipient animal, the

size of the recipient animal, and the formulation(s) of the particular implant(s) used.

Further discussion related to suitable dosing regimens for zeranol and estradiol may be found in, for example, US Patents 4,900,735; and still further discussion related to suitable dosing regimens for estradiol may be found in, for example, 6,498,153 (now
5 US Patent RE 39,592 E).

[51] The combination of this invention also (or alternatively) may be administered as part of a dosing scheme with one or more ionophores. Suitable ionophores include, for example, monensin, lasalocid, laidlomycin propionate, bambermycin, and salts thereof. Sodium monensin, for example, is marketed under
10 the trade name Rumensin® by Elanco Animal Health, and reportedly effective for use in cattle fed in confinement for harvest to prevent and control of coccidiosis due to *Eimeria bovis* and *Eimeria zuernii*. When administered properly, ionophores can, for example, be effective for improving feed utilization efficiency and/or increasing the rate of weight gain. The typical daily dosage for an ionophore will vary, depending
15 on the particular ionophore, route of administration, species of the animal recipient, size of the animal recipient, etc. When, for example, the ionophore is monensin sodium and is administered in feed for cattle, a suitable daily dose is typically less than about 5000 mg/animal. In some such embodiments, for example, the daily dose is from about 10 to about 500 mg/animal, from about 50 to about 480 mg/animal,
20 from about 50 to about 360 mg/animal, or from about 60 to about 350 mg/animal. In other embodiments, the daily dose is less than about 4.2 mg/lb body weight (9.2 mg/kg body weight). In some such embodiments, the daily dose is from about 0.14 to about 0.42 mg/lb body weight (0.31 to about 0.92 mg/kg body weight). Illustrating further, when the ionophore is lasalocid sodium and is administered in feed for cattle,
25 a suitable daily dose is again typically less than about 4000 mg/animal. In some such embodiments, for example, the daily dose is from about 10 to about 400 mg/animal, or from about 100 to about 360 mg/animal.

[52] The combination of this invention also (or alternatively) may be administered as part of a dosing scheme with one or more antibiotics. Suitable
30 antibiotics include, for example, macrolide antibiotics, such as tylosin and salts thereof. Tylosin phosphate and tylosin tartrate, for example, are marketed under the trade name Tylan® by Elanco Animal Health. When administered properly, antibiotics can, for example, be effective for increasing the rate of weight gain, improving feed efficiency, and/or reducing the time for carcass trimming. It is

believed that these effects stem, at least in part, from a reduction of opportunistic bacteria (*e.g.*, *Fusobacterium necrophorum* and *Actinomyces pyogenes*) that can infect the liver. The typical daily dosage for an antibiotic will vary, depending on the antibiotic, route of administration, species of the animal recipient, size of the animal recipient, etc. When, for example, the antibiotic is tylosin phosphate and is administered in feed for cattle, a suitable daily dose is typically less than about 900 mg. In some such embodiments, the daily dose is from about 10 to about 150 mg/animal, from about 50 to about 150 mg/animal, or from about 60 to about 90 mg/animal.

[53] This invention also is directed to kits that are, for example, suitable for use in performing the treatment methods described above.

[54] In some embodiments, the kit comprises a zilpaterol dosage form comprising zilpaterol (or a salt thereof, such as zilpaterol hydrochloride). In some such embodiments, the kit also comprises, for example, at least one of the following:

(a) a melengestrol acetate dosage form comprising melengestrol acetate, (b) instructions for administering zilpaterol (or a salt thereof) and melengestrol acetate to the animal, and (c) instructions for mixing at least a portion of the zilpaterol dosage form with melengestrol acetate. The zilpaterol dosage form may additionally comprise one or more additional components, such as, for example, one or more carrier materials and/or other active ingredients.

[55] In some embodiments, the kit comprises a melengestrol acetate dosage form comprising melengestrol acetate. In some such embodiments, the kit also comprises, for example, at least one of the following: (a) a zilpaterol dosage form comprising zilpaterol (or a salt thereof), (b) instructions for administering melengestrol acetate and zilpaterol (or a salt thereof) to the animal, and (c) instructions for mixing at least a portion of the melengestrol acetate dosage form with zilpaterol (or a salt thereof). The melengestrol acetate dosage form may additionally comprise one or more additional components, such as, for example, one or more carrier materials and/or other active ingredients.

[56] In some embodiments, the kits of this invention comprise instructions for orally administering at least a portion (or all) of the zilpaterol (or salt thereof) and/or melengestrol acetate to the intended recipient animal. In some such embodiments, at least a portion (or all) of the zilpaterol (or salt thereof) and

melengestrol acetate are in the same dosage form. In other embodiments, the zilpaterol (or salt thereof) and melengestrol acetate are in separate dosage forms.

[57] In some embodiments, the kits comprise other components, such as, for example, an apparatus (*e.g.*, a syringe) to administer one or more active ingredients, and/or an apparatus to combine one or more active ingredients and/or carrier materials with one or more other active ingredients and/or carrier materials.

EXAMPLE

[58] This example is merely illustrative, and not intended to be limiting to the remainder of this disclosure in any way.

[59] Angus heifers and Angus cross-brad steers with a starting weight of from 686 lb (311 kg) to 934 lb (424 kg) per animal were randomized to the following treatment groups, as shown in **Table 1**:

- 1) Non-medicated, negative control (steers and heifers)
- 2) Zilmax[®], Rumensin[®], and Tylan[®] (steers and heifers). The Type C feed was formulated to contain 6.8 g/ton (7.5 mg/kg) zilpaterol hydrochloride, 40 g/ton (44 mg/kg) monensin sodium, and 10 g/ton (11 mg/kg) tylosin phosphate on a 90% dry mass basis ("DMB").
- 3) Zilmax[®], Rumensin[®], Tylan[®], and MGA[®] (heifers only). The Type C feed was formulated to contain 6.8 g/ton (7.5 mg/kg) zilpaterol hydrochloride, 40 g/ton (44 mg/kg) monensin sodium, 10 g/ton (11 mg/kg) tylosin phosphate on a 90% DMB; and melengestrol acetate at a level to deliver 0.50 mg melengestrol acetate in 22.5 lb (10.2 kg) of Type C feed.

Table 1
Treatment Groups

Treatment groups	Treatment period (days zero thru 14)	Withdrawal period (days 15 thru 17)
1 (2 steers and 2 heifers)	Non-medicated Type C feed	Non-medicated Type C feed
2 (4 steers and 4 heifers)	Type C feed containing Zilmax [®] , Rumensin [®] , & Tylan [®]	Type C feed containing Rumensin [®] & Tylan [®]
3 (8 heifers)	Type C feed containing Zilmax [®] , Rumensin [®] , Tylan [®] , & MGA [®]	Type C feed containing Rumensin [®] , Tylan [®] , & MGA [®]

The four test articles were administered orally in complete, pelleted Type C feeds with the composition shown in **Table 2**:

5

Table 2
Type C Feed Composition

Ingredient name	Percent As-Fed Basis
Corn (ground) (Bulk)	68.25
Cobs Ground ¼(Bed-O-Cobs)	10.44
Soybean Meal 48% (Bulk)	4.97
PM 42 Kane Lass (50#)	4.70
Ground Steam Rolled Oat Groats (50#)	4.40
Dehy Alfalfa 17% Meal (bulk)	2.20
Beef finisher base(50#)	2.00
Fat liquid (Bulk)	2.00
Calcium Carbonate 38% Ca (bulk)	0.48
Dical 18.5% P (bulk)	0.31
Urea (50#)	0.25
Total	100.00

For the treatment groups (*i.e.*, Groups 2 and 3), the active ingredients were mixed with the Type C feed. The non-medicated and medicated feeds were prepared at a commercial feed manufacturing facility using typical feed manufacturing procedures. The feeds were formulated to meet or exceed 1996 NRC recommendations for finishing beef cattle.

A. Body Weight Gain

[60] As shown in **Table 3** below, the average daily weight gains of the study animals increased by 9%, 92%, and 202% in Treatment Groups 1 (non-medicated), 2 (Zilmax[®], Rumensin[®], and Tylan[®]), and 3 (Zilmax[®], Rumensin[®], Tylan[®], and MGA[®]), respectively, during the treatment period compared to gains observed in the pre-treatment, non-medicated feeding period.

20

Table 3
Individual Weight Gains

Treatment group	Sex	Pre-treatment (non-medicated) average daily gain (per head)	Treatment (medicated) average daily gain (per head)	Change in average daily gain with medicated feeding*
1	Steer	3.03 lb (1.38 kg)	4.67 lb (2.12 kg)	54%

Treatment group	Sex	Pre-treatment (non-medicated) average daily gain (per head)	Treatment (medicated) average daily gain (per head)	Change in average daily gain with medicated feeding*
1	Steer	2.76 lb (1.25 kg)	1.14 lb (0.52 kg)	-59%
1	Heifer	1.90 lb (0.86 kg)	1.19 lb (0.54 kg)	-37%
1	Heifer	1.12 lb (0.51 kg)	2.62 lb (1.19 kg)	134%
2	Steer	2.83 lb (1.28 kg)	5.10 lb (2.32 kg)	80%
2	Steer	2.62 lb (1.19 kg)	5.14 lb (2.33 kg)	96%
2	Steer	2.21 lb (1.00 kg)	4.38 lb (1.99 kg)	99%
2	Steer	3.10 lb (1.41 kg)	4.90 lb (2.22 kg)	58%
2	Heifer	2.16 lb (0.98 kg)	3.33 lb (1.51 kg)	55%
2	Heifer	1.81 lb (0.82 kg)	3.33 lb (1.51 kg)	84%
2	Heifer	1.57 lb (0.71 kg)	5.00 lb (2.27 kg)	219%
2	Heifer	1.21 lb (0.55 kg)	2.38 lb (1.08 kg)	97%
3	Heifer	1.60 lb (0.73 kg)	3.57 lb (1.62 kg)	123%
3	Heifer	1.64 lb (0.74 kg)	2.62 lb (1.19 kg)	60%
3	Heifer	1.21 lb (0.55 kg)	4.05 lb (1.84 kg)	235%
3	Heifer	2.16 lb (0.98 kg)	5.00 lb (2.27 kg)	132%
3	Heifer	1.38 lb (0.63 kg)	4.76 lb (2.16 kg)	245%
3	Heifer	0.90 lb (0.41 kg)	5.48 lb (2.49 kg)	511%
3	Heifer	1.03 lb (0.47 kg)	4.29 lb (1.95 kg)	314%
3	Heifer	1.03 lb (0.47 kg)	3.33 lb (1.51 kg)	222%

*Change in average daily weight gain in treatment (medicated) period compared to pre-treatment (non/medicated) period.

B. Type C Feed Consumption

[61] Average daily non-medicated consumption during the 7 days preceding the initiation of medicated feeding was calculated by pen (See Table 4). During this period, the heifers and steers were offered 22.5 lb (10.2 kg) and 25.0 lb (11.4 kg), respectively, of as-fed feed per animal daily. Based on an average dry matter content of 89.08% for the non-medicated feed offered during this period, these amounts are equivalent to 22.27 lb (10.11 kg) and 24.74 lb (11.23 kg), respectively, of 90% DMB from feed offered. By subtracting unconsumed feed corrected to a 90% DMB from feed offered, the respective average daily feed consumption of the heifers and steers during this 7-day period was 21.47 lb (9.75 kg) and 20.52 lb (9.32 kg), respectively, of 90% dry matter feed per animal daily.

Table 4
Non-Medicated Feed Consumption for
7 Days Before Medicated Feeding Calculated on 90% Dry Matter Basis

Day	Heifers, 4 head/pen					Steers, 5 head/pen	
	Pen 2	Pen 3	Pen 4	Pen 5	Pen 6	Pen 8	Pen 9
Total of 90% Dry Matter Feed Consumed							
1	85.1 lb (38.6 kg)	85.1 lb (38.6 kg)	85.1 lb (38.6 kg)	85.1 lb (38.6 kg)	80.2 lb (36.4 kg)	118.8 lb (53.9 kg)	85.5 lb (38.8 kg)
2	85.1 lb (38.6 kg)	85.1 lb (38.6 kg)	85.1 lb (38.6 kg)	85.1 lb (38.6 kg)	80.2 lb (36.4 kg)	118.8 lb (53.9 kg)	97.9 lb (44.4 kg)
3	87.1 lb (39.5 kg)	87.1 lb (39.5 kg)	87.1 lb (39.5 kg)	87.1 lb (39.5 kg)	82.2 lb (37.3 kg)	94.4 lb (42.8 kg)	98.3 lb (44.6 kg)
4	87.1 lb (39.5 kg)	87.1 lb (39.5 kg)	87.1 lb (39.5 kg)	87.1 lb (39.5 kg)	82.2 lb (37.3 kg)	107.2 lb (48.7 kg)	95.7 lb (43.4 kg)
5	89.1 lb (40.4 kg)	89.1 lb (40.4 kg)	89.1 lb (40.4 kg)	89.1 lb (40.4 kg)	84.1 lb (38.2 kg)	121.2 lb (55.0 kg)	59.9 lb (27.2 kg)
6	89.1 lb (40.4 kg)	89.1 lb (40.4 kg)	89.1 lb (40.4 kg)	89.1 lb (40.4 kg)	86.1 lb (39.1 kg)	121.2 lb (55.0 kg)	106.6 lb (48.4 kg)
7	86.0 lb (39.0 kg)	89.1 lb (40.4 kg)	84.6 lb (38.4 kg)	78.3 lb (35.5 kg)	82.1 lb (37.3 kg)	119.1 lb (54.1 kg)	91.4 lb (41.5 kg)
Total feed consumed in 7 days	608.6 lb (276.3 kg)	611.7 lb (277.7 kg)	607.2 lb (275.7 kg)	600.9 lb (272.8 kg)	576.9 lb (261.9 kg)	800.8 lb (363.6 kg)	635.3 lb (288.4 kg)
7-day average of daily 90% DMB feed consumed (per head) by pen	21.74 lb (9.87 kg)	21.85 lb (9.92 kg)	21.69 lb (9.85 kg)	21.46 lb (9.74 kg)	20.61 lb (9.36 kg)	22.88 lb (10.39 kg)	18.15 lb (8.24 kg)

[62] Average daily feed consumption by treatment group, pen, and sex also was calculated for the Zilmax[®] treatment (15 days) and withdrawal (3 days) periods of the study. The results are shown in Table 5 and 6. During these periods, the heifers and steers were again offered 22.5 lb (10.2 kg) and 25.0 lb (11.4 kg), respectively, of as-fed feed per animal each day. By using the average dry matter

percent of the Type C feeds offered during these periods, the amount of feed offered on a 90% DMB was calculated. Unconsumed feed corrected to a 90% DMB was then subtracted from feed offered to calculate average daily feed consumption on a 90% DMB. Consumptions for treatment groups 1, 2, and 3 during the combined Zilmax[®] treatment period and withdrawal periods were 22.92 lb (10.40 kg), 21.24 lb (9.64 kg), and 21.35 lb (9.69 kg), respectively, of feed (90% DMB) per animal daily.

Table 5
Daily Feed Consumption During the Zilmax[®] Treatment Period (15 days)
and Withdrawal Period (3 days) Calculated on a 90% Dry Matter Basis

	Steers (2 head)	Heifers (2 head)	Steers (4 head)	Heifers (4 head)	Heifers (4 head)	Heifers (4 head)
	Pen 2 Non-medicated control	Pen 3 Non-medicated control	Pen 5 Zilmax [®] , Rumensin [®] , & Tylan [®]	Pen 6 Zilmax [®] , Rumensin [®] , & Tylan [®]	Pen 8 Zilmax [®] , Rumensin [®] , Tylan [®] , & MGA [®]	Pen 9 Zilmax [®] , Rumensin [®] , Tylan [®] , & MGA [®]
Day	Total 90% Dry Matter Feed Consumed					
1	48.1 lb (21.8 kg)	44.5 lb (20.2 kg)	94.9 lb (43.1 kg)	89.0 lb (40.4 kg)	87.7 lb (39.8 kg)	89.1 lb (40.4 kg)
2	48.7 lb (22.1 kg)	44.5 lb (20.2 kg)	78.2 lb (35.5 kg)	89.5 lb (40.6 kg)	61.9 lb (28.1 kg)	89.1 lb (40.4 kg)
3	48.4 lb (22.0 kg)	44.5 lb (20.2 kg)	90.1 lb (40.9 kg)	89.5 lb (40.6 kg)	69.9 lb (31.7 kg)	88.2 lb (40.0 kg)
4	42.9 lb (19.5 kg)	44.5 lb (20.2 kg)	87.1 lb (39.5 kg)	88.7 lb (40.3 kg)	76.6 lb (34.8 kg)	86.0 lb (39.0 kg)
5	45.4 lb (20.6 kg)	43.9 lb (19.9 kg)	87.9 lb (39.9 kg)	84.8 lb (38.5 kg)	82.2 lb (37.3 kg)	83.2 lb (37.8 kg)
6	44.2 lb (20.1 kg)	44.2 lb (20.1 kg)	85.7 lb (38.9 kg)	85.0 lb (38.6 kg)	83.5 lb (37.9 kg)	86.3 lb (39.2 kg)
7	49.5 lb (22.5 kg)	44.5 lb (20.2 kg)	79.2 lb (36.0 kg)	81.7 lb (37.1 kg)	87.0 lb (39.5 kg)	88.7 lb (40.3 kg)
8	49.5 lb (22.5 kg)	44.5 lb (20.2 kg)	83.2 lb (37.8 kg)	83.1 lb (37.7 kg)	85.8 lb (39.0 kg)	89.1 lb (40.4 kg)
9	49.5 lb (22.5 kg)	44.5 lb (20.2 kg)	93.8 lb (42.6 kg)	71.6 lb (32.5 kg)	84.1 lb (38.2 kg)	89.1 lb (40.4 kg)
10	49.5 lb (22.5 kg)	41.1 lb (18.6 kg)	80.3 lb (36.4 kg)	72.9 lb (33.1 kg)	78.2 lb (35.5 kg)	89.1 lb (40.4 kg)
11	49.5 lb (22.5 kg)	44.5 lb (20.2 kg)	96.9 lb (44.0 kg)	79.6 lb (36.1 kg)	84.4 lb (38.3 kg)	89.1 lb (40.4 kg)
12	49.5 lb (22.5 kg)	44.5 lb (20.2 kg)	93.4 lb (42.4 kg)	60.4 lb (27.4 kg)	89.1 lb (40.4 kg)	89.1 lb (40.4 kg)
13	49.5 lb (22.5 kg)	44.5 lb (20.2 kg)	95.9 lb (43.5 kg)	63.7 lb (28.9 kg)	89.1 lb (40.4 kg)	89.1 lb (40.4 kg)
14	49.5 lb (22.5 kg)	44.5 lb (20.2 kg)	99.4 lb (45.1 kg)	87.0 lb (39.5 kg)	89.1 lb (40.4 kg)	89.1 lb (40.4 kg)
15	49.5 lb (22.5 kg)	44.5 lb (20.2 kg)	99.4 lb (45.1 kg)	88.5 lb (40.2 kg)	88.7 lb (40.3 kg)	89.1 lb (40.4 kg)
W1	48.9 lb (22.2 kg)	44.5 lb (20.2 kg)	88.0 lb (40.0 kg)	76.6 lb (34.8 kg)	86.7 lb (39.4 kg)	89.2 lb (40.5 kg)
W2	48.9 lb (22.2 kg)	44.5 lb (20.2 kg)	97.8 lb (44.4 kg)	78.1 lb (35.4 kg)	82.9 lb (37.6 kg)	88.4 lb (40.1 kg)

	Steers (2 head)	Heifers (2 head)	Steers (4 head)	Heifers (4 head)	Heifers (4 head)	Heifers (4 head)
	Pen 2 Non-medicated control	Pen 3 Non-medicated control	Pen 5 Zilmax [®] , Rumensin [®] , & Tylan [®]	Pen 6 Zilmax [®] , Rumensin [®] , & Tylan [®]	Pen 8 Zilmax [®] , Rumensin [®] , Tylan [®] , & MGA [®]	Pen 9 Zilmax [®] , Rumensin [®] , Tylan [®] , & MGA [®]
Day	Total 90% Dry Matter Feed Consumed					
W3	35.3 lb (16.0 kg)	41.4 lb (18.8 kg)	81.5 lb (37.0 kg)	76.2 lb (34.6 kg)	78.4 lb (35.6 kg)	88.2 lb (40.0 kg)

Table 6

Summary of Total Feed Consumption and Average Daily Feed Consumption During the Zilmax[®] Treatment Period and Withdrawal Period on a 90% DMB

	Steers (2 head)	Heifers (2 head)	Steers (4 head)	Heifers (4 head)	Heifers (4 head)	Heifers (4 head)
	Pen 2 Non-medicated control	Pen 3 Non-medicated control	Pen 5 Zilmax [®] , Rumensin [®] , & Tylan [®]	Pen 6 Zilmax [®] , Rumensin [®] , & Tylan [®]	Pen 8 Zilmax [®] , Rumensin [®] , Tylan [®] , & MGA [®]	Pen 9 Zilmax [®] , Rumensin [®] , Tylan [®] , & MGA [®]
Total feed consumed in treatment period	723.2 lb (328.3 kg)	663.7 lb (301.3 kg)	1345.4 lb (610.8 kg)	1215.1 lb (551.6 kg)	1237.3 lb (561.7 kg)	1323.4 lb (600.8 kg)
Total feed consumed in Withdrawal period	133.1 lb (60.4 kg)	130.4 lb (59.2 kg)	267.3 lb (121.4 kg)	230.9 lb (104.8 kg)	248.1 lb (112.6 kg)	265.8 lb (120.7 kg)
Total consumed (all 18 days)	856.3 lb (388.8 kg)	794.1 lb (360.5 kg)	1612.8 lb (732.2 kg)	1446.0 lb (656.5 kg)	1485.4 lb (674.4 kg)	1589.3 lb (721.5 kg)
Average daily feed consumed in treatment period (per head) by pen	24.11 lb (10.94 kg)	22.12 lb (10.04 kg)	22.42 lb (10.18 kg)	20.25 lb (9.19 kg)	20.62 lb (9.36 kg)	22.06 lb (10.02 kg)
Average Daily feed consumed in withdrawal period (per head) by pen	22.19 lb (10.07 kg)	21.74 lb (9.87 kg)	22.28 lb (10.11 kg)	19.24 lb (8.73 kg)	20.67 lb (9.38 kg)	22.15 lb (10.06 kg)
Total average daily feed consumed (per head) by pen (all 18 days)	23.79 lb (10.80 kg)	22.06 lb (10.02 kg)	22.40 lb (10.17 kg)	20.08 lb (9.12 kg)	20.63 lb (9.37 kg)	22.07 lb (10.02 kg)

5

[63] The words "comprise", "comprises", and "comprising" in this patent (including the claims) are to be interpreted inclusively rather than exclusively. This interpretation is intended to be the same as the interpretation that these words are given under United States patent law.

10

[64] The term “pharmaceutically acceptable” is used adjectivally to mean that the modified noun is appropriate for use in a pharmaceutical product. When it is used, for example, to describe a salt or carrier material, it characterizes the salt or carrier material as not being deleterious to the intended recipient animal to the extent
5 that the deleterious effect(s) outweighs the benefit(s) of the salt or carrier material.

[65] This patent sometimes refers to concentrations of various ingredients in an animal feed based on the feed containing approximately 90% dry matter. This stems from the fact that the dry matter content in many typical animal feeds is approximately 90%. This is particularly true for livestock feeds, such as feeds for
10 bovine animals. It should be recognized that such concentrations can be adjusted to account for other dry matter percentages in a feed. To illustrate, if a desired concentration of an ingredient is described as a specific concentration based on the feed having 90% dry matter, the desired concentration in a feed containing no water would be the specific concentration divided by 0.9.

[66] The above detailed description of preferred embodiments is intended
15 only to acquaint others skilled in the art with the invention, its principles, and its practical application so that others skilled in the art may adapt and apply the invention in its numerous forms, as they may be best suited to the requirements of a particular use. This invention, therefore, is not limited to the above embodiments, and may be
20 variously modified.

We claim:

1. A method for improving feed utilization efficiency or increasing the rate of weight gain in an animal, wherein the method comprises administering to the animal
5 an amount of zilpaterol (or a salt thereof) and an amount of melengestrol acetate.

2. A method of claim 1, wherein the amount of the zilpaterol (or salt thereof) and the amount of melengestrol acetate are, when combined, sufficient to improve the animal's average daily feed utilization efficiency over a period of at least about 3 days
10 relative to the average daily feed utilization efficiency that would be realized by administering the amount of the zilpaterol (or salt thereof) without any melengestrol acetate over the same amount of time under otherwise substantially identical feeding conditions.

3. A method of any one of claims 1 and 2, wherein the amount of the zilpaterol (or salt thereof) and the amount of melengestrol acetate are, when combined, sufficient to increase the animal's average daily weight gain over a period of at least about 3 days by at least about 100% the average daily weight gain that is realized during the week immediately before administration of the zilpaterol (or salt thereof)
15 20 and melengestrol acetate under otherwise substantially identical feeding conditions.

4. A method of any one of claims 1 to 3, wherein the amount of the zilpaterol (or salt thereof) and the amount of melengestrol acetate are, when combined, sufficient to increase the animal's average daily weight gain over a period of at least
25 about 3 days relative to the average daily weight gain that would be realized by administering the amount of the zilpaterol (or salt thereof) without any melengestrol acetate over the same amount of time under otherwise substantially identical feeding conditions.

5. A method of any one of claims 1 to 4, wherein the amount of the zilpaterol (or salt thereof) and the amount of melengestrol acetate are, when combined, sufficient to increase the animal's average daily weight gain over a period of at least about 3 days by at least about 1.2 times the average daily weight gain that would be realized by administering the amount of the zilpaterol (or salt thereof) without any
30

melengestrol acetate over the same amount of time under otherwise substantially identical feeding conditions.

5 6. A method of any one of claims 1 to 5, wherein at least a portion of the zilpaterol (or salt thereof) and at least a portion of the melengestrol acetate are administered in a single formulation.

10 7. A method of any one of claims 1 to 5, wherein at least a portion of the zilpaterol (or salt thereof) is administered in a separate formulation than at least a portion of the melengestrol acetate.

15 8. A method of any one of claims 1 to 7, wherein at least a portion of the zilpaterol (or salt thereof) or at least a portion of the melengestrol acetate is administered orally.

 9. A method of any one of claims 1 to 8, wherein the zilpaterol (or salt thereof) and the melengestrol acetate are administered orally.

20 10. A method of any one of claims 1 to 9, wherein the method comprises administering zilpaterol hydrochloride.

 11. A method of any one of claims 1 to 10, wherein the method further comprises administering at least one additional steroid.

25 12. A method of any one of claims 1 to 11, wherein the method further comprises administering trenbolone acetate.

30 13. A method of any one of claims 1 to 12, wherein the method further comprises administering zeranol, estradiol, or estradiol benzoate.

 14. A method of any one of claims 1 to 13, wherein the method further comprises administering at least one ionophore.

15. A method of any one of claims 1 to 14, wherein the method further comprises administering monensin (or a salt thereof).

16. A method of any one of claims 1 to 15, wherein the method further
5 comprises administering monensin sodium.

17. A method of any one of claims 1 to 16, wherein the method further comprises administering at least one antibiotic.

10 18. A method of any one of claims 1 to 17, wherein the method further comprises administering tylosin (or a salt thereof).

19. A method of any one of claims 1 to 18, wherein the method further comprises administering tylosin phosphate or tylosin tartrate.

15 20. A method of any one of claims 1 to 19, wherein the animal is a bovine animal.

20 21. A method of any one of claims 1 to 19, wherein the animal is a swine animal.

22. A method of any one of claims 1 to 19, wherein the animal is a bird.

25 23. A method of any one of claims 1 to 19, wherein the animal is a fish.

24. A composition, wherein the composition comprises zilpaterol (or a salt thereof) and melengestrol acetate.

30 25. A composition of claim 24, wherein the composition comprises zilpaterol hydrochloride.

26. A composition of any one of claims 24 and 25, wherein the composition further comprises one or more additional components selected from the group consisting of steroids, ionophores, and antibiotics.

27. A composition of any one of claims 24 to 26, wherein the composition further comprises one or more additional components selected from the group consisting of trenbolone acetate, zeranol, estradiol, estradiol benzoate, monensin, salts
5 of monensin, tylosin, and salts of tylosin.

28. A ruminant feedstuff for improving feed utilization efficiency or increasing the rate of weight gain in a ruminant animal, wherein the feedstuff comprises a composition of any one of claims 24 to 27 mixed with at least one carrier
10 material.

29. A ruminant feedstuff of claim 28, wherein the zilpaterol (or salt thereof) and melengestrol acetate are present in the feedstuff in amounts that, when combined, are sufficient to improve the animal's average daily feed utilization efficiency over a
15 period of at least about 3 days relative to the average daily feed utilization efficiency that would be realized by administering the amount of the zilpaterol (or salt thereof) without any melengestrol acetate over the same amount of time under otherwise substantially identical feeding conditions.

20 30. A ruminant feedstuff of any one of claims 28 and 29, wherein the zilpaterol (or salt thereof) and melengestrol acetate are present in the feedstuff in amounts that, when combined, increase the average daily weight gain of the animal over a period at least about 3 days relative to the average daily weight gain that would be realized by administering the amount of the zilpaterol (or salt thereof) without any
25 melengestrol acetate over the same amount of time under otherwise substantially identical feeding conditions.

31. A kit for administering zilpaterol (or a salt thereof) and melengestrol acetate to an animal, wherein the kit comprises:
30 a zilpaterol dosage form comprising zilpaterol (or a salt thereof); and
one or more of the following:
a melengestrol acetate dosage form comprising melengestrol
acetate,

instructions for administering zilpaterol (or a salt thereof) and
melengestrol acetate to the animal, and
instructions for mixing at least a portion of the zilpaterol
dosage form with melengestrol acetate.

5

32. A kit of claim 31, wherein:

the kit comprises a melengestrol acetate dosage form comprising melengestrol
acetate,

the amount of zilpaterol (or salt thereof) in the zilpaterol dosage form and the
10 amount of melengestrol acetate in the melengestrol acetate dosage form are, when
combined, sufficient to improve the animal's average daily feed utilization efficiency
over a period of at least about 3 days relative to the average daily feed utilization
efficiency that would be realized by administering the amount of the zilpaterol (or salt
thereof) in the zilpaterol dosage form without any melengestrol acetate over the same
15 amount of time under otherwise substantially identical feeding conditions.

33. A kit of any one of claims 31 to 32, wherein:

the kit comprises a melengestrol acetate dosage form comprising melengestrol
acetate,

20 the amount of the zilpaterol (or salt thereof) in the zilpaterol dosage form and
the amount of melengestrol acetate in the melengestrol acetate dosage form are, when
combined, sufficient to increase the animal's average daily weight gain over a period
of at least about 3 days relative to the average daily weight gain that would be realized
by administering the amount of the zilpaterol (or salt thereof) without any
25 melengestrol acetate over the same amount of time under otherwise substantially
identical feeding conditions.

34. A kit of any one of claims 31 to 33, wherein the kit comprises instructions
for orally administering at least a portion of the zilpaterol dosage form or a
30 melengestrol acetate dosage form comprising melengestrol acetate.

35. A kit of any one of claims 31 to 34, wherein the kit comprises instructions
for orally administering the zilpaterol dosage form and a melengestrol acetate dosage
form comprising melengestrol acetate.

36. A kit of any one of claims 31 to 35, wherein the zilpaterol dosage form comprises zilpaterol hydrochloride.

5 37. A kit of any one of claims 31 to 36, wherein the kit further comprises:
one or more additional components selected from the group consisting of
steroids, ionophores, and antibiotics; or
instructions for administering one or more ingredients selected from the group
consisting of steroids, ionophores, and antibiotics.

10

38. A kit of any one of claims 31 to 37, wherein the kit further comprises:
one or more additional components selected from the group consisting of
trenbolone acetate, zeranol, estradiol, estradiol benzoate, monensin, salts of monensin,
tylosin, and salts of tylosin; or

15

instructions for administering one or more ingredients selected from the group
consisting of trenbolone acetate, zeranol, estradiol, estradiol benzoate, monensin, salts
of monensin, tylosin, and salts of tylosin.

39. A kit of any one of claims 31 to 38, wherein the animal is a bovine animal.

20

40. A kit of any one of claims 31 to 38, wherein the animal is a swine animal.

41. A kit for administering zilpaterol (or a salt thereof) and melengestrol
acetate to an animal, wherein the kit comprises:

25

a melengestrol acetate dosage form comprising melengestrol acetate; and
one or more of the following:

a zilpaterol dosage form comprising zilpaterol (or a salt
thereof),

30

instructions for administering melengestrol acetate and
zilpaterol (or a salt thereof) to the animal, and

instructions for mixing at least a portion of the melengestrol
acetate dosage form with zilpaterol (or a salt thereof).

42. A kit of claim 41, wherein the kit comprises instructions for orally administering at least a portion of the melengestrol acetate dosage form or a zilpaterol dosage form comprising zilpaterol (or a salt thereof).

5 43. A kit of any one of claims 41 and 42, wherein the kit comprises instructions for orally administering the melengestrol acetate dosage form and a zilpaterol dosage form comprising zilpaterol (or a salt thereof).

10 44. A kit of any one of claims 41 to 43, wherein the kit further comprises:
one or more additional components selected from the group consisting of
steroids, ionophores, and antibiotics; or
instructions for administering one or more ingredients selected from the group
consisting of steroids, ionophores, and antibiotics.

15 45. A kit of any one of claims 41 to 44, wherein the kit further comprises:
one or more additional components selected from the group consisting of
trenbolone acetate, zeranol, estradiol, estradiol benzoate, monensin, salts of monensin,
tylosin, and salts of tylosin; or
instructions for administering one or more ingredients selected from the group
20 consisting of trenbolone acetate, zeranol, estradiol, estradiol benzoate, monensin, salts
of monensin, tylosin, and salts of tylosin.

 46. A kit of any one of claims 41 to 45, wherein the kit comprises one or more
of the following:

25 a zilpaterol dosage form comprising zilpaterol hydrochloride,
instructions for administering melengestrol acetate and zilpaterol
hydrochloride to the animal, and
instructions for mixing at least a portion of the melengestrol acetate
dosage form with zilpaterol hydrochloride.

30 47. A kit of any one of claims 41 to 46, wherein the animal is a bovine animal.

 48. A kit of any one of claims 41 to 46, wherein the animal is a swine animal.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2008/085455

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K31/55 A61K31/57 A23K1/16 A23K1/165 A61P43/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4 900 735 A (GRANDADAM JEAN A [FR]) 13 February 1990 (1990-02-13) cited in the application See col. 1, lines 10-15; col. 4, lines 52-56; col. 5, lines 5-10; claims 1, 13-18: compositions for increasing animal weight See col. 1, line 22 - col. 2, line 50; col. 5, line 40-45; claims 1, 2, 4, 7: compositions comprising a beta adrenergic preferably being zilpaterol and a steroid, preferably being trenbolone acetate ----- -/--	1-48

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

27 February 2009

Date of mailing of the international search report

11/03/2009

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Veronese, Andrea

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2008/085455

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SCHIFFER B ET AL: "The fate of trenbolone acetate and melengestrol acetate after application as growth promoters in cattle: environmental studies." ENVIRONMENTAL HEALTH PERSPECTIVES NOV 2001, vol. 109, no. 11, November 2001 (2001-11), pages 1145-1151, XP002517023 ISSN: 0091-6765 See abstract and page 1145, column 1 and 2: both trenbolone acetate and melengestrol acetate are commonly used as growth promoters for farm animals	1-48
Y	HENDRICKS D M ET AL: "SERUM CONCENTRATIONS OF TRENBOLONE-17BETA AND ESTRADIOL-17BETA AND PERFORMANCE OF HEIFERS TREATED WITH TRENBOLONE ACETATE, MELENGESTROL ACETATE, OR ESTRADIOL-17BETA" JOURNAL OF ANIMAL SCIENCE, AMERICAN SOCIETY OF ANIMAL SCIENCE, US, vol. 75, no. 10, 1 October 1997 (1997-10-01), pages 2627-2633, XP001037762 ISSN: 0021-8812 See abstract and results reported in page 2632: both trenbolone acetate and melengestrol acetate are commonly used as growth promoters for farm animals	1-48
Y	AVENDAÑO-REYES L ET AL: "Effects of two beta-adrenergic agonists on finishing performance, carcass characteristics, and meat quality of feedlot steers." JOURNAL OF ANIMAL SCIENCE DEC 2006, vol. 84, no. 12, December 2006 (2006-12), pages 3259-3265, XP002517024 ISSN: 1525-3163 See abstract: zilpaterol improves feedlot performance, and meat quality in steers	1-48
P,X	"New animal drugs for use in animal feed; zilpaterol" FEDERAL REGISTER , 73(70), 19432, 10 APR 2008 CODEN: FEREAC; ISSN: 0097-6326, 2008, XP009112954 See combinations of Zilpaterol, melengestrol acetate and monensin for producing medicated feeds to increase weigh gain in heifers	1-48

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2008/085455

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4900735	A	13-02-1990	DE 3772734 D1 10-10-1991
			EP 0272976 A2 29-06-1988
			HU 47852 A2 28-04-1989
			IE 60964 B1 07-09-1994
			JP 2893520 B2 24-05-1999
			JP 9176016 A 08-07-1997
			JP 2717788 B2 25-02-1998
			JP 63190834 A 08-08-1988
			NZ 222883 A 26-03-1991
