




(86) **Date de dépôt PCT/PCT Filing Date:** 2007/05/02
(87) **Date publication PCT/PCT Publication Date:** 2007/11/15
(45) **Date de délivrance/Issue Date:** 2017/01/17
(85) **Entrée phase nationale/National Entry:** 2008/12/05
(86) **N° demande PCT/PCT Application No.:** US 2007/010587
(87) **N° publication PCT/PCT Publication No.:** 2007/130429
(30) **Priorité/Priority:** 2006/05/03 (US60/797,211)

(51) **Cl.Int./Int.Cl.** **C07D 413/12** (2006.01),
A61K 31/357 (2006.01), **A61K 31/422** (2006.01),
A61K 31/427 (2006.01), **A61K 31/4433** (2006.01),
A61K 31/69 (2006.01), **C07D 319/06** (2006.01),
C07D 405/12 (2006.01), **C07D 417/12** (2006.01)
(72) **Inventeurs/Inventors:**
MAZITSCHK, RALPH, US;
KWIATKOWSKI, NICHOLAS PAUL, US;
BRADNER, JAMES ELLIOTT, US
(73) **Propriétaires/Owners:**
PRESIDENT AND FELLOWS OF HARVARD COLLEGE,
US;
DANA-FARBER CANCER INSTITUTE, INC., US
(74) **Agent:** ROBIC

(54) **Titre : INHIBITEURS D'HISTONE DESACETYLASES ET DE TUBULINE DESACETYLASES**
(54) **Title: HISTONE DEACETYLASE AND TUBULIN DEACETYLASE INHIBITORS**



(57) **Abrégé/Abstract:**

In recognition of the need to develop novel therapeutic agents and efficient methods for the synthesis thereof, the present invention provides novel inhibitors of histone deacetylases, tubulin deacetylases, and/or aggresome inhibitors, and pharmaceutically

(57) Abrégé(suite)/Abstract(continued):

acceptable salts and derivatives thereof. The inventive compounds fall into two classes — "isorubacin" class and "isoisotubacin" class- — all of which include a 1,3-dioxane core. The present invention further provides methods for treating disorders regulated by histone deacetylase activity, tubulin deacetylase activity, and/or the aggresome (e.g., proliferative diseases, cancer, inflammatory diseases, protozoal infections, protein degradation disorders, protein deposition disorders, etc.) comprising administering a therapeutically effective amount of an inventive compound to a subject in need thereof. The present invention also provides methods for preparing compounds of the invention.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
15 November 2007 (15.11.2007)

PCT

(10) International Publication Number
WO 2007/130429 A3

(51) International Patent Classification:

A61K 31/33 (2006.01) C07D 233/00 (2006.01)
 A61K 31/47 (2006.01) C07D 263/00 (2006.01)
 A61K 31/44 (2006.01) C07D 277/68 (2006.01)
 A61K 31/415 (2006.01) C07D 403/00 (2006.01)
 A61K 31/335 (2006.01) C07D 405/00 (2006.01)
 C07D 215/00 (2006.01) C07D 413/00 (2006.01)

02145 (US). **KWIATKOWSKI, Nicholas, Paul** [US/US];
 236 Bryn Mawr Avenue, Auburn, MA 01501 (US). **BRAD-**
NER, James, Elliott [US/US]; 129 Franklin Street, Apt.
 317, Cambridge, MA 02139 (US).

(74) **Agent: BAKER, Hunter, C., M., D.**; Choate, Hall &
 Stewart LLP, Two International Place, Boston, MA 02110
 (US).

(21) International Application Number:

PCT/US2007/010587

(22) International Filing Date: 2 May 2007 (02.05.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/797,211 3 May 2006 (03.05.2006) US

(71) **Applicant (for all designated States except US): THE**
PRESIDENT AND FELLOWS OF HARVARD COL-
LEGE [US/US]; 17 Quincy Avenue, Cambridge, MA
 02139 (US).

(72) Inventors; and


(75) **Inventors/Applicants (for US only): MAZITSCHK,**
Ralph [DE/US]; 43 Glenwood Road, Somerville, MA

(81) **Designated States (unless otherwise indicated, for every**
kind of national protection available): AE, AG, AL, AM,
 AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH,
 CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES,
 FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN,
 IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR,
 LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX,
 MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO,
 RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM,
 TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States (unless otherwise indicated, for every**
kind of regional protection available): ARIPO (BW, GH,
 GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
 ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
 European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
 FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL,
 PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM,
 GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: HISTONE DEACETYLASE AND TUBULIN DEACETYLASE INHIBITORS



(57) **Abstract:** In recognition of the need to develop novel therapeutic agents and efficient methods for the synthesis thereof, the present invention provides novel inhibitors of histone deacetylases, tubulin deacetylases, and/or aggresome inhibitors, and pharmaceutically acceptable salts and derivatives thereof. The inventive compounds fall into two classes — "isorubacin" class and "isoisotubacin" class — all of which include a 1,3-dioxane core. The present invention further provides methods for treating disorders regulated by histone deacetylase activity, tubulin deacetylase activity, and/or the aggresome (e.g., proliferative diseases, cancer, inflammatory diseases, protozoal infections, protein degradation disorders, protein deposition disorders, etc.) comprising administering a therapeutically effective amount of an inventive compound to a subject in need thereof. The present invention also provides methods for preparing compounds of the invention.

WO 2007/130429 A3



Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

(88) Date of publication of the international search report:
3 April 2008

Histone Deacetylase and Tubulin Deacetylase Inhibitors

Government Support

[0002] The work described herein was supported, in part, by grants from the National Institutes of Health (370 31215 017490 614101 0001 00000; 370 31215 133800 341450 0402 44046). The United States government may have certain rights in the invention.

Background of the Invention

[0003] The identification of small organic molecules that affect specific biological functions is an endeavor that impacts both biology and medicine. Such molecules are useful as therapeutic agents and as probes of biological function. In but one example from the emerging field of chemical genetics, in which small molecules can be used to alter the function of biological molecules to which they bind, these molecules have been useful at elucidating signal transduction pathways by acting as chemical protein knockouts, thereby causing a loss of protein function. (Schreiber *et al.*, *J. Am. Chem. Soc.*, **1990**, *112*, 5583; Mitchison, *Chem. and Biol.*, **1994**, *1*, 3) Additionally, due to the interaction of these small molecules with particular biological targets and their ability to affect specific biological function (*e.g.* gene transcription), they may also serve as candidates for the development of new therapeutics. One important class of small molecules, natural products, which are small molecules obtained from nature, clearly have played an important role in the development of biology and medicine, serving as pharmaceutical leads, drugs (Newman *et al.*, *Nat. Prod. Rep.* **2000**, *17*, 215-234), and powerful reagents for studying cell biology (Schreiber, *S.L. Chem. and Eng. News* **1992** (October 26), 22-32).


[0004] One biological target of recent interest is histone deacetylase (see, for example, a discussion of the use of inhibitors of histone deacetylases for the treatment of cancer: Marks *et al. Nature Reviews Cancer* **2001**, *1*, 194; Johnstone *et al. Nature Reviews Drug Discovery* **2002**, *1*, 287). Post-translational modification of proteins through acetylation and deacetylation of lysine residues plays a critical role in regulating their cellular functions.

HDACs are zinc hydrolases that modulate gene expression through deacetylation of the *N*-acetyl-lysine residues of histone proteins and other transcriptional regulators (Hassig *et al. Curr. Opin. Chem. Biol.* **1997**, *1*, 300-308). HDACs participate in cellular pathways that control cell shape and differentiation, and an HDAC inhibitor has been shown effective in treating an otherwise recalcitrant cancer (Warrell *et al. J. Natl. Cancer Inst.* **1998**, *90*, 1621-1625). At this time, eleven human HDACs, which use Zn as a cofactor, have been identified (Taunton *et al. Science* **1996**, *272*, 408-411; Yang *et al. J. Biol. Chem.* **1997**, *272*, 28001-28007. Grozinger *et al. Proc. Natl. Acad. Sci. U.S.A.* **1999**, *96*, 4868-4873; Kao *et al. Genes Dev.* **2000**, *14*, 55-66. Hu *et al. J. Biol. Chem.* **2000**, *275*, 15254-15264; Zhou *et al. Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 10572-10577; Venter *et al. Science* **2001**, *291*, 1304-1351) these members fall into three classes (class I, II, and IV). An additional seven HDACs have been identified which use NAD as a cofactor. To date, no small molecules are known that selectively target any particular class or individual members of this family ((for example ortholog-selective HDAC inhibitors have been reported: (a) Meinke *et al. J. Med. Chem.* **2000**, *14*, 4919-4922; (b) Meinke, *et al. Curr. Med. Chem.* **2001**, *8*, 211-235). There remains a need for preparing structurally diverse HDAC and tubulin deacetylase (TDAC) inhibitors particularly ones that are potent and/or selective inhibitors of particular classes of HDACs or TDACs and individual HDACs and TDACs.

Summary of the Invention



[0005] The present invention provides novel histone deacetylase and tubulin deacetylase inhibitors and methods of preparing and using these compounds. These compounds are particularly useful in the treatment of proliferative diseases such as cancer, inflammatory diseases, infectious diseases, protein degradation disorders, and protein deposition disorders such as Alzheimer's Disease. Certain compounds are particularly useful in specifically inhibiting one class or member of HDACs. Other compounds are particularly useful in specifically inhibiting one class or member of tubulin deacetylases (TDAC). Yet other compounds are useful in inhibiting degradation of proteins by the aggresome.

[0006] The present invention provides novel inhibitors of HDACs and TDACs with a 1,3-dioxane core structure. Compounds of the invention basically fall into two classes, wherein the 1,3-dioxane core of the compound is oriented differently in each class. That is, the 1,3-dioxane core is rotated 120° in each class as compared to tubacin derivatives and as shown in the structures below (see, also, *Figure 1*). The inventive compounds are of the formulae:



wherein


R_1 is of the formula:



Y is NH or O;

L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety; a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety; a substituted or unsubstituted aryl moiety; a substituted or unsubstituted heteroaryl moiety; or a cyclic ring system, wherein the rings may be aryl, heteroaryl, non-aromatic carbocyclic, or non-aromatic heterocyclic; and


A comprises a functional group selected from the group consisting of:



R_2 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-\text{OR}_B$; $-\text{C}(=\text{O})\text{R}_B$; $-\text{CO}_2\text{R}_B$; $-\text{CN}$; $-\text{SCN}$; $-\text{SR}_B$; $-\text{SOR}_B$; $-\text{SO}_2\text{R}_B$; $-\text{NO}_2$; $-\text{N}(\text{R}_B)_2$; $-\text{NHC}(\text{O})\text{R}_B$; or $-\text{C}(\text{R}_B)_3$; wherein each occurrence of R_B is independently a hydrogen, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety; and


R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-\text{OR}_C$; $-\text{C}(=\text{O})\text{R}_C$; $-\text{CO}_2\text{R}_C$; $-\text{CN}$; $-\text{SCN}$; $-\text{SR}_C$; $-\text{SOR}_C$; $-\text{SO}_2\text{R}_C$; $-\text{NO}_2$; $-\text{N}(\text{R}_C)_2$; $-\text{NHC}(\text{O})\text{R}_C$; or $-\text{C}(\text{R}_C)_3$; wherein each occurrence of R_C is independently a hydrogen, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl

moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety; or pharmaceutically acceptable salts thereof. In general, R1 comprises a metal chelating functional group (*e.g.*, hydroxyamic acids, thiols, carboxylic acids, *ortho*-aminoanilides, etc.). The metal chelating




group is thought to bind the active site Zn^{+2} ion of deacetylase enzymes. In certain embodiments, R_2 is a substituted or unsubstituted heteroaliphatic moiety (*e.g.*, a heteroaliphatic moiety substituted with a heteroaryl ring, which may be optionally substituted). In certain embodiments, R_3 is a substituted or unsubstituted aromatic ring system (*e.g.*, a substituted or unsubstituted phenyl).

[0007] Compounds of formula:




that is, tubacin derivatives, have been described in U.S. patent publication no. 2006/0239909, published on October 26, 2006; U.S. patent publication no. 2003/0187027, published on October 2, 2003; and U.S. patent application no. 2004/0072849, published on April 15, 2004. These compounds have been found to be particularly useful in inhibiting HDACs such as HDAC6 and also in the treatment of multiple myeloma by inhibiting HDAC6 known to play a role in the degradation of proteins by the aggresome.

[0008] Inventive compounds of the class of formula:




are compounds of the "isotubacin" class. The 1,3-dioxane core of these compounds has been rotated 120° counter-clockwise as compared to compounds of the "tubacin" class. The compounds of this class are also useful in treating cancer (*e.g.*, multiple myeloma, breast cancer, non-Hodgkin's lymphoma, ovarian cancer, acute myelogenous leukemia), protein degradation disorders (*e.g.*, multiple myeloma, neurodegenerative disorders), protein deposition disorders (*e.g.*, neurodegenerative disorders), infectious diseases, and proliferative

disorders (e.g., diabetic retinopathy, inflammatory diseases, angiogenesis). Pharmaceutical compositions and kits comprising these compounds are also provided by the present invention. In certain particular embodiments, compounds of this class are useful in the treatment of multiple myeloma, leukemia, lymphoma, breast cancer, and prostate cancer. Pharmaceutical composition of the inventive compounds may also comprises other chemotherapeutic agents or other pharmaceutical agents typically administered during the



treatment of cancer (*e.g.*, anti-nausea medications, analgesics, nutritional supplements, *etc.*). The present invention also provides synthetic method for preparing compounds of the “isotubacin” class.

[0009] Inventive compounds of the class of formula:



are compounds of the “isoisotubacin” class. The 1,3-dioxane core of the compounds of this class has been rotated 120° clockwise as compared to compounds of the “tubacin” class. The compounds of this class are also useful in treating cancer (*e.g.*, multiple myeloma, breast cancer, non-Hodgkin’s lymphoma, ovarian cancer, acute myelogenous leukemia), protein degradation disorders (*e.g.*, neurodegenerative disorders, multiple myeloma), protein deposition disorders (*e.g.*, neurodegenerative disorders), infectious diseases, and proliferative disorders (*e.g.*, diabetic retinopathy, inflammatory diseases, angiogenesis). Pharmaceutical compositions and kits comprising these compounds are also included. In certain particular embodiments, compounds of this class are useful in the treatment of multiple myeloma, leukemia, lymphoma, breast cancer, and prostate cancer. Pharmaceutical composition of the inventive compounds may also comprises other chemotherapeutic agents or other pharmaceutical agents typically administered during the treatment of cancer (*e.g.*, anti-nausea medications, analgesics, nutritional supplements, *etc.*). The present invention also provides synthetic method for preparing compounds of the “isoisotubacin” class.

[0010] The inventive compounds are also useful as tools to probe biological function (*e.g.*, the degradation of proteins by the aggresome; inhibition of histone deacetylases, inhibition of tubulin deacetylases). For example, the compounds may be administered to wild type cells or altered cells to understand protein degradation pathways or the effect of acetylation on a protein’s function. In certain embodiments, the compound inhibits a specific histone or tubulin deacetylase. The compounds may also be used to elucidate the cell cycle.

[0011] In another aspect, the present invention provides methods for inhibiting histone deacetylase activity, tubulin deacetylase activity, or aggresome activity in a patient or a biological sample, comprising administering to the patient, or contacting the biological sample with an effective amount of an inventive compound or a pharmaceutical composition thereof.

Definitions

[0012] . Certain compounds of the present invention, and definitions of specific functional groups are also described in more detail below. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed., inside cover, and specific functional groups are generally defined as described therein. Additionally, general principles of organic chemistry, as well as specific functional moieties and reactivity, are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999. Furthermore, it will be appreciated by one of ordinary skill in the art that the synthetic methods, as described herein, utilize a variety of protecting groups.

[0013] It will be appreciated that the compounds, as described herein, may be substituted with any number of substituents or functional moieties. In general, the term "substituted" whether preceded by the term "optionally" or not, and substituents contained in formulas of this invention, refer to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. When more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. As used herein, the term "substituted" is contemplated to include all permissible substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and nonaromatic substituents of organic compounds. For purposes of this invention, heteroatoms such as nitrogen may have hydrogen substituents and/or any permissible substituents of organic compounds described herein which satisfy the valencies of the heteroatoms. Furthermore, this invention is not intended to be limited in any manner by the permissible substituents of organic compounds. Combinations of substituents and variables envisioned by this invention are preferably those that result in the formation of stable compounds useful in the treatment, for example of proliferative disorders, including, but not limited to cancer. The term "stable", as used herein, preferably refers to compounds which possess stability sufficient to allow manufacture and which maintain the integrity of the compound for a sufficient period of time to be detected and preferably for a sufficient period of time to be useful for the purposes detailed herein.

[0014] The term "acyl", as used herein, refers to a carbonyl-containing functionality, *e.g.*, -C(=O)R', wherein R' is an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aryl, heteroaryl, (aliphatic)aryl, (heteroaliphatic)aryl, heteroaliphatic(aryl) or heteroaliphatic(heteroaryl)

moiety, whereby each of the aliphatic, heteroaliphatic, aryl, or heteroaryl moieties is substituted or unsubstituted, or is a substituted (*e.g.*, hydrogen or aliphatic, heteroaliphatic, aryl, or heteroaryl moieties) oxygen or nitrogen containing functionality (*e.g.*, forming a carboxylic acid, ester, or amide functionality).

[0015] The term “aliphatic”, as used herein, includes both saturated and unsaturated, straight chain (*i.e.*, unbranched) or branched aliphatic hydrocarbons, which are optionally substituted with one or more functional groups. As will be appreciated by one of ordinary skill in the art, “aliphatic” is intended herein to include, but is not limited to, alkyl, alkenyl, alkynyl moieties. Thus, as used herein, the term “alkyl” includes straight and branched alkyl groups. An analogous convention applies to other generic terms such as “alkenyl”, “alkynyl” and the like. Furthermore, as used herein, the terms “alkyl”, “alkenyl”, “alkynyl” and the like encompass both substituted and unsubstituted groups. In certain embodiments, as used herein, “lower alkyl” is used to indicate those alkyl groups (substituted, unsubstituted, branched or unbranched) having 1-6 carbon atoms.

[0016] In certain embodiments, the alkyl, alkenyl and alkynyl groups employed in the invention contain 1-20 aliphatic carbon atoms. In certain other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-10 aliphatic carbon atoms. In yet other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-8 aliphatic carbon atoms. In still other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-6 aliphatic carbon atoms. In yet other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-4 carbon atoms. Illustrative aliphatic groups thus include, but are not limited to, for example, methyl, ethyl, n-propyl, isopropyl, allyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, sec-pentyl, isopentyl, tert-pentyl, n-hexyl, sec-hexyl, moieties and the like, which again, may bear one or more substituents. Alkenyl groups include, but are not limited to, for example, ethenyl, propenyl, butenyl, 1-methyl-2-buten-1-yl, and the like. Representative alkynyl groups include, but are not limited to, ethynyl, 2-propynyl (propargyl), 1-propynyl and the like.

[0017] The term “alicyclic”, as used herein, refers to compounds which combine the properties of aliphatic and cyclic compounds and include but are not limited to cyclic, or polycyclic aliphatic hydrocarbons and bridged cycloalkyl compounds, which are optionally substituted with one or more functional groups. As will be appreciated by one of ordinary skill in the art, “alicyclic” is intended herein to include, but is not limited to, cycloalkyl, cycloalkenyl, and cycloalkynyl moieties, which are optionally substituted with one or more

functional groups. Illustrative alicyclic groups thus include, but are not limited to, for example, cyclopropyl, -CH₂-cyclopropyl, cyclobutyl, -CH₂-cyclobutyl, cyclopentyl, -CH₂-cyclopentyl, cyclohexyl, -CH₂-cyclohexyl, cyclohexenylethyl, cyclohexanylethyl, norborboryl moieties and the like, which again, may bear one or more substituents.

[0018] The term “alkoxy” (or “alkyloxy”), or “thioalkyl” as used herein refers to an alkyl group, as previously defined, attached to the parent molecular moiety through an oxygen atom or through a sulfur atom. In certain embodiments, the alkyl group contains 1-20 aliphatic carbon atoms. In certain other embodiments, the alkyl group contains 1-10 aliphatic carbon atoms. In yet other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-8 aliphatic carbon atoms. In still other embodiments, the alkyl group contains 1-6 aliphatic carbon atoms. In yet other embodiments, the alkyl group contains 1-4 aliphatic carbon atoms. Examples of alkoxy, include but are not limited to, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, tert-butoxy, neopentoxy and n-hexoxy. Examples of thioalkyl include, but are not limited to, methylthio, ethylthio, propylthio, isopropylthio, n-butylthio, and the like.

[0019] The term “alkylamino” refers to a group having the structure -NHR' wherein R' is alkyl, as defined herein. The term “aminoalkyl” refers to a group having the structure NH₂R'-, wherein R' is alkyl, as defined herein. In certain embodiments, the alkyl group contains 1-20 aliphatic carbon atoms. In certain other embodiments, the alkyl group contains 1-10 aliphatic carbon atoms. In yet other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-8 aliphatic carbon atoms. In still other embodiments, the alkyl group contains 1-6 aliphatic carbon atoms. In yet other embodiments, the alkyl group contains 1-4 aliphatic carbon atoms. Examples of alkylamino include, but are not limited to, methylamino, ethylamino, iso-propylamino and the like.

[0020] Some examples of substituents of the above-described aliphatic (and other) moieties of compounds of the invention include, but are not limited to aliphatic; heteroaliphatic; aryl; heteroaryl; alkylaryl; alkylheteroaryl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; F; Cl; Br; I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)₂R_x; -NR_x(CO)R_x wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, alicyclic, heteroaliphatic, heterocyclic, aryl, heteroaryl, alkylaryl, or alkylheteroaryl, wherein any of the aliphatic, heteroaliphatic, alkylaryl, or alkylheteroaryl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic,

and wherein any of the aryl or heteroaryl substituents described above and herein may be substituted or unsubstituted. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein.

[0021] In general, the term “aryl”, as used herein, refers to a stable mono- or polycyclic, unsaturated moiety having preferably 3-14 carbon atoms, each of which may be substituted or unsubstituted. In certain embodiments, the term “aryl” refers to a planar ring having p-orbitals perpendicular to the plane of the ring at each ring atom and satisfying the Huckel rule where the number of pi electrons in the ring is $(4n+2)$ wherein n is an integer. A mono- or polycyclic, unsaturated moiety that does not satisfy one or all of these criteria for aromaticity is defined herein as “non-aromatic”, and is encompassed by the term “alicyclic”.

[0022] In general, the term “heteroaryl”, as used herein, refers to a stable mono- or polycyclic, unsaturated moiety having preferably 3-14 carbon atoms, each of which may be substituted or unsubstituted; and comprising at least one heteroatom selected from O, S and N within the ring (*i.e.*, in place of a ring carbon atom). In certain embodiments, the term “heteroaryl” refers to a planar ring comprising at least one heteroatom, having p-orbitals perpendicular to the plane of the ring at each ring atom, and satisfying the Huckel rule where the number of pi electrons in the ring is $(4n+2)$ wherein n is an integer.

[0023] It will also be appreciated that aryl and heteroaryl moieties, as defined herein may be attached via an alkyl or heteroalkyl moiety and thus also include $-(\text{alkyl})\text{aryl}$, $-(\text{heteroalkyl})\text{aryl}$, $-(\text{heteroalkyl})\text{heteroaryl}$, and $-(\text{heteroalkyl})\text{heteroaryl}$ moieties. Thus, as used herein, the phrases “aryl or heteroaryl moieties” and “aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{heteroalkyl})\text{aryl}$, $-(\text{heteroalkyl})\text{heteroaryl}$, and $-(\text{heteroalkyl})\text{heteroaryl}$ ” are interchangeable. Substituents include, but are not limited to, any of the previously mentioned substituents, *i.e.*, the substituents recited for aliphatic moieties, or for other moieties as disclosed herein, resulting in the formation of a stable compound.

[0024] The term “aryl”, as used herein, does not differ significantly from the common meaning of the term in the art, and refers to an unsaturated cyclic moiety comprising at least one aromatic ring. In certain embodiments, “aryl” refers to a mono- or bicyclic carbocyclic ring system having one or two aromatic rings including, but not limited to, phenyl, naphthyl, tetrahydronaphthyl, indanyl, indenyl and the like.

[0025] The term “heteroaryl”, as used herein, does not differ significantly from the common meaning of the term in the art, and refers to a cyclic aromatic radical having from five to ten ring atoms of which one ring atom is selected from S, O and N; zero, one or two ring atoms are additional heteroatoms independently selected from S, O and N; and the

remaining ring atoms are carbon, the radical being joined to the rest of the molecule via any of the ring atoms, such as, for example, pyridyl, pyrazinyl, pyrimidinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, quinolinyl, isoquinolinyl, and the like.

[0026] It will be appreciated that aryl and heteroaryl groups (including bicyclic aryl groups) can be unsubstituted or substituted, wherein substitution includes replacement of one or more of the hydrogen atoms thereon independently with any one or more of the following moieties including, but not limited to: aliphatic; alicyclic; heteroaliphatic; heterocyclic; aromatic; heteroaromatic; aryl; heteroaryl; alkylaryl; heteroalkylaryl; alkylheteroaryl; heteroalkylheteroaryl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; F; Cl; Br; I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)R_x; -S(O)₂R_x; -NR_x(CO)R_x wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, heteroaromatic, aryl, heteroaryl, alkylaryl, alkylheteroaryl, heteroalkylaryl or heteroalkylheteroaryl, wherein any of the aliphatic, alicyclic, heteroaliphatic, heterocyclic, alkylaryl, or alkylheteroaryl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, saturated or unsaturated, and wherein any of the aromatic, heteroaromatic, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl substituents described above and herein may be substituted or unsubstituted. Additionally, it will be appreciated, that any two adjacent groups taken together may represent a 4, 5, 6, or 7-membered substituted or unsubstituted alicyclic or heterocyclic moiety. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein.

[0027] The term "cycloalkyl", as used herein, refers specifically to groups having three to seven, preferably three to ten carbon atoms. Suitable cycloalkyls include, but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like, which, as in the case of aliphatic, alicyclic, heteroaliphatic or heterocyclic moieties, may optionally be substituted with substituents including, but not limited to aliphatic; alicyclic; heteroaliphatic; heterocyclic; aromatic; heteroaromatic; aryl; heteroaryl; alkylaryl; heteroalkylaryl; alkylheteroaryl; heteroalkylheteroaryl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; F; Cl; Br; I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)₂R_x; -NR_x(CO)R_x wherein

each occurrence of R_x independently includes, but is not limited to, aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, heteroaromatic, aryl, heteroaryl, alkylaryl, alkylheteroaryl, heteroalkylaryl or heteroalkylheteroaryl, wherein any of the aliphatic, alicyclic, heteroaliphatic, heterocyclic, alkylaryl, or alkylheteroaryl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, saturated or unsaturated, and wherein any of the aromatic, heteroaromatic, aryl or heteroaryl substituents described above and herein may be substituted or unsubstituted. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein.

[0028] The term “heteroaliphatic”, as used herein, refers to aliphatic moieties in which one or more carbon atoms in the main chain have been substituted with a heteroatom. Thus, a heteroaliphatic group refers to an aliphatic chain which contains one or more oxygen, sulfur, nitrogen, phosphorus or silicon atoms, *e.g.*, in place of carbon atoms. Heteroaliphatic moieties may be linear or branched, and saturated or unsaturated. In certain embodiments, heteroaliphatic moieties are substituted by independent replacement of one or more of the hydrogen atoms thereon with one or more moieties including, but not limited to aliphatic; alicyclic; heteroaliphatic; heterocyclic; aromatic; heteroaromatic; aryl; heteroaryl; alkylaryl; alkylheteroaryl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; F; Cl; Br; I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O) R_x ; -CO₂(R_x); -CON(R_x)₂; -OC(O) R_x ; -OCO₂ R_x ; -OCON(R_x)₂; -N(R_x)₂; -S(O)₂ R_x ; -NR_x(CO) R_x wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, heteroaromatic, aryl, heteroaryl, alkylaryl, alkylheteroaryl, heteroalkylaryl or heteroalkylheteroaryl, wherein any of the aliphatic, alicyclic, heteroaliphatic, heterocyclic, alkylaryl, or alkylheteroaryl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, saturated or unsaturated, and wherein any of the aromatic, heteroaromatic, aryl or heteroaryl substituents described above and herein may be substituted or unsubstituted. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein.

[0029] The term “heterocycloalkyl”, “heterocycle” or “heterocyclic”, as used herein, refers to compounds which combine the properties of heteroaliphatic and cyclic compounds and include, but are not limited to, saturated and unsaturated mono- or polycyclic cyclic ring systems having 5-16 atoms wherein at least one ring atom is a heteroatom selected from O, S and N (wherein the nitrogen and sulfur heteroatoms may be optionally be oxidized), wherein

the ring systems are optionally substituted with one or more functional groups, as defined herein. In certain embodiments, the term "heterocycloalkyl", "heterocycle" or "heterocyclic" refers to a non-aromatic 5-, 6- or 7- membered ring or a polycyclic group wherein at least one ring atom is a heteroatom selected from O, S and N (wherein the nitrogen and sulfur heteroatoms may be optionally be oxidized), including, but not limited to, a bi- or tri-cyclic group, comprising fused six-membered rings having between one and three heteroatoms independently selected from oxygen, sulfur and nitrogen, wherein (i) each 5-membered ring has 0 to 2 double bonds, each 6-membered ring has 0 to 2 double bonds and each 7-membered ring has 0 to 3 double bonds, (ii) the nitrogen and sulfur heteroatoms may be optionally be oxidized, (iii) the nitrogen heteroatom may optionally be quaternized, and (iv) any of the above heterocyclic rings may be fused to an aryl or heteroaryl ring. Representative heterocycles include, but are not limited to, heterocycles such as furanyl, thiofuranyl, pyranal, pyrrolyl, thienyl, pyrrolidinyl, pyrazolinyl, pyrazolidinyl, imidazolinyl, imidazolidinyl, piperidinyl, piperazinyl, oxazolyl, oxazolidinyl, isooxazolyl, isoxazolidinyl, dioxazolyl, thiadiazolyl, oxadiazolyl, tetrazolyl, triazolyl, thiatriazolyl, oxatriazolyl, thiadiazolyl, oxadiazolyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, dithiazolyl, dithiazolidinyl, tetrahydrofuryl, and benzofused derivatives thereof. In certain embodiments, a "substituted heterocycle, or heterocycloalkyl or heterocyclic" group is utilized and as used herein, refers to a heterocycle, or heterocycloalkyl or heterocyclic group, as defined above, substituted by the independent replacement of one, two or three of the hydrogen atoms thereon with but are not limited to aliphatic; alicyclic; heteroaliphatic; heterocyclic; aromatic; heteroaromatic; aryl; heteroaryl; alkylaryl; heteroalkylaryl; alkylheteroaryl; heteroalkylheteroaryl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; F; Cl; Br; I; - OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)₂R_x; -NR_x(CO)R_x wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, heteroaromatic, aryl, heteroaryl, alkylaryl, alkylheteroaryl, heteroalkylaryl or heteroalkylheteroaryl, wherein any of the aliphatic, alicyclic, heteroaliphatic, heterocyclic, alkylaryl, or alkylheteroaryl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, saturated or unsaturated, and wherein any of the aromatic, heteroaromatic, aryl or heteroaryl substituents described above and herein may be substituted or unsubstituted. Additional examples or

generally applicable substituents are illustrated by the specific embodiments shown in the Examples, which are described herein.

[0030] Additionally, it will be appreciated that any of the alicyclic or heterocyclic moieties described above and herein may comprise an aryl or heteroaryl moiety fused thereto. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein. The terms “halo” and “halogen” as used herein refer to an atom selected from fluorine, chlorine, bromine and iodine.

[0031] The terms “halo” and “halogen” as used herein refer to an atom selected from fluorine, chlorine, bromine and iodine.

[0032] The term “haloalkyl” denotes an alkyl group, as defined above, having one, two, or three halogen atoms attached thereto and is exemplified by such groups as chloromethyl, bromoethyl, trifluoromethyl, and the like.

[0033] The term “amino”, as used herein, refers to a primary ($-\text{NH}_2$), secondary ($-\text{NHR}_x$), tertiary ($-\text{NR}_x\text{R}_y$) or quaternary ($-\text{N}^+\text{R}_x\text{R}_y\text{R}_z$) amine, where R_x , R_y and R_z are independently an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aryl, or heteroaryl moiety, as defined herein. Examples of amino groups include, but are not limited to, methylamino, dimethylamino, ethylamino, diethylamino, diethylaminocarbonyl, methylethylamino, isopropylamino, piperidino, trimethylamino, and propylamino.

[0034] The term “alkylidene”, as used herein, refers to a substituted or unsubstituted, linear or branched saturated divalent radical consisting solely of carbon and hydrogen atoms, having from one to n carbon atoms, having a free valence “-” at both ends of the radical. In certain embodiments, the alkylidene moiety has 1 to 6 carbon atoms.

[0035] The term “alkenylidene”, as used herein, refers to a substituted or unsubstituted, linear or branched unsaturated divalent radical consisting solely of carbon and hydrogen atoms, having from two to n carbon atoms, having a free valence “-” at both ends of the radical, and wherein the unsaturation is present only as double bonds and wherein a double bond can exist between the first carbon of the chain and the rest of the molecule. In certain embodiments, the alkenylidene moiety has 2 to 6 carbon atoms.

[0036] The term “alkynylidene”, as used herein, refers to a substituted or unsubstituted, linear or branched unsaturated divalent radical consisting solely of carbon and hydrogen atoms, having from two to n carbon atoms, having a free valence “-” at both ends of the radical, and wherein the unsaturation is present only as triple or double bonds and wherein a

triple or double bond can exist between the first carbon of the chain and the rest of the molecule. In certain embodiments, the alkynylidene moiety has 2 to 6 carbon atoms.

[0037] Unless otherwise indicated, as used herein, the terms “alkyl”, “alkenyl”, “alkynyl”, “heteroalkyl”, “heteroalkenyl”, “heteroalkynyl”, “alkylidene”, “alkenylidene”, “-(alkyl)aryl”, “-(heteroalkyl)aryl”, “-(heteroalkyl)aryl”, “-(heteroalkyl)heteroaryl”, and the like encompass substituted and unsubstituted, and linear and branched groups. Similarly, the terms “aliphatic”, “heteroaliphatic”, and the like encompass substituted and unsubstituted, saturated and unsaturated, and linear and branched groups. Similarly, the terms “cycloalkyl”, “heterocycle”, “heterocyclic”, and the like encompass substituted and unsubstituted, and saturated and unsaturated groups. Additionally, the terms “cycloalkenyl”, “cycloalkynyl”, “heterocycloalkenyl”, “heterocycloalkynyl”, “aromatic”, “heteroaromatic”, “aryl”, “heteroaryl” and the like encompass both substituted and unsubstituted groups.

[0038] The phrase, “pharmaceutically acceptable derivative”, as used herein, denotes any pharmaceutically acceptable salt, ester, or salt of such ester, of such compound, or any other adduct or derivative which, upon administration to a patient, is capable of providing (directly or indirectly) a compound as otherwise described herein, or a metabolite or residue thereof. Pharmaceutically acceptable derivatives thus include among others pro-drugs. A pro-drug is a derivative of a compound, usually with significantly reduced pharmacological activity, which contains an additional moiety, which is susceptible to removal *in vivo* yielding the parent molecule as the pharmacologically active species. An example of a pro-drug is an ester, which is cleaved *in vivo* to yield a compound of interest. Pro-drugs of a variety of compounds, and materials and methods for derivatizing the parent compounds to create the pro-drugs, are known and may be adapted to the present invention. Pharmaceutically acceptable derivatives also include “reverse pro-drugs.” Reverse pro-drugs, rather than being activated, are inactivated upon absorption. For example, as discussed herein, many of the ester-containing compounds of the invention are biologically active but are inactivated upon exposure to certain physiological environments such as a blood, lymph, serum, extracellular fluid, *etc.* which contain esterase activity. The biological activity of reverse pro-drugs and pro-drugs may also be altered by appending a functionality onto the compound, which may be catalyzed by an enzyme. Also, included are oxidation and reduction reactions, including enzyme-catalyzed oxidation and reduction reactions. Certain exemplary pharmaceutical compositions and pharmaceutically acceptable derivatives will be discussed in more detail herein below.

[0039] By the term “protecting group”, has used herein, it is meant that a particular functional moiety, *e.g.*, O, S, or N, is temporarily blocked so that a reaction can be carried out selectively at another reactive site in a multifunctional compound. In preferred embodiments, a protecting group reacts selectively in good yield to give a protected substrate that is stable to the projected reactions; the protecting group must be selectively removed in good yield by readily available, preferably nontoxic reagents that do not attack the other functional groups; the protecting group forms an easily separable derivative (more preferably without the generation of new stereogenic centers); and the protecting group has a minimum of additional functionality to avoid further sites of reaction. As detailed herein, oxygen, sulfur, nitrogen and carbon protecting groups may be utilized. For example, in certain embodiments, as detailed herein, certain exemplary oxygen protecting groups are utilized. These oxygen protecting groups include, but are not limited to methyl ethers, substituted methyl ethers (*e.g.*, MOM (methoxymethyl ether), MTM (methylthiomethyl ether), BOM (benzyloxymethyl ether), PMBM or MPM (p-methoxybenzyloxymethyl ether), to name a few), substituted ethyl ethers, substituted benzyl ethers, silyl ethers (*e.g.*, TMS (trimethylsilyl ether), TES (triethylsilyl ether), TIPS (triisopropylsilyl ether), TBDMS (t-butyldimethylsilyl ether), tribenzyl silyl ether, TBDPS (t-butyldiphenyl silyl ether), to name a few), esters (*e.g.*, formate, acetate, benzoate (Bz), trifluoroacetate, dichloroacetate, to name a few), carbonates, cyclic acetals and ketals. In certain other exemplary embodiments, nitrogen protecting groups are utilized. These nitrogen protecting groups include, but are not limited to, carbamates (including methyl, ethyl and substituted ethyl carbamates (*e.g.*, Troc), to name a few) amides, cyclic imide derivatives, N-Alkyl and N-Aryl amines, imine derivatives, and enamine derivatives, to name a few. Certain other exemplary protecting groups are detailed herein, however, it will be appreciated that the present invention is not intended to be limited to these protecting groups; rather, a variety of additional equivalent protecting groups can be readily identified using the above criteria and utilized in the present invention. Additionally, a variety of protecting groups are described in “Protective Groups in Organic Synthesis” Third Ed. Greene, T.W. and Wuts, P.G., Eds., John Wiley & Sons, New York: 1999.

[0040] The term “solid support”, as used herein, refers to a material having a rigid or semi-rigid surface. Such materials will preferably take the form of small beads, pellets, disks, chips, dishes, multi-well plates, glass slides, wafers, or the like, although other forms may be used. In some embodiments, at least one surface of the substrate will be substantially flat.

The term “surface” refers to any generally two-dimensional structure on a solid substrate and may have steps, ridges, kinks,, terraces, and the like without ceasing to be a surface.

[0041] The term “polymeric support”, as used herein, refers to a soluble or insoluble polymer to which an amino acid or other chemical moiety can be covalently bonded by reaction with a functional group of the polymeric support. Many suitable polymeric supports are known, and include soluble polymers such as polyethylene glycols or polyvinyl alcohols, as well as insoluble polymers such as polystyrene resins. A suitable polymeric support includes functional groups such as those described below. A polymeric support is termed “soluble” if a polymer, or a polymer-supported compound, is soluble under the conditions employed. However, in general, a soluble polymer can be rendered insoluble under defined conditions. Accordingly, a polymeric support can be soluble under certain conditions and insoluble under other conditions.

[0042] The term “linker,” as used herein, refers to a chemical moiety utilized to attach a compound of interest to a solid support to facilitate synthesis of inventive compounds or a linker may attach one portion of a compound to another portion of a compound. Preferably, the linker comprises covalent bonds. Exemplary linkers are described herein. It will be appreciated that other linkers that are known in the art can also be employed for the synthesis of the compounds of the invention.

[0043] “Compound”: The term “compound” or “chemical compound” as used herein can include organometallic compounds, organic compounds, metals, transitional metal complexes, and small molecules. In certain preferred embodiments, polynucleotides are excluded from the definition of compounds. In other preferred embodiments, polynucleotides and peptides are excluded from the definition of compounds. In a particularly preferred embodiment, the term compounds refers to small molecules (*e.g.*, preferably, non-peptidic and non-oligomeric) and excludes peptides, polynucleotides, transition metal complexes, metals, and organometallic compounds.

[0044] “Small Molecule”: As used herein, the term “small molecule” refers to a non-peptidic, non-oligomeric organic compound either synthesized in the laboratory or found in nature. Small molecules, as used herein, can refer to compounds that are “natural product-like”, however, the term “small molecule” is not limited to “natural product-like” compounds. Rather, a small molecule is typically characterized in that it contains several carbon-carbon bonds, and has a molecular weight of less than 1500, although this characterization is not intended to be limiting for the purposes of the present invention. Examples of “small molecules” that occur in nature include, but are not limited to, taxol,

dynemicin, and rapamycin. Examples of “small molecules” that are synthesized in the laboratory include, but are not limited to, compounds described in Tan *et al.*, (“Stereoselective Synthesis of over Two Million Compounds Having Structural Features Both Reminiscent of Natural Products and Compatible with Miniaturized Cell-Based Assays” *J. Am. Chem. Soc.* 120:8565, 1998. In certain other preferred embodiments, natural-product-like small molecules are utilized.

[0045] “Natural Product-Like Compound”: As used herein, the term “natural product-like compound” refers to compounds that are similar to complex natural products which nature has selected through evolution. Typically, these compounds contain one or more stereocenters, a high density and diversity of functionality, and a diverse selection of atoms within one structure. In this context, diversity of functionality can be defined as varying the topology, charge, size, hydrophilicity, hydrophobicity, and reactivity to name a few, of the functional groups present in the compounds. The term, “high density of functionality”, as used herein, can preferably be used to define any molecule that contains preferably three or more latent or active diversifiable functional moieties. These structural characteristics may additionally render the inventive compounds functionally reminiscent of complex natural products, in that they may interact specifically with a particular biological receptor, and thus may also be functionally natural product-like.

[0046] “Metal chelator”: As used herein, the term “metal chelator” refers to any molecule or moiety that is capable of forming a complex (*i.e.*, “chelates”) with a metal ion. In certain exemplary embodiments, a metal chelator refers to any molecule or moiety that “binds” to a metal ion, in solution, making it unavailable for use in chemical/enzymatic reactions. In certain embodiments, the solution comprises aqueous environments under physiological conditions. Examples of metal ions include, but are not limited to, Ca^{2+} , Fe^{3+} , Zn^{2+} , Na^+ , *etc.* In certain embodiments, the metal chelator bind Zn^{2+} , which is found at the active site of HDACs. In certain embodiments, molecules of moieties that precipitate metal ions are not considered to be metal chelators.

[0047] As used herein the term “biological sample” includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from an animal (*e.g.*, mammal) or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof. For example, the term “biological sample” refers to any solid or fluid sample obtained from, excreted by or secreted by any living organism, including single-celled micro-organisms (such as bacteria and yeasts) and multicellular organisms (such as plants and animals, for instance a vertebrate or a mammal, and in particular a healthy or apparently

healthy human subject or a human patient affected by a condition or disease to be diagnosed or investigated). The biological sample can be in any form, including a solid material such as a tissue, cells, a cell pellet, a cell extract, cell homogenates, or cell fractions; or a biopsy, or a biological fluid. The biological fluid may be obtained from any site (e.g. blood, saliva (or a mouth wash containing buccal cells), tears, plasma, serum, urine, bile, cerebrospinal fluid, amniotic fluid, peritoneal fluid, and pleural fluid, or cells therefrom, aqueous or vitreous humor, or any bodily secretion), a transudate, an exudate (e.g. fluid obtained from an abscess or any other site of infection or inflammation), or fluid obtained from a joint (e.g. a normal joint or a joint affected by disease such as rheumatoid arthritis, osteoarthritis, gout or septic arthritis). The biological sample can be obtained from any organ or tissue (including a biopsy or autopsy specimen) or may comprise cells (whether primary cells or cultured cells) or medium conditioned by any cell, tissue or organ. Biological samples may also include sections of tissues such as frozen sections taken for histological purposes. Biological samples also include mixtures of biological molecules including proteins, lipids, carbohydrates and nucleic acids generated by partial or complete fractionation of cell or tissue homogenates. Although the sample is preferably taken from a human subject, biological samples may be from any animal, plant, bacteria, virus, yeast, etc. The term *animal*, as used herein, refers to humans as well as non-human animals, at any stage of development, including, for example, mammals, birds, reptiles, amphibians, fish, worms and single cells. Cell cultures and live tissue samples are considered to be pluralities of animals. In certain exemplary embodiments, the non-human animal is a mammal (e.g., a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a sheep, cattle, a primate, or a pig). An animal may be a transgenic animal or a human clone. If desired, the biological sample may be subjected to preliminary processing, including preliminary separation techniques.

Brief Description of the Drawing

[0048] *Figure 1* shows examples from the three classes of inventive compounds—tubacin, isotubacin, and isoisotubacin.

[0049] *Figure 2* shows a general synthetic scheme for compounds of the “isotubacin” class.

[0050] *Figure 3* shows an exemplary synthesis of isotubacin.

[0051] *Figure 4* shows a general synthetic scheme for compounds of the “isoisotubacin” class.

[0052] *Figure 5* shows an exemplary synthesis of isoisotubacin.

[0053] *Figure 6* shows exemplary epoxide-opening reactions useful in preparing various analogs of the inventive compounds. The scheme illustrates the use of various nucleophiles to open the epoxide group to create the diol functionality later capped to create the isotubacin and isoisotubacin structures.

[0054] *Figure 7* graphically depicts protein degradation pathways and the scientific rationale for combining bortezomib (VELCADE®) with HDAC6 inhibitors (*e.g.*, isotubacin, isoisotubacin) in the treatment of protein degradation disorders. There are two pathways which degrade misfolded/unfolded proteins which are ubiquitinated. The former is the proteasome pathway, and the latter is the aggresome pathway, which requires HDAC 6 activity. Therefore inhibition of both pathways by specific inhibitors, bortezomib (VELCADE®), and isotubacin or isoisotubacin, induced accumulation of cytotoxic misfolded/unfolded proteins..

[0055] *Figure 8* is a schematic of the high-throughput immunofluorescence quantitative assay for acetylated tubulin versus acetylated lysine (as an indicator of acetylated histones) with resulting images.

[0056] *Figure 9* shows the chemical structure for Isotubacin (NKI-93-1). The 1,3-dioxane core in tubacin derivatives is rotated 120° counter-clockwise to yield isotubacin.

[0057] *Figure 10* shows the synergy between isotubacin (NKI-93-1) and bortezomib (VELCADE®) in myeloma cell lines (A) MM.1S, and (B) RPMI cells.

[0058] *Figure 11* demonstrates the specificity of Isotubacin (NKI-93-1) for tubulin acetylation versus lysine acetylation.

[0059] *Figure 12* shows the TDAC inhibitory activity of the compounds—tubacin, NKI-82-1, NKI-81-1, isotubacin (NKI-93-1), NKI-94-1, NKI-59-1-, NKI-60-1, DHM-Tubacin, and MAZ-1428.

[0060] *Figure 13* is a chart showing the HDAC inhibition and TDAC inhibition of the compounds—tubacin, DHM-Tubacin, NKI-59-1, NKI-60-1, NKI-82-1, NKI-84-1, NKI-94-1, NKI-81-1, and isotubacin (NKI-93-1).

[0061] *Figure 14* shows the binding of various compounds including isotubacin (NKI-93-1) to HSA.

Detailed Description of the Invention

[0062] As discussed above, there remains a need for the development of novel inhibitors of histone deacetylases, tubulin histone deacetylases, and the aggresome. In particular, inhibitors that are more potent and/or more specific for their particular target than known

HDAC and TDAC inhibitors. HDAC inhibitors specific for a certain class or member of the HDAC family would be particularly useful both in the treatment of proliferative diseases and protein deposition disorders and in the study of HDACs. Inhibitors that are specific for HDAC versus TDAC and vice versa are also useful in treating disease and probing biological pathways. The present invention provides novel compounds, methods for the synthesis thereof, pharmaceutical compositions thereof, and methods of using these compounds to treat cancer, proliferative diseases, protein degradation disorders, and protein deposition disorders.


Compounds of the Invention

[0063] As discussed above, the present invention provides compounds useful for the treatment of various diseases. In certain embodiments, the compounds of the present invention are useful as inhibitors of histone or tubulin deacetylases and thus are useful as anti-cancer agents, and thus may be useful in the treatment of cancer, by effecting tumor cell death or inhibiting the growth of tumor cells. In certain exemplary embodiments, the inventive anticancer agents are useful in the treatment of cancers and other proliferative disorders, including, but not limited to breast cancer, cervical cancer, colon and rectal cancer, leukemia, lung cancer, melanoma, multiple myeloma, non-Hodgkin's lymphoma, ovarian cancer, pancreatic cancer, prostate cancer, and gastric cancer, to name a few. In certain embodiments, the inventive anticancer agents are active against leukemia cells and melanoma cells, and thus are useful for the treatment of leukemias (*e.g.*, myeloid, lymphocytic, myelocytic and lymphoblastic leukemias) and malignant melanomas. In certain embodiments, the compounds are useful in the treatment of multiple myeloma. Additionally, the inventive compounds may also be useful in the treatment of protozoal infections. The inventive compounds are also useful in the treatment of diseases associated with aberrant protein catabolism, for example, protein degradation disorders, disorders associated with misfolded proteins, and protein deposition disorders. In certain embodiments, the compound are useful in the treatment of the protein deposition disorders, Wilson's disease, spinocerebellar ataxia, prion disease, Parkinson's disease, Huntington's disease, familial amyotrophic lateral sclerosis, amyloidosis, Alzheimer's disease, Alexander's deases, alcoholic liver disease, cystic fibrosis, Pick's disease, and Lewy body dementia. In certain exemplary embodiments, the compounds of the invention are useful for disorders associated with histone deacetylation activity. In certain exemplary embodiments, the compounds of the invention are useful for disorders associated with tubulin deacetylation activity. In other exemplary embodiments, the compounds of the invention are useful for disorders associated

with aggresome activity. In certain embodiments, the compounds, particularly compounds with an ester moiety, are useful in treating skin disorders. Exemplary skin disorders that may be treated using certain of the inventive compounds include cutaneous T-cell lymphoma (CTCL), psoriasis, hair loss, dermatitis, *etc.*

[0064] Compounds of this invention comprise those, as set forth above and described herein, and are illustrated in part by the various classes, subclasses, subgenera, and species disclosed herein. The invention provides compounds, *e.g.*, compounds useful in the methods, pharmaceutical compositions, kits, and packaged compositions of the invention. The inventive compounds are inhibitors of histone deacetylases, tubulin deacetylases, or the aggresome. The compounds of the invention are typically based on a 1,3-dioxane core structure.

[0065] Exemplary classes of compounds of the invention include compounds of the formula:



wherein

R_1 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_A$; $-C(=O)R_A$; $-CO_2R_A$; $-CN$; $-SCN$; $-SR_A$; $-SOR_A$; $-SO_2R_A$; $-NO_2$; $-N(R_A)_2$; $-NHC(O)R_A$; or $-C(R_A)_3$; wherein each occurrence of R_A is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;


R_2 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_B$; $-C(=O)R_B$; $-CO_2R_B$; $-CN$; $-SCN$; $-SR_B$; $-SOR_B$; $-SO_2R_B$; $-NO_2$; $-N(R_B)_2$; $-NHC(O)R_B$; or $-C(R_B)_3$; wherein each occurrence of R_B is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety;

an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety; and

R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety; and pharmaceutically acceptable salts and derivatives thereof. In certain embodiments, R_1 comprises a metal chelating functional group (*e.g.*, hydroxyamic acids, thiols, carboxylic acids, *ortho*-aminoanilides, *etc.*).

[0066] As will be appreciated by one of skill in this art, the 1,3-dioxane core has been rotated 120° in each of the two classes of compounds as compared to the 1,3-dioxane core of tubacin and its derivatives.


[0067] The invention also provides compounds of the isotubacin class. These compounds are of the formula:




A general synthetic scheme for preparing compounds of this class is shown in *Figure 2*.


Figure 3 is a synthetic scheme showing the synthesis of isotubacin.

[0068] In certain embodiments, the compounds of this second class are one of the formulae below with the stereochemistry as shown:




[0069] In certain embodiments, R_1 is a substituted phenyl ring. In certain particular embodiments, R_1 is of the formula:




wherein R_1' is , wherein Y is NH or O; L is a linker moiety; and A comprises a functional group that inhibits histone or tubulin deacetylase.

[0070] In certain embodiments, R_1 is of the formula:




[0071] In other embodiments, R_1 is of the formula:




In certain embodiments, Y is NH. In other embodiments, Y is O. In certain embodiments, L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety; a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety; a substituted or unsubstituted aryl moiety; a substituted or unsubstituted heteroaryl moiety. In certain embodiments, L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety. In certain embodiments, L is C_1 - C_{20} alkylidene, preferably C_1 to C_{12} alkylidene, more preferably C_4 - C_7 alkylidene. In certain embodiments, L is C_1 - C_{20} alkenylidene, preferably C_1 to C_{12} alkenylidene, more preferably C_4 - C_7 alkenylidene. In certain embodiments, L is C_1 - C_{20} alkynylidene, preferably C_1 to C_{12} alkynylidene, more preferably C_4 - C_7 alkynylidene. In certain embodiments, L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety. In certain embodiments, L comprises a cyclic ring system, wherein the rings may be aryl, heteroaryl, non-aromatic carbocyclic, or non-aromatic heterocyclic. In still other


embodiments, L comprises a substituted or unsubstituted heteroaryl moiety. In certain particular embodiments, L comprises a phenyl ring. In certain embodiments, L comprises multiple phenyl rings (*e.g.*, one, two, three, or four phenyl rings).


[0072] In certain embodiments, L is , wherein n is an integer between 1 and 4, inclusive; preferably, between 1 and 3, inclusive; more preferably, 1 or 2; and R₁ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; -OR_A; -C(=O)R_A; -CO₂R_A; -CN; -SCN; -SR_A; -SOR_A; -SO₂R_A; -NO₂; -N(R_A)₂; ; -NHR_A; -NHC(O)R_A; or -C(R_A)₃; wherein each occurrence of R_A is independently a hydrogen, a protecting group, an aliphatic moiety; a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, L is





[0073] In certain embodiments, L is .

[0074] In certain embodiments, L is an unbranched, unsubstituted, acyclic alkyl chain. In


certain embodiments, L is . In other embodiments, L is

. In certain other embodiments, L is


. In other embodiments, L is .


In yet other embodiments, L is .


[0075] In certain embodiments, L is a substituted, acyclic aliphatic chain. In certain


embodiments, L is .


[0076] In certain embodiments, L is an unbranched, unsubstituted, acyclic heteroaliphatic


chain. In certain particular embodiments, L is , wherein n is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; and m is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive. In certain particular


embodiments, L is , wherein n is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; and m is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive. In certain particular embodiments, L is

, wherein n is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; m is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; and R' is hydrogen, C₁-C₆ aliphatic, heteroaliphatic, aryl, heteroaryl, or acyl. In


certain particular embodiments, L is , wherein n is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; and m is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive. In certain embodiments, A comprises a metal chelating functional group. For example, A comprises a Zn²⁺ chelating group. In certain embodiments, A comprises a functional group selected group consisting of:




In certain embodiments, A comprises hydroxamic acid () or a salt thereof. In other embodiments, A comprises the formula:




In certain particular embodiments, A comprises the formula:




In other embodiments, A comprises a carboxylic acid ($-\text{CO}_2\text{H}$). In other embodiments, A





comprises an *o*-aminoanilide (NH_2). In other embodiments, A comprises an





o-hydroxyanilide (OH). In yet other embodiments, A comprises a thiol ($-\text{SH}$).




In certain embodiments, R_1' is , wherein n is an integer between 0 and 15, inclusive; preferably, between 0 and 10, inclusive; more preferably, between 1 and 8, inclusive; even more preferably, 4, 5, 6, 7, or 8. In certain embodiments,




R_1' is , wherein n is an integer between 0 and 15, inclusive; preferably, between 0 and 10, inclusive; more preferably, between 1 and 8, inclusive; even more preferably, 4, 5, 6, 7, or 8. In certain embodiments, R_1' is




. In other particular embodiments, R_1' is





[0077] In certain embodiments, R_2 is hydrogen. In other embodiments, R_2 is hydroxyl or a protected hydroxyl group. In certain embodiments, R_2 is alkoxy. In yet other embodiments, R_2 is a lower alkyl, alkenyl, or alkynyl group. In certain embodiments, R_2 is $-(CH_2)_m-X(R_B)_n$, wherein X is O, S, N, or C, preferably O, S, or N; n is 1, 2, or 3; and m is an integer between 1 and 6, inclusive. In certain embodiments, R_2 is $-CH_2-X(R_B)_n$, wherein X is O, S, N, or C, preferably O, S, or N; and n is 1, 2, or 3. In certain embodiments, R_2 is $-CH_2-OR_B$. In other embodiments, R_2 is $-CH_2-SR_B$. In yet other embodiments, R_2 is $-CH_2-R_B$. In other embodiments, R_2 is $-CH_2-N(R_B)_2$. In still other embodiments, R_2 is $-CH_2-NHR_B$. In certain embodiments of the invention, R_B is one of:






wherein m and p are each independently integers from 0 to 3; q_1 is an integer from 1 to 6; R^{2C} is hydrogen, lower alkyl or a nitrogen protecting group; and each occurrence of R^{2B} is independently hydrogen, halogen, -CN, or WR^{W1} wherein W is O, S, NR^{W2} , -C(=O), -S(=O), -SO₂, -C(=O)O-, -OC(=O), -C(=O)NR^{W2}, -NR^{W2}C(=O); wherein each occurrence of R^{W1} and R^{W2} is independently hydrogen, a protecting group, a prodrug moiety or an alkyl, cycloalkyl, heteroalkyl, heterocyclic, aryl or heteroaryl moiety, or, when W is NR^{W2} , R^{W1} and R^{W2} , taken together with the nitrogen atom to which they are attached, form a heterocyclic or heteroaryl moiety; or any two adjacent occurrences of R^{2B} , taken together with the atoms to which they are attached, form a substituted or unsubstituted, saturated or unsaturated alicyclic or heterocyclic moiety, or a substituted or unsubstituted aryl or heteroaryl moiety. In certain embodiments of the invention, R_B is one of the structures:







wherein m is an integer from 1 to 4; R^{2C} is hydrogen, lower alkyl or a nitrogen protecting group; and each occurrence of R^{2B} is independently hydrogen, halogen, $-CN$, or WR^{W1} wherein W is O , S , NR^{W2} , $-C(=O)$, $-S(=O)$, $-SO_2$, $-C(=O)O-$, $-OC(=O)$, $-C(=O)NR^{W2}$, $-NR^{W2}C(=O)$; wherein each occurrence of R^{W1} and R^{W2} is independently hydrogen, a protecting group, a prodrug moiety or an alkyl, cycloalkyl, heteroalkyl, heterocyclic, aryl or heteroaryl moiety, or, when W is NR^{W2} , R^{W1} and R^{W2} , taken together with the nitrogen atom to which they are attached, form a heterocyclic or heteroaryl moiety; or any two adjacent occurrences of R^{2B} , taken together with the atoms to which they are attached, form a substituted or unsubstituted, saturated or unsaturated alicyclic or heterocyclic moiety, or a substituted or unsubstituted aryl or heteroaryl moiety.

[0078] In certain embodiments, $-X(R_B)_n$ of $-CH_2-X(R_B)_n$ or $-(CH_2)_m-X(R_B)_n$ has one of the structures:









[0079] In certain embodiments, R_2 is

, wherein X is N




and Y is NH, S, or O. In other embodiments, R_2 is

[0080] In certain embodiments, R_2 is selected from one of the following:




[0081] In certain embodiments, R_3 is aliphatic. In other embodiments, R_3 is heteroaliphatic. In certain embodiments, R_3 is a substituted or unsubstituted aryl moiety. In certain embodiments, R_3 is a substituted or unsubstituted heteroaromatic moiety. In certain embodiments, R_3 is a monocyclic moiety. In other embodiments, R_3 is a bicyclic moiety. In yet other embodiments, R_3 is a tricyclic moiety. In yet other embodiments, R_3 is a polycyclic moiety. In certain embodiments, R_3 is a substituted or unsubstituted five- or six-membered aromatic or heteroaromatic moiety. In certain embodiments, R_3 is a substituted or


unsubstituted six-membered aromatic or heteroaromatic moiety. In certain embodiments, R_3 is a substituted or unsubstituted six-membered aromatic moiety. In certain embodiments, R_3 is a substituted or unsubstituted six-membered heteroaromatic moiety. In certain embodiments, R_3 is a substituted or unsubstituted non-aromatic carbocyclic or heterocyclic moiety. In certain embodiments, R_3 is substituted or unsubstituted aryl. In certain embodiments, R_3 is substituted or unsubstituted phenyl. In certain embodiments, R_3 is




. In certain particular embodiments, R_3 is monosubstituted phenyl. In certain embodiments, R_3 is para-substituted phenyl. In certain embodiments, R_3 is



, wherein R_3' is hydrogen, a protecting group, a solid support unit, an alkyl, acyl, cycloalkyl, heteroalkyl, heterocyclic, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety. In certain




embodiments, R_3 is . In certain embodiments, R_3 is not



. In other embodiments, R_3 is substituted or unsubstituted heteroaryl.

[0082] In certain embodiments, the invention provides compounds of the formula:



wherein R_1 and R_2 are defined as above;


n is an integer between 1 and 5, inclusive; and

each occurrence of R_3' is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl;

substituted or unsubstituted, branched or unbranched heteroaryl; $-\text{OR}_C$; $-\text{C}(=\text{O})\text{R}_C$; $-\text{CO}_2\text{R}_C$; $-\text{CN}$; $-\text{SCN}$; $-\text{SR}_C$; $-\text{SOR}_C$; $-\text{SO}_2\text{R}_C$; $-\text{NO}_2$; $-\text{N}(\text{R}_C)_2$; $-\text{NHC}(\text{O})\text{R}_C$; or $-\text{C}(\text{R}_C)_3$; wherein each occurrence of R_C is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety.


[0083] In certain embodiments, n is 0, and the phenyl ring is unsubstituted.

[0084] In other embodiments, n is 1, and the compounds are one of the formulae:



In certain embodiments, the para-substitution pattern is preferred. In other embodiments, the meta-substitution pattern is preferred. And in yet other embodiments, the ortho-substitution pattern is preferred.


[0085] In other embodiments, n is 2. Compounds of the invention include compounds of one of the formulae:







[0086] In other embodiments, n is 3. In still other embodiments, n is 4, and in other embodiments, n is 5.

[0087] In certain embodiments, R_3' is halogen, hydroxyl, protected hydroxyl, alkoxy, amino, alkylamino, dialkylamino, $-\text{NO}_2$, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkenyl, $\text{C}_1\text{-C}_6$ alkynyl, or acyl. In certain embodiments, R_3' is $-\text{NO}_2$. In certain embodiments, R_3' is $-\text{CH}_2\text{OH}$. In certain embodiments, R_3' is $-\text{NH}_2$. In certain embodiments, R_3' is $-\text{H}$. In other embodiments, R_3' is $-\text{OH}$. In other embodiments, R_3' is $-\text{CN}$. In yet other embodiments, R_3' is $-\text{SCN}$. In still other embodiments, R_3' is acyl. In certain embodiments, R_3' is acetyl. In other embodiments, R_3' is $-\text{F}$. In other embodiments, R_3' is $-\text{Cl}$. In other embodiments, R_3' is $-\text{Br}$. In other embodiments, R_3' is $-\text{I}$. In other embodiments, R_3' is methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *tert*-butyl, or *iso*-butyl. In certain embodiments, R_3' is vinyl. In certain embodiments, R_3' is halogen-substituted alkyl (*e.g.*, trifluoromethyl). In certain embodiments, R_3' is methoxy, ethoxy, propoxy, butoxy, or pentoxy.


[0088] Exemplary compounds of this second "isotubacin" class include compounds of the formula:








[0089] The invention also provides compounds of the isoiso-tubacin class. These compounds are of the formula:





A general synthetic scheme for preparing compounds of this class is shown in *Figure 4*.

Figure 5 is a synthetic scheme showing the synthesis of isoisoisotubacin.

[0090] In certain embodiments, the compounds of this second class are one of the formulae below with the stereochemistry as shown:




[0091] In certain embodiments, R₁ is a substituted phenyl ring. In certain particular embodiments, R₁ is of the formula:




wherein R₁' is , wherein Y is NH or O; L is a linker moiety; and A comprises a functional group that inhibits histone deacetylase.

[0092] In certain embodiments, R₁ is of the formula:



[0093] In other embodiments, R₁ is of the formula:



In certain embodiments, Y is NH. In other embodiments, Y is O. In certain embodiments, L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety; a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety; a substituted or unsubstituted aryl moiety; a substituted or unsubstituted heteroaryl moiety. In certain embodiments, L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety. In certain embodiments, L is C₁-C₂₀ alkylidene, preferably C₁ to C₁₂ alkylidene, more preferably C₄-C₇ alkylidene. In certain embodiments, L is C₁-C₂₀ alkenylidene, preferably C₁ to C₁₂ alkenylidene, more preferably C₄-C₇ alkenylidene. In certain embodiments, L is C₁-C₂₀ alkynylidene, preferably C₁ to C₁₂ alkynylidene, more preferably C₄-C₇ alkynylidene. In certain embodiments, L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety. In certain embodiments, L comprises a cyclic ring system, wherein the rings may be aryl, heteroaryl, non-aromatic carbocyclic, or non-aromatic heterocyclic. In still other embodiments, L comprises a substituted or unsubstituted heteroaryl moiety. In certain particular embodiments, L comprises a phenyl ring. In certain embodiments, L comprises multiple phenyl rings (*e.g.*, one, two, three, or four phenyl rings).




[0094] In certain embodiments, L is , wherein n is an integer between 1 and 4, inclusive; preferably, between 1 and 3, inclusive; more preferably, 1 or 2; and R₁ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; -OR_A; -C(=O)R_A; -CO₂R_A; -CN; -SCN; -SR_A; -SOR_A; -SO₂R_A; -NO₂; -N(R_A)₂; -NHR_A; -NHC(O)R_A; or -C(R_A)₃; wherein each occurrence of R_A is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino,


dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, L is



[0095] In certain embodiments, L is




[0096] In certain embodiments, L is an unbranched, unsubstituted, acyclic alkyl chain. In

certain embodiments, L is . In other embodiments, L is



. In certain other embodiments, L is




. In other embodiments, L is





In yet other embodiments, L is .


[0097] In certain embodiments, L is a substituted, acyclic aliphatic chain. In certain

embodiments, L is .


[0098] In certain embodiments, L is an unbranched, unsubstituted, acyclic heteroaliphatic

chain. In certain particular embodiments, L is , wherein n is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; and m is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive. In certain particular


embodiments, L is , wherein n is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; and m is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive. In certain particular embodiments, L is

, wherein n is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; m is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; and R' is hydrogen, C₁-C₆ aliphatic, heteroaliphatic, aryl, heteroaryl, or acyl. In


certain particular embodiments, L is , wherein n is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; and m is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive. In certain embodiments, A comprises a metal chelating functional group. For example, A comprises a Zn²⁺ chelating group. In certain embodiments, A comprises a functional group selected group consisting of:




In certain embodiments, A comprises hydroxamic acid () or a salt thereof. In other embodiments, A comprises the formula:





In certain particular embodiments, A comprises the formula:





In other embodiments, A comprises a carboxylic acid ($-\text{CO}_2\text{H}$). In other embodiments, A


comprises an *o*-aminoanilide (). In other embodiments, A comprises an

o-hydroxyanilide (). In yet other embodiments, A comprises a thiol (-


SH). In certain embodiments, R_1' is  , wherein n is an integer between 0 and 15, inclusive; preferably, between 0 and 10, inclusive; more preferably, between 1 and 8, inclusive; even more preferably, 4, 5, 6, 7, or 8. In certain embodiments,


R_1' is  , wherein n is an integer between 0 and 15, inclusive; preferably, between 0 and 10, inclusive; more preferably, between 1 and 8, inclusive; even more preferably, 4, 5, 6, 7, or 8. In certain embodiments, R_1' is

 . In other particular embodiments, R_1' is




[0099] In certain embodiments, R_2 is hydrogen. In other embodiments, R_2 is hydroxyl or a protected hydroxyl group. In certain embodiments, R_2 is alkoxy. In yet other embodiments, R_2 is a lower alkyl, alkenyl, or alkynyl group. In certain embodiments, R_2 is $-\text{CH}_2-\text{X}(\text{R}_\text{B})_n$, wherein X is O, S, N, or C, preferably O, S, or N; and n is 1, 2, or 3. In certain embodiments, R_2 is $-\text{CH}_2-\text{OR}_\text{B}$. In other embodiments, R_2 is $-\text{CH}_2-\text{SR}_\text{B}$. In yet other embodiments, R_2 is $-\text{CH}_2-\text{R}_\text{B}$. In other embodiments, R_2 is $-\text{CH}_2-\text{N}(\text{R}_\text{B})_2$. In still other embodiments, R_2 is $-\text{CH}_2-\text{NHR}_\text{B}$. In certain embodiments of the invention, R_B is one of:






wherein m and p are each independently integers from 0 to 3; q_1 is an integer from 1 to 6; R^{2C} is hydrogen, lower alkyl or a nitrogen protecting group; and each occurrence of R^{2B} is independently hydrogen, halogen, $-CN$, or WR^{W1} wherein W is O , S , NR^{W2} , $-C(=O)$, $-S(=O)$, $-SO_2$, $-C(=O)O-$, $-OC(=O)$, $-C(=O)NR^{W2}$, $-NR^{W2}C(=O)$; wherein each occurrence of R^{W1} and R^{W2} is independently hydrogen, a protecting group, a prodrug moiety or an alkyl, cycloalkyl, heteroalkyl, heterocyclic, aryl or heteroaryl moiety, or, when W is NR^{W2} , R^{W1} and R^{W2} , taken together with the nitrogen atom to which they are attached, form a heterocyclic or heteroaryl moiety; or any two adjacent occurrences of R^{2B} , taken together with the atoms to which they are attached, form a substituted or unsubstituted, saturated or unsaturated alicyclic or heterocyclic moiety, or a substituted or unsubstituted aryl or heteroaryl moiety. In certain embodiments of the invention, R_B is one of the structures:




wherein m is an integer from 1 to 4; R^{2C} is hydrogen, lower alkyl or a nitrogen protecting group; and each occurrence of R^{2B} is independently hydrogen, halogen, $-CN$, or WR^{W1} wherein W is O , S , NR^{W2} , $-C(=O)$, $-S(=O)$, $-SO_2$, $-C(=O)O-$, $-OC(=O)$, $-C(=O)NR^{W2}$, $-NR^{W2}C(=O)$; wherein each occurrence of R^{W1} and R^{W2} is independently hydrogen, a protecting group, a prodrug moiety or an alkyl, cycloalkyl, heteroalkyl, heterocyclic, aryl or heteroaryl moiety, or, when W is NR^{W2} , R^{W1} and R^{W2} , taken together with the nitrogen atom to which they are attached, form a heterocyclic or heteroaryl moiety; or any two adjacent

[00100] In certain embodiments, $-X(R_B)_n$ has one of the structures:









[00103] In certain embodiments, R_2 is a substituted or unsubstituted aryl or heteroaryl moiety. In certain embodiments, R_2 is a substituted or unsubstituted carbocyclic or heterocyclic moiety. The ring system of R_2 may be monocyclic, bicyclic, tricyclic, or polycyclic. The rings making up the ring system may be three-membered (e.g., oxiranyl, cyclopropyl, aziridinyl); four-membered; five-membered; six-membered; seven-membered; eight-membered; or n-membered. In bicyclic, tricyclic, or polycyclic ring systems, the rings may be fused, spiro-linked, or linked via a covalent bond. In certain embodiments, R_2 is a


substituted or unsubstituted, monocyclic aryl moiety. In certain embodiments, R_2 is a substituted or unsubstituted, monocyclic phenyl moiety. In certain embodiments, R_2 is a substituted, monocyclic phenyl moiety. In certain embodiments, R_2 is unsubstituted phenyl. In certain embodiments, R_2 is a monosubstituted phenyl moiety. In other embodiments, R_2 is a disubstituted phenyl moiety. In yet other embodiments, R_2 is a trisubstituted phenyl moiety. In other embodiments, R_2 is a substituted or unsubstituted, monocyclic heteroaryl moiety. In certain embodiments, R_2 is a substituted or unsubstituted pyridinyl moiety. In certain embodiments, R_2 is a substituted or unsubstituted pyrrolyl moiety. In certain embodiments, R_2 is a substituted or unsubstituted imidazolyl moiety. In certain embodiments, R_2 is a substituted or unsubstituted thiazolyl moiety. In certain embodiments, R_2 is a substituted or unsubstituted oxazolyl moiety. In certain embodiments, R_2 is a substituted or unsubstituted furanyl moiety. In certain embodiments, R_2 is a substituted or unsubstituted thiophenyl moiety. In certain embodiments, R_2 is a substituted or unsubstituted, monocyclic carbocyclic moiety. In certain embodiments, R_2 is a substituted or unsubstituted, cyclopentyl moiety. In certain embodiments, R_2 is a substituted or unsubstituted, cyclohexyl moiety. In certain embodiments, R_2 is a substituted or unsubstituted, monocyclic heterocyclic moiety. In certain embodiments, R_2 is a substituted or unsubstituted, piperidinyl moiety. In certain embodiments, R_2 is a substituted or unsubstituted, pyrrolidinyl moiety.


[00104] In certain embodiments, R_3 is a substituted or unsubstituted aryl moiety. In certain embodiments, R_3 is a substituted or unsubstituted heteroaromatic moiety. In certain embodiments, R_3 is a monocyclic moiety. In other embodiments, R_3 is a bicyclic moiety. In yet other embodiments, R_3 is a tricyclic moiety. In yet other embodiments, R_3 is a polycyclic moiety. In certain embodiments, R_3 is a substituted or unsubstituted five- or six-membered aromatic or heteroaromatic moiety. In certain embodiments, R_3 is a substituted or unsubstituted six-membered aromatic or heteroaromatic moiety. In certain embodiments, R_3 is a substituted or unsubstituted six-membered aromatic moiety. In certain embodiments, R_3 is a substituted or unsubstituted six-membered heteroaromatic moiety. In certain embodiments, R_3 is a substituted or unsubstituted non-aromatic carbocyclic or heterocyclic moiety. In certain embodiments, R_3 is unsubstituted aryl. In certain embodiments, R_3 is substituted aryl. In certain embodiments, R_3 is substituted or unsubstituted phenyl. In certain particular embodiments, R_3 is monosubstituted phenyl. In certain embodiments, R_3 is



. In certain embodiments, R_3 is para-substituted phenyl. In certain


embodiments, R_3 is , wherein R_3' is hydrogen, a protecting group, a solid support unit, an alkyl, acyl, cycloalkyl, heteroalkyl, heterocyclic, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety. In

certain embodiments, R_3 is . In certain embodiments, R_3 is not



. In other embodiments, R_3 is substituted or unsubstituted heteroaryl.

[00105] In certain embodiments, the invention provides compounds of the formula:




wherein R_1 and R_2 are defined as above;

n is an integer between 1 and 5, inclusive; and

each occurrence of R_3' is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety.


[00106] In certain embodiments, n is 0, and the phenyl ring is unsubstituted.

[00107] In other embodiments, n is 1, and the compounds are one of the formulae:



In certain embodiments, the para-substitution pattern is preferred. In other embodiments, the meta-substitution pattern is preferred. And in yet other embodiments, the ortho-substitution pattern is preferred.


[00108] In other embodiments, n is 2. Compounds of the invention include compounds of one of the formulae:




[00109] In other embodiments, n is 3. In still other embodiments, n is 4, and in other embodiments, n is 5.


[00110] In certain embodiments, R₃' is halogen, hydroxyl, protected hydroxyl, alkoxy, amino, alkylamino, dialkylamino, -NO₂, C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl, or acyl. In certain embodiments, R₃' is -NO₂. In certain embodiments, R₃' is -CH₂OH. In certain embodiments, R₃' is -NH₂. In certain embodiments, R₃' is -H. In other embodiments, R₃' is -OH. In other embodiments, R₃' is -CN. In yet other embodiments, R₃' is -SCN. In still

[00111] Exemplary compounds of the “isoisotubacin” class include compounds of the formula:







[00112] In another embodiments, the invention provides dimers, trimers, or multimers of HDAC inhibitors described herein. In certain embodiments, dimers are of the general formula,



wherein


each occurrence of R_1 comprises a functional group that inhibits histone deacetylase or tubulin deacetylase, wherein the two R_1 groups may be the same or different. Dimeric, Trimeric, and multimeric HDAC inhibitors are further described in PCT publication no. WO 2008/091349. The tubacin structures in the dimeric compounds therein may be changed to isotubacin and isoisotubacin structures. The dioxane core may also provide for a trimer wherein R_2 is $-R_1$.

[00113] In certain embodiments, R_1 is a substituted phenyl ring. In certain particular embodiments, R_1 is of the formula:




wherein R_1' is , wherein Y is NH or O; L is a linker moiety; and A comprises a functional group that inhibits histone or tubulin deacetylase.

[00114] In certain embodiments, R_1 is of the formula:




[00115] In other embodiments, R_1 is of the formula:



In certain embodiments, Y is NH. In other embodiments, Y is O. In certain embodiments, L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety; a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety; a substituted or unsubstituted aryl moiety; a substituted or unsubstituted heteroaryl moiety. In certain embodiments, L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety. In certain embodiments, L is C₁-C₂₀ alkylidene, preferably C₁ to C₁₂ alkylidene, more preferably C₄-C₇ alkylidene. In certain embodiments, L is C₁-C₂₀ alkenylidene, preferably C₁ to C₁₂ alkenylidene, more preferably C₄-C₇ alkenylidene. In certain embodiments, L is C₁-C₂₀ alkynylidene, preferably C₁ to C₁₂ alkynylidene, more preferably C₄-C₇ alkynylidene. In certain embodiments, L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety. In certain embodiments, L comprises a cyclic ring system, wherein the rings may be aryl, heteroaryl, non-aromatic carbocyclic, or non-aromatic heterocyclic. In still other embodiments, L comprises a substituted or unsubstituted heteroaryl moiety. In certain particular embodiments, L comprises a phenyl ring. In certain embodiments, L comprises multiple phenyl rings (*e.g.*, one, two, three, or four phenyl rings).


[00116] In certain embodiments, L is , wherein n is an integer between 1 and 4, inclusive; preferably, between 1 and 3, inclusive; more preferably, 1 or 2; and R₁ is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; -OR_A; -C(=O)R_A; -CO₂R_A; -CN; -SCN; -SR_A; -SOR_A; -SO₂R_A; -NO₂; -N(R_A)₂; -NHR_A; -NHC(O)R_A; or -C(R_A)₃; wherein each occurrence of R_A is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, L is




[00117] In certain embodiments, L is .

[00118] In certain embodiments, L is an unbranched, unsubstituted, acyclic alkyl chain. In


certain embodiments, L is . In other embodiments, L is

. In certain other embodiments, L is


. In other embodiments, L is .


In yet other embodiments, L is .


[00119] In certain embodiments, L is a substituted, acyclic aliphatic chain. In certain


embodiments, L is .


[00120] In certain embodiments, L is an unbranched, unsubstituted, acyclic heteroaliphatic


chain. In certain particular embodiments, L is , wherein n is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; and m is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive. In certain particular


embodiments, L is , wherein n is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; and m is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive. In certain particular embodiments, L is

, wherein n is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; m is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; and R' is hydrogen, C₁-C₆ aliphatic, heteroaliphatic, aryl, heteroaryl, or acyl. In


certain particular embodiments, L is , wherein n is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive; and m is an integer between 0 and 10, inclusive; preferably, between 0 and 5, inclusive. In certain embodiments, A comprises a metal chelating functional group. For example, A comprises a Zn²⁺ chelating group. In certain embodiments, A comprises a functional group selected group consisting of:




In certain embodiments, A comprises hydroxamic acid () or a salt thereof. In other embodiments, A comprises the formula:





In certain particular embodiments, A comprises the formula:





In other embodiments, A comprises a carboxylic acid ($-\text{CO}_2\text{H}$). In other embodiments, A





comprises an *o*-aminoanilide () . In other embodiments, A comprises an




o-hydroxyanilide () . In yet other embodiments, A comprises a thiol (-

SH). In certain embodiments, R_1' is , wherein n is an integer between 0 and 15, inclusive; preferably, between 0 and 10, inclusive; more preferably, between 1 and 8, inclusive; even more preferably, 4, 5, 6, 7, or 8. In certain embodiments,


R_1' is , wherein n is an integer between 0 and 15, inclusive; preferably, between 0 and 10, inclusive; more preferably, between 1 and 8, inclusive; even more preferably, 4, 5, 6, 7, or 8. In certain embodiments, R_1' is



. In other particular embodiments, R_1' is



[00121] In certain embodiments of the invention, inventive compounds based on the structure of isotubacin are of the formula:





wherein

each occurrence of R_1' is defined as above and both occurrences of R_1' are the same or different; and


R_2 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or


unbranched heteroaryl; $-OR_B$; $-C(=O)R_B$; $-CO_2R_B$; $-CN$; $-SCN$; $-SR_B$; $-SOR_B$; $-SO_2R_B$; $-NO_2$; $-N(R_B)_2$; $-NHC(O)R_B$; or $-C(R_B)_3$; wherein each occurrence of R_B is independently a hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, R_2 is hydrogen. In other embodiments, R_2 is hydroxyl or a protected hydroxyl group. In certain embodiments, R_2 is alkoxy. In yet other embodiments, R_2 is a lower alkyl, alkenyl, or alkynyl group. In certain embodiments, R_2 is $-CH_2-X(R_B)_n$, wherein X is O, S, N, or C, preferably O, S, or N; and n is 1, 2, or 3. In certain embodiments, R_2 is $-CH_2-OR_B$. In other embodiments, R_2 is $-CH_2-SR_B$. In yet other embodiments, R_2 is $-CH_2-R_B$. In other embodiments, R_2 is $-CH_2-N(R_B)_2$. In still other embodiments, R_2 is $-CH_2-NHR_B$. In certain embodiments of the invention, R_B is one of:






wherein m and p are each independently integers from 0 to 3; q_1 is an integer from 1 to 6; R^{2C} is hydrogen, lower alkyl or a nitrogen protecting group; and each occurrence of R^{2B} is independently hydrogen, halogen, $-CN$, or WR^{W1} wherein W is O , S , NR^{W2} , $-C(=O)$, $-S(=O)$, $-SO_2$, $-C(=O)O-$, $-OC(=O)$, $-C(=O)NR^{W2}$, $-NR^{W2}C(=O)$; wherein each occurrence of R^{W1} and R^{W2} is independently hydrogen, a protecting group, a prodrug moiety or an alkyl, cycloalkyl, heteroalkyl, heterocyclic, aryl or heteroaryl moiety, or, when W is NR^{W2} , R^{W1} and R^{W2} , taken together with the nitrogen atom to which they are attached, form a heterocyclic or heteroaryl moiety; or any two adjacent occurrences of R^{2B} , taken together with the atoms to which they are attached, form a substituted or unsubstituted, saturated or unsaturated alicyclic or heterocyclic moiety, or a substituted or unsubstituted aryl or heteroaryl moiety. In certain embodiments of the invention, R_B is one of the structures:







tt




uu



vv




ww




xx

wherein m is an integer from 1 to 4; R^{2C} is hydrogen, lower alkyl or a nitrogen protecting group; and each occurrence of R^{2B} is independently hydrogen, halogen, $-CN$, or WR^{W1} wherein W is O , S , NR^{W2} , $-C(=O)$, $-S(=O)$, $-SO_2$, $-C(=O)O-$, $-OC(=O)$, $-C(=O)NR^{W2}$, $-NR^{W2}C(=O)$; wherein each occurrence of R^{W1} and R^{W2} is independently hydrogen, a protecting group, a prodrug moiety or an alkyl, cycloalkyl, heteroalkyl, heterocyclic, aryl or heteroaryl moiety, or, when W is NR^{W2} , R^{W1} and R^{W2} , taken together with the nitrogen atom to which they are attached, form a heterocyclic or heteroaryl moiety; or any two adjacent occurrences of R^{2B} , taken together with the atoms to which they are attached, form a substituted or unsubstituted, saturated or unsaturated alicyclic or heterocyclic moiety, or a substituted or unsubstituted aryl or heteroaryl moiety.

[00122] In certain embodiments, $-X(R_B)_n$ of $-CH_2-X(R_B)_n$ has one of the structures:









[00123] In certain embodiments, R_2 is

, wherein X is N




and Y is NH, S, or O. In other embodiments, R_2 is

[00124] In certain embodiments of the invention, the stereochemistry of formula is chosen from one of the following:



[00125] In certain embodiments of the invention, compounds are of the formula:




wherein

A, B, and R_2 are defined as above;

X is O or NH;

n is an integer between 1 and 20, inclusive; preferably, between 1 and 12, inclusive; more preferably between 2 and 8, inclusive. In certain embodiments, n is 2, 3, 4, 5, 6, 7, or 8; preferably, 6. In certain embodiments, X is NH. In other embodiments, X is O.

[00126] In certain embodiments of the invention, compounds are of the formula:




wherein

R_2 is defined as above;

n is an integer between 1 and 20, inclusive; preferably, between 1 and 12, inclusive; more preferably between 2 and 8, inclusive. In certain embodiments, n is 2, 3, 4, 5, 6, 7, or 8; preferably, 6.


[00127] In certain embodiments of the invention, compounds are of the formula:



wherein

R_2 is defined as above.


[00128] In certain embodiments of the invention, compounds are of the formula:




wherein

R_2 is defined as above.

[00129] In certain embodiments of the invention, compounds are of the formula:



[00130] In certain embodiments of the invention, inventive compounds based on the structure of isoisotubacin are of the formula:





wherein

each occurrence of R_1' is independently as defined above and both occurrences of R_1' are the same or different; and


R_2 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted, branched or unbranched aryl; substituted or unsubstituted, branched or unbranched heteroaryl; $-OR_B$; $-C(=O)R_B$; $-CO_2R_B$; $-CN$; $-SCN$; $-SR_B$; $-SOR_B$; $-SO_2R_B$; $-NO_2$; $-N(R_B)_2$; $-NHC(O)R_B$; or $-C(R_B)_3$; wherein each occurrence of R_B is independently a


hydrogen, a protecting group, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety. In certain embodiments, R_2 is hydrogen. In other embodiments, R_2 is hydroxyl or a protected hydroxyl group. In certain embodiments, R_2 is alkoxy. In yet other embodiments, R_2 is a lower alkyl, alkenyl, or alkynyl group. In certain embodiments, R_2 is $-\text{CH}_2-\text{X}(\text{R}_\text{B})_n$, wherein X is O, S, N, or C, preferably O, S, or N; and n is 1, 2, or 3. In certain embodiments, R_2 is $-\text{CH}_2-\text{OR}_\text{B}$. In other embodiments, R_2 is $-\text{CH}_2-\text{SR}_\text{B}$. In yet other embodiments, R_2 is $-\text{CH}_2-\text{R}_\text{B}$. In other embodiments, R_2 is $-\text{CH}_2-\text{N}(\text{R}_\text{B})_2$. In still other embodiments, R_2 is $-\text{CH}_2-\text{NHR}_\text{B}$. In certain embodiments of the invention, R_B is one of:





wherein m and p are each independently integers from 0 to 3; q_1 is an integer from 1 to 6; R^{2C} is hydrogen, lower alkyl or a nitrogen protecting group; and each occurrence of R^{2B} is independently hydrogen, halogen, $-CN$, or WR^{W1} wherein W is O , S , NR^{W2} , $-C(=O)$, $-S(=O)$, $-SO_2$, $-C(=O)O-$, $-OC(=O)$, $-C(=O)NR^{W2}$, $-NR^{W2}C(=O)$; wherein each occurrence of R^{W1} and R^{W2} is independently hydrogen, a protecting group, a prodrug moiety or an alkyl, cycloalkyl, heteroalkyl, heterocyclic, aryl or heteroaryl moiety, or, when W is NR^{W2} , R^{W1} and R^{W2} , taken together with the nitrogen atom to which they are attached, form a heterocyclic or heteroaryl moiety; or any two adjacent occurrences of R^{2B} , taken together with the atoms to which they are attached, form a substituted or unsubstituted, saturated or unsaturated alicyclic or heterocyclic moiety, or a substituted or unsubstituted aryl or heteroaryl moiety. In certain embodiments of the invention, R_B is one of the structures:








XX

wherein m is an integer from 1 to 4; R^{2C} is hydrogen, lower alkyl or a nitrogen protecting group; and each occurrence of R^{2B} is independently hydrogen, halogen, $-CN$, or WR^{W1} wherein W is O , S , NR^{W2} , $-C(=O)$, $-S(=O)$, $-SO_2$, $-C(=O)O-$, $-OC(=O)$, $-C(=O)NR^{W2}$, $-NR^{W2}C(=O)$; wherein each occurrence of R^{W1} and R^{W2} is independently hydrogen, a protecting group, a prodrug moiety or an alkyl, cycloalkyl, heteroalkyl, heterocyclic, aryl or heteroaryl moiety, or, when W is NR^{W2} , R^{W1} and R^{W2} , taken together with the nitrogen atom to which they are attached, form a heterocyclic or heteroaryl moiety; or any two adjacent occurrences of R^{2B} , taken together with the atoms to which they are attached, form a substituted or unsubstituted, saturated or unsaturated alicyclic or heterocyclic moiety, or a substituted or unsubstituted aryl or heteroaryl moiety.

[00131] In certain embodiments, $-X(R_B)_n$ of $-CH_2-X(R_B)_n$ has one of the structures:









[00132] In certain embodiments, R_2 is

, wherein X is N




and Y is NH, S, or O. In other embodiments, R_2 is

[00133] In certain embodiments of the invention, the stereochemistry of formula is chosen from one of the following:



[00134] In certain embodiments of the invention, compounds are of the formula:




wherein

A, B, and R_2 are defined as above;

X is O or NH;

n is an integer between 1 and 20, inclusive; preferably, between 1 and 12, inclusive; more preferably between 2 and 8, inclusive. In certain embodiments, n is 2, 3, 4, 5, 6, 7, or 8; preferably, 6. In certain embodiments, X is NH. In other embodiments, X is O.

[00135] In certain embodiments of the invention, compounds are of the formula:




wherein

R_2 is defined as above;

n is an integer between 1 and 20, inclusive; preferably, between 1 and 12, inclusive; more preferably between 2 and 8, inclusive. In certain embodiments, n is 2, 3, 4, 5, 6, 7, or 8; preferably, 6.


[00136] In certain embodiments of the invention, compounds are of the formula:



wherein

R₂ is defined as above.


[00137] In certain embodiments of the invention, compounds are of the formula:



wherein

R₂ is defined as above.

[00138] In certain embodiments of the invention, compounds are of the formula:



[00139] Some of the foregoing compounds can comprise one or more asymmetric centers, and thus can exist in various isomeric forms, *e.g.*, stereoisomers and/or diastereomers. Thus, inventive compounds and pharmaceutical compositions thereof may be in the form of an individual enantiomer, diastereomer or geometric isomer, or may be in the form of a mixture of stereoisomers. In certain embodiments, the compounds of the invention are enantiopure compounds. In certain other embodiments, mixtures of stereoisomers or diastereomers are provided.

[00140] Furthermore, certain compounds, as described herein may have one or more double bonds that can exist as either the Z or E isomer, unless otherwise indicated. The invention additionally encompasses the compounds as individual isomers substantially free of other isomers and alternatively, as mixtures of various isomers, *e.g.*, racemic mixtures of stereoisomers. In addition to the above-mentioned compounds *per se*, this invention also encompasses pharmaceutically acceptable derivatives of these compounds and compositions comprising one or more compounds of the invention and one or more pharmaceutically acceptable excipients or additives.


[0100] Compounds of the invention may be prepared by crystallization of compound of any of the formula above under different conditions and may exist as one or a combination of polymorphs of compound of any general formula above forming part of this invention. For example, different polymorphs may be identified and/or prepared using different solvents, or different mixtures of solvents for recrystallization; by performing crystallizations at different

temperatures; or by using various modes of cooling, ranging from very fast to very slow cooling during crystallizations. Polymorphs may also be obtained by heating or melting the compound followed by gradual or fast cooling. The presence of polymorphs may be determined by solid probe NMR spectroscopy, IR spectroscopy, differential scanning calorimetry, powder X-ray diffractogram and/or other techniques. Thus, the present invention encompasses inventive compounds, their derivatives, their tautomeric forms, their stereoisomers, their polymorphs, their pharmaceutically acceptable salts their pharmaceutically acceptable solvates and pharmaceutically acceptable compositions containing them.


Synthetic Overview

[0101] As described above, the present invention provides novel compounds, specifically compounds having a 1,3-dioxane core as described above. The synthesis of compounds of the "tubacin" class has been described in previously filed U.S. patent publication no. 2006/0239909, published on October 26, 2006; U.S. patent publication no. 2003/0187027, published on October 2, 2003; and U.S. patent publication no. 2004/0072849, published on April 15, 2004. As would be appreciated by one of skill in this art, the various reactions and synthetic schemes described in these patent applications may be used in preparing the inventive compounds described herein.


[0102] A general synthetic scheme for preparing compounds of the isotubacin class is shown in *Figure 2*. A particular exemplary synthesis of isotubacin is shown in *Figure 3*. It

will be appreciated that for compounds of the formula , a method for the synthesis of the core structure is provided comprising steps of:


providing an epoxy alcohol having the structure:




reacting the epoxy alcohol with a reagent having the structure R_BXH under suitable conditions to generate a diol having the core structure:




reducing the nitro group to generate a diol having the core structure:



reacting the amino group with acylating agent to generate a diol having the core structure:



reacting the diol with a reagent having the structure $\text{R}_3\text{CH}(\text{OMe})_2$ or other acetal under suitable conditions to generate a scaffold having the core structure:




wherein R_1' is $-\text{L}-\text{A}$, wherein L is a linker moiety; and A comprises a functional group that inhibits histone deacetylase as described herein;

R_B is hydrogen, a protecting group, or an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, or heteroaromatic moiety;


X is $-\text{O}-$, $-\text{C}(\text{R}')_2-$, $-\text{S}-$, or $-\text{NR}'-$, wherein R' is hydrogen, a protecting group, or an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic or heteroaromatic moiety; and

R^3 is an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, or heteroaromatic moiety. In certain embodiments rather than reacting the diol with a reagent having the structure $\text{R}_3\text{CH}(\text{OMe})_2$ or other acetal, the diol is reacted with an aldehyde of structure R_3CHO . In other embodiments, the method optionally comprises additional steps of protecting and/or deprotecting functional groups of R_B , R_1 , R_1' , R_2 , or R_3 . In other embodiments, the method optionally comprises additional steps of modifying functional groups of R_B , R_1 , R_1' , R_2 , or R_3 . For example, in *Figure 2* the methyl ester functionality of R_1' is converted to a hydroxamic acid functional group.

[0103] In certain exemplary embodiments, the epoxy alcohol has the structure:




the diol has the structure:




wherein X is S or O;


and the core scaffold has the structure:



[0104] In certain other exemplary embodiments, the epoxy alcohol has the structure:




the diol has the structure:




wherein X is S or O;


and the core scaffold has the structure:



[0105] In certain other exemplary embodiments, the epoxy alcohol has the structure:




the diol has the structure:



wherein X is S or O;

and the core scaffold has the structure:



[0106] In certain exemplary embodiments, A is a hydroxamic acid. In certain embodiments, L is an C₁-C₈ alkylidene moiety. In certain embodiments, R₃ is a substituted or unsubstituted phenyl. In certain particular embodiments, R₃ is an unsubstituted phenyl. In certain embodiments, R₃ is a substituted phenyl. In certain embodiments, X is O. In other embodiments, X is S. In certain embodiments, X is NH or NR_B. In certain embodiments, R_B is substituted or unsubstituted aryl or heteroaryl.

[0107] A general synthetic scheme for preparing compounds of the “isotubacin” class is shown in *Figure 6*. A particular exemplary synthesis of isotubacin is shown in





Figure 7. It will be appreciated that for compounds of the formula R₃, a method for the synthesis of the core structure is provided comprising steps of:


providing an beta-hydroxy ketone having the structure:




reducing the ketone to generate a diol having the core structure:

optionally comprises additional steps of modifying functional groups of R_1 , R_1' , R_2 , or R_3 . For example, in *Figure 4* the protected hydroxamic acid functionality of R_1' is deprotected.


[00141] In certain embodiments, the step of providing a beta-hydroxy ketone having the structure:



comprises reacting an aldehyde of formula:




with a ketone of formula:




under suitable conditions (*e.g.*, basic conditions) to form the beta-hydroxy ketone.

[0108] In certain exemplary embodiments, the epoxy alcohol has the structure:




the diol has the structure:




wherein X is S or O;


and the core scaffold has the structure:



[0109] In certain other exemplary embodiments, the epoxy alcohol has the structure:




the diol has the structure:




wherein X is S or O;


and the core scaffold has the structure:



[0110] In certain other exemplary embodiments, the epoxy alcohol has the structure:




the diol has the structure:



wherein X is S or O;

and the core scaffold has the structure:



[00142] In certain exemplary embodiments, A is a hydroxamic acid. In certain embodiments, L is an C₁-C₈ alkylidene moiety. In certain embodiments, R₃ is a substituted or

unsubstituted phenyl. In certain particular embodiments, R_3 is an unsubstituted phenyl. In certain embodiments, R_3 is a substituted phenyl. In certain embodiments, X is O. In other embodiments, X is S. In certain embodiments, X is NH or NR_B . In certain embodiments, R_B is substituted or unsubstituted aryl or heteroaryl.

Pharmaceutical Compositions

[00143] As discussed above, the present invention provides novel compounds having antitumor, antineurodegenerative, antibiotic, and antiproliferative activity, and thus the inventive compounds are useful for the treatment of cancer, benign neoplasms, neurodegenerative disorders, protein degradation disorders, and protein deposition disorders.

[00144] Accordingly, in another aspect of the present invention, pharmaceutical compositions are provided, which comprise any one of the compounds described herein (or a prodrug, pharmaceutically acceptable salt or other pharmaceutically acceptable derivative thereof), and optionally comprise a pharmaceutically acceptable carrier. In certain embodiments, these compositions optionally further comprise one or more additional therapeutic agents. Alternatively, a compound of this invention may be administered to a patient in need thereof in combination with the administration of one or more other therapeutic agents. For example, additional therapeutic agents for conjoint administration or inclusion in a pharmaceutical composition with a compound of this invention may be an approved chemotherapeutic agent, or it may be any one of a number of agents undergoing approval in the Food and Drug Administration that ultimately obtain approval for the treatment of protozoal infections and/or any disorder associated with cellular hyperproliferation. In certain other embodiments, the additional therapeutic agent is an anticancer agent, as discussed in more detail herein. In certain other embodiments, the compositions of the invention are useful for the treatment of protozoal infections. In the treatment of cancer or protein degradation disorders, the inventive compound may be combined with a proteasome inhibitor (*e.g.*, bortezomib, R115777 FTI, MG132, NPI-0052, *etc.*). In the treatment of cancer or protein degradation disorders, the inventive compound may be combined with protein degradation inhibitor (*e.g.*, another inventive compound, a tubacin-like compound, bortezomib, R115777 FTI, MG132, NPI-0052, SAHA, ^{166}Ho -DOTMP, arsenic trioxide, 17-AAG, MG132, saopjargon, *etc.*).

[00145] As discussed above, the compounds of the present invention are useful as anticancer agents, and thus may be useful in the treatment of cancer, by effecting tumor cell death or inhibiting the growth of tumor cells. In general, the inventive anticancer agents are

useful in the treatment of cancers, including, but not limited to breast cancer, brain cancer, skin cancer, cervical cancer, colon and rectal cancer, leukemia, lung cancer, melanoma, multiple myeloma, non-Hodgkin's lymphoma, ovarian cancer, pancreatic cancer, prostate cancer, and gastric cancer, to name a few. In certain embodiments, the inventive anticancer agents are active against leukemia cells, and thus are useful for the treatment of leukemias (e.g., myeloid, lymphocytic, promyelocytic, myelocytic and lymphoblastic leukemias, whether acute or chronic forms). In still other embodiments, the inventive anticancer agents are active against solid tumors and also kill and/or inhibit the growth of multidrug resistant cells (MDR cells). In certain embodiments, the inventive anticancer agents are active against cancers which are resistant to other known anti-neoplastic agents or which have been found not to respond clinically to other known anti-neoplastic agents.

[00146] It will also be appreciated that the compounds and pharmaceutical compositions of the present invention can be employed in combination therapies, that is, the compounds and pharmaceutical compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed may achieve a desired effect for the same disorder (for example, an inventive compound may be administered concurrently with another anticancer agent), or they may achieve different effects (e.g., control of any adverse effects).

[00147] It will also be appreciated that certain of the compounds of present invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable derivative thereof. According to the present invention, a pharmaceutically acceptable derivative includes, but is not limited to, pharmaceutically acceptable salts, esters, salts of such esters, or a pro-drug or other adduct or derivative of a compound of this invention which upon administration to a patient in need is capable of providing, directly or indirectly, a compound as otherwise described herein, or a metabolite or residue thereof.

[00148] As used herein, the term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts of amines, carboxylic acids, and other types of compounds, are well known in the art. For example, S.M. Berge, *et al.* describe pharmaceutically acceptable salts in

detail in J. Pharmaceutical Sciences, 66: 1-19 (1977). The salts can be prepared *in situ* during the final isolation and purification of the compounds of the invention, or separately by reacting a free base or free acid function with a suitable reagent, as described generally below. For example, a free base function can be reacted with a suitable acid. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may, include metal salts such as alkali metal salts, *e.g.* sodium or potassium salts; and alkaline earth metal salts, *e.g.* calcium or magnesium salts. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, harnisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.


[00149] Additionally, as used herein, the term "pharmaceutically acceptable ester" refers to esters that hydrolyze *in vivo* and include those that break down readily in the human body to leave the parent compound or a salt thereof. Suitable ester groups include, for example,

those derived from pharmaceutically acceptable aliphatic carboxylic acids, particularly alkanoic, alkenoic, cycloalkanoic and alkanedioic acids, in which each alkyl or alkenyl moiety advantageously has not more than 6 carbon atoms. Examples of particular esters include formates, acetates, propionates, butyrates, acrylates and ethylsuccinates.

[00150] Furthermore, the term “pharmaceutically acceptable prodrugs” as used herein refers to those prodrugs of the compounds of the present invention which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals with undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term “prodrug” refers to compounds that are rapidly transformed *in vivo* to yield the parent compound of the above formula, for example by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987.

[00151] As described above, the pharmaceutical compositions of the present invention additionally comprise a pharmaceutically acceptable carrier, which, as used herein, includes any and all solvents, diluents, or other liquid vehicle, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington's Pharmaceutical Sciences, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1980) discloses various carriers used in formulating pharmaceutical compositions and known techniques for the preparation thereof. Except insofar as any conventional carrier medium is incompatible with the compounds of the invention, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutical composition, its use is contemplated to be within the scope of this invention. Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt;

gelatine; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil, sesame oil; olive oil; corn oil and soybean oil; glycols; such as propylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogenfree water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.



[00152] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[00153] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[00154] The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[00155] In order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension or crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release

can be controlled. Examples of other biodegradable polymers include (poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

[00156] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

[00157] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar--agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.


[00158] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[00159] The active compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and

granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose and starch. Such dosage forms may also comprise, as in normal practice, additional substances other than inert diluents, *e.g.*, tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

[00160] The present invention encompasses pharmaceutically acceptable topical formulations of inventive compounds. The term "pharmaceutically acceptable topical formulation," as used herein, means any formulation which is pharmaceutically acceptable for intradermal administration of a compound of the invention by application of the formulation to the epidermis. In certain embodiments of the invention, the topical formulation comprises a carrier system. Pharmaceutically effective carriers include, but are not limited to, solvents (*e.g.*, alcohols, poly alcohols, water), creams, lotions, ointments, oils, plasters, liposomes, powders, emulsions, microemulsions, and buffered solutions (*e.g.*, hypotonic or buffered saline) or any other carrier known in the art for topically administering pharmaceuticals. A more complete listing of art-known carriers is provided by reference texts that are standard in the art, for example, Remington's Pharmaceutical Sciences, 16th Edition, 1980 and 17th Edition, 1985, both published by Mack Publishing Company, Easton, Pa. In certain other embodiments, the topical formulations of the invention may comprise excipients. Any pharmaceutically acceptable excipient known in the art may be used to prepare the inventive pharmaceutically acceptable topical formulations. Examples of excipients that can be included in the topical formulations of the invention include, but are not limited to, preservatives, antioxidants, moisturizers, emollients, buffering agents, solubilizing agents, other penetration agents, skin protectants, surfactants, and propellants, and/or additional therapeutic agents used in combination to the inventive compound. Suitable preservatives include, but are not limited to, alcohols, quaternary amines, organic acids,

parabens, and phenols. Suitable antioxidants include, but are not limited to, ascorbic acid and its esters, sodium bisulfite, butylated hydroxytoluene, butylated hydroxyanisole, tocopherols and



chelating agents like EDTA and citric acid. Suitable moisturizers include, but are not limited to, glycerine, sorbitol, polyethylene glycols, urea, and propylene glycol. Suitable buffering agents for use with the invention include, but are not limited to, citric, hydrochloric, and lactic acid buffers. Suitable solubilizing agents include, but are not limited to, quaternary ammonium chlorides, cyclodextrins, benzyl benzoate, lecithin, and polysorbates. Suitable skin protectants that can be used in the topical formulations of the invention include, but are not limited to, vitamin E oil, allantoin, dimethicone, glycerin, petrolatum, and zinc oxide.

[00161] In certain embodiments, the pharmaceutically acceptable topical formulations of the invention comprise at least a compound of the invention and a penetration enhancing agent. The choice of topical formulation will depend on several factors, including the condition to be treated, the physicochemical characteristics of the inventive compound and other excipients present, their stability in the formulation, available manufacturing equipment, and costs constraints. As used herein the term "penetration enhancing agent" means an agent capable of transporting a pharmacologically active compound through the stratum corneum and into the epidermis or dermis, preferably, with little or no systemic absorption. A wide variety of compounds have been evaluated as to their effectiveness in enhancing the rate of penetration of drugs through the skin. See, for example, *Percutaneous Penetration Enhancers*, Maibach H. I. and Smith H. E. (eds.), CRC Press, Inc., Boca Raton, Fla. (1995), which surveys the use and testing of various skin penetration enhancers, and Buyuktimkin *et al.*, *Chemical Means of Transdermal Drug Permeation Enhancement in Transdermal and Topical Drug Delivery Systems*, Gosh T. K., Pfister W. R., Yum S. I. (Eds.), Interpharm Press Inc., Buffalo Grove, Ill. (1997). In certain exemplary embodiments, penetration agents for use with the invention include, but are not limited to, triglycerides (*e.g.*, soybean oil), aloe compositions (*e.g.*, aloe-vera gel), ethyl alcohol, isopropyl alcohol, octylphenylpolyethylene glycol, oleic acid, polyethylene glycol 400, propylene glycol, N-decylmethylsulfoxide, fatty acid esters (*e.g.*, isopropyl myristate, methyl laurate, glycerol monooleate, and propylene glycol monooleate) and N-methyl pyrrolidone.

[00162] In certain embodiments, the compositions may be in the form of ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. In certain exemplary embodiments, formulations of the compositions according to the invention are creams, which may further contain saturated or unsaturated fatty acids such as stearic acid, palmitic acid, oleic acid, palmito-oleic acid, cetyl or oleyl alcohols, stearic acid being particularly preferred. Creams of the invention may also contain a non-ionic surfactant, for example, polyoxy-40-stearate. In certain embodiments, the active component is admixed under sterile conditions

with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, eardrops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms are made by dissolving or dispensing the compound in the proper medium. As discussed above, penetration enhancing agents can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[00163] It will also be appreciated that the compounds and pharmaceutical compositions of the present invention can be formulated and employed in combination therapies, that is, the compounds and pharmaceutical compositions can be formulated with or administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed may achieve a desired effect for the same disorder (for example, an inventive compound may be administered concurrently with another immunomodulatory agent, anticancer agent or agent useful for the treatment of psoriasis), or they may achieve different effects (*e.g.*, control of any adverse effects).

[00164] For example, other therapies or anticancer agents that may be used in combination with the inventive compounds of the present invention include surgery, radiotherapy (in but a few examples, γ -radiation, neutron beam radiotherapy, electron beam radiotherapy, proton therapy, brachytherapy, and systemic radioactive isotopes, to name a few), endocrine therapy, biologic response modifiers (interferons, interleukins, and tumor necrosis factor (TNF) to name a few), hyperthermia and cryotherapy, agents to attenuate any adverse effects (*e.g.*, antiemetics), and other approved chemotherapeutic drugs, including, but not limited to, alkylating drugs (*e.g.*, mechlorethamine, chlorambucil, Cyclophosphamide, Melphalan, Ifosfamide), antimetabolites (*e.g.*, Methotrexate), purine antagonists and pyrimidine antagonists (*e.g.*, 6-Mercaptopurine, 5-Fluorouracil, Cytarabine, Gemcitabine), spindle poisons (*e.g.*, Vinblastine, Vincristine, Vinorelbine, Paclitaxel), podophyllotoxins (*e.g.*, Etoposide, Irinotecan, Topotecan), antibiotics (Doxorubicin, Bleomycin, Mitomycin), nitrosoureas (*e.g.*, Carmustine, Lomustine), inorganic ions (*e.g.*, Cisplatin, Carboplatin),

enzymes (e.g., Asparaginase), and hormones (e.g., Tamoxifen, Leuprolide, Flutamide, and Megestrol), to name a few. For a more comprehensive discussion of updated cancer therapies see, *The Merck Manual*, Seventeenth Ed. 1999. See also the National Cancer Institute (NCI) website (www.nci.nih.gov) and the Food and Drug Administration (FDA) website for a list of the FDA approved oncology drugs (www.fda.gov/cder/cancer/druglistframe).

[00165] In certain embodiments, the pharmaceutical compositions of the present invention further comprise one or more additional therapeutically active ingredients (e.g., . . . chemotherapeutic and/or palliative). For purposes of the invention, the term "*palliative*" refers to treatment that is focused on the relief of symptoms of a disease and/or side effects of a therapeutic regimen, but is not curative. For example, palliative treatment encompasses painkillers, anti-nausea medications, anti-pyretics, and anti-sickness drugs. In addition, chemotherapy, radiotherapy and surgery can all be used palliatively (that is, to reduce symptoms without going for cure; e.g., for shrinking tumors and reducing pressure, bleeding, pain and other symptoms of cancer).

[00166] Additionally, the present invention provides pharmaceutically acceptable derivatives of the inventive compounds, and methods of treating a subject using these compounds, pharmaceutical compositions thereof, or either of these in combination with one or more additional therapeutic agents.

[00167] It will also be appreciated that certain of the compounds of present invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable derivative thereof. According to the present invention, a pharmaceutically acceptable derivative includes, but is not limited to, pharmaceutically acceptable salts, esters, salts of such esters, or a prodrug or other adduct or derivative of a compound of this invention which upon administration to a patient in need is capable of providing, directly or indirectly, a compound as otherwise described herein, or a metabolite or residue thereof.

Research Uses, Pharmaceutical Uses and Methods of Treatment

Research Uses

[00168] According to the present invention, the inventive compounds may be assayed in any of the available assays known in the art for identifying compounds having antiprotozoal,

HDAC inhibitory, TDAC inhibitory, aggresome inhibitory, and/or antiproliferative activity. For example, the assay may be cellular or non-cellular, *in vivo* or *in vitro*, high- or low-throughput format, *etc.*

[00169] Thus, in one aspect, compounds of this invention which are of particular interest include those which:

- exhibit HDAC-inhibitory activity;
- exhibit TDAC-inhibitory activity;
- exhibit HDAC Class I inhibitory activity (*e.g.*, HDAC1, HDAC2, HDAC3, HDAC8);
- exhibit HDAC Class II inhibitory activity (*e.g.*, HDAC4, HDAC5, HDAC6, HDAC7, HDAC9a, HDAC9b, HDRP/HDAC9c, HDAC10);
- exhibit HDAC Class IV inhibitory activity;
- exhibit the ability to inhibit HDAC1 (Genbank Accession No. NP_004955);
- exhibit the ability to inhibit HDAC2 (Genbank Accession No. NP_001518);
- exhibit the ability to inhibit HDAC3 (Genbank Accession No. O15739);
- exhibit the ability to inhibit HDAC4 (Genbank Accession No. AAD29046);
- exhibit the ability to inhibit HDAC5 (Genbank Accession No. NP_005465);
- exhibit the ability to inhibit HDAC6 (Genbank Accession No. NP_006035);
- exhibit the ability to inhibit HDAC7 (Genbank Accession No. AAP63491);
- exhibit the ability to inhibit HDAC8 (Genbank Accession No. AAF73428, NM_018486, AF245664, AF230097);
- exhibit the ability to inhibit HDAC9 (Genbank Accession No. NM_178425, NM_178423, NM_058176, NM_014707, BC111735, NM_058177);
- exhibit the ability to inhibit HDAC10 (Genbank Accession No. NM_032019);
- exhibit the ability to inhibit HDAC11 (Genbank Accession No. BC009676);

- exhibit the ability to modulate the glucose-sensitive subset of genes downstream of Ure2p;
- exhibit the ability to inhibit the degradation of protein by the aggresome;
- exhibit cytotoxic or growth inhibitory effect on cancer cell lines maintained *in vitro* or in animal studies using a scientifically acceptable cancer cell xenograft model; and/or
- exhibit a therapeutic profile (*e.g.*, optimum safety and curative effect) that is superior to existing chemotherapeutic agents.

[00170] As detailed in the exemplification herein, in assays to determine the ability of compounds to inhibit cancer cell growth certain inventive compounds may exhibit IC_{50} values $\leq 100 \mu M$. In certain other embodiments, inventive compounds exhibit IC_{50} values $\leq 50 \mu M$. In certain other embodiments, inventive compounds exhibit IC_{50} values $\leq 40 \mu M$. In certain other embodiments, inventive compounds exhibit IC_{50} values $\leq 30 \mu M$. In certain other embodiments, inventive compounds exhibit IC_{50} values $\leq 20 \mu M$. In certain other embodiments, inventive compounds exhibit IC_{50} values $\leq 10 \mu M$. In certain other embodiments, inventive compounds exhibit IC_{50} values $\leq 7.5 \mu M$. In certain embodiments, inventive compounds exhibit IC_{50} values $\leq 5 \mu M$. In certain other embodiments, inventive compounds exhibit IC_{50} values $\leq 2.5 \mu M$. In certain embodiments, inventive compounds exhibit IC_{50} values $\leq 1 \mu M$. In certain embodiments, inventive compounds exhibit IC_{50} values $\leq 0.75 \mu M$. In certain embodiments, inventive compounds exhibit IC_{50} values $\leq 0.5 \mu M$. In certain embodiments, inventive compounds exhibit IC_{50} values $\leq 0.25 \mu M$. In certain embodiments, inventive compounds exhibit IC_{50} values $\leq 0.1 \mu M$. In certain other embodiments, inventive compounds exhibit IC_{50} values $\leq 75 \text{ nM}$. In certain other embodiments, inventive compounds exhibit IC_{50} values $\leq 50 \text{ nM}$. In certain other embodiments, inventive compounds exhibit IC_{50} values $\leq 25 \text{ nM}$. In certain other embodiments, inventive compounds exhibit IC_{50} values $\leq 10 \text{ nM}$. In other embodiments, exemplary compounds exhibited IC_{50} values $\leq 7.5 \text{ nM}$. In other embodiments, exemplary compounds exhibited IC_{50} values $\leq 5 \text{ nM}$.

Pharmaceutical Uses and Methods of Treatment

[00171] In general, methods of using the compounds of the present invention comprise administering to a subject in need thereof a therapeutically effective amount of a compound of the present invention. As discussed above, the compounds of the invention are selective inhibitors of histone deacetylases and, as such, are useful in the treatment of disorders

modulated by histone deacetylases. As discussed above, the compounds of the invention are selective inhibitors of tubulin deacetylases and, as such, are useful in the treatment of disorders modulated by tubulin deacetylases. For example, compounds of the invention may be useful in the treatment of cancer (*e.g.*, breast cancer, prostate cancer, multiple myeloma, leukemia, lymphoma, *etc.*). Accordingly, in yet another aspect, according to the methods of treatment of the present invention, tumor cells are killed, or their growth is inhibited by contacting said tumor cells with an inventive compound or composition, as described herein.

[00172] Thus, in another aspect of the invention, methods for the treatment of cancer are provided comprising administering a therapeutically effective amount of an inventive compound, as described herein, to a subject in need thereof. In certain embodiments, a method for the treatment of cancer is provided comprising administering a therapeutically effective amount of an inventive compound, or a pharmaceutical composition comprising an inventive compound to a subject in need thereof, in such amounts and for such time as is necessary to achieve the desired result. In certain embodiments of the present invention a “therapeutically effective amount” of the inventive compound or pharmaceutical composition is that amount effective for killing or inhibiting the growth of tumor cells. The compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for killing or inhibiting the growth of tumor cells. Thus, the expression “amount effective to kill or inhibit the growth of tumor cells,” as used herein, refers to a sufficient amount of agent to kill or inhibit the growth of tumor cells. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular anticancer agent, its mode of administration, and the like.

[00173] In certain embodiments, the method involves the administration of a therapeutically effective amount of the compound or a pharmaceutically acceptable derivative thereof to a subject (including, but not limited to a human or animal) in need of it. In certain embodiments, the inventive compounds as useful for the treatment of cancer (including, but not limited to, glioblastoma, retinoblastoma, breast cancer, cervical cancer, colon and rectal cancer, leukemia (*e.g.*, CML, AML, CLL, ALL), lymphoma, lung cancer (including, but not limited to small cell lung cancer), melanoma and/or skin cancer, multiple myeloma, non-Hodgkin’s lymphoma, ovarian cancer, pancreatic cancer, prostate cancer and gastric cancer, bladder cancer, uterine cancer, kidney cancer, testicular cancer, stomach cancer, brain cancer, liver cancer, or esophageal cancer).

[00174] In certain embodiments, the inventive anticancer agents are useful in the treatment of cancers and other proliferative disorders, including, but not limited to breast cancer, cervical cancer, colon and rectal cancer, leukemia, lung cancer, melanoma, multiple myeloma, non-Hodgkin's lymphoma, ovarian cancer, pancreatic cancer, prostate cancer, and gastric cancer, to name a few. In certain embodiments, the inventive anticancer agents are active against leukemia cells and melanoma cells, and thus are useful for the treatment of leukemias (e.g., myeloid, lymphocytic, myelocytic and lymphoblastic leukemias) and malignant melanomas. In still other embodiments, the inventive anticancer agents are active against solid tumors.

[00175] In certain embodiments, the inventive compounds also find use in the prevention of restenosis of blood vessels subject to traumas such as angioplasty and stenting. For example, it is contemplated that the compounds of the invention will be useful as a coating for implanted medical devices, such as tubings, shunts, catheters, artificial implants, pins, electrical implants such as pacemakers, and especially for arterial or venous stents, including balloon-expandable stents. In certain embodiments inventive compounds may be bound to an implantable medical device, or alternatively, may be passively adsorbed to the surface of the implantable device. In certain other embodiments, the inventive compounds may be formulated to be contained within, or, adapted to release by a surgical or medical device or implant, such as, for example, stents, sutures, indwelling catheters, prosthesis, and the like. For example, drugs having antiproliferative and anti-inflammatory activities have been evaluated as stent coatings, and have shown promise in preventing restenosis (See, for example, Presbitero P. *et al.*, "Drug eluting stents do they make the difference?", *Minerva Cardioangiol*, 2002, **50**(5):431-442; Ruygrok P.N. *et al.*, "Rapamycin in cardiovascular medicine", *Intern. Med. J.*, 2003, **33**(3):103-109; and Marx S.O. *et al.*, "Bench to bedside: the development of rapamycin and its application to stent restenosis", *Circulation*, 2001, **104**(8):852-855. Accordingly, without wishing to be bound to any particular theory, Applicant proposes that inventive compounds having antiproliferative effects can be used as stent coatings and/or in stent drug delivery devices, inter alia for the prevention of restenosis or reduction of restenosis rate. Suitable coatings and the general preparation of coated implantable devices are described in U.S. Patents 6,099,562; 5,886,026; and 5,304,121. The coatings are typically biocompatible polymeric materials such as a hydrogel polymer, polymethyldisiloxane,

polycaprolactone, polyethylene glycol, polylactic acid, ethylene vinyl acetate, and mixtures thereof. The coatings may optionally be further covered by a suitable topcoat of fluorosilicone, polysaccharides, polyethylene glycol, phospholipids or combinations thereof to impart controlled release characteristics in the composition. A variety of compositions and methods related to stent coating and/or local stent drug delivery for preventing restenosis are known in the art (see, for example, U.S. Patent Nos.: 6,517,889; 6,273,913; 6,258,121; 6,251,136; 6,248,127; 6,231,600; 6,203,551; 6,153,252; 6,071,305; 5,891,507; 5,837,313 and published U.S. patent application No.: US2001/0027340). For example, stents may be coated with polymer-drug conjugates by dipping the stent in polymer-drug solution or spraying the stent with such a solution. In certain embodiment, suitable materials for the implantable device include biocompatible and nontoxic materials, and may be chosen from the metals such as nickel-titanium alloys, steel, or biocompatible polymers, hydrogels, polyurethanes, polyethylenes, ethylenevinyl acetate copolymers, etc. In certain embodiments, the inventive compound is coated onto a stent for insertion into an artery or vein following balloon angioplasty.

[00176] The compounds of this invention or pharmaceutically acceptable compositions thereof may also be incorporated into compositions for coating implantable medical devices, such as prostheses, artificial valves, vascular grafts, stents and catheters. Accordingly, the present invention, in another aspect, includes a composition for coating an implantable device comprising a compound of the present invention as described generally above, and in classes and subclasses herein, and a carrier suitable for coating said implantable device. In still another aspect, the present invention includes an implantable device coated with a composition comprising a compound of the present invention as described generally above, and in classes and subclasses herein, and a carrier suitable for coating said implantable device.

[00177] Within other aspects of the present invention, methods are provided for expanding the lumen of a body passageway, comprising inserting a stent into the passageway, the stent having a generally tubular structure, the surface of the structure being coated with (or otherwise adapted to release) an inventive compound or composition, such that the

passageway is expanded. In certain embodiments, the lumen of a body passageway is expanded in order to eliminate a biliary, gastrointestinal, esophageal, tracheal/bronchial, urethral and/or vascular obstruction.

[00178] Methods for eliminating biliary, gastrointestinal, esophageal, tracheal/bronchial, urethral and/or vascular obstructions using stents are known in the art. The skilled practitioner will know how to adapt these methods in practicing the present invention. For example, guidance can be found in U.S. Patent Publication No.: 2003/0004209 in paragraphs [0146]-[0155].

[00179] Another aspect of the invention relates to a method for inhibiting the growth of multidrug resistant cells in a biological sample or a patient, which method comprises administering to the patient, or contacting said biological sample with a compound of the invention or a composition comprising said compound.

[00180] Additionally, the present invention provides pharmaceutically acceptable derivatives of the inventive compounds, and methods of treating a subject using these compounds, pharmaceutical compositions thereof, or either of these in combination with one or more additional therapeutic agents.

[00181] Another aspect of the invention relates to a method of treating or lessening the severity of a disease or condition associated with a proliferation disorder in a patient, said method comprising a step of administering to said patient, a compound of formula I or a composition comprising said compound.

[00182] It will be appreciated that the compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for the treatment of cancer and/or disorders associated with cell hyperproliferation. For example, when using the inventive compounds for the treatment of cancer, the expression "effective amount" as used herein, refers to a sufficient amount of agent to inhibit cell proliferation, or refers to a sufficient amount to reduce the effects of cancer. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the diseases, the particular anticancer agent, its mode of administration, and the like.

[00183] The compounds of the invention are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression "dosage unit form" as used herein refers to a physically discrete unit of therapeutic agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts (see, for example, Goodman and Gilman's, "The Pharmacological Basis of Therapeutics", Tenth Edition, A. Gilman, J. Hardman and L. Limbird, eds., McGraw-Hill Press, 155-173, 2001).

[00184] Another aspect of the invention relates to a method for inhibiting histone deacetylase activity in a biological sample or a patient, which method comprises administering to the patient, or contacting said biological sample with a compound of formula I or a composition comprising said compound.

[00185] Furthermore, after formulation with an appropriate pharmaceutically acceptable carrier in a desired dosage, the pharmaceutical compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, creams or drops), buccally, as an oral or nasal spray, or the like, depending on the severity of the infection being treated. In certain embodiments, the compounds of the invention may be administered at dosage levels of about 0.001 mg/kg to about 50 mg/kg, from about 0.01 mg/kg to about 25 mg/kg, or from about 0.1 mg/kg to about 10 mg/kg of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect. It will also be appreciated that dosages smaller than 0.001 mg/kg or greater than 50 mg/kg (for example 50-100 mg/kg) can be administered to a subject. In certain embodiments, compounds are administered orally or parenterally.

Treatment Kit

[00186] In other embodiments, the present invention relates to a kit for conveniently and effectively carrying out the methods in accordance with the present invention. In general, the pharmaceutical pack or kit comprises one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Such kits are especially suited for the delivery of solid oral forms such as tablets or capsules. Such a kit preferably includes a number of unit dosages, and may also include a card having the dosages oriented in the order of their intended use. If desired, a memory aid can be provided, for example in the form of numbers, letters, or other markings or with a calendar insert, designating the days in the treatment schedule in which the dosages can be administered. Alternatively, placebo dosages, or calcium dietary supplements, either in a form similar to or distinct from the dosages of the pharmaceutical compositions, can be included to provide a kit in which a dosage is taken every day. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceutical products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

Equivalents

[00187] The representative examples which follow are intended to help illustrate the invention, and are not intended to, nor should they be construed to, limit the scope of the invention. Indeed, various modifications of the invention and many further embodiments thereof, in addition to those shown and described herein, will become apparent to those skilled in the art from the full contents of this document, including the examples which follow and the references to the scientific and patent literature cited herein. The following examples contain important additional information, exemplification and guidance which can be adapted to the practice of this invention in its various embodiments and the equivalents thereof.


These and other aspects of the present invention will be further appreciated upon consideration of the following Examples, which are intended to illustrate certain particular embodiments of the invention but are not intended to limit its scope, as defined by the claims.

Examples

[00188] The compounds of this invention and their preparation can be understood further by the examples that illustrate some of the processes by which these compounds are prepared or used. It will be appreciated, however, that these examples do not limit the invention. Variations of the invention, now known or further developed, are considered to fall within the scope of the present invention as described herein and as hereinafter claimed.

Example 1: Synthetic Methods

[00189] The various references cited herein provide helpful background information on preparing compounds similar to the inventive compounds described herein or relevant intermediates, as well as information on formulation, uses, and administration of such compounds which may be of interest.



[00190] Moreover, the practitioner is directed to the specific guidance and examples provided in this document relating to various exemplary compounds and intermediates thereof.

[00191] The compounds of this invention and their preparation can be understood further by the examples that illustrate some of the processes by which these compounds are prepared or used. It will be appreciated, however, that these examples do not limit the invention. Variations of the invention, now known or further developed, are considered to fall within the scope of the present invention as described herein and as hereinafter claimed.

[00192] According to the present invention, any available techniques can be used to make or prepare the inventive compounds or compositions including them. For example, a variety of a variety combinatorial techniques, parallel synthesis and/or solid phase synthetic methods such as those discussed in detail below may be used. Alternatively or additionally, the inventive compounds may be prepared using any of a variety of solution phase synthetic methods known in the art.

[00193] It will be appreciated as described below, that a variety of inventive compounds can be synthesized according to the methods described herein. The starting materials and reagents used in preparing these compounds are either available from commercial suppliers such as Aldrich Chemical Company (Milwaukee, WI), Bachem (Torrance, CA), Sigma (St. Louis, MO), or are prepared by methods well known to a person of ordinary skill in the art following procedures described in such references as Fieser and Fieser 1991, "Reagents for Organic Synthesis", vols 1-17, John Wiley and Sons, New York, NY, 1991; Rodd 1989 "Chemistry of Carbon Compounds", vols. 1-5 and supps, Elsevier Science Publishers, 1989; "Organic Reactions", vols 1-40, John Wiley and Sons, New York, NY, 1991; March 2001, "Advanced Organic Chemistry", 5th ed. John Wiley and Sons, New York, NY; and Larock 1990, "Comprehensive Organic Transformations: A Guide to Functional Group Preparations", 2nd ed. VCH Publishers. These schemes are merely illustrative of some methods by which the compounds of this invention can be synthesized, and various modifications to these schemes can be made and will be suggested to a person of ordinary skill in the art having regard to this disclosure.

[00194] The starting materials, intermediates, and compounds of this invention may be isolated and purified using conventional techniques, including filtration, distillation, crystallization, chromatography, and the like. They may be characterized using conventional methods, including physical constants and spectral data.

Example 2: Biological Assay Procedures

[00195] **Cell culture and Transfections.** TAg-Jurkat cells were transfected by electroporation with 5 µg of FLAG-epitope-tagged pBJ5 constructs for expression of recombinant proteins. Cells were harvested 48 h posttransfection.

[00196] **HDAC assays.** [³H]Acetate-incorporated histones were isolated from butyrate-treated HeLa cells by hydroxyapatite chromatography (as described in Tong, *et al. Nature* 1997, 395, 917-921.) Immunoprecipitates were incubated with 1.4 µg (10,000 dpm) histones for 3 h at 37° C. HDAC activity was determined by scintillation counting of the ethyl acetate-soluble [³H]acetic acid (as described in Taunton, *et al., Science* 1996, 272, 408-411). Compounds were added in DMSO such that final assay concentrations were 1% DMSO. IC₅₀s were calculated using Prism 3.0 software. Curve fitting was done without constraints using the program's Sigmoidal-Dose Response parameters. All data points were acquired in duplicate and IC₅₀s are calculated from the composite results of at least two separate experiments.

Example 3: *In vivo* activity

[00197] Although a variety of methods can be utilized, one exemplary method by which the *in vivo* activity of the inventive compounds is determined is by subcutaneously transplanting a desired tumor mass in mice. Drug treatment is then initiated when tumor mass reaches approximately 100 mm³ after transplantation of the tumor mass. A suitable composition, as described in more detail above, is then administered to the mice, preferably in saline and also preferably administered once a day at doses of 5, 10 and 25 mg/kg, although it will be appreciated that other doses can also be administered. Body weight and tumor size are then measured daily and changes in percent ratio to initial values are plotted. In cases where the transplanted tumor ulcerates, the weight loss exceeds 25-30% of control weight loss, the tumor weight reaches 10% of the body weight of the cancer-bearing mouse, or the cancer-bearing mouse is dying, the animal is sacrificed in accordance with guidelines for animal welfare.

Example 4: Assays to identify potential antiprotozoal compounds by inhibition of histone deacetylase

[00198] As detailed in U.S. Patent 6,068,987, inhibitors of histone deacetylases may also be useful as antiprotozoal agents. Described therein are

assays for histone deacetylase activity and inhibition and describe a variety of known protozoal diseases.

Example 5: High-throughput immunofluorescence Assay


[00199] Compounds of the invention are tested from their HDAC or TDAC specificity using a high-throughput immunofluorescence-based assay. The assay is based on the use of specific antibodies for acetylated tubulin and acetylated lysine (*i.e.*, a marker for acetylated histones).

[00200] Cells (*e.g.*, 293T cells) are incubated the inventive compound over a test range of concentrations after the cells have been allowed to adhere to the cell culture plate overnight. After a determined time of incubation with the inventive compound (*e.g.*, 6-8 hours), the cells are treated with a first primary antibody directed against acetylated tubulin and a second primary antibody directed against acetylated lysine. The cells are then contacted with two secondary antibodies specific for each of the primary antibodies and identifiable by a unique fluorescent signal.

[00201] The plates are then imaged, and the fluorescence signal from each of the secondary antibodies is quantitated. The data gathered is then used to calculate dose-response curves, to calculate IC_{50} values, to establish structure-function relationships, to calculate the ratio of HDAC to TDAC inhibition, and to determine the specificity for HDAC or TDAC.



WHAT IS CLAIMED IS:

1. A compound of the formula:



wherein


R₁ is of the formula:



Y is NH or O;

L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety; a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety; a substituted or unsubstituted aryl moiety; a substituted or unsubstituted heteroaryl moiety; or a cyclic ring system, wherein the rings may be aryl, heteroaryl, non-aromatic carbocyclic, or non-aromatic heterocyclic; and

A comprises a functional group selected from the group consisting of:




R_2 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-\text{OR}_B$; $-\text{C(=O)R}_B$; $-\text{CO}_2\text{R}_B$; $-\text{CN}$; $-\text{SCN}$; $-\text{SR}_B$; $-\text{SOR}_B$; $-\text{SO}_2\text{R}_B$; $-\text{NO}_2$; $-\text{N(R}_B)_2$; $-\text{NHC(O)R}_B$; or $-\text{C(R}_B)_3$; wherein each occurrence of R_B is independently a hydrogen, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety; and


R_3 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-\text{OR}_C$; $-\text{C(=O)R}_C$; $-\text{CO}_2\text{R}_C$; $-\text{CN}$; $-\text{SCN}$; $-\text{SR}_C$; $-\text{SOR}_C$; $-\text{SO}_2\text{R}_C$; $-\text{NO}_2$; $-\text{N(R}_C)_2$; $-\text{NHC(O)R}_C$; or $-\text{C(R}_C)_3$; wherein each occurrence of R_C is independently a hydrogen, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl

moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety; or pharmaceutically acceptable salts thereof.


2. The compound of claim 1, wherein the compound is of one of the formulae:



3. The compound of claim 1 or 2, wherein R_1 is of the formula:



4. The compound of any one of claims 1 to 3, wherein R_1' is of the formula:




wherein n is an integer between 0 and 15, inclusive.

5. The compound of claim 4, wherein n is 5 or 6.


6. The compound of any one of claims 1 to 5, wherein R_2 is of the formula:

R₂' is aliphatic, heteroaliphatic, acyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl.


- 107



13. The compound of any one of claims 1 to 5, wherein R_2 is selected from one of the following:




14. The compound of any one of claims 1 to 5, wherein R_2 is of the formula:



wherein X is N, and Y is NH, S, or O.

15. The compound of any one of claims 1 to 5, wherein R_2 is of the formula:




16. The compound of any one of claims 1 to 15, wherein R_3 is a substituted or unsubstituted aryl.

17. The compound of claim 16, wherein R_3 is an unsubstituted phenyl moiety.

18. The compound of claim 16, wherein R_3 is a substituted phenyl moiety.

- 10 19. The compound of claim 16, of the formula:




wherein



n is an integer between 1 and 5, inclusive; and


each occurrence of R_3' is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently a hydrogen, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety.



20. The compound of claim 1 of the formula:




21. The compound of claim 1 of one of the formulae:












or



22. The compound of claim 1, of formula:





23. A compound of the formula:



wherein


R_1 is of the formula:



R_1' is $\sim$,
Y is NH or O;

L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety; a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety; a substituted or unsubstituted aryl moiety; a substituted or unsubstituted heteroaryl moiety; or a cyclic ring system, wherein the rings may be aryl, heteroaryl, non-aromatic carbocyclic, or non-aromatic heterocyclic; and

A comprises a functional group selected from the group consisting of:




10

R_2 is hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-\text{OR}_B$; $-\text{C(=O)R}_B$; $-\text{CO}_2\text{R}_B$; $-\text{CN}$; $-\text{SCN}$; $-\text{SR}_B$; $-\text{SOR}_B$; $-\text{SO}_2\text{R}_B$; $-\text{NO}_2$; $-\text{N(R}_B)_2$; $-\text{NHC(O)R}_B$; or $-\text{C(R}_B)_3$; wherein each occurrence of R_B is independently a hydrogen, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety; and


R_3 is halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently a hydrogen, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety;

10 or pharmaceutically acceptable salts thereof.


24. The compound of claim 23, wherein the compound is of one of the formulae:



25. The compound of claim 23 or 24, wherein R_1 is of the formula:




26. The compound of any one of claims 23 to 25, wherein R_1' is of the formula:



wherein n is an integer between 0 and 15, inclusive.

27. The compound of claim 26, wherein n is 5 or 6.

28. The compound of any one of claims 23 to 27, wherein R_2 is of the formula:



wherein

m is an integer between 0 and 8, inclusive;

X is O, S, CH_2 , NH, or NR_2' ; and

R_2' is aliphatic, heteroaliphatic, acyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl.

29. The compound of claim 28, wherein X is O.


30. The compound of claim 28, wherein X is S.

31. The compound of any one of claims 28 to 30, wherein m is an integer between 1 and 6, inclusive.


32. The compound of claim 31, wherein m is 1.


33. The compound of any one of claims 28 to 32, wherein R_2' is substituted or unsubstituted heteroaryl.

34. The compound of any one of claims 23 to 27, wherein R_2 is selected from one of the following:




35. The compound of any one of claims 23 to 27, wherein R_2 is selected from one of the following:






36. The compound of any one of claims 23 to 27, wherein R_2 is of the formula:




wherein X is N, and Y is NH, S, or O.

37. The compound of any one of claims 23 to 27, wherein R_2 is of the formula:



38. The compound of any one of claims 23 to 37, wherein R_3 is a substituted or unsubstituted aryl.
39. The compound of claim 38, wherein R_3 is an unsubstituted phenyl moiety.
- 10 40. The compound of claim 38, wherein R_3 is a substituted phenyl moiety.
41. The compound of claim 38 of the formula:




wherein


n is an integer between 1 and 5, inclusive; and



each occurrence of R_3' is independently hydrogen; halogen; cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; substituted or unsubstituted, branched or unbranched acyl; substituted or unsubstituted aryl; substituted or unsubstituted heteroaryl; $-OR_C$; $-C(=O)R_C$; $-CO_2R_C$; $-CN$; $-SCN$; $-SR_C$; $-SOR_C$; $-SO_2R_C$; $-NO_2$; $-N(R_C)_2$; $-NHC(O)R_C$; or $-C(R_C)_3$; wherein each occurrence of R_C is independently a hydrogen, an aliphatic moiety, a heteroaliphatic moiety, an acyl moiety; an aryl moiety; a heteroaryl moiety; alkoxy; aryloxy; alkylthio; arylthio; amino, alkylamino, dialkylamino, heteroaryloxy; or heteroarylthio moiety.



42. The compound of claim 23, of the formula:



43. The compound of claim 23, of one of the formulae:









O=C(O)CCCCCCC(=O)Nc1ccc(cc1)C2COC(C2Cc3ccccc3)SCc4cccnc4

- 125




45. A pharmaceutical composition comprising a compound of any one of claims 1-44 and a pharmaceutically acceptable excipient.
46. The pharmaceutical composition of claim 45, further comprising a cytotoxic agent.
47. The pharmaceutical composition of claim 45, further comprising a proteasome inhibitor.
48. The pharmaceutical composition of claim 47, wherein the proteasome inhibitor is bortezomib.
- 10 49. The pharmaceutical composition of claim 45, further comprising an aggresome inhibitor.
50. Use of a compound of any one of claims 1-44 for inhibiting histone deacetylase.
51. Use of the compound according to claim 50, wherein the histone deacetylase is purified.

52. Use of the compound according to claim 50, wherein the histone deacetylase is in a cell.
53. Use of the compound according to claim 50, wherein the histone deacetylase is HDAC6.
54. Use of a compound of any one of claims 1-44 for inhibiting tubulin deacetylase.
55. Use of a compound of any one of claims 1-44 for inhibiting an aggresome.
56. Use of a compound of any one of claims 1-44 in the manufacture of a medicament for the treatment of a proliferative disorder.
- 10 57. The use according to claim 56, wherein the proliferative disorder is cancer.
58. The use according to claim 56, wherein the proliferative disorder is an inflammatory disease.
59. Use of a compound of any one of claims 1-44 in the manufacture of a medicament for the treatment of a protein degradation disorder.
60. The use according to claim 59, wherein the protein degradation disorder is a protein deposition disorder.
61. The use according to claim 59, wherein the protein degradation disorder is a neurodegenerative disorder.
- 20 62. The use according to claim 59 wherein the medicament further comprises a therapeutically effective amount of a proteasome inhibitor.
63. The use according to claim 62, wherein the proteasome inhibitor is bortezomib (VELCADE®).
64. A method of preparing a compound of the formula:




the method comprising steps of:


providing an epoxy alcohol of formula:



reacting the epoxy alcohol with a reagent having the structure R_BXH under suitable conditions to generate a diol of formula:




reducing the nitro group to generate a diol of formula:




10

reacting the amino group with acylating agent to generate a diol of formula:



reacting the diol with a reagent having the structure $R_3CH(OMe)_2$, or other acetal under suitable conditions to generate a 1,3-dioxane of formula:




wherein

R_1' is $-L-A$;

L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety; a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety; a substituted or unsubstituted aryl moiety; a substituted or unsubstituted heteroaryl moiety; or a cyclic ring system, wherein the rings may be aryl, heteroaryl, non-aromatic carbocyclic, or non-aromatic heterocyclic; and

A comprises a functional group selected from the group consisting of:




R_B is hydrogen, a protecting group, or an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, or heteroaromatic moiety;

X is $-O-$, $-C(R')_2-$, $-S-$, or $-NR'-$, wherein R' is hydrogen, a protecting group, or an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic or heteroaromatic moiety; and


R_3 is an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, or heteroaromatic moiety.

65. A method of preparing a compound of the formula:




the method comprising steps of:


providing an epoxy alcohol of formula:




reacting the epoxy alcohol with a reagent having the structure R_BXH under suitable conditions to generate a diol of formula:




reducing the nitro group to generate a diol of formula:



reacting the amino group with acylating agent to generate a diol of formula:



reacting the diol with an aldehyde of formula R_3CHO under suitable conditions to generate a 1,3-dioxane of formula:




wherein

R_1' is $-L-A$;

L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety; a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety; a substituted or unsubstituted aryl moiety; a substituted or unsubstituted heteroaryl moiety; or a cyclic ring system, wherein the rings may be aryl, heteroaryl, non-aromatic carbocyclic, or non-aromatic heterocyclic; and

10

A comprises a functional group selected from the group consisting of:




R_B is hydrogen, a protecting group, or an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, or heteroaromatic moiety;

X is $-\text{O}-$, $-\text{C}(\text{R}')_2-$, $-\text{S}-$, or $-\text{NR}'-$, wherein R' is hydrogen, a protecting group, or an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic or heteroaromatic moiety; and


R_3 is an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, or heteroaromatic moiety.

66. A method of synthesizing a compound of the formula:




the method comprising steps of:


providing a beta-hydroxy ketone of formula:




reducing the ketone to generate a diol of formula:




reducing the nitro group to generate a diol of formula:



reacting the amino group with acylating agent to generate a diol of formula:



reacting the diol with a reagent having the structure $\text{R}_2\text{CH}(\text{OMe})_2$ or other acetal under suitable conditions to generate a compound of formula:




wherein

R_1' is $-L-A$;

L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety; a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety; a substituted or unsubstituted aryl moiety; a substituted or unsubstituted heteroaryl moiety; or a cyclic ring system, wherein the rings may be aryl, heteroaryl, non-aromatic carbocyclic, or non-aromatic heterocyclic; and

10


A comprises a functional group selected from the group consisting of:



R_2 is an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, or heteroaromatic moiety; and


R_3 is an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, or heteroaromatic moiety.

67. A method of synthesizing a compound of the formula:




the method comprising steps of:


providing a beta-hydroxy ketone of formula:




reducing the ketone to generate a diol of formula:




reducing the nitro group to generate a diol of formula:



reacting the amino group with acylating agent to generate a diol of formula:



reacting the diol with an aldehyde of formula $R_2\text{CHO}$ under suitable conditions to generate a compound of formula:




wherein

R₁' is -L-A;

L is a substituted or unsubstituted, cyclic or acyclic, branched or unbranched aliphatic moiety; a substituted or unsubstituted, cyclic or acyclic, branched or unbranched heteroaliphatic moiety; a substituted or unsubstituted aryl moiety; a substituted or unsubstituted heteroaryl moiety; or a cyclic ring system, wherein the rings may be aryl, heteroaryl, non-aromatic carbocyclic, or non-aromatic heterocyclic; and

10


A comprises a functional group selected from the group consisting of:




R₂ is an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, or heteroaromatic moiety; and

R_3 is an aliphatic, alicyclic, heteroaliphatic, heterocyclic, aromatic, or heteroaromatic moiety.


68. The method of claim 66 or 67, wherein the step of providing a beta-hydroxy ketone having the structure:



comprises reacting an aldehyde of formula:




with a ketone of formula:



10 under suitable conditions to form the beta-hydroxy ketone.

69. The method of any one of claims 66 to 68, wherein R_3 is substituted or unsubstituted aryl or heteroaryl.
70. The method of any one of claims 66 to 68, wherein R_3 is substituted or unsubstituted phenyl.
71. The method of any one of claims 66 to 68, wherein the beta-hydroxy ketone is of the formula:






Fig. 1

General Synthesis Scheme for Isotubacin Analogs




Fig. 2

Scheme Isotubacin

The synthesis of Isotubacin proceeds through the following steps:

- 4-nitrobenzaldehyde reacts with allyl bromide ($\text{BrCH}_2\text{CH=CH}_2$) and zinc (Zn) to form 4-nitro-1-allyl-2-nitrobenzene.
- 4-nitro-1-allyl-2-nitrobenzene is converted to 4-nitro-1-allyl-2-nitrobenzene-1-ol using NaHMDs and Boc_2O .
- 4-nitro-1-allyl-2-nitrobenzene-1-ol is converted to 4-nitro-1-allyl-2-nitrobenzene-1-ol using IBBr in Toluene.
- 4-nitro-1-allyl-2-nitrobenzene-1-ol is converted to 4-nitro-1-allyl-2-nitrobenzene-1-ol using K_2CO_3 .
- 4-nitro-1-allyl-2-nitrobenzene-1-ol is converted to 4-nitro-1-allyl-2-nitrobenzene-1-ol using DIPEA and SH.
- 4-nitro-1-allyl-2-nitrobenzene-1-ol is converted to 4-nitro-1-allyl-2-nitrobenzene-1-ol using NaBH_4 and $\text{Cu}(\text{OAc})_2$ in EtOH.
- 4-nitro-1-allyl-2-nitrobenzene-1-ol is converted to 4-nitro-1-allyl-2-nitrobenzene-1-ol using MeO, DIPEA, and CH_2Cl_2 .
- 4-nitro-1-allyl-2-nitrobenzene-1-ol is converted to 4-nitro-1-allyl-2-nitrobenzene-1-ol using KOH, H_2NOH , and MeOH.
- 4-nitro-1-allyl-2-nitrobenzene-1-ol is converted to Isotubacin using PPTS and a benzylidene acetal.

Fig. 3

Fig. 3

General Synthesis Scheme for Isoisotubacin Analogs




Fig. 4

The reaction scheme illustrates the synthesis of a cyclic nucleoside derivative through several steps:

- Starting Material:** 4-nitrobenzaldehyde reacts with acetophenone in the presence of $\text{Na}_2\text{CO}_3 \cdot \text{H}_2\text{O}$ to form an α,β -unsaturated ketone intermediate.
- Reduction:** The intermediate is reduced using BH_3Py and TiCl_4 to yield a diol with a nitro group.
- Reduction:** The nitro group is reduced to an amino group using NaBH_4 and $\text{Cu}(\text{OAc})_2$.
- Protection:** The diol is protected as an isopropylidene acetal using $i\text{BuOCOC}$ and RCOOH .
- Hydrolysis:** The acetal is hydrolyzed using PPTS to yield a diol with an amino group.
- Reaction with 2,4,6-triphenyl-1,3,5-triazine:** The diol reacts with 2,4,6-triphenyl-1,3,5-triazine in the presence of MeO and S to form a cyclic nucleoside derivative.
- Final Product:** The cyclic nucleoside derivative is further modified using MeO and S to yield the final product, which is a cyclic nucleoside derivative with a triazine ring.

Fig. 5




Fig. 6





Fig. 7



Medicinal Chemistry: Development of HT/IF Quantitative Assay

Fig. 8



10/14




Fig. 10A






Fig. 10B

**Fig. 11**


12/14



TDAC Inhibitory Activity

Fig. 12

Fig. 13



	Average KD (uM)	Average KD (M)	Percent Bound
DHM Tubacin	96.85	9.69E-05	87.5%
Tubacin	5.53	5.53E-06	99.2%
59	100.8	1.01E-04	87.1%
60	67.3	6.73E-05	91.0%
81	244.25	2.44E-04	73.6%
84	80.85	8.09E-05	89.4%
93	152	1.52E-04	81.7%
94	95.8	9.58E-05	87.7%
165	476	4.76E-04	58.8%
201	73.7	7.37E-05	90.2%
207	202.75	2.03E-04	77.0%
208	184.75	1.85E-04	78.6%
233	99.8	9.98E-05	87.2%
234	563.5	5.64E-04	54.7%
235	676	6.76E-04	50.1%
236	108	1.08E-04	86.3%
1428	557	5.57E-04	55.0%
SAHA	333	3.33E-04	67.1%

HSA Binding by SPR

Fig. 14

