

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2012/076601 A1

(43) International Publication Date

14 June 2012 (14.06.2012)

(51) International Patent Classification:

A23F 3/14 (2006.01) *A23L 1/105* (2006.01)
A23F 3/16 (2006.01) *A23L 1/19* (2006.01)
A23F 5/24 (2006.01) *A23L 1/212* (2006.01)
A23F 5/36 (2006.01) *A23L 1/40* (2006.01)
A23F 5/40 (2006.01) *A23L 2/395* (2006.01)
A23F 5/44 (2006.01) *A23L 2/52* (2006.01)
A23L 1/10 (2006.01) *A23G 1/50* (2006.01)

(74) Agent: **LOMHOLT, Stig**; Avenue Nestlé 55, CH-1800 Vevey (CH).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:

PCT/EP2011/072076

(22) International Filing Date:

7 December 2011 (07.12.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

10194212.6 8 December 2010 (08.12.2010) EP

(71) **Applicant** (for all designated States except US): **NESTEC S.A. [CH/CH]**; Avenue Nestlé 55, CH-1800 Vevey (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SCHAFFER-LE-QUART, Christelle** [FR/CH]; Chemin de la Paisible 16, CH-1083 Mezieres (CH). **ROGER, Olivier Yves** [FR/CH]; Chemin de Belflori 7, CH-1510 Moudon (CH). **WAVREILLE, Anne-Sophie** [FR/CH]; Chemin des Rosiers 4, appartement 8, CH-1004 Lausanne (CH). **WEINGAND-ZIADE, Alexandra** [FR/FR]; 257 Rue de la Côte d'Arbère, F-01220 Divonne-les-Bains (FR). **MAR-JANOVIC, Nicolas** [CH/CH]; Chemin de la Damataire 22, CH-1009 Pully (CH). **TE BIESEBEKE, Rob** [NL/CH]; Les Bordes 15, CH-1372 Bavois (CH).

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))

WO 2012/076601 A1

(54) **Title:** INSTANT DRINK POWDERS COMPRISING HYDROLYZED WHOLE GRAIN

(57) **Abstract:** The present invention relates to instant drink powders comprising a primary ingredient of particles or agglomerated particles having a particle size below 500 μm , a hydrolyzed whole grain composition, an alpha-amylase or fragments thereof, which alpha-amylase or fragments thereof show no hydrolytic activity towards dietary fibers when in the active state; and a moisture content of at most 5% (w/w) of the instant drink powder.

Instant drink powders comprising hydrolyzed whole grain

Technical field of the invention

The present invention relates to instant drink powders being supplemented with 5 whole grain. In particular the present invention relates to instant drink powders which are supplemented with hydrolysed whole grain, where neither taste or viscosity nor organoleptic properties of the instant drink powders have been compromised.

10 Background of the invention

There is now extensive evidence emerging mainly from epidemiological studies that a daily intake of three servings of whole grain products, i.e. 48 g of whole grain, is positively associated with decreased risk of cardiovascular diseases, increased insulin sensitivity and decreased risk of type 2 diabetes onset, obesity 15 (mainly visceral obesity) and digestive system cancers. These health benefits of the whole grains are reported to be due to the synergistic role of the dietary fibers and other components, such as vitamins, minerals and bioactive phytochemicals.

The regulatory authorities in Sweden, the US and the UK have already approved 20 specific heart health claims based on the available scientific substantiation.

Food products comprising dietary fibers are also growing in popularity with consumers, not just because whole grain consumption is now included in some national dietary recommendations but also because whole grain products are considered wholesome and natural. Recommendations for whole grain 25 consumption have been set up by government authorities and expert groups to encourage consumers to eat whole grains. For instance, in the U.S.A., recommendations are to consume 45-80 g of whole grain per day. However, data provided by national dietary surveys in the United Kingdom, the U.S.A. and China show that whole grain consumption varies between 0 and 30 g whole grains per 30 day.

The lack of whole grain products offered on the shelves and the poor organoleptic properties of the available whole grain products are generally identified as barriers

for whole grain consumption and restrict the amount of whole grain to be added to e.g. an instant drink powder, because, when increased amounts of whole grain are added the physical and organoleptic properties of the instant drink powder changes dramatically.

5

Whole grains are also a recognised source of dietary fibers, phytonutrients, antioxidants, vitamins and minerals. According to the definition given by the American Association of Cereal Chemists (AACC), whole grains, and food made from whole grains, consist of the entire grain seed. The entire grain seed

10 comprises the germ, the endosperm and the bran. It is usually referred to as the kernel.

Moreover, in recent years, consumers have increased attention to the label of food products, e.g. instant drink powders, and they expect manufactured food products

15 to be as natural and healthy as possible. Therefore, it is desirable to develop food and drink processing technologies and food and drink products that limit the use of non-natural food additives, even when such non-natural food additives have been fully cleared by health or food safety authorities.

20 Given the health benefits of whole grain cereal, it is desirable to provide a whole grain ingredient having as much intact dietary fibers as possible. Instant drink powders are a good vehicle for delivering whole grain and to increase the whole grain content of a product or a serving, it is of course possible to increase the serving size. But this is not desirable as it results in a greater calorie intake.

25 Another difficulty in just increasing the whole grain content of the product is that it usually impacts on physical properties such as the taste, texture and the overall appearance of the instant drink powders (organoleptic parameters), as well as its processability.

30 The consumer is not willing to compromise on instant drink powders organoleptic properties, in order to increase his daily whole grain intake. Taste, texture and overall appearance are such organoleptic properties.

Obviously, industrial line efficiency is a mandatory requirement in the food industry. This includes handling and processing of raw materials, forming of the

instant drink powders, packaging and later storing, in warehouses, on the shelf or at home.

US 4,282,319 relates to a process for the preparation of hydrolyzed products from 5 whole grain, and such derived products. The process includes an enzymatic treatment in an aqueous medium with a protease and an amylase. The obtained product may be added to different types of products. US 4,282,319 describe a complete degradation of the proteins present in the whole grain.

10 US 5,686,123 discloses a cereal suspension generated by treatment with both an alpha-amylase and a beta-amylase both specifically generating maltose units and have no glucanase effect.

Thus, it is an object of the present invention to provide instant drink powders that 15 are rich in whole grains and in dietary fibers, while maintaining a low calorie intake, that provide an excellent consumption experience to the consumer, and that may be easily industrialised at a reasonable cost without compromising the organoleptic parameters.

20 **Summary of the invention**

Accordingly, in a first aspect the invention relates to an instant drink powder comprising

- a primary ingredient of particles or agglomerated particles having a particle size below 500 µm;
- 25 - a hydrolyzed whole grain composition;
- an alpha-amylase or fragment thereof, which alpha-amylase or fragment thereof shows no hydrolytic activity towards dietary fibers when in the active state; and
- a moisture content of at most 5% (w/w) of the instant drink powder.

30

Another aspect of the present invention relates to a process for preparing a instant drink powder according to the present invention, said process comprising:

- 1) preparing a hydrolyzed whole grain composition, comprising the steps of:

- a) contacting a whole grain component with an enzyme composition in water, the enzyme composition comprising at least one alpha-amylase, said enzyme composition showing no hydrolytic activity towards dietary fibers,
- 5 b) allowing the enzyme composition to react with the whole grain component, to provide a whole grain hydrolysate,
- c) providing the hydrolyzed whole grain composition by inactivating said enzymes when said hydrolysate has reached a viscosity comprised between 50 and 5000 mPa.s measured at 65°C;
- 10 2) providing the instant drink powder by mixing the hydrolyzed whole grain composition with a primary ingredient of particles or agglomerated particles having a particle size below 5 mm.

A further aspect relates to an instant drink comprising an instant drink powder according to the present invention reconstituted in a liquid component.

15

Brief description of the drawings

Figure 1 shows a thin layer chromatography analysis of various enzymes put in contact with dietary fibres. The legend for the different tracks is the following:

A0: pure arabinoxylan spot (blank)

20 B0: pure beta-glucan spot (blank)

A: arabinoxylan spot after incubation with the enzyme noted below the track (BAN, Validase HT 425L and Alcalase AF 2.4L)

β: beta-glucan spot after incubation with the enzyme noted below the track (BAN, Validase HT 425L and Alcalase AF 2.4L)

25 E0: enzyme spot (blank)

Figure 2 shows size exclusion chromatography (SEC) of β-Glucan and arabinoxylanmolecular weight profile without enzyme addition (plain line) and after incubation with Alcalase 2.4L (dotted line). A) Oat β-glucan; B) Wheat

30 arabinoxylan.

Figure 3 shows size exclusion chromatography (SEC) of β -Glucan and arabinoxylan molecular weight profile without enzyme addition (plain line) and after incubation with Validase HT 425L (dotted line). A) Oat β -glucan; B) Wheat arabinoxylan.

5

Figure 4 shows size exclusion chromatography (SEC) of β -Glucan and arabinoxylan molecular weight profiles without enzyme addition (plain line) and after incubation with MATS L (dotted line). A) Oat β -glucan; B) Wheat arabinoxylan.

10 Detailed description of the invention

The inventors of the present invention have surprisingly found that by treating the whole grain component with an alpha-amylase and optionally with a protease the whole grain will become less viscous and the following mixing into the instant drink powder may be easier. This results in the possibility to increase the amount 15 of whole grains in the product. Furthermore, the alpha-amylase treatment also results in a reduced need for adding sweetener such as sucrose to the instant drink powder products.

Thus in a first aspect the invention relates to an instant drink powder comprising 20

- a primary ingredient of particles or agglomerated particles having a particle size below 500 μm ;
- a hydrolyzed whole grain composition;
- an alpha-amylase or fragments thereof, which alpha-amylase or fragments thereof show no hydrolytic activity towards dietary fibers when in the 25 active state; and
- a moisture content of at most 5% (w/w) of the instant drink powder.

Several advantages of having an instant drink powder comprising a hydrolyzed whole grain component according to the invention may exist:

30

- I. An increase in whole grain and fiber content may be provided in the final product, while the organoleptic parameters of the product are substantially not affected;
- II. Dietary fibers from the whole grain may be preserved;

III. Greater sense of satiety substantially without affecting the organoleptic parameters of the product and slower digestion. Currently, there are limitations for enriching instant drink powders with whole grain due to non-flowable viscosity, grainy texture, and taste issues. However, the use of 5 hydrolyzed whole grain according to the present invention in instant drink powders allow for providing the desired viscosity, a smooth texture, minimal flavor impact, and added nutritional health and wellness values;

IV. An additional advantage may be to improve the carbohydrate profile of the instant drink powder products by replacing traditional externally supplied 10 sweeteners such as glucose syrup, high fructose corn syrup, invert syrup, maltodextrine, sucrose, fiber concentrate, inulin, etc. with a more wholesome sweetener source.

In the present context the term "instant drink powder" relates to powders 15 containing at least a primary ingredient intended to be reconstituted in a liquid such as milk or water. Examples of a primary ingredient are a tea mixture, a fruit mixture, a coffee mixture, a chicory mixture, a cocoa mixture, a creamer mixture or a soup mixture.

Thus, in an embodiment the instant drink powder is to be reconstituted in a liquid 20 component before consumption. In a further embodiment the liquid component is water, milk, milk components or mixtures hereof. It is to be understood that the liquid may be hot or cold.

A quality parameter of the instant drink powder and an important parameter in 25 respect of the product processability is the viscosity of the hydrolysed whole grain composition. In the present context the term "viscosity" is a measurement of "thickness" or fluidability of a fluid. Thus, viscosity is a measure of the resistance of a fluid which is being deformed by either shear stress or tensile stress. If not indicated otherwise viscosity is given in mPa.s.

30

Viscosity may be measured using a Rapid Visco Analyser from Newport Scientific. The Rapid Visco Analyser measures the resistance of the product to the stirring action of a paddle. The viscosity is measured after 10 minutes stirring, at 65°C and 50 rpm.

35

The whole grain component may be obtained from different sources. Examples of whole grain sources are semolina, cones, grits, flour and micronized grain (micronized flour). The whole grains may be grounded, preferably by dry milling. Such grounding may take place before or after the whole grain component being 5 contacted with the enzyme composition according to the invention. The whole grains may be milled/grounded to a particle size, characterized by the median particle diameter (volume distribution), X50 in the range of 1-500 µm, such as 50-500 µm, such as in the range of 100-300 µm e.g. in the range of 150-250 µm.

10 In an embodiment of the present invention the whole grain component may be heat treated to limit rancidity and microbial count.

Whole grains are cereals of monocotyledonous plants of the Poaceae family (grass family) cultivated for their edible, starchy grains. Examples of whole grain cereals 15 include barley, rice, black rice, brown rice, wild rice, buckwheat, bulgur, corn, millet, oat, sorghum, spelt, triticale, rye, wheat, wheat berries, teff, canary grass, Job's tears and fonio. Plant species that do not belong to the grass family also produce starchy seeds or fruits that may be used in the same way as cereal grains, are called pseudo-cereals. Examples of pseudo-cereals include amaranth, 20 buckwheat, tartar buckwheat and quinoa. When designating cereals, this will include both cereal and pseudo-cereals.

Thus, the whole grain component according to the invention may originate from a cereal or a pseudo-cereal. Thus, in an embodiment the hydrolyzed whole grain composition is obtained from a plant selected from the group consisting of barley, 25 rice, brown rice, wild rice, black rice, buckwheat, bulgur, corn, millet, oat, sorghum, spelt, triticale, rye, wheat, wheat berries, teff, canary grass, Job's tears, fonio, amaranth, buckwheat, tartar buckwheat, quinoa, other variety of cereals and pseudo-cereals and mixtures thereof. In general the source of grain depends on the product type, since each grain will provide its own taste profile.

30 Whole grain components are components made from unrefined cereal grains. Whole grain components comprise the entire edible parts of a grain; i.e. the germ, the endosperm and the bran. Whole grain components may be provided in a variety of forms such as ground, flaked, cracked or other forms, as is commonly 35 known in the milling industry.

In the present context the phrasing "a hydrolyzed whole grain composition" refers to enzymatically digested whole grain components or a whole grain component digested by using at least an alpha-amylase, which alpha-amylase shows no

- 5 hydrolytic activity towards dietary fibers when in the active state. The hydrolyzed whole grain composition may be further digested by the use of a protease, which protease shows no hydrolytic activity towards dietary fibers when in the active state.
- 10 In the present context it is also to be understood that the phrase "a hydrolyzed whole grain composition" is also relating to enzymatic treatment of flour and subsequent reconstitution of the whole grain by blending flour, bran and germ. It is also to be understood that reconstitution may be done before the use in the final product or during mixing in a final product. Thus, reconstitution of whole
- 15 grains after treatment of one or more of the individual parts of the whole grain also forms part of the present invention.

Prior to or after grinding of the whole grain, the whole grain component may be subjected to a hydrolytic treatment in order to breakdown the polysaccharide

- 20 structure and optionally the protein structure of the whole grain component.

The hydrolyzed whole grain composition may be provided in the form of a liquid, a concentrate, a powder, a juice or a puree. If more than one type of enzymes is used it is to be understood that the enzymatic processing of the whole grains may

- 25 be performed by sequential addition of the enzymes, or by providing an enzyme composition comprising more than one type of enzyme.

In the present context the phrase "an enzyme showing no hydrolytic activity towards dietary fibers when in the active state" should be understood as also

- 30 encompassing the enzyme mixture from which the enzyme originates. For example, the proteases, amylases, glucose isomerase and amyloglucosidase described in the present context may be provided as an enzyme mixture before use which is not completely purified and thus, comprise enzymatic activity towards e.g. dietary fibers. However, the activity towards dietary fibers may also
- 35 come from the specific enzyme if the enzyme is multi-functional. As used in here,

the enzymes (or enzyme mixtures) are devoid of hydrolytic activity towards dietary fibers.

The term "no hydrolytic activity" or "devoid of hydrolytic activity towards dietary fibers" may encompass up to 5% degradation of the dietary fibers, such as up to 3%, such as up to 2% and such as up to 1% degradation. Such degradation may be unavoidable if high concentrations or extensive incubation times are used.

The term "In the active state" refers to the capability of the enzyme or enzyme mixture to perform hydrolytic activity, and is the state of the enzyme before it is inactivated. Inactivation may occur both by degradation and denaturation.

In general the weight percentages throughout the application are given as percentage by weight on a dry matter basis unless otherwise stated.

15

The instant drink powder according to the invention may comprise a protease which shows no hydrolytic activity towards dietary fibers when in the active state. The advantage of adding a protease according to the invention is that the viscosity of the hydrolyzed whole grain may be further lowered, which may also 20 result in a decrease in the viscosity of the final product. Thus, in an embodiment according to the invention the instant drink powder comprises said protease or fragment thereof at a concentration of 0.0001 to 5% (w/w) by weight of the total whole grain content, such as 0.01-3%, such as 0.01-1%, such as 0.05-1%, such as 0.1-1%, such as 0.1-0.7%, or such as 0.1-0.5%. The optimal concentration of 25 added proteases depends on several factors. As it has been found that the addition of protease during production of the hydrolyzed whole grain may result in a bitter off-taste, addition of protease may be considered as a tradeoff between lower viscosity and off-taste. In addition the amount of protease may also depend on the incubation time during production of the hydrolyzed whole grain. For 30 example a lower concentration of protease may be used if the incubation time is increased.

Proteases are enzymes allowing the hydrolysis of proteins. They may be used to decrease the viscosity of the hydrolyzed whole grain composition. Alcalase 2.4L 35 (EC 3.4.21.62), from Novozymes is an example of a suitable enzyme.

Depending on the incubation time and concentration of protease a certain amount of the proteins from the hydrolyzed whole grain component may be hydrolyzed to amino acid and peptide fragments. Thus, in an embodiment 1-10% of the proteins 5 from the whole grain composition is hydrolyzed, such as 2-8%, e.g. 3-6%, 10-99%, such as 30-99%, such as 40-99%, such as 50-99%, such as 60-99%, such as 70-99%, such as 80-99%, such as 90-99%, or such as 10-40%, 40-70%, and 60-99%. Again protein degradation may result in a lowered viscosity and improved organoleptic parameters.

10

In the present context the phrase "hydrolyzed protein content" refers to the content of hydrolyzed protein from the whole grain composition unless otherwise defined. The protein may be degraded into larger or smaller peptide units or even into amino acid components. The person skilled in the art will know that during 15 processing and storage small amount of degradation will take place which is not due to external enzymatic degradation.

In general it is to be understood that the enzymes used in the production of the hydrolyzed whole grain composition (and therefore also present in the final 20 product) is different from the corresponding enzymes naturally present in the whole grain component.

Since the instant drink powders according to the invention may also comprise proteins from sources, different from the hydrolyzed whole grain component, 25 which are not degraded, it may be appropriate to evaluate the protein degradation on more specific proteins present in the whole grain composition. Thus, in an embodiment the degraded proteins are whole grain proteins, such as gluten proteins, globulins, albumins and glycoproteins.

30 Amylase (EC 3. 2. 1. 1) is an enzyme classified as a saccharidase: an enzyme that cleaves polysaccharides. It is mainly a constituent of pancreatic juice and saliva, needed for the breakdown of long-chain carbohydrates such as starch, into smaller units. Here, alpha-amylase is used to hydrolyse gelatinized starch in order to decrease the viscosity of the hydrolyzed whole grain composition. Validase HT 35 425L, Validase RA from Valley Research, Fungamyl from Novozymes and MATS

from DSM are examples of alpha-amylases suitable for the present invention. Those enzymes show no activity towards the dietary fibers in the processing conditions used (duration, enzyme concentrations). On the contrary, e.g. BAN from Novozymes degrades dietary fibers besides starch into low molecular weight fibers or oligosaccharides, see also example 3.

In an embodiment of the present invention the enzymes show no activity towards the dietary fibers when the enzyme concentration is below 5% (w/w), such as below, 3% (w/w), e.g. below 1% (w/w), such as below 0.75% (w/w), e.g. below 10 0.5% (w/w).

Some alpha-amylases generate maltose units as the smallest carbohydrate entities, whereas others are also able to produce a fraction of glucose units. Thus, in an embodiment the alpha-amylase or fragments thereof is a mixed sugar 15 producing alpha-amylase, including glucose producing activity, when in the active state. It has been found that some alpha-amylases both comprise glucose producing activity while having no hydrolytic activity towards dietary fibers when in the active state. By having an alpha-amylase which comprises glucose producing activity an increased sweetness may be obtained, since glucose has 20 almost twice the sweetness of maltose. In an embodiment of the present invention a reduced amount of external sugar source needs to be added separately to the instant drink powders when a hydrolysed whole grain composition according to the present invention is used. When an alpha-amylase comprising glucose producing activity is used in the enzyme composition, it may 25 become possible to dispense or at least reduce the use of other external sugar sources or non-sugar sweeteners.

In the present context the term "external sugar source" relates to sugars not originally present or originally generated in the hydrolysed whole grain 30 composition. Examples of such external sugar source could be sucrose, lactose, and artificial sweeteners.

Amyloglucosidase (EC 3.2.1.3) is an enzyme able to release glucose residues from starch, maltodextrins and maltose by hydrolysing glucose units from the non- 35 reduced end of the polysaccharide chain. The sweetness of the preparation

increases with the increasing concentration of released glucose. Thus, in an embodiment the instant drink powder further comprises an amyloglucosidase or fragments thereof. It may be advantageous to add an amyloglucosidase to the production of the hydrolyzed whole grain composition, since the sweetness of the

5 preparation increases with the increasing concentration of released glucose. It may also be advantageous if the amyloglucosidase did not influence health properties of the whole grains, directly or indirectly. Thus, in an embodiment the amyloglucosidase shows no hydrolytic activity towards dietary fibers when in the active state. An interest of the invention, and particularly of the process for

10 preparing the instant drink powders according to the invention, is that it allows reducing the sugar (e.g. sucrose) content of the instant drink powders when compared to products described in the prior art. When an amyloglucosidase is used in the enzyme composition, it may become possible to dispense with other external sugar sources e.g. the addition of sucrose.

15 However, as mentioned above certain alpha-amylases are able to generate glucose units, which may add enough sweetness to the product making the use of amyloglucosidase dispensable. Furthermore, application of amyloglucosidase also increases production costs of the instant drink powders and, hence, it may be desirable to limit the use of amyloglucosidases. Thus, in yet an embodiment the

20 instant drink powder according to the invention does not comprise an amyloglucosidase such as an exogenic amyloglucosidase.

Glucose isomerase (D-glucose ketoisomerase) causes the isomerization of glucose to fructose. Thus, in an embodiment of the present invention the instant drink

25 powder further comprises a glucose isomerase or fragments thereof, which glucose isomerase or fragments thereof show no hydrolytic activity towards dietary fibers when in the active state. Glucose has 70-75% the sweetness of sucrose, whereas fructose is twice as sweet as sucrose. Thus, processes for the manufacture of fructose are of considerable value because the sweetness of the

30 product may be significantly increased without the addition of an external sugar source (such as sucrose or artificial sweetening agents).

A number of specific enzymes or enzyme mixtures may be used for production of the hydrolyzed whole grain composition according to the invention. The

35 requirement is that they show substantially no hydrolytic activity in the process

conditions used towards dietary fibers. Thus, in an embodiment the alpha-amylase may be selected from Validase HT 425L and Validase RA from Valley Research, Fungamyl from Novozymes and MATS from DSM, the protease may be selected from the group consisting of Alcalase, iZyme B and iZyme G

5 (novozymes).

The concentration of the enzymes according to the invention in the instant drink powder product may influence the organoleptic parameters of the instant drink powder. In addition the concentration of enzymes may also be adjusted by

10 changing parameters such as temperature and incubation time. Thus, in an embodiment the instant drink powder comprises 0.0001 to 5% by weight of the total whole grain content in the instant drink powders of at least one of:

- an alpha-amylase or fragments thereof, which alpha-amylase or fragment thereof shows no hydrolytic activity towards dietary fibers when in the 15 active state;
- an amyloglucosidase or fragment thereof, which amyloglucosidase shows no hydrolytic activity towards dietary fibers when in the active state; and
- a glucose isomerase or fragment thereof, which amyloglucosidase shows no hydrolytic activity towards dietary fibers when in the active state.

20 In yet an embodiment the instant drink powder comprises 0.001 to 3% of the alpha-amylase by weight of the total whole grain content in the instant drink powder, such as 0.01-3%, such as 0.01-0.1%, such as 0.01-0.5%, such as 0.01-0.1%, such as 0.03-0.1%, such as 0.04-0.1%. In yet an embodiment the instant drink powder comprises 0.001 to 3% of the amyloglucosidase by weight of the 25 total whole grain content in the instant drink powder, such as 0.001-3%, such as 0.01-1%, such as 0.01-0.5%, such as 0.01-0.5%, such as 0.01-0.1%, such as 0.03-0.1%, such as 0.04-0.1%. In another further embodiment the instant drink powder comprises 0.001 to 3% of the glucose isomerase by weight of the total whole grain content in the instant drink powder, such as 0.001-3%, such as 0.01-30 1%, such as 0.01-0.5%, such as 0.01-0.5%, such as 0.01-0.1%, such as 0.03-0.1%, such as 0.04-0.1%.

Beta-amylases are enzymes which also break down saccharides, however beta-amylases mainly have maltose as the smallest generated carbohydrate entity. Thus, in an embodiment the instant drink powder according to the invention does not comprise a beta-amylase, such as an exogenic beta-amylase. By avoiding

5 beta-amylases a larger fraction of the starches will be hydrolyzed to glucose units since the alpha amylases do have to compete with the beta-amylases for substrates. Thus, an improved sugar profile may be obtained. This is in contrast to US 5,686,123 which discloses a cereal suspension generated by treatment with both an alpha-amylase and a beta-amylase.

10 In certain instances the action of the protease is not necessary, to provide a sufficient low viscosity. Thus, in an embodiment according to the invention, the instant drink powder does not comprise the protease, such as an exogenic protease. As described earlier the addition of protease may generate a bitter off-taste which in certain instances is desirable to avoid. This is in contrast to US

15 4,282,319 which discloses a process including enzymatic treatment with a protease and an amylase.

In general the enzymes used according to the present invention for producing the hydrolyzed whole grain composition show no hydrolytic activity towards dietary

20 fibers when in the active state. Thus, in a further embodiment the hydrolyzed whole grain composition has a substantial intact beta-glucan structure relative to the starting material. In yet a further embodiment the hydrolyzed whole composition has a substantial intact arabinoxylan structure relative to the starting material. By using the one or more enzymes according to the invention for the

25 production of the hydrolyzed whole grain composition, a substantial intact beta-glucan and arabinoxylan structure may be maintained. The degree of degradation of the beta-glucan and arabinoxylan structures may be determined by Size-exclusion chromatography (SEC). This SEC technique has been described in more detail in "Determination of beta-Glucan Molecular Weight Using SEC

30 with Calcofluor Detection in Cereal Extracts Lena Rimsten, Tove Stenberg, Roger Andersson, Annica Andersson, and Per Åman. Cereal Chem. 80(4):485-490", which is hereby incorporated by reference.

In the present context the phrase "substantial intact structure" is to be understood as for the most part the structure is intact. However, due to natural degradation in any natural product, part of a structure (such as beta-glucan structure or arabinoxylan structure) may be degraded although the degradation 5 may not be due to added enzymes. Thus, "substantial intact structure" is to be understood that the structure is at least 95% intact, such as at least 97%, such as at least 98%, or such as at least 99% intact.

In the present context enzymes such as proteases, amylases, glucose isomerases 10 and amyloglucosidases refer to enzymes which have been previously purified or partly purified. Such proteins/enzymes may be produced in bacteria, fungi or yeast, however they may also have plant origin. In general such produced enzymes will in the present context fall under the category "exogenic enzymes". Such enzymes may be added to a product during production to add a certain 15 enzymatic effect to a substance. Similar, in the present context, when an enzyme is disclaimed from the present invention such disclaimer refers to exogenic enzymes. In the present context such enzymes e.g. provide enzymatic degradation of starch and proteins to decrease viscosity. In relation to the process of the invention it is to be understood that such enzymes may both be in solution 20 or attached to a surface, such as immobilized enzymes. In the latter method the proteins may not form part of the final product.

As mentioned earlier, the action of the alpha-amylase results in a useful sugar profile which may affect taste and reduce the amount of external sugar or 25 sweetener to be added to the final product.

In an embodiment of the present invention the hydrolysed whole grain composition has a glucose content of at least 0.25% by weight of the hydrolysed whole grain composition, on a dry matter basis, such as at least 0.35%, e.g. at least 0.5%. 30 Depending on the specific enzymes used the sugar profile of the final product may change. Thus, in an embodiment the instant drink powder has a maltose to glucose ratio below 144:1, by weight in the product, such as below 120:1, such as below 100:1 e.g. below 50:1, such as below 30:1, such as below 20:1 or such as below 10:1.

If the only starch processing enzyme used is a glucose generating alpha-amylase, a larger fraction of the end product will be in the form of glucose compared to the use of an alpha-amylase specifically generating maltose units. Since glucose has a higher sweetness than maltose, this may result in that the addition of a further 5 sugar source (e.g. sucrose) can be dispensed. This advantage may be further pronounced if the ratio is lowered by the conversion of the maltose present in the hydrolyzed whole grain to glucose (one maltose unit is converted to two glucose units).

The maltose to glucose ratio may be further lowered if an amyloglucosidase is 10 included in the enzyme composition since such enzymes also generates glucose units.

If the enzyme composition comprises a glucose isomerase a fraction of the glucose is changed to fructose which has an even higher sweetness than glucose. Thus, in an embodiment the instant drink powder has a maltose to glucose + 15 fructose ratio below 144:1 by weight in the product, such as below 120:1, such as below 100:1 e.g. below 50:1, such as below 30:1, such as below 20:1 or such as below 10:1.

Furthermore, in an embodiment of the present invention the instant drink powder may have a maltose to fructose ratio below 230:1 by weight in the product, such 20 as below 144:1, such as below 120:1, such as below 100:1 e.g. below 50:1, such as below 30:1, such as below 20:1 or such as below 10:1.

In the present context the phrasing "total content of the whole grain" is to be understood as the combination of the content of "hydrolyzed whole grain composition" and "solid whole grain content". If not indicated otherwise, "total 25 content of the whole grain" is provided as % by weight in the final product. In an embodiment the instant drink powder has a total content of the whole grain in the range of 1-35% by weight of the instant drink powder such as 5-35%, such as 5-25%, such as 10-25%, and such as 15-25%, such as 1-20%, such as 1-15%, such as 1-10%, and such as 1-7%.

30

In the present context the phrasing "content of the hydrolyzed whole grain composition" is to be understood as the % by weight of hydrolyzed whole grains

in the final product. Hydrolyzed whole grain composition content is part of the total content of the whole grain composition. Thus, in an embodiment the instant drink powder according to the invention has a content of the hydrolyzed whole grain composition in the range 1-35% by weight of the instant drink powder such as 10-30%, such as 20-28%, such as 1-20%, such as 1-10% and such as 1-5%. The amount of the hydrolyzed whole grain composition in the final product may depend on the type of product. By using the hydrolyzed whole grain composition according to the invention in an instant drink powder, a higher amount of hydrolyzed whole grains may be added (compared to a non-hydrolyzed whole grain composition) without substantially affecting the organoleptic parameters of the product because of the increased amount of soluble fibers in the hydrolysed whole grain.

It would be advantageous to have an instant drink powder comprising a high content of dietary fibers without compromising the organoleptic parameters of the product. Thus, in yet an embodiment the instant drink powder has a content of dietary fibers in the range 0.1-20% by weight of the instant drink powder, such as in the range 0.5-15%, such as in the range 0.5-10%, such as in the range of 0.5-3%, even more preferably in the range of 1-2% (w/w). An instant drink powder according to the invention may be provided with high amounts of dietary fibers by the addition of the hydrolyzed whole grain component provided by the present invention. This may be done due to the unique setup of the process according to the present invention.

Dietary fibers are the edible parts of plants that are not broken down by digestion enzymes. Dietary fibers are fermented in the human large intestine by the microflora. There are two types of fibers: soluble fibers and insoluble fibers. Both soluble and insoluble dietary fibers can promote a number of positive physiological effects, including a good transit through the intestinal tract which helps to prevent constipation, or a feeling of fullness. Health authorities recommend a consumption of between 20 and 35 g per day of fibers, depending on the weight, gender, age and energy intake.

Soluble fibers are dietary fibers that undergo complete or partial fermentation in the large intestine. Examples of soluble fibers from cereals include beta-glucans,

arabinoxylans, arabinogalactans and resistant starch type 2 and 3, and oligosaccharides deriving from the latters. Soluble fibers from other sources include pectins, acacia gum, gums, alginate, agar, polydextrose, inulins and galacto-oligosaccharides for instance. Some soluble fibers are called prebiotics,

- 5 because they are a source of energy for the beneficial bacteria (e.g. Bifidobacteria and Lactobacilli) present in the large intestine. Further benefits of soluble fibers include blood sugar control, which is important in diabetes prevention, control of cholesterol, or risk reduction of cardiovascular disease.
- 10 Insoluble fibers are the dietary fibers that are not fermented in the large intestine or only slowly digested by the intestinal microflora. Examples of insoluble fibers include celluloses, hemicelluloses, resistant starch type 1 and lignins. Further benefits of insoluble fibers include promotion of the bowel function through stimulation of the peristalsis, which causes the muscles of the colon to work more,
- 15 become stronger and function better. There is also evidence that consumption of insoluble fibers may be linked to a reduced risk of gut cancer.

The total solid content of the instant drink powder according to the invention may vary. Thus, in another embodiment the total moisture content is at the most 5%

- 20 by weight of the instant drink powder, such as at the most 3%, such as at the most 2%. Examples of factors influencing the moisture content may be the amount of the hydrolyzed whole grain composition and the degree of hydrolysis in this composition. In the present context the phrasing "total solid content" equals 100 minus moisture content (%) of the product.

25

The instant drink powder according to the invention may be suited for several types of powder product. Thus, in an embodiment the primary ingredient is a coffee mixture, tea mixture, a fruit mixture, a chicory mixture, a cocoa mixture, a creamer mixture or a soup mixture.

30

When used as a coffee mixture, the amount of coffee may vary. Thus, in an embodiment the coffee mixture comprises 5-85% (w/w) coffee relative to the total amount of the instant drink powder, such as 5-70%, such as 5-50% or such as 5-30%.

35

When used as a chicory mixture, the amount of chicory may vary. Thus, in an embodiment the chicory mixture comprises 5-85% (w/w) coffee relative to the total amount of the instant drink powder, such as 5-70%, such as 5-50% or such as 5-30%.

5

It would be advantageously if an instant drink powder with good organoleptic parameters, such as sweetness, could be obtained, without addition of large amounts of external sugar sources. Thus, in another embodiment the instant drink powder has a content of sucrose less than 70% by weight of the instant drink powder, such as less than 50%, less than 30%, such as less than 25%, such as less than 15% and such as less than 10%. Since the hydrolyzed whole grain composition supplements the instant drink powder with a source of carbohydrates, such as glucose and maltose, the instant drink powder is also sweetened from a natural sugar source different from the external sugar source. Thus, the amount of added external sweetener may be limited. In yet an embodiment the sucrose has a crystal size in the range of 100 µm to 450 µm.

Sucrose is a widely used sweetener in food products, however others sugars may also be used. Thus, in a further embodiment the instant drink powder comprises at least one sugar different from sucrose, wherein said sugar different from sucrose is a monosaccharide and/or a disaccharide and/or an oligosaccharide. In yet an embodiment the monosaccharide is glucose, galactose, dextrose, fructose or any combination thereof. In yet another embodiment the disaccharide is maltose, lactose or any combination thereof. It is of course also to be understood that sweeteners different from sucrose may be used. Thus, in an embodiment the sweetener is sugar or artificial intense sweetener or natural intense sweetener or mixtures thereof.

Humectants are often added to products which are to be in a dry or semi-dry state. In an embodiment the instant drink powder does not comprise a humectant. Supplementary ingredients of the instant drink powder include vitamins and minerals, preservatives such as tocopherol, and emulsifiers, such as lecithin, protein powders, cocoa solid, alkylresorcinols, phenolics and other active ingredients, such as DHA, caffeine, and prebiotics.

35

In a further embodiment the instant drink powder has a fat content of less than 20% by weight of the instant drink powder, such as less than 12%, such as less than 10%, such as less than 5% and such as less than 3% by weight of the instant drink powder. The amount of fat may vary depending on the type of

5 product. Fat components are preferably vegetable fats such as cocoa butter, rapeseed oil, sunflower oil or palm oil, preferably not hydrogenated.

In yet an embodiment the instant drink powder may have salt content in the range 0-2% by weight of the instant drink powder. In a more specific embodiment

10 the salt is sodium chloride.

The instant powder according to the invention may comprise one or more further components. Thus, in a further embodiment the instant drink powder is an instant drink powder mix. In yet an embodiment the instant drink powder mix comprises

15 a further ingredient such as a whitener, a sweetener, a flavor component, a bulking agent, and/or a foaming agent.

Depending on the specific type of instant drink powder, different flavor components may be added to provide the desired taste. Thus, in a further

20 embodiment the flavor component is selected from the group consisting of coffee, cocoa, fruit, malt, vegetable, and commercialized flavors. Addition of a flavor component may improve factors such as taste, viscosity and the nutritional profile.

25 Features of the instant drink powder relating the particles in the powder. Thus, in a further embodiment according to the invention the particles or the agglomerated particles are free-flowing. In the present invention the term "free-flowing" relates to adjacent laying particles which do not, or substantially not, stick to each other. The term "substantially not stick to each other" involves at most 5% of the

30 particles stick to each other, such as at most 3%, e.g. at most 1%.

In yet an embodiment the instant drink powder has a particle size, characterized by the median particle diameter (volume distribution), X_{50} in the range of 50-500 μm , such as in the range of 100-300 μm e.g. in the range of 150-250 μm . The

particle size may determine how fast the powder is reconstituted in a liquid component.

For the aspect of providing the product of the present invention a process is
5 provided for preparing an instant drink powder, said process comprising:

1) preparing a hydrolyzed whole grain composition, comprising the steps of:

- a) contacting a whole grain component with an enzyme composition in water, the enzyme composition comprising at least one alpha-amylase, 10 said enzyme composition showing no hydrolytic activity towards dietary fibers,
- b) allowing the enzyme composition to react with the whole grain component, to provide a whole grain hydrolysate,
- c) providing the hydrolyzed whole grain composition by inactivating said enzymes when said hydrolysate has reached a viscosity comprised 15 between 50 and 5000 mPa.s measured at 65°C;

2) providing the instant drink powder by mixing the hydrolyzed whole grain composition with a primary ingredient of particles or agglomerated particles having a particle size below 5 mm.

20 In an embodiment the enzyme composition further comprises a protease or fragment thereof, which protease or fragment thereof shows no hydrolytic activity towards dietary fibers when in the active state. Similar, the enzyme composition may comprise an amyloglucosidase and/or glucose isomerase according to the present invention.

25

Several parameters of the process may be controlled to provide the instant drink powder according to the invention. Thus, in an embodiment step 1b) is performed at 30-100°C, preferably 50 to 85°C. In a further embodiment step 1b) is performed for 1 minute to 24 hours, such as 1 minute to 12 hours, such as 1 30 minute to 6 hours, such as 5-120 minutes. In yet an embodiment step 1b) is performed at 30-100°C for 5-120 minutes.

In yet a further embodiment step 1c) is allowed to proceed at 70-150°C for at least 1 second, such as 1-5 minutes, such as 5-120 minutes, such as 5-60 minutes. In an additional embodiment step 1c) is performed by heating to at least 90°C for 5-30 minutes.

5

In yet an embodiment the reaction in step 1c) is stopped when the hydrolysate has reached a viscosity comprised between 50 and 4000 mPa.s, such as between 50 and 3000 mPa.s, such as between 50 and 1000 mPa.s, such as between 50 and 500 mPa.s. In an additional embodiment viscosity is measured at TS 50.

10 In another embodiment the hydrolyzed whole grain composition in step 1) is provided when said hydrolysate has reached a total solid content of 25-60%. By controlling viscosity and solid content the hydrolyzed whole grain may be provided in different forms.

In an additional embodiment the hydrolyzed whole grain component in step 1c) is

15 provided in the form of a liquid, a concentrate, a powder, a juice or a pure. An advantage of being able to provide the hydrolyzed whole grain component in different forms is that it is possible to add hydrolyzed whole grain in high concentrations without the risk of diluting the product (the instant drink powder).

Similarly, when the hydrolyzed whole grain composition is to be mixed with a

20 powder, it may be advantageously to provide the hydrolyzed whole grain composition in a dry state, such as a powder. Thus, an embodiment of the invention relates to the process wherein the further ingredient(s), the hydrolyzed whole grain composition and the primary ingredient(s) are mixed in a dry state.

25 The above parameters can be adjusted to regulate the degree of starch degradation, the sugar profile, the total solid content and to regulate the overall organoleptic parameters of the final product.

To improve the enzymatic processing of the whole grain component it may be

30 advantageous to process the grains before or after the enzymatic treatment.

Thus, in an embodiment the whole grains are finely grounded to a particle size, characterized by the median particle diameter (volume distribution), X_{50} in the range of 1-1000 μm , such as 5-500 μm , such as in the range of 100-300 μm e.g.

in the range of 150-250 µm. By grounding the grains are larger surface area is made accessible to the enzymes, thereby speeding up the process. In addition the organoleptic parameters are improved by using a smaller particle size of the grains. In an additional embodiment the whole grains are roasted or toasted

5 before or after enzymatic treatment. Roasting and toasting may improve the taste of the final product.

To prolong the storage time of the product several treatment can be performed.

Thus, in an embodiment the process further comprises at least one of the

10 following treatments: UHT, pasteurization, thermal treatment, retort and any other thermal or non-thermal treatments, such as pressure treatment. In a further embodiment the instant drink powder is applied to an enclosure under aseptic conditions. In yet an embodiment the instant drink powder is applied to an enclosure under non-aseptic conditions, such as by retort or hot-for-hold.

15

The instant drink powder according to the invention may be reconstituted in a liquid component. Thus, an aspect of the invention relates to an instant drink comprising the instant drink powder according to the invention reconstituted in a liquid component. In a further embodiment the liquid component is water, milk or

20 a milk component or a combination hereof. In yet an embodiment the milk is selected from the group consisting of whole milk, whey fractions, casein, any combination hereof. The instant drink may at least 1.25 g dietary fibers per serving, such as at least 1.5 g dietary fibers per serving, e.g. at least 2 g dietary fibers per serving.

25

In an embodiment a serving may comprise between 4-20 en% protein by weight of the instant drink; 0-20 en% fat by weight of the instant drink and/or 20-70 en% carbohydrate by weight of the instant drink. A serving size may vary from 100 to 250 ml of liquid.

30

It should be noted that embodiments and features described in the context of one of the aspects or embodiments of the present invention also apply to the other aspects of the invention.

All patent and non-patent references cited in the present application, are hereby incorporated by reference in their entirety.

The invention will now be described in further details in the following non-limiting 5 examples.

EXAMPLES

Example 1 - Preparation of a hydrolyzed whole grain composition

10 Enzyme compositions comprising Validase HT 425L (alpha-amylase) optionally in combination with Alcalase 2.4 L (protease) were used for the hydrolysis of wheat, barley and oats.

15 Mixing may be performed in a double jacket cooker, though other industrial equipment may be used. A scraping mixer works continuously and scraps the inner surface of the mixer. It avoids product burning and helps maintaining a homogeneous temperature. Thus enzyme activity is better controlled. Steam may be injected in the double jacket to increase temperature while cold water is used to decrease it.

20

In an embodiment, the enzyme composition and water are mixed together at room temperature, between 10 and 25°C. At this low temperature, the enzymes of the enzyme composition have a very weak activity. The whole grain component is then added and the ingredients are mixed for a short period of time, usually 25 less than 20 minutes, until the mixture is homogeneous.

The mixture is heated progressively or by thresholds to activate the enzymes and hydrolyse the whole grain component.

30 Hydrolysis results in a reduction of the viscosity of the mixture. When the whole grain hydrolysate has reached a viscosity comprised between 50 and 5000 mPa.s measured at 65°C and e.g. a total solid content of 25 to 60% by weight, the enzymes are inactivated by heating the hydrolysate at a temperature above 100°C, preferably by steam injection at 120°C.

Enzymes are dosed according to the quantity of total whole grain. Quantities of enzymes are different depending on the type of whole grain component, as protein rates are different. The ratio water/whole grain component can be adapted

5 according to required moisture for the final liquid whole grain. Usually, the water/whole grain component ratio is 60/40. Percents are by weight.

Hydrolysed whole wheat	
Whole wheat flour	Substrate
Enzyme amylase	0.10% based on the substrate
Enzyme protease	0.05% based on the substrate

Hydrolysed whole barley	
Whole barley flour	Substrate
Enzyme amylase	0.10% based on the substrate
Enzyme protease	0.05% based on the substrate

Hydrolysed whole oats	
Whole oats flour	Substrate
Enzyme amylase	0.10% based on the substrate
Enzyme protease	0.05% based on the substrate

10

Example 2 - Sugar profile of the hydrolyzed whole grain composition

Hydrolyzed whole grain compositions comprising wheat, barley and oat were prepared according to the method in example 1.

15 Carbohydrates HPAE:

The hydrolyzed whole grain compositions were analysed by HPAE for illustrating the sugar profile hydrolysed whole grain composition.

20 Carbohydrates are extracted with water, and separated by ion chromatography on an anion exchange column. The eluted compounds are detected electrochemically by means of a pulsed amperometric detector and quantified by comparison with the peak areas of external standards.

Total dietary fibres:

Duplicate samples (defatted if necessary) are digested for 16 hours in a manner that simulates the human digestive system with 3 enzymes (pancreatic alpha-5 amylase, protease, and amyloglucosidase) to remove starch and protein. Ethanol is added to precipitate high molecular weight soluble dietary fibre. The resulting mixture is filtered and the residue is dried and weighed. Protein is determined on the residue of one of the duplicates; ash on the other. The filtrate is captured, concentrated, and analyzed via HPLC to determine the value of low molecular 10 weight soluble dietary fibre (LMWSF).

Whole wheat:

	Wheat Reference	Wheat Hydrolysed Alcalase/Validase
Total sugars (% w/w))	2.03	24.36
Glucose	0.1	1.43
Fructose	0.1	0.1
Lactose (monohydrate)	<0.1	<0.1
Sucrose	0.91	0.69
Maltose (monohydrate)	0.91	22.12
Mannitol	<0.02	<0.02
Fucose	<0.02	<0.02
Arabinose	<0.02	0.02
Galactose	<0.02	<0.02
Xylose	<0.02	<0.02
Mannose	<0.02	<0.02
Ribose	<0.02	<0.02
Insoluble and soluble fibers	12.90	12.94
LMW fibers	2.63	2.96
Total fibers	15.53	15.90

Whole oats:

	Oats Reference	Oats Hydrolysed Alcalase/Validase
Total sugars (% w/w))	1.40	5.53
Glucose	0.1	0.58
Fructose	0.1	0.1
Lactose (monohydrate)	<0.1	<0.1
Sucrose	1.09	1.03
Maltose (monohydrate)	0.11	3.83
Mannitol	<0.02	<0.02
Fucose	<0.02	<0.02
Arabinose	<0.02	<0.02
Galactose	<0.02	<0.02
Xylose	<0.02	<0.02
Mannose	<0.02	<0.02
Ribose	<0.02	<0.02
Insoluble and soluble fibers	9.25	11.28
LMW fibers	0.67	1.21
Total fibers	9.92	12.49

Whole Barley:

	Barley Reference	Barley Hydrolysed Alcalase/Validase
Total sugars (% w/w))	1.21	5.24
Glucose	0.1	0.61
Fructose	0.1	0.1
Lactose (monohydrate)	<0.1	<0.1
Sucrose	0.90	0.88
Maltose (monohydrate)	0.11	3.65
Mannitol	<0.02	<0.02
Fucose	<0.02	<0.02
Arabinose	<0.02	<0.02
Galactose	<0.02	<0.02
Xylose	<0.02	<0.02
Mannose	<0.02	<0.02
Ribose	<0.02	<0.02
Glucose	0.1	0.61
Fructose	0.1	0.1
Insoluble and soluble fibers	9.70	10.44
LMW fibers	2.23	2.63
Total fibers	11.93	13.07

5 The results clearly demonstrate that a significant increase in the glucose content is provided by the hydrolysis where the glucose content of the hydrolysed barley

is 0.61% (w/w) on a dry matter basis; the glucose content of the hydrolysed oat is 0.58% (w/w) on a dry matter basis; and the glucose content of the hydrolysed wheat is 1.43% (w/w) on a dry matter basis.

- 5 Furthermore, the results also demonstrates that the maltose:glucose ratio is ranging from about 15:1 to about 6:1.

Thus, based on these results a new sugar profile is provided having a increased sweetness compared to the prior art.

- 10 In conclusion, an increased sweetness may be obtained by using the hydrolyzed whole grain composition according to the invention and therefore the need for further sweetening sources may be dispensed or limited.

In addition, the results demonstrate that the dietary fiber content is kept intact

- 15 and the ratio and amount of soluble and insoluble fibers are substantially the same in the non-hydrolyzed whole grain and in the hydrolyzed whole grain composition.

Example 3 – Hydrolytic activity on dietary fibers

- 20 The enzymes Validase HT 425L (Valley Research), Alcalase 2.4L (Novozymes) and BAN (Novozymes) were analysed using a thin layer chromatography analysis for activity towards arabinoxylan and beta-glucan fibre extracts both components of dietary fibers of whole grain.

- 25 The results from the thin layer chromatography analysis showed that the amylase Validase HT and the protease Alcalase showed no hydrolytic activity on either beta-glucan or arabinoxylan, while the commercial alpha-amylase preparation, BAN, causes hydrolysis of both the beta-glucan and arabinoxylan, see figure 1. See also example 4.

Example 4 - Oat β -Glucan and Arabinoxylan molecular weight profile following enzymatic hydrolysis

Hydrolysis:

A solution of 0.5 % (w/v) of Oat β -Glucan medium viscosity (Megazyme) or Wheat

5 Arabinoxylan medium viscosity (Megazyme) was prepared in water.

The enzyme was added at an enzyme to substrate ratio (E/S) of 0.1 % (v/v). The reaction was allowed to proceed at 50°C for 20 minutes, the sample was then placed at 85°C during 15 min to enable starch gelatinization and hydrolysis. The enzymes were finally inactivated at 95°C for 15 minutes. Different batches of the following enzymes have been evaluated.

Alcalase 2.4L (Valley Research): batch BN 00013

batch 62477

batch 75039

15 Validase HT 425L (Valley Research): batch RA8303A

batch 72044

MATS L (DSM): batch 408280001

Molecular weight analysis

20 Hydrolyzed samples were filtered on a syringe filter (0.22 µm) and 25 µL were
21 injected on a High Pressure Liquid Chromatography Agilent 1200 series equipped
22 with 2 TSKgel columns in serie (G3000PWXL 7,8 x 300 mm), (GMPWXL 7,8X 30
23 mm) and with a guard column (PWXL 6 x 44 mm). (Tosoh Bioscience)
24 Sodium Nitrate 0.1M/ at 0.5ml/min was used as running buffer. Detection was
25 done by reflective index measurement.

Results

On figures 2-4 graphs for both a control (no enzyme) and test with enzymes are plotted. However, since there are substantially no difference between the graphs 30 it may be difficult to differentiate both graphs from each other.

Conclusions

No shift in oat beta glucan and wheat arabinoxylan fibre molecular weight profile was determined following hydrolysis with the Alcalase 2.4 L (figure 2), Validase 35 HT 425 L (figure 3) or MATS L (figure 4).

Example 5 - Preparation of a Coffee Mix containing 6 g of whole grain

A hydrolyzed whole grain composition is prepared according to example 1.

In the following example, the hydrolyzed whole grain composition is mixed with all
 5 the other ingredients in a dry mixing process. This process is done in 1 or 2 steps
 according the type of raw materials used. If minor ingredients are used, they are
 pre-mixed first with part of a major ingredient. This pre-mix is then mixed with all
 the other major ingredients.

The instant powder is then reconstituted in cold or hot water, cold or hot milk,

10 cold or hot milk components or a combination hereof.

Coffee Mix with hydrolyzed whole grain powder:

Ingredient	% by weight
Hydrolyzed whole grain powder	5-35%
Instant coffee powder	5-30%
Whitener powder	0-40%
Sugar	0-65%
Milk or milk component	0-40%
Vitamins & minerals	0-2%
Flavour	0-2%
Total: 12g to 40g to be reconstituted in 100 to 250ml of liquid.	

Claims

1. An instant drink powder comprising:

- a primary ingredient of particles or agglomerated particles having a particle size below 500 µm;
- 5 - a hydrolyzed whole grain composition;
- an alpha-amylase or fragments thereof, which alpha-amylase or fragments thereof show no hydrolytic activity towards dietary fibers when in the active state; and
- a moisture content of at most 5% (w/w) of the instant drink powder.

10

2. The instant drink powder according to claim 1, further comprising a protease or fragment thereof, at a concentration of 0.001-5% by weight of the total whole grain content, which protease or fragment thereof shows no hydrolytic activity towards dietary fibers when in the active state.

15

3. The instant drink powder according to any of the preceding claims, with the proviso that it does not comprise a beta-amylase.

20
20

4. The instant drink powder according to claim 1 or 3, with the proviso that it does not comprise the protease.

25
25

5. The instant drink powder according to any one of the preceding claims, wherein the instant drink powder further comprises an amyloglucosidase or fragments thereof, which amyloglucosidase or fragments thereof show no hydrolytic activity towards dietary fibers when in the active state.

30

6. The instant drink powder according to any one of the preceding claims, wherein the instant drink powder further comprises an glucose isomerase or fragments thereof, which glucose isomerase or fragments thereof show no hydrolytic activity towards dietary fibers when in the active state.

7. The instant drink powder according to any of the preceding claims, wherein the hydrolyzed whole grain composition has a substantial intact beta-glucan structure relative to the starting material.

8. The instant drink powder according to any of the preceding claims, wherein the hydrolyzed whole grain composition has a substantial intact arabinoxylan structure relative to the starting material.

5

9. The instant drink powder according to any of the preceding claims, having a total content of the whole grain in the range of 1-35% by weight of the instant drink powder such as 5-35%, such as 5-25%, such as 10-25%, and such as 15-25%.

10

10. The instant drink powder according to any of the preceding claims, wherein the instant drink powder is an instant drink powder mix.

11. The instant drink powder according to claim 10, wherein the instant drink powder mix comprises a further ingredient such as a whitener, a sweetener, a flavor component, a bulking agent, and/or a foaming agent.

12. The instant drink powder according to any of the preceding claims, wherein the instant drink powder has a maltose to glucose ratio below 144:1 by weight in the product, such as below 120:1, such as below 100:1, such as below 50:1, such as below 30:1, such as below 20:1 and such as below 10:1.

13. The instant drink powder according to any one of the preceding claims, wherein the primary ingredient is a coffee mixture, tea mixture, a fruit mixture, a chicory mixture, a cocoa mixture, a creamer mixture or a soup mixture.

25

14. The instant drink powder according to any one of the preceding claims, wherein the instant drink powder is to be reconstituted in a liquid component before consumption.

30 15. A process for preparing an instant drink powder according to any of claims 1-14, said process comprising:

1) preparing a hydrolyzed whole grain composition, comprising the steps of:

- a) contacting a whole grain component with an enzyme composition in water, the enzyme composition comprising at least one alpha-amylase, said enzyme composition showing no hydrolytic activity towards dietary fibers,
 - 5 b) allowing the enzyme composition to react with the whole grain component, to provide a whole grain hydrolysate,
 - c) providing the hydrolyzed whole grain composition by inactivating said enzymes when said hydrolysate has reached a viscosity comprised between 50 and 5000 mPa.s measured at 65°C;
- 10 2) providing the instant drink powder by mixing the hydrolyzed whole grain composition with a primary ingredient of particles or agglomerated particles having a particle size below 5 mm.

1/4






Fig. 1

2/4

A**B****Fig. 2**

3/4



A**B**

Fig. 3

4/4



A**B**

Fig.4

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2011/072076

A. CLASSIFICATION OF SUBJECT MATTER				
INV.	A23F3/14	A23F3/16	A23F5/24	A23F5/36
	A23F5/44	A23L1/10	A23L1/105	A23L1/19
	A23L1/40	A23L2/395	A23L2/52	A23G1/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A23F A23L A23G

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, FSTA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2010/023351 A1 (RAISIO NUTRITION LTD [FI]; ALHO-LEHTO PIRJO [FI]; KUUSISTO PAEIVI [FI]) 4 March 2010 (2010-03-04) the whole document -----	1-15
X	EP 0 231 729 A1 (BERGKVIST ROLF RAGNAR [SE]; CLAESSEN KARL OLOV [SE]) 12 August 1987 (1987-08-12) page 2, paragraph 3 - page 3, paragraph 1; claims 1-12; examples 6,7 -----	1-15
X	US 4 894 242 A (MITCHELL CHERYL R [US] ET AL) 16 January 1990 (1990-01-16) column 517 - column 8, line 26; claims 1-26; example 6 ----- -/-	1-15

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
30 January 2012	08/02/2012
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Korb, Margit

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2011/072076

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 282 319 A (CONRAD ERNST) 4 August 1981 (1981-08-04) cited in the application the whole document -----	1-15
A	US 2005/202145 A1 (DORR TILLMANN [DE] ET AL DOERR TILLMANN [DE] ET AL) 15 September 2005 (2005-09-15) page 2, paragraph 19; claims 1-34; examples 1-1,2 -----	1-15
A	WO 2009/040249 A1 (NESTEC SA [CH]; KESSLER ULRICH [CH]; DUFFEY JEAN-LOUIS [FR]; DREYER MA) 2 April 2009 (2009-04-02) claims 1-18 -----	1-15
A	EP 0 524 484 A1 (PLANTEXTRAKT PFLANZEN EXTRAKTI [DE]) 27 January 1993 (1993-01-27) column 1, line 1 - column 3, line 51; claims 1-10 -----	1-15
A	EP 0 172 010 A2 (MORINAGA MILK INDUSTRY CO LTD [JP]) 19 February 1986 (1986-02-19) claims 1-7; examples 1,2 -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2011/072076

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 2010023351	A1	04-03-2010	CA CN EP FI US WO	2734386 A1 102137596 A 2343990 A1 121844 B1 2011159145 A1 2010023351 A1		04-02-2010 27-07-2011 20-07-2011 13-05-2011 30-06-2011 04-03-2010
EP 0231729	A1	12-08-1987	DE DE EP ES JP	3688836 D1 3688836 T2 0231729 A1 2059308 T3 S63502360 A		09-09-1993 25-11-1993 12-08-1987 16-11-1994 08-09-1988
US 4894242	A	16-01-1990	NONE			
US 4282319	A	04-08-1981	AR AT AU AU CA CH DE ES FI FR GB GR IE IT JP JP JP LU NL NO NZ PL PT US	214595 A1 365420 B 522176 B2 4063078 A 1120775 A1 644736 A5 2844896 A1 474311 A1 783153 A 2406665 A1 2007960 A 64395 A1 48036 B1 1101253 B 1510048 C 54095760 A 63063179 B 80380 A1 7810395 A 783516 A 188663 A 210359 A1 68667 A 4282319 A		29-06-1979 11-01-1982 20-05-1982 01-05-1980 30-03-1982 31-08-1984 19-04-1979 16-04-1979 19-04-1979 18-05-1979 31-05-1979 21-03-1980 05-09-1984 28-09-1985 26-07-1989 28-07-1979 06-12-1988 19-03-1979 20-04-1979 19-04-1979 11-02-1981 18-06-1979 01-11-1978 04-08-1981
US 2005202145	A1	15-09-2005	AT AU BR CA CN EP JP KR MX US WO ZA	412339 T 2003281488 A1 0312797 A 2493934 A1 1668205 A 1524911 A1 2005536994 A 20050033609 A PA04012958 A 2005202145 A1 2004008870 A1 200501009 A		15-11-2008 09-02-2004 03-05-2005 29-01-2004 14-09-2005 27-04-2005 08-12-2005 12-04-2005 16-05-2005 15-09-2005 29-01-2004 25-10-2006
WO 2009040249	A1	02-04-2009	AR AU CA CN EP	068284 A1 2008303653 A1 2700582 A1 101808530 A 2194795 A1		11-11-2009 02-04-2009 02-04-2009 18-08-2010 16-06-2010

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No
PCT/EP2011/072076

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
		JP 2010539913 A		24-12-2010
		KR 20100072221 A		30-06-2010
		PE 09242009 A1		27-07-2009
		RU 2010116774 A		10-11-2011
		TW 200939968 A		01-10-2009
		US 2010215818 A1		26-08-2010
		UY 31366 A1		30-04-2009
		WO 2009040249 A1		02-04-2009

EP 0524484	A1	27-01-1993	DE 4123124 A1	14-01-1993
			EP 0524484 A1	27-01-1993

EP 0172010	A2	19-02-1986	EP 0172010 A2	19-02-1986
			JP 1472397 C	27-12-1988
			JP 61047143 A	07-03-1986
			JP 63020492 B	27-04-1988
			US 4632832 A	30-12-1986
