



(86) Date de dépôt PCT/PCT Filing Date: 2008/03/14
(87) Date publication PCT/PCT Publication Date: 2008/09/18
(85) Entrée phase nationale/National Entry: 2009/09/09
(86) N° demande PCT/PCT Application No.: GB 2008/000894
(87) N° publication PCT/PCT Publication No.: 2008/110815
(30) Priorités/Priorities: 2007/03/15 (EP07251094.4);
2007/09/27 (US11/905,163); 2007/12/04 (GB0723728.2)

(51) Cl.Int./Int.Cl. *A61K 9/10* (2006.01),
A61K 31/59 (2006.01), *A61K 45/06* (2006.01),
A61P 17/06 (2006.01)

(71) Demandeur/Applicant:
DRUG DELIVERY SOLUTIONS LIMITED, GB

(72) Inventeurs/Inventors:
WHEELER, DEREK ALFRED, GB;
STEELE, DAVID FRASER, GB;
GEORGIU, MICHELLE, GB;
SINDET-PEDERSEN, STEEN, GB

(74) Agent: SMART & BIGGAR

(54) Titre : COMPOSITION TOPIQUE DE POLYAPHRON AVEC VITAMINE D ET CORTICOSTEROIDE
(54) Title: POLYAPHRON TOPICAL COMPOSITION WITH VITAMIN D AND CORTICOSTEROID

(57) **Abrégé/Abstract:**

A composition suitable for topical application comprising a continuous phase and at least one discontinuous phase, said composition comprising at least one polyaphron dispersion, at least one vitamin D or vitamin D analogue and at least one corticosteroid.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 September 2008 (18.09.2008)

PCT

(10) International Publication Number
WO 2008/110815 A1

(51) International Patent Classification:

A61K 9/10 (2006.01) A61P 17/06 (2006.01)
A61K 31/59 (2006.01) A61K 45/06 (2006.01)

(GB). SINDET-PEDERSEN, Steen [DK/DK]; c/o Drug Delivery Solutions Limited, Suite H13, The Leatherhead Enterprise Centre, Randalls Road, Leatherhead Surrey KT22 7RY (GB).

(21) International Application Number:

PCT/GB2008/000894

(74) Agents: HAYES, Adrian, Chetwynd et al.; Boulton Wade Tennant, Verulam Gardens, 70 Gray's Inn Road, London WC1X 8BT (GB).

(22) International Filing Date: 14 March 2008 (14.03.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

07251094.4	15 March 2007 (15.03.2007)	EP
11/905,163	27 September 2007 (27.09.2007)	US
0723728.2	4 December 2007 (04.12.2007)	GB

(71) Applicant (for all designated States except US): DRUG DELIVERY SOLUTIONS LIMITED [GB/GB]; Suite H13, The Leatherhead Enterprise Centre, Randalls Road, Leatherhead Surrey KT22 7RY (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): WHEELER, Derek, Alfred [GB/GB]; c/o Drug Delivery Solutions Limited, Suite H13, The Leatherhead Enterprise Centre, Randalls Road, Leatherhead Surrey KT22 7RY (GB). STEELE, David, Fraser [GB/GB]; c/o Drug Delivery Solutions Limited, Suite H13, The Leatherhead Enterprise Centre, Randalls Road, Leatherhead Surrey KT22 7RY (GB). GEORGIU, Michelle [GB/GB]; c/o Drug Delivery Solutions Limited, Suite H13, The Leatherhead Enterprise Centre, Randalls Road, Leatherhead Surrey KT22 7RY

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(54) Title: POLYAPHRON TOPICAL COMPOSITION WITH VITAMIN D AND CORTICOSTEROID

(57) Abstract: A composition suitable for topical application comprising a continuous phase and at least one discontinuous phase, said composition comprising at least one polyaphron dispersion, at least one vitamin D or vitamin D analogue and at least one corticosteroid.

WO 2008/110815 A1

POLYAPHRON TOPICAL COMPOSITION WITH VITAMIN D AND CORTICOSTEROID

The present invention relates to a composition for topical application comprising at least one vitamin D or
5 vitamin D analogue and at least one corticosteroid.

It is known for compositions comprising vitamin D or vitamin D analogues to be used for the treatment of a number of skin conditions.

For example, EP-B-474,517 discloses the use of
10 compositions containing one or more 1α -hydroxylated-19-nor-vitamin D compounds with a triple bond in the side chain in the treatment of psoriasis.

US 4,871,723 discloses a process for treating psoriasis by topically applying a composition comprising vitamin D and
15 a wax carrier. Specifically, US 4,871,723 discloses a composition comprising a) a pharmaceutically effective amount of an active-type vitamin D₃, b) a solvent selected from fatty acid esters, higher alcohols with 10 or more carbons and propylene carbonate and c) an oily carrier
20 selected from white vaseline, yellow vaseline and liquid paraffin.

US 2005/002546 A1 discloses a pharmaceutical composition comprising an active vitamin D compound in emulsion pre-concentrate formulations, as well as emulsions
25 and sub-micron droplet emulsions produced therefrom. In particular, the pharmaceutical compositions of US 2005/002546 A1 comprise

- (a) a lipophilic phase component;
- (b) one or more surfactants; and
- 30 (c) an active vitamin D compound.

The surfactant or surfactants are suitably present in an amount of 1% to 90% by weight based on the total weight

- 2 -

of the composition, and preferably from about 5% to about 85% by weight based on the total weight of the composition.

Presently available compositions contain relatively high concentrations of vitamin D or vitamin D analogues and surfactants which often lead to skin irritation and worsening of psoriasis. For example, in 1996 the Food and Drug Administration of America required that the label included in Dovonex (calcipotriene) - a product containing 0.005% calcipotriol - be amended to indicate that approximately 25% of patients experienced skin irritation, and approximately 10% worsening of psoriasis. In addition, it has been reported that some patients treated with Dovonex have developed hypercalcaemia (see, for example, Hardman KA, Heath DA, Nelson HM Hypercalcaemia associated with calcipotriol (Dovonex) treatment. BMJ. 1993 Apr 3;306(6882):896-896).

Further, it is known in the prior art to treat a number of skin conditions by applying a combination of two or more pharmacologically active compounds. For example, in the treatment of psoriasis, it is possible to use a combination treatment involving a vitamin D analogue, such as calcipotriol, and a corticosteroid, where each of the active compounds is formulated in a separate preparation (for example in US-B-6,753,013).

Topical pharmaceutical compositions comprising a combination of a vitamin D analogue and a topical corticosteroid are challenging to manufacture. This is because these compounds are stable at different pH values.

For example, the calcipotriol requires a pH value above 8 for maximum stability, whereas corticosteroids, such as betamethasone (9-fluoro-11,17,21-trihydroxy-16-methylpregna-1,4-diene-3,20-dione), require pH values in the range 4 to 6 for maximum stability. It is therefore difficult to combine

- 3 -

the two active components in a single formulation while maintaining good stability of the active compounds if water is present in the formulation.

US-B-6,753,013 discloses a pharmaceutical composition for dermal use, which contains at least one vitamin D or vitamin D analogue, at least one corticosteroid and a solvent selected to enable the two active components to coexist without significant degradation, despite their different stability profiles. The compositions, however, are wax-based and include wax or similar excipients, such as soft white paraffin and paraffin liquid. A disadvantage of this composition is that, in order to mix the vitamin D or vitamin D analogue and the corticosteroid into the wax, the wax must be heated to a temperature of 70°C. Such elevated temperatures may damage the drugs in the composition. Furthermore, wax based compositions tend to be rather oily, which after application leave a greasy film on the skin. This is undesirable and can lead to patient non-compliance.

A further disadvantage of the compositions of US-B-6,753,013 is that in order for the compositions to provide beneficial results in topical treatment, high levels of vitamin D or vitamin D analogue are required in order for sufficient vitamin D or vitamin D analogue to permeate the skin. This is because the compositions exhibit poor diffusion through skin. This is disadvantageous as vitamin D and vitamin D analogues are known skin irritants.

There is a need to formulate an improved composition suitable for topical application which addresses at least some of the problems of the prior art.

The present inventors have now developed a new composition comprising at least one vitamin D or vitamin D analogue and at least one corticosteroid. The present inventors have surprisingly found that such compositions

- 4 -

have an enhanced dermal diffusion rate and/or improved stability compared to known compositions. Such compositions also have an appropriate viscosity such that they are useful for topical application.

5 Accordingly, the present invention provides a composition suitable for topical application comprising a continuous phase and at least one discontinuous phase, said composition comprising at least one polyaphron dispersion, at least one vitamin D or vitamin D analogue and at least
10 one corticosteroid.

According to another aspect of the present invention there is provided a composition as described herein for use in the treatment of psoriasis.

According to another aspect of the present invention
15 there is provided a composition as described herein for use in the manufacture of a medicament for the treatment of psoriasis.

According to a further aspect of the present invention there is provided a composition as described herein for use
20 in a method of treatment of the human or animal body by therapy.

According to a further aspect there is provided a method of treatment or prophylaxis of psoriasis in a subject which comprises topically applying to a subject an effective
25 amount of a composition as herein described.

In the following description, the meaning of the terms used are as follows: by hydrophilic phase or solvent is meant a liquid phase comprising water, comprising water together with other water-miscible liquids, or comprising a
30 non-aqueous liquid which is miscible with water. By hydrophobic phase or solvent is meant a phase comprising pharmaceutically acceptable liquids such as oils that are immiscible or substantially immiscible with the hydrophilic

- 5 -

phase. By immiscible liquids is meant that when mixed together, they separate to form two distinctly separate liquid phases sharing a well-defined interface. By substantially immiscible is meant that two liquids mixed as above having a well defined interface between two phases where each phase may nevertheless contain small quantities of dissolved molecules of the other phase.

According to another aspect of the present invention there is provided a method of making the composition as described herein comprising the following steps:

- (i) providing a hydrophilic solvent, optionally comprising at least one vitamin D or vitamin D analogue, and/or at least one corticosteroid, and/or a surfactant;
- (ii) providing a hydrophobic solvent optionally comprising at least one vitamin D or vitamin D analogue, and/or at least one corticosteroid, and/or a surfactant;
- (iii) mixing the hydrophilic solvent with the hydrophobic solvent under suitable conditions to form the composition comprising at least one polyaphron dispersion, at least one vitamin D or vitamin D analogue and at least one corticosteroid.

According to a further aspect of the present invention there is provided a method of making the composition as described herein comprising the following steps:

- preparing a first polyaphron dispersion comprising a vitamin D or vitamin D analogue;
- preparing a second polyaphron dispersion comprising a corticosteroid;
- and mixing together said first and second polyaphron dispersions to form the composition.

- 6 -

Advantageously, the compositions of the present invention have an enhanced dermal permeation of the active agent compared to known compositions, such as those disclosed in US-B-6,753,013. Thus, lower levels of active agents may be required in the compositions of the present invention in order to achieve beneficial treatment results. As a result of having lower levels of vitamin D and/or vitamin D analogues in the compositions of the present invention the likelihood of causing skin irritation and/or other side-effects is reduced. This may have the effect of increasing patient compliance with the dosage regime. Advantageously, in the compositions of the present invention the vitamin D and/or vitamin D analogue and the corticosteroid can co-exist in an aqueous composition, having the appropriate and controllable viscosity due to the presence of polyaphron dispersions together with any gelling agents included in the composition. A further particular advantage is that the compositions have good long term stability even at elevated temperature (40°C). Furthermore, the presence of water in the compositions may be useful for dissolving water-soluble additives such as water-soluble preservatives, antioxidants, water-soluble permeation enhancers and the like.

A further advantage is that compositions of the present invention are typically manufactured at room temperature without the need to apply heat, making it less likely that actives will be damaged in the composition.

A further advantage of the present composition is that it feels less greasy in use compared to, for example, the compositions of US-B-6,753,013, making it more pleasant to apply and thereby increasing the likelihood of proper patient compliance.

- 7 -

Advantageously the composition of the present invention has low antrophogenic potential (i.e. preferably use of this compound when applied to the skin causes less skin thinning compared to prior art compositions).

5 A yet further advantage of the composition of the present invention is that it need not comprise a high level of surfactant. In high concentrations surfactants are known to cause skin irritation. It is therefore desirable to keep the surfactant level to a minimum when applied to skin, and
10 in particular to damaged skin such as in the case of psoriasis. Preferably the compositions of the present invention comprise less than 4% by weight of surfactant, more preferably less than 3%, more preferably still less than 2% by weight of the total composition.

15 By polyaphron dispersion as used herein is meant a particular kind of hydrophilic liquid-in-hydrophobic liquid or hydrophobic liquid-in-hydrophilic liquid dispersion comprising (a) a hydrophilic liquid miscible phase, (b) a second hydrophobic phase being immiscible or substantially
20 immiscible with the first phase and (c) one or more surfactants, wherein the dispersed or discontinuous phase is in the form of small (e.g. micron to sub-micron diameter, but more usually at least 1 micron diameter) droplets, and the whole having the following characteristics, which
25 distinguish polyaphron dispersions from conventional or common emulsions and other dispersion types:

1. They are capable of existing in a stable form wherein the volume fraction of the dispersed phase (ϕ_{ip}) is greater than 0.7 and can be as high as 0.97. (ϕ_{ip} is the
30 volume ratio of discontinuous to continuous phase expressed as a fraction).

- 8 -

2. The microscopic appearance of polyaphron dispersions where ϕ_{ip} is greater than 0.7 is that of an aggregate of individual droplets, pushed closely together into polyhedral shapes, resembling the appearance of a gas foam. In this form, the dispersion has gel-like properties and is referred to as a Gel Polyaphron Dispersion (GPD).

3. Stable polyaphron dispersions can be formed with a surfactant concentration less than 3% and more typically less than 2% by weight of the total composition.

4. Gel Polyaphron Dispersions (as described in 2 above) can be diluted to any extent by the addition of more continuous phase without the addition of more surfactant, when the gel-like properties disappear. Once ϕ_{ip} has been reduced to below 0.7, the individual droplets of internal phase become separated to take the form of spherical droplets, which remain stable and intact but which may nevertheless join together in loose associations and float to the top or sink to the bottom of the diluted dispersion (depending on the relative densities of the two phases). In this diluted form each droplet is referred to as a Colloidal Liquid Aphron (CLA). Simple shaking of the diluted dispersion instantly causes a homogeneous, stable dispersion of Colloidal Liquid Aphrons to re-form.

Each of the above characteristics and a combination of them clearly differentiate the polyaphron dispersions of the present invention from conventional emulsions and other dispersion types which do not have all of those characteristics. Polyaphron dispersions are disclosed in the following literature references by Sebba: "Biliquid Foams",

- 9 -

J. Colloid and Interface Science, 40 (1972) 468-474 and "The Behaviour of Minute Oil Droplets Encapsulated in a Water Film", Colloid Polymer Sciences, 257 (1979) 392-396, Hicks "Investigating the Generation, Characterisation, and
5 Structure of Biliquid Foams", PhD Thesis, University of Bristol, 2005, Crutchley "The Encapsulation of Oils and Oil Soluble Substances Within Polymer Films", PhD Thesis, The University of Leeds, 2006 and Lye and Stuckey, Colloid and Surfaces, 131 (1998) 119-136. Aphrons are also disclosed in
10 US-A-4,486,333 and WO 97/32559.

Polyaphron dispersions are sometimes referred to as 'Biliquid Foams', 'High Internal Phase Emulsions (HIPEs)', 'High Internal Phase Ratio Emulsions (HIPREs)' and 'Gel Emulsions'. In US 5,573,757 a composition comprising a
15 polyaphron dispersion is described as "a viscoelastic gel". All descriptions that refer to dispersions having the characteristics described above are polyaphron dispersions as used in the present invention.

By "topical application" is meant application to human
20 or animal, preferably to the skin, including for example the face, scalp, feet, limbs or trunk.

The present invention will now be further described. In the following passages different aspects of the invention are defined in more detail. Each aspect so defined may be
25 combined with any other aspect or aspects unless clearly indicated to the contrary. In particular any feature indicated as being preferred or advantageous may be combined with any other feature or features indicated as being preferred or advantageous.

30 As described above polyaphron dispersions comprise a continuous phase, a discontinuous phase and a surfactant. The discontinuous phase is preferably a substantially hydrophobic internal phase, commonly known as an oil

- 10 -

internal phase. Preferably, the discontinuous phase comprises a pharmaceutically acceptable oil phase.

Examples of oils which may be used in the present invention include almond oil, babassu oil, blackcurrant seed oil, borage oil, canola oil, castor oil, coconut oil, cod liver oil, corn oil, cottonseed oil, evening primrose oil, fish oil, grapeseed oil, mustard seed oil, oat oil, olive oil, palm kernel oil, palm oil, peanut oil, rapeseed oil, safflower oil, sesame oil, shark liver oil, squalane, soybean oil, sunflower oil, walnut oil, wheat germ oil, hydrogenated castor oil, hydrogenated coconut oil, hydrogenated cottonseed oil, hydrogenated palm oil, hydrogenated soybean oil, partially hydrogenated soybean oil, hydrogenated vegetable oil, isopropyl myristate, isopropyl isostearate, isopropyl palmitate, modified triglycerides, caprylic/capric glycerides, fractionated triglycerides, glyceryl tricaprinate, glyceryl tricaproate, glyceryl tricaprinate, glyceryl tricaprinate/caprinate, glyceryl tricaprinate/caprinate, glyceryl tricaprinate/caprinate/laurate, glyceryl tricaprinate/caprinate/linoleate, glyceryl tricaprinate/caprinate/stearate, glyceryl trilaurate, glyceryl trilinoleate, glyceryl trilinolenate, glyceryl trioleate, glyceryl triundecanoate, linoleic glycerides, saturated polyglycolized glycerides, synthetic medium chain triglyceride containing primarily C₈-C₁₂ fatty acid chains, medium chain triglycerides, long chain triglycerides, modified triglycerides, fractionated triglycerides, and mixtures thereof.

Suitably the discontinuous phase comprises monoglycerides, diglycerides, or triglycerides.

It will be understood that other suitable oils may be used in the present invention.

- 11 -

In a preferred embodiment, the discontinuous phase comprises or is caprylic/capric triglyceride and/or isopropyl myristate (IPM).

The discontinuous phase may, for example, confer an emollient, occlusive, moisturising, conditioning or other cosmetic or pharmaceutical benefit to the skin. It may also increase the viscosity of the composition and may confer solvency to the active or actives. It may contain materials providing a heating or cooling effect when applied to the skin (for example capsaicin or menthol).

The composition may comprise at least 1 % by weight of the discontinuous phase, more preferably at least 10% by weight, at least 25%, at least 50%, at least 80 % by weight of the discontinuous phase based on the weight of the total composition.

The compositions of the present invention may be non-aqueous, substantially non-aqueous or aqueous.

By the term "non-aqueous composition" as used herein is meant a composition which is effectively free of water and does not contain water that has been deliberately added. A "non-aqueous composition" as used herein preferably has less than 0.5% by weight of water based on the total weight of the composition, more preferably less than 0.2% by weight of water, most preferably less than 0.1% by weight of water based on the total weight of the composition.

By the term "substantially non-aqueous" as used herein is meant a composition comprising less than 5% by weight, more preferably less than 4.5% by weight, of water based on the total weight of the composition.

By the term "aqueous composition" is meant a composition comprising at least 5% by weight of water based on the total weight of the composition. Preferably, the aqueous composition comprises at least 10%, or at least 15%

- 12 -

by weight of water based on the total weight of the composition. The aqueous composition may comprise at least 35%, or at least 40% by weight based on the total weight of the composition. For aqueous compositions, preferably the percentage of water is from 5% to 90% by weight, and more preferably from 5% to 50% by weight and most preferably from 8% to 20% by weight based on the total weight of the composition.

In one embodiment of the present invention there is provided an aqueous composition suitable for topical application comprising a continuous phase and at least one discontinuous phase, said composition comprising at least one polyaphron dispersion, at least one vitamin D or vitamin D analogue and at least one corticosteroid. In this embodiment preferably the composition comprises at least 5%, more preferably at least 6%, more preferably still at least 8% by weight of water based on the total weight of the composition.

Preferably, the aqueous composition suitable for topical application comprises from 60 to 95% by weight of discontinuous phase, and from 5 to 40 % by weight of continuous phase based on the total weight of the composition, wherein preferably the composition comprises from 5 to 40%, or from 5 to 30%, or from 8 to 20% by weight of water based on the total weight of the composition, wherein discontinuous phase and/or the continuous phase comprise at least one the vitamin D or vitamin D analogue and at least one corticosteroid and at least one surfactant. More preferably the aqueous composition suitable for topical application comprises from 70 to 90% by weight of discontinuous phase, and from 10 to 30 % weight of continuous phase based on the total weight of the composition, wherein preferably the composition

- 13 -

comprises at least from 5 to 30%, or from 8 to 20% by weight of water based on the total weight of the composition, wherein discontinuous phase and/or the continuous phase comprise at least one the vitamin D or vitamin D analogue
5 and at least one corticosteroid and at least one surfactant.

Preferably the composition of the present invention is dispersible in water. Preferably the composition of the present invention is dilutable in water. This increases the flexibility of use of the invention, for example in
10 improving the application of the composition to the scalp through hair by leaving the hair wet, or from rinsing the preparation from any topical surface should the desire or need arise, or by the easy removal by rinsing of product from accidental contamination of clothing. These advantages
15 improve the in-use experience of users and improve patient compliance.

The advantages of providing an aqueous composition comprising at least one vitamin D or vitamin D analogue and at least one corticosteroid include improved drug
20 permeation, control of drug permeation by the inclusion of certain permeation enhancers and permeation retarders and improved skin feel.

When the continuous or external phase of the composition comprises water, close control of the pH within
25 suitable limits is desired. Such control is desirable in order to obtain a stable composition. The present inventors have found that stable compositions of the present invention may be obtained when the pH of the composition is adjusted to a pH of from 7.0 to 8.5 , more preferably still to a pH
30 of from 7.25 to 7.75. It will be understood that any suitable acid or base may be used to adjust the pH to the appropriate value or pH range. Typically the pH of the composition will need to be raised by the addition of a

- 14 -

base, which suitably may be triethanolamine. Other suitable bases include, but are not limited to trishydroxymethylaminomethane (tris) and sodium hydroxide. Advantageously and preferably, the pH of the composition may be stabilized by the incorporation of a suitable buffer into the aqueous phase. Suitable buffer systems having a pH within the specified range will be familiar to those skilled in the art.

The continuous phase may comprise or consist essentially of a pharmaceutically acceptable liquid that is miscible or substantially miscible with water, preferably a compound of formula $R_1\text{-OH}$ where R_1 is $C_1\text{-}C_{10}$ alkyl and/or a compound of formula $\text{HO-}R_2\text{-H}$ where R_2 is $-(C_2H_4)_n$ or $-(C_3H_6)_n$ where n is 1 to 100, preferably 1 to 25. R_1 and R_2 may be linear or branched. Preferably R_1 is $C_1\text{-}C_4$ alkyl. n is preferably 1 to 25. Preferably the continuous phase comprises propylene glycol, polyethylene glycol, glycerol, ethanol, isopropyl alcohol, or a mixture thereof. Where the continuous phase comprises polyethylene glycol or polypropylene glycol, the polyethylene or polypropylene glycol is preferably a polyethylene glycol which is liquid at room temperature (20°C). The polyethylene glycol may, for example, contain from 1 to 12 ethylene or propylene oxide units and/or have a molecular weight of up to 600.

In one embodiment of the present invention the composition, and preferably the polyaphron dispersion, comprises from 0 to 60 wt%, preferably from 0 to 20 wt%, more preferably from 0 to 15 wt%, of a $C_1\text{-}C_4$ alcohol, ethylene glycol, a liquid polyethylene glycol, propylene glycol, a liquid polypropylene glycol, diethylene glycol mono ethyl ether or mixtures thereof. One of the advantages of the composition of the present invention over prior art compositions is that high levels of alcohol are not needed

- 15 -

as skin permeation enhancers. Alcohol is known to cause skin irritation when applied to damaged skin. The compositions of the present invention have the advantage that enhanced permeation can be achieved even without the need for alcohol to be present, or without the need for high levels of alcohol. The present inventors have surprisingly found that it is advantageous for the composition of the present invention to comprise from 0 to 25% by weight of alcohol, more preferably from 0 to 15% by weight based on the total weight of the composition. Preferably the alcohol is isopropanol and/or propylene glycol. Such compositions have improved drug delivery properties.

It will be understood that other suitable hydrophilic solvents may be used in the continuous phase of the polyaphrons.

The surfactant used in the present invention may be incorporated into either or both phases of the polyaphron dispersion. Suitable surfactants include an alkyl polyglycol ether, an alkyl polyglycol ester, an ethoxylated alcohol, a polyoxyethylene sorbitan fatty acid ester, a polyoxyethylene fatty acid ester, an ionic or non-ionic surfactant, a hydrogenated castor oil/polyoxyethylene glycol adduct containing from 25 to 60 ethoxy groups a castor oil/polyoxyethylene glycol adduct containing from 25 to 45 ethoxy groups, a sorbitan fatty acid ester (for example Span 20 or Span 80), a block copolymer of ethylene oxide and propylene oxide (for example Pluronic L121 or Pluronic F68), or a mixture thereof.

It will be understood that other suitable surfactants may be used.

Preferably the compositions of the present invention comprise less than 4% by weight of surfactant, more

- 16 -

preferably less than 3%, more preferably still less than 2% by weight of the total composition.

The composition of the present invention may comprise at least one vitamin D or vitamin D analogue predominantly in the continuous phase, or predominantly in the discontinuous phase. Most preferably at least one vitamin D or vitamin D analogue is present predominantly in the discontinuous phase.

The vitamin D analogue employed in the composition of the present invention may, for example, be calcipotriol, seocalcitol, calcitriol, calcipotriol monohydrate, tacalcitol, maxacalcitol, paricalcitol, falecalcitriol, becocalcidiol, $1\alpha,24S$ -dihydroxy-vitamin D₂, 1(S),3(R)-dihydroxy-20(R)-[[(3-(2-hydroxy-2-propyl)-phenyl)-methoxy]-methyl]-9,10-seco-pregna-5(Z),7(E),10(19)-triene, or a mixture thereof. More preferably, the vitamin D analogue is calcipotriol, calcitriol, tacalcitol, maxacalcitol, $1\alpha,24S$ -dihydroxy-vitamin D₂, 1(S),3(R)-dihydroxy-20(R)-[[(3-(2-hydroxy-2-propyl)-phenyl)-methoxy]-methyl]-9,10-seco-pregna-5(Z),7(E),10(19)-triene, or a mixture thereof. Most preferably, the vitamin D analogues are calcipotriol and calcipotriol monohydrate. Other examples of suitable vitamin D analogues are described in US-B-6,753,013.

Synthetic vitamin D analogues are preferred in the compositions of the present invention over naturally occurring vitamin D or vitamin D derivatives, since the therapeutic effects of the latter may be less selective for the treatment of skin diseases, such as psoriasis.

The composition of the present invention may comprise from 0.0001 to 0.05% by weight of vitamin D or vitamin D analogue, preferably from 0.001 to 0.01% by weight and more preferably from 0.0025 to 0.005% by weight of the total composition.

- 17 -

The corticosteroid may be predominantly in the continuous phase, or predominantly in the discontinuous phase. Preferably, the corticosteroid is predominantly in the discontinuous phase. More preferably, both the corticosteroid and the vitamin D or a vitamin D analogue are predominantly in the discontinuous phase. The range of weight ratios of corticosteroid to vitamin D or vitamin D analogue is preferably 4:1 to 50:1 and more preferably 8:1 to 20:1 and most preferably 9:1 to 11:1.

Preferably, the corticosteroid is selected from one or more of Betamethasone (9 -fluoro-11 ,17 ,21-trihydroxy-16 -methylpregna-1,4-diene-3,20-dione) or esters thereof such as the 21-acetate, 17-adamantoate, 17-benzoate, 17-valerate, and 17,21-dipropionate; Alclomethasone or esters thereof such as the 17,21-dipropionate; Clobetasol or esters thereof such as the propionate; Clobetasone or esters thereof such as the 17-butyrate; Desoximetasone; Diflucortolon or esters thereof, Diaflorasone or esters thereof such as the 17,21-diacetate; Fluocinonid; Flumetasone or esters thereof such as the 21-pivalate; Fluocinolone or ethers thereof such as the acetonide; Fluticasone or ester thereof such as the 17-propionate; Fluprednidene or esters thereof such as the 21-acetate; Halcinonide; Hydrocortisone or esters thereof such as the 17-butyrate; Mometasone or esters thereof such as the 17-(2-furoate); and Triamcinolon or ethers or esters thereof such as the acetonide. Nitric oxide donating corticosteroids may also be selected, such as TPI 1020 (NicOx). More preferably the corticosteroid is selected from one or more of Betamethasone or esters thereof such as the 17-valerate or the 17,21-dipropionate; Clobetasol or

- 18 -

esters thereof such as the propionate; Triamcinolon or ethers or esters thereof such as the acetonide or the acetonide-21-N-benzoyl-2-methyl- -alaninate or the acetonide-21-(3,3-dimethylbutyrate); or Hydrocortisone or
5 esters thereof such as the 17-butyrate. Most preferably, the corticosteroid is betamethasone 17,21-dipropionate.

The composition of the present invention preferably comprises from 0.001 to 1.0% by weight of corticosteroid, more preferably from 0.01% to 0.075% by weight and more
10 preferably still from 0.025 to 0.05% by weight of the total composition.

The composition of the present invention may further comprise a gelling agent and/or a rheology modifying agent, such as a viscosity modifier.

15 The gelling agent may, for example, be selected from alginate gums or their salts, guar gum, locust bean gum, xanthan gum, gum acacia, gelatin, hydroxymethyl-cellulose hydroxyethylcellulose, hydroxypropyl-cellulose, carboxymethylcellulose or its salts, bentonites, magnesium
20 aluminium silicates, "Carbomers" (salts of cross-linked polymers of acrylic acid), or glyceryl polymethacrylates or their dispersions in glycols. It will be understood that other suitable gelling agents may be used. Additionally, the inventors have discovered that some of the gelling
25 agents (for example, carbomers) may also function as a chemical buffering agents thus preventing unwanted variation in the pH of the composition during storage and use.

Preferably, the composition of the present invention comprises from 0.05 to 5.0% by weight of a gelling agent,
30 preferably from 0.1 to 2.0% by weight and more preferably from 0.2 to 1.0% by weight of the total composition.

- 19 -

In one embodiment of the present invention the composition has the consistency of a gel. The gel like consistency may be formed from one, two, three or more polyaphron dispersions. Preferably, the composition is an aqueous composition.

The composition of the present invention may be used in a method of treatment of the human or animal body by therapy. Further, the composition of the present invention may be used in the treatment of psoriasis. Also the composition of the present invention may be used in the manufacture of a medicament for treatment of psoriasis.

In one embodiment of the present invention, the composition as described herein may be applied to the scalp or other skin surface through hair. Preferably in this embodiment the hair is wetted (for example by use of water with or without shampoo, and then towel dried). The product may then be applied to the scalp in a suitable amount and then massaged into the scalp through the hair. The hair may then be left to dry naturally or dried using a hair dryer. Advantageously, the water-dispersible form of the formulation enables an even distribution of the actives on the skin using this process. Alternatively, or additionally, the composition may be massaged into the scalp through dry hair and left for a suitable period (which may be 8 to 12 hours) after which the excess or reminder may be rinsed out with water with or without shampoo. Preferably the composition is applied to an animal in unit dosage form.

The compositions of the present invention may also contain other additives such as preservatives (for instance to prevent microbiological spoilage), buffering agents (for the control of pH and to avoid instability and damage to the skin's acid mantle), antioxidants and permeation enhancers.

- 20 -

These additives may be included in the continuous or the discontinuous phase of the polyaphron dispersion.

It will be understood that the inclusion of these additives will be at the levels and with the type of materials which are found to be effective and useful. Care needs to be taken in the choice and amount of these additives to prevent compromise to the other performance advantages of the present invention.

In one embodiment of the present invention, the vitamin D or vitamin D analogue of the present invention is dispersed and/or dissolved in the discontinuous phase of a first polyaphron dispersion. The corticosteroid is dispersed and/or dissolved in the discontinuous phase of a second polyaphron dispersion. The first and the second polyaphron dispersions are then mixed together to form the composition of the present invention. Optionally, a third or further polyaphron dispersion may also be present in the composition of the present invention. The third or further polyaphron dispersion may, for example, comprise agents such as emollient oils (to improve in use 'feel'), occlusive oils to prevent skin dehydration and to enhance skin permeation by the actives, agents that provide a heating or cooling sensation when applied to the skin or sunscreens. Preferably these agents will be present in the discontinuous phase of the polyaphron dispersions.

In one embodiment of the present invention, the vitamin D analogue is calcipotriol or calcipotriol monohydrate.

In a particularly preferred composition the discontinuous phase is a caprylic/capric triglyceride, the continuous phase is demineralised water, the vitamin D analogue is calcipotriol and the corticosteroid is betamethasone dipropionate.

- 21 -

In one embodiment of the present invention, the composition comprises at least one discontinuous phase which comprises at least one of capric/caprylic triglycerides, squalane, dimethicone, cyclomethicone and mixtures of two or more thereof, the vitamin D analogue is calcipotriol and the corticosteroid is betamethasone dipropionate. Preferred surfactants in this embodiment are Laureth-4 and Poloxamer 188. Preferably the continuous phase comprises water. Such an embodiment is exemplified by Example 1. Advantageously such a composition is stable over an extended period of time, for example over 3 months, 6 months or 9 months. This composition provides good solubility for the actives, has an enhanced permeation and has a very pleasant feel on the skin during and after application.

In a further embodiment, the composition of the present invention comprises at least one discontinuous phase which comprises at least one of capric/caprylic triglycerides, squalane, dimethicone, isopropyl myristate, cyclomethicone and mixtures of two or more thereof, the vitamin D analogue is calcipotriol and the corticosteroid is betamethasone dipropionate, preferred surfactants in this embodiment are Laureth-4 and Poloxamer 188. Preferably the continuous phase comprises water and polyacrylic acid which is neutralized with triethanolamine, so as to form a gel to control viscosity. Such an embodiment is exemplified by Example 3. Advantageously this composition uses an aqueous gel phase to provide a greater cream-like viscosity and excellent pH buffering capacity.

In a further embodiment, the composition of the present invention comprises at least one discontinuous phase which comprises at least one of capric/caprylic triglycerides, squalane, dimethicone, isopropyl myristate and cyclomethicone and mixtures of two or more thereof, the

- 22 -

vitamin D analogue is calcipotriol and the corticosteroid is betamethasone dipropionate, preferred surfactants in this embodiment are Laureth-4 and Poloxamer 188. Preferably the continuous phase comprises water and xanthan gum, so as to
5 form a gel to control viscosity. Such an embodiment is exemplified by Example 5. The inventors have found that the use of xanthane produces excellent stability at a lower viscosity which is of particular use as a lotion rather than a cream.

10 In a further embodiment, the composition of the present invention comprises at least one discontinuous phase which comprises at least one of capric/caprylic triglycerides, squalane, dimethicone, silicone elastomer, isopropyl myristate and cyclomethicone and mixtures of two or more
15 thereof, the vitamin D analogue is calcipotriol and the corticosteroid is betamethasone dipropionate, preferred surfactants in this embodiment are Laureth-4 and Poloxamer 188. Preferably the continuous phase comprises water, isopropyl alcohol and polyacrylic acid which is neutralized
20 with triethanolamine, so as to form a gel to control viscosity. Such an embodiment is exemplified by Example 13. The silicone elastomer is added to the discontinuous phase to improve the cosmetic feel of the product. This embodiment provides still further pleasant 'after-feel' (the
25 residual feel of the skin after application) and increased skin barrier properties.

In yet a further embodiment, the composition of the present invention comprises a discontinuous phase which comprises isopropyl myristate and at least one of squalane,
30 dimethicone, silicone elastomer, and cyclomethicone. In this embodiment the corticosteroid is betamethasone dipropionate and the vitamin D analogue is calcipotriol and the preferred surfactants are Laureth-4 and Poloxamer 188. Preferably the

- 23 -

continuous phase comprises water and polyacrylic acid which is neutralized with triethanolamine, so as to form a gel to control viscosity. Such an embodiment is exemplified by Example 17. This embodiment provides greatly enhanced permeation compared to other embodiments.

Accordingly to one aspect of the present invention, there is provided a method of making the composition as described herein comprising the following steps:

- (i) providing a hydrophilic solvent, optionally comprising at least one vitamin D or vitamin D analogue, and/or at least one corticosteroid, and/or a surfactant;
- (ii) providing a hydrophobic solvent optionally comprising at least one vitamin D or vitamin D analogue, and/or at least one corticosteroid, and/or a surfactant;
- (iii) mixing the hydrophilic with the hydrophobic solvent under suitable conditions to form the composition comprising at least one polyaphron dispersion, at least one vitamin D or vitamin D analogue and at least one corticosteroid.

Suitable methods for preparing polyaphron dispersions are described in US-A-4486333. It will be understood by those skilled in the art that other manufacturing methods may be used, as appropriate.

Accordingly to another aspect of the present invention, there is provided a method of making the composition as described herein comprising the following steps:

- preparing a first polyaphron dispersion comprising a vitamin D or vitamin D analogue;
- preparing a second polyaphron dispersion comprising a corticosteroid;

- 24 -

and mixing together said first and second polyaphron dispersions to form the composition.

The method may further comprise:

preparing a third or further polyaphron dispersion
5 comprising an active agent, such as one or more of a
sunscreening agent, a cooling agent, a warming agent, an
antipruritic agent, an aesthetic agent, a cosmetic masking
agent, a foaming agent, a fragrance, a colouring agent, an
antioxidant or an emollient oil and including mixtures
10 thereof;

and mixing said third or further polyaphron with
said first and second polyaphron dispersions to form the
composition. The composition as described herein may be
delivered to the skin in the form of an aerosol or a spray.
15 For example EP 1,575,542 teaches the incorporation of a
biliquid foam into aerosols.

The composition of the present invention may be
incorporated into aerosols by the addition of water and
propellant gas (for example butane).

20 The composition of the present invention may be in the
form of, for example, a lotion or a cream. The composition
may be stored in any suitable jar, tube, bottle, sachet,
aerosol, spray applicator or pump action sealed container.
Advantageously in order for the compositions to remain
25 stable the containers preferably prevent oxygen from
entering the container. Preferably the compositions of the
present invention are sealed in air tight containers in
order to prevent degradation of the compositions prior to
use.

30 The present invention will now be described further, by
way of example only, with reference to the following
figures, in which:

- 25 -

Figure 1 is a HPLC chromatogram of a stable sample (Example 7). Peaks for calcipotriol (3.9 min) and betamethasone dipropionate (BDP) (5.3 min). No evidence of degradation products is observed. HPLC conditions are:

5 Column:NovaPakC18, 4 μ particle size, 3.9 x 150mm column (Waters), Mobile phase: 55% acetonitrile in water. Flow rate: 1ml/minute. Column Temperature: 25°C.

Figure 2 is a HPLC chromatogram of an unstable sample.

10 Degradation of betamethasone dipropionate is observed (extra peak and shoulder at 2.5 minutes) and there is some evidence of calcipotriol degradation (very minor peak at 3.3 min). HPLC conditions are: Column:NovaPakC18, 4 μ particle size, 3.9 x 150mm column (Waters), Mobile phase: 55% acetonitrile
15 in water. Flow rate: 1ml/minute. Column Temperature: 25°C.

Figure 3 is a HPLC chromatogram of the degradation of calcipotriol as evidenced by the presence of extraneous peaks (4.5 min and 3.6 min) either side of the main
20 calcipotriol peak (3.9 min). No betamethasone dipropionate (BDP) was present in this sample, thus these peaks cannot be due to BDP degradation. HPLC conditions are:

Column:NovaPakC18, 4 μ particle size, 3.9 x 150mm column (Waters), Mobile phase: 55% acetonitrile in water. Flow
25 rate: 1ml/minute. Column Temperature: 25°C.

Definition of stability.

For the present invention, a product is considered to be storage stable if it meets with the following criteria.

30

The product is stored in closed, airtight glass containers with headspace comprising no more than 5% by volume of the total usable volume of the container.

- 26 -

The product and the container as defined above are stored at a constant temperature of 40°C in a standard laboratory oven (for example, Heraeus 'Function Line' air circulating oven model UT6, temperature control $\pm 0.3^{\circ}\text{C}$ at 150°C).

The product is examined at the end of the examination period. The examination period is at least 3 months and preferably at least 6 months from the start date of storage.

10

The pass criteria are as follows:

Procedure	Pass Criteria
Visual comparison of the appearance of the stored sample compared to a standard sample stored at 20°C for the same period.	Visual assessment indicates very little, if any, difference between stored sample and standard. In particular, appearance of the sample is uniform throughout with no sign of separation into two or more distinct phases.
A microscopic examination at a magnification of at least 200X comparing the microscopic appearance of the stored sample with a stored image of its appearance at the date of commencement of storage	Examination indicates very little, if any change in the size and size distribution of polyaphron droplets, with no sign of separated phases.
Analysis of the pharmacologically active components of the formulation by the extraction and HPLC method given hereinunder	Each active shall not have diminished by more than 5% by weight of the original content at the date of commencement of the storage test after 3 months of storage. Known decomposition products of the actives, if any such are present, collectively constitute no more than 5% of the original active based upon area under the curve measurements. See

- 27 -

	Figures 1-3 for further clarification.
--	--

Any stored samples that meet the above criteria under the test conditions given above are considered to be storage stable for the purposes of this invention.

5

The FDA's 'Guidance for Industry Q1A (R2) Stability Testing of New Drug Substance and Products' although non-binding, specifies accelerated storage conditions that include storage at specific temperatures (e.g. 40°C) for a specific
10 time (6 months) and at a controlled relative humidity (75% RH). The European Agency for the Evaluation of Medicinal Products, ICH Q1A (R2) 'Stability Testing Guidelines: Stability Testing of New Drug Substances and Products' specifies identical conditions for accelerated storage
15 testing. The stability test method specified above for the purpose of this invention does not include provision for the control of relative humidity since storage takes place in closed glass containers whose walls and closures are impervious to the passage of water vapour.

20

The above definition of storage stable is definitive for the purposes of this invention. However, storage data on some examples were obtained by substituting close-sealed eppendorf tubes for closed glass containers (glass). This
25 method was adopted to cope with relatively small amounts of sample and gives a reasonable approximation to the definitive method since eppendorf tubes also provide an impervious barrier and seal to water vapour and other components of the composition. The examination procedure
30 for eppendorf tubes (plastic) is identical to that of the definitive method.

- 28 -

The following Examples further illustrate the present invention.

EXAMPLE 1

- 5 Three gel polyaphron dispersions of the following compositions were prepared by the following method.

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.0695%) in Caprylic/Capric Triglyceride (Mygliol 812 - Condea)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone Dipropionate solution** (0.3465%) in Caprylic/Capric Triglyceride (Mygliol 812 - Condea)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

10

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	9.00
Dimethicone (Q7-1920, 100 CP - Dow Corning)	44.10
Cyclomethicone (STb5NF - Dow Corning)	36.00
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

FINAL PRODUCT	
GEL POLYAPHRON DISPERSION 1	7.20
GEL POLYAPHRON DISPERSION 2	18.60
GEL POLYAPHRON DISPERSION 3	74.20

- 29 -

*The Calcipotriol content in the final formulation is 50µg/g.

5 **The betamethasone dipropionate content in the final formulation is 643µg/g (equivalent to 500µg/g of betamethasone).

Manufacturing Method

Three polyaphron dispersions were made individually by the following method:

10

A low form, 250ml laboratory beaker (internal diameter 6.5cm) was charged with sufficient aqueous (continuous) phase to make 30 g of gel polyaphron. This was stirred at 200 rpm with a four-bladed impeller having a diameter of 6.0
15 cm whilst adding the oil (discontinuous) phase drop wise from a Pasteur pipette. The rate of addition at the start of the process was slow (approximately one drop every 7 seconds) but was speeded up once 10% of the oil phase had been added so that the total time to make the gel polyaphron
20 was approximately 20 minutes.

Prior to the manufacture of each gel polyaphron dispersion, any active was dissolved in the appropriate phase by gentle stirring overnight with a magnetic stirrer at room
25 temperature in a covered beaker.

To form the final product, the three individual polyaphron dispersions were mixed together.

30

- 30 -

Stability Measurements- conditions 1a

Stability measurements made using the method outlined below.

5 In this example the storage data was obtained in an eppendorf tube.

The calcipotriol and betamethasone were extracted from the composition of Example 1 into isopropanol and assayed by HPLC under the conditions given below.

10

HPLC conditions:

Column: NovaPakC18, 4 μ particle size, 3.9 x 150mm column (Waters)

Mobile phase: 47% acetonitrile in water.

15 Flow rate: 1ml/minute.

Column Temperature: 25°C.

Retention time for calcipotriol was 6.8 minutes

Retention time for betamethasone was 9.9 minutes.

20 The inventors observed that after 2 months storage at 40°C, the levels of calcipotriol and betamethasone were 102% $\pm$ 3% and 99% $\pm$ 1% of the original levels, respectively.

Stability Measurements- conditions 1b

25 Stability measurements made using the method outlined below.

Samples were stored in (airtight) glass jars in a suitable oven maintained at 40°C, as described above.

The calcipotriol and betamethasone were extracted from the composition of Example 1 into isopropanol by mixing the

30 sample in isopropanol followed by sonication and centrifugation. Aliquots of the supernatant were assayed by HPLC under the conditions given below.

- 31 -

HPLC conditions:

Column: NovaPakC18, 4 μ particle size, 3.9 x 150mm column
(Waters)

5 Mobile phase: 55% acetonitrile in water.

Flow rate: 1ml/minute.

Column Temperature: 25°C.

Retention time for calcipotriol was 3.9 minutes

Retention time for betamethasone was 5.3 minutes.

10

The inventors observed that after 2 months storage at 40°C, the levels of calcipotriol and betamethasone were 102% $\pm$ 3% and 99% $\pm$ 1% of the original levels, respectively.

15

Example 2

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.06766%) in Caprylic/Capric Triglyceride (Mygliol 812 - Condea)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone Dipropionate solution** (0.3066%) in a 3:2 w/w blend of isopropyl myristate (Lexol IPM NF - Inolex): Caprylic/Capric Triglyceride (Mygliol 812 - Condea)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

20

- 32 -

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	9.00
Dimethicone (Q7-9120, 100 CP - Dow Corning)	44.10
Cyclomethicone (STb5NF - Dow Corning)	36.00
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

AQUEOUS GEL	
	% w/w
Polyacrylic acid (Ultrez 10 - Noveon)	1.00
Triethanolamine (TEA)	to pH 7.50
Demineralised Water	qs to 100%

5

FINAL PRODUCT	
	% w/w
GEL POLYAPHRON DISPERSION 1	9.19
GEL POLYAPHRON DISPERSION 2	23.61
GEL POLYAPHRON DISPERSION 3	47.27
Propylene glycol	10.05
Aqueous gel	9.88

* final calcipotriol level 55.4 µg/g

** final BDP level 645 µg/g, equivalent to 502 µg/g

10 **Manufacturing Method**

The method used was as described for Example 1 above except that propylene glycol and the neutralised aqueous gel were added to the final mixture of the three gel polyaphron dispersions, 1, 2 and 3 and then mixed in by simple mixing until the product was a homogeneous mixture of the polyaphron dispersions.

By neutralised gel is meant the addition of triethanolamine (a base) to a dispersion of the polyacrylic acid to form a clear gel having a pH value of 7.5 ±0.2. The process of

- 33 -

neutralisation of polyacrylic acid gels is well known to those skilled in the art.

Stability Measurements

5 Stability was tested as in Example 1b.

The inventors observed that after 3.5 months storage at 40°C in glass jars the levels of calcipotriol and betamethasone dipropionate were $104 \pm 12\%$ and $95 \pm 2\%$ of the original
10 levels respectively.

Stability testing is ongoing with this sample.

15 Reference is made to Simonsen et al, Drug Development and Industrial Pharmacy, 30(10) (2004) 1095-1102, wherein the authors state that the inclusion of propylene glycol to a product containing both calcipotriol and betamethasone causes rapid decomposition of one or other of the actives
20 depending upon pH, although propylene glycol is a very good flux rate enhancer. They concluded that it was not possible to include propylene glycol in their formulated product. This Example demonstrates that the inventors of the present invention have overcome this problem. Example 2, table 2 in
25 US-B-6,753,013 illustrates the degradation of calcipotriol in a product comprising calcipotriol, betamethasone dipropionate and propylene glycol after storage for 2.5 months at 40°C.

- 34 -

EXAMPLE 3

Three gel polyaphron dispersions and an aqueous gel of the following compositions were prepared by the following method.

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.0573%) in a 3:2 w/w blend of isopropyl myristate (Lexol IPM NF - Inolex): Caprylic/Capric Triglyceride (Mygliol 812 - Condea)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

10

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone Dipropionate solution** (0.3838%) in Caprylic/Capric Triglyceride (Mygliol 812 - Condea)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	9.00
Dimethicone (Q7-9120, 100 CP - Dow Corning)	44.10
Cyclomethicone (STb5NF - Dow Corning)	36.00
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

AQUEOUS GEL	
	% w/w
Polyacrylic acid (Ultrez 10 - Noveon)	1.00

- 35 -

Triethanolamine (TEA)	to pH 7.50
Demineralised Water	qs to 100%

FINAL PRODUCT	% w/w
GEL POLYAPHRON DISPERSION 1	12.95
GEL POLYAPHRON DISPERSION 2	20.18
GEL POLYAPHRON DISPERSION 3	57.05
AQUEOUS GEL	9.82

*The Calcipotriol content in the final formulation is 66.1µg/g.

5 **The betamethasone dipropionate content in the final formulation is 690µg/g (equivalent to 537µg/g of betamethasone).

Manufacturing Method

10 The method used was as described for Example 1 above except that the neutralised aqueous gel was added to the final mixture of the three gel polyaphron dispersions, 1, 2 and 3 and then mixed in by simple mixing until the product was a homogeneous mixture of the polyaphron dispersions.

Stability Measurements

15 Stability was tested as in Example 1b.

20 The inventors observed that after 9 months storage at 40°C in a sealed glass jar the levels of calcipotriol and betamethasone dipropionate were $97 \pm 7\%$ and $105 \pm 6\%$ of the original levels respectively.

- 36 -

EXAMPLE 4

- 5 Three gel polyaphron dispersions of the following compositions were prepared by the following method.

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.0237%) isopropyl myristate (Lexol IPM NF - Inolex)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Sucrose (Fisher)	1.00
Demineralised Water	8.00

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone Dipropionate solution** (0.3085%) in isopropyl myristate (Lexol IPM NF - Inolex)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Sucrose (Fisher)	1.00
Demineralised Water	8.00

10

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	20.00
Dimethicone (Q7-9120, 20 CP - Dow Corning)	35.00
Cyclomethicone (STb5NF - Dow Corning)	29.10
Elastomer DC10 (Dow Corning)	5.00
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Sucrose (Fisher)	1.00
Demineralised Water	8.00

- 37 -

FINAL PRODUCT	% w/w
GEL POLYAPHRON DISPERSION 1	27.99
GEL POLYAPHRON DISPERSION 2	26.56
GEL POLYAPHRON DISPERSION 3	45.45

*The Calcipotriol content in the final formulation is 59.0µg/g.

5 **The betamethasone dipropionate content in the final formulation is 730µg/g (equivalent to 568µg/g of betamethasone).

Manufacturing Method

The method used was as described for Example 1.

10

Stability Measurements

Stability was tested as in Example 1b

15 The inventors observed that after 4.5 months storage at 40°C in a sealed glass jar the levels of calcipotriol and betamethasone dipropionate were 100% and 119% of the original levels respectively.

20 After 6 months, the levels of calcipotriol and betamethasone dipropionate were 42% and 65% of the original levels respectively.

Stability testing is ongoing with this sample at room temperature.

25

30

- 38 -

5

EXAMPLE 5

Three gel polyaphron dispersions and an aqueous gel of the following compositions were prepared by the following method.

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.0452%) in isopropyl myristate (Lexol IPM NF - Inolex)	89.10
Polyoxyethylene (20) sorbitan monooleate (Surfacare T80)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

10

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone Dipropionate solution** (0.2889%) in isopropyl myristate (Lexol IPM NF - Inolex)	89.10
Polyoxyethylene (20) sorbitan monooleate (Surfacare T80)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	9.00
Dimethicone (Q7-9120, 100 CP - Dow Corning)	44.10
Cyclomethicone (STb5NF - Dow Corning)	36.00
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

- 39 -

AQUEOUS GEL	
	% w/w
Xanthan gum (Thew Arnott)	0.50
NaH ₂ PO ₄ (as pH 7.50 phosphate buffer, 50 mM)	0.60
Demineralised Water	qs to 100%

FINAL PRODUCT	
	% w/w
GEL POLYAPHRON DISPERSION 1	11.66
GEL POLYAPHRON DISPERSION 2	27.54
GEL POLYAPHRON DISPERSION 3	50.87
AQUEOUS GEL	9.93

*The Calcipotriol content in the final formulation is 47.0µg/g.

- 5 **The betamethasone dipropionate content in the final formulation is 709µg/g (equivalent to 551µg/g of betamethasone).

10 **Manufacturing Method**

The method used was precisely as described for Example 1 above except that the neutralised aqueous gel was added to the final mixture of the three gel polyaphron dispersions, 1, 2 and 3 and then mixed in by simple mixing until the product was a homogeneous mixture of the polyaphron dispersions.

EXAMPLE 6

20 Three gel polyaphron dispersions and an aqueous gel of the following compositions were prepared by the following method.

Gel POLYAPHRON DISPERSION 1	
Oil Phase	
Calcipotriol solution* (0.02687%) in isopropyl myristate (Lexol IPM NF - Inolex)	89.00
Laureth-4 (Volpo L4 - Croda)	1.00
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Propan-2-ol	2.70
Demineralised Water	5.40

- 40 -

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone Dipropionate solution** (0.3289%) in isopropyl myristate (Lexol IPM NF - Inolex)	89.00
Laureth-4 (Volpo L4 - Croda)	1.00
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Propan-2-ol (Fisher)	2.70
Demineralised Water	5.40

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	9.00
Dimethicone (Q7-9120, 100 CP - Dow Corning)	44.10
Cyclomethicone (STb5NF - Dow Corning)	36.00
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

5

AQUEOUS GEL	
	% w/w
Polyacrylic acid (Ultrez 10, Noveon)	2.00
Propan-2-ol (Fisher)	30.00
50% triethanolamine (aq)	to pH 7.29
Demineralised Water	qs to 100%

FINAL PRODUCT	
GEL POLYAPHRON DISPERSION 1	% w/w
GEL POLYAPHRON DISPERSION 1	25.00
GEL POLYAPHRON DISPERSION 2	25.00
GEL POLYAPHRON DISPERSION 3	40.00
AQUEOUS GEL	10.00

*The Calcipotriol content in the final formulation is 67.9µg/g.

10 **The betamethasone dipropionate content in the final formulation is 783µg/g (equivalent to 609µg/g of betamethasone).

- 41 -

Manufacturing Method

The method used was as described for Example 1 above except that the neutralised aqueous gel was added to the final mixture of the three gel polyaphron dispersions, 1, 2 and 3 and then mixed in by simple mixing until the product was a homogeneous mixture of the polyaphron dispersions.

Stability Measurements

Stability was tested as in Example 1a, the sample was stored in close-sealed eppendorf tubes.

The inventors observed that after 3 months storage at 40°C the levels of calcipotriol and betamethasone were 101% and 106% of the original levels respectively. Stability testing is ongoing with this sample.

Furthermore, after 6 months storage at 40°C the levels of calcipotriol and betamethasone were 107% and 100% of the original levels respectively.

EXAMPLE 7

Three gel polyaphron dispersions and an aqueous gel of the following compositions were prepared by the following method.

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.0258% in isopropyl myristate (Lexol IPM NF - Inolex)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

- 42 -

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone Dipropionate solution** (0.325%) in isopropyl myristate (Lexol IPM NF - Inolex)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	20.00
Dimethicone (Q7-9120, 20 cSt - Dow Corning)	35.00
Elastomer DC10 (Dow Corning)	5.00
Cyclomethicone (STb5NF - Dow Corning)	26.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

AQUEOUS GEL	
	% w/w
Polyacrylic acid (Carbomer ETD 2020, supplier)	2.00
Disodium ethylenediaminetetracetic acid (Na ₂ EDTA, Fisher)	0.20
Sodium bisulphite (NaHSO ₃ , Fisher)	0.50
NaOH 20% aqueous solution	To pH 7.71
Demineralised Water	qs to 100%

5

FINAL PRODUCT	
GEL POLYAPHRON DISPERSION 1	25.00
GEL POLYAPHRON DISPERSION 2	25.00
GEL POLYAPHRON DISPERSION 3	20.00
AQUEOUS GEL	30.00

*The Calcipotriol content in the final formulation is 57.4µg/g.

- 43 -

**The betamethasone dipropionate content in the final formulation is 723µg/g (equivalent to 562µg/g of betamethasone).

5

Manufacturing Method

The method used was as described for Example 1 above except that the neutralised aqueous gel was added to the final mixture of the three gel polyaphron dispersions, 1, 2 and 3
10 and then mixed in by simple mixing until the product was a homogeneous mixture of the polyaphron dispersions.

Stability Measurements

Stability was tested as in Example 1a, the sample was stored
15 in close-sealed eppendorf tubes.

The inventors observed that after 3 months storage at 40°C the levels of calcipotriol and betamethasone were 96% and 103% of the original levels respectively. Stability testing
20 is ongoing with this sample. Further stability testing after 6 months storage at 40°C indicated levels of calcipotriol and betamethasone were 96% and 103% of the original levels respectively.

- 44 -

Example 8

Commercial product (Dovobet (LEO Pharma), lot No EA5525).

- 5 Dovobet is a commercial product which is also sold under the name Daivobet and Taclonex.

Product	Ingredients (not in order of concentration)	Feel on Skin
Example 1 US 6,753,013 B1	1. Betamethasone dipropionate 2. Calcipotriol (Hydrate), 3. Polyoxypropylene-15-Stearyl Ether, 4. α -Tocopherol, 5. White Soft Paraffin	Very sticky and greasy
Dovobet (as known as Daivobet, Taclonex)	1. Betamethasone dipropionate 2. Calcipotriol (Hydrate), 3. Polyoxypropylene-15-Stearyl Ether, 4. α -Tocopherol, 5. White Soft Paraffin	Very sticky and greasy

10

A sample of the commercial product was stored in a sealed glass jar and stored at 40°C, in the same manner as the Examples above.

- 15 After 3 months, the sample was analysed using the extraction method as used for Example 1. Assay using HPLC (as above) indicated drug levels 91% and 70% for calcipotriol and BDP, respectively, compared with analyses performed on fresh sample.

- 45 -

EXAMPLE 9

Single gel polyaphron dispersion of the following
5 compositions was prepared by the following method.

Gel POLYAPHRON DISPERSION	
Oil Phase	%
Solution of Calcipotriol* (0.0133%) and BDP (0.1554%) in Caprylic/Capric Triglyceride (Mygliol 812 - Condea)	90.00
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
NaOH solution (20% w/w)	to pH 8.50
Demineralised Water	qs to 9.00

*The Calcipotriol content in the final formulation is 120µg/g.

10 **The betamethasone dipropionate content in the final formulation is 1399µg/g (equivalent to 1088µg/g of betamethasone).

Manufacturing Method

15 The polyaphron dispersion was made by the method outlined in Example 1.

Stability Measurements

Stability measurements made using the method outlined in Example 1b.

20

The inventors observed that after storage for 1 month at 40°C in a sealed glass jar, the formulation had separated into a system with a significant oil phase on top of the polyaphron gel. For the sample stored at ambient temperature
25 after 18 months the levels of calcipotriol and betamethasone were 97% ± 3% and 98% ± 3% of the original levels, respectively.

- 46 -

Example 10

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.0671%) in Caprylic/Capric Triglyceride (Mygliol 812 - Condea)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

5

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone Dipropionate solution** (0.3294%) in a 3:2 w/w blend of isopropyl myristate (Lexol IPM NF - Inolex): Caprylic/Capric Triglyceride (Mygliol 812 - Condea)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	9.00
Dimethicone (Q7-9120, 100 CP - Dow Corning)	44.10
Cyclomethicone (STb5NF - Dow Corning)	36.00
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

10

AQUEOUS GEL	
	% w/w
Polyacrylic acid (Ultrez 10 - Noveon)	1.00
Triethanolamine (TEA)	to pH 7.00
Demineralised Water	qs to 100%

- 47 -

FINAL PRODUCT	% w/w
GEL POLYAPHRON DISPERSION 1	8.89
GEL POLYAPHRON DISPERSION 2	23.35
GEL POLYAPHRON DISPERSION 3	57.81
Aqueous gel	9.95

* final calcipotriol level 53.2 µg/g

** final BDP level 692 µg/g (equivalent to 538 µg/g betamethasone)

5

Manufacturing Method

The method used was precisely as described for Example 1 above except that the neutralised aqueous gel was added to the final mixture of the three gel polyaphron dispersions, 1, 2 and 3 and then mixed in by simple mixing until the product was a homogeneous mixture of the polyaphron dispersions.

15 **Stability Measurements**

Stability was tested as in Example 1b

The inventors observed that after 2 months storage at 40°C in a sealed glass jar the levels of calcipotriol and betamethasone dipropionate were 103 ± 2% and 106 ± 3% of the original levels respectively. Furthermore, after 6 months storage at 40°C the levels of calcipotriol and betamethasone dipropionate were 106 ± 2% and 105 ± 3% of the original levels respectively. Stability testing is ongoing with this sample.

25

- 48 -

EXAMPLE 11

Three gel polyaphron dispersions and an aqueous gel of the following compositions were prepared by the following method.

5

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.0581%) in Caprylic/Capric Triglyceride (Mygliol 812 - Condea)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone Dipropionate solution** (0.2908%) in a 3:2 w/w blend of isopropyl myristate (Lexol IPM NF - Inolex): Caprylic/Capric Triglyceride (Mygliol 812 - Condea)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

10

Gel POLYAPHRON DISPERSION 3 64/59/8	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	9.00
Dimethicone (Q7-9120, 100 CP - Dow Corning)	44.10
Cyclomethicone (STb5NF - Dow Corning)	36.00
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

BUFFER	
	% w/w
	1.00
Phosphate buffer salts	to pH 7.50

- 49 -

Demineralised Water	qs to 100%
---------------------	------------

FINAL PRODUCT	% w/w
GEL POLYAPHRON DISPERSION 1	10.15
GEL POLYAPHRON DISPERSION 2	25.32
GEL POLYAPHRON DISPERSION 3	53.54
BUFFER	10.99

*The Calcipotriol content in the final formulation is 52.5µg/g.

5 **The betamethasone dipropionate content in the final formulation is 656µg/g (equivalent to 510µg/g of betamethasone).

Manufacturing Method

10 The method used was precisely as described for Example 1 above except that the buffer was added to the final mixture of the three gel polyaphron dispersions, 1, 2 and 3 and then mixed in by simple mixing until the product was a homogeneous mixture of the polyaphron dispersions.

Stability Measurements

15 Stability was tested as in Example 1b

20 The inventors observed that after 6 months storage at 40°C in a sealed glass jar the levels of calcipotriol and betamethasone were 106 ± 2% and 105 ± 1% of the original levels respectively.

EXAMPLE 12

25 Three gel polyaphron dispersions of the following compositions were prepared by the following method.

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.0283%) isopropyl myristate (Lexol IPM NF - Inolex)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90

- 50 -

Aqueous Phase	
Calcipotriol solution* (0.0400% in aq phase) 29.6% w/w isopropanol in demineralised water	9.00
Poloxamer 188 (Pluronic F68 - BASF)	1.00

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone dipropionate solution** (0.3289%) in isopropyl myristate(Lexol IPM NF - Inolex)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Betamethasone dipropionate solution** (0.2252%) in 33.8% w/w isopropanol in demineralised water	9.00
Poloxamer 188 (Pluronic F68 - BASF)	1.00

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	20.00
Dimethicone (Q7-9120, 20 CP - Dow Corning)	35.00
Cyclomethicone (STb5NF - Dow Corning)	29.10
Elastomer DC10 (Dow Corning)	5.00
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

5

AQUEOUS GEL	
	% w/w
Polyacrylic acid (Ultrez 10 - Noveon)	2.00
Triethanolamine (TEA)	to pH 7.29
Demineralised Water	qs to 100%

FINAL PRODUCT	% w/w
GEL POLYAPHRON DISPERSION 1	25.00
GEL POLYAPHRON DISPERSION 2	25.00
GEL POLYAPHRON DISPERSION 3	30.00
AQUEOUS GEL	20.00

*The Calcipotriol content in the final formulation is 72.0µg/g.

- 51 -

**The betamethasone dipropionate content in the final formulation is 782µg/g (equivalent to 608µg/g of betamethasone).

5 **Manufacturing Method**

The method used was precisely as described for Example 1, except for the inclusion of gel phase.

10 **Stability Measurements**

Stability was tested as in Example 1a.

The inventors observed that after 6 months storage at 40°C in a sealed eppendorf tube the levels of calcipotriol and
15 betamethasone were 106% and 117% of the original levels respectively. Stability testing is ongoing with this sample.

20

EXAMPLE 13

Three gel polyaphron dispersions and an aqueous gel of the following compositions were prepared by the following method.

25

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.0283%) isopropyl myristate (Lexol IPM NF - Inolex)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Calcipotriol solution* (0.0400% in aq phase) 29.6% w/w isopropanol in demineralised water	9.00
Poloxamer 188 (Pluronic F68 - BASF)	1.00

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone dipropionate solution** (0.3289%) in isopropyl myristate (Lexol IPM NF - Inolex)	89.10

- 52 -

Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Betamethasone dipropionate solution** (0.2252%) in 33.8% w/w isopropanol in demineralised water	9.00
Poloxamer 188 (Pluronic F68 - BASF)	1.00

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	20.00
Dimethicone (Q7-9120, 20 CP - Dow Corning)	35.00
Cyclomethicone (STb5NF - Dow Corning)	29.10
Elastomer DC10 (Dow Corning)	5.00
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

AQUEOUS GEL	
	% w/w
Polyacrylic acid (Ultrez 10 - Noveon)	2.00
Triethanolamine (TEA)	to pH 7.29
Isopropanol (30.6% w/w) in demineralised water	qs to 100%

FINAL PRODUCT	% w/w
GEL POLYAPHRON DISPERSION 1	25.00
GEL POLYAPHRON DISPERSION 2	25.00
GEL POLYAPHRON DISPERSION 3	20.00
AQUEOUS GEL	30.00

5 *The Calcipotriol content in the final formulation is 72.0µg/g.

**The betamethasone dipropionate content in the final formulation is 782µg/g (equivalent to 608µg/g of betamethasone).

10

Manufacturing Method

The method used was precisely as described for Example 1 above except that the neutralised aqueous gel was added to the final mixture of the three gel polyaphron dispersions, 1, 2 and 3 and then mixed in by simple mixing until the

15

- 53 -

product was a homogeneous mixture of the polyaphron dispersions.

Stability Measurements

5 Stability was tested as in Example 1a

The inventors observed that after 6 month storage at 40°C in a sealed eppendorf tube the levels of calcipotriol and betamethasone were 104% and 116% of the original levels
10 respectively. Storage stability testing is ongoing.

EXAMPLE 14

Three gel polyaphron dispersions and an aqueous gel of the
15 following compositions were prepared by the following method.

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.0243% w/w) in caprylic capric triglyceride (Mygliol 812- Condea)	89.00
Laureth-4 (Volpo L4 - Croda)	1.00
Aqueous Phase	
Calcipotriol solution* (0.0400% w/w) 33.3% w/w isopropanol in demineralised water	9.00
Poloxamer 188 (Pluronic F68 - BASF)	1.00

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone dipropionate solution** (0.203%w/w) in caprylic capric triglyceride (Mygliol 812- Condea)	89.00
Laureth-4 (Volpo L4 - Croda)	1.00
Aqueous Phase	
Betamethasone dipropionate solution** (0.192%w/w) in 33.3% w/w isopropanol in demineralised water	9.00
Poloxamer 188 (Pluronic F68 - BASF)	1.00

- 54 -

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	19.8
Dimethiconol 20 (- Dow Corning)	34.60
Cyclomethicone (STb5NF - Dow Corning)	28.70
Elastomer DC10 (Dow Corning)	4.90
Laureth-4 (Volpo L4 - Croda)	1.00
Tween 80 (Polysorbate 80 - Surfachem)	1.00
Aqueous Phase	
Isopropanol	3.00
Demineralised Water	6.00
Polyoxyl 40 hydrogenated castor oil (Cremophor RH 40 - BASF)	1.00

AQUEOUS GEL	
	% w/w
Polyacrylic acid (Ultrez 10 - Noveon)	2.00
Triethanolamine (TEA)	to pH 7.29
Isopropanol (30.6% w/w) in demineralised water	qs to 100%

FINAL PRODUCT	
	% w/w
GEL POLYAPHRON DISPERSION 1	15.00
GEL POLYAPHRON DISPERSION 2	27.00
GEL POLYAPHRON DISPERSION 3	40.00
AQUEOUS GEL	18.00

5

*The Calcipotriol content in the final formulation is 37.8 µg/g.

10 **The betamethasone dipropionate content in the final formulation is 534 µg/g (equivalent to 415 µg/g of betamethasone).

Manufacturing Method

15 The method used was precisely as described for Example 1 above except that the neutralised aqueous gel was added to the final mixture of the three gel polyaphron dispersions, 1, 2 and 3 and then mixed in by simple mixing until the product was a homogeneous mixture of the polyaphron dispersions.

20

- 55 -

Stability Measurements

Stability was tested as in Example 1a

The inventors observed that after 3.5 weeks storage at 40°C
 5 in a sealed eppendorf tube the levels of calcipotriol and
 betamethasone dipropionate were 99% and 138% of the original
 levels respectively. Storage stability testing is ongoing.

10

EXAMPLE 15

Three gel polyaphron dispersions and an aqueous gel of the
 following compositions were prepared by the following
 method.

15

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.0243% w/w) in caprylic capric triglyceride (Mygliol 812- Condea)	89.00
Laureth-4 (Volpo L4 - Croda)	1.00
Aqueous Phase	
Calcipotriol solution* (0.0400% w/w) 33.3% w/w isopropanol in demineralised water	9.00
Poloxamer 188 (Pluronic F68 - BASF)	1.00

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone dipropionate solution** (0.203%w/w) in caprylic capric triglyceride (Mygliol 812- Condea)	89.00
Laureth-4 (Volpo L4 - Croda)	1.00
Aqueous Phase	
Betamethasone dipropionate solution** (0.192%w/w) in 33.3% w/w isopropanol in demineralised water	9.00
Poloxamer 188 (Pluronic F68 - BASF)	1.00

- 56 -

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	19.8
Dimethiconol 20 (- Dow Corning)	34.60
Cyclomethicone (STb5NF - Dow Corning)	28.70
Elastomer DC10 (Dow Corning)	4.90
Laureth-4 (Volpo L4 - Croda)	1.00
Tween 80 (Polysorbate 80 - Surfachem)	1.00
Aqueous Phase	
Isopropanol	3.00
Demineralised Water	6.00
Polyoxyl 40 hydrogenated castor oil (Cremophor RH 40 - BASF)	1.00

AQUEOUS GEL	
	% w/w
Polyacrylic acid (Ultrez 10 - Noveon)	2.00
Triethanolamine (TEA)	to pH 7.29
Isopropanol (30.6% w/w) in demineralised water	qs to 100%

FINAL PRODUCT	
	% w/w
GEL POLYAPHRON DISPERSION 1	15.00
GEL POLYAPHRON DISPERSION 2	27.00
GEL POLYAPHRON DISPERSION 3	30.00
AQUEOUS GEL	28.00

*The Calcipotriol content in the final formulation is 37.8 µg/g.

**The betamethasone dipropionate content in the final formulation is 534 µg/g (equivalent to 415 µg/g of betamethasone).

10 **Manufacturing Method**

The method used was precisely as described for Example 1 above except that the neutralised aqueous gel was added to the final mixture of the three gel polyaphron dispersions, 1, 2 and 3 and then mixed in by simple mixing until the product was a homogeneous mixture of the polyaphron dispersions.

- 57 -

Stability Measurements

Stability was tested as in Example 1a

The inventors observed that after 3.5 weeks storage at 40°C in a sealed eppendorf tube the levels of calcipotriol and betamethasone were 95% and 136% of the original levels respectively. Storage stability testing is ongoing.

10

EXAMPLE 16

Three gel polyaphron dispersions and an aqueous gel of the following compositions were prepared by the following method.

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.0243% w/w) in caprylic capric triglyceride (Mygliol 812- Condea)	89.00
Laureth-4 (Volpo L4 - Croda)	1.00
Aqueous Phase	
Calcipotriol solution* (0.0400% w/w) 33.3% w/w isopropanol in demineralised water	9.00
Poloxamer 188 (Pluronic F68 - BASF)	1.00

15

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone dipropionate solution** (0.203%w/w) in caprylic capric triglyceride (Mygliol 812- Condea)	89.00
Laureth-4 (Volpo L4 - Croda)	1.00
Aqueous Phase	
Betamethasone dipropionate solution** (0.192%w/w) in 33.3% w/w isopropanol in demineralised water	9.00
Poloxamer 188 (Pluronic F68 - BASF)	1.00

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	19.8

- 58 -

Dimethiconol 20 (- Dow Corning)	34.60
Cyclomethicone (STb5NF - Dow Corning)	28.70
Elastomer DC10 (Dow Corning)	4.90
Laureth-4 (Volpo L4 - Croda)	1.00
Tween 80 (Polysorbate 80 - Surfachem)	1.00
Aqueous Phase	
Isopropanol	3.00
Demineralised Water	6.00
Polyoxyl 40 hydrogenated castor oil (Cremophor RH 40 - BASF)	1.00

AQUEOUS GEL	
	% w/w
Polyacrylic acid (Ultrez 10 - Noveon)	2.00
Triethanolamine (TEA)	to pH 7.29
Isopropanol (30.6% w/w) in demineralised water	qs to 100%

FINAL PRODUCT	% w/w
GEL POLYAPHRON DISPERSION 1	15.80
GEL POLYAPHRON DISPERSION 2	27.10
GEL POLYAPHRON DISPERSION 3	19.70
AQUEOUS GEL	37.40

*The Calcipotriol content in the final formulation is 39.9 µg/g.

- 5 **The betamethasone dipropionate content in the final formulation is 536 µg/g (equivalent to 417 µg/g of betamethasone).

Manufacturing Method

- 10 The method used was precisely as described for Example 1 above except that the neutralised aqueous gel was added to the final mixture of the three gel polyaphron dispersions, 1, 2 and 3 and then mixed in by simple mixing until the product was a homogeneous mixture of the polyaphron
- 15 dispersions.

Stability Measurements

Stability was tested as in Example 1a

- 59 -

The inventors observed that after 3.5 weeks storage at 40°C in a sealed eppendorf tube the levels of calcipotriol and betamethasone were 101% and 137% of the original levels respectively. Storage stability testing is ongoing.

5

Example 17

Gel POLYAPHRON DISPERSION 1	
Oil Phase	%
Calcipotriol solution* (0.02047%) in isopropyl myristate (Lexol IPM NF - Inolex)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

10

Gel POLYAPHRON DISPERSION 2	
Oil Phase	%
Betamethasone Dipropionate solution** (0.2775%) in isopropyl myristate (Lexol IPM NF - Inolex)	89.10
Laureth-4 (Volpo L4 - Croda)	0.90
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

Gel POLYAPHRON DISPERSION 3	
Oil Phase	% w/w
Squalane (Olive-derived - A & E Connock)	20.00
Dimethicone (Q7-9120, 100 CP - Dow Corning)	35.00
Elastomer DC10 (Dow Corning)	5.00
Cyclomethicone (STb5NF - Dow Corning)	29.00
Laureth-4 (Volpo L4 - Croda)	1.00
Aqueous Phase	
Poloxamer 188 (Pluronic F68 - BASF)	1.00
Demineralised Water	9.00

AQUEOUS GEL	
--------------------	--

- 60 -

	% w/w
Polyacrylic acid (Ultrez 10 - Noveon)	1.00
Triethanolamine (TEA)	to pH 7.25
Demineralised Water	qs to 100%

FINAL PRODUCT	% w/w
GEL POLYAPHRON DISPERSION 1	26.20
GEL POLYAPHRON DISPERSION 2	25.32
GEL POLYAPHRON DISPERSION 3	38.75
Aqueous gel	9.73

* final calcipotriol level 47.8 µg/g

5 ** final BDP level 626 µg/g (equivalent to 487 µg/g
betamethasone)

Manufacturing Method

10 The method used was precisely as described for Example 1
above except that the neutralised aqueous gel was added to
the final mixture of the three gel polyaphron dispersions,
1, 2 and 3 and then mixed in by simple mixing until the
product was a homogeneous mixture of the polyaphron
dispersions.

15

- 61 -

Summary

ID	storage medium	stability (time)
Example 1	Glass	2 months (ongoing)
Example 1	Plastic	2 months ^a
Example 2	Glass	2 months* (ongoing)
Example 3	Glass	9 months
Example 4	Glass	4.5 months**
Example 5	Glass	ongoing
Example 6	Plastic	3 months (ongoing)
Example 7	Plastic	3 months***
Example 8	Glass	3 months**** Unstable
Example 9	Glass	18 months at room temp
Example 10	Glass	6 months
Example 11	Glass	6 months
Example 12	Plastic	6 months
Example 13	Plastic	6 months
Example 14	Plastic	3.5 weeks
Example 15	Plastic	3.5 weeks
Example 16	Plastic	3.5 weeks

5 Stability as defined above; less than 5% loss of either drug compared with initial drug levels. Storage at 40°C, unless otherwise stated.

^a levels of calcipotriol and betamethasone dipropionate 102% and 99% of original levels

10 * Level of betamethasone = 95% of original in latest storage sample.

** After 6 months, levels of calcipotriol and betamethasone dipropionate were 42% and 65% of original, respectively.

15 *** Level of calcipotriol = 86% of original after 4 months at 40°C.

**** After 3 months, levels of calcipotriol and betamethasone dipropionate were 91% and 70% of original, respectively.

Printed: 16-06-2009

CLMSPAMD

15-JAN-2009 14:15 FROM BOULT WADE TENNANT

PCT/GB 2008/000 894

TO 00498923994465

P.12

PCT/GB 2008/000 894 - 15-01-2009

- 62 -

CLAIMS

1. A composition suitable for topical application comprising a continuous phase and at least one discontinuous phase, said composition comprising at least one polyaphron dispersion, at least one vitamin D or vitamin D analogue and at least one corticosteroid, wherein the corticosteroid is predominantly in the discontinuous phase, and the vitamin D or vitamin D analogue is predominantly in the discontinuous phase.
2. A composition according to claim 1 wherein the corticosteroid is betamethasone, clobetasol, clobetasone, desoximethasone, diflucortolon, difluorasona, fluocinoid, flumethasone, fluocinolone, fluticasone, fluprednidene, halcinonide, hydrocortisone, momethasone, triamcinolone, and their esters or a mixture thereof.
3. A composition according to claim 2 wherein the corticosteroid is betamethasone or an ester thereof.
4. A composition according to any one of the preceding claims wherein the vitamin D or vitamin D analogue is vitamin D, calcipotriol, seocalcitol, calcitriol, tacalcitol, maxacalcitol, paricalcitol, falecalcitriol, 1 α ,24S-dihydroxy-vitamin D₂, 1(S),3(R)-dihydroxy-20(R)-[[(3-(2-hydroxy-2-propyl)-phenyl)-methoxy)-methyl]-9,10-seco-pregna-5(Z),7(E),10(19)-triene, or a mixture thereof.
5. A composition according to claim 4 wherein the vitamin D or vitamin D analogue is calcipotriol.

- 63 -

6. A composition according to any one of the preceding claims wherein the discontinuous phase comprises an oil.

7. A composition according to claim 6 wherein the oil comprises a monoglyceride, a diglyceride, a triglyceride, isopropyl myristate or a mixture thereof.

8. A composition according to any one of the preceding claims wherein the composition comprises less than 5% by weight of water, based on the total weight of the composition.

9. A composition according to claim 8 wherein the continuous phase is non-aqueous.

15

10. A composition according to any one of claims 1 to 7 wherein the composition comprises at least 5% by weight of water based on the total weight of the composition.

20

11. A composition according to claim 10 wherein the composition comprises at least 10% by weight of water, based on the total weight of the composition.

25

12. A composition of according to claim 11 wherein the composition comprises at least 15% by weight of water based on the total weight of the composition.

30

13. A composition of any one of claims 1 to 7 or 10 to 12 wherein the composition comprises from 5 to 50% by weight of water based on the total weight of the composition.

- 64 -

14. A composition according to any one of the preceding claims wherein the continuous phase comprises a compound of formula R_1-OH where R_1 is C_1-C_{10} alkyl and/or a compound of formula $HO-R_2-H$ where R_2 is $(C_2H_4)_n$ or $(C_3H_6)_n$ where n is 1 to 100.
15. A composition according to claim 14 wherein the continuous phase comprises propylene glycol, glycerol, ethanol, isopropyl alcohol or a mixture thereof.
16. A composition according to any one of the preceding claims wherein the pH of the composition is from 7.0 to 8.5.
17. A composition according to claim 16 wherein the pH of the aqueous phase from 7.25 to 7.75.
18. A composition according to any one of the preceding claims further comprising a gelling agent.
19. A composition according to any one of the preceding claims further comprising a permeation enhancer.
20. A composition as defined in any one of claims 1 to 19 for use in a method of treatment of the human or animal body by therapy.
21. A composition as defined in any one of claims 1 to 19 for use in the treatment of psoriasis.
22. Use of a composition as defined in any one of claims 1 to 20 in the manufacture of a medicament for treatment of psoriasis.

- 65 -

23. A method of making the composition as defined in any one of claims 1 to 20 comprising the following steps:

5 (i) providing a hydrophilic solvent, optionally comprising at least one vitamin D or a vitamin D analogue, and/or at least one corticosteroid, and/or a surfactant;

10 (ii) providing a hydrophobic solvent optionally comprising at least one vitamin D or a vitamin D analogue, and/or at least one corticosteroid, and/or a surfactant;

15 (iii) mixing the hydrophilic solvent with the hydrophobic solvent under suitable conditions to form the composition comprising at least one polyaphron dispersion, at least one vitamin D or vitamin D analogue and at least one corticosteroid.

24. A method of making the composition as defined in any one of claims 1 to 20 comprising the following steps:

20 preparing a first polyaphron dispersion comprising a vitamin D or vitamin D analogue;

preparing a second polyaphron dispersion comprising a corticosteroid;

25 and mixing together said first and second polyaphron dispersions to form the composition.

25. The method according to any one of claims 23 to 24 further comprising

30 preparing a third or further polyaphron dispersion comprising an active agent;

- 66 -

and mixing said third or further polyaphron dispersion with said first and second polyaphron dispersions.

26. The method of any one of claims 24 to 25 wherein said
5 first and second polyaphron are mixed together with a further hydrophilic solvent, wherein the pH of the hydrophilic solvent is in the range of from 7.25 to 7.75.

27. A composition as defined in any one of claims 1 to 19
10 which is stable, wherein stability is measured as a no more than 5% reduction in the amount of vitamin D or vitamin D analogue and a no more than a 5% reduction in the amount of corticosteroid with respect to the original amounts after 3 months of storage in a sealed glass container at 40°C.

15

1/2




Figure 1




Figure 2




Figure 3