



(51) International Patent Classification:

C07D 487/04 (2006.01) A61K 31/5377 (2006.01)
 C07D 519/00 (2006.01) A61K 31/538 (2006.01)
 A61P 33/00 (2006.01) A61K 31/695 (2006.01)
 A01N 43/90 (2006.01) C07F 7/00 (2006.01)
 A61K 31/5025 (2006.01)

(21) International Application Number:

PCT/US2021/033072

(22) International Filing Date:

19 May 2021 (19.05.2021)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/031,656 29 May 2020 (29.05.2020) US

(71) Applicants: **BOEHRINGER INGELHEIM ANIMAL HEALTH USA INC.** [US/US]; 3239 Satellite Blvd., Duluth, GA 30096 (US). **BOEHRINGER INGELHEIM PHARMA GMBH & CO. KG** [—/DE]; Binger Strasse 173, 55216 Ingelheim am Rhein (DE).

(72) Inventors: **KOOLMAN, Hannes, Fiepko**; Boehringer Ingelheim GmbH, Corporate Patents, Binger Strasse 173, 55216 Ingelheim am Rhein (DE). **HERLE, Bart**; Boehringer Ingelheim GmbH, Corporate Patents, Binger Strasse 173, 55216 Ingelheim am Rhein (DE). **LONG, Alan**; c/o AH Patents, Boehringer Ingelheim AH USA, 3239 Satellite Blvd., Duluth, GA 30096 (US).

(74) Agent: **EZCURRA, John Esteban**; Boehringer Ingelheim Animal Health USA Inc., 3239 Satellite Blvd., Duluth, GA 30096 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,


CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: ANTHELMINTIC HETEROCYCLIC COMPOUNDS



(I)

(57) Abstract: This invention provides for compounds of the formula (I): wherein the variables are defined herein, or salt thereof, compositions comprising these compounds, and method for the treatment, control or prevention of a parasitic infestation or infection in an animal in need thereof by administering an effective amount of these compounds to said animal.



ANTHELMINTIC HETEROCYCLIC COMPOUNDS

FIELD OF THE INVENTION

This patent application relates to new antiparasitic compounds, compositions comprising the compounds, processes for their preparation, and methods of using the compounds to control parasites that harm animals and humans.

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of priority to U.S. Provisional Application No. 63/031,656 filed May 29, 2020, which is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

Animals, such as mammals and birds, are often susceptible to parasite infestations. These parasites may be ectoparasites, such as fleas and ticks. Animals and humans also suffer from endoparasitic infections including, for example, helminthiasis which is most frequently caused by a group of parasitic worms described as nematodes or roundworms. These parasites cause severe economic losses in pigs, sheep, horses, and cattle as well as affecting companion animals (e.g. cats and dogs) and poultry. Other parasites include those which occur in the gastrointestinal tract of animals and humans such as *Ancylostoma*, *Necator*, *Ascaris*, *Strongyloides*, *Trichinella*, *Capillaria*, *Toxocara*, *Toxascaris*, *Trichuris* and *Enterobius*. Other parasites which are found in the blood or other tissues and organs include filarial worms and the extra intestinal stages of *Strongyloides* and *Trichinella*.

One type of endoparasite which seriously harms mammals is *Dirofilaria immitis*, also known as Heartworm. Other filarial endoparasites include *Dirofilaria repens* and *Dirofilaria honkongensis*, which can also infect humans. The most common hosts are dogs and cats but other mammals such as ferrets and raccoons may also be infected. Heartworms go through several life stages before they become adults infecting the pulmonary artery of the host mammal. The worms require the mosquito as an intermediate host to complete their life cycle. The period between the initial infection when the dog is bitten by a mosquito and the maturation of the worms into adults living in the heart and pulmonary arteries is six to seven months in dogs and is known as the "prepatent period." L3 larvae migrate during blood feeding of the mosquito to the tip of the

mosquito's mouth parts (labium), leave the mosquito and are deposited on the skin of the dog where they then migrate through the bite wound into the host. Most L3 larvae molt to fourth-stage larvae (L4s) in canine subcutaneous tissues within 1-3 days after infection. Then, they migrate to the muscles of the chest and abdomen, and 45 to 60 days after infection, molt to the fifth stage (L5, immature adult). Between 75 and 120 days after infection, these immature heartworms then enter the bloodstream and are carried through the heart to reside in the pulmonary artery. Around seven months after infection, *Dirofilaria immitis* adults reach maturity and sexually reproduce in the pulmonary arteries and right ventricle. Adult males are around 15cm in length, and females are around 25cm in length and their normal life span as adults is calculated to be about 5 years.

Heartworm infection is a severe and life-threatening disease. Canine heartworm infection is preventable and prophylaxis treatment is a priority in heartworm endemic areas. Treatment of mature heartworm infection with an adulticide (e.g. melarsomine dihydrochloride) is costly and can cause serious adverse side effects, thus prevention by monthly administration of drugs that interrupt larvae development is widely used. The goal of marketed heartworm preventive therapies in dogs is to prevent the development of the parasite to adult heartworms by interrupting the *Dirofilaria immitis* life cycle post-infection.

The macrocyclic lactones (MLs, e.g. ivermectin, eprinomectin, milbemycin oxime, moxidectin, and selamectin) are the most commonly used chemoprophylaxis agents and are administered at monthly or six-month intervals. These drugs have been effective against *Dirofilaria immitis* infective third-stage larvae (L3) deposited by the mosquito as well as maturing fourth-stage larvae (L4). When administered monthly, MLs kill L3 and L4 larvae acquired within the previous 30 days, and thus prevent disease caused by adult worms. MLs can also be used monthly in infected dogs to suppress reproduction in adult worms and remove microfilariae, thereby reducing transmission and gradually causing the attrition of adult worms (*Vet. Parasitol.* 2005 Oct 24 133(2-3) 197-206).

In recent years, an increased number of lack of efficacy (LOE) cases have been reported, in which dogs develop mature heartworm infections despite receiving monthly prophylactic doses of macrocyclic lactones drugs. For example, Atkins et al., (*Vet. Parasitol.* 206 (2014) 106–113) recently reported that an increasing number of cases of dogs that tested heartworm antigen positive

while receiving heartworm preventive medication which suggests that some populations of *Dirofilaria immitis* have developed selectional resistance to heartworm preventives (American Heartworm Society, 2010. Heartworm Preventive Resistance. Is it Possible, vol. 37. Bulletin of the American Heartworm Society, pp. 5.). Thus, there is an ongoing need to develop new anthelmintic agents with improved activity against *Dirofilaria immitis* and other endoparasites.

WO 2017/178416 A1 provides pyrazolopyrimidine derivatives for the control, treatment and/or prevention of helminths. WO 2018/197401 A1 provides bicyclic pyrazole derivatives for the control, treatment and/or prevention of helminths. WO 2018/087036 A1 provides quinolone-3-carboxamide derivatives for the control, treatment and/or prevention of helminths. WO 2019/025341 provides quinoline compounds for the treatment, control and/or prevention of helminth infections and WO 2019/002132 A1 azaquinone derivatives for the control, treatment and/or prevention of helminths. All these publications are to Bayer Animal Health GmbH and are incorporated herein by reference in their entirety.

More recently WO 2020/014068 A1 (incorporated herein by reference) describes anthelmintic heterocyclic compounds that were found to be active against *Dirofilaria immitis*.


It is expressly noted that citation or identification of any document in this application is not an admission that such document is available as prior art to the present description. Any foregoing applications, and all documents cited therein or during their prosecution (“application cited documents”) and all documents cited or referenced in the application cited documents, and all documents cited or referenced herein (“herein cited documents”), and all documents cited or referenced in herein cited documents, together with any manufacturer’s instructions, descriptions, product specifications, and product sheets for any products mentioned herein or in any document incorporated by reference herein, are hereby incorporated herein by reference, and may be employed in the practice of the description.

SUMMARY OF THE INVENTION

The present application provides for novel anthelmintic and antiparasitic heterocyclic compounds with improved activity against endoparasites and ectoparasites. The application is also directed to compositions comprising the compounds, methods and uses of the compounds for

eradicating, controlling, and/or preventing a parasitic infection and/or infestation in animals including humans. The compounds may be administered to animals, particularly mammals, fish and birds, to prevent and/or treat parasitic infections.

An aspect of the present invention includes a compound of Formula (I):



(I)

a stereoisomer, tautomer, N-oxide, hydrate, solvate, or salt thereof, wherein variables R^1 , R^2 , R^3 , R^9 , $R^{9'}$, Y^1 , Y^2 , Y^3 , Y^4 , Y^5 , Y^6 , L , Q , W , Z , a and q are defined herein, and the dashed bonds (---) signifies a single or double bond.

The invention also includes a veterinarily acceptable composition comprising a compound of Formula (I) and a veterinarily acceptable carrier and a method of controlling parasites, including helminths, comprising administering the compound, or the veterinarily acceptable composition thereof, to an animal in need thereof. An embodiment of the invention also includes the use of the compound of Formula (I) for eradicating, controlling, and/or preventing a parasitic infection and/or infestation in animals or humans. The compounds of the invention may be administered to animals, particularly mammals, fish and birds, to prevent and/or treat parasitic infections and/or infestations.

The compound and compositions comprising the compound are highly effective for the treatment and/or prophylaxis of internal parasites in mammals, fish and birds, and in particular, cats, dogs, horses, chickens, pigs, sheep and cattle, with the aim of substantially ridding these hosts of endoparasites.

In an embodiment, compounds of Formula (I) and compositions comprising said compounds are substantially effective against endoparasites, such as filariae (e.g. heartworm), and hookworms, whipworms and roundworms of the digestive tract of animals and humans. In certain
5 effective against *Dirofilaria immitis* (heartworm) isolates that are less sensitive to treatment with macrocyclic lactones. In another embodiment, the compounds and compositions of the invention are effective for treating and/or preventing infections of animals with nematodes that are less sensitive to treatment with commercially available or known active agents.

In an embodiment, the description includes a combination of a compound of Formula (I)
10 with at least a second active agent, which may broaden the scope of protection afforded to animals against endoparasites and/or ectoparasites.

Another embodiment includes a method for the treatment and/or prevention of a parasitic infection and/or infestation in an animal comprising administering a compound of Formula (I) to the animal. Another embodiment includes a use of a compound of Formula (I) for the treatment
15 and/or prevention of a parasitic infection and/or infestation in an animal and the use of the compound of Formula (I) in the preparation of a medicament for the treatment and/or prevention of a parasitic infection in an animal.

Thus, the invention includes the following non-limiting embodiments:

(a) a compound of Formula (I) or a pharmaceutically or a veterinarily
20 acceptable salt thereof, which is an active endoparasiticide and in some cases is also active against ectoparasites;

(b) a veterinary composition comprising a parasitically effective amount of a compound of Formula (I), or a pharmaceutically or veterinarily acceptable salt thereof, in combination with a pharmaceutically or veterinarily acceptable carrier
25 or diluent;

(c) a veterinary composition comprising a parasitically effective amount of a compound of Formula (I), or a pharmaceutically or veterinarily acceptable salt thereof, in combination with one or more additional active agents (*i.e.*, active

ingredient not encompassed by Formula (I)) and a pharmaceutically or veterinarily acceptable carrier or diluent;

(d) a method for treating a parasitic infection and/or infestation in or on an animal comprising administering a parasitically effective amount a compound of Formula (I), or a pharmaceutically or veterinarily acceptable salt thereof, optionally with one or more additional active agents (*i.e.*, active ingredient not encompassed by Formula (I)), to the animal in need thereof;

(e) a method for the prevention of a parasitic infection and/or infestation of an animal, which comprises administering a parasitically effective amount of a compound of Formula (I), or a pharmaceutically or a veterinarily acceptable salt thereof, optionally with one or more additional active agents (*i.e.*, active ingredient not encompassed by Formula (I)), to the animal in need thereof;

(f) the use of a compound of Formula (I), or a pharmaceutically or a veterinarily acceptable salt thereof, optionally with one or more additional active agents (*i.e.*, active ingredient not encompassed by Formula (I)), for the treatment and/or prevention of a parasitic infection and possibly also a parasitic infestation in an animal;

(g) a use of a compound of Formula (I), or a pharmaceutically or veterinarily acceptable salt thereof, optionally with one or more additional active agents (*i.e.*, active ingredient not encompassed by Formula (I)), for the manufacture of a veterinary medicament for the treatment and/or prevention of a parasitic infection and/or infestation in an animal; and

(h) a process for the preparation of a compound of Formula (I).

These and other embodiments are disclosed or are obvious from and encompassed by, the following Detailed Description.

DEFINITIONS:

It is noted that in this disclosure and particularly in the claims and/or paragraphs, terms such as "comprises", "comprised", "comprising" and the like can be interpreted as "includes", "included", "including", and the like; and that terms such as "consisting essentially of" and "consists essentially of" are interpreted as allowing for elements not explicitly recited, but excluding elements that are found in the prior art or that affect a basic or novel characteristic of the invention.

Terms used herein will have their customary meanings in the art unless otherwise specified. The organic moieties mentioned in the definitions of the variables of the compounds e.g., the compound of formula (I) are like the term halogen – i.e., collective terms for individual listings of the individual group members – fluoro, chloro, bromo and iodo with respect to halogen. The prefix C_n-C_m indicates in each case the possible number of carbon atoms in the group from an integer n to another integer m.

In the present specification and claims the term “including but not limited to” is equivalent to “included”.

The term “compound of Formula (I)” includes any stereoisomer, tautomer, N-oxide, hydrate, solvate, or salt thereof.

By the term “optionally substituted” is meant a radical that is optionally substituted by one or more of the following moieties: halogen, hydroxyl, alkyl, haloalkyl, carboxyl, acyl, acyloxy, alkylcarbonyl, haloalkylcarbonyl, alkoxycarbonyl, haloalkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, haloalkylaminocarbonyl, dihaloalkylaminocarbonyl, amino, alkyl- or dialkylamino, amido, arylamino, alkoxy, haloalkoxy, aryloxy, nitro, cyano, azido, thiol, thioamide, imino, amidine, guanidine, carbonate, silyl, silyl ether, SF₅, sulfonic acid, sulfate, sulfonyl, alkoxysulfonyl, sulfanyl, sulfinyl, sulfamoyl, sulfoximine, sulfinimine, sulfonimidamide, sulfonediimine, ester, phosphonyl, phosphinyl, phosphoryl, phosphine, phosphonamidate, phosphinamidate, phosphinate, phosphine oxide, thioester, thioether, acid halide, anhydride, oxime, hydrazine, carbamate, phosphonic acid, phosphate, phosphonate, aryl, and heteroaryl.

In some embodiments, the term “optionally substituted” includes substitution of a core group with halogen (chloro, fluoro, bromo, iodo), C₁-C₆-alkyl, C₁-C₆-haloalkyl, 3- to 8-membered

cycloalkyl, amino, C₁-C₆-alkylamino, C₁-C₆-dialkylamino, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, cyano, nitro, SF₅, acetyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-dialkylaminocarbonyl, C₁-C₆-haloalkylaminocarbonyl, C₁-C₆-dihaloalkylaminocarbonyl, C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, C₁-C₆-haloalkylthio, C₁-C₆-haloalkylsulfinyl, C₁-C₆-haloalkylsulfonyl, phenyl, 5- or 6-membered heteroaryl, 5- or 6-membered heterocyclyl.

In other embodiments, the term “optionally substituted” includes substitution of a core group with halogen (chloro, fluoro, bromo, iodo), C₁-C₃-alkyl, C₁-C₃-haloalkyl, 3- to 8-membered cycloalkyl, amino, C₁-C₃-alkylamino, C₁-C₃-dialkylamino, C₁-C₃-alkoxy, C₁-C₃-haloalkoxy, cyano, nitro, SF₅, acetyl, C₁-C₃-alkoxycarbonyl, C₁-C₃-haloalkoxycarbonyl, C₁-C₃-alkylcarbonyl, C₁-C₃-haloalkylcarbonyl, aminocarbonyl, C₁-C₃-alkylaminocarbonyl, C₁-C₃-dialkylaminocarbonyl, C₁-C₃-haloalkylaminocarbonyl, C₁-C₃-dihaloalkylaminocarbonyl, C₁-C₃-alkylthio, C₁-C₃-alkylsulfinyl, C₁-C₃-alkylsulfonyl, C₁-C₃-haloalkylthio, C₁-C₃-haloalkylsulfinyl, C₁-C₃-haloalkylsulfonyl, phenyl, 5- or 6-membered heteroaryl, 5- or 6-membered heterocyclyl.

In certain embodiments, the term “optionally substituted” includes substitution by halogen (chloro, fluoro, bromo and iodo), methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, hydroxyl, thiol, amino, methylamino, ethylamino, propylamino, butylamino, dimethylamino, diethylamino, methoxy, ethoxy, propoxy, CF₃, CF₂CF₃, -OCF₃, -OCF₂CF₃, -SCH₃, -SCF₃, -S(O)CH₃, -S(O)CF₃, -S(O)₂CH₃, -S(O)₂CF₃, morpholino, piperidiny, pyridyl and phenyl.

In some embodiments the compounds may be substituted with a viable functional group that does not inhibit the biological activity of the compounds of the description, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in Greene and Wuts, Protective Groups in Organic Synthesis, John Wiley and Sons, Third Edition, 1999, hereby incorporated by reference. For avoidance of doubt, “optionally substituted alkyl” includes haloalkyl and hydroxyalkyl.

Unless otherwise stated, “alkyl” means, either alone or in combination with a heteroatom, e.g., alkoxy, thioalkyl, alkylamino, and the like, saturated straight, branched, primary, secondary or tertiary hydrocarbons, including those having 1 to 12 atoms. In some embodiments, alkyl groups will include C₁-C₁₀, C₁-C₈, C₁-C₆, C₁-C₄ or C₁-C₃ alkyl groups. Examples of C₁-C₁₀ alkyl include, but are not limited to, methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-

ethylpropyl, hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, heptyl, octyl, 2-ethylhexyl, nonyl and decyl and their isomers. C₁-C₄-alkyl means for example methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl or 1,1-dimethylethyl.

Cyclic alkyl groups may be referred to as “cycloalkyl” and include those with 3 to 10 carbon atoms having single or multiple fused rings. Non-limiting examples of cycloalkyl groups include adamantyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl and the like.

“Carbocyclic” groups are cyclic groups composed exclusively of carbon. The carbocyclic groups include both aromatic rings such as phenyl and non-aromatic rings such as cycloalkyl rings including cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like, and include those with 3 to 14 carbon atoms having single or multiple fused rings.

The term “alkenyl” refers to both straight and branched carbon chains which have at least one carbon-carbon double bond. In some embodiments, alkenyl groups may include C₂-C₁₂ alkenyl groups. In other embodiments, alkenyl includes C₂-C₁₀, C₂-C₈, C₂-C₆, C₂-C₄, or C₃-C₄ alkenyl groups. In one embodiment of alkenyl, the number of double bonds is 1-3; in another embodiment of alkenyl, the number of double bonds is one. Other ranges of carbon-carbon double bonds and carbon numbers are also contemplated depending on the location of the alkenyl moiety on the molecule. “Alkenyl” groups may include more than one double bond in the chain. Examples of alkenyl or a specific range thereof include, but are not limited to, ethenyl, 1-propenyl, 2-propenyl, 1-methyl-ethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl; 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-1-butenyl, 2-methyl-1-butenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-1-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-1-propenyl, 1-ethyl-2-propenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-1-pentenyl, 2-methyl-1-pentenyl, 3-methyl-1-pentenyl, 4-methyl-1-pentenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 1-methyl-3-pentenyl, 2-methyl-3-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-

4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-1-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-1-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 2,2-dimethyl-3-butenyl, 2,3-dimethyl-1-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 3,3-dimethyl-1-butenyl, 3,3-dimethyl-2-butenyl, 1-ethyl-1-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-1-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl, 1-ethyl-2-methyl-1-propenyl and 1-ethyl-2-methyl-2-propenyl.

“Alkynyl” refers to both straight and branched carbon chains which have at least one carbon-carbon triple bond. In one embodiment of alkynyl, the number of triple bonds is 1-3; in another embodiment of alkynyl, the number of triple bonds is one. In some embodiments, alkynyl groups include from 2 to 12 carbon atoms. In other embodiments, alkynyl groups may include C₂-C₁₀, C₂-C₈, C₂-C₆ or C₂-C₄ alkynyl groups. Other ranges of carbon-carbon triple bonds and carbon numbers are also contemplated depending on the location of the alkynyl moiety on the molecule. For example, the term “C₂-C₁₀-alkynyl” as used herein refers to a straight-chain or branched unsaturated hydrocarbon group having 2 to 10 carbon atoms and containing at least one triple bond, such as ethynyl, prop-1-yn-1-yl, prop-2-yn-1-yl, n-but-1-yn-1-yl, n-but-1-yn-3-yl, n-but-1-yn-4-yl, n-but-2-yn-1-yl, n-pent-1-yn-1-yl, n-pent-1-yn-3-yl, n-pent-1-yn-4-yl, n-pent-1-yn-5-yl, n-pent-2-yn-1-yl, n-pent-2-yn-4-yl, n-pent-2-yn-5-yl, 3-methylbut-1-yn-3-yl, 3-methylbut-1-yn-4-yl, n-hex-1-yn-1-yl, n-hex-1-yn-3-yl, n-hex-1-yn-4-yl, n-hex-1-yn-5-yl, n-hex-1-yn-6-yl, n-hex-2-yn-1-yl, n-hex-2-yn-4-yl, n-hex-2-yn-5-yl, n-hex-2-yn-6-yl, n-hex-3-yn-1-yl, n-hex-3-yn-2-yl, 3-methylpent-1-yn-1-yl, 3-methylpent-1-yn-3-yl, 3-methylpent-1-yn-4-yl, 3-methylpent-1-yn-5-yl, 4-methylpent-1-yn-1-yl, 4-methylpent-2-yn-4-yl or 4-methylpent-2-yn-5-yl and the like.

The term “haloalkyl” refers to an alkyl group, as defined herein, which is substituted by one or more halogen atoms. For example C₁-C₄-haloalkyl includes, but is not limited to, chloromethyl, bromomethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, 1-chloroethyl, 1-bromoethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl, pentafluoroethyl and the like. The term “fluoroalkyl” as used herein refers to an alkyl in which one or more of the hydrogen atoms is replaced with fluorine atoms, for example difluoromethyl,

trifluoromethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 1,1,2,2-tetrafluoroethyl or pentafluoroethyl.

The term “haloalkenyl” refers to an alkenyl group, as defined herein, which is substituted by one or more halogen atoms.

5 The term “haloalkynyl” refers to an alkynyl group, as defined herein, which is substituted by one or more halogen atoms.

The term “alkoxy” refers to alkyl-O-, wherein alkyl is as defined above. Similarly, the terms “alkenyloxy,” “alkynyloxy,” “haloalkoxy,” “haloalkenyloxy,” “haloalkynyloxy,” “cycloalkoxy,” “cycloalkenyloxy,” “halocycloalkoxy,” and “halocycloalkenyloxy” refer to the
 10 groups alkenyl-O-, alkynyl-O-, haloalkyl-O-, haloalkenyl-O-, haloalkynyl-O-, cycloalkyl-O-, cycloalkenyl-O-, halocycloalkyl-O-, and halocycloalkenyl-O-, respectively, wherein alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, cycloalkyl, cycloalkenyl, halocycloalkyl, and halocycloalkenyl are as defined above. Examples of C₁-C₆-alkoxy include, but are not limited to, methoxy, ethoxy, OCH₂-C₂H₅, OCH(CH₃)₂, n-butoxy, OCH(CH₃)-C₂H₅, OCH₂-CH(CH₃)₂,
 15 OC(CH₃)₃, n-pentoxy, 1-methylbutoxy, 2-methylbutoxy, 3-methylbutoxy, 1,1-dimethylpropoxy, 1,2-dimethylpropoxy, 2,2-dimethyl-propoxy, 1-ethylpropoxy, n-hexoxy, 1-methylpentoxy, 2-methylpentoxy, 3-methylpentoxy, 4-methylpentoxy, 1,1-dimethylbutoxy, 1,2-dimethylbutoxy, 1,3-dimethylbutoxy, 2,2-dimethylbutoxy, 2,3-dimethylbutoxy, 3,3-dimethylbutoxy, 1-ethylbutoxy, 2-ethylbutoxy, 1,1,2-trimethylpropoxy, 1,2,2-trimethylpropoxy, 1-ethyl-1-
 20 methylpropoxy, 1-ethyl-2-methylpropoxy and the like.

The term “aryl” refers to a monovalent aromatic carbocyclic group of from 6 to 14 carbon atoms having a single ring or multiple fused rings. Aryl groups include, but are not limited to, phenyl, biphenyl, and naphthyl. In some embodiments aryl includes tetrahydronaphthyl, phenylcyclopropyl and indanyl. Aryl groups may be unsubstituted or substituted by one or more
 25 moieties selected from halogen, cyano, nitro, hydroxy, mercapto, amino, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy, haloalkenyloxy, haloalkynyloxy, cycloalkoxy, cycloalkenyloxy, halocycloalkoxy, halocycloalkenyloxy, alkylthio, haloalkylthio, cycloalkylthio, halocycloalkylthio, alkylsulfinyl, alkenylsulfinyl, alkynyl-sulfinyl, haloalkylsulfinyl, haloalkenylsulfinyl, haloalkynylsulfinyl, alkylsulfonyl, alkenylsulfonyl, alkynylsulfonyl,
 30 haloalkenylsulfonyl, haloalkynylsulfonyl, alkylsulfonyl, alkenylsulfonyl, alkynylsulfonyl,

haloalkyl-sulfonyl, haloalkenylsulfonyl, haloalkynylsulfonyl, -SF₅, alkylamino, alkenylamino, alkynylamino, di(alkyl)amino, di(alkenyl)-amino, di(alkynyl)amino, or trialkylsilyl.

The term “aralkyl” refers to an aryl group that is bonded to the parent compound through a diradical alkylene bridge, (-CH₂-)_n, where n is 1-12 and where “aryl” is as defined above.

5 The term “heteroaryl” refers to a monovalent aromatic group of from 1 to 15 carbon atoms, preferably from 1 to 10 carbon atoms, having one or more oxygen, nitrogen, and sulfur heteroatoms within the ring, preferably 1 to 4 heteroatoms, or 1 to 3 heteroatoms. The nitrogen and sulfur heteroatoms may optionally be oxidized. Heteroaryl groups will typically include a 5- or 6-membered aromatic ring. Such heteroaryl groups can have a single ring (*e.g.*, pyridyl or furyl) or
10 multiple fused rings provided that the point of attachment is through a heteroaryl ring atom. Examples of heteroaryls include pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, pyrrolyl, indolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, furanyl, thiophenyl, furyl, pyrrolyl, imidazolyl, oxazolyl, isoxazolyl, isothiazolyl, pyrazolyl, benzofuranyl, benzothiophenyl, imidazopyridyl, imidazopyrimidyl, or pyrrolopyrimidyl. Heteroaryl rings may be unsubstituted or
15 substituted by one or more moieties as described for aryl above.

 The term “heterocyclyl,” “heterocyclic” or “heterocyclo” refers to fully saturated or partially unsaturated, but non-aromatic cyclic groups, for example, 3 to 7 membered monocyclic, 7 to 11 membered bicyclic, or 10 to 15 membered tricyclic ring systems, which have one or more oxygen, sulfur, silicon or nitrogen heteroatoms in ring, preferably 1 to 4 or 1 to 3 heteroatoms. The
20 nitrogen and sulfur heteroatoms may optionally be oxidized and the nitrogen heteroatoms may optionally be quaternized. The heterocyclic group may be attached at any heteroatom or carbon atom of the ring or ring system and may be unsubstituted or substituted by one or more moieties as described for aryl groups above.

 Exemplary monocyclic heterocyclic groups include, but are not limited to, aziridinyl, azetidyl, oxetanyl, pyrrolidinyl, pyrrolyl, pyrazolyl, oxetanyl, pyrazolinyl, imidazolyl, imidazolinyl, imidazolidinyl, oxazolyl, oxazolidinyl, isoxazolinyl, isoxazolyl, thiazolyl, thiadiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, furyl, tetrahydrofuryl, thienyl, oxadiazolyl, piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodinyl, 2-oxoazepinyl, azepinyl, 4-piperidonyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl,
30 tetrahydropyranyl, morpholinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, 1,3-dioxolane and tetrahydro-1,1-dioxothienyl, triazolyl, triazinyl, and the like.

Exemplary bicyclic heterocyclic groups include, but are not limited to, indolyl, benzothiazolyl, benzoxazolyl, benzodioxolyl, benzothienyl, quinuclidinyl, quinolinyl, tetrahydroisoquinolinyl, isoquinolinyl, benzimidazolyl, benzopyranyl, indolizinyl, benzofuryl, chromonyl, coumarinyl, benzopyranyl, cinnolinyl, quinoxalinyl, indazolyl, pyrrolopyridyl, furopyridinyl (such as furo[2,3-c]pyridinyl, furo[3,2-b]pyridinyl) or furo[2,3-b]pyridinyl), dihydroisoindolyl, dihydroquinazolinyl (such as 3,4-dihydro-4-oxo-quinazolinyl), tetrahydroquinolinyl and the like.

Bicyclic and tricyclic carbocyclic or heterocyclic ring systems include spirocyclic systems in which at least two of the rings in the system are connected through a single carbon atom. The spirocyclic ring systems will include a combination of from 3- to 8-membered carbocyclic and/or heterocyclic ring systems joined at a common carbon atom. Thus, the spirocyclic ring systems may include a 3-membered ring bonded to another 3-membered ring (either carbocyclic or heterocyclic) to an 8-membered ring bonded to another 8-membered ring and all the combinations of different ring sizes between. The heterocyclic ring component of a spirocyclic ring system will include one or two heteroatoms selected from N, O, Si or S.

The term "alkylthio" refers to alkyl-S-, where "alkyl" is as defined above. In some embodiments, the alkyl component of the alkylthio group will include C₁-C₁₀, C₁-C₈, C₁-C₆, C₁-C₄ or C₁-C₃ alkyl groups. For example, C₁-C₄-alkylthio include, but are not limited to, methylthio, ethylthio, propylthio, 1-methylethylthio, butylthio, 1-methylpropylthio, 2-methylpropylthio or 1,1-dimethylethylthio.

Similarly, the terms "haloalkylthio," "cycloalkylthio", "halocycloalkylthio" refer to the groups -S-haloalkyl, -S-cycloalkyl, and -S-halocycloalkyl, respectively, where the terms "haloalkyl," "cycloalkyl," and "halocycloalkyl" are as defined above.

The term "alkylsulfinyl" refers to the group alkyl-S(=O)-, where "alkyl" is as defined above. In some embodiments, the alkyl component in alkylsulfinyl groups will include C₁-C₁₂, C₁-C₁₀, C₁-C₈, C₁-C₆, C₁-C₄ or C₁-C₃ alkyl groups. Examples include, but are not limited to, -SO-CH₃, -SO-C₂H₅, n-propylsulfinyl, 1-methylethylsulfinyl, n-butylsulfinyl, 1-methylpropylsulfinyl, 2-methylpropylsulfinyl, 1,1-dimethylethylsulfinyl, n-pentylsulfinyl, 1-methylbutylsulfinyl, 2-methylbutylsulfinyl, 3-methylbutylsulfinyl, 1,1-dimethylpropylsulfinyl, 1,2-dimethylpropylsulfinyl, 2,2-dimethylpropylsulfinyl, 1-ethylpropylsulfinyl, n-hexylsulfinyl, 1-methylpentylsulfinyl, 2-methylpentylsulfinyl, 3-methylpentylsulfinyl, 4-methylpentylsulfinyl,

1,1-dimethylbutylsulfinyl, 1,2-dimethylbutylsulfinyl, 1,3-dimethylbutylsulfinyl, 2,2-dimethylbutylsulfinyl, 2,3-dimethylbutylsulfinyl, 3,3-dimethylbutylsulfinyl, 1-ethylbutylsulfinyl, 2-ethylbutylsulfinyl, 1,1,2-trimethylpropylsulfinyl, 1,2,2-trimethylpropylsulfinyl, 1-ethyl-1-methylpropylsulfinyl or 1-ethyl-2-methylpropylsulfinyl.

Similarly, the terms “alkenylsulfinyl,” “alkynylsulfinyl,” “haloalkylsulfinyl,” “haloalkenylsulfinyl,” and “haloalkynylsulfinyl” refer to the groups alkenyl-S(=O)-, alkynyl-S(=O)-, and haloalkyl-S(=O)-, haloalkenyl-S(=O)-, and haloalkynyl-S(=O)-, where the terms “alkenyl,” “alkynyl,” “haloalkyl,” “haloalkenyl,” and “haloalkynyl” are as defined above.

The term “alkylsulfonyl” refers to the group alkyl-S(=O)₂-, where the term “alkyl” is as defined above. In some embodiments, the alkyl component in alkylsulfonyl groups will include C₁-C₁₂, C₁-C₁₀, C₁-C₈, C₁-C₆ or C₁-C₄ alkyl groups. Examples include, but are not limited to, -SO₂-CH₃, -SO₂-C₂H₅, *n*-propylsulfonyl, -SO₂-CH(CH₃)₂, *n*-butylsulfonyl, 1-methylpropylsulfonyl, 2-methylpropylsulfonyl, -SO₂-C(CH₃)₃, *n*-pentylsulfonyl, 1-methylbutylsulfonyl, 2-methylbutylsulfonyl, 3-methylbutylsulfonyl, 1,1-dimethylpropylsulfonyl, 1,2-dimethylpropylsulfonyl, 2,2-dimethylpropylsulfonyl, 1-ethylpropylsulfonyl, *n*-hexylsulfonyl, 1-methylpentylsulfonyl, 2-methylpentylsulfonyl, 3-methylpentylsulfonyl, 4-methylpentylsulfonyl, 1,1-dimethylbutylsulfonyl, 1,2-dimethylbutylsulfonyl, 1,3-dimethylbutylsulfonyl, 2,2-dimethylbutylsulfonyl, 2,3-dimethylbutylsulfonyl, 3,3-dimethylbutylsulfonyl, 1-ethylbutylsulfonyl, 2-ethylbutylsulfonyl, 1,1,2-trimethylpropylsulfonyl, 1,2,2-trimethylpropylsulfonyl, 1-ethyl-1-methylpropylsulfonyl or 1-ethyl-2-methylpropylsulfonyl and the like.


The terms “alkenylsulfonyl,” “alkynylsulfonyl,” “haloalkylsulfonyl,” “haloalkenylsulfonyl,” and “haloalkynylsulfonyl” refer to the groups alkenyl-S(=O)₂-, alkynyl-S(=O)₂-, and haloalkyl-S(=O)₂-, haloalkenyl-S(=O)₂-, and haloalkynyl-S(=O)₂-, where the terms “alkenyl,” “alkynyl,” “haloalkyl,” “haloalkenyl,” and “haloalkynyl” are as defined above.

The terms “alkylamino,” “dialkylamino,” “alkenylamino,” “alkynylamino,” “di(alkenyl)amino,” and “di(alkynyl)amino” refer to the groups -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), -N(alkenyl)₂ and -N(alkynyl)₂, where the terms “alkyl,” “alkenyl,” and “alkynyl” are as defined above. In some embodiments, the alkyl component in alkylamino or dialkylamino groups will include C₁-C₁₂, C₁-C₁₀, C₁-C₈, C₁-C₆ or C₁-C₄ alkyl groups.

The terms “alkylcarbonyl”, “alkoxycarbonyl”, “alkylaminocarbonyl”, and “dialkylaminocarbonyl” refer to alkyl-C(O)-, alkoxy-C(O)-, alkylamino-C(O)- and dialkylamino-C(O)- where alkyl, alkoxy, alkylamino and dialkylamino are as defined above. Similarly, the terms “haloalkylcarbonyl”, “haloalkoxycarbonyl”, “haloalkylaminocarbonyl”, and “dihaloalkylaminocarbonyl” refer to the groups haloalkyl-C(O)-, haloalkoxy-C(O)-, haloalkylamino-C(O)- and dihaloalkylamino-C(O)- where haloalkyl, haloalkoxy, haloalkylamino and dihaloalkylamino are as defined above.

DETAILED DESCRIPTION:








An embodiment of the present invention includes a compound of Formula (I):




(I)

wherein:

L is L1, L2, L3, L4, L5, L6, L7, L8, L9, L10, L11, L12, L13, L14 or L15:





R' is hydrogen, optionally substituted alkyl, optionally substituted cycloalkyl, or optionally substituted aryl;

R¹ is hydrogen, cyano, halo, hydroxyl, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, optionally substituted alkoxyalkyl, optionally substituted aminoalkyl, optionally substituted alkylaminoalkyl, optionally substituted dialkylaminoalkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heteroaryl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted cycloalkoxy, optionally substituted heterocyclyl, optionally substituted alkylcarbonyl, optionally substituted alkoxy carbonyl, aminocarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, --SO_p (optionally substituted alkyl or haloalkyl), --SF_5 , or $\text{--NR}^a\text{R}^b$, wherein R^a and R^b are independently H or optionally substituted alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-

heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

R^2 is hydrogen, cyano, halo, hydroxyl, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted alkoxyalkyl, optionally substituted aminoalkyl, optionally substituted alkylaminoalkyl, optionally substituted dialkylaminoalkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl; optionally substituted aryloxy, optionally substituted heteroaryl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted cycloalkoxy, optionally substituted heterocyclyl, optionally substituted alkylcarbonyl, optionally substituted alkoxy carbonyl, optionally substituted aminocarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, $-SO_p$ (optionally substituted alkyl or haloalkyl), $-SF_5$, or $-NR^aR^b$, wherein R^a and R^b are independently H or optionally substituted alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

R^3 is hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted alkylcarbonyl, optionally substituted alkoxy carbonyl, aminocarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, $-S(O)_p$ (optionally substituted alkyl), $-SF_5$, optionally substituted heterocyclyl, optionally substituted 6- to 10-membered aryl, optionally substituted 5- to 10-membered heteroaryl, a spirocyclic heterocyclyl-carbocyclyl group, a spirocyclic heterocyclyl-heterocyclyl group, a spirocyclic carbocyclyl-carbocyclyl group, a spirocyclic carbocyclyl-heterocyclyl group or $-NR^aR^b$, wherein R^a and R^b are independently H or optionally substituted alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

R^4 and $R^{4'}$ are independently in each occurrence, hydrogen, halogen, cyano, nitro, hydroxyl, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted alkoxyalkyl, optionally substituted aminoalkyl, optionally substituted alkylaminoalkyl, optionally substituted dialkylaminoalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkoxy, optionally substituted alkylcarbonyl, optionally substituted alkoxy carbonyl, optionally substituted aminocarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted di(alkyl)aminocarbonyl, optionally substituted alkylcarbonyloxy, optionally substituted alkylcarbonylamino, optionally substituted aryl, optionally substituted heteroaryl, $-SF_5$, $-SO_p$ (optionally substituted alkyl or haloalkyl); or R^4 together with $R^{4'}$ together form a 2-6-membered chain optionally containing one or two heteroatoms selected from the group consisting of N, O, Si and S to form carbocyclic or heterocyclic ring together with the carbon atom to which they are attached; or $-NR^cR^d$, wherein R^c and R^d are independently H or optionally substituted alkyl; or R^c and R^d may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

R^8 is hydrogen, halogen, alkyl, haloalkyl, cycloalkyl, alkenyl or alkynyl;

R^9 and $R^{9'}$ are independently hydrogen, halo, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy or cycloalkoxy, or R^9 together with $R^{9'}$ form a 2-6-membered chain optionally containing one or two heteroatoms selected from the group consisting of N, O, Si and S to form carbocyclic or heterocyclic ring together with the carbon atom to which they are attached, wherein the carbon or nitrogen atoms in the chain may be optionally substituted;

Q is $C-R^8$ or N;

X is O, S or $N-R^{9'}$;

Y^1 and Y^6 are each independently N, C, or $-CR^4$ -;

Y^2 , Y^3 , Y^4 and Y^5 are each independently N, $NR^{9'}$, S, O, $-CR^4$ - or $CR^4R^{4'}$;

W is CR⁵R⁶, O, SO_p, or N-R⁷,

Z is CR⁵R⁶, O, SO_p, or N-R⁷,

wherein

5 R⁵ and R⁶ are independently in each occurrence hydrogen, halo, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or cycloalkoxy, or R⁵ together with R⁶ form a 2-6-membered chain optionally containing one or two heteroatoms selected from the group consisting of N, O, Si and S to form carbocyclic or heterocyclic ring together with the carbon atom to which they are attached, and wherein each carbon or nitrogen in said carbocyclic or heterocyclic ring may be optionally substituted;

10 R⁷ is hydrogen or C₁-C₄-alkyl; and

wherein at most three of Y¹, Y², Y³, Y⁴, Y⁵ and Y⁶ are heteroatoms;

a is 0 or 1;

q is 0 or 1;

15 p is independently in each occurrence is 0, 1, or 2; and

the dashed bonds (---) signifies a single or double bond;

a stereoisomer, tautomer, N-oxide, hydrate, solvate, or salt thereof.

In another embodiment, the invention provides a compound of Formula (I)

wherein:

20 R' is hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, optionally substituted C₃-C₈-cycloalkyl, or optionally substituted phenyl;

R¹ is hydrogen, cyano, halo, hydroxyl, C₁-C₆-alkyl, C₁-C₆-haloalkyl, hydroxy-C₁-C₆-alkyl, hydroxy-C₁-C₆-haloalkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-haloalkoxy-C₁-C₆-alkyl, amino-C₁-C₆-alkyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-haloalkenyloxy, C₂-C₆-

alkynyloxy, C₂-C₆-haloalkynyloxy, C₁-C₆-haloalkoxy, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-haloalkylaminocarbonyl, di-C₁-C₆-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heteroaryl, optionally substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkenyl, optionally substituted C₃-C₈-cycloalkyloxy, optionally substituted 3- to 7-membered heterocyclyl, -SF₅, -SO_p(optionally substituted C₁-C₆-alkyl or C₁-C₆-haloalkyl), or -NR^aR^b, wherein R^a and R^b are independently H or optionally substituted C₁-C₆-alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

R² is hydrogen, cyano, halo, hydroxyl, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-haloalkoxy-C₁-C₆-alkyl, optionally substituted phenyl; optionally substituted phenyloxy, optionally substituted 5- or 6-membered heteroaryl, optionally substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkenyl, optionally substituted C₃-C₈-cycloalkyloxy, optionally substituted 3- to 7-membered heterocyclyl containing from one to three heteroatoms selected from the group consisting of N, O and S, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-haloalkylaminocarbonyl, di-C₁-C₆-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, -SO_p(optionally substituted C₁-C₆-alkyl or C₁-C₆-haloalkyl), SF₅, or -NR^aR^b, wherein R^a and R^b are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

R³ is C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-haloalkoxy-C₁-C₆-alkyl, optionally

substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkenyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-haloalkylaminocarbonyl, di-C₁-C₆-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, -SF₅, -S(O)_p(C₁-C₆-alkyl or C₁-C₆-haloalkyl), optionally substituted 3- to 7-membered heterocyclyl containing from one to three heteroatoms selected from the group consisting of N, O and S; optionally substituted phenyl, optionally substituted 5- to 10-membered heteroaryl, a 5- to 11-membered spirocyclic heterocyclyl-carbocyclyl group, a 5- to 11-membered spirocyclic heterocyclyl-heterocyclyl group, a 5- to 11-membered spirocyclic carbocyclyl-carbocyclyl group, a 5- to 11-membered spirocyclic carbocyclyl-heterocyclyl group, or -NR^aR^b, wherein R^a and R^b are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

15 R⁴ and R^{4'} are independently in each occurrence, hydrogen, halogen, cyano, nitro, hydroxyl, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-haloalkoxy-C₁-C₆-alkyl, optionally substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkyloxy, optionally substituted C₁-C₆-alkylcarbonyl, optionally substituted C₁-C₆-alkoxycarbonyl, optionally substituted aminocarbonyl, C₁-C₆-alkylaminocarbonyl, di(C₁-C₆-alkyl)aminocarbonyl, optionally substituted C₁-C₆-alkylcarbonyloxy, optionally substituted C₁-C₆-alkylcarbonylamino, optionally substituted phenyl, optionally substituted 5- or 6-membered heteroaryl, -SF₅, -SO_p(optionally substituted C₁-C₆-alkyl or C₁-C₆-haloalkyl); or R⁴ together with R^{4'} together form a 2-6-membered chain optionally containing one or two heteroatoms selected from the group consisting of N, O, Si and S to form carbocyclic or heterocyclic ring together with the carbon atom to which they are attached; or -NR^cR^d, wherein R^c and R^d are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^c and R^d may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional

heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

R⁸ is hydrogen, halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₈-cycloalkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl; and

5 L, Q, X, Y¹, Y², Y³, Y⁴, Y⁵, Y⁶, W, Z, R⁵, R⁶, R⁷, R⁹, R^{9'}, a, q, p and the dashed bonds (-----) are as defined above for the compound of Formula (I).

In one embodiment, L is L1. In another embodiment, L is L2. In another embodiment, L is L3. In another embodiment, L is L4. In another embodiment, L is L5. In another embodiment, L is L6. In another embodiment, L is L7. In another embodiment, L is L8. In another embodiment, L is L9. In another embodiment, L is L10. In another embodiment, L is L11. In another embodiment, L is L12. In another embodiment, L is L13. In another embodiment, L is L14. In another embodiment, L is L15.

In some embodiments:

R¹ is hydrogen, cyano, optionally substituted C₁-C₄-alkyl, optionally substituted C₁-C₄-alkoxy, optionally substituted C₁-C₄-alkenyl, optionally substituted C₁-C₄-alkynyl, optionally substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkenyl, optionally substituted, saturated or partially unsaturated 5-, 6-, or 7-membered heterocycle group, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted aryloxy, optionally substituted C₁-C₄-alkylcarbonyl, optionally substituted C₁-C₄-alkoxycarbonyl, optionally substituted aminocarbonyl, optionally substituted C₁-C₄-alkylaminocarbonyl, optionally substituted C₁-C₄-dialkylaminocarbonyl, optionally substituted alkyl-SO_p-, haloalkyl-SO_p-, amino, -NH-optionally substituted C₁-C₄-alkyl, or -NR^aR^b, wherein R^a and R^b are independently optionally substituted alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O and S and may be optionally substituted;

R' is hydrogen or C₁-C₄-alkyl;

R^2 is hydrogen, halogen, cyano, nitro, -OH, optionally substituted C₁-C₄-alkyl, optionally substituted C₁-C₄-alkoxy, optionally substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkenyl, -amino, NH-optionally substituted C₁-C₄-alkyl, -SF₅, or -NR^aR^b, wherein R^c and R^d are independently optionally substituted C₁-C₄-alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered heterocyclyl group, which may be optionally substituted, SO_p(optionally substituted C₁-C₄-alkyl or haloalkyl);

R^3 is C₁-C₄-alkyl, C₃-C₆-cycloalkyl, optionally substituted C₅-C₇-cycloalkenyl, 4- to 6-membered-heterocyclyl, 6- to 10-membered aryl, 5- to 10-membered heteroaryl, each of which may be optionally substituted with 1, 2, or 3 substituents;

R^4 and $R^{4'}$ are independently hydrogen, halogen, cyano, nitro, -OH, optionally substituted C₁-C₄-alkyl, optionally substituted C₁-C₄-alkoxy, optionally substituted C₃-C₈-cycloalkyl, -amino, NH-optionally substituted C₁-C₄-alkyl, -SF₅; or R^4 together with $R^{4'}$ together form a 2-6-membered chain optionally containing one or two heteroatoms selected from the group consisting of O, Si and S, or containing the group NR', to form carbocyclic or heterocyclic ring together with the carbon atom to which they are attached; or -NR^cR^d, wherein R^c and R^d are independently optionally substituted C₁-C₄-alkyl; or R^c and R^d may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered heterocyclyl group, which may be optionally substituted, SO_p(optionally substituted C₁-C₄-alkyl or haloalkyl).

In some embodiments, R^1 is hydrogen.

In some embodiments, R^1 is C₁-C₄-alkyl, C₁-C₄-haloalkyl, amino, C₁-C₄-alkylamino, or di-(C₁-C₄-alkyl) amino.

In another embodiment, R^1 is halogen.

In another embodiment, R^1 is C₁-C₄-alkyl-SO_p-, C₁-C₄-haloalkyl-SO_p- or -SF₅.

In other embodiments, R^1 is hydroxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkoxy-C₁-C₄-alkyl or C₁-C₄-haloalkoxy-C₁-C₄-haloalkyl.

In another embodiment, R^1 is methyl, ethyl, propyl, butyl, pentyl, isopropyl (*i*-Pr), tert-butyl (*t*-butyl), prop-1-en-2-yl, 2-fluoroprop-2-yl, 1,1-difluoroethyl or 2-hydroxyprop-2-yl.

In another embodiment, R¹ is C₁-C₃-alkoxy or C₁-C₃-haloalkoxy.

In another embodiment, R¹ is OCH₃ or OCH₂CH₃.

In another embodiment, R¹ is OCF₃ or SCF₃.

In another embodiment, R¹ is CF₃, -CH₂CF₃, -CHFCF₃ or -CF₂CF₃.

5 In some embodiments, R¹ is C₂-C₄-alkenyl or C₂-C₄-haloalkenyl.

In some embodiments, R¹ is optionally substituted cyclopentyl or optionally substituted cyclohexyl.

In other embodiments, R¹ is cyclopropyl or cyclobutyl.

10 In some embodiments, R¹ is an optionally substituted, saturated or unsaturated 6-membered heterocyclcyl group.

In one embodiment, R¹ is -NR^aR^b, wherein R^a and R^b are independently hydrogen or C₁-C₆ alkyl. In another embodiment R¹ is -NR^aR^b, wherein R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered heterocyclcyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O and S and may be
15 optionally substituted.

In another embodiment, R¹ is C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-haloalkylaminocarbonyl, di-C₁-C₆-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl.

20 In some embodiments, R¹ is optionally substituted tetrahydrofuryl, dihydrofuryl, morpholino, pyranlyl, dihydropyranlyl, piperidinyl, dihydropiperidinyl, dihydrothiophene, or tetrahydrothiophene.

In some embodiments, R¹ is optionally substituted phenyl.

25 In some embodiments, R¹ is aziridinyl, azetidiny, oxetanyl, pyrrolidinyl, pyrrolyl, pyrazolyl, oxetanyl, pyrazolinyl, imidazolyl, imidazoliny, imidazolidinyl, oxazolyl, oxazolidinyl, isoxazolinyl, isoxazolyl, thiazolyl, thiadiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, furyl, tetrahydrofuryl, thienyl, oxadiazolyl, piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodinyl, 2-oxoazepinyl, azepinyl, 4-piperidonyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyranlyl, dihydropyranlyl, tetrahydropyranlyl, thiopyranlyl, dihydrothiopyranlyl, tetrahydrothiopyranlyl, morpholinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl
30 sulfone, 1,3-dioxolane and tetrahydro-1,1-dioxothienyl, triazolyl or triazinyl.

In some embodiments, R¹ is aziridinyl, azetidiny, oxetanyl, pyrrolidinyl, pyrrolyl, or morpholinyl, all of which are optionally substituted by one or more halogen.

In some embodiments, R² is hydrogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, amino, C₁-C₄-alkylamino, or di-(C₁-C₄alkyl) amino.

5 In another embodiment, R² is hydrogen, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl or *tert*-butyl.

In another embodiment, R² is hydrogen, CF₃, -CH₂CF₃, -CHF₂CF₃ or -CF₂CF₃.

In some embodiments, R² is hydrogen.

In some embodiments, R² is halogen.

10 In another embodiment, R² is fluoro or chloro.

In another embodiment, R² is hydrogen, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or S(O)_p(C₁-C₄-alkyl or C₁-C₄-haloalkyl) where p is 0, 1 or 2.

In another embodiment, R² is methoxy, ethoxy, propoxy or butoxy.

In another embodiment, R² is methylthio, ethylthio, propylthio or butylthio.

15 In another embodiment, R² is -OCF₃ or -SCF₃.

In some embodiments, R² is C₁-C₄-alkenyl or C₁-C₄-haloalkenyl.

In some embodiments, R² is optionally substituted cyclopentyl or optionally substituted cyclohexyl.

20 In some embodiments, R² is an optionally substituted, saturated or unsaturated 6-membered heterocyclyl group.

In some embodiments, R² is optionally substituted tetrahydrofuryl, dihydrofuryl, morpholino, pyranyl, dihydropyranyl, piperidinyl, dihydropiperidinyl, dihydrothiophene, or tetrahydrothiophene.

In some embodiments, R² is optionally substituted phenyl.

25 In other embodiments, R² is phenyl substituted with 1, 2, or 3 substituents, which are independently halo, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkoxy, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy or haloalkenyloxy.

30 In another embodiment, R² is a 5- or 6-membered heteroaryl with 1 or 2 substituents, which are independently halo, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, alkyl, alkenyl, alkynyl,

cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy or haloalkenyloxy.

In one embodiment, R^2 is pyridinyl optionally substituted with halo, cyano, nitro, C_1 - C_3 -alkyl, C_1 - C_3 -haloalkyl, C_1 - C_3 -alkoxy, C_1 - C_3 -haloalkoxy or $(C_1$ - C_3 -alkyl or C_1 - C_3 -haloalkyl) $S(O)_p$.

5 In some embodiments, R^2 is optionally substituted aziridinyl, azetidiny, oxetanyl, pyrrolidinyl, pyrrolyl, pyrazolyl, oxetanyl, pyrazolinyl, imidazolyl, imidazolinyl, imidazolidinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, furyl, tetrahydrofuryl, thienyl, oxadiazolyl, piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodinyl, 2-oxoazepinyl, azepinyl, 4-piperidinyl,
10 pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, tetrahydropyranlyl, morpholinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, 1,3-dioxolane and tetrahydro-1,1-dioxothienyl, triazolyl or triazinyl.

In some embodiments, R^2 is aziridinyl, azetidiny, oxetanyl, pyrrolidinyl, pyrrolyl, or morpholinyl, all of which are optionally substituted by one or more halogen.

15 In some embodiments, R^3 is 6- to 10-membered aryl optionally substituted with 1, 2, 3, 4 or 5 substituents.

In some embodiments, R^3 is C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl.

In some embodiments, R^3 is methyl, ethyl, *n*-propyl, *n*-butyl, *iso*-propyl, *tert*-butyl, *sec*-butyl or *iso*-butyl.

20 In other embodiments, R^3 is CF_3 , $-CH_2CF_3$, $-CHF CF_3$ or $-CF_2CF_3$.

In some embodiments, R^3 is optionally substituted C_3 - C_8 -cycloalkyl. In yet other embodiments, R^3 is optionally substituted C_3 - C_6 -cycloalkyl. In yet other embodiments, R^3 is optionally substituted C_3 - C_8 -cycloalkenyl or C_3 - C_6 -cycloalkenyl. In some embodiments, R^3 is optionally substituted cyclopentyl or cyclohexyl. In other embodiments, R^3 is optionally
25 substituted cyclopropyl or cyclobutyl.


In one embodiment, R^3 is cyclohexyl optionally substituted by one or more halo, C_1 - C_3 -alkyl or C_1 - C_3 -haloalkyl. In another embodiment, R^3 is cyclohexyl substituted by 1 or 2 fluoro, chloro or CF_3 .

In some embodiments, R^3 is optionally substituted piperidinyl, morpholinyl, tetrahydrofuranlyl or dihydrofuranlyl. In some embodiments, R^3 is piperidinyl, morpholinyl, tetrahydrofuranlyl or dihydrofuranlyl substituted with one or more halo, C_1 - C_6 -alkyl or C_1 - C_6 -
30

haloalkyl. In another embodiment, R^3 is piperidiny, morpholinyl, tetrahydrofuranyl or dihydrofuranyl substituted with one or more methyl, chloro or fluoro.

In some embodiments, R^3 is 5- to 10-membered heteroaryl optionally substituted with 1, 2, 3, 4 or 5 substituents. In one embodiment, the 5- to 10-membered heteroaryl is pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, pyrrolyl, indolyl, quinoliny, isoquinoliny, quinazolinyl, quinoxalinyl, furanyl, thiophenyl, furyl, pyrrolyl, imidazolyl, oxazolyl, isoxazolyl, isothiazolyl, pyrazolyl benzofuranyl, benzothiophenyl, imidazopyridyl, imidazopyrimidyl or pyrrolopyrimidyl.

In other embodiments, R^3 is an optionally substituted spirocyclic heterocyclyl-carbocyclyl group, an optionally substituted spirocyclic heterocyclyl-heterocyclyl group, an optionally substituted spirocyclic carbocyclyl-carbocyclyl group or an optionally substituted spirocyclic carbocyclyl-heterocyclyl group. In other embodiments, R^3 is a 5- to 11-membered optionally substituted spirocyclic heterocyclyl-carbocyclyl group, a 5- to 11-membered optionally substituted spirocyclic heterocyclyl-heterocyclyl group, a 5- to 11-membered optionally substituted spirocyclic carbocyclyl-carbocyclyl group or a 5- to 11-membered optionally substituted spirocyclic carbocyclyl-heterocyclyl group. Non-limiting examples of spirocyclic carbocyclyl-carbocyclyl, spirocyclic carbocyclyl-heterocyclyl and spirocyclic heterocyclyl-heterocyclyl groups are shown below for illustration.



However, it will be apparent to persons skilled in the art that the second ring of the spirocyclic group may be joined at any available carbon of the first ring. It will also be understood that the first ring of the spirocyclic group may be bonded to the molecule at any available atom. Thus, the present invention includes 3-, 4-, 5-, 6- and 7-membered carbocyclic or heterocyclic rings as defined herein joined to a second 3-, 4-, 5-, 6- and 7-membered carbocyclic or heterocyclic ring at any available carbon atom of the first ring.

In some embodiments, R^3 is phenyl substituted with 1 to 4 substituents. In another embodiment, R^3 is phenyl substituted by 1 to 3 substituents. In yet another embodiment, R^3 is

phenyl substituted by 1 or 2 substituents. In some embodiments, R³ is phenyl substituted by 1, 2, 3 or 4 substituents which are independently halo, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, phenyl, substituted phenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy or haloalkenyloxy.

In some embodiments, R³ is para-substituted phenyl.

In some embodiments, R³ is meta-substituted phenyl.

In some embodiments, R³ is ortho-substituted phenyl.

In some embodiments, R³ is halophenyl.

In some embodiments, R³ is haloalkylphenyl.

In some embodiments, R³ is haloalkoxyphenyl.

In some embodiments, R³ is phenyl substituted with 2 substituents which are independently halo, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy or haloalkenyloxy.

In some embodiments, R³ is 2,3-disubstituted phenyl.

In some embodiments, R³ is 2,4-disubstituted phenyl.

In some embodiments, R³ is 2,5-disubstituted phenyl.

In some embodiments, R³ is a 2,6-disubstituted phenyl.

In some embodiments, R³ is a 3,5-disubstituted phenyl.

In other embodiments, R³ is a 3,4-disubstituted phenyl.

In other embodiments, R³ is a 3,6-disubstituted phenyl.

In some embodiments, R³ is dihalophenyl, e.g., dichloro; difluoro; or chloro, fluoro.

In some embodiments, R³ is 2,3-dihalophenyl.

In some embodiments, R³ is chlorophenyl. In another embodiment, R³ is fluorophenyl. In another embodiment, R³ is dichlorophenyl. In another embodiment, R³ is difluorophenyl. In yet another embodiment, R³ is 3,5-dichlorophenyl. In another embodiment, R³ is 3,5-difluorophenyl. In another embodiment, R³ is 2,6-dichlorophenyl. In another embodiment, R³ is 2,6-difluorophenyl.

In some embodiments, R³ is phenyl substituted with halo and haloalkyl.

In some embodiments, R³ is phenyl substituted with halo and haloalkoxy.

In some embodiments, R³ is phenyl substituted with haloalkyl and haloalkoxy.

In some embodiments, R³ is phenyl substituted with 3 substituents which are independently halo, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy or haloalkenyloxy.

5 In some embodiments, R³ is trihalophenyl, e.g., trichloro; trifluoro; or chloro, chloro, fluoro, or fluoro, fluoro, chloro.

In some embodiments, R³ is phenyl substituted with 2 halo and haloalkyl.

In some embodiments, R³ is phenyl substituted with 2 halo and haloalkoxy.

In some embodiments, R³ is phenyl substituted with 1 haloalkyl, 1 halo, and 1 haloalkoxy.

10 In some embodiments, R³ is phenyl substituted with 1 halo and 2 haloalkyl.

In some embodiments, R³ is 5-membered heteroaryl optionally substituted with 1 or 2 substituents which are independently halo, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy or haloalkenyloxy.

15 In some embodiments, R³ is 6-membered heteroaryl optionally substituted with 1 or 2 substituents which are independently halo, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy or haloalkenyloxy.

20 In some embodiments, R³ is 2-pyridyl optionally substituted with 1 or 2 substituents which are independently halo, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy or haloalkenyloxy.

25 In some embodiments, R³ is 3-pyridyl optionally substituted with 1 or 2 substituents which are independently halo, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy or haloalkenyloxy.

30 In some embodiments, R³ is 4-pyridyl optionally substituted with 1 or 2 substituents which are independently halo, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy or haloalkenyloxy.

In another embodiment, R³ is 4-pyridyl which is unsubstituted or substituted with 1 or 2 chloro or fluoro. In yet another embodiment R³ is 3-pyridyl which is unsubstituted or substituted with 1 or 2 chloro or fluoro.

In other embodiments, R³ is an optionally substituted 3- to 7-membered heterocycle. In some embodiments, R³ is optionally substituted aziridinyl, azetidiny, oxetanyl, pyrrolidinyl, pyrrolyl, pyrazolyl, oxetanyl, pyrazolinyl, imidazolyl, imidazolinyl, imidazolidinyl, oxazolyl, oxazolidinyl, isoxazoliny, isoxazolyl, thiazolyl, thiadiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, furyl, tetrahydrofuryl, thienyl, oxadiazolyl, piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodiny, 2-oxoazepiny, azepiny, 4-piperidony, pyridiny, pyraziny, pyrimidinyl, pyridazinyl, tetrahydropyranyl, morpholinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, 1,3-dioxolane and tetrahydro-1,1-dioxothienyl, triazolyl or triazinyl.

In another embodiment, R³ may be a heterocyclic, bridged bicyclic group, which may be optionally substituted.

In some embodiments, R⁴ and/or R^{4'} are hydrogen.

In some embodiments, each R⁴ and/or R^{4'} are independently hydrogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, amino, C₁-C₄-alkylamino, or di-(C₁-C₄alkyl) amino.

In another embodiment, each R⁴ and/or R^{4'} are independently hydrogen, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl or *tert*-butyl.

In another embodiment, R⁴ and/or R^{4'} are independently hydrogen, CF₃, -CH₂CF₃, -CHF₂CF₃ or -CF₂CF₃.

In some embodiments, R⁴ and/or R^{4'} are independently hydrogen or halogen.

In another embodiment, R⁴ and/or R^{4'} are independently hydrogen, fluoro or chloro.

In another embodiment, R⁴ and/or R^{4'} are independently hydrogen, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or S(O)_p(C₁-C₄-alkyl or C₁-C₄-haloalkyl), where p is 0, 1 or 2.

In another embodiment, R⁴ and/or R^{4'} are independently hydrogen, methoxy, ethoxy, propoxy or butoxy.

In another embodiment, R⁴ and/or R^{4'} are independently hydrogen, methylthio, ethylthio, propylthio or butylthio.

In another embodiment, R⁴ and/or R^{4'} are independently hydrogen, -OCF₃ or -SCF₃.

In some embodiments, R^4 and/or $R^{4'}$ are independently hydrogen, C₁-C₄-alkenyl or C₁-C₄-haloalkenyl.

In some embodiments, R^4 and/or $R^{4'}$ are independently hydrogen, C₁-C₄-alkylcarbonyl or C₁-C₄-alkoxycarbonyl.

5 In other embodiments, R^4 and/or $R^{4'}$ are independently hydrogen, C₁-C₄-alkylcarbonylamino.

In some embodiments, R^4 and/or $R^{4'}$ are independently hydrogen, optionally substituted cyclopentyl or optionally substituted cyclohexyl.

10 In some embodiments, R^4 and/or $R^{4'}$ are independently hydrogen, optionally substituted tetrahydrofuryl, dihydrofuryl, morpholino, pyranal, dihydropyranal, piperidinyl, dihydropiperidinyl, dihydrothiophene, or tetrahydrothiophene.

In some embodiments, R^4 and/or $R^{4'}$ are independently hydrogen, optionally substituted phenyl.

15 In other embodiments, R^4 and/or $R^{4'}$ are independently hydrogen, phenyl substituted with 1, 2, or 3 substituents, which are independently halo, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkoxy, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy or haloalkenyloxy.

20 In other embodiments, R^4 and/or $R^{4'}$ are independently hydrogen, a 5- or 6-membered heteroaryl with 1 or 2 substituents, which are independently halo, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, halocycloalkyl, halocycloalkenyl, alkoxy, alkenyloxy, alkynyloxy, haloalkoxy or haloalkenyloxy.

25 In some embodiments, R^4 and/or $R^{4'}$ are independently hydrogen, optionally substituted aziridinyl, azetidiny, oxetanyl, pyrrolidinyl, pyrrolyl, pyrazolyl, oxetanyl, pyrazolinyl, imidazolyl, imidazolinyl, imidazolidinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, furyl, tetrahydrofuryl, thienyl, oxadiazolyl, piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, 2-oxoazepinyl, azepinyl, 4-piperidonyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, tetrahydropyranal, morpholinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, 1,3-dioxolane and tetrahydro-1,1-dioxothienyl, triazolyl or triazinyl.

30

In some embodiments, R^4 and $R^{4'}$ are independently hydrogen, aziridinyl, azetidiny, oxetanyl, pyrrolidinyl, pyrrolyl, or morpholinyl, all of which are optionally substituted by one or more halogen.

In one embodiment, R^8 is H. In another embodiment, R^8 is C₁-C₃-alkyl or C₁-C₃-haloalkyl.

5 In one embodiment, R^9 and $R^{9'}$ are each hydrogen. In another embodiment, R^9 and $R^{9'}$ together form a 2- to 6-membered chain to form a spirocyclic ring substituent together with the carbon atom to which they are attached. In another embodiment, R^9 and $R^{9'}$ together form a 2- to 5-membered chain to form a spirocyclic ring substituent together with the carbon atom to which they are attached. In another embodiment, R^9 and $R^{9'}$ together form a 2- to 4-membered chain to
10 form a spirocyclic ring substituent together with the carbon atom to which they are attached. In another embodiment, R^9 and $R^{9'}$ together form a 2- or 3-membered chain to form a spirocyclic ring substituent together with the carbon atom to which they are attached. In another embodiment, R^9 and $R^{9'}$ together form a 2- membered chain to form a spirocyclic ring substituent together with the carbon atom to which they are attached.

15 In some embodiments, a is 0.

In some embodiments, a is 1.

In some embodiments, Q is N.

In other embodiments, Q is C- R^8 .

In some embodiments, X is O.

20 In some embodiments, X is S.

In some embodiments, X is NR' .

In some embodiments, W is CH₂.

In other embodiments, W is C(C₁-C₃-alkyl)₂ or C(C₁-C₃-haloalkyl)₂;

In other embodiments, W is C(CH₃)₂, C(C₂H₅)₂ or C(CF₃)₂

25 In some embodiments, Z is CH₂.

In some embodiments, Z is O.

In some embodiments, Z is SO_p.


In some embodiments, Z is SO₂.

In other embodiments, Z is SO.

30 In some embodiments, Z is NH.

In other embodiments, Z is N(C₁-C₃-alkyl) or N(C₁-C₃-haloalkyl)

In some embodiments, the compound of Formula (I) is the compound of Formula (I-1):




(I-1)

wherein variables L , R^1 , R^2 , R^3 , R^9 , $R^{9'}$, Y^1 , Y^3 , Y^4 , Y^5 , Y^6 , Q , W , Z and a are as defined for formula (I).

In one embodiment of Formula (I-1), W is CH_2 and Z is O . In one embodiment, Q is N . In another embodiment, Q is $C-R^8$. In another embodiment of Formula (I-1), W is CH_2 and Z is CH_2 . In another embodiment of Formula (I-1), W is CR^5R^6 wherein R^5 and R^6 are C_1 - C_3 -alkyl or C_1 - C_3 -haloalkyl and Z is O . In another embodiment of Formula (I-1), W is CR^5R^6 and Z is CR^5R^6 , wherein each R^5 and R^6 are independently C_1 - C_3 -alkyl or C_1 - C_3 -haloalkyl. In another embodiment of Formula (I-1), W is CR^5R^6 wherein R^5 and R^6 together form a 2- to 5-membered chain to form a ring and Z is O . In another embodiment, a is 0 and Z is O . In another embodiment, a is 0, Z is O and W is CH_2 .

In one embodiment of Formula (I-1), Y^3 is S . In another embodiment of Formula (I-1), Y^5 is S . In another embodiment, Y^3 is N . In another embodiment Y^5 is N . In another embodiment of Formula (I-1), Y^5 is N and Y^3 is S . In yet another embodiment of Formula (I-1), Y^5 is S and Y^3 is N . In another embodiment of Formula (I-1), Y^6 and Y^3 are each N . In another embodiment of Formula (I-1), Y^6 is N and Y^3 is N . In another embodiment, Y^1 is N and Y^5 is N .


In some embodiments, the compound of Formula (I) is the compound of formula (I-2):



(I-2)

wherein variables R^1 , R^2 , R^3 , R' , R^8 , R^9 , $R^{9'}$, Y^2 , Y^3 , Y^4 , Y^5 , X , W , Z and a are as defined for formula (I).


5 In other embodiments, the compound of Formula (I) is the compound of Formula (I-3) below:



(I-3)

10 wherein variables R^1 , R^2 , R^3 , R' , R^8 , R^9 , $R^{9'}$, Y^2 , Y^3 , Y^4 , Y^5 , X , W , Z and a are as defined for formula (I).


In other embodiments, the compound of Formula (I) is the compound of Formula (I-4):



(I-4)

15 wherein variables R^1 , R^2 , R^3 , R' , R^8 , R^9 , $R^{9'}$, Y^1 , Y^3 , Y^4 , Y^5 , Y^6 , X , W , Z and a are as defined for formula (I).


In another embodiment, the compound of Formula (I) is the compound of Formula (I-5):



(I-5)

wherein variables R^1 , R^2 , R^3 , R' , R^8 , R^9 , $R^{9'}$, Y^1 , Y^3 , Y^4 , Y^5 , Y^6 , X , W , Z and a are as defined for formula (I).


5 In some embodiments, the compound of formula (I) is the compound of formula (Ia):



(Ia)

wherein variables R^1 , R^2 , R^3 , R' , R^8 , R^9 , $R^{9'}$, W , Z , Y^2 , Y^3 , Y^4 , Y^5 and a are as defined for
10 formula (I).


In some embodiments, the compound of formula (I) is the compound of formula (Ib):



(Ib)

wherein variables R^1 , R^2 , R^3 , R' , R^8 , R^9 , $R^{9'}$, Y^2 , Y^3 , Y^4 , Y^5 , W , Z and a are as defined for formula (I).


In some embodiments, the compound of formula (I) is the compound of formula (Ic):



(Ic)

wherein variables R^1 , R^2 , R^3 , R' , R^8 , R^9 , $R^{9'}$, R^4 , W , Z and a are as defined for formula (I); and o is 0, 1, 2, 3 or 4.


In other embodiments, the compound of formula (I) is the compound of formula (Id):



(Id)

wherein variables R^1 , R^2 , R^3 , R' , R^8 , R^9 , $R^{9'}$, R^4 , W , Z and a are as defined for formula (I); and o is 0, 1, 2, 3 or 4.

In other embodiments, the compound of formula (I) is the compound of formula (Ie):




(Ie)

wherein variables R^1 , R^2 , R^3 , R^8 , R^9 , $R^{9'}$, Y^2 , Y^3 , Y^4 , Y^5 , W , Z and a are as defined for formula (I); m is 0, 1, 2, 3 or 4; and each R^{10} is cyano, halo, hydroxyl, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-haloalkoxy-C₁-C₆-alkyl, optionally substituted phenyl, optionally substituted phenyloxy, optionally substituted 5- or 6-membered heteroaryl, optionally substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkyloxy, optionally substituted 3- to 7-membered heterocyclyl containing from one to three heteroatoms selected from the group consisting of N, O, Si and S, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-haloalkylaminocarbonyl, di-C₁-C₆-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, -SO_p(optionally substituted C₁-C₆-alkyl or C₁-C₆-haloalkyl), where p is 0, 1 or 2, SF₅, or -NR^aR^b, wherein R^a and R^b are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted.

In one embodiment of formula (Ie), R^{10} is halo. In another embodiment, R^{10} is chloro. In yet another embodiment, R^{10} is fluoro. In another embodiment, R^{10} is chloro or fluoro and m is 1, 2 or 3. In yet another embodiment, R^{10} is fluoro and m is 2. In another embodiment, R^{10} is chloro and m is 2. In another embodiment, R^{10} is fluoro or chloro, m is 2 and the fluoro or chloro are substituted are the 3- and 5-positions of the phenyl ring. In another embodiment, R^{10} is fluoro or chloro, m is 2 and the fluoro or chloro are substituted at the 2- and 6-positions.

In other embodiments, the compound of formula (I) is the compound of formula (If):



(If)

wherein variables R^1 , R^2 , R' , R^8 , R^9 , $R^{9'}$, R^4 , $R^{4'}$, Y^2 , Y^3 , Y^4 , Y^5 , W , Z and a are as defined
 5 for formula (I); R^{10} and m are as defined for formula (Ie); b is 0 or 1; the dashed bond (---) signifies a single or double bond; D is N , SiR^{11} , where R^{11} is C_1 - C_6 alkyl or C_1 - C_6 haloalkyl, C or $C-R^4$; D^1 is N , O , $SiR^{11}R^{12}$, where R^{11} and R^{12} are independently C_1 - C_6 alkyl or C_1 - C_6 haloalkyl, $-CR^4R^{4'}$, $S(O)_p$, where p is 0, 1 or 2, or D^1 is $CR^4R^{4'}$, wherein R^4 and $R^{4'}$ together form a 2- to 5-membered chain optionally substituted with one heteroatom in the chain to form a spirocyclic
 10 group.

In some embodiments, the present invention provides compounds of formula (If), wherein the dashed bond is a single bond.

In some embodiments, the present invention provides compounds of formula (If), wherein the dashed bond is a double bond.

15 In some embodiments, the present invention provides compounds of formula (If), wherein D is CH , C -halo or N .

In some embodiments, the present invention provides compounds of formula (If), wherein D is C , CH , $C-F$ or N .

In some embodiments, the present invention provides compounds of formula (If), wherein
 20 D^1 is $CR^4R^{4'}$, wherein R^4 and $R^{4'}$ together form a 2- to 5-membered chain optionally with one heteroatom in the chain to form a spirocyclic group.

In some embodiments, the present invention provides compounds of formula (If), wherein D^1 is CH_2 , independently C -(halo)₂, $CH(C_1$ - C_3 -alkyl) or $CH(C_1$ - C_3 -haloalkyl).

In some embodiments, the present invention provides compounds of formula (If), wherein D^1 is CH_2 , independently CF_2 , $CH(CH_3)$ or $CH(CF_3)$.

In some embodiments, the present invention provides compounds of formula (If), wherein D^1 is O, S, $S(O)$ or $S(O)_2$.

5 In some embodiments, the present invention provides compounds of formula (If), wherein D is CH or C-halo; and D^1 is CH_2 .

In some embodiments, the present invention provides compounds of formula (If), wherein D is N; and D^1 is CH_2 , O or S.

10 In another embodiment, the present invention provides compounds of formula (If), wherein D is N and D^1 is $SiR^{11}R^{12}$. In another embodiment of formula (If), D is CH_2 and D^1 is $SiR^{11}R^{12}$. In yet another embodiment, D is N and D^1 is $Si(CH_3)_2$.

In some embodiments, the present invention provides compounds of formula (If), wherein the dashed line is a double bond; D is C; and D^1 is CH_2 , CF_2 , O or S.


15 In some embodiments, the present invention provides compounds of formula (If), wherein D is N; and D^1 is $CR^4R^{4'}$, wherein R^4 and $R^{4'}$ together form a 2- to 4-membered chain optionally with one oxygen in the chain to form a spirocyclic group.

In some embodiments, the present invention provides compounds of formula (If), wherein D is CH; and D^1 is $CR^4R^{4'}$, wherein R^4 and $R^{4'}$ together form a 2- to 4-membered chain optionally with one oxygen in the chain to form a spirocyclic group.

20 In some embodiments, the present invention provides compounds of formula (If), wherein D is C and the dashed bond signify a double bond; and D^1 is $CR^4R^{4'}$, wherein R^4 and $R^{4'}$ together form a 2- to 4-membered chain optionally with one oxygen in the chain to form a spirocyclic group.

25 It will be appreciated by persons of skill in the art that in Formulae (Ic) and (Id) above, where variable R^4 is indicated to be present as substituents on the aromatic rings (e.g. $(R^4)_o$ groups, where o is 0, 1, 2, 3 or 4), they will represent non-hydrogen substituents since in embodiments where o is 0, R^4 will not be present. The same principle applies to variable R^{10} in the compounds of formula (Ie) and (If).

30 In other embodiments, the invention provides compounds of formula (Ia), wherein variables R^1 , R^2 , R^3 , R^5 , R^6 , R^7 , R^8 and a are as defined for formula (I) above, and Y^2 , Y^3 , Y^4 and Y^5 are as shown in Table 1:




(Ia)

Table 1

Formula	Y ²	Y ³	Y ⁴	Y ⁵
Ia-1	CR ⁴	CR ⁴	CR ⁴	CR ⁴
Ia-2	N	CR ⁴	CR ⁴	CR ⁴
Ia-3	CR ⁴	N	CR ⁴	CR ⁴
Ia-4	CR ⁴	CR ⁴	N	CR ⁴
Ia-5	CR ⁴	CR ⁴	CR ⁴	N
Ia-6	N	N	CR ⁴	CR ⁴
Ia-7	CR ⁴	N	N	CR ⁴
Ia-8	CR ⁴	CR ⁴	N	N
Ia-9	N	CR ⁴	N	CR ⁴
Ia-10	CR ⁴	N	CR ⁴	N
Ia-11	N	CR ⁴	CR ⁴	N

5 In other embodiments, the invention provides compounds of formula (Ib), wherein variables R¹, R², R³, R⁴, R¹, R⁹, R^{9'}, W, Z, R⁸ and a are as defined for formula (I) above, and Y², Y³, Y⁴ and Y⁵ are as shown in Table 2:



(Ib)

Table 2

Formula	Y ²	Y ³	Y ⁴	Y ⁵
Ib-1	CR ⁴	CR ⁴	CR ⁴	CR ⁴
Ib-2	N	CR ⁴	CR ⁴	CR ⁴
Ib-3	CR ⁴	N	CR ⁴	CR ⁴
Ib-4	CR ⁴	CR ⁴	N	CR ⁴
Ib-5	CR ⁴	CR ⁴	CR ⁴	N
Ib-6	N	N	CR ⁴	CR ⁴
Ib-7	CR ⁴	N	N	CR ⁴
Ib-8	CR ⁴	CR ⁴	N	N
Ib-9	N	CR ⁴	N	CR ⁴
Ib-10	CR ⁴	N	CR ⁴	N
Ib-11	N	CR ⁴	CR ⁴	N

5 In some embodiment, the present invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If), wherein each R² is independently H, halo, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or S(O)_p(C₁-C₄-alkyl or C₁-C₄-haloalkyl).

10 In some embodiment, the present invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If), wherein each R² is independently H, chloro, fluoro, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl or *tert*-butyl.

In some embodiment, the present invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If), wherein each R² is independently H, CF₃, -CH₂CF₃, -CHF₂CF₃ or -CF₂CF₃.

In some embodiment, the present invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If), wherein each R^2 is independently H, methoxy, ethoxy, propoxy or butoxy.

5 In some embodiment, the present invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If), wherein each R^2 is independently H, $-OCF_3$ or $-SCF_3$.

10 In some embodiment, the present invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If), wherein each R^4 and/or $R^{4'}$ are independently H, halo, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy or $S(O)_p(C_1$ - C_4 -alkyl or C_1 - C_4 -haloalkyl).

In some embodiment, the present invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If), wherein each R^4 and/or $R^{4'}$ are independently H, chloro, fluoro, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl or *tert*-butyl.

15 In some embodiment, the present invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If), wherein each R^4 and/or $R^{4'}$ are independently H, CF_3 , $-CH_2CF_3$, $-CHF CF_3$ or $-CF_2CF_3$.

In some embodiment, the present invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If) wherein each R^4 and/or $R^{4'}$ are independently H, methoxy, ethoxy, propoxy or butoxy.

20 In some embodiment, the present invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If), wherein each R^4 and/or $R^{4'}$ are independently H, $-OCF_3$ or $-SCF_3$.

In other embodiments, the invention provides compounds of formulae (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If), wherein R^7 and R^8 are independently H or C_1 - C_3 -alkyl.

25 In other embodiments, the invention provides compounds of formulae (Ia) to (If), wherein a is 1, W is CH_2 and Z is O.

30 In other embodiments, the invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If), wherein R^1 is C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, hydroxy- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, C_1 - C_6 -haloalkoxy- C_1 - C_6 -alkyl, amino- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy, C_2 - C_6 -alkenyl, C_2 - C_6 -haloalkenyl, optionally substituted C_3 - C_8 -cycloalkyl, optionally substituted 3- to 7-membered heterocyclyl, or $-NR^aR^b$, wherein R^a and R^b

are independently H or optionally substituted C₁-C₆-alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5- or 6-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O and S and may be optionally substituted.

5 In other embodiments, the invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic) or (Id), wherein R³ is C₁-C₆-alkyl, C₁-C₆-haloalkyl, optionally substituted C₃-C₈- cycloalkyl, optionally substituted 3- to 7-membered heterocyclyl containing from one to three heteroatoms selected from the group consisting of N, O and S; optionally substituted phenyl, optionally substituted 5- to 10-membered heteroaryl, a 5- to 11-membered spirocyclic
10 heterocyclyl-carbocyclyl group, a 5- to 11-membered spirocyclic heterocyclyl-heterocyclyl group, a 5- to 11-membered spirocyclic carbocyclyl-carbocyclyl group, a 5- to 11-membered spirocyclic carbocyclyl-heterocyclyl group, or -NR^aR^b, wherein R^a and R^b are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5- or 6-membered-heterocyclyl group, which may include one to three additional heteroatoms
15 selected from the group consisting of N, O and S and may be optionally substituted.

In other embodiments, the invention provides compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic) or (Id) above, wherein R³ is optionally substituted phenyl. In another embodiment compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic) or (Id) are provided, wherein R³ is phenyl substituted with one or more halogen. In yet another embodiment,
20 compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic) or (Id) are provided, wherein R³ is phenyl substituted with 1 halogen. In another embodiment, compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic) or (Id) are provided, wherein R³ is phenyl substituted with two halogen. In yet another embodiment, compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic) or (Id) are provided, wherein R³ is phenyl substituted with three or four halogen.

25 In another embodiment, compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic) or (Id) are provided, wherein R³ is phenyl substituted with one or more chloro or fluoro. In yet another embodiment, compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic) or (Id) are provided, wherein R³ is phenyl substituted with 1 chloro or fluoro. In another embodiment, compounds of formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic) or (Id) are provided, wherein
30 R³ is phenyl substituted with two chloro or fluoro. In yet another embodiment, compounds of

formulae (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic) or (Id) are provided, wherein R³ is phenyl substituted with three or four chloro or fluoro.

In other embodiments, the invention provides compounds of formulae (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) or (If), wherein R⁷ and R⁸ are independently H or C₁-C₃-alkyl; W is CH₂, Z is O and a is 1.

In other embodiments of formulae (I), (I-2), (I-3), (Ia), (Ib) and (Ie) each of Y², Y³, Y⁴, Y⁵ are CH.

In other embodiments of formulae (I), (I-2), (I-3), (Ia), (Ib) and (Ie) each of Y², Y³, Y⁴, Y⁵ are each independently CH or CR⁴, where R⁴ is a non-hydrogen substituent.

In other embodiments of formulae (I-1), (I-4) and (I-5) each of Y³, Y⁴ and Y⁵ are CH.

In other embodiments of formulae (I), (I-2), (I-3), (Ia), (Ib) and (Ie) each of Y², Y³, Y⁴, Y⁵ are independently CH or C-halogen.


In any of the embodiments of formulae (I), (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) and (If) above, a is 1, W is -CH₂-, and Z is O.

In any of the embodiments of formulae (I), (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) and (If) above, R¹ is C₁-C₄-alkyl, C₁-C₄ alkenyl, C₁-C₄-cycloalkyl, amino, C₁-C₄-alkylamino, di(C₁-C₄-alkyl)amino, morpholino, pyranyl, tetrahydropyranyl, or dihydropyranyl.


In any of the embodiments of formulae (I), (I-1), (I-2), (I-3), (I-4), (I-5), (Ia), (Ib), (Ic), (Id), (Ie) and (If) above, an R⁴ is, independently of other R⁴, a halogen, cyano, C₁-C₄-alkyl, C₁-C₄ haloalkyl, C₁-C₄-cycloalkyl, amino, C₁-C₄-alkylamino, di(C₁-C₄-alkyl)amino, or phenyl optionally substituted 1 or 2 times by halo or C₁-C₄-alkyl.

In other embodiments, the present invention includes the compounds of formula (I), wherein the group:


In other embodiments, the present invention provides compounds of formulae (I) shown in Table 3 below, wherein L, R¹, R² and R³ are defined in the table, X is O, R⁷ is hydrogen, and wherein the group




is one of the following Ring Systems:




Ring System A;




Ring System B;




Ring System C;




Ring System D;




Ring System E;




Ring System F;




Ring System G;




Ring System H;




Ring System I;




Ring System J;








Ring System K;




Ring System L;




Ring System M;







Ring System S;




Ring System T;




Ring System U;




Ring System V;




Ring System W;




Ring System X;




Ring System Y;




Ring System Z;




Ring System AA;




Ring System AB;




Ring System AC;




Ring System AD;




Ring System AE;




Ring System AF;




Ring System AG;




Ring System AH;




Ring System AJ;




Ring System AK;




Ring System AL;




Ring System AM;




Ring System AN;




Ring System AO;




Ring System AP;




Ring System AQ;




Ring System AR;









Ring System AS;




Ring System AT;





Ring System AU;




In Table 3, “Me” represents methyl, the expression “3,5-di-F-Ph” represents the 3,5-difluorophenyl group; “3,5-di-Cl-Ph” represents 3,5-dichlorophenyl; “2,3,5-tri-F-Ph” represents 2,3,5-trifluorophenyl; “3-F-Ph” represents 3-fluorophenyl; “2,6-di-F-Ph” represents 2,6-difluorophenyl; 2,6-di-Cl-Ph” represents 2,6-dichlorophenyl; “2,4-di-F-Ph” represents 2,4-difluorophenyl; “4-F-Ph” represents 4-fluorophenyl; “3-Cl-4-F-Ph” represents 3-chloro-4-fluorophenyl; “3-Cl-Ph” represents 3-chlorophenyl; “2,3-di-F-Ph” represents 2,3-difluorophenyl; and so on;

prop-1-en-2-yl represents the group  ;

2-F-prop-2-yl represents the group 






1,1-difluoroethyl represents the group  .







Formula (I)


Table 3



Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
271	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	A	596 [M+H] ⁺

Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
326	L1	<i>i</i> -Pr	Me		A	502 [M+H] ⁺
327	L1	<i>i</i> -Pr	Me		A	500 [M+H] ⁺
326-0	L1	<i>i</i> -Pr	Me		A	502 [M+H] ⁺
324	L1	<i>i</i> -Pr	Me		A	435 [M+H] ⁺
325	L1	<i>i</i> -Pr	Me		A	437 [M+H] ⁺
323	L1	<i>i</i> -Pr	Me	<i>t</i> -Bu	A	408 [M+H] ⁺
175	L1	<i>i</i> -Pr	Me	3,5-di-F-Ph	A	463 [M+H] ⁺
A407	L1	<i>i</i> -Pr	Me	2,6-di-F-Ph	A	463 [M+H] ⁺



Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
A406	L1	<i>i</i> -Pr	Me	2,6-di-Cl-Ph	A	495 [M+H] ⁺
A413	L1	<i>i</i> -Pr	Me	2,4-di-F-Ph	A	463 [M+H] ⁺
A408	L1		Me	3,5-di-Cl-Ph	A	538 [M+H] ⁺
A412	L1	<i>i</i> -Pr	Me	4-F-Ph	A	445 [M+H] ⁺
A410	L1	<i>i</i> -Pr	Me	3-Cl-4-F-Ph	A	479 [M+H] ⁺
A411	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	C	511 [M+H] ⁺
A409	L1	-N(CH ₃) ₂	Me	3,5-di-Cl-Ph	A	496 [M+H] ⁺
A414	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	D	529 [M+H] ⁺
306	L1	<i>t</i> -Bu	Me	3,5-di-Cl-Ph	A	510 [M+H] ⁺
297	L1	<i>t</i> -Bu	Me	3,5-di-Cl-Ph	B	528 [M+H] ⁺
365	L1	<i>i</i> -Pr	Me	cyclopropyl	A	391 [M+H] ⁺
371	L1	<i>i</i> -Pr	Me		A	470 [M+H] ⁺

Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
370	L1	<i>i</i> -Pr	Me		A	436 [M+H] ⁺
366	L1	<i>i</i> -Pr	Me	CN	A	376 [M+H] ⁺
369	L1	<i>i</i> -Pr	Me		A	447 [M+H] ⁺
308	L1	prop-1-en-2-yl	Me	3,5-di-Cl-Ph	B	512 [M+H] ⁺
364	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	E	470 [M+H] ⁺
352	L1	<i>i</i> -Pr	H	3,5-di-Cl-Ph	A	481 [M+H] ⁺
320-0	L1	<i>i</i> -Pr	Me	3,5-di-F-Ph	F	463 [M+H] ⁺
345	L1	<i>i</i> -Pr	Me	2,3,5-tri-F-Ph	A	481 [M+H] ⁺
344	L1	<i>i</i> -Pr	Me	2,3,5-tri-F-Ph	B	499 [M+H] ⁺
294	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	B	514 [M+H] ⁺
320	L2	<i>i</i> -Pr	Me	3,5-di-F-Ph	A	463 [M+H] ⁺
277	L1	2-F-prop-2-yl	Me	3,5-di-F-Ph	A	481 [M+H] ⁺



Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
323-0	L1	<i>i</i> -Pr	Me	-CH ₂ CH(CH ₃) ₂	A	407 [M+H] ⁺
298-0	L1	<i>t</i> -Bu	CF ₃	3-Cl-Ph	B	548 [M+H] ⁺
299-0	L1	<i>t</i> -Bu	CF ₃	3-Cl-Ph	A	530 [M+H] ⁺
299	L1	<i>t</i> -Bu	CF ₃	3,5-di-Cl-Ph	B	582 [M+H] ⁺
298	L1	<i>t</i> -Bu	CF ₃	3,5-di-Cl-Ph	A	564 [M+H] ⁺
304-0	L1	<i>i</i> -Pr	3,5-di-Cl-Ph	Cl	B	535 [M+H] ⁺
321	L1	<i>i</i> -Pr	4-F-Ph	3,5-di-F-Ph	A	544 [M+H] ⁺
322	L1	<i>i</i> -Pr	Me		A	463 [M+H] ⁺
304	L1	<i>i</i> -Pr	Cl	3,5-di-Cl-Ph	B	535 [M+H] ⁺
307	L1	prop-1-en-2-yl	Me	3,5-di-F-Ph	B	478 [M+H] ⁺
296	L1	<i>i</i> -Pr	CF ₃	3,5-di-Cl-Ph	B	568 [M+H] ⁺
295	L1	<i>i</i> -Pr	CF ₃	3,5-di-Cl-Ph	A	550 [M+H] ⁺
293	L1	<i>i</i> -Pr	Me	2,3-di-Cl-Ph	B	514 [M+H] ⁺
276	L1	<i>i</i> -Pr	CF ₃	3,5-di-F-Ph	A	517

Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
						[M+H] ⁺
274	L1	<i>i</i> -Pr	Me	2,3-di-Cl-Ph	A	496 [M+H] ⁺
273	L1	<i>i</i> -Pr	Me	3-F-Ph	A	446 [M+H] ⁺
272	L1	<i>i</i> -Pr	Me	3-Cl-5-F-Ph	A	480 [M+H] ⁺
275	L1	<i>i</i> -Pr	Me	3,5-di-F-Ph	B	481 [M+H] ⁺
279	L1	prop-1-en-2-yl	Me	3,5-di-F-Ph	A	461 [M+H] ⁺
174	L1	H	H	2,6-di-F-Ph	A	407 [M+H] ⁺
A400	L1	1,1-difluoroethyl	Me	3,5-di-Cl-Ph	A	517 [M+H] ⁺
A401	L1	CF ₃	Me	3,5-di-Cl-Ph	A	521 [M+H] ⁺
373	L1	<i>i</i> -Pr	Me		A	479 [M+H] ⁺
372-0	L1	<i>i</i> -Pr	Me		A	421 [M+H] ⁺
A402	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	G	599 [M+H] ⁺
A403	L1	<i>i</i> -Pr	-CH ₂ OH	3,5-di-Cl-Ph	A	511


Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
						[M+H] ⁺
A404	L1	<i>i</i> -Pr	-CF ₂ CF ₃	3,5-di-Cl-Ph	A	599 [M+H] ⁺
394	L1	-OCH ₃	H	3,5-di-Cl-Ph	A	470 [M+H] ⁺
398	L1	-OCH ₂ CH ₃	H	3,5-di-Cl-Ph	A	484 [M+H] ⁺
A405	L1	-CHF ₂	Me	3,5-di-Cl-Ph	A	503 [M+H] ⁺
573	L1	<i>t</i> -Bu	Cl	2,3,5-tri-F-Ph	A	515 [M+H] ⁺
559	L1	<i>i</i> -Pr	-CN	3,5-di-Cl-Ph	A	504 [M+H] ⁺
614	L1	<i>t</i> -Bu	Me	3,5-di-Cl-Ph	AAA	510 [M+H] ⁺
451	L1	-N(CH ₃) ₂	CF ₃	3,5-di-Cl-Ph	A	550 [M+H] ⁺
572	L1	<i>t</i> -Bu	Cl	2,5-di-Cl-4-F-Ph	A	547 [M+H] ⁺
528	L1	<i>i</i> -Pr	Me	2,5-di-Cl-4-F-Ph	A	513 [M+H] ⁺
571	L1	<i>t</i> -Bu	Cl	2,4,5-tri-F-Ph	A	515 [M+H] ⁺
574	L1	<i>t</i> -Bu	Cl	2,3-di-Cl-5-F-Ph	A	547 [M+H] ⁺
A415	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	H	543 [M+H] ⁺
A416	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	I	543

Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
						[M+H] ⁺
A417	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	J	578 [M+H] ⁺
A418	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	K	578 [M+H] ⁺
A419	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	L	500 [M+H] ⁺
A420	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	M	500 [M+H] ⁺
560	L1	<i>i</i> -Pr	-CHF ₂	3,5-di-Cl-Ph	A	530 [M+H] ⁺
305	L1	<i>i</i> -Pr	-CHF ₂	3,5-di-Cl-Ph	B	548 [M+H] ⁺
A421	L1		Me	2,6-di-Cl-4-F	A	556 [M+H] ⁺
A422	L1		Me	2,3,5-tri-F-Ph	A	524 [M+H] ⁺
420	L1	<i>t</i> -Bu	Cl	3,5-di-Cl-Ph	A	572 [M+H+ CH ₃ CN] ⁺
A423	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	N	557 [M+H] ⁺
A424	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	O	557 [M+H] ⁺
523	L1	<i>t</i> -Bu	Me	2,6-di-Cl-4-F-Ph	A	527

Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
						[M+H] ⁺
A425	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	P	483 [M+H] ⁺
A426	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	Q	483 [M+H] ⁺
526	L1	<i>t</i> -Bu	Me	2,3-di-Cl-5-F-Ph	A	527 [M+H] ⁺
527	L1	<i>i</i> -Pr	Me	2,4,6-tri-F-Ph	A	481 [M+H] ⁺
524	L1	<i>t</i> -Bu	Me	2,4,6-tri-F-Ph	A	495 [M+H] ⁺
525	L1	<i>t</i> -Bu	Me	2,3,5-tri-F-Ph	A	495 [M+H] ⁺
414-0	L2	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	R	496 [M+H] ⁺
514	L2	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	E	496 [M+H] ⁺
A427	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	S	487 [M+H] ⁺
A428	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	T	522 [M+H] ⁺
A429	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	U	487 [M+H] ⁺
A430	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	V	522 [M+H] ⁺
418	L1	<i>t</i> -Bu	H	3,5-di-Cl-Ph	A	495 [M+H] ⁺
513	L2	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	A	495

Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
						[M+H] ⁺
513-0	L2	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	F	496 [M+H] ⁺
511	L1		H	3,5-di-Cl-Ph	A	524 [M+H] ⁺
512	L1	-N(CH ₃) ₂	H	3,5-di-Cl-Ph	A	482 [M+H] ⁺
A431	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	W	513 [M+H] ⁺
450	L1		CF ₃	3,5-di-Cl-Ph	A	592 [M+H] ⁺
A432	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	X	520 [M+H] ⁺
A473	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	Y	520 [M+H] ⁺
A433	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	Z	554 [M+H] ⁺
A434	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AA	554 [M+H] ⁺
A435	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AB	499 [M+H] ⁺
A436	L1	L1	<i>i</i> -Pr	Me	N/A (H)	363 [M+H] ⁺
A437	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AC	499 [M+H] ⁺

Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
A438	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AD	482 [M+H] ⁺
A439	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AF	494 [M+H] ⁺
A440	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AG	494 [M+H] ⁺
A441	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AH	499 [M+H] ⁺
A442	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AE	482 [M+H] ⁺
A443	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AJ	499 [M+H] ⁺
A445	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AK	511 [M+H] ⁺
A446	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AL	511 [M+H] ⁺
A447	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AM	482 [M+H] ⁺
A448	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AO	502 [M+H] ⁺
A449	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AQ	502 [M+H] ⁺
A450	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AR	502 [M+H] ⁺
A451	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AP	502 [M+H] ⁺
A452	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AS	523 [M+H] ⁺

Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
A472	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AT	523 [M+H] ⁺
A453	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AN	482 [M+H] ⁺
A454	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AU	520 [M+H] ⁺
419	L1	<i>i</i> -Pr	Cl	3,5-di-Cl-Ph	A	558 [M+CH ₃ C N] ⁺
397-0	L1	OCH ₂ CH=CH ₂	Me	3,5-di-Cl-Ph	A	495 [M+H] ⁺
A455	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AW	482 [M+H] ⁺
A456	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AX	482 [M+H] ⁺
A457	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AY	515 [M+H] ⁺
A458	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AZ	515 [M+H] ⁺
395	L1	-OCHF ₂	Me	3,5-di-Cl-Ph	A	568 [M+CH ₃ C N+Na] ⁺
A459	L1		Me	3,5-di-Cl-Ph	A	570 [M+H] ⁺
A464	L1	-N(CH ₃) ₂	Me	2,6-di-Cl-4-F-Ph	A	514 [M+H] ⁺

Cmpd. #	L	R ¹	R ²	R ³	Ring System	ESI-MS
A462	L1	-N(CH ₃) ₂	Me	2,4,6-tri-F-Ph	A	481 [M+H] ⁺
A463	L1	-N(CH ₃) ₂	Me	2,3-di-Cl-5-F-Ph	A	514 [M+H] ⁺
A460	L1	-N(CH ₃) ₂	Me	2,3,5-tri-F-Ph	A	481 [M+H] ⁺
A461	L1	-N(CH ₃) ₂	Me	2,3,5-tri-Cl-Ph	A	531 [M+H] ⁺
558	L1	<i>i</i> -Pr	-C(O)CH ₃	3,5-di-Cl-Ph	A	523 [M+H] ⁺

For avoidance of doubt, each of the compounds presented in Table 3 has been prepared.

Stereoisomers and polymorphic forms

It will be appreciated by those of skill in the art that the compound may exist and be isolated in optically active and racemic forms. Compounds having one or more chiral centers, including that at a sulfur atom, may be present as single enantiomers or diastereomers or as mixtures of enantiomers and/or diastereomers. For example, it is well known in the art that sulfoxide compounds may be optically active and may exist as single enantiomers or racemic mixtures. In addition, compounds of the description may include one or more chiral centers, which results in a theoretical number of optically active isomers. In the present case, the compounds of formula (I) include at least one chiral center at the carbon atom bearing variable R⁸ when Q is C-R⁸. Where compounds of the description include n chiral centers, the compounds may comprise up to 2ⁿ optical isomers. Thus, the compounds of the present invention include at least 2 enantiomers which are encompassed by the invention. The description encompasses the specific enantiomers or diastereomers of each compound as well as mixtures of different enantiomers and/or diastereomers of the compounds that possess the useful properties described herein. The optically active forms can be prepared by, for example, resolution of the racemic forms by selective crystallization techniques, by synthesis from optically active precursors, by chiral synthesis, by chromatographic separation using a chiral stationary phase or by enzymatic resolution.

The compound may also be present in different solid forms such as different crystalline forms or in the form of an amorphous solid. The description includes different crystalline forms as well as amorphous forms of the compound.

In addition, the compound may exist as hydrates or solvates, in which a certain stoichiometric amount of water or a solvent is associated with the molecule in the crystalline form. The hydrates and solvates of the compound are also the subject of the description.

Salts

In addition to the neutral compound, salt forms of the compound are also active against endoparasites. The term "veterinarily acceptable salt" is used throughout the specification to describe any salts of the compounds that are acceptable for administration for veterinary applications, and which provides an active compound upon administration.

In cases where compounds are sufficiently basic or acidic to form stable non-toxic acid or base salts, the compound may be in the form of a veterinarily or agriculturally acceptable salt. Veterinarily acceptable salts include those derived from veterinarily or agriculturally acceptable inorganic or organic bases and acids. Suitable salts include those comprising alkali metals such as lithium, sodium or potassium, alkaline earth metals such as calcium, magnesium and barium. Salts comprising transition metals including, but not limited to, manganese, copper, zinc and iron are also suitable. In addition, salts comprising ammonium cations (NH_4^+) as well as substituted ammonium cations, in which one or more of the hydrogen atoms are replaced by alkyl or aryl groups are encompassed by the description.

Salts derived from inorganic acids including, but not limited to, hydrohalide acids (HCl, HBr, HF, HI), sulfuric acid, nitric acid, phosphoric acid, and the like are particularly suitable. Suitable inorganic salts also include, but not limited to, bicarbonate, and carbonate salts. In some embodiments, examples of veterinarily and agriculturally acceptable salts are organic acid addition salts formed with organic acids including, but not limited to, maleate, dimaleate, fumarate, tosylate, methanesulfonate, acetate, citrate, malonate, tartrate, succinate, benzoate, ascorbate, α -ketoglutarate, and α -glycerophosphate. Of course, other acceptable organic acids may be used.


Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of the compound can also be made by reacting a sufficiently acidic residue on the compounds with a hydroxide of the alkali metal or alkaline earth metal.

Veterinarily acceptable salts may be obtained using standard procedures well known in the art, for example by reacting a sufficiently basic compound such as an amine with a suitably acid functional group present in the compound, or by reacting a suitable acid with a suitably basic functional group on the compound of the description.


5 Processes for the Preparation of the Compounds

The compounds of Formula (I) or pharmaceutically or a veterinarily acceptable salts thereof may be prepared by adopting schemes 1 and 2 below and the procedures in the examples:

Scheme 1



Scheme 2



In scheme 2 variables R^1 , R^2 and R^3 represent the groups defined in formula (I) above and may be introduced by a metal-catalyzed cross-coupling reaction. Examples include the Heck reaction, the Negishi coupling reaction, the Stille cross-coupling reaction, the Suzuki reaction, and others well known in the art. Variable R^5 represents the linker L bonded to one of the bicyclic rings shown in formula (I) at this position of the bicyclic core. It is well within the skill level of a person of ordinary skill in the art to adapt these schemes to synthesize a specific compound of the invention. Moreover, the starting materials are either readily available or can be made via known procedures.

Veterinary compositions

The compound and compositions comprising the compound are useful for the prevention and/or treatment of parasitic infections or infestations in animals. The compositions of the description comprise an effective amount of a compound or a veterinarily acceptable salt thereof, in combination with a veterinarily acceptable carrier or diluent and optionally a non-active

excipient. The compositions may be in a variety of solid and liquid forms which are suitable for various forms of application or administration to an animal. For example, the veterinary compositions comprising the compound may be in compositions suitable for oral administration, injectable administration, including subcutaneous and parenteral administration, and topical administration (e.g. spot-on or pour-on), dermal or subdermal administration. The compositions are intended to be administered to an animal including, but not limited to, mammals, birds and fish. Examples of mammals include but are not limited to humans, cattle, sheep, goats, llamas, alpacas, pigs, horses, donkeys, dogs, cats and other livestock or domestic mammals. Examples of birds include turkeys, chickens, ostriches and other livestock or domestic birds. The use of the compound to protect companion animals such as dogs and cats from endoparasites is particularly useful.

As discussed above, the compositions of the description may be in a form suitable for oral use (see, e.g., U.S. Patent No. 4,564,631, which is hereby incorporated by reference in its entirety), dietary supplements, troches, lozenges, chewables, tablets, hard or soft capsules, bolus, emulsions, aqueous or oily suspensions, aqueous or oily solutions, oral drench compositions, dispersible powders or granules, premixes, syrups or elixirs, enteric compositions or pastes. Compositions intended for oral use may be prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more sweetening agents, bittering agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations.

Tablets may contain the active ingredient in admixture with non-toxic, pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be, for example, inert diluents, granulating and disintegrating agents, binding agents, and lubricating agents. The tablets may be uncoated, or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. They may also be coated by the technique described in U.S. Patent Nos. 4,256,108; 4,166,452; and 4,265,874 (all incorporated herein by reference in their entirety) to form osmotic therapeutic tablets for controlled release.

Oral compositions include hard gelatin capsules. Capsules may also be soft gelatin capsules, wherein the active ingredient is mixed with water or water-miscible solvents, or an oil medium.

In one embodiment, the compound may be administered in chewable tablet compositions or soft chewable compositions such as those described in US 2013/0203692 A1, US 2010/0087492, US 2006/0222684, US 2004/0151759, US 7,955,632, all incorporated herein by reference. The veterinary compositions may be in the form of a soft chewable composition (“soft chew”), which is palatable and acceptable to the animal. In addition to the active ingredient(s), the soft chews of the description may include one or more of the following components known in the art for these dosage forms: a solvent or mixture of solvents, one or more fillers, one or more binders, one or more surfactants, one or more humectants, one or more lubricants, one or more disintegrants, one or more colorants, one or more antimicrobial agents, one or more antioxidants, one or more pH modifiers and one or more flavoring agents.

The compositions may also contain other inert ingredients such as antioxidants, preservatives, or pH stabilizers. These compounds are well known in the composition art. Antioxidants may be added to the compositions of the description to inhibit degradation of the active agents.

The compositions of the description may also include one or more lubricants and/or processing aids. In some cases, the lubricant/processing aid may also behave as a solvent, and accordingly, there some of the components of the inventive compositions may have dual functions.

Many flavoring agents may be used in the compositions of the description to improve the palatability of the oral veterinary compositions. Preferred flavoring agents are those that are not derived from animal sources. In various embodiments, flavoring components derived from fruit, meat (including, but not limited to pork, beef, chicken, fish, poultry, and the like), vegetable, cheese, bacon, cheese-bacon and/or artificial flavorings may be used. A flavoring component is typically chosen based upon consideration related to the organism that will be ingesting the soft chew. For example, a horse may prefer an apple flavoring component, while a dog may prefer a meat flavoring component. Although flavoring components derived from non-animal sources are preferred, in some embodiments, natural flavors containing beef or liver extracts, etc., may be used such as braised beef flavor artificial powdered beef flavor, roast beef flavor and corned beef flavor among others.

In another embodiment of the description, the active composition may be administered via a drench, and may be administered either topically or orally. Drench compositions are those in

which the liquid-containing compositions of the description are administered to the mouth or throat of the animal or poured onto the skin or coat of the animal.

The compositions of the description may also be in the form of oil-in-water or water-in-oil emulsions which may include emulsifying agents known in the art. The emulsions may also contain sweetening agents, bittering agents, flavoring agents, and/or preservatives.

In one embodiment, the composition of the description may be in the form of a microemulsion. Microemulsions are well suited as the liquid carrier vehicle. Microemulsions are quaternary systems comprising an aqueous phase, an oily phase, a surfactant and a co-surfactant. They are translucent and isotropic liquids.

Microemulsions are composed of stable dispersions of microdroplets of the aqueous phase in the oily phase or conversely of microdroplets of the oily phase in the aqueous phase.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil. The oily suspensions may contain a thickening agent. Sweetening agents, bittering agents, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant, or other known preservatives.

Aqueous suspensions may contain the active material in admixture with excipients suitable for the manufacture of aqueous suspensions. The aqueous suspensions may also contain one or more preservatives, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents and/or bittering agents.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water may provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Additional excipients, for example, sweetening, bittering, flavoring and coloring agents, may also be present.

Syrups and elixirs may be formulated with sweetening agents. Such compositions may also contain a demulcent, a preservative, flavoring agent(s) and/or coloring agent(s).

In another embodiment of the description, the composition may be in paste form. Examples of embodiments in a paste form include, but are not limited to, those described in U.S. Patent Nos. 6,787,342 and 7,001,889 (each of which are incorporated herein by reference). In addition to the compounds of the description, the paste may further contain fumed silica; a viscosity modifier; a carrier; optionally, an absorbent; and optionally, a colorant, stabilizer, surfactant, or preservative.

In some embodiments, the compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in
5 a non-toxic parenterally-acceptable diluent or solvent.

In addition, sterile, fixed oils may be conventionally employed as a solvent or suspending medium.

Topical, dermal and subdermal compositions may include, by way of non-limiting example, emulsions, creams, ointments, gels, pastes, powders, shampoos, pour-on compositions,
10 ready-to-use compositions, spot-on solutions and suspensions, dips and sprays. Topical application of an inventive compound or of a composition including at least one inventive compound among active agent(s) therein, in the form of a spot-on, spray-on or pour-on composition, may allow for the inventive composition to be absorbed through the skin to achieve systemic levels, distributed through the sebaceous glands or on the surface of the skin achieving levels throughout the coat.

Spot-on compositions are typically applied in a localized region which refers to an area other than
15 the entire animal. In one embodiment, the location may be between the shoulders. In another embodiment the topical composition may be administered as a stripe on the surface of the animal, *e.g.* a stripe from head to tail of the animal.

Pour-on compositions are described in U.S. Patent No. 6,010,710, also incorporated herein
20 by reference. Pour-on compositions may be advantageously oily, and generally comprise a diluent or vehicle and also a solvent (*e.g.* an organic solvent) for the active ingredient if the latter is not soluble in the diluent. In other embodiments, pour-on compositions may comprise water-miscible organic solvents.

The solvent will be used in proportion with the concentration of the active agent compound
25 and its solubility in this solvent. It will be sought to have the lowest possible volume. The vehicle makes up the difference to 100%.

In another embodiment of the description, an emollient and/or spreading and/or film-forming agent may be added to the topical composition.

In another embodiment of the description, the composition may be in ready-to-use solution
30 form as is described in U.S. Patent No. 6,395,765, incorporated herein by reference. In addition to the compounds of the description, the ready-to-use solution may contain a crystallization inhibitor

and an organic solvent or a mixture of organic solvents. In some embodiments, water may be included with the organic solvent.

The composition may also comprise an antioxidizing agent intended to inhibit oxidation in air, this agent may be present in a proportion of about 0.005 to about 1% (w/v), about 0.01 to about 0.1%, or about 0.01 to about 0.05%.

The composition excipients discussed above are well known to the practitioner in this art and may be obtained commercially or through known techniques. These compositions are generally prepared by simple mixing of the constituents as defined above; advantageously, the starting point is to mix the active material in the main solvent and then the other ingredients or adjuvants are added.

The volume of the composition applied will depend on the type of animal and the size of the animal as well as the strength of the composition and the potency of the active agents. In one embodiment, an amount of about 0.1 to about 20 ml of the composition may be applied to the animal. In other embodiment for the volume, the volume may be about 0.1 to about 10 ml, about 0.1 to about 5 ml, about 0.5 ml to about 10 ml, or about 0.3 to about 3 ml.

Spot-on compositions may be prepared by dissolving the active ingredients into the pharmaceutically or veterinary acceptable vehicle. Alternatively, the spot-on composition may be prepared by encapsulation of the active ingredient to leave a residue of the therapeutic agent on the surface of the animal. These compositions will vary with regard to the weight of the therapeutic agent in the combination depending on the species of host animal to be treated, the severity and type of infection and the body weight of the host.

Dosage forms may typically contain from about 0.1 mg to about 5 g. In other embodiments, the dosage form may contain about 0.5 mg to about 5 g of an active agent. In one embodiment of the dosage form, the dosage may contain from about 1 mg to about 500 mg of an active agent, typically about 25 mg, about 50 mg, about 100 mg, about 200 mg, about 300 mg, about 400 mg, about 500 mg, about 600 mg, about 800 mg, or about 1000 mg.

In one embodiment of the description, the compound of Formula (I) may be present in the composition at a concentration of about 0.05 to about 50% weight/weight. In other embodiments, the compound of Formula (I) may be present in a concentration of about 0.1 to about 30% (w/w). In other embodiments, the compound of Formula (I) may be present in a concentration of about 0.5 to about 30% (w/w), about 1 to about 20% (w/w) or about 0.05 to about 10% (w/w). In other

embodiments, the compound of Formula (I) may be present in a concentration of about 10 to about 50% (w/w), about 10 to about 30% (w/w), about 10 to about 20% (w/w). In yet another embodiment, the compound of Formula (I) may be present in a concentration of about 1 to 10% (w/w) or about 5 to about 15% (w/w). In another embodiment of the description, the active agent may be present in the composition as a concentration from about 0.1 to about 2% w/w. In yet another embodiment of the description, the active agent may be present in the composition as a concentration from about 0.25 to about 1.5% w/w. In still another embodiment of the description, the active agent may be present in the composition as a concentration about 1% w/w.

Methods of Treatment

As discussed above, the compound of Formula (I) are effective against endoparasites and may be used to treat and/or prevent parasitic infections in animals. In one embodiment, the present description provides a method of treating and/or preventing an endoparasite infection in or on an animal (*e.g.* a mammal or bird) comprising administering an endoparasitically effective amount of a compound of formula (I), or veterinarily acceptable salts thereof, or a composition of the description, to the animal.

The present description also provides a use of the compound of Formula (I) in the preparation of a medicament for the treatment and/or prevention of parasitic infections in animals. The present description also provides the compound of Formula (I) for use in the treatment and/or prevention of a parasitic infection in animals.

In certain embodiments, the compound of Formula (I) may also effective against ectoparasites and may be used to treat and/or prevent ectoparasitic infestations on animals. In another embodiment, the present description provides a method of treating and/or preventing an ectoparasitic infestation on an animal (*e.g.* a mammal or bird) comprising administering an ectoparasitically effective amount of a compound of formula (I), or veterinarily acceptable salts thereof, or a composition of the description, to the animal.

Also provided is the use of the compound of Formula (I) in the preparation of a medicament for the treatment and/or prevention of an ectoparasitic infestation in animals. The present description also provides a compound of Formula (I) for use in the treatment and/or prevention of an ectoparasitic infestation on animals.

In another embodiment, the description provides a method for treating and/or preventing an endoparasitic infection and an ectoparasitic infestation in and on an animal, comprising

administering a composition comprising an effective amount of a compound of formula (I) in combination with an effective amount of at least a second active agent, or veterinarily acceptable salts thereof, to the animal.

The compound of Formula (I) in combination with at least a second active agent for use in the treatment and/or prevention of an endoparasitic infection and an ectoparasitic infestation is also provided herein. In addition, the use of the compound of Formula (I) in combination with at least a second active agent in the preparation of a medicament for the preparation of a medicament for the treatment and/or prevention of an endoparasitic infection and an ectoparasitic infestation is provided.

In still another embodiment of the description, a method is provided for the treatment and/or prevention of a parasitic infestation at a locus, which comprises administering or applying a parasitically effective amount of a compound of formula (I), or veterinarily acceptable salts thereof, to the locus. With respect to animal health applications, "locus" is intended to mean a habitat, breeding ground, area, material or environment in which a parasite is growing or may grow, excluding in or on an animal.

In another embodiment, the description provides methods and uses of the compound for controlling pests in plants and crops or for protecting wood-containing structures.

In some embodiments, the animals which can be treated are mammals that include but are not limited to humans, cats, dogs, cattle, chickens, cows, bison, deer, goats, horses, llamas, camels, pigs, sheep and yaks. In one embodiment of the description, the mammals treated are humans, cats or dogs.

In one embodiment of the description, the compounds of Formula (I) have been found to have superior efficacy against endoparasites, and in particular against endoparasites that are resistant to active agents of the macrocyclic lactone class. In one embodiment, the compounds and compositions of the description are effective for controlling *Haemonchus contortus*, *Ostertagia circumcincta* and *Trichostrongylus colubriformis* in mammals or birds.

In another embodiment, the description provides a method for the treatment or prevention of a parasitic infestation or infection in an animal, comprising administering an effective amount of an anthelmintic compound of the description in combination with an effective amount of activators of invertebrate GABA receptors including an avermectin or milbemycin to the animal in need thereof.

In another embodiment, the invention provides use of the compound of Formula (I) in the manufacture of a medicament for the treatment or prevention of a parasitic infestation or infection in an animal. In yet another embodiment, the invention provides a compound of Formula (I) for use in treating or preventing a parasitic infection or infestation in an animal.

5 Avermectins that may be used in combination with the compounds of the description include, but are not limited to abamectin, dimadectin, doramectin, emamectin, eprinomectin, ivermectin, latidectin, lepimectin, and selamectin. Milbemycins compounds that may be used in combination with the compounds of the description include, but are not limited to, milbemectin, milbemycin D, moxidectin and nemadectin. Also included are the 5-oxo and 5-oxime derivatives
10 of said avermectins and milbemycins.

In one embodiment, the compounds and compositions of the description may be used for treating and/or preventing an endoparasitic infection of the following parasite: *Anaplocephala* (*Anoplocephala*), *Ancylostoma*, *Necator*, *Ascaris*, *Brugia*, *Bunostomum*, *Capillaria*, *Chabertia*, *Cooperia*, *Cyathostomum*, *Cylicocyclus*, *Cylicodontophorus*, *Cylicostephanus*, *Craterostomum*,
15 *Dictyocaulus*, *Dipetalonema*, *Dipylidium*, *Dirofilaria*, *Dracunculus*, *Echinococcus*, *Enterobius*, *Fasciola*, *Filaroides*, *Habronema*, *Haemonchus*, *Metastrongylus*, *Moniezia*, *Necator*, *Nematodirus*, *Nippostrongylus*, *Oesophagostomum*, *Onchocerca*, *Ostertagia*, *Oxyuris*, *Parascaris*, *Schistosoma*, *Strongylus*, *Taenia*, *Toxocara*, *Strongyloides*, *Toxascaris*, *Trichinella*, *Trichuris*, *Trichostrongylus*, *Triodontophorus*, *Uncinaria*, *Wuchereria*, and combinations thereof.

20 In a particularly preferred embodiment of the description, the compounds and compositions of the description are used to treat and/or prevent an infection by *Dirofilaria immitis*. The compounds have been found to be highly effective against *D. immitis* microfilaria and L4 larvae. Thus, the compounds may be used to protect animals from developing heartworm disease by killing the immature stages of *D. immitis* before they can develop into adult worms. In one
25 embodiment, the compound and compositions comprising the compounds may be used to prevent the development of heartworm disease by killing immature stages of *D. immitis* that are resistant to macrocyclic lactones. In another embodiment the compounds and compositions of the description are used to treat and/or prevent an infection by *Dirofilaria repens* or *Dirofilaria hongkongensis*.

In another embodiment of the description, the parasite is *Haemonchus contortus*, *Ostertagia circumcincta*, *Trichostrongylus axei*, *Trichostrongylus colubrifomis*, *Cooperia curticei*, *Nematodirus battus* and combinations thereof.

In another embodiment for treatment against both endoparasites and ectoparasites when
5 combined with ectoparasitocidal agents, the ectoparasite is one or more insect or arachnid including those of the genera *Ctenocephalides*, *Rhipicephalus*, *Dermacentor*, *Ixodes*, *Boophilus*, *Amblyomma*, *Haemaphysalis*, *Hyalomma*, *Sarcoptes*, *Psoroptes*, *Otodectes*, *Chorioptes*, *Hypoderma*, *Damalinia*, *Linognathus*, *Haematopinus*, *Solenoptes*, *Trichodectes*, and *Felicola*.

In another embodiment for the treatment against ectoparasites, the ectoparasite is from the
10 genera *Ctenocephalides*, *Rhipicephalus*, *Dermacentor*, *Ixodes* and/or *Boophilus*. The ectoparasites treated include but are not limited to fleas, ticks, mites, mosquitoes, flies, lice, blowfly and combinations thereof. Specific examples include but are not limited to cat and dog fleas (*Ctenocephalides felis*, *Ctenocephalides* spp. and the like), ticks (*Rhipicephalus* spp., *Ixodes* spp., *Dermacentor* spp., *Amblyomma* spp. and the like), and mites (*Demodex* spp., *Sarcoptes* spp.,
15 *Otodectes* spp. and the like), lice (*Trichodectes* spp., *Cheyletiella* spp., *Linognathus* spp., and the like), mosquitoes (*Aedes* spp., *Culex* spp., *Anopheles* spp., and the like) and flies (*Haematobia* spp., *Musca* spp., *Stomoxys* spp., *Dermatobia* spp., *Cochliomyia* spp., and the like). In yet another embodiment for the treatment against ectoparasites, the ectoparasite is a flea and/or tick.

Additional examples of ectoparasites include, but are not limited to the tick genus
20 *Boophilus*, especially those of the species *microplus* (cattle tick), *decoloratus* and *annulatus*; myiasis such as *Dermatobia hominis* (known as Berne in Brazil) and *Cochliomyia hominivorax* (greenbottle); sheep myiasis such as *Lucilia sericata*, *Lucilia cuprina* (known as blowfly strike in Australia, New Zealand and South Africa). Flies proper, namely those whose adult constitutes the parasite, such as *Haematobia irritans* (horn fly); lice such as *Linognathus vitulorum*, etc.; and
25 mites such as *Sarcoptes scabiei* and *Psoroptes ovis*. The above list is not exhaustive and other ectoparasites are well known in the art to be harmful to animals and humans. These include, for example migrating dipterous larvae.

In another embodiment of the description, the compounds and compositions of the description are suitable for controlling pests such as insects selected from the group consisting of
30 *Blatella germanica*, *Heliothis virescens*, *Leptinotarsa decemlineata*, *Tetramorium caespitum* and combinations thereof.

The phytoparasitic nematodes include, for example, *Anguina* spp., *Aphelenchoides* spp., *Belonoaimus* spp., *Bursaphelenchus* spp., *Ditylenchus dipsaci*, *Globodera* spp., *Heliocotylenchus* spp., *Heterodera* spp., *Longidorus* spp., *Meloidogyne* spp., *Pratylenchus* spp., *Radopholus similis*, *Rotylenchus* spp., *Trichodorus* spp., *Tylenchorhynchus* spp., *Tylenchulus* spp., *Tylenchulus* *semipenetrans*, *Xiphinema* spp.

In addition, with or without the other pesticidal agents added to the composition, the description can also be used to treat other pests which include but are not limited to pests:

(1) from the order of Isopoda, for example *Oniscus asellus*, *Armadillidium vulgare* and *Porcellio scaber*;

(2) from the order of Diplopoda, for example *Blaniulus guttulatus*;

(3) from the order of Chilopoda, for example *Geophilus carpophagus* and *Scutigera* spp.;

(4) from the order of Symphyla, for example *Scutigera immaculata*;

(5) from the order of Thysanura, for example *Lepisma saccharina*;

(6) from the order of Collembola, for example *Onychiurus armatus*;

(7) from the order of Blattaria, for example *Blatta orientalis*, *Periplaneta americana*, *Leucophaea maderae* and *Blattella germanica*;

(8) from the order of Hymenoptera, for example *Diprion* spp., *Hoplocampa* spp., *Lasius* spp., *Monomorium pharaonis* and *Vespa* spp.;

(9) from the order of Siphonaptera, for example *Xenopsylla cheopis* and *Ceratophyllus* spp.;

(10) from the order of Anoplura (Phthiraptera), for example, *Damalinia* spp., *Haematopinus* spp., *Linognathus* spp., *Pediculus* spp., *Trichodectes* spp.;

(11) from the class of Arachnida, for example, *Acarus siro*, *Aceria sheldoni*, *Aculops* spp., *Aculus* spp., *Amblyomma* spp., *Argas* spp., *Boophilus* spp., *Brevipalpus* spp., *Bryobia praetiosa*, *Chorioptes* spp., *Dermanyssus gallinae*, *Eotetranychus* spp., *Epitrimerus pyri*, *Eutetranychus* spp., *Eriophyes* spp., *Hemitarsonemus* spp., *Hyalomma* spp., *Ixodes* spp., *Latrodectus mactans*, *Metatetranychus* spp., *Oligonychus* spp., *Ornithodoros* spp., *Panonychus* spp., *Phyllocoptura oleivora*, *Polyphagotarsonemus latus*, *Psoroptes* spp., *Rhipicephalus* spp., *Rhizoglyphus* spp., *Sarcoptes* spp., *Scorpio maurus*, *Stenotarsonemus* spp., *Tarsonemus* spp., *Tetranychus* spp., *Vasates lycopersici*;

(12) from the class of Bivalva, for example, *Dreissena* spp.;

(13) from the order of Coleoptera, for example, *Acanthoscelides obtectus*, *Adoretus* spp., *Agelastica alni*, *Agriotes* spp., *Amphimallon solstitialis*, *Anobium punctatum*, *Anoplophora* spp., *Anthonomus* spp., *Anthrenus* spp., *Apogonia* spp., *Atomaria* spp., *Attagenus* spp., *Bruchidius* spp., *Bruchus* spp., *Ceuthorhynchus* spp., *Cleonus mendicus*, *Conoderus* spp., *Cosmopolites* spp., *Costelytra zealandica*, *Curculio* spp., *Cryptorhynchus lapathi*, *Dermestes* spp., *Diabrotica* spp., *Epilachna* spp., *Faustinus cubae*, *Gibbium psylloides*, *Heteronychus arator*, *Hylamorphia elegans*, *Hylotrupes bajulus*, *Hypera postica*, *Hypothenemus* spp., *Lachnosterna consanguinea*, *Leptinotarsa decemlineata*, *Lissorhoptrus oryzophilus*, *Lixus* spp., *Lyctus* spp., *Meligethes aeneus*,
 10 *Melolontha*, *Migdolus* spp., *Monochamus* spp., *Naupactus xanthographus*, *Niptus hololeucus*, *Oryctes rhinoceros*, *Oryzaephilus surinamensis*, *Otiorrhynchus sulcatus*, *Oxycetonia jucunda*, *Phaedon cochleariae*, *Phyllophaga* spp., *Popillia japonica*, *Premnotrypes* spp., *Psylliodes chrysocephala*, *Ptinus* spp., *Rhizobius ventralis*, *Rhizopertha dominica*, *Sitophilus* spp., *Sphenophorus* spp., *Sternechus* spp., *Symphyletes* spp., *Tenebrio molitor*, *Tribolium* spp.,
 15 *Trogoderma* spp., *Tychius* spp., *Xylotrechus* spp., *Zabrus* spp.;

(14) from the order of Diptera, for example, *Aedes* spp., *Anopheles* spp., *Bibio hortulanus*, *Calliphora erythrocephala*, *Ceratitis capitata*, *Chrysomyia* spp., *Cochliomyia* spp., *Cordylobia anthropophaga*, *Culex* spp., *Cuterebra* spp., *Dacus oleae*, *Dermatobia hominis*, *Drosophila* spp., *Fannia* spp., *Gastrophilus* spp., *Hylemyia* spp., *Hyppobosca* spp., *Hypoderma* spp., *Liriomyza* spp., *Lucilia* spp., *Musca* spp., *Nezara* spp., *Oestrus* spp., *Oscinella frit*, *Pegomyia hyoscyami*, *Phorbia* spp., *Stomoxys* spp., *Tabanus* spp., *Tannia* spp., *Tipula paludosa*, *Wohlfahrtia* spp.;

(15) from the class of Gastropoda, for example, *Arion* spp., *Biomphalaria* spp., *Bulinus* spp., *Deroceras* spp., *Galba* spp., *Lymnaea* spp., *Oncomelania* spp., *Succinea* spp.;

(16) from the class of helminths, for example, *Ancylostoma duodenale*, *Ancylostoma ceylanicum*, *Ancylostoma braziliensis*, *Ancylostoma* spp., *Ascaris lubricoides*, *Ascaris* spp., *Brugia malayi*, *Brugia timori*, *Bunostomum* spp., *Chabertia* spp., *Clonorchis* spp., *Cooperia* spp., *Dicrocoelium* spp., *Dictyocaulus filaria*, *Diphyllbothrium latum*, *Dracunculus medinensis*, *Echinococcus granulosus*, *Echinococcus multilocularis*, *Enterobius vermicularis*, *Faciola* spp.,
 25 *Haemonchus* spp., *Heterakis* spp., *Hymenolepis nana*, *Hyostrogylus* spp., *Loa*, *Nematodirus* spp., *Oesophagostomum* spp., *Opisthorchis* spp., *Onchocerca volvulus*, *Ostertagia* spp., *Paragonimus*

spp., *Schistosomen* spp., *Strongyloides fuelleborni*, *Strongyloides stercoralis*, *Strongyloides* spp., *Taenia saginata*, *Taenia solium*, *Trichinella spiralis*, *Trichinella nativa*, *Trichinella britovi*, *Trichinella nelsoni*, *Trichinella pseudopsiralis*, *Trichostrongylus* spp., *Trichuris trichuria*, *Wuchereria bancrofti*;

5 (17) from the order of Heteroptera, for example, *Anasa tristis*, *Antestiopsis* spp., *Blissus* spp., *Calocoris* spp., *Campylomma livida*, *Cavelerius* spp., *Cimex* spp., *Creontiades dilutus*, *Dasynus piperis*, *Dichelops furcatus*, *Diconocoris hewetti*, *Dysdercus* spp., *Euschistus* spp., *Eurygaster* spp., *Heliopeltis* spp., *Horcias nobilellus*, *Leptocorisa* spp., *Leptoglossus phyllopus*, *Lygus* spp., *Macropes excavatus*, *Miridae*, *Nezara* spp., *Oebalus* spp., *Pentomidae*, *Piesma*
10 *quadrata*, *Piezodorus* spp., *Psallus seriatus*, *Pseudacysta persea*, *Rhodnius* spp., *Sahlbergella singularis*, *Scotinophora* spp., *Stephanitis nashi*, *Tibraca* spp., *Triatoma* spp.;

(18) from the order of Homoptera, for example, *Acyrtosipon* spp., *Aeneolamia* spp., *Agonosцена* spp., *Aleurodes* spp., *Aleurolobus barodensis*, *Aleurothrixus* spp., *Amrasca* spp., *Anuraphis cardui*, *Aonidiella* spp., *Aphanostigma piri*, *Aphis* spp., *Arboridia apicalis*, *Aspidiella*
15 spp., *Aspidiotus* spp., *Atanus* spp., *Aulacorthum solani*, *Bemisia* spp., *Brachycaudus helichrysi*, *Brachycolus* spp., *Brevicoryne brassicae*, *Calligypona marginata*, *Carneocephala fulgida*, *Ceratovacuna lanigera*, *Cercopidae*, *Ceroplastes* spp., *Chaetosiphon fragaefolii*, *Chionaspis tegalensis*, *Chlorita omukii*, *Chromaphis juglandicola*, *Chrysomphalus ficus*, *Cicadulina mbila*, *Cocco-mytilus halli*, *Coccus* spp., *Cryptomyzus ribis*, *Dalbulus* spp., *Dialeurodes* spp., *Diaphorina*
20 spp., *Diaspis* spp., *Doralis* spp., *Drosicha* spp., *Dysaphis* spp., *Dysmicoccus* spp., *Empoasca* spp., *Eriosoma* spp., *Erythroneura* spp., *Euscelis bilobatus*, *Geococcus coffeae*, *Homalodisca coagulata*, *Hyalopterus arundinis*, *Icerya* spp., *Idiocerus* spp., *Idioscopus* spp., *Laodelphax striatellus*, *Lecanium* spp., *Lepidosaphes* spp., *Lipaphis erysimi*, *Macrosiphum* spp., *Mahanarva fimbriolata*, *Melanaphis sacchari*, *Metcalfiella* spp., *Metopolophium dirhodum*, *Monellia costalis*,
25 *Monelliopsis pecanis*, *Myzus* spp., *Nasonovia ribisnigri*, *Nephotettix* spp., *Nilaparvata lugens*, *Oncometopia* spp., *Orthezia praelonga*, *Parabemisia myricae*, *Paratrioza* spp., *Parlatoria* spp., *Pemphigus* spp., *Peregrinus maidis*, *Phenacoccus* spp., *Phloeomyzus passerinii*, *Phorodon humuli*, *Phylloxera* spp., *Pinaspis aspidistrae*, *Planococcus* spp., *Protopulvinaria pyriformis*, *Pseudaulacaspis pentagona*, *Pseudococcus* spp., *Psylla* spp., *Pteromalus* spp., *Pyrilla* spp.,
30 *Quadraspidotus* spp., *Quesada gigas*, *Rastrococcus* spp., *Rhopalosiphum* spp., *Saissetia* spp., *Scaphoides titanus*, *Schizaphis graminum*, *Selenaspis articulatus*, *Sogata* spp., *Sogatella*

furcifera, *Sogatodes* spp., *Stictocephala festina*, *Tenalaphara malayensis*, *Tinocallis caryaefoliae*, *Tomaspis* spp., *Toxoptera* spp., *Trialeurodes vaporariorum*, *Trioza* spp., *Typhlocyba* spp., *Unaspis* spp., *Viteus vitifolii*;

(19) from the order of Isoptera, for example, *Reticulitermes* spp., *Odontotermes* spp.;

5 (20) from the order of Lepidoptera, for example, *Acronicta major*, *Aedia leucomelas*, *Agrotis* spp., *Alabama argillacea*, *Anticarsia* spp., *Barathra brassicae*, *Bucculatrix thurberiella*, *Bupalus piniarius*, *Cacoecia podana*, *Capua reticulana*, *Carpocapsa pomonella*, *Cheimatobia brumata*, *Chilo* spp., *Choristoneura fumiferana*, *Clysia ambiguella*, *Cnaphalocerus* spp., *Earias insulana*, *Ephestia kuehniella*, *Euproctis chrysorrhoea*, *Euxoa* spp., *Feltia* spp., *Galleria mellonella*, *Helicoverpa* spp., *Heliothis* spp., *Hofmannophila pseudospretella*, *Homona magnanima*, *Hyponomeuta padella*, *Laphygma* spp., *Lithocolletis blancardella*, *Lithophane antennata*, *Loxagrotis albicosta*, *Lymantria* spp., *Malacosoma neustria*, *Mamestra brassicae*, *Mocis repanda*, *Mythimna separata*, *Oria* spp., *Oulema oryzae*, *Panolis flammea*, *Pectinophora gossypiella*, *Phyllocnistis citrella*, *Pieris* spp., *Plutella xylostella*, *Prodenia* spp., *Pseudaletia* spp.,
10 *Pseudoplusia includens*, *Pyrausta nubilalis*, *Spodoptera* spp., *Thermesia gemmatilis*, *Tinea pellionella*, *Tineola bisselliella*, *Tortrix viridana*, *Trichoplusia* spp.;

(21) from the order of Orthoptera, for example, *Acheta domesticus*, *Blatta orientalis*, *Blattella germanica*, *Gryllotalpa* spp., *Leucophaea maderae*, *Locusta* spp., *Melanoplus* spp., *Periplaneta americana*, *Schistocerca gregaria*;

20 (22) from the order of Thysanoptera, for example, *Baliothrips biformis*, *Enneothrips flavens*, *Frankliniella* spp., *Heliothrips* spp., *Hercinothrips femoralis*, *Kakothrips* spp., *Rhipiphorocthris cruentatus*, *Scirtothrips* spp., *Taeniothrips cardamoni*, *Thrips* spp.;

(23) from the class of Protozoa, for example, *Eimeria* spp.

25 In each aspect of the description, the compounds and compositions of the description can be applied against a single pest or combinations thereof.

Mixtures with other active agents

In another embodiment, the compositions comprising the compounds of Formula (I) may also include other veterinary therapeutic agents. Veterinary pharmaceutical agents that may be included in the compositions of the description are well-known in the art (see *e.g. Plumb's Veterinary Drug Handbook*, 5th Edition, ed. Donald C. Plumb, Blackwell Publishing, (2005) or
30 *The Merck Veterinary Manual*, 9th Edition, (January 2005)) and include but are not limited to

acarbose, acepromazine maleate, acetaminophen, acetazolamide, acetazolamide sodium, acetic acid, acetohydroxamic acid, acetylcysteine, acitretin, acyclovir, albendazole, albuterol sulfate, alfentanil, allopurinol, alprazolam, altrenogest, amantadine, amikacin sulfate, aminocaproic acid, aminopentamide hydrogen sulfate, aminophylline/theophylline, amiodarone, amitriptyline,

5 amlodipine besylate, ammonium chloride, ammonium molybdenate, amoxicillin, clavulanate potassium, amphotericin B desoxycholate, amphotericin B lipid-based, ampicillin, amprolium, antacids (oral), antivenin, apomorphine, apramycin sulfate, ascorbic acid, asparaginase, aspiring, atenolol, atipamezole, atracurium besylate, atropine sulfate, aurnofin, aurothioglucoase, azaperone, azathioprine, azithromycin, baclofen, barbituates, benazepril, betamethasone, bethanechol

10 chloride, bisacodyl, bismuth subsalicylate, bleomycin sulfate, boldenone undecylenate, bromides, bromocriptine mesylate, budenoside, buprenorphine, buspirone, busulfan, butorphanol tartrate, cabergoline, calcitonin salmon, calcitrol, calcium salts, captopril, carbenicillin indanyl sodium, carbimazole, carboplatin, carnitine, carprofen, carvedilol, cefadroxil, cefazolin sodium, cefixime, clorsulon, cefoperazone sodium, cefotaxime sodium, cefotetan disodium, cefoxitin sodium,

15 cefpodoxime proxetil, ceftazidime, ceftiofur sodium, ceftiofur, ceftiaxone sodium, cephalixin, cephalosporins, cephalirin, charcoal (activated), chlorambucil, chloramphenicol, chlordiazepoxide, chlordiazepoxide +/- clidinium bromide, chlorothiazide, chlorpheniramine maleate, chlorpromazine, chlorpropamide, chlortetracycline, chorionic gonadotropin (HCG), chromium, cimetidine, ciprofloxacin, cisapride, cisplatin, citrate salts, clarithromycin, clemastine

20 fumarate, clenbuterol, clindamycin, clofazimine, clomipramine, clonazepam, clonidine, cloprostenol sodium, clorazepate dipotassium, clorsulon, cloxacillin, codeine phosphate, colchicine, corticotropin (ACTH), cosyntropin, cyclophosphamide, cyclosporine, cyproheptadine, cytarabine, dacarbazine, dactinomycin/actinomycin D, dalteparin sodium, danazol, dantrolene sodium, dapsone, decoquinat, deferoxamine mesylate, deracoxib, deslorelin acetate,

25 desmopressin acetate, desoxycorticosterone pivalate, detomidine, dexamethasone, dexpantenol, dexraazoxane, dextran, diazepam, diazoxide (oral), dichlorphenamide, diclofenac sodium, dicloxacillin, diethylcarbamazine citrate, diethylstilbestrol (DES), difloxacin, digoxin, dihydrotachysterol (DHT), diltiazem, dimenhydrinate, dimercaprol/BAL, dimethyl sulfoxide, dinoprost tromethamine, diphenylhydramine, disopyramide phosphate, dobutamine,

30 docusate/DSS, dolasetron mesylate, domperidone, dopamine, doramectin, doxapram, doxepin, doxorubicin, doxycycline, edetate calcium disodium, calcium EDTA, edrophonium chloride,

enalapril/enalaprilat, enoxaparin sodium, enrofloxacin, ephedrine sulfate, epinephrine, epoetin/erythropoietin, eprinomectin, epsiprantel, erythromycin, esmolol, estradiol cypionate, ethacrynic acid/ethacrynate sodium, ethanol (alcohol), etidronate sodium, etodolac, etomidate, euthanasia agents w/pentobarbital, famotidine, fatty acids (essential/omega), felbamate, fentanyl, 5 ferrous sulfate, filgrastim, finasteride, fipronil, florfenicol, fluconazole, flucytosine, fludrocortisone acetate, flumazenil, flumethasone, flunixin meglumine, fluorouracil (5-FU), fluoxetine, fluticasone propionate, fluvoxamine maleate, fomepizole (4-MP), furazolidone, furosemide, gabapentin, gemcitabine, gentamicin sulfate, glimepiride, glipizide, glucagon, glucocorticoid agents, glucosamine/chondroitin sulfate, glutamine, glyburide, glycerine (oral), 10 glycopyrrolate, gonadorelin, griseofulvin, guaifenesin, halothane, hemoglobin glutamer-200 (OXYGLOBIN®), heparin, hetastarch, hyaluronate sodium, hydralazine, hydrochlorothiazide, hydrocodone bitartrate, hydrocortisone, hydromorphone, hydroxyurea, hydroxyzine, ifosfamide, imidacloprid, imidocarb dipropionate, impenem-cilastatin sodium, imipramine, inamrinone lactate, insulin, interferon alfa-2a (human recombinant), iodide (sodium/potassium), ipecac (syrup), 15 ipodate sodium, iron dextran, isoflurane, isoproterenol, isotretinoin, isoxsuprine, itraconazole, ivermectin, kaolin/pectin, ketamine, ketoconazole, ketoprofen, ketorolac tromethamine, lactulose, leuprolide, levamisole, levetiracetam, levothyroxine sodium, lidocaine, lincomycin, liothyronine sodium, lisinopril, lomustine (CCNU), lufenuron, lysine, magnesium, mannitol, marbofloxacin, mechlorethamine, meclizine, meclofenamic acid, medetomidine, medium chain triglycerides, 20 medroxyprogesterone acetate, megestrol acetate, melarsomine, melatonin, meloxicam, melphalan, meperidine, mercaptopurine, meropenem, metformin, methadone, methazolamide, methenamine mandelate/hippurate, methimazole, methionine, methocarbamol, methohexital sodium, methotrexate, methoxyflurane, methylene blue, methylphenidate, methylprednisolone, metoclopramide, metoprolol, metronidazole, mexiletine, mibolerone, midazolam milbemyacin 25 oxime, mineral oil, minocycline, misoprostol, mitotane, mitoxantrone, morphine sulfate, moxidectin, naloxone, mandrolone decanoate, naproxen, narcotic (opiate) agonist analgesics, neomycin sulfate, neostigmine, niacinamide, nitazoxanide, nitenpyram, nitrofurantoin, nitroglycerin, nitroprusside sodium, nizatidine, novobiocin sodium, nystatin, octreotide acetate, olsalazine sodium, omeprazole, ondansetron, opiate antidiarrheals, orbifloxacin, oxacillin sodium, 30 oxazepam, oxibutynin chloride, oxymorphone, oxytetracycline, oxytocin, pamidronate disodium, pancrelipase, pancuronium bromide, paromomycin sulfate, paroxetine, pencillamine, general

information penicillins, penicillin G, penicillin V potassium, pentazocine, pentobarbital sodium, pentosan polysulfate sodium, pentoxifylline, pergolide mesylate, phenobarbital, phenoxybenzamine, pheylbutazone, phenylephrine, phenypropanolamine, phenytoin sodium, pheromones, parenteral phosphate, phytonadione/vitamin K-1, pimobendan, piperazine, pirlimycin, piroxicam, polysulfated glycosaminoglycan, ponazuril, potassium chloride, pralidoxime chloride, prazosin, prednisolone/prednisone, primidone, procainamide, procarbazine, prochlorperazine, propantheline bromide, propionibacterium acnes injection, propofol, propranolol, protamine sulfate, pseudoephedrine, psyllium hydrophilic mucilloid, pyridostigmine bromide, pyrilamine maleate, pyrimethamine, quinacrine, quinidine, ranitidine, rifampin, s-adenosyl-methionine (SAMe), saline/hyperosmotic laxative, selamectin, selegiline /l-deprenyl, sertraline, sevelamer, sevoflurane, silymarin/milk thistle, sodium bicarbonate, sodium polystyrene sulfonate, sodium stibogluconate, sodium sulfate, sodium thiosulfate, somatotropin, sotalol, spectinomycin, spironolactone, stanozolol, streptokinase, streptozocin, succimer, succinylcholine chloride, sucralfate, sufentanil citrate, sulfachlorpyridazine sodium, sulfadiazine/trimethoprim, sulfamethoxazole/trimethoprim, sulfadimentoxine, sulfadimethoxine/ormetoprim, sulfasalazine, taurine, tepoxaline, terbinafine, terbutaline sulfate, testosterone, tetracycline, thiacetarsamide sodium, thiamine, thioguanine, thiopental sodium, thiotepa, thyrotropin, tiamulin, ticarcilin disodium, tiletamine /zolazepam, tilmocsin, tiopronin, tobramycin sulfate, tocanide, tolazoline, telfenamic acid, topiramate, tramadol, trimcinolone acetone, trientine, trilostane, trimeprazine tartrate w/prednisolone, tripeleminamine, tylosin, urdosiol, valproic acid, vanadium, vancomycin, vasopressin, vecuronium bromide, verapamil, vinblastine sulfate, vincristine sulfate, vitamin E/selenium, warfarin sodium, xylazine, yohimbine, zafirlukast, zidovudine (AZT), zinc acetate/zinc sulfate, zonisamide and mixtures thereof.

In one embodiment of the description, arylpyrazole compounds such as phenylpyrazoles may be included in the veterinary compositions of the description. Arylpyrazoles are known in the art and may be suitable for combination with the compound of Formula (I) in the compositions of the description. Examples of such arylpyrazole compounds include but are not limited to those described in U.S. Patent Nos. 6,001,384; 6,010,710; 6,083,519; 6,096,329; 6,174,540; 6,685,954, 6,998,131 and 7,759,381 (all of which are incorporated herein by reference). A particularly preferred arylpyrazole active agent is fipronil.

In another embodiment of the description, one or more macrocyclic lactones, which act as

an acaricide, an anthelmintic agent and/or an insecticide, can be included in the compositions of the description in combination with the compound. For the avoidance of doubt, the term “macrocyclic lactone” as used herein includes both naturally occurring and synthetic or semi-synthetic avermectin and milbemycin compounds.

5 The macrocyclic lactones that may be used in the compositions of the description include, but are not limited to, the naturally produced avermectins (*e.g.* including the components designated as A_{1a}, A_{1b}, A_{2a}, A_{2b}, B_{1a}, B_{1b}, B_{2a} and B_{2b}) and milbemycin compounds, semisynthetic avermectins and milbemycins, avermectin monosaccharide compounds and avermectin aglycone compounds. Examples of macrocyclic lactone compounds that may be used
10 in the compositions include, but are not limited to, abamectin, dimadectin, doramectin, emamectin, eprinomectin, ivermectin, latidectin, lepimectin, selamectin, ML-1,694,554 and milbemycins including, but not limited to, milbemectin, milbemycin D, milbemycin A₃, milbemycin A₄, milbemycin oxime, moxidectin and nemadectin. Also included are the 5-oxo and 5-oxime derivatives of said avermectins and milbemycins.

15 The macrocyclic lactone compounds are known in the art and can easily be obtained commercially or through synthesis techniques known in the art. Reference is made to the widely available technical and commercial literature. For avermectins, ivermectin and abamectin, reference may be made, for example, to the work “Ivermectin and Abamectin”, 1989, by M.H. Fischer and H. Mrozik, William C. Campbell, published by Springer Verlag., or Albers-Schönberg
20 *et al.* (1981), “Avermectins Structure Determination”, J. Am. Chem. Soc., 103, 4216-4221. For doramectin, “Veterinary Parasitology”, vol. 49, No. 1, July 1993, 5-15 may be consulted. For milbemycins, reference may be made, *inter alia*, to Davies H.G. *et al.*, 1986, “Avermectins and Milbemycins”, Nat. Prod. Rep., 3, 87-121, Mrozik H. *et al.*, 1983, Synthesis of Milbemycins from Avermectins, Tetrahedron Lett., 24, 5333-5336, U.S. Patent No. 4,134,973 and EP 0 677 054, both
25 incorporated herein by reference.

 The structure of the avermectins and milbemycins are closely related, *e.g.*, by sharing a complex 16-membered macrocyclic lactone ring. The natural product avermectins are disclosed in U.S. Patent No. 4,310,519 and the 22,23-dihydro avermectin compounds are disclosed in U.S. Patent No. 4,199,569. Mention is also made of U.S. Patent Nos. 4,468,390, 5,824,653, EP 0 007
30 812 A1, U.K. Patent Specification 1 390 336, EP 0 002 916, and New Zealand Patent No. 237 086, *inter alia*. Naturally occurring milbemycins are described in U.S. Patent No. 3,950,360 as well as

in the various references cited in “The Merck Index” 12th ed., S. Budavari, Ed., Merck & Co., Inc. Whitehouse Station, New Jersey (1996). Latidectin is described in the “International Nonproprietary Names for Pharmaceutical Substances (INN)”, *WHO Drug Information*, vol. 17, no. 4, pp. 263- 286, (2003). Semisynthetic derivatives of these classes of compounds are well known in the art and are described, for example, in U.S. Patent Nos. 5,077,308, 4,859,657, 4,963,582, 4,855,317, 4,871,719, 4,874,749, 4,427,663, 4,310,519, 4,199,569, 5,055,596, 4,973,711, 4,978,677, 4,920,148 and EP 0 667 054, all incorporated herein by reference.

In one embodiment, the veterinary compositions of the description comprise an effective amount of at least one of abamectin, dimadectin, doramectin, emamectin, eprinomectin, ivermectin, latidectin, lepimectin, selamectin, milbemectin, milbemycin D, milbemycin A₃, milbemycin A₄, milbemycin oxime, moxidectin or nemadectin, or a combination thereof. In another embodiment, the description provides a veterinary composition comprising an effective amount of at least one of abamectin, emamectin, eprinomectin, ivermectin, doramectin or selamectin, or a combination thereof. In still another embodiment, the veterinary compositions of the description comprise an effective amount of at least one of ivermectin, milbemectin, milbemycin oxime or moxidectin, or a combination thereof.

In another embodiment of the description, a composition comprising a compound of Formula (I) in combination with a class of acaricide or insecticides known as insect growth regulators (IGRs) are provided. Compounds belonging to this group are well known to the practitioner and represent a wide range of different chemical classes. These compounds all act by interfering with the development or growth of the insect pests. Insect growth regulators are described, for example, in U.S. Patent Nos. 3,748,356, 3,818,047, 4,225,598, 4,798,837, 4,751,225, EP 0 179 022 or U.K. 2 140 010 as well as U.S. Patent Nos. 6,096,329 and 6,685,954 (all incorporated herein by reference).

In one embodiment the compositions of the description may include an IGR compound that mimics juvenile hormone or that modulates levels of juvenile hormones in insects. Examples of juvenile hormone mimics include azadirachtin, diofenolan, fenoxycarb, hydroprene, kinoprene, methoprene, pyriproxyfen, tetrahydroazadirachtin and 4-chloro-2-(2-chloro-2-methyl-propyl)-5-(6-iodo-3-pyridylmethoxy)pyridazine-3(2H)-one. In another embodiment, the compositions of the description comprise a compound of formula (I) in combination with methoprene or pyriproxyfen and a pharmaceutically acceptable carrier.

In another embodiment, the compositions of the description include an IGR compound that is a chitin synthesis inhibitor. Chitin synthesis inhibitors include chlorofluazuron, cyromazine, diflubenzuron, fluazuron, flucycloxuron, flufenoxuron, hexaflumoron, lufenuron, tebufenozide, teflubenzuron, triflumoron, 1-(2,6-difluorobenzoyl)-3-(2-fluoro-4-(trifluoromethyl)phenyl)urea, 1-
5 (2,6-difluoro-benzoyl)-3-(2-fluoro-4-(1,1,2,2-tetrafluoroethoxy)-phenyl)urea and 1-(2,6-difluorobenzoyl)-3-(2-fluoro-4-trifluoromethyl)phenylurea.

In some embodiments, the compositions of the description may include one or more antinematodal agents including, but not limited to, active agents in the benzimidazoles, imidazothiazoles, tetrahydropyrimidines and the organophosphate class of compounds. In some
10 embodiments, benzimidazoles including, but not limited to, thiabendazole, cambendazole, parbendazole, oxibendazole, mebendazole, flubendazole, fenbendazole, oxfendazole, albendazole, cyclobendazole, febantel, thiophanate and its *o,o*-dimethyl analogue may be included in the compositions.

In other embodiments, the compositions of the description may include an imidazothiazole
15 compounds including, but not limited to, tetramisole, levamisole and butamisol.

In still other embodiments, the compositions of the description may include tetrahydropyrimidine active agents including, but not limited to, pyrantel, oxantel, and morantel.

Suitable organophosphate active agents include, but are not limited to, coumaphos, trichlorfon, haloxon, naftalofos and dichlorvos, heptenophos, mevinphos, monocrotophos, TEPP,
20 and tetrachlorvinphos.

In other embodiments, the compositions may include the antinematodal compounds phenothiazine, piperazine as the neutral compound and in various salt forms, diethylcarbazine, phenols such as disophenol, arsenicals such as arsenamide, ethanolamines such as bethovenium, thenium closylate, and methyridine; cyanine dyes including pyrvinium chloride, pyrvinium
25 pamoate and dithiazanine iodide; isothiocyanates including bitoscanate, suramin sodium, phthalofyne, and various natural products including, but not limited to, hygromycin B, α -santonin and kainic acid.

In other embodiments, the compositions of the description may include antitrepatodal agents. Suitable antitrepatodal agents include, but are not limited to, the miracils such as miracil
30 D and mirasan; praziquantel, clonazepam and its 3-methyl derivative, oltipraz, lucanthone, hycanthone, oxamniquine, amoscanate, niridazole, nitroxylin, various bisphenol compounds

known in the art including hexachlorophene, bithionol, bithionol sulfoxide and menichlopholan; various salicylanilide compounds including tribromsalan, oxyclozanide, clioxanide, rafoxanide, nitroxylin, brotianide, bromoxanide and closantel; triclabendazole, diamfenetide, clorsulon, hetolin and emetine.

5 Anticestodal compounds may also be advantageously used in the compositions of the description including, but not limited to, arecoline in various salt forms, bunamidine, niclosamide, nitroscanate, paromomycin, paromomycin II, praziquantel and epsiprantel.

In yet other embodiments, the compositions of the description may include other active agents that are effective against arthropod parasites. Suitable active agents include, but are not
10 limited to, bromocyclen, chlordane, DDT, endosulfan, lindane, methoxychlor, toxaphene, bromophos, bromophos-ethyl, carbophenothion, chlorfenvinphos, chlorpyrifos, crotoxyphos, cythioate, diazinon, dichlorenthion, diemthoate, dioxathion, ethion, famphur, fenitrothion, fenthion, fospirate, iodofenphos, malathion, naled, phosalone, phosmet, phoxim, propetamphos, ronnel, stirofos, allethrin, cyhalothrin, cypermethrin, deltamethrin, fenvalerate, flucythrinate,
15 permethrin, phenothrin, pyrethrins, resmethrin, benzyl benzoate, carbon disulfide, crotamiton, diflubenzuron, diphenylamine, disulfiram, isobornyl thiocyanato acetate, methoprene, monosulfiram, pirenonylbutoxide, rotenone, triphenyltin acetate, triphenyltin hydroxide, deet, dimethyl phthalate, and the compounds 1,5a,6,9,9a,9b-hexahydro-4a(4H)-dibenzofurancarboxaldehyde (MGK-11), 2-(2-ethylhexyl)-3a,4,7,7a-tetrahydro-4,7-methano-1H-
20 isoindole-1,3(2H)dione (MGK-264), dipropyl-2,5-pyridinedicarboxylate (MGK-326) and 2-(octylthio)ethanol (MGK-874).

In another embodiment, an antiparasitic agent that can be included in the veterinary composition containing a compound of formula (I) can be a biologically active peptide or protein including, but not limited to, depsipeptides other than the compound. These include PF1022A or
25 analogs thereof and emodepside. Other cyclic depsipeptide compounds that may be included in the compositions comprising a compound of Formula (I) are those described in WO 2016/187534 A1 and WO 2017/116702 A1, both incorporated herein by reference. These compounds act at the neuromuscular junction by stimulating presynaptic receptors belonging to the secretin receptor family resulting in the paralysis and death of parasites. In one embodiment of the depsipeptide, the
30 depsipeptide is emodepside (see Wilson *et al.*, *Parasitology*, Jan. 2003, 126(Pt 1):79-86).

In another embodiment, the compositions of the description may comprise an active agent

from the neonicotinoid class of parasiticides. The neonicotinoids bind and inhibit insect specific nicotinic acetylcholine receptors. In one embodiment, the neonicotinoid insecticidal agent that can be combined with a compound of Formula (I) in a composition of the description is imidacloprid. Agents of this class are described, for example, in U.S. Patent No. 4,742,060 or in EP 0 892 060
5 (both incorporated herein by reference). In another embodiment, the compositions of the description may comprise nitenpyram, another active agent of the neonicotinoid class of pesticides. The use of nitenpyram for controlling fleas is described in U.S. Patent No. 5,750,548, which is incorporated herein by reference in its entirety.

In certain other embodiments of the description, the compound of Formula (I) can be
10 combined with the compositions of the description is a semicarbazone, such as metaflumizone.

In another embodiment, the compositions of the description may advantageously include one or more isoxazoline compounds known in the art. Isoxazoline active agents are highly effective against a variety of ectoparasites and combination with the compound of Formula (I) would expand the scope of efficacy against these parasites. Particularly useful isoxazoline active agents that can
15 be combined with the compound include afoxolaner (including the substantially pure active enantiomer, esafoxolaner), sarolaner, fluralaner (including substantially pure active enantiomer), lotilaner and tigolaner. These active agents are described in US 7,964,204, US 2010/0254960 A1, US2011/0159107, US2012/0309620, US2012/0030841, US2010/0069247, WO 2007/125984, WO 2012/086462, US 8318757, US 8466115, US 8618126, US 8822466, US 8383659, US
20 8853186, US 9221835, US 2011/0144349, US 8,053,452; US 2010/0137612, US 8410153, US 2011/152081, WO 2012/089623, WO 2012/089622, US 8,119,671; US 7,947,715; WO 2102/120135, WO 2012/107533, WO 2011/157748, US 2011/0245274, US 2011/0245239, US 2012/0232026, US 2012/0077765, US 2012/0035122, US 2011/0251247, WO 2011/154433, WO 2011/154434, US 2012/0238517, US 2011/0166193, WO 2011/104088, WO 2011/104087, WO
25 2011/104089, US 2012/015946, US 2009/0143410, WO 2007/123855 A2, US 2011/0118212, US 7951828 & US 7662972, US 2010/0137372 A1, US 2010/0179194 A2, US 2011/0086886 A2, US 2011/0059988 A1, US 2010/0179195 A1, US 2015/0126523, WO 2010/003923, WO 2010/003877, WO 2010/072602, WO 2014/134236, WO 2017/147352, US 7897630, U.S. 7951828, WO 2020/007704 A1, WO 2021/028479 A1, WO 2014/122083 A1, WO 2016/177619
30 A1, WO 2014/012975 A1, WO 2015/078846 A1, WO 2015/078847 A1, WO 2015/150302 A1, WO 2015/181139 A1 and WO 2016/026789 A1, all of which are incorporated herein by reference

in their entirety.

In another embodiment of the description, nodulisporic acid and its derivatives may be added to the compositions of the description. These compounds are used to treat or prevent infections in humans and animals and are described, for example, in U.S. Patent No. 5,399,582, 5,962,499, 6,221,894 and 6,399,786, all of which are hereby incorporated by reference in their entirety. The compositions may include one or more of the known nodulisporic acid derivatives in the art, including all stereoisomers, such as those described in the literature cited above.

In another embodiment, anthelmintic compounds of the amino acetonitrile class (AAD) of compounds such as monepantel (ZOLVIX) and the like may be added to the compositions of the description. These compounds are described, for example, in US 7,084,280 to Ducray *et al.* (incorporated herein by reference); Sager *et al.*, *Veterinary Parasitology*, 2009, 159, 49-54; Kaminsky *et al.*, *Nature* vol. 452, 13 March 2008, 176-181.

The compositions of the description may also include aryloazol-2-yl cyanoethylamino compounds such as those described in US Patent No. 8,088,801 to Soll *et al.*, which is incorporated herein by reference, and thioamide derivatives of these compounds, as described in U.S. Patent No. 7,964,621, which is also incorporated herein by reference. Aryloazol-2-yl cyanoethylamino active agents, which are systemically-acting against endoparasites, may be used in combination with the compound in veterinary compositions of the description.

The compositions of the description may also include paraherquamide compounds and derivatives of these compounds, including derquantel (see Ostlind *et al.*, *Research in Veterinary Science*, 1990, 48, 260-61; and Ostlind *et al.*, *Medical and Veterinary Entomology*, 1997, 11, 407-408). The paraherquamide family of compounds is a known class of compounds that include a spirodioxepino indole core with activity against certain parasites (see *Tett. Lett.* 1981, 22, 135; *J. Antibiotics* 1990, 43, 1380, and *J. Antibiotics* 1991, 44, 492). In addition, the structurally related marcfortine family of compounds, such as marcfortines A-C, are also known and may be combined with the compositions of the description (see *J. Chem. Soc. – Chem. Comm.* 1980, 601 and *Tet. Lett.* 1981, 22, 1977). Further references to the paraherquamide derivatives can be found, for example, in WO 91/09961, WO 92/22555, WO 97/03988, WO 01/076370, WO 09/004432 and US 2010/0197624, U.S. Patent 5,703,078 and U.S. Patent 5,750,695, all of which are hereby incorporated by reference in their entirety.

In another embodiment of the description, the compositions may include a spinosyn active agent produced by the soil actinomycete *Saccharopolyspora spinosa* (see, for example Salgado V.L. and Sparks T.C., “*The Spinosyns: Chemistry, Biochemistry, Mode of Action, and Resistance*,” in *Comprehensive Molecular Insect Science*, vol. 6, pp. 137-173, 2005) or a semi-synthetic spinosoid active agent. The spinosyns are typically referred to as factors or components A, B, C, D, E, F, G, H, J, K, L, M, N, O, P, Q, R, S, T, U, V, W, or Y, and any of these components, or a combination thereof, may be used in the compositions of the description. The spinosyn compound may be a 5,6,5-tricyclic ring system, fused to a 12-membered macro cyclic lactone, a neutral sugar (rhamnose), and an amino sugar (forosamine). These and other natural spinosyn compounds, including 21-butenyl spinosyn produced by *Saccharopolyspora pagona*, which may be used in the compositions of the description, may be produced via fermentation by conventional techniques known in the art. Other spinosyn compounds that may be used in the compositions of the description are disclosed in U.S. Patent Nos. 5,496,931; 5,670,364; 5,591,606; 5,571,901; 5,202,242; 5,767,253; 5,840,861; 5,670,486; 5,631,155 and 6,001,981, all incorporated by reference herein in their entirety. The spinosyn compounds may include, but are not limited to, spinosyn A, spinosyn D, spinosad, spinetoram, or combinations thereof. Spinosad is a combination of spinosyn A and spinosyn D, and spinetoram is a combination of 3'-ethoxy-5,6-dihydro spinosyn J and 3'-ethoxy spinosyn L.

In general, additional active agents (other than the compound of formula (I) described above) is included in the dosage units of the description in an amount of between about 0.1 µg and about 1000 mg. Typically, the active agent may be included in an amount of about 10 µg to about 500 mg, about 10 µg to about 400 mg, about 1 mg to about 300 mg, about 10 mg to about 200 mg or about 10 mg to about 100 mg. More typically the additional active agent will be present in an amount of about 5 mg to about 50 mg in the compositions of the description.

The concentration of the additional active agent in the compositions of the description will typically be from about 0.01% to about 30% (w/w) depending on the potency of the active agent. In certain embodiments for very potent active agents including, but not limited to a macrocyclic lactone active agent, the concentration of the active agent will typically be from about 0.01% to about 10% (w/w), from about 0.01 to about 1% (w/w), from about 0.01% to about 0.5% (w/w), from about 0.1% to about 0.5% (w/w) or from about 0.01% to about 0.1% (w/w). In other embodiments, the concentration of the active agent will typically be from about 0.1% to about 2%

(w/w) or about 0.1% to about 1% (w/w).

In other embodiments, the additional active agent will typically be present at higher concentrations to achieve the desired efficacy. In some embodiments, the active agent will be present in a concentration of about 1% to about 30% (w/w), about 1% to about 20% (w/w) or about 1% to about 15% (w/w). In still other embodiments, the active agent will be present in a concentration of about 5% to about 20% (w/w) or about 5% to about 15% (w/w) in the composition.

In various embodiments of the description, an additional active agent may be included in the composition to deliver a dose of about 0.001 mg/kg to about 50 mg/kg or about 0.5 mg/kg to about 50 mg/kg of body weight of the animal. In other embodiments, the active agent will typically be present in an amount sufficient to deliver a dose of about 0.05 mg/kg to about 30 mg/kg, about 0.1 mg/kg to about 20 mg/kg. In other embodiments, the active agent will be present in an amount sufficient to deliver a dose of about 0.1 mg/kg to about 10 mg/kg, about 0.1 mg/kg to about 1 mg/kg or about 0.5 mg/kg to about 50 mg/kg per body weight of the animal.

In certain embodiments of the description where the additional active agent is a very potent compound such as a macrocyclic lactone or other potent compounds, the active agent will be present in a concentration to provide a dose of about 0.001 mg/kg to about 5 mg/kg, about 0.001 mg/kg to about 0.1 mg/kg or about 0.001 mg/kg to about 0.01 mg/kg. In still other embodiments, the active agent is present in an amount sufficient to deliver a dose of about 0.01 mg/kg to about 2 mg/kg or about 0.1 mg/kg to about 1 mg/kg per body weight of the animal. In still other embodiments, the additional active agent may be present in an amount to deliver a dose of about 1 µg/kg to about 200 µg/kg or about 0.1 mg/kg to about 1 mg/kg of weight of animal.

In addition to the other active agents mentioned above, combinations of two or more active agents may be used with the compounds of the description in a composition to treat a desired spectrum of pests and parasites. It would be well within the skill level of the practitioner to decide which individual compound can be used in the inventive composition to treat a particular infection of an insect.

The invention will now be further described by way of the following non-limiting examples.

EXAMPLES:Preparation Examples

The Examples that follow are intended to only illustrate the present invention without restricting it. The compounds of Formula (I) or pharmaceutically or a veterinarily acceptable salts thereof may be prepared by adopting one of the following reaction schemes. The starting materials for their preparation may be commercially available or can be prepared by methods known by persons of skill in the art and as described in the literature or may be an intermediate in any other of the schemes described herein. It will be appreciated that the following procedures may be modified by persons of skill in the art to prepare additional compounds according to the invention. For example, a person of skill in the art will understand that replacement of certain starting materials or the use of different intermediates will enable the preparation of different compounds of Formula (I).

The terms "ambient temperature" and "room temperature" are used interchangeably and designate a temperature of about 20 °C. Although the following subject matter is described in some detail by way of illustration and example for purposes of clarity of understanding, it will be understood by those skilled in the art that certain changes and modifications can be practiced within the scope of the Examples.

List of abbreviations:

ACN	acetonitrile
AIBN	azobisisobutyronitrile
BINAP	(2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl)
BSA	bovine serum albumin
BOC	<i>tert</i> -butoxycarbonyl
BOP-Cl	Bis(2-oxo-3-oxazolidinyl)phosphinic chloride
DAST	Diethylaminosulfur trifluoride
DCC	<i>N,N'</i> -Dicyclohexylcarbodiimide solution
DCM	dichloromethane
DEAD	Diethyl azodicarboxylate
DIEA	diisopropylethylamine
DMF	<i>N,N</i> -dimethylformamide

	DMAP	4-(Dimethylamino)pyridine
	DMSO	dimethylsulfoxide
	EDAC	<i>N</i> -(3-Dimethylaminopropyl)- <i>N'</i> -ethylcarbodiimide hydrochloride
	EDCI	1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide
5	ES	electrospray
	EtOAc or EA	ethyl acetate
	HATU	1-[bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5 b]pyridinium 3-oxide hexafluorophosphate
	HOBt or HOBt	1-hydroxybenzotriazole
10	KHMDS	potassium hexamethyldisilazide, more precisely potassium bis(trimethylsilyl)amide
	MeOH	methanol
	m-CPBA	m-Chloroperbenzoic acid
	NMO	N-Methylmorpholine-N-oxide
15	o/n	over night
	PE	petroleum ether
	Pd(dtbpf)Cl ₂	Dichloro[1,1'-bis (di-tert-butylphosphino) ferrocene] palladium(II)
	Pd ₂ dba ₃	Tris(dibenzylideneacetone)dipalladium(0)
	Pd(dppf)Cl ₂	[1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane
20	TBAF	<i>tert</i> -butyl ammonium fluoride
	TfO	triflate
	THF	tetrahydrofuran
	TLC	thin-layer chromatography
25		

Some examples of Formula (I) are derived after separation of racemic mixtures and obtained as enantiomerically pure products. The stereochemistry is in some cases arbitrarily assigned and the respective compound characterized by analytical methods as described below:

Method A

Gradient/Solvent Time [min]	% Sol [scCO ₂]	% Sol [MeOH 20mM NH ₃]	Flow [ml/min]	Temp [°C]	Back pressure [PSI]
0.0	75.0	25.0	2.0	40.0	2175.0
4.0	75.0	25.0	2.0	40.0	2175.0

CHIRAL ART® Cellulose SJ_3 x 100 mm_3 µm (Agilent)

Method B

Gradient/Solvent Time [min]	% Sol [scCO ₂]	% Sol [MeOH 20mM NH ₃]	Flow [ml/min]	Temp [°C]	Back pressure [PSI]
0.0	85.0	15.0	2.0	40.0	2175.0
4.0	85.0	15.0	2.0	40.0	2175.0

CHIRAL ART® Cellulose SJ_3 x 100 mm_3 µm (Agilent)

Method C

Gradient/Solvent Time [min]	% Sol [scCO ₂]	% Sol [MeOH 20mM NH ₃]	Flow [ml/min]	Temp [°C]	Back pressure [PSI]
0.0	80.0	20.0	2.0	40.0	2175.0
4.0	80.0	20.0	2.0	40.0	2175.0

CHIRAL ART® Cellulose SJ_3 x 100 mm_3 µm (Agilent)

Method D

Gradient/Solvent Time [min]	% Sol [scCO ₂]	% Sol IPA 20mM NH ₃]	Flow [ml/min]	Temp [°C]	Back pressure [PSI]
0.0	70.0	30.0	4.0	40.0	2175.0
10.0	70.0	30.0	4.0	40.0	2175.0

CHIRAL ART® Cellulose SB_4.6 x 250 mm_5 µm (Agilent)


Method E

Gradient/Solvent Time [min]	% Sol [scCO ₂]	% Sol IPA [MeOH 20mM NH ₃]	Flow [ml/min]	Temp [°C]	Back pressure [PSI]
0.0	70.0	30.0	4.0	40.0	2175.0
10.0	70.0	30.0	4.0	40.0	2175.0


CHIRAL ART® Cellulose SB_4.6 x 250 mm_5 µm (Agilent)

Preparation Example 1: The following example compounds can be synthesized by adopting the subsequent schemes 3 and 4 shown below for compound 175 by someone skilled in the art: 271, 272, 273, 274, 275, 276, 279, 293, 294, 295, 296, 304, 305, 307, 308, 322, 344, 345, 364, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855, 856, 857, 858, 859, 860, 861, 862, 863, 864, 865, 866, 867, 868, 869, 870, 871, 872, 873, 874, 875, 876, 877, 878, 879, 880, 881, 882, 883, 884, 885, 886, 887, 888, 889, 890, 891, 892, 893, 894, 895, 896, 897, 898, 899, 900, 901, 902, 903, 904, 905, 906, 907, 908, 909, 910, 911, 912, 913, 914, 915, 916, 917, 918, 919, 920, 921, 922, 923, 924, 925, 926, 927, 928, 929, 930, 931, 932, 933, 934, 935, 936, 937, 938, 939, 940, 941, 942, 943, 944, 945, 946, 947, 948, 949, 950, 951, 952, 953, 954, 955, 956, 957, 958, 959, 960, 961, 962, 963, 964, 965, 966, 967, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 978, 979, 980, 981, 982, 983, 984, 985, 986, 987, 988, 989, 990, 991, 992, 993, 994, 995, 996, 997, 998, 999, 1000.


Scheme 3



Scheme 4




1. Synthesis of ethyl 1-aminoimidazole-2-carboxylate




Into a 3000-mL 3-necked round-bottom flask, was placed DMF (2000 mL), ethyl 1H-imidazole-2-carboxylate (3-1, 50.0 g, 356.7 mmol, 1.0 equiv). This was followed by the addition of NaH (21.0 g, 875.0 mmol, 2.4 equiv), in portions at room temperature. To this was added amino diphenylphosphinate (119.0 g, 510.2 mmol, 1.4 equiv), in portions at 0 °C. The resulting solution was stirred for 3 hr at room temperature. The resulting mixture was concentrated under vacuum. The solids were filtered off. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with dichloromethane/methanol (10:1). This resulted in 40 g (72.2%) of ethyl 1-aminoimidazole-2-carboxylate (3-2) as a white solid.

2. Synthesis of ethyl 1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-3)



Into a 500-mL round-bottom flask, was placed DMF (200.0 mL), ethyl 1-aminoimidazole-2-carboxylate (3-2, 35.0 g, 225.5 mmol, 1.0 equiv), Boc₂O (63.9 g, 293.2 mmol, 1.3 equiv), DMAP (13.7 g, 112.7 mmol, 0.5 equiv). The resulting solution was stirred for 2 hr at 80 °C. The reaction was then quenched by the addition of 500 mL of water. The resulting solution was extracted with 3x200 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:3). This resulted in 28 g (48.6%) of ethyl 1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-3) as a white solid.


3. Synthesis of ethyl 4-bromo-1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-4)



Into a 250-mL round-bottom flask, was placed DMF (100.0 mL), ethyl 1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-3, 15.0 g, 58.7 mmol, 1.0 equiv), NBS (10.4 g,


58.8 mmol, 1.0 equiv). The resulting solution was stirred for 1 days at room temperature. The reaction was then quenched by the addition of 300 mL of water. The resulting solution was extracted with 3x100 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:4). This resulted in 12 g (61.1%) of ethyl 4-bromo-1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-4) as colorless oil.

4. Synthesis of ethyl 3-[4-bromo-1-[(tert-butoxycarbonyl)amino]imidazol-2-yl]-3-oxopropanoate (3-5)




Into a 1000-mL 3-necked round-bottom flask, was placed THF (400.0 mL), ethyl 4-bromo-1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-4, 12.0 g, 35.9 mmol, 1.0 equiv). This was followed by the addition of t-BuOK (60.0 g, 534.7 mmol, 14.9 equiv), in portions at 0 °C in 30 min. To this was added ethyl acetate (32.0 g, 363.2 mmol, 10.1 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred for 2 hr at room temperature. The reaction was then quenched by the addition of HCl (1M). The resulting solution was extracted with 3x50 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 6.5 g (48.1%) of ethyl 3-[4-bromo-1-[(tert-butoxycarbonyl)amino]imidazol-2-yl]-3-oxopropanoate (3-5) as yellow oil.

5. Synthesis of ethyl 2-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (3-6)




Into a 100-mL round-bottom flask, was placed DCM (30.0 mL), ethyl 3-[4-bromo-1-[(tert-butoxycarbonyl)amino]imidazol-2-yl]-3-oxopropanoate (3-5, 6.5 g, 17.2 mmol, 1.0 equiv), DMF-DMA (5.00 mL, 37.3 mmol, 2.2 equiv). The resulting solution was stirred for 2 hr at room temperature. The resulting mixture was concentrated under vacuum. The crude product was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O:ACN=90:10 increasing to H₂O:ACN=50:50 within 15 min; Detector, 254nm. This resulted in 3.3 g (66.7%) of ethyl 2-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (3-6) as a white solid.

6. Synthesis of ethyl 8-hydroxy-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (3-7)




Into a 250-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed dioxane (60.0 mL), ethyl 2-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (3-6, 3.0 g, 10.5 mmol, 1.0 equiv), trimethyl-1,3,5,2,4,6-trioxatriborinane (2.6 g, 20.9 mmol, 2.0 equiv), Pd(PPh₃)₄ (1.2 g, 1.0 mmol, 0.1 equiv), K₂CO₃ (4.3 g, 31.3 mmol, 3.0 equiv). The resulting solution was stirred for 4 hr at 100 degrees C. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:2). This resulted in 2 g (86.2%) of ethyl 8-hydroxy-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (3-7) as a yellow solid.

7. Synthesis of ethyl 2-methyl-8-(trifluoromethanesulfonyloxy)imidazo[1,2-b]pyridazine-7-carboxylate (4-1)




Into a 50-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed DCM (20.0 mL), ethyl 8-hydroxy-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (3-7, 200.0 mg, 0.9 mmol, 1.0 equiv), TEA (457.0 mg, 4.5 mmol, 5.0 equiv). This was followed by the addition of TiF_4 (765.6 mg, 2.7 mmol, 3.0 equiv) dropwise with stirring at -78 degrees C. The resulting solution was stirred for 1 hr at -50 degrees C. The reaction was then quenched by the addition of water/ice. The resulting solution was extracted with 3x20 mL of dichloromethane and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 200 mg (62.6%) of ethyl 2-methyl-8-(trifluoromethanesulfonyloxy)imidazo[1,2-b]pyridazine-7-carboxylate (4-1) as brown oil.

8. Synthesis of ethyl 2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (4-2)




Into a 50-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed THF (10.0 mL), H_2O (2.0 mL), ethyl 2-methyl-8-(trifluoromethanesulfonyloxy)imidazo[1,2-b]pyridazine-7-carboxylate (4-1, 200.0 mg, 0.6 mmol, 1.0 equiv), 4,4,5,5-tetramethyl-2-(prop-1-en-2-yl)-1,3,2-dioxaborolane (179.0 mg, 1.0 mmol, 1.9 equiv), Pd(dtbpf)Cl_2 (37.2 mg, 0.06 mmol, 0.10 equiv), K_2CO_3 (234.0 mg, 1.7 mmol, 3.0 equiv). The resulting solution was stirred for 2 hr at room temperature. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:4). This resulted in 90 mg (64.8%) of ethyl 2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (4-2) as a white solid.

9. Synthesis of ethyl 8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate




Into a 50-mL round-bottom flask, was placed EA (5.0 mL), ethyl 2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (4-2, 90.0 mg, 0.4 mmol, 1.0 equiv), PtO_2 (30.00 mg, 0.1 mmol, 0.4 equiv), and an atmosphere of H_2 (g) by balloon. The resulting solution was stirred for 1 hr at 50 °C. The solids were filtered out. The resulting mixture was concentrated under vacuum. This resulted in 90 mg (99.2%) of ethyl 8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-3) as a white solid.

10. Synthesis of ethyl 3-bromo-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate




Into a 50-mL round-bottom flask, was placed CHCl_3 (10.0 mL), ethyl 8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-3, 90.0 mg, 0.4 mmol, 1.0 equiv), NBS (90.0 mg, 0.5 mmol, 1.4 equiv). The resulting solution was stirred for 1 hr at 80 degrees C. The reaction was then quenched by the addition of 10 mL of water. The resulting solution was extracted with 3x10 mL of dichloromethane and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 100 mg (crude) of ethyl 3-bromo-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-4) as a white solid.

11. Synthesis of ethyl 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-5)



Into a 50-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed dioxane (10.0 mL), H_2O (3.0 mL), ethyl 3-bromo-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-4, 90.0 mg, 0.3 mmol, 1.0 equiv), 3,5-difluorophenylboronic acid (87.5 mg, 0.5 mmol, 2.0 equiv), $\text{Pd}(\text{dtbpf})\text{Cl}_2$ (18.0 mg, 0.03 mmol, 0.1 equiv), K_2CO_3 (114.6 mg, 0.829 mmol, 3.0 equiv). The resulting solution was stirred for 3 hr at 100 °C. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:3). This resulted in 80 mg (80.7%) of ethyl 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-5) as a yellow solid.


12. Synthesis of 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (4-6)



Into a 50-mL round-bottom flask, was placed H_2O (1.0 mL), i-PrOH (5.0 mL), ethyl 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-5, 80 mg, 0.2 mmol, 1.0 equiv), $\text{LiOH}\cdot\text{H}_2\text{O}$ (28.0 mg, 0.7 mmol, 3.0 equiv). The resulting solution was stirred for 2 hr at 50 °C. The pH value of the solution was adjusted to 4 with HCl (2M). The resulting solution was extracted with 3x20 mL of ethyl acetate and the organic layers combined and dried

over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 60 mg (90%) of 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (4-6) as a white solid.

13. Synthesis of 3-(3,5-difluorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (compound 175)




Into a 50-mL round-bottom flask, was placed DMA (5.0 mL), 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (4-6, 60.0 mg, 0.2 mmol, 1.0 equiv), (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (57.2 mg, 0.4 mmol, 2.1 equiv), HATU (138.0 mg, 0.4 mmol, 2.0 equiv), DIEA (70.0 mg, 0.5 mmol, 3.0 equiv). The resulting solution was stirred for 1 hr at room temperature. The mixture was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O:ACN=60:40 increasing to H₂O:ACN=10:90 within 25 ; Detector, 220nm. This resulted in 38.6 mg (46.0%) of 3-(3,5-difluorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-


methylimidazo[1,2-b]pyridazine-7-carboxamide (175) as a white solid (300 MHz, CD₃OD, ppm) δ 8.33 (s, 1H), 7.41-7.33 (m, 3H), 7.21-7.16 (m, 1H), 7.06-6.97 (m, 1H), 6.95-6.92 (m, 1H), 6.82 (d, J = 8.1 Hz, 1H), 5.32 (t, J = 5.1 Hz, 1H), 4.33-4.25 (m, 2H), 3.70-3.65 (m, 1H), 2.62 (s, 3H), 2.33-2.28 (m, 1H), 2.23-2.19 (m, 1H), 1.62 (t, J = 6.6 Hz, 6H).

Preparation Example 2: Similarly to the process described in Preparation Example 1, the process depicted in Schemes 5-7 below may be used to prepare compound 271:


Scheme 5



Scheme 6




Scheme 7




The process in Schemes 3-7 described above may be modified according to methods known to persons of skill in the art to incorporate different functional groups in the core structure. For example, intermediate 3-6 may be reacted with an alternate coupling partner to introduce a different R^2 substituent. Similarly, intermediates 4-1, 271-1 and 4-4 may be reacted with alternate compounds to introduce different R^1 and R^3 substituents.

1. Synthesis of ethyl 1-aminoimidazole-2-carboxylate (3-2)



Into a 5-L round-bottom flask, was placed DMF (4000.00 mL, 51687.010 mmol, 144.87 equiv), ethyl 1H-imidazole-2-carboxylate (50.00 g, 356.781 mmol, 1.00 equiv). This was followed by the addition of NaH (21.00 g, 875.083 mmol, 2.45 equiv), in portions at room temperature in 30 min. To this was added amino diphenylphosphinate (120.00 g, 514.564 mmol, 1.44 equiv), in portions at room temperature. The resulting solution was stirred for 2 hr at room temperature. The resulting mixture was dried with nitrogen (blowing nitrogen). The residue was dissolved in 2000 mL of EA. The solids were filtered out. The filtrate was concentrated under vacuum. This resulted in 42 g (75.87%) of ethyl 1-aminoimidazole-2-carboxylate (3-2) as a white solid.


2. Synthesis of ethyl 1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-3)



Into a 1000-mL round-bottom flask, was placed DMF (500.00 mL, 6460.876 mmol, 28.64 equiv), ethyl 1-aminoimidazole-2-carboxylate (35.00 g, 225.578 mmol, 1.00 equiv), Boc₂O (73.30 g, 335.858 mmol, 1.49 equiv), DMAP (13.78 g, 112.796 mmol, 0.50 equiv). The resulting solution


was stirred for 2 hrs at 80 degrees C. The reaction was then quenched by the addition of water/ice. The resulting solution was extracted with 3x500 mL of ethyl acetate and the organic layers combined and washed with 2x500 ml of H₂O, 1x500 ml of brine. Organic Layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:2). This resulted in 30 g (52.10%) of ethyl 1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-3) as a white solid.

3. Synthesis of ethyl 4-bromo-1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-4)




Into a 1000-mL round-bottom flask, was placed DMF (400.00 mL, 5168.701 mmol, 26.39 equiv), ethyl 1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (50.00 g, 195.868 mmol, 1.00 equiv), NBS (40.00 g, 0.225 mmol). The resulting solution was stirred overnight at room temperature. The reaction was then quenched by the addition of water/ice. The resulting solution was extracted with 3x100 mL of ethyl acetate and the organic layers combined and washed with 2x500 ml of H₂O, 1x500 ml of brine. Organic Layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1). This resulted in 30 g (45.83%) of ethyl 4-bromo-1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-4) as yellow oil.

4. Synthesis of ethyl 3-[4-bromo-1-[(tert-butoxycarbonyl)amino]imidazol-2-yl]-3-oxopropanoate (3-5)



Into a 1000-mL 3-necked round-bottom flask, was placed THF (500.00 mL, 6171.495 mmol, 58.92 equiv), ethyl 4-bromo-1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (35.00 g, 104.737 mmol, 1.00 equiv), EA (93.50 g, 1061.233 mmol, 10.13 equiv). This was followed by the addition of t-BuOK (170.00 g, 1514.989 mmol, 14.46 equiv), in portions at 0 degrees C. The resulting solution was stirred for 1 hr at room temperature. The reaction was then quenched by the addition of NH₄Cl (aq.). The resulting solution was extracted with 3x300 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was dissolved in 500 ml of hexane. The solids were collected by filtration. This resulted in 22 g (55.83%) of ethyl 3-[4-bromo-1-[(tert-butoxycarbonyl)amino]imidazol-2-yl]-3-oxopropanoate (3-5) as a white solid.


5. Synthesis of ethyl 2-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (3-6)



Into a 100-mL round-bottom flask, was placed DCM (30.00 mL, 471.901 mmol, 27.31 equiv), ethyl 3-[4-bromo-1-[(tert-butoxycarbonyl)amino]imidazol-2-yl]-3-oxopropanoate (6.50 g, 17.278 mmol, 1.00 equiv), DMF-DMA (5.00 mL, 37.344 mmol, 2.16 equiv). The resulting solution was stirred for 2 hr at room temperature. The resulting mixture was concentrated under vacuum. The crude product was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O (0.1% TFA):ACN=90:10 increasing to H₂O

(0.1%TFA):ACN=50:50 within 15 min; Detector, 254 nm. This resulted in 3.3 g (66.76%) of ethyl 2-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (3-6) as a white solid.

6. Synthesis of ethyl 8-hydroxy-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (3-7)



Into a 250-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed dioxane (100.0 mL, 1180.408 mmol, 67.54 equiv), ethyl 2-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (5.00 g, 17.477 mmol, 1.00 equiv), trimethyl-1,3,5,2,4,6-trioxatriborinane (6.60 g, 52.577 mmol, 3.01 equiv), $\text{Pd(PPh}_3)_4$ (2.00 g, 1.731 mmol, 0.10 equiv), K_3PO_4 (7.20 g, 52.096 mmol, 2.98 equiv). The resulting solution was stirred for 4 hrs at room temperature. The resulting mixture was concentrated under vacuum. The crude product was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H_2O (0.1% $\text{NH}_3\cdot\text{H}_2\text{O}$):ACN=100:0 increasing to H_2O (0.1% $\text{NH}_3\cdot\text{H}_2\text{O}$):ACN=50:50 within 10 min; Detector, 254 nm. This resulted in 0.75 g (25.6%) of ethyl 8-hydroxy-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (3-7) as a yellow solid.

7. Synthesis of ethyl 8-chloro-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (271-1)




Into a 100-mL round-bottom flask, was placed CHCl_3 (20.00 mL, 247.952 mmol, 18.28 equiv), ethyl 8-hydroxy-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (3.00 g, 13.561 mmol, 1.00 equiv), oxalyl chloride (6.00 g, 47.274 mmol, 3.49 equiv), DMF (0.10 mL). The resulting solution was stirred for 2 hrs at 80 degrees C. The resulting mixture was concentrated under vacuum. This

resulted in 3.6 g (crude) of ethyl 8-chloro-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (271-1) as a yellow solid.

8. Synthesis of ethyl 2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (4-2)


5



Into a 100-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed THF (30.00 mL, 370.290 mmol, 88.74 equiv), H₂O (5.00 mL, 277.542 mmol, 66.52 equiv), ethyl 8-chloro-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (1.00 g, 4.173 mmol, 1.00 equiv), 4,4,5,5-tetramethyl-2-(prop-1-en-2-yl)-1,3,2-dioxaborolane (2.11 g, 12.557 mmol, 3.01 equiv), Pd(dtbpf)Cl₂ (410.00 mg, 0.629 mmol, 0.15 equiv), K₃PO₄ (2.66 g, 12.531 mmol, 3.00 equiv). The resulting solution was stirred for 2 hrs at 70 degrees C. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:2). This resulted in 500 mg (48.85%) of ethyl 2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (4-2) as yellow oil.

15

9. Synthesis of ethyl 8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-3)




Into a 50-mL round-bottom flask, was placed EA (5.00 mL, 0.057 mmol, 0.03 equiv), ethyl 2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (500.00 mg, 2.038 mmol, 1.00 equiv), PtO₂ (100.00 mg, 0.440 mmol, 0.22 equiv), to the above H₂ (g) was introduced in. The

20


resulting solution was stirred for 2 hrs at 50 degrees C. The solids were filtered out. The filtrate was concentrated under vacuum. This resulted in 350 mg (69.43%) of ethyl 8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-3) as yellow oil.

10. Synthesis of ethyl 3-bromo-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-4)




Into a 50-mL round-bottom flask, was placed CHCl₃ (5.00 mL, 61.988 mmol, 47.90 equiv), ethyl 8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (320.00 mg, 1.294 mmol, 1.00 equiv), NBS (253.70 mg, 1.425 mmol, 1.10 equiv). The resulting solution was stirred for 1 hr at 80 degrees C. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:5). This resulted in 350 mg (82.92%) of ethyl 3-bromo-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-4) as yellow oil.

11. Synthesis of ethyl 3-(3,5-dichlorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (271-2)



Into a 8-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed THF (1.00 mL, 12.343 mmol, 44.74 equiv), H₂O (0.20 mL, 11.102 mmol, 40.24 equiv), ethyl 3-bromo-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (90.00 mg, 0.276 mmol, 1.00 equiv), 3,5-dichlorophenylboronic acid (53.17 mg, 0.279 mmol, 1.01 equiv), Pd(dtbpf)Cl₂ (17.98 mg, 0.028 mmol, 0.10 equiv), K₂CO₃ (75.88 mg, 0.549 mmol, 1.99 equiv). The resulting solution was stirred for 2 hrs at 50° C. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:4). This resulted in 70 mg (64.67%) of ethyl 3-(3,5-dichlorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (271-2) as colorless oil.


12. Synthesis of 3-(3,5-dichlorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (271-3)



Into a 50-mL round-bottom flask, was placed H₂O (0.50 mL, 27.754 mmol, 155.53 equiv), EtOH (2.00 mL, 0.043 mmol, 0.24 equiv), ethyl 3-(3,5-dichlorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (70.00 mg, 0.178 mmol, 1.00 equiv), LiOH.H₂O

(22.50 mg, 0.536 mmol, 3.00 equiv). The resulting solution was stirred for 1 hr at room temperature. The pH value of the solution was adjusted to 5 with HCl (6 mol/L). The resulting solution was extracted with 3x20 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 50 mg (76.93%) of 3-(3,5-dichlorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (271-3) as yellow oil.

Synthesis of 3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (Compound 271)



Into a 50-mL round-bottom flask, was placed DMA (1.00 mL, 10.755 mmol, 87.05 equiv), 3-(3,5-dichlorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (45.00 mg, 0.124 mmol, 1.00 equiv), (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (29.30 mg, 0.196 mmol, 1.59 equiv), HATU (93.90 mg, 0.247 mmol, 2.00 equiv), DIEA (47.80 mg, 0.370 mmol, 2.99 equiv). The resulting solution was stirred for 1 hr at room temperature. The crude mixture was purified by Flash-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O (0.1% NH₃·H₂O):ACN=50:50 increasing to H₂O (0.1% NH₃·H₂O):ACN=10:90 within 20 min; Detector, 254 nm. This resulted in 42.2 mg (68.95%) of 3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (271) as a white solid.

¹H NMR (300 MHz, CDCl₃, *ppm*) δ 8.29 (s, 1H), 7.59 (d, *J* = 1.9 Hz, 2H), 7.43 (t, *J* = 1.9 Hz, 1H), 7.34-7.18 (m, 2H), 6.97 (td, *J* = 7.5, 1.2 Hz, 1H), 6.89 (dd, *J* = 8.2, 1.2 Hz, 1H), 6.15 (d, *J* = 7.6 Hz, 1H), 5.38 (q, *J* = 5.6 Hz, 1H), 4.40-4.36 (m, 1H), 4.28-4.14 (m, 1H), 3.78-3.73 (m, 1H), 2.64 (s, 3H), 2.44-2.41 (m, 1H), 2.26-2.22 (m, 1H), 1.65 (t, *J* = 6.7 Hz, 6H); (ES, *m/z*): 495 [M+H]⁺.

As noted above, the following compounds may be prepared according to schemes 3 to 7:

Compound	¹ H NMR Spectra
272	(300 MHz, CDCl ₃ , ppm) δ 8.30 (s, 1H), 7.49 (s, 1H), 7.37-7.30 (m, 2H), 7.29-7.21 (m, 1H), 7.20-7.15 (m, 1H), 6.99-6.94 (m, 1H), 6.89 (d, J = 8.4 Hz, 1H), 6.19 (d, J = 7.5 Hz, 1H), 5.38 (q, J = 6.9 Hz, 1H), 4.37-4.35 (m, 1H), 4.25-4.14 (m, 1H), 3.79-3.74 (m, 1H), 2.64 (s, 3H), 2.43-2.40 (m, 1H), 2.26-2.21 (m, 1H), 1.64 (t, J = 6.6 Hz, 6H)
273	(300 MHz, CDCl ₃ , ppm) δ 8.31 (s, 1H), 7.53-7.40 (m, 3H), 7.40-7.31 (m, 1H), 7.29-7.17 (m, 2H), 6.99-6.93 (m, 1H), 6.88 (d, J = 8.4 Hz, 1H), 6.26 (d, J = 7.2 Hz, 1H), 5.38 (q, J = 7.2 Hz, 1H), 4.39-4.33 (m, 1H), 4.25-4.20 (m, 1H), 3.78-3.73 (m, 1H), 2.63 (s, 3H), 2.44-2.37 (m, 1H), 2.26-2.23 (m, 1H), 1.63 (t, J = 6.9 Hz, 6H)
274	(300 MHz, CDCl ₃ , ppm) δ 8.24 (d, J = 2.4 Hz, 1H), 7.64 (dd, J = 7.2, 2.1 Hz, 1H), 7.41-7.35 (m, 2H), 7.33-7.29 (m, 1H), 7.27-7.20 (m, 1H), 6.98-6.93 (m, 1H), 6.90-6.87 (m, 1H), 6.20 (d, J = 7.5 Hz, 1H), 5.38 (q, J = 7.2 Hz, 1H), 4.37-4.34 (m, 1H), 4.27-4.13 (m, 1H), 3.79-3.74 (m, 1H), 2.47 (s, 3H), 2.41-2.39 (m, 1H), 2.27-2.20 (m, 1H), 1.69-1.63 (m, 6H)
275	(300 MHz, DMSO-d ₆ , ppm) δ 9.16 (d, J = 8.4 Hz, 1H), 8.53 (s, 1H), 7.51-7.47 (m, 2H), 7.36-7.30 (m, 1H), 7.16 (dd, J = 9.3, 2.7 Hz, 1H), 7.08-7.01 (m, 1H), 6.83 (dd, J = 9.0, 4.8 Hz, 1H), 5.24 (q, J = 7.2 Hz, 1H), 4.28-4.22 (m, 2H), 3.62-3.57 (m, 1H), 2.57 (s, 3H), 2.30-2.16 (m, 1H), 2.11-1.95 (m, 1H), 1.55 (t, J = 7.4 Hz, 6H)
276	(300 MHz, CDCl ₃ , ppm) δ 8.36 (s, 1H), 7.28-7.22 (m, 2H), 7.19-7.16 (m, 2H), 7.03-6.95 (m, 2H), 6.91-6.88 (m, 1H), 6.09 (d, J = 7.2 Hz, 1H), 5.39 (q, J = 7.5 Hz, 1H), 4.41-4.35 (m, 1H), 4.24-4.17 (m, 1H), 3.77-3.72 (m, 1H), 2.44-2.39 (m, 1H), 2.28-2.24 (m, 1H), 1.68 (t, J = 6.9 Hz, 6H)
279	(300 MHz, DMSO-d ₆ , ppm) δ 8.99 (d, J = 8.4 Hz, 1H), 8.60 (s, 1H), 7.58-7.43 (m, 2H), 7.43-7.25 (m, 2H), 7.24-7.12 (m, 1H), 6.93 (t, J = 6.9 Hz, 1H), 6.79 (d, J = 8.1 Hz, 1H), 5.45 (s, 1H), 5.26-5.16 (m, 2H), 4.30-4.17 (m, 2H), 2.55 (s, 3H), 2.30 (s, 3H), 2.23-2.09 (m, 1H), 2.00-1.95 (m, 1H)

293	(300 MHz, CDCl ₃ , ppm) δ 8.16 (d, J = 3.6 Hz, 1H), 7.60 (dd, J = 7.5, 1.5 Hz, 1H), 7.40-7.29 (m, 2H), 6.99-6.89 (m, 2H), 6.83-6.79 (m, 1H), 6.24-6.22 (m, 1H), 5.36-5.33 (m, 1H), 4.34-4.28 (m, 1H), 4.19-4.11 (m, 1H), 3.79-3.74 (m, 1H), 2.46 (s, 3H), 2.39-2.30 (m, 1H), 2.23-2.07 (m, 1H), 1.72-1.58 (m, 6H)
294	(300 MHz, CDCl ₃ , ppm) δ 8.27 (s, 1H), 7.58 (d, J = 1.8 Hz, 2H), 7.42 (t, J = 1.8 Hz, 1H), 7.01-6.90 (m, 2H), 6.84-6.80 (m, 1H), 6.31 (br s, 1H), 5.39-5.34 (m, 1H), 4.36-4.30 (m, 1H), 4.24-4.17 (m, 1H), 3.79-3.75 (m, 1H), 2.63 (s, 3H), 2.42-2.36 (m, 1H), 2.22-2.16 (m, 1H), 1.64 (t, J = 6.6 Hz, 6H)
295	(300 MHz, CDCl ₃ , ppm) δ 8.37 (s, 1H), 7.53-7.51 (m, 3H), 7.26-7.23 (m, 1H), 7.02-6.95 (m, 1H), 6.92-6.89 (m, 1H), 6.06 (d, J = 7.2 Hz, 1H), 5.43-5.37 (s, 1H), 4.42-4.35 (m, 1H), 4.24-4.17 (m, 1H), 3.77-3.70 (m, 1H), 2.47-2.41 (m, 1H), 2.27-2.22 (m, 1H), 1.68 (t, J = 6.6 Hz, 6H)
296	(300 MHz, CDCl ₃ , ppm) δ 8.37 (s, 1H), 7.53-7.51 (m, 3H), 7.00-6.93 (m, 2H), 6.88-6.83 (m, 1H), 6.04 (d, J = 7.8 Hz, 1H), 5.43-5.37 (m, 1H), 4.38-4.33 (m, 1H), 4.26-4.12 (m, 1H), 3.76-3.71 (m, 1H), 1.68 (t, J = 6.9 Hz, 6H)
304	(300 MHz, CDCl ₃ , ppm) δ 8.40 (s, 1H), 8.16 (s, 2H), 7.42 (t, J = 2.1 Hz, 1H), 7.03-6.93 (m, 2H), 6.88-6.83 (m, 1H), 6.06 (d, J = 7.8 Hz, 1H), 5.48-5.32 (m, 1H), 4.39-4.33 (m, 1H), 4.24-4.18 (m, 1H), 3.76-3.69 (m, 1H), 2.46-2.39 (m, 1H), 2.29-2.18 (m, 1H), 1.72-1.68 (m, 6H)
305	(400 MHz, Chloroform-d, ppm) δ : 8.37 (s, 1H), 7.66 (s, 2H), 7.55 (s, 1H), 7.01-6.84 (m, 4H), 6.10 (d, J = 8.3 Hz, 1H), 5.35 (d, J = 8.3 Hz, 1H), 4.39-4.34 (m, 1H), 4.26-4.15 (m, 1H), 3.81-3.74 (m, 1H), 2.44-2.38 (m, 1H), 2.21-2.20 (m, 1H), 1.70-1.55 (m, 6H)
307	(300 MHz, CDCl ₃ , ppm): δ 8.86 (s, 1H), 7.33-7.30 (m, 2H), 7.04-6.80 (m, 5H), 5.69 (s, 1H), 5.50-5.30 (m, 2H), 4.35-4.30 (m, 1H), 4.25-4.00 (m, 1H), 2.68 (s, 3H), 2.45-2.25 (m, 4H), 2.20-2.10 (m, 1H)
308	(300 MHz, Chloroform-d, ppm): δ 8.84 (s, 1H), 7.63 (s, 2H), 7.46 (s, 1H), 7.02-6.78 (m, 4H), 5.68 (s, 1H), 5.45-5.31 (m, 2H), 4.37-4.30 (m, 1H), 4.21-4.07 (m, 1H), 2.66 (s, 3H), 2.45-2.30 (m, 4H), 2.20-2.10 (m, 1H)

322	(300 MHz, Chloroform-d, ppm): δ 8.52 (d, J = 5.1 Hz, 1H), 8.32 (s, 1H), 7.80 (s, 1H), 7.70-7.60 (m, 1H), 7.40-7.20 (m, 2H), 7.05-6.95 (m, 1H), 6.93-6.87 (m, 1H), 6.20-6.00 (m, 1H), 5.45-5.30 (m, 1H), 4.45-4.35 (m, 1H), 4.30-4.15 (m, 1H), 3.83-3.74 (m, 1H), 2.71 (s, 3H), 2.52-2.37 (m, 1H), 2.30-2.15 (m, 1H), 1.75-1.65 (m, 6H)
344	(300 MHz Chloroform-d, ppm): δ 8.23 (s, 1H), 7.14-6.91 (m, 4H), 6.90-6.75 (m, 1H), 6.10 (d, J = 7.8 Hz, 1H), 5.45-5.31 (m, 1H), 4.45-4.32 (m, 1H), 4.30-4.10 (m, 1H), 3.85-3.65 (m, 1H), 2.54 (s, 3H), 2.44-2.30 (m, 1H), 2.24-2.17 (m, 1H), 1.67 (t, J = 6.6 Hz, 6H)
345	(300 MHz, Chloroform-d, ppm): δ 8.23 (s, 1H), 7.28-7.20 (m, 2H), 7.10-7.00 (m, 2H), 6.99-6.93 (m, 1H), 6.90-6.87 (m, 1H), 6.11 (d, J = 7.2 Hz, 1H), 5.40-5.30 (m, 1H), 4.40-4.34 (m, 1H), 4.24-4.15 (m, 1H), 3.85-3.74 (m, 1H), 2.53 (s, 3H), 2.50-2.35 (m, 1H), 2.30-2.15 (m, 1H), 1.66 (t, J = 6.6 Hz, 6H)
364	(300 MHz, Chloroform-d, ppm): δ 8.47 (s, 1H), 8.34-8.24 (m, 2H), 7.60 (d, J = 1.8 Hz, 2H), 7.42 (t, J = 1.8 Hz, 1H), 6.79 (d, J = 5.7 Hz, 1H), 6.55-6.40 (m, 1H), 5.47-5.40 (m, 1H), 4.49-4.41 (m, 1H), 4.40-4.25 (m, 1H), 3.82-3.70 (m, 1H), 2.64 (s, 3H), 2.50-2.25 (m, 2H), 1.68-1.64 (m, 6H)
527	(300 MHz DMSO-d ₆ , ppm): δ 8.08 (s, 1H), 7.15-7.44 (m, 3H), 7.00-6.79 (m, 2H), 6.10 - 6.21 (m, 1H), 5.36 - 5.43 (m, 1H), 4.30 - 4.49 (m, 1H), 4.10 - 4.18 (m, 1H), 2.40-2.38 (m, 4H), 2.28-2.26 (m, 1H), 1.80 (bs, 9H)
528	(300 MHz, DMSO-d ₆ , ppm) δ 9.18 (d, J = 8.1 Hz, 1H), 8.37 (s, 1H), 7.79 (d, J = 8.7 Hz, 2H), 7.32 (d, J = 7.5 Hz, 1H), 7.17 (t, J = 7.5 Hz, 1H), 6.92 (t, J = 7.5 Hz, 1H), 6.79 (d, J = 8.1 Hz, 1H), 5.26-5.20 (m, 1H), 4.30-4.14 (m, 2H), 3.61-3.51 (m, 1H), 2.30 (s, 3H), 2.22-2.17 (m, 1H), 2.08-2.01 (m, 1H), 1.59-1.56 (m, 6H)
A402	(400 MHz, DMSO-d ₆ , ppm) δ 9.19 (d, J = 8.62 Hz, 1H) 8.46 (s, 1H) 7.77 (d, J = 1.90 Hz, 2H) 7.68 (t, J = 1.96 Hz, 1H) 7.45 (d, J = 2.41 Hz, 1H) 7.36 (dd, J = 8.68, 2.47 Hz, 1H) 6.83 (d, J = 8.74 Hz, 1H) 4.81 (d, J = 8.62 Hz, 1H) 4.29 (d, J = 11.28 Hz, 1H) 3.72 - 3.76 (m, 1H) 3.48 - 3.58 (m, 1H) 2.54 (s, 3H) 1.54 (dd, J = 8.36, 7.10 Hz, 6H) 0.89 - 0.99 (m, 1H) 0.75 (dt, J = 9.03, 4.67 Hz, 1H) 0.66 - 0.72 (m, 1H) 0.59 - 0.66 (m, 1H)

A403	(400 MHz, DMSO-d6) δ ppm 9.14 (d, J = 8.11 Hz, 1H) 8.56 (s, 1H) 7.96 (d, J = 1.90 Hz, 2H) 7.69 (t, J = 1.90 Hz, 1H) 7.34 (d, J = 7.10 Hz, 1H) 7.18 (t, J = 7.69 Hz, 1H) 6.93 (t, J = 7.16 Hz, 1H) 6.80 (d, J = 8.24 Hz, 1H) 5.19 - 5.29 (m, 1H) 4.65 (s, 2H) 4.18 - 4.33 (m, 2H) 3.57 - 3.66 (m, 1H) 2.13 - 2.27 (m, 1H) 1.97 - 2.13 (m, 1H) 1.55 (dd, J = 10.77, 6.97 Hz, 6H)
A404	(400 MHz, DMSO-d6) δ ppm 9.18 (d, J=8.11 Hz, 1 H) 8.66 (s, 1 H) 7.84 (t, J=1.90 Hz, 1 H) 7.68 (d, J=1.90 Hz, 2 H) 7.33 (d, J=6.97 Hz, 1 H) 7.17 (t, J=7.77 Hz, 1 H) 6.92 (t, J=7.17 Hz, 1 H) 6.80 (d, J=8.11 Hz, 1 H) 5.21 - 5.27 (m, 1 H) 4.25 - 4.31 (m, 1 H) 4.18 - 4.25 (m, 1 H) 3.58 (dt, J=13.94, 6.97 Hz, 1 H) 2.17 - 2.25 (m, 1 H) 2.00 - 2.08 (m, 1 H) 1.54 (dd, J=11.79)
A406	(400 MHz, DMSO-d6, ppm) δ 9.15 (d, J = 7.98 Hz, 1H), 8.34 (s, 1H), 7.66 - 7.71 (m, 2H), 7.57 - 7.64 (m, 1H), 7.31 (d, J = 6.97 Hz, 1H), 7.16 (t, J = 7.66 Hz, 1H), 6.91 (t, J = 7.15 Hz, 1H), 6.78 (d, J = 8.27 Hz, 1H), 5.19 - 5.25 (m, 1H), 4.18 - 4.30 (m, 2H), 3.35 - 3.61 (m, 3H), 2.56 - 2.66 (m, 1H), 2.15 - 2.32 (m, 1H), 2.01 - 2.09 (m, 1H), 1.51 - 1.59 (m, 6H)
A407	(400 MHz, DMSO-d6, ppm) δ 9.12 (d, J = 8.11 Hz, 1H), 8.39 (s, 1H), 7.62 - 7.70 (m, 1H), 7.14 - 7.36 (m, 4H), 6.91 (t, J = 7.19 Hz, 1H), 6.79 (d, J = 8.11 Hz, 1H), 5.23 (br d, J = 7.48 Hz, 1H), 4.17 - 4.32 (m, 2H), 3.51 - 3.62 (m, 1H), 2.03 - 2.39 (m, 5H), 1.57 - 1.54 (m, 6H)
A410	(400 MHz, DMSO-d6, ppm) δ 9.11 (d, J = 8.11 Hz, 1H), 8.44 - 8.52 (m, 1H), 7.93 (m, 1H), 7.15 - 7.73 (m, 4H), 6.92 (m, 1H), 6.76 - 6.80 (m, 1H), 4.04 - 5.27 (m, 4H), 3.52 - 3.66 (m, 1H), 2.51 - 2.53 (m, 2H), 2.36 - 2.46 (m, 1H), 1.98 - 2.26 (m, 2H), 1.46 - 1.61 (m, 5H)
A411	(400 MHz, DMSO-d6, ppm) δ ppm 9.14 (d, J = 8.24 Hz, 1H) 8.47 (s, 1H) 7.77 (d, J = 1.90 Hz, 2H) 7.68 (t, J = 1.90 Hz, 1H) 7.37 (d, J = 7.73 Hz, 1H) 7.07 - 7.19 (m, 3H) 5.22 - 5.29 (m, 1H) 3.57 - 3.64 (m, 3H) 3.03 - 3.20 (m, 2H) 2.54 (s, 3H) 2.14 - 2.31 (m, 2H) 1.55 (br d, J = 6.97 Hz, 3H) 1.52 (br d, J = 6.97 Hz, 3H)
A412	(400 MHz, DMSO-d6, ppm) δ 9.10 (d, J = 8.11 Hz, 1H), 8.40 (s, 1H), 6.77 - 7.76 (m, 9H), 5.19 - 5.28 (m, 1H), 3.55 - 4.66 (m, 5H), 2.00 - 2.49 (m, 2H), 1.52 - 1.56 (m, 6H)

A413	(400 MHz, DMSO-d ₆ , ppm) δ 9.11 (d, J = 7.98 Hz, 1H), 8.37 (s, 1H), 7.63 - 7.71 (m, 1H), 7.46 (m, 1H), 7.26 - 7.34 (m, 2H), 7.16 (t, J = 7.74 Hz, 1H), 6.92 (t, J = 7.41 Hz, 1H), 6.79 (d, J = 7.60 Hz, 1H), 5.23 (br d, J = 7.73 Hz, 1H), 4.25 (m, 2H), 3.37 - 3.67 (m, 3H), 2.35 - 2.46 (m, 1H), 2.16 - 2.26 (m, 1H), 2.04 (br s, 1H), 1.56 - 1.53 (m, 6H)
A414	(400 MHz, DMSO-d ₆ , ppm) δ 9.15 (d, J = 7.98 Hz, 1H) 8.54 (s, 1H) 7.78 (d, J = 1.90 Hz, 2H) 7.68 (t, J = 1.90 Hz, 1H) 7.35 (d, J = 2.41 Hz, 1H) 7.22 (dd, J = 8.74, 2.66 Hz, 1H) 6.84 (d, J = 8.74 Hz, 1H) 5.23 (q, J = 6.42 Hz, 1H) 3.88 - 4.34 (m, 2H) 3.60 (spt, J = 6.91 Hz, 1H) 2.55 (s, 3 H) 1.94 - 2.24 (m, 2H) 1.55 (dd, J = 10.39, 6.97 Hz, 6H)
A415	(400 MHz, DMSO-d ₆) δ ppm 9.35 (d, J = 8.62 Hz, 1H), 8.57 (s, 1H), 7.57 - 7.85 (m, 7H), 5.52 (m, 1H), 3.57 - 3.81 (m, 3H), 2.53 - 2.69 (m, 4H), 1.55 - 1.50 (m, 7H)
A416	(400 MHz, DMSO-d ₆ , ppm) δ 9.35 (d, J = 8.62 Hz, 1H), 8.57 (s, 1H), 7.57 - 7.85 (m, 7H), 5.52 (m, 1H), 3.57 - 3.81 (m, 3H), 2.53 - 2.69 (m, 4H), 1.55 - 1.50 (m, 7H)
A417	(400 MHz, DMSO-d ₆ , ppm) δ 9.37 (d, J = 8.36 Hz, 1H), 8.61 (s, 1H), 7.78 - 7.89 (m, 3H), 7.61 - 7.71 (m, 3H), 5.48 - 5.55 (m, 1H), 3.57 - 3.84 (m, 3H), 2.54 - 2.68 (m, 5H), 1.58 (d, J = 6.97 Hz, 3H), 1.54 (br d, J = 6.84 Hz, 3H)
A418	(400 MHz, DMSO-d ₆ , ppm) δ 9.37 (d, J = 8.36 Hz, 1H), 8.61 (s, 1H), 7.78 - 7.89 (m, 3H), 7.61 - 7.71 (m, 3H), 5.48 - 5.55 (m, 1H), 3.57 - 3.84 (m, 3H), 2.54 - 2.68 (m, 5H), 1.58 (d, J = 6.97 Hz, 3H), 1.54 (br d, J = 6.84 Hz, 3H)
419	(300 MHz, DMSO-d ₆ , ppm) δ 8.50 (s, 1H), 7.83 (d, J = 7.5 Hz, 2H), 7.45 - 7.50 (m, 1H), 7.10 - 7.28 (m, 1H), 6.98 - 6.89 (m, 2H), 6.10 - 6.05 (m, 1H), 5.45 - 5.40 (m, 1H), 4.50 - 4.10 (m, 2H) 3.80-3.70 (m, 1H), 2.40-2.21 (m, 2H), 1.65-1.58 (m, 6H)
A419	(400 MHz, DMSO-d ₆ , ppm) δ 9.17 (d, J = 7.98 Hz, 1H), 8.97 (s, 1H), 8.39 (s, 1H), 7.67 - 7.78 (m, 3H), 5.28 - 5.35 (m, 1H), 3.58 (m, 1H), 2.68 - 2.80 (m, 2H), 2.52 - 2.55 (m, 3H), 1.51 - 2.16 (m, 10H)

A420	(400 MHz, DMSO-d6, ppm) δ 9.17 (d, J = 7.98 Hz, 1H), 8.97 (s, 1H), 8.39 (s, 1H), 7.67 - 7.78 (m, 3H), 5.28 - 5.35 (m, 1H), 3.58 (m, 1H), 2.68 - 2.80 (m, 2H), 2.52 - 2.55 (m, 3H), 1.51 - 2.16 (m, 10H)
A423	(400 MHz, DMSO-d6, ppm) δ 11.72 - 12.10 (m, 1H), 9.07 (d, J = 8.11 Hz, 1H), 8.37 (s, 1H), 7.66 - 7.79 (m, 3H), 5.21 (d, J = 6.72 Hz, 1H), 3.54 - 3.65 (m, 1H), 2.53 - 2.60 (m, 4H), 1.51 - 2.14 (m, 14H)
A424	(400 MHz, DMSO-d6, ppm) δ 11.72 - 12.10 (m, 1H), 9.07 (d, J = 8.11 Hz, 1H), 8.37 (s, 1H), 7.66 - 7.79 (m, 3H), 5.21 (d, J = 6.72 Hz, 1H), 3.54 - 3.65 (m, 1H), 2.53 - 2.60 (m, 4H), 1.51 - 2.14 (m, 14H)
A425	(400 MHz, DMSO-d6, ppm) δ 9.42 (d, J = 7.86 Hz, 1H), 8.57 (s, 1H), 7.67 - 7.81 (m, 5H), 5.45 (d, J = 6.46 Hz, 1H), 3.56 - 4.22 (m, 3H), 2.54 - 2.57 (m, 3H), 1.50 - 2.32 (m, 10H)
A426	(400 MHz, DMSO-d6, ppm) δ 9.42 (d, J = 7.86 Hz, 1H), 8.57 (s, 1H), 7.67 - 7.81 (m, 5H), 5.45 (d, J = 6.46 Hz, 1H), 3.56 - 4.22 (m, 3H), 2.54 - 2.57 (m, 3H), 1.50 - 2.32 (m, 10H)
A428	(400 MHz, DMSO-d6, ppm) δ 9.02 (d, J = 8.24 Hz, 1H), 8.39 (s, 1H), 7.67 - 7.80 (m, 1H), 7.67 (s, 1H), 5.40 (br d, J = 2.66 Hz, 1H), 3.36 - 3.70 (m, 2H), 2.76 - 3.16 (m, 2H), 2.53 - 2.57 (m, 2H), 2.50 - 2.67 (m, 2H), 2.08 - 2.39 (m, 1H), 1.23 - 1.55 (m, 6H)
A429	400 MHz, DMSO-d6, ppm) δ 9.00 (t, J = 4.06 Hz, 2H), 8.38 (s, 1H), 7.76 (d, J = 1.77 Hz, 1H), 7.67 (s, 1H), 5.45 (d, J = 5.07 Hz, 1H), 3.60 - 3.77 (m, 1H), 2.72 - 3.47 (m, 4H), 2.53 - 2.56 (m, 2H), 2.34 - 2.46 (m, 1H), 1.42 - 1.58 (m, 6H)
A430	(400 MHz, DMSO-d6, ppm) δ 9.02 (d, J = 8.24 Hz, 1H), 8.39 (s, 1H), 7.67 - 7.80 (m, 1H), 7.67 (s, 1H), 5.40 (br d, J = 2.66 Hz, 1H), 3.36 - 3.70 (m, 2H), 2.76 - 3.16 (m, 2H), 2.53 - 2.57 (m, 2H), 2.50 - 2.67 (m, 2H), 2.08 - 2.39 (m, 1H), 1.23 - 1.55 (m, 6H)
A431	(400 MHz, DMSO-d6, ppm) δ 8.82 (br t, J = 5.39 Hz, 1H) 8.40 (s, 1H) 7.77 (d, J = 1.65 Hz, 2H) 7.66 - 7.73 (m, 1H) 7.20 (d, J = 8.36 Hz, 1H) 6.55 - 6.60 (range, 1H) 6.48 - 6.54 (range, 1H) 4.38 (br d, J = 5.45 Hz, 2H) 3.81 (s, 3H) 3.76 (s, 3H) 3.50 (dt, J = 13.88, 6.88 Hz, 1H) 2.53 (s, 3H) 1.50 (d, J = 6.84 Hz, 6H)

A432	(400 MHz, DMSO-d ₆ , ppm) δ 9.15 (d, J = 7.86 Hz, 1H), 8.62 (s, 1H), 7.62 - 7.82 (m, 5H), 6.98 (d, J = 8.49 Hz, 1H), 5.26 (m, 1H), 3.77 - 4.43 (m, 2H), 3.34 - 3.70 (m, 1H), 2.33 - 2.43 (m, 1H), 1.98 - 2.31 (m, 2H), 1.55 (m, 6H)
A433	(400 MHz, DMSO-d ₆ , ppm) δ 9.55 (d, J = 7.60 Hz, 1H), 8.60 (s, 1H), 8.54 (s, 1H), 8.24 (m, 1H), 7.97 (d, J = 8.11 Hz, 1H), 7.68 - 7.80 (m, 3H), 5.89 (d, J = 7.10 Hz, 1H), 3.70 - 4.29 (m, 3H), 2.53 - 2.57 (m, 3H), 1.45 - 1.54 (m, 6H)
A434	(400 MHz, DMSO-d ₆ , ppm) δ 9.55 (d, J = 7.60 Hz, 1H), 8.60 (s, 1H), 8.54 (s, 1H), 8.24 (m, 1H), 7.97 (d, J = 8.11 Hz, 1H), 7.68 - 7.80 (m, 3H), 5.89 (d, J = 7.10 Hz, 1H), 3.70 - 4.29 (m, 3H), 2.53 - 2.57 (m, 3H), 1.45 - 1.54 (m, 6H)
A435	(400 MHz, DMSO-d ₆ , ppm) δ 9.11 (d, J = 8.49 Hz, 1H), 8.37 (s, 1H), 7.38 - 7.79 (m, 4H), 6.82 (d, J = 5.07 Hz, 1H), 5.27 (d, J = 6.59 Hz, 1H), 3.36 - 3.67 (m, 1H), 2.53 - 2.62 (m, 5H), 1.52 - 2.14 (m, 10H)
A436	(400 MHz, DMSO-d ₆) δ ppm 8.44 (s, 1 H) 8.05 (br s, 1 H) 7.80 (br s, 1 H) 7.77 (d, J=1.90 Hz, 2 H) 7.68 (t, J=1.84 Hz, 1 H) 3.56 - 3.67 (m, 1 H) 2.54 (s, 3 H) 1.53 (d, J=6.97 Hz, 6 H)
A437	(400 MHz, DMSO-d ₆ , ppm) δ 9.11 (d, J = 8.49 Hz, 1H), 8.37 (s, 1H), 7.38 - 7.79 (m, 4H), 6.82 (d, J = 5.07 Hz, 1H), 5.27 (d, J = 6.59 Hz, 1H), 3.36 - 3.67 (m, 1H), 2.53 - 2.62 (m, 5H), 1.52 - 2.14 (m, 10H)
A438	(400 MHz, DMSO-d ₆ , ppm) δ 9.32 (d, J = 7.6 Hz, 1H), 8.42 (s, 1H), 8.10 - 8.17 (m, 3H), 7.64 - 7.83 (m, 1H), 7.21 - 7.31 (m, 2H), 5.66 - 5.77 (m, 1H), 4.93 (t, J = 9.4 Hz, 1H), 4.48 (m, 1H), 3.35 - 3.73 (m, 1H), 2.52 - 2.69 (m, 3H), 1.50 - 1.63 (m, 6H)
A439	400 MHz, DMSO-d ₆ , ppm) δ 8.97 (d, J = 8.11 Hz, 1H), 8.44 (s, 1H), 7.66 - 7.79 (m, 3H), 6.94 - 7.17 (m, 2H), 6.52 - 6.63 (m, 2H), 3.88 - 5.17 (m, 1H), 2.53 - 3.60 (m, 7H), 1.44 - 2.33 (m, 8H)
A440	400 MHz, DMSO-d ₆ , ppm) δ 8.97 (d, J = 8.11 Hz, 1H), 8.44 (s, 1H), 7.66 - 7.79 (m, 3H), 6.94 - 7.17 (m, 2H), 6.52 - 6.63 (m, 2H), 3.88 - 5.17 (m, 1H), 2.53 - 3.60 (m, 7H), 1.44 - 2.33 (m, 8H)


A441	(400 MHz, DMSO-d6, ppm) δ 8.94 (d, J = 8.36 Hz, 1H), 8.43 (s, 1H), 7.66 - 7.80 (m, 3H), 6.87 - 7.36 (m, 2H), 5.09 - 5.17 (m, 1H), 3.37 - 3.78 (m, 1H), 2.53 - 2.77 (m, 5H), 2.23 - 2.41 (m, 1H), 1.51 - 2.09 (m, 9H)
A442	(400 MHz, DMSO-d6, ppm) δ 9.32 (d, J = 7.6 Hz, 1H), 8.42 (s, 1H), 8.10 - 8.17 (m, 3H), 7.64 - 7.83 (m, 1H), 7.21 - 7.31 (m, 2H), 5.66 - 5.77 (m, 1H), 4.93 (t, J = 9.4 Hz, 1H), 4.48 (m, 1H), 3.35 - 3.73 (m, 1H), 2.52 - 2.69 (m, 3H), 1.50 - 1.63 (m, 6H)
A443	(400 MHz, DMSO-d6, ppm) δ 8.94 (d, J = 8.36 Hz, 1H), 8.43 (s, 1H), 7.66 - 7.80 (m, 3H), 6.87 - 7.36 (m, 2H), 5.09 - 5.17 (m, 1H), 3.37 - 3.78 (m, 1H), 2.53 - 2.77 (m, 5H), 2.23 - 2.41 (m, 1H), 1.51 - 2.09 (m, 9H)
A445	(400 MHz, DMSO-d6, ppm) δ 9.21 (d, J = 7.35 Hz, 1H), 8.44 (s, 1H), 7.66 - 7.78 (m, 3H), 7.30 (d, J = 8.24 Hz, 1H), 6.51 (m, 1H), 6.47 (d, J = 2.15 Hz, 1H), 5.65 (br d, J = 4.56 Hz, 1H), 4.81 (m, 1H), 4.40 (m, 1H), 3.56 - 3.74 (m, 4H), 2.52 - 2.55 (m, 3H), 1.50 - 1.59 (m, 6H)
A446	(400 MHz, DMSO-d6, ppm) δ 9.21 (d, J = 7.35 Hz, 1H), 8.44 (s, 1H), 7.66 - 7.78 (m, 3H), 7.30 (d, J = 8.24 Hz, 1H), 6.51 (m, 1H), 6.47 (d, J = 2.15 Hz, 1H), 5.65 (br d, J = 4.56 Hz, 1H), 4.81 (m, 1H), 4.40 (m, 1H), 3.56 - 3.74 (m, 4H), 2.52 - 2.55 (m, 3H), 1.50 - 1.59 (m, 6H)
A447	(400 MHz, DMSO-d6, ppm) δ 9.28 (d, J = 7.35 Hz, 1H), 8.46 (s, 1H), 7.66 - 7.79 (m, 3H), 7.43 (d, J = 7.35 Hz, 1H), 7.24 (t, J = 7.41 Hz, 1H), 6.94 (t, J = 7.41 Hz, 1H), 6.87 (d, J = 7.98 Hz, 1H), 5.72 - 5.80 (m, 1H), 4.80 (t, J = 9.19 Hz, 1H), 4.29 - 4.38 (m, 1H), 3.58 (m, 1H), 2.53 - 2.56 (m, 3H), 1.50 - 1.60 (m, 6H)
A448	(400 MHz, DMSO-d6, ppm) δ 9.20 (d, J = 7.35 Hz, 1H), 8.84 (s, 1H), 8.43 (s, 1H), 7.67 - 7.80 (m, 3H), 5.21 - 5.27 (m, 1H), 4.28 - 4.43 (m, 2H), 3.58 (m, 1H), 2.52 - 2.67 (m, 3H), 2.00 - 2.45 (m, 2H), 1.52 (m, 6H)
A451	(400 MHz, DMSO-d6, ppm) δ 9.20 (d, J = 7.35 Hz, 1H), 8.84 (s, 1H), 8.43 (s, 1H), 7.67 - 7.80 (m, 3H), 5.21 - 5.27 (m, 1H), 4.28 - 4.43 (m, 2H), 3.58 (m, 1H), 2.52 - 2.67 (m, 3H), 2.00 - 2.45 (m, 2H), 1.52 (m, 6H)
A452	(400 MHz, DMSO-d6, ppm) δ 9.02 (d, J = 8.49 Hz, 1H), 8.53 (s, 1H), 7.78 (d, J = 1.90 Hz, 2H), 7.68 (t, J = 1.84 Hz, 1H), 7.36 (d, J = 7.73 Hz, 1H), 7.17 (t, J =

	7.71 Hz, 1H), 6.92 (t, J = 7.26 Hz, 1H), 6.76 (d, J = 8.11 Hz, 1H), 5.32 (m, 1H), 3.52 - 3.62 (m, 1H), 2.52 - 2.58 (m, 3H), 2.21 (m, 1H), 1.80 - 1.89 (m, 1H), 1.23 - 1.61 (m, 12H)
A453	(400 MHz, DMSO-d ₆ , ppm) δ 9.28 (d, J = 7.35 Hz, 1H), 8.46 (s, 1H), 7.66 - 7.79 (m, 3H), 7.43 (d, J = 7.35 Hz, 1H), 7.24 (t, J = 7.41 Hz, 1H), 6.94 (t, J = 7.41 Hz, 1H), 6.87 (d, J = 7.98 Hz, 1H), 5.72 - 5.80 (m, 1H), 4.80 (t, J = 9.19 Hz, 1H), 4.29 - 4.38 (m, 1H), 3.58 (m, 1H), 2.53 - 2.56 (m, 3H), 1.50 - 1.60 (m, 6H)
A454	(400 MHz, DMSO-d ₆ , ppm) δ 9.17 (d, J = 8.11 Hz, 1H) 8.54 (s, 1H) 7.78 (d, J = 1.77 Hz, 2H) 7.68 (t, J = 1.90 Hz, 1H) 7.54 (d, J = 7.98 Hz, 1H) 7.36 (dd, J = 7.86, 1.52 Hz, 1H) 7.31 (d, J = 1.52 Hz, 1H) 5.30 (q, J = 6.67 Hz, 1H) 4.24 - 4.39 (m, 2H) 3.50 - 3.65 (m, 1H) 2.55 (s, 3H) 2.15 - 2.31 (m, 1H) 1.99 - 2.15 (m, 1H) 1.54 (dd, J = 11.09, 6.91 Hz, 6H)
A455	(400 MHz, DMSO-d ₆ , ppm) δ 9.41 (d, J = 7.35 Hz, 1H), 8.51 (s, 1H), 8.27 - 8.42 (m, 2H), 7.66 - 7.79 (m, 4H), 5.82 - 5.90 (m, 1H), 4.93 (t, J = 9.38 Hz, 1H), 3.61 - 4.54 (m, 1H), 2.53 - 2.58 (m, 3H), 1.48 - 1.53 (m, 3H), 1.45 - 1.51 (m, 4H)
A456	(400 MHz, DMSO-d ₆ , ppm) δ 9.41 (d, J = 7.35 Hz, 1H), 8.51 (s, 1H), 8.27 - 8.42 (m, 2H), 7.66 - 7.79 (m, 4H), 5.82 - 5.90 (m, 1H), 4.93 (t, J = 9.38 Hz, 1H), 3.61 - 4.54 (m, 1H), 2.53 - 2.58 (m, 3H), 1.48 - 1.53 (m, 3H), 1.45 - 1.51 (m, 4H)
A457	(400 MHz, DMSO-d ₆ , ppm) δ 9.33 (d, J = 7.10 Hz, 1H), 8.52 (s, 1H), 7.66 - 7.79 (m, 3H), 7.48 (s, 1H), 7.20 - 7.28 (m, 1H), 6.90 (d, J = 8.49 Hz, 1H), 5.73 (br d, J = 5.20 Hz, 1H), 4.85 (t, J = 9.25 Hz, 1H), 4.43 (m, 1H), 3.35 - 3.63 (m, 1H), 2.53 - 2.56 (m, 3H), 1.50 - 1.53 (m, 6H)
A458	(400 MHz, DMSO-d ₆ , ppm) δ 9.33 (d, J = 7.10 Hz, 1H), 8.52 (s, 1H), 7.66 - 7.79 (m, 3H), 7.48 (s, 1H), 7.20 - 7.28 (m, 1H), 6.90 (d, J = 8.49 Hz, 1H), 5.73 (br d, J = 5.20 Hz, 1H), 4.85 (t, J = 9.25 Hz, 1H), 4.43 (m, 1H), 3.35 - 3.63 (m, 1H), 2.53 - 2.56 (m, 3H), 1.50 - 1.53 (m, 6H)
A472	(400 MHz, DMSO-d ₆ , ppm) δ 9.02 (d, J = 8.49 Hz, 1H), 8.53 (s, 1H), 7.78 (d, J = 1.90 Hz, 2H), 7.68 (t, J = 1.84 Hz, 1H), 7.36 (d, J = 7.73 Hz, 1H), 7.17 (t, J = 7.71 Hz, 1H), 6.92 (t, J = 7.26 Hz, 1H), 6.76 (d, J = 8.11 Hz, 1H), 5.32 (m, 1H),


	3.52 - 3.62 (m, 1H), 2.52 - 2.58 (m, 3H), 2.21 (m, 1H), 1.80 - 1.89 (m, 1H), 1.23 - 1.61 (m, 12H)
560	(400 MHz, Chloroform-d, ppm) δ : 8.35 (s, 1H), 7.51 (s, 2H), 7.65 (s, 1H), 7.29-7.21 (m, 2H), 7.01-6.84 (m, 3H), 6.20 (d, J = 8.3 Hz, 1H), 6.08-6.07 (m, 1H), 5.40 (d, J = 8.3 Hz, 1H), 4.39-4.34 (m, 1H), 4.21-4.15 (m, 1H), 3.71-3.64 (m, 1H), 2.44-2.39 (m, 1H), 2.26-2.20 (m, 1H), 1.65-1.54 (m, 6H)

Preparation Example 3: The example compounds A400, A401, A405 and A459 were prepared according to scheme 8, with reactions that are adapted from known reactions in the literature. For example, see Campbell, Alison N. et al. Organic Process Research & Development (2013), 17(2), 273-281 and Stanovnik, B. et al., Tetrahedron (1967), 23(6), 2739-46.

Scheme 8



Description of the key step: Synthesis of methyl 3-(3,5-dichlorophenyl)-8-(4,4-difluorocyclohexyl)-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (8-6)



A mixture of 90 mg (0.3 mmol) methyl 3-(3,5-dichlorophenyl)-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (8-5) in 5 mL DMSO was treated with 200 mg (1.0 mmol) of zinc sulfinate and the solution was cooled with an ice bath. 185 μ L (1 mmol) of 2-Methyl-prop-2-yl-hydroperoxide (TBHP) was added dropwise and the reaction mixture was stirred at 50 $^{\circ}$ C for 1 h. The reaction mixture was quenched with sodium carbonate solution and extracted with EE. The organic layers were collected, dried, filtered and evaporated. The mixture was evaporated, purified by column chromatography (silica gel; CyH/EtOAc) and the solvents removed in vacuo to obtain 900 mg (74 %) of the product as yellow oil. (400 MHz, DMSO- d_6) δ ppm 9.11 (d, J =8.11 Hz, 1 H) 8.52 (s, 1 H) 7.76 (d, J =1.77 Hz, 2 H) 7.66 - 7.71 (m, 1 H) 7.39 (d, J =7.35 Hz, 1 H) 7.18 (t, J =7.73 Hz, 1 H) 6.89 - 6.95 (m, 1 H) 6.80 (d, J =8.11 Hz, 1 H) 5.23 - 5.30 (m, 1 H) 4.18 - 4.33 (m, 2 H) 2.61 - 2.80 (m, 3 H) 2.53 (s, 3 H) 2.30 - 2.39 (m, 1 H) 2.09 - 2.30 (m, 3 H) 1.73 - 1.94 (m, 4 H).

The conversion of compound 8-6 into the product may be achieved as in Scheme 4 by hydrolysis of the methyl ester to the carboxylic acid followed by coupling of the acid with the desired amine.


As noted above, compounds A400, A401 and A405 are prepared by adopting the process described in Scheme 8.

Compound	^1H NMR Spectra
A400	(400 MHz, DMSO- d_6) δ ppm 9.12 (d, J =7.98 Hz, 1 H) 8.69 (s, 1 H) 7.79 (d, J =1.90 Hz, 2 H) 7.71 - 7.75 (m, 1 H) 7.34 (d, J =7.73 Hz, 1 H) 7.18 (t, J =7.67 Hz,


	1 H) 6.92 (t, J=7.09 Hz, 1 H) 6.80 (d, J=8.11 Hz, 1 H) 5.14 - 5.22 (m, 1 H) 4.24 - 4.34 (m, 1 H) 4.13 - 4.24 (m, 1 H) 2.67 (dt, J=3.68, 1.84 Hz, 1 H) 2.56 (s, 3 H) 2.27 (t, J=19.33 Hz, 3 H) 2.10 - 2.21 (m, 1 H) 1.96 - 2.08 (m, 1 H)
A401	(400 MHz, DMSO-d ₆) δ ppm 9.34 (d, J=8.11 Hz, 1 H) 8.84 (s, 1 H) 7.80 (d, J=1.90 Hz, 2 H) 7.74 - 7.77 (m, 1 H) 7.33 (d, J=7.10 Hz, 1 H) 7.19 (t, J=7.10 Hz, 1 H) 6.93 (t, J=7.03 Hz, 1 H) 6.81 (d, J=8.11 Hz, 1 H) 5.16 - 5.23 (m, 1 H) 4.23 - 4.33 (m, 1 H) 4.14 - 4.23 (m, 1 H) 2.58 (s, 3 H) 2.16 - 2.27 (m, 1 H) 1.98 - 2.07 (m, 1 H)
A405	(400 MHz, DMSO-d ₆) δ ppm 9.29 (d, J=7.98 Hz, 1 H) 8.77 (s, 1 H) 7.80 (d, J=1.90 Hz, 2 H) 7.73 (d, J=1.90 Hz, 1 H) 7.60 (t, J=52.00 Hz, 1 H) 7.36 (d, J=6.97 Hz, 1 H) 7.16 - 7.21 (m, 1 H) 6.92 (td, J=7.45, 1.08 Hz, 1 H) 6.81 (d, J=8.11 Hz, 1 H) 5.19 - 5.25 (m, 1 H) 4.19 - 4.31 (m, 2 H) 2.58 (s, 3 H) 2.15 - 2.24 (m, 1 H) 2.03 - 2.11 (m, 1 H)

Preparation Example 4: The example 304-0 was prepared according to scheme 9 below. Analogously, compound 321 can be prepared by the one skilled in the art in the same manner.

Scheme 9




1. Synthesis of ethyl 2-(3,5-dichlorophenyl)-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (9-2)



Into a 500-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed ethyl 2-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (3-6, 2.0 g, 0.007 mmol, 1.0 equiv), 3,5-dichlorophenylboronic acid (1.6 g, 0.008 mmol, 1.2 equiv), K₂CO₃ (2.1 g, 0.015 mmol, 2.2 equiv), dioxane (100.0 mL), H₂O (20.0 mL), Pd(dtbpf)Cl₂ (0.27 g, 0.000 mmol, 0.06 equiv). The resulting solution was stirred for 1 hr at 100 °C in an oil bath. The reaction


was then quenched by the addition of 100 mL of water. The solids were collected by filtration. This resulted in 2.1 g (85.3%) of ethyl 2-(3,5-dichlorophenyl)-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (9-2) as an off-white solid. (ES, m/z): 352 [M+H]⁺.

2. Synthesis of ethyl 8-bromo-2-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (9-3)



Into a 100-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed ethyl 2-(3,5-dichlorophenyl)-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (9-2, 2.0 g, 5.7 mmol, 1.0 equiv), CHCl₃ (25.0 mL), POBr₃ (8.14 g, 28.4 mmol, 5.0 equiv). The resulting solution was stirred for 1 overnight at 80 °C. The resulting mixture was cooled to room temperature and concentrated. The residue was diluted with 50 ml of water. The pH value of the solution was adjusted to 7~8 with Na₂CO₃ (saturated). The resulting solution was extracted with 3x50 mL of ethyl acetate. The combined organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:10). This resulted in 810 mg (34.4%) of ethyl 8-bromo-2-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (9-3) as an off-white solid. (ES, m/z): 414 [M+H]⁺.


3. Synthesis of ethyl 2-(3,5-dichlorophenyl)-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (9-4)



Into a 50-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed ethyl 8-bromo-2-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (9-3, 1800.0 mg, 4.4 mmol, 1.0 equiv), K₃PO₄ (2761.5 mg, 13.0 mmol, 3.0 equiv), THF (20.0 mL), H₂O (5.0 mL), Pd(dtbpf)Cl₂ (282.6 mg, 0.4 mmol, 0.1 equiv), 4,4,5,5-tetramethyl-


2-(prop-1-en-2-yl)-1,3,2-dioxaborolane (801.6 mg, 4.8 mmol, 1.1 equiv). The resulting solution was stirred for 1 hr at room temperature. The reaction was then quenched by the addition of 20 mL of water. The resulting solution was extracted with 3x30 mL of ethyl acetate. The combined organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was applied
 5 onto a silica gel column with ethyl acetate/petroleum ether (1:10). This resulted in 0.55 g (33.7%) of ethyl 2-(3,5-dichlorophenyl)-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (9-4) as an off-white solid. (ES, m/z): 376 [M+H]⁺.

4. Synthesis of ethyl 2-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylate (9-5)



Into a 50-mL round-bottom flask purged and maintained with an inert atmosphere of H₂ (g), was placed ethyl 2-(3,5-dichlorophenyl)-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (9-4, 500.0 mg, 1.3 mmol, 1.0 equiv), EA (10.0 mL), PtO₂ (100.0 mg, 0.4 mmol, 0.3 equiv). The resulting solution was stirred for 1 hr at room temperature. The solids were filtered
 15 out. The filtrate was concentrated. This resulted in 450 mg (82.4%) of ethyl 2-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylate as an off-white solid (9-5). (ES, m/z): 378 [M+H]⁺.


5. Synthesis of ethyl 3-chloro-2-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylate (9-6)



Into a 20-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed ethyl 2-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-


carboxylate (9-5, 100.0 mg, 0.3 mmol, 1.0 equiv), CHCl_3 (5.0 mL), NCS (38.8 mg, 0.3 mmol, 1.1 equiv). The resulting solution was stirred for 1 overnight at 50 °C. The resulting mixture was concentrated. The residue was applied on a silica gel column and eluted with EA/PE (1/20). This resulted in 106.6 mg (97.7%) of ethyl 3-chloro-2-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylate (9-6) as an off-white solid. (ES, m/z): 412 $[\text{M}+\text{H}]^+$.

6. Synthesis of 3-chloro-2-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylic acid (9-7)



Into a 20-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed ethyl 3-chloro-2-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylate (9-6, 95.0 mg, 0.2 mmol, 1.0 equiv), THF (5.0 mL), H_2O (1.0 mL), LiOH (27.6 mg, 1.2 mmol, 5.0 equiv). The resulting solution was stirred for 1 overnight at room temperature. The pH value of the solution was adjusted to 3~4 with HCl (1 mol/L). The resulting solution was extracted with 3x10 mL of ethyl acetate and the organic layers combined. The organic phase was dried in an oven under reduced pressure and concentrated. This resulted in 78 mg (88.1%) of 3-chloro-2-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylic acid (9-7) as an off-white solid. (ES, m/z): 384 $[\text{M}+\text{H}]^+$.

7. Synthesis of 3-chloro-2-(3,5-dichlorophenyl)-N-[(4S)-6-fluoro-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropylimidazo[1,2-b]pyridazine-7-carboxamide (Compound 304-0)




Into a 20-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed (4S)-6-fluoro-3,4-dihydro-2H-1-benzopyran-4-amine dihydrochloride (51.0 mg, 0.2 mmol, 1.2 equiv), 3-chloro-2-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylic acid (9-7, 68.0 mg, 0.2 mmol, 1.0 equiv), DMF (5.0 mL), DIEA (45.7 mg, 0.35 mmol, 2.0 equiv), HATU (100.8 mg, 0.3 mmol, 1.5 equiv). The resulting solution was stirred for 1 overnight at room temperature. The crude product was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, ACN:H₂O=72 increasing to ACN:H₂O=95 within 7 ; Detector, 254. This resulted in 67.7 mg (71.7%) of 3-chloro-2-(3,5-dichlorophenyl)-N-[(4S)-6-fluoro-3,4-dihydro-2H-1-benzopyran-4-yl]-8-

isopropylimidazo[1,2-b]pyridazine-7-carboxamide (304-0) as an off-white solid. (300 MHz, CDCl₃, ppm) δ 8.36 (s, 1H), 7.83 (d, J = 1.8 Hz, 2H), 7.45 (t, J = 1.8 Hz, 1H), 7.02-6.93 (m, 2H), 6.87-6.83 (m, 1H), 6.06 (d, J = 7.8 Hz, 1H), 5.41 (t, J = 5.7 Hz, 1H), 4.38-4.32 (m, 1H), 4.24-4.16 (m, 1H), 3.73 (t, J = 6.9 Hz, 1H), 2.45-2.40 (m, 1H), 2.24-2.18 (m, 1H), 1.68-1.64 (m, 6H).


Compound 321: (300 MHz, Chloroform-d, ppm): δ 8.27 (s, 1H), 7.76-7.60 (m, 2H), 7.35-7.30 (m, 1H), 7.28-7.20 (m, 1H), 7.20-7.03 (m, 4H), 7.03-6.80 (m, 3H), 6.20-5.90 (m, 1H), 5.50-5.25 (m, 1H), 4.45-4.30 (m, 1H), 4.30-4.10 (m, 1H), 3.80-3.65 (m, 1H), 2.55-2.35 (m, 1H), 2.33-2.15 (m, 1H), 1.72 (t, J = 6.3 Hz, 6H)

Preparation Example 5: The example 174 was prepared according to scheme 10:

Scheme 10



1. Synthesis of N-(5-chloropyridazin-3-yl)-1,1-diphenylmethanimine (10-2)




5

Into a 40-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 3,5-dichloropyridazine (10-1, 500.0 mg, 3.4 mmol, 1.0 equiv), diphenylmethanimine (675.2 mg, 3.7 mmol, 1.1 equiv), XantPhos (69.9 mg, 0.12 mmol, 0.04 equiv), $\text{Pd}_2(\text{dba})_3$ (34.7 mg, 0.06 mmol, 0.02 equiv), Cs_2CO_3 (2187.1 mg, 6.7 mmol, 2.0 equiv),

dioxane (5 ml). The resulting solution was stirred for 3 hr at 90 degrees C in an oil bath. The solids were filtered out. The resulting mixture was concentrated. This resulted in 3 mL (30.4%) of N-(5-chloropyridazin-3-yl)-1,1-diphenylmethanimine (10-2) as brown oil.


2. Synthesis of 5-chloropyridazin-3-amine



Into a 50-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed N-(5-chloropyridazin-3-yl)-1,1-diphenylmethanimine (10-2, 10.0 mL), HCl (3M) (15.0 mL). The resulting solution was stirred for 1 hr at room temperature. The pH value of the solution was adjusted to 7 with NaHCO₃. The resulting mixture was concentrated. This resulted in 15 mL of 5-chloropyridazin-3-amine (5-1) as brown oil.

10

3. Synthesis of 7-chloroimidazo[1,2-b]pyridazine (10-4)




Into a 250-mL round-bottom flask, was placed 5-chloropyridazin-3-amine (5-1, 15.00 mL), chloroacetaldehyde (17.5 mL), H₂O (17.5 mL), i-PrOH (25 mL). The resulting solution was stirred for 5 hr at 95 °C in an oil bath. The resulting mixture was concentrated. The pH value of the solution was adjusted to 9 with NaOH. The resulting solution was extracted with 3x50 mL of ethyl acetate, the organic layer was washed with 3 x50 ml of brine. The organic phase was collected and dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (2:3). This resulted in 2.2 g (12.4%) of 7-chloroimidazo[1,2-b]pyridazine (10-4) as yellow oil.

15


20

4. Synthesis of 7-chloro-3-iodoimidazo[1,2-b]pyridazine (10-5)




Into a 50-mL round-bottom flask, was placed 7-chloroimidazo[1,2-b]pyridazine (1.0 g, 7.0 mmol, 1.0 equiv), NIS (2.2 g, 10.0 mmol, 1.5 equiv), DMF (10 mL). The resulting solution was stirred for 1 overnight at room temperature. The reaction was then quenched by the addition of 20 mL of water. The resulting solution was extracted with 3x20 mL of ethyl acetate concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (2:1). This resulted in 800 mg (43.9%) of 7-chloro-3-iodoimidazo[1,2-b]pyridazine as yellow oil.

5. Synthesis of 7-chloro-3-(2,6-difluorophenyl)imidazo[1,2-b]pyridazine (10-6)




Into a 40-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 7-chloro-3-iodoimidazo[1,2-b]pyridazine (10-5, 400.0 mg, 1.4 mmol, 1.0 equiv), 2,6-difluorophenylboronic acid (452.0 mg, 2.9 mmol, 2.0 equiv), Pd(dtbpf)Cl₂ (93.3 mg, 0.14 mmol, 0.1 equiv), K₃PO₄ (911.4 mg, 4.3 mmol, 3.0 equiv), THF (10 mL), H₂O (2.5 mL). The resulting solution was stirred for 1 hr overnight at room temperature. The resulting mixture was concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:3). This resulted in 150 mg (39.4%) of 7-chloro-3-(2,6-difluorophenyl)imidazo[1,2-b]pyridazine (10-6) as a white solid.

6. Synthesis of methyl 3-(2,6-difluorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (10-7)




Into a 50-mL pressure tank reactor, was placed 7-chloro-3-(2,6-difluorophenyl)imidazo[1,2-b]pyridazine (10-6) 130.0 mg, 0.5 mmol, 1.0 equiv), Pd(dppf)Cl_2 (35.8 mg, 0.05 mmol, 0.1 equiv), TEA (148.6 mg, 1.5 mmol, 3.0 equiv), CO (20 atm), MeOH (10.00 mL). The resulting solution was stirred for 4 hr at 110 °C in an oil bath. The resulting mixture was concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:3). This resulted in 95 mg (67.1%) of methyl 3-(2,6-difluorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (10-7) as a white solid.

7. Synthesis of 3-(2,6-difluorophenyl)imidazo[1,2-b]pyridazine-7-carboxylic acid (10-8)



Into a 40-mL round-bottom flask, was placed methyl 3-(2,6-difluorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (10-7, 85.0 mg, 0.3 mmol, 1.0 equiv), NaOH (58.7 mg, 1.5 mmol, 5.0 equiv), MeOH (9 mL), H₂O (3 mL). The resulting solution was stirred for 3 hr at room temperature. The resulting solution was diluted with 20 mL of water. The pH value of the solution was adjusted to 3-4 with HCl (3 mol/L). The resulting solution was extracted with 3x20 mL of ethyl acetate, the organic phase was collected and dried over anhydrous sodium sulfate and concentrated. This resulted in 65 mg (80.4%) of 3-(2,6-difluorophenyl)imidazo[1,2-b]pyridazine-7-carboxylic acid (10-8) as a white solid.


8. Synthesis of 3-(2,6-difluorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]imidazo[1,2-b]pyridazine-7-carboxamide (Compound 174)




Into a 40-mL round-bottom flask, was placed 3-(2,6-difluorophenyl)imidazo[1,2-b]pyridazine-7-carboxylic acid (10-8, 60.0 mg, 0.2 mmol, 1.0 equiv), (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (48.8 mg, 0.3 mmol, 1.5 equiv), HATU (165.79 mg, 0.436 mmol, 2 equiv), DIEA (84.5 mg, 0.6 mmol, 3.0 equiv), DMF (3 mL). The resulting solution was stirred for 2 hr at room temperature. The crude product was purified by Prep-HPLC with the following conditions (Waters-2767): Column, X-bridge RP18, 5 μ m, 19*100mm; mobile phase, 0.03% ammonia in water and CH₃CN (30% CH₃CN up to 70% in 15 min); Detector, UV 254 nm. This resulted in 19.4 mg (21.9%) of 3-(2,6-difluorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]imidazo[1,2-b]pyridazine-7-carboxamide (174) as a white solid. (300 MHz, CDCl₃, ppm) δ 8.95 (s, 1H), 8.54 (s, 1H), 7.97 (s, 1H), 7.55-7.45 (m, 1H), 7.25-7.24 (m, 1H), 7.17-7.08 (m, 4H), 6.91-6.81 (m, 2H), 5.43-5.39 (m, 1H), 4.35-4.25 (m, 2H), 2.41-2.31 (m, 1H), 2.28-2.21 (m, 1H).

Preparation Example 6: The example 277 was prepared according to scheme 11 below:

Scheme 11




1. Synthesis of ethyl 8-chloro-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (271-1)




Into a 100-mL round-bottom flask, was placed ethyl 8-hydroxy-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (3-7, 2.8 g, 12.6 mmol, 1.0 equiv), DMF (8.0 uL, 103.4 mmol, 8.2 equiv), CHCl₃ (55.0 mL). This was followed by the addition of (COCl)₂ (8.0 g, 63.1 mmol, 5.0 equiv) dropwise with stirring at room temperature. The resulting solution was stirred for 2 hrs at 80 °C. The resulting mixture was concentrated. The crude product was purified by Prep-Flash with the following conditions: Column, C¹⁸ silica gel; mobile phase, 0.1% FA in water and CH₃CN (10% CH₃CN increasing to 70% within 12 min). Detector, UV 254 nm, 220 nm. This resulted in 537 mg (17.2%) of ethyl 8-chloro-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (271-1) as a yellow solid.

2. Synthesis of ethyl 2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (4-2)




Into a 50-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed ethyl 8-chloro-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (271-1, 480.0 mg, 2.0 mmol, 1.0 equiv), dioxane (19.0 mL), H₂O (4.8 mL), 4,4,5,5-tetramethyl-2-(prop-1-en-2-yl)-1,3,2-dioxaborolane (673.1 mg, 4.0 mmol, 2.0 equiv), Pd(dtbpf)Cl₂ (130.5 mg, 0.20 mmol, 0.1 equiv), K₂CO₃ (553.6 mg, 4.0 mmol, 2.0 equiv). The resulting solution was stirred for overnight at 80 °C. The resulting solution was diluted with 10 mL of water. The resulting solution was extracted with 2x20 mL of ethyl acetate and the organic layers combined. The resulting mixture was washed with 2 x20 ml of brine. The mixture was dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (0-20%). This resulted in 370 mg (73.0%) of ethyl 2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (4-2) as a brown solid.

3. Synthesis of ethyl 3-bromo-2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (11-4)



Into a 40-mL round-bottom flask, was placed ethyl 2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (4-2, 370.0 mg, 1.5 mmol, 1.0 equiv), CHCl₃ (7.0 mL, 86.8 mmol), NBS (295.3 mg, 1.7 mmol, 1.1 equiv). The resulting solution was stirred for 30 min at 65 °C. The resulting solution was diluted with 20 mL of water. The resulting solution was extracted with 2x20 mL of dichloromethane and the organic layers combined. The resulting mixture was washed with 2 x20 ml of water. The mixture was dried over anhydrous magnesium sulfate and concentrated. This resulted in 491 mg (95.4%) of ethyl 3-bromo-2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (11-4) as a brown solid.


4. Synthesis of ethyl 3-(3,5-difluorophenyl)-2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (11-5)



Into a 50-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed ethyl 3-bromo-2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (11-4, 490.0 mg, 1.5 mmol, 1.0 equiv), dioxane (10.0 mL), H₂O (2.5 mL), 3,5-difluorophenylboronic acid (477.4 mg, 3.0 mmol, 2.0 equiv), K₂CO₃ (626.7 mg, 4.535 mmol, 3.0 equiv), Pd(dtbpf)Cl₂ (98.5 mg, 0.15 mmol, 0.1 equiv). The resulting solution was stirred for 30 min at 100 °C. The resulting solution was diluted with 10 mL of water. The resulting solution was extracted with 2x50 mL of ethyl acetate and the organic layers combined and dried over anhydrous


sodium sulfate and concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (0-15%). This resulted in 442 mg (80.2%) of ethyl 3-(3,5-difluorophenyl)-2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (11-5) as a yellow green solid.

5 **5. Synthesis of 3-(3,5-difluorophenyl)-2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylic acid (11-6)**



Into a 50-mL 3-necked round-bottom flask, was placed ethyl 3-(3,5-difluorophenyl)-2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (11-5, 440.0 mg, 1.2 mmol, 1.0 equiv), i-PrOH (15.0 mL), H₂O (8.0 mL), LiOH.H₂O (155.0 mg, 3.7 mmol, 3.0 equiv). The resulting solution was stirred for 1 hr at room temperature. The resulting mixture was concentrated. The pH value of the solution was adjusted to 3 with HCl (2 mol/L). The resulting solution was extracted with 2x20 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated. This resulted in 401 mg (89.0%) of 3-(3,5-difluorophenyl)-2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylic acid (11-6) as a yellow solid.


6. Synthesis of 3-(3,5-difluorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxamide (11-7)



Into a 50-mL 3-necked round-bottom flask, was placed 3-(3,5-difluorophenyl)-2-methyl-

8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylic acid (11-6, 400.0 mg, 1.2 mmol, 1.0 equiv), DMF (12.0 mL), (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (362.4 mg, 2.4 mmol, 2.0 equiv), DIEA (471.0 mg, 3.6 mmol, 3.0 equiv), HATU (692.8 mg, 1.8 mmol, 1.5 equiv). The resulting solution was stirred for 2 hrs at room temperature. The crude product was purified by Prep-Flash with the following conditions: Column, C18 silica gel; mobile phase, 0.1% NH_4HCO_3 in water and CH_3CN (30% CH_3CN increasing to 80% within 10 min). Detector, UV 254 nm, 220 nm. This resulted in 520 mg (92.0%) of 3-(3,5-difluorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxamide (11-7) as a green solid.

7. Synthesis of 3-(3,5-difluorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-(2-hydroxypropan-2-yl)-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (11-8)



Into a 50-mL 3-necked round-bottom flask, was placed 3-(3,5-difluorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-2-methyl-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxamide (11-7, 200.0 mg, 0.4 mmol, 1.0 equiv), ethanol (8.0 mL), toluene (8.0 mL), NaBH_4 (32.9 mg, 0.9 mmol, 2.0 equiv), $\text{Mn}(\text{OAc})_3 \cdot 2\text{H}_2\text{O}$ (9.3 mg, 0.03 mmol, 0.08 equiv), 2-[(1E)-([3-(E)-[(2-hydroxyphenyl)methylidene]amino]-2,2-dimethylpropyl]imino)methyl]phenol (10.8 mg, 0.03 mmol, 0.08 equiv). To the above O_2 (g) was introduced in. The resulting solution was stirred for 4 hrs at room temperature. The resulting solution was diluted with 10 mL of water. The resulting solution was extracted with 2x20 mL of ethyl acetate and the organic layers combined. The resulting mixture was washed with 2 x20 mL of brine. The mixture was dried over anhydrous sodium sulfate and concentrated. This resulted in 250 mg (crude) of 3-(3,5-difluorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-(2-hydroxypropan-2-yl)-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (11-8) as yellow oil.


8. Synthesis of 3-(3,5-difluorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-(2-fluoropropan-2-yl)-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (Compound 277)




Into a 25-mL 3-necked round-bottom flask, was placed 3-(3,5-difluorophenyl)-N-[(4S)-
 5 3,4-dihydro-2H-1-benzopyran-4-yl]-8-(2-hydroxypropan-2-yl)-2-methylimidazo[1,2-
 b]pyridazine-7-carboxamide (11-8, 200.0 mg, 0.4 mmol, 1.0 equiv), DCM (10.0 mL). This was
 followed by the addition of DAST (134.7 mg, 0.8 mmol, 2.0 equiv) dropwise with stirring at room
 temperature. The resulting solution was stirred for 1 hr at room temperature. The resulting mixture
 was concentrated. The crude product was purified by Prep-Flash with the following conditions:
 10 Column, C¹⁸ silica gel; mobile phase, 0.1% TFA in water and CH₃CN (50% CH₃CN increasing to
 100% within 10 min). Detector, UV 254 nm, 220 nm. This resulted in 9.5 mg (4.6%) of 3-(3,5-
 difluorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-(2-fluoropropan-2-yl)-2-
 methylimidazo[1,2-b]pyridazine-7-carboxamide (277) as a white solid. (300 MHz, CDCl₃, ppm)
 δ 8.26 (s, 1H), 7.35-7.33 (m, 1H), 7.29-7.19 (m, 3H), 6.98-6.85 (m, 3H), 5.94 (d, J = 7.5 Hz, 1H),
 15 5.32-5.29 (m, 1H), 4.38-4.33 (m, 1H), 4.23-4.18 (m, 1H), 2.62 (s, 3H), 2.40-2.24 (m, 2H), 2.15 (d,
 J = 6.3 Hz, 3H), 2.08 (d, J = 6.3 Hz, 3H).

Preparation Example 7: Compounds 306, 297, 298, 298-0, 299, 299-0, 418, 420, 523, 524, 525,
 526, 571, 572, 573, 574, A472 may be prepared by utilizing the process shown in Scheme 12
 shown below.

Scheme 12



1. Synthesis of 8-tert-butyl-3-(3,5-dichlorophenyl)-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (12-2)




5

To a stirred mixture of ethyl 8-bromo-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (12-1, 30.0 g, 105.6 mmol, 1.0 equiv) in THF (750.00 mL) was added LiBr (6.9 g, 791.9 mmol, 7.5 equiv) at 0°C under nitrogen atmosphere. To the above mixture was added CuI (150.8 g, 791.9 mmol, 7.5 equiv) in portions over 30 min at 0°C . The resulting mixture was stirred for additional

10 30 min at 0°C . To the above mixture was added tert-butyl(chloro)magnesium (310.5 mL, 527.9


mmol, 5.0 equiv) dropwise over 1 h at 0 °C. The resulting mixture was stirred for additional 5 min at 0 degrees C. The reaction was quenched by the addition of sat. NH₄Cl (aq.) (400 mL) at 0 degrees C. The resulting mixture was extracted with EtOAc (2 x 800 mL). The combined organic layers were washed with brine (2x50 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The resulting mixture ethyl 8-tert-butyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (12-2, 33 g, crude) was used in the next step directly without further purification.

2. Synthesis of ethyl 3-bromo-8-tert-butyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (12-3)




To a stirred solution of ethyl 8-tert-butyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (12-2, 28.0 g, 107.1 mmol, 1.0 equiv) in CHCl₃ (300.0 mL) was added NBS (19.0 g, 107.1 mmol, 1.0 equiv) at room temperature. The resulting mixture was stirred for 1 h at 80 degrees C under nitrogen atmosphere. The mixture was allowed to cool down to room temperature. The resulting mixture was concentrated under vacuum. The residue was purified by silica gel column chromatography, eluted with PE/EtOAc (3:1) to afford ethyl 3-bromo-8-tert-butyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (12-3, 29.5 g, 80.9%) as a yellow solid.

Synthesis of ethyl 8-tert-butyl-3-(3,5-dichlorophenyl)-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (12-4)




To a stirred mixture of ethyl 3-bromo-8-tert-butyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (12-3, 26.0 g, 76.4 mmol, 1.0 equiv) and 3,5-dichlorophenylboronic acid (14.5 g, 76.4 mmol, 1.0 equiv) in THF (240.0 mL) and H_2O (60.0 mL) were added K_2CO_3 (31.7 g, 229.2 mmol, 3.0 equiv) and Pd(dtbpf)Cl_2 (4980.7 mg, 7.6 mmol, 0.1 equiv) at room temperature under nitrogen atmosphere. The resulting mixture was stirred for 2 h at 60 degrees C under nitrogen atmosphere. The mixture was allowed to cool down to room temperature. The resulting mixture was concentrated under vacuum. The resulting mixture was diluted with water (800 mL). The resulting mixture was extracted with EtOAc (2 x 800mL). The combined organic layers were washed with brine (2x50 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE/EtOAc (8:1) to afford ethyl 8-tert-butyl-3-(3,5-dichlorophenyl)-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (12-4, 19 g, 61.2%) as an off-white solid.

3. Synthesis of 8-tert-butyl-3-(3,5-dichlorophenyl)-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (12-5)



Into a 250-mL round-bottom flask, was placed ethyl 8-tert-butyl-3-(3,5-dichlorophenyl)-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (12-4, 18.0 g, 44.3 mmol, 1.0 equiv), MeOH (100.0 mL, 344.3 mmol, 28.0 equiv), THF (100.0 mL), H₂O (200.0 mL) and KOH (53.1 g, 1329.0 mmol, 30.0 equiv). The resulting solution was stirred for 48 hr at 80 °C. The resulting mixture was concentrated under vacuum. The resulting solution was diluted with 50 mL of Water. The pH value of the solution was adjusted to 4 with HCl (2 mol/L). The solids were collected by filtration. The solid was dried in an oven under reduced pressure. This resulted in 8-tert-butyl-3-(3,5-dichlorophenyl)-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (12-5, 18 g, crude) as an off-white solid.

4. Synthesis of 8-tert-butyl-3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (Compound 306)



To a stirred mixture of 8-tert-butyl-3-(3,5-dichlorophenyl)-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (12-5, 17.0 g, 44.9 mmol, 1.0 equiv) and (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (8046.4 mg, 53.9 mmol, 1.2 equiv) in DMF (80.0 mL) were added DIEA (17.4 g, 134.8 mmol, 3.0 equiv) and HATU (20.5 g, 53.9 mmol, 1.2 equiv) at room temperature. The resulting mixture was stirred for 2 h at room temperature. The reaction mixture was added dropwise into 300 mL of H₂O. The precipitated solids were collected by filtration and washed with water (2x20 mL). The residue was dissolved in MeCN (200 mL). Then 800 mL of H₂O was added dropwise. The precipitated solids were collected by filtration. The resulting solid was dried under infrared light to afford 8-tert-butyl-3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (306, 17.2 g, 75.1%) as a light green solid. (300 MHz, CDCl₃, ppm) δ : 8.08 (s, 1H), 7.57 (d, J = 1.8 Hz, 2H), 7.41 (s, 1H), 7.32-7.18 (m, 2H), 6.99-6.95 (m, 1H), 6.88 (dd, J = 8.1, 1.2 Hz, 1H), 6.21 (d, J = 6.6 Hz, 1H),

5.32-5.29 (m, 1H), 4.41-4.34 (m, 1H), 4.24-4.16 (m, 1H), 2.61 (s, 3H), 2.41-2.34 (m, 1H), 2.27-2.21 (m, 1H), 1.78 (s, 9H)


Compound	¹ H NMR Spectra
297	(300 MHz, CDCl ₃ , ppm): δ 8.04 (s, 1H), 7.55 (d, J = 1.8 Hz, 2H), 7.38 (s, 1H), 6.98-6.88 (m, 2H), 6.12 (d, J = 6.9 Hz, 1H), 5.29 (t, J = 5.4 Hz, 1H), 4.34-4.29 (m, 1H), 4.18-4.11 (m, 1H), 2.59 (s, 1H), 2.38-2.31 (m, 1H), 2.19-2.14 (m, 1H), 1.75 (s, 9H)
298	(300 MHz, CDCl ₃ , ppm) δ 8.16 (s, 1H), 7.50-7.46 (m, 3H), 7.26-7.19 (m, 2H), 6.97-6.91 (m, 1H), 6.87 (d, J = 8.4 Hz, 1H), 6.06 (d, J = 7.2 Hz, 1H), 5.33-5.31 (m, 1H), 4.38-4.33 (m, 1H), 4.21-4.13 (m, 1H), 2.39-2.34 (m, 1H), 2.26-2.21 (m, 1H), 1.76 (s, 9H)
298-0	(300 MHz, CDCl ₃ , ppm) δ 8.16 (s, 1H), 7.60 (d, J = 1.8 Hz, 1H), 7.52-7.47 (m, 3H), 6.99-6.93 (m, 2H), 6.90-6.79 (m, 1H), 6.05 (d, J = 7.5 Hz, 1H), 5.34 (q, J = 5.7 Hz, 1H), 4.37-4.34 (m, 1H), 4.19-4.16 (m, 1H), 2.48-2.32 (m, 1H), 2.28-2.12 (m, 1H), 1.79 (s, 9H)
299	(300 MHz, CDCl ₃ , ppm) δ 8.18 (s, 1H), 7.53-7.49 (m, 3H), 6.99-6.93 (m, 2H), 6.87-6.82 (m, 1H), 6.09 (d, J = 7.5 Hz, 1H), 5.37-5.31 (m, 1H), 4.39-4.32 (m, 1H), 4.22-4.14 (m, 1H), 2.42-2.36 (m, 1H), 2.24-2.18 (m, 1H), 1.79 (s, 9H)
299-0	(300 MHz, CDCl ₃ , ppm) 8.16 (s, 1H), 7.60 (s, 1H), 7.57-7.44 (m, 3H), 6.96 (t, J = 8.1 Hz, 2H), 6.92-6.79 (m, 1H), 6.07 (d, J = 7.8 Hz, 1H), 5.35 (q, J = 5.7 Hz, 1H), 4.37-4.34 (m, 1H), 4.24-4.11 (m, 1H), 2.44-2.40 (m, 1H), 2.30-2.13 (m, 1H), 1.79 (s, 9H)
418	(300 MHz, DMSO-d ₆ , ppm) δ 8.33 (s, 1H), 8.11 (s, 1H), 7.95 (s, 1H), 7.35 (s, 1H) 7.25-7.15 (m, 3H), 6.99-6.85 (m, 1H), 6.84-6.80 (m, 1H), 6.25 - 6.20 (m, 1H), 5.35 - 5.30 (m, 1H), 4.36-4.15 (m, 2H), 2.50-2.22 (m, 2H), 1.73 (s, 9H)
420	(300 MHz DMSO-d ₆ , ppm): δ 8.15 (s, 1H), 7.75 (s, 2H) 7.44 (s, 1H), 7.00-6.79 (m, 2H), 6.15 - 6.30 (m, 1H), 5.25 - 5.32 (m, 1H), 4.30 - 4.47 (m, 1H), 4.10 - 4.15 (m, 1H), 2.40-2.38 (m, 1H), 2.25-2.10 (m, 1H), 1.80 (bs, 9H)

523	(300 MHz DMSO-d ₆ , ppm): δ 8.08 (s, 1H), 7.15-7.44 (m, 3H), 7.00-6.79 (m, 2H), 6.10 - 6.21 (m, 1H), 5.36 - 5.43 (m, 1H), 4.30 - 4.49 (m, 1H), 4.10 - 4.18 (m, 1H), 2.40-2.38 (m, 4H), 2.28-2.26 (m, 1H), 1.80 (bs, 9H)
524	(300 MHz DMSO-d ₆ , ppm): δ 9.20 (d, J = 7.9 Hz, 1H), 8.30 (s, 1H), 7.48-7.42 (m, 2H), 7.36-7.34 (m, 1H), 7.20-7.15 (m, 1H), 6.92 (t, J = 7.5, 1H), 6.80 (d, J = 8.2 Hz, 1H), 5.19-5.16 (m, 1H), 4.28-4.19 (m, 2H), 2.35 (s, 3H), 2.20-2.18 (m, 1H), 2.10-2.00 (m, 1H), 1.68 (s, 9H)
525	300 MHz DMSO-d ₆ , ppm): δ 9.20 (d, J = 8.0 Hz, 1H), 8.33 (s, 1H), 7.74-7.68 (m, 1H), 7.47-7.42 (m, 1H), 7.37-7.34 (m, 1H), 7.21-7.15 (m, 1H), 6.94-6.90(m, 1H), 6.80 (d, J = 8.2 Hz, 1H), 5.20-5.17 (m, 1H), 4.28-4.19 (m, 3H), 2.43 (s, 3H), 2.19-2.16 (m, 1H), 2.05-1.99 (m, 1H), 1.68 (s, 9H)
526	(300 MHz DMSO-d ₆ , ppm): δ 8.31 (s, 1H), 7.47-7.44 (m, 2H), 7.26-6.79 (m, 5H), 5.41 (bss, 1H), 4.36 (bs, 2H), 2.55 (bs, 3H), 2.40-2.38 (m, 1H), 2.28-2.26 (m, 1H), 1.80 (bs, 9H)
571	(300 MHz, CDCl ₃ , ppm) δ 8.11 (s, 1H), 7.28-7.21 (m, 2H), 6.97-6.85(m, 4H), 6.12-6.10 (m, 1H), 5.35-5.32 (m, 1H), 4.39-4.33 (m, 1H), 4.22-4.14 (m, 1H), 2.42-2.35 (m, 1H), 2.27-2.21(m, 1H), 1.78 (s, 9H)
572	(400 MHz, CD ₃ OD, ppm) δ 8.21 (s, 1H), 7.48 (d, J = 6.3 Hz, 2H), 7.31 (d, J = 5.7 Hz, 1H), 7.16 (t, J = 5.7 Hz, 1H), 6.94 (t, J = 5.7 Hz, 1H), 6.82 (d, J = 5.7 Hz, 1H), 5.28 (bt, J = 3.6 Hz, 1H), 4.33-4.28 (m, 1H), 4.23-4.17 (m, 1H), 2.30-2.23 (m, 1H), 2.19-2.14 (m, 1H), 1.75 (s, 9H)
573	(300 MHz, CDCl ₃ , ppm) δ 8.11 (s, 1H), 7.24-7.17 (m, 2H), 7.13-7.03 (m, 2H), 6.94 (t, J = 7.5 Hz, 1H), 6.86 (d, J = 8.2 Hz, 1H), 6.09-6.08 (m, 1H), 5.32-5.30 (m, 1H), 4.36-4.33 (m, 1H), 4.19-4.14 (m, J = 10.5 Hz, 1H), 2.39-2.36 (m, 1H), 2.23-2.20 (m, 1H), 1.75 (s, 9H)
574	(300 MHz, CDCl ₃ , ppm) δ 8.12 (d, J = 3.3 Hz, 1H), 7.44-7.40 (m, 1H), 7.25-7.10 (m, 2H), 6.99-6.93 (m, 1H), 6.90-6.87 (m, 1H), 6.10 (m, 1H), 5.35-5.34 (m, 1H), 4.39-4.36 (m, 1H), 4.22-4.15 (m, 1H), 2.43-2.36 (m, 1H), 1.79 (s, 9H)
A472	(400 MHz, DMSO-d ₆ , ppm) δ 9.02 (d, J = 8.49 Hz, 1H), 8.53 (s, 1H), 7.78 (d, J = 1.90 Hz, 2H), 7.68 (t, J = 1.84 Hz, 1H), 7.36 (d, J = 7.73 Hz, 1H), 7.17 (t, J =

7.71 Hz, 1H), 6.92 (t, J = 7.26 Hz, 1H), 6.76 (d, J = 8.11 Hz, 1H), 5.32 (m, 1H), 3.52 - 3.62 (m, 1H), 2.52 - 2.58 (m, 3H), 2.21 (m, 1H), 1.80 - 1.89 (m, 1H), 1.23 - 1.61 (m, 12H)


Preparation Example 8: The following compounds can be synthesized by adopting the process shown in scheme 13 below: 320, 320-0, 513, 513-0, 514, 514-0.

Scheme 13




5

1. Synthesis of ethyl 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-5)




Into a 40-mL round-bottom flask, was placed THF (5.0 mL), H₂O (1.0 mL, 0.06 mmol, 0.18 equiv), ethyl 3-bromo-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-4, 100.0 mg, 0.3 mmol, 1.0 equiv), 3,5-difluorophenylboronic acid (145.0 mg, 0.9 mmol, 3.0 equiv), Pd(dtbpf)Cl₂ (20.0 mg, 0.03 mmol, 0.1 equiv), K₂CO₃ (85.0 mg, 0.6 mmol, 2.0 equiv). The resulting solution was stirred for 2 hr at 70 °C. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:6). This resulted in 90 mg (81.7%) of ethyl 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-5) as a white solid.

2. Synthesis of 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (4-6)




Into a 40-mL round-bottom flask, was placed H₂O (1.0 mL), EtOH (5.0 mL), ethyl 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-5, 90.0 mg, 0.25 mmol, 1.0 equiv), LiOH (60.0 mg, 2.5 mmol, 10.0 equiv). The resulting solution was stirred for 2 hr at 50 °C. HCl (6 mol/L) was employed to adjust the pH to 4. The resulting solution was extracted with 3x20 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 70 mg (84.4%) of 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (4-6) as a white solid.

3. Synthesis of tert-butyl N-[3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazin-7-yl]carbamate (13-1)



Into a 40-mL round-bottom flask, was placed DMF (1.0 mL, 0.01 mmol, 0.06 equiv), t-BuOH (1.0 mL, 0.01 mmol, 0.06 equiv), 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (4-6, 70.0 mg, 0.2 mmol, 1.0 equiv), DPPA (70.0 mg, 0.25 mmol, 1.2 equiv), TEA (24.0 mg, 0.2 mmol, 1.1 equiv). The resulting solution was stirred for 3 hr at 50 degrees C. The mixture was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O:ACN=90:10 increasing to H₂O:ACN=20:80 within 15 min; Detector, 254nm. This resulted in 65 mg (76.4%) of tert-butyl N-[3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazin-7-yl]carbamate (13-1) as a white solid.


4. Synthesis of 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazin-7-amine (13-2)



Into a 50-mL round-bottom flask, was placed HCl (gas) in 1,4-dioxane (4M, 5.00 mL), tert-butyl N-[3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazin-7-yl]carbamate (13-1, 65.0 mg, 0.2 mmol, 1.0 equiv). The resulting solution was stirred for 2 hr at 40 degrees C. The resulting mixture was concentrated under vacuum. This resulted in 40 mg

(81.9%) of 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazin-7-amine (13-2) as a white solid.

5. Synthesis of (4S)-N-[3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazin-7-yl]-3,4-dihydro-2H-1-benzopyran-4-carboxamide and (4R)-N-[3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazin-7-yl]-3,4-dihydro-2H-1-benzopyran-4-carboxamide (320 and 320-0)




Into a 50-mL round-bottom flask, was placed DCM (2.0 mL), 3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazin-7-amine (13-2, 25.0 mg, 0.08 mmol, 1.0 equiv), (4S)-3,4-dihydro-2H-1-benzopyran-4-carbonyl chloride (50.0 mg, 0.2 mmol, 3.0 equiv), DIEA (0.5 mg, 0.004 mmol, 0.05 equiv). The resulting solution was stirred for 1 hr at room temperature. The resulting mixture was concentrated under vacuum. The crude product was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O:ACN=50:50 increasing to H₂O:ACN=10:90 within 20 ; Detector, 254 nm. This resulted in 9.3 mg (24.3%) of (4S)-N-[3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazin-7-yl]-3,4-dihydro-2H-1-benzopyran-4-carboxamide (320) as a white solid and 11 mg (26.0%) of (4R)-N-[3-(3,5-difluorophenyl)-8-isopropyl-2-methylimidazo[1,2-b]pyridazin-7-yl]-3,4-dihydro-2H-1-benzopyran-4-carboxamide (320-0). ¹H NMR for 320: (300 MHz Chloroform-d, ppm): δ 8.93 (s, 1H), 7.50-7.43 (m, 1H), 7.37-7.30 (m, 1H), 7.27-7.25 (m, 2H), 7.08-7.00 (m, 2H), 6.89-6.81 (m, 1H), 4.44-4.40 (m, 1H), 4.16-4.07 (m, 1H), 3.92 (brs, 1H), 3.70-3.66 (m, 1H), 2.70-2.63 (m, 1H), 2.61 (s, 3H), 2.34-2.28 (m, 1H), 1.20-1.14 (m, 6H); ¹H NMR for 320-0: (300 MHz Chloroform-d, ppm): δ 8.94 (s, 1H), 7.50-7.44 (m, 1H), 7.36-7.31 (m, 1H), 7.27-7.20 (m, 2H), 7.15-7.00 (m, 2H), 6.94-6.82 (m, 1H), 4.44-4.40 (m, 1H), 4.14-4.07 (m, 1H), 3.92 (brs, 1H), 3.70-3.60 (m, 1H), 2.70-2.50 (m, 4H), 2.37-2.24 (m, 1H), 1.19-1.14 (m, 6H).

Compound	¹ H NMR Spectra
514	(300 MHz, CDCl ₃ , ppm) δ 8.68 (s, 1H), 8.67 (bs, 1H), 8.44 (bs, 1H), 7.66 (bs, 1H), 7.63 (s, 1H), 7.60 (d, J = 1.8 Hz, 1H), 7.50 (s, 1H), 6.94 (bs, 1H), 4.54-4.50 (m, 1H), 4.35-4.27 (m, 1H), 4.05 (bs, 1H), 3.71-3.62 (m, 1H), 2.74-2.69 (m, 1H), 2.59 (s, 3H), 2.30-2.20 (m, 1H), 1.32-1.29 (m, 6H)


Preparation Example 9: Compounds 323 and 323-0 can be synthesized according to the process depicted in Scheme 14 below.

Scheme 14




5

1. Synthesis of ethyl 3-tert-butyl-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (14-2a) and ethyl 3-isobutyl-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (14-2b)




Into a 8-mL round-bottom flask, was placed THF (3.0 mL), ethyl 3-bromo-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-4, 100.0 mg, 0.3 mmol, 1.00equiv), di-tert-butylzinc (4.0 mL, 2.0 mmol, 6.5 equiv), Pd(PPh₃)₄ (40.0 mg, 0.03 mmol, 0.1 equiv). The resulting solution was stirred for 1 overnight at 80 °C. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:4). This resulted in 20 mg (mixture) of ethyl 3-tert-butyl-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (14-2a) and ethyl 3-isobutyl-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (14-2b) as a white solid.

2. Synthesis of 3-tert-butyl-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (14-3a) and 3-isobutyl-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (14-3b)



Into a 50-mL round-bottom flask, was placed H₂O (1.0 mL), EtOH (5.0 mL), mixture of ethyl 3-tert-butyl-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate and 3-isobutyl-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (mix of 14-2a and 14-2b, 20.0 mg, 0.06mmol, 1.00 equiv), NaOH (40.0 mg, 1.00 mmol, 15.2 equiv). The resulting solution was stirred for 2 hr at 80 °C. HCl (6 mol/L) was employed to adjust the pH to 4. The resulting solution was extracted with 3x20 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 12 mg (mixture) of 3-tert-butyl-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (14-3a) and 3-isobutyl-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (14-3b) as a white solid.


3. Synthesis of 3-tert-butyl-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (323) and N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-2-methyl-3-(2-methylpropyl)-8-(propan-2-yl)imidazo[1,2-b]pyridazine-7-carboxamide (323-0)




Into a 50-mL round-bottom flask, was placed DMF (1.0 mL, 12.9 mmol, 237.2 equiv), a mixture of 3-tert-butyl-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid and 3-isobutyl-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (mix of 14-3a and 14-3b, 15.0 mg, 0.05 mmol, 1.0 equiv), (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (10.0 mg, 0.06 mmol, 1.2 equiv), HATU (35.0 mg, 0.09 mmol, 1.7 equiv), DIEA (17.0 mg, 0.1 mmol, 2.4 equiv). The resulting solution was stirred for 1 hr at room temperature. The crude product was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O:ACN=50:50 increasing to H₂O:ACN=10:90 within 20 min; Detector, 254nm. product was obtained. This resulted in 2.2 mg (9.9%) of 3-tert-butyl-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (323) as a white solid and 5 mg (21%) of N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-2-methyl-3-(2-methylpropyl)-8-(propan-2-yl)imidazo[1,2-b]pyridazine-7-carboxamide (323-0). ¹H NMR for 323: (300 MHz Chloroform-d, ppm): δ 8.22 (s, 1H), 7.40-7.30 (m, 1H), 7.25-7.20 (m, 1H), 6.99-6.96 (m, 1H), 6.91-6.88 (m, 1H), 6.11-5.90 (m, 1H), 5.45-5.30 (m, 1H), 4.41-4.37 (m, 1H), 4.30-4.14 (m, 1H), 3.90-.70 (m, 1H), 2.68 (s, 3H), 2.50-2.30 (s, 1H), 2.30-2.15 (m, 1H), 1.75-1.70 (m, 6H), 1.55 (s, 9H); ¹H NMR for 323-0: (300 MHz Chloroform-d, ppm): δ 8.24 (s, 1H), 7.40-7.31 (m, 1H), 7.27-7.21 (m, 1H), 6.99-6.90 (m, 1H), 6.88-6.85 (m, 1H), 6.10 (brs, 1H), 5.41-5.37 (m, 1H), 4.40-4.34 (m, 1H), 4.25-4.17 (m, 1H), 3.80 (brs, 1H), 2.85 (d, J = 7.2 Hz, 2H), 2.53 (s, 3H), 2.44-2.38 (m, 1H), 2.28-2.10 (m, 2H), 1.60 (t, J = 6.6 Hz, 6H), 0.95 (d, J = 6.6 Hz, 6H).

Preparation Example 10: Compound 324 can be synthesized by the process shown in Scheme 15 shown below. Similarly, compounds 325, 369, 372-0, 373 can be prepared by an analogous methods by persons skilled in the art.

Scheme 15



1. Synthesis of ethyl 8-isopropyl-2-methyl-3-(piperidin-1-yl)imidazo[1,2-b]pyridazine-7-carboxylate (15-1)




5

Into a 8-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed Toluene (2.0 mL), ethyl 3-bromo-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-4, 70.0 mg, 0.2 mmol, 1.0 equiv), piperidine (120.0 mg, 1.4 mmol, 6.6 equiv), Pd₂(dba)₃ (35.0 mg, 0.04 mmol, 0.2 equiv), BINAP (38.0 mg, 0.06 mmol, 0.3 equiv), Cs₂CO₃ (200.0 mg, 0.6 mmol, 2.9 equiv). The resulting solution was stirred for 2 hr at 120 °C. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:5). This resulted in 60 mg (84.6%) of ethyl 8-isopropyl-2-methyl-3-(piperidin-1-yl)imidazo[1,2-b]pyridazine-7-carboxylate (15-1) as a yellow solid.


10

2. Synthesis of 8-isopropyl-2-methyl-3-(piperidin-1-yl)imidazo[1,2-b]pyridazine-7-carboxylic acid (15-2)



Into a 50-mL round-bottom flask, was placed i-PrOH (1.0 mL, 0.02 mmol), THF (1.0 mL),
 5 H₂O (1.0 mL), ethyl 8-isopropyl-2-methyl-3-(piperidin-1-yl)imidazo[1,2-b]pyridazine-7-carboxylate (15-1, 70.0 mg, 0.2 mmol, 1.0 equiv), LiOH.H₂O (30.0 mg, 0.7 mmol, 3.4 equiv). The resulting solution was stirred for 1 hr at 50 °C. HCl (6 mol/L) was employed to adjust the pH to 4. The resulting solution was extracted with 3x20 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted
 10 in 60 mg (93.7%) of 8-isopropyl-2-methyl-3-(piperidin-1-yl)imidazo[1,2-b]pyridazine-7-carboxylic acid (15-2) as a yellow solid.

3. Synthesis of N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methyl-3-(piperidin-1-yl)imidazo[1,2-b]pyridazine-7-carboxamide (324)




15 Into a 50-mL round-bottom flask, was placed DMA (3.0 mL), 8-isopropyl-2-methyl-3-(piperidin-1-yl)imidazo[1,2-b]pyridazine-7-carboxylic acid (15-2, 60.0 mg, 0.2 mmol, 1.0 equiv), (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (44.0 mg, 0.3 mmol, 1.5 equiv), HATU (113.0 mg, 0.3 mmol, 1.5 equiv), DIEA (51.0 mg, 0.4 mmol, 2.0 equiv). The resulting solution was stirred for 1 hr at room temperature. The mixture was purified by Flash-Prep-HPLC with the following
 20 conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O:ACN=50:50 increasing to

H₂O:ACN=10:90 within 20; Detector, 254 nm. This resulted in 36.4 mg (42.3%) of N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methyl-3-(piperidin-1-yl)imidazo[1,2-b]pyridazine-7-carboxamide (324) as a yellow solid. (300 MHz, Chloroform-d, ppm): δ 8.19 (s, 1H), 7.33-7.30 (m, 1H), 7.25-7.20 (m, 1H), 6.97-6.90 (m, 1H), 6.89-6.80 (m, 1H), 6.10-6.02 (m, 1H), 5.40-5.30 (m, 1H), 4.38-4.34 (m, 1H), 4.22-4.10 (m, 1H), 3.79-3.76 (m, 1H), 3.30-3.20 (m, 4H), 2.52 (s, 3H), 2.50-2.35 (m, 1H), 2.30-2.15 (m, 1H), 1.80-1.70 (m, 4H), 1.65-1.58 (m, 8H).


Compound	¹ H NMR Spectra
325	(300 MHz, Chloroform-d, ppm): δ 8.20 (s, 1H), 7.34-7.27 (m, 1H), 7.25-7.21 (m, 1H), 6.98-6.90 (m, 1H), 6.90-6.80 (m, 1H), 6.15-6.00 (m, 1H), 5.40-5.30 (m, 1H), 4.45-4.34 (m, 1H), 4.24-4.15 (m, 1H), 3.91-3.83 (m, 4H), 3.80-3.70 (m, 1H), 3.40-3.20 (m, 4H), 2.52 (s, 3H), 2.49-2.32 (m, 1H), 2.30-2.15 (m, 1H), 1.70-1.50 (m, 6H)
369	(300 MHz, DMSO-d ₆ , ppm) δ 8.98 (d, J = 7.8 Hz, 1H), 8.26 (d, J = 2.7 Hz, 1H), 7.31 (d, J = 7.2 Hz, 1H), 7.17 (t, J = 7.5 Hz, 1H), 6.92 (t, J = 7.2 Hz, 1H), 6.79 (d, J = 8.1 Hz, 1H), 5.30-5.15 (m, 1H), 4.25 (brs, 2H), 3.90 (s, 1H), 3.62-3.50 (m, 1H), 3.40-3.31 (m, 2H), 2.73 (brs, 1H), 2.42 (s, 3H), 2.30-2.15 (m, 1H), 2.10-2.00 (m, 1H), 1.90-1.80 (m, 1H), 1.75-1.40 (m, 11H)
372-0	(300 MHz, Chloroform-d, ppm): δ 8.20 (s, 1H), 7.35-7.30 (m, 1H), 7.26-7.20 (m, 1H), 6.99-6.93 (m, 1H), 6.88 (dd, J = 7.2, 1.2 Hz, 1H), 6.05 (d, J = 7.5 Hz, 1H), 5.45-5.35 (m, 1H), 4.40-4.33 (m, 1H), 4.24-4.18 (m, 1H), 3.80-3.70 (m, 1H), 3.50-3.30 (m, 4H), 2.53 (s, 3H), 2.41-2.32 (m, 1H), 2.27-2.15 (m, 1H), 2.13-1.98 (m, 4H), 1.62 (t, J = 7.2 Hz, 6H)
373	(300 MHz, DMSO-d ₆ , ppm): δ 9.01 (d, J = 8.4 Hz, 1H), 8.32 (s, 1H), 7.32 (d, J = 6.6 Hz, 1H), 7.20-7.14 (m, 1H), 6.95-6.89 (m, 1H), 6.80 (d, J = 6.6 Hz, 1H), 5.25-5.19 (m, 1H), 4.27-4.22 (m, 2H), 3.58-3.53 (m, 1H), 3.41-3.30 (m, 4H), 2.38 (s, 3H), 2.25-2.15 (m, 1H), 2.12-1.96 (m, 1H), 1.52-1.46 (m, 6H), 0.88 (t, J = 6.3 Hz, 4H), 0.14 (s, 6H)

Preparation Example 11: Compounds 327, 326, 326-0, 365, 370, 371 may be prepared by the process shown in Scheme 16 below:

Scheme 16




1. Synthesis of ethyl 8-isopropyl-2-methyl-3-[4-(trifluoromethyl)cyclohex-1-en-1-yl]imidazo[1,2-b]pyridazine-7-carboxylate (16-1)




Into a 8-mL round-bottom flask, was placed dioxane (2.0 mL), H₂O (0.4 mL), ethyl 3-bromo-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (4-4, 100.00mg, 0.3 mmol, 1.0 equiv), 4-(trifluoromethyl)cyclohex-1-en-1-ylboronic acid (120.0 mg, 0.6 mmol, 2.0 equiv), Pd(dtbpf)Cl₂ (20.0 mg, 0.03 mmol, 0.1 equiv), K₂CO₃ (100.0 mg, 0.7 mmol, 2.4 equiv). The resulting solution was stirred for 2 hr at 80 °C. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:5). This resulted in 110 mg (90.7%) of ethyl 8-isopropyl-2-methyl-3-[4-(trifluoromethyl)cyclohex-1-en-1-yl]imidazo[1,2-b]pyridazine-7-carboxylate (16-1) as a yellow solid.

2. Synthesis of 8-isopropyl-2-methyl-3-[4-(trifluoromethyl)cyclohex-1-en-1-yl]imidazo[1,2-b]pyridazine-7-carboxylic acid (16-2)



Into a 50-mL round-bottom flask, was placed i-PrOH (2.0 mL), THF (2.0 mL), H₂O (1.0 mL), ethyl 8-isopropyl-2-methyl-3-[4-(trifluoromethyl)cyclohex-1-en-1-yl]imidazo[1,2-b]pyridazine-7-carboxylate (16-1, 110.0 mg, 0.3 mmol, 1.0 equiv), LiOH (70.0 mg, 2.9 mmol, 10.5 equiv). The resulting solution was stirred for 1 hr at 50 °C. The pH value of the solution was adjusted to 4 with HCl (6 mol/L). The resulting solution was extracted with 3x20 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 90 mg (88.0%) of 8-isopropyl-2-methyl-3-[4-(trifluoromethyl)cyclohex-1-en-1-yl]imidazo[1,2-b]pyridazine-7-carboxylic acid (16-2) as a white solid.


3. Synthesis of N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methyl-3-[4-(trifluoromethyl)cyclohex-1-en-1-yl]imidazo[1,2-b]pyridazine-7-carboxamide (327)



Into a 40-mL round-bottom flask, was placed DMF (5.0 mL), 8-isopropyl-2-methyl-3-[4-(trifluoromethyl)cyclohex-1-en-1-yl]imidazo[1,2-b]pyridazine-7-carboxylic acid (16-2, 130.0 mg, 0.3 mmol, 1.0 equiv), (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (79.2 mg, 0.5 mmol, 1.5 equiv), HATU (200.5 mg, 0.5 mmol, 1.5 equiv), DIEA (137.2 mg, 1.0 mmol, 3.0 equiv). The resulting solution was stirred for 1 hr at room temperature. The mixture was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O:ACN=50:50 increasing to H₂O:ACN=10:90 within 20 min; Detector, 254 nm. This resulted in 130 mg (73.7%) of N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methyl-3-[4-(trifluoromethyl)cyclohex-1-en-1-yl]imidazo[1,2-b]pyridazine-7-carboxamide (327) as a white solid. (300 MHz, Chloroform-d, ppm): δ 8.20 (s, 1H), 7.35-7.30 (m, 1H), 7.25-7.17 (m, 1H), 7.10-6.84 (m, 2H), 6.15-5.90 (m, 2H), 5.45-5.30 (m, 1H), 4.44-4.31 (m, 1H), 4.23-4.19 (m, 1H), 3.83-3.72 (m, 1H), 2.75-2.55 (m, 3H), 2.53 (s, 3H), 2.50-2.35 (m, 3H), 2.29-2.13 (m, 3H), 1.63 (d, J = 6.6 Hz, 6H).

4. Synthesis of N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methyl-3-[(1r,4r)-4-(trifluoromethyl)cyclohexyl]imidazo[1,2-b]pyridazine-7-carboxamide and N-((S)-chroman-4-yl)-8-isopropyl-2-methyl-3-((1s,4R)-4-(trifluoromethyl)cyclohexyl)imidazo[1,2-b]pyridazine-7-carboxamide (326 and 326-0)

5



Into a 50-mL round-bottom flask, was placed EA (5.0 mL), N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methyl-3-[4-(trifluoromethyl)cyclohex-1-en-1-yl]imidazo[1,2-b]pyridazine-7-carboxamide (327, 50.0 mg, 0.10 mmol, 1.0 equiv), aqueous Pd/C (50.0 mg). To the above H₂(g) was introduced at rt. The resulting solution was stirred for 2 hr at room temperature. The solids were filtered out. The resulting mixture was concentrated under vacuum. The crude product was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O:ACN=50:50 increasing to H₂O:ACN=10:90 within 20 min; Detector, 254 nm. The racemic product was purified by Column: XA-YMC Cellulose-SC, 4.6*100mm, 3μm,; Mobile Phase A/ Mobile Phase B: n-Hexane/EtOH=70/30; Flow rate: 1 mL/min; Gradient: 30B to 30 B in 10 min; 254 nm; Injection Volume: 1 ml; This resulted in 19.3 mg (38.4%) of N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methyl-3-[(1r,4r)-4-(trifluoromethyl)cyclohexyl]imidazo[1,2-b]pyridazine-7-carboxamide as a white solid and 8.0 mg (15.8%) of N-((S)-chroman-4-yl)-8-isopropyl-2-methyl-3-((1s,4R)-4-(trifluoromethyl)cyclohexyl)imidazo[1,2-b]pyridazine-7-carboxamide as a white solid (326 and 326-0). The stereochemical depictions are assumed. ¹H NMR spectra for 326: (300 MHz Chloroform-d, ppm): δ 8.23 (s, 1H), 7.35-7.30 (m, 1H), 7.27-7.20 (m, 1H), 7.02-6.94 (m, 1H), 6.90-6.80 (m, 1H), 6.07 (brs, 1H), 5.44-5.34 (m, 1H), 4.44-4.31 (m, 1H), 4.28-4.14 (m, 1H), 3.90-3.80 (m, 1H), 3.45-3.25 (m, 1H), 2.58 (s, 3H), 2.55-2.40 (m, 2H), 2.45-2.15 (m, 4H), 1.85-1.65 (m, 5H), 1.62 (d, J = 6.6 Hz, 6H); ¹H NMR spectra for 326-0: (300 MHz, Chloroform-d, ppm): δ 8.17 (s, 1H), 7.30-7.28 (m, 1H), 7.25-7.15(m, 1H), 6.94 (t, J = 7.5 Hz, 1H), 6.86 (d, J = 8.2 Hz, 1H), 6.15-5.90 (m, 1H), 5.45-5.20 (m, 1H), 4.40-4.30 (m, 1H), 4.25-4.10 (m, 1H), 3.81-3.65 (m, 1H),

3.30-3.10 (m, 1H), 2.53 (s, 3H), 2.46-2.31 (m, 1H), 2.30-2.00 (m, 6H), 2.00-1.85 (m, 2H), 1.65-1.55 (m, 6H), 1.50-1.45 (m, 1H).

Compound	¹ H NMR Spectra
365	(300 MHz, CDCl ₃ , ppm): δ 8.25 (s, 1H), 7.31-7.28 (m, 1H), 7.26-7.20 (m, 1H), 6.99-6.94 (m, 1H), 6.88 (d, J = 8.1 Hz, 1H), 6.06 (d, J = 7.2 Hz, 1H), 5.40-5.30 (m, 1H), 4.40-4.30 (m, 1H), 4.24-4.10 (m, 1H), 3.80-3.70 (m, 1H), 2.54 (s, 3H), 2.49-2.33 (m, 1H), 2.25-2.15 (m, 1H), 2.05-1.90 (m, 1H), 1.64-1.56 (m, 6H), 1.15-1.05 (m, 2H), 0.95-0.80 (m, 2H)
370	(300 MHz, CDCl ₃ , ppm): δ 8.25 (s, 1H), 7.34-7.31 (m, 1H), 7.27-7.22 (m, 1H), 6.99-6.95 (m, 1H), 6.89 (d, J = 8.1 Hz, 1H), 6.16 (brs, 1H), 5.45-5.35 (m, 1H), 4.40-4.34 (m, 1H), 4.28-4.18 (m, 1H), 4.16-4.09 (m, 2H), 3.80 (brs, 1H), 3.63-3.47 (m, 3H), 2.61 (s, 3H), 2.60-2.15 (m, 4H), 1.75-1.70 (m, 2H), 1.62 (t, J = 6.6 Hz, 6H)
371	(300 MHz, CDCl ₃ , ppm): δ 8.22 (s, 1H), 7.31-7.28 (m, 1H), 7.24-7.21 (m, 1H), 6.99-6.94 (m, 1H), 6.90-6.87 (m, 1H), 6.07-6.05 (m, 1H), 5.41-5.35 (m, 1H), 4.41-4.34 (m, 1H), 4.27-4.17 (m, 1H), 3.78-3.74 (m, 1H), 3.38-3.29 (m, 1H), 2.56 (s, 3H), 2.54-2.22 (m, 6H), 2.03-1.86 (m, 4H), 1.62 (t, J = 6.9 Hz, 6H)

Preparation Example 12: Compound 352 was prepared according to the process shown in scheme 17 below:

5

17-1 $\xrightarrow[\text{LiHMDS, THF}]{\text{ethyl acetate}}$ 17-2 $\xrightarrow[\text{DMF, rt, 4 h}]{\text{DMF-DMA, K}_2\text{CO}_3}$ 17-3


17-3 $\xrightarrow[\text{H}_2\text{O, 100}^\circ\text{C, 2h}]{\text{Pd(dtbpf)Cl}_2, \text{Cs}_2\text{CO}_3, \text{NMP}}$ 17-4 $\xrightarrow[\text{DMF, CHCl}_3]{(\text{COCl})_2}$ 17-5

17-5 $\xrightarrow[\text{H}_2\text{O, 80}^\circ\text{C, 2h}]{\text{Pd(dtbpf)Cl}_2, \text{K}_2\text{CO}_3, \text{THF}}$ 17-6 $\xrightarrow{\text{PtO}_2, \text{EA, H}_2(\text{g})}$ 17-7

17-7 $\xrightarrow{\text{LiOH, EtOH, H}_2\text{O}}$ 17-8 $\xrightarrow[\text{H}_2\text{N}-(S)\text{-bicyclo[3.1.0]hexan-6-ol}]{\text{HATU, DIEA, DMF}}$


352

1. Synthesis of ethyl 3-[5-bromo-1-[(tert-butoxycarbonyl)amino]imidazol-2-yl]-3-oxopropanoate (17-2)




Into a 1000-mL round-bottom flask, was placed THF (85.0 g, 1178.8 mmol, 26.3 equiv), ethyl 5-bromo-1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (17-1, 15.0 g, 44.9 mmol, 1.0 equiv), ethyl acetate (50.0 g, 567.5 mmol, 12.6 equiv). This was followed by the addition of t-BuOK (500 mL), in portions at 0 degrees C. The resulting solution was stirred for 2 hr at room temperature. The reaction was then quenched by the addition of NH₄Cl (aq). The resulting solution was extracted with 3x200 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:4). This resulted in 13 g (77.0%) of ethyl 3-[5-bromo-1-[(tert-butoxycarbonyl)amino]imidazol-2-yl]-3-oxopropanoate (17-2) as colorless oil.

2. Synthesis of ethyl 3-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (17-3)




Into a 500-mL round-bottom flask, was placed DCM (100.0 mL, 1573.0 mmol, 59.2 equiv), ethyl 3-[5-bromo-1-[(tert-butoxycarbonyl)amino]imidazol-2-yl]-3-oxopropanoate (17-2, 10.0 g, 26.6 mmol, 1.0 equiv), DMF-DMA (9.0 g, 75.5 mmol, 2.8 equiv). The resulting solution was stirred for 2 hr at 40 °C. The reaction was then quenched by the addition of water/ice. The resulting solution was extracted with 2x100 mL of MTBE and the aqueous layers combined. The pH value of the solution was adjusted to 4 with HCl (4 mol/L). The solids were collected by filtration. This resulted in 5 g (65.7%) of ethyl 3-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (17-3) as a white solid.

3. Synthesis of ethyl 3-(3,5-dichlorophenyl)-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (17-4)



Into a 40-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed H₂O (1.0 mL), NMP (5.0 mL), ethyl 3-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (17-3, 400.0 mg, 1.398 mmol, 1.0 equiv), 3,5-dichlorophenylboronic acid (320.1 mg, 1.7 mmol, 1.2 equiv), Pd(dtbpf)Cl₂ (70.0 mg, 0.1 mmol, 0.08 equiv), Cs₂CO₃ (1.2 g, 3.7 mmol, 2.6 equiv). The resulting solution was stirred for 2 hrs at 100 °C. The mixture was cooled to rt, then added 5ml H₂O. The solids were collected by filtration. This resulted in 300 mg (crude) of ethyl 3-(3,5-dichlorophenyl)-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (17-4) as a yellow solid.


4. Synthesis of ethyl 8-chloro-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (17-5)



Into a 40-mL round-bottom flask, was placed CHCl₃ (5.0 mL, 0.04 mmol, 0.07 equiv), DMF (15.0 mg, 0.2 mmol, 0.4 equiv), ethyl 3-(3,5-dichlorophenyl)-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (17-4, 200.0 mg, 0.6 mmol, 1.0 equiv), (COCl)₂ (400.0 mg, 3.1 mmol, 5.5 equiv). The resulting solution was stirred for 22 hr at 80 degrees C. The resulting mixture was concentrated under vacuum. The resulting mixture was washed with 10x10 mL of ACN:H₂O=1:1. The solids

were collected by filtration. This resulted in 110 mg (52.3%) of ethyl 8-chloro-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (17-5) as a yellow solid.


5. Synthesis of ethyl 3-(3,5-dichlorophenyl)-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (17-6)



Into a 40-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed H₂O (1.00 mL, 55.5 mmol, 187.0 equiv), THF (5.0 mL, 61.7 mmol, 207.9 equiv), ethyl 8-chloro-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (17-5, 110.0 mg, 0.3 mmol, 1.0 equiv), 4,4,5,5-tetramethyl-2-(prop-1-en-2-yl)-1,3,2-dioxaborolane (72.0 mg, 0.43 mmol, 1.4 equiv), Pd(dtbpf)Cl₂ (40.0 mg, 0.06 mmol, 0.2 equiv), K₂CO₃ (160.0 mg, 1.2 mmol, 3.9 equiv). The resulting solution was stirred for 2 hr at 80 °C. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:3). This resulted in 80 mg (71.6%) of ethyl 3-(3,5-dichlorophenyl)-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (17-6) as a yellow solid.

10

6. Synthesis of ethyl 3-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylate (17-7)




Into a 50-mL round-bottom flask, was placed EA (5.0 mL), ethyl 3-(3,5-dichlorophenyl)-8-(prop-1-en-2-yl)imidazo[1,2-b]pyridazine-7-carboxylate (17-6, 80.0 mg, 0.21 mmol, 1.0 equiv), PtO₂ (40.0 mg, 0.2 mmol, 0.8 equiv). To the above H₂(g) was introduced with a balloon. The

20


resulting solution was stirred for 1 hr at 50 °C. The solids were collected by filtration. The resulting mixture was concentrated under vacuum. This resulted in 60 mg of ethyl 3-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylate (17-7) as a yellow solid.

7. Synthesis of 3-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylic acid (17-8)



Into a 8-mL round-bottom flask, was placed EtOH (2.0 mL), H₂O (0.50 mL), ethyl 3-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylate (17-7, 60.0 mg, 0.2 mmol, 1.0 equiv), LiOH (30.0 mg, 1.2 mmol, 7.9 equiv). The resulting solution was stirred for 1 hr at 50 °C. The reaction was then quenched by the addition of water/ice. The pH value of the solution was adjusted to 4 with HCl (6 mol/L). The resulting solution was extracted with 3x20 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 40 mg (72.0%) of 3-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylic acid (17-8) as a white solid.

8. Synthesis of 3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropylimidazo[1,2-b]pyridazine-7-carboxamide (352)




Into a 8-mL round-bottom flask, was placed DMF (4.0 mL), 3-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylic acid (17-8, 40.0 mg, 0.1 mmol, 1.0 equiv), (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (22.0 mg, 0.15 mmol, 1.3 equiv), HATU (70.0 mg, 0.2


mmol, 1.6 equiv), DIEA (46.0 mg, 0.4 mmol, 3.1 equiv). The resulting solution was stirred for 1 hr at room temperature. The crude product was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O:ACN=50:50 increasing to H₂O:ACN=10:90 within 20 min ; Detector, 254nm. This resulted in 15.9 mg (28.9%) of 3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropylimidazo[1,2-b]pyridazine-7-carboxamide (352) as a white solid. (300 MHz, Chloroform-d, ppm): δ 8.41 (s, 1H), 8.11 (s, 1H), 7.99 (d, J = 1.8 Hz, 2H), 7.39 (s, 1H), 7.32-7.28 (m, 1H), 7.28-7.22 (m, 1H), 6.99-6.96 (m, 1H), 6.90 (d, J = 8.4, 1H), 6.13 (d, J = 7.5 Hz, 1H), 5.50-5.30 (m, 1H), 4.46-4.33 (m, 1H), 4.29-4.15 (m, 1H), 3.79-3.70 (m, 1H), 2.50-2.36 (m, 1H), 2.33-2.20 (m, 1H), 1.69-1.65 (m, 6H).

Preparation Example 13: Compound 366 was prepared according to scheme 18 shown below:

Scheme 18




1. Synthesis of ethyl 3-cyano-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate



Into a 20-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed DMAC (5.0 mL), ethyl 3-bromo-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-


carboxylate (4-4, 100.0 mg, 0.3 mmol, 1.0 equiv), dppf (59.3 mg, 0.1 mmol, 0.3 equiv), $\text{Zn}(\text{CN})_2$ (100.1 mg, 0.8 mmol, 2.8 equiv), $\text{Pd}_2(\text{dba})_3$ (50.5 mg, 0.05 mmol, 0.2 equiv), Zn (100.1 mg, 1.5 mmol, 5.0 equiv). The resulting solution was stirred for 2 hr at 100 °C. The reaction was then quenched by the addition of water. The resulting solution was extracted with 3x20 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:5). This resulted in 40 mg (47.9%) of ethyl 3-cyano-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (18-1) as yellow oil.

2. Synthesis of 3-cyano-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (18-2)



Into a 8-mL round-bottom flask, was placed EtOH (1.0 mL), H₂O (0.5 mL), ethyl 3-cyano-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylate (18-1, 40.0 mg, 0.15 mmol, 1.0 equiv), LiOH (30.0 mg, 1.2 mmol, 8.5 equiv). The resulting solution was stirred for 1 hr at room temperature. The pH value of the solution was adjusted to 4 with HCl (4 mol/L). The resulting solution was extracted with 3x20 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 25 mg (69.7%) of 3-cyano-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (18-2) as yellow oil.


3. Synthesis of 3-cyano-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (366)




Into a 8-mL round-bottom flask, was placed DMF (2 mL), 3-cyano-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (18-2, 25.0 mg, 0.1 mmol, 1.0 equiv), (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (16.9 mg, 0.1 mmol, 1.1 equiv), HATU (47.0 mg, 0.1mmol, 1.21 equiv), DIEA (28.0 mg, 0.2 mmol, 2.1 equiv). The resulting solution was stirred for 1 hr at room temperature. The mixture was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O:ACN=70:30 increasing to H₂O:ACN=10:90 within 20 min ; Detector, 254nm. This resulted in 16.5 mg (42.9%) of 3-cyano-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropyl-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (366) as a white solid. (300 MHz, CDCl₃, ppm): δ 8.35 (s, 1H), 7.25-7.20 (m, 2H), 6.97-6.92 (m, 1H), 6.86 (d, J = 8.1 Hz, 1H), 6.20-6.10 (m, 1H), 5.39-5.33 (m, 1H), 4.41-4.32 (m, 1H), 4.27-4.14 (m, 1H), 3.69-3.59 (m, 1H), 2.62 (s, 3H), 2.45-2.35 (m, 1H), 2.29-2.18 (m, 1H), 1.65-1.55 (m, 6H).

Preparation Example 14: Compounds 394, 397-0, 395 and 398 may be synthesized according to scheme 19 shown below:

15 Scheme 19




1. Synthesis of methyl 3-bromo-8-ethoxyimidazo[1,2-b]pyridazine-7-carboxylate (19-1)




Into a 40-mL round-bottom flask, was placed DMF (5.0 mL), ethyl 3-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (17-3, 150.0 mg, 0.5 mmol, 1.0 equiv), methyl iodide (200.0 mg, 1.3 mmol, 2.4 equiv), K_2CO_3 (210.0 mg, 1.5 mmol, 2.9 equiv). The resulting solution was stirred for 1 hr at 70°C . The mixture was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, $\text{H}_2\text{O}:\text{ACN}=90:10$ increasing to $\text{H}_2\text{O}:\text{ACN}=40:60$ within 15 min; Detector, 254nm. This resulted in 80 mg (48.6%) of methyl 3-bromo-8-ethoxyimidazo[1,2-b]pyridazine-7-carboxylate (19-1) as a white solid.

2. Synthesis of methyl 3-(3,5-dichlorophenyl)-8-ethoxyimidazo[1,2-b]pyridazine-7-carboxylate (19-2)




Into a 8-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed THF (5.0 mL), H_2O (1.0 mL), methyl 3-bromo-8-ethoxyimidazo[1,2-b]pyridazine-7-carboxylate (19-1, 80.0 mg, 0.25 mmol, 1.0 equiv), 3,5-dichlorophenylboronic acid (56.0 mg, 0.3 mmol, 1.1 equiv), $\text{Pd}(\text{dtbpf})\text{Cl}_2$ (30.0 mg, 0.05 mmol, 0.2 equiv), K_2CO_3 (100.0 mg, 0.7 mmol, 2.8 equiv). The resulting solution was stirred for 2 hr at 80°C . The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:3). This resulted in 20 mg (20.6%) of methyl 3-(3,5-dichlorophenyl)-8-ethoxyimidazo[1,2-b]pyridazine-7-carboxylate (19-2) as a white solid.

3. Synthesis of 3-(3,5-dichlorophenyl)-8-methoxyimidazo[1,2-b]pyridazine-7-carboxylic acid (19-3)



Into a 40-mL round-bottom flask, was placed EtOH (5.0 mL), H₂O (2.0 mL), ethyl 3-(3,5-dichlorophenyl)-8-methoxyimidazo[1,2-b]pyridazine-7-carboxylate (19-2, 70.0 mg, 0.2 mmol, 1.0 equiv), LiOH (40.0 mg, 1.7 mmol, 8.7 equiv). The resulting solution was stirred for 1 hr at 50 °C. The pH value of the solution was adjusted to 4 with HCl (4 mol/L). The resulting solution was extracted with 3x20 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 40 mg (61.9%) of 3-(3,5-dichlorophenyl)-8-methoxyimidazo[1,2-b]pyridazine-7-carboxylic acid (19-3) as a white solid.

4. Synthesis of 3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-methoxyimidazo[1,2-b]pyridazine-7-carboxamide (394)




Into a 8-mL round-bottom flask, was placed DMF (3.0 mL), 3-(3,5-dichlorophenyl)-8-methoxyimidazo[1,2-b]pyridazine-7-carboxylic acid 19-3, (40.0 mg, 0.1 mmol, 1.0 equiv), (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (19.5 mg, 0.1 mmol, 1.1 equiv), HATU (75.1 mg, 0.2 mmol, 1.7 equiv), DIEA (35.9 mg, 0.3 mmol, 2.3 equiv). The resulting solution was stirred for 1 hr at room temperature. The mixture was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O:ACN=70:30 increasing to

H₂O:ACN=10:90 within 20 min ; Detector, 254nm. This resulted in 12 mg (21.6%) of 3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-methoxyimidazo[1,2-b]pyridazine-7-carboxamide (394) as a white solid. (300 MHz, CDCl₃, ppm): δ 10.08 (d, J = 7.8 Hz, 1H), 9.10 (s, 1H), 7.92 (s, 2H), 7.51 (s, 1H), 7.37 (d, J = 7.8 Hz, 1H), 7.41-7.33 (m, 1H), 7.21-7.15 (m, 1H), 6.94-6.84 (m, 2H), 5.48-5.46 (m, 1H), 4.46 (s, 3H), 4.37-4.29 (m, 2H), 2.39-2.30 (m, 1H), 2.25-2.17 (m, 1H).


Compound	¹ H NMR Spectra
397-0	(300 MHz, DMSO-d ₆ , ppm) δ 10.1 (s, 1H), 9.15 (s, 1H), 7.95 (s, 2H), 7.65 - 7.60 (m, 2H), 7.35 - 7.40 (m, 1H) 7.20-7.15 (m, 1H), 6.99-6.85 (m, 2H), 6.20 - 6.00 (m, 1H), 5.70 - 5.30 (m, 3H), 4.36-4.25 (m, 2H), 3.20 (br, 1H), 2.45-2.20 (m, 2H)
395	(300 MHz, DMSO-d ₆ , ppm) δ 10.05 (d, J = 15 Hz, 1H), 9.25 (s, 1H), 8.95 (s, 1H), 8.25 (s, 2H), 7.85 - 7.80 (m, 1H), 7.35 - 7.15 (m, 1H) 7.00 - 6.80 (m, 2H), 5.25 - 5.18 (m, 1H), 4.36-4.15 (m, 2H), 2.25-2.02 (m, 2H)
398	(300 MHz, CDCl ₃ , ppm): δ 10.11 (d, J = 7.8 Hz, 1H), 9.12 (s, 1H), 7.93 (d, J = 1.9 Hz, 2H), 7.53 (s, 1H), 7.53-7.50 (m, 2H), 7.37 (d, J = 7.8 Hz, 1H), 7.21-7.16 (m, 1H), 6.94-6.85 (m, 2H), 5.48-5.44 (m, 1H), 4.97-4.88 (m, 2H), 4.37-4.30 (m, 2H), 2.37-2.32 (m, 1H), 2.24-2.19(m, 1H), 1.64 (t, J = 7.2 Hz, 3H)

Preparation Example 15: Compounds 450, 451, A408, A409, A421, A422, A460, A461, A462, A463 and A464 can be prepared adopting the process in schemes 20 and 21 shown below:


Scheme 20



Scheme 21




1. Synthesis of ethyl 4-bromo-1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-4)



- 5 Into a 500-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed ethyl 1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-3, 20.0 g, 78.4 mmol, 1.0 equiv), DMF (200.0 mL). This was followed by the addition of NBS (15.3 g, 86.2 mmol, 1.1 equiv) in several batches. The resulting solution was stirred for 1 overnight at room temperature. The reaction was then quenched by the addition of 1 L of water. The resulting


solution was extracted with 3x300 mL of ethyl acetate and the organic layers combined. The resulting mixture was washed with 2 x200 ml of brine. The mixture was dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:8). This resulted in 15 g (56.2%) of ethyl 4-bromo-1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-4) as a white solid.

2. Synthesis of ethyl 3-[4-bromo-1-[(tert-butoxycarbonyl)amino]imidazol-2-yl]-3-oxopropanoate (3-5)




Into a 1-L 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed ethyl 4-bromo-1-[(tert-butoxycarbonyl)amino]imidazole-2-carboxylate (3-4, 14 g, 41.9 mmol, 1.0 equiv), THF (140.0 mL, 1728.0 mmol, 41.2 equiv), ethyl acetate (18.5 g, 209.5 mmol, 5.0 equiv). This was followed by the addition of LiHMDS (209.5 mL, 209.5 mmol, 5.0 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred for 30 min at 0 °C. The reaction was then quenched by the addition of 200 mL of sat. NH₄Cl. The resulting solution was extracted with 3x200 mL of ethyl acetate and the organic layers combined. The resulting mixture was washed with 1 x200 ml of brine. The mixture was dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:3). This resulted in 11.5 g (65.7%) of ethyl 3-[4-bromo-1-[(tert-butoxycarbonyl)amino]imidazol-2-yl]-3-oxopropanoate (3-5) as a yellow solid.

3. Synthesis of ethyl 2-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (3-6)




Into a 250-mL 3-necked round-bottom flask, was placed ethyl 3-[4-bromo-1-[(tert-butoxycarbonyl)amino]imidazol-2-yl]-3-oxopropanoate (3-5, 10.0 g, 26.6 mmol, 1.0 equiv), DMF (100.0 mL), K₂CO₃ (3.7 g, 26.6 mmol, 1.0 equiv), DMF-DMA (7.9 g, 66.5 mmol, 2.5 equiv). The resulting solution was stirred for 1 hr at room temperature. The reaction was then quenched by the addition of 300 mL of water. The solids were collected by filtration. The crude product was re-crystallized from ACN in the ratio of 10 v. This resulted in 7.5 g (98.6%) of ethyl 2-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (3-6) as a white solid.

4. Synthesis of ethyl 2-bromo-8-methoxyimidazo[1,2-b]pyridazine-7-carboxylate (20-1)




Into a 250-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed ethyl 2-bromo-8-hydroxyimidazo[1,2-b]pyridazine-7-carboxylate (3-6, 8.0 g, 28.0 mmol, 1.0 equiv), DMF (80.0 mL), K₂CO₃ (7.7 g, 55.9 mmol, 2.0 equiv), CH₃I (11.9 g, 83.891 mmol, 3.0 equiv). The resulting solution was stirred for 2 hr at 60 °C. The residue was applied onto a C18 column with (25%~45% 8 min ACN in water). This resulted in 2.9 g (34.0%) of ethyl 2-bromo-8-methoxyimidazo[1,2-b]pyridazine-7-carboxylate (20-1) as a white solid.

5. Synthesis of ethyl 8-methoxy-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (20-2)




Into a 40-mL vial purged and maintained with an inert atmosphere of nitrogen, was placed ethyl 2-bromo-8-methoxyimidazo[1,2-b]pyridazine-7-carboxylate (20-1, 300.0 mg, 1.0 mmol, 1.0 equiv), DMF (6.0 mL), methyl 2,2-difluoro-2-sulfoacetate (960.2 mg, 5.0 mmol, 5.0 equiv), HMPA (895.7 mg, 5.0 mmol, 5.0 equiv), CuI (761.5 mg, 4.0 mmol, 4.0 equiv). The resulting solution was stirred for 16 hr at 100 °C. The solids were filtered out. The residue was applied onto a C18 column with (25%~45% 8 min ACN in H₂O (0.1% TFA)). This resulted in 170 mg (56.6%) of ethyl 8-methoxy-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (20-2) as a yellow solid.

6. Synthesis of ethyl 8-hydroxy-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (20-3)




Into a 40-mL vial purged and maintained with an inert atmosphere of nitrogen, was placed ethyl 8-methoxy-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (20-2, 1.8 g, 6.2 mmol, 1.0 equiv), LiCl (2.6 g, 62.2 mmol, 10.0 equiv), DMF (18.0 mL). The resulting solution was stirred for 2 hr at 120 degrees C. The residue was applied onto a C18 column with (75%~83% 8 min ACN in H₂O (0.1% TFA)). This resulted in 900 mg (52.5%) of ethyl 8-hydroxy-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (20-3) as a yellow solid.

7. Synthesis of ethyl 8-bromo-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (20-4)




Into a 40-mL vial purged and maintained with an inert atmosphere of nitrogen, was placed
 5 ethyl 8-hydroxy-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (20-3, 400.0 mg, 1.4 mmol, 1.0 equiv), POBr₃ (3333.7 mg, 11.6 mmol, 8.0 equiv). The resulting solution was stirred for 30 min at 120 °C. The reaction was then quenched by the addition of 10 mL of water/ice. The pH value of the solution was adjusted to 8 with Na₂CO₃. The resulting solution was extracted with 3x10 mL of ethyl acetate and the organic layers combined. The resulting mixture was
 10 washed with 1 x20 ml of brine. The mixture was dried over anhydrous sodium sulfate and concentrated. This resulted in 520 mg (97.3%) of ethyl 8-bromo-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (20-4) as a yellow solid.

8. Synthesis of ethyl 8-(morpholin-4-yl)-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (21-1)




15 Into a 8-mL vial,, was placed ethyl 8-bromo-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (20-4, 150.0 mg, 0.4 mmol, 1.0 equiv), DMF (1.5 mL), morpholine (193.3 mg, 2.2 mmol, 5.0 equiv). The resulting solution was stirred for 1 hr at 100 °C. The residue was applied onto a C18 column with (80%~90% 6 min ACN in H₂O (0.1% FA)). This resulted in 110 mg
 20 (72.0%) of ethyl 8-(morpholin-4-yl)-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (21-1) as a white solid.

9. Synthesis of ethyl 3-(3,5-dichlorophenyl)-8-(morpholin-4-yl)-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (21-2)



Into a 8-mL vial purged and maintained with an inert atmosphere of nitrogen, was placed
 5 ethyl 8-(morpholin-4-yl)-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (21-1, 90.00mg, 0.3 mmol, 1.0 equiv), NMP (1.8 mL), 1,3-dichloro-5-iodobenzene (214.0 mg, 0.8 mmol, 3.0 equiv), Cs_2CO_3 (170.3 mg, 0.5 mmol, 2.0 equiv), $\text{Pd}(\text{AcO})_2$ (11.7 mg, 0.05 mmol, 0.2 equiv), $\text{Ad}_2(\text{n-Bu})\text{P}$ (28.0 mg, 0.08 mmol, 0.3 equiv). The resulting solution was stirred overnight at 100°C . The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:5).
 10 This resulted in 110 mg (72.2%) of ethyl 3-(3,5-dichlorophenyl)-8-(morpholin-4-yl)-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (21-2) as a white solid.


10. Synthesis of 3-(3,5-dichlorophenyl)-8-(morpholin-4-yl)-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylic acid (21-3)



15 Into a 8-mL vial, was placed ethyl 3-(3,5-dichlorophenyl)-8-(morpholin-4-yl)-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylate (21-2, 110.0 mg, 0.2 mmol, 1.0 equiv), $i\text{-PrOH}$ (2.1 mL), H_2O (0.7 mL), $\text{LiOH} \cdot \text{H}_2\text{O}$ (75.4 mg, 1.8 mmol, 8.0 equiv). The resulting

solution was stirred for 5 hr at 60 degrees C. The residue was applied onto a C18 column with (80%~90% 6 min ACN in H₂O (0.1% FA)). This resulted in 70 mg (65.3%) of 3-(3,5-dichlorophenyl)-8-(morpholin-4-yl)-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylic acid (21-3) as a white solid.

5 **11. Synthesis of 3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-(morpholin-4-yl)-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxamide (450)**



Into a 8-mL vial, was placed 3-(3,5-dichlorophenyl)-8-(morpholin-4-yl)-2-(trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxylic acid (21-3, 66.0 mg, 0.1 mmol, 1.0 equiv),
 10 DMF (2.0 mL), (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (32.0 mg, 0.2 mmol, 1.5 equiv),
 DIEA (55.5 mg, 0.4 mmol, 3.0 equiv), HATU (81.6 mg, 0.2 mmol, 1.5 equiv). The resulting
 solution was stirred for 1 hr at room temperature. The residue was applied onto a C18 column
 with (90%~98% 6 min ACN in H₂O (0.1% FA)). This resulted in 35.1 mg (41.4%) of 3-(3,5-
 15 dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-(morpholin-4-yl)-2-
 (trifluoromethyl)imidazo[1,2-b]pyridazine-7-carboxamide (450) as a white solid. (300 MHz,
 DMSO-d₆, ppm): δ 9.08 (d, J = 8.0 Hz, 1H), 8.36 (s, 1H), 7.83 (t, J = 1.9 Hz, 1H), 7.66 (d, J = 1.9
 Hz, 2H), 7.32 (d, J = 7.8 Hz, 1H), 7.21-7.15 (m, 1H), 6.94-6.89 (m, 1H), 6.82-6.79 (m, 1H), 5.23-
 5.16 (m, 1H), 4.30-4.20 (m, 2H), 3.94-3.81 (m, 8H), 2.27-2.16 (m, 1H), 2.08-2.01 (m, 1H).


Compound	¹ H NMR Spectra
451	(300 MHz, DMSO-d ₆ , ppm): δ 9.01 (d, J = 8.1 Hz, 1H), 8.23 (s, 1H), 7.82 (t, J = 1.9 Hz, 1H), 7.64 (d, J = 1.9 Hz, 2H), 7.31-7.28 (m, 1H), 7.20-7.14 (m, 1H), 6.93-6.88 (m, 1H), 6.80 (d, J = 8.1, Hz, 1H), 5.20-5.14 (m, 1H), 4.27-4.21 (m, 2H), 3.43 (s, 6H), 2.27-2.13 (m, 1H), 2.14-1.98 (m, 1H)

A408	(400 MHz, DMSO-d ₆) δ ppm 8.99 (d, J=8.11 Hz, 1 H) 8.20 (s, 1 H) 7.73 (d, J=1.90 Hz, 2 H) 7.62 - 7.69 (m, 1 H) 7.32 (d, J=6.97 Hz, 1 H) 7.12 - 7.21 (m, 1 H) 6.91 (td, J=7.45, 0.95 Hz, 1 H) 6.80 (d, J=8.24 Hz, 1 H) 5.15 - 5.24 (m, 1 H) 4.18 - 4.31 (m, 2 H) 3.80 - 3.94 (m, 4 H) 3.71 - 3.80 (m, 4 H) 2.47 (s, 3 H) 2.11 - 2.27 (m, 1 H) 1.98 - 2.11 (m, 1 H)
A409	(400 MHz, DMSO-d ₆) δ ppm 8.93 (d, J=8.11 Hz, 1 H) 8.12 (s, 1 H) 7.73 (d, J=1.90 Hz, 2 H) 7.64 (t, J=1.90 Hz, 1 H) 7.31 (d, J=7.10 Hz, 1 H) 7.16 (t, J=7.40 Hz, 1 H) 6.91 (td, J=7.48, 1.01 Hz, 1 H) 6.79 (dd, J=8.24, 0.76 Hz, 1 H) 5.14 - 5.21 (m, 1 H) 4.16 - 4.31 (m, 2 H) 3.30 (s, 6 H) 2.47 (s, 3 H) 2.09 - 2.25 (m, 1 H) 1.96 - 2.09 (m, 1 H)
A421	(400 MHz, DMSO-d ₆ , ppm) δ 8.99 (d, J = 7.98 Hz, 1H), 8.06 (s, 1H), 7.74 (d, J = 8.62 Hz, 2H), 7.30 (d, J = 7.60 Hz, 1H), 7.15 (t, J = 7.30 Hz, 1H), 6.89 (t, J = 7.41 Hz, 1H), 6.78 (d, J = 8.11 Hz, 1H), 5.18 (br d, J = 7.73 Hz, 1H), 4.19 - 4.29 (m, 2H), 3.69 - 3.93 (m, 8H), 2.52 - 2.55 (m, 1H), 2.22 (s, 3H), 1.20 - 1.24 (m, 1H)
A422	(400 MHz, DMSO-d ₆ , ppm) δ 8.99 (d, J = 8.11 Hz, 1H), 8.14 (s, 1H), 7.65 - 7.73 (m, 1H), 7.29 - 7.39 (m, 2H), 7.17 (t, J = 7.60 Hz, 1H), 6.91 (t, J = 7.10 Hz, 1H), 6.79 (d, J = 8.11 Hz, 1H), 5.16 - 5.22 (m, 1H), 4.20 - 4.30 (m, 2H), 3.72 - 3.92 (m, 8H), 2.16 - 2.38 (m, 4H), 1.99 - 2.08 (m, 1H)
A460	(400 MHz, DMSO-d ₆ , ppm) δ 8.93 (d, 1H), 8.06 (s, 1H), 7.67 (m, 1H), 7.36 (m, 1H), 7.30 (d, 1H), 7.16 (t, 1H), 6.90 (t, 1H), 6.79 (d, 1H), 5.17 (q, 1H), 4.24 (m, 2H), 3.41 (s, 6H), 2.35 (s, 3H), 1.97-2.20 (m, 2H)
A461	(400 MHz, DMSO-d ₆ , ppm) δ 8.92 (d, 1H), 8.02 (s, 1H), 8.00 (d, 1H), 7.64 (m, 1H), 7.28 (d, 1H), 7.16 (t, 1H), 6.89 (t, 1H), 6.79 (d, 1H), 5.17 (q, 1H), 4.24 (m, 2H), 3.41 (s, 6H), 2.27 (s, 3H), 1.96-2.19 (m, 2H)
A462	400 MHz, DMSO-d ₆) δ ppm 8.91 (d, J=8.1 Hz, 1 H), 8.02 (s, 1 H), 7.27 - 7.44 (m, 3 H), 7.16 (t, J=7.4 Hz, 1 H), 6.89 (td, J=7.5, 1.1 Hz, 1 H), 6.79 (dd, J=8.2, 1.1 Hz, 1 H), 5.14 - 5.21 (m, 1 H), 3.49 (s, 6 H), 2.52 - 2.52 (m, 2 H), 2.21 (s, 3 H), 2.10 - 2.21 (m, 1 H), 1.95 - 2.05 (m, 1 H)


A463	(400 MHz, DMSO-d ₆) δ ppm 8.92 (dd, J=8.1, 2.3 Hz, 1 H), 8.01 (s, 1 H), 7.86 (dd, J=8.3, 3.0 Hz, 1 H), 7.50 (ddd, J=8.6, 7.2, 3.0 Hz, 1 H), 7.28 (d, J=7.7 Hz, 1 H), 7.16 (t, J=7.4 Hz, 1 H), 6.89 (t, J=7.5 Hz, 1 H), 6.79 (d, J=8.3 Hz, 1 H), 5.13 - 5.20 (m, 1 H), 3.52 (s, 6 H), 2.52 - 2.56 (m, 2 H), 2.28 (s, 3 H), 2.10 - 2.21 (m, 1 H), 1.95 - 2.05 (m, 1 H)
A464	(400 MHz, DMSO-d ₆) δ ppm 8.95 (d, J=8.1 Hz, 1 H), 8.00 (s, 1 H), 7.30 - 7.41 (m, 3 H), 7.00 - 7.30 (m, 2H), 6.91 (td, J=7.6, 1.1 Hz, 1 H), 6.75 (dd, J=8.1, 1.1 Hz, 1 H), 5.15 - 5.20 (m, 1 H), 4.20 - 4.25 (m, 2H), 3.51 (s, 6 H), 2.21 (s, 3 H), 2.11 - 2.21 (m, 1 H), 1.95 - 2.05 (m, 1 H)

Preparation Example 16: Compounds 511 and 512 can be prepared adopting the process in scheme 22 below:

Scheme 22



1. **Synthesis of ethyl 3-(3,5-dichlorophenyl)-8-(morpholin-4-yl)imidazo[1,2-b]pyridazine-7-carboxylate (22-1)**




5

Into a 40-mL round-bottom flask, was placed NMP (5.0 mL), ethyl 8-chloro-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (17-5, 100.0 mg, 0.2 mmol, 1.0 equiv), morpholine (0.5 mL, 5.7 mmol, 23.6 equiv). The resulting solution was stirred for 1 hr at 80 °C. The reaction was then quenched by the addition of 10 mL of water. The resulting solution was extracted with 3x20 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 65 mg (64.0%) of ethyl 3-(3,5-

10

dichlorophenyl)-8-(morpholin-4-yl)imidazo[1,2-b]pyridazine-7-carboxylate (22-1) as a yellow solid.

2. Synthesis of 3-(3,5-dichlorophenyl)-8-(morpholin-4-yl)imidazo[1,2-b]pyridazine-7-carboxylic acid (22-2)




5

Into a 40-mL round-bottom flask, was placed EtOH (2.0 mL), H₂O (1.5 mL), ethyl 3-(3,5-dichlorophenyl)-8-(morpholin-4-yl)imidazo[1,2-b]pyridazine-7-carboxylate (22-1, 65.0 mg, 0.1 mmol, 1.0 equiv), LiOH.H₂O (50.0 mg, 1.2 mmol, 7.7 equiv). The resulting solution was stirred for 3 hr at 50 °C. HCl (6 mol/L) was employed to adjust the pH to 4. The resulting solution was extracted with 3x20 mL of ethyl acetate and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 40 mg (65.9%) of 3-(3,5-dichlorophenyl)-8-(morpholin-4-yl)imidazo[1,2-b]pyridazine-7-carboxylic acid (22-2) as a yellow solid.

10

3. Synthesis of 3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-(morpholin-4-yl)imidazo[1,2-b]pyridazine-7-carboxamide (511)

15




Into a 40-mL round-bottom flask, was placed DMF (5.0 mL), 3-(3,5-dichlorophenyl)-8-(morpholin-4-yl)imidazo[1,2-b]pyridazine-7-carboxylic acid (22-2, 40.0 mg, 0.1 mmol, 1.0 equiv), (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (17.0 mg, 0.1 mmol, 1.12equiv), HATU (65.0 mg,

0.2 mmol, 1.7 equiv), DIEA (38.0 mg, 0.3 mmol, 2.9 equiv). The resulting solution was stirred for 1 hr at room temperature. The crude product was purified by Flash-Prep-HPLC with the following conditions (IntelFlash-1): Column, C18 silica gel; mobile phase, H₂O:ACN=50:50 increasing to H₂O:ACN=10:90 within 20 min; Detector, 254nm. product was obtained. This resulted in 37.8 mg (70.9%) of 3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-(morpholin-4-yl)imidazo[1,2-b]pyridazine-7-carboxamide (511) as a white solid. (300 MHz, CDCl₃, ppm) δ 9.03 (d, J = 8.1 Hz, 1H), 8.37 (s, 1H), 8.31 (s, 1H), 8.27 (d, J = 1.8 Hz, 1H), 7.60 (t, J = 1.8 Hz, 1H), 7.34 (d, J = 1.8 Hz, 1H), 7.21-7.15 (m, 1H), 6.95-6.90 (m, 1H), 6.80-6.80 (m, 1H), 5.23-5.20 (m, 1H), 4.27-4.24(m, 2H), 3.93-3.78 (m, 8H), 2.23-2.21 (m, 1H), 2.10-2.07 (m, 1H).


¹H NMR spectra for 512: (300 MHz, CDCl₃, ppm) δ 8.36 (s, 1H), 7.98 (d, J = 1.9 Hz, 2H), 7.90 (s, 1H), 7.37-7.35 (m, 1H), 7.32-7.22 (m, 2H), 7.03-6.86 (m, 2H), 6.67 (d, J = 7.6 Hz, 1H), 5.42-5.34 (m, 1H), 4.43-4.31 (m, 1H), 4.29-4.16 (m, 1H), 3.46 (s, 6H), 2.43-2.36 (m, 1H), 2.27-2.14 (m, 1H)

Preparation Example 17: Compound 558 was prepared according to the scheme 23 shown below. Compound 559 can be prepared by adopting the process described in scheme 23 by someone skilled in the art.

Scheme 23




1. Synthesis of ethyl 7-chloro-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-2-carboxylate (23-2)



To a stirred mixture of ethyl 7-chloroimidazo[1,2-b]pyridazine-2-carboxylate (23-1, 1.0 g, 4.4 mmol, 1.0 equiv) and 1,3-dichloro-5-iodobenzene (1.8 g, 6.6 mmol, 1.5 equiv) in toluene (10 mL, 105.0 mmol, 23.7 equiv) were added K_2CO_3 (1.2 g, 8.9 mmol, 2.0 equiv) and $Pd(OAc)_2$ (0.1 g, 0.4 mmol, 0.1 equiv) and PPh_3 (0.2 g, 0.9 mmol, 0.2 equiv) at room temperature under nitrogen atmosphere. The resulting mixture was stirred for overnight at $100^\circ C$ under nitrogen atmosphere. The mixture was allowed to cool down to room temperature. The resulting mixture was concentrated under vacuum. The residue was purified by silica gel column chromatography, eluted with PE / EA (5:1) to afford ethyl 7-chloro-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-2-carboxylate (23-2, 1.2 g, 73.0%) as a yellow oil.

2. Synthesis of 7-chloro-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-2-carboxylic acid (23-3)




15

To a stirred mixture of ethyl 7-chloro-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-2-carboxylate (23-2, 1.0 g, 2.7 mmol, 1.0 equiv) in EtOH (8 mL) and H_2O (4 mL) was added $LiOH \cdot H_2O$ (1.13 g, 27.0 mmol, 10.0 equiv) at room temperature. The resulting mixture was stirred for 2 h at room temperature. The resulting mixture was concentrated under vacuum. The resulting mixture was diluted with water (50 mL). The mixture was acidified to pH 4 with HCl

20


(aq.). The precipitated solids were collected by filtration and washed with water (2x10 mL). The resulting solid was dried under infrared light.

3. Synthesis of 7-chloro-3-(3,5-dichlorophenyl)-N-methoxy-N-methylimidazo[1,2-b]pyridazine-2-carboxamide (23-4)



To a stirred mixture of 7-chloro-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-2-carboxylic acid (23-3, 1.0 g, 3.0 mmol, 1.0 equiv) and N,O-dimethylhydroxylamine (0.3 g, 4.4 mmol, 1.5 equiv) in DCM (10 mL, 157.3 mmol, 53.9 equiv) were added EDCI (0.6 g, 2.9 mmol, 1.0 equiv) and PyBOP (1.5 g, 2.9 mmol, 1.0 equiv) and DIEA (1.1 g, 8.7 mmol, 3.0 equiv) at room temperature. The resulting mixture was stirred for overnight at room temperature. The resulting mixture was concentrated under vacuum. The residue was purified by reverse flash chromatography with the following conditions: column, silica gel; mobile phase, MeCN in water, 50% to 90% gradient in 10 min; detector, UV 254 nm to afford 400 mg of 7-chloro-3-(3,5-dichlorophenyl)-N-methoxy-N-methylimidazo[1,2-b]pyridazine-2-carboxamide (23-4, 35.5%) as a yellow oil.


4. Synthesis of afford 1-[7-chloro-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazin-2-yl]ethenone (23-5)



To a stirred mixture of 7-chloro-3-(3,5-dichlorophenyl)-N-methoxy-N-methylimidazo[1,2-b]pyridazine-2-carboxamide (23-4, 80.0 mg, 0.2 mmol, 1.0 equiv) in THF (2 mL, 24.7 mmol, 119.0 equiv) was added chloromethylmagnesium (77.6 mg, 1.0 mmol, 5.0 equiv) at -78 °C under


nitrogen atmosphere. The resulting mixture was stirred for 60 min at 50 °C under nitrogen atmosphere. The reaction was quenched with sat. NH_4Cl (aq.) at room temperature. The resulting mixture was extracted with EtOAc (2 x 20 mL). The combined organic layers were washed with brine (2x4 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by Prep-TLC (PE / EA 4:1) to afford 1-[7-chloro-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazin-2-yl]ethanone (23-5, 40 mg, 56.6%) as a light yellow solid.

5. Synthesis of methyl 2-acetyl-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (23-6)




To a solution of 1-[7-chloro-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazin-2-yl]ethanone (23-5, 40.0 mg, 0.1 mmol, 1.0 equiv) in 1 mL MeOH and dioxane (2 mL) was added KOAc (34.6 mg, 0.4 mmol, 3.0 equiv) and $\text{Pd}(\text{dppf})\text{Cl}_2\text{CH}_2\text{Cl}_2$ (9.6 mg, 0.01 mmol, 0.1 equiv) in a pressure tank. The mixture was purged with nitrogen for 2 min and then was pressurized to 30 atm with carbon monoxide at 130 degrees C for overnight. The reaction mixture was cooled to room temperature and filtered to remove insoluble solids. The residue was purified by Prep-TLC (PE / EA 3:1) to afford 20 mg of methyl 2-acetyl-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (23-6, 46.8%) as a yellow solid.

6. Synthesis of afford methyl 2-acetyl-3-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylate (23-7)




To a stirred mixture of methyl 2-acetyl-3-(3,5-dichlorophenyl)imidazo[1,2-b]pyridazine-7-carboxylate (23-6, 20.0 mg, 0.05 mmol, 1.0 equiv) and 2-[(propane-2-sulfonyl)zinciosulfonyl]propane (46.1 mg, 0.2 mmol, 3.0 equiv) in DMSO (2 mL) was added t-BuOOH (28.6 mg, 0.3 mmol, 5.0 equiv) at room temperature. The resulting mixture was stirred for 2 h at 50 °C. The mixture was allowed to cool down to room temperature. The resulting mixture was diluted with water (10 mL). The resulting mixture was extracted with EtOAc (2 x 10 mL). The combined organic layers were washed with brine (2x5 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by Prep-TLC (PE / EA 2:1) to afford methyl 2-acetyl-3-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylate (23-7, 10 mg, 44.8%) as a yellow oil.

7. Synthesis of 2-acetyl-3-(3,5-dichlorophenyl)-8-(propan-2-yl)imidazo[1,2-b]pyridazine-7-carboxylic acid (23-8)



To a stirred mixture of methyl 2-acetyl-3-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylate (23-7, 10.0 mg, 0.02 mmol, 1.0 equiv) in EtOH (1 mL) and H₂O (0.5 mL) was added LiOH.H₂O (10.3 mg, 0.25 mmol, 10.0 equiv) at room temperature. The resulting mixture was stirred for 1 h at room temperature. The resulting mixture was concentrated under vacuum. The resulting mixture was diluted with water (5 mL). The mixture was acidified to pH 4 with HCl (aq.). The resulting mixture was extracted with EtOAc (2 x 5 mL). The combined organic layers were washed with brine (1x5 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The crude product (23-8) was used in the next step directly without further purification.

8. Synthesis of 2-acetyl-3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropylimidazo[1,2-b]pyridazine (558)




To a stirred mixture of 2-acetyl-3-(3,5-dichlorophenyl)-8-isopropylimidazo[1,2-b]pyridazine-7-carboxylic acid (23-8, 10.0 mg, 0.02 mmol, 1.0 equiv) and (4S)-3,4-dihydro-2H-1-benzopyran-4-amine (5.7 mg, 0.04 mmol, 1.5 equiv) in DMF (1 mL, 12.9 mmol, 506.8 equiv) were added DIEA (9.9 mg, 0.07 mmol, 3.0 equiv) and HATU (14.5 mg, 0.04 mmol, 1.5 equiv) at room temperature. The resulting mixture was stirred for 1 h at room temperature. The crude product was purified by Prep-HPLC with the following conditions (Gradient: isocratic) to afford 11 mg of 2-acetyl-3-(3,5-dichlorophenyl)-N-[(4S)-3,4-dihydro-2H-1-benzopyran-4-yl]-8-isopropylimidazo[1,2-b]pyridazine (558, 95.0%) as a solid. (400 MHz, Chloroform-d, ppm) δ : 8.31 (s, 1H), 7.56 (s, 2H), 7.45 (s, 1H), 7.225-7.21 (m, 1H), 6.97-6.93 (m, 1H), 6.88-6.86 (m, 1H), 6.06-6.05 (m, 1H), 5.38-5.36 (m, J = 4.5 Hz, 1H), 4.38-4.34 (m, 1H), 4.21-4.16 (m, 1H), 3.73-3.67 (m, 1H), 2.75 (s, 3H), 2.43-2.38 (m, 1H), 2.23-2.20 (m, 1H), 1.68 (t, J = 7.4 Hz, 6H).


¹H NMR spectra for 559: (400 MHz, Chloroform-d, ppm) δ : 8.44 (s, 1H), 7.88 (s, 2H), 7.50 (s, 1H), 7.26-7.21 (m, 2H), 6.98-6.94 (m, 1H), 6.90 (d, J = 8.3 Hz, 1H), 6.08-6.07 (m, 1H), 5.40-5.36 (m, 1H), 4.39-4.34 (m, 1H), 4.21-4.15 (m, 1H), 3.71-3.64 (m, 1H), 2.44-2.39 (m, 1H), 2.26-2.20 (m, 1H), 1.65-1.54 (m, 6H)

Preparation Example 18: Compound 614 can be prepared according to the process described in Scheme 24.

Scheme 24




1. Synthesis of 4-nitroso-3,4-dihydro-2H-1,4-benzoxazine




- 5 Into a 500mL round-bottom flask, were added methyl 3,4-dihydro-2H-1,4-benzoxazine (24-1, 3.0 g, 22.2 mmol, 1.0 equiv) and 3M/HCl (300 ml) and NaNO₂ (1.8 g, 26.1 mmol, 1.2 equiv) was added dropwise/ in portions at 2 hr at 0 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (10:01) to afford 4-nitroso-3,4-dihydro-2H-1,4-benzoxazine (24-2, 2.7 g, 75.1%) as a yellow oil.
- 10

2. Synthesis of 3,4-dihydro-2H-1,4-benzoxazin-4-amine



Into a 500mL round-bottom flask, were added methyl 4-nitroso-2,3-dihydro-1,4-benzoxazine (24-2, 2.7 g, 19.2 mmol, 1.0 equiv) and THF (300 ml) and LAH (1.35 g, 1.9 mmol, 2.0 equiv) was added dropwise/ in portions at 2 hr at 0 °C under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE / EA (10:01) to afford 3,4-dihydro-2H-1,4-benzoxazin-4-amine (24-3, 2.2 g, 44.9%) as a yellow oil.

3. Synthesis of 8-tert-butyl-3-(3,5-dichlorophenyl)-N-(2,3-dihydro-1,4-benzoxazin-4-yl)-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (614)



To a stirred mixture of 8-tert-butyl-3-(3,5-dichlorophenyl)-2-methylimidazo[1,2-b]pyridazine-7-carboxylic acid (12-5, 100.0 mg, 0.3 mmol, 1.0 equiv) and 2,3-dihydro-1,4-benzoxazin-4-amine (119.1 mg, 0.8 mmol, 3.0 equiv) in DMF (5 mL) were added DIEA (170.8 mg, 1.3 mmol, 5.0 equiv) and HATU (301.6 mg, 0.8 mmol, 3.0 equiv) at room temperature. The resulting mixture was stirred for overnight at 80 degrees C. The mixture was allowed to cool down to room temperature. The residue was purified by reverse flash chromatography with the following conditions: column, C18 silica gel; mobile phase, ACN in water, 40% to 95% gradient in 10 min;

detector, UV 254 nm to afford 8-tert-butyl-3-(3,5-dichlorophenyl)-N-(2,3-dihydro-1,4-benzoxazin-4-yl)-2-methylimidazo[1,2-b]pyridazine-7-carboxamide (614, 43.8 mg, 32.5%) as an off-white solid. (300 MHz, DMSO-d₆, ppm) δ 10.44 (s, 1H), 8.39 (s, 1H), 7.76 (d, J = 1.8 Hz, 2H), 7.61 (t, J = 2.1 Hz, 1H), 7.05-6.96 (m, 1H), 6.88-6.82 (m, 1H), 6.79-6.66 (m, 2H), 4.36-4.33 (m, 2H), 3.65-3.62 (m, 2H), 2.55 (s, 3H), 1.73 (s, 9H).

Biological Examples

The disclosure is further illustrated by the following biological examples, which are not to be construed as limiting this disclosure in scope or spirit to the specific procedures herein described. It is to be understood that the examples are provided to illustrate certain embodiments and that no limitation to the scope of the disclosure is intended thereby. It is to be further understood that resort may be had to various other embodiments, modifications, and equivalents thereof which may suggest themselves to those skilled in the art without departing from the spirit of the present disclosure and/or scope of the appended claims.

Biological Example 1: Screening method to test activity of compounds against microfilaria of *Dirofilaria immitis*.

Four hundred to six hundred microfilariae of *Dirofilaria immitis* were added to wells of a microtiter plate containing RPMI media and the test compound formulated in 100% DMSO. Plates were held for three days at 37°C and 5% CO₂. The efficacy of a compound was determined based on the motility of the microfilaria as compared to average motility of control wells containing DMSO only. A dose response assay was conducted to determine an EC₅₀ value. Compounds 298-0, 304, 295, 296, A412, A406, A405, A411, A400, A401, 325, A419, 513-0, 450, A435, A439 and A442 exhibited EC₅₀ values of between 0.1 μ M and 1 μ M. Compounds 279, 273, 276, 294, 322, 323, 326-0, 323-0, 352, 364, 371, 373, 298, 419, A403, A413, A407, A414, A449, A448, A447, A441, 512, 511, 513, 418, A428, A427, 305, 451 and 558 exhibited EC₅₀ values between 0.01 μ M and 0.1 μ M. Compounds 308, 271, 274, 306, 297, A410, 277, 299-0, 293, 275, 175, 573, 614, 572, 528, 560, 420, A422, 523 and 527 exhibited EC₅₀ values of between 0.001 μ M and 0.01 μ M; and Compounds 307, 324, 345, 524, 526 and A421 exhibited EC₅₀ values of less than 0.001 μ M.

Biological Example 2: Screening method to test activity of compounds against *Haemonchus contortus*.

Twenty L1 *Haemonchus contortus* larvae were added to wells of a microtitre plate containing a nutrient medium and the test compound in DMSO. An analysis was conducted at 4 days to determine the extent of development of the larvae from L1 to L3. Larvae exposed to DMSO alone served as controls. A dose response assay was conducted to determine an EC₅₀ value. Compounds 174, 366, 320-0, 369, 365, 321, 394, 298-0, 323-0, 323, 325, 296, 304-0, 373, 398, A404, 370, 326 and 299 were found to be active with EC₅₀ values of between 1 µM and 10 µM. Compounds 304, A405, A414, A403, A401, 371, 364, 352, 308, 320, 298, 299-0, 327, 324, 279 and 275 exhibited EC₅₀ values of between 0.1 µM and 1 µM. Compounds 175, 272, 273, 295, 326-0, 277, 294, 344, A408, A410, A412, A409, A413 and A400 exhibited EC₅₀ values of between 0.01 µM and 0.1 µM; and compounds 297, 306, 271, 345 and 274 exhibited EC₅₀ values of less than 0.01 µM.

Biological Example 3: Screening method to test activity of compounds against L4 stage larvae of *Dirofilaria immitis*.


Four to six L4 stage *Dirofilaria immitis* worms are added to wells of a microtiter plate containing maintenance nutrient media and the test compound formulated in 100% DMSO. Plates are held at 37°C and 5% CO₂ for three days and then assessed to determine the motility of the larvae. Efficacy of a compound is determined by comparison of the treated L4 motility of the relative to the average motility of worms in control wells containing DMSO only. A dose response assay is conducted to determine an EC₅₀ value. Compounds 366, 320-0, 365, 321, 304-0 and 329 exhibited EC₅₀ values of between 1 µM and 10 µM, compounds 325, 323 and 370 were found to have EC₅₀ values of between 0.1 µM and 1 µM. Compounds 276, 322, 320, 364, 326, 298-0, 299, 323-0, 296, 295 and 307 exhibited EC₅₀ values of from 0.01 µM to 0.1 µM. Compounds 304, 175, 272, 294, 273, 344, 275, 327, 326-0, 299-0, 298, 277, 324 and 297 exhibited EC₅₀ values of between 0.001 µM to 0.01 µM. Compounds 308, 271, 293, 345, 371 and 306 exhibited an EC₅₀ of < 0.001 µM.

* * *

Having thus described in detail preferred embodiments of the present invention, it is to be understood that the invention defined by the above paragraphs is not to be limited to particular details set forth in the above description as many apparent variations thereof are possible without departing from the spirit or scope of the present invention.

WHAT IS CLAIMED IS:










1. A compound of Formula I:









(I)

wherein:

- L is L1, L2, L3, L4, L5, L6, L7, L8, L9, L10, L11, L12, L13, L14 or L15:





R' is hydrogen, optionally substituted alkyl, optionally substituted cycloalkyl, or optionally substituted aryl;

R¹ is hydrogen, cyano, halo, hydroxyl, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, optionally substituted alkoxyalkyl, optionally substituted aminoalkyl, optionally substituted alkylaminoalkyl, optionally substituted dialkylaminoalkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heteroaryl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted cycloalkoxy, optionally substituted heterocyclyl, optionally substituted alkylcarbonyl, optionally substituted alkoxy carbonyl, aminocarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, -SO_p(optionally substituted alkyl or haloalkyl), -SF₅, or -NR^aR^b, wherein R^a and R^b are independently H or optionally substituted alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

R² is hydrogen, cyano, halo, hydroxyl, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl; optionally substituted aryloxy, optionally substituted heteroaryl,

optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted cycloalkoxy, optionally substituted heterocyclyl, optionally substituted alkylcarbonyl, optionally substituted alkoxy carbonyl, optionally substituted aminocarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, $-\text{SO}_p(\text{optionally substituted alkyl or haloalkyl})$, $-\text{SF}_5$, or $-\text{NR}^a\text{R}^b$, wherein R^a and R^b are independently H or optionally substituted alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

R^3 is hydrogen, optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted alkylcarbonyl, optionally substituted alkoxy carbonyl, aminocarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, $-\text{S}(\text{O})_p(\text{optionally substituted alkyl})$, $-\text{SF}_5$, optionally substituted heterocyclyl, optionally substituted 6- to 10-membered aryl, optionally substituted 5- to 10-membered heteroaryl, a spirocyclic heterocyclyl-carbocyclyl group, a spirocyclic heterocyclyl-heterocyclyl group, a spirocyclic carbocyclyl-carbocyclyl group, a spirocyclic carbocyclyl-heterocyclyl group or $-\text{NR}^a\text{R}^b$, wherein R^a and R^b are independently H or optionally substituted alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

R^4 and $\text{R}^{4'}$ are independently in each occurrence, hydrogen, halogen, cyano, nitro, hydroxyl, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted cycloalkyl, optionally substituted cycloalkoxy, optionally substituted alkylcarbonyl, optionally substituted alkoxy carbonyl, optionally substituted aminocarbonyl, alkylaminocarbonyl, or di(alkyl)aminocarbonyl, optionally substituted alkylcarbonyloxy, optionally substituted alkylcarbonylamino, optionally substituted aryl, optionally substituted heteroaryl, $-\text{SF}_5$, $-\text{SO}_p(\text{optionally substituted alkyl or haloalkyl})$; or R^4 together with $\text{R}^{4'}$ together form a 2-6-membered chain optionally containing one or two heteroatoms selected from the group consisting of N, O, Si and S to form carbocyclic or heterocyclic

ring together with the carbon atom to which they are attached; or $-NR^cR^d$, wherein R^c and R^d are independently H or optionally substituted alkyl; or R^c and R^d may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

R^8 is hydrogen, halogen, alkyl, haloalkyl, cycloalkyl, alkenyl or alkynyl;

R^9 and $R^{9'}$ are independently hydrogen, halo, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or cycloalkoxy, or R^9 together with $R^{9'}$ form a 2-6-membered chain optionally containing one or two heteroatoms selected from the group consisting of N, O, Si and S to form carbocyclic or heterocyclic ring together with the carbon atom to which they are attached;

Q is N or C- R^8 ;

X is O, S or N- R^8 ;

Y^1 and Y^6 are each independently N, C, or $-CR^4-$;

Y^2 , Y^3 , Y^4 and Y^5 are each independently N, NR^8 , S, O, $-CR^4-$ or CR^4R^4 ;

W is CR^5R^6 , O, SO_p , or N- R^7 ,

Z is CR^5R^6 , O, SO_p , or N- R^7 ,

wherein

R^5 and R^6 are independently in each occurrence hydrogen, halo, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or cycloalkoxy; or R^5 together with R^6 form a 2-6-membered chain optionally containing one or two heteroatoms selected from the group consisting of N, O, Si and S to form carbocyclic or heterocyclic ring together with the carbon atom to which they are attached;

R^7 is hydrogen or C₁-C₄-alkyl; and

wherein at most three of Y¹, Y², Y³, Y⁴, Y⁵ and Y⁶ are heteroatoms;

a is 0 or 1;

q is 0 or 1;

p is independently in each occurrence is 0, 1, or 2; and

the dashed bonds (---) signifies a single or double bond;

or a pharmaceutically acceptable salt thereof.

2. The compound of Formula (I) of claim 1,

wherein:

- R¹ is hydrogen, cyano, halo, hydroxyl, C₁-C₆-alkyl, C₁-C₆-haloalkyl, hydroxy-C₁-C₆-alkyl, hydroxy-C₁-C₆-haloalkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-haloalkoxy-C₁-C₆-alkyl, amino-C₁-C₆-alkyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-haloalkenyloxy, C₂-C₆-alkynyloxy, C₂-C₆-haloalkynyloxy, C₁-C₆-haloalkoxy, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-haloalkylaminocarbonyl, di-C₁-C₆-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heteroaryl, optionally substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkenyl, optionally substituted C₃-C₈-cycloalkyloxy, optionally substituted 3- to 7-membered heterocyclyl, -SF₅, -SO_p(optionally substituted C₁-C₆-alkyl or C₁-C₆-haloalkyl), or -NR^aR^b wherein R^a and R^b are independently H or optionally substituted C₁-C₆-alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O and S and may be optionally substituted;
- R² is hydrogen, cyano, halo, hydroxyl, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-haloalkoxy-C₁-C₆-alkyl, optionally substituted phenyl;

optionally substituted phenyloxy, optionally substituted 5- or 6-membered heteroaryl, optionally substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkenyl, optionally substituted C₃-C₈-cycloalkyloxy, optionally substituted 3- to 7-membered heterocyclyl containing from one to three heteroatoms selected from the group consisting of N, O and S, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-haloalkylaminocarbonyl, di-C₁-C₆-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, -SO_p(optionally substituted C₁-C₆-alkyl or C₁-C₆-haloalkyl), SF₅, or -NR^aR^b wherein R^a and R^b are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O and S and may be optionally substituted;

R³ is C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-haloalkoxy-C₁-C₆-alkyl, optionally substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkenyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-haloalkylaminocarbonyl, di-C₁-C₆-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, -SF₅, -S(O)_p(C₁-C₆-alkyl or C₁-C₆-haloalkyl), optionally substituted 3- to 7-membered heterocyclyl containing from one to three heteroatoms selected from the group consisting of N, O and S; optionally substituted phenyl, optionally substituted 5- to 10-membered heteroaryl, a 5- to 11-membered spirocyclic heterocyclyl-carbocyclyl group, a 5- to 11-membered spirocyclic heterocyclyl-heterocyclyl group, a 5- to 11-membered spirocyclic carbocyclyl-carbocyclyl group, a 5- to 11-membered spirocyclic carbocyclyl-heterocyclyl group, or -NR^aR^b, wherein R^a and R^b are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O and S and may be optionally substituted;

R⁴ and R^{4'} are independently in each occurrence, hydrogen, halogen, cyano, nitro, hydroxyl, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-

haloalkynyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-haloalkoxy-C₁-C₆-alkyl, optionally substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkyloxy, optionally substituted C₁-C₆-alkylcarbonyl, optionally substituted C₁-C₆-alkoxycarbonyl, optionally substituted aminocarbonyl, C₁-C₆-alkylaminocarbonyl, di(C₁-C₆-alkyl)aminocarbonyl, optionally substituted C₁-C₆-alkylcarbonyloxy, optionally substituted C₁-C₆-alkylcarbonylamino, optionally substituted phenyl, optionally substituted 5- or 6-membered heteroaryl, -SF₅, -SO_p(optionally substituted C₁-C₆-alkyl or C₁-C₆-haloalkyl); or R⁴ together with R^{4'} together form a 2-6-membered chain optionally containing one or two heteroatoms selected from the group consisting of N, O, Si and S to form carbocyclic or heterocyclic ring together with the carbon atom to which they are attached; or -NR^cR^d, wherein R^c and R^d are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^c and R^d may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O and S and may be optionally substituted;

R⁸ is hydrogen, halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₈-cycloalkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl;

R⁹ and R^{9'} are independently hydrogen, halo, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₃-C₈-cycloalkoxy, or R⁹ together with R^{9'} form a 2-6-membered chain optionally containing one or two heteroatoms selected from the group consisting of N, O, Si and S to form carbocyclic or heterocyclic ring together with the carbon atom to which they are attached; and

R['] is hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, optionally substituted C₃-C₈-cycloalkyl, or optionally substituted phenyl.

3. The compound of Formula (I) according to claim 1 or 2, wherein

R¹ is hydrogen, cyano, optionally substituted C₁-C₄-alkyl, optionally substituted C₁-C₄-alkoxy, optionally substituted C₂-C₄-alkenyl, optionally substituted C₂-C₄-alkynyl, optionally substituted C₃-C₈-cycloalkyl, optionally substituted, saturated or unsaturated 5-, 6-, or 7-membered heterocycle, optionally substituted aryl, optionally substituted heteroaryl,

optionally substituted aryloxy, optionally substituted C₁-C₄-alkylcarbonyl, optionally substituted C₁-C₄-alkoxycarbonyl, optionally substituted aminocarbonyl, optionally substituted C₁-C₄-alkylaminocarbonyl, optionally substituted C₁-C₄-dialkylaminocarbonyl, optionally substituted alkyl-SO_p-, haloalkyl-SO_p-, amino, -NH-optionally substituted C₁-C₄-alkyl, or -NR^aR^b, wherein R^a and R^b are independently optionally substituted alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may be optionally substituted;

R' is hydrogen or C₁-C₄-alkyl;

R² is hydrogen, halogen, cyano, nitro, -OH, optionally substituted C₁-C₄-alkyl, optionally substituted C₁-C₄-alkoxy, optionally substituted C₃-C₈-cycloalkyl, amino, NH-optionally substituted C₁-C₄-alkyl, -SF₅, or -NR^aR^b, wherein R^a and R^b are independently optionally substituted C₁-C₄-alkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3, 4, 5, 6, 7, or 8 membered- heterocyclyl group, which may be optionally substituted, -SO_p(optionally substituted C₁-C₄-alkyl or haloalkyl);

R³ is C₁-C₄-alkyl, 6- to 10-membered aryl, 5- to 10-membered heteroaryl, C₃-C₈-cycloalkyl, a 5- to 11-membered spirocyclic heterocyclyl-carbocyclyl group, a 5- to 11-membered spirocyclic heterocyclyl-heterocyclyl group, a 5- to 11-membered spirocyclic carbocyclyl-carbocyclyl group, or a 5- to 11-membered spirocyclic carbocyclyl-heterocyclyl group, each of which may be optionally substituted with 1, 2, or 3 substituents; and

each R⁴ is independently halogen, cyano, nitro, -OH, optionally substituted C₁-C₄-alkyl, optionally substituted C₁-C₄-alkoxy, optionally substituted C₃-C₈-cycloalkyl, amino, NH-optionally substituted C₁-C₄-alkyl, -SF₅, or -NR^cR^d, wherein R^c and R^d are independently optionally substituted C₁-C₄-alkyl; or R^c and R^d may form, with the nitrogen to which they are attached, a 3, 4, 5, 6, 7, or 8 membered- heterocyclyl group, which may be optionally substituted, SO_p(optionally substituted C₁-C₄-alkyl or haloalkyl).

4. The compound of Formula (I) according to any one of claims 1 to 3, wherein Q is N.

5. The compound of Formula (I) according to any one of claims 1 to 3, wherein Q is C-R⁸.
6. The compound of Formula (I) according to any one of claims 1 to 5, wherein X is O.
7. The compound of Formula (I) according to any one of claims 1 to 5, wherein X is S.
8. The compound of Formula (I) according to any one of claims 1 to 7, wherein a is 1 and q is 1.
9. The compound of Formula (I) according to any one of claims 1 to 7, wherein a is 0 and q is 0.
10. The compound of Formula (I) according to any one of claims 1 to 7, wherein a is 1 and q is 0.
11. The compound of Formula (I) according to any one of claims 1 to 10, wherein:

L is L¹;

R' is hydrogen or C₁-C₄-alkyl;

R¹ is halo, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxy-C₁-C₄-alkyl, hydroxy-C₁-C₄-haloalkyl, alkoxy-C₁-C₄-alkyl, alkoxy-C₁-C₄-haloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₂-C₄-alkynyl, C₂-C₄-haloalkynyl, C₁-C₄-alkylcarbonyl, C₁-C₄-haloalkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-haloalkoxycarbonyl, aminocarbonyl, C₁-C₄-alkylaminocarbonyl, C₁-C₄-haloalkylaminocarbonyl, di-C₁-C₄-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, optionally substituted 5- or 6-membered heteroaryl or optionally substituted 5- or 6-membered heterocyclyl containing 1 to 3 N, S or O heteroatoms or -NR^aR^b wherein R^a and R^b are independently H, C₁-C₄-alkyl or C₁-C₄-haloalkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5- or 6-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O and S and may be optionally substituted;

- R^2 is hydrogen, halo, $-SF_5$, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₂-C₄-alkynyl, C₂-C₄-haloalkynyl, optionally substituted C₃-C₆-cycloalkyl, C₁-C₄-alkylcarbonyl, C₁-C₄-haloalkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-haloalkoxycarbonyl, aminocarbonyl, C₁-C₄-alkylaminocarbonyl, C₁-C₄-haloalkylaminocarbonyl, di-C₁-C₄-alkylaminocarbonyl, di-C₁-C₄-haloalkylaminocarbonyl, $-SO_p$ (optionally substituted C₁-C₄-alkyl or C₁-C₄-haloalkyl), or $-NR^aR^b$ wherein R^a and R^b are independently H, C₁-C₄-alkyl or C₁-C₄-haloalkyl;
- R^3 is C₁-C₆-alkyl, C₁-C₆-haloalkyl, optionally substituted C₃-C₈-cycloalkyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-haloalkylaminocarbonyl, di-C₁-C₆-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, $-SO_p$ (optionally substituted C₁-C₆-alkyl or C₁-C₆-haloalkyl), $-SF_5$, optionally substituted 3- to 7-membered heterocyclyl containing from one to three heteroatoms selected from the group consisting of N, O and S; optionally substituted phenyl, optionally substituted 5- or 6-membered heteroaryl, a 5- to 11-membered spirocyclic heterocyclyl-carbocyclyl group, a 5- to 11-membered spirocyclic heterocyclyl-heterocyclyl group, a 5- to 11-membered spirocyclic carbocyclyl-carbocyclyl group, wherein each ring of said spirocyclic group contains 3, 4, 5 or 6 ring atoms; or $-NR^aR^b$ wherein R^a and R^b are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O and S and may be optionally substituted;
- R^4 is independently in each occurrence, hydrogen, halogen, $-SF_5$, C₁-C₆-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, $-SO_p$ (optionally substituted C₁-C₄-alkyl or C₁-C₄-haloalkyl), or $-NR^cR^d$ wherein R^c and R^d are independently H, C₁-C₄-alkyl or C₁-C₄-haloalkyl;
- R^8 is hydrogen, halogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl;
- X is O or S;


Z is O; and

W is CR⁵R⁶.

12. The compound of Formula (I) according to claim 11, wherein a is 1 and q is 1.

13. The compound of Formula (I) according to claim 11, wherein a is 0 or 1 and q is 0.

14. The compound of Formula (I) according to claim 11, which is represented by the compound of formula (Ie):



(Ie)

wherein:

each R¹⁰ is independently halo, cyano, hydroxyl, optionally substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkyloxy, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₃-C₈-halocycloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-haloalkoxy-C₁-C₆-alkyl, optionally substituted phenyl, optionally substituted phenyloxy, optionally substituted 5- or 6-membered heteroaryl, optionally substituted 3- to 7-membered heterocyclyl containing from one to three heteroatoms selected from the group consisting

of N, O, Si and S, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-haloalkylaminocarbonyl, di-C₁-C₆-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, -SO_p(optionally substituted C₁-C₆-alkyl or C₁-C₆-haloalkyl) where p is 0, 1 or 2, SF₅, or -NR^aR^b, wherein R^a and R^b are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted ;

R¹ is halo, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxy-C₁-C₄-alkyl, alkoxy-C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₂-C₄-alkynyl, C₂-C₄-haloalkynyl, C₁-C₄-alkylcarbonyl, C₁-C₄-haloalkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-haloalkoxycarbonyl, aminocarbonyl, C₁-C₄-alkylaminocarbonyl, C₁-C₄-haloalkylaminocarbonyl, di-C₁-C₄-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, optionally substituted 5- or 6-membered heterocyclyl containing 1 to 3 N, S or O heteroatoms or -NR^aR^b wherein R^a and R^b are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5- or 6-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O and S and may be optionally substituted;

R² is hydrogen, halo, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, -SO_p(optionally substituted C₁-C₄-alkyl or C₁-C₄-haloalkyl), -SF₅, or -NR^aR^b wherein R^a and R^b are independently H, C₁-C₄-alkyl or C₁-C₄-haloalkyl;

R⁴ is hydrogen, halo, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, -SO_p(optionally substituted C₁-C₄-alkyl or C₁-C₄-haloalkyl), -SF₅, or -NR^cR^d wherein R^c and R^d are independently H, C₁-C₄-alkyl or C₁-C₄-haloalkyl;

R⁵ and R⁶ are independently hydrogen, halogen, C₁-C₃-alkyl or C₁-C₃-haloalkyl;

R' is hydrogen or C₁-C₃-alkyl;

R⁸ is hydrogen;

R⁹ and R^{9'} are independently hydrogen, halogen, C₁-C₃-alkyl or C₁-C₃-haloalkyl;

Z is CR⁵R⁶ or O;

Y², Y³, Y⁴, Y⁵ are independently CR⁴ or N;

a is 0 or 1; and


m is 0, 1, 2, or 3.

15. The compound according to claim 14, wherein each R¹⁰ is independently halogen and m is 2 or 3.

16. The compound according to claim 14, wherein each R¹⁰ is independently chloro or fluoro and m is 2 or 3.

17. The compound according to claim 14, wherein:
each R¹⁰ is independently chloro or fluoro;
m is 2 or 3;
R¹ is C₁-C₃-alkyl or -NR^aR^b;
R² is hydrogen, halogen, C₁-C₃-alkyl or C₁-C₃-haloalkyl;
W is CH₂;
Z is O; and
R⁹ and R^{9'} are each hydrogen.

18. The compound according to claim 11, which is the compound of formula (If):



(If)

wherein

R¹⁰ is independently cyano, halo, hydroxyl, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₂-C₆-haloalkynyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-haloalkoxy-C₁-C₆-alkyl, optionally substituted phenyl; optionally substituted phenyloxy, optionally substituted 5- or 6-membered heteroaryl, optionally substituted C₃-C₈-cycloalkyl, optionally substituted C₃-C₈-cycloalkyloxy, optionally substituted 3- to 7-membered heterocyclyl containing from one to three heteroatoms selected from the group consisting of N, O, Si and S, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-haloalkylaminocarbonyl, di-C₁-C₆-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, -SO_p(optionally substituted C₁-C₆-alkyl or C₁-C₆-haloalkyl) where p is 0, 1 or 2, SF₅, or -NR^aR^b wherein R^a and R^b are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5-, 6-, 7-, or 8-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O, Si and S and may be optionally substituted;

R¹ is halo, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxy-C₁-C₄-alkyl, alkoxy-C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₂-C₄-alkynyl, C₂-C₄-haloalkynyl,

C₁-C₄-alkylcarbonyl, C₁-C₄-haloalkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-haloalkoxycarbonyl, aminocarbonyl, C₁-C₄-alkylaminocarbonyl, C₁-C₄-haloalkylaminocarbonyl, di-C₁-C₄-alkylaminocarbonyl, di-C₁-C₆-haloalkylaminocarbonyl, optionally substituted 5- or 6-membered heterocyclyl containing 1 to 3 N, S or O heteroatoms or -NR^aR^b wherein R^a and R^b are independently H, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or R^a and R^b may form, with the nitrogen to which they are attached, a 3-, 4-, 5- or 6-membered-heterocyclyl group, which may include one to three additional heteroatoms selected from the group consisting of N, O and S and may be optionally substituted;

R² is hydrogen, halo, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, -SO_p(optionally substituted C₁-C₄-alkyl or C₁-C₄-haloalkyl), -SF₅, or -NR^aR^b wherein R^a and R^b are independently H, C₁-C₄-alkyl or C₁-C₄-haloalkyl;

R⁴ and R^{4'} are independently halo, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, -SO_p(optionally substituted C₁-C₆-alkyl or C₁-C₆-haloalkyl), -SF₅, or -NR^cR^d wherein R^c and R^d are independently H, C₁-C₄-alkyl or C₁-C₄-haloalkyl;

R['] is hydrogen or C₁-C₃-alkyl;

R⁸ is hydrogen;

R⁹ and R^{9'} are each hydrogen;

D is N, SiR¹¹ where R¹¹ is C₁-C₆alkyl or C₁-C₆haloalkyl, C or C-R⁴;

D¹ is N, O, SiR¹¹R¹² where R¹¹ and R¹² are independently C₁-C₆alkyl or C₁-C₆haloalkyl, -CR⁴R^{4'}, S(O)_p where p is 0, 1 or 2; or

D¹ is CR⁴R^{4'} wherein R⁴ and R^{4'} together form a 2- to 5-membered chain optionally substituted with one heteroatom selected from N and O in the chain to form a spirocyclic group;

Y², Y³, Y⁴, Y⁵ are independently CR⁴ or N;


m is 0, 1, 2, or 3; and

b is 0 or 1.


19. The compound according to claim 14, wherein
- R^1 C₁-C₄-alkyl, C₂-C₄ alkenyl, amino, C₁-C₄-alkylamino, di(C₁-C₄-alkyl)amino, morpholino, pyranyl, tetrahydropyranyl, or dihydropyranyl;
- R^2 is hydrogen, halogen, cyano, C₁-C₄-alkyl, C₁-C₄ haloalkyl; and
- R^4 is independently in each occurrence hydrogen, halogen, C₁-C₄-alkyl or C₁-C₄ haloalkyl.
20. The compound according to claim 19, wherein
- R^1 C₁-C₄-alkyl, di(C₁-C₄-alkyl)amino, morpholino, pyranyl, tetrahydropyranyl, or dihydropyranyl;
- R^2 is hydrogen, halogen, C₁-C₃-alkyl or C₁-C₃haloalkyl; and
- R^4 is independently in each occurrence hydrogen, halogen, C₁-C₃-alkyl or C₁-C₃haloalkyl;
- R^{10} is chloro or fluoro; and
- m is 2 or 3
- R.
21. The compound according to claim 18, wherein
- R^1 C₁-C₄-alkyl, C₂-C₄ alkenyl, amino, C₁-C₄-alkylamino, di(C₁-C₄-alkyl)amino, morpholino, pyranyl, tetrahydropyranyl, or dihydropyranyl;
- R^2 is hydrogen, halogen, C₁-C₄-alkyl or C₁-C₄ haloalkyl;
- D is N or CH;
- D^1 is O, Si(CH₃)₂ or CH₂; and
- R^4 is independently in each occurrence hydrogen, halogen, C₁-C₄-alkyl or C₁-C₄ haloalkyl.
22. The compound of any of claims 1 to 21, wherein each of Y², Y³, Y⁴ and Y⁵ are independently -C-R⁴.

23. The compound of any of claims 1 to 21, wherein Y^2 is N; and Y^3 , Y^4 and Y^5 are independently $-C-R^4$.
24. The compound of any of claims 1 to 21, wherein Y^3 is N; and Y^2 , Y^4 and Y^5 are independently $-CR^4$.
25. The compound of any of claims 1 to 21, wherein Y^4 is N and Y^2 , Y^3 and Y^5 are independently $-CR^4$.
26. The compound of any of claims 1 to 21, wherein Y^5 is N and Y^2 , Y^3 and Y^4 are independently $-CR^4$.
27. The compound of any of claims 1 to 21, wherein Y^2 and Y^3 are N; and Y^4 and Y^5 are independently $-CR^4$.
28. The compound of any of claims 1 to 21, wherein Y^3 and Y^4 are N; and Y^2 and Y^5 are independently $-CR^4$.
29. The compound of any of claims 1 to 21, wherein Y^4 and Y^5 are N; and Y^2 and Y^3 are independently $-CR^4$.
30. The compound of any of claims 1 to 21, wherein Y^2 and Y^4 are N; and Y^3 and Y^5 are independently $-CR^4$.
31. The compound of any of claims 1 to 21, wherein Y^3 and Y^5 are N; and Y^2 and Y^4 are independently $-CR^4$.
32. The compound of any of claims 1 to 21, wherein Y^3 and Y^4 are N; and Y^2 and Y^5 are independently $-CR^4$.


33. The compound of claim 1, wherein the compound has the structure presented in the table below, wherein X is oxygen, R' is hydrogen; and the group




represents the following groups:




Ring System A;




Ring System B;




Ring System C;




Ring System D;




Ring System E;




Ring System F;




Ring System G;




Ring System H;




Ring System I;




Ring System J;








Ring System K;




Ring System L;




Ring System M;







Ring System S;




Ring System T;




Ring System U;




Ring System V;




Ring System X;




Ring System Y;




Ring System Z;




Ring System AA;




Ring System AB;




Ring System AC;




Ring System AD;




Ring System AE;




Ring System AF;




Ring System AG;




Ring System AH;




Ring System AJ;




Ring System AK;




Ring System AL;




Ring System AM;




Ring System AN;




Ring System AO;




Ring System AP;




Ring System AQ;




Ring System AR;




Ring System AS;




Ring System AT;




Ring System AU;




Ring System AV;




Ring System AW;




Ring System AX;




Ring System AY;






Ring System AZ;









Ring System AAA;






Formula (I)



Cmpd. #	L	R ¹	R ²	R ³	Ring System
271	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	A
326	L1	<i>i</i> -Pr	Me		A
327	L1	<i>i</i> -Pr	Me		A
326-0	L1	<i>i</i> -Pr	Me		A



Cmpd. #	L	R ¹	R ²	R ³	Ring System
324	L1	<i>i</i> -Pr	Me		A
325	L1	<i>i</i> -Pr	Me		A
323	L1	<i>i</i> -Pr	Me	<i>t</i> -Bu	A
175	L1	<i>i</i> -Pr	Me	3,5-di-F-Ph	A
A407	L1	<i>i</i> -Pr	Me	2,6-di-F-Ph	A
A406	L1	<i>i</i> -Pr	Me	2,6-di-Cl-Ph	A
A413	L1	<i>i</i> -Pr	Me	2,4-di-F-Ph	A
A408	L1		Me	3,5-di-Cl-Ph	A
A412	L1	<i>i</i> -Pr	Me	4-F-Ph	A
A410	L1	<i>i</i> -Pr	Me	3-Cl-4-F-Ph	A
A411	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	C
A409	L1	-N(CH ₃) ₂	Me	3,5-di-Cl-Ph	A
A414	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	D
306	L1	<i>t</i> -Bu	Me	3,5-di-Cl-Ph	A
297	L1	<i>t</i> -Bu	Me	3,5-di-Cl-Ph	B
365	L1	<i>i</i> -Pr	Me	cyclopropyl	A


Cmpd. #	L	R ¹	R ²	R ³	Ring System
371	L1	<i>i</i> -Pr	Me		A
370	L1	<i>i</i> -Pr	Me		A
366	L1	<i>i</i> -Pr	Me	CN	A
369	L1	<i>i</i> -Pr	Me		A
308	L1	prop-1-en-2-yl	Me	3,5-di-Cl-Ph	B
364	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	E
352	L1	<i>i</i> -Pr	H	3,5-di-Cl-Ph	A
320-0	L1	<i>i</i> -Pr	Me	3,5-di-F-Ph	F
345	L1	<i>i</i> -Pr	Me	2,3,5-tri-F-Ph	A
344	L1	<i>i</i> -Pr	Me	2,3,5-tri-F-Ph	B
294	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	B
320	L2	<i>i</i> -Pr	Me	3,5-di-F-Ph	A
277	L1	2-F-prop-2-yl	Me	3,5-di-F-Ph	A
323-0	L1	<i>i</i> -Pr	Me	-CH ₂ CH(CH ₃) ₂	A
298-0	L1	<i>t</i> -Bu	CF ₃	3-Cl-Ph	B
299-0	L1	<i>t</i> -Bu	CF ₃	3-Cl-Ph	A

Cmpd. #	L	R ¹	R ²	R ³	Ring System
299	L1	<i>t</i> -Bu	CF ₃	3,5-di-Cl-Ph	B
298	L1	<i>t</i> -Bu	CF ₃	3,5-di-Cl-Ph	A
304-0	L1	<i>i</i> -Pr	3,5-di-Cl-Ph	Cl	B
321	L1	<i>i</i> -Pr	4-F-Ph	3,5-di-F-Ph	A
322	L1	<i>i</i> -Pr	Me		A
304	L1	<i>i</i> -Pr	Cl	3,5-di-Cl-Ph	B
307	L1	prop-1-en-2-yl	Me	3,5-di-F-Ph	B
296	L1	<i>i</i> -Pr	CF ₃	3,5-di-Cl-Ph	B
295	L1	<i>i</i> -Pr	CF ₃	3,5-di-Cl-Ph	A
293	L1	<i>i</i> -Pr	Me	2,3-di-Cl-Ph	B
276	L1	<i>i</i> -Pr	CF ₃	3,5-di-F-Ph	A
274	L1	<i>i</i> -Pr	Me	2,3-di-Cl-Ph	A
273	L1	<i>i</i> -Pr	Me	3-F-Ph	A
272	L1	<i>i</i> -Pr	Me	3-Cl-5-F-Ph	A
275	L1	<i>i</i> -Pr	Me	3,5-di-F-Ph	B
279	L1	prop-1-en-2-yl	Me	3,5-di-F-Ph	A
174	L1	H	H	2,6-di-F-Ph	A
A400	L1	1,1-difluoroethyl	Me	3,5-di-Cl-Ph	A
A401	L1	CF ₃	Me	3,5-di-Cl-Ph	A

Cmpd. #	L	R ¹	R ²	R ³	Ring System
373	L1	<i>i</i> -Pr	Me		A
372-0	L1	<i>i</i> -Pr	Me		A
A402	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	G
A403	L1	<i>i</i> -Pr	-CH ₂ OH	3,5-di-Cl-Ph	A
A404	L1	<i>i</i> -Pr	-CF ₂ CF ₃	3,5-di-Cl-Ph	A
394	L1	-OCH ₃	H	3,5-di-Cl-Ph	A
398	L1	-OCH ₂ CH ₃	H	3,5-di-Cl-Ph	A
A405	L1	-CHF ₂	Me	3,5-di-Cl-Ph	A
573	L1	<i>t</i> -Bu	Cl	2,3,5-tri-F-Ph	A
559	L1	<i>i</i> -Pr	-CN	3,5-di-Cl-Ph	A
614	L1	<i>t</i> -Bu	Me	3,5-di-Cl-Ph	AAA
451	L1	-N(CH ₃) ₂	CF ₃	3,5-di-Cl-Ph	A
572	L1	<i>t</i> -Bu	Cl	2,5-di-Cl-4-F- Ph	A
528	L1	<i>i</i> -Pr	Me	2,5-di-Cl-4-F- Ph	A
571	L1	<i>t</i> -Bu	Cl	2,4,5-tri-F-Ph	A
574	L1	<i>t</i> -Bu	Cl	2,3-di-Cl-5-F- Ph	A
A415	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	H
A416	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	I
A417	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	J

Cmpd. #	L	R ¹	R ²	R ³	Ring System
A418	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	K
A419	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	L
A420	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	M
560	L1	<i>i</i> -Pr	-CHF ₂	3,5-di-Cl-Ph	A
305	L1	<i>i</i> -Pr	-CHF ₂	3,5-di-Cl-Ph	B
A421	L1		Me	2,6-di-Cl-4-F	A
A422	L1		Me	2,3,5-tri-F-Ph	A
420	L1	<i>t</i> -Bu	Cl	3,5-di-Cl-Ph	A
A423	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	N
A424	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	O
523	L1	<i>t</i> -Bu	Me	2,6-di-Cl-4-F- Ph	A
A425	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	P
A426	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	Q
526	L1	<i>t</i> -Bu	Me	2,3-di-Cl-5-F- Ph	A
527	L1	<i>i</i> -Pr	Me	2,4,6-tri-F-Ph	A
524	L1	<i>t</i> -Bu	Me	2,4,6-tri-F-Ph	A
525	L1	<i>t</i> -Bu	Me	2,3,5-tri-F-Ph	A
414-0	L2	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	R
514	L2	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	E
A427	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	S
A428	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	T

Cmpd. #	L	R ¹	R ²	R ³	Ring System
A429	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	U
A430	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	V
418	L1	<i>t</i> -Bu	H	3,5-di-Cl-Ph	A
513	L2	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	A
513-0	L2	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	F
511	L1		H	3,5-di-Cl-Ph	A
512	L1	-N(CH ₃) ₂	H	3,5-di-Cl-Ph	A
A431	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	W
450	L1		CF ₃	3,5-di-Cl-Ph	A
A432	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	X
A473	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	Y
A433	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	Z
A434	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AA
A435	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AB
A436	L1	L1	<i>i</i> -Pr	Me	N/A (H)
A437	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AC
A438	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AD
A439	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AF
A440	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AG
A441	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AH
A442	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AE
A443	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AJ
A445	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AK

Cmpd. #	L	R ¹	R ²	R ³	Ring System
A446	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AL
A447	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AM
A448	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AO
A449	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AQ
A450	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AR
A451	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AP
A452	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AS
A472	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AT
A453	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AN
A454	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AU
419	L1	<i>i</i> -Pr	Cl	3,5-di-Cl-Ph	A
397-0	L1	OCH ₂ CH=CH ₂	Me	3,5-di-Cl-Ph	A
A455	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AW
A456	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AX
A457	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AY
A458	L1	<i>i</i> -Pr	Me	3,5-di-Cl-Ph	AZ
395	L1	-OCHF ₂	Me	3,5-di-Cl-Ph	A
A459	L1		Me	3,5-di-Cl-Ph	A
A464	L1	-N(CH ₃) ₂	Me	2,6-di-Cl-4-F-Ph	A
A462	L1	-N(CH ₃) ₂	Me	2,4,6-tri-F-Ph	A
A463	L1	-N(CH ₃) ₂	Me	2,3-di-Cl-5-F-Ph	A
A460	L1	-N(CH ₃) ₂	Me	2,3,5-tri-F-Ph	A
A461	L1	-N(CH ₃) ₂	Me	2,3,5-tri-Cl-Ph	A

Cmpd. #	L	R¹	R²	R³	Ring System
558	L1	<i>i</i> -Pr	-C(O)CH ₃	3,5-di-Cl-Ph	A

34. A veterinary composition comprising the compound of any one of claims 1 to 33 or a pharmaceutically or a veterinarily acceptable salt thereof and a veterinarily acceptable carrier.

35. A veterinary composition comprising the compound of any one of claims 1 to 33 or a pharmaceutically or a veterinarily acceptable salt thereof, one or more additional active agent(s), and a veterinarily acceptable carrier.

36. A method for the treatment, control and/or prevention of a parasitic infection or infestation in an animal in need thereof, which comprises administering to said animal an effective amount of the compound of any one of claims 1 to 33 or a pharmaceutically or a veterinarily acceptable salt thereof.

37. Use of a compound of any one of claims 1 to 33 in the preparation of a medicament for the treatment, control and/or prevention of a parasitic infection or infestation in an animal.

38. A compound of Formula (I) of any of claims 1 to 33, for use in a method for the treatment, control and/or prevention of a parasitic infection or infestation in an animal.