

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
18 August 2005 (18.08.2005)

PCT

(10) International Publication Number
WO 2005/076007 A2

(51) International Patent Classification⁷: **G01N 33/68**

(21) International Application Number:
PCT/EP2005/000632

(22) International Filing Date: 22 January 2005 (22.01.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
04002290.7 3 February 2004 (03.02.2004) EP

(71) Applicant (for all designated States except US): **BAYER HEALTHCARE AG** [DE/DE]; 51368 Leverkusen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GOLZ, Stefan** [DE/DE]; Bückmannsmühle 46, 45326 Essen (DE). **BRÜGGEMEIER, Ulf** [DE/DE]; Leysiefen 20, 42799 Leichlingen (DE). **GEERTS, Andreas** [DE/DE]; Schuckertstr. 29, 42113 Wuppertal (DE). **SUMMER, Holger** [DE/DE]; Katernberger Schulweg 3, 42113 Wuppertal (DE).

(74) Common Representative: **BAYER HEALTHCARE AG**; Law and Patents, Patents and Licensing, 51368 Leverkusen (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DIAGNOSTICS AND THERAPEUTICS FOR DISEASES ASSOCIATED WITH G PROTEIN-COUPLED RECEPTOR P2Y₁₂ (P2Y₁₂)

(57) Abstract: The invention provides a human P2Y₁₂ which is associated with the cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases. The invention also provides assays for the identification of compounds useful in the treatment or prevention of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases. The invention also features compounds which bind to and/or activate or inhibit the activity of P2Y₁₂ as well as pharmaceutical compositions comprising such compounds.



WO 2005/076007 A2

Diagnostics and Therapeutics for Diseases Associated with G Protein-Coupled Receptor P2Y₁₂
(P2Y₁₂)

Technical field of the invention

5 The present invention is in the field of molecular biology, more particularly, the present invention relates to nucleic acid sequences and amino acid sequences of a human P2Y₁₂ and its regulation for the treatment of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in mammals.

Background of the invention

10 **G-Protein Coupled Receptors**

P2Y₁₂ is a seven transmembrane G protein coupled receptor (GPCR) [Cattaneo et al. (1992), Nurden et al. (1995), Hollopeter et al. (2001), Oury et al. (2000), Fontana et al. (2003), Jagroop et al. (2003), Zhang et al. (2001), US2001046497, WO0028028, WO0031258]. Many medically significant biological processes are mediated by signal transduction pathways that involve G-
15 proteins [Lefkowitz, (1991)]. The family of G-protein coupled receptors (GPCRs) includes receptors for hormones, neurotransmitters, growth factors, and viruses. Specific examples of GPCRs include receptors for such diverse agents as dopamine, calcitonine, adrenergic hormones, endotheline, cAMP, adenosine, acetylcholine, serotonin, histamine, thrombin, kinine, follicle stimulating hormone, opsins, endothelial differentiation gene-1, rhodopsins, odorants, cytome-
20 galovirus, G-proteins themselves, effector proteins such as phospholipase C, adenylyl cyclase, and phosphodiesterase, and actuator proteins such as protein kinase A and protein kinase C.

GPCRs possess seven conserved membrane-spanning domains connecting at least eight divergent hydrophilic loops. GPCRs, also known as seven transmembrane, 7TM, receptors, have been characterized as including these seven conserved hydrophobic stretches of about 20 to 30 amino
25 acids, connecting at least eight divergent hydrophilic loops. Most GPCRs have single conserved cysteine residues in each of the first two extracellular loops, which form disulfide bonds that are believed to stabilize functional protein structure. The seven transmembrane regions are designated as TM1, TM2, TM3, TM4, TM5, TM6, and TM7. TM3 is being implicated with signal transduction. Phosphorylation and lipidation (palmitoylation or farnesylation) of cysteine residues
30 can influence signal transduction of some GPCRs. Most GPCRs contain potential phosphorylation sites within the third cytoplasmic loop and/or the carboxy terminus. For several GPCRs, such as

the beta-adrenergic receptor, phosphorylation by protein kinase A and/or specific receptor kinases mediates receptor desensitization.

For some receptors, the ligand binding sites of GPCRs are believed to comprise hydrophilic sockets formed by several GPCR transmembrane domains. The hydrophilic sockets are surrounded by hydrophobic residues of the GPCRs. The hydrophilic side of each GPCR transmembrane helix is postulated to face inward and form a polar ligand binding site. TM3 is being implicated with several GPCRs as having a ligand binding site, such as the TM3 aspartate residue. TM5 serines, a TM6 asparagine, and TM6 or TM7 phenylalanines or tyrosines also are implicated in ligand binding.

GPCRs are coupled inside the cell by heterotrimeric G-proteins to various intracellular enzymes, ion channels, and transporters. Different G-protein alpha-subunits preferentially stimulate particular effectors to modulate various biological functions in a cell. Phosphorylation of cytoplasmic residues of GPCRs is an important mechanism for the regulation of some GPCRs. For example, in one form of signal transduction, the effect of hormone binding is the activation of the enzyme, adenylylase, inside the cell. Enzyme activation by hormones is dependent on the presence of the nucleotide GTP. GTP also influences hormone binding. A G-protein connects the hormone receptor to adenylylase. G-protein exchanges GTP for bound GDP when activated by a hormone receptor. The GTP-carrying form then binds to activated adenylylase. Hydrolysis of GTP to GDP, catalyzed by the G-protein itself, returns the G-protein to its basal, inactive form. Thus, the G-protein serves a dual role, as an intermediate that relays the signal from receptor to effector, and as a clock that controls the duration of the signal.

Over the past 15 years, nearly 350 therapeutic agents targeting 7TM receptors have been successfully introduced into the market. This indicates that these receptors have an established, proven history as therapeutic targets. Clearly, there is a need for identification and characterization of further receptors which can play a role in preventing, ameliorating, or correcting dysfunctions or diseases including, but not limited to, infections such as bacterial, fungal, protozoan, and viral infections, particularly those caused by HIV viruses, cancers, allergies including asthma, cardiovascular diseases including acute heart failure, hypotension, hypertension, angina pectoris, myocardial infarction, hematological diseases, genito-urinary diseases including urinary incontinence and benign prostate hyperplasia, osteoporosis, peripheral and central nervous system disorders including pain, Alzheimer's disease and Parkinson's disease, respiratory diseases, metabolic diseases, inflammatory diseases, gastro-enterological diseases, diseases of the endocrine system, dermatological diseases, diseases of muscles or the skeleton, immunological diseases, developmental diseases or diseases of the reproductive system.

TaqMan-Technology / expression profiling

TaqMan is a recently developed technique, in which the release of a fluorescent reporter dye from a hybridisation probe in real-time during a polymerase chain reaction (PCR) is proportional to the accumulation of the PCR product. Quantification is based on the early, linear part of the reaction, and by determining the threshold cycle (CT), at which fluorescence above background is first detected.

Gene expression technologies may be useful in several areas of drug discovery and development, such as target identification, lead optimization, and identification of mechanisms of action. The TaqMan technology can be used to compare differences between expression profiles of normal tissue and diseased tissue. Expression profiling has been used in identifying genes, which are up- or downregulated in a variety of diseases. An interesting application of expression profiling is temporal monitoring of changes in gene expression during disease progression and drug treatment or in patients versus healthy individuals. The premise in this approach is that changes in pattern of gene expression in response to physiological or environmental stimuli (e.g., drugs) may serve as indirect clues about disease-causing genes or drug targets. Moreover, the effects of drugs with established efficacy on global gene expression patterns may provide a guidepost, or a genetic signature, against which a new drug candidate can be compared.

P2Y₁₂

The nucleotide sequence of P2Y₁₂ is accessible in public databases by the accession number NM_022788 and is given in SEQ ID NO: 1. The amino acid sequence of P2Y₁₂ is depicted in SEQ ID NO: 2.

G protein-coupled receptors (GPCRs) are integral membrane proteins containing 7 putative transmembrane domains (TMs). These proteins mediate signals to the interior of the cell via activation of heterotrimeric G proteins that in turn activate various effector proteins, ultimately resulting in a physiologic response.

P2Y₁₂ is published in

Blood platelets fulfill an essential role in primary hemostasis by their capacity to adhere to exposed subendothelium, to aggregate and effect thrombus formation, and to generate procoagulant activity. Aggregation is mediated by a mechanism that involves the binding of fibrinogen, or other adhesive proteins, to the GP IIb-IIIa complexes of platelets (GP2B; GP3A). Among the soluble agonists that stimulate platelet aggregation by reacting with distinct populations of primary receptors in the platelet membrane is adenine diphosphate (ADP). ADP has

at least 2 effects on human platelets: it induces shape change and exposure of fibrinogen-binding determinants, and it inhibits platelet adenylate cyclase, which prevents the accumulation of cAMP. Little is known about the pathology of primary receptors on platelets or of the activation mechanisms that lead to the expression of adhesive protein-binding determinants on GP IIb-IIIa complexes. Cattaneo et al. (1992) reported a congenital defect of platelet function characterized by severe impairment of platelet response to ADP. Nurden et al. (1995) studied a family with what may be a similar abnormality in that ADP induced a small and rapidly reversible platelet aggregation even at high doses. This was associated with a markedly reduced binding of ADP to platelets and a failure of ADP to lower the content of cAMP of platelets preincubated with prostaglandin E1. Electron microscopy showed that the ADP-induced aggregates of platelets in the proband were composed of loosely bound shape-changed platelets with few contact points. Thus, this receptor defect has had a direct influence on the capacity of platelets to bind to each other in response to ADP. The proband was a 61-year-old Caucasian man with a history of episodes of excessive bleeding. The first severe episode occurred at the age of 17 when, after a minor accident, he developed a subretinal hemorrhage that led to the loss of an eye. A second major episode occurred during surgery performed when he was 28 years old. The third episode occurred during middle age when pulmonary bleeding led to hemoptysis. Other affected members of the family included a sister, an uncle who died as a result of severe epistaxis, and his father, who died from bleeding after a motor accident. The parents were cousins. There was no information on the mother, who died during the proband's early childhood. Studies in the sister appeared to indicate that she was similarly involved, whereas the only offspring of the proband appeared normal. Despite the consanguinity, the inheritance of the defect may be autosomal dominant.

ADP mediates platelet aggregation through its action on 2 G protein-coupled receptor subtypes. The P2Y1 receptor (P2RY1) couples to Gq and mobilizes intracellular calcium ions to mediate platelet shape change and aggregation. The second ADP receptor required for aggregation, variously called P2YADP, P2YAC, P2YCYC, or P2TAC, is coupled to the inhibition of adenyl cyclase through Gi. This receptor is the target of efficacious antithrombotic agents, such as ticlopidine and clopidogrel. To identify this Gi-linked platelet ADP receptor, Hollopeter et al. (2001) engineered *Xenopus* oocytes to allow the detection of Gi-linked responses through a sensitive electrophysiologic assay. The human gene cloned, which the authors termed P2Y12, encodes a protein of 342 amino acids that contains 4 extracellular cysteine residues. Northern blot analysis detected abundant P2Y12 expression in human platelets, with lower expression in brain. The predominant transcript of 2.4 kb was absent from all other tissues examined, including peripheral blood leukocytes. A fainter species of approximately 4.5 kb was also detected in platelet and brain, whereas a prominent band of approximately 1.0 kb was observed only in platelet RNA.

Within the brain, the 2.4-kb species was observed in numerous subregions, including the amygdala, caudate nucleus, corpus callosum, hippocampus, substantia nigra, and thalamus. In situ hybridization suggested a glial expression pattern.

By radiation hybrid analysis, Hollopeter et al. (2001) mapped the P2Y₁₂ gene to the interval between DS31279 to DS31280 on 3q24-q25, a region that also includes the P2Y₁ gene and a UDP glucose receptor (KIAA0001). Among G protein-coupled receptors, P2Y₁₂ is most closely related to the UDP glucose receptor (44% identical) but much less so to P2Y₁ (19% identical), suggesting that the UDP glucose receptor and P2Y₁₂ are the products of a relatively recent gene duplication on chromosome 3.

In the patient reported by Nurden et al. (1995), Hollopeter et al. (2001) identified a 2-bp deletion within codon 240 of the P2Y₁₂ gene. Expression was only detected for this allele, although no mutation was found in the coding region for the other allele. Bleeding due to a heterozygous dominant negative mutation in another platelet ADP receptor (P2RX₁) was described by Oury et al. (2000).

In healthy subjects ADP-induced platelet aggregation is associated with a haplotype of the P2Y₁₂ receptor gene. Carriers of the H2 haplotype may have an increased risk of atherothrombosis and less response to platelet inhibitors [Fontana et al. (2003)].

The P2Y₁ purinoceptor and the P2Y₁₂ receptor appear to be involved in ADP-induced platelet shape change, an early phase of platelet activation that precedes aggregation [Jagroop et al. (2003)].

Zhang et al. (2001) et al show that P2Y₁₂ is coupled to G_i and ADP was subsequently shown to inhibit forskolin-stimulated adenylyl cyclase activity through selective activation of SP1999 with an EC₅₀ of 60 nM. Other nucleotides were able to activate P2Y₁₂ with a rank order of potency 2-MeS-ATP = 2-MeS-ADP > ADP = adenosine 5'-O-(thio)diphosphate > 2-Cl-ATP > adenosine 5'-O-(thiotriphosphate).

P2Y₁₂ is published in US2001046497, WO0028028, WO0031258.

The P2Y₁₂ receptor shows the highest homology (49 %) to the human Purinergic receptor P2Y₁₃ as shown in example 1.

Summary of the invention

The invention relates to novel disease associations of P2Y₁₂ polypeptides and polynucleotides.

The invention also relates to novel methods of screening for therapeutic agents for the treatment of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological

diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal. The invention also relates to pharmaceutical compositions for the treatment of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a P2Y₁₂ polypeptide, a P2Y₁₂ polynucleotide, or regulators of P2Y₁₂ or modulators of P2Y₁₂ activity. The invention further comprises methods of diagnosing cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal.

10 **Brief Description of the Drawings**

Fig. 1 shows the nucleotide sequence of a P2Y₁₂ receptor polynucleotide (SEQ ID NO: 1).

Fig. 2 shows the amino acid sequence of a P2Y₁₂ receptor polypeptide (SEQ ID NO: 2).

Fig. 3 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO: 3).

Fig. 4 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO: 4).

15 Fig. 5 shows a nucleotide sequence useful as a probe to detect proteins of the invention (SEQ ID NO: 5).

Detailed description of the invention

Definition of terms

20 An "oligonucleotide" is a stretch of nucleotide residues which has a sufficient number of bases to be used as an oligomer, amplimer or probe in a polymerase chain reaction (PCR). Oligonucleotides are prepared from genomic or cDNA sequence and are used to amplify, reveal, or confirm the presence of a similar DNA or RNA in a particular cell or tissue. Oligonucleotides or oligomers comprise portions of a DNA sequence having at least about 10 nucleotides and as many as about 35 nucleotides, preferably about 25 nucleotides.

25 "Probes" may be derived from naturally occurring or recombinant single- or double-stranded nucleic acids or may be chemically synthesized. They are useful in detecting the presence of identical or similar sequences. Such probes may be labeled with reporter molecules using nick translation, Klenow fill-in reaction, PCR or other methods well known in the art. Nucleic acid probes may be used in southern, northern or in situ hybridizations to determine whether DNA or
30 RNA encoding a certain protein is present in a cell type, tissue, or organ.

A "fragment of a polynucleotide" is a nucleic acid that comprises all or any part of a given nucleotide molecule, the fragment having fewer nucleotides than about 6 kb, preferably fewer than about 1 kb.

5 "Reporter molecules" are radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents which associate with a particular nucleotide or amino acid sequence, thereby establishing the presence of a certain sequence, or allowing for the quantification of a certain sequence.

"Chimeric" molecules may be constructed by introducing all or part of the nucleotide sequence of this invention into a vector containing additional nucleic acid sequence which might be expected to change any one or several of the following P2Y12 characteristics: cellular location, distribution,
10 ligand-binding affinities, interchain affinities, degradation/turnover rate, signaling, etc.

"Active", with respect to a P2Y12 polypeptide, refers to those forms, fragments, or domains of a P2Y12 polypeptide which retain the biological and/or antigenic activity of a P2Y12 polypeptide.

"Naturally occurring P2Y12 polypeptide" refers to a polypeptide produced by cells which have not been genetically engineered and specifically contemplates various polypeptides arising from post-
15 translational modifications of the polypeptide including but not limited to acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation.

"Derivative" refers to polypeptides which have been chemically modified by techniques such as ubiquitination, labeling (see above), pegylation (derivatization with polyethylene glycol), and chemical insertion or substitution of amino acids such as ornithine which do not normally occur in
20 human proteins.

"Conservative amino acid substitutions" result from replacing one amino acid with another having similar structural and/or chemical properties, such as the replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

"Insertions" or "deletions" are typically in the range of about 1 to 5 amino acids. The variation
25 allowed may be experimentally determined by producing the peptide synthetically while systematically making insertions, deletions, or substitutions of nucleotides in the sequence using recombinant DNA techniques.

A "signal sequence" or "leader sequence" can be used, when desired, to direct the polypeptide through a membrane of a cell. Such a sequence may be naturally present on the polypeptides of
30 the present invention or provided from heterologous sources by recombinant DNA techniques.

An "oligopeptide" is a short stretch of amino acid residues and may be expressed from an oligonucleotide. Oligopeptides comprise a stretch of amino acid residues of at least 3, 5, 10 amino acids and at most 10, 15, 25 amino acids, typically of at least 9 to 13 amino acids, and of sufficient length to display biological and/or antigenic activity.

- 5 "Inhibitor" is any substance which retards or prevents a chemical or physiological reaction or response. Common inhibitors include but are not limited to antisense molecules, antibodies, and antagonists.

"Standard expression" is a quantitative or qualitative measurement for comparison. It is based on a statistically appropriate number of normal samples and is created to use as a basis of comparison
10 when performing diagnostic assays, running clinical trials, or following patient treatment profiles.

"Animal" as used herein may be defined to include human, domestic (e.g., cats, dogs, etc.), agricultural (e.g., cows, horses, sheep, etc.) or test species (e.g., mouse, rat, rabbit, etc.).

A "P2Y12 polynucleotide", within the meaning of the invention, shall be understood as being a nucleic acid molecule selected from a group consisting of

- 15 (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2,
- (ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1,
- (iii) nucleic acid molecules having the sequence of SEQ ID NO: 1,
- (iv) nucleic acid molecules the complementary strand of which hybridizes under stringent
20 conditions to a nucleic acid molecule of (i), (ii), or (iii); and
- (v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code;

wherein the polypeptide encoded by said nucleic acid molecule has P2Y12 activity.

A "P2Y12 polypeptide", within the meaning of the invention, shall be understood as being a
25 polypeptide selected from a group consisting of

- (i) polypeptides having the sequence of SEQ ID NO: 2,
- (ii) polypeptides comprising the sequence of SEQ ID NO: 2,

- (iii) polypeptides encoded by P2Y12 polynucleotides; and
- (iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% homology with a polypeptide of (i), (ii), or (iii);

wherein said polypeptide has P2Y12 activity.

- 5 The nucleotide sequences encoding a P2Y12 (or their complement) have numerous applications in techniques known to those skilled in the art of molecular biology. These techniques include use as hybridization probes, use in the construction of oligomers for PCR, use for chromosome and gene mapping, use in the recombinant production of P2Y12, and use in generation of antisense DNA or RNA, their chemical analogs and the like. Uses of nucleotides encoding a P2Y12 disclosed herein
- 10 are exemplary of known techniques and are not intended to limit their use in any technique known to a person of ordinary skill in the art. Furthermore, the nucleotide sequences disclosed herein may be used in molecular biology techniques that have not yet been developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, e.g., the triplet genetic code, specific base pair interactions, etc.
- 15 It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of P2Y12 - encoding nucleotide sequences may be produced. Some of these will only bear minimal homology to the nucleotide sequence of the known and naturally occurring P2Y12. The invention has specifically contemplated each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon
- 20 choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence of naturally occurring P2Y12, and all such variations are to be considered as being specifically disclosed.

- Although the nucleotide sequences which encode a P2Y12, its derivatives or its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring P2Y12
- 25 polynucleotide under stringent conditions, it may be advantageous to produce nucleotide sequences encoding P2Y12 polypeptides or its derivatives possessing a substantially different codon usage. Codons can be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic expression host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the
- 30 nucleotide sequence encoding a P2Y12 polypeptide and/or its derivatives without altering the encoded amino acid sequence include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

Nucleotide sequences encoding a P2Y₁₂ polypeptide may be joined to a variety of other nucleotide sequences by means of well established recombinant DNA techniques. Useful nucleotide sequences for joining to P2Y₁₂ polynucleotides include an assortment of cloning vectors such as plasmids, cosmids, lambda phage derivatives, phagemids, and the like. Vectors of interest include expression vectors, replication vectors, probe generation vectors, sequencing vectors, etc. In general, vectors of interest may contain an origin of replication functional in at least one organism, convenient restriction endonuclease sensitive sites, and selectable markers for one or more host cell systems.

Another aspect of the subject invention is to provide for P2Y₁₂-specific hybridization probes capable of hybridizing with naturally occurring nucleotide sequences encoding P2Y₁₂. Such probes may also be used for the detection of similar GPCR encoding sequences and should preferably show at least 40% nucleotide identity to P2Y₁₂ polynucleotides. The hybridization probes of the subject invention may be derived from the nucleotide sequence presented as SEQ ID NO: 1 or from genomic sequences including promoter, enhancers or introns of the native gene. Hybridization probes may be labelled by a variety of reporter molecules using techniques well known in the art.

It will be recognized that many deletional or mutational analogs of P2Y₁₂ polynucleotides will be effective hybridization probes for P2Y₁₂ polynucleotides. Accordingly, the invention relates to nucleic acid sequences that hybridize with such P2Y₁₂ encoding nucleic acid sequences under stringent conditions.

“Stringent conditions” refers to conditions that allow for the hybridization of substantially related nucleic acid sequences. For instance, such conditions will generally allow hybridization of sequence with at least about 85% sequence identity, preferably with at least about 90% sequence identity, more preferably with at least about 95% sequence identity. Hybridization conditions and probes can be adjusted in well-characterized ways to achieve selective hybridization of human-derived probes. Stringent conditions, within the meaning of the invention are 65°C in a buffer containing 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7 % (w/v) SDS.

Nucleic acid molecules that will hybridize to P2Y₁₂ polynucleotides under stringent conditions can be identified functionally. Without limitation, examples of the uses for hybridization probes include: histochemical uses such as identifying tissues that express P2Y₁₂; measuring mRNA levels, for instance to identify a sample's tissue type or to identify cells that express abnormal levels of P2Y₁₂; and detecting polymorphisms of P2Y₁₂.

PCR provides additional uses for oligonucleotides based upon the nucleotide sequence which encodes P2Y₁₂. Such probes used in PCR may be of recombinant origin, chemically synthesized, or a mixture of both. Oligomers may comprise discrete nucleotide sequences employed under optimized conditions for identification of P2Y₁₂ in specific tissues or diagnostic use. The same
5 two oligomers, a nested set of oligomers, or even a degenerate pool of oligomers may be employed under less stringent conditions for identification of closely related DNAs or RNAs.

Rules for designing polymerase chain reaction (PCR) primers are now established, as reviewed by PCR Protocols. Degenerate primers, i.e., preparations of primers that are heterogeneous at given sequence locations, can be designed to amplify nucleic acid sequences that are highly homologous
10 to, but not identical with P2Y₁₂. Strategies are now available that allow for only one of the primers to be required to specifically hybridize with a known sequence. For example, appropriate nucleic acid primers can be ligated to the nucleic acid sought to be amplified to provide the hybridization partner for one of the primers. In this way, only one of the primers need be based on the sequence of the nucleic acid sought to be amplified.

15 PCR methods for amplifying nucleic acid will utilize at least two primers. One of these primers will be capable of hybridizing to a first strand of the nucleic acid to be amplified and of priming enzyme-driven nucleic acid synthesis in a first direction. The other will be capable of hybridizing the reciprocal sequence of the first strand (if the sequence to be amplified is single stranded, this sequence will initially be hypothetical, but will be synthesized in the first amplification cycle) and
20 of priming nucleic acid synthesis from that strand in the direction opposite the first direction and towards the site of hybridization for the first primer. Conditions for conducting such amplifications, particularly under preferred stringent hybridization conditions, are well known.

Other means of producing specific hybridization probes for P2Y₁₂ include the cloning of nucleic acid sequences encoding P2Y₁₂ or P2Y₁₂ derivatives into vectors for the production of mRNA
25 probes. Such vectors are known in the art, are commercially available and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate reporter molecules.

It is possible to produce a DNA sequence, or portions thereof, entirely by synthetic chemistry. After synthesis, the nucleic acid sequence can be inserted into any of the many available DNA
30 vectors and their respective host cells using techniques which are well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into the nucleotide sequence. Alternately, a portion of sequence in which a mutation is desired can be synthesized and recombined with longer portion of an existing genomic or recombinant sequence.

P2Y12 polynucleotides may be used to produce a purified oligo-or polypeptide using well known methods of recombinant DNA technology. The oligopeptide may be expressed in a variety of host cells, either prokaryotic or eukaryotic. Host cells may be from the same species from which the nucleotide sequence was derived or from a different species. Advantages of producing an
5 oligonucleotide by recombinant DNA technology include obtaining adequate amounts of the protein for purification and the availability of simplified purification procedures.

Quantitative determinations of nucleic acids

An important step in the molecular genetic analysis of human disease is often the enumeration of the copy number of a nucleic acid or the relative expression of a gene in particular tissues.

10 Several different approaches are currently available to make quantitative determinations of nucleic acids. Chromosome-based techniques, such as comparative genomic hybridization (CGH) and fluorescent in situ hybridization (FISH) facilitate efforts to cytogenetically localize genomic regions that are altered in tumor cells. Regions of genomic alteration can be narrowed further using loss of heterozygosity analysis (LOH), in which disease DNA is analyzed and compared with
15 normal DNA for the loss of a heterozygous polymorphic marker. The first experiments used restriction fragment length polymorphisms (RFLPs) [Johnson, (1989)], or hypervariable minisatellite DNA [Barnes, 2000]. In recent years LOH has been performed primarily using PCR amplification of microsatellite markers and electrophoresis of the radio labelled [Jeffreys, (1985)] or fluorescently labelled PCR products [Weber, (1990)] and compared between paired normal and
20 disease DNAs.

A number of other methods have also been developed to quantify nucleic acids [Gergen, (1992)]. More recently, PCR and RT-PCR methods have been developed which are capable of measuring the amount of a nucleic acid in a sample. One approach, for example, measures PCR product quantity in the log phase of the reaction before the formation of reaction products plateaus
25 [Thomas, (1980)].

A gene sequence contained in all samples at relatively constant quantity is typically utilized for sample amplification efficiency normalization. This approach, however, suffers from several drawbacks. The method requires that each sample has equal input amounts of the nucleic acid and that the amplification efficiency between samples is identical until the time of analysis.
30 Furthermore, it is difficult using the conventional methods of PCR quantitation such as gel electrophoresis or plate capture hybridization to determine that all samples are in fact analyzed during the log phase of the reaction as required by the method.

Another method called quantitative competitive (QC)-PCR, as the name implies, relies on the inclusion of an internal control competitor in each reaction [Piatak, (1993), BioTechniques]. The efficiency of each reaction is normalized to the internal competitor. A known amount of internal competitor is typically added to each sample. The unknown target PCR product is compared with the known competitor PCR product to obtain relative quantitation. A difficulty with this general approach lies in developing an internal control that amplifies with the same efficiency than the target molecule.

5' Fluorogenic Nuclease Assays

Fluorogenic nuclease assays are a real time quantitation method that uses a probe to monitor formation of amplification product. The basis for this method of monitoring the formation of amplification product is to measure continuously PCR product accumulation using a dual-labelled fluorogenic oligonucleotide probe, an approach frequently referred to in the literature simply as the "TaqMan method" [Piatak, (1993), Science; Heid, (1996); Gibson, (1996); Holland. (1991)].

The probe used in such assays is typically a short (about 20-25 bases) oligonucleotide that is labeled with two different fluorescent dyes. The 5' terminus of the probe is attached to a reporter dye and the 3' terminus is attached to a quenching dye, although the dyes could be attached at other locations on the probe as well. The probe is designed to have at least substantial sequence complementarity with the probe binding site. Upstream and downstream PCR primers which bind to flanking regions of the locus are added to the reaction mixture. When the probe is intact, energy transfer between the two fluorophors occurs and the quencher quenches emission from the reporter. During the extension phase of PCR, the probe is cleaved by the 5' nuclease activity of a nucleic acid polymerase such as Taq polymerase, thereby releasing the reporter from the oligonucleotide-quencher and resulting in an increase of reporter emission intensity which can be measured by an appropriate detector.

One detector which is specifically adapted for measuring fluorescence emissions such as those created during a fluorogenic assay is the ABI 7700 or 4700 HT manufactured by Applied Biosystems, Inc. in Foster City, Calif. The ABI 7700 uses fiber optics connected with each well in a 96-or 384 well PCR tube arrangement. The instrument includes a laser for exciting the labels and is capable of measuring the fluorescence spectra intensity from each tube with continuous monitoring during PCR amplification. Each tube is re-examined every 8.5 seconds.

Computer software provided with the instrument is capable of recording the fluorescence intensity of reporter and quencher over the course of the amplification. The recorded values will then be used to calculate the increase in normalized reporter emission intensity on a continuous basis. The

increase in emission intensity is plotted versus time, i.e., the number of amplification cycles, to produce a continuous measure of amplification. To quantify the locus in each amplification reaction, the amplification plot is examined at a point during the log phase of product accumulation. This is accomplished by assigning a fluorescence threshold intensity above background and determining the point at which each amplification plot crosses the threshold (defined as the threshold cycle number or Ct). Differences in threshold cycle number are used to quantify the relative amount of PCR target contained within each tube. Assuming that each reaction functions at 100% PCR efficiency, a difference of one Ct represents a two-fold difference in the amount of starting template. The fluorescence value can be used in conjunction with a standard curve to determine the amount of amplification product present.

Non-Probe-Based Detection Methods

A variety of options are available for measuring the amplification products as they are formed. One method utilizes labels, such as dyes, which only bind to double stranded DNA. In this type of approach, amplification product (which is double stranded) binds dye molecules in solution to form a complex. With the appropriate dyes, it is possible to distinguish between dye molecules free in solution and dye molecules bound to amplification product. For example, certain dyes fluoresce only when bound to amplification product. Examples of dyes which can be used in methods of this general type include, but are not limited to, Syber Green.TM. and Pico Green from Molecular Probes, Inc. of Eugene, Oreg., ethidium bromide, propidium iodide, chromomycin, acridine orange, Hoechst 33258, Toto-1, Yoyo-1, DAPI (4',6-diamidino-2-phenylindole hydrochloride).

Another real time detection technique measures alteration in energy fluorescence energy transfer between fluorophors conjugated with PCR primers [Livak, (1995)].

Probe-Based Detection Methods

These detection methods involve some alteration to the structure or conformation of a probe hybridized to the locus between the amplification primer pair. In some instances, the alteration is caused by the template-dependent extension catalyzed by a nucleic acid polymerase during the amplification process. The alteration generates a detectable signal which is an indirect measure of the amount of amplification product formed.

For example, some methods involve the degradation or digestion of the probe during the extension reaction. These methods are a consequence of the 5'-3' nuclease activity associated with some nucleic acid polymerases. Polymerases having this activity cleave mononucleotides or small

oligonucleotides from an oligonucleotide probe annealed to its complementary sequence located within the locus.

The 3' end of the upstream primer provides the initial binding site for the nucleic acid polymerase. As the polymerase catalyzes extension of the upstream primer and encounters the bound probe, the
5 nucleic acid polymerase displaces a portion of the 5' end of the probe and through its nuclease activity cleaves mononucleotides or oligonucleotides from the probe.

The upstream primer and the probe can be designed such that they anneal to the complementary strand in close proximity to one another. In fact, the 3' end of the upstream primer and the 5' end of the probe may abut one another. In this situation, extension of the upstream primer is not
10 necessary in order for the nucleic acid polymerase to begin cleaving the probe. In the case in which intervening nucleotides separate the upstream primer and the probe, extension of the primer is necessary before the nucleic acid polymerase encounters the 5' end of the probe. Once contact occurs and polymerization continues, the 5'-3' exonuclease activity of the nucleic acid polymerase begins cleaving mononucleotides or oligonucleotides from the 5' end of the probe. Digestion of
15 the probe continues until the remaining portion of the probe dissociates from the complementary strand.

In solution, the two end sections can hybridize with each other to form a hairpin loop. In this conformation, the reporter and quencher dye are in sufficiently close proximity that fluorescence from the reporter dye is effectively quenched by the quencher dye. Hybridized probe, in contrast,
20 results in a linearized conformation in which the extent of quenching is decreased. Thus, by monitoring emission changes for the two dyes, it is possible to indirectly monitor the formation of amplification product.

Probes

The labeled probe is selected so that its sequence is substantially complementary to a segment of
25 the test locus or a reference locus. As indicated above, the nucleic acid site to which the probe binds should be located between the primer binding sites for the upstream and downstream amplification primers.

Primers

The primers used in the amplification are selected so as to be capable of hybridizing to sequences
30 at flanking regions of the locus being amplified. The primers are chosen to have at least substantial complementarity with the different strands of the nucleic acid being amplified. When a

probe is utilized to detect the formation of amplification products, the primers are selected in such that they flank the probe, i.e. are located upstream and downstream of the probe.

The primer must have sufficient length so that it is capable of priming the synthesis of extension products in the presence of an agent for polymerization. The length and composition of the primer depends on many parameters, including, for example, the temperature at which the annealing reaction is conducted, proximity of the probe binding site to that of the primer, relative concentrations of the primer and probe and the particular nucleic acid composition of the probe. Typically the primer includes 15-30 nucleotides. However, the length of the primer may be more or less depending on the complexity of the primer binding site and the factors listed above.

10 *Labels for Probes and Primers*

The labels used for labeling the probes or primers of the current invention and which can provide the signal corresponding to the quantity of amplification product can take a variety of forms. As indicated above with regard to the 5' fluorogenic nuclease method, a fluorescent signal is one signal which can be measured. However, measurements may also be made, for example, by monitoring radioactivity, colorimetry, absorption, magnetic parameters, or enzymatic activity. Thus, labels which can be employed include, but are not limited to, fluorophores, chromophores, radioactive isotopes, electron dense reagents, enzymes, and ligands having specific binding partners (e.g., biotin-avidin).

Monitoring changes in fluorescence is a particularly useful way to monitor the accumulation of amplification products. A number of labels useful for attachment to probes or primers are commercially available including fluorescein and various fluorescein derivatives such as FAM, HEX, TET and JOE (all which are available from Applied Biosystems, Foster City, Calif.); lucifer yellow, and coumarin derivatives.

Labels may be attached to the probe or primer using a variety of techniques and can be attached at the 5' end, and/or the 3' end and/or at an internal nucleotide. The label can also be attached to spacer arms of various sizes which are attached to the probe or primer. These spacer arms are useful for obtaining a desired distance between multiple labels attached to the probe or primer.

In some instances, a single label may be utilized; whereas, in other instances, such as with the 5' fluorogenic nuclease assays for example, two or more labels are attached to the probe. In cases wherein the probe includes multiple labels, it is generally advisable to maintain spacing between the labels which is sufficient to permit separation of the labels during digestion of the probe through the 5'-3' nuclease activity of the nucleic acid polymerase.

Patients Exhibiting Symptoms of Disease

A number of diseases are associated with changes in the copy number of a certain gene. For patients having symptoms of a disease, the real-time PCR method can be used to determine if the patient has copy number alterations which are known to be linked with diseases that are associated with the symptoms the patient has.

P2Y₁₂ expression

P2Y₁₂ fusion proteins

Fusion proteins are useful for generating antibodies against P2Y₁₂ polypeptides and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with portions of P2Y₁₂ polypeptides. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such methods are well known in the art and also can be used as drug screens.

A P2Y₁₂ fusion protein comprises two polypeptide segments fused together by means of a peptide bond. The first polypeptide segment can comprise at least 54, 75, 100, 125, 139, 150, 175, 200, 225, 250, or 275 contiguous amino acids of SEQ ID NO: 2 or of a biologically active variant, such as those described above. The first polypeptide segment also can comprise full-length P2Y₁₂.

The second polypeptide segment can be a full-length protein or a protein fragment. Proteins commonly used in fusion protein construction include, but are not limited to β galactosidase, β -glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Additionally, epitope tags are used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, herpes simplex virus (HSV) BP16 protein fusions and G-protein fusions (for example G(α)16, Gs, Gi). A fusion protein also can be engineered to contain a cleavage site located adjacent to the P2Y₁₂.

Preparation of Polynucleotides

A naturally occurring P2Y₁₂ polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides can be made by a cell and isolated

using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the polymerase chain reaction (PCR), or by using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such technique for obtaining a polynucleotide can be used to obtain isolated P2Y12 polynucleotides. For example, restriction enzymes and probes can be used to isolate polynucleotide fragments which comprise P2Y12 nucleotide sequences. Isolated polynucleotides are in preparations which are free or at least 70, 80, or 90% free of other molecules.

P2Y12 cDNA molecules can be made with standard molecular biology techniques, using P2Y12 mRNA as a template. P2Y12 cDNA molecules can thereafter be replicated using molecular biology techniques known in the art. An amplification technique, such as PCR, can be used to obtain additional copies of polynucleotides of the invention, using either human genomic DNA or cDNA as a template.

Alternatively, synthetic chemistry techniques can be used to synthesize P2Y12 polynucleotides. The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which will encode P2Y12 having, for example, an amino acid sequence shown in SEQ ID NO: 2 or a biologically active variant thereof.

Extending Polynucleotides

Various PCR-based methods can be used to extend nucleic acid sequences encoding human P2Y12, for example to detect upstream sequences of P2Y12 gene such as promoters and regulatory elements. For example, restriction-site PCR uses universal primers to retrieve unknown sequence adjacent to a known locus. Genomic DNA is first amplified in the presence of a primer to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region. Primers can be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72°C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA. In this method, multiple restriction enzyme digestions and ligations also can be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries can be useful for extension of sequence into 5' non-transcribed regulatory regions.

Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary sequencing can employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate equipment and software (*e.g.*, GENOTYPER and Sequence NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and electronic data display can be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

Obtaining Polypeptides

P2Y₁₂ can be obtained, for example, by purification from human cells, by expression of P2Y₁₂ polynucleotides, or by direct chemical synthesis.

25 *Protein Purification*

P2Y₁₂ can be purified from any human cell which expresses the receptor, including those which have been transfected with expression constructs which express P2Y₁₂. A purified P2Y₁₂ is separated from other compounds which normally associate with P2Y₁₂ in the cell, such as certain proteins, carbohydrates, or lipids, using methods well-known in the art. Such methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, and preparative gel electrophoresis.

Expression of P2Y12 Polynucleotides

To express P2Y12, P2Y12 polynucleotides can be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art can be used to construct
5 expression vectors containing sequences encoding P2Y12 and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination.

A variety of expression vector/host systems can be utilized to contain and express sequences encoding P2Y12. These include, but are not limited to, microorganisms, such as bacteria
10 transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (e.g., baculovirus), plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids), or animal cell systems.

15 The control elements or regulatory sequences are those non-translated regions of the vector - enhancers, promoters, 5' and 3' untranslated regions -- which interact with host cellular proteins to carry out transcription and translation. Such elements can vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example,
20 when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUESCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and the like can be used. The baculovirus polyhedrin promoter can be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO, and storage protein genes) or from plant viruses (e.g., viral promoters or leader sequences) can be
25 cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of a nucleotide sequence encoding P2Y12, vectors based on SV40 or EBV can be used with an appropriate selectable marker.

Bacterial and Yeast Expression Systems

30 In bacterial systems, a number of expression vectors can be selected. For example, when a large quantity of P2Y12 is needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified can be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT

(Stratagene). In a BLUESCRIPT vector, a sequence encoding P2Y12 can be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced. pIN vectors or pGEX vectors (Promega, Madison, Wis.) also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems can be designed to include heparin, thrombin, or factor Xa protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

10 *Plant and Insect Expression Systems*

If plant expression vectors are used, the expression of sequences encoding P2Y12 can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be used alone or in combination with the omega leader sequence from TMV. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be used. These constructs can be introduced into plant cells by direct DNA transformation or by pathogen-mediated transfection.

An insect system also can be used to express P2Y12. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. Sequences encoding P2Y12 can be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of P2Y12 will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can then be used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which P2Y12 can be expressed.

Mammalian Expression Systems

A number of viral-based expression systems can be used to express P2Y12 in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding P2Y12 can be ligated into an adenovirus transcription/translation complex comprising the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome can be used to obtain a viable virus which is capable of expressing P2Y12 in infected host cells [Engelhard, 1994)]. If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (e.g., liposomes, polycationic amino polymers, or vesicles). Specific initiation signals also can be used to achieve more efficient translation of sequences encoding P2Y₁₂. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding P2Y₁₂, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons can be of various origins, both natural and synthetic.

Host Cells

A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed P2Y₁₂ in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the polypeptide also can be used to facilitate correct insertion, folding and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, VA 20110-2209) and can be chosen to ensure the correct modification and processing of the foreign protein.

Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express P2Y₁₂ can be transformed using expression vectors which can contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells can be allowed to grow for 1-2 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced P2Y₁₂ sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. Any number of selection systems can be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase [Logan, (1984)] and adenine phosphoribosyltransferase [Wigler, (1977)] genes which can be employed in *tk* or *aprt* cells, respectively. Also, antimetabolite, antibiotic, or

herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate [Lowy, (1980)], *npt* confers resistance to the aminoglycosides, neomycin and G-418 [Wigler, (1980)], and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively [Colbere-Garapin, 1981]. Additional selectable genes have been described. For example, *trpB* allows cells to utilize indole in place of tryptophan, or *hisD*, which allows cells to utilize histinol in place of histidine. Visible markers such as anthocyanins, β -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to identify transformants and to quantify the amount of transient or stable protein expression attributable to a specific vector system

10 *Detecting Polypeptide Expression*

Although the presence of marker gene expression suggests that a P2Y12 polynucleotide is also present, its presence and expression may need to be confirmed. For example, if a sequence encoding P2Y12 is inserted within a marker gene sequence, transformed cells containing sequences which encode P2Y12 can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding P2Y12 under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of P2Y12 polynucleotide.

Alternatively, host cells which contain a P2Y12 polynucleotide and which express P2Y12 can be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip-based technologies for the detection and/or quantification of nucleic acid or protein. For example, the presence of a polynucleotide sequence encoding P2Y12 can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides encoding P2Y12. Nucleic acid amplification-based assays involve the use of oligonucleotides selected from sequences encoding P2Y12 to detect transformants which contain a P2Y12 polynucleotide.

A variety of protocols for detecting and measuring the expression of P2Y12, using either polyclonal or monoclonal antibodies specific for the polypeptide, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay using monoclonal antibodies reactive to two non-interfering epitopes on P2Y12 can be used, or a competitive binding assay can be employed.

A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding P2Y12 include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide.

5 Alternatively, sequences encoding P2Y12 can be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to synthesize RNA probes *in vitro* by addition of labeled nucleotides and an appropriate RNA polymerase such as T7, T3, or SP6. These procedures can be conducted using a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical).

10 Suitable reporter molecules or labels which can be used for ease of detection include radio-nuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Expression and Purification of Polypeptides

Host cells transformed with P2Y12 polynucleotides can be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The polypeptide produced by a

15 transformed cell can be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing P2Y12 polynucleotides can be designed to contain signal sequences which direct secretion of soluble P2Y12 through a prokaryotic or eukaryotic cell membrane or which direct the membrane

20 insertion of membrane-bound P2Y12.

As discussed above, other constructions can be used to join a sequence encoding P2Y12 to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals,

25 protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and P2Y12 also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein

30 containing P2Y12 and 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography) Maddox, (1983)], while the enterokinase cleavage site provides a means for purifying P2Y12 from the fusion protein [Porath, (1992)].

Chemical Synthesis

Sequences encoding P2Y12 can be synthesized, in whole or in part, using chemical methods well known in the art. Alternatively, P2Y12 itself can be produced using chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques. Protein synthesis can either be performed using manual techniques or by automation. Automated synthesis can be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Optionally, fragments of P2Y12 can be separately synthesized and combined using chemical methods to produce a full-length molecule.

The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography. The composition of a synthetic P2Y12 can be confirmed by amino acid analysis or sequencing. Additionally, any portion of the amino acid sequence of P2Y12 can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins to produce a variant polypeptide or a fusion protein.

Production of Altered Polypeptides

As will be understood by those of skill in the art, it may be advantageous to produce P2Y12 polynucleotides possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

The nucleotide sequences referred to herein can be engineered using methods generally known in the art to alter P2Y12 polynucleotides for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

Antibodies

Any type of antibody known in the art can be generated to bind specifically to an epitope of P2Y12.

"Antibody" as used herein includes intact immunoglobulin molecules, as well as fragments thereof, such as Fab, F(ab')₂, and Fv, which are capable of binding an epitope of P2Y12.

Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, *e.g.*, at least 15, 25, or 50 amino acid. An antibody which specifically binds to an epitope of P2Y₁₂ can be used therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radio-immunoassays, immunohistochemical assays, immunoprecipitations, or other immunochemical assays known in the art. Various immunoassays can be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays are well known in the art. Such immunoassays typically involve the measurement of complex formation between an immunogen and an antibody which specifically binds to the P2Y₁₂ immunogen.

Typically, an antibody which specifically binds to P2Y₁₂ provides a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in an immunochemical assay. Preferably, antibodies which specifically bind to P2Y₁₂ do not detect other proteins in immunochemical assays and can immunoprecipitate P2Y₁₂ from solution.

P2Y₁₂ can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce polyclonal antibodies. If desired, P2Y₁₂ can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (*e.g.*, aluminum hydroxide), and surface active substances (*e.g.*, lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used in humans, BCG (*bacilli Calmette-Guerin*) and *Corynebacterium parvum* are especially useful.

Monoclonal antibodies which specifically bind to P2Y₁₂ can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique [Roberge, (1995)].

In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. Monoclonal and other antibodies also can be "humanized" to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences by site directed mutagenesis of individual

residues or by grating of entire complementarity determining regions. Antibodies which specifically bind to P2Y₁₂ can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. 5,565,332.

Alternatively, techniques described for the production of single chain antibodies can be adapted
5 using methods known in the art to produce single chain antibodies which specifically bind to P2Y₁₂. Antibodies with related specificity, but of distinct idiotypic composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries. Single-chain antibodies also can be constructed using a DNA amplification method, such as PCR, using hybridoma cDNA as a template. Single-chain antibodies can be mono- or bispecific, and can be bivalent or
10 tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught. A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous
15 phage technology.

Antibodies which specifically bind to P2Y₁₂ also can be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents. Other types of antibodies can be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in
20 WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the "diabodies" described in WO 94/13804, also can be prepared.

Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which P2Y₁₂ is bound.
25 The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

Antisense Oligonucleotides

Antisense oligonucleotides are nucleotide sequences which are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with
30 natural sequences produced by the cell to form complexes and block either transcription or translation. Preferably, an antisense oligonucleotide is at least 11 nucleotides in length, but can be at least 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can

be used. Antisense oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of P2Y12 gene products in the cell.

Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, or a combination of both. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently
5 linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamidate, carboxymethyl esters, carbonates, and phosphate triesters.

Modifications of P2Y12 gene expression can be obtained by designing antisense oligonucleotides
10 which will form duplexes to the control, 5', or regulatory regions of the P2Y12 gene. Oligonucleotides derived from the transcription initiation site, *e.g.*, between positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or
15 chaperons. Therapeutic advances using triplex DNA have been described in the literature [Nicholls, (1993)]. An antisense oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a P2Y12 polynucleotide. Antisense
20 oligonucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a P2Y12 polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent P2Y12 nucleotides, can provide sufficient targeting specificity for P2Y12 mRNA. Preferably, each stretch of complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length.
25 Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular P2Y12 polynucleotide sequence. Antisense oligonucleotides can be modified without affecting their ability to hybridize to a P2Y12 polynucleotide. These modifications can be
30 internal or at one or both ends of the antisense molecule. For example, internucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3', 5'-substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense

oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art.

Ribozymes

Ribozymes are RNA molecules with catalytic activity [Uhlmann, (1987)]. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of specific nucleotide sequences. The coding sequence of a P2Y12 polynucleotide can be used to generate ribozymes which will specifically bind to mRNA transcribed from a P2Y12 polynucleotide. Methods of designing and constructing ribozymes which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art. For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target RNA.

Specific ribozyme cleavage sites within a P2Y12 RNA target can be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target RNA containing the cleavage site can be evaluated for secondary structural features which may render the target inoperable. Suitability of candidate P2Y12 RNA targets also can be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. The nucleotide sequences shown in SEQ ID NO: 1 and its complement provide sources of suitable hybridization region sequences. Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce a ribozyme-containing DNA construct into cells in which it is desired to decrease P2Y12 expression. Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A ribozyme-encoding DNA

construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of ribozymes in the cells (U.S. 5,641,673). Ribozymes also can be engineered to provide an additional level of regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are
5 induced in the cells.

Screening / Screening Assays

Regulators

Regulators as used herein, refer to compounds that affect the activity of a P2Y₁₂ in vivo and/or in vivo. Regulators can be agonists and antagonists of a P2Y₁₂ polypeptide and can be compounds
10 that exert their effect on the P2Y₁₂ activity via the expression, via post-translational modifications or by other means. Agonists of P2Y₁₂ are molecules which, when bound to P2Y₁₂, increase or prolong the activity of P2Y₁₂. Agonists of P2Y₁₂ include proteins, nucleic acids, carbohydrates, small molecules, or any other molecule which activate P2Y₁₂. Antagonists of P2Y₁₂ are
15 molecules which, when bound to P2Y₁₂, decrease the amount or the duration of the activity of P2Y₁₂. Antagonists include proteins, nucleic acids, carbohydrates, antibodies, small molecules, or any other molecule which decrease the activity of P2Y₁₂.

The term "modulate", as it appears herein, refers to a change in the activity of P2Y₁₂ polypeptide. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of P2Y₁₂.

20 As used herein, the terms "specific binding" or "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular structure of the protein recognized by the binding molecule (i.e., the antigenic determinant or epitope). For example, if an antibody is specific for epitope "A" the presence of a polypeptide containing the epitope A, or the presence of free
25 unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The invention provides methods (also referred to herein as "screening assays") for identifying compounds which can be used for the treatment of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological
30 diseases, respiratory diseases, neurological diseases and urological diseases. The methods entail the identification of candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other molecules) which bind to P2Y₁₂ and/or have a stimulatory or inhibitory effect

on the biological activity of P2Y₁₂ or its expression and then determining which of these compounds have an effect on symptoms or diseases regarding the cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in an *in vivo* assay.

Candidate or test compounds or agents which bind to P2Y₁₂ and/or have a stimulatory or inhibitory effect on the activity or the expression of P2Y₁₂ are identified either in assays that employ cells which express P2Y₁₂ on the cell surface (cell-based assays) or in assays with isolated P2Y₁₂ (cell-free assays). The various assays can employ a variety of variants of P2Y₁₂ (e.g., full-length P2Y₁₂, a biologically active fragment of P2Y₁₂, or a fusion protein which includes all or a portion of P2Y₁₂). Moreover, P2Y₁₂ can be derived from any suitable mammalian species (e.g., human P2Y₁₂, rat P2Y₁₂ or murine P2Y₁₂). The assay can be a binding assay entailing direct or indirect measurement of the binding of a test compound or a known P2Y₁₂ ligand to P2Y₁₂. The assay can also be an activity assay entailing direct or indirect measurement of the activity of P2Y₁₂. The assay can also be an expression assay entailing direct or indirect measurement of the expression of P2Y₁₂ mRNA or P2Y₁₂ protein. The various screening assays are combined with an *in vivo* assay entailing measuring the effect of the test compound on the symptoms of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of a membrane-bound (cell surface expressed) form of P2Y₁₂. Such assays can employ full-length P2Y₁₂, a biologically active fragment of P2Y₁₂, or a fusion protein which includes all or a portion of P2Y₁₂. As described in greater detail below, the test compound can be obtained by any suitable means, e.g., from conventional compound libraries. Determining the ability of the test compound to bind to a membrane-bound form of P2Y₁₂ can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the P2Y₁₂-expressing cell can be measured by detecting the labeled compound in a complex. For example, the test compound can be labelled with ¹²⁵I, ³⁵S, ¹⁴C, or ³H, either directly or indirectly, and the radioisotope detected by direct counting of radioemmission or by scintillation counting. Alternatively, the test compound can be enzymatically labelled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product.

In a competitive binding format, the assay comprises contacting P2Y₁₂ expressing cell with a known compound which binds to P2Y₁₂ to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the P2Y₁₂ expressing cell, wherein determining the ability of the test compound to interact with the P2Y₁₂ expressing cell comprises determining the ability of the test compound to preferentially bind the P2Y₁₂ expressing cell as compared to the known compound.

In another embodiment, the assay is a cell-based assay comprising contacting a cell expressing a membrane-bound form of P2Y₁₂ (e.g., full-length P2Y₁₂, a biologically active fragment of P2Y₁₂, or a fusion protein which includes all or a portion of P2Y₁₂) expressed on the cell surface with a test compound and determining the ability of the test compound to modulate (e.g., stimulate or inhibit) the activity of the membrane-bound form of P2Y₁₂. Determining the ability of the test compound to modulate the activity of the membrane-bound form of P2Y₁₂ can be accomplished by any method suitable for measuring the activity of P2Y₁₂, e.g., any method suitable for measuring the activity of a G-protein coupled receptor or other seven-transmembrane receptor (described in greater detail below). The activity of a seven-transmembrane receptor can be measured in a number of ways, not all of which are suitable for any given receptor. Among the measures of activity are: alteration in intracellular Ca²⁺ concentration, activation of phospholipase C, alteration in intracellular inositol triphosphate (IP₃) concentration, alteration in intracellular diacylglycerol (DAG) concentration, and alteration in intracellular adenosine cyclic 3', 5'-monophosphate (cAMP) concentration.

Determining the ability of the test compound to modulate the activity of P2Y₁₂ can be accomplished, for example, by determining the ability of P2Y₁₂ to bind to or interact with a target molecule. The target molecule can be a molecule with which P2Y₁₂ binds or interacts with in nature, for example, a molecule on the surface of a cell which expresses P2Y₁₂, a molecule on the surface of a second cell, a molecule in the extracellular milieu, a molecule associated with the internal surface of a cell membrane or a cytoplasmic molecule. The target molecule can be a component of a signal transduction pathway which facilitates transduction of an extracellular signal (e.g., a signal generated by binding of a P2Y₁₂ ligand, through the cell membrane and into the cell. The target P2Y₁₂ molecule can be, for example, a second intracellular protein which has catalytic activity or a protein which facilitates the association of downstream signaling molecules with P2Y₁₂.

Determining the ability of P2Y₁₂ to bind to or interact with a target molecule can be accomplished by one of the methods described above for determining direct binding. In one embodiment, determining the ability of a polypeptide of the invention to bind to or interact with a target

molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (e.g., intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/-enzymatic activity of the target on an appropriate substrate, detecting the induction of a reporter gene (e.g., a regulatory element that is responsive to a polypeptide of the invention operably linked to a nucleic acid encoding a detectable marker, e.g., luciferase), or detecting a cellular response.

The present invention also includes cell-free assays. Such assays involve contacting a form of P2Y12 (e.g., full-length P2Y12, a biologically active fragment of P2Y12, or a fusion protein comprising all or a portion of P2Y12) with a test compound and determining the ability of the test compound to bind to P2Y12. Binding of the test compound to P2Y12 can be determined either directly or indirectly as described above. In one embodiment, the assay includes contacting P2Y12 with a known compound which binds P2Y12 to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with P2Y12, wherein determining the ability of the test compound to interact with P2Y12 comprises determining the ability of the test compound to preferentially bind to P2Y12 as compared to the known compound.

The cell-free assays of the present invention are amenable to use of either a membrane-bound form of P2Y12 or a soluble fragment thereof. In the case of cell-free assays comprising the membrane-bound form of the polypeptide, it may be desirable to utilize a solubilizing agent such that the membrane-bound form of the polypeptide is maintained in solution. Examples of such solubilizing agents include but are not limited to non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton X-100, Triton X-114, Thesit, Isotridecypoly(ethylene glycol ether)_n, 3-[(3-cholamidopropyl)dimethylamminio]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)-dimethylamminio]-2-hydroxy-1-propane sulfonate (CHAPSO), or N-dodecyl=N,N-dimethyl-3-ammonio-1-propane sulfonate.

In various embodiments of the above assay methods of the present invention, it may be desirable to immobilize P2Y12 (or a P2Y12 target molecule) to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to P2Y12, or interaction of P2Y12 with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase

(GST) fusion proteins or glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical; St. Louis, Mo.) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or P2Y12, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components and complex formation is measured either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of binding or activity of P2Y12 can be determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either P2Y12 or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated polypeptide of the invention or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals; Rockford, Ill.), and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies reactive with P2Y12 or target molecules but which do not interfere with binding of the polypeptide of the invention to its target molecule can be derivatized to the wells of the plate, and unbound target or polypeptide of the invention trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with P2Y12 or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with P2Y12 or target molecule.

The screening assay can also involve monitoring the expression of P2Y12. For example, regulators of expression of P2Y12 can be identified in a method in which a cell is contacted with a candidate compound and the expression of P2Y12 protein or mRNA in the cell is determined. The level of expression of P2Y12 protein or mRNA the presence of the candidate compound is compared to the level of expression of P2Y12 protein or mRNA in the absence of the candidate compound. The candidate compound can then be identified as a regulator of expression of P2Y12 based on this comparison. For example, when expression of P2Y12 protein or mRNA protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of P2Y12 protein or mRNA expression. Alternatively, when expression of P2Y12 protein or mRNA is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of P2Y12 protein or mRNA expression. The level of P2Y12 protein or mRNA expression in the cells can be determined by methods described below.

Binding Assays

For binding assays, the test compound is preferably a small molecule which binds to and occupies the active site of P2Y₁₂ polypeptide, thereby making the ligand binding site inaccessible to substrate such that normal biological activity is prevented. Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules. Potential ligands which bind to a polypeptide of the invention include, but are not limited to, the natural ligands of known P2Y₁₂ GPCRs and analogues or derivatives thereof.

In binding assays, either the test compound or the P2Y₁₂ polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is bound to P2Y₁₂ polypeptide can then be accomplished, for example, by direct counting of radioemission, by scintillation counting, or by determining conversion of an appropriate substrate to a detectable product. Alternatively, binding of a test compound to a P2Y₁₂ polypeptide can be determined without labeling either of the interactants. For example, a microphysiometer can be used to detect binding of a test compound with a P2Y₁₂ polypeptide. A microphysiometer (*e.g.*, CytosensorTM) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a test compound and P2Y₁₂ [Haseloff, (1988)].

Determining the ability of a test compound to bind to P2Y₁₂ also can be accomplished using a technology such as real-time Bimolecular Interaction Analysis (BIA) [McConnell, (1992); Sjolander, (1991)]. BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (*e.g.*, BIAcoreTM). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

In yet another aspect of the invention, a P2Y₁₂-like polypeptide can be used as a "bait protein" in a two-hybrid assay or three-hybrid assay [Szabo, (1995); U.S. 5,283,317], to identify other proteins which bind to or interact with P2Y₁₂ and modulate its activity.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide encoding P2Y₁₂ can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct a DNA sequence that encodes an unidentified protein ("prey" or "sample")

can be fused to a polynucleotide that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact *in vivo* to form a protein-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ), which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein which interacts with P2Y12.

It may be desirable to immobilize either the P2Y12 (or polynucleotide) or the test compound to facilitate separation of the bound form from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either the P2Y12-like polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or particles such as beads (including, but not limited to, latex, polystyrene, or glass beads). Any method known in the art can be used to attach P2Y12-like polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to P2Y12 (or a polynucleotide encoding for P2Y12) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

In one embodiment, P2Y12 is a fusion protein comprising a domain that allows binding of P2Y12 to a solid support. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the non-adsorbed P2Y12; the mixture is then incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.

Other techniques for immobilizing proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either P2Y12 (or a polynucleotide encoding P2Y12) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin.

Biotinylated P2Y₁₂ (or a polynucleotide encoding biotinylated P2Y₁₂) or test compounds can be prepared from biotin-NHS (N-hydroxysuccinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies which specifically bind to P2Y₁₂,
5 polynucleotide, or a test compound, but which do not interfere with a desired binding site, such as the active site of P2Y₁₂, can be derivatized to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies which
10 specifically bind to P2Y₁₂ polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of P2Y₁₂ polypeptide, and SDS gel electrophoresis under non-reducing conditions.

Screening for test compounds which bind to a P2Y₁₂ polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a P2Y₁₂ polypeptide or polynucleotide can
15 be used in a cell-based assay system. A P2Y₁₂ polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test compound to P2Y₁₂ or a polynucleotide encoding P2Y₁₂ is determined as described above.

Functional Assays

Test compounds can be tested for the ability to increase or decrease P2Y₁₂ activity of a P2Y₁₂
20 polypeptide. The P2Y₁₂ activity can be measured, for example, using methods described in the specific examples, below. P2Y₁₂ activity can be measured after contacting either a purified P2Y₁₂, a cell membrane preparation, or an intact cell with a test compound. A test compound which decreases P2Y₁₂ activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for decreasing P2Y₁₂ activity. A test compound
25 which increases P2Y₁₂ activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for increasing P2Y₁₂ activity.

One such screening procedure involves the use of melanophores which are transfected to express P2Y₁₂. Such a screening technique is described in PCT WO 92/01810 published Feb. 6, 1992. Thus, for example, such an assay may be employed for screening for a compound which inhibits
30 activation of the receptor polypeptide of the present invention by contacting the melanophore cells which encode the receptor with both the receptor ligand and a compound to be screened. Inhibition of the signal generated by the ligand indicates that a compound is a potential antagonist for the receptor, i.e., inhibits activation of the receptor. The screen may be employed for

identifying a compound which activates the receptor by contacting such cells with compounds to be screened and determining whether each compound generates a signal, *i.e.*, activates the receptor.

Other screening techniques include the use of cells which express P2Y₁₂ (for example, transfected
5 CHO cells) in a system which measures extracellular pH changes caused by receptor activation [Iwabuchi, (1993)]. For example, compounds may be contacted with a cell which expresses the receptor polypeptide of the present invention and a second messenger response, *e.g.*, signal transduction or pH changes, can be measured to determine whether the potential compound activates or inhibits the receptor. Another such screening technique involves introducing RNA
10 encoding P2Y₁₂ into *Xenopus* oocytes to transiently express the receptor. The receptor oocytes can then be contacted with the receptor ligand and a compound to be screened, followed by detection of inhibition or activation of a calcium signal in the case of screening for compounds which are thought to inhibit activation of the receptor.

Another screening technique involves expressing P2Y₁₂ in cells in which the receptor is linked to
15 a phospholipase C or D. Such cells include endothelial cells, smooth muscle cells, embryonic kidney cells, etc. The screening may be accomplished as described above by quantifying the degree of activation of the receptor from changes in the phospholipase activity.

Gene Expression

In another embodiment, test compounds which increase or decrease P2Y₁₂ gene expression are
20 identified. As used herein, the term "correlates with expression of a polynucleotide" indicates that the detection of the presence of nucleic acids, the same or related to a nucleic acid sequence encoding P2Y₁₂, by northern analysis or real-time PCR is indicative of the presence of nucleic acids encoding P2Y₁₂ in a sample, and thereby correlates with expression of the transcript from the polynucleotide encoding P2Y₁₂. The term "microarray", as used herein, refers to an array of
25 distinct polynucleotides or oligonucleotides arrayed on a substrate, such as paper, nylon or any other type of membrane, filter, chip, glass slide, or any other suitable solid support. A P2Y₁₂ polynucleotide is contacted with a test compound, and the expression of an RNA or polypeptide product of P2Y₁₂ polynucleotide is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test compound is compared to the level of expression of mRNA
30 or polypeptide in the absence of the test compound. The test compound can then be identified as a regulator of expression based on this comparison. For example, when expression of mRNA or polypeptide is greater in the presence of the test compound than in its absence, the test compound is identified as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively,

when expression of the mRNA or polypeptide is less in the presence of the test compound than in its absence, the test compound is identified as an inhibitor of the mRNA or polypeptide expression.

The level of P2Y₁₂ mRNA or polypeptide expression in the cells can be determined by methods well known in the art for detecting mRNA or polypeptide. Either qualitative or quantitative methods can be used. The presence of polypeptide products of P2Y₁₂ polynucleotide can be determined, for example, using a variety of techniques known in the art, including immunochemical methods such as radioimmunoassay, Western blotting, and immunohistochemistry. Alternatively, polypeptide synthesis can be determined *in vivo*, in a cell culture, or in an *in vitro* translation system by detecting incorporation of labelled amino acids into P2Y₁₂.

Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell which expresses P2Y₁₂ polynucleotide can be used in a cell-based assay system. The P2Y₁₂ polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Either a primary culture or an established cell line can be used.

Test Compounds

Suitable test compounds for use in the screening assays of the invention can be obtained from any suitable source, e.g., conventional compound libraries. The test compounds can also be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds [Lam, (1997)]. Examples of methods for the synthesis of molecular libraries can be found in the art. Libraries of compounds may be presented in solution or on beads, bacteria, spores, plasmids or phage.

Modeling of Regulators

Computer modeling and searching technologies permit identification of compounds, or the improvement of already identified compounds, that can modulate P2Y₁₂ expression or activity. Having identified such a compound or composition, the active sites or regions are identified. Such active sites might typically be ligand binding sites, such as the interaction domain of the ligand with P2Y₁₂. The active site can be identified using methods known in the art including, for example, from the amino acid sequences of peptides, from the nucleotide sequences of nucleic acids, or from study of complexes of the relevant compound or composition with its natural ligand.

In the latter case, chemical or X-ray crystallographic methods can be used to find the active site by finding where on the factor the complexed ligand is found.

Next, the three dimensional geometric structure of the active site is determined. This can be done by known methods, including X-ray crystallography, which can determine a complete molecular
5 structure. On the other hand, solid or liquid phase NMR can be used to determine certain intramolecular distances. Any other experimental method of structure determination can be used to obtain partial or complete geometric structures. The geometric structures may be measured with a complexed ligand, natural or artificial, which may increase the accuracy of the active site structure determined.

10 If an incomplete or insufficiently accurate structure is determined, the methods of computer based numerical modeling can be used to complete the structure or improve its accuracy. Any recognized modeling method may be used, including parameterized models specific to particular biopolymers such as proteins or nucleic acids, molecular dynamics models based on computing molecular motions, statistical mechanics models based on thermal ensembles, or combined models.
15 For most types of models, standard molecular force fields, representing the forces between constituent atoms and groups, are necessary, and can be selected from force fields known in physical chemistry. The incomplete or less accurate experimental structures can serve as constraints on the complete and more accurate structures computed by these modeling methods.

Finally, having determined the structure of the active site, either experimentally, by modeling, or
20 by a combination, candidate modulating compounds can be identified by searching databases containing compounds along with information on their molecular structure. Such a search seeks compounds having structures that match the determined active site structure and that interact with the groups defining the active site. Such a search can be manual, but is preferably computer assisted. These compounds found from this search are potential P2Y₁₂ modulating compounds.

25 Alternatively, these methods can be used to identify improved modulating compounds from an already known modulating compound or ligand. The composition of the known compound can be modified and the structural effects of modification can be determined using the experimental and computer modeling methods described above applied to the new composition. The altered structure is then compared to the active site structure of the compound to determine if an improved
30 fit or interaction results. In this manner systematic variations in composition, such as by varying side groups, can be quickly evaluated to obtain modified modulating compounds or ligands of improved specificity or activity.

Therapeutic Indications and Methods

It was found by the present applicant that P2Y₁₂ is expressed in various human tissues.

Neurology

5 CNS disorders include disorders of the central nervous system as well as disorders of the peripheral nervous system.

CNS disorders include, but are not limited to brain injuries, cerebrovascular diseases and their consequences, Parkinson's disease, corticobasal degeneration, motor neuron disease, dementia, including ALS, multiple sclerosis, traumatic brain injury, stroke, post-stroke, post-traumatic brain injury, and small-vessel cerebrovascular disease. Dementias, such as Alzheimer's disease,
10 vascular dementia, dementia with Lewy bodies, frontotemporal dementia and Parkinsonism linked to chromosome 17, frontotemporal dementias, including Pick's disease, progressive nuclear palsy, corticobasal degeneration, Huntington's disease, thalamic degeneration, Creutzfeld-Jakob dementia, HIV dementia, schizophrenia with dementia, and Korsakoff's psychosis, within the meaning of the definition are also considered to be CNS disorders.

15 Similarly, cognitive-related disorders, such as mild cognitive impairment, age-associated memory impairment, age-related cognitive decline, vascular cognitive impairment, attention deficit disorders, attention deficit hyperactivity disorders, and memory disturbances in children with learning disabilities are also considered to be CNS disorders.

Pain, within the meaning of this definition, is also considered to be a CNS disorder. Pain can be
20 associated with CNS disorders, such as multiple sclerosis, spinal cord injury, sciatica, failed back surgery syndrome, traumatic brain injury, epilepsy, Parkinson's disease, post-stroke, and vascular lesions in the brain and spinal cord (e.g., infarct, hemorrhage, vascular malformation). Non-central neuropathic pain includes that associated with post mastectomy pain, phantom feeling, reflex sympathetic dystrophy (RSD), trigeminal neuralgiaradioculopathy, post-surgical pain, HIV/AIDS
25 related pain, cancer pain, metabolic neuropathies (e.g., diabetic neuropathy, vasculitic neuropathy secondary to connective tissue disease), paraneoplastic polyneuropathy associated, for example, with carcinoma of lung, or leukemia, or lymphoma, or carcinoma of prostate, colon or stomach, trigeminal neuralgia, cranial neuralgias, and post-herpetic neuralgia. Pain associated with peripheral nerve damage, central pain (i.e. due to cerebral ischemia) and various chronic pain i.e.,
30 lumbago, back pain (low back pain), inflammatory and/or rheumatic pain. Headache pain (for example, migraine with aura, migraine without aura, and other migraine disorders), episodic and

chronic tension-type headache, tension-type like headache, cluster headache, and chronic paroxysmal hemicrania are also CNS disorders.

Visceral pain such as pancreatitis, intestinal cystitis, dysmenorrhea, irritable Bowel syndrome, Crohn's disease, biliary colic, ureteral colic, myocardial infarction and pain syndromes of the pelvic cavity, e.g., vulvodynia, orchialgia, urethral syndrome and protatodynia are also CNS disorders.

Also considered to be a disorder of the nervous system are acute pain, for example postoperative pain, and pain after trauma.

The human P2Y₁₂ is highly expressed in the following brain tissues: fetal brain, brain, Alzheimer brain, cerebellum, Alzheimer brain cerebellum, cerebellum (left), cerebral cortex, Alzheimer cerebral cortex, frontal lobe, occipital lobe, parietal lobe, temporal lobe, substantia nigra, caudatum, cerebral peduncles, corpus callosum, nucleus accumbens, putamen, hippocampus, thalamus, posteroventral thalamus, dorsomedial thalamus, spinal cord, spinal cord (ventral horn), spinal cord (dorsal horn), neural progenitor cells, astrocytes. The expression in brain tissues and in particular the differential expression between diseased tissue Alzheimer brain and healthy tissue brain, between diseased tissue Alzheimer brain cerebellum and healthy tissue , between diseased tissue Alzheimer cerebral cortex and healthy tissue cerebral cortex demonstrates that the human P2Y₁₂ or mRNA can be utilized to diagnose nervous system diseases. Additionally the activity of the human P2Y₁₂ can be modulated to treat nervous system diseases.

20 *Cardiovascular Disorders*

Heart failure is defined as a pathophysiological state in which an abnormality of cardiac function is responsible for the failure of the heart to pump blood at a rate commensurate with the requirement of the metabolizing tissue. It includes all forms of pumping failures such as high-output and low-output, acute and chronic, right-sided or left-sided, systolic or diastolic, independent of the underlying cause.

Myocardial infarction (MI) is generally caused by an abrupt decrease in coronary blood flow that follows a thrombotic occlusion of a coronary artery previously narrowed by arteriosclerosis. MI prophylaxis (primary and secondary prevention) is included as well as the acute treatment of MI and the prevention of complications.

Ischemic diseases are conditions in which the coronary flow is restricted resulting in a perfusion which is inadequate to meet the myocardial requirement for oxygen. This group of diseases includes stable angina, unstable angina and asymptomatic ischemia.

Arrhythmias include all forms of atrial and ventricular tachyarrhythmias, atrial tachycardia, atrial flutter, atrial fibrillation, atrio-ventricular reentrant tachycardia, preexcitation syndrome, ventricular tachycardia, ventricular flutter, ventricular fibrillation, as well as bradycardic forms of arrhythmias.

- 5 Hypertensive vascular diseases include primary as well as all kinds of secondary arterial hypertension, renal, endocrine, neurogenic, others. The genes may be used as drug targets for the treatment of hypertension as well as for the prevention of all complications arising from cardiovascular diseases.

- 10 Peripheral vascular diseases are defined as vascular diseases in which arterial and/or venous flow is reduced resulting in an imbalance between blood supply and tissue oxygen demand. It includes chronic peripheral arterial occlusive disease (PAOD), acute arterial thrombosis and embolism, inflammatory vascular disorders, Raynaud's phenomenon and venous disorders.

- 15 Atherosclerosis is a cardiovascular disease in which the vessel wall is remodeled, compromising the lumen of the vessel. The atherosclerotic remodeling process involves accumulation of cells, both smooth muscle cells and monocyte/macrophage inflammatory cells, in the intima of the vessel wall. These cells take up lipid, likely from the circulation, to form a mature atherosclerotic lesion. Although the formation of these lesions is a chronic process, occurring over decades of an adult human life, the majority of the morbidity associated with atherosclerosis occurs when a lesion ruptures, releasing thrombogenic debris that rapidly occludes the artery. When such an acute event
20 occurs in the coronary artery, myocardial infarction can ensue, and in the worst case, can result in death.

- The formation of the atherosclerotic lesion can be considered to occur in five overlapping stages such as migration, lipid accumulation, recruitment of inflammatory cells, proliferation of vascular smooth muscle cells, and extracellular matrix deposition. Each of these processes can be shown to
25 occur in man and in animal models of atherosclerosis, but the relative contribution of each to the pathology and clinical significance of the lesion is unclear.

Thus, a need exists for therapeutic methods and agents to treat cardiovascular pathologies, such as atherosclerosis and other conditions related to coronary artery disease.

- 30 Cardiovascular diseases include but are not limited to disorders of the heart and the vascular system like congestive heart failure, myocardial infarction, ischemic diseases of the heart, all kinds of atrial and ventricular arrhythmias, hypertensive vascular diseases, peripheral vascular diseases, and atherosclerosis.

Too high or too low levels of fats in the bloodstream, especially cholesterol, can cause long-term problems. The risk to develop atherosclerosis and coronary artery or carotid artery disease (and thus the risk of having a heart attack or stroke) increases with the total cholesterol level increasing. Nevertheless, extremely low cholesterol levels may not be healthy. Examples of disorders of lipid metabolism are hyperlipidemia (abnormally high levels of fats (cholesterol, triglycerides, or both) in the blood, may be caused by family history of hyperlipidemia), obesity, a high-fat diet, lack of exercise, moderate to high alcohol consumption, cigarette smoking, poorly controlled diabetes, and an underactive thyroid gland), hereditary hyperlipidemias (type I hyperlipoproteinemia (familial hyperchylomicronemia), type II hyperlipoproteinemia (familial hypercholesterolemia), type III hyperlipoproteinemia, type IV hyperlipoproteinemia, or type V hyperlipoproteinemia), hypolipoproteinemia, lipidoses (caused by abnormalities in the enzymes that metabolize fats), Gaucher's disease, Niemann-Pick disease, Fabry's disease, Wolman's disease, cerebrotendinous xanthomatosis, sitosterolemia, Refsum's disease, or Tay-Sachs disease.

Kidney disorders may lead to hypertension or hypotension. Examples for kidney problems possibly leading to hypertension are renal artery stenosis, pyelonephritis, glomerulonephritis, kidney tumors, polycystic kidney disease, injury to the kidney, or radiation therapy affecting the kidney. Excessive urination may lead to hypotension.

The human P2Y₁₂ is highly expressed in the following cardiovascular related tissues: heart atrium (right), heart atrium (left), heart ventricle (left), heart ventricle (right), aorta sclerotic, artery, vein, coronary artery endothel cells, aortic endothel cells, adrenal gland, liver, liver liver cirrhosis, thrombocytes. Expression in the above mentioned tissues and in particular the differential expression between diseased tissue aorta sclerotic and healthy tissue demonstrates that the human P2Y₁₂ or mRNA can be utilized to diagnose of cardiovascular diseases. Additionally the activity of the human P2Y₁₂ can be modulated to treat cardiovascular diseases.

The human P2Y₁₂ is highly expressed in liver tissues: liver, liver liver cirrhosis. Expression in liver tissues demonstrates that the human P2Y₁₂ or mRNA can be utilized to diagnose of dyslipidemia disorders as an cardiovascular disorder. Additionally the activity of the human P2Y₁₂ can be modulated to treat - but not limited to - dyslipidemia disorders.

The human P2Y₁₂ is highly expressed in adrenal gland. Expression in adrenal gland tissues demonstrates that the human P2Y₁₂ or mRNA can be utilized to diagnose of blood pressure disorders as an cardiovascular disorder. Additionally the activity of the human P2Y₁₂ can be modulated to treat - but not limited to - blood pressure disorders as hypertension or hypotension.

Hematological Disorders

Hematological disorders comprise diseases of the blood and all its constituents as well as diseases of organs and tissues involved in the generation or degradation of all the constituents of the blood. They include but are not limited to 1) Anemias, 2) Myeloproliferative Disorders, 3) Hemorrhagic Disorders, 4) Leukopenia, 5) Eosinophilic Disorders, 6) Leukemias, 7) Lymphomas, 8) Plasma Cell Dyscrasias, 9) Disorders of the Spleen in the course of hematological disorders. Disorders according to 1) include, but are not limited to anemias due to defective or deficient hem synthesis, deficient erythropoiesis. Disorders according to 2) include, but are not limited to polycythemia vera, tumor-associated erythrocytosis, myelofibrosis, thrombocythemia. Disorders according to 3) include, but are not limited to vasculitis, thrombocytopenia, heparin-induced thrombocytopenia, thrombotic thrombocytopenic purpura, hemolytic-uremic syndrome, hereditary and acquired disorders of platelet function, hereditary coagulation disorders. Disorders according to 4) include, but are not limited to neutropenia, lymphocytopenia. Disorders according to 5) include, but are not limited to hypereosinophilia, idiopathic hypereosinophilic syndrome. Disorders according to 6) include, but are not limited to acute myeloid leukemia, acute lymphoblastic leukemia, chronic myelocytic leukemia, chronic lymphocytic leukemia, myelodysplastic syndrome. Disorders according to 7) include, but are not limited to Hodgkin's disease, non-Hodgkin's lymphoma, Burkitt's lymphoma, mycosis fungoides cutaneous T-cell lymphoma. Disorders according to 8) include, but are not limited to multiple myeloma, macroglobulinemia, heavy chain diseases. In extension of the preceding idiopathic thrombocytopenic purpura, iron deficiency anemia, megaloblastic anemia (vitamin B12 deficiency), aplastic anemia, thalassemia, malignant lymphoma bone marrow invasion, malignant lymphoma skin invasion, hemolytic uremic syndrome, giant platelet disease are considered to be hematological diseases too.

The human P2Y₁₂ is highly expressed in the following tissues of the hematological system: leukocytes (peripheral blood), Jurkat (T-cells), bone marrow, erythrocytes, thrombocytes, bone marrow CD71+ cells, bone marrow CD34+ cells, bone marrow CD15+ cells, cord blood CD71+ cells, cord blood CD34+ cells, neutrophils cord blood, neutrophils peripheral blood, spleen liver cirrhosis. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue spleen liver cirrhosis and healthy tissue spleen demonstrates that the human P2Y₁₂ or mRNA can be utilized to diagnose of hematological diseases. Additionally the activity of the human P2Y₁₂ can be modulated to treat hematological disorders.

Gastrointestinal and Liver Diseases

Gastrointestinal diseases comprise primary or secondary, acute or chronic diseases of the organs of the gastrointestinal tract which may be acquired or inherited, benign or malignant or metaplastic,

and which may affect the organs of the gastrointestinal tract or the body as a whole. They comprise but are not limited to 1) disorders of the esophagus like achalasia, vigorous achalasia, dysphagia, cricopharyngeal incoordination, pre-esophageal dysphagia, diffuse esophageal spasm, globus sensation, Barrett's metaplasia, gastroesophageal reflux, 2) disorders of the stomach and duodenum like functional dyspepsia, inflammation of the gastric mucosa, gastritis, stress gastritis, chronic erosive gastritis, atrophy of gastric glands, metaplasia of gastric tissues, gastric ulcers, duodenal ulcers, neoplasms of the stomach, 3) disorders of the pancreas like acute or chronic pancreatitis, insufficiency of the exocrine or endocrine tissues of the pancreas like steatorrhea, diabetes, neoplasms of the exocrine or endocrine pancreas like 3.1) multiple endocrine neoplasia syndrome, ductal adenocarcinoma, cystadenocarcinoma, islet cell tumors, insulinoma, gastrinoma, carcinoid tumors, glucagonoma, Zollinger-Ellison syndrome, Vipoma syndrome, malabsorption syndrome, 4) disorders of the bowel like chronic inflammatory diseases of the bowel, Crohn's disease, ileus, diarrhea and constipation, colonic inertia, megacolon, malabsorption syndrome, ulcerative colitis, 4.1) functional bowel disorders like irritable bowel syndrome, 4.2) neoplasms of the bowel like familial polyposis, adenocarcinoma, primary malignant lymphoma, carcinoid tumors, Kaposi's sarcoma, polyps, cancer of the colon and rectum.

Liver diseases comprise primary or secondary, acute or chronic diseases or injury of the liver which may be acquired or inherited, benign or malignant, and which may affect the liver or the body as a whole. They comprise but are not limited to disorders of the bilirubin metabolism, jaundice, syndromes of Gilbert's, Crigler-Najjar, Dubin-Johnson and Rotor; intrahepatic cholestasis, hepatomegaly, portal hypertension, ascites, Budd-Chiari syndrome, portal-systemic encephalopathy, fatty liver, steatosis, Reye's syndrome, liver diseases due to alcohol, alcoholic hepatitis or cirrhosis, fibrosis and cirrhosis, fibrosis and cirrhosis of the liver due to inborn errors of metabolism or exogenous substances, storage diseases, syndromes of Gaucher's, Zellweger's, Wilson's - disease, acute or chronic hepatitis, viral hepatitis and its variants, inflammatory conditions of the liver due to viruses, bacteria, fungi, protozoa, helminths; drug induced disorders of the liver, chronic liver diseases like primary sclerosing cholangitis, alpha₁-antitrypsin-deficiency, primary biliary cirrhosis, postoperative liver disorders like postoperative intrahepatic cholestasis, hepatic granulomas, vascular liver disorders associated with systemic disease, benign or malignant neoplasms of the liver, disturbance of liver metabolism in the new-born or prematurely born.

The human P2Y₁₂ is highly expressed in the following tissues of the gastroenterological system: esophagus, esophagus tumor, stomach, stomach tumor, small intestine, ileum, ileum tumor, rectum, salivary gland, liver, liver liver cirrhosis. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue liver liver cirrhosis and healthy tissue

liver demonstrates that the human P2Y₁₂ or mRNA can be utilized to diagnose of gastroenterological disorders. Additionally the activity of the human P2Y₁₂ can be modulated to treat gastroenterological disorders.

Endocrine System and Hormones

- 5 The endocrine system consists of a group of organs whose main function is to produce and secrete hormones directly into the bloodstream. The major organs of the endocrine system are the hypothalamus, the pituitary gland, thyroid gland, the parathyroid glands, the islets of the pancreas, the adrenal glands, the testes, and the ovaries.

- The hypothalamus secretes several hormones that stimulate the pituitary: Some trigger the release
10 of pituitary hormones; others suppress the release of pituitary hormones.

The pituitary gland coordinates many functions of the other endocrine glands, but some pituitary hormones have direct effects.

- The insulin-secreting cells of the pancreas respond to glucose and fatty acids. Parathyroid cells respond to calcium and phosphate. The adrenal medulla (part of the adrenal gland) responds to
15 direct stimulation by the parasympathetic nervous system.

When endocrine glands malfunction, hormone levels in the blood can become abnormally high or low, disrupting body functions. Many disorders are caused by malfunction of the endocrine system or hormones. Examples of such disorders are presented in the following.

- Diabetes mellitus is a disorder in which blood levels of glucose are abnormally high because the
20 body doesn't release or use insulin adequately.

People with type I diabetes mellitus (insulin-dependent diabetes) produce little or no insulin at all. In type I diabetes more than 90 percent of the insulin-producing cells (beta cells) of the pancreas are permanently destroyed. The resulting insulin deficiency is severe, and to survive, a person with type I diabetes must regularly inject insulin.

- 25 In type II diabetes mellitus (non-insulin-dependent diabetes) the body develops resistance to insulin effects, resulting in a relative insulin deficiency.

- The pancreas has two major functions: to secrete fluid containing digestive enzymes into the duodenum and to secrete the hormones insulin and glucagon. Chronic pancreatitis is a long-standing inflammation of the pancreas. Eventually, the insulin-secreting cells of the pancreas may
30 be destroyed, gradually leading to diabetes. An insulinoma is a rare type of pancreatic tumor that

- secretes insulin. The symptoms of an insulinoma result from low blood glucose levels. A gastrinoma is a pancreatic tumor that produces excessive levels of the hormone gastrin, which stimulates the stomach to secrete acid and enzymes, causing peptic ulcers. The excess gastrin secreted by the gastrinoma causes symptoms, called the Zollinger-Ellison syndrome. A
- 5 glucagonoma is a tumor that produces the hormone glucagon, which raises the level of glucose in the blood and produces a distinctive rash.

Diabetes insipidus is a disorder in which insufficient levels of antidiuretic hormone cause excessive thirst (polydipsia) and excessive production of very dilute urine (polyuria). Diabetes insipidus results from the decreased production of antidiuretic hormone (vasopressin).

- 10 The body has two adrenal glands. The medulla of the adrenal glands secretes hormones such as adrenaline (epinephrine) that affect blood pressure, heart rate, sweating, and other activities also regulated by the sympathetic nervous system. The cortex secretes many different hormones, including corticosteroids (cortisone-like hormones), androgens (male hormones), and mineralocorticoids, which control blood pressure and the levels of salt and potassium in the body.
- 15 A diseases characterized by underactive adrenal glands is Addison's disease (adrenocortical insufficiency).

Several disorders are characterized by overactive Adrenal Glands. The causes can be changes in the adrenal glands themselves or overstimulation by the pituitary gland. Examples of these diseases are listed in the following.

- 20 Overproduction of androgenic steroids (testosterone and similar hormones, leads to virilization), overproduction of corticosteroids (causes could be tumors of the pituitary or the adrenal gland, results in Cushing's syndrome), Nelson's syndrome (developed by people who have both adrenal glands removed, characterized by an enlargement of the pituitary gland), Overproduction of aldosterone (hyperaldosteronism), Conn's syndrome (hyperaldosterism caused by a tumor),
- 25 pheochromocytoma (a tumor that originating from the adrenal gland's chromaffin cells, causing overproduction of catecholamines),

- The thyroid is a small gland located under the Adam's apple. It secretes thyroid hormones, which control the metabolic rate. The thyroid gland traps iodine and processes it into thyroid hormones. The euthyroid sick syndrome is characterized by lack of conversion of the T4 form of thyroid
- 30 hormone to the T3 form. Hyperthyroidism (overactive thyroid gland, production of too much hormone) may have several causes. Thyroiditis (an inflammation of the thyroid gland), typically leads to a phase of hyperthyroidism. The inflammation may damage the thyroid gland, so that in

later stages the disease is characterized by transient or permanent underactivity (hypothyroidism). Toxic thyroid nodules (adenomas) often produce thyroid hormone in large quantities. Toxic multinodular goiter (Plummer's disease) is a disorder in which there are many nodules. Graves' disease (toxic diffuse goiter) is believed to be caused by an antibody that stimulates the thyroid to produce too much thyroid hormone. In toxic nodular goiter, one or more nodules in the thyroid produce too much thyroid hormone and aren't under the control of thyroid-stimulating hormone. Secondary hyperthyroidism may (rarely) be caused by a pituitary tumor that secretes too much thyroid-stimulating hormone, by resistance of the pituitary to thyroid hormone, which results in the pituitary gland secreting too much thyroid-stimulating hormone, or by a hydatidiform mole in women. Thyroid storm is a sudden extreme overactivity of the thyroid gland is a life-threatening emergency requiring prompt treatment.

Hypothyroidism is a condition in which the thyroid gland is underactive and produces too little thyroid hormone. Very severe hypothyroidism is called myxedema. In Hashimoto's thyroiditis (autoimmune thyroiditis) the thyroid gland is often enlarged, and hypothyroidism results because the gland's functioning areas are gradually destroyed. Rarer causes of hypothyroidism include some inherited disorders which are caused by abnormalities of the enzymes in thyroid cells. In other rare disorders, either the hypothalamus or the pituitary gland fails to secrete enough of the hormone needed to stimulate normal thyroid function.

Other examples of Thyroiditis are silent lymphocytic thyroiditis, Hashimoto's thyroiditis, or subacute granulomatous thyroiditis.

Thyroid cancer is any one of four main types of malignancy of the thyroid: papillary, follicular, anaplastic, or medullary.

The pituitary is a pea-sized gland that sits in a bony structure (sella turcica) at the base of the brain. The sella turcica protects the pituitary but allows very little room for expansion. If the pituitary enlarges, it tends to push upward, often pressing on the areas of the brain that carry signals from the eyes, possibly resulting in headaches or impaired vision. The pituitary gland has two distinct parts: the anterior (front) and the posterior (back) lobes. The anterior lobe produces (secretes) hormones that ultimately control the function of the thyroid gland, adrenal glands, and reproductive organs (ovaries and testes); milk production (lactation) in the breasts; and overall body growth. It also produces hormones that cause the skin to darken and that inhibit pain sensations. The posterior lobe produces hormones that regulate water balance, stimulate the let-down of milk from the breasts in lactating women, and stimulate contractions of the uterus.

Examples for disorders of the pituitary gland are Empty Sella Syndrome; hypopituitarism (an underactive pituitary gland); acromegaly, which is excessive growth caused by oversecretion of growth hormone, which is almost always caused by a benign pituitary tumor (adenoma); galactorrhea, which is the production of breast milk in men or in women who aren't breastfeeding, in both sexes, the most common cause of galactorrhea is a prolactin-producing tumor (prolactinoma) in the pituitary gland.

The human P2Y₁₂ is highly expressed in the following tissues of the endocrinological system: adrenal gland, pancreas. The expression in the above mentioned tissues demonstrates that the human P2Y₁₂ or mRNA can be utilized to diagnose of endocrinological disorders. Additionally the activity of the human P2Y₁₂ can be modulated to treat endocrinological disorders.

Cancer Disorders

Cancer disorders within the scope of this definition comprise any disease of an organ or tissue in mammals characterized by poorly controlled or uncontrolled multiplication of normal or abnormal cells in that tissue and its effect on the body as a whole. Cancer diseases within the scope of the definition comprise benign neoplasms, dysplasias, hyperplasias as well as neoplasms showing metastatic growth or any other transformations like e.g. leukoplakias which often precede a breakout of cancer. Cells and tissues are cancerous when they grow more rapidly than normal cells, displacing or spreading into the surrounding healthy tissue or any other tissues of the body described as metastatic growth, assume abnormal shapes and sizes, show changes in their nucleocytoplasmatic ratio, nuclear polychromasia, and finally may cease. Cancerous cells and tissues may affect the body as a whole when causing paraneoplastic syndromes or if cancer occurs within a vital organ or tissue, normal function will be impaired or halted, with possible fatal results. The ultimate involvement of a vital organ by cancer, either primary or metastatic, may lead to the death of the mammal affected. Cancer tends to spread, and the extent of its spread is usually related to an individual's chances of surviving the disease. Cancers are generally said to be in one of three stages of growth: early, or localized, when a tumor is still confined to the tissue of origin, or primary site; direct extension, where cancer cells from the tumour have invaded adjacent tissue or have spread only to regional lymph nodes; or metastasis, in which cancer cells have migrated to distant parts of the body from the primary site, via the blood or lymph systems, and have established secondary sites of infection. Cancer is said to be malignant because of its tendency to cause death if not treated. Benign tumors usually do not cause death, although they may if they interfere with a normal body function by virtue of their location, size, or paraneoplastic side effects. Hence benign tumors fall under the definition of cancer within the scope of this definition as well. In general, cancer cells divide at a higher rate than do normal cells, but the distinction

between the growth of cancerous and normal tissues is not so much the rapidity of cell division in the former as it is the partial or complete loss of growth restraint in cancer cells and their failure to differentiate into a useful, limited tissue of the type that characterizes the functional equilibrium of growth of normal tissue. Cancer tissues may express certain molecular receptors and probably are
5 influenced by the host's susceptibility and immunity and it is known that certain cancers of the breast and prostate, for example, are considered dependent on specific hormones for their existence. The term "cancer" under the scope of the definition is not limited to simple benign neoplasia but comprises any other benign and malign neoplasia like 1) Carcinoma, 2) Sarcoma, 3) Carcinosarcoma, 4) Cancers of the blood-forming tissues, 5) tumors of nerve tissues including the
10 brain, 6) cancer of skin cells. Cancer according to 1) occurs in epithelial tissues, which cover the outer body (the skin) and line mucous membranes and the inner cavitory structures of organs e.g. such as the breast, lung, the respiratory and gastrointestinal tracts, the endocrine glands, and the genitourinary system. Ductal or glandular elements may persist in epithelial tumors, as in adenocarcinomas like e.g. thyroid adenocarcinoma, gastric adenocarcinoma, uterine adeno-
15 carcinoma. Cancers of the pavement-cell epithelium of the skin and of certain mucous membranes, such as e.g. cancers of the tongue, lip, larynx, urinary bladder, uterine cervix, or penis, may be termed epidermoid or squamous-cell carcinomas of the respective tissues and are in the scope of the definition of cancer as well. Cancer according to 2) develops in connective tissues, including fibrous tissues, adipose (fat) tissues, muscle, blood vessels, bone, and cartilage like e.g. osteogenic
20 sarcoma; liposarcoma, fibrosarcoma, synovial sarcoma. Cancer according to 3) is cancer that develops in both epithelial and connective tissue. Cancer disease within the scope of this definition may be primary or secondary, whereby primary indicates that the cancer originated in the tissue where it is found rather than was established as a secondary site through metastasis from another lesion. Cancers and tumor diseases within the scope of this definition may be benign or malign and
25 may affect all anatomical structures of the body of a mammal. By example but not limited to they comprise cancers and tumor diseases of I) the bone marrow and bone marrow derived cells (leukemias), II) the endocrine and exocrine glands like e.g. thyroid, parathyroid, pituitary, adrenal glands, salivary glands, pancreas III) the breast, like e.g. benign or malignant tumors in the mammary glands of either a male or a female, the mammary ducts, adenocarcinoma, medullary
30 carcinoma, comedo carcinoma, Paget's disease of the nipple, inflammatory carcinoma of the young woman, IV) the lung, V) the stomach, VI) the liver and spleen, VII) the small intestine, VIII) the colon, IX) the bone and its supportive and connective tissues like malignant or benign bone tumour, e.g. malignant osteogenic sarcoma, benign osteoma, cartilage tumors; like malignant chondrosarcoma or benign chondroma; bone marrow tumors like malignant myeloma or benign
35 eosinophilic granuloma, as well as metastatic tumors from bone tissues at other locations of the body; X) the mouth, throat, larynx, and the esophagus, XI) the urinary bladder and the internal and

external organs and structures of the urogenital system of male and female like ovaries, uterus, cervix of the uterus, testes, and prostate gland, XII) the prostate, XIII) the pancreas, like ductal carcinoma of the pancreas; XIV) the lymphatic tissue like lymphomas and other tumors of lymphoid origin, XV) the skin, XVI) cancers and tumor diseases of all anatomical structures
5 belonging to the respiration and respiratory systems including thoracic muscles and linings, XVII) primary or secondary cancer of the lymph nodes XVIII) the tongue and of the bony structures of the hard palate or sinuses, XIX) the mouth, cheeks, neck and salivary glands, XX) the blood vessels including the heart and their linings, XXI) the smooth or skeletal muscles and their ligaments and linings, XXII) the peripheral, the autonomous, the central nervous system including
10 the cerebellum, XXIII) the adipose tissue.

The human P2Y₁₂ is highly expressed in the following cancer tissues: esophagus tumor, stomach tumor, ileum tumor, Jurkat (T-cells), ovary tumor, breast tumor, prostate tumor. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue esophagus tumor and healthy tissue esophagus, between diseased tissue stomach tumor and healthy
15 tissue stomach, between diseased tissue ileum tumor and healthy tissue ileum, between diseased tissue Jurkat (T-cells) and healthy tissue, between diseased tissue ovary tumor and healthy tissue ovary, between diseased tissue breast tumor and healthy tissue breast, between diseased tissue prostate tumor and healthy tissue prostate demonstrates that the human P2Y₁₂ or mRNA can be utilized to diagnose of cancer. Additionally the activity of the human P2Y₁₂ can be modulated to
20 treat cancer.

Inflammatory Diseases

Inflammatory diseases comprise diseases triggered by cellular or non-cellular mediators of the immune system or tissues causing the inflammation of body tissues and subsequently producing an acute or chronic inflammatory condition. Examples for such inflammatory diseases are
25 hypersensitivity reactions of type I – IV, for example but not limited to hypersensitivity diseases of the lung including asthma, atopic diseases, allergic rhinitis or conjunctivitis, angioedema of the lids, hereditary angioedema, antireceptor hypersensitivity reactions and autoimmune diseases, Hashimoto's thyroiditis, systemic lupus erythematosus, Goodpasture's syndrome, pemphigus, myasthenia gravis, Grave's and Raynaud's disease, type B insulin-resistant diabetes, rheumatoid
30 arthritis, psoriasis, Crohn's disease, scleroderma, mixed connective tissue disease, polymyositis, sarcoidosis, glomerulonephritis, acute or chronic host versus graft reactions.

The human P2Y₁₂ is highly expressed in the following tissues of the immune system and tissues responsive to components of the immune system as well as in the following tissues responsive to mediators of inflammation: liver liver cirrhosis, leukocytes (peripheral blood), bone marrow, bone

marrow CD15+ cells, neutrophils cord blood, neutrophils peripheral blood, spleen liver cirrhosis. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue liver liver cirrhosis and healthy tissue liver, between diseased tissue spleen liver cirrhosis and healthy tissue spleen demonstrates that the human P2Y12 or mRNA can be
5 utilized to diagnose of inflammatory diseases. Additionally the activity of the human P2Y12 can be modulated to treat inflammatory diseases.

Disorders Related to Pulmology

Asthma is thought to arise as a result of interactions between multiple genetic and environmental factors and is characterized by three major features: 1) intermittent and reversible airway
10 obstruction caused by bronchoconstriction, increased mucus production, and thickening of the walls of the airways that leads to a narrowing of the airways, 2) airway hyperresponsiveness, and 3) airway inflammation. Certain cells are critical to the inflammatory reaction of asthma and they include T cells and antigen presenting cells, B cells that produce IgE, and mast cells, basophils, eosinophils, and other cells that bind IgE. These effector cells accumulate at the site of allergic
15 reaction in the airways and release toxic products that contribute to the acute pathology and eventually to tissue destruction related to the disorder. Other resident cells, such as smooth muscle cells, lung epithelial cells, mucus-producing cells, and nerve cells may also be abnormal in individuals with asthma and may contribute to its pathology. While the airway obstruction of
20 asthma, presenting clinically as an intermittent wheeze and shortness of breath, is generally the most pressing symptom of the disease requiring immediate treatment, the inflammation and tissue destruction associated with the disease can lead to irreversible changes that eventually make asthma a chronic and disabling disorder requiring long-term management.

Chronic obstructive pulmonary (or airways) disease (COPD) is a condition defined physiologically as airflow obstruction that generally results from a mixture of emphysema and peripheral airway
25 obstruction due to chronic bronchitis [Botstein, 1980]. Emphysema is characterised by destruction of alveolar walls leading to abnormal enlargement of the air spaces of the lung. Chronic bronchitis is defined clinically as the presence of chronic productive cough for three months in each of two successive years. In COPD, airflow obstruction is usually progressive and is only partially reversible. By far the most important risk factor for development of COPD is cigarette smoking,
30 although the disease does also occur in non-smokers.

The human P2Y12 is highly expressed in the following tissues of the respiratory system: leukocytes (peripheral blood), bone marrow CD15+ cells, neutrophils cord blood, neutrophils peripheral blood, fetal lung, lung right upper lobe, lung right mid lobe, lung right lower lobe, trachea, primary bronchia, secondary bronchia, bronchial epithelial cells, bronchial smooth muscle

cells. The expression in the above mentioned tissues demonstrates that the human P2Y₁₂ or mRNA can be utilized to diagnose of respiratory diseases. Additionally the activity of the human P2Y₁₂ can be modulated to treat those diseases.

Disorders Related to Urology

- 5 Genitourinary disorders comprise benign and malign disorders of the organs constituting the genitourinary system of female and male, renal diseases like acute or chronic renal failure, immunologically mediated renal diseases like renal transplant rejection, lupus nephritis, immune complex renal diseases, glomerulopathies, nephritis, toxic nephropathy, obstructive uropathies like benign prostatic hyperplasia (BPH), neurogenic bladder syndrome, urinary incontinence like urge-,
10 stress-, or overflow incontinence, pelvic pain, and erectile dysfunction.

The human P2Y₁₂ is highly expressed in the following urological tissues: spinal cord, spinal cord (ventral horn), spinal cord (dorsal horn), prostate, prostate BPH, prostate tumor, bladder, ureter, penis, corpus cavernosum. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue prostate BPH and healthy tissue prostate
15 demonstrates that the human P2Y₁₂ or mRNA can be utilized to diagnose of urological disorders. Additionally the activity of the human P2Y₁₂ can be modulated to treat urological disorders.

The human P2Y₁₂ is highly expressed in spinal cord tissues: spinal cord, spinal cord (ventral horn), spinal cord (dorsal horn). Expression in spinal cord tissues demonstrates that the human P2Y₁₂ or mRNA can be utilized to diagnose of incontinence as an urological disorder. The spinal
20 cord tissues are involved in the neuronal regulation of the urological system. Additionally the activity of the human P2Y₁₂ can be modulated to treat - but not limited to - incontinence.

Metabolic Disorders

Metabolic diseases are defined as conditions which result from an abnormality in any of the chemical or biochemical transformations and their regulating systems essential to producing
25 energy, to regenerating cellular constituents, to eliminating unneeded products arising from these processes, and to regulate and maintain homeostasis in a mammal regardless of whether acquired or the result of a genetic transformation. Depending on which metabolic pathway is involved, a single defective transformation or disturbance of its regulation may produce consequences that are narrow, involving a single body function, or broad, affecting many organs, organ-systems or the
30 body as a whole. Diseases resulting from abnormalities related to the fine and coarse mechanisms that affect each individual transformation, its rate and direction or the availability of substrates like amino acids, fatty acids, carbohydrates, minerals, cofactors, hormones, regardless whether they are

inborn or acquired, are well within the scope of the definition of a metabolic disease according to this application.

Metabolic diseases often are caused by single defects in particular biochemical pathways, defects that are due to the deficient activity of individual enzymes or molecular receptors leading to the regulation of such enzymes. Hence in a broader sense disturbances of the underlying genes, their products and their regulation lie well within the scope of this definition of a metabolic disease. For example, but not limited to, metabolic diseases may affect 1) biochemical processes and tissues ubiquitous all over the body, 2) the bone, 3) the nervous system, 4) the endocrine system, 5) the muscle including the heart, 6) the skin and nervous tissue, 7) the urogenital system, 8) the homeostasis of body systems like water and electrolytes. For example, but not limited to, metabolic diseases according to 1) comprise obesity, amyloidosis, disturbances of the amino acid metabolism like branched chain disease, hyperaminoacidemia, hyperaminoaciduria, disturbances of the metabolism of urea, hyperammonemia, mucopolysaccharidoses e.g. Maroteaux-Lamy syndrom, storage diseases like glycogen storage diseases and lipid storage diseases, glycogenosis diseases like Cori's disease, malabsorption diseases like intestinal carbohydrate malabsorption, oligosaccharidase deficiency like maltase-, lactase-, sucrase-insufficiency, disorders of the metabolism of fructose, disorders of the metabolism of galactose, galactosaemia, disturbances of carbohydrate utilization like diabetes, hypoglycemia, disturbances of pyruvate metabolism, hypolipidemia, hypolipoproteinemia, hyperlipidemia, hyperlipoproteinemia, carnitine or carnitine acyltransferase deficiency, disturbances of the porphyrin metabolism, porphyrias, disturbances of the purine metabolism, lysosomal diseases, metabolic diseases of nerves and nervous systems like gangliosidoses, sphingolipidoses, sulfatidoses, leucodystrophies, Lesch-Nyhan syndrome. For example, but not limited to, metabolic diseases according to 2) comprise osteoporosis, osteomalacia like osteoporosis, osteopenia, osteogenesis imperfecta, osteopetrosis, osteonecrosis, Paget's disease of bone, hypophosphatemia. For example, but not limited to, metabolic diseases according to 3) comprise cerebellar dysfunction, disturbances of brain metabolism like dementia, Alzheimer's disease, Huntington's chorea, Parkinson's disease, Pick's disease, toxic encephalopathy, demyelinating neuropathies like inflammatory neuropathy, Guillain-Barré syndrome. For example, but not limited to, metabolic diseases according to 4) comprise primary and secondary metabolic disorders associated with hormonal defects like any disorder stemming from either an hyperfunction or hypofunction of some hormone-secreting endocrine gland and any combination thereof. They comprise Sipple's syndrome, pituitary gland dysfunction and its effects on other endocrine glands, such as the thyroid, adrenals, ovaries, and testes, acromegaly, hyper- and hypothyroidism, euthyroid goiter, euthyroid sick syndrome, thyroiditis, and thyroid cancer, over- or underproduction of the adrenal steroid hormones, adrenogenital syndrome, Cushing's

syndrome, Addison's disease of the adrenal cortex, Addison's pernicious anemia, primary and secondary aldosteronism, diabetes insipidus, carcinoid syndrome, disturbances caused by the dysfunction of the parathyroid glands, pancreatic islet cell dysfunction, diabetes, disturbances of the endocrine system of the female like estrogen deficiency, resistant ovary syndrome. For example, but not limited to, metabolic diseases according to 5) comprise muscle weakness, myotonia, Duchenne's and other muscular dystrophies, dystrophia myotonica of Steinert, mitochondrial myopathies like disturbances of the catabolic metabolism in the muscle, carbohydrate and lipid storage myopathies, glycogenoses, myoglobinuria, malignant hyperthermia, polymyalgia rheumatica, dermatomyositis, primary myocardial disease, cardiomyopathy. For example, but not limited to, metabolic diseases according to 6) comprise disorders of the ectoderm, neurofibromatosis, scleroderma and polyarteritis, Louis-Bar syndrome, von Hippel-Lindau disease, Sturge-Weber syndrome, tuberous sclerosis, amyloidosis, porphyria. For example, but not limited to, metabolic diseases according to 7) comprise sexual dysfunction of the male and female. For example, but not limited to, metabolic diseases according to 8) comprise confused states and seizures due to inappropriate secretion of antidiuretic hormone from the pituitary gland, Liddle's syndrome, Bartter's syndrome, Fanconi's syndrome, renal electrolyte wasting, diabetes insipidus.

The human P2Y₁₂ is highly expressed in the following metabolic disease related tissues: pancreas, liver, liver liver cirrhosis, spleen liver cirrhosis. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue liver liver cirrhosis and healthy tissue liver, between diseased tissue spleen liver cirrhosis and healthy tissue spleen demonstrates that the human P2Y₁₂ or mRNA can be utilized to diagnose of metabolic diseases. Additionally the activity of the human P2Y₁₂ can be modulated to treat metabolic diseases.

Applications

The present invention provides for both prophylactic and therapeutic methods for cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases.

The regulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of P2Y₁₂. An agent that modulates activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of the polypeptide, a peptide, a peptidomimetic, or any small molecule. In one embodiment, the agent stimulates one or more of the biological activities of P2Y₁₂. Examples of such stimulatory agents include the active P2Y₁₂ and nucleic acid molecules encoding a portion of P2Y₁₂. In another embodiment, the agent inhibits one or more of the biological activities of P2Y₁₂. Examples of such inhibitory

agents include antisense nucleic acid molecules and antibodies. These regulatory methods can be performed in vitro (e.g., by culturing the cell with the agent) or, alternatively, in vivo (e.g, by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by unwanted expression or activity of P2Y12 or a protein in the P2Y12 signaling pathway. In one embodiment, the method involves administering an agent like any agent identified or being identifiable by a screening assay as described herein, or combination of such agents that modulate say upregulate or downregulate the expression or activity of P2Y12 or of any protein in the P2Y12 signaling pathway. In another embodiment, the method involves administering a regulator of P2Y12 as therapy to compensate for reduced or undesirably low expression or activity of P2Y12 or a protein in the P2Y12 signaling pathway.

Stimulation of activity or expression of P2Y12 is desirable in situations in which activity or expression is abnormally low and in which increased activity is likely to have a beneficial effect. Conversely, inhibition of activity or expression of P2Y12 is desirable in situations in which activity or expression of P2Y12 is abnormally high and in which decreasing its activity is likely to have a beneficial effect.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are hereby incorporated by reference.

20 **Pharmaceutical Compositions**

This invention further pertains to novel agents identified by the above-described screening assays and uses thereof for treatments as described herein.

The nucleic acid molecules, polypeptides, and antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

The invention includes pharmaceutical compositions comprising a regulator of P2Y₁₂ expression or activity (and/or a regulator of the activity or expression of a protein in the P2Y₁₂ signaling pathway) as well as methods for preparing such compositions by combining one or more such regulators and a pharmaceutically acceptable carrier. Also within the invention are pharmaceutical compositions comprising a regulator identified using the screening assays of the invention packaged with instructions for use. For regulators that are antagonists of P2Y₁₂ activity or which reduce P2Y₁₂ expression, the instructions would specify use of the pharmaceutical composition for treatment of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases. For regulators that are agonists of P2Y₁₂ activity or increase P2Y₁₂ expression, the instructions would specify use of the pharmaceutical composition for treatment of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases.

An antagonist of P2Y₁₂ may be produced using methods which are generally known in the art. In particular, purified P2Y₁₂ may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind P2Y₁₂. Antibodies to P2Y₁₂ may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies like those which inhibit dimer formation are especially preferred for therapeutic use.

In another embodiment of the invention, the polynucleotides encoding P2Y₁₂, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding P2Y₁₂ may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding P2Y₁₂. Thus, complementary molecules or fragments may be used to modulate P2Y₁₂ activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding P2Y₁₂.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors which will express nucleic acid sequence complementary to the polynucleotides

of the gene encoding P2Y₁₂. These techniques are described, for example, in [Scott and Smith (1990) Science 249:386-390].

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

An additional embodiment of the invention relates to the administration of a pharmaceutical composition containing P2Y₁₂ in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of P2Y₁₂, antibodies to P2Y₁₂, and mimetics, agonists, antagonists, or inhibitors of P2Y₁₂. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EMTM (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, a pharmaceutically acceptable polyol like glycerol, propylene glycol, liquid

polyethylene glycol, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin. Sterile injectable solutions can be prepared by incorporating the active compound (e.g., a polypeptide or antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed.

Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from a pressurized container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal
5 administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal
10 delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be
15 apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those
20 skilled in the art, for example, as described in U.S. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired
25 therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

30 The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration. For pharmaceutical compositions which include an antagonist of P2Y₁₂ activity, a compound which reduces expression of P2Y₁₂, or a compound which reduces expression or activity of a protein in the P2Y₁₂ signaling pathway or any combination thereof, the instructions for administration will specify use of the composition for cardiovascular diseases,

endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases. For pharmaceutical compositions which include an agonist of P2Y₁₂ activity, a compound which increases expression of P2Y₁₂, or a compound which increases expression or activity of a protein in the P2Y₁₂ signaling pathway or any combination thereof, the instructions for administration will specify use of the composition for cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases.

Diagnostics

In another embodiment, antibodies which specifically bind P2Y₁₂ may be used for the diagnosis of disorders characterized by the expression of P2Y₁₂, or in assays to monitor patients being treated with P2Y₁₂ or agonists, antagonists, and inhibitors of P2Y₁₂. Antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for therapeutics. Diagnostic assays for P2Y₁₂ include methods which utilize the antibody and a label to detect P2Y₁₂ in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent joining with a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring P2Y₁₂, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of P2Y₁₂ expression. Normal or standard values for P2Y₁₂ expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to P2Y₁₂ under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, preferably by photometric means. Quantities of P2Y₁₂ expressed in subject samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding P2Y₁₂ may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of P2Y₁₂ may be correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess expression of P2Y₁₂, and to monitor regulation of P2Y₁₂ levels during therapeutic intervention.

Polynucleotide sequences encoding P2Y₁₂ may be used for the diagnosis of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases associated with expression of P2Y₁₂. The polynucleotide sequences encoding P2Y₁₂ may be used in Southern, Northern, or dot-blot analysis, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and ELISA assays; and in microarrays utilizing fluids or tissues from patient biopsies to detect altered P2Y₁₂ expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding P2Y₁₂ may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding P2Y₁₂ may be labelled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the patient sample is significantly altered from that of a comparable control sample, the nucleotide sequences have hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding P2Y₁₂ in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or in monitoring the treatment of an individual patient.

In order to provide a basis for the diagnosis of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases associated with expression of P2Y₁₂, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding P2Y₁₂, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained from normal samples may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO84/03564. In this method, large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test

compounds are reacted with P2Y₁₂, or fragments thereof, and washed. Bound P2Y₁₂ is then detected by methods well known in the art. Purified P2Y₁₂ can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

- 5 In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding P2Y₁₂ specifically compete with a test compound for binding P2Y₁₂. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with P2Y₁₂.

- 10 G-protein coupled receptors are ubiquitous in the mammalian host and are responsible for many biological functions, including many pathologies. Accordingly, it is desirable to find compounds and drugs which stimulate a G-protein coupled receptor on the one hand and which can inhibit the function of a G-protein coupled receptor on the other hand. For example, compounds which activate the G-protein coupled receptor may be employed for therapeutic purposes, such as the treatment of asthma, Parkinson's disease, acute heart failure, urinary retention, and osteoporosis.
- 15 In particular, compounds which activate the receptors of the present invention are useful in treating various cardiovascular ailments such as caused by the lack of pulmonary blood flow or hypertension. In addition these compounds may also be used in treating various physiological disorders relating to abnormal control of fluid and electrolyte homeostasis and in diseases associated with abnormal angiotensin-induced aldosterone secretion.

- 20 In general, compounds which inhibit activation of the G-protein coupled receptor may be employed for a variety of therapeutic purposes, for example, for the treatment of hypotension and/or hypertension, angina pectoris, myocardial infarction, ulcers, asthma, allergies, benign prostatic hypertrophy, and psychotic and neurological disorders including schizophrenia, manic excitement, depression, delirium, dementia or severe mental retardation, dyskinesias, such as
- 25 Huntington's disease or Tourett's syndrome, among others. Compounds which inhibit G-protein coupled receptors have also been useful in reversing endogenous anorexia and in the control of bulimia.

Determination of a Therapeutically Effective Dose

- 30 The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases P2Y₁₂ activity relative to P2Y₁₂ activity which occurs in the absence of the therapeutically effective dose. For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or

pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

Therapeutic efficacy and toxicity, e.g., ED₅₀ (the dose therapeutically effective in 50% of the population) and LD₅₀ (the dose lethal to 50% of the population), can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD₅₀/ED₅₀. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration. The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect. Factors which can be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts can vary from 0.1 micrograms to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc. If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either *ex vivo* or *in vivo* using well-established techniques including, but not limited to, transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, "gene gun", and DEAE- or calcium phosphate-mediated transfection.

If the expression product is mRNA, the reagent is preferably an antisense oligonucleotide or a ribozyme. Polynucleotides which express antisense oligonucleotides or ribozymes can be introduced into cells by a variety of methods, as described above. Preferably, a reagent reduces

expression of P2Y12 gene or the activity of P2Y12 by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism chosen to decrease the level of expression of P2Y12 gene or the activity of P2Y12 can be assessed using methods well known in the art, such as hybridization of nucleotide probes to
5 P2Y12-specific mRNA, quantitative RT-PCR, immunologic detection of P2Y12, or measurement of P2Y12 activity.

In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy can be made by one of ordinary skill in
10 the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects. Any of the therapeutic methods described above can be applied to any subject in need of such therapy, including, for
15 example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

Nucleic acid molecules of the invention are those nucleic acid molecules which are contained in a group of nucleic acid molecules consisting of (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2, (ii) nucleic acid molecules comprising the
20 sequence of SEQ ID NO: 1, (iii) nucleic acid molecules having the sequence of SEQ ID NO: 1, (iv) nucleic acid molecules the complementary strand of which hybridizes under stringent conditions to a nucleic acid molecule of (i), (ii), or (iii); and (v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code, wherein the polypeptide encoded by said nucleic acid molecule
25 has P2Y12 activity.

Polypeptides of the invention are those polypeptides which are contained in a group of polypeptides consisting of (i) polypeptides having the sequence of SEQ ID NO: 2, (ii) polypeptides comprising the sequence of SEQ ID NO: 2, (iii) polypeptides encoded by nucleic acid molecules of the invention and (iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% homology
30 with a polypeptide of (i), (ii), or (iii), wherein said purified polypeptide has P2Y12 activity.

An object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a

mammal comprising the steps of (i) contacting a test compound with a P2Y₁₂ polypeptide, (ii) detect binding of said test compound to said P2Y₁₂ polypeptide. E.g., compounds that bind to the P2Y₁₂ polypeptide are identified potential therapeutic agents for such a disease.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) determining the activity of a P2Y₁₂ polypeptide at a certain concentration of a test compound or in the absence of said test compound, (ii) determining the activity of said polypeptide at a different concentration of said test compound. E.g., compounds that lead to a difference in the activity of the P2Y₁₂ polypeptide in (i) and (ii) are identified potential therapeutic agents for such a disease.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) determining the activity of a P2Y₁₂ polypeptide at a certain concentration of a test compound, (ii) determining the activity of a P2Y₁₂ polypeptide at the presence of a compound known to be a regulator of a P2Y₁₂ polypeptide. E.g., compounds that show similar effects on the activity of the P2Y₁₂ polypeptide in (i) as compared to compounds used in (ii) are identified potential therapeutic agents for such a disease.

Other objects of the invention are methods of the above, wherein the step of contacting is in or at the surface of a cell.

Other objects of the invention are methods of the above, wherein the cell is in vitro.

Other objects of the invention are methods of the above, wherein the step of contacting is in a cell-free system.

Other objects of the invention are methods of the above, wherein the polypeptide is coupled to a detectable label.

Other objects of the invention are methods of the above, wherein the compound is coupled to a detectable label.

Other objects of the invention are methods of the above, wherein the test compound displaces a ligand which is first bound to the polypeptide.

Other objects of the invention are methods of the above, wherein the polypeptide is attached to a solid support.

- 5 Other objects of the invention are methods of the above, wherein the compound is attached to a solid support.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) contacting a test compound with a P2Y₁₂ polynucleotide, (ii) detect binding of said test compound to said P2Y₁₂ polynucleotide. Compounds that, e.g., bind to the P2Y₁₂ polynucleotide are potential therapeutic agents for the treatment of such diseases.

15 Another object of the invention is the method of the above, wherein the nucleic acid molecule is RNA.

Another object of the invention is a method of the above, wherein the contacting step is in or at the surface of a cell.

Another object of the invention is a method of the above, wherein the contacting step is in a cell-free system.

- 20 Another object of the invention is a method of the above, wherein the polynucleotide is coupled to a detectable label.

Another object of the invention is a method of the above, wherein the test compound is coupled to a detectable label.

25 Another object of the invention is a method of diagnosing a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) determining the amount of a P2Y₁₂ polynucleotide in a sample taken from said mammal, (ii) determining the amount of P2Y₁₂ polynucleotide in healthy and/or diseased mammal. A disease is diagnosed, e.g., if there is a substantial similarity in the amount of P2Y₁₂ polynucleotide in said test mammal as compared to a diseased mammal.

30

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a
5 therapeutic agent which binds to a P2Y₁₂ polypeptide.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a
10 therapeutic agent which regulates the activity of a P2Y₁₂ polypeptide.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a
15 therapeutic agent which regulates the activity of a P2Y₁₂ polypeptide, wherein said therapeutic agent is (i) a small molecule, (ii) an RNA molecule, (iii) an antisense oligonucleotide, (iv) a polypeptide, (v) an antibody, or (vi) a ribozyme.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a
20 P2Y₁₂ polynucleotide.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a
25 P2Y₁₂ polypeptide.

Another object of the invention is the use of regulators of a P2Y₁₂ for the preparation of a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal.
30

Another object of the invention is a method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) identifying a regulator of P2Y₁₂, (ii) determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal; and (iii) combining of said regulator with an acceptable pharmaceutical carrier.

Another object of the invention is the use of a regulator of P2Y₁₂ for the regulation of P2Y₁₂ activity in a mammal having a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases.

The expression of human p2y₁₂ in cardiovascular and hematological related tissues (as described above) suggests a particular, but not limited to, utilization of p2y₁₂ for diagnosis and modulation of cardiovascular diseases and hematological diseases. Furthermore the above described expression suggest a, but not limited to utilization of p2y₁₂ to endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, respiratory diseases, neurological diseases and urological diseases.

The examples below are provided to illustrate the subject invention. These examples are provided by way of illustration and are not included for the purpose of limiting the invention.

Examples

Example 1: Search for homologous sequences in public sequence data bases

The degree of homology can readily be calculated by known methods. Preferred methods to determine homology are designed to give the largest match between the sequences tested.

- 5 Methods to determine homology are codified in publicly available computer programs such as BestFit, BLASTP, BLASTN, and FASTA. The BLAST programs are publicly available from NCBI and other sources in the internet.

- For P2Y12 the following hits to known sequences were identified by using the BLAST algorithm [Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ; Nucleic Acids
10 Res 1997 Sep 1; 25(17): 3389-402] and the following set of parameters: matrix = BLOSUM62 and low complexity filter. The following databases were searched: NCBI (non-redundant database) and DERWENT patent database (Geneseq).

The following hits were found:

- >AA2003A:ADA18543 Ada18543 Novel human GPCR, 15625 receptor. 11/2003
15 Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)

- >AA2003A:ABP81734 Abp81734 Human P2Y12 platelet ADP receptor protein SEQ ID NO:643.
3/2003
20 Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)

- >AA2003A:ABU92274 Abu92274 Human G protein-coupled receptor hCHN8. 7/2003
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives =
25 342/342 (100%)

- >AA2002:AAG80236 Aag80236 Human PAFR3 protein. 1/2002
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives =
30 342/342 (100%)

- >AA2002:ABP95608 Abp95608 Human GPCR polypeptide SEQ ID NO 26. 3/2003
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)
- 5 >AA2002:AAU80164 Aau80164 Human ADP-glucose receptor. 7/2002
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)
- >AA2002:ABB05031 Abb05031 Human SP168 receptor protein SEQ ID NO:2. 3/2002
10 Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)
- >AA2002:ABB77868 Abb77868 Amino acid sequence of human ADP receptor P2TAC protein. 9/2002
15 Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)
- >AA2002:AAM48353 Aam48353 Human G protein-coupled receptor, IGPCr17. 4/2002
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)
20
- >AA2001:AAM79249 Aam79249 Human protein SEQ ID NO 1911. 11/2001
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)
25
- >AA2001:AAE04386 Aae04386 Human P2-purinergic receptor subtype, P2Y12. 9/2001
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)
- 30 >AA2000:AAY71306 Aay71306 Human orphan G protein-coupled receptor hCHN8. 11/2000
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)

- >AA2000:AA94444 Aay94444 Human 15625 receptor protein. 8/2000
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)
- 5 >AA2000:AAB02840 Aab02840 Human G protein coupled receptor hCHN8 protein SEQ ID NO:34. 8/2000
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)
- 10 >AA1999:AAW81576 Aaw81576 EBV-induced G-protein coupled receptor (EBI-2) polypeptide. 2/1999
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)
- 15 >emb|CAD23844.1| unnamed protein product [Homo sapiens] emb|CAD35232.1| unnamed protein product [Homo sapiens]
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)
- 20 >ref|NP_073625.1| purinergic receptor P2Y12; platelet ADP receptor; purinergic receptor P2RY12; purinergic receptor P2Y, G-protein coupled 12; G-protein coupled receptor SP1999; Gi-coupled ADP receptor HORK3; putative G-protein coupled receptor; P2Y purinoceptor 12; ADP-glucose receptor [Homo sapiens] ref|NP_795345.1| purinergic receptor P2Y12; platelet ADP receptor; purinergic receptor P2RY12; purinergic receptor P2Y, G-protein coupled 12; G-protein coupled receptor SP1999; Gi-coupled ADP receptor HORK3; putative G-protein coupled receptor; P2Y purinoceptor 12; ADP-glucose receptor [Homo sapiens] sp|Q9H244|P2YC_HUMAN P2Y purinoceptor 12 (P2Y12) (P2Y12 platelet ADP receptor) (P2Y(ADP)) (ADP-glucose receptor) (ADPG-R) (P2Y(AC)) (P2Y(cyc)) (P2T(AC)) (SP1999) gb|AAG48944.1| P2Y12 platelet ADP receptor [Homo sapiens] gb|AAK00948.1| G-protein coupled receptor SP1999 [Homo sapiens]
- 25 dbj|BAB60824.1| Gi-coupled ADP receptor HORK3 [Homo sapiens] gb|AAL32292.1| ADP-glucose receptor [Homo sapiens] gb|AAH17898.1| Purinergic receptor P2Y12 [Homo sapiens] dbj|BAB89309.1| putative G-protein coupled receptor [Homo sapiens] emb|CAC87144.1| platelet ADP-receptor [Homo sapiens] gb|AAN01280.1| purinergic receptor P2RY12 [Homo sapiens]
Length = 342, Score = 689 bits (1778), Expect = 0.0, Identities = 342/342 (100%), Positives = 342/342 (100%)
- 35

- >AA2003A:ADA18545 Ada18545 Novel nonhuman primate GPCR, 15625 receptor. 11/2003
Length = 342, Score = 677 bits (1748), Expect = 0.0, Identities = 335/342 (97%), Positives = 338/342 (98%)
- 5 >AA2000:AA94445 Aay94445 Macaque ortholog of human 15625 receptor protein. 8/2000
Length = 342, Score = 677 bits (1748), Expect = 0.0, Identities = 335/342 (97%), Positives = 338/342 (98%)
- >sp|Q95KC3|P2YC_MACFA P2Y purinoceptor 12 (P2Y12) dbj|BAB60747.1| hypothetical protein
10 [Macaca fascicularis]
Length = 342, Score = 677 bits (1748), Expect = 0.0, Identities = 335/342 (97%), Positives = 338/342 (98%)
- >dbj|BAB33041.1| hypothetical protein [Macaca fascicularis]
15 Length = 342, Score = 677 bits (1748), Expect = 0.0, Identities = 335/342 (97%), Positives = 338/342 (98%)
- >dbj|BAC41779.1| hypothetical protein [Macaca fascicularis]
Length = 342, Score = 675 bits (1741), Expect = 0.0, Identities = 334/342 (97%), Positives = 337/342 (98%)
20
- >AA2001:AAE04385 Aae04385 Human P2-purinergic receptor subtype, P2Y12 protein fragment.
9/2001
Length = 315, Score = 634 bits (1634), Expect = e-180, Identities = 315/315 (100%), Positives = 315/315 (100%)
25
- >AA2003A:ABP70855 Abp70855 Murine P2T receptor. 8/2003
Length = 347, Score = 603 bits (1554), Expect = e-171, Identities = 299/337 (88%), Positives = 315/337 (93%), Gaps = 1/337 (0%)
30
- >AA2002:AAM48354 Aam48354 Murine G protein-coupled receptor, IGPCr17. 4/2002
Length = 347, Score = 603 bits (1554), Expect = e-171, Identities = 299/337 (88%), Positives = 315/337 (93%), Gaps = 1/337 (0%)
- 35 >emb|CAD23845.1| unnamed protein product [Mus musculus]

Length = 347, Score = 603 bits (1554), Expect = e-171, Identities = 299/337 (88%), Positives = 315/337 (93%), Gaps = 1/337 (0%)

5 >ref|NP_081847.2| purinergic receptor P2Y12 [Mus musculus] gb|AAH25428.1| Purinergic receptor P2Y12 [Mus musculus]

Length = 347, Score = 603 bits (1554), Expect = e-171, Identities = 299/337 (88%), Positives = 315/337 (93%), Gaps = 1/337 (0%)

10 >sp|Q9CPV9|P2YC_MOUSE P2Y purinoceptor 12 (P2Y12) dbj|BAB29000.1| unnamed protein product [Mus musculus] dbj|BAB29561.1| unnamed protein product [Mus musculus] gb|AAH27381.1| P2ry12 protein [Mus musculus] emb|CAC85296.1| P2Y12 receptor [Mus musculus]

Length = 347, Score = 603 bits (1554), Expect = e-171, Identities = 299/337 (88%), Positives = 315/337 (93%), Gaps = 1/337 (0%)

15

>dbj|BAC28294.1| unnamed protein product [Mus musculus]

Length = 347, Score = 600 bits (1547), Expect = e-170, Identities = 298/337 (88%), Positives = 314/337 (93%), Gaps = 1/337 (0%)

20 >AA2003A:ABP70856 Abp70856 Rat P2T receptor. 8/2003

Length = 343, Score = 592 bits (1525), Expect = e-168, Identities = 293/337 (86%), Positives = 310/337 (91%), Gaps = 5/337 (1%)

>AA2001:AAE04384 Aae04384 Rat P2-purinergic receptor subtype, P2Y12. 9/2001

25 Length = 343, Score = 592 bits (1525), Expect = e-168, Identities = 293/337 (86%), Positives = 310/337 (91%), Gaps = 5/337 (1%)

30 >ref|NP_073637.1| P2Y12 platelet ADP receptor [Rattus norvegicus] sp|Q9EPX4|P2YC_RAT P2Y purinoceptor 12 (P2Y12) (P2Y12 platelet ADP receptor) gb|AAG48945.1| P2Y12 platelet ADP receptor [Rattus norvegicus]

Length = 343, Score = 592 bits (1525), Expect = e-168, Identities = 293/337 (86%), Positives = 310/337 (91%), Gaps = 5/337 (1%)

- >AA2000:AAAY94498 Aay94498 Rat MP-10 receptor protein. 9/2000
Length = 387, Score = 578 bits (1490), Expect = e-164, Identities = 287/332 (86%), Positives = 304/332 (91%), Gaps = 5/332 (1%)
- 5 >AA2001:AAE04387 Aae04387 Human P2-purinergic receptor subtype, P2Y12 truncated allelic variant. 9/2001
Length = 267, Score = 480 bits (1236), Expect = e-134, Identities = 242/253 (95%), Positives = 245/253 (96%), Gaps = 1/253 (0%)
- 10 >AA2003B:ADC55385 Adc55385 Human purine receptor TG20022 protein sequence. 12/2003
Length = 333, Score = 323 bits (827), Expect = 5e-87, Identities = 155/316 (49%), Positives = 212/316 (67%), Gaps = 2/316 (0%)
- >AA2003A:ADA23279 Ada23279 Human SECX polypeptide, SEC1. 11/2003
- 15 Length = 333, Score = 323 bits (827), Expect = 5e-87, Identities = 155/316 (49%), Positives = 212/316 (67%), Gaps = 2/316 (0%)
- >AA2003A:ABU61800 Abu61800 Human G-protein coupled receptor (GPCR), GPR3/SP174. 9/2003
- 20 Length = 333, Score = 323 bits (827), Expect = 5e-87, Identities = 155/316 (49%), Positives = 212/316 (67%), Gaps = 2/316 (0%)
- >AA2003A:ABP82013 Abp82013 G protein-coupled receptor GPR86/94/P2Y13 protein SEQ ID NO:514. 3/2003
- 25 Length = 333, Score = 323 bits (827), Expect = 5e-87, Identities = 155/316 (49%), Positives = 212/316 (67%), Gaps = 2/316 (0%)
- >AA2003A:ABR40071 Abr40071 Human G protein-coupled receptor GPR86 (P2Y13). 6/2003
Length = 333, Score = 323 bits (827), Expect = 5e-87, Identities = 155/316 (49%), Positives = 212/316 (67%), Gaps = 2/316 (0%)
- 30
- >AA2001:AAB74397 Aab74397 Human G protein coupling receptor. 7/2001
Length = 333, Score = 323 bits (827), Expect = 5e-87, Identities = 155/316 (49%), Positives = 212/316 (67%), Gaps = 2/316 (0%)
- 35
- >AA2001:AAB74494 Aab74494 Novel human G protein coupled receptor. 5/2001

Length = 333, Score = 323 bits (827), Expect = 5e-87, Identities = 155/316 (49%), Positives = 212/316 (67%), Gaps = 2/316 (0%)

>AA2000:AAB23029 Aab23029 Human chemokine receptor-like protein, SECX 2777610. 1/2001

5 Length = 333, Score = 323 bits (827), Expect = 5e-87, Identities = 155/316 (49%), Positives = 212/316 (67%), Gaps = 2/316 (0%)

>sp|Q9BPV8|GPR86_HUMAN Probable G protein-coupled receptor GPR86 (P2Y purinoceptor 13) (P2Y13) gb|AAK01864.1| G-protein coupled receptor GPR86 [Homo sapiens] gb|AAK29068.1|
10 putative G-protein-coupled receptor FKSG77 [Homo sapiens] gb|AAL01038.1| G protein-coupled receptor GPR86 [Homo sapiens] gb|AAL26484.1| G protein-coupled receptor [Homo sapiens] gb|AAH41116.2| Purinergic receptor P2Y13 [Homo sapiens]

Length = 333, Score = 323 bits (827), Expect = 5e-87, Identities = 155/316 (49%), Positives = 212/316 (67%), Gaps = 2/316 (0%).

15

Example 2: Expression profiling

Total cellular RNA was isolated from cells by one of two standard methods: 1) guanidine isothiocyanate/Cesium chloride density gradient centrifugation [Kellogg, (1990)]; or with the Tri-Reagent protocol according to the manufacturer's specifications (Molecular Research Center, Inc.,
20 Cincinnati, Ohio). Total RNA prepared by the Tri-reagent protocol was treated with DNase I to remove genomic DNA contamination.

For relative quantitation of the mRNA distribution of P2Y12, total RNA from each cell or tissue source was first reverse transcribed. 85 µg of total RNA was reverse transcribed using 1 µmole random hexamer primers, 0.5 mM each of dATP, dCTP, dGTP and dTTP (Qiagen, Hilden,
25 Germany), 3000 U RnaseQut (Invitrogen, Groningen, Netherlands) in a final volume of 680 µl. The first strand synthesis buffer and Omniscript reverse transcriptase (2 u/µl) were from (Qiagen, Hilden, Germany). The reaction was incubated at 37°C for 90 minutes and cooled on ice. The volume was adjusted to 6800 µl with water, yielding a final concentration of 12.5 ng/µl of starting RNA.

30 For relative quantitation of the distribution of P2Y12 mRNA in cells and tissues the Applied Biosystems 7900 HT Sequence Detection system or Biorad iCycler was used according to the manufacturer's specifications and protocols. PCR reactions were set up to quantitate P2Y12 and the housekeeping genes HPRT (hypoxanthine phosphoribosyltransferase), GAPDH (glyceraldehyde-3-phosphate dehydrogenase), β-actin, and others. Forward and reverse primers

and probes for P2Y12 were designed using the Perkin Elmer ABI Primer Express™ software and were synthesized by TibMolBiol (Berlin, Germany). The P2Y12 forward primer sequence was: Primer1 (SEQ ID NO: 3). The P2Y12 reverse primer sequence was Primer2 (SEQ ID NO: 4). Probe1 (SEQ ID NO: 5), labelled with FAM (carboxyfluorescein succinimidyl ester) as the reporter dye and TAMRA (carboxytetramethylrhodamine) as the quencher, is used as a probe for P2Y12. The following reagents were prepared in a total of 25 µl : 1x TaqMan buffer A, 5.5 mM MgCl₂, 200 nM of dATP, dCTP, dGTP, and dUTP, 0.025 U/µl AmpliTaq Gold™, 0.01 U/µl AmpErase and Probe1 (SEQ ID NO: 5), P2Y12 forward and reverse primers each at 200 nM, 200 nM P2Y12 FAM/TAMRA-labelled probe, and 5 µl of template cDNA. Thermal cycling parameters were 2 min at 50°C, followed by 10 min at 95°C, followed by 40 cycles of melting at 95°C for 15 sec and annealing/extending at 60°C for 1 min.

Calculation of corrected CT values

The CT (threshold cycle) value is calculated as described in the “Quantitative determination of nucleic acids” section. The CF-value (factor for threshold cycle correction) is calculated as follows :

1. PCR reactions were set up to quantitate the housekeeping genes (HKG) for each cDNA sample.
2. CT_{HKG}-values (threshold cycle for housekeeping gene) were calculated as described in the “Quantitative determination of nucleic acids” section.
- 20 3. CT_{HKG}-mean values (CT mean value of all HKG tested on one cDNAs) of all HKG for each cDNA are calculated (n = number of HKG):

$$CT_{HKG-n}\text{-mean value} = (CT_{HKG1}\text{-value} + CT_{HKG2}\text{-value} + \dots + CT_{HKG-n}\text{-value}) / n$$
4. CT_{panel} mean value (CT mean value of all HKG in all tested cDNAs) =

$$(CT_{HKG1}\text{-mean value} + CT_{HKG2}\text{-mean value} + \dots + CT_{HKG-y}\text{-mean value}) / y$$

(y = number of cDNAs)
- 25 5. CF_{cDNA-n} (correction factor for cDNA n) = CT_{panel}-mean value - CT_{HKG-n}-mean value
6. CT_{cDNA-n} (CT value of the tested gene for the cDNA n) + CF_{cDNA-n} (correction factor for cDNA n) = CT_{cor-cDNA-n} (corrected CT value for a gene on cDNA n)

Calculation of relative expression

Definition : highest $CT_{cor-cDNA-n} \neq 40$ is defined as $CT_{cor-cDNA} [high]$

Relative Expression = $2^{(CT_{cor-cDNA}[high] - CT_{cor-cDNA-n})}$

Tissues

- 5 The expression of P2Y12 was investigated in the tissues listed in table 1.

Expression profile

The results of the the mRNA-quantification (expression profiling) is shown in Table 1.

Table 1: *Relative expression of P2Y12 in various human tissues*

fetal heart	9
heart	92
pericardium	34
heart atrium (right)	82
heart atrium (left)	560
heart ventricle (left)	103
heart ventricle (right)	139
heart apex	25
Purkinje fibers	21
interventricular septum	14
fetal aorta	70
aorta	19
aorta sclerotic	298
artery	124
coronary artery	2
coronary artery sclerotic	13
pulmonary artery	1
carotid artery	1
mesenteric artery	76
vein	161
pulmonic valve	87
coronary artery endothel cells	1075
coronary artery smooth muscle primary cells	64
aortic smooth muscle cells	0

pulmonary artery smooth muscle cells	2
aortic endothel cells	307
HUVEC cells	2
pulmonary artery endothel cells	4
iliac artery endothel cells	6
skin	223
adrenal gland	781
thyroid	76
thyroid tumor	84
pancreas	197
pancreas liver cirrhosis	228
esophagus	175
esophagus tumor	16
stomach	284
stomach tumor	36
colon	67
colon tumor	10
small intestine	516
ileum	815
ileum tumor	41
ileum chronic inflammation	28
rectum	2665
salivary gland	648
fetal liver	65
liver	198
liver liver cirrhosis	2521
liver lupus disease	11
liver tumor	25
HEP G2 cells	19
leukocytes (peripheral blood)	1121
Jurkat (T-cells)	229
bone marrow	461

erythrocytes	352
lymphnode	71
thymus	75
thrombocytes	3236
bone marrow stromal cells	46
bone marrow CD71+ cells	205
bone marrow CD33+ cells	12
bone marrow CD34+ cells	223
bone marrow CD15+ cells	231
cord blood CD71+ cells	115
cord blood CD34+ cells	1278
neutrophils cord blood	989
neutrophils peripheral blood	955
spleen	181
spleen liver cirrhosis	218
skeletal muscle	95
cartilage	186
bone connective tissue	15
adipose	43
fetal brain	1067
brain	6339
Alzheimer brain	7968
cerebellum	4211
Alzheimer brain cerebellum	955
cerebellum (right)	217
cerebellum (left)	750
cerebral cortex	755
Alzheimer cerebral cortex	7332
frontal lobe	1771
Alzheimer brain frontal lobe	653
occipital lobe	5873
parietal lobe	3517
temporal lobe	2557
postcentral gyrus	265

vermis cerebelli	83
pons	377
substantia nigra	461
caudatum	910
cerebral meninges	5
cerebral peduncles	781
corpus callosum	7281
nucleus accumbens	5595
putamen	7643
hippocampus	14462
thalamus	5557
posteroventral thalamus	14263
dorsalmedial thalamus	20595
hypothalamus	161
dorsal root ganglia	282
spinal cord	6562
spinal cord (ventral horn)	28329
spinal cord (dorsal horn)	12077
neuroblastoma SK-N-MC cells	24
neuroblastoma SH-SY5Y cells	43
neuroblastoma IMR32 cells	247
glial tumor H4 cells	105
HEK CNS + APP	55
neural progenitor cells	714
astrocytes	584
retina	234
fetal lung	258
fetal lung fibroblast IMR-90 cells	0
fetal lung fibroblast MRC-5 cells	0
lung	54
lung right upper lobe	163
lung right mid lobe	100
lung right lower lobe	189
lung lupus disease	12
lung tumor	86

lung COPD	97
trachea	292
primary bronchia	1898
secondary bronchia	2034
bronchial epithelial cells	2896
bronchial smooth muscle cells	247
small airway epithelial cells	8
cervix	131
testis	739
HeLa cells (cervix tumor)	3
placenta	197
uterus	48
uterus tumor	10
ovary	251
ovary tumor	75
breast	67
breast tumor	282
MDA MB 231 cells (breast tumor)	6
mammary gland	54
prostate	362
prostate BPH	24
prostate tumor	18951
bladder	335
ureter	362
penis	1698
corpus cavernosum	484
fetal kidney	67
kidney	10
kidney tumor	40
renal epithelial cells	13
HEK 293 cells	12

Example 3: Antisense Analysis

Knowledge of the correct, complete cDNA sequence coding for P2Y12 enables its use as a tool for antisense technology in the investigation of gene function. Oligonucleotides, cDNA or genomic
5 fragments comprising the antisense strand of a polynucleotide coding for P2Y12 are used either in vitro or in vivo to inhibit translation of the mRNA. Such technology is now well known in the art, and antisense molecules can be designed at various locations along the nucleotide sequences. By treatment of cells or whole test animals with such antisense sequences, the gene of interest is effectively turned off. Frequently, the function of the gene is ascertained by observing behavior at
10 the intracellular, cellular, tissue or organismal level (e.g., lethality, loss of differentiated function, changes in morphology, etc.).

In addition to using sequences constructed to interrupt transcription of a particular open reading frame, modifications of gene expression is obtained by designing antisense sequences to intron regions, promoter/enhancer elements, or even to trans-acting regulatory genes.

15 Example 4: Expression of P2Y12

Expression of P2Y12 is accomplished by subcloning the cDNAs into appropriate expression vectors and transfecting the vectors into expression hosts such as, e.g., *E. coli*. In a particular case, the vector is engineered such that it contains a promoter for β -galactosidase, upstream of the cloning site, followed by sequence containing the amino-terminal Methionine and the subsequent
20 seven residues of β -galactosidase. Immediately following these eight residues is an engineered bacteriophage promoter useful for artificial priming and transcription and for providing a number of unique endonuclease restriction sites for cloning.

Induction of the isolated, transfected bacterial strain with Isopropyl- β -D-thiogalactopyranoside (IPTG) using standard methods produces a fusion protein corresponding to the first seven residues
25 of β -galactosidase, about 15 residues of "linker", and the peptide encoded within the cDNA. Since cDNA clone inserts are generated by an essentially random process, there is probability of 33% that the included cDNA will lie in the correct reading frame for proper translation. If the cDNA is not in the proper reading frame, it is obtained by deletion or insertion of the appropriate number of bases using well known methods including in vitro mutagenesis, digestion with exonuclease III or
30 mung bean nuclease, or the inclusion of an oligonucleotide linker of appropriate length.

The P2Y12 cDNA is shuttled into other vectors known to be useful for expression of proteins in specific hosts. Oligonucleotide primers containing cloning sites as well as a segment of DNA

(about 25 bases) sufficient to hybridize to stretches at both ends of the target cDNA is synthesized chemically by standard methods. These primers are then used to amplify the desired gene segment by PCR. The resulting gene segment is digested with appropriate restriction enzymes under standard conditions and isolated by gel electrophoresis. Alternately, similar gene segments are produced by digestion of the cDNA with appropriate restriction enzymes. Using appropriate primers, segments of coding sequence from more than one gene are ligated together and cloned in appropriate vectors. It is possible to optimize expression by construction of such chimeric sequences.

Suitable expression hosts for such chimeric molecules include, but are not limited to, mammalian cells such as Chinese Hamster Ovary (CHO) and human 293 cells, insect cells such as Sf9 cells, yeast cells such as *Saccharomyces cerevisiae* and bacterial cells such as *E. coli*. For each of these cell systems, a useful expression vector also includes an origin of replication to allow propagation in bacteria, and a selectable marker such as the β -lactamase antibiotic resistance gene to allow plasmid selection in bacteria. In addition, the vector may include a second selectable marker such as the neomycin phosphotransferase gene to allow selection in transfected eukaryotic host cells. Vectors for use in eukaryotic expression hosts require RNA processing elements such as 3' polyadenylation sequences if such are not part of the cDNA of interest.

Additionally, the vector contains promoters or enhancers which increase gene expression. Such promoters are host specific and include MMTV, SV40, and metallothionine promoters for CHO cells; trp, lac, tac and T7 promoters for bacterial hosts; and alpha factor, alcohol oxidase and PGH promoters for yeast. Transcription enhancers, such as the rous sarcoma virus enhancer, are used in mammalian host cells. Once homogeneous cultures of recombinant cells are obtained through standard culture methods, large quantities of recombinantly produced P2Y₁₂ are recovered from the conditioned medium and analyzed using chromatographic methods known in the art. For example, P2Y₁₂ can be cloned into the expression vector pcDNA3, as exemplified herein. This product can be used to transform, for example, HEK293 or COS by methodology standard in the art. Specifically, for example, using Lipofectamine (Gibco BRL catalog no. 18324-020) mediated gene transfer.

Example 5: Isolation of Recombinant P2Y₁₂

P2Y₁₂ is expressed as a chimeric protein with one or more additional polypeptide domains added to facilitate protein purification. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals [Appa Rao, 1997] and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Washington). The inclusion of a cleavable linker

sequence such as Factor Xa or enterokinase (Invitrogen, Groningen, The Netherlands) between the purification domain and the P2Y₁₂ sequence is useful to facilitate expression of P2Y₁₂.

Example 6: Testing of Chimeric GPCRs

Functional chimeric GPCRs are constructed by combining the extracellular receptive sequences of a new isoform with the transmembrane and intracellular segments of a known isoform for test purposes. This concept was demonstrated by Kobilka et al. (1988), Science 240:1310-1316) who created a series of chimeric $\alpha 2$ - $\beta 2$ adrenergic receptors (AR) by inserting progressively greater amounts of $\alpha 2$ -AR transmembrane sequence into $\beta 2$ -AR. The binding activity of known agonists changed as the molecule shifted from having more $\alpha 2$ than $\beta 2$ conformation, and intermediate constructs demonstrated mixed specificity. The specificity for binding antagonists, however, correlated with the source of the domain VII. The importance of T7G domain VII for ligand recognition was also found in chimeras utilizing two yeast α -factor receptors and is significant because the yeast receptors are classified as miscellaneous receptors. Thus, functional role of specific domains appears to be preserved throughout the GPCR family regardless of category.

In parallel fashion, internal segments or cytoplasmic domains from a particular isoform are exchanged with the analogous domains of a known GPCRs and used to identify the structural determinants responsible for coupling the receptors to trimeric G-proteins. A chimeric receptor in which domains V, VI, and the intracellular connecting loop from $\beta 2$ -AR were substituted into $\alpha 2$ -AR was shown to bind ligands with $\alpha 2$ -AR specificity, but to stimulate adenylate cyclase in the manner of $\beta 2$ -AR. This demonstrates that for adrenergic-type receptors, G-protein recognition is present in domains V and VI and their connecting loop. The opposite situation was predicted and observed for a chimera in which the V->VI loop from $\alpha 1$ -AR replaced the corresponding domain on $\beta 2$ -AR and the resulting receptor bound ligands with $\beta 2$ -AR specificity and activated G-protein-mediated phosphatidylinositol turnover in the $\alpha 1$ -AR manner. Finally, chimeras constructed from muscarinic receptors also demonstrated that V->VI loop is the major determinant for specificity of G-protein activity.

Chimeric or modified GPCRs containing substitutions in the extracellular and transmembrane regions have shown that these portions of the receptor determine ligand binding specificity. For example, two Serine residues conserved in domain V of all adrenergic and D catecholamine GPCRs are necessary for potent agonist activity. These serines are believed to form hydrogen bonds with the catechol moiety of the agonists within the GPCR binding site. Similarly, an Asp residue present in domain III of all GPCRs which bind biogenic amines is believed to form an ion pair with the ligand amine group in the GPCR binding site.

Functional, cloned GPCRs are expressed in heterologous expression systems and their biological activity assessed. One heterologous system introduces genes for a mammalian GPCR and a mammalian G-protein into yeast cells. The GPCR is shown to have appropriate ligand specificity and affinity and trigger appropriate biological activation (growth arrest and morphological changes) of the yeast cells.

An alternate procedure for testing chimeric receptors is based on the procedure utilizing the purinergic receptor (P_{2u}). Function is easily tested in cultured K562 human leukemia cells because these cells lack P_{2u} receptors. K562 cells are transfected with expression vectors containing either normal or chimeric P_{2u} and loaded with fura-a, fluorescent probe for Ca^{++} .
10 Activation of properly assembled and functional P_{2u} receptors with extracellular UTP or ATP mobilizes intracellular Ca^{++} which reacts with fura-a and is measured spectrofluorometrically.

As with the GPCRs above, chimeric genes are created by combining sequences for extracellular receptive segments of any new GPCR polypeptide with the nucleotides for the transmembrane and intracellular segments of the known P_{2u} molecule. Bathing the transfected K562 cells in
15 microwells containing appropriate ligands triggers binding and fluorescent activity defining effectors of the GPCR molecule. Once ligand and function are established, the P_{2u} system is useful for defining antagonists or inhibitors which block binding and prevent such fluorescent reactions.

Example 7: Production of P2Y12 Specific Antibodies

20 Two approaches are utilized to raise antibodies to P2Y12, and each approach is useful for generating either polyclonal or monoclonal antibodies. In one approach, denatured protein from reverse phase HPLC separation is obtained in quantities up to 75 mg. This denatured protein is used to immunize mice or rabbits using standard protocols; about 100 μ g are adequate for immunization of a mouse, while up to 1 mg might be used to immunize a rabbit. For identifying
25 mouse hybridomas, the denatured protein is radioiodinated and used to screen potential murine B-cell hybridomas for those which produce antibody. This procedure requires only small quantities of protein, such that 20 mg is sufficient for labeling and screening of several thousand clones.

In the second approach, the amino acid sequence of an appropriate P2Y12 domain, as deduced from translation of the cDNA, is analyzed to determine regions of high antigenicity. Oligopeptides
30 comprising appropriate hydrophilic regions are synthesized and used in suitable immunization protocols to raise antibodies. The optimal amino acid sequences for immunization are usually at the C-terminus, the N-terminus and those intervening, hydrophilic regions of the polypeptide

which are likely to be exposed to the external environment when the protein is in its natural conformation.

Typically, selected peptides, about 15 residues in length, are synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry and coupled to keyhole limpet hemocyanin (KLH; Sigma, St. Louis, MO) by reaction with M-maleimidobenzoyl-N-hydroxysuccinimide ester, MBS. If necessary, a cysteine is introduced at the N-terminus of the peptide to permit coupling to KLH. Rabbits are immunized with the peptide-KLH complex in complete Freund's adjuvant. The resulting antisera are tested for antipeptide activity by binding the peptide to plastic, blocking with 1% bovine serum albumin, reacting with antisera, washing and reacting with labeled (radioactive or fluorescent), affinity purified, specific goat anti-rabbit IgG.

Hybridomas are prepared and screened using standard techniques. Hybridomas of interest are detected by screening with labeled P2Y₁₂ to identify those fusions producing the monoclonal antibody with the desired specificity. In a typical protocol, wells of plates (FAST; Becton-Dickinson, Palo Alto, CA) are coated during incubation with affinity purified, specific rabbit anti-mouse (or suitable antiserum 1 g) antibodies at 10 mg/ml. The coated wells are blocked with 1% bovine serum albumin, (BSA), washed and incubated with supernatants from hybridomas. After washing the wells are incubated with labeled P2Y₁₂ at 1 mg/ml. Supernatants with specific antibodies bind more labeled P2Y₁₂ than is detectable in the background. Then clones producing specific antibodies are expanded and subjected to two cycles of cloning at limiting dilution. Cloned hybridomas are injected into pristane-treated mice to produce ascites, and monoclonal antibody is purified from mouse ascitic fluid by affinity chromatography on Protein A. Monoclonal antibodies with affinities of at least

10^8 M^{-1} , preferably 10^9 to 10^{10} M^{-1} or stronger, are typically made by standard procedures.

Example 8: Diagnostic Test Using P2Y₁₂ Specific Antibodies

Particular P2Y₁₂ antibodies are useful for investigating signal transduction and the diagnosis of infectious or hereditary conditions which are characterized by differences in the amount or distribution of P2Y₁₂ or downstream products of an active signaling cascade.

Diagnostic tests for P2Y₁₂ include methods utilizing antibody and a label to detect P2Y₁₂ in human body fluids, membranes, cells, tissues or extracts of such. The polypeptides and antibodies of the present invention are used with or without modification. Frequently, the polypeptides and antibodies are labeled by joining them, either covalently or noncovalently, with a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known

and have been reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent agents, chemiluminescent agents, chromogenic agents, magnetic particles and the like.

5 A variety of protocols for measuring soluble or membrane-bound P2Y₁₂, using either polyclonal or monoclonal antibodies specific for the protein, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). A two-site monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on P2Y₁₂ is preferred, but a competitive binding assay may be employed.

10 **Example 9: Purification of Native P2Y₁₂ Using Specific Antibodies**

Native or recombinant P2Y₁₂ is purified by immunoaffinity chromatography using antibodies specific for P2Y₁₂. In general, an immunoaffinity column is constructed by covalently coupling the anti-TRH antibody to an activated chromatographic resin.

15 Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated Sepharose (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is
20 blocked, and the derivative resin is washed according to the manufacturer's instructions.

Such immunoaffinity columns are utilized in the purification of P2Y₁₂ by preparing a fraction from cells containing P2Y₁₂ in a soluble form. This preparation is derived by solubilization of whole cells or of a subcellular fraction obtained via differential centrifugation (with or without addition of detergent) or by other methods well known in the art. Alternatively, soluble P2Y₁₂
25 containing a signal sequence is secreted in useful quantity into the medium in which the cells are grown.

A soluble P2Y₁₂-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of P2Y₁₂ (e.g., high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions
30 that disrupt antibody/protein binding (e.g., a buffer of pH 2-3 or a high concentration of a chaotrope such as urea or thiocyanate ion), and P2Y₁₂ is collected.

Example 10: Drug Screening

This invention is particularly useful for screening therapeutic compounds by using P2Y₁₂ or binding fragments thereof in any of a variety of drug screening techniques. As P2Y₁₂ is a G protein coupled receptor any of the methods commonly used in the art may potentially be used to identify P2Y₁₂ ligands. For example, the activity of a G protein coupled receptor such as P2Y₁₂ can be measured using any of a variety of appropriate functional assays in which activation of the receptor results in an observable change in the level of some second messenger system, such as adenylylate cyclase, guanylylcyclase, calcium mobilization, or inositol phospholipid hydrolysis. Alternatively, the polypeptide or fragment employed in such a test is either free in solution, affixed to a solid support, borne on a cell surface or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, are used for standard binding assays.

Measured, for example, is the formation of complexes between P2Y₁₂ and the agent being tested. Alternatively, one examines the diminution in complex formation between P2Y₁₂ and a ligand caused by the agent being tested.

Thus, the present invention provides methods of screening for drug candidates, drugs, or any other agents which affect signal transduction. These methods, well known in the art, comprise contacting such an agent with P2Y₁₂ polypeptide or a fragment thereof and assaying (i) for the presence of a complex between the agent and P2Y₁₂ polypeptide or fragment, or (ii) for the presence of a complex between P2Y₁₂ polypeptide or fragment and the cell. In such competitive binding assays, the P2Y₁₂ polypeptide or fragment is typically labeled. After suitable incubation, free P2Y₁₂ polypeptide or fragment is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular agent to bind to P2Y₁₂ or to interfere with the P2Y₁₂-agent complex.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to P2Y₁₂ polypeptides. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with P2Y₁₂ polypeptide and washed. Bound P2Y₁₂ polypeptide is then detected by methods well known in the art. Purified P2Y₁₂ are also coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies are used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding P2Y₁₂ specifically compete with a test compound for binding to P2Y₁₂ polypeptides or fragments thereof. In this manner, the antibodies are used to detect the presence of any peptide which shares one or more antigenic determinants with P2Y₁₂.

5 **Example 11: Rational Drug Design**

The goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of small molecules with which they interact, agonists, antagonists, or inhibitors. Any of these examples are used to fashion drugs which are more active or stable forms of the polypeptide or which enhance or interfere with the function of a polypeptide in vivo.

- 10 In one approach, the three-dimensional structure of a protein of interest, or of a protein-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of a polypeptide is gained by modeling based on the
- 15 structure of homologous proteins. In both cases, relevant structural information is used to design efficient inhibitors. Useful examples of rational drug design include molecules which have improved activity or stability or which act as inhibitors, agonists, or antagonists of native peptides.

It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore

20 upon which subsequent drug design is based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids is expected to be an analog of the original receptor. The anti-id is then used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides then act as the

25 pharmacore.

By virtue of the present invention, sufficient amount of polypeptide are made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the P2Y₁₂ amino acid sequence provided herein provides guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

30 **Example 12: Identification of Other Members of the Signal Transduction Complex**

The inventive purified P2Y₁₂ is a research tool for identification, characterization and purification of interacting G or other signal transduction pathway proteins. Radioactive labels are incorporated

into a selected P2Y₁₂ domain by various methods known in the art and used in vitro to capture interacting molecules. A preferred method involves labeling the primary amino groups in P2Y₁₂ with ¹²⁵I Bolton-Hunter reagent. This reagent has been used to label various molecules without concomitant loss of biological activity.

5 Labeled P2Y₁₂ is useful as a reagent for the purification of molecules with which it interacts. In one embodiment of affinity purification, membrane-bound P2Y₁₂ is covalently coupled to a chromatography column. Cell-free extract derived from synovial cells or putative target cells is passed over the column, and molecules with appropriate affinity bind to P2Y₁₂. P2Y₁₂-complex is recovered from the column, and the P2Y₁₂-binding ligand disassociated and subjected to N-
10 terminal protein sequencing. The amino acid sequence information is then used to identify the captured molecule or to design degenerate oligonucleotide probes for cloning the relevant gene from an appropriate cDNA library.

In an alternate method, antibodies are raised against P2Y₁₂, specifically monoclonal antibodies. The monoclonal antibodies are screened to identify those which inhibit the binding of labeled
15 P2Y₁₂. These monoclonal antibodies are then used therapeutically.

Example 13: Use and Administration of Antibodies, Inhibitors, or Antagonists

Antibodies, inhibitors, or antagonists of P2Y₁₂ or other treatments and compounds that are limiters of signal transduction (LSTs), provide different effects when administered therapeutically. LSTs are formulated in a nontoxic, inert, pharmaceutically acceptable aqueous carrier medium
20 preferably at a pH of about 5 to 8, more preferably 6 to 8, although pH may vary according to the characteristics of the antibody, inhibitor, or antagonist being formulated and the condition to be treated. Characteristics of LSTs include solubility of the molecule, its half-life and antigenicity/-immunogenicity. These and other characteristics aid in defining an effective carrier. Native human proteins are preferred as LSTs, but organic or synthetic molecules resulting from drug
25 screens are equally effective in particular situations.

LSTs are delivered by known routes of administration including but not limited to topical creams and gels; transmucosal spray and aerosol; transdermal patch and bandage; injectable, intravenous and lavage formulations; and orally administered liquids and pills particularly formulated to resist stomach acid and enzymes. The particular formulation, exact dosage, and route of administration
30 is determined by the attending physician and varies according to each specific situation.

Such determinations are made by considering multiple variables such as the condition to be treated, the LST to be administered, and the pharmacokinetic profile of a particular LST. Addi-

tional factors which are taken into account include severity of the disease state, patient's age, weight, gender and diet, time and frequency of LST administration, possible combination with other drugs, reaction sensitivities, and tolerance/response to therapy. Long acting LST formulations might be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular LST.

Normal dosage amounts vary from 0.1 to 10^5 μ g, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. Those skilled in the art employ different formulations for different LSTs. Administration to cells such as nerve cells necessitates delivery in a manner different from that to other cells such as vascular endothelial cells.

It is contemplated that abnormal signal transduction, trauma, or diseases which trigger P2Y₁₂ activity are treatable with LSTs. These conditions or diseases are specifically diagnosed by the tests discussed above, and such testing should be performed in suspected cases of viral, bacterial or fungal infections, allergic responses, mechanical injury associated with trauma, hereditary diseases, lymphoma or carcinoma, or other conditions which activate the genes of lymphoid or neuronal tissues.

Example 14: Production of Non-human Transgenic Animals

Animal model systems which elucidate the physiological and behavioral roles of the P2Y₁₂ are produced by creating nonhuman transgenic animals in which the activity of the P2Y₁₂ is either increased or decreased, or the amino acid sequence of the expressed P2Y₁₂ is altered, by a variety of techniques. Examples of these techniques include, but are not limited to: 1) Insertion of normal or mutant versions of DNA encoding a P2Y₁₂, by microinjection, electroporation, retroviral transfection or other means well known to those skilled in the art, into appropriately fertilized embryos in order to produce a transgenic animal or 2) homologous recombination of mutant or normal, human or animal versions of these genes with the native gene locus in transgenic animals to alter the regulation of expression or the structure of these P2Y₁₂ sequences. The technique of homologous recombination is well known in the art. It replaces the native gene with the inserted gene and hence is useful for producing an animal that cannot express native P2Y₁₂s but does express, for example, an inserted mutant P2Y₁₂, which has replaced the native P2Y₁₂ in the animal's genome by recombination, resulting in underexpression of the transporter. Microinjection adds genes to the genome, but does not remove them, and the technique is useful for producing an animal which expresses its own and added P2Y₁₂, resulting in overexpression of the P2Y₁₂.

One means available for producing a transgenic animal, with a mouse as an example, is as follows: Female mice are mated, and the resulting fertilized eggs are dissected out of their oviducts. The eggs are stored in an appropriate medium such as cesiumchloride M2 medium. DNA or cDNA encoding P2Y12 is purified from a vector by methods well known to the one skilled in the art.

5 Inducible promoters may be fused with the coding region of the DNA to provide an experimental means to regulate expression of the transgene. Alternatively or in addition, tissue specific regulatory elements may be fused with the coding region to permit tissue-specific expression of the transgene. The DNA, in an appropriately buffered solution, is put into a microinjection needle (which may be made from capillary tubing using a piper puller) and the egg to be injected is put in

10 a depression slide. The needle is inserted into the pronucleus of the egg, and the DNA solution is injected. The injected egg is then transferred into the oviduct of a pseudopregnant mouse which is a mouse stimulated by the appropriate hormones in order to maintain false pregnancy, where it proceeds to the uterus, implants, and develops to term. As noted above, microinjection is not the only method for inserting DNA into the egg but is used here only for exemplary purposes.

15 **References**

- U.S. 4,522,811
U.S. 5,283,317
U.S. 5,565,332
U.S. 2001/046497
- 20 WO 84/03564
WO 92/01810
WO 93/03151
WO 94/13804
WO 01/04297
- 25 WO 00/28028
WO 00/31258
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ; *Nucleic Acids Res* 1997 Sep 1; 25(17): 3389-402
- Appa Rao et al., 1997, *Protein Expr Purif* Nov, 11(2): 201-8
- 30 Barnes, 2000, *Chest*, 117:10S14S
- Botstein et al., 1980, *Am J Hum Genet.* 32: 314-31
- Cattaneo et al., *Blood* 80: 2787-2796, 1992
- Colbere-Garapin et al., 1981, *J. Mol. Biol.* 150, 1-14
- Engelhard et al., 1994, *Proc. Nat. Acad. Sci.* 91, 3224-3227
- 35 Fontana et al., *Circulation* 108 (8), 989-995 (2003)

- Gergen and Weiss, 1992, *Am Rev Respir Dis* 146:823-824
- Gibson et al., 1996, *Genome Research* 6: 995-1001
- Haseloff *et al.*, 1988, *Nature* 334, 585-591
- Heid et al., 1996, *Genome Research* 6: 986-994
- 5 Holland et al., 1991, *PNAS* 88: 7276-7280
- Hollopeter et al., *Nature* 409: 202-206, 2001
- Iwabuchi *et al.*, 1993, *Oncogene* 8, 1693-1696
- Jagroop et al., *Platelets* 14 (1), 15-20 (2003)
- Jeffreys et al., 1985, *Nature* 316: 76-9
- 10 Johnson *et al.*, 1989, *Endoc. Rev.* 10, 317-331
- Kellogg et al., 1990, *Anal. Biochem.* 189:202-208
- Lam, 1997, *Anticancer Drug Res.* 12(3):145-67
- Lefkowitz, 1991, *Nature* 351, 353-354
- Livak et al., 1995, *PCR Methods and Applications* 357-362
- 15 Logan, Shenk, 1984, *Proc. Natl. Acad. Sci.* 81, 3655-3659
- Lowy *et al.*, 1980, *Cell* 22, 817-23
- Maddox *et al.*, 1983, *J. Exp. Med.* 158, 1211-1216
- McConnell *et al.*, 1992, *Science* 257, 1906-1912
- Nicholls *et al.*, 1993, *J. Immunol. Meth.* 165, 81-91
- 20 Nurden et al., *J. Clin. Invest.* 95: 1612-1622, 1995
- Oury et al., *J. Biol. Chem.* 275: 22611-22614, 2000
- Piatak et al., 1993, *BioTechniques* 14:70-81
- Piatak et al., 1993, *Science* 259:1749-1754
- Porath *et al.*, 1992, *Prot. Exp. Purif.* 3, 263-281
- 25 Roberge *et al.*, 1995, *Science* 269, 202-204
- Sjolander, Urbaniczky, 1991, *Anal. Chem.* 63, 2338-2345
- Szabo *et al.*, 1995, *Curr. Opin. Struct. Biol.* 5, 699-705
- Thomas, 1980, *Proc. Nat. Acad. Sci.*, 77:5201-5205
- Uhlmann *et al.*, 1987, *Tetrahedron. Lett.* 215, 3539-3542
- 30 Weber et al., 1990, *Genomics* 7: 524-30
- Wigler *et al.*, 1977, *Cell* 11, 223-32
- Wigler *et al.*, 1980, *Proc. Natl. Acad. Sci.* 77, 3567-70
- Zhang et al., *J Biol Chem.* 2001 Mar 16;276(11):8608-15.

Claims

1. A method of screening for therapeutic agents useful in the treatment of a disease
comprised in a group of diseases consisting of cardiovascular diseases, endocrinological
diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hema-
tological diseases, respiratory diseases, neurological diseases and urological diseases in a
mammal comprising the steps of
 - i) contacting a test compound with a P2Y₁₂ polypeptide,
 - ii) detect binding of said test compound to said P2Y₁₂ polypeptide.
2. A method of screening for therapeutic agents useful in the treatment of a disease
comprised in a group of diseases consisting of cardiovascular diseases, endocrinological
diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hema-
tological diseases, respiratory diseases, neurological diseases and urological diseases in a
mammal comprising the steps of
 - i) determining the activity of a P2Y₁₂ polypeptide at a certain concentration of a test
compound or in the absence of said test compound,
 - ii) determining the activity of said polypeptide at a different concentration of said test
compound.
3. A method of screening for therapeutic agents useful in the treatment of a disease
comprised in a group of diseases consisting of cardiovascular diseases, endocrinological
diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation,
hematological diseases, respiratory diseases, neurological diseases and urological diseases
in a mammal comprising the steps of
 - i) determining the activity of a P2Y₁₂ polypeptide at a certain concentration of a test
compound,
 - ii) determining the activity of a P2Y₁₂ polypeptide at the presence of a compound
known to be a regulator of a P2Y₁₂ polypeptide.
4. The method of any of claims 1 to 3, wherein the step of contacting is in or at the surface of
a cell.
5. The method of any of claims 1 to 3, wherein the cell is in vitro.

6. The method of any of claims 1 to 3, wherein the step of contacting is in a cell-free system.
7. The method of any of claims 1 to 3, wherein the polypeptide is coupled to a detectable label.
8. The method of any of claims 1 to 3, wherein the compound is coupled to a detectable label.
- 5 9. The method of any of claims 1 to 3, wherein the test compound displaces a ligand which is first bound to the polypeptide.
10. The method of any of claims 1 to 3, wherein the polypeptide is attached to a solid support.
11. The method of any of claims 1 to 3, wherein the compound is attached to a solid support.
- 10 12. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
 - i) contacting a test compound with a P2Y₁₂ polynucleotide,
 - 15 ii) detect binding of said test compound to said P2Y₁₂ polynucleotide.
13. The method of claim 12 wherein the nucleic acid molecule is RNA.
14. The method of claim 12 wherein the contacting step is in or at the surface of a cell.
15. The method of claim 12 wherein the contacting step is in a cell-free system.
16. The method of claim 12 wherein polynucleotide is coupled to a detectable label.
- 20 17. The method of claim 12 wherein the test compound is coupled to a detectable label.
18. A method of diagnosing a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
 - 25 i) determining the amount of a P2Y₁₂ polynucleotide in a sample taken from said mammal,

- ii) determining the amount of P2Y₁₂ polynucleotide in healthy and/or diseased mammals.
19. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which binds to a P2Y₁₂ polypeptide.
20. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a P2Y₁₂ polypeptide.
21. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a P2Y₁₂ polypeptide, wherein said therapeutic agent is
- i) a small molecule,
- ii) an RNA molecule,
- iii) an antisense oligonucleotide,
- iv) a polypeptide,
- v) an antibody, or
- vi) a ribozyme.
22. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a P2Y₁₂ polynucleotide.
23. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases,

respiratory diseases, neurological diseases and urological diseases in a mammal comprising a P2Y₁₂ polypeptide.

24. Use of regulators of a P2Y₁₂ for the preparation of a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal.
25. Method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
- i) identifying a regulator of P2Y₁₂,
 - ii) determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal; and
 - iii) combining of said regulator with an acceptable pharmaceutical carrier.
26. Use of a regulator of P2Y₁₂ for the regulation of P2Y₁₂ activity in a mammal having a disease comprised in a group of diseases consisting of cardiovascular diseases, endocrinological diseases, metabolic diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases.

Fig. 1

SEQ ID NO: 1

ATCACAATCAGAAGACAGGAGCTGCAGAACAGAACACTTTCTCATGTC
CAGGGTCAGATTACAAGAGCACTCAAGACTTTACTGACGAAAACCTCAG
GAAATCCTCTATCACAAAGAGGTTTGGCAACTAACTAAGACATTAAA
AGGAAAATACCAGATGCCACTCTGCAGGCTGCAATAACTACTACTTAC
TGGATACATTCAAACCCTCCAGAATCAACAGTTATCAGGTAACCAACA
AGAAATGCAAGCCGTCGACAATCTCACCTCTGCGCCTGGGAACACCAG
TCTGTGCACCAGAGACTACAAAATCACCCAGGTCCTCTTCCCACTGCT
CTACACTGTCCTGTTTTTTGTTGGACTTATCACAAATGGCCTGGCGAT
GAGGATTTTCTTTCAAATCCGGAGTAAATCAAACCTTTATTATTTTTCT
TAAGAACACAGTCATTTCTGATCTTCTCATGATTCTGACTTTTCCATT
CAAAATTCTTAGTGATGCCAAACTGGGAACAGGACCACTGAGAACTTT
TGTGTGTCAAGTTACCTCCGTCATATTTTATTTTACAATGTATATCAG
TATTTTCAATCCTGGGACTGATAACTATCGATCGCTACCAGAAGACCAC
CAGGCCATTTAAACATCCAACCCCAAAAATCTCTTGGGGGCTAAGAT
TCTCTCTGTTGTCTGTTGTCATGTTCTTACTCTCTTTGCCTAA
CATGATTCTGACCAACAGGCAGCCGAGAGACAAGAATGTGAAGAAATG
CTCTTTCCTTAAATCAGAGTTCGGTCTAGTCTGGCATGAAATAGTAAA
TTACATCTGTCAAGTCATTTTCTGGATTAAATTTCTTAATTGTTATTGT
ATGTTATACACTCATTACAAAAGAAGTGTACCGGTCTATACGTAAGAAC
GAGGGGTGTAGGTAAAGTCCCCAGGAAAAAGGTGAACGTCAAAGTTTT
CATTATCATTGCTGTATTCTTTATTTGTTTTGTTTCTTTCCATTTTGC
CCGAATTCCTTACACCCTGAGCCAAACCCGGGATGTCTTTGACTGCAC
TGCTGAAAATACTCTGTTCTATGTGAAAGAGAGCACTCTGTGGTTAAC
TTCCTTAAATGCATGCCTGGATCCGTTTCTATTTTTTTCTTTTGCAA
GTCCTTCAGAAATTCCTTGATAAGTATGCTGAAGTGCCCCAATTCTGC
AACATCTCTGTCCCAGGACAATAGGAAAAAAGAACAGGATGGTGGTGA
CCCAAATGAAGAGACTCCAATGTAAACAAATTAATAAGGAAATATTT
CAATCTCTTTGTGTTTCAAGAACTCGTTAAAGCAAAGCGCTAAGTAAAAA
TATTAAGTACGAAGAAGCAACTAAGTTAATAATAATGACTCTAAAGA
AACAAAAA

Fig. 2**SEQ ID NO: 2**

MQAVDNLTSA PGNTSLCTR DYKITQVLFPLLYTVLFFVGLITNGLAMR
IFFQIRSKSNFII FLKNTVISDLLMILTFPFKILSDAKLGTGPLRRTFV
CQVTSVIFYFTMYISISFLGLITIDRYQKTTRPFKTSNPKNLLGAKIL
SVVIWAFMFLLSLPNMILTNRQPRDKNVKKCSFLKSEFGLVWHEIVNY
ICQVIFWINFLIVIVCYTLITKELYRSYVRTRGVGKVPRKKVNVKVFI
IIAVFFICFVPPHFARI PYTSLSQTRDVF DCTAENTLFYVKESTLWLTS
LNACLD PFIYFFLCKSFRNSLISMLKCPNSATSLSQDNRKKEQDGGDP
NEETPM

Fig. 3**SEQ ID NO: 3**

5' - TCGACAATCTCACCTCTGCG - 3'

Fig. 4**SEQ ID NO: 4**

5' - AGAGGACCTGGGTGATTTTGTAGT - 3'

Fig. 5**SEQ ID NO: 5**

5' - TCGACAATCTCACCTCTGCG - 3'

SEQUENCE LISTING

<110> Bayer HealthCare AG

<120> Diagnostics and Therapeutics for Diseases Associated with G
Protein-Coupled Receptor P2Y12 (P2Y12)

<130> BHC 04 01 030

<160> 5

<170> PatentIn version 3.2

<210> 1

<211> 1400

<212> DNA

<213> Homo sapiens

<400> 1

```

atcacaatca gaagacagga gctgcagaac agaacacttt ctcatgtcca gggtcagatt   60
acaagagcac tcaagacttt actgacgaaa actcaggaaa tcctctatca caaagagggt   120
tggcaactaa actaagacat taaaaggaaa ataccagatg ccactctgca ggctgcaata   180
actactactt actggatata ttcaaaccct ccagaatcaa cagttatcag gtaaccaaca   240
agaaatgcaa gccgtcgaca atctcacctc tgcgcctggg aacaccagtc tgtgcaccag   300
agactacaaa atcaccaggg tcctcttccc actgctctac actgtcctgt tttttgttgg   360
acttatcaca aatggccttg cgatgaggat tttctttcaa atccggagta aatcaaactt   420
tattattttt cttaagaaca cagtcatttc tgatcttctc atgattctga cttttccatt   480
caaaattctt agtgatgcca aactgggaac aggaccactg agaacttttg tgtgtcaagt   540
tacctccgtc atattttatt tcacaatgta tatcagtatt tcattcctgg gactgataac   600
tatcgatcgc taccagaaga ccaccaggcc atttaaaaca tccaacccca aaaatctctt   660
gggggctaag attctctctg ttgtcatctg ggcatctcatg ttcttactct ctttgcctaa   720
catgattctg accaacaggc agccgagaga caagaatgtg aagaaatgct ctttccttaa   780
atcagagttc ggtctagtct ggcattgaaat agtaaattac atctgtcaag tcattttctg   840
gattaatttc ttaattgtta ttgtatgtta tacactcatt acaaaagaac tgtaccggtc   900
atacgtaaga acgaggggtg taggtaaagt ccccgagaaa aaggtgaacg tcaaagtttt   960
cattatcatt gctgtattct ttatttggtt tggtcctttc cattttgccc gaattcctta  1020
caccctgagc caaaccggg atgtctttga ctgcactgct gaaaatactc tgttctatgt  1080
gaaagagagc actctgtggt taacttcctt aaatgcatgc ctggatccgt tcattctatt  1140
tttctttgca aagtcttca gaaattcctt gataagtatg ctgaagtgcc ccaattctgc  1200
aacatctctg tcccaggaca ataggaaaaa agaacaggat ggtggtgacc caaatgaaga  1260
gactccaatg taaacaaatt aactaaggaa atattttcaat ctctttgtgt tcagaactcg  1320
ttaaagcaaa gcgctaagta aaaatattaa ctgacgaaga agcaactaag ttaataataa  1380
tgactctaaa gaaacaaaaa                                     1400

```

<210> 2
 <211> 342
 <212> PRT
 <213> Homo sapiens

 <400> 2
 Met Gln Ala Val Asp Asn Leu Thr Ser Ala Pro Gly Asn Thr Ser Leu
 1 5 10 15
 Cys Thr Arg Asp Tyr Lys Ile Thr Gln Val Leu Phe Pro Leu Leu Tyr
 20 25 30
 Thr Val Leu Phe Phe Val Gly Leu Ile Thr Asn Gly Leu Ala Met Arg
 35 40 45
 Ile Phe Phe Gln Ile Arg Ser Lys Ser Asn Phe Ile Ile Phe Leu Lys
 50 55 60
 Asn Thr Val Ile Ser Asp Leu Leu Met Ile Leu Thr Phe Pro Phe Lys
 65 70 75 80
 Ile Leu Ser Asp Ala Lys Leu Gly Thr Gly Pro Leu Arg Thr Phe Val
 85 90 95
 Cys Gln Val Thr Ser Val Ile Phe Tyr Phe Thr Met Tyr Ile Ser Ile
 100 105 110
 Ser Phe Leu Gly Leu Ile Thr Ile Asp Arg Tyr Gln Lys Thr Thr Arg
 115 120 125
 Pro Phe Lys Thr Ser Asn Pro Lys Asn Leu Leu Gly Ala Lys Ile Leu
 130 135 140
 Ser Val Val Ile Trp Ala Phe Met Phe Leu Leu Ser Leu Pro Asn Met
 145 150 155 160
 Ile Leu Thr Asn Arg Gln Pro Arg Asp Lys Asn Val Lys Lys Cys Ser
 165 170 175
 Phe Leu Lys Ser Glu Phe Gly Leu Val Trp His Glu Ile Val Asn Tyr
 180 185 190
 Ile Cys Gln Val Ile Phe Trp Ile Asn Phe Leu Ile Val Ile Val Cys
 195 200 205
 Tyr Thr Leu Ile Thr Lys Glu Leu Tyr Arg Ser Tyr Val Arg Thr Arg
 210 215 220
 Gly Val Gly Lys Val Pro Arg Lys Lys Val Asn Val Lys Val Phe Ile
 225 230 235 240
 Ile Ile Ala Val Phe Phe Ile Cys Phe Val Pro Phe His Phe Ala Arg
 245 250 255
 Ile Pro Tyr Thr Leu Ser Gln Thr Arg Asp Val Phe Asp Cys Thr Ala
 260 265 270
 Glu Asn Thr Leu Phe Tyr Val Lys Glu Ser Thr Leu Trp Leu Thr Ser
 275 280 285
 Leu Asn Ala Cys Leu Asp Pro Phe Ile Tyr Phe Phe Leu Cys Lys Ser
 290 295 300

- 3 -

Phe Arg Asn Ser Leu Ile Ser Met Leu Lys Cys Pro Asn Ser Ala Thr
305 310 315 320
Ser Leu Ser Gln Asp Asn Arg Lys Lys Glu Gln Asp Gly Gly Asp Pro
 325 330 335
Asn Glu Glu Thr Pro Met
 340

<210> 3
<211> 20
<212> DNA
<213> artificial sequence

<220>
<223> forward primer

<400> 3
tcgacaatct cacctctgcg 20

<210> 4
<211> 24
<212> DNA
<213> artificial sequence

<220>
<223> reverse primer

<400> 4
agaggacctg ggtgattttg tagt 24

<210> 5
<211> 20
<212> DNA
<213> artificial sequence

<220>
<223> probe

<400> 5
tcgacaatct cacctctgcg 20