

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 4077307 T3**

-
- (51) Int.Cl.: **C 07 D 401/10 (2006.01)** **A 61 K 31/444 (2006.01)** **A 61 K 31/497 (2006.01)**
A 61 K 31/501 (2006.01) **A 61 K 31/506 (2006.01)** **A 61 P 3/04 (2006.01)**
A 61 P 5/38 (2006.01) **C 07 D 401/12 (2006.01)** **C 07 D 401/14 (2006.01)**
C 07 D 409/14 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2025-05-26**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2025-03-12**
- (86) Europæisk ansøgning nr.: **20903163.2**
- (86) Europæisk indleveringsdag: **2020-12-11**
- (87) Den europæiske ansøgnings publiceringsdag: **2022-10-26**
- (86) International ansøgning nr.: **US2020064493**
- (87) Internationalt publikationsnr.: **WO2021126693**
- (30) Prioritet: **2019-12-18 US 201962949854 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Crinetics Pharmaceuticals, Inc., 6055 Lusk Blvd, San Diego, CA 92121, USA**
- (72) Opfinder: **HAN, Sangdon, , San Diego, California 92121, USA**
KIM, Sun Hee, , San Diego, California 92121, USA
ZHU, Yunfei, , San Diego, California 92121, USA
- (74) Fuldmægtig i Danmark: **NORDIC PATENT SERVICE A/S, Bredgade 30, 1260 København K, Danmark**
- (54) Benævnelse: **GEM-DISUBSTITUERED PIPERIDINMELANOCORTIN SUBTYPE-2-RECEPTOR- (MC2R)-ANTAGONISTER OG ANVENDELSER DERAFT**
- (56) Fremdragne publikationer:
WO-A1-00/74679
WO-A1-03/045918
WO-A1-03/094918
WO-A1-2005/047253
WO-A1-2021/091788
WO-A1-2021/133563
WO-A2-2006/113704
US-A1- 2005 192 286
US-A1- 2019 367 481
SEBHAT, I.K. LAI, Y. BARAKAT, K. YE, Z. TANG, R. KALYANI, R.N. VONGS, A. MACNEIL, T. WEINBERG, D.H. CABELLO, M.A: "Melanocortin subtype 4 receptor agonists: Structure-activity relationships about the 4-alkyl piperidine core", BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, vol. 17, no. 20, 14 September 2007 (2007-09-14), AMSTERDAM, NL, pages 5720 - 5723, XP022249700, ISSN: 0960-894X, DOI:

Fortsættes ...

DESCRIPTION

Description

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Patent Application No. 62/949,854 filed on December 18, 2019.

FIELD OF THE INVENTION

[0002] Described herein are compounds that modulate the activity of one or more melanocortin receptors, methods of making such compounds, pharmaceutical compositions and medicaments comprising such compounds, and such compounds for use in methods of treating conditions, diseases, or disorders that would benefit from modulating melanocortin subtype-2 receptor (MC2R) activity.

BACKGROUND OF THE INVENTION

[0003] The melanocortin receptors form a family of G protein-coupled receptor (GPCRs) (MC1R, MC2R, MC3R, MC4R, and MC5R) that are selectively activated by different melanocortin peptides adrenocorticotrophic hormone (ACTH), and the melanocortin peptides α -, β -, and γ -melanocyte-stimulating hormone (α -MSH, β -MSH, and γ -MSH) that are all derived proteolytically from proopiomelanocortin hormone, or POMC. ACTH is a 39 amino acid peptide that is the primary regulator of adrenal glucocorticoid synthesis and secretion and only has affinity for MC2R. As the central actor in this hypothalamic-pituitary-adrenal (HPA) axis, ACTH is secreted by the pituitary in response to stressful stimuli and acts at the adrenal gland to stimulate the synthesis and secretion of cortisol. Modulation of MC2R is attractive for the treatment of conditions, diseases, or disorders that would benefit from modulating melanocortin receptor activity.

[0004] WO 2021/133563 describes spirocyclic piperidine melanocortin subtype-2 receptor (MC2R) antagonists and uses thereof.

[0005] WO 2021/091788 describes melanocortin subtype-2 receptor (MC2R) antagonists and uses thereof.

[0006] WO 2005/047253 describes melanocortin receptor agonists.


[0007] US 2019/367481 describes melanocortin subtype-2 receptor (MC2R) antagonists and uses thereof.

[0008] Sebhat, I.K., et al., "Melanocortin subtype 4 receptor agonists: Structure-activity relationships about the 4-alkyl piperidine core", BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, AMSTERDAM, NL, (20070914), vol. 17, no. 20, pages 5720 - 5723, describes SAR about the piperidine core in a series of MC4R agonists.

SUMMARY OF THE INVENTION

[0009] Compounds described herein are melanocortin receptor modulator compounds. In some embodiments, compounds described herein modulate one or more of the subtype melanocortin receptor proteins. In some embodiments, compounds described herein modulate two or more of the subtype melanocortin receptor proteins. In some embodiments, compounds described herein modulate MC2R.

[0010] In one aspect, provided herein is a compound of Formula (I), or a pharmaceutically acceptable salt thereof:



Formula (I)

wherein:

R^A unsubstituted or substituted phenyl, unsubstituted or substituted monocyclic 6-membered heteroaryl, or unsubstituted or substituted monocyclic 5-membered heteroaryl, wherein if R^A is substituted then R^A is substituted with 1, 2, 3 or 4 groups selected from R^a , R^b , and R^c ;

R^a , R^b , and R^c are independently selected from the group consisting of hydrogen, halogen, -OR⁴, -CN, -N(R⁴)₂, -C(=O)R⁷, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₂-C₇ heterocycloalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heteroaryl, wherein any substituted group of R^a , R^b , and R^c is substituted with one or more R^6 groups;

or one R^a and one R^b , when present on adjacent atoms of R^A , are taken together with the intervening atoms connecting R^a to R^b to form a 5- to 6-membered monocyclic carbocycle or 5- to 6-membered monocyclic heterocycle, wherein the carbocycle or heterocycle is

unsubstituted or substituted with one or more R^6 groups;

wherein, if R^a , R^b , or R^c is attached to the N atom of a heteroaryl, then it is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1-C_6 alkyl, unsubstituted or substituted C_1-C_6 fluoroalkyl, unsubstituted or substituted C_1-C_6 heteroalkyl, unsubstituted or substituted C_3-C_6 cycloalkyl, or unsubstituted or substituted C_2-C_7 heterocycloalkyl;

R^B is an unsubstituted or substituted phenyl or unsubstituted or substituted monocyclic 6-membered heteroaryl, wherein if R^B is substituted then R^B is substituted with 1, 2, 3 or 4 groups selected from R^d , R^e , and R^f ;

R^d , R^e , and R^f are independently selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, $-N(R^4)_2$, $-C(=O)R^7$, $-C(=O)N(R^4)_2$, unsubstituted or substituted C_1-C_6 alkyl, unsubstituted or substituted C_1-C_6 fluoroalkyl, unsubstituted or substituted C_1-C_6 heteroalkyl, unsubstituted or substituted C_3-C_6 cycloalkyl, unsubstituted or substituted C_2-C_7 heterocycloalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heteroaryl, wherein any substituted group of R^d , R^e , and R^f is substituted with one or more R^6 groups;

wherein, if R^d , R^e , or R^f is attached to the N atom of a heteroaryl, then it is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1-C_6 alkyl, unsubstituted or substituted C_1-C_6 fluoroalkyl, unsubstituted or substituted C_1-C_6 heteroalkyl, unsubstituted or substituted C_3-C_6 cycloalkyl, or unsubstituted or substituted C_2-C_7 heterocycloalkyl;

X^1 is CR^{11} or N;

X^2 is CR^{12} or N;

X^3 is CR^{13} or N;

X^4 is CR^{14} or N;

R^{11} , R^{12} , R^{13} , and R^{14} are each independently hydrogen, halogen, unsubstituted or substituted C_1-C_6 alkyl, unsubstituted or substituted C_1-C_6 fluoroalkyl, unsubstituted or substituted C_1-C_6 heteroalkyl, unsubstituted or substituted C_3-C_6 cycloalkyl, $-CN$, $-OR^4$, $-SR^4$, $-CO_2R^4$, $-C(=O)N(R^4)_2$, or $-N(R^4)_2$;

M is $-(C=O)-$, $-NR^3-$, $-O-$, $-S-$, $-SO_2-$, $^*NR^3-(C=O)-$, $^*-(C=O)-NR^3-$, $^*O-(C=O)NR^3-$, $^*NR^3-(C=O)O-$, $^*NR^3-(C=O)NR^3-$, $^*NR^3(SO_2)-$, $^*SO_2NR^3-$, or 5-membered heterocycle, wherein * indicates the attachment point to R^1 ;

R^1 is unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_2 - C_7 heterocycloalkyl, unsubstituted or substituted $-(C_1$ - C_6 alkyl)-(C_3 - C_6 cycloalkyl), or unsubstituted or substituted $-(C_1$ - C_6 alkyl)-(C_2 - C_7 heterocycloalkyl), wherein any substituted group of R^1 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, $-N(R^4)_2$, $-OR^5$, $-CN$, $-CO_2R^5$, $-C(=O)N(R^4)_2$, $-SR^5$, $-S(=O)R^7$, $-S(=O)_2R^7$, $-NR^4C(=O)R^5$, $-NR^4SO_2R^7$, $-SO_2R^7$, or $-SO_2N(R^4)_2$;

each R^3 is independently hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl;

each R^4 is independently selected from the group consisting of hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted aryl, and unsubstituted or substituted heteroaryl;

or two R^4 are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 6-membered monocyclic heterocycle;

each R^5 is independently selected from the group consisting of hydrogen, substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted aryl, and unsubstituted or substituted heteroaryl;

each R^6 is independently hydrogen, halogen, unsubstituted or substituted C_1 - C_4 alkyl, unsubstituted or substituted C_1 - C_4 alkoxy, unsubstituted or substituted C_1 - C_4 fluoroalkyl, unsubstituted or substituted C_1 - C_4 fluoroalkoxy, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, $-CN$, $-OH$, $-CO_2R^5$, $-CH_2CO_2R^5$, $-C(=O)N(R^4)_2$, $-C(=O)N(R^4)OR^5$, $-CH_2C(=O)N(R^4)_2$, $-N(R^4)_2$, $-CH_2N(R^4)_2$, $-C(R^5)_2N(R^4)_2$, $-NR^4C(=O)R^5$, $-CH_2NR^4C(=O)R^5$, $-NR^4C(=O)N(R^5)_2$, $-NR^4C(=O)N(R^4)_2$, $C(R^5)=N(R^4)-OR^5$, $-SR^5$, $-S(=O)R^7$, $-SO_2R^7$, or $-SO_2N(R^4)_2$; and

each R^7 is independently selected from the group consisting substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heteroaryl.

[0011] Also provided herein, is a compound that is:

- 1-1:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-2:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-pyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-3:** N-(2-aminoethyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxamide;
- 1-4:** N-[(2R)-2-aminopropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxamide;
- 1-5:** N-(3-aminopropyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxamide;
- 1-6:** N-[(2S)-2-aminopropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxamide;
- 1-7:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-8:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxamide;
- 1-9:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-10:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-pyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-11:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-12:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-pyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-13:** N-(2-aminoethyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-14:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-15:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-16:** N-[(2S)-2-aminopropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-17:** N-[(2R)-2-amino-3-hydroxypropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-

[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-18: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(1-methylazetidin-3-yl)methyl]piperidine-4-carboxamide;

1-19: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;

1-20: N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-21: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}-N-[(3R)-pyrrolidin-3-yl]piperidine-4-carboxamide;

1-22: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-23: 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-24: 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-25: 3-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-[(3R)-1-methylpyrrolidin-3-yl]carbamoyl)piperidin-1-yl)-6-(trifluoromethyl)pyridine-2-carboxamide;

1-26: 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-27: 3-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-[(3S)-1-methylpyrrolidin-3-yl]carbamoyl)piperidin-1-yl)-6-(trifluoromethyl)pyridine-2-carboxamide;

1-28: N-[(2S)-2-amino-3-hydroxypropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-29: 1-[2-chloro-6-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-30: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-31: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-32: 1-(2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-33: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]-1-[4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;

1-34: 1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-35: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]-1-[5-(trifluoromethyl)pyridin-2-yl]piperidine-4-carboxamide;

1-36: 1-(4-acetyl-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-37: 1-[2-cyano-4-(difluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-38: 1-[4-cyano-2-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-39: 1-[2-cyano-4-(trifluoromethoxy)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-40: 1-(2,4-dicyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-41: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S,4S)*-4-hydroxy-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-42: 1-[3-cyano-5-(trifluoromethyl)pyridin-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-43: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-44: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(2R)-3-hydroxy-2-(methylamino)propyl]piperidine-4-carboxamide;

1-45: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2R)-2-(dimethylamino)-3-hydroxypropyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-46: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R,4R)*-4-hydroxy-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-47: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-3-hydroxy-2-(methylamino)propyl]piperidine-4-carboxamide;

1-48: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-2-(dimethylamino)-3-hydroxypropyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-51: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-52: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-

methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-53: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2,3-dihydro-1-benzofuran-7-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-54: 4-[6-(1-benzofuran-7-yl)pyridin-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-55: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{6-[2-(hydroxymethyl)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-56: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-57: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-58: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-59: 1-[2-cyano-4-(1,1-difluoroethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-60: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-61: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2,3-difluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-62: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-ethyl-1H-pyrazol-5-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-63: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-[6-(2-propoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-64: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3-fluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-65: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3,5-difluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-66: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-67: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2,2-difluoro-2H-1,3-benzodioxol-4-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-68: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2,5-difluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

- 1-69:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxy-5-fluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-70:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{6-[1-(3-methylbutyl)-1H-pyrazol-5-yl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-71:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-{6-[2-(trifluoromethoxy)phenyl]pyridin-3-yl}piperidine-4-carboxamide;
- 1-72:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxy-3-fluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-73:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-indazol-7-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-74:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-fluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-75:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)-3-fluorophenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-76:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;
- 1-77:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-{6-[2-(trifluoromethyl)phenyl]pyridin-3-yl}piperidine-4-carboxamide;
- 1-78:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-cyanophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-79:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{6-[2-(methoxymethyl)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-80:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3-methoxythiophen-2-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-81:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-82:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-83:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxythiophen-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-84:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{6-[2-(1,1-difluoroethyl)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-85:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2'-(difluoromethoxy)-[2,3'-bipyridin]-5-yl]-N-

[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-86: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{5'-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-87: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3,5-difluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-88: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-89: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-cyclopropoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-90: 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-91: 1-(4-chloro-2-cyanophenyl)-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-92: 1-(4-chloro-2-cyanophenyl)-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;

1-93: 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;

1-94: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;

1-95: 1-(2,4-dichlorophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-96: 1-(2-cyano-4-fluorophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-97: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-98: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-99: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-fluoro-2-methylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-100: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-101: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

- 1-102:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-103:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{5'-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-104:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethylphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-105:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(4-methylpyrimidin-5-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-106:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-107:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-108:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-109:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-110:** 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-111:** 1-(4-chloro-2-cyano-6-fluorophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-112:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-[6-(2-methylthiophen-3-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-113:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-cyclopropylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-114:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{ 6-[2-(difluoromethoxy)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-115:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-116:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methylfuran-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-117:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-[6-(2-propylphenyl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-118:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethyl-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-

methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-119: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxythiophen-3-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-120: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methylfuran-3-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-121: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-cyclopropylphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-122: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]-4-[6-(2-methylthiophen-3-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-123: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-124: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-125: 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-126: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-127: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-128: 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-129: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3-fluoro-2-hydroxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-130: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxy-5-methylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-131: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-132: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-133: 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-134: 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-135: 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;

1-136: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;

1-137: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{6-[2-(cyanomethyl)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-138: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-[6-(4-methylthiophen-3-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-139: 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-140: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;

1-141: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-142: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-143: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]-4-[6-(1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-144: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-145: 4-[6-(2-ethoxyphenyl)pyridin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-146: 4-[6-(2-acetylthiophen-3-yl)pyridin-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-147: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethylthiophen-3-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-148: 4-[6-(2-cyano-3-fluorophenyl)pyridin-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-149: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3-methyl-1,2-oxazol-4-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-150: 1-[3-chloro-5-(trifluoromethyl)pyridin-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-151: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{6-[2-(difluoromethyl)phenyl]pyridin-3-yl}-N-

[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-152: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-pyrrolidin-3-yl]piperidine-4-carboxamide;

1-153: 4-[6-(5-cyano-1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-154: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{5'-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-155: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-156: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethylphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-157: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-methyl-1,2-oxazol-4-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-158: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-cyclopropyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-159: 4-[4-(2-ethoxypyridin-3-yl)phenyl]-1-[2-methoxy-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-160: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-161: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-162: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-163: 1-(4-chloro-2-cyanophenyl)-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-164: 4-[4-(2-ethoxypyridin-3-yl)phenyl]-1-[2-methyl-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-165: 1-[3-chloro-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-166: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-167: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxythiophen-3-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

- 1-168:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-169:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-170:** 1-(4-chloro-2-cyanophenyl)-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-171:** 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-172:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[2'-ethyl-[2,3'-bipyridin]-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-173:** 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-174:** 4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-175:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-176:** 4-[[2,2'-bipyridin]-5-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-177:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[3'-methyl-[2,2'-bipyridin]-5-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-178:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[3'-methoxy-[2,2'-bipyridin]-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-179:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2'-methoxy-[2,3'-bipyridin]-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-180:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2'-cyclopropyl-[2,3'-bipyridin]-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-181:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2'-(difluoromethyl)-[2,3'-bipyridin]-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-182:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-cyclopropyl-1,3-oxazol-4-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-183:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-184:** ethyl (3S)-3-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-

3-yl]piperidine-4-amido}pyrrolidine-1-carboxylate;

1-185: N-[(3S)-1-acetylpyrrolidin-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-186: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-187: N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-188: N-[(3R)-1-azabicyclo[2.2.2]octan-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-189: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyquinolin-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-190: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-indol-7-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-191: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-indol-2-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-192: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-193: 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-194: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-195: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-196: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-197: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[3-(methylamino)propyl]piperidine-4-carboxamide;

1-198: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-199: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[3-(dimethylamino)propyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-200: 4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

- 1-201:** 4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-202:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-203:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-204:** 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-205:** N-(3-aminopropyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-206:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[3-(methylamino)propyl]piperidine-4-carboxamide;
- 1-207:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[3-(dimethylamino)propyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-208:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-ethyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-209:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-210:** 4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-211:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(1-methylazetidin-3-yl)methyl]piperidine-4-carboxamide;
- 1-212:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-213:** N-[(2S)-azetidin-2-yl]methyl-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-214:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-215:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-216:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-217:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(2S)-1-

methylazetididin-2-yl)methyl}piperidine-4-carboxamide;

1-218: N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-219: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-(1-methylazetididin-3-yl)piperidine-4-carboxamide;

1-220: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1-methylazetididin-3-yl)methyl]piperidine-4-carboxamide;

1-221: N-{1-azabicyclo[2.2.1]heptan-4-yl}-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-222: N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;

1-223: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-224: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-225: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-226: N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;

1-227: 4-[6-(2-ethoxyphenyl)pyridin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-228: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-229: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[[2S]-1-methylpyrrolidin-2-yl)methyl]piperidine-4-carboxamide;

1-230: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-231: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[[2R]-pyrrolidin-2-yl)methyl]piperidine-4-carboxamide;

1-232: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[[2R]-1-methylpyrrolidin-2-yl)methyl]piperidine-4-carboxamide;

1-233: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1S,3S)-3-aminocyclobutyl]piperidine-4-carboxamide;

1-234: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1R,3R)-3-aminocyclobutyl]piperidine-4-carboxamide;

1-235: N-{[(2S)-azetidin-2-yl]methyl}-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-236: N-[(2R)-azetidin-2-yl]methyl-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-237: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R,4S)-4-fluoropyrrolidin-3-yl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-238: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R,4R)-4-fluoropyrrolidin-3-yl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-239: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1S,3S)-3-(dimethylamino)cyclobutyl]piperidine-4-carboxamide;

1-240: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1R,3R)-3-(dimethylamino)cyclobutyl]piperidine-4-carboxamide;

1-241: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[[2S]-1-methylazetidin-2-yl]methyl]piperidine-4-carboxamide;

1-242: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R,4S)-4-fluoro-1-methylpyrrolidin-3-yl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-243: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R,4R)-4-fluoro-1-methylpyrrolidin-3-yl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-244: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(2R)-1-methylazetidin-2-yl]methyl]piperidine-4-carboxamide;

1-245: N-[2-(azetidin-1-yl)ethyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-246: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S,4S)-4-fluoropyrrolidin-2-yl]methyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-247: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S,4R)-4-fluoropyrrolidin-2-yl]methyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-248: N-[2-(azetidin-1-yl)ethyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-249: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S,4S)-4-fluoro-1-methylpyrrolidin-2-yl]methyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-250: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S,4R)-4-fluoro-1-methylpyrrolidin-2-

yl)methyl}-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-251: rac-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1S,2S,4R)-7-methyl-7-azabicyclo[2.2.1]heptan-2-yl]piperidine-4-carboxamide;

1-252: rac-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(1S,2S,4R)-7-methyl-7-azabicyclo[2.2.1]heptan-2-yl]piperidine-4-carboxamide;

1-253: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1,3-dimethylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-254: N-[2-(dimethylamino)ethyl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-255: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-256: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-257: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-258: rac-N-[(1R,2S)-2-aminocyclopropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-259: rac-N-[(1R,2R)-2-aminocyclopropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-260: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2R)-1-(dimethylamino)propan-2-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-261: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-262: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-{3-[(dimethylamino)methyl]oxetan-3-yl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-263: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-{[1-(dimethylamino)cyclopropyl]methyl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-264: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S,4S)-4-(dimethylamino)oxolan-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-265: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-{2-[ethyl(methyl)amino]ethyl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-266: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-{2-[cyclopropyl(methyl)amino]ethyl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-267: rac-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S,4R)-4-methoxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-268: rac-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R,4R)-4-methoxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-269: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R,4R)-4-hydroxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-270: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S,4S)-4-hydroxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-271: N-[2-(dimethylamino)ethyl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-272: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-273: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-274: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-275: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2R)-2-(dimethylamino)-3-hydroxypropyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-276: N-[2-(dimethylamino)ethyl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-277: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[[2S,4S]-4-fluoro-1-methylpyrrolidin-2-yl]methyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-278: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N- f [[2S,4R)-4-fluoro-1-methylpyrrolidin-2-yl]methyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-279: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-{ [(2S)-4,4-difluoro-1-methylpyrrolidin-2-yl]methyl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-280: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2R)-4,4-difluoro-1-methylpyrrolidin-2-yl]methyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-281: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-282: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-283: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-

(methylamino)ethyl]piperidine-4-carboxamide;

1-284: N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;

1-285: N-[2-(dimethylamino)ethyl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-286: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-287: 4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-288: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-289: N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;

1-290: 4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-291: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]piperidine-4-carboxamide;

1-292: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-293: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-294: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-295: N-[(2S)-1-(dimethylamino)propan-2-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-296: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-297: N-[(2S)-1-(dimethylamino)propan-2-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-298: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;

1-299: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

- 1-300:** N-[2-(dimethylamino)ethyl]-4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;
- 1-301:** 4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-302:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-303:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-304:** N-[2-(dimethylamino)ethyl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-305:** 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-306:** 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;
- 1-307:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl} - N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;
- 1-308:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-309:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-310:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-311:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-312:** N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;
- 1-313:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-314:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;
- 1-315:** 4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;
- 1-316:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-

[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;

1-317: N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;

1-318: 1-(2,4-dichlorophenyl)-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-319: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-320: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;

1-321: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;

1-322: 1-(2,4-dichlorophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;

1-323: 1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;

1-324: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-325: 1-(4-chloro-2-cyanophenyl)-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-326: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;

1-327: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(2S)-4-methylmorpholin-2-yl]methylpiperidine-4-carboxamide;

1-328: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-4-methylmorpholin-3-yl]methylpiperidine-4-carboxamide;

1-329: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-4-methylmorpholin-3-yl]methylpiperidine-4-carboxamide;

1-330: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(2R)-4-methylmorpholin-2-yl]methylpiperidine-4-carboxamide;

1-331: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidine-4-carboxamide;

1-332: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-2-(dimethylamino)propyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-333: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2R)-2-(dimethylamino)propyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-334: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2R)-2-(dimethylamino)-3-hydroxypropyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-335: N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;

2-1: 2-{4-[2-(dimethylamino)ethoxy]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-1-yl}-5-(trifluoromethyl)benzonitrile;

2-2: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl N-[2-(dimethylamino)ethyl]carbamate;

2-3: 3-amino-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}propanamide;

2-4: 3-amino-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}propanamide;

2-5: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)propanamide;

2-6: (3R)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}pyrrolidine-3-carboxamide;

2-7: (3R)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;

2-8: (3S)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}pyrrolidine-3-carboxamide;

2-9: (3S)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;

2-10: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(dimethylamino)propanamide;

2-11: (2S)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}-1-methylpyrrolidine-2-carboxamide;

2-12: (3R)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamate;

2-13: (3S)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamate;

2-14: 2-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-[(2-hydroxyethyl)amino]piperidin-1-yl)-5-

(trifluoromethyl)benzonitrile;

2-15: 2-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4- { [2-(methylamino)ethyl]amino}piperidin-1-yl)-5-(trifluoromethyl)benzonitrile;

2-16: N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}-1-methylazetidine-3-carboxamide;

2-17: (2R)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}-1-methylpyrrolidine-2-carboxamide;

2-18: (3S)-1-methylpyrrolidin-3-yl N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamate;

2-19: 1-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-[(3R)-1-methylpyrrolidin-3-yl]urea;

2-20: 1-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-[(3S)-1-methylpyrrolidin-3-yl]urea;

2-21: (3R)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)pyrrolidine-1-carboxamide;

2-22: (3R)-1-methylpyrrolidin-3-yl N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamate;

2-23: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)azetidine-1-carboxamide;

2-24: (3S)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)pyrrolidine-1-carboxamide;

2-25: N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-1-methylazetidine-3-carboxamide;

2-26: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-2-(dimethylamino)acetamide;

2-27: (3S)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;

2-28: (3S)-N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-1-methylpyrrolidine-3-carboxamide;

2-29: (3R)-N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-1-methylpyrrolidine-3-carboxamide;

2-30: N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-3-(dimethylamino)propanamide;

- 2-31:** N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-2-(dimethylamino)acetamide;
- 2-32:** N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-1-methylazetidine-3-carboxamide;
- 2-33:** (3R)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;
- 2-34:** N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-2-(dimethylamino)acetamide;
- 2-35:** N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-1-methylazetidine-3-carboxamide;
- 2-36:** N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamide;
- 2-37:** N-[1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl]-1-methylazetidine-3-carboxamide;
- 2-38:** N-[1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl]-2-(dimethylamino)acetamide;
- 2-39:** (3S)-N-[1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl]-1-methylpyrrolidine-3-carboxamide;
- 2-40:** N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-1-methylazetidine-3-carboxamide;
- 2-41:** (3R)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}carbamate;
- 2-42:** (3S)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}carbamate;
- 2-43:** 1-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-3-[(3S)-1-methylpyrrolidin-3-yl]urea;
- 2-44:** (3S)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}pyrrolidine-3-carboxamide;
- 2-45:** N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-1-methylazetidine-3-carboxamide;
- 2-46:** (3S)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide; and
- 2-47:** (3R)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-

yl]piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;

2-48: 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-3-[(3R)-1-methylpyrrolidin-3-yl]urea;

2-49: (3S)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;

2-50: (3R)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;

2-51: 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[(3S)-1-methylpyrrolidin-3-yl]urea;

2-52: 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[(3R)-1-methylpyrrolidin-3-yl]urea;

2-53: (3S)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;

2-54: (3R)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;

2-55: 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[(3S)-1-methylpyrrolidin-3-yl]urea;

2-56: 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[(3R)-1-methylpyrrolidin-3-yl]urea;

2-57: 2-(dimethylamino)ethyl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;

2-58: 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[2-(dimethylamino)ethyl]urea;

2-59: 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(1-methylazetidin-3-yl)urea;

2-60: (3R)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;

2-61: N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-2-(dimethylamino)acetamide;

2-62: N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamide;

2-63: N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(methylamino)propanamide;

2-64: N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamide;

or a pharmaceutically acceptable salt thereof.

[0012] Also provided herein is a pharmaceutical composition comprising a compound as defined in the claims, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable excipient.

[0013] Also provided herein is a compound as defined in the claims, or a pharmaceutically acceptable salt thereof, for use in the treatment of Cushing's syndrome, ectopic Cushing's syndrome, congenital adrenal hyperplasia (CAH), or for reducing the secretion of adrenocorticotrophic hormone (ACTH) in a mammal.

[0014] In some embodiments, the pharmaceutical composition is formulated for administration to a mammal by intravenous administration, subcutaneous administration, oral administration, inhalation, nasal administration, dermal administration, or ophthalmic administration. In some embodiments, the pharmaceutical composition is formulated for administration to a mammal by oral administration. In some embodiments, the pharmaceutical composition is in the form of a tablet, a pill, a capsule, a liquid, a suspension, a gel, a dispersion, a solution, an emulsion, an ointment, or a lotion. In some embodiments, the pharmaceutical composition is in the form of a tablet, a pill, or a capsule.

[0015] In any of the aforementioned aspects are further embodiments in which the effective amount of the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is: (a) systemically administered to the mammal; and/or (b) administered orally to the mammal; and/or (c) intravenously administered to the mammal; and/or (d) administered by inhalation; and/or (e) administered by nasal administration; or and/or (f) administered by injection to the mammal; and/or (g) administered topically to the mammal; and/or (h) administered by ophthalmic administration; and/or (i) administered rectally to the mammal; and/or (j) administered non-systemically or locally to the mammal.

[0016] In any of the aforementioned aspects are further embodiments comprising single administrations of the effective amount of the compound, including further embodiments in which the compound is administered once a day to the mammal or the compound is administered to the mammal multiple times over the span of one day. In some embodiments, the compound is administered on a continuous dosing schedule. In some embodiments, the compound is administered on a continuous daily dosing schedule.

[0017] In any of the embodiments disclosed herein, the mammal is a human.

[0018] In some embodiments, compounds provided herein are orally administered to a human.

[0019] Articles of manufacture, which include packaging material, a compound of Formula (I), or a pharmaceutically acceptable salt thereof, within the packaging material, and a label that indicates that the compound or composition, or pharmaceutically acceptable salt, tautomers, pharmaceutically acceptable N-oxide, is used for modulating one or more subtype melanocortin receptor proteins, or for use in the treatment, prevention or amelioration of one or more symptoms of a disease or condition that would benefit from modulating one or more subtype melanocortin receptor proteins, are provided.

[0020] Other objects, features and advantages of the compounds, methods and compositions described herein will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments, are given by way of illustration only, since various changes and modifications within the scope of the instant disclosure will become apparent to those skilled in the art from this detailed description.

DETAILED DESCRIPTION OF THE INVENTION

[0021] The present invention is defined by the claims. Any subject matter falling outside the scope of the claims is provided for information purposes only. Any references in the description to methods for treatment of the human or animal body by therapy refer to the compounds and pharmaceutical compositions of the present invention for use in methods for the treatment of the human or animal body by therapy.

[0022] Adrenocorticotrophic hormone (ACTH) is a 39 amino acid peptide synthesized by anterior pituitary corticotrophic cells by proteolytic cleavage of the proopiomelanocortin hormone (POMC). ACTH is the primary regulator of adrenal glucocorticoid (GC; cortisol in humans and most other species; corticosterone in rodents) synthesis and secretion. As the central actor in this hypothalamic-pituitary-adrenal (HPA) axis, ACTH is secreted by the pituitary in response to stressful stimuli and acts at the adrenal gland to stimulate the synthesis and secretion of cortisol. This stimulation is mediated through a highly specific G protein-coupled receptor (GPCR) which is expressed almost uniquely in the adrenal cortex. The receptor is the melanocortin 2 receptor (MC2R), and, along with ACTH, is part of the larger melanocortin system.

[0023] The melanocortin system comprises a family of five GPCRs (MC1R, MC2R, MC3R, MC4R, and MC5R); their natural agonists, the melanocortin peptides α -, β -, and γ -melanocyte-stimulating hormone (α -MSH, β -MSH, and γ -MSH) and ACTH; and endogenous melanocortin antagonists agouti and agouti-related protein (AGRP). The melanocortin receptors (MCRs) have different selectivities for endogenous agonist and antagonist peptides and are expressed in diverse tissues where they serve varied and discrete physiological functions (Gantz, I. and T.M. Fong, Am. J. Physiol. Endocrinol. Metab., 284: E468-E474, 2003).

[0024] It is possible to selectively modulate any one of the MCRs, or combinations thereof. In

some embodiments, selectively modulating any one of the MCRs relative to the other MCRs, or combinations thereof, is useful in a variety of clinical applications. In some embodiments, selectively modulating any one of the MCRs relative to the other MCRs, or combinations thereof, reduces unwanted side effects in a variety of clinical applications. In one aspect, compounds described herein are antagonists of MC2R. In some embodiments, compounds described herein are selective antagonists for MC2R relative to other MCRs.

[0025] MC2R is a highly selective receptor for ACTH. Although ACTH can activate all five MCRs, at physiological levels, the sensitivity of the other receptors is not high enough to be activated, and ACTH selectively activates MC2R. Importantly, the other naturally occurring agonists α -MSH, β -MSH, and γ -MSH have no affinity for MC2R (Gantz, I. and T.M. Fong, Am. J. Physiol. Endocrinol. Metab., 284: E468-E474, 2003). The major function of MC2R is to stimulate the fasciculata cells of the adrenal cortex to synthesize and secrete cortisol. MC2R requires the GPCR accessory protein MRAP (melanocortin 2 receptor protein) to be successfully secreted to the cell surface and as well as to function. MRAP is a small protein with a single transmembrane domain that forms an antiparallel homodimer in stable complex with MC2R and is necessary for both cell surface expression of MC2R and its ability to bind ACTH. MRAP can bind to any of the MCRs and affect their activities, but is only essential for MC2R activity. Binding of ACTH to the MC2R/MRAP complex on adrenal cortical cells activates G_s to elevate intracellular cAMP levels which in turn stimulates cortisol synthesis and secretion by regulating multiple steps in the steroidogenic pathway.

[0026] Cushing's syndrome is a rare disorder characterized by chronic, excess glucocorticoid exposure. Clinical signs of Cushing's syndrome include growth of fat pads (collarbone, back of neck, face and trunk), excessive sweating, dilation of capillaries, thinning of the skin, muscle weakness, hirsutism, depression/anxiety, hypertension, osteoporosis, insulin resistance, hyperglycemia, heart disease, and a range of other metabolic disturbances resulting in high morbidity. If inadequately controlled in its severe forms, Cushing's syndrome is associated with high mortality. Although glucocorticoid excess can sometimes be ACTH independent, for example from excessive autonomous secretion of cortisol from a hyperfunctioning adrenal adenoma, carcinoma, or steroid abuse, about 60-80% of all cases are ACTH dependent Cushing's syndrome, known as Cushing's disease. Cushing's disease is caused by microadenomas of pituitary corticotrophic cells that secrete excess ACTH. Corticotroph adenomas are small, usually slow growing, benign tumors that normally come to clinical attention as a result of the effects of glucocorticoid excess, rather than because of the physical effects of an expanding tumor. First line treatments for Cushing's disease are surgical and involve removal of either the ACTH-secreting tumor in the pituitary or the adrenal glands themselves. As surgery is often unsuccessful, contraindicated, or delayed, medical therapy for these patients becomes necessary. Current treatment options include inhibitors of steroid synthesis enzymes that can prevent the production of cortisol and improve symptoms, but these treatments also induce a host of unwanted side effects due to the accumulation of other steroid products. In one aspect, an MC2R antagonist is used in the treatment of Cushing's syndrome. In some embodiments, an MC2R antagonist is used in the treatment of Cushing's disease. In some embodiments, glucocorticoid excess is ACTH independent. In some

embodiments, glucocorticoid excess is ACTH dependent.

[0027] Ectopic ACTH syndrome, or ectopic Cushing's syndrome or disease, is essentially the same as Cushing's disease, except that the underlying tumor expressing ACTH is outside the pituitary gland. In some embodiments, the tumors are small carcinoid tumors that occur anywhere in the lungs or gastrointestinal tract. In some embodiments, an MC2R antagonist is used in the treatment of ectopic ACTH syndrome.

[0028] Congenital adrenal hyperplasia (CAH) is characterized by a reduction or loss of cortisol synthesis and excessive ACTH and corticotropin-releasing hormone. CAH can result from a variety of genetic defects in the adrenal steroidal biosynthesis pathway. In some embodiments, CAH is due to a mutation in 21 β -hydroxylase. The lack of cortisol removes the negative feedback to the pituitary which leads to excessive ACTH secretion. The resulting excessive adrenal stimulation causes overproduction of steroid precursors which also have negative consequences (e.g., hyperandrogenism). Administration of replacement glucocorticoids typically does not adequately suppress ACTH without also causing Cushing's-like symptoms. In some embodiments, an MC2R antagonist is for use in the treatment of CAH.

[0029] In addition to Cushing's disease, Ectopic ACTH syndrome and CAH it has also been hypothesized that there might be a role for an MC2R antagonist in the treatment of ACTH driven adrenal tumors, Functional Adrenal Hyperandrogenism (FAH), stress disorders, psychiatric disorders, type 2 diabetes and septic shock. In some embodiments, an MC2R antagonist is for use in the treatment of ACTH driven adrenal tumors. In some embodiments, an MC2R antagonist is for use in the treatment of Functional Adrenal Hyperandrogenism. In some embodiments, an MC2R antagonist is for use in the treatment of stress disorders. In some embodiments, an MC2R antagonist is for use in the treatment of psychiatric disorders. In some embodiments, an MC2R antagonist is for use in the treatment of type 2 diabetes. In some embodiments, an MC2R antagonist is for use in the treatment of septic shock.

[0030] In some embodiments, an MC2R antagonist is for use in the treatment of septic shock.


[0031] In some embodiments, compounds described herein are amenable to administration to a mammal in need of treatment with an MC2R antagonist.

Compounds

[0032] The compounds are defined in the claims.

[0033] Compounds of Formula (I), including pharmaceutically acceptable salts thereof, are melanocortin receptor modulators. In some embodiments, the compounds of Formula (I), including pharmaceutically acceptable salts, , are MC2R modulators. In some embodiments, the MC2R modulators are MC2R antagonists.

[0034] In one aspect, provided herein is a compound of Formula (I), or a pharmaceutically acceptable salt thereof:



Formula (I)

wherein:

R^A is unsubstituted or substituted phenyl, unsubstituted or substituted monocyclic 6-membered heteroaryl, or unsubstituted or substituted monocyclic 5-membered heteroaryl, wherein if R^A is substituted then R^A is substituted with 1, 2, 3 or 4 groups selected from R^a , R^b , and R^c ;

R^a , R^b , and R^c are independently selected from the group consisting of hydrogen, halogen, -OR⁴, -CN, -N(R⁴)₂, -C(=O)R⁷, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₂-C₇ heterocycloalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heteroaryl, wherein any substituted group of R^a , R^b , and R^c is substituted with one or more R⁶ groups;

or one R^a and one R^b , when present on adjacent atoms of R^A , are taken together with the intervening atoms connecting R^a to R^b to form a 5- to 6-membered monocyclic carbocycle or 5- to 6-membered monocyclic heterocycle, wherein the carbocycle or heterocycle is unsubstituted or substituted with one or more R⁶ groups;

wherein, if R^a , R^b , or R^c is attached to the N atom of a heteroaryl, then it is hydrogen, -C(=O)R⁷, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, or unsubstituted or substituted C₂-C₇ heterocycloalkyl;

R^B is an unsubstituted or substituted phenyl or unsubstituted or substituted monocyclic 6-membered heteroaryl, wherein if R^B is substituted then R^B is substituted with 1, 2, 3 or 4 groups selected from R^d, R^e, and R^f;

R^d, R^e, and R^f are independently selected from the group consisting of hydrogen, halogen, -OR⁴, -CN, -N(R⁴)₂, -C(=O)R⁷, -C(=O)N(R⁴)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₂-C₇ heterocycloalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heteroaryl, wherein any substituted group of R^d, R^e, and R^f is substituted with one or more R⁶ groups;

wherein, if R^d , R^e , or R^f is attached to the N atom of a heteroaryl, then it is hydrogen, -C(=O) R^7 , unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl;

X^1 is CR^{11} or N;

X^2 is CR^{12} or N;

X^3 is CR^{13} or N;

X^4 is CR^{14} or N;

R^{11} , R^{12} , R^{13} , and R^{14} are each independently hydrogen, halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, -CN, -OR⁴, -SR⁴, -CO₂R⁴, -C(=O)N(R⁴)₂, or -N(R⁴)₂;

M is -(C=O)-, -NR³-, -O-, -S-, -SO₂-, *-NR³-(C=O)-, *(C=O)-NR³-, *-O-(C=O)NR³-, *-NR³-(C=O)O-, -NR³-(C=O)NR³-, *-NR³(SO₂)-, *-SO₂NR³-, or 5-membered heterocycle, wherein * indicates the attachment point to R¹;

R¹ is unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_2 - C_7 heterocycloalkyl, unsubstituted or substituted -(C_1 - C_6 alkyl)-(C_3 - C_6 cycloalkyl), or unsubstituted or substituted -(C_1 - C_6 alkyl)-(C₂-C₇ heterocycloalkyl), wherein any substituted group of R¹ is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, -N(R⁴)₂, -OR⁵, -CN, -CO₂R⁵, -C(=O)N(R⁴)₂, -SR⁵, -S(=O)R⁷, -S(=O)₂R⁷, -NR⁴C(=O)R⁵, -NR⁴SO₂R⁷, -SO₂R⁷, or -SO₂N(R⁴)₂;

each R³ is independently hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl;

each R⁴ is independently selected from the group consisting of hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted aryl, and unsubstituted or substituted heteroaryl;

or two R^4 are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 6-membered monocyclic heterocycle;

each R^5 is independently selected from the group consisting of hydrogen, substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted aryl, and unsubstituted or substituted heteroaryl;

each R^6 is independently hydrogen, halogen, unsubstituted or substituted C_1 - C_4 alkyl, unsubstituted or substituted C_1 - C_4 alkoxy, unsubstituted or substituted C_1 - C_4 fluoroalkyl, unsubstituted or substituted C_1 - C_4 fluoroalkoxy, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, $-CN$, $-OH$, $-CO_2R^5$, $-CH_2CO_2R^5$, $-C(=O)N(R^4)_2$, $-C(=O)N(R^4)OR^5$, $-CH_2C(=O)N(R^4)_2$, $-N(R^4)_2$, $-CH_2N(R^4)_2$, $-C(R^5)_2N(R^4)_2$, $-NR^4C(=O)R^5$, $-CH_2NR^4C(=O)R^5$, $-NR^4C(=O)N(R^5)_2$, $-NR^4C(=O)N(R^4)_2$, $C(R^5)=N(R^4)-OR^5$, $-SR^5$, $-S(=O)R^7$, $-SO_2R^7$, or $-SO_2N(R^4)_2$; and

each R^7 is independently selected from the group consisting substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heteroaryl.

[0035] In some embodiments, M is $-C(=O)-$, $-NR^3-$, $-O-$, $-S-$, $-SO_2-$, $^*-NR^3-(C=O)-$, $^*-(C=O)-NR^3-$, $^*-NR^3SO_2-$, or 5-membered heterocycle, wherein * indicates the attachment point to R^1 . In some embodiments, M is $-C(=O)-$, $-NR^3-$, $-O-$, $-S-$, $-SO_2-$, $^*-(C=O)-NR^3-$, $^*-NR^3-(C=O)-$, $^*-NR^3SO_2-$, or 5-membered heteroaryl, wherein * indicates the attachment point to R^1 .

[0036] In some embodiments, M is $-O-$, $^*-NR^3-C(=O)-$, $-S-$, $-SO_2-$, $^*-NR^3SO_2-$, or 5-membered heteroaryl, wherein * indicates the attachment point to R^1 .

[0037] In some embodiments, M is 5-membered heterocycle. In some embodiments, M is 5-membered heteroaryl. In some embodiments, M is furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, triazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, or thiadiazolyl. In some embodiments, M is oxazolyl, imidazolyl, or triazolyl.

[0038] In some embodiments, M is $-NR^3-$, $-O-$, $^*-NR^3-(C=O)-$, $^*-(C=O)-NR^3-$, $^*-O-(C=O)NR^3-$, $^*-NR^3-(C=O)O-$, or $-NR^3-(C=O)NR^3-$, wherein * indicates the attachment point to R^1 . In some embodiments, M is $^*-NR^3-(C=O)-$, $^*-(C=O)-NR^3-$, $^*-O-(C=O)NR^3-$, $^*-NR^3-(C=O)O-$, or $-NR^3-(C=O)NR^3-$, wherein * indicates the attachment point to R^1 . In some embodiments, M is $^*-NR^3-(C=O)-$ or $^*-(C=O)-NR^3-$, wherein * indicates the attachment point to R^1 .


[0039] In some embodiments, M is $^{*}-(C=O)-NR^3-$, wherein * indicates the attachment point to R^1 . In some embodiments, M is $^{*}-NR^3-(C=O)-$, wherein * indicates the attachment point to R^1 .

[0040] In some embodiments, each R^3 is independently hydrogen or unsubstituted or substituted C_1-C_6 alkyl. In some embodiments, each R^3 is independently hydrogen or C_1-C_6 alkyl. In some embodiments, each R^3 is independently hydrogen, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, or $-CH(CH_3)_2$. In some embodiments, each R^3 is independently hydrogen or $-CH_3$. In some embodiments, each R^3 is $-CH_3$. In some embodiments, each R^3 is hydrogen.

[0041] In some embodiments, M is $-NR^3-$, $-O-$, $^{*}-NR^3-(C=O)-$, $^{*}-(C=O)-NR^3-$, $^{*}-O-(C=O)NR^3-$, $^{*}-NR^3-(C=O)O-$, or $-NR^3-(C=O)NR^3-$, wherein * indicates the attachment point to R^1 ; and each R^3 is independently hydrogen, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, or $-CH(CH_3)_2$. In some embodiments, M is $^{*}-NR^3-(C=O)-$ or $^{*}-(C=O)-NR^3-$, wherein * indicates the attachment point to R^1 ; and R^3 is hydrogen.

[0042] In some embodiments, M comprises an N atom and the N atom of M is attached to R^1 , wherein R^1 also comprises an N atom. In some embodiments, the N atom of M is connected to the N atom of R^1 through one carbon atom, two carbon atoms, three carbon atoms, or four carbon atoms. In some embodiments, the N atom of M is connected to the N atom of R^1 through a two carbon atom spacer. In some embodiments, the N atom of R^1 is part of an unsubstituted or substituted aliphatic alkyl chain. In some embodiments, the N atom of R^1 is part of an unsubstituted or substituted cyclic ring. In some embodiments, the N-containing cyclic ring of R^1 is an unsubstituted or substituted monocyclic C_2-C_7 heterocycloalkyl or unsubstituted or substituted bicyclic C_2-C_7 heterocycloalkyl.


[0043] In some embodiments, the compound has the structure of Formula (IIa), or a pharmaceutically acceptable salt thereof:



Formula (IIa).

[0044] In some embodiments, the compound has the structure of Formula (IIb), or a pharmaceutically acceptable salt thereof:

—A—



Formula (IIb).

[0045] R^A is unsubstituted or substituted monocyclic 6-membered heteroaryl, unsubstituted or substituted phenyl, or unsubstituted or substituted monocyclic 5-membered heteroaryl, wherein if R^A is substituted then R^A is substituted with 1, 2, 3 or 4 groups selected from R^a , R^b , and R^c . In some embodiments, R^A is unsubstituted or substituted monocyclic 6-membered heteroaryl, unsubstituted or substituted phenyl, or unsubstituted or substituted monocyclic 5-membered heteroaryl, wherein if R^A is substituted then R^A is substituted with 1, 2, or 3 groups selected from R^a , R^b , and R^c . In some embodiments, R^A is unsubstituted or substituted monocyclic 6-membered heteroaryl, unsubstituted or substituted phenyl, or unsubstituted or substituted monocyclic 5-membered heteroaryl, wherein if R^A is substituted then R^A is substituted with 1 or 2 groups selected from R^a , R^b , and R^c . In some embodiments, R^A is unsubstituted or substituted monocyclic 6-membered heteroaryl, unsubstituted or substituted phenyl, or unsubstituted or substituted monocyclic 5-membered heteroaryl, wherein if R^A is substituted then R^A is substituted with 1 R^a , R^b , or R^c group.

[0046] In some embodiments, R^A is substituted with 1, 2, 3 or 4 groups selected from R^a , R^b , and R^c . In some embodiments, R^A is substituted with 1, 2, or 3 groups selected from R^a , R^b , and R^c . In some embodiments, R^A is substituted with 1 or 2 groups selected from R^a , R^b , and R^c . In some embodiments, R^A is substituted with 1 group selected from R^a , R^b , and R^c . In some embodiments, R^A is substituted with one R^a . In some embodiments, R^A is substituted with one R^c .

[0047] In some embodiments, R^A is unsubstituted or substituted pyrrolyl, unsubstituted or substituted furanyl, unsubstituted or substituted thienyl, unsubstituted or substituted oxazolyl, unsubstituted or substituted thiazolyl, unsubstituted or substituted imidazolyl, unsubstituted or substituted pyrazolyl, unsubstituted or substituted triazolyl, unsubstituted or substituted tetrazolyl, unsubstituted or substituted isoxazolyl, unsubstituted or substituted isothiazolyl, unsubstituted or substituted oxadiazolyl, or unsubstituted or substituted thiadiazolyl, unsubstituted or substituted pyridinyl, unsubstituted or substituted pyrimidinyl, unsubstituted or substituted pyrazinyl, unsubstituted or substituted pyridazinyl, unsubstituted or substituted triazinyl, or unsubstituted or substituted phenyl, wherein if R^A is substituted then R^A is substituted with 1, 2, or 3 groups selected from R^a , R^b , and R^c . In some embodiments, if R^A is

substituted then R^A is substituted with 1 or 2 groups selected from R^a , R^b , and R^c . In some embodiments, if R^A is substituted then R^A is substituted with 1 R^a , R^b , or R^c group.

[0048] In some embodiments, R^A is unsubstituted or substituted monocyclic 5-membered heteroaryl, wherein if R^A is substituted then R^A is substituted with 1, 2 or 3 groups selected from R^a , R^b , and R^c . In some embodiments, R^A is unsubstituted or substituted monocyclic 5-membered heteroaryl, wherein if R^A is substituted then R^A is substituted with 1 or 2 groups selected from R^a , R^b , and R^c . In some embodiments, R^A is unsubstituted or substituted monocyclic 5-membered heteroaryl containing one heteroatom selected from N, O, and S, wherein if R^A is substituted then R^A is substituted with 1 or 2 groups selected from R^a , R^b , and R^c . In some embodiments, R^A is unsubstituted or substituted monocyclic 5-membered heteroaryl containing one heteroatom selected from N, O, and S, wherein if R^A is substituted then R^A is substituted with 1 or 2 groups selected from R^a , R^b , and R^c . In some embodiments, R^A is unsubstituted or substituted monocyclic 5-membered heteroaryl containing one heteroatom selected from N, O, and S, wherein if R^A is substituted then R^A is substituted with 1 R^a , R^b , or R^c group.


[0049] In some embodiments, R^A is unsubstituted or substituted pyrrolyl, unsubstituted or substituted furanyl, unsubstituted or substituted thienyl, unsubstituted or substituted oxazolyl, unsubstituted or substituted thiazolyl, unsubstituted or substituted imidazolyl, unsubstituted or substituted pyrazolyl, unsubstituted or substituted triazolyl, unsubstituted or substituted tetrazolyl, unsubstituted or substituted isoxazolyl, unsubstituted or substituted isothiazolyl, unsubstituted or substituted oxadiazolyl, or unsubstituted or substituted thiadiazolyl, wherein if R^A is substituted then R^A is substituted with 1, 2, or 3 groups selected from R^a , R^b , and R^c . In some embodiments, if R^A is substituted then R^A is substituted with 1 or 2 groups selected from R^a , R^b , and R^c . In some embodiments, if R^A is substituted then R^A is substituted with 1 R^a , R^b , or R^c group.

[0050] In some embodiments, R^A is unsubstituted or substituted pyrrolyl, unsubstituted or substituted furanyl, unsubstituted or substituted thienyl, unsubstituted or substituted pyrazolyl, unsubstituted or substituted isoxazolyl, wherein if R^A is substituted then R^A is substituted with 1, 2, or 3 groups selected from R^a , R^b , and R^c . In some embodiments, if R^A is substituted then R^A is substituted with 1 or 2 groups selected from R^a , R^b , and R^c . In some embodiments, if R^A is substituted then R^A is substituted with 1 R^a , R^b , or R^c group.


[0051] In some embodiments, R^A is unsubstituted or substituted pyrrolyl, unsubstituted or substituted furanyl, or unsubstituted or substituted thienyl, wherein if R^A is substituted then R^A is substituted with 1, 2, or 3 groups selected from R^a , R^b , and R^c . In some embodiments, if R^A

is substituted then R^A is substituted with 1 or 2 groups selected from R^a , R^b , and R^c . In some embodiments, if R^A is substituted then R^A is substituted with 1 R^a , R^b , or R^c group.

[0052] In some embodiments, R^A is




or R^A is




where R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.

[0053] In some embodiments, R^A is




where R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.

[0054] In some embodiments, R^A is




where R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.


[0055] In some embodiments, R^A is




or




In some embodiments, R^A is




[0056] In some embodiments, R^A is




where R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl. In some embodiments, R^A is




In some embodiments, R^A is




In some embodiments, R^A is




In some embodiments, R^A is



[0057] In some embodiments, R^A is




or R^A is




where R^C is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.

[0058] In some embodiments, R^A is




or R^A is



where R^C is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.


[0059] In some embodiments, R^A is




where R^C is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl. In some embodiments, R^C is hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl. In some embodiments, R^C is hydrogen or unsubstituted or substituted C_1 - C_6 alkyl. In some embodiments, R^C is hydrogen or C_1 - C_6 alkyl. In some embodiments, R^C is hydrogen, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)CH_2CH_3$, $-C(CH_3)_3$, $-CH_2CH_2CH_2CH_2CH_3$, or $-CH_2CH_2CH(CH_3)_2$. In some embodiments, R^C is hydrogen, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, or $-CH(CH_3)_2$. In some embodiments, R^C is hydrogen or $-CH_3$. In some embodiments, R^C is $-CH_3$. In some embodiments, R^C is hydrogen. In some embodiments, R^C is unsubstituted or substituted C_1 - C_6 alkyl. In some embodiments, R^C is C_1 -

C₆ alkyl. In some embodiments, R^c is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)CH₂CH₃, -C(CH₃)₃, -CH₂CH₂CH₂CH₂CH₃, or -CH₂CH₂CH(CH₃)₂. In some embodiments, R^c is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, or -CH(CH₃)₂. In some embodiments, R^c is hydrogen, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)CH₂CH₃, -C(CH₃)₃, -CH₂CH₂CH(CH₃)₂, or unsubstituted C₃-C₆ cycloalkyl.


[0060] In some embodiments, R^A is




Y¹ is NR^c, O, or S; Y² and Y³ are independently CH, CR^a, CR^b, or N; and R^c is hydrogen, -C(=O)R⁷, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, or unsubstituted or substituted C₂-C₇ heterocycloalkyl. In some embodiments, R^A is



Y¹ is NR^c; Y² and Y³ are independently CH, CR^a, CR^b, or N; and R^c is hydrogen, -C(=O)R⁷, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, or unsubstituted or substituted C₂-C₇ heterocycloalkyl. In some embodiments, R^A is




Y¹ is NR^c; Y² and Y³ are independently CH, CR^a, or CR^b; and R^c is hydrogen, -C(=O)R⁷, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, or unsubstituted or substituted C₂-C₇ heterocycloalkyl. In some embodiments, R^A is




Y¹ is NR^c; Y² and Y³ are each CH; and R^c is hydrogen, -C(=O)R⁷, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, or unsubstituted or substituted C₂-C₇ heterocycloalkyl.


[0061] In some embodiments, R^A is




In some embodiments, R^A is




[0062] In some embodiments, R^A is




Y^1 is NR^c , O, or S; Y^2 and Y^3 are independently CH, CR^a , CR^b , or N; and R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl. In some embodiments, R^A is



Y^1 is NR^c ; Y^2 and Y^3 are independently CH, CR^a , CR^b , or N; and R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl. In some embodiments, R^A is




Y^1 is NR^c ; Y^2 and Y^3 are independently CH, CR^a , or CR^b ; and R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl. In some embodiments, R^A is



Y^1 is NR^c ; Y^2 and Y^3 are each CH; and R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.


[0063] In some embodiments, the compound has the structure of Formula (III), or a pharmaceutically acceptable salt thereof:




Formula (III)

wherein: Y^1 is NR^c , O, or S; Y^2 and Y^3 are independently CH, CR^a , CR^b , or N; and R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.

[0064] In some embodiments, the compound has the structure of Formula (IIIa), or a pharmaceutically acceptable salt thereof:






Formula (IIIa)

wherein: Y^1 is NR^c , O, or S; Y^2 and Y^3 are independently CH, CR^a , CR^b , or N; and R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.

[0065] In some embodiments, the compound has the structure of Formula (IIIb), or a pharmaceutically acceptable salt thereof:



Formula (IIIb)


wherein: Y^1 is NR^c , O, or S; Y^2 and Y^3 are independently CH, CR^a , CR^b , or N; and R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.

[0066] In some embodiments, R^A is unsubstituted or substituted monocyclic 6-membered heteroaryl or unsubstituted or substituted phenyl, wherein if R^A is substituted then R^A is substituted with 1, 2, 3 or 4 groups selected from R^a , R^b , and R^c . In some embodiments, R^A is unsubstituted or substituted pyridinyl, unsubstituted or substituted pyrimidinyl, unsubstituted or substituted pyrazinyl, or unsubstituted or substituted pyridazinyl, or unsubstituted or substituted phenyl, wherein if R^A is substituted then R^A is substituted with 1, 2, 3 or 4 groups selected from R^a , R^b , and R^c . In some embodiments, if R^A is substituted then R^A is substituted with 1 or 2 groups selected from R^a , R^b , and R^c .


[0067] In some embodiments, R^A is unsubstituted or substituted pyridinyl or unsubstituted or substituted phenyl, wherein if R^A is substituted then R^A is substituted with 1, 2, 3 or 4 groups selected from R^a , R^b , and R^c . In some embodiments, if R^A is substituted then R^A is substituted with 1 or 2 groups selected from R^a , R^b , and R^c .

[0068] In some embodiments, R^A is


$$R^b \sim R^a \quad R^b \sim R^a \quad R^b \sim R^a \quad R^b \sim R^a$$




or




[0069] In some embodiments, R^A is




In some embodiments, R^A is



In some embodiments, R^A is




[0070] In some embodiments, R^A is




where V is CH , CR^a , CR^b , or N . In some embodiments, V is CH or N .

[0071] In some embodiments, R^A is



where V is CH , CR^a , CR^b , or N . In some embodiments, V is CH or N .

[0072] In some embodiments, the compound has the structure of Formula (IV), or a pharmaceutically acceptable salt thereof:




18

Formula (IV)

wherein V is CH, CR^a, CR^b, or N. In some embodiments, V is CH or N.


[0073] In some embodiments, the compound has the structure of Formula (IVa), or a pharmaceutically acceptable salt thereof:



Formula (IVa)

wherein V is CH, CR^a, CR^b, or N. In some embodiments, V is CH or N.

[0074] In some embodiments, the compound has the structure of Formula (IVb), or a pharmaceutically acceptable salt thereof:



Formula (IVb)

wherein V is CH, CR^a, CR^b, or N. In some embodiments, V is CH or N.

[0075] In some embodiments, if R^A is substituted then R^A is substituted with 1 or 2 groups selected from R^a, R^b, and R^c.


[0076] In some embodiments, if R^B is substituted then R^B is substituted with 1, 2, or 3 groups selected from R^d, R^e, and R^f.

[0077] R^B is an unsubstituted or substituted phenyl or unsubstituted or substituted monocyclic 6-membered heteroaryl, wherein if R^B is substituted then R^B is substituted with 1, 2, 3 or 4 groups selected from R^d, R^e, and R^f. In some embodiments, R^B is unsubstituted or substituted phenyl, unsubstituted or substituted pyridinyl, unsubstituted or substituted pyrimidinyl, unsubstituted or substituted pyrazinyl, or unsubstituted or substituted pyridazinyl, wherein if R^B is substituted then R^B is substituted with 1, 2, 3 or 4 groups selected from R^d, R^e, and R^f. In some embodiments, if R^B is substituted then R^B is substituted with 1, 2, or 3 groups selected


from R^d , R^e , and R^f .

[0078] In some embodiments, R^B is unsubstituted or substituted phenyl or unsubstituted or substituted pyridinyl, wherein if R^B is substituted then R^B is substituted with 1, 2, 3 or 4 groups selected from R^d , R^e , and R^f . In some embodiments, if R^B is substituted then R^B is substituted with 1, 2, or 3 groups selected from R^d , R^e , and R^f .


[0079] In some embodiments, R^B is




or




In some embodiments, R^B is




In some embodiments, R^B is



In some embodiments, R^B is




[0080] In some embodiments, R^B is



where W is CH, CR^d , CR^e , or N. In some embodiments, W is CH or N.

[0081] In some embodiments, the compound has the structure of Formula (V), or a pharmaceutically acceptable salt thereof:



Formula (V)

wherein:


Y^1 is NR^c , O, or S;

Y^2 and Y^3 are independently CH, CR^a , CR^b , or N;

R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl; and

W is CH, CR^d , CR^e , or N.

[0082] In some embodiments, the compound of Formula (V) has one of the following structures, or a pharmaceutically acceptable salt thereof:




[0083] In some embodiments, M is $^*-NR^3-(C=O)-$, wherein * indicates the attachment point to R^1 .

[0084] In some embodiments, W is CH or N.

[0085] In some embodiments, Y^1 is NR^c ; and Y^2 and Y^3 are independently CH, CR^a , CR^b , or N. In some embodiments, Y^1 is NR^c ; Y^2 and Y^3 are independently CH, CR^a , or CR^b . In some embodiments; Y^1 is NR^c ; Y^2 and Y^3 are each CH.

[0086] In some embodiments, the compound has the structure of Formula (Va), or a pharmaceutically acceptable salt thereof:



Formula (Va)


wherein: Y^1 is NR^c , O, or S; Y^2 and Y^3 are independently CH, CR^a , CR^b , or N; R^c is hydrogen, -

$C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl; and W is CH, CR^d , CR^e , or N.

[0087] In some embodiments, W is CH or N.

[0088] In some embodiments, Y^1 is NR^c ; and Y^2 and Y^3 are independently CH, CR^a , CR^b , or N. In some embodiments, Y^1 is NR^c ; Y^2 and Y^3 are independently CH, CR^a , or CR^b . In some embodiments; Y^1 is NR^c ; Y^2 and Y^3 are each CH.

[0089] In some embodiments, the compound has the structure of Formula (Vb), or a pharmaceutically acceptable salt thereof:




Formula (Vb)

wherein: R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl; and W is CH, CR^d , CR^e , or N.

[0090] In some embodiments, W is CH or N.

[0091] In some embodiments, R^c is hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl. In some embodiments, R^c is hydrogen or unsubstituted or substituted C_1 - C_6 alkyl. In some embodiments, R^c is hydrogen or C_1 - C_6 alkyl. In some embodiments, R^c is unsubstituted or substituted C_1 - C_6 alkyl. In some embodiments, R^c is C_1 - C_6 alkyl.

[0092] In some embodiments, the compound has the structure of Formula (Vc), or a pharmaceutically acceptable salt thereof:



18


Formula (Vc)

wherein: R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl; and W is CH, CR^d , CR^e , or N.

[0093] In some embodiments, W is CH or N.

[0094] In some embodiments, R^c is hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl. In some embodiments, R^c is hydrogen or unsubstituted or substituted C_1 - C_6 alkyl. In some embodiments, R^c is hydrogen or C_1 - C_6 alkyl. In some embodiments, R^c is unsubstituted or substituted C_1 - C_6 alkyl. In some embodiments, R^c is C_1 - C_6 alkyl.

[0095] In some embodiments, the compound has the structure of Formula (Vd), or a pharmaceutically acceptable salt thereof:




Formula (Vd)


wherein: R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl; and W is CH, CR^d , CR^e , or N.

[0096] In some embodiments, W is CH or N.

[0097] In some embodiments, R^c is hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl. In some embodiments, R^c is hydrogen or unsubstituted or substituted C_1 - C_6 alkyl. In some embodiments, R^c is hydrogen or C_1 - C_6 alkyl. In some embodiments, R^c is unsubstituted or substituted C_1 - C_6 alkyl. In some embodiments, R^c is C_1 - C_6 alkyl.

[0098] In some embodiments, the compound has the structure of Formula (Ve), or a pharmaceutically acceptable salt thereof:






Formula (Ve)

wherein W is CH, CR^d, CR^e, or N. In some embodiments, W is CH or N.


[0099] In some embodiments, the compound has the structure of Formula (Vf), or a pharmaceutically acceptable salt thereof:



Formula (Vf)

wherein W is CH, CR^d, CR^e, or N. In some embodiments, W is CH or N.


[0100] In some embodiments, the compound has the structure of Formula (Vg), or a pharmaceutically acceptable salt thereof:



Formula (Vg)

wherein W is CH, CR^d, CR^e, or N. In some embodiments, W is CH or N.


[0101] In some embodiments, the compound has the structure of Formula (Vh), or a pharmaceutically acceptable salt thereof:



Formula (Vh)

wherein W is CH, CR^d, CR^e, or N. In some embodiments, W is CH or N.


[0102] In some embodiments, the compound has the structure of Formula (Vi), or a pharmaceutically acceptable salt thereof:



Formula (Vi)

wherein W is CH, CR^d, CR^e, or N. In some embodiments, W is CH or N.

[0103] In some embodiments, the compound has the structure of Formula (Vj), or a pharmaceutically acceptable salt thereof:



Formula (Vj)

wherein:

Y¹ is NR^c, O, or S;

Y² and Y³ are independently CH, CR^a, CR^b, or N;


R^c is hydrogen, -C(=O)R⁷, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, or unsubstituted or substituted C₂-C₇ heterocycloalkyl; and

W is CH, CR^d, CR^e, or N.

[0104] In some embodiments, W is CH or N.

[0105] In some embodiments, Y¹ is NR^c; and Y² and Y³ are independently CH, CR^a, CR^b, or N. In some embodiments, Y¹ is NR^c; Y² and Y³ are independently CH, CR^a, or CR^b. In some embodiments; Y¹ is NR^c; Y² and Y³ are each CH.

[0106] In some embodiments, the compound has the structure of Formula (Vk), or a pharmaceutically acceptable salt thereof:



Formula (Vk)

wherein:


R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl; and

W is CH, CR^d , CR^e , or N.

[0107] In some embodiments, W is CH or N.

[0108] In some embodiments, R^c is hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl. In some embodiments, R^c is hydrogen or unsubstituted or substituted C_1 - C_6 alkyl. In some embodiments, R^c is hydrogen or C_1 - C_6 alkyl. In some embodiments, R^c is unsubstituted or substituted C_1 - C_6 alkyl. In some embodiments, R^c is C_1 - C_6 alkyl.

[0109] In some embodiments, the compound has the structure of Formula (VI), or a pharmaceutically acceptable salt thereof:



Formula (VI)

wherein:


R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl; and

W is CH, CR^d, CR^e, or N.

[0110] In some embodiments, W is CH or N.

[0111] In some embodiments, R^c is hydrogen, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, or unsubstituted or substituted C₂-C₇ heterocycloalkyl. In some embodiments, R^c is hydrogen or unsubstituted or substituted C₁-C₆ alkyl. In some embodiments, R^c is hydrogen or C₁-C₆ alkyl. In some embodiments, R^c is unsubstituted or substituted C₁-C₆ alkyl. In some embodiments, R^c is C₁-C₆ alkyl.

[0112] In some embodiments, the compound has the structure of Formula (Vm), or a pharmaceutically acceptable salt thereof:



Formula (Vm)

wherein:

R^c is hydrogen, -C(=O)R⁷, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, or unsubstituted or substituted C₂-C₇ heterocycloalkyl; and


W is CH, CR^d, CR^e, or N.

[0113] In some embodiments, W is CH or N.

[0114] In some embodiments, R^c is hydrogen, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, or unsubstituted or substituted C₂-C₇ heterocycloalkyl. In some embodiments, R^c is hydrogen or unsubstituted or substituted C₁-C₆ alkyl. In some embodiments, R^c is hydrogen or C₁-C₆ alkyl. In some embodiments, R^c is unsubstituted or substituted C₁-C₆ alkyl. In some embodiments, R^c is C₁-C₆ alkyl.

[0115] In some embodiments, the compound has the structure of Formula (Ve), or a


pharmaceutically acceptable salt thereof:



Formula (Vn)

wherein W is CH, CR^d , CR^e , or N. In some embodiments, W is CH or N.


[0116] In some embodiments, the compound has the structure of Formula (Vf), or a pharmaceutically acceptable salt thereof:



Formula (Vn)

wherein W is CH, CR^d , CR^e , or N. In some embodiments, W is CH or N.


[0117] In some embodiments, the compound has the structure of Formula (Vg), or a pharmaceutically acceptable salt thereof:




Formula (Vo)

wherein W is CH, CR^d , CR^e , or N. In some embodiments, W is CH or N.

[0118] In some embodiments, the compound has the structure of Formula (Vh), or a pharmaceutically acceptable salt thereof:






Formula (Vo)

wherein W is CH, CR^d, CR^e, or N. In some embodiments, W is CH or N.


[0119] In some embodiments, the compound has the structure of Formula (Vp), or a pharmaceutically acceptable salt thereof:



Formula (Vp)

wherein W is CH, CR^d, CR^e, or N. In some embodiments, W is CH or N.


[0120] In some embodiments, the compound has the structure of Formula (VI), or a pharmaceutically acceptable salt thereof:



Formula (VI)

wherein V is CH, CR^a, CR^b, or N; and W is CH, CR^d, CR^e, or N.

[0121] In some embodiments, the compound of Formula (VI) has the following structure, or a pharmaceutically acceptable salt thereof:




[0122] In some embodiments, M is ^{*}-NR³-(C=O)-, wherein ^{*} indicates the attachment point to R¹. In some embodiments, M is ^{*}-(C=O)-NR³-, wherein ^{*} indicates the attachment point to R¹.

[0123] In some embodiments, V is CH or N. In some embodiments, W is CH or N. In some

embodiments, V is CH or N; and W is CH or N.

[0124] In some embodiments, the compound has the structure of Formula (VIa), or a pharmaceutically acceptable salt thereof:




Formula (VIa)

wherein V is CH, CR^a, CR^b, or N; and W is CH, CR^d, CR^e, or N.

[0125] In some embodiments, V is CH or N. In some embodiments, W is CH or N. In some embodiments, V is CH or N; and W is CH or N.


[0126] In some embodiments, the compound has the structure of Formula (VIb), or a pharmaceutically acceptable salt thereof:



Formula (VIb)

wherein V is CH, CR^a, CR^b, or N. In some embodiments, V is CH or N.

[0127] In some embodiments, the compound has the structure of Formula (VIc), or a pharmaceutically acceptable salt thereof:




Formula (VIc)

wherein V is CH, CR^a, CR^b, or N; and W is CH, CR^d, CR^e, or N.

[0128] In some embodiments, V is CH or N. In some embodiments, W is CH or N. In some embodiments, V is CH or N; and W is CH or N.

[0129] In some embodiments, the compound has the structure of Formula (VIId), or a pharmaceutically acceptable salt thereof:



Formula (VIId)

wherein V is CH, CR^a, CR^b, or N. In some embodiments, V is CH or N.

[0130] In some embodiments, R¹ is unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₂-C₇ heterocycloalkyl, unsubstituted or substituted -(C₁-C₆ alkyl)-(C₃-C₆ cycloalkyl), or unsubstituted or substituted -(C₁-C₆ alkyl)-(C₂-C₇ heterocycloalkyl), wherein any substituted group of R¹ is substituted with one or more halogen, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted monocyclic heterocycle, -N(R⁴)₂, -OR⁵, -CN, -CO₂R⁵, -C(=O)N(R⁴)₂, -SR⁵, -S(=O)R⁷, -S(=O)₂R⁷, -NR⁴C(=O)R⁵, -NR⁴SO₂R⁷, -SO₂R⁷, or -SO₂N(R⁴)₂; wherein R¹ comprises a basic amine group.

[0131] In some embodiments, R¹ is unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ heteroalkyl containing 1 N atom, unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycloalkyl containing 1-4 N atoms and 0 or 1 O or S atoms, unsubstituted or substituted bridged C₂-C₇ heterocycloalkyl containing 1-2 N atoms, or unsubstituted or substituted -(C₁-C₆ alkyl)-(C₂-C₇ heterocycloalkyl), wherein any substituted group of R¹ is substituted with one or more halogen, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted monocyclic heterocycle, -N(R⁴)₂, -OR⁵, -CN, -CO₂R⁵, -C(=O)N(R⁴)₂, -SR⁵, -S(=O)R⁷, -S(=O)₂R⁷, -NR⁴C(=O)R⁵, -NR⁴SO₂R⁷, -SO₂R⁷, or -SO₂N(R⁴)₂.

[0132] In some embodiments, R¹ is unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ heteroalkyl containing 1 N atom, wherein any substituted group of R¹ is substituted with one or more halogen, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted monocyclic heterocycle, -N(R⁴)₂, -OR⁵, -CN, -CO₂R⁵, -C(=O)N(R⁴)₂, -SR⁵, -S(=O)R⁷, -S(=O)₂R⁷, -NR⁴C(=O)R⁵, -NR⁴SO₂R⁷, -SO₂R⁷, or -SO₂N(R⁴)₂; or R¹ is unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms; or R¹ is unsubstituted or substituted bridged C₂-C₇ heterocycloalkyl containing

1-2 N atoms; or R^1 is unsubstituted or substituted $-(C_1-C_6 \text{ alkyl})-(C_2-C_7 \text{ heterocycloalkyl})$, wherein the heterocycloalkyl is an unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycloalkyl containing 1-4 N atoms and 0 or 1 O or S atoms.

[0133] In some embodiments, R^1 is unsubstituted or substituted C_1-C_6 heteroalkyl containing 1 N atom; or R^1 is unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms; or R^1 is unsubstituted or substituted $-(C_1-C_6 \text{ alkyl})-(C_2-C_7 \text{ heterocycloalkyl})$, wherein the heterocycloalkyl is an unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycloalkyl containing 1-4 N atoms and 0 or 1 O or S atoms.

[0134] In some embodiments, R^1 is unsubstituted or substituted C_1-C_6 alkyl, unsubstituted or substituted C_1-C_6 heteroalkyl containing 1 N atom, wherein any substituted group of R^1 is substituted with one or more halogen, unsubstituted or substituted C_1-C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, $-N(R^4)_2$, $-OR^5$, $-CN$, $-CO_2R^5$, $-C(=O)N(R^4)_2$, $-SR^5$, $-S(=O)R^7$, $-S(=O)_2R^7$, $-NR^4C(=O)R^5$, $-NR^4SO_2R^7$, $-SO_2R^7$, or $-SO_2N(R^4)_2$.

[0135] In some embodiments, R^1 is unsubstituted or substituted C_1-C_6 heteroalkyl containing 1 N atom. In some embodiments, R^1 is unsubstituted or substituted C_1-C_6 heteroalkyl containing 1 N atom; wherein any substituted group of R^1 is substituted with one or more halogen, unsubstituted or substituted C_1-C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, $-N(R^4)_2$, $-OR^5$, $-CN$, $-CO_2R^5$, $-C(=O)N(R^4)_2$, $-SR^5$, $-S(=O)R^7$, $-S(=O)_2R^7$, $-NR^4C(=O)R^5$, $-NR^4SO_2R^7$, $-SO_2R^7$, or $-SO_2N(R^4)_2$.

[0136] In some embodiments, R^1 is unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms.

[0137] In some embodiments, R^1 is unsubstituted or substituted bridged C_2-C_7 heterocycloalkyl containing 1-2 N atoms.

[0138] In some embodiments, R^1 is unsubstituted or substituted $-(C_1-C_6 \text{ alkyl})-(C_2-C_7 \text{ heterocycloalkyl})$, wherein the heterocycloalkyl is an unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycloalkyl containing 1-4 N atoms and 0 or 1 O or S atoms.


[0139] In some embodiments, R^1 is unsubstituted or substituted C_1-C_6 heteroalkyl containing 1 N atom, wherein any substituted group of R^1 is substituted with one or more halogen, C_1-C_4 alkyl, $-N(R^4)_2$, or $-OR^5$; or R^1 is unsubstituted or substituted azetidiny, unsubstituted or

substituted pyrrolidinyl, or unsubstituted or substituted piperidinyl; or R^1 is unsubstituted or substituted $-(C_1-C_6 \text{ alkyl})-(C_2-C_7 \text{ heterocycloalkyl})$, wherein the heterocycloalkyl is an unsubstituted or substituted azetidiny, unsubstituted or substituted pyrrolidinyl, or unsubstituted or substituted piperidinyl.

[0140] In some embodiments, R^1 is unsubstituted or substituted azetidiny, unsubstituted or substituted pyrrolidinyl, or unsubstituted or substituted piperidinyl.

[0141] In some embodiments, R^1 is unsubstituted or substituted C_1-C_6 heteroalkyl containing 1 N atom, wherein any substituted group of R^1 is substituted with one or more halogen, C_1-C_4 alkyl, $-N(R^4)_2$, or $-OR^5$


[0142] In some embodiments, the compound has the structure of Formula (VIIa), or a pharmaceutically acceptable salt thereof:



Formula (VIIa)

wherein W is CH, CR^a , CR^b , or N. In some embodiments, W is CH or N.


[0143] In some embodiments, the compound has the structure of Formula (VIIb), or a pharmaceutically acceptable salt thereof:




Formula (VIIb)

wherein V is CH, CR^a , CR^b , or N; and W is CH, CR^d , CR^e , or N.

[0144] In some embodiments, the compound has the structure of Formula (VIIc), or a pharmaceutically acceptable salt thereof:






Formula (VIIc)

wherein W is CH, CR^a, CR^b, or N. In some embodiments, W is CH or N.

[0145] In some embodiments, V is CH or N. In some embodiments, W is CH or N. In some embodiments, V is CH or N; and W is CH or N.


[0146] In some embodiments, the compound has the structure of Formula (VIId), or a pharmaceutically acceptable salt thereof:



Formula (VIId)

wherein W is CH, CR^a, CR^b, or N. In some embodiments, W is CH or N.


[0147] In some embodiments, the compound has the structure of Formula (VIIe), or a pharmaceutically acceptable salt thereof:



Formula (VIIe)

wherein V is CH, CR^a, CR^b, or N; and W is CH, CR^d, CR^e, or N.

[0148] In some embodiments, the compound has the structure of Formula (VIIf), or a pharmaceutically acceptable salt thereof:




Formula (VIIf)

wherein W is CH, CR^a, CR^b, or N. In some embodiments, W is CH or N.

[0149] In some embodiments, V is CH or N. In some embodiments, W is CH or N. In some embodiments, V is CH or N; and W is CH or N.

[0150] In some embodiments, the compound has the structure of Formula (VIII), or a pharmaceutically acceptable salt thereof:



Formula (VIII)

wherein:

R^A is unsubstituted or substituted phenyl, unsubstituted or substituted pyridinyl, unsubstituted or substituted furanyl, unsubstituted or substituted thienyl, unsubstituted or substituted pyrrolyl, unsubstituted or substituted pyrazolyl, unsubstituted or substituted isoxazolyl, wherein if R^A is substituted then R^A is substituted with 1 or 2 groups selected from R^a, R^b, and R^c;

R^B is unsubstituted or substituted phenyl or unsubstituted or substituted pyridinyl, wherein if R^B is substituted then R^B is substituted with 1, 2, or 3 groups selected from R^d, R^e, and R^f; and

M is -NR³-, -O-, *-NR³-(C=O)-, *(C=O)-NR³-, *-O-(C=O)NR³-, *-NR³-(C=O)O-, or -NR³-(C=O)NR³-, wherein * indicates the attachment point to R¹.

[0151] In some embodiments, X¹ is CR¹¹; X² is CR¹²; X³ is CR¹³; and X⁴ is CR¹⁴. In some embodiments, X¹ is N; X² is CR¹²; X³ is CR¹³; and X⁴ is CR¹⁴. In some embodiments, X¹ is CR¹¹; X² is N; X³ is CR¹³; and X⁴ is CR¹⁴. In some embodiments, X¹ is N; X² is CR¹²; X³ is CR¹³; and X⁴ is N. In some embodiments, X¹ is N; X² is N; X³ is CR¹³; and X⁴ is CR¹⁴. In some embodiments, X¹ is N; X² is CR¹²; X³ is N; and X⁴ is CR¹⁴. In some embodiments, X¹ is CR¹¹; X² is CR¹²; X³ is CR¹³; and X⁴ is CR¹⁴.

[0152] In some embodiments, X¹ is CR¹¹ or N; X² is CR¹²; X³ is CR¹³; and X⁴ is CR¹⁴.

[0153] In some embodiments, X¹ is CR¹¹ or N; X² is CR¹² or N; X³ is CR¹³; and X⁴ is CR¹⁴.

[0154] In some embodiments, R¹¹, R¹², R¹³, and R¹⁴ are each independently hydrogen, halogen, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₃-C₆cycloalkyl, -CN, or -OR⁴. In some embodiments,

R^{11} , R^{12} , R^{13} , and R^{14} are each independently hydrogen, halogen, C_1 - C_6 alkyl, fluoroalkyl, unsubstituted C_3 - C_6 cycloalkyl, -CN, or -OR⁴. In some embodiments, R^{11} , R^{12} , R^{13} , and R^{14} are each independently hydrogen, F, Cl, Br, -CN, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)CH₂CH₃, -C(CH₃)₃, -CH₂F, -CHF₂, -CF₃, or cyclopropyl. In some embodiments, R^{11} , R^{12} , R^{13} , and R^{14} are each independently hydrogen or F.

[0155] In some embodiments, X¹ is CH or CF; X² is CH or CF; X³ is CH or CF; and X⁴ is CH or CF. In some embodiments, X¹ is N; X² is CH or CF; X³ is CH or CF; and X⁴ is CH or CF. In some embodiments, X¹ is CH or CF; X² is N; X³ is CH or CF; and X⁴ is CH or CF. In some embodiments, X¹ is N; X² is CH or CF; X³ is CH or CF; and X⁴ is N. In some embodiments, X¹ is N; X² is N; X³ is CH or CF; and X⁴ is CH or CF. In some embodiments, X¹ is N; X² is CH or CF; X³ is N; and X⁴ is CH or CF. In some embodiments, X¹ is CH or CF; X² is CH or CF; X³ is CR¹³; and X⁴ is CR¹⁴.

[0156] In some embodiments, X¹ is CH, CF, or N; X² is CH or CF; X³ is CH or CF; and X⁴ is CH or CF. In some embodiments, X¹ is CR¹¹ or N; X² is CH; X³ is CH; and X⁴ is CH. In some embodiments, X¹ is CH, CF, or N; X² is CH; X³ is CH; and X⁴ is CH.

[0157] In some embodiments, X¹ is CH, CF, or N; X² is CH, CF, or N; X³ is CH or CF; and X⁴ is CH or CF. In some embodiments, X¹ is CR¹¹ or N; X² is CR¹² or N; X³ is CH; and X⁴ is CH. In some embodiments, X¹ is CH, CF, or N; X² is CH, CF, or N; X³ is CH; and X⁴ is CH.

[0158] In some embodiments, X¹ is CH, CF, or N; X² is CH or CF; X³ is CR¹³; and X⁴ is CH or CF. In some embodiments, X¹ is CH, CF, or N; X² is CH; X³ is CR¹³; and X⁴ is CH. In some embodiments, X¹ is N; X² is CH; X³ is CR¹³; and X⁴ is CH.

[0159] In some embodiments, X¹ is CH or N; X² is CH or N; X³ is CH, CF, or N; and X⁴ is CH or N. In some embodiments, X¹ is CH or N; X² is CH or N; X³ is CH or CF; and X⁴ is CH. In some embodiments, X¹ is N; X² is CH; X³ is CH or CF; and X⁴ is CH. In some embodiments, X¹ is N; X² is CH; X³ is CH; and X⁴ is CH. In some embodiments, X¹ is N; X² is CH; X³ is CF; and X⁴ is CH.

[0160] In some embodiments, R¹³ is hydrogen, halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, -CN, or -OR⁴. In some embodiments, R¹³ is hydrogen, halogen, C_1 - C_6 alkyl, fluoroalkyl, unsubstituted C_3 - C_6 cycloalkyl, -CN, or -OR⁴. In some embodiments, R¹³ is

hydrogen, F, Cl, Br, -CN, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)CH₂CH₃, -C(CH₃)₃, -CH₂F, -CHF₂, -CF₃, or cyclopropyl. In some embodiments, R¹³ is hydrogen or F. In some embodiments, R¹³ is hydrogen. In some embodiments, R¹³ is F.

[0161] In some embodiments, M is *-NR³-(C=O)- or *-(C=O)-NR³-, wherein * indicates the attachment point to R¹.

[0162] In some embodiments, R^A is unsubstituted or substituted pyridinyl or unsubstituted or substituted phenyl, wherein if R^A is substituted then R^A is substituted with 1 or 2 groups selected from R^a, R^b, and R^c.

[0163] In some embodiments, R^B is unsubstituted or substituted phenyl, wherein if R^B is substituted then R^B is substituted with 1, 2, or 3 groups selected from R^d, R^e, and R^f.

[0164] In some embodiments, R^a, R^b, and R^c are independently selected from the group consisting of hydrogen, halogen, -OR⁴, -CN, -N(R⁴)₂, -C(=O)R⁷, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₂-C₇ heterocycloalkyl, unsubstituted or substituted phenyl, unsubstituted or substituted monocyclic heteroaryl, and unsubstituted or substituted bicyclic heteroaryl, wherein any substituted group of R^a, R^b, and R^c is substituted with one or more R⁶ groups; or one R^a and one R^b, when present on adjacent atoms of R^A, are taken together with the intervening atoms connecting R^a to R^b to form a 5- to 6-membered monocyclic carbocycle or 5- to 6-membered monocyclic heterocycle, wherein the carbocycle or heterocycle is unsubstituted or substituted with one or more R⁶ groups; wherein, if R^a, R^b, or R^c is attached to the N atom of a heteroaryl, then it is hydrogen, -C(=O)R⁷, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, or unsubstituted or substituted C₂-C₇ heterocycloalkyl; and R^d, R^e, and R^f are independently selected from the group consisting of hydrogen, halogen, -OR⁴, -CN, -N(R⁴)₂, -C(=O)R⁷, -C(=O)N(R⁴)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₂-C₇ heterocycloalkyl, unsubstituted or substituted phenyl, unsubstituted or substituted monocyclic heteroaryl, and unsubstituted or substituted bicyclic heteroaryl, wherein any substituted group of R^d, R^e, and R^f is substituted with one or more R⁶ groups; wherein, if R^d, R^e, or R^f is attached to the N atom of a heteroaryl, then it is hydrogen, -C(=O)R⁷, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, or unsubstituted or substituted C₂-C₇ heterocycloalkyl.

[0165] In some embodiments, R^a is selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, $-N(R^4)_2$, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl, and unsubstituted or substituted C_3 - C_6 cycloalkyl, wherein any substituted group of R^a is substituted with one or more R^6 groups; and R^b and R^c are independently selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, $-N(R^4)_2$, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted C_1 - C_6 heteroalkyl, wherein any substituted group of R^b and R^c is substituted with one or more R^6 groups; wherein, if R^a , R^b , or R^c is attached to the N atom of a heteroaryl, then it is hydrogen, $-C(=O)R^7$, or unsubstituted or substituted C_1 - C_6 alkyl.

[0166] In some embodiments, R^a is selected from the group consisting of hydrogen, F, Cl, Br, $-CN$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-C(O)CH_3$, $-C(O)CH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CH_2OH$, $-CH_2CH_2CN$, $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, $-CH_2NH_2$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2NHCH_3$, $-CH_2CH_2N(CH_3)_2$, cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl; and R^b and R^c are independently selected from the group consisting of hydrogen, F, Cl, Br, $-CN$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-C(O)CH_3$, $-C(O)CH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CH_2OH$, $-CH_2CH_2CN$, $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, $-CH_2NH_2$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2NHCH_3$, and $-CH_2CH_2N(CH_3)_2$; wherein, if R^a , R^b , or R^c is attached to the N atom of a heteroaryl, then it is hydrogen, $-C(O)CH_3$, $-C(O)CH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CH_2OH$, $-CH_2CH_2CN$, $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, $-CH_2NH_2$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2NHCH_3$, and $-CH_2CH_2N(CH_3)_2$.

[0167] In some embodiments, one R^a and one R^b on adjacent atoms of R^A are taken together with the intervening atoms connecting R^a to R^b to form a 5- to 6-membered monocyclic cycloalkyl or 5- to 6-membered monocyclic heterocycloalkyl, wherein the cycloalkyl or heterocycloalkyl is unsubstituted or substituted with one or more R^6 groups. In some embodiments, one R^a and one R^b on adjacent atoms of R^A are taken together with the intervening atoms connecting R^a to R^b to form a 5-membered monocyclic heterocycloalkyl, wherein the heterocycloalkyl is unsubstituted or substituted with one or more R^6 groups.

[0168] In some embodiments, R^d is selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, $-N(R^4)_2$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_2 - C_7 heterocycloalkyl, unsubstituted or substituted phenyl, unsubstituted or substituted monocyclic heteroaryl, and unsubstituted or substituted bicyclic heteroaryl, wherein any substituted group of R^d is substituted with one or more R^6 groups; and R^e and R^f are independently selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, $-N(R^4)_2$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted C_1 - C_6 heteroalkyl; wherein, if R^d , R^e , or R^f is attached to the N atom of a heteroaryl, then it is hydrogen, $-C(=O)R^7$, or unsubstituted or substituted C_1 - C_6 alkyl.

[0169] In some embodiments, R^d is selected from the group consisting of hydrogen, F, Cl, Br, $-CN$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CH_2OH$, $-CH_2CH_2CN$, $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, $-CH_2NH_2$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2NHCH_3$, $-CH_2CH_2N(CH_3)_2$, unsubstituted or substituted cyclopropyl, unsubstituted or substituted cyclobutyl, unsubstituted or substituted cyclopentyl, unsubstituted or substituted cyclohexyl, unsubstituted or substituted C_2 - C_7 heterocycloalkyl, unsubstituted or substituted phenyl, unsubstituted or substituted monocyclic heteroaryl, and unsubstituted or substituted bicyclic heteroaryl, wherein any substituted group of R^d is substituted with one or more R^6 groups; and R^e and R^f are independently selected from the group consisting of hydrogen, F, Cl, Br, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CN$, $-OH$, $-OCH_3$, and $-OCH_2CH_3$; wherein, if R^d , R^e , or R^f is attached to the N atom of a heteroaryl, then it is hydrogen, $-C(O)CH_3$, $-C(O)CH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CH_2OH$, $-CH_2CH_2CN$, $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, $-CH_2NH_2$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2NHCH_3$, and $-CH_2CH_2N(CH_3)_2$.

[0170] In some embodiments,

R^A is unsubstituted or substituted pyrrolyl, unsubstituted or substituted pyridinyl, or unsubstituted or substituted phenyl, wherein if R^A is substituted then R^A is substituted with 1, or 2 groups selected from R^a , R^b , and R^c ;

R^B is unsubstituted or substituted phenyl or unsubstituted or substituted pyridinyl, wherein if R^B is substituted then R^B is substituted with 1, 2, or 3 groups selected from R^d , R^e , and R^f ;

X^1 is CH or N;

X^2 is CH or N;

X^3 is CR^{13} or N;

X^4 is CH or N;

M is $^*-NR^3-(C=O)-$ or $^*-(C=O)-NR^3-$, wherein * indicates the attachment point to R^1 ;

R^{11} , R^{12} , R^{13} , and R^{14} are each independently hydrogen, F, Cl, $-CH_3$, CF_3 , $-CN$, $-OR^4$, or $-N(R^4)_2$;

R^1 is unsubstituted or substituted C_1 - C_6 heteroalkyl containing 1 N atom, wherein any substituted group of R^1 is substituted with one or more halogen, C_1 - C_4 alkyl, $-N(R^4)_2$, or $-OR^5$;


or R^1 is unsubstituted or substituted azetidiny, unsubstituted or substituted pyrrolidiny, or unsubstituted or substituted piperidiny;

or R^1 is unsubstituted or substituted bridged C_2 - C_7 heterocycloalkyl containing 1-2 N atoms;


or R^1 is unsubstituted or substituted $-(C_1$ - C_6 alkyl)-(C_2 - C_7 heterocycloalkyl), wherein the heterocycloalkyl is an unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycloalkyl containing 1-2 N atoms.

[0171] In some embodiments,


R^A is



or R^A is



or R^A is




where V is CH or N;

R^a is selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted C_3 - C_6 cycloalkyl; and

R^b and R^c are independently selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, unsubstituted or substituted C_1 - C_6 alkyl, and unsubstituted or substituted C_1 - C_6 fluoroalkyl;

R^B is



where W is CH or N;

R^d is selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted C_3 - C_6 cycloalkyl; and

R^e and R^f are independently selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, unsubstituted or substituted C_1 - C_6 alkyl, and unsubstituted or substituted C_1 - C_6 fluoroalkyl;

X^1 is CH, CF, or N;

X^2 is CH, CF, or N;

X^3 is CH, CF, or N;

X^4 is CH, CF, or N;

M is $^*-NH-(C=O)-$ or $^*-(C=O)-NH-$, wherein * indicates the attachment point to R^1 ;

R^1 is unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl containing 1 N atom, wherein any substituted group of R^1 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, $-N(R^4)_2$, $-OR^5$, $-CN$, $-CO_2R^5$, $-C(=O)N(R^4)_2$, $-SR^5$, $-S(=O)R^7$, $-S(=O)_2R^7$, $-NR^4C(=O)R^5$, $-NR^4SO_2R^7$, $-SO_2R^7$, or $-SO_2N(R^4)_2$;


or R^1 is unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms;

or R^1 is unsubstituted or substituted bridged C_2 - C_7 heterocycloalkyl containing 1-2 N atoms;

or R¹ is unsubstituted or substituted -(C₁-C₆ alkyl)-(C₂-C₇ heterocycloalkyl), wherein the heterocycloalkyl is an unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycloalkyl containing 1-2 N atoms.

[0172] In some embodiments,

R^A is




where V is CH or N;

R^a is selected from the group consisting of hydrogen, F, Cl, Br, -CN, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -C(CH₃)₃, -CH₂F, -CHF₂, and -CF₃; and

R^b and R^c are independently selected from the group consisting of hydrogen, F, Cl, -CH₃, -CH₂F, -CHF₂, -CF₃, -CN, and -OCH₃;

R^B is



where W is CH or N;

R^d is selected from the group consisting of hydrogen, F, Cl, Br, -CN, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -C(CH₃)₃, -CH₂F, -CHF₂, and -CF₃;

R^e and R^f are independently selected from the group consisting of hydrogen, F, Cl, Br, -CH₃, -CH₂F, -CHF₂, -CF₃, -CN, -OH, and -OCH₃;

X¹ is CH or N;

X² is CH or N;

X³ is CH, CF, or N;

X⁴ is CH or N;

M is *-NH-(C=O)- or *-(C=O)-NH-, wherein * indicates the attachment point to R¹;

R^1 is unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl containing 1 N atom, wherein any substituted group of R^1 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, $-N(R^4)_2$, $-OR^5$, $-CN$, $-CO_2R^5$, $-C(=O)N(R^4)_2$, $-SR^5$, $-S(=O)R^7$, $-S(=O)_2R^7$, $-NR^4C(=O)R^5$, $-NR^4SO_2R^7$, $-SO_2R^7$, or $-SO_2N(R^4)_2$;

or R^1 is unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms;

or R^1 is unsubstituted or substituted bridged C_2 - C_7 heterocycloalkyl containing 1-2 N atoms;

or R^1 is unsubstituted or substituted $-(C_1$ - C_6 alkyl)-(C_2 - C_7 heterocycloalkyl), wherein the heterocycloalkyl is an unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycloalkyl containing 1-2 N atoms.

[0173] In some embodiments,

R^A is unsubstituted or substituted pyrrolyl, unsubstituted or substituted pyridinyl, or unsubstituted or substituted phenyl, wherein if R^A is substituted then R^A is substituted with 1, or 2 groups selected from R^a , R^b , and R^c ;

wherein, if R^a , R^b , or R^c is attached to the N atom of a heteroaryl, then it is hydrogen, $-C(=O)R^7$, or unsubstituted or substituted C_1 - C_6 alkyl;

R^B is unsubstituted or substituted phenyl or unsubstituted or substituted pyridinyl, wherein if R^B is substituted then R^B is substituted with 1, 2, or 3 groups selected from R^d , R^e , and R^f ;

X^1 is CR^{11} or N;

X^2 is CR^{12} or N;

R^{11} and R^{12} are each independently hydrogen, F, Cl, $-CH_3$, CF_3 , $-CN$, $-OR^4$, or $-N(R^4)_2$;

R^1 is unsubstituted or substituted C_1 - C_6 heteroalkyl containing 1 N atom, wherein any substituted group of R^1 is substituted with one or more halogen, C_1 - C_4 alkyl, $-N(R^4)_2$, or $-OR^5$;

or R^1 is unsubstituted or substituted azetidiny, unsubstituted or substituted pyrrolidinyl, or unsubstituted or substituted piperidinyl;

or R^1 is unsubstituted or substituted bridged C_2 - C_7 heterocycloalkyl containing 1-2 N atoms;

or R^1 is unsubstituted or substituted $-(C_1-C_6 \text{ alkyl})-(C_2-C_7 \text{ heterocycloalkyl})$, wherein the heterocycloalkyl is an unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycloalkyl containing 1-2 N atoms.

[0174] In some embodiments,

R^A is unsubstituted or substituted pyrrolyl, unsubstituted or substituted pyridinyl, or unsubstituted or substituted phenyl, wherein if R^A is substituted then R^A is substituted with 1, or 2 groups selected from R^a , R^b , and R^c ;

wherein, if R^a , R^b , or R^c is attached to the N atom of a heteroaryl, then it is hydrogen, $-C(=O)R^7$, or unsubstituted or substituted C_1-C_6 alkyl;

R^B is unsubstituted or substituted phenyl or unsubstituted or substituted pyridinyl, wherein if R^B is substituted then R^B is substituted with 1, 2, or 3 groups selected from R^d , R^e , and R^f ;

X^1 is CR^{11} or N;

R^{11} is hydrogen, F, Cl, $-CH_3$, CF_3 , $-CN$, $-OR^4$, or $-N(R^4)_2$;


R^1 is unsubstituted or substituted C_1-C_6 heteroalkyl containing 1 N atom, wherein any substituted group of R^1 is substituted with one or more halogen, C_1-C_4 alkyl, $-N(R^4)_2$, or $-OR^5$;

or R^1 is unsubstituted or substituted azetidiny, unsubstituted or substituted pyrrolidinyl, or unsubstituted or substituted piperidinyl;

or R^1 is unsubstituted or substituted bridged C_2-C_7 heterocycloalkyl containing 1-2 N atoms;


or R^1 is unsubstituted or substituted $-(C_1-C_6 \text{ alkyl})-(C_2-C_7 \text{ heterocycloalkyl})$, wherein the heterocycloalkyl is an unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycloalkyl containing 1-2 N atoms.

[0175] In some embodiments, R^A is




or R^A is






or R^A is




where V is CH or N; and R^B is




where W is CH or N.


[0176] In some embodiments, R^A is



or R^A is




where V is CH or N; and R^B is




where W is CH or N.

[0177] In some embodiments, R^A is




and R^B is





, where W is CH or N.

[0178] In some embodiments, R^A is




and R^B is






[0179] In some embodiments, R^A is




where V is CH or N; and R^B is



where W is CH or N.

[0180] In some embodiments, the compound has the structure of Formula (X), or a pharmaceutically acceptable salt thereof:




Formula (X)

wherein:

R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.

[0181] In some embodiments, the compound has the structure of Formula (Xa), or a pharmaceutically acceptable salt thereof:



Formula (Xa)


wherein:

R^a and R^b are independently selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted C_3 - C_6 cycloalkyl;

R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.

[0182] In some embodiments, R^a and R^b are independently selected from the group consisting of hydrogen, F, Cl, Br, -CN, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)CH₂CH₃, -C(CH₃)₃, -CH₂F, -CHF₂, -CF₃, and cyclopropyl; R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl.

[0183] In some embodiments, the compound has the structure of Formula (Xb), or a pharmaceutically acceptable salt thereof:




Formula (Xb)

wherein: X^1 is N; X^2 is CH or N; and R^c is hydrogen, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)CH₂CH₃, -C(CH₃)₃, -CH₂CH₂CH(CH₃)₂, or unsubstituted C_3 - C_6 cycloalkyl.

[0184] In some embodiments, X^2 is N. In some embodiments, X^2 is CH.


[0185] In some embodiments, the compound has the structure of Formula (Xc), or a pharmaceutically acceptable salt thereof:




Formula (Xc)

wherein: R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.

[0186] In some embodiments, the compound has the structure of Formula (Xd), or a pharmaceutically acceptable salt thereof:






Formula (Xd)

wherein: R^a and R^b are independently selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted C_3 - C_6 cycloalkyl; R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, or unsubstituted or substituted C_2 - C_7 heterocycloalkyl.

[0187] In some embodiments, R^a and R^b are independently selected from the group consisting of hydrogen, F, Cl, Br, $-CN$, $-OCH_3$, $-OCH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)CH_2CH_3$, $-C(CH_3)_3$, $-CH_2F$, $-CHF_2$, $-CF_3$, and cyclopropyl; R^c is hydrogen, $-C(=O)R^7$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl.


[0188] In some embodiments, the compound has the structure of Formula (Xe), or a pharmaceutically acceptable salt thereof:



Formula (Xe)

wherein: X^1 is N; and R^c is hydrogen, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)CH_2CH_3$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, or unsubstituted C_3 - C_6 cycloalkyl.


[0189] In some embodiments, the compound has the structure of Formula (XI), or a pharmaceutically acceptable salt thereof:



Formula (XI)

wherein V is CH or N.

[0190] In some embodiments, the compound has the structure of Formula (XIa), or a pharmaceutically acceptable salt thereof:




Formula (XIa)

wherein V is CH or N; R^a is selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted C_3 - C_6 cycloalkyl; and R^b and R^c are independently selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, unsubstituted or substituted C_1 - C_6 alkyl, and unsubstituted or substituted C_1 - C_6 fluoroalkyl.

[0191] In some embodiments, R^a is selected from the group consisting of hydrogen, F, Cl, Br, $-CN$, $-OCH_3$, $-OCH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2F$, $-CHF_2$, and $-CF_3$; R^b and R^c are independently selected from the group consisting of hydrogen, F, Cl, $-CH_3$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CN$, and $-OCH_3$.

[0192] In some embodiments, the compound has the structure of Formula (XIb), or a pharmaceutically acceptable salt thereof:





Formula (XIb)

wherein: V is CH or N; X^1 is N; and X^2 is CH or N.

[0193] In some embodiments, X^2 is N. In some embodiments, X^2 is CH.


[0194] In some embodiments, the compound has the structure of Formula (XIc), or a pharmaceutically acceptable salt thereof:





Formula (XIc)
wherein V is CH or N.

[0195] In some embodiments, the compound has the structure of Formula (XIId), or a pharmaceutically acceptable salt thereof:



Formula (XIId)


wherein V is CH or N;

R^a is selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted C_3 - C_6 cycloalkyl; and

R^b and R^c are independently selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, unsubstituted or substituted C_1 - C_6 alkyl, and unsubstituted or substituted C_1 - C_6 fluoroalkyl.

[0196] In some embodiments, R^a is selected from the group consisting of hydrogen, F, Cl, Br, $-CN$, $-OCH_3$, $-OCH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2F$, $-CHF_2$, and $-CF_3$; and R^b and R^c are independently selected from the group consisting of hydrogen, F, Cl, $-CH_3$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CN$, and $-OCH_3$.


[0197] In some embodiments, the compound has the structure of Formula (XIIe), or a pharmaceutically acceptable salt thereof:



Formula (XIe)


wherein: V is CH or N; and X¹ is N.

[0198] In some embodiments, the compound has the structure of Formula (XII), or a pharmaceutically acceptable salt thereof:



Formula (XII).

[0199] In some embodiments, the compound has the structure of Formula (XIIa), or a pharmaceutically acceptable salt thereof:



Formula (XIIa)

wherein:


R^a is selected from the group consisting of hydrogen, halogen, -OR⁴, -CN, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, and unsubstituted or substituted C₃-C₆ cycloalkyl; and

R^b is selected from the group consisting of hydrogen, halogen, -OR⁴, -CN, unsubstituted or substituted C₁-C₆ alkyl, and unsubstituted or substituted C₁-C₆ fluoroalkyl.

[0200] In some embodiments, R^a is selected from the group consisting of hydrogen, F, Cl, Br, -CN, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -C(CH₃)₃, -CH₂F, -CHF₂, and -CF₃; and R^b is selected from the group consisting of hydrogen, F, Cl, -CH₃, -CH₂F, -CHF₂, -CF₃, -CN, and -OCH₃.


[0201] In some embodiments, X¹ is N; and X² is CH or N. In some embodiments, X² is N. In some embodiments, X² is CH.

[0202] In some embodiments, the compound has the structure of Formula (XIIf), or a pharmaceutically acceptable salt thereof:



Formula (XIIf).

[0203] In some embodiments, the compound has the structure of Formula (XIIf), or a pharmaceutically acceptable salt thereof:



Formula (XIIf).

wherein:


R^a is selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, unsubstituted or substituted C_1-C_6 alkyl, unsubstituted or substituted C_1-C_6 fluoroalkyl, and unsubstituted or substituted C_3-C_6 cycloalkyl; and

R^b is selected from the group consisting of hydrogen, halogen, $-OR^4$, $-CN$, unsubstituted or substituted C_1-C_6 alkyl, and unsubstituted or substituted C_1-C_6 fluoroalkyl.

[0204] In some embodiments, R^a is selected from the group consisting of hydrogen, F, Cl, Br, $-CN$, $-OCH_3$, $-OCH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2F$, $-CHF_2$, and $-CF_3$; and R^b is selected from the group consisting of hydrogen, F, Cl, $-CH_3$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CN$, and $-OCH_3$.


[0205] In some embodiments, X^1 is N; and X^2 is CH or N. In some embodiments, X^2 is N. In some embodiments, X^2 is CH.

[0206] In some embodiments, the compound has the structure of Formula (XIII), or a pharmaceutically acceptable salt thereof:



Formula (XIII)
wherein W is CH or N.

[0207] In some embodiments, the compound has the structure of Formula (XIIIa), or a pharmaceutically acceptable salt thereof:



Formula (XIIIa)

wherein:


W is CH or N:

R^d is selected from the group consisting of hydrogen, halogen, -OR⁴, -CN, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, and unsubstituted or substituted C₃-C₆ cycloalkyl; and

R^e and R^f are independently selected from the group consisting of hydrogen, halogen, -OR⁴, -CN, unsubstituted or substituted C₁-C₆ alkyl, and unsubstituted or substituted C₁-C₆ fluoroalkyl.

[0208] In some embodiments, R^d is selected from the group consisting of hydrogen, F, Cl, Br, -CN, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -C(CH₃)₃, -CH₂F, -CHF₂, and -CF₃; R^e and R^f are independently selected from the group consisting of hydrogen, F, Cl, Br, -CH₃, -CH₂F, -CHF₂, -CF₃, -CN, -OH, and -OCH₃.

[0209] In some embodiments, the compound has the structure of Formula (XIIIb), or a pharmaceutically acceptable salt thereof:




Formula (XIIIb)

wherein: X^1 is N; X^2 is CH or N; W is CH or N; and R^d is selected from the group consisting of F, Cl, -CN, -OCH₃, -CH₃, -CH₂F, -CHF₂, and -CF₃.

[0210] In some embodiments, X^2 is N. In some embodiments, X^2 is CH.


[0211] In some embodiments, the compound has the structure of Formula (XIIIc), or a pharmaceutically acceptable salt thereof:



Formula (XIIIc)

wherein W is CH or N.

[0212] In some embodiments, the compound has the structure of Formula (XIIId), or a pharmaceutically acceptable salt thereof:





Formula (XIIId)

wherein: W is CH or N; R^d is selected from the group consisting of hydrogen, halogen, -OR⁴, -CN, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, and unsubstituted or substituted C₃-C₆ cycloalkyl; and R^e and R^f are independently selected from the group consisting of hydrogen, halogen, -OR⁴, -CN, unsubstituted or substituted C₁-C₆ alkyl, and unsubstituted or substituted C₁-C₆ fluoroalkyl.

[0213] In some embodiments, R^d is selected from the group consisting of hydrogen, F, Cl, Br, -CN, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -C(CH₃)₃, -CH₂F, -CHF₂, and -CF₃; R^e and R^f are independently selected from the group consisting of hydrogen, F, Cl, Br, -CH₃, -CH₂F, -CHF₂, -CF₃, -CN, -OH, and -OCH₃.

[0214] In some embodiments, the compound has the structure of Formula (XIIIe), or a pharmaceutically acceptable salt thereof:





Formula (XIIIe)


wherein: X^1 is N; W is CH or N; and R^d is selected from the group consisting of F, Cl, -CN, -OCH₃, -CH₃, -CH₂F, -CHF₂, and -CF₃.

[0215] In some embodiments, R^1 is unsubstituted or substituted C₁-C₆ heteroalkyl containing 1 N atom, wherein any substituted group of R^1 is substituted with one or more halogen, C₁-C₄ alkyl, -N(R⁴)₂, or -OR⁵; or R^1 is unsubstituted or substituted azetidiny, unsubstituted or substituted pyrrolidiny, or unsubstituted or substituted piperidiny; or R^1 is unsubstituted or substituted -(C₁-C₆ alkyl)-(C₂-C₇ heterocycloalkyl), wherein the heterocycloalkyl is an unsubstituted or substituted azetidiny, unsubstituted or substituted pyrrolidiny, or unsubstituted or substituted piperidiny.

[0216] In some embodiments, R^1 is unsubstituted or substituted azetidiny, unsubstituted or substituted pyrrolidiny, or unsubstituted or substituted piperidiny.

[0217] In some embodiments, R^1 is unsubstituted or substituted C₁-C₆ heteroalkyl containing 1 N atom, wherein any substituted group of R^1 is substituted with one or more halogen, C₁-C₄ alkyl, -N(R⁴)₂, or -OR⁵.


[0218] In some embodiments, compounds described herein have the following structure:




[0219] In some embodiments, R^A , R^B , X^1 , X^2 , X^3 , X^4 , and R^1 are as described herein.


[0220] In some embodiments, R^A , R^B , X^1 , X^2 , X^3 , X^4 , and R^1 are as described in Table 1.


[0221] In some embodiments, R^A is





[0222] In some embodiments, R^B is







[0223] In some embodiments, X¹ is CH or N. In some embodiments, X² is CH or N. In some embodiments, X³ is CH, CF, or N. In some embodiments, X⁴ is CH or N.


[0224] In some embodiments, X¹ is CH or N; X² is CH or N; X³ is CH, or CF; and X⁴ is CH. In some embodiments, X¹ is CH or N; X² is CH; X³ is CH, or CF; and X⁴ is CH.

[0225] In some embodiments, R¹ is






[0226] In some embodiments, compounds described herein have the following structure:




[0227] In some embodiments, R^{A} , R^{B} , X^1 , X^2 , X^3 , M and R^1 are as described herein. In some embodiments, R^{A} , R^{B} , X^1 , X^2 , X^3 , M and R^1 are as described in Table 2.

[0228] In some embodiments, R^{A} is



[0229] In some embodiments, R^B is




[0230] In some embodiments, X¹ is CH or N.

[0231] In some embodiments, X² is CH or N.


[0232] In some embodiments, X³ is CH or CF.

[0233] In some embodiments, M is -O-, -NH-,



wherein * indicates the attachment point to R¹.


























[0234] In some embodiments, R^1 is



[0235] Any combination of the groups described above for the various variables is contemplated herein. Throughout the specification, groups and substituents thereof are chosen by one skilled in the field to provide stable moieties and compounds.

[0236] Exemplary compounds of Formula (I) include the compounds described in the following Tables. Compounds that fall outside the scope of Formula (I) are described as Reference compounds in the Tables:

Table 1:

							
Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-1		CH	CH	CH	CH		
1-2		CH	CH	CH	CH		
1-3		CH	CH	CH	CH		
1-4		CH	CH	CH	CH		
1-5		CH	CH	CH	CH		
1-6		CH	CH	CH	CH		
1-7		CH	CH	CH	CH		
1-8		CH	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-9		N	CH	CH	CH		
1-10		N	CH	CH	CH		
1-11		N	CH	CH	CH		
1-12		N	CH	CH	CH		
1-13		N	CH	CH	CH		
1-14		N	CH	CH	CH		
1-15		N	CH	CH	CH		
1-16		N	CH	CH	CH		
1-17		N	CH	CH	CH		
1-18		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-19		N	CH	CH	CH		
1-20		N	CH	CH	CH		
1-21		CH	N	CH	CH		
1-22		CH	N	CH	CH		
1-23		CH	CH	CH	CH		
1-24		N	CH	CH	CH		
1-25		N	CH	CH	CH		
1-26		N	CH	CH	CH		
1-27		N	CH	CH	CH		
1-28		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
						CN	
1-29		CH	CH	CH	CH		
1-30		N	CH	CH	CH		
1-31		N	CH	CH	CH		
1-32		N	CH	CH	CH		
1-33		N	CH	CH	CH		
1-34		N	CH	CH	CH		
1-35		N	CH	CH	CH		
1-36		N	CH	CH	CH		
1-37		N	CH	CH	CH		
1-38		N	CH	CH	CH		
1-39		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-40		N	CH	CH	CH		
1-41		N	CH	CH	CH		
1-42		N	CH	CH	CH		
1-43		N	CH	CH	CH		
1-44		N	CH	CH	CH		
1-45		N	CH	CH	CH		
1-46		N	CH	CH	CH		
1-47		N	CH	CH	CH		
1-48		N	CH	CH	CH		
1-51		N	CH	CH	N		
































Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-52		N	CH	CH	CH		
1-53 (Ref)		N	CH	CH	CH		
1-54 (Ref)		N	CH	CH	CH		
1-55		N	CH	CH	CH		
1-56		N	N	CH	CH		
1-57		N	N	CH	CH		
1-58		N	CH	CH	N		
1-59		N	CH	CH	CH		
1-60		N	CH	CH	CH		
1-61		N	CH	CH	CH		
1-62		N	CH	CH	CH		






























Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-63		N	CH	CH	CH		
1-64		N	CH	CH	CH		
1-65		N	CH	CH	CH		
1-66		N	CH	CH	CH		
1-67 (Ref)		N	CH	CH	CH		
1-68		N	CH	CH	CH		
1-69		N	CH	CH	CH		
1-70		N	CH	CH	CH		
1-71		N	CH	CH	CH		
1-72	F	N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-73 (Ref)		N	CH	CH	CH		
1-74		N	CH	CH	CH		
1-75		CH	CH	CF	CH		
1-76		N	CH	CH	CH		
1-77		N	CH	CH	CH		
1-78		N	CH	CH	CH		
1-79		N	CH	CH	CH		
1-80		N	CH	CH	CH		
1-81		N	CH	CH	CH		
1-82		N	CH	CH	CH		































Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-83		N	CH	CH	CH		
1-84		N	CH	CH	CH		
1-85		N	CH	CH	CH		
1-86		N	CH	CH	CH		
1-87		N	CH	CH	CH		
1-88		N	CH	CH	CH		
1-89		N	CH	CH	CH		
1-90		N	CH	CH	CH		
1-91		N	CH	CH	CH		
1-92		N	CH	CH	CH		






























Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-93		N	CH	CH	CH		
1-94		N	CH	CH	CH		
1-95		N	CH	CH	CH		
1-96		N	CH	CH	CH		
1-97		N	CH	CH	CH		
1-98		N	CH	CH	CH		
1-99		N	CH	CH	CH		
1-100		N	CH	CH	CH		
1-101		N	CH	CH	CH		
1-102		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
							
1-103		N	CH	CH	CH		
1-104		N	CH	CH	CH		
1-105		N	CH	CH	CH		
1-106		N	CH	N	CH		
1-107		N	CH	N	CH		
1-108		N	CH	CF	CH		
1-109		N	CH	CF	CH		
1-110		N	CH	CF	CH		
1-111		N	CH	CH	CH		
1-112		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-113		N	CH	CH	CH		
1-114		N	CH	CH	CH		
1-115		N	CH	CH	CH		
1-116		N	CH	CH	CH		
1-117		N	CH	CH	CH		
1-118		N	CH	CH	CH		
1-119		N	CH	CH	CH		
1-120		N	CH	CH	CH		
1-121		N	CH	CH	CH		
1-122		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-123		N	CH	CF	CH		
1-124		N	CH	CF	CH		
1-125		N	CH	CF	CH		
1-126		N	CH	CF	CH		
1-127		N	CH	CF	CH		
1-128		N	CH	CF	CH		
1-129		N	CH	CH	CH		
1-130		N	CH	CH	CH		
1-131		N	CH	CF	CH		
1-132		N	CH	CF	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-133		N	CH	CF	CH		
1-134		N	N	CH	CH		
1-135		N	N	CH	CH		
1-136		N	N	CH	CH		
1-137		N	CH	CH	CH		
1-138		N	CH	CH	CH		
1-139		N	N	CH	CH		
1-140		N	N	CH	CH		
1-141		N	CH	CH	CH		
1-142		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-143		N	CH	CH	CH		
1-144		N	CH	CH	CH		
1-145		N	CH	CH	CH		
1-146		N	CH	CH	CH		
1-147		N	CH	CH	CH		
1-148		N	CH	CH	CH		
1-149		N	CH	CH	CH		
1-150		N	CH	CH	CH		
1-151		N	CH	CH	CH		
1-152		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-153		N	CH	CH	CH		
1-154		N	CH	CH	CH		
1-155		N	CH	CH	CH		
1-156		N	CH	CH	CH		
1-157		N	CH	CH	CH		
1-158		N	CH	CH	CH		
1-159		CH	CH	CH	CH		
1-160		N	CH	CH	CH		
1-161		N	CH	CF	CH		
1-162		N	CH	CF	CH		
































Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-163		N	CH	CF	CH		
1-164		CH	CH	CH	CH		
1-165		CH	CH	CH	CH		
1-166		N	CH	CH	CH		
1-167		N	CH	CH	CH		
1-168		N	N	CH	CH		
1-169		N	N	CH	CH		
1-170		N	N	CH	CH		
1-171		N	N	CH	CH		
1-172		N	CH	CH	CH		
1-173		N	CH	CH	CH		



























Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-174		N	N	CH	CH		
1-175		N	N	CH	CH		
1-176		N	CH	CH	CH		
1-177		N	CH	CH	CH		
1-178		N	CH	CH	CH		
1-179		N	CH	CH	CH		
1-180		N	CH	CH	CH		
1-181		N	CH	CH	CH		
1-182		N	CH	CH	CH		
1-183		N	N	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-184		N	CH	CH	CH		
1-185		N	CH	CH	CH		
1-186		N	CH	CH	CH		
1-187		N	CH	CH	CH		
1-188		N	CH	CH	CH		
1-189 (Ref)		N	CH	CH	CH		
1-190 (Ref)		N	CH	CH	CH		
1-191 (Ref)		N	CH	CH	CH		
1-192		N	CH	CF	CH		
1-193		N	N	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
						F	
1-194		N	N	CH	CH		
1-195		N	CH	CF	CH		
1-196		N	CH	CF	CH		
1-197		N	CH	CF	CH		
1-198		N	CH	CF	CH		
1-199		N	CH	CF	CH		
1-200		N	CH	CF	CH		
1-201		N	CH	CF	CH		
1-202		N	CH	CF	CH		
1-203		N	CH	CF	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-204		N	CH	CF	CH		
1-205		N	CH	CH	CH		
1-206		N	CH	CH	CH		
1-207		N	CH	CH	CH		
1-208		N	CH	CH	CH		
1-209		N	CH	CF	CH		
1-210		N	CH	CF	CH		
1-211		N	CH	CH	CH		
1-212		N	CH	CH	CH		
1-213		N	CH	CH	CH		
1-214		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
							
1-215		N	CH	CH	CH		
1-216		N	CH	CH	CH		
1-217		N	CH	CH	CH		
1-218		N	CH	CH	CH		
1-219		N	CH	CH	CH		
1-220		N	CH	CH	CH		
1-221		N	CH	CH	CH		
1-222		N	CH	CH	CH		
1-223		N	CH	CH	CH		
1-224		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
							
1-225		N	CH	CH	CH		
1-226		N	CH	CH	CH		
1-227		N	CH	CH	CH		
1-228		N	CH	CH	CH		
1-229		N	CH	CH	CH		
1-230		N	CH	CH	CH		
1-231		N	CH	CH	CH		
1-232		N	CH	CH	CH		
1-233		N	CH	CH	CH		
1-234		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-235		N	CH	CH	CH		
1-236		N	CH	CH	CH		
1-237		N	CH	CH	CH		
1-238		N	CH	CH	CH		
1-239		N	CH	CH	CH		
1-240		N	CH	CH	CH		
1-241		N	CH	CH	CH		
1-242		N	CH	CH	CH		
1-243		N	CH	CH	CH		
1-244		N	CH	CH	CH		

































Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-245		N	CH	CH	CH		
1-246		N	CH	CH	CH		
1-247		N	CH	CH	CH		
1-248		N	CH	CH	CH		
1-249		N	CH	CH	CH		
1-250		N	CH	CH	CH		
1-251		N	CH	CH	CH		
1-252		N	CH	CH	CH		
1-253		N	CH	CH	CH		
1-254		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-255		N	CH	CH	CH		
1-256		N	CH	CH	CH		
1-257		N	CH	CH	CH		
1-258		N	CH	CH	CH		<i>*cis-</i>
1-259		N	CH	CH	CH		<i>*trans-</i>
1-260		N	CH	CH	CH		
1-261		N	CH	CH	CH		
1-262		N	CH	CH	CH		
1-263		N	CH	CH	CH		
1-264		N	CH	CH	CH		<i>*trans-</i>

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-265		N	CH	CH	CH		
1-266		N	CH	CH	CH		
1-267		N	CH	CH	CH		
1-268		N	CH	CH	CH		
1-269		N	CH	CH	CH		
1-270		N	CH	CH	CH		
1-271		N	CH	CH	CH		
1-272		N	CH	CH	CH		
1-273		N	CH	CH	CH		
1-274		N	CH	CH	CH		






























Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-275		N	CH	CH	CH		
1-276		N	CH	CH	CH		
1-277		N	CH	CH	CH		
1-278		N	CH	CH	CH		
1-279		N	CH	CH	CH		
1-280		N	CH	CH	CH		
1-281		N	CH	CH	CH		
1-282		N	CH	CH	CH		
1-283		N	CH	CH	CH		
1-284		N	CH	CF	CH		




Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-285		N	CH	CF	CH		
1-286		N	CH	CF	CH		
1-287		N	CH	CF	CH		
1-288		N	CH	CF	CH		
1-289		N	CH	CF	CH		
1-290		N	CH	CF	CH		
1-291		N	CH	CF	CH		
1-292		N	CH	CF	CH		
1-293		N	CH	CF	CH		
1-294		N	CH	CF	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
							
1-295		N	CH	CH	CH		
1-296		N	CH	CH	CH		
1-297		N	CH	CH	CH		
1-298		N	CH	CH	CH		
1-299		N	CH	CF	CH		
1-300		N	CH	CF	CH		
1-301		N	CH	CF	CH		
1-302		N	CH	CF	CH		
1-303		N	CH	CF	CH		
1-304		N	CH	CF	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-305		N	CH	CF	CH		
1-306		N	CH	CH	CH		
1-307		N	CH	CH	CH		
1-308		N	CH	CF	CH		
1-309		N	CH	CF	CH		
1-310		N	CH	CF	CH		
1-311		N	CH	CF	CH		
1-312		N	CH	CF	CH		
1-313		N	CH	CH	CH		
1-314		N	CH	CH	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-315		N	CH	CF	CH		
1-316		N	CH	CH	CH		
1-317		N	CH	CH	CH		
1-318		N	CH	CH	CH		
1-319		N	CH	CF	CH		
1-320		N	CH	CF	CH		
1-321		N	CH	CH	CH		
1-322		N	CH	CH	CH		
1-323		N	CH	CF	CH		
1-324		N	CH	CF	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-325		N	CH	CF	CH		
1-326		N	CH	CF	CH		
1-327		N	CH	CH	CH		
1-328		N	CH	CH	CH		
1-329		N	CH	CH	CH		
1-330		N	CH	CH	CH		
1-331		N	N	CH	CH		
1-332		N	CH	CH	CH		
1-333		N	CH	CH	CH		
1-334		N	CH	CF	CH		

Cpd No.	R ^A	X ¹	X ²	X ³	X ⁴	R ^B	R ¹
1-335		N	N	CH	CH		
*absolute stereochemistry has not been determined. ** <i>cis</i> - & <i>trans</i> - were assigned tentatively. (Ref = Reference compound in Table 1)							

[0237] Compounds in Table 1 are named:

1-1: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-2: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-pyrrolidin-3-yl]piperidine-4-carboxamide;

1-3: N-(2-aminoethyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxamide;

1-4: N-[(2R)-2-aminopropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxamide;

1-5: N-(3-aminopropyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxamide;

1-6: N-[(2S)-2-aminopropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxamide;

1-7: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-8: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxamide;

1-9: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-10: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-pyrrolidin-3-yl]piperidine-4-carboxamide;

1-11: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-12: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-pyrrolidin-3-yl]piperidine-4-carboxamide;

- 1-13:** N-(2-aminoethyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-14:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-15:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-16:** N-[(2S)-2-aminopropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-17:** N-[(2R)-2-amino-3-hydroxypropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-18:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(1-methylazetidin-3-yl)methyl]piperidine-4-carboxamide;
- 1-19:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;
- 1-20:** N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-21:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}-N-[(3R)-pyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-22:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-23:** 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-24:** 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-25:** 3-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-[(3R)-1-methylpyrrolidin-3-yl]carbamoyl)piperidin-1-yl)-6-(trifluoromethyl)pyridine-2-carboxamide;
- 1-26:** 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-27:** 3-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-[(3S)-1-methylpyrrolidin-3-yl]carbamoyl)piperidin-1-yl)-6-(trifluoromethyl)pyridine-2-carboxamide;
- 1-28:** N-[(2S)-2-amino-3-hydroxypropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-29:** 1-[2-chloro-6-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-

[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-30: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-31: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-32: 1-(2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-33: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]-1-[4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;

1-34: 1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-35: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]-1-[5-(trifluoromethyl)pyridin-2-yl]piperidine-4-carboxamide;

1-36: 1-(4-acetyl-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-37: 1-[2-cyano-4-(difluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-38: 1-[4-cyano-2-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-39: 1-[2-cyano-4-(trifluoromethoxy)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-40: 1-(2,4-dicyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-41: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S,4S)*-4-hydroxy-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-42: 1-[3-cyano-5-(trifluoromethyl)pyridin-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-43: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-44: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(2R)-3-hydroxy-2-(methylamino)propyl]piperidine-4-carboxamide;

1-45: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2R)-2-(dimethylamino)-3-hydroxypropyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

- 1-46:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R,4R)*-4-hydroxy-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-47:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-3-hydroxy-2-(methylamino)propyl]piperidine-4-carboxamide;
- 1-48:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-2-(dimethylamino)-3-hydroxypropyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-51:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-52:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-53:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2,3-dihydro-1-benzofuran-7-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-54:** 4-[6-(1-benzofuran-7-yl)pyridin-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-55:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-[2-(hydroxymethyl)phenyl]pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-56:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-57:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-58:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-59:** 1-[2-cyano-4-(1,1-difluoroethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-60:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-61:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2,3-difluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-62:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-ethyl-1H-pyrazol-5-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-63:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-[6-(2-propoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-64:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3-fluorophenyl)pyridin-3-yl]-N-[(3R)-1-

methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-65: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3,5-difluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-66: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-67: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2,2-difluoro-2H-1,3-benzodioxol-4-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-68: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2,5-difluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-69: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxy-5-fluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-70: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{6-[1-(3-methylbutyl)-1H-pyrazol-5-yl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-71: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-{6-[2-(trifluoromethoxy)phenyl]pyridin-3-yl}piperidine-4-carboxamide;

1-72: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxy-3-fluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-73: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-indazol-7-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-74: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-fluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-75: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)-3-fluorophenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-76: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;

1-77: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-{6-[2-(trifluoromethyl)phenyl]pyridin-3-yl}piperidine-4-carboxamide;

1-78: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-cyanophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-79: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{6-[2-(methoxymethyl)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-80: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3-methoxythiophen-2-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

- 1-81:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-82:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-83:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxythiophen-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-84:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-[2-(1,1-difluoroethyl)phenyl]pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-85:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2'-(difluoromethoxy)-[2,3'-bipyridin]-5-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-86:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5'-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-87:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3,5-difluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-88:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-89:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-cyclopropoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-90:** 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-91:** 1-(4-chloro-2-cyanophenyl)-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-92:** 1-(4-chloro-2-cyanophenyl)-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;
- 1-93:** 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;
- 1-94:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;
- 1-95:** 1-(2,4-dichlorophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-96:** 1-(2-cyano-4-fluorophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-97:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-

methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-98: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-99: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-fluoro-2-methylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-100: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-101: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-102: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-103: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{5'-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-104: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethylphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-105: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(4-methylpyrimidin-5-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-106: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-107: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-108: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-109: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-110: 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-111: 1-(4-chloro-2-cyano-6-fluorophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-112: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-[6-(2-methylthiophen-3-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-113: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-cyclopropylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

- 1-114:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{6-[2-(difluoromethoxy)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-115:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-116:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methylfuran-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-117:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-[6-(2-propylphenyl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-118:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethyl-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-119:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxythiophen-3-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-120:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methylfuran-3-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-121:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-cyclopropylphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-122:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]-4-[6-(2-methylthiophen-3-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-123:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-124:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-125:** 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-126:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-127:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-128:** 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-129:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3-fluoro-2-hydroxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;
- 1-130:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxy-5-methylphenyl)pyridin-3-yl]-N-

[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-131: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-132: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-133: 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-134: 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-135: 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;

1-136: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;

1-137: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{6-[2-(cyanomethyl)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-138: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-[6-(4-methylthiophen-3-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-139: 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-140: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;

1-141: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-142: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-143: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]-4-[6-(1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-144: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-145: 4-[6-(2-ethoxyphenyl)pyridin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-146: 4-[6-(2-acetylthiophen-3-yl)pyridin-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-147: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethylthiophen-3-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-148: 4-[6-(2-cyano-3-fluorophenyl)pyridin-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-149: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(3-methyl-1,2-oxazol-4-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-150: 1-[3-chloro-5-(trifluoromethyl)pyridin-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-151: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{6-[2-(difluoromethyl)phenyl]pyridin-3-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-152: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-pyrrolidin-3-yl]piperidine-4-carboxamide;

1-153: 4-[6-(5-cyano-1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-154: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{5'-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-155: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-156: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethylphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-157: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-methyl-1,2-oxazol-4-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-158: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-cyclopropyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-159: 4-[4-(2-ethoxypyridin-3-yl)phenyl]-1-[2-methoxy-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-160: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-161: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-162: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-163: 1-(4-chloro-2-cyanophenyl)-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-

[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-164: 4-[4-(2-ethoxypyridin-3-yl)phenyl]-1-[2-methyl-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-165: 1-[3-chloro-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-166: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-167: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxythiophen-3-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-168: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-169: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-170: 1-(4-chloro-2-cyanophenyl)-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-171: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-172: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[2'-ethyl-[2,3'-bipyridin]-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-173: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-174: 4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-175: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-176: 4-[[2,2'-bipyridin]-5-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-177: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[3'-methyl-[2,2'-bipyridin]-5-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-178: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[3'-methoxy-[2,2'-bipyridin]-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-179: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2'-methoxy-[2,3'-bipyridin]-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-180: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-cyclopropyl-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-181: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2'-(difluoromethyl)-[2,3'-bipyridin]-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-182: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-cyclopropyl-1,3-oxazol-4-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-183: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-184: ethyl (3S)-3-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-amido}pyrrolidine-1-carboxylate;

1-185: N-[(3S)-1-acetylpyrrolidin-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-186: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-187: N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-188: N-[(3R)-1-azabicyclo[2.2.2]octan-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-189: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyquinolin-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-190: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-indol-7-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-191: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-indol-2-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-192: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-193: 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-194: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-195: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-196: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-

(methylamino)ethyl]piperidine-4-carboxamide;

1-197: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[3-(methylamino)propyl]piperidine-4-carboxamide;

1-198: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-199: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[3-(dimethylamino)propyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-200: 4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-201: 4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-202: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-203: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-204: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-205: N-(3-aminopropyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-206: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[3-(methylamino)propyl]piperidine-4-carboxamide;

1-207: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[3-(dimethylamino)propyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-208: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-ethyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-209: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-210: 4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide;

1-211: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(1-methylazetidin-3-yl)methyl]piperidine-4-carboxamide;

1-212: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

- 1-213:** N-[(2S)-azetidin-2-yl]methyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-214:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-215:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-216:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-217:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(2S)-1-methylazetidin-2-yl]methyl]piperidine-4-carboxamide;
- 1-218:** N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-219:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide;
- 1-220:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1-methylazetidin-3-yl)methyl]piperidine-4-carboxamide;
- 1-221:** N-{1-azabicyclo[2.2.1]heptan-4-yl}-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-222:** N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;
- 1-223:** 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-224:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-225:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-226:** N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;
- 1-227:** 4-[6-(2-ethoxyphenyl)pyridin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-228:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-229:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-

{[(2S)-1-methylpyrrolidin-2-yl]methyl}piperidine-4-carboxamide;

1-230: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-231: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-{[(2R)-pyrrolidin-2-ylmethyl]piperidine-4-carboxamide};

1-232: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-{[(2R)-1-methylpyrrolidin-2-yl]methyl}piperidine-4-carboxamide;

1-233: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1S,3S)-3-aminocyclobutyl]piperidine-4-carboxamide;

1-234: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1R,3R)-3-aminocyclobutyl]piperidine-4-carboxamide;

1-235: N-{[(2S)-azetidin-2-yl]methyl}-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-236: N-{[(2R)-azetidin-2-yl]methyl}-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-237: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R,4S)-4-fluoropyrrolidin-3-yl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-238: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R,4R)-4-fluoropyrrolidin-3-yl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-239: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1S,3S)-3-(dimethylamino)cyclobutyl]piperidine-4-carboxamide;

1-240: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1R,3R)-3-(dimethylamino)cyclobutyl]piperidine-4-carboxamide;

1-241: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-{[(2S)-1-methylazetidin-2-yl]methyl}piperidine-4-carboxamide;

1-242: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R,4S)-4-fluoro-1-methylpyrrolidin-3-yl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-243: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R,4R)-4-fluoro-1-methylpyrrolidin-3-yl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-244: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-{[(2R)-1-methylazetidin-2-yl]methyl}piperidine-4-carboxamide;

1-245: N-[2-(azetidin-1-yl)ethyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-246: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[[2-(2S,4S)-4-fluoropyrrolidin-2-yl]methyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-247: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[[2-(2S,4R)-4-fluoropyrrolidin-2-yl]methyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-248: N-[2-(azetidin-1-yl)ethyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-249: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[[2-(2S,4S)-4-fluoro-1-methylpyrrolidin-2-yl]methyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-250: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[[2-(2S,4R)-4-fluoro-1-methylpyrrolidin-2-yl]methyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-251: rac-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1S,2S,4R)-7-methyl-7-azabicyclo[2.2.1]heptan-2-yl]piperidine-4-carboxamide;

1-252: rac-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(1S,2S,4R)-7-methyl-7-azabicyclo[2.2.1]heptan-2-yl]piperidine-4-carboxamide;

1-253: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1,3-dimethylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-254: N-[2-(dimethylamino)ethyl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-255: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;

1-256: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-257: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-258: rac-N-[(1R,2S)-2-aminocyclopropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-259: rac-N-[(1R,2R)-2-aminocyclopropyl]-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-260: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2R)-1-(dimethylamino)propan-2-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-261: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-262: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-{3-[(dimethylamino)methyl]oxetan-3-yl}-4-[6-(1-

methyl-1H-pyrrol-2-yl]pyridin-3-yl]piperidine-4-carboxamide;

1-263: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[[1-(dimethylamino)cyclopropyl)methyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-264: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S,4S)-4-(dimethylamino)oxolan-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-265: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-{2-[ethyl(methyl)amino]ethyl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-266: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-{2-[cyclopropyl(methyl)amino]ethyl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-267: rac-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S,4R)-4-methoxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-268: rac-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R,4R)-4-methoxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-269: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R,4R)-4-hydroxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-270: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S,4S)-4-hydroxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-271: N-[2-(dimethylamino)ethyl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-272: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-273: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-274: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-275: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2R)-2-(dimethylamino)-3-hydroxypropyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-276: N-[2-(dimethylamino)ethyl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-277: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[[[(2S,4S)-4-fluoro-1-methylpyrrolidin-2-yl)methyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-278: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[[[(2S,4R)-4-fluoro-1-methylpyrrolidin-2-yl)methyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

- 1-279:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-4,4-difluoro-1-methylpyrrolidin-2-yl)methyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-280:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2R)-4,4-difluoro-1-methylpyrrolidin-2-yl)methyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-281:** 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-282:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-283:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-284:** N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;
- 1-285:** N-[2-(dimethylamino)ethyl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-286:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-287:** 4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-288:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-289:** N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;
- 1-290:** 4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-291:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]piperidine-4-carboxamide;
- 1-292:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-293:** 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-294:** 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;
- 1-295:** N-[(2S)-1-(dimethylamino)propan-2-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-{2'-

methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-296: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-297: N-[(2S)-1-(dimethylamino)propan-2-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-298: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;

1-299: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-300: N-[2-(dimethylamino)ethyl]-4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;

1-301: 4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-302: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-303: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{3-fluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-304: N-[2-(dimethylamino)ethyl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-305: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-306: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;

1-307: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;

1-308: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;

1-309: 1-[2-chloro-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;

1-310: 1-[2-chloro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

1-311: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidine-4-carboxamide;

- 1-312:** N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;
- 1-313:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-314:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;
- 1-315:** 4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;
- 1-316:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;
- 1-317:** N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide;
- 1-318:** 1-(2,4-dichlorophenyl)-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-319:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-320:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;
- 1-321:** 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;
- 1-322:** 1-(2,4-dichlorophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;
- 1-323:** 1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;
- 1-324:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide;
- 1-325:** 1-(4-chloro-2-cyanophenyl)-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;
- 1-326:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidine-4-carboxamide;
- 1-327:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(2S)-4-methylmorpholin-2-yl]methylpiperidine-4-carboxamide;
- 1-328:** 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-4-

methylmorpholin-3-yl)methyl}piperidine-4-carboxamide;

1-329: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-(((3R)-4-methylmorpholin-3-yl)methyl)piperidine-4-carboxamide;

1-330: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-(((2R)-4-methylmorpholin-2-yl)methyl)piperidine-4-carboxamide;

1-331: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidine-4-carboxamide;

















1-332: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2S)-2-(dimethylamino)propyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;





































1-333: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2R)-2-(dimethylamino)propyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;


































1-334: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(2R)-2-(dimethylamino)-3-hydroxypropyl]-4-{2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl}piperidine-4-carboxamide;




































1-335: N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide.



































Table 2:



























							
Cpd No.	R ^A	X ¹	X ²	X ³	R ^B	R ¹	-M-
2-1		CH	CH	CH			-O-
2-2		CH	CH	CH			
2-3		CH	CH	CH			
2-4		N	CH	CH			



















							
Cpd No.	R ^A	X ¹	X ²	X ³	R ^B	R ¹	-M-
							
2-5		N	CH	CH			
2-6		N	CH	CH			
2-7		N	CH	CH			
2-8		N	CH	CH			
2-9		N	CH	CH			
2-10		N	CH	CH			
2-11		CH	CH	CH			
2-12		CH	CH	CH			






































							
Cpd No.	R ^A	X ¹	X ²	X ³	R ^B	R ¹	-M-
2-13		CH	CH	CH			
2-14		N	CH	CH			-NH-
2-15		N	CH	CH			-NH-
2-16		CH	CH	CH			
2-17		CH	CH	CH			
2-18		N	CH	CH			
2-19		N	CH	CH			
2-20		N	CH	CH			
2-21		N	CH	CH			




							
Cpd No.	R ^A	X ¹	X ²	X ³	R ^B	R ¹	-M-
							
2-22		N	CH	CH			
2-23		N	CH	CH			
2-24		N	CH	CH			
2-25		N	CH	CH			
2-26		N	CH	CH			
2-27		N	CH	CH			
2-28		N	CH	CH			
2-29		N	CH	CH			

							
Cpd No.	R ^A	X ¹	X ²	X ³	R ^B	R ¹	-M-
2-30		N	CH	CH			
2-31		N	CH	CH			
2-32		N	CH	CH			
2-33		N	CH	CH			
2-34		N	CH	CH			
2-35		N	CH	CH			
2-36		N	CH	CH			
2-37		N	CH	CH			
2-38		N	CH	CH			

							
Cpd No.	R ^A	X ¹	X ²	X ³	R ^B	R ¹	-M-
							
2-39		N	CH	CH			
2-40		N	CH	CH			
2-41		N	CH	CH			
2-42		N	CH	CH			
2-43		N	CH	CH			
2-44		N	CH	CH			
2-45		N	CH	CH			
2-46		N	CH	CH			
2-47		N	CH	CH			

							
Cpd No.	R ^A	X ¹	X ²	X ³	R ^B	R ¹	-M-
							
2-48		N	CH	CH			
2-49		N	CH	CF			
2-50		N	CH	CF			
2-51		N	CH	CF			
2-52		N	CH	CF			
2-53		N	CH	CH			
2-54		N	CH	CH			
2-55		N	CH	CH			

							
Cpd No.	R ^A	X ¹	X ²	X ³	R ^B	R ¹	-M-
2-56		N	CH	CH			
2-57		N	CH	CH			
2-58		N	CH	CH			
2-59		N	CH	CH			
2-60		N	CH	CH			
2-61		N	CH	CF			
2-62		N	CH	CF			
2-63		N	CH	CF			
2-64		N	N	CH			

							
Cpd No.	R ^A	X ¹	X ²	X ³	R ^B	R ¹	-M-
							
* attachment point to R ¹							

[0238] Compounds in Table 2 are named:

2-1: 2-{4-[2-(dimethylamino)ethoxy]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-1-yl}-5-(trifluoromethyl)benzonitrile;

2-2: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl N-[2-(dimethylamino)ethyl]carbamate;

2-3: 3-amino-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}propanamide;

2-4: 3-amino-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}propanamide;

2-5: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)propanamide;

2-6: (3R)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}pyrrolidine-3-carboxamide;

2-7: (3R)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;

2-8: (3S)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}pyrrolidine-3-carboxamide;

2-9: (3S)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;

2-10: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(dimethylamino)propanamide;

2-11: (2S)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}-1-methylpyrrolidine-2-carboxamide;

2-12: (3R)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamate;

2-13: (3S)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamate;

2-14: 2-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-[(2-hydroxyethyl)amino]piperidin-1-yl)-5-(trifluoromethyl)benzonitrile;

2-15: 2-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-[[2-(methylamino)ethyl]amino]piperidin-1-yl)-5-(trifluoromethyl)benzonitrile;

2-16: N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}-1-methylazetidine-3-carboxamide;

2-17: (2R)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}-1-methylpyrrolidine-2-carboxamide;

2-18: (3S)-1-methylpyrrolidin-3-yl N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamate;

2-19: 1-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-[(3R)-1-methylpyrrolidin-3-yl]urea;

2-20: 1-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-[(3S)-1-methylpyrrolidin-3-yl]urea;

2-21: (3R)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)pyrrolidine-1-carboxamide;

2-22: (3R)-1-methylpyrrolidin-3-yl N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamate;

2-23: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)azetidine-1-carboxamide;

2-24: (3S)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)pyrrolidine-1-carboxamide;

2-25: N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-1-methylazetidine-3-carboxamide;

2-26: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-2-(dimethylamino)acetamide;

2-27: (3S)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;

2-28: (3S)-N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-1-

methylpyrrolidine-3-carboxamide;

2-29: (3R)-N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-1-methylpyrrolidine-3-carboxamide;

2-30: N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-3-(dimethylamino)propanamide;

2-31: N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-2-(dimethylamino)acetamide;

2-32: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-1-methylazetidine-3-carboxamide;

2-33: (3R)-N- { 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;

2-34: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-2-(dimethylamino)acetamide;

2-35: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-1-methylazetidine-3-carboxamide;

2-36: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamide;

2-37: N-[1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl]-1-methylazetidine-3-carboxamide;

2-38: N-[1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl]-2-(dimethylamino)acetamide;

2-39: (3S)-N-[1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl]-1-methylpyrrolidine-3-carboxamide;

2-40: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-1-methylazetidine-3-carboxamide;

2-41: (3R)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}carbamate;

2-42: (3S)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}carbamate;

2-43: 1-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-3-[(3S)-1-methylpyrrolidin-3-yl]urea;

2-44: (3S)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}pyrrolidine-3-carboxamide;

- 2-45:** N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-1-methylazetidine-3-carboxamide;
- 2-46:** (3S)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;
- 2-47:** (3R)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;
- 2-48:** 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-3-[(3R)-1-methylpyrrolidin-3-yl]urea;
- 2-49:** (3S)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;
- 2-50:** (3R)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;
- 2-51:** 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[(3S)-1-methylpyrrolidin-3-yl]urea;
- 2-52:** 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[(3R)-1-methylpyrrolidin-3-yl]urea;
- 2-53:** (3S)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;
- 2-54:** (3R)-1-methylpyrrolidin-3-yl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;
- 2-55:** 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[(3S)-1-methylpyrrolidin-3-yl]urea;
- 2-56:** 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[(3R)-1-methylpyrrolidin-3-yl]urea;
- 2-57:** 2-(dimethylamino)ethyl N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;
- 2-58:** 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[2-(dimethylamino)ethyl]urea;
- 2-59:** 1-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(1-methylazetidin-3-yl)urea;
- 2-60:** (3R)-N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidine-3-carboxamide;
- 2-61:** N-{ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-

yl]piperidin-4-yl}-2-(dimethylamino)acetamide;

2-62: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamide;

2-63: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(methylamino)propanamide;

2-64: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamide.

[0239] In one aspect, compounds described herein are in the form of pharmaceutically acceptable salts. Also disclosed, and not forming part of the invention, are active metabolites of these compounds having the same type of activity. In addition, the compounds described herein can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. The solvated forms of the compounds presented herein are also considered to be disclosed herein.

[0240] "Pharmaceutically acceptable," as used herein, refers a material, such as a carrier or diluent, which does not abrogate the biological activity or properties of the compound, and is relatively nontoxic, i.e., the material is administered to an individual without causing undesirable biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained.

[0241] The term "pharmaceutically acceptable salt" refers to a form of a therapeutically active agent that consists of a cationic form of the therapeutically active agent in combination with a suitable anion, or in alternative embodiments, an anionic form of the therapeutically active agent in combination with a suitable cation. Handbook of Pharmaceutical Salts: Properties, Selection and Use. International Union of Pure and Applied Chemistry, Wiley-VCH 2002. S.M. Berge, L.D. Bighley, D.C. Monkhouse, J. Pharm. Sci. 1977, 66, 1-19. P. H. Stahl and C. G. Wermuth, editors, Handbook of Pharmaceutical Salts: Properties, Selection and Use, Weinheim/Zürich:Wiley-VCH/VHCA, 2002. Pharmaceutical salts typically are more soluble and more rapidly soluble in stomach and intestinal juices than non-ionic species and so are useful in solid dosage forms. Furthermore, because their solubility often is a function of pH, selective dissolution in one or another part of the digestive tract is possible and this capability can be manipulated as one aspect of delayed and sustained release behaviours. Also, because the salt-forming molecule can be in equilibrium with a neutral form, passage through biological membranes can be adjusted.

[0242] In some embodiments, pharmaceutically acceptable salts are obtained by reacting a compound of Formula (I) with an acid. In some embodiments, the compound of Formula (I) (i.e. free base form) is basic and is reacted with an organic acid or an inorganic acid. Inorganic acids include, but are not limited to, hydrochloric acid, hydrobromic acid, sulfuric acid,

phosphoric acid, nitric acid, and metaphosphoric acid. Organic acids include, but are not limited to, 1-hydroxy-2-naphthoic acid; 2,2-dichloroacetic acid; 2-hydroxyethanesulfonic acid; 2-oxoglutaric acid; 4-acetamidobenzoic acid; 4-aminosalicylic acid; acetic acid; adipic acid; ascorbic acid (L); aspartic acid (L); benzenesulfonic acid; benzoic acid; camphoric acid (+); camphor-10-sulfonic acid (+); capric acid (decanoic acid); caproic acid (hexanoic acid); caprylic acid (octanoic acid); carbonic acid; cinnamic acid; citric acid; cyclamic acid; dodecylsulfuric acid; ethane-1,2-disulfonic acid; ethanesulfonic acid; formic acid; fumaric acid; galactaric acid; gentisic acid; glucoheptonic acid (D); gluconic acid (D); glucuronic acid (D); glutamic acid; glutaric acid; glycerophosphoric acid; glycolic acid; hippuric acid; isobutyric acid; lactic acid (DL); lactobionic acid; lauric acid; maleic acid; malic acid (- L); malonic acid; mandelic acid (DL); methanesulfonic acid; naphthalene-1,5-disulfonic acid; naphthalene-2-sulfonic acid; nicotinic acid; oleic acid; oxalic acid; palmitic acid; pamoic acid; phosphoric acid; propionic acid; pyroglutamic acid (- L); salicylic acid; sebacic acid; stearic acid; succinic acid; sulfuric acid; tartaric acid (+ L); thiocyanic acid; toluenesulfonic acid (p); and undecylenic acid.

[0243] In some embodiments, a compound of Formula (I) is prepared as a chloride salt, sulfate salt, bromide salt, mesylate salt, maleate salt, citrate salt or phosphate salt.

[0244] In some embodiments, pharmaceutically acceptable salts are obtained by reacting a compound of Formula (I) with a base. In some embodiments, the compound of Formula (I) is acidic and is reacted with a base. In such situations, an acidic proton of the compound of Formula (I) is replaced by a metal ion, e.g., lithium, sodium, potassium, magnesium, calcium, or an aluminum ion. In some cases, compounds described herein coordinate with an organic base, such as, but not limited to, ethanolamine, diethanolamine, triethanolamine, tromethamine, meglumine, N-methylglucamine, dicyclohexylamine, tris(hydroxymethyl)methylamine. In other cases, compounds described herein form salts with amino acids such as, but not limited to, arginine, lysine, and the like. Acceptable inorganic bases used to form salts with compounds that include an acidic proton, include, but are not limited to, aluminum hydroxide, calcium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium hydroxide, lithium hydroxide, and the like. In some embodiments, the compounds provided herein are prepared as a sodium salt, calcium salt, potassium salt, magnesium salt, meglumine salt, N-methylglucamine salt or ammonium salt.

[0245] It should be understood that a reference to a pharmaceutically acceptable salt includes the solvent addition forms. Also disclosed herein are solvates which contain either stoichiometric or non-stoichiometric amounts of a solvent, and which are formed during the process of crystallization with pharmaceutically acceptable solvents such as water, ethanol, and the like. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. Solvates of compounds described herein are conveniently prepared or formed during the processes described herein. In addition, the compounds provided herein can optionally exist in unsolvated as well as solvated forms.

[0246] The methods and formulations described herein include the use of *N*-oxides (if appropriate), or pharmaceutically acceptable salts of compounds having the structure of

Formula (I), also disclosed are active metabolites of these compounds having the same type of activity.

[0247] In some embodiments, sites on the organic radicals (e.g. alkyl groups, aromatic rings) of compounds of Formula (I) are susceptible to various metabolic reactions. Incorporation of appropriate substituents on the organic radicals will reduce, minimize or eliminate this metabolic pathway. In specific embodiments, the appropriate substituent to decrease or eliminate the susceptibility of the aromatic ring to metabolic reactions is, by way of example only, a halogen, deuterium, an alkyl group, a haloalkyl group, or a deuterioalkyl group.

[0248] In another embodiment, the compounds described herein are labeled isotopically (e.g. with a radioisotope) or by another other means, including, but not limited to, the use of chromophores or fluorescent moieties, bioluminescent labels, or chemiluminescent labels.

[0249] Compounds described herein include isotopically-labeled compounds, which are identical to those recited in the various formulae and structures presented herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into the present compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, sulfur, fluorine chlorine, iodine, phosphorus, such as, for example, ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{35}S , ^{18}F , ^{36}Cl , ^{123}I , ^{124}I , ^{125}I , ^{131}I , ^{32}P and ^{33}P . In one aspect, isotopically-labeled compounds described herein, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. In one aspect, substitution with isotopes such as deuterium affords certain therapeutic advantages resulting from greater metabolic stability, such as, for example, increased *in vivo* half-life or reduced dosage requirements.

[0250] In some embodiments, the compounds of Formula (I) possess one or more stereocenters and each stereocenter exists independently in either the R or S configuration. In some embodiments, the compound of Formula (I) exists in the R configuration. In some embodiments, the compound of Formula (I) exists in the S configuration. The compounds presented herein include all diastereomeric, individual enantiomers, atropisomers, and epimeric forms as well as the appropriate mixtures thereof. The compounds and methods provided herein include all cis, trans, syn, anti, entgegen (E), and zusammen (Z) isomers as well as the appropriate mixtures thereof.

[0251] Individual stereoisomers are obtained, if desired, by methods such as, stereoselective synthesis and/or the separation of stereoisomers by chiral chromatographic columns or the separation of diastereomers by either non-chiral or chiral chromatographic columns or crystallization and recrystallization in a proper solvent or a mixture of solvents. In certain embodiments, compounds of Formula (I) are prepared as their individual stereoisomers by reacting a racemic mixture of the compound with an optically active resolving agent to form a pair of diastereoisomeric compounds/salts, separating the diastereomers and recovering the

optically pure individual enantiomers. In some embodiments, resolution of individual enantiomers is carried out using covalent diastereomeric derivatives of the compounds described herein. In another embodiment, diastereomers are separated by separation/resolution techniques based upon differences in solubility. In other embodiments, separation of stereoisomers is performed by chromatography or by the forming diastereomeric salts and separation by recrystallization, or chromatography, or any combination thereof. Jean Jacques, Andre Collet, Samuel H. Wilen, "Enantiomers, Racemates and Resolutions", John Wiley And Sons, Inc., 1981. In some embodiments, stereoisomers are obtained by stereoselective synthesis.

[0252] Also disclosed, and not forming part of the invention, are compounds described herein are prepared as prodrugs. A "prodrug" refers to an agent that is converted into the parent drug *in vivo*. Prodrugs are often useful because, in some situations, they are easier to administer than the parent drug. They are, for instance, bioavailable by oral administration whereas the parent is not. Further or alternatively, the prodrug also has improved solubility in pharmaceutical compositions over the parent drug. Also disclosed, the design of a prodrug increases the effective water solubility. An example, without limitation, of a prodrug is a compound described herein, which is administered as an ester (the "prodrug") but then is metabolically hydrolyzed to provide the active entity. A further example of a prodrug is a short peptide (polyaminoacid) bonded to an acid group where the peptide is metabolized to reveal the active moiety. Also disclosed, upon *in vivo* administration, a prodrug is chemically converted to the biologically, pharmaceutically or therapeutically active form of the compound. Also disclosed, a prodrug is enzymatically metabolized by one or more steps or processes to the biologically, pharmaceutically or therapeutically active form of the compound.

[0253] Prodrugs of the compounds disclosed herein include, but are not limited to, esters, ethers, carbonates, thiocarbonates, N-acyl derivatives, N-acyloxyalkyl derivatives, N-alkyloxyacyl derivatives, quaternary derivatives of tertiary amines, N-Mannich bases, Schiff bases, amino acid conjugates, phosphate esters, and sulfonate esters. See for example Design of Prodrugs, Bundgaard, A. Ed., Elsevier, 1985 and Method in Enzymology, Widder, K. et al., Ed.; Academic, 1985, vol. 42, p. 309-396; Bundgaard, H. "Design and Application of Prodrugs" in A Textbook of Drug Design and Development, Krosgaard-Larsen and H. Bundgaard, Ed., 1991, Chapter 5, p. 113-191; and Bundgaard, H., Advanced Drug Delivery Review, 1992, 8, 1-38. Also disclosed, a hydroxyl group in the compounds disclosed herein is used to form a prodrug, wherein the hydroxyl group is incorporated into an acyloxyalkyl ester, alkoxycarbonyloxyalkyl ester, alkyl ester, aryl ester, phosphate ester, sugar ester, ether, and the like. Also disclosed, a hydroxyl group in the compounds disclosed herein is a prodrug wherein the hydroxyl is then metabolized *in vivo* to provide a carboxylic acid group. Also disclosed, a carboxyl group is used to provide an ester or amide (i.e. the prodrug), which is then metabolized *in vivo* to provide a carboxylic acid group. Also disclosed, compounds described herein are prepared as alkyl ester prodrugs.

[0254] Prodrug forms of the herein disclosed compounds, wherein the prodrug is metabolized *in vivo* to produce a compound of Formula (I) as set forth herein are included within the scope

of the claims. In some cases, some of the herein-disclosed compounds is a prodrug for another derivative or active compound.

[0255] Also disclosed, any one of the hydroxyl group(s), amino group(s) and/or carboxylic acid group(s) are functionalized in a suitable manner to provide a prodrug moiety. Also disclosed, the prodrug moiety is as described above.

[0256] Also disclosed, the compounds described herein are metabolized upon administration to an organism in need to produce a metabolite that is then used to produce a desired effect, including a desired therapeutic effect.

[0257] A "metabolite" of a compound disclosed herein is a derivative of that compound that is formed when the compound is metabolized. The term "active metabolite" refers to a biologically active derivative of a compound that is formed when the compound is metabolized. The term "metabolized," as used herein, refers to the sum of the processes (including, but not limited to, hydrolysis reactions and reactions catalyzed by enzymes) by which a particular substance is changed by an organism. Thus, enzymes may produce specific structural alterations to a compound. For example, cytochrome P450 catalyzes a variety of oxidative and reductive reactions while uridine diphosphate glucuronyltransferases catalyze the transfer of an activated glucuronic-acid molecule to aromatic alcohols, aliphatic alcohols, carboxylic acids, amines and free sulphydryl groups. Metabolites of the compounds disclosed herein are optionally identified either by administration of compounds to a host and analysis of tissue samples from the host, or by incubation of compounds with hepatic cells in vitro and analysis of the resulting compounds.

Synthesis of Compounds


[0258] Compounds of Formula (I) described herein are synthesized using standard synthetic techniques or using methods known in the art in combination with methods described herein.

[0259] Unless otherwise indicated, conventional methods of mass spectroscopy, NMR, HPLC, protein chemistry, biochemistry, recombinant DNA techniques and pharmacology are employed.

[0260] Compounds are prepared using standard organic chemistry techniques such as those described in, for example, March's Advanced Organic Chemistry, 6th Edition, John Wiley and Sons, Inc. Alternative reaction conditions for the synthetic transformations described herein may be employed such as variation of solvent, reaction temperature, reaction time, as well as different chemical reagents and other reaction conditions.

[0261] In some embodiments, compounds described herein are prepared as described in Scheme A.

Scheme A




a) $R^B X$, Pd(II), or $R^B X$, DIEA; b) $R^A B(OH)_2$, Pd(DtBPF)Cl₂, K₂CO₃, Dioxane/H₂O; c) LiOH, EtOH; d) $R^1 NH_2$, HATU, TEA; e) deprotection

[0262] Commercially available XVIII is converted to compound XIX by either a standard S_NAr reaction with $R^B X$ or Buchwald-Hartwig cross coupling reaction with $R^B X$. Subsequently, introduction of R^A by Suzuki-Miyaura coupling reaction with $R^A B(OH)_2$ or Stille coupling with $R^A SnBu_3$ yield compound XX. Hydrolysis of the ester to the acid XXI and then HATU-mediated amide coupling reaction with $R^1 NH_2$ produce the final compound XXII. If R^1 contains a protecting group, then an additional deprotection step takes place to produce XXII.

[0263] In some embodiments, compounds described herein are prepared as described in Scheme B.

Scheme B




a) *N*-benzyl-2-bromo-*N*-(2-bromoethyl)ethan-1-amine, KOH; b) $R^A B(OH)_2$, Pd(DtBPF)Cl₂, K₂CO₃, Dioxane/H₂O; c) Pd(OH)₂, Pd/C, H₂; d) $R^B X$, DIEA, or $R^B X$, Pd(II); e) LiOH, MeOH; f) $R^1 NH_2$, HATU, TEA; g) deprotection

[0264] Methyl 2-(6-chloropyridin-3-yl)acetate efficiently undergoes a one-pot reaction with *N*-benzyl-2-bromo-*N*-(2-bromoethyl)ethan-1-amine in the presence of a strong inorganic base such as KOH to produce the piperidinyll intermediate **XXIII**. Subsequently, introduction of R^A by Suzuki-Miyaura coupling reaction with $R^AB(OH)_2$ and then de-benzylation yield compound **XXIV**. R^B is introduced by either a standard S_NAr reaction with R^BX or Buchwald-Hartwig cross coupling reaction with R^BX and followed by hydrolysis to yield the acid **XXV**. Variation of R^1 in the final **XXX** is achieved by HATU-mediated coupling reaction with R^1NH_2 . Starting from the intermediate **XXIII**, introduction of various 5- and 6-membered aromatic rings for R^A (**XXVI-XXVII-XXX**) and diverse R^B (**XXVIII-XXIX-XXX**) is accomplished by change of the reaction sequence as described in the scheme. If R^1 contains a protecting group, then an additional deprotection step takes place to produce **XX**.

[0265] In some embodiments, compounds described herein are prepared as described in Scheme C.

Scheme C




[0266] The cyanomethylation of 5-bromo-2-chloro-3-fluoropyridine to **XXXII** efficiently proceeds through Suzuki-Miyaura coupling reaction with isoxazole-4-boronic acid pinacol ester, base-induced fragmentation, and deformylation. The nitrogen-containing heterocycle **XXXIII** is obtained from cyclocondensation of **XXXII** with *N*-benzyl-2-bromo-*N*-(2-bromoethyl)ethan-1-amine in a strong alkaline medium. Starting from the intermediate **XXXIII**, introduction of various R^B and R^1 in the final **XXXVI** is achieved through the intermediate **XXXIV** and **XXXV**, respectively. The reaction conditions for each route are similar to those described in the previous scheme. If R^1 contains a protecting group, then an additional deprotection step takes place to produce **XX**.

deprotection step takes place to produce **XXXVI**.

[0267] In some embodiments, compounds described herein are prepared as described in Scheme D.

Scheme D




a) CS_2CO_3 , DMSO; b) TFA, DCM; c) *N*-benzyl-2-bromo-*N*-(2-bromoethyl)ethan-1-amine, KOH; d) $R^A B(OH)_2$, PdAMPhos, K_2CO_3 , Dioxane/ H_2O ; e) $Pd(OH)_2$, Pd/C, H_2 ; f) $R^B X$, DIEA, or $R^B X$, Pd(II); g) $R^1 NH_2$, LiHMDS; h) deprotection

[0268] Dichloro pyridazine or pyrazine is converted to the acetate **XXXVII** through formation of the malonate adduct and then removal of tert-butyloxycarbonyl group in an acidic condition. Starting from **XXXVII**, the intermediate **XL** is prepared by a similar manner as described in **Scheme B**. Due to chemical instability of the acid of **XL**, the amide bond is directly introduced by S_N2 reaction of the ester with $R^1 NH_2/LiHMDS$. If R^1 contains a protecting group, then an additional deprotection step takes place to produce **XLI**.

[0269] In some embodiments, compounds described herein are prepared as described in Scheme E.

Scheme E




a) $R^A B(OH)_2$, PdAMPhos, K_2CO_3 , Dioxane/ H_2O ; b) MsCl, TEA; c) NaCN, DMSO; d) *N*-benzyl-2-bromo-*N*-(2-bromoethyl)ethan-1-amine, KOH; e) KOH, EtOH; f) $R^1 NH_2$, HATU, TEA; g) $Pd(OH)_2$, Pd/C, H_2 ; h) $R^B X$, DIEA, or $R^B X$, Pd(II); i) deprotection

[0270] The cyanomethyl substituted pyrimidine **XLII** is obtained from a three-step sequence, Suzuki coupling, mesylation, and cyanation. Starting from **XLII**, the final compound **XLVI** is prepared by a similar manner as described in **Scheme C**. If R^1 contains a protecting group,

then an additional deprotection step takes place to produce **XLVI**.

[0271] In some embodiments, compounds described herein are prepared as described in Scheme F.

Scheme F



a) DPPA, TEA, tol, BnOH, 4Å MS; b) Pd(OH)₂, Pd/C, H₂; c) R¹NH₂, HATU, TEA; d) deprotection; e) triphosgene, tol; f) R¹OH, 4Å MS; g) R¹R³NH, TEA, toluene.

[0272] Curtius Rearrangement by the reaction of **XXV** with diphenylphosphoryl azide (DPPA) yields the isocyanate intermediate, which is subjected to *in situ* trapping of benzyl alcohol to produce **XLVII**. Deprotection to **XLVIII** and then HATU-mediated coupling reaction with R¹NH₂ yield the final **XLIX**. The isocyanate **L** derived from **XLVIII** undergoes S_N2 reaction with various nucleophiles such as alcohols (R¹OH) and amines (R¹R³NH) to yield carbamate (**LI**) or urea (**LII**) derivatives, respectively. If R¹ contains a protecting group in **XLIX**, **LI**, and **LII**, then an additional deprotection step is required for the final product.

[0273] In some embodiments, compounds described herein are synthesized as outlined in the Examples.

Certain Terminology

[0274] Unless otherwise stated, the following terms used in this application have the definitions given below. The use of the term "including" as well as other forms, such as "include", "includes," and "included," is not limiting. The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

[0275] As used herein, C₁-C_x includes C₁-C₂, C₁-C₃ . . . C₁-C_x. By way of example only, a group designated as "C₁-C₆" indicates that there are one to six carbon atoms in the moiety, i.e. groups containing 1 carbon atom, 2 carbon atoms, 3 carbon atoms or 4 carbon atoms. Thus, by way of example only, "C₁-C₄ alkyl" indicates that there are one to four carbon atoms in the

alkyl group, *i.e.*, the alkyl group is selected from among methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, and *t*-butyl.

[0276] An "alkyl" group refers to an aliphatic hydrocarbon group. The alkyl group is branched or straight chain. In some embodiments, the "alkyl" group has 1 to 10 carbon atoms, *i.e.* a C₁-C₁₀alkyl. Whenever it appears herein, a numerical range such as "1 to 10" refers to each integer in the given range; *e.g.*, "1 to 10 carbon atoms" means that the alkyl group consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 10 carbon atoms, although the present definition also covers the occurrence of the term "alkyl" where no numerical range is designated. In some embodiments, an alkyl is a C₁-C₆alkyl. In one aspect the alkyl is methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, or *t*-butyl. Typical alkyl groups include, but are in no way limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, tertiary butyl, pentyl, neopentyl, or hexyl.

[0277] An "alkylene" group refers to a divalent alkyl radical. Any of the above mentioned monovalent alkyl groups may be an alkylene by abstraction of a second hydrogen atom from the alkyl. In some embodiments, an alkylene is a C₁-C₆alkylene. In other embodiments, an alkylene is a C₁-C₄alkylene. Typical alkylene groups include, but are not limited to, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂-, and the like. In some embodiments, an alkylene is -CH₂-.

[0278] An "alkoxy" group refers to a (alkyl)O- group, where alkyl is as defined herein.

[0279] The term "alkylamine" refers to the -N(alkyl)_xH_y group, where x is 0 and y is 2, or where x is 1 and y is 1, or where x is 2 and y is 0.

[0280] An "hydroxyalkyl" refers to an alkyl in which one hydrogen atom is replaced by a hydroxyl. In some embodiments, a hydroxyalkyl is a C₁-C₄hydroxyalkyl. Typical hydroxyalkyl groups include, but are not limited to, -CH₂OH, -CH₂CH₂OH, -CH₂CH₂CH₂OH, -CH₂CH₂CH₂CH₂OH, and the like.

[0281] An "aminoalkyl" refers to an alkyl in which one hydrogen atom is replaced by an amino. In some embodiments, aminoalkyl is a C₁-C₄aminoalkyl. Typical aminoalkyl groups include, but are not limited to, -CH₂NH₂, -CH₂CH₂NH₂, -CH₂CH₂CH₂NH₂, -CH₂CH₂CH₂CH₂NH₂, and the like.

[0282] The term "alkenyl" refers to a type of alkyl group in which at least one carbon-carbon double bond is present. In one embodiment, an alkenyl group has the formula -C(R)=CR₂, wherein R refers to the remaining portions of the alkenyl group, which may be the same or different. In some embodiments, R is H or an alkyl. In some embodiments, an alkenyl is selected from ethenyl (*i.e.*, vinyl), propenyl (*i.e.*, allyl), butenyl, pentenyl, pentadienyl, and the like. Non-limiting examples of an alkenyl group include -CH=CH₂, -C(CH₃)=CH₂, -CH=CHCH₃, -

$\text{C}(\text{CH}_3)=\text{CHCH}_3$, and $-\text{CH}_2\text{CH}=\text{CH}_2$.

[0283] The term "alkynyl" refers to a type of alkyl group in which at least one carbon-carbon triple bond is present. In one embodiment, an alkenyl group has the formula $-\text{C}\equiv\text{C}-\text{R}$, wherein R refers to the remaining portions of the alkynyl group. In some embodiments, R is H or an alkyl. In some embodiments, an alkynyl is selected from ethynyl, propynyl, butynyl, pentynyl, hexynyl, and the like. Non-limiting examples of an alkynyl group include $-\text{C}\equiv\text{CH}$, $-\text{C}\equiv\text{CCH}_3$, $-\text{C}\equiv\text{CCH}_2\text{CH}_3$, $-\text{CH}_2\text{C}\equiv\text{CH}$.

[0284] The term "heteroalkyl" refers to an alkyl group in which one or more skeletal atoms of the alkyl are selected from an atom other than carbon, e.g., oxygen, nitrogen (e.g. $-\text{NH}-$, $-\text{N}(\text{alkyl})-$, sulfur, or combinations thereof. A heteroalkyl is attached to the rest of the molecule at a carbon atom of the heteroalkyl. In one aspect, a heteroalkyl is a $\text{C}_1\text{-C}_6$ heteroalkyl.

[0285] The term "aromatic" refers to a planar ring having a delocalized π -electron system containing $4n+2$ π electrons, where n is an integer. The term "aromatic" includes both carbocyclic aryl ("aryl", e.g., phenyl) and heterocyclic aryl (or "heteroaryl" or "heteroaromatic") groups (e.g., pyridine). The term includes monocyclic or fused-ring polycyclic (i.e., rings which share adjacent pairs of carbon atoms) groups.

[0286] The term "carbocyclic" or "carbocycle" refers to a ring or ring system where the atoms forming the backbone of the ring are all carbon atoms. The term thus distinguishes carbocyclic from "heterocyclic" rings or "heterocycles" in which the ring backbone contains at least one atom which is different from carbon. In some embodiments, a carbocycle is a monocyclic carbocycle or a bicyclic carbocycle. In some embodiments, a carbocycle is a monocyclic carbocycle. Carbocycles are non-aromatic or aromatic. Non-aromatic carbocycles are saturated or partially unsaturated. In some embodiments, a carbocycle is a bicyclic carbocycle. In some embodiments, at least one of the two rings of a bicyclic carbocycle is aromatic. In some embodiments, both rings of a bicyclic carbocycle are aromatic. Carbocycles include aryls and cycloalkyls.

[0287] As used herein, the term "aryl" refers to an aromatic ring wherein each of the atoms forming the ring is a carbon atom. In one aspect, aryl is phenyl or a naphthyl. In some embodiments, an aryl is a phenyl. In some embodiments, an aryl is a phenyl, naphthyl, indanyl, indenyl, or tetrahydronaphthyl. In some embodiments, an aryl is a $\text{C}_6\text{-C}_{10}$ aryl. Depending on the structure, an aryl group is a monoradical or a diradical (i.e., an arylene group).

[0288] The term "cycloalkyl" refers to a monocyclic, bicyclic or polycyclic aliphatic, non-aromatic radical, wherein each of the atoms forming the ring (i.e. skeletal atoms) is a carbon atom. In some embodiments, cycloalkyls are spirocyclic or bridged compounds. In some embodiments, cycloalkyls are optionally fused with an aromatic ring, and the point of attachment is at a carbon that is not an aromatic ring carbon atom. Cycloalkyl groups include groups having from 3 to 10 ring atoms. In some embodiments, cycloalkyl groups are selected

from among cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptyl, cyclooctyl, spiro[2.2]pentyl, norbornyl, norbornenyl, bicycle[1.1.1]pentyl, adamantyl, norbornyl, norbornenyl, decalynyl, or 7,7-dimethyl-bicyclo[2.2.1]heptanyl. In some embodiments, a cycloalkyl is a C₃-C₆cycloalkyl. In some embodiments, a cycloalkyl is a monocyclic cycloalkyl. Monocyclic cycloalkyls include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Polycyclic cycloalkyls include, for example, adamantyl, norbornyl (*i.e.*, bicyclo[2.2.1]heptanyl), norbornenyl, decalynyl, 7,7-dimethyl-bicyclo[2.2.1]heptanyl, and the like.

[0289] The term "halo" or, alternatively, "halogen" or "halide" means fluoro, chloro, bromo or iodo. In some embodiments, halo is fluoro, chloro, or bromo.

[0290] The term "fluoroalkyl" refers to an alkyl in which one or more hydrogen atoms are replaced by a fluorine atom. In one aspect, a fluoroalkyl is a C₁-C₆fluoroalkyl.

[0291] The term "heterocycle" or "heterocyclic" refers to heteroaromatic rings (also known as heteroaryls) and heterocycloalkyl rings containing one to four heteroatoms in the ring(s), where each heteroatom in the ring(s) is selected from O, S and N, wherein each heterocyclic group has from 3 to 10 atoms in its ring system, and with the proviso that any ring does not contain two adjacent O or S atoms. Non-aromatic heterocyclic groups (also known as heterocycloalkyls) include rings having 3 to 10 atoms in its ring system and aromatic heterocyclic groups include rings having 5 to 10 atoms in its ring system. The heterocyclic groups include benzo-fused ring systems. Examples of non-aromatic heterocyclic groups are pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothienyl, oxazolidinonyl, tetrahydropyranyl, dihydropyranyl, tetrahydrothiopyranyl, piperidinyl, morpholinyl, thiomorpholinyl, thioxanyl, piperazinyl, aziridinyl, azetidiny, oxetanyl, thietanyl, homopiperidinyl, oxepanyl, thiepanyl, oxazepinyl, diazepinyl, thiazepinyl, 1,2,3,6-tetrahydropyridinyl, pyrrolin-2-yl, pyrrolin-3-yl, indolinyl, 2H-pyranyl, 4H-pyranyl, dioxanyl, 1,3-dioxolanyl, pyrazolinyl, dithianyl, dithiolanyl, dihydropyranyl, dihydrothienyl, dihydrofuranyl, pyrazolidinyl, imidazoliny, imidazolidinyl, 3-azabicyclo[3.1.0]hexanyl, 3-azabicyclo[4.1.0]heptanyl, 3H-indolyl, indolin-2-onyl, isoindolin-1-onyl, isoindoline-1,3-dionyl, 3,4-dihydroisoquinolin-1(2H)-onyl, 3,4-dihydroquinolin-2(1H)-onyl, isoindoline-1,3-dithionyl, benzo[d]oxazol-2(3H)-onyl, 1H-benzo[d]imidazol-2(3H)-onyl, benzo[d]thiazol-2(3H)-onyl, and quinoliziny. Examples of aromatic heterocyclic groups are pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxaliny, naphthyridinyl, and furopyridinyl. The foregoing groups are either C-attached (or C-linked) or *N*-attached where such is possible. For instance, a group derived from pyrrole includes both pyrrol-1-yl (*N*-attached) or pyrrol-3-yl (C-attached). Further, a group derived from imidazole includes imidazol-1-yl or imidazol-3-yl (both *N*-attached) or imidazol-2-yl, imidazol-4-yl or imidazol-5-yl (all C-attached). The heterocyclic groups include benzo-fused ring systems. Non-aromatic heterocycles are optionally substituted with one or two oxo (=O) moieties, such as

pyrrolidin-2-one. In some embodiments, at least one of the two rings of a bicyclic heterocycle is aromatic. In some embodiments, both rings of a bicyclic heterocycle are aromatic.

[0292] The terms "heteroaryl" or, alternatively, "heteroaromatic" refers to an aryl group that includes one or more ring heteroatoms selected from nitrogen, oxygen and sulfur. Illustrative examples of heteroaryl groups include monocyclic heteroaryls and bicyclic heteroaryls. Monocyclic heteroaryls include pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, pyridazinyl, triazinyl, oxadiazolyl, thiadiazolyl, and furazanyl. Monocyclic heteroaryls include indolizine, indole, benzofuran, benzothiophene, indazole, benzimidazole, purine, quinolizine, quinoline, isoquinoline, cinnoline, phthalazine, quinazoline, quinoxaline, 1,8-naphthyridine, and pteridine. In some embodiments, a heteroaryl contains 0-4 N atoms in the ring. In some embodiments, a heteroaryl contains 1-4 N atoms in the ring. In some embodiments, a heteroaryl contains 0-4 N atoms, 0-1 O atoms, and 0-1 S atoms in the ring. In some embodiments, a heteroaryl contains 1-4 N atoms, 0-1 O atoms, and 0-1 S atoms in the ring. In some embodiments, heteroaryl is a C₁-C₉heteroaryl. In some embodiments, monocyclic heteroaryl is a C₁-C₅heteroaryl. In some embodiments, monocyclic heteroaryl is a 5-membered or 6-membered heteroaryl. In some embodiments, bicyclic heteroaryl is a C₆-C₉heteroaryl.

[0293] A "heterocycloalkyl" group refers to a cycloalkyl group that includes at least one heteroatom selected from nitrogen, oxygen and sulfur. In some embodiments, a heterocycloalkyl is fused with an aryl or heteroaryl. In some embodiments, the heterocycloalkyl is oxazolidinonyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, piperidin-2-onyl, pyrrolidine-2,5-dithionyl, pyrrolidine-2,5-dionyl, pyrrolidinonyl, imidazolidinyl, imidazolidin-2-onyl, or thiazolidin-2-onyl. The term heterocycloalkyl also includes all ring forms of the carbohydrates, including but not limited to the monosaccharides, the disaccharides and the oligosaccharides. In one aspect, a heterocycloalkyl is a C₂-C₁₀heterocycloalkyl. In another aspect, a heterocycloalkyl is a C₄-C₁₀heterocycloalkyl. In some embodiments, a heterocycloalkyl contains 0-2 N atoms in the ring. In some embodiments, a heterocycloalkyl contains 0-2 N atoms, 0-2 O atoms and 0-1 S atoms in the ring.

[0294] The term "bond" or "single bond" refers to a chemical bond between two atoms, or two moieties when the atoms joined by the bond are considered to be part of larger substructure. In one aspect, when a group described herein is a bond, the referenced group is absent thereby allowing a bond to be formed between the remaining identified groups.

[0295] The term "moiety" refers to a specific segment or functional group of a molecule. Chemical moieties are often recognized chemical entities embedded in or appended to a molecule.

[0296] The term "optionally substituted" or "substituted" means that the referenced group is optionally substituted with one or more additional group(s) individually and independently

selected from halogen, -CN, -NH₂, -NH(alkyl), -N(alkyl)₂, -OH, -CO₂H, -CO₂alkyl, -C(=O)NH₂, -C(=O)NH(alkyl), -C(=O)N(alkyl)₂, -S(=O)₂NH₂, -S(=O)₂NH(alkyl), -S(=O)₂N(alkyl)₂, alkyl, cycloalkyl, fluoroalkyl, heteroalkyl, alkoxy, fluoroalkoxy, heterocycloalkyl, aryl, heteroaryl, aryloxy, alkylthio, arylthio, alkylsulfoxide, aryl sulfoxide, alkylsulfone, and arylsulfone. In some other embodiments, optional substituents are independently selected from halogen, -CN, -NH₂, -NH(CH₃), -N(CH₃)₂, -OH, -CO₂H, -CO₂(C₁-C₄alkyl), -C(=O)NH₂, -C(=O)NH(C₁-C₄alkyl), -C(=O)N(C₁-C₄alkyl)₂, -S(=O)₂NH₂, -S(=O)₂NH(C₁-C₄alkyl), -S(=O)₂N(C₁-C₄alkyl)₂, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄fluoroalkyl, C₁-C₄heteroalkyl, C₁-C₄alkoxy, C₁-C₄fluoroalkoxy, -SC₁-C₄alkyl, -S(=O)C₁-C₄alkyl, and -S(=O)₂C₁-C₄alkyl. In some other embodiments, optional substituents are independently selected from halogen, -CN, -NH₂, -NH(CH₃), -N(CH₃)₂, -OH, -CO₂H, -CO₂(C₁-C₄alkyl), -C(=O)NH₂, -C(=O)NH(C₁-C₄alkyl), -C(=O)N(C₁-C₄alkyl)₂, -S(=O)₂NH₂, -S(=O)₂NH(C₁-C₄alkyl), -S(=O)₂N(C₁-C₄alkyl)₂, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₂-C₆heterocycloalkyl, C₁-C₄fluoroalkyl, C₁-C₄heteroalkyl, C₁-C₄alkoxy, C₁-C₄fluoroalkoxy, -SC₁-C₄alkyl, -S(=O)C₁-C₄alkyl, and -S(=O)₂C₁-C₄alkyl. In some embodiments, optional substituents are independently selected from halogen, -CN, -NH₂, -OH, -NH(CH₃), -N(CH₃)₂, -CH₃, -CH₂CH₃, -CF₃, -OCH₃, and -OCF₃. In some embodiments, substituted groups are substituted with one or two of the preceding groups. In some embodiments, an optional substituent on an aliphatic carbon atom (acyclic or cyclic) includes oxo (=O).

[0297] The term "acceptable" with respect to a formulation, composition or ingredient, as used herein, means having no persistent detrimental effect on the general health of the subject being treated.

[0298] The term "modulate" as used herein, means to interact with a target either directly or indirectly so as to alter the activity of the target, including, by way of example only, to enhance the activity of the target, to inhibit the activity of the target, to limit the activity of the target, or to extend the activity of the target.

[0299] The term "modulator" as used herein, refers to a molecule that interacts with a target either directly or indirectly. The interactions include, but are not limited to, the interactions of an agonist, partial agonist, an inverse agonist, antagonist, degrader, or combinations thereof. In some embodiments, a modulator is an agonist.

[0300] The terms "administer," "administering," "administration," and the like, as used herein, refer to the methods that may be used to enable delivery of compounds or compositions to the desired site of biological action. These methods include, but are not limited to oral routes, intraduodenal routes, parenteral injection (including intravenous, subcutaneous, intraperitoneal, intramuscular, intravascular or infusion), topical and rectal administration. Those of skill in the art are familiar with administration techniques that can be employed with the compounds and methods described herein. In some embodiments, the compounds and compositions described herein are administered orally.

[0301] The terms "co-administration" or the like, as used herein, are meant to encompass administration of the selected therapeutic agents to a single patient, and are intended to include treatment regimens in which the agents are administered by the same or different route of administration or at the same or different time.

[0302] The terms "effective amount" or "therapeutically effective amount," as used herein, refer to a sufficient amount of an agent or a compound being administered, which will relieve to some extent one or more of the symptoms of the disease or condition being treated. The result includes reduction and/or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. For example, an "effective amount" for therapeutic uses is the amount of the composition comprising a compound as disclosed herein required to provide a clinically significant decrease in disease symptoms. An appropriate "effective" amount in any individual case is optionally determined using techniques, such as a dose escalation study.

[0303] The terms "enhance" or "enhancing," as used herein, means to increase or prolong either in potency or duration a desired effect. Thus, in regard to enhancing the effect of therapeutic agents, the term "enhancing" refers to the ability to increase or prolong, either in potency or duration, the effect of other therapeutic agents on a system. An "enhancing-effective amount," as used herein, refers to an amount adequate to enhance the effect of another therapeutic agent in a desired system.

[0304] The term "pharmaceutical combination" as used herein, means a product that results from the mixing or combining of more than one active ingredient and includes both fixed and non-fixed combinations of the active ingredients. The term "fixed combination" means that the active ingredients, e.g. a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and a co-agent, are both administered to a patient simultaneously in the form of a single entity or dosage. The term "non-fixed combination" means that the active ingredients, e.g. a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and a co-agent, are administered to a patient as separate entities either simultaneously, concurrently or sequentially with no specific intervening time limits, wherein such administration provides effective levels of the two compounds in the body of the patient. The latter also applies to cocktail therapy, e.g. the administration of three or more active ingredients.

[0305] The terms "article of manufacture" and "kit" are used as synonyms.

[0306] The term "subject" or "patient" encompasses mammals. Examples of mammals include, but are not limited to, any member of the Mammalian class: humans, non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. In one aspect, the mammal is a human.

[0307] The terms "treat," "treating" or "treatment," as used herein, include alleviating, abating

or ameliorating at least one symptom of a disease or condition, preventing additional symptoms, inhibiting the disease or condition, e.g., arresting the development of the disease or condition, relieving the disease or condition, causing regression of the disease or condition, relieving a condition caused by the disease or condition, or stopping the symptoms of the disease or condition either prophylactically and/or therapeutically.

Pharmaceutical compositions

[0308] In some embodiments, the compounds described herein are formulated into pharmaceutical compositions. Pharmaceutical compositions are formulated in a conventional manner using one or more pharmaceutically acceptable inactive ingredients that facilitate processing of the active compounds into preparations that are used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. A summary of pharmaceutical compositions described herein is found, for example, in Remington: The Science and Practice of Pharmacy, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Pharmaceutical Dosage Forms and Drug Delivery Systems, Seventh Ed. (Lippincott Williams & Wilkins 1999).

[0309] In some embodiments, the compounds described herein are administered either alone or in combination with pharmaceutically acceptable carriers, excipients or diluents, in a pharmaceutical composition. Administration of the compounds and compositions described herein can be effected by any method that enables delivery of the compounds to the site of action. These methods include, though are not limited to delivery via enteral routes (including oral, gastric or duodenal feeding tube, rectal suppository and rectal enema), parenteral routes (injection or infusion, including intraarterial, intracardiac, intradermal, intraduodenal, intramedullary, intramuscular, intraosseous, intraperitoneal, intrathecal, intravascular, intravenous, intravitreal, epidural and subcutaneous), inhalational, transdermal, transmucosal, sublingual, buccal and topical (including epicutaneous, dermal, enema, eye drops, ear drops, intranasal, vaginal) administration, although the most suitable route may depend upon for example the condition and disorder of the recipient. By way of example only, compounds described herein can be administered locally to the area in need of treatment, by for example, local infusion during surgery, topical application such as creams or ointments, injection, catheter, or implant. The administration can also be by direct injection at the site of a diseased tissue or organ.

[0310] In some embodiments, pharmaceutical compositions suitable for oral administration are presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. In some embodiments, the active ingredient is presented as a bolus, electuary or paste.

[0311] Pharmaceutical compositions which can be used orally include tablets, push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. Tablets may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with binders, inert diluents, or lubricating, surface active or dispersing agents. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. In some embodiments, the tablets are coated or scored and are formulated so as to provide slow or controlled release of the active ingredient therein. All formulations for oral administration should be in dosages suitable for such administration. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In some embodiments, stabilizers are added. Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or Dragee coatings for identification or to characterize different combinations of active compound doses.

[0312] In some embodiments, pharmaceutical compositions are formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. The compositions may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in powder form or in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, saline or sterile pyrogen-free water, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

[0313] Pharmaceutical compositions for parenteral administration include aqueous and non-aqueous (oily) sterile injection solutions of the active compounds which may contain antioxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated

solutions.

[0314] Pharmaceutical compositions may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example, as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[0315] For buccal or sublingual administration, the compositions may take the form of tablets, lozenges, pastilles, or gels formulated in conventional manner. Such compositions may comprise the active ingredient in a flavored basis such as sucrose and acacia or tragacanth.

[0316] Pharmaceutical compositions may be administered topically, that is by non-systemic administration. This includes the application of a compound of the present invention externally to the epidermis or the buccal cavity and the instillation of such a compound into the ear, eye and nose, such that the compound does not significantly enter the blood stream. In contrast, systemic administration refers to oral, intravenous, intraperitoneal and intramuscular administration.

[0317] Pharmaceutical compositions suitable for topical administration include liquid or semiliquid preparations suitable for penetration through the skin to the site of inflammation such as gels, liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose. The active ingredient may comprise, for topical administration, from 0.001% to 10% w/w, for instance from 1% to 2% by weight of the formulation.

[0318] Pharmaceutical compositions for administration by inhalation are conveniently delivered from an insufflator, nebulizer pressurized packs or other convenient means of delivering an aerosol spray. Pressurized packs may comprise a suitable propellant such as dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Alternatively, for administration by inhalation or insufflation, pharmaceutical preparations may take the form of a dry powder composition, for example a powder mix of the compound and a suitable powder base such as lactose or starch. The powder composition may be presented in unit dosage form, in for example, capsules, cartridges, gelatin or blister packs from which the powder may be administered with the aid of an inhalator or insufflator.

[0319] It should be understood that in addition to the ingredients particularly mentioned above, the compounds and compositions described herein may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents.

Methods of Dosing and Treatment Regimens

[0320] In one embodiment, the compounds of Formula (I), or a pharmaceutically acceptable salt thereof, are used in the preparation of medicaments for use in the treatment of diseases or conditions in a mammal that would benefit from modulation of melanocortin receptor activity. Compounds for use in methods for treating any of the diseases or conditions described herein in a mammal in need of such treatment, involves administration of pharmaceutical compositions that include at least one compound of Formula (I) or a pharmaceutically acceptable salt thereof, in therapeutically effective amounts to said mammal.

[0321] In certain embodiments, the compositions containing the compound(s) described herein are administered for prophylactic and/or therapeutic treatments. In certain therapeutic applications, the compositions are administered to a patient already suffering from a disease or condition, in an amount sufficient to cure or at least partially arrest at least one of the symptoms of the disease or condition. Amounts effective for this use depend on the severity and course of the disease or condition, previous therapy, the patient's health status, weight, and response to the drugs, and the judgment of the treating physician. Therapeutically effective amounts are optionally determined by methods including, but not limited to, a dose escalation and/or dose ranging clinical trial.

[0322] In prophylactic applications, compositions containing the compounds described herein are administered to a patient susceptible to or otherwise at risk of a particular disease, disorder or condition. Such an amount is defined to be a "prophylactically effective amount or dose." In this use, the precise amounts also depend on the patient's state of health, weight, and the like. When used in patients, effective amounts for this use will depend on the severity and course of the disease, disorder or condition, previous therapy, the patient's health status and response to the drugs, and the judgment of the treating physician. In one aspect, prophylactic treatments include administering to a mammal, who previously experienced at least one symptom of the disease being treated and is currently in remission, a pharmaceutical composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof, in order to prevent a return of the symptoms of the disease or condition.

[0323] In certain embodiments wherein the patient's condition does not improve, upon the doctor's discretion the administration of the compounds are administered chronically, that is, for an extended period of time, including throughout the duration of the patient's life in order to ameliorate or otherwise control or limit the symptoms of the patient's disease or condition.

[0324] Once improvement of the patient's conditions has occurred, a maintenance dose is administered if necessary. Subsequently, in specific embodiments, the dosage or the frequency of administration, or both, is reduced, as a function of the symptoms, to a level at which the improved disease, disorder or condition is retained. In certain embodiments, however, the patient requires intermittent treatment on a long-term basis upon any recurrence of symptoms.

[0325] The amount of a given agent that corresponds to such an amount varies depending upon factors such as the particular compound, disease condition and its severity, the identity (e.g., weight, sex) of the subject or host in need of treatment, but nevertheless is determined according to the particular circumstances surrounding the case, including, e.g., the specific agent being administered, the route of administration, the condition being treated, and the subject or host being treated.

[0326] In general, however, doses employed for adult human treatment are typically in the range of 0.01 mg-2000 mg per day. In one embodiment, the desired dose is conveniently presented in a single dose or in divided doses administered simultaneously or at appropriate intervals, for example as two, three, four or more sub-doses per day.

[0327] In one embodiment, the daily dosages appropriate for the compound of Formula (I), or a pharmaceutically acceptable salt thereof, described herein are from about 0.01 to about 50 mg/kg per body weight. In some embodiments, the daily dosage or the amount of active in the dosage form are lower or higher than the ranges indicated herein, based on a number of variables in regard to an individual treatment regime. In various embodiments, the daily and unit dosages are altered depending on a number of variables including, but not limited to, the activity of the compound used, the disease or condition to be treated, the mode of administration, the requirements of the individual subject, the severity of the disease or condition being treated, and the judgment of the practitioner.

[0328] Toxicity and therapeutic efficacy of such therapeutic regimens are determined by standard pharmaceutical procedures in cell cultures or experimental animals, including, but not limited to, the determination of the LD₅₀ and the ED₅₀. The dose ratio between the toxic and therapeutic effects is the therapeutic index and it is expressed as the ratio between LD₅₀ and ED₅₀. In certain embodiments, the data obtained from cell culture assays and animal studies are used in formulating the therapeutically effective daily dosage range and/or the therapeutically effective unit dosage amount for use in mammals, including humans. In some embodiments, the daily dosage amount of the compounds described herein lies within a range of circulating concentrations that include the ED₅₀ with minimal toxicity. In certain embodiments, the daily dosage range and/or the unit dosage amount varies within this range depending upon the dosage form employed and the route of administration utilized.

[0329] In any of the aforementioned aspects are further embodiments in which the effective amount of the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is: (a) systemically administered to the mammal; and/or (b) administered orally to the mammal; and/or (c) intravenously administered to the mammal; and/or (d) administered by injection to the mammal; and/or (e) administered topically to the mammal; and/or (f) administered non-systemically or locally to the mammal.

[0330] In any of the aforementioned aspects are further embodiments comprising single administrations of the effective amount of the compound, including further embodiments in

which (i) the compound is administered once a day; or (ii) the compound is administered to the mammal multiple times over the span of one day.

[0331] In any of the aforementioned aspects are further embodiments comprising multiple administrations of the effective amount of the compound, including further embodiments in which (i) the compound is administered continuously or intermittently: as in a single dose; (ii) the time between multiple administrations is every 6 hours; (iii) the compound is administered to the mammal every 8 hours; (iv) the compound is administered to the mammal every 12 hours; (v) the compound is administered to the mammal every 24 hours. In further or alternative embodiments, the method comprises a drug holiday, wherein the administration of the compound is temporarily suspended or the dose of the compound being administered is temporarily reduced; at the end of the drug holiday, dosing of the compound is resumed. In one embodiment, the length of the drug holiday varies from 2 days to 1 year.

Combination Treatments

[0332] In certain instances, it is appropriate to administer at least one compound of Formula (I), or a pharmaceutically acceptable salt thereof, in combination with one or more other therapeutic agents.

[0333] In one embodiment, the therapeutic effectiveness of one of the compounds described herein is enhanced by administration of an adjuvant (*i.e.*, by itself the adjuvant has minimal therapeutic benefit, but in combination with another therapeutic agent, the overall therapeutic benefit to the patient is enhanced). Or, in some embodiments, the benefit experienced by a patient is increased by administering one of the compounds described herein with another agent (which also includes a therapeutic regimen) that also has therapeutic benefit.

[0334] In one specific embodiment, a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is co-administered with a second therapeutic agent, wherein the compound of Formula (I), or a pharmaceutically acceptable salt thereof, and the second therapeutic agent modulate different aspects of the disease, disorder or condition being treated, thereby providing a greater overall benefit than administration of either therapeutic agent alone.

[0335] In any case, regardless of the disease, disorder or condition being treated, the overall benefit experienced by the patient is simply be additive of the two therapeutic agents or the patient experiences a synergistic benefit.

[0336] For combination therapies described herein, dosages of the co-administered compounds vary depending on the type of co-drug employed, on the specific drug employed, on the disease or condition being treated and so forth. In additional embodiments, when co-administered with one or more other therapeutic agents, the compound provided herein is administered either simultaneously with the one or more other therapeutic agents, or

sequentially.

[0337] In combination therapies, the multiple therapeutic agents (one of which is one of the compounds described herein) are administered in any order or even simultaneously. If administration is simultaneous, the multiple therapeutic agents are, by way of example only, provided in a single, unified form, or in multiple forms (e.g., as a single pill or as two separate pills).

[0338] The compounds of Formula (I), or a pharmaceutically acceptable salt thereof, as well as combination therapies, are administered before, during or after the occurrence of a disease or condition, and the timing of administering the composition containing a compound varies. Thus, in one embodiment, the compounds described herein are used as a prophylactic and are administered continuously to subjects with a propensity to develop conditions or diseases in order to prevent the occurrence of the disease or condition. In another embodiment, the compounds and compositions are administered to a subject during or as soon as possible after the onset of the symptoms. In specific embodiments, a compound described herein is administered as soon as is practicable after the onset of a disease or condition is detected or suspected, and for a length of time necessary for the treatment of the disease. In some embodiments, the length required for treatment varies, and the treatment length is adjusted to suit the specific needs of each subject.

Abbreviations:

[0339]

DIEA: N,N-diisopropylethylamine;

DMSO: dimethyl sulfoxide;

CuI: copper(I) iodide;

TBAF: tetra-n-butylammonium fluoride;

P(t-Bu)₃: tri-tert-butylphosphine;

HBF₄: tetrafluoroboric acid;

DBU: 1,8-diazabicyclo[5.4.0]undec-7-ene;

Prep-HPLC: preparative high performance liquid chromatography;

TFA: trifluoroacetic acid;

CH₃CN: acetonitrile;

MeOD: deuterated methanol;

CDCl_3 : deuterated chloroform;

DME: 1,2-dimethoxyethane;

H_2O : water;

KOAc: potassium acetate;

NaOAc: sodium acetate;

Cs_2CO_3 : cesium carbonate;

P-TsOH: p-toluenesulfonic acid;

NaNO_2 : sodium nitrate;

THF: tetrahydrofuran;

NBS: N-bromosuccinimide;

4Å MS: 4Å molecular sieves;

DPPA: diphenyl phosphoryl azide;

Br_2 : bromine;

AgF: silver fluoride;

LiAlH_4 : lithium aluminium hydride;

LiHMDS: lithium bis(trimethylsilyl)amide;

IBX: 2-iodoxybenzoic acid;

CDI: 1,1'-carbonyldiimidazole;

TEA: trimethylamine;

HOBT: hydroxybenzotriazole;

EDCI: 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide;

$\text{Pd}(\text{PPh}_3)_4$: tetrakis(triphenylphosphine)palladium(0);

$\text{Pd}(\text{OH})_2$: palladium hydroxide;

$\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$: bis(triphenylphosphine)palladium(II) dichloride;

$\text{Pd}(\text{dppf})\text{Cl}_2$: [1,1' -Bis(diphenylphosphino)ferrocene]dichloropalladium(II);

PdAMphos or Pd (amphos)Cl₂ or : bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II);

Pd(DTBPF)Cl₂: [1,1'-bis(di-*tert*-butylphosphino)ferrocene]dichloropalladium(II);

Pd₂(dba)₃·CHCl₃: tris(dibenzylideneacetone)dipalladium(0)-chloroform adduct;

XPhos: 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl;

RuPhos-Pd-G3: dicyclohexyl-[2-[2,6-di(propan-2-yloxy)phenyl]phenyl]phosphane methanesulfonic acid palladium 2-phenylaniline;

XPhos-Pd-G2: chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II);

rt: room temperature;

h: hour or hours;

Cpd: compound.


EXAMPLES

[0340] The following examples are provided for illustrative purposes only and not to limit the scope of the claims provided herein.

Synthesis of Compounds

Example 1: N-(2-aminoethyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxamide (cpd 1-3)

[0341]



[0342] Step 1-1, preparation of methyl 4-(4-bromophenyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate: Methyl 4-(4-bromophenyl)piperidine-4-

carboxylate hydrochloride (500 mg, 1.49 mmol), 2-fluoro-5-(trifluoromethyl)benzonitrile (706 mg, 3.74 mmol), DIEA (772 mg, 5.98 mmol), and DMSO (5 mL) were charged into a heavy-wall tube. The tube was sealed, and then the resulting solution was stirred at 125 °C for 16 h and cooled to rt. The reaction was directly purified by reverse phase C18 column chromatography to afford the title compound (600 mg, 86%). LCMS (M+H)⁺ = 467.3.

[0343] Step 1-2. preparation of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxylate: Methyl 4-(4-bromophenyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate (480 mg, 1.03 mmol), (2-ethoxypyridin-3-yl)boronic acid (343 mg, 2.05 mmol), PdAMphos (73 mg, 0.103 mmol), potassium carbonate (284 mg, 2.05 mmol), and dioxane/H₂O (5 mL/0.5 mL) were charged into a heavy-wall tube. The resulting mixture was degassed with N₂ for 5 min, sealed, and stirred at 80 °C for 1 h. The reaction was cooled to rt, and concentrated. The residue was directly purified by reverse phase C18 column chromatography to afford the title compound (450 mg, 86%). LCMS (M+H)⁺ = 510.3.

[0344] Step 1-3. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxylic acid: To a suspension of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxylate (450 mg, 0.883 mmol) in EtOH (8 mL) was added 2N-lithium hydroxide (4 mL, 8.83 mmol) at rt. The resulting mixture was stirred at 85 °C for 3 h and cooled to rt. After removal of the volatile solvent, the aqueous layer was filtered to remove inorganic solid. The aqueous filtrate was acidified to ~pH 4 with 1N-HCl to form ppts. The solid ppts were collected, washed with H₂O, and dried to afford the title compound (270 mg, 62%). LCMS (M+H)⁺ = 496.1.

[0345] Step 1-4. preparation of tert-butyl N-[2-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}formamido)ethyl]carbamate: To a solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxylic acid (30 mg, 0.061 mmol) in ACN (1 mL) was added HATU (25 mg, 0.067 mmol) and TEA (12 mg, 0.12 mmol) at rt. After stirring for 5 min at rt, the reaction was treated with tert-butyl (2-aminoethyl)carbamate (12 mg, 0.073 mmol). The resulting solution was stirred at rt for 30 min, and then directly purified by reverse phase C18 column chromatography to afford the title compound (30 mg, 78%). LCMS (M+H)⁺ = 638.1.


[0346] Step 1-5. preparation of N-(2-aminoethyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-4-carboxamide: To a solution of tert-butyl N-[2-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}formamido)ethyl]carbamate (30 mg, 0.047 mmol) in DCM (0.4 mL) was added TFA (0.1 mL) at rt. The reaction was stirred at rt for 1 h and concentrated under vacuum. The residue was purified by reverse phase C18 column chromatography to afford the title compound (18.5 mg, 73%). LCMS (M+H)⁺ = 538.4.

[0347] The following compounds were prepared similarly to **Example 1** with appropriate substituting reagents and substrates at different steps. Some examples do not require deprotection in the final step.

Compound no.	MS (M+H) ⁺
1-1	578.3
1-2	564.4
1-4	552.3
1-5	552.3
1-6	552.2
1-7	552.4
1-8	566.3
Compound no.	MS (M+H) ⁺
1-23	579.4
1-29	612.3
1-75	596.4
1-159	583.6
1-164	567.4
1-165	587.2

Example 2: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-22)

[0348]



[0349] Step 2-1. preparation of 2-(5-chloropyridin-2-yl)acetonitrile: Into a 250-mL round-bottom flask, were placed 5-chloro-2-(chloromethyl)pyridine (8.0 g, 49 mmol), potassium cyanide (4.5 g, 69 mmol), EtOH (80 mL) and H₂O (60 mL). The resulting mixture was stirred at 100 °C for 1 h and cooled to room temperature. The reaction was quenched with water (200 mL) and then extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was

applied onto a silica gel column with ethyl acetate/petroleum ether (1:1) to afford the title compound (4.3 g, 57%) as a yellow oil. LCMS (M+H)⁺ = 153.2.

[0350] Step 2-2, preparation of 1-benzyl-4-(5-chloropyridin-2-yl)piperidine-4-carbonitrile: To a solution of 2-(5-chloropyridin-2-yl)acetonitrile (2.0 g, 13 mmol) and N-benzyl-2-bromo-N-(2-bromoethyl)ethan-1-amine (5.0 g, 16 mmol) in DMSO (20 mL) was added KOH (2.0 g, 36 mmol) at rt. The resulting mixture was stirred at 30 °C for 1 h and then quenched with water (50 mL). The resulting solution was extracted with ethyl acetate (3x).

The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1). This resulted in the title compound (2.0 g, 49%) as a yellow oil. LCMS (M+H)⁺ = 312.0.

[0351] Step 2-3, preparation of 1-benzyl-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}piperidine-4-carbonitrile: Into a 50 mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, were placed 1-benzyl-4-(5-chloropyridin-2-yl)piperidine-4-carbonitrile (1.0 g, 3.2 mmol), (2-ethoxypyridin-3-yl)boronic acid (1.0 g, 6.0 mmol), Pd(dppf)Cl₂ (0.3 g, 0.4 mmol), K₂CO₃ (1.2 g, 8.7 mmol), and dioxane/H₂O (10 mL/1 mL). The resulting solution was stirred at 80 °C for 1 h under N₂ and cooled to room temperature. The reaction was quenched with water (50 mL) and extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (30:70) to afford the title compound (1.0 g, 78%) as a yellow oil. LCMS (M+H)⁺ = 399.1.

[0352] Step 2-4, preparation of 1-benzyl-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}piperidine-4-carboxylic acid: Into a 50-mL round-bottom flask, were placed 1-benzyl-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}piperidine-4-carbonitrile (1.0 g, 2.5 mmol), KOH (1.0 g, 17.8 mmol), EtOH (10 mL) and H₂O (10 mL). The resulting solution was stirred at 100 °C for 30 h, cooled to rt, and concentrated under vacuum. The residue was diluted with H₂O, and then pH of the solution was adjusted to ~4 with 4M-HCl. The solid ppts were collected, washed with water, and dried to afford the title compound as a white solid (800 mg, 80%). LCMS (M+H)⁺ = 418.3.

[0353] Step 2-5, preparation of tert-butyl (3R)-3-(1-benzyl-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}piperidine-4-amido)pyrrolidine-1-carboxylate: To a solution of 1-benzyl-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}piperidine-4-carboxylic acid (230 mg, 0.55 mmol) in DMF (4 mL) was added HATU (251 mg, 0.66 mmol) and DIEA (427 mg, 3.3 mmol) at rt. After stirring at rt for 5 min, the reaction was treated with tert-butyl (R)-3-aminopyrrolidine-1-carboxylate (123 mg, 0.66 mmol). The resulting solution was stirred at rt for 2 h and directly purified by Prep-HPLC to afford the title compound (230 mg, 71%) as a yellow solid. LCMS (M+H)⁺ = 586.3.

[0354] Step 2-6, preparation of tert-butyl (3R)-3-(4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}piperidine-4-amido)pyrrolidine-1-carboxylate: Into a 50-mL round-bottom flask, were placed tert-butyl

(3R)-3-(1-benzyl-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}piperidine-4-amido)pyrrolidine-1-carboxylate (230 mg, 0.39 mmol), MeOH (10 mL), wet 10%-Pd/C (30 mg), and Pd(OH)₂ (30 mg, 0.21 mmol). The reaction flask was evacuated and flushed three times with nitrogen, followed by flushing with hydrogen. The mixture was stirred at rt for 2 h under hydrogen (balloon). The reaction was diluted with MeOH, and filtered through a pad of Celite. The filtrate was concentrated under vacuum to afford the title compound (180 mg, 92%) as a yellow oil. LCMS (M+H)⁺ = 496.4.


[0355] Step 2-7, preparation of tert-butyl (3R)-3-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}piperidine-4-amido}pyrrolidine-1-carboxylate: A solution of tert-butyl (3R)-3-(4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}piperidine-4-amido)pyrrolidine-1-carboxylate (180 mg, 0.36 mmol), 2-fluoro-5-(trifluoromethyl)benzonitrile (83 mg, 0.44 mmol), and DIEA (142 mg, 1.1 mmol) in DMSO (2 mL) was stirred 70 °C for 2 h. The reaction was cooled to rt and directly purified by Prep-HPLC to afford the title compound (190 mg, 78%) as a yellow solid. LCMS (M+H)⁺ = 665.3.


[0356] Step 2-8, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}-N-[(3R)-pyrrolidin-3-yl]piperidine-4-carboxamide (cpd 2-21): To a solution of tert-butyl (3R)-3-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}piperidine-4-amido}pyrrolidine-1-carboxylate (190 mg, 0.28 mmol) in DCM (2 mL) was added TFA (0.4 mL) at rt. The resulting mixture was stirred at rt for 2h and then diluted with H₂O. The pH of the solution was adjusted to 8-9 with sat-NaHCO₃. The solution was extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, and concentrated to afford the title compound (130 mg, 80%) as a yellow solid. LCMS (M+H)⁺ = 565.2.

[0357] Step 2-9, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide formate: To a solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl }-N-[(3R)-pyrrolidin-3-yl]piperidine-4-carboxamide (55 mg, 0.097 mmol) in MeOH (2 mL) was added formaldehyde (20 mg, 0.67 mmol) and AcOH (6.0 mg, 0.10 mmol), and followed by sodium cyanotrihydroborate (12 mg, 0.19 mmol) at rt. The resulting solution was stirred at rt for 2 h, and then concentrated under vacuum. The residue was purified by Prep-HPLC to afford the title compound (20 mg, 33%) as a white solid. LCMS (M+H)⁺ = 579.3.

Example 3: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl] N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-51)

[0358]





[0359] Step 3-1. preparation of 1-tert-butyl 3-ethyl 2-(5-chloropyrazin-2-yl)propanedioate: Into a 100-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, were placed 2,5-dichloropyrazine (4.0 g, 27 mmol), cesium carbonate (26 g, 80 mmol), tert-butyl ethyl malonate (5.6 g, 30 mmol) and DMSO (40 mL). The resulting mixture was stirred 100 °C for 2 h and cooled to room temperature. The reaction was quenched with water (200 mL) and then extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:3). This resulted in the title compound (4.6 g, 57%) as a yellow oil. LCMS (M+H)⁺ = 301.1.

[0360] Step 3-2. preparation of ethyl 2-(5-chloropyrazin-2-yl)acetate: To a solution of 1-tert-butyl 3-ethyl 2-(5-chloropyrazin-2-yl)propanedioate (4.6 g, 15 mmol) in DCM (40 mL) was added TFA (10 mL) in an ice bath. The reaction was stirred at 0 °C for 1 h and quenched with water (100 mL). The resulting solution was extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:3) to afford the title compound (2.6 g, 85%) as yellow oil. LCMS (M+H)⁺ = 201.1.

[0361] Step 3-3. preparation of ethyl 1-benzyl-4-(5-chloropyrazin-2-yl)piperidine-4-carboxylate: Into a 50 mL round-bottom flask, were placed ethyl 2-(5-chloropyrazin-2-yl)acetate (1.0 g, 5.0 mmol), 18-crown-6 (300 mg, 1.13 mmol) and DMF (10 mL). The reaction mixture was cooled to 0 °C and then treated with 60%-NaH (600 mg, 15 mmol). After stirring at 0 °C for 0.5 h, the reaction was treated with N-benzyl-2-bromo-N-(2-bromoethyl)ethan-1-amine (1.8 g, 5.6 mmol). The resulting mixture was stirred at rt for 2 h and then quenched with water (50 mL). The resulting solution was extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:3). This resulted in the title compound (800 mg, 45%) as a yellow oil. LCMS (M+H)⁺ = 360.2.

[0362] Step 3-4. preparation of ethyl 1-benzyl-4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]piperidine-4-carboxylate: Into a 50 mL three-neck bottle, purged and maintained with an inert atmosphere of nitrogen, were added ethyl 1-benzyl-4-(5-chloropyrazin-2-yl)piperidine-4-carboxylate (800 mg, 2.22 mmol), (2-ethoxypyridin-3-yl)boronic acid (557 mg, 3.34 mmol), Pd(DTBPF)Cl₂ (145 mg, 0.22 mmol), K₂CO₃ (922 mg, 6.67 mmol), and dioxane/H₂O (8 mL/0.8 mL). The resulting mixture was stirred 90 °C for 2 h under N₂ and then cooled to room temperature. The reaction was quenched with water (50 mL) and extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium

sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1: 1) to afford the title compound (800 mg, 80%) as yellow oil. LCMS (M+H)⁺ = 447.5.

[0363] Step 3-5. preparation of ethyl 4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]piperidine-4-carboxylate: To a solution of ethyl 1-benzyl-4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]piperidine-4-carboxylate (800 mg, 1.79 mmol) in MeOH (20 mL) was added 10%-Pd/C (191 mg) and Pd(OH)₂ (252 mg, 1.79 mmol). The flask was evacuated and flushed three times with nitrogen, followed by flushing with hydrogen. The mixture was stirred at rt 40 min under an atmosphere of hydrogen (balloon). The resulting solution was diluted with MeOH and filtered through a pad of Celite. The filtrate was concentrated under vacuum to afford the title compound (580 mg, 91%) as a white oil. LCMS (M+H)⁺ = 357.3.


[0364] Step 3-6. preparation of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]piperidine-4-carboxylate: Into a 8-mL vial, were placed ethyl 4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]piperidine-4-carboxylate (200 mg, 0.56 mmol), DIEA (218 mg, 1.69 mmol), 2-fluoro-5-(trifluoromethyl)benzonitrile (212 mg, 1.12 mmol) and DMSO (2 mL). The resulting mixture was stirred at 60 °C for 2 h and then cooled to room temperature. The crude product was purified by Prep-HPLC to afford the title compound (220 mg, 74%). LCMS (M+H)⁺ = 526.5.

[0365] Step 3-7. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: Into a 8-mL vial purged and maintained with an inert atmosphere of nitrogen, were placed ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]piperidine-4-carboxylate (40 mg, 0.076 mmol), (R)-1-methylpyrrolidine-3-amine (23 mg, 0.23 mmol), LHMDs (130 mg, 0.77 mmol), and THF (1.3 mL). The resulting mixture was stirred at rt for 1 h under N₂ and then quenched with sat-NH₄Cl (20 mL). The resulting solution was extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by Prep-HPLC to afford the title compound (13 mg, 28%). LCMS (M+H)⁺ = 580.2.

[0366] The following compounds were prepared similarly to **Example 3** with appropriate substituting reagents and substrates at different steps.

Compound no.	MS (M+H) ⁺
1-58	580.2

Example 4: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-56)

[0367]

[0368] Step 4-1. preparation of ethyl 1-benzyl-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidine-4-carboxylate: Starting from 3,6-dichloropyridazine, the title compound was prepared by a similar manner described from step 3-1 to 3-4 in **Example 3**. LCMS (M+H)⁺ = 447.3.

[0369] Step 4-2. preparation of 1-tert-butyl 4-ethyl 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidine-1,4-dicarboxylate: Into a 100-mL round bottom flask, was placed ethyl 1-benzyl-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidine-4-carboxylate (240 mg, 0.54 mmol), di-tert-butyl decarbonate (350 mg, 1.60 mmol), MeOH (30 mL), TEA (160 mg, 1.58 mmol), 10%-Pd/C (20 mg) and Pd(OH)₂ (20 mg, 0.14 mmol). The flask was evacuated and flushed three times with nitrogen, followed by flushing with hydrogen. The reaction mixture was stirred at rt for 2h under an atmosphere of hydrogen (balloon). The reaction was filtered through a pad of Celite, and the filtrate was concentrated. The residue was purified by Prep-HPLC to afford the title compound (200 mg, 81%) as a light yellow oil. LCMS (M+H)⁺ = 457.4.

[0370] Step 4-3. preparation of ethyl 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidine-4-carboxylate trifluoroacetate: To a solution of 1-tert-butyl 4-ethyl 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidine-1,4-dicarboxylate (200 mg, 0.44 mmol) in DCM (3 mL) was added TFA (1 mL). The resulting mixture was stirred at rt for 1 h and then concentrated under vacuum. This resulted in the title compound (200 mg, 96%) as yellow oil. LCMS (M+H)⁺ = 357.2.

[0371] Step 4-4. preparation of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidine-4-carboxylate trifluoroacetate: Into a 8-mL vial, was placed ethyl 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidine-4-carboxylate trifluoroacetate (200 mg, 0.42 mmol), 2-fluoro-5-(trifluoromethyl)benzonitrile (96 mg, 51 mmol), DIEA (200 mg, 1.55 mmol) and DMSO (3 mL). The resulting solution was stirred at 60 °C for 1 h and cooled to rt. The reaction was directly purified by Prep-HPLC to afford the title compound (145 mg, 60%) as a light yellow oil. LCMS (M+H)⁺ = 526.3.

[0372] Step 4-5. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: Into a 8-mL vial, were placed ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-


yl]piperidine-4-carboxylate (60 mg, 0.11 mmol), (S)-1-methylpyrrolidine-3-amine (30 mg, 0.30 mmol), and 1M-LiHMDS in THF (1.0 mL, 1.0 mmol) under N₂ atmosphere. The resulting mixture was stirred at rt for 1 h and then quenched with sat-NH₄Cl. The resulting solution was extracted with EtOAc (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by Prep-HPLC to afford the title compound (16 mg, 22%). LCMS (M+H)⁺ = 580.2.

[0373] The following compounds were prepared similarly to **Example 4** with appropriate substituting reagents and substrates at different steps.

Compound no.	MS (M+H) ⁺
1-57	580.2
1-134	545.3
1-135	531.2
1-136	565.3
1-139	546.3
1-140	566.3
1-168	538.2
Compound no.	MS (M+H) ⁺
1-169	547.2
1-170	504.3
1-171	531.2
1-174	572.2
1-175	588.2
1-183	565.5
1-331	568.4

Example 5: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-106)

[0374]



[0375] Step 5-1, preparation of [2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]methanol: Into a 100 mL three-neck bottle, purged and maintained with an inert atmosphere of nitrogen, were added (2-chloropyrimidin-5-yl)methanol (3.0 g, 21 mmol), (2-ethoxypyridin-3-yl)boronic acid (4.0 g, 21 mmol), Pd(DTBPF)Cl₂ (1.4 g, 2.1 mmol), K₂CO₃ (9.0g, 65 mmol) and dioxane (30 mL)/H₂O (3 mL) at rt. The resulting reaction mixture was stirred at 60 °C for 1 h and cooled to room temperature. The reaction was quenched with water and extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (2:1) to afford the title compound (4.2 g, 88%). LCMS (M+H)⁺ = 232.1.

[0376] Step 5-2, preparation of [2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]methyl methanesulfonate: To a solution of [2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]methanol (2.0 g, 8.6 mmol) and TEA (3.0 g, 30 mmol) in DCM (20 mL) was slowly added MsCl (1.0 g, 8.7 mmol) in an ice bath. The reaction was stirred at 0 °C for 1h, quenched with water (30 mL), and extracted with DCM (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in the title compound (2.3 g, 86%) as a yellow oil. LCMS (M+H)⁺ = 310.1.

[0377] Step 5-3, preparation of 2-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]acetonitrile: Into a 50-mL round-bottom flask, were placed [2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]methyl methanesulfonate (2.3 g, 7.4 mmol), sodium cyanide (1.1 g, 22 mmol) and DMSO (20 mL). The resulting mixture was at rt stirred for 1 h, quenched with aq-FeSO₄(~100 mL), and then extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with DCM/MeOH (10:1) to afford the title compound (1.3 g, 73%) as a yellow oil. LCMS (M+H)⁺ = 241.1.

[0378] Step 5-4, preparation of 1-benzyl-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]piperidine-4-carbonitrile: Into a 50-mL round-bottom flask, were placed 2-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]acetonitrile (600 mg, 2.5 mmol), N-benzyl-2-bromo-N-(2-bromoethyl)ethan-1-amine (962 mg, 3.0 mmol), KOH (420 mg, 7.49 mmol), and DMSO (10 mL). The resulting mixture was stirred at rt for 1 h, quenched with water (50 mL) and extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1). This resulted in the title compound (450 mg, 45%) as a yellow oil. LCMS (M+H)⁺ = 400.2.

[0379] Step 5-5, preparation of 1-benzyl-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]piperidine-4-carboxylic acid: Into a 50-mL round-bottom flask, were placed 1-benzyl-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]piperidine-4-carbonitrile (450 mg, 1.13 mmol), KOH (632 mg, 11.3 mmol), and EtOH (5 mL)/H₂O (2.5 mL). The resulting mixture was stirred at 100 °C for 16 h and

cooled to room temperature. The reaction was diluted with water (20 mL), then adjusted to pH 6~7 with 3N-HCl and extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in the title compound (400 mg, 85%) as a yellow oil. LCMS (M+H)⁺ = 419.2.

[0380] Step 5-6, preparation of 1-benzyl-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of 1-benzyl-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]piperidine-4-carboxylic acid (180 mg, 0.43 mmol) in DMF (2 mL) was added HATU (164 mg, 0.43 mmol) and DIEA (167 mg, 1.29 mmol) at rt. After stirring for 10 min at rt, the reaction was treated with (S)-1-methylpyrrolidine-3-amine (52 mg, 0.52 mmol). The resulting mixture was stirred at rt for 1 h and then directly purified by prep-HPLC to afford the title compound (150 mg, 70%) as a white oil. LCMS (M+H)⁺ = 501.3.

[0381] Step 5-7, preparation of tert-butyl 4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-4-[(3S)-1-methylpyrrolidin-3-yl]carbamoyl]piperidine-1-carboxylate: Into a 50-mL round-bottom flask, were placed 1-benzyl-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (160 mg, 0.32 mmol), TEA (98 mg, 0.97 mmol), (Boc)₂O (209 mg, 0.96 mmol), wet 10%-Pd/C (34 mg), Pd(OH)₂ (45 mg, 0.32 mmol) and MeOH (10 mL). The flask was evacuated and flushed three times with nitrogen, followed by flushing with hydrogen. The mixture was stirred at rt 30 min under an atmosphere of hydrogen (balloon). The reaction was diluted with MeOH and filtered through a pad of Celite. The filtrate was concentrated under vacuum. The residue was purified by prep-HPLC to afford the title compound (40 mg, 25%) as a white oil. LCMS (M+H)⁺ = 511.3.

[0382] Step 5-8, preparation of 4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide bistrifluoroacetate: To a solution of tert-butyl 4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-4-[(3S)-1-methylpyrrolidin-3-yl]carbamoyl]piperidine-1-carboxylate (40 mg, 0.078 mmol) in DCM (1 mL) was added TFA (0.3 mL). The resulting solution was stirred at rt for 1 h and then concentrated under vacuum. This resulted in the title compound (30 mg, 60 %) as yellow oil. LCMS (M+H)⁺ = 411.3.


[0383] Step 5-9, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide bistrifluoroacetate: Into a 8-mL vial, were placed 4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide bistrifluoroacetate (30 mg, 0.047 mmol), DIEA (28 mg, 0.22 mmol), 2-fluoro-5-(trifluoromethyl)benzonitrile (28 mg, 0.15 mmol), and DMSO (1 mL). The resulting mixture was stirred at 60 °C for 2 h and cooled to room temperature. The reaction was purified by Prep-HPLC to afford the title compound (14 mg, 38%). LCMS (M+H)⁺ = 580.3.

[0384] The following compounds were prepared similarly to Example 5 with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-107	580.3

Example 6: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-124)

[0385]



[0386] Step 6-1, preparation of 2-chloro-3-fluoro-5-(1,2-oxazol-4-yl)pyridine: Into a 250 mL three-neck bottle, purged and maintained with an inert atmosphere of nitrogen, were added 5-bromo-2-chloro-3-fluoropyridine (5.0 g, 24 mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (5.6 g, 29 mmol), Pd(DTBPF)Cl₂ (1.5 g, 2.3 mmol), KF (4.1 g, 71 mmol), and DMSO (50 mL)/H₂O (5 mL). The resulting mixture was stirred at 100 °C for 1 h and cooled to room temperature. The reaction was diluted with H₂O and extracted with EtOAc (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by prep-HPLC to afford the title compound (3.5 g, 74%) as yellow solid. LCMS (M+H)⁺ = 199.0.

[0387] Step 6-2, preparation of 2-(6-chloro-5-fluoropyridin-3-yl)acetonitrile: To a solution of 2-chloro-3-fluoro-5-(1,2-oxazol-4-yl)pyridine (3.5 g, 18 mmol) in MeOH (35 mL)/H₂O (3.5 mL) was added KF (0.2 g, 3.0 mmol). The resulting solution was stirred at 90 °C for 1 h and cooled to room temperature. The reaction was diluted with water (100 mL) and extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1: 1) to afford the title compound (2.3 g, 77%) as yellow solid. LCMS (M+H)⁺ = 171.1.

[0388] Step 6-3, preparation of 1-benzyl-4-(6-chloro-5-fluoropyridin-3-yl)piperidine-4-carbonitrile: Into a 50-mL round-bottom flask, were placed 2-(6-chloro-5-fluoropyridin-3-yl)acetonitrile (2.3 g, 13 mmol), N-benzyl-2-bromo-N-(2-bromoethyl)ethan-1-amine (4.3 g, 13 mmol), KOH (2.3 g, 41 mmol), and DMSO (25 mL). The resulting mixture was stirred at rt for 1

h, quenched with water (100 mL) and extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1). This resulted in the title compound (2.7 g, 61%) as a yellow oil. LCMS (M+H)⁺ = 330.2.

[0389] Step 6-4, preparation of 1-benzyl-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]piperidine-4-carbonitrile: Into a 50 mL three-neck bottle, purged and maintained with an inert atmosphere of nitrogen, were added 1-benzyl-4-(6-chloro-5-fluoropyridin-3-yl)piperidine-4-carbonitrile (500 mg, 1.52 mmol), (2-ethoxyphenyl)boronic acid (300 mg, 1.81 mmol), Pd(DTBPF)Cl₂ (99 mg, 0.15 mmol), K₂CO₃ (630 mg, 4.56 mmol) and dioxane (5 mL)/H₂O (0.5 mL) at rt. The resulting reaction mixture was stirred at 100 °C for 1 h and cooled to room temperature. The reaction was quenched with water and extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1: 1) to afford the title compound (500 mg, 79%) as light yellow oil. LCMS (M+H)⁺ = 416.3.

[0390] Step 6-5, preparation of 1-benzyl-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]piperidine-4-carboxylic acid: Into a 50-mL round-bottom flask, were placed 1-benzyl-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]piperidine-4-carbonitrile (500 mg, 1.2 mmol), KOH (675 mg, 12.0 mmol), and EtOH (10 mL)/H₂O (5 mL). The resulting mixture was stirred at 100 °C for 24 h and cooled to room temperature. The reaction was diluted with water (20 ml), then adjusted to pH 6~7 with 3N-HCl and extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in the title compound (450 mg, 86%) as a yellow solid. LCMS (M+H)⁺ = 435.2.

[0391] Step 6-6, preparation of 1-benzyl-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of 1-benzyl-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]piperidine-4-carboxylic acid (140 mg, 0.32 mmol) in DMF (2 mL) was added HATU (123 mg, 0.32 mmol) and DIEA (125 mg, 0.97 mmol). After stirring for 10 min at rt, the reaction was treated with (S)-1-methylpyrrolidine-3-amine dihydrochloride (56 mg, 0.32 mmol). The resulting mixture was stirred at rt for 1 h and then directly purified by prep-HPLC to afford the title compound (130 mg, 73%) as a white solid. LCMS (M+H)⁺ = 517.4.

[0392] Step 6-7, preparation of 4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3 S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: Into a 50-mL round-bottom flask, were placed 1-benzyl-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (130 mg, 0.25 mmol), wet 10%-Pd/C (27 mg), Pd(OH)₂ (35 mg, 0.25 mmol) and MeOH (5 mL). The flask was evacuated and flushed three times with nitrogen, followed by flushing with hydrogen. The mixture was stirred at rt 30 min under an atmosphere of hydrogen (balloon). The reaction was diluted with MeOH and filtered through a pad of Celite. The filtrate was concentrated under vacuum. This resulted in the title compound (90 mg, 84%) as a white

oil. LCMS (M+H)⁺ = 427.5.


[0393] Step 6-8. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide formate: Into a 8-mL vial, were placed 4-[6-(2-ethoxyphenyl)-5-fluoropyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (45 mg, 0.11 mmol), DIEA (41 mg, 0.32 mmol), 2-fluoro-5-(trifluoromethyl)benzonitrile (40 mg, 0.21 mmol), and DMSO (1 mL). The resulting mixture was stirred at 60 °C for 2 h and cooled to room temperature. The reaction was purified by Prep-HPLC to afford the title compound (29 mg, 43%) as a white solid. LCMS (M+H)⁺ = 596.3.

[0394] The following compounds were prepared similarly to **Example 6** with appropriate substituting reagents and substrates at different steps. Some examples require deprotection in the final step:

Compound no.	MS (M+H) ⁺
1-108	597.3
1-109	597.3
1-110	598.3
1-123	596.3
1-125	597.3
1-126	582.3
1-127	582.3
1-128	583.3
1-131	583.3
Compound no.	MS (M+H) ⁺
1-132	583.3
1-133	584.3
1-161	555.2
1-162	564.5
1-163	522.2
1-195	600.3
1-196	556.3
1-197	570.3
1-334	615.3

Example 7: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-9)

[0395]



[0396] Step 7-1, preparation of methyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate: To a solution of methyl 6-chloropyridine-3-carboxylate (4.0 g, 22 mmol) and 18-Crown-6 (1.1 g, 4.2 mmol) in DMF (30 mL) at 0°C was added 60 % sodium hydride in mineral oil (2.2 g, 55 mmol) in portions. The mixture was stirred at 0°C for 2 h. To this was added a solution of benzylbis(2-bromoethyl)amine (8.3 g, 26 mmol) in DMF (10 mL). The mixture was stirred at rt for 2 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness to give the title compound (4.8 g, 65%) as a yellow solid. LCMS (M+H)⁺ = 345.2.

[0397] Step 7-2, preparation of methyl 1-benzyl-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate: To a suspension of methyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate (1.8 g, 5.2 mmol) in 1,4-dioxane (20 mL) and water (2 mL) under nitrogen was added potassium carbonate (2.2 g, 16 mmol), (2-ethoxypyridin-3-yl)boronic acid (1.7 g, 10 mmol) and Pd(dtbpf)Cl₂ (0.10 g, 0.15 mmol). The mixture was heated at 90°C for 2 h. The mixture was quenched with water and extracted with EtOAc (2X). The combined organics were concentrated to dryness and the residue was purified by silica gel CC to give the title compound (1.8 g, 80%) as a light yellow solid. LCMS (M+H)⁺ = 432.3.

[0398] Step 7-3, preparation of methyl 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate: To a solution of methyl 1-benzyl-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate (1.8 g, 4.2 mmol) in MeOH (25 mL) was added 10 wt.% of Pd on carbon (100 mg) and 20 wt.% of Pd(OH)₂ on carbon (100 mg). The mixture was charged with hydrogen (g) and stirred at rt for 2 h. The mixture was filtered and the filtrate was concentrated to dryness to give the title compound (1.2 g, 60%) as yellow oil. LCMS (M+H)⁺ = 342.2.

[0399] Step 7-4, preparation of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate: To a solution of methyl 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate (1.2 g, 3.5 mmol) and 2-fluoro-5-(trifluoromethyl)benzonitrile (0.8 g, 4 mmol) in DMSO (15 mL) was added DIEA (1.4 g, 11 mmol). The mixture was heated at 60°C for 2 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were concentrated to dryness and the residue was purified by silica gel CC to give the

title compound (1.1 g, 61%) as a white solid. LCMS (M+H)⁺ = 511.3.

[0400] Step 7-5, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid: To a suspension of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate (1.4 g, 2.7 mmol) in a mixed solvent of EtOH and water (2:1, 12 mL) was added lithium hydroxide (0.60 g, 25 mmol). The mixture was heated at 60°C for 2 h. The mixture was concentrated to remove the organics and the aqueous residue was adjusted to pH 6-7 with 1N HCl (aq). The solution was extracted with EtOAc (3X) and the combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness to give the title compound (1.2 g, 88%) as a light yellow solid. LCMS (M+H)⁺ = 497.3.


[0401] Step 7-6, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid (60 mg, 0.12 mmol) and HATU (55 mg, 0.14 mmol) in DMF (1 mL) was added DIEA (94 mg, 0.73 mmol). After stirring at rt for 5 min, (3R)-1-methylpyrrolidin-3-amine dihydrochloride (31 mg, 0.18 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 1 h and purified by reversed-phase CC to give the title compound (42 mg, 56%) as a white solid. LCMS (M+H)⁺ = 579.3.

[0402] The following compounds were prepared similarly to **Example 7** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-11	579.3
1-15	567.4
1-18	579.3
1-19	565.3
1-20	605.3
1-41	595.3
1-45	597.2
Compound no.	MS (M+H) ⁺
1-46	595.2
1-48	597.2
1-166	564.5
1-176	535.5
1-178	565.7
1-179	565.4

Example 8: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3*R*)-pyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-10)

[0403]



[0404] Step 8-1. preparation of *tert*-butyl (3*R*)-3-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-amido}pyrrolidine-1-carboxylate: To a solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid (60 mg, 0.10 mmol) from Step 7-5 and HATU (58 mg, 0.15 mmol) in DMF (1 mL) was added DIEA (0.10 mL, 0.57 mmol). After stirring at rt for 5 min, *tert*-butyl (3*R*)-3-aminopyrrolidine-1-carboxylate (45 mg, 0.24 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 1 h and purified by reversed-phase CC to give the title compound (69 mg, 100%) as light brown oil. LCMS (M+H)⁺ = 665.3.

[0405] Step 8-2. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3*R*)-pyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of *tert*-butyl (3*R*)-3-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-amido}pyrrolidine-1-carboxylate (69 mg, 0.10 mmol) in DCM (4 mL) was added TFA (1 mL). The mixture was stirred at rt for 1 h. The mixture was concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (32 mg, 51%) as light brown oil. LCMS (M+H)⁺ = 565.3.


[0406] The following compounds were prepared similarly to **Example 8** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-12	565.3
1-13	539.3
1-14	553.3
1-16	553.3
1-17	569.3

Compound no.	MS (M+H) ⁺
1-28	569.2
1-44	583.3
1-47	583.3
1-21	565.3

Example 9: 1-(2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-32)

[0407]



[0408] Step 9-1, preparation of ethyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate: To a solution of ethyl 6-chloropyridine-3-carboxylate (9.5 g, 48 mmol) in DMF (100 mL) at 0°C was added 60 % sodium hydride in mineral oil (3.4 g, 0.14 mol) in portions. The mixture was stirred at 0°C for 0.5 h. To this was added benzylbis(2-bromoethyl)amine (8.3 g, 26 mmol) and the mixture was stirred at rt for 2 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness. The residue was purified by silica gel CC to give the title compound (12 g, 70%) as yellow oil. LCMS (M+H)⁺ = 359.2.

[0409] Step 9-2, preparation of ethyl 1-benzyl-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate: To a suspension of ethyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate (5.0 g, 14 mmol) in 1,4-dioxane (50 mL) and water (5 mL) under nitrogen was added potassium carbonate (5.8 g, 42 mmol), (2-ethoxypyridin-3-yl)boronic acid (3.5 g, 21 mmol) and Pd(dtbpf)Cl₂ (0.91 g, 1.4 mmol). The mixture was heated at 90°C for 2 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were concentrated to dryness and the residue was purified by silica gel CC to give the title compound (5.5 g, 89%) as a yellow solid. LCMS (M+H)⁺ = 446.3.

[0410] Step 9-3, preparation of 1-benzyl-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid: To a suspension of ethyl 1-(4-chloro-2-cyanophenyl)-4-[6-(2-

ethoxyphenyl]pyridin-3-yl]piperidine-4-carboxylate (2.0 g, 4.5 mmol) in a mixed solvent of EtOH and water (2:1, 45 mL) was added lithium hydroxide (1.1 g, 46 mmol). The mixture was heated at 60°C for 1 h. The mixture was diluted with water and adjusted to pH 6-7 with 3N HCl (aq). The mixture was extracted with EtOAc (3X) and the combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness to give the title compound (1.5 g, 80%) as yellow oil. LCMS (M+H)⁺ = 418.1.

[0411] Step 9-4. preparation of 1-benzyl-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of 1-benzyl-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl]piperidine-4-carboxylic acid (1.2 g, 2.9 mmol) and HATU (1.1 g, 2.9 mmol) in DMF (12 mL) was added DIEA (1.5 mL, 8.5 mmol). After stirring at rt for 5 min, (3S)-1-methylpyrrolidin-3-amine (0.30 g, 3.0 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 2 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were concentrated to dryness and the residue was purified by silica gel CC to give the title compound (1.1 g, 77%) as yellow oil. LCMS (M+H)⁺ = 500.4.

[0412] Step 9-5. preparation of 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of 1-benzyl-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (1.1 g, 2.2 mmol) in MeOH (20 mL) was added 10 wt.% of Pd on carbon (0.23 g) and 20 wt.% of Pd(OH)₂ on carbon (0.31 g). The mixture was charged with hydrogen (g) and stirred at rt for 1 h. The mixture was filtered and the filtrate was concentrated to dryness to give the title compound (0.70 g, 78%) as white oil. LCMS (M+H)⁺ = 410.3.

[0413] Step 9-6. preparation of 1-(2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (45 mg, 0.11 mmol) and 2-fluorobenzonitrile (40 mg, 0.33 mmol) in DMSO (1 mL) was added DIEA (28 mg, 0.22 mmol). The mixture was heated at 120°C for 1 h. The mixture was purified by reversed-phase CC to give the title compound (18 mg, 31%) as a white solid. LCMS (M+H)⁺ = 511.2.


[0414] The following compounds were prepared similarly to **Example 9** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-34	545.2
1-35	555.3
1-36	553.2
1-37	561.3
1-38	579.2
1-39	595.2

Compound no.	MS (M+H) ⁺
1-40	536.2
1-42	580.3
1-59	575.3
1-150	588.4
1-308	585.3
1-311	571.3

Example 10: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]-1-[4-(trifluoromethyl)phenyl]piperidine-4-carboxamide (cpd 1-33)

[0415]




[0416] Step 10-1, preparation of 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]-1-[4-(trifluoromethyl)phenyl]piperidine-4-carboxamide formate: To a mixture of 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (40 mg, 0.098 mmol), 1-bromo-4-(trifluoromethyl)benzene (33 mg, 0.115 mmol), XPhos (17 mg, 0.020 mmol), Pd₂(dba)₃ (9.0 mg, 0.0098 mmol), and sodium tert-butoxide (28 mg, 0.29 mmol) under nitrogen was added toluene (1 mL). The mixture was heated at 80°C for 1 h. The mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by reversed-phase CC to give the title compound (25 mg, 43%) as yellow oil. LCMS (M+H)⁺ = 554.2.

[0417] The following compounds were prepared similarly to **Example 10** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-154	592.2
1-155	591.2
1-156	571.2
1-186	572.3

Example 11: 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide & 3-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-[(3R)-1-methylpyrrolidin-3-yl]carbamoyl)piperidin-1-yl)-6-(trifluoromethyl)pyridine-2-carboxamide (cpd 1-24 & 1-25)

[0418]



[0419] Step 11-1, preparation of ethyl 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate: To a solution of ethyl 1-benzyl-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate (2.0 g, 4.5 mmol) in MeOH (100 mL) was added 10 wt.% of Pd on carbon (100 mg) and 20 wt.% of Pd(OH)₂ on carbon (100 mg). The mixture was stirred at rt for 3 h under hydrogen (3 bar). The mixture was filtered and the filtrate was concentrated to dryness to give the title compound (1.0 g, 62%) as light yellow oil. LCMS (M+H)⁺ = 356.2.

[0420] Step 11-2, preparation of ethyl 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate: To a solution of ethyl 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate (0.2 g, 0.6 mmol) and 3-fluoro-6-(trifluoromethyl)pyridine-2-carbonitrile (0.16 g, 0.84 mmol) in DMSO (5 mL) was added DIEA (0.4 mL, 2 mmol). The mixture was heated at 60°C for 2 h. The mixture was purified by reversed-phase CC to give the title compound (0.21 g, 70%) as light yellow oil. LCMS (M+H)⁺ = 526.5.

[0421] Step 11-3, preparation of 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid & 1-[2-carbamoyl-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid: To a suspension of ethyl 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate (0.21 g, 0.40 mmol) in a mixed solvent of EtOH and water (2:1, 6 mL) was added lithium hydroxide (0.10 g, 4.2 mmol). The mixture was heated at 60°C for 1 h. The mixture was adjusted to pH 6-7 with 1N HCl (aq). The solution was extracted with EtOAc (3X) and the combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness to give the mixture of title compounds (0.17 g) in a ratio of 47:53 as a light yellow solid. LCMS (M+H)⁺

= 498.3 & 516.3.


[0422] Step 11-4. preparation of 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3*R*)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide & 3-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-[(3*R*)-1-methylpyrrolidin-3-yl]carbamoyl)piperidin-1-yl)-6-(trifluoromethyl)pyridine-2-carboxamide: To a solution of a mixture of 1-[2-cyano-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid & 1-[2-carbamoyl-6-(trifluoromethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid (85 mg, 0.17 mmol) and HATU (79 mg, 0.21 mmol) in DMF (1 mL) was added DIEA (0.1 mL, 0.8 mmol). After stirring at rt for 5 min, (3*R*)-1-methylpyrrolidin-3-amine (64 mg, 0.64 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 1 h and purified by reversed-phase CC to give the first title compound (32 mg, 33% yield) as a white solid; LCMS (M+H)⁺ = 580.3 and the second title compound (44 mg, 43%) as a white solid; LCMS (M+H)⁺ = 598.3.

[0423] The following compounds were prepared similarly to **Example 1** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-26	580.3
1-27	598.3

Example 12: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1*H*-pyrrol-2-yl)pyridin-3-yl]-N-[(3*R*)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-88)

[0424]



[0425] Step 12-1. preparation of methyl 4-(6-chloropyridin-3-yl)piperidine-4-carboxylate: To a solution of methyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate (2.9 g, 8.4 mmol) from step 7-1 in DCM (30 mL) at 0°C was added chloroethyl chloroformate (2.3 mL, 21 mmol). The mixture was stirred at rt for 2 h and then MeOH (15 mL) was added. The resulting mixture was heated at reflux for 2 h. The mixture was concentrated to dryness and the residue was

trituated with a mixed solvent of hexanes and ether (5:1, 40 mL). The solid was collected by vacuum filtration to give the title compound (1.7 g, 79%) as a white solid. LCMS (M+H)⁺ = 255.0.

[0426] Step 12-2. preparation of methyl 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate: To a solution of methyl 4-(6-chloropyridin-3-yl)piperidine-4-carboxylate (1.7 g, 6.7 mmol) and 2-fluoro-5-(trifluoromethyl)benzonitrile (1.3 g, 7.0 mmol) in DMSO (17 mL) was added DIEA (3.6 mL, 21 mmol). The mixture was heated at 60°C for 2 h. The mixture was diluted with 10% citric acid (aq) (50 mL) and extracted with EtOAc (3X). The combined organics were washed with brine, dried over anhydrous Na₂SO₄ and concentrated to dryness. The residue was trituated with hexanes and the solid was collected by vacuum filtration to give the title compound (1.8 g, 64%) as a white solid. LCMS (M+H)⁺ = 424.0.

[0427] Step 12-3. preparation of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylic acid: To a suspension of methyl 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate (1.8 g, 4.2 mmol) in a mixed solvent of EtOH and water (2:1, 22.5 mL) was added lithium hydroxide (1.0 g, 42 mmol). The mixture was heated at 60°C for 2 h. The mixture was adjusted to pH 6-7 with 1N HCl (aq). The solution was extracted with EtOAc (3X) and the combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness. The residue was purified by reversed-phase CC to give the title compound (1.5 g, 83%) as an off-white solid. LCMS (M+H)⁺ = 409.9.

[0428] Step 12-4. preparation of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylic acid (500 mg, 1.22 mmol) and HATU (650 mg, 1.71 mmol) in DMF (1 mL) was added DIEA (0.43 mL, 2.4 mmol). After stirring at rt for 5 min, (3R)-1-methylpyrrolidin-3-amine (183 mg, 1.83 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 10 min and purified by reversed-phase CC to give the title compound (483 mg, 80.5%) as an off-white solid; LCMS (M+H)⁺ = 492.0.

[0429] Step 12-5. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a mixture of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (30.0 mg, 0.0610 mmol), 1-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole (63.1 mg, 0.305 mmol), Pd(amphos)Cl₂ (17.3 mg, 0.0244 mmol), and potassium carbonate (25.3 mg, 0.183 mmol) in a sealed tube was added 1,4-dioxane (2 mL) and water (0.3 mL). Nitrogen (g) was bubbled through and the resulting mixture was heated at 100°C for 2 h. The mixture was quenched with water and extracted with EtOAc (2x). The combined organics were concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (9.1 mg, 28%) as an off-white solid. LCMS


$(M+H)^+ = 537.5$.

[0430] The following compounds were prepared similarly to **Example 12** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-43	538.5
1-52	564.5
1-80	570.3
1-83	570.3
1-84	598.4
1-89	590.5
1-105	550.7
1-114	600.5
Compound no.	MS (M+H) ⁺
1-115	537.5
1-116	538.6
1-137	573.5
1-153	562.5
1-157	539.5
1-158	563.6
1-160	546.2
1-173	530.2

Example 13: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3*R*)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-81)

[0431]



[0432] Step 13-1, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-*N*-[(3*R*)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a mixture of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-*N*-[(3*R*)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide from Step 12-4 (30.0 mg, 0.0610 mmol), (5-fluoro-2-methoxyphenyl)boronic acid (51.8 mg, 0.305 mmol), Pd(dtbpf)Cl₂ (15.9 mg, 0.0244 mmol), and potassium carbonate (25.3 mg, 0.183 mmol) in a sealed tube was added 1,4-dioxane (2 mL) and water (0.3 mL). Nitrogen (g) was bubbled through and the resulting mixture was heated at 80°C for 3 h. The mixture was quenched with water and extracted with EtOAc (2X). The combined organics were concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (17.2 mg, 48.5%) as a brown solid. LCMS (M+H)⁺ = 582.2.


[0433] The following compounds were prepared similarly to **Example 13** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-31	578.2
1-53	576.3
1-54	574.4
1-55	564.5
1-60	562.5
1-61	570.3
1-62	552.2
1-63	592.5
1-64	552.4
1-65	570.3
1-66	548.7
1-67	614.3
1-68	570.3
1-69	596.5
1-70	594.4
1-71	618.5
1-72	596.3
1-73	588.1
1-74	552.4
Compound no.	MS (M+H) ⁺
1-77	602.4
1-78	559.4

Compound no.	MS (M+H) ⁺
1-79	578.4
1-82	582.3
1-85	601.5
1-86	583.2
1-87	600.4
1-98	565.5
1-99	566.5
1-112	554.4
1-113	574.7
1-117	576.7
1-118	563.5
1-129	568.5
1-130	578.4
1-138	554.1
1-148	577.5
1-151	584.2
1-177	549.5

Example 14: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-100)

[0434]



[0435] Step 14-1. preparation of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylic acid (1.00 g, 2.44 mmol) from Step 12-3 and HATU (1.21 g, 3.17 mmol) in DMF (1 mL) was added DIEA (0.85 mL, 4.9

mmol). After stirring at rt for 5 min, (3S)-1-methylpyrrolidin-3-amine (367 mg, 3.66 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 10 min and purified by reversed-phase CC to give the title compound (1.14 g, 94.6%) as a light brown solid; LCMS (M+H)⁺ = 492.2.


[0436] Step 14-2, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a mixture of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (90.0 mg, 0.183 mmol), 1-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole (189 mg, 0.915 mmol), Pd(amphos)Cl₂ (51.8 mg, 0.0732 mmol), potassium carbonate (75.9 mg, 0.549 mmol) in a sealed tube was added 1,4-dioxane (3 mL) and water (0.3 mL). Nitrogen (g) was bubbled through and the resulting mixture was heated at 100°C for 2 h. The mixture was quenched with water and extracted with EtOAc (2x). The combined organics were concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (23.5 mg, 23.9%) as an off-white solid. LCMS (M+H)⁺ = 537.6.


[0437] The following compounds were prepared similarly to **Example 14** with appropriate substituting reagents and substrates at different steps. Some examples require a deprotection in the final step:

Compound no.	MS (M+H) ⁺
1-119	570.4
1-120	538.6
1-143	523.6
1-146	582.3
Compound no.	MS (M+H) ⁺
1-147	568.3
1-149	539.2
1-187	563.3
1-188	563.2

Example 15: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-102)

[0438]






[0439] Step 15-1, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(5-fluoro-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a mixture of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide from Step 14-1 (60.0 mg, 0.122 mmol), (5-fluoro-2-methoxyphenyl)boronic acid (104 mg, 0.610 mmol), Pd(dtbpf)Cl₂ (31.8 mg, 0.0488 mmol), and potassium carbonate (50.6 mg, 0.366 mmol) in a sealed tube was added 1,4-dioxane (2 mL) and water (0.3 mL). Nitrogen (g) was bubbled through and the resulting mixture was heated at 80°C for 3 h. The mixture was quenched with water and extracted with EtOAc (2X). The combined organics were concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (41.9 mg, 59%) as a brown solid. LCMS (M+H)⁺ = 582.4.

[0440] The following compounds were prepared similarly to **Example 15** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-30	578.2
1-101	578.2
1-103	576.3
1-104	562.4
1-121	574.5
1-122	554.4

Example 16: 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-90)

[0441]





[0442] Step 16-1. preparation of ethyl 1-benzyl-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate: To a suspension of ethyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate (4.2 g, 12 mmol) from Step 9-1 in 1,4-dioxane (50 mL) and water (5 mL) under nitrogen was added potassium carbonate (3.2 g, 23 mmol), (2-ethoxyphenyl)boronic acid (2.9 g, 17 mmol) and Pd(dtbpf)Cl₂ (0.30 g, 0.46 mmol). The mixture was heated at 90°C for 2 h. The mixture was quenched with water and extracted with EtOAc (2X). The combined organics were concentrated to dryness and the residue was purified by silica gel CC to give the title compound (4.0 g, 77%) as a yellow solid. LCMS (M+H)⁺ = 445.2.

[0443] Step 16-2. preparation of ethyl 4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate: To a solution of ethyl 1-benzyl-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (3.5 g, 7.9 mmol) in MeOH (50 mL) was added 10 wt.% of Pd on carbon (160 mg) and 20 wt.% of Pd(OH)₂ on carbon (160 mg). The mixture was charged with hydrogen (g) and stirred at rt for 16 h. The mixture was filtered and the filtrate was concentrated to dryness to give the title compound (2.4 g, 86%) as yellow oil. LCMS (M+H)⁺ = 355.2.

[0444] Step 16-3. preparation of ethyl 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate: To a mixture of ethyl 4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (400 mg, 1.13 mmol), 2-bromo-5-chlorobenzonitrile (320 mg, 1.48 mmol), XPhos (33.3 mg, 0.0699 mmol), Pd₂(dba)₃·CHCl₃ (35.0 mg, 0.0338 mmol), and cesium carbonate (735 mg, 2.26 mmol) under nitrogen was added toluene (5 mL). The mixture was heated at 90°C for 2 h. The mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by silica gel CC to give the title compound (290 mg, 52.4%) as yellow oil. LCMS (M+H)⁺ = 490.3.

[0445] Step 16-4. preparation of 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylic acid: To a suspension of ethyl 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (290 mg, 0.59 mmol) in a mixed solvent of EtOH and water (5:1, 6 mL) was added lithium hydroxide (72 mg, 3.0 mmol). The mixture was heated at 60°C for 2 h. The mixture was diluted with water and adjusted to pH 6-7 with 1N HCl (aq). The mixture was extracted with EtOAc (3X) and the combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness to give the title compound (250 mg, 91.4%) as a yellow solid. LCMS (M+H)⁺ = 462.3.

[0446] Step 16-5. preparation of 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylic acid (50 mg, 0.11 mmol) and HATU (49 mg, 0.13 mmol) in DMF (0.6 mL) was added DIEA (84 mg, 0.65 mmol). After stirring at rt for 5 min, (3S)-1-Methylpyrrolidin-3-amine (14 mg, 0.14 mmol) was added to

the above HATU-activated solution. The mixture was stirred at rt for 1 h and purified by reversed-phase CC to give the title compound (31 mg, 48%) as a white solid. LCMS (M+H)⁺ = 544.3.

[0447] The following compounds were prepared similarly to **Example 16** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-93	530.3
1-141	588.3
1-144	573.2
1-145	571.3
1-167	579.2
1-172	572.2
1-281	504.3
1-282	534.3
1-283	520.3


Compound no.	MS (M+H) ⁺
1-284	578.3
1-285	563.3
1-286	579.3
1-287	564.3
1-288	594.3
1-289	577.3
1-290	563.3
1-291	593.3
1-292	579.3

Compound no.	MS (M+H) ⁺
1-293	549.3
1-294	565.3
1-295	560.3
1-296	576.3
1-297	532.3
1-298	518.3
1-299	580.3
1-300	564.3
1-301	550.3
1-302	580.3
1-303	566.3
1-304	536.3

Compound no.	MS (M+H) ⁺
1-305	522.3
1-306	546.3
1-307	562.3
1-309	552.3
1-310	538.3
1-312	592.3
1-315	578.4
1-317	574.4
1-318	556.3
1-321	560.3
1-322	542.2

Example 17: 1-(2-cyano-4-fluorophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-96)

[0448]



[0449] Step 17-1, preparation of 1-benzyl-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-

carboxylic acid: To a suspension of ethyl 1-benzyl-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (500 mg, 1.12 mmol) from Step 16-1 in a mixed solvent of EtOH and water (5:1, 6 mL) was added lithium hydroxide (269 mg, 11.2 mmol). The mixture was heated at 60°C for 1 h. The mixture was diluted with water and adjusted to pH 6-7 with 1N HCl (aq). The mixture was extracted with EtOAc (3X) and the combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness to give the title compound (450 mg, 96.1%) as yellow oil. LCMS (M+H)⁺ = 417.3.

[0450] Step 17-2, preparation of 1-benzyl-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of 1-benzyl-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylic acid (450 mg, 1.08 mmol) and HATU (430 mg, 1.13 mmol) in DMF (5 mL) was added DIEA (450 mg, 3.48 mmol). After stirring at rt for 5 min, (3S)-1-methylpyrrolidin-3-amine (130 mg, 1.30 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 1 h and purified by reversed-phase CC to give the title compound (450 mg, 83.5%) as a yellow solid. LCMS (M+H)⁺ = 499.4.

[0451] Step 17-3, preparation of 4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of 1-benzyl-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (430 mg, 0.862 mmol) in MeOH (30 mL) was added 10 wt.% of Pd on carbon (20 mg) and 20 wt.% of Pd(OH)₂ on carbon (20 mg). The mixture was charged with hydrogen (g) and stirred at rt for 2 h. The mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by reversed-phase CC to give the title compound (360 mg, 79.9%) as colorless oil. LCMS (M+H)⁺ = 409.3.

[0452] Step 17-4, preparation of 1-(2-cyano-4-fluorophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a mixture of 4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (60 mg, 0.15 mmol), 2-bromo-5-fluorobenzonitrile (50 mg, 0.25 mmol), XPhos (7.0 mg, 0.015 mmol), Pd₂(dba)₃·CHCl₃ (15 mg, 0.014 mmol), and sodium *tert*-butoxide (45 mg, 0.47 mmol) under nitrogen was added toluene (2 mL). The mixture was heated at 100°C for 4 h. The mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by reversed-phase CC to give the title compound (8.5 mg, 10%) as a white solid. LCMS (M+H)⁺ = 528.3.


[0453] The following compounds were prepared similarly to **Example 17** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-95	553.2
1-97	587.3
1-111	562.3

Example 18: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-

(1-methylazetidin-3-yl)piperidine-4-carboxamide (cpd 1-76)

[0454]





[0455] Step 18-1, preparation of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate: To a solution of ethyl 4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (0.38 g, 1.1 mmol) from Step 16-2 and 2-fluoro-5-(trifluoromethyl)benzonitrile (0.31 g, 1.6 mmol) in DMSO (5 mL) was added DIEA (0.55 mL, 3.2 mmol). The mixture was heated at 60°C for 2 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (0.41 g, 73%) as light yellow oil. LCMS (M+H)⁺ = 524.2.

[0456] Step 18-2, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide: To a solution of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (70 mg, 0.13 mmol) and 1-methylazetidin-3-amine (20 mg, 0.23 mmol) in THF (0.5 mL) under nitrogen was added 1M LiHMDS in THF (0.7 mL, 0.7 mmol). The mixture was stirred at rt for 40 min. The mixture was quenched with saturated NH₄Cl (aq) and extracted with EtOAc (3X). The combined organics were concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (29 mg, 36%) as a white solid. LCMS (M+H)⁺ = 564.3.

Example 19: 1-(4-chloro-2-cyanophenyl)-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (cpd 1-91)

[0457]





[0458] Step 19-1, preparation of ethyl 1-benzyl-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate: To a mixture of ethyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate (400 mg, 1.11 mmol) from Step 9-1, (2-methoxyphenyl)boronic acid (200 mg, 1.32 mmol), Pd(amphos)Cl₂ (60 mg, 0.085 mmol), and potassium carbonate (450 mg, 3.26 mmol) under nitrogen was added 1,4-dioxane (5 mL) and water (0.5 mL). The mixture was heated at 90°C for 2 h. The mixture was quenched with water and extracted with EtOAc (2X). The combined organics were concentrated to dryness and the residue was purified by silica gel CC to give the title compound (450 mg, 93.8%) as yellow oil. LCMS (M+H)⁺ = 431.2.

[0459] Step 19-2, preparation of ethyl 4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate: To a solution of ethyl 1-benzyl-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (440 mg, 1.02 mmol) in MeOH (30 mL) was added 10 wt.% of Pd on carbon (40 mg) and 20 wt.% of Pd(OH)₂ on carbon (40 mg). The mixture was charged with hydrogen (g) and stirred at rt for 2 h. The mixture was filtered and the filtrate was concentrated to dryness to give the title compound (260 mg, 74.8%) as a light brown solid. LCMS (M+H)⁺ = 341.2.

[0460] Step 19-3, preparation of 1-(4-chloro-2-cyanophenyl)-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxylic acid: To a mixture of ethyl 4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (180 mg, 0.529 mmol), 2-bromo-5-chlorobenzonitrile (180 mg, 0.832 mmol), XPhos (30 mg, 0.063 mmol), Pd₂(dba)₃·CHCl₃ (45 mg, 0.043 mmol), and sodium *tert*-butoxide (100 mg, 1.04 mmol) under nitrogen was added toluene (2 mL). The mixture was heated at 80°C for 2 h. The mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by reversed-phase CC to give the title compound (170 mg, 71.8%) as a yellow solid. LCMS (M+H)⁺ = 448.2.


[0461] Step 19-4, preparation of 1-(4-chloro-2-cyanophenyl)-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of 1-(4-chloro-2-cyanophenyl)-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxylic acid (60 mg, 0.13 mmol) and HATU (55 mg, 0.14 mmol) in DMF (2 mL) was added DIEA (70 mg, 0.54 mmol). After stirring at rt for 5 min, (3S)-1-methylpyrrolidin-3-amine (25 mg, 0.25 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 1 h and purified by reversed-phase CC to give the title compound (35 mg, 45%) as a white solid. LCMS (M+H)⁺ = 530.3.

[0462] The following compounds were prepared similarly to **Example 19** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-92	516.2
1-142	574.2

Example 20: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide (cpd 1-94)

[0463]



[0464] Step 20-1, preparation of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate: To a solution of ethyl 4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (70 mg, 0.21 mmol) from Step 19-2 and 2-fluoro-5-(trifluoromethyl)benzonitrile (60 mg, 0.32 mmol) in DMSO (2 mL) in a sealed tube was added DIEA (80 mg, 0.62 mmol). The mixture was heated at 60°C for 1 h. The mixture was purified by reversed-phase CC to give the title compound (60 mg, 57%) as a light brown solid. LCMS (M+H)⁺ = 510.3.

[0465] Step 20-2, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxylic acid: To a suspension of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (60 mg, 0.12 mmol) in a mixed solvent of EtOH and water (5:1, 2.4 mL) was added lithium hydroxide (15 mg, 0.63 mmol). The mixture was heated at 60°C for 1 h. The mixture was diluted with water and adjusted to pH 6-7 with 1N HCl (aq). The mixture was extracted with EtOAc (3X) and the combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness to give the title compound (55 mg, 97%) as a yellow solid. LCMS (M+H)⁺ = 482.2.

[0466] Step 20-3, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidine-4-carboxamide: To a solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxylic acid (55 mg, 0.11 mmol) and HATU (50 mg, 0.13 mmol) in DMF (2 mL) was added DIEA (50 mg, 0.39 mmol). After stirring at rt for 5 min, 1-methylazetidin-3-amine (15 mg, 0.17


mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 1 h and purified by reversed-phase CC to give the title compound (47 mg, 69%) as a white solid. LCMS (M+H)⁺ = 550.3.

[0467] The following compounds were prepared similarly to **Example 20** with appropriate substituting reagents and substrates at different steps. Some examples require a deprotection in the final step:

Compound no.	MS (M+H) ⁺
1-313	581.4
1-314	567.4
1-316	525.4
1-319	599.4
1-320	585.3
1-323	551.3
1-324	557.3
Compound no.	MS (M+H) ⁺
1-325	565.3
1-326	543.3
1-327	594.3
1-328	594.3
1-329	594.3
1-330	594.3

Example 21: 2-[4-[2-(dimethylamino)ethoxy]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-1-yl]-5-(trifluoromethyl)benzonitrile (cpd 2-1)

[0468]



[0469] Step 21-1, preparation of 2-[4-(4-bromophenyl)-4-hydroxypiperidin-1-yl]-5-


(trifluoromethyl)benzonitrile: To a solution of 4-(4-bromophenyl)piperidin-4-ol (500 mg, 1.95 mmol) and 2-fluoro-5-(trifluoromethyl)benzonitrile (443 mg, 2.34 mmol) in DMSO (3 mL) in a sealed tube was added DIEA (0.85 mL, 4.9 mmol). The mixture was heated at 130°C for 5 h. The mixture was purified by reversed-phase CC to give the title compound (779 mg, 93.8%) as a light brown gum. LCMS (M+H)⁺ = 425.2.

[0470] Step 21-2, preparation of 2-{4-[4-(2-ethoxypyridin-3-yl)phenyl]-4-hydroxypiperidin-1-yl}-5-(trifluoromethyl)benzonitrile: To a mixture of 2-[4-(4-bromophenyl)-4-hydroxypiperidin-1-yl]-5-(trifluoromethyl)benzonitrile (779 mg, 1.83 mmol), (2-ethoxypyridin-3-yl)boronic acid (459 mg, 2.75 mmol), Pd(amphos)Cl₂ (130 mg, 0.183 mmol), and potassium carbonate (506 mg, 3.66 mmol) in a sealed tube was added 1,4-dioxane (5 mL) and water (0.5 mL). The mixture was charged with N₂ (g) and heated at 100°C for 1 h. The mixture was quenched with water and extracted with EtOAc (2X). The combined organics were concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (777 mg, 90.7%) as an off-white solid. LCMS (M+H)⁺ = 468.1.

[0471] Step 21-3, preparation of 2-{4-[2-(dimethylamino)ethoxy]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-1-yl}-5-(trifluoromethyl)benzonitrile: To a solution of 2-{4-[4-(2-ethoxypyridin-3-yl)phenyl]-4-hydroxypiperidin-1-yl}-5-(trifluoromethyl)benzonitrile (40.0 mg, 0.0856 mmol) and (2-chloroethyl)dimethylamine hydrochloride (24.6 mg, 0.171 mmol) in DMF (1 mL) at 0°C was added 60% sodium hydride in mineral oil (34 mg, 0.86 mmol) in portions. The mixture was stirred at rt for 1 h and 50°C for 1 h. The reaction was not complete but stopped. The mixture was quenched with ice water and extracted with DCM(2X). The combined organics were concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (19.4 mg, 42.1%) as an off-white solid. LCMS (M+H)⁺ = 539.5.

Example 22: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl N-[2-(dimethylamino)ethyl]carbamate (cpd 2-2)

[0472]




[0473] Step 22-1. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl 1*H*-imidazole-1-carboxylate: To a suspension of 2-{4-[4-(2-ethoxypyridin-3-yl)phenyl]-4-hydroxypiperidin-1-yl}-5-(trifluoromethyl)benzonitrile (200 mg, 0.428 mmol) from Step 21-2 in THF (3 mL) was added CDI (208 mg, 1.28 mmol). The mixture was stirred at rt for 16 h. The mixture was concentrated to dryness and the residue was purified by silica gel CC to give the title compound (185 mg, 77.1%) as a white solid. LCMS (M+H)⁺ = 561.7.

[0474] Step 22-2. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl *N*-[2-(dimethylamino)ethyl]carbamate: To a solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl 1*H*-imidazole-1-carboxylate (40.0 mg, 0.0712 mmol) in THF (1 mL) in a sealed tube was added (2-aminoethyl)dimethylamine (12.6 mg, 0.142 mmol) and DIEA (0.037 mL, 0.21 mmol). The mixture was heated at 100°C for 16 h. The mixture was concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (14.9 mg, 36.0%) as a yellow solid. LCMS (M+H)⁺ = 582.4.

Example 23: 3-amino-*N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}propanamide (cpd 2-3)

[0475]



[0476] Step 23-1. preparation of *tert*-butyl 4-amino-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-1-carboxylate: To a mixture of *tert*-butyl 4-amino-4-(4-bromophenyl)piperidine-1-carboxylate (300 mg, 0.844 mmol), (2-ethoxypyridin-3-yl)boronic acid (212 mg, 1.27 mmol), Pd(amphos)Cl₂ (59.8 mg, 0.0844 mmol), and potassium carbonate (233 mg, 1.69 mmol) in a sealed tube was added 1,4-dioxane (5 mL) and water (0.5 mL). The mixture was charged with N₂ (g) and heated at 100°C for 1 h. The mixture was quenched with water and extracted with EtOAc (2X). The combined organics were concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (287 mg, 85.4%) as a brown gum. LCMS (M+H)⁺ = 365.2.

[0477] Step 23-2. preparation of *tert*-butyl 4-(3-[[[(benzyloxy)carbonyl]amino]propanamido]-4-

[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-1-carboxylate: To a solution of 3-[[[(benzyloxy)carbonyl]amino]propanoic acid (44.9 mg, 0.201 mmol) and HATU (68.9 mg, 0.181 mmol) in DMF (0.2 mL) was added DIEA (0.053 mL, 0.30 mmol). After stirring at rt for 5 min, *tert*-butyl 4-amino-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-1-carboxylate (40.0 mg, 0.101 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 10 min and purified by reversed-phase CC to give the title compound (47.5 mg, 78.3%) as a white solid. LCMS (M+H)⁺ = 603.4.


[0478] Step 23-3. preparation of benzyl *N*-[2-({4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamoyl)ethyl]carbamate: To a solution of *tert*-butyl 4-(3-[[[(benzyloxy)carbonyl]amino]propanamido]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-1-carboxylate (47.5 mg, 0.0788 mmol) in DCM (0.1 mL) was added TFA (0.091 mL, 1.2 mmol). The mixture was stirred at rt for 1 h. The mixture was concentrated to dryness and the residue was triturated with saturated NaHCO₃ (aq) to give the title compound (35.9 mg, 90.6%) as an off-white solid. LCMS (M+H)⁺ = 502.9.

[0479] Step 23-4. preparation of benzyl *N*-[2-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamoyl)ethyl]carbamate: To a solution of benzyl *N*-[2-({4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamoyl)ethyl]carbamate (35.9 mg, 0.0714 mmol) and 2-fluoro-5-(trifluoromethyl)benzonitrile (27.0 mg, 0.143 mmol) in DMSO (1 mL) in a sealed tube was added DIEA (0.031 mL, 0.18 mmol). The mixture was heated at 130°C for 16 h and purified by reversed-phase CC to give the title compound (41.2 mg, 85.9%) as a light brown solid. LCMS (M+H)⁺ = 672.1.

[0480] Step 23-5. preparation of 3-amino-*N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}propanamide: To benzyl *N*-[2-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamoyl)ethyl]carbamate (41.2 mg, 0.0613 mmol) was added (methylsulfanyl)benzene (0.030 mL, 0.26 mmol) and TFA (0.15 mL, 2.0 mmol). The mixture was heated at 60°C for 0.5 h and purified by reversed-phase CC to give the title compound (27.9 mg, 84.6%) as an off-white solid. LCMS (M+H)⁺ = 538.0.

Example 24: (2*S*)-*N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}-1-methylpyrrolidine-2-carboxamide (cpd 2-11)

[0481]





[0482] Step 24-1. preparation of *tert*-butyl 4-[[[(benzyloxy)carbonyl]amino]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-1-carboxylate: To a solution of *tert*-butyl 4-amino-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-1-carboxylate (237 mg, 0.595 mmol) from Step 23-1 and benzyl 2,5-dioxopyrrolidin-1-yl carbonate (222 mg, 0.892 mmol) in DCM (5 mL) was added DIEA (0.21 mL, 1.2 mmol). The mixture was stirred at rt for 20 min. The mixture was purified by reversed-phase CC to give the title compound (260 mg, 82.2%) as a white solid. LCMS (M+H)⁺ = 532.3.

[0483] Step 24-2. preparation of benzyl *N*-{4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamate bis(2,2,2-trifluoroacetate): To a solution of *tert*-butyl 4-[[[(benzyloxy)carbonyl]amino]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidine-1-carboxylate (260 mg, 0.489 mmol) in DCM (0.6 mL) was added TFA (0.56 mL, 7.3 mmol). The mixture was stirred at rt for 0.5 h. The mixture was concentrated to dryness to give the title compound (323 mg, 100%) as a brown gum. LCMS (M+H)⁺ = 432.3.

[0484] Step 24-3. preparation of benzyl *N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamate: To a solution of benzyl *N*-{4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamate bis(2,2,2-trifluoroacetate) (323 mg, 0.489 mmol) and 2-fluoro-5-(trifluoromethyl)benzonitrile (185 mg, 0.979 mmol) in DMSO (1 mL) in a sealed tube was added DIEA (0.43 mL, 2.4 mmol). The mixture was heated at 130°C for 16 h. The mixture was purified by reversed-phase CC to give the title compound (208 mg, 70.7%) as a brown solid. LCMS (M+H)⁺ = 601.7.

[0485] Step 24-4. preparation of 2-{4-amino-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-1-yl}-5-(trifluoromethyl)benzonitrile: To benzyl *N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl} carbamate (208 mg, 0.346 mmol) was added (methylsulfanyl)benzene (0.050 mL, 0.43 mmol) and TFA (0.25 mL, 3.3 mmol). The mixture was heated at 60°C for 1 h and purified by reversed-phase CC to give the title compound (121 mg, 75.0%) as an off-white solid. LCMS (M+H)⁺ = 467.4.

[0486] Step 24-5. preparation of (2*S*)-*N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}-1-methylpyrrolidine-2-carboxamide: To a solution of (2*S*)-1-methylpyrrolidine-2-carboxylic acid (16.6 mg, 0.129 mmol) and HATU (44.0 mg, 0.193 mmol) in DMF (0.2 mL) was added DIEA (0.034 mL, 0.19 mmol). After stirring at rt for 5 min, 2-{4-amino-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-1-yl}-5-(trifluoromethyl)benzonitrile (30.0 mg, 0.0643 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 10 min and purified by reversed-phase CC to give the title compound (33.8 mg, 91.0%) as a white solid. LCMS (M+H)⁺ = 578.3.


[0487] The following compounds were prepared similarly to **Example 24** with appropriate

substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
2-16	564.1
2-17	578.2

Example 25: *N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)propanamide (cpd 2-5)

[0488]



[0489] Step 25-1. preparation of benzyl *N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamate: To a solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid from Step 7-5 (1.0 g, 2.0 mmol) in toluene (10 mL) was added TEA (0.3 g, 3 mmol) and DPPA (0.7 g, 3 mmol). The mixture was stirred at rt for 1 h. To this was added benzyl alcohol (2 mL) and 4Å MS (100 mg). The mixture was heated at 100°C for 1 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were concentrated to dryness and the residue was purified by silica gel CC to give the title compound (1.0 g, 83%) as a yellow solid. LCMS (M+H)⁺ = 602.2.

[0490] Step 25-2. preparation of 2-(4-amino-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-1-yl)-5-(trifluoromethyl)benzonitrile: To a solution of benzyl *N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamate (0.98 g, 1.6 mmol) in MeOH (15 mL) was added Pd(OH)₂ (100 mg). The mixture was charged with hydrogen (g) and stirred at rt for 1 h. The mixture was filtered and the filtrate was concentrated to dryness to give the title compound (0.74 g, 97%) as a white solid. LCMS (M+H)⁺ = 468.2.

[0491] Step 25-3. preparation of *tert*-butyl *N*-[2-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamoyl)ethyl]-*N*-methylcarbamate: To a solution of 3-[[*tert*-butoxy]carbonyl](methyl)amino}propanoic acid (30 mg, 0.15 mmol) and HATU (49 mg, 0.13 mmol) in DMF (1 mL) was added DIEA (54 mg, 0.42 mmol). After stirring at rt for 5 min, 2-

(4-amino-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-1-yl)-5-(trifluoromethyl)benzonitrile (60 mg, 0.13 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 2 h and purified by reversed-phase CC to give the title compound (60 mg, 72%) as a white solid. LCMS (M+H)⁺ = 653.3.


[0492] Step 25-4, preparation of *N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)propanamide: To a solution of *tert*-butyl *N*-[2-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamoyl)ethyl]-*N*-methylcarbamate (60 mg, 0.11 mmol) in DCM (2 mL) was added TFA (0.6 mL). The mixture was stirred at rt for 1 h. The mixture was concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (42 mg, 83%) as a white solid. LCMS (M+H)⁺ = 553.2.

[0493] The following compounds were prepared similarly to **Example 25** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
2-4	539.2
2-6	565.2
2-8	565.2
2-26	553.3
2-32	565.3

Example 26: *N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(dimethylamino)propanamide (cpd 2-10)

[0494]



[0495] Step 26-1, preparation of *N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(dimethylamino)propanamide formate: To a solution of *N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)propanamide (55 mg, 0.10 mmol) from Step 25-4 in MeOH (1 mL) was added


formaldehyde (10 mg, 0.33 mmol), sodium cyanoborohydride (20 mg, 0.32 mmol) and acetic acid (0.6 mg, 0.01 mmol). The mixture was stirred at rt for 1 h. The mixture was concentrated and purified by reversed-phase CC to give the title compound (27 mg, 44%) as a white solid. LCMS (M+H)⁺ = 567.3.

[0496] The following compounds were prepared similarly to **Example 26** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
2-7	579.2
2-9	579.2

Example 27: (3R)-1-methylpyrrolidin-3-yl N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamate (cpd 2-12)

[0497]




[0498] Step 27-1, preparation of (3R)-1-methylpyrrolidin-3-yl N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamate: To a solution of (3R)-1-methylpyrrolidin-3-ol (7.8 mg, 0.077 mmol) and DIEA (0.056 mL, 0.32 mmol) in DCM (2 mL) at 0°C was added triphosgene (8.0 mg, 0.027 mmol). The mixture was stirred at 0°C for 1 h. To this mixture was added 2-{4-amino-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-1-yl}-5-(trifluoromethyl)benzonitrile (30.0 mg, 0.0643 mmol) from Step 24-4. The mixture was stirred at rt for 0.5 h, concentrated and purified by reversed-phase CC to give the title compound (9.3 mg, 24%) as a light yellow solid. LCMS (M+H)⁺ = 594.5.

[0499] The following compounds were prepared similarly to **Example 27** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
2-13	594.2

Example 28: 2-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-[(2-hydroxyethyl)amino]piperidin-1-yl)-5-(trifluoromethyl)benzonitrile (cpd 2-14)

[0500]




[0501] Step 28-1. preparation of 2-[4-({2-[(*tert*-butyldimethylsilyl)oxy]ethyl}amino)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-1-yl)-5-(trifluoromethyl)benzonitrile: To a solution of 2-(4-amino-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-1-yl)-5-(trifluoromethyl)benzonitrile (50 mg, 0.11 mmol) from Step 25-2 in MeOH (2 mL) was added 2-[(*tert*-butyldimethylsilyl)oxy]-acetaldehyde (30 mg, 0.17 mmol), sodium cyanoborohydride (20 mg, 0.32 mmol) and acetic acid (2.0 mg, 0.033 mmol). The mixture was stirred at 30°C for 2 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness to give the title compound (30 mg, 45%) as a light yellow solid. LCMS (M+H)⁺ = 626.4.

[0502] Step 28-2. preparation of 2-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-[(2-hydroxyethyl)amino]piperidin-1-yl)-5-(trifluoromethyl)benzonitrile formate: To a solution of 2-[4-({2-[(*tert*-butyldimethylsilyl)oxy]ethyl}amino)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-1-yl)-5-(trifluoromethyl)benzonitrile (30 mg, 0.048 mmol) in DCM (3 mL) was added TFA (1 mL). The mixture was stirred at 30°C for 1 h. The mixture was concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (24 mg, 88%) as a brown gum. LCMS (M+H)⁺ = 512.3.

Example 29: 2-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-{[2-(methylamino)ethyl]amino}piperidin-1-yl)-5-(trifluoromethyl)benzonitrile (cpd 2-15)

[0503]






[0504] Step 29-1, preparation of *tert*-butyl *N*-[2-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}amino)ethyl]-*N*-methylcarbamate: To a solution of 2-(4-amino-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-1-yl)-5-(trifluoromethyl)benzonitrile (50 mg, 0.11 mmol) from Step 25-2 in MeOH (2 mL) was added *tert*-butyl *N*-methyl-*N*-(2-oxoethyl)carbamate (30 mg, 0.17 mmol), sodium cyanoborohydride (20 mg, 0.32 mmol) and acetic acid (5.0 mg, 0.083 mmol). The mixture was stirred at 30°C for 2 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness to give the title compound (30 mg, 45%) as a yellow solid. LCMS (M+H)⁺ = 625.4.

[0505] Step 29-2, preparation of 2-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-{[2-(methylamino)ethyl]amino}piperidin-1-yl)-5-(trifluoromethyl)benzonitrile formate: To a solution of *tert*-butyl *N*-[2-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}amino)ethyl]-*N*-methylcarbamate (30 mg, 0.048 mmol) in DCM (3 mL) was added TFA (1 mL). The mixture was stirred at 30°C for 1 h. The mixture was concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (25 mg, 90%) as a white solid. LCMS (M+H)⁺ = 525.2.

Example 30: (3*S*)-1-methylpyrrolidin-3-yl *N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamate (cpd 2-18)

[0506]



[0507] Step 30-1, preparation of (3*S*)-1-methylpyrrolidin-3-yl *N*-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamate formate: To a solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid (50 mg, 0.10 mmol) from Step 7-5 in toluene (1 mL) was added TEA (0.019 mL, 0.14 mmol) and DPPA (33 mg, 0.12 mmol). The mixture was stirred at rt for 1 h. To this was added (3*S*)-1-methylpyrrolidin-3-ol (20 mg, 0.20 mmol) and 4Å MS (100 mg). The mixture was heated at 100°C for 1 h. The mixture was quenched with water and extracted with EtOAc


(3X). The combined organics were concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (28 mg, 43%) as a white solid. LCMS (M+H)⁺ = 595.3.

[0508] The following compounds were prepared similarly to **Example 30** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
2-22	594.4
2-41	553.2
2-42	553.2
2-49	598.2
Compound no.	MS (M+H) ⁺
2-50	598.2
2-53	580.5
2-54	580.5
2-57	568.3

Example 31: 1-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-[(3*R*)-1-methylpyrrolidin-3-yl]urea (cpd 2-19)

[0509]



[0510] Step 31-1. preparation of 1-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-[(3*R*)-1-methylpyrrolidin-3-yl]ureabis(2,2,2-trifluoroacetate): To a solution of 2-(4-amino-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-1-yl)-5-(trifluoromethyl)benzonitrile (50 mg, 0.11 mmol) from Step 25-2 in toluene (1 mL) was added triphosgene (13 mg, 0.044 mmol). The mixture was stirred at rt for 2 h. To this mixture was added (3*R*)-1-methylpyrrolidin-3-amine (22 mg, 0.22 mmol) and TEA (0.045 mL, 0.33 mmol). The mixture was stirred at rt for 1 h. The mixture was concentrated to dryness and purified by


reversed-phase CC to give the title compound (12 mg, 14%) as yellow oil. LCMS (M+H)⁺ = 594.3.

[0511] The following compounds were prepared similarly to **Example 31** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
2-20	594.3
2-43	552.4
2-48	553.1
2-51	597.3
2-52	597.3
Compound no.	MS (M+H) ⁺
2-55	579.2
2-56	579.2
2-58	567.4
2-59	565.5

Example 32: (3R)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl]-3-(methylamino)pyrrolidine-1-carboxamide (cpd 2-21)

[0512]



[0513] Step 32-1, preparation of *tert*-butyl *N*-[(3*R*)-1-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamoyl)pyrrolidin-3-yl]-*N*-methylcarbamate: To a solution of 2-(4-amino-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-1-yl)-5-(trifluoromethyl)benzonitrile (50 mg, 0.11 mmol) from Step 25-2 in toluene (1 mL) was added triphosgene (13 mg, 0.044 mmol). The mixture was stirred at rt for 2 h. To this mixture was added *tert*-butyl *N*-methyl-*N*-[(3*R*)-pyrrolidin-3-yl]carbamate (22 mg, 0.22 mmol) and the resulting mixture was stirred at rt for 2 h. The mixture was quenched with water and extracted

with EtOAc (3X). The combined organics were concentrated to dryness and purified by silica gel CC to give the title compound (33 mg, 44%) as yellow oil. LCMS (M+H)⁺ = 694.5.


[0514] Step 32-2. preparation of (3R)-N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)pyrrolidine-1-carboxamide formate: To a solution of *tert*-butyl *N*-[(3R)-1-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbonyl)pyrrolidin-3-yl]-*N*-methylcarbamate (33 mg, 0.048 mmol) in DCM (1 mL) was added TFA (0.3 mL). The mixture was stirred at rt for 1 h. The mixture was concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (12 mg, 41%) as a white solid. LCMS (M+H)⁺ = 594.3.

[0515] The following compounds were prepared similarly to **Example 32** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
2-23	580.4
2-24	594.3

Example 33: N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-1-methylazetidine-3-carboxamide (cpd 2-25)

[0516]



[0517] Step 33-1. preparation of ethyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate: To a solution of ethyl 6-chloropyridine-3-carboxylate (9.5 g, 48 mmol) in DMF (100 mL) at 0°C was added 60% sodium hydride in mineral oil (3.4 g, 0.14 mol) in portions. The mixture was stirred at 0°C for 0.5 h. To this was added benzylbis(2-bromoethyl)amine (8.3 g, 26 mmol). The mixture was stirred at rt for 2 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness. The residue was purified by silica gel CC to give the title compound (12 g, 70%) as yellow oil. LCMS (M+H)⁺ = 359.2.

[0518] Step 33-2, preparation of ethyl 1-benzyl-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate: To a suspension of ethyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate (4.2 g, 12 mmol) in 1,4-dioxane (50 mL) and water (5 mL) under nitrogen was added potassium carbonate (3.2 g, 23 mmol), (2-ethoxyphenyl)boronic acid (2.9 g, 17 mmol) and Pd(dtbpf)Cl₂ (0.3 g, 0.5 mmol). The mixture was heated at 90°C for 2 h. The mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by silica gel CC to give the title compound (4.0 g, 77%) as a yellow solid. LCMS (M+H)⁺ = 445.2.

[0519] Step 33-3, preparation of ethyl 4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate: To a solution of ethyl 1-benzyl-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (3.5 g, 7.9 mmol) in MeOH (50 mL) was added 10 wt% of Pd on carbon (160 mg) and Pd(OH)₂ (160 mg). The mixture was charged with hydrogen (g) and stirred at rt for 16 h. The mixture was filtered and the filtrate was concentrated to dryness to give the title compound (2.4 g, 86%) as yellow oil. LCMS (M+H)⁺ = 355.2.

[0520] Step 33-4, preparation of ethyl 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate: To a mixture of ethyl 4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (400 mg, 1.13 mmol), 2-bromo-5-chlorobenzonitrile (320 mg, 1.48 mmol), XPhos (33.3 mg, 0.0699 mmol), Pd₂(dba)₃·CHCl₃ (35.0 mg, 0.0338 mmol), and cesium carbonate (735 mg, 2.26 mmol) under nitrogen was added toluene (5 mL). The mixture was heated at 90°C for 2 h. The mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by silica gel CC to give the title compound (290 mg, 52.4%) as yellow oil. LCMS (M+H)⁺ = 490.3.

[0521] Step 33-5, preparation of 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylic acid: To a suspension of ethyl 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (290 mg, 0.59 mmol) in a mixed solvent of EtOH/water (5:1, 6 mL) was added lithium hydroxide (72 mg, 3.0 mmol). The mixture was heated at 60°C for 2 h. The mixture was diluted with water and adjusted to pH 6-7 with 1N HCl (aq). The mixture was extracted with EtOAc (3X) and the combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness to give the title compound (250 mg, 91.4%) as a yellow solid. LCMS (M+H)⁺ = 462.3.

[0522] Step 33-6, preparation of benzyl N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]carbamate: To a solution of 1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylic acid (730 mg, 1.58 mmol) in toluene (10 mL) was added TEA (208 mg, 2.06 mmol) and DPPA (478 mg, 1.74 mmol). The mixture was stirred at rt for 1 h. To this was added benzyl alcohol (513 mg, 4.74 mmol) and 4Å MS (80 mg). The mixture was heated at 100°C for 1 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were concentrated to dryness and the residue was purified by silica gel CC to give the title compound (550 mg, 61.4%) as a yellow solid. LCMS (M+H)⁺ = 567.3.

[0523] Step 33-7, preparation of 2-{4-amino-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-1-yl}-5-chlorobenzonitrile: To benzyl *N*-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]carbamate (550 mg, 0.97 mmol) was added TFA (3 mL). The mixture was heated at 70°C for 5 h. The mixture was concentrated to dryness and triturated with saturated NaHCO₃ (aq) to give the title compound (420 mg, 90%) as a light yellow solid. LCMS (M+H)⁺ = 433.2.


[0524] Step 33-8, preparation of *N*-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-1-methylazetidine-3-carboxamide formate: To a solution of 1-methylazetidine-3-carboxylic acid (18 mg, 0.16 mmol) and HATU (60 mg, 0.16 mmol) in DMF (2 mL) was added DIEA (0.081 mL, 0.46 mmol). After stirring at rt for 5 min, 2-{4-amino-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-1-yl }-5-chlorobenzonitrile (50 mg, 0.12 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 1 h. The mixture was purified by reversed-phase CC to give the title compound (28 mg, 42%) as a white solid. LCMS (M+H)⁺ = 530.3.

[0525] The following compounds were prepared similarly to **Example 33** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
2-28	544.3
2-29	544.3
2-30	532.3
2-31	518.3

Example 34: *N*-[1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-2-(dimethylamino)acetamide (cpd 2-34)

[0526]



[0527] Step 34-1, preparation of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate: To a solution of ethyl 4-[6-(2-

ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (800 mg, 2.26 mmol) from Step 33-3 in DMSO (10 mL) was added 2-fluoro-5-(trifluoromethyl)benzonitrile (428 mg, 2.26 mmol) and DIEA (875 mg, 6.77 mmol). The mixture was heated at 60°C for 1 h. The mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by reversed-phase CC to give the title compound (600 mg, 50.8%) as a yellow solid. LCMS (M+H)⁺ = 524.2.

[0528] Step 34-2, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylic acid: To a suspension of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylate (600 mg, 1.15 mmol) in a mixed solvent of EtOH/water (5:1, 12 mL) was added lithium hydroxide (138 mg, 5.76 mmol). The mixture was heated at 60°C for 1 h. The mixture was concentrated to remove organics and the aqueous residue was adjusted to pH 6-7 with 1N HCl (aq). The solution was extracted with EtOAc (3X) and the combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness to give the title compound (570 mg, 100%) as a yellow solid. LCMS (M+H)⁺ = 496.1.

[0529] Step 34-3, preparation of benzyl N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate: To a solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidine-4-carboxylic acid (570 mg, 1.15 mmol) in toluene (8 mL) was added TEA (163 mg, 1.61 mmol) and DPPA (380 mg, 1.38 mmol). The mixture was stirred at rt for 1 h. To this was added benzyl alcohol (1.6 mL) and 4Å MS (60 mg). The mixture was heated at 100°C for 1 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were concentrated to dryness and the residue was purified by silica gel CC to give the title compound (550 mg, 79.6%) as a yellow solid. LCMS (M+H)⁺ = 601.2.

[0530] Step 34-4, preparation of 2-{4-amino-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-1-yl}-5-(trifluoromethyl)benzonitrile: To benzyl N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate (544 mg, 0.97 mmol) was added TFA (5 mL). The mixture was heated at 70°C for 7 h. The mixture was concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (370 mg, 87.6%) as a yellow solid. LCMS (M+H)⁺ = 467.1.

[0531] Step 34-5, preparation of N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-2-(dimethylamino)acetamide formate: To a solution of 2-(dimethylamino)acetic acid (31 mg, 0.30 mmol) and HATU (65 mg, 0.17 mmol) in DMF (1 mL) was added DIEA (0.091 mL, 0.52 mmol). After stirring at rt for 5 min, 2-{4-amino-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-1-yl}-5-(trifluoromethyl)benzonitrile (50 mg, 0.11 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 1 h. The mixture was purified by reversed-phase CC to give the title compound (40.1 mg, 63.0%) as an off-white solid. LCMS (M+H)⁺ = 552.3.


[0532] The following compounds were prepared similarly to **Example 34** with appropriate

substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
2-27	578.3
2-33	578.3
2-35	564.3
2-36	566.3
2-40	551.2
2-44	523.4
Compound no.	MS (M+H) ⁺
2-45	523.3
2-46	537.5
2-47	537.5
2-60	564.4
2-61	556.2

Example 35: N-[1-(4-chloro-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-1-methylazetidine-3-carboxamide (cpd 2-37)

[0533]



[0534] Step 35-1. preparation of ethyl 1-benzyl-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate: To a suspension of ethyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate (2.0 g, 5.6 mmol) from Step 33-1 in 1,4-dioxane (20 mL) and water (2 mL) under nitrogen was added potassium carbonate (2.3 g, 17 mmol), (2-ethoxypyridin-3-yl)boronic acid (1.4 g, 8.4 mmol) and Pd(dtbpf)Cl₂ (0.15 g, 0.23 mmol). The mixture was heated at 90°C for 2 h. The mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by silica gel CC to give the title compound (2.1 g, 85%) as yellow oil. LCMS (M+H)⁺ = 446.3.

[0535] Step 35-2. preparation of ethyl 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-

carboxylate: To a solution of ethyl 1-benzyl-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate (2.1 g, 4.7 mmol) in MeOH (30 mL) was added 10 wt% of Pd on carbon (200 mg) and Pd(OH)₂ (200 mg). The mixture was charged with hydrogen (g) and stirred at rt for 2 h. The mixture was filtered and the filtrate was concentrated to dryness to give the title compound (1.6 g, 96%) as yellow oil. LCMS (M+H)⁺ = 356.2.

[0536] Step 35-3, preparation of ethyl 1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate: To a mixture of ethyl 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate (1.59 g, 4.47 mmol), 2-bromo-5-chlorobenzonitrile (1.26 g, 5.82 mmol), XPhos (171 mg, 0.359 mmol), Pd₂(dba)₃·CHCl₃ (185 mg, 0.179 mmol), and cesium carbonate (2.92 g, 8.96 mmol) under nitrogen was added toluene (20 mL). The mixture was heated at 100°C for 2 h. The mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by silica gel CC to give the title compound (1.3 g, 59%) as a yellow solid. LCMS (M+H)⁺ = 491.2.

[0537] Step 35-4, preparation of 1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid: To a suspension of ethyl 1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate (1.29 g, 2.63 mmol) in a mixed solvent of EtOH/water (5:1, 18 mL) was added lithium hydroxide (629 mg, 26.3 mmol). The mixture was heated at 60°C for 1 h. The mixture was adjusted to pH 6-7 with 1N HCl (aq) and extracted with EtOAc (3X) and the combined organics were dried over anhydrous Na₂SO₄ and concentrated to dryness to give the title compound (1.1 g, 90%) as a yellow solid. LCMS (M+H)⁺ = 463.3.

[0538] Step 35-5, preparation of benzyl N-[1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl]carbamate: To a solution of 1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid (800 mg, 1.73 mmol) in toluene (10 mL) was added TEA (245 mg, 2.42 mmol) and DPPA (571 mg, 2.07 mmol). The mixture was stirred at rt for 1 h. To this was added benzyl alcohol (748 mg, 6.92 mmol) and 4Å MS (80 mg). The mixture was heated at 100°C for 1 h. The mixture was quenched with water and extracted with EtOAc (3X). The combined organics were concentrated to dryness and the residue was purified by silica gel CC to give the title compound (820 mg, 83.5%) as a yellow solid. LCMS (M+H)⁺ = 568.4.

[0539] Step 35-6, preparation of 2-(4-amino-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-1-yl)-5-chlorobenzonitrile: To benzyl N-[1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl]carbamate (800 mg, 1.41 mmol) was added TFA (10 mL). The mixture was heated at 70°C for 3 h. The mixture was concentrated to dryness and the residue was purified by reversed-phase CC to give the title compound (450 mg, 73.6%) as a yellow solid. LCMS (M+H)⁺ = 434.3.

[0540] Step 35-7, preparation of N-[1-(4-chloro-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-


yl]piperidin-4-yl]-1-methylazetidine-3-carboxamide formate: To a solution of 1-methylazetidine-3-carboxylic acid (13 mg, 0.11 mmol) and HATU (39 mg, 0.10 mmol) in DMF (0.6 mL) was added DIEA (36 mg, 0.28 mmol). After stirring at rt for 5 min, 2-(4-amino-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-1-yl)-5-chlorobenzonitrile (40 mg, 0.092 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 1 h. The mixture was purified by reversed-phase CC to give the title compound (26 mg, 49%) as a light yellow solid. LCMS $(M+H)^+ = 531.3$.

[0541] The following compounds were prepared similarly to **Example 35** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS $(M+H)^+$
2-38	519.3
2-39	545.3

Example 36: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-cyclopropyl-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (Compound 1-180)

[0542]



[0543] Step 36-1, preparation of 3-bromo-2-cyclopropylpyridine : Into a 100-mL vial, was placed 2,3-dibromopyridine (1.5 g, 6.3 mmol), tetrakis(triphenylphosphine)palladium (0) (0.22 g, 0.19 mmol), and THF (10 mL) under N_2 . The resulting solution was treated with cyclopropylzinc(II) bromide (17 mL, 1.5 Eq) in THF. The reaction was stirred at 65 °C for 2h. The reaction was cooled to rt and concentrated. The residue was purified by a silica gel column chromatography to afford the title compound (150 mg, 12%) as yellow solid. LCMS $(M+H)^+ = 197.9$.

[0544] Step 36-2, preparation of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(tributylstannyl)pyridin-3-yl]piperidine-4-carboxylate: Into a 20-mL vial, were placed methyl 4-(6-chloropyridin-3-yl)-1-(2-cyano-4-(trifluoromethyl)phenyl)piperidine-4-carboxylate (200 mg, 0.457 mmol) from step 12-2 and dioxane (2 mL). 1,1,1,2,2,2-Hexabutyldistannane (291 mg, 0.502 mmol), $Pd_2(dba)_3 \cdot CHCl_3$ (23.6 mg, 0.023 mmol), and tricyclohexylphosphonium

tetrafluoroborate (16.8 mg, 0.046 mmol) were sequentially added into the solution under N₂. The resulting solution was stirred at 120 °C for 4 h. LCMS indicated ~57% desired product which was used in the next step directly without further purification.

[0545] Step 36-3. preparation of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-cyclopropyl-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate: To a solution of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(tributylstannyl)pyridin-3-yl]piperidine-4-carboxylate (316 mg, 0.456 mmol) in dioxane (2 mL) was added 3-bromo-2-cyclopropylpyridine (142 mg, 0.717 mmol), dicyclohexyl(2'4'6'-triisopropyl-[1,1'-biphenyl]-2-yl)phosphane (21.8 mg, 0.046 mmol), and Pd₂(dba)₃ CHCl₃ (23.6 mg, 0.023 mmol) at rt under N₂. The resulting solution was stirred at 120 °C for 1 h. The solid ppts were filtered out. The filtrate was concentrated under vacuum. The residue was purified by a silica gel column chromatography to afford the title compound (96 mg, 40%) as yellow solid. LCMS (M+H)⁺ = 520.9.

[0546] Step 36-4. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-cyclopropyl-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid: Into a 50-mL round-bottom flask purged, were placed ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-cyclopropyl-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate (96 mg, 0.18 mmol) and EtOH (2 mL)/water (0.4 mL). The solution was treated with LiOH (31 mg, 1.3 mmol) and then stirred at 60 °C for 1 h. The reaction was cooled to rt and neutralized to pH 6-7 with 1N-HCl. The resulting solution was extracted with ethyl acetate (3x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated. This resulted in the title compound (85 mg, 94%) as yellow solid. LCMS (M+H)⁺ = 492.9.


[0547] Step 36-5. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-cyclopropyl-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide formate: To a solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{2'-cyclopropyl-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid (45 mg, 0.091 mmol) in DMF (0.6 mL) was added HATU (38 mg, 0.10 mmol) and DIEA (35 mg, 0.27 mmol). After stirring for 5 min, the reaction was treated with (S)-1-methylpyrrolidin-3-amine (10 mg, 0.10 mmol) at rt. The resulting solution was stirred at rt for 1h. The crude product was further purified by Prep-HPLC to afford the title compound (25 mg, 44%) as off-white solid. LCMS (M+H)⁺ = 575.2.

[0548] The following compounds were prepared similarly to **Example 36** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-181	585.2
1-182	565.2

Example 37: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-pyrrolidin-3-yl]piperidine-4-carboxamide (cpd: 1-152)

[0549]



[0550] Step 37-1. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxylic acid: To a mixture of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylic acid (1.00 g, 2.44 mmol) from Step 12-3, 1-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole (1.52 g, 7.32 mmol), Pd(amphos)Cl₂ (518 mg, 0.732 mmol), and potassium carbonate (1.01 g, 7.32 mmol) in a sealed tube was added 1,4-dioxane (20 mL) and water (3 mL). Nitrogen (g) was bubbled through and the resulting mixture was heated at 90°C for 2 h. The mixture was quenched with water, acidified with 1N HCl (aq) to pH 2-3 and extracted with EtOAc (2X). The combined organics were dried over anhydrous MgSO₄ and concentrated to dryness. The residue was purified by reversed-phase CC to give the title compound (911 mg, 82.1% yield) as a brown solid. LCMS (M+H)⁺ = 455.4.

[0551] Step 37-2. preparation of tert-butyl (3S)-3-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-amido}pyrrolidine-1-carboxylate: To a solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxylic acid (30.0 mg, 0.0660 mmol) and HATU (35.1 mg, 0.0924 mmol) in DMF (0.2 mL) was added DIEA (0.023 mL, 0.13 mmol). After stirring at rt for 5 min, tert-butyl (3S)-3-aminopyrrolidine-1-carboxylate (19.7 mg, 0.106 mmol) was added to the above HATU-activated solution. The mixture was stirred at rt for 10 min and purified by reversed-phase CC to give the title compound (23.9 mg, 58.1% yield) as an off-white solid. LCMS (M+H)⁺ = 523.3.

[0552] Step 37-3. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-pyrrolidin-3-yl]piperidine-4-carboxamide: To a solution of tert-butyl (3S)-3-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-amido} pyrrolidine-1-carboxylate (23.9 mg, 0.0384 mmol) in DCM (0.05 mL) was added TFA (0.044 mL). The mixture was stirred at rt for 30 min. The mixture was concentrated and the residue was purified by reversed-phase CC to give the title compound (17.3 mg, 86.3% yield) as a light brown solid. LCMS (M+H)⁺ = 523.3.


[0553] The following compounds were prepared similarly to **Example 37** with appropriate substituting reagents and substrates at different steps. Some examples do not require

deprotection in the final step.

Compound no.	MS (M+H) ⁺
1-206	525.4
1-207	539.2
1-211	564.5
1-212	552.0
1-213	550.7
1-218	590.5
1-219	523.6
1-220	537.4
1-221	549.5
1-229	551.5
1-231	537.4
1-233	523.5
1-234	523.3
1-235	523.3
1-236	550.6
1-237	568.3
Compound no.	MS (M+H) ⁺
1-238	568.3
1-245	564.4
1-246	582.3
1-247	582.3
1-248	537.5
1-253	551.4
1-258	509.4
1-259	509.4
1-260	539.4
1-261	539.2
1-262	567.3
1-263	551.4
1-264	567.2
1-265	539.3
1-266	551.5

Example 38: ethyl (3S)-3-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-amido}pyrrolidine-1-carboxylate (cpd: 1-185)

[0554]




[0555] Step 38-1, preparation of ethyl (3S)-3-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-amido}pyrrolidine-1-carboxylate: To a solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-pyrrolidin-3-yl]piperidine-4-carboxamide (7.0 mg, 0.013 mmol) from Step 37-3 in DCM (1 mL) was added DIEA (0.005 mL, 0.027 mmol) and ethyl chloroformate (0.002 mL, 0.020 mmol). The mixture was stirred at rt for 30 min. The mixture was concentrated and purified by reversed-phase CC to give the title compound (5.6 mg, 70% yield) as an off-white solid. LCMS (M+H)⁺ = 595.4.

[0556] The following compounds were prepared similarly to **Example 38** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-185	565.5

Example 39: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyquinolin-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (Compound: 1-189)

[0557]




[0558] Step 39-1, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyquinolin-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: To a mixture of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (30.0 mg, 0.061 mmol) from Step 12-4, (2-methoxyquinolin-3-yl)boronic acid (62.0 mg, 0.300 mmol), Xphos (12.0 mg, 0.024 mmol), potassium carbonate (25.0 mg, 0.180 mmol), and Xphos-Pd-G2 (9.6 mg, 0.012 mmol) in a sealed tube was added 1,4-dioxane (3.0 mL) and water (0.30 mL). Nitrogen (g) was bubbled through and the resulting mixture was heated at 90°C for 3 h. The mixture was quenched with water and extracted with EtOAc (2X). The organic layers were combined, concentrated and purified by reversed-phase CC to give the title compound (27.3 mg, 73% yield). LCMS (M+H)⁺ = 615.2.

[0559] The following compounds were prepared similarly to **Example 39** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-190	587.5
1-191	587.4

Example 40: 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (Compound 1-192)

[0560]



[0561] Step 40-1, preparation of 1-benzyl-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carbonitrile: A solution of 1-benzyl-4-(6-chloro-5-fluoropyridin-3-yl)piperidine-4-carbonitrile (210 mg, 0.637 mmol) from Step 6-3 in 1,4-dioxane (4 mL) and water (0.4 mL) was stirred in a 50 mL round bottom flask purged and maintained with nitrogen (g). 1-Methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole (185 mg, 0.893 mmol), potassium carbonate (264 mg, 1.91 mmol) and RuPhos-Pd-G3 (26.6 mg, 0.0318 mmol) were added. The mixture was heated at 100 °C for 2 h. The mixture was purified by silica gel column

chromatography with EtOAc/petroleum ether (1:2) to give the title compound (230 mg, 96.5% yield). LCMS (M+H)⁺ = 375.0.

[0562] Step 40-2. preparation of 1-benzyl-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylic acid: A solution of 1-benzyl-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carbonitrile (230 mg, 0.614 mmol) in EtOH (6 mL) and water (3 mL) was stirred in a 50 mL round bottom flask purged with nitrogen (g). Potassium hydroxide (345 mg, 6.15 mmol) was added and the mixture was stirred at 100 °C for 48 h. The EtOH was removed and the pH of the solution was adjusted to 6-7 with 3N-HCl. The solution was extracted with DCM/MeOH (10:1) (3x15 mL). The organic layers were washed with brine (2x20 mL), dried over anhydrous sodium sulfate and concentrated to give the title compound (200 mg, 82.8% yield). LC/MS (M+H)⁺ = 394.0.


[0563] Step 40-3. preparation of 1-benzyl-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: A solution of 1-benzyl-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylic acid (230 mg, 0.508 mmol) in DMF (3 mL) was stirred in a 50 mL round bottom flask purged with Nitrogen (g). HATU (232 mg, 0.610 mmol), DIEA (197 mg, 1.52 mmol), and (3S)-1-methylpyrrolidin-3-amine (57.0 mg, 0.569 mmol) were added, and the reaction was stirred at 25 °C for 1 h. The mixture was purified using reversed-phase CC to give the title compound (210 mg, 86.8% yield). LCMS (M+H)⁺ = 476.2.

[0564] Step 40-4. preparation of 4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: A solution of 1-benzyl-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (210 mg, 0.442 mmol) in MeOH (5 mL) was stirred in a 50 mL round bottom flask purged and maintained with Nitrogen (g). Wet 10% Pd/C (20 mg) and Pd(OH)₂ (20.0 mg, 0.014 mmol) were added. The reaction was bubbled with Hydrogen (g) and stirred at 23 °C for 2 h. The solid was filtered out and the mixture was concentrated to give the title compound (130 mg, 76.4% yield). LCMS (M+H)⁺ = 386.2.

[0565] Step 40-5. preparation of 1-[2-fluoro-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: A solution of 4-[5-fluoro-6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (80 mg, 0.21 mmol) in Toluene (1 mL) was stirred in a 10mL vial. Dicesium carbonate (200 mg, 0.61 mmol), 1-bromo-2-fluoro-4-(trifluoromethyl)benzene (100 mg, 0.41 mmol), Pd₂(dba)₃·CHCl₃ (13.0 mg, 0.013 mmol), and Xphos (9.9 mg, 0.021 mmol) were added and the reaction was stirred at 100 °C for 16 h. The mixture was purified by reversed-phase CC to give the title compound (17 mg, 14% yield). LCMS (M+H)⁺ = 548.3.

Example 41: 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (Compound 1-193)

[0566]



[0567] Step 41-1. preparation of 3-chloro-6-(2-ethoxypyridin-3-yl)pyridazine: A mixture of 3,6-dichloropyridazine (3.00 g, 20.0 mmol), (2-ethoxypyridin-3-yl)boronic acid (5.00 g, 30.0 mmol), potassium carbonate (8.00 g, 60.0 mmol), and Pd(dppf)Cl₂ (1.00 g, 2.00 mmol) were added into a 40mL vial. 1,4-Dioxane (30 mL) and water (3 mL) were added and nitrogen (g) was bubbled through the resulting mixture. The reaction mixture was stirred at 90 °C for 2 h. The mixture was concentrated and purified by silica gel column chromatography using EtOAc/Petroleum Ether (1:1) to give the title compound (3.3 g, 70% yield). LCMS (M+H)⁺ = 236.3.

[0568] Step 41-2. preparation of 1-tert-butyl 4-methyl 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidine-1,4-dicarboxylate: A solution of 3-chloro-6-(2-ethoxypyridin-3-yl)pyridazine (2.00 g, 8.00 mmol) and 1-tert-butyl 4-methyl piperidine-1,4-dicarboxylate (3.00 g, 10.0 mmol) in THF (20 mL) was stirred in a 100 mL round bottom flask. Sodium bis(trimethylsilyl)amide (3.00 g, 20.0 mmol) was added dropwise at 0 °C and the reaction was stirred at 25 °C for 1 h. The reaction was quenched with water (100 mL) and extracted with EtOAc (3x100 mL). The organic layers were combined and washed with brine (2x100 mL). The mixture was dried over anhydrous sodium sulfate, concentrated, and purified by silica gel column chromatography using EtOAc/Petroleum Ether (1:1) to give the title compound (2.0 g, 50% yield). LCMS (M+H)⁺ = 443.2.

[0569] Step 41-3. preparation of tert-butyl 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-4-[(3S)-1-methylpyrrolidin-3-yl]carbamoyl]piperidine-1-carboxylate: A solution of 1-tert-butyl 4-methyl 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidine-1,4-dicarboxylate (500 mg, 1.13 mmol) and (3S)-1-methylpyrrolidin-3-amine (170 mg, 1.69 mmol) in THF (3 mL) was stirred in a 50 mL round bottom flask. Lithium bis(trimethylsilyl)amide (945 mg, 5.65 mmol) was added dropwise and the reaction was stirred at 25 °C for 2 h. The solution was quenched with water (50 mL) and extracted with EtOAc (3x50 mL). The organic layers were combined, washed with brine (2x50 mL), dried over anhydrous sodium sulfate, and concentrated. The mixture was purified with silica gel column chromatography using EtOAc/Petroleum Ether (1:1) to give the title compound (340 mg, 58.9% yield). LCMS (M+H)⁺ = 511.3.

[0570] Step 41-4, preparation of 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide trifluoroacetate: A solution of tert-butyl 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-4-[[[(3S)-1-methylpyrrolidin-3-yl]carbonyl]piperidine-1-carboxylate (340 mg, 0.666 mmol) and TFA (2 mL) in DCM (6 mL) was stirred in a 50 mL round bottom flask at 25 °C for 1 h. The mixture was concentrated to give the title compound (340 mg, 88% yield). LCMS (M+H)⁺ = 411.2.


[0571] Step 41-5, preparation of 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: A solution of 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide trifluoroacetate (80 mg, 0.15 mmol) in toluene (3 mL) was stirred in a 8 mL vial under nitrogen (g). Cesium carbonate (150 mg, 0.46 mmol), 1-bromo-2-fluoro-4-(trifluoromethyl)benzene (74 mg, 0.31 mmol), Pd₂(dba)₃·CHCl₃ (14 mg, 0.015 mmol), and Xphos (15 mg, 0.031 mmol) were added and the reaction was stirred at 100 °C for 16 h. The mixture was concentrated and purified by reversed-phase CC to give the title compound (40.2 mg, 33% yield). LCMS (M+H)⁺ = 573.2.

[0572] The following compounds were prepared similarly to **Example 41** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-194	589.2

Example 42: 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide (Compound 1-198)

[0573]



[0574] Step 42-1, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidine-4-carboxamide: A mixture of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide (47.4 mg, 0.0853 mmol) (compound 1-196) and polyoxymethylene (51.2 mg, 1.71 mmol) were added to a 4 mL vial. A mixture of sodium cyanoborohydride (53.6 mg, 0.853 mmol) in MeOH (1 mL) was added and the reaction was


stirred at 23 °C for 4 h. The reaction was quenched with TFA, neutralized with sat-NaHCO₃, and extracted with EtOAc (3x15 mL). The organic layers were combined, concentrated and purified by reversed-phase CC to give the title compound (20.4 mg, 42.0% yield). LCMS (M+H)⁺ = 570.3.

[0575] The following compounds were prepared similarly to **Example 42** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-199	584.2

Example 43: (S)-4-(2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl)-1-(2-fluoro-4-(trifluoromethyl)phenyl)-N-(1-methylpyrrolidin-3-yl)piperidine-4-carboxamide (Compound 1-201)

[0576]



[0577] Step 43-1. preparation of 1-benzyl-4-(2'-ethoxy-3-fluoro- [2,3'-bipyridin]-5-yl)piperidine-4-carbonitrile: To a mixture 1-benzyl-4-(6-chloro-5-fluoropyridin-3-yl)piperidine-4-carbonitrile (300 mg, 910 μmol) from Step 6-3, (2-ethoxypyridin-3-yl)boronic acid (304 mg, 1.82 mmol), Pd(DTBPF)Cl₂ (59.3 mg, 91.0 μmol), and K₂CO₃ (377 mg, 2.73 mmol) in a sealed tube was added 1,4-dioxane (3 mL) and water (0.3 mL). N₂ (g) was bubbled through the resulting mixture. The mixture was then heated at 100°C for 2 h. The mixture was concentrated in vacuo and the crude product was purified by silica gel column chromatography. Pure fractions were combined and concentrated to dryness to give the title compound (330 mg, 87.1 %) as yellow oil. LCMS (M+H)⁺ = 417.2.

[0578] Step 43-2. preparation of 1-benzyl-4-(2'-ethoxy-3-fluoro- [2,3'-bipyridin]-5-yl)piperidine-4-carboxylic acid: To a solution of 1-benzyl-4-(2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl)piperidine-4-carbonitrile (330 mg, 792 μmol) in EtOH (8 mL) was added KOH (445 mg, 7.92 mmol) and water (4.5 mL). The reaction mixture was heated at 100°C for 24 h. The mixture was acidified with 3 N HCl (aq) to make pH 5. The solid was collected by vacuum filtration, washed with water, and dried under high vacuum to give the title compound (320 mg, 92.7 %) as a white

solid. LCMS (M+H)⁺ = 436.2.

[0579] Step 43-3, preparation of (S)-1-benzyl-4-(2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl)-N-(1-methylpyrrolidin-3-yl)piperidine-4-carboxamide: To a solution of 1-benzyl-4-(2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl)piperidine-4-carboxylic acid (320 mg, 735 μ mol) and HATU (335 mg, 882 μ mol) in DMF (3 mL) was added DIEA (0.39 mL, 2.20 mmol). The reaction mixture was stirred at rt for 15 min. To the above HATU-activated solution was added (S)-1-methylpyrrolidin-3-amine (88.3 mg, 882 μ mol) and the reaction was continued to stir at rt for 15 min. The mixture was purified by reversed-phase CC. Pure fractions were combined and concentrated in vacuo to give the title compound (350 mg, 92.1%) as a yellow solid. LCMS (M+H)⁺ = 518.2.


[0580] Step 43-4, preparation of (S)-4-(2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl)-N-(1-methylpyrrolidin-3-yl)piperidine-4-carboxamide: To a solution of (S)-1-benzyl-4-(2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl)-N-(1-methylpyrrolidin-3-yl)piperidine-4-carboxamide (350 mg, 676 μ mol) in MeOH (40 mL) was added Pd(OH)₂ (475 mg, 3.38 mmol) and Pd/C (360 mg, 3.38 mmol). The reaction mixture was stirred at 25 °C for 2 h under H₂ (g). The mixture was filtered and the filtrate was concentrated to dryness to give the title compound (200 mg, 62 %, 90% purity) as a light yellow solid. LCMS (M+H)⁺ = 428.2.

[0581] Step 43-5, preparation of (S)-4-(2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl)-1-(2-fluoro-4-(trifluoromethyl)phenyl)-N-(1-methylpyrrolidin-3-yl)piperidine-4-carboxamide formate: To a mixture of (S)-4-(2'-ethoxy-3-fluoro-[2,3'-bipyridin]-5-yl)-N-(1-methylpyrrolidin-3-yl)piperidine-4-carboxamide (60 mg, 0.14 mmol), 1-bromo-2-fluoro-4-(trifluoromethyl)benzene (68 mg, 0.28 mmol), XPhos (13 mg, 28 μ mol), Pd₂(dba)₃ (13 mg, 14 μ mol), and cesium carbonate (0.14 g, 0.42 mmol) was added toluene (2 mL). The reaction mixture was heated at 120°C for 16 h under nitrogen condition. The mixture was concentrated in vacuo and the crude product was purified by reversed-phase CC. Pure fractions were combined and concentrated in vacuo to give the title compound (18.3 mg, 20 %) as an off-white solid. LCMS (M+H)⁺ = 590.2.

[0582] The following compounds were prepared similarly to **Example 43** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-200	589.2
1-202	605.2
1-203	606.2
1-204	593.3

Example 44: N-(3-aminopropyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide (Compound 1-205)

[0583]

[0584] Step 44-1, preparation of 4-(6-{1-[(tert-butoxy)carbonyl]-1H-pyrrol-2-yl}pyridine-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylic acid: A mixture of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylic acid (150 mg, 0.366 mmol) from Step 12-3, {1-[(tert-butoxy)carbonyl]-1H-pyrrol-2-yl}boronic acid (155 mg, 0.732 mmol), potassium carbonate (152 mg, 1.10 mmol), and Pd(dtbpf)Cl₂ (47.7 mg, 0.0732 mmol) were placed in a sealed tube. 1,4-Dioxane (3 mL) and water (0.3 mL) were added. Nitrogen (g) was bubbled through and the mixture was stirred at 80 °C for 3 h. The mixture was cooled to room temperature and quenched with water. The solution was extracted with EtOAc and the organic layers were combined and concentrated. The mixture was purified with reversed-phase CC to give the title compound (94.1 mg, 47.6% yield). LCMS (M+H)⁺ = 541.6.

[0585] Step 44-2, preparation of methyl 4-(6-{1-[(tert-butoxy)carbonyl]-1H-pyrrol-2-yl}pyridine-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate: A solution of 4-(6-{1-[(tert-butoxy)carbonyl]-1H-pyrrol-2-yl}pyridine-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylic acid (94.1 mg, 0.174 mmol) in toluene (1.5 mL) and MeOH (1 mL) was stirred. (Diazomethyl)trimethylsilane (39.8 mg, 0.348 mmol) was added dropwise and the mixture was stirred at 25 °C for 30 min. The mixture was quenched with three drops of AcOH, concentrated, and neutralized with sat-NaHCO₃ (aq). The solid was collected, washed with water, and dried to give the title compound (73.6 mg, 76.2% yield). LCMS (M+H)⁺ = 555.4.

[0586] Step 44-3, preparation of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylate: A solution of methyl 4-(6-{1-[(tert-butoxy)carbonyl]-1H-pyrrol-2-yl}pyridine-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate (73.6 mg, 0.133 mmol) in DCM (0.15 mL) was stirred. TFA (227 mg, 1.99 mmol) was added and the reaction was stirred at 25 °C for 1 h. The mixture was concentrated and purified by reversed-phase CC to give the title compound (60.3 mg, 100% yield). LCMS (M+H)⁺ = 455.0.

[0587] Step 44-4, preparation of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylate: A solution of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylate (60.3 mg, 0.133 mmol) and 60% sodium hydride in mineral oil (10.6 mg, 0.265 mmol) in DMF (1 mL) was stirred at 25 °C for 30 min. Methyl iodide (37.7 mg, 0.265 mmol) was added and the mixture was stirred at 25 °C for 10 min. The mixture was quenched with ice water and the solid was

collected, rinsed with water, and dried to give the title compound (57.5 mg, 92.5% yield). LCMS $(M+H)^+ = 469.3$.


[0588] Step 44-5, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylic acid: A solution of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylate (57.5 mg, 0.123 mmol) in THF (1.5 mL), MeOH (0.50 mL), and water (0.50 mL) was stirred. Lithium(I) hydroxide monohydrate (51.5 mg, 1.23 mmol) was added and the mixture was stirred at 25 °C for 16 h. The mixture was concentrated, diluted with water, and acidified using 1N-HCl to pH 3. The solid was collected, rinsed with water, and dried to give the title compound (55.8 mg, 100% yield). LCMS $(M+H)^+ = 455.3$.

[0589] Step 44-6, preparation of tert-butyl N-[3-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidin-4-yl}formamido)propyl]carbamate: A solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylic acid (55.8 mg, 0.123 mmol), HATU (65.4 mg, 0.172 mmol), and DIEA (47.6 mg, 0.368 mmol) in DMF (0.2 mL) was stirred at 25 °C for 5 min. tert-Butyl N-(3-aminopropyl)carbamate (34.2 mg, 0.196 mmol) was added and the mixture was stirred at 25 °C for 10 min. The mixture was purified using reversed-phase CC to give the title compound (43.0 mg, 57.3% yield). LCMS $(M+H)^+ = 611.3$.

[0590] Step 44-7, preparation of N-(3-aminopropyl)-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxamide: A solution of tert-butyl N-[3-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidin-4-yl}formamido)propyl]carbamate (43.0 mg, 0.0704 mmol) in DCM (0.08 mL) was stirred. TFA (120 mg, 1.06 mmol) was added and the reaction was stirred at 25 °C for 1 h. The mixture was concentrated and purified by reversed-phase CC to give the title compound (34.9 mg, 97.1% yield). LCMS $(M+H)^+ = 511.5$.

Example 45: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-ethyl-1H-pyrrol-2-yl)pyridine-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (Compound 1-208)

[0591]



[0592] Step 45-1, preparation of methyl 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate: A solution of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylic acid (150 mg, 0.366 mmol) from Step 12-3 in Toluene (1.5 mL) and MeOH (1 mL) was stirred. (Diazomethyl)trimethylsilane (83.6 mg, 0.732 mmol) was added dropwise, and the mixture was stirred at 25 °C for 30 min. The mixture was quenched with three drops of AcOH, concentrated, and neutralized with sat- NaHCO_3 . The solid was collected, washed with water, and dried to give the title compound (150 mg, 96.4% yield). LCMS $(\text{M}+\text{H})^+ = 424.3$.

[0593] Step 45-2, preparation of methyl 4-(6-{1-[(tert-butoxy)carbonyl]-1H-pyrrol-2-yl}pyridine-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate: A mixture of methyl 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate (150 mg, 0.353 mmol), {1-[(tert-butoxy)carbonyl]-1H-pyrrol-2-yl}boronic acid (149 mg, 0.706 mmol), dipotassium carbonate (146 mg, 1.06 mmol), and $\text{Pd}(\text{dtbpf})\text{Cl}_2$ (46.0 mg, 0.0706 mmol) were placed in a sealed tube. 1,4-dioxane (3 mL) and water (0.3 mL) were added, Nitrogen (g) was bubbled through and the mixture was stirred at 80°C for 3 h. The mixture was cooled to room temperature and quenched with water. The solution was extracted with EtOAc, and the organic layers were combined and concentrated. The mixture was purified with reversed-phase CC to give the title compound (70.9 mg, 36.2% yield). LCMS $(\text{M}+\text{H})^+ = 555.4$.

[0594] Step 45-3, preparation of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylate: A solution of methyl 4-(6-{1-[(tert-butoxy)carbonyl]-1H-pyrrol-2-yl}pyridine-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate (70.9 mg, 0.128 mmol) in DCM (0.2 mL) was stirred. TFA (219 mg, 1.92 mmol) was added and the reaction was stirred at 25 °C for 1 h. The mixture was concentrated and purified by reversed-phase CC to give the title compound (59.5 mg, 102% yield). LCMS $(\text{M}+\text{H})^+ = 455.3$.

[0595] Step 45-4, preparation of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-ethyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylate: A solution of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylate (59.5 mg, 0.131 mmol) and 60% sodium hydride (20.9 mg, 0.524 mmol) in mineral oil in DMF (0.5 mL) was stirred at 25 °C for 30 min. Ethyl iodide (24.5 mg, 0.157 mmol) was added and the mixture was stirred at 25 °C for 10 min. The mixture was quenched with ice water and extracted with EtOAc. The organic layers were combined and concentrated, and the mixture was purified by reversed-phase CC to give the title compound (63.2 mg, 100% yield). LCMS $(\text{M}+\text{H})^+ = 483.1$.


[0596] Step 45-5, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-ethyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylic acid: A solution of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-ethyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylate (63.2 mg, 0.131 mmol) in THF (0.9 mL), MeOH (0.3 mL), and water (0.3 mL) was stirred. Lithium(I) hydrate hydroxide (51.5 mg, 1.23 mmol) was added, and the mixture was stirred at 25 °C for 16 h. The mixture was concentrated, diluted with water, and acidified using 1N-HCl to pH 3.

The solid was collected, rinsed with water, and dried to give the title compound (34.1 mg, 55.6% yield). LCMS (M+H)⁺ = 469.3.

[0597] Step 45-6, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-ethyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: A solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-ethyl-1H-pyrrol-2-yl)pyridine-3-yl]piperidine-4-carboxylic acid (34.1 mg, 0.0728 mmol), HATU (38.7 mg, 0.102 mmol), and DIEA (28.2 mg, 0.218 mmol) in DMF (0.3 mL) were stirred at 25 °C for 5 min. (3S)-1-methylpyrrolidin-3-amine (11.7 mg, 0.116 mmol) was added, and the mixture was stirred at 25 °C for 10 min. The mixture was purified using reversed-phase CC to give the title compound (22.6 mg, 56.4% yield). LCMS (M+H)⁺ = 551.5.

Example 46: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (Compound 1-209)

[0598]



[0599] Step 46-1, preparation of 1-benzyl-4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carbonitrile: A solution of 1-benzyl-4-(6-chloro-5-fluoropyridin-3-yl)piperidine-4-carbonitrile (300 mg, 0.910 mmol) from Step 6-3 in 1,4-dioxane (3 mL) and water (0.3 mL) was stirred in a 50 mL round bottom flask, purged and maintained with nitrogen (g). (5-Fluoro-2-methoxypyridin-3-yl)boronic acid (233 mg, 1.36 mmol), potassium carbonate (377 mg, 2.73 mmol), and RuPhos-Pd-G3 (76 mg, 0.091 mmol) were added. The solution was heated at 100°C for 1 h. The mixture was cooled, quenched with water (40mL), and extracted with EtOAc (3x40 mL). The organic layers were combined, dried over anhydrous sodium sulfate, and concentrated. The mixture was purified by silica gel column chromatography with EtOAc/Petroleum Ether (1:1) to give the title compound (300 mg, 78.4% yield). LCMS (M+H)⁺ = 421.2.

[0600] Step 46-2, preparation of 1-benzyl-4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid: A solution of 1-benzyl-4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carbonitrile (300 mg, 0.714 mmol) in EtOH (6 mL) and water (3 mL) was stirred in a 50 mL round bottom flask. Potassium hydroxide (400 mg, 7.13 mmol) was

added, and the mixture was stirred at 100 °C for 24 h. The reaction was quenched with water (10 mL) and the pH of the solution was adjusted to 6-7 with 3*N*-HCl. The solution was extracted with EtOAc (3x50 mL). The organic layers were dried over anhydrous sodium sulfate and concentrated to give the title compound (300 mg, 95.7% yield). LC/MS (M+H)⁺ = 440.2.


[0601] Step 46-3. preparation of 1-benzyl-4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3*S*)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: A solution of 1-benzyl-4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylic acid (300 mg, 0.683 mmol) and HATU (260 mg, 0.684 mmol) in DMF (3 mL) were stirred at 25 °C for 10 min. (3*S*)-1-Methylpyrrolidin-3-amine (82 mg, 0.82 mmol) and DIEA (265 mg, 2.05 mmol) were added and the mixture was stirred at 25 °C for 1 h. The reaction was quenched with water (200 mL) and extracted with EtOAc (3x200 mL). The organic layers were combined, dried over anhydrous sodium sulfate, and concentrated. The mixture was purified by silica gel column chromatography with EtOAc/Petroleum Ether (1: 1) to give the title compound (280 mg, 78.6% yield). LCMS (M+H)⁺ = 522.3.

[0602] Step 46-4. preparation of tert-butyl 4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-4-[(3*S*)-1-methylpyrrolidin-3-yl]carbamoyl]piperidine-1-carboxylate: A suspension of 1-benzyl-4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3*S*)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (200 mg, 0.383 mmol), TEA (120 mg, 1.19 mmol), wet 10%-Pd/C (41 mg), Pd(OH)₂ (54 mg, 0.38 mmol), and di-tert-butyl dicarbonate (251 mg, 1.15 mmol) in MeOH (10 mL) was stirred in a 50 mL round bottom flask. The flask was flushed with nitrogen (g) three times and then flushed with hydrogen (g). The mixture was stirred at 25 °C for 1 h under hydrogen (g). The solution was diluted with MeOH (10 mL) and filtered. The filtrate was concentrated to give the title compound (180 mg, 73% yield). LCMS (M+H)⁺ = 532.3.

[0603] Step 46-5. preparation of 4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3*S*)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: A solution of tert-butyl 4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-4-[(3*S*)-1-methylpyrrolidin-3-yl]carbamoyl]piperidine-1-carboxylate (180 mg, 0.339 mmol) in DCM (3 mL) was stirred. TFA (0.6 mL) was added and the reaction was stirred at 25 °C for 1 h. The mixture was concentrated and purified by reversed-phase CC to give the title compound (60 mg, 38% yield). LCMS (M+H)⁺ = 432.3.


[0604] Step 46-6. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3*S*)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: A solution of 4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3*S*)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (25 mg, 0.058 mmol) in DMSO (1 mL) was stirred in a 8 mL vial. DIEA (22 mg, 0.017 mmol) and 2-fluoro-5-(trifluoromethyl)benzonitrile (22 mg, 0.12 mmol) were added and the mixture was stirred at 70 °C for 1 h. The mixture was purified by reversed-phase CC to give the title compound (4.1 mg, 11% yield). LCMS (M+H)⁺ = 601.2.

Example 47: 4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3*S*)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide

(Compound 1-210)**[0605]**

[0606] Step 47-1, preparation of 4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide: A solution of 4-{3,5'-difluoro-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidine-4-carboxamide (25 mg, 0.058 mmol) from Step 46-5 in toluene (1 mL) was stirred in a 8mL vial, purged and maintained with nitrogen (g). Cesium carbonate (57 mg, 0.17 mmol), 1-bromo-2-fluoro-4-(trifluoromethyl)benzene (28 mg, 0.12 mmol), Pd₂(dba)₃·CHCl₃ (6 mg, 0.006 mmol), and Xphos (10 mg, 0.012 mmol) were added and the reaction was stirred at 100°C for 16 h. The mixture was purified by reversed-phase CC to give the title compound (6.5 mg, 14% yield). LCMS (M+H)⁺ = 594.2.

Example 48: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide (Compound 1-214)

[0607]

[0608] Step 48-1, preparation of ethyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate: A solution of ethyl 2-(6-chloropyridin-3-yl)acetate (3.7 g, 19 mmol) and 18-crown-6 (1.0 g, 3.8 mmol) in DMF (40 mL) was stirred in a 50 mL round bottom flask. The flask was flushed three times with nitrogen (g) and cooled to 0 °C. 60%-NaH (2.2 mg, 55 mmol) in mineral oil was added and the mixture was stirred at 0 °C for 0.5 h. A solution of N-benzyl-2-bromo-N-(2-bromoethyl)ethan-1-amine (7.2 g, 22 mmol) in DMF was added. The mixture was stirred at 20 °C for 2 h and then quenched with sat-NH₄Cl (50 mL). The resulting solution was

extracted with ethyl acetate (3x60 mL). The combined organic layers were washed with brine (3x30 mL), dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:3). This resulted in the title compound (4.8 g, 72%). LCMS (M+H)⁺ = 359.1.

[0609] Step 48-2, preparation of ethyl 4-(6-chloropyridin-3-yl)piperidine-4-carboxylate hydrochloride: A solution of ethyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate (1.8 g, 5.0 mmol) in DCM (20 mL) was stirred in a 50 mL round bottom flask. 1-Chloroethyl carbonochloridate (0.95 g, 6.6 mmol) was added and the mixture was stirred at 25 °C for 1 h. The mixture was concentrated to remove DCM. MeOH (20 mL) was added and the mixture was stirred at 70 °C for 1 h. The mixture was concentrated to give the title compound (1.5 g, 78% yield). LCMS (M+H)⁺ = 269.1.

[0610] Step 48-3, preparation of ethyl 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate: A solution of ethyl 4-(6-chloropyridin-3-yl)piperidine-4-carboxylate hydrochloride (600 mg, 1.6 mmol) in DMSO (6 mL) was stirred in a 20 mL vial. DIEA (1.0 g, 7.7 mmol) and 2-fluoro-5-(trifluoromethyl)benzonitrile (400 mg, 2.1 mmol) were added and the mixture was stirred at 60 °C for 1 h. The solution was quenched with water (30 mL) and extracted with EtOAc (3x40 mL). The organic layers were combined, washed with brine (2x30 mL), dried over anhydrous sodium sulfate, and concentrated. The mixture was purified by silica gel column chromatography EtOAc/Petroleum Ether (1:3) to give the title compound (480 mg, 70% yield). LCMS (M+H)⁺ = 438.2.

[0611] Step 48-4, preparation of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidine-4-carboxylate: A solution of ethyl 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate (200 mg, 0.457 mmol), (2-methoxyphenyl)boronic acid (85 mg, 0.56 mmol), potassium carbonate (190 mg, 1.37 mmol), and Pd(dtbpf)Cl₂ (30 mg, 0.046 mmol) were placed in a sealed tube. 1,4-Dioxane (1 mL) and water (0.1 mL) were added and the mixture was stirred at 100 °C for 2 h. The mixture was cooled to room temperature and concentrated. The mixture was purified with silica gel column chromatography using EtOAc/Petroleum Ether (1:3) to give the title compound (94.1 mg, 47.6% yield). LCMS (M+H)⁺ = 510.5.

[0612] Step 48-5, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidine-4-carboxylic acid: A solution of ethyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidine-4-carboxylate (150 mg, 0.294 mmol) in EtOH (2 mL) and water (0.40 mL) was stirred in a 8 mL vial. Lithium hydroxide (70 mg, 2.9 mmol) was added and the mixture was stirred at 60 °C for 2 h. The mixture was concentrated, diluted with water (20 mL), and neutralized to pH 6-7 using aq. sodium hydrogen sulfate. The mixture was extracted with EtOAc (3x20 mL) and the organic layers were combined, washed with brine (2x20 mL), dried over anhydrous sodium sulfate, and concentrated to give the title compound (130 mg, 91.7% yield). LCMS (M+H)⁺ = 482.3.

[0613] Step 48-6, preparation of tert-butyl N-[2-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidin-4-yl}formamido)ethyl]-N-methylcarbamate: A solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidine-4-carboxylic acid (70 mg, 0.15 mmol), HATU (60 mg, 0.16 mmol), and DIEA (70 mg, 0.54 mmol) in DMF (2 mL) was stirred in an 8 mL vial at 20 °C for 5 min. tert-Butyl N-(2-aminoethyl)-N-methylcarbamate (35 mg, 0.20 mmol) was added and the mixture was stirred at 20 °C for 2 h. The reaction was purified using reversed-phase CC to give the title compound (55 mg, 59% yield). LCMS (M+H)⁺ = 638.3.


[0614] Step 48-7, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]-N-[2-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidin-4-yl}formamido)ethyl]-N-methylcarbamate: A solution of tert-butyl N-[2-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidin-4-yl}formamido)ethyl]-N-methylcarbamate (55 mg, 0.086 mmol) in DCM (3 mL) was stirred in a 50 mL round bottom flask. TFA (1.49 g, 13.1 mmol) was added and the reaction was stirred at 20 °C for 1 h. The mixture was concentrated and purified by reversed-phase CC to give the title compound (44.8 mg, 89.0% yield). LCMS (M+H)⁺ = 538.2.

[0615] The following compounds were prepared similarly to **Example 48** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-215	511.2
1-216	525.2

Example 49: 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]-N-[(2S)-1-methylazetidin-2-yl]methylpiperidine-4-carboxamide (Compound 1-217)

[0616]



[0617] Step 49-1, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidine-4-carboxylic acid: A mixture of 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylic acid (200.0 mg, 0.4881 mmol) from Step 12-3, (2-methoxyphenyl)boronic acid (111.2 mg, 0.7321 mmol), potassium carbonate (202.3 mg, 1.464 mmol), and Pd(amphos)Cl₂ (69.12 mg, 0.09761 mmol) were

placed in a sealed tube. 1,4-Dioxane (3 mL) and water (0.3 mL) were added. Nitrogen (g) was bubbled through and the mixture was stirred at 90 °C for 2 h. The mixture was cooled to room temperature, quenched with water and EtOAc, and acidified to pH 2-3 using 1N-HCl. The solution was extracted with EtOAc (2x), and the organic layers were combined, dried over anhydrous magnesium sulfate, and concentrated. The mixture was purified with reversed-phase CC to give the title compound (113.8 mg, 48.43% yield). LCMS (M+H)⁺ = 482.3.

[0618] Step 49-2, preparation of tert-butyl (2S)-2-[(1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidin-4-yl]formamido)methyl]azetidine-1-carboxylate: A solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidine-4-carboxylic acid (40.0 mg, 0.0831 mmol), HATU (44.2 mg, 0.116 mmol), and DIEA (53.7 mg, 0.415 mmol) in DMF (0.2 mL) was stirred at 25 °C for 5 min. tert-Butyl (2S)-2-(aminomethyl)azetidine-1-carboxylate (24.8 mg, 0.133 mmol) was added and the mixture was stirred at 25 °C for 10 min. The reaction was purified using reversed-phase CC to give the title compound (33.8 mg, 62.6% yield). LCMS (M+H)⁺ = 650.1.

[0619] Step 49-3, preparation of N-[(2S)-azetidin-2-yl]methyl-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidine-4-carboxamide: A solution of tert-butyl (2S)-2-[(1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidin-4-yl]formamido)methyl]azetidine-1-carboxylate (33.8 mg, 0.0520 mmol) in DCM (0.06 mL) was stirred. TFA (89.0 mg, 0.780 mmol) was added and the reaction was stirred at 25 °C for 1 h. The mixture was concentrated and purified by reversed-phase CC to give the title compound (23.8 mg, 83.2% yield). LCMS (M+H)⁺ = 550.7.

[0620] Step 49-4, preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]-N-[(2S)-1-methylazetidin-2-yl]methyl]piperidine-4-carboxamide: A solution of N-[(2S)-azetidin-2-yl]methyl-1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridine-3-yl]piperidine-4-carboxamide (21.4 mg, 0.0389 mmol) in MeOH (3 mL) was stirred. Polyoxymethylene (24.5 mg, 0.0389 mmol) and sodium cyanoborohydride (24.5 mg, 0.389 mmol) were added, and the reaction was stirred at 25 °C for 1.5 h. The reaction was quenched with TFA, neutralized with sat-NaHCO₃, and extracted with EtOAc (2x). The organic layers were combined, concentrated and purified by reversed-phase CC to give the title compound (4.2 mg, 19% yield). LCMS (M+H)⁺ = 564.3.


[0621] The following compounds were prepared similarly to **Example 49** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-232	551.4
1-239	551.5
1-240	551.4
1-241	537.5

Compound no.	MS (M+H) ⁺
1-242	582.4
1-243	582.2
Compound no.	MS (M+H) ⁺
1-244	564.3
1-249	596.4
1-250	596.4
1-251	563.5
1-252	590.4
1-275	555.4

Example 50: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide (Compound 1-223)

[0622]



[0623] Step 50-1. preparation of 1-tert-butyl 4-ethyl 4-(6-chloropyridin-3-yl)piperidine-1,4-dicarboxylate: A solution of ethyl 4-(6-chloropyridin-3-yl)piperidine-4-carboxylate (3.0 g, 11 mmol) from Step 48-2, TEA (3.5 g, 35 mmol) and di-tert-butyl dicarbonate (5.0 g, 23 mmol) in DCM was stirred in a 50 mL round bottom flask, purged with nitrogen (g). The mixture was stirred at 25 °C for 1 h. The mixture was concentrated and purified by silica gel column chromatography using EtOAc/Petroleum Ether (1:3) to give the title compound (3.5 g, 85% yield). LCMS (M+H)⁺ = 369.2.

[0624] Step 50-2. preparation of 1-tert-butyl 4-ethyl 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-1,4-dicarboxylate: A solution of 1-tert-butyl 4-ethyl 4-(6-chloropyridin-3-yl)piperidine-1,4-dicarboxylate (550 mg, 1.49 mmol) in 1,4-dioxane (10 mL) and water (1 mL) was stirred in a 50 mL round bottom flask, purged and maintained with nitrogen (g). (2-Ethoxypyridin-3-yl)boronic acid (500 mg, 2.99 mmol), potassium carbonate (600 mg, 4.34 mmol), and Pd(dtbpf)Cl₂ (80 mg, 0.082 mmol) were added and the mixture was stirred at 100°C for 1 h. The mixture was concentrated and purified with silica gel column chromatography using

EtOAc/Petroleum Ether (1:3) to give the title compound (550 mg, 81.0% yield). LCMS (M+H)⁺ = 456.3.

[0625] Step 50-3, preparation of ethyl 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate trifluoroacetate: A solution of 1-tert-butyl 4-ethyl 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-1,4-dicarboxylate (550.0 mg, 1.207 mmol) in DCM (5 mL) was stirred in a 50 mL round bottom flask. TFA (1 mL) was added and the reaction was stirred at 25 °C for 1 h. The mixture was concentrated to give the title compound (600.0 mg, 95% yield). LCMS (M+H)⁺ = 356.3.

[0626] Step 50-4, preparation of ethyl 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate: A solution of ethyl 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidine-4-carboxylate trifluoroacetate (550 mg, 1.49 mmol) in toluene (10 mL) was stirred in a 50 mL round bottom flask, purged and maintained with nitrogen (g). Cesium carbonate (1.000 g, 3.069 mmol), 1-bromo-2-fluoro-4-(trifluoromethyl)benzene (300 mg, 1.23 mmol), Pd₂(dba)₃·CHCl₃ (60 mg, 0.066 mmol), and XPhos (60 mg, 0.13 mmol) were added and the reaction was stirred at 100 °C for 16 h. The mixture was concentrated and purified by silica gel column chromatography using EtOAc/Petroleum Ether (1:3) to give the title compound (220 mg, 67% yield). LCMS (M+H)⁺ = 534.2.

[0627] Step 50-5, preparation of 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxylic acid: A solution of ethyl 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate (220 mg, 0.425 mmol) in EtOH (10 mL) and water (5 mL) was stirred in a 50 mL round bottom flask. Lithium hydroxide (100 mg, 4.18 mmol) was added, and the mixture was stirred at 60 °C for 1 h. The mixture was concentrated, diluted with water (10 mL), and neutralized to pH 6-7 using aq. sodium hydrogen sulfate. The mixture was extracted with EtOAc (3x20 mL) and the organic layers were combined, washed with brine (2x30 mL), dried over anhydrous sodium sulfate and concentrated to give the title compound (200 mg, 96.1% yield). LCMS (M+H)⁺ = 490.3.

[0628] Step 50-6, preparation of tert-butyl N-{2-[(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidin-4-yl)formamido]ethyl}-N-methylcarbamate: A solution of 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxylic acid (60 mg, 0.12 mmol), HATU (60 mg, 0.16 mmol) and DIEA (60 mg, 0.46 mmol) in DMF (2 mL) was stirred in a 50 mL round bottom flask at 25 °C for 20 min. tert-Butyl N-(2-aminoethyl)-N-methylcarbamate (40 mg, 0.23 mmol) was added, and the mixture was stirred at 25 °C for 1 h. The reaction was purified using reversed-phase CC to give the title compound (40 mg, 51% yield). LCMS (M+H)⁺ = 646.4.

[0629] Step 50-7, preparation of 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]-N-[2-(methylamino)ethyl]piperidine-4-carboxamide ditrifluoroacetate: A solution of tert-butyl N-{2-[(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidin-4-yl)formamido]ethyl}-N-methylcarbamate (40 mg, 0.062 mmol) in DCM (2 mL) was stirred. TFA (1 mL) was added and the reaction was stirred at 25 °C


for 1 h. The mixture was concentrated, and water and acetonitrile were added. The compound was freeze-dried to give the title compound (34.4 mg, 72.0% yield). LCMS (M+H)⁺ = 546.3.

[0630] The following compounds were prepared similarly to **Example 50** with appropriate substituting reagents and substrates at different steps. Some examples do not require deprotection in the final step.

Compound no.	MS (M+H) ⁺
1-222	560.2
1-224	576.2
1-225	562.3
1-226	559.2
1-227	545.3
1-228	575.2
1-230	561.2
1-254	545.3
Compound no.	MS (M+H) ⁺
1-255	561.3
1-256	531.3
1-257	547.3
1-271	546.3
1-272	532.3
1-273	562.3
1-274	548.3
1-276	518.3

Example 51: rac-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S,4R)-4-methoxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide (Compound 1-267)

[0631]



[0632] Step 51-1. preparation of methyl 4-(6-chloropyridin-3-yl)piperidine-4-carboxylate: A solution of methyl 1-benzyl-4-(6-chloropyridin-3-yl)piperidine-4-carboxylate (6.01 g, 17.4 mmol) from Step 7-1 was stirred in DCM (120 mL) in a 250 mL round bottom flask. 1-chloroethyl carbonochloridate (7.47 g, 52.3 mmol) was added and the mixture was stirred at 25 °C for 4 h. The mixture was concentrated to remove DCM. MeOH (120 mL) was added and the mixture was stirred at 70 °C for 1 h. The mixture was concentrated, diluted with water, neutralized with sat-NaHCO₃, and extracted with EtOAc (3x15 mL). The mixture was then extracted with 25% IPA in DCM (10x20 mL). The layers of 25% IPA in DCM were combined, dried over anhydrous sodium sulfate, and concentrated to give the title compound (4.44 g, 100% yield). LCMS (M+H)⁺ = 255.4.

[0633] Step 51-2. preparation of methyl 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate: A solution of methyl 4-(6-chloropyridin-3-yl)piperidine-4-carboxylate (4.44 g, 17.4 mmol) in DMSO (10 mL) was stirred in a sealed tube. 2-fluoro-5-(trifluoromethyl)benzonitrile (4.94 g, 26.1 mmol) and DIEA (6.76 g, 52.3 mmol) were added, and the mixture was stirred at 90 °C for 1.5 h. The mixture was diluted with water and extracted with DCM (4x20 mL). The organic layers were combined, concentrated, and purified by reversed-phase CC to give the title compound (1.71 g, 23.1% yield). LCMS (M+H)⁺ = 424.3.

[0634] Step 51-3. preparation of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxylate: A solution of methyl 4-(6-chloropyridin-3-yl)-1-[2-cyano-4-(trifluoromethyl)phenyl]piperidine-4-carboxylate (1.00 g, 2.37 mmol), 1-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole (2.21 g, 10.5 mmol), Pd(amphos)Cl₂ (754 mg, 1.05 mmol), and potassium carbonate (1.47 g, 10.5 mmol) in a sealed tube was added 1,4-dioxane (25 mL) and water (2.5 mL). Nitrogen (g) was bubbled through and the mixture was stirred at 90 °C for 4 h. The mixture was quenched with water and extracted with EtOAc (3x). The combined organics were dried over anhydrous MgSO₄ and concentrated. The residue was purified by silica gel column chromatography using EtOAc/Hexanes (1:3) to give the title compound (412 mg, 37.1% yield) as a brown solid. LCMS (M+H)⁺ = 469.5.

[0635] Step 51-4. preparation of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxylic acid: A solution of methyl 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxylate (412 mg, 0.879 mmol) in EtOH (5 mL) and water (2.5 mL) was stirred in a 40 mL vial. Lithium(I) hydrate hydroxide (369 mg, 8.79 mmol) was added, and the mixture was stirred at 60 °C for 1 h. The mixture was concentrated, diluted with water (5 mL), and acidified using 1N-HCl to pH 4. The solution was extracted with EtOAc (3x15 mL) and the organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and concentrated to give the title compound (394 mg, 98.6% yield). LCMS (M+H)⁺ = 455.3.

[0636] Step 51-5, preparation of rac-tert-butyl (3R,4S)-3-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-amido}-4-methoxypyrrolidine-1-carboxylate: A solution of 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxylic acid (40.9 mg, 0.0899 mmol), HATU (47.9 mg, 0.126 mmol) and DIEA (20.7 mg, 0.180 mmol) in DMF (0.25 mL) was stirred at 23 °C for 5 min. rac-tert-butyl (3R,4S)-3-amino-4-methoxypyrrolidine-1-carboxylate (27.3 mg, 0.135 mmol) was added and the reaction was stirred at 23 °C for 30 min. The mixture was purified by reversed-phase CC to give the title compound (49.6 mg, 86.4% yield). LCMS (M+H)⁺ = 653.5.

[0637] Step 51-6, preparation of rac-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S,4R)-4-methoxypyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide: A solution of rac-tert-butyl (3R,4S)-3-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-amido}-4-methoxypyrrolidine-1-carboxylate (49.6 mg, 76.0 mmol) in DCM (0.25 mL) was stirred in a 4 mL vial. TFA (125 mg, 1.14 mmol) was added and the reaction was stirred at 23 °C for 6 h. The mixture was concentrated, neutralized with sat-NaHCO₃, and extracted with EtOAc (3x10 mL). The organic layers were combined, dried over anhydrous magnesium sulfate and concentrated to give the title compound (38.5 mg, 91.7% yield). LCMS (M+H)⁺ = 553.0.


[0638] Step 51-7, preparation of rac-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S,4R)-4-methoxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide: A mixture of rac-1-[2-cyano-4-(trifluoromethyl)phenyl]-N-[(3S,4R)-4-methoxypyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidine-4-carboxamide (38.5 mg, 0.0697 mmol) and polyoxymethylene (41.8 mg, 1.39 mmol) were added to a 4 mL vial. A solution of sodium cyanoborohydride (43.8 mg, 0.697 mmol) in MeOH (1 mL) was added and the reaction was stirred at 23 °C for 2 h. The reaction was quenched with TFA, neutralized with sat-NaHCO₃, and extracted with EtOAc (3x15 mL). The organic layers were combined, concentrated and purified by reversed-phase CC to give the title compound (22.1 mg, 56.0% yield). LCMS (M+H)⁺ = 567.4.

[0639] The following compounds were prepared similarly to **Example 51** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-268	567.3
1-269	553.2
1-270	553.2
1-277	569.4
1-278	569.4
1-279	587.3
1-280	587.4

Example 52: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl]-3-(methylamino)propanamide (Compound 2-63)}

[0640]



[0641] Step 52-1, preparation of 1-benzyl-4-(5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl)piperidine-4-carbonitrile: To a mixture 1-benzyl-4-(6-chloro-5-fluoropyridin-3-yl)piperidine-4-carbonitrile (3.00 g, 9.10 mmol) from Step 6-3, (2-methoxyphenyl)boronic acid (1.66 g, 10.9 mmol), Pd(DTBPF)Cl₂ (592.9 mg, 909.8 μmol), and K₂CO₃ (3.77 g, 27.3 mmol) in a sealed tube was added 1,4-dioxane (30 mL) and water (3 mL). N₂ (g) was bubbled through the resulting mixture. The mixture was then heated at 100°C for 2.5 h. The reaction was not complete and thus more (2-methoxyphenyl)boronic acid (0.3 eq), Pd(DTBPF)Cl₂ (0.03 eq), and K₂CO₃ (0.3 eq) were added. The reaction was continued to heat at 100°C for 1.5 h under nitrogen condition. The mixture was quenched with water and extracted with EtOAc (2X). The combined organics were concentrated in vacuo and the crude product was purified by reversed-phased CC. Pure fractions were combined and concentrated to dryness to give the title compound (1.76 g, 48.1%). LCMS (M+H)⁺ = 402.3.

[0642] Step 52-2, preparation of 1-benzyl-4-(5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl)piperidine-4-carboxylic acid: To a solution of 1-benzyl-4-(5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl)piperidine-4-carbonitrile (1.76 g, 4.38 mmol) in EtOH (20 mL) was added KOH (2.46 g, 43.8 mmol) and water (20 mL). The reaction mixture was heated at 100°C for 23 h. The reaction was not complete and thus more KOH (3.3 eq) was added. The reaction was continued to heat at 100°C for 18 h. The mixture was diluted with water and acidified with 1 N HCl (aq) to make pH 5. The solid was collected by vacuum filtration, washed with water, and dried under high vacuum to give the title compound (1.54 g, 83.7%) as a yellow solid. LCMS (M+H)⁺ = 421.4.

[0643] Step 52-3, preparation of 4-(5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl)piperidine-4-carboxylic acid: To a solution of 1-benzyl-4-(5-fluoro-6-(2-methoxyphenyl)pyridin-3-

yl)piperidine-4-carboxylic acid (500 mg, 1.19 mmol) in MeOH (5 mL) was added ammonium formate (750 mg, 11.9 mmol) and Pd/C (540 mg, 238 μ mol). The reaction mixture was heated at 80°C for 1 h. The mixture was filtered through Celite and the filtrate was concentrated to dryness to give the title compound (185.6 mg, 47.2%). LCMS (M+H)⁺ = 331.3.

[0644] Step 52-4, preparation of 1-(2-cyano-4-(trifluoromethyl)phenyl)-4-(5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl)piperidine-4-carboxylic acid: To a solution of 4-(5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl)piperidine-4-carboxylic acid (185.6 mg, 561.8 μ mol) and 2-fluoro-5-(trifluoromethyl)benzonitrile (159.4 mg, 842.7 μ mol) in DMSO (3.5 mL) in a sealed tube was added DIEA (0.29 mL, 1.69 mmol). The mixture was heated at 90°C for 1.5 h. The mixture was purified by reversed-phase CC. Pure fractions were combined and concentrated in vacuo to give the title compound (101.5 mg, 36.2%) as a white solid. LCMS (M+H)⁺ = 500.1.

[0645] Step 52-5, preparation of benzyl (1-(2-cyano-4-(trifluoromethyl)phenyl)-4-(5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl)piperidin-4-yl)carbamate: To a solution of 1-(2-cyano-4-(trifluoromethyl)phenyl)-4-(5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl)piperidine-4-carboxylic acid (101.5 mg, 203.2 μ mol) in toluene (3 mL) under nitrogen was added TEA (51 μ L, 36.8

[0646] μ mol) and diphenyl phosphorazidate (69.9 μ L, 325.1 μ mol). The mixture was stirred at rt for 1 h. To this mixture was then added phenylmethanol (42.3 μ L, 406.4 μ mol) and the resulting mixture was heated at 100°C for 1 h. The reaction was not complete and thus more phenylmethanol (1 eq) was added. The reaction was continued to heat at 100°C for 1 h. The mixture was quenched with water (20 mL) and extracted with EtOAc (2X). The combined organics were concentrated in vacuo and the crude product was purified by reversed-phase CC. Pure fractions were combined and concentrated in vacuo to give the title compound (62.2 mg, 50.6%). LCMS (M+H)⁺ = 605.2.

[0647] Step 52-6, preparation of 2-(4-amino-4-(5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl)piperidin-1-yl)-5-(trifluoromethyl)benzonitrile: To benzyl (1-(2-cyano-4-(trifluoromethyl)phenyl)-4-(5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl)piperidin-4-yl)carbamate (62.2 mg, 103 μ mol) was added TFA (0.50 mL, 6.5 mmol). The mixture was heated at 70°C for 10 h. The mixture was concentrated to dryness and the residue was purified by reversed-phase CC. Pure fractions were combined and concentrated in vacuo to give the title compound (34.6 mg, 71.5%). LCMS (M+H)⁺ = 471.2.


[0648] Step 52-7, preparation of tert-butyl N-[2-({1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbonyl)ethyl]-N-methylcarbamate: To a solution of 3-{{[(tert-butoxy)carbonyl](methyl)amino}propanoic acid (11 mg, 0.054 mmol) and HATU (19 mg, 0.051 mmol) in DMF (0.5 mL) was added DIEA (14 mg, 0.011 mmol). The mixture was stirred at rt for 5 min. To 2-{4-amino-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-1-yl}-5-(trifluoromethyl)benzonitrile (17 mg, 0.036 mmol) was added the above HATU-activated solution. The resulting mixture was stirred at rt for 30 min. The mixture was purified by reversed-phase CC. Pure fractions were combined and concentrated in vacuo to

give the title compound (9.9 mg, 42%). LCMS (M+H)⁺ = 656.4.

[0649] Step 52-8, preparation of N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(methylamino)propanamide: To a solution of tert-butyl N-[2-({ 1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamoyl)ethyl]-N-methylcarbamate (9.9 mg, 15 μ mol) in DCM (0.25 mL) was added TFA (12 μ L, 0.15 mmol). The mixture was heated at 70°C for 10 h. The mixture was concentrated to dryness and the residue was stirred at rt for 1.5 h. The reaction was not complete. More TFA (12 μ L, 0.15 mmol) was added and the reaction was continued to stir at rt for 2 h. The mixture was concentrated in vacuo to dryness and the residue was purified by reversed-phase CC. Pure fractions were combined and concentrated in vacuo to give the title compound (5.7 mg, 68%). LCMS (M+H)⁺ = 556.3.

Example 53: N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamide (Compound 2-62)


[0650]



[0651] Step 53-1, preparation of N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamide: A mixture of N-{1-[2-cyano-4-(trifluoromethyl)phenyl]-4-[5-fluoro-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(methylamino)propanamide (5.7 mg, 0.010 mmol) from Step 52-8 and polyoxymethylene (6.2 mg, 0.21 mmol) were added to a 4 mL vial. A solution of sodium cyanoborohydride (6.4 mg, 0.10 mmol) in MeOH (1 mL) was added and the reaction was stirred at 23 °C for 2 h. The reaction was quenched with TFA, neutralized with sat-NaHCO₃, and extracted with EtOAc (3x15 mL). The organic layers were combined, concentrated and purified by reversed-phase CC to give the title compound (3.1 mg, 53% yield). LCMS (M+H)⁺ = 570.3.

Example 54: N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-1-[2-fluoro-4-(trifluoromethyl)phenyl]piperidine-4-carboxamide (Compound 1-335)

[0652]



[0653] Step 54-1, preparation of methyl 4-(6-(2-ethoxypyridin-3-yl)pyridazin-3-yl)piperidine-4-carboxylate: To a solution of 1-(tert-butyl) 4-methyl 4-(6-(2-ethoxypyridin-3-yl)pyridazin-3-yl)piperidine-1,4-dicarboxylate (100 mg, 226 μ mol) from Step 41-2 in DCM (1 mL) was added TFA (222 mg, 2.26 mmol). The resulting mixture was stirred at rt for 1 h. The mixture was concentrated in vacuo and the residue was triturated with saturated Na_2CO_3 (aq) (30 mL) and extracted with DCM (2X). The combined organics were dried over anhydrous Na_2SO_4 , filtered, and concentrated in vacuo to dryness to give the crude title compound (96 mg, 120% yield) as yellow oil, which was used in the next step without further purification. LCMS $(\text{M}+\text{H})^+ = 343.4$.

[0654] Step 54-2, preparation of methyl 4-(6-(2-ethoxypyridin-3-yl)pyridazin-3-yl)-1-(2-fluoro-4-(trifluoromethyl)phenyl)piperidine-4-carboxylate: To a mixture of methyl 4-(6-(2-ethoxypyridin-3-yl)pyridazin-3-yl)piperidine-4-carboxylate (80 mg, 0.23 mmol), cesium carbonate (0.23 g, 0.71 mmol), 1-bromo-2-fluoro-4-(trifluoromethyl)benzene (85 mg, 0.35 mmol), dicyclohexyl(2',4',6'-triisopropyl-[1,1'-biphenyl]-2-yl)phosphane (11 mg, 23 μ mol), and $\text{Pd}_2(\text{dba})_3$ (21 mg, 23 μ mol) was added toluene (1 mL) under N_2 (g). The resulting suspension was heated at 100°C for 4 h. The reaction mixture was concentrated in vacuo and the residue was purified by reversed-phase CC to give the title compound (72 mg, 61% yield) as a yellow solid. LCMS $(\text{M}+\text{H})^+ = 505.4$.

[0655] Step 54-3, preparation of N-(2-(dimethylamino)ethyl)-4-(6-(2-ethoxypyridin-3-yl)pyridazin-3-yl)-1-(2-fluoro-4-(trifluoromethyl)phenyl)piperidine-4-carboxamide bis(2,2,2-trifluoroacetate): To a solution of methyl 4-(6-(2-ethoxypyridin-3-yl)pyridazin-3-yl)-1-(2-fluoro-4-(trifluoromethyl)phenyl)piperidine-4-carboxylate (67 mg, 0.13 mmol) in THF (0.05 mL) was added N,N-dimethylethylamine (59 mg, 0.66 mmol). The mixture was cooled to -15°C and then to this was added 1M LiHMDS in THF (2.7 mL, 2.7 mmol). The resulting mixture was stirred at -15°C for 1 h. The mixture was quenched with MeOH (2.5 mL) and the crude was further purified by reversed-phase CC to give the title compound (35.8 mg, 34% yield) as yellow oil. LCMS $(\text{M}+\text{H})^+ = 561.3$.

Example A-1: Parenteral Pharmaceutical Composition

[0656] To prepare a parenteral pharmaceutical composition suitable for administration by injection (subcutaneous, intravenous), 1-100 mg of a water-soluble salt of a compound

Formula (I), or a pharmaceutically acceptable salt thereof, is dissolved in sterile water and then mixed with 10 mL of 0.9% sterile saline. A suitable buffer is optionally added as well as optional acid or base to adjust the pH. The mixture is incorporated into a dosage unit form suitable for administration by injection

Example A-2: Oral Solution

[0657] To prepare a pharmaceutical composition for oral delivery, a sufficient amount of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is added to water (with optional solubilizer(s), optional buffer(s) and taste masking excipients) to provide a 20 mg/mL solution.

Example A-3: Oral Tablet

[0658] A tablet is prepared by mixing 20-50% by weight of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, 20-50% by weight of microcrystalline cellulose, 1-10% by weight of low-substituted hydroxypropyl cellulose, and 1-10% by weight of magnesium stearate or other appropriate excipients. Tablets are prepared by direct compression. The total weight of the compressed tablets is maintained at 100 -500 mg.

Example A-4: Oral Capsule

[0659] To prepare a pharmaceutical composition for oral delivery, 10-500 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is mixed with starch or other suitable powder blend. The mixture is incorporated into an oral dosage unit such as a hard gelatin capsule, which is suitable for oral administration.

[0660] In another embodiment, 10-500 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is placed into Size 4 capsule, or size 1 capsule (hypromellose or hard gelatin) and the capsule is closed.

Example A-5: Topical Gel Composition

[0661] To prepare a pharmaceutical topical gel composition, a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is mixed with hydroxypropyl cellulose, propylene glycol, isopropyl myristate and purified alcohol USP. The resulting gel mixture is then incorporated into containers, such as tubes, which are suitable for topical administration.

Example B-1: MC2R assays

Membrane preparation:

[0662] Crude membrane fractions are prepared from CRE-bla-CHO-K1 cells stably expressing hMC2 receptor and hMRAP accessory protein (Thermo Fisher). The cells are grown to 85 - 100% confluence on standard tissue culture dishes in GlutaMax DMEM growth media (Gibco) with following additives: 10% dialyzed FBS (Gemini), 0.1 mM NEAA (Gibco), 25 mM HEPES (Gibco), 5 µg/mL blasticidin (Goldbio), 100 µg/mL zeocin (Invitrogen), 600 µg/mL hygromycin (Goldbio). To prepare membranes, cells are scraped and collected in 1X Dulbecco's phosphate buffered saline (Corning) and then pelleted at 1000 RPM's. The cell pellet is reconstituted in membrane preparation buffer (20 mM HEPES, 6 mM MgCl₂ and 1 mM EGTA, protease inhibitor tablets (Pierce) adjusted to pH 7.4), homogenized using a dounce homogenizer, and the resulting membrane fraction is pelleted by centrifugation at 12,000 RPM's. The membrane pellet is resuspended in membrane preparation buffer, snap frozen and stored at -80°C for later use.

Binding assay for hMC2 antagonists protocol:

[0663] The hMC2 membrane binding assay utilizes the following components: radiolabel [¹²⁵I]ACTH (1-39) Tyr23 (PerkinElmer), wheatgerm agglutinin coated PVT SPA beads (PerkinElmer), crude hMC2R membranes, and compounds. Briefly hMC2R membranes are incubated with SPA beads in binding assay buffer (50mM HEPES, 5mM MgCl₂, 1mM CaCl₂, 0.2%BSA, protease inhibitor tablets (Pierce) adjusted to pH7.4) prior to assay initiation. A dose response of compound (the final concentration of compound are typically 0-10,000nM), SPA membranes, and [¹²⁵I]ACTH (1-39) Tyr23 at a final concentration of 0.2 nM is plated in a 96-well assay plate and allowed to incubate 1.5 hours at room temperature. Assay plates are read using a Top Count NXT and K_i values for compounds are determined using a GraphPad Prism 6 non-linear regression analysis.

[0664] Illustrative binding affinities of selective compounds are described in **Table A**. The potencies are divided into four criteria: + means that K_i is between 1,000 nM and 10,000 nM; ++ means that K_i is between 100 nM and 1,000 nM; +++ means K_i is between 10 nM and 100 nM; ++++ means K_i is below 10 nM.

Table A.

Cpd No.	MC2R binding potency
1-1	++++
1-2	++++
1-3	+++

Cpd No.	MC2R binding potency
1-4	+++
1-5	+++
1-6	+++
1-7	++++
1-8	++++
1-9	++++
1-10	++++
1-11	++++
1-12	++++
1-13	++++
1-14	++++
1-15	+++
1-16	++++
1-17	++++
1-18	++++
1-19	++++
1-20	++++
1-21	++++
1-22	++++
1-23	++++
1-24	++++
1-25	+++
1-26	++++
1-27	+++
1-28	++++
1-29	+++
1-30	++++
1-31	++++
1-32	+++
1-33	++++
1-34	++++
1-35	++++
1-36	++++
1-37	++++

Cpd No.	MC2R binding potency
1-38	++++
1-39	++++
1-40	++++
1-41	++++
1-42	++++
1-43	++++
1-44	++++
1-45	++++
1-46	++++
1-47	++++
1-48	++++
1-51	++++
1-52	++++
1-53	+++
1-54	+++
1-55	++++
1-56	++++
1-57	++++
1-58	++++
1-59	++++
1-60	++++
1-61	++++
1-62	++++
1-63	++++
1-64	++++
1-65	+++
1-66	++++
1-67	+++
1-68	++++
1-69	++++
1-70	+++
1-71	++++
1-72	++++
1-73	++++
1-74	++++

Cpd No.	MC2R binding potency
1-75	++++
1-76	++++
1-77	++++
1-78	++++
1-79	++++
1-80	++++
1-81	++++
1-82	++++
1-83	++++
1-84	+++
1-85	++++
1-86	++++
1-87	++++
1-88	++++
1-89	++++
1-90	++++
1-91	++++
1-92	++++
1-93	++++
1-94	++++
1-95	++++
1-96	++++
1-97	++++
1-98	++++
1-99	++++
1-100	++++
1-101	++++
1-102	++++
1-103	++++
1-104	++++
1-105	+++
1-106	++++
1-107	++++
1-108	++++

Cpd No.	MC2R binding potency
1-109	++++
1-110	++++
1-111	++++
1-112	++++
1-113	++++
1-114	++++
1-115	+++
1-116	++++
1-117	++++
1-118	++++
1-119	++++
1-120	++++
1-121	++++
1-122	++++
1-123	++++
1-124	++++
1-125	++++
1-126	++++
1-127	++++
1-128	++++
1-129	+++
1-130	++++
1-131	++++
1-132	++++
1-133	++++
1-134	++++
1-135	++++
1-136	++++
1-137	+++
1-138	++++
1-139	++++
1-140	++++
1-141	++++
1-142	++++
1-143	++++

Cpd No.	MC2R binding potency
1-144	++++
1-145	++++
1-146	++++
1-147	++++
1-148	+++
1-149	++++
1-150	+++
1-151	++++
1-152	++++
1-153	+++
1-154	++++
1-155	++++
1-156	++++
1-157	+++
1-158	++++
1-159	+++
1-160	++++
1-161	++++
1-162	++++
1-163	++++
1-164	+++
1-165	+++
1-166	++++
1-167	++++
1-168	++++
1-169	++++
1-170	++++
1-171	++++
1-172	++++
1-173	++++
1-174	++++
1-175	++++
1-176	+++
1-177	++

Cpd No.	MC2R binding potency
1-178	+++
1-179	++++
1-180	++++
1-181	+++
1-182	+++
1-183	++++
1-184	++
1-185	++
1-186	++++
1-187	++++
1-188	+++
1-189	++++
1-190	+++
1-191	+++
1-192	++++
1-193	++++
1-194	++++
1-195	++++
1-196	++++
1-197	++++
1-198	++++
1-199	++++
1-200	++++
1-201	++++
1-202	++++
1-203	++++
1-204	++++
1-205	+++
1-206	+++
1-207	++++
1-208	++++
1-209	++++
1-210	++++
1-211	++++
1-212	++++

Cpd No.	MC2R binding potency
1-213	++++
1-214	++++
1-215	++++
1-216	++++
1-217	++++
1-218	++++
1-219	++++
1-220	++++
1-221	+++
1-222	++++
1-223	++++
1-224	++++
1-225	++++
1-226	++++
1-227	++++
1-228	++++
1-229	+++
1-230	++++
1-231	++++
1-232	++++
1-233	+++
1-234	+++
1-235	+++
1-236	++++
1-237	++++
1-238	++++
1-239	+++
1-240	+++
1-241	++++
1-242	++++
1-243	++++
1-244	++++
1-245	++++
1-246	++++

Cpd No.	MC2R binding potency
1-247	++++
1-248	++++
1-249	++++
1-250	++++
1-251	+++
1-252	+++
1-253	++++
1-254	++++
1-255	++++
1-256	++++
1-257	++++
1-258	++++
1-259	+++
1-260	++++
1-261	++++
1-262	++++
1-263	+++
1-264	+++
1-265	++++
1-266	++++
1-267	++++
1-268	++++
1-269	++++
1-270	++++
1-271	++++
1-272	++++
1-273	++++
1-274	++++
1-275	++++
1-276	++++
1-277	++++
1-278	++++
1-279	++
1-280	++
1-281	++++

Cpd No.	MC2R binding potency
1-282	++++
1-283	++++
1-284	++++
1-285	++++
1-286	++++
1-287	++++
1-288	++++
1-289	++++
1-290	++++
1-291	++++
1-292	++++
1-293	++++
1-294	+++
1-295	++++
1-296	++++
1-297	++++
1-298	++++
1-299	++++
1-300	++++
1-301	++++
1-302	++++
1-303	++++
1-304	++++
1-305	++++
1-306	++++
1-307	++++
1-308	++++
1-309	++++
1-310	++++
1-311	++++
1-312	++++
1-313	++++
1-314	++++
1-315	++++

Cpd No.	MC2R binding potency
1-316	++++
1-317	++++
1-318	++++
1-319	++++
1-320	++++
1-321	++++
1-322	++++
1-323	++++
1-324	++++
1-325	++++
1-326	++++
1-327	+++
1-328	+++
1-329	++++
1-330	+++
1-331	++++
1-332	++++
1-333	++++
1-334	++++
1-335	++++
2-1	+++
2-2	+++
2-3	+++
2-4	++++
2-5	++++
2-6	++++
2-7	++++
2-8	++++
2-9	++++
2-10	++++
2-11	+++
2-12	+++
2-13	+++
2-14	+++
2-15	+++

Cpd No.	MC2R binding potency
2-16	+++
2-17	+++
2-18	++++
2-19	++++
2-20	++++
2-21	++++
2-22	++++
2-23	++++
2-24	++++
2-25	++++
2-26	++++
2-27	++++
2-28	++++
2-29	++++
2-30	++++
2-31	++++
2-32	++++
2-33	++++
2-34	++++
2-35	++++
2-36	++++
2-37	++++
2-38	++++
2-39	++++
2-40	++++
2-41	++++
2-42	++++
2-43	++++
2-44	++++
2-45	++++
2-46	++++
2-47	++++
2-48	++++
2-49	++++

Cpd No.	MC2R binding potency
2-50	++++
2-51	++++
2-52	++++
2-53	++++
2-54	++++
2-55	++++
2-56	++++
2-57	++++
2-58	++++
2-59	+++
2-60	++++
2-61	++++
2-62	++++
2-63	++++

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- [US62949854 \[0001\]](#)
- [WO2021133563A \[0004\]](#)
- [WO2021091788A \[0005\]](#)
- [WO2005047253A \[0006\]](#)
- [US2019367481A \[0007\]](#)

Non-patent literature cited in the description

- **SEBHAT, I.K. et al.** Melanocortin subtype 4 receptor agonists: Structure-activity relationships about the 4-alkyl piperidine core *BIOORGANIC & MEDICINAL CHEMISTRY LETTERS*, AMSTERDAM, NL, 2007, vol. 17, 205720-5723 [0008]
- **GANTZ, I.T.M. FONG** *Am. J. Physiol. Endocrinol. Metab.*, 2003, vol. 284, E468-E474 [0023] [0025]
- *Handbook of Pharmaceutical Salts: Properties, Selection and Use* International Union of Pure and Applied Chemistry, Wiley-VCH 2002 0000 [0241]
- **S.M. BERGEL.D. BIGHLEYD.C. MONKHOUSE** *J. Pharm. Sci.*, 1977, vol. 66, 1-19 [0241]
- *Handbook of Pharmaceutical Salts: Properties, Selection and Use* Wiley-VCH/VHCA 2002 0000 [0241]
- **JEAN JACQUESANDRE COLLETSAMUEL H. WILE** *Enantiomers, Racemates and Resolutions* John Wiley And Sons, Inc. 1981 0000 [0251]
- *Design of Prodrugs*, Bundgaard Elsevier 1985 0000 [0253]
- *Method in Enzymology* Academic 1985 0000 vol. 42, 309-396 [0253]
- *Design and Application of Prodrugs* **BUNDGAARD, H.** *A Textbook of Drug Design and Development* 1991 0000 113-191 [0253]
- **BUNDGAARD, H** *Advanced Drug Delivery Review*, 1992, vol. 8, 1-38 [0253]
- *March's Advanced Organic Chemistry* John Wiley and Sons, Inc [0260]
- *Remington: The Science and Practice of Pharmacy* Mack Publishing Company 1995 0000 [0308]
- **HOOVER, JOHN E.** *Remington's Pharmaceutical Sciences* Mack Publishing Co. 1975 0000 [0308]
- *Pharmaceutical Dosage Forms* Marcel Decker 1980 0000 [0308]
- *Pharmaceutical Dosage Forms and Drug Delivery Systems* Lippincott Williams & Wilkins 1999 0000 [0308]

PATENTKRAV

1. Forbindelse med Formel (I), eller et farmaceutisk acceptabelt salt deraf:



Formel (I)

- 5 hvor:
- R^A er usubstitueret eller substitueret phenyl, usubstitueret eller substitueret, monocyklisk 6-leddet heteroaryl, eller usubstitueret eller substitueret, monocyklisk 5-leddet heteroaryl, hvor, hvis R^A er substitueret, R^A så er substitueret med 1, 2, 3 eller 4 grupper valgt blandt R^a , R^b og R^c ;
- 10 R^a , R^b og R^c uafhængigt vælges fra gruppen bestående af hydrogen, halogen, $-OR^4$, $-CN$, $-N(R^4)_2$, $-C(=O)R^7$, usubstitueret eller substitueret C_1 - C_6 -alkyl, usubstitueret eller substitueret C_1 - C_6 -fluoralkyl, usubstitueret eller substitueret C_1 - C_6 -heteroalkyl, usubstitueret eller substitueret C_3 - C_6 -cycloalkyl, usubstitueret eller substitueret C_2 - C_7 heterocycloalkyl, usubstitueret eller substitueret phenyl og usubstitueret eller substitueret heteroaryl, hvor en
- 15 hvilken som helst substitueret gruppe af R^a , R^b og R^c er substitueret med én eller flere R^6 -grupper;
- eller én R^a og én R^b , ved tilstedeværelse på tilstødende atomer af R^A , tages sammen med de intervenerende atomer, der forbinder R^a med R^b , for at danne en 5- til 6-leddet, monocyklisk carbocyklus eller 5- til 6-leddet monocyklisk heterocyklus, hvor carbocyklussen eller
- 20 heterocyklussen er usubstitueret eller substitueret med én eller flere R^6 -grupper;
- hvor, hvis R^a , R^b eller R^c er bundet til N-atomet af et heteroaryl, den så er hydrogen, $-C(=O)R^7$, usubstitueret eller substitueret C_1 - C_6 -alkyl, usubstitueret eller substitueret C_1 - C_6 -fluoralkyl, usubstitueret eller substitueret C_1 - C_6 -heteroalkyl, usubstitueret eller substitueret C_3 - C_6 -cycloalkyl eller usubstitueret eller substitueret C_2 - C_7 heterocycloalkyl;
- 25 R^B er et usubstitueret eller substitueret phenyl eller usubstitueret eller substitueret, monocyklisk 6-leddet heteroaryl, hvor, hvis R^B er substitueret, R^B så er substitueret med 1, 2, 3 eller 4 grupper valgt blandt R^d , R^e og R^f ;
- R^d , R^e og R^f uafhængigt vælges fra gruppen bestående af hydrogen, halogen, $-OR^4$, $-CN$, $-N(R^4)_2$, $-C(=O)R^7$, $-C(=O)N(R^4)_2$, usubstitueret eller substitueret C_1 - C_6 -alkyl, usubstitueret eller
- 30 substitueret C_1 - C_6 -fluoralkyl, usubstitueret eller substitueret C_1 - C_6 -heteroalkyl, usubstitueret eller substitueret C_3 - C_6 -cycloalkyl, usubstitueret eller substitueret C_2 - C_7 heterocycloalkyl,

usubstitueret eller substitueret phenyl og usubstitueret eller substitueret heteroaryl, hvor en hvilken som helst substitueret gruppe af R^d , R^e og R^f er substitueret med én eller flere R^6 -grupper;

hvor, hvis R^d , R^e eller R^f er bundet til N-atomet af et heteroaryl, den så er hydrogen, -C(=O) R^7 , usubstitueret eller substitueret C₁-C₆-alkyl, usubstitueret eller substitueret C₁-C₆-fluoralkyl, usubstitueret eller substitueret C₁-C₆-heteroalkyl, usubstitueret eller substitueret C₃-C₆-cycloalkyl eller usubstitueret eller substitueret C₂-C₇-heterocycloalkyl;

X^1 er CR¹¹ eller N;

X^2 er CR¹² eller N;

X^3 er CR¹³ eller N;

X^4 er CR¹⁴ eller N;

R^{11} , R^{12} , R^{13} og R^{14} hver uafhængigt er hydrogen, halogen, usubstitueret eller substitueret C₁-C₆-alkyl, usubstitueret eller substitueret C₁-C₆-fluoralkyl, usubstitueret eller substitueret C₁-C₆-heteroalkyl, usubstitueret eller substitueret C₃-C₆-cycloalkyl, -CN, -OR⁴, -SR⁴, -CO₂R⁴, -C(=O)N(R⁴)₂ eller -N(R⁴)₂;

M er -(C=O)-, -NR³-, -O-, -S-, -SO₂-, *-NR³-(C=O)-, *-(C=O)-NR³-, *-O-(C=O)NR³-, *-NR³-(C=O)O-, -NR³-(C=O)NR³-, *-NR³(SO₂)-, *-SO₂NR³- eller 5-leddet heterocyklus, hvor * angiver bindingspunktet til R¹;

R¹ er usubstitueret eller substitueret C₁-C₆-alkyl, usubstitueret eller substitueret C₁-C₆-fluoralkyl, usubstitueret eller substitueret C₁-C₆-heteroalkyl, usubstitueret eller substitueret C₃-C₆-cycloalkyl, usubstitueret eller substitueret C₂-C₇-heterocycloalkyl, usubstitueret eller substitueret -(C₁-C₆-alkyl)-(C₃-C₆-cycloalkyl) eller usubstitueret eller substitueret -(C₁-C₆-alkyl)-(C₂-C₇-heterocycloalkyl), hvor en hvilken som helst substitueret gruppe af R¹ er substitueret med én eller flere halogen, usubstitueret eller substitueret C₁-C₆-alkyl, usubstitueret eller substitueret, monocyklisk heterocyklus, -N(R⁴)₂, -OR⁵, -CN, -CO₂R⁵, -C(=O)N(R⁴)₂, -SR⁵, -S(=O)R⁷, -S(=O)₂R⁷, -NR⁴C(=O)R⁵, -NR⁴SO₂R⁷, -SO₂R⁷ eller -SO₂N(R⁴)₂;

hver R³ uafhængigt er hydrogen, usubstitueret eller substitueret C₁-C₆-alkyl, usubstitueret eller substitueret C₃-C₆-cycloalkyl eller usubstitueret eller substitueret C₂-C₇-heterocycloalkyl;

hver R⁴ uafhængigt vælges fra gruppen bestående af hydrogen, usubstitueret eller substitueret C₁-C₆-alkyl, usubstitueret eller substitueret C₃-C₆-cycloalkyl, usubstitueret eller substitueret C₁-C₆-fluoralkyl, usubstitueret eller substitueret aryl og usubstitueret eller substitueret heteroaryl;

eller to R⁴ tages sammen med det nitrogenatom, hvortil de er bundet, for at danne en usubstitueret eller substitueret 3- til 6-leddet monocyklisk heterocyklus;

hver R^5 uafhængigt vælges fra gruppen bestående af hydrogen, substitueret C_1 - C_6 -alkyl, usubstitueret eller substitueret C_3 - C_6 -cycloalkyl, usubstitueret eller substitueret C_1 - C_6 -fluoralkyl, usubstitueret eller substitueret aryl og usubstitueret eller substitueret heteroaryl;

hver R^6 uafhængigt er hydrogen, halogen, usubstitueret eller substitueret C_1 - C_4 -alkyl, usubstitueret eller substitueret C_1 - C_4 -alkoxy, usubstitueret eller substitueret C_1 - C_4 -fluoralkyl, usubstitueret eller substitueret C_1 - C_4 -fluoralkoxy, usubstitueret eller substitueret, monocyklisk carbocyclylus, usubstitueret eller substitueret, monocyklisk heterocyclylus, -CN, -OH, -CO₂R⁵, -CH₂CO₂R⁵, -C(=O)N(R⁴)₂, -C(=O)N(R⁴)OR⁵, -CH₂C(=O)N(R⁴)₂, -N(R⁴)₂, -CH₂N(R⁴)₂, -C(R⁵)₂N(R⁴)₂, -NR⁴C(=O)R⁵, -CH₂NR⁴C(=O)R⁵, -NR⁴C(=O)N(R⁵)₂, -NR⁴C(=O)N(R⁴)₂, C(R⁵)=N(R⁴)-OR⁵, -SR⁵, -S(=O)R⁷, -SO₂R⁷ eller -SO₂N(R⁴)₂; og


hver R^7 uafhængigt vælges fra gruppen bestående af substitueret C_1 - C_6 -alkyl, usubstitueret eller substitueret C_3 - C_6 -cycloalkyl, usubstitueret eller substitueret C_1 - C_6 -fluoralkyl, usubstitueret eller substitueret phenyl og usubstitueret eller substitueret heteroaryl.

2. Forbindelse ifølge krav 1, eller et farmaceutisk acceptabelt salt deraf, hvor:


M er -NR³-, -O-, *-NR³-(C=O)-, *-(C=O)-NR³-, *-O-(C=O)NR³-, *-NR³-(C=O)O- eller -NR³-(C=O)NR³-, hvor * angiver bindingspunktet til R¹; og

hver R³ uafhængigt er hydrogen, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃ eller -CH(CH₃)₂.

3. Forbindelse ifølge krav 1, eller et farmaceutisk acceptabelt salt deraf, hvor forbindelsen har strukturen i Formel (IIa) eller Formel (IIb), eller et farmaceutisk acceptabelt salt deraf:





Formel (IIa)




Formel (IIb)

hvor:




4. Forbindelse ifølge krav 1, eller et farmaceutisk acceptabelt salt deraf, hvor forbindelsen har strukturen i Formel (IVa), eller et farmaceutisk acceptabelt salt, eller solvat deraf:



Formel (IVa)

5 hvor V er CH, CR^a, CR^b eller N.

5. Forbindelse ifølge krav 1, eller et farmaceutisk acceptabelt salt deraf, hvor forbindelsen har strukturen i Formel (VI), eller et farmaceutisk acceptabelt salt, eller solvat deraf:




Formel (VI)


hvor V er CH, CR^a, CR^b eller N; og W er CH, CR^d, CR^e eller N.

6. Forbindelse ifølge krav 5, eller et farmaceutisk acceptabelt salt deraf, hvor forbindelsen har strukturen i Formel (VIa), Formel (VIc), Formel (VIb), Formel (VIId), Formel (VIIb) eller


15 Formel (VIIe), eller et farmaceutisk acceptabelt salt, eller solvat deraf:



Formel (VIa)




Formel (VIc)



Formel (VIb)

Formel (VIId)



Formel (VIIf)

eller

Formel (VIIe);

5 hvor V er CH, CR^a, CR^b eller N; og W er CH, CR^d, CR^e eller N.

7. Forbindelse ifølge et hvilket som helst af kravene 1-6, eller et farmaceutisk acceptabelt salt deraf, hvor:

10 R¹ er usubstitueret eller substitueret C₁-C₆-alkyl, usubstitueret eller substitueret C₁-C₆-heteroalkyl, der indeholder 1 N-atom, hvor en hvilken som helst substitueret gruppe af R¹ er substitueret med én eller flere halogen, usubstitueret eller substitueret C₁-C₆-alkyl, usubstitueret eller substitueret, monocyklisk heterocyclus, -N(R⁴)₂, -OR⁵, -CN, -CO₂R⁵, -C(=O)N(R⁴)₂, -SR⁵, -S(=O)R⁷, -S(=O)₂R⁷, -NR⁴C(=O)R⁵, -NR⁴SO₂R⁷, -SO₂R⁷ eller -SO₂N(R⁴)₂;

15 eller R¹ er usubstitueret eller substitueret, monocyklisk 4-, 5- eller 6-leddet heterocyclus, der indeholder 1-4 N-atomer og 0 eller 1 O- eller S-atomer;

eller R¹ er usubstitueret eller substitueret, brodannende C₂-C₇-heterocycloalkyl, der indeholder 1-2 N-atomer;

20 eller R¹ er usubstitueret eller substitueret -(C₁-C₆-alkyl)-(C₂-C₇-heterocycloalkyl), hvor heterocycloalkylet er et usubstitueret eller substitueret, monocyklisk 4-, 5- eller 6-leddet heterocycloalkyl, der indeholder 1-4 N-atomer og 0 eller 1 O- eller S-atomer.

8. Forbindelse ifølge krav 7, eller et farmaceutisk acceptabelt salt deraf, hvor:

R¹ er usubstitueret eller substitueret C₁-C₆-heteroalkyl, der indeholder 1 N-atom;

eller R¹ er usubstitueret eller substitueret, monocyklisk 4-, 5- eller 6-leddet heterocyclus,

25 der indeholder 1-4 N-atomer og 0 eller 1 O- eller S-atomer;

eller R^1 er usubstitueret eller substitueret $-(C_1-C_6\text{-alkyl})-(C_2-C_7\text{-heterocycloalkyl})$, hvor heterocycloalkylet er et usubstitueret eller substitueret, monocyklisk 4-, 5- eller 6-leddet heterocycloalkyl, der indeholder 1-4 N-atomer og 0 eller 1 O- eller S-atomer.

- 5 9. Forbindelse ifølge et hvilket som helst af kravene 1-8, eller et farmaceutisk acceptabelt salt deraf, hvor:

X^1 er CH, CF eller N;

X^2 er CH, CF eller N;

X^3 er CH, CF eller N; og

- 10 X^4 er CH, CF eller N;

fortrinsvis hvor:

X^1 er N;

X^2 er CH;

X^3 er CH eller CF; og

- 15 X^4 er CH.

10. Forbindelse ifølge et hvilket som helst af kravene 1-9, eller et farmaceutisk acceptabelt salt deraf, hvor:

- 20 R^a vælges fra gruppen bestående af hydrogen, halogen, $-OR^4$, $-CN$, $-N(R^4)_2$, $-C(=O)R^7$, usubstitueret eller substitueret C_1-C_6 -alkyl, usubstitueret eller substitueret C_1-C_6 -fluoralkyl, usubstitueret eller substitueret C_1-C_6 -heteroalkyl og usubstitueret eller substitueret C_3-C_6 -cycloalkyl, hvor en hvilken som helst substitueret gruppe af R^a er substitueret med én eller flere R^6 -grupper;

- 25 R^b og R^c uafhængigt vælges fra gruppen bestående af hydrogen, halogen, $-OR^4$, $-CN$, $-N(R^4)_2$, $-C(=O)R^7$, usubstitueret eller substitueret C_1-C_6 -alkyl, usubstitueret eller substitueret C_1-C_6 -fluoralkyl og usubstitueret eller substitueret C_1-C_6 -heteroalkyl, hvor en hvilken som helst substitueret gruppe af R^b og R^c er substitueret med én eller flere R^6 -grupper;

hvor, hvis R^a , R^b eller R^c er bundet til N-atomet af et heteroaryl, den så er hydrogen, $-C(=O)R^7$ eller usubstitueret eller substitueret C_1-C_6 -alkyl;

- 30 R^d vælges fra gruppen bestående af hydrogen, halogen, $-OR^4$, $-CN$, $-N(R^4)_2$, usubstitueret eller substitueret C_1-C_6 -alkyl, usubstitueret eller substitueret C_1-C_6 -fluoralkyl, usubstitueret eller substitueret C_1-C_6 -heteroalkyl, usubstitueret eller substitueret C_3-C_6 -cycloalkyl, usubstitueret eller substitueret C_2-C_7 -heterocycloalkyl, usubstitueret eller substitueret phenyl, usubstitueret eller substitueret, monocyklisk heteroaryl og usubstitueret eller substitueret bicyklisk heteroaryl,

hvor en hvilken som helst substitueret gruppe af R^d er substitueret med én eller flere R^6 -grupper;
og

R^e og R^f uafhængigt vælges fra gruppen bestående af hydrogen, halogen, $-OR^4$, $-CN$, $-N(R^4)_2$, usubstitueret eller substitueret C_1 - C_6 -alkyl, usubstitueret eller substitueret C_1 - C_6 -fluoralkyl og usubstitueret eller substitueret C_1 - C_6 -heteroalkyl;

hvor, hvis R^d , R^e eller R^f er bundet til N-atomet af et heteroaryl, den så er hydrogen, $-C(=O)R^7$ eller usubstitueret eller substitueret C_1 - C_6 -alkyl;

fortrinsvis hvor:

R^a vælges fra gruppen bestående af hydrogen, F, Cl, Br, $-CN$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-C(O)CH_3$, $-C(O)CH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CH_2OH$, $-CH_2CH_2CN$, $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, $-CH_2NH_2$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2NHCH_3$, $-CH_2CH_2N(CH_3)_2$, cyclopropyl, cyclobutyl, cyclopentyl og cyclohexyl;

R^b og R^c uafhængigt vælges fra gruppen bestående af hydrogen, F, Cl, Br, $-CN$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-C(O)CH_3$, $-C(O)CH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CH_2OH$, $-CH_2CH_2CN$, $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, $-CH_2NH_2$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2NHCH_3$ og $-CH_2CH_2N(CH_3)_2$;

hvor, hvis R^a , R^b eller R^c er bundet til N-atomet af et heteroaryl, den så er hydrogen, $-C(O)CH_3$, $-C(O)CH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CH_2OH$, $-CH_2CH_2CN$, $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, $-CH_2NH_2$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2NHCH_3$ og $-CH_2CH_2N(CH_3)_2$;

R^d vælges fra gruppen bestående af hydrogen, F, Cl, Br, $-CN$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CH_2OH$, $-CH_2CH_2CN$, $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, $-CH_2NH_2$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2NHCH_3$, $-CH_2CH_2N(CH_3)_2$, usubstitueret eller substitueret cyclopropyl, usubstitueret eller substitueret cyclobutyl, usubstitueret eller substitueret cyclopentyl, usubstitueret eller substitueret cyclohexyl, usubstitueret eller substitueret C_2 - C_7 -heterocycloalkyl, usubstitueret eller substitueret phenyl,

usubstitueret eller substitueret, monocyklisk heteroaryl og usubstitueret eller substitueret, bicyklisk heteroaryl, hvor en hvilken som helst substitueret gruppe af R^d er substitueret med én eller flere R^6 -grupper; og

R^e og R^f uafhængigt vælges fra gruppen bestående af hydrogen, F, Cl, Br, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CN$, $-OH$, $-OCH_3$ og $-OCH_2CH_3$;

hvor, hvis R^d , R^e , eller R^f er bundet til N-atomet af et heteroaryl, den så er hydrogen, $-C(O)CH_3$, $-C(O)CH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2CH_2CH(CH_3)_2$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CH_2OH$, $-CH_2CH_2CN$, $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, $-CH_2NH_2$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2NHCH_3$ og $-CH_2CH_2N(CH_3)_2$.

11. Forbindelse ifølge krav 1, eller et farmaceutisk acceptabelt salt deraf, hvor:

R^A er usubstitueret eller substitueret pyridinyl, usubstitueret eller substitueret phenyl eller usubstitueret eller substitueret pyrrolyl, hvor, hvis R^A er substitueret, R^A så er substitueret med 1 eller 2 grupper valgt blandt R^a , R^b og R^c ;

R^B er usubstitueret eller substitueret phenyl eller usubstitueret eller substitueret pyridinyl, hvor, hvis R^B er substitueret, R^B så er substitueret med 1, 2 eller 3 grupper valgt blandt R^d , R^e og R^f ;

X^1 er CH eller N;

X^2 er CH eller N;

X^3 er CR^{13} eller N;

X^4 er CH eller N;

M er $^*-NR^3-(C=O)-$ eller $^*-(C=O)-NR^3-$, hvor * angiver bindingspunktet til R^1 ;

R^{11} , R^{12} , R^{13} , og R^{14} hver uafhængigt er hydrogen, F, Cl, $-CH_3$, CF_3 , $-CN$, $-OR^4$ eller $-N(R^4)_2$;


R^1 er usubstitueret eller substitueret C_1 - C_6 -heteroalkyl, der indeholder 1 N-atom, hvor en hvilken som helst substitueret gruppe af R^1 er substitueret med én eller flere halogen, C_1 - C_4 -alkyl, $-N(R^4)_2$ eller $-OR^5$;

eller R^1 er usubstitueret eller substitueret azetidiny, usubstitueret eller substitueret pyrrolidinyl eller usubstitueret eller substitueret piperidinyl;

eller R^1 er usubstitueret eller substitueret, brodannende C_2 - C_7 -heterocycloalkyl, der indeholder 1-2 N-atomer;


eller R^1 er usubstitueret eller substitueret $-(C_1-C_6\text{-alkyl})-(C_2-C_7\text{-heterocycloalkyl})$, hvor heterocycloalkylet er et usubstitueret eller substitueret, monocyklisk 4-, 5- eller 6-leddet heterocycloalkyl, der indeholder 1-2 N-atomer;

fortrinsvis hvor:



R^a vælges fra gruppen bestående af hydrogen, F, Cl, Br, -CN, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -C(CH₃)₃, -CH₂F, -CHF₂ og -CF₃; og

R^b og R^c uafhængigt vælges fra gruppen bestående af hydrogen, F, Cl, -CH₃, -CH₂F, -CHF₂, -CF₃, -CN og -OCH₃;



R^d vælges fra gruppen bestående af hydrogen, F, Cl, Br, -CN, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -C(CH₃)₃, -CH₂F, -CHF₂ og -CF₃;

R^e og R^f uafhængigt vælges fra gruppen bestående af hydrogen, F, Cl, Br, -CH₃, -CH₂F, -CHF₂, -CF₃, -CN, -OH og -OCH₃;

X^1 er CH eller N;

X^2 er CH eller N;

X^3 er CH, CF eller N;

X^4 er CH eller N;

M er *-NH-(C=O)- eller -(C=O)-NH-, hvor * angiver bindingspunktet til R^1 ;

R^1 er usubstitueret eller substitueret C_1 - C_6 -alkyl, usubstitueret eller substitueret C_1 - C_6 -heteroalkyl, der indeholder 1 N-atom, hvor en hvilken som helst substitueret gruppe af R^1 er substitueret med én eller flere halogen, usubstitueret eller substitueret C_1 - C_6 -alkyl, usubstitueret eller substitueret, monocyklisk heterocyclus, -N(R^4)₂, -OR⁵, -CN, -CO₂R⁵, -C(=O)N(R^4)₂, -SR⁵, -S(=O)R⁷, -S(=O)₂R⁷, -NR⁴C(=O)R⁵, -NR⁴SO₂R⁷, -SO₂R⁷ eller -SO₂N(R^4)₂;

eller R^1 er usubstitueret eller substitueret, monocyklisk 4-, 5- eller 6-leddet heterocyclus, der indeholder 1-4 N-atomer og 0 eller 1 O- eller S-atomer;

eller R¹ er usubstitueret eller substitueret, brodannende C₂-C₇-heterocycloalkyl, der indeholder 1-2 N-atomer;

eller R¹ er usubstitueret eller substitueret -(C₁-C₆-alkyl)-(C₂-C₇-heterocycloalkyl), hvor heterocycloalkylet er et usubstitueret eller substitueret, monocyklisk 4-, 5- eller 6-leddet heterocycloalkyl, der indeholder 1-2 N-atomer.

12. Forbindelse ifølge et hvilket som helst af kravene 1-11, eller et farmaceutisk acceptabelt salt deraf, hvor:

R¹ er usubstitueret eller substitueret C₁-C₆-heteroalkyl, der indeholder 1 N-atom, hvor en hvilken som helst substitueret gruppe af R¹ er substitueret med én eller flere halogen, C₁-C₄-alkyl, -N(R⁴)₂ eller -OR⁵;

eller R¹ er usubstitueret eller substitueret azetidinyll, usubstitueret eller substitueret pyrrolidinyl eller usubstitueret eller substitueret piperidinyl;

eller R¹ er usubstitueret eller substitueret -(C₁-C₆-alkyl)-(C₂-C₇-heterocycloalkyl), hvor C₂-C₇-heterocycloalkyl er et usubstitueret eller substitueret azetidinyll, usubstitueret eller substitueret pyrrolidinyl eller usubstitueret eller substitueret piperidinyl;

fortrinsvis hvor:

R¹ er usubstitueret eller substitueret azetidinyll, usubstitueret eller substitueret pyrrolidinyl eller usubstitueret eller substitueret piperidinyl;

eller R¹ er usubstitueret eller substitueret C₁-C₆-heteroalkyl, der indeholder 1 N-atom, hvor, hvis R¹ er usubstitueret eller substitueret, den så er substitueret med én eller flere halogen, C₁-C₄-alkyl, -N(R⁴)₂ eller -OR⁵.

13. Forbindelse ifølge krav 1, der er:

1-1: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-2: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-pyrrolidin-3-yl]piperidin-4-carboxamid;

1-3: N-(2-aminoethyl)-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-carboxamid;

1-4: N-[(2R)-2-aminopropyl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-carboxamid;

1-5: N-(3-aminopropyl)-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-carboxamid;

1-6: N-[(2S)-2-aminopropyl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-carboxamid;

1-7: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

5 **1-8:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-carboxamid;

1-9: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

10 **1-10:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-pyrrolidin-3-yl]piperidin-4-carboxamid;

1-11: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-12: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-pyrrolidin-3-yl]piperidin-4-carboxamid;

15 **1-13:** N-(2-aminoethyl)-1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-14: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

20 **1-15:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-16: N-[(2S)-2-aminopropyl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-17: N-[(2R)-2-amino-3-hydroxypropyl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

25 **1-18:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(1-methylazetidin-3-yl)methyl]piperidin-4-carboxamid;

1-19: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-(1-methylazetidin-3-yl)piperidin-4-carboxamid;

30 **1-20:** N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-21: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}-N-[(3R)-pyrrolidin-3-yl]piperidin-4-carboxamid;

1-22: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[3,3'-bipyridin]-6-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-23: 1-[2-cyan-6-(trifluormethyl)pyridin-3-yl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-24: 1-[2-cyan-6-(trifluormethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

5 **1-25:** 3-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}piperidin-1-yl)-6-(trifluormethyl)pyridin-2-carboxamid;

1-26: 1-[2-cyan-6-(trifluormethyl)pyridin-3-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

10 **1-27:** 3-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-4-{[(3S)-1-methylpyrrolidin-3-yl]carbamoyl}piperidin-1-yl)-6-(trifluormethyl)pyridin-2-carboxamid;

1-28: N-[(2S)-2-amino-3-hydroxypropyl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-29: 1-[2-chlor-6-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

15 **1-30:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-31: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

20 **1-32:** 1-(2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-33: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]-1-[4-(trifluormethyl)phenyl]piperidin-4-carboxamid;

1-34: 1-(4-chlor-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

25 **1-35:** 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]-1-[5-(trifluormethyl)pyridin-2-yl]piperidin-4-carboxamid;

1-36: 1-(4-acetyl-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

30 **1-37:** 1-[2-cyan-4-(difluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-38: 1-[4-cyan-2-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-39: 1-[2-cyan-4-(trifluormethoxy)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-40: 1-(2,4-dicyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-41: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S,4S)*-4-hydroxy-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

5 **1-42:** 1-[3-cyan-5-(trifluormethyl)pyridin-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-43: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

10 **1-44:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(2R)-3-hydroxy-2-(methylamino)propyl]piperidin-4-carboxamid;

1-45: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2R)-2-(dimethylamino)-3-hydroxypropyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-46: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3R,4R)*-4-hydroxy-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

15 **1-47:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-3-hydroxy-2-(methylamino)propyl]piperidin-4-carboxamid;

1-48: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2S)-2-(dimethylamino)-3-hydroxypropyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

20 **1-51:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-52: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-53: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2,3-dihydro-1-benzofuran-7-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

25 **1-54:** 4-[6-(1-benzofuran-7-yl)pyridin-3-yl]-1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-55: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{6-[2-(hydroxymethyl)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

30 **1-56:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-57: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-58: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-(2-ethoxypyridin-3-yl)pyrazin-2-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-59: 1-[2-cyan-4-(1,1-difluorethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-60: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

5 **1-61:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2,3-difluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-62: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-ethyl-1H-pyrazol-5-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

10 **1-63:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-[6-(2-propoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-64: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(3-fluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-65: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(3,5-difluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

15 **1-66:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-67: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2,2-difluor-2H-1,3-benzodioxol-4-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

20 **1-68:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2,5-difluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-69: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxy-5-fluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-70: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{6-[1-(3-methylbutyl)-1H-pyrazol-5-yl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

25 **1-71:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-{6-[2-(trifluoromethoxy)phenyl]pyridin-3-yl}piperidin-4-carboxamid;

1-72: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxy-3-fluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

30 **1-73:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-indazol-7-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-74: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-fluorophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-75: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)-3-fluorophenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-76: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidin-4-carboxamid;

1-77: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-{6-[2-(trifluormethyl)phenyl]pyridin-3-yl}piperidin-4-carboxamid;

5 **1-78:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-cyanophenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-79: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{6-[2-(methoxymethyl)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

10 **1-80:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(3-methoxythiophen-2-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-81: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(5-fluor-2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-82: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(3-fluor-2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

15 **1-83:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxythiophen-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-84: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{6-[2-(1,1-difluorethyl)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

20 **1-85:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[2'-(difluormethoxy)-[2,3'-bipyridin]-5-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-86: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{5'-fluor-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-87: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(3,5-difluor-2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

25 **1-88:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-89: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-cyclopropoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

30 **1-90:** 1-(4-chlor-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-91: 1-(4-chlor-2-cyanophenyl)-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-92: 1-(4-chlor-2-cyanophenyl)-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidin-4-carboxamid;

1-93: 1-(4-chlor-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidin-4-carboxamid;

1-94: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidin-4-carboxamid;

5 **1-95:** 1-(2,4-dichlorphenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-96: 1-(2-cyan-4-fluorphenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

10 **1-97:** 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-98: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-99: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(5-fluor-2-methylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

15 **1-100:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-101: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(3-fluor-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

20 **1-102:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(5-fluor-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-103: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{5'-fluor-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-104: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethylphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

25 **1-105:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(4-methylpyrimidin-5-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-106: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

30 **1-107:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[2-(2-ethoxypyridin-3-yl)pyrimidin-5-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-108: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-109: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-110: 1-[2-cyan-6-(trifluormethyl)pyridin-3-yl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-111: 1-(4-chlor-2-cyan-6-fluorphenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

5 **1-112:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-[6-(2-methylthiophen-3-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-113: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-cyclopropylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

10 **1-114:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{6-[2-(difluormethoxy)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-115: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-116: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methylfuran-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

15 **1-117:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-[6-(2-propylphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-118: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethyl-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

20 **1-119:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxythiophen-3-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-120: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methylfuran-3-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-121: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-cyclopropylphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

25 **1-122:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]-4-[6-(2-methylthiophen-3-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-123: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluorpyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

30 **1-124:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluorpyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-125: 1-[2-cyan-6-(trifluormethyl)pyridin-3-yl]-4-[6-(2-ethoxyphenyl)-5-fluorpyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-126: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-127: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-128: 1-[2-cyan-6-(trifluormethyl)pyridin-3-yl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

5 **1-129:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(3-fluor-2-hydroxyphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-130: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxy-5-methylphenyl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

10 **1-131:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{3-fluor-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-132: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{3-fluor-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-133: 1-[2-cyan-6-(trifluormethyl)pyridin-3-yl]-4-{3-fluor-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

15 **1-134:** 1-(4-chlor-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-135: 1-(4-chlor-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-N-(1-methylazetidin-3-yl)piperidin-4-carboxamid;

20 **1-136:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-N-(1-methylazetidin-3-yl)piperidin-4-carboxamid;

1-137: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{6-[2-(cyanomethyl)phenyl]pyridin-3-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-138: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]-4-[6-(4-methylthiophen-3-yl)pyridin-3-yl]piperidin-4-carboxamid;

25 **1-139:** 1-(4-chlor-2-cyanophenyl)-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-140: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-(1-methylazetidin-3-yl)piperidin-4-carboxamid;

30 **1-141:** 1-[2-chlor-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-142: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-143: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]-4-[6-(1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-144: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-145: 4-[6-(2-ethoxyphenyl)pyridin-3-yl]-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

5 **1-146:** 4-[6-(2-acetylthiophen-3-yl)pyridin-3-yl]-1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-147: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethylthiophen-3-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

10 **1-148:** 4-[6-(2-cyan-3-fluorphenyl)pyridin-3-yl]-1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-149: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(3-methyl-1,2-oxazol-4-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-150: 1-[3-chlor-5-(trifluormethyl)pyridin-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

15 **1-151:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{6-[2-(difluormethyl)phenyl]pyridin-3-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-152: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-pyrrolidin-3-yl]piperidin-4-carboxamid;

20 **1-153:** 4-[6-(5-cyan-1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-154: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-{5'-fluor-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-155: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(5-fluor-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

25 **1-156:** 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(2-ethylphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-157: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(5-methyl-1,2-oxazol-4-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

30 **1-158:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-cyclopropyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-159: 4-[4-(2-ethoxypyridin-3-yl)phenyl]-1-[2-methoxy-4-(trifluormethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-160: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-161: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-162: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

5 **1-163:** 1-(4-chlor-2-cyanophenyl)-4-[5-fluor-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-164: 4-[4-(2-ethoxypyridin-3-yl)phenyl]-1-[2-methyl-4-(trifluormethyl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

10 **1-165:** 1-[3-chlor-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-166: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-167: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(2-methoxythiophen-3-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

15 **1-168:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-169: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

20 **1-170:** 1-(4-chlor-2-cyanophenyl)-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-171: 1-[2-fluor-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-172: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-{2'-ethyl-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

25 **1-173:** 1-[2-fluor-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-174: 4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

30 **1-175:** 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-176: 4-{[2,2'-bipyridin]-5-yl}-1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-177: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{3'-methyl-[2,2'-bipyridin]-5-yl}-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-178: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{3'-methoxy-[2,2'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-179: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

5 **1-180:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-cyclopropyl-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-181: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[2'-(difluormethyl)-[2,3'-bipyridin]-5-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

10 **1-182:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(5-cyclopropyl-1,3-oxazol-4-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-183: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-184: ethyl (3S)-3-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-amido}pyrrolidin-1-carboxylat;

15 **1-185:** N-[(3S)-1-acetylpyrrolidin-3-yl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-186: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

20 **1-187:** N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-188: N-[(3R)-1-azabicyclo[2.2.2]octan-3-yl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-189: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyquinolin-3-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

25 **1-190:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-indol-7-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-191: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-indol-2-yl)pyridin-3-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

30 **1-192:** 1-[2-fluor-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-193: 4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-194: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-195: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(5-fluor-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-196: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

5 **1-197:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]-N-[3-(methylamino)propyl]piperidin-4-carboxamid;

1-198: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

10 **1-199:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[3-(dimethylamino)propyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-200: 4-[6-(2-ethoxyphenyl)-5-fluorpyridin-3-yl]-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-201: 4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

15 **1-202:** 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluorpyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-203: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

20 **1-204:** 1-[2-fluor-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(5-fluor-2-methoxyphenyl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-205: N-(3-aminopropyl)-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-206: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[3-(methylamino)propyl]piperidin-4-carboxamid;

25 **1-207:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[3-(dimethylamino)propyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-208: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-ethyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

30 **1-209:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{3,5'-difluor-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-210: 4-{3,5'-difluor-2'-methoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[(3S)-1-methylpyrrolidin-3-yl]piperidin-4-carboxamid;

1-211: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(1-methylazetidin-3-yl)methyl]piperidin-4-carboxamid;

1-212: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-213: N-[[2S]-azetidin-2-yl]methyl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

5 **1-214:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-215: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

10 **1-216:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-217: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[[2S]-1-methylazetidin-2-yl]methyl]piperidin-4-carboxamid;

1-218: N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

15 **1-219:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-(1-methylazetidin-3-yl)piperidin-4-carboxamid;

1-220: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1-methylazetidin-3-yl)methyl]piperidin-4-carboxamid;

20 **1-221:** N-{1-azabicyclo[2.2.1]heptan-4-yl}-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-222: N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]piperidin-4-carboxamid;

1-223: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

25 **1-224:** 1-[2-chlor-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-225: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

30 **1-226:** N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-1-[2-fluor-4-(trifluormethyl)phenyl]piperidin-4-carboxamid;

1-227: 4-[6-(2-ethoxyphenyl)pyridin-3-yl]-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-228: 1-[2-chlor-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-229: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[[2-(2S)-1-methylpyrrolidin-2-yl]methyl]piperidin-4-carboxamid;

1-230: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

5 **1-231:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[[2-(2R)-pyrrolidin-2-yl]methyl]piperidin-4-carboxamid;

1-232: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[[2-(2R)-1-methylpyrrolidin-2-yl]methyl]piperidin-4-carboxamid;

10 **1-233:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1S,3S)-3-aminocyclobutyl]piperidin-4-carboxamid;

1-234: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1R,3R)-3-aminocyclobutyl]piperidin-4-carboxamid;

1-235: N-[[2-(2S)-azetidin-2-yl]methyl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

15 **1-236:** N-[[2-(2R)-azetidin-2-yl]methyl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-237: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R,4S)-4-fluorpyrrolidin-3-yl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

20 **1-238:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R,4R)-4-fluorpyrrolidin-3-yl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-239: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1S,3S)-3-(dimethylamino)cyclobutyl]piperidin-4-carboxamid;

1-240: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1R,3R)-3-(dimethylamino)cyclobutyl]piperidin-4-carboxamid;

25 **1-241:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[[2-(2S)-1-methylazetidin-2-yl]methyl]piperidin-4-carboxamid;

1-242: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R,4S)-4-fluor-1-methylpyrrolidin-3-yl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

30 **1-243:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R,4R)-4-fluor-1-methylpyrrolidin-3-yl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-244: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[[2-(2R)-1-methylazetidin-2-yl]methyl]piperidin-4-carboxamid;

1-245: N-[2-(azetidin-1-yl)ethyl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-246: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-{[(2S,4S)-4-fluorpyrrolidin-2-yl]methyl}-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-247: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-{[(2S,4R)-4-fluorpyrrolidin-2-yl]methyl}-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

5 **1-248:** N-[2-(azetidin-1-yl)ethyl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-249: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-{[(2S,4S)-4-fluor-1-methylpyrrolidin-2-yl]methyl}-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

10 **1-250:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-{[(2S,4R)-4-fluor-1-methylpyrrolidin-2-yl]methyl}-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-251: rac-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(1S,2S,4R)-7-methyl-7-azabicyclo[2.2.1]heptan-2-yl]piperidin-4-carboxamid;

1-252: rac-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[(1S,2S,4R)-7-methyl-7-azabicyclo[2.2.1]heptan-2-yl]piperidin-4-carboxamid;

15 **1-253:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3S)-1,3-dimethylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-254: N-[2-(dimethylamino)ethyl]-1-[2-fluor-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

20 **1-255:** 1-[2-chlor-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-256: 1-[2-fluor-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-257: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

25 **1-258:** rac-N-[(1R,2S)-2-aminocyclopropyl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-259: rac-N-[(1R,2R)-2-aminocyclopropyl]-1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

30 **1-260:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2R)-1-(dimethylamino)propan-2-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-261: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-262: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-{3-[(dimethylamino)methyl]oxetan-3-yl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-263: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-{[1-(dimethylamino)cyclopropyl]methyl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-264: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3S,4S)-4-(dimethylamino)oxolan-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-265: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-{2-[ethyl(methyl)amino]ethyl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-266: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-{2-[cyclopropyl(methyl)amino]ethyl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-267: rac-1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3S,4R)-4-methoxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-268: rac-1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R,4R)-4-methoxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-269: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3R,4R)-4-hydroxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-270: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(3S,4S)-4-hydroxy-1-methylpyrrolidin-3-yl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-271: N-[2-(dimethylamino)ethyl]-1-[2-fluor-4-(trifluormethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-272: 1-[2-fluor-4-(trifluormethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-273: 1-[2-chlor-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-274: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-275: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2R)-2-(dimethylamino)-3-hydroxypropyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-276: N-[2-(dimethylamino)ethyl]-1-[2-fluor-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-277: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-{[(2S,4S)-4-fluor-1-methylpyrrolidin-2-yl]methyl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-278: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-{[(2S,4R)-4-fluor-1-methylpyrrolidin-2-yl]methyl}-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-279: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2S)-4,4-difluor-1-methylpyrrolidin-2-yl]methyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

1-280: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2R)-4,4-difluor-1-methylpyrrolidin-2-yl]methyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

5 **1-281:** 1-[2-fluor-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-282: 1-[2-chlor-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

10 **1-283:** 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-284: N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]piperidin-4-carboxamid;

1-285: N-[2-(dimethylamino)ethyl]-1-[2-fluor-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

15 **1-286:** 1-[2-chlor-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-carboxamid;

1-287: 4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

20 **1-288:** 1-[2-chlor-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-289: N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxyphenyl)-5-fluorpyridin-3-yl]-1-[2-fluor-4-(trifluormethyl)phenyl]piperidin-4-carboxamid;

1-290: 4-[6-(2-ethoxyphenyl)-5-fluorpyridin-3-yl]-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

25 **1-291:** 1-[2-chlor-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxyphenyl)-5-fluorpyridin-3-yl]piperidin-4-carboxamid;

1-292: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)-5-fluorpyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

30 **1-293:** 1-[2-fluor-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-294: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-295: N-[(2S)-1-(dimethylamino)propan-2-yl]-1-[2-fluor-4-(trifluormethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-296: 1-[2-chlor-4-(trifluormethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-[2'-methoxy-[2,3'-bipyridin]-5-yl]piperidin-4-carboxamid;

1-297: N-[(2S)-1-(dimethylamino)propan-2-yl]-1-[2-fluor-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

5 **1-298:** 1-[2-fluor-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidin-4-carboxamid;

1-299: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

10 **1-300:** N-[2-(dimethylamino)ethyl]-4-{3-fluor-2'-methoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]piperidin-4-carboxamid;

1-301: 4-{3-fluor-2'-methoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-302: 1-[2-chlor-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{3-fluor-2'-methoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

15 **1-303:** 1-[2-chlor-4-(trifluormethyl)phenyl]-4-{3-fluor-2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-304: N-[2-(dimethylamino)ethyl]-1-[2-fluor-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

20 **1-305:** 1-[2-fluor-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-306: 1-[2-fluor-4-(trifluormethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidin-4-carboxamid;

1-307: 1-[2-chlor-4-(trifluormethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidin-4-carboxamid;

25 **1-308:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-309: 1-[2-chlor-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[5-fluor-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

30 **1-310:** 1-[2-chlor-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-311: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-N-[2-(methylamino)ethyl]piperidin-4-carboxamid;

1-312: N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]piperidin-4-carboxamid;

1-313: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-314: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidin-4-carboxamid;

5 **1-315:** 4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidin-4-carboxamid;

1-316: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidin-4-carboxamid;

10 **1-317:** N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]piperidin-4-carboxamid;

1-318: 1-(2,4-dichlorophenyl)-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-319: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

15 **1-320:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidin-4-carboxamid;

1-321: 4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-1-[2-fluor-4-(trifluormethyl)phenyl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidin-4-carboxamid;

20 **1-322:** 1-(2,4-dichlorophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidin-4-carboxamid;

1-323: 1-(4-chlor-2-cyanophenyl)-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}-N-[(2S)-1-(methylamino)propan-2-yl]piperidin-4-carboxamid;

1-324: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-[5-fluor-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-carboxamid;

25 **1-325:** 1-(4-chlor-2-cyanophenyl)-N-[(2S)-1-(dimethylamino)propan-2-yl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-326: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]-N-[(2S)-1-(methylamino)propan-2-yl]piperidin-4-carboxamid;

30 **1-327:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[[[(2S)-4-methylmorpholin-2-yl]methyl]piperidin-4-carboxamid;

1-328: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[[[(3S)-4-methylmorpholin-3-yl]methyl]piperidin-4-carboxamid;

1-329: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-[[[(3R)-4-methylmorpholin-3-yl]methyl]piperidin-4-carboxamid;

1-330: 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]-N-
{[(2R)-4-methylmorpholin-2-yl]methyl}piperidin-4-carboxamid;

1-331: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[2-(dimethylamino)ethyl]-4-[6-(2-
ethoxypyridin-3-yl)pyridazin-3-yl]piperidin-4-carboxamid;

5 **1-332:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2S)-2-(dimethylamino)propyl]-4-{2'-
ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-333: 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2R)-2-(dimethylamino)propyl]-4-{2'-
ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

10 **1-334:** 1-[2-cyan-4-(trifluormethyl)phenyl]-N-[(2R)-2-(dimethylamino)-3-
hydroxypropyl]-4-{2'-ethoxy-3-fluor-[2,3'-bipyridin]-5-yl}piperidin-4-carboxamid;

1-335: N-[2-(dimethylamino)ethyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]-1-[2-
fluor-4-(trifluormethyl)phenyl]piperidin-4-carboxamid;

2-1: 2-{4-[2-(dimethylamino)ethoxy]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-1-
yl}-5-(trifluormethyl)benzonitril;

15 **2-2:** 1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-
4-yl N-[2-(dimethylamino)ethyl]carbamate;

2-3: 3-amino-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-
yl)phenyl]piperidin-4-yl}propanamid;

20 **2-4:** 3-amino-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-
yl}piperidin-4-yl}propanamid;

2-5: N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-
yl}piperidin-4-yl}-3-(methylamino)propanamid;

2-6: (3R)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-
yl}piperidin-4-yl}pyrrolidin-3-carboxamid;

25 **2-7:** (3R)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-
yl}piperidin-4-yl}-1-methylpyrrolidin-3-carboxamide;

2-8: (3S)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-
yl}piperidin-4-yl}pyrrolidin-3-carboxamid;

30 **2-9:** (3S)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-
yl}piperidin-4-yl}-1-methylpyrrolidin-3-carboxamid;

2-10: N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-
yl}piperidin-4-yl}-3-(dimethylamino)propanamid;

2-11: (2S)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-
yl)phenyl]piperidin-4-yl}-1-methylpyrrolidin-2-carboxamid;

2-12: (3R)-1-methylpyrrolidin-3-yl N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamate;

2-13: (3S)-1-methylpyrrolidin-3-yl N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}carbamate;

5 **2-14:** 2-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl})-4-[(2-hydroxyethyl)amino]piperidin-1-yl)-5-(trifluormethyl)benzonitril;

2-15: 2-(4-{2'-ethoxy-[2,3'-bipyridin]-5-yl})-4-{[2-(methylamino)ethyl]amino}piperidin-1-yl)-5-(trifluormethyl)benzonitril;

10 **2-16:** N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}-1-methylazetidin-3-carboxamid;

2-17: (2R)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[4-(2-ethoxypyridin-3-yl)phenyl]piperidin-4-yl}-1-methylpyrrolidin-2-carboxamid;

2-18: (3S)-1-methylpyrrolidin-3-yl N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamate;

15 **2-19:** 1-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-[(3R)-1-methylpyrrolidin-3-yl]urea;

2-20: 1-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-[(3S)-1-methylpyrrolidin-3-yl]urea;

20 **2-21:** (3R)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)pyrrolidin-1-carboxamid;

2-22: (3R)-1-methylpyrrolidin-3-yl N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}carbamate;

2-23: N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)azetidin-1-carboxamid;

25 **2-24:** (3S)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-3-(methylamino)pyrrolidin-1-carboxamid;

2-25: N-[1-(4-chlor-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-1-methylazetidin-3-carboxamid;

30 **2-26:** N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-2-(dimethylamino)acetamid;

2-27: (3S)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidin-3-carboxamid;

2-28: (3S)-N-[1-(4-chlor-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-1-methylpyrrolidin-3-carboxamid;

2-29: (3R)-N-[1-(4-chlor-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-1-methylpyrrolidin-3-carboxamid;

2-30: N-[1-(4-chlor-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-3-(dimethylamino)propanamid;

5 **2-31:** N-[1-(4-chlor-2-cyanophenyl)-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl]-2-(dimethylamino)acetamid;

2-32: N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-1-methylazetidin-3-carboxamid;

10 **2-33:** (3R)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidin-3-carboxamid;

2-34: N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-2-(dimethylamino)acetamid;

2-35: N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-1-methylazetidin-3-carboxamid;

15 **2-36:** N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamid;

2-37: N-[1-(4-chlor-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl]-1-methylazetidin-3-carboxamid;

20 **2-38:** N-[1-(4-chlor-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl]-2-(dimethylamino)acetamid;

2-39: (3S)-N-[1-(4-chlor-2-cyanophenyl)-4-{2'-ethoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl]-1-methylpyrrolidin-3-carboxamid;

2-40: N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-{2'-methoxy-[2,3'-bipyridin]-5-yl}piperidin-4-yl}-1-methylazetidin-3-carboxamid;

25 **2-41:** (3R)-1-methylpyrrolidin-3-yl N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}carbamat;

2-42: (3S)-1-methylpyrrolidin-3-yl N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}carbamat;

30 **2-43:** 1-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-3-[(3S)-1-methylpyrrolidin-3-yl]urea;

2-44: (3S)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}pyrrolidin-3-carboxamid;

2-45: N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-1-methylazetidin-3-carboxamid;

2-46: (3S)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidin-3-carboxamid; og

2-47: (3R)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidin-3-carboxamid;

5 **2-48:** 1-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(1-methyl-1H-pyrrol-2-yl)pyridin-3-yl]piperidin-4-yl}-3-[(3R)-1-methylpyrrolidin-3-yl]urea;

2-49: (3S)-1-methylpyrrolidin-3-yl N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;

10 **2-50:** (3R)-1-methylpyrrolidin-3-yl N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;

2-51: 1-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[(3S)-1-methylpyrrolidin-3-yl]urea;

2-52: 1-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[(3R)-1-methylpyrrolidin-3-yl]urea;

15 **2-53:** (3S)-1-methylpyrrolidin-3-yl N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;

2-54: (3R)-1-methylpyrrolidin-3-yl N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;

20 **2-55:** 1-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[(3S)-1-methylpyrrolidin-3-yl]urea;

2-56: 1-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[(3R)-1-methylpyrrolidin-3-yl]urea;

2-57: 2-(dimethylamino)ethyl N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}carbamate;

25 **2-58:** 1-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-[2-(dimethylamino)ethyl]urea;

2-59: 1-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(1-methylazetidin-3-yl)urea;

30 **2-60:** (3R)-N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-1-methylpyrrolidin-3-carboxamid;

2-61: N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-2-(dimethylamino)acetamid;

2-62: N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamid;

2-63: N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(methylamino)propanamid;

2-64: N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[6-(2-ethoxypyridin-3-yl)pyridazin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamid;

5 eller et farmaceutisk acceptabelt salt deraf.

14. Forbindelse, der er N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamid, eller et farmaceutisk acceptabelt salt deraf.

10

15. Forbindelse ifølge krav 14, hvor forbindelsen er N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamid.

16. Forbindelse ifølge krav 14, hvor forbindelsen er et farmaceutisk acceptabelt salt af N-{1-[2-cyan-4-(trifluormethyl)phenyl]-4-[5-fluor-6-(2-methoxyphenyl)pyridin-3-yl]piperidin-4-yl}-3-(dimethylamino)propanamid.

15

17. Farmaceutisk sammensætning, der omfatter en forbindelse ifølge et hvilket som helst af kravene 1-16, eller et farmaceutisk acceptabelt salt deraf, og mindst ét farmaceutisk acceptabelt hjælpestof.

20

18. Forbindelse ifølge et hvilket som helst af kravene 1-16, eller et farmaceutisk acceptabelt salt deraf, til anvendelse i behandling af Cushings syndrom, ektopisk Cushings syndrom, kongenit adrenal hyperplasi (CAH) eller til reduktion af udskillelse af adrenokortikotropt hormon (ACTH) hos et pattedyr.

25