

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date  
3 November 2005 (03.11.2005)

PCT

(10) International Publication Number  
**WO 2005/101979 A2**

(51) International Patent Classification: **Not classified**

(21) International Application Number: **PCT/IL2005/000440**

(22) International Filing Date: 21 April 2005 (21.04.2005)

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
161595 22 April 2004 (22.04.2004) IL  
60/670,290 12 April 2005 (12.04.2005) US

(71) Applicant (for all designated States except US): **MOR RESEARCH APPLICATIONS LTD.** [IL/IL]; 20 HaM-  
agshimim Street, Kiryat-Matalon, 49170 Petach-Tikva  
(IL).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **BARAK, Nir** [IL/IL];  
15 Shalom Ash Street, Hadar-Yosef, 69483 Tel-Aviv (IL).

(74) Agent: **G. E. EHRLICH (1995) LTD.**; 11 Menachem Be-  
gin Street, 52 521 Ramat Gan (IL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**WO 2005/101979 A2**

(54) Title: METHOD OF FOOD INTAKE MANAGEMENT

(57) **Abstract:** Methods for regulating food intake in a human subject; for improving a compliance of a human subject to caloric restriction; and for reducing a desire of a human subject to consume fats, utilizing H1-receptor agonists that have a pharmacological half-life that allows an efficient treatment regime thereof are disclosed. The methods can be efficiently used for treating conditions such as overeating, overweight, obesity, binge eating disorder, night eating syndrome, obsessive eating, compulsive eating and bulimia, as well as conditions associated with metabolic derangement such as dyslipidemia.

## METHOD OF FOOD INTAKE MANAGEMENT

FIELD OF THE INVENTION

5 The present invention relates to novel methods for regulating food intake in human subjects and more particularly to methods of preventing or treating conditions in which weight management is beneficial.

BACKGROUND OF THE INVENTION

10 Obesity is a chronic, complex, multi-factorial disease, involving social, cultural, genetic, physiological and psychological components, and is associated with substantially increased morbidity and mortality. Over-nutrition is attributed as the cause of about 400,000 deaths a year in the USA (*Mokdad, 2004*), and may be considered to be an epidemic. Based on the body-mass index, defined as the ratio of weight and squared height, (ranging normally from 18.5 to 24.9), about one third of the adult 15 population is overweight (an index of from 25 to 29.9), and more than one quarter is obese (index greater than 30) (*National Center for Health Statistics, 2000*). Environmental and behavioral changes brought about by economic development and modernization have been linked to the rise in global obesity. The environmental factors which foster the tendency toward obesity include lack of physical activity 20 combined with high-calorie, low-cost foods. The prevalence of overweight and obesity is increasing worldwide at an alarming rate in both developing and developed countries, in children and adults, men and women. The number of overweight and obese people has continued to increase since 1960, a trend that is not slowing down. Today, 64.5 % of adult Americans – about 127 million – are categorized as being 25 overweight or obese and nearly one-third (30.5 %) – about 60 million – are obese, as reported in the National Health and Nutrition Examination Survey (NHANES) by the Centers for Disease Control and Prevention (CDC).

30 Obesity significantly increases the risk of illness from about thirty serious medical conditions and is associated with increases in deaths from all-causes. Among these are high blood pressure, diabetes, osteoarthritis, heart disease, stroke, gallbladder disease and cancer of the breast, prostate and colon (*National Task Force on the Prevention and Treatment of Obesity, 2000*). Furthermore, each year, obesity causes

at least 300,000 excess deaths in the U.S., being the second leading cause of unnecessary deaths. Healthcare costs of American adults with obesity amount to approximately 100 billion dollars.

There are several different treatment options for management of weight, 5 including: dietary therapy, physical activity, behavior therapy, drug therapy and surgery. For the majority of overweight and obese people, who find they are unable to change their lifestyle, drug therapy is the most favorable and applicable option. Although hundreds of millions of people are seeking drug therapy for the treatment of obesity, current drug therapies do not meet this need due to their undesired side effects 10 and limited efficacy.

Medications for the treatment of obesity are currently approved for use in adults with a body-mass index of 30 or higher, or with a body-mass index of 27 or higher who have obesity-related medical problems (*Physicians' Desk Reference, 2001*). Approximately 10 percent of women and 3 percent of men with a body-mass 15 index of 30 or higher reportedly use weight-loss medications (*Serdula, 1999*).

Medications currently approved for weight loss in the United States fall into two categories: those that decrease food intake by reducing appetite or increasing satiety (appetite suppressants), and those that decrease nutrient absorption. A potentially third category, medications increasing energy expenditure, such as 20 ephedrine, is not currently approved for treating obesity in the United States.

The only FDA-approved medication for obesity that reduces nutrient absorption is orlistat (Xenical), which acts by binding to gastrointestinal lipases in the lumen of the gut, preventing hydrolysis of dietary fat into absorbable free fatty acids and monoacylglycerols.

25 Most appetite suppressants work primarily by increasing the availability of anorexigenic neurotransmitters – notably norepinephrine, serotonin, dopamine, or some combination of these neurotransmitters – in the central nervous system. Noradrenergic drugs available in the United States include phentermine, diethylpropion, phendimetrazine, and benzphetamine. Some of these drugs are 30 considered by the Drug Enforcement Administration (DEA) to have a potential for abuse. Amphetamines, which are considered to have a particularly high potential for abuse are no longer recommended for weight loss for this reason. The Food and Drug

Administration (FDA) approves the medications for use of "a few weeks" only (generally presumed to be 12 weeks or less) for the treatment of obesity.

Side effects of noradrenergic medications include insomnia, dry mouth, constipation, euphoria, palpitations, and hypertension (*Physicians' Desk Reference, 2001*).

Serotonergic agents act by increasing the release of serotonin, inhibiting its reuptake, or both. Fenfluramine (Pondimin) and dexfenfluramine (Redux), medications that both stimulate serotonin release and inhibit its reuptake, were withdrawn from the market in the United States in 1997 because of associations with valvular heart disease and pulmonary hypertension. Some selective serotonin-reuptake inhibitors have induced weight loss in short-term studies, and fluoxetine (Prozac) has undergone considerable evaluation to determine its efficacy for weight loss (*Goldstein, 1993*). However, after initial weight loss, steady regain was observed in later stages of the treatment (*National Task Force on the Prevention and Treatment of Obesity*). Sertraline (Zoloft), evaluated as an adjunct for weight maintenance after a very-low-calorie diet, showed a similar lack of long-term efficacy (*Wadden, 1995*). Sibutramine (Meridia, Reductil), an inhibitor of both norepinephrine reuptake and serotonin reuptake that also weakly inhibits dopamine reuptake, is approved by the FDA for weight loss and weight maintenance in conjunction with a reduction diet. Side effects of sibutramine include increased blood pressure and pulse frequency rate (*McMahon, 2000*).

Rimonabant (Sanofi), which is claimed to stop food cravings, represents a new class of drugs that inhibit the activity of the CB1 receptor. The CB1 receptor forms a part of the endocannabinoid system. The CB1 receptor has been found in the brain, fat cells and other parts of the body, and has been associated with regulating food intake and with tobacco dependency (*Pi-Sunyer et al., 2004*). The endocannabiniod system helps to regulate pleasure, relaxation, and pain tolerance. Little is currently known about the long-term effects of inhibition of this system. Further, neurologists point out that the endocannabinoid system helps to protect the brain under some circumstances (such as stroke and head injury,) such that brain damage in these circumstances might be worse in patients taking drugs that block the endocannabiniod system.

The Rimonabant drug is currently undergoing phase III clinical trials. Presently reported side effects associated therewith include anxiety, nausea and diarrhea.

Hence, although some of the currently approved medications show moderate effects and can help some patients in losing weight, there is a continuing need for efficacious treatment regimes and drugs for alleviating the serious and prevalent disorder – the weight excess.

Histamine, a potent bioactive substance that has been studied for nearly a century, is an aminergic neurotransmitter. Four histamine receptors have been identified: H<sub>1</sub>, H<sub>2</sub>, H<sub>3</sub>, and H<sub>4</sub>, leading to the discovery and therapeutic use of potent receptor antagonists. Activation of the H<sub>1</sub> receptor is associated with effects on smooth muscle and central neurons; activation of the H<sub>2</sub> receptor stimulates acid secretion in the stomach, while activation of the H<sub>3</sub> receptor results in a pre-synaptic autoregulatory effect.

Histamine has been implicated, among others, in the regulation of arousal state (*Lin et al., 1990*), locomotor activity (*Clapham, 1994*), cardiovascular control (*Imamura, 1996*), water intake (*Lecklin, 1998*), food intake (*Leurs, 1998*), and memory formation (*Blandina, 1996*). Some studies have indicated that histamine may suppress appetite by acting on hypothalamic histaminergic neurons that participate in the regulation of food intake (*Sakata, 1997; Bjenning, 2000; Sakata, 1995*). Thus, it was reported that histamine injected intracerebroventricularly acts as an appetite suppressant, and that depletion of histamine stimulates feeding (*Tuomisto, 1994*). Changes in histaminergic tone in the CNS have been associated with genetic models of obesity (*Machidori, 1992*). In addition, intracerebroventricular injection of leptin has been correlated with changes in the turnover rate of hypothalamic neuronal histamine (*Yoshimatsu, 1999*). Since histamine is unable to cross the blood brain barrier, these effects would not be expected to be seen with systemic administration of histamine.

In both humans and rodents, treatment with an H<sub>1</sub> antagonist resulted in hyperphagia (*Fukagawa, 1989*), and administration of H<sub>3</sub> antagonists led to hypophagia (*Attoub, 2001*). However, the long-term effects of H<sub>3</sub> receptors on anorexigenic activities for body-weight homeostasis have not been documented

because of the off-target activity (*Leurs, 1995*) and toxicity profile of H<sub>3</sub> inhibitors (*Onderwater, 1998*).

Betahistine is an orally active histamine-like drug extensively used in the symptomatic treatment of vestibular disorders, mainly vertigo and Menier's disease.

5      Betahistine is a structural analog of histamine, in which the imidazole ring of the histamine is replaced by a pyridine ring. Betahistine is an H<sub>1</sub> receptor agonist, and has been found to exhibit an H<sub>1</sub>-agonism activity of about 0.07 times that of histamine, and to cause hypotensive response, bronchoconstriction, and increased vasopermeability after parenteral administration. Receptor binding studies have also

10     shown that betahistine is a potent H<sub>3</sub>-receptor antagonist. Betahistine is able to cross the blood brain barrier and act centrally by enhancing histamine synthesis in tuberomammillary nuclei of the posterior hypothalamus. Adverse side effects associated with betahistine are typically minor and include skin rashes of various types, urticaria, and itching. Gastric upset, nausea, and headache have also been

15     reported by some patients.

It has been found (*Rossi et al., 1999*) that high doses of betahistine, delivered intraperitoneally, increased water intake and decreased food intake in pygmy goats. This was suggested as being due to stimulation of both H<sub>1</sub> and H<sub>2</sub> receptors, since in addition to its known action as an H<sub>1</sub> receptor agonist, betahistine has been shown to

20     act as a weak partial agonist of peripheral histamine H<sub>2</sub> receptors (*Arrang et al., 1985*). This idea is further supported by recent findings showing that the hypophagic effect of histamine was blocked by the H<sub>2</sub>-receptor antagonist, cimetidine, in pygmy goats. These similarities in the hypophagic effects of histamine and betahistine suggest an involvement of H<sub>2</sub>-receptors in the hypophagic effect of betahistine in

25     pygmy goats. H<sub>2</sub>-receptors are not, however, associated with weight change in humans (*Rasmussen, 1993*).

Szelag et al. (2001) found that betahistine, when given intraperitoneally, decreased food intake in rats, whereas this effect was not seen when betahistine was given intragastrically (*Szelag, 2002*). It was suggested that the effect of betahistine

30     administration on food intake involves increasing histamine synthesis and release as a result of H<sub>3</sub> receptor inhibition. . However, since activation of H<sub>2</sub> receptors is known to stimulate hydrochloric acid secretion (see, for example, *Clayman, 1977*), it was

further suggested that the lack of the influence of betahistine on food intake after intragastrical administration may be due to the fact that betahistine increased hydrochloric acid release by activation of H<sub>2</sub> receptors, thereby abolishing the central anorectic activity of betahistine. It was also suggested that the effects on H<sub>1</sub> receptors in humans may differ significantly from those in rats due to variations in circadian rhythm between the species. Therefore, it appears that the anorectic response of betahistine is dependent upon species and route of administration. Nevertheless, Szelag et al. fails to teach the effect of betahistine administered orally or by any other route of administration, on food intake in humans.

Lecklin et al. (2002) found that inhibition of histamine catabolism by intraperitoneal injection of metoprine, a histamine-N-methyltransferase inhibitor, resulted in suppressed daily energy intake and ingestion of fat in rats.

Pharmacokinetic studies showed that betahistine is transformed, mainly in the liver, to 2-(2-aminoethyl)-pyridine (AEP) and to 2-(2-hydroxyethyl)-pyridine (HEP), (Sternoson, 1974) whereas both betahistine and the metabolites bind to histamine receptors.

The cited references corroborate the complexity of appetite regulation, which includes, among other factors, species specificity and route of administration. It has further been found (Seifert et al., 2003) that multiple differences exist in agonist and antagonist pharmacology of histamine receptors between different species, such as humans and guinea pigs. The prior art does not teach or suggest the use of H<sub>1</sub> agonists for regulating food intake in humans. The prior art further does not teach or suggest the use of such H<sub>1</sub> agonists that have a pharmacological half life that permits an efficient treatment therewith. The prior art further does not teach or suggest the use of orally administered H<sub>1</sub> agonists for regulating food intake in humans.

There is thus a widely recognized need for and it would be highly advantageous to have histamine-related agents for regulation of food intake in humans, devoid of the above limitations.

### 30 SUMMARY OF THE INVENTION

The present invention successfully addresses the shortcomings of the presently known methods for regulating food intake by providing such methods that utilize H<sub>1</sub>

receptor agonists, which are highly efficient in humans, and which have a pharmacological half life of at least 3 hours and which are devoid of the limitations of the currently known methods.

According to one aspect of the present invention there is provided a method of 5 treating of a condition in which regulating a food intake in a human subject is beneficial, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist, which has a pharmacological half-life of at least 3 hours.

According to further features in preferred embodiments of the invention 10 described below, the condition is selected from the group consisting of overeating, overweight, obesity, and a disorder caused or exacerbated thereby. The condition caused or exacerbated by the conditions according to this aspect of the present invention may be selected from the group consisting of a muscoskeletal disorder, a cardiovascular disorder, a dermatological disorder, a sleep disorder, a metabolic 15 condition, diabetes and a diabetes-related condition.

According to still further features in the described preferred embodiments, the condition is associated with high fat consumption.

According to still further features in the described preferred embodiments, the condition is associated with a psychological factor. The condition according to this aspect of the present invention optionally comprises binge eating disorder, night eating syndrome, obsessive eating, compulsive eating or bulimia.

According to still further features in the described preferred embodiments the 20 condition is associated with a drug treatment. The drug may optionally consist of a steroid hormone or a psychoactive drug.

According to another aspect of the present invention, there is provided a method of improving a compliance of a human subject to caloric restriction, the 25 method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist, which has a half-life of at least 3 hours.

According to still another aspect of the present invention, there is provided a method of reducing a desire of a human subject to consume fats, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> 30 agonist, which has a half-life of at least 3 hours.

According to yet another aspect of the present invention, there is provided a method of treating a condition associated with a metabolic derangement in a human subject, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist which has a half-life of at least 3 hours. The metabolic derangement may be, for example, dyslipidemia, such as hypercholesterolemia or lipemia.

As used herein, the term "metabolic derangement" refers to an imbalance in the level of one or more metabolites within a body. A common metabolic derangement is typically associated with an imbalance in the level of metabolites such as cholesterol, including LDL and HDL, triglycerides and the like.

According to still another aspect of the present invention, there is provided a method of reducing total cholesterol level in a human subject, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist which has a half-life of at least 3 hours.

According to still another aspect of the present invention, there is provided a method of reducing low-density lipoprotein (LDL) cholesterol and increasing high-density lipoprotein (HDL) cholesterol levels in a human subject, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist which has a half-life of at least 3 hours.


According to still another aspect of the present invention, there is provided a method of reducing triglyceride level in a human subject, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist which has a pharmacological half-life of at least 3 hours.

According to further features in the described preferred embodiments, the pharmacological half-life ranges from about 3 hours to about 12 hours, preferably from about 3 hours to about 8 hours, more preferably from about 3 hours to about 5 hours..

According to still further features in the described preferred embodiments, the H<sub>1</sub> agonist is further an H<sub>3</sub> antagonist.

According to still further features in the described preferred embodiments, the H<sub>1</sub> agonist is characterized by blood brain barrier permeability.

According to still further features in the described preferred embodiments, the H<sub>1</sub> agonist is selected from the group consisting of betahistine, a betahistine metabolite, a betahistine pharmaceutically acceptable salt, a betahistine prodrug, a betahistine derivative and any combination thereof. Preferably, the betahistine metabolite is 2-(2-aminoethyl)-pyridine or 2-(2-hydroxyethyl)-pyridine. Also preferably, the betahistine salt is betahistine dihydrochloride, betahistine mesilate, or betahistine trimebutine maleate. Also preferably, the betahistine derivative is selected from the group of compounds represented by the general formula I:



10

Formula I

wherein each of R<sub>1</sub>-R<sub>12</sub> is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl and aryl.

According to still further features in the described preferred embodiments, administering is effected by a route selected from the group consisting of the oral, transdermal, intravenous, subcutaneous, intramuscular, intranasal, intraauricular, sublingual, rectal, transmucosal, intestinal, buccal, intramedullar, intrathecal, direct 20 intraventricular, intraperitoneal, and intraocular routes. Preferably, administering is effected by the oral, transdermal, buccal, transmucosal, rectal or sublingual routes. More preferably, administering is effected by the oral, buccal or transdermal routes.

According to still further features in the described preferred embodiments, the therapeutically effective amount ranges from about 2 mg per day to about 96 mg per 25 say, preferably from about 10 mg per day to about 50 mg per day.

According to still further features in the described preferred embodiments, administering is effected from about 1 to about 4 times per day, preferably, twice per day.

According to still further features in the described preferred embodiments, administering is effected according to the development of hunger of the subject.

According to still further features in the described preferred embodiments, administering is performed such that a decrease of the body weight of the subject that 5 ranges from about 1 to about 5 percent is effected, without restricting the food intake of the subject. Preferably, such administering is performed such that no down-regulation of H<sub>1</sub> receptors is effected.

As used herein, the term "down-regulation" with regard to receptors refers to a 10 decrease in the responsiveness of the receptor, or to a decrease in the number or density of receptors. Decreasing the responsiveness of a receptor also includes a complete shutdown of the receptor.

According to still further features in the described preferred embodiments, the H<sub>1</sub> agonist forms a part of a pharmaceutical composition, which further comprises a pharmaceutically acceptable carrier.

15 According to still further features in the described preferred embodiments, each of the methods described herein further comprises administering to the subject a therapeutically effective amount of an additional active agent such as, for example, a weight control agent. The weight control agent may be, for example, an appetite suppressant. Representative examples of suitable appetite suppressants include, 20 without limitation, noradrenergic agents, serotonergic agents, dopamnergic agents, endocannabinoid receptor blockers, or combinations thereof.

According to still further features in the described preferred embodiments, the 25 additional active agent may be an agent for the treatment of a musculoskeletal disorder, a cardiovascular disorder, a dermatological disorder, a sleep disorder, a metabolic condition, diabetes or a diabetes-related condition. Representative examples of such active agents include, without limitation, a non-steroidal anti-inflammatory drug, a muscle relaxant, an antigout agent, an immunosuppressant, a drug affecting bone mineralization, an alpha-adrenergic blocking drugs, an angiotensin-converting enzyme inhibitor, an antiarrhythmic drug, an anticoagulant, an antiplatelet, a 30 thrombolytic, a beta-adrenergic blocking drug, a calcium channel blocker, a centrally acting drug, a digitalis drug, a nitrate, a peripheral adrenergic antagonist, a vasodilator, an acne medication, an antipruritic agent, a corticosteroid, an anti-psoriasis agent, an

5 anti-eczema agent, a hypnotic, an anti-depressant, an anti-histamine, a sulfonylurea, a meglitinide, a biguanide, a thiazolidinedione, an alpha-glucosidase inhibitor, a PPAR (peroxisome proliferator-activated receptor)-gamma antagonist, insulin, a fibrate, an HMG-CoA reductase inhibitor, a bile acid sequestrant, a cholesterol absorption inhibitor, nicotinic acid, a derivative, analog and metabolite thereof, and any mixture thereof.

10 According to still further features in the described preferred embodiments, the methods described herein are used for inducing weight loss. Alternatively, these methods are used for maintaining weight loss or preventing a weight gain after or during a weight reducing diet. Further alternatively, these methods are used for preventing weight gain in a subject having a condition associated with weight gain.

15 Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. In case of conflict, the patent specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

20 As used herein the term "method" refers to manners, means, techniques and procedures for accomplishing a given task including, but not limited to, those manners, means, techniques and procedures either known to, or readily developed from known manners, means, techniques and procedures by practitioners of the chemical, pharmacological, biological, biochemical and medical arts.

25 As used herein, the term "treating" includes abrogating, substantially inhibiting, slowing or reversing the progression of a condition, substantially ameliorating clinical or aesthetical symptoms of a condition or substantially preventing the appearance of clinical or aesthetical symptoms of a condition.

As used herein the term "regulating" with regard to food intake refers to controlling or adjusting the food intake to a desired level.

30 The term "comprising" means that other steps and ingredients that do not affect the final result can be added. This term encompasses the terms "consisting of" and "consisting essentially of".

The phrase "consisting essentially of" means that the composition or method may include additional ingredients and/or steps, but only if the additional ingredients and/or steps do not materially alter the basic and novel characteristics of the claimed composition or method.

5 The term "therapeutically effective amount" or "pharmaceutically effective amount" denotes that dose of an active ingredient or a composition comprising the active ingredient that will provide the therapeutic effect for which the active ingredient is indicated.

10 As used herein, the term "agonist" describes a substance that is capable of binding to a receptor on a cell and thereby initiating a physiological activity or pathway. The phrases "H<sub>1</sub>-receptor agonist" and "H<sub>1</sub> agonist" are used herein interchangeably.

As used herein, the term "antagonist" describes a substance that acts within the body to reduce the physiological activity of another substance.

15 As used herein, the phrase "pharmacological half-life" describes the time required for half the quantity of a drug or other substance deposited in a living organism to be metabolized or eliminated from the plasma by normal biological processes. This phrase is also referred to herein interchangeable as "half life".

20 As used herein a "pharmaceutical composition" refers to a preparation of one or more of the active ingredients described herein, either compounds or physiologically acceptable salts thereof, with other chemical components such as traditional drugs, physiologically suitable carriers and excipients.

25 As used herein, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. Herein, the phrases "physiologically suitable carrier" and "pharmaceutically acceptable carrier" are interchangeably used and refer to an approved carrier or a diluent that does not cause significant irritation to an organism and does not abrogate the biological activity and properties of the administered 30 conjugate.

As used herein, the term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered.

Herein the term "excipient" refers to an inert substance added to a pharmaceutical composition to further facilitate processes and administration of the active ingredients.

As used herein, the singular form "a," "an," and "the" include plural references 5 unless the context clearly dictates otherwise. For example, the term "a compound" or "at least one compound" may include a plurality of compounds, including mixtures thereof.

Throughout this disclosure, various aspects of this invention can be presented 10 in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range 15 should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

Whenever a numerical range is indicated herein, it is meant to include any 20 cited numeral (fractional or integral) within the indicated range. The phrases "ranging/ranges between" a first indicate number and a second indicate number and "ranging/ranges from" a first indicate number "to" a second indicate number are used herein interchangeably and are meant to include the first and second indicated numbers and all the fractional and integral numerals therebetween.

25 **BRIEF DESCRIPTION OF THE DRAWINGS**

The invention is herein described, by way of example only, with reference to 30 the accompanying drawings. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of the preferred embodiments of the present invention only, and are presented in the cause of providing what is believed to be the most useful and readily understood description of the principles and conceptual aspects of the invention. In this regard, no attempt is made to show structural details of the invention

in more detail than is necessary for a fundamental understanding of the invention, the description taken with the drawings making apparent to those skilled in the art how the several forms of the invention may be embodied in practice.

In the drawings:

5 FIGs. 1(a-b) present results obtained in prior art studies, demonstrating the effect of Metoprone treatment on the total caloric intake (Figure 1a) and the fat intake (Figure 1b) in rats;

10 FIGs. 2(a-b) present results obtained in prior art studies, demonstrating the effect of Metoprone treatment on the total caloric intake, compared with the carbohydrate intake (Figure 2a) and the protein intake (Figure 1b) in rats;

FIG. 3 presents plots demonstrating the effect of overweight (expressed as the body mass index) on the relative risk of death from cardiovascular disease (dotted line), cancer (dashed line) and other causes in men (upper plots) and women (bottom plots) ;

15 FIG. 4 is a bar graph presenting the effect of oral administration of betahistine (blue bars) and a placebo (yellow bars) on the total caloric intake of humans following 14 days and 28 days of treatment;

20 FIG. 5 is a bar graph presenting the effect of oral administration of betahistine (blue bars) and a placebo (yellow bars) on the fat, carbohydrate and protein intake of humans following 14 days and 28 days of treatment; and

FIG. 6 is a bar graph presenting the effect of oral administration of betahistine (blue bars) and a placebo (yellow bars) on the weight change in humans following 14 days and 28 days of treatment.

25 DETAILED DESCRIPTION OF THE INVENTION

The present invention is of novel methods for regulating food intake in a human subject; for improving a compliance of a human subject to caloric restriction; and for reducing a desire of a human subject to consume fats, all of which utilize H<sub>1</sub>-receptor agonists that have a pharmacological half-life that allows an efficient treatment regime thereof.

The principles and operation of the compositions and methods according to the present invention may be better understood with reference to the drawings and accompanying descriptions.

Before explaining at least one embodiment of the invention in detail, it is to 5 be understood that the invention is not limited in its application to the details set forth in the following description or exemplified by the Examples. The invention is capable of other embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

10 As is described hereinabove, many of the presently used appetite suppressants are known to have a potential for abuse. Furthermore, these appetite suppressants are associated with various side effects, ranging from insomnia, dry mouth, constipation, euphoria, palpitations and hypertension, to valvular heart disease and pulmonary hypertension. In addition, many of these agents have been found to 15 show a lack of long-term efficacy, with steady weight-gain occurring after initial weight loss.

As is further described hereinabove, studies have shown a relationship 20 between histamine and regulation of food intake. Thus, for example, as shown in Figure 1, inhibition of histamine catabolism by the histamine N-methyltransferase inhibitor, metoprine, which is typically utilized as a cytotoxic, anti-cancer agent, has been shown to suppress total food intake and ingestion of fat by rats (*Lecklin et al.*, 2002). Treatment had no marked effect on intakes of carbohydrates or protein, as shown in Figure 2, indicating that an increase in the brain histamine content may 25 reduce specifically "fat-appetite", namely, the desire to specifically ingest fats, as opposed to other food types.

Systemic administration of histamine, however, is extremely inefficient in its 30 centrally mediated anti-obesity action because exogenous histamine poorly penetrates the blood brain barrier. Histamine has therefore been administered as an appetite suppressant hitherto by intracerebroventricular injection. Similarly, systemic administration of histamine would be expected to have little affect on the desire of a human subject to consume fats.

Furthermore, histamine has a pharmacological half-life of only a few minutes. Therefore, the duration of action of the drug is very short, and it would need to be administered frequently in order to build up and maintain a high enough concentration in the blood to be therapeutically effective. A drug having a longer half life would 5 therefore require less frequent administration, which may lead to increased patient compliance with the dosing regime. This may also result in fewer side effects, as peaks and troughs of the level of the drug in the bloodstream of the patient may be decreased, leading to a more even drug level in the blood over a period of time.

While several studies have suggested a role for H<sub>1</sub>-receptor agonists other than 10 histamine in regulating food intake, none of these studies has established a direct effect of such agonists on food intake in humans.

The present inventors have now surprisingly found that certain H<sub>1</sub>-agonists, which are characterized by relatively long half-life, particularly as compared with histamine, can be efficiently utilized for regulating food intake in humans. More 15 specifically, it was found that betahistine, a presently known and approved drug for treating menier's disease and associated conditions, and which has a half-life of about 3.5 hours, efficiently affects food intake and caloric intake and reduces weight, as well as fat consumption, in obese female subjects. As is exemplified in the Examples section that follows, in a randomized placebo controlled double-blinded study, it was 20 found that betahistine hydrochloride decreased appetite, lowered food intake and particularly affected fat consumption.

It is therefore demonstrated herein that the administration of an H<sub>1</sub> receptor agonist that has a pharmacological half life that permits reasonable and efficient treatment therewith (e.g., of at least 3 hour) can be beneficially used for regulating 25 food intake and caloric intake in a human subject and hence for treating medical and psychological conditions associated with overweight. Without being bound to any particular theory, it is assumed that betahistine, as well as other H<sub>1</sub> receptor agonists, decrease food intake by a different mode of action than the currently used anti-obesity medications.

30 Thus, according to one aspect of the present invention there is provided a method of treating of a condition in which regulating a food intake in a human subject is beneficial, which is effected by administering to the subject a therapeutically

effective amount of an H<sub>1</sub> agonist that has a pharmacological half-life of at least 3 hours.

It should be noted that as used herein the term "treatment" also includes amelioration or alleviation of a pathological condition and/or one or more symptoms thereof, curing such a condition, or preventing the genesis of such a condition.

The pharmacological half-life of the H<sub>1</sub> agonist utilized in this and other aspects of the present invention preferably ranges from about 3 hours to about 12 hours, more preferably from about 3 hours to about 8 hours, and even more preferably from about 3 hours to about 5 hours. Such a pharmacological half life is highly advantageous since, as is discussed hereinabove, such a drug remains in the blood for longer periods than, for example, histamine, achieving steadier blood levels, and therefore fewer side effects. The H<sub>1</sub> agonist of the present invention therefore needs to be administered far less frequently than, for example, histamine, which has a half life of only about 2 minutes. Since, as is known in the art, 97 % of a drug is eliminated after 5 half lives, the administration should be repeated at intervals of less than 5 half lives, and usually less, depending on various parameters, including the clearance rate and the initial concentration administered.

In clinical practice, the choice of the dosage interval usually represents a compromise between the desirability of minimizing the variations of effectiveness between doses and patient inconvenience from too frequent dosing, which results in poor compliance. Dosage regimes of one or two administrations per day are considered optimal.

Using a drug having a half life within the above cited range allows for selecting the administration times so as to ensure maximal effectiveness of the drug at times of day when the effect is most required, such as at known meal times, periods of the day at which the subject most commonly experiences significant levels of hunger, etc., and to be least effective at times of day when no effect is necessary, such as, for example, during periods when the subject is expected to be asleep.

Drugs having a very long half life have the disadvantage that once or twice daily administration is clearly not possible. Less than daily administration is associated with poor patient compliance, as the subject tends to forget the necessity to take a medication which is not part of his daily routine. For example, the histamine-

N-methyltransferase inhibitor, metoprone, which, as discussed above, has been shown to decrease food intake in rats, both by intraperitoneal injection and by central infusion, has a half life of 216 hours. Furthermore, the issue of toxicity must be considered with drugs having such long half lives.

5 In addition, it is known that a prolonged presence of H<sub>1</sub> agonists may lead to downregulation of the H<sub>1</sub> receptor.

Downregulation of a particular receptor after continuous activation can result in decreased response to agonist administration, due to adaptive changes in the receptors that limit their subsequent responsiveness, or to decrease in the number or 10 density of receptors. These receptor-specific changes include a rapid uncoupling of receptors from activation of their cognate G proteins, mediating functional desensitization; a rapid redistribution of receptors into relatively inaccessible compartments in the plasma membrane or inside the cell, variously referred to as sequestration, endocytosis, or internalization; with prolonged agonist exposure, termed 15 down-regulation. Considerable progress has been made in recent years in identifying the receptor modifications involved in these changes (*Krupnick et al.*, 1998), although many of the details of the molecular modifications and protein-protein interactions that are involved in bringing about these changes remain to be determined. Down-regulation of a receptor type may lead to desensitization or tolerance to the agonist for 20 the receptor. It is therefore believed that use of an H<sub>1</sub> receptor agonist having a half-life of less than 12 hours, preferably less than 8 hours and more preferably less than 5 hours prevents continuous exposure of the receptor to the agonist, and thereby down-regulation of the receptor is avoided.

The H<sub>1</sub> agonist of the present invention is preferably also an H<sub>3</sub> antagonist. H<sub>3</sub> 25 antagonists have been shown to have an effect on regulation of food intake. It has been suggested that the inhibition of H<sub>3</sub> receptor activity increases histamine release and synthesis. Histamine subsequently increases histaminergic neuron activity via H<sub>1</sub> receptors and in this way inhibit food intake. H<sub>3</sub> receptors are located not only on histaminergic neurons, but also nonhistaminergic neurons as heteroreceptors, and 30 modulate the release of 5-HT and noradrenaline. Thus, the effects of H<sub>3</sub> ligands on food intake may express through other endogenous substances (*Morimoto et al.*, 2001).

Also preferably, the H<sub>1</sub> agonist is characterized by blood brain barrier permeability, and therefore is able to cross the blood brain barrier and enter brain tissue, thereby acting on central H<sub>1</sub> and H<sub>3</sub> receptors. This enables the agonist to be administered by the systemic route, in contrast to, for example, histamine, which has 5 very poor blood brain barrier permeability and has therefore previously been administered as an appetite suppressant by intracerebroventricular injection.

The H<sub>1</sub> agonist of the present invention can be administered as any pharmaceutically acceptable salt, such as, for example, dihydrochloride, mesilate, or trimebutine maleate. Alternatively, the drug can be administered as any metabolite, 10 prodrug, or derivative, or combination thereof.

The phrase "pharmaceutically acceptable salt" refers to a charged species of the parent compound and its counter ion, which is typically used to modify the solubility characteristics of the parent compound and/or to reduce any significant irritation to an organism by the parent compound, while not abrogating the biological 15 activity and properties of the administered compound.

The term "prodrug" refers to an agent, which is converted into the active compound (the active parent drug) *in vivo*. Prodrugs are typically useful for facilitating the administration of the parent drug. They may, for instance, be bioavailable by oral administration whereas the parent drug is not. The prodrug may 20 also have improved solubility as compared with the parent drug in pharmaceutical compositions.

As used herein, the term "metabolite" describes the actual active moiety of the compound which is formed as a result of metabolic processes that occur *in vivo* upon administration of the compound.

25 The term "derivative" describes the result of chemically altering, modifying or changing a compound or a portion thereof, such that it maintains its original functionality in at least one respect.

According to a preferred embodiment of the present invention the H<sub>1</sub> agonist is betahistine. Betahistine is a structural analog of histamine, which has been shown to 30 have a binding affinity profile which is qualitatively close to that of histamine for both H<sub>1</sub> and H<sub>3</sub> receptors (*Fossati et al., 2001*). Furthermore, it is pharmacologically relevant, since pharmacokinetic data have shown that therapeutic dosages of

betahistine in humans give plasma concentrations which fall in the same range as its affinities for  $H_1$  and  $H_3$  receptors. The pharmacological relevance of betahistine for  $H_1$  receptors is further confirmed *in vivo* in animals and humans, attributable to the increased blood flow in microcirculation in the auditory and internal vestibular system (Meyer *et al.*, 1974). Since betahistine has already been widely used for treating vertigo and other disorders, it has been tested and is approved for use in human medicine.

Betahistine is readily absorbed through the oral route and is converted to at least two metabolites, 2-(2-aminoethyl)-pyridine and 2-(2-hydroxyethyl)-pyridine. 10 Betahistine has an elimination half life time of 3.5 hours, and most of the dose is excreted via the urine as metabolites. Hence, betahistine may be administered according to a convenient dosage regime, as discussed above.

15 The side effects associated with betahistine are minor, consisting mainly of skin rashes of various types, urticaria, itching, gastric upset, nausea, headache, and exacerbation of symptoms in patients with a history of peptic ulcer.


Betahistine can be utilized in this and other aspects of the present invention either *per se* or as a metabolite, a pharmaceutically acceptable salt, a prodrug or a derivative thereof.

20 Betahistine metabolites that can be efficiently used in the context of the present invention include, for example, 2-(2-aminoethyl)-pyridine and 2-(2-hydroxyethyl)-pyridine. As is further described in the art (see, for example, Fossati *et al.*, 2001) these metabolites are also characterized by  $H_1$ -receptor agonist activity and therefore can also be utilized in the various aspects of the present invention.

25 Representative examples of betahistine pharmaceutically acceptable salts that can be efficiently used in the context of the present invention include, without limitation, betahistine hydrochloride, betahistine dihydrochloride, betahistine mesilate, and betahistine trimebutine maleate.

30 Betahistine derivatives that are suitable for use in the context of this and other aspects of the present invention include, for example, compounds having the general Formula I:

21



Formula I

5 In Formula I, each of R<sub>1</sub>-R<sub>12</sub> is preferably independently selected from the group consisting of hydrogen, alkyl, cycloalkyl and aryl and any combination thereof.

10 Optionally, each of R<sub>1</sub>-R<sub>12</sub> can also be selected from other substituents, as long as features such as the blood brain barrier permeability, the half life and the binding to the H<sub>1</sub>-receptor of the compound are not adversely affected. Thus, each of R<sub>1</sub>-R<sub>12</sub> can be further independently selected, for example, from alkenyl, alkynyl, alkoxy, aryloxy, hydroxy, thiohydroxy, thioalkoxy, thioaryloxy, halide, amino, nitro, cyano, carbonyl, C-carboxy, O-carboxy, C-carmabyl, N-carbamyl, sulfonyl, sulfinyl, sulfonamide, urea, thiourea, guanidine, guanyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, N-sulfonamido, and S-sulfonamido or, 15 alternatively, at least two of R<sub>1</sub>-R<sub>4</sub> and/or at least two of R<sub>5</sub>-R<sub>12</sub> form at least one four-, five- or six-membered aromatic, heteroaromatic, alicyclic or heteroalicyclic ring.

20 It will be appreciated by one of skills in the art that the feasibility of each of the substituents (R<sub>1</sub>-R<sub>12</sub>) to be located at the indicated positions depends on the valency and chemical compatibility of the substituent, the substituted position and other substituents. Hence, the present invention is aimed at encompassing all the feasible substituents for any position.

25 As used herein, the term "alkyl" refers to a saturated aliphatic hydrocarbon including straight chain and branched chain groups. Preferably, the alkyl group has 1 to 20 carbon atoms. Whenever a numerical range; *e.g.*, "1-20", is stated herein, it implies that the group, in this case the alkyl group, may contain 1 carbon atom, 2 carbon atoms, 3 carbon atoms, etc., up to and including 20 carbon atoms. More preferably, the alkyl is a medium size alkyl having 1 to 10 carbon atoms. Most

preferably, unless otherwise indicated, the alkyl is a lower alkyl having 1 to 4 carbon atoms. The alkyl group may be substituted or unsubstituted.

A "cycloalkyl" group refers to an all-carbon monocyclic or fused ring (*i.e.*, rings which share an adjacent pair of carbon atoms) group wherein one of more of the 5 rings does not have a completely conjugated pi-electron system. Examples, without limitation, of cycloalkyl groups are cyclopropane, cyclobutane, cyclopentane, cyclopentene, cyclohexane, cyclohexadiene, cycloheptane, cycloheptatriene, and adamantane. A cycloalkyl group may be substituted or unsubstituted.

An "alkenyl" group refers to an alkyl group which consists of at least two 10 carbon atoms and at least one carbon-carbon double bond.

An "alkynyl" group refers to an alkyl group which consists of at least two carbon atoms and at least one carbon-carbon triple bond.

An "aryl" group refers to an all-carbon monocyclic or fused-ring polycyclic (*i.e.*, rings which share adjacent pairs of carbon atoms) groups having a completely 15 conjugated pi-electron system. Examples, without limitation, of aryl groups are phenyl, naphthalenyl and anthracenyl. The aryl group may be substituted or unsubstituted.

A "heteroaryl" group refers to a monocyclic or fused ring (*i.e.*, rings which share an adjacent pair of atoms) group having in the ring(s) one or more atoms, such 20 as, for example, nitrogen, oxygen and sulfur and, in addition, having a completely conjugated pi-electron system. Examples, without limitation, of heteroaryl groups include pyrrole, furane, thiophene, imidazole, oxazole, thiazole, pyrazole, pyridine, pyrimidine, quinoline, isoquinoline and purine.

A "heteroalicyclic" group refers to a monocyclic or fused ring group having in 25 the ring(s) one or more atoms such as nitrogen, oxygen and sulfur. The rings may also have one or more double bonds. However, the rings do not have a completely conjugated pi-electron system. The heteroalicyclic may be substituted or unsubstituted. Representative examples are piperidine, piperazine, tetrahydrofuran, tetrahydropyran, morpholino and the like.

30 A "hydroxy" group refers to an -OH group.

An "alkoxy" group refers to both an -O-alkyl and an -O-cycloalkyl group, as defined herein.

An "aryloxy" group refers to both an -O-aryl and an -O-heteroaryl group, as defined herein.

A "thiohydroxy" group refers to a -SH group.

5 A "thioalkoxy" group refers to both an -S-alkyl group, and an -S-cycloalkyl group, as defined herein.

A "thioaryloxy" group refers to both an -S-aryl and an -S-heteroaryl group, as defined herein.

A "carbonyl" group refers to a -C(=O)-R' group, where R' is hydrogen, alkyl, alkenyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) or heteroalicyclic (bonded through a ring carbon) as defined herein.

10 A "thiocarbonyl" group refers to a -C(=S)-R' group, where R' is as defined herein for R'.

A "C-carboxy" group refers to a -C(=O)-O-R' group, where R' is as defined herein.

15 An "O-carboxy" group refers to an R'C(=O)-O- group, where R' is as defined herein.

A "halide" group refers to fluorine, chlorine, bromine or iodine.

A "sulfinyl" group refers to an -S(=O)-R' group, where R' is as defined herein.

20 A "sulfonyl" group refers to an -S(=O)<sub>2</sub>-R' group, where R' is as defined herein.

An "S-sulfonamido" group refers to a -S(=O)<sub>2</sub>-NR'R" group, with R' is as defined herein and R" is as defined for R'.

An "N-sulfonamido" group refers to an R'S(=O)<sub>2</sub>-NR" group, where R' and R" are as defined herein.

25 An "O-carbamyl" group refers to an -OC(=O)-NR'R" group, where R' and R" are as defined herein.

An "N-carbamyl" group refers to an R"OC(=O)-NR'- group, where R' and R" are as defined herein.

30 An "O-thiocarbamyl" group refers to an -OC(=S)-NR'R" group, where R' and R" are as defined herein.

An "N-thiocarbamyl" group refers to an R"OC(=S)NR'- group, where R' and R" are as defined herein

An "amino" group refers to an  $-NR'R''$  group where  $R'$  and  $R''$  are as defined herein.

A "C-amido" group refers to a  $-C(=O)-NR'R''$  group, where  $R'$  and  $R''$  are as defined herein.

5 An "N-amido" group refers to an  $R'C(=O)-NR''$  group, where  $R'$  and  $R''$  are as defined herein.

A "urea" group refers to an  $-NR'C(=O)-NR''R'''$  group, where  $R'$  and  $R''$  are as defined herein and  $R'''$  is defined as either  $R'$  or  $R''$ .

10 A "guanidino" group refers to an  $-R'NC(=N)-NR''R'''$  group, where  $R'$ ,  $R''$  and  $R'''$  are as defined herein.

A "guanyl" group refers to an  $R'R''NC(=N)-$  group, where  $R'$  and  $R''$  are as defined herein.

A "nitro" group refers to an  $-NO_2$  group.

A "cyano" group refers to a  $-C\equiv N$  group.

15 The term "phosphonyl" describes a  $-O-P(=O)(OR')(OR'')$  group, with  $R'$  and  $R''$  as defined hereinabove.

The term "phosphinyl" describes a  $-PR'R''$  group, with  $R'$  and  $R''$  as defined hereinabove.

20 The term "thiourea" describes a  $-NR'-C(=S)-NR''R'''$  group, with  $R'$  and  $R''$  as defined hereinabove and  $R'''$  as defined herein for  $R'$  and  $R''$ .

25 In any of the methods described herein, the  $H_1$  agonist can be administered by any route selected from the oral, transdermal, intravenous, subcutaneous, intramuscular, intranasal, intraauricular, sublingual, rectal, transmucosal, intestinal, buccal, intramedullar, intrathecal, direct intraventricular, intraperitoneal, or intraocular routes. Preferably, the route of administration is selected from the oral, transdermal, buccal, transmucosal, rectal or sublingual routes. More preferably, the  $H_1$  agonist is administered using the oral, buccal or transdermal route.

30 The  $H_1$  agonist is optionally and preferably administered as a total dose of from about 2 mg to about 96 mg per day. More preferably, the total dose is from about 5 mg to about 50 mg per day, more preferably from about 10 mg to about 50 mg, more preferably from about 16 mg to about 48 mg and most preferably it is from about 24 mg to 48 mg.

The H<sub>1</sub> agonist is preferably administered once or several times a day, for example from about 1 to about 4 times per day, and, more preferably, twice per day. Alternatively, the H<sub>1</sub> agonist may be administered according to the development of hunger of the subject.

5 As is demonstrated in the Examples section that follows, clinical studies have indicated that administering a 16 mg dose of betahistine twice a day for a month, without restricting their food intake, resulted in a significant weight loss, particularly as compared with a control group.

10 Hence, according to an embodiment of the present invention, a treatment regime is performed such that upon repetitious administration, a decrease of the body weight of the subject that ranges from about 1 to about 5 percent is effected, without restricting the food intake of the subject.

15 Since, as is discussed in detail hereinabove, constant presence of a H<sub>1</sub> agonist may result in substantial downregulation of the H<sub>1</sub> receptor, a more preferred treatment regime according to the present invention is performed such that such a decrease in a body weight is achieved without effecting downregulation of the H<sub>1</sub> receptor. Such a treatment regime can be designed by adjusting the dosing and the administration intervals to the half life of the selected H<sub>1</sub> agonist.

20 The dosage may vary depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See e.g., Fingl, et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1). Dosage amount and interval may be adjusted individually to provide plasma levels of the H<sub>1</sub> agonist which are sufficient to maintain the regulating effects.

25 When the H<sub>1</sub> agonist is betahistine, which is commercially available as tablets of 8 mg or 16 mg, the daily dosage range is from 24 to 48 mg, administered orally in divided doses. For example, the 8 mg tablet is administered as 1 to 2 tablets 3 times daily, and the 16 mg tablet is administered as 0.5 to 1 tablet 3 times daily.

30 The H<sub>1</sub> agonist of the present invention may optionally be administered in the form of a slow-release preparation, having a reduced rate of release of the active substance, in order to further increase patient convenience and compliance and optionally the efficiency of the active agent. The slower the rate of release, the less

the blood concentrations fluctuate within a dosing interval. This enables higher doses to be given less frequently, while maintaining therapeutic concentrations over prolonged periods. Furthermore, slow-release preparations are beneficial in reducing potential side-effects of the active ingredient due to transiently high peak blood 5 concentrations being reached soon after administration.

Slow release preparations typically include slow release biodegradable carriers. Slow release biodegradable carriers are well known in the art. These are materials that may form particles that may capture therein an active compound(s) and slowly degrade/dissolve under a suitable environment (e.g., aqueous, acidic, basic, etc.) and 10 thereby degrade/dissolve in body fluids and release the active compound(s) therein. The particles are preferably nanoparticles (i.e., in the nanometer range, e.g., in the range of about 1 to about 500 nm in diameter, preferably about 50-200 nm in diameter, most preferably about 100 nm in diameter).

The rate at which a drug is released is generally dependent on the rate at which 15 the dosage form disintegrates or dissolves. Disintegration greatly increases the drug's surface area in contact with GI fluids, thereby promoting drug dissolution and absorption. Disintegrants and other excipients (e.g., diluents, lubricants, surfactants, binders, dispersants) are often added during manufacture to facilitate these processes. Surfactants increase the dissolution rate by increasing the wettability, solubility, and 20 dispersibility of the drug. Disintegration of solid forms may be retarded by excessive pressure applied during the tableting procedure or by special coatings applied to protect the tablet from the digestive processes of the gut. Hydrophobic lubricants (e.g., magnesium stearate) may bind to the active drug and reduce its bioavailability.

Dissolution rate determines the availability of the drug for absorption. When 25 slower than absorption, dissolution becomes the rate-limiting step. Overall absorption can be controlled by manipulating the formulation. For example, reducing the particle size increases the drug's surface area, thus increasing the rate and extent of GI absorption of a drug whose absorption is normally limited by slow dissolution. Dissolution rate is affected by whether the drug is in salt, crystal, or hydrate form.

30 Oral slow-release forms are often designed to maintain therapeutic drug concentrations for greater than 12 hours. The absorption rate can be controlled by coating drug particles with wax or other water-insoluble material, by embedding the

drug in a matrix from which it is released slowly during transit through the GI tract, or by complexing the drug with ion-exchange resins.

Thus, for example, a slow-release formulation in tablet form, may be based on the use of a hydrophilic polymer which swells in contact with gastrointestinal fluids, to form a gel, which creates a barrier that enrobes the tablet. The barrier limits physical exchanges between the inside of the tablet and the surrounding medium. As a consequence, intrusion of water towards the tablet matrix and diffusion of drug are slowed down, allowing a controlled slow release of the drug.

Various types of polymers may be used as a matrix for the slow-release of drugs, such as polyvinyl chloride, polyethylene polyamides, ethylcellulose, silicone, poly (hydroxyethyl methacrylate), other acrylic co-polymers, and polyvinylacetate-polyvinyl chloride copolymers.

Thus, a slow-release formulation for delivery of the H<sub>1</sub> agonist of the present invention provides for release over a period that ranges from about 2 hour to about 24 hours, preferably from about 4 hours to about 24 hours and hence, for release over a period of at least 4 hour, at least 5 hours, at least 6 hours, at least 7 hours, at least 8 hours, at least 9 hours, at least 10 hours, at least 11 hours, at least 12 hours, at least 13 hours, at least 14 hours, at least 15 hours, at least 16 hours, at least 17 hours, at least 18 hours, at least 19 hours, at least 20 hours, at least 21 hours, at least 22 hours, at least 23 hours, or at least 24 hours. Alternatively, such a slow-release formulation provides for release of the H<sub>1</sub> agonist over a period of more than 24 hours and up to 48 hours.

The method according to this aspect of the present invention can thus be efficiently used for treating any condition in which regulating food intake is beneficial. These include, for example, overeating, overweight, obesity, and disorders caused or exacerbated thereby.

Diseases caused by or exacerbated by these conditions include, for example, muscoskeletal disorders (such as osteoarthritis, spine-related pains, etc.), cardiovascular disorders (such as hypertension, arteriosclerosis, etc.), dermatological disorders (such as fungal and other infections), sleep disorder (such as snoring and obstructive sleep apnea), metabolic conditions (dyslipidemia, lipemia or hypercholesterolemia), diabetes and diabetes-related problems, as well as cancer

(particularly breast, prostate and colon cancer). The effect of overweight, expressed by the body mass index, on the relative risk of death from various disorders caused or exacerbated thereby, is demonstrated in Figure 3. As shown in Figure 3, cardiovascular disease increases significantly in overweight subjects, both male and 5 female, as compared to those within the normal body-mass index range (i.e. from 18.5 to 24.9). A further sharp increase is observed in subjects classified as obese (i.e. having body mass of greater than 30). Similarly, an increase in the relative risk of death from cancer is seen in overweight subjects, particularly in obese subjects, with a greater increase amongst females. The relative risk of death from other causes is seen 10 to increase sharply in obese individuals.

Additional conditions that are treatable by the method according to this aspect of the present invention include, for example, conditions that are associated with a psychological factor, such as binge eating disorder, night eating syndrome, obsessive eating, compulsive eating and bulimia.

15 Alternatively, the condition may be associated with a drug treatment, such as treatment with a steroid hormone or a psychoactive drug (such as traditional antidepressants, benzodiazepines, lithium, and antipsychotic drugs), which is oftentimes associated with weight gain.

As is described in the art, the histaminergic system has been implicated in 20 weight gain associated with steroid-induced and antipsychotic-induced weight gain (*Poyurovsky et al., 2005*). Hence, the H<sub>1</sub>-receptor agonists of the present invention may be co-administered together with a drug treatment which is known to cause appetite stimulation and weight gain in humans due to its histaminergic activity. It is assumed that the H<sub>1</sub> agonist utilized herein interferes with appetite stimulating 25 histaminergic activity of the steroid or anti-psychotic drug mentioned above and thus prevents or reduces the weight gain caused by such drug treatment.

The H<sub>1</sub>-receptor agonists described herein can be further efficiently used for improving a compliance of a human subject to caloric restriction. Thus, according to another aspect of the present invention, there is provided a method of improving a 30 compliance of a human subject to caloric restriction, which is effected by administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist as described herein. The method, according to this aspect of the present invention can be

beneficially practiced with human subject undergoing a weight reducing diet or any other caloric restriction.

By efficiently regulating food intake in human subjects and improving a compliance thereof to caloric restriction, the methods described hereinabove may further be used for inducing weight loss, for maintaining weight loss or preventing a weight gain after or during a weight reducing diet, or for preventing weight gain in a subject having a condition associated with weight gain. For inducing or maintaining weight loss, the H<sub>1</sub> receptor agonist of the present invention may optionally be administered in conjunction with other methods of treatment, such as diet, exercise, behavioral therapy, drug therapy, or surgical therapy.

The H<sub>1</sub>-receptor agonists described herein can be further efficiently used for reducing a desire of a human subject to consume fats. Some neurotransmitters and neuromodulators have been shown to control both amount eaten and selection of food in a macronutrient specific manner (*Lecklin et al.*, 2002). For example, as is discussed hereinabove and is further demonstrated in Figures 1 and 2, metoprine has been shown to suppress daily ingestion of fats in rats, while having no effect on intakes of carbohydrates or proteins, indicating that an increase in the brain histamine content may specifically affect the desire to consume fats.

As is demonstrated in the Examples section that follows, it has been found that betahistine treatment resulted in a significant decrease in fat consumption in human subjects over a 28 day trial, while consumption of carbohydrates was not significantly altered. The increase in brain histamine caused by administration of the H<sub>1</sub>-receptor agonists of the present invention may therefore be effective in reducing the desire of a human subject to consume fats.

Thus, according to still another aspect of the present invention, there is provided a method of reducing the desire of a human subject to consume fats, which is effected by administering to the subject a therapeutically effective amount of the H<sub>1</sub> agonist described herein.

The clinical finding that the H<sub>1</sub> agonist of the present invention reduces fat intake in humans is of unique importance, not only with regard to obese subjects, but also with regard to patients suffering from conditions that are associated with fat consumption and/or in which reduced fat consumption is beneficial.

Thus, according to a further aspect of the present invention, there is provided a method of treating a condition in which reduced fat consumption is beneficial, which is effected by administering to a subject in need thereof a therapeutically effective amount of an H<sub>1</sub> agonist, as described herein.

5 Reducing the fat consumption of a subject can be further utilized for treating conditions associated with metabolic derangement.

Thus, according to still another aspect of the present invention, there is provided a method of treating a condition associated with a metabolic derangement in a human subject, which is effected by administering to the subject a therapeutically effective amount of the H<sub>1</sub> agonist described herein.

10 Such conditions are typically associated with metabolic derangement and more particularly with adverse imbalance of metabolites such as total cholesterol, HDL-cholesterol, LDL-cholesterol, triglycerides and the like and include, for example, dyslipidemia, such as hypercholesterolemia or lipemia.

15 Dyslipidemias are disorders of lipoprotein metabolism, including lipoprotein overproduction or deficiency. These disorders may be manifested by elevation of the serum total cholesterol, low-density lipoprotein (LDL) cholesterol and triglyceride concentrations, and a decrease in the high-density lipoprotein (HDL) cholesterol concentration and therefore include, for example, lipemia and hypercholesterolemia.

20 Lipemia is a condition in which an excess of fats or lipids is found in the blood of subject.

Hypercholesterolemia is a condition in which high levels of cholesterol are found in the blood of a subject.

25 A direct connection has been shown between coronary artery disease and dyslipidemia. Dyslipidemias may be genetic in origin, or may be due to dietary factors (such as excess saturated fats, "trans" fatty acids, cholesterol, excess calories, or alcohol) or drug use (including steroid hormones, diuretics, beta-blockers, cyclosporine and amiodarone, or olanzapine). Dyslipidemias may also be associated with hypothyroidism, diabetes mellitus, hepatobiliary obstruction, nephritic syndrome and chronic renal failure, or with systemic diseases such as porphyries, systemic lupus erythematosus, and lymphomas. The most commonly used options for pharmacologic treatment of dyslipidemia include fibrates, HMG-CoA reductase inhibitors (such as

statins), bile acid sequestrants, cholesterol absorption inhibitors, nicotinic acid and derivatives thereof.

High cholesterol level, as well as other manifestations of dyslipidemia, can cause the formation and accumulation of plaque deposits in the arteries, leading to 5 plaque ruptures and blockages in the arteries, which increase the risk for heart attack, stroke, circulation problems, and death. Patients suffering from metabolic derangement that is manifested by dyslipidemia are therefore typically instructed to follow a low fat diet as a first measure, before being prescribed drugs which affect cholesterol metabolism. However, patient compliance with these instructions is 10 generally not high, and currently there are no available drugs for helping patients to adhere to such a diet. Therefore, the H<sub>1</sub> agonist of the present invention, in reducing the desire of a patient to consume fats, helps such patients to maintain a low fat diet.

Hence, according to still another aspect of the present invention, there is provided a method of reducing total cholesterol level in a human subject, which is 15 effected by administering to the subject a therapeutically effective amount of the H<sub>1</sub> agonist described herein.

Total cholesterol consists of HDL cholesterol and LDL cholesterol. High density lipoprotein (HDL), commonly referred to as "good" cholesterol, tends to carry cholesterol away from the arteries and back to the liver, where it is passed from the 20 body. It is also believed that HDL cholesterol removes excess cholesterol from plaque in arteries, thus slowing the buildup. An increase in HDL cholesterol is therefore considered beneficial. In contrast, LDL cholesterol is considered to be "bad" cholesterol, being responsible for plaque formation and deposits. Reducing LDL-cholesterol is believed to be responsible for reducing or stopping the formation of new 25 cholesterol plaques on the artery walls; reducing existing cholesterol plaques on the artery walls; widening narrowed arteries; preventing the rupture of cholesterol plaques, which initiates blood clot formation; decreasing the risk of heart attacks; and decreasing the risk of strokes.

Thus, according to still another aspect of the present invention, there is 30 provided a method of reducing low-density lipoprotein cholesterol and increasing high-density lipoprotein cholesterol levels in a human subject, which is effected by

administering to the subject a therapeutically effective amount of the H<sub>1</sub> agonist described herein.

Triglycerides are common types of fats (lipids) that are essential for good health when present in normal amounts. They account for about 95 percent of the body's fatty tissue. Triglycerides are both present in food and manufactured by the body. Abnormally high triglyceride levels are associated with a number of diseases and conditions, such as cirrhosis, underactive thyroid, poorly controlled diabetes, and pancreatitis. High triglyceride levels are also associated with known risk factors for heart disease, such as low levels of HDL cholesterol, high levels of LDL cholesterol and obesity. Triglycerides may also contribute to atherosclerosis.

Hence, according to still another aspect of the present invention, there is provided a method of reducing triglyceride level in a human subject, which is effected by administering to the subject a therapeutically effective amount of the H<sub>1</sub> agonist described herein.

In any of the methods described herein, the H<sub>1</sub> agonist may optionally be administered together with a therapeutically effective amount of an additional active agent that may affect the condition being treated.

According to a preferred embodiment of the present invention, the additional active ingredient can be, for example, a weight control agent. Any of the presently known and approved weight control agents or related substances can be used according to this embodiment. Hence, representative examples of such active ingredients include, for example, lipase inhibitors. A non-limiting example of a lipase inhibitor suitable for co-administration with the H<sub>1</sub> agonist of the present invention is orlistat, which acts by binding to gastrointestinal lipases in the lumen of the gut, preventing hydrolysis of dietary fat into absorbable free fatty acids and monoacylglycerols. Orlistat is currently the only FDA-approved medication for obesity that reduces nutrient absorption.

Additional examples of such active ingredients include selective serotonin reuptake inhibitors (SSRIs). A non-limiting example of an SSRI suitable for co-administration with the H<sub>1</sub> agonist of the present invention is Sibutramine, an inhibitor of both norepinephrine reuptake and serotonin reuptake that also weakly

inhibits dopamine reuptake, which is approved by the FDA for weight loss and weight maintenance in conjunction with a reduction diet.

Additional examples of such active ingredients include amphetamines, although the administration thereof might be restricted to a limited time period. A 5 non-limiting example of an amphetamine suitable for co-administration with the H<sub>1</sub> agonist of the present invention is Fentermine.

Additional examples of such active ingredients include cannabinoid receptor antagonists. A non-limiting example of a cannabinoid receptor antagonist suitable for co-administration with the H<sub>1</sub> agonist of the present invention is Rimonabant, a 10 recently developed drug that is currently undergoing Phase III trials and which is claimed to stop food cravings.

Alternatively, the additional active agent may be an agent for the treatment of a musculoskeletal disorder, a cardiovascular disorder, a dermatological disorder, a sleep disorder, a metabolic condition, dyslipidemia (including hypercholesterolemia and 15 lipemia), diabetes or a diabetes-related condition.

Representative examples of agents for the treatment of musculoskeletal disorders include, without limitation, anti-inflammatory agents, including non-steroidal anti-inflammatory drugs, muscle relaxants, anti-gout agents (such as allopurinol, colchicine, and uricosuric drugs), immunosuppressants (such as 20 glucocorticoids, gold, and cytotoxic agents) and drugs affecting bone mineralization (e.g., diphosphonates, calcitonin, estrogen analogs).

Representative examples of non-steroidal anti-inflammatory drugs include, without limitation, piroxicam, isoxicam, tenoxicam, sudoxicam, and CP-14,304; salicylates, such as aspirin, disalcid, benorylate, trilisate, safapryn, solprin, diflunisal, 25 and fendosal; acetic acid derivatives, such as diclofenac, fenclofenac, indomethacin, sulindac, tolmetin, isoxepac, furofenac, tiopinac, zidometacin, acematacin, fentiazac, zomepirac, clindanac, oxepinac, felbinac, and ketorolac; fenamates, such as mefenamic, meclofenamic, flufenamic, niflumic, and tolfenamic acids; propionic acid derivatives, such as ibuprofen, naproxen, benoxaprofen, flurbiprofen, ketoprofen, 30 fenoprofen, fenbufen, indoprofen, pirprofen, carprofen, oxaprozin, pranoprofen, mioprofen, tioxaprofen, suprofen, alminoprofen, and tiaprofenic; pyrazoles, such as phenylbutazone, oxyphenbutazone, feprazole, azapropazone, and trimethazone.

Non-limiting examples of steroid anti-inflammatory drugs include, without limitation, corticosteroids such as hydrocortisone, hydroxyltriamcinolone, alpha-methyl dexamethasone, dexamethasone-phosphate, beclomethasone dipropionate, clobetasol valerate, desonide, desoxymethasone, desoxycorticosterone acetate, 5 dexamethasone, dichlorisone, diflorasone diacetate, diflucortolone valerate, fludrenolone, fluclorolone acetonide, fludrocortisone, flumethasone pivalate, fluosinolone acetonide, fluocinonide, flucortine butylesters, fluocortolone, fluprednidene (fluprednylidene) acetate, flurandrenolone, halcinonide, hydrocortisone acetate, hydrocortisone butyrate, methylprednisolone, triamcinolone acetonide, 10 cortisone, cortodoxone, flucetonide, fludrocortisone, difluorosone diacetate, fluradrenolone, fludrocortisone, difluorosone diacetate, fluradrenolone acetonide, medrysone, amcinafel, amcinafide, betamethasone and the balance of its esters, chloroprednisone, chlorprednisone acetate, clocortelone, clescinolone, dichlorisone, 15 diflurprednate, flucoronide, flunisolide, fluoromethalone, fluperolone, fluprednisolone, hydrocortisone valerate, hydrocortisone cyclopentylpropionate, hydrocortamate, meprednisone, paramethasone, prednisolone, prednisone, beclomethasone dipropionate, triamcinolone, and mixtures thereof.

Representative examples of agents for the treatment of cardiovascular disorder include, without limitation, alpha-adrenergic blocking drugs (such as doxazocin, 20 prazocin or terazosin); angiotensin-converting enzyme inhibitors (such as captopril, enalapril, or lisinopril); antiarrhythmic drugs (such as amiodarone); anticoagulants, antiplatelets or thrombolytics (such as aspirin); beta-adrenergic blocking drugs (such as acebutolol, atenolol, metoprolol, nadolol, pindolol or propanolol); calcium channel blockers (such as diltiazem, nicardipine, verapamil or nimopidipine); centrally acting 25 drugs (such as clonidine, guanfacine or methyldopa); digitalis drugs (such as digoxin); diuretics (such as chlorthalidone); nitrates (such as nitroglycerin); peripheral adrenergic antagonists (such as reserpine); and vasodilators (such as hydralazine).

Representative examples of agents for the treatment of sleep disorders include, without limitation, hypnotic medications, such as benzodiazepines (including 30 flurazepam, estazolam, temazepam and triazolam); zaleplon and zolpidem; eszopiclone; antidepressants, such as trazodone, antihistamines (such as diphenhydramine products).

Representative examples of agents for the treatment of diabetes include, without limitation, a sulfonylurea (such as chlorpropamide, glipizide, glyburide, and glimepiride) a meglitinide (such as repaglinide or nateglinide), a biguanide (such as metformin), a thiazolidinedione (such as rosiglitazone, troglitazone or pioglitazone), 5 and an alpha-glucosidase inhibitor (such as acarbose or meglitol) and insulin.

Non-limiting examples of agents for the treatment of dyslipidemia include fibrates, HMG-CoA reductase inhibitors, bile acid sequestrants, cholesterol absorption inhibitors, cholesterol biosynthesis inhibitors, nicotinic acid and derivatives thereof.

HMG-CoA reductase inhibitors (statins) are well known drugs that effectively 10 reduce LDL-cholesterol levels by inhibiting the enzyme that regulates the rate of cholesterol production and increasing the clearance of LDL-cholesterol present in the blood by the liver. Representative examples of commonly prescribed statins include Atorvastatin, Fluvastatin, Lovastatin, Pravastatin and Simvastatin.

Proliferative Activated Receptor (PPAR) agonists, also known as fibrates, 15 are fatty acid-activated members of the nuclear receptor superfamily that play important roles in lipid and glucose metabolism, and have been implicated in obesity-related metabolic diseases such as hyperlipidemia, insulin resistance, and coronary artery disease. Fibrates are generally effective in lowering elevated plasma triglycerides and cholesterol and act as PPAR agonists. The most pronounced effect 20 of fibrates includes a decrease in plasma triglyceride-rich lipoproteins (TRLs). Levels of LDL cholesterol (LDL-C) generally decrease in individuals with elevated baseline plasma concentrations, and HDL cholesterol (HDL-C) levels are usually increased when baseline plasma concentrations are low. Non-limiting examples of commonly prescribed fibrates include bezafibrate, gemfibrozil and fenofibrate.

25 Representative examples of cholesterol absorption inhibitors include ezetimibe. Ezetimibe is the first of a new class of cholesterol absorption inhibitors that potently and selectively inhibits dietary and biliary cholesterol absorption at the brush border of the intestinal epithelium, without affecting the absorption of triglyceride or fat-soluble vitamins. Ezetimibe thus reduces overall cholesterol 30 delivery to the liver, secondarily inducing increased expression of LDL receptors, resulting in an increased removal of LDL-C from the plasma.

Cholesterol absorption may also be affected by Cholesteryl Ester Transfer Protein (CETP) inhibitors, which play a major role in atherogenesis, by reducing cholesteryl ester accumulation within macrophages and the arterial wall, and thus reducing foam cell formation and affecting the cholesterol absorption. The most 5 promising presently known CETP inhibitor is avisimibe.

Representative examples of cholesterol biosynthesis inhibitors include squalene inhibitors (such as monooxygenase and synthase). Squalene is an isoprenoid compound structurally similar to beta-carotene, is an intermediate metabolite in the synthesis of cholesterol. In humans, about 60 percent of dietary squalene is absorbed. 10 It is transported in serum generally in association with very low density lipoproteins and is distributed ubiquitously in human tissues, with the greatest concentration in the skin, where it is one of the major components of skin surface lipids. Squalene inhibitors (e.g., monooxygenase and synthase) serve as cholesterol biosynthesis inhibitors.

15 Nicotinic acid is a known agent that lowers total cholesterol, LDL-cholesterol, and triglyceride levels, while raising HDL-cholesterol levels. There are three types of nicotinic acid drugs: immediate release, timed release, and extended release. Nicotinic acid or niacin, the water-soluble B vitamin, improves all lipoproteins when given in doses well above the vitamin requirement.

20 Additional active agents that can be utilized according to this embodiment of the present invention include, for example, analgesics, growth factors and toxins.

Non-limiting examples of analgesics (pain relievers) include aspirin and other salicylates (such as choline or magnesium salicylate), ibuprofen, ketoprofen, naproxen sodium, and acetaminophen.

25 Growth factors are hormones which have numerous functions, including regulation of adhesion molecule production, altering cellular proliferation, increasing vascularization, enhancing collagen synthesis, regulating bone metabolism and altering migration of cells into given area. Non-limiting examples of growth factors include insulin-like growth factor-1 (IGF-1), transforming growth factor- $\beta$  (TGF- $\beta$ ), 30 a bone morphogenic protein (BMP) and the like.

Non-limiting examples of toxins include the cholera toxin, which also serves as an adjuvant.

When utilized in each of the methods and aspects described above, the H<sub>1</sub> agonist of the present invention optionally and preferably forms a part of a pharmaceutical composition. The pharmaceutical composition comprises, in addition to the H<sub>1</sub> agonist, a pharmaceutically acceptable carrier, and may optionally further 5 comprise one or more components selected from binding agents, stabilizers, diluents, excipients, surfactants, flavors, and odorants.

The purpose of a pharmaceutical composition is to facilitate administration of an active ingredient (herein the H<sub>1</sub> agonist described above) to an organism (herein, a human being). Pharmaceutical compositions of the present invention may be 10 manufactured by processes well known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, 15 soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions.

Suitable pharmaceutical excipients include without limitation, calcium 20 carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils, polyethylene glycols, sodium stearate, glycerol monostearate, talc, sodium chloride, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of 25 solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like.

Further techniques for formulation and administration of active ingredients may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., 30 Easton, PA, latest edition, which is incorporated herein by reference as if fully set forth herein.

The pharmaceutical compositions herein described may also comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients

include, but are not limited to, calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin and polymers such as polyethylene glycols.

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more pharmaceutically acceptable carriers comprising excipients and auxiliaries, which facilitate processing of the active ingredients into preparations which, can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

Formulations for oral delivery can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E. W. Martin. Such compositions will contain a therapeutically effective amount of the compound, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should be suitable for the mode of administration.

For oral administration, the active ingredients can be formulated readily by combining the active ingredients with pharmaceutically acceptable carriers well known in the art. Such carriers enable the active ingredients of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for oral ingestion by a patient. Pharmacological preparations for oral use can be made using a solid excipient, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carbomethylcellulose; and/or physiologically acceptable polymers such as polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used which may optionally contain gum arabic,

talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active ingredient doses.

5 Pharmaceutical compositions, which can be used orally, include push-fit capsules made of gelatin as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules may contain the active ingredients in admixture with filler such as lactose, binders such as starches, lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft 10 capsules, the active ingredients may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for the chosen route of administration.

15 For transdermal administration, the composition can be formulated in a form of a gel, a cream, an ointment, a paste, a lotion, a milk, a suspension, an aerosol, a spray, a foam, a serum, a swab, a pledget, a pad or a patch. Formulations for transdermal delivery can typically include carriers such as water, liquid alcohols, liquid glycols, liquid polyalkylene glycols, liquid esters, liquid amides, liquid protein hydrolysates, liquid alkylated protein hydrolysates, liquid lanolin, lanolin derivatives, 20 glycerin, mineral oil, silicone, petroleum jelly, lanolin, fatty acids, vegetable oils, parabens, waxes, and like materials commonly employed in topical compositions. Various additives, known to those skilled in the art, may be included in the transdermal formulations of the invention. For example, solvents may be used to solubilize certain active ingredients substances. Other optional additives include skin 25 permeation enhancers, opacifiers, anti-oxidants, gelling agents, thickening agents, stabilizers, and the like.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

30 For administration by inhalation, the active ingredients for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from a pressurized pack or a nebulizer with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichloro-

tetrafluoroethane or carbon dioxide. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the active ingredient and a suitable powder base such as 5 lactose or starch.

The active ingredients described herein may be formulated for parenteral administration, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multidose containers with optionally, an added preservative. The compositions may be 10 suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical compositions for parenteral administration include aqueous solutions of the active preparation in water-soluble form. Additionally, suspensions of the active ingredients may be prepared as appropriate oily injection suspensions. 15 Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acids esters such as ethyl oleate, triglycerides or liposomes. Aqueous injection suspensions may contain substances, which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the 20 solubility of the active ingredients to allow for the preparation of highly concentrated solutions.

The composition can be formulated as rectal compositions such as suppositories or retention enemas, using, e.g., conventional suppository bases such as cocoa butter or other glycerides.

25 The pharmaceutical compositions herein described may also comprise suitable solid of gel phase carriers or excipients. Examples of such carriers or excipients include, but are not limited to, calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin and polymers such as polyethylene glycols.

Compositions of the present invention may, if desired, be presented in a pack or 30 dispenser device, such as an FDA approved kit, which may contain one or more unit dosage forms containing the active ingredient. The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be

accompanied by instructions for administration. The pack or dispenser may also be accompanied by a notice associated with the container in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the compositions or 5 human or veterinary administration. Such notice, for example, may be of labeling approved by the U.S. Food and Drug Administration for prescription drugs or of an approved product insert. Compositions comprising the H<sub>1</sub> agonist of the invention, and optionally other active ingredients, formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for 10 treatment of an indicated condition, as is detailed herein.

Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various 15 embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

### **EXAMPLES**

Reference is now made to the following example, which together with the 20 above descriptions, illustrate the invention in a non limiting fashion.

#### **EXAMPLE 1**

The following study was conducted to evaluate the effect of oral administration of betahistine on food intake:

25 Twenty obese but otherwise healthy persons were recruited. Their characteristics upon recruitment are shown in Table 1. Exclusion criteria for the study were age younger than 18, active diseases, medication use, known hypersensitivity or contra-indication for the use of betahistine.

Each subject was randomly allocated to receive betahistine 16 mg at 10:00 and 30 16:00 or placebo. Weight, caloric intake (24 hour recall) and appetite during the day (VAS, Visual Analogue Score) were obtained on day 0, 14 and 28 of the study. Subjects were instructed to eat according to their appetite without limitations.

Statistical significance was assessed with t-test. BMI stands for body mass index.

**Table 1. Patients' characteristics**

	Treatment	<b>betahistine</b>	<b>Placebo</b>
5	Age	48 $\pm$ 9	38 $\pm$ 15
	Weight (kg)	93 $\pm$ 17	90 $\pm$ 4
	BMI	35.1 $\pm$ 7.3	32.7 $\pm$ 1.7
	Mean caloric intake (kcal)	975 $\pm$ 472	1397 $\pm$ 693
10	Mean appetite score (VAS)	45 $\pm$ 15	50 $\pm$ 12

NS denotes no statistical significance (p value >0.05)

Of the 20 subjects recruited, 8 did not complete the study and were excluded from the final analysis:

15 Subject No. 3 (placebo) dropped out on day 3 due to side effects – weakness.

Subject No. 5 (betahistine) reported on day 5 food aversion that did not allow her to consume any food. She was instructed to reduce the dose by half and was excluded from the results analysis.

Subject No. 9 (placebo) was excluded for protocol violation.

20 Subject No. 11 (placebo) dropped out on week 4 due to flu.

Subject No. 13 (placebo) was lost to follow up.

Subject No. 15 (betahistine) dropped out on week 4 due to flu.

Subject No. 16 (placebo) dropped out on week 3 due to dyspnea.

Subject No. 17 (betahistine) was lost to follow up.

25

The effect of betahistine treatment on the total caloric intake, and on the specific consumption of fat, carbohydrates and protein of the participants in this study was also studied. The obtained data are presented in Figures 4 and 5.

As can be seen in Figure 4, treatment with betahistine was found to reduce 30 total caloric intake as compared to the placebo. The caloric intake decreased to 80 percents of the pre-treatment level at day 14 of treatment, and to 68 percents at day 28. No decrease was seen with the subjects receiving placebo.

As can be seen in Figure 5, while only a small reduction in carbohydrate consumption occurred with betahistine treatment (6 %), fat and protein consumption were reduced by 49 % and 35 %, respectively. At the same time patients treated with placebo increased their carbohydrates, fats and protein consumption by 24 %, 3 % and 5 37 % respectively.

It is therefore clearly shown that betahistine significantly reduces the caloric intake and fat consumption of a human subject. Therefore, administration of betahistine may be efficiently utilized for both improving compliance of a human subject to caloric restriction, and for reducing the desire of a human subject to 10 consume fat.

The effect of betahistine administration on weight change was also studied. The results are presented in Figure 6. Four out of seven subjects in the treatment group lost more than 1 kg during the study period, in comparison to only one out of five in the placebo group.

15 These clinical data clearly show that betahistine is an efficient medication for weight management.

#### ***EXAMPLE 2***

In a further study, a healthy, overweight woman was treated twice daily with 20 betahistine 16 mg for a month without any dietary changes. The level of certain metabolites of the woman, as observed in blood tests, was measured before and after the betahistine treatment and are presented in Table 2 below.

**Table 2**

	Day 0	Day 30	Change
Total cholesterol	167	155	-7%
HDL-cholesterol	54	58	+7%
LDL-cholesterol	99	84	-15%
Triglycerides	69	62	-10%
Fructoseamine	195	202	+4%

As seen in Table 2, the results show that during the 30-day time period studied, the total cholesterol level of the subject decreased, with a corresponding decrease in 5 the level of LDL-cholesterol and an increase in the level of HDL-cholesterol. The level of triglycerides decreased, while the level of fructoseamine increased very slightly. It is therefore concluded that the subject restricted her fat intake, (as shown by the decrease in LDL-cholesterol), without reducing her carbohydrate intake, (as shown by the slight increase in fructosamine level), indicating that betahistine has a 10 specific effect on reducing fat intake in a human subject.

It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention, 15 which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable subcombination.

Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations 20 will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims.

All publications, patents and patent applications mentioned in this specification are herein incorporated in their entirety by reference into the specification, to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated herein by reference. In addition, citation 5 or identification of any reference in this application shall not be construed as an admission that such reference is available as prior art to the present invention.

**Bibliography**

Arrang, J.-M., et al.. *Eur. J. Pharmacol.* 111: 73-84 (1985).

Attoub, S. *Life Sci.* 69: 469-78 (2001).

Bjenning, C. *Int. Sendai Histamine Symp.* P39-40 (2000).

5 Blandina, P. *Br. J. Pharmacol.* 119 : 1656-64 (1996).

Clapham, J. *Eur. J. Pharmacol.* 259 : 107-14 (1994).

Clayman, C.B. *J. Amer. Med. Assoc.* 238:1289-1290 (1977).

Fossati, A. et al. *Pharmacol. Res.* 43:389-392 (2001).

Fukagawa, K. *Am. J. Physiol.* 256: R605-R611 (1989).

10 Goldstein, D.J. *Obes. Res.* 2: 92-98 (1993).

Imamura, M. *Circ. Res.* 78 : 863-9 (1996).

Krupnick J.G. et al., *Ann. Rev. Pharmacol. Toxicol.* 38: 289-319 (1998).

Lecklin, A. *Brain Res.* 793 : 279-88 (1998).

Lecklin et al. *Inflamm. Res.* 51: suppl. 1, S53-S54 (2002)

15 Leurs, R. *Br. J. Pharmacol.* 116 : 2315-21 (1995).

Leurs, R. *Trends Pharmacol. Sci.* 19 : 177-84 (1998).

Lin, J.S. et al. *Brain Res.* 523 : 325-30 (1990).

Machidori, H. *Brain Res.* 590: 180-6 (1992).

McMahon, F.G. *Arch. Intern. Med.* 160: 2185-91 (2000).

20 Meyer, J.S., et al. *J. Clin. Pharmacol.* 14: 280-9 (1974).

Mokdad, A.H., *J. Amer. Med. Assoc.* 291: 1238-45 (2004).

Morimoto, T. *Behav. Brain Res.* 124: 145-150 (2001).

National Center for Health Statistics, Health E-Stats, Hyattsville, Md., (2000).

National Task Force on the Prevention and Treatment of Obesity, *J. Amer. Med. Assoc.* 276: 1907-15 (1996).

25 National Task Force on the Prevention and Treatment of Obesity, *Arch. Intern. Med.* 160: 898-904 (2000).

Onderwater, R.C. *Toxicology* 125: 117-29 (1998).

Physicians' Desk Reference, 55<sup>th</sup> edition, Montvale, N.J. (2001).

30 Pi-Sunyer et al. *American Heart Association (AHA) Scientific Sessions*, New Orleans, Louisiana (2004).

Poyurovsky, M. et al. *Int. Clin. Psychopharmacol.* 20 : 101-103 (2005).

Rasmussen, M.H. *Brit. Med. J.* 306: 1093-6 (1993).

Rossi, R., et al. *Physiol. and Behav.* 66: 517-521 (1999). Sakata, T. *Obes. Res. (Suppl.* 4) S541-S548 (1995).

Sakata, T. *Nutrition* 13: 403-11 (1997).

5 Saljoughian et al., *U.S Pharmacist*, 29: 2 (2004).

Seifert et al., *J. Pharmacol. Exp. Therap.* 305(3) : 1104, 2003.

Serdula, M.K. *J. Amer. Med. Assoc.* 282: 1353-8 (1999).

Sternoson, L.A. *Drug Metab. Dispos.* 2; 123-8 (1974).

Szelag, A., Malgorzata, T., and Merwid-Lad, A. *Polish J. Pharmacol.* 53: 701-707

10 (2001).

Szelag, A. *Adv. Clin. Exp. Med.* 11(3): 293-300 (2002).

Tuomisto, L. *Meth. Find. Exp. Clin. Pharmacol.* 16: 355-9 (1994).

Wadden, T.A. *Obes. Res.* 3: 549-57 (1995).

Yoshimatsu, H. *Diabetes* 48: 2286-91 (1999).

## WHAT IS CLAIMED IS:

1. A method of treating of a condition in which regulating a food intake in a human subject is beneficial, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist, said H<sub>1</sub> agonist having a pharmacological half-life of at least 3 hours.
2. A method of improving a compliance of a human subject to caloric restriction, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist, said H<sub>1</sub> agonist having a half-life of at least 3 hours.
3. A method of reducing a desire of a human subject to consume fats, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist, said H<sub>1</sub> agonist having a half-life of at least 3 hours.
4. A method of treating a condition associated with a metabolic derangement in a human subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist, said H<sub>1</sub> agonist having a pharmacological half-life of at least 3 hours.
5. A method of reducing total cholesterol level in a human subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist, said H<sub>1</sub> agonist having a pharmacological half-life of at least 3 hours.
6. A method of reducing low-density lipoprotein cholesterol and increasing high-density lipoprotein cholesterol levels in a human subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist, said H<sub>1</sub> agonist having a pharmacological half-life of at least 3 hours.

7. A method of reducing triglyceride level in a human subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of an H<sub>1</sub> agonist, said H<sub>1</sub> agonist having a pharmacological half-life of at least 3 hours.

8. The method of any of claims 1-3, wherein said pharmacological half-life ranges from about 3 hours to about 5 hours.

9. The method of any of claims 1-8, wherein said H<sub>1</sub> agonist is further an H<sub>3</sub> antagonist.


10. The method of any of claims 1-9, wherein said H<sub>1</sub> agonist is characterized by blood brain barrier permeability.

11. The method of any of claims 1-10, wherein said H<sub>1</sub> agonist is selected from the group consisting of betahistine, a betahistine metabolite, a betahistine pharmaceutically acceptable salt, a betahistine prodrug, a betahistine derivative and any combination thereof.

12. The method of claim 11, wherein said betahistine metabolite is selected from the group consisting of 2-(2-aminoethyl)-pyridine and 2-(2-hydroxyethyl)-pyridine.

13. The method of claim 11, wherein said betahistine salt is selected from the group consisting of betahistine dihydrochloride, betahistine mesilate, and betahistine trimebutine maleate.

14. The method of claim 11, wherein said betahistine derivative is selected from the group of compounds represented by the general formula I:



Formula I

wherein each of R<sub>1</sub>-R<sub>12</sub> is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl and aryl.

15. The method of any of claims 1-14, wherein said administering is effected by a route selected from the group consisting of the oral, transdermal, intravenous, subcutaneous, intramuscular, intranasal, intraauricular, sublingual, rectal, transmucosal, intestinal, buccal, intramedullar, intrathecal, direct intraventricular, intraperitoneal, and intraocular routes.

16. The method of claim 15, wherein said administering is effected orally.

17. The method of claim 15, wherein said administering is effected transdermally.

18. The method of any of claims 1-14, wherein said administering is effected by a route selected from the group consisting of the oral, transdermal, buccal, transmucosal, rectal and sublingual routes.

19. The method of any of claims 1-18, wherein said therapeutically effective amount ranges from about 2 mg per unit dosage to about 96 mg per unit dosage.

20. The method of claim 19, wherein said therapeutically effective amount ranges from about 10 mg per day to about 50 mg per day.

21. The method of any of claims 1-20, wherein said administering is effected from about 1 to about 4 times per day.

22. The method of claim 21, wherein said administering is effected twice per day.

23. The method of claim any of claims 1-18, wherein said administering is effected according to the development of hunger of the subject.

24. The method of any of claims 1-18, wherein said administering is performed such that a decrease of the body weight of the subject that ranges from about 1 to about 5 percent is effected, without restricting the food intake of the subject.

25. The method of claim 24, wherein said administering is performed such that no down-regulation of H<sub>1</sub> receptors is effected.

26. The method of claim 1, wherein said condition is selected from the group consisting of overeating, overweight, obesity, and a disorder caused or exacerbated thereby.

27. The method of claim 1, wherein said condition is associated with a psychological factor.

28. The method of claim 27, wherein said condition comprises binge eating disorder, night eating syndrome, obsessive eating, compulsive eating and bulimia.

29. The method of claim 26, wherein said condition is associated with a drug treatment.

30. The method of claim 29, wherein said drug is selected from the group consisting of a steroid hormone and a psychoactive drug.

31. The method of claim 26, wherein said disorder caused by or exacerbated by said condition is selected from the group consisting of a muscoskeletal disorder, a cardiovascular disorder, a dermatological disorder, a sleep disorder, a metabolic condition, diabetes and a diabetes-related condition.

32. The method of claim 4, wherein said condition is associated with high fat consumption.

33. The method of claim 4, wherein said condition is dyslipidemia.

34. The method of claim 33, wherein said condition is hypercholesterolemia.

35. The method of any of claims 1-31, wherein said H<sub>1</sub> agonist forms a part of a pharmaceutical composition, said pharmaceutical composition further comprising a pharmaceutically acceptable carrier.

36. The method of claim 35, wherein said pharmaceutical composition is a slow-release composition.

37. The method of any of claims 1-35, further comprising administering to the subject a therapeutically effective amount of an additional active agent.

38. The method of claim 37, wherein said additional active agent is selected from the group consisting of a non-steroidal anti-inflammatory drug, a muscle relaxant, an antigout agent, an immunosuppressant, a drug affecting bone mineralization, an alpha-adrenergic blocking drugs, an angiotensin-converting enzyme inhibitor, an antiarrhythmic drug, an anticoagulant, an antiplatelet, a thrombolytic, a beta-adrenergic blocking drug, a calcium channel blocker, a centrally acting drug, a

digitalis drug, a nitrate, a peripheral adrenergic antagonist, a vasodilator, an acne medication, an antipruritic agent, a corticosteroid, an anti-psoriasis agent, an anti-eczema agent, a hypnotic, an anti-depressant, an anti-histamine, a sulfonylurea, a meglitinide, a biguanide, a thiazolidinedione, an alpha-glucosidase inhibitor, a PPAR-gamma antagonist, insulin, a fibrate, an HMG-CoA reductase inhibitor, a bile acid sequestrant, a cholesterol absorption inhibitor, nicotinic acid, a derivative, analog and metabolite thereof, and any mixture thereof.

39. The method of claim 37, wherein said additional active agent is a weight control agent.

40. The method of claim 39, wherein said weight control agent is an appetite suppressant.

41. The method of claim 40, wherein said appetite suppressant is selected from the group consisting of noradrenergic agents, serotonergic agents, dopamnergic agents, endocannabinoid receptor blockers, or combinations thereof.

42. The method of any of claims 1-39, wherein said administering is affected according to the development of a hunger of the subject.

43. The method of any of claims 1-42, being used for inducing weight loss.

44. The method of any of claims 1-42, being used for maintaining weight loss or preventing a weight gain after or during a weight reducing diet.

45. The method of any of claims 1-42, being used for preventing weight gain in a subject having a condition associated with weight gain.

46. The method of claim 34, further comprising administering to the subject a therapeutically effective amount of an additional active agent selected from the group consisting of a fibrate, an HMG-CoA reductase inhibitor, a bile acid

54

sequestrant, a cholesterol absorption inhibitor, nicotinic acid, a derivative, analog and metabolite thereof, and any mixture thereof.

1/5




Fig. 1a (prior art)




Fig. 1b (prior art)

2/5




Fig. 2a (prior art)




Fig. 2b (prior art)




Fig. 3 (prior art)

4/5




Fig. 4




Fig. 5

5/5




Fig. 6