



(86) Date de dépôt PCT/PCT Filing Date: 2001/09/28

(87) Date publication PCT/PCT Publication Date: 2003/04/11

(85) Entrée phase nationale/National Entry: 2003/04/11

(86) N° demande PCT/PCT Application No.: EP 2001/011243

(87) N° publication PCT/PCT Publication No.: 2002/032898

(30) Priorité/Priority: 2000/10/14 (100 50 995.9) DE

(51) Cl.Int.⁷/Int.Cl.⁷ C07D 451/10, A61K 31/439, A61P 11/00

(71) Demandeur/Applicant:

BOEHRINGER INGELHEIM PHARMA GMBH & CO. KG,
DE


(72) Inventeurs/Inventors:

MEISSNER, HELMUT, DE;
MORSCHHAUSER, GERD, DE;
PIEPER, MICHAEL PAUL, DE;
POHL, GERALD, DE;
REICHL, RICHARD, DE;
SPECK, GEORG, DE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : NOUVEAUX ANTICHOLINERGIQUES, LEURS PROCEDES DE PRODUCTION ET LEUR UTILISATION EN
TANT QUE MEDICAMENTS

(54) Title: NOVEL ANTICHOLINERGICS, METHOD FOR THE PRODUCTION THEREOF AND USE THEREOF AS
MEDICAMENTS




(57) Abrégé/Abstract:

The invention relates to novel anticholinergics of general formula (1) wherein A, X⁻ and the radical R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ can have the meaning as cited in the claims and the description. The invention also relates to the production thereof in addition to the use thereof as medicaments.



Abstract

5 The present invention relates to new anticholinergics of general formula 1




wherein A, X⁻ and the groups R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ may have the
10 meanings given in the claims and in the specification, processes for preparing them
and their use as pharmaceutical compositions.

74402fft.204

**New Anticholinergics, Processes for Preparing them
and their use as Pharmaceutical Compositions**

5 The present invention relates to new anticholinergics of general formula 1



wherein A, X⁻ and the groups R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ may have the
10 meanings given in the claims and in the specification, processes for preparing them
and their use as pharmaceutical compositions.

Background to the Invention

Anticholinergics may be used therapeutically in a variety of complaints. Particular
15 mention should be made here, for example, of the treatment of asthma or COPD
(chronic obstructive pulmonary disease). For treating these diseases WO 92/16528
proposes anticholinergics which have a scopolamine, tropenol or tropine basic structure.

The objective on which WO 92/16528 is based is to prepare compounds with an
20 anticholinergic activity which are characterised by a long-lasting activity. To solve
this problem WO 92/16528 discloses, *inter alia*, benzoic acid esters of scopolamine,
tropenol or tropine.

For treating chronic diseases it is often desirable to prepare pharmaceutical
25 compositions having a fairly long duration of activity. As a rule, this ensures that the
concentration of the active substance needed in the body to achieve the therapeutic
effect is provided over a fairly long period of time without having to administer the
drug too frequently. Moreover, administering an active substance at fairly long time
intervals makes a major contribution to the patient's well being. It is particularly
30 desirable to provide a drug which can be used in a therapeutically beneficial manner
by a single application per day (single dose). The use of a drug once a day has the

advantage that the patient can become accustomed relatively quickly to regularly taking the drug at certain times of the day.

In order to be used as a medicament for use once a day, the active substance to be
5 given must meet particular requirements. First of all, the onset of the desired activity should take place relatively quickly after administration of the drug and ideally should have as constant an effect as possible over a subsequent fairly long period of time. On the other hand, the duration of activity of the drug should not substantially exceed a period of about one day. Ideally, an active substance has an activity profile such
10 that the preparation of a drug for administration once a day, which contains the active substance in therapeutically beneficial doses, can be deliberately controlled.

It has been found that the benzilic acid esters of scopolamine, tropenol and tropine disclosed in WO 92/16528 do not meet these stringent requirements. Because of
15 their extremely long period of activity, which significantly exceeds the above-mentioned period of about one day, they cannot be used therapeutically for administration in a single dose per day.

The aim of the present invention is therefore to provide new anticholinergics which,
20 by virtue of their activity profile, make it possible to prepare a drug for administration once a day. A further objective of the invention is to prepare compounds characterised by a relative rapid onset of activity. The invention further sets out to provide compounds which, after a rapid onset of activity, have as constant an activity as possible over a subsequent lengthy period of time. A further aim of the invention
25 is to provide compounds whose duration of activity does not substantially exceed a period of about one day in therapeutically beneficial doses. Finally, the invention sets out to provide compounds which have an activity profile which ensures good control of the therapeutic effect (i.e. total therapeutic effect without side effects caused by a build-up of the substance in the body).


30

Detailed Description of the Invention

Surprisingly, it has been found that the above objectives are achieved by means of compounds of general formula 1 wherein at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ does not represent hydrogen.


35

Accordingly, the present invention relates to compounds of general formula 1



wherein

A denotes a double-bonded group selected from among




X⁻ denotes an anion with a single negative charge,

R¹ and R² denote C₁-C₄-alkyl which may optionally be substituted by hydroxy or halogen;

R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, C₁-C₄-alkyl, C₁-C₄-alkyloxy, hydroxy, CF₃, CN, NO₂ or halogen, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

Preferred compounds of general formula 1 are those wherein

A denotes a double bonded group selected from among




X⁻ denotes an anion with a single negative charge selected from among chloride, bromide, methylsulphate, 4-toluenesulphonate and methanesulphonate, preferably bromide,

R¹ and R² which may be identical or different, denote a group selected from among methyl, ethyl, n-propyl and iso-propyl, which may optionally be substituted by hydroxy or fluorine, preferably unsubstituted methyl;

R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, methyl, ethyl, methyloxy, ethyloxy, hydroxy, fluorine, chlorine, bromine, CN, CF₃ or NO₂, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

Particularly preferred are compounds of general formula 1, wherein

A denotes a double-bonded group selected from among



X - denotes bromide;


5 R¹ and R², which may be identical or different, denote methyl or ethyl, preferably methyl;

R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, methyl, methyloxy, fluorine, chlorine or bromine, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

10

Of particular importance according to the invention are compounds of general formula 1, wherein

A denotes a double-bonded group selected from among



15 X - denotes bromide;


R¹ and R² which may be identical or different denote methyl or ethyl, preferably methyl;

R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, fluorine, chlorine or bromine, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

20

Also of particular importance according to the invention are compounds of general formula 1, wherein

A denotes a double-bonded group selected from among



25

X - denotes bromide;

R¹ and R² which may be identical or different denote methyl or ethyl, preferably methyl;


R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, fluorine, chlorine or bromine, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

30

Of outstanding importance according to the invention are the compounds of general formula 1, wherein

5

A denotes a double-bonded group selected from among



X - denotes bromide;


R¹ and R² denote methyl;

5 R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen or fluorine, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

The invention relates to the compounds of formula 1, optionally in the form of the
10 individual optical isomers, mixtures of the individual enantiomers or racemates thereof.

In the compounds of general formula 1 the group R³, R⁴, R⁵, R⁶, R⁷ and R⁸, provided that they do not denote hydrogen, may each be in the ortho, meta or para
15 position relative to the bond to the "-C-OH"-group. Provided that none of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ denotes hydrogen, R³ and R⁵ are preferably linked in the para-position and R⁴, R⁶, R⁷ and R⁸ are preferably linked in the ortho- or meta-position, most preferably in the meta-position. Particularly preferred are compounds of formula 1 wherein the groups R⁷ and R⁸ denote hydrogen. If in this case one of
20 the groups R³ and R⁴ and one of the groups R⁵ and R⁶ denotes hydrogen, the other group in each case is preferably linked in the meta- or para-position, most preferably in the para-position. If none of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ denotes hydrogen, according to the invention the compounds of general formula 1 wherein the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ have the same meaning are
25 particularly preferred.

Of particular importance according to the invention are the compounds of general formula 1 wherein the ester-substituent on the nitrogen-bicyclic group is in the α -configuration. These compounds correspond to general formula 1- α



The following compounds are of particular significance according to the invention:

- tropenol 3,3',4,4'-tetrafluorobenzilate-methobromide;
- 5 - scopine 3,3',4,4'-tetrafluorobenzilate-methobromide;
- scopine 4,4'-dichlorobenzilate-methobromide;
- scopine 4,4'-difluorobenzilate-methobromide;
- tropenol 3,3'-difluorobenzilate-methobromide;
- scopine 3,3'-difluorobenzilate-methobromide;
- 10 - tropenol 4,4'-difluorobenzilate-ethyl bromide.

Unless otherwise stated, the alkyl groups are straight-chained or branched alkyl groups having 1 to 4 carbon atoms. The following are mentioned by way of example: methyl, ethyl, propyl or butyl. In some cases the abbreviations Me, Et, Prop or Bu are used to denote the groups methyl, ethyl, propyl or butyl. Unless otherwise stated, the definitions propyl and butyl include all the possible isomeric forms of the groups in question. Thus, for example, propyl includes n-propyl and iso-propyl, butyl includes iso-butyl, sec.butyl and tert.-butyl, etc.

Unless otherwise stated, the term alkyloxy groups denotes branched and unbranched alkyl groups having 1 to 4 carbon atoms which are linked via an oxygen atom. Examples of these include: methyloxy, ethyloxy, propyloxy or butyloxy. The abbreviations MeO-, EtO-, PropO- or BuO- are used in some cases to denote the groups methyloxy, ethyloxy, propyloxy or butyloxy. Unless otherwise stated, the definitions propyloxy and butyloxy include all possible isomeric forms of the groups in question. Thus, for example, propyloxy includes n-propyloxy and iso-propyloxy, butyloxy includes iso-butyloxy, sec.butyloxy and tert.-butyloxy, etc. In some cases, within the scope of the present invention, the term alkoxy is used instead of the term alkyloxy. Accordingly, the terms methoxy, ethoxy, propoxy or butoxy may also be used to denote the groups methyloxy, ethyloxy, propyloxy or butyloxy

Halogen within the scope of the present invention denotes fluorine, chlorine, bromine or iodine. Unless stated otherwise, bromine is the preferred halogen.

- 5 The compounds according to the invention may partly be prepared, as illustrated below, analogously to procedures which are already known from the prior art (Diagram 1).




Diagram 1:


Starting from the compounds of formula 2 the esters of general formula 4 may be obtained by reacting with the benzilic acid esters of formula 3, wherein R denotes a C₁-C₄-alkyl group, for example. This reaction may be carried out, for example, in a sodium melt at elevated temperature, preferably at about 50-150°C, particularly preferably at about 90-100°C at low pressure, preferably below 500mbar, most preferably below 75mbar. Alternatively, instead of the benzilic acid esters 3, the corresponding α -chlorine compounds may be used (Cl instead of OH). In this case the reaction may be carried out analogously to the examples of synthesis disclosed in WO 92/16528, to which reference is hereby made in its entirety. The compounds of formula 4 thus obtained can be converted into the target compounds of formula 1 by reacting with the compounds R²-X wherein R² and X may be as hereinbefore defined. This synthesis step may also be carried out analogously to the examples of synthesis disclosed in WO 92/16528.

Alternatively to the method of synthesising the compounds of formula 4 illustrated in Diagram 1, the derivatives 4 wherein the nitrogen bicyclic group denotes a scopine-derivative may be obtained by oxidising compounds of formula 4 wherein the bicyclic

nitrogen group is a tropanyl group. This may be done as follows, according to the invention.


The compound 4 wherein A denotes -CH=CH-, is suspended in a polar organic solvent, preferably in a solvent selected from among N-methyl-2-pyrrolidone (NMP), dimethylacetamide and dimethylformamide, preferably dimethylformamide, and then heated to a temperature of about 30-90°C, preferably 40-70°C. Then a suitable oxidising agent is added and the mixture is stirred at constant temperature for 2 to 8 hours, preferably 3 to 6 hours. The preferred oxidising agent is vanadium pentoxide mixed with H₂O₂, most preferably H₂O₂-urea complex combined with vanadium pentoxide. The mixture is worked up in the usual way. The products may be purified by crystallisation or chromatography, depending on their crystallisation tendencies.

As shown in Diagram 1, the intermediate products of general formula 4 are of crucial importance. Accordingly, in another aspect, the present invention relates to the intermediates of formula 4



wherein

A denotes a double-bonded group selected from among




R¹ denotes C₁-C₄-alkyl which may optionally be substituted by hydroxy or halogen;

R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, C₁-C₄-alkyl, C₁-C₄-alkyloxy, hydroxy, CF₃, CN, NO₂ or halogen, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

The intermediates of general formula 4 are preferred wherein

9

A denotes a double-bonded group selected from among




R¹ methyl, ethyl, n-propyl and iso-propyl, which may optionally be substituted by hydroxy or fluorine, preferably unsubstituted methyl;

5 R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, methyl, ethyl, methyloxy, ethyloxy, hydroxy, fluorine, chlorine, bromine, CN, CF₃ or NO₂, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

10 Particularly preferred are the intermediates of general formula 4, wherein

A denotes a double-bonded group selected from among




R¹ denotes methyl or ethyl, preferably methyl;

15 R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, methyl, methyloxy, fluorine, chlorine or bromine, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

Of particular importance according to the invention are the intermediates of general formula 4, wherein

20 A denotes a double-bonded group selected from among




R¹ denotes methyl or ethyl, preferably methyl;

25 R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, fluorine, chlorine or bromine, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

Also of particular importance according to the invention are the intermediates of formula 4, wherein

A denotes a double-bonded group selected from among




30

R¹ denotes methyl or ethyl, preferably methyl;

R^3, R^4, R^5, R^6, R^7 and R^8 , which may be identical or different, denote hydrogen, fluorine, chlorine or bromine, with the proviso that at least one of the groups R^3, R^4, R^5, R^6, R^7 and R^8 is not hydrogen.

- 5 Of exceptional importance according to the invention are the intermediates of general formula 4, wherein

A denotes a double-bonded group selected from among




R^1 denotes methyl;

- 10 R^3, R^4, R^5, R^6, R^7 and R^8 , which may be identical or different, denote hydrogen or fluorine, with the proviso that at least one of the groups R^3, R^4, R^5, R^6, R^7 and R^8 is not hydrogen.

As in the compounds of general formula 1, in the intermediates of formula 4 the
 15 groups R^3, R^4, R^5, R^6, R^7 and R^8 , if they do not represent hydrogen, may be in the ortho, meta or para position relative to the bond to the "-C-OH" group. If none of the groups R^3, R^4, R^5, R^6, R^7 and R^8 denotes hydrogen, R^3 and R^5 are preferably linked in the para-position and R^4, R^6, R^7 and R^8 are preferably linked in the ortho- or meta-position, most preferably in the meta-position. Particularly preferred are
 20 intermediates of formula 4 wherein R^7 and R^8 denote hydrogen. If in this case one of the groups R^3 and R^4 and one of the groups R^5 and R^6 denote hydrogen, the other group in each case is preferably linked in the meta- or para-position, most preferably in the para-position. If none of the groups R^3, R^4, R^5, R^6, R^7 and R^8 denotes hydrogen, according to the invention the intermediates of general formula 4
 25 wherein the groups R^3, R^4, R^5, R^6, R^7 and R^8 have the same meaning are particularly preferred.

In order to prepare the intermediate products of general formula 4, correspondingly substituted benzilic acid derivatives of general formula 3 are used. These are
 30 obtained, analogously to methods known in the art, by reacting Grignard reagents generated *in situ* from the bromides 5 with the aromatic α -carbonyl-carboxylic acid esters 6.

11


Diagram 2:

The reaction is carried out in anhydrous organic solvents, preferably in ethereal
 5 solvents, most preferably in a solvent selected from among diethylether, dioxane and tetrahydrofuran (THF), the latter being particularly significant. The Grignard reagent is generated from the bromides 5 by reacting with Mg-chips. In some cases it may be necessary to add a reaction starter such as iodine or dibromomethane. In order to complete the formation of Grignard-reagent it may be necessary to heat the reaction
 10 mixture over a period of 0.5 to 2 hours, preferably to above 30°C, most preferably to above 50°C. The upper limit of the temperature range which may be used will naturally be determined from the boiling temperature of the solvent used. The Grignard-reagent thus obtained is then slowly added dropwise to a solution of 6 in one of the above-mentioned solvents. The work may be done at ambient
 15 temperature, but is preferably done at a temperature in the range from 0-15°C. The reaction is generally complete after 1 to 4 hours. The mixture is worked up by conventional methods. The products are purified by crystallisation or column chromatography, depending on the crystallisation tendencies of the compounds 3. Alternatively, the compounds of formula 3 may also be obtained analogously to other
 20 methods of synthesis known from the prior art. Where, for example, suitably substituted benzilic acids are already known in the art and are commercially available, the compounds of formula 3 may also be obtained directly from them by acid- or base-catalysed esterification with the corresponding alcohols R-OH. Where suitably substituted benzils are already known from the art and are commercially
 25 available, the compounds of formula 3 may also be obtained directly from them by benzilic acid rearrangement and subsequent acid- or base-catalysed esterification with the corresponding alcohols R-OH.

As can be seen from Diagram 1, the benzilic acid derivatives of general formula 3
 30 have a central importance in preparing the compounds of general formula 4 and thus those of formula 1 as well.

Accordingly, in another aspect, the present invention relates to benzilic acid derivatives of formula 3

12



wherein

- 5 R denotes C₁-C₄-alkyl, preferably methyl or ethyl;
 R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, C₁-C₄-alkyl, C₁-C₄-alkyloxy, hydroxy, CN, CF₃, NO₂ or halogen, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

10

Preferred are benzoic acid derivatives of general formula 3, wherein

- R denotes methyl or ethyl;
 R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, methyl, ethyl, methoxy, ethoxy, hydroxy, fluorine, chlorine, bromine,
 15 CN, CF₃ or NO₂, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

Particularly preferred are benzoic acid derivatives of general formula 3, wherein

- R denotes methyl or ethyl;
 20 R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, methyl, methoxy, fluorine, chlorine or bromine, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

- Of particular importance according to the invention are benzoic acid derivatives of
 25 general formula 3, wherein

- R denotes methyl or ethyl;
 R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, fluorine, chlorine or bromine, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen.

30

Of exceptional importance according to the invention are the intermediates of general formula 3, wherein

- R denotes methyl or ethyl;

R^3 , R^4 , R^5 , R^6 , R^7 and R^8 , which may be identical or different, denote hydrogen or fluorine, with the proviso that at least one of the R^3 , R^4 , R^5 , R^6 , R^7 and R^8 is not hydrogen.

- 5 As in the compounds of general formula 4 and 1, in the benzilic acid derivatives of formula 3 the groups R^3 , R^4 , R^5 , R^6 , R^7 and R^8 , provided that they are not hydrogen, may each be in the ortho, meta or para position relative to the bond to the "-C-OH" group. If none of the groups R^3 , R^4 , R^5 , R^6 , R^7 and R^8 denotes hydrogen, R^3 and R^5 are preferably linked in the para-position and R^4 , R^6 , R^7 and R^8 are
- 10 preferably linked in the ortho- or meta-position, most preferably in the meta-position. Particularly preferred are benzilic acids of formula 3 wherein R^7 and R^8 denote hydrogen. If one of the groups R^3 and R^4 and one of the groups R^5 and R^6 denotes hydrogen, the other group in each case is preferably linked in the meta- or para-position, most preferably in the para-position. If none of the groups R^3 , R^4 , R^5 , R^6 ,
- 15 R^7 and R^8 denotes hydrogen, according to the invention the benzilic acid derivatives of general formula 3 wherein the groups R^3 , R^4 , R^5 , R^6 , R^7 and R^8 have the same meaning are particularly preferred.

The examples of synthesis described hereinafter serve to illustrate the present invention still further. However, they are intended only as examples of procedures as an illustration of the invention without restricting the invention to the subject-matter described by way of example.

I. Synthesis of the benzilic acid derivatives of formula 3:

I.1.: ethyl 3,3',4,4'-tetrafluorobenzilate 3a:.

25 The Grignard reagent is prepared from 2.24 g (0.092 mol) of magnesium chips, a few granules of iodine and 17.80 g (0.092 mol) of 1-bromo-3,4-difluoro-benzene in 100 ml of THF at 50° C. After the halide has all been added, the mixture is stirred for another hour. The Grignard reagent thus obtained is added dropwise to 18.81 g

30 (0.088 mol) of ethyl 3,4-difluorophenylglyoxylate in 80 ml of THF at 10°-15° C and the mixture obtained is stirred for 2 hours at 5°C.

The white suspension is poured onto ice/sulphuric acid for working up, extracted with ethyl acetate, the organic phase is washed with water, dried over $MgSO_4$ and evaporated to dryness. The crude product is purified by column chromatography

35 (eluant: toluene).

Yield: 10.80 g of oil 1 (= 38 % of theory)

I.2.: methyl 2,2'-dichlorobenzilate 3b:.

A solution of 10.0 g (0.034 mol) of 2,2'-dichlorobenzilic acid in 50 ml of ethanol was added dropwise to freshly prepared sodium ethoxide solution of 0.78 g (0.034 mol)

of sodium and 100 ml of ethanol at 20° C and stirred overnight. The solution was evaporated to dryness, the residue was dissolved in DMF and at 20°C 9.57 g (0.0674 mol) of methyl iodide were added dropwise and the mixture was stirred for a further 72 hours. 300 ml of water were added dropwise to the resulting suspension while cooling with ice, it was extracted with diethyl ether, the organic phase was washed with water, dried over Na₂SO₄ and evaporated to dryness. Yield: 10.48 g 8 (= 100% of theory)

I.3.: methyl 4,4'-difluorobenzilate 3c:.

10 I.3.1.: 4,4'-difluorobenzilic acid:

At about 100° C a solution of 24.62 g (0.1 mol) of 4,4'-difluorobenzil in 250 ml of dioxane is added dropwise to a solution of 49.99 g (1.25 mol) of NaOH flakes in 300 ml of water and stirred for 2 h. The dioxane is largely distilled off and the aqueous solution remaining is extracted with dichloromethane. When the aqueous solution is acidified with sulphuric acid, a precipitate is formed which is suction filtered, washed and dried. The filtrate is extracted with dichloromethane, the organic phase is dried over Na₂SO₄ and evaporated to dryness. Yield: 25,01 g (= 95 % of theory); melting point:133°-136° C

20 I.3.2.: methyl 4,4'-difluorobenzilate:

25.0 g (0.095 mol) of 4,4'-difluorobenzilic acid are added to freshly prepared sodium ethoxide solution containing 2.17 g (0.095 mol) of sodium and 200 ml of ethanol at 20° C and stirred for 3 h. The solution is evaporated to dryness, the residue is dissolved in DMF, 22.57 g (0.16 mol) of methyl iodide are added dropwise at 20° C and the mixture is stirred for 24 h. It is worked up and purified analogously to compound **3b**. Yield: 21.06 g of 11 (= 80 % of theory)

I.4.: methyl 2,2',4,4'-tetrafluorobenzilate 3d:.

I.4.1: methyl 2,4-difluorophenylglyoxylate:

30 50 g (0.44 mol) of 1,3-difluorobenzene are dissolved at 20° C in 135 ml of carbon disulphide with 73.5 g (0.55 mol) AlCl₃ and then 55.15 g (0.45 mol) of methyl oxalate chloride are added and the mixture is stirred for 2.5 h at 20° C. Ice-cold 2 N aqueous hydrochloric acid is added dropwise while cooling, extracted with ethyl acetate, the organic phase is washed with water and 10 % aqueous Na₂CO₃ solution, dried over 35 MgSO₄ and evaporated to dryness. Yield: 38.0 g of oil 14 (= 43 % of theory)

I.4.2.: methyl 2,2',4,4'-tetrafluorobenzilate:

15

Starting from the methyl 2,4-difluorophenylglyoxylate obtained according to the above procedure, the title compound is prepared analogously to 3a;

Yield: 7.55 g of 15 (= 13 % of theory).

5

I.5.: methyl 4,4'-dimethylbenzilate 3e:.

At 20° C, 12.43 g (0.048 mol) of 4,4'-dimethylbenzilic acid in 50 ml ethanol are added dropwise to freshly prepared sodium ethoxide solution containing 1.1 g (0.045 mol) of sodium and 100 ml of ethanol and the mixture is stirred for 30 min. The solution is
10 evaporated to dryness, the residue dissolved in 50 ml of DMF, 9.08 g (0.064 mol) of methyl iodide are added dropwise at 20° C and stirring is continued for a further 24 h. 300 ml of water are added dropwise to the resulting suspension while cooling with ice, the mixture is extracted with diethylether, the organic phase is washed with water, dried over Na₂SO₄ and evaporated to dryness.

15 Yield: 8.6 g (= 99 % of theory); melting point: 83-84°C.

I.6.: methyl 3,3',4,4'-tetramethoxybenzilate 3f:.

I.6.1: methyl 3,4-dimethoxyglyoxylate:

14.00 g (0.11 mol) of AlCl₃ are placed in 100 ml of dichloromethane and at 5°C
20 combined with 12.86 g (0.11 mol) of monomethyl oxalate chloride. 1,2-dimethoxybenzene is added dropwise to the organic solution at 0°C, the mixture is stirred for 1 h at 0°C, then for 24 h at 20° C, poured onto ice/hydrochloric acid, extracted, the organic phase is washed with water and NaHCO₃ solution, dried over MgSO₄ and evaporated to dryness. The residue is crystallised from
25 diethylether/petroleum ether.

Yield: 13.55 g (= 60 % of theory); melting point: 65°-66° C.

I.6.2: methyl 3,3',4,4'-tetramethoxybenzilate:

The Grignard reagent is prepared from 1.58 g (0.065 mol) of magnesium chips,
30 some iodine and 14.10 g (0.065 mol) of bromoveratrol in 50 ml of THF at 50°C. Stirring is continued for 1 h. 11.28 g (0.05 mol) of methyl 3,4-dimethoxyglyoxylate are placed in 80 ml of THF and the Grignard reagent is added dropwise at 10°-15° C, the mixture is stirred for 2 h at 20° C, poured onto ice/sulphuric acid and extracted with ethyl acetate. Then the organic phase is washed with water, dried over MgSO₄
35 and evaporated to dryness.

The product is purified by crystallisation from acetone/diethylether.

Yield: 7.62 g (= 42 % of theory); melting point: 129°-130° C.

I.7.: methyl 4,4'-dimethoxybenzilate 3g:.

I.7.1: methyl 4-methoxyglyoxylate:

The compound is prepared analogously to step I.6.1 starting from 21.65 g (0.20 mol) of anisol; Yield: 16.45 g (= 60 % of theory); melting point: 52° C.

5

I.7.2: methyl 3,4-dimethoxybenzilate:

The compound is prepared analogously to step I.6.2, starting from 16.45 g (0.085 mol) of methyl 4-methoxyglyoxylate; the product is purified by crystallisation from isopropanol; Yield: 12.28 g (= 48 % of theory); melting point: 111°C.

10

I.8.: methyl 3,3'-dimethyl-4,4'-dimethoxybenzilate 3h:.**I.8.1: methyl 3-methyl-4-methoxyglyoxylate:**

The compound is prepared analogously to step I.6.1 starting from 26.88 g (0.22 mol) of 2-methylanisol; Yield: 21.0 g (= 46 % of theory); melting point: 49° C.

15

I.8.2: methyl 3,4-dimethoxybenzilate:

The compound is prepared analogously to step I.6.2, starting from 21.0 g (0.1 mol) of methyl 3-methyl-4-methoxyglyoxylate; the product is purified by crystallisation from petroleum ether/diethylether; Yield: 11.1 g (= 33 % of theory); melting point: 134°C.

20

I.9.: ethyl 4,4'-dichlorobenzilate 3i:.

The product may be synthesised analogously to step I.5.

I.10: methyl 3,3',5,5'-tetrafluorobenzilate 3j:.

25

I.10.1: 3,3',5,5'-tetrafluorobenzil:

110 ml of ethanol are taken at ambient temperature and 50.0 g (0.352 mol) of 3,5-difluorobenzaldehyde and 444 g (0.018 mol) of 3-ethyl-5-(2-hydroxyethyl)-4-methylthiazolium bromide are added. Then 10.7 g (0.11 mol) of triethylamine are added dropwise. The mixture is refluxed for 3 h and evaporated to dryness. The residue is taken up in ethyl acetate and extracted with water, sodium pyrosulphite in water and Na₂CO₃ solution. After drying over MgSO₄ the mixture is evaporated to dryness.

30

Yield: 39.91 g of yellow crystals (= 80% of theory).

35

39.91 g of the acyloin thus obtained are dissolved in 1700 ml of acetonitrile at ambient temperature, 0.2 ml of vanadium-(V)-oxytrichloride are added and oxygen is piped in. After 1.5 h the solution is evaporated to dryness, extracted with ethyl

acetate and water, as well as Na_2CO_3 solution, dried over MgSO_4 and evaporated to dryness. The residue remaining is stirred with petroleum ether/ethyl acetate 95:5. Yield: 26.61 g yellowish-green crystals (= 67% of theory); melting point.: 136-138°C.

5 I.10.2: 3,3',5,5'-Tetrafluorobenzilic acid:

46.98 g (1.747 mol) of sodium hydroxide in 330 ml water are placed in a boiling water bath with thorough stirring and a solution of 26.61 g (0.094 mol) of 3,3',5,5'-tetrafluorobenzil in 330 ml dioxane is added dropwise and then stirred for another 1 h. After cooling the dioxane is evaporated, the residue is diluted with water and
10 extracted with diethylether. The organic phase is acidified, extracted with dichloromethane, dried over MgSO_4 , evaporated to dryness. Yield: 20.15 g of yellowish crystals (= 71% of theory).

I.10.3: methyl 3,3',5,5'-tetrafluorobenzilate:

15 20.15 g (0.0671 mol) of 3,3',5,5'-tetrafluorobenzilic acid are placed in 250 ml acetonitrile, 18.6 g (0.121 mol) of DBU and 34.4 g (0.2426 mol) of methyl iodide are added and then stirred for 6 h at ambient temperature. The reaction mixture is evaporated to dryness, the residue extracted with ethyl acetate and water, the organic phase dried over MgSO_4 , evaporated to dryness.
20 The product is recrystallised from cyclohexane.
Yield: 15.11 g of beige crystals (= 68% of theory); melting point.: 113-114° C

I.11: methyl 3,3'-dichlorobenzilate 3k:

I.11.1.: 3,3'-dichlorobenzil:

25 100 ml of ethanol are used at ambient temperature and 50.0 g (0.356 mol) of 3-chlorobenzaldehyde and 4.54 g (0.018 mol) of 3-ethyl-5-(2-hydroxyethyl)-4-methylthiazolium bromide are added. Then 10.7 g (0.11 mol) of triethylamine are added dropwise. The mixture is refluxed for 3 hours and evaporated to dryness. The residue is taken up in ethyl acetate and extracted with water, sodium pyrosulphite in
30 water and Na_2CO_3 solution. After drying over MgSO_4 it is evaporated to dryness. The product obtained is recrystallised from isopropanol and petroleum ether.
Yield: 13.2 g of white crystals (= 13% of theory); melting point.: 69-70°C.

13.0 g of the acyloin thus obtained are dissolved in 460 ml acetonitrile at RT, 0.0867
35 g of vanadium-(V)-oxytrichloride are added and oxygen is piped in. After 1.5 h the solution is evaporated to dryness, extracted with ethyl acetate and water, as well as Na_2CO_3 solution, dried over MgSO_4 and evaporated to dryness. The residue remaining is stirred with petroleum ether/ethyl acetate 95:5.
Yield: 12.59 g of yellow crystals (= 97% of theory); melting point.: 116-117°C.

I.11.2.: 3,3'-dichlorobenzilic acid:

51.45 g (1.286 mol) of sodium hydroxide in 1000 ml water are placed in a bath of boiling water with thorough stirring and a solution of 28.5 g (0.102 mol) of 3,3'-dichlorobenzil in 700 ml dioxane is added dropwise and then stirred for another 1 h. After cooling the dioxane is evaporated down, the residue is diluted with water and extracted with diethylether. The organic phase is acidified, extracted with dichloromethane, dried over MgSO_4 , evaporated to dryness.
Yield: 32.7 g (= 71% of theory).

10

I.11.3.: methyl 3,3'-dichlorobenzilate:

From 100 ml of ethanol and 1.97 g (0.0855 mol) of sodium, a sodium ethoxide solution is prepared to which 26.6 g (0.0855 mol) of 3,3'-dichlorobenzilic acid in 50 ml of ethanol are added dropwise. The mixture is then stirred for 4 h at ambient temperature. After the solvent has been distilled off the residue is dissolved in 150 ml DMF and 24.27 g (0.171 mol) of methyl iodide are added dropwise, then stirred for another 24 h. While cooling with ice, 300 ml of water and 200 ml of diethylether are added dropwise, the phases are separated, the aqueous phase is extracted with diethylether, then the organic phases are washed with Na_2CO_3 solution and shaken with water till neutral. After drying over Na_2SO_4 the mixture is evaporated to dryness.
Yield: 22.91 g of yellow oil (= 82% of theory).

I.12: methyl 4,4'-di(trifluoromethyl)benzilate 3I:.**I.12.1.: 4,4'-di(trifluoromethyl)benzil:**

25 The reaction to form the acyloin derivative is carried out analogously to the process according to step I.11.1 starting from 174.1 g of 4-(trifluoromethyl)-benzaldehyde.
Yield: 150.2 g of whitish-yellow crystals (= 86% of theory).

150.2 g of the acyloin thus obtained are reacted analogously to the method according to step I.11.1 to obtain the benzil.
Yield: 93.5 g of yellow crystals (= 63% of theory); melting point.: 141-142°C.

I.12.2.: 4,4'-di(trifluoromethyl)benzilic acid:

10.00 g (0.0289 mol) of 4,4'-di(trifluoromethyl)benzil are reacted analogously to the method according to step I.11.2 to obtain the corresponding benzilic acid.
Yield: 8.15 g of yellowish crystals (= 77% of theory).

I.12.3.: methyl 4,4'-di(trifluoromethyl)benzilate:

38.5 g (0.115 mol) of 4,4'-di(trifluoromethyl)benzilic acid, 30.5 g (0.20 mol) of DBU and 56.8 g (0.40 mol) of methyl iodide are reacted in 400 ml acetonitrile analogously to step I.10.3. The product is purified by flash chromatography (eluant cyclohexane/ethyl acetate 95:5). Yield: 20.05 g of white crystals (= 46% of theory)
5 melting point.: 68° C

I.13: methyl 3,3'-di(trifluoromethyl)benzilate 3m:.

I.13.1.: 3,3'-di(trifluoromethyl)benzil:

The reaction to form the acyloin derivative is carried out analogously to the process
10 according to step I.11.1 starting from 30.0 g of 3-(trifluoromethyl)-benzaldehyde.
Yield: 25.96 g of yellow oil (= 87% of theory).

25,96 g of the acyloin thus obtained are reacted analogously to the method according to step I.11.1 to obtain the benzil.
15 Yield: 10.5 g of light yellow crystals (= 24% of theory).

I.13.2.: 3,3'-di(trifluoromethyl)benzilic acid:

10.5 g (0.0182 mol) of 3,3'-di(trifluoromethyl)benzil are reacted analogously to the method according to step I.11.2 to obtain the corresponding benzilic acid.
20 Yield: 10.55 g of yellowish oil.

I.13.3.: methyl 3,3'-di(trifluoromethyl)benzilate:

10.55 g (0.0289 mol) of 4,4'-di(trifluoromethyl)benzilic acid, 8.82 g (0.0579 mol) of DBU and 16.44 g (0.1158 mol) of methyl iodide are reacted in 110 ml acetonitrile
25 analogously to step I.10.3. The product is purified by recrystallisation from cyclohexane.
Yield: 6.02 g of white crystals (= 57% of theory); melting point.: 69-70°C.

I.14: methyl 3,3'-dichloro-4,4'-difluorobenzilate 3n:.

30 **I.14.1.: 3,3'-dichloro-4,4'-difluorobenzil:**

The reaction to form the acyloin derivative is carried out analogously to the process according to step I.11.1 starting from 30.0 g of 3-chloro-4-fluoro-benzaldehyde.
Yield: 29.49 g of orange oil.

35 29.49 g of the acyloin thus obtained are reacted analogously to the method according to step I.11.1 to obtain the benzil.
Yield: 24.88 g of yellow crystals.

I.14.2.: 3,3'-dichloro-4,4'-difluorobenzilic acid:

24,88 g (0.079 mol) of 3,3'-dichloro-4,4'-difluorobenzil are reacted analogously to the method according to step I.11.2 to obtain the corresponding benzilic acid.

Yield: 17.07g of orange solid.

5 I.14.3.: methyl 3,3'-dichloro-4,4'-difluorobenzilate:

17.07 g (0.0512 mol) of 3,3'-dichloro-4,4'-difluorobenzilic acid, 14.10 g (0.0926 mol) of DBU and 26.29 g (0.1852 mol) of methyl iodide are reacted in 200 ml acetonitrile analogously to step I.10.3. Yield: 6.77 g (= 38% of theory).

10 **I.15: methyl 2,2',5,5'-tetrafluorobenzilate 3o:**

I.15.1.: 2,2',5,5'-tetrafluorobenzil:

The reaction to form the acyloin derivative is carried out analogously to the process according to step I.11.1 starting from 50.0 g 2,5-difluoro-benzaldehyde.

Yield: 45.5 g of yellow crystals.

15

45.5 g of the acyloin thus obtained are reacted analogously to the method according to step I.11.1 to obtain the benzil.

Yield: 39,75 g of yellow crystals (=88% of theory).

20 I.15.2.: 2,2',5,5'-tetrafluorobenzilic acid:

39.75 g (0.14 mol) of 2,2',5,5'-tetrafluorobenzil are reacted analogously to the method according to step I.11.2 to obtain the corresponding benzilic acid.

Yield: 44.76 g of yellowish oil.

25 I.15.3.: methyl 2,2',5,5'-tetrafluorobenzilate:

44.76 g (0.14 mol) of 2,2',5,5'-tetrafluorobenzilbenzilic acid are reacted analogously to step I.10.3. Yield: 29.4 g of oil.

I.16: methyl 2,2',3,3'-tetrafluorobenzilate 3p:

30 I.16.1.: 2,2',3,3'-tetrafluorobenzil:

The reaction to form the acyloin derivative is carried out analogously to the process according to step I.11.1 starting from 30.0 g 2,3-difluoro-benzaldehyde.

Yield: 29.85 g of yellow crystals.

35 29.85 g of the acyloin thus obtained are reacted analogously to the method according to step I.11.1 to obtain the benzil.

Yield: 25.22 g of orange oil.

I.16.2.: 2,2',3,3'-tetrafluorobenzilic acid:

25.22 g (0.0894 mol) of 2,2',3,3'-tetrafluorobenzil are reacted analogously to the method according to step I.11.2 to obtain the corresponding benzilic acid.

Yield: 29.13 g of orange solid.

5 I.16.3.: methyl 2,2',3,3'-tetrafluorobenzilate:

29.13 g (0.097 mol) of 2,2',3,3'-tetrafluorobenzilbenzilic acid are reacted analogously to step I.10.3. Yield: 15.78 g of beige crystals; melting point.: 102-103°C.

I.17: methyl 3,3'-difluorobenzilate 3q:

10 I.17.1.: 3,3'-difluorobenzil:

The reaction to form the acyloin derivative is carried out analogously to the process according to step I.11.1 starting from 50.0 g 3-fluoro-benzaldehyde.

Yield: 49.45 g.

15 49.45 g of the acyloin thus obtained are reacted analogously to the method according to step I.11.1 to obtain the benzil.

Yield: 42.01 g of light yellow crystals; melting point.: 104-105°C.

I.17.2.: 3,3'-difluorobenzilic acid:

20 42,01 g (0.171 mol) of 3,3'-difluorobenzil are reacted analogously to the method according to step I.11.2 to obtain the corresponding benzilic acid.

Yield: 33.07 g (=75% of theory); melting point.: 121-122°C.

I.17.3.: methyl 3,3'-difluorobenzilate:

25 33.7 g (0.128 mol) of 3,3'-difluorobenzilbenzilic acid are reacted analogously to step I.10.3. Yield: 34.78 g beige crystals (=98% of theory); melting point.: 84-85°C.

I.18: methyl 4,4'-dichloro-3,3'-difluorobenzilate 3r:

I.18.1.: 4,4'-dichloro-3,3'-difluorobenzil:

30 The reaction to form the acyloin derivative is carried out analogously to the process according to step I.11.1 starting from 33,13 g 4-chloro-3-fluoro-benzaldehyde.

Yield: 30.07 g of oil.

30.07 g of the acyloin thus obtained are reacted analogously to the method

35 according to step I.11.1 to obtain the benzil.

Yield: 19.32 g of light yellow powder.

I.18.2.: 4,4'-dichloro-3,3'-difluorobenzilic acid:

19.32 g (0.0613 mol) of 4,4'-dichloro-3,3'-difluorobenzil are reacted analogously to the method according to step I.11.2 to obtain the corresponding benzilic acid.

Yield: 25.3 g of oil.

5 I.18.3.: methyl 4,4'-dichloro-3,3'-difluorobenzilate:

25.3 g (0.075 mol) of 4,4'-dichloro-3,3'-difluorobenzilic acid are reacted analogously to step I.10.3. Yield: 13.07 g of yellow oil (= 50% of theory).

I.19: methyl 3,3',4,4'-tetrachlorobenzilate 3s:

10 I.19.1.: 3,3',4,4'-tetrachlorobenzil:

The reaction to form the acyloin derivative is carried out analogously to the process according to step I.11.1 starting from 100 g of 3,4-dichlorobenzaldehyde.

Yield: 60.89 g of oil.

15 60.89 g of the acyloin thus obtained are reacted analogously to the method according to step I.11.1 to obtain the benzil.

Yield: 42.45 g of yellow crystals.

I.19.2.: 3,3',4,4'-tetrachlorobenzilic acid:

20 44.75 g (0.128 mol) of 3,3',4,4'-tetrachlorobenzil are reacted analogously to the method according to step I.11.2 to obtain the corresponding benzilic acid.

Yield: 42 g of yellowish powder; melting point.: 224°C.

I.19.3.: methyl 3,3',4,4'-tetrachlorobenzilate:

25 42 g (0.114 mol) of 3,3',4,4'-tetrachlorobenzilic acid are reacted analogously to step I.10.3. Yield: 15.84 g (= 37% of theory); melting point.: 69°C.

I.20: methyl 3,3',4,4',5,5'-hexafluorobenzilate 3t:

I.20.1.: 3,3',4,4',5,5'-hexafluorobenzil:

30 The reaction to form the acyloin derivative is carried out analogously to the process according to step I.11.1 starting from 31.38 g of 3,4,5-trifluorobenzaldehyde.

Yield: 24.92 g.

24.92 g of the acyloin thus obtained are reacted analogously to the method

35 according to step I.11.1 to obtain the benzil.

Yield: 14.65 g of yellow crystals.

I.20.2.: 3,3',4,4',5,5'-hexafluorobenzilic acid:

14,62 g (0.046 mol) of 3,3',4,4',5,5'-hexafluorobenzil are reacted analogously to the method according to step I.11.2 to obtain the corresponding benzilic acid.

Yield: 15.77 g of yellow crystals.

5 I.20.3.: methyl 3,3',4,4',5,5'-hexafluorobenzilbenzilate:

42 g (0.114 mol) of 3,3',4,4'-tetrachlorobenzilic acid are reacted analogously to step I.10.3. Yield: 5.65 g (= 35% of theory); melting point.: 82-83°C.

II. Synthesis of the compounds of general formula 4:

10

II.1.: tropenol 3,3',4,4'-tetrafluorobenzilate 4.1:

4.27 g (0.013 mol) of ethyl 3,3',4,4'-tetrafluorobenzilate 3a, 1.81 g (0.013 mol) of tropenol and 0.03 g of sodium are heated for 4 h as a melt at 75 mbar over a bath of boiling water with occasional shaking. After cooling, the sodium residues are
15 dissolved with acetonitrile, the solution is evaporated to dryness and the residue is extracted with dichloromethane/water. The organic phase is washed with water, dried over MgSO₄ and evaporated to dryness.

The residue remaining is combined with diethylether/petroleum ether 1:9, suction filtered and washed. Yield: 2.50 g (= 46 % of theory);

20 TLC: R_f value: 0.29 (eluant: sec. butanol/formic acid/water 75:15:10);
melting point: 147°-148°C.

II.2.: scopine 3,3',4,4'-tetrafluorobenzilate 4.2:

4.2 is prepared analogously to the method according to II.1;

25 Yield: 1.75 g (= 36 % of theory) starting from 3.61 g (0.011 mol) of 3a and 1.71 g (0.011 mol) of scopine; TLC: R_f value: 0.22 (eluant as in step II.1);
melting point: 178-179°C;

II.3.: tropenol 4,4'-dichlorobenzilate 4.3:

30 4.3 is prepared analogously to the method according to II.1;

Yield: 6.95 g (= 83 % of theory) starting from 6.5 g (0.02 mol) of 3i and 2.78 g (0.02 mol) of tropenol; TLC: R_f value: 0.30 (eluant as in step II.1); melting point: 197-199°C;

II.4.: tropenol 2,2'-dichlorobenzilate 4.4:

4.4 is prepared analogously to the method according to II.1; the product was precipitated as the hydrochloride and recrystallised from acetonitrile;

Yield: 1.13 g (= 8 % of theory) starting from 9.3 g (0.03 mol) of **3b** and 8.32 g (0.06 mol) of tropenol; TLC: R_f value: 0.26 (eluant as in step II.1); melting point: 253-256°C (hydrochloride);

5 **II.5.: tropenol 4,4'-difluorobenzilate 4.5:**

4.5 is prepared analogously to the method according to II.1;

Yield: 8.71 g (= 69 % of theory) starting from 8.35 g (0.03 mol) of **3c** and 4.18 g (0.03 mol) of tropenol; TLC: R_f value: 0.34 (eluant as in step II.1); melting point: 167-169°C;

10

II.6.: tropenol 2,2',4,4'-tetrafluorobenzilate 4.6:

4.6 is prepared analogously to the method according to II.1;

Yield: 1.80 g (= 27 % of theory) starting from 4.00 g (0.013 mol) of **3d** and 3.54 g (0.036 mol) of tropenol; melting point: 190°C;

15

II.7.: scopine 4,4'-dichlorobenzilate 4.7:

3.78 g (0.01 mol) of tropenol 4,4'-dichlorobenzilate **4.3** are suspended in 40 ml of DMF and heated to 60°C until a clear solution is formed. At an internal temperature of about 40°C a solution of 1.92 g (0.0216 mol) of H₂O₂-urea in 10 ml of water, and
 20 0.183 g (0.0011 mol) of vanadium-(V)-oxide are added and the mixture is stirred for 4.5 h at 60°C. After cooling to 20°C the precipitate formed is suction filtered, the filtrate is adjusted to pH3 with 4 N hydrochloric acid and combined with 0.437 g (0.0023 mol) of Na₂S₂O₇ in 10 ml of water. The green solution thus formed is evaporated to dryness, the residue is extracted with dichloromethane/water. The
 25 acidic aqueous phase is made basic with Na₂CO₃, extracted with dichloromethane and the organic phase is dried over Na₂SO₄ and concentrated. Then 0.5 ml of acetyl chloride is added at about 15°C and the mixture is stirred for 1.5 h. After extraction with 0.1 N hydrochloric acid the aqueous phase is made basic, extracted with dichloromethane, the organic phase is dried over Na₂SO₄ and evaporated to dryness.
 30 The hydrochloride is precipitated from the residue and recrystallised from methanol/diethylether.

Yield: 1.92 g of 18 (= 45 % of theory); TLC: R_f value: 0.29 (eluant as in II.1); melting point: 238-239°C (hydrochloride).

35 **II.8.: scopine 4,4'-difluorobenzilate 4.8:**

4.8 is prepared analogously to the method according to II.7;

Yield: 2.60 g (= 70 % of theory) starting from 3.27 g (0.09 mol) of **4e**;

TLC: R_f value: 0.25 (eluant as in step II.1); melting point: 243-244°C (hydrochloride).

II.9.: tropenol 4,4'-dibromobenzilate 4.9:

4.9 is prepared analogously to the method according to II.1;

Yield: 2.36 g (= 19 % of theory) starting from 9.9 g (0.023 mol) of commercially obtainable isopropyl 4,4'-dibromobenzilate and 3.21 g (0.023 mol) of tropenol; to
5 purify it, the hydrochloride was precipitated and recrystallised from acetonitrile; TLC:
R_f value: 0.30 (eluant as in step II.1);
melting point: 205-207°C (hydrochloride).

II.10.: tropenol 4,4'-dimethylbenzilate 4.10:

10 **4.10** is prepared analogously to the method according to II.1;

Yield: 3.55 g (= 81 % of theory) starting from 2.87 g (0.01 mol) of **3e** and 1.48 g (0.01 mol) of tropenol; to purify it, the hydrochloride was precipitated and recrystallised from acetonitrile; melting point: 232-233°C (hydrochloride).

15 **II.11.: scopine 4,4'-dimethylbenzilate 4.11:**

4.11 is prepared analogously to the method according to II.1;

Yield: 1.02 g (= 24 % of theory) starting from 2.87 g (0.01 mol) of **3e** and 1.65 g (0.01 mol) of scopine; to purify it, the hydrochloride was precipitated;
melting point: 181-183°C (hydrochloride).

20

II.12.: tropine 3,3',4,4'-tetrafluorobenzilate 4.12:

4.12 is prepared analogously to the method according to II.1;

Yield: 2.35 g (= 53 % of theory) starting from 3.45 g (0.01 mol) of **3a** and 1.49 g (0.01 mol) of tropine; melting point: 142-144°C.

25

II.13.: tropenol 3,3',4,4'-tetramethoxybenzilate 4.13:

2.60 g (0.007 mol) of **3f**, 1.00 g (0.007 mol) of tropenol and 0.03 g of sodium are placed in 15 ml of toluene and refluxed for 4 h. After cooling, the mixture is diluted with about 100 ml of toluene, extracted with water, the organic phase is dried over
30 MgSO₄ and evaporated to dryness. Yield: 1.60 g of oil (= 47 % of theory)

II.14.: tropenol 4,4'-dimethoxybenzilate 4.14:

4.14 is prepared analogously to the method according to II.1;

Yield: 3.3 g (= 78 % of theory) starting from 3.0 g (0.01 mol) of **3g** and 1.39 g (0.01
35 mol) of tropenol; melting point: 146-147°C.

II.15.: tropine 3,3',4,4'-tetramethoxybenzilate 4.15:

4.15 is prepared analogously to the method according to II.1;

Yield: 1.65 g (= 32 % of theory; oil) starting from 4.0 g (0.02 mol) of 3f and 3.12 g (0.02 mol) of tropine.

II.16.: scopine 3,3',4,4'-tetramethoxybenzilate 4.16:

5 4.16 is prepared analogously to the method according to II.7;

Yield: 0.8 g (= 41 % of theory, oil) starting from 1.857 g (0.004 mol) of 4.13.

II.17.: tropenol 3,3'-dimethyl-4,4'-dimethoxybenzilate 4.17:

10 4.17 is prepared analogously to the method according to II.1; the product is purified by recrystallising from diethylether; Yield: 2.30 g (= 35 % of theory) starting from 5.0 g (0.015 mol) of 3h and 4.21 g (0.03 mol) of tropenol; melting point: 126°C.

II.18.: scopine 3,3'-dimethyl-4,4'-dimethoxybenzilate 4.18:

4.18 is prepared analogously to the method according to II.7;

15 Yield: 0.6 g (= 44 % of theory, oil) starting from 1.3 g (0.003 mol) of 4q.

II.19.: tropenol 3,3',5,5'-tetrafluorobenzilate 4.19:

1,53 g (0.0636 mol) of NaH are placed in 30 ml of toluene, a solution of 14 g (0.0446 mol) of 3j and 8.85 g (0.0636 mol) of tropenol in 80 ml of toluene is added dropwise
20 at 10°C /860 mbar. The alcohol produced during the reaction is distilled off while at the same time toluene is added dropwise. After 3 h the mixture is cooled and extracted, with dichloromethane and water. The combined organic phases were dried over MgSO₄ and evaporated to dryness. The residue was stirred with petroleum ether/ethyl acetate 95:5.

25 Yield: 11.21 g of light beige crystals (= 60% of theory); melting point.: 168°-170° C.

II.20.: scopine 2,2',4,4'-tetrafluorobenzilate 4.20:

The preparation of 4.20 is carried out analogously to the process according to II.7;

Yield: 1.05 g of white crystals (= 89 % of theory) starting from 1.15 g (0.0027 mol) of
30 4.6.

II.21.: scopine 3,3',5,5'-tetrafluorobenzilate 4.21:

The preparation of 4.21 is carried out analogously to the process according to II.7;

Yield: 1.13 g of white crystals (= 55 % of theory) starting from 2.0 g (0.0047 mol) of
35 4.19. melting point.: 199-200°C.

II.22.: tropenol 3,3'-dichlorobenzilate 4.22:

22.9 g (0.074 mol) of methyl 3,3'-dichlorobenzilate 3k, 15.37 g (0.11 mol) of tropenol and 0.17 g of sodium are heated for 4 h as a melt over a bath of boiling water at 75

mbar with occasional shaking. After cooling the sodium residues are dissolved with acetonitrile, the solution is evaporated to dryness and the residue is extracted with dichloromethane/water. The organic phase is washed with water, dried over MgSO_4 and evaporated to dryness.

- 5 The product is recrystallised from acetonitrile in the form of its hydrochloride. Yield: 16.83 g of white crystals (= 50 % of theory);
melting point.: 184-185°C.

II.23.: tropenol 4,4'-di(trifluoromethyl)benzilate 4.23:

- 10 Starting from 10.0 g (0.0264 mol) of **3l** the reaction is carried out analogously to step II.1. Yield: 4.70 g of beige crystals (= 37 % of theory);
melting point.: 155°C.

II.24.: tropenol 3,3'-di(trifluoromethyl)benzilate 4.24:

- 15 Starting from 6.01 g (0.0159 mol) of **3m** the reaction is carried out analogously to step II.1. Yield: 3.03 g of white crystals (= 39 % of theory);
melting point.: 124-125°C.

II.25.: scopine 4,4'-di(trifluoromethyl)benzilate 4.25:

- 20 The preparation of **4.25** is carried out analogously to the process according to II.7;
Yield: 0.95 g of clear oil (= 46 % of theory) starting from 2.0 g (0.0041 mol) of **4.23**.

II.26.: scopine 3,3'-di(trifluoromethyl)benzilate 4.26:

- The preparation of **4.26** is carried out analogously to the process according to II.7;
25 Yield: 1.02 g of white crystals (= 51 % of theory) starting from 1.94 g (0.0039 mol) of **4.24**.

II.27.: tropenol 3,3'-dichloro-4,4'-difluorobenzilate 4.27:

- Starting from 6.77 g (0.0195 mol) of **3n** the reaction is carried out analogously to
30 step II.1. The crude product obtained is recrystallised from acetonitrile. Yield: 4.05 g
of light beige crystals (= 46 % of theory);
melting point.: 177-179°C.

II.28.: scopine 3,3'-dichlorobenzilate 4.28:

- 35 The preparation of **4.28** is carried out analogously to the process according to II.7;
Yield: 7.86 g (= 75 % of theory) starting from 10.37 g (0.024 mol) of **4.22**.
melting point.: 174-175°C.

II.29.: tropenol 2,2',5,5'-tetrafluorobenzilate 4.29:

Starting from 6.29 g (0.02 mol) of 3o the reaction is carried out analogously to step II.1. The crude product obtained is recrystallised from acetone in the form of the hydrochloride. Yield: 0.89 g of white crystals (= 10 % of theory); melting point.: 177-179°C.

5

II.30.: tropenol 2,2',3,3'-tetrafluorobenzilate 4.30:

Starting from 8.0 g (0.0255 mol) of 3p the reaction is carried out analogously to step II.1. The crude product obtained is recrystallised from acetonitrile. Yield: 0.96 g of beige crystals (= 9 % of theory);

10 melting point.: 176-177°C.

II.31.: tropenol 3,3'-difluorobenzilate 4.31:

Starting from 11.13 g (0.04 mol) of 3q the reaction is carried out analogously to step II.1. The crude product obtained is recrystallised from acetonitrile in the form of the hydrochloride. Yield: 7.98 g (= 47 % of theory); melting point.: 245-246°C.

15

II.32.: scopine 3,3'-difluorobenzilate 4.32:

The preparation of 4.32 is carried out analogously to the process according to II.7; Yield: 4.2 g (= 76 % of theory) starting from 4.89 g (0.013 mol) of 4.31.

20 melting point.: 216-218°C.

II.33.: scopine 3,3'-dichloro-4,4'-difluorobenzilate 4.33:

The preparation of 4.33 is carried out analogously to the process according to II.7; Yield: 0.78 g of white crystals (= 34 % of theory) starting from 4.6 g (0.0098 mol) of

25 4.27; melting point.: 216-218°C.

II.34.: tropenol 4,4'-dichloro-3,3'-difluorobenzilate 4.34:

Starting from 12.0 g (0.0345 mol) of 3r the reaction is carried out analogously to step II.1. The crude product obtained is extracted from petroleum ether.

30 Yield: 6.65 g of creamy-white powder (= 42 % of theory);

melting point.: 180-181°C.

II.35.: scopine 4,4'-dichloro-3,3'-difluorobenzilate 4.35:

The preparation of 4.35 is carried out analogously to the process according to II.7;

35 Yield: 2.58 g of white crystals (= 62 % of theory) starting from 4.0 g (0.0088 mol) of 4.34; melting point.: 150-151°C.

II.36.: tropenol 3,3',4,4'-tetrachlorobenzilate 4.36:

Starting from 14.24 g (0.0375 mol) of **3s** the reaction is carried out analogously to step II.1. The crude product obtained is recrystallised from acetonitrile. Yield: 4.81 g of white crystals (= 26 % of theory);
melting point.: 149-150°C.

5

II.37.: tropenol 3,3',4,4',5,5'-hexafluorobenzilate **4.37:**


Starting from 5.0 g (0.0143 mol) of **3t** the reaction is carried out analogously to step II.1. The crude product obtained is recrystallised from acetonitrile. Yield: 3.49 g of white crystals (= 53 % of theory);

10 melting point.: 164-165°C.

III. Synthesis of the compounds of general formula **1:**

Example 1: tropenol 3,3',4,4'-tetrafluorobenzilate methobromide :

15



2.00 g (0.0048 mol) of **4.1**, 30 ml of acetonitrile, 10 ml of dichloromethane and 2.88 g (0.0143 mol) of 46.92 % bromomethane in acetonitrile are combined at 20°C and left to stand for 3 days. The solution is evaporated to dryness and the residue is

20 recrystallised from acetonitrile. Yield: 1.95 g (= 80 % of theory)

TLC: R_f value: 0.12 (eluant as in step II.1); melting point: 238°C.


C₂₃H₂₂F₄NO₃xBr (516.33);

Elemental analysis: calculated: C (53.50) H (4.29) N (2.71)

found: C (53.52) H (4.30) N (2.65).

25

Example 2: scopine 3,3',4,4'-tetrafluorobenzilate -methobromide :



1.5 g (0.0034 mol) of 4.2, 20 ml of acetonitrile, 20 ml of dichloromethane and 2.08 g (0.01 mol) of 46.92 % bromomethane in acetonitrile are reacted analogously to Example 1.

5 Yield: 1.40 g (= 77 % of theory)

TLC: R_f value: 0.16 (eluant as in step II.1); melting point: 227°C.


C₂₃H₂₂F₄NO₄xBr (532.33);

Elemental analysis: calculated: C (51.90) H (4.17) N (2.63)

found: C (51.91) H (4.16) N (2.60).

10

Example 3: tropenol 4,4'-dichlorobenzilate methobromide :



2.09 g (0.005 mol) of 4.3 are reacted analogously to Example 1. The crystals formed
15 are suction filtered, washed with dichloromethane, dried and then recrystallised from methanol/diethylether.

Yield: 1.72 g (=67 % of theory)


C₂₃H₂₄Cl₂NO₃xBr (513.26);

TLC: R_f value: 0.12 (eluant as in step II.1); melting point: 195-196°C.

20 Elemental analysis: calculated: C (53.82) H (4.71) N (2.73)

found: C (53.54) H (4.80) N (2.73).

Example 4: tropenol 2,2'-dichlorobenzilate methobromide :



0.86 g (0.0021 mol) of the free base of 4.4 are reacted analogously to Example 1. The crystals formed are suction filtered, washed with acetone, dried and then recrystallised from methanol/diethylether.

5 Yield: 0.99 g (= 94 % of theory)

TLC: R_f value: 0.14 (eluant as in step II.1); melting point: 260-261°C.


C₂₃H₂₄Cl₂NO₃xBr (513.26);

Elemental analysis: calculated: C (53.82) H (4.71) N (2.73)

found: C (53.62) H (4.76) N (2.69).

10

Example 5: tropenol 4,4'-difluorobenzilate methobromide :



1.9 g (0.005 mol) of the free base of 4.5 are reacted analogously to Example 1.

Yield: 2.1 g (= 89 % of theory)


15 TLC: R_f value: 0.14 (eluant as in step II.1); melting point: 219-220°C.

C₂₃H₂₄F₂NO₃xBr (480.35);

Elemental analysis: calculated: C (57.51) H (5.04) N (2.92)

found: C (57.33) H (4.86) N (2.90).

20 **Example 6: tropenol 2,2',4,4'-tetrafluorobenzilate methobromide :**



1.60 g (0.004 mol) of 4.6 are reacted analogously to Example 1. The crystals formed are suction filtered and recrystallised from acetone.

Yield: 1.70 g (= 87 % of theory)

- 5 TLC: R_f value: 0.13 (eluant: n-butanol/water/formic acid(conc.)/acetone/dichloromethane 36:15:15:15:5); melting point: 241-242°C.


C₂₃H₂₂F₄NO₃xBr (516.33);

Elemental analysis: calculated: C (53.50) H (4.29) N (2.71)

found: C (53.55) H (4.33) N (2.73).

10

Example 7: scopine 4,4'-dichlorobenzilate methobromide :



1.54 g (0.0035 mol) of the free base of 4.7 are reacted analogously to Example 1.

- 15 The crystals formed are suction filtered, washed with acetone, dried and then recrystallised from methanol/diethylether.

Yield: 1.68 g (= 90 % of theory)


TLC: R_f value: 0.22 (eluant as in step II.1); melting point: 209-210°C.

C₂₃H₂₄Cl₂NO₄xBr (529.26);

- 20 Elemental analysis: calculated: C (52.20) H (4.57) N (2.65)

found: C (51.25) H (4.83) N (2.49).

Example 8: scopine 4,4'-difluorobenzilate methobromide :



1,50 g (0.003 mol) of the free base of 4.8 are reacted analogously to Example 1. The crystals formed are suction filtered, washed with acetone, dried and then recrystallised from methanol/diethylether.

5 Yield: 1.73 g (= 93 % of theory)

TLC: R_f value: 0.19 (eluant as in step II.1); melting point: 224-225°C.


C₂₃H₂₄F₂NO₄xBr (496.35);

Elemental analysis: calculated: C (55.66) H (4.87) N (2.82)

found: C (55.20) H (4.81) N (2.82).

10

Example 9: tropenol 2,2',4,4'-tetrafluorobenzilate methobromide :



15 1.50 g (0.003 mol) of the free base of 4.9 are reacted analogously to Example 1. The crystals formed are recrystallised from methanol.

Yield: 1.53 g (= 86 % of theory)


TLC: R_f value: 0.14 (eluant as in step II.1); melting point: 175-177°C.

C₂₃H₂₄Br₂NO₃xBr (602.16);

20 Elemental analysis: calculated: C (45.88) H (4.02) N (2.33)

found: C (45.46) H (4.42) N (2.18).

Example 10: tropenol 4,4'-dimethylbenzilate methobromide :



2.60 g (0.007 mol) of the free base of 4.10 are reacted analogously to Example 1.
The crystals formed are recrystallised from ethanol.

Yield: 3.16 g (= 97 % of theory)


5 TLC: R_f value: 0.14 (eluant as in step II.1);

C₂₅H₃₀NO₃·Br (472.42);

Elemental analysis: calculated: C (63.56) H (6.40) N (2.96)

found: C (62.88) H (6.87) N (2.74).

10 **Example 11: scopine 4,4'-dimethylbenzilate -methobromide :**



0.56 g (0.0014 mol) of the free base of 4.11 are reacted analogously to Example 1.
The crystals formed are recrystallised from methanol/diethylether.

15 Yield: 0.55 g (= 80 % of theory)

TLC: R_f value: 0.19 (eluant as in step II.1); melting point: 221-222°C;


C₂₅H₃₀NO₄·Br (488.42);

Elemental analysis: calculated: C (61.48) H (6.19) N (2.87)

found: C (60.61) H (6.31) N (2.80).

20

Example 12: tropine 3,3',4,4'-tetrafluorobenzilate methobromide :



1.80 g (0.004 mol) of the free base of **4.12** are reacted analogously to Example 1.

Yield: 1.73 g (=78 % of theory)

TLC: R_f value: 0.10 (eluant as in step II.1); melting point: 157°C;


5 $C_{23}H_{24}F_4NO_3 \cdot Br$ (518.34);

Elemental analysis: calculated: C (53.30) H (4.67) N (2.70)

found: C (53.39) H (4.53) N (2.73).

Example 13: tropenol 3,3',4,4'-tetramethoxybenzilate -methobromide :

10



1.20 g (0.003 mol) of **4.13** are reacted analogously to Example 1. The crystals formed are recrystallised from acetonitrile/diethylether.

Yield: 1.05 g (=73 % of theory)


15 TLC: R_f value: 0.10 (eluant as in step II.1); melting point: 212°C;

$C_{27}H_{34}NO_7 \cdot Br$ (564.47);

Elemental analysis: calculated: C (57.45) H (6.07) N (2.48)

found: C (56.91) H (6.05) N (2.45).

20 **Example 14: tropenol 4,4'-dimethoxybenzilate methobromide :**



1,13 g (0.003 mol) of 4.14 are reacted analogously to Example 1. The crystals formed are recrystallised from acetonitrile/diethylether.

Yield: 1.21 g (= 87 % of theory)


5 $C_{25}H_{30}NO_5 \times Br$ (504.42);

TLC: R_f value: 0.01 (eluant as in step II.1); melting point: 180-181°C;

Elemental analysis: calculated: C (59.53) H (5.99) N (2.78)

found: C (59.29) H (6.24) N (2.84).

10 Example 15: tropine 3,3',4,4'-tetramethoxybenzilate methobromide :



1.65 g (0.004 mol) of 4.15 are reacted analogously to Example 1. The crystals formed are recrystallised from diethylether.


15 Yield: 1.70 g (= 86 % of theory); melting point: 148-150°C;

$C_{27}H_{36}NO_7 \times Br$ (566.49);

Elemental analysis: calculated: C (57.25) H (6.41) N (2.47)

found: C (56.41) H (6.75) N (2.45).

20 Example 16: scopine 3,3',4,4'-tetramethoxybenzilate methobromide :



0.80 g (0.0017 mol) of **4.16** are reacted analogously to Example 1. The crystals formed are recrystallised from acetone.

Yield: 0.35 g (= 37 % of theory); melting point: 211-212°C;


5 $C_{27}H_{34}NO_8 \times Br$ (580.47);

Elemental analysis: calculated: C (55.87) H (5.90) N (2.41)

found: C (55.62) H (6.09) N (2.53).

Example 17: tropenol 3,3'-dimethyl-4,4'-dimethoxybenzilate methobromide :

10



1.00 g (0.003 mol) of **4.17** are reacted analogously to Example 1.


Yield: 0.85 g (=70 % of theory); melting point: 217°C;

$C_{27}H_{34}NO_5 \times Br$ (532.47);

15 Elemental analysis: calculated: C (60.90) H (6.51) N (2.63)

found: C (59.83) H (6.51) N (2.93).

Example 18: scopine 3,3'-dimethyl-4,4'-dimethoxybenzilate :



0.60 g (0.0013 mol) of **4.18** are reacted analogously to Example 1.

The product is purified by crystallisation from acetone;


Yield: 0.40 g (= 56 % of theory);

5 $C_{27}H_{34}NO_6 \cdot Br$ (548.47);

Elemental analysis: calculated: C (59.13) H (6.25) N (2.55)

found: C (58.69) H (6.54) N (2.61).

Example 19: tropenol 3,3',4,4'-tetrafluorobenzilate -ethylbromide :



10

2.00 g (0.0048 mol) of **4.1** is dissolved in 20 ml dichloromethane and 20 ml acetonitrile, combined with 2.59 g (0.0238 mol) of bromoethane and the reaction vessel is sealed. It is left to stand at about 20°C for 3 weeks with the exclusion of light. The solution is evaporated to dryness and the residue is recrystallised from

15 acetonitrile.

Yield: 1.96 g of 3 (= 78 % of theory)


TLC: R_f value: 0.11 (eluant as in step II.1); melting point: 247°C.

Elemental analysis: calculated: C (54.35) H (4.56) N (2.64)

found: C (53.93) H (4.59) N (2.60).

20

Example 20: tropenol 3,3',5,5'-tetrafluorobenzilate methobromide :



2.0 g (0.0047 mol) of **4.19** are reacted analogously to Example 1.

Yield: 2.22 g of white crystals (=92 % of theory); melting point.: 262-264°C;


$C_{23}H_{22}F_4NO_3 \times Br$ (516.33);

5 Elemental analysis: calculated: C (53,50) H (4,29) N
(2,71)

found : C (53,48) H (4,30) N (2,65).

Example 21: scopolamine 2,2',4,4'-tetrafluorobenzilate methobromide :

10



1,05 g (0.0024 mol) of **4.20** are reacted analogously to Example 1.


Yield: 0.54 g of white crystals (=42 % of theory); melting point.: 208-209°C;

$C_{23}H_{22}F_4NO_4 \times Br$ (532.33);

15 Elemental analysis: calculated: C (51.90) H (4.17) N
(2.63)

found : C (50.84) H (4.39) N (2.50).

Example 22: scopolamine 3,3',5,5'-tetrafluorobenzilate methobromide :



1,13 g (0.00258 mol) of **4.21** are reacted analogously to Example 1.


Yield: 1.07 g of white crystals (=78 % of theory); melting point.: 238-239°C;

$C_{23}H_{22}F_4NO_4 \cdot Br$ (532.33);

5 Elemental analysis: calculated: C (51.90) H (4.17) N (2.63)

found : C (51.85) H (4.29) N (2.70).

Example 23: tropenol 3,3'-dichlorobenzilate methobromide :



10

2.09 g (0.005 mol) of **4.22** are reacted analogously to Example 1.

Yield: 2.42 g of white crystals (=94 % of theory); melting point.: 200-201°C;


C₂₃H₂₄Cl₂NO₃xBr (513.26);

Elemental analysis: calculated: C (53.82) H (4.71) N (2.73)

15

found : C (53.73) H (4.74) N (2.78).

Example 24: tropenol 4,4'-di(trifluoromethyl)benzilate methobromide :




1.2 g (0.00247 mol) of **4.23** are reacted analogously to Example 1.

Yield: 1.05 g of white crystals (=73 % of theory); melting point.: 140-141°C;

$C_{25}H_{24}F_6NO_3 \cdot Br$ (580.36);

5

Example 25: tropenol 3,3'-di(trifluoromethyl)benzilate methobromide :



1.09 g (0.00225 mol) of **4.24** are reacted analogously to Example 1.


10 Yield: 0.84 g of white crystals (=65 % of theory); melting point.: 228-229°C;

$C_{25}H_{24}F_6NO_3 \cdot Br$ (580.36);

Elemental analysis: calculated: C (51.74) H (4.17) N (2.41)

found : C (51.40) H (4.24) N (2.42).

15 **Example 26:** scopine 4,4'-di(trifluoromethyl)benzilate methobromide :




42

0.95 g (0.0019 mol) of **4.25** are reacted analogously to Example 1. The product is recrystallised from diethylether. Yield: 0.98 g of white crystals (=87 % of theory); melting point.: 158-160°C; $C_{25}H_{24}F_6NO_4 \cdot Br$ (596.36);

Elemental analysis: calculated: C (50.35) H (4.06) N (2.35)

5 found : C (50.34) H (4.03) N (2.36).

Example 27: scopine 3,3'-di(trifluoromethyl)benzilate methobromide :




10 1.0 g (0.002 mol) of **4.26** are reacted analogously to Example 1. The product is recrystallised from acetonitrile. Yield: 0.7 g of white crystals (=59 % of theory); melting point.: 220-221°C; $C_{25}H_{24}F_6NO_4 \cdot Br$ (596.36);

Elemental analysis: calculated: C (50.35) H (4.06) N (2.35)

found : C (50.24) H (4.17) N (2.40).

15

Example 28: tropenol 4,4'-difluorobenzilate -ethyl bromide :




1,54 g (0.004 mol) of **4.5** are reacted analogously to Example 19. The product is recrystallised from ethanol. Yield: 1.72 g of white crystals (=87 % of theory); melting point.: 228-229°C; $C_{24}H_{26}F_2NO_3 \cdot Br$ (494.37);

20 Elemental analysis: calculated: C (58.31) H (5.30) N (2.83)


found : C (58.25) H (5.29) N (2.83).

Example 29: tropenol 3,3'-dichloro-4,4'-difluorobenzilate methobromide :




1.5 g (0.0033 mol) of **4.27** are reacted analogously to Example 1. The product is
5 recrystallised from acetonitrile. Yield: 1.49 g of white crystals (=82 % of theory);
melting point.: 245-246°C; C₂₃H₂₂Cl₂F₂NO₃xBr (549.24);
Elemental analysis: calculated: C (50.30) H (4.04) N (2.55)
found : C (50.44) H (4.19) N (2.51).

10 Example 30: scopine 3,3'-dichlorobenzilate methobromide :



2,61 g (0.006 mol) of **4.28** are reacted analogously to Example 1. The product is recrystallised from ethanol. Yield: 2.13 g of white crystals (=67 % of theory);
15 melting point.: 221-222°C; C₂₃H₂₄Cl₂NO₄xBr (529.26);
Elemental analysis: calculated: C (52,20) H (4.57) N (2.65)
found : C (52,25) H (4.61) N (2.70).

Example 31: 2. tropenol 2',5,5'-tetrafluorobenzilate methobromide :

0,67 g (0.0016 mol) of **4.29** are reacted analogously to Example 1. The product is
5 recrystallised from methanol/diethylether.


Yield: 0.70 g of white crystals (=86 % of theory);

melting point.: 269-270°C; C₂₃H₂₂F₄NO₃xBr (516,33);

Elemental analysis: calculated: C (53.50) H (4.29) N (2.71)

found : C (53.30) H (4.52) N (2.76).

10

Example 32: 2. tropenol 2',3,3'-tetrafluorobenzilate methobromide :

0,96 g (0.002 mol) of **4.30** are reacted analogously to Example 1. The product is
15 recrystallised from acetonitrile.

Yield: 0.61 g of white crystals (=59 % of theory);


melting point.: 268-269°C; C₂₃H₂₂F₄NO₃xBr (516.33);

Elemental analysis: calculated: C (53.50) H (4.29) N (2.71)

found : C (53.56) H (4.38) N (2.75).


20

Example 33: scopine 4,4'-difluorobenzilate -ethyl bromide :




1.2 g (0.003 mol) of 4.8 are reacted analogously to Example 19. The product is
5 recrystallised from ethanol. Yield: 0.93 g of white crystals (=61 % of theory);
melting point.: 162-163°C; C₂₄H₂₆F₂NO₄xBr (510.38);
Elemental analysis: calculated: C (56.48) H (5.13) N (2.74)
found : C (55.96) H (5.30) N (2.75).

10 **Example 34:** tropenol 3,3'-difluorobenzilate methobromide :



1,61 g (0.004 mol) of **4.31** are reacted analogously to Example 1. The product is recrystallised from ethanol/diethylether.

15 Yield: 1.93 g of white crystals (=96 % of theory);
melting point.: 227-228°C; C₂₃H₂₄F₂NO₃xBr (480.35);
Elemental analysis: calculated: C (57.51) H (5.04) N (2.92)
found : C (57.38) H (5.14) N (2.95).

Example 35: scopine 3,3'-difluorobenzilate methobromide :

1.61 g (0.004 mol) of **4.32** are reacted analogously to Example 1. The product is
5 recrystallised from ethanol.


Yield: 1.83 g of white crystals (=92 % of theory);

melting point.: 221-222°C; C₂₃H₂₄F₂NO₄xBr (496.35);

Elemental analysis: calculated: C (55.66) H (4.87) N (2.82)

found : C (55.49) H (4.78) N (2.73).

10

Example 36: scopine 3,3'-dichloro-4,4'-difluorobenzilate methobromide :

0,78 g (0.002 mol) of **4.33** are reacted analogously to Example 1. The product is
15 recrystallised from acetonitrile.


Yield: 0.67 g of white crystals (=59 % of theory);

C₂₃H₂₂Cl₂F₂NO₄xBr (565,24);


Elemental analysis: calculated: C (48,87) H (3,92) N (2.48)

found : C (48,87) H (3,81) N (2.46).

20

Example 37: scopine 4,4'-dichloro-3,3'-difluorobenzilate methobromide :

2.4 g (0.0051 mol) of **4.35** are reacted analogously to Example 1. The product is
 5 recrystallised from acetonitrile. Yield: 2.45 g of white crystals (=85 % of theory);
 melting point.: 211-212°C; $C_{23}H_{22}Cl_2F_2NO_4 \cdot Br$ (565.24).

Example 38: tropenol 3,3',4,4'-tetrachlorobenzilate methobromide :

10

1.8 g (0.00369 mol) of **4.36** are reacted analogously to Example 1. The product is
 recrystallised from methanol/diethylether.


Yield: 2.01 g of white crystals (=93 % of theory);

melting point.: 245-246°C; $C_{23}H_{22}Cl_4NO_3 \cdot Br$ (582.15);

15 Elemental analysis: calculated: C (47.45) H (3.81) N (2.41)

found : C (47.27) H (3.82) N (2.36).


Example 39: tropenol 4,4'-dichloro-3,3'-difluorobenzilate methobromide :



2.5 g (0.0055 mol) of 4.34 are reacted analogously to Example 1. The product is recrystallised from acetonitrile. Yield: 1.53 g of white crystals (=51 % of theory); melting point.: 229-231°C; $C_{23}H_{22}Cl_2F_2NO_3 \cdot Br$ (549.24).

5

Example 40: tropenol 3,3',4,4',5,5'-tetrafluorobenzilate methobromide :



1.7 g (0.0037 mol) of 4.37 are reacted analogously to Example 1. The product is extracted from acetone. Yield: 1.9 g of white crystals (=92 % of theory); melting point.: 241-242°C; $C_{23}H_{20}F_6NO_3 \cdot Br$ (552.32).

Elemental analysis: calculated: C (50.02) H (3.65) N (2.54)

found : C (50.09) H (3.61) N (2.49).

- 15 As has been found, the compounds of general formula 1 are characterised by their versatility in therapeutic use. Particular mention should be made of those applications for which the compounds of formula 1 according to the invention are preferably used on the basis of their pharmaceutical activity as anticholinergic agents.
- 20 These include, for example, the treatment of asthma or COPD (chronic obstructive pulmonary disease). The compounds of general formula 1 may also be used to treat vagally induced sinus bradycardia and to treat heart rhythm disorders. In general, the compounds according to the invention may also be used to treat spasms, e.g. in

the gastrointestinal tract, with therapeutic benefit. They may also be used in the treatment of spasms in the urinary tract and in menstrual disorders, for example. Of the ranges of indications mentioned above, the treatment of asthma and COPD using the compounds of formula 1 according to the invention is of particular importance.

The compounds of general formula 1 may be used on their own or combined with other active substances of formula 1 according to the invention.

- 10 The compounds of general formula 1 may optionally also be combined with other pharmacologically active substances. These include, in particular, betamimetics, antiallergic agents, PAF-antagonists, leukotriene-antagonists and corticosteroids and combinations of these active substances.
- 15 Examples of betamimetics which may be used in conjunction with the compounds of formula 1 according to the invention include compounds selected from among bambuterol, bitolterol, carbuterol, clenbuterol, fenoterol, formoterol, hexoprenaline, ibuterol, pirbuterol, procaterol, reproterol, salmeterol, sulphonterol, terbutaline, tulobuterol, 4-hydroxy-7-[2-{{2-{{3-(2-phenylethoxy)propyl}sulphonyl}ethyl}-
- 20 amino}ethyl]-2(3H)-benzothiazolone, 1-(2-fluoro-4-hydroxyphenyl)-2-[4-(1-benzimidazolyl)-2-methyl-2-butylamino]ethanol, 1-[3-(4-methoxybenzyl-amino)-4-hydroxyphenyl]-2-[4-(1-benzimidazolyl)-2-methyl-2-butylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-N,N-dimethylaminophenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-
- 25 methoxyphenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-n-butyloxyphenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-{{4-{{3-(4-methoxyphenyl)-1,2,4-triazol-3-yl}-2-methyl-2-butylamino}ethanol, 5-hydroxy-8-(1-hydroxy-2-isopropylaminobutyl)-2H-1,4-benzoxazin-3-(4H)-on, 1-(4-amino-3-chloro-5-trifluoromethylphenyl)-2-*tert.*-
- 30 butylamino)ethanol and 1-(4-ethoxycarbonylamino-3-cyano-5-fluorophenyl)-2-(*tert.*-butylamino)ethanol, optionally in the form of their racemates, their enantiomers, their diastereomers, as well as optionally their pharmacologically acceptable acid addition salts and hydrates. It is particularly preferable to use, as betamimetics, active substances of this kind, combined with the compounds of formula 1 according to the
- 35 invention, selected from among fenoterol, formoterol, salmeterol, 1-[3-(4-methoxybenzyl-amino)-4-hydroxyphenyl]-2-[4-(1-benzimidazolyl)-2-methyl-2-butylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-N,N-dimethylaminophenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-methoxyphenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-

- 5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-n-butyloxyphenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-{4-[3-(4-methoxyphenyl)-1,2,4-triazol-3-yl]-2-methyl-2-butylamino}ethanol, optionally in the form of their racemates, their enantiomers, their diastereomers, as well as optionally
- 5 their pharmacologically acceptable acid addition salts and hydrates. Of the betamimetics mentioned above, the compounds formoterol and salmeterol, optionally in the form of their racemates, their enantiomers, their diastereomers, as well as optionally their pharmacologically acceptable acid addition salts and hydrates, are particularly important.
- 10 The acid addition salts of the betamimetics selected from among the hydrochloride, hydrobromide, sulphate, phosphate, fumarate, methanesulphonate and xinafoate are preferred according to the invention. In the case of salmeterol, the salts selected from among the hydrochloride, sulphate and xinafoate are particularly preferred, especially the sulphates and xinafoates. Of outstanding importance according to the
- 15 invention are salmeterol x $\frac{1}{2}$ H₂SO₄ and salmeterol xinafoate. In the case of formoterol, the salts selected from among the hydrochloride, sulphate and fumarate are particularly preferred, especially the hydrochloride and fumarate. Of outstanding importance according to the invention is formoterol fumarate.
- 20 Within the scope of the present invention, the term corticosteroids, which may optionally be used in conjunction with the compounds of formula 1, denotes compounds selected from among flunisolide, beclomethasone, triamcinolone, budesonide, fluticasone, mometasone, ciclesonide, rofleponide, GW 215864, KSR 592, ST-126 and dexamethasone. The preferred corticosteroids within the scope of
- 25 the present invention are those selected from among flunisolide, beclomethasone, triamcinolone, budesonide, fluticasone, mometasone, ciclesonide and dexamethasone, while budesonide, fluticasone, mometasone and ciclesonide, especially budesonide and fluticasone, are of particular importance. The term steroids may be used on its own, within the scope of the present patent application,
- 30 instead of the term corticosteroids. Any reference to steroids within the scope of the present invention also includes a reference to salts or derivatives which may be formed from the steroids. Examples of possible salts or derivatives include: sodium salts, sulphobenzoates, phosphates, isonicotinates, acetates, propionates, dihydrogen phosphates, palmitates, pivalates or furoates. The corticosteroids may
- 35 optionally also be in the form of their hydrates.

Within the scope of the present invention, the term dopamine agonists, which may optionally be used in conjunction with the compounds of formula 1, denotes compounds selected from among bromocriptine, cabergolin, alpha-

dihydroergocryptine, lisuride, pergolide, pramipexol, roxindol, ropinirol, talipexol, tergurid and viozan. It is preferable within the scope of the present invention to use, as combination partners with the compounds of formula 1, dopamine agonists selected from among pramipexol, talipexol and viozan, pramipexol being of particular
 5 importance. Any reference to the abovementioned dopamine agonists also includes, within the scope of the present invention, a reference to any pharmacologically acceptable acid addition salts and hydrates thereof which may exist. By the physiologically acceptable acid addition salts thereof which may be formed by the abovementioned dopamine agonists are meant, for example, pharmaceutically
 10 acceptable salts selected from among the salts of hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, methanesulphonic acid, acetic acid, fumaric acid, succinic acid, lactic acid, citric acid, tartaric acid and maleic acid.

Examples of antiallergic agents which may be used according to the invention as a
 15 combination with the compounds of formula 1 include epinastin, cetirizin, azelastin, fexofenadin, levocabastin, loratadine, mizolastin, ketotifen, emedastin, dimetinden, clemastine, bamipin, cexchlorpheniramine, pheniramine, doxylamine, chlorphenoxamine, dimenhydrinate, diphenhydramine, promethazine, ebastin, desloratidine and meclizine. Preferred antiallergic agents which may be used within
 20 the scope of the present invention in combination with the compounds of formula 1 according to the invention are selected from among epinastin, cetirizin, azelastin, fexofenadin, levocabastin, loratadine, ebastin, desloratidine and mizolastin, epinastin and desloratidine being particularly preferred. Any reference to the abovementioned antiallergic agents also includes, within the scope of the present invention, a
 25 reference to any pharmacologically acceptable acid addition salts thereof which may exist.

The following are examples of PAF antagonists which may be used in conjunction with the compounds of formula 1 according to the invention:

30 4-(2-chlorophenyl)-9-methyl-2-[3(4-morpholinyl)-3-propanon-1-yl]-6H-thieno-[3,2-f] [1,2,4]triazolo[4,3-a][1,4]diazepine,
 6-(2-chlorophenyl)-8,9-dihydro-1-methyl-8-[(4-morpholinyl)carbonyl]-4H,7H-cyclopenta-[4,5]thieno-[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine.

35 Suitable preparations for administering the compounds of formula 1 include tablets, capsules, suppositories, solutions, etc.

Of particular importance according to the invention (particularly when treating asthma or COPD) is the administration of the compounds according to the invention by inhalation. The proportion of pharmaceutically active compound or compounds

should be in the range from 0.05 to 90 % by weight, preferably 0.1 to 50 % by weight of the total composition. Suitable tablets may be obtained, for example, by mixing the active substance(s) with known excipients, for example inert diluents such as calcium carbonate, calcium phosphate or lactose, disintegrants such as corn starch
5 or alginic acid, binders such as starch or gelatine, lubricants such as magnesium stearate or talc and/or agents for delaying release, such as carboxymethyl cellulose, cellulose acetate phthalate, or polyvinyl acetate. The tablets may also comprise several layers.

- 10 Coated tablets may be prepared accordingly by coating cores produced analogously to the tablets with substances normally used for tablet coatings, for example collidone or shellac, gum arabic, talc, titanium dioxide or sugar. To achieve delayed release or prevent incompatibilities the core may also consist of a number of layers. Similarly the tablet coating may consist of a number of layers to achieve delayed
15 release, possibly using the excipients mentioned above for the tablets.

Syrups or elixirs containing the active substances or combinations thereof according to the invention may additionally contain a sweetener such as saccharine, cyclamate, glycerol or sugar and a flavour enhancer, e.g. a flavouring such as
20 vanillin or orange extract. They may also contain suspension adjuvants or thickeners such as sodium carboxymethyl cellulose, wetting agents such as, for example, condensation products of fatty alcohols with ethylene oxide, or preservatives such as p-hydroxybenzoates.

- 25 Solutions are prepared in the usual way, e.g. with the addition of isotonic agents, preservatives such as p-hydroxybenzoates, or stabilisers such as alkali metal salts of ethylenediamine tetraacetic acid, optionally using emulsifiers and/or dispersants, whilst if water is used as the diluent, for example, optionally organic solvents may optionally be used as solvating agents or dissolving aids, and transferred into
30 injection vials or ampoules or infusion bottles.

Capsules containing one or more active substances or combinations of active substances may for example be prepared by mixing the active substances with inert carriers such as lactose or sorbitol and packing them into gelatine capsules.

- 35 Suitable suppositories may be made for example by mixing with carriers provided for this purpose, such as neutral fats or polyethyleneglycol or the derivatives thereof. Excipients which may be used include, for example, water, pharmaceutically acceptable organic solvents such as paraffins (e.g. petroleum fractions), vegetable oils (e.g. groundnut or sesame oil), mono- or polyfunctional alcohols (e.g. ethanol or

glycerol), carriers such as e.g. natural mineral powders (e.g. kaolins, clays, talc, chalk), synthetic mineral powders (e.g. highly dispersed silicic acid and silicates), sugars (e.g. cane sugar, lactose and glucose), emulsifiers (e.g. lignin, spent sulphite liquors, methylcellulose, starch and polyvinylpyrrolidone) and lubricants (e.g. magnesium stearate, talc, stearic acid and sodium lauryl sulphate).

The preparations are administered by the usual methods, preferably by inhalation in the treatment of asthma or COPD. For oral administration the tablets may, of course, contain, apart from the abovementioned carriers, additives such as sodium citrate, calcium carbonate and dicalcium phosphate together with various additives such as starch, preferably potato starch, gelatine and the like. Moreover, lubricants such as magnesium stearate, sodium lauryl sulphate and talc may be used at the same time for the tableting process. In the case of aqueous suspensions the active substances may be combined with various flavour enhancers or colourings in addition to the excipients mentioned above.

The dosage of the compounds according to the invention is naturally greatly dependent on the route of administration and the complaint to be treated. When administered by inhalation the compounds of formula 1 are characterised by high efficacy even at doses in the μg range. The compounds of formula 1 can also be used effectively above the μg range. The dosage may then be in the gram range, for example. Particularly when administered by a method other than inhalation, the compounds according to the invention may be given in higher doses (in the range from 1 to 1000 mg, for example, although this does not imply any limitation).

The examples of formulations which follow illustrate the present invention without restricting its scope:

Examples of pharmaceutical formulations

30	A)	<u>Tablets</u>	<u>per tablet</u>
		active substance	100 mg
		lactose	140 mg
35		corn starch	240 mg
		polyvinylpyrrolidone	15 mg
		magnesium stearate	5 mg
			500 mg

The finely ground active substance, lactose and some of the corn starch are mixed together. The mixture is screened, then moistened with a solution of polyvinylpyrrolidone in water, kneaded, wet-granulated and dried. The granules, the remaining corn starch and the magnesium stearate are screened and mixed
 5 together. The mixture is compressed to produce tablets of suitable shape and size.

B) <u>Tablets</u>		<u>per tablet</u>
10	active substance	80 mg
	lactose	55 mg
	corn starch	190 mg
	microcrystalline cellulose	35 mg
	polyvinylpyrrolidone	15 mg
15	sodium-carboxymethyl starch	23 mg
	magnesium stearate	2 mg
		400 mg

20 The finely ground active substance, some of the corn starch, lactose, microcrystalline cellulose and polyvinylpyrrolidone are mixed together, the mixture is screened and worked with the remaining corn starch and water to form a granulate which is dried and screened. The sodium carboxymethyl starch and the magnesium stearate are added and mixed in and the mixture is compressed to form tablets of a
 25 suitable size.

C) <u>Ampoule solution</u>	
	active substance 50 mg
30	sodium chloride 50 mg
	water for inj. 5 ml

The active substance is dissolved in water at its own pH or optionally at pH 5.5 to 6.5 and sodium chloride is added to make it isotonic. The solution obtained is filtered
 35 free from pyrogens and the filtrate is transferred under aseptic conditions into ampoules which are then sterilised and sealed by fusion. The ampoules contain 5 mg, 25 mg and 50 mg of active substance.

D) Metering aerosol

	Active substance	0.005
	Sorbitan trioleate	0.1
5	Monofluorotrichloromethane and difluorodichloromethane 2 : 3	ad 100

The suspension is transferred into a conventional aerosol container with a metering valve. Preferably, 50 µl of suspension are delivered per spray. The active substance may also be metered in higher doses if desired (e.g. 0.02 % by weight).

10

E) Solutions (in mg/100ml)

	Active substance	333.3 mg
	Formoterol fumarate	333.3 mg
15	Benzalkonium chloride	10.0 mg
	EDTA	50.0 mg
	HCl (1n)	ad pH 3.4

This solution may be prepared in the usual manner.

20

F) Powder for inhalation

	Active substance	6 µg
	Formoterol fumarate	6 µg
25	Lactose monohydrate	ad 25 mg


The powder for inhalation is produced in the usual way by mixing the individual ingredients together.

30 G) Powder for inhalation

	Active substance	10 µg
	Lactose monohydrate	ad 5 mg


The powder for inhalation is produced in the usual way by mixing the individual
35 ingredients together.

74402amendedclaims

New Claims**1) Compounds of general formula 1**

wherein


A denotes a double-bonded group selected from among

X⁻ denotes an anion with a single negative charge,R¹ and R² denote C₁-C₄-alkyl which may optionally be substituted by hydroxy or halogen;R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, C₁-C₄-alkyl, C₁-C₄-alkyloxy, hydroxy, CF₃, CN, NO₂ or halogen, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen,

optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates.

2) Compounds of general formula 1 according to claim 1, wherein

A denotes a double bonded group selected from among


X⁻ denotes an anion with a single negative charge selected from among chloride, bromide, methylsulphate, 4-toluenesulphonate and methanesulphonate,

R^1 and R^2 , which may be identical or different, denote a group selected from among methyl, ethyl, n-propyl and iso-propyl, which may optionally be substituted by hydroxy or fluorine;

R^3 , R^4 , R^5 , R^6 , R^7 and R^8 , which may be identical or different, denote hydrogen, methyl, ethyl, methoxy, ethoxy, hydroxy, fluorine, chlorine, bromine, CN, CF_3 or NO_2 , with the proviso that at least one of the groups R^3 , R^4 , R^5 , R^6 , R^7 and R^8 is not hydrogen, optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates.

3) Compounds of general formula 1 according to claim 1 or 2, wherein

A denotes a double-bonded group selected from among




X - denotes bromide;

R^1 and R^2 , which may be identical or different, denote methyl or ethyl;

R^3 , R^4 , R^5 , R^6 , R^7 and R^8 , which may be identical or different, denote hydrogen, methyl, methoxy, fluorine, chlorine or bromine, with the proviso that at least one of the groups R^3 , R^4 , R^5 , R^6 , R^7 and R^8 is not hydrogen, optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates.

4) Compounds of general formula 1 according to claim 1, 2 or 3, wherein

A denotes a double-bonded group selected from among



X - denotes bromide;


R^1 and R^2 which may be identical or different denote methyl or ethyl;

R^3 , R^4 , R^5 , R^6 , R^7 and R^8 , which may be identical or different, denote hydrogen, fluorine, chlorine or bromine, with the proviso that at least one of the groups R^3 , R^4 , R^5 , R^6 , R^7 and R^8 is not hydrogen, optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates.

5) Compounds of general formula 1 according to claim 1, 2, 3 or 4,

wherein

A denotes a double-bonded group selected from among



X - denotes bromide;

R¹ and R² which may be identical or different denote methyl or ethyl;


R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen, fluorine, chlorine or bromine, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen,

optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates.

6) Compounds of general formula 1 according to one of claims 1 to 5,

wherein

A denotes a double-bonded group selected from among



X - denotes bromide;

R¹ and R² denote methyl;

R³, R⁴, R⁵, R⁶, R⁷ and R⁸, which may be identical or different, denote hydrogen or fluorine, with the proviso that at least one of the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is not hydrogen,

optionally in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates.

7) Use of a compound of general formula 1 according to one of claims 1 to 6 as a pharmaceutical composition.


8) Use of a compound of general formula 1 according to one of claims 1 to 6 for preparing a pharmaceutical composition for treating diseases in which anticholinergics may develop a therapeutic benefit.

9) Use of a compound of general formula 1 according to one of claims 1 to 6 for preparing a pharmaceutical composition for the treatment of asthma, COPD, vagally induced sinus bradycardia, heart rhythm disorders, spasms in the gastrointestinal tract, spasms in the urinary tract and menstrual disorders.


10) Pharmaceutical preparations containing as active substance one or more compounds of general formula 1 according to one of claims 1 to 6 or the physiologically acceptable salts thereof, optionally combined with conventional excipients and/or carriers.

11) Pharmaceutical preparations according to claim 10, characterised in that they contain, in addition to one or more of the compounds of formula 1, at least one further active substance which is selected from among the betamimetics, antiallergic agents, PAF-antagonists, leukotriene-antagonists and steroids.


12) Process for preparing a compound of general formula 1




wherein A, X⁻ and the groups R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ may have the meanings given in claims 1 to 6, characterised in that in a first step a compound of general formula 3



wherein the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ may have the meanings given in claims 1 to 6 and R denotes C₁-C₄-alkyl, is reacted with a compound of formula 2




wherein A and R¹ may have the meanings given in claims 1 to 6, to obtain a compound of formula 4




wherein A and the groups R¹, R³, R⁴, R⁵, R⁶, R⁷ and R⁸ may have the meanings given in claims 1 to 6, and this is subsequently quaternised by reacting with a compound R²-X, wherein R² and X may have the meanings given in claims 1 to 6, to obtain a compound of formula 1.

13) Intermediate products of general formula 4



wherein A and the groups R¹, R³, R⁴, R⁵, R⁶, R⁷ and R⁸ may have the meanings given in claims 1 to 6.

14) Intermediate products of general formula 3



wherein R denotes C₁-C₄-alkyl and the groups R³, R⁴, R⁵, R⁶, R⁷ and R⁸ may have the meanings given in claim 6.

