



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) **Date de dépôt PCT/PCT Filing Date:** 2022/12/07
(87) **Date publication PCT/PCT Publication Date:** 2023/06/15
(85) **Entrée phase nationale/National Entry:** 2024/05/30
(86) **N° demande PCT/PCT Application No.:** EP 2022/084829
(87) **N° publication PCT/PCT Publication No.:** 2023/104902
(30) **Priorité/Priority:** 2021/12/08 (EP21213084.3)

(51) **Cl.Int./Int.Cl.** **C07K 14/37** (2006.01),
C07K 14/39 (2006.01), **C12N 15/80** (2006.01),
C12N 9/90 (2006.01), **C12P 21/00** (2006.01)

(71) **Demandeurs/Applicants:**
LESAFFRE ET COMPAGNIE, FR;
EUROPEAN MOLECULAR BIOLOGY LABORATORY,
DE

(72) **Inventeurs/Inventors:**
DESFOUGERES, THOMAS, FR;
DULERMO, THIERRY, FR;
PIGNEDE, GEORGES, FR;
STEINMETZ, LARS, DE;
JOHANSSON, S. ANDREAS, DE

(74) **Agent:** GOWLING WLG (CANADA) LLP

(54) **Titre : PRODUCTION AMELIOREE DE PROTEINES SECRETEES DANS DES CELLULES DE LEVURE**

(54) **Title: IMPROVED PRODUCTION OF SECRETED PROTEINS IN YEAST CELLS**

(57) **Abrégé/Abstract:**

The present invention relates to a yeast cell producing at least one secreted protein of interest, wherein said cell comprises at least one additional fungal gene showing increased expression and/or overexpression, showing reduced expression and/or inactivation, wherein said gene improves the production of the at least one secreted protein of interest. The present invention further relates to respective methods for production and uses of the yeast cell.



Date Submitted: 2024/05/30

CA App. No.: 3239731

Abstract:

The present invention relates to a yeast cell producing at least one secreted protein of interest, wherein said cell comprises at least one additional fungal gene showing increased expression and/or overexpression, showing reduced expression and/or inactivation, wherein said gene improves the production of the at least one secreted protein of interest. The present invention further relates to respective methods for production and uses of the yeast cell.

IMPROVED PRODUCTION OF SECRETED PROTEINS IN YEAST CELLS

The present invention relates to a yeast or filamentous fungal cell producing at least one secreted protein of interest, wherein said cell comprises at least one additional fungal gene showing increased expression and/or overexpression, showing reduced expression and/or inactivation, wherein said gene improves the production of the at least one secreted protein of interest. The present invention further relates to respective methods for production and uses of the yeast or filamentous fungal cells.

Background of the invention

The production of recombinant enzymes is growing rapidly and is estimated to generate several tens of billions of dollars (Martinez et al., 2012). Almost 60% of the enzymes used in detergents, the food industry and biofuel alcohol are recombinant enzymes, i.e. produced by an organism other than that of origin of the protein (COWAN, 1996). The expression of enzymes in a heterologous host allows (i) the production of enzymes of interest from slow growing or even non-cultivable organisms, (ii) the much higher production of the enzyme of interest, (iii) the production of proteins from pathogenic or toxin-producing organisms, and (iv) the increase of the stability or activity of an enzyme by protein engineering (Falch, 1991; Demain and Vaishnav, 2009).

Many microorganisms, including filamentous fungi (*Aspergillus sp.*, *Trichoderma sp.*), yeasts (for example *Pichia pastoris*, *Saccharomyces cerevisiae*, *Yarrowia lipolytica*) or bacteria (for example *Escherichia coli*, *Bacillus sp.*), are used to produce recombinant proteins (Demain and Vaishnav, 2009).

The production of recombinant proteins is dependent on the expression cassette (promoters and terminators used, signal sequence, codon bias), on the cellular machinery involved in the synthesis and degradation of proteins, intracellular trafficking and/or secretion, but also the energy level and/or redox of the cell as well as the culture conditions and the availability of nutrients (Zahrl et al., 2019).

Compared to other organisms conventionally used to produce recombinant proteins, *S. cerevisiae* has the advantage of rapid growth, easy manipulation both at the genetic level and at the level of production in bioreactors, and having Generally Recognized As Safe (GRAS) status. The production of a heterologous target protein in yeast host cells is further advantageous in that it allows the target proteins to be folded and secreted through the cellular secretory machinery.

Yeast is already widely used for many industrial applications (breadmaking, production of drinking alcohol and biofuels, etc. Parapouli *et al.*, 2020) where it may be advantageous to have it produce heterologous enzymes. For example, in the field of biofuel alcohol, the commercialized yeast strains of *S. cerevisiae* secrete enzymatic activities allowing the degradation of industrial mashes containing starch derivatives. This allows bioethanol manufacturers to limit their intake of exogenous enzymes and reduce their production costs.

US 2011-0129872A1 relates to a method for producing a recombinant protein, comprising culturing a yeast transformed with a recombinant gene construct comprising a yeast promoter, a gene coding a signal sequence and a gene coding a target protein; and also with one or more genes coding folding accessory protein selected from the group consisting of *PDI1* (protein disulfide isomerase 1), *SEC23* (secretory 23), *TRX2* (thioredoxin 2) *AHA1* (activator of heat shock protein 90 ATPase), and *SCJ1* (*S. cerevisiae* DnaJ), followed by culturing the transformed yeast.

US 2013-0011875 relates to a method and the production of higher titers of recombinant protein in a modified yeast host cell, for example *Pichia pastoris*, wherein the modified yeast cell lacks vacuolar sorting activity or has decreased vacuolar sorting activity relative to an unmodified yeast host cell of the same species.

US 2014-0335622 discloses an expression vector for secreting a protein (Z) to be recovered or a fusion protein having the protein (Z) moiety therein; a method for producing a transformant using the expression vector; the transformant; and a method for producing a protein using the transformant. It is disclosed that co-expression of a foreign secretory protein with *PDI1* increases the secretory production amount.

US 2016-0186192 describes a method for producing a desired protein comprising: (a) providing a host cell comprising a first recombinant gene encoding a protein comprising the sequence of a first chaperone protein, a second recombinant gene encoding a protein comprising the sequence of a second chaperone protein and a third gene, such as a third recombinant gene, encoding a desired protein (such as a desired heterologous protein), wherein the first and second chaperones are different; and (b) culturing the host cell in a culture medium to obtain expression of the first, second and third genes.

US 2018-0022785 claims a method for producing a heterologous protein, said method comprising: culturing a *Saccharomyces cerevisiae* yeast host cell or a culture thereof to produce the heterologous protein, wherein said *Saccharomyces cerevisiae* yeast host cell comprises a modified Not4 protein, and wherein said heterologous protein is an albumin, or a variant, fragment and/or fusion thereof.

Eun Jung Thak et al. (in: Yeast synthetic biology for designed cell factories producing secretory recombinant proteins, FEMS Yeast Research, Volume 20, Issue 2, March 2020, foaa009, <https://doi.org/10.1093/femsyr/foaa009>) disclose that yeasts are prominent hosts for the production of recombinant proteins from industrial enzymes to therapeutic proteins. Particularly, the similarity of protein secretion pathways between these unicellular eukaryotic microorganisms and higher eukaryotic organisms has made them a preferential host to produce secretory recombinant proteins. However, there are several bottlenecks, in terms of quality and quantity, restricting their use as secretory recombinant protein production hosts. They discuss recent developments in synthetic biology approaches to constructing yeast cell factories endowed with enhanced capacities of protein folding and secretion as well as designed targeted post-translational modification process functions, and focus on the new genetic tools for optimizing secretory protein expression, such as codon-optimized synthetic genes, combinatory synthetic signal peptides and copy number-controllable integration systems, and the advanced cellular engineering strategies, including endoplasmic reticulum and protein trafficking pathway engineering, synthetic glycosylation, and cell wall engineering, for improving the quality and yield of secretory recombinant proteins.

Zihe Liu, et al. (in: Improved Production of a Heterologous Amylase in *Saccharomyces cerevisiae* by Inverse Metabolic Engineering, Applied and Environmental Microbiology Aug 2014, 80 (17) 5542-5550; DOI: 10.1128/AEM.00712-14) disclose that the increasing demand

for industrial enzymes and biopharmaceutical proteins relies on robust production hosts with high protein yield and productivity. Being one of the best-studied model organisms and capable of performing posttranslational modifications, the yeast *Saccharomyces cerevisiae* is widely used as a cell factory for recombinant protein production. However, many recombinant proteins are produced at only 1% (or less) of the theoretical capacity due to the complexity of the secretory pathway, which has not been fully exploited. They applied the concept of inverse metabolic engineering to identify novel targets for improving protein secretion. Screening that combined UV-random mutagenesis and selection for growth on starch was performed to find mutant strains producing heterologous amylase 5-fold above the level produced by the reference strain. Genomic mutations that could be associated with higher amylase secretion were identified through whole-genome sequencing. Several single-point mutations, including an S196I point mutation in the VTA1 gene coding for a protein involved in vacuolar sorting, were evaluated by introducing these to the starting strain. By applying this modification alone, the amylase secretion could be improved by 35%. As a complement to the identification of genomic variants, transcriptome analysis was also performed in order to understand on a global level the transcriptional changes associated with the improved amylase production caused by UV mutagenesis.

Huang, M., et al. (in: Efficient protein production by yeast requires global tuning of metabolism. Nat Commun 8, 1131 (2017). <https://doi.org/10.1038/s41467-017-00999-2>) describe that the biotech industry relies on cell factories for production of pharmaceutical proteins, of which several are among the top-selling medicines. There is, therefore, considerable interest in improving the efficiency of protein production by cell factories. Protein secretion involves numerous intracellular processes with many underlying mechanisms still remaining unclear. They used RNA-seq to study the genome-wide transcriptional response to protein secretion in mutant yeast strains, and find that many cellular processes have to be attuned to support efficient protein secretion. In particular, altered energy metabolism resulting in reduced respiration and increased fermentation, as well as balancing of amino-acid biosynthesis and reduced thiamine biosynthesis seem to be particularly important. They confirmed their findings by inverse engineering and physiological characterization and show that by tuning metabolism cells are able to efficiently secrete recombinant proteins.

Huang M, et al. (In: Microfluidic screening and whole-genome sequencing identifies mutations associated with improved protein secretion by yeast. Proc Natl Acad Sci U S A. 2015 Aug

25;112(34):E4689-96. doi: 10.1073/pnas.1506460112. Epub 2015 Aug 10. PMID: 26261321; PMCID: PMC4553813) disclose that there is an increasing demand for biotech-based production of recombinant proteins for use as pharmaceuticals in the food and feed industry and in industrial applications, that the yeast *Saccharomyces cerevisiae* is among preferred cell factories for recombinant protein production, and there is increasing interest in improving its protein secretion capacity. Due to the complexity of the secretory machinery in eukaryotic cells, it is said to be difficult to apply rational engineering for construction of improved strains. They used high-throughput microfluidics for the screening of yeast libraries, generated by UV mutagenesis. Several screening and sorting rounds resulted in the selection of eight yeast clones with significantly improved secretion of recombinant α -amylase. Efficient secretion was genetically stable in the selected clones. They performed whole-genome sequencing of the eight clones and identified 330 mutations in total. Gene ontology analysis of mutated genes revealed many biological processes, including some that had not been identified before in the context of protein secretion. Mutated genes identified are disclosed to be potentially used for reverse metabolic engineering, with the objective to construct efficient cell factories for protein secretion. The combined use of microfluidics screening and whole-genome sequencing to map the mutations associated with the improved phenotype can easily be adapted for other products and cell types to identify novel engineering targets, and this approach could broadly facilitate design of novel cell factories.

Bao et al. (in: Moderate Expression of SEC16 Increases Protein Secretion by *Saccharomyces cerevisiae*, APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 83, no. 14, 15 July 2017) discloses that a moderate overexpression of the gene SEC16 increases protein secretion by *S. cerevisiae*. SEC16 is involved in protein translocation from the endoplasmic reticulum to the Golgi apparatus. The data also show that a high-level expression of SEC76 could be harmful for the cell due to higher accumulation of reactive oxygen species (ROS) and thus for recombinant protein production. Qi et al (in: Different Routes of Protein Folding Contribute to Improved Protein Production in *Saccharomyces cerevisiae*, mBio, 10 November 2020 (2020-11-10), xP055932697, Retrieved from the Internet: URL:<https://doi.org/10.1128/mBio.02743-20>) discloses that overexpression of Cwh41p improves protein production as seen by an increased α -amylase productivity.

WO2019027364 discloses recombinant *S. cerevisiae* allowing increased production of secreted proteins. It is suggested to overexpress PDI1 and Sec3, and/or downregulate the expression of YPS7, and VSP27.

WO200607511 discloses the use of chaperones to improve the production of a desired protein (secreted). One chaperone used is CCT3. JP2009240185 discloses the promotion of protein production by disrupting for example the VHS2 gene or the VSP27. WO094/08024 discloses recombinant yeast and filamentous fungi transformed with SSO genes, showing increased capacity to produce secreted foreign or endogenous proteins.

Finally, Huang M, et al. (in: Engineering the protein secretory pathway of *Saccharomyces cerevisiae* enables improved protein production. Proc Natl Acad Sci U S A. 2018 Nov 20;115(47):E11025-E11032. doi: 10.1073/pnas.1809921115. Epub 2018 Nov 5. PMID: 30397111; PMCID: PMC6255153) describe that baker's yeast *Saccharomyces cerevisiae* is one of the most important and widely used cell factories for recombinant protein production. Many strategies have been applied to engineer this yeast for improving its protein production capacity, but productivity is still relatively low, and with increasing market demand, it is important to identify new gene targets, especially targets that have synergistic effects with previously identified targets. Despite improved protein production, previous studies rarely focused on processes associated with intracellular protein retention. They identified genetic modifications involved in the secretory and trafficking pathways, the histone deacetylase complex, and carbohydrate metabolic processes as targets for improving protein secretion in yeast. Especially modifications of endosome-to-Golgi trafficking was found to effectively reduce protein retention besides increasing protein secretion. Through combinatorial genetic manipulations of several of the newly identified gene targets, they enhanced the protein production capacity of yeast by more than fivefold, and the best engineered strains could produce 2.5 g/L of a fungal α -amylase with less than 10% of the recombinant protein retained within the cells, using fed-batch cultivation.

Cryptic unstable transcripts (CUTs) are a subset of non-coding RNAs (ncRNAs) that are produced from intergenic and intragenic regions. Additionally, stable uncharacterized transcripts, or SUTs, have also been detected in cells and bear many similarities to CUTs but are not degraded through the same pathways.

Genetic engineering strategies to overcome bottlenecks in the yeast protein secretion pathway have to consider that protein secretion in yeast involves multiple complex steps, such as protein translocation, folding, post-translational modification and vesicle trafficking between several membrane organelles and plasma membranes. The secretion of proteins synthesized inside cells can be hampered by low secretion efficiency, abnormal post-translational modifications, retention within the secretion pathway or the cell wall space as a cell-associated form. The development of engineering strategies targeted to each step of the secretion pathway in a modular fashion is required in order to design cell factories producing secretory recombinant proteins. Today, despite its obvious qualities, *S. cerevisiae* remains relatively limited in its ability to secrete proteins compared to organisms such as filamentous fungi or *P. pastoris* (Demain and Vaishnav, 2009). It is therefore an object of the present invention to provide new factors to improve recombinant protein production and secretion in yeast. Other objects and advantages will become apparent to the person of skill when studying the present description of the present invention.

In a first aspect of the present invention, the above object is solved in accordance with the claims, preferably by providing a cell of *Saccharomyces cerevisiae*, producing at least one secreted protein of interest, wherein said cell comprises at least one fungal gene selected from the group consisting of ENO2, NMA2, PRY2, SUT074, TFG2, AVT2, TRM10, BNA7, and TOM22, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, MNT2, TPO2, ATG33, THR4, INP51, CUT901, YDR262W, MRP10, NDC1, and CMC1, wherein said at least one fungal gene shows reduced expression and/or inactivation, and optionally further comprising the fungal gene *HDA2* and/or *PDII*, showing an increased expression and/or overexpression.

Preferred is the yeast cell according to the present invention, wherein said cell comprises at least one fungal gene selected from the groups consisting of ENO2, NMA2, PRY2, SUT074, and TFG2, or AVT2, TRM10, PRY2, SUT074, BNA7, and TOM22, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the groups consisting of TLG2, CUT901, ATG33, THR4, YDR262W, and CMC1, or MRP10, TLG2, CUT901, ATG33, THR4, YDR262W, CMC1, MNT2, TPO2, and NDC1, preferably MNT2 and TPO2, wherein said at least one fungal gene shows reduced expression and/or inactivation, and optionally further

comprising the fungal genes *HDA2* and/or *PDH1*, showing an increased expression and/or overexpression, and/or *INP51* showing an reduced expression and/or inactivation.

The above object is further solved according to the present invention by providing a yeast or filamentous fungal cell producing at least one secreted protein of interest, wherein said cell comprises at least one fungal gene selected from the group consisting of MIC19, TOM22, NKP1, DML1, CUT859, GAL80, APM3, COQ10, BLM10, MDH1, VHS2, ASA1, TRP4, YPS7, CUT824, YOR318C, PRM7, ERV46, FIT2, GPM3, CUT892, SRN2, SUT643, CUT461, THR4, GMH1, SOL1, NAB6, YPR148C, ALP1, CUT097, ATG33, YOR316C-A, SOG2, MCM6, SUT230, SUT419, TIF11, TAF5, PHO91, AIM32, ENO2, UBA2, PUS5, ERG1, SUT311, KSS1, MRP10, CUT598, CUT188, YOR238W, EMW1, BNA7, SNR63, CCT3, PRY2, MAL11, KRS1 RAI1, SUT784, YPR148C, YEL1, CUT832, NMA2, VPS27, SUT428, PEX29, YLR446W, WBP1, AVT2, CUT854, TRM10, SLX9, YPL077C, PET122, TFG2, PUN1, CUT152, AIR2, CUT571, RPS26B, RRT6, RPC19, URA3, YGR045C, SMC3, PNG1, THI6, MEU1, CUT239, NSE4, SUT074, AAH1, RMD5, CUT607, ACS1, MNN1, ARH1, YHR140W, CET1, RRB1, YLR342W-A, RPS22B, CHS5, YIL165C, SUT093, LPX1, NCA3, EFG1, NBP35, CUT765, MSL1, SCD6, ATG42, CHS6, COQ2, RPO31, MKK1, HED1, PBP2, BET5, CUT678, YGR021W, SUT474, YGL159W, IRC21, VHR1, SPP1, PRP43, ZRT1, YLR041W, SUT711, COX18, CBP6, SUT575, CLG1, CUT213, QCR10, SNR3, MSS2, CUT505, YOS1, SUT073, UTP21, ACA1, CUT632, RIP1, HUL5, CUT727, RPL35B, CUT184, CUT420, YFL041W-A, SUT460, ATG10, MFA1, UGX2, TRK2, CUT704, SUT083, TRE1, RVS161, LEA1, EBP2, THI80, CTI6, CUT322, XPT1, MRPL35, YPL025C, SUT737, PGA2, ULP2, MRX16, EST1, NUP100, IES3, ATG39, YMR084W, SUT428, YPL119C-A, MIN8, CUT490, SUT287, KEL3, SUT678, SEC3, SOL4, SIS2, CUT915, RRP3, ESA1, PCL8, TRX3, YKL115C, EMP65, ZDS1, CUT167, SOD1, UBR2, LSP1, SNR81, RGD3, YTP1, SMY2, CUT449, FIN1, YKL106C-A, YAR019W-A, CCH1, AYR1, SUT573, VNX1, FOL3, SUT511, GIS4, CUT743, RPL24A, HMT1, SUT333, SPP2, SUT128, SMC6, PHR1, RPS15, CUT642, GYP7, tK(CUU)K, CUT896, SLM5, CUT586, CUT158, CUT276, CUT480, SUT751, SUT251, CUT643, and RRP12, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, CUT901, ATG33, THR4, YDR262W, CMC1, MRP17, YPT52, CUT312, MRPS5, RDR1, DAL7, RPL20A, YBR137W, RPL36B, YEL008C-A, RAX1, CUT729, INP51, UBP8, CUT258, YLR342W-A, SUT568, PEX7, MSD1, CUT136, TIM10, CUT361, snR51, TAL1, RIP1,

MRP10, SUT078, MRP51, GLO3, EHD3, HER1, NMA111, PBP4, MFB1, IKI3, NDL1, SUT433, YOR238W, SUT750, QDR2, RDI1, SUT014, CUT437, MSC6, SUT497, YCR051W, MRPL33, RPL14A, TRM7, RNH202, RTC5, SUT027, CDC5, SUT729, YOR131C, CUT665, GLG2, SUT268, SUT705, MED4, RCR2, EFB1, RXT2, KGD1, TUP1, RNH203, YDR338C, SED1, CUT522, HIS2, SUT145, MET17, APC4, NKP2, MKK2, NDC1, PET100, NIP7, VHT1, SUT685, BNI5, SNA3, EGH1, MRP4, POB3, PIB2, SUT317, YKL024C, YGL116W, YLR118C, YFR031C-A, YGL190C, YDL108W, YMR128W, YBR253W, YJR113C, YIL031W, YGR109C, YBR282W, YMR125W, YMR236W, YDR411C, YML029W, YDL033C, YPL050C, YHR171W, YDR352W, and NTO1, wherein said at least one fungal gene shows reduced expression and/or inactivation.

Preferably, said cell comprises at least one fungal gene selected from the group consisting of MIC19, TOM22, NKP1, DML1, CUT859, GAL80, APM3, COQ10, BLM10, MDH1, EMW1, BNA7, SNR63, CCT3, PRY2, MAL11, KRS1, RAI1, SUT784, YPR148C, YEL1, CUT832, NMA2, VPS27, SUT428, PEX29, YLR446W, and WBP1, preferably ENO2, NMA2, PRY2, SUT074, and TFG2, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, CUT901, ATG33, THR4, NDC1, PET100, NIP7, VHT1, and SUT685, preferably MNT2, and TPO2, wherein said at least one fungal gene shows reduced expression and/or inactivation..

More preferred is the yeast or filamentous fungal cell according to the present invention, wherein said genes or SUTs or CUTs are furthermore selected from the group of genes or SUTs or CUTs having a value of $\log_{FC}/FDR$ $\log_{FC}/FDR$ of more than 40, preferably of more than 200, more preferred of more than 300, and most preferred of more than 500, based on the values as determined herein.

More preferred is the yeast or filamentous fungal cell according to the present invention, further comprising a fungal gene selected from the group consisting of THR4, MRP10, RIP1, YLR342W-A, ATG33, and YOR238W, either showing an increased expression and/or overexpression or reduced expression and/or inactivation, depending on the experimental conditions, such as, without wanting to be bound by theory, for example, the impact of CRISPRa and CRISPRi on gene expression due to the position of the gRNA in the promoting region.

Even more preferred is the yeast or filamentous fungal cell according to the present invention, further comprising the fungal gene *HDA2* and/or *PDII*, showing an increased expression and/or overexpression.

Advantageously, the yeast or filamentous fungal cell according to the present invention produces the at least one secreted protein to about 20% or more about, or about 30% or more, or about 40% or more, preferably about 50% or more, more preferably to about 75% or more, when compared to a control yeast or filamentous fungal cell.

In a second aspect of the present invention, the above object is solved by a method for producing a secreted protein in a yeast or filamentous fungal cell, comprising the steps of i) providing a yeast or filamentous fungal cell producing at least one secreted protein of interest according to the present invention, ii) culturing said yeast or filamentous fungal cell in suitable culture medium, and iii) isolating said secreted protein from said culture medium. Preferred is the method according to the present invention, further comprising suitably inducing the increased expression and/or overexpression or reduced expression and/or inactivation of the at least one fungal gene.

Further preferred is the method according to the present invention, wherein about 30% or more, or about 40% or more, preferably about 50% or more, more preferably to about 75% or more of said at least one secreted protein is produced, when compared to the production of a control yeast or filamentous fungal cell.

In a third aspect of the present invention, the above object is solved by a method for producing a yeast or filamentous fungal cell producing at least one secreted protein of interest according to the present invention, comprising introducing into said cell producing at least one secreted protein of interest at least one fungal gene selected from the group consisting of MIC19, TOM22, NKP1, DML1, CUT859, GAL80, APM3, COQ10, BLM10, MDH1, VHS2, ASA1, TRP4, YPS7, CUT824, YOR318C, PRM7, ERV46, FIT2, GPM3, CUT892, SRN2, SUT643, CUT461, THR4, GMH1, SOL1, NAB6, YPR148C, ALP1, CUT097, ATG33, YOR316C-A, SOG2, MCM6, SUT230, SUT419, TIF11, TAF5, PHO91, AIM32, ENO2, UBA2, PUS5, ERG1, SUT311, KSS1, MRP10, CUT598, CUT188, YOR238W, EMW1, BNA7, SNR63, CCT3, PRY2, MAL11, KRS1 RAI1, SUT784, YPR148C, YEL1, CUT832, NMA2, VPS27,

SUT428, PEX29, YLR446W, WBP1, AVT2, CUT854, TRM10, SLX9, YPL077C, PET122, TFG2, PUN1, CUT152, AIR2, CUT571, RPS26B, RRT6, RPC19, URA3, YGR045C, SMC3, PNG1, THI6, MEU1, CUT239, NSE4, SUT074, AAH1, RMD5, CUT607, ACS1, MNN1, ARH1, YHR140W, CET1, RRB1, YLR342W-A, RPS22B, CHS5, YIL165C, SUT093, LPX1, NCA3, EFG1, NBP35, CUT765, MSL1, SCD6, ATG42, CHS6, COQ2, RPO31, MKK1, HED1, PBP2, BET5, CUT678, YGR021W, SUT474, YGL159W, IRC21, VHR1, SPP1, PRP43, ZRT1, YLR041W, SUT711, COX18, CBP6, SUT575, CLG1, CUT213, QCR10, SNR3, MSS2, CUT505, YOS1, SUT073, UTP21, ACA1, CUT632, RIP1, HUL5, CUT727, RPL35B, CUT184, CUT420, YFL041W-A, SUT460, ATG10, MFA1, UGX2, TRK2, CUT704, SUT083, TRE1, RVS161, LEA1, EBP2, THI80, CTI6, CUT322, XPT1, MRPL35, YPL025C, SUT737, PGA2, ULP2, MRX16, EST1, NUP100, IES3, ATG39, YMR084W, SUT428, YPL119C-A, MIN8, CUT490, SUT287, KEL3, SUT678, SEC3, SOL4, SIS2, CUT915, RRP3, ESA1, PCL8, TRX3, YKL115C, EMP65, ZDS1, CUT167, SOD1, UBR2, LSP1, SNR81, RGD3, YTP1, SMY2, CUT449, FIN1, YKL106C-A, YAR019W-A, CCH1, AYR1, SUT573, VNX1, FOL3, SUT511, GIS4, CUT743, RPL24A, HMT1, SUT333, SPP2, SUT128, SMC6, PHR1, RPS15, CUT642, GYP7, tK(CUU)K, CUT896, SLM5, CUT586, CUT158, CUT276, CUT480, SUT751, SUT251, CUT643, and RRP12, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, CUT901, ATG33, THR4, YDR262W, CMC1, MRP17, YPT52, CUT312, MRPS5, RDR1, DAL7, RPL20A, YBR137W, RPL36B, YEL008C-A, RAX1, INP51, CUT729, UBP8, CUT258, YLR342W-A, SUT568, PEX7, MSD1, CUT136, TIM10, CUT361, snR51, TAL1, RIP1, MRP10, SUT078, MRP51, GLO3, EHD3, HER1, NMA111, PBP4, MFB1, IKI3, NDL1, SUT433, YOR238W, SUT750, QDR2, RDI1, SUT014, CUT437, MSC6, SUT497, YCR051W, MRPL33, RPL14A, TRM7, RNH202, RTC5, SUT027, CDC5, SUT729, YOR131C, CUT665, GLG2, SUT268, SUT705, MED4, RCR2, EFB1, RXT2, KGD1, TUP1, RNH203, YDR338C, SED1, CUT522, HIS2, SUT145, MET17, APC4, NKP2, MKK2, NDC1, PET100, NIP7, VHT1, SUT685, BNI5, SNA3, EGH1, MRP4, POB3, PIB2, SUT317, YKL024C, YGL116W, YLR118C, YFR031C-A, YGL190C, YDL108W, YMR128W, YBR253W, YJR113C, YIL031W, YGR109C, YBR282W, YMR125W, YMR236W, YDR411C, YML029W, YDL033C, YPL050C, YHR171W, YDR352W, and NTO1, wherein said at least one fungal gene shows reduced expression and/or inactivation..

Preferred is a method of the present invention for producing a yeast or filamentous fungal cell producing at least one secreted protein of interest according to the present invention, comprising introducing into said cell producing at least one secreted protein of interest at least one fungal gene selected from the group consisting of MIC19, TOM22, NKP1, DML1, CUT859, GAL80, APM3, COQ10, BLM10, MDH1, EMW1, BNA7, SNR63, CCT3, PRY2, MAL11, KRS1, RAI1, SUT784, YPR148C, YEL1, CUT832, NMA2, VPS27, SUT428, PEX29, YLR446W, and WBP1, preferably ENO2, NMA2, PRY2, SUT074, and TFG2, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, CUT901, ATG33, THR4, NDC1, PET100, NIP7, VHT1, and SUT685, preferably MNT2, and TPO2, wherein said at least one fungal gene shows reduced expression and/or inactivation.

Furthermore, the method according to the invention may include further introducing into said cell a fungal gene selected from the group consisting of THR4, MRP10, RIP1, YLR342W-A, ATG33, and YOR238W, either showing an increased expression and/or overexpression or reduced expression and/or inactivation, depending on the experimental conditions. Furthermore, the method may include further introducing into said cell the fungal gene *HDA2* and/or *PDH1*, showing an increased expression and/or overexpression.

In a fourth aspect of the present invention, the above object is solved by the use of a yeast or filamentous fungal cell according to the present invention for producing at least one secreted protein of interest.

As mentioned above, the analysis of UV *S. cerevisiae* mutants expressing an α -amylase has revealed improved strains for secretion (Huang et al., 2015; Huang et al., 2018). Coupling microfluidics with a phenotypic screening using a starch complexed with BODIPY (which becomes fluorescent when it is released), the authors had selected the mutants secreting the most enzyme into the extracellular medium. The sequencing of eight hypersecretory clones (x1.5 to x6) revealed 330 mutations potentially involved in improving α -amylase production and secretion (Huang et al., 2015). A more in-depth analysis led to the identification of – amongst others as disclosed herein – a role of the known *PDH1* gene in the production and secretion of α -amylase in *S. cerevisiae*.

The purpose of the present invention was to discover new factors and genes involved in protein secretion in order to improve protein production and secretion, as exemplified in the industrial Ethanol Red ® strain of *S. cerevisiae*.

As mentioned above, in the first aspect of the present invention, a yeast or filamentous fungal cell is provided that produces at least one secreted protein of interest. In addition, the cell comprises at least one fungal gene selected from the group consisting of MIC19, TOM22, NKP1, DML1, CUT859, GAL80, APM3, COQ10, BLM10, MDH1, VHS2, ASA1, TRP4, YPS7, CUT824, YOR318C, PRM7, ERV46, FIT2, GPM3, CUT892, SRN2, SUT643, CUT461, THR4, GMH1, SOL1, NAB6, YPR148C, ALP1, CUT097, ATG33, YOR316C-A, SOG2, MCM6, SUT230, SUT419, TIF11, TAF5, PHO91, AIM32, ENO2, UBA2, PUS5, ERG1, SUT311, KSS1, MRP10, CUT598, CUT188, YOR238W, EMW1, BNA7, SNR63, CCT3, PRY2, MAL11, KRS1 RAI1, SUT784, YPR148C, YEL1, CUT832, NMA2, VPS27, SUT428, PEX29, YLR446W, WBP1, AVT2, CUT854, TRM10, SLX9, YPL077C, PET122, TFG2, PUN1, CUT152, AIR2, CUT571, RPS26B, RRT6, RPC19, URA3, YGR045C, SMC3, PNG1, THI6, MEU1, CUT239, NSE4, SUT074, AAH1, RMD5, CUT607, ACS1, MNN1, ARH1, YHR140W, CET1, RRB1, YLR342W-A, RPS22B, CHS5, YIL165C, SUT093, LPX1, NCA3, EFG1, NBP35, CUT765, MSL1, SCD6, ATG42, CHS6, COQ2, RPO31, MKK1, HED1, PBP2, BET5, CUT678, YGR021W, SUT474, YGL159W, IRC21, VHR1, SPP1, PRP43, ZRT1, YLR041W, SUT711, COX18, CBP6, SUT575, CLG1, CUT213, QCR10, SNR3, MSS2, CUT505, YOS1, SUT073, UTP21, ACA1, CUT632, RIP1, HUL5, CUT727, RPL35B, CUT184, CUT420, YFL041W-A, SUT460, ATG10, MFA1, UGX2, TRK2, CUT704, SUT083, TRE1, RVS161, LEA1, EBP2, THI80, CTI6, CUT322, XPT1, MRPL35, YPL025C, SUT737, PGA2, ULP2, MRX16, EST1, NUP100, IES3, ATG39, YMR084W, SUT428, YPL119C-A, MIN8, CUT490, SUT287, KEL3, SUT678, SEC3, SOL4, SIS2, CUT915, RRP3, ESA1, PCL8, TRX3, YKL115C, EMP65, ZDS1, CUT167, SOD1, UBR2, LSP1, SNR81, RGD3, YTP1, SMY2, CUT449, FIN1, YKL106C-A, YAR019W-A, CCH1, AYR1, SUT573, VN1, FOL3, SUT511, GIS4, CUT743, RPL24A, HMT1, SUT333, SPP2, SUT128, SMC6, PHR1, RPS15, CUT642, GYP7, tK(CUU)K, CUT896, SLM5, CUT586, CUT158, CUT276, CUT480, SUT751, SUT251, CUT643, and RRP12, wherein these at least one fungal gene shows increased expression and/or overexpression.

In the context of the present invention, the terms “increased expression” or “overexpression” indicate that the amount of protein as produced by the cell is higher when compared to the

expression in a control cell showing normal, unaltered or baseline expression. The change in expression can be achieved in any suitable way, and examples include mutated promoters, cloning of the gene under the control of a heterologous “strong” promoter, either inducible or constitutive, codon optimization, and mutations that stabilize the structure of the protein, and the like. In the context of the present invention, a preferred example of how to detect “increased expression” or “overexpression” is a change in logFC (log fold change, see the tables below), more preferably a statistically relevant change (FDR) in the logFC. Examples are a value of logFC/FDR of more than 40, preferably of more than 200, more preferred of more than 300, and most preferred of more than 500, based on the values as determined herein.

Alternatively or in addition, the cell comprises at least one fungal gene selected from the group consisting of TLG2, CUT901, ATG33, THR4, YDR262W, CMC1, MRP17, YPT52, CUT312, MRPS5, RDR1, DAL7, RPL20A, YBR137W, RPL36B, YEL008C-A, RAX1, INP51, CUT729, UBP8, CUT258, YLR342W-A, SUT568, PEX7, MSD1, CUT136, TIM10, CUT361, snR51, TAL1, RIP1, MRP10, SUT078, MRP51, GLO3, EHD3, HER1, NMA111, PBP4, MFB1, IKI3, NDL1, SUT433, YOR238W, SUT750, QDR2, RDI1, SUT014, CUT437, MSC6, SUT497, YCR051W, MRPL33, RPL14A, TRM7, RNH202, RTC5, SUT027, CDC5, SUT729, YOR131C, CUT665, GLG2, SUT268, SUT705, MED4, RCR2, EFB1, RXT2, KGD1, TUP1, RNH203, YDR338C, SED1, CUT522, HIS2, SUT145, MET17, APC4, NKP2, MKK2, NDC1, PET100, NIP7, VHT1, SUT685, BNI5, SNA3, EGH1, MRP4, POB3, PIB2, SUT317, YKL024C, YGL116W, YLR118C, YFR031C-A, YGL190C, YDL108W, YMR128W, YBR253W, YJR113C, YIL031W, YGR109C, YBR282W, YMR125W, YMR236W, YDR411C, YML029W, YDL033C, YPL050C, YHR171W, YDR352W, and NTO1, wherein said at least one fungal gene shows reduced expression and/or inactivation, wherein said at least one fungal gene shows reduced expression and/or inactivation. In the context of the present invention, the terms “reduced expression” or “inactivation” indicate that the amount of protein as produced by the cell is lower when compared to the expression in a control cell showing normal, unaltered or baseline expression. The change in expression can be achieved in any suitable way, and examples include mutated promoters, cloning of the gene under the control of a heterologous “weak” promoter, either inducible or constitutive, codon changes, and mutations that de-stabilize the structure of the protein, and the like.

Systematic studies of the effects on protein secretion from gene perturbations are challenging, primarily due to the size of the readout, yeast encodes around 6300 genes, in addition to other

genetic elements, including long non-coding RNAs, such as cryptic untranslated transcripts (CUTs) and stable uncharacterized transcripts (SUTs) that are not transcribed into proteins, but instead affect and modulate gene expression in the nucleus or the cytosol.

Preferably, said yeast or filamentous fungal cell as provided comprises at least one fungal gene selected from the group consisting of MIC19, TOM22, NKP1, DML1, CUT859, GAL80, APM3, COQ10, BLM10, MDH1, EMW1, BNA7, SNR63, CCT3, PRY2, MAL11, KRS1, RAI1, SUT784, YPR148C, YEL1, CUT832, NMA2, VPS27, SUT428, PEX29, YLR446W, and WBP1, preferably ENO2, NMA2, PRY2, SUT074, and TFG2, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, CUT901, ATG33, THR4, NDC1, PET100, NIP7, VHT1, and SUT685, preferably MNT2, and TPO2, wherein said at least one fungal gene shows reduced expression and/or inactivation.

More preferred is the yeast or filamentous fungal cell according to the present invention, wherein said genes or SUTs or CUTs are furthermore selected from the group of genes or SUTs or CUTs having a value of logFC/FDR of more than 40, preferably of more than 200, more preferred of more than 300, and most preferred of more than 500, based on the values as determined herein.

More preferred is the yeast or filamentous fungal cell according to the present invention, further comprising a fungal gene selected from the group consisting of THR4, MRP10, RIP1, YLR342W-A, ATG33, and YOR238W, either showing an increased expression and/or overexpression or reduced expression and/or inactivation, depending on the experimental conditions.

Even more preferred is the yeast or filamentous fungal cell according to the present invention, further comprising the fungal gene *HDA2* and/or *PDII*, showing an increased expression and/or overexpression.

It is expected that a combination of genes as mentioned herein can lead to an even further increased production of the protein of interest, even having synergistic effects. Examples for these combinations are all of *TLG2*, *YDR262W*, and *TRM10*, optionally further comprising

HDA2 and/or *PDII*. Other examples are *ATG33* and *MRP10*, *NDC1* and *TRM10*, or *PRY2*, and *TOM22*, again each pair optionally further comprising *HDA2* and/or *PDII*.

Most preferred are either *AVT2*, *PRY2*, *SUT074*, *BNA7*, *TOM22* or *TRM10*. The overexpression of *AVT2*, *TRM10*, *PRY2*, *SUT074*, *BNA7*, or *TOM22*, and the inactivation of *INP51* is further preferred. Further examples are *TLG2*, *CUT901*, *ATG33*, *THR4*, *YDR262W*, and *CMC1*, optionally further comprising *HDA2* and/or *PDII*. Also preferred is *ENO2*, *NMA2*, *PRY2*, *SUT074*, and *TFG2* (increased expression and/or overexpression), *MNT2*, and *TPO2* (reduced expression and/or inactivation), optionally further comprising *HDA2* and/or *PDII*.

The fungal gene(s) and/or SUTs or CUTs as used are preferably derived from *S. cerevisiae*, or a related yeast. The fungal gene(s) and/or SUTs or CUTs and their reference numbers are according to the Saccharomyces Genome Database (*SGD*) (<https://www.yeastgenome.org/>), as of November 15, 2021. Related genes that may be used as well encode for proteins sharing the same biological effect (increased secretion) in the yeast or filamentous fungal cell with the genes as above, and/or have an amino acid identity of about 80% or more, preferably about 90% or more, more preferably about 95% or more with the polypeptide sequence as encoded by a genes as above.

Advantageously, preferably the yeast or filamentous fungal cell according to the present invention produces the at least one secreted protein to about 30% or more or 40% or more, preferably about 50% or more, more preferably to about 75% or more, when compared to a control yeast or filamentous fungal cell, preferably one that does not contain a gene as mentioned above leading to increased secretion of the protein of interest.

As the protein of interest, any protein can be chosen that can be suitably produced by the yeast or filamentous fungal cell according to the present invention, e.g. expressed, folded, glycosylated and/or secreted. The gene of the protein of interest can be codon optimized, and preferably show an increased expression and/or overexpression, as explained above for the fungal gene according to the present invention. Examples of preferred proteins of interest are human serum albumin (HSA), amylase, human insulin, and components of hepatitis vaccines, human papillomavirus (HPV) vaccines, interferon(s), or epidermal growth factor (hEGF), and proteins used in food production, such as cellulase, glucoamylase, xylanase, and the like.

In order to identify new genes involved in the production and secretion of recombinant and heterologous proteins in yeast or filamentous fungal cells, such as *S. cerevisiae*, the inventors have developed CRISPRi and CRISPRa libraries allowing the overexpression or the repression of all genes as well as previously identified Stable Unannotated Transcripts (SUT's) and (Cryptic Unstable Transcripts CUT's) of this yeast (see Xu, Z. *et al.* Bidirectional promoters generate pervasive transcription in yeast. *Nature* **457**, 1033–1037 (2009)). These libraries utilize an inactivated Cas9 (dCas9) able to bind DNA at the CRISPR site but unable to cleave the DNA molecule, fused to a transcriptional activation (CRISPRa) (e.g. the VP64-p65-Rta (VPR) tripartite activation domain described in Chavez, A. *et al.* Highly efficient Cas9-mediated transcriptional programming. *Nat Methods* **12**, 326–328 (2015)) or repression domain (CRISPRi) (Dominguez *et al.*, 2015).

The industrial Ethanol Red® (ER) yeast strain overexpressing an α -amylase (*Amy6* from *A. niger*) was used as a model for the present invention (Lesaffre, Marcq-en-Barœul, France). A 40,890 gRNA library targeting the promoters of 7,247 yeast genes, SUT's and CUT's at an average of 5.8 positions per gene, SUT or CUT was developed and cloned into replicative vectors allowing their expression as well as the expression of dCas9-VP64-p65-Rta (CRISPRa) or dCas9-Mxi1 (CRISPRi). The ER + α -amylase strain was then transformed using the CRISPRa and CRISPRi libraries, and the cell population as obtained was screened by microfluidics on the basis of its capacity to degrade a starch substrate labelled with BODIPY FL dye which fluoresces in green when the starch is degraded by α -amylase (e.g. EnzChek® Ultra Amylase Assay Kit: <https://www.thermofisher.com/order/catalog/product/E33651#/E33651>).

Clones presenting high fluorescence were sorted, and gRNA regions from replicative vectors were analyzed by Illumina sequencing. Data analysis revealed that 320 activated or repressed genes favor α -amylase secretion. These genes were manually selected further, and the genes MIC19, TOM22, NKP1, DML1, CUT859, GAL80, APM3, COQ10, BLM10, MDH1, VHS2, ASA1, TRP4, YPS7, CUT824, YOR318C, PRM7, ERV46, FIT2, GPM3, CUT892, SRN2, SUT643, CUT461, THR4, GMH1, SOL1, NAB6, YPR148C, ALP1, CUT097, ATG33, YOR316C-A, SOG2, MCM6, SUT230, SUT419, TIF11, TAF5, PHO91, AIM32, ENO2, UBA2, PUS5, ERG1, SUT311, KSS1, MRP10, CUT598, CUT188, YOR238W, EMW1, BNA7, SNR63, CCT3, PRY2, MAL11, KRS1, RAI1, SUT784, YPR148C, YEL1, CUT832, NMA2, VPS27, SUT428, PEX29, YLR446W, WBP1, AVT2, CUT854, TRM10, SLX9,

YPL077C, PET122, TFG2, PUN1, CUT152, AIR2, CUT571, RPS26B, RRT6, RPC19, URA3, YGR045C, SMC3, PNG1, THI6, MEU1, CUT239, NSE4, SUT074, AAH1, RMD5, CUT607, ACS1, MNN1, ARH1, YHR140W, CET1, RRB1, YLR342W-A, RPS22B, CHS5, YIL165C, SUT093, LPX1, NCA3, EFG1, NBP35, CUT765, MSL1, SCD6, ATG42, CHS6, COQ2, RPO31, MKK1, HED1, PBP2, BET5, CUT678, YGR021W, SUT474, YGL159W, IRC21, VHR1, SPP1, PRP43, ZRT1, YLR041W, SUT711, COX18, CBP6, SUT575, CLG1, CUT213, QCR10, SNR3, MSS2, CUT505, YOS1, SUT073, UTP21, ACA1, CUT632, RIP1, HUL5, CUT727, RPL35B, CUT184, CUT420, YFL041W-A, SUT460, ATG10, MFA1, UGX2, TRK2, CUT704, SUT083, TRE1, RVS161, LEA1, EBP2, THI80, CTI6, CUT322, XPT1, MRPL35, YPL025C, SUT737, PGA2, ULP2, MRX16, EST1, NUP100, IES3, ATG39, YMR084W, SUT428, YPL119C-A, MIN8, CUT490, SUT287, KEL3, SUT678, SEC3, SOL4, SIS2, CUT915, RRP3, ESA1, PCL8, TRX3, YKL115C, EMP65, ZDS1, CUT167, SOD1, UBR2, LSP1, SNR81, RGD3, YTP1, SMY2, CUT449, FIN1, YKL106C-A, YAR019W-A, CCH1, AYR1, SUT573, VNX1, FOL3, SUT511, GIS4, CUT743, RPL24A, HMT1, SUT333, SPP2, SUT128, SMC6, PHR1, RPS15, CUT642, GYP7, tK(CUU)K, CUT896, SLM5, CUT586, CUT158, CUT276, CUT480, SUT751, SUT251, CUT643, and RRP12, were overexpressed using common techniques (integration of overexpression cassette into the genome and/or overexpression through a replicative plasmid), and genes TLG2, CUT901, ATG33, THR4, YDR262W, CMC1, MRP17, YPT52, CUT312, MRPS5, RDR1, DAL7, RPL20A, YBR137W, RPL36B, YEL008C-A, RAX1, INP51, CUT729, UBP8, CUT258, YLR342W-A, SUT568, PEX7, MSD1, CUT136, TIM10, CUT361, snR51, TAL1, RIP1, MRP10, SUT078, MRP51, GLO3, EHD3, HER1, NMA111, PBP4, MFB1, IKI3, NDL1, SUT433, YOR238W, SUT750, QDR2, RDI1, SUT014, CUT437, MSC6, SUT497, YCR051W, MRPL33, RPL14A, TRM7, RNH202, RTC5, SUT027, CDC5, SUT729, YOR131C, CUT665, GLG2, SUT268, SUT705, MED4, RCR2, EFB1, RXT2, KGD1, TUP1, RNH203, YDR338C, SED1, CUT522, HIS2, SUT145, MET17, APC4, NKP2, MKK2, NDC1, PET100, NIP7, VHT1, SUT685, BNI5, SNA3, EGH1, MRP4, POB3, PIB2, SUT317, YKL024C, YGL116W, YLR118C, YFR031C-A, YGL190C, YDL108W, YMR128W, YBR253W, YJR113C, YIL031W, YGR109C, YBR282W, YMR125W, YMR236W, YDR411C, YML029W, YDL033C, YPL050C, YHR171W, YDR352W, and NTO1, were inactivated by gene deletion. Then, α -amylase activity was evaluated in the respective strains. The overexpression of *BNB7*, *SUT074*, *TOM22*, *TLG2*, *YDR262W*, *ALP1*, *ENO2*, *NMA2*, *PRY2*, and *INP51* were identified as preferred for the exemplary α -amylase secretion in the Ethanol Red® strain.

In the context of the present invention, any suitable cell of a yeast or filamentous fungus can be used for the production of the protein of interest according to the present invention. Preferably, said yeast or filamentous fungal cell is selected from the group consisting of *Aspergillus* spp., *Trichoderma* spp., *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* spp., *Pichia* spp., *Hansenula polymorpha*, *Fusarium* spp., *Neurospora* spp., and *Penicillium* spp., preferably *Saccharomyces cerevisiae*.

In the yeast or filamentous fungal cell according to the present invention, the at least one fungal gene showing increased expression and/or overexpression and/or showing reduced expression and/or inactivation is a native gene and/or is a recombinant gene, i.e. a modified gene of the yeast or filamentous fungal cell itself, or at least one gene that is recombinantly introduced and may be a heterologous gene, i.e. coming from a different strain or fungal species. Preferably, the recombinant gene is integrated into the genome as an expression cassette. Respective expression cassettes for fungal expression are known, and basically consist of a promoter, the fungal gene, and a terminator. Alternatively or in additionally, the gene can be extrachromosomally expressed, preferably using a replicative expression vector, such as a shuttle vector. Promoters used in yeast and fungal expression systems are usually either inducible or constitutive.

The folding and glycosylation of the secretory proteins in the endoplasmatic reticulum (ENDR) is assisted by numerous ENDR-resident proteins. The chaperones like Bip (GRP78), GRP94 or yeast Lhs1p help the secretory protein to fold by binding to exposed hydrophobic regions in the unfolded states and preventing unfavourable interactions (Blond-Elguindi et al., 1993, *Cell* 75:717-728). The chaperones are also important for the translocation of the proteins through the ENDR membrane. The proteins like protein disulphide isomerase and its homologs and prolyl-peptidyl cis-trans isomerase assist in formation of disulphide bridges and formation of the right conformation of the peptide chain adjacent to proline residues, respectively. A machinery including many protein components also resides in the ENDR for the addition of the N-linked core glycans to the secretory protein and for the initial trimming steps of the glycans.

Preferred is therefore the yeast or filamentous fungal cell according to the present invention, wherein the cell furthermore comprises at least one additional recombinant secretion promoting gene, for example a fungal gene for a chaperone, for a foldase and/or for a glycosylation-

promoting protein. Like the other genes as disclosed herein, these proteins may be controllably expressed, inducible, constitutive, and even overexpressed.

Therefore, preferred is the yeast or filamentous fungal cell according to the present invention, wherein the increased expression and/or overexpression or reduced expression and/or inactivation of the at least one fungal gene or the at least one additional recombinant secretion promoting gene is constitutive or inducible.

Another important aspect of the present invention relates to a method for producing a secreted protein in a yeast or filamentous fungal cell, comprising the steps of i) providing a yeast or filamentous fungal cell producing at least one secreted protein of interest according to the present invention as above, ii) suitably culturing said yeast or filamentous fungal cell in suitable culture medium, and iii) isolating said secreted protein from said culture medium. Methods for isolating proteins from cultures are known by the person of skill.

Culturing methods for producing proteins in yeast or filamentous fungal cells are known by the person of skill, and can be readily adjusted to the present invention. Culturing can be continuous or in batches or fed-batches. Preferred is the method according to the present invention, further comprising suitably inducing the increased expression and/or overexpression or reduced expression and/or inactivation of the at least one fungal gene. Induction can be achieved based on the promotor(s) as used, e.g. by adding inducers, or switching conditions, e.g. temperature.

There are many examples of engineering of *S. cerevisiae* for improved protein production, including optimizing of fermentation process, selecting the expression vectors systems, choosing the signal sequence for extracellular targeting and engineering host strains for better folding and post-translational modification (Tohda H., Kumagai H., Takegawa, K, (2010) Engineering of protein secretion in yeast: strategies and impact on protein production. Appl Microbiol Biotechnol 86: 403–417).

Preferred is the method according to the present invention, wherein about 30% or more or 40% or more, preferably about 50% or more, more preferably to about 75% or more of said at least one secreted protein is produced, when compared to the production of a control yeast or filamentous fungal cell, preferably one that does not contain a gene as mentioned above leading to increased secretion of the protein of interest

Another important aspect of the present invention relates to a method for producing a yeast or filamentous fungal cell producing at least one secreted protein of interest, comprising introducing into said cell producing at least one secreted protein of interest at least one fungal gene selected from the group consisting of MIC19, TOM22, NKP1, DML1, CUT859, GAL80, APM3, COQ10, BLM10, MDH1, VHS2, ASA1, TRP4, YPS7, CUT824, YOR318C, PRM7, ERV46, FIT2, GPM3, CUT892, SRN2, SUT643, CUT461, THR4, GMH1, SOL1, NAB6, YPR148C, ALP1, CUT097, ATG33, YOR316C-A, SOG2, MCM6, SUT230, SUT419, TIF11, TAF5, PHO91, AIM32, ENO2, UBA2, PUS5, ERG1, SUT311, KSS1, MRP10, CUT598, CUT188, YOR238W, EMW1, BNA7, SNR63, CCT3, PRY2, MAL11, KRS1 RAI1, SUT784, YPR148C, YEL1, CUT832, NMA2, VPS27, SUT428, PEX29, YLR446W, WBP1, AVT2, CUT854, TRM10, SLX9, YPL077C, PET122, TFG2, PUN1, CUT152, AIR2, CUT571, RPS26B, RRT6, RPC19, URA3, YGR045C, SMC3, PNG1, THI6, MEU1, CUT239, NSE4, SUT074, AAH1, RMD5, CUT607, ACS1, MNN1, ARH1, YHR140W, CET1, RRB1, YLR342W-A, RPS22B, CHS5, YIL165C, SUT093, LPX1, NCA3, EFG1, NBP35, CUT765, MSL1, SCD6, ATG42, CHS6, COQ2, RPO31, MKK1, HED1, PBP2, BET5, CUT678, YGR021W, SUT474, YGL159W, IRC21, VHR1, SPP1, PRP43, ZRT1, YLR041W, SUT711, COX18, CBP6, SUT575, CLG1, CUT213, QCR10, SNR3, MSS2, CUT505, YOS1, SUT073, UTP21, ACA1, CUT632, RIP1, HUL5, CUT727, RPL35B, CUT184, CUT420, YFL041W-A, SUT460, ATG10, MFA1, UGX2, TRK2, CUT704, SUT083, TRE1, RVS161, LEA1, EBP2, THI80, CTI6, CUT322, XPT1, MRPL35, YPL025C, SUT737, PGA2, ULP2, MRX16, EST1, NUP100, IES3, ATG39, YMR084W, SUT428, YPL119C-A, MIN8, CUT490, SUT287, KEL3, SUT678, SEC3, SOL4, SIS2, CUT915, RRP3, ESA1, PCL8, TRX3, YKL115C, EMP65, ZDS1, CUT167, SOD1, UBR2, LSP1, SNR81, RGD3, YTP1, SMY2, CUT449, FIN1, YKL106C-A, YAR019W-A, CCH1, AYR1, SUT573, VNX1, FOL3, SUT511, GIS4, CUT743, RPL24A, HMT1, SUT333, SPP2, SUT128, SMC6, PHR1, RPS15, CUT642, GYP7, tK(CUU)K, CUT896, SLM5, CUT586, CUT158, CUT276, CUT480, SUT751, SUT251, CUT643, and RRP12, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, CUT901, ATG33, THR4, YDR262W, CMC1, MRP17, YPT52, CUT312, MRPS5, RDR1, DAL7, RPL20A, YBR137W, RPL36B, YEL008C-A, RAX1, INP51, CUT729, UBP8, CUT258, YLR342W-A, SUT568, PEX7, MSD1, CUT136, TIM10, CUT361, snR51, TAL1, RIP1, MRP10, SUT078, MRP51, GLO3, EHD3, HER1, NMA111, PBP4, MFB1, IKI3, NDL1, SUT433, YOR238W, SUT750, QDR2, RDI1, SUT014, CUT437,

MSC6, SUT497, YCR051W, MRPL33, RPL14A, TRM7, RNH202, RTC5, SUT027, CDC5, SUT729, YOR131C, CUT665, GLG2, SUT268, SUT705, MED4, RCR2, EFB1, RXT2, KGD1, TUP1, RNH203, YDR338C, SED1, CUT522, HIS2, SUT145, MET17, APC4, NKP2, MKK2, NDC1, PET100, NIP7, VHT1, SUT685, BNI5, SNA3, EGH1, MRP4, POB3, PIB2, SUT317, YKL024C, YGL116W, YLR118C, YFR031C-A, YGL190C, YDL108W, YMR128W, YBR253W, YJR113C, YIL031W, YGR109C, YBR282W, YMR125W, YMR236W, YDR411C, YML029W, YDL033C, YPL050C, YHR171W, YDR352W, and NTO1, wherein said at least one fungal gene shows reduced expression and/or inactivation. Preferably, said at least one fungal gene is integrated into the genome as an expression cassette and/or extrachromosomally expressed, preferably using a replicative expression vector.

Preferred is a method of the present invention for producing a yeast or filamentous fungal cell producing at least one secreted protein of interest according to the present invention, comprising introducing into said cell producing at least one secreted protein of interest at least one fungal gene selected from the group consisting of MIC19, TOM22, NKP1, DML1, CUT859, GAL80, APM3, COQ10, BLM10, MDH1, EMW1, BNA7, SNR63, CCT3, PRY2, MAL11, KRS1, RAI1, SUT784, YPR148C, YEL1, CUT832, NMA2, VPS27, SUT428, PEX29, YLR446W, and WBP1, preferably ENO2, NMA2, PRY2, SUT074, and TFG2, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, CUT901, ATG33, THR4, NDC1, PET100, NIP7, VHT1, and SUT685, preferably MNT2, and TPO2, wherein said at least one fungal gene shows reduced expression and/or inactivation.

Furthermore, the method according to the invention may include further introducing into said cell a fungal gene selected from the group consisting of THR4, MRP10, RIP1, YLR342W-A, ATG33, and YOR238W, either showing an increased expression and/or overexpression or reduced expression and/or inactivation, depending on the experimental conditions.

In a preferred embodiment according to the method according to the present invention, said method further comprises introducing into said cell the fungal gene *HDA2* and/or *PDII*, showing an increased expression and/or overexpression. Preferably, said at least one fungal gene is also integrated into the genome as an expression cassette and/or extrachromosomally expressed, preferably using a replicative expression vector

Finally, another important aspect of the present invention relates to the use of a yeast or filamentous fungal cell according to the present invention for producing at least one secreted protein of interest, preferably using a method according to the present invention.

In the context of the present invention, the inventors deploy genome-wide CRISPRi (repression, Smith, J. D. et al. Quantitative CRISPR interference screens in yeast identify chemical-genetic interactions and new rules for guide RNA design. *Genome Biol* **17**, 45 (2016)) and CRISPRa (activation, Chavez, A. et al. Highly efficient Cas9-mediated transcriptional programming. *Nat Methods* **12**, 326–328 (2015)) libraries to systematically probe the effects from perturbations of gene expression on the protein secretion machinery; by targeting the transcription of all identified genes, SUT's and CUTs in *S. cerevisiae* on a per gene basis. The application of CRISPR/Cas9 in combination with high throughput screening and next-generation sequencing (NGS) allowed the inventors to maintain a genome-wide scope with single gene precision. This is, to the inventor's knowledge, the first systematic attempt at interrogating the effects from gene activation and repression on the protein secretion machinery across all genes in yeast.

In summary, the present invention provides the following items.

Item 1. A yeast or filamentous fungal cell producing at least one secreted protein of interest, wherein said cell comprises at least one fungal gene selected from the group consisting of MIC19, TOM22, NKP1, DML1, CUT859, GAL80, APM3, COQ10, BLM10, MDH1, VHS2, ASA1, TRP4, YPS7, CUT824, YOR318C, PRM7, ERV46, FIT2, GPM3, CUT892, SRN2, SUT643, CUT461, THR4, GMH1, SOL1, NAB6, YPR148C, ALP1, CUT097, ATG33, YOR316C-A, SOG2, MCM6, SUT230, SUT419, TIF11, TAF5, PHO91, AIM32, ENO2, UBA2, PUS5, ERG1, SUT311, KSS1, MRP10, CUT598, CUT188, YOR238W, EMW1, BNA7, SNR63, CCT3, PRY2, MAL11, KRS1, RAI1, SUT784, YPR148C, YEL1, CUT832, NMA2, VPS27, SUT428, PEX29, YLR446W, WBP1, AVT2, CUT854, TRM10, SLX9, YPL077C, PET122, TFG2, PUN1, CUT152, AIR2, CUT571, RPS26B, RRT6, RPC19, URA3, YGR045C, SMC3, PNG1, THI6, MEU1, CUT239, NSE4, SUT074, AAH1, RMD5, CUT607, ACS1, MNN1, ARH1, YHR140W, CET1, RRB1, YLR342W-A, RPS22B, CHS5, YIL165C, SUT093, LPX1, NCA3, EFG1, NBP35, CUT765, MSL1, SCD6, ATG42, CHS6, COQ2, RPO31, MKK1, HED1, PBP2, BET5, CUT678, YGR021W, SUT474, YGL159W, IRC21, VHR1, SPP1, PRP43, ZRT1, YLR041W, SUT711, COX18, CBP6, SUT575, CLG1, CUT213, QCR10, SNR3, MSS2, CUT505, YOS1, SUT073, UTP21, ACA1, CUT632, RIP1, HUL5, CUT727, RPL35B, CUT184, CUT420, YFL041W-A, SUT460, ATG10, MFA1, UGX2,

TRK2, CUT704, SUT083, TRE1, RVS161, LEA1, EBP2, THI80, CTI6, CUT322, XPT1, MRPL35, YPL025C, SUT737, PGA2, ULP2, MRX16, EST1, NUP100, IES3, ATG39, YMR084W, SUT428, YPL119C-A, MIN8, CUT490, SUT287, KEL3, SUT678, SEC3, SOL4, SIS2, CUT915, RRP3, ESA1, PCL8, TRX3, YKL115C, EMP65, ZDS1, CUT167, SOD1, UBR2, LSP1, SNR81, RGD3, YTP1, SMY2, CUT449, FIN1, YKL106C-A, YAR019W-A, CCH1, AYR1, SUT573, VNX1, FOL3, SUT511, GIS4, CUT743, RPL24A, HMT1, SUT333, SPP2, SUT128, SMC6, PHR1, RPS15, CUT642, GYP7, tK(CUU)K, CUT896, SLM5, CUT586, CUT158, CUT276, CUT480, SUT751, SUT251, CUT643, and RRP12, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, CUT901, ATG33, THR4, YDR262W, CMC1, MRP17, YPT52, CUT312, MRPS5, RDR1, DAL7, RPL20A, YBR137W, RPL36B, YEL008C-A, RAX1, INP51, CUT729, UBP8, CUT258, YLR342W-A, SUT568, PEX7, MSD1, CUT136, TIM10, CUT361, snR51, TAL1, RIP1, MRP10, SUT078, MRP51, GLO3, EHD3, HER1, NMA111, PBP4, MFB1, IKI3, NDL1, SUT433, YOR238W, SUT750, QDR2, RDI1, SUT014, CUT437, MSC6, SUT497, YCR051W, MRPL33, RPL14A, TRM7, RNH202, RTC5, SUT027, CDC5, SUT729, YOR131C, CUT665, GLG2, SUT268, SUT705, MED4, RCR2, EFB1, RXT2, KGD1, TUP1, RNH203, YDR338C, SED1, CUT522, HIS2, SUT145, MET17, APC4, NKP2, MKK2, NDC1, PET100, NIP7, VHT1, SUT685, BNI5, SNA3, EGH1, MRP4, POB3, PIB2, SUT1317, YKL024C, YGL116W, YLR118C, YFR031C-A, YGL190C, YDL108W, YMR128W, YBR253W, YJR113C, YIL031W, YGR109C, YBR282W, YMR125W, YMR236W, YDR411C, YML029W, YDL033C, YPL050C, YHR171W, YDR352W, and NTO1, wherein said at least one fungal gene shows reduced expression and/or inactivation.

Item 2. The yeast or filamentous fungal cell according to Item 1, wherein said cell comprises at least one fungal gene selected from the group consisting of MIC19, TOM22, NKP1, DML1, CUT859, GAL80, APM3, COQ10, BLM10, MDH1, EMW1, BNA7, SNR63, CCT3, PRY2, MAL11, KRS1, RAI1, SUT784, YPR148C, YEL1, CUT832, NMA2, VPS27, SUT428, PEX29, YLR446W, and WBP1, preferably ENO2, NMA2, PRY2, SUT074, and TFG2, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, CUT901, ATG33, THR4, NDC1, PET100, NIP7, VHT1, and SUT685, preferably MNT2, and TPO2, wherein said at least one fungal gene shows reduced expression and/or inactivation.

Item 3. The yeast or filamentous fungal cell according to Item 1 or 2, wherein said genes or SUTs or CUTs are furthermore selected from the group of genes or SUTs or CUTs having a value of logFC/FDR of more than 40, preferably of more than 200, more preferred of more than 300, and most preferred of more than 500, based on the values as determined herein.

Item 4. The yeast or filamentous fungal cell according to any one of Items 1 to 3, further comprising a fungal gene selected from the group consisting of THR4, MRP10, RIP1, YLR342W-A, ATG33, and YOR238W, either showing an increased expression and/or overexpression or reduced expression and/or inactivation, depending on the experimental conditions.

Item 5. The yeast or filamentous fungal cell according to any one of Items 1 to 4, further comprising the fungal gene *HDA2* and/or *PDII*, showing an increased expression and/or overexpression.

Item 6. The yeast or filamentous fungal cell according to any one of Items 1 to 5, wherein said yeast or filamentous fungal cell is selected from the group consisting of *Aspergillus* spp., *Trichoderma* spp., *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* spp., *Pichia* spp., *Hansenula polymorpha*, *Fusarium* spp., *Neurospora* spp., and *Penicillium* spp., preferably *Saccharomyces cerevisiae*.

Item 7. The yeast or filamentous fungal cell according to any one of Items 1 to 6, wherein said at least one secreted protein of interest also shows an increased expression and/or overexpression.

Item 8. The yeast or filamentous fungal cell according to any one of Items 1 to 7, wherein said at least one fungal gene showing increased expression and/or overexpression and/or showing reduced expression and/or inactivation is a native gene and/or is a recombinant gene, wherein preferably said recombinant gene is integrated into the genome as an expression cassette and/or extrachromosomally expressed, preferably using a replicative expression vector.

Item 9. The yeast or filamentous fungal cell according to any one of Items 1 to 8, wherein the cell furthermore comprises at least one additional recombinant secretion promoting gene, for example a gene for a chaperone, for a foldase and/or for a glycosylation-promoting protein.

Item 10. The yeast or filamentous fungal cell according to any one of Items 1 to 9, wherein the increased expression and/or overexpression or reduced expression and/or inactivation of the at least one fungal gene or the at least one additional recombinant secretion promoting gene is constitutive or inducible.

Item 11. The yeast or filamentous fungal cell according to any one of Items 1 to 10, wherein the cell produces the at least one secreted protein to about 30% or more, or to about 40% or more, preferably about 50% or more, more preferably to about 75% or more, when compared to a control yeast or filamentous fungal cell.

Item 12. A method for producing a secreted protein in a yeast or filamentous fungal cell, comprising the steps of i) providing a yeast or filamentous fungal cell producing at least one secreted protein of interest according to any one of Items 1 to 11, ii) culturing said yeast or filamentous fungal cell in suitable culture medium, and iii) isolating said secreted protein from said culture medium.

Item 13. The method according to Item 12, further comprising suitably inducing the increased expression and/or overexpression or reduced expression and/or inactivation of the at least one fungal gene.

Item 14. The method according to Item 11 or 12, wherein about 30% or more, or about 40% or more, preferably about 50% or more, more preferably to about 75% or more of said at least one secreted protein is produced, when compared to the production of a control yeast or filamentous fungal cell.

Item 15. A method for producing a yeast or filamentous fungal cell producing at least one secreted protein of interest, comprising introducing into said cell producing at least one secreted protein of interest at least one fungal gene selected from the group consisting of MIC19, TOM22, NKP1, DML1, CUT859, GAL80, APM3, COQ10, BLM10, MDH1, EMW1, BNA7, SNR63, CCT3, PRY2, MAL11, KRS1, RAI1, SUT784, YPR148C, YEL1, CUT832, NMA2,

VPS27, SUT428, PEX29, YLR446W, and WBP1, preferably ENO2, NMA2, PRY2, SUT074, and TFG2, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, CUT901, ATG33, THR4, NDC1, PET100, NIP7, VHT1, and SUT685, preferably MNT2, and TPO2, wherein said at least one fungal gene shows reduced expression and/or inactivation.

Item 16. The method according to Item 15, further introducing into said cell a fungal gene selected from the group consisting of THR4, MRP10, RIP1, YLR342W-A, ATG33, and YOR238W, either showing an increased expression and/or overexpression or reduced expression and/or inactivation, depending on the experimental conditions.

Item 17. The method according to Item 15 or 16, further introducing into said cell the fungal gene *HDA2* and/or *PDII*, showing an increased expression and/or overexpression.

Item 18. The method according to any one of Items 15 to 17, wherein said at least one fungal gene is integrated into the genome as an expression cassette and/or extrachromosomally expressed, preferably using a replicative expression vector.

Item 19. Use of a yeast or filamentous fungal cell according to any one of Items 1 to 10 for producing at least one secreted protein of interest.

In summary, the present invention in particular provides the following items.

Item 20. A cell of *Saccharomyces cerevisiae*, producing at least one secreted protein of interest, wherein said cell comprises at least one fungal gene selected from the group consisting of ENO2, NMA2, PRY2, SUT074, TFG2, AVT2, TRM10, BNA7, and TOM22, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, MNT2, TPO2, ATG33, THR4, INP51, CUT901, YDR262W, MRP10, NDC1, and CMC1, wherein said at least one fungal gene shows reduced expression and/or inactivation, and optionally further comprising the fungal gene *HDA2* and/or *PDII*, showing an increased expression and/or overexpression.

Item 21. The yeast cell according to Item 20, wherein said cell comprises at least one fungal gene selected from the groups consisting of ENO2, NMA2, PRY2, SUT074, and TFG2, or AVT2, TRM10, PRY2, SUT074, BNA7, and TOM22, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the groups consisting of TLG2, CUT901, ATG33, THR4, YDR262W, and CMC1, or MRP10, TLG2, CUT901, ATG33, THR4, YDR262W, CMC1, MNT2, TPO2, and NDC1, preferably MNT2 and TPO2, wherein said at least one fungal gene shows reduced expression and/or inactivation, and optionally further comprising the fungal genes *HDA2* and/or *PDH1*, showing an increased expression and/or overexpression, and/or *INP51* showing an reduced expression and/or inactivation.

Item 23. The yeast cell according to Item 21 or 22, wherein said genes or SUTs or CUTs are furthermore selected from the group of genes or SUTs or CUTs having a value of logFC/FDR logFC/FDR of more than 40, preferably of more than 200, more preferred of more than 300, and most preferred of more than 500, based on the values as determined herein.

Item 24. The yeast cell according to any one of Items 21 to 23, wherein said yeast cell is from *Saccharomyces cerevisiae* strain ER.sec2.

Item 25. The yeast cell according to any one of Items 21 to 24, wherein said at least one secreted protein of interest also shows an increased expression and/or overexpression.

Item 26. The yeast cell according to any one of Items 21 to 25, wherein said at least one fungal gene showing increased expression and/or overexpression and/or showing reduced expression and/or inactivation is a native gene and/or is a recombinant gene, wherein preferably said recombinant gene is integrated into the genome as an expression cassette and/or extrachromosomally expressed, preferably using a replicative expression vector.

Item 27. The yeast cell according to any one of Items 21 to 26, wherein the cell furthermore comprises at least one additional recombinant secretion promoting gene, for example a gene for a chaperone, for a foldase and/or for a glycosylation-promoting protein.

Item 28. The yeast cell according to any one of Items 21 to 27, wherein the increased expression and/or overexpression or reduced expression and/or inactivation of the at least one fungal gene or the at least one additional recombinant secretion promoting gene is constitutive or inducible.

Item 29. The yeast cell according to any one of Items 21 to 28, wherein the cell produces the at least one secreted protein to about 30% or more, or about 40% or more, preferably about 50% or more, more preferably to about 75% or more, when compared to a control yeast or filamentous fungal cell.

Item 30. A method for producing a secreted protein in a yeast cell, comprising the steps of i) providing a cell of *Saccharomyces cerevisiae* producing at least one secreted protein of interest according to any one of Items 21 to 29, ii) culturing said yeast cell in suitable culture medium, and iii) isolating said secreted protein from said culture medium, and optionally further comprising suitably inducing the increased expression and/or overexpression or reduced expression and/or inactivation of the at least one fungal gene.

Item 31. The method according to Item 30, wherein preferably about 30% or more, or about 40% or more, preferably about 50% or more, more preferably to about 75% or more of said at least one secreted protein is produced, when compared to the production of a control yeast cell.

Item 32. A method for producing a yeast cell producing at least one secreted protein of interest, comprising introducing into said cell producing at least one secreted protein of interest at least one fungal gene selected from the group consisting of ENO2, NMA2, PRY2, SUT074, TFG2, AVT2, TRM10, BNA7, and TOM22, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, MNT2, TPO2, ATG33, THR4, INP51, CUT901, YDR262W, MRP10, NDC1, and CMC1, preferably MNT2, and TPO2, wherein said at least one fungal gene shows reduced expression and/or inactivation, and optionally further introducing into said cell a fungal gene selected from the group consisting of RIP1, YLR342W-A, and YOR238W, either showing an increased expression and/or overexpression or reduced expression and/or inactivation, depending on the experimental conditions, and/or optionally further introducing into said cell the fungal gene *HDA2* and/or *PDII*, showing an increased expression and/or overexpression.

Item 33. The method according to any one of Items 30 to 32, wherein said at least one fungal gene is integrated into the genome as an expression cassette and/or extrachromosomally expressed, preferably using a replicative expression vector.

Item 34. Use of a yeast cell according to any one of Items 21 to 29 for producing at least one secreted protein of interest.

The present invention will now be described further in the following examples with reference to the accompanying Figure, nevertheless, without being limited thereto. For the purposes of the present invention, all references as cited herein are incorporated by reference in their entireties.

Figure 1 shows the map of plasmid pLI410-062 as used in the methods according to the present invention.

Figure 2 A and B shows the results of the α -amylase secretion measurements relative to baseline for selected genes of the present invention as box plots in % control over time (4, 24, 48, and 120 hours). Genes are ALP1, BNA7, GMH1, SUT074, TFG2, ENO2, NMA2, PRY2, and TOM22. HAC1 is control.

Figure 3 A and B shows the results of the α -amylase secretion measurements per cell for selected genes of the present invention as box plots in % control over time (4, 24, 48, and 120 hours). Genes are ALP1, BNA7, GMH1, SUT074, TFG2, ENO2, NMA2, PRY2, and TOM22. HAC1 is control.

Figure 4 A and B shows the results of the α -amylase secretion measurements (total amylase) for selected genes of the present invention as box plots in % control over time (4, 24, and 48 hours). Genes are INP51, MNT2, TLG2, TPO2, and YDR262W. HAC1, HDA2 and ER.sec2 are controls.

Figure 5 A and B shows the results of the α -amylase secretion measurements per cell for selected genes of the present invention as box plots in % control over time (4, 24, and 48 hours). Genes are INP51, MNT2, TLG2, TPO2, and YDR262W. HAC1, HDA2 and ER.sec2 are controls.

Examples

Materials and Methods

Selection of guide-RNA and oligo design

Guide RNA covering all known genes, SUTs, CUTs (for simplicity referred to as genes from here on) in *S. cerevisiae* were selected using Azimuth (Listgarten, J. et al. Prediction of off-target activities for the end-to-end design of CRISPR guide RNAs. *Nat Biomed Eng* **2**, 38–47 (2018).) and chosen to be as evenly distributed as possible in 5 bins of 100 bp each from 400 bp upstream to 100 bp downstream of the predicted transcription start site (TSS). This resulted in a library of 40890 guides for an average of approximately six guides per feature. The potential for off-target effects was minimized by blasting the individual guide RNAs (gRNA) against each other guide and all potential gRNA binding sites (4.7 M in total) throughout the genome and removing any guide with less than three mismatches. Oligos were ordered from Agilent using a design that optimizes the number of guides per oligo, each 190 bp oligo contains four individual 20 bp guide-RNA sequences interspersed with spacer sequences containing double Type II-S recognition sites, enabling restriction digest and release using BspQI with subsequent removal of the recognition site.

Construction of yeast overexpression strains

For overexpression of target genes using genome integration, candidate genes were cloned into plasmid pLI410-062 between the *AscI* and *SbfI* restriction sites, which was then linearized by *NotI* enzyme, and transformed into yeast strain ER.sec2. The plasmid integrates into the yeast chromosome at the *BUD5* locus (Figure 1). For plasmid based overexpression of target genes, native candidate genes were cloned into plasmid p427-TEF between *SpeI* and *SalI* and transformed into yeast strain ER.sec2.

Construction of yeast deletion strains

Deletion strains were constructed by golden gate assembly of annealed oligos with gRNA sequences targeting the start and end position of the target gene, into sgRNA expression vector pWS082. The assembled plasmid and Cas9 expression vector pWS173 were linearized using *EcoRV* or *BsmBI* and co-transformed with annealed repair fragments, consisting of the joined 60 bp flanking regions of each target gene, which upon successful homology directed repair, resulted in the deletion of the target gene in ER.sec2.

The industrial Ethanol Red® (ER) yeast strain overexpressing an α -amylase (*Amy6* from *A. niger*) was used as a model for the present invention. The person of skill in the art will be able to adapt the principles of the present invention to other fungal/yeast strains as shall be used, and – if required – to select suitable genes from the lists as disclosed in order to achieve the changes in expression(s) as disclosed herein.

α -Amylase activity measurement

Preculture of YPD (Yeast extract Peptone Dextrose) was performed, either with 22h of culture on SD-2xSCAA, or 22h and 96 h of culture on YPD. SD-2xSCAA medium was prepared as described previously (Hackel et al. 2006; Tyo et al. 2012), and the composition of SD-2xSCAA was as follows: 10 g/L glucose, 6.7 g/L yeast nitrogen base without amino acids, 2 g/L, KH₂PO₄ (pH 6.0 by NaOH), and 1 g/L BSA, containing filter sterilized SCAA solution (190 mg/L arginine, 108 mg/L methionine, 52 mg/L tyrosine, 290 mg/L isoleucine, 440 mg/L lysine, 200 mg/L phenylalanine, 1,260 mg/L, glutamic acid, 400 mg/L aspartic acid, 380 mg/L valine, 220 mg/L threonine, 130 mg/L glycine, 400 mg/L leucine, 40 mg/L tryptophan, and 140 mg/L histidine) (see Liu et al., 2013 – Correlation of cell growth and heterologous protein production by *Saccharomyces cerevisiae*).

The initial OD_{600nm} was 0.1, and flasks of 250 ml + 50 ml of medium were used. Culture density was measured at OD_{600nm}.

For the assay, 100 μ L of supernatant + 900 μ L of acetate buffer 50mM pH5.5 were combined, and 10 μ L of sample were incubated for 5 min at 40°C in a PCR well plate.

Afterwards, 10 μ L of BPNPG7 substrate was added, followed by incubation for 10 min at 40°C. The reaction was stopped by adding 150 μ L of Trizma base 1%, followed by vortexing. The result was read at an OD of 400nm, which generally required a prior step of 10 or 20-fold dilution.

Calculation of α -amylase activity

The activity U was calculated as $U = (\Delta E_{400}/10) \times (0.17/0.01) \times (1/18.1) \times D$

E_{400} : Sample absorbance – blank absorbance, 10 : time of reaction, 0.17 : total volume of reaction, 0.01 : volume of sample, 18.1 : E_{mM} p-nitrophenol in Trizma base 1%, D : Dilution of sample. Normalization of α -amylase activity was performed with respective $OD_{600\text{nm}}$.

Results

Previous studies using microfluidics platforms, which screened for strains with an increased ability to secrete protein, using yeast cells treated with a mutagen, and encapsulated in a droplet with a suitable substrate, identified several strains that over-expressed α -amylase compared to the wild-type strain. These screens efficiently identified over-secretion strains by screening and sorting for increased protein secretion, but were to some degree hampered by the lack of a direct read-out of the affected genes, which necessitated whole-genome sequencing to identify the affected locus or loci.

The inventors utilized CRISPR with nuclease-null dCas9 to perturb a single gene per cell in a pooled format across the genome, coupled with microfluidic sorting of high fluorescence droplets using the same α -amylase assay described in the previous studies (Sjostrom, S. L. et al. High-throughput screening for industrial enzyme production hosts by droplet microfluidics. *Lab Chip* **14**, 806–813 (2013), Huang, M. et al. Microfluidic screening and whole-genome sequencing identifies mutations associated with improved protein secretion by yeast. *Proc National Acad Sci* **112**, E4689–E4696 (2015)), and a previously established chip design (Chaipan, C. et al. Single-Virus Droplet Microfluidics for High-Throughput Screening of Neutralizing Epitopes on HIV Particles. *Cell Chem Biol* **24**, 751-757.e3 (2017)); the guide RNA in this design also serves as a barcode, which allowed to directly identify genes for which an increase or decrease in expression is beneficial for improved protein secretion. As the background strain, a commercially available strain (Ethanol Red) was used, commonly used to produce bioethanol. The strain was engineered to express α -amylase by insertion of an expression cassette containing the codon-optimized α -amylase gene from (*Aspergillus niger*) in the HO-locus and then transformed with plasmid activation or repression libraries. The microfluidic system was used to create droplets containing cells from the transformed protein secreting strain, together with the fluorescent substrate, growth medium and a Tc to induce expression of the guide RNA, these droplets were incubated off chip, before sorting, with gating using thresholds adjusted to capture droplets of average size with the 2-5% highest fluorescence signal into a high fluorescence fraction with the remaining droplets passed passively into a low

fluorescence fraction. Sequencing of the plasmid guide region from the sorted cells allowed to identify the guide population in each fraction.

Sequencing of the original assembled and transformed libraries identified a surviving gRNA representation of 72 and 86 percent, respectively, for the activation and the repression libraries following assembly, and 49 and 69 percent following re-transformation into yeast.

The activation screen identified 71 SUTs or CUTs as significantly enriched, SUTs generate stable transcripts that are thought to interact with other transcripts in both the nucleus and the cytosol, while CUTs are more unstable and quickly degraded upon transcription. An enrichment analysis of genes in the local genomic environment (1kb interval centered on the SUT or CUT guide) identified genes from vacuolar, endosomal, and Golgi and related cellular components as the five most overrepresented cellular components within the range.

Validation of identified genes

A set of genes identified as enriched, were selected for follow-up experimental validation of amylase over-secretion. Genes identified from the activation screens were validated via plasmid-based overexpression of the native gene, while genes from repression screens were validated via gene deletion in both alleles. The units of secreted α -amylase and cell density (OD 600) were measured at several time points after 4, 24, and 48 hours of growth. Overexpression of *ENO2*, *NMA2*, *PRY2*, *SUT074* and *TFG2* resulted in 20-40% increases in total α -amylase secretion after 24 and 48 hours, with even higher increases (35-60%) in the exponential phase after 4 hours of growth, while for *BNA7* and *TOM2* the relative amount of secreted protein per cell was instead significantly increased after 24 and 48 hours and 48 hours of growth respectively. Gene deletions of a smaller set of genes, resulted in increased total protein secretion for *HDA2* (included as a positive control) *MNT2*, *TPO2* after 4 hours, for *INP51* protein secretion was initially significantly decreased after 4 hours, but increased over time and resulted in a significant increase after 48 hours. Deletion of *INP51* also resulted in a significant increase in the secreted protein per cell during all measurements, while for *HDA2* the increase was only significant for the first 24 hours.

The following genes and SUTs (stable uncharacterized transcripts) or CUTs (cryptic unstable transcripts) were identified as being of relevance, and relevance was defined as at least 2% increase of amylase activity (see above). See also Figure 2.

1. Genes that were activated/overexpressed (integration of overexpression cassette into the genome and/or overexpression through a replicative plasmid) after statistical and enrichment analysis – preferred selection. logFC (log fold change) indicates the measure of enrichment, a higher value, equals a higher enrichment in the experiments as performed. FDR (false discovery rate) indicates the corrected p-value, a lower value means less variance between replicates as performed.

Gene (common name, SUT or CUT or systematic designation)	Name and function (if known)	logFC	FDR
MIC19	Component of the MICOS complex	13.883	0.036
TOM22	Translocase of the Outer Mitochondrial membrane; responsible for initial import of mitochondrially directed proteins	13.781	0.008
NKP1	Non-essential Kinetochores Protein	13.389	0.012
DML1	Drosophila melanogaster Misato-Like protein, Essential protein involved in mtDNA inheritance	13.307	0.014
CUT859	SUT or CUT	13.152	0.033
GAL80	GALactose metabolism, Transcriptional regulator involved in the repression of GAL genes	12.170	0.008
APM3	clathrin Adaptor Protein complex Medium chain	12.088	0.020
COQ10	COenzyme Q, Coenzyme Q (ubiquinone) binding protein	12.048	0.025
BLM10	BLeoMycin resistance, Proteasome activator	12.008	0.030
MDH1	Malate DeHydrogenase, Mitochondrial malate dehydrogenase	11.915	0.008
VHS2	Viable in a Hal3 Sit4 background, Regulator of septin dynamics	11.838	0.032
ASA1	AStra Associated protein, Subunit of the ASTRA complex	11.801	0.015

TRP4	TRyPtophan, Anthranilate phosphoribosyl transferase	11.698	0.019
YPS7	YaPSin, Putative GPI-anchored aspartic protease	11.620	0.030
CUT824	SUT or CUT	11.529	0.041
YOR318C	Gene of unknown function	11.515	0.013
PRM7	Pheromone-Regulated Membrane protein	11.485	0.023
ERV46	ER Vesicle, Protein localized to COPII-coated vesicles	11.350	0.010
FIT2	Facilitator of Iron Transport, Mannoprotein that is incorporated into the cell wall	11.287	0.034
GPM3	Glycerate PhosphoMutase	11.062	0.019
CUT892	SUT or CUT	10.972	0.050
SRN2	Suppressor of Rna mutations, Number 2	10.938	0.021
SUT643	SUT or CUT	10.910	0.039
CUT461	SUT or CUT	10.901	0.042
THR4	THReonine requiring, Threonine synthase	10.840	0.047
GMH1	Gea1-6 Membrane-associated High-copy suppressor; Golgi membrane protein of unknown function	10.780	0.055
SOL1	Suppressor Of Los1-1, Protein with a possible role in tRNA export	10.725	0.026
NAB6	Nucleic Acid Binding protein, Putative RNA-binding protein	10.674	0.013
YPR148C	Gene of unknown function	10.614	0.027
ALP1	Arginine transporter	10.598	0.046
CUT097	SUT or CUT	10.597	0.046
ATG33	AuTophagy related, Mitochondrial mitophagy-specific protein	10.585	0.030
YOR316C-A	Gene of unknown function	10.547	0.025
SOG2	Key component of the RAM signaling network; required for proper cell morphogenesis and cell separation after mitosis	10.546	0.039
MCM6	MiniChromosome Maintenance, Protein involved in DNA replication	10.531	0.019

SUT230	SUT or CUT	10.507	0.010
SUT419	SUT or CUT	10.398	0.027
TIF11	Translation Initiation Factor	10.334	0.024
TAF5	TATA binding protein-Associated Factor, involved in RNA polymerase II transcription initiation and in chromatin modification	10.328	0.027
PHO91	PHOSphate metabolism, Low-affinity vacuolar phosphate transporter	10.303	0.024
AIM32	Altered Inheritance rate of Mitochondria, 2Fe-2S mitochondrial protein involved in redox quality control	10.271	0.042
ENO2	ENOLase, Enolase II, a phosphopyruvate hydratase	10.260	0.050
UBA2	UBiquitin Activating, Subunit of heterodimeric nuclear SUMO activating enzyme E1 with Aosl1p	10.215	0.030
PUS5	PseudoUridine Synthase	10.197	0.030
ERG1	ERGosterol biosynthesis, Squalene epoxidase	10.139	0.013
SUT311	SUT or CUT	10.130	0.012
KSS1	Kinase Suppressor of Sst2 mutations, Mitogen-activated protein kinase (MAPK)	10.116	0.039
MRP10	Mitochondrial Ribosomal Protein, Mitochondrial ribosomal protein of the small subunit	10.099	0.023
CUT598	SUT or CUT	10.099	0.046
CUT188	SUT or CUT	10.073	0.026
YOR238W	Gene of unknown function	10.023	0.025
EMW1	Essential for Maintenance of the cell Wall, Essential conserved protein with a role in cell wall integrity	15.549	0.071
BNA7	Biosynthesis of NAD, Formylkynurenine formamidase	14.863	0.071
SNR63	Small Nucleolar RNA, C/D box small nucleolar RNA (snoRNA)	14.717	0.071
CCT3	Chaperonin Containing TCP-1, Subunit of the cytosolic chaperonin Cct ring complex	14.647	0.071

PRY2	Pathogen Related in Yeast, Sterol binding protein involved in the export of acetylated sterols	14.548	0.071
MAL11	MALtose fermentation, High-affinity maltose transporter (alpha-glucoside transporter)	14.484	0.071
KRS1	Lysyl (K) tRNA Synthetase	14.290	0.072
RAI1	Rat1p Interacting Protein, Nuclear decapping endonuclease	14.254	0.071
SUT784	SUT or CUT	13.682	0.071
YPR148C	Gene of unknown function	13.572	0.071
YEL1	Yeast EFA6-Like, Guanine nucleotide exchange factor specific for Arf3p	13.417	0.096
CUT832	SUT or CUT	13.118	0.071
NMA2	Nicotinamide Mononucleotide Adenylyltransferase	13.116	0.071
VPS27	Vacuolar Protein Sorting, Endosomal protein that forms a complex with Hse1p	12.963	0.071
SUT428	SUT or CUT	12.841	0.089
PEX29	PEroXisome related, ER-resident protein involved in peroxisomal biogenesis	12.477	0.071
YLR446W	Gene of unknown function	12.369	0.071
WBP1	Wheat germ agglutinin-Binding Protein, Beta subunit of the oligosaccharyl transferase glycoprotein complex	12.078	0.087
AVT2	Amino acid Vacuolar Transport, Putative transporter	10.965	0.071
CUT854	SUT or CUT	10.873	0.093
TRM10	Transfer RNA Methyltransferase, methylates the N-1 position of guanine at position 9 in tRNAs	10.442	0.099
SLX9	Protein required for pre-rRNA processing	9.996	0.012
YPL077C	Gene of unknown function	9.994	0.038
PET122	PETite colonies, Mitochondrial translational activator specific for the COX3 mRNA	9.982	0.039
TFG2	Transcription Factor G; involved in both transcription initiation and elongation of RNA polymerase II	9.973	0.090

PUN1	Plasma membrane protein Upregulated during Nitrogen stress	9.950	0.027
CUT152	SUT or CUT	9.936	0.020
AIR2	Arginine methyltransferase-Interacting RING finger protein, involved in nuclear RNA processing and degradation	9.886	0.044
CUT571	SUT or CUT	9.799	0.033
RPS26B	Protein component of the small (40S) ribosomal subunit	9.789	0.023
RRT6	Regulator of rDNA Transcription	9.749	0.012
RPC19	RNA Polymerase C, RNA polymerase subunit AC19	9.715	0.047
URA3	URAcil requiring, Orotidine-5'-phosphate (OMP) decarboxylase	9.687	0.046
YGR045C	Gene of unknown function	9.679	0.039
SMC3	Stability of MiniChromosomes, Subunit of the multiprotein cohesin complex	9.669	0.025
PNG1	Peptide N-Glycanase	9.654	0.019
THI6	THIamine biosynthesis, Thiamine-phosphate diphosphorylase and hydroxyethylthiazole kinase	9.653	0.033
MEU1	Multicopy Enhancer of UAS2, Methylthioadenosine phosphorylase (MTAP)	9.558	0.031
CUT239	SUT or CUT	9.531	0.032
NSE4	Non-SMC Element, Component of the SMC5-SMC6 complex	9.502	0.023
SUT074	SUT or CUT	9.478	0.019
AAH1	Adenine AminoHydrolase, Adenine deaminase (adenine aminohydrolase)	9.454	0.044
RMD5	Required for Meiotic nuclear Division, Component of GID Complex that confers ubiquitin ligase (U3) activity	9.452	0.024
CUT607	SUT or CUT	9.313	0.020
ACS1	Acetyl CoA Synthetase, Acetyl-coA synthetase isoform	9.305	0.036

MNN1	MaNNosyltransferase, Alpha-1,3-mannosyltransferase	9.265	0.019
ARH1	Adrenodoxin Reductase Homolog, Oxidoreductase of the mitochondrial inner membrane	9.244	0.039
YHR140W	Gene of unknown function	9.220	0.021
CET1	Capping Enzyme Triphosphatase, RNA 5'-triphosphatase involved in mRNA 5' capping	9.203	0.019
RRB1	Regulator of Ribosome Biogenesis, Specific assembly chaperone for ribosomal protein Rpl3p	9.185	0.030
YLR342W-A	Gene of unknown function	9.166	0.010
RPS22B	Ribosomal Protein of the Small subunit, Protein component of the small (40S) ribosomal subunit	9.154	0.024
CHS5	CHitin Synthase-related, Component of the exomer complex	9.143	0.027
YIL165C	Gene of unknown function	9.140	0.040
SUT093	SUT or CUT	9.139	0.030
LPX1	Lipase of PeroXisomes, Peroxisomal matrix-localized lipase	9.114	0.039
NCA3	Nuclear Control of ATPase, Protein involved in mitochondrion organization	9.078	0.026
EFG1	Exit From G1, Ribosome biogenesis factor required for maturation of 18S rRNA	9.063	0.040
NBP35	Nucleotide Binding Protein, Essential cytoplasmic iron-sulfur cluster binding protein	9.055	0.042
CUT765	SUT or CUT	9.038	0.037
MSL1	MUD Synthetic Lethal	9.019	0.015
SCD6	Suppressor of Clathrin Deficiency, Repressor of translation initiation	9.004	0.025
ATG42	AuTophagy, Vacuolar serine-type carboxypeptidase	9.001	0.028
CHS6	CHitin Synthase-related, Member of the ChAPs (Chs5p-Arf1p-binding proteins) family	8.974	0.020
COQ2	COenzyme Q, Para hydroxybenzoate polyprenyl transferase	8.973	0.045

RPO31	RNA Polymerase, RNA polymerase III largest subunit C160	8.969	0.044
MKK1	Mitogen-activated protein Kinase-Kinase, MAPKK involved in the protein kinase C signaling pathway	8.958	0.030
HED1	High copy suppressor of rED1, Meiosis-specific protein	8.903	0.025
PBP2	Pbp1p Binding Protein, RNA binding protein; has similarity to mammalian heterogeneous nuclear RNP K protein	8.891	0.027
BET5	Blocked Early in Transport, Core component of transport protein particle (TRAPP) complexes I-III	8.890	0.019
CUT678	SUT or CUT	8.876	0.045
YGR021W	Gene of unknown function	8.823	0.012
SUT474	SUT or CUT	8.811	0.042
YGL159W	Gene of unknown function	8.802	0.014
IRC21	Increased Recombination Centers, unknown function	8.795	0.027
VHR1	VHt1 Regulator, Transcriptional activator	8.760	0.046
SPP1	Set1c, Phd finger Protein, Subunit of COMPASS (Set1C)	8.721	0.025
PRP43	Pre-mRNA Processing, RNA helicase in the DEAH-box family	8.707	0.042
ZRT1	Zinc-Regulated Transporter, High-affinity zinc transporter of the plasma membrane; responsible for the majority of zinc uptake	8.705	0.039
YLR041W	Gene of unknown function	8.687	0.044
SUT711	SUT or CUT	8.686	0.039
COX18	Cytochrome c OXidase, Protein required for membrane insertion of C-terminus of Cox2p	8.685	0.046
CBP6	Cytochrome B Protein synthesis, Mitochondrial protein required for translation of the COB mRNA	8.678	0.043
SUT575	SUT or CUT	8.651	0.042
CLG1	Cyclin-Like Gene, Cyclin-like protein that interacts with Pho85p	8.651	0.047

CUT213	SUT or CUT	8.610	0.036
QCR10	ubiQuinol-cytochrome C oxidoReductase, Subunit of the ubiquinol-cytochrome c oxidoreductase complex	8.604	0.019
SNR3	Small Nucleolar RNA, H/ACA box small nucleolar RNA (snoRNA)	8.571	0.044
MSS2	Mitochondrial Splicing, Peripherally bound inner membrane protein of the mitochondrial matrix	8.559	0.023
CUT505	SUT or CUT	8.557	0.039
YOS1	Yip One Suppressor, Integral membrane protein required for ER to Golgi transport	8.540	0.023
SUT073	SUT or CUT	8.519	0.033
UTP21	U Three Protein, Subunit of U3-containing 90S preribosome and SSU processome complexes	8.511	0.039
ACA1	ATF/CREB Activator, ATF/CREB family basic leucine zipper (bZIP) transcription factor	8.478	0.045
CUT632	SUT or CUT	8.475	0.039
RIP1	Rieske Iron-sulfur Protein, Ubiquinol-cytochrome-c reductase	8.466	0.037
HUL5	Hect Ubiquitin Ligase, Multiubiquitin chain assembly factor (E4)	8.383	0.042
CUT727	SUT or CUT	8.373	0.030
RPL35B	Ribosomal 60S subunit protein L35B	8.360	0.019
CUT184	SUT or CUT	8.304	0.039
CUT420	SUT or CUT	8.300	0.023
YFL041W-A	Gene of unknown function	8.290	0.010
SUT460	SUT or CUT	8.248	0.023
ATG10	AuTophagy related, Conserved E2-like conjugating enzyme	8.244	0.019
MFA1	Mating Factor A, Mating pheromone a-factor	8.231	0.023
UGX2	Protein of unknown function	8.226	0.023
TRK2	TRansport of potassium (K), Component of the Trk1p-Trk2p potassium transport system	8.218	0.027

CUT704	SUT or CUT	8.201	0.041
SUT083	SUT or CUT	8.189	0.025
TRE1	Transferrin REceptor like, Transferrin receptor-like protein	8.183	0.046
RVS161	Reduced Viability on Starvation, Amphiphysin-like lipid raft protein	8.110	0.034
LEA1	Looks Exceptionally like U2A, Component of U2 snRNP complex	8.092	0.044
EBP2	EBNA1-binding protein (homolog), Required for 25S rRNA maturation and 60S ribosomal subunit assembly;	8.089	0.030
THI80	THIAMINE metabolism, Thiamine pyrophosphokinase	8.071	0.012
CTI6	Cyc8-Tup1 Interacting protein, Component of the Rpd3L histone deacetylase complex	8.065	0.019
CUT322	SUT or CUT	8.002	0.027
XPT1	Xanthine Phosphoribosyl Transferase, Xanthine-guanine phosphoribosyl transferase	7.984	0.036
MRPL35	Mitochondrial Ribosomal Protein, Large subunit	7.963	0.031
YPL025C	Gene of unknown function	7.962	0.037
SUT737	SUT or CUT	7.950	0.025
PGA2	Processing of Gas1p and ALP, Essential protein required for maturation of Gas1p and Pho8p	7.941	0.046
ULP2	UbL-specific Protease, Peptidase that deconjugates Smt3/SUMO-1 peptides from proteins	7.935	0.033
MRX16	Mitochondrial oRganization of gene eXpression (MIOREX), Protein that associates with the large mitoribosomal subunit	7.917	0.044
EST1	Ever Shorter Telomeres, TLC1 RNA-associated factor involved in telomere length regulation	7.911	0.042
NUP100	NUclear Pore, FG-nucleoporin component of central core of the nuclear pore complex	7.902	0.021
IES3	Ino Eighty Subunit, Subunit of the INO80 chromatin remodeling complex	7.880	0.031

ATG39	AuTophagy related, Autophagy receptor with a role in degradation of the ER and nucleus	7.876	0.038
YMR084W	Gene of unknown function	7.850	0.027
SUT428	SUT or CUT	7.827	0.030
YPL119C-A	Gene of unknown function	7.791	0.031
MIN8	mitochondrial MINi protein of 8 kDa	7.783	0.027
CUT490	SUT or CUT	7.779	0.045
SUT287	SUT or CUT	7.708	0.027
KEL3	KELch	7.705	0.027
SUT678	SUT or CUT	7.699	0.025
SEC3	SECretery, Subunit of the exocyst complex	7.691	0.045
SOL4	Suppressor Of Los1-1, 6-phosphogluconolactonase	7.678	0.030
SIS2	SIt4 Suppressor, Negative regulatory subunit of protein phosphatase 1 (Ppz1p)	7.650	0.026
CUT915	SUT or CUT	7.649	0.044
RRP3	Ribosomal RNA Processing, Protein involved in rRNA processing	7.635	0.034
ESA1	Catalytic subunit of the histone acetyltransferase complex (NuA4)	7.612	0.031
PCL8	Pho85 CycLin, Cyclin	7.581	0.046
TRX3	ThioRedoXin, Mitochondrial thioredoxin	7.579	0.033
YKL115C	Gene of unknown function	7.530	0.043
EMP65	ER Membrane Protein of 65 kDa, Integral membrane protein of the ER	7.520	0.029
ZDS1	Zillion Different Screens, Protein with a role in regulating Swel p-dependent polarized growth	7.488	0.049
CUT167	SUT or CUT	7.486	0.016
SOD1	SuperOxide Dismutase, Cytosolic copper-zinc superoxide dismutase	7.471	0.019
UBR2	Cytoplasmic ubiquitin-protein ligase (E3	7.470	0.044
LSP1	Long chain bases Stimulate Phosphorylation, Eisosome core component	7.391	0.031
SNR81	H/ACA box small nucleolar RNA (snoRNA)	7.389	0.030

RGD3	GTPase activating protein (GAP) for Rho3p	7.370	0.032
YTP1	Yeast putative Transmembrane Protein, Probable type-III integral membrane protein of unknown function	7.365	0.042
SMY2	Suppressor of MYo2-66, involved in COPII vesicle formation	7.352	0.034
CUT449	SUT or CUT	7.340	0.024
FIN1	Filaments In between Nuclei, Spindle pole body-related intermediate filament protein	7.335	0.039
YKL106C-A	Gene of unknown function	7.293	0.021
YAR019W-A	Gene of unknown function	7.280	0.019
CCH1	Calcium Channel Homolog, Voltage-gated high-affinity calcium channel	7.270	0.031
AYR1	1-Acyldihydroxyacetone-phosphate Reductase, ifunctional triacylglycerol lipase and 1-acyl DHAP reductase	7.243	0.012
SUT573	SUT or CUT	7.234	0.042
VNX1	Vacuolar Na ⁺ /H ⁺ exchanger, Calcium/H ⁺ antiporter localized to the endoplasmic reticulum membrane	7.232	0.010
FOL3	FOLic acid synthesis, Dihydrofolate synthetase	7.215	0.032
SUT511	SUT or CUT	7.212	0.026
GIS4	GIg1-2 Suppressor, proposed to be involved in the RAS/cAMP signaling pathway	7.196	0.027
CUT743	SUT or CUT	7.171	0.034
RPL24A	Ribosomal 60S subunit protein L24A	7.169	0.039
HMT1	HnRNP MethylTransferase, Nuclear SAM-dependent mono- and asymmetric methyltransferase	7.163	0.026
SUT333	SUT or CUT	7.141	0.031
SPP2	Suppressor of PrP, Essential protein that promotes the first step of splicing	7.137	0.027
SUT128	SUT or CUT	7.120	0.049
SMC6	Structural Maintenance of Chromosomes, Subunit of the SMC5-SMC6 complex	7.120	0.047

PHR1	PHotoreactivation Repair deficient, DNA photolyase involved in photoreactivation	7.119	0.030
RPS15	Protein component of the small (40S) ribosomal subunit	7.072	0.012
CUT642	SUT or CUT	7.066	0.025
GYP7	Gtpase-activating protein for Ypt7 Protein, GTPase-activating protein for yeast Rab family members	7.063	0.021
tK(CUU)K	Lysine tRNA (tRNA-Lys)	7.034	0.041
CUT896	SUT or CUT	7.026	0.041
SLM5	Synthetic Lethal with Mss4, Mitochondrial asparaginyl-tRNA synthetase	7.024	0.039
CUT586	SUT or CUT	7.020	0.038
CUT158	SUT or CUT	7.003	0.030
RRP12	Ribosomal RNA Processing, Protein required for export of the ribosomal subunits	7.002	0.031
CUT276	SUT or CUT	6.84	0.026
CUT480	SUT or CUT	6.81	0.030
SUT751	SUT or CUT	6.75	0.023
SUT251	SUT or CUT	6.30	0.035
CUT643	SUT or CUT	5.48	0.021

1a. Gene to be preferably combined with the preferred selection

PDI1	Protein Disulfide Isomerase	12.524	0.072
------	-----------------------------	--------	-------

2. Genes or SUTs or CUTs that were inactivated/repressed after statistical and enrichment analysis – preferred selection. logFC (log fold change) indicates the measure of enrichment, a higher value, equals a higher enrichment in the experiments as performed. FDR (false discovery rate) indicates the corrected p-value, a lower value means less variance between replicates as performed.

Gene (common name, SUT or CUT or	Name and function (if known)	logFC	FDR
--	------------------------------	-------	-----

systematic designation)			
TLG2	T-snare affecting a Late Golgi compartment, Syntaxin-like t-SNARE	13.51	0.010
CUT901	SUT or CUT	11.72	0.009
ATG33	AuTophagy related, Mitochondrial mitophagy-specific protein	11.53	0.009
THR4	THReonine requiring, Threonine synthase	11.49	0.009
YDR262W	Gene of unknown function	10.92	0.009
CMC1	Cx9C Mitochondrial protein necessary for full assembly of Cytochrome c oxidase, Copper-binding protein of the mitochondrial intermembrane space	10.86	0.009
MRP17	Mitochondrial ribosomal protein of the small subunit	10.20	0.019
YPT52	Yeast Protein Two, Endosomal Rab family GTPase; required for vacuolar protein sorting	8.91	0.043
CUT312	SUT or CUT	8.90	0.014
MRPS5	Mitochondrial Ribosomal Protein, Small subunit	8.87	0.022
RDR1	Repressor of Drug Resistance, Transcriptional repressor involved in regulating multidrug resistance	8.65	0.042
DAL7	Degradation of Allantoin, Malate synthase	8.55	0.009
RPL20A	Ribosomal 60S subunit protein L20A	8.19	0.025
YBR137W	Gene of unknown function	8.11	0.056
RPL36B	Ribosomal 60S subunit protein L36B	8.02	0.028
YEL008C-A	Gene of unknown function	7.85	0.062
RAX1	Revert to Axial, Protein involved in establishing bud site selection	7.65	0.019
INP51	INositol polyphosphate 5-Phosphatase	7.51	0.102
CUT729	SUT or CUT	7.27	0.066
UBP8	UBiquitin-specific processing Protease, Ubiquitin-specific protease component of the SAGA acetylation complex	7.18	0.066
CUT258	SUT or CUT	7.10	0.089
YLR342W-A	Gene of unknown function	7.09	0.025

SUT568	SUT or CUT	7.04	0.027
PEX7	PEroXin, Peroxisomal signal receptor for peroxisomal matrix proteins	7.00	0.024
MSD1	Mitochondrial aminoacyl-tRNA Synthetase, Aspartate (D)	6.97	0.089
CUT136	SUT or CUT	6.88	0.039
TIM10	Translocase of the Inner Membrane, Essential protein of the mitochondrial intermembrane space	6.84	0.064
CUT361	SUT or CUT	6.83	0.037
snR51	Small Nucleolar RNA	6.80	0.085
TAL1	TransALdolase, Transaldolase, enzyme in the non-oxidative pentose phosphate pathway	6.74	0.069
RIP1	Rieske Iron-sulfur Protein, Ubiquinol-cytochrome-c reductase	6.65	0.058
MRP10	Mitochondrial Ribosomal Protein, Mitochondrial ribosomal protein of the small subunit	6.63	0.051
SUT078	SUT or CUT	6.52	0.074
MRP51	Mitochondrial Ribosomal Protein, Mitochondrial ribosomal protein of the small subunit	6.51	0.065
GLO3	GLyOxalase, ADP-ribosylation factor GTPase activating protein (ARF GAP); involved in ER-Golgi transport	6.51	0.053
EHD3	3-hydroxyisobutyryl-CoA hydrolase	6.50	0.025
HER1	Hmg2p ER Remodeling, Protein of unknown function	6.48	0.051
NMA111	Nuclear Mediator of Apoptosis, Serine protease and general molecular chaperone	6.45	0.041
PBP4	Pbp1p binding protein	6.28	0.044
MFB1	Mitochondria-associated F-box protein; involved in maintenance of normal mitochondrial morphology	6.25	0.098
IKI3	Insensitive to Killer toxin, Subunit of Elongator complex	6.21	0.031
NDL1	NuDeL homolog, Homolog of nuclear distribution factor NudE	6.15	0.057
SUT433	SUT or CUT	5.99	0.022
YOR238W	Gene of unknown function	5.91	0.054

SUT750	SUT or CUT	5.86	0.016
QDR2	QuiniDine Resistance, Plasma membrane transporter of the major facilitator superfamily	5.84	0.020
RDI1	Rho GDP Dissociation Inhibitor	5.79	0.023
SUT014	SUT or CUT	5.76	0.059
CUT437	SUT or CUT	5.75	0.045
MSC6	Meiotic Sister-Chromatid recombination, Multicopy suppressor of HER2 involved in mitochondrial translation	5.66	0.055
SUT497	SUT or CUT	5.54	0.072
YCR051W	Gene of unknown function	5.52	0.076
MRPL33	Mitochondrial Ribosomal Protein, Large subunit	5.47	0.024
RPL14A	Ribosomal 60S subunit protein L14A	5.46	0.077
TRM7	2'-O-ribose methyltransferase	5.43	0.081
RNH202	Ribonuclease H2 subunit; required for RNase H2 activity	5.43	0.083
RTC5	Restriction of Telomere Capping, Protein of unknown function	5.38	0.060
SUT027	SUT or CUT	5.34	0.058
CDC5	Cell Division Cycle, Polo-like kinase essential for mitotic cell cycle	5.33	0.070
SUT729	SUT or CUT	5.30	0.076
YOR131C	Gene of unknown function	5.28	0.078
CUT665	SUT or CUT	5.12	0.097
GLG2	Glycogenin-Like Gene, Glycogenin glucosyltransferase	5.12	0.079
SUT268	SUT or CUT	4.89	0.087
SUT705	SUT or CUT	4.87	0.086
MED4	MEDIator complex, Subunit of the RNA polymerase II mediator complex	4.61	0.093
RCR2	Resistance to Congo Red, Vacuolar ubiquitin ligase-substrate adaptor	4.59	0.055
EFB1	Elongation Factor Beta, Translation elongation factor 1 beta	4.58	0.036
RXT2	Component of the histone deacetylase Rpd3L complex	4.49	0.073

KGD1	alpha-KetoGlutarate Dehydrogenase, Subunit of the mitochondrial alpha-ketoglutarate dehydrogenase complex	4.42	0.093
TUP1	dTMP-UPtake, General repressor of transcription	4.35	0.080
RNH203	Ribonuclease H2 subunit	4.31	0.096
YDR338C	Gene of unknown function	3.92	0.029
SED1	Suppression of Exponential Defect, Major stress-induced structural GPI-cell wall glycoprotein	3.81	0.089
CUT522	SUT or CUT	3.75	0.092
HIS2	HIStidine requiring, Histidinolphosphatase	3.74	0.090
SUT145	SUT or CUT	3.67	0.072
MET17	METHionine requiring, O-acetyl homoserine-O-acetyl serine sulfhydrylase	3.58	0.063
APC4	Anaphase Promoting, Subunit of the Anaphase-Promoting Complex/Cyclosome (APC/C)	3.58	0.077
NKP2	Non-essential Kinetochoe Protein, Central kinetochoe protein and subunit of the Ctf19 complex	3.54	0.022
MKK2	Mitogen-activated Kinase Kinase, MAPKK involved in the protein kinase C signaling pathway	3.05	0.042
NDC1	Nuclear Division Cycle, Subunit of the transmembrane ring of the nuclear pore complex (NPC)	14.18	0.079
PET100	PETite colonies, Chaperone that facilitates the assembly of cytochrome c oxidase	12.69	0.086
NIP7	Nuclear ImPort, Nucleolar protein required for 60S ribosome subunit biogenesis	12.54	0.086
VHT1	Vitamin H Transporter, High-affinity plasma membrane H ⁺ -biotin (vitamin H) symporter	12.31	0.086
SUT685	SUT or CUT	12.07	0.086
BNI5	Bud Neck Involved, Linker protein responsible for recruitment of myosin to the bud neck	11.96	0.086
SNA3	Sensitivity to NA ⁺ , Protein involved in efficient MVB sorting of proteins to the vacuole	11.93	0.086
EGH1	Cryptococcus neoformans EGCp2 Homolog, Steryl-beta-glucosidase with broad specificity for aglycones	11.81	0.086

MRP4	Mitochondrial ribosomal protein of the small subunit	11.67	0.086
POB3	POI1 Binding, Subunit of the heterodimeric FACT complex (Spt16p-Pob3p)	10.87	0.086
PIB2	PtdIns(3)p-Binding, Phosphatidylinositol 3-phosphate binding protein	10.80	0.086
SUT317	SUT or CUT	10.74	0.086
NTO1	NuA Three Orf, Subunit of the NuA3 histone acetyltransferase complex	10.62	0.086
YKL024C	URA6 Uridylate kinase; catalyzes the seventh enzymatic step in the de novo biosynthesis of pyrimidines	7.08	0.102
YGL116W	CDC20 Activator of anaphase-promoting complex/cyclosome (APC/C)	6.44	0.140
YLR118C	TML25 Acyl-protein thioesterase responsible for depalmitoylation of Gpa1p	6.24	0.149
YFR031C-A	RPL2A Ribosomal 60S subunit protein L2A	6.16	0.104
YGL190C	CDC55 Regulatory subunit B of protein phosphatase 2A (PP2A)	6.02	0.134
YDL108W	KIN28 Ser/Thr protein kinase and subunit of TFI _{IK} , a TFI _{IIH} subassembly	5.81	0.101
YMR128W	ECM16 Essential DEAH-box ATP-dependent RNA helicase specific to U3 snoRNP	5.57	0.118
YBR253W	SRB6 Subunit of the RNA polymerase II mediator complex	5.33	0.142
YJR113C	RSM7 Mitochondrial ribosomal protein of the small subunit	5.28	0.134
YIL031W	ULP2 Peptidase that deconjugates Smt3/SUMO-1 peptides from proteins	5.20	0.140
YGR109C	CLB6 B-type cyclin involved in DNA replication during S phase	5.15	0.141
YBR282W	MRPL27 Mitochondrial ribosomal protein of the large subunit	5.14	0.136
YMR125W	STO1 Large subunit of the nuclear mRNA cap-binding protein complex	4.97	0.106
YMR236W	TAF9 Subunit (17 kDa) of TFI _{IID} and SAGA complexes	4.96	0.125
YDR411C	DFM1 Endoplasmic reticulum (ER) localized protein	4.71	0.108

YML029W	USA1 Scaffold subunit of the Hrd1p ubiquitin ligase	4.55	0.106
YDL033C	SLM3 tRNA-specific 2-thiouridylase	4.50	0.131
YPL050C	MNN9 Subunit of Golgi mannosyltransferase complex	4.42	0.102
YHR171W	ATG7 Autophagy-related protein and dual specificity member of the E1 family	4.32	0.143
YDR352W	YPQ2 Putative vacuolar membrane transporter for cationic amino acids	4.27	0.137

3. Genes that were either overexpressed or inactivated/repressed depending on experimental conditions after statistical and enrichment analysis preferred selection. logFC (log fold change) indicates the measure of enrichment, a higher value, equals a higher enrichment in the experiments as performed. FDR (false discovery rate) indicates the corrected p-value, a lower value means less variance between replicates as performed.

Gene (common name, SUT or CUT or designation)	Name and function (if known)	logFC activation	logFC repression	FDR activation	FDR repression
THR4	THReonine requiring, Threonine synthase	10.84	11.49	0.047	0.009
MRP10	Mitochondrial Ribosomal Protein, Mitochondrial ribosomal protein of the small subunit	10.10	6.63	0.023	0.051
RIP1	Rieske Iron-sulfur Protein, Ubiquinol- cytochrome-c reductase	8.47	6.65	0.037	0.058
YLR342W-A	Gene of unknown function	9.17	7.09	0.010	0.025
ATG33	AuTophagy related, Mitochondrial	10.59	11.53	0.030	0.009

	mitophagy-specific protein				
YOR238W	Gene of unknown function	10.02	5.91	0.025	0.054

3. Genes that were overexpressed – particularly preferred selection

Gene (common name, SUT or CUT or systematic designation)	Name and function (if known)	logFC	FDR
MIC19	Component of the MICOS complex	13.883	0.036
TOM22	Translocase of the Outer Mitochondrial membrane; responsible for initial import of mitochondrially directed proteins	13.781	0.008
NKP1	Non-essential Kinetochore Protein	13.389	0.012
DML1	Drosophila melanogaster Misato-Like protein, Essential protein involved in mtDNA inheritance	13.307	0.014
CUT859	SUT or CUT	13.152	0.033
GAL80	GALactose metabolism, Transcriptional regulator involved in the repression of GAL genes	12.170	0.008
APM3	clathrin Adaptor Protein complex Medium chain	12.088	0.020
COQ10	COenzyme Q, Coenzyme Q (ubiquinone) binding protein	12.048	0.025
BLM10	BLeoMycin resistance, Proteasome activator	12.008	0.030
MDH1	Malate DeHydrogenase, Mitochondrial malate dehydrogenase	11.915	0.008
VHS2	Viable in a Hal3 Sit4 background, Regulator of septin dynamics	11.838	0.032
ASA1	AStra Associated protein, Subunit of the ASTRA complex	11.801	0.015
TRP4	TRyPtophan, Anthranilate phosphoribosyl transferase	11.698	0.019
YPS7	YaPSin, Putative GPI-anchored aspartic protease	11.620	0.030

CUT824	SUT or CUT	11.529	0.041
YOR318C	Gene of unknown function	11.515	0.013
PRM7	Pheromone-Regulated Membrane protein	11.485	0.023
ERV46	ER Vesicle, Protein localized to COPII-coated vesicles	11.350	0.010
FIT2	Facilitator of Iron Transport, Mannoprotein that is incorporated into the cell wall	11.287	0.034
GPM3	Glycerate PhosphoMutase	11.062	0.019
CUT892	SUT or CUT	10.972	0.050
SRN2	Suppressor of Rna mutations, Number 2	10.938	0.021
SUT643	SUT or CUT	10.910	0.039
CUT461	SUT or CUT	10.901	0.042
THR4	THReonine requiring, Threonine synthase	10.840	0.047
GMH1	Geal-6 Membrane-associated High-copy suppressor; Golgi membrane protein of unknown function	10.780	0.055
SOL1	Suppressor Of Los1-1, Protein with a possible role in tRNA export	10.725	0.026
NAB6	Nucleic Acid Binding protein, Putative RNA-binding protein	10.674	0.013
YPR148C	Gene of unknown function	10.614	0.027
ALP1	Arginine transporter	10.598	0.046
CUT097	SUT or CUT	10.597	0.046
ATG33	AuTophagy related, Mitochondrial mitophagy-specific protein	10.585	0.030
YOR316C-A	Gene of unknown function	10.547	0.025
SOG2	Key component of the RAM signaling network; required for proper cell morphogenesis and cell separation after mitosis	10.546	0.039
MCM6	MiniChromosome Maintenance, Protein involved in DNA replication	10.531	0.019
SUT230	SUT or CUT	10.507	0.010
SUT419	SUT or CUT	10.398	0.027

TIF11	Translation Initiation Factor	10.334	0.024
TAF5	TATA binding protein-Associated Factor, involved in RNA polymerase II transcription initiation and in chromatin modification	10.328	0.027
PHO91	PHOSphate metabolism, Low-affinity vacuolar phosphate transporter	10.303	0.024
AIM32	Altered Inheritance rate of Mitochondria, 2Fe-2S mitochondrial protein involved in redox quality control	10.271	0.042
ENO2	ENOlase, Enolase II, a phosphopyruvate hydratase	10.260	0.050
UBA2	UBiquitin Activating, Subunit of heterodimeric nuclear SUMO activating enzyme E1 with Aosl p	10.215	0.030
PUS5	PseudoUridine Synthase	10.197	0.030
ERG1	ERGosterol biosynthesis, Squalene epoxidase	10.139	0.013
SUT311	SUT or CUT	10.130	0.012
KSS1	Kinase Suppressor of Sst2 mutations, Mitogen-activated protein kinase (MAPK)	10.116	0.039
MRP10	Mitochondrial Ribosomal Protein, Mitochondrial ribosomal protein of the small subunit	10.099	0.023
CUT598	SUT or CUT	10.099	0.046
CUT188	SUT or CUT	10.073	0.026
YOR238W	Gene of unknown function	10.023	0.025
EMW1	Essential for Maintenance of the cell Wall, Essential conserved protein with a role in cell wall integrity	15.549	0.071
BNA7	Biosynthesis of NAD, Formylkynurenine formamidase	14.863	0.071
SNR63	Small Nucleolar RNA, C/D box small nucleolar RNA (snoRNA)	14.717	0.071
CCT3	Chaperonin Containing TCP-1, Subunit of the cytosolic chaperonin Cct ring complex	14.647	0.071
PRY2	Pathogen Related in Yeast, Sterol binding protein involved in the export of acetylated sterols	14.548	0.071
MAL11	MALtose fermentation, High-affinity maltose transporter (alpha-glucoside transporter)	14.484	0.071

KRS1	Lysyl (K) tRNA Synthetase	14.290	0.072
RAI1	Rat1p Interacting Protein, Nuclear decapping endonuclease	14.254	0.071
SUT784	SUT or CUT	13.682	0.071
YPR148C	Gene of unknown function	13.572	0.071
YEL1	Yeast EFA6-Like, Guanine nucleotide exchange factor specific for Arf3p	13.417	0.096
CUT832	SUT or CUT	13.118	0.071
NMA2	Nicotinamide Mononucleotide Adenylyltransferase	13.116	0.071
VPS27	Vacuolar Protein Sorting, Endosomal protein that forms a complex with Hse1p	12.963	0.071
SUT428	SUT or CUT	12.841	0.089
PEX29	PEroXisome related, ER-resident protein involved in peroxisomal biogenesis	12.477	0.071
YLR446W	Gene of unknown function	12.369	0.071
WBP1	Wheat germ agglutinin-Binding Protein, Beta subunit of the oligosaccharyl transferase glycoprotein complex	12.078	0.087
AVT2	Amino acid Vacuolar Transport, Putative transporter	10.965	0.071
CUT854	SUT or CUT	10.873	0.093
TRM10	Transfer RNA Methyltransferase, methylates the N-1 position of guanine at position 9 in tRNAs	10.442	0.099

Gene to be preferably combined with the particularly preferred selection

PDI1	Protein Disulfide Isomerase	12.524	0.072
------	-----------------------------	--------	-------

4. Genes or SUTs or CUTs that were inactivated/repressed after statistical and enrichment analysis – particularly preferred selection

Gene (common name, SUT or CUT or systematic designation)	Name and function (if known)	logFC	FDR
--	------------------------------	-------	-----

TLG2	T-snare affecting a Late Golgi compartment, Syntaxin-like t-SNARE	13.51	0.010
CUT901	SUT or CUT	11.72	0.009
ATG33	AuTophagy related, Mitochondrial mitophagy-specific protein	11.53	0.009
THR4	THReonine requiring, Threonine synthase	11.49	0.009
YDR262W	Gene of unknown function	10.92	0.009
CMC1	Cx9C Mitochondrial protein necessary for full assembly of Cytochrome c oxidase, Copper-binding protein of the mitochondrial intermembrane space	10.86	0.009
MRP17	Mitochondrial ribosomal protein of the small subunit	10.20	0.019
NDC1	Nuclear Division Cycle, Subunit of the transmembrane ring of the nuclear pore complex (NPC)	14.18	0.079
PET100	PETite colonies, Chaperone that facilitates the assembly of cytochrome c oxidase	12.69	0.086
NIP7	Nuclear ImPort, Nucleolar protein required for 60S ribosome subunit biogenesis	12.54	0.086
VHT1	Vitamin H Transporter, High-affinity plasma membrane H ⁺ -biotin (vitamin H) symporter	12.31	0.086
SUT685	SUT or CUT	12.07	0.086
BNI5	Bud Neck Involved, Linker protein responsible for recruitment of myosin to the bud neck	11.96	0.086
SNA3	Sensitivity to NA ⁺ , Protein involved in efficient MVB sorting of proteins to the vacuole	11.93	0.086
EGH1	Cryptococcus neoformans EGCp2 Homolog, Steryl-beta-glucosidase with broad specificity for aglycones	11.81	0.086
MRP4	Mitochondrial ribosomal protein of the small subunit	11.67	0.086
POB3	POI1 Binding, Subunit of the heterodimeric FACT complex (Spt16p-Pob3p)	10.87	0.086
PIB2	PtdIns(3)p-Binding, Phosphatidylinositol 3-phosphate binding protein	10.80	0.086
SUT317	SUT or CUT	10.74	0.086

NT01	NuA Three Orf, Subunit of the NuA3 histone acetyltransferase complex	10.62	0.086
------	--	-------	-------

5. Genes that were overexpressed – most preferred selection

Gene (common name, SUT or CUT or systematic designation)	Name and function (if known)	logFC	FDR
MIC19	Component of the MICOS complex	13.883	0.036
TOM22	Translocase of the Outer Mitochondrial membrane; responsible for initial import of mitochondrially directed proteins	13.781	0.008
NKP1	Non-essential Kinetochore Protein	13.389	0.012
DML1	Drosophila melanogaster Misato-Like protein, Essential protein involved in mtDNA inheritance	13.307	0.014
CUT859	SUT or CUT	13.152	0.033
GAL80	GALactose metabolism, Transcriptional regulator involved in the repression of GAL genes	12.170	0.008
APM3	clathrin Adaptor Protein complex Medium chain	12.088	0.020
COQ10	COenzyme Q, Coenzyme Q (ubiquinone) binding protein	12.048	0.025
BLM10	BLeoMycin resistance, Proteasome activator	12.008	0.030
MDH1	Malate DeHydrogenase, Mitochondrial malate dehydrogenase	11.915	0.008
EMW1	Essential for Maintenance of the cell Wall, Essential conserved protein with a role in cell wall integrity	15.549	0.071
BNA7	Biosynthesis of NAD, Formylkynurenine formamidase	14.863	0.071
SNR63	Small Nucleolar RNA, C/D box small nucleolar RNA (snoRNA)	14.717	0.071
CCT3	Chaperonin Containing TCP-1, Subunit of the cytosolic chaperonin Cct ring complex	14.647	0.071

PRY2	Pathogen Related in Yeast, Sterol binding protein involved in the export of acetylated sterols	14.548	0.071
MAL11	MALtose fermentation, High-affinity maltose transporter (alpha-glucoside transporter)	14.484	0.071
KRS1	Lysyl (K) tRNA Synthetase	14.290	0.072
RAI1	Rat1p Interacting Protein, Nuclear decapping endonuclease	14.254	0.071
SUT784	SUT or CUT	13.682	0.071
YPR148C	Gene of unknown function	13.572	0.071
YEL1	Yeast EFA6-Like, Guanine nucleotide exchange factor specific for Arf3p	13.417	0.096
CUT832	SUT or CUT	13.118	0.071
NMA2	Nicotinamide Mononucleotide Adenylyltransferase	13.116	0.071
VPS27	Vacuolar Protein Sorting, Endosomal protein that forms a complex with Hse1p	12.963	0.071
SUT428	SUT or CUT	12.841	0.089
PEX29	PEroXisome related, ER-resident protein involved in peroxisomal biogenesis	12.477	0.071
YLR446W	Gene of unknown function	12.369	0.071
WBP1	Wheat germ agglutinin-Binding Protein, Beta subunit of the oligosaccharyl transferase glycoprotein complex	12.078	0.087

Gene to be preferably combined with the most preferred selection

PDI1	Protein Disulfide Isomerase	12.524	0.072
------	-----------------------------	--------	-------

6. Genes or SUTs or CUTs that were inactivated/repressed after statistical and enrichment analysis – most preferred selection

Gene (common name, SUT or CUT or systematic designation)	Name and function (if known)	logFC	FDR
TLG2	T-snare affecting a Late Golgi compartment, Syntaxin-like t-SNARE	13.51	0.010
CUT901	SUT or CUT	11.72	0.009

ATG33	AuTophagy related, Mitochondrial mitophagy-specific protein	11.53	0.009
THR4	THReonine requiring, Threonine synthase	11.49	0.009
NDC1	Nuclear Division Cycle, Subunit of the transmembrane ring of the nuclear pore complex (NPC)	14.18	0.079
PET100	PETite colonies, Chaperone that facilitates the assembly of cytochrome c oxidase	12.69	0.086
NIP7	Nuclear ImPort, Nucleolar protein required for 60S ribosome subunit biogenesis	12.54	0.086
VHT1	Vitamin H Transporter, High-affinity plasma membrane H ⁺ -biotin (vitamin H) symporter	12.31	0.086
SUT685	SUT or CUT	12.07	0.086

Preferred are further genes or SUTs or CUTs that are selected from the group of genes or SUTs or CUTs having a value of logFC/FDR logFC/FDR of more than 40, preferably of more than 200, more preferred of more than 300, and most preferred of more than 500, based on the values herein.

References as cited

1. Martinez Ruiz, J.; Liu, L.; Petranovic, D. (2012) "Pharmaceutical protein production by yeast: towards production of human blood proteins by microbial fermentation". Current Opinion in Biotechnology, vol. 23(6), pp. 965-971.
2. Falch EA. Industrial enzymes--developments in production and application. Biotechnol Adv. 1991;9(4):643-58. doi: 10.1016/0734-9750(91)90736-f. PMID: 14542053.
3. Demain AL, Vaishnav P. Production of recombinant proteins by microbes and higher organisms. Biotechnol Adv. 2009 May-Jun;27(3):297-306. doi: 10.1016/j.biotechadv.2009.01.008. Epub 2009 Jan 31. PMID: 19500547.
4. Zahrl RJ, Gasser B, Mattanovich D, Ferrer P. Detection and Elimination of Cellular Bottlenecks in Protein-Producing Yeasts. Methods Mol Biol. 2019;1923:75-95. doi: 10.1007/978-1-4939-9024-5_2. PMID: 30737735
5. Parapouli M, Vasileiadis A, Afendra AS, Hatziloukas E. *Saccharomyces cerevisiae* and its industrial applications. AIMS Microbiol. 2020 Feb 11;6(1):1-31. doi: 10.3934/microbiol.2020001. PMID: 32226912; PMCID: PMC7099199.

6. Dominguez AA, Lim WA, Qi LS. Beyond editing: repurposing CRISPR-Cas9 for precision genome regulation and interrogation. *Nat Rev Mol Cell Biol.* 2016 Jan;17(1):5-15. doi: 10.1038/nrm.2015.2. Epub 2015 Dec 16. PMID: 26670017; PMCID: PMC4922510.

Claims

1. A cell of *Saccharomyces cerevisiae*, producing at least one secreted protein of interest, wherein said cell comprises at least one fungal gene selected from the group consisting of ENO2, NMA2, PRY2, SUT074, TFG2, AVT2, TRM10, BNA7, and TOM22, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, MNT2, TPO2, ATG33, THR4, INP51, CUT901, YDR262W, MRP10, NDC1, and CMC1, wherein said at least one fungal gene shows reduced expression and/or inactivation, and optionally further comprising the fungal gene *HDA2* and/or *PDH1*, showing an increased expression and/or overexpression.
2. The yeast cell according to claim 1, wherein said cell comprises at least one fungal gene selected from the groups consisting of ENO2, NMA2, PRY2, SUT074, and TFG2, or AVT2, TRM10, PRY2, SUT074, BNA7, and TOM22, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the groups consisting of TLG2, CUT901, ATG33, THR4, YDR262W, and CMC1, or MRP10, TLG2, CUT901, ATG33, THR4, YDR262W, CMC1, MNT2, TPO2, and NDC1, preferably MNT2 and TPO2, wherein said at least one fungal gene shows reduced expression and/or inactivation, and optionally further comprising the fungal genes *HDA2* and/or *PDH1*, showing an increased expression and/or overexpression, and/or *INP51* showing an reduced expression and/or inactivation.
3. The yeast cell according to claim 1 or 2, wherein said genes or SUTs or CUTs are furthermore selected from the group of genes or SUTs or CUTs having a value of logFC/FDR logFC/FDR of more than 40, preferably of more than 200, more preferred of more than 300, and most preferred of more than 500, based on the values as determined herein.
4. The yeast cell according to any one of claims 1 to 3, wherein said yeast cell is from *Saccharomyces cerevisiae* strain ER.sec2.
5. The yeast cell according to any one of claims 1 to 4, wherein said at least one secreted protein of interest also shows an increased expression and/or overexpression.

6. The yeast cell according to any one of claims 1 to 5, wherein said at least one fungal gene showing increased expression and/or overexpression and/or showing reduced expression and/or inactivation is a native gene and/or is a recombinant gene, wherein preferably said recombinant gene is integrated into the genome as an expression cassette and/or extrachromosomally expressed, preferably using a replicative expression vector.
7. The yeast cell according to any one of claims 1 to 6, wherein the cell furthermore comprises at least one additional recombinant secretion promoting gene, for example a gene for a chaperone, for a foldase and/or for a glycosylation-promoting protein.
8. The yeast cell according to any one of claims 1 to 7, wherein the increased expression and/or overexpression or reduced expression and/or inactivation of the at least one fungal gene or the at least one additional recombinant secretion promoting gene is constitutive or inducible.
9. The yeast cell according to any one of claims 1 to 8, wherein the cell produces the at least one secreted protein to about 30% or more, or about 40% or more, preferably about 50% or more, more preferably to about 75% or more, when compared to a control yeast or filamentous fungal cell.
10. A method for producing a secreted protein in a yeast cell, comprising the steps of i) providing a cell of *Saccharomyces cerevisiae* producing at least one secreted protein of interest according to any one of claims 1 to 9, ii) culturing said yeast cell in suitable culture medium, and iii) isolating said secreted protein from said culture medium, and optionally further comprising suitably inducing the increased expression and/or overexpression or reduced expression and/or inactivation of the at least one fungal gene.
11. The method according to claim 10, wherein preferably about 30% or more, or about 40% or more, preferably about 50% or more, more preferably to about 75% or more of said at least one secreted protein is produced, when compared to the production of a control yeast cell.
12. A method for producing a yeast cell producing at least one secreted protein of interest, comprising introducing into said cell producing at least one secreted protein of interest at least one fungal gene selected from the group consisting of ENO2, NMA2, PRY2, SUT074, TFG2,

AVT2, TRM10, BNA7, and TOM22, wherein said at least one fungal gene shows increased expression and/or overexpression, and/or wherein said cell comprises at least one fungal gene selected from the group consisting of TLG2, MNT2, TPO2, ATG33, THR4, INP51, CUT901, YDR262W, MRP10, NDC1, and CMC1, preferably MNT2, and TPO2, wherein said at least one fungal gene shows reduced expression and/or inactivation, and optionally further introducing into said cell a fungal gene selected from the group consisting of RIP1, YLR342W-A, and YOR238W, either showing an increased expression and/or overexpression or reduced expression and/or inactivation, depending on the experimental conditions, and/or optionally further introducing into said cell the fungal gene *HDA2* and/or *PDII*, showing an increased expression and/or overexpression.

13. The method according to any one of claims 10 to 12, wherein said at least one fungal gene is integrated into the genome as an expression cassette and/or extrachromosomally expressed, preferably using a replicative expression vector.

14. Use of a yeast cell according to any one of claims 1 to 9 for producing at least one secreted protein of interest.

Figure 1




Figure 2 - A




Figure 2 (continued) - B




Figure 3 – A




Figure 3 – B




Figure 4 – A




Figure 4 (continued) – B




Figure 5 – A




Figure 5 (continued) – B

