

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 February 2007 (08.02.2007)

PCT

(10) International Publication Number
WO 2007/014851 A2

(51) International Patent Classification:
C07D 401/06 (2006.01) C07D 209/22 (2006.01)
C07D 405/14 (2006.01) C07D 209/42 (2006.01)

(74) Agent: WASNER, Marita; Grenzacherstrasse 124,
CH-4070 Basel (CH).

(21) International Application Number:
PCT/EP2006/064395

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date: 19 July 2006 (19.07.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
05107044.9 29 July 2005 (29.07.2005) EP
05111072.4 22 November 2005 (22.11.2005) EP

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): **F. HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH).

(72) Inventors; and


(75) Inventors/Applicants (for US only): **BISSANTZ, Caterina** [DE/FR]; 3, Rue Du Palmrain, F-68128 Village Neuf (FR). **GRUNDSCHÖBER, Christophe** [CH/CH]; Landskronstrasse 18, CH-4118 Rodersdorf (CH). **RATNI, Hasane** [FR/FR]; 4, Rue Louis Pasteur, F-68440 Habsheim (FR). **ROGERS-EVANS, Mark** [GB/CH]; Im Zehnten-frei 19, CH-4102 Binningen (CH). **SCHNIDER, Patrick** [CH/CH]; Birnbaumweg 15, CH-4103 Bottmingen (CH).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INDOL-3-YL-CARBONYL-PIPERIDIN AND PIPERAZIN DERIVATIVES




(57) Abstract: The present invention relates to indol-3-yl-carbonyl-piperidin and piperazin derivatives which act as V1a receptor antagonists and which are represented by Formula (I) wherein the residues R¹ to R³ are as defined herein. The invention further relates to pharmaceutical compositions containing such compounds, their use in medicaments against dysmenorrhea, hypertension, chronic heart failure, inappropriate secretion of vasopressin, liver cirrhosis, nephrotic syndrome, obsessive compulsive disorder, anxious and depressive disorders, and methods of preparation thereof.

WO 2007/014851 A2

Case 23237INDOL-3-YL-CARBONYL-PIPERIDIN AND PIPERAZIN DERIVATIVES


The present invention relates to compounds of general formula (I)



wherein

- R^1 is H,
- 5 or is C_{1-6} -alkyl substituted by CN, C_{1-6} -alkoxy, OH, halo, or by NR^iR^{ii} ,
 or is C_{2-6} -alkyl,
 or is aryl, 5 or 6 membered heteroaryl or sulfonylaryl which are optionally
 substituted by one or more B,
 or is $-(CH_2)_m-R^a$ wherein R^a is:
- 10 CN,
 OR^i ,
 NR^iR^{ii} ,
 C_{3-6} -cycloalkyl, 4 to 7 membered-heterocycloalkyl, aryl, or 5 or 6 membered
 heteroaryl which are optionally substituted by one or more B,
- 15 or is $-(CH_2)_n-(CO)-R^b$ or $-(CH_2)_n-(SO_2)-R^b$, wherein R^b is:
 C_{1-6} -alkyl,
 C_{1-6} -alkoxy,
 C_{3-6} -cycloalkyl,
 $-(CH_2)_m-NR^{iii}R^{iv}$,
- 20 NR^iR^{ii} ,
 C_{3-6} -cycloalkyl, 4 to 7 membered-heterocycloalkyl, aryl, or 5 or 6 membered
 heteroaryl which are optionally substituted by one or more B,
 or R^1 and R^3 together with the indole ring to which they are attached form a 5 or
 6 membered heterocycloalkyl which can be substituted by =O, $C(O)O-C_{1-6}$ -
 25 alkyl or C_{1-6} -alkyl;

- R^2 is one or more of H, OH, halo, CN, nitro, C_{1-6} -alkoxy, $-O-CH_2-C_{2-6}$ -alkenyl, benzyloxy, C_{1-6} -haloalkoxy, C_{1-6} -alkyl optionally substituted by $-NR^{iii}R^{iv}$ or halo, or two R^2 may form an oxo or dioxo bridge together with the indole ring to which they are attached;
- 5 R^3 is H,
or is F,
or is $-(CO)-R^c$, wherein R^c is:
- C_{1-6} -alkyl,
 $-(CH_2)_n-NR^iR^{ii}$,
10 $-(CH_2)_n-NR^{iii}R^{iv}$,
- 5 or 6 membered heterocycloalkyl optionally substituted by C_{1-6} -alkyl,
or is C_{1-6} -alkyl which is optionally substituted by
- halo,
 NR^iR^{ii} ,
15 $NR^{iii}R^{iv}$,
- $-O(CO)-C_{1-6}$ -alkyl,
or by $-NH(CO)R^d$, wherein R^d is C_{1-6} -alkyl optionally substituted by halo or nitro, or R^d is aryl or a 5 or 6 membered heteroaryl, which are optionally substituted by halo, nitro, C_{1-6} -alkyl or C_{1-6} -haloalkyl;
- 20 A is selected from the group consisting of (a), (a'), (b), (c) and (d):



wherein

- R^4 is $-NH(CO)R^e$, wherein R^e is C_{1-6} -alkoxy, or aryl optionally substituted by halo, or is C_{1-6} -alkoxy, CN,
- 25 or is aryl, 5 or 6 membered heteroaryl, benzyl, aryloxy or a 9 or 10-membered bicyclic heteroaryl ring, which are optionally substituted by CN, halo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -haloalkyl, nitro, hydroxyl, NR^iR^{ii} , $NR^{iii}R^{iv}$, C_{1-6} -alkoxy- C_{1-6} -alkylene, $S(O)_2-C_{1-6}$ -alkyl, C_{1-6} -haloalkoxy, or by a oxo or dioxo bridge; and
- 30 R^5 is H, OH, CN, $COOR^{iii}$ or $CONR^{iii}R^{iv}$;
 R^6 is C_{2-6} -alkyl,

or is $-C(O)-R^f$ wherein R^f is an aryl group substituted by halo, C_{1-6} -alkoxy, or CN,

or is aryl, 5 or 6 membered heteroaryl, or a 9 or 10-membered bicyclic heteroaryl ring which are optionally substituted by halo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -haloalkyl, CN, nitro, NR^iR^{ii} , $NR^{iii}R^{iv}$, C_{1-6} -alkoxy- C_{1-6} -alkylene, COOH, $S(O)_2$ - C_{1-6} -alkyl, hydroxyl, C_{1-6} -haloalkoxy, or by a oxo or dioxo bridge, or is benzyl substituted by halo, C_{1-6} -alkyl, C_{1-6} -haloalkyl or by a oxo – or dioxo bridge;

5
10
15
20
B is halo, CN, NR^iR^{ii} , C_{1-6} -alkyl optionally substituted by CN, halo or C_{1-6} -alkoxy, C_{1-6} -haloalkyl, C_{1-6} -alkoxy, C_{1-6} -haloalkoxy, C_{3-6} -cycloalkyl, $-C(O)O-C_{1-6}$ -alkyl, $-C(O)NR^iR^{ii}$, $-C(O)-C_{1-6}$ -alkyl, $-S(O)_2-C_{1-6}$ -alkyl, $-S(O)_2-NR^iR^{ii}$, $(CR^{iii}R^{iv})_n$ -phenyl, or $(CR^{iii}R^{iv})_n$ -5 or 6 membered heteroaryl wherein the phenyl or 5 or 6 membered heteroaryl moiety is optionally substituted by one or more substituent(s) selected from the group consisting of:

halo, CN, NR^iR^{ii} , C_{1-6} -alkyl optionally substituted by CN or C_{1-6} -alkoxy, C_{1-6} -alkoxy, C_{1-6} -haloalkoxy, C_{3-6} -cycloalkyl, $-C(O)O-C_{1-6}$ -alkyl, $-C(O)-NR^iR^{ii}$, $-C(O)-C_{1-6}$ -alkyl, $-S(O)_2-C_{1-6}$ -alkyl, $-S(O)_2-NR^iR^{ii}$;

R^i and R^{ii} are H, C_{1-6} -alkyl, C_{1-6} -alkyl- $NR^{iii}R^{iv}$, $-(CO)O-C_{1-6}$ -alkyl, $-C(O)-NR^{iii}R^{iv}$, $-C(O)-C_{1-6}$ -alkyl, $-S(O)_2-C_{1-6}$ -alkyl or $-S(O)_2-NR^{iii}R^{iv}$;

20 R^{iii} and R^{iv} are H or C_{1-6} -alkyl;

m is 1 to 6;

n is 0 to 4;

as well as pharmaceutically acceptable salts thereof.

25 The compounds of formula (I) may contain some asymmetric carbon atoms. Accordingly, the present invention includes all stereoisomeric forms of the compounds of formula I, including each of the individual enantiomers and mixtures thereof.

30 It has been found that the compounds of formula (I) have a good activity on the V1a receptor. Therefore, the invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in the manufacture of medicaments for the treatment of dysmenorrhea, hypertension, chronic heart failure, inappropriate secretion of vasopressin, liver cirrhosis, nephrotic syndrome, obsessive compulsive disorder, anxious and depressive disorders.

35 Vasopressin is a 9 amino acid peptide mainly produced by the paraventricular nucleus of the hypothalamus. Three vasopressin receptors, all belonging to the class I G-

protein coupled receptors, are known. The V1a receptor is expressed in the brain, liver, vascular smooth muscle, lung, uterus and testis, the V1b or V3 receptor is expressed in the brain and pituitary gland, the V2 receptor is expressed in the kidney where it regulates water excretion and mediates the antidiuretic effects of vasopressin.

5 In the periphery vasopressin acts as a neurohormone and stimulates vasoconstriction, glycogenolysis and antidiuresis. In the brain vasopressin acts as a neuromodulator and is elevated in the amygdala during stress (Ebner, K., C. T. Wotjak, et al. (2002). "Forced swimming triggers vasopressin release within the amygdala to modulate stress-coping strategies in rats." Eur J Neurosci 15(2): 384-8). The V1a receptor
10 is extensively expressed in the brain and particularly in limbic areas like the amygdala, lateral septum and hippocampus which are playing an important role in the regulation of anxiety. Indeed V1a knock-out mouse show a reduction in anxious behavior in the plus-maze, open field and light-dark box (Bielsky, I. F., S. B. Hu, et al. (2003). "Profound Impairment in Social Recognition and Reduction in Anxiety-Like Behavior in
15 Vasopressin V1a Receptor Knockout Mice." Neuropsychopharmacology). The downregulation of the V1a receptor using antisense oligonucleotide injection in the septum also causes a reduction in anxious behavior (Landgraf, R., R. Gerstberger, et al. (1995). "V1 vasopressin receptor antisense oligodeoxynucleotide into septum reduces vasopressin binding, social discrimination abilities, and anxiety-related behavior in rats."
20 Regul Pept 59(2): 229-39).

The V1a receptor is also mediating the cardiovascular effects of vasopressin in the brain by centrally regulating blood pressure and heart rate in the solitary tract nucleus (Michelini, L. C. and M. Morris (1999). "Endogenous vasopressin modulates the cardiovascular responses to exercise." Ann N Y Acad Sci 897: 198-211). In the periphery it
25 induces the contraction of vascular smooth muscles and chronic inhibition of the V1a receptor improves hemodynamic parameters in myocardial infarcted rats (Van Kerckhoven, R., I. Lankhuizen, et al. (2002). "Chronic vasopressin V(1A) but not V(2) receptor antagonism prevents heart failure in chronically infarcted rats." Eur J Pharmacol 449(1-2): 135-41).

30 Thus vasopressin receptor antagonists are useful as therapeutics in the conditions of dysmenorrhea, hypertension, chronic heart failure, inappropriate secretion of vasopressin, liver cirrhosis, nephrotic syndrome, obsessive compulsive disorder, anxious and depressive disorders.

The preferred indications with regard to the present invention are the treatment of anxious and depressive disorders.

As used herein, the term "aryl" means a monovalent cyclic aromatic hydrocarbon moiety consisting of a mono-, bi- or tricyclic aromatic ring. Examples of aryl moieties include, but are not limited to, optionally substituted phenyl, naphthyl, phenanthryl, fluorenyl, indenyl, pentalenyl, azulenyl, oxydiphenyl, biphenyl, methylenediphenyl, aminodiphenyl, diphenylsulfidyl, diphenylsulfonyl, diphenylisopropylidenyl, benzodioxanyl, benzofuranyl, benzodioxyl, benzopyranyl, benzoxazinyl, benzoxazinonyl, benzopiperadiny, benzopiperazinyl, benzopyrrolidiny, benzomorpholinyl, methylenedioxyphenyl, ethylenedioxyphenyl, as well as those specifically illustrated by the examples herein below. Substituents for aryl include but are not limited to halogen, C₁₋₆-alkyl, C₁₋₆-alkoxy. Preferred aryl are phenyl and naphthyl and still preferably phenyl.

The term "aryloxy" denotes a group wherein the aryl residue is as defined hereinabove, and which is attached via an oxygen atom. The preferred aryloxy group is phenoxy, optionally substituted by halo, e.g. F, as well as those specifically illustrated by the examples herein below.

The term "C₁₋₆-alkyl" denotes a saturated straight- or branched-chain group containing from 1 to 6 carbon atoms, for example, methyl, ethyl, propyl, isopropyl, n-butyl, i-butyl, 2-butyl, t-butyl as well as those specifically illustrated by the examples herein below. Preferred C₁₋₆-alkyl groups are C₁₋₄-groups, i.e. with 1 - 4 carbon atoms.

The term "C₁₋₆-alkoxy" denotes a group wherein the alkyl residues are as defined above, and which is attached via an oxygen atom. Preferred C₁₋₆-alkoxy groups are methoxy and ethoxy as well as those specifically illustrated by the examples herein below.

The term "C₂₋₆-alkenyl" denotes a carbon chain of 2 to 6 carbon atoms comprising a double bond in its chain. C₂₋₆-alkenyl groups include ethenyl, propen-1-yl, propen-2-yl, buten-1-yl, buten-3-yl, penten-1-yl, penten-2-yl, penten-3-yl, penten-4-yl, hexen-1-yl, hexen-2-yl, hexen-3-yl, hexen-4-yl and hexen-5-yl, as well as those specifically illustrated by the examples herein below.

The term "benzyloxy" denotes a benzyl group attached via an oxygen atom.

The term "halogen" or "halo" denotes chlorine (Cl), iodine (I), fluorine (F) and bromine (Br).

The term "C₁₋₆-haloalkyl" denotes a C₁₋₆-alkyl group as defined above which is substituted by one or more halogen. Examples of C₁₋₆-haloalkyl include but are not limited to methyl, ethyl, propyl, isopropyl, isobutyl, sec-butyl, tert-butyl, pentyl or n-hexyl substituted by one or more Cl, F, Br or I atom(s) as well as those groups specifically
5 illustrated by the examples herein below. Preferred C₁₋₆-haloalkyl are difluoro- or trifluoro-methyl or ethyl.

"C₁₋₆-haloalkoxy" denotes a C₁₋₆-alkoxy group as defined above which is substituted by one or more halogen. Examples of C₁₋₆-haloalkoxy include but are not limited to methoxy or ethoxy, substituted by one or more Cl, F, Br or I atom(s) as well as those
10 groups specifically illustrated by the examples herein below. Preferred C₁₋₆-haloalkoxy are difluoro- or trifluoro-methoxy or ethoxy.

The term "C₃₋₆-cycloalkyl" denotes a monovalent or divalent saturated carbocyclic moiety consisting of a monocyclic ring. Cycloalkyl can optionally be substituted with one, two, three or four substituents, wherein each substituent is independently hydroxy, C₁₋₆-
15 alkyl, C₁₋₆-alkoxy, halogen, amino, unless otherwise specifically indicated. Examples of cycloalkyl moieties include optionally substituted cyclopropyl, optionally substituted cyclobutyl, optionally substituted cyclopentyl and optionally substituted cyclohexyl as well as those specifically illustrated by the examples herein below.

The term "4 to 7 membered heterocycloalkyl" means a monovalent saturated
20 moiety, consisting of one ring of 4 to 7 atoms as ring members, including one, two, or three heteroatoms chosen from nitrogen, oxygen or sulfur, the rest being carbon atoms. 4 to 7 membered heterocycloalkyl can optionally be substituted with one, two, three or four substituents, wherein each substituent is independently hydroxy, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-thioalkyl, halo, C₁₋₆-haloalkyl, C₁₋₆-hydroxyalkyl, alkoxycarbonyl, amino, C₁₋₆-
25 alkylamino, di(C₁₋₆)alkylamino, aminocarbonyl, or carbonylamino, unless otherwise specifically indicated. Examples of heterocyclic moieties include, but are not limited to, optionally substituted oxetane, optionally substituted tetrahydro-furanyl, optionally substituted piperidinyl, optionally substituted pyrrolidinyl, optionally substituted morpholinyl, optionally substituted piperazinyl, and the like or those which are
30 specifically exemplified herein. Substituents can be selected from C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-haloalkyl, halo, CN, OH, NH₂, as well as those substituents which are specifically illustrated in the examples hereinafter. Preferred 4 to 7 membered heterocycloalkyl are 5 to 6 membered heterocycloalkyl.


The term "5 or 6 membered heteroaryl" means an aromatic ring of 5 or 6 ring
35 atoms as ring members containing one, two, or three ring heteroatoms selected from N,

O, or S, the rest being carbon atoms. 5 or 6 heteroaryl can optionally be substituted with one, two, three or four substituents, wherein each substituent is independently hydroxy, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-thioalkyl, halo, C₁₋₆-haloalkyl, C₁₋₆-hydroxyalkyl, alkoxy carbonyl, amino, C₁₋₆-alkylamino, di(C₁₋₆)alkylamino, aminocarbonyl, or carbonylamino, unless otherwise specifically indicated. Examples of heteroaryl moieties include, but are not limited to, optionally substituted imidazolyl, optionally substituted oxazolyl, optionally substituted thiazolyl, optionally substituted pyrazinyl, optionally substituted pyrrolyl, optionally substituted pyrazinyl, optionally substituted pyridinyl, optionally substituted pyrimidinyl, optionally substituted furanyl, and those which are specifically exemplified herein.

The term "sulfonylaryl" denotes an aryl group as defined hereinabove which is attached via a sulfonyl group.


The term "9 or 10 membered bicyclic heteroaryl" means an aromatic bicyclic ring of 9 or 10 ring atoms as ring members containing one, two, or three ring heteroatoms selected from N, O, or S, the rest being carbon atoms. 9 or 10 membered bicyclic heteroaryl can optionally be substituted with one, two, three or four substituents, wherein each substituent is independently hydroxy, -C(O), C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-thioalkyl, halo, C₁₋₆-haloalkyl, C₁₋₆-hydroxyalkyl, alkoxy carbonyl, amino, C₁₋₆-alkylamino, di(C₁₋₆)alkylamino, aminocarbonyl, or carbonylamino, unless otherwise specifically indicated. Examples of 9 membered bicyclic heteroaryl moieties include, but are not limited to, optionally substituted indolyl, optionally substituted thieno[2,3-c]pyridinyl, thieno[3,2-c]pyridinyl, benzoxazolyl, benzisoxazolyl as well as those 9 membered bicyclic heteroaryl which are specifically exemplified herein.

The expression "two R² may form an oxo or dioxo bridge together with the indole ring to which they are attached" denotes an oxo or dioxo bridge of the following formulae:




which bind two adjacent carbon atoms of the phenyl or indole ring of the compound of formula (I) to which R² is binding.

Analogously, the expression “oxo or dioxo bridge” denotes an oxo or dioxo bridge of the following formulae:



5 Examples of group illustrating the expression “R¹ and R³ together with the indole ring to which they are attached form a 5 or 6 membered heterocycloalkyl which can be substituted by =O” are:



as well as those specifically illustrated by the examples.


10 The term “pharmaceutically acceptable acid addition salts” embraces salts with inorganic and organic acids, such as hydrochloric acid, nitric acid, sulfuric acid, phosphoric acid, citric acid, formic acid, fumaric acid, maleic acid, acetic acid, succinic acid, tartaric acid, methane-sulfonic acid, p-toluenesulfonic acid, as well as those specifically illustrated by the examples herein below.

15 In certain embodiments of the invention, the compounds of formula (I) are those compounds wherein:

R¹ is H,
 or is C₁₋₆-alkyl optionally substituted by CN or by C₁₋₆-alkoxy,
 or is aryl, 5 or 6 membered heteroaryl or sulfonylaryl which are optionally substituted by one or more B,
 20 or is -(CH₂)_m-R^a wherein R^a is:
 CN,
 ORⁱ,
 NRⁱRⁱⁱ,
 C₃₋₆-cycloalkyl, 4 to 7 membered-heterocycloalkyl, aryl, or 5 or 6 membered
 25 heteroaryl which are optionally substituted by one or more B,
 or is -(CH₂)_n-(CO)-R^b or -(CH₂)_n-(SO₂)-R^b, wherein R^b is:

- C_{1-6} -alkyl,
 C_{1-6} -alkoxy,
 C_{3-6} -cycloalkyl,
 $-(CH_2)_m-NR^{iii}R^{iv}$,
 NR^iR^{ii} ,
 C_{3-6} -cycloalkyl, 4 to 7 membered-heterocycloalkyl, aryl, or 5 or 6 membered heteroaryl which are optionally substituted by one or more B, or R^1 and R^3 together with the indole ring to which they are attached form a 5 or 6 membered heterocycloalkyl which can be substituted by (CO);
 R^2 is one or more of H, OH, halo, CN, nitro, C_{1-6} -alkyl optionally substituted by $NR^{iii}R^{iv}$, C_{1-6} -alkoxy, $-O-CH_2-C_{2-6}$ -alkenyl, benzyloxy, or two R^2 may form an oxo or dioxo bridge together with the indole ring to which they are attached;
 R^3 is H,
or is halo,
or is $-(CO)-R^c$, wherein R^c is:
 C_{1-6} -alkyl,
 $-(CH_2)_n-NR^iR^{ii}$,
 $-(CH_2)_n-NR^{iii}R^{iv}$,
5 or 6 membered heterocycloalkyl optionally substituted by C_{1-6} -alkyl, or is C_{1-6} -alkyl or aryl, which are optionally substituted by halo,
 $-O(CO)-C_{1-6}$ -alkyl,
or by $-NH(CO)R^d$, wherein R^d is C_{1-6} -alkyl optionally substituted by halo or nitro, or R^d is aryl or a 5 or 6 membered heteroaryl, which are optionally substituted by halo, nitro, C_{1-6} -alkyl or C_{1-6} -haloalkyl;

A is selected from the group consisting of (a), (b), (c) and (d):



wherein

- R^4 is $-NH(CO)R^e$, wherein R^e is C_{1-6} -alkoxy, or aryl optionally substituted by halo, or is C_{1-6} -alkoxy, CN,

- or is aryl, 5 or 6 membered heteroaryl, benzyl, aryloxy or a 9 or 10-membered bicyclic heteroaryl ring which are optionally substituted by CN, halo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-haloalkyl, or by a dioxo bridge; and
- R⁵ is H, OH, CN, COORⁱⁱⁱ or CONRⁱⁱⁱR^{iv};
- 5 R⁶ is C₁₋₆-alkyl,
 or is -C(O)-R^f wherein R^f is an aryl group optionally substituted by halo, C₁₋₆-alkoxy, or CN,
 or is aryl, 5 or 6 membered heteroaryl, benzyl, or a 9 or 10-membered bicyclic heteroaryl ring which are optionally substituted by halo, C₁₋₆-alkyl, C₁₋₆-alkoxy,
 10 C₁₋₆-haloalkyl, CN or by a dioxo bridge;
- B is halo, CN, NRⁱRⁱⁱ, C₁₋₆-alkyl optionally substituted by CN, halo or C₁₋₆-alkoxy, C₁₋₆-alkoxy, C₁₋₆-haloalkoxy, C₃₋₆-cycloalkyl, -C(O)O-C₁₋₆-alkyl, -C(O)NRⁱRⁱⁱ, -C(O)-C₁₋₆-alkyl, -S(O)₂-C₁₋₆-alkyl, -S(O)₂-NRⁱRⁱⁱ, (CRⁱⁱⁱR^{iv})_n-phenyl, or
 15 heteroaryl moiety is optionally substituted by one or more substituent(s) selected from the group consisting of:
 halo, CN, NRⁱRⁱⁱ, C₁₋₆-alkyl optionally substituted by CN or C₁₋₆-alkoxy, C₁₋₆-alkoxy, C₁₋₆-haloalkoxy, C₃₋₆-cycloalkyl, -C(O)O-C₁₋₆-alkyl, -C(O)-NRⁱRⁱⁱ, -C(O)-C₁₋₆-alkyl, -S(O)₂-C₁₋₆-alkyl, -S(O)₂-NRⁱRⁱⁱ;
- 20 Rⁱ and Rⁱⁱ are H, C₁₋₆-alkyl, C₁₋₆-alkyl-NRⁱⁱⁱR^{iv}, -(CO)O-C₁₋₆-alkyl, -C(O)-NRⁱⁱⁱR^{iv}, -C(O)-C₁₋₆-alkyl, -S(O)₂-C₁₋₆-alkyl or -S(O)₂-NRⁱⁱⁱR^{iv};
- Rⁱⁱⁱ and R^{iv} are H or C₁₋₆-alkyl;
 m is 1 to 6;
 n is 0 to 4;
- 25 as well as pharmaceutically acceptable salts thereof.

In certain embodiments of the invention, the compounds of formula (I) are those compounds wherein:

- R¹ is H,
 30 or is C₁₋₆-alkyl optionally substituted by CN or by C₁₋₆-alkoxy,
 or is aryl,
 or is 5 or 6 membered heteroaryl,
 or is sulfonylaryl,
 or is -(CH₂)_m-R^a wherein R^a is C₃₋₆-cycloalkyl, 5 or 6 membered-
 35 heterocycloalkyl, aryl, or 5 or 6 membered heteroaryl which are optionally substituted by one or more substituents selected from the group consisting of:

halo, CN, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-haloalkoxy, -C(O)O-C₁₋₆-alkyl and phenyl optionally substituted by halo, C₁₋₆-alkyl, C₁₋₆-haloalkyl or C₁₋₆-alkoxy,

or is -(CH₂)_m-NRⁱRⁱⁱ,

5 or is -(CH₂)_n-(CO)-R^b, wherein R^b is aryl or 5 or 6 membered-heterocycloalkyl;

R² is one or more of H, halo, CN, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, -O-CH₂-C₂₋₆-alkenyl, benzyloxy, or two R² may form an oxo or dioxo bridge together with the indole ring to which they are attached;

R³ is H,

10 or is halo,

or is -(CO)-R^c, wherein R^c is C₁₋₆-alkyl, 5 or 6 membered heterocycloalkyl optionally substituted by C₁₋₆-alkyl, or R^c is -(CH₂)_n-NRⁱRⁱⁱ or,

or is C₁₋₆-alkyl or aryl, which are optionally substituted by

-O(CO)-C₁₋₆-alkyl,


15 or by -NH(CO)R^d, wherein R^d is C₁₋₆-alkyl optionally substituted by halo or nitro, or R^d is aryl or a 5 or 6 membered heteroaryl, which are optionally substituted by halo, nitro, C₁₋₆-alkyl or C₁₋₆-haloalkyl;

Rⁱ and Rⁱⁱ are independently selected from H, C₁₋₆-alkyl or -(CO)O-C₁₋₆-alkyl;

m is 1 to 6;

20 n is 0 to 4;

A is selected from the group consisting of (a), (b), (c) and (d):



wherein

25 R⁴ is is -NH(CO)R^e, wherein R^e is C₁₋₆-alkoxy, or aryl optionally substituted by halo, or is aryl, benzyl, aryloxy or a 9 or 10-membered bicyclic heteroaryl ring which are optionally substituted by halo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-haloalkyl, or by a dioxo bridge; and

R⁵ is H, OH or CN;

R⁶ is C₁₋₆-alkyl,

30 or is -C(O)-R^f wherein R^f is an aryl group optionally substituted by halo,

or is aryl, benzyl, or a 9 or 10-membered bicyclic heteroaryl ring which are optionally substituted by halo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-haloalkyl, or by a dioxo bridge;

as well as pharmaceutically acceptable salts thereof.

5

In certain embodiments of the invention, the compounds of formula (I) are those compounds wherein:

R¹ is H,

or is C₁₋₆-alkyl substituted by NRⁱRⁱⁱ,

10 or is -(CH₂)_m-R^a wherein R^a is:

NRⁱRⁱⁱ,

5 to 6 membered-heterocycloalkyl, aryl, or 5 or 6 membered heteroaryl which are optionally substituted by one or more B,

or is -(CH₂)_n-(CO)-R^b or -(CH₂)_n-(SO₂)-R^b, wherein R^b is:

15 C₁₋₆-alkyl,

C₁₋₆-alkoxy,

NRⁱRⁱⁱ,

5 to 6 membered-heterocycloalkyl, aryl, or 5 or 6 membered heteroaryl which are optionally substituted by one or more B,


20 or R¹ and R³ together with the indole ring to which they are attached form a 5 or 6 membered heterocycloalkyl which can be substituted by =O, C(O)O-C₁₋₆-alkyl or C₁₋₆-alkyl;

R² is one or more of H, halo, C₁₋₆-alkyl;

R³ is H,

25 or is C₁₋₆-alkyl;

A is selected from the group consisting of (a), (a'), (b), (c) and (d):



wherein


30 R⁴ is -NH(CO)R^e, wherein R^e is C₁₋₆-alkoxy, or aryl optionally substituted by halo, or is aryl, 5 or 6 membered heteroaryl, benzyl, aryloxy or a 9 or 10-membered bicyclic heteroaryl ring, which are optionally substituted by halo, C₁₋₆-

alkyl, C₁₋₆-alkoxy, C₁₋₆-haloalkyl, nitro, hydroxyl, C₁₋₆-haloalkoxy, or by a oxo or dioxo bridge; and

- R⁵ is H, OH, CN;
 R⁶ is aryl, 5 or 6 membered heteroaryl, or a 9 or 10-membered bicyclic heteroaryl ring
 5 which are optionally substituted by halo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-haloalkyl, CN, nitro, NRⁱRⁱⁱ, NRⁱⁱⁱR^{iv}, C₁₋₆-alkoxy-C₁₋₆-alkylene, COOH, S(O)₂-C₁₋₆-alkyl, or by a oxo or dioxo bridge;
 B is halo, C₁₋₆-alkyl optionally substituted by CN, halo or C₁₋₆-alkoxy, C₁₋₆-alkoxy;
 Rⁱ and Rⁱⁱ are H, C₁₋₆-alkyl, C₁₋₆-alkyl-NRⁱⁱⁱR^{iv}, -C(O)-C₁₋₆-alkyl or -S(O)₂-C₁₋₆-alkyl;
 10 Rⁱⁱⁱ and R^{iv} are H or C₁₋₆-alkyl;
 m is 1 to 6;
 n is 0 to 4;
 as well as pharmaceutically acceptable salts thereof.



- 15 Preferably, the invention encompasses compounds according to the general formula (I) as described herein, with the proviso that R¹, R² and R³ are not simultaneously H, as well as pharmaceutical acceptable salts thereof.

- 20 Preferably, the invention encompasses compounds according to the general formula (I) as described herein, with the proviso that R⁴ is not an 1H-benzimidazole derivative:



- 25 optionally substituted by CN, halo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-haloalkyl, nitro, hydroxyl, NRⁱRⁱⁱ, NRⁱⁱⁱR^{iv}, C₁₋₆-alkoxy-C₁₋₆-alkylene, S(O)₂-C₁₋₆-alkyl, C₁₋₆-haloalkoxy, or by a oxo or dioxo bridge.

Also encompassed by the compounds of formula (I) are the following compounds of formula (I-a) or (I-a') according to the invention:



wherein R^1 to R^4 are as defined hereinabove for formula (I).

Preferred compounds of formula (I-a) are those compounds wherein,

- R^1 is H or $-(CH_2)_m-R^a$ wherein R^a is aryl and m is 1 to 6;
 5 R^2 is one or more of H or halo;
 R^3 is H or C_{1-6} -alkyl;
 R^4 is aryl which is optionally substituted by halo or C_{1-6} -alkoxy;
 as well as pharmaceutically acceptable salts thereof.

Again, it is preferred that R^1 , R^2 and R^3 are not simultaneously H.

10 Examples of compounds according to the invention are:

(1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-(4-fluoro-phenyl)-3,6-dihydro-2*H*-pyridin-1-yl]-methanone;

(1-Benzyl-2-methyl-1*H*-indol-3-yl)-(4-phenyl-3,6-dihydro-2*H*-pyridin-1-yl)-methanone;

15 (1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-3,6-dihydro-2*H*-pyridin-1-yl]-methanone;

(6-Chloro-1*H*-indol-3-yl)-(4-phenyl-3,6-dihydro-2*H*-pyridin-1-yl)-methanone;

(6-Chloro-1*H*-indol-3-yl)-[4-(4-fluoro-phenyl)-3,6-dihydro-2*H*-pyridin-1-yl]-methanone;


20 (6-Chloro-1*H*-indol-3-yl)-(4-phenyl-3,6-dihydro-2*H*-pyridin-1-yl)-methanone;

(6-Chloro-1*H*-indol-3-yl)-[4-(4-fluoro-phenyl)-3,6-dihydro-2*H*-pyridin-1-yl]-methanone;

(6-Chloro-1*H*-indol-3-yl)-[4-(4-chloro-phenyl)-3,6-dihydro-2*H*-pyridin-1-yl]-methanone; and

25 (1-Benzyl-2-methyl-1*H*-indol-3-yl)-(5-methyl-4-phenyl-3,6-dihydro-2*H*-pyridin-1-yl)-methanone.

Also encompassed by the compounds of formula (I) are the following compounds of formula (I-b) according to the invention:



wherein R^1 to R^5 are as defined hereinabove for formula (I).

5 Preferred compounds of formula (I-b) are those compounds wherein,

R^1 is H,

or is $-(CH_2)_m-R^a$ wherein R^a is:

NR^iR^{ii} ,

aryl, or 5 or 6 membered heteroaryl which are optionally substituted by one

10 or more halo, C_{1-6} -alkyl, C_{1-6} -alkoxy,

or is $-(CH_2)_n-(CO)-R^b$ or $-(CH_2)_n-(SO_2)-R^b$, wherein R^b is:

C_{1-6} -alkyl,

C_{1-6} -alkoxy,

NR^iR^{ii} ,

15 5 to 6 membered-heterocycloalkyl, or 5 or 6 membered heteroaryl which are optionally substituted by one or more halo, C_{1-6} -alkyl, C_{1-6} -alkoxy,

or R^1 and R^3 together with the indole ring to which they are attached form a 5 or

6 membered heterocycloalkyl which can be substituted by $C(O)O-C_{1-6}$ -alkyl or C_{1-6} -alkyl;

20 R^2 is one or more of H, halo or C_{1-6} -alkyl;

R^3 is H or C_{1-6} -alkyl;

R^4 is $-NH(CO)R^e$, wherein R^e is C_{1-6} -alkoxy or aryl which is optionally substituted by halo,

or is aryl, benzyl, aryloxy or a 9 or 10-membered bicyclic heteroaryl ring which

25 are optionally substituted by halo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -haloalkyl, nitro, hydroxyl, C_{1-6} -haloalkoxy or by a oxo or dioxo bridge; and

R^5 is H, OH or CN;

R^i and R^{ii} are H, C_{1-6} -alkyl, C_{1-6} -alkyl- $NR^{iii}R^{iv}$;

Rⁱⁱⁱ and R^{iv} are H or C₁₋₆-alkyl;

m is 1 to 6;

n is 0 to 4;

as well as pharmaceutically acceptable salts thereof.

5 Again, it is preferred that R¹, R² and R³ are not simultaneously H.

Moreover, it is preferred that R⁴ is not an 1H-benzimidazole derivative as described hereinabove.

Examples of compounds of formula (I-b) according to the invention are:

- (4-Benzyl-4-hydroxy-piperidin-1-yl)-(1-benzyl-2-methyl-1*H*-indol-3-yl)-
10 methanone;
(4-Benzyl-4-hydroxy-piperidin-1-yl)-[1-(2-chloro-benzyl)-2-methyl-1*H*-indol-3-yl]-methanone;
(1-Benzyl-5-chloro-2-methyl-1*H*-indol-3-yl)-(4-benzyl-4-hydroxy-piperidin-1-yl)-methanone;
15 (4-Benzyl-4-hydroxy-piperidin-1-yl)-[5-chloro-1-(3-methoxy-benzyl)-2-methyl-1*H*-indol-3-yl]-methanone;
(4-Benzyl-4-hydroxy-piperidin-1-yl)-[1-(2-methoxy-benzyl)-2-methyl-1*H*-indol-3-yl]-methanone;
(1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-(3-fluoro-phenoxy)-piperidin-1-yl]-
20 methanone;
(1-Benzyl-2-methyl-1*H*-indol-3-yl)-(4-hydroxy-4-phenyl-piperidin-1-yl)-methanone;
(1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-(4-fluoro-phenyl)-4-hydroxy-piperidin-1-yl]-methanone;
25 (1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-(4-chloro-3-trifluoromethyl-phenyl)-4-hydroxy-piperidin-1-yl]-methanone;
(4-Benzo[1,3]dioxol-5-yl-4-hydroxy-piperidin-1-yl)-(1-benzyl-2-methyl-1*H*-indol-3-yl)-methanone;
(1-Benzyl-2-methyl-1*H*-indol-3-yl)-(4-hydroxy-4-*p*-tolyl-piperidin-1-yl)-
30 methanone;
(1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-hydroxy-4-(3-trifluoromethyl-phenyl)-piperidin-1-yl]-methanone;
(6-Chloro-1*H*-indol-3-yl)-[4-hydroxy-4-(3-trifluoromethyl-phenyl)-piperidin-1-yl]-methanone;
35 1-(6-Chloro-1*H*-indole-3-carbonyl)-4-phenyl-piperidine-4-carbonitrile;
(6-Chloro-1*H*-indol-3-yl)-(4-hydroxy-4-*p*-tolyl-piperidin-1-yl)-methanone;

- (6-Chloro-1*H*-indol-3-yl)-[4-(4-chloro-phenyl)-4-hydroxy-piperidin-1-yl]-methanone;
- (6-Chloro-1*H*-indol-3-yl)-[4-(4-chloro-3-trifluoromethyl-phenyl)-4-hydroxy-piperidin-1-yl]-methanone;
- 5 [4-(4-Bromo-phenyl)-4-hydroxy-piperidin-1-yl]-(6-chloro-1*H*-indol-3-yl)-methanone;
- [1-(6-Chloro-1*H*-indole-3-carbonyl)-piperidin-4-yl]-carbamic acid *tert*-butyl ester;
- N*-[1-(6-Chloro-1*H*-indole-3-carbonyl)-piperidin-4-yl]-2-fluoro-benzamide;
- 10 (6-Chloro-1*H*-indol-3-yl)-[4-(4-chloro-phenyl)-4-hydroxy-piperidin-1-yl]-methanone;
- (6-Chloro-1*H*-indol-3-yl)-[4-(4-nitro-phenyl)-piperidin-1-yl]-methanone;
- (6-Chloro-1*H*-indol-3-yl)-(4-methoxy-4-phenyl-piperidin-1-yl)-methanone;
- (6-Chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone;
- 15 (6-Chloro-1*H*-indol-3-yl)-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone;
- N*-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide;
- 2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-methylamino-ethyl)-acetamide;
- 20 (6-Chloro-1*H*-indol-3-yl)-[4-(2-trifluoromethoxy-phenyl)-piperidin-1-yl]-methanone;
- (6-Chloro-1*H*-indol-3-yl)-[4-(2-trifluoromethyl-phenyl)-piperidin-1-yl]-methanone;
- 25 2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide;
- 2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide;
- 2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-methylamino-ethyl)-acetamide;
- 30 *N*-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide;
- 2-{6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide;
- 35 2-{6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-methylamino-ethyl)-acetamide or hydrochloride salt thereof;

N-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide or hydrochloride salt thereof;

2-{6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-N-(2-dimethylamino-ethyl)-acetamide;

5 [1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone;

[1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-trifluoromethoxy-phenyl)-piperidin-1-yl]-methanone;

10 [1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone;

[1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperidin-1-yl]-methanone;

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide;

15 (6-Chloro-1*H*-indol-3-yl)-(3,4,5,6-tetrahydro-2*H*-[4,4']bipyridinyl-1-yl)-methanone;

(6-Chloro-2-methyl-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone;

20 (6-Chloro-1-methanesulfonyl-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone;

10-[4-(2-Methoxy-phenyl)-piperidine-1-carbonyl]-3,4-dihydro-1*H*-pyrazino[1,2-*a*]indole-2-carboxylic acid *tert*-butyl ester;

[4-(2-Methoxy-phenyl)-piperidin-1-yl]-(1,2,3,4-tetrahydro-pyrazino[1,2-*a*]indol-10-yl)-methanone or hydrochloride salt thereof; and

25 [4-(2-Methoxy-phenyl)-piperidin-1-yl]-(2-methyl-1,2,3,4-tetrahydro-pyrazino[1,2-*a*]indol-10-yl)-methanone.

Preferred examples of compounds of formula (I-b) according to the invention are:

30 (1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-(4-chloro-phenyl)-4-hydroxy-piperidin-1-yl]-methanone;

(1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-(4-bromo-phenyl)-4-hydroxy-piperidin-1-yl]-methanone;

(6-Chloro-1*H*-indol-3-yl)-(4-phenyl-piperidin-1-yl)-methanone;

35 (4-Benzo[1,3]dioxol-5-yl-4-hydroxy-piperidin-1-yl)-(6-chloro-1*H*-indol-3-yl)-methanone;

(1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-(4-fluoro-phenyl)-piperidin-1-yl]-methanone;

- (6-Chloro-1*H*-indol-3-yl)-[4-(3-chloro-phenyl)-piperidin-1-yl]-methanone;
(6-Chloro-1*H*-indol-3-yl)-[4-(2-hydroxy-phenyl)-piperidin-1-yl]-methanone;
(6-Chloro-1*H*-indol-3-yl)-[4-(4-fluoro-phenyl)-piperidin-1-yl]-methanone;
(6-Chloro-1*H*-indol-3-yl)-[4-(2-isopropoxy-phenyl)-piperidin-1-yl]-methanone;
- 5 2-{3-[4-(2,6-Dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-
piperazin-1-yl)-ethanone;
2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-
(2-dimethylamino-ethyl)-acetamide;
2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-
- 10 piperazin-1-yl)-ethanone;
2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-
yl}-*N*-(2-dimethylamino-ethyl)-acetamide;
2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-
yl}-1-piperazin-1-yl)-ethanone;
- 15 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-5-methyl-indol-1-
yl}-*N,N*-dimethyl-acetamide;
2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-
yl}-acetamide;
2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-
- 20 (2-methylamino-ethyl)-acetamide;
2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-
yl}-1-piperazin-1-yl)-ethanone;
2-{6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-
1-piperazin-1-yl)-ethanone or hydrochloride salt thereof;
- 25 [6-Chloro-1-(2-methylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-
piperidin-1-yl]-methanone;
2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-
dimethylamino-ethyl)-acetamide;
2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-
- 30 methylamino-ethyl)-acetamide or hydrochloride salt thereof;
N-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-
indol-1-yl}-acetamide or hydrochloride salt thereof;
2-{6-Chloro-3-[4-(5-fluoro-benzo[*d*]isoxazol-3-yl)-piperidine-1-carbonyl]-indol-
1-yl}-*N*-methyl-acetamide;
- 35 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-4-hydroxy-piperidine-1-carbonyl]-indol-1-
yl}-*N*-methyl-acetamide; and

2-[6-Chloro-3-(3,4,5,6-tetrahydro-2*H*-[4,4']bipyridinyl-1-carbonyl)-indol-1-yl]-*N*-methyl-acetamide.


Particularly preferred examples of compounds of formula (I-b) according to the
5 invention are:

- (1-Benzyl-2-methyl-1*H*-indol-3-yl)-(4-phenyl-piperidin-1-yl)-methanone;
(6-Chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-
methanone;
2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-
10 yl}-*N,N*-dimethyl-acetamide;
2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-
yl}-1-pyridin-2-yl-ethanone;
(6-Chloro-1-pyridin-4-ylmethyl-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-
piperidin-1-yl]-methanone;
15 (6-Chloro-1-pyridin-3-ylmethyl-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-
piperidin-1-yl]-methanone;
(6-Chloro-1-pyridin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-
piperidin-1-yl]-methanone;
[6-Chloro-1-(6-chloro-pyridin-3-ylmethyl)-1*H*-indol-3-yl]-[4-(2,6-
20 dimethoxy-phenyl)-piperidin-1-yl]-methanone;
{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-
acetic acid *tert*-butyl ester;
(6-Chloro-1-pyrazin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-
phenyl)-piperidin-1-yl]-methanone;
25 [6-Chloro-1-(3,5-difluoro-benzyl)-1*H*-indol-3-yl]-[4-(2,6-dimethoxy-
phenyl)-piperidin-1-yl]-methanone;
[6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2,6-dimethoxy-
phenyl)-piperidin-1-yl]-methanone;
2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-
30 yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
[6-Chloro-1-(3,5-difluoro-benzyl)-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-
piperidin-1-yl]-methanone;
2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-
N,N-dimethyl-acetamide;
35 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-
1-(4-methyl-piperazin-1-yl)-ethanone;

- {6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid *tert*-butyl ester;
- 2-{6-Chloro-3-[4-cyano-4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 5 2-{6-Chloro-3-[4-cyano-4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2-{3-[4-(2,6-Dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- [6-Chloro-1-(2-methyl-pyridin-4-ylmethyl)-1*H*-indol-3-yl]-[4-(2,6-
- 10 dimethoxy-phenyl)-piperidin-1-yl]-methanone;
- 2-{6-Chloro-3-[4-(2-isopropoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2-{6-Chloro-3-[4-(2-isopropoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- 15 2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone;
- 2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- 2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-
- 20 indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- 2-{5,6-Dichloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 25 2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide;
- 2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- N*-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-
- 30 carbonyl]-indol-1-yl}-acetamide;
- [6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone;
- 2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- 35 2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;

- 2- {6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2- {6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- 5 2- {6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- 2- {6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2- {6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- 10 [6-Chloro-1-(2-methylamino-ethyl)-1*H*-indol-3-yl]-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone;
- [6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone;
- 15 2- {6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- 2- {6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- [6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperidin-1-yl]-methanone;
- 20 2- {6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- 2- {6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone or hydrochloride salt thereof; and
- 25 2-[6-Chloro-3-(3,4,5,6-tetrahydro-2*H*-[4,4']bipyridinyl-1-carbonyl)-indol-1-yl]-*N,N*-dimethyl-acetamide.

Also encompassed by the compounds of formula (I) are the following compounds
30 of formula (I-c) according to the invention:



wherein R^1 to R^3 and R^6 are as defined hereinabove for formula (I).

Preferred compounds of formula (I-c) are those compounds wherein,

- R^1 is H or
- 5 or is C_{1-6} -alkyl substituted by NR^iR^{ii} ,
 or is $-(CH_2)_m-R^a$ wherein R^a is:
 NR^iR^{ii} ,
 5 to 6 membered-heterocycloalkyl, aryl, or 5 or 6 membered heteroaryl which
 are optionally substituted by one or more C_{1-6} -alkyl,
- 10 or is $-(CH_2)_n-(CO)-R^b$, wherein R^b is:
 NR^iR^{ii} ,
 5 to 7 membered-heterocycloalkyl which is optionally substituted by one or
 more C_{1-6} -alkyl;
- R^2 is one or more of H, halo or C_{1-6} -alkyl;
- 15 R^3 is H or C_{1-6} -alkyl;
- R^6 is aryl, 5 or 6 membered heteroaryl, or a 9 or 10-membered bicyclic heteroaryl
 ring which are optionally substituted by halo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -
 haloalkyl, CN, nitro, NR^iR^{ii} , $NR^{iii}R^{iv}$, C_{1-6} -alkoxy- C_{1-6} -alkylene, COOH, $S(O)_2$ -
 C_{1-6} -alkyl, or by a oxo or dioxo bridge;
- 20 R^i and R^{ii} are H, C_{1-6} -alkyl, C_{1-6} -alkyl- $NR^{iii}R^{iv}$, $-C(O)-C_{1-6}$ -alkyl, or $-S(O)_2-C_{1-6}$ -alkyl;
 R^{iii} and R^{iv} are H or C_{1-6} -alkyl;
- m is 1 to 6;
- n is 0 to 4;
- as well as pharmaceutically acceptable salts thereof.
- 25 It is preferred that R^1 , R^2 and R^3 are not simultaneously H.

Examples of compounds of formula (I-c) according to the invention are:

(1-Benzyl-2-methyl-1*H*-indol-3-yl)-(4-phenyl-piperazin-1-yl)-methanone;

(1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone;

- (1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-(4-methoxy-phenyl)-piperazin-1-yl]-methanone;
 (1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-(2-chloro-phenyl)-piperazin-1-yl]-methanone;
 (1-Benzyl-2-methyl-1*H*-indol-3-yl)-[4-(4-chloro-phenyl)-piperazin-1-yl]-methanone;
 (6-Chloro-1*H*-indol-3-yl)-(4-phenyl-piperazin-1-yl)-methanone;
 5 (6-Chloro-1*H*-indol-3-yl)-[4-(2-chloro-6-nitro-phenyl)-piperazin-1-yl]-methanone;
 (6-Chloro-1*H*-indol-3-yl)-[4-(2,6-dichloro-phenyl)-piperazin-1-yl]-methanone;
 (6-Chloro-1*H*-indol-3-yl)-[4-(2,3-dihydro-benzo[1,4]dioxin-5-yl)-piperazin-1-yl]-
 methanone;
 (6-Chloro-1*H*-indol-3-yl)-[4-(2-nitro-4-trifluoromethyl-phenyl)-piperazin-1-yl]-
 10 methanone;
 (6-Chloro-1*H*-indol-3-yl)-[4-(2-chloro-phenyl)-piperazin-1-yl]-methanone;
 [4-(2-Amino-6-chloro-phenyl)-piperazin-1-yl]-(6-chloro-1*H*-indol-3-yl)-methanone;
 (6-Chloro-1*H*-indol-3-yl)-[4-(4-methoxy-phenyl)-piperazin-1-yl]-methanone;
 (6-Chloro-1*H*-indol-3-yl)-[4-(3-methoxy-phenyl)-piperazin-1-yl]-methanone;
 15 (6-Chloro-1*H*-indol-3-yl)-[4-(2-nitro-phenyl)-piperazin-1-yl]-methanone;
 (6-Chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone;
 (6-Chloro-1*H*-indol-3-yl)-[4-(4-fluoro-phenyl)-piperazin-1-yl]-methanone;
 (6-Chloro-1*H*-indol-3-yl)-[4-(3-fluoro-phenyl)-piperazin-1-yl]-methanone;
 3-Chloro-4-[4-(6-chloro-1*H*-indole-3-carbonyl)-piperazin-1-yl]-benzonnitrile;
 20 (6-Chloro-1*H*-indol-3-yl)-[4-(2-ethoxy-phenyl)-piperazin-1-yl]-methanone;
 2-{6-Chloro-3-[4-(2-ethoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-(4-methyl-
 piperazin-1-yl)-ethanone;
 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-(2-
 dimethylamino-ethyl)-acetamide;
 25 2-{6-Chloro-3-[4-(2-ethoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-(2-
 dimethylamino-ethyl)-acetamide;
 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetamide;
 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-5-methyl-indol-1-yl}-*N,N*-
 dimethyl-acetamide;
 30 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-piperazin-1-
 yl-ethanone or hydrochloride salt thereof;
 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-(2-
 methylamino-ethyl)-acetamide or hydrochloride salt thereof;
N-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-
 35 1-yl}-acetamide or hydrochloride salt thereof;
 [1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperazin-1-yl]-
 methanone;
 [6-Chloro-1-(2-methylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperazin-1-
 yl]-methanone;

- [6-Chloro-1-(2-methylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-ethoxy-phenyl)-piperazin-1-yl]-methanone;
- 2-[6-Chloro-3-[4-(2-ethoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-(2-methylamino-ethyl)-acetamide;
- 5 [6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-ethoxy-phenyl)-piperazin-1-yl]-methanone;
- (6-Chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- [1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- 10 2-[6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-(2-dimethylamino-ethyl)-acetamide;
- 2-[6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-(2-methylamino-ethyl)-acetamide;
- N*-(2-Amino-ethyl)-2-[6-chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl]-acetamide;
- 15 2-[6-Chloro-3-[4-(2-methoxymethyl-phenyl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-(2-dimethylamino-ethyl)-acetamide;
- 2-[6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-(2-dimethylamino-ethyl)-acetamide;
- 20 2-[6-Chloro-3-(4-pyridin-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N*-(2-dimethylamino-ethyl)-acetamide;
- (6-Chloro-1*H*-indol-3-yl)-[4-(2,4-difluoro-phenyl)-piperazin-1-yl]-methanone;
- (6-Chloro-1*H*-indol-3-yl)-[4-(3-methyl-pyridin-2-yl)-piperazin-1-yl]-methanone;
- (6-Chloro-1*H*-indol-3-yl)-[4-(3,5-dichloro-pyridin-4-yl)-piperazin-1-yl]-methanone;
- 25 (6-Chloro-1*H*-indol-3-yl)-[4-(3-trifluoromethyl-pyridin-2-yl)-piperazin-1-yl]-methanone;
- 2-[4-(6-Chloro-1*H*-indole-3-carbonyl)-piperazin-1-yl]-nicotinonitrile;
- (6-Chloro-1*H*-indol-3-yl)-(4-pyridin-2-yl-piperazin-1-yl)-methanone;
- (6-Chloro-1*H*-indol-3-yl)-(4-thiazol-2-yl-piperazin-1-yl)-methanone;
- 30 2-[6-Chloro-3-(2,3,5,6-tetrahydro-[1,2']bipyrazinyl-4-carbonyl)-indol-1-yl]-*N*-methylacetamide;
- 2-[6-Chloro-3-[4-(3-methyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methylacetamide;
- 2-[6-Chloro-3-[4-(3,5-dichloro-pyridin-4-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methylacetamide;
- 35 2-[6-Chloro-3-(4-pyridin-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N*-methylacetamide;
- 2-[6-Chloro-3-(4-thiazol-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N*-methylacetamide;
- 2-[6-Chloro-3-[4-(3-cyano-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methylacetamide;

- (6-Chloro-1-(S)-1-piperidin-3-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- (6-Chloro-1-(RS)-1-pyrrolidin-3-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- 5 [6-Chloro-1-((S)-1-methyl-piperidin-3-ylmethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- [6-Chloro-1-((RS)-1-methyl-pyrrolidin-3-ylmethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- 10 2-[6-Chloro-3-(2,3,5,6-tetrahydro-[1,2']bipyrazinyl-4-carbonyl)-indol-1-yl]-*N,N*-dimethyl-acetamide;
- 2-{6-Chloro-3-[4-(3-methyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2-{6-Chloro-3-[4-(3,5-dichloro-pyridin-4-yl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 15 2-[6-Chloro-3-(4-pyridin-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N,N*-dimethyl-acetamide;
- 2-[6-Chloro-3-(4-thiazol-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N,N*-dimethyl-acetamide;
- 2-[6-Chloro-3-[4-(3-cyano-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N,N*-dimethyl-acetamide;
- 20 2-[6-Chloro-3-(4-pyrimidin-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N,N*-dimethyl-acetamide;
- (6-Chloro-2-methyl-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone;
- 2-[6-Chloro-3-[4-(6-chloro-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methyl-
- 25 acetamide;
- 2-[6-Chloro-3-[4-(5-trifluoromethyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methyl-acetamide;
- 2-[6-Chloro-3-(4-thieno[2,3-*c*]pyridin-7-yl-piperazine-1-carbonyl)-indol-1-yl]-*N*-methyl-acetamide;
- 30 2-[6-Chloro-3-[4-(3-chloro-5-trifluoromethyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methyl-acetamide;
- 2-[4-(6-Chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carbonyl)-piperazin-1-yl]-nicotinic acid;
- 2-[6-Chloro-3-[4-(2,4-difluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-(2-
- 35 dimethylamino-ethyl)-acetamide; and
- 2-[6-Chloro-3-[4-(4-fluoro-2-methanesulfonyl-phenyl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methyl-acetamide.

40 Preferred examples of compounds of formula (I-c) according to the invention are:

- 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2-{6-Chloro-3-[4-(2-ethoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 5 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- [6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone;
- 10 [6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- 15 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetamide;
- 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- 20 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone;
- 2-{6-Chloro-3-[4-(2,4-difluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- 25 2-{6-Chloro-3-[4-(3-trifluoromethyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- [1-((*S*)-2-Amino-propyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- (6-Chloro-1-(*S*)-1-pyrrolidin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- 30 [6-Chloro-1-((*S*)-1-methyl-pyrrolidin-2-ylmethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- 2-{6-Chloro-3-[4-(2,4-difluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 35 2-{6-Chloro-3-[4-(3-trifluoromethyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;

(6-Chloro-1-(R)-1-pyrrolidin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;

[6-Chloro-1-((R)-1-methyl-pyrrolidin-2-ylmethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;

5 *N*-(2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-ethyl)-acetamide;

N-(2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-ethyl)-methanesulfonamide;


10 *N*-(2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-ethyl)-*N*-methyl-acetamide;

N-(2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-ethyl)-*N*-methyl-methanesulfonamide;

2-[6-Chloro-3-(4-thieno[3,2-*c*]pyridin-4-yl-piperazine-1-carbonyl)-indol-1-yl]-*N*-methyl-acetamide; and

15 2-{6-Chloro-3-[4-(3-iodo-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide.

Also encompassed by the compounds of formula (I) are the following compounds of formula (I-d) according to the invention:



20 wherein R¹ to R³ and R⁶ are as defined hereinabove for formula (I).

Again, it is preferred that R¹, R² and R³ are not simultaneously H.

25 Preferred compounds of formula (I-d) are those compounds wherein the substitution pattern is analogously to that of the preferred compounds of formula (I-c), and further preferred are those compounds wherein

R¹ is H;

R² is one or more halo;
R³ is H;
R⁶ is aryl substituted by C₁₋₆-alkoxy;
as well as pharmaceutically acceptable salts thereof.

- 5 As an example, (6-chloro-1H-indol-3-yl)-[4-(2-methoxy-phenyl)-4-oxy-piperazin-1-yl]-methanone is mentioned.


It may be stated that combinations of the substitution patterns of compounds of formula (I), (I-a), (I-a'), (I-b), (I-c) and (I-d) according to the embodiments as given above are also encompassed by the invention.

- 10 The invention also encompasses the compounds of formulae (I), (I-a), (I-a'), (I-b), (I-c) and (I-d) for a use in the prevention or treatment of dysmenorrhea, hypertension, chronic heart failure, inappropriate secretion of vasopressin, liver cirrhosis, nephrotic syndrome, obsessive compulsive disorder, anxious and depressive disorders.

- The invention also encompasses a pharmaceutical composition comprising a
15 compound of formula (I), (I-a), (I-a'), (I-b), (I-c) or (I-d) which pharmaceutical composition is useful against dysmenorrhea, hypertension, chronic heart failure, inappropriate secretion of vasopressin, liver cirrhosis, nephrotic syndrome, obsessive compulsive disorder, anxious and depressive disorders.


- The invention further encompasses the use of a compound of formula (I), (I-a), (I-
20 a'), (I-b), (I-c) or (I-d) for the preparation of a medicament which is useful against dysmenorrhea, hypertension, chronic heart failure, inappropriate secretion of vasopressin, liver cirrhosis, nephrotic syndrome, obsessive compulsive disorder, anxious and depressive disorders.

- In a certain embodiment, the compounds of formula (I) of the invention can be
25 manufactured according to a process comprising the step of reacting a compound of formula (II):




with a compound of formula A-H, to obtain a compound of formula (I) wherein A, R¹, R² and R³ are as defined hereinabove for formula (I).

In another embodiment, the compounds of formula (I) of the invention can be manufactured according to a process comprising the step of reacting a compound of
5 formula (I-1), wherei R¹ equals H:




with a compound of formula R¹-X, to obtain a compound of formula (I) wherein
wherein A, R¹, R² and R³ are as defined hereinabove for formula (I) and X is halo.

In another embodiment, the compounds of formula (I) of the invention can be
10 manufactured according to a process comprising the step of hydrolyzing a compound of
formula (V):




to obtain a compound of formula (II):



15 and then reacting the compound of formula (II) with a compound of formula A-H, to
obtain a compound of formula (I) wherein R¹, R² and R³ are as defined hereinabove for
formula (I).

These processes are described more in details with the following general schemes and procedures A to C.


General scheme A



General scheme A

- 5 Compounds of formula (I) can be prepared via an amide coupling between an indole 3-carboxylic acid (II) and a compound of formula (A-H), wherein A is defined as hereinabove. The usual reagents and protocols known in the art can be used to effect the amide coupling. Indole 3-carboxylic acids (II) are either commercially available or readily prepared using a procedure described in *J.Med.Chem.* 1991, 34, 140. Alternatively, they
- 10 can be prepared following the general scheme C as described hereinafter. The compounds of formula (A-H) are either commercially available or prepared using methods known in the art starting from commercially available materials. General scheme A is hereinafter further illustrated with general procedure I.


General scheme B



- 15 General procedure B

- Compounds of formula (I) with R¹ different from H can be prepared using methods known in the art, e.g. by N-deprotonation of a compound of formula (I-1) (compounds of formula (I) wherein R¹ is H) followed by treatment with an electrophilic reactant R¹-X (wherein X is a leaving group, e.g. halo) which is either commercially
- 20 available or easily prepared according to methods well known in the art and commercially available starting materials. General scheme B is hereinafter further illustrated with general procedure II.

General scheme C



General procedure C

- 5 The treatment of an indole derivative (III-1) with trifluoroacetic anhydride in DMF affords intermediate (IV) which can be hydrolysed with an aqueous sodium hydroxide solution to give the 3-carboxylic acid indole derivative (II-1). Alternatively, (IV) could react with an electrophilic reactant R¹-X to give (V), which is then converted to the corresponding carboxylic acid derivative (II) with NaH/H₂O in DMF (see *J. Org Chem.*,
- 10 1993, 10, 2862). Intermediate (V) can alternatively be obtained by treatment of an indole derivative (III-2) with trifluoroacetic anhydride in a suitable solvent, e.g. DMF, dichloromethane or 1,2-dichloroethane. Addition of a suitable base may be advantageous.

15

Results – V1a activity

Material & Method:

The human V1a receptor was cloned by RT-PCR from total human liver RNA. The coding sequence was subcloned in an expression vector after sequencing to confirm the identity of the amplified sequence. To demonstrate the affinity of the compounds from

20 the present invention to the human V1a receptor binding studies were performed. Cell

membranes were prepared from HEK293 cells transiently transfected with the expression vector and grown in 20 liter fermenters with the following protocol.

50g of cells are resuspended in 30ml freshly prepared ice cold Lysis buffer (50mM HEPES, 1mM EDTA, 10mM MgCl₂ adjusted to pH= 7.4 + complete cocktail of protease inhibitor (Roche Diagnostics)). Homogenized with Polytron for 1min and sonicated on ice for 2x 2 minutes at 80% intensity (Vibracell sonicator). The preparation is centrifuged 20 min at 500 g at 4°C, the pellet is discarded and the supernatant centrifuged 1hour at 43'000g at 4°C (19'000rpm). The pellet is resuspended in 12.5 ml Lysis buffer+12.5ml Sucrose 20% and homogenized using a Polytron for 1-2 min. The protein concentration is determined by the Bradford method and aliquots are stored at -80°C until use. For binding studies 60mg Yttrium silicate SPA beads (Amersham) are mixed with an aliquot of membrane in binding buffer (50 mM Tris, 120mM NaCl, 5 mM KCl, 2 mM CaCl₂, 10 mM MgCl₂) for 15 minutes with mixing. 50ul of bead/membrane mixture is then added to each well of a 96 well plate, followed by 50ul of 4 nM 3H-Vasopressin (American Radiolabeled Chemicals). For total binding measurement 100ul of binding buffer are added to the respective wells, for non-specific binding 100ul of 8.4mM cold vasopressin and for compound testing 100ul of a serial dilution of each compound in 2%DMSO. The plate is incubated 1h at room temperature, centrifuged 1 min at 1000g and counted on a Packard Top-Count. Non-specific binding counts are subtracted from each well and data is normalized to the maximum specific binding set at 100%. To calculate an IC 50 the curve is fitted using a non-linear regression model (XLfit) and the Ki is calculated using the Cheng-Prussoff equation.

Ex.No.	Ki (nM)	Ex.No.	Ki (nM)	Ex.No.	Ki (nM)
3	14	71	5	153	23
24	10	72	3	162	27
40	8	76	8	163	22
43	2	80	6	171	7
44	3	81	13	172	10
45	7	82	2	174	3
46	7	84	11	175	32
47	4	87	4	176	8
48	7	88	7	177	8
49	3	90	8	190	19
50	4	95	13	196	26
51	0	96	11	198	20
52	4	97	6	200	21
55	2	102	7	203	2
56	2	106	5	205	10
57	2	107	12	211	13
58	9	108	8	214	16
59	2	111	7	215	6
60	8	114	3	217	21
61	13	115	11	218	16
62	11	117	6	219	15
63	2	118	12	220	17
66	4	122	10	224	27
67	5	151	14	225	12
69	6	152	14		

The compounds of formulae (I), (I-a), (I-a'), (I-b), (I-c) and (I-d) as well as their pharmaceutically usable acid addition salts can be used as medicaments, e.g. in the form of pharmaceutical preparations. The pharmaceutical preparations can be administered orally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions. The administration can, however, also be effected rectally, e.g. in the form of suppositories, or parenterally, e.g. in the form of injection solutions.

The compounds of formulae (I), (I-a), (I-a'), (I-b), (I-c) and (I-d) and their pharmaceutically usable acid addition salts can be processed with pharmaceutically inert, inorganic or organic excipients for the production of tablets, coated tablets, dragees and hard gelatine capsules. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts etc can be used as such excipients e.g. for tablets, dragées and hard gelatine capsules.

Suitable excipients for soft gelatine capsules are e.g. vegetable oils, waxes, fats, semi-solid and liquid polyols etc.

Suitable excipients for the manufacture of solutions and syrups are e.g. water, polyols, saccharose, invert sugar, glucose etc.

Suitable excipients for injection solutions are e.g. water, alcohols, polyols, glycerol, vegetable oils etc.

Suitable excipients for suppositories are e.g. natural or hardened oils, waxes, fats, semi-liquid or liquid polyols etc.

- 5 Moreover, the pharmaceutical preparations can contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They can also contain still other therapeutically valuable substances.

10 The dosage can vary within wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, in the case of oral administration a daily dosage of about 10 to 1000 mg per person of a compound of general formula (I), (I-a), (I-b); (I-c) or (I-d) should be appropriate, although the above upper limit can also be exceeded when necessary.

15 The following Examples illustrate the present invention without limiting it. All temperatures are given in degrees Celsius.

Example A

Tablets of the following composition are manufactured in the usual manner:

	<u>mg/tablet</u>
Active substance	5
20 Lactose	45
Corn starch	15
Microcrystalline cellulose	34
Magnesium stearate	1
Tablet weight	100

Example B

Capsules of the following composition are manufactured:

	<u>mg/capsule</u>
Active substance	10
5 Lactose	155
Corn starch	30
Talc	5
Capsule fill weight	200

The active substance, lactose and corn starch are firstly mixed in a mixer and then
10 in a comminuting machine. The mixture is returned to the mixer, the talc is added
thereto and mixed thoroughly. The mixture is filled by machine into hard gelatine
capsules.

Example C

Suppositories of the following composition are manufactured:

	<u>mg/supp.</u>
Active substance	15
Suppository mass	1285
Total	1300

The suppository mass is melted in a glass or steel vessel, mixed thoroughly and
20 cooled to 45°C. Thereupon, the finely powdered active substance is added thereto and
stirred until it has dispersed completely. The mixture is poured into suppository moulds
of suitable size, left to cool; the suppositories are then removed from the moulds and
packed individually in wax paper or metal foil.

EXAMPLES

25 General procedure I: Amide coupling:

To a stirred solution of an indole-3-carboxylic acid derivative (1 mmol) in 10 ml CH₂Cl₂
were added (1.3 mmol) EDC, (1.3 mmol) HOBt, (1.3 mmol) Et₃N and (1 mmol) of the
amine derivative. The mixture was stirred overnight at RT and then poured onto water
and extracted with CH₂Cl₂. The combined organic phases were dried over Na₂SO₄ and

concentrated in vacuo. Flash chromatography or preparative HPLC afforded the title compound.

General procedure II: Alkylation:

To a stirred solution of (6-chloro-1*H*-indol-3-yl)-(4-phenyl-piperidin-1-yl)-methanone
5 in DMF were added 2.1eq. NaH (60% in oil). The mixture was stirred at RT for 30 min.
and then the electrophilic reagent R¹-X (1.1eq.) was added. The mixture was stirred an
additional 14 hours at 60 °C and then poured onto water and extracted with ethyl acetate.
The combined organic phases were dried over Na₂SO₄ and concentrated in vacuo.
Purification by preparative HPLC afforded the corresponding derivatives.

10

Acid intermediates of formula II and II-1

Acid 1

6-Chloro-1*H*-indole-3-carboxylic acid

Using a procedure described in *J. Med. Chem.* 1991, 34, 140, from 7.0 g (0.046 mmol) of
15 6-chloro-1*H*-indole were prepared 5.80 g (64%) of 6-chloro-1*H*-indole-3-carboxylic acid
as a light brown solid.

ES-MS m/e (%): 194 (M-H⁺).

Acid 2

6-Chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid

20 a) 1-(6-Chloro-1*H*-indol-3-yl)-2,2,2-trifluoro-ethanone

Using a procedure described in *J. Med. Chem.* 1991, 34, 140, from 0.50 g (0.004 mol) of 6-
chloro-1*H*-indole were prepared 0.76 g (95%) of 1-(6-chloro-1*H*-indol-3-yl)-2,2,2-
trifluoro-ethanone as a white solid.

b) 2-[6-Chloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-*N,N*-dimethyl-acetamide

25 To a stirred solution of 1-(6-chloro-1*H*-indol-3-yl)-2,2,2-trifluoro-ethanone (0.75 g) in
20 ml of DMF at 0 °C, were added 128 mg (1.1 eq.) of NaH (60% in oil). The mixture was
stirred for 30 min. and then 0.32 ml (1.1 eq.) of dimethylamino-acetyl chloride were
added. The mixture was stirred an additional hour and then poured onto water and
extracted with ethyl acetate. The combined organic phases were dried over Na₂SO₄ and
30 concentrated in vacuo to afford 598 mg (61%) of 2-[6-chloro-3-(2,2,2-trifluoro-acetyl)-
indol-1-yl]-*N,N*-dimethyl-acetamide as a white solid.

c) 6-Chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid

Using a procedure similar to that described in *J. Med. Chem.* 1991, 34, 140, from 0.50 g of
2-[6-chloro-5-methyl-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-*N,N*-dimethyl-acetamide
35 were prepared 0.38 g (76%) of 6-chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-
carboxylic acid as a white solid

Acid 3

6-Chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acida) 2-[6-Chloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-*N*-methyl-acetamide

Following general procedure II, the alkylation of 1-(6-chloro-1*H*-indol-3-yl)-2,2,2-trifluoro-ethanone, with (commercially available) 2-chloro-*N*-methyl-acetamide gave the title compound.

ES-MS *m/e* (%): 319.3 (M+H⁺).

b) 6-Chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid

2-[6-Chloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-*N*-methyl-acetamide was suspended in DCE and treated with (2.2 eq.) of sodium trimethylsilanolate. After shaking at room temperature for 20 min, the mixture was concentrated in vacuo and purified by prep. HPLC to give the title compound in 27% yield.

ES-MS *m/e* (%): 265.0 (M-H⁺).

Acid 4

6-Chloro-1-[(2-dimethylamino-ethylcarbamoyl)-methyl]-1*H*-indole-3-carboxylic acid

a) [6-Chloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-acetic acid methyl ester

To a stirred solution of 6.65 g (0.0265 mol, 1eq.) of 1-(6-chloro-1*H*-indol-3-yl)-2,2,2-trifluoro-ethanone in 25 ml of DMF at 0 °C were added 1.23 g of NaH (55% in oil, 0.0282 mol, 1.05 eq.). After 30 minutes, 4.31 g (0.0282 mol, 1.05 eq.) of bromo-acetic acid methyl ester were added and the temperature raised to RT, and stirring was continued overnight. The reaction was quenched by the addition of aq.HCl 0.1 M, and the product was extracted with EtOAc and the combined organic phases dried over Na₂SO₄.

Recrystallisation in Et₂O / Heptane afforded 6.90 g (80%) of [6-chloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-acetic acid methyl ester as white crystals.

b) [6-Chloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-acetic acid

To a stirred solution of 2 g (6.3 mmol) of [6-chloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-acetic acid methyl ester in 45 ml of THF / MeOH / H₂O 1/1/1, at 40°C, were added 0.79 g (18.9 mmol, 3 eq.) of LiOH.H₂O. After 1 hour, the reaction mixture was diluted in EtOAc, acidified with aq.HCl 1M. The organic phase were dried over Na₂SO₄, and concentrated under vacuo to afford 1.9 g (99%) of [6-chloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-acetic acid as a white solid.

c) 2-[6-Chloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-*N*-(2-dimethylamino-ethyl)-acetamide

An amide coupling between 0.99 g of [6-Chloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-acetic and *N*,*N*-dimethyl-ethane-1,2-diamine according to the general procedure I, afforded 78 mg (6%) of 2-[6-chloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-*N*-(2-dimethylamino-ethyl)-acetamide as a white solid.

d) 6-Chloro-1-[(2-dimethylamino-ethylcarbamoyl)-methyl]-1H-indole-3-carboxylic acid

Using a similar procedure described in *J. Med. Chem.* 1991, 34, 140, from 78 mg of of 2-[6-chloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-N-(2-dimethylamino-ethyl)-acetamide were prepared 65 mg (96%) of 6-chloro-1-[(2-dimethylamino-ethylcarbamoyl)-methyl]-1H-indole-3-carboxylic acid as a white solid.

Acid 5

6-Chloro-1-dimethylcarbamoylmethyl-5-methyl-1H-indole-3-carboxylic acid

a) 1-(6-Chloro-5methyl-1H-indol-3-yl)-2,2,2-trifluoro-ethanone

Using a procedure described in *J. Med. Chem.* 1991, 34, 140, from 0.250 g (0.002 mol) of 6-chloro-5-methyl-1H-indole were prepared 0.38 g (96%) of 1-(6-chloro-5methyl-1H-indol-3-yl)-2,2,2-trifluoro-ethanone as a white solid.

b) 2-[6-Chloro-5-methyl-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-N,N-dimethyl-acetamide

To a stirred solution of 1-(6-chloro-5methyl-1H-indol-3-yl)-2,2,2-trifluoro-ethanone (0.38 g) in 10 ml of DMF at 0 °C, were added 64 mg (1.1 eq.) of NaH (60% in oil). The mixture was stirred for 30 min. and then 0.16 ml (1.1 eq.) of dimethylamino-acetyl chloride were added. The mixture was stirred an additional hour and then poured onto water and extracted with ethyl acetate. The combined organic phases were dried over Na₂SO₄ and concentrated in vacuo to afford 300 mg (60%) of 2-[6-chloro-5-methyl-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-N,N-dimethyl-acetamide as a white solid.

c) 6-Chloro-1-dimethylcarbamoylmethyl-5-methyl-1H-indole-3-carboxylic acid

Using a similar procedure described in *J. Med. Chem.* 1991, 34, 140, from 0.280 g of 2-[6-chloro-5-methyl-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-N,N-dimethyl-acetamide were prepared 0.18 g (76%) of 6-chloro-1-dimethylcarbamoylmethyl-5-methyl-1H-indole-3-carboxylic acid as a white solid

Acid 6

5,6-Dichloro-1-dimethylcarbamoylmethyl-1H-indole-3-carboxylic acid

a) 1-(5,6-Dichloro-1H-indol-3-yl)-2,2,2-trifluoro-ethanone

Using a procedure described in *J. Med. Chem.* 1991, 34, 140, from 0.120 g (0.64 mmol) of 5,6-dichloro-1H-indole were prepared 0.11 g (59%) of 1-(5,6-dichloro-1H-indol-3-yl)-2,2,2-trifluoro-ethanone as a white solid.

b) 2-[5,6-Dichloro-3-(2,2,2-trifluoro-acetyl)-indol-1-yl]-N,N-dimethyl-acetamide

To a stirred solution of 1-(5,6-dichloro-1H-indol-3-yl)-2,2,2-trifluoro-ethanone (0.11 g) in 3 ml of DMF at 0 °C, were added 18 mg (1.05 eq.) of NaH (60% in oil). The mixture was stirred for 30 min. and then 0.04 ml (1.0 eq.) of dimethylamino-acetyl chloride were added. The mixture was stirred an additional hour and then poured onto water and extracted with ethyl acetate. The combined organic phases were dried over Na₂SO₄ and

concentrated in vacuo to afford 112 mg (78%) of 2-[5,6-dichloro-3-(2,2,2-trifluoroacetyl)-indol-1-yl]-N,N-dimethyl-acetamide as a white solid.

c) 5,6-Dichloro-1-dimethylcarbamoylmethyl-1H-indole-3-carboxylic acid

- Using a similar procedure described in *J. Med. Chem.* 1991, 34, 140, from 0.112 g of 2-[5,6-dichloro-3-(2,2,2-trifluoroacetyl)-indol-1-yl]-N,N-dimethyl-acetamide were prepared 0.047 g (49%) of 5,6-dichloro-1-dimethylcarbamoylmethyl-1H-indole-3-carboxylic acid as a white solid.

Acid 7

6-Chloro-2-methyl-1H-indole-3-carboxylic acid

10 a) (6-Chloro-1H-indol-2-yl)-methanol

To a solution of 2.00 g (8.94 mmol) 6-chlorindole-2-carboxylic acid ethyl ester in 50 ml diethyl ether were added 0.475 g (12.5 mmol) lithium aluminum hydride at 0 °C. The reaction mixture was heated at reflux for 45 min and quenched by consecutive addition of 10 ml water, 10 ml aqueous 2 M sodium hydroxide solution and 10 ml water at 0 °C.

- 15 The aqueous layer was extracted with tert-butyl methyl ether (3 x 100 ml). The combined organic layers were dried over sodium sulfate and concentrated in vacuo to give the crude title compound (1.64 g; 100%) as a white solid.

MS m/e (%): 180 (M-H⁺, 100).

b) 6-Chloro-2-methyl-1H-indole

- 20 A solution of 1.60 g (8.81 mmol) (6-chloro-1H-indol-2-yl)-methanol in 5 ml 1,2-dichloroethane was added to a mixture of 80.0 ml trifluoroacetic acid and 32.0 ml triethylsilane at 65 °C. After 5 min, the reaction mixture was cooled to room temperature and quenched with water. The pH was adjusted to 14 by the addition of aqueous sodium hydroxide solution (32 %). The aqueous layer was extracted with tert-butyl methyl ether (3 x 200 ml). The combined organic layers were dried over sodium sulfate and concentrated in vacuo. The residue was purified by flash-chromatography (aminopropyl-modified silica gel, n-heptane / ethyl acetate) to give the title compound (0.39 g; 27%) as a white solid.

MS m/e (%): 164 (M-H⁺, 100).

30 c) 1-(6-Chloro-2-methyl-1H-indol-3-yl)-2,2,2-trifluoro-ethanone

- To a solution of 0.38 g (2.3 mmol) 6-chloro-2-methyl-1H-indole in 20 ml 1,2-dichloroethane at 0 °C were added 0.35 ml (2.5 mmol) trifluoroacetic anhydride. The reaction mixture was quenched with aqueous 2 M sodium carbonate solution after 30 min and extracted with dichloromethane (3 x 100 ml). The combined organic layers were dried over sodium sulfate and concentrated in vacuo to give the title compound (0.57 g; 95%) as an off-white solid.

MS m/e (%): 260 (M-H⁺, 100).

d) 6-Chloro-2-methyl-1H-indole-3-carboxylic acid

A solution of 0.57 g (2.2 mmol) 1-(6-chloro-2-methyl-1H-indol-3-yl)-2,2,2-trifluoroethanone in 21.7 ml (86.8 mmol) aqueous 4 M sodium hydroxide solution was heated at reflux for 45 min. After cooling to room temperature the reaction mixture was diluted with water and extracted with tert-butyl methyl ether (2 x 50 ml). The aqueous layer was cooled to 0-5 °C, acidified (pH 1-2) with concentrated aqueous hydrochloric acid solution and extracted with ethyl acetate (3 x 100 ml). The combined ethyl acetate layers were dried over sodium sulfate and concentrated in vacuo to give the title compound (0.14 g, 31%) as an off-white solid.

MS m/e (%): 208 (M-H⁺, 100).

Acid 8

5-Chloro-2-methyl-1H-indole-3-carboxylic acid

Using a procedure described in *J. Med. Chem.* 1991, 34, 140, from the 5-chloro-2-methyl-1H-indole was prepared 5-chloro-2-methyl-1H-indole-3-carboxylic acid.

Acid 9

1-Benzyl-2-methyl-1H-indole-3-carboxylic acid

To a stirred solution of 0.50 g (3.1 mmol) 2-methyl-1H-indole-3-carboxylic acid (described in *J.Heterocyclic Chem.* 1977, 14, 1123) in 5 ml DMF were added 0.27 g (6.75 mmol) of NaH (60% in oil). The mixture was stirred at RT for 30 min. and then 0.39 ml (3.28 mmol) of benzyl bromid were added. The mixture was stirred an additional hour and then poured onto water and extracted with ethyl acetate. The combined organic phases were dried over Na₂SO₄ and concentrated in vacuo. Crystallization in Et₂O afforded 0.61 g (78%) of 1-benzyl-2-methyl-1H-indole-3-carboxylic acid as a white solid.

Acid 10

3,4-Dihydro-1H-pyrazino[1,2-a]indole-2,10-dicarboxylic acid 2-tert-butyl ester
a) 10-(2,2,2-Trifluoro-acetyl)-3,4-dihydro-1H-pyrazino[1,2-a]indole-2-carboxylic acid tert-butyl ester

To a stirred solution of 0.21 ml (1.5 mmol) trifluoroacetic anhydride in 7 ml 1,2-dichloroethane was added at 0 °C a solution of 0.37 g (1.4 mmol) 3,4-dihydro-1H-pyrazino[1,2-a]indole-2-carboxylic acid tert-butyl ester and a solution of 0.23 ml (1.63 mmol) triethylamine in 3 ml 1,2-dichloroethane. After stirring for 30 min the reaction mixture was quenched with saturated aqueous sodium hydrogen carbonate solution and extracted with dichloromethane (2 x 100 ml). The combined organic layers were dried over sodium sulfate, concentrated in vacuo and purified by flash chromatography (n-heptane / ethyl acetate) to give the title compound (0.288 g, 58%) as a light yellow solid.

MS m/e (%): 369 (M+H⁺, 27).


b) 3,4-Dihydro-1H-pyrazino[1,2-a]indole-2,10-dicarboxylic acid 2-tert-butyl ester

To a solution of 0.29 g (0.77 mmol) 10-(2,2,2-trifluoro-acetyl)-3,4-dihydro-1H-pyrazino[1,2-a]indole-2-carboxylic acid tert-butyl ester in 7 ml N,N-dimethylformamide were subsequently added 0.22 g (4.6 mmol) sodium hydride (50% in oil) and a solution of 0.070 ml (3.9 mmol) water in 1 ml N,N-dimethylformamide at room temperature. The reaction mixture was diluted with tert-butyl methyl ether after 2 h and extracted with 1 M sodium hydroxide solution (2 x 30 ml). The combined aqueous layers were acidified (pH 1-2) with 2 M hydrochloric acid at 0 °C and extracted with tert-butyl methyl ether (3 x 50 ml). The combined organic layers were dried over sodium sulfate and concentrated in vacuo to give the title compound (0.21 g, 86%) as a light brown solid. MS m/e (%): 315 (M-H⁺, 100).

Examples of compounds of formula I-a and I-a'

Example 1

(1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(4-fluoro-phenyl)-3,6-dihydro-2H-pyridin-1-yl]-methanone




Amide coupling according to general procedure I:

- Amine: 4-(4-Fluoro-phenyl)-1,2,3,6-tetrahydro-pyridine (commercially available),
 - Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,
- ES-MS m/e (%): 425.5 (M+H⁺).

Example 2

(1-Benzyl-2-methyl-1H-indol-3-yl)-(4-phenyl-3,6-dihydro-2H-pyridin-1-yl)-methanone

- 43 -




Amide coupling according to general procedure I:

- Amine: 4-Phenyl-1,2,3,6-tetrahydro-pyridine (commercially available),
- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,

5 ES-MS m/e (%): 407.5 (M+H⁺).

Example 3

(1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(2-methoxy-phenyl)-3,6-dihydro-2H-pyridin-1-yl]-methanone




10 Amide coupling according to general procedure I:

- Amine: 4-(2-Methoxy-phenyl)-1,2,3,6-tetrahydro-pyridine (described in US6326381),
- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,

ES-MS m/e (%): 437.6 (M+H⁺).

Example 4

15 (6-Chloro-1H-indol-3-yl)-(4-phenyl-3,6-dihydro-2H-pyridin-1-yl)-methanone



Amide coupling according to general procedure I:

- 44 -


- Amine: 4-Phenyl-1,2,3,6-tetrahydro-pyridine (commercially available),

- Acid: 6-chloro-1H-indole-3-carboxylic acid,

ES-MS m/e (%): 337.4 (M+H⁺).

Example 5

5 (6-Chloro-1H-indol-3-yl)-[4-(4-fluoro-phenyl)-3,6-dihydro-2H-pyridin-1-yl]-methanone



Amide coupling according to general procedure I:


- Amine: 4-(4-Fluoro-phenyl)-1,2,3,6-tetrahydro-pyridine (commercially available),

10 - Acid: 6-chloro-1H-indole-3-carboxylic acid,

ES-MS m/e (%): 355.4 (M+H⁺).

Example 6

(6-Chloro-1H-indol-3-yl)-(4-phenyl-3,6-dihydro-2H-pyridin-1-yl)-methanone



15 Following general procedure I, the coupling of (commercially available) 4-phenyl-1,2,3,6-tetrahydro-pyridine, with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.


ES-MS m/e (%): 337.4 (M+H⁺).

Example 7

(6-Chloro-1H-indol-3-yl)-[4-(4-fluoro-phenyl)-3,6-dihydro-2H-pyridin-1-yl]-methanone

20

- 45 -




Following general procedure I, the coupling of (commercially available) 4-(4-fluorophenyl)-1,2,3,6-tetrahydro-pyridine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

5 ES-MS m/e (%): 355.4 (M+H⁺).

Example 8

(6-Chloro-1H-indol-3-yl)-[4-(4-chloro-phenyl)-3,6-dihydro-2H-pyridin-1-yl]-methanone




10 Following general procedure I, the coupling of (commercially available) 4-(4-chlorophenyl)-1,2,3,6-tetrahydro-pyridine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 371.4 (M+H⁺).

Example 9

15 (1-Benzyl-2-methyl-1H-indol-3-yl)-(5-methyl-4-phenyl-3,6-dihydro-2H-pyridin-1-yl)-methanone




Following general procedure I, the coupling of (commercially available) 5-methyl-4-phenyl-1,2,3,6-tetrahydro-pyridine (described in WO 2005077912) with 1-benzyl-2-methyl-1*H*-indole-3-carboxylic acid gave the title compound.

- 5 ES-MS *m/e* (%): 421.6 ($M+H^+$).

Examples of compounds of formula I-b

Example 10


(4-Benzyl-4-hydroxy-piperidin-1-yl)-(1-benzyl-2-methyl-1*H*-indol-3-yl)-methanone



- 10 Amide coupling according to general procedure I:
 - Amine: 4-Benzyl-piperidin-4-ol (commercially available),
 - Acid: 1-Benzyl-2-methyl-1*H*-indole-3-carboxylic acid,
 ES-MS *m/e* (%): 439.5 ($M+H^+$).

Example 11

- 15 (4-Benzyl-4-hydroxy-piperidin-1-yl)-[1-(2-chloro-benzyl)-2-methyl-1*H*-indol-3-yl]-methanone



a) (4-Benzyl-4-hydroxy-piperidin-1-yl)-(2-methyl-1H-indol-3-yl)-methanone

To a stirred solution of 0.1 g (0.57 mmol) 2-methyl-1H-indole-3-carboxylic acid in 10 ml CH_2Cl_2 were added 0.14 g (0.73 mmol) EDC, 0.10 g (0.73 mmol) HOBt, 90 μl (0.63 mmol) Et_3N and 0.11 g (0.57 mmol) 4-benzyl-piperidin-4-ol. The mixture was stirred overnight at RT and then poured onto water and extracted with CH_2Cl_2 . The combined organic phases were dried over Na_2SO_4 and concentrated in vacuo. Flash chromatography (ethyl acetate) afforded 0.15 g (75%) of (4-benzyl-4-hydroxy-piperidin-1-yl)-(2-methyl-1H-indol-3-yl)-methanone as a white solid.

ES-MS m/e (%): 349 ($\text{M}+\text{H}^+$).


b) (4-Benzyl-4-hydroxy-piperidin-1-yl)-[1-(2-chloro-benzyl)-2-methyl-1H-indol-3-yl]-methanone

To a stirred solution of 40 mg (0.11 mmol) of (4-benzyl-4-hydroxy-piperidin-1-yl)-(2-methyl-1H-indol-3-yl)-methanone in 5 ml DMF were added 5 mg (0.11 mmol) NaH (60% in oil). The mixture was stirred at RT for 30 min. and then 28 mg (0.13 mmol) of 1-bromomethyl-2-chloro-benzene were added. The mixture was stirred an additional hour and then poured onto water and extracted with ethyl acetate. The combined organic phases were dried over Na_2SO_4 and concentrated in vacuo. Preparative HPLC (30% $\text{CH}_3\text{CN} / \text{H}_2\text{O}$) afforded 33 mg (61%) of (4-benzyl-4-hydroxy-piperidin-1-yl)-[1-(2-chloro-benzyl)-2-methyl-1H-indol-3-yl]-methanone as a white solid.

ES-MS m/e (%): 473.4 ($\text{M}+\text{H}^+$).

Example 12

(1-Benzyl-5-chloro-2-methyl-1H-indol-3-yl)-(4-benzyl-4-hydroxy-piperidin-1-yl)-methanone



- 5 a) (4-benzyl-4-hydroxy-piperidin-1-yl)-(5-chloro-2-methyl-1H-indol-3-yl)-methanone
 Using the procedure described for the preparation of (4-benzyl-4-hydroxy-piperidin-1-yl)-(2-methyl-1H-indol-3-yl)-methanone, from 40 mg (0.19 mmol) of 5-chloro-2-methyl-1H-indole-3-carboxylic acid and 40 mg (0.21 mmol) of 4-benzyl-piperidin-4-ol were prepared 45 mg (62%) of (4-benzyl-4-hydroxy-piperidin-1-yl)-(5-chloro-2-methyl-1H-indol-3-yl)-methanone as a white solid.

ES-MS m/e (%): 383 (M+H⁺).

b) (1-benzyl-5-chloro-2-methyl-1H-indol-3-yl)-(4-benzyl-4-hydroxy-piperidin-1-yl)-methanone


- Using the procedure described for the preparation of (4-benzyl-4-hydroxy-piperidin-1-yl)-[1-(2-chloro-benzyl)-2-methyl-1H-indol-3-yl]-methanone, from 13 mg (0.034 mmol) of (4-benzyl-4-hydroxy-piperidin-1-yl)-(5-chloro-2-methyl-1H-indol-3-yl)-methanone and 5.8 mg (0.034 mmol) of benzyl bromide were prepared 14 mg (87%) of (1-benzyl-5-chloro-2-methyl-1H-indol-3-yl)-(4-benzyl-4-hydroxy-piperidin-1-yl)-methanone as a white solid.

- 20 ES-MS m/e (%): 473.3 (M+H⁺).

Example 13

(4-Benzyl-4-hydroxy-piperidin-1-yl)-[5-chloro-1-(3-methoxy-benzyl)-2-methyl-1H-indol-3-yl]-methanone

- 49 -




Using the procedure described for the preparation of (4-benzyl-4-hydroxy-piperidin-1-yl)-[1-(2-chloro-benzyl)-2-methyl-1H-indol-3-yl]-methanone, from 13 mg (0.034 mmol) of (4-benzyl-4-hydroxy-piperidin-1-yl)-(5-chloro-2-methyl-1H-indol-3-yl)-methanone and 6.8 mg (0.034 mmol) of 1-bromomethyl-3-methoxy-benzene were prepared 11 mg (65%) of (4-benzyl-4-hydroxy-piperidin-1-yl)-[5-chloro-1-(3-methoxy-benzyl)-2-methyl-1H-indol-3-yl]-methanone as a white solid.

ES-MS m/e (%): 503.5 ($M+H^+$).

Example 14

10 (4-Benzyl-4-hydroxy-piperidin-1-yl)-[1-(2-methoxy-benzyl)-2-methyl-1H-indol-3-yl]-methanone




Using the procedure described for the preparation of (4-benzyl-4-hydroxy-piperidin-1-yl)-[1-(2-chloro-benzyl)-2-methyl-1H-indol-3-yl]-methanone, from 28 mg (0.080 mmol) of (4-benzyl-4-hydroxy-piperidin-1-yl)-(2-methyl-1H-indol-3-yl)-methanone and 15 mg (0.096 mmol) of 1-chloromethyl-2-methoxy-benzene were prepared 29 mg (77%) of (4-benzyl-4-hydroxy-piperidin-1-yl)-[1-(2-methoxy-benzyl)-2-methyl-1H-indol-3-yl]-methanone as a white solid.

ES-MS m/e (%): 469.6 (M+H⁺).

Example 15

(1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(3-fluoro-phenoxy)-piperidin-1-yl]-methanone



5 Amide coupling according to general procedure I:


- Amine: 4-(3-Fluoro-phenoxy)-piperidine (described in US260294700),

- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,

ES-MS m/e (%): 443.6 (M+H⁺).

Example 16

10 (1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(4-chloro-phenyl)-4-hydroxy-piperidin-1-yl]-methanone



Amide coupling according to general procedure I:

- Amine: 4-(4-Chloro-phenyl)-piperidin-4-ol (commercially available),


15 - Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid

ES-MS m/e (%): 459.5 (M+H⁺).

Example 17

(1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(4-bromo-phenyl)-4-hydroxy-piperidin-1-yl]-methanone

- 51 -



Amide coupling according to general procedure I:


- Amine: 4-(4-Bromo-phenyl)-piperidin-4-ol (commercially available),

- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,

5 ES-MS m/e (%): 505.5 (M+H⁺).

Example 18

(1-Benzyl-2-methyl-1H-indol-3-yl)-(4-hydroxy-4-phenyl-piperidin-1-yl)-methanone



Amide coupling according to general procedure I:

10 - Amine: 4-Phenyl-piperidin-4-ol (commercially available),

- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,


ES-MS m/e (%): 425.6 (M+H⁺).

Example 19

(1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(4-fluoro-phenyl)-4-hydroxy-piperidin-1-yl]-

15

methanone



Amide coupling according to general procedure I:


- Amine: 4-(4-Fluoro-phenyl)-piperidin-4-ol (commercially available),

- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,

20 ES-MS m/e (%): 443.6 (M+H⁺).

Example 20

(1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(4-chloro-3-trifluoromethyl-phenyl)-4-hydroxy-piperidin-1-yl]-methanone



5 Amide coupling according to general procedure I:

- Amine: 4-(4-Chloro-3-trifluoromethyl-phenyl)-piperidin-4-ol (commercially available),


- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid

ES-MS m/e (%): 527.5 (M+H⁺).

10

Example 21

(4-Benzo[1,3]dioxol-5-yl-4-hydroxy-piperidin-1-yl)-(1-benzyl-2-methyl-1H-indol-3-yl)-methanone



Amide coupling according to general procedure I:

15 - Amine: 4-Benzo[1,3]dioxol-5-yl-piperidin-4-ol (described in WO9709311),


- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,

ES-MS m/e (%): 469.6 (M+H⁺).

Example 22

(1-Benzyl-2-methyl-1H-indol-3-yl)-(4-hydroxy-4-p-tolyl-piperidin-1-yl)-methanone

- 53 -



Amide coupling according to general procedure I:


- Amine: 4-p-Tolyl-piperidin-4-ol (described in EP445701),

- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,

5 ES-MS m/e (%): 439.6 (M+H⁺).

Example 23

(1-Benzyl-2-methyl-1H-indol-3-yl)-[4-hydroxy-4-(3-trifluoromethyl-phenyl)-piperidin-1-yl]-methanone



10 Amide coupling according to general procedure I:


- Amine: 4-(3-Trifluoromethyl-phenyl)-piperidin-4-ol (commercially available),

- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,

ES-MS m/e (%): 493.6 (M+H⁺).

Example 24

15 (1-Benzyl-2-methyl-1H-indol-3-yl)-(4-phenyl-piperidin-1-yl)-methanone



Amide coupling according to general procedure I:


- Amine: 4-Phenyl-piperidine (commercially available),

- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid

ES-MS m/e (%): 409.5 (M+H⁺).

Example 25


(6-Chloro-1H-indol-3-yl)-(4-phenyl-piperidin-1-yl)-methanone



- 5 Amide coupling according to general procedure I:
- Amine: 4-Phenyl-piperidine (commercially available),
- Acid: 6-chloro-1H-indole-3-carboxylic acid,
ES-MS m/e (%): 339.4 (M+H⁺).

Example 26


- 10 (6-Chloro-1H-indol-3-yl)-[4-hydroxy-4-(3-trifluoromethyl-phenyl)-piperidin-1-yl]-
methanol



- Amide coupling according to general procedure I:
- Amine: 4-(3-Trifluoromethyl-phenyl)-piperidin-4-ol (commercially available),
15 - Acid: 6-chloro-1H-indole-3-carboxylic acid,
ES-MS m/e (%): 423.4 (M+H⁺).

Example 27

1-(6-Chloro-1H-indole-3-carbonyl)-4-phenyl-piperidine-4-carbonitrile




Amide coupling according to general procedure I:

- Amine: 4-Phenyl-piperidine-4-carbonitrile (commercially available),
- Acid: 6-chloro-1H-indole-3-carboxylic acid,

5 ES-MS m/e (%): 364.4 (M+H⁺).

Example 28

(6-Chloro-1H-indol-3-yl)-(4-hydroxy-4-p-tolyl-piperidin-1-yl)-methanone




Amide coupling according to general procedure I:

- 10
- Amine: 4-p-Tolyl-piperidin-4-ol (described in EP445701),
 - Acid: 6-chloro-1H-indole-3-carboxylic acid

ES-MS m/e (%): 369.4 (M+H⁺).

Example 29

(6-Chloro-1H-indol-3-yl)-[4-(4-chloro-phenyl)-4-hydroxy-piperidin-1-yl]-methanone




Amide coupling according to general procedure I:

- Amine: 4-(4-Chloro-phenyl)-piperidin-4-ol (commercially available),
- Acid: 6-chloro-1H-indole-3-carboxylic acid,

5 ES-MS m/e (%): 389.4 (M^+).

Example 30

(6-Chloro-1H-indol-3-yl)-[4-(4-chloro-3-trifluoromethyl-phenyl)-4-hydroxy-piperidin-1-yl]-methanone



10 Amide coupling according to general procedure I:


- Amine: 4-(4-Chloro-3-trifluoromethyl-phenyl)-piperidin-4-ol (commercially available),
- Acid: 6-chloro-1H-indole-3-carboxylic acid,

ES-MS m/e (%): 457.4 (M^+).

15

Example 31

(4-Benzo[1,3]dioxol-5-yl-4-hydroxy-piperidin-1-yl)-(6-chloro-1H-indol-3-yl)-methanone



Amide coupling according to general procedure I:


- Amine: 4-Benzo[1,3]dioxol-5-yl-piperidin-4-ol (described in WO9709311),

- Acid: 6-chloro-1H-indole-3-carboxylic acid,

5 ES-MS m/e (%): 399.4 (M+H⁺).

Example 32

[4-(4-Bromo-phenyl)-4-hydroxy-piperidin-1-yl]-(6-chloro-1H-indol-3-yl)-methanone



Amide coupling according to general procedure I:


10 - Amine: 4-(4-Bromo-phenyl)-piperidin-4-ol (commercially available),

- Acid: 6-chloro-1H-indole-3-carboxylic acid,

ES-MS m/e (%): 435.3 (M⁺).

Example 33

[1-(6-chloro-1H-indole-3-carbonyl)-piperidin-4-yl]-carbamic acid tert-butyl ester




Amide coupling according to general procedure I:

- 5 - Amine: Piperidin-4-yl-carbamic acid tert-butyl ester (commercially available),
 - Acid: 6-chloro-1H-indole-3-carboxylic acid,
 ES-MS m/e (%):.378.3 (M+H⁺).

Example 34

N-[1-(6-Chloro-1H-indole-3-carbonyl)-piperidin-4-yl]-2-fluoro-benzamide



10

a) (4-Amino-piperidin-1-yl)-(6-chloro-1H-indol-3-yl)-methanone

- To a stirred solution of 0.50 g (1.32 mmol) of [1-(6-chloro-1H-indole-3-carbonyl)-piperidin-4-yl]-carbamic acid tert-butyl ester (the preparation of which has been described in example 31) in CH₂Cl₂ (10 ml), were added 2 ml of TFA. The mixture was stirred 2 hours, then poured onto an aqueous solution of sat. NaHCO₃ and then extracted with ethyl acetate. The combined organic phases were dried over Na₂SO₄ and concentrated in vacuo, to afford 0.25 g (68%) of (4-amino-piperidin-1-yl)-(6-chloro-1H-indol-3-yl)-methanone as a light brown solid.

ES-MS m/e (%): 78.1 (M+H⁺).20 b) N-[1-(6-Chloro-1H-indole-3-carbonyl)-piperidin-4-yl]-2-fluoro-benzamide


To a stirred solution of 50 mg (0.36 mmol) 2-fluoro-benzoic acid in 5 ml CH₂Cl₂ were added 90 mg (0.47 mmol) EDC, 63 mg (0.47 mmol) HOBt, 60 μl (0.47 mmol) Et₃N and

100 mg (0.36 mmol) (4-amino-piperidin-1-yl)-(6-chloro-1H-indol-3-yl)-methanone. The mixture was stirred overnight at RT and then poured onto water and extracted with CH_2Cl_2 . The combined organic phases were dried over Na_2SO_4 and concentrated in vacuo. Flash chromatography (ethyl acetate) afforded 82 mg (57%) N-[1-(6-Chloro-1H-
5 indole-3-carbonyl)-piperidin-4-yl]-2-fluoro-benzamide of as a white solid.

ES-MS m/e (%): 400.3 ($\text{M}+\text{H}^+$).

Example 35

1-[1-(1-Cyclohexylmethyl-1H-indole-3-carbonyl)-piperidin-4-yl]-1,3-dihydro-
benzoimidazol-2-one



10


Following general procedure I, the coupling of (commercially available) 4-(4-fluoro-phenyl)-piperidine with 1-benzyl-2-methyl-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 427.6 ($\text{M}+\text{H}^+$).

15

Example 36

(6-Chloro-1H-indol-3-yl)-[4-(4-chloro-phenyl)-4-hydroxy-piperidin-1-yl]-methanone




Following general procedure I, the coupling of (commercially available) 4-(4-chloro-phenyl)-piperidin-4-ol with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 389.4 ($M+H^+$).

5

Example 37

(6-Chloro-1H-indol-3-yl)-[4-(4-nitro-phenyl)-piperidin-1-yl]-methanone




Following general procedure I, the coupling of (commercially available) 4-(4-nitro-phenyl)-piperidine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

10 ES-MS m/e (%): 384.4 ($M+H^+$).

Example 38

(6-Chloro-1H-indol-3-yl)-[4-(3-chloro-phenyl)-piperidin-1-yl]-methanone




15 Following general procedure I, the coupling of (commercially available) 4-(3-chloro-phenyl)-piperidine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 373.3 ($M+H^+$).

Example 39

(6-Chloro-1H-indol-3-yl)-[4-(2-hydroxy-phenyl)-piperidin-1-yl]-methanone

- 61 -




Following general procedure I, the coupling of (commercially available) 2-piperidin-4-yl-phenol with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 355.4 ($M+H^+$).

5

Example 40

(6-Chloro-1H-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone



Following general procedure I, the coupling of (commercially available) 4-(2,6-
10 dimethoxy-phenyl)-piperidine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.


ES-MS m/e (%): 399.2 ($M+H^+$).

Example 41

15

(6-Chloro-1H-indol-3-yl)-[4-(4-fluoro-phenyl)-piperidin-1-yl]-methanone

- 62 -




Following general procedure I, the coupling of (commercially available) 4-(4-fluorophenyl)-piperidine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 357.4 ($M+H^+$).

5

Example 42

(6-Chloro-1H-indol-3-yl)-(4-methoxy-4-phenyl-piperidin-1-yl)-methanone



Following general procedure I, the coupling of 4-methoxy-4-phenyl-piperidine (described in WO 9800400) with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.


10

ES-MS m/e (%): 369.4 ($M+H^+$).

Example 43

2-[6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl]-*N,N*-dimethyl-acetamide

- 63 -




Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with commercially available 2-chloro-*N,N*-dimethyl-acetamide gave the title compound.

5 ES-MS *m/e* (%): 484.3 ($M+H^+$).

Example 44


2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-pyridin-2-yl-ethanone



10 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with commercially available 2-chloro-1-pyridin-2-yl-ethanone gave the title compound.
ES-MS *m/e* (%): 518.4 ($M+H^+$).

Example 45

15 (6-Chloro-1-pyridin-4-ylmethyl-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone




Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with commercially available 4-chloromethyl-pyridine gave the title compound.

ES-MS *m/e* (%): 490.3 (M+H⁺).

5

Example 46

(6-Chloro-1-pyridin-3-ylmethyl-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone



Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with commercially available 3-chloromethyl-pyridine gave the title compound.


10

ES-MS *m/e* (%): 490.3 (M+H⁺).

Example 47

(6-Chloro-1-pyridin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone

15




Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with methanesulfonic acid pyridin-2-ylmethyl ester (described in WO 9955318) gave the title compound.

20

ES-MS *m/e* (%): 490.3 (M+H⁺).

Example 48

[6-Chloro-1-(6-chloro-pyridin-3-ylmethyl)-1*H*-indol-3-yl]-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone




- 5 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with methanesulfonic acid 6-chloro-pyridin-3-ylmethyl ester (described in Journal of Organic Chemistry (1999), 64(23), 8576-8581) gave the title compound.
ES-MS *m/e* (%): 524.2 (M+H⁺).

10

Example 49

{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid *tert*-butyl ester




- 15 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) bromo-acetic acid *tert*-butyl ester gave the title compound.
ES-MS *m/e* (%): 513.3 (M+H⁺).

Example 50

6-Chloro-1-pyrazin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone

20

- 66 -




Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with methanesulfonic acid pyrazin-2-ylmethyl ester (described in WO 2002064574) gave the

5 title compound.

ES-MS *m/e* (%): 491.1 (M+H⁺).

Example 51

[6-Chloro-1-(3,5-difluoro-benzyl)-1*H*-indol-3-yl]-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone



10

Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 1-chloromethyl-3,5-difluoro-benzene gave the title compound.


ES-MS *m/e* (%): 525.3 (M+H⁺).

15

Example 52

[6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone

- 67 -




Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) (2-chloro-ethyl)-dimethyl-amine gave the title compound.

5 ES-MS *m/e* (%): 470.5 ($M+H^+$).

Example 53

(6-Chloro-1*H*-indol-3-yl)-[4-(2-isopropoxy-phenyl)-piperidin-1-yl]-methanone




Following general procedure I, the coupling of 4-(2-isopropoxy-phenyl)-piperidine
10 (described in Journal of Medicinal Chemistry (1998), 41(12), 1997-2009) with 6-chloro-1*H*-indole-3-carboxylic acid gave the title compound.

ES-MS *m/e* (%): 397.1 ($M+H^+$).

Example 54

(6-Chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone




Following general procedure I, the coupling of (commercially available) 4-(2-methoxy-phenyl)-piperidine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 367.0 ($M+H^+$).

Example 55

- 5 2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone




- 10 Following general procedure II, the alkylation of (6-chloro-1H-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-1-(4-methyl-piperazin-1-yl)-ethanone gave the title compound.

ES-MS m/e (%): 539.6 ($M+H^+$).

Example 56

- 15 [6-Chloro-1-(3,5-difluoro-benzyl)-1H-indol-3-yl]-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone




Following general procedure II, the alkylation of (6-chloro-1H-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 1-chloromethyl-3,5-difluoro-benzene gave the title compound.

ES-MS m/e (%): 495.2 (M+H⁺).

Example 57

2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide




5

Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-*N,N*-dimethyl-acetamide gave the title compound. ES-MS m/e (%): 454.3 (M+H⁺).

10

Example 58

2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone



15

Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-1-(4-methyl-piperazin-1-yl)-ethanone gave the title compound.


ES-MS m/e (%): 509.3 (M+H⁺).

20

- 70 -

Example 59


{6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid *tert*-butyl ester



- 5 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) bromo-acetic acid *tert*-butyl ester gave the title compound. ES-MS *m/e* (%): 483.3 (M+H⁺).

Example 60

- 10 2-{6-Chloro-3-[4-cyano-4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-N,N-dimethyl-acetamide




Amide coupling according to general procedure I:

- Amine: 4-(2-Methoxy-phenyl)-piperidine-4-carbonitrile (described in
15 WO2003053361),
- Acid: 6-Chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid,
ES-MS *m/e* (%): 479.1 (M+H⁺).

Example 61

- 20 2-{6-Chloro-3-[4-cyano-4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-N,N-dimethyl-acetamide

- 71 -




Amide coupling according to general procedure I:

- Amine: 4-(2-Fluoro-phenyl)-piperidine-4-carbonitrile (described in *Tetrahedron* 2004, 4875),

- 5 - Acid: 6-Chloro-1-dimethylcarbamoylmethyl-1H-indole-3-carboxylic acid, ES-MS m/e (%): 467.1 (M+H⁺).

Example 62

2-{3-[4-(2,6-Dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-N,N-dimethyl-acetamide



10


Following general procedure II, the alkylation of (6-chloro-1H-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-N,N-dimethyl-acetamide gave the title compound. ES-MS m/e (%): 450.6 (M+H⁺).

15

Example 63

[6-Chloro-1-(2-methyl-pyridin-4-ylmethyl)-1H-indol-3-yl]-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone


- 72 -



Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with methanesulfonic acid 2-methyl-pyridin-4-ylmethyl ester (prepared by mesylation of the
 5 commercially available (2-methyl-pyridin-4-yl)-methanol) gave the title compound.
 ES-MS *m/e* (%): 504.1 (M+H⁺).

Example 64

2-{3-[4-(2,6-Dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone



10

Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-1-(4-methyl-piperazin-1-yl)-ethanone gave the title compound.

15 ES-MS *m/e* (%): 505.4 (M+H⁺).

Example 65


2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide

a) {6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid:
 20

Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) bromo-acetic acid gave the title compound.

ES-MS *m/e* (%): 457.5 (M+H⁺).

- 5 b) 2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide




- 10 Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid with (commercially available) *N*¹,*N*¹-dimethyl-ethane-1,2-diamine gave the title compound.

ES-MS *m/e* (%): 527.3 (M+H⁺).

Example 66

- 2-{6-Chloro-3-[4-(2-isopropoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide



15


Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-isopropoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-*N,N*-dimethyl-acetamide gave the title compound.

ES-MS *m/e* (%): 482.6 (M+H⁺).

20

Example 67

2-{6-Chloro-3-[4-(2-isopropoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone



- 5 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-isopropoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-1-(4-methyl-piperazin-1-yl)-ethanone gave the title compound.

ES-MS *m/e* (%): 537.6 (M+H⁺).

10

Example 68


2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone

a) 6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid

- 15 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) bromo-acetic acid gave the title compound.

ES-MS *m/e* (%): 427.5 (M+H⁺).

b) 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone




20

Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid with (commercially available) piperazine-1-carboxylic acid *tert*-butyl ester gave, after treatment with TFA and neutralisation, the title compound.

- 5 ES-MS *m/e* (%): 495.5 (M+H⁺).

Example 69

2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone



- 10 Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with (commercially available) piperazine-1-carboxylic acid *tert*-butyl ester gave, after treatment with TFA and neutralisation, the title compound.

ES-MS *m/e* (%): 525.5 (M+H⁺).

- 15

Example 70

(6-Chloro-1*H*-indol-3-yl)-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone
 a) 4-(5-Bromo-2,3-dihydro-benzofuran-7-yl)-pyridine

- 4-Bromopyridine or hydrochloride salt thereof was dissolved in DME under argon and Pd(Ph₃P)₄ (3mol%) was added. The mixture was stirred for 10 min at 50 °C. To this
 20 solution was added (commercially available) 5-bromo-2,3-dihydrobenzo[*b*]furan-7-boronic acid dissolved in a minimum amount of EtOH/DME 1:2 followed by 2N Na₂CO₃. The reaction mixture was refluxed (110 °C) under stirring for 2h. The flask was cooled to RT, the mixture was treated with sat. aq. NH₄Cl solution and extracted with CHCl₃. Evaporation and purification by SiO₂ gel chromatography (hexane/ethyl acetate:
 25 4/1) gave the title compound in 76% yield.

ES-MS *m/e* (%): 275.9 (M+H⁺).

b) 1-Benzyl-4-(5-bromo-2,3-dihydro-benzofuran-7-yl)-1,2,3,6-tetrahydro-pyridine

To a solution of 4-(5-bromo-2,3-dihydro-benzofuran-7-yl)-pyridine in toluene was added benzyl bromide (1eq.) and the reaction mixture was stirred under reflux for 14h. Reaction was monitored by TLC (CH₂Cl₂ / MeOH 95:5) and revealed total conversion to
5 an intermediate. Complete evaporation of the solvent gave a white solid. The solid was dissolved in MeOH under argon and cooled down to 0 °C, NaBH₄ (2.05 eq.) was added portion-wise (exothermic reaction) and the reaction mixture was stirred for 3h at RT. Evaporation of MeOH, redissolution in CH₂Cl₂ and sequential washing with 1N NaHCO₃ and brine, followed by purification by SiO₂ gel chromatography (CH₂Cl₂/MeOH: 98/2)
10 gave 71% yield of the title compound.


ES-MS m/e (%): 370.0 (M+H⁺).

c) 4-(2,3-Dihydro-benzofuran-7-yl)-piperidine

To a solution of 1-benzyl-4-(5-bromo-2,3-dihydro-benzofuran-7-yl)-1,2,3,6-tetrahydro-pyridine in EtOH and 4N HCl was added 40wt% of 10% Pd/C catalyst. The reaction flask
15 was charged with H₂ (3.5 bar) and then stirred overnight at 50 °C. The reaction mixture was filtered over Celite under argon. The solvents were evaporated and the product was partitioned between CH₂Cl₂ and aq. K₂CO₃. The organic phase was dried over Na₂SO₄ and evaporated to give the title compound in 81% yield.

ES-MS m/e (%): 204.3 (M+H⁺).

20 d) (6-Chloro-1H-indol-3-yl)-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone




Following general procedure I, the coupling of 4-(2,3-dihydro-benzofuran-7-yl)-piperidine (described herein below) with 6-chloro-1H-indole-3-carboxylic acid gave the
25 title compound.

ES-MS m/e (%): 381.1 (M+H⁺).

Example 71

2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone




- 5 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-1-(4-methyl-piperazin-1-yl)-ethanone gave the title compound.

ES-MS *m/e* (%): 521.6 (M+H⁺).

10

Example 72

2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide



- 15 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-*N,N*-dimethyl-acetamide gave the title compound.

ES-MS *m/e* (%): 466.6 (M+H⁺).

Example 73


- 20 2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide

a) {6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid

Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) bromo-acetic acid gave the title compound.

ES-MS *m/e* (%): 437.1 (M-H⁺).

- 5 b) 2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid with (commercially available) *N*¹,*N*¹-dimethyl-ethane-1,2-diamine gave the title compound.

ES-MS *m/e* (%): 509.6 (M+H⁺).

Example 74

- 2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone




15 Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared described herein) with (commercially available) piperazine-1-carboxylic acid *tert*-butyl ester gave, after treatment with TFA and neutralisation, the title compound.

20 ES-MS *m/e* (%): 507.0 (M+H⁺).

Example 75


2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-5-methyl-indol-1-yl}-
N,N-dimethyl-acetamide



- 5 Amide coupling according to general procedure I:
 - Amine: 4-(2-Methoxy-phenyl)-piperidine (commercially available),
 - Acid: 6-Chloro-1-dimethylcarbamoylmethyl-5-methyl-1H-indole-3-carboxylic acid,
 ES-MS m/e (%): 468.3 (M+H⁺).

Example 76


- 10 2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-N-
methyl-acetamide



- Following general procedure II, the alkylation of (6-chloro-1H-indol-3-yl)-[4-(2,3-
dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone (preparation described herein),
 15 with (commercially available) 2-chloro-N-methyl-acetamide gave the title compound.
 ES-MS m/e (%): 451.7 (M+H⁺).

Example 77

N-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-
carbonyl]-indol-1-yl}-acetamide




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with (commercially available) (2-amino-ethyl)-carbamic acid *tert*-butyl ester gave, after
5 treatment with HCl and neutralisation, the title compound.

ES-MS m/e (%): 481.3 ($M+H^+$).

Example 78

2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide



10

Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with (commercially available) ammonia in THF, gave the title compound.


ES-MS m/e (%): 438.2 ($M+H^+$).

15

Example 79

2-{6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-methylamino-ethyl)-acetamide

- 81 -




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2,3-dihydro-
benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with
(commercially available) (2-amino-ethyl)-methyl-carbamic acid tert-butyl ester gave,
5 after treatment with TFA and neutralisation, the title compound.

ES-MS m/e (%): 495.5 (M+H⁺).

Example 80

2-{5,6-Dichloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-N,N-
dimethyl-acetamide



10

Amide coupling according to general procedure I:

- Amine: 4-(2-Methoxy-phenyl)-piperidine (commercially available),
- Acid: 5,6-Dichloro-1-dimethylcarbamoylmethyl-1H-indole-3-carboxylic acid,


ES-MS m/e (%): 488.5 (M+H⁺).

15

Example 81

2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide

- 82 -




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with (commercially available) ammonia in THF gave the title compound.

- 5 ES-MS m/e (%): 456.2 ($M+H^+$).

Example 82

2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide




- 10 Analogous to general procedure II, the alkylation of (6-chloro-1H-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone (prepared herein) with (commercially available) 2-chloro-*N*-methyl-acetamide gave the title compound.

ES-MS m/e (%): 470.2 ($M+H^+$).

Example 83

- 15 2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-methylamino-ethyl)-acetamide

- 83 -




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with
 (commercially available) (2-amino-ethyl)-methyl-carbamic acid tert-butyl ester gave,
 5 after treatment with TFA and neutralisation, the title compound.

ES-MS m/e (%): 513.5 ($M+H^+$).

Example 84

N-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-
 indol-1-yl}-acetamide



10

Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with
 (commercially available) (2-amino-ethyl)-carbamic acid tert-butyl ester gave, after
 treatment with HCl and neutralisation, the title compound.

15 ES-MS m/e (%): 499.5 ($M+H^+$).

Example 85

(6-Chloro-1*H*-indol-3-yl)-[4-(2-trifluoromethoxy-phenyl)-piperidin-1-yl]-methanone
 a) 4-(2-Trifluoromethoxy-phenyl)-pyridine:

To a solution of (commercially available) (1-bromo-2-trifluoromethoxy-benzene) in DME (first degased with argon) under argon was added 3mol% Pd(PPh₃)₄, pyridyl-4-boronic acid and 2N Na₂CO₃, the resulting reaction mixture was vigorously stirred under reflux. After 5h at reflux, reaction as monitored by TLC showed completion. Two phases were separated, evaporation of most of the DME, redissolution of the residue in EtOAc and washing with aq. NaOH followed by evaporation gave a yellow oil. Flash Chromatography on SiO₂ gel with a mixture of CH₂Cl₂/MeOH gave the title compound in 51% yield.

ES-MS m/e (%): 239.9 (M+H⁺).

10 b) 1-Benzyl-4-(2-trifluoromethoxy-phenyl)-1,2,3,6-tetrahydro-pyridine

To a solution of 4-(2-trifluoromethoxy-phenyl)-pyridine in toluene was added (1 eq.) benzyl bromide, and reaction mixture was stirred under reflux for 2h. Reaction was monitored by TLC (CH₂Cl₂/MeOH 95:5) and revealed total conversion to the first intermediate. Complete evaporation of solvent gave a white solid which was pure enough to carry out the next step. The white solid was dissolved in MeOH under argon and cooled down to 0 °C, (4eq.) NaBH₄ was added portion-wise (exothermic reaction). The reaction mixture was then stirred for 3h at RT, reaction again monitored by TLC (CH₂Cl₂/MeOH 95:5) and revealed total conversion. Evaporation of MeOH, redissolution in CH₂Cl₂, and washing with 1N NaHCO₃ then brine, and evaporation, followed by SiO₂ gel chromatography (CH₂Cl₂/MeOH) gave the title compound in 61% yield.

ES-MS m/e (%): 334.3 (M+H⁺).


c) 4-(2-Trifluoromethoxy-phenyl)-piperidine

To a solution of 1-benzyl-4-(2-trifluoromethoxy-phenyl)-1,2,3,6-tetrahydro-pyridine (described herein below) in EtOH was added (0.2wt%) 10%Pd/C, followed by (5eq.) TFA in a sealed tube. The reaction mixture was stirred at 50 °C under 3.0bar of H₂ for 12h. The reaction mixture was filtered over Celite and the filtrate evaporated down to dryness. Redissolution in EtOAc, washing with 1N NaHCO₃ and concentration gave the crude product which was directly used for the next step.

30 ES-MS m/e (%): 246.6 (M+H⁺).

d) (6-Chloro-1H-indol-3-yl)-[4-(2-trifluoromethoxy-phenyl)-piperidin-1-yl]-methanone

- 85 -




Following general procedure I, the coupling of 4-(2-trifluoromethoxy-phenyl)-piperidine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 423.3 ($M+H^+$).

5

Example 86

(6-Chloro-1H-indol-3-yl)-[4-(2-trifluoromethyl-phenyl)-piperidin-1-yl]-methanone




Following general procedure I, the coupling of (commercially available) 4-(2-trifluoromethyl-phenyl)-piperidine (described herein) with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 407.1 ($M+H^+$).

Example 87

[6-Chloro-1-(2-dimethylamino-ethyl)-1H-indol-3-yl]-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone



15


Analogous to general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone (prepared herein) with (commercially available) (2-chloro-ethyl)-dimethyl-amine gave the title compound.

ES-MS *m/e* (%): 452.2 (M+H⁺).

5

Example 88

2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone



Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-trifluoromethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-1-(4-methyl-piperazin-1-yl)-ethanone gave the title compound.

ES-MS *m/e* (%): 563.3 (M+H⁺).

Example 89

2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide


a) {6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid

Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-trifluoromethoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) bromo-acetic acid gave the title compound.

ES-MS *m/e* (%): 481.3 (M+H⁺).

b) 2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide

- 87 -




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with (commercially available) N^1,N^1 -dimethyl-ethane-1,2-diamine gave the title compound.

5 ES-MS m/e (%): 551.5 ($M+H^+$).

Example 90

2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}- N -methyl-acetamide




10 Analogous to general procedure I, the coupling of (6-chloro-1*H*-indol-3-yl)-[4-(2-trifluoromethoxy-phenyl)-piperidin-1-yl]-methanone (prepared herein) with (commercially available) 2-chloro- N -methyl-acetamide gave the title compound.

ES-MS m/e (%): 494.5 ($M+H^+$).

Example 91

15 2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide

- 88 -




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with (commercially available) ammonia in THF, gave the title compound.

5 ES-MS m/e (%): 480.3 ($M+H^+$).

Example 92

2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone



10 Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared described herein) with (commercially available) piperazine-1-carboxylic acid *tert*-butyl ester gave, after treatment with TFA and neutralisation, the title compound.


ES-MS m/e (%): 548.7 ($M+H^+$).

15

Example 93

2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-methylamino-ethyl)-acetamide

- 89 -




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared described herein) with (commercially available) (2-amino-ethyl)-methyl-carbamic acid tert-butyl ester gave, after treatment with TFA and neutralisation, the title compound.

ES-MS m/e (%): 536.7 ($M+H^+$).

Example 94

N-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide



10


Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared described herein) with (commercially available) (2-amino-ethyl)-carbamic acid tert-butyl ester gave, after treatment with HCl and neutralisation, the title compound.

ES-MS m/e (%): 522.7 ($M+H^+$).

Example 95

2-{6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide

- 90 -




Analogous to general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-trifluoromethoxy-phenyl)-piperidin-1-yl]-methanone (prepared herein) with (commercially available) 2-chloro-*N,N*-dimethyl-acetamide gave the title compound.

5 ES-MS *m/e* (%): 508.4 ($M+H^+$).

Example 96

2-{6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide




10 Analogous to general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-trifluoromethyl-phenyl)-piperidin-1-yl]-methanone (prepared herein) with (commercially available) 2-chloro-*N*-methyl-acetamide gave the title compound.

ES-MS *m/e* (%): 477.7 ($M+H^+$).

Example 97

15 2-{6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone

- 91 -



Analogous to general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-trifluoromethyl-phenyl)-piperidin-1-yl]-methanone (prepared herein) with (commercially available) 2-chloro-1-(4-methyl-piperazin-1-yl)-ethanone gave the title compound.

ES-MS *m/e* (%): 546.8 (M+H⁺).

Example 98


2-{6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide

10 a) {6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid

Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-trifluoromethyl-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) bromo-acetic acid gave the title compound.

15 ES-MS *m/e* (%): 481.3 (M+H⁺).

b) 2-{6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid with (commercially available) ammonia in THF, gave the title compound.

ES-MS *m/e* (%): 464.6 (M+H⁺).

Example 99

2-{6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone hydrochloride




- 5 Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with (commercially available) piperazine-1-carboxylic acid *tert*-butyl ester gave, after treatment with HCl, the title compound.

ES-MS *m/e* (%): 532.7 (M+H⁺).

10

Example 100

2-{6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-methylamino-ethyl)-acetamide hydrochloride




- 15 Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with (commercially available) (2-amino-ethyl)-methyl-carbamic acid *tert*-butyl ester gave, after treatment with HCl the title compound.

ES-MS *m/e* (%): 521.3 (M+H⁺).

Example 101

- 20 *N*-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide hydrochloride

- 93 -




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with
 5 (commercially available) (2-amino-ethyl)-carbamic acid tert-butyl ester gave, after treatment with HCl, the title compound.

ES-MS m/e (%): 507.3 (M+H⁺).

Example 102

2-{6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide



10


Analogous to general procedure I, the coupling of (6-chloro-1*H*-indol-3-yl)-[4-(2-trifluoromethyl-phenyl)-piperidin-1-yl]-methanone (prepared herein) with
 (commercially available) 2-chloro-*N,N*-dimethyl-acetamide gave the title compound.

ES-MS m/e (%): 491.7 (M+H⁺).

15

Example 103

N-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide hydrochloride




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with (commercially available) N^1,N^1 -dimethyl-ethane-1,2-diamine, gave the title compound.

5 ES-MS m/e (%): 534.8 ($M+H^+$).

Example 104


[1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone



10 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-ethylamine gave the title compound.
ES-MS m/e (%): 424.5 ($M+H^+$).

Example 105

15 [1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-trifluoromethoxy-phenyl)-piperidin-1-yl]-methanone




Analogous to general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-trifluoromethoxy-phenyl)-piperidin-1-yl]-methanone (prepared herein) with (commercially available) 2-chloro-ethylamine gave the title compound.

ES-MS *m/e* (%): 466.4 (M+H⁺).

5

Example 106

2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-N-methyl-acetamide



Following general procedure II, the alkylation of (6-Chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-N-methyl-acetamide gave the title compound.


10

ES-MS *m/e* (%): 440.4 (M+H⁺).

Example 107

[6-Chloro-1-(2-methylamino-ethyl)-1*H*-indol-3-yl]-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone

15



Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) (2-chloro-ethyl)-methyl-amine gave the title compound.


ES-MS *m/e* (%): 438.4 (M+H⁺).

20

- 96 -

Example 108


[6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone



- 5 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) (2-chloro-ethyl)-dimethyl-amine gave the title compound. ES-MS *m/e* (%): 440.3 (M+H⁺).

Example 109

- 10 [6-Chloro-1-(2-methylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone




- Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with
15 (commercially available) (2-chloro-ethyl)-methyl-amine gave the title compound. ES-MS *m/e* (%): 426.3 (M+H⁺).

Example 110

[1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone

20

- 97 -



Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-ethylamine gave the title compound.

5 ES-MS *m/e* (%): 412.3 ($M+H^+$).

Example 111


2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide

a) (6-Chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperidin-1-yl]-methanone

10 Following general procedure I, the coupling of (commercially available) 4-(2-fluoro-phenyl)-piperidine (described herein below) with 6-chloro-1*H*-indole-3-carboxylic acid gave the title compound.

ES-MS *m/e* (%): 357.3 ($M+H^+$).

b) 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-
15 acetamide




Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperidin-1-yl]-methanone, with (commercially available) 2-chloro-*N*-methyl-acetamide gave the title compound.

20 ES-MS *m/e* (%): 428.2 ($M+H^+$).

Example 112

[1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperidin-1-yl]-methanone



- 5 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-ethylamine gave the title compound.

ES-MS m/e (%): 400.1 ($M+H^+$).


Example 113

- 10 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide
a) 6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid:

Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially
15 available) bromo-acetic acid gave the title compound.

ES-MS m/e (%): 413.0 ($M-H^+$).

b) 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide




- Following general procedure I, the coupling of {6-chloro-3-[4-(2-fluoro-phenyl)-
20 piperidine-1-carbonyl]-indol-1-yl}-acetic acid, with (commercially available) ammonia
in THF gave the title compound.

ES-MS m/e (%): 414.2 (M+H⁺).

Example 114

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide




5

Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperidin-1-yl]-methanone (preparation described herein below), with (commercially available) 2-chloro-*N,N*-dimethyl-acetamide gave the title compound.
ES-MS m/e (%): 442.3 (M+H⁺).

10

Example 115

[6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperidin-1-yl]-methanone



15


Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) (2-chloro-ethyl)-dimethyl-amine gave the title compound.
ES-MS m/e (%): 428.2 (M+H⁺).

Example 116

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone

20

- 100 -




Following general procedure I, the coupling of {6-chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (preparation described herein), with (commercially available) *N*¹,*N*¹-dimethyl-ethane-1,2-diamine gave the title compound.

5 ES-MS *m/e* (%): 485.2 (*M*+*H*⁺).

Example 117

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone




10 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperidin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-1-(4-methyl-piperazin-1-yl)-ethanone gave the title compound.

ES-MS *m/e* (%): 497.2 (*M*+*H*⁺).

Example 118

15 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone hydrochloride

- 101 -




Following general procedure I, the coupling of {6-chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (preparation described herein), with (commercially available) piperazine-1-carboxylic acid *tert*-butyl ester gave, after

5 treatment with HCl, gave the title compound.

ES-MS *m/e* (%): 483.2 (M+H⁺).

Example 119

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-(2-methylamino-ethyl)-acetamide hydrochloride



10


Following general procedure I, the coupling of {6-chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (preparation described herein), with (commercially available) (2-amino-ethyl)-methyl-carbamic acid *tert*-butyl ester gave, after treatment with HCl, the title compound.

15 ES-MS *m/e* (%): 471.2 (M+H⁺).

Example 120

N-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide hydrochloride

- 102 -




Following general procedure I, the coupling of {6-chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid (preparation described herein), with (commercially available) (2-amino-ethyl)-carbamic acid tert-butyl ester gave, after
5 treatment with HCl the title compound.

ES-MS m/e (%): 457.2 ($M+H^+$).

Example 121

(6-Chloro-1*H*-indol-3-yl)-(3,4,5,6-tetrahydro-2*H*-[4,4']bipyridinyl-1-yl)-methanone




10 Following general procedure I, the coupling of (commercially available) 1,2,3,4,5,6-hexahydro-[4,4']bipyridinyl with 6-chloro-1*H*-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 340.1 ($M+H^+$).

Example 122

15 2-[6-Chloro-3-(3,4,5,6-tetrahydro-2*H*-[4,4']bipyridinyl-1-carbonyl)-indol-1-yl]-*N,N*-dimethyl-acetamide

- 103 -




Following general procedure I, the coupling of (commercially available) 2-piperazin-1-yl-nicotinonitrile with 6-chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid gave the title compound.

5 ES-MS *m/e* (%): 425.2 ($M+H^+$).

Example 123

(6-Chloro-2-methyl-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone




To a solution of 0.040 g (0.19 mmol) 6-chloro-2-methyl-1*H*-indole-3-carboxylic acid,
 10 0.069 ml (0.40 mmol) *N,N*-diisopropylethylamine and 0.061 g (0.19 mmol) 2-(1*H*-
 benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate in 1 ml dry *N,N*-
 dimethylformamide were added 0.038 g (0.20 mmol) 4-(2-methoxy-phenyl)-piperidine
 at room temperature. After stirring for 1 h the reaction mixture was quenched with 0.5 M
 aqueous sodium hydroxide solution (20 ml) and extracted with ethyl acetate (2 x 30 ml).
 15 The combined organic layers were washed with water (2 x 30 ml) and brine (1 x 30 ml),
 dried over sodium sulfate and concentrated in vacuo. The residue was purified by flash
 chromatography (*n*-heptane / ethyl acetate) to give the title compound (0.050 g; 69%) as
 a white solid.

MS *m/e* (%): 381 ($M-H^+$, 100).

20

Example 124

(6-Chloro-1-methanesulfonyl-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-
 methanone




To a solution of 0.035 g (0.09 mmol) (6-chloro-1H-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone in 2 ml N,N-dimethylformamide were added 0.005 g (0.10 mmol) sodium hydride (50 % in oil). After 45', 0.008 ml (0.10 mmol) methanesulfonyl chloride were added. The reaction mixture was quenched with water after 3 h and extracted with ethyl acetate (2 x 50 ml). The combined organic layers were washed with water (2 x 30 ml) and brine (1 x 30), dried over sodium sulfate and concentrated to dryness. The residue was chromatographed (loaded as a solution in toluene; Flashpac 5 g; n-heptane / ethyl acetate 100 : 0 -> 75 : 25) to give the title compound (0.007 g; 17%) as a light yellow solid.

MS m/e (%): 447 (M+H⁺, 100).

Example 125

2-{6-Chloro-3-[4-(5-fluoro-benzo[d]isoxazol-3-yl)-piperidine-1-carbonyl]-indol-1-yl}-N-methyl-acetamide



15

Following general procedure I, the coupling of (commercially available) 5-fluoro-3-piperidin-4-yl-benzo[d]isoxazole with 6-chloro-1-methylcarbamoylmethyl-1H-indole-3-carboxylic acid (prepared as described herein) gave the title compound.


ES-MS m/e (%): 469.1 (M+H⁺).

20

Example 126

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-4-hydroxy-piperidine-1-carbonyl]-indol-1-yl}-N-methyl-acetamide

- 105 -




Following general procedure I, the coupling of 4-(2-fluoro-phenyl)-piperidin-4-ol (described in WO 2005118587) with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

- 5 ES-MS *m/e* (%): 444.1 (M+H⁺).

Example 127

2-[6-Chloro-3-(3,4,5,6-tetrahydro-2*H*-[4,4']bipyridinyl-1-carbonyl)-indol-1-yl]-*N*-methyl-acetamide




- 10 Following general procedure I, the coupling of (commercially available) 2-piperazin-1-yl-nicotinonitrile with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

ES-MS *m/e* (%): 411.2 (M+H⁺).

Example 128

- 15 10-[4-(2-Methoxy-phenyl)-piperidine-1-carbonyl]-3,4-dihydro-1*H*-pyrazino[1,2-*a*]indole-2-carboxylic acid tert-butyl ester

- 106 -




To a solution of 0.10 g (0.32 mmol) 3,4-dihydro-1H-pyrazino[1,2-a]indole-2,10-dicarboxylic acid 2-tert-butyl ester, 0.067 g (0.35 mmol) 4-(2-methoxyphenyl)-piperidine and 0.051 g (0.38 mmol) 1-hydroxybenzotriazole in 3.5 ml N,N-dimethylformamide
 5 were added 0.073 g (0.38 mmol) N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride at room temperature. After stirring for 3 h the reaction mixture was diluted with saturated aqueous ammonium chloride solution and extracted with tert-butyl methyl ether (2 x 50 ml). The combined organic layers were washed with 1 M sodium hydroxide solution (1 x 30 ml) and water (1 x 30 ml), dried over sodium sulfate,
 10 concentrated in vacuo and purified by flash-chromatography (aminopropyl-modified silica gel, n-heptane / ethyl acetate) to give the title compound (0.087 g, 56%) as a light yellow solid.

MS m/e (%): 490 (M+H⁺, 47).

Example 129

15 [4-(2-Methoxy-phenyl)-piperidin-1-yl]-(1,2,3,4-tetrahydro-pyrazino[1,2-a]indol-10-yl)-methanone hydrochloride




A mixture of 0.085 g (0.17 mmol) 10-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-3,4-dihydro-1H-pyrazino[1,2-a]indole-2-carboxylic acid tert-butyl ester and 1.4 ml of a
 20 1.25 M solution of hydrochloric acid (1.7 mmol) in methanol was stirred for 15 min at 50

°C. The reaction mixture was concentrated in vacuo to give the title compound (0.072 g, 97%) as a light yellow solid.

MS m/e (%): 390 (M+H⁺, 100).

Example 130

- 5 [4-(2-Methoxy-phenyl)-piperidin-1-yl]-(2-methyl-1,2,3,4-tetrahydro-pyrazino[1,2-a]indol-10-yl)-methanone



- A solution of 0.040 g (0.094 mmol) [4-(2-methoxy-phenyl)-piperidin-1-yl]-(1,2,3,4-tetrahydro-pyrazino[1,2-a]indol-10-yl)-methanone hydrochloride, 0.026 ml (0.19 mmol) triethylamine and 0.023 g (0.77 mmol) paraformaldehyde in 2 ml methanol was heated at reflux for 1 h. The reaction mixture was cooled to 0 °C on an ice-water bath and treated with 0.0089 g (0.14 mmol) sodium cyanoborohydride. After completed addition the mixture was allowed to warm to room temperature and stirred for 2 h. Quenching with water and dilution with 2 M aqueous sodium carbonate solution was followed by extraction with dichloromethane (2 x 50 ml). The combined organic layers were dried over sodium sulfate, concentrated in vacuo and purified by flash-chromatography (aminopropyl-modified silica gel, n-heptane / ethyl acetate) to give the title compound (0.031 g, 82%) as an off-white solid.

MS m/e (%): 404 (M+H⁺, 100).


20

Examples of compounds of formula I-c

Example 131

(1-Benzyl-2-methyl-1H-indol-3-yl)-(4-phenyl-piperazin-1-yl)-methanone

- 108 -




Amide coupling according to general procedure I:

- Amine: 1-Phenyl-piperazine (commercially available),
 - Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,
- 5 ES-MS m/e (%): 410.6 (M+H⁺).

Example 132

(1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone




Amide coupling according to general procedure I:

- 10 - Amine: 1-(2-Methoxy-phenyl)-piperazine (commercially available),
 - Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,
- ES-MS m/e (%): 440.6 (M+H⁺).

Example 133

(1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(4-methoxy-phenyl)-piperazin-1-yl]-methanone



15


Amide coupling according to general procedure I:

- Amine: 1-(4-Methoxy-phenyl)-piperazine (commercially available),
- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,

ES-MS m/e (%): 440.6 (M+H⁺).

Example 134

(1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(2-chloro-phenyl)-piperazin-1-yl]-methanone



5 Amide coupling according to general procedure I:


- Amine: 1-(2-Chloro-phenyl)-piperazine (commercially available),

- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,

ES-MS m/e (%): 444.5 (M+H⁺).

Example 135

10 (1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(4-chloro-phenyl)-piperazin-1-yl]-methanone



Amide coupling according to general procedure I:


- Amine: 1-(4-Chloro-phenyl)-piperazine (commercially available),

- Acid: 1-Benzyl-2-methyl-1H-indole-3-carboxylic acid,

15 ES-MS m/e (%): 444.5 (M+H⁺).

Example 136

(6-Chloro-1H-indol-3-yl)-(4-phenyl-piperazin-1-yl)-methanone



Amide coupling according to general procedure I:


Amine: 1-Phenyl-piperazine (commercially available),

- Acid: 6-chloro-1H-indole-3-carboxylic acid,

5 ES-MS m/e (%): 340.4 ($M+H^+$).

Example 137

(6-Chloro-1H-indol-3-yl)-[4-(2-chloro-6-nitro-phenyl)-piperazin-1-yl]-methanone




Following general procedure I, the coupling of (commercially available) 1-(2-chloro-6-
10 nitro-phenyl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title
compound.

ES-MS m/e (%): 419.4 ($M+H^+$).

Example 138

(6-Chloro-1H-indol-3-yl)-[4-(2,6-dichloro-phenyl)-piperazin-1-yl]-methanone




Following general procedure I, the coupling of (commercially available) 1-(2,6-dichloro-phenyl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 408.4 (M+H⁺).

Example 139

- 5 (6-Chloro-1H-indol-3-yl)-[4-(2,3-dihydro-benzo[1,4]dioxin-5-yl)-piperazin-1-yl]-methanone




- Following general procedure I, the coupling of (commercially available) 1-(2,3-dihydro-benzo[1,4]dioxin-5-yl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the
10 title compound.

ES-MS m/e (%): 398.4 (M+H⁺).

Example 140

- (6-Chloro-1H-indol-3-yl)-[4-(2-nitro-4-trifluoromethyl-phenyl)-piperazin-1-yl]-
methanone




- 15 Following general procedure I, the coupling of (commercially available) 1-(2-nitro-4-trifluoromethyl-phenyl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 453.4 (M+H⁺).

- 20 Example 141

(6-Chloro-1H-indol-3-yl)-[4-(2-chloro-phenyl)-piperazin-1-yl]-methanone

- 112 -




Following general procedure I, the coupling of (commercially available) 1-(2-chloro-phenyl)-piperazine, with 6-chloro-1H-indole-3-carboxylic acid gave the title compound

ES-MS m/e (%): 374.4 ($M+H^+$).

5

Example 142

[4-(2-Amino-6-chloro-phenyl)-piperazin-1-yl]-(6-chloro-1H-indol-3-yl)-methanone




Following general procedure I, the coupling of 3-chloro-2-piperazin-1-yl-phenylamine (described in Tetrahedron Letters (2001), 42(9), 1645-1646), with 6-chloro-1H-indole-3-carboxylic acid gave the title compound

10

ES-MS m/e (%): 389.4 ($M+H^+$).

Example 143

(6-Chloro-1H-indol-3-yl)-[4-(4-methoxy-phenyl)-piperazin-1-yl]-methanone




Following general procedure I, the coupling of (commercially available) 1-(4-methoxy-phenyl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound

ES-MS m/e (%): 370.4 ($M+H^+$).

Example 144

5 (6-Chloro-1H-indol-3-yl)-[4-(3-methoxy-phenyl)-piperazin-1-yl]-methanone




Following general procedure I, the coupling of (commercially available) 1-(3-methoxy-phenyl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound

ES-MS m/e (%): 370.4 ($M+H^+$).

10

Example 145

(6-Chloro-1H-indol-3-yl)-[4-(2-nitro-phenyl)-piperazin-1-yl]-methanone




Following general procedure I, the coupling of (commercially available) 1-(2-nitro-phenyl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

15 ES-MS m/e (%): 385.4 ($M+H^+$).

Example 146

(6-Chloro-1H-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone

- 114 -




Following general procedure I, the coupling of (commercially available) 1-(2-methoxyphenyl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 370.4 ($M+H^+$).

5

Example 147

(6-Chloro-1H-indol-3-yl)-[4-(4-fluoro-phenyl)-piperazin-1-yl]-methanone




Following general procedure I, the coupling of (commercially available) 1-(4-fluorophenyl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound

10 ES-MS m/e (%): 358.4 ($M+H^+$).

Example 148

(6-Chloro-1H-indol-3-yl)-[4-(3-fluoro-phenyl)-piperazin-1-yl]-methanone




15 Following general procedure I, the coupling of (commercially available) 1-(3-fluorophenyl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 358.4 ($M+H^+$).

Example 149

3-Chloro-4-[4-(6-chloro-1*H*-indole-3-carbonyl)-piperazin-1-yl]-benzotrile




- 5 Following general procedure I, the coupling of (commercially available) 3-chloro-4-piperazin-1-yl-benzotrile with 6-chloro-1*H*-indole-3-carboxylic acid gave the title compound

ES-MS m/e (%): 399.4 ($M+H^+$).

Example 150

- 10 (6-Chloro-1*H*-indol-3-yl)-[4-(2-ethoxy-phenyl)-piperazin-1-yl]-methanone




Amide coupling according to general procedure I:

- Amine: 1-(2-Ethoxy-phenyl)-piperazine (commercially available),
 - Acid: 6-Chloro-1*H*-indole-3-carboxylic acid,
- 15 ES-MS m/e (%): 384.0 ($M+H^+$).

Example 151

2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide

- 116 -




Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-*N,N*-dimethylacetamide gave the title compound.

5 ES-MS *m/e* (%): 455.2 (*M*+*H*⁺).

Example 152

2-{6-Chloro-3-[4-(2-ethoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethylacetamide



10 Amide coupling according to general procedure I:


- Amine: 1-(2-Ethoxy-phenyl)-piperazine (commercially available),

- Acid: 6-Chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid,

ES-MS *m/e* (%): 469.1 (*M*+*H*⁺).

Example 153

15 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-(4-methylpiperazin-1-yl)-ethanone




Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-1-(4-methyl-piperazin-1-yl)-ethanone gave the title compound.

- 5 ES-MS *m/e* (%): 510.6 (M+H⁺).

Example 154


2-{6-Chloro-3-[4-(2-ethoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone



- 10 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-ethoxy-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-1-(4-methyl-piperazin-1-yl)-ethanone gave the title compound.
ES-MS *m/e* (%): 524.6 (M+H⁺).

Example 155

- 15 2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide




- Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with (commercially available) *N*¹,*N*¹-dimethyl-ethane-1,2-diamine gave the title compound.
20

ES-MS m/e (%): 499.6 ($M+H^+$).

Example 156

2-{6-Chloro-3-[4-(2-ethoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide



5


Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-ethoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with (commercially available) *N,N*-dimethyl-ethane-1,2-diamine gave the title compound.

ES-MS m/e (%): 512.6 ($M+H^+$).

10

Example 157

2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetamide



15


Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid (prepared described herein) with (commercially available) ammonia in THF, gave the title compound.

ES-MS m/e (%): 427.5 ($M+H^+$).

Example 158

2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-5-methyl-indol-1-yl}-*N,N*-dimethyl-acetamide

- 119 -




Amide coupling according to general procedure I:

- Amine: 1-(2-Methoxy-phenyl)-piperazine (commercially available),
- Acid: 6-Chloro-1-dimethylcarbamoylmethyl-5-methyl-1H-indole-3-carboxylic acid,

5 ES-MS m/e (%): 469.3 (M+H⁺).

Example 159

2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone hydrochloride



- 10 Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid (prepared described herein) with (commercially available) piperazine-1-carboxylic acid *tert*-butyl ester gave, after treatment with HCl, the title compound.


ES-MS m/e (%): 496.5 (M+H⁺).

15

Example 160

2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-N-(2-methylamino-ethyl)-acetamide hydrochloride

- 120 -



Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid (prepared herein) with (commercially available) (2-amino-ethyl)-methyl-carbamic acid *tert*-butyl ester gave, after treatment
5 with HCl, the title compound.

ES-MS *m/e* (%): 484.5 (M+H⁺).

Example 161


N-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetamide hydrochloride

10 a) {6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid

Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) bromo-acetic acid gave the title compound.

ES-MS *m/e* (%): 428.5 (M+H⁺).

15 b) *N*-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetamide hydrochloride




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid with (commercially available) (2-amino-ethyl)-carbamic acid *tert*-butyl ester gave, after treatment with HCl, the title compound.

ES-MS *m/e* (%): 470.6 (M+H⁺).

5

Example 162

2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide




Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-*N*-methyl-acetamide gave the title compound.
ES-MS *m/e* (%): 441.5 (M+H⁺).

10

Example 163

[6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone




Analogous to general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone (prepared herein) with (commercially available) (2-chloro-ethyl)-dimethyl-amine gave the title compound.

ES-MS *m/e* (%): 441.5 (M+H⁺).

20

Example 164

[1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone




Analogous to general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone (prepared herein) with (commercially available) 2-chloro-ethylamine gave the title compound.

ES-MS m/e (%): 413.4 ($M+H^+$).

Example 165

[6-Chloro-1-(2-methylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone




Analogous to general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone (prepared herein) with (commercially available) (2-chloro-ethyl)-methyl-amine gave the title compound.

ES-MS m/e (%): 427.5 ($M+H^+$).

Example 166

[6-Chloro-1-(2-methylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-ethoxy-phenyl)-piperazin-1-yl]-methanone

- 123 -



Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-ethoxyphenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) (2-chloro-ethyl)-methyl-amine gave the title compound.

5 ES-MS *m/e* (%): 441.5 ($M+H^+$).

Example 167

2-{6-Chloro-3-[4-(2-ethoxyphenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-(2-methylamino-ethyl)-acetamide


a) {6-Chloro-3-[4-(2-ethoxyphenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid

10 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-ethoxyphenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) bromo-acetic acid gave the title compound.

ES-MS *m/e* (%): 440.0 ($M+H^+$).

b) 2-{6-Chloro-3-[4-(2-ethoxyphenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-(2-

15 methylamino-ethyl)-acetamide




Analogous to general procedure I, the coupling of {6-chloro-3-[4-(2-ethoxyphenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid with (commercially available) (2-aminoethyl)-methyl-carbamic acid *tert*-butyl ester gave, after treatment with HCl and

20 neutralisation, the title compound.

ES-MS *m/e* (%): 498.5 ($M+H^+$).

Example 168

[6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-ethoxy-phenyl)-piperazin-1-yl]-methanone




- 5 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-ethoxy-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) (2-chloro-ethyl)-dimethyl-amine gave the title compound.

ES-MS m/e (%): 455.3 ($M+H^+$).

Example 169

10 (6-Chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone




Following general procedure I, the coupling of (commercially available) 1-(2-fluoro-phenyl)-piperazine (described herein) with 6-chloro-1*H*-indole-3-carboxylic acid gave the title compound.

- 15 ES-MS m/e (%): 358.0 ($M+H^+$).

Example 170

[1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone

- 125 -




Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluorophenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-ethylamine gave the title compound.

5 ES-MS m/e (%): 401.2 ($M+H^+$).

Example 171

[6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-fluorophenyl)-piperazin-1-yl]-methanone




10 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluorophenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) (2-chloro-ethyl)-dimethyl-amine gave the title compound.

ES-MS m/e (%): 429.3 ($M+H^+$).

Example 172

15 2-{6-Chloro-3-[4-(2-fluorophenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone




Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-1-(4-methyl-piperazin-1-yl)-ethanone gave the title compound.

ES-MS *m/e* (%): 498.3 (M+H⁺).

5

Example 173

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide




Following general procedure I, the coupling of {6-chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid, with (commercially available) *N*¹,*N*¹-dimethyl-ethane-1,2-diamine gave the title compound.

ES-MS *m/e* (%): 486.3 (M+H⁺).

15

Example 174

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide



Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-*N,N*-dimethyl-acetamide gave the title compound.


ES-MS *m/e* (%): 443.2 (M+H⁺).

20

Example 175

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetamide

- 127 -




Following general procedure I, the coupling of {6-chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid (preparation described herein), with (commercially available) ammonia in THF gave the title compound.

- 5 ES-MS m/e (%): 415.2 (M+H⁺).

Example 176

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-N-methyl-acetamide



- 10 Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) 2-chloro-*N*-methyl-acetamide gave the title compound.

ES-MS m/e (%): 429.2 (M+H⁺).

Example 177


- 15 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone

a) {6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid

- Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (commercially available) bromo-acetic acid gave the title compound.

20 ES-MS m/e (%): 414.0 (M-H⁺).

b) 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone




Following general procedure I, the coupling of {6-chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid, with (commercially available) piperazine-1-carboxylic acid *tert*-butyl ester gave, after treatment with HCl and neutralisation, gave the title compound.

ES-MS *m/e* (%): 484.2 (M+H⁺).

Example 178

10 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-(2-methylamino-ethyl)-acetamide




Following general procedure II, the alkylation of {6-chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid (preparation described herein), with (commercially available) (2-amino-ethyl)-methyl-carbamic acid *tert*-butyl ester gave, after treatment with HCl and neutralisation, the title compound.

ES-MS *m/e* (%): 472.2 (M+H⁺).

Example 179

20 *N*-(2-Amino-ethyl)-2-{6-chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetamide



Following general procedure I, the coupling of {6-chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetic acid (preparation described herein), with (commercially available) (2-amino-ethyl)-carbamic acid *tert*-butyl ester gave, after
 5 treatment with HCl and neutralisation, the title compound.

ES-MS *m/e* (%): 458.2 (M+H⁺).

Example 180

2-{6-Chloro-3-[4-(2-methoxymethyl-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide

10 a) 4-(2-Methoxymethyl-phenyl)-piperazine-1-carboxylic acid *tert*-butyl ester

A solution of (commercially available) 4-(2-hydroxymethyl-phenyl)-piperazine-1-carboxylic acid *tert*-butyl ester and (1eq) MeI dissolved in THF was treated with NaH (2eq) and stirred at RT for 2h. After work-up, the crude solid was purified by SiO₂ chromatography (Ethyl acetate- Heptane 1:4) to give the title compound.


15 ES-MS *m/e* (%): 307.5 (M+H⁺).

b) 1-(2-Methoxymethyl-phenyl)-piperazine

A solution of 4-(2-methoxymethyl-phenyl)-piperazine-1-carboxylic acid *tert*-butyl ester (prepared as described herein below) in dioxan was treated with 4eq. of 4M HCl in dioxane and stirred at RT for 16h. After neutralization, the mixture was completely
 20 evaporated to give the crude title compound.

ES-MS *m/e* (%): 207.1 (M+H⁺).

c) 2-{6-Chloro-3-[4-(2-methoxymethyl-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide




Following general procedure I, the coupling of 1-(2-methoxymethyl-phenyl)-piperazine, with 6-chloro-1-[(2-dimethylamino-ethylcarbamoyl)-methyl]-1*H*-indole-3-carboxylic acid (preparation described herein) gave the title compound.

5 ES-MS *m/e* (%): 512.3 ($M+H^+$).

Example 181

2-[6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-(2-dimethylamino-ethyl)-acetamide




10 Following general procedure I, the coupling of (commercially available) 1-(2-trifluoromethyl-phenyl)-piperazine, with 6-chloro-1-[(2-dimethylamino-ethylcarbamoyl)-methyl]-1*H*-indole-3-carboxylic acid (preparation described herein) gave the title compound.

ES-MS *m/e* (%): 536.2 ($M+H^+$).

15

Example 182

2-[6-Chloro-3-(4-pyridin-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N*-(2-dimethylamino-ethyl)-acetamide




Following general procedure I, the coupling of (commercially available) 1-pyridin-2-yl-piperazine, with 6-chloro-1-[(2-dimethylamino-ethylcarbamoyl)-methyl]-1*H*-indole-3-carboxylic acid (preparation described herein) gave the title compound.

- 5 ES-MS *m/e* (%): 469.2 ($M+H^+$).

Example 183

2-[6-Chloro-3-(4-pyridin-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N*-(2-dimethylamino-ethyl)-acetamide




- 10 Following general procedure I, the coupling of (commercially available) 1-(2,4-difluorophenyl)-piperazine with 6-chloro-1*H*-indole-3-carboxylic acid gave the title compound.

ES-MS *m/e* (%): 376.1 ($M+H^+$).

Example 184

(6-Chloro-1*H*-indol-3-yl)-[4-(3-methyl-pyridin-2-yl)-piperazin-1-yl]-methanone

- 132 -




Following general procedure I, the coupling of (commercially available) 1-(3-methylpyridin-2-yl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

5 ES-MS m/e (%): 355.1 (M+H⁺).

Example 185

(6-Chloro-1H-indol-3-yl)-[4-(3,5-dichloro-pyridin-4-yl)-piperazin-1-yl]-methanone




10 Following general procedure I, the coupling of (commercially available) 1-(3,5-dichloropyridin-4-yl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 409.0 (M+H⁺).

Example 186

(6-Chloro-1H-indol-3-yl)-[4-(3-trifluoromethylpyridin-2-yl)-piperazin-1-yl]-methanone

15




Following general procedure I, the coupling of (commercially available) 1-(3-trifluoromethyl-pyridin-2-yl)-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 409.1 ($M+H^+$).

5

Example 187

2-[4-(6-Chloro-1H-indole-3-carbonyl)-piperazin-1-yl]-nicotinonitrile




Following general procedure I, the coupling of (commercially available) 2-piperazin-1-yl-nicotinonitrile with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

10 ES-MS m/e (%): 366.1 ($M+H^+$).

Example 188

(6-Chloro-1H-indol-3-yl)-(4-pyridin-2-yl-piperazin-1-yl)-methanone




15 Following general procedure I, the coupling of (commercially available) 1-pyridin-2-yl-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 341.1 ($M+H^+$).

Example 189

(6-Chloro-1H-indol-3-yl)-(4-thiazol-2-yl-piperazin-1-yl)-methanone

- 134 -




Following general procedure I, the coupling of (commercially available) 1-thiazol-2-yl-piperazine with 6-chloro-1H-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 347.0 ($M+H^+$).

5

Example 190

2-[6-Chloro-3-[4-(2,4-difluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methylacetamide




Following general procedure I, the coupling of (commercially available) 1-(2,4-difluoro-phenyl)-piperazine, with 6-chloro-1-methylcarbamoylmethyl-1H-indole-3-carboxylic acid (prepared as described herein below) gave the title compound.

ES-MS m/e (%): 447.1 ($M+H^+$).

Example 191

2-[6-Chloro-3-(2,3,5,6-tetrahydro-[1,2']bipyrazinyl-4-carbonyl)-indol-1-yl]-*N*-methylacetamide

15




Following general procedure I, the coupling of (commercially available) 3,4,5,6-tetrahydro-2*H*-[1,2']bipyrazinyl, with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

ES-MS *m/e* (%): 413.1 (M+H⁺).

5

Example 192

2-[6-Chloro-3-[4-(3-methyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methyl-acetamide




Following general procedure I, the coupling of (commercially available) 1-(3-methyl-
10 pyridin-2-yl)-piperazine with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

ES-MS *m/e* (%): 426.2 (M+H⁺).

Example 193

15 2-[6-Chloro-3-[4-(3,5-dichloro-pyridin-4-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methyl-acetamide




Following general procedure I, the coupling of (commercially available) 1-(3,5-dichloro-
20 pyridin-4-yl)-piperazine with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

ES-MS *m/e* (%): 480.1 (M+H⁺).

Example 194

2-[6-Chloro-3-(4-pyridin-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N*-methyl-acetamide

- 136 -




Following general procedure I, the coupling of (commercially available) 1-pyridin-2-yl-piperazine with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

5 ES-MS *m/e* (%): 412.1 ($M+H^+$).

Example 195

2-[6-Chloro-3-(4-thiazol-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N*-methyl-acetamide




Following general procedure I, the coupling of (commercially available) 1-thiazol-2-yl-piperazine with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

10 ES-MS *m/e* (%): 418.1 ($M+H^+$).

Example 196

2-[6-Chloro-3-[4-(3-trifluoromethyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methyl-acetamide

15




Following general procedure I, the coupling of (commercially available) 1-(3-trifluoromethyl-pyridin-2-yl)-piperazine with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

ES-MS *m/e* (%): 480.1 (M+H⁺).

5

Example 197

2-{6-Chloro-3-[4-(3-cyano-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide




Following general procedure I, the coupling of (commercially available) 2-piperazin-1-yl-nicotinonitrile with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

ES-MS *m/e* (%): 437.1 (M+H⁺).

Example 198

[1-((*S*)-2-Amino-propyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone

15




Following general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with methanesulfonic acid (*S*)-2-*tert*-butoxycarbonylamino-propyl ester (described in WO 2005100321) gave, after treatment with TFA and subsequent neutralization, the title compound.

ES-MS *m/e* (%): 415.2 (M+H⁺).

Example 199


(6-Chloro-1-(S)-1-piperidin-3-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone



- 5 In analogy to the reaction conditions used in general procedure II, the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with methanesulfonic acid (S)-3-methanesulfonyloxymethyl-piperidine-1-carboxylic acid *tert*-butyl ester (described in JP 2001278872), followed by treatment with TFA and subsequent neutralisation, gave the title compound in 56% yield.
- 10 ES-MS *m/e* (%): 455.3 (M+H⁺).

Example 200


(6-Chloro-1-(S)-1-pyrrolidin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone



- 15 In analogy to the preparation of (6-chloro-1-(S)-1-piperidin-3-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (described herein), the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (S)-2-methanesulfonyloxymethyl-pyrrolidine-1-carboxylic acid *tert*-butyl ester (described in Tetrahedron: Asymmetry (1997), 8(13), 2209-2213),
- 20 followed by treatment with TFA and subsequent neutralisation, gave the title compound in 55% yield.
- ES-MS *m/e* (%): 441.3 (M+H⁺).

Example 201


(6-Chloro-1-(RS)-1-pyrrolidin-3-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone



- 5 In analogy to the preparation of (6-chloro-1-(*S*)-1-piperidin-3-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (described herein), the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (*RS*)-3-methanesulfonyloxymethyl-pyrrolidine-1-carboxylic acid *tert*-butyl ester (described in WO 9742189), followed by treatment with TFA and
- 10 subsequent neutralisation, gave the title compound in 51% yield.
ES-MS *m/e* (%): 441.3 (*M*+*H*⁺).

Example 202


[6-Chloro-1-((*S*)-1-methyl-piperidin-3-ylmethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone



- 15 A solution of (6-chloro-1-(*S*)-1-piperidin-3-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein below) in MeOH was treated with aq. H₂CO (1.5 eq) and AcOH (1.1eq) and stirred for 15min at RT, then treated with NaCNBH₃ (1.1 eq) and stirred at RT for 1h. Concentration and purification
- 20 by prep HPLC gave the desired product.in 71% yield.
ES-MS *m/e* (%): 469.3 (*M*+*H*⁺).

Example 203


[6-Chloro-1-((S)-1-methyl-pyrrolidin-2-ylmethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone



- 5 (6-Chloro-1-(S)-1-pyrrolidin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein), was treated with a 37% aq. solution of formaldehyde (1.05eq.), acetic acid (1.05eq.) and sodium cyanoborohydride (1.0 eq.) in MeOH at RT for 2h to give after purification by prep. HPLC the title compound in 62% yield.
- 10 ES-MS m/e (%): 455.3 (M+H⁺).

Example 204

[6-Chloro-1-((RS)-1-methyl-pyrrolidin-3-ylmethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone




- 15 Following the procedure described in the preparation of (6-chloro-1-(S)-1-pyrrolidin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (described herein), the alkylation of (6-chloro-1-(RS)-1-pyrrolidin-3-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein) gave the title compound in 64% yield.
- 20 ES-MS m/e (%): 455.3 (M+H⁺).

Example 205

2-{6-Chloro-3-[4-(2,4-difluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide

- 141 -




Following general procedure I, the coupling of (commercially available) 1-(2,4-difluorophenyl)-piperazine, with 6-chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid gave the title compound.

- 5 ES-MS *m/e* (%): 461.2 (M+H⁺).

Example 206


2-[6-Chloro-3-(2,3,5,6-tetrahydro-[1,2']bipyrazinyl-4-carbonyl)-indol-1-yl]-*N,N*-dimethyl-acetamide



- 10 Following general procedure I, the coupling of (commercially available) 3,4,5,6-tetrahydro-2*H*-[1,2']bipyrazinyl, with 6-chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.
ES-MS *m/e* (%): 427.2 (M+H⁺).

Example 207

- 15 2-[6-Chloro-3-[4-(3-methyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N,N*-dimethyl-acetamide




Following general procedure I, the coupling of (commercially available) 1-(3-methylpyridin-2-yl)-piperazine with 6-chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

- 5 ES-MS m/e (%): 440.2 ($M+H^+$).

Example 208

2-[6-Chloro-3-[4-(3,5-dichloro-pyridin-4-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N,N*-dimethyl-acetamide




- 10 Following general procedure I, the coupling of (commercially available) 1-(3,5-dichloro-pyridin-4-yl)-piperazine with 6-chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

ES-MS m/e (%): 494.1 ($M+H^+$).

Example 209

- 15 2-[6-Chloro-3-(4-pyridin-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N*-methyl-acetamide

- 143 -




Following general procedure I, the coupling of (commercially available) 1-pyridin-2-yl-piperazine with 6-chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

5 ES-MS *m/e* (%): 426.2 ($M+H^+$).

Example 210

2-[6-Chloro-3-(4-thiazol-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N,N*-dimethylacetamide




10 Following general procedure I, the coupling of (commercially available) 1-thiazol-2-yl-piperazine with 6-chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

ES-MS *m/e* (%): 432.2 ($M+H^+$).

Example 211

15 2-{6-Chloro-3-[4-(3-trifluoromethyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethylacetamide




Following general procedure I, the coupling of (commercially available) 1-(3-trifluoromethyl-pyridin-2-yl)-piperazine with 6-chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid gave the title compound.

ES-MS m/e (%): 494.2 ($M+H^+$).

5

Example 212

2-{6-Chloro-3-[4-(3-cyano-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide




Following general procedure I, the coupling of (commercially available) 2-piperazin-1-yl-nicotinonitrile with 6-chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

ES-MS m/e (%): 451.2 ($M+H^+$).

15

Example 213

2-[6-Chloro-3-(4-pyrimidin-2-yl-piperazine-1-carbonyl)-indol-1-yl]-*N,N*-dimethyl-acetamide




Following general procedure I, the coupling of (commercially available) 2-piperidin-4-yl-pyrimidine with 6-chloro-1-dimethylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

ES-MS m/e (%): 427.2 ($M+H^+$).

20

Example 214

(6-Chloro-1-(R)-1-pyrrolidin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone




- 5 In analogy to the preparation of (6-chloro-1-(*S*)-1-piperidin-3-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (described herein), the alkylation of (6-chloro-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein), with (R)-2-methanesulfonyloxymethyl-pyrrolidine-1-carboxylic acid *tert*-butyl ester (described in Tetrahedron: Asymmetry (1997), 8(13), 2209-2213),
 10 followed by treatment with TFA and subsequent neutralisation, gave the title compound in 26% yield.

ES-MS *m/e* (%): 441.3 (M+H⁺).

Example 215

15 [6-Chloro-1-((R)-1-methyl-pyrrolidin-2-ylmethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone




- Following the procedure described in the preparation of [6-chloro-1-((*S*)-1-methyl-pyrrolidin-2-ylmethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone, the alkylation of (6-chloro-1-(*R*)-1-pyrrolidin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein) gave the title
 20 compound.

ES-MS *m/e* (%): 455.3 (M+H⁺)

Example 216

(6-Chloro-2-methyl-1H-indol-3-yl)-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone



The title compound was obtained as a white solid in 16% yield according to the procedure described for the preparation of (6-chloro-2-methyl-1H-indol-3-yl)-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone using 1-(2-methoxy-phenyl)-piperazine instead of 4-(2-methoxy-phenyl)-piperidine.

MS m/e (%): 382 (M-H⁺, 100).

Example 217

N-(2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-ethyl)-acetamide

a) [1-(2-Amino-ethyl)-6-chloro-1H-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone:


To a solution of (6-chloro-1H-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone in dry DMF under argon at 0 °C was added (1.1eq) NaH. The reaction mixture was stirred for 1 h at 0 °C. 2,2-Dioxo-2λ⁶-[1,2,3]oxathiazolidine-3-carboxylic acid *tert*-butyl ester (1.1eq) (described in WO 2003037327) was added, and the reaction mixture was stirred for 2h at RT. After evaporation of DMF, redissolution in dioxan, addition of 5eq. of HCl (4.0M solution in dioxane) and a few drops of water, the reaction mixture was stirred for 2h at 50 °C; LC-MS showed complete conversion to the crude product.

Concentration in vacuo, redissolution in EtOAc, and washing with 1N NaHCO₃, gave after concentration, the crude product which was purified by SiO₂ gel chromatography with CH₂Cl₂/MeOH to give the title compound in 55% yield.

ES-MS m/e (%): 401.2 (M+H⁺).

b) *N*-(2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-ethyl)-acetamide

- 147 -




[1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone, was treated with acetylchloride (1.05eq) and triethylamine (1.05eq) in CH₂Cl₂ under argon at RT for 2h to give after purification by prep. HPLC the title compound.

ES-MS m/e (%): 443.2 (M+H⁺).

Example 218

N-(2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-ethyl)-methanesulfonamide



10


[1-(2-Amino-ethyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone (preparation described herein), was treated with mesylchloride (1.05eq) and triethylamine (1.05eq) in CH₂Cl₂ under argon at RT.to give after purification by prep. HPLC the title compound.

ES-MS m/e (%): 479.1 (M+H⁺).

Example 219

N-(2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-ethyl)-*N*-methyl-acetamide

- 148 -




N-(2-(6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl)-ethyl)-acetamide (preparation described herein) was treated with NaH (1.05eq), MeI (1.05eq) in dry DMF under argon at RT for 2h to give after purification by prep. HPLC the title compound.

ES-MS *m/e* (%): 457.1 (M+H⁺).

Example 220

N-(2-(6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl)-ethyl)-*N*-methyl-acetamide



10


N-(2-(6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl)-ethyl)-methanesulfonamide (preparation described herein), was treated with NaH (1.05eq), MeI (1.05eq) in dry DMF under argon at RT for 2h to give after purification by prep. HPLC the title compound.

ES-MS *m/e* (%): 493.1 (M+H⁺).

Example 221

2-(6-Chloro-3-[4-(6-chloro-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl)-*N*-methyl-acetamide

- 149 -




Following general procedure I, the coupling of (commercially available) 1-(6-chloro-pyridin-2-yl)-piperazine, with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

5 ES-MS *m/e* (%): 446.1 ($M+H^+$).

Example 222

2-[6-Chloro-3-[4-(5-trifluoromethyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methyl-acetamide




10 Following general procedure I, the coupling of (commercially available) 1-(5-trifluoromethyl-pyridin-2-yl)-piperazine, with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.
ES-MS *m/e* (%): 480.1 ($M+H^+$).

Example 223

15 2-[6-Chloro-3-(4-thieno[2,3-*c*]pyridin-7-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methyl-acetamide

- 150 -




Following general procedure I, the coupling of (commercially available) 7-piperazin-1-yl-thieno[2,3-*c*]pyridine, with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

5 ES-MS *m/e* (%): 468.1 ($M+H^+$).

Example 224

2-[6-Chloro-3-(4-thieno[3,2-*c*]pyridin-4-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methyl-acetamide




10 Following general procedure I, the coupling of (commercially available) 4-piperazin-1-yl-thieno[3,2-*c*]pyridine with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

ES-MS *m/e* (%): 468.1 ($M+H^+$).

Example 225

15 2-[6-Chloro-3-[4-(3-iodo-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methyl-acetamide

- 151 -




Following general procedure I, the coupling of (commercially available) 1-(3-iodo-pyridin-2-yl)-piperazine with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid gave the title compound.

- 5 ES-MS *m/e* (%): 538.0 ($M+H^+$).

Example 226

2-[6-Chloro-3-[4-(3-chloro-5-trifluoromethyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl]-*N*-methylacetamide




- 10 Following general procedure I, the coupling of (commercially available) 1-(3-chloro-5-trifluoromethyl-pyridin-2-yl)-piperazine with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

ES-MS *m/e* (%): 514.1 ($M+H^+$).

Example 227

- 15 2-[4-(6-Chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carbonyl)-piperazin-1-yl]-nicotinic acid

- 152 -




Following general procedure I, the coupling of (commercially available) 2-piperazin-1-yl-nicotinic acid with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

5 ES-MS *m/e* (%): 456.1 (M+H⁺).

Example 228


2-{{6-Chloro-3-[4-(2,4-difluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-(2-dimethylamino-ethyl)-acetamide



10 Following general procedure I, the coupling of (commercially available) 1-(2,4-difluoro-phenyl)-piperazine with 6-chloro-1-[(2-dimethylamino-ethylcarbamoyl)-methyl]-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.
ES-MS *m/e* (%): 504.3 (M+H⁺).

Example 229

15 2-{{6-Chloro-3-[4-(4-fluoro-2-methanesulfonyl-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide




Following general procedure I, the coupling of (commercially available) 1-(4-fluoro-2-methanesulfonyl-phenyl)-piperazine with 6-chloro-1-methylcarbamoylmethyl-1*H*-indole-3-carboxylic acid (prepared as described herein) gave the title compound.

5 ES-MS *m/e* (%): 506.9 ($M+H^+$).

Example of a compound of formula I-d

Example 230

(6-Chloro-1*H*-indol-3-yl)-[4-(2-methoxy-phenyl)-4-oxy-piperazin-1-yl]-methanone



10

Amide coupling according to general procedure I:

- Amine: 1-(2-Methoxy-phenyl)-piperazine 1-oxide (described in EP126480),


- Acid: 6-chloro-1*H*-indole-3-carboxylic acid,

ES-MS *m/e* (%): 386.4 ($M+H^+$).

15

Claims

1. A compound of the following general formula I:



wherein

- 5 R^1 is H,
 or is C_{1-6} -alkyl substituted by CN, C_{1-6} -alkoxy, OH, halo, or by NR^iR^{ii} ,
 or is C_{2-6} -alkyl,
 or is aryl, 5 or 6 membered heteroaryl or sulfonylaryl which are optionally
 substituted by one or more B,
 10 or is $-(CH_2)_m-R^a$ wherein R^a is:
 CN,
 OR^i ,
 NR^iR^{ii} ,
 C_{3-6} -cycloalkyl, 4 to 7 membered-heterocycloalkyl, aryl, or 5 or 6 membered
 15 heteroaryl which are optionally substituted by one or more B,
 or is $-(CH_2)_n-(CO)-R^b$ or $-(CH_2)_n-(SO_2)-R^b$, wherein R^b is:
 C_{1-6} -alkyl,
 C_{1-6} -alkoxy,
 C_{3-6} -cycloalkyl,
 20 $-(CH_2)_m-NR^{iii}R^{iv}$,
 NR^iR^{ii} ,
 C_{3-6} -cycloalkyl, 4 to 7 membered-heterocycloalkyl, aryl, or 5 or 6 membered
 heteroaryl which are optionally substituted by one or more B,
 or R^1 and R^3 together with the indole ring to which they are attached form a 5 or
 25 6 membered heterocycloalkyl which can be substituted by =O, $C(O)O-C_{1-6}$ -
 alkyl or C_{1-6} -alkyl;
 R^2 is one or more of H, OH, halo, CN, nitro, C_{1-6} -alkoxy, $-O-CH_2-C_{2-6}$ -alkenyl,
 benzyloxy, C_{1-6} -haloalkoxy, C_{1-6} -alkyl optionally substituted by $-NR^{iii}R^{iv}$ or halo,
 or two R^2 may form an oxo or dioxo bridge together with the indole ring to
 30 which they are attached;
 R^3 is H,

or is F,

or is $-(\text{CO})-\text{R}^c$, wherein R^c is:

C_{1-6} -alkyl,

$-(\text{CH}_2)_n-\text{NR}^i\text{R}^{ii}$,

5 $-(\text{CH}_2)_n-\text{NR}^{iii}\text{R}^{iv}$,

5 or 6 membered heterocycloalkyl optionally substituted by C_{1-6} -alkyl,

or is C_{1-6} -alkyl which is optionally substituted by

halo,

NR^iR^{ii} ,

10 $\text{NR}^{iii}\text{R}^{iv}$,


$-\text{O}(\text{CO})-\text{C}_{1-6}$ -alkyl,

or by $-\text{NH}(\text{CO})\text{R}^d$, wherein R^d is C_{1-6} -alkyl optionally substituted by halo

or nitro, or R^d is aryl or a 5 or 6 membered heteroaryl, which are

optionally substituted by halo, nitro, C_{1-6} -alkyl or C_{1-6} -haloalkyl;

15 A is selected from the group consisting of (a), (a'), (b), (c) and (d):



wherein

R^4 is $-\text{NH}(\text{CO})\text{R}^e$, wherein R^e is C_{1-6} -alkoxy, or aryl optionally substituted by halo,
or is C_{1-6} -alkoxy, CN,

20 or is aryl, 5 or 6 membered heteroaryl, benzyl, aryloxy or a 9 or 10-membered bicyclic heteroaryl ring, which are optionally substituted by CN, halo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -haloalkyl, nitro, hydroxyl, NR^iR^{ii} , $\text{NR}^{iii}\text{R}^{iv}$, C_{1-6} -alkoxy- C_{1-6} -alkylene, $\text{S}(\text{O})_2-\text{C}_{1-6}$ -alkyl, C_{1-6} -haloalkoxy, or by a oxo or dioxo bridge; and

25 R^5 is H, OH, CN, COOR^{iii} or $\text{CONR}^{iii}\text{R}^{iv}$;

R^6 is C_{2-6} -alkyl,

or is $-\text{C}(\text{O})-\text{R}^f$ wherein R^f is an aryl group substituted by halo, C_{1-6} -alkoxy, or CN,

30 or is aryl, 5 or 6 membered heteroaryl, or a 9 or 10-membered bicyclic heteroaryl ring which are optionally substituted by halo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -

haloalkyl, CN, nitro, $\text{NR}^{\text{i}}\text{R}^{\text{ii}}$, $\text{NR}^{\text{iii}}\text{R}^{\text{iv}}$, C_{1-6} -alkoxy- C_{1-6} -alkylene, COOH , $\text{S}(\text{O})_2$ - C_{1-6} -alkyl, hydroxyl, C_{1-6} -haloalkoxy, or by a oxo or dioxo bridge,
or is benzyl substituted by halo, C_{1-6} -alkyl, C_{1-6} -haloalkyl or by a oxo – or dioxo
bridge;

5 B is halo, CN, $\text{NR}^{\text{i}}\text{R}^{\text{ii}}$, C_{1-6} -alkyl optionally substituted by CN, halo or C_{1-6} -alkoxy,
 C_{1-6} -haloalkyl, C_{1-6} -alkoxy, C_{1-6} -haloalkoxy, C_{3-6} -cycloalkyl, $-\text{C}(\text{O})\text{O}-\text{C}_{1-6}$ -alkyl, $-\text{C}(\text{O})\text{NR}^{\text{i}}\text{R}^{\text{ii}}$, $-\text{C}(\text{O})-\text{C}_{1-6}$ -alkyl, $-\text{S}(\text{O})_2-\text{C}_{1-6}$ -alkyl, $-\text{S}(\text{O})_2-\text{NR}^{\text{i}}\text{R}^{\text{ii}}$, $(\text{CR}^{\text{iii}}\text{R}^{\text{iv}})_n$ -
phenyl, or $(\text{CR}^{\text{iii}}\text{R}^{\text{iv}})_n$ -5 or 6 membered heteroaryl wherein the phenyl or 5 or 6
membered heteroaryl moiety is optionally substituted by one or more
10 substituent(s) selected from the group consisting of:

halo, CN, $\text{NR}^{\text{i}}\text{R}^{\text{ii}}$, C_{1-6} -alkyl optionally substituted by CN or C_{1-6} -alkoxy,
 C_{1-6} -alkoxy, C_{1-6} -haloalkoxy, C_{3-6} -cycloalkyl, $-\text{C}(\text{O})\text{O}-\text{C}_{1-6}$ -alkyl, $-\text{C}(\text{O})-$
 $\text{NR}^{\text{i}}\text{R}^{\text{ii}}$, $-\text{C}(\text{O})-\text{C}_{1-6}$ -alkyl, $-\text{S}(\text{O})_2-\text{C}_{1-6}$ -alkyl, $-\text{S}(\text{O})_2-\text{NR}^{\text{i}}\text{R}^{\text{ii}}$;

R^{i} and R^{ii} are H, C_{1-6} -alkyl, C_{1-6} -alkyl- $\text{NR}^{\text{iii}}\text{R}^{\text{iv}}$, $-(\text{CO})\text{O}-\text{C}_{1-6}$ -alkyl, $-\text{C}(\text{O})-\text{NR}^{\text{iii}}\text{R}^{\text{iv}}$, $-\text{C}(\text{O})-$
15 C_{1-6} -alkyl, $-\text{S}(\text{O})_2-\text{C}_{1-6}$ -alkyl or $-\text{S}(\text{O})_2-\text{NR}^{\text{iii}}\text{R}^{\text{iv}}$;

R^{iii} and R^{iv} are H or C_{1-6} -alkyl;

m is 1 to 6;

n is 0 to 4;

as well as pharmaceutically acceptable salts thereof.

20

2. A compound of the following general formula I according to claim 1,

wherein

R^{i} is H,

or is C_{1-6} -alkyl optionally substituted by CN or by C_{1-6} -alkoxy,

25 or is aryl, 5 or 6 membered heteroaryl or sulfonylaryl which are optionally
substituted by one or more B,

or is $-(\text{CH}_2)_m-\text{R}^{\text{a}}$ wherein R^{a} is:

CN,

OR^{i} ,

30 $\text{NR}^{\text{i}}\text{R}^{\text{ii}}$,


C_{3-6} -cycloalkyl, 4 to 7 membered-heterocycloalkyl, aryl, or 5 or 6 membered
heteroaryl which are optionally substituted by one or more B,

or is $-(\text{CH}_2)_n-\text{CO}-\text{R}^{\text{b}}$ or $-(\text{CH}_2)_n-\text{SO}_2-\text{R}^{\text{b}}$, wherein R^{b} is:

C_{1-6} -alkyl,

35 C_{1-6} -alkoxy,

C_{3-6} -cycloalkyl,



C₃₋₆-cycloalkyl, 4 to 7 membered-heterocycloalkyl, aryl, or 5 or 6 membered heteroaryl which are optionally substituted by one or more B,
 5 or R¹ and R³ together with the indole ring to which they are attached form a 5 or 6 membered heterocycloalkyl which can be substituted by (CO);

R² is one or more of H, OH, halo, CN, nitro, C₁₋₆-alkyl optionally substituted by –NRⁱⁱⁱR^{iv}, C₁₋₆-alkoxy, –O-CH₂-C₂₋₆-alkenyl, benzyloxy,
 10 or two R² may form an oxo or dioxo bridge together with the indole ring to which they are attached;


R³ is H,
 or is halo,
 or is –(CO)-R^c, wherein R^c is:

C₁₋₆-alkyl,
 15 –(CH₂)_n-NRⁱRⁱⁱ,
 –(CH₂)_n-NRⁱⁱⁱR^{iv},

5 or 6 membered heterocycloalkyl optionally substituted by C₁₋₆-alkyl,
 or is C₁₋₆-alkyl or aryl, which are optionally substituted by halo,

20 –O(CO)-C₁₋₆-alkyl,
 or by –NH(CO)R^d, wherein R^d is C₁₋₆-alkyl optionally substituted by halo or nitro, or R^d is aryl or a 5 or 6 membered heteroaryl, which are optionally substituted by halo, nitro, C₁₋₆-alkyl or C₁₋₆-haloalkyl;

A is selected from the group consisting of (a), (b), (c) and (d):




25 wherein

R⁴ is –NH(CO)R^e, wherein R^e is C₁₋₆-alkoxy, or aryl optionally substituted by halo, or is C₁₋₆-alkoxy, CN,
 30 or is aryl, 5 or 6 membered heteroaryl, benzyl, aryloxy or a 9 or 10-membered bicyclic heteroaryl ring which are optionally substituted by CN, halo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-haloalkyl, or by a dioxo bridge; and

- R⁵ is H, OH, CN, COORⁱⁱⁱ or CONRⁱⁱⁱR^{iv};
- R⁶ is C₁₋₆-alkyl,
 or is -C(O)-R^f wherein R^f is an aryl group optionally substituted by halo, C₁₋₆-alkoxy, or CN,
 5 or is aryl, 5 or 6 membered heteroaryl, benzyl, or a 9 or 10-membered bicyclic heteroaryl ring which are optionally substituted by halo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-haloalkyl, CN or by a dioxo bridge;
- B is halo, CN, NRⁱRⁱⁱ, C₁₋₆-alkyl optionally substituted by CN, halo or C₁₋₆-alkoxy, C₁₋₆-alkoxy, C₁₋₆-haloalkoxy, C₃₋₆-cycloalkyl, -C(O)O-C₁₋₆-alkyl, -C(O)NRⁱRⁱⁱ, -
 10 C(O)-C₁₋₆-alkyl, -S(O)₂-C₁₋₆-alkyl, -S(O)₂-NRⁱRⁱⁱ, (CRⁱⁱⁱR^{iv})_n-phenyl, or (CRⁱⁱⁱR^{iv})_n-5 or 6 membered heteroaryl wherein the phenyl or 5 or 6 membered heteroaryl moiety is optionally substituted by one or more substituent(s) selected from the group consisting of:
- halo, CN, NRⁱRⁱⁱ, C₁₋₆-alkyl optionally substituted by CN or C₁₋₆-alkoxy,
 15 C₁₋₆-alkoxy, C₁₋₆-haloalkoxy, C₃₋₆-cycloalkyl, -C(O)O-C₁₋₆-alkyl, -C(O)-NRⁱRⁱⁱ, -C(O)-C₁₋₆-alkyl, -S(O)₂-C₁₋₆-alkyl, -S(O)₂-NRⁱRⁱⁱ;
- Rⁱ and Rⁱⁱ are H, C₁₋₆-alkyl, C₁₋₆-alkyl-NRⁱⁱⁱR^{iv}, -(CO)O-C₁₋₆-alkyl, -C(O)-NRⁱⁱⁱR^{iv}, -C(O)-C₁₋₆-alkyl, -S(O)₂-C₁₋₆-alkyl or -S(O)₂-NRⁱⁱⁱR^{iv};
- Rⁱⁱⁱ and R^{iv} are H or C₁₋₆-alkyl;
- 20 m is 1 to 6;
 n is 0 to 4;
- as well as pharmaceutically acceptable salts thereof.

3. The compound of formula (I) according to claim 1 or 2, wherein
- 25 R¹ is H,
 or is C₁₋₆-alkyl optionally substituted by CN or by C₁₋₆-alkoxy,
 or is aryl,
 or is 5 or 6 membered heteroaryl,
 or is sulfonylaryl,
 30 or is -(CH₂)_m-R^a wherein R^a is C₃₋₆-cycloalkyl, 5 or 6 membered-heterocycloalkyl, aryl, or 5 or 6 membered heteroaryl which are optionally substituted by one or more substituents selected from the group consisting of:
- halo, CN, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-haloalkoxy, -C(O)O-C₁₋₆-alkyl
 35 and phenyl optionally substituted by halo, C₁₋₆-alkyl, C₁₋₆-haloalkyl or C₁₋₆-alkoxy,
 or is -(CH₂)_m-NRⁱRⁱⁱ,

- or is $-(\text{CH}_2)_n-(\text{CO})-\text{R}^b$, wherein R^b is aryl or 5 or 6 membered-heterocycloalkyl;
- R^2 is one or more of H, halo, CN, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-\text{O}-\text{CH}_2-\text{C}_{2-6}$ -alkenyl, benzyloxy, or two R^2 may form an oxo or dioxo bridge together with the indole ring to which they are attached;
- 5 R^3 is H,
or is halo,
or is $-(\text{CO})-\text{R}^c$, wherein R^c is C_{1-6} -alkyl, 5 or 6 membered heterocycloalkyl optionally substituted by C_{1-6} -alkyl, or R^c is $-(\text{CH}_2)_n-\text{NR}^i\text{R}^{ii}$ or,
or is C_{1-6} -alkyl or aryl, which are optionally substituted by
- 10 $-\text{O}(\text{CO})-\text{C}_{1-6}$ -alkyl,
or by $-\text{NH}(\text{CO})\text{R}^d$, wherein R^d is C_{1-6} -alkyl optionally substituted by halo or nitro, or R^d is aryl or a 5 or 6 membered heteroaryl, which are optionally substituted by halo, nitro, C_{1-6} -alkyl or C_{1-6} -haloalkyl;
- R^i and R^{ii} are independently selected from H, C_{1-6} -alkyl or $-(\text{CO})\text{O}-\text{C}_{1-6}$ -alkyl;
- 15 m is 1 to 6;
n is 0 to 4;
- A is selected from the group consisting of (a), (b), (c) and (d):



- wherein
- 20 R^4 is $-\text{NH}(\text{CO})\text{R}^e$, wherein R^e is C_{1-6} -alkoxy, or aryl optionally substituted by halo, or is aryl, benzyl, aryloxy or a 9 or 10-membered bicyclic heteroaryl ring which are optionally substituted by halo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -haloalkyl, or by a dioxo bridge; and
- R^5 is H, OH or CN;
- 25 R^6 is C_{1-6} -alkyl,
or is $-\text{C}(\text{O})-\text{R}^f$ wherein R^f is an aryl group optionally substituted by halo, or is aryl, benzyl, or a 9 or 10-membered bicyclic heteroaryl ring which are optionally substituted by halo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -haloalkyl, or by a dioxo bridge;
- 30 as well as pharmaceutically acceptable salts thereof.

4. A compound of the following general formula I according to any one of claims 1 to 3, wherein:

R^1 is H,

or is C_{1-6} -alkyl substituted by NR^iR^{ii} ,

5 or is $-(CH_2)_m-R^a$ wherein R^a is:

NR^iR^{ii} ,

5 to 6 membered-heterocycloalkyl, aryl, or 5 or 6 membered heteroaryl which are optionally substituted by one or more B,

or is $-(CH_2)_n-(CO)-R^b$ or $-(CH_2)_n-(SO_2)-R^b$, wherein R^b is:

10 C_{1-6} -alkyl,

C_{1-6} -alkoxy,

NR^iR^{ii} ,

5 to 6 membered-heterocycloalkyl, aryl, or 5 or 6 membered heteroaryl which are optionally substituted by one or more B,


15 or R^1 and R^3 together with the indole ring to which they are attached form a 5 or 6 membered heterocycloalkyl which can be substituted by =O, $C(O)O-C_{1-6}$ -alkyl or C_{1-6} -alkyl;

R^2 is one or more of H, halo, C_{1-6} -alkyl;

R^3 is H,

20 or is C_{1-6} -alkyl;

A is selected from the group consisting of (a), (a'), (b), (c) and (d):



wherein

25 R^4 is is $-NH(CO)R^e$, wherein R^e is C_{1-6} -alkoxy, or aryl optionally substituted by halo, or is aryl, 5 or 6 membered heteroaryl, benzyl, aryloxy or a 9 or 10-membered bicyclic heteroaryl ring, which are optionally substituted by halo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -haloalkyl, nitro, hydroxyl, C_{1-6} -haloalkoxy, or by a oxo or dioxo bridge; and


R^5 is H, OH, CN;

- R^6 is aryl, 5 or 6 membered heteroaryl, or a 9 or 10-membered bicyclic heteroaryl ring which are optionally substituted by halo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -haloalkyl, CN, nitro, NR^iR^{ii} , $NR^{iii}R^{iv}$, C_{1-6} -alkoxy- C_{1-6} -alkylene, COOH, $S(O)_2$ - C_{1-6} -alkyl, or by a oxo or dioxo bridge;
- 5 B is halo, C_{1-6} -alkyl optionally substituted by CN, halo or C_{1-6} -alkoxy, C_{1-6} -alkoxy; R^i and R^{ii} are H, C_{1-6} -alkyl, C_{1-6} -alkyl- $NR^{iii}R^{iv}$, $-C(O)-C_{1-6}$ -alkyl or $-S(O)_2-C_{1-6}$ -alkyl; R^{iii} and R^{iv} are H or C_{1-6} -alkyl;
- m is 1 to 6;
- n is 0 to 4;
- 10 as well as pharmaceutically acceptable salts thereof.

5. A compound of the general formula I according to any one of claims 1 to 4, with the proviso that R^1 , R^2 and R^3 are not simultaneously H, as well as pharmaceutical acceptable salts thereof.



15

6. A compound of the general formula I according to any one of claims 1 to 5, with the proviso that R^4 is not



optionally substituted by CN, halo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -haloalkyl, nitro, hydroxyl, NR^iR^{ii} , $NR^{iii}R^{iv}$, C_{1-6} -alkoxy- C_{1-6} -alkylene, $S(O)_2$ - C_{1-6} -alkyl, C_{1-6} -haloalkoxy, or by a oxo or dioxo bridge.


7. The compound of formula (I-a) or (I-a') according to any one of claims 1 to 6,



25

wherein R^1 to R^4 are as defined in any one of claims 1 to 6, as well as pharmaceutically acceptable salts thereof.

8. The compound of formula (I-a) or (I-a') according to claim 7,
wherein
- R¹ is H or -(CH₂)_m-R^a wherein R^a is aryl and m is 1 to 6;
- 5 R² is one or more of H or halo;
- R³ is H or C₁₋₆-alkyl;
- R⁴ is aryl which is optionally substituted by halo or C₁₋₆-alkoxy;
as well as pharmaceutically acceptable salts thereof.
- 10 9. The compound of formula (I-a) or formula (I-a') according to claim 1,
wherein it is selected from the group consisting of:
- (1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(4-fluoro-phenyl)-3,6-dihydro-2H-pyridin-1-yl]-
methanone;
- (1-Benzyl-2-methyl-1H-indol-3-yl)-(4-phenyl-3,6-dihydro-2H-pyridin-1-yl)-
15 methanone;
- (1-Benzyl-2-methyl-1H-indol-3-yl)-[4-(2-methoxy-phenyl)-3,6-dihydro-2H-pyridin-1-
yl]-methanone;
- (6-Chloro-1H-indol-3-yl)-(4-phenyl-3,6-dihydro-2H-pyridin-1-yl)-methanone;
- (6-Chloro-1H-indol-3-yl)-[4-(4-fluoro-phenyl)-3,6-dihydro-2H-pyridin-1-yl]-
20 methanone;
- (6-Chloro-1H-indol-3-yl)-(4-phenyl-3,6-dihydro-2H-pyridin-1-yl)-methanone; and
(1-Benzyl-2-methyl-1H-indol-3-yl)-(5-methyl-4-phenyl-3,6-dihydro-2H-pyridin-1-yl)-
methanone.
- 25 10. The compound of formula (I-b) according to anyone of claims 1 to 6,



wherein R¹ to R⁵ are as defined in any one of claims 1 to 6,
as well as pharmaceutically acceptable salts thereof.

11. The compound of formula (I-b) according to claim 10, wherein:
- R¹ is H,
 or is -(CH₂)_m-R^a wherein R^a is:
 NRRⁱRⁱⁱ,
 5 aryl, or 5 or 6 membered heteroaryl which are optionally substituted by one
 or more halo, C₁₋₆-alkyl, C₁₋₆-alkoxy,
 or is -(CH₂)_n-(CO)-R^b or -(CH₂)_n-(SO₂)-R^b, wherein R^b is:
 C₁₋₆-alkyl,
 C₁₋₆-alkoxy,
 10 NRRⁱRⁱⁱ,
 5 to 6 membered-heterocycloalkyl, or 5 or 6 membered heteroaryl which are
 optionally substituted by one or more halo, C₁₋₆-alkyl, C₁₋₆-alkoxy,
 or R¹ and R³ together with the indole ring to which they are attached form a 5 or
 6 membered heterocycloalkyl which can be substituted by C(O)O-C₁₋₆-alkyl
 15 or C₁₋₆-alkyl;
 R² is one or more of H, halo or C₁₋₆-alkyl;
 R³ is H or C₁₋₆-alkyl;
 R⁴ is is -NH(CO)R^e, wherein R^e is C₁₋₆-alkoxy or aryl which is optionally substituted
 by halo,
 20 or is aryl, benzyl, aryloxy or a 9 or 10-membered bicyclic heteroaryl ring which
 are optionally substituted by halo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-haloalkyl,
 nitro, hydroxyl, C₁₋₆-haloalkoxy or by a oxo or dioxo bridge; and
 R⁵ is H, OH or CN;
 Rⁱ and Rⁱⁱ are H, C₁₋₆-alkyl, C₁₋₆-alkyl-NRⁱⁱⁱR^{iv};
 25 Rⁱⁱⁱ and R^{iv} are H or C₁₋₆-alkyl;
 m is 1 to 6;
 n is 0 to 4;
 as well as pharmaceutically acceptable salts thereof.

12. The compound of formula (I-b) according to claim 10 or 11, wherein it is
 30 selected from the group consisting of:
 (1-Benzyl-2-methyl-1*H*-indol-3-yl)-(4-phenyl-piperidin-1-yl)-methanone;
 (6-Chloro-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-
 methanone;
 2-{6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-
 35 yl}-*N,N*-dimethyl-acetamide;

- 2- {6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-pyridin-2-yl-ethanone;
- (6-Chloro-1-pyridin-4-ylmethyl-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone;
- 5 (6-Chloro-1-pyridin-3-ylmethyl-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone;
- (6-Chloro-1-pyridin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone;
- [6-Chloro-1-(6-chloro-pyridin-3-ylmethyl)-1*H*-indol-3-yl]-[4-(2,6-
- 10 dimethoxy-phenyl)-piperidin-1-yl]-methanone;
- {6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid *tert*-butyl ester;
- (6-Chloro-1-pyrazin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone;
- 15 [6-Chloro-1-(3,5-difluoro-benzyl)-1*H*-indol-3-yl]-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone;
- [6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone;
- 2- {6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-
- 20 yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- [6-Chloro-1-(3,5-difluoro-benzyl)-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone;
- 2- {6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 25 2- {6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- {6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetic acid *tert*-butyl ester;
- 2- {6-Chloro-3-[4-cyano-4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-
- 30 indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2- {6-Chloro-3-[4-cyano-4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2- {3-[4-(2,6-Dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 35 [6-Chloro-1-(2-methyl-pyridin-4-ylmethyl)-1*H*-indol-3-yl]-[4-(2,6-dimethoxy-phenyl)-piperidin-1-yl]-methanone;

- 2- {6-Chloro-3-[4-(2-isopropoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2- {6-Chloro-3-[4-(2-isopropoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- 5 2- {6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone;
- 2- {6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- 2- {6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 10 indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2- {6-Chloro-3-[4-(2,3-dihydro-benzofuran-7-yl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- 2- {5,6-Dichloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 15 2- {6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide;
- 2- {6-Chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- N*-(2-Amino-ethyl)-2- {6-chloro-3-[4-(2,6-dimethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-acetamide;
- 20 carbonyl]-indol-1-yl}-acetamide;
- [6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone;
- 2- {6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- 25 2- {6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- 2- {6-Chloro-3-[4-(2-trifluoromethoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 2- {6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- 30 1-yl}-*N*-methyl-acetamide;
- 2- {6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;
- 2- {6-Chloro-3-[4-(2-trifluoromethyl-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 35 2- {6-Chloro-3-[4-(2-methoxy-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;

[6-Chloro-1-(2-methylamino-ethyl)-1*H*-indol-3-yl]-[4-(2,3-dihydro-benzofuran-7-yl)-piperidin-1-yl]-methanone;

[6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperidin-1-yl]-methanone;

5 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;


10 [6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperidin-1-yl]-methanone;

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperidine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone or hydrochloride salt thereof; and

15 2-[6-Chloro-3-(3,4,5,6-tetrahydro-2*H*-[4,4']bipyridinyl-1-carbonyl)-indol-1-yl]-*N,N*-dimethyl-acetamide.

13. The compound of formula (I-c) according to any one of claims 1 to 6,



20 wherein R^1 to R^3 and R^6 are as defined in any one of claims 1 to 6, as well as pharmaceutically acceptable salts thereof.

14. The compound of formula (I-c) according to claim 13, wherein R^1 is H or

25 or is C_{1-6} -alkyl substituted by NR^iR^{ii} ,

or is $-(CH_2)_m-R^a$ wherein R^a is:

NR^iR^{ii} ,

5 to 6 membered-heterocycloalkyl, aryl, or 5 or 6 membered heteroaryl which are optionally substituted by one or more C_{1-6} -alkyl,

30 or is $-(CH_2)_n-(CO)-R^b$, wherein R^b is:

NR^iR^{ii} ,

5 to 7 membered-heterocycloalkyl which is optionally substituted by one or more C_{1-6} -alkyl;

R^2 is one or more of H, halo or C_{1-6} -alkyl;

5 R^3 is H or C_{1-6} -alkyl;

R^6 is aryl, 5 or 6 membered heteroaryl, or a 9 or 10-membered bicyclic heteroaryl ring which are optionally substituted by halo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -haloalkyl, CN, nitro, NR^iR^{ii} , $\text{NR}^{iii}\text{R}^{iv}$, C_{1-6} -alkoxy- C_{1-6} -alkylene, COOH, $\text{S}(\text{O})_2$ - C_{1-6} -alkyl, or by a oxo or dioxo bridge;

10 R^i and R^{ii} are H, C_{1-6} -alkyl, C_{1-6} -alkyl- $\text{NR}^{iii}\text{R}^{iv}$, $-\text{C}(\text{O})-\text{C}_{1-6}$ -alkyl, or $-\text{S}(\text{O})_2-\text{C}_{1-6}$ -alkyl;

R^{iii} and R^{iv} are H or C_{1-6} -alkyl;

m is 1 to 6;

n is 0 to 4;

as well as pharmaceutically acceptable salts thereof.

15 15. The compound of formula (I-c) according to claim 13, wherein it is selected from the group consisting of:

2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;

20 2-{6-Chloro-3-[4-(2-ethoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;

2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;

2-{6-Chloro-3-[4-(2-methoxy-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;

25 [6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-methoxy-phenyl)-piperazin-1-yl]-methanone;

[6-Chloro-1-(2-dimethylamino-ethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;

30 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-(4-methyl-piperazin-1-yl)-ethanone;


2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;

2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-acetamide;

35 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;

- 2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-1-piperazin-1-yl-ethanone;
- 2-{6-Chloro-3-[4-(2,4-difluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- 5 2-{6-Chloro-3-[4-(3-trifluoromethyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide;
- [1-((*S*)-2-Amino-propyl)-6-chloro-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- (6-Chloro-1-(*S*)-1-pyrrolidin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- 10 [6-Chloro-1-((*S*)-1-methyl-pyrrolidin-2-ylmethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- 2-{6-Chloro-3-[4-(2,4-difluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- 15 2-{6-Chloro-3-[4-(3-trifluoromethyl-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl}-*N,N*-dimethyl-acetamide;
- (6-Chloro-1-(*R*)-1-pyrrolidin-2-ylmethyl-1*H*-indol-3-yl)-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- [6-Chloro-1-((*R*)-1-methyl-pyrrolidin-2-ylmethyl)-1*H*-indol-3-yl]-[4-(2-fluoro-phenyl)-piperazin-1-yl]-methanone;
- 20 *N*-(2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-ethyl)-acetamide;
- N*-(2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-ethyl)-methanesulfonamide;
- 25 *N*-(2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-ethyl)-*N*-methyl-acetamide;
- N*-(2-{6-Chloro-3-[4-(2-fluoro-phenyl)-piperazine-1-carbonyl]-indol-1-yl}-ethyl)-*N*-methyl-methanesulfonamide;
- 2-[6-Chloro-3-(4-thieno[3,2-*c*]pyridin-4-yl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide; and
- 30 2-{6-Chloro-3-[4-(3-iodo-pyridin-2-yl)-piperazine-1-carbonyl]-indol-1-yl}-*N*-methyl-acetamide.

16. The compound of formula (I-d) according to any one of claims 1 to 6,



wherein R^1 to R^3 and R^6 are as defined in any one of claims 1 to 6, as well as pharmaceutically acceptable salts thereof.

5 17. The compound of formula (I-d) according to claim 16, wherein:

R^1 is H;

R^2 is one or more halo;

R^3 is H;


R^6 is aryl substituted by C_{1-6} -alkoxy;

10 as well as pharmaceutically acceptable salts thereof.

18. The compound of formula (I-d) according to claim 16, wherein it is


(6-Chloro-1H-indol-3-yl)-[4-(2-methoxy-phenyl)-4-oxy-piperazin-1-yl]-methanone.

15 19. A process for the preparation of compounds of formula (I) according to any one of claims 1 to 18, comprising the step of reacting a compound of formula (II):




with a compound of formula A-H, in order to obtain a compound of formula (I) wherein A, R^1 , R^2 and R^3 are as defined in claim 1.

20. A process for the preparation of compounds of formula (I) according to any one of claims 1 to 18 comprising the step of reacting a compound of formula (I-1):




5 with a compound of formula R¹-X, to obtain a compound of formula (I) wherein R¹ to R³ and A are as defined in claim 1 and X is halo.

21. A process for the preparation of compounds of formula (I) according to any one of claims 1 to 18 comprising the step of hydrolyzing a compound of formula (V):



10

to obtain a compound of formula (II):



15 and then reacting the compound of formula (II) with a compound of formula A-H, to obtain a compound of formula (I) wherein A, R¹, R² and R³ are as defined in claim 1.

22. A compound formula (I-a), (I-a'), (I-b), (I-c) or (I-d), obtainable by a process according to claim 19 to 21.
23. A compound of formula (I-a), (I-a'), (I-b), (I-c) or (I-d) according to any one of claims 1 to 18 for a use in the prevention or treatment of dysmenorrhea, hypertension, chronic heart failure, inappropriate secretion of vasopressin, liver cirrhosis, nephrotic syndrome, obsessive compulsive disorder, anxious and depressive disorders.
24. A pharmaceutical composition comprising a compound of formula (I-a), (I-a'), (I-b), (I-c) or (I-d) according to any one of claims 1 to 18.
25. A pharmaceutical composition according to claim 24, wherein it is useful against dysmenorrhea, hypertension, chronic heart failure, inappropriate secretion of vasopressin, liver cirrhosis, nephrotic syndrome, obsessive compulsive disorder, anxious and depressive disorders.
26. Use of a compound of formula (I-a), (I-a'), (I-b), (I-c) or (I-d) according to any one of claims 1 to 18 for the preparation of a medicament.
27. Use according to claim 26, wherein the medicament is useful against dysmenorrhea, hypertension, chronic heart failure, inappropriate secretion of vasopressin, liver cirrhosis, nephrotic syndrome, obsessive compulsive disorder, anxious and depressive disorders.
28. The invention as described hereinabove.