




(86) **Date de dépôt PCT/PCT Filing Date:** 2012/09/13
(87) **Date publication PCT/PCT Publication Date:** 2013/03/21
(45) **Date de délivrance/Issue Date:** 2020/03/24
(85) **Entrée phase nationale/National Entry:** 2014/03/12
(86) **N° demande PCT/PCT Application No.:** US 2012/055248
(87) **N° publication PCT/PCT Publication No.:** 2013/040269
(30) **Priorité/Priority:** 2011/09/13 (US61/534,194)

(51) **Cl.Int./Int.Cl.** **A61K 31/575** (2006.01),
A61P 17/02 (2006.01), **A61P 31/00** (2006.01)
(72) **Inventeurs/Inventors:**
GENBERG, CARL, US;
SAVAGE, PAUL B., US
(73) **Propriétaire/Owner:**
BRIGHAM YOUNG UNIVERSITY, US
(74) **Agent:** OSLER, HOSKIN & HARCOURT LLP

(54) **Titre : PROCÉDES ET PRODUITS POUR AUGMENTER LA VITESSE DE GUÉRISON DE PLAIES TISSULAIRES**
(54) **Title: PRODUCTS FOR HEALING OF TISSUE WOUNDS**



(57) **Abrégé/Abstract:**

Disclosed are methods for increasing the rate of healing of a tissue wound by administering a composition including a therapeutically effective amount of at least one cationic steroid antimicrobial (CSA). Also disclosed herein are methods of promoting wound healing in a subject in need of such promotion, comprising administering a composition comprising a therapeutically effective amount of at least one CSA. Additionally, disclosed herein are compounds and compositions comprising at least one CSA, or a pharmaceutically acceptable salt thereof, for use in the treatment of a tissue wound. Kits comprising such compositions and instructions on such methods are also contemplated herein.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



WIPO | PCT



(10) International Publication Number
WO 2013/040269 A1

(43) International Publication Date
21 March 2013 (21.03.2013)

(51) International Patent Classification:

A61K 31/575 (2006.01) *A61P 31/00* (2006.01)
A61P 17/02 (2006.01)

(21) International Application Number:

PCT/US2012/055248

(22) International Filing Date:

13 September 2012 (13.09.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/534,194 13 September 2011 (13.09.2011) US

(71) Applicant (for all designated States except US):

BRIGHAM YOUNG UNIVERSITY [US/US]; Technology Commercialization Office, 3760 Harold B. Lee Library, Provo, Utah 84602-6844 (US).

(72) Inventors; and

(75) **Inventors/Applicants (for US only): GENBERG, Carl** [US/US]; Las Vegas, Nevada 89120 (US). **SAVAGE, Paul, B.** [US/US]; Mapleton, Utah 84664 (US).

(74) **Agent: HANSEN, Andrew, S.**; Workman Nydegger, 60 E South Temple, Suite 1000, Salt Lake City, Utah, 84111 (US).

(81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) **Title:** PRODUCTS FOR HEALING OF TISSUE WOUNDS

(57) **Abstract:** Disclosed are methods for increasing the rate of healing of a tissue wound by administering a composition including a therapeutically effective amount of at least one cationic steroid antimicrobial (CSA). Also disclosed herein are methods of promoting wound healing in a subject in need of such promotion, comprising administering a composition comprising a therapeutically effective amount of at least one CSA. Additionally, disclosed herein are compounds and compositions comprising at least one CSA, or a pharmaceutically acceptable salt thereof, for use in the treatment of a tissue wound. Kits comprising such compositions and instructions on such methods are also contemplated herein.



WO 2013/040269 A1

PRODUCTS FOR HEALING OF TISSUE WOUNDS

[0001] This Application claims the benefit of US Provisional Patent Application No. 61/534194, filed September 13, 2011 and titled "Compositions and Methods For Wound Healing".

BACKGROUND OF THE INVENTION

1. Field of the Invention

[0002] This invention relates to compositions and methods for increasing the rate of healing of tissue wounds in animals using cationic steroid antimicrobials (CSAs).

2. Relevant Technology

[0003] Wound healing is the complex and dynamic process of self-repair to restore cellular structures and tissue layers after an injury has occurred. The human wound healing process is divided into three distinct phases: the inflammatory phase, the proliferative phase, and the remodeling phase. The wound healing process may be further broken down beyond the three broad phases to a complex and coordinated series of biochemical events that include chemotaxis, phagocytosis, neocollagenesis, collagen degradation, and collagen remodeling to repair the damage. In addition, angiogenesis, epithelization, and the production of new glycosaminoglycans (GAGs) and proteoglycans are vital to the wound healing process. The culmination of these biological processes results in the replacement of normal skin structures with fibroblastic mediated scar tissue.

[0004] Prior to the initiation of the inflammatory phase, the clotting cascade begins to stop blood loss via clotting. Once clotting has begun, various soluble factors, including chemokines and cytokines, are released to attract cells to the site of injury. Growth factors, which are cytokines released by platelets, stimulate cells to accelerate their rate of division leading to increased healing rates. Platelets also release other proinflammatory factors like serotonin, bradykinin, prostaglandins, prostacyclins, thromboxane, and histamine, which serve a number of purposes, including to increase cell proliferation and migration to the area and to cause blood vessels to become dilated and porous.

[0005] The inflammation phase can be further broken down into four biochemical processes. First, immediately after a blood vessel is breached, ruptured cell membranes release inflammatory factors like thromboxanes and prostaglandins that result in vasoconstriction to prevent blood loss and to collect inflammatory cells and factors. Second, polymorphonuclear neutrophils (PMNs) arrive at the wound site, phagocytose debris and bacteria. Neutrophils

also cleanse the wound by secreting proteases that break down damaged tissue. Third, neutrophils cleanse the injured area via the phagocytosis of debris, bacteria, neutrophils, damaged cells and tissue, and any other foreign cells or material. Macrophages also secrete a number of factors that push wound healing toward the next phase. Finally, inflammation dies
5 down, fewer inflammatory factors are secreted, existing ones are broken down, and numbers of neutrophils and macrophages are reduced at the wound site.

[0006] The proliferative phase begins with the arrival of fibroblasts at the wound site, marking the onset of the proliferative phase, which biologically begins with angiogenesis. Angiogenesis or neovascularization is the process which occurs concurrently with fibroblast
10 proliferation when endothelial cells migrate to the area of the wound. Fibroplasia and granulation begin during angiogenesis as a result of fibroblast recruitment and extracellular matrix formation. Then collagen and fibronectin are deposited by the fibroblast at the site of tissue granulation. Once collagen deposition has occurred, epithelialization takes place as a result of the presence of keratinocytes arriving at the newly deposited extracellular matrix.

15 The final step in the proliferative phase is contraction, which requires the presence of neomyoblasts to connect the scar tissue to the surrounding healthy tissue and to pull the edges of the scar tissue together, restoring tensile strength to the skin surrounding the previously injured area.

[0007] A need exists to develop compositions and methods to enhance wound healing.

BRIEF SUMMARY

[0008] Disclosed herein are methods and compositions for treating a wound in a subject by increasing the rate of healing. The increased rate of healing is achieved using a cationic steroidal antimicrobial (CSA) compound. The CSA compounds include a sterol backbone and a plurality of cationic groups attached thereto.


[0009] The treatment of tissue wounds with CSAs has the surprising and unexpected result of increasing the rate of wound healing. Experimental data indicates that the increased rate of wound healing is distinct from the known anti-microbial effects of CSAs. When applied to wound sites in living subjects, the CSA compounds described herein have a profound and surprising effect on the rate of wound healing. The increased rates of healing is in many cases several times faster than healing rates using traditional antimicrobials. Studies indicate that this increase in wound healing is at least in part distinct from antimicrobial benefits. CSAs have been found to increase the rate of healing even in wounds where antimicrobial load is very low or in cases where infection is not an issue.

[0010] The increased rate of healing results in healthier and more natural tissue formation as compared to treatments with traditional antimicrobial compounds. The mechanism of action for the increased rate of healing is presently being studied. While the present invention is not limited to any particular mechanism, it is believed that the increased rate of tissue healing is caused by increases in fibroblastic migration and enhanced epithelial growth factors at the wound site. Subjects have also exhibited a significantly reduced sensitivity to pain. These benefits are unexpected for CSAs, which were thought to only be active against foreign microbes.

[0011] In some embodiments, a method is described for increasing the rate of wound healing in a subject. The method includes (i) providing a tissue treatment composition including a cationic steroidal anti-microbial (CSA) compound, the CSA compound including a steroidal group and a plurality of cationic groups attached thereto; (ii) identifying a subject in need of accelerated healing of a tissue wound; and (iii) contacting the tissue wound with the tissue treatment composition to increase the rate of healing thereof.


[0012] In some embodiments, the invention relates to a CSA for use in a novel method of treating wounds. In particular, the invention relates to a pharmaceutically acceptable CSA compound having a sterol backbone and a plurality of cationic groups attached thereto, wherein the CSA compound is for use in increasing the rate of wound healing in a subject.

[0013] In some embodiments, the CSA is a compound of Formula (V) or a pharmaceutically acceptable salt thereof:



(V).


[0014] In some embodiments, the CSA, or a pharmaceutically acceptable salt thereof, is selected from the compound of Formula (I):



(I).

5

[0015] In some embodiments, the CSA, or a pharmaceutically acceptable salt thereof, is selected from the compound of Formula (Ia):



(Ia).

[0016] In some embodiments rings A, B, C, and D are independently saturated, or are fully or partially unsaturated, provided that at least two of rings A, B, C, and D are saturated; m, n, p, and q are independently 0 or 1; R₁ through R₄, R₆, R₇, R₁₁, R₁₂, R₁₅, R₁₆, and R₁₈ are independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted alkyl, a substituted or unsubstituted hydroxyalkyl, a substituted or unsubstituted alkyloxyalkyl, a substituted or unsubstituted alkylcarboxyalkyl, a substituted or unsubstituted alkylaminoalkyl, a substituted or unsubstituted alkylaminoalkylamino, a substituted or unsubstituted aminoalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted arylaminoalkyl, a substituted or unsubstituted haloalkyl, a substituted or unsubstituted alkenyl, a substituted or

unsubstituted alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted aminoalkoxy, a substituted or unsubstituted aminoalkoxyalkyl, a substituted or unsubstituted aminoalkylcarboxy, a substituted or unsubstituted aminoalkylaminocarbonyl, a substituted or unsubstituted aminoalkylcarboxamido, a substituted or unsubstituted di(alkyl)aminoalkyl, a substituted or unsubstituted C-carboxyalkyl, $\text{H}_2\text{N}-\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{O}-$, $\text{H}_2\text{N}-\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{N}(\text{H})-$, a substituted or unsubstituted azidoalkoxy, a substituted or unsubstituted cyanoalkoxy, $\text{P.G.}-\text{HN}-\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{O}-$, a substituted or unsubstituted guanidinoalkoxy, a substituted or unsubstituted quaternaryammoniumalkylcarboxy, and a substituted or unsubstituted guanidinoalkyl carboxy, where Q_5 is a side chain of any amino acid (including a side chain of glycine, i.e., H), and P.G. is an amino protecting group; and R_5 , R_8 , R_9 , R_{10} , R_{13} , R_{14} and R_{17} are independently deleted when one of rings A, B, C, or D is unsaturated so as to complete the valency of the carbon atom at that site, or R_5 , R_8 , R_9 , R_{10} , R_{13} , and R_{14} are independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted alkyl, a substituted or unsubstituted hydroxyalkyl, a substituted or unsubstituted alkyloxyalkyl, a substituted or unsubstituted aminoalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted haloalkyl, a substituted or unsubstituted alkenyl, a substituted or unsubstituted alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted aminoalkoxy, a substituted or unsubstituted aminoalkylcarboxy, a substituted or unsubstituted aminoalkylaminocarbonyl, a substituted or unsubstituted di(alkyl)aminoalkyl, a substituted or unsubstituted C-carboxyalkyl, $\text{H}_2\text{N}-\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{O}-$, $\text{H}_2\text{N}-\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{N}(\text{H})-$, azidoalkoxy, cyanoalkoxy, $\text{P.G.}-\text{HN}-\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{O}-$, guanidinoalkoxy, and guanidinoalkylcarboxy, where Q_5 is a side chain of any amino acid, P.G. is an amino protecting group, provided that at least two or three of R_{1-4} , R_6 , R_7 , R_{11} , R_{12} , R_{15} , R_{16} , R_{17} , and R_{18} are independently selected from the group consisting of a substituted or unsubstituted aminoalkyl, a substituted or unsubstituted aminoalkoxy, a substituted or unsubstituted alkylcarboxyalkyl, a substituted or unsubstituted alkylaminoalkylamino, a substituted or unsubstituted alkylaminoalkylaminoalkylamino, a substituted or unsubstituted aminoalkylcarboxy, a substituted or unsubstituted arylaminoalkyl, a substituted or unsubstituted aminoalkoxyaminoalkylaminocarbonyl, a substituted or unsubstituted aminoalkylaminocarbonyl, a substituted or unsubstituted aminoalkylcarboxyamido, a quaternaryammoniumalkylcarboxy, a substituted or unsubstituted di(alkyl)aminoalkyl, a substituted or unsubstituted C-carboxyalkyl, $\text{H}_2\text{N}-\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{O}-$, $\text{H}_2\text{N}-\text{HC}(\text{Q}_5)-\text{C}(\text{O})-$

N(H)-, azidoalkyloxy, cyanoalkyloxy, P.G.-HN-HC(Q5)-C(O)—O—, a substituted or unsubstituted guanidinoalkyloxy, and a substituted or unsubstituted guanidinoalkylcarboxy.

[0017] In some embodiments, R₁ through R₄, R₆, R₇, R₁₁, R₁₂, R₁₅, R₁₆, and R₁₈ are independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted (C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈) hydroxyalkyl, a substituted or unsubstituted (C₁-C₁₈) alkyloxy-(C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈) alkylcarboxy-(C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino, a substituted or unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino- (C₁-C₁₈) alkylamino, a substituted or unsubstituted (C₁-C₁₈) aminoalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted arylamino-(C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈) haloalkyl, a substituted or unsubstituted C₂-C₆ alkenyl, a substituted or unsubstituted C₂-C₆ alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted (C₁-C₁₈) aminoalkyloxy, a substituted or unsubstituted (C₁-C₁₈) aminoalkyloxy-(C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkylcarboxy, a substituted or unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkylcarboxamido, a substituted or unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, a substituted or unsubstituted C-carboxy(C₁-C₁₈)alkyl, H₂N-HC(Q5)-C(O)-O—, H₂N—HC(Q5)-C(O)—N(H)—, a substituted or unsubstituted (C₁-C₁₈) azidoalkyloxy, a substituted or unsubstituted (C₁-C₁₈) cyanoalkyloxy, P.G.-HN—HC(Q5)-C(O)—O—, a substituted or unsubstituted (C₁-C₁₈) guanidinoalkyloxy, a substituted or unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, and a substituted or unsubstituted (C₁-C₁₈) guanidinoalkyl carboxy, where Q₅ is a side chain of any amino acid (including a side chain of glycine, i.e., H), and P.G. is an amino protecting group; R₅, R₈, R₉, R₁₀, R₁₃, R₁₄ and R₁₇ are independently deleted when one of rings A, B, C, or D is unsaturated so as to complete the valency of the carbon atom at that site, or R₅, R₈, R₉, R₁₀, R₁₃, and R₁₄ are independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted (C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈) hydroxyalkyl, a substituted or unsubstituted (C₁-C₁₈) alkyloxy-(C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted (C₁-C₁₈) haloalkyl, a substituted or unsubstituted (C₂-C₆) alkenyl, a substituted or unsubstituted (C₂-C₆) alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted (C₁-C₁₈) aminoalkyloxy, a substituted or unsubstituted (C₁-C₁₈) aminoalkylcarboxy, a substituted or unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl, a substituted or unsubstituted di(C₁-C₁₈

alkyl)aminoalkyl, a substituted or unsubstituted C-carboxy(C₁-C₁₈)alkyl, H₂N—HC(Q₅)-C(O)—O—, H₂N—HC(Q₅)-C(O)—N(H)—, a substituted or unsubstituted (C₁-C₁₈) azidoalkyloxy, a substituted or unsubstituted (C₁-C₁₈) cyanoalkyloxy, P.G.-HN—HC(Q₅)-C(O)—O—, a substituted or unsubstituted (C₁-C₁₈) guanidinoalkyloxy, and (C₁-C₁₈)
 5 guanidinoalkylcarboxy, where Q₅ is a side chain of any amino acid, and P.G. is an amino protecting group; provided that at least two or three of R₁₋₄, R₆, R₇, R₁₁, R₁₂, R₁₅, R₁₆, R₁₇, and R₁₈ are independently selected from the group consisting of a substituted or unsubstituted (C₁-C₁₈) aminoalkyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkyloxy, a substituted or unsubstituted (C₁-C₁₈) alkylcarboxy-(C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈)
 10 alkylamino-(C₁-C₁₈) alkylamino, a substituted or unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino (C₁-C₁₈) alkylamino, a substituted or unsubstituted (C₁-C₁₈) aminoalkylcarboxy, a substituted or unsubstituted arylamino (C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkyloxy (C₁-C₁₈) aminoalkylaminocarbonyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkylcarboxyamido,
 15 a substituted or unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, a substituted or unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, a substituted or unsubstituted C-carboxy(C₁-C₁₈)alkyl, H₂N-HC(Q₅)-C(O)-O—, H₂N-HC(Q₅)-C(O)—N(H)—, a substituted or unsubstituted (C₁-C₁₈) azidoalkyloxy, a substituted or unsubstituted (C₁-C₁₈) cyanoalkyloxy, P.G.-HN-HC(Q₅)-C(O)—O—, a substituted or unsubstituted (C₁-C₁₈) guanidinoalkyloxy,
 20 and a substituted or unsubstituted (C₁-C₁₈) guanidinoalkylcarboxy.

[0018] In some embodiments, R₁ through R₄, R₆, R₇, R₁₁, R₁₂, R₁₅, R₁₆, and R₁₈ are independently selected from the group consisting of hydrogen, hydroxyl, an unsubstituted (C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) hydroxyalkyl, unsubstituted (C₁-C₁₈) alkyloxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylcarboxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino, (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino- (C₁-C₁₈) alkylamino, an unsubstituted (C₁-C₁₈) aminoalkyl, an unsubstituted aryl, an unsubstituted arylamino-(C₁-C₁₈) alkyl, oxo, an unsubstituted (C₁-C₁₈) aminoalkyloxy, an unsubstituted (C₁-C₁₈) aminoalkyloxy-(C₁-C₁₈) alkyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxy, an unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl, an
 25 unsubstituted (C₁-C₁₈) aminoalkylcarboxamido, an unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, a substituted or unsubstituted C-carboxy(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) guanidinoalkyloxy, unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, and unsubstituted (C₁-C₁₈) guanidinoalkyl carboxy; R₅, R₈, R₉, R₁₀, R₁₃, R₁₄ and R₁₇ are independently deleted when one of rings A, B, C, or D is unsaturated so as to complete the

valency of the carbon atom at that site, or R₅, R₈, R₉, R₁₀, R₁₃, and R₁₄ are independently selected from the group consisting of hydrogen, hydroxyl, an unsubstituted (C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) hydroxyalkyl, unsubstituted (C₁-C₁₈) alkyloxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylcarboxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈)alkyl, (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino- (C₁-C₁₈) alkylamino, an unsubstituted (C₁-C₁₈) aminoalkyl, an unsubstituted aryl, an unsubstituted arylamino-(C₁-C₁₈) alkyl, oxo, an unsubstituted (C₁-C₁₈) aminoalkyloxy, an unsubstituted (C₁-C₁₈) aminoalkyloxy-(C₁-C₁₈) alkyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxy, an unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxamido, an unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, a substituted or unsubstituted C-carboxy(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) guanidinoalkyloxy, unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, and unsubstituted (C₁-C₁₈) guanidinoalkyl carboxy; provided that at least two or three of R₁₋₄, R₆, R₇, R₁₁, R₁₂, R₁₅, R₁₆, R₁₇, and R₁₈ are independently selected from the group consisting of hydrogen, hydroxyl, an unsubstituted (C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) hydroxyalkyl, unsubstituted (C₁-C₁₈) alkyloxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylcarboxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino- (C₁-C₁₈) alkylamino, an unsubstituted (C₁-C₁₈) aminoalkyl, an unsubstituted aryl, an unsubstituted arylamino-(C₁-C₁₈) alkyl, oxo, an unsubstituted (C₁-C₁₈) aminoalkyloxy, an unsubstituted (C₁-C₁₈) aminoalkyloxy-(C₁-C₁₈) alkyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxy, an unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxamido, an unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, a substituted or unsubstituted C-carboxy(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) guanidinoalkyloxy, unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, and unsubstituted (C₁-C₁₈) guanidinoalkyl carboxy.

[0019] In some embodiments, R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of hydrogen, an unsubstituted (C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) hydroxyalkyl, unsubstituted (C₁-C₁₈) alkyloxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylcarboxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino- (C₁-C₁₈) alkylamino, an unsubstituted (C₁-C₁₈) aminoalkyl, an unsubstituted arylamino-(C₁-C₁₈) alkyl, an unsubstituted (C₁-C₁₈) aminoalkyloxy, an unsubstituted (C₁-C₁₈) aminoalkyloxy-(C₁-C₁₈) alkyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxy, an unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxamido, an unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, a substituted or unsubstituted C-carboxy(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) guanidinoalkyloxy, unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, and unsubstituted (C₁-C₁₈) guanidinoalkyl carboxy.

aminoalkylaminocarbonyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxamido, an unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, a substituted or unsubstituted C-carboxy(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) guanidinoalkyloxy, unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, and unsubstituted (C₁-C₁₈) guanidinoalkyl carboxy; and
 5 R₁, R₂, R₄, R₅, R₆, R₈, R₉, R₁₀, R₁₁, R₁₃, R₁₄, R₁₅, R₁₆, and R₁₇ are independently selected from the group consisting of hydrogen and unsubstituted (C₁-C₆) alkyl.

[0020] In some embodiments, R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of hydrogen, an unsubstituted (C₁-C₆) alkyl, unsubstituted (C₁-C₆) hydroxyalkyl, unsubstituted (C₁-C₁₆) alkyloxy-(C₁-C₅) alkyl, unsubstituted (C₁-C₁₆) alkylcarboxy-(C₁-C₅)
 10 alkyl, unsubstituted (C₁-C₁₆) alkylamino-(C₁-C₅)alkyl, (C₁-C₁₆) alkylamino-(C₁-C₅) alkylamino, unsubstituted (C₁-C₁₆) alkylamino-(C₁-C₁₆) alkylamino-(C₁-C₅) alkylamino, an unsubstituted (C₁-C₁₆) aminoalkyl, an unsubstituted arylamino-(C₁-C₅) alkyl, an unsubstituted (C₁-C₅) aminoalkyloxy, an unsubstituted (C₁-C₁₆) aminoalkyloxy-(C₁-C₅) alkyl, an unsubstituted (C₁-C₅) aminoalkylcarboxy, an unsubstituted (C₁-C₅)
 15 aminoalkylaminocarbonyl, an unsubstituted (C₁-C₅) aminoalkylcarboxamido, an unsubstituted di(C₁-C₅ alkyl)amino-(C₁-C₅) alkyl, a substituted or unsubstituted C-carboxy(C₁-C₁₈)alkyl, unsubstituted (C₁-C₅) guanidinoalkyloxy, unsubstituted (C₁-C₁₆) quaternaryammoniumalkylcarboxy, and unsubstituted (C₁-C₁₆) guanidinoalkylcarboxy;

[0021] In some embodiments, R₁, R₂, R₄, R₅, R₆, R₈, R₁₀, R₁₁, R₁₄, R₁₆, and R₁₇ are each
 20 hydrogen; and R₉ and R₁₃ are each methyl.

[0022] In some embodiments, R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of aminoalkyloxy; aminoalkylcarboxy; alkylaminoalkyl; alkoxycarbonylalkyl; alkylcarbonylalkyl; di(alkyl)aminoalkyl; C-carboxyalkyl, alkoxycarbonylalkyl; and alkylcarboxyalkyl.





[0023] In some embodiments, R₃, R₇, and R₁₂ are independently selected from the group consisting of aminoalkyloxy and aminoalkylcarboxy; and R₁₈ is selected from the group consisting of alkylaminoalkyl; alkoxycarbonylalkyl; alkylcarbonyloxyalkyl; di(alkyl)aminoalkyl; C-carboxyalkyl, alkylaminoalkyl; alkyoxycarbonylalkyl; and alkylcarboxyalkyl.


[0024] In some embodiments, R₃, R₇, and R₁₂ are the same. In some embodiments, R₃, R₇, and R₁₂ are aminoalkyloxy. In some embodiments, R₃, R₇, and R₁₂ are aminoalkylcarboxy.

[0025] In some embodiments, R₁₈ is alkylaminoalkyl. In some embodiments, alkoxycarbonylalkyl. In some embodiments, R₁₈ is di(alkyl)aminoalkyl. In some embodiments, R₁₈ is alkylcarboxyalkyl. In some embodiments R₁₈ is C-carboxyalkyl.


[0026] In some embodiments, R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of amino-C₃-alkyloxy; amino-C₃-alkyl-carboxy; C₈-alkylamino-C₅-alkyl; C₈-alkoxy-carbonyl-C₄-alkyl; C₈-alkyl-carbonyl-C₄-alkyl; di-(C₅-alkyl)amino-C₅-alkyl; C₁₃-alkylamino-C₅-alkyl; C₆-alkoxy-carbonyl-C₄-alkyl; C-carboxy-C₄-alkyl and C₆-alkyl-carboxy-C₄-alkyl.

[0027] In some embodiments, the CSA, or a pharmaceutically acceptable salt thereof, is:





[0028] In some embodiments, the CSA, or a pharmaceutically acceptable salt thereof, is



[0029] In some embodiments, the pharmaceutically acceptable salt is a hydrochloride salt. In some embodiments, the pharmaceutically acceptable salt is a tri-hydrochloride salt.

[0030] The present invention may be embodied in other specific forms without departing from its spirit or essential characteristics. The described embodiments are to be considered in all respects only as illustrative and not restrictive. The scope of the invention is, therefore, indicated by the appended claims rather than by the foregoing description. All changes which come within the meaning and range of equivalency of the claims are to be embraced within their scope.

Detailed Description

Definitions

[0031] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which these
5 embodiments belong. The terminology used in the description herein is for describing particular embodiments only and is not intended to be limiting of the embodiments. As used in the specification and the appended claims, the singular forms “a,” “an,” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise.

[0032] Terms and phrases used in this application, and variations thereof, especially in the
10 appended claims, unless otherwise expressly stated, should be construed as open ended as opposed to limiting. As examples of the foregoing, the term ‘including’ should be read to mean ‘including, without limitation,’ ‘including but not limited to,’ or the like; the term ‘comprising’ as used herein is synonymous with ‘including,’ ‘containing,’ or ‘characterized by,’ and is inclusive or open-ended and does not exclude additional, unrecited elements or method steps;
15 the term ‘having’ should be interpreted as ‘having at least;’ the term ‘includes’ should be interpreted as ‘includes but is not limited to;’ the term ‘example’ is used to provide exemplary instances of the item in discussion, not an exhaustive or limiting list thereof; and use of terms like ‘preferably,’ ‘preferred,’ ‘desired,’ or ‘desirable,’ and words of similar meaning should not be understood as implying that certain features are critical, essential, or even important to the
20 structure or function of the invention, but instead as merely intended to highlight alternative or additional features that may or may not be utilized in a particular embodiment. In addition, the term “comprising” is to be interpreted synonymously with the phrases "having at least" or "including at least". When used in the context of a process, the term "comprising" means that the process includes at least the recited steps, but may include additional steps. When used in
25 the context of a compound, composition or device, the term "comprising" means that the compound, composition or device includes at least the recited features or components, but may also include additional features or components. Likewise, a group of items linked with the conjunction ‘and’ should not be read as requiring that each and every one of those items be present in the grouping, but rather should be read as ‘and/or’ unless expressly stated otherwise.
30 Similarly, a group of items linked with the conjunction ‘or’ should not be read as requiring mutual exclusivity among that group, but rather should be read as ‘and/or’ unless expressly stated otherwise.

[0033] It is understood that, in any compound described herein having one or more chiral centers, if an absolute stereochemistry is not expressly indicated, then each center may independently be of R-configuration or S-configuration or a mixture thereof. Thus, the compounds provided herein may be enantiomerically pure, enantiomerically enriched, racemic mixture, diastereomerically pure, diastereomerically enriched, or a stereoisomeric mixture. In addition it is understood that, in any compound described herein having one or more double bond(s) generating geometrical isomers that can be defined as E or Z, each double bond may independently be E or Z a mixture thereof.

[0034] Likewise, it is understood that, in any compound described, all tautomeric forms are also intended to be included.

[0035] It is to be understood that where compounds disclosed herein have unfilled valencies, then the valencies are to be filled with hydrogens or isotopes thereof, e.g., hydrogen-1 (protium) and hydrogen-2 (deuterium).

[0036] It is understood that the compounds described herein can be labeled isotopically.

Substitution with isotopes such as deuterium may afford certain therapeutic advantages resulting from greater metabolic stability, such as, for example, increased *in vivo* half-life or reduced dosage requirements. Each chemical element as represented in a compound structure may include any isotope of said element. For example, in a compound structure a hydrogen atom may be explicitly disclosed or understood to be present in the compound. At any position of the compound that a hydrogen atom may be present, the hydrogen atom can be any isotope of hydrogen, including but not limited to hydrogen-1 (protium) and hydrogen-2 (deuterium). Thus, reference herein to a compound encompasses all potential isotopic forms unless the context clearly dictates otherwise.

[0037] It is understood that the methods and combinations described herein include crystalline forms (also known as polymorphs, which include the different crystal packing arrangements of the same elemental composition of a compound), amorphous phases, salts, solvates, and hydrates. In some embodiments, the compounds described herein exist in solvated forms with pharmaceutically acceptable solvents such as water, ethanol, or the like. In other embodiments, the compounds described herein exist in unsolvated form. Solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and may be formed during the process of crystallization with pharmaceutically acceptable solvents such as water, ethanol, or the like. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. In addition, the compounds provided herein can exist in unsolvated as well as solvated forms. In general, the solvated forms are considered

equivalent to the unsolvated forms for the purposes of the compounds and methods provided herein.

[0038] Unless otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term “about.” Accordingly, unless indicated to the contrary, the numerical parameters set forth in the specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained by the present embodiments. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should be construed in light of the number of significant digits and ordinary rounding approaches.

[0039] Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the embodiments are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. Any numerical value, however, inherently contains certain errors necessarily resulting from the standard deviation found in their respective testing measurements. Every numerical range given throughout this specification and claims will include every narrower numerical range that falls within such broader numerical range, as if such narrower numerical ranges were all expressly written herein. Where a range of values is provided, it is understood that the upper and lower limit, and each intervening value between the upper and lower limit of the range is encompassed within the embodiments.

[0040] As used herein, any "R" group(s) such as, without limitation, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, and R¹⁸ represent substituents that can be attached to the indicated atom. Unless otherwise specified, an R group may be substituted or unsubstituted.

[0041] A “ring” as used herein can be heterocyclic or carbocyclic. The term “saturated” used herein refers to a ring having each atom in the ring either hydrogenated or substituted such that the valency of each atom is filled. The term “unsaturated” used herein refers to a ring where the valency of each atom of the ring may not be filled with hydrogen or other substituents. For example, adjacent carbon atoms in the fused ring can be doubly bound to each other. Unsaturation can also include deleting at least one of the following pairs and completing the valency of the ring carbon atoms at these deleted positions with a double bond; such as R₅ and R₉, R₈ and R₁₀; and R₁₃ and R₁₄.

[0042] Whenever a group is described as being “substituted” that group may be substituted with one, two, three or more of the indicated substituents, which may be the same or

different, each replacing a hydrogen atom. If no substituents are indicated, it is meant that the indicated "substituted" group may be substituted with one or more group(s) individually and independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl, acylalkyl, alkoxyalkyl, aminoalkyl, amino acid, aryl, heteroaryl, heteroalicycyl, aralkyl, heteroaralkyl, (heteroalicycyl)alkyl, hydroxy, protected hydroxyl, alkoxy, aryloxy, acyl, mercapto, alkylthio, arylthio, cyano, halogen (e.g., F, Cl, Br, and I), thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, protected C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, nitro, oxo, silyl, sulfenyl, sulfinyl, sulfonyl, haloalkyl, haloalkoxy, trihalomethanesulfonyl, trihalomethanesulfonamido, an amino, a mono-substituted amino group and a di-substituted amino group, $R^aO(CH_2)_mO-$, $R^b(CH_2)_nO-$, $R^cC(O)O(CH_2)_pO-$, and protected derivatives thereof. The substituent may be attached to the group at more than one attachment point. For example, an aryl group may be substituted with a heteroaryl group at two attachment points to form a fused multicyclic aromatic ring system. Biphenyl and naphthalene are two examples of an aryl group that is substituted with a second aryl group.

[0043] As used herein, "C_a" or "C_a to C_b" in which "a" and "b" are integers refer to the number of carbon atoms in an alkyl, alkenyl or alkynyl group, or the number of carbon atoms in the ring of a cycloalkyl, cycloalkenyl, cycloalkynyl, aryl, heteroaryl or heteroalicycyl group. That is, the alkyl, alkenyl, alkynyl, ring of the cycloalkyl, ring of the cycloalkenyl, ring of the cycloalkynyl, ring of the aryl, ring of the heteroaryl or ring of the heteroalicycyl can contain from "a" to "b", inclusive, carbon atoms. Thus, for example, a "C₁ to C₄ alkyl" group refers to all alkyl groups having from 1 to 4 carbons, that is, CH₃-, CH₃CH₂-, CH₃CH₂CH₂-, (CH₃)₂CH-, CH₃CH₂CH₂CH₂-, CH₃CH₂CH(CH₃)- and (CH₃)₃C-. If no "a" and "b" are designated with regard to an alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl, aryl, heteroaryl or heteroalicycyl group, the broadest range described in these definitions is to be assumed.

[0044] As used herein, "alkyl" refers to a straight or branched hydrocarbon chain that comprises a fully saturated (no double or triple bonds) hydrocarbon group. The alkyl group may have 1 to 25 carbon atoms (whenever it appears herein, a numerical range such as "1 to 25" refers to each integer in the given range; e.g., "1 to 25 carbon atoms" means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 25 carbon atoms, although the present definition also covers the occurrence of the term "alkyl" where no numerical range is designated). The alkyl group may also be a medium size alkyl having 1 to 15 carbon atoms. The alkyl group could also be a lower alkyl

having 1 to 6 carbon atoms. The alkyl group of the compounds may be designated as “C₄” or “C₁-C₄ alkyl” or similar designations. By way of example only, “C₁-C₄ alkyl” indicates that there are one to four carbon atoms in the alkyl chain, i.e., the alkyl chain is selected from methyl, ethyl, propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and t-butyl. Typical alkyl groups include, but are in no way limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tertiary butyl, pentyl and hexyl. The alkyl group may be substituted or unsubstituted.

[0045] As used herein, “alkenyl” refers to an alkyl group that contains in the straight or branched hydrocarbon chain one or more double bonds. The alkenyl group may have 2 to 25 carbon atoms (whenever it appears herein, a numerical range such as “2 to 25” refers to each integer in the given range; e.g., “2 to 25 carbon atoms” means that the alkenyl group may consist of 2 carbon atom, 3 carbon atoms, 4 carbon atoms, *etc.*, up to and including 25 carbon atoms, although the present definition also covers the occurrence of the term “alkenyl” where no numerical range is designated). The alkenyl group may also be a medium size alkenyl having 2 to 15 carbon atoms. The alkenyl group could also be a lower alkenyl having 1 to 6 carbon atoms. The alkenyl group of the compounds may be designated as “C₄” or “C₂-C₄ alkyl” or similar designations. An alkenyl group may be unsubstituted or substituted.

[0046] As used herein, “alkynyl” refers to an alkyl group that contains in the straight or branched hydrocarbon chain one or more triple bonds. The alkynyl group may have 2 to 25 carbon atoms (whenever it appears herein, a numerical range such as “2 to 25” refers to each integer in the given range; e.g., “2 to 25 carbon atoms” means that the alkynyl group may consist of 2 carbon atom, 3 carbon atoms, 4 carbon atoms, *etc.*, up to and including 25 carbon atoms, although the present definition also covers the occurrence of the term “alkynyl” where no numerical range is designated). The alkynyl group may also be a medium size alkynyl having 2 to 15 carbon atoms. The alkynyl group could also be a lower alkynyl having 2 to 6 carbon atoms. The alkynyl group of the compounds may be designated as “C₄” or “C₂-C₄ alkyl” or similar designations. An alkynyl group may be unsubstituted or substituted.

[0047] As used herein, “aryl” refers to a carbocyclic (all carbon) monocyclic or polycyclic aromatic ring system (including fused ring systems where two carbocyclic rings share a chemical bond) that has a fully delocalized pi-electron system throughout all the rings. The number of carbon atoms in an aryl group can vary. For example, the aryl group can be a C₆-C₁₄ aryl group, a C₆-C₁₀ aryl group, or a C₆ aryl group (although the definition of C₆-C₁₀ aryl covers the occurrence of “aryl” when no numerical range is designated). Examples of aryl groups include, but are not limited to, benzene, naphthalene and azulene. An aryl group may be substituted or unsubstituted.

[0048] As used herein, “aralkyl” and “aryl(alkyl)” refer to an aryl group connected, as a substituent, via a lower alkylene group. The aralkyl group may have 6 to 20 carbon atoms (whenever it appears herein, a numerical range such as “6 to 20” refers to each integer in the given range; *e.g.*, “6 to 20 carbon atoms” means that the aralkyl group may consist of 6
5 carbon atom, 7 carbon atoms, 8 carbon atoms, *etc.*, up to and including 20 carbon atoms, although the present definition also covers the occurrence of the term “aralkyl” where no numerical range is designated). The lower alkylene and aryl group of an aralkyl may be substituted or unsubstituted. Examples include but are not limited to benzyl, 2-phenylalkyl, 3-phenylalkyl, and naphthylalkyl.

10 [0049] “Lower alkylene groups” refer to a C₁-C₂₅ straight-chained alkyl tethering groups, such as -CH₂- tethering groups, forming bonds to connect molecular fragments via their terminal carbon atoms. Examples include but are not limited to methylene (-CH₂-), ethylene (-CH₂CH₂-), propylene (-CH₂CH₂CH₂-), and butylene (-CH₂CH₂CH₂CH₂-). A lower alkylene group can be substituted by replacing one or more hydrogen of the lower alkylene
15 group with a substituent(s) listed under the definition of “substituted.”

[0050] As used herein, “cycloalkyl” refers to a completely saturated (no double or triple bonds) mono- or multi- cyclic hydrocarbon ring system. When composed of two or more rings, the rings may be joined together in a fused fashion. Cycloalkyl groups can contain 3 to 10 atoms in the ring(s) or 3 to 8 atoms in the ring(s). A cycloalkyl group may be
20 unsubstituted or substituted. Typical cycloalkyl groups include, but are in no way limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

[0051] As used herein, “cycloalkenyl” refers to a mono- or multi- cyclic hydrocarbon ring system that contains one or more double bonds in at least one ring; although, if there is more than one, the double bonds cannot form a fully delocalized pi-electron system throughout all
25 the rings (otherwise the group would be “aryl,” as defined herein). When composed of two or more rings, the rings may be connected together in a fused fashion. A cycloalkenyl group may be unsubstituted or substituted.

[0052] As used herein, “cycloalkynyl” refers to a mono- or multi- cyclic hydrocarbon ring system that contains one or more triple bonds in at least one ring. If there is more than one
30 triple bond, the triple bonds cannot form a fully delocalized pi-electron system throughout all the rings. When composed of two or more rings, the rings may be joined together in a fused fashion. A cycloalkynyl group may be unsubstituted or substituted.

[0053] As used herein, “alkoxy” or “alkyloxy” refers to the formula -OR wherein R is an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl or a cycloalkynyl as defined above.

A non-limiting list of alkoxys are methoxy, ethoxy, n-propoxy, 1-methylethoxy (isopropoxy), n-butoxy, iso-butoxy, sec-butoxy and tert-butoxy. An alkoxy may be substituted or unsubstituted.

[0054] As used herein, “acyl” refers to a hydrogen, alkyl, alkenyl, alkynyl, aryl, or heteroaryl connected, as substituents, via a carbonyl group. Examples include formyl, acetyl, propanoyl, benzoyl, and acryl. An acyl may be substituted or unsubstituted.

[0055] As used herein, “alkoxyalkyl” or “alkyloxyalkyl” refers to an alkoxy group connected, as a substituent, via a lower alkylene group. Examples include alkyl-O-alkyl- and alkoxy-alkyl- with the terms alkyl and alkoxy defined herein.

[0056] As used herein, “hydroxyalkyl” refers to an alkyl group in which one or more of the hydrogen atoms are replaced by a hydroxy group. Exemplary hydroxyalkyl groups include but are not limited to, 2-hydroxyethyl, 3-hydroxypropyl, 2-hydroxypropyl, and 2,2-dihydroxyethyl. A hydroxyalkyl may be substituted or unsubstituted.

[0057] As used herein, “haloalkyl” refers to an alkyl group in which one or more of the hydrogen atoms are replaced by a halogen (e.g., mono-haloalkyl, di-haloalkyl and tri-haloalkyl). Such groups include but are not limited to, chloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl and 1-chloro-2-fluoromethyl, 2-fluoroisobutyl. A haloalkyl may be substituted or unsubstituted.

[0058] The term “amino” as used herein refers to a -NH_2 group.

[0059] As used herein, the term “hydroxy” refers to a -OH group.

[0060] A “cyano” group refers to a -CN group.

[0061] A “carbonyl” or an “oxo” group refers to a C=O group.

[0062] The term “azido” as used herein refers to a -N_3 group.


[0063] As used herein, “aminoalkyl” refers to an amino group connected, as a substituent, via a lower alkylene group. Examples include $\text{H}_2\text{N-alkyl-}$ with the term alkyl defined herein.

[0064] As used herein, “alkylcarboxyalkyl” refers to an alkyl group connected, as a substituent, to a carboxy group that is connected, as a substituent, to an alkyl group. Examples include $\text{alkyl-C(=O)O-alkyl-}$ and $\text{alkyl-O-C(=O)-alkyl-}$ with the term alkyl as defined herein.

[0065] As used herein, “C-carboxyalkyl” refers to a carboxy group connected, as a substituent, to an alkyl group. Examples include HO-(C=O)-alkyl , with the term alkyl as defined herein.

[0066] As used herein, “alkylaminoalkyl” refers to an alkyl group connected, as a substituent, to an amino group that is connected, as a substituent, to an alkyl group. Examples include alkyl-NH-alkyl-, with the term alkyl as defined herein.

[0067] As used herein, “dialkylaminoalkyl” or “di(alkyl)aminoalkyl” refers to two alkyl groups connected, each as a substituent, to an amino group that is connected, as a substituent,



to an alkyl group. Examples include with the term alkyl as defined herein.

[0068] As used herein, “alkylaminoalkylamino” refers to an alkyl group connected, as a substituent, to an amino group that is connected, as a substituent, to an alkyl group that is connected, as a substituent, to an amino group. Examples include alkyl-NH-alkyl-NH-, with the term alkyl as defined herein.

[0069] As used herein, “alkylaminoalkylaminoalkylamino” refers to an alkyl group connected, as a substituent, to an amino group that is connected, as a substituent, to an alkyl group that is connected, as a substituent, to an amino group that is connected, as a substituent, to an alkyl group. Examples include alkyl-NH-alkyl-NH-alkyl-, with the term alkyl as defined herein.

[0070] As used herein, “arylaminoalkyl” refers to an aryl group connected, as a substituent, to an amino group that is connected, as a substituent, to an alkyl group. Examples include aryl-NH-alkyl-, with the terms aryl and alkyl as defined herein.

[0071] As used herein, “aminoalkyloxy” refers to an amino group connected, as a substituent, to an alkyloxy group. Examples include H₂N-alkyl-O- and H₂N-alkoxy- with the terms alkyl and alkoxy as defined herein.

[0072] As used herein, “aminoalkyloxyalkyl” refers to an amino group connected, as a substituent, to an alkyloxy group connected, as a substituent, to an alkyl group. Examples include H₂N-alkyl-O-alkyl- and H₂N-alkoxy-alkyl- with the terms alkyl and alkoxy as defined herein.

[0073] As used herein, “aminoalkylcarboxy” refers to an amino group connected, as a substituent, to an alkyl group connected, as a substituent, to a carboxy group. Examples include H₂N-alkyl-C(=O)O- and H₂N-alkyl-O-C(=O)- with the term alkyl as defined herein.

[0074] As used herein, “aminoalkylaminocarbonyl” refers to an amino group connected, as a substituent, to an alkyl group connected, as a substituent, to an amino group connected, as a substituent, to a carbonyl group. Examples include H₂N-alkyl-NH-C(=O)- with the term alkyl as defined herein.


[0075] As used herein, “aminoalkylcarboxamido” refers to an amino group connected, as a substituent, to an alkyl group connected, as a substituent, to a carbonyl group connected, as a substituent to an amino group. Examples include $\text{H}_2\text{N-alkyl-C(=O)-NH-}$ with the term alkyl as defined herein.

5 [0076] As used herein, “azidoalkyloxy” refers to an azido group connected as a substituent, to an alkyloxy group. Examples include $\text{N}_3\text{-alkyl-O-}$ and $\text{N}_3\text{-alkoxy-}$ with the terms alkyl and alkoxy as defined herein.

[0077] As used herein, “cyanoalkyloxy” refers to a cyano group connected as a substituent, to an alkyloxy group. Examples include NC-alkyl-O- and NC-alkoxy- with the terms alkyl and alkoxy as defined herein.

[0078] As used herein, “guanidinoalkyloxy” refers to a guanidiny group connected, as a

substituent, to an alkyloxy group. Examples include



with the terms alkyl and alkoxy as defined herein.

[0079] As used herein, “guanidinoalkylcarboxy” refers to a guanidiny group connected, as a substituent, to an alkyl group connected, as a substituent, to a carboxy group. Examples

include

The image shows two chemical structures. The first structure is a guanidinoalkylcarboxy group: $\text{H}_2\text{N}-\text{C}(=\text{NH})-\text{NH}-\text{Alkyl}-\text{O}-\text{C}(=\text{O})-$ with a wavy line indicating the point of attachment. The second structure is a guanidinoalkylcarboxylate group: $\text{H}_2\text{N}-\text{C}(=\text{NH})-\text{NH}-\text{Alkyl}-\text{C}(=\text{O})-\text{O}-$ with a wavy line indicating the point of attachment.

and with the term alkyl as defined herein.

[0080] As used herein, “quaternaryammoniumalkylcarboxy” refers to a quaternized amino group connected, as a substituent, to an alkyl group connected, as a substituent, to a carboxy

group. Examples include

The image shows two chemical structures. The first structure is a quaternaryammoniumalkylcarboxy group: $\text{Alkyl}-\text{N}^+(\text{Alkyl})_3-\text{Alkyl}-\text{O}-\text{C}(=\text{O})-$ with a wavy line indicating the point of attachment. The second structure is a quaternaryammoniumalkylcarboxylate group: $\text{Alkyl}-\text{N}^+(\text{Alkyl})_3-\text{Alkyl}-\text{C}(=\text{O})-\text{O}-$ with a wavy line indicating the point of attachment.

and with the term alkyl as defined herein.

[0081] The term “halogen atom” or “halogen” as used herein, means any one of the radio-stable atoms of column 7 of the Periodic Table of the Elements, such as, fluorine, chlorine, bromine and iodine.

25 [0082] Where the numbers of substituents is not specified (e.g. haloalkyl), there may be one or more substituents present. For example “haloalkyl” may include one or more of the same or different halogens.

[0083] As used herein, the term “amino acid” refers to any amino acid (both standard and non-standard amino acids), including, but not limited to, α -amino acids, β -amino acids, γ -amino acids and δ -amino acids. Examples of suitable amino acids include, but are not limited to, alanine, asparagine, aspartate, cysteine, glutamate, glutamine, glycine, proline, serine, tyrosine, arginine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan and valine. Additional examples of suitable amino acids include, but are not limited to, ornithine, hypusine, 2-aminoisobutyric acid, dehydroalanine, gamma-aminobutyric acid, citrulline, beta-alanine, alpha-ethyl-glycine, alpha-propyl-glycine and norleucine.

[0084] A linking group is a divalent moiety used to link one steroid to another steroid. In some embodiments, the linking group is used to link a first CSA with a second CSA (which may be the same or different). An example of a linking group is (C₁-C₁₀) alkyloxy-(C₁-C₁₀) alkyl.

[0085] The terms “P.G.” or “protecting group” or “protecting groups” as used herein refer to any atom or group of atoms that is added to a molecule in order to prevent existing groups in the molecule from undergoing unwanted chemical reactions. Examples of protecting group moieties are described in T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 3. Ed. John Wiley & Sons, 1999, and in J.F.W. McOmie, *Protective Groups in Organic Chemistry* Plenum Press, 1973, both of which disclose suitable protecting groups. The protecting group moiety may be chosen in such a way, that they are stable to certain reaction conditions and readily removed at a convenient stage using methodology known from the art.

A non-limiting list of protecting groups include benzyl; substituted benzyl; alkylcarbonyls and alkoxy carbonyls (e.g., t-butoxycarbonyl (BOC), acetyl, or isobutyryl); arylalkylcarbonyls and arylalkoxy carbonyls (e.g., benzyloxycarbonyl); substituted methyl ether (e.g. methoxymethyl ether); substituted ethyl ether; a substituted benzyl ether; tetrahydropyranyl ether; silyls (e.g., trimethylsilyl, triethylsilyl, triisopropylsilyl, t-butyldimethylsilyl, tri-isopropylsilyloxymethyl, [2-(trimethylsilyl)ethoxy]methyl or t-butyldiphenylsilyl); esters (e.g. benzoate ester); carbonates (e.g. methoxymethylcarbonate); sulfonates (e.g. tosylate or mesylate); acyclic ketal (e.g. dimethyl acetal); cyclic ketals (e.g., 1,3-dioxane, 1,3-dioxolanes, and those described herein); acyclic acetal; cyclic acetal (e.g., those described herein); acyclic hemiacetal; cyclic hemiacetal; cyclic dithioketals (e.g., 1,3-dithiane or 1,3-dithiolane); orthoesters (e.g., those described herein) and triarylmethyl groups (e.g., trityl; monomethoxytrityl (MMTr); 4,4'-dimethoxytrityl (DMTr); 4,4',4''-trimethoxytrityl (TMTr); and those described herein). Amino-protecting groups are known to those skilled in the art. In


general, the species of protecting group is not critical, provided that it is stable to the conditions of any subsequent reaction(s) on other positions of the compound and can be removed at the appropriate point without adversely affecting the remainder of the molecule. In addition, a protecting group may be substituted for another after substantive synthetic transformations are complete. Clearly, where a compound differs from a compound disclosed herein only in that one or more protecting groups of the disclosed compound has been substituted with a different protecting group, that compound is within the disclosure.

CSA Compounds

[0086] Compounds useful in accordance with this disclosure are described herein, both generically and with particularity, and in U.S. Patent Nos. 6,350,738, 6,486,148, 6,767,904, 7,598,234, and 7,754,705. Compounds include steroid derivatives, such as cationic steroid antimicrobials ("CSAs") that exhibit one or more wound healing activities or functions.

[0087] CSA compounds, are synthetically produced small molecules that include a sterol backbone having various charged groups (e.g., amine and cationic groups) attached to the backbone. The backbone can be used to orient the amine or guanidine groups on one face, or plane, of the sterol backbone.


[0088] CSAs are cationic and amphiphilic, based upon the functional groups attached to the backbone. They are facially amphiphilic with a hydrophobic face and a polycationic face. For example, a scheme showing a compound having primary amino groups on one face, or plane, of a backbone is shown below in Scheme I:



Scheme I

[0089] The charged groups are believed to be responsible for antimicrobial properties. For example, the charged groups may disrupt the bacterial cellular membrane to cause cell death or sensitization.

[0090] In some embodiments disclosed herein the CSA compound may have a formula as set for in Formula (V) or a pharmaceutically acceptable salt thereof:



(V)

[0091] Where m , n , p , and q are independently 0 or 1; R^1 - R^{18} represent substituents that are attached to the indicated atom on the steroid backbone (i.e., steroid group); and at least two, preferably at least three, of R^1 - R^{18} each include a cationic group.

- 5 [0092] In one embodiment, rings A, B, C, and D are independently saturated, or are fully or partially unsaturated, provided that at least two of rings A, B, C, and D are saturated; m , n , p , and q are independently 0 or 1; R_1 through R_4 , R_6 , R_7 , R_{11} , R_{12} , R_{15} , R_{16} , and R_{18} are independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted alkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted alkoxyalkyl, substituted or unsubstituted alkylcarboxyalkyl, substituted or unsubstituted alkylaminoalkyl, substituted or unsubstituted alkylaminoalkylamino, substituted or unsubstituted alkylaminoalkylaminoalkylamino, a substituted or unsubstituted aminoalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted arylaminoalkyl, substituted or unsubstituted haloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted aminoalkyloxy, a substituted or unsubstituted aminoalkyloxyalkyl, a substituted or unsubstituted aminoalkylcarboxy, a substituted or unsubstituted aminoalkylaminocarbonyl, a substituted or unsubstituted aminoalkylcarboxamido, a substituted or unsubstituted di(alkyl)aminoalkyl, a substituted or unsubstituted C-carboxyalkyl, $H_2N-HC(Q_5)-C(O)-O-$, $H_2N-HC(Q_5)-C(O)-N(H)-$, substituted or unsubstituted azidoalkyloxy, substituted or unsubstituted cyanoalkyloxy, P.G.-HN-HC(Q₅)-C(O)-O-, substituted or unsubstituted guanidinoalkyloxy, substituted or unsubstituted quaternaryammoniumalkylcarboxy, and substituted or unsubstituted guanidinoalkyl carboxy, where Q_5 is a side chain of any amino acid (including a side chain of glycine, i.e., H), and P.G. is an amino protecting group; and
- 10 15 20 25 R_5 , R_8 , R_9 , R_{10} , R_{13} , R_{14} and R_{17} are independently deleted when one of rings A, B, C, or D is unsaturated so as to complete the valency of the carbon atom at that site, or R_5 , R_8 , R_9 , R_{10} , R_{13} , and R_{14} are independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted alkyl, substituted or unsubstituted hydroxyalkyl, substituted or

unsubstituted alkyloxyalkyl, a substituted or unsubstituted aminoalkyl, a substituted or unsubstituted aryl, substituted or unsubstituted haloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted aminoalkyloxy, a substituted or unsubstituted aminoalkylcarboxy, a substituted or unsubstituted aminoalkylaminocarbonyl, a substituted or unsubstituted di(alkyl)aminoalkyl, a substituted or unsubstituted C-carboxyalkyl, $\text{H}_2\text{N}-\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{O}-$, $\text{H}_2\text{N}-\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{N}(\text{H})-$, substituted or unsubstituted azidoalkyloxy, substituted or unsubstituted cyanoalkyloxy, P.G.-HN- $\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{O}-$, substituted or unsubstituted guanidinoalkyloxy, and substituted or unsubstituted guanidinoalkylcarboxy, where Q5 is a side chain of any amino acid, P.G. is an amino protecting group; provided that at least two or three of R₁₋₄, R₆, R₇, R₁₁, R₁₂, R₁₅, R₁₆, R₁₇, and R₁₈ are independently selected from the group consisting of a substituted or unsubstituted aminoalkyl, a substituted or unsubstituted aminoalkyloxy, substituted or unsubstituted alkylcarboxyalkyl, substituted or unsubstituted alkylaminoalkylamino, substituted or unsubstituted alkylaminoalkylaminoalkylamino, a substituted or unsubstituted aminoalkylcarboxy, a substituted or unsubstituted arylaminoalkyl, a substituted or unsubstituted aminoalkyloxyaminoalkylaminocarbonyl, a substituted or unsubstituted aminoalkylaminocarbonyl, a substituted or unsubstituted aminoalkylcarboxyamido, a substituted or unsubstituted quaternaryammoniumalkylcarboxy, a substituted or unsubstituted di(alkyl)aminoalkyl, a substituted or unsubstituted C-carboxyalkyl, $\text{H}_2\text{N}-\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{O}-$, $\text{H}_2\text{N}-\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{N}(\text{H})-$, substituted or unsubstituted azidoalkyloxy, substituted or unsubstituted cyanoalkyloxy, P.G.-HN- $\text{HC}(\text{Q}_5)-\text{C}(\text{O})-\text{O}-$, substituted or unsubstituted guanidinoalkyloxy, and a substituted or unsubstituted guanidinoalkylcarboxy.

[0093] In some embodiments, R₁ through R₄, R₆, R₇, R₁₁, R₁₂, R₁₅, R₁₆, and R₁₈ are independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted (C₁-C₁₈) alkyl, substituted or unsubstituted (C₁-C₁₈) hydroxyalkyl, substituted or unsubstituted (C₁-C₁₈) alkyloxy-(C₁-C₁₈) alkyl, substituted or unsubstituted (C₁-C₁₈) alkylcarboxy-(C₁-C₁₈) alkyl, substituted or unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkyl, substituted or unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino, substituted or unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino, a substituted or unsubstituted (C₁-C₁₈) aminoalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted arylamino-(C₁-C₁₈) alkyl, substituted or unsubstituted (C₁-C₁₈) haloalkyl, substituted or unsubstituted (C₂-C₆) alkenyl, substituted or unsubstituted (C₂-C₆) alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted (C₁-C₁₈)


aminoalkyloxy, a substituted or unsubstituted (C₁-C₁₈) aminoalkyloxy-(C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkylcarboxy, a substituted or unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkylcarboxamido, a substituted or unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, a substituted or unsubstituted C-carboxy(C₁-C₁₈)alkyl, H₂N-HC(Q₅)-C(O)-O—, H₂N—HC(Q₅)-C(O)—N(H)—, substituted or unsubstituted (C₁-C₁₈) azidoalkyloxy, substituted or unsubstituted (C₁-C₁₈) cyanoalkyloxy, P.G.-HN—HC(Q₅)-C(O)—O—, substituted or unsubstituted (C₁-C₁₈) guanidinoalkyloxy, substituted or unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, and substituted or unsubstituted (C₁-C₁₈) guanidinoalkyl carboxy, where Q₅ is a side chain of any amino acid (including a side chain of glycine, i.e., H), and P.G. is an amino protecting group; and R₅, R₈, R₉, R₁₀, R₁₃, R₁₄ and R₁₇ are independently deleted when one of rings A, B, C, or D is unsaturated so as to complete the valency of the carbon atom at that site, or R₅, R₈, R₉, R₁₀, R₁₃, and R₁₄ are independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted (C₁-C₁₈) alkyl, substituted or unsubstituted (C₁-C₁₈) hydroxyalkyl, substituted or unsubstituted (C₁-C₁₈) alkyloxy-(C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkyl, a substituted or unsubstituted aryl, substituted or unsubstituted (C₁-C₁₈) haloalkyl, substituted or unsubstituted (C₂-C₆) alkenyl, substituted or unsubstituted (C₂-C₆) alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted (C₁-C₁₈) aminoalkyloxy, a substituted or unsubstituted (C₁-C₁₈) aminoalkylcarboxy, a substituted or unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl, di(C₁-C₁₈ alkyl)aminoalkyl, a substituted or unsubstituted C-carboxy(C₁-C₁₈)alkyl, H₂N—HC(Q₅)-C(O)—O—, H₂N—HC(Q₅)-C(O)—N(H)—, substituted or unsubstituted (C₁-C₁₈) azidoalkyloxy, substituted or unsubstituted (C₁-C₁₈) cyanoalkyloxy, P.G.-HN—HC(Q₅)-C(O)—O—, substituted or unsubstituted (C₁-C₁₈) guanidinoalkyloxy, and substituted or unsubstituted (C₁-C₁₈) guanidinoalkylcarboxy, where Q₅ is a side chain of any amino acid, and P.G. is an amino protecting group; provided that at least two or three of R₁₋₄, R₆, R₇, R₁₁, R₁₂, R₁₅, R₁₆, R₁₇, and R₁₈ are independently selected from the group consisting of a substituted or unsubstituted (C₁-C₁₈) aminoalkyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkyloxy, substituted or unsubstituted (C₁-C₁₈) alkylcarboxy-(C₁-C₁₈) alkyl, substituted or unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino, substituted or unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino (C₁-C₁₈) alkylamino, a substituted or unsubstituted (C₁-C₁₈) aminoalkylcarboxy, a substituted or unsubstituted arylamino (C₁-C₁₈) alkyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkyloxy (C₁-C₁₈) aminoalkylaminocarbonyl, a substituted or unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl,

a substituted or unsubstituted (C₁-C₁₈) aminoalkylcarboxyamido, a substituted or unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, substituted or unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, a substituted or unsubstituted C-carboxy(C₁-C₁₈)alkyl, H₂N-HC(Q₅)-C(O)-O—, H₂N-HC(Q₅)-C(O)—N(H)—, substituted or unsubstituted (C₁-C₁₈) azidoalkyloxy, substituted or unsubstituted (C₁-C₁₈) cyanoalkyloxy, P.G.-HN-HC(Q₅)-C(O)—O—, substituted or unsubstituted (C₁-C₁₈) guanidinoalkyloxy, and a substituted or unsubstituted (C₁-C₁₈) guanidinoalkylcarboxy.

[0094] In some embodiments, R₁ through R₄, R₆, R₇, R₁₁, R₁₂, R₁₅, R₁₆, and R₁₈ are independently selected from the group consisting of hydrogen, hydroxyl, an unsubstituted (C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) hydroxyalkyl, unsubstituted (C₁-C₁₈) alkyloxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylcarboxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino- (C₁-C₁₈) alkylamino, an unsubstituted (C₁-C₁₈) aminoalkyl, an unsubstituted aryl, an unsubstituted arylamino-(C₁-C₁₈) alkyl, oxo, an unsubstituted (C₁-C₁₈) aminoalkyloxy, an unsubstituted (C₁-C₁₈) aminoalkyloxy-(C₁-C₁₈) alkyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxy, an unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxamido, an unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, an unsubstituted C-carboxy(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) guanidinoalkyloxy, unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, and unsubstituted (C₁-C₁₈) guanidinoalkyl carboxy; and R₅, R₈, R₉, R₁₀, R₁₃, R₁₄ and R₁₇ are independently deleted when one of rings A, B, C, or D is unsaturated so as to complete the valency of the carbon atom at that site, or R₅, R₈, R₉, R₁₀, R₁₃, and R₁₄ are independently selected from the group consisting of hydrogen, hydroxyl, an unsubstituted (C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) hydroxyalkyl, unsubstituted (C₁-C₁₈) alkyloxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylcarboxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino- (C₁-C₁₈) alkylamino, an unsubstituted (C₁-C₁₈) aminoalkyl, an unsubstituted aryl, an unsubstituted arylamino-(C₁-C₁₈) alkyl, oxo, an unsubstituted (C₁-C₁₈) aminoalkyloxy, an unsubstituted (C₁-C₁₈) aminoalkyloxy-(C₁-C₁₈) alkyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxy, an unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxamido, an unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, an unsubstituted C-carboxy(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) guanidinoalkyloxy, unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, and unsubstituted (C₁-C₁₈) guanidinoalkyl carboxy;

provided that at least two or three of R₁₋₄, R₆, R₇, R₁₁, R₁₂, R₁₅, R₁₆, R₁₇, and R₁₈ are independently selected from the group consisting of hydrogen, hydroxyl, an unsubstituted (C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) hydroxyalkyl, unsubstituted (C₁-C₁₈) alkyloxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylcarboxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino- (C₁-C₁₈) alkylamino, an unsubstituted (C₁-C₁₈) aminoalkyl, an unsubstituted aryl, an unsubstituted arylamino-(C₁-C₁₈) alkyl, oxo, an unsubstituted (C₁-C₁₈) aminoalkyloxy, an unsubstituted (C₁-C₁₈) aminoalkyloxy-(C₁-C₁₈) alkyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxy, an unsubstituted (C₁-C₁₈) aminoalkylaminocarbonyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxamido, an unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, an unsubstituted C-carboxy(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) guanidinoalkyloxy, unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, and unsubstituted (C₁-C₁₈) guanidinoalkyl carboxy.

[0095] In some embodiments, the compounds or pharmaceutically acceptable salts thereof of Formula (V) can be also represented by Formula (I):



wherein fused rings A, B, C, and D are independently saturated or fully or partially unsaturated; and each of R₁ through R₄, R₆, R₇, R₁₁, R₁₂, R₁₆, R₁₇, and R₁₈ is independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted (C₁-C₁₀) alkyl, (C₁-C₁₀) hydroxyalkyl, (C₁-C₁₀) alkyloxy-(C₁-C₁₀) alkyl, (C₁-C₁₀) alkylcarboxy-(C₁-C₁₀) alkyl, (C₁-C₁₀) alkylamino-(C₁-C₁₀) alkyl, (C₁-C₁₀) alkylamino-(C₁-C₁₀) alkylamino, (C₁-C₁₀) alkylamino-(C₁-C₁₀) alkylamino-(C₁-C₁₀) alkylamino, a substituted or unsubstituted (C₁-C₁₀) aminoalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted arylamino-(C₁-C₁₀) alkyl, (C₁-C₁₀) haloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted (C₁-C₁₀) aminoalkyloxy, a substituted or unsubstituted (C₁-C₁₀) aminoalkyloxy-(C₁-C₁₀) alkyl, a substituted or unsubstituted (C₁-C₁₀) aminoalkylcarboxy, a substituted or unsubstituted (C₁-C₁₀) aminoalkylaminocarbonyl, a substituted or unsubstituted (C₁-C₁₀) aminoalkylcarboxamido, a substituted or unsubstituted C-carboxy(C₁-C₁₀)alkyl, H₂N-HC(Q5)-C(O)—O—, H₂N—HC(Q5)-C(O)—N(H)—, (C₁-C₁₀) azidoalkyloxy, (C₁-C₁₀)

cyanoalkyloxy, P.G.-HN—HC(Q5)-C(O)—O—, (C₁-C₁₀) guanidinoalkyloxy, (C₁-C₁₀) quaternaryammoniumalkylcarboxy, and (C₁-C₁₀) guanidinoalkyl carboxy, where Q5 is a side chain of any amino acid (including the side chain of glycine, i.e., H), PG. is an amino protecting group, and R₅, R₈, R₉, R₁₀, R₁₃, and R₁₄ is each independently: deleted when one of
 5 fused rings A, B, C, or D is unsaturated so as to complete the valency of the carbon atom at that site, or selected from the group consisting of hydrogen, hydroxyl, 10 a substituted or unsubstituted (C₁-C₁₀) alkyl, (C₁-C₁₀) hydroxyalkyl, (C₁-C₁₀) alkyloxy-(C₁-C₁₀) alkyl, a substituted or unsubstituted (C₁-C₁₀) aminoalkyl, a substituted or unsubstituted aryl, C₁-C₁₀ haloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, oxo, a linking group attached to a second steroid, a
 10 substituted or unsubstituted (C₁-C₁₀) aminoalkyloxy, a substituted or unsubstituted (C₁-C₁₀) aminoalkylcarboxy, a substituted or unsubstituted (C₁-C₁₀) aminoalkylaminocarbonyl, H₂N—HC(Q5)-C(O)—O—, H₂N—HC(Q5)-C(O)—N(H)—, (C₁-C₁₀) azidoalkyloxy, (C₁-C₁₀) cyanoalkyloxy, P.G.-HN—HC(Q5)-C(O)—O—, (C₁-C₁₀) guanidinoalkyloxy, and (C₁-C₁₀) guanidinoalkylcarboxy, where Q5 is a side chain of any amino acid, PG. is an amino
 15 protecting group, and provided that at least two of R₁ through R₁₄ are independently selected from the group consisting of a substituted or unsubstituted (C₁-C₁₀) aminoalkyloxy, (C₁-C₁₀) alkylcarboxy-(C₁-C₁₀) alkyl, (C₁-C₁₀) alkylamino-(C₁-C₁₀) alkylamino, (C₁-C₁₀) alkylamino-(C₁-C₁₀) alkylamino-(C₁-C₁₀) alkylamino, a substituted or unsubstituted (C₁-C₁₀) aminoalkylcarboxy, a substituted or unsubstituted arylamino(C₁-C₁₀) alkyl, a substituted or
 20 unsubstituted (C₁-C₁₀) aminoalkyloxy-(C₁-C₁₀) alkyl, a substituted or unsubstituted (C₁-C₁₀) aminoalkylaminocarbonyl, (C₁-C₁₀) quaternary ammonium alkylcarboxy, H₂N-HC(Q5)-C(O)—O—, H₂N-HC(Q5)-C(O)—N(H)—, (C₁-C₁₀) azidoalkyloxy, (C₁-C₁₀) cyanoalkyloxy, PG.-HN-HC(Q5)-C(O)—O—, (C₁-C₁₀) guanidinoalkyloxy, and (C₁-C₁₀) guanidinoalkylcarboxy; or a pharmaceutically acceptable salt thereof.

25 **[0096]** In some embodiments, rings A, B, C, and D are independently saturated, heterocyclic, and/or non-heterocyclic.

[0097] In some embodiments, R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of hydrogen, an unsubstituted (C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) hydroxyalkyl, unsubstituted (C₁-C₁₈) alkyloxy-(C₁-C₁₈) alkyl, unsubstituted (C₁-C₁₈) alkylcarboxy-(C₁-C₁₈)
 30 alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino, unsubstituted (C₁-C₁₈) alkylamino-(C₁-C₁₈) alkylamino- (C₁-C₁₈) alkylamino, an unsubstituted (C₁-C₁₈) aminoalkyl, an unsubstituted arylamino-(C₁-C₁₈) alkyl, an unsubstituted (C₁-C₁₈) aminoalkyloxy, an unsubstituted (C₁-C₁₈) aminoalkyloxy-(C₁-C₁₈) alkyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxy, an unsubstituted (C₁-C₁₈)

aminoalkylaminocarbonyl, an unsubstituted (C₁-C₁₈) aminoalkylcarboxamido, an unsubstituted di(C₁-C₁₈ alkyl)aminoalkyl, an unsubstituted C-carboxy(C₁-C₁₈)alkyl, unsubstituted (C₁-C₁₈) guanidinoalkyloxy, unsubstituted (C₁-C₁₈) quaternaryammoniumalkylcarboxy, and unsubstituted (C₁-C₁₈) guanidinoalkyl carboxy; and
 5 R₁, R₂, R₄, R₅, R₆, R₈, R₉, R₁₀, R₁₁, R₁₃, R₁₄, R₁₅, R₁₆, and R₁₇ are independently selected from the group consisting of hydrogen and unsubstituted (C₁-C₆) alkyl.


[0098] In some embodiments, R₁, R₂, R₄, R₅, R₆, R₈, R₁₀, R₁₁, R₁₄, R₁₆, and R₁₇ are each hydrogen; and R₉ and R₁₃ are each methyl.

[0099] In some embodiments, R₃, R₇, R₁₂, and R₁₈ are independently selected from the group
 10 consisting of aminoalkyloxy; aminoalkylcarboxy; alkylaminoalkyl; alkoxy-carbonylalkyl; alkylcarbonylalkyl; di(alkyl)aminoalkyl; C-carboxyalkyl; alkoxy-carbonylalkyl; and alkylcarboxyalkyl.






[00100] In some embodiments, R₃, R₇, and R₁₂ are independently selected from the group consisting of aminoalkyloxy and aminoalkylcarboxy; and R₁₈ is selected from the group
 15 consisting of alkylaminoalkyl; alkoxy-carbonylalkyl; alkylcarbonyloxyalkyl; di(alkyl)aminoalkyl; alkylaminoalkyl; alkoxy-carbonylalkyl; and alkylcarboxyalkyl.

[00101] In some embodiments, R₃, R₇, and R₁₂ are the same substituent or different substituents and/or may be independently an aminoalkyloxy and/or an aminoalkylcarboxy. In some embodiments, R₁₈ is alkylaminoalkyl, alkoxy-carbonylalkyl, di(alkyl)aminoalkyl, C-carboxyalkyl, or an alkylcarboxyalkyl. In some embodiments, R₃, R₇, R₁₂, and R₁₈ are
 20 independently selected from the group consisting of amino-C₃-alkyloxy; amino-C₃-alkyl-carboxy; C₈-alkylamino-C₅-alkyl; C₈-alkoxy-carbonyl-C₄-alkyl; C₈-alkyl-carbonyl-C₄-alkyl; di-(C₅-alkyl)amino-C₅-alkyl; C-carboxy-C₄-alkyl; C₁₃-alkylamino-C₅-alkyl; C₆-alkoxy-carbonyl-C₄-alkyl; and C₆-alkyl-carboxy-C₄-alkyl.

[00102] In some embodiments, the compounds or pharmaceutically acceptable salts thereof
 25 of Formula (V) can be also represented by Formula (Ia):



[00103] In some embodiments, the compounds or pharmaceutically acceptable salts thereof
 30 of Formula (Ia) may be one of the following or selected from the following group:



[00104] The foregoing compounds may be provided as a pharmaceutically acceptable salt such as, but not limited to, a hydrochloride salt or a tri-hydrochloride salt.

[00105] In some embodiments, compounds comprise a ring system of at least 4 fused rings, where each of the rings has from 5-7 atoms. The ring system has two faces, and contains 3 chains attached to the same face. Each of the chains contains a nitrogen-containing group that is separated from the ring system by at least one atom; the nitrogen-containing group is an amino group, e.g., a primary amino group, or a guanidino group.

[00106] The compound can also contain a hydrophobic group. In some embodiments, the hydrophobic group is a substituted (C₃₋₁₀) aminoalkyl group, a (C₁-C₁₀) alkyloxy (C₃₋₁₀) alkyl group, or a (C₁-C₁₀) alkylamino (C₃₋₁₀) alkyl group, attached to the steroid backbone. In some embodiments, the hydrophobic group is a substituted, branched, or unbranched substituent with greater than 12, 16, 18, 20, or 22 carbons. In some embodiments, the hydrophobic group may include a hydrocarbon chain of at least 9, 11, or 13 carbons distal to a heteroatom. In some embodiments, a compound having a structure according to Formula (V) includes a hydrophobic group at R¹⁸.

[00107] In some embodiments, the compounds set forth herein preserve certain stereochemical and electronic characteristics found in steroids. The term “same configuration” as used herein refers to substituents on the fused steroid having the same stereochemical orientation. For example, in some embodiments, substituents R₃, R₇ and R₁₂ are all β-substituted or α-substituted.

[00108] In some embodiments, compounds include, but are not limited to, compounds having amine or guanidine groups covalently attached to a steroid backbone or scaffold at any carbon position, e.g., cholic acid. In various embodiments, a group is covalently attached at any one, or more, of positions C₃, C₇ and C₁₂ of the steroid backbone or scaffold. In additional embodiments, a group is absent from anyone, or more, of positions C₃, C₇ and C₁₂ of the steroid backbone or scaffold. Compounds that include such groups can include a tether, the tether having variable chain length or size. As used herein, the terms “tether” or “tethered,” when used in reference to a compound, refers to the chain of atoms between the steroid backbone or scaffold and a terminal amino or guanidine group. In various embodiments, a tether is covalently attached at anyone, or more, of positions C₃, C₇ and C₁₂. In additional embodiments, a tether is lacking at anyone, or more, of positions C₃, C₇ and C₁₂. A tether length may include the heteroatom (O or N) covalently attached to the steroid backbone. The tether may include a hydrolysable linkage such as an ester linkage.

[00109] In some embodiments, other ring systems can also be used, e.g., 5-member fused rings. Compounds with backbones having a combination of 5- and 6-membered rings are also contemplated. Amine or guanidine groups can be separated from the backbone by at least one, two, three, four or more atoms. The backbone can be used to orient the amine or
5 guanidine groups on one face, or plane, of the steroid. For example, a scheme showing a compound having primary amino groups on one face, or plane, of a backbone is shown in Scheme I above.

[00110] The compounds and compositions disclosed herein are optionally prepared as pharmaceutically acceptable salts. The term “pharmaceutically acceptable salt” as used
10 herein is a broad term, and is to be given its ordinary and customary meaning to a skilled artisan (and is not to be limited to a special or customized meaning), and refers without limitation to a salt of a compound that does not cause significant irritation to an organism to which it is administered and does not abrogate the biological activity and properties of the compound. In some embodiments, the salt is an acid addition salt of the compound.
15 Pharmaceutical salts can be obtained by reacting a compound with inorganic acids such as hydrohalic acid (*e.g.*, hydrochloric acid or hydrobromic acid), sulfuric acid, nitric acid, and phosphoric acid. Pharmaceutical salts can also be obtained by reacting a compound with an organic acid such as aliphatic or aromatic carboxylic or sulfonic acids, for example formic acid, acetic acid, propionic acid, glycolic acid, pyruvic acid, malonic acid, maleic acid,
20 fumaric acid, trifluoroacetic acid, benzoic acid, cinnamic acid, mandelic acid, succinic acid, lactic acid, malic acid, tartaric acid, citric acid, ascorbic acid, nicotinic acid, methanesulfonic acid, ethanesulfonic acid, *p*-toluenesulfonic acid, salicylic acid, stearic acid, muconic acid, butyric acid, phenylacetic acid, phenylbutyric acid, valproic acid, 1,2-ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 2-naphthalenesulfonic acid, or
25 naphthalenesulfonic acid. Pharmaceutical salts can also be obtained by reacting a compound with a base to form a salt such as an ammonium salt, an alkali metal salt, such as a lithium, sodium or a potassium salt, an alkaline earth metal salt, such as a calcium, magnesium or aluminum salt, a salt of organic bases such as dicyclohexylamine, *N*-methyl-*D*-glucamine, tris(hydroxymethyl)methylamine, C₁-C₇ alkylamine, cyclohexylamine, dicyclohexylamine,
30 triethanolamine, ethylenediamine, ethanolamine, diethanolamine, triethanolamine, tromethamine, and salts with amino acids such as arginine and lysine; or a salt of an inorganic base, such as aluminum hydroxide, calcium hydroxide, potassium hydroxide, sodium carbonate, sodium hydroxide, or the like.

[00111] The forgoing compositions may be used for the novel use of increasing the rate of wound healing in a subject with a tissue wound.

Tissue Treatment Compositions

5 [00112] While it is possible for the compounds described herein to be administered alone, it may be preferable to formulate the compounds as pharmaceutical compositions suitable for application to the tissue wound. As such, in yet another aspect, tissue treatment compositions useful in the methods and uses of the disclosed embodiments are provided. More particularly, the tissue treatment compositions described herein may be useful, *inter alia*, for
10 treating or promoting wound healing in a subject. A tissue treatment composition is any composition that may be administered *in vitro* or *in vivo* or both to a subject in order to promote or enhance wound healing. In a preferred embodiment, a tissue treatment compositions may be administered *in vivo*.

[00113] As used herein the terms “pharmaceutically acceptable” and “physiologically acceptable” mean a biologically compatible formulation, gaseous, liquid or solid, or mixture thereof, which is suitable for one or more routes of administration, *in vivo* delivery, or contact. A formulation is compatible in that it does not destroy activity of an active ingredient therein (e.g., a CSA), or induce adverse side effects that far outweigh any prophylactic or therapeutic effect or benefit.

20 [00114] In an embodiment, the tissue treatment compositions may be formulated with pharmaceutically acceptable excipients such as carriers, solvents, stabilizers, adjuvants, diluents, etc., depending upon the particular mode of administration and dosage form. The tissue treatment compositions should generally be formulated to achieve a physiologically compatible pH, and may range from a pH of about 3 to a pH of about 11, preferably about pH
25 3 to about pH 7, depending on the formulation and route of administration. In alternative embodiments, it may be preferred that the pH is adjusted to a range from about pH 5.0 to about pH 8. More particularly, the tissue treatment compositions may comprise a therapeutically or prophylactically effective amount of at least one compound as described herein, together with one or more pharmaceutically acceptable excipients. Optionally, the
30 tissue treatment compositions may comprise a combination of the compounds described herein, or may include a second active ingredient useful in the treatment or prevention of bacterial infection (e.g., anti-bacterial or anti-microbial agents).

[00115] Formulations, e.g., for parenteral or oral administration, are most typically solids, liquid solutions, emulsions or suspensions, while inhalable formulations for pulmonary

administration are generally liquids or powders, with powder formulations being generally preferred. A preferred tissue treatment composition may also be formulated as a lyophilized solid that is reconstituted with a physiologically compatible solvent prior to administration. Alternative tissue treatment compositions may be formulated as syrups, creams, ointments, tablets, and the like.

[00116] In general, formulations are prepared by uniformly and intimately associating the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product. For example, a tablet may be made by compression or molding. Compressed tablets may be prepared by compressing, in a suitable machine, an active ingredient (*e.g.*, a CSA) in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface-active or dispersing agent. Molded tablets may be produced by molding, in a suitable apparatus, a mixture of powdered compound (*e.g.*, CSA) moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide a slow or controlled release of the active ingredient therein.

[00117] Cosolvents and adjuvants may be added to the formulation. Non-limiting examples of cosolvents contain hydroxyl groups or other polar groups, for example, alcohols, such as isopropyl alcohol; glycols, such as propylene glycol, polyethyleneglycol, polypropylene glycol, glycol ether; glycerol; polyoxyethylene alcohols and polyoxyethylene fatty acid esters. Adjuvants include, for example, surfactants such as, soya lecithin and oleic acid; sorbitan esters such as sorbitan trioleate; and polyvinylpyrrolidone.

[00118] Additionally, the tissue treatment compositions may be in the form of a sterile injectable preparation, such as a sterile injectable aqueous emulsion or oleaginous suspension. This emulsion or suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents, which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, such as a solution in 1,2-propane-diol.

[00119] The sterile injectable preparation may also be prepared as a lyophilized powder. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile fixed oils may be employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid may likewise be used in the preparation of injectables.

[00120] To obtain a stable water-soluble dose form of a tissue treatment compositions, a pharmaceutically acceptable salt of a compound described herein may be dissolved in an aqueous solution of an organic or inorganic acid, such as 0.3 M solution of succinic acid, or more preferably, citric acid. If a soluble salt form is not available, the compound may be dissolved in a suitable co-solvent or combination of co-solvents. Examples of suitable co-solvents include alcohol, propylene glycol, polyethylene glycol 300, polysorbate 80, glycerin and the like in concentrations ranging from about 0 to about 60% of the total volume. In one embodiment, the active compound is dissolved in DMSO and diluted with water.

[00121] The pharmaceutical composition may also be in the form of a solution of a salt form of the active ingredient in an appropriate aqueous vehicle, such as water or isotonic saline or dextrose solution. Also contemplated are compounds which have been modified by substitutions or additions of chemical or biochemical moieties which make them more suitable for delivery (e.g., increase solubility, bioactivity, palatability, decrease adverse reactions, etc.), for example by esterification, glycosylation, PEGylation, etc.

[00122] In one embodiment, the compounds described herein may be formulated for oral administration in a lipid-based formulation suitable for low solubility compounds. Lipid-based formulations can generally enhance the oral bioavailability of such compounds.

[00123] In some exemplary embodiments, a CSA comprises a multimer (e.g., a dimer, trimer, tetramer, or higher order polymer). In some exemplary embodiments, the CSAs can be incorporated into pharmaceutical compositions or formulations. Such compositions/formulations are useful for administration to a subject, in vivo or ex vivo. Tissue treatment compositions and formulations include carriers or excipients for administration to a subject.

[00124] Such formulations include solvents (aqueous or non-aqueous), solutions (aqueous or non-aqueous), emulsions (e.g., oil-in-water or water-in-oil), suspensions, syrups, elixirs, dispersion and suspension media, coatings, isotonic and absorption promoting or delaying agents, compatible with pharmaceutical administration or in vivo contact or delivery. Aqueous and non-aqueous solvents, solutions and suspensions may include suspending agents and thickening agents. Such pharmaceutically acceptable carriers include tablets (coated or uncoated), capsules (hard or soft), microbeads, powder, granules and crystals. Supplementary active compounds (e.g., preservatives, antibacterial, antiviral and antifungal agents) can also be incorporated into the compositions.

[00125] Cosolvents and adjuvants may be added to the formulation. Non-limiting examples of cosolvents contain hydroxyl groups or other polar groups, for example, alcohols, such as

isopropyl alcohol; glycols, such as propylene glycol, polyethyleneglycol, polypropylene glycol, glycol ether; glycerol; polyoxyethylene alcohols and polyoxyethylene fatty acid esters. Adjuvants include, for example, surfactants such as, soya lecithin and oleic acid; sorbitan esters such as sorbitan trioleate; and polyvinylpyrrolidone.

- 5 [00126] Additionally, it may be desirable to include other therapeutically beneficial agents in the formulation. For example, the vehicles or carriers may also include humectants or moisturizers to maintain a desired moisture level in the treated area. Other possibilities include drugs such as anesthetics or antibiotics, which provide other desired effects. Again, the possibilities are unlimited and are left to the practitioner.
- 10 [00127] A tissue treatment composition contains a total amount of the active ingredient(s) sufficient to achieve an intended therapeutic effect. In some embodiments, the tissue treatment composition includes the CSA in a weight/weight concentration of at least 0.001%, 0.01%, 0.1%, 1.0%, and/or less than 80%, 50%, 25%, 15%, 10%, or 5%, or within a range between any of the foregoing lower and upper endpoints. In some embodiments, the tissue
- 15 treatment composition includes the CSA in a weight/weight concentration of about 0.04%. In other embodiments, the tissue treatment composition includes the CSA in a weight/weight concentration of less than 0.1%. In some embodiments, the tissue treatment composition includes the CSA in a weight/weight concentration of between about 0.01% and about 2.0%.

20 Methods and Uses

- [00128] Methods disclosed herein include identifying a subject with a tissue wound and administering a tissue treatment composition including a CSA, thereby increasing the rate of wound healing. In some embodiments, the method includes (i) providing a tissue treatment composition as described above; (ii) identifying a subject in need of accelerated healing of a
- 25 tissue wound; and (iii) contacting the tissue wound with the tissue treatment composition to increase the rate of healing thereof.

- [00129] In some embodiments, the subject is an animal. In some embodiments, the animal is a mammal. The mammal may be a human or primate in some embodiments. A mammal includes any mammal, such as by way of non-limiting example, cattle, pigs, sheep, goats,
- 30 horses, camels, buffalo, cats, dogs, rats, mice, and humans. Preferably the subject is a human, horse, or dog. In some embodiments, the subject is a vertebrate. In other embodiments, the subject is a non-human animal.

[00130] The subject has a tissue wound in need of healing and/or accelerated healing. Unless specified, the term “wound” is used herein in its generic sense, meaning that it encompasses

all types of wounds and injuries. The term “wound” encompasses burns, ulcers, lacerations, incisions, etc. “Wound” and “lesion” may be used interchangeably herein, and unless the context specifically dictates otherwise, no distinction is intended. Lesions/wounds can be acute or chronic. Wounds can be full thickness, i.e., penetrating all layers of skin, or partial

5 thickness, i.e., penetrating less than all layers of skin. Examples of acute wounds include, but are not limited to, surgical wounds (i.e., incisions), penetrating wounds, avulsion injuries, crushing injuries, shearing injuries, burn injuries, lacerations, and bite wounds. Examples of chronic wounds include, but are not limited to, ulcers, such as arterial ulcers, venous ulcers, pressure ulcers, and diabetic ulcers. Of course, acute wounds can become chronic wounds.

10 [00131] In some embodiments, the tissue wound can be a navel of a newborn subject (i.e., a newborn with navel ill).

[00132] The tissue treatment composition can be applied to an open wound or a closed wound. In some embodiments the composition is applied to an open wound for a period of sufficient time to cause closure of the wound. The composition can be applied at least once

15 or twice daily at least 2, 4, 8, or 16 times and/or for over a period of at least 2, 4, 8, or 16 days.

[00133] The compositions disclosed herein can be administered to any wound, anywhere it is desirable to promote wound healing. The compositions are also useful to reduce scarring after a wound is closed and/or healed. The compositions can be applied in any form of vehicle or

20 carrier, including but not limited to, liquids, gels, lotions, creams, pastes, and ointments. The means of application will depend upon what form the composition takes: liquids can be sprayed or poured, for example; gels, lotions, creams, pastes, and ointments can be rubbed or massaged, for example. These and other forms, and/or carriers/vehicles, for composition delivery, are described in publications such as Remington’s Pharmaceutical Science, and

25 other similar publications.

[00134] The delivery forms can be homogeneous, *e.g.*, forms in which the composition is in solution, or heterogeneous, *e.g.*, forms in which the composition is contained within liposomes or microspheres. The forms can produce an immediate effect, and can alternatively, or additionally, produce an extended effect. For example, liposomes, or

30 microspheres, or other similar means of providing an extended release of the composition, can be used to extend the period during which the composition is exposed to the lesion; non-encapsulated compositions can also be provided for an immediate effect.

[00135] The delivery forms can also take the form of devices, which can deliver the composition to a lesion for a desired period of time. Devices include, but are not limited to,

bandages, surgical dressings, gauzes, adhesive strips, surgical staples, clips, hemostats, intrauterine devices, sutures, trocars, catheters, tubes, collagen sponges, and implants. Implants include, but are not limited to, pills, pellets, rods, wafers, discs, and tablets.

[00136] Devices according to the disclosure can be prepared according to known methods, and can include, or be made from, polymeric material. In some instances, the polymeric material will be an absorbable material and in other instances, a non-absorbable material, or in other instances a resorbable material. Devices can include absorbable, non-absorbable, resorbable materials, and combinations thereof.

[00137] Absorbable materials can be synthetic materials and non-synthetic materials. Absorbable synthetic materials include, but are not limited to, cellulosic polymers, glycolic acid polymers, methacrylate polymers, ethylene vinyl acetate polymers, ethylene vinyl alcohol copolymers, polycaprolactam, polyacetate, copolymers of lactide and glycolide, polydioxanone, polyglactin, poliglecaprone, polyglyconate, polygluconate, and combinations thereof. Absorbable non-synthetic materials include, but are not limited to, catgut, cargile membrane, fascia lata, gelatin, collagen, and combinations thereof.

[00138] Nonabsorbable synthetic materials include, but are not limited to nylons, rayons, polyesters, polyolefins, and combinations thereof. Non-absorbable non-synthetic materials include, but are not limited to, silk, dermal silk, cotton, linen, and combinations thereof.

[00139] Combinations of the foregoing devices and carriers/vehicles are also envisioned. For example, a CSA gel or ointment can be impregnated into a bandage or wound dressing for delivery of the CSA to the desired location. As another example, an implantable absorbable device can be loaded with a CSA solution and release the solution from the device over a period as desired.

[00140] It may be desirable to provide for other conditions in the practice of the present methods. For example, it may be desirable to ensure that the target region of the lesion is sufficiently oxygenated; generally, it is sufficient that atmospheric oxygen be present. It also may be desirable to maintain a desired level of moisture and a particular temperature; in some embodiments, a warm, moist environment is desirable. While not required, it may also be desirable to establish or maintain a sterile environment.

[00141] In some embodiments, the composition may be incorporated into a medical device coating.

[00142] One of ordinary skill in the art to which these exemplary embodiments belong will understand that the compositions may be administered in numerous ways. For example, administration may mean simply applying the compositions to a wound directly. In some

exemplary embodiments, administration may be enteral, parenteral, or topical. Other exemplary routes of administration for contact or *in vivo* delivery which a compound can optionally be formulated include inhalation, respiration, intubation, intrapulmonary instillation, oral (buccal, sublingual, mucosal), intrapulmonary, rectal, vaginal, intrauterine, intradermal, topical, dermal, parenteral (*e.g.*, subcutaneous, intramuscular, intravenous, intradermal, intraocular, intratracheal and epidural), intranasal, intrathecal, intraarticular, intracavity, transdermal, iontophoretic, ophthalmic, optical (*e.g.*, corneal), intraglandular, intraorgan, intralymphatic.

10 Dosage

[00143] Compounds (*e.g.*, CSAs), including pharmaceutical formulations can be packaged in unit dosage forms for ease of administration and uniformity of dosage. A “unit dosage form” as used herein refers to a physically discrete unit suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of compound optionally in association with a pharmaceutical carrier (excipient, diluent, vehicle or filling agent) which, when administered in one or more doses, is calculated to produce a desired effect (*e.g.*, prophylactic or therapeutic effect or benefit). Unit dosage forms can contain a daily dose or unit, daily sub-dose, or an appropriate fraction thereof, of an administered compound (*e.g.*, CSA). Unit dosage forms also include, for example, capsules, troches, cachets, lozenges, tablets, ampules and vials, which may include a composition in a freeze-dried or lyophilized state; a sterile liquid carrier, for example, can be added prior to administration or delivery *in vivo*. Unit dosage forms additionally include, for example, ampules and vials with liquid compositions disposed therein. Unit dosage forms further include compounds for transdermal administration, such as “patches” that contact with the epidermis of the subject for an extended or brief period of time. The individual unit dosage forms can be included in multi-dose kits or containers. Pharmaceutical formulations can be packaged in single or multiple unit dosage forms for ease of administration and uniformity of dosage.

[00144] Compounds (*e.g.*, CSAs) can be administered in accordance with the methods at any frequency as a single bolus or multiple dose *e.g.*, one, two, three, four, five, or more times hourly, daily, weekly, monthly, or annually or between about 1 to 10 days, weeks, months, or for as long as appropriate. Exemplary frequencies are typically from 1-7 times, 1-5 times, 1-3 times, 2-times or once, daily, weekly or monthly. Timing of contact, administration *ex vivo* or *in vivo* delivery can be dictated by the infection, pathogenesis, symptom, pathology or adverse side effect to be treated. For example, an amount can be administered to the subject

substantially contemporaneously with, or within about 1-60 minutes or hours of the onset of a symptom or adverse side effect, pathogenesis, or vaccination.

[00145] Doses may vary depending upon whether the treatment is therapeutic or prophylactic, the onset, progression, severity, frequency, duration, probability of or susceptibility of the symptom, the type pathogenesis to which treatment is directed, clinical endpoint desired, previous, simultaneous or subsequent treatments, general health, age, gender or race of the subject, bioavailability, potential adverse systemic, regional or local side effects, the presence of other disorders or diseases in the subject, and other factors that will be appreciated by the skilled artisan (e.g., medical or familial history). Dose amount, frequency or duration may be increased or reduced, as indicated by the clinical outcome desired, status of the infection, symptom or pathology, any adverse side effects of the treatment or therapy. The skilled artisan will appreciate the factors that may influence the dosage, frequency and timing required to provide an amount sufficient or effective for providing a prophylactic or therapeutic effect or benefit.

[00146] In instances where animal and/or human dosages for compounds have been established for at least some condition, those same dosages may be used, or dosages that are between about 0.1% and 500%, more preferably between about 25% and 250% of the established animal and/or human dosage. Where no animal and/or human dosage is established, as will be the case for newly-discovered pharmaceutical compositions, a suitable animal and/or human dosage can be inferred from ED₅₀ or ID₅₀ values, or other appropriate values derived from *in vitro* or *in vivo* studies, as qualified by toxicity studies and efficacy studies in animals.

[00147] In cases of administration of a pharmaceutically acceptable salt, dosages may be calculated as the free base. As will be understood by those of skill in the art, in certain situations it may be necessary to administer the compounds disclosed herein in amounts that exceed, or even far exceed, the above-stated, preferred dosage range in order to effectively and aggressively treat particularly aggressive diseases or conditions.

[00148] From preliminary studies, it found that toxicity for CSA compounds was greatly reduced at 5µM and lower doses. An optimal standalone dose of CSAs may be 10µM or less at, on, or near the tissue wound, but smaller amounts may be used to minimize toxicity as needed.

Kits


[00149] Kits comprising the tissue treatment compositions and instructions for performing such methods are also disclosed. The disclosure also provides kits including compounds (*e.g.*, CSA), combination compositions and pharmaceutical compositions/formulations thereof, packaged into a suitable packaging material. In one embodiment, a kit includes packaging material, a CSA, and instructions. In various aspects, the instructions are for administering the CSA to enhance wound healing in a subject with a tissue wound. Where the tissue treatment composition is to be sprayed, the kit may include a spray container.

[00150] The term “packaging material” refers to a physical structure housing one or more components of the kit. The packaging material can maintain the components sterily, and can be made of material commonly used for such purposes (*e.g.*, paper, corrugated fiber, glass, plastic, foil, ampules, vials, tubes, etc.). A kit can contain a plurality of components, *e.g.*, two or more compounds alone or in combination with a wound healing agent or treatment or drug, optionally sterile.

[00151] A kit optionally includes a label or insert including a description of the components (type, amounts, doses, etc.), instructions for use *in vitro*, *in vivo*, or *ex vivo*, and any other components therein. Labels or inserts include “printed matter,” *e.g.*, paper or cardboard, or separate or affixed to a component, a kit or packing material (*e.g.*, a box), or attached to an ampule, tube or vial containing a kit component. Labels or inserts can additionally include a computer readable medium, such as a disk (*e.g.*, floppy diskette, hard disk, ZIP disk), optical disk such as CD- or DVD-ROM/RAM, DVD, MP3, magnetic tape, or an electrical storage media such as RAM and ROM or hybrids of these such as magnetic/optical storage media, FLASH media or memory type cards.

Examples

[00152] Example 1: A horse with a chronic non-healing oral cutaneous fistula was first treated with two prior art treatments with no success. The first treatment was with an aqueous spray of Vetericyn (0.007% hypochlorite) for a period of 7 days. The second unsuccessful treatment used Furacin Salve rubbed onto the open wound and applied daily for a period of 5 days with no appreciable improvement in wound healing. The wound was successfully treated in 7 days by spraying a composition of 0.04% CSA in water to the wound once daily for 7 days. The CSA-44 had the following structure



[00153] Closer of the fistula and healthy tissue were achieved in 7 days.

[00154] Example 2: A horse with a torn right front shoulder on a T post had a wound that was approximately 4 inches by 6 inches and 3 inches deep. This wound occurs commonly in horses and usually takes 3 to 6 months to heal. Typically this wound will result in a significant loss of muscle in the region. This wound was treated twice a day with a 0.04% solution of CSA-44 in water. The tissue treatment composition was applied by spraying sufficient composition to wet the entire surface of the wound. The wound closed and completely healed in 28 days without any apparent muscle loss of muscle. This result is surprising and unexpected given the location and size of wound and the typical length of time typically need for this type of wound to heal.

[00155] Example 3: A reining broodmare in Texas was involved in an accident resulting in a laceration to her left hind cannon region. The wound was 3 inches in length and 2 inches wide. The laceration extended all the way to the cannon bone, exposing the periosteum. This wound was also treated with a 0.04% solution of CSA-44 in water twice daily. No wrap was applied (i.e., no immobilization). The wound closed in 11 days with normal hair growth at 30 days. This type of wound would usually require aggressive wraps for 3 months with the possibility of other procedures being necessary for the excessive granulation tissue that commonly develops.

[00156] Example 4: A horse with an extreme case of scratches was successfully healed using CSA. Scratches is a fungal infection of the lower extremities in horses. This infection usually results in not only a fungal but also a secondary bacterial infection. In this case, the horse developed severe and debilitating granulation tissue of the ankle resulting in poor mobility due to cracking and bleeding. Prior to treatment with CSA, the wound was unsuccessfully treated with an oral antifungal, antibiotic (Naxcel), and Veterycin. The chronic infection persisted for over two years despite the multiple attempts to treat with an antimicrobial. Due to the debilitating wound the horse was slated for slaughter. However, the wound was successfully treated with a 0.04% of CSA-44 in water sprayed twice daily for four weeks. Normal hair growth and tissue was present in 4 weeks.

[00157] Example 5: A show mare developed an uncontrollable fungal infection of the skin. This malady is commonly referred to as summer itch. A 0.04% solution of CSA-44 in water was used twice daily for 3 weeks resulting in complete regrowth of normal skin and hair. Normal treatment time periods for this problem usually range from 3 to 4 months.

5 [00158] Example 6: Genetically diabetic 8-week-old female mice (db/db, BKS.Cg-m +/+ Leprdb/J) and heterozygous nondiabetic littermates (db/-) were purchased from The Jackson Laboratory (Bar Harbor, Me.). Strain and age of mouse were chosen because these mutant mice exhibit severe diabetic conditions, with hyperglycemia peaking between 8 and 12 weeks. They have also been shown to have delayed wound healing with occlusive dressings. Mice were
10 ordered to arrive at 7 weeks of age in order to decrease the chance of travel stress effect as a confounding factor in our experiment. Mice were housed individually in the Veteran's Affairs Medical Center – San Francisco Animal Research Facility, maintained on a 12-hour light/dark cycle, and allowed ad libitum access to rodent chow and water. Mice were anesthetized with an intraperitoneal injection of diluted chloral hydrate syrup (7 ½ gr/5ml). Approximately 0.2
15 and 0.1 ml of a solution containing 5ml of syrup diluted with 7ml of ddH₂O was given to each db/db and db/- mouse respectively. For each mouse, the dorsal skin was shaved, and cleansed with alcohol swabs. Mice were kept warm during anesthesia and surgery by being huddled together until they regained consciousness. Two full-thickness 6-mm punch biopsy (Acuderm, Inc., Ft. Lauderdale, FL.) wounds were created on the dorsal surface of the mice. Wounds were
20 placed approximately 1 – 2 cm apart. Areas that were chosen were free of anagen hair follicles. A piece of CSA impregnated film, approximately 6mm in diameter, was placed in the wound. This film lay adjacent, touching the wound edges. Mastisol was applied to the edges of each wound, which was then covered with a 1 x 1 cm Tegaderm semioclusive dressing. To analyze the effect of the CSA impregnated film, digital photographs of wounds, including a metric
25 ruler, were taken on day zero and then every fourth day until complete wound closure was achieved. Prior to postoperative day 15, if Tegaderm dressing and/or impregnated film fell off, they were promptly replaced with a new dressing and/or film. On postoperative day 15, Tegaderm dressings were removed and wounds were allowed to close by contraction. On subsequent days, scabs were gently removed from the wounds not covered because they
30 decrease the rate of wound contraction and impair assessment of wound closure. All wounds were compared to their original wound size using ImageJ (Rasband WS, NIH, 1997-2006).

[00159] Example 7: To determine the role of synthetic Ceragenins CSA-13, 44 and 90 in wound healing using mesenchymal stem cells (MSC), targeted mRNA panels from

SABiosciences, and primary cells from Lonza were selected. Cells were purchased from Lonza.com and used fresh for each test using recommended media and culture conditions. After treatment, mRNA was isolated using Qiagen RNeasy Mini Kit®, and quantified using a NanoDrop 2000® by UV at 260 nm and 260/280 ratio for purity. cDNA was made using a

5 First Strand Kit® from SABiosciences and processed for real time PCR using a kit from the same company for selected analysis of wound healing pathways. Results from q-PCR were uploaded to the SABiosciences site and to Ingenuity.com web site for analysis and pathway mapping. SABiosciences wound healing array plates (Cat# PAHS-121) and innate/acquired immune response plates (PAHS-052) were used. These arrays are fully validated when used

10 as recommended. On day 1, primary human MSC cells were plated at 200,000 cells/well using 6-well plates with 3ml of recommended media—hMSC Basal Medium + BulletKit (50ml Growth Supplement, 10ml L-Glutamine and 0.5ml Gentamicin Sulfate Amphotercin-B) for 24 hours. Only early passages of cells were used, and never from frozen stock. On day 2, cells were treated with compounds dissolved in DMSO diluted 1:1000 or more to

15 avoid effects of the solvent. Final testing concentration for CSA-13 was 5.0 µM. Treatment lasted 8 hours, and was followed by RNA isolation using QIAGEN RNeasy Mini Kit® (74104). RNA was measured at 260/280 nm using a NanoDrop 2000® and normalized to 2.4 ng per well, cDNA preparation was done using QIAGEN First Strand kit 330401. q-PCR was run as absolute quantification and threshold set at 0.1 units. Dendritic cells were plated

20 at 500,000 cells/well using 24-well plate with 500ul of Lonza LGM-3 Complete Growth Medium with and without compound. Treatment lasted 8 hours, and was followed by RNA isolation using QIAGEN RNeasy Mini Kit® (74104). RNA was measured at 260/280 nm using NanoDrop2000® and normalized to 2.4 ng per well, cDNA preparation was done using QIAGEN First Strand kit 330401. PCR was run as absolute quantification and threshold set at

25 0.1 units. For the wound healing array, strong upregulation of growth factors such as HB-EGF and cell migration factors such as MMP1 and CXCL2 were found, indicating a clear potential for CSA as a modulator of wound healing. Additional gene expression data is provided in Tables 1-3 for CSA-13, 44, and 90, respectively

[00160] Table 1: Gene Expression Results for CSA-13

Gene Symbol	Fold Regulation
CCL7	1.6632
CXCL1	1.6181
CXCL2	4.873
CXCL5	2.0582

F13A1	2.0916
FGF10	3.8659
HBEGF	3.255
IL2	1.865
IL6	3.1692
ITGA2	3.5659
MMP1	4.4172
PLAU	1.7849
PLAUR	1.6286
PTGS2	3.3333
VEGFA	1.7274
VTN	2.0612
ANGPT1	-2.0046
CSF2	-2.4867
F3	-3.3945
FGF2	-1.633
IL10	-1.6166
IL4	-1.9944
ITGB3	-1.5243
PLAT	-2.1487

[00161] Table 2: Gene Expression Results for CSA-44

Gene Symbol	Fold Regulation
CCL7	2.1961
COL1A2	1.5483
COL3A1	1.7385
CTSK	1.6388
CTSL2	1.7924
CXCL2	14.3964
EGFR	1.5364
F13A1	2.0963
FGF10	2.2811
FGF7	4.84

HBEGF	3.5463
HGF	3.1098
IGF1	1.6877
IL2	2.1928
IL6	4.0387
ITGA2	16.0648
ITGB6	1.6323
MMP1	68.9688
MMP9	1.5543
PLAU	1.6131
PLAUR	2.5454
PTGS2	48.6907
TIMP1	1.6126
VEGFA	4.6052
ACTA2	-1.9377
ANGPT1	-2.0857
CCL2	-3.1925
CDH1	-2.7158
COL4A3	-2.4845
CSF2	-1.8551
CTGF	-24.5295
FGF2	-1.6016
IL10	-2.0128
ITGB3	-1.5802
PLAT	-1.754
SERPINE1	-2.9618
TGFBR3	-2.0462
WISP1	-1.9722
ACTB	-1.7981

[00162] Table 2: Gene Expression Results for CSA-90

Gene Symbol	Fold Regulation
CCL7	2.2874


CTSK	1.5366
CTSL2	1.6306
CXCL1	3.1083
CXCL2	36.5878
EGFR	1.6212
F13A1	2.1032
FGF10	1.9842
FGF7	5.1689
HBEGF	3.5988
HGF	2.3334
IFNG	1.839
IGF1	2.2724
IL2	1.8522
IL6	7.2299
ITGA2	14.3637
ITGB6	1.9381
MMP1	45.3753
MMP9	2.5652
PLAUR	1.9924
PTGS2	66.6189
TIMP1	1.8212
VEGFA	3.3257
ACTA2	-2.5014
ANGPT1	-1.915
CCL2	-1.5793
COL4A3	-1.6195
CTGF	-15.7171
IL10	-1.7382
ITGA3	-1.5127
ITGB3	-1.538
PLAT	-2.14
SERPINE1	-3.7307
TGFBR3	-1.5938

WISP1	-2.3543
ACTB	-2.0745

[00163] Furthermore, although the foregoing has been described in some detail by way of illustrations and examples for purposes of clarity and understanding, it will be understood by those of skill in the art that numerous and various modifications can be made without departing from the spirit of the present disclosure. Therefore, it should be clearly understood that the forms disclosed herein are illustrative only and are not intended to limit the scope of the present disclosure, but rather to also cover all modification and alternatives coming with the true scope and spirit of the invention.

The embodiments of the present invention for which an exclusive property or privilege is claimed are defined as follows:

1. A composition comprising: a cationic steroidal anti-microbial (CSA) compound of Formula (Ia), or a pharmaceutically acceptable salt thereof, for use in increasing the rate of healing of a tissue wound:



where,

R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted alkyl, a substituted or unsubstituted hydroxyalkyl, a substituted or unsubstituted alkyloxyalkyl, a substituted or unsubstituted alkylcarboxyalkyl, a substituted or unsubstituted alkylaminoalkyl, a substituted or unsubstituted alkylaminoalkylamino, a substituted or unsubstituted alkylaminoalkylaminoalkylamino, a substituted or unsubstituted aminoalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted arylaminoalkyl, a substituted or unsubstituted haloalkyl, a substituted or unsubstituted alkenyl, a substituted or unsubstituted alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted aminoalkyloxy, a substituted or unsubstituted aminoalkyloxyalkyl, a substituted or unsubstituted aminoalkylcarboxy, a substituted or unsubstituted aminoalkylaminocarbonyl, a substituted or unsubstituted aminoalkylcarboxamido, a substituted or unsubstituted di(alkyl)aminoalkyl, a substituted or unsubstituted C-carboxyalkyl, H₂N-HC(Q₅)-C(O)-O-, H₂N-HC(Q₅)-C(O)-N(H)-, a substituted or unsubstituted azidoalkyloxy, a substituted or unsubstituted cyanoalkyloxy, P.G.-HN-HC(Q₅)-C(O)-O-, a substituted or unsubstituted guanidinoalkyloxy, a substituted or unsubstituted quaternary ammonium alkylcarboxy, and a substituted or unsubstituted guanidinoalkyl carboxy, where Q₅ is a side chain of any amino acid, and P.G. is an amino protecting group;

provided that at least two or three of R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of a substituted or unsubstituted aminoalkyl, a substituted or unsubstituted aminoalkyloxy, a substituted or unsubstituted alkylcarboxyalkyl, a substituted or unsubstituted alkylaminoalkylamino, a substituted or unsubstituted alkylaminoalkylaminoalkylamino, a substituted or unsubstituted aminoalkylcarboxy, a substituted or unsubstituted arylaminoalkyl, a substituted or unsubstituted aminoalkyloxyaminoalkylaminocarbonyl, a substituted or unsubstituted aminoalkylaminocarbonyl, a substituted or unsubstituted aminoalkylcarboxyamido, a quaternary ammonium alkylcarboxy, a substituted or unsubstituted di(alkyl)aminoalkyl, a substituted or unsubstituted C-carboxyalkyl, H₂N-HC(Q₅)-C(O)-O-, H₂N-HC(Q₅)-C(O)-N(H)-, azidoalkyloxy, cyanoalkyloxy, P.G.-HN-HC(Q₅)-C(O)-O-, a substituted or unsubstituted guanidinoalkyloxy, and a substituted or unsubstituted guanidinoalkylcarboxy; and

a pharmaceutically acceptable excipient.

2. The composition of claim 1, wherein Q₅ is a side chain of glycine.
3. The composition of claim 1 or claim 2, wherein the tissue wound is an open wound.
4. The composition of claim 1 or claim 2, wherein the tissue wound is a closed wound.
5. The composition of any one of claims 1 to 4, wherein the tissue wound is a tissue laceration, an incision site, or a chronic infection.
6. The composition of any one of claims 1 to 5, wherein the composition is formulated for application to the tissue wound in at least four separate applications and/or over a period of at least 4 days.
7. The composition of any one of claims 1 to 6, wherein the composition is formulated for application to the tissue wound by spraying.

8. The composition of any one of claims 1 to 7, wherein the composition is formulated for administration from a pharmaceutically acceptable device selected from the group consisting of bandages, surgical dressings, gauzes, adhesive strips, surgical staples, clips, hemostats, intrauterine devices, sutures, trocars, catheters, tubes, and implants.

9. The composition of any one of claims 1 to 8, wherein the composition is formulated for application to a tissue wound of a vertebrate.

10. The composition of any one of claims 1 to 9, wherein the composition is formulated for application to a tissue wound of a mammal.

11. The composition of any one of claims 1 to 10, wherein the composition is formulated for application to a tissue wound of a non-human mammal.

12. The composition of any one of claims 1 to 10, wherein the composition is formulated for application to a tissue wound of a human, horse, or dog.

13. The composition of any one of claims 1 to 12, wherein the composition is formulated for application to a tissue wound of a newborn mammal and the tissue wound is a navel of the newborn mammal.

14. The composition of any one of claims 1 to 13, wherein the composition is formulated for veterinary use.

15. The composition of any one of claims 1 to 14, wherein the tissue wound is not a burn.

16. The composition of any one of claims 1 to 15, wherein the tissue wound includes a chronic infection.

17. The composition of any one of claims 1 to 16, wherein the composition includes the CSA compound, or a pharmaceutically acceptable salt thereof, in a concentration in a range between 0.01-5% wt/wt.

18. The composition of any one of claims 1 to 17, wherein the composition includes the CSA compound, or a pharmaceutically acceptable salt thereof, in a concentration in a range between about 0.01-2.0% wt/wt.

19. The composition of any one of claims 1 to 18, wherein the composition includes the CSA compound, or a pharmaceutically acceptable salt thereof, in a concentration of about 0.04% wt/wt.

20. The composition of any one of claims 1 to 19, wherein the plurality of cationic groups are each attached to the steroidal group through a hydrolysable ester linkage.





21. The composition of any one of claims 1 to 20, wherein:

R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ hydroxyalkyl, a substituted or unsubstituted C₁-C₁₈ alkyloxy-C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ alkylcarboxy-C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ alkylamino-C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ alkylamino-C₁-C₁₈ alkylamino, a substituted or unsubstituted C₁-C₁₈ alkylamino-C₁-C₁₈ alkylamino, a substituted or unsubstituted C₁-C₁₈ aminoalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted arylamino-C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ haloalkyl, a substituted or unsubstituted C₂-C₆ alkenyl, a substituted or unsubstituted C₂-C₆ alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted C₁-C₁₈ aminoalkyloxy, a substituted or unsubstituted C₁-C₁₈ aminoalkyloxy-C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ aminoalkylcarboxy, a substituted or unsubstituted C₁-C₁₈ aminoalkylaminocarbonyl, a substituted or unsubstituted C₁-C₁₈ aminoalkylcarboxamido, a substituted or unsubstituted di C₁-C₁₈ alkyl aminoalkyl, a substituted or unsubstituted C-carboxy C₁-C₁₈ alkyl, H₂N-HC(Q₅)-C(O)-O-, H₂N-HC(Q₅)-C(O)-N(H)-, a substituted or unsubstituted C₁-C₁₈ azidoalkyloxy, a substituted or unsubstituted C₁-C₁₈ cyanoalkyloxy, P.G.-HN-HC(Q₅)-C(O)-O-, a substituted or unsubstituted C₁-C₁₈ guanidinoalkyloxy, a substituted or unsubstituted C₁-C₁₈ quaternary ammonium alkylcarboxy, and a substituted or unsubstituted C₁-C₁₈ guanidinoalkyl carboxy, where

provided that at least two or three of R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of a substituted or unsubstituted C₁-C₁₈ aminoalkyl, a substituted or unsubstituted C₁-C₁₈ aminoalkyloxy, a substituted or unsubstituted C₁-C₁₈ alkylcarboxy-C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ alkylamino-C₁-C₁₈ alkylamino, a substituted or unsubstituted C₁-C₁₈ alkylamino-C₁-C₁₈ alkylamino C₁-C₁₈ alkylamino, a substituted or unsubstituted C₁-C₁₈ aminoalkylcarboxy, a substituted or unsubstituted arylamino C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ aminoalkyloxy C₁-C₁₈ aminoalkylaminocarbonyl, a substituted or unsubstituted C₁-C₁₈ aminoalkylaminocarbonyl, a substituted or unsubstituted C₁-C₁₈ aminoalkylcarboxyamido, a substituted or unsubstituted C₁-C₁₈ quaternary ammonium alkylcarboxy, a substituted or unsubstituted di C₁-C₁₈ alkyl aminoalkyl, a substituted or unsubstituted C-carboxy C₁-C₁₈ alkyl, H₂N-HC(Q₅)-C(O)-O-, H₂N-HC(Q₅)-C(O)-N(H)-, a substituted or unsubstituted C₁-C₁₈ azidoalkyloxy, a substituted or unsubstituted C₁-C₁₈ cyanoalkyloxy, P.G.-HN-HC(Q₅)-C(O)-O-, a substituted or unsubstituted C₁-C₁₈ guanidinoalkyloxy, and a substituted or unsubstituted C₁-C₁₈ guanidinoalkylcarboxy.


23. The composition of any one of claims 1 to 22, wherein R₁₈ is alkylaminoalkyl, di(alkyl)aminoalkyl, alkoxycarbonylalkyl, alkylcarboxyalkyl, or C-carboxyalkyl.

The chemical structure shows a steroid nucleus with several modifications. At the 3-position, there is a primary amine group (H₂N) attached via an ether linkage (O). At the 17-position, there is a long alkyl chain (CH₂)₁₀ attached via an amine group (NH). At the 20-position, there is a primary amine group (H₂N) attached via an ether linkage (O). The structure also includes a methyl group at the 13-position and a hydroxyl group at the 14-position. Stereochemistry is indicated with wedges and dashes.




pharmaceutically acceptable salts thereof.

25. The composition of any one of claims 1 to 24, wherein the CSA, or a pharmaceutically acceptable salt thereof, is:



or a pharmaceutically acceptable salt thereof.

26. Use of a cationic steroid antimicrobial (CSA) compound of Formula (Ia), or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for increasing the rate of healing of a tissue wound:



where

R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted alkyl, a substituted or unsubstituted hydroxyalkyl, a substituted or unsubstituted alkyloxyalkyl, a substituted or unsubstituted alkylcarboxyalkyl, a substituted or unsubstituted alkylaminoalkyl, a substituted or unsubstituted alkylaminoalkylamino, a substituted or unsubstituted aminoalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted arylaminoalkyl, a substituted or unsubstituted haloalkyl, a substituted or unsubstituted alkenyl, a substituted or unsubstituted alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted aminoalkyloxy, a substituted or unsubstituted aminoalkyloxyalkyl, a substituted or unsubstituted aminoalkylcarboxy, a substituted or unsubstituted aminoalkylaminocarbonyl, a substituted or unsubstituted aminoalkylcarboxamido, a substituted or unsubstituted di(alkyl)aminoalkyl, a substituted or unsubstituted C-carboxyalkyl, H₂N-HC(Q₅)-C(O)-O-, H₂N-HC(Q₅)-

C(O)-N(H)-, a substituted or unsubstituted azidoalkyloxy, a substituted or unsubstituted cyanoalkyloxy, P.G.-HN-HC(Q₅)-C(O)-O-, a substituted or unsubstituted guanidinoalkyloxy, a substituted or unsubstituted quaternary ammonium alkylcarboxy, and a substituted or unsubstituted guanidinoalkyl carboxy, where Q₅ is a side chain of any amino acid, and P.G. is an amino protecting group;

provided that at least two or three of R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of a substituted or unsubstituted aminoalkyl, a substituted or unsubstituted aminoalkyloxy, a substituted or unsubstituted alkylcarboxyalkyl, a substituted or unsubstituted alkylaminoalkylamino, a substituted or unsubstituted alkylaminoalkylaminoalkylamino, a substituted or unsubstituted aminoalkylcarboxy, a substituted or unsubstituted arylaminoalkyl, a substituted or unsubstituted aminoalkyloxyaminoalkylaminocarbonyl, a substituted or unsubstituted aminoalkylaminocarbonyl, a substituted or unsubstituted aminoalkylcarboxyamido, a quaternary ammonium alkylcarboxy, a substituted or unsubstituted di(alkyl)aminoalkyl, a substituted or unsubstituted C-carboxyalkyl, H₂N-HC(Q₅)-C(O)-O-, H₂N-HC(Q₅)-C(O)-N(H)-, azidoalkyloxy, cyanoalkyloxy, P.G.-HN-HC(Q₅)-C(O)-O-, a substituted or unsubstituted guanidinoalkyloxy, and a substituted or unsubstituted guanidinoalkylcarboxy.

27. The use according to claim 26, wherein Q₅ is a side chain of glycine.
28. The use of claim 26 or claim 27, wherein the tissue wound is an open wound.
29. The use of claim 26 or claim 27, wherein the tissue wound is a closed wound.
30. The use of any one of claims 26 to 29, wherein the tissue wound is a tissue laceration, an incision site, or a chronic infection.
31. The use of any one of claims 26 to 30, wherein the medicament is formulated for application to the tissue wound in at least four separate applications and/or over a period of at least 4 days.
32. The use of any one of claims 26 to 31, wherein the medicament is formulated

for application to the tissue wound by spraying.

33. The use of any one of claims 26 to 31, wherein the medicament is formulated for administration from a pharmaceutically acceptable device selected from the group consisting of bandages, surgical dressings, gauzes, adhesive strips, surgical staples, clips, hemostats, intrauterine devices, sutures, trocars, catheters, tubes, and implants.

34. The use of any one of claims 26 to 33, wherein the medicament is formulated for application to a tissue wound of a vertebrate.

35. The use of any one of claims 26 to 34, wherein the medicament is formulated for application to a tissue wound of a mammal.

36. The use of any one of claims 26 to 35, wherein the medicament is formulated for application to a tissue wound of a non-human mammal.

37. The use of any one of claims 26 to 35, wherein the medicament is formulated for application to a tissue wound of a human, horse, or dog.

38. The use of any one of claims 26 to 37, wherein the medicament is formulated for application to a tissue wound of a newborn mammal and the tissue wound is a navel of the newborn mammal.

39. The use of any one of claims 26 to 38, wherein the medicament is formulated for veterinary use.

40. The use of any one of claims 26 to 39, wherein the tissue wound is not a burn.

41. The use of any one of claims 26 to 40, wherein the tissue wound includes a chronic infection.

42. The use of any one of claims 26 to 41, wherein the composition includes the CSA compound, or a pharmaceutically acceptable salt thereof, in a concentration in a range

between 0.01-5% wt/wt.

43. The use of any one of claims 26 to 42, wherein the composition includes the CSA compound, or a pharmaceutically acceptable salt thereof, in a concentration in a range between about 0.01-2.0% wt/wt.

44. The use of any one of claims 26 to 43, wherein the composition includes the CSA compound, or a pharmaceutically acceptable salt thereof, in a concentration of about 0.04% wt/wt.

45. The use of any one of claims 26 to 44, wherein the plurality of cationic groups are each attached to the steroidal group through a hydrolysable ester linkage.

46. The use of any one of claims 26 to 45, wherein:

R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of hydrogen, hydroxyl, a substituted or unsubstituted C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ hydroxyalkyl, a substituted or unsubstituted C₁-C₁₈ alkyloxy-C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ alkylcarboxy-C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ alkylamino-C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ alkylamino-C₁-C₁₈ alkylamino, a substituted or unsubstituted C₁-C₁₈ alkylamino-C₁-C₁₈ alkylamino, a substituted or unsubstituted C₁-C₁₈ aminoalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted arylamino(C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ haloalkyl, a substituted or unsubstituted C₂-C₆ alkenyl, a substituted or unsubstituted C₂-C₆ alkynyl, oxo, a linking group attached to a second steroid, a substituted or unsubstituted C₁-C₁₈ aminoalkyloxy, a substituted or unsubstituted C₁-C₁₈ aminoalkyloxy-C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ aminoalkylcarboxy, a substituted or unsubstituted C₁-C₁₈ aminoalkylaminocarbonyl, a substituted or unsubstituted C₁-C₁₈ aminoalkylcarboxamido, a substituted or unsubstituted di C₁-C₁₈ alkyl aminoalkyl, a substituted or unsubstituted C-carboxy C₁-C₁₈ alkyl, H₂N-HC(Q₅)-C(O)-O-, H₂N-HC(Q₅)-C(O)-N(H)-, a substituted or unsubstituted C₁-C₁₈ azidoalkyloxy, a substituted or unsubstituted C₁-C₁₈ cyanoalkyloxy, P.G.-HN-HC(Q₅)-C(O)-O-, a substituted or unsubstituted C₁-C₁₈ guanidinoalkyloxy, a substituted or unsubstituted C₁-C₁₈ quaternary ammonium


alkylcarboxy, and a substituted or unsubstituted C₁-C₁₈ guanidinoalkyl carboxy, where Q₅ is a side chain of any amino acid, and P.G. is an amino protecting group,

provided that at least two or three of R₃, R₇, R₁₂, and R₁₈ are independently selected from the group consisting of a substituted or unsubstituted C₁-C₁₈ aminoalkyl, a substituted or unsubstituted C₁-C₁₈ aminoalkyloxy, a substituted or unsubstituted C₁-C₁₈ alkylcarboxy-C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ alkylamino-C₁-C₁₈ alkylamino, a substituted or unsubstituted C₁-C₁₈ alkylamino-C₁-C₁₈ alkylamino C₁-C₁₈ alkylamino, a substituted or unsubstituted C₁-C₁₈ aminoalkylcarboxy, a substituted or unsubstituted arylamino C₁-C₁₈ alkyl, a substituted or unsubstituted C₁-C₁₈ aminoalkyloxy C₁-C₁₈ aminoalkylaminocarbonyl, a substituted or unsubstituted C₁-C₁₈ aminoalkylaminocarbonyl, a substituted or unsubstituted C₁-C₁₈ aminoalkylcarboxyamido, a substituted or unsubstituted C₁-C₁₈ quaternary ammonium alkylcarboxy, a substituted or unsubstituted di C₁-C₁₈ alkyl aminoalkyl, a substituted or unsubstituted C-carboxy C₁-C₁₈ alkyl, H₂N-HC(Q₅)-C(O)-O-, H₂N-HC(Q₅)-C(O)-N(H)-, a substituted or unsubstituted C₁-C₁₈ azidoalkyloxy, a substituted or unsubstituted C₁-C₁₈ cyanoalkyloxy, P.G.-HN-HC(Q₅)-C(O)-O-, a substituted or unsubstituted C₁-C₁₈ guanidinoalkyloxy, and a substituted or unsubstituted C₁-C₁₈ guanidinoalkylcarboxy.





47. The use of any one of claims 26 to 46, wherein at least R₃, R₇, and R₁₂ are independently an aminoalkyloxy or an aminoalkylcarboxy.

48. The use of any one of claims 26 to 47, wherein R₁₈ is alkylaminoalkyl, di(alkyl)aminoalkyl, alkoxycarbonylalkyl, alkylcarboxyalkyl, or C-carboxyalkyl.

49. The use of any one of claims 26 to 48, wherein the CSA, or a pharmaceutically acceptable salt thereof, is selected from the group consisting of:




;



pharmaceutically acceptable salts thereof.

50. The use of any one of claims 26 to 49, wherein the CSA, or a pharmaceutically acceptable salt thereof, is:



or a pharmaceutically acceptable salt thereof.

