



(51) International Patent Classification:

C07K 14/37 (2006.01) *CUD 3/386* (2006.01)
CUD 3/38 (2006.01) *C12N 9/20* (2006.01)

(21) International Application Number:

PCT/IB20 12/05 1660

(22) International Filing Date:

4 April 2012 (04.04.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

PCT/CN201 1/000614 8 April 201 1 (08.04.201 1) CN
61/561,044 17 November 201 1 (17. 11.201 1) US

(71) Applicant (for all designated States except US): **DAN-ISCO US, INC.** [US/US]; 925 Page Mill Road, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SHIPOVSKOV, Stepan** [RU/DK]; Kystparkvej 18, DK-8250 Ega (DK). **JENSEN, Lene Bojsen** [DK/DK]; Provsteparkvej 26, DK-8270 Højbjerg (DK). **QIAN, Zhen** [CN/CN]; Room 502, No 19, Lane 47 Yanzhong Road, Shanghai 200125 (CN).

(74) Agent: **DUNCAN, Garreth**; D Young & Co LLP, 120 Holborn, London EC1N 2DY (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

— with sequence listing part of description (Rule 5.2(a))

(54) Title: COMPOSITIONS

(57) Abstract: A composition comprising: (a) a lipolytic enzyme; (b) a hydrophobin, as defined herein; and optionally (c) a detergent; is provided. The composition is useful as a cleaning composition for removing lipid-based stains from surfaces.



WO 2012/137147 A1

COMPOSITIONSField of the invention

5 This invention relates to a composition, particularly although not exclusively for use as a detergent. The invention also relates to methods of cleaning surfaces and items, such as clothing items and tableware items, using the composition.

Background to the Invention

10

As described in Wosten, *Annu. Rev. Microbiol.* 2001, 55, 625-646, hydrophobins are proteins generally of fungal origin that play a broad range of roles in the growth and development of filamentous fungi. For example, they are involved in the formation of aerial structures and in the attachment of hyphae to hydrophobic surfaces.

15

The mechanisms by which hydrophobins perform their function are based around their property to self-assemble at hydrophobic-hydrophilic interfaces (particularly air-water interfaces) into an amphipathic film.

20

Typically, hydrophobins are divided into Classes I and II. As described in more detail herein, the assembled amphipathic films of Class II hydrophobins are capable of redissolving in a range of solvents (particularly although not exclusively an aqueous ethanol) at room temperature. In contrast, the assembled amphipathic films of Class I hydrophobins are much less soluble, redissolving only in strong acids such as

25

trifluoroacetic acid or formic acid.

Detergent compositions containing hydrophobins are known in the art. For example, US 2009/0101 167 (corresponding to WO 2007/014897) describes the use of hydrophobins, particularly fusion hydrophobins, for washing textiles and washing compositions containing them.

30

There remains a need in the art for detergent compositions containing surfactants capable of being used in smaller quantities and thereby minimising impact on the environment.

35

Summary of the Invention

According to one aspect of the invention, there is provided a composition comprising:

- (a) a lipolytic enzyme; and
- 5 (b) a hydrophobin, as defined herein.

According to another aspect of the invention, there is provided a composition comprising:

- (a) a lipolytic enzyme;
- 10 (b) a hydrophobin, as defined herein; and
- (c) a detergent.

According to one aspect of the invention, there is provided a composition comprising:

- (a) a GX lipolytic enzyme, wherein G is glycine and X is an oxyanion hole-forming
- 15 amino acid residue, wherein the GX lipolytic enzyme belongs to an alpha/beta hydrolase superfamily selected from the group consisting of abH23, abH25, and abH15; and
- (b) a hydrophobin, as defined herein.

20 According to another aspect of the invention, there is provided a composition comprising:

- (a) a GX lipolytic enzyme, wherein G is glycine and X is an oxyanion hole-forming amino acid residue;
- (b) a hydrophobin, as defined herein; and
- 25 (c) a detergent.

According to a yet further aspect of the invention, there is provided a method of removing a lipid-based stain from a surface by contacting the surface with a composition as defined herein.

30

According to a still further aspect of the invention, there is provided the use of a composition as defined herein to reduce or remove lipid stains from a surface.

35 According to a further aspect of the invention, there is provided a method of cleaning a surface, comprising contacting the surface with a composition as defined herein.

According to a further aspect of the invention, there is provided a method of cleaning an item, in particular a clothing item or a tableware item, comprising contacting the item **with** a composition as defined herein.

5 Advantages

It has surprisingly been found **that** the combination of hydrophobin, lipolytic enzyme and, optionally, detergent is capable of removing oily soils from surfaces, such as textile, clothing or tableware surfaces: it is generally problematic to remove such soils
10 using existing commercial detergents. This effect confers the potential for using the combination in washing compositions.

In particular, it has surprisingly been found that the combination of hydrophobin and GX lipolytic enzyme selected from the abH superfamilies referred to above exhibits a
15 greatly improved cleaning effect than would be expected from an additive effect of either of these proteins when used alone. These properties confer the potential for using the combination as a replacement for detergent in washing compositions, thereby minimising the environmental impact of such compositions.

It has also surprisingly been found that the combination of hydrophobin, GX lipolytic enzyme and detergent exhibits a greatly improved cleaning effect than would be expected from an additive effect of any of these three components when used alone. These properties confer the potential for using the combination to minimise the amount of detergent required in washing compositions, thereby minimising the
20 environmental impact of such compositions.
25

Brief Description of the Drawings

Fig. 1a shows the % change in Stain Removal index (SRI) as a function of the detergent concentration at various specified hydrophobin concentrations in the presence of heat-inactivated liquid detergent ARIEL™ Color, but in the absence of a lipolytic enzyme;
30

Fig. 1b shows the % change in SRI as a function of the hydrophobin concentration at various specified detergent concentrations in the presence of heat-inactivated liquid detergent ARIEL™ Color, but in the absence of a lipolytic enzyme;
35

- Fig. 1c shows the % change in SRI as a function of the detergent concentration at various specified hydrophobin concentrations in the presence of heat-inactivated powder detergent ARIEL™ Color, but in the absence of a lipolytic enzyme;
- Fig. 2a shows the % change in SRI as a function of the detergent concentration at various specified hydrophobin concentrations in the presence of the lipolytic enzyme LIPEX™ and the heat-inactivated liquid detergent ARIEL™ Color;
- Fig. 2b shows the % change in SRI as a function of the hydrophobin concentration at various specified detergent concentrations in the presence of the lipolytic enzyme LIPEX™ and the heat-inactivated liquid detergent ARIEL™ Color;
- Fig. 2c shows the % change in SRI as a function of the detergent concentration at various specified hydrophobin concentrations in the presence of the lipolytic enzyme LIPEX™ and the heat-inactivated powder detergent ARIEL™ Color;
- Fig. 2d shows the % change in SRI as a function of the hydrophobin concentration at various specified detergent concentrations in the presence of the lipolytic enzyme LIPEX™ and the heat-inactivated powder detergent ARIEL™ Color;
- Fig. 2e shows the % change in SRI as a function of the hydrophobin concentration in the presence of the lipolytic enzyme LIPEX™ but in the absence of detergent;
- Fig. 3a shows the % change in SRI as a function of the detergent concentration at various specified hydrophobin concentrations in the presence of the lipolytic enzyme LIPOMAX™ and the heat-inactivated liquid detergent ARIEL™ Color;
- Fig. 3b shows the % change in SRI as a function of the hydrophobin concentration at various specified detergent concentrations in the presence of the lipolytic enzyme LIPOMAX™ and the heat-inactivated liquid detergent ARIEL™ Color;
- Fig. 3c shows the % change in SRI as a function of the detergent concentration at various specified hydrophobin concentrations in the presence of the lipolytic enzyme LIPOMAX™ and the heat-inactivated powder detergent ARIEL™ Color;
- Fig. 3d shows the % change in SRI as a function of the hydrophobin concentration at various specified detergent concentrations in the presence of the lipolytic enzyme LIPOMAX™ and the heat-inactivated powder detergent ARIEL™ Color;
- Fig. 3e shows the % change in SRI as a function of the hydrophobin concentration in the presence of the lipolytic enzyme LIPOMAX™ but in the absence of detergent;
- Fig. 4a shows the % change in SRI as a function of the detergent concentration at various specified hydrophobin concentrations in the presence of the lipolytic enzyme SprLip2 and the heat-inactivated liquid detergent ARIEL™ Color;
- Fig. 4b shows the % change in SRI as a function of the hydrophobin concentration at various specified detergent concentrations in the presence of the lipolytic enzyme SprLip2 and the heat-inactivated liquid detergent ARIEL™ Color;

- Fig. 4c shows the % change in SRI as a function of the detergent concentration at various specified hydrophobin concentrations in the presence of the lipolytic enzyme SprUp2 and the heat-inactivated powder detergent ARIEL™ Color;
- Fig. 4d shows the % change in SRI as a function of the hydrophobin concentration at various specified detergent concentrations in the presence of the lipolytic enzyme SprLip2 and the heat-inactivated powder detergent ARIEL™ Color;
- Fig. 4e shows the % change in SRI as a function of the hydrophobin concentration in the presence of the lipolytic enzyme SprLip2 but in the absence of detergent;
- Fig. 5a shows the % change in SRI as a function of the detergent concentration at various specified hydrophobin concentrations in the presence of the lipolytic enzyme TfuLip2 and the heat-inactivated liquid detergent ARIEL™ Color;
- Fig. 5b shows the % change in SRI as a function of the hydrophobin concentration at various specified detergent concentrations in the presence of the lipolytic enzyme TfuLip2 and the heat-inactivated liquid detergent ARIEL™ Color;
- Fig. 5c shows the % change in SRI as a function of the detergent concentration at various specified hydrophobin concentrations in the presence of the lipolytic enzyme TfuLip2 and the heat-inactivated powder detergent ARIEL™ Color;
- Fig. 5d shows the % change in SRI as a function of the hydrophobin concentration at various specified detergent concentrations in the presence of the lipolytic enzyme TfuLip2 and the heat-inactivated powder detergent ARIEL™ Color;
- Fig. 5e shows the % change in SRI as a function of the hydrophobin concentration in the presence of the lipolytic enzyme TfuLip2 but in the absence of detergent;
- Fig. 6 shows SEQ ID NO: 1, the DNA sequence encoding the hydrophobin *Trichoderma reesei* HFBII (Y1 1894.1);
- Fig. 7 shows SEQ ID NO: 2, the amino acid sequence of the hydrophobin *Trichoderma reesei* HFBII (P79073.1);
- Fig. 8 shows SEQ ID NO: 3, the DNA sequence encoding the hydrophobin *Trichoderma reesei* HFBI (Z68124.1);
- Fig. 9 shows SEQ ID NO: 4, the amino acid sequence of the hydrophobin *Trichoderma reesei* HFBI (P52754.1);
- Fig. 10 shows SEQ ID NO: 5, the DNA sequence encoding the hydrophobin *Schizophyllum commune* SC3 (M32329.1);
- Fig. 11 shows SEQ ID NO: 6, the amino acid sequence of the hydrophobin *Schizophyllum commune* SC3 (AAA96324.1);
- Fig. 12 shows SEQ ID NO: 7, the DNA sequence encoding the hydrophobin *Neurospora crassa* EAS (X67339.1);

- Fig. 13 shows SEQ ID NO: 8, the amino acid sequence of the hydrophobin *Neurospora crassa* EAS (AAB24462.1);
- Fig. 14 shows SEQ ID NO: 9, *Talaromyces thermophilus* TT1 (the DNA sequence encoding the precursor TT1 hydrophobin, SEQ ID NO: 4 of US 7241734);
- 5 Fig. 15 shows SEQ ID NO: 10, *Talaromyces thermophilus* TT1 (the amino acid sequence of the precursor TT1 hydrophobin, SEQ ID NO: 3 of US 7241734);
- Fig. 16 shows SEQ ID NO: 11 the mature amino acid sequence of LIPEX™;
- Fig. 17 shows SEQ ID NO: 12 the full amino acid sequence for Sprl_ip2 (*Streptomyces pristinaespiralis* ATCC 25486 Uniprot B5H9Q8, NCB!:
- 10 ZP_06912654.1) with the signal sequence shown in bold;
- Fig. 18 shows SEQ ID NO: 13 the mature amino acid sequence of the *Fusarium heterosporum* phospholipase (disclosed in WO 2005/087918 and available from Danisco A/S as GRINDAMYL POWERBAKE 4100™);
- Fig. 19 shows SEQ ID NO: 29 the full amino acid sequence of Lipase 3 disclosed in
- 15 WO 98/45453, residues 1 to 270 comprise the mature sequence referred to herein as SEQ ID NO: 14;
- Fig. 19a shows SEQ ID NO: 14 the mature amino acid sequence of Lipase 3;
- Fig. 20 shows SEQ ID NO: 15 the mature amino acid sequence of LIPOMAX™;
- Fig. 21 shows SEQ ID NO: 16 the mature amino acid sequence of TfuLip2;
- 20 Fig. 22 shows SEQ ID NO: 17 the mature amino acid sequence of SprLip2;
- Fig. 23 shows SEQ ID NO: 18 the full amino acid sequence of LIPEX, including the signal sequence (amino acid residues 1 to 17), propeptide (amino acid residues 18 to 22) and mature sequence (amino acid residues 23 to 291 - shown in Fig. 16 as SEQ ID NO: 11);
- 25 Fig. 24 shows SEQ ID NO: 19 the full amino acid sequence of LIPOMAX, including the signal sequence (amino acid residues 1 to 24) and mature sequence (amino acid residues 25 to 313 - shown in Fig. 20 as SEQ ID NO: 15);
- Fig. 25 shows SEQ ID NO: 20 the full amino acid sequence of TfuLip2, including the signal sequence (amino acid residues 1 to 40) and mature sequence (amino acid
- 30 residues 41 to 301 - shown in Fig. 21 as SEQ ID NO: 16);
- Fig. 26 shows a protein preprosequence SEQ ID NO: 21 of a lipolytic enzyme from *Fusarium heterosporum* CBS 782.83 (wild type) disclosed in WO 2005/087918 - the preprosequence undergoes translational modification such that the mature form of the enzyme preferably comprises the enzyme shown in Fig. 18 as SEQ ID NO: 13; in
- 35 some host organisms the protein may be N-terminally processed such that a number of additional amino acids are added to the N or C terminus;

- Fig. 27 shows SEQ ID NO: 22 the nucleotide sequence of the synthesized *SprUp2* gene;
- Fig. 28 shows SEQ ID NO: 23 the nucleotide sequence of the *SprLip2* gene from expression plasmid pZQ205 (*celA* signal sequence is underlined);
- 5 Fig. 29 shows SEQ ID NO: 24 the amino acid sequence of *SprLip2* produced from plasmid pZQ205 (signal sequence is underlined);
- Fig. 30 shows the plasmid map of pZQ205 expression vector;
- Fig. 31 shows pNB hydrolysis by *SprLip2*;
- Fig. 32 shows pNPP hydrolysis by *SprLip2*;
- 10 Fig. 33 shows trioctanoate hydrolysis in the absence of detergent by *SprLip2*;
- Fig. 34 shows trioctanoate hydrolysis in the presence of detergent by *SprLip2*;
- Fig. 35 shows the performance of *SprLip2* in the presence and absence of detergent;
- Fig. 36 shows SEQ ID NO: 25, the amino acid sequence of a lipase from *Geobacillus stearothermophilus* strain T1 (GeoT1) which is available on the NCBI database as
- 15 accession number JC8061 (signal sequence is underlined);
- Fig. 37 shows SEQ ID NO: 26 the amino acid sequence of the BCE-GeoT1 fusion protein which is a fusion of SEQ ID NO: 25 and the carboxy-terminus of the catalytic domain of a bacterial cellulase;
- Fig. 38 shows SEQ ID NO: 27 the amino acid sequence of a lipase from *Bacillus subtilis* 168 (LipA) which is available as GENBANK Accession No. P37957 (signal
- 20 sequence is underlined);
- Fig. 39 shows SEQ ID NO: 28 the amino acid sequence of the BCE-LipA fusion protein which is a fusion of SEQ ID NO: 27 and the carboxy-terminus of the catalytic domain of a bacterial cellulase; and
- 25 Fig. 40 shows SEQ ID NO: 30 the nucleotide sequence of the *NsiI*-*MluI*-*HpaI* enzyme restriction sites before the *Bam*HI site.

Detailed Description of Preferred Embodiments

30 HYDROPHOBINS

In this specification the term "hydrophobin" is defined as meaning a polypeptide capable of self-assembly at a hydrophilic / hydrophobic interface, and having the general formula (I):

35

$$(Y_1)_n - B_1 - (X_1)_a - B_2 - (X_2)_{b-B3} - (X_3)_{c-B4-(X_4)} - (X_5)_{d-B5-(X_5)} - B_6 - (X_6)_r B_7 - (X_7)_g - B_8 - (Y_2)_m \quad (1)$$

wherein:

m and n are independently 0 to 2000;

B₁, B₂, B₃, B₄, B₅, B₆, B₇ and B₈ are each independently amino acids selected from Cys, Leu, Ala, Pro, Ser, Thr, Met or Gly, at least 6 of the residues B₁ through B₈ being Cys;

5 X₁, X₂, X₃, X₄, X₅, X₆, X₇, Y₁ and Y₂ independently represent any amino acid;

a is 1 to 50;

b is 0 to 5;

c is 1 to 100;

d is 1 to 100;

10 e is 1 to 50;

f is 0 to 5; and

g is 1 to 100.

Suitably, the hydrophobin has a sequence of between 40 and 120 amino acids in the
15 hydrophobin core. More preferably, the hydrophobin has a sequence of between 45
and 100 amino acids in the hydrophobin core. In one embodiment, the hydrophobin
has a sequence of between 50 and 90, preferably 50 to 75, and more preferably 55
to 65 amino acids in the hydrophobin core. In this specification the term "the
hydrophobin core" means the *sequence beginning with the* residue B₁ and
20 *terminating with the* residue B₈.

In the formula (I), at least 6, preferably at least 7, and most preferably all 8 of the
residues B₁ through B₈ are Cys.

25 In the formula (I), in one embodiment m is suitably 0 to 500, preferably 0 to 200,
more preferably 0 to 100, still more preferably 0 to 20, yet more preferably 0 to 10,
still more preferably 0 to 5, and most preferably 0.

In the formula (I), in one embodiment n is suitably 0 to 500, preferably 0 to 200, more
30 preferably 0 to 100, still more preferably 0 to 20, yet more preferably 0 to 10, and
most preferably 0 to 3.

In the formula (I), a is preferably 3 to 25, more preferably 5 to 15. In one
embodiment, a is 5 to 9.

35

In the formula (I), b is preferably 0 to 2, more preferably 0.

in the formula (i), c is preferably 5 to 50, more preferably 5 to 40. In one embodiment, c is 1 to 39.


in the formula (i), d is preferably 2 to 35, more preferably 4 to 23. In one embodiment, d is 8 to 23.

in the formula (i), e is preferably 2 to 15, more preferably 5 to 12. In one embodiment, e is 5 to 9.

In the formula (i), f is preferably 0 to 2, more preferably 0.

in the formula (i), g is preferably 3 to 35, more preferably 6 to 21. In one embodiment, g is 6 to 18.

Preferably, the hydrophobins used in the present invention have the general formula (II):



wherein:

m and n are independently 0 to 20;

B₁, B₂, B₃, B₄, B₅, B₆, B₇ and B₈ are each independently amino acids selected from Cys, Leu, Ala, Pro, Ser, Thr, Met or Gly, at least 7 of the residues B₁ through B₈ being Cys;

a is 3 to 25;

b is 0 to 2;

c is 5 to 50;

d is 2 to 35;


e is 2 to 15;

f is 0 to 2; and

g is 3 to 35.

In the formula (II), at least 7, and preferably all 8 of the residues B₁ through B₈ are Cys.

More preferably, the hydrophobins used in the present invention have the general formula (III):



wherein:

m and n are independently 0 to 20;

5 B₁, B₂, B₃, B₄, B₅, B₆, B₇ and B₈ are each independently amino acids selected from Cys, Leu, Ala, Pro, Ser, Thr, Met or Gly, at least 7 of the residues B₁ through B₈ being Cys;

a is 5 to 15;

c is 5 to 40;

d is 4 to 23;

10 e is 5 to 12; and

g is 6 to 21.

In the formula (III), at least 7, and preferably 8 of the residues B₁ through B₈ are Cys.

15 In the formulae (I), (II) and (III), when 6 or 7 of the residues B₁ through B₈ are Cys, it is preferred that the residues B₃ through B₇ are Cys.

In the formulae (I), (II) and (III), when 7 of the residues B₁ through B₈ are Cys, it is preferred that: (a) B₁ and B₃ through B₈ are Cys and B₂ is other than Cys; (b) B₁ through B₇ are Cys and B₈ is other than Cys, (c) B₁ is other than Cys and B₂ through B₈ are Cys. When 7 of the residues B₁ through B₈ are Cys, it is preferred that the other residue is Ser, Pro or Leu. In one embodiment, B₁ and B₃ through B₈ are Cys and B₂ is Ser. In another embodiment, B₁ through B₇ are Cys and B₈ is Leu. In a further embodiment, B₁ is Pro and B₂ through B₈ are Cys.

25

The cysteine residues of the hydrophobins used in the present invention may be present in reduced form or form disulfide (-S-S-) bridges with one another in any possible combination. In one particularly preferred embodiment, when all 8 of the residues B₁ through B₈ are Cys, disulfide bridges may be formed between one or more (preferably at least 2, more preferably at least 3, most preferably all 4) of the following pairs of cysteine residues: B₁ and B₆; B₂ and B₅; B₃ and B₄; B₇ and B₈. In one alternative preferred embodiment, when all 8 of the residues B₁ through B₈ are Cys, disulfide bridges may be formed between one or more (preferably at least 2, more preferably at least 3, most preferably all 4) of the following pairs of cysteine residues: B₁ and B₂; B₃ and B₄; B₅ and B₆; B₇ and B₈.

35

Examples of specific hydrophobic useful in the present invention include those described and exemplified in the following publications: Under *et al.*, *FEMS Microbiology Rev.* 2005, 29, 877-896; Kubicek *et al.*, *BMC Evolutionary Biology*, 2008, 8, 4; Sunde *et al.*, *Micron*, 2008, 39, 773-784; Wessels, *Adv. Micr. Physiol.* 1997, 33, 1-45; Wosten, *Annu. Rev. Microbiol.* 2001, 55, 625-646; Hektor and Scholtmeijer, *Curr. Opin. Biotech.* 2005, 16, 434-439; Szilvay *et al.*, *Biochemistry*, 2007, 46, 2345-2354; Kisko *et al.* *Langmuir*, 2009, 25, 1612-1619; Blijdenstein, *Soft Matter*, 2010, 6, 1799-1808; Wosten *et al.*, *EMBO J.* 1994, 13, 5848-5854; Hakanpaa *et al.*, *J. Biol. Chem.*, 2004, 279, 534-539; Wang *et al.*; *Protein Sci.*, 2004, 13, 810-821; De Vocht *et al.*, *Biophys. J.* 1998, 74, 2059-2068; Askolin *et al.*, *Biomacromolecules* 2006, 7, 1295-1301; Cox *et al.*; *Langmuir*, 2007, 23, 7995-8002; Linder *et al.*, *Biomacromolecules* 2001, 2, 511-517; Kallio *et al.* *J. Biol. Chem.*, 2007, 282, 28733-28739; Scholtmeijer *et al.*, *Appl. Microbiol. Biotechnol.*, 2001, 56, 1-8; Lumsdon *et al.*, *Colloids & Surfaces B: Biosurfaces*, 2005, 44, 172-178; Palomo *et al.*, *Biomacromolecules* 2003, 4, 204-210; Kirkland and Keyhani, *J. Ind. Microbiol. Biotechnol.*, July 17 2010 (e-publication); Stubner *et al.*, *Int. J. Food Microbiol.*, 30 June 2010 (e-publication); Laaksonen *et al.* *Langmuir*, 2009, 25, 5185-5192; Kwan *et al.* *J. Mol. Biol.* 2008, 382, 708-720; Yu *et al.* *Microbiology*, 2008, 154, 1677-1685; Lahtinen *et al.* *Protein Expr. Purif.*, 2008, 59, 18-24; Szilvay *et al.*, *FEBS Lett*, 2007, 5811, 2721-2726; Hakanpaa *et al.*, *Acta Crystallogr. D. Biol. Crystallogr.* 2006, 62, 356-367; Scholtmeijer *et al.*, *Appl. Environ. Microbiol.*, 2002, 68, 1367-1373; Yang *et al.*, *BMC Bioinformatics*, 2006, 7 Suppl. 4, S16; WO 01/57066; WO 01/57528; WO 2006/082253; WO 2006/103225; WO 2006/103230; WO 2007/014897; WO 2007/087967; WO 2007/087968; WO 2007/030966; WO 2008/019965; WO 2008/107439; WO 2008/1 10456; WO 2008/1 16715; WO 2008/120310; WO 2009/050000; US 2006/0228484; and EP 2042 156A; the contents of which are incorporated herein by reference.

In one embodiment, the hydrophobin is a polypeptide selected from SEQ ID NOs: 2, 4, 6 8 or 10, or a polypeptide having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, or at least 99% sequence identity in the hydrophobin core to any thereof and retaining the above-described self-assembly property of hydrophobins.

Sources of hydrophobins

In one embodiment, the hydrophobin is obtained or obtainable from a microorganism. The microorganism may preferably be a bacteria or a fungus, more preferably a
5 fungus. In a preferred embodiment, the hydrophobin is obtained or obtainable from a filamentous fungus.

In one embodiment, the hydrophobin is obtained or obtainable from fungi of the phyla Basidiomycota or Ascomycota.

10

In one embodiment, the hydrophobin is obtained or obtainable from fungi of the genera *Cladosporium* (particularly *C. fulvum* or *C. herbarum*), *Ophistoma* (particularly *O. ulmi*), *Cryphonectria* (particularly *C. parasitica*), *Trichoderma* (particularly *T. harzianum*, *T. longibrichiatum*, *T. asperellum*, *T. Koningiopsis*, *T. aggressivum*, *T. stromaticum* or *T. reesei*), *Gibberella* (particularly *G. moniliformis*), *Neurospora* (particularly *N. crassa*), *Maganaporthe* (particularly *M. grisea*), *Hypocrea* (particularly *H. jecorina*, *H. atroviridis*, *H. virens* or *H. lixii*), *Xanthoria* (particularly *X. ectanoides* and *X. parietina*), *Emericella* (particularly *E. nidulans*), *Aspergillus* (particularly *A. fumigatus*, *A. oryzae*), *Paracoccidioides* (particularly *P. brasiliensis*), *Metarhizium*
15 (particularly *M. anisoplaie*), *Pleurotus* (particularly *P. ostreatus*), *Coprinus* (particularly *C. cinereus*), *Dicotyonema* (particularly *D. glabratum*), *Flammulina* (particularly *F. velutipes*), *Schizophyllum* (particularly *S. commune*), *Agaricus* (particularly *A. bisporus*), *Pisolithus* (particularly *P. tinctorius*), *Tricholoma* (particularly *T. terreum*), *Pholioka* (particularly *P. nameko*), *Talaromyces* (particularly
20 *T. thermophilus*) or *Agrocybe* (particularly *A. aegerita*).

25

Assays

One property of the hydrophobins used in the present invention is the self-assembly
30 property of the hydrophobins at a hydrophilic / hydrophobic interface.

In accordance with the definition of the present invention, self-assembly can be detected by adsorbing the protein to polytetrafluoroethylene (TEFLON®) and using Circular Dichroism (CD) to establish the change in secondary structure exemplified
35 by the occurrence of motifs in the CD spectrum corresponding to a newly formed α -helix (De Vocht et al., *Biophys. J.* 1998, 74, 2059-2068). A full procedure for

carrying out the CD spectral analysis can be found in Askolin *et al.*
Biomacromolecules, 2006, 7, 1295-1301 .

in one embodiment, the hydrophobins used in the present invention are
5 characterised by their effect on the surface properties at an interface, particularly
although not exclusively at an air/water interface. The surface property may be
surface tension (especially equilibrium surface tension) or surface shear rheology,
particularly the surface shear elasticity (storage modulus).

10 In one embodiment, the hydrophobin may cause the equilibrium surface tension at a
water/air interface to reduce to below 45 mN/m, preferably below 40 mN/m, and more
preferably below 35 mN/m. In contrast, the surface tension of pure water is 72 mN/m
room temperature. Typically, such a reduction in the equilibrium surface tension at a
water/air interface may be achieved using a hydrophobin concentration of between 5
15 $\times 10^{-8}$ M and 2×10^{-6} M, more preferably between 1×10^{-7} M and 1×10^{-6} M.
Typically such a reduction in the equilibrium surface tension at a water/air interface
may be achieved at a temperature ranging from 0°C to 50°C, especially room
temperature. The change in equilibrium surface tension can be measured using a
tensiometer following the method described in Cox *et al.*, *Langmuir*, 2007, 23, 7995-
20 8002.

In another embodiment, the hydrophobin may cause the surface shear elasticity at a
water/air interface to increase to 300-700 mN/m, preferably 400-600 mN/m. Typically,
such a surface shear elasticity at a water/air interface may be achieved using a
25 hydrophobin concentration of between 1×10^{-4} M and 0.01 M, preferably between $5 \times$
 10^{-4} M and 2×10^{-3} M, especially 1×10^{-3} M. Typically, such a surface shear elasticity
at a water/air interface may be achieved at a temperature ranging from 0°C to 50°C,
especially room temperature. The change in equilibrium surface tension can be
measured using a rheometer following the method described in Cox *et al.*, *Langmuir*,
30 2007, 23, 7995-8002.

In some embodiments, the hydrophobins used in the present invention are
biosurfactants. Biosurfactants are surface-active substances synthesised by living
cells. They have the properties of reducing surface tension, stabilising emulsions,
35 promoting foaming and are generally non-toxic and biodegradable.

Examples of specific hydrophobins useful in the compositions of the present invention are listed in Table 1 below.

Table 1

Organism	Gene, Protein name	NCBI accession code and version number
<i>Agaricus bisporus</i>	ABH3	Y14602.1
<i>Agaricus bisporus</i>	HYPB	Y15940.1
<i>Aspergillus fumigatus</i>	HYP1/RODA	L25258.1, U06121.1
<i>Aspergillus fumigatus</i>	RODB	AY057385.1
<i>Aspergillus niger</i>	A_NIGi	XM_0G1394993.1
<i>Aspergillus oryzae</i>	HYPB	AB097448.1
<i>Aspergillus oryzae</i>	ROLA	AB094496.1
<i>Aspergillus terreus</i>	A_TER	XM_001213908.1
<i>Cladosporium fulvum</i>	HCF-5	AJ133703.1
<i>Cladosporium fulvum</i>	HCF-6	AJ251294.1
<i>Cladosporium fulvum</i>	HCF-3	AJ566186.1
<i>Cladosporium fulvum</i>	HCF-1	X98578.1
<i>Cladosporium fulvum</i>	HCF-2	AJ133700.1
<i>Cladosporium fulvum</i>	HCF-4	AJ566187.1
<i>Cladosporium herbarum</i>	HCH-1	AJ496190.1
<i>Claviceps fusiformis</i>	CFTH1J-III	AJ133774.1
<i>Claviceps fusiformis</i>	CLF	CAB61236.1
<i>Claviceps purpurea</i>	CLP	CAD10781.1
<i>Claviceps purpurea</i>	CPPH1J-V	AJ418045.1
<i>Coprinus cinereus</i>	COH1	Y10627.1
<i>Coprinus cinereus</i>	COH2	Y10628.1
<i>Cryphonectria parasitica</i>	CRP	L09559.1
<i>Dictyonema glabratum</i>	DGH3	AJ320546.1
<i>Dictyonema glabratum</i>	DGH2	AJ320545.1
<i>Dictyonema glabratum</i>	DGH1	AJ320544.1
<i>Emericella nidulans</i>	RODA	M61113.1
<i>Emericella nidulans</i>	DEWA	U07935.1
<i>Flammulina velutipes</i>	FVH1	AB026720.1
<i>Flammulina velutipes</i>	FvHYDI	AB126686.1
<i>Gibberella moniliformis</i>	HYD5, GIM	AY158024.1
<i>Gibberella moniliformis</i>	HYD4	AY155499.1
<i>Gibberella moniliformis</i>	HYD1	AY155496.1
<i>Gibberella moniliformis</i>	HYD2	AY155497.1
<i>Gibberella moniliformis</i>	HYD3	AY155498.1
<i>Gibberella zeae</i>	GIZ, FG01831.1	XP_382007.1
<i>Lentinula edodes</i>	Le.HYDI	AF217807.1
<i>Lentinula edodes</i>	Le.HYD2	AF217808.1
<i>Magnaporthe grisea</i>	MGG4	XM_364289.1
<i>Magnaporthe grisea</i>	MGG2	XM_001522792.1
<i>Magnaporthe grisea</i>	MHP1, MGG1	AF126872.1
<i>Magnaporthe grisea</i>	MPG1	L20685.2
<i>Metarhizium anisopliae</i>	SSGA	M85281.1
<i>Neurospora crassa</i>	NCU08192.1	AABX01000408.1
<i>Neurospora crassa</i>	EAS	AAB24462.1

<i>Ophiostoma ulmi</i>	CU	U00963 .1
<i>Paracoccidioides brasiliensis</i>	PbHYD2	AY427793 .1
<i>Paracoccidioides brasiliensis</i>	PbHYDI	AF526275.1
<i>Passalora fulva</i>	PF3	CAC27408. 1
<i>Passalora fulva</i>	PF1	CAC27407.1
<i>Passalora fulva</i>	PP2	CAB3931 2.1
<i>Pholiota narneko</i>	PNH2	AB0791 29.1
<i>Phoiiota narneko</i>	PNH1	AB0791 28.1
<i>Pisolithus linctorius</i>	HYDPt-1	U29605.1
<i>Pisolithus iinctorius</i>	HYDPt-2	U29606.1
<i>Pisolithus tinctorius</i>	HYDPt-3	AF09751 6.1
<i>Pleurotus ostreatus</i>	P0H2	Y14657. 1
<i>Pleurotus ostreatus</i>	P0H3	Y16881 .1
<i>Pleurotus ostreatus</i>	VMH3	AJ2381 48.1
<i>Pleurotus ostreatus</i>	P0H1	Y14656. 1
<i>Pleurotus ostreatus</i>	FBHI	AJ004883.1
<i>Schizophyllum commune</i>	SC4	M32330. 1
<i>Schizophyllum commune</i>	SC1 , 1G2	X00788. 1
<i>Schizophyllum commune</i>	SC6	AJ007504. 1
<i>Schizophyllum commune</i>	SC3	AAA96324.1
<i>Talaromyces thermophilus</i>	TT1	
<i>Trichoderma harzianum</i>	QID3	X71 9 13.1
<i>Trichoderma harzianum</i>	SRH1	Y11841 .1
<i>Trichoderma reesei</i>	HFB!!	P79073. 1
<i>Trichoderma reesei</i>	HFBI	P52754. 1
<i>Tricholoma terreum</i>	HYD1	AY048578. 1
<i>Verticillium dahliae</i>	VED	AAY891 0 1.1
<i>Xanthoria ectaneoides</i>	XEH1	AJ250793.1
<i>Xanthoria parietina</i>	XPH1	AJ250794. 1

Fusion Proteins

The definition of hydrophobin in the context of the present invention includes fusion
 5 proteins of a hydrophobin and another polypeptide as well as conjugates of
 hydrophobin and other molecules such as polysaccharides.

In one embodiment, the hydrophobin is a hydrophobin fusion protein. In this
 specification the term "fusion protein" means a hydrophobin sequence (as defined
 10 and exemplified above) bonded to a further peptide sequence (described herein as "a
 fusion partner") which does not occur naturally in a hydrophobin.

In one embodiment, the fusion partner may be bonded to the amino terminus of the
 hydrophobin core, thereby forming the group $(Y_i)_m$. In this embodiment, m may
 15 range from 1 to 2000, preferably 2 to 1000, more preferably 5 to 500, even more
 preferably 10 to 200, still more preferably 20 to 100.

in one embodiment, the fusion partner may be bonded to the carboxyl terminus of the hydrophobin core, thereby forming the group $(Y_2)_n$. In this embodiment, n may range from 1 to 2000, preferably 2 to 1000, more preferably 5 to 500, even more preferably 10 to 200, still more preferably 20 to 100.

In another embodiment, fusion partners may be bonded to both the amino and carboxyl termini of the hydrophobin core. In this embodiment, the fusion partners may be the same or different, and preferably have amino acid sequences having the number of amino acids defined above by the preferred values of m and n.

In one embodiment, the hydrophobin is not a fusion protein and m and n are 0.

Class I and II hydrophobins

In the art, hydrophobins are divided into Classes I and II. It is known in the art that hydrophobins of Classes I and II can be distinguished on a number of grounds, including solubility. As described herein, hydrophobins self-assemble at an interface (especially a water/air interface) into amphipathic interfacial films. The assembled amphipathic films of Class I hydrophobins are generally re-solubilised only in strong acids (typically those having a pK_a of lower than 4, such as formic acid or trifluoroacetic acid), whereas those of Class II are soluble in a wider range of solvents.

In one embodiment, the hydrophobin is a Class II hydrophobin. In another embodiment, the hydrophobin is a Class I hydrophobin.

In one embodiment, the term "Class II hydrophobin" means a hydrophobin (as defined and exemplified herein) having the above-described self-assembly property at a water/air interface, the assembled amphipathic films being capable of redissolving to a concentration of at least 0.1% (w/w) in an aqueous ethanol solution (60% v/v) at room temperature. In contrast, in this embodiment, the term "Class I hydrophobin" means a hydrophobin (as defined and exemplified herein) having the above-described self-assembly property but which does not have this specified redissolution property.

In another embodiment the term "Class *si* hydrophobin" means a hydrophobin (as defined and exemplified herein) having the above-described self-assembly property at a water/air interface and the assembled amphipathic films being capable of redissolving to a concentration of at least 0.1% (w/w) in an aqueous sodium dodecyl sulphate solution (2% w/w) at room temperature. In contrast, in this embodiment, the term "Class *ii* hydrophobin" means a hydrophobin (as defined and exemplified herein) having the above-described self-assembly property but which does not have this specified redissolution property.

Hydrophobins of Classes *i* and *ii* may also be distinguished by the hydrophobicity / hydrophilicity of a number of regions of the hydrophobin protein.

In one embodiment, the term "Class *ii* hydrophobin" means a hydrophobin (as defined and exemplified herein) having the above-described self-assembly property and in which the region between the residues B_3 and B_4 , *i.e.* the moiety $(X_3)_c$, is predominantly hydrophobic. In contrast, in this embodiment, the term "Class *i* hydrophobin" means a hydrophobin (as defined and exemplified herein) having the above-described self-assembly property but in which the region between the residues B_3 and B_4 , *i.e.* the group $(X_3)_c$, is predominantly hydrophilic.

In one embodiment, the term "Class *ii* hydrophobin" means a hydrophobin (as defined and exemplified herein) having the above-described self-assembly property and in which the region between the residues B_7 and B_8 , *i.e.* the moiety $(X_7)_g$, is predominantly hydrophobic. In contrast, in this embodiment, the term "Class *i* hydrophobin" means a hydrophobin (as defined and exemplified herein) having the above-described self-assembly property but in which the region between the residues B_7 and B_8 , *i.e.* the moiety $(X_7)_g$, is predominantly hydrophilic.

The relative hydrophobicity / hydrophilicity of the various regions of the hydrophobin protein can be established by comparing the hydropathy pattern of the hydrophobin using the method set out in Kyte and Doolittle, *J. Mol. Biol.*, 1982, 157, 105-132. According to the teaching of this reference, a computer program can be used to progressively evaluate the hydrophilicity and hydrophobicity of a protein along its amino acid sequence. For this purpose, the method uses a hydropathy scale (based on a number of experimental observations derived from the literature) comparing the hydrophilic and hydrophobic properties of each of the 20 amino acid side-chains. The program uses a moving-segment approach that continuously determines the


average hydrophathy within a segment of predetermined length as it advances through the sequence. The consecutive scores are plotted from the amino to the carboxy terminus. At the same time, a midpoint line is printed that corresponds to the grand average of the hydrophathy of the amino acid compositions found in most of the sequenced proteins. The method is further described for hydrophobins in Wessels, *Adv. Microbial Physiol.* 1997, 38, 1-45.

In one embodiment, the term "Class II hydrophobin" means a hydrophobin (as defined and exemplified herein) having the above-described self-assembly property and in which the region between the residues B₃ and B₄, *i.e.* the moiety (X₃)_c, is predominantly hydrophobic. In contrast, in this embodiment, the term "Class I hydrophobin" means a hydrophobin (as defined and exemplified herein) having the above-described self-assembly property but in which the region between the residues B₃ and B₄, *i.e.* the group (X₃)_c, is predominantly hydrophilic.

In one embodiment, the term "Class II hydrophobin" means a hydrophobin (as defined and exemplified herein) having the above-described self-assembly property and in which the region between the residues B₇ and B₈, *i.e.* the moiety (X₇)_g, is predominantly hydrophobic. In contrast, in this embodiment, the term "Class I hydrophobin" means a hydrophobin (as defined and exemplified herein) having the above-described self-assembly property but in which the region between the residues B₇ and B₈, *i.e.* the moiety (X₇)_g, is predominantly hydrophilic.

The relative hydrophobicity / hydrophilicity of the various regions of the hydrophobin protein can be established by comparing the hydrophathy pattern of the hydrophobin using the method set out in Kyte and Doolittle, *J. Mol. Biol.*, **1982**, 157, 105-132 and described for hydrophobins in Wessels, *Adv. Microbial Physiol.* **1997**, 38, 1-45.

Class II hydrophobins may also be characterised by their conserved sequences. In one embodiment, the Class II hydrophobins used in the present invention have the general formula (IV):



wherein:
m and n are independently 0 to 200;

B₁, B₂, B₃, B₄, B₅, B₆, B₇ and B₈ are each independently amino acids selected from Cys, Leu, Asa, Ser, Thr, Met or Gly, at least 6 of the residues B₁ through B₈ being Cys;

a is 6 to 12;

5 c is 8 to 16;

d is 2 to 20;

e is 4 to 12; and

g is 5 to 15.

10 in the formula (IV), a is preferably 7 to 11.

In the formula (IV), c is preferably 10 to 12, more preferably 11.


In the formula (IV), d is preferably 4 to 18, more preferably 4 to 16.

15

In the formula (IV), e is preferably 6 to 10, more preferably 9 or 10.

In the formula (IV), g is preferably 6 to 12, more preferably 7 to 10.

20 In one embodiment, the Class II hydrophobins used in the present invention have the general formula (V):



25 wherein:

m and n are independently 0 to 10;

B₁, B₂, B₃, B₄, B₅, B₆, B₇ and B₈ are each independently amino acids selected from Cys, Leu or Ser, at least 7 of the residues B₁ through B₈ being Cys;

a is 7 to 11;

30 c is 11;

d is 4 to 18;

e is 6 to 10; and

g is 7 to 10.

35 In the formulae (IV) and (V), at least 7, and preferably all 8 of the residues B₁ through B₈ are Cys.

In the formulae (IV) and (V), when 7 of the residues B₁ through B₈ are Cys, it is preferred that the residues B₃ through B₇ are Cys.

In the formulae (IV) and (V), when 7 of the residues B₁ through B₈ are Cys, it is preferred that: (a) B₁ and B₃ through B₈ are Cys and B₂ is other than Cys; (b) B₁ through B₇ are Cys and B₈ is other than Cys, or (c) B₁ is other than Cys and B₂ through B₈ are Cys. When 7 of the residues B₁ through B₈ are Cys, it is preferred that the other residue is Ser, Pro or Leu. In one embodiment, B₁ and B₃ through B₈ are Cys and B₂ is Ser. In another embodiment, B₁ through B₇ are Cys and B₈ is Leu. In a further embodiment, B₁ is Pro and B₂ through B₈ are Cys.

In the formulae (IV) and (V), preferably the group (X₃)_c comprises the sequence motif ZZXZ, wherein Z is an aliphatic amino acid; and X is any amino acid. In this specification the term "aliphatic amino acid" means an amino acid selected from the group consisting of glycine (G), alanine (A), leucine (L), isoleucine (I), valine (V) and proline (P).

More preferably, the group (X₃)_c comprises the sequence motif selected from the group consisting of LLXV, ILXV, 1LXL, VLXL and VLXV. Most preferably, the group (X₃)_c comprises the sequence motif VLXV.

In the formulae (IV) and (V), preferably the group (X₃)_c comprises the sequence motif ZZZZZXZ, wherein Z is an aliphatic amino acid; and X is any amino acid. More preferably, the group (X₃)_c comprises the sequence motif VLZVZXL, wherein Z is an aliphatic amino acid; and X is any amino acid.

In one embodiment, the hydrophobin is a polypeptide selected from SEQ ID NOs: 2, 4, 6, 8 or 10, or a polypeptide having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, or at least 99% sequence identity in the hydrophobin core to any thereof. By "the hydrophobin core" is meant the sequence beginning with the residue B₁ and terminating with the residue B₈.

In one embodiment, the hydrophobin is obtained or obtainable from fungi of the phylum *Ascomycota*. In one embodiment, the hydrophobin is obtained or obtainable from fungi of the genera *Cladosporium* (particularly *C. fulvum*), *Ophistoma* (particularly *O. ulmi*), *Cryphonectria* (particularly *C. parasitica*), *Trichoderma*

(particularly *T. harzianum*, *T. longibrichiatum*, *T. asperellum*, *T. Koningiopsis*, *T. aggressivum*, *T. stromaticum* or *T. reesei*), *Gibberella* (particularly *G. moniliformis*), *Neurospora* (particularly *N. crassa*), *Maganaporthe* (particularly *M. grisea*) or *Hypocrea* (particularly *H. jecorina*, *H. atroviridis*, *H. virens* or *H. lixii*).

5

In a preferred embodiment, the bicyrophofain is obtained or obtainable from fungi of the genus *Trichoderma* (particularly *T. harzianum*, *T. longibrichiatum*, *T. asperellum*, *T. Koningiopsis*, *T. aggressivum*, *T. stromaticum* or *T. reesei*). In a particularly preferred embodiment, the hydrophobin is obtained or obtainable from fungi of the species *T. reesei*.

10

In a more preferred embodiment, the hydrophobin is the protein selected from the group consisting of:

- (a) HFBII (SEQ ID NO: 2; obtainable from the fungus *Trichoderma reesei*);
- 15 (b) HFB; (SEQ ID NO: 4; obtainable from the fungus *Trichoderma reesei*);
- (c) SC3 (SEQ ID NO: 6; obtainable from the fungus *Schizophyllum commune*);
- (d) EAS (SEQ ID NO: 8; obtainable from the fungus *Neurospora crassa*); and
- (e) TT1 (SEQ ID NO: 10; obtainable from the fungus *Talaromyces thermophilus*); or a protein having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, or at least 99% sequence identity in the hydrophobin core to any thereof.

20

In a more preferred embodiment, the hydrophobin is the protein encoded by the polynucleotide selected from the group consisting of:

- 25 (a) HFBII (SEQ ID NO: 1; obtainable from the fungus *Trichoderma reesei*);
- (b) HFBi (SEQ ID NO: 3; obtainable from the fungus *Trichoderma reesei*);
- (c) SC3 (SEQ ID NO: 5; obtainable from the fungus *Schizophyllum commune*);
- (d) EAS (SEQ ID NO: 7; obtainable from the fungus *Neurospora crassa*); and
- (e) TT1 (SEQ ID NO: 9; obtainable from the fungus *Talaromyces thermophilus*);
- 30 or the protein encoded by a polynucleotide which is degenerate as a result of the genetic code to the polynucleotides defined in (a) to (e) above.

30

In an especially preferred embodiment, the hydrophobin is the protein "HFBII" (SEQ ID NO: 2; obtainable from *Trichoderma reesei*) or a protein having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, or at least 99% sequence identity in the hydrophobin core thereof.

35

In one embodiment, the hydrophobin may be present as an initial component of the composition. In another embodiment, the hydrophobin may be generated *in situ* in the composition (for example, by *in situ* hydrolysis of a hydrophobin fusion protein).

5

In an alternative embodiment, the hydrophobin may be replaced wholly or partially with a chaplin. Chaplins are hydrophobin-like proteins which are also capable of self-assembly at a hydrophobic-hydrophilic interface, and are therefore functional equivalents to hydrophobins. Chaplins have been identified in filamentous fungi and bacteria such as Actinomyces and Streptomyces. Unlike hydrophobins, they may have only two cysteine residues and may form only one disulphide bridge. Examples of chaplins are described in WO 01/74864, US 2010/0151525 and US 2010/0099844 and in Talbot, *Curr. Biol.* 2003, 13, R696-R698.

10

15 LIPOLYTIC ENZYME

In this specification the term 'lipolytic enzyme' is defined as an enzyme capable of acting on a lipid substrate to liberate a free fatty acid molecule. Preferably, the lipolytic enzyme is an enzyme capable of hydrolysing an ester bond in a lipid substrate (particularly although not exclusively a triglyceride, a glycolipid and/or a phospholipid) to liberate a free fatty acid molecule. Examples of possible lipid substrate are described below.

20

The lipolytic enzyme used in the present invention preferably has activity on both non-polar and polar lipids. The term "polar lipids" as used herein means phospholipids and/or glycolipids. Preferably, the term "polar lipids" as used herein means both phospholipids and glycolipids. Polar and non-polar lipids are discussed in Eliasson and Larsson, "Cereals in Breadmaking: A Molecular Colloidal Approach", publ. Marcel Dekker, 1993.

30

In particular, the lipolytic enzyme used in the present invention preferably has activity on the following classes of lipids: triglycerides; phospholipids, particularly but not exclusively phosphatidylcholine (PC) and/or N-acylphosphatidylethanolamine (APE); and glycolipids, particularly although not exclusively digalactosyl diglyceride (DGDG).

35

In this specification the term 'free fatty acid' means a compound of the formula R-C(=O)-OH wherein R is a straight- or branched chain, saturated or unsaturated,

hydrocarbonyl group, the compound having a total of 4 to 40 carbon atoms, preferably 6 to 40 carbon atoms, such as at least 10 to 40 carbon atoms, for example 12 to 40, such as 14 to 40, 16 to 40, 18 to 40, 20 to 40 or 22 to 40 carbon atoms, more preferably 10 to 24, especially 12 to 22, particularly 14 to 18, for example 16 or 18 carbon atoms. In one particular embodiment, such an acyl group is an alkanoyl group. Alternatively, such an acyl group comprises an alkenoyl group, which may have, for example, 1 to 5 double bonds, preferably 1, 2 or 3 double bonds.

Suitably, the lipolytic enzyme for use in the present invention may have one or more of the following activities selected from the group consisting of: phospholipase activity (such as phospholipase A1 activity (E.C. 3.1.1.32) or phospholipase A2 activity (E.C. 3.1.1.4); glycolipase activity (E.C. 3.1.1.26), triacylglycerol hydrolysing activity (E.C. 3.1.1.3), lipid acyltransferase activity (generally classified as E.C. 2.3.1.x in accordance with the Enzyme Nomenclature Recommendations (1992) of the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology), and any combination thereof. Such lipolytic enzymes are well known within the art.

Suitably, the lipolytic enzyme for use in the present invention may be a phospholipase (such as a phospholipase A1 (E.C. 3.1.1.32) or phospholipase A2 (E.C. 3.1.1.4); glycolipase or galactolipase (E.C. 3.1.1.26), triacylglyceride lipase (E.C. 3.1.1.3). Such enzyme may exhibit additional side activities such as lipid acyltransferase side activity.

Preferably, the lipolytic enzyme for use in the present invention has triacylglycerol hydrolysing activity (E.C. 3.1.1.3).

A lipolytic enzyme may be categorised as belonging to one of three classes (GX, GGGX or Y) based on structure and sequence analysis of the oxyanion hole of the enzyme.

A "GX lipolytic enzyme" is one where the oxyanion hole-forming residue X of the enzyme is structurally well conserved and is preceded by a strictly conserved glycine.

A "GGGX enzyme" is one where there is a well conserved GGG pattern, followed by a conserved hydrophobic amino acid X and the backbone amide of glycine preceding the residue X forms the oxyanion hole.

A "Y lipolytic enzyme" in one in which the oxyanion hole is not formed by a backbone amide but by the hydroxy! group of a tyrosine side chain.

5 In one aspect, the present invention relates to the use of a GX lipolytic enzyme.

Suitably, the oxyanion hole forming residue X may be M, Q, F, S, T, A, L or 1.

Preferably, the oxyanion hole forming residue X may be M, Q, F, S or T.

10 in one embodiment, the lipolytic enzyme may belong to one of the following alpha/beta hydrolase superfamilies abH23 (preferably abH23.01), abH25 (preferably 25.01), abH16 (preferably 16.01), abH18 (preferably abH18.01) and abH15 (preferably 15.01 or 15.02).

15 In one embodiment, the lipolytic enzyme may belong to one of the following alpha/beta hydrolase superfamilies abH23 (preferably abH23.01), abH25 (preferably 25.01), abH16 (preferably 16.01) and abH15 (preferably 15.02).

In one embodiment, preferably the lipolytic enzyme is classified as a member of the
20 abH23 superfamily, preferably as a member of the abH23.01 homologous family in the Lipase Engineering Database.

Details regarding these superfamilies may be found on the Lipase Engineering Database (<http://www.led.uni-stuttgart.de/>). When referring to the Lipase Engineering
25 database herein reference is made to version 3.0 of the database released on 10 December 2009.

In particular, in one embodiment a lipolytic enzyme may be considered to belong to the abH23 superfamily if it is a GX lipolytic enzyme from a filamentous fungus.
30 Preferably, a lipolytic enzyme is a GX lipolytic enzyme if the catalytic triad of the enzyme aligns with that of a lipase from *Rhizopus miehei*, such as swissprot P19515.

Examples of lipolytic enzymes belonging to the abH23 superfamily include those indicated in Table 2.

35

Table 2

abH23	Organism	NCBI accession code and version number* OR gi number
abH23.01 <i>(Rhizomucor miehei</i> lipase like)	<i>Arabidopsis thaliana</i>	NP_197365.1 AAL24204.1 42570528 145362642
	<i>Aspergillus awamori</i>	BAA92937.3 84028205
	<i>Aspergillus clavatus</i>	121719262
	<i>Aspergillus flavus</i>	27525628
	<i>Aspergillus fumigatus</i>	70985264
		70987066
	<i>Aspergillus nidulans</i>	67902118
		67537354
	<i>Aspergillus niger</i>	AAK60631.1
		042807.1
		1UWC_A
		2HL6_A
		1USW_A
		2BJH_A
		145252728
		110431975
		145241772
		109677003
		145251976
	110431973	
	<i>Aspergillus oryzae</i>	83766610
		169771817
		169768448
		169780130
		169774351
		BAA1291.2.1
	<i>Aspergillus parasiticus</i>	27525626
	<i>Aspergillus iamarii</i>	124108031

<i>Aspergillus terreus</i>	115402833
	115385463
	115400761
	115443274
<i>Aspergillus tubingensis</i>	04281 5.1
<i>Brugia malayi</i>	17059251 1
<i>Caenorhabditis briggsae</i>	157761 233
	157761 241
	157755883
	157771 698
	1577631 72
	157747253
	1577591 79
	1577591 77
	157772997
	1577731 05
	157773031
	15777461 3
	15777461 7
	157772605
	15777461 9
	157774601
<i>Caenorhabditis elegans</i>	115534096
	17552584
	71983228
	71983230
	71983236
	193207843
	115534067
	15851 8 185
	865751 43
	115534303
	72000668
	AAF60431 .2
	71994497
	T27056
	71994547

	CAB61 137.3
	193247829
<i>Chaetomium globosum</i>	116206442
<i>Cyanobium</i> sp.	19762731 0
<i>Cyanothece</i> sp.	172037675
	17766391 5
	196246404
<i>Dictyostelium discoideum</i>	60463496
	66825791
	AAM43784.1
<i>Dictyostelium discoideum</i> AX4	66802624
<i>Fusarium oxysporum</i>	143791 375
<i>Gibberella zeae</i>	33621 223
	461 23057
<i>Magnaporthe grisea</i>	39978263
<i>Nectria haematococca</i>	CAC1 9602.1
<i>Neosartorya fischeri</i>	1194991 43
	119480389
<i>Neurospora crassa</i>	CAC28687.1
<i>Neurospora crassa</i> QR74A	EAA32 130.1
<i>Oryza sativa</i>	115463525
	125552085
	125577937
	115486491
	115473965
	125586239
	125543854
	1255351 66
	125559538
	115442095
	115453007
	BAB64204.1
	125529023
<i>Penicillium allii</i>	3 1872092
<i>Penicillium camemberti</i>	P25234
	1Ti A
	1TIA_A

<i>Penicillium cyclopium</i>	48429006
	AAF82375.1
<i>Penicillium expansum</i>	AAG22769.1
<i>Phaeosphaeria nodorum</i>	169595748
	169606904
<i>Physcomitrella patens</i>	168020609
	168040480
	168037728
<i>Podospora anserina</i>	17 1693635
<i>Populus frichocarpa</i>	118482274
<i>Pyrenophora tntici-repentis</i>	1891 9251 6
	189202058
<i>Rhizomucor miehei</i>	P 1951 5.2
	3TGL
	5TGL
	4TGL TM
	1TGL
	5TGL_A
	4TGL_A
	1TGL_A
	3TGL_A
<i>Rhizopus arrhizus</i>	1TIC_A
	AAF32408.1
	1TIC_B
<i>Rhizopus javanicus</i>	73621 144
<i>Rhizopus microsporus</i>	156470335
	166078592
<i>Rhizopus niveus</i>	P21 8 11
	1LGY_A
	BAA31 548.1
	1LGY_B
<i>Rhizopus oryzae</i>	1LGY_C
	AAS84458.1
	P61 872. 1
	1TIC_A
	94962082
7 13901 09	
<i>Rhizopus stolonifer</i>	AAZ66864.1

<i>Synechococcus</i> sp.	87301 494
<i>Thermomyces lanuginosus</i>	059952.1
	1Tt B
	1DTE_A
	1DT5_D
	1DU4_B
	1DT3_A
	1EIN_B
	1DT3_B
	1DT5_E
	1DT5_B
	1DT5_G
	1DT5_F
	1DT5_H
	1DT5_A
	1DT5_C
	1DTE_B
	1DU4_A
	1DU4_D
	1DU4_C
	1EIN_C
1EIN_A	
1GT6_A	
<i>Triticum aestivum</i>	CAD32696.1
	CAD32695.1
<i>Vitis vinifera</i>	157336329
<i>Zea mays</i>	194691 896
	194690642
	194706432
	194694588
	19469421 0

In this embodiment, preferably the oxyanion hole forming residue is a serine or threonine.

- 5 Preferably, the lipolytic enzyme belongs to the *Rhizopus miehei* like homologous family abH23.01 . Suitably, particularly preferred enzymes for use in the present invention may include any lipolytic enzymes classified in homologous family

abH23.01 from *Thermomyces* (preferably, *T. lanuginosus*), *Fusarium* (preferably *F. heterosporum*), *Aspergillus* (preferably *A. tubiengisis* and/or *A. fumigatus*) and *Rhizopus* (preferably, *R. arrizus*), preferably from *Thermomyces* (preferably, *T. lanuginosus*), *Fusarium* (preferably *F. heterosporum*), or *Aspergillus* (preferably *A. tubiengisis*). Examples of such lipolytic enzymes include LIPEX™ (a *Thermomyces lanuginosus* lipolytic enzyme disclosed in WO 94/026 17 and shown herein as SEQ ID NO: 11, the *Fusarium heterosporum* lipolytic enzyme disclosed in WO 2005/08791 8 and shown herein as SEQ ID NO: 13 (available from Danisco A/S as Grindamyl POWERBAKE 4100™) and Lipase 3 (an *Aspergillus tubigensis* lipolytic enzyme disclosed in WO 98/45453 and shown herein as SEQ ID NO: 14).

In one embodiment of the present invention, a lipolytic enzyme may be considered to belong to the abH25 superfamily if the catalytic triad aligns with that of the *Moraxella* lipase 1 like lipolytic enzyme as shown in the swissprot protein knowledge base (<http://www.expasy.org/sprot/> and <http://www.ebi.ac.uk/swissprot/>) under accession number P19833 - version of 26 July 2005.

Examples of lipolytic enzymes belonging to this family include those listed in Table 3.

20 Table 3

abH25	Organism	NCBi accession code and version number* OR gi number
abH25.01 (<i>Moraxella</i> lipase 1 like)	<i>Acidovorax delafieldii</i>	BAB86909.1
	<i>Kineococcus radiotolerans</i>	152967773
	<i>Kineococcus radiotolerans</i> SRS3021 6	EAM75386.1
	<i>Moraxella</i> sp.	P19833.1
	<i>Streptomyces albus</i>	AAA53485.1
	<i>Streptomyces ambofaciens</i>	117164910
	<i>Streptomyces coelicolor</i>	AAD09315.1
		CAB69685.1
	<i>Streptomyces exfoliatus</i>	1JFR_B
		1JFR_A
	<i>Streptomyces griseus</i>	182439251
	<i>Thermobifida fusca</i>	72161287
		72161286

	<i>Thermobifida fusca</i> DSM 43793	CAH1 7553.1
		CAH 17554.1

In this embodiment, preferably the oxyanion hole forming residue is M, Q, A, F, L or I.

in one embodiment of the present invention, a lipolytic enzyme may be considered to belong to the abH16 superfamily if the catalytic triad aligns with that of *Streptomyces*.

Examples of lipolytic enzymes belonging to this family include those indicated in Table 4.

10 Table 4

abH16	Organism	NCBI accession code and version number* OR gi number	
	<i>Arthrobacter chlorophenolicus</i>	169176591	
	<i>Arthrobacter</i> sp. FB24	116669612	
	<i>Corynebacterium diphtheriae</i>	38232746	
	<i>Corynebacterium efficiens</i>		25026650
			25026649
	<i>Corynebacterium efficiens</i> YS-314		BAC16904.1
			BAC16903.1
	<i>Corynebacterium glutamicum</i>		19551331
			145294142
			19551330
			145294141
	<i>Frankia</i> sp.	158312565	
	<i>Frankia</i> sp. EAN1pec	EAN12331.1	
	<i>Nocardia farcinica</i>	54025580	
	<i>Nocardioides</i> sp.	119715399	
	<i>Nocardioides</i> sp. JS614	EAO07564.1	
<i>Propionibacterium acnes</i>		50843543	
		50843256	
	<i>Propionibacterium acnes</i> P-37	CAA67627.1	
abH16.01 (<i>Streptomyces</i> lipases)	<i>Rhodococcus</i> sp.	111021394	
		111024112	
		111025204	

		111025876
		111022422
		111024917
		40787231
	<i>Rubrobacter xylanophilus</i>	108805093
	<i>Rubrobacter xylanophilus</i> DSM 9941	EAN36909.1
	<i>Streptomyces avermitilis</i>	29833101
	<i>Streptomyces avermitilis</i> MA-4680	BAC74270.1
	<i>Streptomyces cinnamoneus</i>	AAB71210.1
	<i>Streptomyces coelicolor</i>	NP606008
	<i>Streptomyces fradiae</i>	148832709
	<i>Streptomyces griseus</i>	182439565
	<i>Streptomyces pristinaespiralis</i>	YP002199726
	<i>Streptomyces</i> sp.	197333608
	<i>Streptomyces sviceps</i>	197781872
	<i>Synthetic construct</i>	AA092397.1

In this embodiment, preferably the oxyanion hole forming residue is T or Q.

In one embodiment of the present invention, a lipolytic enzyme may be considered to belong to the abH15 superfamily if the catalytic triad aligns with that of a GX *Burkholderia* lipase.

Examples of lipolytic enzymes belonging to this family include those indicated in Table 5 and LIPOMAX as shown herein as SEQ ID NO: 15.

10

Table 5

abH1 5	Organism	NCBI accession code and version number* OR gi number
abH1 5.02 (<i>Burkholderia cepacia</i> lipase like)	<i>Acidovorax avenae</i>	12061 2825
	<i>Acinetobacter baumannii</i>	16979451 5
		1266431 75
		193078538

	15851 7002
<i>Acinetobacter calcoaceticus</i>	AAD29441 .1
<i>Acinetobacter schindleri</i>	1581 20326
	1581 20327
<i>Acinetobacter</i> sp.	50086294
<i>Acinetobacter</i> sp. SY-01	AAP44577. 1
<i>Aeromonas hydrophila</i>	1176 18653
<i>Aeromonas salmonicida</i>	145300587
<i>Alcanivorax borkumensis</i>	110834836
<i>Alcanivorax</i> sp.	1961 94968
	1961 931 33
<i>Alteromonas macleodii</i>	88795738
<i>Azotobacter vinelandii</i> AvOP	EAM0521 4. 1
<i>Burkholderia ambifaria</i>	115358044
	118695660
	17 13 16092
	170702796
	17 1320247
<i>Burkholderia cenocepacia</i>	124875244
	107026795
	11871 3500
	84354072
	198038844
<i>Burkholderia cenocepacia</i> AU 1054	190607421
<i>Burkholderia cenocepacia</i> AU 1054	EAM08623. 1
<i>Burkholderia cenocepacia</i> HI2424	EAM 18550. 1
<i>Burkholderia cepacia</i>	AAY86757.2
	1167391 50
	16 1406799
	10IL_B
	1HQD_A
	4LIPJD
	P22088.2
	10IL_A
	4LIP_E
	1YS2_X
<i>Burkholderia cepacia</i> KCTC 2966	AAT85572. 1
<i>Burkholderia cepacia</i> R1808	4631 9469
	4631 9468

<i>Burkholderia cepacia</i> R 18 194	4631 2540
<i>Burkholderia cepacia</i> ST-200	BAD1 3379.1
<i>Burkholderia dolosa</i>	8436031 3
	1TAH_A
	1TAH_C
	1TAH_B
	1TAHJD
	1QGE_E
<i>Burkholderia glumae</i>	2ES4_A
	8361 8505
	53715898
	8361 8339
	167003692
<i>Burkholderia mallei</i>	67636935
<i>Burkholderia mallei</i> 10399	67635666
<i>Burkholderia mallei</i> FMH	69987887
<i>Burkholderia mallei</i> GB8 horse 4	67640408
	67642620
<i>Burkholderia mallei</i> JHU	70001349
<i>Burkholderia mallei</i> NCTC 10247	67645935
<i>Burkholderia multivorans</i>	16 1521 2 10
	16 15251 17
<i>Burkholderia multivorans</i> RG2	AAW30 196. 1
<i>Burkholderia multivorans</i> Uwc 10	AAZ39650.1
<i>Burkholderia oklahomensis</i>	167573565
	167568063
	167567050
	1675741 27
	53722762
	126445060
	9991 1132
	1001 26424
	16791 581 5
	126442397
	157806477
	134281 779
	7681 8459
	100231 475
	9990851 5
<i>Burkholderia pseudomallei</i>	

	100059930
	53723336
	1001 2 1879
	167744369
	184212969
	167908322
	167725450
	67671 904
<i>Burkholderia pseudomallei</i> 1655	67670022
	67684997
<i>Burkholderia pseudomallei</i> 1710a	67681 352
<i>Burkholderia pseudomallei</i> 668	677351 59
	67755633
<i>Burkholderia pseudomallei</i> Pasteur	67753658
<i>Burkholderia pseudomallei</i> S13	67759470
<i>Burkholderia</i> sp. 383	78063020
<i>Burkholderia</i> sp. HY-10	154091354
<i>Burkholderia</i> sp. 99-2-1	AAV34204. 1
<i>Burkholderia</i> sp. MC16-3	AAV34203.1
	8371 7248
	167577201
	8371 6483
	167579206
	16761 7325
<i>Burkholderia thailandensis</i>	167840423
<i>Burkholderia ubonensis</i>	167583926
	134293086
<i>Burkholderia vietnamiensis</i>	134293087
	EAM26790. 1
	67548784
<i>Burkholderia vietnamiensis</i> G4	EAM26789. 1
<i>Chromobacterium violaceum</i>	344981 69
<i>Chromobacterium violaceum</i> ATCC 12472	AAQ60384. 1
<i>Burkholderia glumae</i>	1CVL_A
<i>Cupriavidus taiwanensis</i>	194289366
<i>Dehalococcoides</i> sp.	16381 3742
	1982621 10
<i>Gamma proteobacterium</i>	1982621 37

<i>Hahella chejuensis</i>	83646958
<i>Listonella anguillarum</i>	19731 3280
<i>Listonella anguillarum</i> M93Sm	AA Y261 46.2
	1493761 15
<i>Marinobacter algicola</i>	149378244
<i>Marinomonas</i> sp.	871 19903
	149908369
	14991 1484
<i>Moritella</i> sp.	149909327
<i>Myxococcus xanthus</i>	108756922
	945001 83
<i>Oceanobacter</i> sp.	94501 726
	90409701
<i>Photobactenum profundum</i>	5430361 2
<i>Photobacterium profundum</i> ss9	CAG23805.1
<i>Photobactenum</i> sp.	89072072
<i>Plesiocystis pacifica</i>	149921 436
<i>Proteus mirabilis</i>	197284877
<i>Proteus</i> sp.	1841 91073
<i>Proteus vulgaris</i>	AAB01 071 .1
	AAC34733. 1
	P26876.2
	BAA091 35.1
	AAF641 56.1
	BAA231 28.1
	1EX9_A
	1071 0241 1
	152989672
<i>Pseudomonas aeruginosa</i>	152983830
<i>Pseudomonas aeruginosa</i> KCTC 1637	AAT85570.1
<i>Pseudomonas entomophila</i>	104783837
	77456799
	77459293
	AAC1 5585. 1
<i>Pseudomonas fluorescens</i>	707341 19
	23058245
<i>Pseudomonas fluorescens</i> PfO-1	23061 908
<i>Pseudomonas fragi</i>	CAC071 91.1

	P08658.2
	AAA25879. 1
<i>Pseudomonas luteola</i>	AAC0551 0.1
	146307587
	146306794
<i>Pseudomonas mendocina</i>	AAM1 4701 .1
	167035900
	119858840
	170723807
	26991534
<i>Pseudomonas putida</i>	148549934
<i>Pseudomonas putida</i> KT2440	AAN 70423. 1
	4LIP_E
	1891 7871 1
<i>Pseudomonas</i> sp.	18917871 3
<i>Pseudomonas</i> sp. 109	P26877. 1
<i>Pseudomonas</i> sp. KFCC1 081 8	AAD22078. 1
<i>Pseudomonas</i> sp. KWI-56	P25275. 1
<i>Pseudomonas</i> sp. SW-3	AAG47649.2
<i>Pseudomonas stutzeri</i>	146282376
<i>Pseudomonas wisconsinensis</i>	AAB53647.1
<i>Psychrobacter cryohalolentis</i>	93005273
<i>Psychrobacter cryohalolentis</i> K5	EA0 10600.1
<i>Psychrobacter</i> sp.	148652775
<i>Ralstonia eutropha</i>	113867341
<i>Ralstonia metallidurans</i>	22979988
	153885935
<i>Ralstonia pickettii</i>	12 1531 370
<i>Ralstonia</i> sp. M1	AAR1 3272. 1
<i>Rhodoferax ferrireducens</i>	899021 27
<i>Shewanella denitrificans</i>	9 1792458
<i>Shewanella denitrificans</i> OS-21 7	69944965
<i>Shewanella denitrificans</i> OS21 7	EAN69301 .1
<i>Shewanella frigidimarina</i>	114564999
<i>Shewanella frigidimarina</i> NCIMB 400	EAN741 11.1
<i>Shewanella woodyi</i>	118073371
<i>Sorangium cellulosum</i>	162451 743
	AAT51 282.1
<i>Synthetic construct</i>	AAT51 165. 1

<i>Vibrio alginolyticus</i>	9 1225988
<i>Vibrio angustum</i>	90580697
<i>Vibrio campbellii</i>	163801 15 1
	P 15493.2
	AAA1 7487. 1
	150423294
	11621 9797
	153801 593
	15321 5 150
<i>Vibrio cholerae</i>	11621 4571
<i>Vibrio cholerae</i> M0 10	75830993
<i>Vibrio cholerae</i> RC385	75821 182
<i>Vibrio cholerae</i> V51	75819240
<i>Vibrio cholerae</i> V52	7581 6524
	156974975
<i>Vibrio harveyi</i>	1538341 78
	28897955
<i>Vibrio parahaemolyticus</i>	153837472
<i>Vibrio shilonii</i>	149187907
	116 184955
<i>Vibrio</i> sp.	861 44587
<i>Vibrio</i> sp. Ex25	75855688
<i>Vibrio splendidus</i>	84385385
	376801 74
<i>Vibrio vulnificus</i>	27365668
<i>Vibrio vulnificus</i> CKM-1	AAQ04476. 1
<i>Vibrio vulnificus</i> CMCP6	AA01 0723. 1
<i>Vibrionales bacterium</i>	148974047
	22996002
<i>Xylella fastidiosa</i>	281 98381
<i>Xylella fastidiosa</i> Ann-1	EA03 1309.1
<i>Xylella fastidiosa</i> Temeculal	AA028344.1
<i>Yersinia enterocolitica</i>	1234421 25
<i>Yersinia mollaretii</i> ATCC 43969	77961 583

abH15	Organism	NCBI accession coda and version number* OR gi number
abH1 5.01 (<i>Staphylococcus aureus</i> lipase like)	<i>Ailuropoda melano!euca</i>	6251 1068
	<i>Alouaita seniculus</i>	583391 72
		533391 74
		583391 76
		583391 78
		583391 80
	<i>Arabidopsis thaliana</i>	AAF1 7667. 1
		AAF8701 2.1
		D86367
		26451 003
		AAD31 339. 1
	<i>Ateles geoffroyi</i>	42571 431
	<i>Ateles geoffroyi</i>	1846251 2
		1846251 4
	<i>Bacillus anthracis</i>	30262592
	<i>Bacillus anthracis Ames</i>	AAP26455. 1
	<i>Bacillus cereus</i>	521 42888
		42781 684
		1681 39359
		1681 341 90
		167938472
1681 58861		
166993225		
19604361 8		
196040277		

	<i>Bacillus cereus</i> G924 1	EAL 12983. 1
	<i>Bacillus</i> sp. 42	AAV351 02. 1
	<i>Bacillus</i> sp. L2	AAW47928.1
	<i>Bacillus</i> sp. TP10A. 1	AAF63229.1
	<i>Bacillus</i> sp. Tosh	AAW21775. 1
		757641 33
		49477789
	<i>Bacillus thuringiensis</i>	118477999
	<i>Bacillus thuringiensis</i> A TCC 35646	EA051 633.1
	<i>Bacillus welhenstephanensis</i>	163940476
	<i>Balaenoptera borealis</i>	081 2 180A
		55583872
	<i>Balaenoptera physalus</i>	1104245A
		164597876
	<i>Bos frontalis</i>	116256079
		6251 1051
	<i>Bos grunniens</i>	119675392
		270861 1
		6063098
	<i>Bos indicus</i>	164597854
		8341 6245
		8341 6247
		30794288
		134244277
		164597862
		8341 6249
		59797396
	<i>Bos taurus</i>	12663221 3

	6063096
	83416241
	60651 145
	13431390
<i>Bubalus bubalis</i>	296143
	58339182
	58339184
<i>Callicebus moloch</i>	58339183
	17368913
	21449837
<i>Callithrix jacchus</i>	21449839
	6251 1040
<i>Camelus dromedarius</i>	126567081
	312196
<i>Canis lupus</i>	50978904
	190683030
	83416243
	155183991
	6063094
	15 10 157A
	60687495
<i>Capra hircus</i>	126632219
	6251 1092
<i>Cavia porcellus</i>	7677454
<i>Cebus albifrons</i>	1 16634246
	3024641
<i>Cervus elaphus</i>	70909960
<i>Cloning vector</i>	12584848

		153941 353
		168178255
		187932762
		1681 79769
		153940345
		1681 85824
		170759344
		188588446
		1681 86291
		170756926
		14838001 8
		170758348
		1681 83734
		188590654
		187935767
		188587698
		148378855
		1681 84078
	<i>Clostridium botulinum</i>	170757848
		118443364
	<i>Clostridium novyi</i>	11844321 1
		187777968
	<i>Clostridium sporogenes</i>	187779336
	<i>Clostridium tetani</i>	2821 0658
	<i>Clostridium tetani Massachusetts</i>	AA035539.1
	<i>Deinococcus radiodurans</i>	C75533
	<i>Delphinus delphis</i>	6251 1070
	<i>Elephantidae gen.</i>	1509285A

	126352373
	170931 OA
	156723467
	56786671
	168693409
	197941 001
	11 1606634
<i>Equus caballus</i>	11 1606636
	571 63879
<i>Felis catus</i>	567042
<i>Galago senegalensis</i>	17368901
<i>Geobacillus kaustophilus</i>	56420521
	67906830
<i>Geobacillus sp. (Strain T1)</i>	JC8061
<i>Geobacillus sp. T1</i>	AAO92067.2
	AAF4021 7.1
	1JI3_B
	1JI3_A
	AAL28099. 1
	117373028
	JW0068
	1KU0_A
	1KU0_B
<i>Geobacillus stearothermophilus</i>	AAX1 1388.1
<i>Geobacillus thermocatenulatus</i>	CAA64621 .1
	AAD30278. 1
	113431 924
<i>Geobacillus thermoleovorans</i>	AAM21 774. 1

	83939852
<i>Geobacillus thermoleovorans IHI-91</i>	AAN7241 7.1
	1102651 50
	2DSN.._A
<i>Geobacillus zalihae</i>	2Z5G_A
<i>Giraffa camelopardalis</i>	6251 1039
<i>Hippopotamus amphibius</i>	6251 1038
	1AXI_A
	1HGU_A
	1KF9_A
	71 1074A
	10334861
	4503083
	1Z7C_A
	34784701
	18 1 127
	731 144A
	36544
	12545376
	12545381
	1302781 2
	1HWG_A
	11961 4650
	471 2 1568
	3HHR_A
	471 2 1579
	1HWH_A
<i>Homo sapiens</i>	1403262B

		3 1905
		11961 4648
		1302781 4
		1403262A
		1302781 6
		4503991
		49456759
		49456803
		1831 77
		11961 4662
		13027822
		11961 4661
		11961 4666
	<i>Lactobacillus casei CL96</i>	AAP02960.1
		110338953
	<i>Lama pacos</i>	58601 0
	<i>Loxodonta africana</i>	134706
		538541 58
		541 24352
		538541 63
	<i>Macaca assamensis</i>	538541 65
		112293303
		2931 11
		112293293
		681 36596
		1 14052777
		11405271 7
	<i>Macaca mulatta</i>	114052929

	1 12293289
	1 12293299
	681 36594
	2500855
	1091 16855
	1091 49084
	1091 48991
<i>Mesocricetus auratus</i>	58601 2
<i>Monodelphis domestica</i>	741 36533
	6679997
<i>Mus musculus</i>	4096656
<i>Nannospalax ehrenbergi</i>	62510957
	134709
	4684921 5
<i>Neovison vison</i>	164254
	538541 3 1
	53854129
	53854133
	53854135
	53854137
<i>Nomascus leucogenys</i>	53854139
<i>Nycticebus pygmaeus</i>	17368910
<i>Oryctolagus cuniculus</i>	1174399
	115463847
<i>Oryza sativa</i>	125552313
	94183527
	94406690
<i>Ovis aries</i>	94183483

		941 8351 9
		155001 235
		941 83467
		1666694
		941 83402
		941 83398
		941 83424
		126632207
		941 83444
		18051 46A
		941 83426
		941 83523
		10051 82A
		941 83400
		941 8351 1
		941 8341 0
		12663221 1
		941 83452
		165887
		1167351 58
		941 83438
		57527824
		941 83495
		941 83507
		941 8351 5
		941 83475
		126632209
		941 83420

		94183432
		83955026
		94183430
	<i>Paenibacillus larvae</i>	167465325
		20140016
		20140015
		114669972
		114669970
		114669980
		114669998
		114669984
		114669978
		114669976
		114669996
		114669982
		114670000
		114669918
		114669948
		114669944
		114669938
		571 13881
		114669920
		114669930
		114669994
		114669992
		114669990
		114670016
<i>Pan troglodytes</i>	114670014	

	55645705
	114669905
	114669936
	571 13891
	114669942
	114669934
	114669940
	571 13885
	281 88745
	11466991 5
	114669922
	114669932
	114670004
<i>Physcomitrella patens</i>	162691 248
	583391 90
	583391 92
<i>Pithecia pithecia</i>	583391 95
	538541 4 1
	541 24350
	538541 46
<i>Pygathrix nemaus</i>	538541 48
	13471 7
	77861 9 10
	149054569
<i>Rattus norvegicus</i>	149054567
	538541 50
	538541 52
<i>Rhinopithecus roxellana</i>	538541 54

		53854156
	<i>Saimiri boliviensis</i>	17368174
	<i>Shuttle vector</i>	2342750
		153104
		88193885
		1314205A
		49482354
		57652458
		83682315
		120864890
		83682355
		586027
		83682335
		15923101
		154736704
		83682395
		83682375
		83682371
		120864986
		120865151
		83682327
		120865143
		120864794
		120865004
		120864887
		120865236
		46695
	<i>Staphylococcus aureus</i>	82750020

		154736702
		120865077
		83682365
		83682377
		120865094
		120865232
		83682345
		1208651 40
		83682333
		83682369
		83682331
		83682339
		120865030
		120864975
		1208651 0 1
		120865021
		8368231 1
		15 1220267
		148266538
		133853458
		83682383
		1891 69989
		16 1508379
		120864978
		1905280A
		83682307
		2 1281 8 13
		83682309

		83682363
		83682397
		120864800
		1208651 83
		120864824
		154736696
		83682379
		120864797
		120864834
		83682337
		120865080
		83682389
		154736698
		154736692
		1208651 23
		83682385
		83682359
		83682351
		BAB96455.1
		BAB43769. 1
		S68970
		AAD52059. 1
		P65289.2
		57651 062
		8402821 8
		P 10335. 1
		AAK291 27.1
		B89797

	871 621 30
	2 1232026
	57651 244
	148266743
	158347635
	49484866
	84029334
	49482552
	1480567
	82752249
<i>Staphylococcus aureus</i> MW2	Q8NYC2.1
<i>Staphylococcus aureus</i> Mu50	Q99QX0
	643453
<i>Staphylococcus carnosus</i>	643451
	274671 03
	193888386
	Q0251 0
	82654954
	AAC38597. 1
	AAC67547. 1
	57865775
	57865971
	27469321
	274671 63
<i>Staphylococcus epidermidis</i>	57865673
<i>Staphylococcus epidermidis</i> 9	AAA1 9729.1
	AAO06046.1
	AAO03782.1
<i>Staphylococcus epidermidis</i> A TCC 12228	AAO03878. 1
	AAO03842. 1
	707251 69
<i>Staphylococcus haemolyticus</i>	AAF21 294. 1
	2HIHA_A
<i>Staphylococcus hyicus</i>	P04635. 1

	AAT34964. 1
	73663604
<i>Staphylococcus saprophyticus</i>	73661 8 1 1
<i>Staphylococcus simulans</i>	CAC83747.1
	AAG35723.1
	BAD90561 .1
	BAD90565. 1
<i>Staphylococcus warneri</i>	BAD90562. 1
	551 988
	551 987
	AAG35726.1
<i>Staphylococcus xylosus</i>	52854061
	124268
<i>Streptococcus sp.</i>	47072
	46361 729
	164478
	166835929
	5723331 1
	16081 12A
	13 12298A
	5723331 3
	57233321
	475231 20
<i>Sus scrofa</i>	9 12486
	33341 802
	6671 284
	14582904
	6081 0 1 19
	6 1364449
	6082741 2
	6081 5489
	305841 4 1
	60655785
<i>Synthetic construct</i>	6671 282
	12964200
<i>Tragulus javanicus</i>	129641 98
<i>Trichosurus vulpecula</i>	391 5004

	<i>Uncultured bacterium</i>	145965989
	<i>Uncultured bacterium</i>	145965991
	<i>Vitis vinifera</i>	15732981 9
		158346762
		16634381 4
	<i>Vulpes lagopus</i>	JS0429
	<i>Vuipes vulpes</i>	134722

Throughout the specification examples of enzymes falling into a particular superfamily and/or homologous family in accordance with the Lipase Engineering Database version 3.0 are provided. In one embodiment of the present invention, the lipolytic enzyme of the present invention may be selected from any one or more of the lipolytic enzymes in these exemplified groups.

In another embodiment, the lipolytic enzyme for use in the present invention may be from one or more of the following genera: *Thermomyces* (preferably *T. lanuginosus*), *Thermobifida* (preferably, *T. fusea*), *Pseudomonas* (preferably *P. alcaligenes*) and *Streptomyces* (preferably *S. pristinaespiralis*).

Suitably, the lipolytic enzyme may comprise one of more of the following amino acid sequences:

- a) SEQ ID NO: 11;
- b) SEQ ID NO: 15;
- c) SEQ ID NO: 16;
- d) SEQ ID MO: 17;
- e) an amino acid sequence having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to any one of the amino acid sequences defined in a) to d); or
- f) an amino acid sequence as set forth in any one of a) to d) except for one or several modifications (*i.e.* deletions, substitutions and/or insertions), such as 2, 3, 4, 5, 6, 7, 8, 9 amino acid modifications, or more amino acid modifications such as 10 and having lipolytic enzyme activity.

Suitably, the lipolytic enzyme may belong to the abH 15 superfamily, preferably the abH 15.01 superfamily.

5 Suitably, the lipolytic enzyme may comprise one of more of the following amino acid sequences

- a) SEQ ID NO. 25;
- b) SEQ ID NO: 26;
- 10 c) SEQ ID NO.25 lacking the signal peptide as indicated in Figure 36;
- d) an amino acid sequence having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably-at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%,
15 preferably at least 98%, or preferably at least 99% identity to any one of the amino acid sequences defined in a) to c); or
- e) an amino acid sequence as set forth in any one of a) to c) except for one or several modifications (*i.e.* deletions, substitutions and/or insertions), such as 2, 3, 4, 5, 6, 7, 8, 9 amino acid modifications, or more amino acid
20 modifications such as 10 and having lipolytic enzyme activity.

Suitably, the lipolytic enzyme may comprise a lipase cloned from *Geobacillus* species, preferably *G. stearothermophilus* strain T 1 (GeoT1), such as that shown in SEQ ID NO: 25. In some embodiments the lipolytic enzyme, such as GeoT1, is fused
25 to the carboxy-terminus of the catalytic domain of a bacterial cellulase such as that shown in SEQ ID NO: 26. In some embodiments, the bacterial cellulase is derived from a *Bacillus* strain deposited as CBS 670.93 (referred to as BCE103) with the Central Bureau voor Schimmelcultures, Baam, The Netherlands. In some
embodiments the lipolytic enzyme, such as GeoT1, is connected to the BCE103
30 cellulase by a cleavable linker. Thus in some embodiments the lipolytic enzyme, such as GeoT1, is not a fusion protein.

Suitably, the lipolytic enzyme may belong to the abH 18 superfamily, preferably the abH 18.01 superfamily.

35

Suitably, the lipolytic enzyme may comprise one of more of the following amino acid sequences

- f) SEQ ID NO: 27;
- g) SEQ ID NO: 28;
- h) SEQ ID NO: 27 lacking the signal peptide as indicated in Figure 36;
- 5 i) an amino acid sequence having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to any one of the
- 10 amino acid sequences defined in a) to c); or
- j) an amino acid sequence as set forth in any one of a) to c) except for one or several modifications (*i.e.* deletions, substitutions and/or insertions), such as 2, 3, 4, 5, 6, 7, 8, 9 amino acid modifications, or more amino acid modifications such as 10 and having lipolytic enzyme activity.

15

Suitably, the lipolytic enzyme may comprise a lipase cloned from *Bacillus subtilis*, preferably a lipaseA (LipA) from *Bacillus subtilis* such as that shown in SEQ ID NO: 27. In some embodiments, the lipolytic enzyme, such as LipA, is fused to the carboxy-terminus of the catalytic domain of a bacterial cellulase such as that shown

20 in SEQ ID NO:28. In some embodiments, the bacterial cellulase is derived from a *Bacillus* strain deposited as CBS 670.93 (referred to as BCE103) with the Central Bureau voor Schimmelcultures, Baam, The Netherlands. In some embodiments the lipolytic enzyme, such as LipA, is connected to the BCE103 cellulase by a cleavable linker. Thus in some embodiments the lipolytic enzyme, such as LipA, is not a fusion

25 protein.

In one aspect, as used herein, a "lipase", "lipase enzyme", "lipolytic enzymes", "lipolytic polypeptides", or "lipolytic proteins" refers to an enzyme, polypeptide, or protein exhibiting a lipid degrading capability such as a capability of degrading a

30 triglyceride or a phospholipid. The lipolytic enzyme may be, for example, a lipase, a phospholipase, an esterase or a cutinase. As used herein, lipolytic activity may be determined according to any procedure known in the art (see, *e.g.*, Gupta *et al.*, *Biotechnol. Appl. Biochem.*, **2003**, 37:63-71 ; U.S. Pat. No. 5,990,069; and International Publication No. WO 96/18729).

35

In one aspect, the present invention provides a detergent or cleaning composition comprising:

- a) a polypeptide as shown in SEQ ID NO: 17 or a fragment thereof having lipase activity;
- 5 b) a polypeptide having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to the amino acid sequence shown as SEQ ID
10 NO: 17 and having lipase activity; or
- c) a polypeptide as set forth in SEQ ID NO: 17 except for one or several modifications (*i.e.* deletions, substitutions and/or insertions), such as 2, 3, 4, 5, 6, 7, 8, 9 amino acid modifications, or more amino acid modifications such as
10 and having lipase activity;
- 15 d) a polypeptide encoded by the nucleotide sequence of SEQ ID NO: 23 or by a nucleic acid which is related to the nucleotide sequence of SEQ ID NO: 23 by the degeneration of the genetic code;
- e) a polypeptide having lipase activity encoded by a nucleic acid sequence having at least 70%, preferably at least 80%, preferably at least 85%,
20 preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to the amino acid sequence shown as SEQ ID
25 NO: 23 or to a nucleic acid which is related to the nucleotide sequence of SEQ ID NO: 23 by the degeneration of the genetic code;
- f) a polypeptide having lipase activity encoded by a nucleic acid sequence which hybridizes under stringent conditions to the complement of the nucleic acid sequence of SEQ ID NO: 23; or
- 30 g) a polypeptide obtainable (preferably obtained) from *Streptomyces* (preferably *S. pristinaespiralis*) having lipase activity.

Suitably, the polypeptide may be present in a concentration of 0.01 to 2 ppm by weight of the total weight of the composition. The composition may further comprise one or more enzymes selected from the group consisting of a protease, an amylase,
35 a glucoamylase, a maltogenic amylase, a non-maltogenic amylase, a lipase, a cutinase, a carbohydrase, a cellulase, a pectinase, a mannanase, an arabinase, a

galactanase, a xylianase, an oxidase, a laccase, a peroxidase, and an acyl transferase.

5 Suitably, the composition may comprise one or more surfactants, such as one or more surfactants selected from the group consisting of non-ionic (including semi-polar), anionic, cationic and zwitterionic.

Suitably, the composition may be in powder form or may be in liquid form.

10 The present invention further provides a method of removing a lipid-based stain from a surface by contacting the surface with a composition comprising:

- a) a polypeptide as shown in SEQ ID NO: 17 or a fragment thereof having lipase activity;
- 15 b) a polypeptide having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to the amino acid sequence shown as SEQ ID NO: 17 and having lipase activity; or
- 20 c) a polypeptide as set forth in SEQ ID NO: 17 except for one or several modifications (*i.e.* deletions, substitutions and/or insertions), such as 2, 3, 4, 5, 6, 7, 8, 9 amino acid modifications, or more amino acid modifications such as 10 and having lipase activity;
- d) a polypeptide encoded by the nucleotide sequence of SEQ ID NO: 23 or by a 25 nucleic acid which is related to the nucleotide sequence of SEQ ID NO: 23 by the degeneration of the genetic code;
- e) a polypeptide having lipase activity encoded by a nucleic acid sequence having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, 30 preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to the amino acid sequence shown as SEQ ID NO: 23 or to a nucleic acid which is related to the nucleotide sequence of SEQ ID NO: 23 by the degeneration of the genetic code;

- f) a polypeptide having lipase activity encoded by a nucleic acid sequence which hybridizes under stringent conditions to the complement of the nucleic acid sequence of SEQ ID NO: 23; or
- g) a polypeptide obtainable (preferably obtained) from *Streptomyces* (preferably *S. pristinaespiralis*) having lipase activity.

In another aspect, the present invention provides the use of a composition comprising:

- a) a polypeptide as shown in SEQ ID NO: 17 or a fragment thereof having lipase activity;
- b) a polypeptide having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to the amino acid sequence shown as SEQ ID NO: 17 and having lipase activity; or
- c) a polypeptide as set forth in SEQ ID NO: 17 except for one or several modifications (*i.e.* deletions, substitutions and/or insertions), such as 2, 3, 4, 5, 6, 7, 8, 9 amino acid modifications, or more amino acid modifications such as 10 and having lipase activity;
- d) a polypeptide encoded by the nucleotide sequence of SEQ ID NO: 23 or by a nucleic acid which is related to the nucleotide sequence of SEQ ID NO: 23 by the degeneration of the genetic code;
- e) a polypeptide having lipase activity encoded by a nucleic acid sequence having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to the amino acid sequence shown as SEQ ID NO: 23 or to a nucleic acid which is related to the nucleotide sequence of SEQ ID NO: 23 by the degeneration of the genetic code;
- f) a polypeptide having lipase activity encoded by a nucleic acid sequence which hybridizes under stringent conditions to the complement of the nucleic acid sequence of SEQ ID NO: 23; or
- g) a polypeptide obtainable (preferably obtained) from *Streptomyces* (preferably *S. pristinaespiralis*) having lipase activity,

in cleaning and/or in a detergent. For example, such use may be to reduce or remove lipid stains from a surface.

In another aspect, the present invention provides a method of cleaning a surface, comprising contacting the surface with a composition comprising:

- a) a polypeptide as shown in SEQ ID NO: 17 or a fragment thereof having lipase activity;
- b) a polypeptide having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to the amino acid sequence shown as SEQ ID NO: 17 and having lipase activity; or
- c) a polypeptide as set forth in SEQ ID NO: 17 except for one or several modifications (*i. e.* deletions, substitutions and/or insertions), such as 2, 3, 4, 5, 6, 7, 8, 9 amino acid modifications, or more amino acid modifications such as 10 and having lipase activity;
- d) a polypeptide encoded by the nucleotide sequence of SEQ ID NO: 23 or by a nucleic acid which is related to the nucleotide sequence of SEQ ID NO: 23 by the degeneration of the genetic code;
- e) a polypeptide having lipase activity encoded by a nucleic acid sequence having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to the amino acid sequence shown as SEQ ID NO: 23 or to a nucleic acid which is related to the nucleotide sequence of SEQ ID NO: 23 by the degeneration of the genetic code;
- f) a polypeptide having lipase activity encoded by a nucleic acid sequence which hybridizes under stringent conditions to the complement of the nucleic acid sequence of SEQ ID NO: 23; or
- g) a polypeptide obtainable (preferably obtained) from *Streptomyces* (preferably *S. pristinaespiralis*) having lipase activity.

In a further aspect, the present invention provides a method of cleaning an item, comprising contacting the item with a composition comprising:

- a) a polypeptide as shown in SEQ ID NO: 17 or a fragment thereof having lipase activity;
- b) a polypeptide having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to the amino acid sequence shown as SEQ ID NO: 17 and having lipase activity; or
- c) a polypeptide as set forth in SEQ ID NO: 17 except for one or several modifications (*i.e.* deletions, substitutions and/or insertions), such as 2, 3, 4, 5, 6, 7, 8, 9 amino acid modifications, or more amino acid modifications such as 10 and having lipase activity;
- d) a polypeptide encoded by the nucleotide sequence of SEQ ID NO: 23 or by a nucleic acid which is related to the nucleotide sequence of SEQ ID NO: 23 by the degeneration of the genetic code;
- e) a polypeptide having lipase activity encoded by a nucleic acid sequence having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to the amino acid sequence shown as SEQ ID NO: 23 or to a nucleic acid which is related to the nucleotide sequence of SEQ ID NO: 23 by the degeneration of the genetic code;
- f) a polypeptide having lipase activity encoded by a nucleic acid sequence which hybridizes under stringent conditions to the complement of the nucleic acid sequence of SEQ ID NO: 23; or
- g) a polypeptide obtainable (preferably obtained) from *Streptomyces* (preferably *S. pristinaespiralis*) having lipase activity.

Suitably, the item may be a clothing item or a tableware item.

The present invention provides many applications, methods and uses of a composition comprising a lipolytic enzyme and a hydrophobin. For the avoidance of doubt, each of these applications, methods and uses may be applied to a composition comprising:

- a) a polypeptide as shown in SEQ ID NO: 17 or a fragment thereof having lipase activity;
- b) a polypeptide having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to the amino acid sequence shown as SEQ ID NO: 17 and having lipase activity; or
- c) a polypeptide as set forth in SEQ ID NO: 17 except for one or several modifications (*i.e.* deletions, substitutions and/or insertions), such as 2, 3, 4, 5, 6, 7, 8, 9 amino acid modifications, or more amino acid modifications such as 10 and having lipase activity;
- d) a polypeptide encoded by the nucleotide sequence of SEQ ID NO: 23 or by a nucleic acid which is related to the nucleotide sequence of SEQ ID NO: 23 by the degeneration of the genetic code;
- e) a polypeptide having lipase activity encoded by a nucleic acid sequence having at least 70%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 91%, preferably at least 92%, preferably at least 93%, preferably at least 94%, preferably at least 95%, preferably at least 96%, preferably at least 97%, preferably at least 98%, or preferably at least 99% identity to the amino acid sequence shown as SEQ ID NO: 23 or to a nucleic acid which is related to the nucleotide sequence of SEQ ID NO: 23 by the degeneration of the genetic code;
- f) a polypeptide having lipase activity encoded by a nucleic acid sequence which hybridizes under stringent conditions to the complement of the nucleic acid sequence of SEQ ID NO: 23; or
- g) a polypeptide obtainable (preferably obtained) from *Streptomyces* (preferably *S. pristinaespiralis*) having lipase activity.

30 HOST CELL

The term "host cell" - in relation to the present invention includes any cell that comprises either the nucleotide sequence or an expression vector as described above and which is used in the recombinant production of an enzyme having the specific properties as defined herein.

Thus, a further embodiment of the present invention provides host cells transformed or transfected with a nucleotide sequence that expresses the enzyme of the present invention. The cells will be chosen to be compatible with the said vector and may for example be prokaryotic (for example bacterial), fungal, yeast or plant cells.

5 Preferably, the host cells are not human cells.

Examples of suitable bacterial host organisms are gram positive or gram negative bacterial species.

10 Depending on the nature of the nucleotide sequence encoding the enzyme of the present invention, and/or the desirability for further processing of the expressed protein, eukaryotic hosts such as yeasts or other fungi may be preferred. However, some proteins are either poorly secreted from the yeast cell, or in some cases are not processed properly {e.g., hyper-glycosylation in yeast). In these instances, a
15 different fungal host organism should be selected.

The use of suitable host cells - such as yeast, fungal and plant host cells - may provide for post-translational modifications (e.g., myristoylation, glycosylation, truncation, lipidation and tyrosine, serine or threonine phosphorylation, or N-terminal
20 acetylation as may be needed to confer optimal biological activity on recombinant expression products of the present invention.

The host cell may be a protease deficient or protease minus strain.

25 The genotype of the host cell may be modified to improve expression.

Examples of host cell modifications include protease deficiency, supplementation of rare tRNAs, and modification of the reductive potential in the cytoplasm to enhance disulphide bond formation.

30

For example, the host cell *E. coli* may overexpress rare tRNAs to improve expression of heterologous proteins as exemplified/described in Kane (*Curr Opin Biotechnol* (1995), 6, 494-500 "Effects of rare codon clusters on high-level expression of heterologous proteins in *E. coli*"). The host cell may be deficient in a number of
35 reducing enzymes thus favouring formation of stable disulphide bonds as exemplified/described in Bessette (*Proc Natl Acad Sci USA* (1999), 96, 13703-13708

"Efficient folding of proteins with multiple disulphide bonds in the *Escherichia coli* cytoplasm").

ISOLATED

5

In one aspect, the enzymes for use in the present invention may be in an isolated form.

The term "isolated" means that the sequence or protein is at least substantially free from at least one other component with which the sequence or protein is naturally associated in nature and as found in nature.

10

PURIFIED

In one aspect, the enzymes for use in the present invention may be used in a purified form.

15

The term "purified" means that the sequence is in a relatively pure state – e.g., at least about 51% pure, or at least about 75%, or at least about 80%, or at least about 90% pure, or at least about 95% pure or at least about 98% pure.

20

CLONING A NUCLEOTIDE SEQUENCE ENCODING A POLYPEPTIDE ACCORDING TO THE PRESENT INVENTION

A nucleotide sequence encoding either a polypeptide which has the specific properties as defined herein or a polypeptide which is suitable for modification may be isolated from any cell or organism producing said polypeptide. Various methods are well known within the art for the isolation of nucleotide sequences.

25

For example, a genomic DNA and/or cDNA library may be constructed using chromosomal DNA or messenger RNA from the organism producing the polypeptide. If the amino acid sequence of the polypeptide is known, labelled oligonucleotide probes may be synthesised and used to identify polypeptide-encoding clones from the genomic library prepared from the organism. Alternatively, a labelled oligonucleotide probe containing sequences homologous to another known

30

35

polypeptide gene could be used to identify polypeptide-encoding clones. In the latter case, hybridisation and washing conditions of lower stringency are used.

5 Alternatively, polypeptide-encoding clones could be identified by inserting fragments of genomic DNA into an expression vector, such as a plasmid, transforming enzyme-negative bacteria with the resulting genomic DNA library, and then plating the transformed bacteria onto agar containing an enzyme inhibited by the polypeptide, thereby allowing clones expressing the polypeptide to be identified.

10 In a yet further alternative, the nucleotide sequence encoding the polypeptide may be prepared synthetically by established standard methods, *e.g.*, the phosphoroamidite method described by Beaucage S.L. *et al.* (1981) *Tetrahedron Letters* 22, 1859-1869, or the method described by Matthes *et al.* (1984) *EMBO J.* 3, 801-805. In the phosphoroamidite method, oligonucleotides are synthesised, *e.g.*, in an automatic
15 DNA synthesiser, purified, annealed, ligated and cloned in appropriate vectors.

The nucleotide sequence may be of mixed genomic and synthetic origin, mixed synthetic and cDNA origin, or mixed genomic and cDNA origin, prepared by ligating
20 fragments of synthetic, genomic or cDNA origin (as appropriate) in accordance with standard techniques. Each ligated fragment corresponds to various parts of the entire nucleotide sequence. The DNA sequence may also be prepared by polymerase chain reaction (PGR) using specific primers, for instance as described in US 4,683,202 or in Saiki R K *et al.* (*Science* (1988) **239**, 487-491).

25 NUCLEOTIDE SEQUENCES

The present invention also encompasses nucleotide sequences encoding polypeptides having the specific properties as defined herein. The term "nucleotide
30 sequence" as used herein refers to an oligonucleotide sequence or polynucleotide sequence, and variant, homologues, fragments and derivatives thereof (such as portions thereof). The nucleotide sequence may be of genomic or synthetic or recombinant origin, which may be double-stranded or single-stranded whether representing the sense or antisense strand.

35 The term "nucleotide sequence" in relation to the present invention includes genomic DNA, cDNA, synthetic DNA, and RNA. Preferably it means DNA, more preferably cDNA for the coding sequence.

In a preferred embodiment, the nucleotide sequence *per se* encoding a polypeptide having the specific properties as defined herein does not cover the native nucleotide sequence in its natural environment when it is linked to its naturally associated sequence(s) that is/are also in its/their natural environment. For ease of reference, we shall call this preferred embodiment the "non-native nucleotide sequence". In this regard, the term "native nucleotide sequence" means an entire nucleotide sequence that is in its native environment and when operatively linked to an entire promoter with which it is naturally associated, which promoter is also in its native environment.

However, the amino acid sequence encompassed by scope of the present invention can be isolated and/or purified post expression of a nucleotide sequence in its native organism. Preferably, however, the amino acid sequence encompassed by scope of the present invention may be expressed by a nucleotide sequence in its native organism but wherein the nucleotide sequence is not under the control of the promoter with which it is naturally associated within that organism.

Preferably the polypeptide is not a native polypeptide. In this regard, the term "native polypeptide" means an entire polypeptide that is in its native environment and when it has been expressed by its native nucleotide sequence.

Typically, the nucleotide sequence encoding polypeptides having the specific properties as defined herein is prepared using recombinant DNA techniques (*i.e.*, recombinant DNA). However, in an alternative embodiment of the invention, the nucleotide sequence could be synthesised, in whole or in part, using chemical methods well known in the art (see Caruthers **MH** *et al.* (1980) *Nuc Acids Res Symp Ser* 2 15-23 and Horn T *et al.* (1980) *Nuc Acids Res Symp Ser* 225-232).

MOLECULAR EVOLUTION

Once an enzyme-encoding nucleotide sequence has been isolated, or a putative enzyme-encoding nucleotide sequence has been identified, it may be desirable to modify the selected nucleotide sequence, for example it may be desirable to mutate the sequence in order to prepare an enzyme in accordance with the present invention.

Mutations may be introduced using synthetic oligonucleotides. These oligonucleotides contain nucleotide sequences flanking the desired mutation sites.

A suitable method is disclosed in Morinaga *et al.* (*Biotechnology* (1984) 2, 646-649).

5 Another method of introducing mutations into enzyme-encoding nucleotide sequences is described in Nelson and Long (*Analytical Biochemistry* (1989), 180, 147-151).

10 Instead of site directed mutagenesis, such as described above, one can introduce mutations randomly for instance using a commercial kit such as the GeneMorph PGR mutagenesis kit from Stratagene, or the Diversify PGR random mutagenesis kit from Clontech. EP 0 583 265 refers to methods of optimising PGR based mutagenesis, which can also be combined with the use of mutagenic DNA analogues such as those described in EP 0 866 796. Error prone PGR technologies are suitable for the
15 production of variants of lipid acyl transferases with preferred characteristics. WO 02/06457 refers to molecular evolution of lipases.

A third method to obtain novel sequences is to fragment non-identical nucleotide sequences, either by using any number of restriction enzymes or an enzyme such as
20 Dnase I, and reassembling full nucleotide sequences coding for functional proteins. Alternatively one can use one or multiple non-identical nucleotide sequences and introduce mutations during the reassembly of the full nucleotide sequence. DNA shuffling and family shuffling technologies are suitable for the production of variants of lipid acyl transferases with preferred characteristics. Suitable methods for
25 performing 'shuffling' can be found in EP 0 752 008, EP 1 138 763, EP 1 103 606. Shuffling can also be combined with other forms of DNA mutagenesis as described in US 6,180,406 and WO 01/34835.

30 Thus, it is possible to produce numerous site directed or random mutations into a nucleotide sequence, either *in vivo* or *in vitro*, and to subsequently screen for improved functionality of the encoded polypeptide by various means. Using *in silico* and exo-mediated recombination methods (see, e.g., WO 00/58517, US 6,344,328, US 6,361,974), for example, molecular evolution can be performed where the variant produced retains very low homology to known enzymes or proteins. Such variants
35 thereby obtained may have significant structural analogy to known transferase enzymes, but have very low amino acid sequence homology.

As a **non-limiting** example, in addition, mutations or natural variants of a polynucleotide sequence can be recombined with either the wild type or other mutations or natural variants to produce new variants. Such new variants can also be screened for improved functionality of the encoded polypeptide.

5

The application of the above-mentioned and similar molecular evolution methods allows the identification and selection of variants of the enzymes of the present invention which have preferred characteristics without any prior knowledge of protein structure or function, and allows the production of non-predictable but beneficial mutations or variants. There are numerous examples of the application of molecular evolution in the art for the optimisation or alteration of enzyme activity, such examples include, but are not limited to one or more of the following: optimised expression and/or activity in a host cell or in vitro, increased enzymatic activity, altered substrate and/or product specificity, increased or decreased enzymatic or structural stability, altered enzymatic activity/specificity in preferred environmental conditions, e.g., temperature, pH, substrate.

10

15

As will be apparent to a person skilled in the art, using molecular evolution tools an enzyme may be altered to improve the functionality of the enzyme.

20

Suitably, the nucleotide sequence encoding a lipolytic enzyme used in the invention may encode a variant, *i.e.*, the lipolytic enzyme may contain at least one amino acid substitution, deletion or addition, when compared to a parental enzyme. Variant enzymes retain at least **70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 99%** homology with the parent enzyme.

25

Variant lipolytic enzymes may have decreased activity on triglycerides, and/or monoglycerides and/or diglycerides compared with the parent enzyme.

Suitably the variant enzyme may have no activity on triglycerides and/or monoglycerides and/or diglycerides.

30

Alternatively, the variant enzyme may have increased thermostability.

The variant enzyme may have increased activity on one or more of the following, polar lipids, phospholipids, lecithin, phosphatidylcholine, glycolipids, digalactosyl monoglyceride, monogalactosyl monoglyceride.

35

Variants of lipid acyltransferases are known, and one or more of such variants may be suitable for use in the methods and uses according to the present invention and/or in the enzyme compositions according to the present invention. By way of example only, variants of lipid acyltransferases are described in the following references may be used in accordance with the present invention: Hilton & Buckley *J. Biol. Chem.* 1991 Jan 15; 266 : 997-1000; Robertson *et al. J. Biol. Chem.* 1994 Jan 21; 269: 2146-50; Brumlik *et al. J. Bacteriol.* 1996 Apr; 178 : 2060-4; Peelman *et al. Protein Sci.* 1998 Mar; 7:587-99.

10

AMINO ACID SEQUENCES

The present invention also encompasses the use of amino acid sequences encoded by a nucleotide sequence which encodes an enzyme for use in any one of the methods and/or uses of the present invention.

15

As used herein, the term "amino acid sequence" is synonymous with the term "polypeptide" and/or the term "protein". In some instances, the term "amino acid sequence" is synonymous with the term "peptide". In some instances, the term "amino acid sequence" is synonymous with "enzyme".

20

The amino acid sequence may be prepared/isolated from a suitable source, or it may be made synthetically or it may be prepared by use of recombinant DNA techniques.

25

Suitably, the amino acid sequences may be obtained from the isolated polypeptides taught herein by standard techniques.

One suitable method for determining amino acid sequences from isolated polypeptides is as follows:

30

Purified polypeptide may be freeze-dried and 100 µg of the freeze-dried material may be dissolved in 50 µl of a mixture of 8 M urea and 0.4 M ammonium hydrogen carbonate, pH 8.4. The dissolved protein may be denatured and reduced for 15 minutes at 50°C following overlay with nitrogen and addition of 5 µl of 45 mM dithiothreitol. After cooling to room temperature, 5 µl of 100 mM iodoacetamide may be added for the cysteine residues to be derivatized for 15 minutes at room temperature in the dark under nitrogen.

35

135 μ l of water and 5 μ g of endoproteinase Lys-C in 5 μ l of water may be added to the above reaction mixture and the digestion may be carried out at 37°C under nitrogen for 24 hours.

5

The resulting peptides may be separated by reverse phase HPLC on a VYDAC C18 column (0.46x1 5cm; 10 μ m; The Separation Group, California, USA) using solvent A: 0.1% TFA in water and solvent B: 0.1% TFA in acetonitrile. Selected peptides may be re-chromatographed on a Develosil C18 column using the same solvent system, prior to N-terminal sequencing. Sequencing may be done using an Applied Biosystems 476A sequencer using pulsed liquid fast cycles according to the manufacturer's instructions (Life Technologies, California, USA).

10

SEQUENCE IDENTITY OR SEQUENCE HOMOLOGY

15

Here, the term "homologue" means an entity having a certain homology with the subject amino acid sequences and the subject nucleotide sequences. Here, the term "homology" can be equated with "identity".

20

The homologous amino acid sequence and/or nucleotide sequence should provide and/or encode a polypeptide which retains the functional activity and/or enhances the activity of the enzyme.

25

In the present context, a homologous sequence is taken to include an amino acid sequence which may be at least 50%, 55%, 60%, 70%, 71%, 72%, 73%, 74%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical, preferably at least 95%, 96%, 97%, 98%, or 99% identical to the subject sequence. Typically, the homologues will comprise the same active sites etc. as the subject amino acid sequence. Although homology can also be considered in terms of similarity (*i.e.*, amino acid residues having similar chemical properties/functions), in the context of the present invention it is preferred to express homology in terms of sequence identity.

30

35

In the present context, a homologous sequence is taken to include a nucleotide sequence which may be at least 75, 85 or 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical, preferably at least 95%, 96%, 97%, 98%, or 99% identical to a nucleotide sequence encoding a polypeptide of the present invention

(the subject sequence). Typically, the homologues will comprise the same sequences that code for the active sites etc. as the subject sequence. Although homology can also be considered in terms of similarity (*i.e.*, amino acid residues having similar chemical properties/functions), in the context of the present invention it is preferred to express homology in terms of sequence identity.

Homology comparisons can be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs can **calculate** % homology between two or more sequences.

% homology may be calculated over contiguous sequences, *i.e.*, one sequence is aligned with the other sequence and each amino acid in one sequence is directly compared with the corresponding amino acid in the other sequence, one residue at a time. This is called an "ungapped" alignment. Typically, such ungapped alignments are performed only over a relatively short number of residues.

Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion will cause the following amino acid residues to be put out of alignment, thus potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into consideration possible insertions and deletions without penalising unduly the overall homology score. This is achieved by inserting "gaps" in the sequence alignment to try to maximise local homology.

However, these more complex methods assign "gap penalties" to each gap that occurs in the alignment so that, for the same number of identical amino acids, a sequence alignment with as few gaps as possible - reflecting higher relatedness between the two compared sequences - will achieve a higher score than one with many gaps. "Affine gap costs" are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties will of course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, it is preferred to use the default values when using such software for sequence comparisons.

Calculation of maximum % homology therefore firstly requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for carrying out such an alignment is the Vector NTI (Invitrogen Corp.).

5 Examples of other software that can perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel *et al.* 1999 Short Protocols in Molecular Biology, 4th Ed - Chapter 18), and FASTA (Altschui *et al.* 1990 *J. Mol. Biol.* 403-410). Both BLAST and FASTA are available for offline and online searching (see Ausubel *et al.* 1999, pages 7-58 to 7-60). However, for some applications, it is preferred to use the Vector NTI program. A new tool, called BLAST 2 Sequences is also available for comparing protein and nucleotide sequence (see *FEMS Microbiol Lett* 1999 174: 247-50; *FEMS Microbiol Lett* 1999 177: 187-8 and tatiana@ncbi.nlm.nih.gov).

15 Although the final % homology can be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each pairwise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix - the default matrix for the BLAST suite of programs. Vector NTI programs generally use either the public default values or a custom symbol comparison table if supplied (see user manual for further details). For some applications, it is preferred to use the default values for the Vector NTI ADVANCE™ 10 package.

25 Alternatively, percentage homologies may be calculated using the multiple alignment feature in Vector NTI ADVANCE™ 10 (Invitrogen Corp.), based on an algorithm, analogous to CLUSTAL (Higgins DG & Sharp PM (1988), *Gene* 73, 237-244).

30 Once the software has produced an optimal alignment, it is possible to calculate % homology, preferably % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

Suitably, the degree of identity with regard to a nucleotide sequence is determined over at least 20 contiguous nucleotides, preferably over at least 30 contiguous nucleotides, preferably over at least 40 contiguous nucleotides, preferably over at least 50 contiguous nucleotides, preferably over at least 60 contiguous nucleotides, preferably over at least 100 contiguous nucleotides.

Suitably, the degree of identity with regard to a nucleotide sequence may be determined over the whole sequence.

- 5 Should Gap Penalties be used when determining sequence identity, then preferably the default parameters for the programme are used for pairwise alignment. For example, the following parameters are the current default parameters for pairwise alignment for BLAST 2:

FOR BLAST2	DNA	PROTEIN
EXPECT THRESHOLD	10	10
WORD SIZE	11	3
SCORING PARAMETERS		
Match/Mismatch Scores	2, -3	n/a
Matrix	n/a	BLOSUM62
Gap Costs	Existence: 5 Extension: 2	Existence: 11 Extension: 1

- 10 In one embodiment, preferably the sequence identity for the nucleotide sequences and/or amino acid sequences may be determined using BLAST2 (blastn) with the scoring parameters set as defined above.

For the purposes of the present invention, the degree of identity is based on the
 15 number of sequence elements which are the same. The degree of identity in accordance with the present invention for amino acid sequences may be suitably determined by means of computer programs known in the art such as Vector NTI ADVANCE™ 11 (Invitrogen Corp.). For pairwise alignment the scoring parameters used are preferably BLOSUM62 with Gap existence penalty of 11 and Gap
 20 extension penalty of 1.

Suitably, the degree of identity with regard to an amino acid sequence is determined over at least 20 contiguous amino acids, preferably over at least 30 contiguous amino acids, preferably over at least 40 contiguous amino acids, preferably over at
 25 least 50 contiguous amino acids, preferably over at least 60 contiguous amino acids, preferably over at least 100 contiguous amino acids.

Suitably, the degree of identity with regard to an amino acid sequence may be determined over the whole sequence.

The sequences may also have deletions, insertions or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent substance. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the secondary binding activity of the substance is retained. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine, valine, glycine, alanine, asparagine, glutamine, serine, threonine, phenylalanine, and tyrosine.

Conservative substitutions may be made, for example according to the Table below. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

ALIPHATIC	Non-polar	G A P
		I L V
	Polar – uncharged	C S T M
		N Q
	Polar – charged	D E
		K R
AROMATIC		H F W Y

The present invention also encompasses homologous substitution (substitution and replacement are both used herein to mean the interchange of an existing amino acid residue, with an alternative residue) that may occur, *i.e.*, like-for-like substitution such as basic for basic, acidic for acidic, polar for polar etc. Non-homologous substitution may also occur, *i.e.*, from one class of residue to another or alternatively involving the inclusion of unnatural amino acids such as ornithine (hereinafter referred to as Z), diaminobutyric acid ornithine (hereinafter referred to as B), norleucine ornithine (hereinafter referred to as O), pyridylalanine, thienylalanine, naphthylalanine and phenylglycine.

Replacements may also be made by unnatural amino acids.

Variant amino acid sequences may include suitable spacer groups that may be inserted between any two amino acid residues of the sequence including alkyl groups such as methyl, ethyl or propyl groups in addition to amino acid spacers such as glycine or β -alanine residues. A further form of variation, involves the presence of one or more amino acid residues in peptoid form, will be well understood by those skilled in the art. For the avoidance of doubt, "the peptoid form" is used to refer to variant amino acid residues wherein the α -carbon substituent group is on the residue's nitrogen atom rather than the α -carbon. Processes for preparing peptides in the peptoid form are known in the art, for example Simon RJ *et al.*, *PNAS* (1992) 89, 9367-9371 and Horwell DC, *Trends Biotechnol.* (1995) 13, 132-134.

Nucleotide sequences for use in the present invention or encoding a polypeptide having the specific properties defined herein may include within them synthetic or modified nucleotides. A number of different types of modification to oligonucleotides are known in the art. These include methylphosphonate and phosphorothioate backbones and/or the addition of acridine or polylysine chains at the 3' and/or 5' ends of the molecule. For the purposes of the present invention, it is to be understood that the nucleotide sequences described herein may be modified by any method available in the art. Such modifications may be carried out in order to enhance the *in vivo* activity or life span of nucleotide sequences.

The present invention also encompasses the use of nucleotide sequences that are complementary to the sequences discussed herein, or any derivative, fragment or derivative thereof. If the sequence is complementary to a fragment thereof then that sequence can be used as a probe to identify similar coding sequences in other organisms etc.

Polynucleotides which are not 100% homologous to the sequences of the present invention but fall within the scope of the invention can be obtained in a number of ways. Other variants of the sequences described herein may be obtained for example by probing DNA libraries made from a range of individuals, for example individuals from different populations. In addition, other viral/bacterial, or cellular homologues particularly cellular homologues found in mammalian cells (*e.g.*, rat, mouse, bovine and primate cells), may be obtained and such homologues and fragments thereof in general will be capable of selectively hybridising to the sequences shown in the sequence listing herein. Such sequences may be obtained

by probing cDNA libraries made from or genomic DNA libraries from other animal species, and probing such libraries with probes comprising all or part of any one of the sequences in the attached sequence listings under conditions of medium to high stringency. Similar considerations apply to obtaining species homologues and allelic variants of the polypeptide or nucleotide sequences of the invention.

Variants and strain/species homologues may also be obtained using degenerate PGR which will use primers designed to target sequences within the variants and homologues encoding conserved amino acid sequences within the sequences of the present invention. Conserved sequences can be predicted, for example, by aligning the amino acid sequences from several variants/homologues. Sequence alignments can be performed using computer software known in the art. For example the GCG Wisconsin PileUp program is widely used.

The primers used in degenerate PGR will contain one or more degenerate positions and will be used at stringency conditions lower than those used for cloning sequences with single sequence primers against known sequences.

Alternatively, such polynucleotides may be obtained by site directed mutagenesis of characterised sequences. This may be useful where for example silent codon sequence changes are required to optimise codon preferences for a particular host cell in which the polynucleotide sequences are being expressed. Other sequence changes may be desired in order to introduce restriction polypeptide recognition sites, or to alter the property or function of the polypeptides encoded by the polynucleotides.

Polynucleotides (nucleotide sequences) of the invention may be used to produce a primer, e.g., a PGR primer, a primer for an alternative amplification reaction, a probe e.g., labelled with a revealing label by conventional means using radioactive or non-radioactive labels, or the polynucleotides may be cloned into vectors. Such primers, probes and other fragments will be at least 15, preferably at least 20, for example at least 25, 30 or 40 nucleotides in length, and are also encompassed by the term polynucleotides of the invention as used herein.

Polynucleotides such as DNA polynucleotides and probes according to the invention may be produced recombinantly, synthetically, or by any means available to those of skill in the art. They may also be cloned by standard techniques.

In general, primers will be produced by synthetic means, involving a stepwise manufacture of the desired nucleic acid sequence one nucleotide at a time. Techniques for accomplishing this using automated techniques are readily available in the art.

5

Longer polynucleotides will generally be produced using recombinant means, for example using a PGR (polymerase chain reaction) cloning techniques. This will involve making a pair of primers (e.g., of about 15 to 30 nucleotides) flanking a region of the lipid targeting sequence which it is desired to clone, bringing the primers into contact with mRNA or cDNA obtained from an animal or human cell, performing a polymerase chain reaction under conditions which bring about amplification of the desired region, isolating the amplified fragment (e.g., by purifying the reaction mixture on an agarose gel) and recovering the amplified DNA. The primers may be designed to contain suitable restriction enzyme recognition sites so that the amplified DNA can be cloned into a suitable cloning vector.

10
15

HYBRIDISATION

The present invention also encompasses the use of sequences that are complementary to the sequences of the present invention or sequences that are capable of hybridising either to the sequences of the present invention or to sequences that are complementary thereto.

20

The term "hybridisation" as used herein shall include "the process by which a strand of nucleic acid joins with a complementary strand through base pairing" as well as the process of amplification as carried out in polymerase chain reaction (PGR) technologies.

25

The present invention also encompasses the use of nucleotide sequences that are capable of hybridising to the sequences that are complementary to the subject sequences discussed herein, or any derivative, fragment or derivative thereof.

30

The present invention also encompasses sequences that are complementary to sequences that are capable of hybridising to the nucleotide sequences discussed herein.

35

Hybridisation conditions are based on the melting temperature (T_m) of the nucleotide binding complex, as taught in Berger and Kimmei (1987, Guide to Molecular Cloning Techniques, Methods in Enzymology. Vol 152, Academic Press, San Diego CA), and confer a defined "stringency" as explained below.

5

Maximum stringency typically occurs at about $T_m - 5^\circ\text{C}$ (5°C below the T_m of the probe); high stringency at about 5°C to 10°C below T_m ; intermediate stringency at about 10°C to 20°C below T_m ; and low stringency at about 20°C to 25°C below T_m . As will be understood by those of skill in the art, a maximum stringency hybridisation can be used to identify or detect identical nucleotide sequences while an intermediate (or low) stringency hybridisation can be used to identify or detect similar or related polynucleotide sequences.

10

Preferably, the present invention encompasses the use of sequences that are complementary to sequences that are capable of hybridising under high stringency conditions or intermediate stringency conditions to nucleotide sequences encoding polypeptides having the specific properties as defined herein.

15

More preferably, the present invention encompasses the use of sequences that are complementary to sequences that are capable of hybridising under high stringency conditions (e.g., 65°C and $0.1\times\text{SSC}$ ($1\times\text{SSC} = 0.15\text{ M NaCl}, 0.015\text{ M Na-citrate pH } 7.0$)) to nucleotide sequences encoding polypeptides having the specific properties as defined herein.

20

The present invention also relates to the use of nucleotide sequences that can hybridise to the nucleotide sequences discussed herein (including complementary sequences of those discussed herein).

25

The present invention also relates to the use of nucleotide sequences that are complementary to sequences that can hybridise to the nucleotide sequences discussed herein (including complementary sequences of those discussed herein).

30

Also included within the scope of the present invention are the use of polynucleotide sequences that are capable of hybridising to the nucleotide sequences discussed herein under conditions of intermediate to maximal stringency.

35

In a preferred aspect, the present invention covers the use of nucleotide sequences that can hybridise to the nucleotide sequences discussed herein, or the complement thereof, under stringent conditions {e.g., 50°C and 0.2 x SSC).

5 In a more preferred aspect, the present invention covers the use of nucleotide sequences that can hybridise to the nucleotide sequences discussed herein, or the complement thereof, under high stringency conditions (e.g., 65°C and 0.1 x SSC).

BIOLOGICALLY ACTIVE

10

Preferably, the variant sequences etc. are at least as biologically active as the sequences presented herein.

As used herein "biologically active" refers to a sequence having a similar structural
15 function (but not necessarily to the same degree), and/or similar regulatory function (but not necessarily to the same degree), and/or similar biochemical function (but not necessarily to the same degree) of the naturally occurring sequence.

RECOMBINANT

20

In one aspect the sequence for use in the present invention is a recombinant sequence - *i.e.*, a sequence that has been prepared using recombinant DNA techniques.

25 These recombinant DNA techniques are within the capabilities of a person of ordinary skill in the art. Such techniques are explained in the literature, for example, J. Sambrook, E. F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory Manual*, Second Edition, Books 1-3, Cold Spring Harbor Laboratory Press.

30 SYNTHETIC

In one aspect the sequence for use in the present invention is a synthetic sequence - *i.e.*, a sequence that has been prepared by *in vitro* chemical or enzymatic synthesis. It includes, but is not limited to, sequences made with optimal codon usage for host
35 organisms - such as the methylotrophic yeasts *Pichia* and *Hansenula*.

EXPRESSION OF POLYPEPTIDES

A nucleotide sequence for use in the present invention or for encoding a polypeptide
5 having the specific properties as defined herein can be incorporated into a
recombinant replicable vector. The vector may be used to replicate and express the
nucleotide sequence, in polypeptide form, in and/or from a compatible host cell.
Expression may be controlled using control sequences which include
promoters/enhancers and other expression regulation signals. Prokaryotic promoters
10 and promoters functional in eukaryotic cells may be used. Tissue specific or stimuli
specific promoters may be used. Chimeric promoters may also be used comprising
sequence elements from two or more different promoters described above.

The polypeptide produced by a host recombinant cell by expression of the nucleotide
15 sequence may be secreted or may be contained intracellularly depending on the
sequence and/or the vector used. The coding sequences can be designed with
signal sequences which direct secretion of the substance coding sequences through
a particular prokaryotic or eukaryotic cell membrane.

20 EXPRESSION VECTOR

The term "expression vector" means a construct capable of *in vivo* or *in vitro*
expression.

25 Preferably, the expression vector is incorporated into the genome of a suitable host
organism. The term "incorporated" preferably covers stable incorporation into the
genome.

The nucleotide sequence encoding an enzyme for use in the present invention may
30 be present in a vector in which the nucleotide sequence is operably linked to
regulatory sequences capable of providing for the expression of the nucleotide
sequence by a suitable host organism.

The vectors for use in the present invention may be transformed into a suitable host
35 cell as described below to provide for expression of a polypeptide of the present
invention.

The choice of vector *e.g.*, a plasmid, cosmid, or phage vector will often depend on the host cell into which it is to be introduced.

5 The vectors for use in the present invention may contain one or more selectable marker genes such as a gene which confers antibiotic resistance *e.g.*, ampicillin, kanamycin, chloramphenicol or tetracycline resistance. Alternatively, the selection may be accomplished by co-transformation (as described in WO 91/17243).

10 Vectors may be used *in vitro*, for example for the production of RNA or used to transfect, transform, transduce or infect a host cell.

The vector may further comprise a nucleotide sequence enabling the vector to replicate in the host cell in question. Examples of such sequences are the origins of replication of plasmids pUC1 9, pACYC1 77, pUB1 10, pE1 94, pAMB1 and pIJ702.

REGULATORY SEQUENCES

In some applications, the nucleotide sequence for use in the present invention is operably linked to a regulatory sequence which is capable of providing for the expression of the nucleotide sequence, such as by the chosen host cell. By way of example, the present invention covers a vector comprising the nucleotide sequence of the present invention operably linked to such a regulatory sequence, *i.e.*, the vector is an expression vector.

25 The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A regulatory sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under condition compatible with the control sequences.

The term "regulatory sequences" includes promoters and enhancers and other expression regulation signals.

35 The term "promoter" is used in the normal sense of the art, *e.g.* an RNA polymerase binding site.

Enhanced expression of the nucleotide sequence encoding the enzyme of the present invention may also be achieved by the selection of heterologous regulatory regions, e.g., promoter, secretion leader and terminator regions.

5

Preferably, the nucleotide sequence according to the present invention is operably linked to at least a promoter.

10

Examples of suitable promoters for directing the transcription of the nucleotide sequence in a bacterial, fungal or yeast host are well known in the art.

CONSTRUCTS

15

The term "construct" - which is synonymous with terms such as "conjugate", "cassette" and "hybrid" - includes a nucleotide sequence encoding a polypeptide having the specific properties as defined herein for use according to the present invention directly or indirectly attached to a promoter. An example of an indirect attachment is the provision of a suitable spacer group such as an intron sequence, such as the Sh1-intron or the ADH intron, intermediate the promoter and the nucleotide sequence of the present invention. The same is true for the term "fused" in relation to the present invention which includes direct or indirect attachment. In some cases, the terms do not cover the natural combination of the nucleotide sequence coding for the protein ordinarily associated with the wild type gene promoter and when they are both in their natural environment.

20

The construct may even contain or express a marker which allows for the selection of the genetic construct.

25

For some applications, preferably the construct comprises at least a nucleotide sequence of the present invention or a nucleotide sequence encoding a polypeptide having the specific properties as defined herein operably linked to a promoter.

ORGANISM

30

The term "organism" in relation to the present invention includes any organism that could comprise a nucleotide sequence according to the present invention or a

nucleotide sequence encoding for a polypeptide having the specific properties as defined herein and/or products obtained therefrom.

5 The term "transgenic organism" in relation to the present invention includes any organism that comprises a nucleotide sequence coding for a polypeptide having the specific properties as defined herein and/or the products obtained therefrom, and/or wherein a promoter can allow expression of the nucleotide sequence coding for a polypeptide having the specific properties as defined herein within the organism. Preferably the nucleotide sequence is incorporated in the genome of the organism.

10

Suitable organisms include a prokaryote, fungus yeast or a plant.

The term "transgenic organism" does not cover native nucleotide coding sequences in their natural environment when they are under the control of their native promoter
15 which is also in its natural environment.

Therefore, the transgenic organism of the present invention includes an organism comprising any one of, or combinations of, a nucleotide sequence coding for a polypeptide having the specific properties as defined herein, constructs as defined
20 herein, vectors as defined herein, plasmids as defined herein, cells as defined herein, or the products thereof. For example the transgenic organism can also comprise a nucleotide sequence coding for a polypeptide having the specific properties as defined herein under the control of a promoter not associated with a sequence encoding a lipid acyltransferase in nature.

25

TRANSFORMATION OF HOST CELLS/ORGANISM

The host organism can be a prokaryotic or a eukaryotic organism.

30 Examples of suitable prokaryotic hosts include bacteria such as *E. coli* and *Bacillus licheniformis*, preferably *B. licheniformis*.

Teachings on the transformation of prokaryotic hosts is well documented in the art, for example see Sambrook *et al.* (Molecular Cloning: A Laboratory Manual, 2nd

edition, 1989, Cold Spring Harbor Laboratory Press). If a prokaryotic host is used then the nucleotide sequence may need to be suitably modified before transformation - such as by removal of introns.

5 In another embodiment the transgenic organism can be a yeast.

Filamentous fungi cells may be transformed using various methods known in the art - such as a process involving protoplast formation and transformation of the protoplasts followed by regeneration of the cell wall in a manner known. The use of
10 *Aspergillus* as a host microorganism is described in EP 0 238 023. In one embodiment, preferably *T. reesei* is the host organism.

Another host organism can be a plant. A review of the general techniques used for transforming plants may be found in articles by Potrykus (*Annu Rev Plant Physiol*
15 *Plant Mol Biol* (1991) 42:205-225) and Christou (Agro-Food-Industry Hi-Tech March/April 1994 17-27). Further teachings on plant transformation may be found in EP-A-0449375.

General teachings on the transformation of fungi, yeasts and plants are presented in
20 following sections.

TRANSFORMED FUNGUS

A host organism may be a fungus - such as a filamentous fungus. Examples of
25 suitable such hosts include any member belonging to the genera *Fusarium*, *Thermomyces*, *Acremonium*, *Aspergillus*, *Penicillium*, *Mucor*, *Neurospora*, *Trichoderma* and the like. In one embodiment, *Trichoderma* is the host organism, preferably *T. reesei*.

30 Teachings on transforming filamentous fungi are reviewed in US-A-5741665 which states that standard techniques for transformation of filamentous fungi and culturing the fungi are well known in the art. An extensive review of techniques as applied to *N. crassa* is found, for example in Davis and de Serres, *Methods Enzymol* (1971)
17A: 79-143.

35

Further teachings on transforming filamentous fungi are reviewed in US-A-5674707.

In one aspect, the host organism can be of the genus *Aspergillus*, such as *Aspergillus niger*.

5 A transgenic *Aspergillus* according to the present invention can also be prepared by following, for example, the teachings of Turner G. 1994 (Vectors for genetic manipulation. In: Martinelli S.D., Kinghorn J.R. (Editors) *Aspergillus: 50 years on*. Progress in industrial microbiology vol 29. Elsevier Amsterdam 1994. pp. 641-666).

10 Gene expression in filamentous fungi has been reviewed in Punt *et al. Trends Biotechnol.* (2002); 20(5):200-6, Archer & Peberdy *Crit. Rev. Biotechnol.* (1997) 17:273-306.

TRANSFORMED YEAST

15 In another embodiment, the transgenic organism can be a yeast.

A review of the principles of heterologous gene expression in yeast are provided in, for example, *Methods Mol Biol* (1995), 49:341-54, and *Curr Opin Biotechnol* (1997); 8:554-60.

20 In this regard, yeast - such as the species *Saccharomyces cerevisi* or *Pichia pastoris* or *Hansenula polymorpha* (see *FEMS Microbiol Rev* (2000 24:45-66), may be used as a vehicle for heterologous gene expression.

25 A review of the principles of heterologous gene expression in *Saccharomyces cerevisiae* and secretion of gene products is given by E Hinchcliffe E Kenny (1993, "Yeast as a vehicle for the expression of heterologous genes", *Yeasts*, Vol 5, Anthony H Rose and J. Stuart Harrison, eds, 2nd edition, Academic Press Ltd.).

30 For the transformation of yeast, several transformation protocols have been developed. For example, a transgenic *Saccharomyces* according to the present invention can be prepared by following the teachings of Hinnen *et al.*, (1978, *Proceedings of the National Academy of Sciences of the USA* 75, 1929); Beggs, J D (1978, *Nature*, London, 275, 104); and Ito, H *et al.* (1983, *J Bacteriology* 153, 163-168).

35

The transformed yeast cells may be selected using various selective markers - such as auxotrophic markers dominant antibiotic resistance markers.

5 A suitable yeast host organism can be selected from the biotechnologically relevant yeasts species such as, but not limited to, yeast species selected from *Pichia* spp., *Hansenula* spp., *Kluyveromyces*, *Yarrowinia* spp., *Saccharomyces* spp., including *S. cerevisiae*, or *Schizosaccharomyces* spp., including *Schizosaccharomyces pombe*.

10 A strain of the methylotrophic yeast species *Pichia pastoris* may be used as the host organism.

In one embodiment, the host organism may be a *Hansenula* species, such as *H. polymorpha* (as described in WO 01/39544).

15 TRANSFORMED PLANTS/PLANT CELLS

A host organism suitable for the present invention may be a plant. A review of the general techniques may be found in articles by Potrykus (*Annu Rev Plant Physiol Plant Mol Biol* (1991) 42:205-225) and Christou (*Agro-Food-Industry Hi-Tech* March/April 1994 17-27), or in WO 01/16308. The transgenic plant may produce 20 enhanced levels of phytosterol esters and phytostanol esters, for example.

CULTURING AND PRODUCTION

25 Host cells transformed with the nucleotide sequence of the present invention may be cultured under conditions conducive to the production of the encoded enzyme and which facilitate recovery of the enzyme from the cells and/or culture medium.

30 The medium used to cultivate the cells may be any conventional medium suitable for growing the host cell in question and obtaining expression of the enzyme.

The protein produced by a recombinant cell may be displayed on the surface of the cell.

35 The enzyme may be secreted from the host cells and may conveniently be recovered from the culture medium using well-known procedures.

SECRETION

Often, it is desirable for the polypeptide to be secreted from the expression host into the culture medium from where the enzyme may be more easily recovered.

5 According to the present invention, the secretion leader sequence may be selected on the basis of the desired expression host. Hybrid signal sequences may also be used with the context of the present invention.

10 Typical examples of secretion leader sequences not associated with a nucleotide sequence encoding a lipid acyltransferase in nature are those originating from the fungal amyloglucosidase (AG) gene (*g/aA* - both 18 and 24 amino acid versions *e.g.*, from *Aspergillus*), the *a*-factor gene (yeasts *e.g.*, *Saccharomyces*, *Kluyveromyces* and *Hansenula*) or the *a*-amylase gene (*Bacillus*).

15 DETECTION

A variety of protocols for detecting and measuring the expression of the amino acid sequence are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting
20 (FACS).

A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic and amino acid assays.

25 A number of companies such as Pharmacia Biotech (Piscataway, NJ, USA), Promega (Madison, WI, USA), and US Biochemical Corp (Cleveland, OH, USA) supply commercial kits and protocols for these procedures.

30 Suitable reporter molecules or labels include those radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles and the like. Patents teaching the use of such labels include US-A-3,817,837; US-A-3,850,752; US-A-3,939,350; US-A-3,996,345; US-A-4,277,437; US-A-4,275,149 and US-A-4,366,241 .

35 Also, recombinant immunoglobulins may be produced as shown in US-A-4,816,567.

FUSION PROTEINS

An enzyme for use in the present invention may be produced as a fusion protein, for example to aid in extraction and purification thereof. Examples of fusion protein partners include glutathione-S-transferase (GST), 6xHis, GAL4 (DNA binding and/or transcriptional activation domains) and β -galactosidase. It may also be convenient to include a proteolytic cleavage site between the fusion protein partner and the protein sequence of interest to allow removal of fusion protein sequences. Preferably the fusion protein will not hinder the activity of the protein sequence,

Gene fusion expression systems in *E. coli* have been reviewed in *Curr. Opin. Biotechnol.* (1995) 6:501-6.

The amino acid sequence of a polypeptide having the specific properties as defined herein may be ligated to a non-native sequence to encode a fusion protein. For example, for screening of peptide libraries for agents capable of affecting the substance activity, it may be useful to encode a chimeric substance expressing a non-native epitope that is recognised by a commercially available antibody.

ADDITIONAL POIs

20

The sequences for use according to the present invention may also be used in conjunction with one or more additional proteins of interest (POIs) or nucleotide sequences of interest (NOIs).

Non-limiting examples of POIs include: proteins or enzymes involved in starch metabolism, proteins or enzymes involved in glycogen metabolism, acetyl esterases, aminopeptidases, amylases, arabinases, arabinofuranosidases, carboxypeptidases, catalases, cellulases, chitinases, chymosin, cutinase, deoxyribonucleases, epimerases, esterases, α -galactosidases, β -galactosidases, α -glucanases, glucan lysases, endo-(3-glucanases, glucoamylases, glucose oxidases, α -glucosidases, β -glucosidases, glucuronidases, hemicellulases, hexose oxidases, hydrolases, invertases, isomerases, laccases, lipases, lyases, mannosidases, oxidases, oxidoreductases, pectate lyases, pectin acetyl esterases, pectin depolymerases, pectin methyl esterases, pectinolytic enzymes, peroxidases, phenoloxidases, phytases, polygalacturonases, proteases, rhamno-galacturonases, ribonucleases, thaumatin, transferases, transport proteins, transglutaminases, xylanases, hexose

35

oxidase (D-hexose: O_2 -oxidoreductase, EC 1.1.3.5) or combinations thereof. The NO₂ may even be an antisense sequence for any of those sequences.

5 The POI may even be a fusion protein, for example to aid in extraction and purification.

The POI may even be fused to a secretion sequence.

DETERGENT

10

The compositions of the present invention may form a component of a cleaning and/or detergent composition. In particular, certain embodiments of the present invention may additionally include a detergent.

15 In general, cleaning and detergent compositions are well described in the art and reference is made to WO 96/34946; WO 97/07202; and WO 95/30011 for further description of suitable cleaning and detergent compositions.

20 The compounds of the invention may for example be formulated as a hand or machine laundry detergent composition, including a laundry additive composition suitable for pretreatment of stained fabrics, and a rinse added fabric softener composition, or be formulated as a detergent composition for use in general household hard surface cleaning operations (including car washing or cleaning compositions), or be formulated for hand or machine dishwashing operations. It may
25 also be formulated for use as a personal hygiene product, including but not limited to hand soaps, shampoos and shower gels.

In one embodiment the laundry composition of the present invention may comprise the lipolytic enzyme, hydrophobin and, optionally, detergent in combination with one
30 or more enzymes, such as a protease, a carboxypeptidase, an aminopeptidase, an amylase, a glucoamylase, a maltogenic amylase, a non-maltogenic amylase, an α -galactosidase, a β -galactosidase, an α -glucosidase, a β -glucosidase, a phospholipase, a glycosyltransferase, a chitinase, a cutinase, a carbohydrase, a cellulase, a pectinase, a mannanase, a mannosidase, an arabinase, a galactanase,
35 xylanase, an oxidase, a polyesterase, a laccase, a cyclodextrin esterase, a phytase, a catalase, a haloperoxidase, and/or a peroxidase, a pectinolytic enzyme, a

peptidoglutaminase, a polyphenoloxidase, a transglutaminase, a deoxyribonuclease, a ribonuclease, and/or combinations thereof, in general the properties of the chosen enzyme(s) should be compatible with the selected detergent, (e.g., phi-optimum, compatibility with other enzymatic and non-enzymatic ingredients, etc.), and the enzyme(s) should be present in effective amounts.

Proteases: suitable proteases include those of animal, vegetable or microbial origin. Chemically modified or protein engineered mutants are also suitable. The protease may be a serine protease or a metalloprotease, e.g., an alkaline microbial protease or a trypsin-like protease. Examples of alkaline proteases are subtilisins, especially those derived from *Bacillus* sp., e.g., subtilisin Novo, subtilisin Carlsberg, subtilisin 309 (see, e.g., U.S. Patent No. 6,287,841), subtilisin 147, and subtilisin 168 (see, e.g., WO 89/06279). Examples of trypsin-like proteases are trypsin (e.g., of porcine or bovine origin), and *Fusarium* proteases (see, e.g., WO 89/06270 and WO 94/25583). Examples of useful proteases also include but are not limited to the variants described in WO 92/19729 and WO 98/20115. Suitable commercially available protease enzymes include ALCALASE®, SAVINASE®, LIQUANASE®, OVOZYME®, POLARZYME®, ESPERASE®, EVERLASE®, and KANNASE® (Novozymes, formerly Novo Nordisk A/S); EXCELLASE™, MAXATASE®, MAXACAL™, MAXAPEM™, PROPERASE®, PROPERASE L®, PURAFECT®, PURAFECT L®, PURAFAST™, OXP™, FN2™, and FN3™ (Genencor - a division of Danisco A/S).

Polyesterases: Suitable polyesterases include, but are not limited to, those described in WO 01/34899 (Genencor) and WO 01/14629 (Genencor), and can be included in any combination with other enzymes discussed herein.

Amylases: The compositions can comprise amylases such as α -amylases (EC 3.2.1.1), G4-forming amylases (EC 3.2.1.60), β -amylases (EC 3.2.1.2) and γ -amylases (EC 3.2.1.3). These can include amylases of bacterial or fungal origin, chemically modified or protein engineered mutants are included. Commercially available amylases, such as, but not limited to, DURAMYL®, TERMAMYL™, FUNGAMYL® and BAN™ (Novozymes, formerly Novo Nordisk A/S), RAPIDASE®, and PURASTAR® (Danisco USA, Inc.), LIQUEZYME™, NATALASE™, SUPRAMYL™, STAINZYME™, FUNGAMYL and BAN™ (Novozymes A/S), RAPIDASE™, PURASTAR™, PURASTAROXAM™ and POWERASE™ (from

Danisco USA Inc.), GRINDAMYL™ PowerFresh, POWERFlex™ and GRINDAMYL PowerSoft (from Danisco A/S).

5 Peroxidases/Oxidases: Suitable peroxidases/oxidases contemplated for use in the compositions include those of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from *Coprinus*, e.g., from *C. cinereus*, and variants thereof as those described in WO 93/24618, WO 95/10602, and WO 98/15257. Commercially available peroxidases include GUARDZYME® (Novozymes A/S).

10

Cellulases: Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Suitable cellulases include cellulases from the genera *Bacillus*, *Pseudomonas*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*, e.g., the fungal cellulases produced from *Humicola insolens*,
15 *Myceliophthora thermophila* and *Fusarium oxysporum* disclosed in U.S. Patent Nos. 4,435,307; 5,648,263; 5,691,178; 5,776,757; and WO 89/09259, for example. Exemplary cellulases contemplated for use are those having colour care benefit for the textile. Examples of such cellulases are cellulases described in EP 0495257; EP531372; WO 99/25846 (Genencor International, Inc.), WO 96/34108 (Genencor
20 International, Inc.), WO 96/11262; WO 96/29397; and WO 98/08940, for example. Other examples are cellulase variants, such as those described in WO 94/07998; WO 98/12307; WO 95/24471; WO 99/01544; EP 531 315; U.S. Patent Nos. 5,457,046; 5,686,593; and 5,763,254. Commercially available cellulases include CELLUZYME®, CAREZYME® and ENDOLASE® (Novozymes, formerly Novo
25 Nordisk A/S); CLAZINASE™ and PURADAX® HA (Genencor); and KAC-50Q(B)™ (Kao Corporation).

30

Examples of commercially available mannanases include MANNAWAY™ (Novozymes, Denmark) and MANNASTAR™ (Genencor).

35

The composition of the invention can be formulated as either a solid or a liquid. Examples of formulations include granulates, pellets, slurries, bars, pastes, foams, gels, strips, etc. Preferred detergent additive formulations are granulates, in particular non-dusting granulates, liquids, in particular stabilized liquids, or slurries. A liquid
detergent may be aqueous, typically containing up to 70% water and 0-30% organic solvent, or non-aqueous.

Non-dusting granulates may be produced, e. g., as disclosed in US 4,106,991 and 4,661,452 and may optionally be coated by methods known in the art. Examples of waxy coating materials are poly (ethylene oxide) products (polyethylene glycol, PEG) with mean molar weights of 1000 to 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB 1483591. Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid according to established methods. Protected enzymes may be prepared according to the method disclosed in EP-A-238216.

The detergent composition may also comprise one or more further surfactants, which may be non-ionic including semi-polar and/or anionic and/or cationic and/or zwitterionic. The surfactants are typically present at a level of from 0.1% to 60% by weight.

When included therein the detergent will usually contain from about 1% to about 40% of an anionic surfactant such as linear alkylbenzenesulfonate, alpha-olefinsulfonate, alkyl sulfate (fatty alcohol sulfate), alcohol ethoxysulfate, secondary alkanesulfonate, alpha-sulfo fatty acid methyl ester, alkyl- or alkenylsuccinic acid or soap.

When included therein the detergent will usually contain from about 0.2% to about 40% of a non-ionic surfactant such as alcohol ethoxylate, nonylphenol ethoxylate, alkylpolyglycoside, alkyldimethylamineoxide, ethoxylated fatty acid monoethanolamide, fatty acid monoethanolamide, polyhydroxy alkyl fatty acid amide, or other N-acyl or N-alkyl derivatives of glucosamine.

The detergent may contain 0-65% of a detergent builder or complexing agent such as zeolite, diphosphate, triphosphate, phosphonate, carbonate, citrate, nitrilotriacetic acid, ethylenediaminetetraacetic acid, diethylenetriaminepentaacetic acid, alkyl- or alkenylsuccinic acid, soluble silicates or layered silicates (e. g. SKS-6 from Hoechst).

The detergent may comprise one or more polymers. Examples are carboxymethylcellulose, poly (vinylpyrrolidone), poly (ethylene glycol), poly (vinyl alcohol), poly (vinylpyridine-N-oxide), poly (vinylimidazole), polycarboxylates such as

polyacrylates, maleic/acrylic acid copolymers and lauryl methacrylate / acrylic acid copolymers.

5 The detergent may contain a bleaching system which may comprise a hydrogen peroxide source such as perborate or percarbonate which may be combined with a peracid-forming bleach activator such as tetraacetythyenediamine or nonanoyloxybenzenesulfonate. Alternatively, the bleaching system may comprise peroxyacids of e.g., the amide, imide, or sulfone type.

10 The enzyme(s) of the detergent composition of the invention may be stabilized using conventional stabilizing agents, e. g., a polyol such as propylene glycol or glycerol, a sugar or sugar alcohol, lactic acid, boric acid, or a boric acid derivative, e. g., an aromatic borate ester, or a phenyl boronic acid derivative such as 4-formylphenyl boronic acid, and the composition may be formulated as described in e.g., WO
15 92/1 9709 and WO 92/1 9708.

The detergent may also contain other conventional detergent ingredients such as fabric conditioners including clays, foam boosters, suds suppressors, anti-corrosion agents, soil-suspending agents, anti-soil redeposition agents, dyes, bactericides,
20 optical brighteners, hydrotropes, tarnish inhibitors, or perfumes.

DOSAGE

In the compositions of the present invention, the hydrophobin may be present in any
25 concentration sufficient to enable it to exhibit the effects described herein. Suitably, the hydrophobin is present in a concentration of between 0.001% and 5%, preferably 0.002% to 2.5%, more preferably 0.005% to 1%, even more preferably 0.01% to 0.5% by weight of the total weight of the composition. In particularly preferred examples, the hydrophobin is present in a concentration of 0.01 , 0.05, 0.1 , 0.25 or
30 0.4% by weight of the total weight of the composition.

In the compositions of the present invention, the lipolytic enzyme may be present in any concentration sufficient to enable it to exhibit the effects described herein.

35 Suitably, the lipolytic enzyme is present in a concentration of 0.001 to 400 ppm, preferably 0.002 to 200 ppm, more preferably 0.005 to 100 ppm, even more

preferably 0.01 to 50 ppm, still more preferably 0.02 to 25 ppm, of pure enzyme protein by weight of the total weight of the composition.

Suitably, the lipolytic enzyme is present in a concentration of 0.025 to 25, preferably 5 0.05 to 10, more preferably 0.1 to 5, units of enzyme activity per g of the composition. The activity is measured according to the trioctanoate assay described below, wherein 1 unit of activity represents 1 μmol of the free fatty acid produced by 1 g of enzyme solution in 1 minute.

10 Where the compositions of the present invention include a detergent, the detergent may be present in any concentration sufficient to enable it to exhibit the effects described herein. Suitably, the detergent is present in a concentration of between 0.001 and 20 g/L, preferably 0.01 to 10 g/L, more preferably 0.05 to 5 g/L, even more preferably 0.1 to 2.5 g/L by Do the litres refer to the volume of the washing solution!n
15 particularly preferred examples, the detergent is present in a concentration of 0.01, 0.05, 0.1, 0.25 or 0.4 g/L of the washing solution.

Trioctanoate assay

20 Reaction emulsions of trioctanoate in the compositions was prepared from 0.4% trioctanoate pre-suspended in ethanol (5%), in one of two buffers: 0.05M 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) adjusted to pH 8.2, or 0.05M N-cyclohexyl-3-aminopropanesulfonic acid (CAPS) adjusted to pH 10. For both buffers water hardness adjusted to 240 ppm. The final assay mixtures contained
25 varying amounts of detergents, to aid in the emulsification of the triglyceride.

The reaction emulsions were made by applying high shear mixing for 2 minutes (24000 m^{-1} , Ultra Turrax T25, Janke & Kunkel), and then transferring 150 μL to 96-well microtiter plate wells already containing 30 μL enzyme samples. Free fatty acid generation was measured using an *in vitro* enzymatic colorimetric assay for the
30 quantitative determination of non-esterified fatty acids (NEFA). This method is specific for free fatty acids, and relies upon the acylation of coenzyme A (CoA) by the fatty acids in the presence of added acyl-CoA synthetase. The acyl-CoA thus produced is oxidized by added acyl-CoA oxidase with generation of hydrogen peroxide, in the presence of peroxidase. This permits the oxidative condensation of
35 3-methyl -N-ethyl -N-(P-hydroxyethyl)-aniline with 4-aminoantipyrine to form a purple colored adduct which can be measured colorimetrically. The amount of free fatty

acids generated after a 6 minute incubation at 30°C was determined using the materials in a NEFA HR(2) kit (Wako Chemicals GmbH, Germany) by transferring 30 µL of the hydrolysis solution to 96-well microtiter plate wells already containing 120 µL NEFA A solution. Incubation for 3 min at 30°C was followed by addition of 60 µL NEFA B solution. After incubation for 4.5 min at 30°C OD at 520 nm was measured.

LAUNDRY COMPOSITIONS

The hydrophobins used in the present invention may be generated *in situ* in a laundry composition, for example by hydrolysis of hydrophobin precursor (such as a hydrophobin fusion protein) in the laundry composition.

The hydrophobin precursor (such as a hydrophobin fusion protein) is required in order to generate *in situ* the hydrophobins used in the present invention. It may be present as an initial component of the laundry composition. Alternatively, if no or insufficient hydrophobin precursor is initially present, this component can be added to the composition.

If required, a catalyst (particularly an enzyme, especially a protease enzyme) may be present. It may be present as an initial component of the laundry composition. Alternatively, if no or insufficient catalyst is initially present, this component can be added to the composition.

The laundry composition may further comprise a stain, which may be a lipid (in particular, a triglyceride and/or a diglyceride and/or a monoglyceride). The stain may be on a surface, for example a fabric. The laundry composition of the present invention may therefore comprise a surface for example a fabric.

Converting a hydrophobin precursor into a hydrophobin used in the present invention may help remove a stain comprising a lipid from a fabric.

CLEANING METHODS

The present invention further comprises a method of removing a lipid-based stain from a surface by contacting the surface with a composition according to the invention. In addition, the present invention comprises a method of cleaning a

surface, comprising contacting the surface with a composition according to the invention. Furthermore, the present invention comprises a method of cleaning an item (particularly although not exclusively a clothing item or a tableware item), comprising contacting the item with a composition according to the invention.

5

In another aspect, methods for removing oily stains from fabrics are provided. The methods generally involve identifying fabrics having oily stains, contacting the fabrics with a composition of the invention, and rinsing the fabric to remove the oily stain from the fabrics.

10

In some embodiments, the lipolytic enzyme, the hydrophobin and, optionally, the detergent are present together in a single composition. In some embodiments, the lipolytic enzyme, the hydrophobin and, optionally, the detergent are separate in different compositions that are combined prior to contacting the fabric, or mixed together on the fabric. Therefore, application of the lipase and the adjuvant may be simultaneous or sequential. In some embodiments, the contacting occurs in a wash pretreatment step, *i.e.*, prior to hand or machine-washing a fabric. In some embodiments, the contacting occurs at the time of hand or machine-washing the fabric. The contacting may occur as a result of mixing the present compositions with wash water, spraying, pouring, or dripping the composition on the fabric, or applying the composition using an applicator.

15

20

The methods are effective for removing a variety of oil stains, or portions of oily stains, which typically include esters of fatty acids, such as triglycerides.

25

It will be appreciated that rinsing may occur some time after the washing, and that in some aspects the present method of cleaning is essentially complete following the contacting of the fabric with the composition.

30

FOODSTUFF

The compositions of the present invention may be used as a component of a foodstuff. The term "foodstuff" as used herein means a substance which is suitable for human and/or animal consumption.

35

Suitably, the term "foodstuff" as used herein may mean a foodstuff in a form which is ready for consumption. Alternatively or in addition, however, the term foodstuff as

used herein may mean one or more food materials which are used in the preparation of a foodstuff. By way of example only, the term foodstuff encompasses both baked goods produced from dough as well as the dough used in the preparation of said baked goods.

5

The foodstuff may be in the form of a solution or as a solid - depending on the use and/or the mode of application and/or the mode of administration.

When used as - or in the preparation of - a food - such as functional food - the composition of the present invention may be used in conjunction **with** one or more of: a nutritionally acceptable carrier, a nutritionally acceptable diluent, a nutritionally acceptable excipient, a nutritionally acceptable adjuvant, a nutritionally active ingredient.

15 In a preferred aspect the present invention provides a foodstuff as defined above wherein the foodstuff is selected from one or more of the following: eggs, egg-based products, including but not limited to mayonnaise, salad dressings, sauces, ice
creams, egg powder, modified egg yolk and products made therefrom; baked goods, including breads, cakes, sweet dough products, laminated doughs, liquid batters,
20 muffins, doughnuts, biscuits, crackers and cookies; confectionery, including chocolate, candies, caramels, halawa, gums, including sugar free and sugar sweetened gums, bubble gum, soft bubble gum, chewing gum and puddings; frozen products including sorbets, preferably frozen dairy products, including ice cream and ice **milk**; dairy products, including cheese, butter, milk, coffee cream, whipped cream,
25 custard cream, milk drinks and yoghurts; mousses, whipped vegetable creams, meat products, including processed meat products; edible oils and fats, aerated and non-aerated whipped products, oil-in-water emulsions, water-in-oil emulsions, margarine, shortening and spreads including low fat and very low fat spreads; dressings, mayonnaise, dips, cream based sauces, cream based soups, beverages, spice
30 emulsions and sauces.

Suitably the foodstuff in accordance with the present invention may be a "fine food", including cakes, pastry, confectionery, chocolates, fudge and the like.

35 In one aspect the foodstuff in accordance with the present invention may be a dough product or a baked product, such as bread, a fried product, a snack, cakes, pies,

brownies, cookies, noodles, snack items such as crackers, graham crackers, pretzels, and potato chips, and pasta.

5 in another aspect the foodstuff in accordance with the present invention may be a convenience food, such as a part-baked or part-cooked product. Examples of such part-baked or part-cooked product include part-baked versions of the dough and baked products described above.

10 in a further aspect, the foodstuff in accordance with the present invention may be a plant derived food product such as flours, pre-mixes, oils, fats, cocoa butter, coffee whitener, salad dressings, margarine, spreads, peanut butter, shortenings, ice cream, cooking oils.

15 In another aspect, the foodstuff in accordance with the present invention may be a dairy product, including butter, milk, cream, cheese such as natural, processed, and imitation cheeses in a variety of forms (including shredded, block, slices or grated), cream cheese, ice cream, frozen desserts, yoghurt, yoghurt drinks, butter fat, anhydrous milk fat, other dairy products. The enzyme according to the present invention may improve fat stability in dairy products.

20

In another aspect, the foodstuff in accordance with the present invention may be a food product containing animal derived ingredients, such as processed meat products, cooking oils, shortenings.

25 In a further aspect, the foodstuff in accordance with the present invention may be a beverage, a fruit, mixed fruit, a vegetable, a marinade or wine.

In one aspect, the foodstuff in accordance with the present invention is a plant derived oil (*i.e.* a vegetable oil), such as olive oil, sunflower oil, peanut oil or rapeseed oil. The oil may be a degummed oil.

30

EXAMPLES

EXAMPLE 1

5

The following experiments were carried out to test whether the cleaning performance of a lipase is enhanced by adding hydrophobin in the presence or absence of commercially available heat inactivated detergent.

- 10 The lipases used were as follows (each dosed in a single dose): -
LIPEX™ (abH23.1, fungal) (SEQ ID NO: 11) (commercially available from
Novozymes A/S), 1.25 mg in 1 mL
LIPOMAX™ (abH15.2, family I-1) (SEQ ID NO: 15) (commercially available from
Danisco A/S), 6 mg in 1 mL
15 SprLip2 (abH16, family I-7) (SEQ ID NO: 17), 258 µL in 1 mL
TfuLip2 (abH25.1, family III) (SEQ ID NO: 16), 30.8 pL in 1 mL

The hydrophobin used was hydrophobin HFBII (SEQ ID NO: 2; obtainable from the
fungus *Trichoderma reesei*). 26.6 g HFBII (containing 150 mg/g hydrophobin protein)
20 was dissolved in 100 mL water to give a solution containing 40 g/L hydrophobin
protein. The solution was diluted as appropriate to give a hydrophobin dose of 0.01,
0.05, 0.1, 0.25 and 0.40% by weight of the total weight of the composition.

The detergents used were heat inactivated liquid detergent (ARIEL™ colour liquid)
25 and heat inactivated powder detergent (ARIEL™ colour powder). These are
commercially available from Procter & Gamble. The detergents were diluted as
appropriate to give a dose of 0, 0.1, 0.25 and 0.4 g/L.

The detergents were heat-inactivated as follows: the liquid detergents were placed in
30 a water bath at 95°C for 2 hours, while 0.1 g/mL preparations in water of the powder
detergents were boiled on a hot plate for 1 hour. Heat treatments inactivate the
enzymatic activity of any protein components in commercial detergent formulas,
while retaining the properties of the nonenzymatic detergent components. Following
heating, the detergents are diluted and assayed for lipase enzyme activity.

35

Cleaning performance of lipase and hydrophobin on stained fabrics was tested in a
microswatch assay format. Stain removal experiments were carried out using a lipid-

containing technical stain (CS-61 swatches: cotton, beef fat with colorant, purchased from Center for Testmaterials, Netherlands) set in a 24-well plate format (Nunc, Denmark). Each assay well was set to contain a pre-cut 13 mm piece of CS-61 swatch. Swatches were pre-read using a scanner (Microtek Scan Maker 900) and placed in the 24-well plate.

The buffers used were 20 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) (0.2M, pH 8.2) for testing liquid detergents, and 20 mM *N*-cyclohexyl-3-aminopropanesulfonic acid (CAPS) (0.2M, pH 10.0) for testing powder detergents. Wafer hardness was adjusted to 24 degrees French (FH - one degree French is defined as 10 milligrams of calcium carbonate per litre of water) using 15000 ppm 2/1 Ca²⁺/Mg²⁺ diluted to 2400 ppm (dilution factor 6.25) for both buffers.

A 24 well plate was used, each well containing 1 ml solution. The hydrophobin concentration in each row was as follows: zero; 0.01%; 0.05%; 0.1%; 0.25%; and 0.4% by weight of the total weight of the composition. The detergent concentration in each column was as follows: zero; 0.1 g/L; 0.25 g/L; and 0.4 g/L.

900 μ L of the appropriate buffer described above was added to each swatch-containing well of the 24-well plate. 100 μ L hydrophobin solution was added into each well. To initiate the reaction, enzyme samples were added at a volume of 100 μ L into each well. The plates were shaken for 30 minutes at 200 rpm at 37°C. After incubation, the reaction buffer was removed and the fabric in each well was rinsed with 1 mL distilled water three times. After removing the rinse the swatches were dried at 50°C for 4 hours and reflectance was measured. Cleaning performance was quantified after a single wash cycle. Stain removal was calculated as the difference of the post- and pre-cleaning RGB colour measurements for each swatch. RGB measurements were taken with a scanner (Microtek Scan Maker 900).

The difference in Stain Removal Index (ASRI) values of the washed fabric were calculated in relation to the unwashed fabrics using the formula:

$$\% \text{ Soil Removal}_{(\text{RGB})} = (\text{Soil removal } \Delta E_{(\text{RGB})} / \text{Initial soil } \Delta E_{(\text{RGB})}) \times 100\%$$

Where:

$$\text{Soil removal } \Delta E_{(\text{RGB})} = \sqrt{((R_{\text{after}} - R_{\text{before}})^2 + (G_{\text{after}} - G_{\text{before}})^2 + (B_{\text{after}} - B_{\text{before}})^2)}$$

And:

$$\text{Initial soil } \Delta E_{(\text{RGB})} = \sqrt{((R_{\text{ref}} - R_{\text{before}})^2 + (G_{\text{ref}} - G_{\text{before}})^2 + (B_{\text{ref}} - B_{\text{before}})^2)}$$

RGB_{ref} values are the values of the unsoiled cotton (white).

The results are shown in Figures (Figs). 1a through 5e, as follows:

- 5 Figs. 1a through 1c: no lipolytic enzyme (control)
- Figs. 2a through 2e: the lipolytic enzyme LIPEX™ (abH23.1)
- Figs. 3a through 3e: the lipolytic enzyme LIPOMAX™ (abH15.2)
- Figs. 4a through 4e: the lipolytic enzyme SprLip2 (abH16)
- Figs. 5a through 5e: the lipolytic enzyme TfuLip2 (abK25.1)

10

In particular, Figs. 2e, 4e and 5e illustrate the effects of hydrophobin on the presence of lipases in the system in the absence of a detergent. These Figures show that, for these lipases at least, a synergistic effect superior to the additive effect of each component when used individually can be observed.

15

In addition, Figure 2b illustrates that, when a combination of hydrophobin, the lipase LIPEX® and the detergent ARIEL® Color Liquid is used, as the concentration of the detergent increases, the system reaches a performance plateau at lower concentrations of hydrophobin (0.05% instead of 0.4%) compared with when no detergent is used. Furthermore, Figure 5b shows that, using a combination of hydrophobin, the lipase TfuLip2 and the detergent ARIEL® Color Liquid, by increasing the concentration of detergent and the concentration of hydrophobin, an improved washing effect can be achieved (in particular with 0.4 g/L detergent and 0.4% hydrophobin).

25

In addition, Figure 2d illustrates that, when a combination of hydrophobin, the lipase LIPEX® and the detergent ARIEL® Color Powder is used, the performance pattern is not affected by lower levels of detergent (the system reaches plateau at 0.05% hydrophobin). However, at higher concentrations of the detergent, the higher SRI value can be reached (30% at 0.4 g/L detergent). Furthermore, Figure 5d illustrates that, when a combination of hydrophobin, the lipase TfuLip2 and the detergent ARIEL® Color Powder is used, the overall performance of the system improves with increase of the concentration of detergent in the system.

35

Finally, Figure 1b shows that, when a combination of hydrophobin and the detergent ARIEL® Color Liquid is used in the absence of lipases, there is a small synergistic

effect at low concentrations of hydrophobin (0.01-0.1%) and detergent (below 0.25 g/L).

EXAMPLE 2 - Cloning and expression of *Streptomyces pristinaespiralis* ATCC 2548 lipase (SprUp2)

The SprLip2 gene was synthesized by GeneRay (Shanghai, China). The SprLip2 synthetic gene was cloned into expression plasmid pKB128 by NheI/BamHI double digestion and ligation. Plasmid pKB128 is a derivative of plasmid pKB105 (described in U.S. Patent Application Publication No. 2006/0154843) and is the source of the A4 promoter-CeIA signal sequence. Plasmid pKB128 contains the NsiI-MluI-HpaI restriction sites (atgcataegcgtgtaac; SEQ ID No 30) before the BamHI site. The *A. niger* A4 promoter and the CeIA truncated signal sequences were at the 5' end of the SprLip2 gene sequence (corresponding to the predicted mature protein), and the 11AGS terminator sequence was fused to the 3' end of the SprLip2 gene sequence. The pZQ205 expression vector (Figure 30) was constructed by ligation of pKB128 after digestion with the restriction enzymes NheI and BamHI, to a similarly digested SprLip2 synthetic gene, followed by transformation of *E. coli* cells. The correct sequence of SprLip2 gene was confirmed by DNA sequencing.

Plasmid DNA of pZQ205 was transformed into host *Streptomyces lividans* TK23 protoplast cells (described in U.S. Patent Application Publication No. 2006/0154843). Three transformants were picked and transferred into a seed shake flask (15 ml of TSG medium containing 50 ug/ml of thiostrepton in dimethyl sulfoxide), grown for 2 days at 30°C with shaking at 200 rpm. 3 ml of the two-day culture from seed shake flask were transferred to 30 ml of *Streptomyces* modified production medium II for protein production. The production cultures were grown for 2 days at 30°C with shaking at 200 rpm. The protein was secreted into the extracellular medium and filtered culture medium was used to perform the cleaning assay and for biochemical characterization experiments. The dosing was based on total protein determined by a Bradford type assay using the Biorad protein assay (500-0006EDU) and corrected for purity determined by SDS-PAGE using a Criterion stain free system from Bio-Rad.

EXAMPLE 3 - Biochemical characterization of SprLip2

The lipase/esterase activity of SprLip2 was tested using para-nitrophenyl butyrate ester (pNB) and para-nitrophenyl palmitate (pNPP) as substrates. A 20mM stock

solution of each substrate (p-nitrophenyl butyrate, pNB, Sigma, CAS 2635-84-9, catalog number N9876) dissolved in dimethyl sulfoxide (Pierce, 20688, Water content <0.2%) and p-nitrophenyl palmitate, pNPP; Sigma, CAS 1492-30-4, catalog number N2752 dissolved in dimethyl sulfoxide) was prepared and stored at -80°C for long term storage. Filtered culture supernatant from SprLip2 expressing cells was serially diluted in assay buffer [50mM HEPES pH 8.2, containing 0.75 mM CaCl₂ and 0.25mM MgCl₂) containing 2% Polyvinyl Alcohol (PVA) (Sigma)] in 96-well microtiter plates and equilibrated at 25°C. 100 μl of 1:20 diluted substrate (in assay buffer) was added to another microtiter plate. The plate was equilibrated to 25°C for 10 minutes with shaking at 500rpm. 10 μl of enzyme solution from dilution plate was added to the substrate containing plate to initiate reaction. The plate was immediately transferred to a spectrophotometer capable of kinetic measurements equilibrated at 25°C. The absorbance change in kinetic mode was read for 5 minutes at 410nm. The background rate (with no enzyme) was subtracted from the rate of the test samples. Sample concentration was determined as:

$$\text{Sample concentration} = (\text{unknown Rate} \times \text{standard concentration}) / \text{standard rate}$$

Results are shown in Figures 32 (pNB hydrolysis) and 33 (pNPP hydrolysis). (relative rates of hydrolysis.)]

EXAMPLE 4 - Triglyceride hydrolysis by SprLip2

This assay was designed to measure release of fatty acids from triglyceride substrate by lipases. The assay consists of a hydrolysis reaction where incubation of lipase with a triglyceride emulsion results in liberation of fatty acids and thus a reduction in the turbidity of the emulsified substrate. The triglyceride substrate used for the assay was glyceryl trioctanoate (Sigma, CAS 538-23-8, catalog number T9126-100ML). Emulsified trioctanoate (0.75% (v/v or w/v)) was prepared by mixing 50 ml of the gum arabic (Sigma, CAS 9000-01-5, catalog number G9752; 10 mg/ml gum arabic solution made in 50 mM HEPES pH8.2) or detergent solution (0.1% heat inactivated Tide® Cold Water detergent, Procter & Gamble, Cincinnati, OH, USA, (containing 0.75 mM CaCl₂ and 0.25mM MgCl₂) in 50 mM HEPES pH8.2) with 375 μl of triglyceride. The solutions were mixed and sonicated for at least 2 minutes to prepare a stable emulsion. 200 μl of emulsified substrate was added to a 96-well microtiter plate. 20 μl of serially diluted enzyme sample (filtered culture supernatant from cells expressing SprLip2) were added to the substrate containing plate. The plate was

covered with a plate sealer and incubated at 20°C for 20 minutes. After incubation, the presence of fatty acids in solution was detected as increase in absorbance at 550nm using the HR Series NEFA-HR (2) NEF.A kit (Wako Chemicals GmbH, Germany) as indicated by the manufacturer. Results are shown in Figures 34 (no detergent) and 35 (with detergent).

EXAMPLE 5 - Cleaning performance of SprLip2

The cleaning performance of SprUp2 was tested in the presence and absence of commercially available heat inactivated detergents. Stock solution of lipase was prepared by diluting 258 μ l of the enzyme into 1 ml by distilled water. The detergents used were heat inactivated liquid detergent (ARIEL™ color liquid) and heat inactivated powder detergent (ARIEL™ color powder) from Procter & Gamble, Cincinnati, OH, USA.

Stain removal experiments were carried out using a lipid-containing technical stain (CS-61 swatches: cotton, beef fat with colorant, purchased from Center for Testmaterials, Netherlands) in a 24-well plate format (Nunc, Denmark). Each assay well was set to contain a pre-cut 13 mm piece of CS-61 swatch. Swatches were pre-read using a scanner (Microtek Scan Maker 900) and placed in the 24-well plate. The buffers used were 20 mM HEPES pH 8.2 for liquid detergent and 20mM CAPS pH 10.0 for powder detergent. Water hardness was adjusted to 24 degrees French using 15000 ppm 2/1 Ca^{2+}/Mg^{2+} diluted to 2400 ppm for both buffers. The detergents were tested at a concentration of zero; 0.1 g/L; 0.25 g/L; and 0.4 g/L. 1 ml of the appropriate buffer described above was added to each swatch-containing well of the 24-well plate. To initiate the reaction, enzyme samples were added at a volume of 100 μ L into each well. The plates were shaken for 30 minutes at 200 rpm at 37°C. After incubation, the reaction buffer was removed and the fabric in each well was rinsed three times with 1 mL distilled water. The rinsed swatches were dried at 50°C for 4 hours and their reflectance was measured. Cleaning performance was quantified after a single wash cycle. Stain removal was calculated as the difference of the post- and pre-cleaning RGB measurements for each swatch. RGB measurements were taken with a scanner (Microtek Scan Maker 900). Stain Removal Index values (SRI) of the washed fabric were calculated in relation to the unwashed fabrics using the formula:

$$\% \text{ Soil Removal(RGB)} = (\text{Soil removal } A E_{(RGB)} / \text{Initial soil } \Delta E_{(RGB)}) \times 100\%$$

Where:

$$\text{Soil removal } \Delta E_{(RGB)} = \sqrt{\left((R_{after} - R_{before})^2 + (G_{after} - G_{before})^2 + (B_{after} - B_{before})^2 \right)}$$

And:

$$\text{Initial soil } \Delta E_{(RGB)} = \sqrt{\left((R_{ref} - R_{before})^2 + (G_{ref} - G_{before})^2 + (B_{ref} - B_{before})^2 \right)}$$

- 5 RGB_{ref} values are the values of the unsoiled cotton (white).

Results are shown in Figure 36.


All publications mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described methods and
 10 system of the present invention will be apparent to those skilled in the art without departing from the scope and spirit of the present invention. Although the present invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes
 15 for carrying out the invention which are obvious to those skilled in chemistry, biochemistry and biotechnology or related fields are intended to be within the scope of the following claims.

CLAIMS

1. A composition comprising:

(a) a lipolytic enzyme; and

(fa) a hydrophobin having the general formula (I):



wherein:

m and n are independently 0 to 2000;

B₁, B₂, B₃, B₄, B₅, B₆, B₇ and B₈ are each independently amino acids selected from Cys, Leu, Ala, Pro, Ser, Thr, Met or Gly, at least 6 of the residues B₁ through B₈ being Cys;

X₁, X₂, X₃, X₄, X₅, X₆, X₇, Y₁ and Y₂ independently represent any amino acid;

a is 1 to 50;

b is 0 to 5;

c is 1 to 100;

d is 1 to 100;

e is 1 to 50;

f is 0 to 5; and

g is 1 to 100.

2. A composition according to claim 1, wherein the lipolytic enzyme has triacylglycerol hydrolysing activity (E.G. 3.1.1.3).


3. A composition according to claim 1 or claim 2, wherein the lipolytic enzyme is a GX lipolytic enzyme, wherein G is glycine and X is an oxyanion hole-forming amino acid residue, wherein the GX lipolytic enzyme belongs to an alpha/beta hydrolase superfamily selected from the group consisting of abH23, abH25, and abH15.

4. A composition according to any one of claims 1 to 3, additionally comprising:
(c) a detergent.

5. A composition comprising:

(a) a GX lipolytic enzyme, wherein G is glycine and X is an oxyanion hole-forming amino acid residue;

(b) a hydrophobin having the general formula (I):



wherein:

m and n are independently 0 to 2000;

B₁, **B**₂, **B**₃, **B**₄, **B**₅, **B**₆, **B**₇ and **B**₈ are each independently amino acids selected from Cys, Leu, Ala, Pro, Ser, Thr, Met or Gly, at least 6 of the residues **B**₁ through **B**₈ being Cys;

X₁, **X**₂, **X**₃, **X**₄, **X**₅, **X**₆, **X**₇, **Y**₁ and **Y**₂ independently represent any amino acid;

a is 1 to 50;

b is 0 to 5;

c is 1 to 100;

d is 1 to 100;

e is 1 to 50;

f is 0 to 5; and

g is 1 to 100; and

(c) a detergent.

6. A composition according to claim 5, wherein the GX lipolytic enzyme belongs to an alpha/beta hydrolase superfamily selected from the group consisting of abH23, abH25, abH16 and abH15.
7. A composition according to claim 6, wherein the GX lipolytic enzyme belongs to an alpha/beta hydrolase superfamily selected from the group consisting of abH23.01, abH 25.01, abH16.01 and abH15.02.
8. A composition according to any one of claims 3 to 7, wherein the oxyanion hole forming residue X is selected from the group consisting of M, Q, F, S, T, A, L and I.
9. A composition according to claim 3 or claim 4, wherein the GX lipolytic enzyme belongs to an alpha/beta hydrolase superfamily selected from the group consisting of abH23.01, abH 25.01 and abH15.02.

10. A composition according to any one of claims 3 to 9, wherein the GX lipolytic enzyme is obtained or obtainable from a filamentous fungus.
11. A composition according to any one of claims 3 to 10, wherein the GX lipolytic enzyme belongs to the *Rhizopus meihei* like homologous family abH23.01 .
12. A composition according to any one of claims 3 to 11, wherein the GX lipolytic enzyme is classified in homologous family abH23.01 and is obtained or obtainable from a fungus of a genus selected from the group consisting of *Thermomyces*, *Fusarium*, *Aspergillus* and *Rhizopus*.
13. A composition according to claim 12, wherein the GX lipolytic enzyme is classified in homologous family abH23.01 and is obtained or obtainable from a fungal species selected from the group consisting of *Thermomyces lanuginosus*, *Fusarium heterosporum*, *Aspergillus tubiengisis*, *Aspergillus fumigatus* and *Rhizopus arrizus*.
14. A composition according to any preceding claim, wherein the lipolytic enzyme is present in a concentration of 0.001 to 20 ppm by weight of the total weight of the composition.
15. A composition according to any preceding claim, wherein the lipolytic enzyme is present in a concentration of 0.01 to 2 ppm by weight of the total weight of the composition.
16. A composition according to any preceding claim, wherein the hydrophobin has a sequence of between 40 and 120 amino acids in the hydrophobin core.
17. A composition according to any preceding claim, wherein the hydrophobin has the general formula (II):



wherein:

m and n are independently 0 to 20;

B_1 , B_2 , B_3 , B_4 , B_5 , B_6 , B_7 and B_8 are each independently amino acids selected from Cys, Leu, Ala, Pro, Ser, Thr, Met or Gly, at least 7 of the residues B_1 through B_8 being Cys;

a is 3 to 25;

b is 0 to 2;

c is 5 to 50;


d is 2 to 35;

e is 2 to 15;

f is 0 to 2; and

g is 3 to 35.

18. A composition according to any preceding claim, wherein the hydrophobin has the general formula (III):



wherein:

m and n are independently 0 to 20;

B_1 , B_2 , B_3 , B_4 , B_5 , B_6 , B_7 and B_8 are each independently amino acids selected from Cys, Leu, Ala, Pro, Ser, Thr, Met or Gly, at least 7 of the residues B_1 through B_8 being Cys;

a is 5 to 15;

c is 5 to 40;

d is 4 to 23;

e is 5 to 12; and


g is 6 to 21.

19. A composition according to any preceding claim, wherein all 8 of the residues B_1 through B_8 are Cys.

20. A composition according to any preceding claim, wherein the hydrophobin is a hydrophobin fusion protein.

21. A composition according to any preceding claim, wherein the hydrophobin is obtained or obtainable from a filamentous fungus.

22. A composition according to claim 21, wherein the hydrophobin is obtained or obtainable from a fungus of genus selected from the group consisting of *Cladosporium*, *Ophistoma*, *Cryphonectria*, *Trichoderma*, *Gibberella*, *Neurospora*, *Maganaporthe*, *Hypocrea*, *Xanioria*, *Emericella*, *Aspergillus*, *Paracocciooides*, *Metarhizium*, *Pleurotus*, *Coprinus*, *Dicotyonema*, *Flammulina*, *Schizophyllum*, *Agaricus*, *Pisolithus*, *Tricholoma*, *Pholioka*, *Talaromyces* and *Agrocybe*.
23. A composition according to any preceding claim, wherein the hydrophobin is generated *in situ* in the composition.
24. A composition according to any preceding claim, wherein, in use, the hydrophobin causes the equilibrium surface tension at a water/air interface to reduce to below 45 mN/m.
25. A composition according to any preceding claim, wherein, in use, the hydrophobin causes the surface shear elasticity at a water/air interface to increase to 300-700 mN/m.
26. A composition according to any preceding claim, wherein the hydrophobin is a Class I hydrophobin.
27. A composition according to claim 26, wherein the hydrophobin is a Class II hydrophobin having the general formula (IV):



wherein:

m and n are independently 0 to 200;

B₁, B₂, B₃, B₄, B₅, B₆, B₇ and B₈ are each independently amino acids selected from Cys, Leu, Ala, Ser, Thr, Met or Gly, at least 6 of the residues B₁ through B₈ being Cys;

a is 6 to 12;


c is 8 to 16;

d is 2 to 20;

e is 4 to 12; and

g is 5 to 15.

28. A composition according to claim 26 or claim 27, wherein the hydrophobin is a Class II hydrophobin having the general formula (V):



wherein:

m and n are independently 0 to 10;

B₁, B₂, B₃, B₄, B₅, B₆, B₇ and B₈ are each independently amino acids selected from Cys, Leu or Ser, at least 7 of the residues B₁ through B₈ being Cys;

a is 7 to 11;

c is 11;

d is 4 to 18;

e is 6 to 10; and

g is 7 to 10.

29. A composition according to any one of claims 26 to 28, wherein all 8 of the residues B₁ through B₈ are Cys.
30. A composition according to any one of claims 26 to 29, wherein the group (X₃)_c comprises the sequence motif ZZXZ, wherein Z is an aliphatic amino acid; and X is any amino acid.
31. A composition according to any preceding claim, wherein the hydrophobin is present in a concentration of 0.001% to 5% by weight of the total weight of the composition.
32. A composition according to claim 31, wherein the hydrophobin is present in a concentration of 0.01 % to 0.5% by weight of the total weight of the composition.
33. A composition according to any one of claims 4 to 32, wherein the detergent is present in a concentration of between 0.001 and 5 g/L.
34. A composition according to claim 33, wherein the detergent is present in a concentration of between 0.01 to 0.5 g/L.
35. A composition according to any preceding claim, additionally containing one or more enzymes selected from the group consisting of a protease, an amylase, a

glucoamylase, a maitogenic amylase, a non-maitogenic amylase, a lipase, a cuiinase, a carbohydrase, a ce'uiase, a pectinase, a mannanase, an arabinase, a galactanase, a xylanase, an oxidase, a laccase, and a peroxidase.

36. A composition according to any one of claims 4 to 35, wherein the detergent comprises one or more surfactants.
37. A composition according to claim 36, wherein the surfactants are selected from the group consisting of non-ionic (including semi-polar), anionic, cationic and zwitterionic.
38. A composition according to any one of claims 1 to 37, in powder form.
39. A composition according to any one of claims 1 to 37, in liquid form.
40. A method of removing a lipid-based stain from a surface by contacting the surface with a composition according to any one of claims 1 to 39.
41. The use of composition according to any one of claims 1 to 39 to reduce or remove lipid stains from a surface.
42. A method of cleaning a surface, comprising contacting the surface with a composition according to any one of claims 1 to 39.
43. A method of cleaning an item, comprising contacting the item with a composition according to any one of claims 1 to 39.
44. A method according to claim 43, wherein the item is a clothing item.
45. A method according to claim 43, wherein the item is a tableware item.




Fig. 1a




Fig. 1b




Fig. 1c




Fig. 2a




Fig. 2b




Fig. 2c




Fig. 2d




Fig. 2e




Fig. 3a




Fig. 3b




Fig. 3c




Fig. 3d




Fig. 3e




Fig. 4a




Fig. 4b




Fig. 4c




Fig. 4d




Fig. 4e




Fig. 5a




Fig. 5b




Fig. 5c




Fig. 5d




Fig. 5e

Fig. 6**SEQ ID NO: 1***Trichoderma reesei* HFBII (Y11894.1)

```
1 cacattcact caactcctct ttctcaactc tccaaacaca aacattcttt gttgaatacc
61 aaccatcacc acctttcaag atgcagttct tcgccgtcgc cctcttcgcc accagcgccc
121 tggctgctgt ctgccctacc ggctctttct ccaacctctc gtgctgtgcc accaacgtcc
181 tcgacctcat tggcgttgac tgcaagaccg gtatggtgaa ttccaatctc tgggcatcct
241 gacattggac gatacagttg acttacacga tgctttacag ctaccatcgc cgtcgcacact
301 ggcgccatct tccaggctca ctgtgccagc aagggtcca agcctctttg ctgcgttgct
361 cccgtggtaa gtagtgctcg caatggcaaa gaagtaaaaa gacatttggg cctgggatcg
421 ctaactcttg atatcaaggc cgaccaggct ctctgtgcc agaaggccat cggcaccttc
481 taaagcaatg gcttgcttta ctgccggcag tctttgagaa ctctgggctc acaaaagacg
541 acttgcatgt atcatggggg ctgcgcaaatg ggaggatttg gaggggattg aggctggggt
601 tggcctatta gaggattgca taatggaaga tttgcgagca ggacatagac gtatctagag
661 ttctagt
```

Fig. 7**SEQ ID NO: 2***Trichoderma reesei* HFBII (P79073.1)

```
1 MQFFAVALFA TSALAAVCPT GLFSNPLCCA TNVLDLIGVD CKTPTIAVDT GAIFQAHCAS
61 KGSKPLCCVA PVADQALLCQ KAIGTF
```

Fig. 8

SEQ ID NO: 3

Trichoderma reesei HFBI (Z68124.1)

1 tttgtatggc tggatctcga aaggcccttg tcatcgccaa gcggtggctaa tatcgaatga
 61 gggacaccga gttgcatatc tcttgatcat tcaaacgaca agtgtgaggt aggcaatcct
 121 cgtatcccat tgctgggctg aaagcttcac acgtatcgca taagcgtctc caaccagtgc
 181 ttaggtgacc cttaaggata cttacagtaa gactgtatta agtcagtcac tctttcactc
 241 gggctttgaa tacgatcctc aatactcccg ataacagtaa gaggatgata cagcctgcag
 301 ttggcaaatg taagcgtaat taaactcagc tgaacggccc ttggtgaaag tctctctcga
 361 tcaaagcaaa gctatccaca gacaaggggt aagcaggctc actcttccta cgccttggat
 421 atgcagcttg gccagcatcg cgcattggcca atgatgcacc cttcacggcc caacggatct
 481 cccgttaaac tcccctgtaa cttggcatca ctcatctgtg atoccaacag actgagttgg
 541 gggctcggc tggcggatgt cggagcaaaag gatcacttca agagcccaga tccggttggg
 601 ccattgccaa tggatctaga ttcggcacct tgatctcgat cactgagaca tgggtgagttg
 661 cccggacgca ccacaactcc cctgtgtgca ttgagtcccc atatgctct tctcagcgtg
 721 caactctgag acggattagt cctcacgatg aaattaactt ccagcttaag ttcgtagcct
 781 tgaatgagtg aagaaatttc aaaaacaaac tgagttagagg tcttgagcag ctgggggtgg
 841 acgcccctcc tcgactcttg ggacatcgta cggcagagaa tcaacggatt cacacetttg
 901 ggtcgagatg agctgatctc gacagatacg tgcttcacca cagctgcagc tacctttgcc
 961 caaccattgc gttccaggat cttgatctac atcaccgag caccggagcc aggacggaga
 1021 gaacaatccg gccacagagc agcaccgcct tccaactctg ctccctggca cgtcacacaa
 1081 cctgatatta gatatccacc tgggtgattg ccattgcaga gagggtggcag ttggtgatac
 1141 cgactggcca tgcaagacgc ggcggggcta gctgaaatgt ccccgagagg acaattggga
 1201 gcgtctatga cggcgtggag acgacgggaa aggactcagc cgtcatgttg tgttgccaat
 1261 ttgagattgt tgaccgggaa aggggggacg aagaggatgg ctgggtgagg tggattggg
 1321 ctgatgcatc attcgactca gtgagcgtatg tagagctcca agaataaaa tatccttct
 1381 ctgtcttctc aaaaatctcct tccatcttgt ccttcatcag caccgagcc agcctgaaca
 1441 cctccagtca acttccctta ccagtacatc tgaatcaaca tccattcttt gaaatctcac
 1501 cacaaccacc atcttcttca aatgaagt tctcgccatc gccgctctct ttgcccgcgc
 1561 tgccgttggc cagcctctcg aggaccgag caacggcaac ggcaatggtt gccctcccgg
 1621 cctcttcagc aacccccagt gctgtgccac ccaagtctt ggctcatcg gccttgactg
 1681 caaagtcctg aagttgagcc ataacataag aatcctcttg acggaaatat gccttctcac
 1741 tcttttacc ctgaacagcc tcccagaacg tttacgacgg caccgacttc cgcaacgtct
 1801 gcgcaaaaac cggcgcaccg cctctctgct gcggtggccc cgttgaagt tgatgccc
 1861 gctcaagctc cagtctttgg caaacccatt ctgacacca gactgcaggc cggccaggct
 1921 cttctgtgcc agaccgcctg cgggtgcttga gatgcccgc cggggtaag gtgtgcccgt
 1981 gagaaagccc acaaatggtt gatgaggacc atttccggtg ctgggaaagt tggctccacg
 2041 tgtttgggca ggtttgggca agttgtgtag atattccatt cgtacgcat tcttattctc
 2101 caatatttca gtacactttt cttcataaat caaaaagact gctattctct ttgtgacatg
 2161 ccggaaggga acaattgctc ttggtctctg ttatttgcaa gtaggagtg gagattcgcc
 2221 ttagagaaag tagagaagct gtgcttgacc gtggtgtgac tcgacgagga tggactgaga
 2281 gtgttaggat taggtcgaac gttgaagtgt atacaggatc gtctggcaac ccacggatcc
 2341 tatgacttga tgcaatgggt aagatgaatg acagtgtgag aggaaaagga aatgtccgcc
 2401 ttcagctgat atccacgcca atgatacagc gatatactc caatatctgt gggaacgaga
 2461 catgacatat ttgtgggaac aacttcaaac agcgagcca gacctcaata tgcacatcca
 2521 aagccaaaca ttggcaagac gagagacagt cacattgtcg tcgaaagat gcatcgtacc
 2581 caaatcatca gctctcatta tcgcctaac cacagattgt ttgcccctcc ccaactccaa
 2641 aacgttacta caaaagacat ggcgaatgc aaagacctga aagcaaac tttttgagc
 2701 tcaattccct cctttgtcct cggaaatgat atccttcacc aagtaaaaga aaaagaagat
 2761 tgagataata catgaaaagc acaacggaaa cgaaagaacc aggaaaagaa taaatctatc
 2821 acgcaccttg tccccacact aaaaagcaaca gggggggtaa aatgaaat

Fig. 9**SEQ ID NO: 4***Trichoderma reesei* HFBI (P52754.1)

1 MKFFAIAALF AAAAVAQPLE DRSNGNGNVC PPGLFSNPQC CATQVLGLIG LDCKVPSQNV
 61 YDGTDFRNVC AKTGAQPLCC VAPVAGQALL CQTAVGA

Fig. 10**SEQ ID NO: 5***Schizophyllum commune* SC3 (M32329.1)

1 agtcgaacac ccagttcaa ctaccccagc ccttccttcc ttcgctatcc ttccttacia
 61 cctgctegcc atgttegcc gtctcccctc cgtgttcctc taegccttcg tcegttcgg
 121 cgcctcgtc gctgccctcc caggtggcca cccgggcaag acgtacgtcg acctctcacc
 181 gtccctctaat gtcttgctga tgaagccccc tatagcacgc cgcgggttac gacgacggtg
 241 acggtgacca cgggtgagtag ctttctcgcc gtcgacgact cgaacgcatt ggctaatttt
 301 tgctcatagc cgccctcgac gacgaccatc gccccgggtg gcacgtgtac tacgggggtg
 361 ctctcttctg gcaaccaggt tcaatcggta cgtacatcaa agcggcacga ccaggcatct
 421 cagctgacgg ccacatcgta caggcgagca gcagccctgt taccgccctc ctccggcctgc
 481 tcggcatgtg cctcagcgac ctcaacgttc tcgttggcat cagctgtctc cccctcactg
 541 tgagatcttt ttgttccactg tcccaattac tgcgactga cagactttgc caggtcactg
 601 gtgtcggagg cagcggctgt tcggcgcaga ccgtctgtct cgaaaacacc caattcgtat
 661 gtatactttc catgcgtgtc ctttctccg ctaatcatct gtagaacggg ctgatcaaca
 721 tcggttgca ccccatcaac atcctctgag caggtgaacg ccctgttcgg tgggatattc
 781 gggcgacggg agcctcgggc aatctgagcc tcgttactgc ctagcaaatt cggaatccct
 841 tcgatgtcat agggtcgogg acaagtgatc gtcttgctac atactccaag gtgttgactc
 901 attccctcag ataatgaaca ttgtgttgtg tgtgtttgt tctct

Fig. 11**SEQ ID NO: 6***Schizophyllum commune* SC3 (AAA96324.1)

1 MFARLPVVFL YAFVAFGALV AALPGGHPGT TTPPVTTT VTTPPSTTTI AAGGTCTTGS
 61 LSCCNQVQSA SSSPVTALLG LLGIVLSDLN VLVGISCSP LTVIGVGGSGC SAQTVCCENT
 121 QFNGLINIGC TPINIL

Fig. 12**SEQ ID NO: 7***Neurospora crassa* EAS (X67339.1)

1 atcatcagca tcaacatctt cacttcacaa catcttctca accttccaac tcacettcca
 61 aaccaccttc aaaaccaact cccagcttct ttcagcaaac cccaaccgc caaaatgcag
 121 ttcaccageg tcttcacat cctcgccatt gccatgaccg cgcgtgcggc cccggctgag
 181 gttgttcccc gcgccaccac catcggcccc aacacctgct ccatcgacga ctacaagcct
 241 tactgctgcc agtctatgtc cggccccgcc ggctccctg gtctcctcaa cctcatcccc
 301 gtcgacctca gcgcctcgct cggctgcggt gtcgggtgca tcggctcca atgtggtgcc
 361 agcgtcaagt gctgcaagga cgatgttacc aacaccggca actccttctt catcatcaac
 421 gctgccaact gcgttgccca agtgtttacg cggcaacagc gcaaagtcta ggcaatgcct
 481 tgtttcaac gctgctgcca gtccagcacc ccccttctgc agcaaggagc ccccttctgc
 541 tggactggca gcacaacgag ctgctactac aacacaagca tcatgctgg agcaacaga
 601 agccgataat cttggggttt ggttttgggg gatgaagggt atgagttgat ggattggatc
 661 gatattctac aatgcgtgto tcttctggtt aagatctgct ttactatctt cctattttct
 721 ttacacata gctatgtatc actaaggcct ggtgattaat acactctctt aacct

Fig. 13**SEQ ID NO: 8***Neurospora crassa* EAS (AAB24462.1)

1 MQFTSVFTIL AIAMTAAAP AEVVPRTTI GPNTCSIDY KPYCCQMSG PAGSPGLLNL
 61 IPVDLSASLG CVVGVIGSQC GASVKCKDD VNTGNSFLI INAANCVA

Fig. 14**SEQ ID NO: 9***Talaromyces thermophilus* TT1

(DNA sequence encoding the precursor TT1 hydrophobin, SEQ ID NO:4 of US 7241734)

1 atgaagttcg cgggtgtett gcttgctgto gccgctgcgg cgactgcctt gccaaacgtc
 61 ggtcccagtg ggaagacggc tcacaagcog caccaggagc ctttctggcc tgtgcagcag
 121 gacgtgaccg tggaaacaggc caaggctatc tgtggtgaag gcaaccaggt cgcttgctgc
 181 aacgaggtca gtaacgcggg cgacaccaac gaaatcgga cggccccctt ggctggcacc
 241 ctcaaggacc tgctggcgg caagaacggc gccaaaggcc tgggtctctt cgacaagtgc
 301 tcgctctca atgtgatct cctgcttggc ctgtcgagcc tcatcaacca agaatgcaag
 361 cagcacattg cctgctgcca gggcaacgag gccgattcct ccaacgacct catcggtctc
 421 aacattcctt gcattgcctt tggtcgctg ctg

Fig. 15**SEQ ID NO: 10***Talaromyces thermophilus* TT1

(amino acid sequence of the precursor TT1 hydrophobin, SEQ ID NO:3 US 7241734)

1 MKFAGVLLAV AAAATALPNV GPSGKTAHKP HQEPFWPVQQ DVTVEQAKAI CGEGNQVACC
 61 NEVSYAGDIT EIATGPLAGT LKDLLGGKNG AKGLGLFDKC SRLNVDLLG LSSLINQECK
 121 QHIACCQNE ADSSNDLIGL NIPCIALGSL L

Fig. 16**SEQ ID NO: 11**

EVSQDLFNQFNLFQAQYSAAAYCGKNNDAPAGTNTCTGNACPEVEKADATFLYSFEDSGVGD
VTGFLALDNTNKLIVLSFRGSRSIENWIGNLNFDLKEINDICSGCRGHDGFTSSWRSVADTLRQ
KVEDAVREHPDYRVVFTGHSLGGALATVAGADLRNGYDIDVFSYGAPRVGNRAFAEFLTV
QTGGTLYRITHTNDIVPRLPPREFGYSHSSPEYWIKSGTLVPVRRRDIVKIEGIDATGGNNQPNIP
DIPAHLWYFGLIGTCL

Fig. 17**SEQ ID NO: 12**

MLPWKRALRPLSALMLAVAVALTPAATATADTTTAAAPSSGWNDYDCKPSAAHPRPVVLV
HGTGLNSVDNWLVLAPYLVKRGYCVFSLDYGQLPGVPPFFHGLGPVDKSAEQLDAYVDKVL
ATGAPEADIVGHSQGGMMPRYYLKFLGGAAKVNALVGIAPSNHGTDLNGFTALLPYFPGAAD
LLGRHTPALADQVTGS AFLTRLNADGDTVAGVRYTVIATRYDEVVTPWRSQYLSGPNVRNVL
LQDLCPLDLSEHVAIGVFDLIAAYHEVANALDPAHATPTTCASVFG

Fig. 18**SEQ ID NO: 13**

Ala Val Gly Val Thr Ser Thr Asp Phe Thr Asn Phe Lys Phe Tyr Ile Gln His Gly Ala Ala Ala Tyr
Cys Asn Ser Gly Thr Ala Ala Gly Ala Lys Ile Thr Cys Ser Asn Asn Gly Cys Pro Thr Ile Glu Ser
Asn Gly Val Thr Val Val Ala Ser Phe Thr Gly Ser Lys Thr Gly Ile Gly Gly Tyr Val Ser Thr Asp
Ser Ser Arg Lys Glu Ile Val Val Ala Ile Arg Gly Ser Ser Asn Ile Arg Asn Trp Leu Thr Asn Leu
Asp Phe Asp Gln Ser Asp Cys Ser Leu Val Ser Gly Cys Gly Val His Ser Gly Phe Gln Asn Ala
Trp Ala Glu Ile Ser Ala Gln Ala Ser Ala Ala Val Ala Lys Ala Arg Lys Ala Asn Pro Ser Phe Lys
Val Val Ala Thr Gly His Ser Leu Gly Gly Ala Val Ala Thr Leu Ser Ala Ala Asn Leu Arg Ala Ala
Gly Thr Pro Val Asp Ile Tyr Thr Tyr Gly Ala Pro Arg Val Gly Asn Ala Ala Leu Ser Ala Phe Ile
Ser Asn Gln Ala Gly Gly Glu Phe Arg Val Thr His Asp Lys Asp Pro Val Pro Arg Leu Pro Pro
Leu Ile Phe Gly Tyr Arg His Thr Thr Pro Glu Tyr Trp Leu Ser Gly Gly Gly Gly Asp Lys Val Asp
Tyr Ala Ile Ser Asp Val Lys Val Cys Glu Gly Ala Ala Asn Leu Met Cys Asn Gly Gly Thr Leu
Gly Leu Asp Ile Asp Ala His Leu His Tyr Phe Gln Ala Thr Asp Ala Cys Asn Ala Gly Gly Phe Ser
Trp Arg

Fig. 19

Lipase 3 – SEQ ID NO. 29

Met Phe Ser Gly Arg Phe Gly Val Leu Leu Thr Ala Leu Ala Ala Leu
 -27 -25 -20 -15

Gly Ala Ala Ala Pro Ala Pro Leu Ala Val Arg Ser Val Ser Thr Ser
 -10 -5 1 5

Thr Leu Asp Glu Leu Gln Leu Phe Ala Gln Trp Ser Ala Ala Ala Tyr
 10 15 20

Cys Ser Asn Asn Ile Asp Ser Lys Asp Ser Asn Leu Thr Cys Thr Ala
 25 30 35

Asn Ala Cys Pro Ser Val Glu Glu Ala Ser Thr Thr Met Leu Leu Glu
 40 45 50

Phe Asp Leu Thr Asn Asp Phe Gly Gly Thr Ala Gly Phe Leu Ala Ala
 55 60 65

Asp Asn Thr Asn Lys Arg Leu Val Val Ala Phe Arg Gly Ser Ser Thr
 70 75 80 85

Ile Glu Asn Trp Ile Ala Asn Leu Asp Phe Ile Leu Glu Asp Asn Asp
 90 95 100

Asp Leu Cys Thr Gly Cys Lys Val His Thr Gly Phe Trp Lys Ala Trp
 105 110 115

Glu Ser Ala Ala Asp Glu Leu Thr Ser Lys Ile Lys Ser Ala Met Ser
 120 125 130

Thr Tyr Ser Gly Tyr Thr Leu Tyr Phe Thr Gly His Ser Leu Gly Gly
 135 140 145

Ala Leu Ala Thr Leu Gly Ala Thr Val Leu Arg Asn Asp Gly Tyr Ser
 150 155 160 165

Val Glu Leu Tyr Thr Tyr Gly Cys Pro Arg Ile Gly Asn Tyr Ala Leu
 170 175 180

Ala Glu His Ile Thr Ser Gln Gly Ser Gly Ala Asn Phe Arg Val Thr
 185 190 195

His Leu Asn Asp Ile Val Pro Arg Val Pro Pro Met Asp Phe Gly Phe
 200 205 210

Ser Gln Pro Ser Pro Glu Tyr Trp Ile Thr Ser Gly Asn Gly Ala Ser
 215 220 225

Val Thr Ala Ser Asp Ile Glu Val Ile Glu Gly Ile Asn Ser Thr Ala
 230 235 240 245

Gly Asn Ala Gly Glu Ala Thr Val Ser Val Val Ala His Leu Trp Tyr
 250 255 260

Phe Phe Ala Ile Ser Glu Cys Leu Leu *
 265 270

Fig. 19a

SEQ ID No: 14

```

Ser Val Ser Thr Ser
 1           5

Thr Leu Asp Glu Leu Gln Leu Phe Ala Gln Trp Ser Ala Ala Ala Tyr
           10           15           20

Cys Ser Asn Asn Ile Asp Ser Lys Asp Ser Asn Leu Thr Cys Thr Ala
           25           30           35

Asn Ala Cys Pro Ser Val Glu Glu Ala Ser Thr Thr Met Leu Leu Glu
           40           45           50

Phe Asp Leu Thr Asn Asp Phe Gly Gly Thr Ala Gly Phe Leu Ala Ala
           55           60           65

Asp Asn Thr Asn Lys Arg Leu Val Val Ala Phe Arg Gly Ser Ser Thr
 70           75           80           85

Ile Glu Asn Trp Ile Ala Asn Leu Asp Phe Ile Leu Glu Asp Asn Asp
           90           95           100

Asp Leu Cys Thr Gly Cys Lys Val His Thr Gly Phe Trp Lys Ala Trp
           105           110           115

Glu Ser Ala Ala Asp Glu Leu Thr Ser Lys Ile Lys Ser Ala Met Ser
           120           125           130

Thr Tyr Ser Gly Tyr Thr Leu Tyr Phe Thr Gly His Ser Leu Gly Gly
           135           140           145

Ala Leu Ala Thr Leu Gly Ala Thr Val Leu Arg Asn Asp Gly Tyr Ser
           150           155           160           165

Val Glu Leu Tyr Thr Tyr Gly Cys Pro Arg Ile Gly Asn Tyr Ala Leu
           170           175           180

Ala Glu His Ile Thr Ser Gln Gly Ser Gly Ala Asn Phe Arg Val Thr
           185           190           195

His Leu Asn Asp Ile Val Pro Arg Val Pro Pro Met Asp Phe Gly Phe
           200           205           210

Ser Gln Pro Ser Pro Glu Tyr Trp Ile Thr Ser Gly Asn Gly Ala Ser
           215           220           225

Val Thr Ala Ser Asp Ile Glu Val Ile Glu Gly Ile Asn Ser Thr Ala
           230           235           240           245

Gly Asn Ala Gly Glu Ala Thr Val Ser Val Val Ala His Leu Trp Tyr
           250           255           260

Phe Phe Ala Ile Ser Glu Cys Leu Leu *
           265           270
    
```


Fig. 20

SEQ ID NO: 15

VYITEVSQLNTSELERGEELLEQVEEIAAISGK GKVNLVGHSHGGPTVRYVAAVRPDLVASVTS
VGAPHKGSDTADFIRQIPPGSAGEAIVAGIVNGLGALINFLSGSSSTSPQNALGALES LNSEGAA
AFNAKYPQGIPTSACGEGAYKVNQVSVSYSSWSGTSPLTNVLDVSDLLLGASSLTFDEPN DGLVG
RCSSHLGK VIRDDYRMNHLDEVNQTFGLTSLFETDPVTVYRQQANRLKLAGL

Fig. 21

SEQ ID NO: 16

ANPYERGPNTDALLEASSGPFSVSEENVSRLSASGFGGGTIYYPRENNTYGAVAISPGYTGTE
ASIAWLGERIASHGFVVITIDTITITLDQPD SRAEQLNAALNHMINRASSTVRSRIDSSRLAVMGH
SMGGGGTLRLASQRPD LKAAIPLTPWHLNKNWSSVTVP TLIIGADLDTIAPVATHAKPFYNSLP
SSISKAYLELDGATHFAPNIPNKIIGKYSVAWLKRFVDNDTRYTQFLCPGPRDGLFGEVEEYRS
TCPF

Fig. 22

SEQ ID NO: 17

DTTTAAPSSGWNDYDCKPSAAHPRPVVLVHGT LGNSVDNWLVLAPYLVKRGYCVFSLDYGQ
LPGVPPFFHGLGPVDKSAEQLDAYVDKVLAA TGAPEADIVGHSQGGMMPRYYLKFLGGAAKV
NALVGIAPSNHGTDLNGFTALLPYFPGAADLLGRHTPALADQVTGSAFLTRLNADGDTVAGV
RYTVIATRYDEVVTPWRSQYLSGPNVRNVLLQDLCPLDLSEHVAIGVFDLIAYHEVANALDPA
HATPTTCASVFG

Fig. 23

SEQ ID NO: 18

MRSSLVFFVSAWTALASPIRREVSQDLFNQFN LFAQYSAAA YCGKNN DAPAGTNITCTGNAC
PEVEKADATFLYSFEDSGVDVTGFLALDNTNKLIVLSFRGSR SIENWIGNLNFDLKEINDICSG
CRGHDGFTSSWRSVADTLRQKVEDAVREHPDYR VVFTGHSLGGALATVAGADLRGNGYDID
VFSYGAPRVGNRAFAEFLTVQTGGTLYRITH TNDIVPRLPPREFGYSHSSPEYWIKSGTLVPVR
RRDIVKIEGIDATGGNNQPNIPDIP AHLWYFGLIGTCL

Fig. 24

SEQ ID NO: 19

MNNKKTLLALCIGSSLLLSGPAEAGLFGSTGYTKTKYPIVLTHGLLGFDSILGVDYWYGIPSSL
RSDGASVYITEVSQLNTSELERGEELLEQVEEIAAISGK GKVNLVGHSHGGPTVRYVAAVRPDL
VASVTSVGAPHKGSDTADFIRQIPPGSAGEAIVAGIVNGLGALINFLSGSSSTSPQNALGALES L
NSEGAAAFNAKYPQGIPTSACGEGAYKVNQVSVSYSSWSGTSPLTNVLDVSDLLLGASSLTFDEP
NDGLVGRCSSHLGK VIRDDYRMNHLDEVNQTFGLTSLFETDPVTVYRQQANRLKLAGL

Fig. 25

SEQ ID No. 20

MAVMTPRRERSLLSRALQVTAATAALVTAVSLAAPAHAANPYERGPNPDTALLEASSGPFS
 VSEENVSRLSASGFGGGTIYYPRENNTYGAVAISPGYTGTEASIAWLGERIASHGFVVITIDTITT
 LDQPDSRAEQLNAALNHMINRASSTVRSRIDSSRLAVMGHSMGGGGTLRLASQRPDLKAAIPL
 TPWHLNKNWSSVTVPTLIIGADLDTIAPVATHAKPFYNSLPSSISKAYLELDGATHFAPNIPNKII
 GKYSVAWLKRFVDNDTRYTQFLCPGPRDGLFGEVEEYRSTCPF

Fig. 26

SEQ ID No. 21

1	EAEAAVGVTS	TDFTNFKFYI	QHGAAYCNS	GTAAGAKITC	SNNGCPTIES
51	NGVTVVASFT	GSKTGIGGYV	STDSSRKEIV	VAIRGSSNIR	NWLTNLDLFDQ
101	SDCSLVSGCG	VHSGFQNAWA	EISAQASAAV	AKARKANPSF	KVVATGHSLG
151	GAVATLSAAN	LRAAGTPVDI	YTYGAPRVGN	AALSAFISNQ	AGGEFRVTHD
201	KDPVPRLPPL	IFGYRHITPE	YWLSGGGGDK	VDYAISDVKV	CEGAANLMCN
251	GGTLGLDIDA	HLHYFQATDA	CNAGGFSWR		

Fig. 27

SEQ ID No. 22

GACACCACGACCGCGGCACCCTCCTCGGGCTGGAACGACTACGACTGCAAG
 CCGTCCGCGCGCACCCCCGCCCCGTGGTCCCTCGTCCACGGCACGCTCGGC
 AACAGCGTGGACAACCTGGCTGGTCCCTGGCCCCGTACCTCGTCAAGCGCGGC
 TACTGCGTGTCTCCCTGGACTACGGCCAGCTGCCGGGGCGTGCCCTTCTTCC
 ACGGCCTGGGCCCGGTGGACAAGAGCGCCGAGCAGCTGGACGCCTACGTGG
 ACAAGGTGCTCGCCGCCACCGGCGCCCCGGAGGGCGGACATCGTCGGGCACT
 CGCAGGGGGGCATGATGCCCCGGTACTACCTGAAGTTCCTCGGCGGGGGCGG
 CCAAGGTCAACGCCCTGGTGGGCATCGCCCCCTCGAACCACGGGACGGACC
 TCAACGGCTTCAACGCCCTCCTGCCGTAATTCCCAGGGCGCCGCCGACCTCCT
 CGGCCGGCACACCCCGGCGCTGGCCGACCAGGTACCGGGAGCGCGTTCCT
 GACCCGCTGAACGCGGACGGCGACACGGTCGCGGGGGTCCGCTACACCGT
 CATCGCCACGCGCTACGACGAGGTCGTCACCCCCTGGCGGTCCCAGTACCT
 GAGCGGCCCGAACGTCCGGAACGTGCTGCTCCAGGACCTGTGCCCCCTCGA
 CTTGAGCGAACACGTGGCCATCGGCGTGTTCGACCTCATCGCATAACCACGAG
 GTCGCCAACGCCCTGGACCCGGCGCACGCCACCCCCACGACCTGCGCGTCC
 GTCTTCGGC

Fig. 28

SEQ ID No. 23

ATGGGCTTTGGGAGCGCTCCCATCGCGTTGTGTCCGCTTCGCACGAGGAGGAACGCTTTGA
AACGCCTTTGGCCCTGCTCGCGACCGGCGTGTGATCGTCGGCCTGACTGCGCTAGCCGG
CCCCCGGCACAGGCCGACACCACGACCGCGGCACCCTCCTCGGGCTGGAACGACTACGA
CTGCAAGCCGTCCGCCGCGCACCCCCGCCCGTGGTCCCTCGTCCACGGCACGCTCGGCAAC
AGCGTGGACAACCTGGCTGGTCCCTGGCCCCGTACCTCGTCAAGCGCGGCTACTGCGTGTCT
CCCTGGACTACGGCCAGCTGCCGGGCGTGCCCTTCTTCCACGGCCTGGGCCCCGGTGGACA
AGAGCGCCGAGCAGCTGGACGCCTACGTGGACAAGGTGCTCGCCGCCACCGGCGCCCCGG
AGGCGGACATCGTCGGGCACTCGCAGGGGGGCATGATGCCCCGGTACTACCTGAAGTTCC
TCGGCGGGGCGGCCAAGGTCAACGCCCTGGTGGGCATCGCCCCCTCGAACCACGGGACGG
ACCTCAACGGCTTCAACGCCCTCCTGCCGTACTTCCCGGGCGCCCGACCTCCTCGGCCG
GCACACCCCGCGCTGGCCGACCAAGGTCACCGGGAGCGCGTTCCTGACCCGCCTGAACGC
GGACGGCGACACGGTCCGCGGGGTCCGCTACACCGTCATCGCCACGCGCTACGACGAGGT
CGTACCCCCCTGGCGGTCCCAGTACCTGAGCGGCCCGAACGTCCGGAACGTGCTGCTCCA
GGACCTGTGCCCCCTCGACTTGAGCGAACACGTGGCCATCGGCGTGTTCGACCTCATCGCA
TACCACGAGGTCGCCAACGCCCTGGACCCGGCGCACGCCACCCCCACGACCTGCGCGTCC
GTCTTCGGC

Fig. 29

SEQ ID No. 24

MGFGSAPIALCPLRTRRNALKRLLALLATGVSIVGLTALAGPPAQADTTTAAPSSGWN DYDCK
PSAAHPRPVVLVHGTLGNSVDNWLVLAPYLVKRGYCVFSLDYGQLPGVPPFHGLGPVDKSAE
QLDAYVDKVLAAATGAPEADIVGHSQGGMMPRYYLKFLLGGAAKVNALVGIAPSNHGTDLNGF
TALLPYFPGAADLLGRHTPALADQVTGSAFLTRLNADGDTVAGVRYTVIATRYDEVVTPWRS
QYLSGPNVRNVLLQDLCPDLSEHVAIGVFDLIAYHEVANALDPAHATPTTCASVFG

Fig. 30




Fig. 31




Fig. 32




Fig. 33




Fig. 34




Fig. 35




Fig. 36

SEQ ID No. 25

MKCCRIMFVLLGLWVFLGSLVPGGRTEAASLRANDAPIVLLHGFTGWGREEMFGFKYWGGVVRGDI EQWLN
 DNGYRTFTLAVGLPSSNWDRA CEAYAQLVGGTVDYGA AHA AKHG HARFGR TYPGLLPELKRGGRIH IIAH
 SQGGQTARMLVSLLENGSQEEREYAKAHNVSLSPLEGGHHFVLSVTTIATPHDGTTLVNMVDFTRDFD
 LQKAVLEAAA VASNPYTSQVYDFKLDQWGLRRQPGESFDHYFERLKRSPVWTSTDTARYDLSVSGAEKL
 NQWVQASPNTYYLSFSTERTYRGAL TGNHYPELGMNAFSAVVCAPFLG SYRNPTLGIDDRWLENDGIVNT
 VSMNGPKRGSSDRIVPYDGTLLKKG VWN DMGTYNV D HLEIIGVDPNPSFDIRAFYLR LAEQ L ASLQP

Fig. 37

SEQ ID No. 26

DDYSVVEEHGQLSISNGELVNERGEQVQLKGMSSHGLQWYGQFVNYESMKWLRDDWGITVFRAMYTSSG
 GYIDDP SVKEKVKETVEAAIDLGIYVIDWHILSDNDPNYKKEAKDFFDEMSELYGDYPNVIYEIANEP
 NGS DVTW DNIKPYAEEVIPVIRDNDPNNIIVIGTGTWSQDVHHAADNQLADPNV MYAFHFYAGTHGQNL
 RDQVDYALDQGA AIFVSEWGTSAATGDGGVFLDEAQVWIDFMDERNLSWANWSLTHKDES SAALMPGANP
 TGGWTEAELSPSGTFVREKIREASDNNDPI PDPDDEASLRANDAPIVLLHGFTGWGREEMFGFKYWGGV
 RGDIEQWLN DNGYRTFTLAVGLPSSNWDRA CEAYAQLVGGTVDYGA AHA AKHG HARFGR TYPGLLPELKR
 GGRIH IIAHSQGGQTARMLVSLLENGSQEEREYAKAHNVSLSPLEGGHHFVLSVTTIATPHDGTTLVNM
 VDFTRDFDLQKAVLEAAA VASNPYTSQVYDFKLDQWGLRRQPGESFDHYFERLKRSPVWTSTDTARYD
 LSVSGAEKLNQWVQASPNTYYLSFSTERTYRGAL TGNHYPELGMNAFSAVVCAPFLG SYRNPTLGIDDRW
 LE NDGIVNTVSMNGPKRGSSDRIVPYDGTLLKKG VWN DMGTYNV D HLEIIGVDPNPSFDIRAFYLR LAEQ L
 ASLQP

Fig. 38

SEQ ID No. 27

MKFVKRRIIALV TILMLSVTSLFALQPSAKAAEHN PVVMVHGIGGASFN FAGIKSYLVSQGWSRDKLYAV
 DFWDKTGTNYNNGPVL SRFVQKVLDETGA KKVDI VAHSMGGANTLYYIKNLDGGNKVANVVTLGGANRLT
 TGKALPGTDPNQKILYTSIYSSADMIVMNYLSRLDGARNVQIHGVGHIGLLYSSQVNSLIKEGLNGGGQNTN

Fig. 39

SEQ ID No. 28

DDYSVVEEHGQLSISNGELVNERGEQVQLKGMSSHGLQWYGQFVNYESMKWLRDDWGITVFRAM
 YTSSGGYIDDP SVKEKVKETVEAAIDLGIYVIDWHILSDNDPNYKKEAKDFFDEMSELYGDY
 NVIYEIANEPNGSDVTW DNIKPYAEEVIPVIRDNDPNNIIVIGTGTWSQDVHHAADNQLADPNV
 MYAFHFYAGTHGQNL RDQVDYALDQGA AIFVSEWGTSAATGDGGVFLDEAQVWIDFMDERNLSWA
 NWSLTHKDESSAALMPGANPTGGWTEAELSPSGTFVREKIREASDNNDPI PDPDDEAEHNPVVM
 VHGIGGASFN FAGIKSYLVSQGWSRDKLYAVDFWDKTGTNYNNGPVL SRFVQKVLDETGA KKVDI
 VAHSMGGANTLYYIKNLDGGNKVANVVTLGGANRLT TGKALPGTDPNQKILYTSIYSSADMIVM
 NYSRLDGARNVQIHGVGHIGLLYSSQVNSLIKEGLNGGGQNTN

Fig. 40

SEQ ID No. 30

atgcatacgctgttaac

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2012/051660

A. CLASSIFICATION OF SUBJECT MATTER INV. C07K14/37 C11D3/38 C11D3/386 C12N9/20 ADD.				
According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07K C11D C12N				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal , BIOSIS, EMBASE, WPI Data				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
X	us 2009/101167 AI (BOECKH DI ETER [DE] ET AL) 23 April 2009 (2009-04-23) cited in the appl icati on p. 2, paragraph [0028] - p. 9, paragraph [0150]	1-45		
X	----- JOSÉ M. PALOMO ET AL: "Sol id-Phase Handl ing of Hydrophobi ns: Immobi lized Hydrophobi ns as a New Tool To Study Li pases" , BIOMACROMOLECULES, vol . 4, no. 2, 1 March 2003 (2003-03-01) , pages 204-210, XP55030962 , ISSN: 1525-7797, DOI : 10. 1021/bm0200711 the whole document ----- - / - -	1,2,4, 16-19 , 21-25 , 31,32 , 36-39		
<table border="0" style="width:100%;"> <tr> <td style="width:50%;"><input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.</td> <td style="width:50%;"><input checked="" type="checkbox"/> See patent family annex.</td> </tr> </table>			<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.	<input checked="" type="checkbox"/> See patent family annex.
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.	<input checked="" type="checkbox"/> See patent family annex.			
* Special categories of cited documents :				
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family			
Date of the actual completion of the international search 29 June 2012	Date of mailing of the international search report 26/07/2012			
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer R. von Eggel kraut-G.			

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2012/051660

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>EP 1 595 949 A1 (TOHOKU TECHNO ARCH CO LTD [JP] ; NAT INST OF ADVANCED IND SCI EN [JP]) 16 November 2005 (2005-11-16) p. 6, paragraph [0030] - p. 10, paragraph [0084] , pp. 21-26, Ex. 4-9</p> <p>-----</p>	1, 16-32, 35,38,39
A	<p>LINDER M B ET AL: "Hydrophobi ns: the protei n-amphi phi les of fi lamentous fungi ", FEMS MICROBIOLOGY REVI EWS, ELSEVI ER, AMSTERDAM, NL, vol . 29, no. 5, 1 November 2005 (2005-11-01) , pages 877-896, XP027666169 , ISSN: 0168-6445 [retri eved on 2005-11-01] the whole document</p> <p>-----</p>	1-45
A	<p>HASAN F ET AL: "Enzymes used i n detergents: Li pases" , AFRICAN JOURNAL OF BIOTECHNOLOGY, ACADEMIC PRESS, US, vol . 9, no. 31, 2 August 2010 (2010-08-02) , pages 4836-4844, XP003027509, ISSN: 1684-5315 the whole document</p> <p>-----</p>	1-45
A	<p>MESSA0UDI ABDELMONAEM ET AL: "Classi ficati on of EC 3.1.1.3 bacteri al true li pases using phylogenet i c analysi s", AFRICAN JOURNAL OF BIOTECHNOLOGY, vol . 9, no. 48, November 2010 (2010-11) , pages 8243-8247, XP002678951 , ISSN: 1684-5315 the whole document</p> <p>-----</p>	1-45

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/IB2012/051660

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2009101167	AI	23-04-2009	AT 485359 T 15--11--2010
			AU 2006274836 AI 08--02--2007
			BR PIQ614703 A2 23--08--2011
			CA 2617092 AI 08--02--2007
			CN 101233220 A 30--07--2008
			EP 1913123 AI 23--04--2008
			ES 2352970 T3 24--02--2011
			JP 2009503280 A 29--01--2009
			KR 20080041228 A 09--05--2008
			US 2009101167 AI 23--04--2009
			Wo 2007014897 AI 08--02--2007
			ZA 200801881 A 26--08--2009
EP 1595949	AI	16-11-2005	EP 1595949 AI 16--11--2005
			JP 4273504 B2 03--06--2009
			US 2006106120 AI 18--05--2006
			Wo 2004038016 AI 06--05--2004