



(86) Date de dépôt PCT/PCT Filing Date: 1995/11/13
(87) Date publication PCT/PCT Publication Date: 1996/05/30
(45) Date de délivrance/Issue Date: 2007/11/06
(85) Entrée phase nationale/National Entry: 1996/07/04
(86) N° demande PCT/PCT Application No.: JP 1995/002315
(87) N° publication PCT/PCT Publication No.: 1996/016034
(30) Priorité/Priority: 1994/11/24 (JP6/289763)

(51) Cl.Int./Int.Cl. *C07C 319/06* (2006.01),
C07C 319/24 (2006.01), *C07C 321/24* (2006.01)
(72) Inventeurs/Inventors:
GODA, HIROSHI, JP;
YAMAMOTO, MIKIO, JP;
SAKAMOTO, JUN-ICHI, JP;
KARINO, HITOSHI, JP
(73) Propriétaire/Owner:
SUMITOMO SEIKA CHEMICALS CO., LTD., JP
(74) Agent: MARKS & CLERK

(54) Titre : PROCEDE DE PRODUCTION DE COMPOSES DE SOUFRE AROMATIQUES ET HETEROAROMATIQUES
(54) Title: PROCESS FOR PRODUCING AROMATIC OR HETEROAROMATIC SULFUR COMPOUNDS

(57) **Abrégé/Abstract:**

A process for producing an aromatic or heteroaromatic thiol represented by the following general formula (2): $\text{Ar}-(\text{-SH})_n$ by hydrolyzing an aromatic or heteroaromatic halomethyl sulfide represented by the following general formula (1): $\text{Ar}-(\text{-SCH}_{3-m}\text{X}_m)_n$, in an industrial manner readily at a low cost, wherein Ar represents an unsubstituted or arbitrarily substituted aromatic or heteroaromatic group; X represents a halogen atom; m represents an integer of 1 to 3; and n represents 1 or 2.



ABSTRACT OF THE DISCLOSURE

A process for producing an aromatic or heteroaromatic thiol represented by the following general formula (2): $\text{Ar}-(\text{-SH})_n$ by hydrolyzing an aromatic or heteroaromatic halomethyl sulfide represented by the following general formula (1): $\text{Ar}-(\text{-SCH}_3\text{-}_m\text{X}_m)_n$, in an industrial manner readily at a low cost, wherein Ar represents an unsubstituted or arbitrarily substituted aromatic or heteroaromatic group; X represents a halogen atom; m represents an integer of 1 to 3; and n represents 1 or 2.

DESCRIPTION

Field of the Invention

5 The present invention relates to a novel process
for preparing an aromatic or heteroaromatic thiol or an
aromatic or heteroaromatic disulfide and a novel process
for preparing an aromatic or heteroaromatic halogenated
methyl sulfide which is useful as the starting material for
10 said thiol or disulfide. The aromatic or heteroaromatic
thiol and disulfide are useful compounds which are used for
various purposes as in pharmaceutical compositions,
agricultural compositions, functional materials or the
like.


15

Background Art

Various processes have been known for preparing
an aromatic or heteroaromatic thiol or disulfide. Among
them, the conventional processes for preparing an aromatic
or heteroaromatic thiol by bond cleavage in alkyl sulfide
20 are classified into the following three processes:

- (A) Process comprising reacting an aromatic alkyl sulfide
with alkyl thiolate in hexamethylphosphoric triamide
(abbreviation HMPA) used as a solvent
(Tetrahedron Letters 21 3099 (1980))
25 (Synthesis Communications 478 (1982))


-2-



(B) Process comprising reacting an aromatic alkyl sulfide with metal sodium in liquid ammonia


(Org. Synth. Coll. Vol.5 419 (1973))

(J. Am. Chem. Soc. 81 4939 (1959))



(C) Process comprising oxidizing an aromatic methyl sulfide with perbenzoic acid, reacting the oxide with trifluoroacetic acid, and causing triethylamine to act on the reaction mixture

(Tetrahedron Letters 25 1753 (1984))



However, these conventional processes pose the following drawbacks when commercially conducted.

The process (A) is defective in that the HMPA used as a solvent is not easily available and expensive and

that the sodium mercaptide used as a reagent is difficult to handle in a nonaqueous system. Further problematic is the disposal of dialkyl sulfide produced as a by-product.

The process (B) is industrially difficult to
5 carry out because of high risks involved in handling, use of an expensive metal sodium and low-yield production. The process (C) requires a number of reaction procedures and necessitates the use of an expensive, highly dangerous reaction reagent such as perbenzoic acid, trifluoroacetic
10 acid, etc.

As described above, none of the conventional processes for preparing an aromatic or heteroaromatic thiol by bond cleavage in alkyl sulfide are satisfactory from the viewpoint of industrial application.

15 Disclosure of the Invention


An object of the present invention is to provide a process for preparing an aromatic or heteroaromatic thiol at a commercially low cost and with ease.

Another object of the invention is to provide
20 a process for preparing an aromatic or heteroaromatic disulfide at a commercially low cost and with ease.

The present inventors conducted extensive research on the reaction for bond cleavage in alkyl sulfide to overcome the drawbacks of said known processes and to
25 provide processes for preparing aromatic or heteroaromatic

thiol and disulfide at commercially low costs and with ease.


The inventors' research found that the aromatic or heteroaromatic halogenated methyl sulfide represented by the formula (1) can be easily hydrolyzed to give the corresponding aromatic or heteroaromatic thiol represented by the formula (2) in a high yield. The finding led to the completion of the present invention. Stated more specifically, the first invention is directed to a process for preparing an aromatic or heteroaromatic thiol represented by the formula (2), the process comprising hydrolyzing an aromatic or heteroaromatic halogenated methyl sulfide represented by the formula (1) as shown below:



wherein Ar is an aromatic or heteroaromatic ring which has no substituent or which has an optional substituent or substituents, X is a halogen atom, m is an integer of 1 to 3 and n is 1 or 2.

Since thiol can be easily converted to disulfide by oxidation, the corresponding disulfide can be produced by executing an oxidation step subsequent to the above-mentioned reaction step. The second invention is directed to a process for preparing an aromatic or heteroaromatic

disulfide represented by the formula (3), the process comprising the steps of hydrolyzing an aromatic or heteroaromatic halogenated methyl sulfide represented by the formula (1), and oxidizing the reaction mixture as shown below:



wherein Ar is an aromatic or heteroaromatic ring which has no substituent or which has an optional substituent or substituents, X is a halogen atom, m is an integer of 1 to 3 and n is 1 or 2.

The present invention will be described below in more detail.

The contemplated aromatic or heteroaromatic thiol represented by the formula (2) is prepared in high yields by hydrolyzing an aromatic or heteroaromatic halogenated methyl sulfide represented by the formula (1).

In the formulas, the aromatic or heteroaromatic rings represented by Ar are not specifically limited and include a wide variety of aromatic or heteroaromatic rings which have no substituent or which have an optional substituent or substituents. Examples of the aromatic or heteroaromatic ring represented by Ar are a benzene ring, naphthalene ring, pyridine ring, pyrazole ring, pyrazine ring, triazole ring, oxazole ring, isooxazole ring,

thiazole ring, isothiazole ring, thiophene ring, benzothiophene ring, furan ring, benzofuran ring, pyrrole ring, indole ring, etc.

Examples of optional substituents are halogen,
5 cyano, formyl, amino, carboxyl or its ester, carbamoyl, alkyl carbonyl, nitro, sulfonic acid, alkyl, alkoxyl, hydroxyl, substituted phenylthio, etc.

Among the aromatic or heteroaromatic rings represented by Ar, preferred are a benzene ring, pyridine
10 ring, thiazole ring, and isothiazole ring which have an optional substituent or substituents, and more preferred are benzene rings each having, in an optional position or positions, halogen, cyano, formyl, amino, carboxyl or its ester, carbamoyl, alkyl carbonyl, nitro, sulfonic acid,
15 alkyl, alkoxyl, hydroxyl or substituted phenylthio.

In the aromatic or heteroaromatic halogenated methyl sulfide represented by the formula (1), X is chlorine, bromine or iodine and m is 1, 2 or 3. Among these substances, chlorine is preferred as X. For
20 hydrolysis to make smooth progress, desirably m is 2 or 3 or a mixture of sulfides wherein m is 2 or 3 is used.

The hydrolysis takes place and proceeds merely by adding water and heating, thereby producing the desired aromatic or heteroaromatic thiol of the formula (2). The
25 hydrolysis reaction makes smooth progress if effected in

2180533

-7-

the presence of an acid. While such acids are not specifically limited, mineral acids such as hydrochloric acid, sulfuric acid, nitric acid and phosphoric acid are used for economical advantage. Among them, hydrochloric
5 acid and sulfuric acid are more preferred. The acid is sufficient if its amount is 0.01 to 1 times the weight of halogenated methyl sulfide of the formula (1).

In this case, the addition of an alcohol, particularly a lower alcohol, causes the hydrolysis
10 reaction to more smoothly proceed. Examples of useful lower alcohols are methanol, ethanol, isopropanol, n-propanol, isobutanol, n-butanol, sec-butanol, etc. Among them, methanol is preferred from an economical viewpoint. The amount of the alcohol, although not specifically
15 limited, is usually 0.5 to 10 times the weight of halogenated methyl sulfide of the formula (1).

Useful solvents are not specifically limited and even said reaction proceeds without a solvent. Examples of useful solvents are hydrocarbons such as hexane,
20 cyclohexane and heptane, halogenated hydrocarbons such as dichloroethane, dichloromethane and chloroform, aromatic hydrocarbons such as benzene, toluene, xylene, chlorobenzene, dichlorobenzene and trichlorobenzene. The amount of the solvent used is not specifically limited but
25 is usually 0.1 to 10 times the weight of the halogenated

methyl sulfide of the formula (1).

The reaction temperature is usually in a range of about 20 to about 100°C, preferably about 50 to about 90°C. Too low a reaction temperature retards the reaction, whereas too high a reaction temperature causes a side reaction, which results in the decrease of yields. The reaction time is usually in a range of about 1 to about 10 hours.

The aromatic or heteroaromatic thiol produced according to the present invention can be isolated by conventional distillation or crystallization.

The aromatic or heteroaromatic thiols obtainable in the present invention include various compounds such as thiophenol, 4-chlorothiophenol, 2-chlorothiophenol, 2,4-dichlorothiophenol, 2,5-dichlorothiophenol, 3,5-dichlorothiophenol, 2,6-dichlorothiophenol, 4-bromothiophenol, 2,4-dibromothiophenol, 1,4-benzenedithiol, 1,2-benzenedithiol, 4-butyl-1,2-benzenedithiol, 4-cyanothiophenol, 2-cyanothiophenol, 4-methylthiophenol, 2-methylthiophenol, 4-hydroxythiophenol, 2-hydroxythiophenol, 4-acetylthiophenol, 4-nitrothiophenol, 2-nitrothiophenol, 4-aminothiophenol, 2-aminothiophenol, 4-mercaptobenzoic acid, 2-mercaptobenzoic acid, 2-formylthiophenol, 4-formylthiophenol, 4-mercaptobenzenesulfonic acid, 2-

cyano-3-chlorothiophenol, 4-carbamoylthiophenol, 4,4'-
thiodibenzenethiol, 4,4'-oxydibenzenethiol, 4-
(phenylthio)benzenethiol, 4-(phenylsulfonyl)benzenethiol,
2-pyridinethiol, 4-pyridinethiol, 2-mercaptothiophene,
5 2,6-dimercaptothiophene, 1-naphthalenethiol, 2-
naphthalenethiol, 2-mercaptopyrazine, 4-mercaptotriazole,
5-mercaptotriazole, 2-mercaptooxazole, 4-mercaptooxazole,
4-mercaptoisooxazole, 2-mercaptothiazole, 4-
mercaptothiazole, and 4-mercaptoisothiazole.

10 To obtain the aromatic or heteroaromatic
disulfide of the formula (3) as the contemplated product,
the object can be easily accomplished by additionally
executing an oxidation step after said hydrolysis reaction.
In other words, this can be done by adding an oxidizer
15 after the hydrolysis.

A specific oxidizer is not necessary and
conventional known oxidizers can be used. Oxidation
methods employable herein are oxygen oxidation, air
oxidation, oxidation involving a halogen such as chlorine,
20 and bromine, oxidation involving a peroxide such as
hydrogen peroxide and peracetic acid, and oxidation
involving an alkali metal salt of hypohalous acid such as
sodium hypochlorite and sodium hypobromite.

The reaction temperature is variable depending on
25 the oxidation method, and can not be specifically defined.


It is usually in a range of about 0 to about 100°C, preferably about 30 to about 90°C. Too low a reaction temperature results in the reduction of reaction rate, whereas too high a reaction temperature gives rise to a side reaction, thereby lowering the yield. The reaction time is usually in a range of about 0.5 to about 15 hours.

The aromatic or heteroaromatic disulfide thus formed can be easily isolated by crystallization or the like.

10 The aromatic or heteroaromatic disulfide produced by the present invention includes a disulfide derived from said aromatic or heteroaromatic thiol.

As stated above, the aromatic or heteroaromatic thiol represented by the formula (2) can be easily prepared by hydrolyzing the aromatic or heteroaromatic halogenated methyl sulfide represented by the formula (1), and the aromatic or heteroaromatic disulfide represented by the formula (3) can be easily prepared by hydrolyzing the aromatic or heteroaromatic halogenated methyl sulfide represented by the formula (1), followed by oxidation.

20 The aromatic or heteroaromatic halogenated methyl sulfide of the formula (1) to be used herein can be produced by halogenating the corresponding aromatic or heteroaromatic methyl sulfide represented by the formula (4) (the third invention of the present application).



5 wherein Ar is an aromatic or heteroaromatic ring which has no substituent or which has an optional substituent or substituents, X is a halogen atom, m is an integer of 1 to 3 and n is 1 or 2.

10 When the aromatic or heteroaromatic halogenated methyl sulfide of the formula (1) is prepared by halogenating the aromatic or heteroaromatic methyl sulfide of the formula (4), chlorine, sulfuryl chloride, bromine, sulfuryl bromide, etc. can be used as a halogenating agent. Among them, chlorine is preferred from an economical
15 viewpoint.

The aromatic or heteroaromatic halogenated methyl sulfide of the formula (1) is a compound wherein m is 1, 2 or 3. If the compound of the formula (1) wherein m is 2 or 3, or a mixture of these compounds is used, the hydrolysis
20 reaction in the subsequent step can proceed smoothly.

The amount of the halogenating agent to be used is 1.5 to 7 moles, preferably 2 to 5 moles, per mole of the aromatic or heteroaromatic methyl sulfide of the formula (4).

25 Useful solvents are not specifically limited and even said reaction proceeds without a solvent. Examples of

useful solvents are hydrocarbons such as hexane, cyclohexane and heptane, halogenated hydrocarbons such as dichloroethane, dichloromethane and chloroform, and aromatic hydrocarbons such as benzene, toluene, xylene, chlorobenzene, dichlorobenzene and trichlorobenzene. The amount of the solvent used is not specifically limited but is usually 0.1 to 10 times the weight of aromatic or heteroaromatic methyl sulfide of the formula (4).

The halogenation reaction temperature is usually in a range of about -20 to about 100°C, preferably about -5 to about 60°C, although variable with the contemplated compound. Too low a reaction temperature retards the reaction, whereas too high a reaction temperature brings about a side reaction, which leads to the decrease of the yield.

The aromatic or heteroaromatic halogenated methyl sulfide of the formula (1) thus formed can be isolated by conventional distillation or crystallization.

To obtain the aromatic or heteroaromatic thiol of the formula (2), or the aromatic or heteroaromatic disulfide of the formula (3) as the contemplated product, the reaction mixture of the halogenation reaction itself can be used in the subsequent hydrolysis reaction or oxidation reaction further to hydrolysis reaction without isolation of the aromatic or heteroaromatic halogenated

methyl sulfide of the formula (1).

In other words, the present invention provides a process for preparing the aromatic or heteroaromatic thiol of the formula (2), the process comprising the steps of
5 halogenating the aromatic or heteroaromatic methyl sulfide of the formula (4) to give the aromatic or heteroaromatic halogenated methyl sulfide of the formula (1), and hydrolyzing the compound of the formula (1) (the fourth invention of the present application).

10 Also the present invention provides a process for preparing the aromatic or heteroaromatic disulfide of the formula (3), the process comprising the steps of halogenating the aromatic or heteroaromatic methyl sulfide of the formula (4) to give the aromatic or heteroaromatic
15 halogenated methyl sulfide of the formula (1), subsequently hydrolyzing the compound of the formula (1) and further oxidizing the hydrolyzate (the fifth invention of the present application).

In these processes, by combining said reactions,
20 the aromatic or heteroaromatic thiol of the formula (2) or the aromatic or heteroaromatic disulfide of the formula (3) is produced from the aromatic or heteroaromatic methyl sulfide of the formula (4) in one pot in a high yield.

The present invention provides a novel process
25 for preparing an aromatic or heteroaromatic thiol or

disulfide which is used for various applications as in pharmaceutical compositions, agricultural compositions, functional materials or the like and a novel process for preparing an aromatic or heteroaromatic halogenated methyl sulfide useful as the starting material for preparing said thiol or disulfide. According to the processes of the present invention, the end product can be simply prepared in high yields by halogenating, hydrolyzing and oxidizing a commercially available aromatic or heteroaromatic methyl sulfide. Thus the processes of the present invention have pronouncedly high economical and commercial values.

Best Mode for Carrying Out the Invention

The present invention will be described below in more detail with reference to Examples to which the present invention is not limited at all.

Example 1

A 1-liter, 4-necked flask equipped with a stirrer, thermometer, condenser and gas inlet tube was charged with 227.5 g (1.00 mole) of 4-chlorophenyl dichloromethyl sulfide, 50 g of water and 400 g of methanol. The mixture was heated to 70°C for 5 hours to complete hydrolysis. After completion of the reaction, water was added to separate the oil layer. Then distillation gave 136.5 g of 4-chlorothiophenol. The yield based on 4-chlorophenyl dichloromethyl sulfide was 94.5%.

Examples 2 to 42

The same procedure as in Example 1 was carried out with exception of using the aromatic or heteroaromatic halogenated methyl sulfides listed below in Tables 1, 2 and 5 3 as the starting material in place of the compound used in Example 1, whereby the corresponding aromatic or heteroaromatic thiols were produced.

Table 1

Ex.	Starting Material	Reaction Product	Yield (%)
2	2-Chlorophenyl dichloromethyl sulfide	2-Chlorothiophenol	93.8
3	2-Chlorophenyl trichloromethyl sulfide	2-Chlorothiophenol	93.5
4	2,4-Dichlorophenyl dichloromethyl sulfide	2,4-Dichlorothiophenol	94.7
5	2,5-Dichlorophenyl dichloromethyl sulfide	2,5-Dichlorothiophenol	94.6
6	3,5-Dichlorophenyl dichloromethyl sulfide	3,5-Dichlorothiophenol	93.1
7	2,6-Dichlorophenyl dichloromethyl sulfide	2,6-Dichlorothiophenol	95.0
8	4-Bromophenyl dichloromethyl sulfide	4-Bromothiophenol	93.3
9	4-Bromophenyl trichloromethyl sulfide	4-Bromothiophenol	94.5
10	1,4-Di(dichloromethylthio)benzene	1,4-Benzenedithiol	92.5
11	1,4-Di(trichloromethylthio)benzene	1,4-Benzenedithiol	92.2
12	1,2-Di(dichloromethylthio)benzene	1,2-Benzenedithiol	91.4
13	1,2-Di(dichloromethylthio)-4-butylbenzene	4-Butyl-1,2-benzenedithiol	91.8
14	4-Cyanophenyl dichloromethyl sulfide	4-Cyanothiophenol	94.5
15	2-Cyanophenyl dichloromethyl sulfide	2-Cyanothiophenol	94.6

Table 2

Ex.	Starting Material	Reaction Product	Yield (%)
16	2-Cyanophenyl trichloromethyl sulfide	2-Cyanothiophenol	95.1
17	4-Methylphenyl dichloromethyl sulfide	4-Methylthiophenol	93.8
18	4-Hydroxyphenyl dichloromethyl sulfide	4-Hydroxythiophenol	94.2
19	4-Acetylphenyl dichloromethyl sulfide	4-Acetylthiophenol	92.4
20	4-Nitrophenyl dichloromethyl sulfide	4-Nitrothiophenol	94.9
21	2-Aminophenyl dichloromethyl sulfide	2-Aminothiophenol	94.2
22	4-(Dichloromethylthio)benzoic acid	4-Mercaptobenzoic acid	93.6
23	4-(Trichloromethylthio)benzoic acid	4-Mercaptobenzoic acid	94.2
24	4-Dichloromethylbenzenesulfonic acid	4-Mercaptobenzenesulfonic acid	91.5
25	2-Cyano-3-chlorophenyl dichloromethyl sulfide	2-Cyano-3-chlorothiophenol	93.5
26	4-Carbamoylphenyl dichloromethyl sulfide	4-Carbamoylthiophenol	94.1
27	4,4'-(Dichloromethylthio)diphenyl sulfide	4,4'-Thiodibenzenethiol	91.9
28	4,4'-(Trichloromethylthio)diphenyl sulfide	4,4'-Thiodibenzenethiol	92.9
29	4,4'-(Dichloromethylthio)diphenyl ether	4,4'-Oxydibenzenethiol	92.1

Table 3

Ex.	Starting Material	Reaction Product	Yield (%)
30	4-(Phenylthio)phenyl dichloromethyl sulfide	4-(Phenylthio)benzenethiol	95.2
31	4-(Phenylsulfonyl)phenyl dichloromethyl sulfide	4-(Phenylsulfonyl)benzenethiol	92.9
32	2-Pyridyl dichloromethyl sulfide	2-Pyridinethiol	93.1
33	2-Pyridyl trichloromethyl sulfide	2-Pyridinethiol	94.1
34	4-Pyridyl dichloromethyl sulfide	4-Pyridinethiol	93.3
35	2-Thienyl dichloromethyl sulfide	2-Mercaptothiophene	92.1
36	1-Naphthyl dichloromethyl sulfide	1-Naphthalenethiol	94.5
37	1-Naphthyl trichloromethyl sulfide	1-Naphthalenethiol	95.2
38	2-Naphthyl dichloromethyl sulfide	2-Naphthalenethiol	94.8
39	2-Pyrazyl dichloromethyl sulfide	2-Mercaptopyrazine	93.6
40	4-Triazyl dichloromethyl sulfide	4-Mercaptotriazole	94.7
41	2-Oxazyl dichloromethyl sulfide	2-Mercaptooxazole	92.8
42	4-Isothiazyl dichloromethyl sulfide	4-Mercaptoisothiazole	93.1

Example 43

A 1-liter, 4-necked flask equipped with a stirrer, thermometer, condenser and gas inlet tube was charged with 227.5 g (1.00 mole) of 4-chlorophenyl dichloromethyl sulfide, 50 g of water and 500 g of methanol. The mixture was heated to 70°C for 5 hours to complete hydrolysis. After completion of the reaction, 357 g (1.05 moles) of 10% aqueous solution of hydrogen peroxide was added dropwise at 40°C over a period of 1 hour. The mixture was stirred for 1 hour at the same temperature. The precipitated crystals were filtered, giving 134.7 g of 4,4'-dichlorodiphenyl disulfide in a yield of 93.9% based on 4-chlorophenyl dichloromethyl sulfide.

Example 44

The same procedure as in Example 43 was carried out with the exception of using 2-cyanophenyl dichloromethyl sulfide in place of 4-chlorophenyl dichloromethyl sulfide as an aromatic or heteroaromatic halogenated methyl sulfide as the starting material, giving 2,2'-dicyanodiphenyl disulfide in a yield of 92.1% based on 2-cyanophenyl dichloromethyl sulfide.

Example 45

The same procedure as in Example 43 was carried out with the exception of using 4-bromophenyl trichloromethyl sulfide in place of 4-chlorophenyl

-20-

dichloromethyl sulfide as an aromatic or heteroaromatic halogenated methyl sulfide as the starting material, giving 4,4'-dibromodiphenyl disulfide in a yield of 90.3% based on 4-bromophenyl trichloromethyl sulfide.

5 Example 46

A 1-liter, 4-necked flask equipped with a stirrer, thermometer, condenser and dropping funnel was charged with 158.5 g (1.00 mole) of 4-chlorothioanisole and 300 g of monochlorobenzene. Chlorine (149.1 g, 2.1 moles) was blown into the mixture at 5°C over 2 hours. Thereafter 50 g of water and 500 g of methanol were added, and the mixture was heated at 70°C for 5 hours to complete hydrolysis. After completion of the reaction, water was added to separate the oil layer, and monochlorobenzene was distilled off. Thereafter, distillation gave 133.3 g of 4-chlorothiophenol. The yield based on 4-chlorothioanisole was 92.2%.

Examples 47 to 70

The same procedure as in Example 46 was carried out with exception of using the aromatic or heteroaromatic thioanisoles as the starting material listed below in Tables 4 and 5 in place of the compound used in Example 46, whereby the corresponding aromatic or heteroaromatic thiols were produced.

Table 4

Ex.	Starting Material	Reaction Product	Yield (%)
47	2-Chlorothiophenol	2-Chlorothiophenol	91.5
48	2,4-Dichlorothiophenol	2,4-Dichlorothiophenol	92.4
49	2,5-Dichlorothiophenol	2,5-Dichlorothiophenol	92.1
50	3,5-Dichlorothiophenol	3,5-Dichlorothiophenol	93.0
51	2,6-Dichlorothiophenol	2,6-Dichlorothiophenol	91.5
52	4-Bromothiophenol	4-Bromothiophenol	92.5
53	1,4-(Dimethylthio)benzene	1,4-Benzenedithiol	90.0
54	1,2-(Dimethylthio)benzene	1,2-Benzenedithiol	89.5
55	4-Cyanothiophenol	4-Cyanothiophenol	92.5
56	2-Cyanothiophenol	2-Cyanothiophenol	92.4
57	4-Hydroxythiophenol	4-Hydroxythiophenol	93.5
58	4-Acetylthiophenol	4-Acetylthiophenol	90.8
59	4-Nitrothiophenol	4-Nitrothiophenol	91.5

Table 5

Ex.	Starting Material	Reaction Product	Yield (%)
60	4-(Methylthio)benzoic acid	4-Mercaptobenzoic acid	92.4
61	4-Methylthiobenzenesulfonic acid	4-Mercaptobenzenesulfonic acid	89.2
62	4-Carbamoylthioanisole	4-Carbamoylthiophenol	89.8
63	4,4'-(Dimethylthio)diphenyl sulfide	4,4'-Thiodibenzenethiol	89.9
64	4,4'-(Dimethylthio)diphenyl ether	4,4'-Oxydibenzenethiol	90.4
65	4-(Phenylthio)thioanisole	4-(Phenylthio)benzenethiol	91.0
66	4-(Phenylsulfonyl)thioanisole	4-(Phenylsulfonyl)-benzenethiol	90.5
67	2-Pyridylthioanisole	2-Pyridinethiol	92.5
68	4-Pyridylthioanisole	4-Pyridinethiol	93.0
69	2-Thienylthioanisole	2-Mercaptothiophene	92.0
70	1-Naphthylthioanisole	1-Naphthalenethiol	95.1

Example 71

A 1-liter, 4-necked flask equipped with a stirrer, thermometer, condenser and dropping funnel was charged with 158.5 g (1.00 mole) of 4-chlorothioanisole and 300 g of monochlorobenzene. Chlorine (149.1 g, 2.1 moles) was blown into the mixture at 5°C over 2 hours. Thereafter 50 g of water and 500 g of methanol were added, and the mixture was heated at 70°C for 5 hours to complete hydrolysis. After completion of the reaction, 357 g (1.05 moles) of 10% aqueous solution of hydrogen peroxide was added dropwise at 40°C over a period of 1 hour. The mixture was stirred for 1 hour at the same temperature. The precipitated crystals were filtered, giving 132.0 g of 4,4'-dichlorodiphenyl disulfide in a yield of 92.0% based on 4-chlorothioanisole.

Example 72

The same procedure as in Example 71 was carried out with the exception of using 2-cyanothioanisole in place of 4-chlorothioanisole as an aromatic or heteroaromatic thioanisole as the starting material, giving 2,2'-dicyanodiphenyl disulfide in a yield of 92.1% based on 2-cyanothioanisole.

Example 73

A 1-liter, 4-necked flask equipped with a stirrer, thermometer, condenser and dropping funnel was

-24-

charged with 158.5 g (1.00 mole) of 4-chlorothioanisole and 300 g of monochlorobenzene. Chlorine (149.1 g, 2.1 moles) was blown into the mixture at 5°C over 2 hours. The reaction mixture was washed with water and dehydrated with 5 magnesium sulfate, followed by distilling off monochlorobenzene. Thereafter, distillation gave 220.3 g of 4-chlorophenyl dichloromethyl sulfide in a yield of 96.8% based on 4-chlorothioanisole.

- 25 -

The embodiments of the present invention in which an exclusive property or privilege is claimed are defined as follows:

1. A process for preparing an aromatic or heteroaromatic thiol represented by formula (2):



the process comprising hydrolyzing, in the presence of sulfuric acid and a C₁₋₄ alcohol, an aromatic or heteroaromatic halogenated methyl sulfide represented by formula (1):



wherein Ar is an aromatic or heteroaromatic ring, X is a halogen atom, m is an integer of 1 to 3 and n is 1 or 2.

2. The process according to claim 1, wherein X in the formula (1) is chlorine.
3. The process according to claim 1 or 2, wherein m in the formula (1) is 2 or 3.
4. The process according to claim 1, 2 or 3, wherein the C₁₋₄ alcohol is methanol.
5. The process according to any one of claims 1 to 4, wherein Ar in the formulas (1) and (2) is selected from the group consisting of a benzene ring, a pyridine ring, a thiazole ring, and an isothiazole ring.

- 26 -

6. A process for preparing an aromatic thiol represented by formula (2):




the process comprising hydrolyzing, in the presence of sulfuric acid and a C₁₋₄ alcohol, an aromatic halogenated methyl sulfide represented by formula (1):




wherein Ar is a benzene ring having at least one substituent selected from the group consisting of halogen, cyano, formyl, amino, carboxyl, esterified carboxyl, carbamoyl, alkyl carbonyl, nitro, sulfonic acid, alkyl, alkoxyl, hydroxyl, and phenylthio groups; X is a halogen atom; m is an integer of 1 to 3; and n is 1 or 2.

7. A process for preparing an aromatic or heteroaromatic disulfide represented by formula (3):



the process comprising the steps of hydrolyzing, in the presence of sulfuric acid and a C₁₋₄ alcohol, an aromatic or heteroaromatic halogenated methyl sulfide represented by formula (1):



and oxidizing the reaction mixture;

wherein Ar is an aromatic or heteroaromatic ring, X is a halogen atom, m is an integer of 1 to 3 and n is 1 or 2.

8. The process according to claim 7, wherein X in formula (1) is chlorine.

9. The process according to claim 7 or 8, wherein m in formula (1) is 2 or 3.

- 27 -

10. The process according to claim 7, 8 or 9, wherein the C₁₋₄ alcohol is methanol.

11. The process according to any one of claims 7 to 10, wherein Ar in formulas (1) and (3) is selected from the group consisting of a benzene ring, a pyridine ring, a thiazole ring, and an isothiazole ring.

12. A process for preparing an aromatic disulfide represented by formula (3):



the process comprising the steps of hydrolyzing, in the presence of sulfuric acid and a C₁₋₄ alcohol, an aromatic halogenated methyl sulfide represented by formula (1):



and oxidizing the reaction mixture,

wherein Ar is a benzene ring having at least one substituent selected from the group consisting of halogen, cyano, formyl, amino, carboxyl, esterified carboxyl, carbamoyl, alkyl carbonyl, nitro, sulfonic acid, alkyl, alkoxy, hydroxyl, and phenylthio groups; X is a halogen atom; m is an integer of 1 to 3; and n is 1 or 2.

13. A process for preparing an aromatic or heteroaromatic thiol represented by formula (2):



the process comprising the steps of halogenating an aromatic or heteroaromatic methyl sulfide represented by formula (4):



- 28 -

to give an aromatic or heteroaromatic halogenated methyl sulfide represented by formula (1):



and hydrolyzing, in the presence of sulfuric acid and a C₁₋₄ alcohol, the compound of formula (1);

wherein Ar is an aromatic or heteroaromatic ring, X is a halogen atom, m is an integer of 1 to 3 and n is 1 or 2.

14. A process for preparing an aromatic or heteroaromatic disulfide represented by formula (3):



the process comprising the steps of halogenating an aromatic or heteroaromatic methyl sulfide represented by formula (4):



to give an aromatic or heteroaromatic halogenated methyl sulfide represented by formula (1):



hydrolyzing, in the presence of sulfuric acid and a C₁₋₄ alcohol, the compound of formula (1) and oxidizing the hydrolyzate;

wherein Ar is an aromatic or heteroaromatic ring, X is a halogen atom, m is an integer of 1 to 3 and n is 1 or 2.